



(86) Date de dépôt PCT/PCT Filing Date: 1999/08/04  
 (87) Date publication PCT/PCT Publication Date: 2000/02/17  
 (45) Date de délivrance/Issue Date: 2010/11/23  
 (85) Entrée phase nationale/National Entry: 2001/02/05  
 (86) N° demande PCT/PCT Application No.: JP 1999/004197  
 (87) N° publication PCT/PCT Publication No.: 2000/008170  
 (30) Priorités/Priorities: 1998/08/05 (JP10/232382);  
 1999/06/28 (JP11/182362)

(51) Cl.Int./Int.Cl. *C12N 15/53* (2006.01),  
*C12N 1/21* (2006.01), *C12N 15/54* (2006.01),  
*C12N 9/00* (2006.01), *C12N 9/02* (2006.01),  
*C12N 9/04* (2006.01), *C12N 9/10* (2006.01),  
*C12P 13/04* (2006.01), *C12P 13/14* (2006.01)

(72) Inventeurs/Inventors:  
 FUJII, TADASHI, JP;  
 NARITA, TAKAO, JP;  
 NAKATA, KUNIHO, JP;  
 AGEMATU, HITOSI, JP;  
 TSUNEKAWA, HIROSHI, JP;  
 ...

(73) Propriétaire/Owner:

(54) Titre : GENE PARTICIPANT A LA PRODUCTION D'ACIDE HOMOGLUTAMIQUE, ET UTILISATION ASSOCIEE  
 (54) Title: GENE PARTICIPATING IN THE PRODUCTION OF HOMOGLUTAMIC ACID AND ITS USE

(57) Abrégé/Abstract:

Provided are an isolated gene capable of participating in the production of L-homoglutamic acid, and a production system of L-homoglutamic acid by using this gene. The gene is derived from the genome of Flavobacterium lutescens.

(72) Inventeurs(suite)/Inventors(continued): ISSHIKI, KUNIO, JP; YOSHIOKA, TAKEO, JP

(73) Propriétaires(suite)/Owners(continued):MERCIAN CORPORATION, JP

(74) Agent: SMART & BIGGAR

ABSTRACT

Provided are an isolated gene capable of participating in the production of L-homoglutamic acid, and a production system of L-homoglutamic acid by using this gene. The gene is derived from the genome of Flavobacterium lutescens.

## DESCRIPTION

GENE PARTICIPATING IN THE PRODUCTION OF  
HOMOGLUTAMIC ACID AND ITS USE

5

**Technical Field**

This invention relates to gene manipulation, and more specifically, relates to a DNA containing a gene participating in the production of L-homoglutamic acid (also referred to as L-2-amino-  
10 adipic acid or L- $\alpha$ -aminoadipic acid), and a production system of L-homoglutamic acid (hereinafter, merely referred to as homoglutamic acid) using it.

**Background Art**

15 Homoglutamic acid is found widely in organisms such as plants including Cholera vibrio as a bacterium and corn (Zea mays), the embryos of frogs. Homoglutamic acid acts as an intermediate of lysine biosynthesis in fungi, etc. and as a precursor in biosynthesis of  $\beta$ -lactam antibiotics. Further, homoglutamic acid is also useful as a  
20 synthetic intermediate of various medicines including methotrexate derivatives (WO 92/09436).


Since preparation of homoglutamic acid by chemical synthesis needs optical resolution and multistage reaction, it is not a useful means from the aspect of costs. On the other hand, a process  
25 of preparing homoglutamic acid from L-lysine using a microorganism belonging to the genus Agrobacterium, Klebsiella, Alcaligenes, Brevibacterium or Bacillus is known (Japanese Laid-open Patent Publication No. 6-181787). Part of the present inventors also proposed a process of preparing homoglutamic acid from L-lysine using a micro-  
30 organism belonging to the genus Flavobacterium (WO 96/ 31616). However, even in the process using such a microorganism, a process

capable of preparing homoglutamic acid more efficiently is desired earnestly.

Thus, the present inventors aimed to reinforce the production system of homoglutamic acid in any of the above microorganisms, for example by gene manipulation. When a review of helpful information is made on the manipulation, for example, as part of researches of biosynthetic pathway of cephamycin C, are confirmed the presence of lysine-6-aminotransferase and L- $\Delta^1$ -piperidine-carboxylate dehydrogenase participating in conversion from L-lysine to  $\alpha$ -amino-adipic acid (or homoglutamic acid) of Streptomyces clavuligerus as a cephamycin C-producing actinomycetes, and as to the former, the presence position of the gene encoding the enzyme, etc. (Fuente et al., Biochem. J. (1997) 327, 59-64).

As to Flavobacterium lutescens (which was re-identified from Flavobacterium fuscum) IFO 3084 used in bioassay of L-lysine, it is known that 2-oxoglutarate 6-aminotransferase [or lysine 6-aminotransferase (hereinafter also referred to as LAT)] catalyzing the following pathway is present (Soda et al., Biochemistry 7 (1968), 4102-4109, ibid. 4110-4119).

20



In the above bioassay, the absorbance of the product obtained by reacting piperidine-6-carboxylic acid (hereinafter, also

30

referred to as P6C) with o-aminobenzaldehyde is measured. In another bioassay of L-lysine, the L-lysine 6-dehydrogenase activity of Agrobacterium tumefaciens is utilized (Misono et al., J. Biochem. (Tokyo) 105 (1989), 1002-1008).

5           The above IFO 3084 strain is commonly used in bioassay of L-lysine as mentioned above, and its use method is also established. Therefore, if the IFO 3084 strain had a gene encoding a protein having P6C (or, the 2-aminoadipic acid semialdehyde which is said to be in a quantitatively equilibrium state with P6C in a living body) de-  
10 hydrogenase (hereinafter, also merely referred to as dehydrogenase) activity, in addition to LAT, the strain would be a candidate bacterium for gene cloning meeting the object of the present invention, namely the object to provide a gene participating in the production of homo-  
glutamic acid.

15

#### Disclosure of Invention

The present inventors have tried cloning of the lysine-6-aminotransferase (LAT) gene (lat) of Flavobacterium lutescens and, according to circumstances, a gene encoding a protein having de-  
20 hydrogenase activity on P6C of the bacterium. However, as cloning methods regularly used for such a case, a method of obtaining a targeted gene from DNA consensus sequences between amino-  
transferases of other bacteria, and a method utilizing information obtained from the result of amino acid sequencing of a purified pro-  
25 tein, and the like have all failed in their early researches.

However, unexpectedly, they have found that when the host-vector system finally selected by the inventor is used, a gene at least capable of participating in the production of homoglutamic acid, more specifically a gene encoding a protein having dehydrogenase  
30 activity on P6C can be cloned by shotgun cloning. They have also

found that a modifier having a certain homology (or identity) to the gene also functions similarly.

On the other hand, the above Soda et al., *Biochemistry* 7 (1968), 4110-4119 discloses a process of obtaining crystalline LAT of a molecular weight of 116,000 from *Achromobacter liquidum* (= *Flavobacterium lutescence*), and Yagi et al., *J. Biochem.* 87 (1980), 1395-1402 discloses that LAT from *Flavobacterium lutescens* is composed of four nonidentical subunits of A, B1, B2 and C. Their early researches of cloning a gene encoding a protein having LAT activity utilizing the information obtained from the amino acid sequencing of the purified LAT protein, based on these descriptions, have failed. However, using a process entirely different from the processes described in these prior art references, the present inventors have purified proteins having LAT activity from *Flavobacterium lutescens*, have determined the amino acid sequences of the obtained proteins, and have cloned the objective genes utilizing these sequence informations, and as a result they have succeeded in cloning a gene encoding LAT (lat). The invention is based on the above findings.

Thus, according to the invention is provided an isolated pure DNA containing a gene participating in the production of homoglutamic acid which gene can be obtained from a bacterium belonging to the genus *Flavobacterium lutescens*, or a modifier which hybridizes with the gene under a stringent condition and has a function capable of recovering the homoglutamic acid-producing ability of a mutant which lacks the producing ability.

More specifically, the gene participating in the production of homoglutamic acid is a DNA encoding partly or wholly at least one protein selected from the group consisting of a protein having LAT activity and a protein having dehydrogenase activity, or a modifier thereof.

67566-1434

5

The invention also relates to an autonomously replicative or integration replicative recombinant plasmid carrying the DNA, and a transformant obtained by transformation with the recombinant plasmid, and a process  
5 of producing homoglutamic acid using the transformant.

In one aspect, the invention relates to an isolated pure DNA containing a gene participating in the production of L-homoglutamic acid, derived from a bacterium belonging to Flavobacterium lutescens, or a DNA molecule  
10 which hybridizes with the complement of the gene under a stringent condition and has a function of recovering the L-homoglutamic acid-producing ability of a mutant of Flavobacterium lutescens which lacks the L-homoglutamic acid-producing ability, wherein the gene participating in  
15 the production of L-homoglutamic acid is a DNA encoding at least one protein selected from the group consisting of (a) a protein having L-lysine: 2-oxoglutaric acid 6-aminotransferase (LAT) activity, and (b) a protein having L- $\Delta^1$ -piperidine-6-carboxylic acid (P6C) dehydrogenase  
20 activity, wherein the DNA encoding the protein having the LAT activity (a) contains a continuous base sequence from base 801 to base 2276 of SEQ ID NO: 1 and the DNA encoding the protein having the P6C dehydrogenase activity (b) contains a continuous base sequence from base 2855 to  
25 base 4384 of SEQ ID NO: 3, and wherein the hybridization under the stringent condition involves washing at 60°C in 0.5 x SSC.

In another aspect, the invention relates to an autonomously replicative or integration replicative  
30 recombinant plasmid carrying the DNA as defined herein.



67566-1434

5a

In another aspect, the invention relates to a transformant obtained by transforming a bacterium belonging to the genus Flavobacterium as a host with the recombinant plasmid as defined herein.

5 In another aspect, the invention relates to a process for producing L-homoglutamic acid, which comprises: culturing in a medium a transformant obtained by transformation with the recombinant plasmid as defined herein; during or after the culture, contacting the grown  
10 transformant with L-lysine or L- $\Delta^1$ -1-piperidine-6-carboxylic acid to convert it to L-homoglutamic acid; and recovering the thus produced L-homoglutamic acid.

#### **Brief Description of Drawings**

Figure 1 is a drawing showing the analytical  
15 results by thin layer chromatography of homoglutamic acid production by mutants of F. lutescens. St is standard homoglutamic acid (HG), Lanes 1 to 4, Lanes 5 to 7, Lanes 8 to 10, Lane 11, and Lanes 12 and 13 show the analytical results of the first mutants, the second mutants,  
20 the third mutants, the wild type strain and the first mutants having plasmid pCF704, respectively.

Figure 2 is a graph showing the lysine  
6-aminotransferase (LAT) activity of mutants of  
F. lutescens. Wild, 1st, 2nd and 3rd show the  
25 LAT activities of the wild type strain, the first mutant, the second mutant and the third mutant, respectively.

67566-1434

5b

Figure 3 shows the results of analyses by thin layer chromatography showing complementarity of homoglutamic acid productivity of homoglutamic acid productivity-lacking mutants by plasmid pCF213.

5 HG and Lys show the moved position of homoglutamic acid and the moved position of L-lysine, and St; 1st pCF213, 2nd pCF213 and 3rd pCF213; Wild pCF213 and Wild pCF704; 1st pCF704 and 2nd pCF704; and 1st pCF111 are the results of TLC analyses of homoglutamic acid standard substance;

10 culture broths of the first, the second and the third mutants having pCF213, respectively; culture broths of wild type strains having pCF213 and pCF704, respectively; culture broths of the first and second mutants having pCF704,

respectively; and culture broths of the first mutant having pCF 111; respectively.

Figure 4 is a graph showing the productivity of homoglutamic acid with time lapse of *F. lutescens* IFO 3084 (pCF213) (in the drawing, represented by pCF213) and *F. lutescens* IFO 3084 (pCF704) (in the drawing, represented by pCF704).

Figure 5 is a graph showing the presence position ORF found based on the base sequence of the pCF213 insert region.

Figure 6 is a graph showing relations between the elution fractions by the MonoQ HR5/5 column treatment in 3(6) of Example 2 and the relative LAT activities.

Figure 7 is a photograph in place of a drawing showing the results of Native PAGE (A) and SDS-PAGE (B) of the LAT active fractions using Multigel 4/20 and 10/20, in 3(7) of Example 2. In the drawing, M is a molecular weight marker, C represents the ultrafiltrate obtained in 3 (5) of Example 2, and the figures represent the respective fraction numbers.

Figure 8 is a graph showing relative LAT activities in homoglutamic acid productivity-lacking mutants and wild type strains by various plasmids.

Figure 9 is a graph showing the productivity of homoglutamic acid with time lapse of *F. lutescens* IFO 3084 transformed with various plasmids.

## 25 **Specific Embodiments of Invention**

As to origins of genes according to the invention, any strains of *Flavobacterium lutescens* (hereinafter, also referred to as *F. lutescens*) including spontaneous mutants so long as they can provide a gene participating in the production of homoglutamic acid which gene can be expressed, for example, in *F. lutescens* as a host.

However, mentioned as preferred is the IFO 3084 strain which is easy to obtain and whose suitable handling conditions such as culture are established.

The gene participating in the production of homoglutamic acid in the invention means any gene capable of participating in the two-stage conversion system from L-lysine to homoglutamic acid via P6C or 2-aminoadipic acid-6-semialdehyde which is chemically in an equilibrium relation with P6C (the former stage: LAT activity, the latter stage: dehydrogenase activity). First of all, as specific examples of genes encoding a protein having dehydrogenase activity which is the latter conversion system, there can be mentioned genes which can be obtained using the host-vector system established by the present inventors based on the following strategy.

Establishment of a suitable host-vector system of F. lutescens is necessary for carrying out the gene manipulation of F. lutescens, but therefor it is needed to solve the following three problems.

(1) Obtain a replicon which can autonomously replicate in F. lutescens.

(2) Obtain a drug resistance marker which can be expressed and function in F. lutescens.

(3) Establish a method of introducing a DNA into F. lutescens.

Fortunately, the above problems (1) and (2) could be solved by finding that pBBR122, lately put on the market by Mo Bi Tec corporation, which autonomously replicates in a wide range of Gram-negative bacteria and has kanamycin and chloramphenicol resistance can be used. For solution of the above problem (3), first, it becomes a prerequisite that a method of introducing the plasmid pBBR122 into F. lutescens is established. However, examination was

made based on the method of DNA introduction into E. coli by the electroporation method, as a result a colony of F. lutescens grew in an L plate containing 20 µg/ml kanamycin, and by liquid culturing this and extracting plasmids by the alkali SDS method, it was confirmed  
5 that pBBR122 was stably held in F. lutescens. Thus, the problem (3) was also solved. As to this host-vector system, it has itself been known that when other bacteria were used as a host, (a) transformation efficiency is very high and (b) a DNA fragment of a suitable size can be inserted into pBBR122 (J. Bac. 178 (1996), 1053- 1060), but it  
10 was revealed that the above (a) and (b) are possible also in F. lutescens, and further it was made possible to amplify the obtained gene in F. lutescens, and more over, it was also made possible to obtain a gene encoding a protein having dehydrogenase activity on P6C by shotgun cloning. For facilitating the operation, pCF704 in which the  
15 multicloning site of pUC19 was introduced in place of the chloramphenicol resistance gene of pBBR122 was prepared, and this was then used as a vector.

Then, in order to establish a system for evaluating an obtained and amplified gene, mutation was induced in F. lutescens  
20 IFO 3084 with N-methyl-N'-nitro-N-nitrosoguanidine (NTG), and screening was made using an MEM plate (pH 7.0) containing eosin Y.

Thus, the first mutant not producing homoglutamic acid at all, and the second and third mutants only slightly producing homoglutamic acid were obtained. In the first mutant not producing  
25 homoglutamic acid at all, lat activity equal to the wild type strain was confirmed, and in the second and third mutants only slightly producing homoglutamic acid, only slight lat activity was confirmed. Namely, there is a possibility that the first mutant is suffering some injuries to gene(s) other than lat participating in the production of homo-  
30 glutamic acid, and on the other hand the second and third mutants

are suffering some injuries at least to lat.

Then, the genome DNA of the wild type strain was partly digested with SauIII<sup>AI</sup>, and the 6-8 kbp fragments were inserted into the BamHI site of pCF704, respectively, to prepare a DNA library.

5 These plasmids were introduced into the first, second and third mutants, respectively, and strains which recovered homoglutamic acid-producing ability were screened. In this occasion, a method was used which comprises collecting colonies blackened in a MEM plate (pH 7.0) containing eosin Y, used for the screening of the mutants,  
10 and confirming homoglutamic acid-producing ability thereof by TLC. As representable ones of these mutants, the second mutant (Flavo-  
bacterium lutescens 2nd mutant) was deposited on July 6, 1998 with National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, and has been assigned an acces-  
15 sion number of FERM P-16874, and the first mutant (Flavobacterium  
lutescens 1st mutant) was deposited on June 10, 1999 with the Institute, and has been assigned an accession number of FERM P-17419, and these strains are kept there. These FERM P-16874 strain and FERM P-17419 strain were transferred on July 26, 1999 on their  
20 deposition to the international deposition authority on Budapest Treaty in the Institute, and have been assigned accession numbers of FERM BP-6798 and FERM BP-6799, respectively.

As a result, a strain having a plasmid complementing the productivity of homoglutamic acid of the first mutant and a strain  
25 having a plasmid partly complementing the productivity of homoglutamic acid of the second mutant were obtained. However, the plasmids of these strains, particularly plasmid of the strain complementing the second mutant were liable to be deleted, and further screening for obtaining a stable plasmid has been needed. As a result  
30 of DNA fragment analysis with restriction enzyme treatment, it was

revealed that the thus obtained plasmid designated pCF111 which complements the first mutant and partly complements the second mutant and the plasmid designated pCF213 were apparently quite the same plasmid.

5           On the other hand, pCF111 and pCF213 were re-transformed into the first, second and third mutants, respectively, and homoglutamic acid-producing ability was checked. As a result, both plasmids complemented the first mutant, but only partly complemented the second and third mutant.

10           Based on the complementation test, it was revealed that in a plasmid sufficiently recovering the homoglutamic acid-producing ability of a homoglutamic acid productivity-lacking mutant, a gene participating at least in the production of homoglutamic acid, more specifically some gene other than lat is present.

15           Thus, not limited thereto, but as one of the "genes participating in the production of homoglutamic acid", there can be mentioned a gene which is contained in the insert part of plasmid pCF213 and encoding a protein having dehydrogenase activity, the protein comprising the amino acid sequence shown in SEQ ID NO: 4. For example, this gene is present in the sequence  
20 shown in SEQ ID NO: 3, in the continuous base sequence from base 2077 to base 4578, particularly in the continuous base sequence from base 2855 to base 4384.

On the other hand, a gene participating in the former conversion, namely encoding a protein having LAT activity according to the invention can be cloned as follows.

25           F. lutescens is cultured under a certain culture condition, the obtained strain is fractured, the fracture dispersion is centrifuged to remove the fractured cells, and from the thus obtained cell extract, the desired protein is isolated and purified by ultracentrifugation treatment, ammonium sulfate precipitation, desalting, ion exchange  
30 column chromatography, affinity column chromatography, ultra-filtration, electrophoresis, etc.

From the analytical results of the N-terminus amino acid

67566-1434

11

sequence of the purified protein, DNA primers are designed, and PCR is carried out on the genome DNA of *F. lutescens* (IFO 3084) strain. Based on the DNA fragment amplified by PCR further PCR is carried out, and thereby the neighborhood region of both outer sides of the  
5 DNA fragment is obtained. Thus, a DNA encoding the desired protein of the invention is obtained.

Thus, it becomes possible to provide a DNA encoding a protein having LAT activity as another gene participating in the production of L-homoglutamic acid. Namely, as another gene of the  
10 invention, there can, for example, be mentioned one having a sequence composing the coding region of the base sequence of SEQ ID NO: 1, i.e., in the continuous base sequence from 801 to base 2276 of SEQ ID NO: 1. Practically, the gene may be the continuous sequence from base 545 to base 2658 of SEQ ID NO: 1. The N-terminus of the corresponding purified protein  
15 consisting of the amino acid sequence as shown in SEQ ID NO: 2 is Ser as shown in SEQ ID NO: 1, but it is considered that N-terminal Met is processed after translation.

Further, the DNA containing a gene participating in the production of homoglutamic acid according to the invention includes a DNA containing at least one of the gene encoding a protein  
20 having dehydrogenase activity and the gene encoding a protein having LAT activity.

In addition, the gene referred to in the invention also includes a DNA molecule of the above-mentioned genes, which has a base sequence hybridizing with the complement of one of the genes under a certain hybridization  
25 condition, for example, washing under a stringent condition, at 60°C in 2 x SSC (in standard citric acid saline), preferably at 60°C in 0.5 x SSC, particularly preferably at 60°C in 0.2 x SSC, and has a function of recovering the homoglutamic acid-producing ability of a mutant of *F. lutescens* lacking the producing ability.

30 More specifically, a modifier of the gene encoding the protein having dehydrogenase activity is one showing at least 70 % of identity with the base sequence of from base 2855 to base 4384 in SEQ ID NO:



3, and a modifier of a gene encoding a protein having LAT activity is one showing at least 50 %, preferably 70 %, more preferably 95 % of identity with the base sequence of from base 545 to base 2658 (coding region) in SEQ ID NO: 1.

5           Such modifiers include one wherein base(s) is/are removed or added or part of the bases is replaced with other base(s), at the 5'-terminus or 3'-terminus or halfway of one of both the above sequences. The modifier wherein part of the bases is replaced with other base(s) also includes a modifier which encodes the same protein  
10 but has a base sequence different from those of both the above genes because of degeneracy of genetic code.

          It is recommended to make the substitution of base other than substitution followed by degeneracy of genetic code, considering estimated amino acid sequences encoded by both the above genes, so  
15 as to have a similar shape as the whole of protein, based on similarity of the side chain of each amino acid, for example hydrophobicity, hydrophilicity, charge, size, etc. Thus, a modifier having a function equal to the function of one of both the above genes, namely a function capable of recovering the homoglutamic acid-producing ability of  
20 a mutant of *F. lutescens* which lacks the producing ability will be obtained in a considerably high probability.

          The modifier according to the invention can be synthesized using a nucleic acid synthesizer or prepared by per se known point mutagenesis or site-directed mutagenesis, considering the base  
25 sequences of both the above genes or estimated amino acid sequences encoded by them.

          According to the invention, a recombinant plasmid carrying the above gene or modifier can also be provided. Such a plasmid can be autonomously replicative one containing, besides the above gene or  
30 modifier, an autonomously replicative sequence, a promoter se-

quence, a terminator sequence, a drug resistance gene, etc. Further, the plasmid can be integration type plasmid containing a sequence homologous to a certain region of the genome of the host intended to be used. As an example of the autonomously replicative recombinant  
5 plasmid carrying a DNA containing a gene encoding a protein having dehydrogenase activity, there can be mentioned a plasmid pBBR122, or one comprising plasmid pBBR122 having inserted in a certain site thereof a multicloning site or having substituted a multicloning site for the certain site or region and having inserted the above gene or  
10 modifier using the multicloning site. As specific examples of such plasmids, there can be mentioned ones designated plasmids pCF111 and pCF213 in the specification. pCF213 can be obtained by a per se known plasmid isolation method from Flavobacterium lutescens IFO 3084 (pCF213) which was deposited on March 11, 1998 with National  
15 Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, and has been assigned an accession number of FERM P-16699, and then transferred to international deposition on Budapest Treaty, and has been assigned an accession number of FERM BP-6797. A recombinant plasmid carrying a DNA containing a  
20 gene encoding a protein having LAT activity and a recombinant plasmid carrying a DNA containing both genes can also be constructed in the same manner as in the pCF213.

According to the invention, there can further also be provided a transformant obtained by transforming a bacterium  
25 belonging to the genus Flavobacterium as a host with the above recombinant plasmid. As the host bacterium belonging to the genus Flavobacterium, any strain of any species can be used so long as it meets the object of the invention, but as preferred ones, there can be mentioned F. lutescens IFO 3084 and F. lutescens SP.7-1 (FERM  
30 BP-5457).

Thus, as a specific example of the above transformant, there can be mentioned one obtained by transforming F. lutescens IFO 3084 or F. lutescens SP.7-1 with pCF213, and F. lutescens IFO 3084 (pCF213) is deposited as the FERM BP-6797 with the international deposition authority of National Institute of Bioscience and Human Technology.

According to the invention, a process of producing homoglutamic acid using the transformant is also provided. In the process, the transformant in a medium grown by culture is contacted with L-lysine or in some case P6C (or 2-aminoadipic 6-semialdehyde) as a starting material, or the starting material is contacted with a grown transformant or treated cells thereof (e.g., cells treated with an organic solvent, a cell extract, immobilized treated cells) to convert the starting material to homoglutamic acid.

As carbon sources of the medium, any carbon sources can be used so long as they are utilizable by the transformant, and when F. lutescens is used as a host, there can, for example, be used saccharides such as glucose, fructose, sucrose and dextrin, sugar alcohols such as glycerol and sorbitol, and organic acids such as fumaric acid and citric acid, and it is desirable that the addition amount of these carbon sources is, usually, on the order of 0.1 to 10 % by weight (hereinafter, abbreviated as %).

As nitrogen sources of the medium, there can, for example, be used ammonium salts of inorganic acids such as ammonium chloride, ammonium sulfate and ammonium phosphate, ammonium salts of organic acids such as ammonium fumarate and ammonium citrate, and further natural nitrogen sources such as meat extract, yeast extract, corn steep liquor and casein hydrolyzate, and it is desirable that the addition amount of these nitrogen sources is, usually, on the order of 0.1 to 10 %.

As inorganic salts, there can, for example, be used alkaline metal salts of phosphoric acid such as potassium phosphate and sodium phosphate, alkaline metal chlorides such as potassium chloride and sodium chloride, and metal salts of sulfuric acid such as magnesium sulfate and ferrous sulfate, and it is desirable that the addition amount of these inorganic salts is, usually, on the order of 0.001 to 1 %.

Among them, liquid culture using a usual growth medium for bacteria is preferred, and glucose, maltose, starch, etc., as carbon sources and ammonium sulfate, peptone, yeast extract, soybean meal, etc., as nitrogen sources are particularly effective. In addition, potassium phosphate, magnesium sulfate, table salt, etc., are usually used as inorganic salts.

It is recommended that the culture of the microorganism is carried out in such a medium at 20 to 40°C, preferably 28 to 37°C and at a pH of 5 to 9, preferably 6 to 8 under an aerobic condition.

The contact during the culture of the grown transformant with the starting material is carried out by previously adding the starting material in the medium or appropriately adding the starting material during the culture. The contact can also be carried out, after completion of the culture, by stirring or shaking the collected cells or treated cells and the starting material in a medium or a suitable buffer, if necessary with addition of suitable coenzymes, etc., in a reactor, or by flowing a starting material-containing matter onto immobilized cells.

The case where the transformant and L-lysine are contacted during the culture is taken as an example, and it is more specifically described below. The transformant is inoculated into a medium and cultured, for example, at 20 to 40°C for 12 to 120 hours to obtain a culture broth of the strain containing  $10^6$  to  $10^{10}$  microor-

ganisms as the transformant per ml. The starting material L-lysine as a solution in water or an auxiliary solvent or L-lysine as such without being dissolved is added so that the final concentration may usually be 0.5 to 30 mg/ml, and reaction is carried out usually at 20 to 5 40°C for 18 hours to 7 days, preferably 18 hours to 5 days. Then, homoglutamic acid can be obtained by ordinary purification methods, for example, various ion exchange chromatography using cation exchange resins, anion exchange resins, etc., adsorption chromatography using HP20, etc., precipitation or crystallization utilizing solvents 10 and temperature, and the like.

The shape and addition time of L-lysine to be added is not particularly limited, but preferably L-lysine is used as monohydrochloride in view of solubility, and it can be added at the start of culture or during the culture, e.g. in 1st to 5th day.

15 According to the invention is provided a DNA containing a gene participating in the production of homoglutamic acid which gene converts L-lysine to homoglutamic acid. This DNA is useful in a microbiological production process of homoglutamic acid. According to the invention are also provided a process of producing homo- 20 glutamic acid by a transformant capable of producing homoglutamic acid efficiently, and its use.

Hereinafter, the invention is further detailedly described by specific examples. These specific examples are provided for facilitating the understanding of the invention, and it is not intended 25 to restrict the invention to them.

### Example 1

Cloning of a gene encoding a protein having dehydrogenase activity, etc.

1. Obtention of a homoglutamic acid-not producing strain

F. lutescens IFO 3084 strain was inoculated into 3 ml of L medium (1.0 % polypeptone, 0.5 % yeast extract, 0.5 % NaCl, 0.1 % glucose, pH 7.2), and shaking cultured at 32°C overnight. 100 µl of the culture broth as an inoculum was inoculated into 50 ml of L medium, and shaking cultured at 32°C for 4.5 hours. The cells were collected from this culture broth by centrifugation of 5,000 × g for 10 minutes, washed once with 0.2 M phosphate buffer (pH6.0), and suspended in 6 ml of 0.2 M phosphate buffer (pH6.0). 50 µl of 80 mg/ml NTG was added to this cell suspension, and shaking culture was carried out at 32°C for 20 minutes. Cells collected from this culture broth were washed once with 0.2 M phosphate buffer (pH 6.0), and the whole amount was inoculated into 50 ml of L medium and shaking cultured at 32°C overnight. 500 µl portions of this culture broth were poured, respectively, 500 µl portions of 60 % glycerol solution were added, and the mixtures were well mixed, respectively, and then freeze stored at -70°C. The freeze stored mixtures are referred to as mutant storage suspensions.

This mutant storage suspension was 10<sup>6</sup>-fold diluted with 0.85 % NaCl, and 100 µl portions of the dilution were smeared on MEM agar media (0.5 % polypeptone, 0.2 % yeast extract, 1.0 % lysine-HCl, 0.006 % Methylene Blue, 0.04 % eosin Y and 1.5 % agar, pH 7.2) in 8-cm Petri dishes, and culture was carried out at 32°C for 3 days. White colonies among the grown colonies were inoculated into 1 ml portions of a screening medium (1.0 % polypeptone, 0.2 % yeast extract, 1.0 % lysine-HCl, pH 7.2), and shaking cultured at 32°C for 2 days. 3 µl of each culture was transferred to a silica gel TLC plate, and dried. This plate was developed with a solvent system consisting of butanol, acetic acid and water (3:1:1), and subjected to ninhydrin coloring, and thereby each lane was checked for the presence or

absence of homoglutamic acid. Thus, from the mutants were separated the first mutant (FERM BP-6799) not producing homoglutamic acid at all, and the second mutant (FERM BP-6798) and the third mutant producing just a bit amount of homoglutamic acid. The results obtained by checking these mutants for the ability of conversion of from L-lysine to homoglutamic acid (or productivity of homoglutamic acid) by TLC analysis are shown in Figure 1. In Figure 1, homoglutamic acid is represented by HG (this is also the case with other drawings). The results of assay of LAT activity on these mutants are shown in Figure 2.

## 2. Construction of a host-vector system and a transformation system

*F. lutescens* IFO 3084 strain was inoculated into 3 ml of L medium, and shaking cultured at 32°C overnight. 100 µl of the culture broth as an inoculum was inoculated into 50 ml of L medium, and shaking cultured at 32°C for 4.5 hours. The cells were collected from this culture broth by centrifugation of 5,000 × g for 10 minutes, washed once with 10 % glycerol solution, and suspended in 3 ml of 10 % glycerol solution. 200 µl portions of this suspension were poured, and freeze stored at -70°C. The freeze stored suspensions are referred to as Electrocell storage suspensions. This storage suspension was thawed on ice, and 1 µl of a solution of 200 µg/ml of Broad Host Range Vector pBBR122 (Mo Bi Tec incorporation) in TE was added. The mixture was put in 0.2-cm Electrocuvette (BIORAD incorporation), electric pulse was once given under a condition of 2.4 kV, 200 Ω and 25 µF using Gene Pulser II (BIORAD incorporation). Then the cells were put in a Falcon tube, 1 ml of ice-cooled L medium was added, and shaking culture was carried out at 32°C for 2 hours. The culture broth was smeared on L agar medium (1.0 % polypeptone, 0.5 % yeast extract, 0.5 % NaCl, 0.1 % glucose, 1.5 % agar, pH 7.2) containing 20 µg/ml kanamycin, and cultured at 32°C for 3 days. A transformant of

a number of  $2.4 \times 10^5$  was obtained.

### 3. Construction of a plasmid pCF704

A primer having an EcoRI site and a primer having an NcoI site were synthesized (Pharmacia incorporation), and the  
5 muticloning site and 95 bp of its neighborhood region of pUC18 were amplified, using Taq polymerase (Pharmacia incorporation) and PCR Thermal Cycler PERSONAL (Takara company). This DNA fragment was digested with restriction enzymes EcoRI and NcoI, and the  
10 digested product was ligated to the EcoRI and NcoI sites of pBBR122 using Ligation Kit version 2 (Takara company). An E. coli competent cell JM109 (Takara company) was transformed with this reaction mixture, and from the resulting transformant, a plasmid pCF704 was prepared using QIAGEN Plasmid Midi Kit.

### 4. Construction of a plasmid pCF213

15 The genome DNA of F. lutescens IFO 3084 strain was extracted and purified according to QIAGEN Blood and Cell Culture DNA Kit. This genome DNA was partly decomposed with a restriction enzyme SauIIAI, and the resulting 6 to 8 kbp fragments were cut out from agarose gel, and DNAs were recovered and purified  
20 using Ultrafree C3 Unit 0.45  $\mu\text{m}$  (Millipore corporation) and dissolved in TE solution. The resulting solution is referred to as Insert DNA solution. On the other hand, pCF704 was digested with a restriction enzyme BamHI, and the digest was dephosphorylated with alkaline phosphatase. The resulting digest and Insert DNA solution were  
25 subjected to ligation reaction using Ligation Kit version 2 (Takara company), and the reaction mixture was used as a DNA library.

This DNA library was added to the Electrocell storage suspension of the second mutant, and electric pulse was given. The resulting cells were put in a Falcon tube, 1 ml of ice-cooled L medium  
30 was added, and shaking culture was carried out at 32°C for 2 hours.



The whole amount of this culture broth was inoculated into 50 ml of L medium containing 20 µg/ml kanamycin, and shaking culture was carried out at 32°C for 2 days. 500 µl portions of the culture broth were poured, respectively, and 500 µl portions of 60 % glycerol solution were added and well mixed, respectively, and the mixtures were freeze stored at -70°C. The freeze stored mixtures are referred to as complementary strain storage suspensions.

This complementary strain storage suspension was  $10^3$  - fold diluted with 0.85 % NaCl, and 100 µl portions of the dilution were smeared on MEM agar media of pH 7.0 (0.5 % polypeptone, 0.2 % yeast extract, 1.0 % lysine-HCl, 0.006 % Methylene Blue, 0.04 % eosin Y and 1.5 % agar, pH 7.0) in 8-cm Petri dishes, and culture was carried out at 32°C for 3 days. The black parts of the cells grown on the whole surfaces are referred to as complementary strain mixture cells. The respective complementary strain mixture cells were inoculated into 3 ml portions of the screening medium, and shaking cultured at 32°C for 2 days. 3 µl portions of each of the culture broths were added dropwise on each lane of a silica gel TLC plate, and dried. This plate was developed with a solvent system consisting of butanol, acetic acid and water (3:1:1), and subjected to ninhydrin coloring, and thereby each lane was checked for the presence or absence of homoglutamic acid. Thus, complementary strain mixture cells recovering homoglutamic acid-producing ability were selected and separated into single colonies, and strains recovering homoglutamic acid-producing ability were selected, and they were referred to as complementary strains. One of plasmids prepared from these complementary strains using QIAGEN Plasmid Midi Kit was named pCF213. About 6.5 kbp of an insert DNA was inserted into pCF213. Together with the complementarity of a separately obtained plasmid pCE111 on each mutant, the complementarity of the above pCF213 was examined, and



Solution B 70 % acetonitrile

Flow rate: 2.0 ml/min

Elution condition: gradient of a fixed flow rate, linear  
gradient of from 2 % to 40 % of Solution B in from  
0 to 7 minutes, 100 % of Solution B in more than 7  
minutes

Detection: UV 254 nm

Temperature: 40°C

Under these conditions, the retention time of homo-  
glutamic acid was 1.3 minutes, and that of lysine was 7.7 minutes.

As is seen from the results shown in Figure 4, the wild type  
pCF213 strain has homoglutamic acid-producing ability 1.5 times  
higher than that of the wild type pCF704 strain.

6. Determination of the gene base sequence of the pCF 213 insert  
region

The base sequence of the pCF 213 insert region was deter-  
mined according to the primer walking method using ABIPRISM  
377XL DNA Sequencer (Perkin Elmer corporation). This base se-  
quence is shown in SEQ ID NO: 3.

The open reading frame (ORF) on the determined base  
sequence was determined using the method of Bibb et al.(Gene 30,  
157 (1984)). As a result, ORF shown in Figure 5 was found.

7. Analysis of the NotI site of about 2.5 kbp in the pCF213 insert  
region

Analysis of the NotI site of about 2.5 kbp (the base se-  
quence of from 2077 to 4578 in SEQ ID NO: 3) in the pCF213 insert  
region was carried out. This NotI site of about 2.5 kbp was cut out  
from the agarose gel, and the DNA was recovered and purified using  
Ultrafree C3 Unit 0.45  $\mu$ m (Millipore corporation) and dissolved in TE  
solution, and the termini were blunted according to DNA Blunting Kit

(Takara company), and the resulting solution was referred to as Insert DNA solution. On the other hand, pCF704 was digested with a restriction enzyme HincII and then dephosphorylated with alkaline phosphatase. This and Insert DNA solution were subjected to ligation  
5 reaction using Ligation Kit version 1 (Takara company). F. lutescens IFO 3084 strain was transformed with this reaction mixture, and a plasmid pCF235 was prepared from the transformant using QUI-AGEN Plasmid Midi Kit.

The first mutant transform with pCF235 was inoculated  
10 into 3 ml of the screening medium, and shaking cultured at 32°C for 2 days. 3 µl portions of this culture broth were added dropwise on each lane of TLC silica gel plate and dried. This plate was developed with a solvent system consisting of butanol, acetic acid and water (3:1:1) and subjected to ninhydrin coloring, and each lane was checked for the  
15 presence or absence of homoglutamic acid. As a result, it was revealed that the first mutant transformed with pCF235 recovered homoglutamic acid-producing ability.

In the DNA sequence of about 2.5 kbp integrated into pCF235 was present an ORF encoding 510 amino acids starting from  
20 ATG of 2855th of the base sequence of SEQ ID NO: 3 and ending in TAA of 4387th. This amino acid sequence was subjected to homology search by BLAST, and as a result, showed high homology with various aldehyde dehydrogenases, and further showed high homology with the amino acid sequence of piperidine-6-carboxylic acid dehydrogenase of  
25 Streptomyces dlavuligerus lately registered with database (J. Bac., Vol.180, No.17, 4753-4756 (1998)) over the whole amino acid sequence. Taking it into account that the first mutant transformed with pCF235 recovered homoglutamic acid-producing ability and that the homoglutamic acid-producing ability of the wild type pCF213 strain was  
30 heightened, the protein encoded by this ORF can be regarded as

having piperidine-6-carboxylic acid dehydrogenase.

### Example 2

Cloning of a gene encoding a protein having LAT activity, etc.

#### 1. Assay of LAT activity

5           Lysine-HCl (73 mg) and 59 mg of 2-ketoglutaric acid were dissolved in 1 ml of 0.2 M phosphate buffer (pH 7.3) containing 0.5 mM pyridoxal phosphate, and the solution was referred to as reaction solution. The reaction solution (28.75  $\mu$ l) was added to 260  $\mu$ l of the enzyme solution, and the mixture was left alone at 32°C for 1 hour.

10   151.8  $\mu$ l of a solution of 5 % trichloroacetic acid in ethanol was added to discontinue the reaction, the reaction mixture was centrifuged, 90  $\mu$ l of 0.2 M phosphate buffer (pH 7.3) containing 4 mM o-aminobenzaldehyde was added to 60  $\mu$ l of the supernatant, and the mixture was left alone at 37°C for 1 hour. The mixture was assayed for A465, and

15   the fractions having relatively high A465 were referred to as LAT active fractions.

#### 2. Culture of strain

F. lutescens IFO 3084 strain was shaking cultured at 32°C overnight. The culture broth (50 ml) as an inoculum was inoculated

20   into 10 L of flavo-M9 medium (0.6 %  $\text{Na}_2\text{HPO}_4$ , 0.3 %  $\text{KH}_2\text{PO}_4$ , 0.1 %  $\text{NH}_4\text{Cl}$ , 0.2 %  $\text{NaCl}$ , 1.0 % polypeptone, 0.5 % yeast extract, 0.5 % lysine-HCl, 0.005 % silicone KM75, 0.025 %  $\text{MgSO}_4$ , 0.0015 %  $\text{CaCl}_2$ , pH 7.2) in 30 L jar fermenter, and aeration stirring cultured for 17 hours. The resulting culture broth (5 L) was centrifuged (1,000  $\times$  g, 10

25   minutes) to collect the cells, and the cells were washed twice with 0.01 M phosphate buffer (pH 7.2). The cells were suspended in the same buffer and subjected to ultrasonic fracture. The fractured cells were removed by centrifugation (1,000  $\times$  g, 10 minutes) to obtain a cell extract. The cell extract was ultracentrifuged (16,000  $\times$  g, 90 min-

30   utes), and the supernatant fraction was subjected to the following

purification operations.

### 3. Purification of enzyme

All the following purification operations were carried out at 4°C, unless otherwise noted.

#### 5 (1) Ammonium sulfate fractionation

The supernatant fraction (600 ml) obtained in Example 1 was purified by ammonium sulfate precipitation. The precipitates formed in the fractions of from 30 % saturation to 80 % saturation were collected by centrifugation (1,000 × g, 30 minutes), and dissolved  
10 in 0.01 M phosphate buffer (pH 7.2), and the solution was dialyzed against the same buffer.

#### (2) Desalting

The dialyzed enzyme solution (10 ml) was poured on 4 PD10 columns (Amasham Pharmacia) and eluted and desalted with  
15 0.1 M Tris-HCl buffer (pH 7.4) containing 0.5 mM pyridoxal phosphate.

#### (3) QAE-TOYOPEAL550C column chromatography

The desalted enzyme solution was poured on QAE-TOYOPEAL550C (TOSOH) column (φ 5.5 × 6.0 cm) previously equi-  
20 brated with 0.1 M Tris-HCl buffer (pH 7.4) containing 0.5 mM pyridoxal phosphate, washed with the same buffer, and eluted by 2 L of sodium chloride linear gradient (0 to 1.0 M) using the same buffer, and LAT active fractions were collected.

#### (4) Phenyl-TOYOPERL650S column chromatography

25 1 M ammonium sulfate was added to the LAT active fractions, and the mixture was poured on Phenyl-TOYOPERL650S (TOSOH) column (φ 5.5 × 3.0 cm) previously equilibrated with 0.01 M phosphate buffer (pH 7.2) containing 0.5 mM pyridoxal phosphate and 1 M ammonium sulfate, and eluted with 1,200 ml of ammonium  
30 sulfate gradient (0.8 to 0 M) using the same buffer, and LAT active

fractions were collected.

(5) Ultrafiltration

The LAT active fractions (150 ml) were ultrafiltered with ADVANTEC UP-20 to make the volume 15 ml. This concentrate (2.5  
5 ml) was poured on PD10 column (Amasham Pharmacia), and eluted and desalted with 0.1 M Tris-HCl buffer (pH 7.4).

(6) AKTA MonoQ HR5/5 column chromatography

The desalted enzyme solution (3.5 ml) was poured on MonoQ HR5/5 column of AKTAexplorer 10S System (Amasham  
10 Pharmacia) previously equilibrated with 0.1 M Tris-HCl buffer (pH 7.4), washed with the same buffer, and eluted with 40 ml of sodium chloride linear gradient (0 to 0.4 M) using the same buffer, and LAT active fractions were collected. The LAT active fractions (5 ml) were desalted with PD10 column, and subjected to MonoQ HR5/5 column of  
15 AKTAexplorer 10S System, and LAT active fractions were collected. Relations between each fraction and relative LAT activity are shown in Figure 6.

(7) Electrophoresis

The LAT active fractions were subjected to Multigel 4/20  
20 and 10/20 (Daiichi Kagaku Yakuhin Co., Ltd.) and Native-PAGE and SDS-PAGE were carried out, and the results are shown in Figure 7. As to the LAT active fractions, a band was observed at a molecular weight of around 100,000 in Native-PAGE and a band was observed at a molecular weight of around 55,000 in SDS-PAGE. A PVDF mem-  
25 brane was blotted with the band of a molecular weight of around 55,000 in SDS-PAGE using PhastTransfer (Amasham Pharmacia).

4. Analysis of N-terminus amino acid sequence

Analysis of N-terminus amino acid sequence of the band subjected to the blotting was carried out by Edman degradation  
30 method using HP G1005A Protein Sequencing System (HEWLETT

PACKARD). As a result, it was revealed that the N-terminus amino acid sequence was

SLLAPLAPLRAHAGTRLTQG (SEQ ID NO: 9)

Based on this, DNA primers (N = 1)

5 NmaRout CCYTGNGTNARNCKNGTNCNGCRTCNGGCNCG (SEQ ID NO: 10)

NmaRin CCNGCRTCNGGCNCGNARNGGNGCNARNGGNGC (SEQ ID NO: 11)

were designed, and PCR was carried out on the genome DNA of *F. lutescens* IFO 3084 strain using LA PCR in vitro cloning KIT (Takara Company). The PCR reaction condition was 30 cycles of 94°C, 30  
 10 seconds → 55°C, 2 minutes → 72°C, 1 minute. As a result, a PCR amplification fragment of about 400 bp containing the above terminus and its upstream region was obtained. Based on this sequence, its neighborhood region was obtained using PCR. Namely, the genome DNA of *F. lutescens* IFO 3084 strain was digested with restriction  
 15 enzymes *Pst*I and *Sal*I, respectively, and the digests were subjected, respectively, to self-ligation reaction using Ligation Kit version 2 (Takara Company), and the resulting DNAs were used as template DNAs.

Based on these template DNAs, DNA primers

20 NIFout ttgatttgag cagattcgca ctgccattt (SEQ ID NO: 5)

NIRout aaggttttcg acaaagtgac catttcca (SEQ ID NO: 6)

were designed, and PCR was carried out using LA Taq (Takara Company). The PCR reaction condition was 30 cycles of 98°C, 20 seconds → 68°C, 6 minutes. As a result, a PCR amplification fragment of  
 25 about 2 kbp was obtained from the *Pst*I template and a PCR amplification fragment of about 8 kbp from the *Sal*I template. The base sequence was determined by the primer walking method using ABIPRISM 377XL DNA Sequencer (Perkin Elmer corporation) on these PCR amplification fragments. This base sequence is shown in  
 30 SEQ ID NO: 1.



## 5. Construction of plasmids pCF301 and pCF335

The following DNA primers wherein the PstI sites of base 545 and base 2658 of SEQ ID NO: 1 were converted to KpnI and SacI sites, respectively,

5                   ctggtaccgc tcgatccggc tctgcaccgt           (SEQ ID NO: 7)

                  ctggagctca ggcaggtgcg ggccacgtgt           (SEQ ID NO: 8)

were prepared, and PCR reaction was carried out using these primers to amplify the lat gene region. The amplified fragment of about 2.1 kbp was digested with restriction enzymes KpnI and SacI, and the  
10 resulting solution was referred to as Insert DNA solution. On the other hand, pCF704 was digested with restriction enzymes KpnI and SacI, and the digest and Insert DNA solution were subjected to ligation reaction using Ligation Kit version 2 (Takara company), and the resulting plasmid was referred to as pCF301. Further, pCF301  
15 was digested with restriction enzymes KpnI and SacI, and the 2.1 kbp fragment was cut out from agarose gel, and this and the digest of pCF235 with restriction enzymes KpnI and SacI were subjected to ligation reaction, and the resulting plasmid was named pCF335.

## 6. Complementation of LAT activity by plasmid pCF301

20                   A mutant obtained by transforming the second mutant with pCF704 was designated 2nd pCF704 strain, and a mutant obtained by transforming the second mutant with pCF301 was designated 2nd pCF301 strain. These strains were shaking cultured at 32°C overnight. Each (30 µl) of the culture broths as an inoculum was  
25 inoculated into 3 ml of a production medium (1.5 % polypeptone, 0.5 % yeast extract, 2.0 % lysine-HCl, pH not adjusted) in a centrifugation tube, and aeration stirring cultured for 17 hours. The resulting culture broth (1 ml) was centrifuged (1,000 × g, 10 minutes) to collect the cells, and the cells were washed with 10 ml of 0.2 M  
30 phosphate buffer (pH 7.3) containing 0.5 mM pyridoxal phosphate.

The cells were suspended in 1 ml of the same buffer and ultrasonically fractured. The fractured cells were removed by centrifugation (1,000 × g, 10 minutes) to obtain a cell extract. This cell extract was assayed for LAT activity. The results are shown in Figure 8. pCF301  
5 complemented the lat mutation in the second mutant.

#### 7. Heightening of homoglutamic acid-producing ability by pCF335

A transformant obtained by transforming the wild type E. lutescens IFO 3084 strain with pCF704 was designated wild type pCF704 strain, and transformants obtained by transforming the IFO  
10 3084 strain with plasmids pCF301 and pCF335 were designated wild type pCF301 strain and wild type pCF335 strain, respectively. These strains were inoculated into 3 ml portions of the screening medium containing 20 µg/ml kanamycin, respectively, and shaking cultured at 32°C overnight. 100 µl portions of each of the culture broths as an  
15 inoculum were inoculated into 25 ml portions of a production medium (1.5 % polypeptone, 0.5 % yeast extract, 2.0 % lysine-HCl, pH not adjusted), and shaking cultured at 32°C for 24 hours, 48 hours and 72 hours, respectively. The amount of homoglutamic acid in the supernatant of each of the culture broths was measured by HPLC. Namely,  
20 each of the culture broths was diluted with distilled water so that the total amino acid concentration could be on the order of 1,000 mg/L, and 50 µl of the dilution was transferred into a test tube and concentrated to dryness. To the residue was added 50 µl of a mixed solution of phenyl isothiocyanate, triethylamine, ethanol and distilled water  
25 (1:1:7:2), and the mixture was stirred to make a solution, left alone at room temperature for 10 minutes and concentrated to dryness under reduced pressure. The residue was dissolved in 500 µl of Solution A as a mobile phase of HPLC, and 5 µl thereof was injected. The HPLC condition is as described in 5 of Example 1.

30 As a result, as shown in Figure 9, the wild type pCF335

strain had homoglutamic acid-producing ability about twice higher than that of the wild type pCF704 strain.

## SEQUENCE LISTING

<110> MERCIAN CORPORATION

<120> GENE PARTICIPATING IN THE PRODUCTION OF HOMO-GLUTAMIC  
ACID AND UTILIZATION THEREOF

<130> 67566-1434

<140> CA 2,337,981

<141> 1999-08-04

<150> JP10/232382

<151> 1998-08-05

<150> JP11/182362

<151> 1999-06-28

<160> 11

<170> PatentIn Ver. 2.1

<210> 1

<211> 2663

<212> DNA

<213> Flavobacterium lutescens

<220>

<221> CDS

<222> (801)..(2276)

<400> 1

```

cccgggtgtc attgaatacc agcagggtcgc caggttgcag cagctgggtcc agatcgcgca 60
cctggcgatc ctccagcgcga gccgggtgccg gcggcaccag cagcaggcgg ctggccgaac 120
gctccggcag cggcgcctgg gcaatcagtt cgggaggcag gtggtaggca aaatcggact 180
tcttcaacgc cggcagctcg atacaacggg ggcgtcagtt tacgcccctg taccgcctgt 240
gccctcaccg ctcgaacttg gtgcccagga tcaccgccgt ggtggtgcgc tcgaccccat 300
cagtggcgcc gatggcatcg gtcagctcgt ccategccgc cagccatcg acggcggcca 360
tcgccaccag gtcatgcgcg cactgaccg aatgcaggct gcgcaccgca gcaatggcct 420
gcagcgcccg cagcaccgcc ggcattttct tcggcatcac ggtgatggag atatgcgcgc 480
ggacctgctg gcgctccatc gcctggccaa ggcgcacggg ttagccggcg attattccgc 540
tgtgctgcag ccgctcgatc cggctctgca ccgtgggtccg cgacaccccg agccggcgcg 600
ccagcgccgc ggtcgaggcg cgcgcacccg cacgcaacag gtcaagcaac tgtgcatccg 660
cctgggaaat ggtcactttg tcgaaaacct ttcgtcaatc cgccgaaacc ggccattgat 720
ttgagcagat tcgcactgcc atttgtagtg cgtaaaccgt tacaactaac actagacaca 780

```

```

atcagcacgg attcagcatg tcc ctt ctt gcc ccg ctc gcc ccg ctc cgc gcc 833
                Ser Leu Leu Ala Pro Leu Ala Pro Leu Arg Ala
                1                5                10

```

```

cat gcc ggc acc cgc ctt acc cag ggc ctg tct gac ccg cag gtc gag 881
His Ala Gly Thr Arg Leu Thr Gln Gly Leu Ser Asp Pro Gln Val Glu
                15                20                25

```

```

cag ctg gcc gcc aac cac cct gac ctg cgc gcc gcc atc gac gcc gct 929
Gln Leu Ala Ala Asn His Pro Asp Leu Arg Ala Ala Ile Asp Ala Ala
                30                35                40

```

gcc gac gaa tac gcg cgc atc aaa ccg cag gcc gcg gca ttg ctg gac Ala Asp Glu Tyr Ala Arg Ile Lys Pro Gln Ala Ala Ala Leu Leu Asp 45 50 55	977
ctg gat gaa agc gcg cag atc gcc gcc gtg cag gat ggc ttc gtc aac Leu Asp Glu Ser Ala Gln Ile Ala Ala Val Gln Asp Gly Phe Val Asn 60 65 70 75	1025
ttc tat gcc gat gat gcg gtg gtg ccc tat atc gcc ctg gcc gcc cgc Phe Tyr Ala Asp Asp Ala Val Val Pro Tyr Ile Ala Leu Ala Ala Arg 80 85 90	1073
ggg ccg tgg gtg gtc agc ctg aag ggc gcg gtg ctg tat gac gcc ggc Gly Pro Trp Val Val Ser Leu Lys Gly Ala Val Leu Tyr Asp Ala Gly 95 100 105	1121
ggc tac gcc atg ctc gcc ttc gcc cat acc ccg gcc gat atc ctg gag Gly Tyr Gly Met Leu Gly Phe Gly His Thr Pro Ala Asp Ile Leu Glu 110 115 120	1169
gcg gtc gcc aag ccg cag gtg atg gcc aac atc atg act ccc tcg ctg Ala Val Gly Lys Pro Gln Val Met Ala Asn Ile Met Thr Pro Ser Leu 125 130 135	1217
gcc cag gcc cgc ttc att gcc gca atg cgc cgc gaa atc gcc cat acc Ala Gln Gly Arg Phe Ile Ala Ala Met Arg Arg Glu Ile Gly His Thr 140 145 150 155	1265
cgc gcc gcc tgc ccg ttc tcg cac ttc atg tgc ctg aac tcc gcc tcc Arg Gly Gly Cys Pro Phe Ser His Phe Met Cys Leu Asn Ser Gly Ser 160 165 170	1313
gaa gcg gtc ggg ctg gcc gcg cgc atc gcc gac atc aac gcc aag ctg Glu Ala Val Gly Leu Ala Ala Arg Ile Ala Asp Ile Asn Ala Lys Leu 175 180 185	1361
atg acc gac ccg gcc gcc cgg cat gcc gcc gcc acg atc aag cgc gtg Met Thr Asp Pro Gly Ala Arg His Ala Gly Ala Thr Ile Lys Arg Val 190 195 200	1409
gtg atc aag gcc agt ttc cac gcc cgt acc gac cgt ccg gcg ctg tat Val Ile Lys Gly Ser Phe His Gly Arg Thr Asp Arg Pro Ala Leu Tyr 205 210 215	1457
tcc gat tcc acc cgc aag gcc tac gat gcg cat ctg gcc agc tac cgc Ser Asp Ser Thr Arg Lys Ala Tyr Asp Ala His Leu Ala Ser Tyr Arg 220 225 230 235	1505
gac gag cac agc gtc att gcc atc gcc ccg tat gac cag cag gcc ctg Asp Glu His Ser Val Ile Ala Ile Ala Pro Tyr Asp Gln Gln Ala Leu 240 245 250	1553
cgc cag gtg ttt gcc gat gcc cag gcc aac cac tgg ttc atc gag gcg Arg Gln Val Phe Ala Asp Ala Gln Ala Asn His Trp Phe Ile Glu Ala 255 260 265	1601
gtg ttc ctg gag ccg gtg atg gcc gaa gcc gac ccg gcc cgt gcg gtg Val Phe Leu Glu Pro Val Met Gly Glu Gly Asp Pro Gly Arg Ala Val 270 275 280	1649

ccg gtg gac ttc tac cgc ctg gcc cgt gag ctg acc cgc gaa cac ggc 1697  
 Pro Val Asp Phe Tyr Arg Leu Ala Arg Glu Leu Thr Arg Glu His Gly  
 285 290 295

agc ctg ctg ctg atc gat tcg atc cag gcc gcg ctg cgc gtg cac ggc 1745  
 Ser Leu Leu Leu Ile Asp Ser Ile Gln Ala Ala Leu Arg Val His Gly  
 300 305 310 315

acc ctg tcc ttc gtc gac tac ccc ggc cac cag gag ctg gag gca ccg 1793  
 Thr Leu Ser Phe Val Asp Tyr Pro Gly His Gln Glu Leu Glu Ala Pro  
 320 325 330

gac atg gag acc tac tcc aag gcc ctg aac ggc gcc cag ttc ccg ctg 1841  
 Asp Met Glu Thr Tyr Ser Lys Ala Leu Asn Gly Ala Gln Phe Pro Leu  
 335 340 345

tcg gta gtg gcc gtg acc gag cac gcc gcc gcg ctg tac cgc aag ggc 1889  
 Ser Val Val Ala Val Thr Glu His Ala Ala Ala Leu Tyr Arg Lys Gly  
 350 355 360

gtg tac ggc aac acc atg acc acc aac ccg cgg gcg ctg gac gtg gcc 1937  
 Val Tyr Gly Asn Thr Met Thr Thr Asn Pro Arg Ala Leu Asp Val Ala  
 365 370 375

tgc gcc acc ctg gca cgc ctg gat gag ccg gtc cgc aac aat atc cgc 1985  
 Cys Ala Thr Leu Ala Arg Leu Asp Glu Pro Val Arg Asn Asn Ile Arg  
 380 385 390 395

ctg cgt ggc cag cag gcg atg cag aag ctg gaa gca ttg aag gaa cgg 2033  
 Leu Arg Gly Gln Gln Ala Met Gln Lys Leu Glu Ala Leu Lys Glu Arg  
 400 405 410

ctg ggg ggc gcg atc acc aag gtg cag ggc acc ggc ctg ctg ttc tcc 2081  
 Leu Gly Gly Ala Ile Thr Lys Val Gln Gly Thr Gly Leu Leu Phe Ser  
 415 420 425

tgc gag ctg gcc ccg cag tac aag tgc tac ggg gcc ggc tcc acc gag 2129  
 Cys Glu Leu Ala Pro Gln Tyr Lys Cys Tyr Gly Ala Gly Ser Thr Glu  
 430 435 440

gag tgg ctg cgc atg cac ggg gtc aat gtg atc cac ggc ggc gag aat 2177  
 Glu Trp Leu Arg Met His Gly Val Asn Val Ile His Gly Gly Glu Asn  
 445 450 455

tcg ctg cgc ttc acc ccg cac ttc ggc atg gac gag gcc gaa ctg gac 2225  
 Ser Leu Arg Phe Thr Pro His Phe Gly Met Asp Glu Ala Glu Leu Asp  
 460 465 470 475

ctg ctg gtg gag atg gtc ggg cgt gcg ctg gtc gaa ggc cca cgc cgg 2273  
 Leu Leu Val Glu Met Val Gly Arg Ala Leu Val Glu Gly Pro Arg Arg  
 480 485 490

gcc tgatccgcac ccgcaggacg gaaggcacga gccaccgtg aggcgggctc 2326  
 Ala

tttgctgccc ggcaccagcg gcaacaggcc gcgctgtcac cggccaggcg gggcgccggc 2386  
 agtgggttcc agccgcaggg gtccgccctg ccagcgcctg cggcggggca caggcttgcg 2446  
 ggcattgcgg cctctgccac gggcacgcag ccggagatca ggctgacaag ggggctgccc 2506  
 cgggtggcag tacacgacca gccagttgac tgccggtatt tgcttgatca gcgctgcatc 2566

cagaacagca ccatacgggttg cgtgactgac ggcgccgctgg ccgttgcggg acagcagcct 2626  
 ttgcgtcaca cgtggccccgc acctgcctgc actgcag 2663

<210> 2  
 <211> 492  
 <212> PRT  
 <213> Flavobacterium lutescens

<400> 2  
 Ser Leu Leu Ala Pro Leu Ala Pro Leu Arg Ala His Ala Gly Thr Arg  
 1 5 10 15  
 Leu Thr Gln Gly Leu Ser Asp Pro Gln Val Glu Gln Leu Ala Ala Asn  
 20 25 30  
 His Pro Asp Leu Arg Ala Ala Ile Asp Ala Ala Ala Asp Glu Tyr Ala  
 35 40 45  
 Arg Ile Lys Pro Gln Ala Ala Ala Leu Leu Asp Leu Asp Glu Ser Ala  
 50 55 60  
 Gln Ile Ala Ala Val Gln Asp Gly Phe Val Asn Phe Tyr Ala Asp Asp  
 65 70 75 80  
 Ala Val Val Pro Tyr Ile Ala Leu Ala Ala Arg Gly Pro Trp Val Val  
 85 90 95  
 Ser Leu Lys Gly Ala Val Leu Tyr Asp Ala Gly Gly Tyr Gly Met Leu  
 100 105 110  
 Gly Phe Gly His Thr Pro Ala Asp Ile Leu Glu Ala Val Gly Lys Pro  
 115 120 125  
 Gln Val Met Ala Asn Ile Met Thr Pro Ser Leu Ala Gln Gly Arg Phe  
 130 135 140  
 Ile Ala Ala Met Arg Arg Glu Ile Gly His Thr Arg Gly Gly Cys Pro  
 145 150 155 160  
 Phe Ser His Phe Met Cys Leu Asn Ser Gly Ser Glu Ala Val Gly Leu  
 165 170 175  
 Ala Ala Arg Ile Ala Asp Ile Asn Ala Lys Leu Met Thr Asp Pro Gly  
 180 185 190  
 Ala Arg His Ala Gly Ala Thr Ile Lys Arg Val Val Ile Lys Gly Ser  
 195 200 205  
 Phe His Gly Arg Thr Asp Arg Pro Ala Leu Tyr Ser Asp Ser Thr Arg  
 210 215 220  
 Lys Ala Tyr Asp Ala His Leu Ala Ser Tyr Arg Asp Glu His Ser Val  
 225 230 235 240  
 Ile Ala Ile Ala Pro Tyr Asp Gln Gln Ala Leu Arg Gln Val Phe Ala  
 245 250 255  
 Asp Ala Gln Ala Asn His Trp Phe Ile Glu Ala Val Phe Leu Glu Pro  
 260 265 270

Val Met Gly Glu Gly Asp Pro Gly Arg Ala Val Pro Val Asp Phe Tyr  
 275 280 285

Arg Leu Ala Arg Glu Leu Thr Arg Glu His Gly Ser Leu Leu Leu Ile  
 290 295 300

Asp Ser Ile Gln Ala Ala Leu Arg Val His Gly Thr Leu Ser Phe Val  
 305 310 315 320

Asp Tyr Pro Gly His Gln Glu Leu Glu Ala Pro Asp Met Glu Thr Tyr  
 325 330 335

Ser Lys Ala Leu Asn Gly Ala Gln Phe Pro Leu Ser Val Val Ala Val  
 340 345 350

Thr Glu His Ala Ala Ala Leu Tyr Arg Lys Gly Val Tyr Gly Asn Thr  
 355 360 365

Met Thr Thr Asn Pro Arg Ala Leu Asp Val Ala Cys Ala Thr Leu Ala  
 370 375 380

Arg Leu Asp Glu Pro Val Arg Asn Asn Ile Arg Leu Arg Gly Gln Gln  
 385 390 395 400

Ala Met Gln Lys Leu Glu Ala Leu Lys Glu Arg Leu Gly Gly Ala Ile  
 405 410 415

Thr Lys Val Gln Gly Thr Gly Leu Leu Phe Ser Cys Glu Leu Ala Pro  
 420 425 430

Gln Tyr Lys Cys Tyr Gly Ala Gly Ser Thr Glu Glu Trp Leu Arg Met  
 435 440 445

His Gly Val Asn Val Ile His Gly Gly Glu Asn Ser Leu Arg Phe Thr  
 450 455 460

Pro His Phe Gly Met Asp Glu Ala Glu Leu Asp Leu Leu Val Glu Met  
 465 470 475 480

Val Gly Arg Ala Leu Val Glu Gly Pro Arg Arg Ala  
 485 490

<210> 3  
 <211> 6357  
 <212> DNA  
 <213> Flavobacterium lutescens

<220>  
 <221> CDS  
 <222> (2855)..(4384)

<400> 3  
 ggatcgggcc actgggctca ctgctggacg caatccgagt gccgggatgg ctcggggtga 60  
 aggtgttgcg gatcacgac gccatctgcc gggcgatggc cgggctcatc gtctgcgggt 120  
 gcaccacctt ggcgccgaaa taggccagtt cgcaggcctc gtcatagctg agcgtggcca 180  
 gggtcaccgc ctcgggcacc acccgcgggt cggccgacag cacaccgtcg acatcgggtcc 240  
 agatgtgcag ctcggccgcc tcgaacagcg cggcaaagat cgccccggaa taatcgctgc 300  
 cgttgcggcc caggggtggtg atcctgcctt ggccatcacg ggcgacaaaa ccggtgacca 360



ccacccgcga ctgCGGGttg tccacacgcc aggcggccag gttggccgca ctgcgttccc 420  
 agtcgacgct gacccccagc tcgCCgtgtg cgaccaccag cacatcgcgg gcatcgagca 480  
 ccgCGcaggg gtggccgagc cggttgaaat agcggcccag cagctggggc gagaacacct 540  
 cgcccagccc ctgcaccctt tcaagcacct cctcggggcag gccgcccgat accgcccagcg 600  
 cttccagcaa cccggccagc ttgtcaaagc gtccatccag ccaactgcagc aggtcggcag 660  
 aatcctcgcc cagcagttcg gtggccgctt catggtggcg ctggcgcagg gcctgccagg 720  
 catcacgcca gcgcggctga ccgtggggcgg ccagggtagc cagctcgatc aaggcatcgg 780  
 tgacaccctt catcgccgag accaccacca cctgggtggg ttccggggcg tgcagcagca 840  
 actcggcgac atggcggtag cgctgcgccc aggccaccga ggtgccgccc aacttgtggg 900  
 cgatgacctg ggcacggggc gcgggagcgg gagcgggtgc agcggcaggc gatgacatca 960  
 caacagacct ctggggttga ggcccggcac cgcaggttgc gaagtcccgc aacctggtcg 1020  
 gtgcgggggc gttgttttcg ggggttagac gaatacgacg ggccgcacca gccaaagtgg 1080  
 ggtggtaatg atggtcatgc cggtgacgcc agcaggcggc agcaggggcg cagtggaatc 1140  
 aacggtggcg cggcagatcg acatgcagcg agcagaccgc acagcgcctg ctgctgtcaa 1200  
 ctggtgcatt gcaaaataat tttccgcgca tcatcggcga acatgcaccg atttggttgc 1260  
 aatgtgatc gtcagcgatc ttctgtcaaa acccgcggat caagcggcca cagccgctgc 1320  
 ggcagccgcg gaccaccgcg cgcgatgcc agcgcggggc ggcagagcaa gccgcccagcg 1380  
 caaccggcca ttaccgcggc caggcgcggc gcctgcgcg ctcaaccgtg gattttttcc 1440  
 cagcgggCGT gggcctgcgc ggcagcacc acccgcgcca ccaacagcgc aatggccagc 1500  
 agtccagca gggtcggggc acgctgctgc cagatgaagc cataaagcaa cgCGaacagg 1560  
 gtttcaaaCa cgatcagctg cccgcccagg ctCagcggca ggctgcgCGT ggcccggttc 1620  
 cagcaggcat tgcccagcac cgaggcaccg acggccagca gcgcacagat gccggcaaag 1680  
 tgcagccact ggccctggct ctgcccgagc ggcccagcc acagcgcagc cggcagcaac 1740  
 agcagggcga tggcccctgt ggccaccccg gtcaacaacg accaggcatg cccggacagg 1800  
 tgcggatagc gccgcatcca caccacattg gcgatcgagt agccactcca ggCGgcccagc 1860  
 gcggccagcg cgcagagcag gccagcacc cgtgaccga tgtccttgcc agcatcgcCC 1920  
 gccgcgcccg cgcCGTggcc gagtgCagcc caggccacca gcagcagacc cagcacacac 1980  
 aggcacagcg ccggtgccag ctgacgcaac ggcagggcgg ttggccgccc cgcatccacc 2040  
 gccgcccacca ccaccggcac catgcccacg atcagcgcgg ccgcccgacc gccagcccag 2100  
 tgcacggcca tcgcccagaaa cacgaaatag accaggttgc cgagcaggct cagcccggcc 2160  
 agggccagcc aggcgcggcg atcgacctgc gcacgcagcg ccggccacaa cggcagcagc 2220  
 aacgcacagg ccaccgcacc gtacagcagg tagcggccca cggccagctg cagcgcagaa 2280  
 aatgcggTgg tcaaggccgg cgcCaggaac accatgcccc acagggcacc ggCGagcacg 2340  
 ccgTtgaaca gtccccacgc ggtctggttG ttgcgctgga tcacgctgca aggccctgca 2400  
 atgaacaaca ggccggggcg gcgcagcGca tgggcgctgg cagctctccg acctgtgcaa 2460  
 aggtggtggc cccgacacga ttcgaacgtg cgacctgtcc cttaggaggg gaccgctcta 2520  
 tccagctgag ctacggagcc atgaggccgg cgattctagc atccgctctc cgttcacggc 2580  
 catcgcCCcg agccgcagtt cacagtgcag ggcaaccgca gcaagcccc gccccgctgc 2640  
 aacctcgcg cccgcgcgca acgtgaccgg cgcCGcggca ggcccggccc ccacggccac 2700  
 tggcgcCGgt tccgcaccac gccaccggca acacgcccc agccctgccc aacgtggtgt 2760  
 ttcagcGctc tgTtaagatg gcatgcccac atgccacctc ccccCGgacg cgcCGcgggt 2820

gcgtgacctt ttcgtaacgt aatctggagt ttcc atg tcg ttt gaa ctg ctc aag 2875  
 Met Ser Phe Glu Leu Leu Lys

1

5

gcc tta ggg ctg gac gcc acc aat tcc ggc acc tac ctg ggt gat gga 2923  
 Ala Leu Gly Leu Asp Ala Thr Asn Ser Gly Thr Tyr Leu Gly Asp Gly  
 10 15 20

gaa tgg tcc agc gct acc ggt gcc ggg acc atc agc ccg cgc aac ccg 2971  
 Glu Trp Ser Ser Ala Thr Gly Ala Gly Thr Ile Ser Pro Arg Asn Pro  
 25 30 35

acc acc ggc gag gtc att gcc cag gtc cag gcc acc acc gag gcg gac 3019  
 Thr Thr Gly Glu Val Ile Ala Gln Val Gln Ala Thr Thr Glu Ala Asp  
 40 45 50 55

tac gaa acc atc ctg gcc cgc gcc cag cag gcc ttc aag gtc tgg cgc	3067
Tyr Glu Thr Ile Leu Ala Arg Ala Gln Gln Ala Phe Lys Val Trp Arg	
60 65 70	
acc acc ccg gca ccg cgc cgc ggc gag gcc atc cgc ctg tgt ggc gag	3115
Thr Thr Pro Ala Pro Arg Arg Gly Glu Ala Ile Arg Leu Cys Gly Glu	
75 80 85	
gcc ctg cgc cgc cac aag gac gcg ctg ggt tcg ctg gtc gcg ctg gaa	3163
Ala Leu Arg Arg His Lys Asp Ala Leu Gly Ser Leu Val Ala Leu Glu	
90 95 100	
atg ggc aag tcc aag ccg gaa ggc gac ggc gaa gtc cag gaa atg atc	3211
Met Gly Lys Ser Lys Pro Glu Gly Asp Gly Glu Val Gln Glu Met Ile	
105 110 115	
gac atc gcc gac ttt gcc gtc ggc cag agc cgc atg ctg tat ggc tac	3259
Asp Ile Ala Asp Phe Ala Val Gly Gln Ser Arg Met Leu Tyr Gly Tyr	
120 125 130 135	
acc atg cac agc gag cgc ccc ggc cac cgc atg tac gag cag tac cag	3307
Thr Met His Ser Glu Arg Pro Gly His Arg Met Tyr Glu Gln Tyr Gln	
140 145 150	
ccg ctg ggc atc gtc ggc atc atc tcg gcc ttc aac ttc ccg gtc gcg	3355
Pro Leu Gly Ile Val Gly Ile Ile Ser Ala Phe Asn Phe Pro Val Ala	
155 160 165	
gtc tgg gcc tgg aac agc ttc ctg gcc gcg atc tgt ggt gat gtc tgc	3403
Val Trp Ala Trp Asn Ser Phe Leu Ala Ala Ile Cys Gly Asp Val Cys	
170 175 180	
atc tgg aag ccg tcc aac aag acc ccg ctg acc gcg atc gcg tcc atg	3451
Ile Trp Lys Pro Ser Asn Lys Thr Pro Leu Thr Ala Ile Ala Ser Met	
185 190 195	
cgc atc tgc aac gaa gca ctg cgc gaa ggc ggc ttc ccg gac atc ttc	3499
Arg Ile Cys Asn Glu Ala Leu Arg Glu Gly Gly Phe Pro Asp Ile Phe	
200 205 210 215	
ttc ctg atc aac gac gcc ggc acc gcg ttg tcg gag aag ctg gtc gag	3547
Phe Leu Ile Asn Asp Ala Gly Thr Ala Leu Ser Glu Lys Leu Val Glu	
220 225 230	
gac aag cgc gtg ccg ctg atc tcc ttc acc ggc tcg acc cag gtc ggg	3595
Asp Lys Arg Val Pro Leu Ile Ser Phe Thr Gly Ser Thr Gln Val Gly	
235 240 245	
cgc atc gtc aac cag aag gtc gcc gcc cgc ctg ggc cgc tgc ctg ctc	3643
Arg Ile Val Asn Gln Lys Val Ala Ala Arg Leu Gly Arg Cys Leu Leu	
250 255 260	
gag ctg ggc ggc aac aac gcg atc atc ctg gac gaa acc gcc gac ctg	3691
Glu Leu Gly Gly Asn Asn Ala Ile Ile Leu Asp Glu Thr Ala Asp Leu	
265 270 275	
aag ctg gcc gtg ccg ggc atc gtc ttc ggc gcc gtc ggc acc gcc ggc	3739
Lys Leu Ala Val Pro Gly Ile Val Phe Gly Ala Val Gly Thr Ala Gly	
280 285 290 295	

cag cgc tgc acc acc acc cgc cgc ctg atc gtg cac gaa tcg atc tac 3787  
 Gln Arg Cys Thr Thr Thr Arg Arg Leu Ile Val His Glu Ser Ile Tyr  
 300 305 310

gac aac gtg ctg gcc acc ttg atc aag gcc tac aag cag gtc gaa ggc 3835  
 Asp Asn Val Leu Ala Thr Leu Ile Lys Ala Tyr Lys Gln Val Glu Gly  
 315 320 325

aag atc ggc gat ccg ctg gat gcc gcc aac ctg atg ggc ccg ctc aac 3883  
 Lys Ile Gly Asp Pro Leu Asp Ala Ala Asn Leu Met Gly Pro Leu Asn  
 330 335 340

agc ccc gaa gcg gtg cag cag ttc ctg gcc tcg atc gag aaa gcc aag 3931  
 Ser Pro Glu Ala Val Gln Gln Phe Leu Ala Ser Ile Glu Lys Ala Lys  
 345 350 355

gcc gct ggc ggc acc gtt caa acc ggt ggt acc gcg atc gac cgc ccg 3979  
 Ala Ala Gly Gly Thr Val Gln Thr Gly Gly Thr Ala Ile Asp Arg Pro  
 360 365 370 375

ggc aac ttc gtg ctg ccg gcc atc gtc acc ggc ctg aag aac agc gat 4027  
 Gly Asn Phe Val Leu Pro Ala Ile Val Thr Gly Leu Lys Asn Ser Asp  
 380 385 390

gag gtg gtc cag cac gag acc ttc gcc ccg atc ctg tac gta atg aag 4075  
 Glu Val Val Gln His Glu Thr Phe Ala Pro Ile Leu Tyr Val Met Lys  
 395 400 405

tac tcc acc ctg gac gaa gcc atc gag atg cag aac ggc gtg ccg cag 4123  
 Tyr Ser Thr Leu Asp Glu Ala Ile Glu Met Gln Asn Gly Val Pro Gln  
 410 415 420

ggc ctg tcc tcg tcg atc ttc acc acg aac ctg aag gca gcc gag aag 4171  
 Gly Leu Ser Ser Ser Ile Phe Thr Thr Asn Leu Lys Ala Ala Glu Lys  
 425 430 435

ttc ctg tcg gcg gcc ggc agc gac tgc ggc att gcc aac gtc aac atc 4219  
 Phe Leu Ser Ala Ala Gly Ser Asp Cys Gly Ile Ala Asn Val Asn Ile  
 440 445 450 455

ggc act tcc ggt gcc gag atc ggc ggc gcc ttc ggt ggc gag aag gaa 4267  
 Gly Thr Ser Gly Ala Glu Ile Gly Gly Ala Phe Gly Gly Glu Lys Glu  
 460 465 470

acc ggc ggt ggc cgt gag tcc ggc tcg gat gcg tgg aag gtc tac atg 4315  
 Thr Gly Gly Gly Arg Glu Ser Gly Ser Asp Ala Trp Lys Val Tyr Met  
 475 480 485

cgc cgc cag acc aac acc atc aac tac tcc gac tcg ctg ccg ctg gcc 4363  
 Arg Arg Gln Thr Asn Thr Ile Asn Tyr Ser Asp Ser Leu Pro Leu Ala  
 490 495 500

cag ggc atc aag ttc gac ctg taagccgctc gccacggccc gccttccccg 4414  
 Gln Gly Ile Lys Phe Asp Leu  
 505 510

gaagcaggcc gtggctgttg caccagccag aggagtgact gcatgactgc aattgaatcg 4474  
 actgccgcac gcaccaccaa cacttgccgc atcctgtcgc tggactggc actgctgggc 4534  
 tggaatcttt tgccggtgat tggctttgtc ggcgccatca tctgcggccg catcgcccag 4594

```

cgccagctca agcagcccgg caatacccag gacggtcacg gcctggcaag ggcgggcatc 4654
tggatcagtt ggatcagcct gatcctgggt gcgctgctga tcggcgctcg gatcccgtgg 4714
ttgaccgccc cgatcacgat caacctgccc gtttccacct gaccctcctc cctgccagtc 4774
gcccattgcg tgacaggcca acccgtttcc tgctggacc agaccatgct cccgcccgac 4834
catccggctc caccatcgcc cattgccggc accacaacct cgaccaatgg ctatgcggtg 4894
gcctcgctgg tgatgggcat ccttggctgg tcgatgatcc cgctggtggg cagcatcggc 4954
gccatcggtg tcgggcatct ggcccgggcg cagatccgtc gccagcccca gcagggcgat 5014
ggcctggcac tggccgggct gatcctgggc tggatctcga ttgcgctgtg gatcctcggg 5074
atcctggcgt ttttctctt ctttggcggg ctggccatgc tgctcagcct gaacgcctga 5134
cccagacctt gccgtatgta ttccctgctc cgtcccgcct tgttctgcat ggatgccgag 5194
cgcgcccatt gcgcccggct gcgcccctg gatcttgct accgcagcgg tacgctgggg 5254
ctgctggcca gccggccagc accgcttcca acccgcgctt tcggcctgga attccccaac 5314
ccgggtgggccc tggcggccgg cctggacaag aacggcgagc atatcgatgc actgttcgcg 5374
ctgggctttg gctatgtcga aatcggcacg gtgaccccgc gcccgcaggc cggcaatccg 5434
cagccacggc tgttccgcgt gcccgagcac ctgggctgta tcaaccgcat gggtttcaac 5494
aatgccggcg tcgatgcgct ggtggccaat gtgcgcgcgg cacggcgtga ccgcggcac 5554
ctcggcattc acatcggcaa gaacaaggac accccaacg agctggccca taccgattac 5614
ctgacctgcc tggaaaagggt gtacgcgctg gccgactaca tcaccgtcaa catctcctcg 5674
cccaacaccg ccgggctgcg cgagctgcag gaagaacagg ccctgcgcga gctggtcagc 5734
cgctgcgcg agggccagga aaccctggcc gcacgccatg gcaagcgggt gccgatgctg 5794
gtcaaggctc cgccggacct gagcgatgcc gatgtcgatg ccgcccggcg tgtgctggca 5854
gagctgcagg tggacgggggt gatcgccacc aacaccacca tcgcgcgctg gggcatggaa 5914
aaccaccac tggccagcga ggccggcggc ctgtccgggg caccggtgat ggcgcgctcc 5974
accgcggtgc tgcgcccct gcgcaccgg ctgcccggagt cgatcccgt gatcggcgtc 6034
ggcggcatct gttccggggc tgatgcggcg gccaaatga gtgccggcgc gaccatggtg 6094
cagctctaca gcgggctggt ttaccgcggc ccggcactgg tcggcgaatg cgtcgaatcg 6154
atccgcccgc ggcgcgaagc gccctccagc ggggtagccc atctgtgagt acgcccgggt 6214
ggcagctgca ccacgatgtc gcaactgcaat caatgaacac cttcggggta gcggccaccg 6274
cgccgcgct gctgcgctg cacgacagcc aggcctgcc ggcggcgctg gcgcaccgg 6334
aagtagccgg acagccgctg atc 6357

```

&lt;210&gt; 4

&lt;211&gt; 510

&lt;212&gt; PRT

<213> *Flavobacterium lutescens*

&lt;400&gt; 4

```

Met Ser Phe Glu Leu Leu Lys Ala Leu Gly Leu Asp Ala Thr Asn Ser
  1           5           10           15

Gly Thr Tyr Leu Gly Asp Gly Glu Trp Ser Ser Ala Thr Gly Ala Gly
          20           25           30

Thr Ile Ser Pro Arg Asn Pro Thr Thr Gly Glu Val Ile Ala Gln Val
          35           40           45

Gln Ala Thr Thr Glu Ala Asp Tyr Glu Thr Ile Leu Ala Arg Ala Gln
          50           55           60

Gln Ala Phe Lys Val Trp Arg Thr Thr Pro Ala Pro Arg Arg Gly Glu
          65           70           75           80

Ala Ile Arg Leu Cys Gly Glu Ala Leu Arg Arg His Lys Asp Ala Leu
          85           90           95

Gly Ser Leu Val Ala Leu Glu Met Gly Lys Ser Lys Pro Glu Gly Asp
          100          105          110

```

Gly Glu Val Gln Glu Met Ile Asp Ile Ala Asp Phe Ala Val Gly Gln  
 115 120 125  
 Ser Arg Met Leu Tyr Gly Tyr Thr Met His Ser Glu Arg Pro Gly His  
 130 135 140  
 Arg Met Tyr Glu Gln Tyr Gln Pro Leu Gly Ile Val Gly Ile Ile Ser  
 145 150 155 160  
 Ala Phe Asn Phe Pro Val Ala Val Trp Ala Trp Asn Ser Phe Leu Ala  
 165 170 175  
 Ala Ile Cys Gly Asp Val Cys Ile Trp Lys Pro Ser Asn Lys Thr Pro  
 180 185 190  
 Leu Thr Ala Ile Ala Ser Met Arg Ile Cys Asn Glu Ala Leu Arg Glu  
 195 200 205  
 Gly Gly Phe Pro Asp Ile Phe Phe Leu Ile Asn Asp Ala Gly Thr Ala  
 210 215 220  
 Leu Ser Glu Lys Leu Val Glu Asp Lys Arg Val Pro Leu Ile Ser Phe  
 225 230 235 240  
 Thr Gly Ser Thr Gln Val Gly Arg Ile Val Asn Gln Lys Val Ala Ala  
 245 250 255  
 Arg Leu Gly Arg Cys Leu Leu Glu Leu Gly Gly Asn Asn Ala Ile Ile  
 260 265 270  
 Leu Asp Glu Thr Ala Asp Leu Lys Leu Ala Val Pro Gly Ile Val Phe  
 275 280 285  
 Gly Ala Val Gly Thr Ala Gly Gln Arg Cys Thr Thr Thr Arg Arg Leu  
 290 295 300  
 Ile Val His Glu Ser Ile Tyr Asp Asn Val Leu Ala Thr Leu Ile Lys  
 305 310 315 320  
 Ala Tyr Lys Gln Val Glu Gly Lys Ile Gly Asp Pro Leu Asp Ala Ala  
 325 330 335  
 Asn Leu Met Gly Pro Leu Asn Ser Pro Glu Ala Val Gln Gln Phe Leu  
 340 345 350  
 Ala Ser Ile Glu Lys Ala Lys Ala Ala Gly Gly Thr Val Gln Thr Gly  
 355 360 365  
 Gly Thr Ala Ile Asp Arg Pro Gly Asn Phe Val Leu Pro Ala Ile Val  
 370 375 380  
 Thr Gly Leu Lys Asn Ser Asp Glu Val Val Gln His Glu Thr Phe Ala  
 385 390 395 400  
 Pro Ile Leu Tyr Val Met Lys Tyr Ser Thr Leu Asp Glu Ala Ile Glu  
 405 410 415  
 Met Gln Asn Gly Val Pro Gln Gly Leu Ser Ser Ser Ile Phe Thr Thr  
 420 425 430

Asn Leu Lys Ala Ala Glu Lys Phe Leu Ser Ala Ala Gly Ser Asp Cys  
 435 440 445

Gly Ile Ala Asn Val Asn Ile Gly Thr Ser Gly Ala Glu Ile Gly Gly  
 450 455 460

Ala Phe Gly Gly Glu Lys Glu Thr Gly Gly Gly Arg Glu Ser Gly Ser  
 465 470 475 480

Asp Ala Trp Lys Val Tyr Met Arg Arg Gln Thr Asn Thr Ile Asn Tyr  
 485 490 495

Ser Asp Ser Leu Pro Leu Ala Gln Gly Ile Lys Phe Asp Leu  
 500 505 510

<210> 5

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 5

ttgatttgag cagattcgca ctgccattt

29

<210> 6

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 6

aaggttttcg acaaagtgac catttccca

29

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 7

ctgggtaccgc tcgatccggc tctgcaccgt

30

<210> 8

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 8  
ctggagctca ggcaggtgcg ggccacgtgt 30

<210> 9  
<211> 20  
<212> PRT  
<213> F. lutescens IFO 3084

<400> 9  
Ser Leu Leu Ala Pro Leu Ala Pro Leu Arg Ala His Ala Gly Thr Arg  
1 5 10 15

Leu Thr Gln Gly  
20

<210> 10  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer; N = I

<400> 10  
ccytgngtna rnckngtncc ngcrtgngcn cg 32

<210> 11  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer; N = I

<400> 11  
ccngcrtgng cncgnarngg ngcnarnggn gc 32

67566-1434

31

CLAIMS:

1. An isolated pure DNA containing a gene participating in the production of L-homoglutamic acid, derived from a bacterium belonging to Flavobacterium  
5 lutescens, or a DNA molecule which hybridizes with the complement of the gene under a stringent condition and has a function of recovering the L-homoglutamic acid-producing ability of a mutant of Flavobacterium lutescens which lacks the L-homoglutamic acid-producing ability,

10 wherein the gene participating in the production of L-homoglutamic acid is a DNA encoding at least one protein selected from the group consisting of (a) a protein having L-lysine: 2-oxoglutaric acid 6-aminotransferase (LAT) activity, and (b) a protein having L- $\Delta^1$ -piperidine-6-  
15 carboxylic acid (P6C) dehydrogenase activity,

wherein the DNA encoding the protein having the LAT activity (a) contains a continuous base sequence from base 801 to base 2276 of SEQ ID NO: 1 and the DNA encoding the protein having the P6C dehydrogenase activity  
20 (b) contains a continuous base sequence from base 2855 to base 4384 of SEQ ID NO: 3, and

wherein the hybridization under the stringent condition involves washing at 60°C in 0.5 x SSC.

2. The DNA according to claim 1 comprising the  
25 continuous base sequence from base 545 to base 2658 of SEQ ID NO: 1.

3. The DNA according to claim 1 consisting of the base sequence of SEQ ID NO: 1.



67566-1434

32

4. The DNA according to claim 1 comprising the continuous base sequence from base 2077 to base 4578 of SEQ ID NO: 3.
5. The DNA according to claim 1 consisting of the 5 base sequence of SEQ ID NO: 3.
6. The DNA according to claim 1 comprising the continuous base sequence from base 801 to base 2276 of SEQ ID NO: 1 and the continuous base sequence from base 2855 to base 4384 of SEQ ID NO: 3.
- 10 7. The DNA according to claim 1 comprising the continuous base sequence from base 545 to base 2658 of SEQ ID NO: 1 and the continuous base sequence from 2077 to base 4578 of SEQ ID NO: 3.
8. The DNA according to claim 1 consisting of the 15 base sequence of SEQ ID NO: 1 and the base sequence of SEQ ID NO: 3.
9. An autonomously replicative or integration replicative recombinant plasmid carrying the DNA as defined in any one of claims 1 to 8.
- 20 10. The recombinant plasmid according to claim 9 derived from the plasmid pBBR122.
11. The recombinant plasmid according to claim 10 which is contained in Flavobacterium lutescens IFO 3084 (pCF213) (FERM BP-6797).
- 25 12. A transformant obtained by transforming a bacterium belonging to the genus Flavobacterium as a host with the recombinant plasmid as defined in any one of claims 9 to 11.

67566-1434

33

13. The transformant according to claim 12, wherein the bacterium is Flavobacterium lutescens IFO 3084 or Flavobacterium lutescens SP.7-1 (FERM BP-5457).

14. Flavobacterium lutescens IFO 3084 (pCF213)  
5 (FERM BP-6797).

15. A process for producing L-homoglutamic acid, which comprises:

culturing in a medium a transformant obtained by transformation with the recombinant plasmid as defined in  
10 any one of claims 9 to 11;

during or after the culture, contacting the grown transformant with L-lysine or L- $\Delta^1$ -1-piperidine-6-carboxylic acid to convert it to L-homoglutamic acid; and

recovering the thus produced L-homoglutamic acid.

15 16. The process according to claim 15, wherein the transformant is the transformant as defined in claim 12 or 13.

17. The process according to claim 15 wherein the transformant is Flavobacterium lutescens IFO 3084 (pCF213)  
20 (FERM BP-6797).

SMART &amp; BIGGAR

OTTAWA, CANADA

PATENT AGENTS

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7





Fig. 8






Fig. 9

