



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
14.06.2006 Bulletin 2006/24

(51) Int Cl.:
C12N 15/86^(2006.01) C12N 7/01^(2006.01)
C12N 5/10^(2006.01)

(21) Application number: **98962352.5**

(86) International application number:
PCT/EP1998/007380

(22) Date of filing: **17.11.1998**

(87) International publication number:
WO 1999/025860 (27.05.1999 Gazette 1999/21)

(54) **VECTOR FOR TISSUE-SPECIFIC REPLICATION AND GENE EXPRESSION**

VEKTOR ZUR GEWEBESPEZIFISCHEN REPLIKATION UND EXPRESSION

VECTEUR POUR REPLICATION ET EXPRESSION GENIQUE A SPECIFICITE TISSULAIRE

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

(56) References cited:
WO-A-95/05835 WO-A-96/14875
WO-A-96/17053 WO-A-96/34969
WO-A-96/36365 WO-A-97/01358
WO-A-98/35028 WO-A-98/39465
WO-A-98/39466 WO-A-98/39467

(30) Priority: **19.11.1997 US 974391**

(43) Date of publication of application:
06.09.2000 Bulletin 2000/36

(73) Proprietor: **Cell Genesys Inc.
South San Francisco, CA 94080 (US)**

(72) Inventors:
• **CHANG, Yung-Nien**
Cockeysville, MD 21030 (US)
• **HALLENBECK, Paul, L.**
Gaithersburg, MD 20882 (US)
• **HAY, Carl, M.**
Damascus, MD 20872 (US)
• **STEWART, David, A.**
Eldersburg, MD 21784 (US)

- **HALLENBECK, P. L. ET AL: "Novel tumor specific replication competent adenoviral vectors for gene therapy of cancer." CANCER GENE THERAPY, (1996) VOL. 3, NO. 6 CONF. SUPPL., PP. S19-S20. MEETING INFO.: FIFTH INTERNATIONAL CONFERENCE ON GENE THERAPY OF CANCER SAN DIEGO, CALIFORNIA, USA NOVEMBER 14-16, 1996 ISSN: 0929-1903., XP002099752**
- **HALLENBECK, P. L. ET AL: "Novel tumor specific replication restricted adenoviral vector for gene therapy of liver cancer." CANCER GENE THERAPY, (NOV.-DEC., 1997) VOL. 4, NO. 6 CONF. SUPPL., PP. S22. MEETING INFO.: SIXTH INTERNATIONAL CONFERENCE ON GENE THERAPY OF CANCER SAN DIEGO, CALIFORNIA, USA NOVEMBER 20-22, 1997 ISSN: 0929-1903., XP002099753**

(74) Representative: **Hallybone, Huw George et al**
Carpmaels and Ransford,
43 Bloomsbury Square
London WC1A 2RA (GB)

EP 1 032 696 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionBackground of the Invention

5 **[0001]** The invention generally relates to cell-specific expression vectors. It particularly relates to targeted gene therapy using recombinant expression vectors and particularly adenovirus vectors. The invention specifically relates to modu-
latable replication-conditional expression vectors and methods for using them. Such vectors are able to selectively
10 replicate in a target cell or tissue to provide a therapeutic benefit in a tissue from the presence of the vector per se or from one or more heterologous gene products expressed from the vector and distributed throughout the tissue, and
which vectors are designed so that replication and gene expression from the vector can be modulated.

[0002] In such vectors, a gene essential for replication is placed under the control of a heterologous tissue-specific transcriptional regulatory sequence. Thus, replication is conditioned on the presence of a factor(s) that induces tran-
scription or the absence of a factor(s) that inhibits transcription of the gene by means of the transcriptional regulatory
15 sequence.

[0003] Preferred vectors contain a heterologous gene that produces a product that increases or inhibits viral replication. Such genes are useful for modulating viral replication and thus also for modulating expression of genes in the vector. With these vectors, therefore, the vector can be expressed in a desirable cell, target tissue can be selectively treated,
and replication and expression modulated.

[0004] The invention also relates to cells and/or methods to produce multiple heterologous gene products in high
20 quantity from essentially one promoter element.

[0005] The invention also relates to cells for producing high titers of the recombinant replication-conditional vector and/or high titers of a gene product expressed from the vector, useful for making clinically relevant gene products or
vector preparations for targeted gene therapy.

25 Background Art

[0006] The introduction of exogenous genes into cells in vitro or in vivo, systemically or in situ, has been of limited use for compositions in which it would be disadvantageous for non-target cells to take up the exogenous gene. One
30 strategy to overcome this problem is to develop administration procedures or vectors that target a specific cell-type. Using systemic administration, attempts have been made to direct exogenous genes to myocytes and muscle cells by
direct injection of DNA, to direct the exogenous DNA to hepatocytes using DNA-protein complexes, and to endothelial
cells using liposomes.

[0007] Using in situ administration, retroviral replication functions have been utilized to target cells that are actively
35 replicating.

[0008] Thus far, the ability to target cells has been limited, however, by the lack of cell-type specificity and low gene transfer efficiencies. The limited ability to target an exogenous gene to diseased cells in an organism, while avoiding
(eliminating) uptake of the gene by normal, untargeted cells, particularly has been an obstacle to developing effective
gene-transfer-based therapies for diseases in animals and humans.

[0009] One especially difficult challenge is targeting tumor cells. Many seemingly promising strategies for these cells,
40 moreover, are limited to one or a few cell-types.

[0010] The present invention, in one aspect, provides a way to deliver an exogenous gene efficiently, with high dis-
tribution in a tumor and in a controlled manner.

[0011] Adenoviruses are nonenveloped, regular icosohedrons. The protein coat (capsid) is composed of 252 cap-
somereres of which 240 are hexons and 12 are pentons. Most of the detailed structural studies of the adenovirus polypep-
45 tides have been done for adenovirus types 2 and 5. The viral DNA is 23.85×10^6 daltons for adenovirus 2 and varies slightly in size depending on serotype. The DNA has inverted terminal repeats and the length of these varies with the
serotype.

[0012] The replicative cycle is divided into early (E) and late (L) phases. The late phase defines the onset of viral DNA
replication. Adenovirus structural proteins are generally synthesized during the late phase. Following adenovirus infection,
50 host DNA and protein synthesis is inhibited in cells infected with most serotypes. The adenovirus lytic cycle with adenovirus 2 and adenovirus 5 is very efficient and results in approximately 10,000 virions per infected cell along with the synthesis
of excess viral protein and DNA that is not incorporated into the virion. Early adenovirus transcription is a complicated
sequence of interrelated biochemical events, but it entails essentially the synthesis of viral RNAs prior to the onset of
viral DNA replication.

[0013] The organization of the adenovirus genome is similar in all of the adenovirus groups and specific functions are
generally positioned at identical locations for each serotype studied. Early cytoplasmic messenger RNAs are comple-
mentary to four defined, noncontiguous regions on the viral DNA. These regions are designated (E1-E4). The early
transcripts have been classified into an array of immediate early (E1a), delayed early (E1 b, E2a, E2b, E3 and E4), and
55

intermediate (IVa2.IX) regions.

[0014] E1a is a transactivator of multiple gene products in adenovirus through activation of the E1b, E2, E3 and E4 promoters. The E1a region is involved in transcriptional transactivation of viral and cellular genes as well as transcriptional repression of other sequences. The E1a gene exerts an important control function on all of the other early adenovirus messenger RNAs. In normal tissues, in order to transcribe regions E1b, E2a, E2b, E3, or E4 efficiently, active E1 a product is required.

[0015] The E1b region is required for the normal progression of viral events late in infection. The E1b product acts in the host nucleus. Mutants generated within the E1b sequences exhibit diminished late viral mRNA accumulation as well as impairment in the inhibition of host cellular transport normally observed late in adenovirus infection (Berkner, K.L., *Biotechniques* 6:616-629 (1988)). E1 b is required for altering functions of the host cell such that processing and transport are shifted in favor of viral late gene products. These products then result in viral packaging and release of virions. E1 b produces a 19 kD protein that prevents apoptosis. E1 b also produces a 55 kD protein that binds to p53.

[0016] For a complete review on adenoviruses and their replication, see Horwitz, M.S., *Virology* 2d ed., Fields, B.N., eds., Raven Press Limited, New York (1990), Chapter 60, pp. 1679-1721.

[0017] Adenovirus provides advantages as a vector for adequate gene delivery for the following reasons. It is a double stranded DNA nonenveloped virus with tropism for the human respiratory system and gastrointestinal tract. It causes a mild flu-like disease. Adenoviral vectors enter cells by receptor mediated endocytosis. The large (36 kilobase) genome allows for the removal of genes essential for replication and nonessential regions so that foreign DNA may be inserted and expressed from the viral genome. Adenoviruses infect a wide variety of cell types in vivo and in vitro. Adenoviruses have been used as vectors for gene therapy and for expression of heterologous genes. The expression of viral or foreign genes from the adenovirus genome does not require a replicating cell. Adenovirus vectors rarely integrate into the host chromosome; the adenovirus genome remains as an extrachromosomal element in the cellular nucleus. There is no association of human malignancy with adenovirus infection; attenuated strains have been developed and have been used in humans for live vaccines.

[0018] For a more detailed discussion of the use of adenovirus vectors for gene therapy, see Berkner, K.L., *Biotechniques* 6:616-629 (1988); Trapnell, B.C., *Advanced Drug Delivery Reviews* 12:185-199 (1993).

[0019] Adenovirus vectors are generally deleted in the E1 region of the virus. The E1 region may then be substituted with the DNA sequences of interest. It was pointed out in a recent article on human gene therapy, however, that "the main disadvantage in the use of adenovirus as a gene transfer vector is that the viral vector generally remains episomal and does not replicate, thus, cell division leads to the eventual loss of the vector from the daughter cells" (Morgan, R.A., et al., *Annual Review of Biochemistry* 62:191-217 (1993)) (emphasis added).

[0020] Non-replication of the vector leads not only to eventual loss of the vector without expression in most or all of the target cells but also leads to insufficient expression in the cells that do take up the vector, because copies of the gene whose expression is desired are insufficient for maximum effect. The insufficiency of gene expression is a general limitation of all non-replicating delivery vectors. Thus, it is desirable to introduce a vector that can provide multiple copies of a gene and hence greater amounts of the product of that gene. The present invention overcomes the disadvantages discussed above by providing a tissue-specific, and especially a tumor-specific replicating vector, multiple DNA copies, and thus increased amounts of gene product.

[0021] Adenoviral vectors for recombinant gene expression have been produced in the human embryonic kidney cell line 293 (Graham, F.L. et al., *J. Gen. Virol.* 36:59-72 (1977)). This cell line, initially transformed with human adenovirus 5, now contains the left end of the adenovirus 5 genome and expresses E1. Therefore, these cells are permissive for growth of adenovirus 2 and adenovirus 5 mutants defective in E1 functions. They have been extensively used for the isolation and propagation of E1 mutants. Therefore, 293 cells have been used for helper-independent cloning and expression of adenovirus vectors in mammalian cells. E1 genes integrated in cellular DNA of 293 cells are expressed at levels which permit deletion of these genes from the viral vector genome. The deletion provides a nonessential region into which DNA may be inserted. For a review, see, Young, C.S.H., et al. in *The Adenoviruses*, Ginsberg, H.S., ed., Plenum Press, New York and London (1984), pp. 125-172.

[0022] However, 293 cells are subject to severe limitations as producer cells for adenovirus vectors. Growth rates are low. Titres are limited, especially when the vector produces a heterologous gene product that proves toxic for the cells. Recombination with the viral E1 sequence in the genome can lead to the contamination of the recombinant defective virus with unsafe wild-type virus. The quality of certain viral preparations is poor with regard to the ratio of virus particle to plaque forming unit. Further, the cell line does not support growth of more highly deleted mutants because the expression of E1 in combination with other viral genes in the cellular genome (required to complement the further deletion), such as E4, is toxic to the cells. Therefore, the amount of heterologous DNA that can be inserted into the viral genome is limited in these cells. It is desirable, therefore, to produce adenovirus vectors for gene therapy in a cell that cannot produce wild-type recombinants and can produce high titres of high-quality virus.

[0023] It is also desirable to control the replication of a therapeutic vector by other than endogenous cellular factors present within the cells to be treated. A high degree of replication could be disadvantageous to the patient, in that vector

particles could be released into the bloodstream. This could have toxic side effects in tissues clearing the virus, such as kidneys, liver, lung and spleen, even if the vector does not replicate in non-treated tissue. Furthermore, while current therapeutic vectors have not been shown to replicate at low exposure to normal cells, a high exposure, as potentially caused from a high degree of replication and release of virus at the site of the treated tissue, may cause replication even

in normal cells. For example, adenoviruses devoid of E1a, given in a sufficiently high dose, will replicate in normal cells. [0024] Accordingly, it would be highly advantageous to be able to control the level of replication by adding a compound that could dampen replication if replication were excessive.

[0025] Another problem in the art is providing more than one gene product controlled by a heterologous regulatory sequence, for example a promoter, on a viral vector (such as cytokines, TK, or other cytotoxic genes, or heat-shock proteins). Thus, in a specific milieu, a particular combination of genes may offer therapeutic advantage. Accordingly, it would be desirable to have several genes in one vector and specifically expressed in the cells to be treated, for example, in a tumor cell. A limitation, however, is that one cannot provide multiple copies of the same promoter to drive each gene, because space would be wasted and the excess nucleotide stretches might not even be accommodated by the virus. Furthermore, during the cloning and replication, homologous recombination could produce deletions between identical promoters and therefore destroy the vector.

[0026] WO 97/01358 discloses host cell-specific adenoviral vectors in which at least one essential replication function and at least one transgene are under the control of a target-cell-specific response element.

[0027] WO 96/34969 discloses adenoviral vectors that are replication-competent in certain tumor cells in which at least one gene essential for viral replication is placed under the control of a tissue/tumor-specific regulatory element that may further comprise a transgene which may also be regulated by the tissue/tumor-specific regulatory element.

[0028] WO 96/17053 discloses adenoviral vectors in which at least one gene essential for viral replication is placed under the control of a heterologous tissue-specific transcriptional regulatory sequence that may further comprise a transgene which is under the control of a promoter other than a tissue/tumor-specific regulatory element.

[0029] Hallenbeck et al., Cancer Gene Therapy 3: Conf Suppl. pp. S19-S20 (1996) discloses replication-competent, adenoviral vectors for killing hepatic tumor cells in which the E1a gene is placed under the control of a modified alpha-fetoprotein which is expressed primarily in hepatic cells.

Summary of the Invention

[0030] In view of the limitations discussed above, a general object of the invention is to provide novel expression vectors for tissue-specific vector replication and gene expression from the replicating vector.

[0031] The invention provides a replication conditional adenoviral vector comprising:

(a) a heterologous transcriptional regulatory sequence (TRS) operably linked to the coding sequence of an adenoviral gene essential for replication (GER), and

(b) at least one additional coding sequence encoding a heterologous gene product, wherein said additional coding sequence is operably linked to a homologous transcriptional regulatory sequence (TRS) that is transactivated by the gene product of said adenoviral gene essential for replication.

[0032] Other aspects of the invention are set out in the appended claims.

[0033] Accordingly, an expression vector is provided in which the expression of multiple genes on the vector can be controlled and coordinated through the expression of a gene that is essential for replication, so that the expression of each of the multiple genes is conditional upon vector replication, replication depending upon the tissue-specific expression of the gene product.

[0034] Another object of the invention is to provide tissue-specific treatment of abnormal tissue. Thus, a further object of the invention is to provide a method to selectively distribute a vector in vivo in a target tissue, such that a greater number of cells contain the vector than would with a non-replicating vector, and spread of the vector is avoided or significantly reduced in non-target tissue.

[0035] Accordingly, there is disclosed herein a method for selectively distributing a vector in a target tissue by introducing the replication-conditional vector of the present invention into a target tissue that allows modulatable replication of the vector.

[0036] For providing tissue-specific treatment, another object of the invention is to selectively distribute a polynucleotide in a target tissue in vivo.

[0037] Accordingly, there is disclosed herein a method for selectively distributing a polynucleotide in a target tissue in vivo by introducing the replication-conditional vectors of the present invention, containing the polynucleotide, into the target tissue that allows modulatable replication of the vector. The polynucleotide includes the entire vector or parts thereof, such as one or more heterologous genes.

[0038] For providing tissue-specific treatment, a further object of the invention is to selectively distribute one or more

heterologous gene products in a target tissue.

[0039] Accordingly, the replication-conditional vectors of the present invention are constructed so that they contain one or more heterologous DNA sequences encoding a gene product that is expressed from the vector. When the vectors replicate in the target tissue, effective quantities of the desired gene product are also produced in the target tissue.

[0040] Another object of the invention, especially where tissue-specific treatment is involved, is to be able to modulate vector replication, that is, control the level of vector replication. When vector replication proceeds at levels that are undesirable, and particularly levels that may interfere with treatment, an object of the invention is to dampen or decrease the levels of replication by a desired degree. If the levels fall below the desirable amount, an object of the invention is also to be able to allow replication to increase to desirable levels by desired degrees.

[0041] Accordingly, there is disclosed herein a method to modulate vector replication during treatment or otherwise (e.g., in producer cells), by providing a gene encoding a gene product that has the ability to interfere with vector replication. When replication is undesirably high, it can be decreased by means of this gene product.

[0042] Thus, the invention is further directed to vectors further containing a gene encoding a gene product that can be used to modulate vector replication.

[0043] Modulating replication also modulates the expression of heterologous genes contained in the vector. Thus, the expression of such genes can be induced, decreased, increased, or eliminated using the methods and vectors described herein.

[0044] Another object of the invention is to provide a method to identify abnormal tissue that can then be treated by the vectors of the present invention. Therefore, a further object of the invention is to identify a tissue in which the replication-conditional vectors of the present invention can be replicated by means of the heterologous transcriptional regulatory sequence contained on the vector so that the tissue can subsequently be treated with the vector, and in which tissue the replication can be modulated.

[0045] Accordingly, there is also disclosed herein a method wherein the replication-conditional vectors of the present invention are exposed to a given abnormal tissue. If that tissue allows replication and modulation, then replication of the vector will occur and can be detected. Following identification of such a tissue, targeted treatment of that tissue can be effected by tissue-specific transcription and the consequent vector replication in that tissue in vivo.

[0046] Thus, a method is disclosed for assaying vector utility for tissue treatment comprising the steps of removing a tissue biopsy from a patient, explanting the biopsy into tissue culture, introducing a replication-conditional vector into the cells of the biopsy, and assaying for modulatable vector replication in the cells.

[0047] Another object of the invention is to provide producer cell lines for vector production. Preferably, the cell lines have one or more of the following characteristics: high titer virus production, resistance to toxic effects due to heterologous gene products expressed in the vector, lack of production of wild-type virus contaminating the virus preparation and resulting from recombination between integrated viral sequences and vector sequences, growth to high density and in suspension, unlimited passage potential, high growth rate, and by permitting the growth of highly deleted viruses that are impaired for viral functions and able to accommodate large pieces of heterologous DNA.

[0048] Accordingly, in a further embodiment of the invention, cell lines are provided containing the replication-conditional vector of the present invention, the cells of which allow modulatable replication of the vector or is deficient in a transcription-inhibiting factor(s) that prevents replication of the vector.

[0049] In further embodiments of the invention, the cell lines contain nucleic acid copies of the replicated vector. In other embodiments, the cell lines contain virions produced in the cell by replication in the cell of the replication-conditional vector.

[0050] In further embodiments, a method is provided for producing a replication-conditional vector or virion comprising the steps of culturing a producer cell line described above and recovering the vector or virion from the cells.

[0051] There is also disclosed a method for producing replication-conditional virions free of wild-type virions or viral vectors free of wild-type vectors, comprising the steps of culturing a producer cell line described above and recovering the replication-deficient virions or vectors from the cells.

[0052] In a preferred methods of treatment and diagnosis, the tissue is abnormally proliferating, and especially is tumor tissue. However, the methods are also relevant to other abnormal tissue as described herein.

[0053] The replication-conditional vector of the invention is a DNA tumor viral vector, in particular an adenovirus vector.

[0054] However, it is to be understood that potentially any vector source is useful if it contains a gene essential for replication that can be operably linked to a tissue-specific transcriptional regulatory sequence.

[0055] In further methods of treatment and diagnosis, the vector is introduced into the cell or tissue by infection.

[0056] Replication can be vector nucleic acid replication alone or can also include virus replication (i.e., virion production). Thus, either DNA or virions or both may be distributed in the target tissue.

[0057] In a further preferred embodiment of the invention, a gene in the adenovirus E1 region is operably linked to the tissue-specific heterologous transcriptional regulatory sequence. Preferably, the E1a, E1b or E2a gene is operably linked to the tissue-specific transcriptional regulatory sequence.

[0058] The vector encodes one or more heterologous gene products. These heterologous gene products are expressed

from the vector replicating in the target tissue. The heterologous gene product may be operably linked to its own promoter or may be operably linked to a transcriptional regulatory sequence from another gene. Regardless, the expression of the heterologous gene product may be controlled by E1 a or another transactivator which in turn is regulated by the desired tumor-specific promoter.

5 **[0059]** Expression of one or more heterologous gene products depends upon expression of the gene essential for replication, which in turn is operably linked to the tissue-specific transcriptional regulatory sequence. Accordingly, when the gene essential for replication is expressed, it causes expression of one or more heterologous gene products, by transactivation. In this configuration, such a heterologous gene is activated/induced only when the vector replicates. This is because activation of the tissue-specific regulatory sequence causes expression of the gene product essential for replication which then causes both replication and activation of the expression of the one or more heterologous genes by its transactivation function. Accordingly, not only is expression of the heterologous gene activated, but expression is also amplified because when the vector replicates, each additional copy of the vector also contains a copy of the heterologous gene.

10 **[0060]** In a highly preferred embodiment, the E1 a gene, being operably linked to a heterologous tissue-specific transcriptional regulatory sequence, controls the expression of one or more heterologous genes under the control of promoters that are transactivated by the E1a gene product. Thus, when the E1a gene is expressed, viral replication occurs and gene expression of one or more heterologous genes also occurs. Thus, expression of those genes is controlled at the level of replication and transactivation, such that an expression-amplifying effect is obtained.

15 **[0061]** In a further embodiment, one or more of the heterologous genes is operably linked to a tissue-specific regulatory sequence, such as the same transcriptional regulatory sequence to which the gene essential for replication is operably linked. Accordingly, the one or more heterologous genes are then activated only in a specific tissue. When the tissue-specific transcriptional regulatory sequence is the same one as that to which the gene essential for replication is operably linked, activation of the one or more heterologous genes occurs only when the vector replicates. Accordingly, as above, the one or more heterologous genes is both activated and amplified.

20 **[0062]** In other embodiments, the one or more heterologous genes is amplified when the vector replicates but not necessarily activated. This is when these genes are under the control of their own or other constitutive promoters. In this case, there is always a basal level of expression, but when the vector replicates, this expression is amplified because of expression from each new copy of the vector.

25 **[0063]** In the context of coordinate control (one or more heterologous genes under the control of the same transcriptional regulatory sequence, the same scenario of activation and amplification as discussed above applies.

30 **[0064]** In a further embodiment, the vector is not used to selectively destroy a cell or tissue type, such as a tumor, but is used to provide a gene product or products at specific levels that are physiologically desirable. An example of such gene product is insulin. Thus, using replication to control gene expression in order to achieve a very specific amount of gene expression is provided by the invention. Accordingly, the vector encodes a gene product which regulates the degree of virus replication, and hence gene expression of other gene products whose amounts must be controlled. An example of such product is one that would inhibit DNA polymerase or nucleotide synthesis. Accordingly, an excess of this product will down-regulate the vector replication. Gene products are also encompassed that will increase the level of vector replication if sufficient gene product is not produced.

35 **[0065]** Thus, according to the invention, one or more of the heterologous gene products is useful for modulating viral replication. It may directly or indirectly control viral replication as, for example, in the case of thymidine kinase in which viral replication is negatively affected and modulated by the addition of ganciclovir.

40 **[0066]** With respect to modulating gene expression, the vector provides for modulating the expression of heterologous genes in any of the configurations described above by modulating replication of the vector. This is preferably done via thymidine kinase or a gene product functioning as thymidine kinase does.

45 **[0067]** In a highly preferred embodiment, the thymidine kinase coding region is operably linked to the E3 promoter in an adenoviral vector. The E1a coding region is under the control of a heterologous tissue-specific promoter. Accordingly, when the tissue-specific promoter is activated, the E1a gene product is produced. This gene product then transactivates the E3 promoter, so that the TK gene is expressed. Further, since the E1a gene activates viral replication, and the E3 gene is not essential for replication, more copies of the viral vector are present to allow greater expression of the TK gene. Ganciclovir can then be added to modulate vector replication. If after adding the ganciclovir, replication falls below a desirable level, the amount of ganciclovir can be decreased to allow an increase in vector replication and so forth, so that desirable levels are permitted within any particular time frame. One example of desirable modulation is to decrease vector replication in a non-target cell.

50 **[0068]** In a further embodiment of the invention, a heterologous gene product is toxic for the target tissue.

55 **[0069]** In a further embodiment of the invention, the toxic heterologous gene product acts on a non-toxic prodrug, converting the non-toxic prodrug into a form that is toxic for the target tissue. Preferably, the toxin has anti-tumor activity or eliminates cell proliferation.

[0070] In preferred embodiments of the invention, the transcriptional regulatory sequence is a promoter. Preferred

promoters include, but are not limited to, CEA, DF3, α -fetoprotein, Erb-B2, surfactant, and especially lung surfactant, tyrosinase promoter, and endothelial-specific promoters. However, any genetic control region that controls transcription of the essential gene can be used to activate (or derepress) the gene. Thus, other genetic control elements, such as enhancers, repressible sequences, and silencers, can be used to regulate replication of the vector in the target cell. The only requirement is that the genetic element be activated, derepressed, enhanced, or otherwise genetically regulated by factors in the host cell and, with respect to methods of treatment, not in the non-target cell.

[0071] Preferred enhancers include the DF3 breast cancer-specific enhancer and enhancers from viruses and the steroid receptor family. Other preferred transcriptional regulatory sequences include NF1, SP1, AP1, and FOS/JUN.

[0072] In further embodiments, promoters are not necessarily activated by factors in the target tissue, but are derepressed by factors present in the target tissue. Thus, in the target tissue, repression is lifted.

[0073] Transcriptional regulatory factors include, but are not limited to, transactivating factors produced by endogenous viral sequences such as from CMV, HIV, EBV, HSV, SV40, and other such viruses that are pathogenic in mammals and, particularly, in humans.

[0074] Methods for making such vectors are well known to the person of ordinary skill in the art. The art adequately teaches the construction of recombinant vectors with deletions or modifications in specific coding sequences and operable linkage to a heterologous transcription control sequence such that expression of a desired coding region is under control of the heterologous transcriptional regulatory sequence. Many viral sequences have been adequately mapped such that it is routine to identify a gene of choice and use appropriate well known techniques (such as homologous recombination of the virus with deleted or otherwise modified plasmids, or ligation of the two) to construct the vectors for tissue-specific replication and expression.

Brief Description of the Figures

[0075]

Figure 1 A. Cloning of pAVE1a02i.

Figure 1 B. Diagram of pAF(AB)2(S)6CAT. The map shows the enhancer regions (AB) in opposite orientation.

Figure 1C. Diagram of pAvE1a04i as predicted from a plasmid map and sequence information describing pAF(AB)2(S)6CAT, and pAvE1a06i as determined by restriction digestion mapping and sequencing to be the correct structure.

Figure 2A. Diagram of Clal digestion of Av1 LacZ.

Figure 2B. Diagram of co-transfection of 293 cells with pAvE1 a06i and Av1LacZ Clal digest (30,884 bp), to produce AvE1a06i.

Figure 2C. Amplification of AvE1 a06i virus on S8 cells deleted one direct enhancer (AB) from the promoter. The new construct is designated AvE1a04i.

Figure 3A. Cloning of the plasmid pREpacTK.

Figure 3B. Cloning of the plasmid pREpacTKm.

Figure 3C. Construction of AV5E1aTK01i.

Figure 4A. Digestion of Av1 E1 a04i.

Figure 4B. Digestion of Av5E1aTK01i.

Figure 4C. Construction of Av15E1aTK04i using the digestion products of Figures 4A and 4B.

Figure 5. Cytopathic effect of Av15E1aTK04i.

Figure 6. GCV inhibition of Av5E1aTk01i Replication In Vitro. A549 cells were transduced with either Add1327 or Add1327Tk01i. Cells were then treated with 10 μ M GCV for 5 days. Following the 5 day incubation, cells were harvested, washed in HBSS, and then freeze/thawed to prepare a crude viral lysate. Titers were then performed by standard TCID50 assays on 293 cells to determine viral titer. The experiment was repeated three times. Results

show that while GCV had no effect on an adenovirus not carrying the HSV-TK gene, it caused a greater than 4 order of magnitude drop in titer of a vector carrying the HSV-TK gene.

Figure 7. Diagram of Possible Configurations of Heterologous Genes in Expression Vectors.

5

Detailed Description of the Preferred Embodiments

[0076] The term "abnormally proliferating" is intended to mean a cell having a higher mitotic index than its normally-functioning counterpart, such that there is an abnormal accumulation of such cells.

10 **[0077]** The term "anti-tumor activity" is intended to mean any activity which inhibits, prevents, or destroys the growth of a tumor.

[0078] As used herein, the term "cytotoxic gene" refers to a gene that encodes a protein which either alone or in combination with other agents is lethal to cell viability. Examples of cytotoxic genes which alone are lethal include toxins such as pertussis toxin, diphtheria toxin and the like.

15 **[0079]** Examples of cytotoxic genes which are used in combination with other agents to achieve cell lethality include, for example, herpes simplex-1 thymidine kinase and cytosine deaminase. The subject is then administered an effective amount of a therapeutic agent, which in the presence of the -cytotoxic gene is toxic to the cell. In the specific case of thymidine kinase, the therapeutic agent is a thymidine kinase substrate such as ganciclovir (GCV), 6-methoxypurine arabinonucleoside (araM), or a functional equivalent thereof. Both the thymidine kinase gene and the thymidine kinase metabolite must be used concurrently to be toxic to the host cell. However, in its presence, GCV is phosphorylated and becomes a potent inhibitor of DNA synthesis whereas araM gets converted to the cytotoxic anabolite araATP. Other genes can be used as well in combination with the corresponding therapeutic agent. Such other gene and therapeutic agent combinations are known by one skilled in the art. Another example would be the vector of this invention expressing the enzyme cytosine deaminase. Such vector would be used in conjunction with administration of the drug 5-fluorouracil (Austin and Huber, 1993), or the recently described E. coli Deo Δ gene in combination with 6-methyl-purine-2'-deosribo-
20 nucleoside (Sorscher et al., 1994).

25 **[0080]** The term "distributing" is intended to mean the spreading of a vector and its attendant heterologous gene (product) (when present on the vector) throughout a target tissue, and especially throughout abnormally proliferating tissue (non-malignant or malignant). The object of the distribution is to deliver the vector, gene product or the effects of the gene product (as by a bystander effect, for example) to substantially all or a significant number of cells of the target tissue, so as to treat substantially the entire target tissue.

30 **[0081]** The term "enhancer" is used according to its art-recognized meaning. It is intended to mean a sequence found in eukaryotes and certain eukaryotic viruses which can increase transcription from a gene when located (in either orientation) up to several kilobases from the gene being studied. These sequences usually act as enhancers when on the 5' side (upstream) of the gene in question. However, some enhancers are active when placed on the 3' side (down-
35 stream) of the gene. In some cases, enhancer elements can activate transcription from a gene with no (known) promoter.

[0082] The term "functional inactivation" is intended to mean a genetic lesion that prevents the normal activity of a gene product. Thus, functional inactivation could result from a mutation in the gene encoding the gene product. Such a lesion includes insertions, deletions, and base changes. Alternatively, functional inactivation may occur by the abnormal interaction of the normal gene product with one or more other cellular gene products which bind to or otherwise prevent the functional activity of said gene product. Thus, the gene product may be a protein produced from a normal gene but which cannot perform its ordinary and normal function because of an interaction with a second factor.

40 **[0083]** The term "gene essential for replication" refers to a genetic sequence whose transcription is required for the vector to replicate in the target cell.

45 **[0084]** The term "gene product" is intended to mean DNA, RNA, protein, peptides, or viral particles. Thus, the distribution, for the purposes of the invention, is of any of these components.

[0085] The term "heterologous" means a DNA sequence not found in the native vector genome. Thus, for example, when the vector is based on the adenovirus genome, heterologous DNA sequences are those that are not found in the native adenovirus genome. With respect to a "heterologous transcriptional regulatory sequence," "heterologous" indicates that the transcriptional regulatory sequence is not naturally ligated to the DNA sequence for the gene essential for replication of the vector.

[0086] The term "promoter" is used according to its art-recognized meaning. It is intended to mean the DNA region, usually upstream to the coding sequence of a gene or operon, which binds RNA polymerase and directs the enzyme to the correct transcriptional start site.

55 **[0087]** The term "replication" means duplication of a vector. This duplication, in the case of viruses, can occur at the level of nucleic acid, or at the level of infectious viral particle. In the case of DNA viruses, replication at the nucleic acid level is DNA replication. In the case of RNA viruses, nucleic acid replication is replication into plus or minus strand (or both). In the case of retroviruses, replication at the nucleic acid level includes the production of cDNA as well as the

further production of RNA viral genomes. The essential feature is nucleic acid copies of the original viral vector. However, replication also includes the formation of infectious DNA or RNA viral particles. Such particles may successively infect cells in a given target tissue thus distributing the vector through all or a significant portion of the target tissue.

5 [0088] The term "replication-conditional vector" refers to a vector which when introduced into a tissue will not replicate unless a transcriptional regulatory sequence in that vector is activated or derepressed in that tissue. That is, replication depends upon transcription by means of that transcriptional regulatory sequence. Such a vector is replication-conditional as described because it has been modified in the following manner. A gene that is essential for replication has been modified by replacing the transcriptional regulatory sequence on which transcription of that gene normally depends with a heterologous transcriptional regulatory sequence. This transcriptional regulatory sequence depends upon the presence of transcriptional regulatory factors or the absence of transcriptional regulatory inhibitors. The presence of these factors in a given tissue or the absence of such inhibitors in a given tissue provides the replication-conditional. Accordingly, the native transcriptional regulatory sequence may be replaced with the heterologous transcriptional regulatory sequence. Alternatively, the native transcriptional regulatory sequence may be disabled or rendered dysfunctional by partial removal (deletion) or other mutation (one or more base changes, insertions, inversions, etc.).

10 [0089] The gene sequence may be a coding sequence. It may contain one or more open reading frames, as well as intron sequences. However, such a sequence is not limited to a coding sequence, but includes sequences that are transcribed into RNA, which RNA is itself essential for vector replication. The essential feature is that the transcription of the gene sequences does not depend on the native transcriptional regulatory sequences.

15 [0090] The term "silencer," used in its art-recognized sense, means a sequence found in eucaryotic viruses and eucaryotes which can decrease or silence transcription of a gene when located within several kilobases of that gene.

20 [0091] The term "tissue-specific" is intended to mean that the transcriptional regulatory sequence to which the gene essential for replication is operably linked functions specifically in that tissue so that replication proceeds in that tissue. This can occur by the presence in that tissue, and absence in non-target tissues, of positive transcription factors that activate the transcriptional regulatory sequence. It can also occur by the absence of transcription inhibiting factors that normally occur in non-target tissues and prevent transcription as a result of the transcription regulatory sequence. Thus, when transcription occurs, it proceeds into the gene essential for replication such that in that target tissue, replication of the vector and its attendant functions occur.

25 [0092] As described herein, tissue specificity is particularly relevant in the treatment of the abnormal counterpart of a normal tissue. Such counterparts include, but are not limited to, liver tissue and liver cancer, breast tissue and breast cancer, melanoma and normal skin tissue. Tissue specificity also includes the presence of an abnormal tissue type interspersed with normal tissue of a different tissue type, as for example in the case of metastases of colon cancer, breast cancer, and the like, into tissue such as liver. In this case, the target tissue is the abnormal tissue, and tissue specificity reflects the restriction of vector replication to the abnormal tissue interspersed in the normal tissue. It is also to be understood that tissue specificity, in the context of treatment, is particularly relevant in vivo. However, as described herein, ex vivo treatment and tissue replacement also falls within the concept of tissue specificity according to the present invention.

30 [0093] The term "transcriptional regulatory function" or "transcriptional regulatory factor" is intended to mean any cellular function whose presence activates or represses the heterologous transcriptional regulatory sequence described herein or whose absence permits transcription as a result of the transcriptional regulatory sequences described herein. It is understood that in the given target tissue, a tissue that "lacks the transcriptional regulatory factor" or is "deficient" in the transcriptional regulatory factor could refer to either the absence of the factor or the functional inactivation of the factor in the target tissue.

35 [0094] The term "transcriptional regulatory sequence" is used according to its art-recognized meaning. It is intended to mean any DNA sequence which can, by virtue of its sequence, cause the linked gene to be either up- or down-regulated in a particular cell. In one embodiment of the present invention, the native transcriptional regulatory sequence is completely deleted from the vector and replaced with a heterologous transcriptional regulatory sequence. The transcriptional regulatory sequence may be adjacent to the coding region for the gene that is essential for replication, or may be removed from it. Accordingly, in the case of a promoter, the promoter will generally be adjacent to the coding region. In the case of an enhancer, however, an enhancer can be found at some distance from the coding region such that there is an intervening DNA sequence between the enhancer and the coding region. In some cases, the native transcriptional regulatory sequence remains on the vector but is non-functional with respect to transcription of the gene essential for replication.

40 [0095] Various combinations of transcriptional regulatory sequences can be included in a vector. One or more may be heterologous. Further, one or more may have the tissue-specificity. For example, a single transcriptional regulatory sequence could be used to drive replication by more than one gene essential for replication. This is the case, for example, when the gene product of one of the genes drives transcription of the further gene(s). An example is a heterologous promoter linked to a cassette containing an E1a coding sequence (E1a promoter deleted) and the entire E1b gene. In such a cascade, only one heterologous transcriptional regulatory sequence may be necessary. Thus, a single transcrip-

tional regulatory sequence may drive transcription of the further gene(s) as one mRNA, as the genes are linked by an internal ribosome entry site (IRES). When genes are individually (separately) controlled, however, more than one transcriptional regulatory sequence can be used if more than one such gene is desired to control replication.

5 **[0096]** The vectors of the present invention, therefore, also include transcriptional regulatory sequence combinations wherein there is more than one heterologous transcriptional regulatory sequence, but wherein one or more of these is not tissue-specific. For example, one transcriptional regulatory sequence can be a basal level constitutive transcriptional regulatory sequence. For example, a tissue-specific enhancer can be combined with a basal level constitutive promoter.

10 **[0097]** The term "tissue-specific gene regulatory region" or "tissue-specific regulatory region" or "tissue-specific promoter" or "tissue-specific promoter/enhancer" refers to transcription and/or translation regulatory regions that function selectively or preferentially in a specific cell type. Selective or preferential function confers specificity to the gene therapy treatment since the therapeutic gene will be primarily expressed in a targeted or specific cell type. Specific regulatory regions include transcriptional, mRNA maturation signals and translational regulatory regions that are cell type specific. Transcriptional regulatory regions for tumors include, for example, promoters, enhancers, silencers, or artificial control elements added to the vector. Examples are steroid hormone receptor and/or response elements controlled by steroid hormones. Specific examples of such transcriptional regulatory regions for tumors include the promoter/enhancer elements for alpha-fetoprotein, carcinoembryonic antigen and prostate specific antigen. RNA processing signals include, for example, tissue-specific intron splicing signals, whereas translational and regulatory signals can include, for example, mRNA stability signals and translation initiation signals. Thus, specific regulatory regions include all elements that are essential for the production of a mature gene product in a specific cell type.

20 **[0098]** The invention is particularly directed to neoplastic cells. The neoplastic phenotype is characterized by altered morphology, faster growth rates, higher saturation density, growth in soft agar and tumorigenicity. The therapeutic genes described herein encode proteins which exhibit this activity. "Tumorigenicity" is intended to mean having the ability to form tumors or capable of causing tumor formation and is synonymous with neoplastic growth. "Malignancy" is intended to describe a tumorigenic cell having the ability to metastasize and endanger the life of the host organism. "Hyperproliferative phenotype" is intended to describe a cell growing and dividing at a rate beyond the normal limitations of growth for that cell type. "Neoplastic" also is intended to include cells lacking endogenous functional tumor suppressor protein or the inability of the cell to express endogenous nucleic acid encoding a functional tumor suppressor protein.

25 **[0099]** The invention can provide tumor-specific replication competent vectors wherein the gene regulatory regions include, but are not limited to alpha-fetoprotein promoter/enhancer, the carcinoembryonic antigen promoter/enhancer, the tyrosinase promoter/enhancer and the prostate-specific antigen promoter/enhancer. It is to be understood that any regulatory or tumor-specific sequence can be used. For other diseases such as inflammatory conditions, the inducer could be TNF- α and the responding regulatory element the interleukin-6 (IL-6) promoter. The therapeutic gene can encode interleukin-10 (IL-10) or another anti-inflammatory cytokine.

30 **[0100]** The vectors useful in the methods of this invention can replicate specifically in specific tumor cells. The tumor specificity results from the incorporation of tumor-specific gene regulatory regions which drive the expression of one or more genes which are essential for replication. Such elements include, for example, the alpha-fetoprotein promoter/enhancer, the carcinoembryonic antigen promoter/enhancer, the tyrosine promoter/enhancer and the prostate-specific antigen promoter/enhancer. Each of these gene regulatory regions functions preferentially in specific tumor cell types. For example, the alpha-fetoprotein promoter/enhancer functions preferentially in hepatocellular carcinoma tumor cells. The carcinoembryonic antigen promoter/enhancer functions preferentially in colon cancer and breast tumor cells. The prostate-specific antigen promoter/enhancer functions in prostate tumor cells. The tyrosine promoter enhancer preferentially functions in melanoma tumor cells. Thus, the invention provides for the treatment of cancers including, for example, breast cancer, colorectal cancer, hepatocellular carcinoma and melanoma cancer.

35 **[0101]** The present invention is generally directed to an expression vector capable of expressing one or more heterologous genes in a modulated and tissue-specific manner wherein a first coding sequence, derived from a gene that is essential for vector replication is operably linked to a tissue-specific transcriptional regulatory sequence, and wherein the vector contains one or more additional heterologous coding sequences.

40 **[0102]** The first coding sequence for a gene product essential for replication and the tissue-specific transcriptional regulatory sequence are not derived from the same gene (i.e., are heterologous to one another). The tissue-specific transcriptional regulatory sequence, in other words, is not derived from native vector sequences. For example, when the vector is an adenovirus vector, the tissue-specific transcriptional regulatory sequence is not derived from adenovirus.

45 **[0103]** The additional heterologous coding sequences can be expressed from various locations on the vector. For example, the coding sequence could be linked to the first coding sequence so that expression is directly dependent on transcription from that regulatory sequence through the first coding sequence and then into the additional coding sequence. Alternatively, the additional coding sequence could be expressed by being operably linked to a separate transcriptional regulatory sequence that is activated by the gene product of the first coding sequence (as in transactivation, for example). As a further alternative, the additional coding sequence could be placed under the control of a separate copy of the tissue-specific transcriptional regulatory sequence to which the first coding sequence is linked, although not

located in proximity to the first coding sequence. Finally, the additional coding sequence could be, in some instances, transcribed from its native transcriptional regulatory sequence (e.g., promoter) or other different transcriptional regulatory sequence, different from the one to which the first coding sequence is operably linked and not activated by the gene product from the first coding sequence.

5 **[0104]** When there are multiple heterologous genes expressed, the vector could contain various permutations of the above arrangement. For example, in the case where there is only one additional gene, this gene could be under the control of the same transcriptional regulatory sequence controlling the first coding sequence, under control of a trans-activatable transcriptional regulatory sequence, or transcriptionally linked to the first coding sequence so that they are transcribed as a unit from the tissue-specific transcriptional regulatory sequence controlling the first coding sequence, 10 or, finally, under control of another transcriptional regulatory sequence, for example its own native transcriptional regulatory sequence or another constitutive transcriptional regulatory sequence. When a second additional heterologous gene is expressed from the vector, the permutations increase but follow the same general strategy. For example, all three coding sequences can be transcribed from the same unit linked to a single tissue-specific transcriptional regulatory sequence. Alternatively, two of these may be transcribed from the unit and a third transcribed from the same transcriptional regulatory sequence but located in a different proximity. Alternatively, the first coding sequence may be transcribed separately from the second two sequences but both transcribed from the same tissue-specific transcriptional regulatory sequence, etc. Alternatively, the first coding sequence can be used to transactivate the second and third genes, which can be linked under the control of one transcriptional regulatory sequence or under separate transactivatable transcriptional regulatory sequences. Figure 7 illustrates some of the possible permutations that are encompassed in the present invention. The person of ordinary skill in the art would appreciate the possible permutations even if not explicitly described herein.

15 **[0105]** In a preferred embodiment, multiple genes on the vector can be controlled through one transactivator. Accordingly, where the first coding sequence has transactivator function, activation at the tissue-specific transcriptional regulatory sequence results in expression in that function and thus transactivation of any genes placed under control of transactivatable transcriptional regulatory sequences at one or more sites in the vector. For example, in an adenoviral-based vector, by linking a heterologous regulatory element to the E1a gene in an otherwise replication competent adenoviral vector, the invention provides the ability to place genes in other transcriptional regulatory sequences, such as E3, E4, E1 regions transactivated by E1a, or multiple copies of these promoters, such as two or more E3 promoters. The design will depend upon the intended use. For example, for cancer therapy a tumor-specific transcriptional regulatory sequence would be desirable to drive E1 a expression while therapeutic genes could be placed under transcriptional regulatory sequences transactivated by E1a, such as one or more E3 promoters, one or more E4 promoters, E1 promoters, or combinations thereof, that contain genes such as HSV-TK, GM-CSF, and IL-2.

20 **[0106]** In highly preferred embodiments, vectors are designed to express a heterologous gene that is toxic for viral replication. Through this gene one is able to modulate the amount of replication and thus expression of other heterologous gene products, therapeutic or otherwise, for maximal benefit and specificity. For example, in adenoviral vectors having an E1 a coding sequence linked to a tissue-specific promoter and having the HSV-TK gene under control of the E3 promoter, replication could be controlled by the addition of ganciclovir.

25 **[0107]** The preferred vectors of the present invention are adenoviral vectors. In a preferred embodiment of the invention, an adenovirus vector contains a tissue-specific transcriptional regulatory sequence linked to a gene in the E1 region.

30 **[0108]** In one embodiment, both E1 a and E1 b are operably linked to heterologous tissue-specific transcriptional regulatory sequences. In an alternative embodiment, only E1a is linked; E1b remains intact. In still another embodiment, E1b is linked, and E1a remains intact or is deleted. In any case, one or more tissue-specific and promoter-specific cellular transcriptional regulatory factors allows virus replication to proceed by transcribing the E1 a and/or E1 b gene functionally linked to the promoter. Further, either one or both of the E1 b functions may be linked to the transcriptional regulatory sequence.

35 **[0109]** In alternative embodiments, adenovirus vectors are provided with any of the other genes essential for replication, such as E2-E4, under control of a heterologous transcriptional regulatory sequence.

40 **[0110]** The invention further embodies the use of plasmids and vectors having only the essential regions of adenovirus needed for replication with either E1a, E1b 19kDa gene, or E1b 55kDa gene, or some combination thereof, modified. Such a plasmid, lacking any structural genes, would be able to undergo DNA replication. Accordingly, the vectors of the invention may consist essentially of the transcriptional regulatory sequence and one or more genes essential for replication of the vector. In the case of viral vectors, the vectors may consist essentially of the transcriptional regulatory sequence and the gene or genes essential for replication or life-cycle functions of the virus. It is also understood that these vectors may also further consist essentially of a DNA sequence encoding one or more toxic heterologous gene products when such vectors are intended as expression vectors for treatment.

45 **[0111]** Vectors as disclosed herein may also be derived from another DNA tumor virus. Such viruses generally include, but are not limited to, Herpesviruses (such as Epstein-Barr virus, cytomegalovirus, Herpes zoster, and Herpes simplex), papillomaviruses, papovaviruses (such as polyoma and SV40), and hepatitis viruses, parvoviruses, and picomaviruses.

[0112] The alternative viruses preferably are selected from any group of viruses in which the essential genes for replication of the virus can be placed under the control of a tissue-specific transcriptional regulatory sequence. All serotypes are included. The only common property of such viruses, therefore, is that they are transducible into target tissue, are genetically manipulatable, and are non-toxic when not replicating.

5 [0113] The relevant viral gene(s) are those that are essential for replication of the viral vector or of the virus. Examples of genes include, but are not limited to, the E6 and E7 regions of human papilloma virus, 16 and 18, T antigen of SV40, and CMV immediate early genes, polymerases from retroviruses and the like. Essentially, these include any gene that is necessary for the life cycle of the virus.

10 [0114] Vectors as disclosed herein may also be derived from an RNA virus or derived from a retrovirus. It is understood, however, that potentially any replicating vector can be made and used according to the essential design disclosed herein.

[0115] The vectors described herein can be constructed using standard molecular biological techniques. Standard techniques for the construction of such vectors are well-known to those of ordinary skill in the art, and can be found in references such as Sambrook et al., in *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York (1989), or any of the myriad of laboratory manuals on recombinant DNA technology that are widely available. A variety of strategies are available for ligating fragments of DNA, the choice of which depends on the nature of the termini of the DNA fragments and can be readily determined by the skilled artisan.

15 [0116] An adenovirus vector, in a preferred embodiment, is constructed first by constructing, according to standard techniques, a shuttle plasmid which contains, beginning at the 5' end, the "critical left end elements," which include an adenoviral 5' ITR, an adenoviral encapsidation signal, and an E1 a enhancer sequence; a promoter (which may be an adenoviral promoter or a foreign promoter); a tripartite leader sequence, a multiple cloning site (which may be as herein described); a poly A signal; and a DNA segment which corresponds to a segment of the adenoviral genome. Such DNA segment serves as a substrate for homologous recombination with a modified or mutated adenovirus. The plasmid may also include a selectable marker and an origin of replication. The origin of replication may be a bacterial origin of replication. Representative examples of such shuttle plasmids include pAVS6, as discussed herein and see Trapnell, B. et al., *Adv. Drug Deliv. Rev* 12:185-189 (1994). A desired DNA sequence containing a heterologous gene may then be inserted into the multiple cloning site to produce a plasmid vector.

20 [0117] This construct then is used to produce an adenoviral vector. Homologous recombination then is effected with a modified or mutated adenovirus in which one or more of the native transcriptional regulatory sequences have been deleted and replaced with the desired transcriptional regulatory sequence. Such homologous recombination may be effected through co-transfection of the plasmid vector and the modified adenovirus into a helper cell line by CaPO₄ precipitation.

25 [0118] Through such homologous recombination, a vector is formed which includes adenoviral DNA free of one or more of the native transcriptional regulatory sequences. This vector may then be transfected into a helper cell line for viral replication and to generate infectious viral particles. Transfections may take place by electroporation, calcium phosphate precipitation, microinjection, or through proteoliposomes.

30 [0119] The vector may include a multiple cloning site to facilitate the insertion of DNA sequence(s) containing the heterologous gene into the cloning vector. In general, the multiple cloning site includes "rare" restriction enzyme sites; i.e., sites which are found in eukaryotic genes at a frequency of from about one in every 10,000 to about one in every 100,000 base pairs. An appropriate vector is thus formed by cutting the cloning vector by standard techniques at appropriate restriction sites in the multiple cloning site, and then ligating the DNA sequence containing the heterologous gene into the cloning vector.

35 [0120] The coding sequence whose gene product is essential for vector replication is under the control of a suitable tissue-specific transcriptional regulatory sequence, which may be a promoter or an enhancer. A tissue-specific promoter may be, but is not limited to, AFP, PSA, CEA, DE3, α -fetoprotein, Erb-B2, surfactant, and the tyrosinase promoter. A tissue-specific enhancer may be, but is not limited to, DF3 breast cancer-specific enhancer, enhancers from viruses, and the steroid receptor family.

40 [0121] Suitable promoters for expressing the DNA sequence encoding the heterologous gene product include, but are not limited to, viral promoters, such as the adenoviral major late promoter, cytomegalovirus (CMV) promoter, Rous sarcoma virus promoter, inducible promoters, such as the MMTV promoter, the metallothionein promoter, heat shock promoters, the albumin promoter, the ApoE promoter, and the ApoA1 promoter. It is to be understood, however, the scope of the present invention is not limited to specific heterologous genes or promoters.

45 [0122] Suitable native promoters (already located on the viral vector) include, but are not limited to, adenovirus E2, E3 and E4. Further, as discussed, the tissue-specific transcriptional regulatory sequence used to activate vector replication can also be used to control transcription of one or more heterologous genes on locations separate from the first coding sequence in the vector genome.

50 [0123] In one embodiment, the adenovirus may be constructed by using a yeast artificial chromosome (or YAC) containing an adenoviral genome according to the method described in Ketner, et al., *Proc. Nat. Acad. Sci.* 91:6186-6190 (1994), in conjunction with the teachings contained herein. In this embodiment, the adenovirus yeast artificial chromosome

is produced by homologous recombination in vivo between adenoviral DNA and yeast artificial chromosome plasmid vectors carrying segments of the adenoviral left and right genomic termini. A DNA sequence containing the heterologous gene then may be cloned into the adenoviral DNA. The modified adenoviral genome then is excised from the adenovirus yeast artificial chromosome in order to be used to generate infectious adenoviral particles.

5 **[0124]** The infectious viral particles may then be administered in vivo to a host. The host may be an animal host, including mammalian, non-human primate, and human hosts.

[0125] The viral particles may be administered in combination with a pharmaceutically acceptable carrier suitable for administration to a patient. The carrier may be a liquid carrier (for example, a saline solution), or a solid carrier, such as, for example, microcarrier beads.

10 **[0126]** For the methods and particularly treatment methods described herein, modulating the amount of vector replication is a highly preferred aspect of the invention. This provides controlled amounts of a heterologous gene product.

[0127] The vectors herein provide a method for amplifying the expression of the heterologous gene. In addition to providing constitutive expression of a heterologous gene (as from a native promoter or other transcriptional regulatory sequence or from a tissue-specific promoter in the target tissue), via replication, the vector provides further copies of the heterologous gene for amplified expression. Including a toxic gene on the vector provides a way to control or modulate amplification and thus, expression.

15 **[0128]** Expression of a gene, particularly a toxic gene, can be absolutely controlled by conditioning the transcription of the gene completely on replication. This is the case when expression of such a gene depends upon transactivation from a gene product produced from a coding sequence of a gene essential for replication operably linked to a tissue-specific transcriptional regulatory sequence.

20 **[0129]** Vectors expressing gene products, such as HSV-TK, allow the control of replication by drugs such as ganciclovir. This expression in conjunction with tissue-specific control of the expression of a gene essential for replication allows the control of replication by the administration of an external drug. For example, tumor cells can be transduced with a vector expressing E1 a in a tissue-specific manner, which vector contains the HSV-TK gene. Upon replication, HSV-TK is either expressed or expression is amplified by replication of the vector. If in the course of monitoring vector replication, there is undesirable spread beyond the tumor edges into surrounding normal tissues, or release into the bloodstream which can be monitored, then viral replication can be dampened by adding ganciclovir to protect the normal cells. At high doses of ganciclovir (for example, 10-50 μ M), replication can be eliminated completely. At lower doses, replication can be dampened. In addition, since E1 a expression is dependent upon tissue-specific transcriptional regulatory sequences, in normal cells, E1 a expression would be prevented. Therefore, TK expression would also be diminished. Further, in those embodiments in which the TK coding sequence is operably linked to either the tissue-specific promoter or a promoter dependent on E1a transactivation, TK expression should cease entirely.

25 **[0130]** Also, since most normal cells are not actively dividing, the addition of ganciclovir should shut down viral replication but should not harm cells which would not incorporate this analogue into cellular DNA. Further, even if the cell were dividing, the elimination of TK expression would prevent the phosphorylation of ganciclovir and thus, no deleterious on the cell should result.

30 **[0131]** In some embodiments, the gene such as TK does not need to be expressed from the vector itself, but can be expressed from an unlinked location such as a separate vector or a cellular genome.

35 **[0132]** It should be appreciated that the ability to modulate viral replication allows the temporary expression of a given gene, and particularly a cytotoxic gene. This is useful, for example, in a treatment context where temporary expression of a gene is necessary to provide gene therapy. A gene can be expressed at a certain level, maintained at that level, and then expression can be increased and the cell eliminated.

40 **[0133]** In addition, it may not be desirable to eliminate a cell, but may be desirable to simply express a gene. In this case, it is desirable to express a gene at certain levels. Control of gene expression thus could be effected by a combination of, for example, E1a, TK, and GCV.

45 **[0134]** Expression of heterologous genes and/or treatment thereby are possible using the vectors described herein. Therefore, there are disclosed herein methods of replicating the vectors described herein.

50 **[0135]** In preferred embodiments, the methods are specifically directed to the introduction into a target tissue of a replication-conditional adenoviral vector. This vector selectively replicates in the cells of the target tissue. The replication is conditioned upon the function of a transcriptional regulatory sequence to which a viral gene is operably linked, which gene is necessary for vector replication. Thus, in the target tissue, replication can occur because either a cellular function in the target tissue allows transcription. Alternatively, there is a deficiency in a cellular function in the target tissue that normally prevents or inhibits transcription. The presence or absence of such functions provides the selectivity that allows the treatment of a specific tissue with minimum effect on the surrounding tissue(s).

55 **[0136]** Thus, there are disclosed herein methods for selectively distributing a polynucleotide in a given tissue in vivo, significantly reducing or avoiding distribution in non-target tissue. The polynucleotide is provided in the replication-conditional vector which is selectively distributed in the given tissue.

[0137] There are also disclosed herein methods for selectively expressing a gene product in a given tissue, avoiding

or significantly reducing expression in non-target or non-tumor tissue. In preferred methods, the gene expressed from the vector is a gene that has the potential of being cytotoxic to the host cell and/or toxic to viral replication in the cell. Thus, there is the option of obliterating the cell in which the virus is replicating, or simply decreasing or eliminating viral replication and avoiding cell killing. Accordingly, there is disclosed herein method for expressing such genes, for example

5 thymidine kinase, in a cell. This provides the possibility of treatment of diseases in which tissue-specific viral replication and gene expression is desired. Thus, also disclosed is a method for killing a cell in the case of conditions in which cell killing is desirable, such as cancer and other diseases involving abnormal cellular proliferation such as restenosis.

[0138] The invention allows for distribution of the above-mentioned vectors to a greater number of target cells than would be reached using a non-replicating vector. Successive infection provides a "domino effect" so that all or substantially

10 all of the cells in the target tissue are reached. Cells in addition to those first exposed to the polynucleotide, vector, or gene product, are thus potentially reached by the methods.

[0139] Such treatment is particularly necessary in cases in which surgical intervention is not feasible. For example, in patients with abnormal tissue intimately associated with neural tissue, surgery may be precluded or highly dangerous. Further, in the case of multiple metastases or microscopic metastases, surgery is not feasible.

15 [0140] In the target tissue, DNA replication alone may occur. Late viral functions that result in packaging of vector DNA into virions may also occur.

[0141] The vector may be introduced into the target tissue as naked DNA or by means of encapsidation (as an infectious virus particle or virion). In the latter case, the distribution is accomplished by successive infections of cells in the tissue by the virus such that substantially all or a significant number of the daughter cells are infected.

20 [0142] Tissue specificity is particularly relevant with respect to targeting an abnormal counterpart of a particular tissue type while avoiding the normal counterpart of the tissue, or avoiding surrounding tissue of a different type than the abnormal tissue, while treating the abnormal tissue. For example, the vectors of the present invention are useful for treating metastases to the liver. One specific example is colon cancer, which often metastasizes into the liver. It has been found that even when colon cancer metastasizes into the liver, the CEA promoter is active in the cells of the

25 metastases but not in normal liver cells. Accordingly, normal human adult liver should not support replication of a virus that has viral genes essential for replication linked to the colon cancer CEA-specific promoter. Replication should occur in the primary cancer cells. Another example is breast cancer, which also metastasizes to the liver. In this case, the DF3 mucin enhancer is linked to a gene essential for replication such as both E1a and E2a. Replication should occur in breast cancer but not in normal liver. A further example is the α -fetoprotein promoter, which is active in hepatocellular carcinoma.

30 This promoter is linked to a gene essential for replication. It has been found that the promoter is active only in the hepatocellular carcinoma. Accordingly, a virus is used that has a gene essential for replication linked to this promoter. Replication should be limited to hepatocellular carcinoma. A further example is the tyrosinase promoter. This promoter is linked to a gene essential for replication. Replication should occur in melanoma and not in normal skin. In each case, replication is expected in the abnormal but not the normal cells.

35 [0143] The vector of the invention encodes a heterologous gene product which is expressed from the vector in the tissue cells. The heterologous gene product can be toxic for the cells in the targeted tissue or confer another desired property.

[0144] A gene product produced by the vector can be distributed throughout the tissue, because the vector itself is distributed throughout the tissue. Alternatively, although the expression of the gene product may be localized, its effect

40 may be more far-reaching because of a bystander effect or the production of molecules which have long-range effects such as chemokines. The gene product can be RNA, such as antisense RNA or ribozyme, or protein. Examples of toxic products include, but are not limited to, thymidine kinase in conjunction with ganciclovir.

[0145] A wide range of toxic effects is possible. Toxic effects can be direct or indirect. Indirect effects may result from the conversion of a prodrug into a directly toxic drug. For example, Herpes simplex virus thymidine kinase phosphorylates

45 ganciclovir to produce the nucleotide toxin ganciclovir phosphate. This compound functions as a chain terminator and DNA polymerase inhibitor, prevents DNA synthesis, and thus is cytotoxic. Another example is the use of cytosine deaminase to convert 5-fluorocytosine to the anti-cancer drug 5-fluorouracil. For a discussion of such "suicide" genes, see Blaese, R.M. et al., Eur. J. Cancer 30A:1190-1193 (1994).

[0146] Direct toxins include, but are not limited to, diphtheria toxin (Brietman et al., Mol. Cell Biol. 10:474-479 (1990)), pseudomonas toxin, cytokines (Blankenstein, T., et al., J. Exp. Med. 173:1047-1052 (1991), Colombo, M.P., et al., J. Exp. Med. 173:889-897 (1991), Leone, A., et al., Cell 65:25-35 (1991)), antisense RNAs and ribozymes (Zaia, J.A. et al., Ann. N.Y. Acad. Sci. 660:95-106 (1992)), tumor vaccination genes, and DNA encoding for ribozymes.

50 [0147] In accordance with the present invention, the agent which is capable of providing for the inhibition, prevention, or destruction of the growth of the target tissue or tumor cells upon expression of such agent can be a negative selective marker; i.e., a material which in combination with a chemotherapeutic or interaction agent inhibits, prevents or destroys the growth of the target cells.

[0148] Thus, upon introduction to the cells of the negative selective marker, an interaction agent is administered to the host. The interaction agent interacts with the negative selective marker to prevent, inhibit, or destroy the growth of

the target cells.

[0149] Negative selective markers which may be used include, but are not limited to, thymidine kinase and cytosine deaminase. In one embodiment, the negative selective marker is a viral thymidine kinase selected from the group consisting of Herpes simplex virus thymidine kinase, cytomegalovirus thymidine kinase, and varicella-zoster virus thymidine kinase. When viral thymidine kinases are employed, the interaction or chemotherapeutic agent preferably is a nucleoside analogue, for example, one selected from the group consisting of ganciclovir, acyclovir, and 1-2-deoxy-2-fluoro- β -D-arabinofuranosil-5-iodouracil (FIAU). Such interaction agents are utilized efficiently by the viral thymidine kinases as substrates, and such interaction agents thus are incorporated lethally into the DNA of the tumor cells expressing the viral thymidine kinases, thereby resulting in the death of the target cells.

[0150] When cytosine deaminase is the negative selective marker, a preferred interaction agent is 5-fluorocytosine. Cytosine deaminase converts 5-fluorocytosine to 5-fluorouracil, which is highly cytotoxic. Thus, the target cells which express the cytosine deaminase gene convert the 5-fluorocytosine to 5-fluorouracil and are killed.

[0151] The interaction agent is administered in an amount effective to inhibit, prevent, or destroy the growth of the target cells. For example, the interaction agent is administered in an amount based on body weight and on overall toxicity to a patient. The interaction agent preferably is administered systemically, such as, for example, by intravenous administration, by parenteral administration, by intraperitoneal administration, or by intramuscular administration.

[0152] When the vectors of the present invention induce a negative selective marker and are administered to a tissue or tumor in vivo, a "bystander effect" may result, i.e., cells which were not originally transduced with the nucleic acid sequence encoding the negative selective marker may be killed upon administration of the interaction agent. Although the scope of the present invention is not intended to be limited by any theoretical reasoning, the transduced cells may be producing a diffusible form of the negative selective marker that either acts extracellularly upon the interaction agent, or is taken up by adjacent, non-target cells, which then become susceptible to the action of the interaction agent. It also is possible that one or both of the negative selective marker and the interaction agent are communicated between target cells.

[0153] In one embodiment, the agent which provides for the inhibition, prevention, or destruction of the growth of the tumor cells is a cytokine. In one embodiment, the cytokine is an interleukin. Other cytokines which may be employed include interferons and colony-stimulating factors, such as GM-CSF. Interleukins include, but are not limited to, interleukin-1, interleukin-1 β , and interleukins-2-15. In one embodiment, the interleukin is interleukin-2.

[0154] In a preferred embodiment of the invention, the target tissue is abnormally proliferating, and preferably tumor tissue. The vector or virus is distributed throughout the tissue or tumor mass.

[0155] All tumors are potentially amenable to treatment with the methods of the invention. Tumor types include, but are not limited to hematopoietic, pancreatic, neurologic, hepatic, gastrointestinal tract, endocrine, biliary tract, sino-pulmonary, head and neck, soft tissue sarcoma and carcinoma, dermatologic, reproductive tract, and the like. Preferred tumors for treatment are those with a high mitotic index relative to normal tissue. Preferred tumors are solid tumors, and especially, tumors of the brain, most preferably glioma.

[0156] The methods can also be used to target other abnormal cells, for example, any cells which are harmful or otherwise unwanted in vivo. Broad examples include cells causing autoimmune disease, restenosis, and scar tissue formation, abnormal angiogenesis, PRD, arthritis, chronic diabetes, and ARMD.

[0157] Further, treatment can be ex vivo. Ex vivo transduction of tumor cells would overcome many of the problems with current viral delivery systems. Tissue is harvested under sterile conditions, dissociated mechanically and/or enzymatically and cultured under sterile conditions in appropriate media. Vector preparations demonstrated to be free of endotoxins and bacterial contamination are used to transduce cells under sterile conditions in vitro using standard protocols. The accessibility of virus to cells in culture is currently superior to in vivo injection and permits introduction of vector viral sequences into essentially all cells. Following removal of virus-containing media cells are immediately returned to the patient or are maintained for several days in culture while testing for function or sterility is performed.

[0158] For example, patients with hypercholesterolemia have been treated successfully by removing portions of the liver, explanting the hepatocytes in culture, genetically modifying them by exposure to retrovirus, and re-infusing the corrected cells into the liver (Grossman et al., Nature Genetics 6:335-341 (1994)).

[0159] Viral transduction also has potential applications in the area of experimental medicine. Transient expression of biological modifiers of immune system function such as IL-2, IFN- γ , GM-CSF or the B7 co-stimulatory protein has been proposed as a potential means of inducing anti-tumor responses in cancer patients.

[0160] Vectors as disclosed herein may also be derived from another DNA tumor virus. Such viruses generally include, but are not limited to, Herpesviruses (such as Epstein-Barr virus, cytomegalovirus, Herpes zoster, and Herpes simplex), papillomaviruses, papovaviruses (such as polyoma and SV40), and hepatitis viruses.

[0161] The relevant viral gene(s) are those that are essential for replication of the viral vector or of the virus. Examples of genes include, but are not limited to, the E6 and E7 regions of human papilloma virus, 16 and 18, T antigen of SV40, and CMV immediate early genes, polymerases from retroviruses and the like. Essentially, these include any gene that is necessary for the life cycle of the virus.

[0162] Vectors as disclosed herein may also be derived from an RNA virus or derived from a retrovirus. It is understood, however, that potentially any replicating vector can be made and used according to the essential design disclosed herein.

[0163] It is important to know whether the vectors of the invention will replicate in a specific tissue from a patient. If vector replication is found to be beneficial for therapy, then a screen is provided for those patients who best respond to the therapy disclosed herein. If it is found to be harmful, then there is a screen for prevention of the treatment of patients who would have an adverse response to the treatment. Currently, the only non-biological assays that are commonly used are expression screening, PCR, and sequencing. These often result in false negatives, are time-consuming, expensive, and yield only information in the best of cases about the status of the genes and not their biological function.

[0164] Accordingly, a method is disclosed for identifying an abnormal tissue, the cells of which contain a transcription factor that allows replication of a replication-conditional vector, or are deficient for an inhibitory factor for transcription.

[0165] In this method, a tissue biopsy is explanted, a replication-conditional vector is introduced into the cells of the biopsy, and vector DNA replication in the cells is quantitated. Accordingly, a method is disclosed for screening tissue for the presence of factors that allow vector replication, or for a deficiency of a factor that inhibits transcription. Such a screen is useful, among other things, for identifying tissue, prior to treatment, which will be amenable to treatment with a particular vector to be replicated in the tissue.

[0166] Therefore, a method is disclosed for assaying vector utility for treatment by removing a tissue biopsy from a patient, explanting the biopsy into tissue culture, introducing the replication-conditional vector into the biopsy, and assaying vector replication in the cells of the biopsy.

[0167] Testing or screening of tissues includes an assay for vector nucleic acid replication or for virus replication, when the vector is capable of forming infectious virions.

[0168] Thus, there is disclosed a method for screening a tumor for transcription regulatory functions that allow vector replication or for the absence of these functions which would normally prevent the replication of a virus vector.

[0169] However, any abnormal tissue can be screened for the functions described above by an assay for nucleic acid or virus replication.

[0170] In a further embodiment of the invention, a cell is provided which contains a virion produced in the cell by replication in the cell of the replication-conditional vectors of the present invention. Thus, the invention provides "producer cells" for the efficient and safe production of recombinant replication-conditional vectors for further use for targeted gene therapy *in vivo*.

[0171] One of the major problems with the currently available producer cells is that such cells contain, in the genome, viral sequences that provide complementing functions for the replicating vector. Because the cell contains such sequences, homologous recombination can occur between the viral sequence in the genome and the viral vector sequences. Such recombination can regenerate recombinant wild-type viruses which contaminate the vector or virus preparation produced in the producer cell. Such contamination is undesirable, as the wild-type viruses or vectors can then replicate in non-target tissue and thereby impair or kill non-target cells. Therefore, one of the primary advantages of the producer cells of the present invention is that they do not contain endogenous viral sequences homologous to sequences found in the vector to be replicated in the cells. The absence of such sequences avoids homologous recombination and the production of wild-type viral recombinants that can affect non-target tissue.

[0172] Accordingly, the invention embodies methods for constructing and producing replication-conditional virions in a cell comprising introducing the replication-conditional vector of the present invention into the cell wherein the genome of the cell is devoid of vector sequences, replicating the vector in the cell, forming the virion, and purifying the virion from the cell. The virion is an adenoviral virion and the vector is an adenoviral vector. In further embodiments of the invention, the cell is a tumor cell.

[0173] The vector encodes a heterologous gene product such that the virion also encodes the gene product, and when the vector or virion are used for gene therapy, the therapy is facilitated by expression of the heterologous gene product. Alternatively, the producer cell can be used for the production of a heterologous gene product *per se* encoded by the vector. When the vector replicates in the producer cell, the gene product is expressed from the multiple copies of the gene encoding the gene product. Following expression, the gene product can be purified from the producer cells by conventional lysis procedures, or secreted from the producer cell by appropriate secretion signals linked to the heterologous gene by known methods. The transduction of cells by adenoviral vectors has been described. Transfection of plasmid DNA into cells by calcium phosphate (Hanahan, D., *J. Mol. Biol.* 166:577 (1983)), lipofection (Feigner et al., *PNAS* 84:7413 (1987)), or electroporation (Seed, B., *Nature* 329:840 ()) has been described. DNA, RNA, and virus purification procedures are described (Graham et al., *J. Gen. Virol.* 36:59-72 (1977)).

[0174] Preferred hosts for producer cell lines include but are not limited to HuH7, SW480, BIGF10, HepG2, MCF-7, and SK-MEL2. Primary tumors from which cell lines can be derived, or existing cell lines, can be tested for the ability to allow replication by means of the tissue-specific transcriptional regulatory sequence. Any primary tumor could be explanted and developed into producer cells for the vectors of the present invention. As long as the cell does not contain endogenous vector or viral sequences that could recombine with the vector or virus to produce wild-type vector or virus, the cell is potentially useful as a host. It is understood that any cell is potentially useful, not only tumor cells.

[0175] The ultimate goal for a producer cell line, and particularly an adenoviral producer line, is to produce the highest yield of vector with the least possibility of contamination by wild-type vector. Yield depends upon the number of cells infected. Thus, the more cells that it is possible to grow and infect, the more virus it is possible to generate. Accordingly, candidate cells would have a high growth rate and will grow to a high density. The cell should also have a high amount of viral receptor so that the virus can easily infect the cell. Another characteristic is the quality of the vector produced (i.e., the preparation should not include a high amount of non-infectious viral particles). Accordingly, candidate producer cells would have a low particle-to-plaque-forming-unit ratio. Thus, these cells are a preferred cell type for deriving a producer cell line. Primary explants or the known cell lines can be used.

[0176] Thus, such obtainable cells can serve as producer cells for recombinant replication-conditional vectors, viruses, and gene products.

[0177] A variety of ways have been developed to introduce vectors into cells in culture, and into cells and tissues of an animal or a human patient. Methods for introducing vectors into mammalian and other animal cells include calcium phosphate transfection, the DEAE-dextran technique, microinjection, liposome mediated techniques, cationic lipid-based techniques, transfection using polybrene, protoplast fusion techniques, electroporation and others. These techniques are well known to those of skill, are described in many readily available publications and have been extensively reviewed. Some of the techniques are reviewed in Transcription and Translation, A Practical Approach, Hames, B.D. and Higgins, S.J., eds., IRL Press, Oxford (1984), and Molecular Cloning, Second Edition, Maniatis et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989).

[0178] Several of these techniques have been used to introduce vectors into tissues and cells in animals and human patients. Chief among these have been systemic administration and direct injection into sites in situ. Depending on the route of administration and the vector, the techniques have been used to introduce naked DNA, DNA complexed with cationic lipid, viral vectors and vector producer cell lines into normal and abnormal cells and tissues, generally by direct injection into a targeted site.

[0179] The aforementioned techniques for introducing polynucleotide, viral and other vectors into cells in culture, in animals and in patients can be used to develop, test and produce, as well as use vectors in accordance with the invention. For instance, cells containing a vector introduced by these methods can be used for producing the vector. In addition, cells containing a vector can be used as producer-cells and introduced into cells or tissues of an animal to produce the vector in situ.

[0180] Replication of a polynucleotide, viral or other vector can be assayed by well-known techniques. Assays for replication of a vector in a cell generally involve detecting a polynucleotide, virions or infective virus. A variety of well-known methods that can be used for this purpose involve determining the amount of a labeled substrate incorporated into a polynucleotide during a given period in a cell.

[0181] When replication involves a DNA polynucleotide, ³H-thymidine often is used as the labelled substrate. In this case, the amount of replication is determined by separating DNA of the vector from the bulk of cellular DNA and measuring the amount of tritium incorporate specifically into vector DNA.

[0182] Other methods to assay replication, however, include, but are not limited to, hexon immunohistochemistry or another late gene immunoassay that is linked to DNA or viral replication.

[0183] Replication of a polynucleotide vector also may be detected by lysing or permeating cells to release the polynucleotide, then isolating the polynucleotide and quantitating directly the DNA or RNA that is recovered. Polynucleotide replication also may be detected by quantitative PCR using primers that are specific for the assay polynucleotide.

[0184] Virions may be assayed by EM counting techniques well known to the art, by isolating the virions and determining protein and nucleic acid content, and by labeling viral genomic polynucleotides or virion proteins and determining the amount of virion from the amount of polynucleotide or protein.

[0185] It is well known that virions may not all be viable and where infectivity is important, infectious titer may be determined by cytopathic effect or plaque assay.

[0186] Any of these well-known techniques, among others, can be employed to assay replication of a vector in a cell or tissue in accordance with the invention. It will be appreciated that different techniques will be better suited to some vectors than others and to some cells or tissues than others.

[0187] Having thus described herein the invention in general terms, the following non-limiting examples are presented to illustrate the invention.

Example 1

The hepatoma-specific promoter, alpha-fetoprotein promoter, linked to E1 a

[0188] The alpha-fetoprotein (AFP) promoter has been previously shown to be highly active in hepatoma cells and silent in adult hepatocytes and other adult tissues. A 4.9 kb alpha-fetoprotein promoter-containing construct was used to derive the promoter. Alternatively, the promoter could also be made based on available references.

[0189] pAVE1a02i (Figure 1 A) which places the E1a/E1b genes under the control of the alpha-fetoprotein promoter in an adenovirus shuttle plasmid was cloned by purifying a restriction fragment which contains the E1 a coding region only and all of E1 b gene by cleaving the plasmid 380-280E1 (Figure 1 A) with SpeI and MunI and ligating this to pAVSAFP.TK1 (Figure 1 A) cleaved with MunI and NheI.

[0190] The adenovirus shuttle plasmid pAVSAFP.TK1 (Figure 1 A), which has the TK gene under the control of the native 4.9 kb alpha-fetoprotein promoter, was made exactly as described in Figures 11 and 12 of U.S. patent application 08/444,284, Chiang et al., "Gene therapy of hepatocellular carcinoma through cancer-specific gene expression," filed on May 18, 1995, which is incorporated herein by reference for its relevant teaching. This shuttle plasmid contains the left ITR, packaging signal, the native AFP promoter, the HSV TK gene, and a homologous recombination fragment. Digestion of this plasmid with MunI and NheI removes the TK gene and part of the E1a gene. What remains is a 10,667 base pair fragment. This fragment is added to the E1 a and E1b open reading frames from plasmid 380-280E1 (see below).

[0191] Plasmid 380-280E1 contains the E1 a ORF and all of E1b. A SpeI/MunI restriction fragment of 3397 base pairs as described in Figure 1 A from plasmid 380-280E1 was used to ligate to the MunI/NheI fragment from pAVSAFP.TK1 to construct pAVE1a02i. The SpeI/MunI fragment from 380-280E1 can be found by reference to plasmid SE280-E1 which contains the same fragment as found in 380-280E1; SE280-E1 can be found in U.S. patent application 08/458,403, incorporated herein by reference for its relevant teaching.

[0192] The shuttle plasmid pAVE1a04i (Figure 1 C) was cloned by digesting pAVE1a02i (Figure 1 A) with AscI followed by DNA polymerase I large fragment (Klenow) filling of the 3' recessed end (Figure 1 A). The linearized pAVE1a02i was then digested with SpeI (Figure 1 A) and the resulting 9104 base pair fragment was isolated by agarose gel electrophoresis. This procedure removes the AFP promoter so that the 9104 base pair fragment contains the above-mentioned components but no promoter.

[0193] The plasmid pAF(AB)2(S)6-CAT (Figure 1A, 1 B) was digested with XbaI followed by Klenow filling of the 3' recessed ends. pAF(AB)2(S)6-CAT contains a shortened AFP promoter with six silencer elements and two enhancer regions, AB2 that are responsible for enhancing the activity in hepatoma cells and repressing the activity in adult liver cells.

[0194] pAF(AB)2(S)6-CAT was constructed by placing six copies of the distal silencer immediately upstream of the basal 200 base pair AFP promoter. Two copies of the enhancer AB region, originally thought to be in opposite orientation, were placed immediately upstream of the silencer elements. The distal silencer element, the basal promoter, and the enhancer elements are as described in Nakabayashi et al. (Molec. & Cell. Biol. 11:5885-5893 (1991)).

[0195] The linearized pAF(AB)2(S)6-CAT was then digested with SpeI (Figure 1 A) and the 1986 base pair fragment containing the shortened AFP promoter was isolated by gel electrophoresis. The 9104 base pair fragment of pAVE1 a02i and the 1986 base pair fragment of pAF(AB)2(S)6-CAT containing the shortened synthetic AFP promoter, were ligated to make the plasmid pAVE1a04i (Figure 1C) which places the adenovirus E1a gene under the control of the shortened AFP promoter.

[0196] The pAF(AB)2(S)6-CAT was reported to have two copies of the enhancer AB region in opposite orientation but was determined by the inventors to have the enhancer region as a tandem repeat. The plasmid map of pAVE1 a04i shows the enhancer (AB) region of the AFP promoter as an inverted repeat (Figure 1A). However, the map was later corrected to show the true orientation of the enhancer regions. The corrected plasmid map was designated as pAVE1a06i.

Construction of a virus with the hepatoma-specific AFP promoter operably linked to the Ela gene

[0197] The adenovirus AvE1 a04i (Figure 2C) was constructed by homologous recombination of the shuttle plasmid, pAVE1 a06i (see Figures 1C and 2B), with the large (C1a1) fragment of Av1lacZ DNA (Figure 2A) in 293 cells. Resulting recombinant virus containing the AFP promoter with the two direct repeat enhancers was isolated and initially designated as AvE1a06i (Figure 2C, top of page). In the process of plaque purifying the recombinant virus, a single enhancer deletion mutant was isolated and designated as AvE1 a04i (Figure 2C, bottom of page).

[0198] AvE1a04i was shown to replicate specifically in cell lines as described in PCT App. No. US 95/15455 (U.S. 08/849,117).

Example 2

AV5E1aTK01i Construction

[0199] pAVS10TK1 (U.S. Appl. No. 08/444,284) (the source of the TK gene in Av15EKa04i below) was digested with XbaI and SpeI and the resulting 1230 bp fragment, containing the HSV-TK open reading frame, was ligated into the partially digested XbaI site at base pair position 6162 of prepac (Figure 3A) (described in U.S. patent application 08/852, 924). This shuttle plasmid contains the last 8886 base pairs from 25171 through 34057 of the Add1327 genome (Thimmapaya, Cell 31:543 (1983)) cloned into pBluescript SKII(+) (Stratagene). Prepac (Figure 3A) is a large plasmid that contains the adenoviral genomic DNA from the right ITR through the E3 region. The TK gene derived from pAVS10TKI

was inserted into the XbaI site in prepac, putting this gene under the control of the E3 promoter. The resulting ligated plasmid DNA was transformed into *E. coli* STBL2 cells (Life Technologies) and labeled as prepacTK. The plasmid prepacTK was then used to make the plasmid prepacTKbnl (Figure 3B) by ligating the following two annealed oligonucleotides (5'-GGCCGCATGCATGTTTAAACG-3' and

5'-GATCCGTTTAAACATGCATGC-3') into the BamHI and NotI digested sites of prepacTK (Figures 3A, 3B). The ligation of this oligonucleotide creates another BamHI site without destroying the open reading frame.

[0200] The plasmid prepacTKm was then cloned by digesting prepacTKbnl with BamHI and SpeI and isolating the resulting 11148 base pair fragment (Figure 3B). This removes a portion of the E2a gene and allows it to be replaced with a modified E2a gene containing a mutation that permits replication in monkey cells (from pG1MBS, see below). This 11198 base pair fragment was ligated to the 5520 base pair fragment obtained by digesting the plasmid pG1MBS with BamHI and SpeI (Figure 3B). The final plasmid is designated prepacTKm. This plasmid has the TK gene in the E3 region, and also contains the E2a mutation as explained below.

[0201] The plasmid pG1MBS contains the adenoviral type 5 (Ad5) region between the BamHI site and the SpeI site (Ad5 genome base pair positions 21562 to 27082, respectively, Figure 3B). Plasmid pG1MBS has a point mutation in the adenoviral sequences at base pair 11670 which allows for increased adenoviral replication in monkeys (as described by Kruijen, Nucl. Acids Res. 9:4439 (1981)).

[0202] To prepare the adenoviral vector Av5E1aTK01i the plasmid pREpacTKm (Figure 3C) was digested with the restriction enzymes BamHI and Sall and the 13748 base pair fragment was isolated by agarose gel electrophoresis. Adenovirus Addl327 DNA was prepared by digesting cesium density gradient centrifugation purified virus with Proteinase K. The Proteinase K digested Addl327 viral DNA was then extracted first with phenol/chloroform followed by chloroform and finally with buffer saturated ether. The DNA was recovered after equilibrating in water using an Amicon Centricon 100 unit. The purified Addl327 DNA was then digested with BamHI and the 21562 base pair fragment was isolated by agarose gel electrophoresis. The 13748 base pair BamHI/Sall fragment from pREpacTKm and the 21562 base pair BamHI fragment from Addl327 were then ligated together. The resulting ligated fragments were then transfected into A549.A30.S8 cells using Lipofectamine (Life Technologies Inc.) and incubated at 37°C in a 5% CO₂, humidified, incubator until cytopathic effects (CPE) were observed. The resulting recombinant Av5E1aTK01i (Figure 3C) virus was then plaque purified on A549.A30.S8 cells. Av5E1aTK01i plaques were screened by PCR for the presence of HSV-TK sequences using the following primers: (LMC11: 5'-AGCAAGAAGCCACGG AAGTC-3' and LMC12: 5'-AGGTCGCAGATCGTCG-GTAT-3').

Av15E1 a04i Construction

[0203] Av1 E1 a04i was completely digested with SrfI (Figure 4A) confirmed digestion on 1% agarose gel in 1x TAE, organic extracted digest with buffered phenol, buffered phenol/chloroform/isoamyl alcohol (25:24:1), and chloroform, and then added 1/10 volume 3M NaOAc and 2.5 volume 95% EtOH to precipitate. The precipitate was pelleted by 20 minute centrifugation at 12000 g, washed 1x with 70% EtOH, and then air dried for 30 minutes. The pellet was resuspended in dH₂O and digested with BamHI. The complete digestion was confirmed and purified as before.

[0204] 293 cells were cotransfected with 2.5 ug Av5E1aTk01i (digested with Bst 1107 and ClaI (Figure 4B) and 2.5 ug Av1 E1a04i (digested with Bam HI and Srf I, Figure 4A) using Promega CaCl₂ Transfection Kit (Figure 4C). Transfections were washed 1 x with Richters media supplemented with 10% FBS (R10) and then overlaid with 1x MEM/10% FBS/0.5 % Pen-Strep/1% Fungizone. Plaques were identified by microscopic examination and picked with 1000ul pipet tip into 500 ul R5 media. Plaques were lysed by freeze-thaw (4x) and then infected into S8 cells stimulated with 0.3 um dexamethasone. When early CPE was evident, the cells were washed 1 x with PBS, lysed with 200 ul 1 N NaOH and neutralized with 30 ul of 7.5 M ammonium acetate. The CVL was diluted 1:50 in dH₂O and 5 ul of the diluted lysate was used in a 50 ul PCR reaction using the BMB master mix reagents.

[0205] E1a promoter primer pair detects both Av15E1aTk04i (1653 bp) and Addl327 (405 bp) CH12 5'-GACCGTT-TACGTGGAGACTCGC -3' bp 367 in ITR CH13 5'-ACCGCCAACATTACAGAGTCG-3' bp 772 or bp 2020 in E1a gene of Addl327 or Av15E1aTk04i, respectively.

HSV-TK primer pair 990 bp product in either Av15E1 aTk04i or Av5E1aTk01i

LMC11 5'-AGCAAGAAGCCACGGAAGTC -3' bp 100 of HSV-Tk ORF

LMC12 5'-AGGTCGCAGATCGTCGGTAT -3' bp 1090 of HSV-Tk ORF

[0206] Three primary plaques were identified for additional plaque purification on S8 cells using same PCR of CVL to identify positive plaques. One plaque was subsequently purified three times by plaque purification on S8 cells. A bulk preparation of the tertiary plaque for Av15E1 aTk04i has a titer of 9 x 10¹⁰ particles/ml, ratio 26.

[0207] Av15E1 aTk04i (Figure 4C) has the AFP promoter controlling E1 a expression, and thus, only AFP positive cells are permissive for vector replication. Av15E1 aTk04i also has an E2a mutation (hr404) which makes monkey cells permissive for replication, potentially expanding the range of permissive species to additional restricted species. Finally, Av15E1 aTk04i has HSV-TK under the control of the E3 promoter which is positively regulated by E1a; therefore, there

should be no expression in the absence of E1a, which will only be present in AFP positive cells. As a safety feature in the event that replication occurs in nontarget cells, then only the nontarget and target cells that are replicating vector would be sensitive to GCV treatment. Any infected, nontarget cells that are not replicating vector will be insensitive to GCV.

5 Example 3

[0208] Av15E1aTk04i should have the same tissue specific replication restricted adenoviral profile as Av1E1a04i, specifically, replication only in AFP positive cells (as described in PCT Appl. No. US 95/15455 (U.S. 08/849,117)). Infection of Hep3B and HepG2 (AFP positive hepatocellular carcinoma cell lines) and S8 (E1a positive cell line) with Addl327, Av1E1a04i, or Av15E1aTk04i results in visible CPE at infections of 0.5, 5, and 50 particle per cell (Figure 5). Av1nBg02v (E1 deleted virus) infection at the same input particle numbers exhibit no visible CPE except in S8 cells. Conversely, Chang and HeLa (AFP negative cell lines) exhibit visible CPE only when infected with Addl327 (wild type) when infected at the same input particles. Av1E1a04i, Av15E1aTk04i, and Av1nBg02v all manifest no CPE at infections of 0.5, 5, and 50 particle per cell in Chang and HeLa cells (Figure 5). This indicates that AV15E1aTK04i also replicates specifically in AFP-positive cell lines. Therefore, the E1a gene product, which controls the E3 promoter, must also be induced specifically in AFP-positive cell lines. Therefore, HSV-TK, which is controlled by the E3 promoter, is also induced specifically in AFP-positive cell lines.

20 Example 4

Replacement of other therapeutic toxic genes into the tumor-specific replication competent vectors

[0209] Genes such as TK, cytokines, or any therapeutic genes can be placed into the E3 region of the vector backbone by standard plasmid construction and homologous recombination. Those genes can be placed under the control of an E1 α -dependent promoter, or a constitutive promoter such as RSV or CMV.

In vitro control of replication by inclusion of HSV-TK into the E3 region (Figure 6).

[0210] HSV-TK was placed into the E3 region of Addl327 to form Av5E1aTK01i. A549 cells were transduced with either Addl327 or Av5E1aTK01i. Cells were then treated with 10 μ M GCV for 5 days. Following the 5 day incubation cells were harvested, washed in HBSS, and then freeze/thawed to prepare a crude viral lysate. Titers were then performed by standard TCID50 assays on 293 cells to determine viral titer. The experiment was repeated three times. Results show that while GCV had no effect on an adenovirus not carrying the HSV-TK gene, it caused a greater than 4 order of magnitude drop in titer of a vector carrying the HSV-TK gene.

In vivo control of replication by inclusion of HSV-TK into E3 region.

[0211] Subcutaneous tumors were formed by injecting 1×10^7 A549 cells into the subcutaneous space of the right flank of nude mice. After tumors formed, 1×10^9 pfu of Av5E1aTK01i was injected into several animals containing tumors. After 5 days half the animals received 5 days of IP GCV treatment at 75 mg/kg once a day. At the end of this time frame all animals were sacrificed. Immunohistochemistry was performed on all samples for HSV-TK expression and hexon expression, the latter which is only expressed when the vector is replicating. Results show that only hexon is severely diminished when the animals were treated with GCV.

45 Example 5

GCV Inhibition of Av5E1aTk01i Replication In Vivo.

[0212] Subcutaneous tumors were formed by injecting 1×10^7 A549 cells into the subcutaneous space of the right flank of nude mice. After tumors formed 1×10^9 pfu of Addl327Tk01i was injected into several animals containing tumors. After 5 days, half of the animals received 5 days of IP GCV treatment at 75 mg/kg once a day. At the end of this time frame all animals were sacrificed. Immunohistochemistry was performed on all samples for HSV-TK expression and hexon expression, the latter which is only expressed when the vector is replicating. Results show that only hexon is severely diminished when the animals were treated with GCV.

TABLE 1: Oligonucleotide Primers for Constructing Tissue-Specific Promoters

1. DF3 (BREAST CANCER)

5' GGG CGC GCC CTG GAA AGT CCG GCT GGG GCG
 5 GGG ACT GTG GGT TTC AGG GTA GAA CTG CGT
 GTG GAA 3'

5' CGG GAC AGG GAG CGG TTA GAA GGG TGG GGC
 10 TAT TCC GGG AAG TGG TGG GGG GAG GGA ACT
 AGT A 3'

5' GAT CTA CTA GTT CCC TCC CCC CAC CAC TTC
 15 CCG GAA TAG CCC CAC CCT TCT AAC CGC TCC
 CTG 3'

5' TCC CGT TCC ACA CGC AGT TCT ACC CTG AAA
 20 CCC ACA GTC CCC GCC CCA GCC GGA CTT TCC
 AGG GCG CGC CC 3'

2. TYROSINASE (MELANOMA)

5' GAC CCG GGC GCG CCG GAG CAG TGC TAT TCA
 25 AAC CAT CCA G 3'

5' CGA GAT CTA CTA GTT CTG CAC CAA TAG GTT
 30 AAT GAG TGT C 3'

3. CEA PROMOTER (HEPATOCELLULAR CARCINOMA)

5' GAC CCG GGC GCG CCT CTG TCA CCT TCC TGT
 35 TGG 3'

5' CGA GAT CTA CTA GTT CTC TGC TGT CTG CTC
 TGT C 3'

40 SEQUENCE LISTING

[0213]

(1) GENERAL INFORMATION:

45

(i) APPLICANT:

(A) NAME: Novartis AG

50

(B) STREET: Schwarzwaldallee 215

(C) CITY: Basel

(E) COUNTRY: Switzerland

55

(F) POSTAL CODE (ZIP): 4058

EP 1 032 696 B1

(G) TELEPHONE: +41 61 324 11 11

(H) TELEFAX: + 41 61 322 75 32

5 (ii) TITLE OF INVENTION: Vector for tissue-specific replication and gene expression

(iii) NUMBER OF SEQUENCES: 8

10 (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

15

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 66 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: NO

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

35

GGGCGCGCCC TGGAAAGTCC GGCTGGGGCG GGGACTGTGG GTTTCAGGGT AGAACTGCGT 60

GTGGAA 66

40

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

45

(A) LENGTH: 64 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: NO

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CGGGACAGGG AGCGGTTAGA AGGGTGGGGC TATTCCGGGA AGTGGTGGGG GGAGGGA ACT 60

5 AGTA 64

(2) INFORMATION FOR SEQ ID NO: 3:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 63 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

20 (A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

25 GATCTACTAG TTCCCTCCCC CCACCACTTC CCGGAATAGC CCCACCCTTC TAACCGCTCC 60

30 CTG 63

(2) INFORMATION FOR SEQ ID NO: 4:

35 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 71 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

45 (A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

50 TCCCGTTCCA CACGCAGTTC TACCCTGAAA CCCACAGTCC CCGCCCCAGC CGGACTTTCC 60

55 AGGGCGCGCC C 71

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

EP 1 032 696 B1

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

10

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5: GACCCGGGCG CGCCGGAGCA GTGCTATTCA AACCATC-CAG 40

15

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 40 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: other nucleic acid

25

(A) DESCRIPTION: /desc = "oligonucleotide"

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6: CGAGATCTAC TAGTTCTGCA CCAATAGGTT AATGAGT-GTC 40

30

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

45

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7: GACCCGGGCG CGCCTCTGTC ACCTTCCTGT TGG 33

50

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

5 (iii) HYPOTHETICAL: NO
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8: CGAGATCTAC TAGTTCTCTG CTGTCTGCTC TGTC
34

10 **Claims**

1. A replication conditional adenoviral vector comprising:

15 (a) a heterologous transcriptional regulatory sequence (TRS) operably linked to the coding sequence of an adenoviral gene essential for replication (GER) and (b) at least one additional coding sequence encoding a heterologous gene product, wherein said additional coding sequence is operably linked to a homologous transcriptional regulatory sequence (TRS) that is transactivated by the gene product of said adenoviral gene essential for replication.

20 2. The adenoviral vector of claim 1, wherein said adenoviral gene essential for replication (GER) is an adenoviral E1a gene

3. The adenoviral vector of claim 2, wherein said heterologous TRS is tissue-specific.

25 4. The adenoviral vector of claim 2 or 3, further comprising a second additional coding sequence operably linked to a separate, heterologous tissue specific transcriptional regulatory sequence.

5. The adenoviral vector of claim 2 or 3, wherein said homologous TRS is selected from the group consisting of an adenoviral E1, E2, E3 and E4 promoter and is transactivated by the gene product of said adenoviral E1a gene.

30 6. The adenoviral vector of claim 5, wherein said homologous TRS is an adenoviral E3 promoter.

7. The adenoviral vector of claim 1, wherein said heterologous gene is a cytokine gene.

35 8. The adenoviral vector of claim 7, wherein said cytokine gene is granulocyte macrophage - colony stimulating factor (GM-CSF) or interleukin-2 (IL-2).

9. The adenoviral vector of claim 1, wherein the expression product of said heterologous gene is toxic for the target tissue.

40 10. The adenoviral vector of claim 9, wherein said heterologous gene is thymidine kinase or cytosine deaminase.

11. The adenoviral vector of claim 5, wherein an Herpes Simplex Virus thymidine kinase (HSV-TK), GM-CSF or IL-2 gene is expressed under the control of an E3 promoter.

45 12. The adenoviral vector of claim 5, wherein an HSV-TK, GM-CSF or IL-2 gene is expressed under the control of an E4 promoter.

50 13. The adenoviral vector of claim 5, wherein an HSV-TK, GM-CSF or IL-2 gene is expressed under the control of an E1 promoter.

14. The adenoviral vector of claim 11, wherein said GM-CSF gene is expressed under the control of an E3 promoter.

15. The adenoviral vector of claim 11, wherein said IL-2 gene is expressed under the control of an E3 promoter.

55 16. The adenoviral vector of claim 11, wherein said HSV-TK gene is expressed under the control of an E3 promoter.

17. The adenoviral vector of claim 1, wherein said adenoviral E1a gene and an adenoviral E1b gene are transcribed

from a single heterologous transcriptional regulatory sequence.

18. The adenoviral vector of any of claims 1 to 6, wherein said heterologous TRS comprises a promoter or an enhancer.

5 19. A cell containing the vector according to any of claims 1-18.

20. The cell of claim 19, wherein the cell is a tumor cell or an abnormally proliferating cell.

21. A virion containing the adenoviral vector of any of claims 1 to 18.

10

22. A cell containing the virion of claim 21, wherein said cell is a producer cell for said virion.

23. A cell containing the virion of claim 21, wherein said cell is a tumor cell.

15

24. A cell containing the virion of claim 23, wherein said heterologous gene product provides anti-tumor activity in said tumor cell.

25. A method of producing the adenoviral vector of claim 1, comprising culturing the cell of claim 19 and recovering the vector from said cell.

20

26. A method of producing the virion of claim 21, comprising culturing the cell of claim 22 and recovering said virion from said cell.

25 **Patentansprüche**

1. Replikations-bedingter adenoviraler Vektor, umfassend:

30

(a) eine heterologe regulatorische Transkriptionssequenz (TRS), die mit der Kodierungssequenz eines adenoviralen, für die Replikation essentiellen Gens (GER) operabel verknüpft ist, und (b) mindestens eine zusätzliche Kodierungssequenz, die ein heterologes Genprodukt kodiert, wobei die zusätzliche kodierende Sequenz mit einer homologen regulatorischen Transkriptionssequenz (TRS) operabel verknüpft ist, die durch das Genprodukt des für die Replikation essentiellen adenoviralen Gens transaktiviert wird.

35

2. Adenoviraler Vektor nach Anspruch 1, wobei das für die Replikation essentielle adenovirale Gen (GER) ein adenovirales E1 a-Gen ist.

3. Adenoviraler Vektor nach Anspruch 2, wobei die heterologe TRS Gewebe-spezifisch ist.

40

4. Adenoviraler Vektor nach Anspruch 2 oder 3, der außerdem eine zweite zusätzliche Kodierungssequenz umfasst, die mit einer getrennten heterologen Gewebe-spezifischen regulatorischen Transkriptionssequenz operabel verknüpft ist.

45

5. Adenoviraler Vektor nach Anspruch 2 oder 3, wobei die homologe TRS aus der Gruppe, bestehend aus einem adenoviralen E1-, E2-, E3- und E4-Promotor, ausgewählt ist und durch das Genprodukt des adenoviralen E1a-Gens transaktiviert wird.

6. Adenoviraler Vektor nach Anspruch 5, wobei die homologe TRS ein adenoviraler E3-Promotor ist.

50

7. Adenoviraler Vektor nach Anspruch 1, wobei das heterologe Gen ein Cytokin-Gen ist.

8. Adenoviraler Vektor nach Anspruch 7, wobei das Cytokin-Gen Granulocyten-Macrophagen koloniestimulierender Faktor (GM-CSF) oder Interleukin-2 (IL-2) ist.

55

9. Adenoviraler Vektor nach Anspruch 1, wobei das Expressionsprodukt des heterologen Gens für das Zielgewebe toxisch ist.

10. Adenoviraler Vektor nach Anspruch 9, wobei das heterologe Gen Thymidinkinase oder Cytosindeaminase ist.

EP 1 032 696 B1

11. Adenoviraler Vektor nach Anspruch 5, wobei ein Herpes Simplex-Virus-Thymidinkinase (HSV-TK)-, ein GM-CSF- oder ein IL-2-Gen unter der Kontrolle eines E3-Promotors exprimiert wird.
- 5 12. Adenoviraler Vektor nach Anspruch 5, wobei ein HSV-TK-, ein GM-CSF- oder ein IL-2-Gen unter der Kontrolle eines E4-Promotors exprimiert wird.
13. Adenoviraler Vektor nach Anspruch 5, wobei ein HSV-TK-, ein GM-CSF- oder ein IL-2-Gen unter der Kontrolle eines E1-Promotors exprimiert wird.
- 10 14. Adenoviraler Vektor nach Anspruch 11, wobei das GM-CSF-Gen unter der Kontrolle eines E3-Promotors exprimiert wird.
- 15 15. Adenoviraler Vektor nach Anspruch 11, wobei das IL-2-Gen unter der Kontrolle eines E3-Promotors exprimiert wird.
16. Adenoviraler Vektor nach Anspruch 11, wobei das HSV-TK-Gen unter der Kontrolle eines E3-Promotors exprimiert wird.
17. Adenoviraler Vektor nach Anspruch 1, wobei das adenovirale E1a-Gen und ein adenovirales E1b-Gen aus einer einzigen heterologen regulatorischen Transkriptionssequenz transkribiert werden.
- 20 18. Adenoviraler Vektor nach einem der Ansprüche 1 bis 6, wobei die heterologe TRS einen Promotor oder einen Enhancer umfasst.
- 25 19. Zelle, die den Vektor nach einem der Ansprüche 1 bis 18 enthält.
20. Zelle nach Anspruch 19, wobei die Zelle eine Tumorzelle oder eine anormal proliferierende Zelle ist.
21. Virion, das den adenoviralen Vektor nach einem der Ansprüche 1 bis 18 enthält.
- 30 22. Zelle, die das Virion nach Anspruch 21 enthält, wobei die Zelle eine Producerzelle für das Virion ist.
23. Zelle, die das Virion nach Anspruch 21 enthält, wobei die Zelle eine Tumorzelle ist.
- 35 24. Zelle, die das Virion nach Anspruch 23 enthält, wobei das heterologe Genprodukt in der Tumorzelle Antitumoraktivität bereitstellt.
25. Verfahren zur Herstellung des adenoviralen Vektors nach Anspruch 1, das das Kultivieren der Zelle nach Anspruch 19 und das Gewinnen des Vektors aus der Zelle umfasst.
- 40 26. Verfahren zur Herstellung des Virions nach Anspruch 21, das das Kultivieren der Zelle nach Anspruch 22 und das Gewinnen des Virions aus der Zelle umfasst.

Revendications

- 45 1. Un vecteur adénoviral à réplication conditionnelle comprenant :
- (a) une séquence de régulation transcriptionnelle (TRS) hétérologue liée de manière fonctionnelle à la séquence codante d'un gène adénoviral essentiel pour la réplication (GER) et (b) au moins une séquence codante supplémentaire codant un produit de gène hétérologue, dans lequel la séquence codante supplémentaire est liée de manière fonctionnelle à une séquence de régulation transcriptionnelle (TRS) homologue qui est transactivée par le produit de gène de ce gène adénoviral essentiel pour la réplication.
- 50 2. Le vecteur adénoviral de la revendication 1, dans lequel le gène adénoviral essentiel pour la réplication (GER) est un gène adénoviral E1a.
- 55 3. Le vecteur adénoviral de la revendication 2, dans lequel la séquence TRS hétérologue a une spécificité tissulaire.

EP 1 032 696 B1

4. Le vecteur adénoviral de la revendication 2 ou 3, qui inclut de plus une seconde séquence codante supplémentaire liée de manière fonctionnelle à une séquence hétérologue de régulation transcriptionnelle à spécificité tissulaire.
5. Le vecteur adénoviral de la revendication 2 ou 3, dans lequel la séquence TRS homologue est sélectionnée parmi le groupe consistant en un promoteur adénoviral E1, E2, E3 et E4 et est transactivée par le produit génique du gène adénoviral E1a.
6. Le vecteur adénoviral de la revendication 5, dans lequel la séquence homologue TRS est un promoteur adénoviral E3.
7. Le vecteur adénoviral de la revendication 1, dans lequel le gène hétérologue est un gène de cytokine.
8. Le vecteur adénoviral de la revendication 7, dans lequel le gène de cytokine est le facteur stimulant la formation de colonies de granulocytes ou de macrophages (GM-CSF) ou l'interleukine-2 (IL-2).
9. Le vecteur adénoviral de la revendication 1, dans lequel le produit d'expression du gène hétérologue est toxique pour le tissu cible.
10. Le vecteur adénoviral de la revendication 9, dans lequel le gène hétérologue est la thymidine kinase ou la cytosine désaminase.
11. Le vecteur adénoviral de la revendication 5, dans lequel un gène de thymidine kinase du virus herpès simplex (HSV-TK), GM-CSF ou IL-2 est exprimé sous le contrôle d'un promoteur E3.
12. Le vecteur adénoviral de la revendication 5, dans lequel un gène HSV-TK, GM-CSF ou IL-2 est exprimé sous le contrôle d'un promoteur E4.
13. Le vecteur adénoviral de la revendication 5, dans lequel un gène HSV-TK, GM-CSF ou IL-2 est exprimé sous le contrôle d'un promoteur E1.
14. Le vecteur adénoviral de la revendication 11, dans lequel le gène GM-CSF est exprimé sous le contrôle d'un promoteur E3.
15. Le vecteur adénoviral de la revendication 11, dans lequel le gène IL-2 est exprimé sous le contrôle d'un promoteur E3.
16. Le vecteur adénoviral de la revendication 11, dans lequel le gène HSV-TK est exprimé sous le contrôle d'un promoteur E3.
17. Le vecteur adénoviral de la revendication 1, dans lequel le gène adénoviral E1a et un gène adénoviral E1b sont transcrits à partir d'une seule séquence de régulation transcriptionnelle hétérologue.
18. Le vecteur adénoviral selon l'une quelconque des revendications 1 à 6, dans lequel la séquence TRS hétérologue comprend un promoteur ou une séquence amplificatrice.
19. Une cellule contenant le vecteur selon l'une quelconque des revendications 1 à 18.
20. La cellule selon la revendication 19 dans laquelle la cellule est une cellule tumorale ou une cellule proliférant anormalement.
21. Un virion contenant le vecteur adénoviral de l'une quelconque des revendications 1 à 18.
22. Une cellule contenant le virion de la revendication 21, dans laquelle la cellule est une cellule productrice du virion.
23. Une cellule contenant le virion de la revendication 21, dans laquelle la cellule est une cellule tumorale.
24. Une cellule contenant le virion de la revendication 23, dans laquelle le produit du gène hétérologue a une activité anti-tumorale dans la cellule tumorale.
25. Un procédé de production du vecteur adénoviral de la revendication 1, comprenant la culture de la cellule de la

EP 1 032 696 B1

revendication 19 et le recueil du vecteur à partir de cette cellule.

- 26.** Un procédé de production du virion de la revendication 21, comprenant la culture de la cellule de la revendication 22 et le recueil du virion à partir de cette cellule.

5

10

15

20

25

30

35

40

45

50

55






Fig. 1B





==




Fig. 1C




Fig. 2A





Fig. 2C






Fig. 3A




Fig. 3B







Fig. 3C





Av5E1aTk01i (35286 bps)

digest with ClaI (bp 917)
and BstI107 (bp 5764)




Av5 digested with ClaI and BstI107 (29520 bps)

Fig. 4B


Fig. 4C

Av1E1a04i digested with BamHI and SrfI (22800 bps)



Av5E1aTk01i digested with ClaI and BstI107 (29520 bps)

Co-transfect 293 cells with
 Av1E1a04i digested with BamHI/SrfI and
 Av5E1aTk01i digested with ClaI/BstI107



Av15E1aTk04i Cytopathic Effect




Fig. 5






Fig. 6

Vector	Av5E1aTk01i	Av5E1aTk01i	Add1327	Add1327
GCV	+	-	+	-

*Limit of sensitivity = 7.4×10^4



TSR = tissue specific regulatory sequence
 GER = gene essential for replication
 TAR = transactivatable regulatory sequence
 AR, BR etc. = homologous promoter of gene A, B, etc.

Fig. 7

etc.