



DEMAND.

AU9889895

f)

(51) Classification internationale des brevets⁶ : C07H 21/00, C12N 15/11, C12Q 1/68	A1	(11) Numéro de publication internationale: WO 99/10361 (43) Date de publication internationale: 4 mars 1999 (04.03.99)
(21) Numéro de la demande internationale: PCT/FR98/01767 (22) Date de dépôt international: 7 août 1998 (07.08.98) (30) Données relatives à la priorité: 97/10643 26 août 1997 (26.08.97) FR (71) Déposant (pour tous les Etats désignés sauf US): SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATIONS SCIENTIFIQUES (S.C.R.A.S.) [FR/FR]; 51/53, rue du Docteur Blanche, F-75016 Paris (FR). (72) Inventeurs; et (75) Inventeurs/Déposants (US seulement): MARTINEZ, Jean [FR/FR]; Chemin des Romains, F-34570 Saussan (FR). GOZE, Catherine [FR/FR]; Résidence Château d'Alco, Bâtiment 8, 33, rue des Avants-Monts, F-34080 Montpellier (FR). (74) Mandataire: BOURGOUIN, André; Beaufour IPSEN - S.C.A.F., Direction de la Propriété Industrielle, 42, rue du Docteur Blanche, F-75016 Paris (FR).		(81) Etats désignés: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, brevet ARIPO (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), brevet eurasien (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), brevet européen (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), brevet OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Publiée <i>Avec rapport de recherche internationale. Avant l'expiration du délai prévu pour la modification des revendications, sera republiée si des modifications sont reçues.</i>
(54) Title: OLIGONUCLEOTIDES FOR IDENTIFYING PRECURSORS OF AMIDATED POLYPEPTIDE HORMONES		
(54) Titre: OLIGONUCLEOTIDES PERMETTANT L'IDENTIFICATION DE PRECURSEURS D'HORMONES POLYPEPTIDIQUES AMIDEES		
(57) Abstract		
<p>The invention concerns novel oligonucleotides and their application as probes for identifying RNAm's coding for precursors of amidated polypeptide hormones and, thereby, identifying novel amidated polypeptide hormones. The invention also concerns oligonucleotides of the disclosed nucleotide sequence and a method for identifying precursors of amidated hormones.</p>		
(57) Abrégé		
<p>L'invention a pour objet de nouveaux oligonucléotides et leur application comme sondes pour l'identification des ARNm codant pour les précurseurs d'hormones polypeptidiques amidées et, ainsi, l'identification de nouvelles hormones polypeptidiques amidées. L'invention concerne également des oligonucléotides de séquence nucléotidique décrite et une méthode d'identification des précurseurs d'hormones amidées.</p>		

Oligonucleotides which allow identification of precursors of amidated polypeptide hormones

5 The present invention relates to new oligonucleotides and their use as probes for identification of the mRNA which codes for precursors of amidated polypeptide hormones, and thus, to the identification of new amidated polypeptide hormones. The invention thus relates to oligonucleotides of which the nucleotide sequence is described below and a method for identification of precursors of amidated hormones.

10 Amidated polypeptide hormones are synthesized in the form of a precursor which undergoes maturation. This maturation consists of an amidation reaction.

15 The amidation reaction of the C-terminal end is a characteristic reaction of amidated polypeptide hormones. This reaction, which occurs on the precursor of one or more hormones, allows maturation of the hormone and also ensures its biostability in the physiological medium: the amide group formed is less vulnerable than the free acid function. The hormone is therefore more resistant to carboxypeptidases, it remains active in the cell for longer and retains an optimum affinity for its receptor site.

20 Amidation has been widely described ("Peptide amidation", Alan F. Bradbury and Derek G. Smyth, TIBS 16 : 112-115, March 1991 and "Functional and structural characterization of peptidylamidoglycolate lyase, the enzyme catalysing the second step in peptide amidation", A. G. Katopodis, D. S. Ping, C. E. Smith and S. W. May, Biochemistry, 30(25) : 6189-6194, June 1991), and its mechanism is as follows:

25

1 - cleavage of the precursor polypeptide chain of the hormone by an endoprotease at the two basic amino acids, that is to say arginine and/or lysine,

30

2 - subsequently two cleavages by carboxypeptidase result, which lead to the extended glycine intermediate,

35

3 - the enzyme PAM (peptidyl-glycine- α -amidating monooxygenase) comprises two distinct enzymatic activities: firstly, it converts the extended glycine intermediate into an α -hydroxyglycine derivative, the subunit of the enzyme PAM involved is PHM (peptidyl- α -glycine- α -hydroxylating monooxygenase). The derivative obtained serves as the substrate for the second subunit of PAM (called PAL: peptidyl- α -hydroxyglycine- α -amidating lyase), which fixes the amine function of the glycine on to the amino acid immediately adjacent to the N-terminal side and liberates glyoxylate.

This reaction involves the presence of a recognition site on the precursor of the hormone or hormones, a site which always comprises the sequence: glycine and two basic amino acids (arginine or lysine).

5

The amidated polypeptide hormones which are to be secreted outside the endoplasmic reticulum are known to comprise a consensus signal sequence of about fifteen to thirty amino acids, this sequence being present at the N-terminal end of the polypeptide chain. It is cut later by a signal peptidase enzyme such that it is no longer found in the protein once secreted.

10

At the present time, the discovery of a new protein is not easy. Proteins can be isolated and purified by various techniques: precipitation at the isoelectric point, selective extraction by certain solvents, then purification by crystallization, counter-current distribution, adsorption, partition or ion exchange chromatography, electrophoresis.... However, these techniques imply knowledge of the properties of the protein to be isolated. Furthermore, if a pure sample of a new protein of interest at the therapeutic level is available, there are still several stages before a genetically modified microorganism capable of synthesizing it is available.

15

20

The method proposed by the present invention offers the advantage, by using a characteristic of the peptide sequence of the precursor of all amidated hormones known to date, of allowing simultaneous detection of several new hormones of this category. This search is affected by direct identification of the nucleotide sequence which codes for the said precursors in cDNA banks prepared from tissues in which the precursors of these hormones can be synthesized.

25

The search by this method is much less restricting than the abovementioned conventional techniques of biochemistry, since:

30

- it can lead to the isolation of several distinct precursors present in the same tissue by the same principle;

35

- it allows detection, under the same technical conditions, of precursors corresponding to hormones which have very different biochemical and biological properties;

- it allows concomitant identification of all the peptide hormones which can be contained in the same precursor.

As a result, this invention allows a not insignificant saving in time and money in a sector where the costs of research and development represent a very high proportion of turnover.

5

The present invention will also allow pharmacological study of active substances having a fundamental physiological roll in the mammalian organism: hormones and more particularly amidated polypeptide neurohormones. Having available for the first time cDNA corresponding to active substances, it will then be possible to introduce the cloned vector by genetic engineering to lead to synthesis of hormones having a therapeutic use by means of microorganisms.

15 The invention first relates to a single-stranded oligonucleotide OX which can hybridize under mild conditions with an oligonucleotide OY of the sequence Y1-Y2-Y3-Y4-Y5, in which Y1 represents a nucleotide sequence of 1 to 12 nucleotides or Y1 is suppressed, Y2 represents a trinucleotide which codes for Gly, Y3 and Y4 independently represent a trinucleotide which codes for Arg or Lys and Y5 represents a nucleotide sequence of 1 to 21 nucleotides or Y5 is suppressed.

20 Nucleotide is understood as meaning a monomeric unit of RNA or DNA having the chemical structure of a nucleoside phosphoric ester. A nucleoside results from bonding of a purine base (purine, adenine, guanine or analogues) or of a pyrimidine base (pyrimidine, cytosine, uracil or analogues) with ribose or deoxyribose. An oligonucleotide is a polymer of nucleotides designating a primer sequence, a probe or a fragment of RNA or DNA.

30 The oligonucleotides mentioned can be obtained by synthesis, and there is a reference automated method which is described in the following publications: "DNA synthesis" by S. A. Narang, Tetrahedron, 39, 3 (1983) and "Synthesis and use of synthetic oligonucleotides" by K. Itakura, J. J. Rossi and R. B. Wallace, Annu. Rev. Biochem., 53, 323 (1984).

Preferably, OX can hybridize with OY under stringent conditions.

35 More preferably, OX can hybridize with an oligonucleotide OY of the sequence Y2-Y3-Y4-Y5.

Still more preferably, OX can hybridize with an oligonucleotide OY of the sequence Y1-Y2-Y3-Y4 or Y2-Y3-Y4.

5 In particular, OX can hybridize with an oligonucleotide OY such that Y5 represents a nucleotide sequence Y6-Y7-Y8-Y9, in which Y6 represents a trinucleotide which codes for Ser, Thr or Tyr, Y7 represents a trinucleotide which codes for any amino acid, Y8 represents a trinucleotide which codes for Glu or Asp and Y9 represents a nucleotide sequence comprising 1 to 12 nucleotides. More particularly, OX can hybridize with an oligonucleotide OY such that Y1 and Y9 are suppressed.

10

Especially particularly, OX can hybridize with an oligonucleotide OY in which Y2 represents a trinucleotide which codes for Gly, Y3 represents a trinucleotide which codes for Lys, Y4 represents a trinucleotide which codes for Arg and Y5 represents a sequence of 3 trinucleotides which codes for Ser-Ala-Glu.

15

This sequence was determined with the aid of a statistical study of 27 known amidation sites and led to the definition of a given pattern of amino acids over 6 positions: Gly-Lys-Arg-Ser-Ala-Glu.

20 Because of the degeneration of the genetic code and the high number of codons corresponding to Gly (4 codons), Arg (6 codons) and Ser (6 codons), the oligonucleotide sequence was constructed with the aid of two procedures which allow this degeneration to be taken into account:

25 - use of certain positions of inosine, a nucleotide in which the nitrogen base hypoxanthine pairs indiscriminately with the 4 nitrogen bases which make up the DNA,

- variation at certain positions of the nature of the nitrogen base incorporated, thus generating a number of combinations of oligonucleotides proportional to the number of different bases introduced.

30

The present invention also relates to an oligonucleotide OY comprising 9 to 42 nucleotides of the sequence Y1-Y2-Y3-Y4-Y5, in which Y1 represents a nucleotide sequence of 1 to 12 nucleotides or Y1 is suppressed, Y2 represents a trinucleotide which codes for Gly, Y3 and Y4 independently represent a trinucleotide which codes for Arg or Lys and Y5 represents a nucleotide sequence of 1 to 21 nucleotides or Y5 is suppressed.

35

Preferably, the invention relates to an oligonucleotide OY such that Y1 is suppressed or such that Y5 is suppressed.

5 The invention particularly relates to an oligonucleotide OY such that Y5 represents a nucleotide sequence Y6-Y7-Y8-Y9, in which Y6 represents a trinucleotide which codes for Ser, Thr or Tyr, Y7 represents a trinucleotide which codes for any amino acid, Y8 represents a trinucleotide which codes for Glu or Asp and Y9 represents a nucleotide sequence comprising 1 to 12 nucleotides.

10 The invention more particularly relates to an oligonucleotide OY such that Y1 and Y9 are suppressed.

15 The invention especially particularly relates to an oligonucleotide OY, characterized in that Y1 is suppressed, Y2 represents a trinucleotide which codes for Gly, Y3 represents a trinucleotide which codes for Lys, Y4 represents a trinucleotide which codes for Arg and Y5 represents a sequence of three trinucleotides which codes for Ser-Ala-Glu.

20 The present invention also relates to a single-stranded oligonucleotide OZ, characterized in that it comprises 15 to 39 nucleotides and is capable of hybridizing with a consensus signal sequence characteristic of amidated polypeptide hormones, the said sequence having Z1-Z2-Z3-Z4-Z5-Z6-Z7 as formula, in which Z1 represents a nucleotide sequence of 1 to 12 nucleotides or Z1 is suppressed, Z2 and Z3 represent two trinucleotides which code for Leu, Z4 and Z5 represent two trinucleotides which code for any two amino acids, Z6 represents a trinucleotide which codes for Leu and Z7
25 represents a nucleotide sequence of 1 to 12 nucleotides or Z7 is suppressed.

In this invention, hormone will be understood as meaning amidated polypeptide hormones of the endocrine system, and more particularly neurohormones.

30 The consensus signal sequence is a sequence carried by the precursors of proteins which are secreted by cells after their maturation.

Finally, the present invention relates to a group of oligonucleotides OX or OZ such as constitutes a combinatorial library.

35

In the invention described, combinatorial library is understood as meaning a group of oligonucleotides synthesized by taking as the model a nucleotide sequence which codes

for a sequence of amino acids of which some can be varied. Because of the degeneration of the genetic code, a group of different oligonucleotides will be obtained.

5 The invention also relates to a method for identification of the precursor of a peptide having an amidated C-terminal end, characterized by the following successive stages:

1 - obtaining of a DNA bank;

10 2 - hybridization of one or more oligonucleotides OX with the said DNA bank;

3 - identification of the DNA sequence or sequences of the said bank which hybridizes with an oligonucleotide OX;

15 4 - identification in this sequence or sequences of one or more peptides with a possible amidated C-terminal end.

A method such that the DNA bank is a cDNA bank will be preferred.

20 Complementary DNA (cDNA) is a nucleotide chain of which the sequence is complementary to that of an mRNA, the reaction leading to monocatenated cDNA being catalysed by reverse transcriptase. Bicatenated cDNA can be obtained by the action of DNA polymerase, and is then inserted with the aid of a ligase into a plasmid or a vector derived from λ bacteriophage.

25

A cDNA bank contains the cDNA corresponding to the cytoplasmic mRNA extracted from a given cell. The bank is called complete if it comprises at least one bacterial clone for each starting mRNA.

30 Hybridization takes place if two oligonucleotides have substantially complementary nucleotide sequences, they can combine over their length by establishing hydrogen bonds between complementary bases.

35 A method such that the oligonucleotide OX can be detected with the aid of a marking agent, such as ^{32}P or digoxigenin, will be particularly preferred.

The agents for radioactive marking of nucleotides most usually used are the elements which emit β -rays, for example ^3H , ^{12}C , ^{32}P , ^{33}P and ^{35}S .

Marking of the oligonucleotide is effected by addition of a phosphate group carried by (γ -³²P)-ATP on to its 5' end, this reaction being catalysed by the enzyme T4-polynucleotide kinase. Marking by digoxigenin is immunoenzymatic, the digoxigenin being combined with a nitrogen base and incorporated into the oligonucleotide. Its presence is revealed by using an antibody directed against digoxigenin and coupled to an alkaline phosphatase. The presence is revealed using the colour developed by a substrate hydrolysed by the alkaline phosphatase.

10 Other marking techniques can be employed: oligonucleotides modified chemically so that they contain a metal-complexing agent (complexes of lanthanide are often used), a group containing biotin or acridine ester, a fluorescent compound (fluorescein, rhodamine, Texas red) or others.

15 A method for identification of the precursor of the amidated polypeptide hormone such that the hybridization stage uses a combinatorial library of oligonucleotides OX will be especially particularly preferred.

The invention also relates to a method for identification of the precursor of a peptide having an amidated C-terminal end, which comprises the following stages:

1 - obtaining of a DNA bank;

25 2 - use of the PCR technique to amplify the fragment of interest with the aid of a group of oligonucleotides OX and another group of oligonucleotides OZ;

3 - identification of the DNA sequence of the said bank which hybridizes with the oligonucleotide OX and which has been amplified by the PCR reaction;

30 4 - identification in this sequence of one or more peptides with a possible amidated C-terminal end.

Fragment of interest is understood as meaning the cDNA sequence which codes for the precursor of one or more amidated polypeptide hormones.

35

The reaction of amplification of the DNA by a PCR (polymerase chain reaction) requires a DNA preparation denatured by heating at 95°C. This preparation is then paired with an excess of two complementary oligonucleotides at opposite strands of the

DNA, on both sides of the sequence to be amplified. Each oligonucleotide then serves as a primer for a DNA polymerase (extracted from thermophilic bacteria of the type *Thermus aquatilis*: Taq polymerase) for copying each of the strands of the DNA. This cycle can be repeated in an automated manner by successive denaturations-
5 renaturations.

There are numerous references detailing PCR protocols: US Patents no. 4,683,192, 4,683,202, 4,800,159 and 4,965,188, "PCR technology : principles and applications for DNA amplification", H. Erlich, ed. Stockton Press, New York (1989) and "PCR
10 protocols : a guide to methods and applications", Innis et al., eds. Academic Press, San Diego, California (1990).

Preferably, the said DNA bank is a cDNA bank.

15 More preferably, the said oligonucleotide OX can be detected with the aid of a marking agent, such as ³²P or digoxigenin.

A method for identification of the precursor of an amidated polypeptide hormone such that the amplification stage uses a combinatorial library of oligonucleotides OX and
20 another combinatorial library of oligonucleotides OZ will be particularly preferred.

The invention also relates to a method for identification of the precursor of a peptide having an amidated C-terminal end, which comprises the following stages:

25 1 - obtaining of a DNA bank;

2 - use of the PCR technique to amplify the fragment of interest with the aid of a group of oligonucleotides OX;

30 3 - identification of the DNA sequence of the said bank which hybridizes with the oligonucleotide OX and which has been amplified by the PCR reaction;

4 - identification in this sequence of one or more peptides with a possible amidated C-terminal end.

35

The aim of this method is to characterize the nucleotide sequences which code for precursors having more than one amidation site.

Preferably, the said DNA bank is a cDNA bank.

More preferably, the said oligonucleotide OX can be detected with the aid of a marking agent, such as ³²P or digoxigenin.

5

A method for the identification of the precursor of an amidated polypeptide hormone such that the amplification stage uses a combinatorial library of oligonucleotides OX will be particularly preferred.

10 Another method proposed by the present invention for identification of the precursor of a polypeptide having an amidated C-terminal end is characterized by the following stages:

1 - obtaining of a DNA bank;

15

2 - use of the PCR technique to amplify the fragment of interest with the aid of an oligonucleotide OX and another single-stranded oligonucleotide capable of hybridizing under mild or stringent conditions with a universal consensus sequence contained in the sequence of the plasmid vector in which the DNA of the said DNA bank are cloned, such as the primers T3, T7, KS, SK, M13, Reverse;

20

3 - identification of the DNA sequence of the said bank which hybridizes with an oligonucleotide OX;

25

4 - identification in this sequence of one or more peptides with a possible amidated C-terminal end.

30

The universal consensus sequence is a sequence carried by the vector in which the DNA of the bank is cloned. This sequence can serve as a primer for the sequencing. The nucleotide sequences of these primers are available in: Sambrook, J., Fritsch, E. F., Maniatis, T., *"Molecular cloning, a laboratory manual"*, 2nd edition, 1989, Colel Spring Harbor Laboratory Press.

35

The PCR reaction requires that two oligonucleotides are fixed on to the cDNA cloned in a vector for its amplification to have taken place. In the case where only a single sequence belonging to the DNA fragment to be amplified is known, a solution to overcome this problem is to use an oligonucleotide which could hybridize with a

nucleotide sequence belonging to the vector in which the cDNA has been cloned, such as a universal consensus sequence.

Preferably, the said DNA bank is a cDNA bank.

5

An oligonucleotide OY which can be detected with aid of a marking agent, such as ^{32}P or digoxigenin, will be preferred.

10 An amplification stage using a combinatorial library of oligonucleotides OX will be more particularly preferred.

EXAMPLE :

The method described by the invention has been validated by its application to a hormone which has already been isolated. The neurohormone chosen is cholecystokinin (CCK), which is the neuromediator which quantitatively is represented the most in the brain.

1.1. Preparation of the DNA matrix used for PCR reactions from a commercial bank, Lambda Zapp II (Rat Brain cDNA Library Vector, ref. 936 501) of STRATAGENE (Lafolla, USA).

This Stratagene cDNA bank contains the cloning of the cDNA of the cells of the rat brain.

1.1.1. Release of the cloned cDNA in the form of Bluescript phagemids (Stratagene, Lajolla, U.S.A.).

This is carried out in accordance with the following protocol: 250 μ l of the cDNA bank at $2 \cdot 10^8$ PFU/ml, 200 μ l of XL₁ blue bacteria (genotype: recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac [F' proAB lacI^qZ Δ M15 Tn10 (Tet^r)]^c - cf. Bullock, Fernandez, Short, *Biotechniques*, **5**, 376-379 (1987) - optical density at 600 nm: OD = 2.5) and 1 μ l of the phage ExAssistTM (cf. Hay, B., Short, J., *Strategies*, **5**, 16-18 (1992)) at 10^{10} PFU/ml are brought into contact for 15 minutes at 37°C. The entire system is then incubated on 50 ml of LB medium (composition: 10 g NaCl, 5 g yeast extract and 10 g Bactotryptone per 1 litre of sterile physiological water are mixed) for 3 hours while stirring at 37°C. The culture broth is centrifuged and the supernatant is then inactivated by heating at 70°C for 20 minutes.

1.1.2. Obtaining of the cDNA in the form of a double-stranded plasmid bank.

This stage requires 15 minutes of incubation at 37°C of 100 μ l of the inactivated supernatant and 200 μ l of SOLRTM bacteria (genotype : e14⁻(McrA⁻) Δ (mcrCB-hsdSMR-mrr)171 sbcC recB recJ uvrC umuC::Tn5 (Kan^r) lac gyrA96 relA1 thi-1 endA1 λ^R [F' proAB lacI^qZ Δ M15]^c Su⁻ (nonsuppressing) - cf. Hay, B., Short, J. M., *Strategies*, **5**(1), 16-18 (1992) - OD = 1 to 600 nm). After addition of 50 μ l ampicillin (at 100 mg/ml) and 50 ml of LB medium, the entire system is incubated at 37°C while stirring for one night. The plasmids are prepared from 50 ml of culture with the QIAGEN Plasmid Midi Kit protocol and columns from QIAGEN (the QIAGEN

columns contain an anion exchange resin with positively charged diethylaminoethanol groups on its surface which interact with the phosphates of the DNA skeleton). A DNA solution at 1.37 µg/µl was thus obtained.

5 1.2. Amplification of a portion of the precursor of CCK from the plasmid bank thus prepared.

1.2.1. Establishing the sequences of the two oligonucleotides necessary for the PCR reaction.

10

One of these two nucleotides will contain the sequence complementary to that which codes for the amidation site of CCK, which site is known and has as the sequence Gly-Arg-Arg-Ser-Ala-Glu. This oligonucleotide, which will be called *oligo CCK amide*, has as its nucleotide sequence:

15

5' CTCAGCACTGCGCCGGCC 3'

The second oligonucleotide, called *oligo CCK 5'*, corresponds to the consensus signal sequence:

20

5' GTGTGTCTGTGCGTGGTG 3'

The size of the expected amplification product is 315 base pairs, which is the distance between the sequences corresponding to these two oligonucleotides on the precursor sequence of the CCK.

25

1.2.2. PCR reaction.

30

A dilution D1 containing 1 µl of the enzyme Taq polymerase Goldstar 5 U/µl (cf. Reynier, P., Pellissier, J. F., Harle, J. R., Malthiéry, Y., *Biochemical and Biophysical Research Communications*, **205**(1), 375-380 (1994)), 1 µl of a buffer concentrated 10-fold in standard Taq polymerase and 8 µl water is prepared.

35

1 µl *oligo CCK 5'* at 250 ng/µl, 1 µl *oligo CCK amide* at 250 ng/µl, 1 µl dNTP at 10 mM each, 1 µl of the cDNA bank at 250 ng/µl, 5 µl of buffer concentrated 10-fold in the enzyme Taq polymerase, 2 µl MgCl₂ at 25 mM, 1 µl of the dilution D1 and 37 µl water are then mixed.

The amplification conditions are the following: heat treatment is first carried out for 5 minutes at 95°C, and 30 cycles are then repeated. The denaturations are carried out at 95°C for 45 seconds, the hybridization at 60°C for 30 seconds and the elongation at 72°C for 1 minute. Finally, a supplementary cycle is conducted with an elongation at 72°C for 10 minutes.

1.2.3. Results.

The results are read by migration on agarose gel at 0.8% of 1/10 of the product of the PCR reaction. In the presence of 3,8-diamino-5-ethyl-6-phenylphenanthridinium bromide (ethidium bromide), a single intense band of a size slightly greater than the marker of molecular weight 300 is visualized.

1.3. Subcloning of the PCR product into a vector which allows sequencing.

The vector used is pGEM T-easy Vector (marketed by PROGEMA Corporation, Madison, USA, ref. A 1380 - sequence given in appendix I). The stages are the following:

- purification of the band corresponding to the PCR product by electroelution,
- ligation for one night at 16°C with 1 µl of the vector pGEM T-easy at 50 ng/µl and 1 µl of ligase buffer concentrated 10-fold,
- 3 µl of product extracted from the purified band, estimated at 20 ng/µl,
- topped up to 10 µl with water.

JM 109 bacteria (genotype: e14⁻(McrA⁻) recA1 endA1 gyrA96 thi-1 hsdR17(r_K-m_K⁺) supE44 relA1 Δ(lac-proAB) [F' traD36 proAB lacI^qZΔM15] - cf. Yanish-Perron, C., Viera, J., Messing, J., *Gene*, **33**, 103-199 (1985)) are rendered competent by a treatment beforehand with CaCl₂ and are then transformed by a thermal shock of 45 seconds at 42°C with 1/5 of the ligation. The cells are then cultured on LB-ampicillin medium in a Petri dish overnight at 37°C.

The plasmid DNA of some recombinant clones are prepared. The subcloning is then verified by enzymatic digestion with Eco RI.

1.4 Sequencing

5 This is carried out by the conventional technique of dideoxynucleotides of SANGER on the vector pGEM T-easy Vector, the PCR product of 315 base pairs having been incorporated (prepared on a large scale using the QIAGEN tip 100 kit). The primer used for the sequencing is the universal oligonucleotide T7 present on the pGEM T-easy Vector plasmid.

10 1.5. Result.

The following crude sequence is obtained:

GTG TGT CTG TGC GTG GTG ATG GCA GTC CTG GCA GCA GGC GCC CTG
GCG CAG CCG GTA GTC CCT GTA GAA GCT GTG GAC CCT ATG GAG CAG
15 CGG GCG GAG GAG GCG CCC CGA AGG CAG CTG AGG GCT GTG CTC CGA
CCG GAC AGC GAG CCC CGA GCG CGC CTG GGC GCA CTG CTA GCC CGA
TAC ATC CAG CAG GTC CGC AAA GCT CCC TCT GGC CGC ATG TCC GTT
CTT AAG AAC CTG CAG GGC CTG GAC CCT AGC CAC AGG ATA AGT GAC
CGG GAC TAC ATG GGC TGG ATG GAT TTC GGC CGG CGC AGT GCT GAG

20

Translation of the sequence obtained into amino acids results in:

VCLCVV	MAVLAAGALA	QPVVPVEAVD	PMEQRAEEAP
RRQLRAVLRP	DSEPRARLGA	LLARYIQQVR	KAPSGRMSVL
KNLQGLDPSH	RISDRDYMGW	MDFGRRSAE	

25 which enables the nucleotide sequence of the precursor of CCK (the sequence of which has been provided by the Swiss Prot databank no. p01355) to be easily found.

The amino acids have the following abbreviations:

Alanine	A	Leucine	L
Argine	R	Lysine	K
Aspartic acid	D	Methionine	M
Asparagine	N	Phenylalanine	F
Cysteine	C	Proline	P
Glutamic acid	E	Serine	S
Glutamine	Q	Threonine	T
Glycine	G	Tryptophan	W
Histidine	H	Tyrosine	Y
Isoleucine	I	Valine	V

Claims

1. Single-stranded oligonucleotide OX, characterized in that it comprises 9 to
5 42 nucleotides and is capable of hybridizing under mild conditions with an
oligonucleotide OY of the sequence Y1-Y2-Y3-Y4-Y5, in which Y1 represents a
nucleotide sequence of 1 to 12 nucleotides or Y1 is suppressed, Y2 represents a
trinucleotide which codes for Gly, Y3 and Y4 independently represent a trinucleotide
which codes for Arg or Lys and Y5 represents a nucleotide sequence of 1 to 21
10 nucleotides or Y5 is suppressed.
2. Oligonucleotide OX according to claim 1, characterized in that it comprises 9 to
42 nucleotides and is capable of hybridizing under stringent conditions with an
oligonucleotide OY of the sequence Y1-Y2-Y3-Y4-Y5, in which Y1 represents a
15 nucleotide sequence of 1 to 12 nucleotides or Y1 is suppressed, Y2 represents a
trinucleotide which codes for Gly, Y3 and Y4 independently represent a trinucleotide
which codes for Arg or Lys and Y5 represents a nucleotide sequence of 1 to 21
nucleotides or Y5 is suppressed.
- 20 3. Oligonucleotide OX according to claim 1 or 2, characterized in that Y1 is suppressed
in the oligonucleotide OY.
4. Oligonucleotide OX according to claim 1, 2 or 3, characterized in that Y5 is
suppressed in the oligonucleotide OY.
- 25 5. Oligonucleotide OX according to claim 1, 2 or 3, characterized in that, in OY, Y5
represents a nucleotide sequence Y6-Y7-Y8-Y9, in which Y6 represents a trinucleotide
which codes for Ser, Thr or Tyr, Y7 represents a trinucleotide which codes for any
amino acid, Y8 represents a trinucleotide which codes for Glu or Asp and Y9 represents
30 a nucleotide sequence comprising 1 to 12 nucleotides.
6. Oligonucleotide OX according to claim 5, characterized in that Y1 and Y9 are
suppressed in the oligonucleotide OY.
- 35 7. Oligonucleotide OX according to claim 6, characterized in that it can hybridize with
the said oligonucleotide OY in which Y2 represents a trinucleotide which codes for Gly,
Y3 represents a trinucleotide which codes for Lys, Y4 represents a trinucleotide which

codes for Arg and Y5 represents a sequence of 3 trinucleotides which codes for Ser-Ala-Glu.

5 8. Single-stranded oligonucleotide OY, characterized in that it comprises 9 to 42 nucleotides of the sequence Y1-Y2-Y3-Y4-Y5, in which Y1 represents a nucleotide sequence of 1 to 12 nucleotides or Y1 is suppressed, Y2 represents a trinucleotide which codes for Gly, Y3 and Y4 independently represent a trinucleotide which codes for Arg or Lys and Y5 represents a nucleotide sequence of 1 to 21 nucleotides or Y5 is suppressed.

10

9. Oligonucleotide OY according to claim 8, characterized in that Y1 is suppressed.

10. Oligonucleotide OY according to claim 8 or 9, characterized in that Y5 is suppressed.

15

11. Oligonucleotide OY according to claim 8 or 9, characterized in that Y5 represents a nucleotide sequence Y6-Y7-Y8-Y9, in which Y6 represents a trinucleotide which codes for Ser, Thr or Tyr, Y7 represents a trinucleotide which codes for any amino acid, Y8 represents a trinucleotide which codes for Glu or Asp and Y9 represents a nucleotide sequence comprising 1 to 12 nucleotides.

20

12. Oligonucleotide OY according to claim 11, characterized in that Y1 and Y9 are suppressed.

25 13. Oligonucleotide OY according to claim 12, characterized in that Y2 represents a trinucleotide which codes for Gly, Y3 represents a trinucleotide which codes for Lys, Y4 represents a trinucleotide which codes for Arg and Y5 represents a sequence of 3 trinucleotides which codes for Ser-Ala-Glu.

30 14. Single-stranded oligonucleotide OZ, characterized in that it comprises 15 to 39 nucleotides and is capable of hybridizing under mild or stringent conditions with a consensus signal sequence characteristic of amidated polypeptide hormones, the said sequence having Z1-Z2-Z3-Z4-Z5-Z6-Z7 as formula, in which Z1 represents a nucleotide sequence of 1 to 12 nucleotides or Z1 is suppressed, Z2 and Z3 represent two trinucleotides which code for Leu, Z4 and Z5 represent two trinucleotides which code for any two amino acids, Z6 represents a trinucleotide which codes for Leu and Z7 represents a nucleotide sequence of 1 to 12 nucleotides or Z7 is suppressed.

35

15. Group of oligonucleotides OX according to any one of claims 1 to 7 or of oligonucleotides OZ according to claim 14, characterized in that it constitutes a combinatorial library.

5 **16.** Method for identification of the precursor of a peptide having an amidated C-terminal end, characterized by the following successive stages:

- obtaining of a DNA bank;

10 - hybridization of one or more oligonucleotides according to any one of claims 1 to 7 with the said DNA bank;

- identification of the DNA sequence or sequences of the said bank which hybridizes with an oligonucleotide according to any one of claims 1 to 7;

15

- identification in this sequence or sequences of one or more peptides with a possible amidated C-terminal end.

20 **17.** Method according to claim 16, characterized in that the hybridization stage uses a combinatorial library according to claim 15.

18. Method for identification of the precursor of a peptide having an amidated C-terminal end, characterized by the following successive stages:

25

- obtaining of a DNA bank;

- use of the PCR technique to amplify the fragment of interest with the aid of a group of oligonucleotides according to any one of claims 1 to 7 and another group of oligonucleotides according to claim 14;

30

- identification of the DNA sequence or sequences of the said bank which hybridizes with an oligonucleotide according to any one of claims 1 to 7;

35 - identification in this sequence or sequences of one or more peptides with a possible amidated C-terminal end.

19. Method according to claim 18, characterized in that the amplification stage uses a combinatorial library according to claim 15.

20. Method for identification of the precursor of a peptide having an amidated C-terminal end, characterized by the following successive stages:

- 5 - obtaining of a DNA bank;
- use of the PCR technique to amplify the fragment of interest with the aid of a group of oligonucleotides according to any one of claims 1 to 7;
- 10 - identification of the DNA sequence or sequences of the said bank which hybridizes with an oligonucleotide according to any one of claims 1 to 7;
- identification in this sequence or sequences of one or more peptides with a possible amidated C-terminal end.

15 **21.** Method according to claim 20, characterized in that the amplification stage uses a combinatorial library according to claim 15.

22. Method for identification of the precursor of a polypeptide having an amidated C-terminal end, characterized by the following stages:

- 20 - obtaining of a DNA bank;
- use of the PCR technique to amplify the fragment of interest with the aid of an oligonucleotide according to any one of claims 1 to 7 and another single-stranded oligonucleotide capable of hybridizing under mild or stringent conditions with a universal consensus sequence contained in the sequence of the plasmid vector in which the cDNA of the said DNA bank are cloned, such as the primers T3, T7, KS, SK, M13, Reverse;
- 25 - identification of the DNA sequence of the said bank which hybridizes with an oligonucleotide according to any one of claims 1 to 7;
- 30 - identification in this sequence of one or more peptides with a possible amidated C-terminal end.
- 35

23. Method according to claim 22, characterized in that the amplification stage uses a combinatorial library according to claim 15.

24. Method according to any one of claims 16 to 23, characterized in that the DNA bank is cDNA bank.

- 5 **25.** Method according to any one of claims 16 to 24, characterized in that the single-stranded oligonucleotide can be detected with the aid of a marking agent, such as ^{32}P or digoxigenin.

APPENDIX 1

Sequence of the pGEM®-T Easy Vector plasmid

5

The pGEM®-T Easy Vector plasmid, the sequence of which is reproduced below, was linearized with *EcoR* V at base 60 of this sequence (indicated by an asterisk). A T with two 3' ends was added to it. The T added is not included in this sequence. The sequence reproduced below corresponds to the RNA synthesized by T7 RNA polymerase and is complementary to the RNA synthesized with SP6 RNA polymerase.

10

```
1  GGGCGAATTG  GGCCCGACGT  CGCATGCTCC  CGGCCGCCAT  GGC GGCCGCG
51  GGAATTCGAT* ATCACTAGTG  AATTCGCGGC  CGCCTGCAGG  TCGACCATAT
101 GGGAGAGCTC  CCAACGCGTT  GGATGCATAG  CTTGAGTATT  CTATAGTGTC
151 ACCTAAATAG  CTTGGCGTAA  TCATGGTCAT  AGCTGTTTCC  TGTGTGAAAT
201 TGTTATCCGC  TCACAATTCC  ACACAACATA  CGAGCCGGAA  GCATAAAGTG
251 TAAAGCCTGG  GGTGCCTAAT  GAGTGAGCTA  ACTCACATTA  ATTGCGTTGC
301 GCTCACTGCC  CGCTTTCAG  TCGGGAAACC  TGTCGTGCCA  GCTGCATTAA
351 TGAATCGGCC  AACGCGCGGG  GAGAGGCGGT  TTGCGTATTG  GGC GCTCTTC
401 CGCTTCCTCG  CTCACTGACT  CGCTGCGCTC  GGTCGTTCGG  CTGCGGCGAG
451 CGGTATCAGC  TCACTCAAAG  GCGGTAATAC  GGTTATCCAC  AGAATCAGGG
501 GATAACGCAG  GAAAGAACAT  GTGAGCAAAA  GGCCAGCAAA  AGGCCAGGAA
551 CCGTAAAAG  GCCGCGTTGC  TGGCGTTTTT  CCATAGGCTC  CGCCCCCTG
601 ACGAGCATCA  CAAAATCGA  CGCTCAAGTC  AGAGGTGGCG  AAACCCGACA
```

651	GGACTATAAA	GATACCAGGC	GTTTCCCCCT	GGAAGCTCCC	TCGTGCGCTC
701	TCCTGTTCCG	ACCCTGCCGC	TTACCGGATA	CCTGTCCGCC	TTTCTCCCTT
751	CGGGAAGCGT	GGCGCTTTCT	CATAGCTCAC	GCTGTAGGTA	TCTCAGTTCG
801	GTGTAGGTCG	TTCGCTCCAA	GCTGGGCTGT	GTGCACGAAC	CCCCCGTTCA
851	GCCCGACCGC	TGCGCCTTAT	CCGTAACTA	TCGTCTTGAG	TCCAACCCGG
901	TAAGACACGA	CTTATCGCCA	CTGGCAGCAG	CCACTGGTAA	CAGGATTAGC
951	AGAGCGAGGT	ATGTAGGCGG	TGCTACAGAG	TTCTTGAAGT	GGTGGCCTAA
1001	CTACGGCTAC	ACTAGAAGGA	CAGTATTTGG	TATCTGCGCT	CTGCTGAAGC
1051	CAGTTACCTT	CGGAAAAGA	GTTGGTAGCT	CTTGATCCGG	CAAACAAACC
1101	ACCGCTGGTA	GCGGTGGTTT	TTTTGTTTGC	AAGCAGCAGA	TTACGCGCAG
1151	AAAAAAAGGA	TCTCAAGAAG	ATCCTTTGAT	CTTTTCTACG	GGGTCTGACG
1201	CTCAGTGGAA	CGAAAACCTCA	CGTTAAGGGA	TTTTGGTCAT	GAGATTATCA
1251	AAAAGGATCT	TCACCTAGAT	CCTTTTAAAT	TAAAAATGAA	GTTTTAAATC
1301	AATCTAAAGT	ATATATGAGT	AAACTTGGTC	TGACAGTTAC	CAATGCTTAA
1351	TCAGTGAGGC	ACCTATCTCA	GCGATCTGTC	TATTTGTTTC	ATCCATAGTT
1401	GCCTGACTCC	CCGTCGTGTA	GATAACTACG	ATACGGGAGG	GCTTACCATC
1451	TGGCCCCAGT	GCTGCAATGA	TACCGCGAGA	CCCACGCTCA	CCGGCTCCAG
1501	ATTTATCAGC	AATAAACCCAG	CCAGCCGGAA	GGGCCGAGCG	CAGAAGTGGT
1551	CCTGCAACTT	TATCCGCCTC	CATCCAGTCT	ATTAATTGTT	GCCGGGAAGC
1601	TAGAGTAAGT	AGTTCGCCAG	TTAATAGTTT	GCGCAACGTT	GTTGGCATTG
1651	CTACAGGCAT	CGTGGTGTC	CGCTCGTCGT	TTGGTATGGC	TTCATTCAGC
1701	TCCGGTTCCC	AACGATCAAG	GCGAGTTACA	TGATCCCCCA	TGTTGTGCAA
1751	AAAAGCGGTT	AGCTCCTTCG	GTCCTCCGAT	CGTTGTCAGA	AGTAAGTTGG

1801 CCGCAGTGTT ATCACTCATG GTTATGGCAG CACTGCATAA TTCTCTTACT
1851 GTCATGCCAT CCGTAAGATG CTTTTCTGTG ACTGGTGAGT ACTCAACCAA
1901 GTCATTCTGA GAATAGTGTA TGC GGCGACC GAGTTGCTCT TGCCCGGCGT
1951 CAATACGGGA TAATACCGCG CCACATAGCA GAACTTTAAA AGTGCTCATC
2001 ATTGGAAAAC GTTCTTCGGG GCGAAAAC TC AAGGATCT TACCGCTGTT
2051 GAGATCCAGT TCGATGTAAC CCACTCGTGC ACCCAACTGA TCTTCAGCAT
2101 CTTTTACTTT CACCAGCGTT TCTGGGTGAG CAAAACAGG AAGGCAAAAT
2151 GCCGCAAAA AGGGAATAAG GCGACACGG AAATGTTGAA TACTCATACT
2201 CTTCCTTTTT CAATATTATT GAAGCATTTA TCAGGGTTAT TGTCTCATGA
2251 GCGGATACAT ATTTGAATGT ATTTAGAAAA ATAAACAAAT AGGGGTCCG
2301 CGCACATTC CCCGAAAAGT GCCACCTGTA TGCGGTGTGA AATACCGCAC

2351 AGATGCGTAA GGAGAAAATA CCGCATCAGG CGAAATTGTA AACGTTAATA
2401 TTTTGTAAA ATTGCGGTTA AATATTTGTT AAATCAGCTC ATTTTTTAAC
2451 CAATAGGCCG AAATCGGCAA AATCCCTTAT AAATCAAAG AATAGACCGA
2501 GATAGGGTTG AGTGTTGTTT CAGTTTGAA CAAGAGTCCA CTATTAAGA
2551 ACGTGGACTC CAACGTCAA GGGCGAAAA CCGTCTATCA GGGCGATGGC
2601 CCACTACGTG AACCATCACC CAAATCAAGT TTTTTCGGT CGAGGTGCCG
2651 TAAAGCTCTA AATCGGAACC CTAAAGGGAG CCCCCGATTT AGAGCTTGAC
2701 GGGGAAAGCC GCGAACGTG GCGAGAAAGG AAGGGAAGAA AGCGAAAGGA
2751 GCGGGCGCTA GGGCGCTGGC AAGTGTAGCG GTCACGCTGC GCGTAACCAC
2801 CACACCCGCC GCGCTTAATG CGCCGCTACA GGGCGCGTCC ATTCGCCATT
2851 CAGGCTGCGC AACTGTTGGG AAGGGCGATC GGTGCGGGCC TCTTCGCTAT
2901 TACGCCAGCT GCGGAAAGG GGATGTGCTG CAAGGCGATT AAGTTGGGTA
2951 ACGCCAGGGT TTTCCAGTC ACGACGTTGT AAAACGACGG CCAGTGAATT
3001 GTAATACGAC TCACTATA