

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 April 2006 (06.04.2006)

PCT

(10) International Publication Number
WO 2006/036175 A2

(51) International Patent Classification:
G01N 33/574 (2006.01)

(21) International Application Number:
PCT/US2005/000267

(22) International Filing Date: 5 January 2005 (05.01.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/612,098 21 September 2004 (21.09.2004) US

(71) Applicant (for all designated States except US): **RHODE ISLAND HOSPITAL, A LIFESPAN-PARTNER** [US/US]; 593 Eddy Street, Providence, RI 02902 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WANDS, Jack, R.** [US/US]; 55 Claverick Street, Providence, RI 02903 (US). **KIM, Miran** [KR/KR]; 940 Quaker Lane, Apartment #1709, Warwick, RI 02818 (US).

(74) Agent: **FRASER, Janis, K.**; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: WNT PROTEINS AND DETECTION AND TREATMENT OF CANCER

(57) Abstract: The present specification provides, *inter alia*, methods of using Wnt and FZD proteins, genes, FZD and Wnt-specific antibodies and probes in diagnosis and treatment of cancer and for screening test compounds for an ability to treat cancer. Also disclosed are compounds useful for treating cancer such as liver cancer.

WO 2006/036175 A2

Wnt Proteins and Detection and Treatment of Cancer

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

This invention was made with Government support under National Institutes of Health Grant Nos. CA035711 and AA002666. The Government has certain rights in this invention.

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/612,098, filed September 21, 2004, which is incorporated herein by reference in its entirety.

TECHNICAL FIELD

This invention relates to detection and treatment of liver cancer.

BACKGROUND

Hepatocellular carcinoma (HCC) is the major primary malignant tumor of the liver. Although viral etiological factors have been identified, the molecular mechanisms that contribute to tumor progression during hepatocarcinogenesis remain largely unknown. The Frizzled family of proteins is composed of ten or more seven-transmembrane proteins that act as receptors for Wnt proteins. The Wnt/Frizzled signaling network influences diverse biological processes ranging from cell fate determination to cell motility and proliferation.

β -catenin is a multifactorial protein with a role in cell-cell adhesion that involves strengthening the linkage of cadherin and α -catenin to the actin cytoskeleton. In the absence of Wnt/Frizzled signaling, β -catenin is phosphorylated by interactions with glycogen synthase kinase (GSK)- 3β , and forms a complex with axin and the adenomatous polyposis coli protein (APC). Subsequently β -catenin is targeted for degradation by the ubiquitinproteasome system. In contrast, binding of a Wnt ligand to its Frizzled receptor stabilizes intracellular β -catenin through the inhibition of GSK- 3β enzymatic activity. Subsequently, β -catenin translocates into the nucleus in association with high mobility group domain factors such as Tcf/Lef. This complex is

associated with transcriptional up-regulation of growth regulatory and cell migration related genes.

SUMMARY

The present invention is based, in part, on the discovery that Wnt 3, 8b and 11 are ligands for Frizzled 7, which is commonly overexpressed at the mRNA and protein level in HCC, for example, in hepatitis B virus (HBV) related HCC. Liver cancer cells that overexpress Frizzled 7 exhibit enhanced cell motility and migration. Overexpression appears to be an early event during the multi-step process of hepatocyte transformation, and therefore Frizzled 7 and Wnt 3, 8b and 11 are novel molecular targets for therapy of liver cancer.

Accordingly, in one aspect, the invention provides a method for identifying an anti-cancer agent. The method includes selecting a test compound that binds to a polypeptide comprising the amino acid sequence of a Wnt 3, Wnt 8b or Wnt 11 protein or a FZD-binding fragment thereof; and determining whether the test compound is capable of (i) reducing Wnt/FZD 7 signaling in a cell, (ii) reducing liver cancer cell motility, reducing β -catenin accumulation in a liver cancer cell; or (iv) treating liver cancer in vitro or in vivo; wherein a test compound that is capable of at least one of (i) to (iv) is an anti-cancer agent. In some embodiments, selecting a test compound can include providing a polypeptide comprising the amino acid sequence of a Wnt 3, Wnt 8b or Wnt 11 protein or a FZD-binding fragment thereof, contacting the polypeptide with a test compound, detecting binding between the polypeptide and the test compound; and selecting the test compound if it binds to the polypeptide. The polypeptide to which a test compound binds can be a (i) naturally occurring polypeptide, a (ii) recombinant polypeptide, (iii) a polypeptide expressed on the surface of a cell or (iv) an isolated polypeptide. Where the polypeptide includes the amino acid sequence of a Wnt 3 protein, the polypeptide can include any one of SEQ ID NO:8 to 12. Where the the polypeptide includes the amino acid sequence of a Wnt 8b protein, the polypeptide can include any one of SEQ ID NO: 15 to 19. Where the polypeptide includes the amino acid sequence of a Wnt 11 protein, the polypeptide can include any one of SEQ ID NO: 22 to 26. In certain embodiments, the polypeptide can include of any one of SEQ ID NO: 7 to 27 and at least one non-Wnt sequence.

Anti-cancer agents identified by the methods of identifying a cancer agent described herein include, but are not limited to, an anti-Wnt antibody, e.g., a monoclonal antibody, FZD7 receptors, Wnt-binding fragments of FZD7 receptors, and other Wnt-binding compounds. Anti-cancer agents identified by these methods can be used in the treatment of cancer, e.g., liver cancer. Additionally, an anti-cancer agent identified by these methods can be used to manufacture a medicament for (i) treating liver cancer or (ii) reducing the motility of liver cancer cells in a patient.

In another aspect, the invention includes a method of treating cancer (e.g., liver cancer) in a patient. The method includes administering to the patient an effective amount of compound that binds Wnt 3, 8b or 11 and thereby reducing Wnt/ FZD 7 signaling in cells of the patient.

In still another aspect, the invention includes a method of reducing motility in a cancer cell. The method includes administering to the cell an effective amount of a compound that binds Wnt 3, 8b or 11 thereby reducing Wnt/FZD 7 signaling in the cell.

In yet another aspect, the invention includes the use of one or more compounds that bind to Wnt 3, 8b or 11 and reduce Wnt/ FZD 7 signaling in cells in a patient, in the manufacture of a medicament for treating liver cancer or reducing the motility of liver cancer cells.

In another aspect, the invention includes a method of identifying a candidate anti-cancer agent. The method includes (a) providing a first polypeptide that: (i) comprises a FZD polypeptide (e.g., a FZD 7 polypeptide) or a fragment thereof; and (ii) displays Wnt (e.g., Wnt 3, 8b or 11)-binding ability; (b) providing a second polypeptide that: (i) comprises a Wnt polypeptide (e.g., a Wnt 3, 8b or 11 polypeptide) or a fragment thereof; and (ii) displays FZD (e.g., FZD 7) binding ability; (c) contacting the first and second polypeptides in the presence of a test compound; and (d) comparing the level of binding between the first and second polypeptides in the presence of the test compound with the level of binding in the absence of the test compound, wherein a reduced level of binding in the presence of the test compound than in its absence indicates that the test compound is a candidate anti-cancer agent. The method can further include: (e) determining whether the candidate anti-cancer agent is capable of: (i) reducing Wnt/FZD 7 signaling in a cell; (ii) reducing cancer cell motility; (iii) reducing β -catenin accumulation in a cancer cell; or (iv) treating cancer *in vitro* or *in vivo*; wherein a candidate that is capable of at least one of (i) to (iv) is an anti-cancer

agent. The test compound can be selected from the group consisting of polypeptides, ribonucleic acids, small molecules, and deoxyribonucleic acids. The Wnt polypeptide can include, e.g., any one or more of SEQ ID NO:8 to 12, 15 to 19, and/or 22 to 26. The FZD polypeptide can include, e.g., any one or more of SEQ ID NO:1 to 3.

In certain embodiments, the first polypeptide is a first fusion protein comprising a FZD polypeptide (e.g., FZD 7 polypeptide) fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor; the second polypeptide is a second fusion protein comprising a Wnt polypeptide (e.g., a Wnt 3, 8b, or 11 polypeptide) fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor, wherein the Wnt polypeptide is fused to a domain different from that fused to the Wnt polypeptide; and binding of the first and second polypeptides is detected as reconstitution of a transcription factor.

Candidate anti-cancer agents identified by the methods for identifying a candidate anti-cancer agent described herein can be used in the treatment of cancer, for example, liver cancer. Additionally, candidate anti-cancer agents identified by the methods for identifying a candidate anti-cancer agent described herein can be used in the manufacture of a medicament for treating cancer, for example, liver cancer.

In certain aspects, the invention includes the use of any of the compounds described herein in the preparation of a pharmaceutical composition for the treatment or prevention of a condition described herein, e.g., cancer, e.g., liver cancer. The composition can be used in a method for treating cancer in accordance with the methods described herein. The composition can be in any form described herein, e.g., a liquid or solid composition.

Also included within the invention are nucleic acids described herein (e.g., a primer described in Table 1, below) that are useful for detecting Wnt proteins.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Suitable methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will

control. The materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

DESCRIPTION OF DRAWINGS

Fig. 1 is a picture of a silver-stained SDS-PAGE gel of proteins from Heparin-agarose affinity chromatography. Heparan sulfate proteoglycans (HSPGs) of Huh7 cells with or without heparin-treated were applied to Heparin agarose affinity resin. Eluted fractions were subjected to SDS-PAGE and stained with silver. Heparin-untreated fraction (0.25 M NaCl) showed a distinctive protein bands (*).

Fig. 2 is a set of pictures of two-dimensional gels illustrating the pattern of fractionated HSPG from a Huh7 cell line. 0.25 M NaCl eluted fractions of Heparin agarose affinity chromatography were separated onto first-dimensional pH 3-10 nonlinear IPG gels and second-dimensional 4-12% gradient NuPAGE gels. Protein spots 1 and 2 (circled) from Heparin-untreated fraction showed increased protein level.

Fig. 3 is a mass spectrum of tryptic peptides obtained from a single spot. Spot 1 from the silver stained 2-D gel of Fig. 2 was excised and destained, followed by trypsin digestion. The resulting peptide fragments were analyzed by PBSII instrument (Ciphergen). The peaks (*) represent matched peptides from human Wnt 11.

Fig. 4 are pictures of agarose gels that illustrate the detection of Wnt ligand mRNA in various hepatocellular carcinoma cell lines using RT-PCR. Wnt3 mRNA was detected in all of HCC cell lines. Wnt 8b mRNA was expressed in Hep3B. Wnt 11 mRNA was detected in 3 hepatoma cell lines except for Focus cell line. mRNA of other Wnt ligand genes were not detectable by RT-PCR method.

Figs. 5A-5E illustrate exemplary FZD7, FZD8, Wnt 3, Wnt 8b and Wnt 11 human and mouse amino acid sequences, including putative binding motifs.

DETAILED DESCRIPTION

This invention is based, at least in part, on the discovery that particular Frizzled (FZD) proteins, e.g., FZD 7 and 8, are associated with certain cancers, such as liver cancer, and that Wnt 3, 8b and 11 are Wnt 7 ligands. Accordingly, the present

specification provides, *inter alia*, methods of using Wnt and FZD proteins, genes, FZD-specific antibodies and probes in diagnosis and treatment of cancer and for screening test compounds for an ability to treat cancer. Also disclosed are compounds useful for treating cancer such as liver cancer.

5

I. Nucleic Acids, Proteins, Vectors, and Host Cells

The terms “Frizzled,” “FZD,” “Frizzled protein” and “Frizzled receptor” refer to a family of mammalian proteins related to the *Drosophila* Frizzled genes, which play a role in the development of tissue polarity. The Frizzled family comprises at least 10 mammalian genes. Exemplary human Frizzled receptors include Frizzled 1, Frizzled 2, Frizzled 3, Frizzled 4, Frizzled 5, Frizzled 6, Frizzled 7, Frizzled 8, Frizzled 9 and Frizzled 10. Frizzled receptors are involved in a dynamic model of transmembrane signal transduction analogous to G-protein-coupled receptors with amino-terminal ligand binding domains.

The terms “Wnt protein,” “Wnt ligand” and “Wnt” refer to a family of mammalian proteins related to the *Drosophila* segment polarity gene, *wingless*. In humans, the Wnt family of genes typically encode 38 to 43 kDa cysteine rich glycoproteins having hydrophobic signal sequence and a conserved asparagine-linked oligosaccharide consensus sequence (see e.g., Shimizu et al., *Cell Growth Differ* 8:1349-1358 (1997)). The Wnt family contains at least 19 mammalian members. Exemplary Wnt proteins include Wnt-1, Wnt-2, Wnt-2b (also known as Wnt-13), Wnt-3, Wnt-3A, Wnt-4, Wnt-5A, Wnt-5B, Wnt-6, Wnt-7A, Wnt-7B, Wnt-8A, Wnt-8B, Wnt-10A, Wnt-10B, Wnt-11, Wnt 14, Wnt 15, and Wnt 16.

In addition to Wnt ligands, a family of secreted Frizzled-related proteins (sFRPs) has been isolated. sFRPs appear to function as soluble endogenous modulators of Wnt signaling by competing with the membrane-spanning Frizzled receptors for the binding of secreted Wnt ligands. sFRPs can either antagonize Wnt function by binding the protein and blocking access to its cell surface signaling receptor, or they can enhance Wnt activity by facilitating the presentation of ligand to the Frizzled receptors.

The term “Wnt/FZD signaling pathway” refers to an intracellular signal transduction pathway that is initiated by an interaction between a Frizzled receptor, e.g., FZD 7, and one or more of its ligands, e.g., a Wnt protein, e.g., Wnt 3, 8b or 11. Typically, a Wnt/FZD interaction involves binding of a Wnt protein, e.g., Wnt 3, 8b or

11, to a Frizzled receptor, e.g., FZD 7, leading to activation of a signal transduction pathway. In some instances, activation of the Wnt/Frizzled signaling pathway will lead to induction of downstream-Wnt and/or FZD-inducible genes. A “downstream Wnt/FZD regulated gene product” is a protein or RNA that is regulated (e.g., up- or
5 down-regulated) as a result of signaling by a Wnt/FZD signaling pathway.

The invention includes the use of certain FZD and Wnt nucleic acids. For example, the present invention includes the use of certain FZD 7 and 8 nucleic acids, such as those that encode the amino acid sequences of the exemplary human and mouse FZD 7 (SEQ ID NOs:1 and 3, respectively) and 8 (SEQ ID NO:4 and 6, respectively)
10 receptors set forth in Figs. 5A to 5E. As another example, the invention includes the use of certain Wnt 3, 8b, and 11 nucleic acids, such as those that encode the amino acid sequences of the exemplary human and mouse Wnt 3 (SEQ ID NOs:7 and 13, respectively), 8b (SEQ ID NOs:14 and 20, respectively), and 11 (SEQ ID NOs:21 and 27, respectively) proteins set forth in Figs. 8A to 8E.

Also included within the present invention are the use of certain fragments of FZD and Wnt nucleic acids, e.g., a fragment of a nucleic acid sequence that encodes SEQ ID NOs:1, 3, 4, 6, 7, 13, 14, 20, 21, or 27. Fragments of FZD or Wnt nucleic acids encode at least one useful fragment of a FZD or Wnt polypeptide (e.g., a human or rodent polypeptide), respectively, such as a binding domain (e.g., a CRD domain) or
20 other useful fragment. For example, a useful fragment of a FZD nucleic acid may encode a fragment of a FZD receptor having binding activity, e.g., a fragment corresponding to SEQ ID NO:3 or 5. As another example, a useful fragment of an Wnt nucleic acid may encode a fragment of a Wnt polypeptide having binding activity, e.g., a fragment corresponding to any one or more of SEQ ID NOs:8 to 12, 15 to 19 and 22
25 to 26.

FZD and Wnt nucleic acids described herein include both RNA and DNA, including genomic DNA and synthetic (e.g., chemically synthesized) DNA. Nucleic acids can be double-stranded or single-stranded. Where single-stranded, the nucleic acid can be a sense strand or an antisense strand. Nucleic acids can be synthesized
30 using oligonucleotide analogs or derivatives (e.g., inosine or phosphorothioate nucleotides). Such oligonucleotides can be used, for example, to prepare nucleic acids that have altered base-pairing abilities or increased resistance to nucleases.

An "isolated nucleic acid" is a nucleic acid the structure of which is not identical to that of any naturally occurring nucleic acid or to that of any spanning more than three separate genes. The term therefore covers, for example, (a) a DNA which has the sequence of part of a naturally occurring genomic DNA molecule but is not flanked by both of the coding sequences that flank that part of the molecule in the genome of the organism in which it naturally occurs; (b) a nucleic acid incorporated into a vector or into the genomic DNA of a prokaryote or eukaryote in a manner such that the resulting molecule is not identical to any naturally occurring vector or genomic DNA; (c) a separate molecule such as a cDNA, a genomic fragment, a fragment produced by polymerase chain reaction (PCR), or a restriction fragment; and (d) a recombinant nucleotide sequence that is part of a hybrid gene, i.e., a gene encoding a fusion protein. Specifically excluded from this definition are nucleic acids present in mixtures of (i) DNA molecules, (ii) transfected cells; and (iii) cell clones, e.g., as these occur in a DNA library such as a cDNA or genomic DNA library.

In some embodiments, the invention includes the use of nucleic acid sequences that are substantially homologous to a FZD or Wnt nucleic acid. A nucleic acid sequence that is "substantially homologous" to a FZD or Wnt nucleic acid is at least 75% homologous to FZD or Wnt nucleic acid sequences that encode any one of SEQ ID NOs:1 to 27. For example, substantially homologous nucleic acid sequences can be at least about 80%, 85%, 90%, 95%, 98%, or at least about 99% homologous to sequences that encode SEQ ID NOs:1 to 27. For purposes of comparison of nucleic acids, the length of the reference nucleic acid sequence will be at least 50 nucleotides, but can be longer, e.g., at least 60 nucleotides, or more nucleotides.

As used herein, "percent homology" of two amino acid sequences or two nucleic acid sequences is determined using the algorithm of Karlin and Altschul (1990) *Proc. Nat'l Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993) *Proc. Nat'l Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990); *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches are performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to FZD or Wnt nucleic acid molecules used in the invention. BLAST protein searches are performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to a reference polypeptide. To obtain gapped alignments for comparison

purposes, Gapped BLAST is utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) are used. See the World Wide Web at address ncbi.nlm.nih.gov.

5 The invention also includes the use of nucleic acids that hybridize under stringent hybridization conditions (as defined herein) to all or a portion of nucleotide sequences that encode any of SEQ ID NOs:1 to 27, or to a complement of such nucleic acid sequences. The hybridizing portion of the hybridizing nucleic acids is typically at least 15 (e.g., 20, 25, 30, or 50) nucleotides in length. The hybridizing portion of the
10 hybridizing nucleic acid is at least about 75% (e.g., at least 80%, 90%, 95% or 98%) identical to the sequence of a portion or all of a nucleic acid encoding an FZD or Wnt polypeptide, or to its complement. Hybridizing nucleic acids of the type described herein can be used, for example, as a cloning probe, a primer (e.g., a PCR primer), or a diagnostic probe. Hybridization of the oligonucleotide probe to a nucleic acid sample
15 typically is performed under stringent conditions. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions. If sequences are to be identified that are related and substantially identical to the probe, rather than identical, then it is useful to
20 first establish the lowest temperature at which only homologous hybridization occurs with a particular concentration of salt (e.g., SSC or SSPE).

 Then, assuming that 1% mismatching results in a 1°C decrease in the T_m , the temperature of the final wash in the hybridization reaction is reduced accordingly (for example, if sequences having > 95% identity with the probe are sought, the final wash
25 temperature is decreased by 5°C). In practice, the change in T_m can be between 0.5°C and 1.5°C per 1% mismatch. Stringent conditions involve hybridizing at 68°C in 5x SSC/5x Denhardt's solution/1.0% SDS, and washing in 0.2x SSC/0.1% SDS at room temperature. Moderately stringent conditions include washing in 3x SSC at 42°C. The parameters of salt concentration and temperature can be varied to achieve the optimal
30 level of identity between the probe and the target nucleic acid. Additional guidance regarding such conditions is readily available in the art, for example, by Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, N.Y.;

and Ausubel et al. (eds.), 1995, Current Protocols in Molecular Biology, (John Wiley & Sons, N.Y.) at Unit 2.10.

Nucleic acids that hybridize to nucleotide sequence that encode any of SEQ ID NOs:1 to 27 are considered "antisense oligonucleotides."

5 Also included in the invention are genetic constructs (e.g., vectors and plasmids) that include a FZD and/or Wnt nucleic acid described herein, operably linked to a transcription and/or translation sequence to enable expression, e.g., expression vectors. A selected nucleic acid, e.g., a DNA molecule encoding a FZD or Wnt polypeptide, is "operably linked" to another nucleic acid molecule, e.g., a promoter,
10 when it is positioned in such a way that the other molecule can direct transcription and/or translation of the selected nucleic acid. For example, the selected nucleic acid can be positioned adjacent to the other nucleic acid molecule.

Also included in the invention are various engineered cells which contain a FZD and/or Wnt nucleic acid described herein. For example, the invention includes
15 transformed host cells, i.e., cells into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a nucleic acid encoding a FZD and/or Wnt polypeptide. Both prokaryotic and eukaryotic cells are included, e.g., mammalian cells (e.g., liver cells), fungi, and bacteria (such as *Escherichia coli*), and the like. An engineered cell exemplary of the type included in the invention is a liver
20 cell that overexpresses a FZD 7 transgene.

A cell that "overexpresses FZD" is a cancer cell and/or transgenic cell in which expression of a particular FZD protein, such as FZD 7 and/or 8, is at least about 1.5 times, e.g., at least about 2, 3, 4 or 5 times, the level of expression in a non-cancer cell or non-transgenic cell, respectively, from the same tissue type. In some embodiments,
25 FZD expression in a cell can be compared to expression in a non-cancer or non-transgenic cell of a different tissue-type or a panel of non-cancer or non-transgenic cells of a different tissue type. In addition, expression of one type of FZD protein (e.g., FZD 7) can be compared to other FZD proteins in the same cell. Methods for determining the level of expression of a particular gene are well known in the art. Such methods
30 include, but are not limited to, RT-PCR, real time PCR and use of antibodies against the gene products.

The use of certain FZD and Wnt polypeptides are also included within the present invention. Examples of FZD polypeptides used in the present invention are

human and mouse FZD polypeptides, such as those shown in SEQ ID NOs:1 and 3, respectively, and SEQ ID NOs:4 and 6, respectively. Examples of Wnt polypeptides used in the present invention are human and mouse Wnt 3, 8b and 11 polypeptides, such as those shown in SEQ ID NOs:7, 13, 14, 20, 21 and 27. Also included used in the present invention are certain fragments of FZD and Wnt polypeptides, e.g., fragments of SEQ ID NOs:1, 3, 4, 6, 7, 13, 14, 20, 21 and 27. Fragments of FZD and Wnt polypeptides may include at least one binding domain, or other useful portion of a full-length FZD and Wnt polypeptide. For example, useful fragments of FZD and Wnt polypeptides include, but are not limited to, fragments having binding activity (e.g., SEQ ID NOs: 2, 5, 8 to 12, 15 to 19, and 22 to 26).

The terms "protein" and "polypeptide" both refer to any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation). Thus, the terms "Frizzled protein," "Wnt protein," "Frizzled polypeptide," and "Wnt polypeptide" include full-length naturally occurring isolated proteins, as well as recombinantly or synthetically produced polypeptides that correspond to the full-length naturally occurring proteins, or to a fragment of the full-length naturally occurring or synthetic polypeptide.

As discussed above, the terms "Frizzled polypeptide," and "Wnt polypeptide" include biologically active fragments of naturally occurring or synthetic FZD and Wnt polypeptides, respectively. Fragments of a protein can be produced by any of a variety of methods known to those skilled in the art, e.g., recombinantly, by proteolytic digestion, or by chemical synthesis. Internal or terminal fragments of a polypeptide can be generated by removing one or more nucleotides from one end (for a terminal fragment) or both ends (for an internal fragment) of a nucleic acid that encodes the polypeptide. Expression of such mutagenized DNA can produce polypeptide fragments. Digestion with "end-nibbling" endonucleases can generate DNAs that encode an array of fragments. DNAs that encode fragments of a protein can also be generated, e.g., by random shearing, restriction digestion, chemical synthesis of oligonucleotides, amplification of DNA using the polymerase chain reaction, or a combination of the above-discussed methods. Fragments can also be chemically synthesized using techniques known in the art, e.g., conventional Merrifield solid phase Fmoc or t-Boc chemistry.

A purified or isolated compound is a composition that is at least 60% by weight the compound of interest, e.g., a FZD polypeptide, Wnt polypeptide, or antibody. For example, the preparation can be at least 75% (e.g., at least 90%, 95%, or even 99%) by weight the compound of interest. Purity can be measured by any appropriate method
5 known in the art, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

In certain embodiments, FZD and Wnt polypeptides include sequences substantially identical to all or portions of a naturally occurring FZD and Wnt polypeptides. Polypeptides “substantially homologous” to the FZD and Wnt
10 polypeptide sequences described herein have an amino acid sequence that is at least 65% (e.g., at least 75%, 80%, 85%, 90%, 95% or 99%, e.g., 100%), homologous to an amino acid sequence represented by SEQ ID NOs:1 to 27 (measured as described herein). For purposes of comparison, the length of the reference FZD and Wnt polypeptide sequence can be at least 16 amino acids, e.g., at least 20 or 25 amino acids.

A “conservative amino acid substitution” is one in which the amino acid residue
15 is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine,
20 asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine).

The invention also includes the use of fusion proteins (and nucleic acids that
25 encode such fusion proteins) in which a portion of a FZD (e.g., FZD 7 and/or 8) or Wnt (e.g., Wnt 3, 8b and/or 11) polypeptide is fused to an unrelated polypeptide (e.g., a marker polypeptide or a fusion partner) to create a fusion protein. For example, the polypeptide can be fused to a hexa-histidine tag or a FLAG tag to facilitate purification of bacterially expressed polypeptides or to a hemagglutinin tag or a FLAG tag to
30 facilitate purification of polypeptides expressed in eukaryotic cells. The invention also includes, for example, the use of isolated polypeptides (and the nucleic acids that encode these polypeptides) that include a first portion and a second portion, wherein the first portion includes, e.g., a FZD or Wnt polypeptide, and the second portion includes

an unrelated polypeptide, e.g., an immunoglobulin constant (Fc) region or a detectable marker.

The fusion partner can be, for example, a polypeptide that facilitates secretion, e.g., a secretory sequence. Such a fused polypeptide is typically referred to as a preprotein. The secretory sequence can be cleaved by the host cell to form the mature protein. Also within the invention are nucleic acids that encode a FZD and/or Wnt polypeptide fused to a polypeptide sequence to produce an inactive preprotein. Preproteins can be converted into the active form of the protein by removal of the inactivating sequence.

II. Methods for Detecting Cancer

Without being bound by theory, it appears that various FZD proteins, e.g., FZD 7 and 8, are important in cancer, e.g., liver cancer. In particular, hepatocytes appear to overexpress FZD 7 early during the process of transformation, e.g., prior to the development of HCC. Similarly, such cells often underexpress FZD 8. Further, it appears that Wnt 3, 8b and 11 are FZD 7 ligands.

Accordingly, the present invention provides methods of detecting cancer cells, facilitating the diagnosis of the presence and severity (e.g., tumor grade, tumor burden, and the like) of cancer in a patient, facilitating a determination of the prognosis of a patient and assessing the responsiveness of the patient to therapy (e.g., by providing a measure of therapeutic effect through, for example, assessing tumor burden during or following a chemotherapeutic regimen).

Detection can be based on detection of a polynucleotide (e.g., a FZD 7 and/or 8 polynucleotide) that is differentially expressed in a cancer cell (e.g., as compared to a non-cancer cell) and/or detection of a polypeptide (e.g., a FZD 7 and/or 8 polypeptide) encoded by a polynucleotide that is differentially expressed in a cancer cell. The detection methods of the invention can be conducted *in vitro* or *in vivo*, on a biological sample, e.g., isolated cells and/or whole tissues.

A "biological sample" as used herein is a sample of biological tissue or fluid that contains nucleic acids or polypeptides, e.g., a FZD 7 protein, polynucleotide or transcript. Such samples include, but are not limited to, tissue obtained from, e.g., liver, lung, lymph nodes, colon, stomach, pancreas, bile duct, small bowel and/or esophagus. Biological samples may also include sections of tissues such as biopsy and

autopsy samples, frozen sections taken for histologic purposes, blood, plasma, serum, sputum, stool, tears, mucus, bile, saliva, lymph, hair, skin, etc. Biological samples also include explants and primary and/or transformed cell cultures derived from patient tissues. A biological sample is typically obtained from a eukaryotic organism, e.g., a
5 primate such as a chimpanzee or human; cow; horse; goat; sheep; dog; cat; a rodent, e.g., guinea pig, rat or mouse; rabbit; bird; reptile; or fish. A sample is usually provided by removing a sample of cells from an animal, but can also be accomplished by providing previously isolated cells (e.g., isolated by another person, at another time and/or for another purpose), or by performing the methods of the invention *in vivo*.
10 Archival tissues, having treatment or outcome history, can be used.

In some embodiments, methods are provided for detecting a cancer cell by detecting expression in the cell of a transcript (e.g., a FZD 7 and/or 8 transcript) that is differentially expressed in a cancer cell. Any of a variety of known methods can be used for detection including but not limited to, detection of a transcript by hybridization
15 of mRNA with an appropriate hybridization probe; detection of a transcript by a polymerase chain reaction using specific oligonucleotide primers; and *in situ* hybridization using an appropriate hybridization probe. The methods can be used to detect and/or measure mRNA levels of a gene that is differentially expressed in a cancer cell. In some embodiments, the methods comprise: a) contacting a sample with
20 a polynucleotide that corresponds to a differentially expressed gene described herein under conditions that allow hybridization; and b) detecting hybridization, if any.

Detection of differential hybridization, when compared to a suitable control, is an indication of the presence in the sample of a polynucleotide that is differentially
25 expressed in a cancer cell. Appropriate controls include, for example, a sample that is not a cancer cell, a sample that is known not to contain a polynucleotide that is differentially expressed in a cancer cell, and use of a labeled polynucleotide of the same "sense" as the polynucleotide that is differentially expressed in the cancer cell. Conditions that allow hybridization are known in the art and have been described in more detail above. Detection can also be accomplished by any known method,
30 including, but not limited to, *in situ* hybridization, PCR (polymerase chain reaction) and/or RT-PCR (reverse transcription-PCR), or combinations of known techniques. A variety of labels and labeling methods for polynucleotides are known in the art and can

be used in the assay methods of the invention. Specificity of hybridization can be determined by comparison to appropriate controls.

Polynucleotides generally comprising at least 10 nt, at least 12 nt or at least 15 contiguous nucleotides of a polynucleotide described herein, such as those having the sequence as depicted herein, can be used for a variety of purposes, such as probes or PCR primers for detection and/or measurement of transcription levels of a polynucleotide that is differentially expressed in a cancer cell. As will be appreciated by the skilled artisan, the probe can be detectably labeled and contacted with, for example, an array comprising immobilized polynucleotides obtained from a test sample (e.g., mRNA). Alternatively, the probe can be immobilized on an array and the test sample detectably labeled. The use of these and other variations of the methods of the invention are well within the skill in the art and are within the scope of the invention.

Nucleotide probes can be used to detect expression of a gene corresponding to the provided polynucleotide. In Northern blots, mRNA is separated electrophoretically and contacted with a probe. A probe is detected as hybridizing to an mRNA species of a particular size. The amount of hybridization can be quantified to determine relative amounts of expression. Probes can be used for *in situ* hybridization to cells to detect expression. Probes can also be used *in vivo* for diagnostic detection of hybridizing sequences. Probes can be labeled with a radioactive isotope or other types of detectable labels, e.g., chromophores, fluorophores and/or enzymes. Other examples of nucleotide hybridization assays are described in WO92/02526 and U.S. Pat. No. 5,124,246.

PCR is another means for detecting small amounts of target nucleic acids (see, e.g., Mullis et al., Meth. Enzymol. (1987) 155:335; U.S. Pat. No. 4,683,195; and U.S. Pat. No. 4,683,202). Two primer oligonucleotides that hybridize with the target nucleic acids can be used to prime the reaction. The primers can be composed of sequence within or 3' and 5' to the polynucleotides described herein. After amplification of the target by standard PCR methods, the amplified target nucleic acids can be detected by methods known in the art, e.g., Southern blot. mRNA or cDNA can also be detected by traditional blotting techniques (e.g., Southern blot, Northern blot, etc.) described in Sambrook et al., "Molecular Cloning: A Laboratory Manual" (New York, Cold Spring Harbor Laboratory, 1989) (e.g., without PCR amplification). In general, mRNA or cDNA generated from mRNA using a polymerase enzyme can be purified and separated using gel electrophoresis, and transferred to a solid support, such as

nitrocellulose. The solid support can be exposed to a labeled probe and washed to remove any unhybridized probe. Duplexes containing the labeled probe can then be detected.

5 Methods using PCR amplification can be performed on the DNA from one or more cells. The use of the polymerase chain reaction is described in Saiki et al. (1985) Science 239:487, and a review of techniques may be found in Sambrook et al., "Molecular Cloning: A Laboratory Manual" (New York, Cold Spring Harbor Laboratory, 1989; pp.14.2-14.33). A detectable label may be included in the amplification reaction. Suitable detectable labels include fluorochromes, (e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, 10 allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein, 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4,4',5',7,7' - hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA)), radioactive labels, (e.g., ³²P, ³⁵S, ³H, etc.), and the like. The label may be a two stage system, where the polynucleotide is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the 20 amplification product.

In one embodiment, expression level is assessed by using real time PCR. RNA is isolated from a sample of interest. PCR primers are designed to amplify the specific gene of interest. PCR product accumulation is measured using a dual-labeled fluorescent oligonucleotide probe. The probe is labeled with two different fluorescent dyes, the 5' terminus reporter dye and the 3' terminus quenching dye. The 25 oligonucleotide probe is selected to be homologous to an internal target sequence present in the PCR amplicon. When the probe is intact, energy transfer occurs between the two fluorophores, and the fluorescent emission is quenched. During the extension phase of PCR, the probe is cleaved by 5' nuclease activity of Taq polymerase. Therefore, the reporter is no longer in proximity to the quencher, and the increase in 30 emission intensity is measured. An exemplary method for detecting FZD expression using real time PCR is provided in the Examples section, below. The primers can also

be used in other methods, for example RT-PCR. This assay provides a quantitative measure of nucleic acid.

In other embodiments, methods are provided for detecting a cancer cell by detecting expression of a protein (e.g., a FZD 7 and/or 8 protein) that is differentially expressed by the cell. Any of a variety of known methods can be used for detection, including but not limited to methods that employ binding compounds, e.g., antibodies, e.g., as is useful in ELISA and/or Western blotting methods. Such antibodies can be polyclonal or monoclonal, or antigen binding fragment thereof, and can be labeled with a detectable marker (e.g., fluorophore, chromophore or isotope, etc). Where appropriate, the compound can be attached to a solid support such as a bead, plate, filter, resin, etc. Determination of formation of the compound/target complex can be effected by contacting the complex with a further compound (e.g., a secondary antibody) that specifically binds to the first compound (or complex). Like the first compound, the further compound can be attached to a solid support and/or can be labeled with a detectable marker.

The materials needed to perform the detection methods described herein can be provided as part of a kit. Thus, the invention further provides kits for detecting the presence and/or a level of a polynucleotide that is differentially expressed in a cancer cell (e.g., by detection of an mRNA encoded by the differentially expressed gene of interest), and/or a polypeptide encoded thereby, in a biological sample. Procedures using these kits can be performed by clinical laboratories, experimental laboratories, medical practitioners or private individuals. The kits of the invention for detecting a polypeptide encoded by a polynucleotide that is differentially expressed in a cancer cell may comprise a moiety, such as an antibody, that specifically binds the polypeptide. The kits of the invention used for detecting a polynucleotide that is differentially expressed in a cancer cell may comprise a moiety that specifically hybridizes to such a polynucleotide. The kit may optionally provide additional components that are useful in the procedure including, e.g., buffers, developing reagents, labels, reacting surfaces, means for detection, control samples, standards, instructions, and interpretive information.

The present invention further relates to methods of detecting/diagnosing a neoplastic or preneoplastic condition in a mammal (for example, a human). "Diagnosis" as used herein generally includes determination of a patient's susceptibility

to a disease or disorder, determination as to whether a subject is presently affected by a disease or disorder, prognosis of a subject affected by a disease or disorder (e.g., identification of pre-metastatic or metastatic cancerous states, stages of cancer, or responsiveness of cancer to therapy), and therametrics (e.g., monitoring a subject's condition to provide information as to the effect or efficacy of therapy).

One exemplary detection/diagnostic method includes: (a) obtaining from a mammal (e.g., a human) a biological sample (e.g., liver tissue), (b) detecting in the sample the presence of a FZD 7 and/or 8 gene product (e.g., protein or mRNA), and (c) comparing the amount of FZD 7 and/or 8 gene product present with that in a control sample. In accordance with this method, the presence in the sample of elevated levels of FZD 7 gene product and/or reduced levels of FZD 8 gene product indicates that the subject has a neoplastic or preneoplastic condition, e.g., liver cancer or a risk for developing liver cancer.

The identification of elevated levels of FZD 7 protein and/or reduced levels of FZD 8 protein in accordance with the present invention makes possible the identification of patients that are likely to benefit from specialized therapy. For example, a biological sample from a post primary therapy subject (e.g., subject having undergone surgery) can be screened for the presence of elevated levels of the protein, determined by studies of normal populations, being indicative of residual tumor tissue. Similarly, tissue surrounding the cut site of a surgically removed tumor (e.g., peritumorous tissue) can be examined (e.g., by immunofluorescence), the presence of elevated levels of FZD 7 or reduced levels of FZD 8 (relative to the surrounding tissue) being indicative of potential development of the disease in this tissue or incomplete removal of the tumor. The ability to identify such patients makes it possible to tailor therapy to the needs of the particular patient. Subjects undergoing non-surgical therapy, e.g., chemotherapy or radiation therapy, can also be monitored, the presence in samples from such subjects of elevated levels of FZD 7 or reduced levels of FZD 8 being indicative of the need for continued treatment. Skilled practitioners will also appreciate that staging of cancer (e.g., liver cancer) for purposes of optimizing treatment regimens can be performed using the methods described herein.

III. Methods for Identifying Compounds Capable of Treating Cancer

The invention provides methods for screening test compounds for an ability to treat cancer, e.g., liver cancer. A “test compound” as described herein is any compound that can be screened using the methods described herein. For example, a
5 test compound can be, e.g., a small organic or inorganic molecule (M.W. less than 1,000 Da). Alternatively or in addition, the test compound can be a polypeptide (e.g., a polypeptide having a random or predetermined amino acid sequence or a naturally-occurring or synthetic polypeptide) or a nucleic acid, such as a DNA or RNA molecule. A test compound can be naturally occurring (e.g., an herb or a natural product), or
10 synthetic, or can include both natural and synthetic components. A test compound can have a formula weight of less than about 10,000 grams per mole, less than 5,000 grams per mole, less than 1,000 grams per mole, or less than about 500 grams per mole. The test compound can be, for example, any organic or inorganic compound (e.g., heteroorganic or organometallic compound), an amino acid, amino acid analog,
15 polypeptide, peptidomimetic (e.g., peptoid), oligopeptide (e.g., from about 5 to about 25 amino acids in length, preferably from about 10 to 20 or 12 to 18 amino acids in length, preferably 12, 15, or 18 amino acids in length), nucleotide, nucleotide analog, polynucleotide, polynucleotide analog, ribonucleic acid, deoxyribonucleic acid, antisense oligonucleotide, ribozyme, saccharide, lipid (e.g., a sphingolipid), and/or a
20 fatty acid, or any combination thereof.

The terms “antagonist” or “inhibitor” of Wnt/FZD signaling (e.g., Wnt/FZD 7 signaling) refer to compounds that, e.g., bind to Wnt proteins (e.g., Wnt 3, 8, and/or 11) and/or FZD receptors (e.g., FZD 7) and/or partially or totally block or inhibit Wnt/FZD signaling (e.g., Wnt/FZD 7 signaling) as measured in known assays for Wnt/FZD
25 signaling (e.g., measurement of β -catenin levels, oncogene expression controlled by Tcf and Lef transcription factors or other downstream Wnt/Frizzled regulated gene products). Inhibitors include, e.g., antibodies directed against Wnt or FZD proteins, modified versions of Wnt or FZD proteins, naturally occurring and synthetic ligands, antagonists, agonists, antibodies, small chemical molecules, and the like. Assays for
30 detecting inhibitors or antagonists are described in more detail below.

Libraries of Test Compounds

In certain embodiments, screens of the present invention utilize libraries of test compounds. A "library" is a collection of compounds (e.g., as a mixture or as physically separated individual compounds) synthesized from various combinations of one or more starting components. At least some of the compounds must differ from at least some of the other compounds in the library. A library can include, e.g., 5, 10, 50, 100, 1000, or even 10,000, 50,000, or 100,000, or more different compounds (i.e., not simply multiple copies of the same compounds, although some compounds in the library may be duplicated or represented more than once). Each of the different compounds will be present in an amount such that its presence can be determined by some means, e.g., can be isolated, analyzed, and/or detected with a receptor or suitable probe. The actual quantity of each different compound needed so that its presence can be determined will vary due to the actual procedures used and may change as the technologies for isolation, detection, and analysis advance. When the compounds are present in a mixture in substantially equimolar amounts, for example, an amount of 100 picomoles of each compound can often be detected. Libraries can include both libraries of individual compounds (e.g., present substantially as a single type of compound-per-well, made via parallel synthesis or the pool and split pool method) and mixtures containing substantially equimolar amounts of each desired compound (i.e., wherein no single compound dominates). Either library format can allow identification of an active compound discovered in an assay.

Test compounds can be screened individually or in parallel. An example of parallel screening is a high throughput drug screen of large libraries of chemicals. Such libraries of candidate compounds can be generated or purchased, e.g., from Chembridge Corp., San Diego, CA. Alternatively, prior experimentation and anecdotal evidence can suggest a class or category of compounds of enhanced potential. A library can be designed and synthesized to cover such a class of chemicals.

The synthesis of combinatorial libraries is well known in the art and has been reviewed (see, e.g., E.M. Gordon *et al.*, *J. Med. Chem.* (1994) 37:1385-1401 ; DeWitt, S. H.; Czarnik, A. W. *Acc. Chem. Res.* (1996) 29:114; Armstrong, R. W.; Combs, A. P.; Tempest, P. A.; Brown, S. D.; Keating, T. A. *Acc. Chem. Res.* (1996) 29:123; Ellman, J. A. *Acc. Chem. Res.* (1996) 29:132; Gordon, E. M.; Gallop, M. A.; Patel, D. V. *Acc. Chem. Res.* (1996) 29:144; Lowe, G. *Chem. Soc. Rev.* (1995) 309, Blondelle *et al.* *Trends Anal. Chem.* (1995) 14:83; Chen *et al.* *J. Am. Chem. Soc.* (1994) 116:2661;

U.S. Patents 5,359,115, 5,362,899, and 5,288,514; PCT Publication Nos. WO92/10092, WO93/09668, WO91/07087, WO93/20242, WO94/08051).

Libraries of compounds can be prepared according to a variety of methods, some of which are known in the art. For example, a "split-pool" strategy can be implemented in the following way: beads of a functionalized polymeric support are placed in a plurality of reaction vessels; a variety of polymeric supports suitable for solid-phase peptide synthesis are known, and some are commercially available (for examples, see, e.g., M. Bodansky "Principles of Peptide Synthesis", 2nd edition, Springer-Verlag, Berlin (1993)). To each aliquot of beads is added a solution of a different activated amino acid, and the reactions are allowed to proceed to yield a plurality of immobilized amino acids, one in each reaction vessel. The aliquots of derivatized beads are then washed, "pooled" (i.e., recombined), and the pool of beads is again divided, with each aliquot being placed in a separate reaction vessel. Another activated amino acid is then added to each aliquot of beads. The cycle of synthesis is repeated until a desired peptide length is obtained. The amino acid residues added at each synthesis cycle can be randomly selected; alternatively, amino acids can be selected to provide a "biased" library, e.g., a library in which certain portions of the inhibitor are selected non-randomly, e.g., to provide an inhibitor having known structural similarity or homology to a known peptide capable of interacting with an antibody, e.g., the an anti-idiotypic antibody antigen binding site. It will be appreciated that a wide variety of peptidic, peptidomimetic, or non-peptidic compounds can be readily generated in this way.

The "split-pool" strategy can result in a library of peptides, e.g., modulators, which can be used to prepare a library of test compounds of the invention. In another illustrative synthesis, a "diversomer library" is created by the method of Hobbs DeWitt *et al.* (*Proc. Natl. Acad. Sci. U.S.A.* 90:6909 (1993)). Other synthesis methods, including the "tea-bag" technique of Houghten (see, e.g., Houghten *et al.*, *Nature* 354:84-86 (1991)) can also be used to synthesize libraries of compounds according to the subject invention.

Libraries of compounds can be screened to determine whether any members of the library have a desired activity, and, if so, to identify the active species. Methods of screening combinatorial libraries have been described (see, e.g., Gordon *et al.*, *J Med. Chem.*, *supra*). Soluble compound libraries can be screened by affinity

chromatography with an appropriate receptor to isolate ligands for the receptor, followed by identification of the isolated ligands by conventional techniques (e.g., mass spectrometry, NMR, and the like). Immobilized compounds can be screened by contacting the compounds with a soluble receptor; preferably, the soluble receptor is conjugated to a label (e.g., fluorophores, colorimetric enzymes, radioisotopes, luminescent compounds, and the like) that can be detected to indicate ligand binding. Alternatively, immobilized compounds can be selectively released and allowed to diffuse through a membrane to interact with a receptor. Exemplary assays useful for screening libraries of test compounds are described above.

Screening Methods

The invention provides methods for identifying compounds capable of treating cancer, e.g., liver cancer. Although applicants do not intend to be bound by any particular theory as to the biological mechanism involved, such compounds are thought to modulate specifically (1) Wnt/FZD signaling (e.g., by binding to FZD 7, Wnt 3, Wnt 8b and/or Wnt 11 polypeptides and/or reducing (e.g., preventing) Wnt/FZD-mediated transcription) and/or (2) expression of FZD 7 and/or FZD 8.

In certain aspects of the present invention, screening for such compounds is accomplished by (i) identifying from a group of test compounds those that bind to FZD 7, Wnt 3, Wnt 8b and/or Wnt 11 polypeptide, modulate (i.e., increase or decrease) an interaction between FZD 7 and its ligand (e.g., Wnt 3, Wnt 8b and/or Wnt 11) and/or modulate (i.e., increase or decrease) transcription and/or translation of FZD 7 and/or FZD 8; and, optionally, (ii) further testing such compounds for their ability to modulate Wnt/FZD signaling, reduce cancer cell motility, reduce β -catenin accumulation in cancer cells and/or to treat cancer *in vitro* or *in vivo*. Test compounds that bind to FZD 7, Wnt 3, Wnt 8b and/or Wnt 11 polypeptides, modulate an interaction between FZD 7 and its ligand (e.g., Wnt 3, Wnt 8b and/or Wnt 11), or modulate transcription and/or translation of FZD 7 and/or FZD 8, are referred to herein as "candidate anti-cancer agents." Candidate anti-cancer agents further tested and found to be capable of modulating *in vitro* or *in vivo* Wnt/FZD signaling, reducing cancer cell motility, reduce β -catenin accumulation in cancer cells and/or treating cancer are considered "anti-cancer agents." In the screening methods of the present invention, candidate anti-cancer agents can be, but do not necessarily have to be, tested to determine whether

they are anti-cancer agents. Assays of the present invention may be carried out in biological samples, whole cell preparations and/or *ex vivo* cell-free systems.

In one aspect, the invention includes methods for screening test compounds to identify compounds that bind to FZD polypeptides, e.g., FZD 7 polypeptides, and/or to Wnt polypeptides, e.g., Wnt 3, 8b and/or 11 polypeptides. Binding of a test compound to a FZD or Wnt polypeptide can be detected, for example, *in vitro* by reversibly or irreversibly immobilizing the test compound(s) or the Wnt or FZD polypeptide on a substrate, e.g., the surface of a well of a 96-well polystyrene microtitre plate. Methods for immobilizing polypeptides and other small molecules are well known in the art. For example, microtitre plates can be coated with a FZD or Wnt polypeptide by adding the polypeptide in a solution (typically, at a concentration of 0.05 to 1 mg/ml in a volume of 1-100 μ l) to each well, and incubating the plates at room temperature to 37°C for a given amount of time, e.g., for 0.1 to 36 hours. Polypeptides not bound to the plate can be removed by shaking excess solution from the plate, and then washing the plate (once or repeatedly) with water or a buffer. Typically, the polypeptide is in water or a buffer. The plate can then be washed with a buffer that lacks the bound polypeptide. To block the free protein-binding sites on the plates, plates can be blocked with a protein that is unrelated to the bound polypeptide. For example, 300 μ l of bovine serum albumin (BSA) at a concentration of 2 mg/ml in Tris-HCl can be used. Suitable substrates include those substrates that contain a defined cross-linking chemistry (e.g., plastic substrates, such as polystyrene, styrene, or polypropylene substrates from Corning Costar Corp. (Cambridge, MA), for example). If desired, a particle, e.g., beaded agarose or beaded sepharose, can be used as the substrate. Test compounds can then be added to the coated plate and allowed to bind to the FZD or Wnt polypeptide (e.g., at 37°C for 0.5 - 12 hours). The plate can then be rinsed as described above. Skilled practitioners will appreciate that many variations of this method are possible. For example, the method can include coating a substrate with a test compound and adding Wnt or FZD polypeptides to the substrate-bound compound.

Binding of FZD or Wnt to a second compound, e.g., the test compound described above or to a binding partner (e.g., FZD 7 to Wnt 3, 8b and/or 11; discussed in further detail below), can be detected by any of a variety of art-known methods. For example, an antibody that specifically binds to a FZD or Wnt polypeptide (i.e., an anti-FZD antibody, e.g., the polyclonal antibody described in the Examples section, or an

anti-Wnt antibody) can be used in an immunoassay. If desired, the antibody can be labeled (e.g., fluorescently or with a radioisotope) and detected directly (*see*, e.g., West and McMahon, *J. Cell Biol.* 74:264, 1977). Alternatively, a second antibody can be used for detection (e.g., a labeled antibody that binds to the Fc portion of the anti-FZD or anti-Wnt antibody). In an alternative detection method, the FZD or Wnt polypeptide is labeled (e.g., with a radioisotope, fluorophore, chromophore, or the like), and the label is detected. In still another method, a FZD or Wnt polypeptide is produced as a fusion protein with a protein that can be detected optically, e.g., green fluorescent protein (which can be detected under UV light). In an alternative method, the polypeptide is produced as a fusion protein with an enzyme having a detectable enzymatic activity, such as horseradish peroxidase, alkaline phosphatase, β -galactosidase, or glucose oxidase. Genes encoding all of these enzymes have been cloned and are available for use by skilled practitioners. If desired, the fusion protein can include an antigen or epitope that can be detected and measured with a polyclonal or monoclonal antibody using conventional methods. Suitable antigens include enzymes (e.g., horse radish peroxidase, alkaline phosphatase, and β -galactosidase) and non-enzymatic polypeptides (e.g., serum proteins, such as BSA and globulins, and milk proteins, such as caseins).

In various methods for identifying polypeptides (e.g., test polypeptides) that bind to a FZD or Wnt polypeptides, the conventional two-hybrid assays of protein/protein interactions can be used (*see* e.g., Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578, 1991; Fields et al., U.S. Pat. No. 5,283,173; Fields and Song, *Nature*, 340:245, 1989; Le Douarin et al., *Nucleic Acids Research*, 23:876, 1995; Vidal et al., *Proc. Natl. Acad. Sci. USA*, 93:10315-10320, 1996; and White, *Proc. Natl. Acad. Sci. USA*, 93:10001-10003, 1996). Generally, two-hybrid methods involve reconstitution of two separable domains of a transcription factor. One fusion protein includes the FZD or Wnt polypeptide fused to either a transactivator domain or DNA binding domain of a transcription factor (e.g., of Gal4). The other fusion protein contains a test polypeptide or a binding partner for the polypeptide included in the first fusion protein, fused to either the DNA binding domain or a transactivator domain of a transcription factor. Binding of the FZD or Wnt polypeptide to the test polypeptide or binding partner reconstitutes the transcription factor. Reconstitution of the transcription factor can be detected by detecting expression of a gene (i.e., a reporter gene) that is operably linked

to a DNA sequence that is bound by the DNA binding domain of the transcription factor. Kits for practicing various two-hybrid methods are commercially available (e.g., from Clontech; Palo Alto, CA).

5 In another aspect, the invention includes methods for screening test compounds to identify a compound that modulates a protein-protein interaction between FZD and Wnt polypeptides. A method useful for high throughput screening of compounds capable of modulating protein-protein interactions between transcriptional regulators is described in Lepourcelet et al., *Cancer Cell* 5: 91-102 (2004), which is incorporated herein by reference in its entirety. Typically, a first compound is provided. The first
10 compound is a FZD (e.g., FZD 7) or Wnt (e.g., Wnt 3, 8b, or 11) polypeptide or biologically active fragment thereof. A second compound is provided that is different from the first compound and is labeled. The second compound is a FZD (e.g., FZD 7) or Wnt (e.g., Wnt 3, 8b, or 11) polypeptide or biologically active fragment thereof. A test compound is provided. The first compound, second compound and test compound
15 are contacted with each other. The amount of label bound to the first compound is then determined. A change in protein-protein interaction between the first compound and the second compound as assessed by label bound is indicative of the usefulness of the test compound in modulating a protein-protein interaction between the FZD and Wnt polypeptide.

20 In certain embodiments, the first compound provided is attached to a solid support. Solid supports include, e.g., resins (e.g., agarose and beads) and multiwell plates. In certain embodiments, the method includes a washing step after the contacting step, so as to separate bound and unbound label.

In certain embodiments, a plurality of test compounds is contacted with the first
25 compound and second compound. The different test compounds can be contacted with the other compounds in groups or separately. In certain embodiments, each of the test compounds is contacted with both the first compound and the second compound in an individual well. For example, the method can screen libraries of test compounds. Libraries of test compounds are discussed in detail above. Libraries can include, e.g.,
30 natural products, organic chemicals, peptides, and/or modified peptides, including, e.g., D-amino acids, unconventional amino acids, and N-substituted amino acids. Typically, the libraries are in a form compatible with screening in multiwell plates, e.g., 96-well plates. The assay is particularly useful for automated execution in a multiwell format

in which many of the steps are controlled by computer and carried out by robotic equipment. The libraries can also be used in other formats, e.g., synthetic chemical libraries affixed to a solid support and available for release into microdroplets.

In certain embodiments, the first compound is a FZD 7 polypeptide or fragment thereof and the second compound is a Wnt polypeptide, such as Wnt 3, 8b, or 11, or fragment thereof. In other embodiments, the first compound is a Wnt polypeptide, such as Wnt 3, 8b, or 11 polypeptide or fragment thereof, and the second compound is a FZD 7 polypeptide or fragment thereof. The solid support to which the first compound is attached can be, e.g., sepharose beads, SPA beads (microspheres that incorporate a scintillant) or a multiwell plate. SPA beads can be used when the assay is performed without a washing step, e.g., in a scintillation proximity assay. Sepharose beads can be used when the assay is performed with a washing step. The second compound can be labeled with any label that will allow its detection, e.g., a radiolabel, a fluorescent agent, biotin, a peptide tag, or an enzyme fragment. The second compound can also be radiolabeled, e.g., with ^{125}I or ^3H .

In certain embodiments, the enzymatic activity of an enzyme chemically conjugated to, or expressed as a fusion protein with, the first or second compound, is used to detect bound protein. A binding assay in which a standard immunological method is used to detect bound protein is also included. In certain other embodiments, the interaction of Wnt and FZD polypeptides or fragments thereof is detected by fluorescence resonance energy transfer (FRET) between a donor fluorophore covalently linked to a FZD or Wnt polypeptide (e.g., a fluorescent group chemically conjugated to FZD or Wnt, or a variant of green fluorescent protein (GFP) expressed as an FZD or Wnt-GFP chimeric protein) and an acceptor fluorophore covalently linked to a substrate protein, where there is suitable overlap of the donor emission spectrum and the acceptor excitation spectrum to give efficient nonradiative energy transfer when the fluorophores are brought into close proximity through the protein-protein interaction of FZD and Wnt polypeptides.

In other embodiments, the protein-protein interaction is detected by reconstituting domains of an enzyme, e.g., beta-galactosidase (see Rossi et al, Proc. Natl. Acad. Sci. USA 94:8405-8410 (1997)).

In still other embodiments, the protein-protein interaction is assessed by fluorescence ratio imaging (Bacsikai et al, Science 260:222-226 (1993)) of suitable

chimeric constructs of FZD and Wnt polypeptides in cells, or by variants of the two-hybrid assay (Fearon et al, Proc Natl Acad Sci USA 89:7958-7962 (1992); Takacs et al, Proc Natl Acad Sci USA 90:10375-10379 (1993); Vidal et al, Proc Natl Acad Sci USA 93:10321-10326 (1996)) employing suitable constructs of FZD and Wnt polypeptides and tailored for a high throughput assay to detect compounds that inhibit the FZD/Wnt interaction. These embodiments have the advantage that the cell permeability of the test compounds is assured.

For example, in one assay, a FZD or Wnt polypeptide or fragment thereof is adsorbed to ELISA plates. The FZD or Wnt polypeptides are then exposed to test compounds, followed by a glutathione-S-transferase (GST)-binding partner fusion protein, e.g., a GST-FZD or -Wnt polypeptide fusion protein. Bound protein is detected with goat anti-GST antibody, alkaline phosphatase (AP)-coupled anti-goat IgG, and AP substrate. Compounds that interfere with protein-protein interactions yield reduced AP signals in the ELISA plates.

In still another aspect, the invention provides methods of identifying test compounds that modulate (e.g., increase or decrease) expression of a FZD polypeptide. The method includes contacting a FZD nucleic acid with a test compound and then measuring expression of the encoded FZD polypeptide. In a related aspect, the invention features a method of identifying compounds that modulate (e.g., increase or decrease) the expression of FZD polypeptides by measuring expression of a FZD polypeptide in the presence of the test compound or after the addition of the test compound in: (a) a cell line into which has been incorporated a recombinant construct including the FZD nucleic acid sequence or fragment or an allelic variation thereof; or (b) a cell population or cell line that naturally selectively expresses FZD, and then measuring the expression of the FZD protein.

Since the FZD nucleic acids described herein have been identified, they can be cloned into various host cells (e.g., mammalian cells, insect cells, bacteria or fungi) for carrying out such assays in whole cells.

In certain embodiments, an isolated nucleic acid molecule encoding a FZD polypeptide is used to identify a compound that modulates (e.g., increases or decreases) the expression of FZD *in vivo* (e.g., in a FZD-producing cell). In such embodiments, cells that express a FZD (e.g., FZD 7 and/or 8) are cultured, exposed to a test compound (or a mixture of test compounds), and the level of FZD expression is

compared with the level of FZD expression or activity in cells that are otherwise identical but that have not been exposed to the test compound(s). Standard quantitative assays of gene expression can be used.

5 Expression of FZD can be measured using art-known methods, for example, by Northern blot PCR analysis or RNase protection analyses using a nucleic acid molecule of the invention as a probe. Other examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). The level of expression in the presence of the test molecule, compared with the level of expression in its absence, will indicate whether or not the test
10 compound modulates the expression of a FZD polypeptide.

In still another aspect, the invention provides methods of screening test compounds utilizing cell systems that are sensitive to perturbation of one or several transcriptional/translational components.

15 In certain embodiments, the methods include identifying candidate compounds that interfere with steps in FZD translational accuracy, such as maintaining a proper reading frame during translation and terminating translation at a stop codon. This method involves constructing cells in which a detectable reporter polypeptide can only be produced if the normal process of staying in one reading frame or of terminating translation at a stop codon has been disrupted. This method further involves contacting
20 the cell with a test compound to examine whether it increases or decreases the production of the reporter polypeptide.

In other embodiments, the cell system is a cell-free extract and the method involves measuring transcription or translation *in vitro*. Conditions are selected so that transcription or translation of the reporter is increased or decreased by the addition of a
25 transcription modifier or a translation modifier to the cell extract.

One method for identifying candidate compounds relies upon a transcription-responsive gene product. This method involves constructing a cell in which the production of a reporter molecule changes (i.e., increases or decreases) under conditions in which cell transcription of a FZD nucleic acid changes (i.e., increases or
30 decreases). Specifically, the reporter molecule is encoded by a nucleic acid transcriptionally linked to a sequence constructed and arranged to cause a relative change in the production of the reporter molecule when transcription of a FZD nucleic acid changes. A gene sequence encoding the reporter may, for example, be fused to

part or all of the gene encoding the transcription-responsive gene product and/or to part or all of the genetic elements that control the production of the gene product.

Alternatively, the transcription-responsive gene product may stimulate transcription of the gene encoding the reporter, either directly or indirectly. The method further
5 involves contacting the cell with a test compound, and determining whether the test compound increases or decreases the production of the reporter molecule in the cell.

Alternatively, the method for identifying candidate compounds can rely upon a translation-responsive gene product. This method involves constructing a cell in which cell translation of a FZD nucleic acid changes (i.e., increases or decreases).

Specifically, the reporter molecule is encoded by nucleic acid translationally linked to a
10 sequence constructed and arranged to cause a relative increase or decrease in the production of the reporter molecule when transcription of a FZD nucleic acid changes.

A gene sequence encoding the reporter may, for example, be fused to part or all of the gene encoding the translation-responsive gene product and/or to part or all of the

15 genetic elements that control the production of the gene product. Alternatively, the translation-responsive gene product may stimulate translation of the gene encoding the reporter, either directly or indirectly. The method further involves contacting the cell with a test compound, and determining whether the test compound increases or decreases the production of the first reporter molecule in the cell.

20 For these and any method described herein, a wide variety of reporters may be used, with typical reporters providing conveniently detectable signals (e.g., by spectroscopy). By way of example, a reporter gene may encode an enzyme that catalyses a reaction that alters light absorption properties.

Examples of reporter molecules include but are not limited to β -galactosidase,
25 invertase, green fluorescent protein, luciferase, chloramphenicol acetyltransferase, beta-glucuronidase, exo-glucanase, glucoamylase and radiolabeled reporters. For example, the production of the reporter molecule can be measured by the enzymatic activity of the reporter gene product, such as β -galactosidase.

Any of the methods described herein can be used for high throughput screening
30 of numerous test compounds to identify candidate anti-cancer agents. By high-throughput screening is meant that the method can be used to screen a large number of candidate compounds relatively easily and quickly.

Having identified a test compound as a candidate anti-cancer agent, the compound can be further tested *in vivo* or *in vitro* using techniques known in the art to confirm whether it is an anti-cancer agent, i.e., to determine whether it can modulate Wnt/FZD signaling, cancer cell motility and/or FZD expression *in vitro* (e.g., using isolated cells or cell-free systems) or *in vivo* (e.g., using an animal, e.g., rodent, model system) if desired.

In vitro testing of a candidate compound can be accomplished by means known to those in the art, such as assays involving the use of cells, e.g., wild type, cancerous and/or transgenic liver cells. Exemplary assays for monitoring Wnt/FZD signaling, FZD expression and cancer cell motility, as well as useful cells that can be used in such assays, are described in the Examples section, below.

Alternatively or in addition, *in vivo* testing of candidate compounds can be performed by means known to those in the art. For example, the candidate compound(s) can be administered to a mammal, such as a rodent (e.g., mouse) or rabbit. Such animal model systems are art-accepted for testing potential pharmaceutical agents to determine their therapeutic efficacy in patients, e.g., human patients. Animals that are particularly useful for *in vivo* testing are wild type animals or non-wild type animals (e.g., mice) that over-produce FZD polypeptides, e.g., animals that overexpress a FZD transgene (e.g., a FZD 7 transgene) and/or that display reduced production of FZD 8 polypeptides. Other animals that are useful for *in vivo* testing are animals bred to develop liver cancer. Certain particularly useful transgenic mice that develop liver cancer are described in the Examples section and are included in the present invention.

In a typical *in vivo* assay, an animal (e.g., a wild type or transgenic mouse) is administered, by any route deemed appropriate (e.g., by injection), a dose of a candidate compound. Conventional methods and criteria can then be used to monitor animals for the desired activity. If needed, the results obtained in the presence of the candidate compound can be compared with results in control animals that are not treated with the test compound.

Medicinal Chemistry

Once a compound (or agent) of interest has been identified, standard principles of medicinal chemistry can be used to produce derivatives of the compound for further

rounds of testing. Derivatives can be screened for improved pharmacological properties, for example, efficacy, pharmaco-kinetics, stability, solubility, and clearance. The moieties responsible for a compound's activity in the assays described above can be delineated by examination of structure-activity relationships (SAR) as is commonly practiced in the art. A person of ordinary skill in pharmaceutical chemistry could modify moieties on a candidate compound or agent and measure the effects of the modification on the efficacy of the compound or agent to thereby produce derivatives with increased potency. For an example, see Nagarajan *et al.* (1988) *J. Antibiot.* 41: 1430-8. Furthermore, if the biochemical target of the compound (or agent) is known or determined, the structure of the target and the compound can inform the design and optimization of derivatives. Molecular modeling software is commercially available (e.g., Molecular Simulations, Inc.) for this purpose.

IV. Antibodies

The invention features purified or isolated antibodies that bind, e.g., specifically bind, to a FZD and/or Wnt polypeptide, i.e., anti-FZD and anti-Wnt antibodies. An antibody "specifically binds" to a particular antigen, e.g., a FZD 7 and/or 8 polypeptide, when it binds to that antigen, but recognizes and binds to a lesser extent (e.g., does not recognize and bind) to other molecules in a sample. Antibodies of the invention include monoclonal antibodies, polyclonal antibodies, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, and molecules produced using a Fab expression library.

An example of a type of antibody included in the present invention is the polyclonal anti-FZD 7 antibody described in the Examples section, below. Methods for producing polyclonal antibodies are well known to those of skill in the art.

As used herein, the term "antibody" refers to a protein comprising at least one, e.g., two, heavy (H) chain variable regions (abbreviated herein as VH), and at least one, e.g., two light (L) chain variable regions (abbreviated herein as VL). The VH and VL regions can be further subdivided into regions of hypervariability, termed "complementarity determining regions" ("CDR"), interspersed with regions that are more conserved, termed "framework regions" (FR). The extent of the framework region and CDR's has been precisely defined (see, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and

Human Services, NIH Publication No. 91-3242, and Chothia, C. et al. (1987) *J. Mol. Biol.* 196:901-917). Each VH and VL is composed of three CDR's and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

5 An anti-FZD or -Wnt antibody can further include a heavy and light chain constant region, to thereby form a heavy and light immunoglobulin chain, respectively. The antibody can be a tetramer of two heavy immunoglobulin chains and two light immunoglobulin chains, wherein the heavy and light immunoglobulin chains are inter-connected by, e.g., disulfide bonds. The heavy chain constant region is comprised of
10 three domains, CH1, CH2, and CH3. The light chain constant region is comprised of one domain, CL. The variable region of the heavy and light chains contains a binding domain that interacts with an antigen. The constant regions of the antibodies typically mediate the binding of the antibody to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (Clq) of the classical
15 complement system.

A "FZD binding fragment" and "Wnt binding fragment" of an antibody refers to one or more fragments of a full-length antibody that retain the ability to specifically bind to FZD or Wnt polypeptides, respectively, or to portions thereof. Examples of polypeptide binding fragments of an antibody include, but are not limited to: (i) a Fab
20 fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546), which consists
25 of a VH domain; and (vi) an isolated complementarity determining region (CDR).

Furthermore, although the two domains of the Fv fragment, VL and VH, are encoded by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also encompassed within the terms
30 "FZD binding fragment" and "Wnt binding fragment" of an antibody. These antibody

fragments can be obtained using conventional techniques known to those with skill in the art.

To produce antibodies, polypeptides (or antigenic fragments (e.g., fragments of a polypeptide that appear likely to be antigenic by criteria such as high frequency of charged residues) or analogs of such polypeptides), e.g., those produced by recombinant or peptide synthetic techniques (see, e.g., *Solid Phase Peptide Synthesis*, supra; Ausubel et al., supra), can be used. In general, the polypeptides can be coupled to a carrier protein, such as KLH, as described in Ausubel et al., supra, mixed with an adjuvant, and injected into a host mammal. A "carrier" is a substance that confers stability on, and/or aids or enhances the transport or immunogenicity of, an associated molecule. For example, FZD or Wnt proteins, or fragments thereof, can be generated using standard techniques of PCR, and can be cloned into a pGEX expression vector (Ausubel et al., supra). Fusion proteins can be expressed in *E. coli* and purified using a glutathione agarose affinity matrix as described in Ausubel et al., supra.

Typically, various host animals are injected with FZD and/or Wnt polypeptides. Examples of suitable host animals include rabbits, mice, guinea pigs, and rats. Various adjuvants can be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete adjuvant), adjuvant mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Such procedures result in the production of polyclonal antibodies, i.e., heterogeneous populations of antibody molecules derived from the sera of the immunized animals. Antibodies can be purified from blood obtained from the host animal, for example, by affinity chromatography methods in which FZD and/or Wnt polypeptide antigens are immobilized on a resin.

The present invention also includes anti-FZD and anti-Wnt monoclonal antibodies. Monoclonal antibodies (mAbs), which are homogeneous populations of antibodies specific for a particular antigen, can be prepared using FZD or Wnt polypeptides and standard hybridoma technology (see, e.g., Kohler et al., *Nature*, 256:495, 1975; Kohler et al., *Eur. J. Immunol.*, 6:511, 1976; Kohler et al., *Eur. J. Immunol.*, 6:292, 1976; Hammerling et al., In Monoclonal Antibodies and T Cell Hybridomas, Elsevier, NY, 1981; Ausubel et al., supra).

Typically, monoclonal antibodies are produced using any technique that provides for the production of antibody molecules by continuous cell lines in culture, such as those described in Kohler et al., *Nature*, 256:495, 1975, and U.S. Patent No. 4,376,110; the human B-cell hybridoma technique (Kosbor et al., *Immunology Today*, 4:72, 1983; Cole et al., *Proc. Natl. Acad. Sci. USA*, 80:2026, 1983); and the EBV-hybridoma technique (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96, 1983). Such antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD, and any subclass thereof. The hybridomas producing the mAbs of this invention can be cultivated *in vitro* or *in vivo*.

Once produced, polyclonal or monoclonal antibodies can be tested for recognition, e.g., specific recognition, of FZD or Wnt polypeptides in an immunoassay, such as a Western blot or immunoprecipitation analysis using standard techniques, e.g., as described in Ausubel et al., *supra*. Antibodies that specifically bind to FZD or Wnt polypeptides (e.g., FZD 7, FZD 8, Wnt 3, Wnt 8b and/or Wnt 11) are useful in the invention. For example, such antibodies can be used in an immunoassay to detect the polypeptide in a sample, e.g., a tissue sample.

Alternatively or in addition, a monoclonal antibody can be produced recombinantly, e.g., produced by phage display or by combinatorial methods as described in, e.g., Ladner et al. U.S. Patent No. 5,223,409; Kang et al. International Publication No. WO 92/18619; Dower et al. International Publication No. WO 91/17271; Winter et al. International Publication WO 92/20791; Markland et al. International Publication No. WO 92/15679; Breitling et al. International Publication WO 93/01288; McCafferty et al. International Publication No. WO 92/01047; Garrard et al. International Publication No. WO 92/09690; Ladner et al. International Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum Antibod Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J* 12:725-734; Hawkins et al. (1992) *J Mol Biol* 226:889-896; Clackson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrard et al. (1991) *Bio/Technology* 9:1373-1377; Hoogenboom et al. (1991) *Nuc Acid Res* 19:4133-4137; and Barbas et al. (1991) *PNAS* 88:7978-7982.

Anti-FZD and -Wnt antibodies can be fully human antibodies (e.g., an antibody made in a mouse which has been genetically engineered to produce an antibody from a human immunoglobulin sequence), or non-human antibodies, e.g., rodent (mouse or

rat), rabbit, horse, cow, goat, primate (e.g., monkey), camel, donkey, pig, or bird antibodies.

An anti-FZD and anti-Wnt antibody can be one in which the variable region, or a portion thereof, e.g., the CDRs, is generated in a non-human organism, e.g., a rat or mouse. The anti-FZD and anti-Wnt antibody can also be, for example, chimeric, CDR-grafted, or humanized antibodies. The anti-FZD and anti-Wnt antibody can also be generated in a non-human organism, e.g., a rat or mouse, and then modified, e.g., in the variable framework or constant region, to decrease antigenicity in a human.

Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci.*, 81:6851, 1984; Neuberger et al., *Nature*, 312:604, 1984; Takeda et al., *Nature*, 314:452, 1984) can be used to splice the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; and U.S. Patents 4,946,778 and 4,704,692) can be adapted to produce single chain antibodies specific for a FZD or Wnt polypeptide. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments that recognize and bind to specific epitopes can be generated by known techniques. For example, such fragments can include but are not limited to F(ab')₂ fragments, which can be produced by pepsin digestion of the antibody molecule, and Fab fragments, which can be generated by reducing the disulfide bridges of F(ab')₂ fragments. Alternatively, Fab expression libraries can be constructed (Huse et al., *Science*, 246:1275, 1989) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Polyclonal and monoclonal antibodies (or fragments thereof) that specifically bind to a FZD and/or Wnt polypeptides can be used, for example, to detect expression of FZD and/or Wnt in various tissues of a patient. For example, a FZD 7 and/or 8 polypeptide can be detected in conventional immunoassays of biological tissues or

extracts. Examples of suitable assays include, without limitation, Western blotting, ELISAs, radioimmunoassays, and the like.

V. Pharmaceutical Compositions

5 Any pharmaceutically active compound, agent, nucleic acid, polypeptide, or antibody (all of which can be referred to herein as “active compounds”), can be incorporated into pharmaceutical compositions. Such compositions typically include the active compound and a pharmaceutically acceptable carrier. A “pharmaceutically acceptable carrier” can include solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible
10 with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition can be formulated to be compatible with its intended route of administration. Examples of routes of administration include enteral (e.g., oral or rectal) and parenteral, e.g., intravenous (e.g., into the portal vein of the
15 liver), intradermal, subcutaneous, transdermal, transmucosal, and pulmonary administration. Administration may be directly into the liver, e.g., by injection or by topical administration during surgery. Solutions or suspensions used for injection can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic
20 solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The
25 parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For
30 intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy

syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol or sorbitol and sodium chloride. Prolonged absorption of the injectable compositions can be achieved by including an agent which delays absorption, e.g., aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying, which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, e.g., gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon

dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressured container or dispenser that contains a suitable
5 propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and
10 fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides). For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

15 In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid.
20 Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

25 It may be advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required
30 pharmaceutical carrier.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the

dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio LD50/ED50. Compounds which exhibit high therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue, e.g., liver, in order to minimize potential damage to healthy cells and, thereby, reduce side effects.

The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The terms "effective amount" and "effective to treat," as used herein, refer to an amount or concentration of a compound described herein utilized for a period of time (including acute or chronic administration and periodic or continuous administration) that is effective within the context of its administration for causing an intended effect or physiological outcome. For compounds described herein, an effective amount, e.g., of a polypeptide (i.e., an effective dosage), ranges from about 0.001 to 500 mg/kg body weight, e.g. about 0.01 to 50 mg/kg body weight, e.g. about 0.1 to 20 mg/kg body weight. The polypeptide can be administered one time per week for between about 1 to 10 weeks, e.g. between 2 to 8 weeks, about 3 to 7 weeks, or for about 4, 5, or 6 weeks. The skilled artisan will appreciate that certain factors influence the dosage and timing required to effectively treat a patient, including but not limited to the type of patient to be treated, the severity of the disease or disorder, previous treatments, the general health and/or age of the patient, and other diseases present. Moreover, treatment of a patient with a therapeutically effective amount of a compound can include a single treatment or, preferably, can include a series of treatments.

With respect to antibodies, partially human antibodies and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration are possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake and tissue penetration. A method for lipidation of antibodies is described by Cruikshank *et al.* ((1997) *J. Acquired Immune Deficiency Syndromes and Human Retrovirology* 14:193).

If the compound is a small molecule, exemplary doses include milligram or microgram amounts of the small molecule per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is furthermore understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. When one or more of these small molecules is to be administered to an animal (e.g., a human) to modulate expression or activity of a FZD or Wnt polypeptide or nucleic acid, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

Nucleic acid molecules (e.g., FZD, e.g., FZD 7, DNA) can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see, e.g., U.S. Patent 5,328,470) or by stereotactic injection (see, e.g., Chen *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system. Exemplary constructs that can

potentially be used in gene therapy methods are described in the Examples section, below.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

5

VI. Cancer and Treatments Therefor

The term "cancer" refers to animal cells having the capacity for autonomous growth. Examples of such cells include cells having an abnormal state or condition characterized by rapidly proliferating cell growth. The term is meant to include cancerous growths, e.g., tumors; oncogenic processes, metastatic tissues, and malignantly transformed cells, tissues, or organs, irrespective of histopathologic type or stage of invasiveness. Also included are malignancies of the various organ systems, such as respiratory, cardiovascular, renal, reproductive, hematological, neurological, hepatic, gastrointestinal, and endocrine systems; as well as adenocarcinomas which include malignancies such as most colon cancers, renal-cell carcinoma, prostate cancer and/or testicular tumors, non-small cell carcinoma of the lung, cancer of the small intestine, and cancer of the esophagus. Cancer that is "naturally arising" includes any cancer that is not experimentally induced by implantation of cancer cells into a subject, and includes, for example, spontaneously arising cancer, cancer caused by exposure of a patient to a carcinogen(s), cancer resulting from insertion of a transgenic oncogene or knockout of a tumor suppressor gene, and cancer caused by infections, e.g., viral infections. The term "carcinoma" is art recognized and refers to malignancies of epithelial or endocrine tissues. The term includes carcinosarcomas, which include malignant tumors composed of carcinomatous and sarcomatous tissues. An "adenocarcinoma" refers to a carcinoma derived from glandular tissue or in which the tumor cells form recognizable glandular structures. The term "hepatocellular carcinoma" (HCC) refers to cancer that arises from hepatocytes, the major cell type of the liver.

The term "patient" is used throughout the specification to describe an animal, human or non-human, rodent or non-rodent, to whom treatment according to the methods of the present invention is provided. Veterinary and human clinical applications are contemplated. The term "patient" includes, but is not limited to, birds, reptiles, amphibians, and mammals, e.g., humans, other primates, pigs, rodents such as

30

mice and rats, rabbits, guinea pigs, hamsters, cows, horses, cats, dogs, sheep and goats. Preferred subjects are humans, farm animals, and domestic pets such as cats and dogs. The term "treat(ment)," is used herein to denote delaying the onset of, inhibiting, alleviating the effects of, or prolonging the life of a patient suffering from, a condition, e.g., cancer.

Cancers that may be treated using the methods and compositions of the present invention include, but are not limited to, cancers of the liver, stomach, colon, rectum, mouth/pharynx, esophagus, larynx, pancreas, lung, small bowel, and bile ducts, among others.

Individuals considered at risk for developing cancer may benefit particularly from the invention, primarily because prophylactic treatment can begin before there is any evidence of a tumor. Individuals "at risk" include, e.g., individuals exposed to carcinogens, e.g., by consumption, e.g., by inhalation and/or ingestion, at levels that have been shown statistically to promote cancer in susceptible individuals. Also included are individuals exposed to a virus, e.g., a hepatitis virus, e.g., hepatitis B virus (HBV). Also included are individuals at risk due to exposure to ultraviolet radiation, or their environment, occupation, and/or heredity, as well as those who show signs of a precancerous condition. Similarly, individuals in very early stages of cancer or development of metastases (i.e., only one or a few aberrant cells are present in the individual's body or at a particular site in an individual's tissue) may benefit from such prophylactic treatment.

Skilled practitioners will appreciate that a patient can be diagnosed by a physician (or veterinarian, as appropriate for the patient being diagnosed) as suffering from or at risk for cancer using the methods described herein, optionally using additional methods, e.g., assessing a patient's medical history, performing other diagnostic tests and/or by employing imaging techniques.

One strategy for treating patients suffering from or at risk for cancer is to modulate Wnt/FZD signaling in the patient. The goal is to increase signaling where signaling is too low and to decrease signaling where signaling is too high. Modulation of Wnt/FZD signaling falls into two basic categories: decreasing (i.e., reducing, e.g., eliminating) Wnt/FZD signaling and increasing (i.e., supplementing or providing) Wnt/FZD signaling where there is insufficient or no activity. Whether Wnt/FZD signaling should be inhibited or increased depends upon the intended application.

Wnt/FZD signaling can be modulated using the active compounds (e.g., candidate compounds and/or anti-cancer agents) described herein. Compounds that decrease Wnt/FZD signaling activity, e.g., by decreasing expression of FZD 7 and/or interfering with an interaction between FZD 7 and its ligand (e.g., Wnt 3, 8b and/or 11) can be used, e.g., as treatments for cancer, e.g., liver cancer. Compounds that increase activity, e.g., by increasing expression of FZD 8 can also be used, e.g., as treatments for cancer, e.g., liver cancer.

Decreasing Wnt/FZD Signaling

Art-known methods for decreasing the expression of a particular protein in a patient can be used to decrease Wnt/FZD signaling. For example, an antisense nucleic acid effective to inhibit expression of an endogenous FZD gene, e.g., FZD 7 gene, can be utilized. As used herein, the term “antisense oligonucleotide” or “antisense” describes an oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that mRNA.

Antisense molecules are designed so as to interfere with transcription or translation of a target gene (e.g., a gene encoding FZD 7 or Wnt 3, 8b or 11) upon hybridization with the target gene or transcript. The antisense nucleic acid can include a nucleotide sequence complementary to an entire FZD or Wnt RNA or only a portion of the RNA. On one hand, the antisense nucleic acid needs to be long enough to hybridize effectively with FZD or Wnt RNA. Therefore, the minimum length is approximately 12 to 25 nucleotides. On the other hand, as length increases beyond about 150 nucleotides, effectiveness at inhibiting translation may increase only marginally, while difficulty in introducing the antisense nucleic acid into target cells may increase significantly. Accordingly, an appropriate length for the antisense nucleic acid may be from about 15 to about 150 nucleotides, e.g., 20, 25, 30, 35, 40, 45, 50, 60, 70, or 80 nucleotides. The antisense nucleic acid can be complementary to a coding region of FZD or Wnt mRNA or a 5' or 3' non-coding region of a FZD mRNA, or both. One approach is to design the antisense nucleic acid to be complementary to a region on both sides of the translation start site of the FZD or Wnt mRNA.

Based upon the sequences disclosed herein, one of skill in the art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. For example, a "gene walk" comprising a series of oligonucleotides of 15-30 nucleotides complementary to and spanning the length of a FZD mRNA can be prepared, followed by testing for inhibition of FZD or Wnt expression. Optionally, gaps of 5-10 nucleotides can be left between the oligonucleotides to reduce the number of oligonucleotides synthesized and tested.

The antisense nucleic acid can be chemically synthesized, e.g., using a commercial nucleic acid synthesizer according to the vendor's instructions.

Alternatively, the antisense nucleic acids can be produced using recombinant DNA techniques. An antisense nucleic acid can incorporate only naturally occurring nucleotides. Alternatively, it can incorporate variously modified nucleotides or nucleotide analogs to increase its *in vivo* half-life or to increase the stability of the duplex formed between the antisense molecule and its target RNA. Examples of nucleotide analogs include phosphorothioate derivatives and acridine-substituted nucleotides. Given the description of the targets and sequences, the design and production of suitable antisense molecules is within ordinary skill in the art. For guidance concerning antisense nucleic acids, see, e.g., Goodchild, "Inhibition of Gene Expression by Oligonucleotides," in *Topics in Molecular and Structural Biology, Vol. 12: Oligodeoxynucleotides* (Cohen, ed.), MacMillan Press, London, pp. 53-77 (1989).

Delivery of antisense oligonucleotides can be accomplished by any method known to those of skill in the art. For example, delivery of antisense oligonucleotides for cell culture and/or *ex vivo* work can be performed by standard methods such as the liposome method or simply by addition of membrane-permeable oligonucleotides.

Delivery of antisense oligonucleotides for *in vivo* applications can be accomplished, for example, via local injection of the antisense oligonucleotides at a selected site, e.g., a liver. This method has previously been demonstrated for psoriasis growth inhibition and for cytomegalovirus inhibition. See, for example, Wraight et al., (2001). *Pharmacol Ther.* 90(1):89-104; Anderson et al., (1996) *Antimicrob Agents Chemother* 40: 2004-2011; and Crooke et al., (1996) *J Pharmacol Exp Ther* 277: 923-937.

Similarly, RNA interference (RNAi) techniques can be used to inhibit FZD or Wnt expression, in addition or as an alternative to the use of antisense techniques. For

example, small interfering RNA (siRNA) duplexes directed against FZD or Wnt nucleic acids could be synthesized and used to prevent expression of the encoded protein(s).

Another approach to inhibiting Wnt/FZD signaling involves administering to a patient a candidate compound or anti-cancer agent that binds to FZD polypeptides (e.g., FZD 7 polypeptides) and/or their binding partners (e.g., Wnt 3, 8b and/or 11), thereby preventing interaction between the two. Such compounds and agents may, for example, bind to the FZD polypeptide (e.g., to the CRD domain of the FZD polypeptide) and/or to the Wnt polypeptide (e.g., to a binding domain of the Wnt polypeptide) in such a way that interaction between the proteins is prevented. Such candidate compounds and anti-cancer agents can be identified using screening methods described herein. Examples of a compound that can bind to a Wnt polypeptide, e.g., Wnt 3, 8b and/or 11, is a FZD 7 receptor or truncated form thereof, as described in the Examples section, below and an anti-Wnt antibody.

Yet another approach to inhibiting Wnt/FZD signaling involves administering to a patient a vector (e.g., a gene therapy vector) that encodes a mutated (e.g., truncated) form of a FZD receptor, e.g., a FZD 7 receptor. Expression of the mutated form of the receptor by the patient's cells that incorporate the construct can interfere with Wnt/FZD signaling in the cells. For example, a construct that encodes a secreted and soluble form of a FZD receptor (e.g., a FZD 7 receptor) can be used. Expression of such a construct by target cells would cause the cells to secrete a soluble form of the FZD receptor that would bind Wnt polypeptides, rendering them unable to bind to intact FZD receptors on the cell surface. Alternatively or in addition, a construct that encodes a membrane bound but inactive form of a FZD receptor (i.e., a mutant FZD receptor unable to perform some function performed by a counterpart wild-type FZD receptor) can be used. Expression of such a construct by target cells may bind up Wnt polypeptides or interfere with Wnt/FZD signaling via an internal mechanism not involving Wnt polypeptides. The vector can be derived from a non-replicating linear or circular DNA or RNA vector, or from an autonomously replicating plasmid or viral vector. Methods for constructing suitable expression vectors are known in the art, and useful materials are commercially available. Exemplary expression vectors that encode useful mutated FZD 7 polypeptides are described in the Examples section, below.

Increasing Wnt/FZD Signaling

New or supplemental Wnt/FZD signaling can be provided *in vivo* by increasing expression of FZD polypeptides (e.g., FZD 8 polypeptides) in the patient. For example, a FZD polypeptide can be generated directly within an organism, e.g., a human, by
5 expressing within the cells of the organism a nucleic acid construct containing a nucleotide sequence encoding a FZD polypeptide (e.g., a FZD 8 polypeptide). Any appropriate expression vector suitable for transfecting the cells of the organism of interest can be used for such purposes.

10 VII. Transgenic Animals

The present invention also features transgenic animals that develop liver cancer and overexpress FZD 7 in their liver cells. Such animals represent model systems for the study of liver cancer and for the development of therapeutic agents that can modulate Wnt/FZD signaling and treat cancer.

15 Transgenic animals can be, for example, farm animals (pigs, goats, sheep, cows, horses, rabbits, and the like), rodents (such as rats, guinea pigs, and mice), non-human primates (for example, baboons, monkeys, and chimpanzees), and domestic animals (for example, dogs and cats).

Any technique known in the art can be used to introduce transgenes into
20 animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., *Proc. Natl. Acad. Sci., USA* 82:6148, 1985); gene targeting into embryonic stem cells (Thompson et al., *Cell* 56:313, 1989); and electroporation of embryos (Lo, *Mol. Cell. Biol.* 3:1803, 1983).

25 Especially useful are the methods described in Yang et al. (*Proc. Natl. Acad. Sci. USA* 94:3004-3009, 1997).

For a review of techniques that can be used to generate and assess transgenic animals, skilled artisans can consult Gordon (*Intl. Rev. Cytol.* 115:171-229, 1989), and may obtain additional guidance from, for example: Hogan et al. Manipulating the
30 Mouse Embryo, Cold Spring Harbor Press, Cold Spring Harbor, NY, 1986); Krimpenfort et al. (*Bio/Technology* 9:86, 1991), Palmiter et al. (*Cell* 41:343, 1985), Kraemer et al. (Genetic Manipulation of the Early Mammalian Embryo, Cold Spring Harbor Press, Cold Spring Harbor, NY, 1985), Hammer et al. (*Nature* 315:680, 1985),

Purcel et al. (*Science*, 244:1281, 1986), Wagner et al. (U.S. Patent No. 5,175,385), and Krimpenfort et al. (U.S. Patent No. 5,175,384).

EXAMPLES

5 The invention is illustrated in part by the following examples, which are not to be taken as limiting the invention in any way.

Example 1. Identification of natural Wnt ligands for Inhibition of HCC growth

10 The Wnt signaling pathway is involved in various differentiation events during embryonic development and leads to tumor formation when aberrantly activated. Molecular studies have pinpointed activating mutations of the Wnt signaling pathway as the cause of approximately 90% of colorectal cancer and somewhat less frequently in cancers at other sites, such as hepatocellular carcinoma (HCC). HCC is one of the most frequent fatal malignancies worldwide (Murray et al., 1997; Caselmann and Alt, 15 1996; Anthony, 2001; El-Serag and Mason, 1999).

 Aberrant activation of Wnt/ β -catenin pathway can be due to mutations of the β -catenin gene (de La Costa et al., 1998; Calvisi et al., 2001). However, additional studies estimate that 46% of hepatic carcinoma and between 35 to 80% of HCC have aberrant β -catenin cellular accumulation not associated with mutations. In addition, 20 Axin1 and APC gene mutations are rare in HCC and the mechanisms of the wild-type β -catenin accumulation in the majority of HCC have yet to be defined (Torbenon et al., 2002; Hsu et al., 2000; Devereux et al., 2001; Wong et al., 2001; Calvisi et al., 2004). Recent studies showed that other upstream components of Wnt signaling pathway stabilize β -catenin in the cytosol to allow nuclear translocation and up- 25 regulation of genes associated with the malignant phenotype (Merle et al., 2004). Moreover, FZD7, the receptor of Wnt/ β -catenin signal transduction pathway, was overexpressed in most human HCC tumors and associated with downstream activation of the Wnt/ β -catenin pathway in human HCC tumors (Merle et al., 2004). Although it is clear that up-regulation of FZD7 receptor involved HCC development, the molecular mechanisms that promote up-regulation of FZD7 gene remain unknown. 30

 Numerous studies have demonstrated that proteoglycans can regulate growth factor activity *in vitro*. Heparan sulfate proteoglycans (HSPGs) are a major component

of the extracellular matrix that regulate transmission of developmental signals and also are implicated in the pathophysiology of diseases, including cancer, in which signals and tissue interactions malfunction (Selva and Perrimon, 2001; Nybakken and Perrimon, 2002). Typically, the ligand binds to the glycosaminoglycan chains on cell surface proteoglycans, influences their activities in FGF (Rapraeger et al., 1994; Lundin et al., 2000; Lin et al., 1999) and Wnt signaling (Lin and Perrimon, 1999). One HSPG, Glypican1, is required for Wnt signaling through Wnt-Heparan sulfate (HS) binding. The modulation of this binding affinity can promote the HS-mediated presentation of Wnt ligand to its Frizzled receptor to initiate Wnt signal transduction (Ai, X et al., 2003). Since biochemical and genetic studies suggest that Wnt ligand activity is regulated extracellularly, there are two general ways in which this process can be controlled. One is to modulate the concentration gradient of the ligand, the other is to adjust the affinity of receptor-ligand binding (Cumberledge and Reichsman, 1997; Lin et al., 1999).

Molecular biological methods such as RT-PCR are widely used to search for specific gene expression in tissues or cells. However, these methods detect mRNA expression levels, not proteins. It is difficult to identify Wnt ligands as natural released proteins for at least the following reasons: 1) the amount of protein is too small to detect using typical methods (e.g., ELISA, Western blot); 2) there is currently no available antibody to detect human Wnt ligands proteins in tissues; and 3) Wnt ligand proteins are often associated with extracellular matrix and cell surfaces and are known to be highly hydrophobic.

The present study employed two different approaches to identify Wnt ligands. First, proteomic tools were used to identify Wnt ligands as secreted proteins. It is known that wingless (WG) and Wnt proteins are associated with cell surface and extracellular matrix as secretory glycoproteins. Indeed, in S2 cells, the majority of secreted WG is bound to the cell surface and extracellular matrix through specific, noncovalent interactions, and WG binds directly to heparin agarose beads with high affinity (Reichsman et al., 1996). Wnt1 is also associated tightly with the cell surface and extracellular matrix and can be released from the cell surface by treatment with exogenous heparin (Bradley and Brown, 1990). Moreover, Wnt1 induction is inhibited by heparin, which potentially binds Wnt1 and competes for its interaction with HSPGs

(Dhoot et al., 2001). Interestingly, proteoglycans are required for maintenance of Wnt11 expression in the ureter tips (Kispert et al, 1996).

Accordingly, Wnt ligand proteins were isolated from prepared cell surface heparan sulfate polypeptide glycan from hepatocellular carcinoma cell line (Huh7) following peptide mapping. mRNA expression of 19 known Wnt ligand genes was also analyzed using RT-PCR from HCC cell lines.

The natural ligands for activation of Wnt signaling cascade in HCC was unknown until the present study. Identification of specific Wnt ligand (s) in HCC provides molecular targets against HCC development. Recent studies have shown that blockage of certain Wnt ligands with a specific antibody can induce apoptosis and inhibit tumor growth (He et al., 2004; You et al., 2004).

Cell Culture

HepG2, Hep3B, Huh7, and Focus cell lines were cultured in Eagle's minimal essential medium (MEM) supplemented with 10% fetal bovine calf serum (FBS). The cells were harvested after washing with phosphated buffered saline twice, when the confluency reached about 70%.

Preparation of cell surface heparan sulfate proteoglycan (HSPG) and fractionation by Heparin-agarose chromatography

Huh7 cells were grown until 40% confluent in 10% FBS MEM, then the media was changed to 0% FBS MEM. 24 hours after incubation, cells were treated with or without heparin (50 $\mu\text{g}/\text{ml}$) in 0.1% FBS MEM for 12 hours. Cells were washed 5 times with ice-cold PBS and cell layers were incubated with crystallized trypsin (20 $\mu\text{g}/\text{ml}$) in TBS/EDTA for 10 minutes at on ice. Trypsin activity was stopped by the addition of soybean trypsin inhibitor at a final concentration of 100 $\mu\text{g}/\text{ml}$. Contaminated cells were removed by centrifugation at 400 x g for 5 minutes at 4 °C from trypsin.

The trypsin was subjected to Heparin-agarose chromatography. In brief, trypsin was incubated with heparin (4%) agarose beads overnight at 4°C. Beads were collected by centrifugation at 2000 x g for 4 minutes and washed with 20 volumes of 0.1 M NaCl in PBS. Eluted fractions were collected with 0.25, 0.5, 0.75, and 1.0 M NaCl in PBS.

Two-dimensional gel electrophoresis, In-gel digestion and Peptide Mapping

Fractionated samples from heparin-agarose chromatography were prepared by precipitation and rehydration with IPG buffer for isoelectric focusing (IEF). IEF was carried out according to the manufacturer's protocol (ZOOM IPGRunner, Invitrogen). ZOOM strips (pH 3-10) were rehydrated with samples overnight, then a step voltage ramping method was applied as follows: 200 Volts (V) for 20 minutes, 450 V for 15 minutes, 750 V for 15 minutes, and 2000 V for 120 minutes. Focused gels were performed using SDS-PAGE using ZOOM gels (Invitrogen) as second dimensional electrophoresis. Following electrophoresis, gels were stained using SilverQuest Silver Staining kit (Invitrogen).

Protein spots excised from silver-stained gels were destained and dried before enzymatic digestion with sequence grade modified trypsin (Promega, Madison, WI). Tryptic peptides were desalted and concentrated with ZipTip_{C18} (Millipore; Bedford, MA). The concentrated tryptic peptides were applied to SEND ProteinChip and performed peptide mapping using PBSII (CIPHERGEN, Fremont, CA). Peptide mass fingerprinting was conducted with the database search tool MS-fit in the program Protein Prospector, available on the internet at address (<http://prospector.ucsf.edu>). A number of restrictions were applied to the initial search based on localization of the spot in the 2-D gel: species = homo sapiens, pI range = 6-9.5, mass range = 35 – 50 kDa.

RT-PCR Analysis

Total cellular RNA was extracted by using TRIzol[®] Reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. Isolated RNA was then reverse transcribed by using First Strand cDNA Synthesis Kit for RT-PCR (AMV) (Roche Diagnostics Co., Indianapolis, IN, USA).

Polymerase chain reaction (PCR) was performed in a thermocycler (MJ Research Inc., Waltham, MA, USA) using 50 ng of cDNA and High Fidelity PCR Master (Roche Diagnostics Co.). The primer pairs for each Wnt ligand are listed in Table 1, below. The final concentration of each primer was 0.25 μ M. After initial denaturation at 94°C for 4 minutes, reactions were subjected to 35 cycles of the following thermal program: 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1

minute, followed by a final elongation step at 72°C for 10 minutes. The amplified products were analyzed on ethidium bromide-stained 2% agarose gel.

RESULTS

Because Wnt proteins are generally not secreted in a soluble form, an attempt was made to purify cell surface heparan sulfate proteoglycans (HSPGs) bound to Wnt protein(s) as an associated form in the hepatocellular carcinoma cell line, Huh7. Huh7-HSPG prepared by trypsin was subjected to Heparin-agarose affinity resin for pre-fractionation. Eluted fractions were applied to SDS-PAGE and compared the protein bands between heparin-treated and non-treated samples by SDS-PAGE. From 0.25 M NaCl eluted fraction, an approximately 45 kDa protein band was distinguishable at heparin-untreated fraction (Fig. 1). As the estimated molecular weight of Wnt proteins is in the range of 35–45 kDa, this protein band was suspected of containing Wnt proteins. To define this protein band, 2-D electrophoresis was performed. As shown in Fig. 2, several silver-stained protein spots showed the different protein level between heparin-treated and untreated samples. The range for protein spots was narrowed to 35–55 kDa and pI 5-9.5, as it is known that Wnt proteins are basic. Two up-regulated protein spots in heparin-untreated samples were of interest since Wnt proteins can be released by heparin treatment.

Excised protein spots from silver-stained 2-D gels were performed in-gel digestion following peptide mapping by mass spectrometry. Using a database search, one protein spot was determined to be human Wnt 11 by matching peptide fragments with calculated fragments (Fig. 3).

Wnt 3 mRNA could be detected in all of the 4 hepatocellular carcinoma cell lines. Wnt 11 mRNA was positive in HepG2, Hep3B, and Huh7 cell lines. Wnt8b mRNA was detected in the Hep3B cell line. mRNA for other Wnt ligands could not be detected.

REFERENCES

1. Ai, X., Do, A-T, Lozynska, O., Kusche-Gullberg, M., Lindahl, U., and Emerson, Jr. C.P. (2003). Qsulf1 remodels the 6-O sulfation states of cell surface
5 heparan proteoglycans to promote Wnt signaling. *J Cell Biol* 162, 341-351.
2. Anthony, P.P. (2001). Hepatocellular carcinoma: an overview. *Histopathology*
39, 109-118.
3. Bradley, R.S., and Brown, A.M. (1990). The proto-oncogene int-1 encodes a
secreted protein associated with the extracellular matrix. *EMBO J* 9 (5), 1569-1575.
- 10 4. Calvisi, D.F., Factor, V.M, Ladu, S., Conner, E.A., and Thorgeirsson, S.S.
(2004). Disruption of b-catenin pathway or genomic instability define two distinct
categories of liver cancer in transgenic mice. *Gastroenterology* 126, 1374-1386.
5. Calvisi, D.F., Factor, V.M, Loi, R., and Thorgeirsson, S.S. (2001). Activation of
 β -catenin during hepatocarcinogenesis in transgenic mouse models. *Cancer Res.* 61,
15 2085-2091.
6. Caselmann, W.H., and Alt, M. (1996). Hepatitis C virus infection as a major
risk factor for hepatocellular carcinoma. *J Hepatol* 24, 61-66.
7. de la Coste, A., Romagnolo, B., Billuart, P., Renard, C.A., Buendia, M.A.,
Soubrane, O., Fabre, M., Chelly, J., Beldjord, C., Kahn, A., and Perret, C. (1998).
20 Somatic mutations of the β -catenin gene are frequent in mouse and human
hepatocellular carcinoma. *Proc Natl Acad Sci USA* 95, 8847-8851.
8. Cumberledge, S., and Reichsman, F. (1997). Glycosaminoglycans and WNTs:
just a spoonful of sugar helps the signal go down. *TIG* 13 (11), 421-423.
9. Devereux, T.R., Stern, M.C., Flake, G.P., Yu, M.C., Zhang, Z.Q., London, S.J.,
25 and Tayler, J.A., (2001). CTNNB1 mutations and beta-catenin protein accumulation in
human hepatocellular carcinomas associated with high exposure to aflatoxin B1. *Mol*
Carcinog 31, 68-73.
10. Dhoot, G.K., Gustafsson, M.K., Ai, X., Sun, W., Standiford, D.M., and
Emerson Jr., C.P. (2001). Regulation of Wnt signaling and embryo patterning by an
30 extracellular sulfatase. *Science* 293, 1663-1666.
11. El-Serag, H.B., and Mason, A.C. (1999). Rising incidence of hepatocellular
carcinoma in the United States. *N Engl J Med* 340 (10), 745-750.

12. He, B., You, L., Uematsu, K., Xu, Z., Lee, A.Y., Matsangou, M., McCormick, F., and Jablons, D.M. (2004). A monoclonal antibody against Wnt-1 induces apoptosis in human cancer cells. *Neoplasia* 6 (1), 7-14
13. Hsu, H.C., Jeng, Y.M, Mao, T.L., Chu, J.S., Lai, P.L., and Peng, S.Y. (2000).
5 B-catenin mutations are associated with a subset of low-stage hepatocellular carcinoma negative for hepatitis B virus and with favorable prognosis. *Am J Pathol* 157, 763-770.
14. Kispert, A., Vainio, S., Shen, L., Rowitch, D.H., and McMahon, A.P. (1996).
Proteoglycans are required for maintenance of Wnt-11 expression in the ureter tips. *Development* 122 (11), 3627-3637.
- 10 15. Lin, X., Buff, E.M., Perrimon, N., and Michelson, A.M. (1999). Heparan sulfate proteoglycans are essential for FGF receptor signaling during *Drosophila* embryonic development. *Development* 126 (17), 3715-3723.
16. Lin, X., and Perrimon, N. (1999). Dally cooperates with *Drosophila* Frizzled 2 to transduce Wingless signaling. *Nature* 400, 281-284.
- 15 17. Lundin, L., Larsson, H., Kreuger, L., Kanda, S., Lindahl, U., Salmivirta, M., and Claesson-Welsh, L. (2000). Selectively desulfated heparin inhibits fibroblast growth factor-induced mitogenicity and angiogenesis. *J Biol Chem* 275 (32), 24653-24660.
18. Merle, P., de la Monte, S.M., Kim, M., Herrmann, M., Tanaka, S., von dem
20 Bussche, A., Kew, M., Trepo, C., and Wands, J.R. (2004). Functional consequences of frizzled-7 receptor overexpression in human hepatocellular carcinoma. *Gastroenterology* (in print).
19. Murray, C.J.L., and Lopez, A.D. (1997). Mortality by cause for eight regions of the world: global burden of disease study. *Lancet* 349, 1269-1276.
- 25 20. Nybakken, K., and Perrimon, N. (2002). Heparan sulfate proteoglycan modulation of developmental signaling in *Drosophila*. *Biochim Biophys Acta* 1573, 280-291.
21. Rapraeger, A.C., Guimond, S., Krufka, A., and Olwin, B.B. (1994). Regulation by heparan sulfate in fibroblast growth factor signaling. *Methods Enzymol* 245, 219-
30 240.
22. Reichsman F, Smith L, and Cumberland S. (1996). Glycosaminoglycans can modulate extracellular localization of the wingless protein and promote signal transduction. *J Cell Biol* 135 (3), 819-827.

23. Selva, E.M, and Perrimon, N. (2001). Role of heparan sulfate proteoglycans in cell signaling and cancer. *Adv Cancer Res* 83, 67-80.
24. Torbenson, M, Lee, J.H., Choti, M., Gag, W., Abraham, S.C., Montgomery, E., Boitnott, J., and Wu, T.T. (2002). Hepatic adenomas: analysis of sex steroid receptor status and the Wnt signaling pathway. *Mod Pathol* 15 (3), 189-196.
- 5 25. Wong, C.M., Fan, S.T., and Ng, I.O. (2001). β -catenin mutation and overexpression in hepatocellular carcinoma: clinicopathologic and prognostic significance. *Cancer* 92, 136-145.
26. You, L., He, B., Xu, Z., Uematsu, K., Mazieres, J., Fujii, N., Mikami, I.,
10 Reguart, N., McIntosh, J.K., Kashani-Sabet, M., McCormick, F., and Jablons, D.M. (2004). An anti-Wnt-2 monoclonal antibody induces apoptosis in malignant melanoma cells and inhibits tumor growth. *Cancer Res* 64 (15), 5385-5389.

Table 1. Primer Pairs Used for the Detection of Wnt Ligand mRNA

Wnt	Sequence	SEQ ID NO
GAPDH	Sense: 5'-GAAATCCCATCACCATCTTCCAG-3'	28
	Anti-sense: 5'-ATGAGTCCTTCCACGATACCAAAG-3'	29
Wnt1	Sense: 5'-TGTTGCCTGGCTGGGTTTC-3'	30
	Anti-sense: 5'-CTGTAAGCAGGTTTCGTGGAG-3'	31
Wnt2	Sense: 5'-GTGGATGCAAAGGAAAGGAA-3'	32
	Anti-sense: 5'-AGCCAGCATGTCTGAGAGT-3'	33
Wnt2b	Sense: 5'-ACCCAAGATGGTGCCAACTTC-3'	34
	Anti-sense: 5'-CACAACCGTCTGTTCTTTTGATG-3'	35
Wnt3	Sense: 5'-GGAGTGTATTCGCATCTACGACG-3'	36
	Anti-sense: 5'-CGAGTTGGGTCTGGGTCATTTAC-3'	37
Wnt3a	Sense: 5'-CCCCACTCGGATACTTCTTACTCC-3'	38
	Anti-sense: 5'-CTCCTGGATGCCAATCTTGATG-3'	39
Wnt4	Sense: 5'-TTTGTGGATGTGCGGGAGAG-3'	40
	Anti-sense: 5'-ATCTGTGTGCGGCTTGAAGT-3'	41
Wnt5a	Sense: 5'-ACACCTCTTTCCAAACAGGCC-3'	42
	Anti-sense: 5'-GGATTGTTAAACTCAACTCTC-3'	43
Wnt5b	Sense: 5'-GGAGCGAGAGAAGAAGTGGCC-3'	44
	Anti-sense: 5'-GAAGCAGCACCAGTGGAAGTGG-3'	45
Wnt6	Sense: 5'-CTTGGTTATGGACCCTACCCAGGCATC-3'	46
	Anti-sense: 5'-CACTGCAGCAGCTCGCCCATAGAA-3'	47
Wnt7a	Sense: 5'-GCTGCCTGGGCCACCTCTTTCTCA-3'	48
	Anti-sense: 5'-CCCGGTGGTACAGGCCTTGCTTCT-3'	49
Wnt7b	Sense: 5'-TCAACGAGTGCCAGTACCAG-3'	50
	Anti-sense: 5'-CCCTCGGCTTGGTTGTAGTA-3'	51
Wnt8a	Sense: 5'-TCCAGTTTGCTTGGGAACGC-3'	52
	Anti-sense: 5'-CCATCACAGCCACAGTTTTCG-3'	53
Wnt8b	Sense: 5'-CATCTGTCTTTTACCTGTGTCCTC-3'	54
	Anti-sense: 5'-AATGCTGTCTCCGATTGGC-3'	55
Wnt10a	Sense: 5'-TCTGGGTGCTCCTGTTCTTCCCTAC-3'	56
	Anti-sense: 5'-ATTGGTGTGGCATTCTGGTGG-3'	57
Wnt10b	Sense: 5'-ACTGTCCCGAGGCAAGAGTTTC-3'	58
	Anti-sense: 5'-GCATTTCCGCTTCAGGTTTTC-3'	59
Wnt11	Sense: 3'-TGCTGACCTCAAGACCCGATAC-3'	60
	Anti-sense: 3'-TGTCGCTTCCGTTGGATGTC-3'	61
Wnt14	Sense: 5'-TGCCAGTTCAGTTCGCTTTG-3'	62
	Anti-sense: 5'-TTCACACCCACGAGGTTGTTG-3'	63
Wnt15	Sense: 5'-TGAGTGCCAGTTTCAGTTCGG-3'	64
	Anti-sense: 5'-CTTGTTTCTCTTGGACCCC-3'	65
Wnt16	Sense: 5'-CTGCTCCGATGATGTCCAGTATG-3'	66
	Anti-sense: 5'-CATTCTCTGCCTTGTGTCCCTG-3'	67

A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A method for identifying an anti-cancer agent, the method comprising:
selecting a test compound that binds to a polypeptide comprising the amino acid sequence of a Wnt 3, Wnt 8b or Wnt 11 protein or a FZD-binding fragment thereof;
and
5 determining whether the test compound is capable of:
 - (i) reducing Wnt/FZD 7 signaling in a cell;
 - (ii) reducing liver cancer cell motility;
 - (iii) reducing β -catenin accumulation in a liver cancer cell; or
 - (iv) treating liver cancer in vitro or in vivo;10 wherein a test compound that is capable of at least one of (i) to (iv) is an anti-cancer agent.

2. The method of claim 1, wherein the selection of the test compound comprises:
contacting the test compound with the polypeptide comprising the amino acid
15 sequence of a Wnt 3, Wnt 8b or Wnt 11 protein or a FZD-binding fragment thereof;
detecting binding between the polypeptide and the test compound; and
selecting the test compound if it binds to the polypeptide.

3. The method of claim 1 or 2, wherein the polypeptide is
 - (i) a naturally occurring polypeptide;
 - (ii) a recombinant polypeptide;
 - (iii) provided as a polypeptide expressed on the surface of a cell; or
 - (iv) provided as an isolated polypeptide.

4. The method of claim 1 or 2, wherein the polypeptide comprises
 - (i) the amino acid sequence of a Wnt 3 protein;
 - (ii) the amino acid sequence of a Wnt 8b protein; or
 - (iii) the amino acid sequence of a Wnt 11 protein;

5. The method of claim 1 or 2, wherein the polypeptide comprises
 - (i) any one of SEQ ID NOs:8 to 12;

- (ii) any one of SEQ ID NOs: 15 to 19;
- (iii) any one of SEQ ID NO: 22 to 26; or
- (iv) any one of SEQ ID NO:7 to 27 and at least one non-Wnt sequence.

6. A compound identified by the method of any one of claims 1-5 for use in the treatment of liver cancer.

7. The compound of claim 6 for use in the treatment of liver cancer, wherein the compound is

- (i) an anti-Wnt antibody; or
- (ii) an FZD 7 receptor or Wnt-binding fragment thereof.

8. Use of a compound identified by the method of any one of claims 1-5 in the manufacture of a medicament for

- (i) treating liver cancer or
- (ii) reducing motility of liver cancer cells in a patient.

9. Use of a compound that binds Wnt 3, 8b or 11 and thereby reduces Wnt/ FZD 7 signaling in cells of a patient in the manufacture of a medicament for treating liver cancer or reducing the motility of liver cancer cells in a patient.

5

10. A method for identifying a candidate anti-cancer agent, the method comprising:

(a) providing a first polypeptide that:

- (i) comprises a FZD 7 polypeptide or a fragment thereof; and
- (ii) displays Wnt-binding ability;

10

(b) providing a second polypeptide that:

- (i) comprises a Wnt 3, 8b or 11 polypeptide or a fragment thereof; and
- (ii) displays FZD 7 binding ability;

(c) contacting the first and second polypeptides in the presence of a test compound; and

15

(d) comparing the level of binding between the first and second polypeptides in the presence of the test compound with the level of binding in the absence of the test

compound, wherein a reduced level of binding in the presence of the test compound than in its absence indicates that the test compound is a candidate anti-cancer agent.

11. The method of claim 10, further comprising:

(e) determining whether the candidate anti-cancer agent is capable of:

(i) reducing Wnt/FZD 7 signaling in a cell that expresses FZD 7;

(ii) reducing cancer cell motility;

(iii) reducing β -catenin accumulation in a cancer cell; or

(iv) treating cancer *in vitro* or *in vivo*;

5 wherein a candidate that is capable of at least one of (i) to (iv) is an anti-cancer agent.

12. The method of claim 10, wherein

10 (i) the test compound is selected from the group consisting of polypeptides, ribonucleic acids, small molecules, and deoxyribonucleic acids;

(ii) the first polypeptide is a first fusion protein comprising a FZD polypeptide fused to (a) a transcription activation domain of a transcription factor or (b) a DNA-binding domain of a transcription factor; the second polypeptide is a second fusion protein comprising a Wnt polypeptide fused to (c) a transcription activation domain of
15 a transcription factor or (d) a DNA-binding domain of a transcription factor, wherein the Wnt polypeptide is fused to a domain different from that fused to the Wnt polypeptide; and binding of the first and second polypeptides is detected as reconstitution of a transcription factor;

(iii) the Wnt polypeptide comprises the amino acid sequence of a Wnt 3 protein;

20 (iv) the Wnt polypeptide comprises the amino acid sequence of a Wnt 8b protein;

or

(v) the Wnt polypeptide comprises the amino acid sequence of a Wnt 11 protein.

25 13. The method of claim 10, wherein the Wnt polypeptide comprises

(i) any one of SEQ ID NO:8 to 12;

(ii) any one of SEQ ID NO: 15 to 19; or

(iii) any one of SEQ ID NO: 22 to 26.

14. A candidate anti-cancer agent or an anti-cancer agent identified by the method of any one of claims 10-13 for use in the treatment of cancer.

5 15. Use of a candidate anti-cancer agent or an anti-cancer agent identified by the method of any one of claims 10-13 in the manufacture of a medicament for the treatment of cancer.

Figure 1




Figure 2




Figure 3





Figure 4



Human Frizzled 7

Accession: O75084

Protein name: Frizzled 7 [Precursor]

DEFINITION: frizzled 7; Frizzled, drosophila, homolog of, 7; frizzled(Drosophila) homolog 7 [Homo sapiens].

ACCESSION NP_003498

ORIGIN (574 aa)

MRDPGAAVPL SSLGFCALVL ALLGALSAGA GAQPYHGEKG ISVDPDHGFCQ PISIPLECTDI
 AYNQTILPNL LGHTNQEDAG LEVHQFYPLV KVQCSPELRF FLCSMYAPVC TVLDQAIPPC
 RSLCERARQG CEALMNKFGF QWPERLRCEN FPVHGAGEIC VGQNTSDGSG GPGGGPTAYP
 TAPYLPDLPF TALPPGASDG KGRPAFPFSC PRQLKVPPYL GYRFLGERDC GAPCEPGRAN
 GLMYFKEEER RFARLWVGWV SVLCCASTLF TVLTYLVDMR RFSYPERPII FLSGCYFMVA
 VAHVAGFFLE DRAVCVERFS DDGYRTVAQG TKKEGCTILF MVLYFFGMAS SIWWVILSLT
 WFLAAGMKWG HEAIEANSQY FHAAWAVPA VKTITILAMG QVDGDLNGV CYVGFSSVDA
 LRGFVLAPLF VYFFIGTSEL LAGFVSFFRI RTIMKHDGK TEKLEKLMVR IGVSFLYTV
 PATIVLACYF YEQAFREHWE RTWLLQTCCKS YAVPCPPGHF PPMSPDFTVF MIKCLMTMIV
 GITTGFWIWS GKTLSQWRRF YHRLSHSSKG ETAV (SEQ ID NO:1)

Human Frizzled 7 Putative ligand binding site

* Cystein-rich domain (CRD)

CPISIPLECTDIAYNQITILPNLLGHTNQEDAGLEVHQFYPLVKVQCSPELRF FLCSMYAPVCRSLCERARQ
GCEALMNKFGFQWPERLRCENFP (49-152) (SEQ ID NO:2)

Mouse Frizzled 7

Accession: Q61090

Protein name: Frizzled 7 [Precursor]

DEFINITION: frizzled 7 [Mus musculus].

ACCESSION NP_032083

ORIGIN (572 aa)

MRGPGTAASH SPLGLCALVL ALLGALPTDT RAQPYHGEKG ISVDPDHGFCQ PISIPLECTDI
 AYNQTILPNL LGHTNQEDAG LEVHQFYPLV KVQCSPELRF FLCSMYAPVC TVLDQAIPPC
 RSLCERARQG CEALMNKFGF QWPERLRCEN FPVHGAGEIC VGQNTSDGSG GAGGSPTAYP
 TAPYLPDPPF TAMSPDGRG RLSFPFSCPR QLKVPPYLYG RFLGERDCGA PCEPGRANGL
 MYFKEEERF ARLWVGWVSV LSCASTLFTV LTYLVDMRRF SYPERPIIFL SGCYFMVAVA
 HVAGFLEDR AVCVERFSDD GYRTVAQGTK KEGCTILEMV LYFFGMASII WWVILSLTWF
 LAAGMKWGHE AIEANSQYFH LAAWAVPAVK TITILAMGQV DGDLLSGVCY VGLSSVDALR
 GFVLAPLFVY LFIGTSFLLA GFVSLFRIRT IMKHDGKTE KLEKLMVRIG VFSVLYTVPA
 TIVLACYFYE QAFREHWERT WLLQTCCKSYA VPCPPRHFSF MSPDFTVFMI KYLMTMIVGI
 TTGFWIWSGK TLQSWRRFYH RLSHSSKGET AV(SEQ ID NO:3)

Human Frizzled 8

Accession: Q9H461

Protein name: Frizzled 8 [Precursor]

Origin (694 aa)

MEWGYLLEVT SLLAALALLQ RSSGAAAASA KELACQEITV PLCKGIGYNY TYMPNQFNHD
 TQDEAGLEVH QFWPLVEIQC SPDLKFFLCS MYTPICLEDY KKPLPPCRSV CERAKAGCAP
 LMRQYGFAPW DRMRCRLPE QGNPDTLCMD YNRDILTAA PSPRRRLPPP PPGEQPPSGS
 GHGRPPGAR PHRGGGRGGG GGDAAPPAR GGGGGKARP PGGGAAPCEP GCQCRAPMVS
 VSSERHPLYN RVKTGQIANC ALPCHNPFPS QDERAFTVFW IGLWSVLCFV STFATVSTFL
 IDMERFKYPE RPIIFLSACY LFSVGYLVR LVAGHEKVAC SGGAPGAGGA GGAGGAAAGA
 GAAGAGAGGP GGRGEYEELG AVEQHVRVET TGPALCTVVF LLVYFFGMAS SIWWVILSLT
 WFLAAGMKWG NEAIAGYSQY FHAAWLVPV VKSIAVLALS SVDGDPVAGI CYVGNQSLDN
 LRGFVLAPLV IYLFIGTMFL LAGFVSLFRI RSVIKQQDGP TKTHKLEKLM IRLGLFTVLY
 TVPAAVVVAC LFYEQHNRPR WEATHNCPCL RDLQPDQARR PDYAVFMLKY FMCLVVGITS
 GVWVWSGKTL ESWRSLCTRC CWASKGAAVG GGAGATAAGG GGGPGGGGGG GPGGGGGPGG
 GGGSLYSDVS TGLTWRSGTA SSVSYPKQMP LSQV (SEQ ID NO:4)

Human FZD8 Putative ligand binding site

* Cystein-rich domain (CRD)

CQEITVPLCKGIGYNYTYMPNQFNHDTQDEAGLEVHQFWPLVEIQCSPLDKFFLCSMYTPICLEDD
YKKPLPPCRSVCERAKAGCAPLMRQYGFAPWDRMRCRLP (35-139) ((SEQ ID NO:5)

Mouse Frizzled 8

Accession: Q61091

Protein name: Frizzled 8 [Precursor]

Origin (685 aa)

MEWGYLLEVT SLLAALAVLQ RSSGAAAASA KELACQEITV PLCKGIGYNY TYMPNQFNHD
 TQDEAGLEVH QFWPLVEIQC SPDLKFFLCS MYTPICLEDY KKPLPPCRSV CERAKAGCAP
 LMRQYGFAPW DRMRCRLPE QGNPDTLCMD YNRDILTAA PSPRRRLPPP PPPGEQPPSG
 SGHSRPPGAR PPHRGGSSRG SGDAAPPAS RGGKARPPGG GAAPCEPGCQ CRAPMVSVSS
 ERHPLYNRVK TGQIANCALP CHNPFPSQDE RAFTVFWIGL WSVLFCFVSTF ATVSTFLIDM
 ERFKYPERP IFLSACYLFRV SVGYLVRLVA GHEKVACSSG APGAGGRGGA GGAAAAGAGA
 AGRGASSPGA RGEYEELGAV EQHVRVETTG PALCTVVFLV VYFFGMASSI WWVILSLTWF
 LAAGMKWGNE AIAGYSQYFH LAAWLVPSVK SIAVLALSSV DGDVPVAGICY VGNQSLDNLR
 GFVLAPLVIIY LFIGTMFLLA GFVSLFRIRS VIKQQGGPTK THKLEKLMIR LGLFTVLYTV
 PAAVVVACLF YEQHNRPRWE ATHNCPCLRD LQPDQARRPD YAVFMLKYFM CLVVGITSGV
 WVWWSGKTLES WRALCTRCCW ASKGAAVGAG AGGSGPGGSG PPGGGGGHG GGGGSLYSDV
 STGLTWRSGT ASSVSYPKQM PLSQV (SEQ ID NO:6)

Human Wnt3

Accession P56703

Protein name: Wnt-3 proto-oncogene protein [Precursor]

ACCESSION NP_110380

Definition: wingless-type MMTV integration site family, member 3; WNT-3 proto-oncogene protein precursor [Homo sapiens]

ORIGIN (355 aa)

MEPHLLGLLL GLLLGTRVL AGYPIWWSLA LGQOYTSLS QPLLCGSIPG LVPKQLRFCR
 NYIEIMPSVA EGVKLGIOEC QHQFRGRRW CTTIDDSLAI FGPVLDKATR ESAFVHAIAS
 AGVAVAVTRS CAEGTSTICG CDSHHKGPPG EGWKWGGCSE DADFGVLVSR EFADARENRP
 DARSAMNKHN NEAGRITILD HMHLKCKCHG LSGSCEVKTC WWAQPDFRAI GDFLKDKYDS
 ASEMVVEKHR ESRGWVETLR AKYSLFKPPT ERDLVYYENS PNFCEPNPET GSFGRDRRTC
 NVTSHGIDGC DLLCCGRGHN TRTEKRKEKC HCIFHWCCYV SCQECIRIYD VHTCK(SEQ ID
 NO:7)

Putative binding motifsHuman Wnt3

* Secreted growth factor protein (motif or domain)

- 1) RESAFVHAIASAGVA (110-124) (SEQ ID NO:8)
- 2) RSCAEGTSTICGCD (129-142) (SEQ ID NO:9)
- 3) WKWGGCSEDADFG (153-165) (SEQ ID NO:10)
- 4) **CKCHGLSGSCEVKTCW (206-221) (SEQ ID NO:11)**
- 5) DLVYYENSPNFC (273-284) (SEQ ID NO:12)

Mouse Wnt3

Accession: P17553

Protein name: Wnt-3 proto-oncogene protein [Precursor]

DEFINITION:wingless-related MMTV integration site 3 [Mus musculus].

ACCESSION NP_033547

ORIGIN (355 aa)

MEPHLLGLLL GLLLSGTRVL AGYPIWWSLA LGQOYTSLS QPLLCGSIPG LVPKQLRFCR
 NYIEIMPSVA EGVKLGIOEC QHQFRGRRW CTTIDDSLAI FGPVLDKATR ESAFVHAIAS
 AGVAVAVTRS CAEGTSTICG CDSHHKGPPG EGWKWGGCSE DADFGVLVSR EFADARENRP
 DARSAMNKHN NEAGRITILD HMHLKCKCHG LSGSCEVKTC WWAQPDFRAI GDFLKDKYDS
 ASEMVVEKHR ESRGWVETLR AKYALFKPPT ERDLVYYENS PNFCEPNPET GSFGRDRRTC
 NVTSHGIDGC DLLCCGRGHN TRTEKRKEKC HCVFHWCCYV SCQECIRIYD VHTCK(SEQ ID
 NO:13)

Human Wnt8B

Accession: Q93098

Protein name: Wnt-8b protein [Precursor]

DEFINITION: wingless-type MMTV integration site family, member 8B precursor [Homo sapiens].

ACCESSION: NP_003384

ORIGIN (351 aa)

MFLSKPSVYI CLFTCVLQLS HSWSVNNFLM TGPKAYLIYS SVAAGAQSG IEECKYQFAW
 DRWNCPERAL QLSSHGGLRS ANRETAHVHA ISSAGVMYTL TRNCSLGDFD NCGCDDSRNG
 QLGGQGWLWG GCSDNVGFGE AISKQFVDAL ETGQDARAAM NLHNNEAGRK AVKGTMKRTC
 KCHGVSGSCT TQTCWLQQLPE FREVGAHLKE KYHAALKVDL LQGAGNSAAA RGAIADTFRS
 ISTRELVHLE DSPDYCLENK TLGLLGTEGR ECLRRGRALG RWELRSCRRL CGDCGLAVEE
 RRAETVSSCN CKFHWCACVR CEQCRRRVTK YFCSRAERPR GAAHKPGRK P (SEQ ID NO:14)

Putative binding motifsHuman Wnt8b

* Secreted growth factor protein (motif or domain)

- 1) RETAFVHAISSAGVM (83-97) (SEQ ID NO:15)
- 2) RNCSLGDFDNCGCD (102-115) (SEQ ID NO:16)
- 3) WLWGGCSDNVGFGE (127-139) (SEQ ID NO:17)
- 4) **CKCHGVSGSCTTQTCW (180-195)** (SEQ ID NO:18)
- 5) ELVHLEDSPDYC (245-256) (SEQ ID NO:19)

Mouse Wnt8B

Accession: Q9WUD6

Protein name: Wnt-8b protein [Precursor]

DEFINITION: wingless related MMTV integration site 8b [Mus musculus].

ACCESSION NP_035850

ORIGIN (350 aa)

MFLMKPVCVL LVTCVLHRSH AWSVNNFLMT GPKAYLVYSS SVAAGAQSGI EECKYQFAWD
 RWNCPERALQ LSSHGGLRSA NRETAHVHAI SSAGVMYTLT RNCSLGDFDN CGCDDSRNGQ
 LGGQGWLWGG CSDNVGFGEA ISKQFVDALE TGQDARAAMN LHNNEAGRKA VKGTMKRTCK
 CHGVSGSCTT QTCWLQQLPEF REVGAHLKEK YHAALKVDLL QGAGNSAAGR GAIADTFRSI
 STRELVHLED SPDYCLENKT LGLLGTEGRE CLRRGRALGR WERRSCRRLC GDCGLAVEER
 RAETVSSCNC KFHWCACVRC EQCRRRVTKY FCSRAERPPR GAAHKPGKNS (SEQ ID NO:20)

FIG. 5D

Human Wnt 11

Accession: O96014

Protein name: Wnt-11 protein [Precursor]

DEFINITION: wingless-type MMTV integration site family,
member 11 precursor [Homo sapiens]

ACCESSION NP_004617

ORIGIN (354 aa)

MRARPQVCEA LLFALALQTG VCYGIKWLAL SKTPSALALN QTQHCKQLEG LVSAQVQLCR
 SNLELMHTTV HAAREVMKAC RRAFADMRWN CSSIELAPNY LLDLERTRE SAFVYALSAA
 AISHAIARAC TSGDLPGCSC GPVPGPEPPG GNRWGGCADN LSYGLLMGAK FSDAPMKVKK
 TGSQANKLMR LHNSEVGRQA LRASLEMKCK CHGVSGSCSI RTCWKGLQEL QDVAADLKTR
 YLSATKVVHR PMGTRKHLVP KDLDIRPVKD SELVYLQSSP DFCMKNEKVG SHGTQDRQCN
 KTSNGSDSCD LMCCGRGYNP YTDRVVERCH CKYHWCCYVT CRRCERTVER YVCK (SEQ ID
 NO:21)

Putative binding motifsHuman Wnt11

* Secreted growth factor protein (motif or domain)

- 1) RESAFVYALSAAAIS (109-123) (SEQ ID NO:22)
- 2) RACTSGDLPGCSCG (128-141) (SEQ ID NO:23)
- 3) NRWGGCADNLSYG (152-164) (SEQ ID NO:24)
- 4) **CKCHGVSGSCSIRTCW (209-224) (SEQ ID NO:25)**
- 5) ELVYLQSSPDFC (272-283) (SEQ ID NO:26)

Mouse Wnt 11

Accession: P48615

Protein name: Wnt-11 protein [Precursor]

DEFINITION: wingless-related MMTV integration site 11 [Mus
musculus].

ACCESSION NP_033545

ORIGIN (354 aa)

MRARPQVCEA LLFALALHTG VCYGIKWLAL SKTPAALALN QTQHCKQLEG LVSAQVQLCR
 SNLELMRTIV HAARGAMKAC RRAFADMRWN CSSIELAPNY LLDLERTRE SAFVYALSAA
 TISHTIARAC TSGDLPGCSC GPVPGPEPPG GNRWGGCADN LSYGLLMGAK FSDAPMKVKK
 TGSQANKLMR LHNSEVGRQA LRASLETKCK CHGVSGSCSI RTCWKGLQEL QDVAADLKTR
 YLSATKVVHR PMGTRKHLVP KDLDIRPVKD SELVYLQSSP DFCMKNEKVG SHGTQDRQCN
 KTSNGSDSCD LMCCGRGYNP YTDRVVERCH CKYHWCCYVT CRRCERTVER YVCK (SEQ ID
 NO:27)