

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2003 (10.04.2003)

PCT

(10) International Publication Number
WO 03/028636 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US02/30574
- (22) International Filing Date:
26 September 2002 (26.09.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/966,451 28 September 2001 (28.09.2001) US
- (71) Applicant: **ISIS PHARMACEUTICALS, INC.**
[US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).
- (72) Inventors: **BENNETT, Frank, C.**; 1347 Cassins Street, Carlsbad, CA 92008 (US). **FREIER, Susan, M.**; 2946 Renault Street, San Diego, CA 92112 (US).
- (74) Agents: **LICATA, Jane, Massey** et al.; Licata & Tyrell P.C., 66 E. Main Street, Marlton, NJ 08053 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 03/028636 A2

(54) Title: ANTISENSE MODULATION OF IL-1 RECEPTOR-ASSOCIATED KINASE-4 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of IL-1 receptor-associated kinase-4. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding IL-1 receptor-associated kinase-4. Methods of using these compounds for modulation of IL-1 receptor-associated kinase-4 expression and for treatment of diseases associated with expression of IL-1 receptor-associated kinase-4 are provided.

-1-

**ANTISENSE MODULATION OF IL-1 RECEPTOR-ASSOCIATED KINASE-4
EXPRESSION**

FIELD OF THE INVENTION

The present invention provides compositions and methods for modulating the expression of IL-1 receptor-associated kinase-4. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding IL-1 receptor-associated kinase-4. Such compounds have been shown to modulate the expression of IL-1 receptor-associated kinase-4.

BACKGROUND OF THE INVENTION

The proinflammatory cytokine interleukin-1 (IL-1) is a central regulator in immune and inflammatory responses, involved in generating systemic and local response to infection, injury, and immunologic challenges. IL-1 is produced mainly by activated macrophages and monocytes, and participates in lymphocyte activation, induction of fever, leukocyte trafficking, the acute phase response, and cartilage remodeling. The expression of more than 90 genes is affected by IL-1, including genes that encode other cytokines, cytokine receptors, acute-phase reactants, growth factors, tissue-remodeling enzymes, extracellular matrix components, and cell adhesion molecules. IL-1 is a critical cytokine in the pathogenesis of viral infections and inflammatory diseases such as rheumatoid arthritis (O'Neill and Greene, *J. Leukoc. Biol.*, **1998**, 63, 650-657).

Cellular responses are transduced through the type I IL-1 receptor (IL-1RI), located on the plasma membrane of a variety of IL-1-responsive cells. Binding of IL-1 to IL-1RI ultimately triggers activation of transcription factors in the NF- κ B family, which are bound by inhibitory proteins (IKBs) and remain anchored

-2-

in the cytoplasm until the inhibitory proteins are degraded. In response to IL-1, tumor necrosis factor (TNF), or other extracellular stimuli such as lipopolysaccharides, double-stranded RNA, or oxidative stress, and once unbound and activated, NF- κ B is then transported to the nucleus, where it influences the activity of many genes (O'Neill and Greene, *J. Leukoc. Biol.*, **1998**, 63, 650-657).

A family of proteins has been described that share significant homology with the type I IL-1 receptor in their signaling domains. This family includes IL-1 receptor accessory protein (IL1-RACp), which does not bind IL-1, but is essential for IL-1 signaling; a *Drosophila* receptor protein, Toll; a number of human Toll-like receptors (hTLRs); the interferon- γ -inducing factor/IL-1 γ / IL-18 receptor-related protein (IL-1Rrp), a number of plant proteins, and the IL-1 receptor-associated kinases (IRAK-1, IRAK-2, and IRAK-M). All members of this family appear to be involved in host responses to injury and infection (O'Neill and Greene, *J. Leukoc. Biol.*, **1998**, 63, 650-657; Wesche et al., *J. Biol. Chem.*, **1999**, 274, 19403-19410).

The IL-1 signaling pathway in mammals is analogous to the Toll pathway in *Drosophila melanogaster*. Homologues of the IL-1 receptor-associated kinases are found in *D. melanogaster* (Pelle) and in plants (Pto), and in these systems, the kinases have been shown to be components of a signal transduction system leading to the activation of NF- κ B. A model for the signaling pathway in which IL-1 receptor associated kinase-4 is likely to function is as follows: when cells receive the extracellular IL-1 signal, a complex between IL-1RI and IL-1RACp is formed (Huang et al., *Proc. Natl. Acad. Sci. U. S. A.*, **1997**, 94, 12829-12832), the cytosolic adapter protein MyD88 interacts with IL-1RACp in the receptor complex (Burns et al., *J. Biol. Chem.*, **1998**, 273, 12203-12209), and MyD88 rapidly recruits a IL-1 receptor-associated kinase into the complex. Tollip also interacts with IL-1RACp and is believed to block autophosphorylation of the IL-1 receptor-associated kinase or its association with another kinase; thus, the association of Tollip with the IL-1 receptor-associated kinase is inhibitory (Burns et al., *Nat. Cell Biol.*, **2000**, 2, 346-351). At some point after its IL-1-dependent

-3-

association with the receptor complex, the IL-1 receptor-associated kinase may be extensively phosphorylated and its own serine/threonine kinase catalytic activity activated (Cao et al., *Science*, **1996**, 271, 1128-1131). IL-1 receptor-associated kinase-1 is known to interact with an adapter protein, TRAF6, a protein critical for IL-1-dependent activation of NF- κ B, which then dissociates from the receptor complex. TRAF6 relays a signal via NF- κ B-inducing kinase (NIK) to two I- κ B kinases (IKK-1 and -2), culminating in activation of NF- κ B (Bacher et al., *FEBS Lett.*, **2001**, 497, 153-158; Jefferies et al., *Mol. Cell. Biol.*, **2001**, 21, 4544-4552; O'Neill and Greene, *J. Leukoc. Biol.*, **1998**, 63, 650-657).

IL-1 receptor-associated kinase-4 (also known as interleukin 1 receptor-associated kinase 4, IRAK4, IRAK-4, DD-1, and NY-REN-64 antigen) was originally identified as NY-REN-64, one of 65 human tumor antigens recognized by autologous antibodies from patients with renal cell carcinoma using serological analysis of recombinant cDNA expression libraries (SEREX). Sequence analysis of the NY-REN-64 cDNA clone identified in the SEREX screen revealed a novel gene encoding a transcript of 2.8 kilobases and a predicted protein of 460 amino acids (Genbank Accession AF155118) noted to bear a protein kinase motif (Scanlan et al., *Int. J. Cancer*, **1999**, 83, 456-464). Based on its homology to the other IL-1 receptor-associated kinases, this gene has more recently been placed in the IRAK family and given the name IL-1 receptor-associated kinase-4.

There is evidence that the IRAKs are functionally redundant *in vivo*. Members of this family of IL-1 receptor-associated kinases may also have overlapping yet distinct signaling roles. IRAK-2 and IRAK-M can reconstitute the IL-1 response in human embryonic kidney 293 mutant cell lines lacking IRAK-1 (Wesche et al., *J. Biol. Chem.*, **1999**, 274, 19403-19410). Furthermore, at least IL-1 receptor-associated kinase-1 plays a role in regulation of multiple signaling pathways. In addition to transducing the IL-1 signal, IL-1 receptor-associated kinase-1 also transduces a signal initiated by binding of tumor necrosis factor (TNF)- α to its receptor, again leading to activation of

-4-

NF- κ B (Vig et al., *J. Biol. Chem.*, **1999**, 274, 13077-13084; Vig et al., *J. Biol. Chem.*, **2001**, 276, 7859-7866). In another signal transduction pathway separate from its activation of NF- κ B, IL-1 receptor-associated kinase-1 has also been implicated in activation of the Jun amino-terminal kinase (JNK) and the transcription factor AP-1 (Bacher et al., *FEBS Lett.*, **2001**, 497, 153-158).

Disclosed and claimed in PCT Publication WO 00/73801 are methods of diagnosing a disorder and methods for treating a pathological cell condition characterized by aberrant expression of a human cancer associated antigen encoded by a nucleic acid molecule, wherein the nucleic acid molecule is IL-1 receptor-associated kinase-4 or a fragment thereof, and wherein the agent used in said method of treatment is an antisense nucleic acid molecule. Also claimed is an isolated nucleic acid molecule of sufficient length to represent a sequence unique within the human genome, identifying a nucleic acid encoding IL-1 receptor-associated kinase-4. Further claimed is a method for treating a subject having a condition characterized by IL-1 receptor-associated kinase-4 expression (Obata, **2000**).

Disclosed and claimed in PCT Publication WO 01/51641 are isolated nucleic acids and expression vector constructs encoding an IL-1 receptor-associated kinase-4 polypeptide as well as isolated polypeptides with at least about 98% amino acid sequence identity to IL-1 receptor-associated kinase-4. Further claimed is a method for identifying a compound useful in the treatment or prevention of inflammatory diseases by expressing IL-1 receptor-associated kinase-4 and truncated or mutant forms of the IL-1 receptor-associated kinase-4 protein. Also claimed is a method of treating an inflammatory disease in a patient by administering a compound that modulates IL-1 receptor-associated kinase-4, as well as a method of inhibiting the transduction of a signal resulting from the activation of an IL-1R/Toll receptor in a cell, comprising introducing into said cell an inhibitor of the activity or expression of IL-1 receptor-associated kinase-4 (Wesche and Li, *WO Application*, **2001**).

IL-1 receptor-associated kinase-4 is involved in tumorigenesis, and the pharmacological modulation of IL-1

-5-

receptor-associated kinase-4 activity and/or expression is therefore believed to be an appropriate point of therapeutic intervention in pathological conditions such as cancer, as well as viral infections, rheumatoid arthritis, and other inflammatory disease and immune disorders.

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of IL-1 receptor-associated kinase-4. Consequently, there remains a long felt need for agents capable of effectively inhibiting IL-1 receptor-associated kinase-4 function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of IL-1 receptor-associated kinase-4 expression.

The present invention provides compositions and methods for modulating IL-1 receptor-associated kinase-4 expression.

SUMMARY OF THE INVENTION

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding IL-1 receptor-associated kinase-4, and which modulate the expression of IL-1 receptor-associated kinase-4. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of IL-1 receptor-associated kinase-4 in cells or tissues comprising contacting said cells or tissues with one or more of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of IL-1 receptor-associated kinase-4 by administering a therapeutically or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric compounds,

-6-

particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding IL-1 receptor-associated kinase-4, ultimately modulating the amount of IL-1 receptor-associated kinase-4 produced. This is accomplished by providing antisense compounds which specifically hybridize with one or more nucleic acids encoding IL-1 receptor-associated kinase-4. As used herein, the terms "target nucleic acid" and "nucleic acid encoding IL-1 receptor-associated kinase-4" encompass DNA encoding IL-1 receptor-associated kinase-4, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of IL-1 receptor-associated kinase-4. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a

-7-

nucleic acid molecule encoding IL-1 receptor-associated kinase-4. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding IL-1 receptor-associated kinase-4, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50

-8-

contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with

-9-

sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense and other compounds of the invention which hybridize to the target and inhibit expression of the target are identified through experimentation, and the sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The target sites to which these preferred sequences are complementary are hereinbelow referred to as

-10-

"active sites" and are therefore preferred sites for targeting. Therefore another embodiment of the invention encompasses compounds which hybridize to these active sites.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with other antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

Expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds which affect expression patterns.

Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, **2000**, 480, 17-24; Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., *Drug Discov. Today*, **2000**, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, **1999**, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U. S. A.*, **2000**, 97, 1976-81), protein arrays and proteomics (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16; Jungblut, et al., *Electrophoresis*, **1999**, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, **2000**, 480, 2-16;

-11-

Larsson, et al., *J. Biotechnol.*, **2000**, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., *Anal. Biochem.*, **2000**, 286, 91-98; Larson, et al., *Cytometry*, **2000**, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, **2000**, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, **1998**, 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, **1999**, 35, 1895-904) and mass spectrometry methods (reviewed in (To, *Comb. Chem. High Throughput Screen*, **2000**, 3, 235-41).

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes, have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 50 nucleobases (i.e. from about 8 to about 50 linked nucleosides). Particularly preferred

-12-

antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates,

-13-

5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Representative United States patents that teach the

-14-

preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH₂-NH-O-CH₂-, -CH₂-N(CH₃)-O-CH₂- [known as a methylene (methylimino) or MMI backbone], -CH₂-O-N(CH₃)-CH₂-, -CH₂-N(CH₃)-N(CH₃)-CH₂- and -O-N(CH₃)-CH₂-CH₂- [wherein the native phosphodiester backbone is represented as -O-P-O-CH₂-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise

-15-

one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, **1995**, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₃)₂, also described in examples hereinbelow.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH₂-)_n group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂), 2'-allyl (2'-CH₂-CH=CH₂), 2'-O-allyl (2'-O-CH₂-CH=CH₂) and 2'-fluoro (2'-F). The 2' modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on

-16-

the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl ($-C\equiv C-CH_3$) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-

-17-

pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, **1990**, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, **1991**, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B. , ed., CRC Press, **1993**. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, **1993**, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one

-18-

or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. The compounds of the invention can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992 the entire disclosure of which is incorporated herein by reference. Conjugate moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, **1989**, *86*, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1994**, *4*, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, **1992**, *660*, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, **1993**, *3*, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, **1992**, *20*, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, **1991**, *10*, 1111-1118; Kabanov et al., *FEBS Lett.*, **1990**, *259*, 327-330; Svinarchuk et al., *Biochimie*, **1993**, *75*, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, **1995**, *36*, 3651-3654; Shea et al., *Nucl. Acids Res.*, **1990**, *18*, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, **1995**, *14*, 969-973), or adamantane acetic acid

-19-

(Manoharan et al., *Tetrahedron Lett.*, **1995**, 36, 3651-3654), a palmitoyl moiety (Mishra et al., *Biochim. Biophys. Acta*, **1995**, 1264, 229-237), or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, **1996**, 277, 923-937. Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are

-20-

antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or

-21-

alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO

-22-

94/26764 and U.S. 5,770,713 to Imbach et al.

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, **1977**, *66*, 1-19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid,

-23-

glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of IL-1

-24-

receptor-associated kinase-4 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding IL-1 receptor-associated kinase-4, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding IL-1 receptor-associated kinase-4 can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of IL-1 receptor-associated kinase-4 in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments,

-25-

lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C₁₋₁₀ alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitables. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts

-26-

include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucholic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate, . Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprate, tricaprata, monoolein, dilaurin, glyceryl 1-monocaprata, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pollulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines, protamine, polyvinylpyridine, polythiodiethylamino-methylethylene P(TDAE), polyaminostyrene (e.g. p-amino), poly(methylcyanoacrylate), poly(ethylcyanoacrylate), poly(butylcyanoacrylate), poly(isobutylcyanoacrylate), poly(isohexylcynaoacrylate), DEAE-methacrylate, DEAE-hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran, polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid), poly(DL-lactic-co-glycolic acid (PLGA), alginate, and polyethyleneglycol (PEG). Oral formulations for oligonucleotides and their preparation are described in detail in United States applications 08/886,829 (filed July 1, 1997), 09/108,673 (filed July 1, 1998), 09/256,515 (filed February 23,

-27-

1999), 09/082,624 (filed May 21, 1998) and 09/315,298 (filed May 20, 1999) each of which is incorporated herein by reference in their entirety.

Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations

-28-

vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, **1985**, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not.

-29-

Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin

-30-

and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration

-31-

of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., **1989**, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three

-32-

to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, **1985**, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), **1988**, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprinate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited

-33-

to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, **1993**, *13*, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, **1994**, *11*, 1385; Ho et al., *J. Pharm. Sci.*, **1996**, *85*, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as

-34-

belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

-35-

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang *et al.*, *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou *et al.*, *Journal of Controlled Release*, **1992**, 19, 269-274).

-36-

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner *et al.*, *Journal of Drug Targeting*, **1992**, *2*, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis *et al.*, *Antiviral Research*, **1992**, *18*, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu *et al.* *S.T.P. Pharma. Sci.*, **1994**, *4*, *6*, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one

-37-

or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{M1} , or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen *et al.*, *FEBS Letters*, **1987**, 223, 42; Wu *et al.*, *Cancer Research*, **1993**, 53, 3765).

Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos *et al.* (*Ann. N.Y. Acad. Sci.*, **1987**, 507, 64) reported the ability of monosialoganglioside G_{M1} , galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon *et al.* (*Proc. Natl. Acad. Sci. U.S.A.*, **1988**, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen *et al.*, disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G_{M1} or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb *et al.*) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-*sn*-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim *et al.*).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto *et al.* (*Bull. Chem. Soc. Jpn.*, **1980**, 53, 2778) described liposomes comprising a nonionic detergent, 2C₁₂15G, that contains a PEG moiety. Illum *et al.* (*FEBS Lett.*, **1984**, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klibanov *et al.* (*FEBS*

-38-

Lett., 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume *et al.* (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, *e.g.*, DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle *et al.* (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin *et al.* (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin *et al.*) and in WO 94/20073 (Zalipsky *et al.*) Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi *et al.*). U.S. Patent Nos. 5,540,935 (Miyazaki *et al.*) and 5,556,948 (Tagawa *et al.*) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.* describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love *et al.* discloses liposomes comprising antisense oligonucleotides targeted to the *raf* gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, *e.g.* they are self-

-39-

optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

-40-

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced.

-41-

In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, **1988**, *40*, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein (1-monooleoyl-*rac*-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C₁₋₁₀ alkyl esters thereof (e.g., methyl, isopropyl and *t*-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, *7*, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, **1992**, *44*, 651-654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, **1996**, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and

-42-

polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, page 92; Swinyard, Chapter 39 *In: Remington's Pharmaceutical Sciences*, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, **1990**, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, **1992**, 263, 25; Yamashita et al., *J. Pharm. Sci.*, **1990**, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, **1993**, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, 7, 1-33; Buur et al., *J. Control Rel.*, **1990**, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, **1990**, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, **1991**, page 92); and non-steroidal anti-inflammatory agents such as diclofenac

-43-

sodium, indomethacin and phenylbutazone (Yamashita *et al.*, *J. Pharm. Pharmacol.*, **1987**, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi *et al.*, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo *et al.*, PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (*i.e.*, does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'-isothiocyanostilbene-2,2'-disulfonic acid (Miyao *et al.*, *Antisense Res. Dev.*, **1995**, 5, 115-121; Takakura *et al.*, *Antisense & Nucl. Acid Drug Dev.*, **1996**, 6, 177-183).

-44-

Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

-45-

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone,

-46-

tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. **1987**, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., **1987**, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the

-47-

course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC_{50} s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

-48-

EXAMPLES**Example 1****Nucleoside Phosphoramidites for Oligonucleotide Synthesis
Deoxy and 2'-alkoxy amidites**

2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, **1993**, *21*, 3197-3203] using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

2'-Fluoro amidites**2'-Fluorodeoxyadenosine amidites**

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, **1993**, *36*, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

-49-

2'-Fluorodeoxyguanosine

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutyryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

2'-Fluorodeoxycytidine

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, **1995**, 78, 486-504.

2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenyl-carbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced

-50-

pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

2'-O-Methoxyethyl-5-methyluridine

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/acetone/MeOH (20:5:3) containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added

-51-

and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product). The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set aside. Triethylamine (189 mL, 1.44

-52-

M) was added to a solution of triazole (90 g, 1.3 M) in CH₃CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl₃ was added dropwise, over a 30 minute period, to the stirred solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO₃ and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M) in dioxane (500 mL) and NH₄OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was evaporated and the residue azeotroped with MeOH (2x200 mL). The residue was dissolved in MeOH (300 mL) and transferred to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH₃ gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl₃ (700 mL) and extracted with saturated NaHCO₃ (2x300 mL) and saturated NaCl

-53-

(2x300 mL), dried over $MgSO_4$ and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et_3NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-cytidine-3'-amidite

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-cytidine (74 g, 0.10 M) was dissolved in CH_2Cl_2 (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra(isopropyl)-phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction mixture was extracted with saturated $NaHCO_3$ (1x300 mL) and saturated $NaCl$ (3x300 mL). The aqueous washes were back-extracted with CH_2Cl_2 (300 mL), and the extracts were combined, dried over $MgSO_4$ and concentrated. The residue obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

5'-O-tert-Butyldiphenylsilyl-0²-2'-anhydro-5-methyluridine

0²-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at

-54-

ambient temperature under an argon atmosphere and with mechanical stirring. *tert*-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (Rf 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

5'-O-*tert*-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-*tert*-Butyldiphenylsilyl-0²-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (Rf 0.67 for desired product and Rf 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was purified by column chromatography (2kg silica gel,

-55-

ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine

5'-O-tert-Butylidiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P₂O₅ under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

5'-O-tert-butylidiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine

2'-O-([2-phthalimidoxy)ethyl]-5'-t-butylidiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH₂Cl₂ (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH₂Cl₂, and the combined organic phase was washed with water, brine and dried over anhydrous Na₂SO₄. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting

-56-

mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy) ethyl]-5-methyluridine as white foam (1.95 g, 78%).

5'-O-*tert*-Butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine

5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH₂Cl₂). Aqueous NaHCO₃ solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na₂SO₄, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO₃ (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH₂Cl₂ to get 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then

-57-

added to 5'-O-*tert*-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH₂Cl₂). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH₂Cl₂ to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P₂O₅ under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine (26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH₂Cl₂ (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxylethyl)-5-methyluridine (1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P₂O₅ under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,Nⁱ,Nⁱ-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO₃ (40mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and concentrated. Residue

-58-

obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminoxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

2'-(Aminoxyethoxy) nucleoside amidites

2'-(Aminoxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminoxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

The 2'-O-aminoxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-([2-phthalamidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-

-59-

CH₂-O-CH₂-N(CH₂)₂, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O²-,2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH₂Cl₂ (2x200 mL). The combined CH₂Cl₂ layers are washed with saturated NaHCO₃ solution, followed by saturated NaCl solution and dried over anhydrous sodium sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH₂Cl₂:Et₃N (20:1, v/v, with 1% triethylamine) gives the title compound.

5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite

-60-

Diisopropylaminotetrazolidide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH₂Cl₂ (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

Example 2**Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and

-61-

PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

Example 3

Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, herein incorporated by reference.

Example 4

PNA Synthesis

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, **1996**, *4*, 5-23. They may also be prepared in accordance with U.S. Patents

-62-

5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Example 5

Synthesis of Chimeric Oligonucleotides

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate Oligonucleotides

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

-63-

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)]
Chimeric Phosphorothioate Oligonucleotides**

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotides

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites, oxidization with iodine to generate the phosphodiester internucleotide linkages within the wing portions of the chimeric structures and sulfurization utilizing 3,4-dihydro-2H-benzodithiole-3-one 1,1 dioxide (Beaucage Reagent) to generate the phosphorothioate internucleotide linkages for the center gap.

Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States patent 5,623,065, herein incorporated by reference.

Example 6

Oligonucleotide Isolation

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides were analyzed by polyacrylamide gel electrophoresis on denaturing gels and judged to be at least 85% full length material. The relative amounts of phosphorothioate and phosphodiester linkages obtained in synthesis were periodically

-64-

checked by ^{31}P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* **1991**, *266*, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

Example 7**Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated NH_4OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

Example 8**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by

-65-

mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

Example 9**Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following 4 cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, Ribonuclease protection assays, or RT-PCR.

T-24 cells:

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA).

-66-

A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200 μ L OPTI-MEMTM-1 reduced-serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEMTM-1 containing 3.75 μ g/mL LIPOFECTINTM (Gibco BRL) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control

-67-

oligonucleotide is ISIS 13920, **TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to human H-ras. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCCAAGGA**, SEQ ID NO: 2, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments.

Example 10

Analysis of oligonucleotide inhibition of IL-1 receptor-associated kinase-4 expression

Antisense modulation of IL-1 receptor-associated kinase-4 expression can be assayed in a variety of ways known in the art. For example, IL-1 receptor-associated kinase-4 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems,

-68-

Foster City, CA and used according to manufacturer's instructions.

Protein levels of IL-1 receptor-associated kinase-4 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to IL-1 receptor-associated kinase-4 can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., **1997**. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., **1997**.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., **1998**. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., **1997**. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., **1991**.

Example 11

Poly(A)+ mRNA isolation

Poly(A)+ mRNA was isolated according to Miura et al., *Clin. Chem.*, **1996**, 42, 1758-1764. Other methods for poly(A)+ mRNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., **1993**. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM

-69-

vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55 μ L of lysate was transferred to Oligo d(T) coated 96-well plates (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200 μ L of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60 μ L of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C hot plate for 5 minutes, and the eluate was then transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

Example 12

Total RNA Isolation

Total RNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 μ L cold PBS. 100 μ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100 μ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a

-70-

collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 μ L water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 μ L water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

Example 13

Real-time Quantitative PCR Analysis of IL-1 receptor-associated kinase-4 mRNA Levels

Quantitation of IL-1 receptor-associated kinase-4 mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE, FAM, or VIC, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of

-71-

Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 μ L PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl₂, 300 μ M each of dATP, dCTP and dGTP, 600 μ M of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 μ L total RNA solution. The RT reaction was carried out by incubation for 30

-72-

minutes at 48°C. Following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA using RiboGreen™ (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen™ RNA quantification reagent from Molecular Probes. Methods of RNA quantification by RiboGreen™ are taught in Jones, L.J., et al, *Analytical Biochemistry*, **1998**, 265, 368-374.

In this assay, 175 µL of RiboGreen™ working reagent (RiboGreen™ reagent diluted 1:2865 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 25uL purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

Probes and primers to human IL-1 receptor-associated kinase-4 were designed to hybridize to a human IL-1 receptor-associated kinase-4 sequence, using published sequence information (GenBank accession number NM_016123, incorporated herein as SEQ ID NO:3). For human IL-1 receptor-associated kinase-4 the PCR primers were:
forward primer: CAGTTGAAGCTATGTACTCTGGTGCTA (SEQ ID NO: 4)
reverse primer: AGCTGGTGAACCTTCTTAATGTCTG (SEQ ID NO: 5) and the PCR probe was: FAM-CCAATGTCGGCATGAAAAGAAAATAAGAGC-TAMRA (SEQ ID NO: 6) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye. For human GAPDH the PCR primers were:
forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO:7)
reverse primer: GAAGATGGTGATGGGATTTTC (SEQ ID NO:8) and the PCR probe was: 5' JOE-CAAGCTTCCCCTTCTCAGCC- TAMRA 3' (SEQ ID NO: 9) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems,

-73-

Foster City, CA) is the quencher dye.

Example 14**Northern blot analysis of IL-1 receptor-associated kinase-4 mRNA levels**

Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA) and then probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions.

To detect human IL-1 receptor-associated kinase-4, a human IL-1 receptor-associated kinase-4 specific probe was prepared by PCR using the forward primer CAGTTGAAGCTATGTACTCTGGTGCTA (SEQ ID NO: 4) and the reverse primer AGCTGGTGAACCTTCTTAATGTCTG (SEQ ID NO: 5). To normalize for variations in loading and transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

-74-

Example 15**Antisense inhibition of human IL-1 receptor-associated kinase-4 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human IL-1 receptor-associated kinase-4 RNA, using published sequences (GenBank accession number NM_016123, incorporated herein as SEQ ID NO: 3, and residues 116001- 147000 from GenBank accession number AC016143, the complement of which is incorporated herein as SEQ ID NO: 10). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human IL-1 receptor-associated kinase-4 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 1

Inhibition of human IL-1 receptor-associated kinase-4 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	%_INHIB	SEQ ID NO
156448	Start Codon	3	31	tcttctattcctgcccgggc	27	11
156449	Start Codon	3	36	gttcatccttctattcctgcc	70	12
156450	Coding	3	62	acatatgttgatggtggttat	49	13
156451	Coding	3	85	ttagtccaacattgaggcag	72	14
156452	Coding	3	94	gcttcctaattagtccaaca	71	15
156453	Coding	3	122	ccttcttgaggatcaataaa	59	16
156454	Coding	3	149	ttaatagctacagctaactt	24	17
156455	Coding	3	180	ctgattgtatctatcatcac	60	18

-75-

156456	Coding	3	238	attcagaagtgggacttttt	38	19
156457	Coding	3	245	aacagtaattcagaagtggg	8	20
156458	Coding	3	260	gtgggtgccccagtcaaacag	46	21
156459	Coding	3	270	tgtgcaatttgtgggtgcccc	62	22
156460	Coding	3	347	ggaacagcatctgggagcaa	19	23
156461	Coding	3	371	gaaggtagtgtatttagcagt	73	24
156462	Coding	3	394	gctgaactgttatagcttct	74	25
156463	Coding	3	426	cctgtctttgtcacagaaag	65	26
156464	Coding	3	431	aatgtcctgtctttgtcaca	68	27
156465	Coding	3	440	gggtgcatcaatgtcctgtc	65	28
156466	Coding	3	561	tgtgacattcttcaattcat	7	29
156467	Coding	3	639	gcctttatatacaactccaa	84	30
156468	Coding	3	683	attgtgcaagcttcttcac	61	31
156469	Coding	3	703	cttcagtagtaatgtcaacc	60	32
156470	Coding	3	777	aagtagttctactaagtttt	53	33
156471	Coding	3	800	tcatctccatcacttgagaa	72	34
156472	Coding	3	841	gcaatgaaccattaggcatg	67	35
156473	Coding	3	891	catgtgccaagaaagtgggtg	62	36
156474	Coding	3	914	gcacctgagcaatcttgca	75	37
156475	Coding	3	922	cattagctgcacctgagca	81	38
156476	Coding	3	941	tcatgtagaaaattgatgcc	55	39
156477	Coding	3	950	tgatgattttcatgtagaaa	52	40
156478	Coding	3	966	aatatctctatgaatatgat	18	41
156479	Coding	3	1014	agatatttttagcagtaaaag	16	42
156480	Coding	3	1081	ttcccacaattctgctagtc	69	43
156481	Coding	3	1086	tgttgttcccacaattctgc	32	44
156482	Coding	3	1113	acgcaaagcttctggtgcca	27	45
156483	Coding	3	1190	acagctggaagtccagttat	63	46
156485	Coding	3	1220	agcaataactgaggttcacg	60	47
156486	Coding	3	1227	aatatctagcaataactgag	7	48
156488	Coding	3	1228	taatatctagcaataactga	79	49
156490	Coding	3	1399	tcctctgcagcagctgggtga	75	50
156492	Stop Codon	3	1422	aataaagttttaagaagctg	11	51
156494	3'UTR	3	1548	atgctttaataaccactgtc	65	52
156496	3'UTR	3	1560	gaagttcaaccatgcttta	78	53
156498	3'UTR	3	1646	atttctcagggattactgta	73	54
156500	3'UTR	3	1674	aaactgtgtttgggtgatgct	66	55
156502	3'UTR	3	1711	tacagcccaggctctttttg	25	56
156504	3'UTR	3	1720	caccctacatacagcccagg	55	57
156506	3'UTR	3	1737	tcagatcagagtgtttccac	64	58
156508	3'UTR	3	1756	tagtggagtcagctgggctt	63	59
156510	3'UTR	3	1810	ttattagtggctcacagcag	55	60
156512	3'UTR	3	1829	agcagatattagcccaatgt	82	61
156514	3'UTR	3	1846	acctgtcagagaagcacagc	72	62
156516	3'UTR	3	1854	tcatgactacctgtcagaga	71	63
156518	3'UTR	3	1894	atttacaagtgctttgtata	47	64
156520	3'UTR	3	1951	tgactaataggattttgtaa	55	65
156522	3'UTR	3	1985	taaatgattgctgtgaacac	63	66
156524	3'UTR	3	2066	ccttcaactttgtcatgtaa	83	67
156526	3'UTR	3	2089	accttaactgcatctgccaa	73	68
156528	3'UTR	3	2137	tggattaggtcaggcccttt	64	69
156530	3'UTR	3	2191	ctcacatacttcttcaaggc	0	70
156532	3'UTR	3	2211	gttttagccaatgtggccct	66	71
156534	3'UTR	3	2218	cctttagggttttagccaatg	53	72
156536	3'UTR	3	2223	ggccaccttttaggttttagc	57	73
156538	3'UTR	3	2236	ctcatctccttagaggccacc	35	74
156540	3'UTR	3	2256	ctgacaactggaaggtaggt	71	75
156542	3'UTR	3	2267	tttctgtctgtgacaact	68	76
156544	3'UTR	3	2364	gtggttatcatctgaagctt	11	77

-76-

156546	3'UTR	3	2377	gtcagcccaggctgtgggta	45	78
156548	3'UTR	3	2419	tagattctcactgaggat	55	79
156550	3'UTR	3	2469	tgaccattatctcacagtt	67	80
156552	Exon 1	10	755	aatgccttacctgcccgggc	21	81
156554	Coding	10	2898	cattatgaacaaaagctggt	51	82
156556	Coding	10	16646	agggttagacagacaatgca	34	83
156558	Coding	10	18506	gtggcagcatctgtggtaag	56	84
156560	Exon 9	10	24229	cggaacttaccacaccaaag	24	85
156562	Coding	10	25400	ctagtaaacctatgaagac	20	86
156564	Coding	10	27190	gtgcttctaatagcgcaatt	35	87
156566	Exon 11	10	28296	aaatgcataccttcttaatg	8	88

As shown in Table 1, SEQ ID NOs 12, 13, 14, 15, 16, 18, 19, 21, 22, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 43, 44, 46, 47, 49, 50, 52, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 76, 78, 79, 80, 82, 83, 84 and 87 demonstrated at least 30% inhibition of human IL-1 receptor-associated kinase-4 expression in this assay and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

Example 16

Western blot analysis of IL-1 receptor-associated kinase-4 protein levels

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to IL-1 receptor-associated kinase-4 is used, with a radiolabelled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

-77-

What is claimed is:

1. A compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding IL-1 receptor-associated kinase-4, wherein said compound specifically hybridizes with said nucleic acid molecule encoding IL-1 receptor-associated kinase-4 and inhibits the expression of IL-1 receptor-associated kinase-4.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 12, 13, 14, 15, 16, 18, 19, 21, 22, 24, 25, 26, 27, 28, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 43, 44, 46, 47, 49, 50, 52, 53, 54, 55, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 76, 78, 79, 80, 82, 83, 84 or 87.

4. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

5. The compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.

6. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

7. The compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

9. The compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.

10. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

11. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding IL-1 receptor-associated kinase-4.

12. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

-78-

13. The composition of claim 12 further comprising a colloidal dispersion system.

14. The composition of claim 12 wherein the compound is an antisense oligonucleotide.

15. A method of inhibiting the expression of IL-1 receptor-associated kinase-4 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of IL-1 receptor-associated kinase-4 is inhibited.

16. A method of treating an animal having a disease or condition associated with IL-1 receptor-associated kinase-4 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of IL-1 receptor-associated kinase-4 is inhibited.

17. The method of claim 16 wherein the disease or condition is cancer.

18. The method of claim 17 wherein the cancer is renal.

19. The method of claim 16 wherein the disease or condition is an inflammatory disease.

20. The method of claim 16 wherein the disease or condition is an infection.

SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.
C. Frank Bennett
Susan M. Freier

<120> ANTISENSE MODULATION OF IL-1 RECEPTOR-ASSOCIATED KINASE-4
EXPRESSION

<130> RTSP-0420

<150> 09/966,451
<151> 2001-09-28

<160> 88

<210> 1
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1
tccgtcatcg ctcctcaggg 20

<210> 2
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2
atgcattctg cccccaagga 20

<210> 3
<211> 2817
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (50)...(1432)

<400> 3
gttcttctgt cgccggcttc agcagcccgc gcccgggcag gaatagaag atg aac aaa 58
Met Asn Lys
1

ccc ata aca cca tca aca tat gtg cgc tgc ctc aat gtt gga cta att 106
Pro Ile Thr Pro Ser Thr Tyr Val Arg Cys Leu Asn Val Gly Leu Ile

5	10	15	
agg aag ctg tca gat ttt att gat cct caa gaa gga tgg aag aag tta			154
Arg Lys Leu Ser Asp Phe Ile Asp Pro Gln Glu Gly Trp Lys Lys Leu			
20	25	30	35
gct gta gct att aaa aaa cca tct ggt gat gat aga tac aat cag ttt			202
Ala Val Ala Ile Lys Lys Pro Ser Gly Asp Asp Arg Tyr Asn Gln Phe			
40	45	50	
cac ata agg aga ttt gaa gca tta ctt caa act gga aaa agt ccc act			250
His Ile Arg Arg Phe Glu Ala Leu Leu Gln Thr Gly Lys Ser Pro Thr			
55	60	65	
tct gaa tta ctg ttt gac tgg ggc acc aca aat tgc aca gtt ggt gat			298
Ser Glu Leu Leu Phe Asp Trp Gly Thr Thr Asn Cys Thr Val Gly Asp			
70	75	80	
ctt gtg gat ctt ttg atc caa aat gaa ttt ttt gct cct gcg agt ctt			346
Leu Val Asp Leu Leu Ile Gln Asn Glu Phe Phe Ala Pro Ala Ser Leu			
85	90	95	
ttg ctc cca gat gct gtt ccc aaa act gct aat aca cta cct tct aaa			394
Leu Leu Pro Asp Ala Val Pro Lys Thr Ala Asn Thr Leu Pro Ser Lys			
100	105	110	115
gaa gct ata aca gtt cag caa aaa cag atg cct ttc tgt gac aaa gac			442
Glu Ala Ile Thr Val Gln Gln Lys Gln Met Pro Phe Cys Asp Lys Asp			
120	125	130	
agg aca ttg atg aca cct gtg cag aat ctt gaa caa agc tat atg cca			490
Arg Thr Leu Met Thr Pro Val Gln Asn Leu Glu Gln Ser Tyr Met Pro			
135	140	145	
cct gac tcc tca agt cca gaa aat aaa agt tta gaa gtt agt gat aca			538
Pro Asp Ser Ser Ser Pro Glu Asn Lys Ser Leu Glu Val Ser Asp Thr			
150	155	160	
cgt ttt cac agt ttt tca ttt tat gaa ttg aag aat gtc aca aat aac			586
Arg Phe His Ser Phe Ser Phe Tyr Glu Leu Lys Asn Val Thr Asn Asn			
165	170	175	
ttt gat gaa cga ccc att tct gtt ggt ggt aat aaa atg gga gag gga			634
Phe Asp Glu Arg Pro Ile Ser Val Gly Gly Asn Lys Met Gly Glu Gly			
180	185	190	195
gga ttt gga gtt gta tat aaa ggc tac gta aat aac aca act gtg gca			682
Gly Phe Gly Val Val Tyr Lys Gly Tyr Val Asn Asn Thr Thr Val Ala			
200	205	210	
gtg aag aag ctt gca gca atg gtt gac att act act gaa gaa ctg aaa			730
Val Lys Lys Leu Ala Ala Met Val Asp Ile Thr Thr Glu Glu Leu Lys			
215	220	225	
cag cag ttt gat caa gaa ata aaa gta atg gca aag tgt caa cat gaa			778
Gln Gln Phe Asp Gln Glu Ile Lys Val Met Ala Lys Cys Gln His Glu			
230	235	240	
aac tta gta gaa cta ctt ggt ttc tca agt gat gga gat gac ctc tgc			826

gaacttccaa aatataaaaa tagagccacc atatcaacac ttagccctac ccattagtat 1632
 cacccccagt tcttacagta atccctgaga aatctccttc aagcatcacc aaacacagtt 1692
 tgaaaattac agggtttagca aaaagagcct gggctgtatg taggggtggaa aactctgat 1752
 ctgaagccca gctgactcca ctactaattt gctgtaaagc tttggacata cacttagctg 1812
 ctgtgagcca ctaataacat tgggctaata tctgtctgtc ttctctgaca ggtagtcatg 1872
 aaaatcaaat gatgcaaat atatacaagc actttgtaaa ttgtaaaatg atacaaaatt 1932
 taaagtttat agagccagtt acaaaatcct attagtcata tatttataga ttgtgttcac 1992
 agcaatcatt taaccacaaa taaaatatcc cttgatgata ctgccataat gatatgtcca 2052
 ttattagatt atgttacatg acaaagttga aggaatttgg cagatgcagt taaggttcct 2112
 aaacaactca ctttgagact gttgaaaggg cctgacctaa tccaagtga ccccttgcaa 2172
 gaagaattct ccttgtaagc cttgaagaag tatgtgagag ggccacattg gctaaaaact 2232
 aaaggtggcc tctaggagat gagacctacc ttccagttgt cagcaagcag gaaaaaaaaa 2292
 ttgggacctc agttgcaacc acaaggaact gaattctgcc aaaaatctga gtcagcttag 2352
 aagagtactc caagcttcag atgataacca cagcctgggc tgacacctgg atttcagctt 2412
 tgcagatcc tcagtatgag aatctatctg ttctgtgctg gacttctaata atatagaact 2472
 gtgagataat gggtcacatt ggctggatgt ggtggctcat acctgtaaat cccagcactt 2532
 tgggaggccg aggcaggcag atcacctgag gtcaagagtt caagaccggc ctggccaaca 2592
 tggtgaaacc ccgtctctac taaaaataca aaaattagac gagcgtggtg gtggacacct 2652
 gtagtcccag ctgcttggga ggctgaggca ggagactagc tggaaccagg gaggtagagg 2712
 ttgcagtgag ctgagatcgt gccactgcac tccagcctgg gtgacagagt gagactccat 2772
 cataaataaa taaataaata aatgggtccc attaagcctt taaaa 2817

<210> 4
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 4
 cagttgaagc tatgtactct ggtgcta

27

<210> 5
 <211> 25
 <212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5

agctggtgaa ccttcttaat gtctg

25

<210> 6

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 6

ccaatgctcg catgaaaaga aaaataagag c

31

<210> 7

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 7

gaaggtgaag gtcggagtc

19

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 8

gaagatggtg atgggatttc

20

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 9

caagcttccc gttctcagcc 20

<210> 10

<211> 31000

<212> DNA

<213> Homo sapiens

<220>

<400> 10

attccagcat cctggtagaa gagtggaact gacctggagg acagcaaagg agcccacgtg 60

gcgtaagcc tgaagggtgca gtattcccat tcaacagaga ctgctgctga cagtgggagg 120

gggagtagtt tgcttggccc gtggttgagg aattcgttgg cggaggaggg atcctgagaa 180

ggtactagaa atactgaacc cagtccctgca gaagctactt aacttcacag aaaactaggg 240

cgtcggtgcc ctccagcacg tccaaagggtg tccccgacaa ggccagcagc tctacctgtt 300

tcttctccct cctcttactc tgccctgtcca aaggcacgac actagtccac gtcacctttc 360

tttgggcgca aagggtatgg atcctcaatg ccaacctcgt ccccccccc cccacacac 420

acacacacac acacacatac aagcccacca tcgcgcaaga aactcgacc cgtctacctc 480

ggttcagtgg aaggcattca tttgcacaac gctgtgcgca tgcccggaag ccttaagtgt 540

ttcagcctcc gacaggggag taattaaagg ggcaggcgcc cacgtaaaga cgcgccctac 600

gcaatacgtg gagtttctca ctttgtggag ccgcctggag gcgccaagat gaagcccatc 660

ggggaagtgg ggcggggcca acccggcagg ccccgcccct tcgcggcgct tcctagtctg 720

gctggttctt ctgtcgccgg cttcagcagc ccgcgcccgg gcaggtaagg cattccccgc 780

cttaatgcct cgggtcaaacg tcgtccgagg tccgcttcca ggccctggca cctctgcccc 840

agggttgctg gggtcggctg cggaaacggaa atctctttat catccgtgag aaaagttaga 900

gaggaagtgg gttcctttga cagagcttac tccagaagaa agacttgggt aggtccccgg 960

taatcaagaa gttcgtcggc cctgaggaag agcctgaggc ttggaggggac gtcaccagc 1020

tgttctgggg cgcagtcgga cctcaagttt ctcacctctc aggcctgga ttccgtgaac 1080

ccgactgcga aatgaagtct cttttcccca gaaccaggaa gtcaggccag cacctgacag 1140

atgcctgtgg cgccagcagc tcgctgagcc tccacacgat tgtaattcgc ttgtttctcc 1200

ggcctccaag cttccaggga attagaaagt ggaggcggct tatgggatca gatgagtaca 1260

ttcgcagatg ggggaaaaaa gtgtacctgt cagcttgaa gccagagga gacaaagga 1320

taaaagcaaa ccctgtccag gtgcggtggc tcatgcttgt aatcccagca ctttgggagg 1380

cccaggcggg cggatcgctt gaggtcagga gttcgagacc agcctggcca acatggtgaa 1440

accocgtgtct ctactaagat acaaaattta gctggcatag tggcgggccc ccgtaatgcc 1500
agctactccg ggggctgagg caggagaatc gcttgaaccc aggaggcaga ggttgcagtg 1560
agccgagatc gtgccactac actccagcct gggcaacaga gggagactcc gtctcaaaag 1620
aaaaaaaaaa aaaaaaaagc aaaccctgag aaagcgagaa agcactttga gcatttagaa 1680
tgggaacaaa cggcgctatt agtgcttcaa aaacgtaact gcccaaggcc cctggtaagc 1740
agaggatact aaaatctgcc ttttcccttt cttggagtgg tccttgaaga gggctatcgg 1800
gaagttcaaa tttcttctgt tttcctgcag agcgttcatt tgaagtttgc tttctacctc 1860
tccaaactga gcttctgata ttccaggcca aacttgctcc ttctcagcc ttctctatct 1920
cagtaaatat cggtaaaactt ggttatgctg tcttctcact ctccacatca gcccatagct 1980
ttctattggc tccacctcag aatcctagga cttctgcact tgctctgcta cgaccctggc 2040
cccagcctgc agcaogtccc acctggctta ttggtaggct ctctaaatg cttctttctt 2100
gcttctctgt ccttactcca acatccttta ttcccagcag ccccagtatc cttctaagac 2160
atgagtacag agcatgtctc ttttctcaaa acttccagca tctacatctc acccagagga 2220
taaacaaggt caatacaaaa actctaaggc ccttgaagac ctggcccctt tacctttctg 2280
acctcatctg ctacttgccct gttccctcct ctgcagtctc accatgtcct tgctgttgcc 2340
cagtgggccc agttccaacc tttgcattta catttcgttt tgtccacaaa cttttccac 2400
agatagtcca gtggctggct ttctccttct ttttgacctt taatcaaagc gcaccttctt 2460
attgaggctt cctgtgacta ccatatztat aattacaacc ttttctctat ttctgtttgt 2520
tctttcattt cacttggcac catctaacat actatagagt ttatttattt gcttattgcc 2580
ttccttcctc cattagaata taaactccat gaattcaggt tttggttttg ttttgttttt 2640
tttctotaat tttcctaagc cttcatctcc aggggcctgg agcagtatct agcacaaggt 2700
agacactcag taaatatttg gtgagtcaat ggtattatat tttgtttttt aaatatatgc 2760
aaagcatttt caattactta ctgttggtt ggatcaatgt tcaagaaaga tatgttaagt 2820
ttaatttatg gtacaccagc acatatocca gtttgtaata ggcagcccta tctaatttaa 2880
tagctctgga agccctcacc agcttttggt cataatgatg atgttaaagt atctaaaaat 2940
aaagattaat ggttatacag catattttgc taagaatata tggaagtaaa tagtgtattt 3000
tagaaaggat ataaatacat taaaattacc atggttgatc aaataagaca gccaccatct 3060
caaaagctta gctattttct cagacttggt atttgtacgc tttcagtaga tcctcactac 3120
tgttctggaa attgaacctt aatacagctt gcacatatct ttaaaaataa tgtgatataa 3180

gaataatrtc tgccaggaaa taataattgc tccttttttt ttttgaccac ctattattat 3240
gtgacaagat ttggcccttt gcatgtgttg attttaattc ttacagcaat tcttaggtat 3300
tactatctct gtttttcaga taaagaaaag caaggtgcag agaaatgtat ttactagctc 3360
tggtcacagt tggcttttgg tagaacttag gcctgtgcaa cctcaaagtt catcttccct 3420
ccccccctgc acaatctttt cctctgatca aagtgcataat gaagcagtct taaaacttta 3480
ttaacacatt tgacaagaat cttttgccct cctgttttgg gcacagtact ttgggaaggc 3540
attggattcc ataagcaggt gcttatttta atatatctct caatcaccac cccacctttt 3600
acacttgtct gaattgtctt acttgaaaat atgaatgggt tttctttact atgagagtta 3660
gtgggtatgag gaaacagtca tgaaggattt tacttatttg taagcaacat atcctgttcc 3720
acctgtgtta atcctaagtg tactaaagta aatcttgaag aatttgaaag agatgacact 3780
ttttaagttt ttttttgccc agtctgtttt cattattgaa aattcagaaa agcaaaaatga 3840
aaacaaaact atcataaaaag tcatcaccaa gggataacca cgttaatatt ttggcttatg 3900
ttcttgtaag tatgtgggtg ggagtggtat atgaatatac atttaaagta aaattttaat 3960
ttctgggatt ttgaacttct gggcacatgt gtggaggagt tgcagaaggc cttctttttc 4020
ctagtatccc tacacttatc acctctgctg gaagcctggg tgcaggagag ctaccaaatt 4080
tatccctggc ctttagatag tctgggtttt ttctcaatt ccttgtattt tgctctaatt 4140
tataagacta ctaagcagtt ctatcttgtc tagctttgtc aatcatgttg taccttacag 4200
tagagagaaa tcttagtaag gaaaaataac ttgaagagag caactgcaga gtctataggc 4260
aacctagata caatttctta attaaagtta aaaagtcatac agcctttcat tcaacacaaa 4320
tgttgtgagt cgttctgag gtgaatttgg ttgggtccta gctcaggaga gcacctatct 4380
aacagagaaa gcaaacacat aaacagtaaa acataataat tattataaca gaagaatgac 4440
taggtgacat tttggagcaa aagagatgac aatcagggga agctaacagt gccttttgaa 4500
ttgggcttga aggaataata ggcatttacc aggtggcctt taaggatgtt gggctattct 4560
tttacctggc cctagatgac ttcatctctt gtttcacaga aaaaataaca gtatgtggga 4620
actcttctcc cttcaaataat atctgtattt aaacctctgc ttccttcttt ctagattcaa 4680
aagaggtttc tctctattca aggtgattt cttcacccta tatgctattc attatcccat 4740
ttgtattgcc aaacagctta agtcttaaac ccttaacacc aacagttctc aaaagtctgg 4800
ttggagtccc taatgctctt tcaggagatt tgtaaggcca aatcatgtt tagttaaag 4860
attcattcag agtgcaatgg attttaatgt tccaagttaa tttatacagc ttcagattcc 4920
tcattgcagc ttaccttag gaagttattt cttgtcaaat tttagtgtag tatcaaagag 4980

aaatatctac tacttttgtga aaaggctatt aaaatacacc tttctttata actatatgtc 5040
tgtgtgaagc tggattttct tcatatacat caacccaaaac aacatgctgc agtgggttga 5100
gtgcagaagc agatacaaga attcagctct cttctattaa gttaaataatt gaagagatta 5160
gcaaaaacttt aaaacaatgt tgttcttctg tttttgagga aaaggtaatt attttattaa 5220
aatatataatg ctaacatcta tatttatgat tattttaaat gaattaataa gtaattttta 5280
agtttctcag catcaattta ttatggtaaa tattggtaga tataaccaac ataaacaaaa 5340
gccctttggg gtctacaata gtttttaaga gggtaaaagg tcctgagacc aaatttgaaa 5400
actgctactc tcttgcacgc tctcctccct ctctttccct acccctccac ccccattttc 5460
cctccoctca tactcctctc tctctctttc tccttccctt cataaccaga cttcttggaa 5520
aatctgtttt tgccattcac tcactttttc actatagctg ttccaagtc accaatggcc 5580
ttctgatttc agagtctaac atatccatca gtgtctatct gacctctctg agtcttttat 5640
accaacctgc caccttttct agaagtctct ggtatgctac catcctgggt ttttcttgc 5700
cttcatgact aagcagtatt ttttaaaatt cttctttcac atcattcctt tccattctaa 5760
gcattcctaa taggcaacct cattttattg ttgcttcaac tgtatatgtg tatatgctct 5820
taatttgtat gagggcattt caaaaagttt gtggaaaaat tgaattaaaa gataaaaaat 5880
gtaaactttg gctcagcact atggctcata cctgtaatcc cagcactttg ggaggccgag 5940
gtgggcagat cacctgaggt caggagtcca agaccagcct ggtcaacatg gcgaaacccc 6000
atctctacca aaaatacaag aattagctgg gtgtgggtggg gcacacctgt aatcccagct 6060
acttgggagg ctgaggcagg agaatcgctt aaacctggga ggcagaggtt gcagtaagcc 6120
aaaattgcac cactgcacta cagcctggat gatagagcaa gactccaaaa acaaaaacaa 6180
aactgtaaac tttatttctg aacataagcc catgaagttc aagacacttt tgtaagcgat 6240
aacaccagcc atttgggcca tccctaaaga actgagagtc ctgggaattt aaccacatca 6300
atgaagtctt ttttacatta ttaactgaag aaaaatgggt gctctatgaa gatttttcaa 6360
gattgggaaa caaaaagaag tcagaaggag acaaatcagg actgtaaggt ggatgcctaa 6420
tgatttccag ttgaaactca cagaattgcc cttgtttgat gagagaaaag agcaggggca 6480
ttgtgatgga gaaggactct gtggtgaagc tttcccaggt gtttttctgc taacatttg 6540
actaaagctg taacagaaaa gctttagttt agaaacactt agaaaagcta ctgtttctca 6600
aaaccctctc taataagcag atgttaacct tctttttccc tccagaaagt cagcaagcaa 6660
aataccttga gcatcccaa aacctgcagc catgcccttg cttttgactg tctgcttttg 6720

ctttaattgg accactgcca ctttggggga gccattgctt tgattgtgct ttaggatcat 6780
actcgtaaag tcatgtttca tctcctgtta caatTTTTTg aggaaatggt ttaggatcctt 6840
catcccactt gtttaaagtt tccattgaaa gctctgctcg catctgcagc tgatctgggt 6900
acaacagttt gggcagccat caagtggaaa gtttgcTcaa ctttaatttt cagtcagaat 6960
tgtataggct gaactagttg tctatggtgt tCactgttgg ttctgctgTt aattataggt 7020
actcttcatt tagggcacia acgagatgaa tgTTTTctc gaaaatcgat gtggatggTc 7080
tgccgcagca ggcttcatct tcaacatcat ctcatccctt aaaatgagtt actgatttgt 7140
aactgctga ttttctgggg gcattgtccc cataaataaa gcatcagtga tttcaccatt 7200
cttccaacca acctccacca taaatgtggt gtttgttctt gctttagttt tagcagaatt 7260
catgttgctg tggtagaggt tcttttcaaa ctgtcttatt cttctcagtg tctcaactag 7320
atcctgttta gacatgaatg ataagttagt atgagtttat tttggaaata cttttgcaa 7380
aaaaaaaaa attgaaatcc atgcatgggt tttgcacgat atgcattttc catgaaTct 7440
ttgaaaacc cttatatatg agaatactta cataatgagc attctttgct gaaacctctc 7500
tctaagctc cagaccacat atataagtac ttggatgaa ggctcctaga tagtgccaaa 7560
ggtagatatt tcacatttaa catgagccaa aaaaaaaaaa aaaacttcat tctatctccc 7620
tttcttctaa tcgctcatct ctagtgaaat tgTTTTgatt tggcctccta aatgttttga 7680
cgTTTTacct ctccattccc ctacctata ctttttccaa gctctccttt ctgctcacct 7740
ggattattgc agaagTTTTc aaactaattt tcttgcctca atatacatcc ctttgatttt 7800
tatccagagg aatccactta accatataag attatTTaac atccctccta aaaatcctaa 7860
agtggctctc ctgtaacttt tagtttataa gttcaaactc tttgtcttag cacatgaggt 7920
acatgatgat ctggccctgc tacctttcca ccctcatctc ttctcactta atcacaggca 7980
tcctgtggTc cagccatagc aagctctttg cctgagacac acatgctctt accttttccc 8040
atgctattcc ctctgcctgg aattttcttt tgggcttata ctcatccct ttcattcatc 8100
cttccaacct cctactccaa ccaaactgtt cttgatatcc ctattccaga ctgcctctgc 8160
ttgatgcccc ttttctgggc ccataacat tcttgcagca tttatcactc cctgtcactg 8220
tttgtatccc ctggTatact gaacagcttg gaaacaaaga ttttctcttc tctgtgctcc 8280
cacttgTtag taciaagact gacataaata ttttctgaat gtattcatga atgaaaggaa 8340
aggaatggTc tttctgaaat agagaatggc atgaacaatg cattgtgtta taaaagcaga 8400
gttgtatggc atgactTTTTg gagtcagaca taattgtaag tcctggatct ctcaTtacc 8460
acctttgtaa aatagagatg ataaaactgt actgggaaat aatttagttg ttccaaaagt 8520

aaatactgaa caataccatt atctacagta tgcaagtcac taggtgtata ttcatgaata 8580
aaattagcat ggtcctgaac tcatggatct taaaatcgaa tatgggagac agtaaacagg 8640
taaacaaata agtatatgat taaaaatata ttgtgcagtg atgaaagata attaggtgaa 8700
gggcagctag acctatntag atgtgggtcaa agaagtccaa agagaagaag tgaataactaa 8760
catagaagcc totgatacag agaagctttt cgctttttac tctttttgat agtttttagaa 8820
tgttgttcca tatgaacaca ttatctactc taaaattgta tttttaaaat atatataagg 8880
aagcaaaaat aattgacaga agaatatctg agtctatctg tcttaattgg aaggaaagat 8940
catttcctat tgagaatgag ggacacgcag aacaaataaa atccacatag aaagaagtaa 9000
aggtatgaaa tagtcactaa ggactatgag agaagacaag tggaaggatt aacggcagtg 9060
ttgaaggccc tgtggaaagg gggttacat aaatttgtat tcacatcagc aacctggtg 9120
tacaagtaaa gttgttagat ttatccctgg ctgcagctct ccaggatgga gtagcagaag 9180
taaaggggga ttgcagaaat aaagattctg tctagagagt agttgaagtg gtgaagcaca 9240
ggttttggcg tagtagcaaa tgaattgtgg tcaggcagat ctcatggact caaaaaatgt 9300
gcagaggtca aggaagagca aagtatgtga aaagtgggtg ggtggacat atctgttaat 9360
gctacacaaa tcatttcagt acgttttttt tttttacttc taccaaagaa tgttgcataa 9420
attttaattg taaataagta ctaattgaga aacataaatt gaaaggtaaa tgtttttatt 9480
aatagatttg atttctgaat atgctaattc tgatagtttg ttgatattac tcttaagaat 9540
ttttctggta agttttttgc ccctttctat aaattttact aggaaaatat cagtgtcaga 9600
aaaaagtct ccgattttta ttgctattaa tgggtggaaa aaggaagcaa acccagagga 9660
taattggaca aatgaattac aattttaatt tcatcttagg atatcaatga cttgactttg 9720
gaatgtttct ttattatggg ataactcagtt gctgacattt catatgatag agcaaagtgt 9780
gatataataag cttacagatg tcatctctaa aagtactgta aaattttaat gagatttttc 9840
catattttag gaatagaaga tgaacaaacc cataacacca tcaacatag tgcgctgcct 9900
caatgttgga ctaattagga agctgtcaga ttttattgat cctcaagaag gatggaagaa 9960
gtagctgta gctattaaaa aaccatctgg tgatgataga tacaatcagt ttcacataag 10020
gtaacagata aaattctttg ttttttaaa ttcttacatc acaatatgga atgattaatt 10080
ggataagta ctgtctaattg tggcagttta gaaagtaaac ctctccacta tggaggagac 10140
attggtagta ggtaacctct ttttgattgt tccttttgca gttttgtctt ctttgtccca 10200
tatgtcaata gggctaatac tgtgatctct caaataagtt ttttgggtta gaatcaagat 10260

tcccaccttt gctgtgggtc cctgtttggc aagccaagtg aagttcactc tgcctcacac 10320
ttctcatttc cagattgaag gaaaataaga atgttacaag aatgtctggt aaaaatctag 10380
agcttaaatt tcaaagtgtg tgaatctcta acacactcta acaatgtatt ttgggacttt 10440
ttcaatttga cgttttaaag ttgttgctta caagtgtaca cttctgtgca cagaattctt 10500
aaatcttagc taggatttct cagtgtgtc ttcagataat ttcaattttt cttttttact 10560
catctctgct caatttagtg taaaatatgc atgcatatth tattttttgtc agtcattgca 10620
gttaaaacct atgtcaaat gtttctttgg tagcaaaagg aaaagaagac tatttgttat 10680
tttcagcatt aatctgggaa agaacctaaa agacttggct cccttgcagg aaatatttcc 10740
tcataagaag agcgaacatg ttaaattact tgaagaagc ttaagttaa actgtgttaa 10800
gccaatgttt aaagataatt ttttgatatt actgttaaaa acttatggaa gataattggg 10860
gcagtgggtg caggaagata aagctcttct cttttttttt tcttttttcc aagactcatt 10920
cctgtgacag aaaaagcttt tctttacaga ataatactg caaataattg tgaaaagaat 10980
ggtagaatga gaaaaatacc tatttgcaac caccaatgag taactgattc atgcaaggct 11040
catcactgaa ctaaaaagca taaggtgaaa ggtggttgtt ctctgtcatg tgagcaggat 11100
atgcacaaga ttttaaagta ttataggctg agcacagtgg ctcatgctg taatcctaac 11160
acttgtggag accgaggtgg gcagattgct tgagcccagg agttcaagac cagcctggac 11220
aacatggtaa aaccccatca ctacaaaaaa ttagctgggt gtggtagtgc acgcctgtag 11280
tcccagctac ttgggaggtt gaggtgggag gatcacctga gccaggaag tcaaggctgc 11340
agtgagccat gatcatgcca ctgcactcca gcctgagtga cagagcaaga cccctcaat 11400
agatagagaa aaagtattgt acacaaatta cttgtgaagt atgaaagaga aggttacatt 11460
tttaatgagg aaatctgaag tacacttctt tcaccaagtg atcaaacttt agcaccaata 11520
ataatggtac aaaatgctaa tatgtacccc tggatgtgat acaatgggaa aggcacatta 11580
tacctatgta gtatttctac caaaatgctt aatctgaatt taatcatgag aaagtagtca 11640
gataaatcta ggctgttaga ccatttctat aagacatctg gccctattc taaaacacac 11700
acaccacag acacagattg aaagagagaa tgcacaagag cccagatatg gcacaatttt 11760
atcaattatt caattgggtg atctagggtg atggttgttc attctatcat tttttcaact 11820
tttctgtaag ttgaaattt ttaaaaataa aatctttaga ggaaaacag aaagaacaa 11880
tattggtaa atttttaaat gttacttttc aaaactaatt tttgatttta aagaaatgga 11940
aagaaatggg ttgaaagccg actttgtgat ttttttcaat ttaattgaat tacagttaag 12000
atcaaaggaa taatttctta agtttttgtg tgttctgttt ctttaattaa gcaggtgcat 12060

ttattaaaaa ttatatagct atcagagtct taagtcatgt gggaaagtaat attgctagta 12120
gattaagaat acactctttt gtcaaatgag atctgaattt gtacctcaac ttggatacat 12180
catattggct gtgtggctat gggctggta ctctacctct ccaagcctgg ctttttccact 12240
agtaaattca tatgataatc ataatatctt tgaacatgct catgtgtgtg catcctataa 12300
cccactaagt aaatatccag taagtaccca gtaaattatt atcatcatct ttggcatctt 12360
ttttcctcct atctttcctg tctctctgta attcttgctg gttctcttgt tgtccctttt 12420
ctttaatggt gtaggagatg ctgtgggcaa aatgggcagg ggagatcaac acctattttc 12480
atatgacctc agtttataaa tgctattggt tgaatgtgtc cccaaaagtt catgtgttag 12540
aaacttaata ccattgcaa cagggttgat aggtgggacc tttaaaggtt agttaggcca 12600
tgagagcttt gcccttatga gtggattaat gtcgtcatgg caggagtggg ttagttatct 12660
agggagtggg tttctgataa aaggataagt taggccctat tttctttttt ctctctctct 12720
cttttttttt ctctttcaca tgcacgtgtg catgctcttt tgcccttctg cccttccacc 12780
atgagatgat gtagcaagaa ggcctaacc agatgcagcc tctcgatctt ggacttccca 12840
gcctccagaa ccgtgagcca aattaactat tatttataaa ttaccctgtc tgtggtattc 12900
tattattgca gcacaaaatg gactaagaca atagactaat tttgtttctt tgttacttac 12960
ttttaaggag atttgaagca ttacttcaaa ctggaaaaag tcccacttct gaattactgt 13020
ttgactgggg caccacaaat tgcacagttg gtgatcttgt ggatcttttg atccaaaatg 13080
aattttttgc tectgagagt cttttgctcc caggtaaact gattgtgacc aggggtgcca 13140
caattaggtt ggaaagacaa atggcagaaa tataaatggt tcttcttact ctctcttttt 13200
tctcatagta gatgaagctt acatttgaga gtccctttct ttcagcactc ctcaactctt 13260
taaaaagcag cacagacaaa ggacactgtg gactctgctg ctaaggatgat agaagcttcg 13320
taagagttag atagttttgt gccaacagga agtttagaag gaaagacttc atacttctgg 13380
cttaggctgt aagaaagtaa ttataatfff gagttcttcc tttttttcat cttcaacttc 13440
tacctgatg ggactctata atcataatff taaaaaattg tattctgatg tggtagtgtc 13500
tagttgcctt ccttataagc tttctttttt tcttttgatg cttttcacac cagtggttct 13560
caacctttat catgcataag aatcacttag gttctgggta aacatgtagt ttcttagggt 13620
ctgttccaaa tctatgtaat cagaattctg gaattggggt taaggctctat attcttaaac 13680
agggtcttca gtgcctctga tatagggtggc tcatggatca ctgtttgaaa aactgcctt 13740
actctcttta cccatcttca ttataactta cagattctta ctaggcagct tcttgtgtgt 13800

gctgtgagaa tatgagacca acctgtagaa actggaatga tattaaatga accaagtttc 13860
tagtttaact ttttcacaac cactttttct tactgaaaaa ccacttgtat cttacttcat 13920
ttgttagatg ctgttcccaa aactgctaata aactacctt ctaaagaagc tataacagtt 13980
cagcaaaaac agatgccttt ctgtgacaaa gacaggacat tgatgacacc tgtgcagaat 14040
cttgaacaaa gctatatgcc acctgactcc tcaagtcag aaaataaaag tttagaagtt 14100
agtatacac gtaagtaaca ttttcagtgc tttccactag ggatttgtca ttaagactac 14160
cagtgcttta aaagaaagct cttgctcttt tgtttgtgca gcaatcacag gcacactggc 14220
aatagctctt ttgtgagttg tttcctctg atatattaag aacctcttc attgtattaa 14280
tcagtgatta gaggcaatag gacatgcaaa caaccagaga agctatgaaa aaaaataagt 14340
aaatatttac ataacttggg aaggtcactt tttaaaataa aatatgtgac tagagttttg 14400
ggtaggtaga ctagaccac tcaaatgtgt gactagaatt ttgggtgggt gactacagtg 14460
ttcagagggg aggatcacca aacagagttc tcaagaaaaa cttataattt gtattttaga 14520
aatggtttta tcttctctca tcttgtctaa ttcaaatca taaatatttg catatatgtg 14580
aaatcttcaa atgagaaaat attataatta ttaaacgaa tttttaaagt ttaagcatgt 14640
ttttcttatt ttgacatagg ttttcacagt ttttcatttt atgaattgaa gaatgtcaca 14700
aataactttg atgaacgacc catttctgtt ggtggttaata aaatgggaga gggaggattt 14760
ggagttgtat ataaaggcta cgtaaataac acaactgtgg cagtgaagaa gcttgcagca 14820
gtaagttata ttttcaggaa taaaaagaaa gagttgcttc atagtgtgcc ctataatagg 14880
ttttaaagtt aatctttagg aaatacatta tttcagtag tattttcttt aaataaacat 14940
cattctaaat agtagttctt aaaattttta aatctgttga aactatgga ccctaccct 15000
agaaaaatgt ggttatgtcc atgtacacaa gatcatacat acaattttgg gagattcaca 15060
aactaaagtt gagaagccct gctccacata gtgcttcatg tgtaatcaag attaaaatgg 15120
taagatacca ggaaaactat ttttaaaaaa ccttgtttgc attacttttc ccatttaaaa 15180
ataaaattta ggaagtatac attaagaaaa aacattaaaa tttttttat ctgtgtcctt 15240
tgttgccatt tctactgagg tgtttacctt ttacttactt gttatattct aaatcttttg 15300
ttagagccta aaaatatcaa tagaatgcat atgtatgtga taaagatagt cactctttgt 15360
catatatgtt gcagggcttt tgtagggttt tttttccat ttgacaccta aattgcttat 15420
gttttctgtt tcctagaatt tttttcaaat tttttgtgt agtgaaattt cgtgttattt 15480
cattttgttt ccacaactca gtgaaattta tatatatttt tcattccact aaccagccat 15540
gggccttgag caaatcacta gtatttctag atctcagtta ttctgtagaa ttgggagaat 15600

aatacactat tgttttaaag aaagggagat gagccagctg atctcttgat ccctctgagc 15660
attaggaacc ttagaattgt gtagtattac attgtatatt ttatTTTTTc agatgggtga 15720
cattactact gaagaactga aacagcagtt tgatcaagaa ataaaagtaa tggcaaagta 15780
agtcttaatc tggcagtgcg gtgtagtgga aagaaaaaag acaaggagta aagaacctgg 15840
ttcactctag agtatgccat gaactagtga tgtttgcaca tatatgaaat taaaataaag 15900
tgtttagatt agagaaattg aaagtacttt tggctccata attctatgat tacatatact 15960
catatgcctg gagattaaat agcagtcatt tttatttatt tttttttttt gagacagagt 16020
ctcgcctctt tgcccaagct ggagtgcagt ggcattgatct cggctcactg caacctctgc 16080
ctcccagggt caagcgattc tcctgcctca gcctcccagag tagctggacc tacaggcgtg 16140
tgccaccatg cctggctaatt tttttctgt atttttagta gagacggggt ttcaccgtgt 16200
tagccaggat agtctcgatc tcctgacctt gtgatccacc tgcctcagcc tcccaaagag 16260
ctgggattac aggcgtgacc acgcctggcc gcagtcatta ttttaagtca caacaaagtt 16320
acgtgaaatt tcactatcat gtccttcatt atcatgtttt ccacttccag aattcatgag 16380
agatcaattg tagagatact taaaaacaac tatctataga gtaataggcc cttatatata 16440
aagtagcatc tatagagcaa ctagagtcatt gttatctgta aggatgctta gaaggcattg 16500
atcctgtgtg ctggaaaatt gttggacatc atcaggaaaa gtaggagggga caaagtagga 16560
aatgtattct aattattata gtgattaata gtgattgata tttgagtgtg tgcaacatgc 16620
cagacattgt gctatgtact ttatgtgcat tgtctgtcta acccttacia ctgttaciaac 16680
tgcccatttt gtttgttaac tgagatttag aaacattaaa tagcttgtcc aagatcacat 16740
agccagtaag tgttagagat gggattgaaa tccagatctg tttgattgag gagcctgaac 16800
tttagatctg aatcgtaatg ggatttacca aattataccc attaaaccca ctgctgtaaa 16860
tcaccaatca tagttaccta acctcaagta agaagaatag actagaattg atttatgcac 16920
ggtaaaaaatg atacatgctg aaaaaagatc aaactttttt tttcaggatt tgtcttcttc 16980
actcctaccc ctactctcca attctaaaag taacttgttc acagtaggtc atctggaaaa 17040
tacaggaaaa catttttaaa atatacaagc ctatatgcat gcaggactta atacgtagggt 17100
gatgggttga taggtgcagc aaaccactat ggtacacggt tacctgtatg acaaacctgc 17160
acattctgta catgtatccc ggaacttaat aatacaaac atctctcaaa tatatcactc 17220
agagataacc actgctaata tttttatata cttgcttcat gatgtgtgtg tagcattttc 17280
ccatgtcact aaaaataatt tgaaaacatt ttaaattatt gtacaacagt ctagcatatg 17340

tgaaaatcat attttattta accatagcta ggtttgtttc taattttcag tattaanaaat 17400
aaagcttctt taaagttctt tgcacataaa tctttaaatg cagttatact tactttatca 17460
tctagggtaa agtcctagaa aaagaattac taggtaggaa tgcatagaact ttttaaagac 17520
tttctaaata cacattcaaa tggcttttga gataagttac taccagcaat gtattacaat 17580
ttcaaacttt actggttcct taccttcatt taatatcatt actttttttt tttttttttt 17640
ttttttggag acggaatctc gctctgttgc ccaggctgga atgcagtggt gcgatcctgg 17700
ctcactgcaa gctccgctc ccaggttcac gccattctcc cgctcagcc tccagagtag 17760
ctgggactac aggcgaccgt caccatgccc ggctaatttt gtttttgtat ttttagtaga 17820
gacggggttt caccacgta gccaggatgg tttagatctg acgtcgtgat ctgcctgcct 17880
cggctgggaa tacaggcgtg agtcaactgc cccggcctat tattacaatt ttttaataag 17940
tgccagtctc tggatcttac tagcagaatg aacttggcca agttttttaa ctttotatcc 18000
gttaattttc tcatctataa aatgggacaa tagtaactac ttaagttatt gtgaggatta 18060
aataaaacag ataaatcact tgccacactg cctcgtttat agtataaatt cagcaaactg 18120
taactactca tgtttaaaat gtgccagttt ggtggaaaat gggacactta ctgttttaat 18180
ttgcacttct tcaaataatt atgagactag aagaaattga tataatgaac tgctagaatt 18240
gtggatcatt ttgaatgaac aaaatgaata agcacttaaa taataaaaca cttatcatgt 18300
gccagatagc tttctaggct gtcatgttta catttgcctc ttttaattgtc tcagcatcca 18360
tattacatac tgttttctgc atttagcaga taaaggagct gaggctcagt aaggctgtta 18420
acttgcggta ggtcacatat ccaggagggg gcagaactag actttaaatt taggatgtat 18480
aactctgagc cctgttttct ttcgtcttac cacagatgct gccactttat tagattgtag 18540
gtctctttag ttttattact ctcaagtata taaagacact caaattaggt tataaataat 18600
ccagtgtgat taaagatcaa aaacatgtaa attagatgtg attcaagatc atgtctgtaa 18660
atcagtgctt tgagtgtgtt agaaaattct tgacagtatt agcataaaaa ttcaacggac 18720
ttcttgaaat cttatttgtg ccatcattga aaattgggct aaagaaagtg agcatcagcc 18780
tagtccttaa ccaaacttat cacaagttcc aaaaatctga attatttagc ttttactgta 18840
aaatagaaag aggtttggcc agctgtagtg gctcacacct gtaatcccaa tactttggga 18900
ggccaaagtg gtaggagcac ttgaggccag cagtttgaga ccagtctggg aaacacagtg 18960
agacctcatc tttacaaaaa aatttaaaaa gtagctgagt gtggtgtaca ccagtagtcc 19020
aagctacttg ggaggatgag ttaggaggat cacttgagtc cagatggctg aggccgagc 19080
gagctatgat tacaccactg cactccagcc tgggcaacag agttagacct tgtctcaaaa 19140

aaaagaaaa gaggttttac ttctaagtgt gattaatact ggctgaaaag agaagtat 19200
gcagaaaatt ataaacagtt acaataaaca tatataaaca taaacatcaa gttagaaata 19260
atatataaca tactatTTTT ttgctctttt tttctattaa aactttgaat aacttccaac 19320
ctatagctga atatatatTTT attataataa ttttgcatga aaaattatTTT gtcacaggtg 19380
tcaacatgaa aacttagtag aactacttgg tttctcaagt gatggagatg acctctgctt 19440
agtatatggt tacatgccta atggttcatt gctagacaga ctctcttgct tggtaagcta 19500
tttgttcatc agattgTTTt gctTTTTgtt tatatgctgg aatattaata ttcattTTTgt 19560
tactggatta tttaaacc aaattcatttc atatttttcc tgctctatga aaattataca 19620
gttgatatgt caacaaatta tacagttgat atgtcaacaa ctgggttgcc attttattag 19680
gtggtaatct taaatgtaga attcagttaa gactatacta ggaagatgct tttatatatt 19740
tttatcttca aataacaaat gtaaaccacc atttgagctt attagatat tattttta 19800
aaactaaaaa tgtagtctta aattattaat gctataacat catcttcagt tgttgcttag 19860
aaaaatattc tgtgttatat attctctgta gattttttat tattctttct cattttta 19920
ttggtttatt tctttataag gatggctctc caccactttc ttggcacatg agatgcaaga 19980
ttgctcaggg tgcagctaat ggcacatcaatt ttctacatga aaatcatcat attcatagag 20040
atattaaaag gtaaagtcta ctgtttaaaa gtttttgga agctgtcttt aaagaataat 20100
ttgctgtcac tctattcacg attcatatgc aggtcttaac ctagagcatc tggacttttt 20160
tcccatcact ccatgaaagc tcttctcaat aaggacatct ataatctcct ttattccaag 20220
cccagtgacc tacttccagt actcatccta cttgaccttt tttgTTTgtt ttaatagtag 20280
tttattctgt tcttcttgac acttttgctt tatggaaatt agttgTTTTc ttagtttatg 20340
gatcactatt cttttctgaa tcaccttcat cctccatgag tacttttcag ttctctttct 20400
ttcttgtgta tttatctagt taaatattta cctactataa gtaacatatt catatataga 20460
gctcaacatc aatcaaataa gcacaaaata attatagaat gttaatatgt acagttcaca 20520
tatatatatt atgtattatt attatatagt atagtacata ggattaatat aaagaaggg 20580
ttccaagaaa gtgacattta agcagagatt caaagaatgg gcaggaaata agtagacaca 20640
atTTTgaggg aactgtagaa tggatgatgg ggagaagaac attctaggca gaaggaactg 20700
catttGtaag accagtgggt agacagaata tgggagcatg atgagttaca agaaatgaaa 20760
tatgctTTTgg gagccaagat aggaggatca tttgaggcca ggagtagcag acctgactat 20820
gttccctgaa caacatagca agacctcatc totacaaaaa aactagtcaa acgtggatt 20880

gcatgcctac agttccagac actagggcag ctgaggaggg aggaggcttg agcccaggag 20940
ttcaagggtg aagtgagcta tgatcgtgcc agtgctcccc agcctgggta ataaagtaag 21000
actctgtcta aaaaaaaaaatg aagtaaatgc agtatgactg agatgcagaa agttataggg 21060
tggtggattg gagataaaac tacagagaag tgtaggaagc taagtcctat aggacttaat 21120
agatcaagggt aaggctttta gttttatcac aagtgtaatg agatgccggt gaaaatttta 21180
agtaggaata tggcatcata ttttaagtttt gcaaagatga ttttaactct agtgtagaga 21240
acagtttgta agaagaggca gttagtaggc tatcataggt ggcagttcat aagaagagag 21300
gggagcagat tagaaagatt aaaatggcaa aacttggtga taccttggct atgcaggggt 21360
tcaggaagag cgaagtatca ggaatgactc ctggaataat ggcatccaga cttgcattct 21420
ggaatgatta tggcatgatt cacttatgta gaaaggactg aaaaaagact tggttaggtt 21480
ggggttagag agatggttag aaaatgtgaa ttctcttttg tacatgttgt gtttggggtg 21540
cttttaagat gtagtagtag agatggtaga cagtgtgata tacagatttg gagctggagg 21600
tctgagctga agatataaat gtgtattact tgcatacagg tgataattga agtgactgta 21660
ttcacttgag gggagatgta gattaagaaa aggtcacaag attaagtctt ggggaaccct 21720
aagacttaat ggccaaatag aagagatgat cacaaaaagg aatctaagga agtacacata 21780
cagagaaggg taagatgaaa acctggagag tgtggagtcc ctgaaaacga ggaaagaaag 21840
tgtttcccaa gagggagtag tcaaaaagag tcaagggtt taatcattgg attagcatca 21900
ccatggtctt tggtgacttt atggaaacct atttccaggt agtagttggt ttagaaatca 21960
cattgtagtt gagagagtga gagctagaga actgagatag gagtatagac aagtctttta 22020
agacatttta ctagtgaagg ggacagctag aaatagataa gtcattcagg aaactttttt 22080
taagctggca gagattaagc tgtgcttatg tgcctatagg aaggatccag attaaaaggg 22140
agaggatagt atggaagaga gaaatgagat gatgtgttca gtagggttac tgagaatgta 22200
gaaagcagct caagcagagg aactgacct agatgagaaa agaaacgcca tctcctatcc 22260
ttcatggatt ggtactttct tgtgttctta tatcattgtc tcttttctcc ctgtatgtc 22320
tttttttttt tttttttgag acaagttca ctcttggtgc ccaggctgga gtgcatggc 22380
gcggtcttg ctcactgcaa ccgtcgctc ccgggttcaa gagattctcg agcctcagcc 22440
tctgagtag cttggattac aggcatccac caccagctc tgctaatttt gtatttttag 22500
tagagatggg gtttctccat gttgatcagg ttggtcttga actcccagcc tcaggtgatc 22560
cgcccgcctt ggcctcccaa agtgctggga ttacaggcat tagccaccg cctggcctct 22620
ctctgtatgt ttttgctgag tgatattatg ttaattttaa ctgctactta tatgtgagt 22680

tcttttgaat ctgtattatt agcttccaat cctggatttt tagtatcccc gtaccaaaaa 22740
cacttcatct gggtatccca cagtagcctt taagtcagaa cattagaaat caaactcatc 22800
agctcttccc aacctaaagt cttcccacat ttcctttctg atccagttaa gtcactgttc 22860
attctgtctc tcaaactgaa cacgttctca ttctctatat catgctaatt acaaaattct 22920
gttgtgtgag cctcaaaatt ttcactttcc atttatccca tgttatgacc atttaaacc 22980
tcattatctc ccatttgctt tgttacaaac cttcgtagtc aacctaaatc cagcctatca 23040
gttctcatct attcactacc aactactag gtcatgggcc taaaagttaa gtctaataac 23100
atcagtttcc tactgtgaaa acttttagctg gtatcctggt gctcacagga taaaatgcaa 23160
attatctcca tgtttcttgg aactcgttc cttgaagttg accatgtata ccatatcatg 23220
ttttgtcac tttgtacctg ttactccatt tgctggagt gcctttcttt ccctcatata 23280
actattaact tctaataatc cttcaatact aagcacactt ttcacctct gtttgaaatt 23340
tttctactt tttccaggca catggttgtt ctctcagcat tttgtacctata atattaccat 23400
tgttttaaaa aattatttat gcctttgtca cagacaagat ttttaagactc cgccatgccc 23460
agcagtgagc ttaaatacta cacaacatag ggagttctca ataagattgt gttgaattaa 23520
agtatgaaag cacataatga attaattgag atattttggc catatTTTTT tctcttcaaa 23580
agccagatag tacagaacaa aagaataaag aaattcataa ttgagatact ttggttgga 23640
atgtgttttg aatgaattg atacatacta agaaaggaat taataaagggt gccttggaag 23700
actttgtagt aatttaccct ataataaatg atttgtggaa taatgaaaat atctcttact 23760
aatgtagaca gttttatttc atttttattc tcttaaatgt tttctcaag aaacataaaa 23820
aaagaggaca gttgcttctc tttagctata atctgaattc taggtttatt tctataaact 23880
ttaggctttt atgatgcata ctataaaacg ttacactctg taaatatgta tgtacatatg 23940
catgtatgca tatacatata tacatgcata catacgtaca tctagctaag gaactgtttg 24000
acttttttgg ggtgggaaaa acattttttt cttcaaactt tacatttttt tcagtgcaaa 24060
tatcttactg gatgaagctt ttactgctaa aatatctgac tttggccttg cacgggcttc 24120
tgagaagttt gccagacag tcatgactag cagaattgtg ggaacaacag cttatatggc 24180
accagaagct ttgcgtggag aaataacacc caaatctgat atttacagct ttggtgtgg 24240
aagttccgta tacataatta ttaaaaataa tcattctgct ataattgtga aatgaagag 24300
aatattttt aatacaccca tcttgtttat ctctcttatg taacaatata agtttttcac 24360
attaacctct acttatacca gaaagtttat aaaaacttct tccaatgtgt aaaacttttg 24420

agttgatttt tgaatgact acaagaaatg attttcttgc catctgtctt tatgaattag 24480
catgacacct aatgcattaa atgaatcatg tcatcacagta tatacttttag tctcacattt 24540
tttgacattt actaatctac ctctctattc ccaataactt ttaagataga gcaagagata 24600
gggtgtgggtg gttgaagaag ccgggggctt tttcttcatt ttatTTTTTc atacgtgttt 24660
catatatgga aatttaactt catagcactg aaaataaggc atagggctat agaatatgcc 24720
taatgttttag aaaataatat tctcctaaga gccagagtaa attattcttg tgaagaatcc 24780
ccaagaggca ggaaaaggat tgggagtcct tagaacatat gttgagtggg gctgactcct 24840
tagtaagttg ccaatagaaa agagctgctt atggatcatg tgggggtccaa gatgtccctg 24900
aaggagggat ggtgagaaat gcccaaagct catgttcttt ccactttaat atctaaaaga 24960
catagctttg tatagaatga ttaatcaaat tttgtaagca agattcatat tcattctagt 25020
ttttgtcttg agtattatat atataattta ttaatatcat gtataatata tatatTTTTT 25080
agagagacaa ggtctcactc agtcaccag actgaagtgc agtgggtggaa ttatggctcg 25140
ctgtagcctc aacctcccag gttctagtga tcctctcact tcagcctccc aagtagctgg 25200
gactacaggt gtgtgccacc acaccagcc aatgtTTTTT taaaattTTTT tgtagagaca 25260
gggtctcact gttgccagc ctggctcgtga agtccggggc tcaagttatc ctcccacatc 25320
tgtctcccaa agtgctggga ttacaggcgt tagccactgt gccagccta tcttgtgtat 25380
tatattaatg atTTTTTTTg tcttcatagg ttttactaga aataataact ggacttccag 25440
ctgtggatga acaccgtgaa cctcagttat tggtaaataa aatattcatt ttcttcaatc 25500
ctTTTTTctc tgctTTTTgta gtctaaattt atatggataa atttgacatc taaaaataac 25560
atTTTctatt ggcaatcttt ccattggcta ttacacgaca gcatatggaa aaagcattaa 25620
atTTTTcat tctcagctct cccatgaagt atttgtgaag cctagaaaag tcacttatat 25680
attctttggt ctgatttctt tgtgggtaaa atgagaatag aactaaatga tttctgaggt 25740
cctcttttagc ttatgtttat tttgtgtcac ttttggggtg gctactgtta ataacagtgt 25800
ttccagcatt ttcagagggg ggaaaaaaaa ggcttcatt caaatttctt tcagaataaaa 25860
attgctagta aaagttatca cttatttcta tcatggggta ttttatttac aaagttacaa 25920
taaaaaacat accaaatttg ctattgttag gaaaaattca tatcatgttt ttcttttgct 25980
ctcacaccac aacaacaatc aacacagaag atttctgtga ccaaatgcga ctgggaaggg 26040
tttctcccca ccaacaagca ggcaatctac tttgcagcgg aagtcagctg gatctcctcc 26100
aatttaattc tgacactaca ttcttggaaca tagcatcaga gtcagatccc acaggttaag 26160
agctcagtc ccaagactgc catcccacca gacaccagtt gtaagtttag gcctccagaa 26220

cttctgacca actgccttca agttggagtt cctatgacct cctctgtggg tatggagggc 26280
tcaagattga acccagattg attcactaag ttgcagcata tactttacct aagttttcca 26340
tcgttttgcc taagactttt tccttcaaaa tatataaaag cctcaaaact ttggccttgag 26400
taagaatgaa atgcagaaaa acataagctg tatctcagtc acaagttttg catttcatca 26460
tataatagtg aaaatcataa atctttggag aaggttttac tctttcacco attcttaagg 26520
tttgactgtg atgtgactct aaagtaataa gattttcata caagcacagt ttttaaaaaa 26580
gtgatcatct gaagaaccgt tatgtcacta gattctgccg ttgtcaaaca ttttcagaat 26640
ttatctttga aagtggtttt gaagagttaa catcctacct ctcatgaata gtttttttca 26700
tctgtattag cttcctagga ctgttataaa aatataccaa aaactgtgtg acttaaagca 26760
acagaaattt atttgcctcat agttctggag gccagagtct gaaatcaagg tatgggcaag 26820
actatgctct ctctggagac tccagagaga atcttgcctt gctcattgga tgtagggctt 26880
acctccagga ttagccataa tctggaatga tctcatctca ggataactta ttacacctaa 26940
aagggccctt cttttcctaa taagatcaca ttcacagggt ccaggaatta tgacatagac 27000
atgttttggg ggaccatgat tcaccaact ccacgatctt attcctacac ttctctccct 27060
gttaattttt ttttatcatg gcctgctccc tattttttcc cctccttttt tctctttgat 27120
tcaacaacct tagtaccat agagaaattt gttgggtaca gggaggagga gtcagggtcc 27180
aaaagttgga attgcgctat tagaagcaca tcaagaagca gaccacctg tgctgcccac 27240
ctagtaaagtg ctgctaagtg gaacatataa gcaattagct cataacatag atataaaaat 27300
tggaaggcaa atgggtggaa gtatctagga ggtataaata attaaatata catttgtata 27360
tatatatgtg tgtatatata tatatttata tatatatata tacacatacc cagccatgtg 27420
ttgcctaagtg acaggaatac attccgagaa atgtgttgtt aggcagtttc attgtgtgaa 27480
cctcatagag tgtacttaca caaacctaga tgatatgacc tacgacatac ctgggctata 27540
tggtattgct cctaggctac aaacgtatgc tatacagcat aatactctcc tgaatactgt 27600
aagcaattgt aacacactgg taagtgtttg tgtatgtaaa cataaaagag gtacagtaaa 27660
aatacggatata tataatttta tgggaccacc atctatatgt gatctgtcat tgaccaattg 27720
accaaaatgt ctttttgtgg tggatggctg tacttattta cataggattt gtcctgtatg 27780
tcttcaattt tttttttttt ttttgagaca gagtctcgct gtgtcaccta ggctggagtg 27840
gcgtgatctc ggctcagtgc aacctctgcc tcctgggttc aaacgattct cctgccagcc 27900
ttccaagtag ctaagattac aggcattgcac caccatgcct ggctaatttt ttgtattttt 27960

agtagaaaca gggtttcacc atggtgccca ggctgggtct tcaattctta ataaggtttt 28020
aagataagat agtaaaatga gagcacatgt tattacaaag tagattaatt taaataatta 28080
aaatataat tctatcttct ttcactatga tttgaagctc ttaaagtttt aacactcatt 28140
ttaaagctag atattaaaga agaaattgaa gatgaagaaa agacaattga agattatatt 28200
gataaaaaga tgaatgatgc tgattccact tcagttgaag ctatgtactc tgttgctagt 28260
caatgtctgc atgaaaagaa aaataagaga ccagacatta agaaggtatg cattttttat 28320
acttatataa aaagtgaag gggtaggggt catcatatct tccagagtgt atattttaaa 28380
gcaactgtat aatgtgggtc tttgtttttt ttctttcttt ttaaagggtt caacagctgc 28440
tgcaagagat gacagcttct taaaacttta ttggaaaaga ctcttgactt tttatataca 28500
cctatctcaa ccattttttt aactgatttt tttcctaaat attcttcttt accttaaca 28560
aggcataggc tggtgcagga cagtgggtat taaagcatgg gttgaacttc caaaatataa 28620
aaatagagcc accatatcaa cacttagccc taccattag tatcaccccc agttcttaca 28680
gtaatccctg agaaatctcc ttcaagcatc accaaacaca gtttgaaaat tacagggtta 28740
gcaaaaagag cctgggctgt atgtagggtg gaaacactct gatctgaagc ccagctgact 28800
ccactactaa tttgctgtaa agctttggac atacacttag ctgctgtgag ccactaataa 28860
cattgggcta atatctgctg tgcttctctg acaggtagtc atgaaaatca aatgatgcaa 28920
aatatataca agcactttgt aaattgtaa atgatacaaa atttaaagtt tatagagcca 28980
gttacaaaat cctattagtc atatatttat agattgtgtt cacagcaatc atttaaccac 29040
aaataaaata tcccttgatg atactgcat aatgatatgt ccattattag attatgttac 29100
atgacaaagt tgaaggaatt tggcagatgc agttaagggt cctaaacaac tcactttgag 29160
actgttgaaa gggcctgacc taatcaagtg aaccctgca agaagaattc tccttgtaag 29220
ccttgagaa gtatgtgaga gggccacatt ggctaaaacc taaagggtggc ctctaggaga 29280
tgagacctac cttccagttg tcagcaagca ggaaaaaaaa attgggacct cagttgcaac 29340
cacaaggaac tgaattctgc caaaaatctg agtcagctta gaagagtact ccaagcttca 29400
gatgataacc acagcctggg ctgacacctg gatttcagct ttgcatgatc ctgagatga 29460
gaatctatct gttctgtgct ggacttctaa tatatagaac tgtgagataa tgggtcacat 29520
tggttgatg tgggtgctca tacctgtaa tcccagcact ttgggaggcc gaggcaggca 29580
gatcacctga ggtcaagagt tcaagaccgg cctggccaac atgggtgaaac cccgtctcta 29640
ctaaaaatac aaaaattaga cgagcgtggg ggtggacacc tgtagtccca gctacttggg 29700
aggctgaggc aggagactag ctggaaccag ggaggtagag gttgcagtga gctgagatcg 29760

tgccactgca ctccagcctg ggtgacagag tgagactcca tcataaataa ataaataaat 29820
 aaatgggtca cattaagcct ttaagtttgt ggtaatttat tattcagtaa tagaaaacaa 29880
 atacagatac tctcccatga tgtttttccc atgatgattt cccatgatat ttacaggttt 29940
 tgcccacatt tgaggggtat gtggaaatta tacagagcat gtacagcggg aggcttatag 30000
 tgtacgtact gaaatgtggg gttggagccc caacacagag accccagcag gacactgcct 30060
 agtagagcta tgggaagggg gctgccaccc tccagacttg agaattgtag agccaccagc 30120
 agcttgcaact ctgagcttgg aaaagccaca ggcactcaac ttcaaccatg agggaagcca 30180
 cgcaccctgc aaagccacag gagtggagct gccacgggcc tcgagggccc accccttgca 30240
 ccagtgtgcc aggatgtggg acatggaatc aaggaatatg ttagggcttt tttttttttt 30300
 tttttgagac ggagtcttgc tctgttggcc aggctggagt gcagtggcac gatctcggct 30360
 cactgcaagc tctgcctccc aggttcacgc cattcccctg cctcagcctc cccagtagct 30420
 gggactacag gtgcccgcca ccatgcccag ctaatttttt ttgtattttt agtagagatg 30480
 gggtttcaact gtgttagcca ggatggctct gatctcctga cctcgtgatc acccgcctcg 30540
 gcctcccaaa gtgccagtat ttaaagtta atgtcttccc agctggggtt cagacttgcc 30600
 aggatcctgt tgcccctttc tttagccaat ttctcccttt tgggacaaga atgttttact 30660
 tattgcctgt accaccaact gtatcttggg aataaataac ttatatttta tttcagagggc 30720
 tcataggcgg caggaactta ccttgagtct caaatgagac ttaggacttt tgagtgatgc 30780
 tagaatgagt taagactttg ggaagggatg attatatttt gcaatgtgag aaagacatta 30840
 gatttggggg gctgggggta gaatgacatt gtttagatgt ttgtctcctt caaatttcat 30900
 gtttaaagt aatccccagt gttgggggtg gaggtggggc ctgatgggaa gtgtttgggt 30960
 catggtggat gatccctcat gaatggctta gagccactgg 31000

<210> 11

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 11

tcttctattc ctgcccgggc

20

<210> 12

<211> 20

<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 12
gttcaccttc tattcctgcc 20

<210> 13
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 13
acatatggtg atggtggtat 20

<210> 14
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 14
ttagtccaac attgaggcag 20

<210> 15
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 15
gcttcctaatt tagtccaaca 20

<210> 16
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 16
ccttcttgag gatcaataaa 20

<210> 17
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 17
ttaatagcta cagctaactt 20

<210> 18
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 18
ctgattgtat ctatcatcac 20

<210> 19
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 19
attcagaagt gggacttttt 20

<210> 20
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 20
aacagtaatt cagaagtggg 20

<210> 21
<211> 20
<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 21

gtggtgcccc agtcaaacag

20

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 22

tgtgcaattt gtggtgcccc

20

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23

ggaacagcat ctgggagcaa

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 24

gaaggtagtg tattagcagt

20

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 25

gctgaactgt tatagcttct 20

<210> 26
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 26
cctgtctttg tcacagaaag 20

<210> 27
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 27
aatgtcctgt ctttgtcaca 20

<210> 28
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 28
ggtgtcatca atgtcctgtc 20

<210> 29
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 29
tgtgacattc ttcaattcat 20

<210> 30
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 30

gcctttatat acaactccaa

20

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 31

attgctgcaa gcttcttcac

20

<210> 32

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 32

cttcagtagt aatgtcaacc

20

<210> 33

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 33

aagtagttct actaagtttt

20

<210> 34

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 34

tcatctccat cacttgagaa

20

<210> 35
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 35
gcaatgaacc attaggcatg 20

<210> 36
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 36
catgtgccaa gaaagtgggtg 20

<210> 37
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 37
gcaccctgag caatcttgca 20

<210> 38
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 38
cattagctgc accctgagca 20

<210> 39
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 39

tcatgtagaa aattgatgcc

20

<210> 40

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 40

tgatgatttt catgtagaaa

20

<210> 41

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 41

aatatctcta tgaatgat

20

<210> 42

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 42

agatatttta gcagtaaaag

20

<210> 43

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 43

ttcccacaat tctgctagtc

20

<210> 44
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 44
tgttgttccc acaattctgc 20

<210> 45
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 45
acgcaaagct tctggtgcca 20

<210> 46
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 46
acagctggaa gtccagttat 20

<210> 47
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 47
agcaataact gaggttcacg 20

<210> 48
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 48

aatatctagc aataactgag

20

<210> 49

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 49

taatatctag caataactga

20

<210> 50

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 50

tctcttgag cagctggtga

20

<210> 51

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 51

aataaagttt taagaagctg

20

<210> 52

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 52

atgctttaat aaccactgtc

20

<210> 53
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 53
gaagttcaac ccatgcttta 20

<210> 54
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 54
atttctcagg gattactgta 20

<210> 55
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 55
aaactgtggt tggtgatgct 20

<210> 56
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 56
tacagcccag gctctttttg 20

<210> 57
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 57
caccctacat acagcccagg 20

<210> 58
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 58
tcagatcaga gtgtttccac 20

<210> 59
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 59
tagtggagtc agctgggctt 20

<210> 60
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 60
ttattagtagg ctcacagcag 20

<210> 61
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 61
agcagatatt agcccaatgt 20

<210> 62

<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 62
acctgtcaga gaagcacagc 20

<210> 63
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 63
tcatgactac ctgtcagaga 20

<210> 64
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 64
atttacaag tgcttgata 20

<210> 65
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 65
tgactaatag gattttgtaa 20

<210> 66
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 66
taaatgattg ctgtgaacac 20

<210> 67
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 67
ccttcaactt tgtcatgtaa 20

<210> 68
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 68
accttaactg catctgccaa 20

<210> 69
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 69
tggattaggt caggcccttt 20

<210> 70
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 70
ctcacatact tcttcaaggc 20

<210> 71
<211> 20

<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 71
gttttagcca atgtggcct 20

<210> 72
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 72
cctttagggt ttagccaatg 20

<210> 73
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 73
ggccaccttt aggttttagc 20

<210> 74
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 74
ctcatctcct agaggccacc 20

<210> 75
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 75
ctgacaactg gaaggtaggt 20

<210> 76
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 76
tttcctgctt gctgacaact 20

<210> 77
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 77
gtgggtatca tctgaagctt 20

<210> 78
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 78
gtcagcccag gctgtggtta 20

<210> 79
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 79
tagattctca tactgaggat 20

<210> 80
<211> 20
<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 80

tgaccatta tctcacagtt

20

<210> 81

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 81

aatgccttac ctgcccgggc

20

<210> 82

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 82

cattatgaac aaaagctggt

20

<210> 83

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 83

agggttagac agacaatgca

20

<210> 84

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 84

gtggcagcat ctgtggtaag 20

<210> 85
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 85
cggaacttac cacaccaaag 20

<210> 86
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 86
ctagtaaaac ctatgaagac 20

<210> 87
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 87
gtgcttctaa tagcgaatt 20

<210> 88
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 88
aaatgcatac cttcttaatg 20