



(72) ANNEFORS, Staffan, SE

(72) BAUER, Carl-Axel, SE

(72) NILSSON, Hans, SE

(71) ASTRA AKTIEBOLAG, SE

(51) Int.Cl.⁶ A61K 31/27, A61K 9/08

(30) 1996/12/20 (9604752-7) SE

(54) **FORMULATION AQUEUSE COMPRENANT DU BAMBUTEROL
ET SON UTILISATION**

(54) **AN AQUEOUS FORMULATION COMPRISING BAMBUTEROL
AND THE USE THEREOF**

(57) L'invention concerne de nouvelles formulations aqueuses de bambutérol et de leurs sels pharmaceutiquement acceptables ainsi que leur utilisation pédiatrique.

(57) New aqueous formulations of, and the paediatric use of, bambuterol, and of its pharmaceutically acceptable salts, are described.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 31/27, 9/08</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/27981 (43) International Publication Date: 2 July 1998 (02.07.98)</p>
<p>(21) International Application Number: PCT/SE97/02053 (22) International Filing Date: 9 December 1997 (09.12.97) (30) Priority Data: 9604752-7 20 December 1996 (20.12.96) SE (71) Applicant (for all designated States except US): ASTRA AKTIEBOLAG (publ) [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): ANNEFORS, Staffan [SE/SE]; Stilgjutaregatan 9, S-227 36 Lund (SE). BAUER, Carl-Axel [SE/SE]; Kruthornsgränd 16, S-226 52 Lund (SE). NILSSON, Hans [SE/SE]; Hjalmar Gullbergs väg 24, S-224 66 Lund (SE). (74) Agent: ASTRA AKTIEBOLAG; Patent Dept., S-151 85 Södertälje (SE).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: AN AQUEOUS FORMULATION COMPRISING BAMBUTEROL AND THE USE THEREOF

(57) Abstract

New aqueous formulations of, and the paediatric use of, bambuterol, and of its pharmaceutically acceptable salts, are described.

AN AQUEOUS FORMULATION COMPRISING BAMBUTEROL AND THE USE THEREOF

This invention relates to a new pharmaceutical formulation and to its manufacture and use. It also relates to a new paediatric use of bambuterol.

5

Bambuterol, and its pharmaceutically acceptable salts, is known as a bronchodilator from EP 43807. Bambuterol is sold under the Trade Mark Bambec[®] in the form of tablets of its hydrochloride salt. Bambuterol is a pro-drug of the adrenergic selective β_2 -receptor agonist terbutaline and is slowly metabolised in the body to active terbutaline.

10

There has been, and continues to be, an increase in the incidence of asthma amongst children. The administration of drugs to children, e.g. by way of inhalation or by means of tablets and capsules which are to be swallowed, can be difficult as the children, and especially younger children, do not always, or readily, co-operate. Furthermore some other patients find that a tablet is difficult to swallow, and if the tablet is chewed the drug in it can produce a bitter taste. Thus patient compliance when a tablet is prescribed is not always good.

15

We have now found that bambuterol can be formulated and administered in a more convenient form than a tablet.

20

Thus according to the invention we provide an aqueous formulation of bambuterol, or of a pharmaceutically acceptable salt thereof, wherein the formulation has a pH in the range 3.7 to 4.2.

25

We prefer the formulation to be adapted to be administered by swallowing.

We prefer the bambuterol to be in the form of its hydrochloride salt. The formulation is preferably a solution and preferably contains from about 0.5 to 5, e.g. about 1.0 mg, of bambuterol, measured as the hydrochloride, per ml.

30

The density of the formulation is preferably between about 1.03 and 1.09 g/ml.

We also prefer the pH of the formulation to be about 3.9. The pH of the formulation may
5 be adjusted to the desired range or value by means of a suitable buffering agent, e.g. citric
acid and if necessary sodium hydroxide. The skilled person will readily be able to find the
appropriate quantity of buffering agent necessary to achieve the desired pH, but in general
this will be in the range 3.0 to 5.0, e.g. about 4.0 mg/ml of citric acid, and from 0.4 to 0.8,
e.g. about 0.6, mg/ml of sodium hydroxide.

10

The formulation may also contain a sweetening agent, e.g. sorbitol and/or glycerol, or a
glucose polymer, for example that known as Lycasin[®]. The proportion of sweetening
agent in the formulation will depend on the particular sweetening agent(s) used, but should
be sufficient, together with any flavouring agent which is used, to cover the bitter taste of
15 the bambuterol. We prefer the sweetening agent not to be such as to encourage dental
caries. We prefer the formulation to contain from 100 to 200 mg, more preferably about
150 mg, per ml of sorbitol. We also prefer the formulation to contain from 75 to 125 mg,
and more preferably about 100 mg, of glycerol per ml.

20 The formulation may also, e.g. when it is to be put up in a multi-dose form, contain a
preservative, e.g. sodium benzoate. The preservative should be present in such a quantity
as to produce a satisfactory preservative effect during its expected period of use. Thus we
prefer the formulation to contain from 0.75 to 1.25 mg, and more preferably about 1.0 mg,
per ml of sodium benzoate.

25

We have surprisingly found that the narrow pH range of the formulations according to the
invention gives the optimal combination of stability of the active agent and of preservative
action.

We have found that certain flavouring agents are incompatible with the desired formulation and/or with the containers to be used for the formulation. Surprisingly however we have found that no such incompatibility exists when essence of blackcurrent is used. The essence of blackcurrent contains natural and synthetic flavours (which are identical to the natural
5 flavours) in propylene glycol as solvent.

The unit and daily dosage of bambuterol to be used will depend on the patient and the type and severity of the condition to be treated.

10 One would expect that the dosage required for a child would be considerably lower than for an adult, i.e. a lower dosage proportionate to the child's, as compared to an adult's, body weight. Surprisingly we have found that this is not the case and that the dosage for children aged 6 and above, e.g. aged 6-12, is substantially the same as for an adult. The dosage for children aged below 6, e.g. from 2-5, is about one half of the adult dose. In both instances
15 this is much more on a mg/kg basis than the corresponding adult dose.

We believe that the higher dose required for children is explained by an unexpected higher metabolic rate of the active ingredient in children as compared to adults.

20 Thus according to a further aspect of the invention we provide a method of treatment of a child in need of treatment with a bronchodilator, which comprises administering to the child a higher dose, as measured on an mg/kg basis, of bambuterol than the corresponding adult dose.

25 We prefer to administer a daily dose of from about 0.5 mg/kg to about 1.0 mg/kg of bambuterol, measured as the hydrochloride, to children aged from 2 to 5, and a dose of from about 0.14 mg/kg to about 1.0 mg/kg measured on the same basis to children aged from 6 to 12.

An adult unit dose of bambuterol comprises from 10 to 20mg measured as the hydrochloride, i.e. 0.14 or 0.28 mg/kg for a 70 kg adult.

The dosage is generally given once a day for both children and adults.

5

According to the invention we also provide the use of bambuterol, or a pharmaceutically acceptable salt thereof, for the preparation of a paediatric medicament.

The paediatric medicament is preferably an aqueous formulation according to the invention, and may be put up as unit doses of the quantities given above or may be in the form of multiple doses, e.g. 100 or 300 ml units. The multiple doses may be packaged in any suitable container with which the formulation is compatible, e.g. a high density polyethylene bottle. The container is preferably fitted with a child resistant closure, which may also, or alternatively, be tamper evident.

15

The formulations according to the invention may be made by conventional pharmaceutical means, e.g. by simple mixing of the ingredients in the desired proportions.

The invention is illustrated, but in no way limited, by the following Examples

20

Example 1

Formulation in parts by weight

	Bambuterol hydrochloride	1.0
	Sorbitol 70% (non crystallising)	150.0
25	Glycerol	100.0
	Sodium benzoate	1.0
	Citric acid	4.0
	Sodium hydroxide	0.6
	Blackcurrent essence	0.5
30	Water purified	to 1,000

The formulation may be made by dissolving the sorbitol and glycerol in a portion of the water. The citric acid, sodium hydroxide, sodium benzoate and bambuterol hydrochloride are dissolved in a further portion of the water and the two solutions are then mixed. The
5 blackcurrent essence is then added to the solution and the whole mixed, filtered and filled into containers. The pH of the solution is about 3.9.

Example 2

Using the formulation of Example 1 several asthmatic children aged between 2 and 5 years
10 were treated once daily, given in the evening, at a dose of 10mg measured as bambuterol hydrochloride. The treatment was continued for up to 3 months. Effectiveness was evaluated from a daily diary (morning and evening) which included details re. asthma symptoms, use of rescue medication (inhaled beta-agonist), no. of awakenings due to asthma and PEF (peak expiratory flow). The treatment was effective.

Claims

1. An aqueous formulation of bambuterol, or of a pharmaceutically acceptable salt thereof, wherein the formulation has a pH in the range 3.7 to 4.2.
5
2. A formulation according to claim 1, wherein the bambuterol is in the form of its hydrochloride salt.
3. A formulation according to either of claims 1 or 2, wherein the formulation is a solution
10 containing from 0.5 to 5 mg of bambuterol, measured as the hydrochloride, per ml.
4. A formulation according to any one of the preceding claims comprising a buffering agent.
- 15 5. A formulation according to any one of the preceding claims comprising a sweetening agent.
6. A formulation according to any one of the preceding claims comprising a preservative.
- 20 7. A formulation according to any one of the preceding claims comprising blackcurrent flavouring.
8. A formulation according to any one of the preceding claims having a density of from 1.03 to 1.09 g/ml.
25
9. A method of treatment of a child in need of treatment with a bronchodilator, which comprises administering to that child a higher dose, as measured on an mg/kg basis, of bambuterol, or of a pharmaceutically acceptable salt thereof, than the corresponding adult dose.
30

10. A method according to claim 9, wherein a dose of from 10 to 20 mg of bambuterol, measured as the hydrochloride salt, is administered once a day to a child aged 2 to 5.
11. A method according to claim 10, wherein a daily dose of from 0.5 mg/kg to 1.0 mg/kg
5 of bambuterol, measured as the hydrochloride salt, is administered to the child.
12. A method according to any one of claims 9 to 11, wherein the bambuterol is administered as a formulation according to any one of claims 1 to 8.
- 10 13. Use of bambuterol, or of a pharmaceutically acceptable salt thereof, for the preparation of a paediatric medicament.
14. Use according to claim 13, wherein the medicament is such that a dose of 10 to 20 mg of bambuterol, measured as the hydrochloride salt, is to be administered once a day to a
15 child aged 2 to 5.