



US005118689A

United States Patent [19][11] **Patent Number:** **5,118,689**

Oinuma et al.

[45] **Date of Patent:** **Jun. 2, 1992**[54] **PIPERIDINE DERIVATIVE AND PHARMACEUTICAL COMPOSITION CONTAINING THE SAME**[75] **Inventors:** Hitoshi Oinuma, Sakuramura; Motosuke Yamanaka, Abiko; Kazutoshi Miyake, Ushiku; Tomonori Hoshiko, Tsuchiura; Norio Minami, Sakura; Tadao Shoji, Kukizakimachi; Yoshiharu Daiku, Sakuramura; Kohei Sawada, Toride; Kenichi Nomoto, Tsuchiura, all of Japan[73] **Assignee:** Eisai Co., Ltd., Tokyo, Japan[21] **Appl. No.:** 594,079[22] **Filed:** Oct. 9, 1990**Related U.S. Application Data**

[62] Division of Ser. No. 408,106, Sep. 15, 1989, Pat. No. 4,996,215, which is a division of Ser. No. 16,035, Feb. 18, 1987, Pat. No. 4,876,262.

[30] **Foreign Application Priority Data**


Feb. 26, 1986 [JP] Japan 61-39270

[51] **Int. Cl.⁵** **A61K 31/47; A61K 31/52; A61K 31/495; C07D 473/04**[52] **U.S. Cl.** **514/300; 514/252; 514/265; 514/312; 514/314; 544/267; 544/271; 544/353; 544/354; 544/355; 546/121; 546/156; 546/157; 546/168; 546/172**[58] **Field of Search** 546/121, 157, 156, 172, 546/168; 514/300, 252, 265, 312, 314; 544/353, 354, 355, 267, 271[56] **References Cited PUBLICATIONS**

Wong et al., C.A. 107:32609z (1987).

Primary Examiner—Joseph Paul Brust
Attorney, Agent, or Firm—Flynn, Thiel, Boutell & Tanis[57] **ABSTRACT**

A new piperidine compound is pharmacologically effective for treatment of arrhythmia and has the formula:

in which R^1 is lower alkyl or tolyl; R^2 is hydrogen, lower alkyl, lower alkenyl, cycloalkyl or cycloalkylalkyl; X is $-CO-$, $-CH_2-$ or $-CHOH-$; the sum of g plus h equals the integer 3 or 4 with g and h being 1, 2 or 3; A is substituted or unsubstituted alkylene, alkenylene, $-(CH_2)_k-S-$, wherein k is an integer of 2 to 5, or $-(CH_2)_pCO-$, wherein p is an integer of 1 to 4; and B is a compound having at least one heterocyclic ring containing a nitrogen atom.**9 Claims, No Drawings**

**PIPERIDINE DERIVATIVE AND
PHARMACEUTICAL COMPOSITION
CONTAINING THE SAME**

This is a division of U.S. Ser. No. 07/408,106, filed Sep. 15, 1989, now U.S. Pat. No. 4,996,215, patented Feb. 26, 1991, which is a division of U.S. Ser. No. 07/016,035, filed Feb. 18, 1987, now U.S. Pat. No. 4,876,262, issued Oct. 24, 1989.

The present invention relates to piperidine derivatives and pharmacologically acceptable salts thereof having excellent medicinal effects, processes for producing them and medicines containing them.

PRIOR ART

Arrhythmias often occur as complications to cardiac diseases such as myocardial infarction and heart failure. In a serious case, arrhythmias give rise to a ventricular fibrillation to cause sudden death.

Although various antiarrhythmic agents are now available on the market, those having both satisfactory effects and high safety have not been obtained yet. For example, antiarrhythmic agents of Class I according to the classification of Vaughan-Williams which cause a selective inhibition of the maximum velocity of the upstroke of the action potential (V_{max}) have only insufficient effects for preventing the ventricular fibrillation. In addition, they have problems regarding safety, namely, they cause a depression of the myocardial contractility and have a tendency to induce arrhythmias due to an inhibition of the impulse conduction. Beta-adrenoceptor blockers and calcium antagonists which belong to Class II and IV respectively, have a defect that their effects are either limited to a certain type of arrhythmia or are unreliable, though their safety is higher than that of the antiarrhythmic agents of Class I.


Antiarrhythmic agents of Class III are drugs which cause a selective prolongation of the duration of the action potential without a significant depression of the V_{max} . Although there has been no available drug which possess pure and potent properties of Class III antiarrhythmics, drugs of this class are expected to be effective in preventing ventricular fibrillations. Moreover, they are, by definition, not considered to cause a myocardial depression or an induction of arrhythmias due to the inhibition of the action potential conduction as seen with Class I antiarrhythmic agents.

Under these circumstances, the development of pure and potent antiarrhythmic agents of Class III has been awaited.


SUMMARY OF THE INVENTION

An object of the present invention is to provide new piperidine derivatives and pharmacologically acceptable salts thereof, processes for producing the piperidine derivatives and pharmacologically acceptable salts thereof and medicines containing said piperidine derivatives or pharmacologically acceptable salts thereof as the active ingredient.

The invention provides a new piperidine derivative, an analogous compound thereto, however having a heterocyclic ring other than the piperidine ring and a pharmacologically acceptable salt thereof. These compounds have the formula shown below.



in which R1 is a lower alkyl or a tolyl, R2 is hydrogen, hydroxyl, a lower alkoxy or a lower alkyl, R3 is hydrogen, a lower alkyl, a lower alkenyl, a cycloalkyl or a cycloalkylalkyl, X is $-\text{CO}-$, $-\text{CH}_2-$ or $-\text{CHOH}-$, g is an integer of 1 to 3, h is an integer of 1 to 3, Y is hydrogen, a lower alkyl, a lower alkenyl, cyano, $-\text{CH}_2\text{COOR}$, R being hydrogen or a lower alkyl, a cycloalkyl, a cycloalkylalkyl,




25 1 being 1 or 2, -A-B, A being $-(\text{CH}_2)_n-$, n being an integer of 1 to 5,


a straight-chain alkylene group having 1 to 5 carbon atoms which is a divalent group derived from a straight-chain alkane having lower alkyl, phenyl or hydroxyl group(s) bonded directly to one or more carbon atoms constituting said alkane by removing a hydrogen atom bonded to each of the carbon atoms located at both ends thereof,

30 a straight-chain alkenylene group having 2 to 5 carbon atoms which is a divalent group derived from a straight-chain alkene having a double bond formed between carbon atoms adjacent to each other by removing a hydrogen atom bonded to each of the carbon atoms located at both ends thereof,


35 $-(\text{CH}_2)_k-\text{S}-$, k being an integer of 2 to 5, $-(\text{CH}_2)_p\text{CO}-$, p being an integer of 1 to 4, B being cyano, $-\text{NR}^4\text{R}^5$,




50



60




65




5

-continued



Production process A




Step 2 ↓ hydrolysis




Step 3 ↓ Z-Y' (wherein Z represents a halogen atom such as a chlorine, bromine or iodine atom or a free group such as a methanesulfonyloxy or p-toluenesulfonyloxy group and n is as defined above)
Y' is defined above except for hydrogen excluded.



step 4 of reduction ↓



step 6 of reduction ↓





Production Process B

The intended compounds of the above general formula in which R³ is a lower alkyl, a lower alkenyl, or a cycloalkyl can be produced as shown below.


Process 1

6


-continued

10 step 7 of alkylation ↓ R³.Hal (X) (Hal is a halogen)



reduction ↓



reduction ↓



Process 2

45 step 8 of alkylation ↓ R³.Hal (X)

Production Process C


The compounds of the above general formula (I) in which Y is H and X is a group of the formula:

60




can be produced by directly reducing the above-mentioned compound (IV), omitting the step 3, to obtain the compound (XIII). The reduction is conducted in the same way as shown in the step 4.

65




Process D


The compound in which X is —CH₂— in the formula (I) is produced below.




step 9 of reduction



step 10 of hydrolysis




step 11 of alkylation




Process E


The process A provides the invention compound in the step 3. The invention compound is also produced by each below shown method when Y is one of the followings. method E-1







A compound in which A is —CH₂— and B is



is produced. A compound in which A is




is produced. This B is called Y².




method E-2

A compound in which A is —(CH₂)₂— and B is




is produced below. This —A—B is called Y³.



Each above shown step is conducted as described below in detail.

Step 1

A sulfonamide derivative (II) is reacted, according to the Friedel-Crafts reaction, with a reactive acid derivative such as a halogenate or anhydride of a carboxylic acid having the following formula:



in which R' is a lower alkyl or phenyl and g and h are defined above, in the presence of a Lewis acid such as aluminum chloride, tin chloride and zinc chloride in an inert solvent such as carbon disulfide, dichloromethane, chloroform and nitrobenzene to form a corresponding anilide derivative (III).


Step 2

The acetyl group of the compound (III) obtained in the step 1 is hydrolyzed in this step. The hydrolysis is effected in, for example, a dilute aqueous alkali solution or dilute aqueous mineral acid solution. In a preferred embodiment, the hydrolysis is effected in 2 to 6N hydrochloric acid under reflux or in a 0.5 to 3N aqueous solution of sodium hydroxide under reflux.

Step 3

(1) in case Y is other than hydrogen, that is, Y is Y'


The compound (IV) of the step 2 is normally condensed with a compound having the formula: Z-Y'(V), such as Z-A-B and




10 in which Z is a group to eliminate and is a halogen such as chlorine, bromine and iodine, methanesulfonyloxy and p-toluenesulfonyloxy.

In a preferred embodiment of this process, the reaction is carried out in the presence of a deacidifying agent such as potassium carbonate or sodium carbonate and potassium iodide (used when Z is not iodine) in a solvent such as N,N-dimethylformamide, dimethyl sulfoxide, acetonitrile, acetone, butanol, propanol, ethanol or methanol at a temperature of about 50° to 120° C. to obtain the compound (VII).

(2) A compound in which A is $-(CH_2)_2-$ and B is



30 is produced below. The reaction in detail is illustrated below.



More particularly, an unsubstituted or substituted vinylpyridine (VI) is reacted with the compound (free base) (IV) obtained in the above-mentioned step 2 or pharmacologically acceptable acid-addition salt thereof in a lower alkyl alcohol such as methanol, ethanol or propanol or a mixture thereof with water at a temperature ranging from room temperature to about 100° C. to obtain the intended compound (XXVI). When a free base is used as the starting material in this process, preferred results are obtained by using an acidic catalyst such as acetic or hydrochloric acid or an alkali metal catalyst such as sodium.

In the steps 4 to 6, the compound (VII) obtained in the step 3 is reduced to produce the compounds (VIII) or (IX).

11

Step 4

The compound (VII) prepared in the step 3 is reduced in this step. The reduction is effected by an ordinary method wherein, for example, the compound (VII) is treated with sodium borohydride or lithium borohydride in a solvent such as methanol, ethanol, 2-propanol, dimethylformamide or dimethyl sulfoxide at a temperature ranging from about -10° C. to room temperature to obtain the intended alcohol (VIII) of the present invention.

Step 5

An arylketone compound (VII) is reacted with two or more equivalents of a trialkylsilane, preferably triethylsilane, in an excess of trifluoroacetic acid, at a temperature ranging from the room temperature to 100° C. for several days to produce the compound (IX). Dichloroethane, carbon tetrachloride, chloroform, dichloromethane and acetonitrile are also used as a solvent.

Step 6

An alcohol compound (VIII) obtained in the same way as shown in the step 4 is treated with an acid, preferably with 20% sulfuric acid-acetic acid for a short time to obtain a dehydrate and the dehydrate is catalytically hydrogenated to produce the compound (IX).

Step 7

The compound (VII)' of the above general formula (I) wherein R^3 is H is N-alkylated in this step. The compound (VII)' is reacted with an alkyl halide of the formula (X) in the presence of a base in a solvent such as dimethylformamide, dimethyl sulfoxide, a lower alkyl alcohol, e.g. methanol, ethanol or propanol, or acetone at a reaction temperature of about 50° to 120° C. to obtain the intended compound (XI). The bases used in this step include, for example, potassium carbonate, sodium carbonate, sodium hydrogencarbonate, sodium ethoxide, sodium methoxide and sodium hydride.

When 2 mole equivalents or more of R^3Hal (X) is used, dialkylation, alkenylation for di-lower alkenyl and alkylation for dicycloalkyl may be carried out as far as Y is hydrogen. Similarly the step 8 of the process B can be conducted for N-alkylation.

The compound (XI) obtained is reduced in the same way as shown in the steps 4 to 6 to further obtain the compound (VIII)' or (IX)'.

Steps 9 to 11

Each step is conducted in the same way as shown the step 5 (reduction), the step 2 (hydrolysis) and the steps 7 and 8 (alkylation).

Step 12

A secondary amine (XVII) is converted to a tertiary amine (XXI) by the Mannich reaction. An active hydrogen-containing compound, such as a furane compound, a pyrrole compound and a nitrogen-including heterocyclic compound having methyl, is condensed with an amine and formaldehyde or para-formaldehyde, preferably in a solvent such as water and an alcohol, under the acidic condition with acetic acid or hydrochloric acid, at the room temperature to 100° C.

Step 13

A secondary amine is alkylated by the Mannich reaction. The reaction proceeds in the same way as shown in

12

the step 12. It is also possible that the secondary amine is used in the form of hydrochloric acid salt, not a free acid, to produce the compound (XXV).

The piperidine derivative obtained above can be converted to a pharmacologically acceptable salt thereof by a conventional method.

To facilitate the understanding of the present invention, typical examples of the compounds of the present invention will be shown below, which by no means limit the invention. The compounds are shown in their free form.

1. 4-(4-methylsulfonylaminobenzoyl)-1-[2-(3-pyridyl)ethyl]piperidine,
2. 4-(4-methylsulfonylaminobenzoyl)-1-(4-pyridyl)methylpiperidine,
3. 4-(4-methylsulfonylaminobenzoyl)-1-[3-(4-pyridyl(propyl))piperidine,
4. 1-(6-methyl-3-pyridyl)methyl-4-(4-methylsulfonylaminobenzoyl)piperidine
5. 4-(4-methylsulfonylaminobenzoyl)-1-[2-(4-pyridyl)ethyl]piperidine,
6. 1-[2-(3,4-dimethoxyphenyl)ethyl]-4-(4-methylsulfonylaminobenzoyl)piperidine,
7. 4-(4-methylsulfonylaminobenzoyl)-1-[4-(3-pyridyl)butyl]piperidine,
8. 4-(4-methylsulfonylaminobenzoyl)-1-[2-(4-pyridyl)thioethyl]piperidine,
9. 4-(4-methylsulfonylaminobenzoyl)piperidine,
10. 4-(4-methylsulfonylaminobenzoyl)-1-[3-(3-pyridyl)propyl]piperidine,
11. 4-(4-methylsulfonylaminobenzoyl)-1-[5-(3-pyridyl)pentyl]piperidine,
12. 1-(6-chloro-3-pyridyl)methyl-4-(4-methylsulfonylaminobenzoyl)piperidine,
13. 4-(4-methylsulfonylaminobenzoyl)-1-[2-(2-pyridyl)ethyl]piperidine,
14. 4-(4-methylsulfonylaminobenzoyl)-1-(2-phenylethyl)piperidine,
15. 4-(2-hydroxy-4-methylsulfonylaminobenzoyl)-1-(4-pyridyl)methylpiperidine,
16. 4-(2-hydroxy-4-methylsulfonylaminobenzoyl)-1-(2-phenylethyl)piperidine,
17. 4-(4-methylsulfonylaminobenzoyl)-1-(3-pyridyl)methylpiperidine,
18. 4-(4-methylsulfonylaminobenzoyl)-1-(2-pyridyl)methylpiperidine,
19. 4-(4-methylsulfonylaminobenzoyl)-1-nicotinoylmethylpiperidine,
20. 4-(4-methylsulfonylaminobenzoyl)-1-[2-(2-thienyl)ethyl]piperidine
21. 4-(4-methylsulfonylaminophenyl)-hydroxymethyl-1-(2-(2-pyridyl)ethyl)piperidine
22. 4-(2-methoxy-4-methylsulfonylaminobenzoyl)-1-(4-pyridyl)methylpiperidine,
23. 1-[2-(4-chlorophenyl)ethyl]-4-(4-methylsulfonylaminobenzoyl)piperidine,
24. 1-[2-(4-methoxyphenyl)ethyl]-4-(4-methylsulfonylaminobenzoyl)piperidine,
25. 4-(4-methylsulfonylaminobenzoyl)-1-[3-(3-pyridyl)-2-propenyl]piperidine,
26. 4-(4-ethylsulfonylaminobenzoyl)-1-(4-pyridyl)methylpiperidine,
27. 4-(4-ethylsulfonylaminobenzoyl)-1-(2-phenylethyl)piperidine,
28. 1-benzyl-4-(4-methylsulfonylaminobenzoyl)piperidine,
29. 1-[2-(4-fluorophenyl)ethyl]-4-(4-methylsulfonylaminobenzoyl)piperidine,

30. 4-(4-methylsulfonylamino benzoyl)-1-(3-phenylpropyl)piperidine,
31. 4-(4-methylsulfonylamino benzoyl)-1-(2-thienylmethyl)methylpiperidine,
32. 1-[2-(4-hydroxyphenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
33. 1,4-di(4-methylsulfonylamino benzoyl)piperidine,
34. 1-[6,7-dihydro-5H-7-cyclopenta[b]pyridinyl]methyl-4-(4-methylsulfonylamino benzoyl)piperidine,
35. N-methyl-4-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine,
36. N-butyl-4-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine,
37. 4-(4-methylsulfonylamino benzoyl)-1-[1-(4-pyridyl)ethyl]piperidine,
38. 4-(4-methylsulfonylamino benzoyl)-1-[1-phenyl-1-(4-pyridyl)methyl]piperidine,
39. 1-[2-(4-methylphenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
40. 4-(4-methylsulfonylamino benzoyl)-1-(1-naphthyl)methylpiperidine,
41. 1-[2-hydroxy-2-(4-methylsulfonylamino phenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
42. 1-[2-hydroxy-2-(3-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
43. 1-(2-hydroxy-2-phenylethyl)-4-(4-methylsulfonylamino benzoyl)piperidine,
44. 1-(2-chlorophenyl)methyl-4-(4-methylsulfonylamino benzoyl)piperidine,
45. 1-[2-(5-ethyl-2-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
46. 1-[2-(6-methyl-2-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
47. 1-[2-(6-methyl-3-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
48. 1-[3-(6-methyl-3-pyridyl)propyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
49. 1-[1-(3-methoxy-2-pyridyl)methyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
50. 1-[2-(5-methyl-4-pyridylthio)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
51. 1-[1-(2-methoxy-5-pyridyl)methyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
52. 1-[1-(3-hydroxy-2-pyridyl)methyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
53. 1-[2-(6-ethyl-2-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
54. 1-[3-(5-ethyl-2-pyridyl)propyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
55. 1-[4-(5-ethyl-2-pyridyl)butyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
56. 1-[2-(5-butyl-2-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
57. 1-[2-(2-methyl-4-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
58. 1-[2-(2-ethyl-4-pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine,
59. 1-[2-(2-chloro-5-pyridyl)ethyl]-4-[4-methylsulfonylamino benzoyl]piperidine,
60. 1-[3-(2-chloro-5-pyridyl)propyl]-4-(4-methylsulfonylamino benzoyl)piperidine, and
61. 4-(4-methylsulfonylamino benzoyl)-1-[2-(4-methyl-5-thiazole)ethyl]piperidine.
62. 1-[4-(1-imidazolyl)benzoylmethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
63. 1-[2-[3-(2-cyano)pyridyl]-ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine

64. 1-[3-[3-(2-cyano)pyridyl]propyl]-4-(4-methylsulfonylamino benzoyl)piperidine
65. 1-[3-[4-(2-cyano)pyridyl]propyl]-4-(4-methylsulfonylamino benzoyl)piperidine
- 5 66. 1-[2-(1-imidazolyl)-3-pyridylmethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
67. 1-(5-methyl-2-furanyl)methyl-4-(4-methylsulfonylamino benzoyl)piperidine
68. 1-(1-methyl-2-pyrrolyl)methyl-4-(4-methylsulfonylamino benzoyl)piperidine
- 10 69. 1-(1-imidazolyl-3-propyl)-4-(4-methylsulfonylamino benzoyl)piperidine
70. 4-(4-methylsulfonylamino)-1-[2-(3-pyridazinylethyl)piperidine
- 15 71. 4-(4-methylsulfonylamino benzoyl)-1-[2-(4-pyrimidinyl)-2-propenyl]piperidine
72. 4-(4-methylsulfonylamino benzoyl)-1-(2-pyrazinylmethyl)piperidine
73. 4-(4-methylsulfonylamino benzoyl)-1-[2-(2-pyrazinyl)ethyl]piperidine
- 20 74. 1-[2-(1,2-dihydro-6-methyl-2-oxo-3-pyridinecarbonitril-5-yl)-2-oxoethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
- 25 75. 4-(4-methylsulfonylamino benzoyl)-1-(6-uracilmethyl)piperidine
76. 1-[2-(3-indolyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
77. 4-(4-methylsulfonylamino benzoyl)-1-(2-phthalimidoethyl)piperidine
- 30 78. 4-(4-methylsulfonylamino benzoyl)-1-(2-quinolylmethyl)piperidine
79. 4-(4-methylsulfonylamino benzoyl)-1-(3-quinolylmethyl)piperidine
- 35 80. 1-(1-imidazo[1,2-a]pyridylmethyl)-4-(4-methylsulfonylamino benzoyl)piperidine
81. 1-[2-(1-imidazo[1,2-a]pyridyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
82. 1-(6-imidazo[1,2-a]pyridylmethyl)-4-(4-methylsulfonylamino benzoyl)piperidine
- 40 83. 1-[2-(3-imidazo[1,2-a]pyridyl)-2-oxoethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
84. 1-(2-benzimidazolylmethyl)-4-(4-methylsulfonylamino benzoyl)piperidine
- 45 85. 4-(4-methylsulfonylamino benzoyl)-1-[2-(2-quinoxaliny)ethyl]piperidine
86. 4-(4-methylsulfonylamino benzoyl)-1-[2-(7-theophilynyl)ethyl]piperidine
87. 1-(9-fluorenyl)-4-(4-methylsulfonylamino benzoyl)piperidine
- 50 88. 1-ethyl-4-(4-methylsulfonylamino benzoyl)piperidine
89. 1-n-butyl-4-(4-methylsulfonylamino benzoyl)piperidine
- 55 90. 1-cyclohexylmethyl-4-(4-methylsulfonylamino benzoyl)piperidine
91. 1-(2-methyl-2-propenyl)-4-(4-methylsulfonylamino benzoyl)piperidine
92. 1-ethoxycarbonylmethyl-4-(4-methylsulfonylamino benzoyl)piperidine
- 60 93. 1-cyano-4-(4-methylsulfonylamino benzoyl)piperidine
94. 1-cyanomethyl-4-(4-methylsulfonylamino benzoyl)piperidine
95. 1-(3-cyanopropyl)-4-(4-methylsulfonylamino benzoyl)piperidine
96. 1-[2-(N',N'-diethylamino)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine

- 97. 4-(4-methylsulfonylamino benzoyl)-1-[2-(1-pyrrolidinyl)ethyl]piperidine
- 98. 4-(4-methylsulfonylamino benzoyl)-1-[2-(1-piperidinyl)ethyl]piperidine
- 99. 4-(4-methylsulfonylamino benzoyl)-1-[2-(4-morpholinyl)ethyl]piperidine
- 100. 4-(4-methylsulfonylamino benzoyl)-1-[3-(1-piperidinyl)propyl]piperidine
- 101. 1-[3-(4-pyridyl)propyl]-4-[4-(p-toluenesulfonylamino)benzoyl]piperidine
- 102. 1-[2-(6-methyl-2-pyridyl)ethyl]-4-[4-(p-toluenesulfonylamino)benzoyl]piperidine
- 103. 4-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine
- 104. 1-[2-(3,4-dimethoxyphenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine
- 105. 3-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine
- 106. 1-[2-(6-methyl-2-pyridyl)ethyl]-3-(4-methylsulfonylamino benzoyl)piperidine
- 107. 3-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]pyrrolidine
- 108. 1-[2-(6-methyl-2-pyridyl)ethyl]-3-(4-methylsulfonylamino benzoyl)pyrrolidine
- 109. 1-[2-(3,4-dimethoxyphenyl)ethyl]-3-(4-methylsulfonylamino benzoyl)piperidine
- 110. 1-ethyl-4-(N-ethyl-4-methylsulfonylamino benzoyl)piperidine
- 111. 1-n-butyl-4-(N-n-butyl-4-methylsulfonylamino benzoyl)piperidine
- 112. 1-cyclohexylmethyl-4-(N-cyclohexylmethyl-4-methylsulfonylamino benzoyl)piperidine
- 113. 1-(2-methyl-2-propenyl)-4-[N-(2-methyl-2-propenyl)-4-methylsulfonylamino benzoyl]piperidine
- 114. 4-(4-methylsulfonylamino benzoyl)-1-(2-(4-methylsulfonylamino phenyl)-2-oxoethyl)piperidine

The piperidine derivatives obtained according to the present invention prolong the refractory period by specifically prolonging the action potential duration to prevent arrhythmia without exerting any influence on the myocardial conduction velocity. These derivatives are antiarrhythmic agents of Class III of the above-mentioned Vaughan-Williams classification.

The following experimental examples will further illustrate the effects of the compounds of the present invention.

EXPERIMENTAL EXAMPLE 1

Effects on the action potential duration in the isolated myocardium of guinea-pigs

Right ventricular papillary muscles were isolated from male guinea-pigs of Hartley strain weighing 300 to 400 g and fixed at the bottom of an acrylic bath with pins. They were perfused with Tyrode solution kept at 37° C. and saturated with a mixture of 95% O₂ and 5% CO₂. The muscles were stimulated at 1 Hz with rectangular pulses of 1 msec duration and supramaximal voltage. Action potentials were recorded using conventional glass microelectrodes filled with 3M KCl. The duration of the action potential and the maximum velocity of the upstroke of the action potential (V_{max}) were determined. Each of the test compounds was included in Tyrode solution at 10⁻⁶ or 10⁻⁵ M and perfused. The effects of the 10⁻⁶ M solution was observed for the first 10 min, then those of the 10⁻⁵ M solution were observed for another 10 min.

The results are shown in Table 1. The test compounds shown in Example 1 were as follows. Sotalol, a

beta-adrenoceptor antagonist was employed as the reference drug because this compound is known to prolong the duration of the myocardial action potential.

Test compound	
Compound A:	1-benzyl-4-(4-methylsulfonylamino benzoyl)piperidine hydrochloride
Compound B:	4-(4-methylsulfonylamino benzoyl)-1-(2-phenylethyl)piperidine hydrochloride
Compound C:	1-[2-(4-chlorophenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine hydrochloride
Compound D:	1,4-di(4-methylsulfonylamino benzoyl)-piperidine hydrochloride
Compound E:	1-[2-(3,4-dimethoxyphenyl)ethyl]-4-(4-methylsulfonylamino benzoyl)piperidine hydrochloride
Compound F:	4-(4-methylsulfonylamino benzoyl)-1-[2-(2-thienyl)ethyl]piperidine methylsulfonate
Compound G:	4-(4-methylsulfonylamino benzoyl)-1-(4-pyridyl)methylpiperidine dihydrochloride
Compound H:	4-(4-methylsulfonylamino benzoyl)-1-[2-(2-pyridyl)ethyl]piperidine dihydrochloride
Compound I:	4-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine dihydrochloride
Compound J:	4-(4-methylsulfonylamino benzoyl)-1-[2-(4-pyridyl)ethyl]piperidine dihydrochloride, and
Compound K:	4-(4-methylsulfonylamino benzoyl)-1-nicotinylpiperidine.
Compound L: (28)	4-(4-methylsulfonylamino benzoyl)-1-(2-quinolylmethyl)piperidine dihydrochloride
Compound M: (29)	4-(4-methylsulfonylamino benzoyl)-1-(3-quinolylmethyl)piperidine dihydrochloride
Compound N: (7)	1-[2-(3-imidazo[1,2-a]pyridyl-1-oxoethyl)-4-(4-methylsulfonylamino benzoyl)-piperidine dihydrochloride
Compound O: (30)	1-(1-imidazo[1,2-a]pyridylmethyl)-4-(4-methylsulfonylamino benzoyl)piperidine dihydrochloride
Compound P: (12)	1-ethyl-4-(4-methylsulfonylamino benzoyl)-piperidine hydrochloride
Compound Q: (31)	1-(6-imidazo[1,2-a]pyridylmethyl)-4-(4-methylsulfonylamino benzoyl)piperidine dihydrochloride
Compound R:	4-(4-methylsulfonylamino benzoyl)-1-(3-(4-pyridyl)propyl)piperidine dihydrochloride
Compound S:	1-(2-(6-methyl-2-pyridyl)ethyl)-4-(4-methylsulfonylamino benzoyl)piperidine dihydrochloride

TABLE 1

Test Compound	10 ⁻⁶ M		10 ⁻⁵ M	
	APD ₉₀ prolongation (%)	V _{max} inhibition (%)	APD ₉₀ prolongation (%)	V _{max} inhibition (%)
A	0	0	7	0
B	26	0	34	19
C	7	0	15	0
D	3	0	10	0
E	12	0	14	0
F	8	0	11	0
G	18	0	33	0
H	2	0	5	0
I	10	0	17	0
J	4	0	10	0
K	9	0	24	14
R	18	0	27	0
S	20	0	30	0
Sotalol	0	0	7	0
L	18	0	26	2
M	3	0	5	0
N	8	0	8	0
O	6	0	12	5
P	7	0	23	2

TABLE 1-continued

Test Compound	10 ⁻⁶ M		10 ⁻⁵ M	
	APD ₉₀ prolongation (%)	V _{max} inhibition (%)	APD ₉₀ prolongation (%)	V _{max} inhibition (%)
Q	4	0	23	0

EXPERIMENTAL EXAMPLE 2

Effects on QTc-interval of ECG in anesthetized dogs

Mongrel dogs were anesthetized with enflurane. The chest was opened at the fifth intercosta and the pericardium was incised to expose the left ventricle. A monopolar electrode fixed on an acrylic plate was sutured to a ventricular surface of the region where the left anterior descending branch of the coronary artery was dominant. The electrocardiogram was recorded from the surface of the left ventricle through the electrode. Test compounds were injected through a catheter inserted into a forearm vein.

Compound B caused a 51% prolongation of QTc-interval (i.e. from 435 to 665 msec) at 0.1 mg/kg. Compound G produced 17, 27 and 35% prolongation of QTc-interval at 0.1, 0.3 and 1 mg/kg, respectively. When the test compound J was injected at 0.1 and 0.3 mg/kg, QTc-interval was prolonged by 21 and 42%, respectively.

When the test compound L was used, the QTc prolongation of 31% with 0.1 mg/kg and 56% with 0.3 mg/kg thereof were recognized. When the test compound M was used, the QTc prolongation of 7% with 0.1 mg/kg, 13% with 0.3 mg/kg and 21% with 1.0 mg/kg thereof were recognized. When the test compound P was used, the QTc prolongation of 7% with 0.1 mg/kg and 14% with 0.3 mg/kg thereof were recognized. The administrations of 0.03 mg/kg and 0.1 mg/kg of the compound R were found to provide 13% prolongation and 21% prolongation of QTc-interval, respectively. As to the compound S, the administrations of 0.01 mg/kg and 0.03 mg/kg were found to provide 30% prolongation and 42%. With 1.0 mg/kg of Sotalol used as a control, 12% prolongation was observed.

EXPERIMENTAL EXAMPLE 3

Acute toxicity in mice

Make ddy mice weighing 20 to 30 g were used for the acute toxicity test. LD₅₀ values were calculated by the up-and-down method. The compound G, J, M, N, O, R and sotalol were dissolved in saline to obtain 16 mg/ml stock solution. Administration of 0.1 ml of the solution to 10 g body weight of the animal corresponds to 160 mg/kg of the sample. The compounds B, F, L and S (the free body) were each dissolved in 20% polyethyleneglycol to obtain stock solutions having a concentration of 8 mg/ml. The dose to be administered was determined on the basis of the volume of the stock solution. The solutions were injected into a tail vein by means of a 1-ml tuberculin syringe. The judgement of survival or death was made 30 min after each injection. The results are shown in Table 2.

TABLE 2

Compound	LD ₅₀ (iv) mg/kg
B	100
F	112
G	108
J	195

TABLE 2-continued

Compound	LD ₅₀ (iv) mg/kg
L	112
M	112
N	224
O	302
R	138
S	112
Sotalol	113

It is apparent from the above Experimental Examples 1 and 2 that the compounds of the present invention have the pharmacological properties required for the antiarrhythmic agents of Class III, namely the prolongation of the myocardial action potential in vitro without a significant depression of the V_{max} and the prolongation of QTc-interval in anesthetized dogs. Moreover, their effects were much more potent than the reference drug, sotalol.

It is expected that the compounds of the present invention are effective in treating and preventing all types of arrhythmias including ventricular and atrial (supraventricular) arrhythmias. The compounds of the present invention are especially expected to control recurrent arrhythmias and prevent sudden death due to the ventricular fibrillation.

The compounds of the present invention can be used either orally or parenterally (intramuscular, subcutaneous or intravenous injection). The dose is not particularly limited, since it varies depending on the type of the arrhythmia, symptoms, age, condition and body weight of the patient. In case when used in combination with other drugs or treatments, it also depends on the kind, frequency and intended effects of the drug or the treatment. The usual oral dose for the adults dose is estimated to be 1 to 100 mg/day, preferably 5 to 50 mg/day and particularly 5 to 15 mg/day. The administration will be made once a day or more. In the case of injection, the dose is estimated to be 0.01 or 1 mg/kg, preferably 0.03 to 0.1 mg/kg.

The compound of the present invention can be given in the form of, for example, powders, finely divided particles, granules, tablets, capsules suppositories and injections. The preparations are produced by an ordinary process by using an ordinary carrier.

More particularly, for example, an oral solid preparation is produced by adding an excipient and, if necessary, a binder, disintegrator, lubricant, colorant, corrigent, etc. to the active ingredient and the mixture is shaped into tablets, coated tablets, granules, powder, capsules, etc.

Examples of the excipients include lactose, corn starch, white sugar, glucose, sorbitol, crystalline cellulose and silicon dioxide. Examples of the binders include polyvinyl alcohol, polyvinyl ethers, ethylcellulose, methylcellulose, acacia gum, tragacanth, gelatin, shellac, hydroxypropylcellulose, hydroxypropylstarch and polyvinylpyrrolidone. Examples of the disintegrators include starch, agar, gelatin powder, crystalline cellulose, calcium carbonate, sodium hydrogencarbonate, calcium citrate, dextrin and pectin. Examples of the lubricants include magnesium stearate, talc, polyethylene glycol, silica and hardened vegetable oils. The colorants are those acceptable as additives for medicines. Examples of the corrigents include cocoa powder, menthol, aromatic acids, peppermint oil, borneol and cinnamon powder. As a matter of course, these tablets and

granules may be coated suitably with sugar, gelatin or the like.



In the production of the injection, a pH adjustor, buffer, stabilizer, solubilizer, etc. are added, if necessary, to the active ingredient and as intravenous injection is produced by an ordinary method.

The following examples will further illustrate the present invention, which by no means limit the present invention.

The final step in the production of the intended compound of the present invention will be shown in the following examples and the production of the starting materials used in the examples will be shown in the following referential examples.

Step I

REFERENCE EXAMPLE 1



21

20.0 g (155 mmol) of (\pm)-nipecotic acid was dissolved in 33 ml of a 20% aqueous sodium hydroxide solution. 23.84 g of benzoyl chloride was added dropwise thereto at such a rate that the reaction temperature would not exceed 20° C. Then 60 ml of a 20% aqueous sodium hydroxide solution was added dropwise thereto and the obtained mixture was stirred at 0° C. for 1 h, acidified with concentration hydrochloric acid and extracted with dichloromethane. The organic layer was washed with water and a saturated aqueous common salt solution, dried over magnesium sulfate and concentrated. The solid residue was recrystallized from ethanol to obtain 18.0 g (yield: 40%) of the intended compound in the form of white crystals.

Melting point: 187° to 188° C.

NMR (90° C., DMSO- d_6) δ 1.3-2.2 (4H, m) 2.2-4.4 (5H, m) 7.42 (5H, S) 12.0-12.6 (1H, br).

The same procedure as above was repeated except that (\pm)-nipecotic acid was replaced with (\pm)- β -proline to obtain the following compound:


(\pm)-N-benzoyl- β -proline

Melting point: 111° to 113° C.

NMR (90 MHz, CDCl₃) δ 2.18 (2H, q-like, J=7 Hz) 2.8-3.3 (1H, m) 3.35-4.00 (4H, m) 7.36 (5H, m)

Step II

REFERENTIAL EXAMPLE 2



10.0 g (42.9 mmol) of (\pm)-N-benzoylnipecotic acid was dissolved in 15 ml of thionyl chloride. Several drops of dimethylformamide were added to the solution and the mixture was stirred at room temperature for 2 h. Excess thionyl chloride was distilled off under reduced pressure to obtain the intended compound in the form of a colorless oil almost quantitatively.


The same procedure as above was repeated except that (\pm)-N-benzoylnipecotic acid was repeated with (\pm)-N-benzoyl- β -proline to obtain the following compound.

(\pm)-N-Benzoyl- β -prolyl chloride

The acid chloride thus obtained in Referential Example 2 was used in the subsequent reaction (Referential Example 3) without purification.


Step III

REFERENTIAL EXAMPLE 3



22

-continued



14.40 g (0.108 mol) of aluminum chloride was suspended in 25 ml of methylene chloride. 5.50 g (0.029 mol) of 1-acetylisonipecotoyl chloride and 5.00 g (0.029 mol) of methanesulfonamide were added to the suspension under stirring and the obtained mixture was refluxed for 2 h. After cooling, the liquid reaction mixture was poured into 100 ml of ice/water and the mixture was stirred violently. Crystals thus formed were recovered by filtration and dried to obtain 7.22 g of the intended compound.

Melting point: 210° to 211.5° C.

NMR (90 MHz, DMSO- d_6) δ 1.20-2.00 (4H, m), 2.00 (3H, s), 2.60-4.00 (4H, m), 3.10 (3H, s), 4.36 (1H, broad), 7.28 (2H, d, J=8 Hz), 10.34 (1H, s, D₂O exchange).

The same procedure as above was repeated except that methanesulfonamide was replaced with p-toluenesulfonamide or 1-acetylisonipecotoyl chloride was replaced with (\pm)-N-benzoylnipecotoyl chloride was replaced with (\pm)-N-benzoyl- β -prolyl chloride prepared in Referential Example 2 to obtain the following compounds:

1-acetyl-4-(4-ethylsulfonylamino)benzoyl)piperidine

NMR (90 MHz, CDCl₃) δ : 1.35 (3H, t, J=7 Hz), 1.5-2.10 (4H, m), 2.11 (3H, s), 2.65-3.70 (3H, m), 3.16 (2H, q, J=7 Hz), 3.88 (1H, brd, J=12 Hz), 4.51 (1H, br, J=12 Hz), 7.28 (2H, d, J=8 Hz), 7.83 (2H, d, J=8 Hz), 8.60 (1H, brs, D₂O exchange).

1-acetyl-4-(2-hydroxy-4-methylsulfonylamino)benzoyl)piperidine

NMR (90 MHz, CDCl₃) δ : 1.2-2.0 (4H, m), 1.99 (3H, s), 3.10 (3H, s), 4.16 (1H, brd, J=13 Hz), 6.62-6.80 (2H, m), 7.90 (1H, d, J=8 Hz), 10.34 (1H, s, D₂O exchange), 13.22 (1H, s, D₂O exchange).

1-acetyl-4-(2-methoxy-4-methylsulfonylamino)benzoyl)piperidine

NMR (90 MHz, DMSO- d_6) δ : 1.2-2.0 (4H, m), 2.00 (3H, s), 3.12 (3H, s), 3.88 (3H, s), 6.86 (1H, dd, J=8.2 Hz), 6.96 (1H, d, J=2 Hz), 7.57 (1H, d, J=8 Hz), 10.34 (1H, s, D₂O exchange).

1-acetyl-4-(4-p-toluenesulfonylamino)benzoyl)piperidine

NMR (90 MHz, CDCl₃) δ 1.4-2.0 (4H, m) 2.14 (3H, s) 2.37 (3H, s) 2.5-3.6 (3H, m) 3.92 (1H, brd, J=14 Hz) 4.57 (1H, brd, J=14 Hz) 7.23 (4H, d, J=8 Hz) 7.75 (2H, d, J=8 Hz) 7.83 (2H, J=8 Hz) 8.80 (1H, br).

(\pm)-1-benzoyl-3-(4-methylsulfonylamino)benzoyl)piperidine

NMR (90 MHz, DMSO- d_6) δ 1.5-2.2 (4H, m) 3.00 (3H, s) 7.24 (2H, m) 7.42 (5H, s) 7.88 (2H, m) etc.

(\pm)-1-benzoyl-3-(4-methylsulfonylamino)benzoyl)pyrrolidine

NMR (90 MHz, CDCl₃) δ 1.9-3.0 (3H, m) 3.01 (3H, s) 3.4-4.2 (4H, m) 7.34 (7H, m) 7.84 (2H, m)

REFERENTIAL EXAMPLE 4

1-Acetyl-N-methyl-4-(4-methylsulfonylaminobenzoyl)-piperidine


3.24 g (10.0 mmol) of 1-acetyl-4-(4-methylsulfonylaminobenzoyl)piperidine was added to a suspension of 0.29 g (12.0 mmol) of sodium hydride in 30 ml of dimethylformamide in a nitrogen atmosphere and the mixture was stirred at 60° C. for 20 min. 1.7 g (12.0 mmol) of methyl iodide was added to the liquid mixture and the obtained mixture was stirred at 60° C. for 1 h. 100 ml of chloroform was added thereto and the mixture was washed with water and saturated aqueous common salt solution. The chloroform layer was concentrated and the residual oil was subjected to column chromatography (chloroform:methanol=99:1) to concentrate the intended fraction. 2.0 g of the intended compound was obtained.

Melting point (°C.): 162 to 163.

NMR (90 MHz, CDCl₃)δ: 1.5~2.1 (4H, m) 2.11 (3H, s), 2.95 (3H, s), 3.37 (3H, s), 3.92 (1H, br, J=13 Hz), 4.56 (1H, d, J=13 Hz), 7.50 (2H, d, J=8 Hz), 7.96 (2H, d, J=8 Hz).

Step IV

REFERENTIAL EXAMPLE 5



7.50 g (23.1 mmol) of 1-acetyl-4-(4-methylsulfonylaminobenzoyl)piperidine obtained in Referential Example 3 was dissolved in 110 ml of dichloroethane. 11.0 ml of triethylsilane and 17.8 ml of trifluoroacetic acid were added to the solution and the mixture was refluxed for 50 h. After cooling, a 20% aqueous sodium hydroxide solution was added thereto to neutralize the same. After extraction with dichloromethane, the organic layer was washed with water and a saturated aqueous common salt solution, dried over magnesium sulfate and concentrated. The obtained oily residue was purified according to silica gel column chromatography (chloroform: methanol=98:2) to obtain 3.30 g (yield: 46%) of the intended compound in the form of white crystals.


Melting point: 145° to 146° C.

NMR (90 MHz, CDCl₃)δ: 1.4~2.0 (3H, m) 2.08 (3H, s) 2.2~3.1 (4H, m) 3.00 (3H, s) 3.78 (1H, brd, J=13 Hz) 4.56 (1H, brd, J=13 Hz) 7.12 (4H, m).

Elementary analysis for C ₁₅ H ₂₂ N ₂ O ₃ S:			
	C	H	N
calculated (%)	58.04	7.14	9.02
found (%)	57.64	6.93	9.00

Step V. Part 1

EXAMPLE 1



43.4 g (0.142 mol) of 1-acetyl-4-(4-methylsulfonylaminobenzoyl)piperidine was suspended in 1 l of 3N hydrochloric acid and the suspension was stirred under reflux for 3 h. After completion of the reaction, the liquid reaction mixture was cooled and white crystals thus formed were filtered, washed with water and dried to obtain 37.8 g (yield: 84%) of the intended compound.

Melting point: >265° C. (decomp.).

NMR (90 MHz, DMSO-d₆)δ: 1.6~2.1 (4H, m), 3.12 (3H, s) 7.33 (2H, d, J=8 Hz), 8.01 (2H, d, J=8 Hz), 8.8~9.5 (2H, br, D₂O exchange), 10.46 (1H, s, D₂O exchange).

Elementary analysis for C ₁₅ H ₁₈ N ₂ O ₃ S.HCl:			
	C	H	N
calculated (%)	48.95	6.01	8.79
found (%)	48.64	5.77	8.65

The same procedure as above was repeated except that the same starting materials as in the above referential examples were used to obtain the following compounds.

EXAMPLE 2

4-(4-Ethylsulfonylaminobenzoyl)piperidine hydrochloride

Melting point (°C.): >220 (decomp).

NMR (90 MHz, DMSO-d₆)δ: 1.22 (3H, t, 7 Hz), 1.62~2.1 (4H, m), 2.8~3.9 (4H, m), 3.21 (3H, q, J=7 Hz), 7.34 (2H, d, J=8 Hz), 8.01 (2H, d, J=8 Hz), 8.8~9.5 (2H, br), 10.38 (1H, s, D₂O exchange).

Elementary analysis for C ₁₄ H ₂₀ N ₂ O ₃ S.HCl:			
	C	H	N
calculated (%)	50.52	6.06	8.42
found (%)	50.31	6.30	8.29

EXAMPLE 3

4-(2-Hydroxy-4-methylsulfonylaminobenzoyl)piperidine hydrochloride

Melting point (°C.): >250.

NMR (90 MHz, DMSO-d₆)δ: 1.6~2.1 (4H, m), 3.10 (3H, s), 6.65~6.87 (2H, m), 7.89 (1H, d, J=8 Hz), 8.6~9.4 (2H, br, D₂O exchange), 10.40 (1H, s, D₂O exchange), 12.05 (1H, s, D₂O exchange).

Elementary analysis for $C_{13}H_{18}N_2O_4S.HCl$:			
	C	H	N
calculated (%)	46.64	5.72	8.37
found (%)	46.71	5.97	8.30

EXAMPLE 4

4-(2-Methoxy-4-methylsulfonylaminobenzoyl)piperidine hydrochloride


Melting point ($^{\circ}C$): >220 (decomp.).

NMR (90 MHz, $DMSO-d_6$) δ : 1.5~2.1 (4H, m), 3.12 (3H, s), 3.88 (3H, s), 6.86 (1H, dd, $J=8, 2$ Hz), 6.96 (1H, d, $J=2$ Hz), 7.58 (1H, d, $J=8$ Hz), 9.0 (2H, br, D_2O exchange), 10.32 (1H, s, D_2O exchange).

Elementary analysis for $C_{14}H_{20}ON_2O_4S.HCl$:			
	C	H	N
calculated (%)	48.20	5.78	8.03
found (%)	48.32	5.93	7.81

The same procedure as above was repeated except that 1-acetyl-4-(4-methylsulfonylaminobenzoyl)piperidine was replaced with 1-acetyl-4-(4-p-toluenesulfonylaminobenzoyl)piperidine or 1-acetyl-4-(4-methylsulfonylaminobenzoyl)piperidine to obtain the following compounds:

EXAMPLE 5




4-(4-p-toluenesulfonylaminobenzoyl)piperidine hydrochloride

Melting point: 240° to $242^{\circ}C$.

NMR (90 MHz, $DMSO-d_6$) δ : 1.6~2.1 (4H, m) 2.5~3.8 (5H, m) 2.36 (3H, s) 7.24 (2H, d, $J=8$ Hz) 7.36 (2H, d, $J=8$ Hz) 7.76 (2H, d, $J=8$ Hz) 7.90 (2H, d, $J=8$ Hz) 9.0 (2H, br) 10.97 (1H, s).

Elementary analysis for $C_{19}H_{22}N_2O_3S.HCl.H_2O$:			
	C	H	N
calculated (%)	55.27	5.61	6.78
found (%)	55.25	5.68	6.85

EXAMPLE 6



4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride



Melting point: 255° to $257^{\circ}C$.

NMR (90 MHz, $DMSO-d_6$) δ : 1.2~2.0 (5H, m) 2.94 (3H, s) 7.14 (4H, s) 9.0 (2H, br) 9.67 (1H, s).

Elementary analysis for $C_{13}H_{20}N_2O_2S.HCl$:			
	C	H	N
calculated (%)	51.22	6.94	9.19
found (%)	51.26	6.86	9.16

Step V, Part 2

EXAMPLE 7

 (\pm) -3-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride

5.70 g (14.8 mmol) of (\pm) -1-benzoyl-3-(4-methylsulfonylaminobenzoyl)piperidine obtained in Referential Example 3 was dissolved in a mixture of 120 ml of 5N hydrochloric acid and 80 ml of methanol and the obtained solution was refluxed for 8 h. The reaction solution was concentrated and the obtained solid residue was recrystallized from ethanol to obtain 2.61 g (yield: 55%) of the intended compound in the form of white crystals.

Melting point: 235° to $237^{\circ}C$.

NMR (90 MHz, $DMSO-d_6$) δ : 1.4~2.2 (4H, m) 2.6~4.1 (5H, m) 3.11 (3H, s) 7.35 (2H, d, $J=8$ Hz) 7.98 (2H, d, $J=8$ Hz) 8.0~8.5 (2H, br) 10.48 (1H, brs).

Elementary analysis for $C_{13}H_{18}N_2O_3S.HCl$:			
	C	H	N
calculated (%)	48.98	6.01	8.79
found (%)	48.86	5.87	8.77

EXAMPLE 8

The same procedure as in Example 7 was followed except that (\pm) -1-benzoyl-3-(4-methylsulfonylaminobenzoyl)piperidine was replaced with (\pm) -1-benzoyl-3-(4-methylsulfonylaminobenzoyl)pyrrolidine to obtain the following compound:

 (\pm) -3-(4-methylsulfonylaminobenzoyl)pyrrolidine hydrochloride

Melting point: 198° to $200^{\circ}C$.

NMR (90 MHz, $DMSO-d_6$) δ : 1.7~2.5 (2H, m), 3.0~3.8 (4H, m) 3.14 (3H, s) 4.20 (1H, q-like, $J=7$ Hz) 7.36 (2H, d, $J=8$ Hz) 8.01 (2H, d, $J=8$ Hz) 9.5 (2H, br) 10.26 (1H, s).

Elementary analysis for $C_{12}H_{16}N_2O_3S.HCl$:			
	C	H	N
calculated (%)	47.29	5.62	9.19
found (%)	47.17	5.49	9.11

EXAMPLE 9

N-Methyl-4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride

1.43 g (yield: 90%) of the intended compound was obtained from 1.90 g of 1-acetyl-N-methyl-4-(4-methylsulfonylaminobenzoyl)piperidine prepared in Referential Example 4.

Melting point (°C.): 254 to 255.

NMR (90 MHz, DMSO-d₆)δ: 1.5~2.1 (4H, m), 3.04 (3H, s), 3.32 (3H, s), 7.57 (2H, d, J=8Hz), 8.06 (2H, d, J=8Hz), 8.8~9.6 (2H, br, D₂O exchange).

Elementary analysis for C ₁₄ H ₂₀ N ₂ O ₃ ·HCl:			
	C	H	N
calculated (%)	50.50	6.37	8.42
found (%)	50.43	6.42	8.39

EXAMPLE 10

4-(4-Methylsulfonylaminobenzoyl)-1-(4-pyridylmethyl)piperidine dihydrochloride

1.13 g (18.8 mmol) of sodium methoxide was added to a suspension of 3.0 g (9.4 mmol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride and 1.55 g (9.4 mmol) of 4-chloromethylpyridine hydrochloride in 90 ml of acetonitrile. The mixture was stirred at room temperature for 10 min. 2.88 g of potassium carbonate was added to the mixture and the obtained mixture was refluxed for 3 h. After cooling, the liquid reaction mixture was filtered and the filtrate was concentrated, while the residue was purified according to silica gel chromatography (chloroform:methanol=95.5). The product was converted into its hydrochloride with ethanolic hydrogen chloride and recrystallized from methanol/isopropanol to obtain 1.4 g of the intended compound:

Melting point: ~207° C. (decomp.).

NMR (400 MHz, DMSO-d₆)δ: 1.85~2.30 (4H, m), 3.11 (3H, s), 4.53 (2H, s), 7.31 (2H, d, J=8.8 Hz), 7.98 (2H, d, J=8.8 Hz), 8.17 (2H, d, J=4.9 Hz), 8.92 (2H, d, J=5.9 Hz), 10.41 (1H, s, D₂O exchange), 11.6~12.0 (1H, brs, D₂O exchange).

Elementary analysis for C ₁₉ H ₂₃ N ₃ O ₃ ·2HCl:			
	C	H	N
calculated (%)	51.12	5.64	9.41
found (%)	51.04	5.41	9.28

EXAMPLE 11

4-(4-Methylsulfonylaminobenzoyl)-1-[2-(3-pyridyl)ethyl]piperidine dihydrochloride

35 g (0.101 mol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride and 55 g of potassium carbonate were suspended in 700 ml of dimethylformamide and the suspension was stirred at 40° C. for 20 min. 19.6 g (0.101 mol) of 3-(2-chloroethyl)pyridine hydrochloride and 6.0 g (0.036 mol) of potassium iodide were added to the suspension and the mixture was stirred at 90° C. for 3.5 h. The liquid reaction mixture was filtered and the filtrate was concentrated, while the residue was purified according to silica gel chromatography (chloroform:methanol=93.7). The purified product was converted into its hydrochloride with ethanolic

hydrogen chloride and recrystallized from methanol/isopropanol to obtain 13.4 g of the intended compound.

Melting point (°C.): 200 to 203.

NMR (100 MHz, DMSO-d₆)δ: 1.8~2.3 (4H, m), 3.11 (3H, s), 7.32 (2H, d, J=8 Hz), 7.90~8.10 (3H, m), 8.50 (1H, dt, J=6.2 Hz), 8.8 (1H, d, J=6 Hz), 8.93 (1H, d, J=2 Hz), 10.43 (1H, s, D₂O exchange), 10.90~11.40 (1H, br, D₂O exchange).

Elementary analysis for C ₂₀ H ₂₅ N ₃ O ₃ ·2HCl:			
	C	H	N
calculated (%)	52.17	5.91	9.13
found (%)	52.00	5.86	8.83

EXAMPLE 12

4-(4-Methylsulfonylaminobenzoyl)-1-[3-(4-pyridyl)propyl]piperidine dihydrochloride

0.295 g (0.926 mmol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride and 0.380 g (4.52 mmol) of sodium hydrogencarbonate were suspended in 4 ml of dimethylformamide and the suspension was stirred at 85° C. for 40 min. 0.20 g (1.04 mmol) of 4-(3-chloropropyl)pyridine hydrochloride and 0.31 g (1.87 mmol) of potassium iodide were added to the suspension and the mixture was stirred at 85° C. for 1.5 h. The liquid reaction mixture was filtered and the filtrate was concentrated. The obtained residue was purified according to silica gel column chromatography (chloroform:methanol:aqueous ammonia=96:4:0.4). The purified product was converted into its hydrochloride with ethanolic hydrogen chloride and recrystallized from ethanol to obtain 0.288 g (66%) of the intended compound.

Melting point: 230° C. (decomp.).

NMR (100 MHz, DMSO-d₆)δ: 1.8~2.4 (6H, m), 3.10 (3H, s), 7.30 (2H, d, J=8 Hz), 7.95 (2H, d, J=8 Hz), 7.97 (2H, d, J=6 Hz), 8.83 (2H, d, J=6 Hz), 10.44 (1H, brs, D₂O exchange), 10.9~11.4 (1H, br, D₂O exchange).

Elementary analysis for C ₂₁ H ₂₇ N ₃ O ₃ ·2HCl:			
	C	H	N
calculated (%)	53.16	6.16	8.86
found (%)	52.95	6.10	8.73

EXAMPLE 13

1-[2-(6-Methyl-2-pyridyl)ethyl]-4-(4-methylsulfonylaminobenzoyl)piperidine dihydrochloride

0.254 g (0.797 mmol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride, 0.22 g (1.88 mmol) of 6-methyl-2-vinylpyridine and 0.15 g of sodium acetate were suspended in 3 ml of a mixture of methanol and water (1:1) and the suspension was refluxed for 2 h. The liquid reaction mixture was filtered and the filtrate was concentrated. The obtained residue was purified according to silica gel column chromatography (chloroform:methanol:aqueous ammonia=96:4:0.4). The purified product was converted into its hydrochloride with ethanolic hydrogen chloride and recrystallized from ethanol to obtain 0.285 g (yield: 81%) of the intended compound.

Melting point: 219° C. (decomp.).

NMR (90 MHz, DMSO-d₆)δ: 1.6~2.4 (4H, m), 2.74 (3H, s), 3.12 (3H, s), 7.33 (2H, d, J=8 Hz), 7.70 (1H, d,


J=7 Hz) 7.78 (1H, d, J=7 Hz) 8.02 (2H, d, J=8 Hz) 8.33 (1H, t, J=7 Hz) 10.47 (1H, s, D₂O exchange) 11.2 (1H, br, D₂O exchange).

5

drochloride and 4-(2-chloroethyl)pyridine in Example 29 in the same manner as in Example 13.

Step VI. Part 1

EXAMPLE 15



1-[2-(2-Imidazo[1,2-a]pyridyl)ethyl]-4-(4-methyl-sulfonylaminobenzoyl)piperidine

Elementary analysis for C₂₁H₂₇N₃O₃S·2HCl:

	C	H	N
calculated (%)	53.16	6.16	8.86
found (%)	52.94	6.16	8.73

EXAMPLE 14

4-(4-Methylsulfonylaminobenzoyl)-1-[2-(4-pyridyl)ethyl]piperidine dihydrochloride

another process for synthesizing the compound of Example 29

10.0 g (31.4 mmol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride was suspended in 20 ml of an aqueous solution of 1.32 g of sodium hydroxide and the suspension was stirred at room temperature for 1 h. The formed solid was filtered, washed with water and dried to obtain 8.28 g of crystals, which were suspended in a solution comprising 15 ml of water, 15 ml of methanol and 0.2 ml of acetic acid. 3.39 g of 4-vinylpyridine was added to the suspension and the mixture was refluxed for 10 h. After cooling, the formed crystals were filtered and converted into its hydrochloride with ethanolic hydrogen chloride to obtain 7.54 g of the intended compound.

The melting point and NMR signals of this compound coincided with those of the compound produced from 4-(4-methylsulfonylaminobenzoyl)piperidine hy-

A mixture of 1.02 g (3.2 mmol) of 4-(4-methylsulfonylaminobenzoyl)piperidine hydrochloride, 1.34 g of sodium hydrogencarbonate and 10 ml of dimethylformamide was stirred at 80° C. for 1 h. 0.48 g of 2-(2-chloroethyl)imidazo[1,2-a]pyridine hydrochloride and 0.53 g of potassium iodide were added thereto and the mixture was stirred at 80° C. for 2 h. The mixture was filtered and the filtrate was concentrated to obtain a solid residue, which was then purified according to silica gel column chromatography (chloroform:methanol:aqueous ammonia=190:9:1). The fraction of the intended compound was concentrated to obtain a solid residue, which was then recrystallized from ethyl acetate to obtain 0.25 g (yield: 18%) of the intended compound.


Melting point: 190° to 191° C.

NMR (90 MHz, DMSO-d₆)δ1.4-1.9 (4H, m) 2.18 (2H, m) 2.4-3.6 (7H, m) 3.10 (3H, s) 6.80 (1H, dt, J=5.2 Hz) 7.04-7.34 (4H, m) 7.72 (1H, s) 7.95 (2H, d, J=8 Hz) 8.45 (1H, d, J=7 Hz).


Elementary analysis for C₂₂H₂₆N₄O₃S:

	C	H	N
calculated (%)	61.95	6.14	13.14
found (%)	61.92	6.10	12.92

EXAMPLE 16



-continued




1-[2-(3-Imidazo[1,2-a]pyridyl)-1-oxoethyl]-4-(4-methylsulfonylamino)benzoyl piperidine dihydrochloride

i) 22.1 g of 3-acetylimidazo[1,2-a]pyridine was dissolved in 220 ml of acetic acid. 35.1 ml of a 30% hydrogen bromide/acetic acid solution was added dropwise to the solution at 0° C. Then 28.6 g of bromine was added dropwise thereto at 40° C. The mixture was stirred at 40° C. for 2 h and crystals thus formed were filtered. The crystals were dissolved in 100 ml of water. The solution was made alkaline with an excess of an aqueous sodium hydrogencarbonate solution and then extracted with ethyl acetate. The organic layer was

Elementary analysis for $C_{22}H_{24}N_4O_4S \cdot 2HCl \cdot 1.2H_2O$:			
	C	H	N
calculated (%)	49.39	5.35	10.47
found (%)	49.46	5.09	10.41

Step VI, Part 3
EXAMPLE 17



4-(4-Methylsulfonylamino)benzoyl-1-[2-(1-pyrrolidinyl)ethyl]piperidine dioxalate

concentrated and the obtained brown solid was purified according to silica gel column chromatography (eluted with ethyl acetate) to obtain 13.5 g (yield: 40%) of 3-bromoacetylimidazo[1,2-a]pyridine (yield: 40%) in the form of yellow crystals. ii) A suspension comprising 1.91 g (6.0 mmol) of 4-(4-methylsulfonylamino)benzoyl piperidine hydrochloride, 3.0 g of potassium carbonate and 40 ml of dimethylformamide was stirred at 80° C. for 1 h. After cooling to room temperature, 1.99 g of 3-bromoacetylimidazo[1,2-a]pyridine prepared in the above step i) was added thereto and the mixture was stirred at room temperature for 6 h. The reaction mixture was filtered and the filtrate was concentrated to obtain a solid residue, which was then purified according to silica gel column chromatography (chloroform:methanol = 96:4). The purified product was converted into its dihydrochloride with ethanolic hydrogen chloride and recrystallized from methanol/acetone to obtain 1.75 g (yield: 58%) of the intended compound.

Melting point: 176° to 178° C.

NMR (400 MHz, $DMSO-d_6$) δ 1.95-2.10 (4H, m) 3.12 (3H, s) 3.33 (2H, m) 3.48-3.90 (3H, m) 4.96 (2H, s) 7.34 (2H, d, $J=8.8$ Hz) 7.51 (1H, t, $J=7.1$ Hz) 7.89 (1H, dd, $J=7.8, 7.3$ Hz) 8.02 (2H, d, $J=8.8$ Hz) 9.01 (1H, s) 9.54 (1H, d, $J=6.8$ Hz) 10.50 (1H, s, D_2O exchange) 10.72 (1H, br, D_2O exchange).


10.0 g (31.4 mmol) of 4-(4-methylsulfonylamino)benzoyl piperidine hydrochloride was suspended in 20 ml of an aqueous solution of 1.32 g of sodium hydroxide and the suspension was stirred at room temperature for 1 h. The formed crystals were filtered, washed with water and dried to obtain 8.28 g of 4-(4-methylsulfonylamino)benzoyl piperidine in free form. A mixture of 2.0 g (7.09 mmol) of the obtained crystals, 1.57 g of chloroethylpyrrolidine hydrochloride, 2.35 g of potassium iodide and 40 ml of dimethylformamide was stirred at 80° C. for 3 h. The reaction mixture was filtered and the filtrate was concentrated to obtain a solid residue, which was then purified according to silica gel column chromatography (chloroform:methanol:aqueous ammonia = 90:9:1). 0.68 g of the purified product was converted into its dioxalate with 0.32 g of oxalic acid in ethanol. After recrystallization from methanol/ethanol, 0.40 g (yield: 10% of the intended compound) was obtained.

Melting point: 214° to 216° C.

NMR (90 MHz, $DMSO-d_6$) δ 1.4-2.3 (8H, m) 2.4-3.6 (13H, m) 3.08 (3H, s) 7.21 (2H, d, $J=8$ Hz) 7.88 (2H, d, $J=8$ Hz).

Elementary analysis for $C_{19}H_{20}N_2O_3S_2(CCOH)_2$:			
	C	H	N
calculated (%)	49.37	5.94	7.51
found (%)	49.40	5.85	7.37

Step VI, Part 4



1.88 ml of formalin and 1.07 g of methylfuran were added to a mixture of 4.43 g (15.7 mmol) of 4-(4-methylsulfonylamino)benzoyl piperidine in free form obtained in Example 17, 1.57 ml of glacial acetic acid and 10 ml of water and the obtained mixture was stirred at 90° C. for 2 h. After cooling, the mixture was neutralized with a 20% aqueous sodium hydroxide solution and extracted with dichloromethane. The organic layer was washed with water and a saturated aqueous common salt solution, dried over magnesium sulfate and concentrated. The solid residue was recrystallized from ethanol/methanol to obtain 4.16 g (yield: 70%) of the intended compound.

Melting point: 181° to 182° C.

NMR (90 MHz, DMSO- d_6) δ 1.3–1.9 (4H, m) 1.9–2.3 (2H, m) 2.23 (3H, d, J=1 Hz) 2.6–3.4 (3H, m) 3.10 (3H, s) 3.43 (2H, s) 5.97 (1H, m) 6.13 (1H, d, J=3 Hz) 7.28 (2H, d, J=8 Hz) 7.94 (2H, d, J=8 Hz).


Elementary analysis for $C_{19}H_{24}N_2O_4S$:			
	C	H	N
calculated (%)	60.62	6.43	7.44

-continued

Elementary analysis for $C_{19}H_{24}N_2O_4S$:			
	C	H	N
found (%)	60.43	6.46	7.44

Step VI, Part 5

EXAMPLE 19



4-(4-Methylsulfonylamino)benzoyl-1-[2-(2-quinoxaliny)ethyl]piperidine


5.0 g (15.7 mmol) of 4-(4-methylsulfonylamino)benzoyl piperidine hydrochloride was suspended in 5 ml of ethanol. 2.49 g of 2-methylquinoxaline and 7.0 ml of formalin were added to the suspension and the mixture was stirred at 90° C. for 1 h. After cooling, the mixture was neutralized with a 20% aqueous sodium hydroxide solution and the formed crystals were recrystallized from ethyl acetate to obtain 0.32 g (yield: 5%) of the intended compound.

Melting point: 156° to 157° C.

NMR (90 MHz, DMSO- d_6) δ 1.4–2.0 (4H, m) 2.0–2.4 (2H, m) 2.6–3.5 (7H, m) 3.13 (3H, s) 7.31 (2H, d, J=8 Hz) 7.73–8.15 (6H, m) 8.91 (1H, s)

Elementary analysis for $C_{23}H_{26}N_4O_5S$:			
	C	H	N
calculated (%)	62.99	5.98	12.78
found (%)	62.83	5.95	12.61

Step VI, Part 6



(Fraction 2)

(Fraction 1)

EXAMPLE 20 AND 21

1-Ethyl-4-(4-methylsulfonylamino benzoyl)piperidine
and
1-ethyl-4-(N-ethyl-4-methylsulfonylamino benzoyl)-
piperidine

A suspension comprising 2.54 g (7.97 mmol) of 4-(4-methylsulfonylamino benzoyl)piperidine hydrochloride, 5.0 g of potassium carbonate and 40 ml of dimethylformamide was stirred at 80° C. for 1 h. 1.3 g (8.3 mmol) of ethyl iodide was added thereto and the mixture was stirred at 80° C. for 12 h. The mixture was filtered and the filtrate was concentrated to obtain a solid residue, which was then purified according to silica gel column chromatography (chloroform:methanol:aqueous ammonia = 190:9:1). The respective fractions were concentrated to obtain a solid residue, which was then converted into its hydrochloride with ethanolic hydrochloric acid and recrystallized from methanol/ethyl acetate to obtain the intended compound.

EXAMPLE 20

1-Ethyl-4-(N-ethyl-4-methylsulfonylamino benzoyl)-
piperidine hydrochloride: 0.23 g

Melting point: 188° to 191° C.

NMR (90 MHz, DMSO-d₆) 1.04 (3H, t, J=7H2) 1.28 (3H, t, J=7H2) 1.65-2.30 (4H, m) 2.60-3.95 (7H, m) 3.05 (3H, s) 3.78 (2H, q, J=7H2) 7.57 (2H, d, J=8H2) 8.06 (2H, d, J=8H2).

Elementary analysis for C ₁₇ H ₂₆ N ₂ O ₃ S.HCl:			
	C	H	N
calculated (%)	54.45	7.27	7.47
found (%)	54.20	7.09	7.24

EXAMPLE 21

1-Ethyl-4-(4-methylsulfonylamino benzoyl)piperidine
hydrochloride: 1.70 g

Melting point: 204° to 207° C.

NMR (90 MHz, DMSO-d₆) 1.27 (3H, t, J=7H2) 1.64-2.23 (4H, m) 2.62-3.90 (7H, m) 3.13 (3H, s) 7.34 (2H, d, J=8H2) 8.01 (2H, d, J=8H2) 10.42 (1H, brs).

Elementary analysis for C ₁₅ H ₂₂ N ₂ O ₃ S.HCl:			
	C	H	N
calculated (%)	51.93	6.70	8.08
found (%)	51.76	6.57	7.86

EXAMPLE 22

N-Methyl-4-(4-methylsulfonylamino benzoyl)-1-[2-(3-
pyridyl)ethyl]piperidine dihydrochloride

1.5 g (3.26 mmol) of 4-(4-methylsulfonylamino benzoyl)-1-[2-(3-pyridyl)ethyl]piperidine dihydrochloride was added to a suspension of 0.242 g (10.08 mmol) of sodium hydride in 30 ml of dimethylformamide and the mixture was stirred at 60° C. for 5 h. 0.56 g (3.94 mmol) of methyl iodide was added thereto at room temperature and the obtained mixture was stirred at that temperature for 2 h. The liquid reaction mixture was filtered and the filtrate was concentrated to obtain a residue, which was purified according to silica gel chromatography (chloroform:methanol=97:3). This product was converted into its hydrochloride with ethanolic

hydrogen chloride to obtain 0.5 g of the intended compound.

Melting point (°C.): ~182

NMR (90 MHz, DMSO-d₆)δ: 1.8~2.4 (4H, m), 3.02 (3H, s), 3.31 (3H, s), 7.48 (2H, d, J=8 Hz), 7.98 (3H, m), 8.45 (1H, brd, J=7 Hz), 8.80 (2H, m).

Elementary analysis for C ₂₁ H ₂₇ N ₃ O ₃ S.2HCl:			
	C	H	N
calculated	53.16	6.16	8.86
found	53.37	6.12	8.65

EXAMPLE 23

N-Butyl-4-(4-methylsulfonyl benzoyl)-1-[2-(3-pyridyl)-
ethyl]piperidine dihydrochloride

The same procedure as in Example 22 was repeated except that methyl iodide was replaced with n-butyl iodide and the obtained product was purified according to silica gel chromatography to obtain the intended compound having the following physical properties:

Melting point (°C.): 110 to 111.

NMR (90 MHz, CDCl₃)δ: 0.88 (3H, t), 1.2~3.3 (19H, m), 2.88 (3H, s), 3.73 (2H, t, J=8 Hz), 7.2 (1H, q, J=6.6 Hz), 7.46 (2H, d, J=8 Hz), 7.5~7.64 (1H, m), 7.98 (2H, d, J=8 Hz), 8.46 (1H, dd, J=2, 6 Hz), 8.5 (1H, d, J=2 Hz).

Elementary analysis for C ₂₄ H ₃₃ N ₃ O ₃ S:			
	C	H	N
calculated (%)	64.97	7.51	9.47
found (%)	64.90	7.41	9.43

EXAMPLE 2

4-(4-methylsulfonylamino phenyl)hydroxymethyl-1-(2-
(2-pyridyl)ethyl)piperidine dihydrochloride

2.0 g (5.16 mmol) of 4-(4-methylsulfonylamino benzoyl)-1-[2-(2-pyridyl)ethyl]piperidine was dissolved in 150 ml of methanol. 0.39 g of sodium borohydride was added to the solution under cooling with ice and the mixture was stirred at 0° C. for 1 h. The liquid reaction mixture was acidified by adding ethanolic hydrogen chloride dropwise thereto at 0° C. and an inorganic matter thus formed was filtered out. The filtrate was concentrated and the resulting residual oil was dissolved in ethanol. The solution was made alkaline with aqueous ammonia and an inorganic matter thus formed was further filtered out. The filtrate was concentrated and the obtained residual oil was purified according to silica gel chromatography (chloroform:methanol:aqueous ammonia = 90:9:1). The oil thus obtained was converted into its hydrochloride in an ordinary manner to obtain 0.76 g of the intended compound.

Melting point (°C.): ~182.

NMR (90 MHz, DMSO-d₆)δ: 1.3~2.4 (4H, m), 2.92 (3H, s), 4.22 (1H, br), 7.23 (4H, m), 7.6~7.9 (2H, m), 8.28 (1H, dt, J=1, 7 Hz), 8.65 (1H, d, J=5 Hz), 9.64 (1H, s, D₂O exchange), 10.8 (1H, br, D₂O exchange).

Elementary analysis for C ₂₁ H ₂₆ N ₃ O ₃ S.2HCl:			
	C	H	N
calculated (%)	52.94	6.56	8.82

-continued

Elementary analysis for $C_{21}H_{26}N_2O_3S \cdot 2HCl$:			
	C	H	N
found (%)	53.16	6.83	8.61

EXAMPLE 25

N-Methyl-4-(4-methylsulfonylaminobenzoyl)-1-[2-(3-pyridyl)ethyl]piperidine dihydrochloride

another process for synthesizing the compound of Example 22

0.8 g (2.7 mmol) of N-methyl-4-(4-methylsulfonylaminobenzoyl)piperidine dihydrochloride, 0.52 g (2.7 mmol) of 3-(2-chloroethyl)pyridine hydrochloride, 3.0 g of potassium carbonate and 0.2 g of potassium iodide were dissolved in 15 ml of dimethylformamide and the solution was stirred at 90° C. for 3.5 h. After

cooling, an inorganic matter was filtered out and the filtrate was concentrated to obtain a residue, which was purified according to silica gel chromatography (chloroform:methanol=97.3). The purified product was converted into its hydrochloride in an ordinary manner to obtain 0.6 g of the intended compound. The melting point and NMR signals of this product coincided with those of the compound obtained in Example 22.

EXAMPLE 26 TO 115

Each compound listed in Tables 3 to 12 was obtained from the piperidine compound, the pyrrolidine compound and a halide compound thereof shown in Referential Examples 3 to 5 in the same way as shown in Examples 1 to 25.

A variety of compounds of the invention was obtained, including a heterocyclic ring or a condensed aromatic ring.

5

10

15

20

25

30

35

40

45

50

55

60

65

TABLE 3








Example	Y	m.p. (°C.)	Molecular formula	Elementary analysis calculated: (upper column) found: (lower column)	NMR		
					C	H	
26		~198	$C_{10}H_{12}N_3O_3 \cdot 2HCl \cdot 3H_2O$	49.14 48.96	5.86 5.84	9.05 8.89	(400MHz, DMSO-d ₆) δ: 1.9-2.1(4H, m) 3.11(3H, s) 3.21(2H, m) 3.46(2H, m) 3.66(1H, m) 4.48(2H, s) 7.31(2H, d, J=8.8) 7.52(1H, dd, J=7.8, 4.9) 7.68(1H, d, J=7.8) 7.95(1H, dd, J=7.5, 1.5) 7.98(2H, d, J=8.8) 8.70(1H, d, J=4.4) 10.41(1H, s)
27		~230	$C_{10}H_{12}N_3O_3 \cdot 2HCl \cdot 3H_2O$	49.78 49.81	5.79 5.85	9.17 9.02	(400MHz, DMSO-d ₆) δ: 1.85-2.25(4H, m) 3.11(3H, s) 4.48(2H, s) 7.31(2H, d, J=8.8) 7.88(1H, dd, J=5.7, 7.6) 7.99(2H, d, J=8.8) 8.55(1H, d, J=7.3) 9.05(1H, s) 10.41(1H, s) 11.41(1H, br)
28		~180	$C_{20}H_{25}N_3O_3 \cdot 2HCl \cdot 3H_2O$	50.85 50.88	6.04 6.02	8.89 8.81	(400MHz, DMSO-d ₆) δ: 1.9-2.1(4H, m) 3.12(3H, s) 3.18(2H, m) 7.32(2H, d, J=8.8) 7.66(1H, t, J=6.2) 7.75(1H, d, J=7.8) 7.99(2H, d, J=8.8) 8.21(1H, t, J=7.3) 8.71(1H, d, J=4.9) 10.42(1H, s, D ₂ O exchange) 10.84 (1H, br, D ₂ O exchange)
29		~260	$C_{20}H_{25}N_3O_3 \cdot 2HCl$	52.17 52.25	5.91 5.91	9.13 9.05	(100MHz, DMSO-d ₆) δ: 1.8-2.3(4H, m) 3.13(3H, s) 7.32(2H, d, J=8) 7.99(4H, m) 8.88(2H, d, J=6) 10.46(1H, s, D ₂ O exchange) 11.2(1H, br, D ₂ O exchange)
30		~200	$C_{21}H_{27}N_3O_3 \cdot 2HCl$	53.16 52.95	6.16 6.02	8.86 8.84	(90MHz, DMSO-d ₆) δ: 1.7-2.4(6H, m) 3.10(3H, s) 7.24(2H, d, J=8) 7.90(2H, d, J=8) 7.95(1H, m) 8.42(1H, d, J=6) 8.71(1H, d, J=4) 8.81(1H, br) 10.36(1H, s, D ₂ O exchange) 10.9(1H, br, D ₂ O exchange)
31		~230	$C_{21}H_{27}N_3O_3 \cdot 2HCl$	53.16 52.95	6.16 6.10	8.86 8.73	(100MHz, DMSO-d ₆) δ: 1.8-2.4(6H, m) 3.10(3H, s) 7.30(2H, d, J=8) 7.95(2H, d, J=8) 7.97(2H, d, J=6) 8.83(2H, d, J=6) 10.44(1H, br, D ₂ O exchange) 10.9-11.4(1H, br, D ₂ O exchange)
32		~190	$C_{22}H_{29}N_3O_3 \cdot 2HCl \cdot H_2O$	52.17 52.15	6.57 6.34	8.30 8.27	(90MHz, DMSO-d ₆) δ: 1.5-2.2(8H, m) 3.10(3H, s) 7.27(2H, d, J=8) 7.82-8.06(3H, m) 8.50(1H, d, J=7) 8.74(1H, d, J=5) 8.84(1H, br)

TABLE 3-continued







Example	Y	m.p. (°C)	Molecular formula	Elementary analysis		NMR	
				calculated: found:	found: (lower column)	C	H
33		217 ~	C ₂₃ H ₃₁ N ₃ O ₃ ·2HCl	54.96	6.63	8.36	(90MHz, DMSO-d ₆) δ: 1.1-2.3(10H, m) 3.12(3H, s)
				54.70	6.54	8.21	7.34(2H, d, J = 7) 7.85-8.10(1H, m) 8.00(2H, d, J = 7) 8.46(1H, d, t, J = 2.7) 8.79(1H, dt, J = 2.5) 8.86(1H, d, J = 2) 10.48(1H, s, D ₂ O exchange) 10.90(1H, bs, D ₂ O exchange)
34		~140	C ₂₀ H ₂₃ N ₃ O ₄ S·2HCl·1.2H ₂ O	48.23	5.95	8.43	(90MHz, DMSO-d ₆) δ: 1.6-2.3(4H, m) 3.12(3H, s)
				48.07	5.41	8.10	7.32(2H, d, J = 8) 8.01(2H, d, J = 8) 7.97(1H, m) 8.54(1H, br, J = 6) 8.88(2H, m) 10.44(1H, br, D ₂ O exchange)
35			C ₂₀ H ₂₁ N ₃ O ₄ S	59.83	5.77	10.47	(90MHz, CDCl ₃) δ: 1.6-2.2(4H, m) 3.06(3H, s)
				60.15	5.90	10.18	3.80(2H, s) 7.30(2H, d, J = 8) 7.41(1H, m) 7.92(2H, d, J = 8) 8.32(1H, dt, J = 2.7) 8.66(1H, dd, J = 5.2) 9.24(1H, d, J = 2)
36		~175	C ₂₀ H ₂₅ N ₃ O ₃ S·2HCl·3H ₂ O	50.85	6.04	8.89	(90MHz, DMSO-d ₆) δ: 1.75(3H, d, J = 7) 1.7-2.4(4H, m)
				50.70	5.78	8.54	3.11(3H, s) 4.76(1H, br) 7.24(2H, d, J = 8) 7.89(2H, d, J = 8) 8.20(2H, d, J = 5) 8.90(2H, d, J = 5) 10.33(1H, s, D ₂ O exchange) 11.9(1H, br, D ₂ O exchange)
37		~195	C ₂₅ H ₂₇ N ₃ O ₃ S·2HCl	57.47	5.59	8.04	(90MHz, DMSO-d ₆) δ: 1.7-2.4(4H, m) 3.10(3H, s)
				57.33	5.62	8.31	6.13(1H, br, D ₂ O sharpen) 7.24(2H, d, J = 8) 7.30(3H, m) 7.93(4H, d, J = 8) 8.57(2H, d, J = 5) 8.90(2H, d, J = 5) 10.40(1H, s, D ₂ O exchange)
38		~140	C ₂₂ H ₂₇ N ₃ O ₃ S·2HCl	54.32	5.59	8.64	(90MHz, DMSO-d ₆) δ: 1.7-2.4(4H, m) 3.12(3H, s)
				54.29	5.61	8.49	7.32(2H, d, J = 8) 7.48(1H, dt, J = 6.8) 8.00(3H, d, J = 8) 8.52(1H, d, J = 6) 10.43(1H, s)

TABLE 3-continued








Example	Y	m.p. (°C.)	Molecular formula	Elementary analysis calculated: (upper column) found: (lower column)		NMR		
				C	H	N	C	H
39		~200	C ₂₁ H ₂₅ N ₃ O ₃ S ₂ ·2HCl	53.39 53.41	5.76 5.65	8.89 8.70	(90MHz, DMSO-d ₆) δ: 1.7-2.4(4H, m) 3.09(3H, s) 6.92(2H, m) 7.25(2H, d, J=8) 7.92(3H, m) 8.57(1H, m) 8.82(1H, d, J=5) 9.00(1H, s)	
40		~195	C ₂₀ H ₂₅ N ₃ O ₃ S ₂ ·2HCl	47.06 47.20	5.73 5.51	8.23 8.05	(90MHz, DMSO-d ₆) δ: 1.7-2.4(4H, m) 3.09(3H, s) 7.26(2H, d, J=8) 7.94(2H, d, J=8) 8.02(2H, d, J=6) 8.60(2H, d, J=6)	
41		178~180	C ₂₀ H ₂₅ N ₃ O ₄ S	61.99 61.69	6.50 6.34	10.84 10.57	(90MHz, CDCl ₃) δ: 1.52-2.32(5H, m) 2.55(3H, s) 2.80-3.25(4H, m) 3.09(3H, s) 3.51(2H, s) 7.14(1H, d, J=8), 7.31(2H, d, J=8.4) 7.60(1H, dd, J=8, 2.2) 7.91(2H, d, J=8.4) 8.39(1H, d, J=2.2)	
42		181~183	C ₁₉ H ₂₂ ClN ₃ O ₃ S	55.94 55.88	5.44 5.47	10.30 9.97	(90MHz, CDCl ₃) δ: 1.60-2.30(6H, m) 2.74-3.38(3H, m) 3.10(3H, s) 3.52(2H, s) 7.26(2H, d, J=8.8) 7.30(1H, d, J=8) 7.69(1H, dd, J=8, 2.2) 7.94(2H, d, J=8.8) 8.30(1H, d, J=2.2)	
43		~150	C ₂₀ H ₂₄ N ₂ O ₃ S ₂ ·HCl·3H ₂ O	57.07 56.83	6.30 6.20	6.65 6.45	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 3.12(3H, s) 4.29(2H, s) 7.30(2H, d, J=8) 7.28-7.70(5H, m, D ₂ O sharpen) 7.99(2H, d, J=8) 10.42(1H, br, D ₂ O exchange) 10.80(1H, br, D ₂ O exchange)	
44		222~224	C ₂₁ H ₂₆ N ₂ O ₃ S ₂ ·HCl	59.63 59.61	6.43 6.36	6.62 6.55	(100MHz, DMSO-d ₆) δ: 1.8-2.3(4H, m) 3.12(3H, s) 7.28(5H, s) 7.32(2H, d, J=8) 7.96(2H, d, J=8) 10.3-11.0(2H, br, D ₂ O exchange)	
45		215~217	C ₂₂ H ₂₈ N ₂ O ₃ S ₂ ·C ₇ H ₂ O ₄	58.76 58.97	6.16 6.16	5.71 5.82	(100MHz, DMSO-d ₆) δ: 1.6-2.2(4H, m) 3.12(3H, s) 7.24(5H, m) 7.31(2H, d, J=8) 7.97(2H, d, J=8) 8.35(2H, br)	

TABLE 3-continued

Example	Y	m.p. (°C.)	Molecular formula	Elementary analysis			NMR
				calculated: found:	calculated: found:	calculated: found:	
				C	H	N	
46		~193	$C_{20}H_{25}N_2O_3S.HCl$	54.18 54.31	5.40 5.48	6.32 5.92	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 3.12(3H, s) 4.45(2H, s, D ₂ O sharpen) 7.30(2H, d, J=8) 7.40-7.50 (4H, m) 7.99(3H, m) 10.45(1H, br, D ₂ O exchange) 11.2(1H, br, D ₂ O exchange)
47		~220	$C_{12}H_{25}ClN_2O_3S.HCl$	55.14 55.43	5.73 5.68	6.12 5.74	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 3.12(3H, s) 7.20-7.50(6H, m, D ₂ O sharpen → 7.39(4H, s) & 7.52 (2H, d, J=8) 8.00(2H, d, J=8) 10.0-11.0(1H, br, D ₂ O exchange)
48		~220	$C_{21}H_{25}FN_2O_3S.HCl.H_2O$	56.43 56.43	5.86 5.99	6.27 6.42	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 3.12(3H, s) 7.00-7.45(5H, m) 8.00(2H, d, J=8) 10.0-11.5(2H, br, D ₂ O exchange)
49		~230	$C_{22}H_{26}N_2O_3S.HCl$	60.47 60.78	6.69 6.72	6.41 5.84	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 2.27(3H, s) 3.12(3H, s) 7.16(4H, s) 7.34(2H, d, J=8.0) 8.01(2H, d, J=8) 10.50(1H, br, D ₂ O exchange) 10.5-11.5(1H, br, D ₂ O exchange)
50		~225	$C_{22}H_{26}N_2O_4S.HCl$	58.33 58.48	6.45 6.40	6.18 5.68	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m) 3.12(3H, s) 3.74(3H, s) 6.88(2H, d, J=8) 7.20(2H, d, J=8) 7.32(2H, d, J=8) 8.01(2H, d, J=8) 10.0-11.0(2H, br, D ₂ O exchange)
51		190~191	$C_{21}H_{30}N_2O_5S.HCl$	57.19 57.27	6.47 6.43	5.80 5.77	(90MHz, DMSO-d ₆) δ: 1.8-2.4(4H, m) 3.12(3H, s) 3.72(3H, s) 3.75(3H, s) 6.7-7.0(3H, m) 7.32(2H, d, J=8) 8.00(2H, d, J=8) 10.6(1H, br)

TABLE 3-continued

Example	Y	Molecular formula	m.p. (°C.)	Elementary analysis calculated: (upper column) found: (lower column)	NMR		
					C	H	N
52		$C_{21}H_{26}N_2O_4S \cdot HCl$	219-221	57.46 57.60	6.20 6.19	6.38 6.46	(100MHz, DMSO-d ₆) δ: 1.8-2.2(4H, m), 3.15(3H, s), 6.63(2H, d, J=8), 7.08(2H, d, J=8), 7.33(2H, d, J=8), 8.00(2H, d, J=8), 9.43(1H, s, D ₂ O exchange)
53		$C_{21}H_{26}N_2O_4S \cdot CH_3SO_3H \cdot 3H_2O$	133-135	52.37 51.99	6.13 5.65	5.55 5.51	(100MHz, DMSO-d ₆) δ: 1.7-2.3(4H, m), 3.12(3H, s), 7.31(2H, d, J=8), 7.42(5H, s), 8.00(2H, d, J=8), 10.38(1H, s)
54		$C_{23}H_{28}N_2O_6S_2 \cdot HCl$	223-226	49.66 49.73	5.68 5.60	7.90 7.81	(90MHz, DMSO-d ₆) δ: 1.7-2.3(4H, m), 3.04(3H, s), 3.08(3H, s), 7.20(4H, d, J=8), 7.52(2H, d, J=8), 7.90(2H, d, J=8), 9.98(1H, s, D ₂ O exchange), 10.2(1H, br, D ₂ O exchange), 10.3(1H, s, D ₂ O exchange)
55		$C_{22}H_{27}N_2O_6S_2 \cdot HCl \cdot H_2O$	~200	48.21 48.13	5.52 5.23	7.67 7.68	(400MHz, DMSO-d ₆) δ: 1.9-2.2(4H, m), 3.12(3H, s), 3.15(3H, s), 5.03(2H, s), 7.34(2H, d, J=8), 7.38(2H, d, J=8), 7.99(2H, d, J=8), 8.02(2H, d, J=8), 10.18(1H, br, D ₂ O exchange), 10.47(1H, s, D ₂ O exchange), 10.64(1H, s, D ₂ O exchange)
56		$C_{18}H_{22}N_2O_3S_2 \cdot CH_3SO_3H$	173-175	48.08 48.24	5.52 5.46	5.90 5.86	(100MHz, DMSO-d ₆) δ: 1.7-2.2(4H, m), 2.38(3H, s), 3.12(3H, s), 4.60(2H, s, D ₂ O sharpen), 7.10-7.40(4H, m), 7.74(1H, d, J=4), 8.00(2H, d, J=8), 9.7(1H, br, D ₂ O exchange), 10.39(1H, br, D ₂ O exchange)
57		$C_{18}H_{24}N_2O_3S_2 \cdot CH_3SO_3H$	219-221	49.16 49.18	5.78 5.66	5.73 5.72	(100MHz, DMSO-d ₆) δ: 1.7-2.2(4H, m), 2.37(3H, s), 3.12(3H, s), 7.01(2H, d, J=4), 7.32(2H, d, J=8), 7.40(1H, t, J=4), 8.01(2H, d, J=8), 10.2(1H, br, D ₂ O exchange)

TABLE 3-continued







Example	Y	m.p. (°C.)	Molecular formula	Elementary analysis		NMR		
				calculated: (upper column) found: (lower column)	C	H	N	
58		220~222	$C_{24}H_{26}N_2O_3S \cdot CH_3SO_3H$	57.90 58.09	5.83 5.72	5.40 4.94		(100MHz, DMSO-d ₆) δ: 1.7-2.1(4H, m) 2.39(3H, s) 3.12(3H, s) 4.93(2H, s) 7.32(2H, d, J = 8) 7.50-8.20 (8H, m) 8.32(1H, brd, J = 7)
59		208~	$C_{22}H_{29}N_3O_3S \cdot 2HCl$	54.10 53.95	6.40 6.39	8.60 8.48		(90MHz, DMSO-d ₆) δ: 1.13(3H, t, J = 8) 2.77) 2.77 (2H, q, J = 8) 3.12(3H, s) 7.34(2H, d, J = 8) 7.87(1H, d, J = 7) 8.02(2H, d, J = 8) 8.33(1H, dd, J = 2.7) 8.70(1H, d, J = 2)
60		172~174	$C_{21}H_{27}N_3O_3S$	62.80 62.88	6.79 6.76	10.47 10.37		(90MHz, CDCl ₃) δ: 2.53(3H, s) 3.10(3H, s) 6.15(1H, s, D ₂ O exchange) 6.97(2H, dd, J = 2.7) 7.27(2H, d, J = 8) 7.48(1H, t, J = 7) 7.94(2H, d, J = 8)
61		~167	$C_{10}H_{25}N_3O_3S \cdot 2HCl$	47.49 47.69	5.67 5.78	8.7 8.27		(90MHz, DMSO-d ₆) δ: 1.7-2.3(4H, m) 2.39(3H, s) 3.10(3H, s) 7.26(2H, d, J = 8) 7.93(2H, d, J = 8) 9.07 (1H, s) 10.37(1H, s, D ₂ O exchange) 11.2(1H, br, D ₂ O exchange)

TABLE 4



Ex- am- ple	R ¹	R ³	X	Y	m.p. (°C.)	Molecular formula	Elementary analysis calculated: (upper column)			NMR
							found: (lower column)			
							C	H	N	
62	C ₂ H ₅	H			~217	C ₂₀ H ₂₈ N ₃ O ₃ S. 2HCl.½H ₂ O	51.50 51.50	5.96 6.15	9.00 8.94	(100MHz, DMSO-d ₆)δ: 1.18 (3H, t, J=7)1.7-2.3(4H, m) 3.18(2H, q, J=7)4.59(2H, s) 7.30(2H, d, J=8)7.94(2H, d, J=8)8.30(2H, d, J=6)8.96(2H, d, J=6)10.41(1H, s, D ₂ O exchange)11.5-12.5(1H, br, D ₂ O exchange)
63	C ₂ H ₅	H				C ₂₂ H ₂₈ N ₂ O ₃ S. HCl.H ₂ O	58.07 57.57	6.87 6.55	6.16 6.49	(100MHz, DMSO-d ₆)δ: 1.20 (3H, t, J=7)1.6-2.2(4H, m) 3.20(2H, q, J=7)7.2-7.6(6H, m) 7.96(2H, d, J=8)10.4(1H, br, D ₂ O exchange)

TABLE 5



Ex- am- ple	R ²	Y	m.p. (°C.)	Molecular formula	Elementary analysis calculated: (upper column)			NMR
					found: (lower column)			
					C	H	N	
64	-OH		~208	C ₁₉ H ₂₃ N ₃ O ₃ S. 2HCl	49.35 49.30	5.45 5.39	9.09 8.95	(100MHz, DMSO-d ₆)δ: 1.8-2.2(4H, m)3.15 (3H, s)4.64(2H, s)6.78(2H, m)7.95(1H, d, J=8)8.27(2H, d, J=6)8.97(2H, d, J=6) 10.55(1H, s, D ₂ O exchange)12.1(2H, br, D ₂ O exchange)
65	-OH		~240	C ₂₁ H ₂₆ N ₂ O ₄ S. HCl.½H ₂ O	56.68 56.82	6.27 6.07	6.29 6.25	(100MHz, DMSO-d ₆)δ: 1.8-2.2(4H, m) 3.16(3H, s)6.77(1H, d, J=8)6.82(1H, s) 7.32(5H, s)7.94(1H, d, J=8)10.54(1H, s, D ₂ O exchange)11.0(1H, br, D ₂ O exchange) 12.13(1H, s, D ₂ O exchange)
66	-OCH ₃		~205	C ₂₀ H ₂₁ N ₃ O ₂ S. 2HCl.½H ₂ O	49.19 49.01	5.80 5.58	8.60 8.11	(100MHz, DMSO-d ₆)δ: 1.7-2.3(4H, m) 3.13(3H, s)3.88(3H, s)4.61(2H, s)6.98(1H, dd, J=7.1)7.97(1H, d, J=1)8.58(1H, d, J=7) 9.35(2H, d, J=6)10.00(2H, d, J=6)

55

60

65

TABLE 6






Example	Y	Molecular formula (m.p. °C.)	Elementary analysis		¹ H-NMR δ (90MHz unless otherwise stated)				
			calculated: (upper column)	found: (lower column)	C	H	N		
67		C ₂₄ H ₂₆ N ₄ O ₄ S·H ₂ O (148-150)	59.49 59.14	5.82 5.69	11.56 11.27				(DMSO-d ₆) 2.2-3.7(5H, m) 3.10(3H, s) 3.85(2H, s) 7.15(1H, brs) 7.28(2H, d, J = 8) 7.75-8.05(5H, m) 8.16(2H, d, J = 8) 8.43(1H, brs)
68		C ₂₁ H ₂₄ N ₄ O ₃ S·HCl (182-184)	56.18 56.15	5.61 5.54	12.48 12.31				(DMSO-d ₆) 1.8-2.4(4H, m) 2.6-4.0(9H, m) 3.12(3H, s) 7.32(2H, d, J = 8) 7.76(1H, dd, J = 8, 5) 8.00(2H, d, J = 8) 8.10(1H, brd, J = 8) 8.68(1H, brd, J = 5) 10.41(1H, brs) 10.7(1H, br)
69		C ₂₂ H ₂₆ N ₄ O ₃ S·HCl·0.2H ₂ O (194-196)	56.63 56.62	5.91 5.88	12.01 11.68				(DMSO-d ₆) 1.8-2.4(4H, m) 2.6-4.0(11H, m) 3.12(3H, s) 7.34(2H, d, J = 8) 7.72(1H, dd, J = 8.5) 7.98(2H, d, J = 8) 8.08(1H, dd, J = 8, 2) 8.65 (1H, dd, J = 5, 2) 10.49(1H, brs) 10.8(1H, br)
70		C ₂₂ H ₂₆ N ₄ O ₃ S·HCl (221-224)	57.07 57.23	5.88 5.77	12.10 11.76				(DMSO-d ₆) 1.8-2.4(4H, m) 2.6-3.9(11H, m) 3.12(3H, s) 7.33(2H, d, J = 8) 7.67(1H, dd, J = 5, 2) 7.99(2H, d, J = 8) 8.02(1H, brs) 8.68(1H, d, J = 5) 10.47(1H, br) 10.80(1H, br)
71		C ₂₂ H ₂₅ N ₅ O ₃ S·3HCl (207-210)	48.12 47.86	5.15 5.33	12.76 12.71				(CDCl ₃ ; free form) 1.62-2.38(6H, m) 2.78-3.60(3H, m) 3.02(3H, s) 3.42 (2H, s) 7.17(1H, s), 7.30(3H, m) 7.32(2H, d, J = 9.0) 7.64(1H, s), 7.88(1H, dd, J = 2.0, 4.5), 7.90(1H, d, J = 9.0) 8.22(1H, s) 8.46 (1H, dd, J = 2.0, 4.2)

TABLE 6-continued








72		$C_{10}H_{25}N_3O_3S$ (144-145)	60.78 60.65	6.71 6.62	11.19 11.17	(DMSO-d ₆) 1.4-2.3(6H, m) 2.86(2H, m) 3.0-3.3(1H, m) 3.09(3H, s) 3.39(2H, s) 3.57(3H, s) 5.85(2H, d-like, J = 2) 6.63(1H, t, J = 2) 7.28(2H, d, J = 8) 7.94(2H, d, J = 8)
73		$C_{10}H_{26}N_4O_3S \cdot 2HCl$ (187-190)	49.23 49.37	6.10 6.31	12.09 12.13	(CDCl ₃ , free form) 1.70-2.40(10H, m) 2.75-3.32(3H, m) 3.04(3H, s) 4.00(2H, t, J = 7.2) 6.02(1H, s) 6.08(1H, s) 7.00(1H, t, J = 8) 7.28(2H, d, J = 8.7) 7.53(1H, s) 7.83(2H, d, J = 8.7)
74		$C_{10}H_{24}N_4O_3S \cdot 0.2H_2O$ (156-157)	58.20 58.24	6.17 6.16	14.29 14.20	(DMSO-d ₆) 1.4-2.0(4H, m) 2.16(2H, m) 2.6-3.6(7H, m) 3.10(3H, s) 7.28(2H, d, J = 8) 7.59(2H, d, J = 3) 7.94(2H, d, J = 8) 8.95(6H, t, J = 3)
75		$C_{20}H_{24}N_4O_3S$ (157-158)	59.98 59.94	6.04 6.01	13.99 13.78	(DMSO-d ₆) 1.4-2.0(4H, m) 2.0-2.4(2H, m) 2.94(2H, m) 3.10(3H, s) 3.32(1H, m) 3.45(2H, s) 5.63(1H, br) 6.33(1H, d, J = 2) 7.28(2H, d, J = 8) 7.74(1H, dd, J = 1, 5) 7.94(2H, d, J = 8) 8.77(1H, t, J = 5) 9.12(1H, d, J = 1)
76		$C_{18}H_{22}N_4O_3S \cdot 2HCl$ (210-212)	48.31 48.46	5.42 5.29	12.53 12.49	(DMSO-d ₆) 1.75-2.35(4H, m) 3.12(3H, s) 2.80-3.85(5H, m) 4.52(2H, s) 7.33(2H, d, J = 8) 7.99(2H, d, J = 8) 8.77(2H, s) 9.00(1H, s) 10.43(1H, s)
77		$C_{10}H_{24}N_4O_3S \cdot 4(COOH)_2$ (195-196)	55.40 55.13	5.82 5.82	12.93 12.75	(DMSO-d ₆) 1.5-2.1(4H, m) 2.2-3.8(9H, m) 3.11(3H, s) 7.30(2H, d, J = 8) 7.96(2H, d, J = 8) 8.43-8.63(3H, m)
78		$C_{22}H_{24}N_4O_3S \cdot HCl \cdot 1.2H_2O$ (210-decomp.)	51.35 51.30	5.37 5.27	10.89 10.70	(DMSO-d ₆) 1.8-2.2(4H, m) 2.66(3H, s) 3.10(3H, s) 4.84(2H, s) 7.32(2H, d, J = 8) 8.00(2H, d, J = 8) 8.68(1H, s) 10.4(1H, s) 12.5-13.5(1H, br)

TABLE 6-continued






Example	Y	Structural formula	Melting point	Elementary analysis (C,H,N) calculated: (upper column) found: (lower column)	¹ H-NMR (90MHz, unless otherwise stated)	
79		$C_{18}H_{22}N_4O_5S$ (232-234)	53.19 53.25	5.46 5.44	13.78 13.75	(DMSO-d ₆) 1.4-2.0(4H, m) 2.30(2H, m) 2.88(2H, m) 3.10(3H, s) 3.26(2H, s) 5.45(1H, s) 7.28(2H, d, J = 8) 7.95(2H, d, J = 8)
80		$C_{23}H_{26}N_3O_3S \cdot HCl \cdot 1.5H_2O$ (165-167)	56.61 56.63	6.19 5.85	8.61 8.42	(100MHz, DMSO-d ₆) 1.8-2.2(4H, m) 3.12 (3H, s) 6.9-7.15(2H, m) 7.15-7.45 (4H, m) 7.63(1H, dd-like, J = 6, 2) 7.99(2H, d, J = 8) 10.5(1H, br) 10.97(1H, br) etc.
81		$C_{23}H_{25}N_3O_3S$ (204-205)	60.64 60.71	5.53 5.56	9.22 9.24	(DMSO-d ₆) 1.64(4H, m) 2.14(2H, m) 2.5-3.6(5H, m) 3.10(3H, s) 3.71(2H, t, J = 7) 7.28(2H, d, J = 8) 7.85 (4H, s) 7.93(2H, d, J = 8)
82		$C_{23}H_{25}N_3O_3S \cdot 2HCl$ (-205 decomp.)	55.65 55.76	5.48 5.57	8.46 8.59	(400MHz, DMSO-d ₆) 2.02(4H, m) 3.11 (3H, s) 3.30(2H, m) 3.50-3.70(3H, m) 4.68(2H, s) 7.31(2H, d, J = 8.8) 7.70(1H, t, J = 6.8) 7.82-7.88(2H, m) 7.98(2H, d, J = 8.8) 8.07(1H, d, J = 8.3) 8.10(1H, d, J = 8.8) 8.53(1H, d, J = 8.3) 10.42(1H, s) 10.83(1H, br)
83		$C_{23}H_{25}N_3O_3S \cdot 2HCl \cdot 0.7H_2O$ (-205 decomp.)	54.27 54.10	5.62 5.54	8.25 8.09	(100MHz, DMSO-d ₆) 1.7-2.4(4H, m) 3.12(3H, s) 7.30(2H, d, J = 8) 7.68-8.44(6H, m) 9.19(1H, brs) 9.48(1H, d, J = 1) 10.47(1H, s) 11.6(1H, br)

TABLE 6-continued






84	176-179		$C_{21}H_{24}N_4O_3 \cdot 2HCl$	51.96	5.40	11.54	(DMSO-d ₆) 1.8-2.4(4H, m) 3.12(3H, s) 4.63(2H, s) 7.32(2H, d, J = 8.7) 4.5 (1H, m) 7.95(4H, m) 8.57(1H, s) 9.00(1H, d, J = 7) etc.
85	220-222		$C_{21}H_{24}N_4O_3 \cdot 2HCl$	51.96 51.74	5.40 5.41	11.54 11.29	(400MHz, DMSO-d ₆) 1.75-2.20(4H, m) 3.11(3H, s) 3.20-3.90(5H, m) 4.50(2H, s) 7.31(2H, d, J = 8.5) 7.97(2H, d, J = 8.5) 8.07(1H, d, J = 9.3) 8.20-8.32(2H, m) 8.46(1H, s) 9.14(1H, s) 10.42(1H, s) 11.6(1H, br)
86	225-257		$C_{21}H_{24}N_4O_3 \cdot S$	61.15 61.10	5.86 6.00	13.58 13.29	(DMSO-d ₆) 1.69(4H, m) 2.0-3.6(5H, m) 3.09(3H, s) 3.75(2H, s) 7.12(2H, dd, J = 6, 3) 7.28(2H, d, J = 8) 7.50 (2H, dd, J = 6, 3) 7.95(2H, d, J = 8)
87	171-173		$C_{22}H_{28}N_4O_3 \cdot S$	54.09 54.17	5.78 5.80	17.20 16.95	(DMSO-d ₆) 1.4-2.0(4H, m) 2.0-3.6(7H, m) 3.10(3H, s) 3.22(3H, s) 3.42(3H, s) 4.36(2H, t, J = 7) 7.28(2H, d, J = 8) 7.94(2H, d, J = 8) 8.03(1H, s)
88	238-240		$C_{26}H_{26}N_2O_3 \cdot S$	69.93 69.59	5.87 5.93	6.27 6.27	(CDCl ₃) 1.60-1.92(4H, m) 2.50-3.18 (5H, m) 3.07(1H, s) 4.87(1H, s) 7.19(2H, d, J = 8.8) 7.20-7.40(4H, m) 7.59-7.76(4H, m)
89	259-262	$-(CH_2)_3CH_3$	$C_{17}H_{26}N_2O_3 \cdot HCl$	54.45 54.54	7.27 7.12	7.47 7.40	(DMSO-d ₆) 0.93(3H, t, J = 7) 1.07-2.35(8H, m) 2.60-3.95(7H, m) 3.12(3H, s) 7.32(2H, d, J = 8) 7.99(2H, d, J = 8) 10.22(1H, brs)

TABLE 6-continued












90		-260 (decomp.)	C ₂₀ H ₃₀ N ₂ O ₃ S ₂ HCl	57.87 57.62	7.54 7.41	6.75 6.54	(DMSO-d ₆)0.162-2.43(15H, m)2.60-3.95 (7H, m)3.12(3H, s)7.33(2H, d, J = 8) 8.00(2H, d, J = 8)10.43(1H, brs)
91		221-224	C ₁₇ H ₂₄ N ₂ O ₃ S ₂ HCl	54.74 54.81	6.77 6.75	7.51 7.50	(DMSO-d ₆)1.80-2.38(4H, m)1.88(3H, s) 2.60-3.80(5H, m)3.09(3H, s) 3.61(2H, s)4.19(2H, s)7.24(2H, d, J = 8)7.90(2H, d, J = 8)10.32(1H, brs)
92		159-162	C ₁₇ H ₂₄ N ₂ O ₃ S ₂ (COOH) ₂	49.76 50.11	5.73 5.71	6.11 6.14	(DMSO-d ₆)1.21(3H, t, J = 7)1.38-2.02 (4H, m)2.50-3.78(5H, m)3.08(3H, s) 3.61(2H, s)4.11(2H, q, J = 7) 7.88(2H, d, J = 8) 7.88(2H, d, J = 8)
93		205-207	C ₁₄ H ₁₇ N ₃ O ₃ S	54.71 54.61	5.57 5.51	13.67 13.51	(DMSO-d ₆)1.5-2.0(4H, m)2.95-3.70 (5H, m)3.08(3H, s)7.21(2H, d, J = 8) 7.88(2H, d, J = 8)10.22(1H, brs)
94		178-180	C ₁₅ H ₁₉ N ₃ O ₃ S	56.06 55.78	5.96 5.83	13.07 12.83	(DMSO-d ₆)1.4-2.0(4H, m)2.2-3.6(5H, m)3.12(3H, s)3.74(2H, s)7.33(2H, d, J = 8)8.02(2H, d, J = 8)
95		201-203	C ₁₇ H ₂₃ N ₃ O ₃ S ₂ HCl	52.91 52.80	6.27 6.13	10.89 10.68	(DMSO-d ₆)1.7-2.4(4H, m)2.4-4.0 (11H, m)3.12(3H, s)7.36(2H, d, J = 8)8.02(2H, d, J = 8)10.48(1H, brs) 10.9(1H, br)
96		194-196	C ₁₉ H ₃₁ N ₃ O ₃ S ₂ (COOH) ₂	48.68 48.60	6.33 6.40	7.40 7.32	(DMSO-d ₆)1.16(6H, t, J = 7)1.4-2.0 (4H, m)3.08(3H, s)7.22(2H, d, J = 8) 7.92(2H, d, J = 8)etc.
97		198-202	C ₂₀ H ₃₁ N ₃ O ₃ S ₂ (COOH) ₂	50.25 50.15	6.15 6.13	7.33 7.21	(DMSO-d ₆)1.3-2.0(10H, m)2.4-3.6 (13H, m)3.08(3H, s)7.21(2H, d, J = 8)7.84(2H, d, J = 8)
98		217-219	C ₁₉ H ₂₉ N ₃ O ₄ S ₂ (COOH) ₂ ·1.3H ₂ O	46.12 46.18	5.99 6.10	7.01 6.77	(DMSO-d ₆)1.6-2.1(4H, m)3.12(3H, s) 7.30(2H, d, J = 8)7.07(2H, d, J = 8)etc.
99		223-225	C ₂₁ H ₃₃ N ₃ O ₃ S ₂ (COOH) ₂	51.10 51.38	6.35 6.53	7.15 7.54	(DMSO-d ₆)1.3-2.4(12H, m)3.12(3H, s) 7.29(2H, d, J = 8)7.96(2H, d, J = 8)etc.


TABLE 7



Structural formula
Elementary analysis
(C.H.N.)
calculated: (upper column)
found: (lower column)


Example	Y	Melting point	Structural formula Elementary analysis	¹ H NMR
100		162-165	C ₂₇ H ₃₁ N ₃ O ₃ S.2HCl.3H ₂ O 53.64 6.50 6.95 53.73 6.14 6.74	(DMSO-d ₆)1.7-2.4(6H, m)2.33(3H, s) 2.7-3.9(9H, m)7.25(2H, d, J=8) 7.36(2H, d, J=8)7.76(2H, d, J=8)7.89(2H, d, J=8)7.99(2H, d, J=6)8.86(2H, d, J=6)11.01 (1H, brs)11.20(1H, br)
101		192-195	C ₂₇ H ₃₁ N ₃ O ₃ S.2HCl.0.2H ₂ O 58.52 6.07 7.58 58.55 6.02 7.40	(DMSO-d ₆)1.8-2.1(4H, m)2.33(3H, s) 2.73(3H, s)2.2-3.9(9H, m)7.24 (2H, d, J=8)7.35(2H, d, J=8) 7.64-7.81(2H, m)7.75(2H, d, J=8) 7.89(2H, d, J=8)8.31(1H, t, J=7) 10.77(1H, brs)11.20(1H, br)

TABLE 8



Example	Y	Melting point	Structural formula Elementary analysis	¹ H NMR
102		156-157	C ₂₀ H ₂₇ N ₃ O ₂ S 64.31 7.29 11.25 64.48 7.22 11.21	(DMSO-d ₆)0.9-2.0(5H, m)2.90(3H, s) 7.18(1H, dd, J=7.5)7.02(4H, s) 7.53(1H, brd, J=7)8.28(2H, m) etc.
103		202-204	C ₂₃ H ₃₂ N ₂ O ₄ S.(COOH) ₂ 57.50 6.56 5.36 57.17 6.50 5.35	(DMSO-d ₆)1.2-2.0(5H, m)2.91(3H, s) 3.67(3H, s)3.69(3H, s)6.56-6.90 (3H, m)7.06(4H, s)etc.

TABLE 9



Example	Y	Melting point	Structural formula Elementary analysis	¹ H NMR
104		133-134	C ₂₀ H ₂₅ N ₃ O ₃ S 61.99 6.50 10.84 61.87 6.47 10.72	(DMSO-d ₆)1.0-2.4(4H, m)2.4-3.8 (9H, m)3.12(3H, s)7.22-7.42(3H, m) 7.48(1H, dt, J=8, 2)7.98(2H, d, J=8)8.43(2H, m)
105		189-191	C ₂₁ H ₂₇ N ₃ O ₃ S.2HCl 53.16 6.16 8.86 52.88 6.10 8.77	(DMSO-d ₆)2.72(3H, s)3.15(3H, s) 7.37(2H, d, J=8)7.71(1H, d, J= 8)7.75(1H, d, J=8)8.04(2H, d, J=8)8.31(1H, t, J=8)10.55 (1H, s)11.40(1H, br)etc.

TABLE 10

Example	Y	Melting point	Structural formula Elementary analysis	¹ H NMR
106		182-184	C ₁₉ H ₂₃ N ₃ O ₃ S 61.11 6.21 11.25 61.13 6.25 11.08	(CDCl ₃ -CD ₃ OD)2.0-2.32(2H, m) 2.4-3.2(8H, m)3.06(3H, s) 3.72-4.08(1H, m)7.16-7.38(3H, m) 7.60(1H, dt, J=8, 2)7.92(2H, d, J=8)8.37(2H, m)
107		151-153	C ₂₀ H ₂₅ N ₃ O ₃ S.2(COOH) ₂ 50.79 5.15 7.40 50.76 5.09 7.19	(DMSO-d ₆)2.47(3H, s)3.14(3H, s) 7.13(2H, d, J=7)7.32(2H, d, J=8) 7.65(1H, t, J=7)8.00(2H, d, J=8)etc.
108		142-143	C ₂₂ N ₂₈ N ₂ O ₅ S 61.09 6.52 6.48 61.02 6.53 6.43	(DMSO-d ₆)1.8-2.2(2H, m)2.4-3.5 (8H, m)3.11(3H, s)3.70(3H, s) 3.72(3H, s) 3.80-4.08(1H, m)6.60-6.90(3H, m) 7.27(2H, d, J=8)7.94(2H, d, J=8)

TABLE 11

Example	Y	Melting point	Structural formula Elementary analysis	¹ H NMR
109	-(CH ₂) ₃ CH ₃	201-204	C ₂₁ H ₃₄ N ₂ O ₃ S.HCl 58.50 8.20 6.50 58.65 8.07 6.43	(DMSO-d ₆)0.65-1.10(6H, m)1.05-2.30 (12H, m)2.60-3.95(9H, m)2.99(3H, s) 7.48(2H, d, J=8)7.96(2H, d, J=8)10.45(1H, brs)
110		-257 (decomp.)	C ₂₇ H ₄₂ N ₂ O ₃ S 63.43 8.50 5.48 63.31 8.38 5.19	(DMSO-d ₆)0.63-2.40(26H, m) 2.60-4.00(9H, m)2.97(3H, s) 7.57(2H, d, J=8)8.05(2H, d, J=8)9.95(1H, brs)
111		138-141	C ₂₁ H ₃₀ N ₂ O ₃ S.(COOH) ₂ 57.47 6.72 5.83 57.68 6.80 5.38	(DMSO-d ₆)1.63(3H, s)1.97(3H, s) 1.60-2.13(4H, m)2.50-3.78(5H, m) 3.04(3H, s)3.43(2H, s)4.28(2H, s) 4.67-4.90(2H, m)5.08(2H, s) 5.38(2H, brs)7.47(2H, d, J=8) 7.92(2H, d, J=8)

TABLE 12


Example	Y	melting point	structural formula	elementary analysis	¹ H NMR
112		~ 193 (decomp)	C ₂₁ H ₂₅ ClN ₂ O ₃ S.HCl	55.14 5.73 6.12 55.40 5.62 6.16	(90MHz, DMSO-d ₆)δ 1.73-2.23(4H, m) 2.67-3.91(9H, m)3.13(3H, s)7.17-7.49 (6H, m)8.00(2H, d, J=8.4)

TABLE 12-continued


Example	Y	melting point	structural formula	elementary analysis	¹ H NMR
113		~205 (decomp)		C ₂₂ H ₂₈ N ₂ O ₃ S.HCl 60.47 6.69 6.41 60.33 6.63 6.52	(90MHz, DMSO-d ₆)δ 1.63-2.20(4H, m) 2.30(3H, s)2.61-3.90(9H, m)3.12(3H, s) 6.96-7.25(4H, m)7.34(2H, d, J=8.8) 8.01(2H, d, J=8.8)
114		178-180	C ₂₄ H ₃₁ N ₃ O ₃ S	65.28 7.08 9.52 64.95 6.87 9.62	(90MHz, DMSO-d ₆)δ 1.35-2.03(8H, m) 2.03-3.40(10H, m)2.98(3H, s)6.97 (1H, dd, J=7.7, 4.6)7.12-7.38(3H, m) 7.81(2H, d, J=8.8)8.26(1H, dd, J=4.6, 1.6)
115		257-259	C ₂₃ H ₂₆ N ₄ O ₅ ·½H ₂ O	57.61 5.67 11.68 57.54 5.47 11.67	(90MHz, DMSO-d ₆)δ 1.38-1.93(4H, m) 1.99-2.37(2H, m) 2.43-3.51(5H, m) 3.11(3H, s)4.04(2H, brt, J=7)7.08-7.32 (4H, m)7.66(1H, t, J=7.5)7.94(3H, d, J=8.8)

What is claimed is:

1. A compound having the formula



in which R¹ is a lower alkyl or a tolyl; R² is hydrogen, hydroxyl, a lower alkoxy or a lower alkyl; R³ is hydrogen, a lower alkyl, a lower alkenyl, a cycloalkyl or a cycloalkylalkyl; X is —CO—, —CH₂— or —CHOH—; g is an integer of 1, 2 or 3; h is an integer of 1, 2 or 3, with the proviso that the sum of g plus h equals the integers 3 or 4; A is (1) alkylene having 1 to 5 carbon atoms, (2) straight-chain alkylene having 1 to 5 carbon atoms in the chain and substituted with lower alkyl, phenyl or hydroxy, (3) straight-chain alkenylene having 2 to 5 carbon atoms, (4) —(CH₂)_k—S—, wherein k is an integer of 2 to 5, or (5) —(CH₂)_pCO—, wherein p is an integer of 1 to 4, and B is



or pharmaceutically acceptable salts thereof.


2. A compound according to claim 1, wherein X is a group of the formula: —CO—.

3. A compound according to claim 1, wherein X is a group of the formula: —CHOH—.


4. A compound according to claim 1, wherein X is a group of the formula: —CH₂—.

5. A compound according to claim 1, wherein A is a group of the formula (5): —(CH₂)_pCO—, wherein p is an integer of 1 to 4.

6. A compound according to claim 1, wherein B is a group of the formula:



7. A compound according to claim 1, wherein X is a group of the formula: —CO—, A is a group of the formula (5): —(CH₂)_pCO—, wherein p is an integer of 1 to 4, and B is a group of the formula:



8. A compound according to claim 1, which is 1-[2-(3-imidazo[1,2-a]pyridyl)-1-oxoethyl]-4-(4-methylsulfonylamino benzoyl)-piperidine dihydrochloride.

9. A pharmaceutical composition which comprises a pharmacologically effective amount of the compound defined in claim 2 or a pharmacologically acceptable salt thereof and a pharmacologically acceptable carrier.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5 118 689
DATED : June 2, 1992
INVENTOR(S) : Hitoshi OINUMA et al

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, item [75] Inventors:

line 5, change "Sakura" to ---Sakuramura---

Column 67, line 41; change "akkoxy" to ---alkoxy---

Column 67, line 49; change "hydroxy" to ---hydroxyl---

Signed and Sealed this

Twenty-ninth Day of March, 1994

Attest:



BRUCE LEHMAN

Attesting Officer

Commissioner of Patents and Trademarks