



US 20150098999A1

(19) **United States**

(12) **Patent Application Publication**
Lulla et al.

(10) **Pub. No.: US 2015/0098999 A1**

(43) **Pub. Date: Apr. 9, 2015**

(54) **PHARMACEUTICAL PRODUCTS AND COMPOSITION COMPRISING SPECIFIC ANTI-CHOLINERGIC AGENTS, BETA-2 AGONISTS AND CORTICOSTEROIDS**

(71) Applicant: **Cipla Limited**, Mumbai Central (IN)

(72) Inventors: **Amar Lulla**, Mumbai (IN); **Geena Malhotra**, Mumbai (IN)

(21) Appl. No.: **14/570,679**

(22) Filed: **Dec. 15, 2014**

Related U.S. Application Data

(63) Continuation of application No. 10/525,736, filed on Jun. 6, 2005, now Pat. No. 8,962,601, filed as application No. PCT/GB2003/003751 on Aug. 29, 2003.

(30) **Foreign Application Priority Data**

Aug. 29, 2002 (GB) 02200954
Nov. 22, 2002 (GB) 02273423

Publication Classification

(51) **Int. Cl.**
A61K 31/58 (2006.01)
A61K 9/00 (2006.01)
A61K 31/439 (2006.01)
A61K 31/573 (2006.01)
A61K 31/56 (2006.01)
A61K 31/138 (2006.01)
A61K 31/46 (2006.01)
A61K 9/16 (2006.01)
A61K 31/137 (2006.01)

(52) **U.S. Cl.**
CPC . *A61K 31/58* (2013.01); *A61K 9/16* (2013.01);
A61K 9/0078 (2013.01); *A61K 31/439*
(2013.01); *A61K 31/137* (2013.01); *A61K*
31/56 (2013.01); *A61K 31/138* (2013.01);
A61K 31/46 (2013.01); *A61K 31/573* (2013.01)

(57) **ABSTRACT**
This invention relates to pharmaceutical products and compositions for use in the treatment of asthma and related disorders, and especially but not exclusively for the treatment of chronic obstructive pulmonary disease (COPD). More particularly, the invention provides pharmaceutical products and compositions comprising specific anticholinergic agents, β -2 agonists and corticosteroids.

**PHARMACEUTICAL PRODUCTS AND
COMPOSITION COMPRISING SPECIFIC
ANTIOCHOLINERGIC AGENTS, BETA-2
AGONISTS AND CORTICOSTEROIDS**

[0001] This invention relates to pharmaceutical products and compositions for use in the treatment of asthma and related disorders, and especially but not exclusively for the treatment of chronic obstructive pulmonary disease (COPD).

[0002] The pathophysiology of asthma and related disorders involves various distinct symptoms, including bronchoconstriction, inflammation of the airways, and increased mucous secretion, which results in wheezing, coughing and shortness of breath. A persistent or recurrent cough may exacerbate the problem by causing further irritation and inflammation of the airways. The causes of asthma are wide-ranging and not yet fully understood.

[0003] Bronchoconstriction occurs due to bronchial smooth muscle spasm and airway inflammation with mucosal edema. Asthma and other related disorders, have been known to be treated with β -2 adrenergic receptor agonists (β -2 agonist) as they provide a bronchodilator effect to the patients, resulting in relief from the symptoms of breathlessness. β -2 Agonists can be short acting for immediate relief, or long acting for long-term prevention, of asthma symptoms. Short acting β -2 agonists currently available include: salbutamol, biltolterol, pirbuterol and terbutaline. Long acting β -2 agonists currently available include salmeterol and formoterol.

[0004] Whilst it is also known that β -2 agonists provide symptomatic relief of bronchoconstriction in patients, another component of asthma, i.e. inflammation, often requires separate treatment. Typically this involves treatment with a steroid. Indeed, treatment with a corticosteroid is considered one of the most potent and effective therapies currently available for persistent asthma. Currently available corticosteroids include: beclomethasone, budesonide, flunisolide, fluticasone, mometasone and triamcinolone.

[0005] Bronchoconstriction and inflammation are also associated with bronchial plugging with secretions, which may be treated with anti-cholinergic agents, such as troventol, ipratropium, oxitropium and tiotropium.

[0006] These medicaments can be administered in different ways, such as in MDIs (metered-dosage inhalers), in DPIs (dry powder inhalers), and in oral and liquid formulations. Treatment in these different ways calls for the patient to comply with different dosage regimens, different frequencies of administration, etc. Also, since most of the medications are in the form of aerosols, the patient is required to carry several formulations and dispensers, one for each of these medicaments.

[0007] To assist patient compliance, combination products are known, e.g. an inhalation combination medication of fluticasone propionate and salmeterol, the combination being provided in one easy-to-use device. This combination product provides simultaneous treatment of airway constriction by means of the β -2 agonist (salmeterol), and treatment of inflammation by means of the steroid (fluticasone propionate).

[0008] A combination of ipratropium bromide and salbutamol is also known. This combination therapy provides an anti-cholinergic (ipratropium bromide) to reduce the bronchial secretions and a β -2 agonist (salbutamol) to reduce constriction. Other described combinations include ipratropium and salbutamol (WO 01/76601) and tiotropium and formoterol (WO 00/47200).

[0009] It would be highly desirable, however, to provide a combination therapy suitable to reduce bronchial inflammation, bronchial constriction and bronchial secretions in a single product or dosage form. It would also be desirable to provide such a combination product or composition in a form whereby the correct dosage of the various components is easily and safely administered.

[0010] We have now found that certain therapeutic three-in-one combinations comprising specific β -2 agonists, anticholinergics and steroids surprisingly provide an enhanced, synergistic, effect in terms of treatment of bronchoconstriction, inflammation and mucous secretions of airways. Also the three-in-one combination therapy as provided by the present invention is an extremely patient-friendly combination, which results in maximum patient compliance and better control of asthma than the known combinations or single therapies.

[0011] The present invention further provides, therefore, a pharmaceutical product comprising any one of the following combinations of therapeutic agents, as a combined preparation for simultaneous, separate or sequential use in the treatment of conditions for which administration of one or more of the therapeutic agents is indicated:

- [0012]** (i) salmeterol, ciclesonide and tiotropium;
- [0013]** (ii) formoterol, budesonide and ipratropium;
- [0014]** (iii) formoterol, ciclesonide and tiotropium;
- [0015]** (iv) formoterol, budesonide and oxitropium;
- [0016]** (v) salbutamol, beclomethasone and ipratropium;
- [0017]** (vi) salbutamol, budesonide and tiotropium;
- [0018]** (vii) terbutaline, fluticasone and tiotropium;
- [0019]** (viii) terbutaline, fluticasone and ipratropium;
- [0020]** (ix) salbutamol, budesonide and ipratropium;
- [0021]** (x) salmeterol, fluticasone and ipratropium;
- [0022]** (xi) salmeterol, budesonide and ipratropium;
- [0023]** (xii) salmeterol, fluticasone and tiotropium; and
- [0024]** (xiii) formoterol, budesonide and tiotropium.

[0025] It will also be appreciated from the above, that the respective therapeutic agents of the combined preparations can be administered simultaneously, either in the same or different pharmaceutical formulations, or separately or sequentially. If there is separate or sequential administration, it will also be appreciated that the subsequently administered therapeutic agents should be administered to a patient within a time scale so as to achieve, or more particularly optimise, the above referred to advantageous synergistic therapeutic effect of a combined preparation as present in a pharmaceutical product according to the present invention.

[0026] The present invention also provides a pharmaceutical composition comprising any one of the following combinations of therapeutic agents:

- [0027]** (i) salmeterol, ciclesonide and tiotropium;
- [0028]** (ii) formoterol, budesonide and ipratropium;
- [0029]** (iii) formoterol, ciclesonide and tiotropium;
- [0030]** (iv) formoterol, budesonide and oxitropium;
- [0031]** (v) salbutamol, beclomethasone and ipratropium;
- [0032]** (vi) salbutamol, budesonide and tiotropium;
- [0033]** (vii) terbutaline, fluticasone and tiotropium;
- [0034]** (viii) terbutaline, fluticasone and ipratropium;
- [0035]** (ix) salbutamol, budesonide and ipratropium;
- [0036]** (x) salmeterol, fluticasone and ipratropium;
- [0037]** (xi) salmeterol, budesonide and ipratropium;
- [0038]** (xii) salmeterol, fluticasone and tiotropium; and
- [0039]** (xiii) formoterol, budesonide and tiotropium.

together with a pharmaceutically acceptable carrier or excipient therefor.

[0040] The abovementioned compounds may exist, and be used in the present invention, in various active forms, whilst retaining the same physiological function. For example, the anticholinergic agents, β -2 agonists and corticosteroids may exist as various acid addition salts, such as those formed from hydrochloric, hydrobromic, sulphuric, acetic, lactic, maleic, tartaric, oxalic, methanesulphonic, p-toluenesulphonic and benzenesulphonic acids. The skilled person will also appreciate that the abovementioned compounds may also exist as esters and (R) and (S) enantiomers and provided the desired activity is maintained, they may be used in the present invention.

[0041] Specific triple combinations of the invention as illustrated by the Examples are:

[0042] (i) salmeterol, ciclesonide and tiotropium bromide;

[0043] (ii) formoterol, budesonide and ipratropium;

[0044] (iii) formoterol, ciclesonide and tiotropium bromide;

[0045] (iv) formoterol, budesonide and oxitropium;

[0046] (v) salbutamol sulphate, beclomethasone and ipratropium;

[0047] (vi) salbutamol sulphate, budesonide and tiotropium bromide;

[0048] (vii) terbutaline sulphate, fluticasone and tiotropium bromide;

[0049] (viii) terbutaline sulphate, fluticasone and ipratropium bromide;

[0050] (ix) salbutamol sulphate, budesonide and ipratropium bromide;

[0051] (x) salmeterol, fluticasone propionate and ipratropium bromide;

[0052] (xi) salmeterol, budesonide and ipratropium bromide;

[0053] (xii) salmeterol, fluticasone propionate and tiotropium bromide; and

[0054] (xiii) formoterol, budesonide and tiotropium bromide.

[0055] The pharmaceutical composition may be provided in any suitable dosage form. Preferably the composition is in the form of a suspension, a particulate suspension or a clear solution. The pharmaceutical composition of the present invention may alternatively be provided as an inhalation powder.

[0056] The pharmaceutical composition of the present invention may be administered by any suitable administration method and it may be preferred that the composition is administered as an aerosol. The composition of the present invention can comprise a propellant-containing dosage aerosol, an inhalation powder or a propellant free inhalation solution or suspension. The compositions of the present invention may thus be provided by, for example, a metered dose inhaler (MDI), dry powder inhaler (DPI), nebulers, nebuliser or nasal spray.

[0057] In the case of a propellant-containing dosage aerosol, typically the propellant can be selected from the group consisting of 1,1,1,2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoroethane, monofluorotrichloromethane and dichlorodifluoromethane.

[0058] In the case of a particulate suspension or a solution for the propellant-containing aerosol composition, the composition may further comprise one or more co-solvents. The composition may comprise both a propellant and a co-sol-

vent, in which case it is desirable that the co-solvent has a greater polarity than the propellant. The co-solvent used may be any suitable solvent. Typically the co-solvent is ethanol. Generally the ratio of propellant to solvent is between 50:50 to 99:1.

[0059] If aerosolized, the formulation may consist of a surface-active agent to stabilize the formulation and for the lubrication of a valve system in the inhaler/nebuliser/nasal spray.

[0060] Some of the most commonly used surface-active agents in the aerosol formulations are oils derived from natural sources, such as corn oil, olive oil, cotton seed oil and sunflower seed oil and phospholipids. Preferred surfactants for use according to the present invention are oleic acid, lecithin or sorbitol trioleate. In these embodiments, the surface-active agents are preferably used in the formulations in the ratio of 0.00002 wt % wt to 20 wt % of the active ingredients. The surface-active agents may exceed this weight ratio in cases where drug concentration in the formulation is very low.

[0061] The active ingredients in all the above aerosol formulations are preferably in the concentration of 0.001 wt % to 5 wt % of the total formulation.

[0062] The active ingredients are provided in an appropriate particle size, generally in the range from nano-size to about 12 μ m. Preferably, approximately 95% are below 5 or 6 μ m (micrometers), with the all particles being below 12 μ m (when measured by laser), or approximately 95% below 2.5 μ m and the rest of the particles between 2.5-5 μ m (when measured by microscope).

[0063] According to another aspect of the invention there is provided an aerosol device, comprising a housing containing a composition as described above, and a dispensing mechanism for dispensing the composition from the housing in a metered dose.

[0064] The dispensing mechanism may include a valve capable of releasing a metered dosage of the composition. Preferably the housing is sealed and pressurized at a pressure exceeding atmospheric pressure.

[0065] **[text missing or illegible when filed]** housing may be metallic, preferably aluminum. Preferably, the housing is plastic-coated, lacquer-coated or anodised. The composition of the present invention may be placed in the housing through a suitable metering device.

[0066] Preferred combinations of active ingredients for administration by way of a propellant-containing dosage aerosols according to the present invention are further illustrated by the Examples and can include any one of the following combinations:

[0067] (i) salmeterol, ciclesonide and tiotropium;

[0068] (ii) formoterol, budesonide and ipratropium;

[0069] (iii) formoterol, ciclesonide and tiotropium;

[0070] (iv) formoterol, budesonide and oxitropium;

[0071] (v) salbutamol, beclomethasone and ipratropium;

[0072] (vi) salbutamol, budesonide and tiotropium;

[0073] (vii) terbutaline, fluticasone and tiotropium;

[0074] (viii) salmeterol, fluticasone and tiotropium; and

[0075] (ix) formoterol, budesonide and tiotropium.

[0076] More specifically the following may be administered by way of propellant-containing dosage aerosols according to the present invention:

[0077] (i) salmeterol, ciclesonide and tiotropium bromide;

[0078] (ii) formoterol, budesonide and ipratropium;

[0079] (iii) formoterol, ciclesonide and tiotropium bromide;

[0080] (iv) formoterol, budesonide and oxitropium;

[0081] (v) salbutamol sulphate, beclomethasone and ipratropium;

[0082] (vi) salbutamol sulphate, budesonide and tiotropium bromide;

[0083] (vii) terbutaline sulphate, fluticasone and tiotropium bromide;

[0084] (viii) salmeterol, fluticasone propionate and tiotropium bromide;

[0085] (ix) formoterol, budesonide and tiotropium bromide.

[0086] In the case where the triple combinations of the present invention are provided as inhalation powders, typically suitable pharmaceutically acceptable excipients may be selected from monosaccharides, disaccharides, oligosaccharides, polysaccharides or the like, with the use of lactose as the excipient in the inhalation powders of the present invention being preferred. The inhalation powders of the present invention can typically be administered by means of dry powder inhalers known in the art. Preferred combinations of active ingredients for administration by way of inhalation powders according to the present invention are further illustrated by the Examples and can include any one of the following combinations:

[0087] (i) salmeterol, ciclesonide and tiotropium;

[0088] (ii) formoterol, budesonide and ipratropium;

[0089] (iii) formoterol, ciclesonide and tiotropium;

[0090] (iv) salbutamol, beclomethasone and ipratropium;

[0091] (v) salbutamol, budesonide and tiotropium; and

[0092] (vi) terbutaline, fluticasone and tiotropium.

[0093] More specifically the following may be administered by way of inhalation powders according to the present invention:

[0094] (i) salmeterol, ciclesonide and tiotropium bromide;

[0095] (ii) formoterol, budesonide and ipratropium;

[0096] (iii) formoterol, ciclesonide and tiotropium bromide;

[0097] (iv) salbutamol sulphate, beclomethasone and ipratropium;

[0098] (v) salbutamol sulphate, budesonide and tiotropium bromide; and

[0099] (vi) terbutaline sulphate, fluticasone and tiotropium bromide.

[0100] The propellant free inhalation solutions of the present invention are typically suitable for administration by way of nebulisation, employing nebulisers well known in the art that can advantageously be employed to produce inhalable aerosols comprising the triple combinations of active ingredients according to the present invention. Preferred combinations of active ingredients for administration by way of propellant free inhalation solutions according to the present invention are further illustrated by the Examples and can include any one of the following combinations:

[0101] (i) terbutaline, fluticasone and ipratropium;

[0102] (ii) salbutamol, budesonide and ipratropium;

[0103] (iii) salmeterol, fluticasone and ipratropium; and

[0104] (iv) salmeterol, budesonide and ipratropium.

[0105] More specifically the following may be administered by way of propellant free inhalation solutions according to the present invention:

[0106] (i) terbutaline sulphate, fluticasone and ipratropium bromide;

[0107] (ii) salbutamol sulphate, budesonide and ipratropium bromide;

[0108] (iii) salmeterol, fluticasone propionate and ipratropium bromide; and

[0109] (iv) salmeterol, budesonide and ipratropium bromide.

[0110] According to another aspect of the invention, there is provided the use of any one of the following combinations in the manufacture of a medicament for the prophylaxis or treatment of conditions for which administration of one or more of the therapeutic agents is indicated:

[0111] (i) salmeterol, ciclesonide and tiotropium;

[0112] (ii) formoterol, budesonide and ipratropium;

[0113] (iii) formoterol, ciclesonide and tiotropium;

[0114] (iv) formoterol, budesonide and oxitropium;

[0115] (v) salbutamol, beclomethasone and ipratropium;

[0116] (vi) salbutamol, budesonide and tiotropium;

[0117] (vii) terbutaline, fluticasone and tiotropium;

[0118] (viii) terbutaline, fluticasone and ipratropium;

[0119] (ix) salbutamol, budesonide and ipratropium;

[0120] (x) salmeterol, fluticasone and ipratropium;

[0121] (xi) salmeterol, budesonide and ipratropium;

[0122] (xii) salmeterol, fluticasone and tiotropium; and

[0123] (xiii) formoterol, budesonide and tiotropium.

[0124] The triple combinations as provided by the present invention are useful for the treatment of inflammatory or respiratory tract diseases, especially asthma and/or chronic obstructive pulmonary disease (COPD), by simultaneous or successive administration.

[0125] According to another aspect of the invention there is provided a method for the prophylaxis or treatment of inflammatory or respiratory tract diseases, said method comprising administering either sequentially or simultaneously, to a patient in need thereof, a therapeutically effective amount of any one of the following combinations:

[0126] (i) salmeterol, ciclesonide and tiotropium;

[0127] (ii) formoterol, budesonide and ipratropium;

[0128] (iii) formoterol, ciclesonide and tiotropium;

[0129] (iv) formoterol, budesonide and oxitropium;

[0130] (v) salbutamol, beclomethasone and ipratropium;

[0131] (vi) salbutamol, budesonide and tiotropium;

[0132] (vii) terbutaline, fluticasone and tiotropium;

[0133] (viii) terbutaline, fluticasone and ipratropium;

[0134] (ix) salbutamol, budesonide and ipratropium;

[0135] (x) salmeterol, fluticasone and ipratropium;

[0136] (xi) salmeterol, budesonide and ipratropium;

[0137] (xii) salmeterol, fluticasone and tiotropium; and

[0138] (xiii) formoterol, budesonide and tiotropium.

[0139] Any of the following specific triple combinations of the invention as illustrated by the Examples are suitable to be employed in the manufacture of a medicament, or a method of treatment, as referred to above:

[0140] (i) salmeterol, ciclesonide and tiotropium bromide;

[0141] (ii) formoterol, budesonide and ipratropium;

[0142] (iii) formoterol, ciclesonide and tiotropium bromide;

[0143] (iv) formoterol, budesonide and oxitropium;

[0144] (v) salbutamol sulphate, beclomethasone and ipratropium;

[0145] (vi) salbutamol sulphate, budesonide and tiotropium bromide;

[0146] (vii) terbutaline sulphate, fluticasone and tiotropium bromide;

[0147] (viii) terbutaline sulphate, fluticasone and ipratropium bromide;

[0148] (ix) salbutamol sulphate, budesonide and ipratropium bromide;

[0149] (x) salmeterol, fluticasone propionate and ipratropium bromide;

[0150] (xi) salmeterol, budesonide and ipratropium bromide;

[0151] (xii) salmeterol, fluticasone propionate and tiotropium bromide; and

[0152] (xiii) formoterol, budesonide and tiotropium bromide.

[0153] These products or compositions as provided by the present invention are especially useful in the treatment of COPD. They will normally be administered by inhalation, once or twice daily. By way of example, a preferred dosage for twice daily administrations would be:

[0154] a) formoterol (6 mcg)/budesonide (200 mcg)/ipratropium (40 mcg)

[0155] b) formoterol (6 mcg)/budesonide (200 mcg)/oxitropium (200 mcg).

[0156] The invention will now be described with reference to the following examples, which do not limit the scope of the invention in any way.

EXAMPLE 1

[0157]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Salbutamol	24 mg
Budesonide	24 mg
1,1,1,2-Tetrafluoroethane	18.2 gms

EXAMPLE 2

[0158]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Terbutaline Sulphate	60 mg
Fluticasone	12 mg
1,1,1,2-Tetrafluoroethane	18.2 gms

[0159] In the above formulations (Examples 1 and 2), the active ingredients were initially weighed in an aluminum can. Then a metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA 134a was charged through the metering valve.

EXAMPLE 3

[0160]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Salbutamol	24 mg
Budesonide	24 mg
Ethanol	2.73 gms
1,1,1,2-Tetrafluoroethane	15.47 gms

In the above formulation the active ingredients were first weighed in an aluminum can, then the ethanol was added and

the solution was sonicated for 5 min. The metering valve was placed on the can and crimped with a vacuum crimper, and then propellant HFA 134a was charged through the metering valve.

EXAMPLE 4

[0161]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Salbutamol	24 mg
Budesonide	24 mg
Ethanol	2.73 gms
Oleic acid (10%)	5.04 mg
1,1,1,2-Tetrafluoroethane	15.47 gms

In the above formulation the active ingredients were first weighed in an aluminum can then the ethanol and the surfactant were added and solution was sonicated for 5 min. The metering valve was placed on the can and crimped with a vacuum crimper and then the HFA 134a was charged through the metering valve.

EXAMPLE 5

[0162]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Terbutaline Sulphate	60 mg
Fluticasone	12 mg
Ethanol	0.364 gms
1,1,1,2-Tetrafluoroethane	18.2 gms

EXAMPLE 6

[0163]

Per aerosol housing	
Tiotropium bromide	2.4 mg
Terbutaline Sulphate	60 mg
Fluticasone	12 mg
Ethanol	0.364 gms
Oleic acid (0.02%)	0.014 mg
1,1,1,2-Tetrafluoroethane	17.83 gms

[0164] The above formulations (Examples 5 and 6) were weighed in an aluminum can, and a metering valve was crimped on the can and the propellant added.

EXAMPLE 7

[0165]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Ciclesonide	8.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 8

[0166]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Ciclesonide	8.0 mg
Ethanol (2%)	0.244 g
1,1,1,2-Tetrafluoroethane	12 g

[0167] The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 9

[0168]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Ciclesonide	8.0 mg
Ethanol (2%)	0.244 g
Oleic acid (0.02%)	0.003 mg
1,1,1,2-Tetrafluoroethane	12 g

The active ingredients were initially weighed in an aluminum can, then ethanol and oleic acid were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 10

[0169]

Ingredients	Quantity (Per aerosol housing)
Fomoterol	1 mg
Ipratropium	3.2 mg
Budesonide	8.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 11

[0170]

Ingredients	Quantity (Per aerosol housing)
Fomoterol	1 mg
Ipratropium	3.2 mg

-continued

Ingredients	Quantity (Per aerosol housing)
Budesonide	8.0 mg
Ethanol (15%)	1.83 g
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 12

[0171]

Ingredients	Quantity (Per aerosol housing)
Fomoterol	1.0 mg
Ipratropium	3.2 mg
Budesonide	8.0 mg
Ethanol	1.83 g
Lecithin (1%)	0.122 mg
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can, then ethanol and lecithin were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 13

[0172]

Ingredients	Quantity (Per aerosol housing)
Tiotropium Bromide	1.8 mg
Fomoterol	1.0 mg
Ciclesonide	8.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 14

[0173]

Ingredients	Quantity (Per aerosol housing)
Tiotropium Bromide	1.8 mg
Fomoterol	1.0 mg
Ciclesonide	8.0 mg
Ethanol (15%)	1.83 g
1,1,1,2-Tetrafluoroethane	10.4 g

[0174] The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the

can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 15

[0175]

Ingredients	Quantity (Per aerosol housing)
Tiotropium Bromide	1.8 mg
Formoterol	1.0 mg
Ciclesonide	8.0 mg
Ethanol	1.83 g
Lecithin (1%)	0.10 mg
1,1,1,2-Tetrafluoroethane	10.4 g

The active ingredients were initially weighed in an aluminum can, then ethanol and lecithin were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 16

[0176]

Ingredients	Quantity (Per aerosol housing)
Oxitropium	4.8 mg
Formoterol	1.0 mg
Budesonide	8.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 17

[0177]

Ingredients	Quantity (Per aerosol housing)
Oxitropium	4.8 mg
Formoterol	1.0 mg
Budesonide	8.0 mg
Ethanol (15%)	1.83 g
1,1,1,2-Tetrafluoroethane	10.4 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 18

[0178]

Ingredients	Quantity (Per aerosol housing)
Oxitropium	4.8 mg
Formoterol	1.0 mg
Budesonide	8.0 mg
Ethanol (15%)	1.83 g
Lecithin (1%)	0.138 mg
1,1,1,2-Tetrafluoroethane	10.4 g

The active ingredients were initially weighed in an aluminum can, then ethanol and lecithin were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 19

[0179]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Beclomethasone	12.0 mg
Ipratropium	4.8 mg
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 20

[0180]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Beclomethasone	12.0 mg
Ipratropium	4.8 mg
Ethanol (2.5%)	0.455 g
1,1,1,2-Tetrafluoroethane	17.75 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 21

[0181]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Beclomethasone	12.0 mg
Ipratropium	4.8 mg

-continued

Ingredients	Quantity (Per aerosol housing)
Ethanol (2.5%)	0.455 g
Oleic acid (0.02%)	0.009 mg
1,1,1,2-Tetrafluoroethane	17.75 g

The active ingredients were initially weighed in an aluminum can, then ethanol and oleic acid were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 22

[0182]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Tiotropium bromide	2.7 mg
Budesonide	12 mg
1,1,1,2-Tetrafluoroethane	18.2 g

[0183] The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 23

[0184]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Tiotropium bromide	2.7 mg
Budesonide	12 mg
Ethanol (2%)	0.364 g
1,1,1,2-Tetrafluoroethane	17.83 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 24

[0185]

Ingredients	Quantity (Per aerosol housing)
Salbutamol sulphate	28.8 mg
Tiotropium bromide	2.7 mg
Budesonide	12 mg
Ethanol (2%)	0.364 g
Oleic acid (0.02%)	0.0087 mg
1,1,1,2-Tetrafluoroethane	17.83 g

The active ingredients were initially weighed in an aluminum can, then ethanol and oleic acid were added and the solution

was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 25

[0186]

Ingredients	Quantity (Per aerosol housing)
Terbutaline sulphate	40 mg
Tiotropium bromide	1.8 mg
Fluticasone	8.0 mg
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can. The metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 26

[0187]

Ingredients	Quantity (Per aerosol housing)
Terbutaline sulphate	40 mg
Tiotropium bromide	1.8 mg
Fluticasone	8.0 mg
Ethanol	2.73 g
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 27

[0188]

Ingredients	Quantity (Per aerosol housing)
Terbutaline sulphate	40 mg
Tiotropium bromide	1.8 mg
Fluticasone	8.0 mg
Ethanol	2.73 g
Lecithin	1.38 mg
1,1,1,2-Tetrafluoroethane	15.83 g

The active ingredients were initially weighed in an aluminum can, then ethanol and lecithin were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 28

[0189]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Fluticasone	20.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. Metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 29

[0190]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Fluticasone	20.0 mg
Ethanol (2%)	0.244 g
1,1,1,2-Tetrafluoroethane	12 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 30

[0191]

Ingredients	Quantity (Per aerosol housing)
Tiotropium bromide	1.8 mg
Salmeterol	5.81 mg
Fluticasone	20.0 mg
Ethanol (2%)	0.244 g
Oleic acid (0.02%)	0.003 mg
1,1,1,2-Tetrafluoroethane	12 g

The active ingredients were initially weighed in an aluminum can, then ethanol and the surfactant were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 31

[0192]

Ingredients	Quantity (Per aerosol housing)
Formoterol	1 mg
Tiotropium Bromide	1.8 mg
Budesonide	8.0 mg
1,1,1,2-Tetrafluoroethane	12.2 g

The active ingredients were initially weighed in an aluminum can. Metering valve was then placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 32

[0193]

Ingredients	Quantity (Per aerosol housing)
Formoterol	1 mg
Tiotropium Bromide	1.8 mg
Budesonide	8.0 mg
Ethanol (15%)	1.83 g
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can, then ethanol was added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

EXAMPLE 33

[0194]

Ingredients	Quantity (Per aerosol housing)
Formoterol	1.0 mg
Tiotropium Bromide	1.8 mg
Budesonide	8.0 mg
Ethanol	1.83 g
Lecithin (1%)	0.122 mg
1,1,1,2-Tetrafluoroethane	18.2 g

The active ingredients were initially weighed in an aluminum can, then ethanol and the surfactant were added and the solution was sonicated for 5 minutes. The metering valve was placed on the can and crimped with a vacuum crimper and then the propellant HFA134a was charged through the metering valve.

[0195] The following Examples are inhalation powders suitable for DPIs, and were prepared by techniques well known in the art.

EXAMPLE 34

[0196]

Ingredients	Quantity (in mg per capsule)
Tiotropium bromide	0.018 mg
Salmeterol	0.050 mg
Ciclesonide	0.1-0.4 mcg
Lactose	q.s. 5-25 mg

EXAMPLE 35

[0197]

Ingredients	Quantity (in mg per capsule)
Formoterol	0.025-0.4 mg
Ipratropium	0.048 mg
Budesonide	0.1-0.4 mg
Lactose	q.s. 5-25 mg

EXAMPLE 36

[0198]

Ingredients	Quantity (in mg per capsule)
Tiotropium Bromide	0.018 mg
Formoterol	0.025-0.4 mg
Ciclesonide	0.1-0.4 mg
Lactose	q.s. 5-25 mg

EXAMPLE 37

[0199]

Ingredients	Quantity (in mg per capsule)
Salbutamol sulphate	0.2 mg
Beclomethasone	0.1-0.4 mg
Ipratropium	0.048 mg
Lactose	q.s. 5-25 mg

EXAMPLE 38

[0200]

Ingredients	Quantity (in mg per capsule)
Salbutamol sulphate	0.2 mg
Tiotropium bromide	0.018 mg
Budesonide	0.1-0.4 mg
Lactose	q.s. 5-25 mg

EXAMPLE 39

[0201]

Ingredients	Quantity (in mg per capsule)
Terbutaline sulphate	0.1-0.2 mg
Tiotropium bromide	0.018 mg
Fluticasone	0.025-0.4 mg
Lactose	q.s. 5-25 mg

The following Examples are for propellant free inhalation solutions, and were prepared by techniques well known in the art.

EXAMPLE 40

[0202]

Ingredients	Quantity (% w/v)
Terbutaline sulphate	0.25
Ipratropium bromide	0.025
Fluticasone	0.025-0.1
Polysorbat-80	0.1
Sodium chloride	0.9
Anhydrous citric acid	q.s. to pH 4.5
Water, purified	100 ml

EXAMPLE 41

[0203]

Ingredients	Quantity (% w/v)
Ipratropium bromide	0.025
Salbutamol sulphate	0.125
Budesonide	0.05
Polysorbat-80	0.1
Sodium chloride	0.9
Water, purified	100 ml

EXAMPLE 42

[0204]

Ingredients	Quantity (% w/v)
Ipratropium bromide	0.025
Salmeterol	0.005
Fluticasone propionate	0.025-0.1
Polysorbat-80	0.1
Sodium chloride	0.9
Water, purified	100 ml

EXAMPLE 43

[0205]

Ingredients	Quantity (% w/v)
Ipratropium bromide	0.025
Salmeterol	0.005
Budesonide	0.05
Polysorbat-80	0.1
Sodium chloride	0.9
Water, purified	100 ml

1. (canceled)

2. A pharmaceutical composition comprising any one of the following combinations of therapeutic agents in a form suitable for administration by inhalation for use in the treatment of inflammatory or respiratory diseases:

- (i) salmeterol, ciclesonide and tiotropium;
- (ii) formoterol, budesonide and ipratropium;
- (iv) formoterol, budesonide and oxitropium;
- (v) salbutamol, beclomethasone and ipratropium;
- (vi) salbutamol, budesonide and tiotropium;
- (vii) terbutaline, fluticasone and tiotropium;

- (viii) terbutaline, fluticasone and ipratropium;
- (ix) salbutamol, budesonide and ipratropium;
- (x) salmeterol, fluticasone and ipratropium;
- (xi) salmeterol, budesonide and ipratropium;
- (xii) salmeterol, fluticasone and tiotropium; and

wherein the therapeutic agents are each provided in particulate form, each of the therapeutic agents individually has a particle size from nano-size up to about 12 μm , and approximately 95% of the therapeutic agent particles have a particle size of below about 5 μm ; and

wherein each of the therapeutic agents can optionally independently be present as a pharmaceutically acceptable salt or ester thereof, and each of the therapeutic agents can optionally independently be present in enantiomerically pure form or as a racemic mixture, together with a pharmaceutically acceptable carrier or excipient therefor.

3. A composition according to claim 2, wherein each of the therapeutic agents of the composition is present in an amount

0.001 wt % to 0.5wt % based on the weight of the total composition.

4. A composition according to claim 2, in the form of an aerosol.

5. A composition according to claim 4, in the form of a propellant free inhalation solution or suspension.

6. A composition according to claim 2, in the form of an inhalation powder

7. A composition according to claim 6, which comprises lactose as the excipient.

8. A composition according to claim 2, in a form suitable for use with a nebuliser.

9. A metered dose inhaler which contains a composition as defined in claim 2.

10. A dry powder inhaler which contains a composition as defined in claim 2.

11. A nebuliser which contains a composition as defined in claim 2.

* * * * *