



US005834607A

United States Patent [19][11] **Patent Number:** **5,834,607****Manoharan et al.**[45] **Date of Patent:** **Nov. 10, 1998**[54] **AMINES AND METHODS OF MAKING AND USING THE SAME**[75] Inventors: **Muthiah Manoharan; P. Dan Cook**, both of Carlsbad, Calif.[73] Assignee: **ISIS Pharmaceuticals, Inc.**, Carlsbad, Calif.[21] Appl. No.: **361,858**[22] Filed: **Dec. 22, 1994****Related U.S. Application Data**

[63] Continuation of Ser. No. 943,516, Sep. 11, 1992, abandoned, which is a continuation-in-part of Ser. No. 558,663, Jul. 27, 1990, Pat. No. 5,138,045, and a continuation-in-part of Ser. No. 844,845, Mar. 3, 1992, Pat. No. 5,218,105.

[51] **Int. Cl.⁶** **C12Q 1/68**[52] **U.S. Cl.** **536/22.1**; 435/6; 435/810; 436/501; 514/44; 536/23.1; 536/24.1; 536/24.3; 536/24.31; 536/24.32; 536/24.33; 935/77; 935/78[58] **Field of Search** 435/6, 860; 436/501; 514/44; 536/22.1, 23.1, 24.1, 24.3–24.33; 935/77, 78[56] **References Cited****U.S. PATENT DOCUMENTS**

4,515,781	5/1985	Torrence et al.	514/46
4,840,892	6/1989	Adams et al.	435/5
5,034,506	7/1991	Summerton et al.	528/391
5,138,045	8/1992	Cook et al.	536/27
5,218,105	6/1993	Cook et al.	536/25.31

FOREIGN PATENT DOCUMENTS

PCT/US91/00243	1/1991	WIPO .
PCT/US91/04086	6/1991	WIPO .

OTHER PUBLICATIONS


Wilson, D.B. *Ann. Rev. Biochem.* 47:933–965 (1978).
 Wu, G.Y. et al., *Biochemistry* 27:887–892 (1988).
 Yoshinari et al., *J. Am. Chem. Soc.* 113 5899–5901 (1991).
 Zon, G. in *Oligodeoxynucleotides: Antisense Inhibitors of Gene Expression* 234–247, ed. J.S. Cohen (CRC Press, Boca Raton FL, 1989).
 Uhlmann, E. and Peyman, “Antisense Oligonucleotides: A New Therapeutic Principle”, *Chemical Reviews* 1990, 90 (40), 544–584.
 Bayard, et al., *Biochemistry* 25: 3730 (1986).
 Behr, *Tetrahedron Letters* vol. 27 No. 48 5861–5864 (1986).
 Ceruzzi, M. et al., *Nucleosides & Nucleotides* 8:815–818 (1989).
 Doetsch and Cunningham, *Mutation Research*, 236, 173 (1990).
 Friedberg, *DNA Repair* (W.H. Freeman and Company, NY, 1985) p. 153.
 Groehke et al., *Helvetica Chimica Acta* vol. 73 608–617 (1990).
 Horn, et al., *Nucleosides and Nucleotides* 10:299 (1991).
 Inoue et al., *Nucleic Acids Research* vol. 15 No. 15 6131 (1987).
 Iacono et al., *Tetrahedron Letters* vol. 31 No. 2 175–178 (1990).
 Le Doan, et al., *Nucleic Acids Research* 15, 8643 (1987).
 Lemaitre, M., et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:648–652 (1987).
 Letsinger, et al, *Proc. Natl. Acad. Sci. U.S.A.* 86:653–656 (1989).
 Lindahl et al., *J. Biol. Chem.* 252: 3286 (1977).

(List continued on next page.)

Primary Examiner—George C. Elliott
Assistant Examiner—Ardin H. Marschel
Attorney, Agent, or Firm—Woodcock Washburn Kurtz Mackiewicz & Norris LLP

[57] **ABSTRACT**

Novel amine compounds are provided by the present invention. Methods of preparing and using said novel amine compounds are also provided.

5 Claims, 3 Drawing Sheets

OTHER PUBLICATIONS

- Loke, S.L. et al., *Curr. Top Microbiol. Immunol.* 141:282–289 (1988).
- Iyer, et al., *Nucleic Acids Research* 18: 2855–2859 (1990).
- Manoharan, et al., *JACS*, 110, 2690 (1988).
- Marcus–Sekura, C.H. et al, *Nuc. Acids Res.* 15:5749–5763 (1987).
- Miller, P.S. et al, *Biochemistry* 20:1874–1880 (1981).
- Miller, P.S. et al., *Biochemistry* 16:1988–1996 (1977).
- Nielsen, et al., *Science* 254 1497 (1991).
- Péoc'h, et al., *Tetrahedron Letters* 32: 207 (1991).
- Robins et al., *J. Org. Chem.* vol. 39 No. 13 1891–(1974).
- Robins et al., *Can. J. Chem.* 59 3360 (1981).
- Sakumi et al., *Mutation Research* 236 161–172 (1990).
- Smith–Jones et al., *Bioconjugate Chem.* 2 415–421 (1991).
- Stevenson, M. et al., *J. Gen. Virol* 70:2673–2682 (1989).
- Studer et al., *Helvetica Chimica Acta* vol. 69 2081–2086 (1986).
- Suh et al., *J. Am. Chem. Soc.* 113 4198–4202 (1991).
- Takagi et al., *Tetrahedron Letters* vol. 32 No. 9 1207–1208 (1991).
- Vasseur, et al., *Nucleosides and Nucleotides*, 10, 107 (1991).
- Wilde et al., *J. Am. Chem. Soc.* 111 1894–1896 (1989).




Fig. 1





Fig. 2



AMINES AND METHODS OF MAKING AND USING THE SAME

This application is a continuation of application Ser. No. 07/943,516, filed Sep. 11, 1992, now abandoned, which is a continuation-in-part of application Ser. No. 07/558,663, files Jul. 27, 1990 (now U.S. Pat. No. 5,138,045) and of application Ser. No. 07/844,845, filed Mar. 3, 1992 (now U.S. Pat. No. 5,218,105).

FIELD OF THE INVENTION

This invention relates to novel amine-containing compounds useful for therapeutics and methods of making and using the same.

BACKGROUND OF THE INVENTION

It is well known that most of the bodily states in mammals including most disease states, are effected by proteins. Such proteins, either acting directly or through their enzymatic functions, contribute in major proportion to many diseases in animals and man. Classical therapeutics has generally focused upon interactions with such proteins in efforts to moderate their disease causing or disease potentiating functions. Recently, however, attempts have been made to moderate the actual production of such proteins by interactions with molecules that direct their synthesis, intracellular RNA. These interactions involved the binding of complementary "antisense" oligonucleotides or their analogs to the trans-cellular RNA in a sequence specific fashion such as by Watson-Crick base pairing interactions.

The pharmacological activity of antisense compounds, as well as other therapeutics, depends on a number of factors that influence the effective concentration of these agents at specific intracellular targets. One important factor is the ability of antisense compounds to traverse the plasma membrane of specific cells involved in the disease process.

Cellular membranes consist of lipid protein bilayers that are freely permeable to small, nonionic, lipophilic compounds and inherently impermeable to most natural metabolites and therapeutic agents. Wilson, D. B. *Ann. Rev. Biochem.* 47:933-965 (1978). The biological and antiviral effects of natural and modified oligonucleotides in cultured mammalian cells have been well documented, so it appears that these agents can penetrate membranes to reach their intracellular targets. Uptake of antisense compounds into a variety of mammalian cells, including HL-60, Syrian Hamster fibroblast, U937, L929, CV-1, and ATH8 cells has been studied using natural oligonucleotides and nuclease resistant analogs, such as alkyl triesters, Miller, P. S., Braiterman, L. T. and Ts'O, P. O. P., *Biochemistry* 16:1988-1996 (1977); methylphosphonates, Marcus-Sekura, C. H., Woerner, A. M., Shinozuka, K. Zon, G., and Quinman, G. V., *Nuc. Acids Res.* 15:5749-5763 (1987) and Miller, P. S., McParland, K. B., Hayerman, K. and Ts'O, P. O. P., *Biochemistry* 20:1874-1880 (1981); and phosphorothioates, Ceruzzi, M. and Draper, K. *Nucleosides & Nucleotides* 8:815-818 (1989); Miller, P. S., Braiterman, L. T. and Ts'O, P. O. P. *Biochemistry* 16:1988-1996 (1977) and Loke, S. L., Stein, C., Zhang, X. H. Avigan, M., Cohen, J. and Neckers, L. M. *Curr. Top. Microbiol. Immunol.* 141:282-289 (1988).

Enhanced cellular uptake has previously been achieved by attachment of functional groups to the 3' and 5' end of oligonucleotides to enhance cellular uptake in specific cell types. Previous studies have shown that plasmid DNA complexed with an (asialo)glycoprotein-poly(L-lysine) conjugate, could be targeted to hepatocytes, which contain

unique cell surface receptors for galactose-terminal (asialo) glycoproteins. Wu, G. Y. and Wu, C. H. *Biochemistry* 27:887-892 (1988). Other groups have synthesized oligodeoxyribonucleotides that have a 5'-attached alkylating agent and a 3' attached cholesterol moiety and determined that these modified oligonucleotides were taken up into cells more efficiently than control compounds without the steroid moiety. Zon, G. in *Oligodeoxynucleotides: Antisense Inhibitors of Gene Expression* 234-247, ed. J. S. Cohen (CRC Press, Boca Raton Fla., 1989). Letsinger, et al. *Proc. Natl. Acad. Sci. U.S.A.* 86:653-656 (1989), have also synthesized cholesteryl-conjugated phosphorothioates whose anti-HIV activity is significantly greater than natural oligonucleotides with the same sequence. Additional modifications include conjugation of oligonucleotides to poly(L-lysine) alone. Stevenson, M. and Iversen, P. L. *J. Gen. Virol* 70:2673-2682 (1989) and Lemaitre, M., Baynard, B. and LeBleu, B. *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652 (1987). This modification enhanced the antiviral activity of the compound studied presumably due to increased cellular uptake imparted by the polycationic poly(L-lysine).

The conjugation of polyamines to oligonucleotides have been found to enhance cellular uptake of oligonucleotides, increased lipophilicity, cause greater cellular retention and increased distribution of the compound. Vasseur, et al., *Nucleosides and Nucleotides*, 1991, 10, 107 prepared abasic sites at different sites of oligothymidylates by acid hydrolysis. Thereafter the abasic sites were functionalized with functionalities such as 3-amino carbazole, 9-amino elipticine and psoralen. Vasseur, et al. also refers to unpublished results in which the functionalities spermidine and proflavin were employed. Le Doan, et al., *Nucleic Acids Research* 1987, 15, 8643 teaches oligothymidylates covalently linked to porphyrins at their 3' end via one of the linkers —O—CH₂—CO—NH—(CH₂)₂—NH or PO₄—(CH₂)₆—NH—. Le Doan, et al. also used the linker PO₄—(CH₂)₆—NH— to link porphyrins to the 5' end of oligothymidylates. Another group, Summerton, et al., U.S. Pat. No. 5,034,506 issued Jul. 23, 1991 teaches morpholino subunits, linked together by uncharged, achiral linkages such as amides. As described in PCT/US91/04086 filed Jun. 10, 1991, polyamines have also been linked at the 5' end of an oligonucleotide at the 5' site of the sugar moiety of the terminal nucleoside and at the 2-position carbon of the heterocyclic base of 2'-deoxyadenosine, 2'-deoxyguanosines and other purines and purine analogs by known procedures as described in PCT/US/91/00243 filed Jan. 11, 1991.

Novel amines and methods of preparing the same are greatly needed in order to enhance cellular uptake of oligonucleotides, increase lipophilicity, cause greater cellular retention and increase distribution of the compound within the cell. The present invention fulfills this need.

OBJECTS OF THE INVENTION

It is one object of the present invention to provide novel amine-containing compounds useful in therapeutics.

It is a further object of the present invention to provide methods of producing said novel compounds.

It is another object of the present invention to provide methods of modulating the production of a protein by an organism.

It is still a further object of the present invention to provide methods of treating a mammal suffering from a disease characterized by the undesired production of a protein.

It is yet a further object of the present invention to provide methods of diagnosing the presence of an RNA in a biological sample.


3

These and other objects will become apparent from the following description and accompanying claims.

SUMMARY OF THE INVENTION


The present invention provides compounds which may have enhanced efficacy as an antisense-based therapy. Compounds of the present invention can have enhanced cellular uptake, increased lipophilicity, cause greater cellular retention and demonstrate increased distribution. Furthermore the present invention provides simple methods for synthesis of these novel compounds.

In accordance with some embodiments of the present invention, compounds having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, oligonucleotide, or polyamine and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic non-aromatic polyamine species, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkylaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide, are provided.


In accordance with still other embodiments of the present invention, compounds having the structure:



wherein R_4 is an oligonucleotide and M is a pendent group having a polyamine species attached thereto are provided.


Methods of preparing such compounds utilizing enzymatic reagents are also provided in some aspects of the invention. Thus compounds of Formula I may be prepared by methods comprising the steps of providing a synthon having the structure:

4




wherein R_1 and R_2 are independently H, a nucleotide, oligonucleotide or polyamine, and at least one of R_1 and R_2 is a purine containing oligonucleotide, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkylaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide. Thereafter the synthon is reacted with R_3 , wherein R_3 is a linear or cyclic non-aromatic polyamine species, under reducing conditions to yield the final product.

Compounds of Formula II may also be prepared enzymatically by providing a starting material having the structure:



wherein R_4 is an oligonucleotide, R_{12} is an oligonucleotide and B is urea or a heterocyclic base having a corresponding glycosylase and reacting the starting material with an endonuclease to generate a conjugated α,β -unsaturated system in the sugar residue of the 3' terminal nucleotide. Thereafter the compound having a conjugated α,β -unsaturated system is reacted with a pendent group containing a nucleophile functionality thereon. Following addition of the pendent group the double bond of the α,β system is reduced with a reducing agent. A polyamine species may then be attached to the pendent group via an alkylation reaction. Alternatively, a polyamine species may be attached to a pendent group which is a bifunctional linker.

In accordance with still other embodiments of the present invention compounds having the structure:




III

5

wherein R_4 is an oligonucleotide, R_5 is a linear or cyclic non-aromatic polyamine species containing at least one non-amide nitrogen atom, and R_6 is H, a purine heterocycle or a pyrimidine heterocycle, are provided. Methods of preparing compounds of Formula III are also provided in some aspects of the present invention comprising the steps of reacting an oligonucleotide having a 3' ribofuranosyl sugar with an oxidizing agent to produce an activated dialdehyde-terminated oligonucleotide and reacting said activated oligonucleotide with a linear or cyclic non-aromatic polyamine species under reducing conditions to yield said compound.

In accordance with other aspects of the invention compounds having the structure:



wherein B is a purine or pyrimidine heterocyclic base, R_8 and R_9 are independently H, PO_2 , a nucleotide, oligonucleotide or polyamine species and at least one of R_8 and R_9 is a purine containing oligonucleotide, and at least one of R_8 , R_9 and A is a species comprising the formula L_1-L_2 -polyamine wherein L_1 is an amino linker and L_2 is a heterobifunctional linker; and wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2 ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH are provided.

Therapeutic and diagnostic methods are also encompassed by the present invention. Methods of modulating the production of protein by an organism comprising contacting an organism with a compound having the structure of Formula I, Formula II, Formula III or Formula IV are encompassed by some embodiments of the present invention. In other aspects of the invention, methods of treating an animal having a disease characterized by undesired production of protein comprising contacting an animal with a compound having the structure of Formula I, Formula II, Formula III, or Formula IV in a pharmaceutically acceptable carrier are provided. Still other methods of the present invention provide methods for detecting the presence or absence of an RNA in a biological sample suspected of containing said RNA are provided comprising contacting a sample with a compound having the structure of Formula I, Formula II, Formula III or Formula IV wherein the compound is specifically hybridizable with the RNA and detecting the presence or absence of hybridization of the compound to the sample wherein hybridization is indicative of the presence of RNA in the sample.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a schematic representation of one preferred syntheses of compounds of Formula I.


FIG. 2 is a schematic representation of one preferred syntheses of compounds of Formula II.

FIG. 3 is a schematic representation of one preferred syntheses of compounds of Formula III.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention provides novel amine compounds useful for antisense therapy. In one embodiment of the present invention compounds having the structure:

6



wherein R_1 and R_2 are independently H, a nucleotide, oligonucleotide, or polyamine and at least one of R_1 and R_2 is a purine containing oligonucleotide, R_3 is a linear or cyclic non-aromatic polyamine species, and X is H, $O-R_{11}$, $S-R_{11}$, F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkynyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkylaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide, are provided. In some embodiments of the present invention both R_1 and R_2 are oligonucleotides, at least one of which includes at least one purine nucleotide.

In the context of this invention, the term "oligonucleotide" refers to a polynucleotide formed from naturally occurring bases, such as purine and pyrimidine heterocycles, and furanyl groups joined by native phosphodiester bonds. This term effectively refers to naturally occurring species or synthetic species formed from naturally occurring subunits or their close homologs. The term "oligonucleotide" may also refer to moieties which have portions similar to naturally occurring oligonucleotides but which have non-naturally occurring portions. Thus, oligonucleotides may have altered sugar moieties or inter-sugar linkages. Exemplary among these are the phosphorothioate and other sulfur-containing species which are known for use in the art. In accordance with some preferred embodiments, at least some of the phosphodiester bonds of the oligonucleotide have been substituted with a structure which functions to enhance the stability of the oligonucleotide or the ability of the oligonucleotide to penetrate into the region of cells where the viral RNA is located. It is preferred that such substitutions comprise phosphorothioate bonds, phosphotriesters, methyl phosphonate bonds, short chain alkyl or cycloalkyl structures or short chain heteroatomic or heterocyclic structures. Most preferred are $CH_2-NH-O-CH_2$, $CH_2-N(CH_3)-O-CH_2$, $CH_2-O-N(CH_3)-CH_2$, $CH_2-N(CH_3)-N(CH_3)-CH_2$ and $O-N(CH_3)-CH_2-CH_2$ structures where phosphodiester is $O-P-O-CH_2$). Also preferred are morpholino structures. Summerton, J. E. and Weller, D. D., U.S. Pat. No. 5,034,506 issued Jul. 23, 1991. In other preferred embodiments, such as the protein-nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide may be replaced with a polyamide backbone, the bases being bound directly or indirectly to the nitrogen atoms of the polyamide backbone. P. E. Nielsen, et al., *Science* 1991 254 1497. In accordance with other preferred embodiments, the phosphodiester bonds are substi-

tuted with other structures which are, at once, substantially non-ionic and non-chiral, or with structures which are chiral and enantiomerically specific. Still other linkages include the those disclosed in U.S. patent applications Ser. No. 566,836, filed Aug. 13, 1990, entitled Novel Nucleoside Analogs; Ser. No. 703,619, filed May 21, 1991, entitled Backbone Modified Oligonucleotide Analogs; Ser. No. 903,160, filed Jun. 24, 1992, entitled Heteroatomic Oligonucleoside Linkages; Ser. No. PCT/US92/04294, filed May 21, 1992, entitled Backbone Modified Oligonucleotides; and Ser. No. PCT/US92/04305, all assigned to the assignee of this invention. Persons of ordinary skill in the art will be able to select other linkages for use in practice of the invention.

Oligonucleotides may also include species which include at least some modified base forms. Thus, purines and pyrimidines other than those normally found in nature may be so employed. For example, deaza or aza purines and pyrimidines may be used in place of naturally purine or pyrimidine bases and pyrimidine bases having substituent groups at the 5- or 6-positions; purine bases having altered or replacement substituent groups at the 2-, 6- or 8-positions are also provided in some aspects of the present invention. Similarly, modifications on the furanosyl portion of the nucleotide subunits may also be effected, as long as the essential tenets of this invention are adhered to. Examples of such modifications are 2'-O-alkyl- and 2'-halogen-substituted nucleotides. Some specific examples of modifications at the 2' position of sugar moieties which are useful in the present invention are OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. Sugar mimetics such as cyclobutyls may also be used in place of the pentofuranosyl group. Exemplary modifications are disclosed in U.S. patent applications: Ser. No. 463,358, filed Jan. 11, 1990, entitled Compositions And Methods For Detecting And Modulating RNA Activity; Ser. No. 566,977, filed Aug. 13, 1990, entitled Sugar Modified Oligonucleotides That Detect And Modulate Gene Expression; Ser. No. 558,663, filed Jul. 27, 1990, entitled Novel Polyamine Conjugated Oligonucleotides; Ser. No. 558,806, filed Jul. 27, 1991, entitled Nuclease Resistant Pyrimidine Modified Oligonucleotides That Detect And Modulate Gene Expression; and Ser. No. PCT/US91/00243, filed Jan. 11, 1991, entitled Compositions and Methods For Detecting And Modulating RNA Activity; Ser. No. 777,670, filed Oct. 15, 1991, entitled Oligonucleotides Having Chiral Phosphorus Linkages; Ser. No. 814,961, filed Dec. 24, 1991, entitled Gapped 2' Modified Phosphorothioate Oligonucleotides; Ser. No. 808,201, filed Dec. 13, 1991, entitled Cyclobutyl Oligonucleotide Analogs; and Ser. No. 782,374, filed 782,374, entitled Derivatized Oligonucleotides Having Improved Uptake & Other Properties, all assigned to the assignee of this invention. The disclosures of all of the above noted patent applications are incorporated herein by reference. Oligonucleotides may also comprise other modifications consistent with the spirit of this invention. Such oligonucleotides are best described as being functionally interchangeable with yet structurally distinct from natural oligonucleotides. All such oligonucleotides are compre-

hended by this invention so long as they effectively function as subunits in the oligonucleotide. Thus, purine containing oligonucleotide are oligonucleotides comprising at least one purine base or analog thereof. In other embodiments of the present invention compounds of the present invention may be "subunits" of a species comprising two or more compounds of the present invention which together form a single oligonucleotide.

Oligonucleotides of the present invention may be naturally occurring or synthetically produced and may range in length from about 8 to about 50 nucleotides. In more preferred embodiments of the present invention said oligonucleotides may be from 8 to 40 nucleotides in length. Most preferably, oligonucleotides of the present invention may be from 12 to about 20 nucleotides in length.

The phrase polyamine species as used throughout the specification refers to species that have a plurality of nitrogen atoms thereon. Polyamines include primary amines, hydrazines, semicarbazines, thiosemicarbazines and similar nitrogenous species. Such species can be symmetrical species such as polyamine containing polymers or they can be unsymmetrical wherein the amine functionalities of the polyamine are separated in space by different moieties. In addition to carbon atoms other atomic species such as nitrogen and sulfur may also be incorporated into the polyamine species. In some preferred embodiments of the invention, at least one nitrogen atom of the polyamine has a free electron pair.

Preferred as polyamine species are species that range in length from about 3 to about 20 units. More preferably species having at least one nitrogen atom have the general formula H₂N[(CH₂)_nNH]_m— wherein n is an integer between 2 and 8 and m is an integer between 1 and 10. These species can be linear or cyclic. Cyclic amines would include crown amines and mixed crown amines/crown ethers.


Other suitable nitrogen-containing compound suitable for the formation of polyamine species include C₁-C₂₀ straight chain alkylamine, C₁-C₂₀ straight chain substituted alkylamine, C₂-C₅₀ branched chain alkylamine, C₂-C₅₀ branched chain substituted alkylamine, C₃-C₅₀ cyclic alkylamine, C₃-C₅₀ cyclic substituted alkylamine, C₂-C₂₀ straight chain alkenylamine, C₂-C₂₀ straight chain substituted alkenylamine, C₃-C₅₀ branched chain alkenylamine, C₃-C₅₀ branched chain substituted alkenylamine, C₃-C₅₀ cyclic alkenylamine, C₃-C₅₀ cyclic substituted alkenylamine, C₂-C₂₀ straight chain alkynylamine, C₂-C₂₀ straight chain substituted alkynylamine, C₃-C₅₀ branched chain alkynylamine, C₃-C₅₀ branched chain substituted alkynylamine, C₃-C₅₀ cyclic alkynylamine, C₃-C₅₀ cyclic substituted alkynylamine, C₁-C₂₀ straight chain alkyldiazine, C₁-C₅₀ straight chain substituted alkyldiazine, C₂-C₅₀ branched chain alkyldiazine, C₂-C₅₀ branched chain substituted alkyldiazine, C₃-C₅₀ cyclic hydrazoalkane, C₃-C₅₀ cyclic substituted hydrazoalkane, C₂-C₂₀ straight chain alkenylhydrazine, C₂-C₂₀ straight chain substituted alkenylhydrazine, C₃-C₅₀ branched chain alkenylhydrazine, C₃-C₅₀ branched chain substituted alkenylhydrazine, C₃-C₅₀ cyclic hydrazoalkene, C₃-C₅₀ cyclic substituted hydrazoalkene, C₂-C₂₀ straight chain alkynylhydrazine, C₂-C₂₀ straight chain substituted alkynylhydrazine, C₃-C₅₀ branched chain alkynylhydrazine, C₃-C₅₀ branched chain substituted alkynylhydrazine, C₃-C₅₀ cyclic hydrazoalkyne, C₃-C₅₀ cyclic substituted hydrazoalkyne, C₁-C₂₀ straight chain alkyldihydroxyamine, C₁-C₂₀ straight chain substituted alkyldihydroxyamine, C₂-C₅₀ branched chain alkyldihydroxyamine, C₂-C₅₀ branched chain substituted

alkylhydroxyamine, C₃-C₅₀ cyclic oxyalkylamine, C₃-C₅₀ cyclic substituted oxyalkylamine, C₂-C₂₀ straight chain alkenylhydroxyamine, C₂-C₂₀ straight chain substituted alkenylhydroxyamine, C₃-C₅₀ branched chain alkenylhydroxyamine, C₃-C₅₀ cyclic oxyalkenylamine, C₃-C₅₀ cyclic substituted oxyalkenylamine, C₂-C₂₀ straight chain alkynylhydroxyamine, C₂-C₂₀ straight chain substituted alkynylhydroxyamine, C₃-C₅₀ branched chain alkynylhydroxyamine, C₃-C₅₀ cyclic oxyalkynylamine, C₃-C₅₀ cyclic substituted oxyalkynylamine, C₁-C₂₀ straight chain alkylsemicarbazide, C₁-C₂₀ straight chain substituted alkylsemicarbazide, C₂-C₅₀ branched chain alkylsemicarbazide, C₂-C₅₀ branched chain substituted alkylsemicarbazide, C₃-C₅₀ cyclic alkylsemicarbazide, C₃-C₅₀ cyclic substituted alkylsemicarbazide, C₂-C₂₀ straight chain alkenylsemicarbazide, C₂-C₂₀ straight chain substituted alkenylsemicarbazide, C₃-C₅₀ branched chain alkenylsemicarbazide, C₃-C₅₀ branched chain substituted alkenylsemicarbazide, C₃-C₅₀ cyclic alkenylsemicarbazide, C₃-C₅₀ cyclic substituted alkenylsemicarbazide, C₂-C₂₀ straight chain alkynylsemicarbazide, C₂-C₂₀ straight chain substituted alkynylsemicarbazide, C₃-C₅₀ branched chain alkynylsemicarbazide, C₃-C₅₀ branched chain substituted alkynylsemicarbazide, C₃-C₅₀ cyclic alkynylsemicarbazide, C₃-C₅₀ cyclic substituted alkynylsemicarbazide, C₁-C₂₀ straight chain alkylthiosemicarbazide, C₁-C₂₀ straight chain substituted alkylthiosemicarbazide, C₂-C₅₀ branched chain alkylthiosemicarbazide, C₂-C₅₀ branched chain substituted alkylthiosemicarbazide, C₃-C₅₀ cyclic alkylthiosemicarbazide, C₃-C₅₀ cyclic substituted alkylthiosemicarbazide, C₂-C₂₀ straight chain alkenylthiosemicarbazide, C₂-C₂₀ straight chain substituted alkenylthiosemicarbazide, C₃-C₅₀ branched chain alkenylthiosemicarbazide, C₃-C₅₀ branched chain substituted alkenylthiosemicarbazide, C₃-C₅₀ cyclic alkenylthiosemicarbazide, C₃-C₅₀ cyclic substituted alkenylthiosemicarbazide, C₂-C₂₀ straight chain alkynylthiosemicarbazide, C₂-C₂₀ straight chain substituted alkynylthiosemicarbazide, C₃-C₅₀ branched chain alkynylthiosemicarbazide, C₃-C₅₀ branched chain substituted alkynylthiosemicarbazide, C₃-C₅₀ cyclic alkynylthiosemicarbazide, C₃-C₅₀ cyclic substituted alkynylthiosemicarbazide, C₁-C₂₀ straight chain alkylhydrazone, C₁-C₂₀ straight chain substituted alkylhydrazone, C₂-C₅₀ branched chain alkylhydrazone, C₂-C₅₀ branched chain substituted alkylhydrazone, C₃-C₅₀ cyclic hydrazoalkane, C₃-C₅₀ cyclic substituted hydrazoalkane, C₂-C₂₀ straight chain alkenylhydrazone, C₂-C₂₀ straight chain substituted alkenylhydrazone, C₃-C₅₀ branched chain alkenylhydrazone, C₃-C₅₀ branched chain substituted alkenylhydrazone, C₃-C₅₀ cyclic hydrazoalkene, C₃-C₅₀ cyclic substituted hydrazoalkene, C₂-C₂₀ straight chain alkynylhydrazone, C₂-C₂₀ straight chain substituted alkynylhydrazone, C₃-C₅₀ branched chain alkynylhydrazone, C₃-C₅₀ branched chain substituted alkynylhydrazone, C₃-C₅₀ cyclic hydrazoalkyne, C₃-C₅₀ cyclic substituted hydrazoalkyne, C₁-C₂₀ straight chain alkylhydrazide, C₁-C₂₀ straight chain substituted alkylhydrazide, C₃-C₅₀ branched chain alkylhydrazide, C₃-C₅₀ branched chain substituted alkylhydrazide, C₃-C₅₀ cyclic alkylhydrazide, C₃-C₅₀ cyclic substituted alkylhydrazide, C₂-C₂₀ straight chain alkenylhydrazide, C₂-C₂₀ straight chain substituted alkenylhydrazide, C₃-C₅₀ branched chain alkenylhydrazide, C₃-C₅₀ branched chain substituted alkenylhydrazide, C₃-C₅₀ cyclic

alkenylhydrazide, C₃-C₅₀ cyclic substituted alkenylhydrazide, C₂-C₂₀ straight chain alkynylhydrazide, C₂-C₂₀ straight chain substituted alkynylhydrazide, C₃-C₅₀ branched chain alkynylhydrazide, C₃-C₅₀ branched chain substituted alkynylhydrazide, C₃-C₅₀ cyclic alkynylhydrazide and C₃-C₅₀ cyclic substituted alkynylhydrazide.


In accordance with preferred embodiments of the present invention polyamine species are linear or cyclic non-aromatic polyamine species. In still more preferred embodiments of the present invention polyamine species are linear or cyclic non-aromatic comprising non-amide nitrogen atoms. By non-amide is meant a nitrogen which is not adjacent to a carbonyl group (i.e. C=O or C=S).

In still other embodiments of the present invention compounds having the structure:




wherein R₄ is an oligonucleotide and M is a pendent group having a polyamine species attached thereto are provided. The pendent group may be any group to which a polyamine may be attached. In preferred embodiments the pendent group is a R₁₀S or R₁₀NH, wherein R₁₀ is any of a broad range of reactive groups effective for subsequent attachment of polyamine species to the pendent group. Suitable for R₁₀ are substituted and un-substituted, straight chain or branched chained C₁-C₂₀ alkyl groups or substituted or un-substituted C₇-C₁₄ aryl groups having the nucleophile in one position thereon and a further functional group in a further position thereon. The pendent group may thus, subsequently functionalized with a bifunctional linker group amendable for attachment of a polyamine species to the pendent group. Alternatively the polyamine species may be directly attached to a pendent group such as by alkylation.

Further in accordance with the present invention are provided compounds having the structure:



wherein R₄ is an oligonucleotide, R₅ is a linear or cyclic non-aromatic polyamine species containing non-amide nitrogen atoms, and R₆ is H, a purine heterocycle or a pyrimidine heterocycle.

The present invention also provides novel polyamine containing compounds having the structure:



wherein B is a purine or pyrimidine heterocycle, R₈ and R₉ are independently H, PO₂⁻, a nucleotide, oligonucleotide or polyamine species and at least one of R₈ and R₉ is a purine containing oligonucleotide, and at least one of R₈, R₉ and A is a species comprising the formula L₁-L₂-polyamine


wherein L_1 is an amino linker and L_2 is a heterobifunctional linker; and wherein if R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- ; if R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide; and if A is not a polyamine species then A is H or OH.

Thus R_8 and R_9 may be oligonucleotides and A may be a species comprising the formula L_1 - L_2 -polyamine, or alternatively, R_8 may be an oligonucleotide and one or both of R_9 and A may be a species comprising the formula L_1 - L_2 -polyamine; or R_9 may be an oligonucleotide and one or both of R_8 and A may be a species comprising the formula L_1 - L_2 -polyamine. Furthermore, when R_8 is not a purine containing oligonucleotide or polyamine species, then R_8 is a nucleotide or PO_2^- . If R_9 is not a purine containing oligonucleotide or polyamine species, then R_9 is H or a nucleotide, and if A is not a polyamine species then A is H or OH.

In preferred embodiments of the present invention commercially available amino linkers may be used. For example, the 3' amino modifiers having the trade names C3 CPG and C7 CPG available through Glen Research may be employed. 5' amino modifiers may also be used such as C3 and C7 5' branched modifiers available through Glen Research. Similarly, 2' amino modifiers are also envisioned for use in some aspects of the present invention. The amino linkers are designed to functionalize a target oligonucleotide by the introduction of a primary amine at a designated site, be it 2', 3' or 5'. As will be apparent to one skilled in the art, any linker which meets this end is encompassed by the present invention.

Likewise, bifunctional linkers effective for purposes of the present invention are available commercially. For example, bis-(maleimido)-methyl ether (BMME), disuccinimidyl suberate (DSS), 3-maleimidobenzoyl-N-hydroxysuccinimide (MBS), maleimidohexanoyl-N-hydroxysuccinimide (MHS) and N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP) may be useful in some embodiments of the present invention. Other useful bifunctional linkers will be apparent to one skilled in the art as for instance from Pierce, Rockford, Ill.

Compounds of the present invention may be prepared by providing an oligonucleotide comprising one or more abasic sites. In the context of the present invention "abasic site" refers to a nucleotide unit in which the purine or pyrimidine group has been removed or replaced by a non-naturally occurring group such as a hydroxyl group. One or more abasic sites may be incorporated into one or more nucleotide bases of an oligonucleotide to form a synthon having the structure:



wherein R_1 and R_2 are independently H, a nucleotide, oligonucleotide or polyamine species and at least one of R_1 and R_2 is a purine containing oligonucleotide, and X is H, O- R_{11} , S- R_{11} , F, Cl, Br, CN, CF_3 , OCF_3 , OCN, $SOCH_3$, SO_2CH_3 , ONO_2 , N_3 , HN_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, a reporter molecule, an RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacody-

amic properties of an oligonucleotide wherein R_{11} is H, C_1 to C_{10} straight or branched chain lower alkyl or substituted lower alkyl, C_2 to C_{10} straight or branched chain lower alkenyl or substituted lower alkenyl, C_3 to C_{10} straight or branched chain lower alkynyl or substituted lower alkynyl, a ^{14}C containing lower alkyl, lower alkenyl or lower alkynyl, C_7 to C_{14} substituted or unsubstituted alkaryl or aralkyl, a ^{14}C containing C_7 to C_{14} alkaryl or aralkyl, alicyclic, heterocyclic, a reporter molecule, a RNA cleaving group, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide.

An enzymatic process may be used to produce such a synthon having abasic sites by reaction of a DNA glycosylase with an oligonucleotide starting material. For example, uracil DNA glycosylase act on uracil bases within an oligonucleotide to create abasic sites. Of course it should be recognized that enzymatic methods using DNA glycosylase may be less effective for oligonucleotides more closely resembling RNA such as oligonucleotides having 2' modifications.


Enzymes, as employed in the present invention, may be derived from naturally occurring sources or may be prepared by recombinant techniques. Many useful enzymes are available commercially.

Such synthons may alternatively be prepared by incorporation of abasic sites into an oligonucleotide via abasic sugar precursors. For example, 5-O-(4,4'-dimethoxytrityl)-1,2-dideoxy-1-(o-nitrobenzoyl)-D-ribofuranose-3-O-(2-Cyanoethyl-N,N'-diisopropyl) phosphoramidite may be prepared by modification of the procedures of Lyer, et al., *Nucleic Acids Research* 18: 2855 (1990) and Didier, et al., *Tetrahedron Letters* 32: 207 (1991).

Phosphoramidites having a 2' substitutions and abasic sites may also be prepared. For example, a synthon may have 2'-O-methyl or 2'-fluoro substitutions. Such phosphoramidite may be incorporated into an oligonucleotide by standard procedures. An o-nitrobenzyldeoxyfuranose containing oligonucleotide can be synthesized in accordance with these procedures. Post synthesis photolysis utilizing a high intensity Hg lamp generates the corresponding abasic site-containing polymer. In addition, other methods of introducing abasic sites at the 3', 5' and internal positions of an oligonucleotide to form a synthon are known to those skilled in the art. Thereafter the synthon may be reacted with a polyamine species under reducing conditions. As illustrated in FIG. 1, Step A, a compound may be prepared wherein B is uridine and an enzymatic process may be used to produce a synthon having abasic sites at one or more uridine sites by digestion of the compound with an enzyme such as uracil-DNA glycosylase. Other glycosylases will be effective for different targets. As described above, a glycosylase may be determined by the combined sequence of R_1 , R_2 and B. Some useful glycosylases and their respective targets are described for example, by Friedberg, DNA Repair (W. H. Freeman and Company, NY, 1985) p. 153, incorporated by reference herein in its entirety. These enzymes are commercially available or may be prepared from known procedures in the art.

In other embodiments of the present invention as exemplified in FIG. 2 compound of Formula II may be prepared by providing starting material having the structure:

13



wherein R_4 is an oligonucleotide, R_{12} is an oligonucleotide and B is urea or a heterocyclic base having a corresponding glycosylase and reacting the compound with an endonuclease to produce the compound 2 as described by Manoharan, et al., *JACS*, 1988, 110, 2690. Thereafter, the compound 2 is contacted with a pendent group such as $R_{10}S$, and reduced with the reducing agent $NaCNBH_4$ to stabilize the product 5. A polyamine species may then be added such as by alkylation to provide the final product 7. A polyamine species may alternatively be added directly to a bifunctional pendent group. Some endonucleases which will be useful in embodiments of the present invention are described, for example, in Doetsch and Cunningham, *Mutation Research*, 1990, 236, 173, incorporated by reference herein in its entirety. The endonuclease chosen will depend upon the identity of B and the sequence of R_4 and/or R_{12} . Thus, if B is a pyrimidine heterocycle, and the sequence of R_{12} begins with a pyrimidine, then an endonuclease such as T4 or *M. luteus* UV endonuclease may be chosen. Following digestion by T4 or *M. luteus* UV endonuclease, B and R_{12} are removed, resulting in a 3' terminal α,β unsaturated aldehydic species. In some instances, it may be desirable to engineer the sequence of the species so as to provide a endonuclease digestion site at a desired location.

Thus, in one preferred embodiment of the present invention R_4 may be TGGGAGCCATAGCGAGGC (SEQ ID NO: 1), B may be the pyrimidine thymine and R_{12} may be a thymidine dinucleotide. The net result of digestion of this species with T4 UV endonuclease will be TGGGAGCCATAGCGAGGCN (SEQ ID NO: 2) wherein N represents the aldehydic species.

Treatment of the digested compound with pendent group comprising a linker bearing a nucleophile results in the addition of the pendent group at the 3' terminus of the compound to join the linker to the digested compound. Suitable nucleophilic species include thiols and amines moieties as described above. In preferred embodiments of the present invention the pendent group is $R_{10}S^-$ or $R_{10}NH$. A polyamine species such as $NH_2(CH_2)_nNH_2$ wherein n is an integer from 1 to about 10 could be used as the attacking nucleophile by suitably blocking one end thereof and utilizing the other end as the attacking nucleophilic species. R_{10} can be further selected to provide a linkage or bridge between the nucleophile and a polyamine. Suitable for R_{10} are substituted and un-substituted, straight chain or branched chained C_1-C_{20} alkyl groups or substituted or un-substituted C_7-C_{14} aryl groups having the nucleophile in one position thereon and a further functional group in a further position thereon. After attachment of the pendent group via nucleophilic attack on compound 2, for attachment of the polyamine species the further functional group is then derivitized either via a bifunctional linking group, an alkylation type reaction or other derivation reaction known to those skilled in the art.

Upon addition of the pendent group to the digested compound, the double bond remaining on the digested compound is reduced to stabilize the product. Reducing agents effective to stabilize the end product of such a reaction are well known in the art. Some suitable reducing

14

agents include sodium cyanoborohydride, lithium cyanoborohydride and sodium borohydride.

Thereafter a polyamine may be added via an alkylation reaction or directly to a pendent group which is a bifunctional linker. The compound may further be derivitized by attaching one or more reactive groups to at least one of the nitrogen atoms of the polyamine species. Reactive groups include, but are not limited to reporter groups, alkylating agents, intercalating agents, RNA cleaving moieties, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins and cross-linking agents.

In accordance with other methods of the present invention compounds of Formula III may be prepared by reacting an oligonucleotide having a 3' ribofuranosyl sugar with an oxidizing agent to produce an dialdehyde-terminated activated oligonucleotide. Suitable oxidants include periodate solution, lead tetraacetate, activated MnO_2 , thallium (III) salts, pyridinium chlorochromate and O_2 catalyzed by Co (III) salts.

Thereafter the dialdehyde-terminated activated oligonucleotide is reacted with a polyamine species under reducing conditions. Reducing agents are known to those skilled in the art. Preferably, the activated oligonucleotide and species containing at least one nitrogen atom will be reacted in the presence of a solution of sodium cyanoborohydride, lithium cyanoborohydride or sodium borohydride.

In preferred embodiments of the present invention compounds may be produced as illustrated by FIG. 3, by preparation of an oligonucleotide having a 3' ribofuranosyl end followed by attack of the 3' ribofuranosyl ring by an oxidant such as m-periodate solution in 0.1M NaOAc buffer pH5, as described by Bayard, Bisbal and Lebleu in *Biochemistry* 25: 3730 (1986) to produce a dialdehyde-terminated activated oligonucleotide (FIG. 3, Step A). The activated oligonucleotide and a species containing four nitrogen atoms, spermine, can be reacted in the presence of the reducing agent, sodium cyanoborohydride (FIG. 3, Step B).

Compounds of the present invention are preferably specifically hybridizable with a target region. By "specifically hybridizable" herein is meant capable of forming a stable duplex with a target DNA or RNA. It is believed that oligonucleotides which form Watson-Crick base pairs, i.e. are complementary with target DNA or RNA and which specifically hybridize with target DNA or RNA inhibit the flow of genetic information from DNA to protein. In some embodiments of the present invention the oligonucleotide portions of compounds of the present invention are at least 70% complementary to a target sequence. In preferred embodiments of the present invention the oligonucleotide portions of compounds of the present invention are at least 80% complementary to a target sequence. 100% complementarity of the oligonucleotide portions of compounds of the present invention to a target sequence is most preferred. In preferred embodiments of the present invention, the oligonucleotide portions may be specifically hybridizable with DNA or RNA from papilloma virus, herpes viruses, human immunodeficiency virus, *Candida*, cytomegaloviruses, and influenza viruses. In addition, the oligonucleotide portions may also be specifically hybridizable with endogenous DNA or RNA of a cell. By oligonucleotide portions is meant R_1 and/or R_2 of Formula I, R_4 of Formula II, R_4 and/or R_6 of Formula III, or R_8 and/or R_9 of Formula IV.


For therapeutics, an animal suspected of having a disease characterized by excessive or abnormal production of a protein is treated by administering a compound having the

structure set forth in Formula I, Formula II, Formula III, or Formula IV in a pharmaceutically acceptable carrier. Most preferable, the compound is hybridizable with an RNA coding for the protein. Persons of ordinary skill in the art can easily determine optimum dosages, dosing methodologies and repetition rates. Such treatment is generally continued until either a cure is effected or a diminution in the diseased state is achieved. Long term treatment is likely for some diseases.

The compounds of the present invention will also be useful as a research reagent useful for the modulation of the production of a protein by an organism. Modulation may be accomplished by contacting the organism with compounds of the present invention having structures as set forth in Formula I, Formula II, Formula III, or Formula IV. Preferably the compounds are hybridizable with RNA coding for the protein.

Diagnostic applications include the detection of the presence or absence of an RNA in a sample suspected of containing RNA comprising contacting the sample with a compound having structures as set forth in Formula I, Formula II, Formula III or Formula IV wherein the compound is specifically hybridizable with the RNA and detecting the presence or absence of hybridization of the compound to the sample wherein hybridization is indicative of the present of the RNA in the sample.

It is also envisioned by the present invention to provide compounds in which at least one of the nitrogen atoms of the polyamine are derivatized with one or more of the group consisting of functionalities such as reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, crown amines, porphyrins and cross-linking agents. Therapeutic, diagnostic and research reagent applications are equally, or even more effective when the polyamine species further comprises such groups. Such compounds allow greater numbers of functionalities to be delivered to a target. For example, reporter groups such as biotin, fluorescent molecules and various fluorophores may be attached to compounds of the present invention to effect diagnostic ends, resulting in signal amplification as compared to conventional oligonucleotide-reporter group combinations. In a preferred embodiment of the present invention, biotin may be used to functionalize compounds of the present invention by reacting a compound with D-biotin-N-hydroxysuccinimide ester. In a further preferred embodiment, the polyamine species may be further functionalized by reacting the compound containing the polyamine species with an activated ester having the structure:



to form a compound with repeating imidazole catalytic cleaver units useful as an antisense therapeutic agents. Heterobifunctional linkers also can be utilized for attachment of intercalators, RNA cleaving agents including imidazoles, cell receptor binding molecules, steroids, alkylating agents, crown amines, porphyrins and cross-linkers to the polyamine species.

The following examples are illustrative but are not meant to be limiting of the present invention.

EXAMPLES

Example 1

Preparation of an Abasic Site Containing Oligonucleotide via Enzymatic Reaction

A. Synthesis of an Oligonucleotide Containing a Single Uridine Site

An oligonucleotide having the sequence CGC AGU CAG CC (SEQ ID NO:3) wherein U represents a 2' deoxyuridine nucleotide, was prepared by standard solid phase synthesis. The deoxyuridine nucleotide in the middle of the sequence was added during synthesis utilizing deoxyuridine phosphoramidite (Glen Research, Sterling, Va.). The oligonucleotide was prepared utilizing standard synthesis cycles. It was deprotected by normal deprotection at 55° C. utilizing ammonium hydroxide, 30%, for 16 hours. The solvent was evaporated and the residue was purified by HPLC and detritylated. Final purification was effected on Sephadex G-25.

B. Preparation of Enzyme Stock Solution

Uracil-DNA glycosylase was isolated from *E. Coli* M5219 cells transformed with the expression plasmid pBD396 containing the ung gene. The enzyme was purified by electrophoretic homogeneity as described by Lindahl et al., *J. Biol. Chem.* 252: 3286 (1977) and stored in 30 mM HEPES-NaOH, pH 7.4, containing 5% glycerol, 2 mM DTT and 1 mM EDTA.

C. Preparation of Oligonucleotide Containing Single Abasic Site

An abasic oligonucleotide of the sequence CGC AGN CAG CC (SEQ ID NO:4) wherein N represents an abasic site, was prepared by treating 237 O.D. units of an oligonucleotide having SEQ ID NO:1 of Example 1A in 0.5 ml water with 200 μ l of the stock solution of Example 1B (200 μ g of uracil DNA-glycosylase) and incubating at room temperature overnight. HPLC analysis showed quantitative removal of uracil as indicated by a 1:10 ratio between uracil and the abasic dodecamer oligonucleotide. The uracil retention time was 2.43 minutes and the abasic oligonucleotide was 21.68 minutes. The solution was lyophilized and stored in the freezer until further use.

D. Preparation of Oligonucleotide Containing Multiple Uridine Sites

In the manner of Example 1A the following oligonucleotide was prepared GAC AGA GGU AGG AGA AGU GA (SEQ ID NO: 5) wherein U represents a 2'-deoxyuridine nucleotide. The oligonucleotide is treated in accordance with the procedure of Example 1C resulting in an oligonucleotide of the sequence GAC AGA GGN AGG AGA AGN GA (SEQ ID NO: 6) wherein N represents an abasic site within the oligonucleotide.

Example 2

Preparation of an Abasic Site Containing Oligonucleotide via a Abasic Sugar Precursor

A. Preparation of 5-O-4,4'-Dimethoxytrityl-1,2-Dideoxy-1-(o-nitrobenzoyl)-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

5-O-4,4'-dimethoxytrityl-1,2-dideoxy-D-ribofuranose-3-O-(2-cyanoethyl-N,N'-diisopropyl) phosphoramidite is prepared in accordance with modification of the procedures of Lyer, et al. *Nucleic Acids Research* 18: 2855 (1990) and Didier, et al., *Tetrahedron Letters* 32: 207 (1991) incorporated by reference herein in their entireties.

B. Preparation of Oligonucleotide Containing Abasic Site

Oligonucleotide having the sequence CGC AGN CAG CC wherein N represents an abasic site (SEQ ID NO:4) from Example 1C can be prepared in accordance with modifications of the procedures of Lyer, et al. *Nucleic Acids Research* 18: 2855 (1990) and Didier, et al., *Tetrahedron Letters* 32: 207 (1991). In accordance with these procedures, an o-nitrobenzyl deoxyfuranose containing oligonucleotide is synthesized using the oligonucleotide synthetic methods of Lyer, et al. and Didier, et al. Photolysis utilizing a high intensity Hg lamp (300 nm) generates the corresponding abasic site containing oligonucleotide. Such abasic oligonucleotides are also described in Horn, et al., *Nucleosides and Nucleotides* 10:299 (1991).

Example 3

Preparation of Modified Abasic Sugar Precursors

A. Preparation of 5-O-(4,4'-Dimethoxytrityl)-2-O-Methyl-1,2-Dideoxy-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

1-O-methyl-D-ribofuranose is 3,5 protected with TIPS-Cl₂. It is then 2-position methylated with either diazomethane or methyl iodide/silver oxide (CH₃I/Ag₂O). The composition is then treated with an acetic anhydride/acetic acid/sulfuric acid mixture to give a 1-O-acetyl, 2-O-methyl 3,5 protected sugar. The 1-O-acetyl, 2-O-methyl 3,5 protected sugar is deprotected with tetrabutyl ammonium fluoride, 5-position dimethoxytritylated, and 3-position phosphitylated. Thereafter, this phosphoramidite may be incorporated into an oligonucleotide by standard phosphoramidite procedures and ammonia deprotected to form a 2'-O-methyl, 1' abasic site containing oligonucleotide.

B. Preparation of 5-O-4,4'-Dimethoxytrityl-2-O-Methyl-1,2-Dideoxy-1-(o-nitrobenzoyl)-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

1-O-acetyl 2,3,5-tri-O-benzoyl-D-ribofuranose is condensed with o-nitrobenzyl alcohol under Vorbruggen conditions. The resultant 1-O-(ortho-nitrobenzyl)-2,3,5-tri-O-benzoyl (α,δ)-D-ribofuranose is deprotected with ammonia and subsequently treated with TIPS-Cl₂. The resultant 3,5-silyl protected 1-O-(ortho-nitro benzyl) D-ribofuranose is reacted with diazomethane or CH₃I/Ag₂O to give the required 2-O-methyl compound. Subsequent 3,5-deprotection, 5-dimethoxytritylation and 3-phosphitylation gives the named phosphoramidite. The phosphoramidite can be incorporated into an oligonucleotide via standard phosphoramidite procedures.

C. Preparation of 5-O-(4,4'-Dimethoxytrityl)-2-Fluoro-1,2-Dideoxy-D-Ribofuranose-3-O-(2-Cyanoethyl-N,N'-Diisopropyl) Phosphoramidite.

1-O-(ortho-nitrobenzyl)-2,3,5-tri-O-benzoyl-D-ribofuranose is deprotected at 2,3,5 positions using ammonia. Tritylation with excess trityl chloride/pyridine/4-dimethylaminopyridine gives 3-5-ditrityl-1-O-nitrobenzyl-D-ribofuranose. Oxidation at 2 position with CrO₃ followed by NaBH₄ reduction inverts the configuration at 2 position yielding an arabino sugar. The arabino sugar is converted to its triflate at 2 position and the triflate is displaced with fluoride ion to yield the 2-fluoride modified sugar which can be 5 position protected and phosphitylated to incorporate the sugar into an oligonucleotide via standard oligonucleotide synthesis.

Example 4

Oligonucleotides conjugated in the following example are set forth in Table 2.

TABLE II

OLIGOMER (SEQ ID NO.)	TARGET	SEQUENCE	LINKER (L)
A (SEQ ID NO: 7)	ICAM	TGG GAG CCA TAG CGA GGC-L	3-carbon amino
B (SEQ ID NO: 7)	ICAM	TGG GAG CCA TAG CGA GGC-L	3-carbon amino
C (SEQ ID NO: 8)	BPV	CTG TCT CCA* TCC TCT TCA CT	2'aminopentoxo
D (SEQ ID NO: 9)	BPV	CTG TCT CCA TCC TCT TCA	3-carbon amino
E (SEQ ID NO: 9)	BPV	CTG TCT CCA TCC TCT TCA	6-carbon amino
F (SEQ ID NO: 10)	CMV	GGC GUC UCC AGG CGA UCU GAC*	2'-OMe
G (SEQ ID NO: 11)	ICAM	TCT GAG TAG CAG AGG AGC TC*	2'-OMe
H (SEQ ID NO: 12)		GGA UGG CGU CUC CAG GCG AUC*	2'-OMe
I (SEQ ID NO: 13)		GGA UGG CGU CUC CAG GCG AUC-L	3-carbon amino
J (SEQ ID NO: 13)		GGA UGG CGU CUC CAG GCG AUC-L	6-carbon amino
K (SEQ ID NO: 7)		F-TGG GAG CCA TAG CGA GGC-L	3-carbon amino

30 A* = 2'-O-aminopentoxo-2'-deoxyadenosine
C* = 2'-aminopropoxy cytosine
F = Fluorocsein

A. 3' Terminus Polyamine End Labeled Oligonucleotide

1. 3'-Terminus Polyamine Oligonucleotide I

35 Polyamines were attached to the 3'-terminus end of a phosphodiester oligonucleotide having the sequence D-polyamine [(SEQ ID NO: 9)-polyamine], wherein the polyamine is one of the following:

TABLE III

1,6 Diaminohexane	Oligomer D (i)
Diethylenetriamine	Oligomer D (ii)
Triethylenetetramine	Oligomer D (iii)
Spermine	Oligomer D (iv)
Pentaethylenehexamine	Oligomer D (v)

a. Preparation of the Intermediate Linker

The oligonucleotide sequence having a 3'-terminus amino group was synthesized using 3'-amino modifier (with a three carbon linker) controlled pore glass (CPG) from Glen Research as the solid support. The synthesis was conducted with an Applied Biosystems 380B or 994 in the "Trityl-Off" mode. The resultant oligonucleotide was cleaved from the solid support and deprotected with concentrated NH₄OH for 16 hrs at 55° C. Purification on a Sephadex G-25 column yielded a 3'-amino modified oligonucleotide of the specified sequence.

b. Preparation of Polyamine Functionalized Oligonucleotide

60 The crude 3'-aminolinker-oligonucleotide (SEQ ID NO:9) (15 O.D. units, approximately 85 nmols) was dissolved in freshly prepared NaHCO₃ buffer (150 ul, 0.2M, pH 8.1) and treated with a solution of disuccinimidyl suberate (DSS) (approximately 5 mgs) dissolved in 150 ul of methyl sulfoxide (DMSO). The reaction mixture was left to react for 20 minutes at room temperature. The mixture was then passed over a Sephadex G-25 column (0.7x45 cm) to separate the

activated oligonucleotide-DSS from the excess DSS. The oligonucleotide-DSS was then frozen immediately and lyophilized to dryness. A solution of polyamine in 0.33M NaOAc (approximately 6 mg polyamine in 300 ul 0.33M NaOAc, pH 5.2, final solution pH 6–8.0) was added to the dried oligonucleotide-DSS, and this mixture was allowed to react overnight at room temperature. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 4.

TABLE IV

Oligomer	Retention Time
unreacted D	26.44 mins
Oligomer D (i)	27.48 mins
Oligomer D (ii)	27.23 mins
Oligomer D (iii)	27.27 mins
Oligomer D (iv)	27.54 mins
Oligomer D (v)	27.36 mins

In a second test run under the same conditions the HPLC gradient was 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 15% solvent B in 60 minutes. HPLC retention times were as set forth in Table 5.

TABLE V

Oligomer	Retention Time
unreacted D	60.74 mins
Oligomer D (ii)	62.37 mins
Oligomer D (v)	65.24 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313–107)

c. Nuclease stability of 3' polyamine conjugates in Fetal Calf Serum

Polyamine conjugates of the invention are assessed for their resistance to serum nucleases by incubation of the oligonucleotides in media containing various concentrations of fetal calf serum. Labeled oligonucleotides are incubated for various times, treated with protease K and then analyzed by gel electrophoresis on 20% polyacrylamide-urea denaturing gels and subsequent autoradiography or phosphorimaging. Autoradiograms are quantitated by laser densitometry. Based upon the location of the modifications and the known length of the oligonucleotide it is possible to determine the effect of the particular modification on nuclease degradation. For the cytoplasmic nucleases, a HL60 cell line is used. A post-mitochondrial supernatant is prepared by differential centrifugation and the labeled oligonucleotides are incubated in this supernatant for various times. Following the incubation, oligonucleotides are assessed for degradation as outlined above for serum nucleolytic degradation. Autoradiography results are quantitated for comparison of the unmodified and the modified oligonucleotides. The $t_{1/2}$ are set forth below.

TABLE VI

Oligonucleotide	$t_{1/2}$ (hours)
wild type oligomer D	0.5 (no aminolinker)
unreacted oligomer D	22 (with aminolinker)
oligomer D (ii)	48
oligomer D (v)	>50

2. 3'-Terminus Polyamine Conjugate II

Polyamines were attached to the 3'-terminus end of a phosphodiester oligonucleotide having the sequence E-polyamine [(SEQ ID NO: 9)-polyamine] wherein the polyamine is one of the following:

TABLE VII

Diethylenetriamine	Oligomer E (i)
Pentaethylenehexamine	Oligomer E (ii)

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a substituting a 3' amino modifier with a six carbon linker (Clontech, Palo Alto, Calif.) for the 3'-amino modifier (with a three carbon linker).

3. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide was prepared in accordance with Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 25% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 8.

TABLE VIII

Oligomer	Retention Time
unreacted E	41.38 mins
Oligomer E (i)	43.29 mins
Oligomer E (ii)	43.43 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 353-35).

4. 3' -Terminus Polyamine Conjugate III

Polyamines were attached to the 3'-terminus end of a phosphorothioate oligonucleotide having the sequence A-polyamine [(SEQ ID NO:7)-polyamine] where the polyamine is one of the following:

TABLE IX

1,6 Diaminohexane	Oligomer A (i)
Diethylenetriamine	Oligomer A (ii)
Triethylenetetramine	Oligomer A (iii)
Spermine	Oligomer A (iv)
Pentaethylenehexamine	Oligomer A (v)

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a utilizing the Beaucage reagent (3H-1,2-benzodithioate-3-one 1,1-dioxide, see Radhakrishnan, P. I., Egan, W., Regan, J. B. and Beaucage, S. L., (1990) *J. Am. Chem. Soc.*, 112:1253) to form the phosphothioate inter-

21

nucleotide backbone. The 3'-aminolinker was introduced as described in example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

Oligonucleotides were functionalized as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 10.

TABLE X

Oligomer	Retention Time
unreacted A	30.77 mins
Oligomer A (iii)	31.31 mins
Oligomer A (v)	31.02 mins

In a second test run under the same conditions, the HPLC gradient was 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 15% solvent B in 60 minutes. Retention times were as set forth in Table 11.

TABLE XI

Oligomer	Retention Time
unreacted A	68.62 mins
Oligomer A (i)	68.70 mins
Oligomer A (ii)	68.69 mins

In a second test run under the same conditions, HPLC retention times were as set forth in Table 12.

TABLE XII

Oligomer	Retention Time
unreacted A	30.34 mins
Oligomer A (iv)	30.57 mins
Oligomer A (v)	30.72 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Test run 1 Gel, 313-82; Test run 2 Gel, 285–138; Test run 3 Gel, 353-57)

C. Preparation of Biotin Functionalized Oligonucleotide Polyamine Conjugate

To further characterize the oligonucleotide polyamine conjugate, biotin was attached to the free amines made available by the polyamines attached in Example 4-A-4-b. About 10 O.D. units (A₂₆₀) of Oligomers A(i) and A(ii) (approximately 58 nmoles) were dried in a microfuge tube. The oligonucleotide polyamine conjugate was rehydrated in 400 ul of 0.2M NaHCO₃ (pH 8.1) buffer and D-biotin-N-hydroxysuccinimide ester (approximately 5.0 mgs biotin for the 1,6 Diaminohexane conjugate, 8.0 mgs for the Diethylenetriamine) (Sigma) was added followed by 200 ul of DMF. The solution was left to react overnight at room temperature. The solution was then passed over a NAP-25 column and analyzed by reverse phase HPLC. Solvent A was 50 mM TEAA and solvent B was CH₃CN. The HPLC gradient was 0–10 mins, 95% A, 5% B; linear increase to 40% B in the next 50 minutes using a Water's Delta-Pak

22

C-18, reverse phase column. The HPLC retention times were as set forth in Table 13.

TABLE XIII

Oligomer	Retention Time
unreacted A	30.77 mins
Oligomer A (i)	31.31 mins
Oligomer A (i)-Biotin	35.56 mins
Oligomer A (ii)	31.02 mins
Oligomer A (ii)-Biotin	36.23 mins

5. 3' -Terminus Polyamine Conjugate IV

Polyamines were attached to the 3'-terminus end of the phosphodiester oligonucleotide having the sequence B-polyamine [(SEQ ID NO: 7)-polyamine] wherein the polyamine is one of the following:

TABLE XIV

	Oligomer B (i)
Diethylenetriamine	Oligomer B (i)
Triethylenetetramine	Oligomer B (ii)
Spermine	Oligomer B (iii)
Pentaethylenehexamine	Oligomer B (iv)

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

The oligonucleotide was functionalized with polyamines as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 15.

TABLE XV

Oligomer	Retention Time
untreated B	25.71 mins
Oligomer B (i)	26.11 mins
Oligomer B (ii)	25.26 mins
Oligomer B (iii)	25.10 mins
Oligomer B (iv)	25.12 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313-112)

B. 2' Internal Polyamine Labeled Oligonucleotide

1. 2'-Internal Polyamine oligonucleotide I

Polyamines were attached to the 2'-internal linker site of a phosphodiester oligonucleotide having the sequence C-polyamine [(SEQ ID NO: 8)-polyamine] wherein the polyamine is one of the following:

TABLE XVI

	Oligomer C (i)
Diethylenetriamine	Oligomer C (i)
Triethylenetetramine	Oligomer C (ii)
Pentaethylenehexamine	Oligomer C (iii)

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a incorporating a modified adenosine phos-

phoramidite (with a 2'-aminolinker) at position #9. This oligonucleotide and the 2'-amino linker have been described in Manoharan, M., Guinosso, C. J., Cook, P. D., *Tetrahedron Letters*, 32, (pgs. 7171-7174), 1991.

b. Preparation of Polyamine Functionalized Oligonucleotide

The oligonucleotide was functionalized as described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using a Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 17.

TABLE XVII

Oligomer	Retention Time
untreated C	26.20 mins
Oligomer C (i)	27.52 mins
Oligomer C (ii)	27.50 mins
Oligomer C (iii)	27.59 mins

Gel analysis showed progressively slower migration times for the polyamine conjugates (the larger the polyamine, the slower the migration) versus the oligonucleotide alone. (Gel: 313-97)

C. 3' Terminus Polyamine End Labeled Oligonucleotide, Using a 2'-aminolinker

1. 3' Terminus Polyamine Labeled Oligonucleotide I, Using a 2'-aminolinker

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'-aminolinker having the sequence F-polyamine [(SEQ ID NO:10)-polyamine] wherein the polyamine is pentaethyl-enehexamine (oligomer F(i)).

a. Preparation of the Intermediate Linker

The intermediate linker was prepared as described in Example 4-A-1-a, except that a modified cytosine CPG (with a 2'-propylaminolinker) was introduced at the 3' end. The 2'-modification can be prepared by modification of the procedure previously described in application Ser. No. 918, 362 filed Jul. 23, 1992. The CPG containing 2'-w-phthalimido-propoxy-cytidine was synthesized according to the standard protocols reported in the literature. See, for example, B. S. Sproat and A. I. Lamond, in "Oligonucleotides and Analogues" edited by F. Eckstein, IRL Press at Oxford University Press (1991) p71-72.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide was prepared in accordance with Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 18.

TABLE XVIII

Oligomer	Retention Time
unreacted F	28.53 mins
oligomer F (i)	29.47 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

2. 3' Terminus Polyamine Labeled Oligonucleotide II, Using a 2'-aminolinker

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'-aminolinker having the sequence G-polyamine [(SEQ ID NO:11)-polyamine] wherein the polyamine is pentaethyl-enehexamine (oligomer G(i)).

a. Preparation of the Intermediate Linker

The intermediate linker was prepared in accordance with the method described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide was prepared in accordance with the procedures described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 19.

TABLE XIX

Oligomer	Retention Time
unreacted G	28.43 mins
oligomer G (i)	29.06 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

3. 3' Terminus Polyamine Labeled Oligonucleotide III Using a 2'-aminolinker

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide via a 2'-aminolinker having the sequence H-polyamine [(SEQ ID NO:12)-polyamine] wherein the polyamine is pentaethyl-enehexamine.

a. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0-10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 20.

TABLE XX

Oligomer	Retention Time
unreacted H	28.49 mins
oligomer H (i)	30.36 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

25

Example 5

Polyamine Labeled 2'-OMe Oligonucleotides and Other RNA Mimics

1. Polyamine Labeled 2'-OMe Oligonucleotide I

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 3 carbon linker) having the sequence I-polyamine [(SEQ ID NO:13)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer I(i)).

a. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 21.

TABLE XXI

Oligomer	Retention Time
unreacted I	28.93 mins
oligomer I (i)	29.59 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 353-156)

2. Polyamine Labeled 2'-OMe Oligonucleotide II

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 6 carbon linker) having the sequence J-polyamine [(SEQ ID NO:13)-polyamine] wherein the polyamine is pentaethylenehexamine (oligomer J(i)).

a. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-b. The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 22.

TABLE XXII

Oligomer	Retention Time
unreacted J	28.76 mins
oligomer J (i)	29.39 mins

Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 397-85)

3. Polyamine Labeled 2'-OMe oligonucleotide III

Polyamines were attached to the 3'-terminus end of a phosphodiester (2'-OMe) oligonucleotide (via a 3 carbon linker) having another reporter group (such as biotin,

26

fluorescein) at the other end in the sequence K-polyamine [(SEQ ID NO:7)-polyamine]. Fluorescein at 5' end was added using the required amidite commercially available from Clontech. The polyamine is one of the following

5 pentaethylenehexamine oligomer K(i)

spermine oligomer K(ii)

a. Preparation of the Intermediate Linker

The intermediate linker is prepared in accordance with methods described in Example 4-A-1-a.

10 b. Preparation of Polyamine Functionalized Oligonucleotide

The polyamine functionalized oligonucleotide is prepared in accordance with methods described in Example 4-A-1-b.

15 The resulting polyamine-oligonucleotide conjugate was characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 23.

TABLE XXIII

Oligomer	Retention Time
unreacted K	31.35 mins
oligomer K (i)	31.96 mins
oligomer K (ii)	32.15 mins

30 Gel analysis showed progressively slower migration times for the polyamine conjugate versus the oligonucleotide alone. (Gel: 353-149)

Example 6

5' Terminus Polyamine End Labeled Oligonucleotide

1. 5'-Terminus Polyamine Oligonucleotide I

Polyamines were attached to the 5'-terminus end of a phosphodiester oligonucleotide having the following sequences:

40 5'-aminolinker-TCAG (oligomer L)

5'-aminolinker-CGCACGC (oligomer M) to provide the polyamine oligonucleotides:

5'-polyamine-TCAG (oligomer L(i))

5'-polyamine-CGCACGC (oligomer M(i)) wherein the polyamine is pentaethylenehexamine.

a. Preparation of the Intermediate Linker

The oligonucleotide sequence having a 5'-terminus amino group was synthesized using Aminolink-II (with a six carbon linker) phosphoramidite from Applied Biosystems in the last round of synthesis. The synthesis was conducted with an Applied Biosystems 380B or 994 in the "Trityl-On" mode. The resultant oligonucleotide was cleaved from the solid support and deprotected with concentrated NH₄OH for 16 hrs at 55° C. Purification on a Sephadex G-25 column yielded a 5'-amino modified oligonucleotide of the specified sequence.

b. Preparation of Polyamine Functionalized Oligonucleotide L(i)

60 The crude 5'-aminolinker-oligonucleotide (150 O.D. units, approximately 3.75 mmols) was dissolved in freshly prepared NaHCO₃ buffer (900 ul, 0.2M, pH 8.1) and treated with a solution of disuccinimidyl suberate (DSS) (approximately 30 mgs) dissolved in 750 ul of methyl sulfoxide (DMSO). The reaction mixture was left to react for 20 minutes at room temperature. The mixture was divided into three portions and then passed over a Sephadex G-25 column (0.7x45 cmx3columns) to separate the activated

oligonucleotide-DSS from the excess DSS. The oligonucleotide-DSS was then frozen immediately and lyophilized to dryness. A solution of polyamine in 0.33M NaOAc (approximately 60 mL polyamine in 1950 ul 0.33M NaOAc, pH 5.2, final solution pH 6–8.0) was added to the dried oligonucleotide-DSS, and this mixture was allowed to react overnight at room temperature.

c. Preparation of Polyamine Functionalized Oligonucleotide M(i)

The crude 5'-aminolinker-oligonucleotide (oligomer M) (150 O.D. units, approximately 2.50) was reacted as described in Example 6(b).

d. Characterization of 5' Polyamine Functionalized Oligonucleotides

The resulting polyamine-oligonucleotide conjugates were characterized by reverse phase HPLC and a 20% denaturing gel. Solvent A was 50 mM TEAA, solvent B was CH₃CN. The HPLC gradient was from 0–10 mins, 95% solvent A, 5% solvent B; linear increase to 40% solvent B in the next 50 minutes using Water's Delta-Pak C-18 reverse phase column. HPLC retention times were as set forth in Table 24.

TABLE XXIV

Oligomer	DNA TARGET			RNA TARGET		
	T _m (°C.)	ΔT _m (°C.)	ΔΔG° _{37°C.} (°C.)	(°C.)	(°C.)	ΔΔG° _{37°C.} (°C.)
wild type	60.6	—	—	64.9	—	—
oligomer D	60.3	-0.3	+0.3	64.6	-0.3	0.0
oligomer D + 5'-6-carbon amino linker)	60.8	+0.2	0.0	65.1	+0.2	0.0
oligomer E	60.8	+0.2	-0.8	65.8	+0.9	-1.0
oligomer E(i)	61.2	+0.6	-1.4	66.3	+1.4	-1.9
oligomer E + spermine	61.5	+0.9	-1.7	67.1	+2.2	-2.1
oligomer E(ii)	61.2	+0.6	-1.3	67.5	+2.6	-2.6

Example 7

Preparation of a Reactive Site containing Oligonucleotide

An oligonucleotide having the sequence TGGGAGC-CATAGCGAGGUCT (SEQ ID NO: 14) is treated with uracil DNA glycosylase followed by T4 endonuclease. The product is then treated with 1-phthalimidobutyl-4-thiol. Nucleophilic attack by the thiol adds to the protected aminobutyl moiety to what was the 3' position of the opened nucleotide. Treatment of this composition with hydrazine will deblock the phthalimide yielding an amino species which is then treated with bifunctional linker followed by treatment with an appropriate polyamine species as per Example 4-A-1-b.

Example 8

Preparation of Polyamine Conjugated Oligonucleotide

An oligonucleotide is prepared as described in Example 7 treating the product with NH₂—CH₂—CH₂—SH. The thiol group will attack the double bond of the opened nucleotide. This species may then be further derivatized with a reactive group.

Example 9

Thermodynamic Parameters of Oligoamine-Oligonucleotide Conjugates with DNA and RNA Targets

The ability of the functionalized oligonucleotides of the invention to hybridize to their complementary RNA or DNA sequences is determined by thermal melting analysis. The RNA complement is synthesized from T7 RNA polymerase and a template-promoter of DNA synthesized with an Applied Biosystems, Inc. 380B nucleic acid synthesizer. The RNA species is purified by ion exchange using FPLC (LKB Pharmacia, Inc.) or by denaturing urea-PAGE. Natural antisense oligonucleotides or those containing functionalization at specific locations are added to either the RNA or DNA complement at stoichiometric concentrations to form hybrid duplexes. The absorbance (260 nm) hyperchromicity dependence on temperature upon duplex to random coil transition is monitored using a Gilford Response II spectrophotometer. These measurements are performed in a buffer of 10 mM Na-phosphate, pH 7.4, 0.1 mM EDTA, and NaCl to yield an ionic strength of either 0.1M or 1.0M. Data are analyzed by a graphic representation of 1/T_m vs ln[Ct], where [Ct] is the total oligonucleotide concentration. From this analysis the thermodynamic parameters are determined. Based upon the information gained concerning the stability of the duplex or hetero-duplex formed, the placement of the polyamines into oligonucleotides is assessed for its effects on helix stability. Modifications that drastically alter the stability of the hybrid exhibit reductions or enhancements in the free energy (delta G) and decisions concerning their usefulness in antisense oligonucleotides are made.

Example 10

Conjugation of Polyamines to Abasics Sites Containing Oligonucleotides

To 15.2 ODS of an abasic oligonucleotide (SEQ ID NO: 4) in 100 μl water was added 25 μl 1M NaOAc (pH 5.0) solution. The final concentration of the acetate buffer was 0.2M. 5.3 mg of triethylenetetramine was dissolved in 500 μl of 1M NaOAc (pH 5.0) solution. 50 μl of the resulting solution was added to the oligonucleotide solution followed by 50 μl of NaCNBH₃ (57 MM solution). The pH of the resulting solution was below 8.0. The solution was vortexed and left to stand overnight. HPLC and Gel analysis indicated conjugation of the triethylenetetramine to the oligonucleotide. The conjugated oligonucleotide was purified by G-25 and HPLC. HPLC retention times are set forth in Table 25.

TABLE XXV

Oligomer	Retention time (mins)
parent oligonucleotide (SEQ ID NO: 3)	26.66
abasic oligonucleotide (SEQ ID NO: 4)	26.16
(SEQ ID NO: 4)-triethylenetetramine conjugate	26.04

-continued

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:1:

T G G G A G C C A T A G C G A G G C

1 8

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 19
- (D) OTHER INFORMATION: /note= "abasic, aldehydic species"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:2:

T G G G A G C C A T A G C G A G G C N

1 9

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "2'deoxyuridine residue"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:3:

C G C A G U C A G C C

1 1

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 6
- (D) OTHER INFORMATION: /note= "abasic residue"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:4:

C G C A G N C A G C C

1 1

(2) INFORMATION FOR SEQ ID NO:5:

-continued

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 9
 (D) OTHER INFORMATION: /note= "2'-deoxyuridine residue"

(i x) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 18
 (D) OTHER INFORMATION: /note= "2'-deoxyuridine residue"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:
 G A C A G A G G U A G G A G A A G U G A 2 0

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 9
 (D) OTHER INFORMATION: /note= "abasic residue"

(i x) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 18
 (D) OTHER INFORMATION: /note= "abasic residue"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:6:
 G A C A G A G G N A G G A G A A G N G A 2 0

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:7:
 T G G G A G C C A T A G C G A G G C 1 8

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 9
 (D) OTHER INFORMATION: /note= " 2'-O-aminopentoxo-2'-deoxyadenosine"

-continued

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CTGTCTCCAT CCTCTTCACT

2 0

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTGTCTCCAT CCTCTTCACT

2 0

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 21
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGCGUCUCCA GGCGAUCUGA C

2 1

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 20
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TCTGAGTAGC AGAGGAGCTC

2 0

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 21
- (D) OTHER INFORMATION: /note= "2'-aminopropoxy cytosine"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:12:

-continued

GGAUGGCGUC UCCAGGCGAU C

2 1

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGAUGGCGUC UCCAGGCGAU C

2 1

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (genomic)


(x i) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TGGGAGCCAT AGCGAGGUCT

2 0

What is claimed is:

1. A compound having the structure:



wherein R₄ is an oligonucleotide and M is a pendent group having a polyamine species attached thereto.

30

2. The compound of claim 1 wherein M is R₇S or R₇NH wherein R₇ is a polyamine species.

3. The compound of claim 2 wherein the polyamine species comprises at least one nitrogen atom having a free electron pair.

35 4. The compound of claim 1 further including one or more of reporter groups, alkylating agents, intercalating agents, cell receptor binding molecules, steroids, peptides, crown amines, porphyrins, or cross-linking agents attached to at least one of the nitrogen atoms of said polyamine.

40 5. The compound of claim 1 in a pharmaceutically acceptable carrier.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5,834,607
DATED : November 10, 1998
INVENTOR(S) : Muthiah Manoharan

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 3,

Line 42, delete "alkylaryl" and insert therefor -- alkyaryl --;

Column 3,

2nd compound, please insert -- II --.

Column 4,

Line 14, delete "heterocycoalkyl" and insert therefor -- heterocycloalkyl --;

Line 25, delete "alkylaryl" and insert therefor -- alkyaryl --.

Column 6,

Line 36, delete "furanyl" and insert therefor -- furanosyl --.

Column 8,

Line 53, delete "₃-C₅₀" and insert therefor-- C₃-C₅₀ --.

Column 9,

Line 26, delete "cyclicalkynylsemicarbazide" and insert therefor -- cyclic alkynylsemicarbazide --

Signed and Sealed this

Second Day of October, 2001

Attest:

Nicholas P. Godici

Attesting Officer

NICHOLAS P. GODICI
Acting Director of the United States Patent and Trademark Office