




(86) Date de dépôt PCT/PCT Filing Date: 2005/05/27
 (87) Date publication PCT/PCT Publication Date: 2005/12/15
 (85) Entrée phase nationale/National Entry: 2006/11/16
 (86) N° demande PCT/PCT Application No.: US 2005/018949
 (87) N° publication PCT/PCT Publication No.: 2005/117846
 (30) Priorité/Priority: 2004/05/27 (US60/574,993)

(51) Cl.Int./Int.Cl. *A61K 31/401* (2006.01),
A61K 31/551 (2006.01), *A61P 29/00* (2006.01)
 (71) Demandeur/Applicant:
VERTEX PHARMACEUTICALS INCORPORATED, US
 (72) Inventeur/Inventor:
RANDLE, JOHN C. R., US
 (74) Agent: SMART & BIGGAR

(54) Titre : TRAITEMENT DE MALADIES A L'AIDE D'INHIBITEURS DE L'ICE
 (54) Title: ICE INHIBITORS FOR THE TREATMENT OF AUTOINFLAMMATORY DISEASES



(57) Abrégé/Abstract:

This invention relates to methods and compositions for treating autoinflammatory diseases. The invention also assays for evaluating the ability of an ICE inhibitor to treat autoinflammatory diseases.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 December 2005 (15.12.2005)

PCT

(10) International Publication Number
WO 2005/117846 A2


- (51) International Patent Classification⁷: **A61K 31/00**
- (21) International Application Number: PCT/US2005/018949
- (22) International Filing Date: 27 May 2005 (27.05.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/574,993 27 May 2004 (27.05.2004) US
- (71) Applicant (for all designated States except US): **VERTEX PHARMACEUTICALS INCORPORATED** [US/US];
130 Waverly Street, Cambridge, MA 02139 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **RANDLE, John, C., R.** [CA/US]; 73 Windsor Road, Brookline, MA 02445 (US).
- (74) Agents: **HALEY, James, F., Jr.** et al.; Fish & Neave IP Group, Ropes & Gray LLP, 1251 Avenue of the Americas, New York, NY 10020 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,

[Continued on next page]

(54) Title: TREATMENT OF DISEASES USING ICE INHIBITORS



(57) Abstract: This invention relates to methods and compositions for treating autoinflammatory diseases. The invention also assays for evaluating the ability of an ICE inhibitor to treat autoinflammatory diseases.

WO 2005/117846 A2

WO 2005/117846 A2



CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS,

LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

TREATMENT OF DISEASES USING ICE INHIBITORS

Cross-Reference to Related Application

[0001] This application claims benefit under 35
U.S.C. § 119(e) of United States provisional patent
5 application no. 60/574,993, filed May 27, 2004, the
disclosure of which is herein incorporated by
reference.

Background of the Invention

[0002] Autoinflammatory diseases are a group of
10 related conditions that include Muckle-Wells Syndrome
(MWS), Familial Cold Autoinflammatory Syndrome (FCAS)
(also known as Familial Cold Urticaria or FCU),
Familial Mediterranean Fever (FMF), Chronic Infantile
Neurological Cutaneous and Articular Syndrome (CINCAS),
15 a.k.a. Neonatal Onset Multisystem Inflammatory Disease
(NOMID), TNFR1-Associated Periodic Syndrome (TRAPS),
Hyper-IgD periodic fever Syndrome (HIDS), and Blau's
syndrome (Gumucio et al., Clin. Exp. Rheumatol. 20: S-
45-S-53, 2002).

20 [0003] These conditions have distinct genetic and
phenotypic characteristics, but share certain features
of their clinical presentation, including fever,

- 2 -

urticaria, arthralgia, serositis, systemic amyloidosis as common complication, robust acute phase response and prominent neutrophilia in affected tissues.

[0004] These conditions are caused by mutations in the genes encoding various proteins involved in cellular signaling, notably those involved in regulation of activation of the interleukin-1 β converting enzyme (ICE, a.k.a. caspase-1). ICE is a protease that converts the inactive precursor, pro-interleukin-1 β (pro-IL-1 β) to the active cytokine interleukin-1 β (IL-1 β), as well as pro-IL-18 to mature, active IL-18. Among these modulatory genes & proteins, for example, is the gene *CAIS1*, which codes for the protein cryopyrin. Mutations of this gene and alteration of cryopyrin alters ICE activation, the consequent processing of pro-IL-1 β to IL-1 β and causes MWS and FCAS (Hoffman et al., Nat. Genet. 29: 301-305, 2001).

[0005] The critical role of IL-1 β in the etiology of MWS (and by inference in FCAS, FMF, CINCAS, NOMID, TRAPS, HIDS and other autoinflammatory diseases and conditions) is illustrated by the demonstration that patients with MWS respond to treatment with recombinant human interleukin-1 receptor antagonist (rhuIL-1Ra, anakinra) with a dramatic, rapid and sustained reduction in clinical inflammatory symptoms, reduction of serum amyloid A (an acute phase inflammatory reactant) and urinary excretion of amyloid. This treatment, although effective, is limited in its use because it requires daily injection of the therapeutic agent.

- 3 -

[0006] Profound clinical improvement in response to anakinra may be an indicator that IL-1 is the key inflammatory cytokine underlying disease states, and indeed that such disease states are in fact
5 autoinflammatory syndromes whose underlying genetic basis has yet to be determined. Examples include systemic-onset juvenile idiopathic arthritis (soJIA), also known as Still's disease, and the macrophage activation syndrome. The present invention also
10 encompasses these indications.

[0007] The present invention describes the use of inhibitors of ICE/caspase-1, whether selective for ICE/caspase-1, or broadly active on a range of other caspases (2-14). This treatment, by inhibiting ICE and
15 inhibiting IL-1 β production will reduce the symptoms and other disease conditions of MWS, FCAS, FMF, CINCAS, NOMID, TRAPS, HIDS and other autoinflammatory diseases, conditions or syndromes.

[0008] Although anakinra has been shown to be
20 effective in treating Muckle-Wells syndrome, Familial Cold Autoinflammatory Syndrome, Hyper-IgD syndrome and systemic onset Juvenile idiopathic Arthritis, anakinra is a protein and it therefore does not possess ideal pharmaceutical properties. Therefore, there is a need
25 for small molecule, orally-active ICE inhibitors for treating certain diseases. Such compounds would be extremely useful in treating the aforementioned disease states where ICE plays a role.

Summary of the Invention

30 [0009] The present invention relates to compounds for treating certain diseases associated with autoinflammatory diseases (sometimes referred to as

- 4 -

autoinflammatory fever syndromes), Cryopyrin-associated
Periodic Syndromes, and related syndromes such as
Muckle-Wells Syndrome (MWS), Familial Cold
Autoinflammatory Syndrome (FCAS) (also known as
5 Familial Cold Urticaria or FCU), Familial Mediterranean
Fever (FMF), Chronic Infantile Neurological Cutaneous
and Articular Syndrome (CINCAS), a.k.a. Neonatal Onset
Multisystem Inflammatory Disease (NOMID), TNFR1-
Associated Periodic Syndrome (TRAPS), Hyper-IgD
10 periodic fever Syndrome (HIDS), and Blau's syndrome.
The invention also relates to compounds for treating
certain diseases exhibiting profound clinical
improvement in response to anakinra. The invention
also relates to methods for treating the above
15 diseases.

[0010] The invention also relates to methods for
identifying agents useful for treating these diseases.

Brief Description of the Figures

20 [0011] FIGS. 1A-1C depict the *in vitro* inhibition by
compound 7 of IL-1 β and IL-18 production by whole blood
from patients with FCAS compared with healthy
volunteers. See Example 18.

[0012] FIG. 2 depicts the *in vitro* inhibition by
25 compound 7 of IL-1 β production by whole blood from
patients with MWS. See Example 19.

[0013] FIGS. 3A-3C depict the clinical response of
MWS patients to 14-day treatment with 900 mg compound 7
administered three times daily. In FIG. 3A, serum
30 interleukin-18 was reduced on average by 55% at day 14.
In FIG. 3B, serum acute phase reactant proteins serum
amyloid A (SAA) and C-reactive protein (CRP) was
reduced on average by 75% and 65%, respectively. In

- 5 -

FIG. 3C, self-reported patient total symptoms were reduced on average by 60% during treatment. See Example 21.

5 Detailed Description of the Invention

[0014] This invention provides compounds disclosed herein and pharmaceutically acceptable derivatives thereof that are particularly effective for ameliorating, treating and/or preventing certain
10 diseases associated with autoinflammatory syndromes and related syndromes, including, but not limited to, Muckle-Wells Syndrome (MWS), Familial Cold Autoinflammatory Syndrome (FCAS) (also known as Familial Cold Urticaria or FCU), Familial Mediterranean
15 Fever (FMF), Chronic Infantile Neurological Cutaneous and Articular Syndrome (CINCAS), a.k.a. Neonatal Onset Multisystem Inflammatory Disease (NOMID), TNFR1-Associated Periodic Syndrome (TRAPS), Hyper-IgD periodic fever Syndrome (HIDS), and Blau's syndrome, as
20 well as systemic onset juvenile idiopathic arthritis (also known as Still's disease) and macrophage activation syndrome. Symptoms associated with these diseases include, but are not limited to fever,
25 urticaria, arthritis, serositis, systemic amyloidosis, a robust acute phase response, and a prominent neutrophilia in affected tissues.

[0015] The compounds according to this invention may be used to treat these disease states in mammals. Any compound that inhibits ICE may be used in the methods
30 and compositions of this invention. Such compounds include those compounds that inhibit ICE selectively and those that inhibit one or more enzyme in the caspase or ICE/CED-3 family.

- 6 -

[0016] The compounds of this invention inhibit ICE and/or decrease IL-1, particularly IL-1 β and IL-18 levels. These compounds can be assayed, for example, for their ability to inhibit ICE, the production of
5 IL-1 β and/or IL-18, the regulation of IL-1 and/or IL-18 levels, and/or affect IL-1 β and/or IL-18 activity. Assays for testing each of these activities are known in the art (see Examples herein, WO 95/35308, WO 97/22619, WO 99/47545, or WO 01/90063). Accordingly,
10 these compounds are capable of targeting and inhibiting events in the ICE and/or IL-1 mediated diseases set forth herein.

[0017] Compounds that may be used in connection with this invention include, but are not limited to, the
15 compounds of the following documents: WO 04/058718, WO 04/002961, WO 03/088917, WO 03/068242, WO 03/042169, WO 98/16505, WO 93/09135, WO 03/106460, WO 03/103677, WO 03/104231, WO 02/085899, WO 00/55114, WO 00/55127, WO 00/61542, WO 01/05772, WO 01/10383, WO 01/16093,
20 WO 01/42216, WO 01/72707, WO 01/90070, WO 01/94351, WO 02/094263, WO 02/42278, US Patent 6,184,210, US Patent 6,184,244, US Patent 6,187,771, US Patent 6,197,750, US Patent 6,242,422, April 2001 American Chemical Society (ACS) meeting in San Diego,
25 California, USA, WO 02/22611, US 2002/0058630, WO 02/12638, WO 95/35308, US Patent 5,716,929, WO 97/22619, US Patent 6,204,261, WO 99/47545, WO 01/90063, US Patent Publication 2004/0014753, US Patent Publication 2004/0009966, US Patent Publication
30 2003/0236296, US Patent Publication 2003/0096737, US Patent Publication 2003/0092703, US Patent Publication 2002/0169177, US Patent 6,693,096, US Patent 6,610,683,

- 7 -

US Patent 6,531,467, US Patent 6,528,506, US Patent 6,200,969, WO 2003/072528, WO 2003/032918, WO 01/00658, WO 98/10778, US Patent 6,716,818, US Patent 6,620,782, US Patent 6,566,338, US Patent 6,495,522, US Patent 6,355,618, 6,153,591, WO 2005/003100, WO 2004/002401, WO 00/61542, WO 00/55114, WO 99/47154, US Patent 6,083,981, US Patent 5,932,549, US Patent 5,919,790, US Patent 5,744,451, WO 2002/089749, WO 99/36426, WO 98/16505, WO 98/16504, WO 98/16502, US Patent 6,316,415, US Patent 5,932,549, US Patent 5,919,790, US Patent 5,744,451, EP 1082127, EP 1049703, EP 0932600, EP 0932598, WO 99/56765, WO 93/05071, EP 0600880, and EP 1378573 (which, as set forth herein, are all incorporated by reference herein). In one embodiment, compounds for use in this invention include those of WO 00/55114, WO 00/55127, WO 00/61542, WO 00/61542, WO 01/05772, WO 01/10383, WO 01/16093, WO 01/42216, WO 01/72707, WO 01/90070, WO 01/94351, US Publication 2003/0092703, WO 02/094263, US Publication 2002/0169177, US Patent 6,184,210, US Patent 6,184,244, US Patent 6,187,771, US Patent 6,197,750, US Patent 6,242,422, April 2001 American Chemical Society (ACS) meeting in San Diego, California, USA, WO 02/22611, US Publication 2002/0058630, US Publication 2003/0096737, WO 95/35308, WO 97/22619, WO 99/47545, and WO 01/90063. In another embodiment, compounds for use in this invention include those of WO 04/058718, WO 04/002961, WO 95/35308, WO 97/22619, WO 99/47545, and WO 01/90063. Alternately, compounds for use in this invention include those of WO 95/35308, WO 97/22619, WO 99/47545, and WO 01/90063. Preferred compounds are those recited in the claims herein. These compounds may be obtained

- 8 -

by methods known to skilled practitioners and the methods disclosed in documents cited herein.

[0018] The pharmaceutical compositions and methods of this invention, therefore, will be useful for
5 controlling the levels of IL-1 β , IL-18 and other biomarkers and/or activity in vitro or in vivo. The compositions and methods of this invention will thus be useful for controlling the levels of IL-1 β , IL-18 and other biomarkers in vivo and for treating or reducing
10 the advancement, severity or effects of certain conditions, including diseases, disorders, or effects as set forth herein.

[0019] According to another embodiment, the invention provides a composition comprising a compound
15 of this invention or a pharmaceutically acceptable derivative (e.g., salt) thereof, as described above, and a pharmaceutically acceptable carrier.

[0020] According to another embodiment, the compositions of this invention may further comprise
20 another therapeutic agent. Such agents include, but are not limited to, a thrombolytic agent such as tissue plasminogen activator and streptokinase, an anti-inflammatory agent, a matrix metalloprotease inhibitor, a lipoxigenase inhibitor, a cytokine antagonist,
25 (including, but not limited to, an antibody or binding protein against TNF, IL-1, IL-6, IL-12, IL-18 or their receptors), a chemokine antagonist, an immunosuppressant, an anti-cancer agent, an anti-viral agent, a cytokine (including, but not limited to, the
30 interleukin-1 receptor antagonist or anakinra), a growth factor, an immunomodulator (e.g., bropirimine, anti-human alpha interferon antibody, IL-2, GM-CSF, methionine enkephalin, interferon alpha,

- 9 -

diethyldithiocarbamate, tumor necrosis factor, naltrexone and rEPO), a prostaglandin, or an anti-vascular hyperproliferation compound.

[0021] The term "pharmaceutically acceptable carrier" refers to a non-toxic carrier that may be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof.

[0022] Pharmaceutically acceptable carriers that may be used in these compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0023] In pharmaceutical compositions comprising only a compound of this invention as the active component, methods for administering these compositions may additionally comprise the step of administering to the subject an additional agent. Such agents include, but are not limited to, a thrombolytic agent such as tissue plasminogen activator and streptokinase, an anti-inflammatory agent, a matrix metalloprotease inhibitor, a lipoxigenase inhibitor, a cytokine antagonist (including, but not limited to, an antibody

- 10 -

or binding protein against TNF, IL-1, IL-6, IL-12, IL-18 or their receptors), a chemokine antagonist, an immunosuppressant, an anti-cancer agent, an anti-viral agent, a cytokine (including, but not limited to, the interleukin-1 receptor antagonist or anakinra), a growth factor, an immunomodulator (e.g., bropirimine, anti-human alpha interferon antibody, IL-2, GM-CSF, methionine enkephalin, interferon alpha, diethyldithiocarbamate, tumor necrosis factor, naltrexone and rEPO), a prostaglandin, or an anti-vascular hyperproliferation compound. When a second agent is used, the second agent may be administered either as a separate dosage form or as part of a single dosage form with the compounds or compositions of this invention.

[0024] The amount of compound present in the above-described compositions should be sufficient to cause a detectable decrease in the severity of the disease, or in ICE inhibition, the levels of IL-1 β , IL-18 or other biomarkers, or the activity of IL-1 β , IL-18 or other biomarkers.

[0025] If pharmaceutically acceptable salts of the compounds of this invention are utilized in these compositions, those salts are preferably derived from inorganic or organic acids and bases. Included among such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzene sulfonate, bisulfate, butyrate, citrate, camphorate, camphor sulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate,

- 11 -

maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenyl-propionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and
5 undecanoate. Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, and
10 salts with amino acids such as arginine, lysine, and so forth.

[0026] Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl
15 chlorides, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; aralkyl
halides, such as benzyl and phenethyl bromides and
20 others. Water or oil-soluble or dispersible products are thereby obtained.

[0027] The compounds utilized in the compositions and methods of this invention may also be modified by appending appropriate functionalities to enhance
25 selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system (e.g., blood, lymphatic system, or central nervous system), increase oral availability, increase
30 solubility to allow administration by injection, alter metabolism and/or alter rate of excretion.

[0028] According to a preferred embodiment, the compositions of this invention are formulated for

- 12 -

pharmaceutical administration to a subject or patient, e.g., a mammal, preferably a human being. Such pharmaceutical compositions are used to ameliorate, treat or prevent autoinflammatory diseases in a subject and comprise a compound that inhibits ICE and a pharmaceutically acceptable carrier.

[0029] Such pharmaceutical compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection and infusion techniques. Preferably, the compositions are administered orally or intravenously.

[0030] Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride

- 13 -

derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil and castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[0031] If a solid carrier is used, the preparation can be in tablet form, placed in a hard gelatin capsule in powder or pellet form, or in the form of a troche or lozenge. The amount of solid carrier will vary, e.g., from about 25 to 800 mg, preferably about 25 mg to 400 mg. When a liquid carrier is used, the preparation can be, e.g., in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension. Where the composition is in the form of a capsule, any routine encapsulation is suitable, for example, using the aforementioned carriers in a hard gelatin capsule shell.

[0032] A syrup formulation can consist of a suspension or solution of the compound in a liquid carrier for example, ethanol, glycerin, or water with a flavoring or coloring agent. An aerosol preparation can consist of a solution or suspension of the compound

- 14 -

in a liquid carrier such as water, ethanol or glycerin; whereas in a powder dry aerosol, the preparation can include e.g., a wetting agent.

[0033] Formulations of the present invention
5 comprise an active ingredient together with one or more acceptable carrier(s) thereof and optionally any other therapeutic ingredient(s). The carrier(s) should be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not
10 deleterious to the recipient thereof.

[0034] The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, and aqueous suspensions or
15 solutions. In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents
20 include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[0035] Alternatively, the pharmaceutical
25 compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room
30 temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

- 15 -

[0036] The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[0037] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[0038] For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetaryl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0039] For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative

- 16 -

such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

[0040] The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents known in the art.

[0041] It will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration, and other well-known variables.

[0042] The above-described compounds and compositions are also useful in therapeutic applications relating to certain diseases.

[0043] Such certain diseases include autoinflammatory syndromes such as cryopyrin-associated Periodic Syndromes, (including Muckle-Wells syndrome, Familial Cold Autoinflammatory Syndrome (FCAS, also known as Familial Cold Urticaria or FCU), chronic infantile neurological cutaneous and articular syndrome (a.k.a. neonatal onset multisystem inflammatory disease)), familial mediterranean fever, TNFR1-Associated Periodic Syndrome (TRAPS), Hyper-IgD periodic fever Syndrome (HIDS), and Blau's syndrome. Other autoinflammatory syndromes whose underlying

- 17 -

genetic basis has yet to be determined, may include systemic-onset juvenile idiopathic arthritis (soJIA), also known as Still's disease, and the macrophage activation syndrome.

5 [0044] The compounds of this invention can inhibit the release of IL-1, particularly IL-1 β and IL-18, and thus can be useful for inhibiting or blocking several pathophysiological effects of certain diseases as set forth herein.

10 [0045] This invention also relates to a therapeutic method for treating certain diseases, specifically autoinflammatory diseases, by (1) inhibiting IL-1, particularly IL-1 β and IL-18 release from cells and/or
15 (2) preventing the untoward, toxic or lethal effects of excessively high tissue levels of IL-1, particularly IL-1 β and IL-18 in a mammal, including a human. This method comprises administering to a mammal an effective ICE inhibiting quantity of one or more ICE/CED-3 inhibitors. This method also can be used for the
20 prophylactic treatment or prevention of certain diseases amenable thereto, including Muckle-Wells Syndrome (MWS), Familial autoinflammatory fever syndromes and related syndromes such as Cryopyrin-associated Periodic Syndromes, Cold Autoinflammatory
25 Syndrome (FCAS, also known as Familial Cold Urticaria or FCU), Familial Mediterranean Fever (FMF), Chronic Infantile Neurological Cutaneous and Articular Syndrome (CINCAS), a.k.a. Neonatal Onset Multisystem Inflammatory Disease (NOMID), TNFR1-Associated Periodic
30 Syndrome (TRAPS), Hyper-IgD periodic fever Syndrome (HIDS), Blau's syndrome, systemic onset juvenile idiopathic arthritis, Still's disease and macrophage

- 18 -

activation syndrome. The invention provides a method for the treating these disorders by administering to a mammal, including a human, in need thereof an effective amount of such compounds.

5 [0046] The compounds, by inhibiting ICE and blocking the release of IL-1, particularly IL-1 β and IL-18 or decreasing IL-1, particularly IL-1 β and IL-18 levels and activity, as well as the pathophysiologic actions of excessive levels of IL-1, particularly IL-1 β and IL-18
10 in each of these circumstances, directly facilitate the arrest or resolution of certain diseases, and facilitates the restoration of normal function. Together, these actions relate their novel use in treating certain diseases.

15 [0047] ICE inhibition may be measured by methods known in the art and as described more fully herein.

[0048] The phrase "inhibiting the release of IL-1, particularly IL-1 β , and IL-18" means:

a) a decrease of in vivo IL-1 β and/or IL-18 levels in a
20 mammal such as a human;

b) a down regulation of IL-1, particularly IL-1 β , and IL-18 levels in vitro or in-vivo; or

c) a down regulation of IL-1, particularly IL-1 β , and IL-18 activity, by direct inhibition of the synthesis
25 of IL-1 β and IL-18 or a post-translation event in vivo or in vitro as well as indirect inhibition of IL-1 α .

[0049] The compounds may be useful in inhibiting the release of IL-1, particularly IL-1 β , and IL-18 release by monocytes, macrophages, neuronal cells, endothelial
30 cells, epidermal cells, mesenchymal cells (for example: fibroblasts, skeletal myocytes, smooth muscle myocytes, cardiac myocytes) and many other types of cells.

- 19 -

[0050] The term "condition" or "state" refers to any disease, disorder, or effect that produces deleterious biological consequences in a subject.

[0051] The level of IL-1, particularly IL-1 β , and
5 IL-18 protein in the blood or cell of a patient or a cell culture (i.e., within the cell or the cell culture media) can be determined by for example, assaying for immunospecific binding to IL-1 β or IL-18 or to other proteins known to be produced as a result of the
10 presence of active IL-1 β and IL-18, including, but not limited to, IL-1 α . Such methods are known in the art. For example, immunoassays which can be used include, but are not limited to competitive and non-competitive assay systems, western blots, radioimmunoassays, ELISA
15 (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays,
20 fluorescent immunoassays, protein A immunoassays and FACS analysis with labeled antibodies. Such assays well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is
25 incorporated by reference herein in its entirety).

[0052] Competitive binding assays can also be used to determine the level of IL-1, particularly IL-1 β , and IL-18. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled
30 proteins from cells expressing IL-1, particularly IL-1 β , and IL-18 (e.g., ^3H or ^{125}I) with an IL-1 β or IL-18 antibody in the presence of increasing amounts of

- 20 -

unlabeled IL-1 β or IL-18, and the detection of the IL-1 β or IL-18 antibody bound to the labeled IL-1 β or IL-18. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by Scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ^3H or ^{125}I) in the presence of increasing amounts of an unlabeled second antibody.

[0053] IL-1 β or IL-18 levels can also be assayed by activity, for example, IL-1 levels can be assayed by a cell line that is capable of detecting bioactive levels of cytokines like IL-1 β or IL-18 or a growth factor. According to one embodiment, the level of bioactive IL-1 β in a biological sample is detected by incubating a cell line genetically engineered with isopropyl- β -D-thiogalactopyranoside. The cell line is incubated with the sample to be tested and cell death in the cell line is monitored by determining the intensity of blue color, which is indicative of a bioactive cytokine or growth factor in the sample tested. See also, e.g., Burns (1994) 20(1):40-44 for IL-1 activity assay of serum of patients.

[0054] Dosage levels of between about 0.01 and about 100 mg/kg body weight per day, preferably between about 0.5 and about 75 mg/kg body weight per day and most preferably between about 1 and about 50 mg/kg body weight per day of the active ingredient compound are useful in a monotherapy.

- 21 -

[0055] In another embodiment, dosage levels of between about 2 and about 200 mg/kg body weight per day, preferably between about 6 and about 100 mg/kg body weight per day and most preferably between about 5 25 and about 100 mg/kg body weight per day of the active ingredient compound are useful in a monotherapy.

[0056] According to one embodiment of this invention, an active ingredient compound of this invention is administered to a subject at a dose of 10 between about 20 mg to about 10,000 mg per administration. In another embodiment, an active ingredient compound of this invention is administered to a subject at a dose of between about 300 mg to about 2,400 mg per administration. In yet another embodiment 15 of this invention, an active ingredient compound of this invention is administered to a subject at a dose of between about 600 mg to about 1,800 mg per administration. In a preferred embodiment, a compound of this invention is administered to a subject at a 20 dose of about 900 mg per administration.

[0057] Typically, the pharmaceutical compositions of this invention will be administered from about 1 to 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic 25 or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 30 95% active compound (w/w). Preferably, such preparations contain from about 20% to about 80% active compound.

- 22 -

[0058] When the compositions of this invention comprise a combination of a compound of this invention and one or more additional therapeutic agents, both the compound and the additional agent should be present at dosage levels of between about 10% to about 80% of the dosage normally administered in a monotherapy regime.

[0059] Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained. When the symptoms have been alleviated to the desired level, treatment may cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence or disease symptoms.

[0060] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of active ingredients will also depend upon the particular compound and other therapeutic agent, if present, in the composition.

[0061] Accordingly, a method for ameliorating, treating or preventing a disease of this invention in a subject comprises the step of administering to the subject any compound, pharmaceutical composition, or combination described herein.

- 23 -

[0062] In a preferred embodiment, the invention provides a method of treating a mammal, having one of the aforementioned diseases, comprising the step of administering to said mammal a pharmaceutically acceptable composition described above. In this embodiment, if the patient is also administered another therapeutic agent, it may be delivered together with the compound of this invention in a single dosage form, or, as a separate dosage form. When administered as a separate dosage form, the other therapeutic agent may be administered prior to, at the same time as, or following administration of a pharmaceutically acceptable composition comprising a compound of this invention.

[0063] The methods for identifying a compound or composition for treating a disease according to this invention include methods for screening of a plurality of compounds or compositions for their ability to ameliorate the effects of certain disease(s) and/or improve the condition of a patient having certain disease(s) of this invention. According to one embodiment of this invention, high throughput screening can be achieved by having cells in culture in a plurality of wells in a microtiter plate, adding a different compound or composition to each well and comparing the ICE inhibition and/or IL-1 levels and/or activity in each cell culture to the levels or activity present in a cell culture in a control well. Controls that are useful for the comparison step according to this invention include cells or subjects that have not been treated with a compound or composition and cells or subjects have been treated with a compound or composition that is known to have no effect on ICE

- 24 -

inhibition or activity. According to one embodiment of this invention, the high throughput screening is automated so that the steps including the addition of the cells to the plate up to the data collection and analysis after addition of the compound or composition are done by machine. Instruments that are useful in the comparison step of this invention, e.g., instruments that can detect labeled objects (e.g., radiolabelled, fluorescent or colored objects) or objects that are themselves detectable, are commercially available and/or known in the art. Accordingly, compounds and compositions according to this invention that are useful for treating the certain disease disclosed herein can be quickly and efficiently screened.

[0064] In another embodiment of this invention, a method for identifying a compound or composition for ameliorating, treating or preventing an autoinflammatory disease in a subject comprises administering to said subject a compound that inhibits ICE or a pharmaceutical composition comprising the compound and comparing the ICE inhibition in the subject before and after treatment with the compound. In an alternate embodiment, this invention provides a method for identifying a compound or composition for ameliorating, treating or preventing an autoinflammatory disease in a subject which comprises administering to said subject a compound that inhibits ICE or a pharmaceutical composition comprising the compound and comparing a biomarker for autoinflammatory diseases in said subject before and after treatment with said compound.

- 25 -

[0065] The term "biomarker" is a physical, functional, or biochemical indicator, e.g., the presence of a particular metabolite, of a physiological or disease process. According to this invention, certain

5 biomarkers of inflammation may be used to evaluate the response of patients having autoinflammatory diseases listed herein to compounds that inhibit ICE. These inflammatory biomarkers include, but are not limited to, interleukin-1, interleukin-6 (all autoinflammatory

10 diseases), interleukin-8 (all autoinflammatory diseases), interleukin-18, serum amyloid A, C-reactive protein, erythrocyte sedimentation rate (ESR), haptoglobin, TNFalpha, immunoglobins (particularly IgD), ferritin, leukocyte counts, platelet counts and

15 hemoglobin.

[0066] Accordingly, this invention also provides a method for reduction of IL-1 β , IL-18 or other biomarker in a subject having an autoinflammatory disease comprising the step of administering to said subject a

20 compound that inhibits ICE or a pharmaceutical composition comprising said compound. In one embodiment, the plasma concentration of the compound or composition in the subject is about equal to or greater than an IC₅₀ value taught according to this invention

25 (see, e.g., Examples 18 and 19). In another embodiment, the plasma concentration of the compound or composition in the subject is between about 0.2 μ M to about 50 μ M, or about 0.8 μ M to about 6 μ M. In yet another embodiment, the reduction of IL-1 β , IL-18 or

30 other biomarker in a subject is measured by comparing (a) IL-1 β , IL-18 or other biomarker concentrations in said subject before or after treatment with said

- 26 -


compound or composition to (b) IL-1 β , IL-18 or other biomarker concentrations in said subject during treatment with said compound or composition. According to this invention, the percent reduction of IL-1 β ,
 5 IL-18 or other biomarker in a subject is selected from the group consisting of (a) at least about 50% to about 100% reduction; (b) at least about 50% to about 90% reduction; and (c) at least about 60% to about 90% reduction.

10 **[0067]** All applications, patents and references disclosed herein are incorporated by reference. In order that this invention be more fully understood, the following preparative and testing examples are set forth. These examples are for the purpose of
 15 illustration only and are not to be construed as limiting the scope of the invention in any way.


Examples

Example 1

20 [3S/R, (2S)]-5-Fluoro-4-oxo-3-{[1-(phenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-pentanoic acid




Method A: (S)-(1-Phenothiazine-10-carbonyl)piperidine-2-carboxylic acid methyl ester



- 27 -

[0068] To a stirred solution of methyl pipercolate hydrochloride (1g, 5.57 mmol) in THF (10 ml) was added phenothiazine carbonyl chloride (1.457g, 5.57 mmol) followed by diisopropylethylamine (2.02 ml, 11.68 mmol). The resulting solution was stirred for 16h before being partitioned between ethyl acetate and aq. sat. NH₄Cl. The organic layer was washed with brine, dried (MgSO₄), filtered and evaporated. The residue was purified by flash chromatography (15% ethyl acetate in hexane) to afford the sub-title compound as a colorless oil which crystallized upon standing (1.823g, 89%): ¹H NMR (400MHz, CDCl₃) δ 1.13-1.48 (3H, m), 2.57-2.69 (2H, m), 2.16 (1H, m), 3.00 (1H, m), 3.74 (4H, s+m), 5.00 (1H, m), 7.11 (2H, t), 7.22-7.34 (4H, m), 7.76 (2H, d); ¹³C NMR (100MHz, CDCl₃) δ 21.3 (CH₂), 24.8 (CH₂), 27.3 (CH₂), 44.9 (CH₂), 52.5 (CH₃), 55.9 (CH), 122.8 (CH), 125.5 (CH), 127.8 (CH), 128.0 (CH), 129.2 (C), 141.7 (C), 158.4 (C), 172.2 (C).

20 Method B: (S)-(1-Phenothiazine-10-carbonyl)piperidine-2-carboxylic acid




[0069] To a stirred solution of (S)-(1-phenothiazine-10-carbonyl)piperidine-2-carboxylic acid methyl ester (0.912g) in THF (15 ml) and water (8 ml) was added 2M NaOH (3.71 ml) and the reaction mixture was stirred for 16 hours. The reaction mixture was poured into sodium hydrogen carbonate solution (50 ml) and extracted with ethyl acetate (40 ml). Aqueous

- 28 -

phase was made acidic and extracted with ethyl acetate (2x75 ml). Organic extracts were combined, dried (MgSO₄) and concentrated to give the sub-title compound as a white solid (0.709g, 81%): ¹H NMR (400MHz, CDCl₃) δ 0.99-1.72 (5H, m), 2.23 (1H, m), 2.97 (1H, m), 3.58 (1H, m), 4.93 (1H, m), 7.16 (2H, t), 7.28 (2H, t), 7.37 (2H, d), 7.78 (2H, d); ¹³C NMR (100MHz, CDCl₃) δ 21.0 (CH₂), 24.2 (CH₂), 26.7 (CH₂), 45.7 (CH₂), 56.0 (CH), 123.8 (CH), 126.0 (CH), 127.9 (CH), 128.1 (CH), 130.3 (C), 141.2 (C), 160.1 (C), 175.9 (C).

Method C: [3S/R, 4S/R (2S)]-5-Fluoro-4-hydroxy-3-{[1-(phenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-pentanoic acid *tert*-butyl ester




15 [0070] A stirred mixture of (S)-(1-phenothiazine-10-carbonyl)piperidine-2-carboxylic acid (233mg, 0.658mmol), 3-amino-5-fluoro-4-hydroxy-pentanoic acid *tert*-butyl ester (150mg, .724mmol), HOBt (98mg, 0.724mmol), DMAP (88mg, 0.724mmol) and anhydrous THF (10ml) was cooled to 0°C before EDC (139mg, 0.724mmol) was added. The mixture was allowed to warm to room temperature during 16h then concentrated under reduced pressure. The residue was purified by flash chromatography (50% ethyl acetate in hexane) to afford the sub-title compound as a pale pink foam (294mg, 82%): ¹H NMR (400MHz, CDCl₃) δ 1.96 (1H, m), 1.18-1.60 (13H, m), 2.10-2.25 (1H, m), 2.48-2.70 (2H, m), 2.78-2.94 (1H, m), 3.51-4.72 (7H, m), 7.03-7.36 (7H, m),

- 29 -

7.71-7.76 (2H, m); ^{19}F (376MHz, CDCl_3) δ -228.9 (t), -229.3 (t), -230.1 (t), -230.2 (t).

Method D: [3S/R, (2S)]-5-Fluoro-4-oxo-3-{ [1-
5 (phenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-pentanoic acid tert-butyl ester




[0071] A stirred solution of [3S/R, 4S/R (2S)]-5-
Fluoro-4-hydroxy-3-{ [1-(phenothiazine-10-
10 carbonyl)piperidine-2-carbonyl]amino}-pentanoic acid
tert-butyl ester (294mg, 0.541mmol) in anhydrous DCM
(10mL) was treated with 1,1,1-triacetoxy-1,1-dihydro-
1,2-benziodoxol-3(1H)-one (344mg, 0.812mmol) at 0°C.
The resulting mixture was allowed to warm to room
15 temperature over 2h, diluted with ethyl acetate, then
poured into a 1:1 mixture of saturated aqueous sodium
hydrogen carbonate and saturated aqueous sodium
thiosulphate. The organic layer was removed and the
aqueous layer was re-extracted with ethyl acetate. The
20 combined organic extracts were dried (Na_2SO_4) and
concentrated. The residue was purified by flash
chromatography (30% ethyl acetate in hexane) to afford
the sub-title compound as a pale pink foam (220mg,
75%): ^1H NMR (400MHz, CDCl_3) δ 0.84-0.96 (1H, m), 1.20-
25 1.40 (10H, m+2s), 1.51-1.56 (3H, m), 2.20-2.27 (1H, m),
2.70-2.98 (3H, m), 3.49-3.63 (1H, m), 4.74-5.24 (4H,
m), 7.14-7.18 (2H, m), 7.28-7.38 (4H, m), 7.48-7.79
(3H, m); ^{13}C (100MHz, CDCl_3) δ 20.8/21.0 (CH_2), 23.7/23.9
(CH_2), 25.8/25.9 (CH_2), 28.2/28.3 (CH_3), 36.8/36.9 (CH_2),

- 30 -

46.0/46.1 (CH₂), 52.9 (CH), 56.8 (CH), 82.6 (C),
 84.4/84.5 (2d, J 184.0/183.3, CH₂F), 123.7/123.8 (CH),
 126.1 (CH), 128.0/128.1 (CH), 128.2/128.3 (CH),
 130.4/130.5 (C), 141.4 (C), 160.0 (C), 170.0 (C), 171.7
 5 (C), 202.9 (C); ¹⁹F (376MHz, CDCl₃) δ-231.9 (t), -232.2
 (t).

Method E: [3S/R, (2S)]-5-Fluoro-4-oxo-3-{[1-
 (phenothiazine-10-carbonyl)piperidine-2-
 10 carbonyl]amino}-pentanoic acid




[0072] Trifluoroacetic acid (5mL) was added to a
 stirred ice cold solution of [3S/R, (2S)]-5-fluoro-4-
 oxo-3-{[1-(phenothiazine-10-carbonyl)piperidine-2-
 15 carbonyl]amino}-pentanoic acid tert-butyl ester (130mg,
 0.24mmol) in anhydrous DCM (5mL). The mixture was
 stirred at 0°C for 0.5h then at room temperature for
 0.5h. The mixture was concentrated under reduced
 pressure and then the residue was dissolved in dry DCM.
 20 This process was repeated several times in order to
 remove excess trifluoroacetic acid. The gum was
 lyophilized twice from HPLC grade water to afford the
 title compound as a white powder (77mg, 66%): IR
 (solid) 1670, 1716, 1782 cm⁻¹; ¹H NMR (400MHz, d₆-DMSO) δ
 25 0.96-0.99 (1H, m), 1.23-1.26 (2H, m), 1.42-1.44 (1H,
 m), 1.60 (1H, m), 1.91-1.98 (1H, m), 2.51-2.89 (2H, m),
 3.11-3.22 (1H, m), 3.57-3.60 (1H, m), 4.30-4.72 and
 5.05-5.29 (4H, 2m), 7.11-7.17 (2H, m), 7.24-7.30 (2H,
 m), 7.34-7.38 (2H, m), 7.57-7.63 (2H, m), 8.07-8.61

- 31 -

(1H, m); ^{13}C NMR (100MHz, DMSO) δ (DMSO+TFA) 18.8/18.9, (CH₂), 22.2/22.3 (CH₂), 25.8/26.1 (CH₂), 31.5/33.2 (CH₂), 43.2 (CH₂), 50.6/51.1 (CH), 54.4/54.5 (CH), 82.8/82.9 (2d, J 178.6/178.1, CH₂F), 119.9/120.0 (CH), 120.4/120.5 (CH), 124.0/124.1 m(CH), 125.9/126.0 (C), 126.4/126.5 (CH), 139.6/139.7 (C), 156.0/156.4 (CO), 170.3 (CO), 170.7/170.8 (CO), 202.2/202.3 (2d, J 14.6/15.1, CO).; ^{19}F (376MHz, DMSO) δ chemical shift (multiplicity, relative intensity) -226.7 (t, 3), -226.9 (t, 3), -230.4 (t, 1), -231.2 (t, 1), -232.7 (t, 10), -233.0 (t, 10).

Example 2

[3S/R, (2S)]-3-{[1-(2-Chlorophenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-4-fluoro-4-oxo-pentanoic acid




[0073] This was prepared from 2-chlorophenothiazine carbonyl chloride using procedures similar to those described above in Methods A-E (73mg, 69%): IR (solid, cm⁻¹) 1738, 1660, 1555, 1363, 1224 ; ^1H NMR (400MHz, d₆-DMSO+TFA) δ 0.98-1.61 (4H, m), 1.94-2.03 (1H, m), 2.53-2.89 (2H, m), 3.12-3.24 (1H, m), 3.51-3.61 (1H, m), 4.31-4.73 and 5.10-5.24 (4H, 2m), 7.15-7.49 (6H, m), 7.77-7.81 (1H, m), 8.13-8.64 (1H, m); ^{13}C NMR (100MHz, DMSO+TFA) δ 18.7/18.8 (CH₂), 22.3/22.6 (CH₂), 25.9/26.2 (CH₂), 31.5/33.2 (CH₂), 43.0/43.2 (CH₂), 50.6/51.1 (CH), 54.4/54.5 (CH), 82.8/82.9 (2d, J 178.7/178.3, CH₂F), 119.3/119.8 (CH), 120.2/120.3 (CH), 123.6/123.7 (CH),

- 32 -

124.4/124.5 (CH), 124.6/124.8 (C), 126.6 (CH), 126.9 (CH), 127.5 (CH), 131.0 (C), 139.2/139.2 (C), 140.7/140.7 (C), 155.5/155.9 (C), 170.1/170.2 (C), 170.7/170.8 (C), 201.2/201.3 (2d, J 14.3/13.9, CO); ¹⁹F NMR (376MHz, DMSO+TFA) δ -226.7 (t), -226.9 (t), -230.3 (t), -232.7 (t), -233.0 (t).

Example 3

[3S/R, (2S)]-3-{[1-(3-Chlorophenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-4-fluoro-4-oxo-
pentanoic acid




[0074] This was prepared from 3-chlorophenothiazine carbonyl chloride using procedures similar to those described above in Methods A-E (108mg, 65%): IR (solid, cm⁻¹) 1737, 1655, 1455, 1373, 1224 ; ¹H NMR (400MHz, d₆-DMSO+TFA) δ 0.99-1.61 (5H, m), 1.91-2.04 (1H, m), 2.54-2.90 (2H, m), 3.12-3.24 (1H, m), 3.48-3.60 (1H, m), 4.26-5.28 (4H, m), 7.15-7.68 (7H, m), 8.10-8.62 (1H, m); ¹³C NMR (100MHz, DMSO+TFA) δ 18.8 (CH₂), 22.2/22.3 (CH₂), 25.8 (CH₂), 33.1/33.2 (CH₂), 43.2 (CH₂), 50.6/51.0 (CH), 54.3/54.4 (CH), 82.7/82.8 (2d, CH₂F), 120.2/120.3 (CH), 121.3/121.4 (CH), 124.2/124.3 (CH), 124.8/125.0 (C), 125.7 (CH), 126.3 (CH), 126.6 (CH), 126.8 (CH), 127.7/127.9 (C), 127.9/128.0 (C), 138.5 (C), 139.3 (C), 156.0 (CO), 170.1 (CO), 170.6/170.7 (CO), 201.1/201.2 (2d, CO); ¹⁹F NMR (376MHz, DMSO+TFA) δ -226.6 (t), -226.9 (t), -232.6 (t), -232.9 (t).

- 33 -

Example 4

[3S/R, (2S)]-3-{[1-(3,4-Dichlorophenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-4-fluoro-4-oxo-pentanoic acid



5


[0075] This was prepared from 3,4-dichlorophenothiazine carbonyl chloride using procedures similar to those described above in Methods A-E (91mg, 66%): IR (solid, cm^{-1}) 1737, 1439, 1363, 1219; ^1H NMR (400MHz, $\text{d}_6\text{-DMSO+TFA}$) δ 1.03-1.62 (5H, m), 1.97-2.06 (1H, m), 2.54-2.86 (2H, m), 3.14-3.28 (1H, m), 3.59-3.66 (1H, m), 4.30-5.26 (4H, m), 7.15-7.68 (6H, m), 8.14-8.96 (1H, m); ^{13}C NMR (100MHz, DMSO+TFA) δ 20.2 (CH_2), 23.8 (CH_2), 27.3 (CH_2), 34.6/34.7 (CH_2), 44.5 (CH_2), 52.1/52.5 (CH), 55.7/55.9 (CH), 84.2/84.3 (2d, CH_2F), 120.2/120.3 (CH), 120.8/120.9 (CH), 124.2/124.4 (C), 125.9 (CH), 127.7/127.8 (C), 128.2 (CH), 128.4/128.5 (C), 128.8 (CH), 128.9 (CH), 140.0 (C), 140.1 (C), 140.6 (C), 156.8/156.8 (CO), 171.5 (CO), 172.1/172.1 (CO), 202.6/202.7 (2d, CO); ^{19}F NMR (376MHz, DMSO+TFA) δ -226.6 (t), -226.8 (t), -232.6 (t), -232.9 (t).

25

Example 5

[3S/R, (2S)]-3-{[1-(2,6-Dichlorophenothiazine-10-carbonyl)piperidine-2-carbonyl]amino}-4-fluoro-4-oxo-pentanoic acid


- 34 -



[0076] This was prepared from 2.7 g dichlorophenothiazine carbonyl chloride using procedures similar to those described above in Methods
 5 A-E (91mg, 70%): IR (solid, cm^{-1}) 1737, 1660, 1555, 1363, 1224; ^1H NMR (400MHz, d_6 -DMSO+TFA) δ 1.02-1.62 (5H, m), 1.91-2.02 (1H, m), 2.53-2.90 (2H, m), 3.13-3.25 (1H, m), 3.51-3.62 (1H, m), 4.31-5.29 (4H, m), 7.22-7.75 (6H, m), 8.18-8.65 (1H, m); ^{13}C NMR (100MHz,
 10 DMSO+TFA) δ 20.2 (CH_2), 23.8 (CH_2), 27.3 (CH_2), 34.6 (CH_2), 44.7 (CH_2), 52.5 (CH), 55.8 (CH), 84.3 (d, J 178.2, CH_2F), 120.7/121.2 (CH), 122.7/122.8 (CH), 124.7/125.1 (C), 125.3/125.4 (CH), 127.4 (CH), 128.1 (CH), 128.7/128.9 (C), 129.1 (CH), 129.8 (C), 132.7
 15 (C), 139.5/139.6 (C), 141.8/141.9 (C), 157.0 (CO), 171.5 (CO), 172.1 (CO), 202.6 (d, J 14.3, CO); ^{19}F NMR (376MHz, DMSO+TFA) δ -226.6 (t), -226.9 (t), -232.6 (t), -232.9 (t).

Example 6

20 [3S/R, (2S)]-3-{[1-(Carbazole-9-carbonyl)piperidine-2-carbonyl]amino}-4-fluoro-4-oxo-pentanoic acid




- 35 -

[0077] This was prepared from 9-carbazole carbonyl chloride using procedures similar to those described above in Methods A-E (180mg, 75%): IR (solid, cm^{-1}) 1737, 1655, 1419, 1373, 1224 ; ^1H NMR (400MHz, d_6 -DMSO+TFA) δ 1.36-1.65 (6H, m), 1.94-1.99 (1H, m), 2.12-2.21 (1H, m), 2.59-2.89 (2H, m), 4.32-5.27 (4H, m), 7.30-7.36 (2H, m), 7.48-7.54 (2H, m), 7.63-7.76 (2H, m), 8.17-8.72 (3H, m); ^{13}C NMR (100MHz, DMSO+TFA) δ 19.0 (CH_2), 23.7/23.8 (CH_2), 26.5/26.8 (CH_2), 33.3/33.5 (CH_2), 44.1 (br, CH_2), 50.9/51.4 (CH), 54.5 (br, CH), 82.9/83.1 (2d, J 178.7/178.7, CH_2F), 111.0/111.1 (CH), 111.9 (CH), 119.5/119.7 (CH), 120.6/120.7 (CH), 122.5/122.7 (C), 125.8/125.9 (CH), 137.1/137.4 (C), 153.2/153.3 (C), 170.3/170.4 (C), 170.8/170.9 (C), 201.4/201.5 (2d, J 14.6/14.6, CO); ^{19}F NMR (376MHz, DMSO+TFA) δ d (J, %I) -226.6 (t, 3), -226.8 (t, 3), -230.0 (t, 1), -232.7 (t, 10), -232.7 (t, 10).

Example 7

[3S/R, (2S)]-5-Fluoro-4-oxo-3-{[1-(6H-phenanthridine-5-carbonyl)-piperidine-2-carbonyl]amino}-pentanoic acid



[0078] This was prepared from 9,10-dihydrophenanthridine carbonyl chloride using procedures similar to those described above in Methods A-E (115mg, 61%): IR (solid, cm^{-1}) 1731, 1419, 1363, 1219; ^1H NMR (400MHz, d_6 -DMSO+TFA) δ 1.27-1.69 (5H, m), 1.90-2.06 (1H, m), 2.55-2.87 (2H, m), 3.13-3.21 (2H, m), 4.31-5.26 (6H, m), 7.12-7.48 (6H, m), 7.84-7.86


- 36 -

(2H, m), 8.08-8.58 (1H, m) ; ^{13}C NMR (100MHz, DMSO+TFA) δ 20.5 (CH₂), 24.2 (CH₂), 27.73 (CH₂), 34.6/34.8 (CH₂), 44.9 (CH₂), 48.5/48.7 (CH), 52.1/52.5 (CH), 55.4/55.7 (CH), 84.2 (d, CH₂F), 120.2 (CH), 123.3 (CH), 123.6 (CH), 124.7 (CH), 126.1 (C), 126.3 (CH), 128.0 (CH), 128.3 (CH), 128.7 (CH), 131.6 (C), 134.6 (C), 140.2 (C), 172.1/172.2 (CO), 172.4/172.4 (CO), 203.0 (d, CO); ^{19}F NMR (376MHz, DMSO+TFA) δ -226.8 (t), -226.9 (t), -232.7 (t), -232.9 (t).

10

Example 8


[3S/R, (2S)]-5-Fluoro-3-{2-[(1H-imidazole-2-carbonyl)-amino]-propionylamino}-4-oxo-pentanoic acid, trifluoroacetate salt (Compound 1)



15

Method A:

(2S)-2-[(1H-Imidazole-2-carbonyl)-amino]-propionic acid tert-butyl ester



[0079] To a solution of 1H-imidazole-2-carboxylic acid (0.17g) in *N,N*-dimethylformamide (DMF) (3mL) was added alanine tert-butyl ester hydrochloride (0.22g), diisopropylethyl amine (0.27mL) and HOBT (0.41g) before cooling to 0° C and the reaction mixture was then treated with EDC HCl (0.32g). The cooling bath was removed and the reaction mixture was stirred at ambient temperature for 18 hrs before being diluted with ethyl acetate and washed with water and brine, dried (MgSO₄) and concentrated at reduced pressure. The residue was


25

- 37 -

purified by silica gel chromatography (30% ethyl acetate in hexanes) to afford the sub-title compound as a colourless oil (0.263g, 73%): ¹H NMR 400 MHz CDCl₃ 1.50 (9H, s), 1.51 (3H, d, J 7.2), 3.70 (1H, m), 7.28 (2H, s), 7.78 (1H, d, J 7.6), 11.49 (1H, br s).

Method B:

[3S/R, 4S/R, (2S)]-5-Fluoro-4-hydroxy-3-{2-[(1H-imidazole-2-carbonyl)-amino]-propionylamino}-pentanoic acid tert-butyl ester



[0080] A solution of (2S)-2-[(1H-imidazole-2-carbonyl)-amino]-propionic acid tert-butyl ester (0.257g) in dichloromethane (2ml) was cooled to 0° C before dropwise addition of trifluoroacetic acid and the reaction mixture was warmed to room temperature and stirred for 2hr before evaporation under reduced pressure. The residue was co-evaporated with dichloromethane (twice) and toluene (twice) to leave the required (2S)-2-[(1H-imidazole-2-carbonyl)-amino]-propionic acid that was used without further purification (0.40g).


[0081] A solution of (2S)-2-[(1H-imidazole-2-carbonyl)-amino]-propionic acid and 3-amino-5-fluoro-4-hydroxy-pentanoic acid tert-butyl ester (0.254g) in THF (7mL) was cooled to 0° C before addition of DMAP (0.151g), diisopropylethyl amine (0.56mL), HOBT (0.16g) and EDC HCl (0.23g). The reaction mixture was stirred at ambient temperature for 18 hrs before being concentrated at reduced pressure. The residue was

- 38 -

purified by silica gel chromatography (5% methanol in dichloromethane) to afford the sub-title compound as a colourless solid (0.386g, 97%): ^1H NMR 400MHz $\text{CDCl}_3/\text{CD}_3\text{OD}$ 1.40 (12H, m), 3.92 (1H, m), 4.20-4.55 (4H, m), 7.11 (2H, d, J 15); ^{19}F NMR CDCl_3 -229.74 (m), -229.84 (m), -230.54 (m), -230.87 (m).

Method C:

[3S/R, (2S)]-5-Fluoro-3-{2-[(1H-imidazole-2-carbonyl)-amino]-propionylamino}-4-oxo-pentanoic acid tert-butyl ester




[0082] A solution of [3S/R, (2S)]-5-fluoro-4-hydroxy-3-{2-[(1H-imidazole-2-carbonyl)-amino]-propionylamino}-pentanoic acid tert-butyl ester (0.381g) in dichloromethane was cooled to 0°C before addition of 1,1,1 triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one (0.476g). The mixture was stirred at room temperature for 2h before addition of an additional portion of 1,1,1 triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one (0.05g) and reaction mixture was then stirred for 90 min before being concentrated at reduced pressure. The residue was dissolved in ethyl acetate and washed with a 1:1 mixture of aqueous NaHSO_4 and aqueous $\text{Na}_2\text{S}_2\text{O}_3$. The organic layer was collected, dried (MgSO_4) and concentrated. The residue was purified by flash chromatography (5% methanol in dichloromethane) to give the sub-title compound as a colourless foam (319mg, 84%): ^1H NMR 400MHz CDCl_3 1.37+1.43 (9H, 2xs), 1.54 (3H, m), 2.85 (1H, m), 3.03 (1H, m), 4.85-5.30 (4H, m), 7.18

- 39 -

(2H, d, J 16), 7.90 (1H, m), 7.98 (1H, m), 11.37+11.45 (1H, 2xs); ^{19}F NMR 376 MHz CDCl_3 -231.85 (t, J 48), -232.12 (t, J 48).

5 Method D:

Compound 1




[0083] A solution of [3S/R, (2S)]-5-Fluoro-3-{2-[(1H-
imidazole-2-carbonyl)-amino]-propionylamino}-4-oxo-
10 pentanoic acid tert-butyl ester (0.31g) in
dichloromethane (2ml) was cooled to 0° C before
dropwise addition of trifluoroacetic acid and the
reaction mixture was warmed to room temperature and
stirred for 2hr before evaporation under reduced
15 pressure. The residue was co-evaporated with
dichloromethane (twice) and triturated under ether to
give the title compound as a colourless solid (0.35g):
IR 1785.7, 1730.1, 1653.7, 1538.1, 1458.2, 1384.2,
1268.7, 1188.4, 1150.9, 1053.3, 992.13, 931.8, 867.9,
20 847.0, 768.5 cm^{-1} ; ^1H NMR 400MHz DMSO-d_6 1.37 (3H, d),
2.40-2.85 (2H, m, asp CH_2), 4.34-4.75 (2.5H, m, 2xCH+0.5 CH_2F),
5.13-5.41 (1.5H, m, CH_2F), 7.50 (2H, s, imidazole CHs), 8.58-
8.79 (2H, m, NHs), ^{13}C NMR 100 MHz DMSO-d_6 18.13, 18.85 (ala
25 CH_3); 33.13, 34.75 (asp CH_2), 48.68, 52.41 (CHs), 83.46,
85.21 (CH_2F), 123.67 (CH imidazole),
139.57, 158.86, 172.35 (m) (C=Os), 202.70 (5 peaks ketone);
 ^{19}F NMR 376 MHz DMSO-d_6 decoupled -75.19 (3F, s, CF_3COOH),
-(226.89, 226.96, 230.80, 231.59, 232.95, 233.06 (1F, 6xs,
 COCH_2F ring opened and ring closed).

- 40 -

Example 9

[3S/R, (2S)]-3-{2-[(1H-Benzoimidazole-2-carbonyl)-amino]-propionylamino}-5-fluoro-4-oxo-pentanoic acid, trifluoroacetate salt (Compound 2)



5

[0084] This was prepared from 1H-benzoimidazole-2-carboxylic acid using procedures similar to those described in Methods A-D above (142 mg, 90% for final step): (compound isolated as the TFA salt) off-white solid; IR (solid, cm^{-1}) 3277.9, 1654.6, 1526.6, 1188.6, 1142.5, 1050.4, 927.5, 748.2, 712.4; ^1H NMR (DMSO- d_6) 1.42 (3H, d), 2.51-2.95 (2H, m), 4.21-4.75 (2H, m), 4.76-5.60 (3H, brm), 7.41 (2H, m), 7.65 (2H, m), 8.21-9.05 (2H, m); ^{13}C NMR (DMSO- d_6) 18.0, 18.7, 18.8 (Ala CH_3), 37.2, 34.6, 34.7 (Asp CH_2), 47.6, 48.8, 48.85, 49.1 (Asp CH), 52.0, 52.5 (Ala CH), 83.5, 85.2, 85.3, 103.8, 106.0 (CH_2F), 116.6, 123.9 (Aryl CH), 145.3, 145.4, (Aryl C), 158.4, 158.7, 158.8, 172.1, 172.2, 172.4, 172.5, 172.6, 172.7, 173.2 (NC=O), 202.6, 202.7, 202.8, 202.9 (C=O); Found M^+ 364.1177. $\text{C}_{16}\text{H}_{17}\text{FN}_4\text{O}_5$ requires M^+ 364.1183 (1.8 ppm).

10

15


20

Example 10

[3S/R, (2S)]-5-Fluoro-3-{2-[(1H-imidazole-2-carbonyl)-amino]-butyrylamino}-4-oxo-pentanoic acid, trifluoroacetate salt (Compound 3)

25


- 41 -



[0085] This was prepared from 1*H*-benzimidazole-2-carboxylic acid using procedures similar to those described in Methods A-D above (147 mg, 64% for final
 5 step): IR(cm^{-1}) 3280.0, 1659.5, 157.9, 1192.5, 1141.6, 784.7, 721.1; ^1H NMR 400MHz DMSO- d_6 0.95 (3H, m), 1.78 (2H, m), 2.58-2.98 (2H, m), 4.30-4.78 (2.5H, m), 5.10-5.42 (1.5H, m), 7.41 (2H, s), 8.44+8.75 (2H, 2xm); ^{13}C NMR 100 MHz DMSO- d_6 10.19, 10.29, 15.52 (CH_3), 25.42,
 10 25.49, 26.03, 33.06, 33.13, 34.65, 34.80 (CH_2), 47.45, 47.53, 52.0, 53.96, 54.13 (CH) 65.27 (CH_2), 84.36 (d, J 177, CH_2F), 103.81, 104.00 (C), 123.89 (CH), 139.74 (C=O), 156.90, 158.39, 158.74, 171.51, 171.80, 171.83, 172.02, 173.11 (C=O), 202.51, 202.66, 202.76, 202.90
 15 ($\text{CH}_2\text{FC=O}$); ^{19}F NMR 376 MHz DMSO- d_6 -226.82 (t, J 45), -226.84 (t, J 45), -230.67 (t, J 45), -231.43 (t, J 45), -232.79 (t, J 45), -232.82 (t, J 45).

Example 11

[3*S*/*R*, (2*S*)]-5-Fluoro-3-{2-[(1*H*-imidazole-2-carbonyl)-
 20 amino]-3-methylbutyrylamino}-4-oxo-pentanoic acid
 (Compound 4)




[0086] This was prepared from 1*H*-benzimidazole-2-carboxylic acid using procedures similar to those

- 42 -

described in Methods A-D above (80g, 85% for final step): white powder, IR (solid, cm^{-1}) 1736, 1649, 1557, 1511, 1455, 1434, 1393; ^1H NMR (DMSO+TFA) 0.92-0.95 (6H, m), 2.06-2.15 (1H, m), 2.56-2.90 (2H, m), 4.33-5.36 (4H, m), 7.79 (2H, s), 8.58-8.90 (2H, m); ^{19}F NMR (DMSO+TFA) -226.8 (t), -230.6 (t), -231.0 (t), -232.5 (t), -232.6 (t); ^{13}C NMR (DMSO+TFA) 18.1/18.4 (CH_3), 19.2/19.3 (CH_3), 34.5/34.8 (CH_2), 51.9/52.2 (CH), 58.5/58.8 (CH), 84.3/84.4 (2d, J 178.7/178.7, CH_2F), 122.0 (CH), 137.5 (C), 153.7 (C), 170.6 (C), 171.9/172.0 (C), 202.5/202.8 (2d, J 14.6/14.6, CO).

Example 12


[3S/R, (2S)]-3-{2-[(1H-Benzoimidazole-2-carbonyl)-amino]-3-methylbutyrylamino}-5-fluoro-4-oxo-pentanoic acid (Compound 5)



[0087] This was prepared from 1H-benzoimidazole-2-carboxylic acid using procedures similar to those described in Methods A-D above (90mg, 87% for final step): white powder, IR (solid, cm^{-1}) 1737, 1665, 1527, 1373, 1194, 1137; ^1H NMR (DMSO- d_6) 0.90-0.95 (6H, m), 2.15-2.18 (1H, m), 2.59-2.92 (2H, m), 4.33-4.76 and 5.12-5.38 (4H, 2m), 7.31-7.35 (2H, m), 7.66-7.68 (2H, m), 8.36-8.82 (2H, m); ^{19}F NMR (DMSO+TFA) -226.7 (t), -226.9 (t), -232.4 (t), -232.6 (t); ^{13}C NMR (DMSO- d_6) 18.3/18.4/18.5/18.7 (CH_3), 19.4/19.5 (CH_3), 31.0/31.1/31.6 (CH), 34.7/34.8 (CH_2), 51.8/52.1 (CH), 57.9/58.3/58.6 (CH), 84.3/84.4 (2d, J 178.7/178.7, CH_2F), 124.0 (CH), 145.2/145.2 (C),

- 43 -


158.4/158.5/158.7/158.8 (C), 170.9/171.1/171.2 (C),
172.0/172.0 (C), 173.1 (C), 173.9 (C), 202.06/202.6
(2d, J 13.8, CO).

Example 13

5

[3S/R(2S)]-3-[2-(Carbazol-9-yl-2-oxo-ethyl)-
pentanoylamino]-5-fluoro-4-oxo-pentanoic acid

Method A: (4S)-Benzyl-3-pentanoyl-oxazolidin-2-one




10

[0088] A solution of 4(S)-(-)-benzyl-2-oxazolidinone
(10g, 56.43mmol) in anhydrous THF (200ml) at -78°C was
treated with a 2.5M solution of n-butyl lithium in
hexanes (23.70ml, 59.26mmol) with stirring. The
15 reaction mixture was allowed to stir at -78°C for 30min
before valeryl chloride (7.57ml, 62.10mmol) was added.
The reaction mixture was then allowed to warm to
ambient temperature over 15h after which it was diluted
with NH₄Cl solution, diluted with ethyl acetate and
20 washed with brine. The organic phase was dried (Na₂SO₄)
and concentrated to give a gum. This was purified by
flash chromatography (10% EtOAc in 40/60 hexanes) to
give the sub-title compound (14.61g, 99%) as a
colourless oil: ¹H NMR (400MHz, CDCl₃) δ 0.94-1.20 (3H,
25 m), 1.35-1.50 (2H, m), 1.62-1.80 (2H, m), 2.74-2.84

- 44 -

(1H, m), 2.86-3.08 (2H, m), 3.27-3.39 (1H, m), 4.11-4.26 (2H, m), 4.62-4.76 (1H, m), 7.18-7.40 (5H, m).

Method B: [4S(3R)]-3-(4-Benzyl-2-oxo-oxazolidine-3-carbonyl)-hexanoic acid *tert*-butyl ester




[0089] A solution of (4S)-benzyl-3-pentanoyl-oxazolidin-2-one (14.20g, 54.34mmol) in THF (100ml) at -78°C was treated over 10min with a 1M solution of sodium bis(trimethylsilyl)amide in THF (59.80ml, 59.77mmol) with stirring. The reaction mixture was allowed to stir at -78°C for 30min before *tert*-butyl bromoacetate (10.43ml, 70.64mmol) was added. The reaction mixture was then allowed to stir for a further 3.5h at -78°C after which it was diluted with NH₄Cl solution, diluted with ethyl acetate and washed sequentially with NaHCO₃ solution and brine. The organic phase was dried (Na₂SO₄) and concentrated to give a gum. On standing a white solid was formed and this was recrystallized from 40/60 DCM/hexanes to give the sub-title compound (14.62g, 72%) as a white solid:

¹H NMR (400MHz, CDCl₃) δ 0.81-1.20 (3H, m), 1.21-1.76 (13H, m), 2.41-2.55 (1H, m), 2.66-2.92 (2H, m), 3.27-3.40 (1H, m), 4.05-4.26 (2H, m), 4.61-4.72 (1H, m), 7.12-7.40 (5H, m).

Method C: (2R)-2-Propyl-succinic acid 1-benzyl ester 4-*tert*-butyl ester


- 45 -



[0090] A solution of benzyl alcohol (4.62ml, 44.64mmol) in THF (80ml) at -20°C was treated with a
 5 2.5M solution of n-butyl lithium in hexanes (13.36ml, 33.48mmol) with stirring. The reaction mixture was allowed to warm to -5°C over 40min before a solution of [4S(3R)]-3-(4-benzyl-2-oxo-oxazolidine-3-carbonyl)-hexanoic acid tert-butyl ester (8.38g, 22.32mmol) in
 10 THF (20ml) was added. The reaction mixture was warmed to ambient temperature over 15h after which it was diluted with NH₄Cl solution and ethyl acetate and washed with brine. The organic phase was dried (Na₂SO₄) and concentrated to give a gum. This was purified by flash
 15 chromatography (11% EtOAc in 40/60 hexanes) to give the sub-title compound (4.56g, 67%) as a colourless oil: ¹H NMR (400MHz, CDCl₃) δ 0.83-1.00 (3H, m), 1.21-1.71 (13H, m), 2.34-2.45 (1H, m), 2.75-2.95 (1H, m), 5.09-5.25 (2H, m), 7.30-7.43 (5H, m).

20

Method D: (2R)-2-Propyl-succinic acid 1-benzyl ester




[0091] A stirred solution of (2R)-2-propyl-succinic
 25 acid 1-benzyl ester 4-tert-butyl ester (4.56g, 14.88mmol) in anhydrous DCM (20ml) at 0°C was treated with a solution of trifluoroacetic acid (10ml) in

- 46 -

anhydrous DCM (10ml). The reaction mixture was allowed to warm to ambient temperature over 3h before being concentrated under reduced pressure. The residue was dissolved in dry DCM, before concentrating again. This process was repeated several times in order to remove excess trifluoroacetic acid to leave the sub-title compound (3.70g, 99%) as a gum: ^1H NMR (400MHz, CDCl_3) δ 0.82-0.99 (3H, m), 1.21-1.76 (4H, m), 2.45-2.60 (1H, m), 2.76-3.00 (2H, m), 5.10-5.21 (2H, m), 7.28-7.43 (5H, m), 7.83-8.18 (1H, m).

Method E: (2R)-2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoic acid benzyl ester



[0092] A stirred solution of carbazole (2.49g, 14.88mmol) in anhydrous THF (30ml) at -78°C was treated with a 1.0M solution of lithium bis(trimethylsilyl)amide in THF (14.88ml, 14.88mmol). The reaction mixture was allowed to warm to ambient temperature over 2h before being re-cooled to -78°C .


[0093] A solution of (2R)-2-propyl-succinic acid 1-benzyl ester (3.70g, 14.78mmol) in anhydrous DCM (20ml), stirring at 0°C , was treated with oxalyl chloride (1.43ml, 16.37mmol) and DMF (14 drops). The reaction mixture was stirred at 0°C for 1h before being concentrated *in vacuo*. The residue was dissolved in anhydrous THF (10mL) and added to the lithium anion of carbazole previously prepared at -78°C . The reaction

- 47 -

mixture was warmed to ambient temperature over 40h after which it was diluted with NH₄Cl solution, and ethyl acetate and washed sequentially with 2N HCl, NaHCO₃ solution and brine. The organic phase was dried (Na₂SO₄) and concentrated to give a gum which was purified by flash chromatography (10% EtOAc in 40/60 hexanes) to give the sub-title compound (4.50g, 76%) as a semi-solid/oil which also contained carbazole: ¹H NMR (400MHz, CDCl₃) δ 0.82-1.05 (3H, m), 1.11-1.99 (4H, m), 3.18-3.38 (2H, m), 3.56-3.71 (1H, m), 5.10-5.30 (2H, m), 7.11-7.60 (9H, m), 7.92-8.29 (4H, m).

Method F: (2R)-2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoic acid

15




[0094] A stirred solution of (2R)-2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoic acid benzyl ester (4.50g, 11.26mmol) in EtOAc (60ml) was treated with 10% Pd on carbon (~400mg) and the reaction mixture was then placed under an atmosphere of hydrogen. After 1h further 10% Pd on carbon (~300mg) was added and the reaction mixture was placed under hydrogen, with stirring, for a further 3h after which the reaction mixture was filtered through a celite pad and concentrated to give the sub-title compound (2.94g, 84%) as a white solid which also contained carbazole: ¹H NMR (400MHz, CDCl₃) δ 0.92-1.04 (3H, m), 1.32-2.00 (4H,

- 48 -

m), 3.19-3.34 (2H, m), 3.58-3.70 (1H, m), 7.30-7.53 (4H, m), 8.00-8.30 (4H, m).

Method G: [3S/R, 4S/R, (2R)]-3-[2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoylamino]-5-fluoro-4-hydroxy-pentanoic acid *tert*-butyl ester




[0095] A stirred mixture of (2R)-2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoic acid (2.94g, 9.50mmol), 3-amino-5-fluoro-4-hydroxy-pentanoic acid *tert*-butyl ester (2.07g, 9.99mmol), HOBT (1.41g, 10.43mmol), DMAP (1.34g, 10.97mmol) and anhydrous THF (40ml) was cooled to 0°C before EDC (2.00g, 10.43mmol) was added. The mixture was allowed to warm to room temperature during 16h then concentrated under reduced pressure. The residue purified by flash chromatography (33% EtOAc in 40/60 hexanes) to give the sub-title compound (2.51g, 53%) as a foam: ¹H NMR (400MHz, CDCl₃) δ 0.90-1.03 (3H, m), 1.20-1.90 (13H, m), 2.50-3.00 (3H, m), 3.12-3.26 (1H, m), 3.59-3.80 (2H, m), 4.00-4.68 (3H, m), 6.53-6.89 (1H, m), 7.30-7.52 (4H, m), 7.95-8.05 (2H, m), 8.15-8.26 (2H, m); ¹⁹F NMR (376 MHz, CDCl₃) -229.10, -229.34, -230.95, -231.09.

25

Method H: [3S/R, (2R)]-3-[2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoylamino]-5-fluoro-4-oxo-pentanoic acid *tert*-butyl ester


- 49 -



[0096] A stirred solution of [3S/R, 4S/R, (2R)]-3-
 5 [2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoylamino]-5-
 fluoro-4-hydroxy-pentanoic acid tert-butyl ester
 (2.51g, 5.03mmol) in anhydrous DCM (60ml) was treated
 with 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-
 3(1H)-one (2.35g, 5.53mmol) at 0°C . The resulting
 10 mixture was kept at 0°C for 3h, diluted with DCM, and
 then washed sequentially with saturated aqueous sodium
 thiosulphate, NaHCO₃ solution and brine. The organics
 were dried (Na₂SO₄) and concentrated. The residue was
 purified by flash chromatography (25% ethyl acetate in
 40/60 hexanes) to afford the sub-title compound as an
 15 off white solid(1.437g, 57%): IR (solid, cm⁻¹) 1722,
 1689, 1636, 1531, 1441, 1365, 1279, 1155; ¹H NMR
 (400MHz, CDCl₃) δ 0.85-1.50 (3H, m), 1.35-1.54 (11H, m),
 1.55-1.69 (1H, m), 1.78-1.95 (1H, m), 2.67-3.28 (4H,
 m), 3.60-3.79 (1H, m), 4.80-5.59 (3H, m), 6.89-7.04
 20 (1H, m), 7.33-7.54 (4H, m), 7.98-8.04 (2H, m), 8.15-
 8.28 (2H, m); ¹³C (100MHz, CDCl₃) δ 14.12, 14.40, 14.47,
 14.60, 20.78, 20.84, 21.47, 28.32, 28.42, 28.48, 29.77,
 33.63, 34.58, 34.91, 40.05, 43.05, 43.26, 43.29, 52.60,
 53.00, 53.64, 66.90, 66.99, 82.62, 82.69, 85.53,
 25 116.88, 116.94, 120.28, 120.31, 124.27, 127.76, 127.86,
 128.69, 128.77, 128.99, 138.80, 171.21, 171.29, 172.21,
 172.25, 175.53, 176.03, 203.04, 203.20, 203.30, 203.46;
¹⁹F (376MHz, CDCl₃) δ -232.12 , -233.24.


- 50 -

Method I: [3S/R, (2R)]-3-[2-(2-Carbazol-9-yl-2-oxo-ethyl)-pentanoylamino]-5-fluoro-4-oxo-pentanoic acid



5 [0097] A solution of [3S/R, (2R)]-3-[2-(2-carbazol-9-yl-2-oxo-ethyl)-pentanoylamino]-5-fluoro-4-oxo-pentanoic acid *tert*-butyl ester (1.43g, 2.88mmol) in anhydrous DCM(20ml) was treated with a solution of TFA (10ml) in anhydrous DCM (10ml) with stirring. The
 10 mixture was stirred at 0°C for 2h then at room temperature for 2h. The mixture was concentrated under reduced pressure and then the residue was dissolved in dry DCM. This process was repeated several times in order to remove excess trifluoroacetic acid. The off-
 15 white solid was recrystallized from Et₂O / 40/60 hexanes to give the title compound as a white powder (71mg): IR (solid, cm⁻¹) 1746, 1689, 1641, 1541, 1436, 1374, 1284, 1207, 1160cm⁻¹; ¹H NMR (400MHz, d₆-DMSO) δ 0.80-1.00 (3H, m), 1.20-1.76 (4H, m), 2.30-2.90 (2H, m), 2.95-
 20 3.24 (1H, m), 3.26-3.59 (2H, m), 4.25-4.79 (1.5H, m), 5.02-5.43 (1.5H, m), 7.36-7.58 (4H, m), 8.10-8.30 (4H, m), 8.54-8.91 (1H, m); ¹³C NMR (100MHz, DMSO) δ 14.31, 20.03, 20.13, 21.92, 22.51, 34.36, 34.77, 41.20, 41.62, 44.06, 51.77, 52.84, 83.45, 85.22, 116.70, 120.54,
 25 123.91, 124.01, 127.85, 126.01, 138.20, 172.15, 172.36, 172.96, 173.00, 175.32, 175.48, 202.60, 203.10; ¹⁹F (376MHz, DMSO) δ -226.68, -226.73, -231.21, -232.95, -233.38, -233.52.

- 51 -

Example 14

5 [3S/R(2S)]-3-[2-(2-Carbazol-9-yl-2-oxo-ethyl)-3-methyl-
butyrylamino]-5-fluoro-4-oxo-pentanoic acid

[0098] This was prepared using procedures similar to those described in Methods A-I. The product was isolated as a white powder (71% for final step): IR (solid, cm^{-1}) 1739, 1682, 1646, 1545, 1447, 1381, 1290,
 10 1209, 1170 cm^{-1} ; ^1H NMR (400MHz, DMSO + TFA) δ 0.79-1.08 (6H, m), 1.89-2.15 (1H, m), 2.31-3.60 (5H, m), 4.21-4.78 (1.25H, m), 4.98-5.45 (1.75H, m), 7.38-7.60 (4H, m), 8.14-8.35 (4H, m), 8.56-8.90 (1H, m); ^{13}C NMR (100MHz, DMSO) δ 20.46, 20.84, 21.04, 21.21, 30.77,
 15 30.85, 33.37, 34.83, 35.24, 38.16, 38.89, 47.67, 48.23, 52.19, 53.43, 83.96, 84.01, 85.72, 85.77, 117.16, 121.02, 124.43, 126.42, 126.52, 128.42, 138.75, 172.64, 172.90, 173.85, 173.90, 174.74, 174.93, 175.16, 202.91, 203.04, 203.51, 203.65; ^{19}F (376MHz, DMSO) δ -226.63, -
 20 226.68, -231.24, -233.16, -233.38, -233.55.

Biological MethodsExample 15

[0099] ICE Inhibition may be measured by methods
 25 known in the art.

[0100] The effectiveness of ICE inhibitors in the treatment of MWS, FCU, FMF, CINCAS, NOMID, TRAPS, HIDS, systemic onset Juvenile Idiopathic Arthritis, Still's disease and Macrophage Activation Syndrome and other

- 52 -

autoinflammatory diseases, conditions or syndromes may be demonstrated by two studies:

Example 16

5 [0101] Blood samples are collected from patients with MWS, FCU, FMF, CINCAS, NOMID, TRAPS, HIDS, systemic onset Juvenile Idiopathic Arthritis, Still's disease and Macrophage Activation Syndrome or other autoinflammatory disease or condition and cultured in
10 *vitro* as whole blood or processed for the preparation of peripheral blood mononuclear cells (PBMCs). The production of IL-1 β is evaluated under normal, unstimulated conditions or in the presence of a stimulus such as lipopolysaccharide, *Staphylococcus*
15 *aureus*, nigericin or phorbol myristate. ICE inhibitors are added to the cultures at concentrations of 1-100,000 nM to demonstrate their ability to inhibit the production of IL-1 β and/or IL-18 by cells from patients with MWS, FCU, FMF, CINCAS, NOMID, TRAPS, HIDS,
20 systemic onset Juvenile Idiopathic Arthritis, Still's disease and Macrophage Activation Syndrome or other autoinflammatory disease, condition or syndrome.

Example 17

25 [0102] Patients with MWS, FCU, FMF, CINCAS, NOMID, TRAPS, HIDS, systemic onset Juvenile Idiopathic Arthritis, Still's disease and Macrophage Activation Syndrome or other autoinflammatory disease or condition are enrolled into a clinical trial to receive treatment
30 with an ICE inhibitor. The drug is administered at a dose of 10-10,000 mg by an appropriate route (oral, intravenous, subcutaneous, etc.) at a dosing frequency of 1-4 times/day or as needed to control the symptoms

- 53 -

and other manifestations of the disease, condition or syndrome. Treatment duration may range from periodic use of one or more days to daily use over the life-time of the patient. The efficacy is demonstrated by virtue of their ability to reduce the clinical signs and symptoms, such as fever, urticaria, arthralgia, serositis, systemic amyloidosis, robust acute phase response and prominent neutrophilia in affected tissues.

10

Example 18

Inhibition by compound 7 of LPS-stimulated IL-1 β and IL-18 release from PBMCs of patients with FCAS compared with healthy controls.

15 [0103] Cryopyrin, the protein encoded by the *CIAS1* gene that is mutated in FCAS, modulates ICE/caspase-1 activation and its processing of pro-IL-1 β to active IL-1 β , as well as pro-IL-18 to active IL-18. It may be expected that the mutated cryopyrin protein would
20 interact differently with ICE/caspase-1 and could alter the ICE/caspase-1 conformation in a way that would alter the interaction of the ICE inhibitor compound with ICE/caspase-1, thus reducing the ability to the compound to inhibit ICE activation and IL-1 β and IL-18
25 processing. Therefore, the experiments of this example were conducted to confirm that compound 7 is able to inhibit ICE in cells of FCAS patients.

[0104] FCAS patients from 3 families were studied along with healthy controls. The FCAS patients
30 expressed 3 different *CIAS1* gene mutations. Five related patients (mother, two daughters, granddaughter, and a distant male relative) possessed the

- 54 -

common L353P ancestral mutation in cryopyrin that has been described previously (Johnstone RF, Dolen WK, Hoffman HM. A large kindred with familial cold autoinflammatory syndrome. *Ann Allergy Asthma Immunol.* 2003; 90(2): 233-7). The other patients possessed novel mutations that have not been reported. Two patients (mother and daughter) had a T-to-A transition at nucleotide 1976 in the CIAS1 gene, leading to a M659K substitution in cryopyrin. One patient had a G-to-A transition at nucleotide 1573 in the CIAS1 gene, generating a E525K amino acid substitution that has not been reported, as well as the V198M variation described previously in several patients with diverse phenotypes, but also in normal controls. All of the mutations are located in exon 3, which encodes the NBS (NACHT) domain or NBS associated domain, and all are within 3 amino acids of other reported mutations, consistent with studies suggesting mutational hot spots. (Neven B, Callebaut I, Prieur AM, Feldmann J, Bodemer C, Lepore L, Derfalvi B, Benjaponpitak S, Vesely R, Sauvain MJ, Oertle S, Allen R, Morgan G, Borkhardt A, Hill C, Gardner-Medwin J, Fischer A, de Saint Basile G. Molecular basis of the spectral expression of CIAS1 mutations associated with phagocytic cell-mediated autoinflammatory disorders CINCA/NOMID, MWS, and FCU. *Blood.* 2004;103(7): 2809-15).

[0105] Whole blood samples were collected from patients with FCAS and from healthy volunteer controls and cultured *in vitro*. The production of IL-1 β was evaluated under normal, unstimulated conditions and in the presence of lipopolysaccharide at a range of concentrations from 10-10,000 ng/mL. Compound 7 was added to the cultures at concentrations of 30-30,000 nM

- 55 -

to observe its ability to inhibit the production of IL-1 β and IL-18 by cells from patients with FCAS or healthy volunteers. Figure 1 illustrates the results of these experiments.

5 [0106] More specifically, in Figure 1A, PBMCs from FCAS patients and normal controls were incubated with 10 ng/ml LPS in the presence of increasing concentrations of compound 7 (0.03-30 μ M) for 24 hours. IL-1 β in cell culture media was measured by ELISA and
10 the results for each subject were normalized to the level of IL-1 β release in the absence of compound 7. Results were reported in mean \pm SEM, for 3 FCAS patients and 5 control patients. LPS-stimulated IL-1 β was inhibited by compound 7 with potency very similar
15 to that of PBMCs from normal subjects (IC₅₀ value ~1 μ M).

[0107] Figure 1B illustrates the release of IL-1 β by PBMCs from FCAS patients or control subjects after exposure to LPS for 24 hours in the absence or presence
20 of 10 μ M compound 7. Results were reported in mean \pm SEM for 7 FCAS patients and 5 control patients. LPS stimulated greater amounts of IL-1 β release by PBMCs from FCAS patients than normal subjects and FCAS PBMCs responded to lower concentrations of LPS.

25 Nevertheless, LPS-stimulated IL-1 β was inhibited by 10 μ M of compound 7 similar to that of PBMCs from normal subjects.

[0108] Figure 1C illustrates the release of IL-18 by PBMCs from FCAS patients or control subjects after 24
30 hrs exposure to LPS in the absence or presence of 10 μ M compound 7. Results were reported in mean \pm SEM for 3

- 56 -

FCAS patients and 3 control patients. LPS stimulated greater amounts of IL-18 release by PBMCs from FCAS patients than normal subjects and FCAS PBMCs responded to lower concentrations of LPS. Nevertheless, LPS-stimulated IL-18 was inhibited by 10 μ M compound 7 to a similar extent as that of PBMCs from normal subjects.

[0109] These data demonstrate that FCAS patients exhibited a marked hyper-responsiveness of both IL-1 β and IL-18 secretion to LPS stimulation, but no evidence of increased basal secretion of these cytokines, or alterations in basal or stimulated pro-IL-1 β levels. Compound 7 blocked IL-1 β and IL-18 secretion (Figure 1) with equal potency in LPS-stimulated cells from FCAS and control subjects. Furthermore, the FCAS patient data presented herein suggests that the mutations of the *CAIS1* gene and its encoded cryopyrin protein in FCAS patients do not alter the sensitivity of ICE/caspase-1 to ICE inhibitors like compound 7.

Example 19

Inhibition by compound 7 of LPS-stimulated IL-1 β release from PBMCs from 4 MWS patients incubated with 10 ng/ml LPS in the presence of increasing concentrations of compound 7 (0.03-10 μ M) for 24 hour.

[0110] As described in Example 18, it may be expected that the mutated cryopyrin protein would interact differently with ICE/caspase-1 and could alter the ICE/caspase-1 conformation in a way that would alter the interaction of the ICE inhibitor compound with ICE/caspase-1, thus reducing the ability to the compound to inhibit ICE activation and IL-1 β and IL-18 processing. Therefore, the experiments of this example

- 57 -

were conducted to confirm that compound 7 is able to inhibit ICE in cells of MWS patients.

[0111] Four MWS patients were studied, three of whom possessed the arginine to tryptophan replacement in position 262 (R262W) and one the threonine to methionine mutation in position 350 (T350M).

[0112] Whole blood samples were collected from patients with MWS and cultured *in vitro*. The production of IL-1 β was evaluated under normal, unstimulated conditions and in the presence of lipopolysaccharide at a range of concentrations from 10-10,000 ng/mL. Compound 7 was added to the cultures at concentrations of 30-10,000 nM to observe its ability to inhibit the production of IL-1 β by cells from patients with MWS or healthy volunteers. Figure 2 illustrates the results of these experiments.

[0113] More specifically, in Figure 2, IL-1 β in cell culture media was measured by ELISA and the results for each subject were normalized to the level of IL-1 β release in the absence of compound 7. Results were reported individually for each of the 4 MWS patients. LPS-stimulated IL-1 β release by PBMCs from MWS patients was inhibited by compound 7 with potency similar to that of PBMCs from normal subjects (as observed in example 18). This indicates that the cryopyrin mutation in these MWS patients does not alter the ICE/caspase-1 activation in a way that it alters the potency of the interaction of the active metabolite (compound 8) of compound 7 with ICE/caspase-1.

30

- 58 -

Example 20Tablet Formation

[0114] The composition of compound 7 tablets used in Example 21 below is provided in Table 1. The drug product was formulated to provide 300 mg of compound 7 per tablet.

Table 1: Composition of Compound 7 300 mg Tablets

Component	Quantity (mg/tablet)	Function
Compound 7	300	Active Ingredient
Microcrystalline Cellulose (NF)	277.50	Filler
Pregelatinized Starch (NF)	131.25	Disintegrant
Sodium Starch Glycolate (NF)	15.00	Disintegrant
Colloidal Silicon Dioxide (NF)	11.25	Glidant
Talc (USP)	7.50	Glidant
Magnesium Stearate (NF)	7.50	Lubricant
Total	750	

Example 21

10 Clinical response of 4 MWS patients to 900 mg compound 7 administered orally three times daily for 14 days.

[0115] Four patients with MWS were enrolled into an open-label clinical trial to receive treatment with compound 7. Three of the MWS patients possessed the arginine to tryptophan replacement in position 262 (R262W) and one patient the threonine to methionine mutation in position 350 (T350M).

[0116] Compound 7 was administered orally at a dose of 900 mg 3 times daily for 14 days. The anti-inflammatory activity of compound 7 in MWS patients was demonstrated by marked reduction in levels of serum interleukin-18, amyloid A and C-reactive protein during the period of treatment (see Figure 3). The clinical

- 59 -

efficacy of compound 7 was demonstrated by virtue of the reduction in the clinical symptoms reported by the patients, illustrated in Figure 3 as the sum total of symptoms including rash, fever/chills,

5 arthralgia/myalgia, eye discomfort, fatigue (see Figure 3). Compound 7 was also well tolerated by the four patients during the 14-day treatment period.

[0117] Specifically, in Figure 3A, serum interleukin-18 (IL-18) was measured before (day -9 and
10 day 0 (first point)), during (day 0 (second point), day 7 and day 14, (first point)) and after (day 14 (second point) and day 21) treatment with compound 7. On average, serum IL-18 levels were reduced by 55% at day 14. Data were reported in mean \pm SEM from 4 patients.
15 Baseline levels of serum IL-18 were 170-370 pg/mL and were only slightly elevated relative to those typically observed in healthy control subjects.

[0118] In Figure 3B, Serum amyloid A (SAA) and C-reactive protein (CRP) were measured before, during
20 and after treatment with compound 7, and the amounts of SAA and CRP were reduced on average by 75% and 65%, respectively. Baseline levels of SAA were 150-650 mg/L and of CRP were 20-90 mg/L, both considerably elevated relative to healthy control subjects.

25 [0119] Figure 3C illustrates that the self-reported patient total symptom scores were reduced on average by 60% during treatment. Individual symptoms (rash, fever/chills, arthralgia/myalgia, eye discomfort, fatigue) were rated daily by patients to be absent (0),
30 minimal (1), mild (2), moderate (3) or severe (4) and the total score was calculated as the sum of individual symptom scores (range 0-20). Patients were experiencing mild-moderate symptoms at the beginning of

- 60 -

treatment (scores 10-15), which were reduced to a minimal level (scores about 5) during treatment with compound 7.

[0120] Thus, in this example we have demonstrated
5 that an ICE/caspase-1 inhibitor, compound 7, is effective and beneficial in the treatment of an autoinflammatory disease, specifically Muckle-Wells Syndrome, since administration of compound 7 decreases markers of inflammation, including serum interleukin-
10 18, serum amyloid A and serum C-reactive protein and reduces the symptoms of MWS experienced by the patients.

[0121] While we have described a number of
15 embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments, which utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific
20 embodiments, which have been represented by way of example.

25

- 61 -

What is Claimed is:

1. A method for ameliorating, treating, or preventing an autoinflammatory disease in a subject, comprising administering a compound that inhibits ICE or a pharmaceutical composition comprising said compound to said subject.

2. A method for ameliorating, treating, or preventing Cryopyrin-associated Periodic Syndromes, Muckle-Wells Syndrome (MWS), Familial Cold Autoinflammatory Syndrome (FCAS, also known as Familial Cold Urticaria or FCU), Familial Mediterranean Fever (FMF), Chronic Infantile Neurological Cutaneous and Articular Syndrome (CINCAS), a.k.a. Neonatal Onset Multisystem Inflammatory Disease (NOMID), TNFR1-Associated Periodic Syndrome (TRAPS), Hyper-IgD periodic fever Syndrome (HIDS), Blau's syndrome, systemic onset juvenile idiopathic arthritis (Still's disease), or macrophage activation syndrome in a subject, comprising administering a compound that inhibits ICE or a pharmaceutical composition comprising said compound to said subject.


3. The method according to claim 1 or claim 2, wherein the compound is according to any of WO 00/55114, WO 00/55127, WO 00/61542, WO 00/61542, WO 01/05772, WO 01/10383, WO 01/16093, WO 01/42216, WO 01/72707, WO 01/90070, WO 01/94351, US Publication 2003/0092703, WO 02/094263, US Publication 2002/0169177, US Patent 6,184,210, US Patent 6,184,244, US Patent 6,187,771, US Patent 6,197,750, US Patent 6,242,422, April 2001 American Chemical Society (ACS) meeting in San Diego, California, USA, WO 02/22611, US

- 62 -


Publication 2002/0058630, US Publication 2003/0096737, WO 95/35308, WO 97/22619, WO 99/47545, and WO 01/90063.

4. The method according to claim 3, wherein the compound is according to any of WO 95/35308, WO 97/22619, WO 99/47545 and WO 01/90063.

5. The method according to claim 1, wherein the compound is:




and each stereoisomer thereof, including:




6. The method according to claim 1, wherein the compound is:


- 63 -




and each stereoisomer thereof, including:



7. The method according to claim 1, wherein the compound is:




and each stereoisomer thereof, including:




(compound 7)

8. The method according to claim 1, wherein the compound is:

- 64 -



and each stereoisomer thereof, including:



9. A method for identifying a compound for ameliorating, treating or preventing an autoinflammatory disease in a subject comprising administering a compound that inhibits ICE or a pharmaceutical composition comprising said compound and comparing the ICE inhibition in the subject before and after treatment with said compound.

10. A method for identifying a compound for ameliorating, treating, or preventing an autoinflammatory disease in a subject comprising administering a compound that inhibits ICE or a pharmaceutical composition comprising said compound and comparing a biomarker for said autoinflammatory disease in said subject before and after treatment with said compound.

11. The method according to claim 10, wherein said biomarker is selected from the group consisting of: interleukin-1 β , interleukin-6, interleukin-8, interleukin-18, serum amyloid A, C-reactive protein, erythrocyte sedimentation rate

- 65 -

(ESR), haptoglobin, TNFalpha, immunoglobins, ferritin, leukocyte counts and platelet counts and hemoglobin.

12. A pharmaceutical composition for ameliorating, treating, or preventing an autoinflammatory disease in a subject, comprising a compound that inhibits ICE and a pharmaceutically acceptable carrier.

13. The pharmaceutical composition according to claim 12 or the methods according to any one of claims 9-11, wherein the compound is as recited in any one of claims 3-8.

14. The method according to claim 1, wherein the compound is administered to the subject at a dose selected from the group consisting of:

(a) between about 300 mg to about 2,400 mg per administration;

(b) between about 600 mg to about 1,800 mg per administration; and

(c) about 900 mg per administration.

15. The method according to claim 1, wherein the compound is administered to the subject at a dose selected from the group consisting of:

(a) between about 2 mg per kg body weight to about 200 mg per kg body weight per day;

(b) between about 6 mg per kg body weight to about 100 mg per kg body weight per day; and

(c) between about 25 mg per kg body weight to about 75 mg per kg body weight per day.

16. A method for reduction of IL-1 β , IL-18 or other biomarker in a subject having an

- 66 -

autoinflammatory disease, comprising the step of administering to said subject a compound that inhibits ICE or a pharmaceutical composition comprising said compound.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C

