



US 20120015375A1

(19) **United States**

(12) **Patent Application Publication**  
**Malinowski et al.**

(10) **Pub. No.: US 2012/0015375 A1**

(43) **Pub. Date: Jan. 19, 2012**

(54) **MCM6 AND MCM7 MONOCLONAL ANTIBODIES AND METHODS FOR THEIR USE IN THE DETECTION OF CERVICAL DISEASE**

**Publication Classification**

(51) **Int. Cl.**  
**G01N 33/577** (2006.01)

(75) Inventors: **Douglas P. Malinowski**, Hillsborough, NC (US); **Timothy J. Fischer**, Raleigh, NC (US); **Adriann J. Taylor**, Durham, NC (US)

(52) **U.S. Cl.** ..... **435/7.1; 436/501**

(73) Assignee: **TriPath Imaging, Inc.**, Burlington, NC (US)

(57) **ABSTRACT**

(21) Appl. No.: **13/247,614**

(22) Filed: **Sep. 28, 2011**

Compositions and methods for diagnosing high-grade cervical disease in a patient sample are provided. The compositions include novel monoclonal antibodies, and variants and fragments thereof, that specifically bind to MCM6 or MCM7. Monoclonal antibodies having the binding characteristics of an MCM6 or MCM7 antibody of the invention are further provided. Hybridoma cell lines that produce an MCM6 or MCM7 monoclonal antibody of the invention are also disclosed herein. The compositions find use in practicing methods for diagnosing high-grade cervical disease comprising detecting overexpression of MCM6, MCM7, or both MCM6 and MCM7 in a cervical sample from a patient. Kits for practicing the methods of the invention are further provided. Polypeptides comprising the amino acid sequence for an MCM6 or an MCM7 epitope and methods of using these polypeptides in the production of antibodies are also encompassed by the present invention.

**Related U.S. Application Data**

(62) Division of application No. 12/177,481, filed on Jul. 22, 2008, now Pat. No. 8,058,404, which is a division of application No. 11/501,391, filed on Aug. 9, 2006, now Pat. No. 7,632,498.

(60) Provisional application No. 60/751,495, filed on Dec. 19, 2005.

**MCM6 AND MCM7 MONOCLONAL  
ANTIBODIES AND METHODS FOR THEIR  
USE IN THE DETECTION OF CERVICAL  
DISEASE**

**CROSS REFERENCE TO RELATED  
APPLICATIONS**

[0001] This application is a divisional of U.S. application Ser. No. 12/177,481, filed Jul. 22, 2008, which is a divisional of U.S. application Ser. No. 11/501,391, filed Aug. 9, 2006, now U.S. Pat. No. 7,632,498, issued Dec. 15, 2009, which claims the benefit of U.S. Provisional Application Ser. No. 60/751,495, filed Dec. 19, 2005, each of which is incorporated herein by reference in its entirety.

**REFERENCE TO A SEQUENCE LISTING  
SUBMITTED AS A TEXT FILE VIA EFS-WEB**

[0002] The official copy of the sequence listing is submitted concurrently with the specification as a text file via EFS-Web, in compliance with the American Standard Code for Information Interchange (ASCII), with a file name of 410688SequenceListing.txt, a creation date of Sep. 28, 2011, and a size of 47 KB. The sequence listing filed via EFS-Web is part of the specification and is hereby incorporated in its entirety by reference herein.

**FIELD OF THE INVENTION**

[0003] The invention relates to antibodies capable of binding to MCM6 or MCM7 and methods of using these antibodies, particularly in the diagnosis of cervical disease.

**BACKGROUND OF THE INVENTION**

[0004] Carcinoma of the cervix is the second most common neoplasm in women, accounting for approximately 12% of all female cancers and causing approximately 250,000 deaths per year. Baldwin et al. (2003) *Nature Reviews Cancer* 3:1-10. In many developing countries where mass screening programs are not available, the clinical problem is more serious. Cervical cancer in these countries is the number one cause of cancer deaths in women.

[0005] The majority of cases of cervical cancer represent squamous cell carcinoma, although adenocarcinoma is also seen. Cervical cancer can be prevented by population screening as it evolves through well-defined noninvasive intraepithelial stages, which can be distinguished morphologically. Williams et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:14932-14937. While it is not understood how normal cells become transformed, the concept of a continuous spectrum of histopathological change from normal, stratified epithelium through cervical intraepithelial neoplasia (CIN) to invasive cancer has been widely accepted for years. The precursor to cervical cancer is dysplasia, also known in the art as CIN or squamous intraepithelial lesions (SIL). Squamous intraepithelial abnormalities may be classified by using the three-tiered (CIN) or two-tiered (Bethesda) system. Under the Bethesda system, low-grade squamous intraepithelial lesions (LSIL), corresponding to CIN I and HPV infection, generally represent productive HPV infections with a relatively low risk of progression to invasive disease. High-grade squamous intraepithelial lesions (HSIL), corresponding to CIN II and CIN III in the three-tiered system, show a higher risk of progression to cervical cancer than do LSIL, although both LSIL and HSIL are viewed as potential precursors of malignancy.

Patient samples may also be classified as ASCUS (atypical squamous cells of unknown significance) or AGUS (atypical glandular cells of unknown significance) under this system.

[0006] A strong association of cervical cancer and infection by high-risk types of human papilloma virus (HPV), such as types 16, 18, and 31, has been established. In fact, a large body of epidemiological and molecular biological evidence has established HPV infection as a causative factor in cervical cancer. Moreover, HPV is found in 85% or more of the cases of high-grade cervical disease. However, HPV infection is very common, possibly occurring in 5-15% of women over the age of 30, but few HPV-positive women will ever develop high-grade cervical disease or cancer. The presence of HPV alone is indicative only of infection, not of high-grade cervical disease, and, therefore, testing for HPV infection alone results in many false positives. See, for example, Wright et al. (2004) *Obstet. Gynecol.* 103:304-309.

[0007] Current literature suggests that HPV infects the basal stem cells within the underlying tissue of the uterine-cervix. Differentiation of the stem cells into mature keratinocytes, with resulting migration of the cells to the stratified cervical epithelium, is associated with HPV viral replication and re-infection of cells. During this viral replication process, a number of cellular changes occur that include cell-cycle de-regulation, active proliferation, DNA replication, transcriptional activation and genomic instability (Crum (2000) *Modern Pathology* 13:243-251; Middleton et al. (2003) *J. Virol.* 77:10186-10201; Pett et al., (2004) *Cancer Res.* 64:1359-1368).

[0008] Most HPV infections are transient in nature, with the viral infection resolving itself within a 12-month period. For those individuals who develop persistent infections with one or more oncogenic subtypes of HPV, there is a risk for the development of neoplasia in comparison to patients without an HPV infection. Given the importance of HPV in the development of cervical neoplasia, the clinical detection of HPV has become an important diagnostic tool in the identification of patients at risk for cervical neoplasia development. The clinical utility of HPV-based screening for cervical disease is in its negative predictive value. An HPV negative result in combination with a history of normal Pap smears is an excellent indicator of a disease-free condition and a low risk of cervical neoplasia development during the subsequent 1-3 years. However, a positive HPV result is not diagnostic of cervical disease; rather it is an indication of infection. Although the majority of HPV infections is transient and will spontaneously clear within a 12-month period, a persistent infection with a high-risk HPV viral subtype indicates a higher risk for the development of cervical neoplasia. To supplement HPV testing, the identification of molecular markers associated with cervical neoplasia is expected to improve the clinical specificity for cervical disease diagnosis.

[0009] Cytological examination of Papanicolaou-stained cervical smears (Pap smears) currently is the method of choice for detecting cervical cancer. The Pap test is a subjective method that has remained substantially unchanged for 60 years. There are several concerns, however, regarding its performance. The reported sensitivity of a single Pap test (the proportion of disease positives that are test-positive) is low and shows wide variation (30-87%). The specificity of a single Pap test (the proportion of disease negatives that are test-negative) might be as low as 86% in a screening population and considerably lower in the ASCUS PLUS population for the determination of underlying high-grade disease. See,

Baldwin et al., supra. A significant percentage of Pap smears characterized as LSIL or CIN1 are actually positive for high-grade lesions. Furthermore, up to 10% of Pap smears are classified as ASCUS (atypical squamous cells of undetermined significance), i.e., it is not possible to make a clear categorization as normal, moderate or severe lesion, or tumor. However, experience shows that up to 10% of this ASCUS population has high-grade lesions, which are consequently overlooked. See, for example, Manos et al. (1999) *JAMA* 281:1605-1610. Therefore, molecular biomarkers that are selectively overexpressed in high-grade cervical disease and compositions for the detection of these biomarkers are needed to practice reliable methods for diagnosing high-grade cervical disease.

**[0010]** Minichromosome maintenance (MCM) proteins play an essential part in eukaryotic DNA replication. The minichromosome maintenance (MCM) proteins function in the early stages of DNA replication through loading of the prereplication complex onto DNA and functioning as a helicase to help unwind the duplex DNA during de novo synthesis of the duplicate DNA strand. Each of the MCM proteins has DNA-dependent ATPase motifs in their highly conserved central domain. Levels of MCM proteins generally increase in a variable manner as normal cells progress from G0 into the G1/S phase of the cell cycle. In the G0 phase, MCM2 and MCM5 proteins are much less abundant than are the MCM7 and MCM3 proteins. MCM6 forms a complex with MCM2, MCM4, and MCM7, which binds histone H3. In addition, the subcomplex of MCM4, MCM6, and MCM7 has helicase activity, which is mediated by the ATP-binding activity of MCM6 and the DNA-binding activity of MCM4. See, for example, Freeman et al. (1999) *Clin. Cancer Res.* 5:2121-2132; Lei et al. (2001) *J. Cell Sci.* 114:1447-1454; Ishimi et al. (2003) *Eur. J. Biochem.* 270:1089-1101, all of which are herein incorporated by reference in their entirety.

**[0011]** Early publications have shown that the MCM proteins, and in particular, MCM5, are useful for the detection of cervical disease (Williams et al. (1998) *Proc Natl Acad Sci U.S.A.* 95:14932-14937), as well as other cancers (Freeman et al. (1999) *Clin Cancer Res.* 5:2121-2132). The published literature indicates that antibodies to MCM5 are capable of detecting cervical neoplastic cells. The specificity for detection of high-grade cervical disease has not been demonstrated for MCM5 (Williams et al. (1998) *Proc Natl Acad Sci U.S.A.* 95:14932-14937). The detection of MCM5 expression is not restricted to high-grade cervical disease but is also detected in identified low-grade dysplasia and proliferative cells that have re-entered the cell cycle following infection with high-risk HPV. In addition to MCM5, other members from the MCM family, including MCM2 and MCM7 have been shown to be potentially useful markers for the detection of cervical neoplasia in tissue samples (Freeman et al. (1999) *Clin Cancer Res.* 5:2121-2132; Brake et al. (2003) *Cancer Res.* 63:8173-8180). Recent results have shown that MCM7 appears to be a specific marker for the detection of high-grade cervical disease using immunochemistry formats (Brake et al. (2003) *Cancer Res.* 63:8173-8180; Malinowski et al. (2004) *Acta Cytol.* 43:696).

**[0012]** Therefore, there is a need in the art for antibodies that are capable of detecting expression of a biomarker that is selectively overexpressed in high-grade cervical disease. Such antibodies could be used in methods for differentiating

high-grade disease from conditions that are not considered clinical disease, such as early-stage HPV infection and mild dysplasia.

#### SUMMARY OF THE INVENTION

**[0013]** Compositions and methods for diagnosing high-grade cervical disease are provided. Compositions include monoclonal antibodies capable of binding to nuclear biomarker proteins of the invention, particularly MCM proteins, more particularly MCM6 and MCM7. Antigen-binding fragments and variants of these monoclonal antibodies, hybridoma cell lines capable of producing these antibodies, and kits comprising the monoclonal antibodies of the invention are also encompassed herein.

**[0014]** The compositions of the invention find use in methods for diagnosing high-grade cervical disease. The methods comprise detecting expression of at least one nuclear biomarker, wherein overexpression of the nuclear biomarker is indicative of high-grade cervical disease. Specifically, the methods comprise using the antibodies of the invention to detect overexpression of MCM6 or MCM7 in a cervical sample.

**[0015]** Compositions of the invention further include isolated polypeptides that comprise an epitope capable of binding an MCM6 or MCM7 monoclonal antibody. These polypeptides find use in methods for producing MCM6 or MCM7 antibodies. Isolated nucleic acid molecules encoding the amino acid sequences of the MCM6 or MCM7 epitopes are also provided.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0016]** Compositions and methods for diagnosing high-grade cervical disease are provided. Compositions include monoclonal antibodies that are capable of binding to nuclear biomarker proteins that are selectively overexpressed in high-grade cervical disease, particularly MCM proteins, more particularly MCM6 and MCM7. Hybridoma cell lines that produce the monoclonal antibodies of the present invention are also disclosed. Kits comprising the monoclonal antibodies described herein are further provided. The present compositions find use in methods for diagnosing high-grade cervical disease in a patient.

**[0017]** The compositions of the invention include monoclonal antibodies that specifically bind to MCM6 or MCM7, or to a variant or fragment thereof. The amino acid and nucleotide sequences for MCM6 are set forth in SEQ ID NO:3 (Accession No. NP\_005906) and SEQ ID NO:4 (Accession No. NM\_005915), respectively. The amino acid and nucleotide sequences for MCM7 are set forth in SEQ ID NO:1 (Accession No. NP\_005907) and SEQ ID NO:2 (Accession No. NM\_005916), respectively. In particular embodiments, the MCM6 monoclonal antibody designated as 9D4.3 and the MCM7 monoclonal antibody designated as 2E6.2 are provided. A hybridoma cell line that produces MCM7 monoclonal antibody 2E6.2 was deposited with the Patent Depository of the American Type Culture Collection (ATCC), Manassas, Va., 20110-2209 on Apr. 14, 2005 and assigned Patent Deposit No. PTA-6669. A hybridoma cell line that produces MCM6 monoclonal antibody 9D4.3 was deposited with the Patent Depository of the American Type Culture Collection (ATCC), Manassas, Va., 20110-2209 on Aug. 9, 2005 and assigned Patent Deposit No. PTA-6911. These deposits will be maintained under the terms of the Budapest

Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. This deposit was made merely as a convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

**[0018]** Antibodies that have the binding characteristics of monoclonal antibody 9D4.3 and 2E6.2 are also disclosed herein. Such antibodies include, but are not limited to, antibodies that compete in competitive binding assays with these antibodies, as well as antibodies that bind to an epitope capable of binding monoclonal antibody 9D4.3 or 2E6.2. Variants and fragments of monoclonal antibody 9D4.3 and 2E6.2 that retain the ability to specifically bind to MCM6 or MCM7, respectively, are also provided. Compositions further include hybridoma cell lines that produce the monoclonal antibodies of the present invention and kits comprising at least one monoclonal antibody disclosed herein.

**[0019]** “Antibodies” and “immunoglobulins” (Igs) are glycoproteins having the same structural characteristics. While antibodies exhibit binding specificity to an antigen, immunoglobulins include both antibodies and other antibody-like molecules that lack antigen specificity. Polypeptides of the latter kind are, for example, produced at low levels by the lymph system and at increased levels by myelomas.

**[0020]** The terms “antibody” and “antibodies” broadly encompass naturally occurring forms of antibodies and recombinant antibodies such as single-chain antibodies, chimeric and humanized antibodies and multi-specific antibodies as well as fragments and derivatives of all of the foregoing, which fragments and derivatives have at least an antigenic binding site. Antibody derivatives may comprise a protein or chemical moiety conjugated to the antibody. The term “antibody” is used in the broadest sense and covers fully assembled antibodies, antibody fragments that can bind antigen (e.g., Fab', F'(ab)<sub>2</sub>, Fv, single chain antibodies, diabodies), and recombinant peptides comprising the foregoing. As used herein, “MCM6 antibody” or “MCM7 antibody” refers to any antibody that specifically binds to MCM6 (SEQ ID NO:3) or MCM7 (SEQ ID NO:1), or to a variant or fragment thereof, and includes monoclonal antibodies, polyclonal antibodies, single-chain antibodies, and fragments thereof which retain the antigen binding function of the parent antibody.

**[0021]** The MCM6 and MCM7 antibodies of the invention are optimally monoclonal antibodies. The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

**[0022]** “Native antibodies” and “native immunoglobulins” are usually heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light (L) chains and two identical heavy (H) chains. Each light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies among the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (VH) followed by a number of constant domains. Each light chain has a variable domain at one end (V<sub>L</sub>) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of

the heavy chain. Particular amino acid residues are believed to form an interface between the light and heavy-chain variable domains.

**[0023]** The term “variable” refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called complementarity determining regions (CDRs) or hypervariable regions both in the light chain and the heavy-chain variable domains. The more highly conserved portions of variable domains are called the framework (FR) regions. The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a p-sheet configuration, connected by three CDRs, which form loops connecting, and 15 in some cases forming part of, the p-sheet structure. The CDRs in each chain are held together in close proximity: by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site: of antibodies (see Kabat et al., NIH Publ. No. 91-3242, Vol. I, pages 647-669 (1991)).

**[0024]** The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

**[0025]** The term “hypervariable region” when used herein refers to the amino acid residues of an antibody which: are responsible for antigen-binding. The hypervariable region comprises amino acid residues from a “complementarily determining region” or “CDR” (i.e., residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed, Public Health Service, National Institute of Health, 25 Bethesda, Md. [1991]) and/or those residues from a “hypervariable loop” (i.e., residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 2632(H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Clothia and Lesk, J. Mol. Biol., 196:901-917 [1987]). Framework” or “FR” residues are those variable domain residues other than the hypervariable region residues as herein deemed.

**[0026]** “Antibody fragments” comprise a portion of an intact antibody, preferably the antigen-binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata et al. (1995) *Protein Eng.* 8(10):1057-1062); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments. Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, each with a single antigen-binding site, and a residual “Fc” fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

**[0027]** “Fv” is the minimum antibody fragment that contains a complete antigen recognition and binding site. In a two-chain Fv species, this region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. In a single-chain Fv species, one heavy- and one light-chain variable domain can be covalently linked by flexible peptide linker such that the light and heavy chains can associate in a “dimeric” structure analogous to that in a

two-chain Fv species. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the  $V_H$ - $V_L$  dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

**[0028]** The Fab fragment also contains the constant domain of the light chain and the first constant domain ( $C_{H1}$ ) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy-chain  $C_{H1}$  domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them.

**[0029]** Fragments of the MCM6 and MCM7 antibodies are encompassed by the invention so long as they retain the desired affinity of the full-length antibody. Thus, for example, a fragment of an MCM6 antibody will retain the ability to bind to an MCM6 antigen. Similarly, a fragment of an MCM7 antibody will retain the ability to bind to an MCM7 antigen. Such fragments are characterized by properties similar to the corresponding full-length antibody, that is, the fragments will specifically bind MCM6 or MCM7. Such fragments are referred to herein as "antigen-binding" fragments.

**[0030]** Suitable antigen-binding fragments of an antibody comprise a portion of a full-length antibody, generally the antigen-binding or variable region thereof. Examples of antibody fragments include, but are not limited to, Fab, F(ab')<sub>2</sub>, and Fv fragments and single-chain antibody molecules. By "Fab" is intended a monovalent antigen-binding fragment of an immunoglobulin that is composed of the light chain and part of the heavy chain. By F(ab')<sub>2</sub> is intended a bivalent antigen-binding fragment of an immunoglobulin that contains both light chains and part of both heavy chains. By "single-chain Fv" or "sFv" antibody fragments is intended fragments comprising the  $V_H$  and  $V_L$  domains of an antibody, wherein these domains are present in a single polypeptide chain. See, for example, U.S. Pat. Nos. 4,946,778, 5,260,203, 5,455,030, and 5,856,456, herein incorporated by reference. Generally, the Fv polypeptide further comprises a polypeptide linker between the  $V_H$  and  $V_L$  domains that enables the sFv to form the desired structure for antigen binding. For a review of sFv see Pluckthun (1994) in *The Pharmacology of Monoclonal Antibodies*, Vol. 113, ed. Rosenburg and Moore (Springer-Verlag, New York), pp. 269-315.

**[0031]** Antibodies or antibody fragments can be isolated from antibody phage libraries generated using the techniques described in, for example, McCafferty et al. (1990) *Nature* 348:552-554 (1990) and U.S. Pat. No. 5,514,548. Clackson et al. (1991) *Nature* 352:624-628 and Marks et al. (1991) *J. Mol. Biol.* 222:581-597 describe the isolation of murine and human antibodies, respectively, using phage libraries. Subsequent publications describe the production of high affinity (nM range) human antibodies by chain shuffling (Marks et al. (1992) *Bio/Technology* 10:779-783), as well as combinatorial infection and in vivo recombination as a strategy for constructing very large phage libraries (Waterhouse et al. (1993) *Nucleic. Acids Res.* 21:2265-2266). Thus, these techniques

are viable alternatives to traditional monoclonal antibody hybridoma techniques for isolation of monoclonal antibodies.

**[0032]** Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al. (1992) *Journal of Biochemical and Biophysical Methods* 24:107-117 (1992) and Brennan et al. (1985) *Science* 229:81). However, these fragments can now be produced directly by recombinant host cells. For example, the antibody fragments can be isolated from the antibody phage libraries discussed above. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')<sub>2</sub> fragments (Carter et al. (1992) *Bio/Technology* 10:163-167). According to another approach, F(ab')<sub>2</sub> fragments can be isolated directly from recombinant host cell culture. Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

**[0033]** Preferably antibodies of the invention are monoclonal in nature. As indicated above, "monoclonal antibody" is intended an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. The term is not limited regarding the species or source of the antibody. The term encompasses whole immunoglobulins as well as fragments such as Fab, F(ab')<sub>2</sub>, Fv, and others which retain the antigen binding function of the antibody. Monoclonal antibodies are highly specific, being directed against a single antigenic site, i.e., a particular epitope within the MCM6 or MCM7 protein, as defined herein below. Furthermore, in contrast to conventional (polyclonal) antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler et al. (1975) *Nature* 256:495, or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in, for example, Clackson et al. (1991) *Nature* 352:624-628; Marks et al. (1991) *J. Mol. Biol.* 222:581-597; and U.S. Pat. No. 5,514,548.

**[0034]** Monoclonal antibodies can be prepared using the method of Kohler et al. (1975) *Nature* 256:495-496, or a modification thereof. Typically, a mouse is immunized with a solution containing an antigen. Immunization can be performed by mixing or emulsifying the antigen-containing solution in saline, preferably in an adjuvant such as Freund's complete adjuvant, and injecting the mixture or emulsion parenterally. Any method of immunization known in the art may be used to obtain the monoclonal antibodies of the invention. After immunization of the animal, the spleen (and optionally, several large lymph nodes) are removed and dissociated into single cells. The spleen cells may be screened by applying a cell suspension to a plate or well coated with the antigen of interest. The B cells expressing membrane bound immunoglobulin specific for the antigen (i.e., antibody-pro-

ducing cells) bind to the plate and are not rinsed away. Resulting B cells, or all dissociated spleen cells, are then induced to fuse with myeloma cells to form monoclonal antibody-producing hybridomas, and are cultured in a selective medium. The resulting cells are plated by serial dilution and are assayed for the production of antibodies that specifically bind the antigen of interest (and that do not bind to unrelated antigens). The selected monoclonal antibody (mAb)-secreting hybridomas are then cultured either in vitro (e.g., in tissue culture bottles or hollow fiber reactors), or in vivo (as ascites in mice). Monoclonal antibodies can also be produced using Repetitive Immunizations Multiple Sites technology (RIMMS). See, for example, Kilpatrick et al. (1997) *Hybridoma* 16(4):381-389; Wring et al. (1999) *J. Pharm. Biomed. Anal.* 19(5):695-707; and Bynum et al. (1999) *Hybridoma* 18(5):407-411, all of which are herein incorporated by reference in their entirety.

**[0035]** As an alternative to the use of hybridomas, antibody can be produced in a cell line such as a CHO cell line, as disclosed in U.S. Pat. Nos. 5,545,403; 5,545,405; and 5,998,144; incorporated herein by reference. Briefly the cell line is transfected with vectors capable of expressing a light chain and a heavy chain, respectively. By transfecting the two proteins on separate vectors, chimeric antibodies can be produced. Another advantage is the correct glycosylation of the antibody. A monoclonal antibody can also be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with a biomarker protein to thereby isolate immunoglobulin library members that bind the biomarker protein. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP I Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, U.S. Pat. No. 5,223,409; PCT Publication Nos. WO 92/18619; WO 91/17271; WO 92/20791; WO 92/15679; 93/01288; WO 92/01047; 92/09690; and 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J.* 12:725-734.

**[0036]** In some aspects of the invention, antibodies may be selected on the basis of desirable staining of cytological, rather than histological, samples. That is, in particular embodiments the antibodies are selected with the end sample type (e.g., cytology preparations) in mind and for binding specificity. Antibodies directed to specific biomarkers of interest, such as MCM6 or MCM7, are selected and purified via a multi-step screening process. Such methods for antibody selection are described in pending U.S. application Ser. No. 11/087,227, entitled "Methods and Compositions for the Detection of Cervical Disease," filed Mar. 23, 2005, which is herein incorporated by reference in its entirety.

**[0037]** Antibodies having the binding characteristics of a monoclonal antibody of the invention are also provided. "Binding characteristics" or "binding specificity" when used in reference to an antibody means that the antibody recognizes the same or similar antigenic epitope as a comparison antibody. Examples of such antibodies include, for example, an antibody that competes with a monoclonal antibody of the invention in a competitive binding assay. One of skill in the art

could determine whether an antibody competitively interferes with another antibody using standard methods.

**[0038]** By "epitope" is intended the part of an antigenic molecule to which an antibody is produced and to which the antibody will bind. An "MCM6 epitope" comprises the part of the MCM6 protein to which an MCM6 monoclonal antibody binds. An "MCM7 epitope" comprises the part of the MCM7 protein to which an MCM7 monoclonal antibody binds. Epitopes can comprise linear amino acid residues (i.e., residues within the epitope are arranged sequentially one after another in a linear fashion), nonlinear amino acid residues (referred to herein as "nonlinear epitopes"; these epitopes are not arranged sequentially), or both linear and nonlinear amino acid residues. Typically epitopes are short amino acid sequences, e.g. about five amino acids in length. Systematic techniques for identifying epitopes are known in the art and are described, for example, in U.S. Pat. No. 4,708,871 and in the examples set forth below. Briefly, in one method, a set of overlapping oligopeptides derived from the antigen may be synthesized and bound to a solid phase array of pins, with a unique oligopeptide on each pin. The array of pins may comprise a 96-well microtiter plate, permitting one to assay all 96 oligopeptides simultaneously, e.g., for binding to a biomarker-specific monoclonal antibody. Alternatively, phage display peptide library kits (New England BioLabs) are currently commercially available for epitope mapping. Using these methods, the binding affinity for every possible subset of consecutive amino acids may be determined in order to identify the epitope that a given antibody binds. Epitopes may also be identified by inference when epitope length peptide sequences are used to immunize animals from which antibodies are obtained.

**[0039]** The invention also encompasses isolated polypeptides comprising an epitope for binding an MCM6 or MCM7 monoclonal antibody. These polypeptides correspond to a portion of the antigen (i.e., MCM6 or MCM7) that binds to a monoclonal antibody. Such polypeptides find use in methods for producing antibodies that bind selectively to MCM6 or MCM7. The ability of a polypeptide to be used in the production of antibodies is referred to herein as "antigenic activity." For example, the amino acid sequence set forth in SEQ ID NO:5 (corresponding to residues 760-772 in the MCM6 amino acid sequence set forth in SEQ ID NO:3) comprise an epitope recognized by an MCM6 monoclonal antibody, more particularly monoclonal antibody 9D4.3. The amino acid sequence set forth in SEQ ID NO:6 (corresponding to residues 127-138 in the MCM7 amino acid sequence set forth in SEQ ID NO:1) comprise an epitope recognized by an MCM7 monoclonal antibody, more particularly monoclonal antibody 2E6.2. Variants and fragments of the MCM6 and MCM7 epitope sequences set forth in SEQ ID NOs:5 and 6 that retain the antigenic activity of the original polypeptide are also provided. The invention further includes isolated nucleic acid molecules that encode polypeptides that comprise MCM6 or MCM7 epitopes, and variants and fragments thereof.

**[0040]** The polypeptides of the invention comprising MCM6 or MCM7 epitopes can be used in methods for producing monoclonal antibodies that specifically bind to MCM6 or MCM7, as described herein above. Such polypeptides can also be used in the production of polyclonal MCM6 or MCM7 antibodies. For example, polyclonal antibodies can be prepared by immunizing a suitable subject (e.g., rabbit, goat, mouse, or other mammal) with a polypeptide compris-

ing an MCM6 or MCM7 epitope (i.e., an immunogen). The antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized biomarker protein. At an appropriate time after immunization, e.g., when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497, the human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985) in *Monoclonal Antibodies and Cancer Therapy*, ed. Reisfeld and Sell (Alan R. Liss, Inc., New York, N.Y.), pp. 77-96) or trioma techniques. The technology for producing hybridomas is well known (see generally Coligan et al., eds. (1994) *Current Protocols in Immunology* (John Wiley & Sons, Inc., New York, N.Y.); Galfre et al. (1977) *Nature* 266:550-52; Kenneth (1980) in *Monoclonal Antibodies: A New Dimension In Biological Analyses* (Plenum Publishing Corp., NY; and Lerner (1981) *Yale J. Biol. Med.*, 54:387-402).

**[0041]** Amino acid sequence variants of a monoclonal antibody or a polypeptide comprising an MCM6 or MCM7 epitope described herein are also encompassed by the present invention. Variants can be prepared by mutations in the cloned DNA sequence encoding the antibody of interest. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York); Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel et al. (1987) *Methods Enzymol.* 154:367-382; Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor, N.Y.); U.S. Pat. No. 4,873,192; and the references cited therein; herein incorporated by reference. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the polypeptide of interest may be found in the model of Dayhoff et al. (1978) in *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred. Examples of conservative substitutions include, but are not limited to, Gly $\leftrightarrow$ Ala, Val $\leftrightarrow$ Ile $\leftrightarrow$ Leu, Asp $\leftrightarrow$ Glu, Lys $\leftrightarrow$ Arg, Asn $\leftrightarrow$ Gln, and Phe $\leftrightarrow$ Trp $\leftrightarrow$ Tyr.

**[0042]** In constructing variants of the polypeptide of interest, modifications are made such that variants continue to possess the desired activity, i.e., similar binding affinity to the biomarker. Obviously, any mutations made in the DNA encoding the variant polypeptide must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See EP Patent Application Publication No. 75,444.

**[0043]** Preferably, variants of a reference polypeptide have amino acid sequences that have at least 70% or 75% sequence identity, preferably at least 80% or 85% sequence identity, more preferably at least 90%, 91%, 92%, 93%, 94% or 95% sequence identity to the amino acid sequence for the reference antibody molecule, or to a shorter portion of the reference antibody molecule. More preferably, the molecules share at least 96%, 97%, 98% or 99% sequence identity. For purposes of the present invention, percent sequence identity is determined using the Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty

of 12 and a gap extension penalty of 2, BLOSUM matrix of 62. The Smith-Waterman homology search algorithm is taught in Smith and Waterman (1981) *Adv. Appl. Math.* 2:482-489. A variant may, for example, differ from the reference antibody by as few as 1 to 15 amino acid residues, as few as 1 to 10 amino acid residues, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

**[0044]** With respect to optimal alignment of two amino acid sequences, the contiguous segment of the variant amino acid sequence may have additional amino acid residues or deleted amino acid residues with respect to the reference amino acid sequence. The contiguous segment used for comparison to the reference amino acid sequence will include at least 20 contiguous amino acid residues, and may be 30, 40, 50, or more amino acid residues. Corrections for sequence identity associated with conservative residue substitutions or gaps can be made (see Smith-Waterman homology search algorithm).

**[0045]** The MCM6 and MCM7 monoclonal antibodies of the invention may be labeled with a detectable substance as described below to facilitate biomarker protein detection in the sample. Such antibodies find use in practicing the methods of the invention. The antibodies and antibody fragments of the invention can be coupled to a detectable substance to facilitate detection of antibody binding. The word "label" when used herein refers to a detectable compound or composition that is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable. Examples of detectable substances for purposes of labeling antibodies include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$ , or  $^3\text{H}$ .

**[0046]** Kits comprising at least one MCM6 or one MCM7 monoclonal antibody of the invention are further provided. By "kit" is intended any manufacture (e.g., a package or a container) comprising at least one reagent, i.e., an antibody, for specifically detecting the expression of MCM6 or MCM7. The kit may be promoted, distributed, or sold as a unit for performing the methods of the present invention. Additionally, the kits may contain a package insert describing the kit and methods for its use.

**[0047]** Kits of the invention generally comprise at least one monoclonal antibody directed to MCM6 or MCM7, chemicals for the detection of antibody binding, a counterstain, and, optionally, a bluing agent to facilitate identification of positive staining cells. Any chemicals that detect antigen-antibody binding may be used in the kits of the invention. In some embodiments, the detection chemicals comprise a labeled polymer conjugated to a secondary antibody. For example, a secondary antibody that is conjugated to an enzyme that catalyzes the deposition of a chromogen at the antigen-anti-

body binding site may be provided. Such enzymes and techniques for using them in the detection of antibody binding are well known in the art. In one embodiment, the kit comprises a secondary antibody that is conjugated to an HRP-labeled polymer. Chromogens compatible with the conjugated enzyme (e.g., DAB in the case of an HRP-labeled secondary antibody) and solutions, such as hydrogen peroxide, for blocking non-specific staining may be further provided. In other embodiments, antibody binding to a biomarker protein is detected through the use of a mouse probe reagent that binds to monoclonal antibodies, followed by addition of a dextran polymer conjugated with HRP that binds to the mouse probe reagent. Such detection reagents are commercially available from, for example, Biocare Medical.

**[0048]** The kits of the present invention may further comprise a peroxidase blocking reagent (e.g., hydrogen peroxide), a protein blocking reagent (e.g., purified casein), and a counterstain (e.g., hematoxylin). A bluing agent (e.g., ammonium hydroxide or TBS, pH 7.4, with Tween-20 and sodium azide) may be further provided in the kit to facilitate detection of positive staining cells. Kits may also comprise positive and negative control samples for quality control purposes.

**[0049]** In another embodiment, the kits of the invention comprise at least two monoclonal antibodies. In certain aspects of the invention, the kits comprise an MCM6 and an MCM7 antibody, more particularly the MCM6 monoclonal antibody 9D4.3 and the MCM7 antibody 2E6.2. When multiple antibodies are present in the kit, each antibody may be provided as an individual reagent or, alternatively, as an antibody cocktail comprising all of the antibodies of interest. Furthermore, any or all of the kit reagents may be provided within containers that protect them from the external environment, such as in sealed containers. The kits of the invention are useful in the diagnosis of high-grade cervical disease and may further include reagents for Pap staining (e.g., EA50 and Orange G).

**[0050]** The compositions of the invention find use in methods for diagnosing high-grade cervical disease in a patient such as those disclosed in pending U.S. application Ser. No. 11/087,227, entitled "Methods and Compositions for the Detection of Cervical Disease," filed Mar. 23, 2005, which is herein incorporated by reference in its entirety. "Diagnosing high-grade cervical disease" is intended to include, for example, diagnosing or detecting the presence of cervical disease, monitoring the progression of the disease, and identifying or detecting cells or samples that are indicative of high-grade cervical disease. The terms diagnosing, detecting, and identifying high-grade cervical disease are used interchangeably herein. By "high-grade cervical disease" is intended those conditions classified by colposcopy as pre-malignant pathology, malignant pathology, moderate to severe dysplasia, and cervical cancer. Underlying high-grade cervical disease includes histological identification of CINII, CINIII, HSIL, carcinoma in situ, adenocarcinoma, and cancer (FIGO stages I-IV).

**[0051]** The methods of the invention comprise detecting overexpression of at least one nuclear biomarker that is selectively overexpressed in high-grade cervical disease. By "nuclear biomarker" is intended any gene or protein that is predominantly expressed in the nucleus of the cell. A nuclear biomarker may be expressed to a lesser degree in other parts of the cell. By "selectively overexpressed in high-grade cervical disease" is intended that the nuclear biomarker of interest is overexpressed in high-grade cervical disease but is not

overexpressed in conditions classified as LSIL, CINI, HPV-infected samples without any dysplasia present, immature metaplastic cells, and other conditions that are not considered to be clinical disease. Thus, detection of the nuclear biomarkers of the invention permits the differentiation of samples indicative of underlying high-grade cervical disease from samples that are indicative of benign proliferation, early-stage HPV infection, or mild dysplasia. Nuclear biomarkers of particular interest include MCM proteins, particularly MCM6 and MCM7.

**[0052]** In a particular aspect of the invention, the methods comprise obtaining a cervical sample from a patient, contacting the sample with at least one MCM6 or MCM7 monoclonal antibody of the invention, and detecting binding of the antibody to the MCM protein. In other embodiments, the sample is contacted with at least two monoclonal antibodies, a first antibody that specifically binds to MCM6, particularly monoclonal antibody 9D4.3, and a second antibody that specifically binds to MCM7, particularly monoclonal antibody 2E6.2. Techniques for detecting antibody binding are well known in the art. Antibody binding to a biomarker of interest may be detected through the use of chemical reagents that generate a detectable signal that corresponds to the level of antibody binding and, accordingly, to the level of biomarker protein expression. Any method for detecting antibody-antigen binding may be used to practice the methods of the invention.

**[0053]** As used herein, "cervical sample" refers to any sampling of cells, tissues, or bodily fluids from the cervix in which expression of a biomarker can be detected. Examples of such body samples include but are not limited to gynecological fluids, biopsies, and smears. Cervical samples may be obtained from a patient by a variety of techniques including, for example, by scraping or swabbing an area or by using a needle to aspirate bodily fluids. Methods for collecting cervical samples are well known in the art. In particular embodiments, the cervical sample comprises cervical cells, particularly in a liquid-based preparation. In one embodiment, cervical samples are collected according to liquid-based cytology specimen preparation guidelines such as, for example, the liquid based Pap test, SUREPATH® (TriPath Imaging, Inc.) or the modified Pap test, THINPREP® preparation (CYTYC, Inc.). Cervical samples may be transferred to a glass slide for viewing under magnification. Fixative and staining solutions may be applied to the cells on the glass slide for preserving the specimen and for facilitating examination. In one embodiment the cervical sample will be collected and processed to provide a monolayer sample, as set forth in U.S. Pat. No. 5,346,831, herein incorporated by reference.

**[0054]** One of skill in the art will appreciate that any or all of the steps in the methods of the invention could be implemented by personnel in a manual or automated fashion. Thus, the steps of cervical sample preparation, antibody, and detection of antibody binding may be automated. The methods of the invention may also be combined with conventional Pap staining techniques to permit a more accurate diagnosis of high-grade cervical disease.

**[0055]** The following examples are offered by way of illustration and not by way of limitation:

## EXPERIMENTAL

### Example 1

#### Production of Mouse Monoclonal Antibodies to MCM7

**[0056]** Mouse monoclonal antibodies specific for MCM7 were generated. The antigen (an immunogenic polypeptide)



was a recombinant hexahistidine-tagged N-terminal fragment of the MCM7 protein. The antigen was expressed using a baculovirus expression system in Tni cells. Specifically, the coding sequence for the hexahistidine-tagged MCM7 N-terminal fragment (SEQ ID NO:7) was cloned into the pFast-Bac1 plasmid (Invitrogen) for expression in Tni cells. Methods for producing recombinant proteins using baculovirus expression systems are well known in the art. The tagged MCM7 fragment was purified using a chelating agarose charged with Ni+2 ions (Ni-NTA from Qiagen) and used as an immunogen. The amino acid sequence of the immunogenic MCM7 N-terminal polypeptide fragment is provided in SEQ ID NO:8.

**[0057]** Mouse immunizations and hybridoma fusions were performed essentially as described in Kohler et al. (1975) *Nature* 256:495-496. Mice were immunized with the immunogenic tagged-MCM7 fragment in solution. Antibody-producing cells were isolated from the immunized mice and fused with myeloma cells to form monoclonal antibody-producing hybridomas. The hybridomas were cultured in a selective medium. The resulting cells were plated by serial dilution and assayed for the production of antibodies that specifically bind MCM7 (and that do not bind to unrelated antigens). To confirm that the monoclonal antibodies of interest reacted with the MCM7 protein only and not with the hexahistidine tag, selected hybridomas were screened against an MCM7-FLAG-tagged protein. Selected monoclonal antibody (mAb)-secreting hybridomas were then cultured.

**[0058]** Antibodies were purified from the culture media supernatants of "exhausted" hybridoma cells (i.e., cells grown until viability drops to between 0-15%) using recombinant Protein A-coated resin (STREAMLINE®, Amersham, Inc.). Antibodies were eluted using low pH followed by immediate neutralization of pH. Fractions with significant absorbances at 280 nM were pooled. The resultant pool was dialyzed against PBS. Purified antibodies were subjected to further characterization. MCM7 monoclonal antibody 2E6.2 was determined to be an IgG<sub>1</sub> isotype. Details of the epitope mapping of this antibody are described below.

#### Example 2

##### Production of Mouse Monoclonal Antibodies to MCM6

**[0059]** Mouse monoclonal antibodies specific for MCM6 were generated. The antigen (an immunogenic polypeptide) was a recombinant FLAG-tagged MCM6 protein. The antigen was expressed using a proprietary expression vector from Cell & Molecular Technology, Inc. in HEK293 cells or alternatively expressed using a baculovirus expression system in Tni cells. The coding sequence for the FLAG-tagged MCM6 is set forth in SEQ ID NO:9. FLAG-tagged MCM6 was purified from cell lysates using the anti-FLAG M2 Affinity Gel matrix and the FLAG peptide for elution (Sigma Chemical Co., St. Louis, Mo.). The FLAG-tagged MCM6 protein used as an immunogen. The amino acid sequence of the immunogenic FLAG-tagged MCM6 polypeptide is provided in SEQ ID NO:10.

**[0060]** Mouse immunizations and lymphocyte fusion were performed by RIMMS technology, essentially as described in Kilpatrick et al. (1997) *Hybridoma* 16(4):381-389. Mice were immunized with the immunogenic FLAG-tagged-MCM6. Primary screening of uncloned hybridoma supernatants was performed using recombinant MCM6 protein. Sec-

ondary screening and screening of cloned hybridoma supernatants was performed using a separate batch of recombinant MCM6 protein. Selected monoclonal antibody (mAb)-secreting hybridomas were then cultured.

**[0061]** Antibodies were purified from the culture media supernatants of "exhausted" hybridoma cells (i.e., cells grown until viability drops to between 0-15%) using recombinant Protein A-coated resin (STREAMLINE®, Amersham, Inc.). Antibodies were eluted using low pH followed by immediate neutralization of pH. Fractions with significant absorbances at 280 nM were pooled. The resultant pool was dialyzed against PBS. Purified antibodies were subjected to further characterization. MCM6 monoclonal antibody 9D4.3 was determined to be an IgG<sub>2a</sub> isotype. Details of the epitope mapping of this antibody are described below.

#### Example 3

##### Isolation of Monoclonal Antibodies from Hybridoma Cells

**[0062]** The following procedure is used to isolate monoclonal antibodies from hybridoma cells:

##### Media Preparation

**[0063]** To a sterile 1,000 ml storage bottle, add 100 ml Hyclone Fetal Bovine Serum (FBS).

**[0064]** Add 10 ml of MEM Non-Essential Amino Acids Solution.

**[0065]** Add 10 ml of Penicillin-Streptomycin-L-Glutamine Solution.

**[0066]** QS to approximately 1000 ml with ExCell 610-HSF media.

**[0067]** Place sterile cap on bottle and secure tightly. Swirl gently to mix.

**[0068]** Connect a 1000 ml sterile acetate vacuum filter unit (0.2 µm) to a vacuum pump system.

**[0069]** Gently pour approximately half of the media solution into sterile acetate vacuum filter unit and turn on the vacuum.

**[0070]** Once the first half of the media has been filtered, pour the remaining media into the filter unit and continue filtering.

**[0071]** After all the media has been filtered, disconnect the vacuum hose from the vacuum filter unit and turn off the vacuum pump. Remove the receiver portion of the filter unit from the filter bottle. Place a new sterile bottle cap on the bottle.

**[0072]** Store at 2° C. to 10° C. Protect from light.

##### Initial Hybridoma Cell Culture

**[0073]** Thaw vial of stock hybridoma frozen culture in a pre-warmed 37° C. H<sub>2</sub>O bath.

**[0074]** Spray the outside of the freeze vial with 70% ethanol.

**[0075]** Move the thawed vial into the Biological Safety Cabinet.

**[0076]** Remove the cells from the freeze vial and transfer the cells to a 15 ml centrifuge tube.

**[0077]** Add 7 ml of cell culture media drop-wise to the 15 ml centrifuge tube containing the thawed cells.

**[0078]** Centrifuge the 15 ml centrifuge tube containing the thawed cells and culture media for 5 minutes at 200 g force.

- [0079] While the cells are in the centrifuge, add 45 ml of cell culture media to a sterile T-225 flask.
- [0080] After centrifugation, visually inspect the tube for the presence of a cell pellet.
- [0081] Remove the media from the centrifuge tube being careful not to dislodge the cell pellet. Note: If the cell pellet is disturbed, repeat the centrifugation step.
- [0082] Add 5 ml of cell culture media to the 15 ml centrifuge tube containing the pelleted cells. Pipette to resuspend the cell pellet into the media.
- [0083] Transfer the entire contents of the resuspended cells and culture media into the T-225 flask containing the 45 ml of media.
- [0084] Cap the T-225 flask.
- [0085] Observe for presence of intact cells under the microscope. Place the T-225 flask immediately into a CO<sub>2</sub> incubator and allow the cells to incubate overnight.

#### Expansion of Hybridoma Cell Line

- [0086] Continue to monitor the cell culture for viability, concentration, and presence of contamination.
- [0087] Monitor and adjust the cell suspension from the initial T-225 flask until the concentration is approximately 600,000 cells/ml to 800,000 cells/ml and a total of 200 to 250 ml of media.
- [0088] Dislodge cells and add additional media as needed to meet minimum cell density requirements. Divide and transfer cell suspension into one new sterile T-225 flask. Place the 2×T-225 flasks into the CO<sub>2</sub> incubator.
- [0089] Monitor the cells from the 2×T-225 flasks until the concentration is approximately 600,000 cells/ml to 800,000 cells/ml, and a total of between 200 to 250 ml of media for each flask.
- [0090] Dislodge cells and add additional media as needed to meet minimum cell density requirements. Divide and transfer the cell suspensions into 2 additional new sterile T-225 flasks for a total of 4×T-225 flasks. Return all flasks to the CO<sub>2</sub> incubator.
- [0091] Monitor the cells, and adjust volume in the 4×T-225 flasks until the cell concentration is approximately 600,000 cells/ml to 800,000 cells/ml with a total volume of approximately 250 ml per T-225 flask (or approximately 1000 ml total).
- [0092] Continue to monitor the cells from the 4×T-225 flasks until the cells have grown to exhaustion, with a final viability of 0%-15%. The cell culture supernatant is now ready for the Clarification Process.

#### Clarification of Supernatant

- [0093] Turn on the tabletop centrifuge. Place the 500 ml tube adapters into the rotor buckets, close the lid and set the temperature to 4° C. (+/-) 4° C.
- [0094] Using aseptic technique, pour the media from all four of the now exhausted T-225 flasks into 2×500 ml conical centrifuge tubes.
- [0095] Make sure the 2×500 ml tubes are balanced. Transfer supernatant from one tube to the other as necessary to balance them.
- [0096] Centrifuge the exhausted supernatant at 1350 g (+/-40 g) for 15 minutes at 2° C. to 10° C.

- [0097] After centrifugation is complete, aseptically decant the supernatant into a sterile 1000 ml storage bottle and secure with a sterile cap.
- [0098] Aseptically transfer 1 ml to the microfuge tube. Store microfuge tube with sample at 2° C. to 10° C. (Protect from light).
- [0099] The clarified supernatant sample is ready for IgG evaluation using the antibody assay, EASY-TITER® Assay.

#### Buffer Preparation

##### Binding Buffer:

- [0100] Add approximately 600 ml of DI H<sub>2</sub>O to a clean beaker.
- [0101] Add 77.28 ml of Boric Acid solution (4% WN). Stir at room temperature with a clean stir bar.
- [0102] Weigh out 233.76 g of Sodium Chloride and place into the solution while continuing to stir.
- [0103] Bring solution up to approximately 950 ml with DI H<sub>2</sub>O and continue to stir.
- [0104] When the Sodium Chloride has dissolved and the solution is clear, adjust the pH to 9.0±0.2 with Sodium Hydroxide.
- [0105] Remove the solution to a clean 1000 ml graduated cylinder and QS to 1000 ml with DI H<sub>2</sub>O.
- [0106] Transfer the completed buffer to an appropriate storage bottle. This buffer may be stored for up to 7 days before use.
- [0107] Repeat this entire process to prepare an additional 0.2 liters to 1.0 liter of Binding Buffer.

##### Elution Buffer

- [0108] Weigh out 1.725 g of sodium phosphate, monobasic and place into a clean 250 ml beaker with a clean stir bar.
- [0109] Weigh out 3.676 g of sodium citrate and place into the same clean 250 ml beaker.
- [0110] Add approximately 175 ml of DI H<sub>2</sub>O and stir at room temperature until dissolved.
- [0111] Weigh out 4.38 g of Sodium Chloride and place into the solution while continuing to stir.
- [0112] Bring solution up to approximately 225 ml with DI H<sub>2</sub>O and continue to stir.
- [0113] When the Sodium Chloride has dissolved and the solution is clear, adjust the pH to 3.5±0.2 with Hydrochloric Acid.
- [0114] Remove the solution to a clean 250 ml graduated cylinder and QS to 250 ml with DI H<sub>2</sub>O.
- [0115] Connect a 500 ml sterile acetate vacuum filter unit (0.2 μm) to a vacuum pump system and filter sterilize the solution.
- [0116] Remove the filter and close the container with a sterile cap.

##### Antibody Adsorption

- [0117] Pour the Clarified Supernatant (~1 L) into a clean 4000 ml plastic beaker with a clean stir bar.
- [0118] Add an approximately equal amount (~1 L) of the Binding Buffer to the clean 4000 ml plastic beaker containing the clarified supernatant. Add a clean stir bar.
- [0119] Cover the beaker with clean plastic wrap and label "Antibody Binding."
- [0120] Calculate the approximate amount of STREAMLINE® Protein A that will be needed using the data in Table 1.

TABLE 1

Volume of Protein A Resin Required	
Quantity IgG ( $\mu\text{g/ml}$ ) in Supernatant	Volume of Protein A Resin Required in Milliliters (ml)
>180- $\leq$ 200	12.0
>160- $\leq$ 180	11.0
>140- $\leq$ 160	10.0
>120- $\leq$ 140	9.0
>100- $\leq$ 120	8.0
>80- $\leq$ 100	7.0
>60- $\leq$ 80	6.0
>40- $\leq$ 60	4.5
>20- $\leq$ 40	3.5
$\leq$ 20	2.0

**[0121]** Secure a clean Disposable Column and stopcock assembly to a ring stand and clamp. Close the stopcock.

**[0122]** Mix appropriate amount of STREAMLINE Protein A beads by inverting the bottle several times. Withdraw the required volume and place into the Disposable Column.

**[0123]** Wash the STREAMLINE Protein A beads with 10 ml of DI H<sub>2</sub>O. Open the stopcock and allow the DI H<sub>2</sub>O to drain. Close the stopcock. Repeat with an additional 10 ml of DI H<sub>2</sub>O.

**[0124]** Wash the STREAMLINE Protein A beads with 10 ml of Binding Buffer. Open the stopcock and allow the Binding Buffer to drain. Close the stopcock. Repeat with an additional 10 ml of Binding Buffer.

**[0125]** Resuspend the STREAMLINE Protein A beads in ~10 ml of the Clarified Supernatant and Binding Buffer solution (from the 4000 ml beaker) and transfer the beads into the 4000 ml beaker containing the Clarified Supernatant and Binding Buffer solution. Repeat as required to transfer any remaining beads. When completed, discard the column and stopcock.

**[0126]** Allow the mixture to mix vigorously at 2° C. to 10° C. for approximately 18 hours.

**[0127]** When mixing is complete, turn off the stir plate and remove the "Antibody Binding" beaker with the buffered supernatant and bead suspension back to the lab bench area. Allow the STREAMLINE Protein A beads to settle to the bottom of the beaker (approximately 5 minutes).

**[0128]** Secure a clean Disposable Column and stopcock assembly to a ring stand and clamp. Close the stopcock.

**[0129]** Label a clean, 250 ml bottle or suitable container "Column Wash-Post Binding."

**[0130]** Label a clean plastic beaker "Supernatant-Post Binding."

**[0131]** Decant the supernatant from the 4000 ml beaker into the clean, labeled, 2 liter plastic beaker, leaving the beads in the bottom of the 4000 ml beaker. Cover the 2000 ml beaker containing the "Supernatant-Post Binding" solution with clean plastic wrap and store at 2° C. to 10° C.

**[0132]** Add approximately 15 ml of Binding Buffer into the decanted 4000 ml "Antibody Binding" beaker. Resuspend the STREAMLINE Protein A beads and transfer them to the column. Open the stopcock and allow the Binding Buffer to drain into the "Column Wash-Post binding" container. Close the stopcock when drained.

**[0133]** Transfer any remaining STREAMLINE Protein A beads in the "Antibody Binding" beaker by adding

additional Binding Buffer, mixing, and transferring to the column as in the preceding steps. Close the stopcock when drained.

**[0134]** Calculate the approximate amount of Binding Buffer needed to wash the STREAMLINE Protein A beads in the column using the data in Table 2.

TABLE 2

Binding Buffer Volume for Column Wash	
Quantity IgG ( $\mu\text{g/ml}$ ) in Supernatant	Volume of Binding Buffer Required in Milliliters (ml)
>180- $\leq$ 200	5 column washes total with 15.0 ml each
>160- $\leq$ 180	5 column washes total with 15.0 ml each
>140- $\leq$ 160	5 column washes total with 12.5 ml each
>120- $\leq$ 140	5 column washes total with 12.5 ml each
>100- $\leq$ 120	5 column washes total with 12.5 ml each
>80- $\leq$ 100	5 column washes total with 10.0 ml each
>60- $\leq$ 80	5 column washes total with 10.0 ml each
>40- $\leq$ 60	5 column washes total with 7.5 ml each
>20- $\leq$ 40	5 column washes total with 5.0 ml each
$\leq$ 20	5 column washes total with 5.0 ml each

**[0135]** Wash the STREAMLINE Protein A beads in the column with the appropriate volume of Binding Buffer for the appropriate number of washes, continuing to collect the effluent into the "Column Wash-Post Binding" container.

**[0136]** When completed, close the stopcock. Store the "Column Wash-Post Binding" container at 2° C. to 10° C.

**[0137]** Determine the Total Volumes of Elution Buffer and Neutralization Buffer needed to elute the STREAMLINE Protein A beads in the column from Table 3.

TABLE 3

Determination of Amount of Elution Buffer and Neutralization Buffer				
Quantity IgG ( $\mu\text{g/ml}$ ) in Supernatant	Total Volume of Elution Buffer Required (ml)	Total Volume of Neutralization Buffer Required (ml)	Volume of Elution Buffer Required per fraction (ml)	Volume of Neutralization Buffer Required per fraction (ml)
>180- $\leq$ 200	72	7.2	12	1.2
>160- $\leq$ 180	66	6.6	11	1.1
>140- $\leq$ 160	60	6.0	10	1.0
>120- $\leq$ 140	54	5.4	9	0.9
>100- $\leq$ 120	48	4.8	8	0.8
>80- $\leq$ 100	42	4.2	7	0.7
>60- $\leq$ 80	36	3.6	6	0.6
>40- $\leq$ 60	27	2.7	4.5	0.45
>20- $\leq$ 40	21	2.1	3.5	0.35
$\leq$ 20	12	1.2	2	0.2

**[0138]** Label 9 sterile conical centrifuge tubes "Eluted Antibody", Fraction # (1 through 9).

**[0139]** Place the appropriate volume of Neutralization Buffer required per fraction (as determined from Table

“C” above) into each of the 9 “Eluted Antibody” fraction tubes and place securely under the column stopcock outlet.

**[0140]** Elute the STREAMLINE Protein A beads in the column fraction by fraction with the appropriate volume of Elution Buffer required per fraction (as determined from Table 3 above) while collecting the eluate into each of the “Eluted Antibody” tubes containing Neutralization Buffer.

**[0141]** When the elutions are complete, mix each “Eluted Antibody” fraction tube gently by swirling several times. Remove approximately 50 µl of fraction # 3 and place on a pH test paper strip to ensure that the eluate has been neutralized to an approximate pH between 6.5 to 8.5. If required, add additional Neutralizing Buffer or Elution Buffer as needed to bring pH into range.

**[0142]** When pH evaluation is completed, perform an Absorbance Scan of a sample from each fraction at 280 nm-400 nm to determine the approximate concentration of IgG in the eluate prior to proceeding to the Dialysis Process.

**[0143]** Accept fractions as part of the Eluate Pool if the A280-A400 value is  $\geq 0.200$ .

**[0144]** Reject fractions as part of the Eluate Pool if the A280-A400 value is  $< 0.200$ .

**[0145]** Label a sterile conical centrifuge tube “Eluted Antibody,” “Eluate Pool,” and combine all fractions that were Accepted as part of the pool.

**[0146]** Perform an Absorbance Scan of a sample of the Eluate Pool to determine the approximate concentration of IgG in the eluate prior to proceeding to the Dialysis Process.

**[0147]** Estimate the volume of the Eluate Pool and calculate the approximate total mgs of IgG.

**[0148]** Volume of Eluate Pool: \_\_\_\_\_ mls  $\times$  \_\_\_\_\_ IgG mg/ml = \_\_\_\_\_ Total mgs of IgG

Antibody Dialysis

**[0149]** Remove the “Eluted Antibody” tube from 2° C. to 10° C.

**[0150]** Calculate the approximate length of Dialysis Tubing that will be needed to dialyze the antibody eluate using the approximate volume of eluate and the data in Table 4.

**[0151]** Cut the appropriate length of dialysis tubing required. (multipurpose dialysis tubing, SPECTRA/POR® 2 Regenerated Cellulose Membrane, 12,000-14,000

**[0152]** Dalton Molecular Weight Cutoff (MWCO), 16 mm Diameter, Spectrum Laboratories Inc., Cat. No. 132678)

**[0153]** Hydrate the dialysis membrane tubing in 1000 ml of DIH<sub>2</sub>O for >30 minutes.

**[0154]** Calculate the approximate volume of Dialysis Buffer needed to dialyze the antibody eluate using the data in Table 5.

TABLE 5

Volume of Dialysis Buffer Required			
Quantity IgG (µg/ml) in Supernatant	Final Volume of Eluted Antibody in Milliliters (ml)	Length of Dialysis Tubing Needed (cm)	Volume of Dialysis Buffer (1 x PBS) Needed in Liters
>180-≤200	39.6 ml	63 cm	3 complete changes of 4.0 Liters
>160-≤180	36.3 ml	59 cm	3 complete changes of 3.6 Liters
>140-≤160	33.0 ml	55 cm	3 complete changes of 3.3 Liters
>120-≤140	29.7 ml	51 cm	3 complete changes of 3.0 Liters
>100-≤120	26.4 ml	47 cm	3 complete changes of 2.6 Liters
>80-≤100	23.1 ml	43 cm	3 complete changes of 2.3 Liters
>60-≤80	19.8 ml	39 cm	3 complete changes of 1.9 Liters
>40-≤60	14.85 ml	33 cm	3 complete changes of 1.5 Liters
>20-≤40	11.55 ml	29 cm	3 complete changes of 1.2 Liters
≤20	6.6 ml	23 cm	3 complete changes of 0.7 Liters

**[0155]** Place the appropriate amount of Dialysis Buffer into a suitable sized plastic beaker. Label the beaker “Dialyzed Antibody.” Add a clean stir bar and place the beaker on a stir plate inside a refrigerator or cold room at 2° C. to 10° C.

**[0156]** Rinse the dialysis tubing thoroughly in D<sub>1</sub>-H<sub>2</sub>O. Tie two end knots approximately 7 cm from one end of the dialysis tubing and secure tightly.

TABLE 4

Calculation of Length of Dialysis Tubing Needed						
Approximate Volume of Eluent (ml)	Volume/length Ratio of Dialysis Tubing	Approximate Length Needed for Eluent Sample (cm)	Head Space of 20% (cm)	Approximate Length Needed for Sample plus Headspace (cm)	Approximate Length Needed for Tie Off of Tubing (cm)	Approximate Total Length of Dialysis Tubing Needed (cm)
39.6	2	20	4	24	15	63
36.3	2	18	4	22	15	59
33.0	2	17	3	20	15	55
29.7	2	15	3	18	15	51
26.4	2	13	3	16	15	47
23.1	2	12	2	14	15	43
19.8	2	10	2	12	15	39
14.85	2	7	1	9	15	33
11.55	2	6	1	7	15	29
6.6	2	3	1	4	15	23

- [0157] Add approximately 5 ml of D<sub>1</sub>-H<sub>2</sub>O into the dialysis tubing.
- [0158] Fill the dialysis tubing with the eluted antibody from the "Eluted Antibody" collection tube.
- [0159] Tie two end knots approximately 7 cm from the remaining open end of the dialysis tubing and secure tightly. Ensure that the headspace is approximately that as derived from Table 4.
- [0160] Place the filled and closed dialysis tubing into the dialysis reservoir with the appropriate volume of 1×PBS (from Table 5).
- [0161] Cover the beaker with clean plastic wrap. Adjust the speed on the stir plate such that the dialysis sample spins freely, but is not pulled down into the vortex of the dialysate. Dialysis should take place at 2° C. to 10° C. with 3 buffer exchanges in total within a 24 hour period.

#### Antibody Filtration

- [0162] Label a sterile collection tube "Dialyzed Antibody."
- [0163] Remove the dialyzed sample tubing from the dialysis beaker. Cut the dialysis tubing open at one end and transfer the dialyzed sample into the "Dialyzed Antibody" centrifuge tube.
- [0164] Label another sterile collection tube "Dialyzed Antibody."
- [0165] Select a sterile Luer Lok syringe with adequate capacity to hold the final dialyzed volume.
- [0166] Attach a syringe filter, ACRODISC® Syringe Filter to the opening of the syringe (0.2 µm HT membrane filter, TUFFRYN® Membrane, Low Protein binding, Gelman Laboratories, Cat. No. 4192). Remove the plunger from the syringe and while holding the syringe upright, transfer the dialyzed monoclonal antibody from the "Dialyzed Antibody" tube into the syringe. Replace the plunger.
- [0167] Hold the syringe filter, ACRODISC® Syringe Filter over the opened, sterile, labeled "Purified Antibody" collection tube, and depress the syringe plunger to filter the purified antibody into the "Purified Antibody" tube.
- [0168] When filtration is complete, cap the "Purified Antibody" tube and store at 2° C. to 10° C.
- [0169] Determine concentration of purified monoclonal antibody using A280 procedure.

#### Example 4

##### General Method for Epitope Mapping

#### General Approach

[0170] Epitope mapping is performed to identify the linear amino acid sequence within an antigenic protein (i.e., the epitope) that is recognized by a particular monoclonal antibody. A general approach for epitope mapping requires the expression of the full-length protein, as well as various fragments (i.e., truncated forms) of the protein, generally in a heterologous expression system. These various recombinant proteins are then used to determine if the specific monoclonal antibody is capable of binding one or more of the truncated forms of the target protein. Through the use of reiterative truncation and the generation of recombinant proteins with overlapping amino acid regions, it is possible to identify the region that is recognized by the monoclonal antibody under

investigation. Western blot analysis or ELISA is employed to determine if the specific monoclonal antibody under investigation is capable of binding one or more of the recombinant protein fragments. This approach can ultimately identify the peptide regions that contains the epitope and, in some cases, to refine the epitope precisely to an 8-15 amino acid sequence. An epitope can be a continuous linear sequence of 8-15 amino acids or it can be discontinuous with the antibody binding to a site on the protein composed of different sections of the peptide chain. Discontinuous epitopes generally cannot be mapped.

#### Construct Design and Creation

[0171] The first step in epitope mapping is the design of nested gene truncations. Frequently, the gene is divided into four equal parts for further analysis.

#### Gene Cloning Strategy

[0172] The general cloning strategy begins with PCR-based generation of the cloned gene fragments. In order to efficiently express the cloned fragment, especially when using small amino acid regions, the cloned fragment is expressed as a fusion protein, i.e. fused to another carrier protein that is stably expressed in the system. Green fluorescent protein (GFP) is frequently used as the carrier protein. GFP is included as a fusion partner to stabilize the truncation fragments and improve expression during the subsequent in vitro protein expression step. GFP also permits the tracking of fusion-protein expression using anti-GFP antibodies.

[0173] Cloning to create the GFP-protein construct is performed using either the mega-priming approach or through the use of plasmid cloning into the pScreen-GFP vector. Generally, the truncation fragments are fused to GFP and control sequences necessary for protein expression using a technique called megapriming.

[0174] Megapriming is the joining of two or more DNA fragments by annealing homologous regions at the end of the respective fragments and extending the annealed single-stranded DNA with a thermostable DNA polymerase. This process creates one large DNA fragment from two or more smaller fragments, linking them by their shared sequence. This large fragment is then amplified using standard PCR.

[0175] If megapriming cannot be used successfully, the truncation fragments can be cloned into a plasmid containing GFP and protein-expression control sequences. This cloning creates the GFP/fragment fusions necessary for epitope mapping. The remainder of the protocol can then proceed as described below.

#### Protein Expression

[0176] The expression constructs created by, for example, megapriming are then introduced into the Rapid Translation System (RTS). RTS is a cell-free protein expression system derived from *E. coli* lysates. This system permits rapid (3-4 hour) expression of proteins from DNA templates.

[0177] If RTS does not produce adequate levels of protein expression, then the truncation fragments will be cloned into the GFP protein-expression plasmid. These fusion plasmids are then transformed into an *E. coli* strain optimized for protein expression. Protein expression is induced in a growing culture of bacteria and, following outgrowth, the cells are lysed. The proteins in the complex cell lysate are then sepa-

rated by polyacrylamide gel electrophoresis (PAGE), and the remainder of the protocol is the same as below.

#### Protein Detection and Epitope Mapping

**[0178]** Protein fragments produced by RTS are separated using PAGE and transferred onto nitrocellulose membranes. The membrane-bound proteins are then exposed to the antibody under investigation in solution. Antibody/protein binding is identified using colorimetric techniques known in the art.

**[0179]** Antibody binding of the full-length protein and some subset of the truncated protein fragments constitutes a positive result. If the absence of a particular section of the protein eliminates antibody binding, then the epitope lies on this fragment.

**[0180]** If the antibody to be mapped does not recognize protein bound to nitrocellulose membranes, then alternative methods for detecting antibody/protein interactions, such as, for example, ELISA or immunoprecipitation are used. Methods for detecting antibody/protein interactions are well known in the art.

#### Refining the Epitope Location

**[0181]** Since the above-described protocol will only narrow the location of the epitope down to approximately one-quarter of the protein, it is necessary to repeat the process on the quarter of the protein determined to contain the epitope in order to further resolve the location of the epitope. For a very large protein, it may be necessary to repeat this process two to three times to narrow the epitope down to 8-15 amino acids.

#### Example 5

##### Characterization of Epitope for MCM6 Monoclonal Antibody 9D4.3

**[0182]** Epitope mapping for MCM6 monoclonal antibody 9D4.3 was carried out essentially as described above in Example 4. Specifically, PCR was used to create four MCM6 gene truncations of the full-length MCM6 protein, followed by RTS to generate recombinant MCM6 protein fragments as GFP fusion proteins, and finally western blotting to detect antibody binding to specific MCM6 fragments. GFP was joined with the MCM6 gene truncations in a second round of PCR to ensure robust and stable expression in RTS.

**[0183]** The MCM6 protein fragments were analyzed by western blotting to identify fragment(s) that bind the 9D4.3 antibody. The western blot was probed directly with the 9D4.3 monoclonal antibody and a GFP antibody. A positive

band was detected with the MCM6 truncation product designated as fragment 4. Fragment 4 was divided into five smaller fragments and the above process repeated to narrow the epitope.

**[0184]** The second set of MCM6 protein fragments was also analyzed by western blotting to identify fragment(s) that bind the 9D4.3 antibody. The western blot was probed directly with the 9D4.3 monoclonal antibody and a GFP antibody. Monoclonal antibody 9D4.3 was shown to bind to the region of MCM6 designated as 4-4. This fragment was again divided into six smaller fragments and the above process repeated to narrow the epitope.

**[0185]** The MCM6 protein fragments were again analyzed by western blotting as before. The western blot was probed directly with the 9D4.3 monoclonal antibody and a GFP antibody. A positive band was detected with the MCM6 fragment designated as 4-4-1. Additional fragments were generated to narrow the epitope region. Western blot analysis indicated that the epitope for the MCM6 antibody 9D4.3 comprises the amino acid sequence IDSEEELINKKRI (SEQ ID NO:5).

#### Results

**[0186]** Initial results showed that the epitope for the MCM6 monoclonal antibody 9D4.3 is located within the C-terminal region of the MCM6 protein. Continued truncations of the MCM6 protein showed that the epitope recognized by 9D4.3 is located within a thirteen amino acid region, specifically corresponding to amino acid residues 760-772 of SEQ ID NO:3 (IDSEEELINKKRI (SEQ ID NO:5)). Additional rounds of RTS may be able to refine the epitope location further.

#### Example 6

##### Characterization of Epitope for MCM7 Monoclonal Antibody 2E6.2

**[0187]** Epitope mapping for MCM7 monoclonal antibody 2E6.2 was performed essentially as described above in Example 5. The full-length MCM7 gene sequence (SEQ ID NO:1) was used as the starting sequence for designing gene fragments.

#### Results

**[0188]** The epitope for MCM7 monoclonal antibody 2E6.2 was determined to be located within the protein region comprising amino acid residues 127-138 of SEQ ID NO:1 (PAELMRRFELYF (SEQ ID NO:6)). Additional rounds of RTS may be able to refine the epitope location further.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 10

<210> SEQ ID NO 1

<211> LENGTH: 719

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Ala Leu Lys Asp Tyr Ala Leu Glu Lys Glu Lys Val Lys Lys Phe

-continued

1	5	10	15
Leu	Gln	Glu	Phe Tyr Gln Asp Asp Glu Leu Gly Lys Lys Gln Phe Lys
	20		25 30
Tyr	Gly	Asn	Gln Leu Val Arg Leu Ala His Arg Glu Gln Val Ala Leu
	35		40 45
Tyr	Val	Asp	Leu Asp Asp Val Ala Glu Asp Asp Pro Glu Leu Val Asp
	50		55 60
Ser	Ile	Cys	Glu Asn Ala Arg Arg Tyr Ala Lys Leu Phe Ala Asp Ala
	65		70 75 80
Val	Gln	Glu	Leu Leu Pro Gln Tyr Lys Glu Arg Glu Val Val Asn Lys
		85	90 95
Asp	Val	Leu	Asp Val Tyr Ile Glu His Arg Leu Met Met Glu Gln Arg
	100		105 110
Ser	Arg	Asp	Pro Gly Met Val Arg Ser Pro Gln Asn Gln Tyr Pro Ala
	115		120 125
Glu	Leu	Met	Arg Arg Phe Glu Leu Tyr Phe Gln Gly Pro Ser Ser Asn
	130		135 140
Lys	Pro	Arg	Val Ile Arg Glu Val Arg Ala Asp Ser Val Gly Lys Leu
	145		150 155 160
Val	Thr	Val	Arg Gly Ile Val Thr Arg Val Ser Glu Val Lys Pro Lys
		165	170 175
Met	Val	Val	Ala Thr Tyr Thr Cys Asp Gln Cys Gly Ala Glu Thr Tyr
	180		185 190
Gln	Pro	Ile	Gln Ser Pro Thr Phe Met Pro Leu Ile Met Cys Pro Ser
	195		200 205
Gln	Glu	Cys	Gln Thr Asn Arg Ser Gly Gly Arg Leu Tyr Leu Gln Thr
	210		215 220
Arg	Gly	Ser	Arg Phe Ile Lys Phe Gln Glu Met Lys Met Gln Glu His
	225		230 235 240
Ser	Asp	Gln	Val Pro Val Gly Asn Ile Pro Arg Ser Ile Thr Val Leu
		245	250 255
Val	Glu	Gly	Glu Asn Thr Arg Ile Ala Gln Pro Gly Asp His Val Ser
	260		265 270
Val	Thr	Gly	Ile Phe Leu Pro Ile Leu Arg Thr Gly Phe Arg Gln Val
	275		280 285
Val	Gln	Gly	Leu Leu Ser Glu Thr Tyr Leu Glu Ala His Arg Ile Val
	290		295 300
Lys	Met	Asn	Lys Ser Glu Asp Asp Glu Ser Gly Ala Gly Glu Leu Thr
	305		310 315 320
Arg	Glu	Glu	Leu Arg Gln Ile Ala Glu Glu Asp Phe Tyr Glu Lys Leu
		325	330 335
Ala	Ala	Ser	Ile Ala Pro Glu Ile Tyr Gly His Glu Asp Val Lys Lys
		340	345 350
Ala	Leu	Leu	Leu Leu Val Gly Gly Val Asp Gln Ser Pro Arg Gly
	355		360 365
Met	Lys	Ile	Arg Gly Asn Ile Asn Ile Cys Leu Met Gly Asp Pro Gly
	370		375 380
Val	Ala	Lys	Ser Gln Leu Leu Ser Tyr Ile Asp Arg Leu Ala Pro Arg
	385		390 395 400
Ser	Gln	Tyr	Thr Thr Gly Arg Gly Ser Ser Gly Val Gly Leu Thr Ala
		405	410 415

-continued

---

Ala Val Leu Arg Asp Ser Val Ser Gly Glu Leu Thr Leu Glu Gly Gly  
 420 425 430

Ala Leu Val Leu Ala Asp Gln Gly Val Cys Cys Ile Asp Glu Phe Asp  
 435 440 445

Lys Met Ala Glu Ala Asp Arg Thr Ala Ile His Glu Val Met Glu Gln  
 450 455 460

Gln Thr Ile Ser Ile Ala Lys Ala Gly Ile Leu Thr Thr Leu Asn Ala  
 465 470 475 480

Arg Cys Ser Ile Leu Ala Ala Ala Asn Pro Ala Tyr Gly Arg Tyr Asn  
 485 490 495

Pro Arg Arg Ser Leu Glu Gln Asn Ile Gln Leu Pro Ala Ala Leu Leu  
 500 505 510

Ser Arg Phe Asp Leu Leu Trp Leu Ile Gln Asp Arg Pro Asp Arg Asp  
 515 520 525

Asn Asp Leu Arg Leu Ala Gln His Ile Thr Tyr Val His Gln His Ser  
 530 535 540

Arg Gln Pro Pro Ser Gln Phe Glu Pro Leu Asp Met Lys Leu Met Arg  
 545 550 555 560

Arg Tyr Ile Ala Met Cys Arg Glu Lys Gln Pro Met Val Pro Glu Ser  
 565 570 575

Leu Ala Asp Tyr Ile Thr Ala Ala Tyr Val Glu Met Arg Arg Glu Ala  
 580 585 590

Trp Ala Ser Lys Asp Ala Thr Tyr Thr Ser Ala Arg Thr Leu Leu Ala  
 595 600 605

Ile Leu Arg Leu Ser Thr Ala Leu Ala Arg Leu Arg Met Val Asp Val  
 610 615 620

Val Glu Lys Glu Asp Val Asn Glu Ala Ile Arg Leu Met Glu Met Ser  
 625 630 635 640

Lys Asp Ser Leu Leu Gly Asp Lys Gly Gln Thr Ala Arg Thr Gln Arg  
 645 650 655

Pro Ala Asp Val Ile Phe Ala Thr Val Arg Glu Leu Val Ser Gly Gly  
 660 665 670

Arg Ser Val Arg Phe Ser Glu Ala Glu Gln Arg Cys Val Ser Arg Gly  
 675 680 685

Phe Thr Pro Ala Gln Phe Gln Ala Ala Leu Asp Glu Tyr Glu Glu Leu  
 690 695 700

Asn Val Trp Gln Val Asn Ala Ser Arg Thr Arg Ile Thr Phe Val  
 705 710 715

<210> SEQ ID NO 2  
 <211> LENGTH: 2821  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (511) ... (2670)

<400> SEQUENCE: 2

```
cgccccctcc cagcccccaag ggtctaggat acagtctttg tagatgagcg ggtccccctt    60
ggaggacaga atgaagaatt gggaaatcat ggccgttctg gagagtagac aagaagacgg    120
cgaaagtcgg gcctgccccg cctgcggcc ccggaacaaa agaacgcgtg tgcgctggcc    180
ctttaagagc gattctctc cgccccgcc agctcggacc gcgggaaacc cggcgcctgc    240
```



-continued

actaccccgc cggagattc ccttccgacg cccgcaccgc ctccccgtca ctctattctag	300
gccccgcacgg tgattggctt gcggctagcg ggaggtgaag aaggccgcct tgtccgattg	360
gccccgcacgc agtggcgccg gtcacgtggg gggcgacgtt tcgcgccaat ttcggttggc	420
cgccacacagt ccaccgcgcg gagattctca gcttcccag gagcaagacc tctgagcccg	480
ccaagcgcgg ccgcacggcc ctccggcagcg atg gca ctg aag gac tac gcg cta	534
Met Ala Leu Lys Asp Tyr Ala Leu	
1 5	
gag aag gaa aag gtt aag aag ttc tta caa gag ttc tac cag gat gat	582
Glu Lys Glu Lys Val Lys Lys Phe Leu Gln Glu Phe Tyr Gln Asp Asp	
10 15 20	
gaa ctc ggg aag aag cag ttc aag tat ggg aac cag ttg gtt cgg ctg	630
Glu Leu Gly Lys Lys Gln Phe Lys Tyr Gly Asn Gln Leu Val Arg Leu	
25 30 35 40	
gct cat cgg gaa cag gtg gct ctg tat gtg gac ctg gac gac gta gcc	678
Ala His Arg Glu Gln Val Ala Leu Tyr Val Asp Leu Asp Asp Val Ala	
45 50 55	
gag gat gac ccc gag ttg gtg gac tca att tgt gag aat gcc agg cgc	726
Glu Asp Asp Pro Glu Leu Val Asp Ser Ile Cys Glu Asn Ala Arg Arg	
60 65 70	
tac gcg aag ctc ttt gct gat gcc gta caa gag ctg ctg cct cag tac	774
Tyr Ala Lys Leu Phe Ala Asp Ala Val Gln Glu Leu Leu Pro Gln Tyr	
75 80 85	
aag gag agg gaa gtg gta aat aaa gat gtc ctg gac gtt tac att gag	822
Lys Glu Arg Glu Val Val Asn Lys Asp Val Leu Asp Val Tyr Ile Glu	
90 95 100	
cat cgg cta atg atg gag cag cgg agt cgg gac cct ggg atg gtc cga	870
His Arg Leu Met Met Glu Gln Arg Ser Arg Asp Pro Gly Met Val Arg	
105 110 115 120	
agc ccc cag aac cag tac cct gct gaa ctc atg cgc aga ttt gag ctg	918
Ser Pro Gln Asn Gln Tyr Pro Ala Glu Leu Met Arg Arg Phe Glu Leu	
125 130 135	
tat ttt caa ggc cct agc agc aac aag cct cgt gtg atc cgg gaa gtg	966
Tyr Phe Gln Gly Pro Ser Ser Asn Lys Pro Arg Val Ile Arg Glu Val	
140 145 150	
cgg gct gac tct gtg ggg aag ttg gta act gtg cgt gga atc gtc act	1014
Arg Ala Asp Ser Val Gly Lys Leu Val Thr Val Arg Gly Ile Val Thr	
155 160 165	
cgt gtc tct gaa gtc aaa ccc aag atg gtg gtg gcc act tac act tgt	1062
Arg Val Ser Glu Val Lys Pro Lys Met Val Val Ala Thr Tyr Thr Cys	
170 175 180	
gac cag tgt ggg gca gag acc tac cag ccg atc cag tct ccc act ttc	1110
Asp Gln Cys Gly Ala Glu Thr Tyr Gln Pro Ile Gln Ser Pro Thr Phe	
185 190 195 200	
atg cct ctg atc atg tgc cca agc cag gag tgc caa acc aac cgc tca	1158
Met Pro Leu Ile Met Cys Pro Ser Gln Glu Cys Gln Thr Asn Arg Ser	
205 210 215	
gga ggg cgg ctg tat ctg cag aca cgg ggc tcc aga ttc atc aaa ttc	1206
Gly Gly Arg Leu Tyr Leu Gln Thr Arg Gly Ser Arg Phe Ile Lys Phe	
220 225 230	
cag gag atg aag atg caa gaa cat agt gat cag gtg cct gtg gga aat	1254
Gln Glu Met Lys Met Gln Glu His Ser Asp Gln Val Pro Val Gly Asn	
235 240 245	
atc cct cgt agt atc acg gtg ctg gta gaa gga gag aac aca agg att	1302
Ile Pro Arg Ser Ile Thr Val Leu Val Glu Gly Glu Asn Thr Arg Ile	
250 255 260	

-continued

gcc cag cct gga gac cac gtc agc gtc act ggt att ttc ttg cca atc	1350
Ala Gln Pro Gly Asp His Val Ser Val Thr Gly Ile Phe Leu Pro Ile	
265 270 275 280	
ctg cgc act ggg ttc cga cag gtg gta cag ggt tta ctc tca gaa acc	1398
Leu Arg Thr Gly Phe Arg Gln Val Val Gln Gly Leu Leu Ser Glu Thr	
285 290 295	
tac ctg gaa gcc cat cgg att gtg aag atg aac aag agt gag gat gat	1446
Tyr Leu Glu Ala His Arg Ile Val Lys Met Asn Lys Ser Glu Asp Asp	
300 305 310	
gag tct ggg gct gga gag ctc acc agg gag gag ctg agg caa att gca	1494
Glu Ser Gly Ala Gly Glu Leu Thr Arg Glu Glu Leu Arg Gln Ile Ala	
315 320 325	
gag gag gat ttc tac gaa aag ctg gca gct tca atc gcc cca gaa ata	1542
Glu Glu Asp Phe Tyr Glu Lys Leu Ala Ala Ser Ile Ala Pro Glu Ile	
330 335 340	
tac ggg cat gaa gat gtg aag aag gca ctg ctg ctc ctg cta gtc ggg	1590
Tyr Gly His Glu Asp Val Lys Lys Ala Leu Leu Leu Leu Val Gly	
345 350 355 360	
ggt gtg gac cag tct cct cga ggc atg aaa atc cgg ggc aac atc aac	1638
Gly Val Asp Gln Ser Pro Arg Gly Met Lys Ile Arg Gly Asn Ile Asn	
365 370 375	
atc tgt ctg atg ggg gat cct ggt gtg gcc aag tct cag ctc ctg tca	1686
Ile Cys Leu Met Gly Asp Pro Gly Val Ala Lys Ser Gln Leu Leu Ser	
380 385 390	
tac att gat cga ctg gcg cct cgc agc cag tac aca aca ggc cgg ggc	1734
Tyr Ile Asp Arg Leu Ala Pro Arg Ser Gln Tyr Thr Thr Gly Arg Gly	
395 400 405	
tcc tca gga gtg ggg ctt acg gca gct gtg ctg aga gac tcc gtg agt	1782
Ser Ser Gly Val Gly Leu Thr Ala Ala Val Leu Arg Asp Ser Val Ser	
410 415 420	
gga gaa ctg acc tta gag ggt ggg gcc ctg gtg ctg gct gac cag ggt	1830
Gly Glu Leu Thr Leu Glu Gly Gly Ala Leu Val Leu Ala Asp Gln Gly	
425 430 435 440	
gtg tgc tgc att gat gag ttc gac aag atg gct gag gcc gac cgc aca	1878
Val Cys Cys Ile Asp Glu Phe Asp Lys Met Ala Glu Ala Asp Arg Thr	
445 450 455	
gcc atc cac gag gtc atg gag cag cag acc atc tcc att gcc aag gcc	1926
Ala Ile His Glu Val Met Glu Gln Gln Thr Ile Ser Ile Ala Lys Ala	
460 465 470	
ggc att ctc acc aca ctc aat gcc cgc tgc tcc atc ctg gct gcc gcc	1974
Gly Ile Leu Thr Thr Leu Asn Ala Arg Cys Ser Ile Leu Ala Ala Ala	
475 480 485	
aac cct gcc tac ggg cgc tac aac cct cgc cgc agc ctg gag cag aac	2022
Asn Pro Ala Tyr Gly Arg Tyr Asn Pro Arg Arg Ser Leu Glu Gln Asn	
490 495 500	
ata cag cta cct gct gca ctg ctc tcc cgg ttt gac ctc ctc tgg ctg	2070
Ile Gln Leu Pro Ala Ala Leu Leu Ser Arg Phe Asp Leu Leu Trp Leu	
505 510 515 520	
att cag gac cgg ccc gac cga gac aat gac cta cgg ttg gcc cag cac	2118
Ile Gln Asp Arg Pro Asp Arg Asp Asn Asp Leu Arg Leu Ala Gln His	
525 530 535	
atc acc tat gtg cac cag cac agc cgg cag ccc ccc tcc cag ttt gaa	2166
Ile Thr Tyr Val His Gln His Ser Arg Gln Pro Pro Ser Gln Phe Glu	
540 545 550	
cct ctg gac atg aag ctc atg agg cgt tac ata gcc atg tgc cgc gag	2214
Pro Leu Asp Met Lys Leu Met Arg Arg Tyr Ile Ala Met Cys Arg Glu	
555 560 565	

-continued

```

aag cag ccc atg gtg cca gag tct ctg gct gac tac atc aca gca gca 2262
Lys Gln Pro Met Val Pro Glu Ser Leu Ala Asp Tyr Ile Thr Ala Ala
570 575 580

tac gtg gag atg agg cga gag gct tgg gct agt aag gat gcc acc tat 2310
Tyr Val Glu Met Arg Arg Glu Ala Trp Ala Ser Lys Asp Ala Thr Tyr
585 590 595 600

act tct gcc cgg acc ctg ctg gct atc ctg cgc ctt tcc act gct ctg 2358
Thr Ser Ala Arg Thr Leu Leu Ala Ile Leu Arg Leu Ser Thr Ala Leu
605 610 615

gca cgt ctg aga atg gtg gat gtg gtg gag aaa gaa gat gtg aat gaa 2406
Ala Arg Leu Arg Met Val Asp Val Val Glu Lys Glu Asp Val Asn Glu
620 625 630

gcc atc agg cta atg gag atg tca aag gac tct ctt cta gga gac aag 2454
Ala Ile Arg Leu Met Glu Met Ser Lys Asp Ser Leu Leu Gly Asp Lys
635 640 645

ggg cag aca gct agg act cag aga cca gca gat gtg ata ttt gcc acc 2502
Gly Gln Thr Ala Arg Thr Gln Arg Pro Ala Asp Val Ile Phe Ala Thr
650 655 660

gtc cgt gaa ctg gtc tca ggg ggc cga agt gtc cgg ttc tct gag gca 2550
Val Arg Glu Leu Val Ser Gly Gly Arg Ser Val Arg Phe Ser Glu Ala
665 670 675 680

gag cag cgc tgt gta tct cgt ggc ttc aca ccc gcc cag ttc cag gcg 2598
Glu Gln Arg Cys Val Ser Arg Gly Phe Thr Pro Ala Gln Phe Gln Ala
685 690 695

gct ctg gat gaa tat gag gag ctc aat gtc tgg cag gtc aat gct tcc 2646
Ala Leu Asp Glu Tyr Glu Glu Leu Asn Val Trp Gln Val Asn Ala Ser
700 705 710

cgg aca cgg atc act ttt gtc tga ttccagcctg cttgcaacc tggggctctc 2700
Arg Thr Arg Ile Thr Phe Val
715

ttgttccctg ctggcctgcc ccttgggaag gggcagtgat gcctttgagg ggaaggagga 2760

gcccctcttt ctcccatgct gcacttactc cttttgctaa taaaagtgtt tgtagattgt 2820

c 2821
    
```

```

<210> SEQ ID NO 3
<211> LENGTH: 821
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 3

```

Met Asp Leu Ala Ala Ala Ala Glu Pro Gly Ala Gly Ser Gln His Leu
1 5 10 15

Glu Val Arg Asp Glu Val Ala Glu Lys Cys Gln Lys Leu Phe Leu Asp
20 25 30

Phe Leu Glu Glu Phe Gln Ser Ser Asp Gly Glu Ile Lys Tyr Leu Gln
35 40 45

Leu Ala Glu Glu Leu Ile Arg Pro Glu Arg Asn Thr Leu Val Val Ser
50 55 60

Phe Val Asp Leu Glu Gln Phe Asn Gln Gln Leu Ser Thr Thr Ile Gln
65 70 75 80

Glu Glu Phe Tyr Arg Val Tyr Pro Tyr Leu Cys Arg Ala Leu Lys Thr
85 90 95

Phe Val Lys Asp Arg Lys Glu Ile Pro Leu Ala Lys Asp Phe Tyr Val
100 105 110
    
```

-continued

---

Ala	Phe	Gln	Asp	Leu	Pro	Thr	Arg	His	Lys	Ile	Arg	Glu	Leu	Thr	Ser
		115					120				125				
Ser	Arg	Ile	Gly	Leu	Leu	Thr	Arg	Ile	Ser	Gly	Gln	Val	Val	Arg	Thr
	130					135					140				
His	Pro	Val	His	Pro	Glu	Leu	Val	Ser	Gly	Thr	Phe	Leu	Cys	Leu	Asp
145				150					155						160
Cys	Gln	Thr	Val	Ile	Arg	Asp	Val	Glu	Gln	Gln	Phe	Lys	Tyr	Thr	Gln
				165					170					175	
Pro	Asn	Ile	Cys	Arg	Asn	Pro	Val	Cys	Ala	Asn	Arg	Arg	Arg	Phe	Leu
			180					185					190		
Leu	Asp	Thr	Asn	Lys	Ser	Arg	Phe	Val	Asp	Phe	Gln	Lys	Val	Arg	Ile
		195				200						205			
Gln	Glu	Thr	Gln	Ala	Glu	Leu	Pro	Arg	Gly	Ser	Ile	Pro	Arg	Ser	Leu
	210					215					220				
Glu	Val	Ile	Leu	Arg	Ala	Glu	Ala	Val	Glu	Ser	Ala	Gln	Ala	Gly	Asp
225				230					235						240
Lys	Cys	Asp	Phe	Thr	Gly	Thr	Leu	Ile	Val	Val	Pro	Asp	Val	Ser	Lys
			245					250						255	
Leu	Ser	Thr	Pro	Gly	Ala	Arg	Ala	Glu	Thr	Asn	Ser	Arg	Val	Ser	Gly
			260					265					270		
Val	Asp	Gly	Tyr	Glu	Thr	Glu	Gly	Ile	Arg	Gly	Leu	Arg	Ala	Leu	Gly
		275				280						285			
Val	Arg	Asp	Leu	Ser	Tyr	Arg	Leu	Val	Phe	Leu	Ala	Cys	Cys	Val	Ala
	290					295					300				
Pro	Thr	Asn	Pro	Arg	Phe	Gly	Gly	Lys	Glu	Leu	Arg	Asp	Glu	Glu	Gln
305				310					315						320
Thr	Ala	Glu	Ser	Ile	Lys	Asn	Gln	Met	Thr	Val	Lys	Glu	Trp	Glu	Lys
			325					330						335	
Val	Phe	Glu	Met	Ser	Gln	Asp	Lys	Asn	Leu	Tyr	His	Asn	Leu	Cys	Thr
		340						345					350		
Ser	Leu	Phe	Pro	Thr	Ile	His	Gly	Asn	Asp	Glu	Val	Lys	Arg	Gly	Val
	355						360						365		
Leu	Leu	Met	Leu	Phe	Gly	Gly	Val	Pro	Lys	Thr	Thr	Gly	Glu	Gly	Thr
	370				375						380				
Ser	Leu	Arg	Gly	Asp	Ile	Asn	Val	Cys	Ile	Val	Gly	Asp	Pro	Ser	Thr
385				390					395						400
Ala	Lys	Ser	Gln	Phe	Leu	Lys	His	Val	Glu	Glu	Phe	Ser	Pro	Arg	Ala
			405					410						415	
Val	Tyr	Thr	Ser	Gly	Lys	Ala	Ser	Ser	Ala	Ala	Gly	Leu	Thr	Ala	Ala
		420						425					430		
Val	Val	Arg	Asp	Glu	Glu	Ser	His	Glu	Phe	Val	Ile	Glu	Ala	Gly	Ala
	435						440					445			
Leu	Met	Leu	Ala	Asp	Asn	Gly	Val	Cys	Cys	Ile	Asp	Glu	Phe	Asp	Lys
	450				455						460				
Met	Asp	Val	Arg	Asp	Gln	Val	Ala	Ile	His	Glu	Ala	Met	Glu	Gln	Gln
465				470					475						480
Thr	Ile	Ser	Ile	Thr	Lys	Ala	Gly	Val	Lys	Ala	Thr	Leu	Asn	Ala	Arg
			485					490						495	
Thr	Ser	Ile	Leu	Ala	Ala	Ala	Asn	Pro	Ile	Ser	Gly	His	Tyr	Asp	Arg
		500						505					510		
Ser	Lys	Ser	Leu	Lys	Gln	Asn	Ile	Asn	Leu	Ser	Ala	Pro	Ile	Met	Ser

-continued

515	520	525
Arg Phe Asp Leu Phe Phe Ile Leu Val Asp Glu Cys Asn Glu Val Thr 530 535 540		
Asp Tyr Ala Ile Ala Arg Arg Ile Val Asp Leu His Ser Arg Ile Glu 545 550 555 560		
Glu Ser Ile Asp Arg Val Tyr Ser Leu Asp Asp Ile Arg Arg Tyr Leu 565 570 575		
Leu Phe Ala Arg Gln Phe Lys Pro Lys Ile Ser Lys Glu Ser Glu Asp 580 585 590		
Phe Ile Val Glu Gln Tyr Lys His Leu Arg Gln Arg Asp Gly Ser Gly 595 600 605		
Val Thr Lys Ser Ser Trp Arg Ile Thr Val Arg Gln Leu Glu Ser Met 610 615 620		
Ile Arg Leu Ser Glu Ala Met Ala Arg Met His Cys Cys Asp Glu Val 625 630 635 640		
Gln Pro Lys His Val Lys Glu Ala Phe Arg Leu Leu Asn Lys Ser Ile 645 650 655		
Ile Arg Val Glu Thr Pro Asp Val Asn Leu Asp Gln Glu Glu Glu Ile 660 665 670		
Gln Met Glu Val Asp Glu Gly Ala Gly Gly Ile Asn Gly His Ala Asp 675 680 685		
Ser Pro Ala Pro Val Asn Gly Ile Asn Gly Tyr Asn Glu Asp Ile Asn 690 695 700		
Gln Glu Ser Ala Pro Lys Ala Ser Leu Arg Leu Gly Phe Ser Glu Tyr 705 710 715 720		
Cys Arg Ile Ser Asn Leu Ile Val Leu His Leu Arg Lys Val Glu Glu 725 730 735		
Glu Glu Asp Glu Ser Ala Leu Lys Arg Ser Glu Leu Val Asn Trp Tyr 740 745 750		
Leu Lys Glu Ile Glu Ser Glu Ile Asp Ser Glu Glu Glu Leu Ile Asn 755 760 765		
Lys Lys Arg Ile Ile Glu Lys Val Ile His Arg Leu Thr His Tyr Asp 770 775 780		
His Val Leu Ile Glu Leu Thr Gln Ala Gly Leu Lys Gly Ser Thr Glu 785 790 795 800		
Gly Ser Glu Ser Tyr Glu Glu Asp Pro Tyr Leu Val Val Asn Pro Asn 805 810 815		
Tyr Leu Leu Glu Asp 820		

<210> SEQ ID NO 4  
<211> LENGTH: 3769  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens  
<220> FEATURE:  
<221> NAME/KEY: CDS  
<222> LOCATION: (77)...(2542)

<400> SEQUENCE: 4

aaagctgcag cgtctgga aaagcgactt gtggcggtcg agcgtggcgc aggcgaatcc 60

tgggcactaa gcaaat atg gac ctc gcg gca gca gca gag ccg ggc gcc gcc 112  
Met Asp Leu Ala Ala Ala Ala Glu Pro Gly Ala Gly  
1 5 10

-continued

agc Ser	cag Gln	cac His	ctg Leu	gag Glu	gtc Val	cgc Arg	gac Asp	gag Glu	gtg Val	gcc Ala	gag Glu	aag Lys	tgc Cys	cag Gln	aaa Lys	160
	15					20						25				
ctg Leu	ttc Phe	ctg Leu	gac Asp	ttc Phe	ttg Leu	gag Glu	gag Glu	ttt Phe	cag Gln	agc Ser	agc Ser	gat Asp	gga Gly	gaa Glu	att Ile	208
	30					35					40					
aaa Lys	tac Tyr	ttg Leu	caa Gln	tta Leu	gca Ala	gag Glu	gaa Glu	ctg Leu	att Ile	cgt Arg	cct Pro	gag Glu	aga Arg	aac Asn	aca Thr	256
	45				50					55					60	
ttg Leu	gtt Val	gtg Val	agt Ser	ttt Phe	gtg Val	gac Asp	ctg Leu	gaa Glu	caa Gln	ttt Phe	aac Asn	cag Gln	caa Gln	ctt Leu	tcc Ser	304
				65					70					75		
acc Thr	acc Thr	att Ile	caa Gln	gag Glu	gag Glu	ttc Phe	tat Tyr	aga Arg	gtt Val	tac Tyr	cct Pro	tac Tyr	ctg Leu	tgt Cys	egg Arg	352
			80					85					90			
gcc Ala	ttg Leu	aaa Lys	aca Thr	ttc Phe	gtc Val	aaa Lys	gac Asp	cgt Arg	aaa Lys	gag Glu	atc Ile	cct Pro	ctt Leu	gcc Ala	aag Lys	400
		95					100					105				
gat Asp	ttt Phe	tat Tyr	ggt Val	gca Ala	ttc Phe	caa Gln	gac Asp	ctg Leu	cct Pro	acc Thr	aga Arg	cac His	aag Lys	att Ile	cga Arg	448
	110				115						120					
gag Glu	ctc Leu	acc Thr	tca Ser	tcc Ser	aga Arg	att Ile	ggt Gly	ttg Leu	ctc Leu	act Thr	cgc Arg	atc Ile	agt Ser	ggg Gly	cag Gln	496
	125				130					135				140		
gtg Val	gtg Val	cgg Arg	act Thr	cac His	cca Pro	gtt Val	cac His	cca Pro	gag Glu	ctt Leu	gtg Val	agc Ser	gga Gly	act Thr	ttt Phe	544
				145					150					155		
ctg Leu	tgc Cys	ttg Leu	gac Asp	tgt Cys	cag Gln	aca Thr	gtg Val	atc Ile	agg Arg	gat Asp	gta Val	gaa Glu	cag Gln	cag Gln	ttc Phe	592
			160					165					170			
aaa Lys	tac Tyr	aca Thr	cag Gln	cca Pro	aac Asn	atc Ile	tgc Cys	cga Arg	aat Asn	cca Pro	gtt Val	tgt Cys	gcc Ala	aac Asn	agg Arg	640
		175					180					185				
agg Arg	aga Arg	ttc Phe	tta Leu	ctg Leu	gat Asp	aca Thr	aat Asn	aaa Lys	tca Ser	aga Arg	ttt Phe	gtt Val	gat Asp	ttt Phe	caa Gln	688
		190			195					200						
aag Lys	gtt Val	cgt Arg	att Ile	caa Gln	gag Glu	acc Thr	caa Gln	gct Ala	gag Glu	ctt Leu	cct Pro	cga Arg	ggg Gly	agt Ser	atc Ile	736
	205			210						215				220		
ccc Pro	cgc Arg	agt Ser	tta Leu	gaa Glu	gta Val	att Ile	tta Leu	agg Arg	gct Ala	gaa Glu	gct Ala	gtg Val	gaa Glu	tca Ser	gct Ala	784
			225					230						235		
caa Gln	gct Ala	ggt Gly	gac Asp	aag Lys	tgt Cys	gac Asp	ttt Phe	aca Thr	ggg Gly	aca Thr	ctg Thr	att Leu	gtt Ile	gtg Val	cct Pro	832
		240					245						250			
gac Asp	gtc Val	tcc Ser	aag Lys	ctt Leu	agc Ser	aca Thr	cca Pro	gga Gly	gca Ala	cgt Arg	gca Ala	gaa Glu	act Thr	aat Asn	tcc Ser	880
		255					260					265				
cgt Arg	gtc Val	agt Ser	ggt Gly	gtt Val	gat Asp	gga Gly	tat Tyr	gag Glu	aca Thr	gaa Glu	ggc Gly	att Ile	cga Arg	gga Gly	ctc Leu	928
	270				275						280					
egg Arg	gcc Ala	ctt Leu	ggt Gly	gtt Val	agg Arg	gac Asp	ctt Leu	tct Ser	tat Tyr	agg Arg	ctg Leu	gtc Val	ttt Phe	ctt Leu	gcc Ala	976
	285				290					295				300		
tgc Cys	tgt Cys	gtt Val	gcg Ala	cca Pro	acc Thr	aac Asn	cca Pro	agg Arg	ttt Phe	ggg Gly	ggg Gly	aaa Lys	gag Glu	ctc Leu	aga Arg	1024
				305					310					315		

-continued

gat gag gaa cag aca gct gag agc att aag aac caa atg act gtg aaa	1072
Asp Glu Glu Gln Thr Ala Glu Ser Ile Lys Asn Gln Met Thr Val Lys	
320 325 330	
gaa tgg gag aaa gtg ttt gag atg agt caa gat aaa aat cta tac cac	1120
Glu Trp Glu Lys Val Phe Glu Met Ser Gln Asp Lys Asn Leu Tyr His	
335 340 345	
aat ctt tgt acc agc ctg ttc cct act ata cat ggc aat gat gaa gta	1168
Asn Leu Cys Thr Ser Leu Phe Pro Thr Ile His Gly Asn Asp Glu Val	
350 355 360	
aaa cgg ggt gtc ctg ctg atg ctc ttt ggt ggc gtt cca aag aca aca	1216
Lys Arg Gly Val Leu Leu Met Leu Phe Gly Gly Val Pro Lys Thr Thr	
365 370 375 380	
gga gaa ggg acc tct ctt cga ggg gac ata aat gtt tgc att gtt ggt	1264
Gly Glu Gly Thr Ser Leu Arg Gly Asp Ile Asn Val Cys Ile Val Gly	
385 390 395	
gac cca agt aca gct aag agc caa ttt ctc aag cac gtg gag gag ttc	1312
Asp Pro Ser Thr Ala Lys Ser Gln Phe Leu Lys His Val Glu Glu Phe	
400 405 410	
agc ccc aga gct gtc tac acc agt ggt aaa gcg tcc agt gct gct ggc	1360
Ser Pro Arg Ala Val Tyr Thr Ser Gly Lys Ala Ser Ser Ala Ala Gly	
415 420 425	
tta aca gca gct gtt gtg aga gat gaa gaa tct cat gag ttt gtc att	1408
Leu Thr Ala Ala Val Val Arg Asp Glu Glu Ser His Glu Phe Val Ile	
430 435 440	
gag gct gga gct ttg atg ttg gct gat aat ggt gtg tgt tgt att gat	1456
Glu Ala Gly Ala Leu Met Leu Ala Asp Asn Gly Val Cys Cys Ile Asp	
445 450 455 460	
gaa ttt gat aag atg gac gtg cgg gat caa gtt gct att cat gaa gct	1504
Glu Phe Asp Lys Met Asp Val Arg Asp Gln Val Ala Ile His Glu Ala	
465 470 475	
atg gaa cag cag acc ata tcc atc act aaa gca gga gtg aag gct act	1552
Met Glu Gln Gln Thr Ile Ser Ile Thr Lys Ala Gly Val Lys Ala Thr	
480 485 490	
ctg aac gcc cgg acg tcc att ttg gca gca gca aac cca atc agt gga	1600
Leu Asn Ala Arg Thr Ser Ile Leu Ala Ala Ala Asn Pro Ile Ser Gly	
495 500 505	
cac tat gac aga tca aaa tca ttg aaa cag aat ata aat ttg tca gct	1648
His Tyr Asp Arg Ser Lys Ser Leu Lys Gln Asn Ile Asn Leu Ser Ala	
510 515 520	
ccc atc atg tcc cga ttc gat ctc ttc ttt atc ctt gtg gat gaa tgt	1696
Pro Ile Met Ser Arg Phe Asp Leu Phe Phe Ile Leu Val Asp Glu Cys	
525 530 535 540	
aat gag gtt aca gat tat gcc att gcc agg cgc ata gta gat ttg cat	1744
Asn Glu Val Thr Asp Tyr Ala Ile Ala Arg Arg Ile Val Asp Leu His	
545 550 555	
tca aga att gag gaa tca att gat cgt gtc tat tcc ctc gat gat atc	1792
Ser Arg Ile Glu Glu Ser Ile Asp Arg Val Tyr Ser Leu Asp Asp Ile	
560 565 570	
aga aga tat ctt ctc ttt gca aga cag ttt aaa ccc aag att tcc aaa	1840
Arg Arg Tyr Leu Leu Phe Ala Arg Gln Phe Lys Pro Lys Ile Ser Lys	
575 580 585	
gag tca gag gac ttc att gtg gag caa tat aaa cat ctc cgc cag aga	1888
Glu Ser Glu Asp Phe Ile Val Glu Gln Tyr Lys His Leu Arg Gln Arg	
590 595 600	
gat ggt tct gga gtg acc aag tct tca tgg agg att aca gtg cga cag	1936
Asp Gly Ser Gly Val Thr Lys Ser Ser Trp Arg Ile Thr Val Arg Gln	
605 610 615 620	

-continued

ctt gag agc atg att cgt ctc tct gaa gct atg gct cgg atg cac tgc	1984
Leu Glu Ser Met Ile Arg Leu Ser Glu Ala Met Ala Arg Met His Cys	
625 630 635	
tgt gat gag gtc caa cct aaa cat gtg aag gaa gct ttc cgg tta ctg	2032
Cys Asp Glu Val Gln Pro Lys His Val Lys Glu Ala Phe Arg Leu Leu	
640 645 650	
aat aaa tca atc atc cgt gtg gaa aca cct gat gtc aat cta gat caa	2080
Asn Lys Ser Ile Ile Arg Val Glu Thr Pro Asp Val Asn Leu Asp Gln	
655 660 665	
gag gaa gag atc cag atg gag gta gat gag ggt gct ggt ggc atc aat	2128
Glu Glu Glu Ile Gln Met Glu Val Asp Glu Gly Ala Gly Gly Ile Asn	
670 675 680	
ggt cat gct gac agc cct gct cct gtg aac ggg atc aat ggc tac aat	2176
Gly His Ala Asp Ser Pro Ala Pro Val Asn Gly Ile Asn Gly Tyr Asn	
685 690 695 700	
gaa gac ata aat caa gag tct gct ccc aaa gcc tcc tta agg ctg ggc	2224
Glu Asp Ile Asn Gln Glu Ser Ala Pro Lys Ala Ser Leu Arg Leu Gly	
705 710 715	
ttc tct gag tac tgc cga atc tct aac ctt att gtg ctt cac ctc aga	2272
Phe Ser Glu Tyr Cys Arg Ile Ser Asn Leu Ile Val Leu His Leu Arg	
720 725 730	
aag gtg gaa gaa gaa gag gac gag tca gca tta aag agg agc gag ctt	2320
Lys Val Glu Glu Glu Glu Asp Glu Ser Ala Leu Lys Arg Ser Glu Leu	
735 740 745	
gtt aac tgg tac ttg aag gaa atc gaa tca gag ata gac tct gaa gaa	2368
Val Asn Trp Tyr Leu Lys Glu Ile Glu Ser Glu Ile Asp Ser Glu Glu	
750 755 760	
gaa ctt ata aat aaa aaa aga atc ata gag aaa gtt att cat cga ctc	2416
Glu Leu Ile Asn Lys Lys Arg Ile Ile Glu Lys Val Ile His Arg Leu	
765 770 775 780	
aca cac tat gat cat gtt cta att gag ctc acc cag gct gga ttg aaa	2464
Thr His Tyr Asp His Val Leu Ile Glu Leu Thr Gln Ala Gly Leu Lys	
785 790 795	
ggc tcc aca gag gga agt gag agc tat gaa gaa gat ccc tac ttg gta	2512
Gly Ser Thr Glu Gly Ser Glu Ser Tyr Glu Glu Asp Pro Tyr Leu Val	
800 805 810	
gtt aac cct aac tac ttg ctc gaa gat tga gatagtgaaa gtaactgacc	2562
Val Asn Pro Asn Tyr Leu Leu Glu Asp	
815 820	
agagctgagg aactgtggca cagcacctcg tggcctggag cctggctgga gctctgctag	2622
ggacagaagt gtttctggaa gtgatgcttc caggatttgt tttcagaaac aagaattgag	2682
ttgatggtcc tatgtgtcac attcatcaca ggtttcatcac caacacaggc ttcagcactt	2742
cctttggtgt gtttccctgtc ccagtgaagt tggaaacaaa taatgtgtag tctctataac	2802
caataaccttt gttttcatgt gtaagaaaag gccattact ttttaaggat gtgctgtcct	2862
attgagcaaaa taactttttt tcaattgccca gctactgctt ttattcatca aaataaata	2922
acttgttctg aagttgtota ttggatttct ttctactgta ccctgattat tacttccatc	2982
tacttttgaa tgtgagactt tccctttttg cttaacctgg agtgaagagg tagaactgtg	3042
gtattatgga tgaggtttct atgagaagga gtcattagag aactcatatg aaagctagag	3102
gccttagaga tgactttoca aggttaattc cagttgtttt tttttttttt taagtttata	3162
aaagtttatt atactttttt aaaattactc ttttagtaatt tattttactt ctgtgtocta	3222
agggtaattt ctcaggattg ttttcaaatt gcttttttag gggaaatagg tcatttgcta	3282



-continued

---

```
tattacaagc aatcccaaaa ttttatggtc ttccaggaag agttattacc gtttatgata 3342
ctaacagttc ctgagactta gctatgatca gtatgttcat gaggtggagc agttcctgtg 3402
ttgcagcttt taacaacaga tggcattcat taaatcaca agtatgttaa aggtcacaaa 3462
agcaaaaataa ctgtctgagg ctaaggccca cgtgggacag tctaataccc atgagtactc 3522
aacttgccctt gatgtctgag ctttccagtg caatgtgaat ttgagcagcc agaaatctat 3582
tagtagaaag caagacagat taatataggt taaaacaatg atttaaatat gtttctccca 3642
ataattatct ctttccctgg aatcaacttg tatgaaacct tgtcaaaatg tactccacaa 3702
gtatgtacaa ttaagtatct taaaaataaa tggcaaacat taaaacaaa aaaaaaaaaa 3762
aaaaaaaaa 3769
```

```
<210> SEQ ID NO 5
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Epitope for MCM6 monoclonal antibody 9D4.3

<400> SEQUENCE: 5
```

```
Ile Asp Ser Glu Glu Leu Ile Asn Lys Lys Arg Ile
1           5           10
```

```
<210> SEQ ID NO 6
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Epitope for MCM7 monoclonal antibody 2E6.2

<400> SEQUENCE: 6
```

```
Pro Ala Glu Leu Met Arg Arg Phe Glu Leu Tyr Phe
1           5           10
```

```
<210> SEQ ID NO 7
<211> LENGTH: 555
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Nucleotide sequence for a hexahistidine-tagged
N-terminal fragment of MCM7

<400> SEQUENCE: 7
```

```
atggcactga aggactacgc gctagagaag gaaaaggtta agaagttctt acaagagttc 60
taccaggatg atgaactcgg gaagaagcag ttcaagtatg ggaaccagtt ggttcggctg 120
gctcatcggg aacaggtggc tctgtatgtg gacctggacg acgtagccga ggatgacccc 180
gagttggtgg actcaatttg tgagaatgcc aggcgctacg cgaagctctt tgctgatgcc 240
gtacaagagc tgctgcctca gtacaaggag agggaagtgg taaataaaga tgcctcggac 300
gtttacattg agcatcggct aatgatggag cagcggagtc gggaccctgg gatggtcgca 360
agccccaga accagtaacc tgctgaactc atgcgcagat ttgagctgta ttttcaaggc 420
cctagcagca acaagcctcg tgtgatccgg gaagtgcggg ctgactctgt ggggaagtgg 480
gtaactgtgc gtggaatcgt cactcgtgtc tctgaagtca aaccaaggg tggatcatcat 540
catcatcatc attga 555
```

-continued

---

<210> SEQ ID NO 8  
 <211> LENGTH: 184  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Amino acid sequence for the  
 hexahistidine-tagged N-terminal fragment of MCM7 encoded by SEQ ID  
 NO:7

&lt;400&gt; SEQUENCE: 8

Met Ala Leu Lys Asp Tyr Ala Leu Glu Lys Glu Lys Val Lys Lys Phe  
 1 5 10 15  
 Leu Gln Glu Phe Tyr Gln Asp Asp Glu Leu Gly Lys Lys Gln Phe Lys  
 20 25 30  
 Tyr Gly Asn Gln Leu Val Arg Leu Ala His Arg Glu Gln Val Ala Leu  
 35 40 45  
 Tyr Val Asp Leu Asp Asp Val Ala Glu Asp Asp Pro Glu Leu Val Asp  
 50 55 60  
 Ser Ile Cys Glu Asn Ala Arg Arg Tyr Ala Lys Leu Phe Ala Asp Ala  
 65 70 75 80  
 Val Gln Glu Leu Leu Pro Gln Tyr Lys Glu Arg Glu Val Val Asn Lys  
 85 90 95  
 Asp Val Leu Asp Val Tyr Ile Glu His Arg Leu Met Met Glu Gln Arg  
 100 105 110  
 Ser Arg Asp Pro Gly Met Val Arg Ser Pro Gln Asn Gln Tyr Pro Ala  
 115 120 125  
 Glu Leu Met Arg Arg Phe Glu Leu Tyr Phe Gln Gly Pro Ser Ser Asn  
 130 135 140  
 Lys Pro Arg Val Ile Arg Glu Val Arg Ala Asp Ser Val Gly Lys Leu  
 145 150 155 160  
 Val Thr Val Arg Gly Ile Val Thr Arg Val Ser Glu Val Lys Pro Lys  
 165 170 175  
 Gly Gly His His His His His His  
 180

<210> SEQ ID NO 9  
 <211> LENGTH: 2590  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Nucleotide sequence for a FLAG-tagged MCM6  
 polypeptide

&lt;400&gt; SEQUENCE: 9

cgagctcgga tccactagta acggccgcca gtgtgctgga attcgccctt ttgctagccc 60  
 accatggaac tcgcgggcgc agcggagccg ggcgccggca gccagcacct ggaggctcgc 120  
 gacgaggtgg ccgagaagtg ccagaaactg ttcctggact tcttgaggga gtttcagagc 180  
 agcgatggag aaattaataa cttgcaatta gcagaggaac tgattcgtcc tgagagaaac 240  
 acattggttg tgagttttgt ggacctggaa caatttaacc agcaactttc caccaccatt 300  
 caagaggagt tctatagagt ttacccttac ctgtgtcggg ccttgaaaac attcgtcaaa 360  
 gaccgtaaag agatccctct tgccaaggat ttttatgttg cattocaaga cctgcctacc 420  
 agacacaaga ttcgagagct cacctcatcc agaattggtt tgctcaactcg catcagtggtg 480  
 caggtggtgc ggactcacc agttcaccca gagcttgtga gcggaacttt tctgtgcttg 540

-continued

---

gactgtcaga cagtgatcag ggatgtagaa cagcagttca aatacacaca gccaaacatc	600
tgccgaaatc cagtttgtgc caacaggagg agattcttac tggatacaaa taaatcaaga	660
tttgttgatt ttcaaaagg tctgattcaa gagacccaag ctgagcttcc tcgagggagt	720
atcccccgca gtttagaagt aattttaagg gctgaagctg tggaaatcagc tcaagctggt	780
gacaagtgtg actttacagg gacactgatt gttgtgcctg acgtctccaa gcttagcaca	840
ccaggagcac gtgcagaaac taattcccgt gtcagtggtg ttgatggata tgagacagaa	900
ggcattcgag gactccgggc ccttggtgtt agggaccttt cttataggct ggtctttctt	960
gcctgctgtg ttgcgccaac caaccaagg ttgggggga aagagctcag agatgaggaa	1020
cagacagctg agagcattaa gaaccaaatg actgtgaaag aatgggagaa agtgtttgag	1080
atgagtcaag ataaaaatct ataccacaat ctttgtacca gcctgttccc tactatacat	1140
ggcaatgatg aagtaaaacg ggggtgcctg ctgatgctct ttggtggcgt tccaaagaca	1200
acaggagaag ggacctctct tcgaggggac ataaatgttt gcattgttgg tgaccaagt	1260
acagctaaga gccaatctct caagcacgtg gaggagtcca gcccagagc tgtctacacc	1320
agtggtaag cgtccagtgc tgctggctta acagcagctg ttgtgagaga tgaagaatct	1380
catgagtttg tcattgaggc tggagctttg atgttggctg ataatggtgt gtgttgatt	1440
gatgaatttg ataagatgga cgtgcccgat caagtgtcta ttcataagc tatggaacag	1500
cagaccatat ccatcactaa agcaggagtg aaggctactc tgaacgcccg gacgtccatt	1560
ttggcagcag caaaccaat cagtggacac tatgacagat caaaatcatt gaaacagaat	1620
ataaatttgt cagctcccat catgtcccga ttcgatctct tctttatcct tgtggatgaa	1680
tgtaatgagg ttacagatta tgccattgcc aggcgcatag tagatttgca ttcaagaatt	1740
gaggaaatcaa ttgatcgtgt ctattccctc gatgatatca gaagatatct tctctttgca	1800
agacagttta aacccaagat ttccaaagag tcagaggact tcattgtgga gcaatataaa	1860
catctccgcc agagagatgg ttctggagtg accaagtctt catggaggat tacagtgcga	1920
cagcttgaga gcatgattcg tctctctgaa gctatggctc ggatgcactg ctgtgatgag	1980
gtccaaccta aacatgtgaa ggaagcttc cggttactga ataaatcaat catccgtgtg	2040
gaaacacctg atgtcaatct agatcaagag gaagagatcc agatggagggt agatgagggt	2100
gctggtggca tcaatggtca tgctgacagc cctgctcctg tgaacgggat caatggctac	2160
aatgaagaca taaatcaaga gctgctccc aaagcctcct taaggctggg cttctctgag	2220
tactgccgaa tctctaacct tattgtgctt cacctcagaa aggtggaaga agaagaggac	2280
gagtcagcat taaagaggag cgagcttgtt aactggtact tgaaggaaat cgaatcagag	2340
atagactctg aagaagaact tataaataaa aaaagaatca tagagaaagt tattcatcga	2400
ctcacacact atgatcatgt tctaattgag ctcaccagc ctggattgaa aggctccaca	2460
gagggaagtg agagctatga agaagatccc tacttggtag ttaaccctaa ctacttgctc	2520
gaagatttgc aactgcagaa aagggcgaat tctgcagata tccatcacac tggcggccgc	2580
tcgagcatgc	2590

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 821

&lt;212&gt; TYPE: PR1

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

-continued

---

 <223> OTHER INFORMATION: Amino acid sequence for the FLAG-tagged MCM6 polypeptide encoded by SEQ ID NO:9

&lt;400&gt; SEQUENCE: 10

Met Asp Leu Ala Ala Ala Ala Glu Pro Gly Ala Gly Ser Gln His Leu  
 1 5 10 15  
 Glu Val Arg Asp Glu Val Ala Glu Lys Cys Gln Lys Leu Phe Leu Asp  
 20 25 30  
 Phe Leu Glu Glu Phe Gln Ser Ser Asp Gly Glu Ile Lys Tyr Leu Gln  
 35 40 45  
 Leu Ala Glu Glu Leu Ile Arg Pro Glu Arg Asn Thr Leu Val Val Ser  
 50 55 60  
 Phe Val Asp Leu Glu Gln Phe Asn Gln Gln Leu Ser Thr Thr Ile Gln  
 65 70 75 80  
 Glu Glu Phe Tyr Arg Val Tyr Pro Tyr Leu Cys Arg Ala Leu Lys Thr  
 85 90 95  
 Phe Val Lys Asp Arg Lys Glu Ile Pro Leu Ala Lys Asp Phe Tyr Val  
 100 105 110  
 Ala Phe Gln Asp Leu Pro Thr Arg His Lys Ile Arg Glu Leu Thr Ser  
 115 120 125  
 Ser Arg Ile Gly Leu Leu Thr Arg Ile Ser Gly Gln Val Val Arg Thr  
 130 135 140  
 His Pro Val His Pro Glu Leu Val Ser Gly Thr Phe Leu Cys Leu Asp  
 145 150 155 160  
 Cys Gln Thr Val Ile Arg Asp Val Glu Gln Gln Phe Lys Tyr Thr Gln  
 165 170 175  
 Pro Asn Ile Cys Arg Asn Pro Val Cys Ala Asn Arg Arg Arg Phe Leu  
 180 185 190  
 Leu Asp Thr Asn Lys Ser Arg Phe Val Asp Phe Gln Lys Val Arg Ile  
 195 200 205  
 Gln Glu Thr Gln Ala Glu Leu Pro Arg Gly Ser Ile Pro Arg Ser Leu  
 210 215 220  
 Glu Val Ile Leu Arg Ala Glu Ala Val Glu Ser Ala Gln Ala Gly Asp  
 225 230 235 240  
 Lys Cys Asp Phe Thr Gly Thr Leu Ile Val Val Pro Asp Val Ser Lys  
 245 250 255  
 Leu Ser Thr Pro Gly Ala Arg Ala Glu Thr Asn Ser Arg Val Ser Gly  
 260 265 270  
 Val Asp Gly Tyr Glu Thr Glu Gly Ile Arg Gly Leu Arg Ala Leu Gly  
 275 280 285  
 Val Arg Asp Leu Ser Tyr Arg Leu Val Phe Leu Ala Cys Cys Val Ala  
 290 295 300  
 Pro Thr Asn Pro Arg Phe Gly Gly Lys Glu Leu Arg Asp Glu Glu Gln  
 305 310 315 320  
 Thr Ala Glu Ser Ile Lys Asn Gln Met Thr Val Lys Glu Trp Glu Lys  
 325 330 335  
 Val Phe Glu Met Ser Gln Asp Lys Asn Leu Tyr His Asn Leu Cys Thr  
 340 345 350  
 Ser Leu Phe Pro Thr Ile His Gly Asn Asp Glu Val Lys Arg Gly Val  
 355 360 365  
 Leu Leu Met Leu Phe Gly Gly Val Pro Lys Thr Thr Gly Glu Gly Thr  
 370 375 380

-continued

---

Ser Leu Arg Gly Asp Ile Asn Val Cys Ile Val Gly Asp Pro Ser Thr  
 385 390 395 400  
 Ala Lys Ser Gln Phe Leu Lys His Val Glu Glu Phe Ser Pro Arg Ala  
 405 410 415  
 Val Tyr Thr Ser Gly Lys Ala Ser Ser Ala Ala Gly Leu Thr Ala Ala  
 420 425 430  
 Val Val Arg Asp Glu Glu Ser His Glu Phe Val Ile Glu Ala Gly Ala  
 435 440 445  
 Leu Met Leu Ala Asp Asn Gly Val Cys Cys Ile Asp Glu Phe Asp Lys  
 450 455 460  
 Met Asp Val Arg Asp Gln Val Ala Ile His Glu Ala Met Glu Gln Gln  
 465 470 475 480  
 Thr Ile Ser Ile Thr Lys Ala Gly Val Lys Ala Thr Leu Asn Ala Arg  
 485 490 495  
 Thr Ser Ile Leu Ala Ala Ala Asn Pro Ile Ser Gly His Tyr Asp Arg  
 500 505 510  
 Ser Lys Ser Leu Lys Gln Asn Ile Asn Leu Ser Ala Pro Ile Met Ser  
 515 520 525  
 Arg Phe Asp Leu Phe Phe Ile Leu Val Asp Glu Cys Asn Glu Val Thr  
 530 535 540  
 Asp Tyr Ala Ile Ala Arg Arg Ile Val Asp Leu His Ser Arg Ile Glu  
 545 550 555 560  
 Glu Ser Ile Asp Arg Val Tyr Ser Leu Asp Asp Ile Arg Arg Tyr Leu  
 565 570 575  
 Leu Phe Ala Arg Gln Phe Lys Pro Lys Ile Ser Lys Glu Ser Glu Asp  
 580 585 590  
 Phe Ile Val Glu Gln Tyr Lys His Leu Arg Gln Arg Asp Gly Ser Gly  
 595 600 605  
 Val Thr Lys Ser Ser Trp Arg Ile Thr Val Arg Gln Leu Glu Ser Met  
 610 615 620  
 Ile Arg Leu Ser Glu Ala Met Ala Arg Met His Cys Cys Asp Glu Val  
 625 630 635 640  
 Gln Pro Lys His Val Lys Glu Ala Phe Arg Leu Leu Asn Lys Ser Ile  
 645 650 655  
 Ile Arg Val Glu Thr Pro Asp Val Asn Leu Asp Gln Glu Glu Ile  
 660 665 670  
 Gln Met Glu Val Asp Glu Gly Ala Gly Gly Ile Asn Gly His Ala Asp  
 675 680 685  
 Ser Pro Ala Pro Val Asn Gly Ile Asn Gly Tyr Asn Glu Asp Ile Asn  
 690 695 700  
 Gln Glu Ser Ala Pro Lys Ala Ser Leu Arg Leu Gly Phe Ser Glu Tyr  
 705 710 715 720  
 Cys Arg Ile Ser Asn Leu Ile Val Leu His Leu Arg Lys Val Glu Glu  
 725 730 735  
 Glu Glu Asp Glu Ser Ala Leu Lys Arg Ser Glu Leu Val Asn Trp Tyr  
 740 745 750  
 Leu Lys Glu Ile Glu Ser Glu Ile Asp Ser Glu Glu Glu Leu Ile Asn  
 755 760 765  
 Lys Lys Arg Ile Ile Glu Lys Val Ile His Arg Leu Thr His Tyr Asp  
 770 775 780

-continued

---

His	Val	Leu	Ile	Glu	Leu	Thr	Gln	Ala	Gly	Leu	Lys	Gly	Ser	Thr	Glu
785					790					795					800
Gly	Ser	Glu	Ser	Tyr	Glu	Glu	Asp	Pro	Tyr	Leu	Val	Val	Asn	Pro	Asn
				805					810					815	
Tyr	Leu	Leu	Glu	Asp											
			820												

---

That which is claimed:

1. A method for diagnosing high-grade cervical disease in a patient, the method comprising:

- a) contacting a cervical sample obtained from said patient with at least one monoclonal antibody that specifically binds to MCM7 and is selected from the group consisting of:
  - i) the monoclonal antibody produced by the hybridoma cell line 2E6.2, deposited with the ATCC as Patent Deposit No. PTA-6669;
  - ii) a monoclonal antibody that binds to an epitope capable of binding the monoclonal antibody produced by the hybridoma cell line 2E6.2;
  - iii) a monoclonal antibody that binds to an epitope consisting of the amino acid sequence set forth in SEQ ID NO: 6;
  - iv) a monoclonal antibody that competes in a competitive binding assay with the monoclonal antibody produced by the hybridoma cell line 2E6.2; and,
  - v) a monoclonal antibody that is an antigen binding fragment of a monoclonal antibody of (i)-(iv),

wherein the fragment retains the capability of specifically binding to MCM7; and,

- b) detecting binding of the antibody to MCM7 to determine if MCM7 is overexpressed in said cervical sample compared to a normal cervical sample, wherein overexpression is indicative of high-grade cervical disease.

2. The method of claim 1, wherein the monoclonal antibody is the monoclonal antibody produced by the hybridoma cell line 2E6.2, deposited with the ATCC as Patent Deposit No. PTA-6669.

3. The method of claim 1, further comprising contacting the sample with at least one monoclonal antibody that specifically binds to MCM6.

4. The method of claim 3, wherein the monoclonal antibody that specifically binds to MCM6 is the monoclonal antibody produced by the hybridoma cell line 9D4.3, deposited with the ATCC as Patent Deposit No. PTA-6911.

5. The method according to claim 3, wherein the antibodies are contacted with the sample sequentially as individual antibody reagents or as an antibody cocktail.

\* \* \* \* \*