

(19)



(11)

**EP 4 342 895 A1**

(12)

**EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 153(4) EPC

(43) Date of publication:

**27.03.2024 Bulletin 2024/13**

(21) Application number: **22793620.0**

(22) Date of filing: **16.05.2022**

(51) International Patent Classification (IPC):

**C07D 413/04** <sup>(2006.01)</sup>    **A61K 31/506** <sup>(2006.01)</sup>  
**A61K 31/5377** <sup>(2006.01)</sup>    **A61K 31/541** <sup>(2006.01)</sup>  
**A61P 35/00** <sup>(2006.01)</sup>    **C07D 413/14** <sup>(2006.01)</sup>  
**C07D 471/04** <sup>(2006.01)</sup>

(52) Cooperative Patent Classification (CPC):

**C07D 413/04; A61K 31/495; A61K 31/506;**  
**A61K 31/5377; A61K 31/541; A61K 31/55;**  
**A61P 35/00; C07D 401/14; C07D 413/14;**  
**C07D 471/04; C07D 471/10; C07D 487/04;**  
**C07D 487/08; C07D 487/10; C07D 498/04;** (Cont.)

(86) International application number:

**PCT/KR2022/006994**

(87) International publication number:

**WO 2022/245085 (24.11.2022 Gazette 2022/47)**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB  
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO  
PL PT RO RS SE SI SK SM TR**

Designated Extension States:

**BA ME**

Designated Validation States:

**KH MA MD TN**

(30) Priority: **17.05.2021 KR 20210063733**

(71) Applicant: **Voronoi Inc.**

**Incheon 21984 (KR)**

(72) Inventors:

- **LEE, Youn Ho**  
**Incheon 21984 (KR)**
- **HWANG, Seon Ah**  
**Incheon 21984 (KR)**
- **SHIM, In Seob**  
**Incheon 21984 (KR)**

- **JEON, Hyeon Ho**  
**Incheon 21984 (KR)**
- **DO, Woo Mi**  
**Incheon 21984 (KR)**
- **RYU, Hee Sun**  
**Incheon 21984 (KR)**
- **SON, Jung Beom**  
**Incheon 21984 (KR)**
- **KIM, Nam Doo**  
**Incheon 21984 (KR)**
- **KIM, Sung Hwan**  
**Incheon 21984 (KR)**
- **JUNG, Hong Ryul**  
**Incheon 21984 (KR)**
- **LEE, Young Yi**  
**Incheon 21984 (KR)**

(74) Representative: **Viering, Jentschura & Partner**

**mbB**  
**Patent- und Rechtsanwälte**  
**Am Brauhaus 8**  
**01099 Dresden (DE)**

(54) **HETEROARYL DERIVATIVE COMPOUND AND USE THEREOF**

(57) The present disclosure relates to a heteroaryl derivative and uses thereof. The heteroaryl derivative of the present disclosure exhibits excellent inhibitory activ-

ity against EGFR and/or HER2, and thus may be usefully employed as a therapeutic agent for EGFR- and/or HER2-related diseases.

**EP 4 342 895 A1**

(52) Cooperative Patent Classification (CPC): (Cont.)  
**C07D 498/08; C07F 9/6584**

**Description**

[Technical field]

5 **[0001]** The present disclosure relates to a heteroaryl derivative compound and medicinal uses thereof. Specifically, the present disclosure relates to a heteroaryl derivative compound having EGFR and/or HER2 inhibitory activity.

[Background Art]

10 **[0002]** Protein kinases are involved in signaling pathways by acting as molecular switches, and the transition between the activity and inactivity of the target protein by kinase in the cell must be smoothly regulated. If the transition between the active and inactive states is abnormally regulated, intracellular signal transmission is excessively activated or deactivated, leading to uncontrollable cell division and proliferation. In particular, abnormal activation by mutation, amplification and/or overexpression of protein kinase genes causes the development and progression of various tumors or plays a  
15 decisive role in the onset of various diseases such as inflammatory diseases, degenerative brain diseases, and autoimmune diseases, and the like.

**[0003]** An epidermal growth factor receptor (EGFR), a receptor tyrosine kinase in the ErbB family, is abnormally active in many epithelial cell tumors, including non-small cell lung cancer (NSCLC), breast cancer, glioma, squamous cell carcinoma of the head and neck, colorectal cancer, rectal adenocarcinoma, head and neck cancer, gastric and prostate  
20 cancer. It has been known that the activation of the EGFR-tyrosine kinase causes continuous cell proliferation, invasion into surrounding tissues, remote metastasis, and angiogenesis, and increases cell survival.

**[0004]** In addition, EGFR Del19 or EGFR L858R, an EGFR mutation, has been known to be a major cause of non-small cell lung cancer and head and neck cancer, and Iressa and Tarceva, which are therapeutic agents for these mutations, have been developed and are currently being used in clinical practice. However, when these drugs were  
25 used in patients, acquired resistance that caused EGFR secondary mutations based on the structure of the drug was observed, and it was also found that this is a real major cause of drug resistance. When the first-generation EGFR inhibitors are used for an average of about 10 months, acquired resistance called the T790M mutation located in the gatekeeper of the EGFR kinase occurs, and thus the first-generation EGFR inhibitors are not effective. In other words,  
30 EGFR Del19/T790M or EGFR L858R/T790M double mutation occurs, which prevents existing therapeutic agents from exhibiting drug efficacy. Osimertinib, a third-generation EGFR-TKI target drug that exhibits high reactivity to drug resistance according to EGFR T790M mutation, has been developed, but it has also been reported to cause drug resistance (Niederst MJ. et al., Clin Cancer Res, 2015, 17(21):3924-3933). EGFR C797S mutation has been suggested as one of the main mechanisms causing drug resistance to osimertinib, and it has been reported that about 40% of clinical trial patients have EGFR C797S mutation (Thress KS. et al., Nature Medicine, 2015, 21:560-562). Thus, EGFR Del19/C797S  
35 (EGFR DC) or EGFR L858R/C797S (EGFR LC) may be the main target.

**[0005]** Further, L861Q, G719A, S768I, L718Q, G724S, or the like, expressing EGFR rare (or uncommon) and drug-resistant mutations may also be potential targets.

**[0006]** Meanwhile, HER2 (human epidermal growth factor receptor 2; also called ErbB2), which is a receptor tyrosine kinase of the ErbB family, forms homodimers or heterodimers with other EGFR receptors, HER1 (EGFR, ErbB1), HER3  
40 (ErbB3) or HER4 (ErbB4), and is activated by autophosphorylation at intracellular tyrosine residues, which plays an important role in cell proliferation, differentiation and survival in normal cells and cancer cells (Di Fiore PP. et al., Science. 1987, 237(481):178-182). HER2 is known to be overexpressed in several carcinomas such as breast, gastric and ovarian cancer (Hardwick RH. et al., Eur. J Surg Oncol. 1997, 23(1):30-35; Korkaya H. et al., Oncogene. 2008, 27(47):6120-6130;).

45 **[0007]** As described above, there is an increasing unmet need for a novel compound that is capable of being usefully employed for the treatment of EGFR- and/or HER2-related diseases by regulating EGFR activities (particularly C797S mutations such as EGFR Del19/C797S and EGFR L858R/C797S, EGFR rare mutations, drug resistance mutations, or the like) and/or HER2.

50 [Disclosure]

[Technical Problem]

55 **[0008]** An object of the present disclosure is to provide a heteroaryl derivative having a novel structure, an optical isomer thereof, or a pharmaceutically acceptable salt thereof.

**[0009]** Another object of the present disclosure is to provide a method for preparing the heteroaryl derivative compound.

**[0010]** Still another object of the present disclosure is to provide a pharmaceutical use of the heteroaryl derivative compound, and specifically, to a pharmaceutical composition for the treatment or prevention of EGFR- and/or HER2-

related diseases comprising the heteroaryl derivative compound as an active ingredient, use of the compound for the treatment or prevention of EGFR- and/or HER2-related diseases, or a method for treating or preventing EGFR- and/or HER2-related diseases comprising administering the compound.

5 [Technical Solution]

[0011] In order to achieve the above-described objects, the present inventors made efforts to study, and as a result, found that the following heteroaryl derivative compounds represented by Chemical Formula 1 inhibited the proliferation of EGFR- and/or HER2-activated cells, and completed the present disclosure.

10

### Heteroaryl derivative compound

[0012] The present disclosure provides a compound represented by the following Chemical Formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof:


15

[Chemical Formula 1]

20

25

30



35

in the Chemical Formula 1,

$X_1$  to  $X_3$  are each independently CH or N;

$R_X$  is -H,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ aminoalkyl,  $-NH_2$ ,  $-NH(-C_{1-6}alkyl)$ , or  $-N(-C_{1-6}alkyl)(-C_{1-6}alkyl)$ ;

40

Y is  $-C_{1-6}alkyl$ ,  $-(CH_2)_n$ aryl,  $-(CH_2)_n$ hydroaryl,  $-(CH_2)_n$ heteroaryl, or  $-(CH_2)_n$ hydroheteroaryl in which at least one H of the  $-(CH_2)_n$ aryl,  $-(CH_2)_n$ hydroaryl,  $-(CH_2)_n$ heteroaryl, or  $-(CH_2)_n$ hydroheteroaryl ring may be substituted with  $-C_{1-6}alkyl$ ,  $-C_{1-6}aminoalkyl$ ,  $-C_{1-6}hydroxyalkyl$ ,  $-C_{1-6}haloalkyl$ ,  $-C_{1-6}alkyl-O-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-CN$ ,  $-(C=O)NR_1R_2$ ,  $-(C=O)OR_3$ ,  $-NR_4R_5$ ,  $-OR_6$ , -halo, =O, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the cycloalkyl, heterocycloalkyl, aryl, or heteroaryl may be substituted with  $-C_{1-6}alkyl$ ,  $-C_{1-6}haloalkyl$ , or -halo;

45

n is 0, 1, 2, 3, or 4;

$R_1$  to  $R_3$  are each independently -H,  $-C_{1-6}alkyl$ , or cycloalkyl;

$R_4$  and  $R_5$  are each independently -H or  $-C_{1-6}alkyl$ ;

$R_6$  is -H,  $-C_{1-6}alkyl$ , or phenyl in which at least one H of the phenyl ring may be substituted with  $-C_{1-6}alkyl$ ,  $-C_{1-6}haloalkyl$ , or halo;

50

$R_{Y1}$  to  $R_{Y5}$  are each independently -H or  $-C_{1-6}alkyl$ , or  $R_{Y2}$  and  $R_{Y3}$  may be linked to each other to form cycloalkyl or heterocycloalkyl,  $R_{Y4}$  and  $R_{Y5}$  may be linked to each other to form cycloalkyl or heterocycloalkyl, and  $R_{Y3}$  and  $R_{Y4}$  may be linked to each other to form aryl or heteroaryl;

L is  $-(CH_2)_m-$ ,  $-C(=O)-$ , or null;

m is 0, 1, 2, 3, or 4;

55

a ring Z is aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be substituted with  $-C_{1-6}alkyl$ ,  $-C_{1-6}cyanoalkyl$ ,  $-C_{1-6}hydroxyalkyl$ ,  $-C_{1-6}aminoalkyl$ ,  $-C_{1-6}haloalkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-CN$ ,  $-NR_7R_8$ ,  $-OH$ ,  $-O-C_{1-6}alkyl$ ,  $-O-C_{1-6}haloalkyl$ ,  $-S-C_{1-6}alkyl$ ,  $-S-C_{1-6}haloalkyl$ ,  $-C(=O)-C_{1-6}alkyl$ ,  $-C(=O)-C_{1-6}haloalkyl$ ,  $-C(=O)O-$

C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or Z<sub>1</sub>, or two or more substituents of the aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -O-C<sub>1-6</sub>alkyl, -halo, or Z<sub>1</sub>;

R<sub>7</sub> and R<sub>8</sub> are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -(C=O)-C<sub>1-6</sub>alkyl, or -(C=O)-C<sub>1-6</sub>haloalkyl;

Z<sub>1</sub> is cycloalkyl, heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -(C=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-C(=O)-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>9</sub>R<sub>10</sub>, -halo, cycloalkyl, or Z<sub>2</sub>;

R<sub>9</sub> and R<sub>10</sub> are each independently -H or -C<sub>1-6</sub>alkyl;

Z<sub>2</sub> is heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -(C=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>11</sub>R<sub>12</sub>, cycloalkyl, or Z<sub>3</sub>;

R<sub>11</sub> and R<sub>12</sub> are each independently -H or -C<sub>1-6</sub>alkyl; and

Z<sub>3</sub> is heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or cycloalkyl.

**[0013]** According to an embodiment of the present disclosure, the compound represented by Chemical Formula 1 above, an optical isomer thereof, or a pharmaceutically acceptable salt thereof may be included in the following scope:

X<sub>1</sub> to X<sub>3</sub> are each independently CH or N;

R<sub>X</sub> is -H, -NH<sub>2</sub>, -NH(-C<sub>1-6</sub>alkyl), or -N(-C<sub>1-6</sub>alkyl)(-C<sub>1-6</sub>alkyl);

Y is -C<sub>1-6</sub>alkyl, -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl in which at least one H of the -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -(C=O)NR<sub>1</sub>R<sub>2</sub>, -(C=O)OR<sub>3</sub>, -NR<sub>4</sub>R<sub>5</sub>, -OR<sub>6</sub>, -halo, =O, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the heterocycloalkyl, aryl, or heteroaryl may be substituted with -halo;

n is 0, 1, or 2;

R<sub>1</sub> to R<sub>3</sub> are each independently -H, -C<sub>1-6</sub>alkyl, or cycloalkyl;

R<sub>4</sub> and R<sub>5</sub> are each independently -H or -C<sub>1-6</sub>alkyl;

R<sub>6</sub> is -C<sub>1-6</sub>alkyl or phenyl in which at least one H of the phenyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, or halo;

R<sub>Y1</sub> to R<sub>Y5</sub> are each independently -H or -C<sub>1-6</sub>alkyl, or R<sub>Y2</sub> and R<sub>Y3</sub> may be linked to each other to form cycloalkyl, and R<sub>Y3</sub> and R<sub>Y4</sub> may be linked to each other to form aryl;

L is -(CH<sub>2</sub>)<sub>m</sub>-, -C(=O)-, or null;

m is 0, 1, or 2;

a ring Z is aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>cyanoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -NR<sub>7</sub>R<sub>8</sub>, -O-C<sub>1-6</sub>alkyl, -O-C<sub>1-6</sub>haloalkyl, -S-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>haloalkyl, -C(=O)O-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or Z<sub>1</sub>, or two or more substituents of the aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl or Z<sub>1</sub>;

R<sub>7</sub> and R<sub>8</sub> are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -(C=O)-C<sub>1-6</sub>alkyl, or -(C=O)-C<sub>1-6</sub>haloalkyl;

Z<sub>1</sub> is cycloalkyl, heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocyclo-

cloalkyl, -NH-heterocycloalkyl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>9</sub>R<sub>10</sub>, -halo, cycloalkyl, or Z<sub>2</sub>;

R<sub>9</sub> and R<sub>10</sub> are each independently -H or -C<sub>1-6</sub>alkyl;

Z<sub>2</sub> is heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>11</sub>R<sub>12</sub>, cycloalkyl, or Z<sub>3</sub>;

R<sub>11</sub> and R<sub>12</sub> are each independently -H or -C<sub>1-6</sub>alkyl; and

Z<sub>3</sub> is heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or cycloalkyl.

**[0014]** According to an embodiment of the present disclosure, the compound represented by Chemical Formula 1 above, an optical isomer thereof, or a pharmaceutically acceptable salt thereof may be included in the following scope:

X<sub>1</sub> is N;

X<sub>2</sub> and X<sub>3</sub> are each independently CH or N; and

R<sub>X</sub> is -H, -NH<sub>2</sub>, or -NH(-C<sub>1-6</sub>alkyl).

**[0015]** According to an embodiment of the present disclosure, the compound represented by Chemical Formula 1 above, an optical isomer thereof, or a pharmaceutically acceptable salt thereof may be included in the following scope:

Y is -C<sub>1-6</sub>alkyl, -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl in which at least one H of the -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkynyl, -CN, -(C=O)NH-cycloalkyl, -(C=O)O-C<sub>1-6</sub>alkyl, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -O(C<sub>1-6</sub>alkyl), -O-phenyl, -halo, =O, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the heterocycloalkyl, aryl, or heteroaryl may be substituted with -halo;

n is 0 or 1; and

R<sub>Y1</sub> to R<sub>Y5</sub> are each independently -H or -C<sub>1-6</sub>alkyl, or R<sub>Y2</sub> and R<sub>Y3</sub> may be linked to each other to form 3- to 6-membered cycloalkyl, and R<sub>Y3</sub> and R<sub>Y4</sub> may be linked to each other to form phenyl.

**[0016]** According to an embodiment of the present disclosure, the compound represented by Chemical Formula 1 above, an optical isomer thereof, or a pharmaceutically acceptable salt thereof may be included in the following scope:

L is -(CH<sub>2</sub>)<sub>m</sub>-, -C(=O)-, or null;

m is 0 or 1;

a ring Z is aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>cyanoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -NR<sub>7</sub>R<sub>8</sub>, -O-C<sub>1-6</sub>alkyl, -O-C<sub>1-6</sub>haloalkyl, -S-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>haloalkyl, -C(=O)O-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or Z<sub>1</sub>, wherein at least one H of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or -halo, or two or more substituents of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl or Z<sub>1</sub>;

R<sub>7</sub> and R<sub>8</sub> are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -(C=O)-C<sub>1-6</sub>alkyl, or -(C=O)-C<sub>1-6</sub>haloalkyl;

Z<sub>1</sub> is 3- to 7-membered cycloalkyl, 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the 5- to 7-membered

heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -halo, cycloalkyl, or Z<sub>2</sub>;

Z<sub>2</sub> is 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, and S, and at least one H of the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), 3- to 7-membered cycloalkyl, or Z<sub>3</sub>; and

Z<sub>3</sub> is 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, and S, and at least one H of the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or 3- to 7-membered cycloalkyl.

**[0017]** According to an embodiment of the present disclosure, the compound represented by Chemical Formula 1 above may be selected from the group consisting of compounds listed in Table 1 below.

**[0018]** In the present disclosure, unless otherwise specified, the term "alkyl" may refer to a straight or branched chain acyclic, cyclic, or saturated hydrocarbon to which they are bonded. For example, "C<sub>1-6</sub>alkyl" may indicate an alkyl containing 1 to 6 carbon atoms. As an example, acyclic alkyl may include, but is not limited to, methyl, ethyl, n-propyl, n-butyl, isopropyl, sec-butyl, isobutyl, tert-butyl, or the like. Cyclic alkyl may be used interchangeably with "cycloalkyl" as used herein, and as an example, may include, but is not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, or the like.

**[0019]** In the present disclosure, "alkoxy" may indicate -(O-alkyl) as an alkyl ether group, wherein alkyl is the same as defined above. For example, "C<sub>1-6</sub>alkoxy" may mean alkoxy containing C<sub>1-6</sub>alkyl, that is, -(O-C<sub>1-6</sub>alkyl), and as an example, may include, but is not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, or the like.

**[0020]** In the present disclosure, "halo" may be F, Cl, Br, or I.

**[0021]** As used herein, "haloalkyl" may mean a straight or branched chain alkyl (hydrocarbon) having one or more halo-substituted carbon atoms as defined herein. Examples of the haloalkyl may include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl or n-butyl independently substituted with one or more halogens, such as F, Cl, Br, or I.

**[0022]** As used herein, "hydroxyalkyl" may indicate a straight or branched chain alkyl (hydrocarbon) having a carbon atom substituted with hydroxy (OH). Examples of the hydroxyalkyl may include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl or n-butyl independently substituted with one or more -OH.

**[0023]** As used herein, "aminoalkyl" may mean a straight or branched chain alkyl (hydrocarbon) having a carbon atom substituted with amino (NR'R"). Here, R' and R" may be each independently selected from the group consisting of hydrogen and C<sub>1-6</sub>alkyl, and the selected R' and R" may be each independently substituted or unsubstituted.

**[0024]** As used herein, "cyanoalkyl" may indicate a straight or branched alkyl (hydrocarbon) having a carbon atom substituted with cyano (CN).

**[0025]** In the present disclosure, "heterocycloalkyl" may mean a ring containing at least one selected from N, O, P, P(=O), and S in the ring, and may be saturated or partially unsaturated. Here, when unsaturated, it may be referred to as a heterocycloalkene. Unless otherwise stated, heterocycloalkyl may be a single ring or a multiple ring such as a spiro ring, a bridged ring or a fused ring. In addition, "3- to 12-membered heterocycloalkyl" may indicate a heterocycloalkyl containing 3 to 12 atoms forming a ring. As an example, the heterocycloalkyl may include, but is not limited to, pyrrolidine, piperidine, imidazolidine, pyrazolidine, butyrolactam, valerolactam, imidazolidinone, hydantoin, dioxolane, phthalimide, piperidine, pyrimidin-2,4(1H,3H)-dione, 1,4-dioxane, morpholine, thiomorpholine, thiomorpholine-S-oxide, thiomorpholine-S,S-oxide, piperazine, pyran, pyridone, 3-pyrroline, thiopyran, pyrone, tetrahydrofuran, tetrahydrothiophene, quinclidine, tropane, 2-azaspiro[3.3]heptane, (1R,5S)-3-azabicyclo[3.2.1]octane, (1s,4s)-2-azabicyclo[2.2.2]octane, or (1R,4R)-2-oxa-5-azabicyclo[2.2.2]octane, and the like.



**[0026]** In the present disclosure, "arene" may mean an aromatic hydrocarbon ring. The arene may be a monocyclic arene or a polycyclic arene. The number of ring-forming carbons in the arene may be 5 or more and 30 or less, 5 or more and 20 or less, or 5 or more and 15 or less. Examples of the arene may include, but are not limited to, benzene, naphthalene, fluorene, anthracene, phenanthrene, bibenzene, terbenzene, quaterbenzene, quinquebenzene, sexibenzene, triphenylene, pyrene, benzofluoranthene, chrysene, and the like. In the present specification, the residue obtained by removing one hydrogen atom from "arene" is referred to as "aryl".

**[0027]** In the present disclosure, "heteroarene" may be a ring containing at least one of O, N, P, Si, and S as a heterogeneous element. The number of ring-forming carbons in the heteroarene may be 2 or more and 30 or less, or 2

or more and 20 or less. The heteroarene may be a monocyclic heteroarene or a polycyclic heteroarene. The polycyclic heteroarene may have, for example, a bicyclic or tricyclic structure. Examples of the heteroarene may include thiophene, purine, pyrrole, pyrazole, imidazole, thiazole, oxazole, isothiazole, oxadiazole, triazole, pyridine, bipyridyl, triazine, acridyl, pyridazine, pyrazine, quinoline, quinazoline, quinoxaline, phenoxazine, phthalazine, pyrimidine, pyridopyrimidine, pyridopyrazine, pyrazinopyrazine, isoquinoline, indole, carbazole, imidazopyridazine, imidazopyridine, imidazopyrimidine, pyrazolopyrimidine, imidazopyrazine or pyrazolopyridine, N-arylcarbazole, N-heteroarylcarbazole, N-alkylcarbazole, benzoxazole, benzoimidazole, benzothiazole, benzocarbazole, benzothiophene, dibenzothiophene, thienothiophene, benzofuran, phenanthroline, isoxazole, oxadiazole, thiadiazole, benzothiazole, tetrazole, phenothiazine, dibenzosilole, dibenzofuran, and the like, but are not limited thereto. In an embodiment of the present disclosure, heteroarene may also include bicyclic heterocyclo-arene containing heteroarene fused to an arene ring or a cycloalkyl ring fused to heterocycloalkyl rings. In the present specification, the residue obtained by removing one hydrogen atom from the "heteroarene" is referred to as "heteroaryl".

**[0028]** In the present disclosure, "hydroaryl" means that one or more double bonds present in "aryl" are substituted with a single bond.

**[0029]** In the present disclosure, "hydroheteroaryl" means that one or more double bonds present in "heteroaryl" are substituted with a single bond.

**[0030]** In the present disclosure, the term "optical isomers (enantiomers)" mean compounds of the present disclosure or salts thereof that have the same chemical formula or molecular formula but are different in stereostructure. Each of these enantiomers and mixtures thereof are also included within the scope of the present disclosure. Unless otherwise specified, the straight solid-line bond (-) connecting an asymmetric carbon atom may include a wedge-shaped solid-line bond () or a wedge-shaped dashed-line bond () indicating the absolute configuration of the stereocenter.

**[0031]** In the present disclosure, the term "cis" refers to a case in which the binding directions of two substituents in a ring are the same, and the term "trans" refers to a case in which the binding directions of two substituents in the ring are different.

**[0032]** The compound of Chemical Formula 1 of the present disclosure may exist in the form of a "pharmaceutically acceptable salt". As the salt, an acid addition salt formed by a pharmaceutically acceptable free acid is useful. The term "pharmaceutically acceptable salt" as used herein means any and all organic or inorganic acid addition salts of the compound represented by Chemical Formula 1 of which side effects caused by the salt do not reduce the beneficial efficacy of the compound at concentrations having an effective action that is relatively non-toxic and harmless to a patient.

**[0033]** Acid addition salts are prepared by conventional methods, for example by dissolving the compound in an excess of aqueous acid solution and precipitating the salt using a water-miscible organic solvent such as methanol, ethanol, acetone or acetonitrile. An acid or alcohol in an equimolar amount of the compound and water may be heated, and the mixture may then be evaporated to dryness, or the precipitated salt may be filtered off with suction.

**[0034]** Here, an organic acid and an inorganic acid may be used as the free acid, wherein the inorganic acid may be hydrochloric acid, phosphoric acid, sulfuric acid, nitric acid, or the like, and the organic acid may be methanesulfonic acid, p-toluenesulfonic acid, acetic acid, trifluoroacetic acid, maleic acid, succinic acid, oxalic acid, benzoic acid, tartaric acid, fumaric acid, mandelic acid, propionic acid, citric acid, lactic acid, glycolic acid, gluconic acid, galacturonic acid, glutamic acid, glutaric acid, glucuronic acid, aspartic acid, ascorbic acid, carbonic acid, vanillic acid, hydroiodic acid, or the like. However, the present disclosure is not limited thereto.

**[0035]** In addition, it is possible to prepare a pharmaceutically acceptable metal salt using a base. The alkali metal salt or alkaline earth metal salt is obtained, for example, by dissolving a compound in an excess of alkali metal hydroxide or alkaline earth metal hydroxide solution, filtering the undissolved compound salt, and then evaporating and drying the filtrate. Here, it is pharmaceutically suitable to prepare a sodium, potassium, or calcium salt as the metal salt, but the present disclosure is not limited thereto. Further, the corresponding silver salt may be obtained by reacting an alkali metal or alkaline earth metal salt with a suitable silver salt (for example, silver nitrate).


**[0036]** Unless otherwise indicated, the pharmaceutically acceptable salt of the present disclosure includes salts of acidic or basic groups that may be present in the compound of Chemical Formula 1. For example, the pharmaceutically acceptable salt may include sodium, calcium and potassium salts of hydroxyl groups, and the like, and as other pharmaceutically acceptable salts of amino groups, may include hydrobromide, sulfate, hydrogen sulfate, phosphate, hydrogen phosphate, dihydrogen phosphate, acetate, succinate, citrate, tartrate, lactate, mandelate, methanesulfonate (mesylate), and p-toluenesulfonate (tosylate) salts, and the like, and may be prepared by a method for preparing a salt known in the art.

### Use of Heteroaryl Derivative Compound

**[0037]** The present disclosure provides use of a compound represented by the following Chemical Formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof:



[Chemical Formula 1]



**[0038]** The compound represented by Chemical Formula 1 of the present disclosure, an optical isomer thereof, or a pharmaceutically acceptable salt thereof exhibits inhibitory activity against various kinases.

**[0039]** According to an embodiment of the present disclosure, the heteroaryl derivative represented by Chemical Formula 1 exhibits excellent inhibitory activity against EGFR and/or HER2 kinase, and thus may be usefully employed for the treatment or prevention of EGFR- and/or HER2-related diseases, in particular, cancer. Specifically, the compound of Chemical Formula 1 is able to inhibit EGFR and/or HER2 wild-type or mutant kinase, which is supported by the Experimental Examples to be described below. The EGFR mutation may be a C797S mutation such as EGFR Del19/C797S (EGFR DC) or EGFR L858R/C797S (EGFR LC), but is not limited thereto. In addition, the EGFR mutation may be EGFR L861Q, EGFR G719A, EGFR S768I, EGFR L718Q, or EGFR G724S, but is not limited thereto. Further, the EGFR mutation may be EGFR d746-750, EGFR d746-750/C797A, EGFR d746-750/C797S, EGFR d746-750/T790M/C797S, EGFR D761Y, EGFR G719C, EGFR G719D, EGFR G719S, EGFR L747S, EGFR L792F, EGFR L858R, or EGFR L792F/L858R, but is not limited thereto.

**[0040]** In the present disclosure, the cancer includes any cancer capable of exhibiting therapeutic or prophylactic efficacy due to inhibition of EGFR and/or HER2 kinase activity, and may be a solid cancer or a hematologic cancer. As a non-limiting example, the cancer may be one or more selected from the group consisting of pseudomyxoma, intrahepatic biliary tract cancer, hepatoblastoma, liver cancer, thyroid cancer, colon cancer, testicular cancer, myelodysplastic syndrome, glioblastoma, oral cancer, labial cancer, mycosis fungoides, acute myeloid leukemia, acute lymphocytic leukemia, basal cell cancer, ovarian epithelial cancer, ovarian germ cell cancer, male breast cancer, brain cancer, pituitary adenoma, multiple myeloma, gallbladder cancer, biliary tract cancer, colorectal cancer, chronic myelogenous leukemia, chronic lymphocytic leukemia, retinoblastoma, choroidal melanoma, ampulla of Vater cancer, bladder cancer, peritoneal cancer, parathyroid cancer, adrenal cancer, sinus cancer, non-small cell lung cancer, tongue cancer, astrocytoma, small cell lung cancer, pediatric brain cancer, pediatric lymphoma, pediatric leukemia, small intestine cancer, meningioma, esophageal cancer, glioma, renal pelvic cancer, kidney cancer, heart cancer, duodenal cancer, malignant soft tissue cancer, malignant bone cancer, malignant lymphoma, malignant mesothelioma, malignant melanoma, eye cancer, vulvar cancer, ureter cancer, urethral cancer, cancer of unknown primary site, gastric lymphoma, gastric cancer, gastric carcinoma, gastrointestinal stromal cancer, Wilms cancer, breast cancer, sarcoma, penile cancer, pharyngeal cancer, gestational choriocarcinoma, cervical cancer, endometrial cancer, uterine sarcoma, prostate cancer, metastatic bone cancer, metastatic brain cancer, mediastinal cancer, rectal cancer, rectal carcinoma, vaginal cancer, spinal cord cancer, acoustic tumor, pancreatic cancer, salivary gland cancer, Kaposi's sarcoma, Paget's disease, tonsil cancer, squamous cell carcinoma, lung adenocarcinoma, lung cancer, lung squamous cell carcinoma, skin cancer, anal cancer, rhabdomyosarcoma, laryngeal cancer, pleural cancer, blood cancer, and thymus cancer. The cancer includes not only primary cancer but also metastatic cancer.

**[0041]** According to an embodiment of the present disclosure, the present disclosure provides a pharmaceutical composition for treatment or prevention of EGFR- and/or HER2-related diseases containing the compound represented by Chemical Formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof as an active ingredient. Specifically, the EGFR- and/or HER2-related disease may be cancer. The types of cancer are the same as described above.

**[0042]** The pharmaceutical composition of the present disclosure may further include one or more active ingredients

exhibiting the same or similar drug efficacy in addition to the compound represented by Chemical Formula 1 above, the optical isomer thereof, or the pharmaceutically acceptable salt thereof.

[0043] The pharmaceutical composition of the present disclosure may be used in clinical administration, and may be prepared to be administered in various oral and parenteral formulations.

[0044] Further, according to an embodiment of the present disclosure, there is provided a method for treating or preventing EGFR- and/or HER2-related diseases, comprising: administering to a subject in need thereof a therapeutically effective amount of the compound represented by Chemical Formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof. The subject may be a mammal including a human.

[0045] The term "therapeutically effective amount" as used herein refers to an amount of the compound represented by Chemical Formula 1 that is effective for the treatment or prevention of EGFR- and/or HER2-related diseases. Specifically, "therapeutically effective amount" indicates an amount sufficient to treat a disease with a reasonable benefit/risk ratio applicable to medical treatment, and the effective dose level may be determined depending on factors including the subject type and severity, age, sex, type of disease, drug activity, drug sensitivity, administration time, administration route and excretion rate, treatment period, drugs used at the same time, and other factors well-known in medical fields.

The pharmaceutical composition of the present disclosure may be administered as an individual therapeutic agent or may be administered in combination with other therapeutic agents, and may be administered sequentially or simultaneously with commercially available therapeutic agents. In addition, the pharmaceutical composition of the present disclosure may be administered in a single dose or multiple doses. It is important to administer the minimum amount capable of obtaining the maximum effect without side effects in consideration of all of the above factors, and the amount may be readily determined by those skilled in the art. The dosage of the pharmaceutical composition of the present disclosure may be determined by a medical specialist according to various factors such as the patient's condition, age, sex, complications, and the like. Since the active ingredient of the pharmaceutical composition of the present disclosure has excellent safety, it may be used at a dose higher than the determined dosage.

[0046] Further, according to an embodiment of the present disclosure, the present disclosure provides use of the compound represented by Chemical Formula 1, the optical isomer thereof, or the pharmaceutically acceptable salt thereof for use in preparation of a medicament to treat or prevent EGFR- and/or HER2-related diseases. The compound represented by Chemical Formula 1 for preparing the medicament may be mixed with acceptable adjuvants, diluents, carriers, and the like, and may have a synergistic effect of active ingredients by being prepared as a complex formulation with other active agents.

[0047] Matters mentioned in the uses, compositions and treatment methods of the present disclosure are applied equally except to the extent that they are inconsistent with each other.

[Advantageous Effects]

[0048] The heteroaryl derivative compound of the present disclosure exhibits excellent inhibitory activity against EGFR and/or HER2, and thus may be usefully employed for the treatment or prevention of EGFR- and/or HER2-related diseases.

[Best Mode]

[0049] Hereinafter, the present disclosure will be described in more detail through Examples and Experimental Examples. However, the following Examples and Experimental Examples are merely presented to illustrate the present disclosure, and the content of the present disclosure is not limited thereto.

#### <Analysis and purification conditions>

[0050] Compounds synthesized in the Examples of the present disclosure were purified by the following HPLC conditions or subjected to structural analysis.

#### 1. HPLC conditions

##### Analytical HPLC Condition (ACQUITY UPLC H-Class Core System)

[0051] Waters UPLC system (ACQUITY UPLC PDA Detector) equipped with a mass QDa detector manufactured by Waters corporation was used. The column used was Waters ACQUITY UPLC® BEH C18 (1.7  $\mu$ m, 2.1  $\times$  50 mm), and performed at 30°C.

[0052] Water containing 0.1% formic acid was used as the mobile phase A and acetonitrile containing 0.1% formic acid was used as the mobile phase B.

[0053] Gradient conditions (10-100% B over 3 minutes and flow rate = 0.6 ml/min)

Prep-LCMS for purification (Preparative-Liquid chromatography mass spectrometry)

[0054] Waters Autopurification HPLC system (2767 sample manager, 2545 binary gradient module, 2998 Photodiode Array Detector) equipped with a mass QDa detector manufactured by Waters corporation was used. The column used was Waters SunFire®Prep C18 OBD™ (5 µm, 19 × 50mm), and performed at room temperature.

[0055] Water containing 0.035% trifluoroacetic acid was used as the mobile phase A and methanol containing 0.035% trifluoroacetic acid was used as the mobile phase B.

[0056] Gradient conditions (15-100% B over 3 minutes and flow rate = 25 ml/min)

Prep-150 LC System for purification (Preparative-Liquid chromatography UV spectrometry)

[0057] Waters Prep 150 LC system (2545 Quaternary gradient module, 2998 Photodiode Array Detector, Fraction collector III) was used. The column used was Waters XTERRA®Prep RP18 OBD™ (10 µm, 30 × 300 mm) and performed at room temperature.

[0058] Gradient conditions (3-100% B over 120 minutes and flow rate = 40 ml/min)

Preparative HPLC System for purification (Preparative-Liquid chromatography UV spectrometry)

[0059] Teledyne ACCQPrep HP150 was used. The column used was Waters XTERRA®Prep RP18 OBD™ (10 µm, 30 × 300 mm) and performed at room temperature.

[0060] Gradient conditions (10-100% B over 120 minutes and flow rate = 42 ml/min)


**2. NMR analysis**

[0061] NMR analysis was performed using AVANCE III 400 or AVANCE III 400 HD manufactured by Bruker, and data are expressed in parts per million (δ) (ppm).

[0062] Commercially available reagents were used without further purification. In the present disclosure, room temperature or ordinary temperature refers to a temperature of about 5°C to 40°C, for example, 10°C to 30°C, as another example, 20°C to 27°C, which is not strictly limited to the above scope. Concentration under reduced pressure or solvent distillation was performed using a rotary evaporator.

**<Preparation Example 1> Preparation of (S)-3-phenylisoxazolidine**

[0063]

Step 1: Preparation of tert-butyl (R)-(3-hydroxy-3-phenylpropoxy) carbamate

[0064] Tert-butyl hydroxycarbamate (7.8 g, 58.6 mmol) was dissolved in dimethylformamide (140 ml), then sodium hydride (2.58 g, 64.5 mmol) was added at 0°C and the reaction mixture was reacted for 30 min. Next, (R)-3-chloro-1-phenylpropan-1-ol (5 g, 29.3 mmol) dissolved in dimethylformamide (DMF; 10 ml) was slowly added dropwise at 0°C for 10 min, and stirred at room temperature for 72 hours. The reaction was terminated by adding an aqueous ammonium chloride solution to the reaction mixture, followed by extraction with ethyl acetate and brine to combine the organic layers. The organic layer was dried over sodium sulfate and concentrated under reduced pressure. The title compound (2.8 g, 68%) was obtained by purification using medium pressure liquid chromatography (ethyl acetate/n-hexane).

[0065] MS (m/z): 150.17 [M+1]<sup>+</sup>, UPLC r.t. (min): 1.51

Step 2 : Preparation of tert-butyl (S)-3-phenylisoxazolidin-2-carboxylate

**[0066]** Tert-Butyl (R)-(3-hydroxy-3-phenylpropoxy)carbamate (2.55 g, 9.54 mmol) obtained in Step 1 above and triethylamine (3.13 ml, 22.44 mmol) were dissolved in dichloromethane (250 ml) and then cooled to 0°C. Next, methanesulfonyl chloride (1 ml, 13 mmol) was added dropwise and the reaction mixture was reacted at 0°C for 2 hours. The reaction mixture was extracted with brine and dichloromethane, and the organic layers were combined. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound, which was used in the next reaction without purification.

**[0067]** MS (m/z): 194.13 [M+1]<sup>+</sup>, UPLC r.t. (min): 1.69


Step 3: Preparation of (S)-3-phenylisoxazolidine

**[0068]** Tert-butyl (S)-3-phenylisoxazolidin-2-carboxylate (2.3 g) obtained in Step 2 was dissolved in dichloromethane (90 ml), then trifluoroacetic acid (14 ml) was added and the reaction mixture was reacted at room temperature for 1 hour. The reaction mixture was neutralized with aqueous sodium bicarbonate solution, and then the organic layers were combined. The organic layer was dried over sodium sulfate, followed by concentration under reduced pressure. The title compound (1.3 g, 94 %) was obtained by purification using medium pressure liquid chromatography (tetrahydrofuran/n-hexane).

**[0069]** MS (m/z): 150.08 [M+1]<sup>+</sup>, UPLC r.t. (min): 0.72

**<Preparation Example 2> Preparation of (R)-3-phenylisoxazolidine**

**[0070]**




**[0071]** A compound of Preparation Example 2 was prepared in a manner similar to that of Preparation Example 1, and was used for the synthesis of the Example compound shown in Table 1 below.

**[0072]** MS (m/z): 150.08 [M+1]<sup>+</sup>, UPLC r.t. (min): 0.72

**<Preparation Example 3> Preparation of (R)-3-(3-fluorophenyl)isoxazolidine**

**[0073]**



#### Step 1: Preparation of 3-fluoro-N-methoxy-N-methylbenzamide

**[0074]** 3-Fluorobenzoic acid (90 g, 642.35 mmol, 1 eq.) was dissolved in pyridine (150 mL), and N-methoxy methanamine (75.19 g, 770.81 mmol, 1.2 eq, HCl) was added thereto. Then, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI; 147.77 g, 770.81 mmol, 1.2 eq.) was added at 15°C. The reaction mixture was stirred at 50°C for 30 min. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 3:1), all of the starting materials disappeared, and a new spot with low polarity was detected. The pyridine solvent was removed by concentration under reduced pressure, and the organic layer was extracted using dichloromethane (DCM; 500 mL), hydrochloric acid (500 mL, 2N), and brine (200 mL). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow oil (110 g, 600.50 mmol, 93.49% yield).

**[0075]**  $^1\text{H NMR}$  (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.47-7.40 (m, 1H), 7.39-7.38 (m, 2H), 7.14-7.13 (m, 1H), 3.54 (s, 3H), 3.45 (s, 3H).

#### Step 2: Preparation of 1-(3-fluorophenyl)prop-2-en-1-one

**[0076]** 3-Fluoro-N-methoxy-N-methyl-benzamide (110 g, 600.50 mmol, 1 eq.) obtained in Step 1 above was dissolved in tetrahydrofuran (THF; 1 L), then bromo(vinyl)magnesium (1M, 630.53 mL, 1.05 eq.) was added dropwise at 0°C. Next, the reaction mixture was stirred at 0°C for 30 min. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 4:1), all of the starting materials disappeared, and a new spot with low polarity was detected. The reaction was terminated by adding hydrochloric acid (4N, 500 mL), and the organic layer was extracted using methyl tert-butyl ether (MTBE; 2000 mL) and brine (500 mL). The organic layer was dried over sodium sulfate, followed by concentration under reduced pressure. The concentrated compound was purified by chromatography (petroleum ether/ethyl acetate = 30/1) to obtain the title compound as a yellow oil (80 g, 532.80 mmol, 88.73% yield).

**[0077]**  $^1\text{H NMR}$  (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.65 (m, 1H), 7.58-7.52 (m, 1H), 7.39 (m, 1H), 7.24-7.17 (m, 1H), 7.04 (dd,  $J$  = 17.2, 10.4 Hz, 1H), 6.39 (dd,  $J$  = 17.2, 1.6 Hz, 1H), 5.90 (dd,  $J$  = 10.4, 1.6 Hz, 1H).

#### Step 3: Preparation of 3-chloro-1-(3-fluorophenyl)propan-1-one

**[0078]** 1-(3-Fluorophenyl)prop-2-en-1-one (71 g, 472.86 mmol, 1.0 eq.) obtained in Step 2 above was dissolved in dichloromethane (DCM; 71 mL), and HCl/dioxane (4M, 295.54 mL, 2.5 eq.) was added at 0°C. Next, the reaction mixture was stirred at 15°C for 1.5 hours. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 10:1), all of the starting materials disappeared, and the title compound was detected. The reaction mixture was concentrated under reduced pressure, and dichloromethane (DCM; 450 mL) and water (200 mL \* 5) were added to extract the organic layer. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow solid (73 g, 391.19 mmol, 82.73% yield).

**[0079]**  $^1\text{H NMR}$  (400MHz, CHLOROFORM- $d$ )  $\delta$  7.78-7.72 (m, 1H), 7.69-7.60 (m, 1H), 7.53-7.44 (m, 1H), 7.37-7.24 (m, 1H), 3.93 (t,  $J$  = 6.8 Hz, 2H), 3.46 (t,  $J$  = 6.8 Hz, 2H).

## EP 4 342 895 A1

### Step 4: Preparation of (S)-3-chloro-1-(3-fluorophenyl)propan-1-ol

- 5 **[0080]** (3aR)-1-methyl-3,3-diphenyl-3a,4,5,6-tetrahydropyrrolo[1,2-c] [1,3,2]oxazabolol (1M, 32.15 mL, 0.1 eq.) was dissolved in tetrahydrofuran (THF; 1.2 L), then borane tetrahydrofuran (BH<sub>3</sub>-THF; 1M, 186.48 mL, 0.6 eq.) was added dropwise at 0°C under a nitrogen atmosphere. The reaction mixture was stirred at 0°C for 30 min. Next, 3-chloro-1-(3-fluorophenyl)propan-1-one (60 g, 309.02 mmol, 1 eq.) obtained in Step 3 above diluted in tetrahydrofuran was added dropwise to the reaction mixture at 0°C. The reaction mixture was stirred at 0°C for 30 min. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 5:1), all of the starting materials disappeared, and a spot of the title compound was detected. The reaction was terminated by adding methanol (100 mL) at 0°C, and then the solvent was distilled off under reduced pressure. The organic layer was extracted from the concentrated compound using dichloromethane (DCM; 100 mL \* 3) and ammonium chloride (NH<sub>4</sub>Cl) solution (300 mL). The organic layer was dried over sodium sulfate, followed by concentration under reduced pressure. The concentrated compound was purified using silica gel chromatography (petroleum ether (PE):ethyl acetate (EA) = 50:1 to 5:1) to obtain the title compound as a colorless oil (140 g, 664.2 mmol, 71.65% yield, 89.49 % purity, 65.5 % e.e).
- 15 **[0081]** <sup>1</sup>H NMR (400 MHz, CHLOROFORM-d) δ 7.33 (m, 1H), 7.16-7.07 (m, 2H), 7.02-6.96 (m, 1H), 4.96 (m, 1H), 3.75 (m, 1H), 3.57 (m, 1H), 2.26-2.15(m, 2H).

### Step 5: Preparation of tert-butyl (S)-(3-(3-fluorophenyl)-3-hydroxypropoxy) carbamate

- 20 **[0082]** Tert-butyl hydroxycarbamate (50.4 g, 378.52 mmol, 1.05 eq.) was dissolved in dimethylformamide (DMF; 500 mL), and then sodium hydride (NaH; 15.86 g, 396.55 mmol, 60% purity, 1.1 eq.) was added at 0°C under a nitrogen atmosphere. The reaction mixture was stirred at 10°C for 1 hour, and (S)-3-chloro-1-(3-fluorophenyl)propan-1-ol (68 g, 360.5 mmol, 1 eq.) obtained in step 4 above diluted in dimethylformamide (DMF; 180 mL) was added dropwise at 0°C and stirred at 10°C for 16 hours. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 2:1), all of the starting materials disappeared, and the title compound was detected. After the reaction was terminated by adding an aqueous ammonium chloride solution (3 L), the organic layer was extracted using ethyl acetate (2000 mL) and brine (2000 mL). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a light yellow solid (176 g, 616.87 mmol, 85.56% yield).
- 25 **[0083]** <sup>1</sup>H NMR (400 MHz, CHLOROFORM-d) δ 7.67-7.64 (m, 1H), 7.23-7.17 (m, 1H), 7.08-7.03 (m, 2H), 6.88-6.81 (m, 1H), 4.99-4.84 (m, 1H), 4.02-3.97 (m, 1H), 3.96-3.89 (m, 1H), 1.95-1.89 (m, 1H), 1.88-1.78 (m, 1H), 1.42-1.39 (m, 9H).
- 30

### Step 6: Preparation of tert-butyl (R)-3-(3-fluorophenyl)isoxazolidin-2-carboxylate

- 35 **[0084]** Tert-butyl (S)-(3-(3-fluorophenyl)-3-hydroxypropoxy)carbamate (88 g, 308.44 mmol, 1 eq.) obtained in step 5 above and triethylamine (93.63 g, 925.31 mmol, 128.79 mL, 3 eq.) were dissolved in dichloromethane (DCM; 1 L), and then methanesulfonic anhydride (80.59 g, 462.65 mmol, 1.5 eq.) was added slowly at 0°C. The reaction mixture was stirred at 20°C for 12 hours. As a result of TLC analysis (petroleum ether (PE): ethyl acetate (EA) = 3:1), all of the starting materials disappeared, and a new spot was detected. After the reaction was terminated by adding water (2000 mL), the organic layer was extracted using dichloromethane (DCM; 200 mL \* 3). The organic layer was dried over sodium sulfate, followed by concentration under reduced pressure. The concentrated compound was purified by chromatography (petroleum ether (PE):ethyl acetate (EA) = 50:1 to 5:1) to extract 88 g of the title compound having an 82.5% e.e value. The title compound was purified through SFC (column: DAICEL CHIRALPAK AD (250 mm \* 50 mm, 10 μm); mobile phase: [Neu-MeOH]; B%: 15%-15%, 3.4 min; 380 min) to obtain the title compound as a white solid (51 g, 189.66 mmol, 30.74% yield, 99.4% purity).
- 40 **[0085]** The purity of the optical isomer of tert-butyl (R)-3-(3-fluorophenyl)isoxazolidin-2-carboxylate obtained in step 6 above was analyzed under SFC conditions as follows.

Instrument: CAS-WH-ANA-SFC-C(SHIMADZU LC-30ADsf)

Column: Amycoat 50 × 4.6 mm I.D., 3 μm

50 Mobile phase: Phase A for CO<sub>2</sub>, and Phase B for MeOH(0.05 % DEA) ;

Gradient elution: MeOH(0.05 % DEA) in CO<sub>2</sub> from 5% to 40%

Flow rate: 3 mL/min; Detector: PDA;

Column Temp: 35 °C; Back Pressure: 100 Bar

- 55 **[0086]** When the purity of the optical isomer of tert-butyl (R)-3-(3-fluorophenyl)isoxazolidin-2-carboxylate obtained in step 6 was low, the desired optical isomer as a yellow liquid was obtained by performing purification under SFC conditions as follows.

## EP 4 342 895 A1

Column: DAICEL CHIRALPAK AD-H(250 mm \* 30 mm, 5 um);  
Mobile phase: [0.1% NH<sub>3</sub>H<sub>2</sub>O MeOH]; B%: 15%-15%, 3.8 min; 600 min

### Step 7: Preparation of (R)-3-(3-fluorophenyl)isoxazolidine

[0087] Tert-butyl (R)-3-(3-fluorophenyl)isoxazolidin-2-carboxylate (50 g, 185.94 mmol, 1 eq.) obtained in step 6 above was dissolved in ethyl acetate (EA; 200 mL), and then HCl/EtOAc (4M, 300 mL, 6.45 eq.) was added at 0°C. Next, the reaction mixture was stirred at 10°C for 1 hour. As a result of LCMS analysis, all of the starting materials disappeared and the reaction mixture was concentrated under reduced pressure to obtain a solid to thereby yield the title compound as a white solid (32 g, 150.26 mmol, 80.81% yield, 95.62 % purity, 100 % e.e. HCl).

[0088] MS: m/z 168.2 [M+H]<sup>+</sup>

[0089] <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.53-7.43 (m, 2H), 7.39 (d, J = 7.8 Hz, 1H), 7.30-7.23 (m, 1H), 5.01 (t, J = 8.0 Hz, 1H), 4.47 (m, 1H), 4.27 (m, 1H), 2.87 (m, 1H), 2.62-2.52 (m, 1H).

[0090] The following conditions were employed for purification or analysis of optical isomers of the compound in step 7 above.


Instrument: CAS-WH-ANA-SFC-C(SHIMADZU LC-30ADsf)  
Column: Chiralpak AY-3 50 × 4.6 mm I.D., 3 um;  
Mobile phase: Phase A for CO<sub>2</sub>, and Phase B for IPA(0.05 % DEA) ;  
Gradient elution: B in A from 5% to 40%;  
Flow rate: 3mL/min; Detector: PDA;  
Column Temp: 35°C; Back Pressure: 100 Bar

### <Preparation Examples 4 to 44>

[0091] The following compounds of Preparation Examples 4 to 44 were prepared in a manner similar to those of Preparation Examples 1 to 3 above, and the compounds of Preparation Examples 1 to 44 were employed to prepare the Example compounds of the present disclosure.

### <Preparation Example 4> Preparation of (R)-3-(3,5-difluorophenyl)isoxazolidine


[0092]



[0093] <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.36-7.27 (m, 3H), 5.04-4.98 (t, J = 7.6 Hz, 1H), 4.46-4.36 (m, 1H), 4.25-4.19 (dd, J = 7.6, 15.2 Hz, 1H), 2.90-2.78 (m, 1H), 2.56-2.51 (m, 1H).

### <Preparation Example 5> Preparation of (R)-3-(2,5-difluorophenyl)isoxazolidine


[0094]



### <Preparation Example 6> Preparation of (R)-3-(4-fluorophenyl)isoxazolidine

[0095]

5




<Preparation Example 7> Preparation of (R)-3-(3-chloro-4-fluorophenyl)isoxazolidine

10

[0096]

15



20


[0097]  $^1\text{H NMR}$  (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  7.82-7.89 (dd,  $J = 2, 7.2$ , 1H), 7.56-7.51 (s,  $J = 15.6$ , 2H), 5.00-4.96 (m, 1H), 4.46-4.40 (m, 1H), 4.24-4.20 (m, 1H), 2.85-2.82 (m, 1H), 2.54-2.52 (m, 1H).

<Preparation Example 8> Preparation of (R)-3-(3-chloro-2-fluorophenyl)isoxazolidine

25

[0098]

30



35


[0099]  $^1\text{H NMR}$  (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  7.49-7.42 (m, 2H), 7.20-7.16 (m, 1H), 6.56 (s, 1H), 4.66-4.65 (m, 1H), 3.96-3.91 (m, 1H), 3.67-3.65 (m, 1H), 2.66-2.61 (m, 1H), 2.08-2.01 (m, 1H).

<Preparation Example 9> Preparation of (R)-3-(3-methoxyphenyl)isoxazolidine

40

[0100]

45



50

[0101]  $^1\text{H NMR}$  (400 MHz, Chloroform-d)  $\delta$  7.25-7.20 (m, 2H), 7.11-7.09 (m, 1H), 6.88-6.86 (m, 1H), 4.80-4.76 (m, 1H), 4.46-4.44 (m, 1H), 4.17-4.15 (m, 1H), 3.76 (s, 3H), 2.69-2.66 (m, 2H).


<Preparation Example 10> Preparation of (R)-3-(6-methylpyridin-3-yl)isoxazolidine

55

[0102]




5



<Preparation Example 11> Preparation of (R)-3-(3-ethynylphenyl)isoxazolidine

10 [0103]

15


20 [0104] <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.49 (s, 1H), 7.43-7.37 (m, 1H), 7.36-7.29 (m, 2H), 6.41 (s, 1H), 4.38 (s, 1H),

4.15 (s, 1H), 3.90 (m, 1H), 3.71 (s, 1H), 2.65-2.53 (m, 1H), 2.11-2.00 (m, 1H).

<Preparation Example 12> Preparation of (R)-3-methyl-3-phenylisoxazolidine

25 [0105]

25



30


35 [0106] <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 12.88 (br s, 1H), 7.56-7.46 (m, 2H), 7.44-7.36 (m, 2H), 7.34-7.26 (m, 1H), 3.74-3.62 (m, 1H), 3.46-3.28 (m, 1H), 2.72-2.54 (m, 2H), 1.64 (s, 3H).

35

<Preparation Example 13> Preparation of (R)-3-(2,4-difluorophenyl)isoxazolidine

40 [0107]

40



45


50 [0108] <sup>1</sup>H NMR (400 MHz, CHLOROFORM-d) δ 7.52-7.47 (m, 1H), 6.87-6.75 (m, 2H), 5.30 (s, 1H), 4.71-4.68 (m, 1H), 4.09-4.04 (m, 1H), 3.91-3.85 (m, 1H), 2.73-2.64 (3, 1H), 2.24-2.20 (m, 1H).

50

<Preparation Example 14> Preparation of (R)-3-(naphthalen-1-yl)isoxazolidine

55 [0109]


55



10 [0110]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  8.13 (br s, 1H), 7.89-7.88 (m, 1H), 7.87-7.78 (m, 2H), 7.55-7.48 (m, 3H), 5.54 (br s, 1H), 5.23-5.20 (m, 1H,  $J = 6.4$  Hz), 4.15-4.03 (m, 2H), 2.90-2.81 (m, 1H), 2.44-2.41 (m, 1H).

15 <Preparation Example 15> Preparation of (R)-3-(thiophen-2-yl)isoxazolidine


[0111]



25 [0112]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.23 (d,  $J = 5.0$  Hz, 1H), 7.04-6.94 (m, 2H), 4.97-4.58 (m, 2H), 4.11-3.96 (m, 2H), 2.75-2.58 (m, 1H), 2.44-2.33 (m, 1H).

30 <Preparation Example 16> Preparation of (R)-3-(3-(trifluoromethyl)phenyl)isoxazolidine


[0113]



40 [0114]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.65 (s, 1H), 7.59 (d,  $J = 7.7$  Hz, 1H), 7.53 (d,  $J = 7.8$  Hz, 1H), 7.46 (t,  $J = 7.7$  Hz, 1H), 5.64-5.19 (m, 1H), 4.58 (t,  $J = 7.2$  Hz, 1H), 4.11 (td,  $J = 8.2, 5.2$  Hz, 1H), 3.94 (s, 1H), 2.80-2.67 (m, 1H), 2.36-2.23 (m, 1H).

45 <Preparation Example 17> Preparation of (R)-3-(naphthalen-2-yl)isoxazolidine

[0115]




55 [0116]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.91-7.81 (m, 4H), 7.56-7.46 (m, 3H), 5.80-5.00 (m, 1H), 4.68 (t,  $J = 7.2$  Hz, 1H), 4.19-3.99 (m, 2H), 2.80-2.72 (m, 1H), 2.45-2.37 (m, 1H).

<Preparation Example 18> Preparation of (R)-3-(3,4-difluorophenyl)isoxazolidine

[0117]

5




10 [0118]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  7.24-7.19 (m, 1H), 7.12-7.06 (m, 2H), 5.24 (s, 1H), 4.46 (dd,  $J_1 = 8.4$  Hz,  $J_2 = 5.6$  Hz, 1H), 4.05 (dt,  $J_1 = 8.0$  Hz,  $J_2 = 5.2$  Hz, 1H), 3.91-3.85 (m, 1H), 2.70-2.61 (m, 1H), 2.25-2.17 (m, 1H).

<Preparation Example 19> Preparation of (R)-3-(2,3-difluorophenyl)isoxazolidine

15

[0119]



20


25 [0120]  $^1\text{H}$  NMR (CHLOROFORM- $d$ , 400 MHz)  $\delta$  7.29-7.27 (m, 1H), 7.06-7.02 (m, 2H), 5.44 (br s, 1H), 4.75 (dd,  $J_1 = 4.4$  Hz,  $J_2 = 8.4$  Hz, 1H), 4.08 (dt,  $J_1 = 5.2$  Hz,  $J_2 = 8.0$  Hz, 1H), 3.86 (q,  $J = 8.0$  Hz, 1H), 2.76-2.66 (m, 1H), 2.27-2.19 (m, 1H).

25

<Preparation Example 20> Preparation of (R)-3-(3-chloro-2,4-difluorophenyl)isoxazolidine

[0121]

30



35


40 [0122]  $^1\text{H}$  NMR (DMSO- $d_6$ , 400 MHz)  $\delta$  7.51 (dt,  $J = 6.8, 8.4$  Hz, 1H), 7.28 (dt,  $J = 2.0, 8.8$  Hz, 1H), 6.60 (br s, 1H), 4.64 (br s, 1H), 3.94 (dt,  $J = 5.2, 8.0$  Hz, 1H), 3.76-3.57 (m, 1H), 2.68-2.61 (m, 1H), 2.10-2.01 (m, 1H).

40

<Preparation Example 21> Preparation of (R)-3-(4-chloro-2-fluorophenyl)isoxazolidine

[0123]

45



50


55 [0124]  $^1\text{H}$  NMR (400 MHz, DEUTERIUM OXIDE)  $\delta$  7.48-7.38 (m, 1H), 7.34-7.22 (m, 2H), 5.29-5.20 (m, 1H), 4.58-4.50 (m, 1H), 4.36-4.27 (m, 1H), 2.96-2.84 (m, 1H), 2.79-2.66 (m, 1H).

55

<Preparation Example 22> Preparation of (R)-3-(4-chloro-3-fluorophenyl)isoxazolidine

[0125]

5




<Preparation Example 23> Preparation of (R)-3-(isoxazolidin-3-yl)-N,N-dimethylaniline

10

[0126]

15




20

<Preparation Example 24> Preparation of (S)-3-(5-fluoropyridin-3-yl)isoxazolidine

[0127]

25




30

<Preparation Example 25> Preparation of (R)-3-(5-fluoropyridin-3-yl)isoxazolidine

[0128]

35




40

<Preparation Example 26> Preparation of (R)-3-fluoro-5-(isoxazolidin-3-yl)benzotrile

[0129]

45




50

<Preparation Example 27> Preparation of (S)-3-fluoro-5-(isoxazolidin-3-yl)benzotrile

55

[0130]

5




<Preparation Example 28> Preparation of (R)-3-(1-methyl-1H-pyrazol-4-yl)isoxazolidine

10

[0131]

15




20

<Preparation Example 29> Preparation of (R)-3-(furan-2-yl)isoxazolidine

[0132]

25




30

<Preparation Example 30> Preparation of (R)-3-(5-chloropyridin-3-yl)isoxazolidine

[0133]

35




40

<Preparation Example 31> Preparation of (S)-3-(5-chloropyridin-3-yl)isoxazolidine

45

[0134]

50




55

<Preparation Example 32> Preparation of (R)-3-(3-difluoromethyl)phenyl)isoxazolidine

[0135]

5




<Preparation Example 33> Preparation of (S)-3-(3-(difluoromethyl)phenyl)isoxazolidine

10

[0136]

15




20

<Preparation Example 34> Preparation of (R)-3-(2-fluoro-3-(trifluoromethyl)phenyl)isoxazolidine

[0137]

25




30

<Preparation Example 35> Preparation of (S)-3-benzylisoxazolidine

35

[0138]

40




[0139]  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $d$ )  $\delta$  12.7-12.4 (m, 1H), 7.28-7.18 (m, 5H), 4.42-4.32 (m, 1H), 4.25-4.10 (m, 2H), 3.50 (dd,  $J$  = 4.8, 13.6 Hz, 1H), 3.03 (dd,  $J$  = 10.4, 13.2 Hz, 1H), 2.44-2.33 (m, 1H), 2.32-2.20 (m, 1H).

45

<Preparation Example 36> Preparation of isopropyl (R)-3-(isoxazolidin-3-yl)benzoate

[0140]

50




55

[0141] The desired title compound was prepared in a manner similar to that of <Preparation Example 3> using 3-(isopropoxycarbonyl)benzoic acid prepared in <Preparation Example 36-1> below as an intermediate.

<Preparation Example 36-1> Preparation of 3-(isopropoxycarbonyl)benzoic acid

[0142]




Step 1: Preparation of 3-(isopropoxycarbonyl)benzoic acid

[0143] Isophthalic acid (40 g, 1 eq.) was dissolved in isopropyl alcohol (150 mL) and tetrahydrofuran (THF; 450 mL), and then sulfuric acid (concentrated H<sub>2</sub>SO<sub>4</sub>; 38.5 mL, 3 eq.) was added. The reaction mixture was stirred at 75°C for 48 hours. After the reaction was completed, the organic solvent was subjected to concentration under reduced pressure, and the organic layer was extracted using ethyl acetate (EA; 500 mL) and water (200 mL). The organic layer was dried over sodium sulfate, followed by concentration under reduced pressure. The concentrated compound was purified by chromatography (dichloromethane/methanol = 8/1) to obtain the title compound as a clear oil (23.25 g, 46.5% yield).

<Preparation Example 37> Preparation of (R)-3-(isoxazolidin-3-yl)benzoic acid


[0144]



[0145] The desired title compound was prepared by hydrolyzing the isopropyl (R)-3-(isoxazolidin-3-yl)benzoate obtained in <Preparation Example 36> with an aqueous base solution.

<Preparation Example 38> Preparation of (R)-N-cyclohexyl-3-(isoxazolidin-3-yl)benzamide


[0146]



[0147] The desired title compound was prepared by introducing an amide functional group by a method such as HATU employing the (R)-3-(isoxazolidin-3-yl)benzoic acid obtained in <Preparation Example 37>.

<Preparation Example 39> Preparation of (R)-3-(3-(benzyloxy)phenyl)isoxazolidine


[0148]



[0149] The desired title compound was prepared in a manner similar to that of <Preparation Example 3> using 3-(benzyloxy)-N-methoxy-N-methylbenzamide prepared in <Preparation Example 39-1> below as an intermediate.

#### <Preparation Example 39-1> Preparation of 3-(benzyloxy)-N-methoxy-N-methylbenzamide

[0150]



#### Step 1: Preparation of methyl 3-(benzyloxy)benzoate

[0151] Methyl 3-hydroxybenzoate (20 g, 1.0 eq.) was dissolved in acetone (260 mL), and then (bromomethyl)benzene (18.76 ml 1.2 eq.) and potassium carbonate (54.5 g, 3 eq.) were added. The reaction mixture was stirred at 60°C for 16 hours. As a result of TLC analysis (hexane:ethyl acetate = 3:2), all of the starting materials disappeared and the title compound was detected. The reaction mixture was concentrated under reduced pressure, dichloromethane (DCM; 300 mL \* 2) and water (200 mL) were added to extract the organic layer. The organic layer was dried over sodium sulfate, and then concentrated under reduced pressure. The concentrated compound was recrystallized using hexane to obtain the title compound as a white solid (29.27 g, 92% yield).

#### Step 2: Preparation of 3-(benzyloxy)benzoic acid


[0152] Methyl 3-(benzyloxy)benzoate (29 g, 1.0 eq.) obtained in step 1 was dissolved in methanol (MeOH; 300 ml), and potassium hydroxide (KOH; 6M, 4.5 eq.) was added. Next, the reaction mixture was stirred at 80°C for 3 hours. As a result of TLC analysis (hexane:ethyl acetate = 7:3), all of the starting materials disappeared and a new spot with low polarity was detected. The reaction mixture was concentrated under reduced pressure, water (100 ml) was added and hydrochloric acid (3N) was added dropwise to acidify the reaction solution to pH 1. The resulting precipitate was filtered under reduced pressure and dried to obtain the title compound as a white solid (27 g, 99% yield).

#### Step 3: Preparation of 3-(benzyloxy)-N-methoxy-N-methylbenzamide

[0153] 3-(Benzyloxy)benzoic acid (20 g, 1.0 eq.) obtained in step 2 was dissolved in dichloromethane (DCM; 700 ml), and 1,1-carboxyldiimidazole (9.40 g, 1.1 eq.) was slowly added. The reaction mixture was stirred at room temperature for 2 hours, then N, O-dimethylhydroxylamine hydrochloride (15.63 g, 1.1 eq.) was added and the resulting mixture was stirred at 40°C for 18 hours. UPLC/MS analysis was performed, and as a result, all of the starting material disappeared, and the title compound was detected. The reaction mixture was washed with hydrochloric acid (1N, 500 ml) and a saturated aqueous sodium hydrogen carbonate solution (500 ml), dried over sodium sulfate, and then concentrated under reduced pressure to obtain the title compound as a pale yellow oil (90 g, 85% yield).

#### <Preparation Example 40> Preparation of (R)-3-(3-phenoxyphenyl)isoxazolidine

[0154]






[0155] The desired title compound was prepared in a manner similar to that of <Preparation Example 39>.

<Preparation Example 41> Preparation of (R)-3-(3-bromo-5-fluorophenyl)isoxazolidine hydrochloride

[0156]



Step 1: Preparation of 3-bromo-5-fluoro-N-methoxy-N-methylbenzamide

[0157] 3-Bromo-5-fluorobenzoic acid (10 g, 1 eq.) was dissolved in dichloromethane (110 mL), and then N,O-dimethylhydroxylamine hydrochloride (5.4 g, 1.2 eq.), triethylamine (TEA; 5.7 mL, 0.9 eq.), and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI; 11.5 g, 1.2 eq.) were added sequentially at room temperature. The reaction mixture was stirred at room temperature for 3 hours. As a result of TLC analysis (dichloromethane), all of the starting materials disappeared and a new spot with low polarity was detected. The organic layer was extracted using ethyl acetate (EA; 300 mL) and a saturated aqueous sodium hydrogen carbonate solution (400 mL \* 2). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow oil (11.2 g, 94.0% yield).

Step 2: Preparation of 1-(3-bromo-5-fluorophenyl)prop-2-en-1-one

[0158] 3-Bromo-5-fluoro-N-methoxy-N-methylbenzamide (11.2 g, 1 eq.) obtained in step 1 was dissolved in tetrahydrofuran (THF; 220 mL), and then bromo(vinyl)magnesium (0.7M, 93 mL, 1.5 eq.) was added dropwise at 0°C. Next, the reaction mixture was stirred at 0°C for 1 hour. As a result of TLC analysis (hexane:dichloromethane = 1:1), all of the starting materials disappeared and a new spot with low polarity was detected. The reaction was terminated by adding hydrochloric acid (1N, 50 mL), and then the organic layer was extracted using ethyl acetate (EA; 300 mL) and hydrochloric acid (1N, 400 mL \* 2). The organic layer was dried over sodium sulfate, and then concentrated under reduced pressure. The concentrated compound was purified by chromatography (hexane:dichloromethane = 1:1) to obtain the title compound as a colorless oil (7.9 g, 76% yield).

Step 3: Preparation of 1-(3-bromo-5-fluorophenyl)-3-chloropropan-1-ol

[0159] 1-(3-Bromo-5-fluorophenyl)prop-2-en-1-one (7.9 g, 1.0 eq.) obtained in Step 2 was dissolved in dichloromethane (DCM; 13 mL), and then HCl/dioxane (4M, 13 mL, 1.2 eq.) was added at 0°C. Then, the reaction mixture was stirred at room temperature for 12 hours. As a result of TLC analysis (hexane:dichloromethane = 1:1), all of the starting materials disappeared, and the title compound was detected. The reaction mixture was concentrated under reduced pressure, and ethyl acetate (EA; 300 mL) and a saturated aqueous sodium hydrogen carbonate solution (400 mL \* 2) were added to extract the organic layer. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow oil (8.9 g, 97% yield).

Step 4: Preparation of (S)-1-(3-bromo-5-fluorophenyl)-3-chloropropan-1-ol

[0160] (3aR)-1-methyl-3,3-diphenyl-3a,4,5,6-tetrahydropyrrolo[1,2-c][1,3,2]oxazabolol (1M, 6.7 mL, 0.2 eq.) was dissolved in tetrahydrofuran (THF; 84 mL), and then borane dimethylsulfide (BH<sub>3</sub>·Me<sub>2</sub>S; 1M, 21.8 mL, 1.3 eq.) was added

## EP 4 342 895 A1

dropwise at 0°C under a nitrogen atmosphere. The reaction mixture was stirred at 0°C for 30 min, and then 1-(3-bromo-5-fluorophenyl)-3-chloropropan-1-one (8.9 g, 1 eq.) obtained in step 3 and diluted in tetrahydrofuran was added dropwise at 0°C. The reaction mixture was stirred at 0°C for 2 hours. As a result of TLC analysis (hexane:dichloromethane = 1:1), all of the starting materials disappeared, and a spot of the title compound was detected. The reaction was terminated by adding methanol (20 mL) at 20°C, and then the solvent was distilled off under reduced pressure. The concentrated compound was treated with ethyl acetate (EA; 300 mL) and hydrochloric acid (1N, 400 mL \* 2) to extract the organic layer. The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow oil (8.5 g, 95% yield).

### Step 5: Preparation of tert-butyl (S)-(3-(3-bromo-5-fluorophenyl)-3-hydroxypropoxy)carbamate

**[0161]** Tert-butyl hydroxycarbamate (9.3 g, 2.2 eq.) was dissolved in dimethylformamide (DMF; 80 mL), and then sodium hydride (NaH; 3.1 g, 60% purity, 2.4 eq.) was added at 0°C under a nitrogen atmosphere. The reaction mixture was stirred at 0°C for 30 min. (S)-1-(3-bromo-5-fluorophenyl)-3-chloropropan-1-ol (8.5 g, 1 eq.) obtained in step 4 and diluted in dimethylformamide (DMF; 10 mL) was added dropwise at 0°C and stirred at room temperature for 12 hours. As a result of TLC analysis (dichloromethane:EA = 9:1), all of the starting materials disappeared, and the title compound was detected. The reaction was terminated by adding brine (50 mL), and then the organic layer was extracted using ethyl acetate (EA; 300 mL) and a saturated aqueous sodium hydrogen carbonate solution (400 mL \* 3). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a yellow oil (9.2 g, 79% yield).

### Step 6: Preparation of tert-butyl (R)-3-(3-bromo-5-fluorophenyl)isoxazolidin-2-carboxylate

**[0162]** Tert-butyl (S)-(3-(3-bromo-5-fluorophenyl)-3-hydroxypropoxy)carbamate (9.2 g, 1 eq.) obtained in step 5 and triphenylphosphine (Ph<sub>3</sub>P; 8.6 g, 1.3 eq.) were dissolved in dichloromethane (DCM; 110 mL), and then diisopropyl azodicarboxylate (DIAD; 6.6 g, 1.3 eq.) diluted in dichloromethane (DCM; 20 mL) was added slowly at 0°C. The reaction mixture was stirred at 0°C for 2 hours. As a result of TLC analysis (dichloromethane (DCM):ethyl acetate (EA) = 9:1), all of the starting materials disappeared, and a new spot was detected. The reaction mixture was concentrated under reduced pressure, and the concentrated compound was purified by chromatography (dichloromethane:ethyl acetate = 10:0 to 9:1) to obtain the title compound as a yellow oil (7.7 g, 88% yield).

### Step 7: Preparation of (R)-3-(3-bromo-5-fluorophenyl)isoxazolidine hydrochloride

**[0163]** Tert-butyl (R)-3-(3-bromo-5-fluorophenyl)isoxazolidin-2-carboxylate (7.7 g, 1 eq.) obtained in step 6 was dissolved in dichloromethane (DCM; 40 mL), and then HCl/dioxane (4M, 28 mL, 5 eq.) was added at room temperature. Then, the reaction mixture was stirred at room temperature for 2 hours. As a result of LCMS analysis, all of the starting materials disappeared, diethyl ether (200 mL) was added to obtain a solid, and the resulting precipitate was filtered and dried to obtain the title compound as a white solid (5.3 g, 84% yield).


**[0164]** The following conditions were employed for purification or analysis of optical isomers of the compound in step 7 above.

Instrument: CAS-WH-ANA-SFC-C(SHIMADZU LC-30ADsf)  
Column: Chiralpak AY-3 50 × 4.6mm I.D., 3 μm;  
Mobile phase: Phase A for CO<sub>2</sub>, and Phase B for IPA(0.05 % DEA) ;  
Gradient elution: B in A from 5% to 40%;  
Flow rate: 3 mL/min; Detector: PDA;  
Column Temp: 35 °C; Back Pressure: 100 Bar

**[0165]** (R)-3-(3-Bromo-5-fluorophenyl)isoxazolidine hydrochloride (5.3 g) obtained in step 7 was purified under SFC conditions as follows to obtain the desired optical isomer (100% purity, 100% e.e.).

Column: DAICEL CHIRALPAK AD-H(250 mm \* 30 mm, 5 μm);  
Mobile phase: [0.1% NH<sub>3</sub>H<sub>2</sub>O MeOH]; B%: 15%-15%, 3.8 min; 600 min

**[0166]** <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 7.62-7.58 (m, 2H), 7.43 (dt, J = 9.8, 2.0 Hz, 1H), 4.93 (t, J = 7.4 Hz, 1H), 4.36 (td, J = 8.1, 4.4 Hz, 1H), 4.12 (q, J = 7.8 Hz, 1H), 2.81 (dtd, J = 12.4, 7.9, 4.4 Hz, 1H), 2.49-2.41 (m, 1H).

**<Preparation Example 42> Preparation of (R)-3-(3',5-difluoro-[1,1'-biphenyl]-3-yl)isoxazolidine hydrochloride****[0167]****Step 1: Preparation of tert-butyl (R)-3-(3-bromo-5-fluorophenyl)isoxazolidin-2-carboxylate**

**[0168]** (R)-3-(3-Bromo-5-fluorophenyl)isoxazolidine hydrochloride (1 g, 1 eq.) obtained in <Preparation Example 41> and triethylamine (TEA; 1.5 mL, 3 eq.) were dissolved in tetrahydrofuran (7 mL), and then di-tert-butyl dicarbonate (Boc<sub>2</sub>O; 1.0 mL, 1.2 eq.) was slowly added at 0°C. Next, the reaction mixture was stirred at 50°C for 2 hours. As a result of TLC analysis (DCM), all of the starting materials disappeared and a new spot with different polarity was detected. The organic layer was extracted using ethyl acetate (EA; 70 mL) and a saturated aqueous sodium hydrogen carbonate solution (100 mL \* 2). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to obtain the title compound as a pale yellow oil (1.1 g, 95% yield).

15

20

**Step 2: Preparation of tert-butyl (R)-3-(3',5-difluoro-[1,1'-biphenyl]-3-yl)isoxazolidin-2-carboxylate**

**[0169]** Tert-butyl (R)-3-(3-bromo-5-fluorophenyl)isoxazolidin-2-carboxylate obtained in step 1 (350 mg, 1 eq.), (3-fluorophenyl)boronic acid (170 mg, 1.2 eq.), and K<sub>2</sub>CO<sub>3</sub>(280 mg, 2 eq.) were dissolved in 1,4-dioxane (5 mL) at room temperature under a nitrogen atmosphere. Next, tetrakis(triphenylphosphine)palladium(0) (110 mg, 0.1 eq.) was added to the reaction mixture at 80°C and stirred for 3 hours. As a result of TLC analysis (DCM), all of the starting materials disappeared and a new spot with different polarity was detected. The organic layer was extracted using ethyl acetate (EA; 70 mL) and a saturated aqueous sodium hydrogen carbonate solution (100 mL \* 2). The organic layer was dried over sodium sulfate, and then concentrated under reduced pressure. The concentrated compound was purified by chromatography (hexane:dichloromethane = 5:5 to 0:10) to obtain the title compound as a colorless oil (320 mg, 88% yield).

25


30

**Step 3: Preparation of (R)-3-(3',5-difluoro-[1,1'-biphenyl]-3-yl)isoxazolidine hydrochloride**

**[0170]** Tert-butyl (R)-3-(3',5-difluoro-[1,1'-biphenyl]-3-yl)isoxazolidin-2-carboxylate (320 mg, 1 eq.) obtained in step 2 was dissolved in dichloromethane (DCM; 40 mL), and then HCl/dioxane (4M, 1 mL, 5 eq.) was added at room temperature. Then, the reaction mixture was stirred at room temperature for 2 hours. As a result of LCMS analysis, all of the starting materials disappeared, diethyl ether (10 mL) was added to obtain a solid, and the resulting precipitate was filtered and dried to obtain the title compound as a white solid (240 mg, 91% yield).


35

40

**<Preparation Example 43> Preparation of (R)-3-(3-([1,2,4]triazolo[1,5-a]pyridin-7-yl)phenyl)isoxazolidine****[0171]**

**[0172]** The desired title compound was prepared in a manner similar to that of <Preparation Example 42>.


**<Preparation Example 44> Preparation of (R)-3-(3-fluoro-5-thiomorpholinophenyl)isoxazolidine****[0173]**



[0174] A compound was prepared by a method such as  $S_NAr$ , or the like, using the compound of <Preparation Example 41>.

<Example 1> Preparation of (R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine

[0175]



Step 1: Preparation of (R)-2-(6-chloropyrimidin-4-yl)-3-phenylisoxazolidine

[0176] 4,6-Dichloropyrimidine (600 mg, 1 eq.) and (R)-3-phenylisoxazolidine (631 mg, 1.05 eq.) were dissolved in dimethyl sulfoxide (DMSO, 7 ml), and then N,N-diisopropylethylamine (DIPEA; 1.41 mL, 2 eq.) was added. The reaction solution was stirred at 60°C for 30 min. After the reaction was completed, the reaction solution was extracted using ethyl acetate and water. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, concentrated under reduced pressure, and purified by MPLC (ethyl acetate/hexane) to obtain the title compound (810 mg, 77% yield) as a clear liquid.

Step 2: Preparation of (R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine

[0177] (R)-2-(6-chloropyrimidin-4-yl)-3-phenylisoxazolidine (139 mg, 1 eq.) obtained in step 1, 4-(4-methylpiperazin-1-yl)aniline (152 mg, 1.5 eq.), and potassium carbonate (220 mg, 3 eq.) were added to and dissolved in sec-butanol (1.8 ml), followed by sonication for 5 min under nitrogen to degas. Tris(dibenzylideneacetone)dipalladium(O)(Pd<sub>2</sub>(dba)<sub>3</sub>; 47 mg, 0.1 eq.) and Xphos (51 mg, 0.2 eq.) were added to the reaction mixture and stirred at 100°C for 1 hour. After the reaction was completed, the mixture was filtered through celite and washed with dichloromethane. The resulting filtrate was concentrated and purified by Prep-HPLC to obtain the title compound (71 mg, 32%).

[0178] <sup>1</sup>H NMR (400 MHz, Chloroform-d) δ 8.23 (s, 1H), 7.42 (d, J = 7.4 Hz, 2H), 7.34 (dd, J = 8.4, 6.7 Hz, 2H), 7.20-7.16 (m, 2H), 6.97-6.91 (m, 2H), 6.38 (s, 1H), 5.64 (dd, J = 8.6, 4.7 Hz, 1H), 4.10 (td, J = 7.8, 4.5 Hz, 1H), 3.85 (q, J = 7.8 Hz, 1H), 3.23 (t, J = 5.1 Hz, 4H), 2.71 (dtd, J = 12.2, 7.9, 4.4 Hz, 1H), 2.64 (t, J = 5.0 Hz, 4H), 2.38 (s, 3H), 2.37-2.32 (m, 1H), 2.04 (s, 1H).

<Examples 2 to 237>

[0179] All of the Example compounds (Example compounds Nos. 1 to 237) of the present disclosure were prepared in a manner similar to that of Example 1. The compound names, chemical structures, and NMR and LCMS analysis results of the respective Example compounds are summarized and shown in Table 1 below.

[Table 1]

5	Ex			
	am			
	pl			
10	e	Structure	Compound Name	$^1\text{H}$ NMR; MS [M+H] <sup>+</sup>
	Co			
15	mp			
	ou			

20

25

30



35




40



45

50



55

nd			
5  10  15  20  25	1  	(R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform-d) δ 8.23 (s, 1H), 7.42 (d, <i>J</i> = 7.4 Hz, 2H), 7.34 (dd, <i>J</i> = 8.4, 6.7 Hz, 2H), 7.20–7.16 (m, 2H), 6.97–6.91 (m, 2H), 6.38 (s, 1H), 5.64 (dd, <i>J</i> = 8.6, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.85 (q, <i>J</i> = 7.8 Hz, 1H), 3.23 (t, <i>J</i> = 5.1 Hz, 4H), 2.71 (dtd, <i>J</i> = 12.2, 7.9, 4.4 Hz, 1H), 2.64 (t, <i>J</i> = 5.0 Hz, 4H), 2.38 (s, 3H), 2.37–2.32 (m, 1H), 2.04 (s, 1H); 417.4[M+H] <sup>+</sup>
30  35  40  45  50	2  	(S)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform-d) δ 8.23 (s, 1H), 7.42 (d, <i>J</i> = 7.4 Hz, 2H), 7.34 (dd, <i>J</i> = 8.4, 6.7 Hz, 2H), 7.20–7.16 (m, 2H), 6.97–6.91 (m, 2H), 6.38 (s, 1H), 5.64 (dd, <i>J</i> = 8.6, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.85 (q, <i>J</i> = 7.8 Hz, 1H), 3.23 (t, <i>J</i> = 5.1 Hz, 4H), 2.71 (dtd, <i>J</i> = 12.2, 7.9, 4.4 Hz, 1H), 2.64 (t, <i>J</i> = 5.0 Hz, 4H), 2.38 (s, 3H), 2.37–2.32 (m, 1H), 2.04 (s, 1H); 417.3[M+H] <sup>+</sup>

5 3 10		(S)-6-(3-benzylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	431.4 [M+H] <sup>+</sup>
15 20 25 30 35		(R)-N-(3-methoxy-4-(4-(4-methylpiperidin-1-yl)phenyl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30-8.26 (m, 1H), 7.46-7.41 (m, 2H), 7.34 (t, <i>J</i> = 7.7 Hz, 2H), 7.26-7.22 (m, 1H), 6.96-6.90 (m, 2H), 6.85-6.80 (m, 2H), 6.47-6.43 (m, 1H), 5.66 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.11 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.89-3.82 (m, 4H), 3.54 (d, <i>J</i> = 11.2 Hz, 2H), 2.76-2.65 (m, 4H), 2.63-2.33 (m, 9H), 2.30 (s, 3H), 1.92 (d, <i>J</i> = 12.3 Hz, 2H), 1.85-1.75 (m, 2H); 530.4 [M+H] <sup>+</sup>
40 45 50 55		(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)phenyl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (s, 1H), 7.51 (d, <i>J</i> = 8.5 Hz, 1H), 7.45 (d, <i>J</i> = 7.6 Hz, 2H), 7.34 (t, <i>J</i> = 7.5 Hz, 2H), 7.26-7.23 (m, 1H), 6.70 (s, 1H), 6.57-6.51 (m, 2H), 6.41 (s, 1H), 5.68 (dd, <i>J</i> = 8.7, 4.5 Hz, 1H), 4.11 (td, <i>J</i> = 7.8, 4.3 Hz, 1H),



5		amine	3.92-3.87 (m, 1H), 3.84 (s, 3H), 3.71 (d, $J = 12.0$ Hz, 2H), 2.78-2.64 (m, 7H), 2.58-2.47 (m, 3H), 2.44-2.35 (m, 3H), 2.32 (s, 3H), 1.96 (d, $J = 12.0$ Hz, 2H), 1.74-1.67 (m, 2H); 530.4 [M+H] <sup>+</sup>
6		(R)-N-(4-(4-(4-cyclopropylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine	592.3 [M+H] <sup>+</sup>
7		(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-morpholinophenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Methanol-d <sub>4</sub> ) $\delta$ 8.47 (s, 1H), 8.39 (s, 1H), 8.22 (s, 1H), 7.47 (s, 1H), 7.06-6.99 (m, 3H), 6.94 (s, 1H), 5.56 (s, 1H), 4.44 (dt, $J = 7.6, 3.8$ Hz, 1H), 4.21 (s, 1H), 4.14 (s, 3H), 3.94 (s, 3H), 3.88-3.82 (m, 4H), 3.56-3.51 (m, 1H), 3.11-3.00 (m, 5H), 2.54-2.44 (m, 1H); 550.4 [M+H] <sup>+</sup>








5 10 15 20 25	<p>8</p> 	<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Methanol-d<sub>4</sub>) δ 8.50 (s, 1H), 8.31 (s, 1H), 8.19 (s, 1H), 7.45 (s, 1H), 7.01 (dt, <i>J</i> = 6.8, 2.1 Hz, 2H), 6.96-6.88 (m, 1H), 6.00 (s, 1H), 5.52 (d, <i>J</i> = 10.1 Hz, 1H), 4.42 (d, <i>J</i> = 4.4 Hz, 1H), 4.17 (s, 4H), 4.06 (s, 1H), 3.91 (s, 5H), 3.81-3.47 (m, 7H), 3.37 (s, 2H), 3.07 (s, 4H), 2.84 (t, <i>J</i> = 11.8 Hz, 2H), 2.48 (d, <i>J</i> = 5.9 Hz, 1H), 2.32 (d, <i>J</i> = 11.7 Hz, 2H), 2.13 (d, <i>J</i> = 9.4 Hz, 2H); 646.5 [M+H]<sup>+</sup></p>
30 35 40 45	<p>9</p> 	<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-5-yl)-4-(4-(4-morpholinophenyl)pyrimidin-4-amine</p>	<p>550.4 [M+H]<sup>+</sup></p>

50



55

5 10 15 20 25		<p>(R)-1'-[4-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-methoxy-2-(1-methyl-1H-pyrazol-4-yl)phenyl)-N,N-dimethyl-[1,4'-bipiperidin]-4-amine</p>	674.6 [M+H] <sup>+</sup>
30 35 40 45 50		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-5-(methylsulfonyl)phenyl)pyrimidin-4-amine</p>	614.3 [M+H] <sup>+</sup>

5 10 12 15 20		(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(methylsulfonyl)-5-(4-morpholinopiperidin-1-yl)phenyl)pyrimidin-4-amine	601.3 [M+H] <sup>+</sup>
25 30 13 35 40		(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(4-methylpiperazin-1-yl)-5-(methylsulfonyl)phenyl)pyrimidin-4-amine	531.3 [M+H] <sup>+</sup>
45 50 14 55		(R)-2-(3-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-	534.3 [M+H] <sup>+</sup>



5  10		yl amino)-5-(4-methylpiperazin-1-yl)methyl)phenyl)-2-methylpropanenitrile	
15  20  25  30  35	15  	(R)-2-(3-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-(4-methylpiperazin-1-yl)phenyl)-2-methylpropanenitrile	520.3 [M+H] <sup>+</sup>
40  45  50  55	16  	(R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-(3-phenoxyphenyl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform-d) δ 8.25 (d, <i>J</i> = 1.1 Hz, 1H), 7.12 (dt, <i>J</i> = 4.1, 2.0 Hz, 2H), 7.11-7.05 (m, 3H), 7.01 (dt, <i>J</i> = 5.3, 1.8 Hz, 3H), 7.00-6.98 (m, 1H), 6.96-6.92 (m, 3H), 6.90-6.85 (m, 2H), 5.63 (dt, <i>J</i> = 8.6, 5.6 Hz, 2H), 4.22 (td, <i>J</i> = 7.8, 4.2 Hz, 1H), 3.93-3.79 (m, 2H), 3.24-3.20

## EP 4 342 895 A1

5			(m, 4H), 2.61 (t, $J = 5.0$ Hz, 4H), 2.48–2.41 (m, 1H), 2.37 (s, 3H), 2.33 (dd, $J = 7.8, 4.4$ Hz, 1H); 509.4 [M+H] <sup>+</sup>
10 15 20 25 30	17 	(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.20 (s, 1H), 8.17 (d, $J = 1.0$ Hz, 1H), 7.20–7.14 (m, 2H), 6.99–6.92 (m, 4H), 6.69 (tt, $J = 8.8, 2.4$ Hz, 1H), 6.38 (d, $J = 1.0$ Hz, 1H), 5.61 (dd, $J = 8.8, 4.7$ Hz, 1H), 4.10 (td, $J = 7.9, 4.2$ Hz, 1H), 3.83 (q, $J = 8.0$ Hz, 1H), 3.77–3.69 (m, 2H), 2.82 (s, 7H), 2.77–2.69 (m, 4H), 2.54 (tt, $J = 11.5, 3.6$ Hz, 1H), 2.45 (s, 3H), 2.36–2.27 (m, 1H), 2.01–1.94 (m, 2H), 1.76–1.64 (m, 2H); 536.4 [M+H] <sup>+</sup>
35 40 45 50 55	18 	(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.18 (s, 1H), 8.11 (s, 1H), 7.19 (d, $J = 8.8$ Hz, 2H), 6.99–6.92 (m, 4H), 6.69 (tt, $J = 8.9, 2.4$ Hz, 1H), 6.39 (d, $J = 1.0$ Hz, 1H), 5.62 (dd, $J = 8.8, 4.8$ Hz, 1H), 4.10 (td, $J = 7.9, 4.2$ Hz, 1H), 3.83 (q, $J = 7.9$ Hz, 1H), 3.31–3.23 (m, 4H), 2.75 (t, $J = 5.2$ Hz, 5H), 2.43 (s, 3H), 2.33 (ddd, $J$



EP 4 342 895 A1

			= 12.4, 8.3, 4.7 Hz, 1H); 453.3[M+H] <sup>+</sup>
5		(R)-N <sup>1</sup> -(6-(3-(3,5-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.20
10		difluorophenyl)isox	(s, 1H), 7.52 (s, 1H), 7.17-7.10 (m,
		azolidin-2-	2H), 7.00-6.92 (m, 2H), 6.75-6.66 (m,
		yl)pyrimidin-4-yl)-	3H), 6.33-6.31 (m, 1H), 5.63 (dd, <i>J</i>
15	19	N <sup>4</sup> -(2-	= 8.8, 4.7 Hz, 1H), 4.08 (td, <i>J</i> = 7.9,
		(dimethylamino)ethy	4.2 Hz, 1H), 3.83 (q, <i>J</i> = 7.9 Hz, 1H),
		l)-N <sup>4</sup> -	3.57-3.52 (m, 2H), 2.98 (s, 3H),
20		methylbenzene-1,4-	2.78-2.62 (m, 3H), 2.40 (s, 6H),
		diamine	2.36-2.27 (m, 1H); 455.3[M+H] <sup>+</sup>
25			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.51
		(R)-6-(3-(3,5-	(s, 1H), 8.26 (d, <i>J</i> = 1.0 Hz, 1H),
30		difluorophenyl)isox	7.42 (s, 1H), 7.01-6.95 (m, 2H), 6.69
		azolidin-2-yl)-N-	(tt, <i>J</i> = 8.9, 2.4 Hz, 1H), 6.62 (s,
35		(2-methoxy-5-	1H), 6.40-6.37 (m, 1H), 5.64 (dd, <i>J</i>
		methyl-4-(4-(4-	= 8.8, 4.7 Hz, 1H), 4.11 (td, <i>J</i> = 7.9,
40	20	methylpiperazin-1-	4.2 Hz, 1H), 3.90-3.84 (m, 1H), 3.83
		yl)piperidin-1-	(s, 3H), 3.19 (d, <i>J</i> = 11.3 Hz, 2H),
		yl)phenyl)pyrimidin	2.86 (s, 7H), 2.78-2.62 (m, 4H),
45		-4-amine	2.61-2.52 (m, 1H), 2.45 (s, 3H),
			2.39-2.28 (m, 1H), 2.24 (s, 3H), 1.97
50			(d, <i>J</i> = 12.4 Hz, 2H), 1.74 (qd, <i>J</i> =
			11.9, 3.8 Hz, 2H); 580.4[M+H] <sup>+</sup>



<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p>	<p>21</p> 	<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.28 (s, 1H), 7.48 (d, <i>J</i> = 9.1 Hz, 1H), 7.01-6.95 (m, 2H), 6.93 (s, 1H), 6.73-6.65 (m, 1H), 6.57-6.52 (m, 2H), 6.39 (s, 1H), 5.65 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.13-4.06 (m, 1H), 3.90-3.81 (m, 4H), 3.74-3.68 (m, 2H), 2.78-2.67 (m, 8H), 2.66-2.57 (m, 3H), 2.49-2.40 (m, 1H), 2.38-2.28 (m, 4H), 2.01-1.93 (m, 2H), 1.77-1.65 (m, 2H); 566.5 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p>	<p>22</p> 	<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(dimethylamino)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.30 (s, 1H), 7.52 (d, <i>J</i> = 9.3 Hz, 1H), 7.02-6.96 (m, 2H), 6.81 (s, 1H), 6.72-6.66 (m, 1H), 6.58-6.53 (m, 2H), 6.41 (s, 1H), 5.65 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.14-4.07 (m, 1H), 3.90-3.82 (m, 4H), 3.74-3.68 (m, 2H), 2.79-2.68 (m, 3H), 2.48-2.28 (m, 8H), 2.02-1.96 (m, 2H), 1.77-1.65 (m, 2H); 511.5 [M+H]<sup>+</sup></p>



50

55



23		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-3-(trifluoromethyl)phenyl)pyrimidin-4-amine</p>	<p><math>^1\text{H NMR}</math> (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.57 (s, 1H), 8.25–8.22 (m, 1H), 7.55 (d, <math>J = 2.6</math> Hz, 1H), 7.49 (dd, <math>J = 8.6, 2.6</math> Hz, 1H), 7.33 (d, <math>J = 8.6</math> Hz, 1H), 7.00–6.92 (m, 2H), 6.70 (tt, <math>J = 8.8, 2.3</math> Hz, 1H), 6.50–6.48 (m, 1H), 5.62 (dd, <math>J = 8.7, 4.8</math> Hz, 1H), 4.19–4.13 (m, 1H), 3.88 (q, <math>J = 8.0</math> Hz, 1H), 3.48 (s, 3H), 3.16–3.08 (m, 2H), 2.88 (s, 5H), 2.81–2.71 (m, 3H), 2.66–2.55 (m, 1H), 2.48 (s, 3H), 2.40–2.30 (m, 1H), 1.96–1.88 (m, 2H), 1.75 (tt, <math>J = 11.9, 6.0</math> Hz, 2H); 604.4[M+H]<sup>+</sup></p>
24		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-(methylsulfonyl)phenyl)piperidin-1-yl)-3-(methylsulfonyl)phenyl)pyrimidin-4-amine</p>	<p><math>^1\text{H NMR}</math> (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.30–8.26 (m, 1H), 8.02 (s, 1H), 7.98 (d, <math>J = 2.7</math> Hz, 1H), 7.70 (dd, <math>J = 8.6, 2.7</math> Hz, 1H), 7.37 (d, <math>J = 8.6</math> Hz, 1H), 7.00–6.94 (m, 2H), 6.70 (tt, <math>J = 8.8, 2.4</math> Hz, 1H), 6.53–6.51 (m, 1H), 5.63 (dd, <math>J = 8.7, 4.8</math> Hz, 1H), 4.17 (td, <math>J = 7.9, 4.2</math> Hz, 1H), 3.93 (q, <math>J = 8.0</math> Hz, 1H), 3.35–3.28 (m, 4H), 2.82–2.70 (m, 9H), 2.53–2.44 (m, 1H), 2.41 (s, 3H), 2.40–2.32 (m, 1H), 2.05 (s,</p>






5			3H), 2.00 (d, $J = 12.7$ Hz, 2H), 1.75 (q, $J = 11.9$ Hz, 2H); 614.4 [M+H] <sup>+</sup>
25		(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (s, 1H), 7.49 (s, 1H), 7.12 (s, 1H), 7.02-6.95 (m, 2H), 6.72-6.67 (m, 2H), 6.49 (s, 1H), 5.65 (dd, $J = 8.8, 4.7$ Hz, 1H), 4.11 (td, $J = 7.9, 4.1$ Hz, 1H), 3.90-3.79 (m, 7H), 3.13 (d, $J = 11.4$ Hz, 2H), 2.79-2.62 (m, 12H), 2.51-2.44 (m, 1H), 2.39 (s, 3H), 2.36-2.30 (m, 1H), 2.17 (s, 1H), 1.96 (d, $J = 12.2$ Hz, 2H), 1.73 (qd, $J = 11.5, 3.2$ Hz, 2H); 594.4 [M+H] <sup>+</sup>
26		(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.25-8.20 (m, 1H), 7.65-7.60 (m, 1H), 7.00-6.89 (m, 3H), 6.86-6.78 (m, 2H), 6.69 (tt, $J = 8.7, 2.4$ Hz, 1H), 6.46 (d, $J = 0.9$ Hz, 1H), 5.62 (dd, $J = 8.8, 4.8$ Hz, 1H), 4.10 (td, $J = 7.9, 4.2$ Hz, 1H), 3.87 (s, 3H), 3.86-3.82 (m, 1H), 3.55 (d, $J = 11.4$ Hz, 2H), 2.82-2.65 (m, 8H), 2.64-2.47 (m, 4H), 2.38 (s, 3H), 2.37-2.31 (m, 1H), 1.94 (d, $J = 12.2$ Hz, 2H), 1.82 (tt, $J =$

			12.0, 6.0 Hz, 2H); 566.4[M+H] <sup>+</sup>
5			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.28
10		(R)-N-(2-methoxy-4-	( <i>s</i> , 1H), 7.71 ( <i>s</i> , 1H), 7.64 ( <i>d</i> , <i>J</i> =
15	27	(4-(4-	7.5 Hz, 1H), 7.48 ( <i>dt</i> , <i>J</i> = 15.8, 7.7
20		methylpiperazin-1-	Hz, 3H), 6.95 ( <i>s</i> , 1H), 6.59-6.49 ( <i>m</i> ,
25		yl)piperidin-1-	2H), 6.41 ( <i>s</i> , 1H), 5.74 ( <i>dd</i> , <i>J</i> = 8.8,
		yl)phenyl)-6-(3-(3-	4.6 Hz, 1H), 4.11 ( <i>q</i> , <i>J</i> = 8.0 Hz, 1H),
		(trifluoromethyl)ph	3.93-3.80 ( <i>m</i> , 4H), 3.71 ( <i>d</i> , <i>J</i> = 12.5
		enyl)isoxazolidin-	Hz, 2H), 2.81-2.67 ( <i>m</i> , 7H), 2.62 ( <i>s</i> ,
		2-yl)pyrimidin-4-	4H), 2.48-2.31 ( <i>m</i> , 5H), 1.97 ( <i>d</i> , <i>J</i> =
		amine	12.4 Hz, 2H), 1.74-1.66 ( <i>m</i> , 2H);
			598.44[M+H] <sup>+</sup>
30			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.28
35		(R)-6-(3-(2-fluoro-	( <i>s</i> , 1H), 7.79 ( <i>t</i> , <i>J</i> = 7.4 Hz, 1H),
40	28	3-	7.51 ( <i>t</i> , <i>J</i> = 9.5 Hz, 2H), 7.21 ( <i>t</i> , <i>J</i>
45		(trifluoromethyl)ph	= 7.9 Hz, 1H), 6.99 ( <i>s</i> , 1H), 6.55 ( <i>d</i> ,
50		enyl)isoxazolidin-	<i>J</i> = 7.5 Hz, 2H), 6.42 ( <i>s</i> , 1H), 5.93
55		2-yl)-N-(2-methoxy-	( <i>dd</i> , <i>J</i> = 8.8, 4.6 Hz, 1H), 4.12-4.05
		4-(4-(4-	( <i>m</i> , 1H), 3.92-3.81 ( <i>m</i> , 4H), 3.72 ( <i>d</i> ,
		methylpiperazin-1-	<i>J</i> = 12.2 Hz, 2H), 2.84 ( <i>d</i> , <i>J</i> = 8.7
		yl)piperidin-1-	Hz, 1H), 2.80-2.51 ( <i>m</i> , 10H), 2.50-
		yl)phenyl)pyrimidin	2.41 ( <i>m</i> , 1H), 2.36 ( <i>s</i> , 3H), 2.32-2.25
		-4-amine	( <i>m</i> , 1H), 1.97 ( <i>d</i> , <i>J</i> = 12.6 Hz, 2H),
			1.71 ( <i>q</i> , <i>J</i> = 11.4, 10.9 Hz, 2H);

			616.35[M+H] <sup>+</sup>
5			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30
10			( <i>s</i> , 1H), 7.50 ( <i>d</i> , <i>J</i> = 8.9 Hz, 1H),
15		(R)-6-(3-(3-	7.34-7.28 ( <i>m</i> , 1H), 7.19 ( <i>dd</i> , <i>J</i> =
20	29	fluorophenyl)isoxaz	13.5, 8.8 Hz, 2H), 6.94 ( <i>td</i> , <i>J</i> = 8.4,
25		olidin-2-yl)-N-(2-	1.9 Hz, 1H), 6.80 ( <i>s</i> , 1H), 6.57-6.51
30		methoxy-4-(4-(4-	( <i>m</i> , 2H), 6.40 ( <i>d</i> , <i>J</i> = 1.1 Hz, 1H),
35		methylpiperazin-1-	5.67 ( <i>dd</i> , <i>J</i> = 8.7, 4.6 Hz, 1H), 4.10
40		yl)piperidin-1-	( <i>td</i> , <i>J</i> = 7.8, 4.2 Hz, 1H), 3.91-3.82
45		yl)phenyl)pyrimidin	( <i>m</i> , 4H), 3.71 ( <i>d</i> , <i>J</i> = 12.2 Hz, 2H),
50		-4-amine	2.78-2.53 ( <i>m</i> , 11H), 2.41-2.38 ( <i>m</i> ,
55			1H), 2.37-2.34 ( <i>m</i> , 1H), 2.32 ( <i>d</i> , <i>J</i> =
			1.3 Hz, 3H), 1.96 ( <i>d</i> , <i>J</i> = 12.7 Hz,
			2H), 1.70 ( <i>dt</i> , <i>J</i> = 11.5, 7.9 Hz, 2H);
			548.39[M+H] <sup>+</sup>
35		(R)-6-(3-(2,5-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30
40		difluorophenyl)isox	( <i>s</i> , 1H), 7.51 ( <i>d</i> , <i>J</i> = 8.9 Hz, 1H),
45		azolidin-2-yl)-N-	7.30 ( <i>d</i> , <i>J</i> = 3.3 Hz, 1H), 7.00 ( <i>td</i> ,
50		(2-methoxy-4-(4-(4-	<i>J</i> = 9.0, 4.0 Hz, 1H), 6.94-6.86 ( <i>m</i> ,
55		methylpiperazin-1-	1H), 6.83 ( <i>s</i> , 1H), 6.58-6.52 ( <i>m</i> , 2H),
		yl)piperidin-1-	6.43 ( <i>s</i> , 1H), 5.87 ( <i>dd</i> , <i>J</i> = 8.7, 4.4
		yl)phenyl)pyrimidin	Hz, 1H), 4.10-4.05 ( <i>m</i> , 1H), 3.85 ( <i>s</i> ,
		-4-amine	4H), 3.71 ( <i>d</i> , <i>J</i> = 12.3 Hz, 2H), 2.80-
			2.66 ( <i>m</i> , 7H), 2.56 ( <i>s</i> , 4H), 2.42 ( <i>t</i> ,

5			$J = 11.6 \text{ Hz, 1H), 2.33 (s, 3H), 2.31-2.23 (m, 1H), 2.05-2.01 (m, 2H), 1.70 (dd, } J = 20.7, 11.3 \text{ Hz, 2H); 566.36 [M+H]^+$
10 15 20 25 30 35	<p>31</p> 	<p>(R)-6-(3-(3-chloro-4-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	$^1\text{H NMR (400 MHz, Chloroform-}d) \delta 8.28 (s, 1H), 7.50 (t, J = 7.4 \text{ Hz, 2H), 7.33-7.28 (m, 1H), 7.10 (t, } J = 8.6 \text{ Hz, 1H), 6.87 (s, 1H), 6.54 (d, } J = 7.8 \text{ Hz, 2H), 6.39 (s, 1H), 5.63 (dd, } J = 8.3, 4.5 \text{ Hz, 1H), 4.15-4.06 (m, 1H), 3.86 (d, } J = 11.0 \text{ Hz, 4H), 3.71 (d, } J = 12.0 \text{ Hz, 2H), 2.78-2.68 (m, 5H), 2.58 (s, 6H), 2.46-2.40 (m, 1H), 2.37-2.26 (m, 4H), 1.97 (d, } J = 12.4 \text{ Hz, 2H), 1.70 (dd, } J = 22.3, 10.3 \text{ Hz, 2H); 582.36[M+H]^+$
40 45 50 55	<p>32</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	$^1\text{H NMR (400 MHz, Chloroform-}d) \delta 8.27 (s, 1H), 7.47 (d, J = 8.5 \text{ Hz, 1H), 7.31 (t, } J = 7.0 \text{ Hz, 1H), 7.12-7.01 (m, 3H), 6.54 (d, } J = 7.3 \text{ Hz, 2H), 6.39 (d, } J = 1.0 \text{ Hz, 1H), 5.90 (dd, } J = 8.8, 4.7 \text{ Hz, 1H), 4.09 (dt, } J = 7.8, 3.8 \text{ Hz, 1H), 3.90-3.85 (m, 1H), 3.84 (s, 3H), 3.71 (d, } J = 12.3 \text{ Hz, 2H); 582.36[M+H]^+$



			2H), 2.86-2.60 (m, 11H), 2.51-2.44 (m, 1H), 2.38 (s, 3H), 2.34-2.28 (m, 1H), 1.97 (d, $J = 12.6$ Hz, 2H), 1.76-1.65 (m, 2H); 566.31[M+H] <sup>+</sup>
33		(R)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.27 (d, $J = 1.0$ Hz, 1H), 7.46 (d, $J = 8.3$ Hz, 1H), 7.43-7.37 (m, 2H), 7.06 (s, 1H), 7.02 (t, $J = 8.7$ Hz, 2H), 6.53 (d, $J = 8.1$ Hz, 2H), 6.36 (d, $J = 1.0$ Hz, 1H), 5.64 (dd, $J = 8.5, 4.6$ Hz, 1H), 4.11 (dt, $J = 7.9, 4.0$ Hz, 1H), 3.91-3.83 (m, 4H), 3.71 (d, $J = 12.3$ Hz, 2H), 2.88-2.55 (m, 11H), 2.50-2.43 (m, 1H), 2.42-2.29 (m, 4H), 1.97 (d, $J = 12.4$ Hz, 2H), 1.76-1.65 (m, 2H); 548.31[M+H] <sup>+</sup>
34		(R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) $\delta$ 8.44 (s, 1H), 8.11 (d, $J = 1.0$ Hz, 1H), 7.49 (q, $J = 8.6, 8.1$ Hz, 1H), 7.34-7.22 (m, 2H), 7.07 (t, $J = 8.8$ Hz, 1H), 6.62 (d, $J = 2.7$ Hz, 1H), 6.49 (dd, $J = 8.8, 2.6$ Hz, 1H), 6.17 (s, 1H), 5.68 (dd, $J = 8.7, 4.7$ Hz, 1H), 4.16-4.10 (m, 1H), 3.83-3.75 (m, 4H), 3.72

5  10			(d, $J = 12.1$ Hz, 2H), 3.43–3.36 (m, 5H), 2.80–2.72 (m, 1H), 2.67 (t, $J = 12.1$ Hz, 2H), 2.38–2.25 (m, 4H), 2.21–2.09 (m, 4H), 1.83 (d, 2H), 1.51 (q, $J = 11.4$ Hz, 2H); 566.27[M+H] <sup>+</sup>
15  20  25  30  35  40	35  	(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(thiophen-2-yl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.32 (d, $J = 1.0$ Hz, 1H), 7.50 (d, $J = 8.5$ Hz, 1H), 7.20 (dd, $J = 5.0, 1.2$ Hz, 1H), 7.08–7.04 (m, 1H), 7.02–6.94 (m, 2H), 6.55–6.49 (m, 2H), 6.38 (d, $J = 1.1$ Hz, 1H), 5.98 (dd, $J = 8.3, 3.5$ Hz, 1H), 4.17 (td, $J = 7.9, 5.2$ Hz, 1H), 3.93 (q, $J = 7.8$ Hz, 1H), 3.84 (s, 3H), 3.75–3.67 (m, 2H), 2.93 (s, 6H), 2.75 (td, $J = 12.2, 2.3$ Hz, 3H), 2.69–2.62 (m, 2H), 2.60 (s, 3H), 2.55–2.44 (m, 2H), 1.98 (d, $J = 12.4$ Hz, 2H), 1.79–1.66 (m, 2H); 536.4[M+H] <sup>+</sup>

45




50

55



<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p>	<p>36</p> 	<p>(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(naphthalen-1-yl)isoxazolidin-2-yl)pyrimidin-4-amine</p>	<p>1H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.28 (s, 1H), 8.12 (d, <i>J</i> = 8.3 Hz, 1H), 7.89 (d, <i>J</i> = 8.0 Hz, 1H), 7.78 (d, <i>J</i> = 8.2 Hz, 1H), 7.73 (d, <i>J</i> = 7.2 Hz, 1H), 7.58-7.43 (m, 4H), 6.54 (d, <i>J</i> = 8.0 Hz, 2H), 6.45 (s, 1H), 6.39 (dd, <i>J</i> = 8.7, 4.0 Hz, 1H), 4.16-4.09 (m, 1H), 4.00 (q, <i>J</i> = 7.8 Hz, 1H), 3.86 (s, 3H), 3.71 (d, <i>J</i> = 12.3 Hz, 2H), 2.97-2.87 (m, 4H), 2.76 (t, <i>J</i> = 12.0 Hz, 3H), 2.63 (s, 4H), 2.45-2.35 (m, 2H), 2.08 (s, 3H), 1.97 (d, 2H), 1.75 (ddd, 2H); 580.39 [M+H]<sup>+</sup></p>
<p>35</p> <p>40</p> <p>45</p>	<p>37</p> 	<p>(R)-6-(3-(3-ethynylphenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>554.3 [M+H]<sup>+</sup></p>

50

55




5 10 15 20 25	<p>38</p> 	<p>(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(6-methylpyridin-3-yl)isoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.75–8.49 (m, 2H), 8.22–8.12 (m, 1H), 8.08–7.94 (m, 1H), 7.57–7.27 (m, 2H), 6.68 (s, 1H), 6.61–6.50 (m, 1H), 6.25–6.12 (m, 1H), 5.65–5.53 (m, 1H), 4.20–4.13 (m, 1H), 3.99–3.93 (m, 1H), 3.91–3.67 (m, 6H), 3.50–3.27 (m, 4H), 3.19–3.14 (m, 2H), 2.83–2.66 (m, 6H), 2.60–2.55 (m, 2H), 2.38–2.19 (m, 2H), 2.04–1.92 (m, 2H), 1.92–1.79 (m, 2H), 1.68–1.50 (m, 2H); 545.4 [M+H]<sup>+</sup></p>
30 35 40	<p>39</p> 	<p>(R)-6-(3-(3-((1,2,4)triazolo[1,5-a]pyridin-7-yl)phenyl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>534.3 [M+H]<sup>+</sup></p>
45 50 55	<p>40</p> 	<p>(R)-6-(3-(3-fluoro-5-thiomorpholinophenyl)isoxazolidin-2-yl)-N-(4-(4-</p>	<p>536.3 [M+H]<sup>+</sup></p>







5		methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	
10 15 20 25 30		(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(3-methoxyphenyl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 10.83 (s, 1H), 8.04 (s, 1H), 7.31-7.26 (m, 1H), 7.15 (d, <i>J</i> = 8.5 Hz, 1H), 6.89-6.79 (m, 3H), 6.62-6.52 (m, 2H), 5.75 (s, 1H), 5.54 (dd, <i>J</i> = 8.6, 4.5 Hz, 1H), 4.24 (td, <i>J</i> = 7.5, 5.5 Hz, 1H), 4.03 (q, <i>J</i> = 7.5 Hz, 1H), 3.89-3.77 (m, 8H), 3.66 (s, 8H), 3.34-3.22 (m, 1H), 2.94-2.81 (m, 6H), 2.49-2.37 (m, 1H), 2.26-2.15 (m, 2H), 2.05-1.93 (m, 2H); 560.4 [M+H] <sup>+</sup>
35 40 45		(R)-6-(3-methyl-3-phenylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	431.3 [M+H] <sup>+</sup>




50



55



5 10 15		<p>isopropyl (R)-3-(2- (6-((4-(4- methylpiperazin-1- yl)phenyl)amino)pyr imidin-4- yl)isoxazolidin-3- yl)benzoate</p>	503.3 [M+H] <sup>+</sup>
20 25 30		<p>(R)-N-cyclohexyl-3- (2-(6-((4-(4- methylpiperazin-1- yl)phenyl)amino)pyr imidin-4- yl)isoxazolidin-3- yl)benzamide</p>	542.3 [M+H] <sup>+</sup>
35 40 45 50		<p>(R)-6-(3-(3',5- difluoro-[1,1'- biphenyl]-3- yl)isoxazolidin-2- yl)-N-(4-(4- methylpiperazin-1- yl)phenyl)pyrimidin -4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.27 (d, <i>J</i> = 1.0 Hz, 1H), 7.45--7.10 (m, 9H), 7.09--6.90 (m, 3H), 6.40 (d, <i>J</i> = 1.0 Hz, 1H), 5.71 (dd, <i>J</i> = 8.7, 4.7 Hz, 1H), 4.11 (td, <i>J</i> = 7.9, 4.1 Hz, 1H), 3.85 (q, <i>J</i> = 7.9 Hz, 1H), 3.26- 3.19 (m, 4H), 2.76 (dtt, <i>J</i> = 12.2, 7.6, 4.2 Hz, 1H), 2.63-2.54 (m, 4H), 2.44-2.31 (m, 4H); 529.4 [M+H]<sup>+</sup></p>



5 10 15		(R)-N,N-dimethyl-7-(2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-7-azaspiro[3.5]nonan-2-amine	499.3 [M+H] <sup>+</sup>
20 25 30		(R)-N,N-dimethyl-2-(2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-2-azaspiro[3.5]nonan-7-amine	499.3 [M+H] <sup>+</sup>
35 40 45 50 55		(R)-3-(1-methylpiperidin-4-yl)-N-(6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-1,2,3,4,4a,5-hexahydrobenzo[b]pyrazino[1,2-d][1,4]oxazin-8-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.17 (d, <i>J</i> = 1.0 Hz, 1H), 8.07 (s, 1H), 7.44-7.40 (m, 2H), 7.35 (t, <i>J</i> = 7.6 Hz, 2H), 7.26-7.22 (m, 1H), 6.79-6.69 (m, 3H), 6.42 (d, <i>J</i> = 1.0 Hz, 1H), 5.64 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.20 (dd, <i>J</i> = 10.6, 2.7 Hz, 1H), 4.12 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 4.01 (dd, <i>J</i> = 10.6, 8.9 Hz, 1H), 3.88 (q, <i>J</i> = 7.8 Hz, 1H), 3.68 (dt, <i>J</i> = 11.6, 2.8 Hz,

5 10 15 20			1H), 3.30 (d, $J = 11.5$ Hz, 2H), 3.16 (ddt, $J = 11.5, 9.4, 2.9$ Hz, 1H), 3.02 (d, $J = 11.2$ Hz, 1H), 2.88 (dt, $J = 10.5, 2.6$ Hz, 1H), 2.80 (td, $J = 11.8, 3.4$ Hz, 1H), 2.73 (ddt, $J = 8.8, 7.6, 4.5$ Hz, 1H), 2.54 (s, 3H), 2.47 (ddt, $J = 11.6, 8.9, 3.8$ Hz, 3H), 2.38 (dtd, $J = 12.4, 7.9, 4.7$ Hz, 1H), 2.07 (s, 1H), 2.02 (s, 1H), 1.98–1.86 (m, 4H); 528.37 [M+H] <sup>+</sup>
25 30 35 40 45 50	49 	(6aR,8S)-8-(4-cyclopropylpiperazin-1-yl)-2-methoxy-N-(6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-6,6a,7,8,9,10-hexahydrobenzo[b]pyrido[1,2-d][1,4]oxazin-3-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (d, $J = 1.1$ Hz, 1H), 7.45 (d, $J = 7.4$ Hz, 2H), 7.35 (t, $J = 7.6$ Hz, 2H), 7.24 (t, $J = 7.1$ Hz, 1H), 7.20 (s, 1H), 6.80 (s, 1H), 6.45 (s, 1H), 6.41 (d, $J = 1.1$ Hz, 1H), 5.68 (dd, $J = 8.6, 4.5$ Hz, 1H), 4.20–4.07 (m, 2H), 3.98 (dd, $J = 10.8, 8.2$ Hz, 1H), 3.90 (q, $J = 8.0$ Hz, 2H), 3.79 (s, 3H), 3.02 (dd, $J = 11.4, 8.4$ Hz, 1H), 2.79–2.47 (m, 10H), 2.43–2.34 (m, 1H), 2.07–1.99 (m, 2H), 1.88–1.83 (m, 1H), 1.69–1.61 (m, 7H); 584.45 [M+H] <sup>+</sup>

5 10 15 20		<p>N-(4-(4-(5-ethylhexahydroindolo[3,4-c]pyrrolo[2,1-h]indolizin-2(1H)-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	570.3 [M+H] <sup>+</sup>
25 30 35 40		<p>(R)-N-(2-methoxy-4-(4-((1-methylpiperidin-4-yl)amino)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	544.3 [M+H] <sup>+</sup>
45 50 55		<p>(R)-N-(2-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.40 (s, 1H), 8.14 (s, 1H), 7.40–7.31 (m, 4H), 7.24 (s, 1H), 7.13 (d, <i>J</i> = 8.6 Hz, 1H), 6.85–6.73 (m, 2H), 5.97 (s, 1H), 5.62 (dd, <i>J</i> = 8.7, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.85

5  10  15		2-yl)pyrimidin-4-amine	(q, $J = 7.8$ Hz, 1H), 3.76 (d, $J = 12.2$ Hz, 2H), 3.02 (d, $J = 30.2$ Hz, 8H), 2.81-2.66 (m, 4H), 2.63 (s, 3H), 2.42-2.31 (m, 1H), 2.22 (s, 3H), 1.98 (d, $J = 12.4$ Hz, 2H), 1.72 (qd, $J = 12.1, 4.0$ Hz, 2H); 514.34[M+H] <sup>+</sup>
20  25 53 30  35  40		(R)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29-8.26 (m, 1H), 7.46-7.42 (m, 2H), 7.37-7.31 (m, 2H), 7.26-7.23 (m, 1H), 7.12-7.06 (m, 2H), 7.02-6.98 (m, 1H), 6.63 (s, 1H), 6.45-6.43 (m, 1H), 5.67 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.11 (td, $J = 7.9, 4.5$ Hz, 1H), 3.91-3.83 (m, 1H), 3.21-3.13 (m, 2H), 2.83-2.61 (m, 8H), 2.60-2.45 (m, 3H), 2.43-2.36 (m, 2H), 2.34 (s, 3H), 2.29 (s, 3H), 1.97 (d, $J = 12.2$ Hz, 2H), 1.75-1.69 (m, 2H); 514.3[M+H] <sup>+</sup>
45 54 50  55		(R)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.20 (s, 1H), 7.61 (s, 1H), 7.42 (d, $J = 7.5$ Hz, 2H), 7.34 (t, $J = 7.5$ Hz, 2H), 7.26-7.22 (m, 1H), 7.19-7.12 (m, 2H), 6.94 (d, $J = 8.8$ Hz, 2H), 6.37 (s, 1H), 5.65 (dd, $J = 8.7, 4.6$ Hz, 1H),



5 10 15		amine	4.10 (td, $J = 7.8, 4.4$ Hz, 1H), 3.86 (q, $J = 7.8$ Hz, 1H), 3.73 (d, $J = 12.2$ Hz, 2H), 2.79–2.63 (m, 11H), 2.48–2.39 (m, 2H), 2.37 (s, 3H), 1.97 (d, $J = 12.5$ Hz, 2H), 1.69 (qd, $J = 12.1, 4.0$ Hz, 2H); 500.3[M+H] <sup>+</sup>
55 20 25 30		(R)-N-(2-ethoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	544.3 [M+H] <sup>+</sup>
56 35 40 45 50 55		N-(4-(4-(4-(2-(trifluoromethyl)oxy)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.31 (d, $J = 1.0$ Hz, 1H), 7.57 (d, $J = 8.8$ Hz, 1H), 7.47–7.42 (m, 2H), 7.35 (dd, $J = 8.4, 6.9$ Hz, 2H), 7.26–7.23 (m, 1H), 6.70 (s, 1H), 6.67 (dd, $J = 8.9, 2.6$ Hz, 1H), 6.58 (d, $J = 2.6$ Hz, 1H), 6.37 (d, $J = 1.0$ Hz, 1H), 5.67 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.56 (p, $J = 6.3$ Hz, 1H), 4.13 (td, $J = 7.8, 4.4$ Hz, 1H), 3.68 (d, $J = 12.1$ Hz, 2H),



5  10			2.84–2.46 (m, 11H), 2.40 (ddd, $J = 12.5, 8.3, 4.5$ Hz, 2H), 2.32 (s, 3H), 1.96 (s, 2H), 1.68 (qd, $J = 12.2, 4.0$ Hz, 2H), 1.49–1.46 (m, 3H), 1.33–1.27 (m, 1H); 612.38 [M+H] <sup>+</sup>
15  20  25 57		N-(2-methoxy-4-(4-((1R,4R)-5-methyl-2,5-diazabicyclo[2.2.1]heptan-2-yl)piperidin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 10.95 (s, 1H), 8.01 (s, 1H), 7.35 (d, $J = 7.2$ Hz, 2H), 7.29 (d, $J = 7.4$ Hz, 3H), 7.15 (d, $J = 8.5$ Hz, 1H), 6.62 (s, 1H), 6.57 (d, $J = 8.7$ Hz, 1H), 5.75 (s, 1H), 5.57 (dd, $J = 8.7, 4.6$ Hz, 1H), 4.32 (s, 1H), 4.25 (q, $J = 7.0$ Hz, 1H), 4.09 (s, 1H), 4.03 (q, $J = 7.6$ Hz, 1H), 3.82 (s, 3H), 3.77–3.69 (m, 2H), 3.59 (s, 2H), 3.16 (s, 2H), 2.95–2.81 (m, 7H), 2.45–2.39 (m, 2H), 2.31 (d, $J = 11.9$ Hz, 1H), 2.05 (s, 1H), 1.94 (s, 3H); 542.40 [M+H] <sup>+</sup>
45  50 58		(R)-N-(2-methoxy-4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-	489.3 [M+H] <sup>+</sup>



55








		amine	
5			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.28
10			(d, <i>J</i> = 1.0 Hz, 1H), 7.47-7.37 (m,
15			2H), 7.34 (t, <i>J</i> = 7.6 Hz, 2H), 7.25-
20	59	N-(4-((1 <i>R</i> ,4 <i>R</i> )-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-2-methoxyphenyl)-6-(( <i>R</i> )-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	7.21 (m, 2H), 6.58 (s, 1H), 6.35 (d, <i>J</i> = 1.0 Hz, 1H), 6.20 (dd, <i>J</i> = 8.6, 2.5 Hz, 1H), 6.16 (d, <i>J</i> = 2.5 Hz, 1H), 5.68 (dd, <i>J</i> = 8.7, 4.5 Hz, 1H), 4.67 (s, 1H), 4.40 (s, 1H), 4.14-4.07 (m, 1H), 3.97 (d, <i>J</i> = 7.4 Hz, 1H), 3.92-3.85 (m, 2H), 3.83 (s, 3H), 3.58 (dd, <i>J</i> = 9.0, 1.6 Hz, 1H), 3.20 (d, <i>J</i> = 9.1 Hz, 1H), 2.76-2.66 (m, 1H), 2.43-2.33 (m, 1H), 2.07-1.96 (m, 2H); 446.3[M+H] <sup>+</sup>
35		( <i>R</i> )-1-(4-(1-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperidin-4-yl)piperazin-1-yl)ethan-1-one	
40	60		558.3 [M+H] <sup>+</sup>
45			
50			



61		<p>(R)-N-(4-(4-(4-cyclopropyl-3,3-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.23 (s, 1H), 7.71 (s, 1H), 7.47-7.39 (m, 3H), 7.38-7.31 (m, 2H), 7.29-7.25 (m, 1H), 6.55-6.48 (m, 2H), 6.27 (s, 1H), 5.65 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.13 (td, $J = 7.7, 4.6$ Hz, 1H), 3.91 (q, $J = 7.8$ Hz, 1H), 3.83 (s, 3H), 3.70 (d, $J = 12.4$ Hz, 2H), 3.20 (s, 3H), 3.08-2.86 (m, 3H), 2.82-2.63 (m, 6H), 2.46-2.32 (m, 2H), 2.14 (s, 1H), 2.06 (s, 1H), 2.03-1.94 (m, 2H), 1.73 (d, $J = 12.3$ Hz, 2H), 1.44 (s, 6H); 584.5 [M+H] <sup>+</sup>
62		<p>(R)-N-(2-methoxy-4-(4-(pyrrolidin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 12.12 (s, 1H), 10.75 (s, 1H), 8.02 (s, 1H), 7.39-7.27 (m, 4H), 7.14 (d, $J = 8.5$ Hz, 1H), 6.67-6.54 (m, 2H), 5.75 (s, 1H), 5.57 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.71 (s, 1H), 4.29-4.21 (m, 1H), 4.04 (q, $J = 7.5$ Hz, 1H), 3.93-3.35 (m, 7H), 3.28-3.13 (m, 1H), 3.00-2.80 (m, 4H), 2.48-2.39 (m, 1H), 2.25-1.89 (m, 8H); 501.3 [M+H] <sup>+</sup>



<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p>	<p>63</p> 	<p>(R)-N-(4-(4-(4-</p> <p>cyclopropylpiperazi</p> <p>n-1-yl)-[1,4'-</p> <p>bipiperidin]-1'-</p> <p>yl)-2-</p> <p>methoxyphenyl)-6-</p> <p>(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.22</p> <p>(s, 1H), 7.71 (s, 1H), 7.43 (d, <i>J</i> =</p> <p>7.6 Hz, 2H), 7.35 (t, <i>J</i> = 7.5 Hz, 2H),</p> <p>7.24 (s, 1H), 6.91 (d, <i>J</i> = 8.2 Hz,</p> <p>1H), 6.82 (d, <i>J</i> = 7.6 Hz, 2H), 6.46</p> <p>(s, 1H), 5.64 (dd, <i>J</i> = 8.7, 4.7 Hz,</p> <p>1H), 4.12 (td, <i>J</i> = 7.8, 4.3 Hz, 1H),</p> <p>3.86 (s, 4H), 3.55 (d, <i>J</i> = 11.2 Hz,</p> <p>2H), 3.17 (d, <i>J</i> = 11.0 Hz, 2H), 2.75-</p> <p>2.55 (m, 12H), 2.49-2.35 (m, 5H),</p> <p>1.98-1.63 (m, 12H); 639.56 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p>	<p>64</p> 	<p>(R)-N-(4-(2-</p> <p>(dimethylamino)etho</p> <p>xy)-2-</p> <p>methoxyphenyl)-6-</p> <p>(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.30</p> <p>(d, <i>J</i> = 1.0 Hz, 1H), 7.55 (d, <i>J</i> = 8.6</p> <p>Hz, 1H), 7.47-7.40 (m, 2H), 7.35 (t,</p> <p><i>J</i> = 8.4, 6.9 Hz, 2H), 7.26-7.23 (m,</p> <p>1H), 6.90 (s, 1H), 6.58 (d, <i>J</i> = 2.6</p> <p>Hz, 1H), 6.51 (dd, <i>J</i> = 8.7, 2.6 Hz,</p> <p>1H), 6.38 (d, <i>J</i> = 1.0 Hz, 1H), 5.67</p> <p>(dd, <i>J</i> = 8.6, 4.5 Hz, 1H), 4.17-4.07</p> <p>(m, 3H), 3.89 (q, <i>J</i> = 7.9 Hz, 1H),</p> <p>3.83 (s, 3H), 2.81 (t, <i>J</i> = 5.6 Hz,</p> <p>2H), 2.72 (tdd, <i>J</i> = 8.7, 7.9, 4.4 Hz,</p> <p>1H), 2.43-2.32 (m, 7H); 436.24 [M+H]<sup>+</sup></p>

5 10 15 20 25	<p>65</p> 	<p>(R)-N-(4-((2-(dimethylamino)ethyl)thio)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.36 (d, <i>J</i> = 1.0 Hz, 1H), 7.89 (d, <i>J</i> = 8.3 Hz, 1H), 7.46 (d, <i>J</i> = 7.3 Hz, 2H), 7.36 (t, <i>J</i> = 7.6 Hz, 2H), 7.29–7.25 (m, 1H), 7.10 (s, 1H), 7.02 (dd, <i>J</i> = 8.3, 2.0 Hz, 1H), 6.96 (d, <i>J</i> = 2.0 Hz, 1H), 6.52 (d, <i>J</i> = 1.0 Hz, 1H), 5.68 (dd, <i>J</i> = 8.6, 4.7 Hz, 1H), 4.16 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.96–3.87 (m, 4H), 3.06–2.98 (m, 2H), 2.81–2.70 (m, 1H), 2.62–2.55 (m, 2H), 2.46–2.36 (m, 1H), 2.29 (s, 6H); 452.21 [M+H]<sup>+</sup></p>
30 35 40 45 50	<p>66</p> 	<p>(R)-N-(2-methoxy-4-thiomorpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.30 (d, <i>J</i> = 1.0 Hz, 1H), 7.56 (d, <i>J</i> = 8.3 Hz, 1H), 7.45 (d, <i>J</i> = 7.5 Hz, 2H), 7.35 (t, <i>J</i> = 7.6 Hz, 2H), 7.26–7.23 (m, 1H), 6.89 (s, 1H), 6.56–6.48 (m, 2H), 6.42 (d, <i>J</i> = 1.0 Hz, 1H), 5.68 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.12 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.89 (q, <i>J</i> = 7.8 Hz, 1H), 3.85 (s, 3H), 3.53–3.49 (m, 4H), 2.81–2.67 (m, 5H), 2.44–2.34 (m, 1H); 450.22 [M+H]<sup>+</sup></p>



5 10 15 20 25 30	<p>67</p> 	<p>(R)-N-(2-methoxy-4-(4-methyl-1,4-diazepan-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.26 (d, <math>J</math> = 1.0 Hz, 1H), 7.47-7.42 (m, 2H), 7.34 (td, <math>J</math> = 6.9, 6.3, 3.0 Hz, 3H), 7.24 (t, <math>J</math> = 7.2 Hz, 1H), 6.88 (s, 1H), 6.32 (d, <math>J</math> = 1.0 Hz, 1H), 6.31-6.23 (m, 2H), 5.68 (dd, <math>J</math> = 8.6, 4.5 Hz, 1H), 4.10 (td, <math>J</math> = 7.8, 4.4 Hz, 1H), 3.88 (q, <math>J</math> = 7.9 Hz, 1H), 3.83 (s, 3H), 3.67-3.62 (m, 2H), 3.50 (t, <math>J</math> = 6.4 Hz, 2H), 2.91-2.86 (m, 2H), 2.78-2.68 (m, 4H), 2.48 (s, 3H), 2.42-2.33 (m, 1H), 2.13 (p, <math>J</math> = 6.2 Hz, 1H); 461.21[M+H]<sup>+</sup></p>
35 40 45 50 55	<p>68</p> 	<p>(R)-N-(4-(4-allylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.81 (s, 1H), 8.15 (s, 1H), 7.44-7.31 (m, 4H), 7.29-7.17 (m, 3H), 6.97-6.91 (m, 2H), 6.26 (s, 1H), 6.03-5.91 (m, 1H), 5.62 (dd, <math>J</math> = 8.6, 4.6 Hz, 1H), 5.47-5.36 (m, 2H), 4.17 (td, <math>J</math> = 7.7, 4.9 Hz, 1H), 3.94 (q, <math>J</math> = 7.7 Hz, 1H), 3.45 (d, <math>J</math> = 7.0 Hz, 2H), 3.39 (t, <math>J</math> = 5.1 Hz, 4H), 3.04 (t, <math>J</math> = 5.0 Hz, 4H), 2.83-2.73 (m, 1H), 2.45-2.35 (m, 1H); 443.3[M+H]<sup>+</sup></p>



5 10 15 20	<p>69</p> 	<p>N-(4-((R)-3- (dimethylamino)pyrr olidin-1- yl)phenyl)-6-((R)- 3- phenylisoxazolidin- 2-yl)pyrimidin-4- amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 10.89 (s, 1H), 8.01 (s, 1H), 7.39-7.27 (m, 6H), 7.18-7.13 (m, 2H), 6.60 (d, <i>J</i> = 8.7 Hz, 1H), 6.02 (s, 1H), 5.58 (dd, <i>J</i> = 8.7, 4.5 Hz, 1H), 4.29-4.20 (m, 1H), 4.04 (q, <i>J</i> = 7.5 Hz, 1H), 3.90- 3.83 (m, 1H), 3.75-3.55 (m, 3H), 3.38 (q, <i>J</i> = 8.3 Hz, 1H), 2.86 (s, 3H), 2.68 (s, 3H), 2.50-2.34 (m, 4H); 431.3 [M+H]<sup>+</sup></p>
25 30 35 40	<p>70</p> 	<p>N-(4-((S)-2- (dimethylamino)met hyl)pyrrolidin-1- yl)phenyl)-6-((R)- 3- phenylisoxazolidin- 2-yl)pyrimidin-4- amine</p>	<p>445.3 [M+H]<sup>+</sup></p>
45 50 55	<p>71</p> 	<p>(R)-N-(1-(1-(2- (dimethylamino)ethy l)piperidin-4-yl)- 1H-pyrazol-4-yl)-6- (3- phenylisoxazolidin-</p>	<p>463.3 [M+H]<sup>+</sup></p>



5		2-yl)pyrimidin-4-amine	
72		(R)-4-methylpiperazin-1-yl (4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)methanone	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.66 (s, 1H), 8.24 (s, 1H), 7.47-7.40 (m, 4H), 7.39-7.33 (m, 4H), 7.31-7.27 (m, 1H), 6.57 (s, 1H), 5.63 (dd, <i>J</i> = 8.7, 4.8 Hz, 1H), 4.20 (td, <i>J</i> = 7.7, 4.5 Hz, 1H), 3.93 (q, <i>J</i> = 7.8 Hz, 1H), 3.74 (s, 4H), 2.83-2.74 (m, 1H), 2.63 (s, 4H), 2.48-2.37 (m, 4H); 445.24 [M+H] <sup>+</sup>
73		(R)-6-(3-phenylisoxazolidin-2-yl)-N-(3-(trifluoromethyl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.36 (s, 1H), 7.65 (s, 1H), 7.58 (d, <i>J</i> = 8.5 Hz, 1H), 7.48 (t, <i>J</i> = 8.8 Hz, 1H), 7.45 (d, <i>J</i> = 7.4 Hz, 2H), 7.36 (t, <i>J</i> = 7.7 Hz, 3H), 7.28 (d, <i>J</i> = 7.7 Hz, 1H), 6.94 (s, 1H), 6.52 (d, <i>J</i> = 1.0 Hz, 1H), 5.67 (dd, <i>J</i> = 8.7, 4.8 Hz, 1H), 4.18 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.91 (q, <i>J</i> = 7.9 Hz, 1H), 2.77 (dtd, <i>J</i> = 12.2, 8.0, 4.4 Hz, 1H), 2.46-2.36 (m, 1H); 387.17 [M+H] <sup>+</sup>




5 10 15 20 25 30	74 	N-(5-((1S,4S)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-2-fluoropyridin-3-yl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.37 (d, <i>J</i> = 1.0 Hz, 1H), 7.92 (dd, <i>J</i> = 8.6, 2.9 Hz, 1H), 7.48-7.41 (m, 2H), 7.37 (t, <i>J</i> = 7.6 Hz, 2H), 7.30-7.27 (m, 1H), 7.17-7.09 (m, 1H), 6.90 (s, 1H), 6.46 (d, <i>J</i> = 0.8 Hz, 1H), 5.66 (dd, <i>J</i> = 8.6, 4.8 Hz, 1H), 4.67 (d, <i>J</i> = 2.2 Hz, 1H), 4.38 (s, 1H), 4.19 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.97-3.86 (m, 3H), 3.59 (dd, <i>J</i> = 9.1, 1.6 Hz, 1H), 3.18 (d, <i>J</i> = 9.2 Hz, 1H), 2.78 (dddd, <i>J</i> = 12.0, 8.6, 7.5, 4.4 Hz, 1H), 2.43 (dtd, <i>J</i> = 12.5, 7.9, 4.7 Hz, 1H), 2.02 (ddd, <i>J</i> = 22.7, 9.8, 2.4 Hz, 2H); 435.27 [M+H] <sup>+</sup>
35 40 45 50 55	75 	(R)-1-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)pyrrolidin-2-one	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.26 (s, 1H), 7.63 (d, <i>J</i> = 8.7 Hz, 2H), 7.50 (s, 1H), 7.44 (d, <i>J</i> = 7.7 Hz, 2H), 7.39-7.27 (m, 5H), 6.49 (d, <i>J</i> = 1.1 Hz, 1H), 5.65 (dd, <i>J</i> = 8.7, 4.8 Hz, 1H), 4.14 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.88 (q, <i>J</i> = 7.2 Hz, 3H), 2.75 (dtd, <i>J</i> = 12.5, 8.0, 4.3 Hz, 1H), 2.64 (t, <i>J</i> = 8.1 Hz, 2H), 2.44-2.35 (m,







5			1H), 2.20 (dd, $J = 13.7, 6.3$ Hz, 2H); 402.21 [M+H] <sup>+</sup>
10 15 20 25 30	76 	(R)-2-methyl-5-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)isoindolin-1-one	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.32 (d, $J = 1.0$ Hz, 1H), 7.81 (d, $J = 8.2$ Hz, 1H), 7.71 (s, 1H), 7.59 (d, $J = 1.2$ Hz, 1H), 7.47-7.41 (m, 2H), 7.37 (td, $J = 6.9, 1.6$ Hz, 2H), 7.33-7.27 (m, 2H), 6.58 (d, $J = 1.0$ Hz, 1H), 5.65 (dd, $J = 8.6, 4.8$ Hz, 1H), 4.38 (s, 2H), 4.18 (td, $J = 7.8, 4.5$ Hz, 1H), 3.91 (q, $J = 7.9$ Hz, 1H), 3.20 (s, 3H), 2.83-2.74 (m, 1H), 2.42 (dtd, $J = 12.5, 7.9, 4.8$ Hz, 1H); 388.23 [M+H] <sup>+</sup>
35 40 45 50 55	77 	(R)-6-(3-phenylisoxazolidin-2-yl)-N-(4-(piperazin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.19 (d, $J = 1.0$ Hz, 1H), 7.97 (s, 1H), 7.44-7.40 (m, 2H), 7.35 (dd, $J = 8.5, 6.8$ Hz, 2H), 7.28-7.24 (m, 1H), 7.22-7.17 (m, 2H), 6.98-6.90 (m, 2H), 6.39 (d, $J = 1.1$ Hz, 1H), 5.64 (dd, $J = 8.6, 4.7$ Hz, 1H), 4.12 (td, $J = 7.7, 4.4$ Hz, 1H), 3.87 (q, $J = 7.9$ Hz, 1H), 3.25 (dd, $J = 6.5, 3.5$ Hz, 4H), 3.17 (dd, $J = 6.5, 3.5$ Hz, 4H), 2.74 (dddd,



5			$J = 12.1, 8.7, 7.6, 4.5 \text{ Hz, 1H}), 2.38$ $(\text{dtd, } J = 12.4, 7.9, 4.7 \text{ Hz, 1H});$ $403.23 [\text{M}+\text{H}]^+$
10 15 20 25 30	78	 <p>(R)-N-(6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-1,2,3,4-tetrahydroisoquinolin-6-amine</p>	$^1\text{H NMR (400 MHz, Chloroform-}d) \delta 8.23$ $(s, 1\text{H}), 7.81 (s, 1\text{H}), 7.46\text{--}7.30 (m,$ $4\text{H}), 7.26\text{--}7.22 (m, 1\text{H}), 7.20\text{--}7.12 (m,$ $2\text{H}), 7.04 (d, } J = 8.1 \text{ Hz, 1H}), 6.46$ $(s, 1\text{H}), 5.61 (dd, } J = 8.6, 4.8 \text{ Hz,}$ $1\text{H}), 5.13 (s, 1\text{H}), 4.23 (s, 2\text{H}), 4.14$ $(\text{td, } J = 7.8, 4.4 \text{ Hz, 1H}), 3.88 (q,$ $J = 7.8 \text{ Hz, 1H}), 3.43\text{--}3.33 (m, 2\text{H}),$ $3.09\text{--}2.98 (m, 2\text{H}), 2.80\text{--}2.67 (m, 1\text{H}),$ $2.43\text{--}2.33 (m, 1\text{H}); 374.2 [\text{M}+\text{H}]^+$
35 40 45 50 55	79	 <p>(R)-N-(5-(4-(dimethylamino) piperidin-1-yl)pyridin-2-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	$^1\text{H NMR (400 MHz, Chloroform-}d) \delta 8.76$ $(s, 1\text{H}), 8.30 (d, } J = 1.0 \text{ Hz, 1H),$ $7.96 (d, } J = 3.0 \text{ Hz, 1H), 7.47 (dd,$ $J = 9.0, 7.1 \text{ Hz, 3H}), 7.36 (dd, } J =$ $8.4, 6.8 \text{ Hz, 2H}), 7.32\text{--}7.28 (m, 1\text{H}),$ $7.26\text{--}7.24 (m, 2\text{H}), 5.67 (dd, } J = 8.6,$ $4.7 \text{ Hz, 1H}), 4.21 (\text{td, } J = 7.9, 4.6$ $\text{Hz, 1H}), 3.99 (q, } J = 7.8 \text{ Hz, 2H),$ $3.66 (d, } J = 12.1 \text{ Hz, 2H), 2.87\text{--}2.71$ $(m, 4\text{H}), 2.55 (s, 6\text{H}), 2.42 (\text{ddt, } J$ $= 11.5, 7.5, 3.8 \text{ Hz, 1H}), 2.07 (s,$

5			2H), 1.80 (qd, $J = 12.1, 4.1$ Hz, 1H); 446.30 [M+H] <sup>+</sup>
80		(R)-N-(8-(1-methyl-1H-pyrazol-5-yl)-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.39 (d, $J = 1.0$ Hz, 1H), 7.63 (d, $J = 8.4$ Hz, 1H), 7.53 (d, $J = 1.9$ Hz, 1H), 7.48-7.44 (m, 2H), 7.39-7.33 (m, 2H), 7.30-7.27 (m, 1H), 7.01 (s, 1H), 6.86 (d, $J = 8.4$ Hz, 1H), 6.61 (d, $J = 1.0$ Hz, 1H), 6.27 (d, $J = 1.9$ Hz, 1H), 5.69 (dd, $J = 8.6, 4.7$ Hz, 1H), 4.40-4.29 (m, 4H), 4.24-4.15 (m, 1H), 3.95 (q, $J = 7.8$ Hz, 1H), 3.79 (s, 3H), 2.84-2.71 (m, 1H), 2.48-2.37 (m, 1H); 457.2 [M+H] <sup>+</sup>
81		(R)-1-cyclopropyl-4-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-1,4-azaphosphinane 4-oxide	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 10.67 (s, 1H), 8.13 (s, 1H), 7.68-7.60 (m, 1H), 7.53-7.43 (m, 2H), 7.41-7.25 (m, 5H), 6.03 (s, 1H), 5.61 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.32-4.25 (m, 1H), 4.08 (q, $J = 7.5$ Hz, 1H), 4.03-3.79 (m, 7H), 3.03 (td, $J = 14.7, 4.9$ Hz, 2H), 2.94-2.82 (m, 1H), 2.64-2.57 (m, 1H), 2.50-2.40 (m, 1H), 2.18 (t, $J = 17.1$ Hz, 2H), 1.42-1.34 (m, 2H), 1.00-0.91

			(m, 2H); 506.3 [M+H] <sup>+</sup>
5			
10	82	 <p>N-(4-((2R,6S)-2,6-dimethylmorpholino)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	432.3 [M+H] <sup>+</sup>
15			
20			
25	83	 <p>6-((R)-3-phenylisoxazolidin-2-yl)-N-((R)-7-(pyrrolidin-1-yl)-6,7,8,9-tetrahydro-5H-benzo[7]annulen-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 12.65 (s, 1H), 11.67 (s, 1H), 8.01 (s, 1H), 7.40-7.29 (m, 4H), 7.20-7.04 (m, 3H), 6.15 (s, 1H), 5.59 (dd, <i>J</i> = 8.6, 4.4 Hz, 1H), 4.29 (q, <i>J</i> = 7.0 Hz, 1H), 4.11 (q, <i>J</i> = 7.4 Hz, 1H), 3.83-3.70 (m, 2H), 3.41 (t, <i>J</i> = 11.7 Hz, 1H), 2.98-2.87 (m, 5H), 2.78 (t, <i>J</i> = 13.1 Hz, 2H), 2.51-2.36 (m, 3H), 2.22-2.10 (m, 2H), 2.05-1.95 (m, 2H), 1.70-1.56 (m, 2H); 456.3 [M+H] <sup>+</sup>
30			
35			
40			
45	84	 <p>(R)-N,N-dimethyl-1'-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-[1,4'-bipiperidin]-</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.27 (d, <i>J</i> = 1.0 Hz, 1H), 7.46-7.40 (m, 2H), 7.38-7.31 (m, 2H), 7.24 (d, <i>J</i> = 7.2 Hz, 1H), 7.19-7.15 (m, 2H), 6.94 (d, <i>J</i> = 8.9 Hz, 2H), 6.53 (s, 1H), 6.36 (d, <i>J</i> = 1.1 Hz, 1H), 5.67 (dd,
50			
55			


5		4-amine	$J = 8.6, 4.6 \text{ Hz}, 1\text{H}$ ), 4.14-4.06 (m, 1H), 3.86 (q, $J = 7.9 \text{ Hz}, 1\text{H}$ ), 3.74 (d, $J = 12.1 \text{ Hz}, 2\text{H}$ ), 3.05 (d, $J = 11.3 \text{ Hz}, 2\text{H}$ ), 2.78-2.67 (m, 3H), 2.52-2.35 (m, 3H), 2.33 (s, 6H), 2.29-2.19 (m, 3H), 1.98-1.83 (m, 5H), 1.72 (d, $J = 3.7 \text{ Hz}, 2\text{H}$ ); 528.3 [M+H] <sup>+</sup>
20 25 30 35 40	85 	(R)-N-(5-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-2-(4-(dimethylamino)-[1,4'-bipiperidin]-1'-yl)-4-methoxyphenyl)propanamide	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.77 (s, 1H), 8.35 (s, 1H), 8.30 (s, 1H), 7.04-6.96 (m, 2H), 6.94 (s, 1H), 6.75-6.64 (m, 3H), 5.71-5.61 (m, 1H), 4.19-4.11 (m, 1H), 4.05 (q, $J = 8.1 \text{ Hz}, 1\text{H}$ ), 3.83 (s, 3H), 3.17-3.01 (m, 4H), 2.82-2.66 (m, 3H), 2.46-2.29 (m, 11H), 2.24-2.16 (m, 3H), 1.94-1.86 (m, 2H), 1.75-1.54 (m, 5H), 1.31-1.23 (m, 3H); 665.6 [M+H] <sup>+</sup>
45 50 55	86 	3-chloro-N-(5-((6-((R)-3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-2-(4-(dimethylamino)-[1,4'-bipiperidin]-1'-yl)-4-methoxyphenyl)propanamide	698.3 [M+H] <sup>+</sup>

5 10		(hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)piperidin-1-yl)-4-(methoxyphenyl)propionamide	
15 20 25 30 35 40	87 	(R)-N-(5-((6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)-[1,4'-bipiperidin]-1'-yl)phenyl)propionamide	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.72 (s, 1H), 8.36-8.24 (m, 2H), 7.55 (q, <i>J</i> = 7.9 Hz, 1H), 6.85-6.77 (m, 2H), 6.71 (d, <i>J</i> = 10.1 Hz, 2H), 5.86 (dd, <i>J</i> = 8.8, 4.4 Hz, 1H), 4.14-4.00 (m, 3H), 3.85-3.80 (m, 3H), 3.21 (d, <i>J</i> = 11.1 Hz, 2H), 3.11-2.99 (m, 3H), 2.82-2.66 (m, 11H), 2.63-2.53 (m, 2H), 2.48-2.41 (m, 3H), 2.39 (s, 3H), 2.35-2.22 (m, 4H), 1.96-1.88 (m, 3H), 1.81-1.63 (m, 5H); 720.7[M+H] <sup>+</sup>
45 50 55	88 	(R)-7-(5-methoxy-2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-N,N-dimethyl-7-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.29-8.27 (m, 1H), 7.44 (s, 1H), 7.11 (s, 1H), 7.03-6.96 (m, 2H), 6.73-6.65 (m, 1H), 6.61 (s, 1H), 6.43-6.39 (m, 1H), 5.65 (dd, <i>J</i> = 8.8, 4.6 Hz, 1H), 4.11 (td, <i>J</i> = 7.9, 4.1 Hz, 1H), 3.90-3.84

5  10		azaspiro[3.5]nonan-2-amine	(m, 1H), 3.83 (s, 3H), 3.43-3.34 (m, 5H), 3.02-2.92 (m, 1H), 2.83-2.70 (m, 5H), 2.35 (s, 6H), 2.24 (s, 3H), 2.15-2.10 (m, 2H), 2.03-1.98 (m, 2H); 565.3[M+H] <sup>+</sup>
15  20  25  30  35  40	89  	(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(naphthalen-2-yl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.28 (s, 1H), 7.91 (d, <i>J</i> = 1.6 Hz, 1H), 7.86-7.78 (m, 3H), 7.54 (dd, <i>J</i> = 8.5, 1.8 Hz, 1H), 7.49-7.41 (m, 3H), 7.08 (s, 1H), 6.57-6.48 (m, 2H), 6.40 (s, 1H), 5.82 (dd, <i>J</i> = 8.6, 4.7 Hz, 1H), 4.15 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.93 (q, <i>J</i> = 7.9 Hz, 1H), 3.84 (s, 3H), 3.71 (d, <i>J</i> = 12.2 Hz, 2H), 2.76 (dt, <i>J</i> = 19.9, 12.4 Hz, 1H), 2.51-2.42 (m, 2H), 2.39 (s, 3H), 1.97 (d, <i>J</i> = 12.5 Hz, 2H), 1.71 (ddd, <i>J</i> = 24.0, 12.1, 3.8 Hz, 2H); 580.44[M+H] <sup>+</sup>
45  50  55	90  	(R)-6-(3-(3,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.29 (s, 1H), 7.49 (d, <i>J</i> = 8.8 Hz, 1H), 7.32-7.27 (m, 1H), 7.18-7.08 (m, 2H), 6.84 (s, 1H), 6.54 (d, <i>J</i> = 7.2 Hz, 2H), 6.39 (d, <i>J</i> = 1.1 Hz, 1H), 5.63 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.10 (td,

EP 4 342 895 A1



5		yl)phenyl)pyrimidin-4-amine	$J = 7.9, 4.3 \text{ Hz, 1H}$ ), 3.92-3.80 (m, 4H), 3.71 (d, $J = 12.3 \text{ Hz, 2H}$ ), 2.84-2.47 (m, 11H), 2.42 (dq, $J = 11.5, 3.4 \text{ Hz, 1H}$ ), 2.34 (s, 4H), 1.97 (d, $J = 14.1 \text{ Hz, 2H}$ ), 1.75-1.65 (m, 2H); 566.31 [M+H] <sup>+</sup>
15 20 25 30 35 40 45	91	(R)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (d, $J = 2.9 \text{ Hz, 1H}$ ), 7.48 (d, $J = 2.9 \text{ Hz, 1H}$ ), 7.42 (ddd, $J = 8.5, 5.3, 2.8 \text{ Hz, 2H}$ ), 7.16 (s, 1H), 7.03 (td, $J = 8.7, 2.9 \text{ Hz, 2H}$ ), 6.67 (d, $J = 3.0 \text{ Hz, 1H}$ ), 6.48 (d, $J = 2.9 \text{ Hz, 1H}$ ), 5.65 (dd, $J = 8.4, 4.5 \text{ Hz, 1H}$ ), 4.11 (td, $J = 7.8, 4.1 \text{ Hz, 1H}$ ), 3.92-3.86 (m, 1H), 3.83 (d, $J = 2.9 \text{ Hz, 3H}$ ), 3.13 (d, $J = 11.3 \text{ Hz, 2H}$ ), 2.84-2.59 (m, 13H), 2.50 (d, $J = 12.7 \text{ Hz, 1H}$ ), 2.40 (d, $J = 2.9 \text{ Hz, 3H}$ ), 2.35 (dt, $J = 8.0, 4.3 \text{ Hz, 1H}$ ), 1.96 (d, $J = 12.1 \text{ Hz, 2H}$ ), 1.78-1.68 (m, 2H), 1.22 (t, $J = 7.5 \text{ Hz, 3H}$ ); 576.42 [M+H] <sup>+</sup>




50

55





<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>92</p>		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.31 (s, 1H), 7.52 (s, 1H), 7.33 (t, <i>J</i> = 7.4 Hz, 1H), 7.05 (t, <i>J</i> = 6.2 Hz, 2H), 6.91 (s, 1H), 6.67 (s, 1H), 6.53 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.10 (dq, <i>J</i> = 7.6, 4.2 Hz, 1H), 3.89 (q, <i>J</i> = 8.2, 7.8 Hz, 1H), 3.84 (s, 3H), 3.13 (d, <i>J</i> = 11.4 Hz, 2H), 2.88-2.49 (m, 13H), 2.43-2.36 (m, 1H), 2.33 (s, 3H), 2.32-2.27 (m, 1H), 1.95 (d, <i>J</i> = 13.0 Hz, 2H), 1.78-1.68 (m, 2H), 1.25-1.21 (m, 3H); 594.33 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p> <p>93</p>		<p>(R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.32 (d, <i>J</i> = 1.0 Hz, 1H), 7.58-7.50 (m, 2H), 6.88-6.78 (m, 3H), 6.67 (s, 1H), 6.54 (d, <i>J</i> = 1.0 Hz, 1H), 5.85 (dd, <i>J</i> = 8.7, 4.5 Hz, 1H), 4.08 (td, <i>J</i> = 7.9, 4.4 Hz, 1H), 3.90 (q, <i>J</i> = 7.9 Hz, 1H), 3.84 (s, 3H), 3.13 (d, <i>J</i> = 11.5 Hz, 2H), 2.82-2.50 (m, 12H), 2.40-2.24 (m, 5H), 1.96 (d, <i>J</i> = 12.1 Hz, 2H), 1.72 (qd, <i>J</i> = 12.0, 3.8 Hz, 3H), 1.23 (t, <i>J</i> = 7.5 Hz, 3H); 594.33 [M+H]<sup>+</sup></p>


5 10 15 20 25 30	<p>94</p>	<p>(R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.30 (d, <i>J</i> = 1.0 Hz, 1H), 7.49 (s, 1H), 7.29 (td, <i>J</i> = 6.0, 3.0 Hz, 1H), 7.17 (s, 1H), 7.00 (td, <i>J</i> = 9.1, 4.3 Hz, 1H), 6.96-6.85 (m, 1H), 6.67 (s, 1H), 6.51 (d, <i>J</i> = 1.1 Hz, 1H), 5.87 (dd, <i>J</i> = 9.0, 4.5 Hz, 1H), 4.08 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.88 (q, <i>J</i> = 7.9 Hz, 1H), 3.84 (s, 3H), 3.13 (d, <i>J</i> = 11.6 Hz, 2H), 2.90-2.59 (m, 13H), 2.54-2.44 (m, 1H), 2.40 (s, 3H), 2.34-2.24 (m, 1H), 1.96 (d, <i>J</i> = 12.1 Hz, 2H), 1.73 (qd, <i>J</i> = 11.8, 3.7 Hz, 2H), 1.23 (t, <i>J</i> = 7.5 Hz, 3H); 594.28 [M+H]<sup>+</sup></p>
35 40 45 50 55	<p>95</p>	<p>(R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) <math>\delta</math> 9.26 (s, 1H), 8.24 (s, 1H), 7.50 (dd, <i>J</i> = 8.8, 6.6 Hz, 1H), 7.27 (ddd, <i>J</i> = 11.2, 9.2, 2.6 Hz, 1H), 7.16 (d, <i>J</i> = 2.4 Hz, 1H), 7.07 (ddt, <i>J</i> = 11.0, 5.5, 2.4 Hz, 2H), 6.84 (d, <i>J</i> = 8.5 Hz, 1H), 6.43 (d, <i>J</i> = 1.0 Hz, 1H), 5.70 (dd, <i>J</i> = 8.7, 4.9 Hz, 1H), 4.18 (td, <i>J</i> = 7.9, 3.9 Hz, 1H), 3.89 (t, <i>J</i> = 8.1 Hz, 1H), 3.76</p>



5			(s, 3H), 2.77 (dq, $J = 8.1, 4.1$ Hz, 1H), 2.38-2.18 (m, 6H), 2.15 (s, 3H), 1.80 (d, $J = 12.1$ Hz, 2H), 1.76-1.64 (m, 1H), 1.54 (dt, $J = 12.1, 5.9$ Hz, 2H), 1.32-1.21 (m, 2H), 1.19-0.98 (m, 1H), 0.95-0.80 (m, 2H); 566.4 [M+H] <sup>+</sup>
15 20 25 30 35 40 45	96	 <p>(R)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-2-amine</p>	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) $\delta$ 9.22 (s, 1H), 8.23 (d, $J = 1.0$ Hz, 1H), 7.44 (ddd, $J = 8.8, 5.6, 2.6$ Hz, 2H), 7.21-7.12 (m, 3H), 7.05 (dd, $J = 8.5, 2.3$ Hz, 1H), 6.84 (d, $J = 8.5$ Hz, 1H), 6.40 (d, $J = 1.1$ Hz, 1H), 5.53 (dd, $J = 8.6, 5.0$ Hz, 1H), 4.16 (td, $J = 7.9, 4.0$ Hz, 1H), 3.85 (d, $J = 8.1$ Hz, 1H), 3.76 (s, 3H), 2.75 (dq, $J = 8.3, 4.3$ Hz, 1H), 2.46 (s, 2H), 2.37-2.22 (m, 6H), 2.15 (s, 3H), 2.09 (d, $J = 1.0$ Hz, 1H), 1.80 (d, $J = 12.0$ Hz, 2H), 1.54 (dt, $J = 11.6, 5.9$ Hz, 2H), 1.25 (d, $J = 3.4$ Hz, 2H), 1.05-0.80 (m, 2H); 548.3 [M+H] <sup>+</sup>

50

55

<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p> <p>35</p>	<p>97</p> 	<p>N-(4-(4-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.24 (s, 1H), 7.46 (d, <i>J</i> = 8.5 Hz, 1H), 7.42 (d, <i>J</i> = 7.3 Hz, 2H), 7.35 (t, <i>J</i> = 7.6 Hz, 2H), 7.28 (d, <i>J</i> = 1.2 Hz, 1H), 7.24 (t, <i>J</i> = 1.4 Hz, 1H), 6.55 (d, <i>J</i> = 2.5 Hz, 1H), 6.51 (dd, <i>J</i> = 8.6, 2.5 Hz, 1H), 6.29 (s, 1H), 5.65 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.61 (s, 1H), 4.45 (s, 1H), 4.22-4.09 (m, 2H), 3.90 (q, <i>J</i> = 7.8 Hz, 1H), 3.84 (s, 4H), 3.72 (d, <i>J</i> = 12.3 Hz, 3H), 2.98 (s, 2H), 2.74 (ddd, <i>J</i> = 11.9, 7.9, 3.9 Hz, 4H), 2.39 (ddt, <i>J</i> = 11.7, 7.7, 3.9 Hz, 2H), 2.29 (d, <i>J</i> = 11.1 Hz, 1H), 2.16-2.10 (m, 2H), 1.99 (d, <i>J</i> = 12.8 Hz, 1H); 529.39 [M+H]<sup>+</sup></p>
<p>40</p> <p>45</p> <p>50</p> <p>55</p>	<p>98</p> 	<p>(R)-N-(4-(4-(diethylamino)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.26 (d, <i>J</i> = 1.0 Hz, 1H), 7.50 (d, <i>J</i> = 8.3 Hz, 1H), 7.45-7.41 (m, 2H), 7.35 (dd, <i>J</i> = 8.4, 6.7 Hz, 3H), 7.27-7.22 (m, 1H), 6.52 (d, <i>J</i> = 8.6 Hz, 2H), 6.35 (d, <i>J</i> = 1.0 Hz, 1H), 5.66 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.12 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.89 (q, <i>J</i> = 7.8 Hz, 1H),</p>



5 10 15			3.84 (s, 3H), 3.76 (d, $J = 13.3$ Hz, 2H), 3.39 (tt, $J = 12.2, 3.7$ Hz, 1H), 3.16 (q, $J = 7.3$ Hz, 4H), 2.80 (t, $J = 11.3$ Hz, 2H), 2.72 (tt, $J = 8.0, 4.4$ Hz, 1H), 2.43-2.32 (m, 1H), 2.16 (d, $J = 9.3$ Hz, 2H), 1.98 (q, $J = 12.4, 4.1$ Hz, 2H), 1.39 (t, $J = 7.3$ Hz, 6H); 503.40 [M+H] <sup>+</sup>
20 25 30 35 40 45 50	99	 <p>N-(2-methoxy-4-((R)-2-methyl-4-(1-methylpiperidin-4-yl)piperazin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	1H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.27 (s, 1H), 7.54 (d, $J = 6.6$ Hz, 1H), 7.43 (d, $J = 7.6$ Hz, 2H), 7.35 (t, $J = 7.6$ Hz, 2H), 7.28-7.24 (m, 1H), 6.56 (d, $J = 6.1$ Hz, 2H), 6.34 (s, 1H), 5.66 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.14 (td, $J = 7.8, 4.5$ Hz, 1H), 3.91 (q, $J = 7.7$ Hz, 1H), 3.84 (s, 3H), 3.74 (s, 1H), 3.25-3.10 (m, 3H), 2.89-2.84 (m, 1H), 2.80-2.70 (m, 5H), 2.67-2.61 (m, 2H), 2.44-2.35 (m, 2H), 2.24-2.19 (m, 1H), 2.08 (s, 3H), 2.03-1.98 (m, 2H), 1.31 (s, 1H), 1.06 (d, $J = 6.1$ Hz, 3H), 0.87-0.80 (m, 1H); 544.39 [M+H] <sup>+</sup>

<p>5</p> <p>10</p> <p>15</p> <p>20</p>		<p>isopropyl (R)-3-(2-</p> <p>(6-((2-methoxy-4-</p> <p>(4-(4-</p> <p>methylpiperazin-1-</p> <p>yl)piperidin-1-</p> <p>yl)phenyl)amino)pyr</p> <p>imidin-4-</p> <p>yl)isoxazolidin-3-</p> <p>yl)benzoate</p>	<p>616.4 [M+H]<sup>+</sup></p>
<p>25</p> <p>30</p> <p>35</p> <p>40</p>		<p>(R)-N-(2-methoxy-4-</p> <p>(4-(1-</p> <p>methylpiperidin-4-</p> <p>yl)piperazin-1-</p> <p>yl)phenyl)-6-(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.21</p> <p>(s, 1H), 8.00 (s, 1H), 7.44-7.31 (m,</p> <p>5H), 7.29-7.26 (m, 1H), 6.55-6.49 (m,</p> <p>2H), 6.24 (s, 1H), 5.64 (dd, <i>J</i> = 8.6,</p> <p>4.6 Hz, 1H), 4.18-4.09 (m, 1H), 3.90</p> <p>(q, <i>J</i> = 7.8 Hz, 1H), 3.84 (s, 3H),</p> <p>3.50 (s, 2H), 3.24 (t, <i>J</i> = 4.8 Hz,</p> <p>4H), 2.89-2.76 (m, 5H), 2.75 (s, 3H),</p> <p>2.74-2.61 (m, 2H), 2.44-2.34 (m, 1H),</p> <p>2.18-2.06 (m, 5H); 530.4 [M+H]<sup>+</sup></p>

45



50

55

<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p>	<p>10</p> <p>2</p> 	<p>(R)-N-(3-methoxy-4-(4-(1-methylpiperidin-4-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.29 (d, <i>J</i> = 1.0 Hz, 1H), 7.46-7.41 (m, 2H), 7.38-7.31 (m, 2H), 7.26-7.22 (m, 1H), 6.95-6.90 (m, 1H), 6.86-6.80 (m, 2H), 6.60 (s, 1H), 6.45 (d, <i>J</i> = 1.0 Hz, 1H), 5.66 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.16-4.07 (m, 1H), 3.91-3.82 (m, 4H), 3.11 (s, 4H), 2.98 (d, <i>J</i> = 11.2 Hz, 2H), 2.82-2.76 (m, 4H), 2.71 (s, 1H), 2.44-2.34 (m, 2H), 2.32 (s, 3H), 2.11-1.96 (m, 2H), 1.88 (d, <i>J</i> = 12.4 Hz, 2H), 1.76-1.68 (m, 2H); 530.3 [M+H]<sup>+</sup></p>
<p>35</p> <p>40</p> <p>45</p>	<p>10</p> <p>3</p> 	<p>(R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperidin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>550.44 [M+H]<sup>+</sup></p>



50

55



<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p>	<p>10</p> <p>4</p> 	<p>(R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.23 (d, <i>J</i> = 1.0 Hz, 1H), 7.51 (q, <i>J</i> = 8.3 Hz, 1H), 7.41 (s, 1H), 7.12-7.06 (m, 2H), 7.00 (d, <i>J</i> = 9.2 Hz, 1H), 6.88-6.77 (m, 2H), 6.47 (d, <i>J</i> = 1.0 Hz, 1H), 5.83 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.08 (td, <i>J</i> = 7.9, 4.4 Hz, 1H), 3.86 (q, <i>J</i> = 7.9 Hz, 1H), 3.18 (d, <i>J</i> = 11.7 Hz, 2H), 2.78-2.69 (m, 11H), 2.44 (dtd, <i>J</i> = 11.7, 7.6, 3.8 Hz, 1H), 2.37 (s, 3H), 2.30 (s, 3H), 2.28-2.23 (m, 1H), 1.96 (d, <i>J</i> = 12.2 Hz, 2H), 1.72 (qd, <i>J</i> = 12.2, 3.3 Hz, 2H);</p> <p>550.44 [M+H]<sup>+</sup></p>
<p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p>	<p>10</p> <p>5</p> 	<p>(R)-6-(3-(4-chloro-2-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.26 (s, 1H), 7.73 (s, 1H), 7.49 (t, <i>J</i> = 8.4 Hz, 1H), 7.13-7.04 (m, 2H), 6.92 (d, <i>J</i> = 8.9 Hz, 1H), 6.86-6.79 (m, 2H), 6.48 (s, 1H), 5.82 (dd, <i>J</i> = 8.7, 4.6 Hz, 1H), 4.06 (td, <i>J</i> = 7.9, 4.3 Hz, 1H), 3.86 (s, 3H), 3.85-3.79 (m, 1H), 3.55 (d, <i>J</i> = 11.1 Hz, 2H), 2.83-2.75 (m, 1H), 2.72-2.39 (m, 11H), 2.31 (s, 3H), 2.29-2.18 (m, 1H), 1.96-1.88 (m,</p>







EP 4 342 895 A1



5			2H), 1.80 (qd, $J = 12.0, 3.7$ Hz, 2H); 582.3[M+H] <sup>+</sup>
10 15 20 25 30		(R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.28 (d, $J = 1.1$ Hz, 1H), 7.29 (dd, $J = 5.9, 3.2$ Hz, 1H), 7.05–6.80 (m, 6H), 6.48 (d, $J = 1.1$ Hz, 1H), 5.85 (dd, $J = 8.8, 4.6$ Hz, 1H), 4.07 (td, $J = 7.8, 4.2$ Hz, 1H), 3.87 (s, 3H), 3.85–3.81 (m, 1H), 3.55 (d, $J = 11.6$ Hz, 2H), 2.86–2.76 (m, 1H), 2.73–2.55 (m, 7H), 2.54–2.39 (m, 4H), 2.31 (s, 3H), 2.30–2.21 (m, 1H), 1.92 (d, $J = 12.3$ Hz, 2H), 1.81 (qd, $J = 11.9, 3.8$ Hz, 2H); 566.4[M+H] <sup>+</sup>
35 40 45 50 55		(R)-N-(3-methyl-4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (d, $J = 1.1$ Hz, 1H), 7.44 (d, $J = 7.3$ Hz, 2H), 7.35 (t, $J = 7.7$ Hz, 2H), 7.25–7.22 (m, 1H), 7.14–7.02 (m, 3H), 6.59 (s, 1H), 6.45 (d, $J = 1.1$ Hz, 1H), 5.67 (dd, $J = 8.7, 4.7$ Hz, 1H), 4.12 (td, $J = 7.8, 4.4$ Hz, 1H), 3.88 (q, $J = 7.8$ Hz, 1H), 3.06–2.95 (m, 4H), 2.80–2.59 (m, 5H), 2.43 (s, 3H), 2.42–2.37 (m, 1H), 2.31 (s, 3H);



EP 4 342 895 A1



			431.3 [M+H] <sup>+</sup>
5  10  15  20  25		(R)-N-(2-methyl-4-morpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.26 (s, 1H), 7.46-7.30 (m, 4H), 7.25-7.22 (m, 2H), 6.84-6.76 (m, 2H), 6.36 (s, 1H), 6.04 (d, <i>J</i> = 1.2 Hz, 1H), 5.66 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.07 (td, <i>J</i> = 7.9, 4.4 Hz, 1H), 3.87 (q, <i>J</i> = 7.0, 5.9 Hz, 4H), 3.84-3.79 (m, 1H), 3.18 (t, <i>J</i> = 4.9 Hz, 4H), 2.78-2.65 (m, 1H), 2.42-2.30 (m, 1H), 2.23 (s, 3H); 418.3 [M+H] <sup>+</sup>
30  35  40  45  50  55		(R)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 10.93 (s, 1H), 8.07-8.02 (m, 1H), 7.41-7.27 (m, 5H), 7.11 (s, 1H), 6.65 (s, 1H), 5.88-5.80 (m, 1H), 5.59 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.24 (td, <i>J</i> = 7.6, 5.3 Hz, 1H), 4.02 (q, <i>J</i> = 7.5 Hz, 1H), 3.80 (s, 3H), 3.40 (s, 7H), 3.22 (d, <i>J</i> = 11.7 Hz, 2H), 3.10-2.98 (m, 1H), 2.91-2.82 (m, 1H), 2.79 (s, 3H), 2.77-2.69 (m, 2H), 2.59 (q, <i>J</i> = 7.5 Hz, 2H), 2.49-2.39 (m, 1H), 2.13-2.05 (m, 3H), 1.95-1.83 (m, 2H), 1.20 (t, <i>J</i> = 7.5 Hz, 3H); 558.4 [M+H] <sup>+</sup>



5 10 15 20 25 30	 <p>11 0</p>	<p>(R)-N-(2-methoxy-4-morpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.29 (d, <math>J = 1.0</math> Hz, 1H), 7.54 (d, <math>J = 9.3</math> Hz, 1H), 7.44 (d, <math>J = 7.5</math> Hz, 2H), 7.35 (dd, <math>J = 8.4, 6.8</math> Hz, 2H), 7.27-7.24 (m, 1H), 6.98 (s, 1H), 6.53 (dd, <math>J = 6.3, 2.6</math> Hz, 2H), 6.40 (d, <math>J = 1.0</math> Hz, 1H), 5.67 (dd, <math>J = 8.7, 4.6</math> Hz, 1H), 4.12 (td, <math>J = 7.8, 4.4</math> Hz, 1H), 3.93-3.86 (m, 5H), 3.85 (s, 3H), 3.20-3.11 (m, 4H), 2.72 (dtd, <math>J = 12.3, 8.0, 4.4</math> Hz, 1H), 2.39 (ddt, <math>J = 11.7, 7.7, 3.9</math> Hz, 1H); 434.25 [M+H]<sup>+</sup></p>
35 40 45 50 55	 <p>11 1</p>	<p>(R)-N4-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-2,4-diamine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 7.44-7.39 (m, 3H), 7.34 (dd, <math>J = 8.4, 6.7</math> Hz, 2H), 7.24 (d, <math>J = 7.2</math> Hz, 2H), 7.15 (s, 1H), 6.55-6.47 (m, 2H), 5.61 (dd, <math>J = 8.6, 4.4</math> Hz, 1H), 5.07 (s, 2H), 4.06 (td, <math>J = 7.8, 4.6</math> Hz, 1H), 3.86 (q, <math>J = 7.9</math> Hz, 1H), 3.83 (s, 3H), 3.70 (d, <math>J = 12.2</math> Hz, 2H), 2.80-2.61 (m, 11H), 2.38 (s, 3H), 2.33-2.30 (m, 2H), 1.97 (d, <math>J = 12.5</math> Hz, 2H), 1.72 (q, <math>J = 13.1, 9.2</math> Hz, 2H);</p>

			545.40 [M+H] <sup>+</sup>
5			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.33
10		(R)-N-(2-methoxy-4-	(s, 1H), 7.59-7.28 (m, 7H), 6.55-6.45
11		(4-(4-	(m, 1H), 6.26 (s, 1H), 5.69-5.45 (m,
15		methylpiperazin-1-	1H), 4.31 (s, 1H), 4.07 (q, <i>J</i> = 7.7
20	yl)piperidin-1-	Hz, 1H), 3.83 (s, 3H), 3.64 (d, <i>J</i> =	
25	yl)phenyl)-4-(3-	11.9 Hz, 2H), 2.91-2.83 (m, 1H), 2.69	
	phenylisoxazolidin-	(t, <i>J</i> = 11.4 Hz, 6H), 2.58 (s, 3H),	
	2-yl)-1,3,5-	2.47-2.37 (m, 3H), 2.34 (s, 3H), 1.95	
	triazin-2-amine	(d, <i>J</i> = 12.5 Hz, 2H), 1.78-1.68 (m,	
		2H); 531.3 [M+H] <sup>+</sup>	
30			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 12.30
35		(R)-2-methoxy-N4-	(s, 1H), 10.70-10.51 (m, 1H), 8.02-
40		(1-methylpiperidin-	7.96 (m, 1H), 7.40-7.26 (m, 5H), 6.99
45		4-yl)-N1-(6-(3-	(d, <i>J</i> = 8.4 Hz, 1H), 6.32-6.18 (m,
50	phenylisoxazolidin-	1H), 5.71 (s, 1H), 5.56 (dd, <i>J</i> = 8.6,	
	2-yl)pyrimidin-4-	4.6 Hz, 1H), 4.24 (q, <i>J</i> = 7.1 Hz, 1H),	
	yl)benzene-1,4-	4.03 (q, <i>J</i> = 7.5 Hz, 1H), 3.79-3.68	
	diamine	(m, 3H), 3.67-3.49 (m, 3H), 3.36 (d,	
		<i>J</i> = 12.3 Hz, 1H), 3.25-3.12 (m, 1H),	
		2.95-2.75 (m, 5H), 2.48-2.37 (m, 1H),	
		2.24 (d, <i>J</i> = 14.3 Hz, 2H), 1.99 (q,	
		<i>J</i> = 13.8, 11.8 Hz, 2H); 461.3 [M+H] <sup>+</sup>	



5 10 15 20		<p>(R)-N-(2-methoxy-5-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 9.13 (s, 1H), 8.15 (s, 1H), 7.39-7.17 (m, 6H), 6.61 (s, 1H), 6.08 (s, 1H), 5.62 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.21-4.12 (m, 1H), 3.95 (q, <i>J</i> = 7.7 Hz, 1H), 3.81 (s, 3H), 3.42-3.19 (m, 3H), 3.09 (s, 5H), 2.86-2.57 (m, 7H), 2.46-2.36 (m, 1H), 2.21 (s, 3H), 2.09-1.97 (m, 4H), 1.85-1.72 (m, 2H); 544.4 [M+H] <sup>+</sup>
25 30 35 40 45 50		<p>(R)-1'-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-N,N-dimethyl-[1,4'-bipiperidin]-4-amine</p>	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 10.21 (s, 1H), 9.83 (s, 1H), 9.07 (s, 1H), 8.19 (d, <i>J</i> = 0.9 Hz, 1H), 7.38-7.35 (m, 3H), 7.29-7.26 (m, 1H), 6.69 (d, <i>J</i> = 2.5 Hz, 1H), 6.57 (dd, <i>J</i> = 8.8, 2.5 Hz, 1H), 6.05 (s, 1H), 5.51 (dd, <i>J</i> = 8.6, 5.2 Hz, 1H), 4.20 (td, <i>J</i> = 7.7, 4.2 Hz, 1H), 3.97-3.87 (m, 3H), 3.79 (s, 3H), 3.69 (d, <i>J</i> = 12.0 Hz, 2H), 3.48-3.35 (m, 2H), 3.11-2.99 (m, 2H), 2.88-2.70 (m, 8H), 2.50 (s, 6H), 2.32-2.24 (m, 2H), 2.14-2.08 (m, 2H); 558.5 [M+H] <sup>+</sup>

<p>5</p> <p>11</p> <p>6</p> <p>15</p> <p>20</p>		<p>N-(4-(4-((1R,4R)-5-ethyl-2,5-diazabicyclo[2.2.1]heptan-2-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 7.98 (s, 1H), 7.37-6.98 (m, 7H), 6.64-6.44 (m, 2H), 5.73 (s, 1H), 5.56-5.40 (m, 1H), 4.65-4.54 (m, 1H), 4.44-4.30 (m, 1H), 4.23 (q, <i>J</i> = 7.1 Hz, 1H), 4.13-3.93 (m, 2H), 3.91-3.56 (m, 6H), 3.54-3.04 (m, 5H), 2.95-2.69 (m, 3H), 2.57-2.25 (m, 3H), 2.19-1.74 (m, 5H), 1.34-1.23 (m, 2H); 556.4 [M+H]<sup>+</sup></p>
<p>25</p> <p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p>		<p>(R)-N-(3-ethyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.20 (d, <i>J</i> = 1.0 Hz, 1H), 8.10 (s, 1H), 7.45-7.39 (m, 2H), 7.35 (dd, <i>J</i> = 8.5, 6.8 Hz, 2H), 7.26-7.22 (m, 1H), 7.12 (d, <i>J</i> = 2.4 Hz, 1H), 7.10-7.02 (m, 2H), 6.49 (d, <i>J</i> = 1.0 Hz, 1H), 5.65 (dd, <i>J</i> = 8.6, 4.7 Hz, 1H), 4.13 (td, <i>J</i> = 7.8, 4.4 Hz, 1H), 3.88 (q, <i>J</i> = 7.9 Hz, 1H), 3.12 (d, <i>J</i> = 12.0 Hz, 2H), 2.87 (s, 6H), 2.71 (dq, <i>J</i> = 26.2, 8.1, 7.3, 4.3 Hz, 7H), 2.60-2.52 (m, 1H), 2.48 (s, 3H), 2.39 (dtd, <i>J</i> = 12.5, 7.9, 4.7 Hz, 1H), 1.99-1.92 (m, 2H), 1.74 (qd, <i>J</i> =</p>

5			11.7, 3.7 Hz, 2H), 1.23 (t, $J = 7.5$ Hz, 3H); 528.42 [M+H] <sup>+</sup>
11 8		(R)-N-(4-(4-(4-ethylpiperazin-1-yl)piperidin-1-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.30 (s, 1H), 7.51 (d, $J = 8.5$ Hz, 1H), 7.45 (d, $J = 7.6$ Hz, 2H), 7.34 (t, $J = 7.6$ Hz, 2H), 7.25-7.22 (m, 1H), 6.70 (s, 1H), 6.57-6.52 (m, 2H), 6.41 (s, 1H), 5.68 (dd, $J = 8.7, 4.5$ Hz, 1H), 4.16-4.07 (m, 1H), 3.92-3.86 (m, 1H), 3.84 (s, 3H), 3.71 (d, $J = 12.2$ Hz, 2H), 2.78-2.65 (m, 6H), 2.54 (s, 2H), 2.51-2.35 (m, 5H), 1.97 (d, $J = 12.6$ Hz, 2H), 1.75-1.68 (m, 4H), 1.11 (t, $J = 7.2$ Hz, 3H); 544.3 [M+H] <sup>+</sup>
11 9		(R)-N-(4-(4-(dimethylamino)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.23 (s, 1H), 7.83 (s, 1H), 7.48 (d, $J = 8.6$ Hz, 1H), 7.43-7.38 (m, 2H), 7.34 (td, $J = 8.4, 7.9, 2.8$ Hz, 2H), 7.27-7.23 (m, 1H), 6.55-6.47 (m, 2H), 6.26 (d, $J = 2.8$ Hz, 1H), 5.64 (dd, $J = 8.6, 4.5$ Hz, 1H), 4.14 (td, $J = 7.7, 4.5$ Hz, 1H), 3.95-3.88 (m, 1H), 3.83 (s, 3H), 3.80-3.73 (m, 2H), 3.30-3.18 (m, 1H), 2.86-2.71 (m, 9H), 2.45-2.34



5			(m, 1H), 2.19–2.12 (m, 2H), 1.90 (qd, $J = 12.2, 3.8$ Hz, 2H); 475.3 [M+H] <sup>+</sup>
10 15 20 25 30		(R)-N-(2-methoxy-4-((4-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine)morpholinopiperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.28 (s, 1H), 7.50–7.41 (m, 3H), 7.34 (t, $J = 7.7$ Hz, 2H), 7.26–7.22 (m, 1H), 6.98 (s, 1H), 6.57–6.51 (m, 2H), 6.40–6.36 (m, 1H), 5.67 (dd, $J = 8.7, 4.5$ Hz, 1H), 4.11 (td, $J = 7.8, 4.5$ Hz, 1H), 3.92–3.86 (m, 1H), 3.84 (s, 3H), 3.76 (t, $J = 4.6$ Hz, 3H), 3.71 (d, $J = 12.1$ Hz, 2H), 2.79–2.70 (m, 3H), 2.63 (t, $J = 4.7$ Hz, 3H), 2.43–2.34 (m, 3H), 2.07 (s, 1H), 1.98 (d, $J = 12.4$ Hz, 2H), 1.70 (qd, $J = 12.1, 3.9$ Hz, 2H); 517.3 [M+H] <sup>+</sup>
35 40 45 50		(R)-2-(4-(1-(2-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperidin-4-yl)piperazin-1-yl)ethan-1-ol	560.4 [M+H] <sup>+</sup>





<p>5</p> <p>10</p> <p>12</p> <p>2</p> <p>15</p> <p>20</p> <p>25</p>		<p>(R)-1-(4-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperazin-1-yl)ethan-1-one</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.21-8.10 (m, 2H), 7.45-7.38 (m, 2H), 7.34 (t, <math>J</math> = 7.5 Hz, 2H), 7.26-7.17 (m, 3H), 6.94 (d, <math>J</math> = 8.8 Hz, 2H), 6.39 (d, <math>J</math> = 1.1 Hz, 1H), 5.64 (dd, <math>J</math> = 8.7, 4.7 Hz, 1H), 4.16-4.07 (m, 1H), 3.87 (q, <math>J</math> = 7.8 Hz, 1H), 3.79 (t, <math>J</math> = 5.3 Hz, 2H), 3.64 (t, <math>J</math> = 5.2 Hz, 2H), 3.22-3.11 (m, 4H), 2.80-2.67 (m, 1H), 2.44-2.32 (m, 1H), 2.15 (s, 3H); 445.3 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>12</p> <p>3</p> <p>40</p> <p>45</p>		<p>(R)-N-(3,5-difluoro-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.27 (s, 1H), 7.66 (s, 1H), 7.46-7.33 (m, 4H), 7.31-7.26 (m, 1H), 6.92 (d, <math>J</math> = 9.9 Hz, 2H), 6.40 (s, 1H), 5.65 (dd, <math>J</math> = 8.7, 4.7 Hz, 1H), 4.25-4.17 (m, 1H), 3.97 (q, <math>J</math> = 7.7 Hz, 1H), 3.27 (d, <math>J</math> = 11.9 Hz, 3H), 3.12 (t, <math>J</math> = 11.8 Hz, 4H), 2.90 (s, 3H), 2.85-2.75 (m, 3H), 2.66 (s, 3H), 2.48-2.38 (m, 2H), 1.99-1.84 (m, 4H); 536.3 [M+H]<sup>+</sup></p>

50



55



5 10 12 4 15 20		(R)-N-(4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 10.52 (s, 1H), 8.48 (s, 1H), 8.04 (s, 1H), 7.36-7.27 (m, 4H), 7.22-7.13 (m, 2H), 6.97-6.87 (m, 2H), 6.14 (s, 1H), 5.58 (dd, $J$ = 8.6, 4.6 Hz, 1H), 4.74-4.64 (m, 4H), 4.25-4.15 (m, 1H), 4.04-3.95 (m, 1H), 3.66-3.56 (m, 1H), 3.27 (t, $J$ = 5.1 Hz, 4H), 2.88-2.77 (m, 1H), 2.57 (t, $J$ = 5.0 Hz, 4H), 2.45-2.35 (m, 1H); 459.3 [M+H] <sup>+</sup>
25 30 35 12 5 40 45 50		(R)-6-(3-(3-chloro-2-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.27 (s, 1H), 7.50-7.41 (m, 1H), 7.34-7.25 (m, 2H), 7.05 (t, $J$ = 7.9 Hz, 1H), 6.93 (d, $J$ = 9.0 Hz, 1H), 6.87-6.79 (m, 2H), 6.48 (d, $J$ = 1.1 Hz, 1H), 5.87 (dd, $J$ = 8.9, 4.8 Hz, 1H), 4.07 (tt, $J$ = 9.3, 4.6 Hz, 1H), 3.86-3.81 (m, 4H), 3.55 (d, $J$ = 11.1 Hz, 2H), 2.87-2.77 (m, 1H), 2.74-2.54 (m, 7H), 2.43 (ddt, $J$ = 11.1, 7.3, 3.8 Hz, 4H), 2.31 (s, 3H), 2.29-2.23 (m, 1H), 1.92 (d, $J$ = 12.2 Hz, 2H), 1.81 (q, $J$ = 12.0, 3.8 Hz, 2H); 582.3 [M+H] <sup>+</sup>



<p>5</p> <p>12</p> <p>10</p> <p>6</p> <p>15</p>		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>550.39 [M+H]<sup>+</sup></p>
<p>20</p> <p>25</p> <p>30</p> <p>12</p> <p>7</p> <p>35</p> <p>40</p> <p>45</p>		<p>(R)-N-(4-(4-(dimethylamino)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.27 (d, <i>J</i> = 1.1 Hz, 1H), 7.46-7.41 (m, 2H), 7.35 (dd, <i>J</i> = 8.4, 6.8 Hz, 2H), 7.28 (t, <i>J</i> = 1.5 Hz, 1H), 7.23-7.19 (m, 2H), 6.97-6.92 (m, 2H), 6.37 (d, <i>J</i> = 1.1 Hz, 1H), 5.67 (dd, <i>J</i> = 8.7, 4.7 Hz, 1H), 4.12 (td, <i>J</i> = 7.8, 4.5 Hz, 1H), 3.88 (q, <i>J</i> = 7.9 Hz, 1H), 3.79 (d, <i>J</i> = 12.6 Hz, 2H), 3.23-3.09 (m, 1H), 2.85-2.67 (m, 9H), 2.44-2.33 (m, 1H), 2.15 (d, <i>J</i> = 11.6 Hz, 2H), 1.89 (qd, <i>J</i> = 12.1, 4.1 Hz, 3H);</p> <p>445.24 [M+H]<sup>+</sup></p>

50

55

5 10 15 20 25 30		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.28 (d, <i>J</i> = 1.0 Hz, 1H), 7.35–7.28 (m, 1H), 7.13–7.01 (m, 2H), 7.00–6.90 (m, 1H), 6.89–6.79 (m, 3H), 6.47 (d, <i>J</i> = 1.1 Hz, 1H), 5.89 (dd, <i>J</i> = 8.8, 4.8 Hz, 1H), 4.09 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.87 (s, 3H), 3.86–3.81 (m, 1H), 3.55 (d, <i>J</i> = 11.3 Hz, 2H), 2.86–2.77 (m, 1H), 2.75–2.51 (m, 9H), 2.49–2.40 (m, 2H), 2.33 (s, 3H), 2.30–2.27 (m, 1H), 1.93 (d, <i>J</i> = 12.3 Hz, 2H), 1.81 (qd, <i>J</i> = 11.9, 3.8 Hz, 2H); 566.4 [M+H]<sup>+</sup></p>
35 40 45 50 55		<p>(R)-6-(3-(4-chloro-3-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.27 (s, 1H), 7.47 (d, <i>J</i> = 8.5 Hz, 1H), 7.36 (t, <i>J</i> = 7.8 Hz, 1H), 7.28 (d, <i>J</i> = 1.8 Hz, 1H), 7.16 (dd, <i>J</i> = 8.3, 1.9 Hz, 1H), 6.95 (s, 1H), 6.54 (d, <i>J</i> = 7.3 Hz, 2H), 6.37 (s, 1H), 5.63 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.09 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.90–3.80 (m, 4H), 3.71 (d, <i>J</i> = 12.2 Hz, 2H), 2.78–2.65 (m, 10H), 2.48–2.40 (m, 2H), 2.37–2.25 (m, 4H), 1.97 (d, <i>J</i> = 12.5 Hz,</p>

5			2H), 1.70 (ddd, $J = 24.4, 12.4, 3.3$ Hz, 2H); 582.28 [M+H] <sup>+</sup>
10 15 20 25 30		(R)-6-(3-(3-chloro-2,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.26 (s, 1H), 7.45 (td, $J = 8.8, 5.6$ Hz, 2H), 7.14 (s, 1H), 6.95 (td, $J = 8.6, 1.8$ Hz, 1H), 6.54 (d, $J = 7.8$ Hz, 2H), 6.38 (s, 1H), 5.83 (dd, $J = 8.9, 4.7$ Hz, 1H), 4.08 (td, $J = 8.0, 4.1$ Hz, 1H), 3.84 (s, 4H), 3.73 (s, 2H), 2.77 (td, $J = 16.1, 12.7, 6.9$ Hz, 11H), 2.49 (ddt, $J = 11.5, 7.5, 3.6$ Hz, 1H), 2.40 (s, 3H), 2.25 (dtd, $J = 12.7, 8.0, 4.8$ Hz, 1H), 1.98 (d, $J = 12.7$ Hz, 2H), 1.71 (qtd, $J = 12.1, 3.9$ Hz, 2H); 600.28 [M+H] <sup>+</sup>
35 40 45 50 55		(R)-6-(3-(3-(dimethylamino)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.26 (s, 1H), 7.44 (d, $J = 8.3$ Hz, 1H), 7.21 (t, $J = 7.9$ Hz, 1H), 7.16 (s, 1H), 6.82 (t, $J = 2.1$ Hz, 1H), 6.77 (d, $J = 7.5$ Hz, 1H), 6.63 (dd, $J = 8.3, 2.6$ Hz, 1H), 6.55-6.48 (m, 2H), 6.35 (s, 1H), 5.59 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.11 (td, $J = 7.8, 4.5$ Hz, 1H), 3.89-3.81 (m, 4H), 3.70 (d, $J =$

5  10			11.9 Hz, 2H), 2.95 (s, 6H), 2.83–2.65 (m, 11H), 2.52–2.45 (m, 1H), 2.40 (s, 4H), 1.97 (d, $J = 12.7$ Hz, 2H), 1.71 (qd, $J = 12.0, 4.0$ Hz, 2H); 573.42 [M+H] <sup>+</sup>
15  20  25  30  35  40	13  2	 <p>(R)-N-(5-chloro-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.36 (d, $J = 1.0$ Hz, 1H), 7.98 (s, 1H), 7.47–7.43 (m, 2H), 7.35 (t, $J = 7.6$ Hz, 2H), 7.24 (s, 1H), 6.80 (s, 1H), 6.62 (s, 1H), 6.43 (d, $J = 1.0$ Hz, 1H), 5.68 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.16 (td, $J = 7.9, 4.5$ Hz, 1H), 3.92 (q, $J = 7.8$ Hz, 1H), 3.86 (s, 3H), 3.44 (d, $J = 11.4$ Hz, 2H), 2.80–2.61 (m, 7H), 2.51 (s, 3H), 2.46–2.35 (m, 3H), 2.31 (s, 3H), 1.94 (d, $J = 12.2$ Hz, 2H), 1.85–1.74 (m, 2H); 564.3 [M+H] <sup>+</sup>
45  50  55	13  3	 <p>(R)-N-(1-methyl-1H-pyrazol-4-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 11.36 (s, 1H), 7.98 (s, 1H), 7.49 (d, $J = 26.4$ Hz, 2H), 7.41–7.26 (m, 5H), 6.03 (s, 1H), 5.59 (dd, $J = 8.7, 4.5$ Hz, 1H), 4.30 (q, $J = 6.8$ Hz, 1H), 4.10 (q, $J = 7.4$ Hz, 1H), 3.92 (s, 3H),

EP 4 342 895 A1

5


10

15

4

20

25

			2.96–2.85 (m, 1H), 2.52–2.41 (m, 1H); 323.2 [M+H] <sup>+</sup>
	<p>tert-butyl (R)-7-</p> <p>((6-(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>yl)amino)-3,4-</p> <p>dihydroisoquinoline</p> <p>-2(1H)-carboxylate</p> 	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 11.65</p> <p>(s, 1H), 8.04–7.97 (m, 1H), 7.37 (dt,</p> <p><i>J</i> = 7.9, 4.5 Hz, 2H), 7.34–7.27 (m,</p> <p>3H), 7.25–6.98 (m, 3H), 6.19–6.09 (m,</p> <p>1H), 5.63–5.54 (m, 1H), 4.59 (s, 1H),</p> <p>4.32–4.25 (m, 1H), 4.16–4.07 (m, 1H),</p> <p>3.75–3.54 (m, 2H), 3.18–3.00 (m, 1H),</p> <p>2.97–2.76 (m, 3H), 2.50–2.40 (m, 1H),</p> <p>1.67–1.20 (m, 9H); 474.3 [M+H]<sup>+</sup></p>	


30

35



40

45



50

	<p>(R)-N-(4-(9-methyl-</p> <p>3,9-</p> <p>diazaspiro[5.5]unde-</p> <p>can-3-yl)phenyl)-6-</p> <p>(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p> 	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.23</p> <p>(d, <i>J</i> = 1.1 Hz, 1H), 7.46–7.41 (m,</p> <p>2H), 7.37–7.32 (m, 2H), 7.26–7.21 (m,</p> <p>2H), 7.20–7.13 (m, 2H), 6.98–6.92 (m,</p> <p>2H), 6.36 (d, <i>J</i> = 1.1 Hz, 1H), 5.65</p> <p>(dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.09 (td,</p> <p><i>J</i> = 7.8, 4.5 Hz, 1H), 3.85 (q, <i>J</i> =</p> <p>7.8 Hz, 1H), 3.20–3.13 (m, 4H), 3.05</p> <p>(s, 2H), 2.77–2.68 (m, 1H), 2.64 (s,</p> <p>3H), 2.42 (s, 3H), 2.40–2.33 (m, 1H),</p> <p>1.72–1.65 (m, 7H); 485.4 [M+H]<sup>+</sup></p>
--	---	---



55

5 10 13 15 20 25		<p>N-(4-(6-methyl-3,6-diazabicyclo[3.1.1]heptan-3-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.21 (d, <i>J</i> = 1.0 Hz, 1H), 7.54 (s, 1H), 7.46–7.30 (m, 4H), 7.27–7.18 (m, 3H), 6.79–6.72 (m, 2H), 6.33 (d, <i>J</i> = 1.1 Hz, 1H), 5.67 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.14–4.04 (m, 1H), 3.95 (d, <i>J</i> = 6.0 Hz, 2H), 3.90–3.81 (m, 2H), 3.62 (d, <i>J</i> = 11.4 Hz, 2H), 3.47 (d, <i>J</i> = 11.3 Hz, 2H), 2.80 (q, <i>J</i> = 6.5 Hz, 1H), 2.78–2.66 (m, 1H), 2.37 (dt, <i>J</i> = 7.9, 4.5 Hz, 1H), 2.25 (s, 2H), 1.71 (d, <i>J</i> = 9.0 Hz, 1H); 429.3 [M+H] <sup>+</sup>
30 35 40 45 50		<p>(R)-3-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)oxazolidin-2-one</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.36 (d, <i>J</i> = 1.0 Hz, 1H), 7.91 (d, <i>J</i> = 8.7 Hz, 1H), 7.66 (d, <i>J</i> = 2.4 Hz, 1H), 7.48–7.43 (m, 2H), 7.38–7.32 (m, 2H), 7.25–7.21 (m, 1H), 6.97 (s, 1H), 6.73 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H), 6.50 (d, <i>J</i> = 1.0 Hz, 1H), 5.68 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.56–4.46 (m, 2H), 4.20–4.05 (m, 3H), 3.96–3.87 (m, 4H), 2.80–2.69 (m, 1H), 2.47–2.35 (m, 1H); 434.2 [M+H] <sup>+</sup>





<p>5</p> <p>10</p> <p>13</p> <p>15</p> <p>20</p> <p>25</p>		<p>(R)-N-(5-(4-</p> <p>(dimethylamino)piper</p> <p>ridin-1-yl)pyridin-</p> <p>2-yl)-6-(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.34</p> <p>(s, 1H), 8.31 (d, <i>J</i> = 1.0 Hz, 1H),</p> <p>7.97 (d, <i>J</i> = 2.9 Hz, 1H), 7.50-7.42</p> <p>(m, 3H), 7.40-7.34 (m, 2H), 7.32-7.27</p> <p>(m, 1H), 7.26-7.23 (m, 1H), 7.19 (s,</p> <p>1H), 5.67 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H),</p> <p>4.23-4.17 (m, 1H), 4.01-3.95 (m, 1H),</p> <p>3.64 (d, <i>J</i> = 12.0 Hz, 2H), 2.83-2.68</p> <p>(m, 3H), 2.62-2.51 (m, 1H), 2.48-2.34</p> <p>(m, 7H), 2.01 (d, <i>J</i> = 12.5 Hz, 2H),</p> <p>1.73 (qd, <i>J</i> = 12.1, 4.1 Hz, 2H); 446.3</p> <p>[M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p>		<p>(R)-N-(4-(4-</p> <p>methylpiperazin-1-</p> <p>yl)-3-(2,2,2-</p> <p>trifluoroethoxy)phe</p> <p>nyl)-6-(3-</p> <p>phenylisoxazolidin-</p> <p>2-yl)pyrimidin-4-</p> <p>amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 13.44</p> <p>(s, 1H), 11.75 (s, 1H), 8.01 (d, <i>J</i> =</p> <p>0.8 Hz, 1H), 7.41-7.26 (m, 4H), 7.07-</p> <p>6.94 (m, 2H), 6.78 (d, <i>J</i> = 2.0 Hz,</p> <p>1H), 6.11-6.03 (m, 1H), 5.59 (dd, <i>J</i></p> <p>= 8.6, 4.4 Hz, 1H), 4.40 (q, <i>J</i> = 8.0</p> <p>Hz, 2H), 4.36-4.29 (m, 1H), 4.17-4.10</p> <p>(m, 1H), 3.67 (d, <i>J</i> = 11.5 Hz, 2H),</p> <p>3.57-3.46 (m, 2H), 3.39-3.25 (m, 2H),</p> <p>3.06 (t, <i>J</i> = 11.4 Hz, 2H), 2.96-2.85</p> <p>(m, 4H), 2.52-2.42 (m, 1H); 515.3</p> <p>[M+H]<sup>+</sup></p>

EP 4 342 895 A1



<p>5</p> <p>10</p> <p>14</p> <p>0</p> <p>15</p> <p>20</p>		<p>6-( (R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(3-fluoro-4-(4-methylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 10.84 (s, 1H), 8.05 (s, 1H), 7.18-6.98 (m, 4H), 6.56-6.47 (m, 2H), 5.83-5.75 (m, 2H), 4.28-4.19 (m, 1H), 4.11-3.96 (m, 2H), 3.89 (d, <i>J</i> = 12.8 Hz, 1H), 3.81 (s, 3H), 3.75-3.28 (m, 8H), 3.17-2.99 (m, 2H), 2.98-2.89 (m, 2H), 2.85 (s, 3H), 2.46-2.34 (m, 1H), 2.32-2.19 (m, 1H), 2.05 (s, 1H), 2.00 (d, <i>J</i> = 12.5 Hz, 1H); 584.4 [M+H]<sup>+</sup></p>
<p>25</p> <p>30</p> <p>35</p> <p>40</p> <p>45</p>		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(1-methylpiperidin-4-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 11.01 (s, 1H), 8.07 (s, 1H), 7.21 (d, <i>J</i> = 7.9 Hz, 1H), 7.17-6.98 (m, 3H), 6.89-6.82 (m, 2H), 5.86 (s, 1H), 5.80 (dd, <i>J</i> = 8.7, 5.0 Hz, 1H), 4.27 (td, <i>J</i> = 7.6, 5.0 Hz, 1H), 4.06 (q, <i>J</i> = 7.6 Hz, 1H), 3.84 (s, 3H), 3.74-3.66 (m, 2H), 3.01-2.83 (m, 6H), 2.81-2.72 (m, 1H), 2.47-2.37 (m, 1H), 2.28 (q, <i>J</i> = 12.3 Hz, 2H), 2.12-2.03 (m, 2H); 482.3 [M+H]<sup>+</sup></p>



50



55

<p>5</p> <p>10</p> <p>14</p> <p>15</p> <p>2</p> <p>20</p> <p>25</p>		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-isopropyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 11.18 (s, 1H), 8.07 (s, 1H), 7.17–7.01 (m, 4H), 6.67 (s, 1H), 5.93 (s, 1H), 5.81 (dd, <i>J</i> = 8.7, 5.0 Hz, 1H), 4.23 (td, <i>J</i> = 7.6, 4.7 Hz, 1H), 4.01 (q, <i>J</i> = 7.6 Hz, 1H), 3.81 (s, 3H), 3.67 (s, 7H), 3.34–3.18 (m, 4H), 3.01–2.91 (m, 1H), 2.89 (s, 3H), 2.81 (t, <i>J</i> = 11.8 Hz, 2H), 2.40 (td, <i>J</i> = 12.7, 7.4 Hz, 1H), 2.18 (d, <i>J</i> = 11.9 Hz, 2H), 2.09 (s, 1H), 2.06–1.92 (m, 2H), 1.18 (dd, <i>J</i> = 6.9, 2.2 Hz, 6H); 608.4 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>14</p> <p>3</p> <p>45</p> <p>50</p> <p>55</p>		<p>(R)-N-(5-cyclopropyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 10.86 (s, 1H), 8.04 (s, 1H), 7.41–7.26 (m, 5H), 6.68 (s, 1H), 6.59 (s, 1H), 5.74 (s, 1H), 5.58 (dd, <i>J</i> = 8.6, 4.6 Hz, 1H), 4.25 (td, <i>J</i> = 7.6, 5.5 Hz, 1H), 4.03 (q, <i>J</i> = 7.5 Hz, 1H), 3.97–3.91 (m, 1H), 3.79 (s, 3H), 3.78–3.64 (m, 7H), 3.56 (d, <i>J</i> = 11.6 Hz, 2H), 3.30 (t, <i>J</i> = 12.2 Hz, 1H), 2.91 (s, 3H), 2.89–2.85 (m, 1H), 2.84–2.75 (m, 2H), 2.50–2.40 (m, 1H), 2.22 (d, <i>J</i> = 11.4 Hz, 2H), 2.09–1.97 (m, 3H), 1.03–0.97</p>



EP 4 342 895 A1



5			(m, 2H), 0.66-0.60 (m, 2H); 570.5 [M+H] <sup>+</sup>
10 15 20 25 30 35	<p>14 4</p> 	<p>(R)-N-(4-(4-methylpiperazin-1-yl)-2-(3-((methylsulfonyl)methyl)azetidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.18 (d, <i>J</i> = 1.0 Hz, 1H), 7.45-7.40 (m, 2H), 7.37-7.31 (m, 2H), 7.25-7.22 (m, 1H), 7.00 (d, <i>J</i> = 8.6 Hz, 1H), 6.95 (s, 1H), 6.42 (dd, <i>J</i> = 8.6, 2.6 Hz, 1H), 6.05 (d, <i>J</i> = 2.7 Hz, 1H), 6.03 (d, <i>J</i> = 1.1 Hz, 1H), 5.63 (dd, <i>J</i> = 8.7, 4.7 Hz, 1H), 4.14-4.08 (m, 3H), 3.84 (q, <i>J</i> = 7.9 Hz, 1H), 3.73-3.65 (m, 2H), 3.29 (d, <i>J</i> = 7.4 Hz, 2H), 3.24 (t, <i>J</i> = 5.1 Hz, 3H), 3.19-3.10 (m, 1H), 2.88 (s, 3H), 2.77-2.70 (m, 2H), 2.65 (t, <i>J</i> = 5.0 Hz, 4H), 2.39 (s, 3H), 2.38-2.33 (m, 1H); 564.4 [M+H]<sup>+</sup></p>
40 45 50 55	<p>14 5</p> 	<p>(R)-1-(5-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)phenyl)acetyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.34 (s, 1H), 8.04 (s, 1H), 7.36-7.30 (m, 1H), 7.11-7.02 (m, 2H), 6.76 (s, 1H), 6.62 (s, 1H), 6.48 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.13 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.98-3.93 (m, 1H), 3.91 (s, 3H), 3.33-3.24 (m, 2H),</p>

5		methylpiperazin-1-yl)piperidin-1-yl)phenyl)ethan-1-one	2.89–2.73 (m, 4H), 2.67 (s, 6H), 2.58–2.40 (m, 4H), 2.37–2.25 (m, 5H), 2.00 (d, $J = 12.4$ Hz, 2H), 1.81–1.68 (m, 2H); 608.4 [M+H] <sup>+</sup>
10 15 20 25 30	<p>14 6</p> 	(R,E)-1-(5-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)ethan-1-one O-methyl oxime	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 9.46–9.18 (m, 1H), 8.11 (s, 1H), 7.33 (s, 1H), 7.18–6.98 (m, 3H), 6.61 (s, 1H), 5.99 (s, 1H), 5.89–5.74 (m, 1H), 4.31–4.22 (m, 1H), 4.12–4.02 (m, 1H), 3.96 (s, 3H), 3.91–3.71 (m, 6H), 3.61–3.34 (m, 6H), 3.01–2.69 (m, 7H), 2.50–2.37 (m, 2H), 2.36–2.24 (m, 2H), 2.23–2.14 (m, 3H), 2.04–1.87 (m, 2H); 637.4 [M+H] <sup>+</sup>
35 40 45 50 55	<p>14 7</p> 	(R)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-6-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.32 (d, $J = 1.0$ Hz, 1H), 8.07 (s, 1H), 7.82 (d, $J = 0.7$ Hz, 1H), 7.75 (s, 1H), 7.47–7.42 (m, 2H), 7.38–7.31 (m, 2H), 7.25–7.20 (m, 1H), 6.85 (s, 1H), 6.37 (d, $J = 1.1$ Hz, 1H), 5.68 (dd, $J = 8.7, 4.6$ Hz, 1H), 4.15 (td, $J = 7.9, 4.6$ Hz, 1H), 3.99–3.88 (m, 6H), 3.55 (d, $J = 12.4$ Hz, 2H), 2.78–2.62

5		amine	(m, 9H), 2.46–2.38 (m, 1H), 2.35 (s, 4H), 2.06 (s, 3H), 1.92 (d, $J = 12.4$ Hz, 2H), 1.64–1.54 (m, 2H); 611.5 [M+H] <sup>+</sup>
10 15 20 25 30 35	14 8	 <p>(R)-6-(3-(4-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine)-4-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-2-azolidin-2-yl)-N-(2-fluorophenyl)isoxazolidin-2-yl</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.33 (d, $J = 1.0$ Hz, 1H), 7.90–7.84 (m, 2H), 7.78 (s, 1H), 7.46–7.37 (m, 2H), 7.08–6.97 (m, 2H), 6.84 (s, 1H), 6.70 (s, 1H), 6.51 (d, $J = 1.1$ Hz, 1H), 5.66 (dd, $J = 8.6, 4.6$ Hz, 1H), 4.13 (td, $J = 7.8, 4.3$ Hz, 1H), 3.95 (s, 3H), 3.94–3.89 (m, 1H), 3.87 (s, 3H), 3.23 (d, $J = 10.9$ Hz, 2H), 2.84–2.42 (m, 11H), 2.40–2.34 (m, 1H), 2.31 (s, 3H), 2.28–2.20 (m, 1H), 1.99–1.92 (m, 2H), 1.61 (dt, $J = 11.9, 3.8$ Hz, 2H); 628.4 [M+H] <sup>+</sup>
40 45 50 55	14 9	 <p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-2-azolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.33 (d, $J = 1.0$ Hz, 1H), 7.88 (s, 1H), 7.87 (s, 1H), 7.78 (s, 1H), 7.38–7.29 (m, 1H), 7.13–6.99 (m, 2H), 6.90 (s, 1H), 6.71 (s, 1H), 6.53 (d, $J = 1.1$ Hz, 1H), 5.92 (dd, $J = 8.8, 4.7$ Hz, 1H), 4.11 (td, $J = 7.9, 4.2$ Hz, 1H),



EP 4 342 895 A1



<p>5 10</p>		<p>yl)piperidin-1- yl)phenyl)pyrimidin -4-amine</p>	<p>3.95 (s, 3H), 3.93-3.89 (m, 1H), 3.88 (s, 3H), 3.24 (dt, <math>J = 11.0, 3.5</math> Hz, 2H), 2.87-2.36 (m, 11H), 2.32 (s, 3H), 2.30-2.20 (m, 2H), 2.00-1.92 (m, 2H), 1.68-1.53 (m, 2H); 646.3 [M+H]<sup>+</sup></p>
<p>15 20 25 30</p>	<p>15 0</p> 	<p>(R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-5'-methoxyspiro[cyclopropane-1,3'-indoline]-6'-amine</p>	<p>452.1 [M+H]<sup>+</sup></p>
<p>35 40 45 50 55</p>	<p>15 1</p> 	<p>(R)-6'-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5'-methoxyspiro[cyclopropane-1,3'-indolin]-2'-one</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 10.35 (s, 1H), 8.68 (s, 1H), 8.27-8.20 (m, 1H), 7.57 (s, 1H), 7.13 (ddd, <math>J = 6.7, 5.3, 2.9</math> Hz, 2H), 6.81 (s, 1H), 6.54 (s, 1H), 5.55 (dd, <math>J = 8.7, 5.0</math> Hz, 1H), 4.16 (td, <math>J = 7.9, 3.9</math> Hz, 1H), 3.76 (s, 3H), 3.39 (dt, <math>J = 10.9, 6.4</math> Hz, 2H), 2.76 (ddd, <math>J = 12.2, 8.3, 4.4</math> Hz, 1H), 2.26 (ddt, <math>J = 11.8, 7.7, 4.0</math> Hz, 1H), 1.55 (q, <math>J = 3.6, 3.2</math> Hz, 2H), 1.42 (q, <math>J = 3.5</math> Hz, 2H);</p>

			466.3 [M+H] <sup>+</sup>
5		(R)-1-(6-((6-(3-	
10		(3,5-	
		difluorophenyl)isox	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.78 (s,
		azolidin-2-	1H), 8.56 (s, 1H), 8.21 (d, <i>J</i> = 1.0
		yl)pyrimidin-4-	Hz, 1H), 7.13 (dtd, <i>J</i> = 9.6, 5.2, 2.1
15		yl)amino)-5-	Hz, 3H), 6.53 (s, 1H), 4.18-4.13 (m,
2		methoxy-1'-	2H), 3.86 (s, 3H), 3.84-3.78 (m, 2H),
20		methylspiro[indolin	2.93 (s, 2H), 2.82-2.74 (m, 2H),
		-3,4'-piperidin]-1-	2.40-2.32 (m, 4H), 2.30-2.22 (m, 2H),
		yl)-2,2,2-	2.02 (d, <i>J</i> = 12.3 Hz, 3H), 1.71 (s,
25		trifluoroethan-1-	2H); 605.4 [M+H] <sup>+</sup>
30		one	
		(R)-6-((6-(3-(3,5-	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 10.58 (s,
35		difluorophenyl)isox	1H), 8.64 (s, 1H), 7.31 (s, 1H),
		azolidin-2-	7.16-7.08 (m, 3H), 6.69 (s, 1H), 6.38
		yl)pyrimidin-4-	(s, 1H), 5.55 (dd, <i>J</i> = 8.8, 5.1 Hz,
40		yl)amino)-7-	1H), 4.15 (td, <i>J</i> = 7.9, 3.9 Hz, 1H),
3		methoxyspiro[benzo[	3.83 (q, <i>J</i> = 7.9 Hz, 1H), 3.73 (s,
45		b][1,4]oxazin-2,1'-	3H), 2.78-2.71 (m, 2H), 2.26 (ddd, <i>J</i>
		cyclopropane]-	= 12.7, 8.4, 5.0 Hz, 1H), 1.25 (q, <i>J</i>
50		3(4H)-one	= 4.1, 3.2 Hz, 2H), 1.16-1.12 (m,
			2H); 482.3 [M+H] <sup>+</sup>





EP 4 342 895 A1



<p>5</p> <p>10</p> <p>15</p> <p>4</p> <p>15</p> <p>20</p>		<p>(R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-5-methoxy-1'-methylspiro[indolin-3,4'-piperidin]-6-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.48 (s, 1H), 8.18 (d, <i>J</i> = 1.0 Hz, 1H), 7.13 (td, <i>J</i> = 7.4, 2.9 Hz, 3H), 7.01 (s, 1H), 6.79 (s, 1H), 6.43 (s, 1H), 4.14 (td, <i>J</i> = 7.8, 3.8 Hz, 1H), 3.72 (s, 3H), 3.28 (s, 3H), 2.88-2.69 (m, 4H), 2.30 (s, 3H), 2.24 (ddd, <i>J</i> = 12.9, 8.4, 5.1 Hz, 2H), 2.17 (d, <i>J</i> = 6.9 Hz, 2H), 1.91-1.80 (m, 2H), 1.60 (d, <i>J</i> = 12.9 Hz, 2H); 509.3 [M+H]<sup>+</sup></p>
<p>25</p> <p>30</p> <p>35</p> <p>15</p> <p>5</p> <p>40</p> <p>45</p> <p>50</p>		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.82 (s, 1H), 8.16 (d, <i>J</i> = 1.0 Hz, 1H), 7.29 (d, <i>J</i> = 8.3 Hz, 1H), 7.11 (dd, <i>J</i> = 7.6, 2.9 Hz, 3H), 6.70 (d, <i>J</i> = 8.3 Hz, 1H), 6.28 (s, 1H), 4.55 (t, <i>J</i> = 9.0 Hz, 2H), 4.14 (td, <i>J</i> = 7.9, 3.8 Hz, 2H), 3.81 (q, <i>J</i> = 7.9 Hz, 2H), 3.22 (t, <i>J</i> = 8.7 Hz, 2H), 2.87 (dt, <i>J</i> = 11.7, 3.2 Hz, 2H), 2.76 (dt, <i>J</i> = 8.4, 4.5 Hz, 2H), 2.47-2.37 (m, 1H), 2.30-2.21 (m, 1H), 2.19 (s, 3H), 1.96 (td, <i>J</i> = 11.0, 4.4 Hz, 2H), 1.68 (td, <i>J</i> = 9.5, 3.1 Hz, 2H); 494.4 [M+H]<sup>+</sup></p>



<p>5</p> <p>15</p> <p>10</p> <p>6</p> <p>15</p>		<p>(R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-3-methoxy-7-(4-methylpiperazin-1-yl)-9H-carbazol-2-amine</p>	<p>572.3 [M+H]<sup>+</sup></p>
<p>20</p> <p>25</p> <p>30</p> <p>15</p> <p>7</p> <p>35</p> <p>40</p> <p>45</p>		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.82 (s, 1H), 8.17 (d, <i>J</i> = 0.9 Hz, 1H), 7.30 (d, <i>J</i> = 8.2 Hz, 1H), 7.12 (qd, <i>J</i> = 6.9, 3.0 Hz, 3H), 6.69 (d, <i>J</i> = 8.3 Hz, 1H), 6.28 (s, 1H), 4.55 (t, <i>J</i> = 9.1 Hz, 2H), 4.14 (td, <i>J</i> = 7.8, 3.8 Hz, 2H), 3.81 (q, <i>J</i> = 7.9 Hz, 2H), 3.22 (t, <i>J</i> = 8.7 Hz, 2H), 3.06 (d, <i>J</i> = 11.8 Hz, 2H), 2.76 (ddp, <i>J</i> = 12.1, 7.6, 3.8 Hz, 1H), 2.68–2.54 (m, 3H), 2.24 (dtd, <i>J</i> = 16.2, 8.2, 4.1 Hz, 1H), 1.68 (d, <i>J</i> = 12.3 Hz, 2H), 1.63–1.49 (m, 2H); 480.4 [M+H]<sup>+</sup></p>



50



55



5 10 15 8 20 25 30		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-methyl-1H-pyrazol-5-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine</p>	$^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$ ) $\delta$ 9.06 (s, 1H), 8.25 (d, $J = 0.9$ Hz, 1H), 7.73 (d, $J = 8.3$ Hz, 1H), 7.49 (d, $J = 1.9$ Hz, 1H), 7.13 (dtd, $J = 7.6, 4.2, 2.2$ Hz, 3H), 6.91 (d, $J = 8.3$ Hz, 1H), 6.53 (s, 1H), 6.39 (d, $J = 1.9$ Hz, 1H), 5.57 (dd, $J = 8.7, 5.0$ Hz, 1H), 4.61 (t, $J = 8.7$ Hz, 2H), 4.18 (td, $J = 7.9, 3.9$ Hz, 1H), 3.85 (q, $J = 7.9$ Hz, 1H), 3.79 (s, 3H), 3.21 (t, $J = 8.7$ Hz, 2H), 2.78 (dq, $J = 8.2, 4.2, 3.6$ Hz, 1H), 2.27 (dt, $J = 12.6, 3.9$ Hz, 1H); 477.3 [M+H] <sup>+</sup>
35 40 45 50 55		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine</p>	$^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$ ) $\delta$ 8.92 (s, 1H), 8.20 (d, $J = 0.9$ Hz, 1H), 7.41 (d, $J = 8.0$ Hz, 1H), 7.12 (dq, $J = 9.5, 3.2$ Hz, 3H), 7.03 (dd, $J = 7.4, 1.2$ Hz, 1H), 6.82 (t, $J = 7.6$ Hz, 1H), 6.35 (s, 1H), 5.56 (dd, $J = 8.7, 5.0$ Hz, 1H), 4.56 (t, $J = 8.8$ Hz, 2H), 4.15 (td, $J = 7.9, 3.8$ Hz, 1H), 3.86-3.76 (m, 1H), 3.24 (t, $J = 8.7$ Hz, 2H), 2.81-2.72 (m, 1H), 2.25 (ddt, $J = 11.8, 7.8, 4.0$ Hz, 1H); 397.2 [M+H] <sup>+</sup>

5 10 15 20 25		<p>(R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-(1-methylpiperidin-4-yl)-1H-pyrazol-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.92 (s, 1H), 8.22 (d, <i>J</i> = 1.0 Hz, 1H), 8.14 (s, 1H), 7.88 (s, 1H), 7.46 (d, <i>J</i> = 8.3 Hz, 1H), 7.18-7.04 (m, 4H), 6.37 (s, 1H), 5.57 (dd, <i>J</i> = 8.7, 5.0 Hz, 1H), 4.63 (t, <i>J</i> = 8.8 Hz, 2H), 4.48 (t, <i>J</i> = 7.5 Hz, 1H), 4.20-4.09 (m, 1H), 3.83 (d, <i>J</i> = 8.1 Hz, 1H), 3.48-3.36 (m, 4H), 3.11-3.00 (m, 2H), 2.76 (s, 2H), 2.26 (dt, <i>J</i> = 9.1, 4.7 Hz, 4H), 1.92 (s, 3H); 560.4 [M+H]<sup>+</sup></p>
30 35 40 45 50 55		<p>(R)-N-(2-(4-(4-cyclopropylpiperazin-1-yl)piperidin-1-yl)-5-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxyphenyl)propionamide</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.68 (s, 1H), 8.62 (s, 1H), 8.04 (s, 1H), 7.15-7.07 (m, 3H), 6.81 (s, 1H), 6.33 (s, 1H), 5.55 (dd, <i>J</i> = 8.7, 5.0 Hz, 1H), 4.13 (dt, <i>J</i> = 8.6, 4.2 Hz, 1H), 3.82 (q, <i>J</i> = 6.9, 5.7 Hz, 1H), 3.78 (s, 3H), 3.08 (s, 2H), 2.77 (ddd, <i>J</i> = 12.3, 8.2, 4.1 Hz, 2H), 2.70 (d, <i>J</i> = 13.7 Hz, 4H), 2.50 (d, <i>J</i> = 1.9 Hz, 6H), 2.38 (q, <i>J</i> = 7.6 Hz, 2H), 2.30-2.17 (m, 2H), 1.99 (q, <i>J</i> = 6.0, 5.5 Hz, 1H), 1.89 (d, <i>J</i> = 17.8 Hz, 2H), 1.70 (s, 2H), 1.10 (t, <i>J</i> = 7.5 Hz,</p>

5			3H), 0.45 (s, 2H), 0.32 (s, 2H); 663.4 [M+H] <sup>+</sup>
10 15 16 20 25 30		(R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-methylpiperazin-1-yl)-2,3-dihydrobenzofuran-7-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.80 (s, 1H), 8.17 (s, 1H), 7.97 (d, <i>J</i> = 6.5 Hz, 1H), 7.73 (s, 1H), 7.40 (d, <i>J</i> = 7.1 Hz, 3H), 7.35 (t, <i>J</i> = 7.5 Hz, 3H), 7.29–7.22 (m, 1H), 6.31 (s, 1H), 5.53 (dd, <i>J</i> = 8.6, 5.0 Hz, 1H), 4.56 (t, <i>J</i> = 8.9 Hz, 2H), 4.15 (td, <i>J</i> = 7.8, 3.9 Hz, 1H), 3.88 (s, 3H), 3.82 (q, <i>J</i> = 7.9 Hz, 1H), 3.40 (t, <i>J</i> = 8.6 Hz, 2H), 3.10 (d, <i>J</i> = 11.1 Hz, 7H), 2.74 (s, 4H), 2.24 (dtd, <i>J</i> = 12.8, 8.1, 5.0 Hz, 1H); 539.4 [M+H] <sup>+</sup>
35 40 45 50 55		(R)-N-(2-(4-(4-cyclopropylpiperazin-1-yl)piperidin-1-yl)-4-methoxy-5-((6-(3-(thiophene)-2-yl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)propionamide	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.66 (s, 1H), 8.61 (s, 1H), 8.04 (s, 1H), 7.40 (dd, <i>J</i> = 5.1, 1.3 Hz, 1H), 7.08 (dt, <i>J</i> = 3.5, 1.2 Hz, 1H), 6.98 (dd, <i>J</i> = 5.1, 3.5 Hz, 1H), 6.81 (s, 1H), 6.28 (s, 1H), 5.81 (dd, <i>J</i> = 8.3, 3.9 Hz, 1H), 4.14 (td, <i>J</i> = 7.9, 4.7 Hz, 1H), 3.86 (q, <i>J</i> = 7.9 Hz, 1H), 3.77 (s, 3H), 3.04 (s, 2H), 2.65 (dd, <i>J</i> = 12.2, 3.6 Hz, 4H), 2.53 (d, <i>J</i> = 6.6 Hz, 8H),

5			2.43–2.32 (m, 4H), 1.85 (s, 2H), 1.67 (s, 2H), 1.58 (s, 1H), 1.10 (t, $J = 7.6$ Hz, 3H), 0.40 (s, 2H), 0.28 (s, 2H); 633.4 [M+H] <sup>+</sup>
16 4		(R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.58 (s, 1H), 8.08 (s, 1H), 7.87 (s, 1H), 7.48 (s, 1H), 7.43–7.37 (m, 2H), 7.34 (dd, $J = 8.5, 6.7$ Hz, 2H), 7.28–7.23 (m, 1H), 6.90 (s, 1H), 6.23 (s, 1H), 5.53 (dd, $J = 8.6, 4.9$ Hz, 1H), 4.74 (q, $J = 8.9$ Hz, 2H), 4.12 (td, $J = 7.9, 4.1$ Hz, 1H), 3.87 (s, 3H), 3.80 (q, $J = 7.9$ Hz, 1H), 3.12 (d, $J = 11.2$ Hz, 2H), 2.73 (qt, $J = 7.9, 4.1$ Hz, 2H), 2.64–2.50 (m, 9H), 2.24 (hept, $J = 7.9$ Hz, 5H), 1.84 (dd, $J = 13.2, 7.5$ Hz, 2H), 1.61–1.43 (m, 3H); 678.5 [M+H] <sup>+</sup>
16 5		(R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-morpholinopiperidin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.59 (s, 1H), 8.08 (s, 1H), 7.87 (s, 1H), 7.48 (s, 1H), 7.44–7.38 (m, 2H), 7.34 (dd, $J = 8.5, 6.8$ Hz, 2H), 7.30–7.23 (m, 1H), 6.91 (s, 1H), 6.24 (s, 1H), 5.53 (dd, $J = 8.6, 4.9$ Hz, 1H), 4.75 (q,


5 10 15		nyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	$J = 9.0 \text{ Hz, 2H}$ ), 4.11 (tt, $J = 9.1, 4.6 \text{ Hz, 1H}$ ), 3.87 (s, 3H), 3.80 (q, $J = 7.9 \text{ Hz, 1H}$ ), 3.60 (s, 3H), 3.12 (s, 2H), 2.73 (dtd, $J = 12.1, 7.9, 4.1 \text{ Hz, 1H}$ ), 2.57 (t, $J = 12.6 \text{ Hz, 8H}$ ), 2.29-2.13 (m, 2H), 1.85 (s, 2H), 1.53 (s, 2H); 665.4 [M+H] <sup>+</sup>
20 25 30 35 40	16 6 	(R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-methylpiperazin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.66 (s, 1H), 8.15 (t, $J = 1.1 \text{ Hz, 2H}$ ), 7.92 (s, 1H), 7.56 (s, 1H), 7.42-7.38 (m, 2H), 7.37-7.32 (m, 2H), 7.28-7.22 (m, 1H), 6.92 (s, 1H), 6.29 (s, 1H), 5.54 (dd, $J = 8.6, 4.9 \text{ Hz, 1H}$ ), 4.79 (q, $J = 9.0 \text{ Hz, 2H}$ ), 4.13 (d, $J = 4.1 \text{ Hz, 1H}$ ), 3.89 (s, 3H), 3.81 (d, $J = 8.0 \text{ Hz, 1H}$ ), 3.26 (s, 4H), 3.05 (s, 4H), 2.81 (s, 3H), 2.73 (dd, $J = 8.2, 4.1 \text{ Hz, 1H}$ ), 2.25 (dt, $J = 8.6, 4.7 \text{ Hz, 1H}$ ); 595.4 [M+H] <sup>+</sup>
45 50 55	16 7 	(R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-morpholino-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 9.36 (s, 1H), 8.22 (s, 1H), 8.16 (s, 1H), 7.46 (s, 1H), 7.38-7.33 (m, 4H), 7.27 (tt, $J = 5.3, 2.4 \text{ Hz, 1H}$ ), 6.96 (s, 1H), 6.12 (s, 1H), 5.54 (dd, $J = 8.5, 5.1$



5  10  15		nyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine	Hz, 1H), 4.79 (dt, $J = 8.7, 4.4$ Hz, 2H), 4.24-4.18 (m, 1H), 3.93 (d, $J = 7.8$ Hz, 1H), 3.88 (s, 3H), 3.74 (t, $J = 4.5$ Hz, 4H), 2.89-2.84 (m, 4H), 2.82 (t, $J = 4.6$ Hz, 1H), 2.28 (dt, $J = 8.0, 4.8$ Hz, 1H); 582.4 [M+H] <sup>+</sup>
16  20  25  30  35  40		(S)-6-(3-(5-fluoropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.50 (s, 1H), 8.37 (s, 1H), 8.29 (s, 1H), 7.55 (d, $J = 9.4$ Hz, 1H), 7.49 (d, $J = 8.5$ Hz, 1H), 6.96 (s, 1H), 6.58-6.51 (m, 2H), 6.41 (s, 1H), 5.76 (dd, $J = 8.8, 4.6$ Hz, 1H), 4.12 (q, $J = 6.9$ Hz, 1H), 3.94-3.86 (m, 1H), 3.84 (s, 3H), 3.72 (d, $J = 12.0$ Hz, 2H), 3.21 (s, 3H), 2.75 (t, $J = 12.1$ Hz, 4H), 2.68-2.61 (m, 3H), 2.57-2.42 (m, 4H), 2.39-2.34 (m, 2H), 1.96 (d, $J = 12.4$ Hz, 2H), 1.70 (q, $J = 13.0, 12.5$ Hz, 2H); 549.3 [M+H] <sup>+</sup>
45  50  55		(R)-6-(3-(5-fluoropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.50 (d, $J = 1.8$ Hz, 1H), 8.37 (d, $J = 2.8$ Hz, 1H), 8.29 (s, 1H), 7.55 (dt, $J = 9.4, 2.3$ Hz, 1H), 7.50 (d, $J = 8.8$ Hz, 1H), 6.85 (s, 1H), 6.58-6.51 (m,



EP 4 342 895 A1




5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55



		<p>methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>2H), 6.44-6.39 (m, 1H), 5.76 (dd, <math>J = 8.8, 4.6</math> Hz, 1H), 4.14-4.10 (m, 1H), 3.93-3.86 (m, 1H), 3.84 (s, 3H), 3.72 (dt, <math>J = 12.1, 3.4</math> Hz, 2H), 2.84-2.34 (m, 13H), 2.31 (s, 3H), 1.96 (d, <math>J = 11.6</math> Hz, 2H), 1.70 (qd, <math>J = 12.1, 3.9</math> Hz, 2H); 549.3 [M+H]<sup>+</sup></p>
<p>17 0</p>		<p>(R)-3-fluoro-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzotrile</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) <math>\delta</math> 8.29 (s, 1H), 7.56 (s, 1H), 7.50 (d, <math>J = 9.1</math> Hz, 1H), 7.44 (dt, <math>J = 9.4, 2.1</math> Hz, 1H), 7.26-7.23 (m, 1H), 6.79 (s, 1H), 6.58-6.51 (m, 2H), 6.41 (d, <math>J = 0.9</math> Hz, 1H), 5.70 (dd, <math>J = 8.8, 4.8</math> Hz, 1H), 4.12 (td, <math>J = 8.0, 3.9</math> Hz, 1H), 3.90-3.85 (m, 4H), 3.73 (dd, <math>J = 9.5, 6.3</math> Hz, 2H), 2.84-2.33 (m, 12H), 2.31 (s, 3H), 2.30-2.23 (m, 1H), 1.97 (d, <math>J = 11.5</math> Hz, 2H), 1.70 (qd, <math>J = 12.0, 3.8</math> Hz, 2H); 573.3 [M+H]<sup>+</sup></p>



5 10 15 20 25		<p>(S)-3-fluoro-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzonitrile</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.29 (s, 1H), 7.56 (s, 1H), 7.51 (d, <i>J</i> = 9.1 Hz, 1H), 7.44 (dt, <i>J</i> = 9.4, 2.0 Hz, 1H), 7.26-7.23 (m, 1H), 6.77 (s, 1H), 6.58-6.51 (m, 2H), 6.41 (s, 1H), 5.70 (dd, <i>J</i> = 8.8, 4.8 Hz, 1H), 4.12 (td, <i>J</i> = 8.1, 3.9 Hz, 1H), 3.92-3.85 (m, 4H), 3.72 (d, <i>J</i> = 12.5 Hz, 2H), 2.84-2.32 (m, 13H), 2.31 (s, 3H), 1.96 (d, <i>J</i> = 11.0 Hz, 2H), 1.70 (q, <i>J</i> = 12.2, 4.1 Hz, 2H); 573.3 [M+H]<sup>+</sup></p>
30 35 40 45		<p>tert-butyl (R)-3-(4-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.32 (d, <i>J</i> = 1.0 Hz, 1H), 7.40-7.36 (m, 2H), 7.34-7.27 (m, 3H), 7.10-7.03 (m, 2H), 6.73 (s, 1H), 6.55 (d, <i>J</i> = 1.1 Hz, 1H), 5.91 (dd, <i>J</i> = 8.7, 4.8 Hz, 1H), 5.22 (dd, <i>J</i> = 8.7, 5.5 Hz, 1H), 4.20 (q, <i>J</i> = 8.1, 3.8 Hz, 1H), 4.12 (td, <i>J</i> = 8.0, 4.2 Hz, 1H), 3.94-3.84 (m, 2H), 2.88-2.73 (m, 2H), 2.37-2.26 (m, 2H), 1.48 (s, 9H); 526.25 [M+H]<sup>+</sup></p>



50



55



5 10 15 20	17 3 	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((R)-2-methylisoxazolidin-3-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.32 (d, <i>J</i> = 1.1 Hz, 1H), 7.43-7.37 (m, 2H), 7.34-7.28 (m, 3H), 7.11-7.02 (m, 2H), 6.85 (s, 1H), 6.56 (d, <i>J</i> = 1.0 Hz, 1H), 5.91 (dd, <i>J</i> = 8.7, 4.8 Hz, 1H), 4.16-4.06 (m, 3H), 3.89 (q, <i>J</i> = 8.0 Hz, 1H), 3.55 (s, 1H), 2.89-2.79 (m, 1H), 2.76-2.67 (m, 1H), 2.64 (s, 3H), 2.37-2.26 (m, 2H); 440.16 [M+H] <sup>+</sup>
25 30 35 40	17 4 	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((R)-isoxazolidin-3-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (d, <i>J</i> = 1.1 Hz, 1H), 7.43-7.39 (m, 2H), 7.33-7.29 (m, 3H), 7.10-7.01 (m, 4H), 6.55 (d, <i>J</i> = 1.1 Hz, 1H), 5.90 (dd, <i>J</i> = 8.8, 4.8 Hz, 1H), 4.50 (t, <i>J</i> = 10.0 Hz, 1H), 4.16-4.09 (m, 2H), 3.88 (q, <i>J</i> = 8.2 Hz, 1H), 3.35 (t, <i>J</i> = 10.1 Hz, 1H), 2.88-2.80 (m, 1H), 2.74-2.64 (m, 1H), 2.36-2.27 (m, 2H); 426.22 [M+H] <sup>+</sup>
45 50 55	17 5 	tert-butyl ((S)-3-(4-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)phenyl)pyrimidin-4-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.31 (d, <i>J</i> = 1.1 Hz, 1H), 7.40-7.27 (m, 5H), 7.09-7.03 (m, 2H), 6.78 (s, 1H), 6.55 (d, <i>J</i> = 1.1 Hz, 1H), 5.91 (dd, <i>J</i> = 8.9, 4.8 Hz, 1H), 5.22 (dd, <i>J</i> =

5  10		yl)amino)phenyl)iso xazolidin-2- carboxylate	8.7, 5.4 Hz, 1H), 4.23-4.16 (m, 1H), 4.12 (td, $J = 7.8, 5.6$ Hz, 1H), 3.94- 3.84 (m, 2H), 2.88-2.75 (m, 2H), 2.37-2.26 (m, 2H), 1.48 (s, 9H); 526.20 [M+H] <sup>+</sup>
15  20 25 30	17 6 	6-((R)-3-(2,3- difluorophenyl)isox azolidin-2-yl)-N- (4-((S)-2- methylisoxazolidin- 3- yl)phenyl)pyrimidin -4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.31 (d, $J = 1.0$ Hz, 1H), 7.42-7.36 (m, 2H), 7.35-7.27 (m, 3H), 7.12-6.99 (m, 2H), 6.94 (s, 1H), 6.57 (d, $J = 1.1$ Hz, 1H), 5.91 (dd, $J = 8.8, 4.8$ Hz, 1H), 4.17-4.04 (m, 3H), 3.89 (q, $J =$ 8.0 Hz, 1H), 3.56 (s, 1H), 2.84 (dtd, $J = 12.3, 8.1, 4.1$ Hz, 1H), 2.77-2.66 (m, 1H), 2.64 (s, 3H), 2.37-2.26 (m, 2H); 440.16 [M+H] <sup>+</sup>
35  40 45 50 55	17 7 	6-((R)-3-(2,3- difluorophenyl)isox azolidin-2-yl)-N- (4-((S)- isoxazolidin-3- yl)phenyl)pyrimidin -4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.31 (d, $J = 1.1$ Hz, 1H), 7.44-7.39 (m, 2H), 7.34-7.29 (m, 3H), 7.13-7.02 (m, 3H), 6.87 (s, 1H), 6.55 (d, $J = 1.1$ Hz, 1H), 5.90 (dd, $J = 8.8, 4.9$ Hz, 1H), 4.50 (t, $J = 10.1$ Hz, 1H), 4.17- 4.09 (m, 2H), 3.88 (q, $J = 8.2$ Hz, 1H), 3.35 (t, $J = 10.1$ Hz, 1H), 2.88- 2.79 (m, 1H), 2.69 (ddd, $J = 14.1,$

5			12.3, 8.1 Hz, 1H), 2.33 (ddd, $J = 12.7, 8.4, 5.4$ Hz, 2H); 426.22 [M+H] <sup>+</sup>
10			<sup>1</sup> H NMR (400 MHz, MeOD) $\delta$ 8.18 (s, 1H), 7.61 (s, 1H), 7.49 (s, 1H), 7.34 (d, $J = 8.9$ Hz, 2H), 7.09-6.97 (m, 2H), 6.34 (d, $J = 0.6$ Hz, 1H), 5.54 (dd, $J = 8.1, 3.3$ Hz, 1H), 4.14 (tt, $J = 8.8, 4.4$ Hz, 1H), 4.05-3.95 (m, 1H), 3.86 (d, $J = 5.6$ Hz, 3H), 3.35 (dd, $J = 8.5, 3.7$ Hz, 4H), 3.13-3.03 (m, 4H), 2.69 (s, 3H), 2.60 (dtd, $J = 12.2, 8.4, 5.8$ Hz, 1H), 2.38 (tdd, $J = 7.9, 6.7, 3.4$ Hz, 1H); 421.19 [M+H] <sup>+</sup>
15		(R)-6-(3-(1-methyl-1H-pyrazol-4-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	
17	8		
20			
25			
30			
35		(R)-6-(3-(furan-2-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	
40	9		
45			
50			<sup>1</sup> H NMR (400 MHz, MeOD) $\delta$ 8.19 (s, 1H), 7.47 (dd, $J = 1.6, 0.7$ Hz, 1H), 7.33 (d, $J = 8.8$ Hz, 2H), 7.03 (d, $J = 8.9$ Hz, 2H), 6.42-6.31 (m, 3H), 5.64 (t, $J = 6.1$ Hz, 1H), 4.19 (dt, $J = 13.6, 6.7$ Hz, 1H), 3.96 (q, $J = 7.8$ Hz, 1H), 3.32 (d, $J = 1.6$ Hz, 4H), 3.08-2.99 (m, 4H), 2.65 (s, 3H), 2.57 (dt, $J = 12.9, 6.5$ Hz, 2H); 407.20 [M+H] <sup>+</sup>

5 10 15 20 25		(R)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 9.20 (s, 1H), 8.60 (d, <i>J</i> = 1.9 Hz, 1H), 8.55 (d, <i>J</i> = 2.4 Hz, 1H), 8.22 (d, <i>J</i> = 0.9 Hz, 1H), 7.91–7.90 (m, 1H), 7.38 (d, <i>J</i> = 8.5 Hz, 2H), 6.94–6.89 (m, 2H), 6.36 (d, <i>J</i> = 1.1 Hz, 1H), 3.87 (q, <i>J</i> = 7.9 Hz, 1H), 3.11 (t, <i>J</i> = 5.1 Hz, 4H), 2.79 (dtt, <i>J</i> = 12.2, 7.8, 4.0 Hz, 1H), 2.53 (t, <i>J</i> = 5.5 Hz, 4H), 2.33 (ddd, <i>J</i> = 12.6, 8.5, 4.8 Hz, 2H), 2.27 (s, 2H), 1.91 (s, 3H); 452.1 [M+H] <sup>+</sup>
30 35 40 45 50		(S)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.60 (d, <i>J</i> = 1.9 Hz, 1H), 8.55 (d, <i>J</i> = 2.4 Hz, 1H), 8.22 (s, 1H), 7.90 (t, <i>J</i> = 2.2 Hz, 1H), 7.37 (d, <i>J</i> = 8.5 Hz, 2H), 6.93–6.89 (m, 2H), 6.36 (s, 1H), 5.63–5.59 (m, 1H), 3.87 (q, <i>J</i> = 8.0 Hz, 2H), 3.08 (t, <i>J</i> = 5.0 Hz, 4H), 2.78 (dtd, <i>J</i> = 12.2, 7.9, 3.9 Hz, 2H), 2.45 (t, <i>J</i> = 5.0 Hz, 4H), 2.37–2.28 (m, 2H), 2.22 (s, 3H), 1.90 (s, 2H); 452.1 [M+H] <sup>+</sup>



5 10 15 18 2 20 25 30		<p>(R)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.59 (d, <i>J</i> = 1.9 Hz, 1H), 8.54 (d, <i>J</i> = 2.3 Hz, 1H), 8.47 (s, 1H), 8.13 (d, <i>J</i> = 1.0 Hz, 1H), 7.89–7.88 (m, 1H), 7.30 (d, <i>J</i> = 8.5 Hz, 1H), 6.62 (d, <i>J</i> = 2.5 Hz, 1H), 6.49 (dd, <i>J</i> = 8.8, 2.5 Hz, 1H), 5.59 (dd, <i>J</i> = 8.7, 5.0 Hz, 1H), 4.15 (td, <i>J</i> = 7.8, 3.9 Hz, 1H), 3.82 (t, <i>J</i> = 8.0 Hz, 1H), 3.77 (s, 3H), 3.72 (d, <i>J</i> = 12.4 Hz, 2H), 2.77 (qt, <i>J</i> = 7.9, 3.9 Hz, 2H), 2.67 (td, <i>J</i> = 12.2, 2.4 Hz, 3H), 2.53 (s, 2H), 2.35–2.25 (m, 4H), 2.17 (s, 3H), 1.91 (s, 5H), 1.85 (d, <i>J</i> = 12.5 Hz, 2H), 1.51 (qd, <i>J</i> = 12.1, 3.9 Hz, 3H); 565.2 [M+H]<sup>+</sup></p>
35 40 45 3 50 55		<p>(S)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>565.3 [M+H]<sup>+</sup></p>



5 10 15 18 4 20 25 30		(R)-6-(3-(3-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-2-yl)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-2-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, MeOD) δ 8.11 (d, <i>J</i> = 0.6 Hz, 1H), 7.66–7.55 (m, 2H), 7.50–7.43 (m, 2H), 7.32 (d, <i>J</i> = 8.6 Hz, 1H), 6.77 (d, <i>J</i> = 3.1 Hz, 1H), 6.70 (d, <i>J</i> = 2.4 Hz, 1H), 6.63–6.58 (m, 1H), 6.24 (s, 1H), 5.58 (dd, <i>J</i> = 8.5, 4.8 Hz, 1H), 4.13 (tt, <i>J</i> = 10.5, 5.3 Hz, 1H), 3.97–3.88 (m, 1H), 3.83 (d, <i>J</i> = 6.4 Hz, 3H), 3.80 (d, <i>J</i> = 12.7 Hz, 2H), 2.98 (d, <i>J</i> = 22.3 Hz, 7H), 2.87–2.71 (m, 4H), 2.68–2.65 (m, 4H), 2.39–2.29 (m, 1H), 2.05 (d, <i>J</i> = 11.9 Hz, 2H), 1.78–1.65 (m, 2H); 580.25 [M+H] <sup>+</sup>
35 40 45 50 55		(S)-6-(3-(3-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-2-yl)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-2-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, MeOD) δ 8.11 (s, 1H), 7.65–7.56 (m, 2H), 7.51–7.43 (m, 2H), 7.32 (d, <i>J</i> = 8.6 Hz, 1H), 6.77 (d, <i>J</i> = 3.2 Hz, 1H), 6.70 (d, <i>J</i> = 2.5 Hz, 1H), 6.66–6.57 (m, 1H), 6.24 (s, 1H), 5.58 (dd, <i>J</i> = 8.5, 4.8 Hz, 1H), 4.13 (td, <i>J</i> = 7.8, 4.3 Hz, 1H), 3.91 (q, <i>J</i> = 7.9 Hz, 1H), 3.85 (d, <i>J</i> = 6.4 Hz, 3H), 3.80 (d, <i>J</i> = 12.6 Hz, 2H), 2.99 (d, <i>J</i> = 29.1 Hz, 7H), 2.79 (dq, <i>J</i> =



5			10.1, 7.5, 4.4 Hz, 4H), 2.68–2.63 (m, 4H), 2.40–2.26 (m, 1H), 2.04 (d, $J = 12.2$ Hz, 2H), 1.73 (tt, $J = 11.9, 6.1$ Hz, 2H); 580.30 [M+H] <sup>+</sup>
10 15 20 25 30 35	6 18 6	tert-butyl (R)-3-(3-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (d, $J = 1.1$ Hz, 1H), 7.35 (t, $J = 7.8$ Hz, 1H), 7.33–7.29 (m, 2H), 7.26–7.20 (m, 2H), 7.16–7.12 (m, 1H), 7.09–7.03 (m, 2H), 6.59 (d, $J = 1.0$ Hz, 1H), 5.90 (dd, $J = 8.8, 4.8$ Hz, 1H), 5.23 (dd, $J = 8.7, 5.4$ Hz, 1H), 4.19 (td, $J = 7.9, 3.7$ Hz, 1H), 4.12 (td, $J = 8.0, 4.3$ Hz, 1H), 3.89 (qd, $J = 8.1, 7.7, 1.9$ Hz, 2H), 2.89–2.74 (m, 2H), 2.37–2.25 (m, 2H), 1.48 (s, 9H); 526.25 [M+H] <sup>+</sup>
40 45 50 55	7 18 7	tert-butyl (S)-3-(3-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.29 (d, $J = 1.0$ Hz, 1H), 7.36 (t, $J = 7.8$ Hz, 1H), 7.32 (p, $J = 2.7$ Hz, 2H), 7.26–7.22 (m, 2H), 7.16–7.12 (m, 1H), 7.06 (qt, $J = 8.4, 5.8$ Hz, 2H), 6.60 (d, $J = 1.0$ Hz, 1H), 5.91 (dd, $J = 8.8, 4.8$ Hz, 1H), 5.24 (dd, $J = 8.7, 5.3$ Hz, 1H), 4.18 (dt, $J = 7.9, 4.0$




## EP 4 342 895 A1


			Hz, 1H), 4.12 (td, $J = 8.0, 4.2$ Hz, 1H), 3.95–3.85 (m, 2H), 2.89–2.75 (m, 2H), 2.38–2.26 (m, 2H), 1.47 (s, 9H); 526.20 [M+H] <sup>+</sup>
5 10 15 20 25 30	18 8 	(R)-N-(6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)cyclopropanecarboxamide	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.38 (d, $J = 1.1$ Hz, 1H), 8.26 (s, 1H), 7.91 (d, $J = 1.1$ Hz, 1H), 7.31–7.27 (m, 1H), 7.12–6.98 (m, 2H), 5.86 (dd, $J = 8.8, 5.1$ Hz, 1H), 4.18 (td, $J = 7.9, 3.9$ Hz, 1H), 3.93 (td, $J = 8.4, 7.3$ Hz, 1H), 2.92–2.83 (m, 1H), 2.38–2.28 (m, 1H), 1.55 (dt, $J = 7.8, 4.7$ Hz, 1H), 1.18–1.10 (m, 2H), 0.97–0.90 (m, 2H); 347.19 [M+H] <sup>+</sup>
35 40 45 50 55	18 9 	(R)-N-(6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)benzamide	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.51 (s, 1H), 8.42 (d, $J = 1.1$ Hz, 1H), 8.12 (d, $J = 1.1$ Hz, 1H), 7.95–7.90 (m, 2H), 7.64–7.59 (m, 1H), 7.56–7.50 (m, 2H), 7.34–7.29 (m, 1H), 7.13–7.03 (m, 2H), 5.91 (dd, $J = 8.8, 5.1$ Hz, 1H), 4.24 (td, $J = 8.0, 4.0$ Hz, 1H), 3.99 (td, $J = 8.4, 7.4$ Hz, 1H), 2.96–2.85 (m, 1H), 2.43–2.32 (m, 1H); 383.20 [M+H] <sup>+</sup>

5 10 15 20 25		(R)-N-(cyclopentylmethyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.19 (s, 1H), 7.34 (qd, <i>J</i> = 4.8, 1.9 Hz, 1H), 7.12–6.97 (m, 2H), 6.13 (d, <i>J</i> = 1.0 Hz, 1H), 5.92 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 5.00–4.84 (m, 1H), 4.14 (ddd, <i>J</i> = 8.3, 7.5, 4.3 Hz, 1H), 3.93 (q, <i>J</i> = 8.0 Hz, 1H), 3.19 (s, 2H), 2.88–2.79 (m, 1H), 2.32 (tdd, <i>J</i> = 12.5, 6.5, 3.0 Hz, 1H), 2.16 (hept, <i>J</i> = 7.6 Hz, 1H), 1.89–1.77 (m, 2H), 1.71–1.59 (m, 3H), 1.32–1.20 (m, 3H); 361.22 [M+H] <sup>+</sup>
30 35 40 45		(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-fluorobenzyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.24 (d, <i>J</i> = 1.0 Hz, 1H), 7.35–7.28 (m, 3H), 7.08–7.00 (m, 4H), 6.14 (d, <i>J</i> = 1.1 Hz, 1H), 5.89 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 5.21 (s, 1H), 4.50 (d, <i>J</i> = 5.9 Hz, 2H), 4.10 (tt, <i>J</i> = 7.8, 3.5 Hz, 1H), 3.84 (q, <i>J</i> = 7.9 Hz, 1H), 2.87–2.76 (m, 1H), 2.36–2.24 (m, 1H); 387.17 [M+H] <sup>+</sup>

50



55



5 10 15 20	<p>19 2</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(1-(methylsulfonyl)piperidin-4-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.22 (s, 1H), 7.36-7.27 (m, 1H), 7.12-6.99 (m, 3H), 6.13 (d, <i>J</i> = 1.0 Hz, 1H), 5.89 (dd, <i>J</i> = 8.8, 4.8 Hz, 1H), 4.14 (td, <i>J</i> = 7.9, 4.4 Hz, 1H), 3.92 (q, <i>J</i> = 7.9 Hz, 1H), 3.77 (d, <i>J</i> = 12.8 Hz, 2H), 2.94 (t, <i>J</i> = 12.0 Hz, 2H), 2.88-2.78 (m, 5H), 2.37-2.27 (m, 1H), 2.21-2.11 (m, 2H), 1.68-1.61 (m, 2H); 440.21 [M+H]<sup>+</sup></p>
25 30 35 40 45	<p>19 3</p> 	<p>(R)-6-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-methoxy-2-methylisoindolin-1-one</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.38 (d, <i>J</i> = 1.0 Hz, 1H), 8.37 (s, 1H), 7.35-7.27 (m, 2H), 7.12-7.02 (m, 2H), 6.97 (s, 1H), 6.58 (d, <i>J</i> = 1.0 Hz, 1H), 5.92 (dd, <i>J</i> = 8.9, 4.8 Hz, 1H), 4.33 (s, 2H), 4.13 (ddd, <i>J</i> = 8.3, 7.5, 4.2 Hz, 1H), 3.96 (s, 3H), 3.92 (q, <i>J</i> = 8.0 Hz, 1H), 3.20 (s, 3H), 2.89-2.79 (m, 1H), 2.37-2.26 (m, 1H); 454.13 [M+H]<sup>+</sup></p>
50 55	<p>19 4</p> 	<p>(R)-3-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-cyano-1H-indazole</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 9.26 (t, <i>J</i> = 1.1 Hz, 1H), 8.59 (d, <i>J</i> = 1.0 Hz, 1H), 7.65 (dd, <i>J</i> = 8.2, 0.8 Hz, 1H), 7.49 (d, <i>J</i> = 1.1 Hz, 1H), 7.46</p>



5  10  15		yl)amino)-1H-indazol-6-carbonitrile	(dd, $J = 8.2, 1.3$ Hz, 1H), 7.33 (td, $J = 7.0, 6.2, 3.7$ Hz, 1H), 7.15-6.98 (m, 2H), 5.95 (dd, $J = 8.8, 5.1$ Hz, 1H), 4.48 (s, 2H), 4.25 (td, $J = 7.9, 4.0$ Hz, 1H), 4.00 (q, $J = 8.2$ Hz, 1H), 2.97-2.87 (m, 1H), 2.45-2.33 (m, 1H); 420.15 [M+H] <sup>+</sup>
19  20  25  30  35  40  45		(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(9-(1-fluoro-2-methylpropan-2-yl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) $\delta$ 8.60 (s, 1H), 8.15 (d, $J = 1.0$ Hz, 1H), 8.05 (s, 1H), 7.85 (s, 1H), 7.57 (s, 1H), 7.35 (q, $J = 8.7$ Hz, 2H), 7.27 (q, $J = 6.8, 6.2$ Hz, 2H), 7.20 (dd, $J = 8.3, 5.2$ Hz, 1H), 6.84 (s, 1H), 6.28 (s, 1H), 4.16 (td, $J = 8.0, 3.8$ Hz, 1H), 3.94 (q, $J = 8.0$ Hz, 1H), 3.85 (s, 3H), 3.81 (s, 3H), 2.80 (t, $J = 5.7$ Hz, 4H), 2.45 (d, $J = 6.5$ Hz, 4H), 2.38 (s, 1H), 2.21 (tt, $J = 8.7, 4.3$ Hz, 2H), 1.91 (s, 2H), 1.53 (s, 8H), 1.32 (s, 3H), 1.27 (s, 3H); 691.3 [M+H] <sup>+</sup>

50



55




<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p>	<p>19</p> <p>6</p> 	<p>(R)-N-(2,5-dichloro-4-(9-(1-fluoro-2-methylpropan-2-yl)-3,9-diazaspiro[5.5]undecan-3-yl)phenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine</p>	<p>649.3 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p>	<p>19</p> <p>7</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-5-(3-((methylsulfonyl)methyl)azetidino-1-yl)-4-(4-morpholinopiperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p>684.3 [M+H]<sup>+</sup></p>



5 10 15 20 25		<p>(R)-N-(5-(3-chloro-1-methyl-1H-pyrazol-4-yl)-2-methoxy-4-(4-methylpiperazin-1-yl)phenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine</p>	597.3 [M+H] <sup>+</sup>
30 35 40 45 50		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-methylpiperazin-1-yl)-5-vinylphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.62 (s, 1H), 8.17 (d, <i>J</i> = 1.0 Hz, 1H), 7.76 (s, 1H), 7.39–7.30 (m, 2H), 6.90 (dd, <i>J</i> = 17.8, 11.0 Hz, 1H), 6.73 (s, 1H), 6.32 (s, 1H), 5.75 (dd, <i>J</i> = 8.9, 5.1 Hz, 1H), 5.57 (dd, <i>J</i> = 17.7, 1.6 Hz, 1H), 5.15 (dd, <i>J</i> = 10.9, 1.5 Hz, 1H), 4.17 (td, <i>J</i> = 8.0, 3.8 Hz, 2H), 3.82 (s, 3H), 2.94 (s, 4H), 2.86–2.76 (m, 2H), 2.59 (s, 2H), 2.32 (s, 3H), 2.26–2.16 (m, 3H), 1.91 (s, 2H);</p> <p>509.3 [M+H]<sup>+</sup></p>

5 10 15 20 25		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethynyl-2-methoxy-4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.64 (d, <i>J</i> = 1.2 Hz, 1H), 7.38 (dtd, <i>J</i> = 10.3, 8.0, 1.9 Hz, 2H), 7.28 (t, <i>J</i> = 6.9 Hz, 1H), 7.23 (dd, <i>J</i> = 8.1, 5.1 Hz, 1H), 7.14 (d, <i>J</i> = 1.2 Hz, 1H), 6.80 (s, 1H), 6.56 (s, 1H), 5.74 (dd, <i>J</i> = 8.8, 5.8 Hz, 1H), 4.31 (td, <i>J</i> = 7.8, 3.2 Hz, 1H), 3.94 (q, <i>J</i> = 8.0 Hz, 1H), 3.83 (s, 3H), 3.06 (s, 4H), 2.98–2.89 (m, 2H), 2.53 (d, <i>J</i> = 5.5 Hz, 2H), 2.25 (s, 3H), 1.91 (s, 4H); 507.3 [M+H]<sup>+</sup></p>
30 35 40 45 50		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-6-(1-methyl-1H-pyrazol-4-yl)-5-(4-methylpiperazin-1-yl)pyridin-2-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, DMSO-<i>d</i><sub>6</sub>) δ 8.31 (t, <i>J</i> = 1.3 Hz, 2H), 8.29 (d, <i>J</i> = 1.1 Hz, 1H), 8.22 (s, 1H), 8.17 (s, 1H), 7.41–7.35 (m, 1H), 7.34–7.31 (m, 1H), 7.25–7.19 (m, 1H), 5.80 (dd, <i>J</i> = 8.7, 5.4 Hz, 1H), 4.01 (q, <i>J</i> = 7.9 Hz, 1H), 3.94 (s, 3H), 3.93 (s, 3H), 2.97–2.93 (m, 4H), 2.89 (dd, <i>J</i> = 12.3, 3.9 Hz, 2H), 2.74 (s, 2H), 2.42 (s, 3H), 2.35–2.26 (m, 2H), 1.91 (s, 2H); 564.3 [M+H]<sup>+</sup></p>



5 10 20 15 20		(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-6-(4-methylpiperazin-1-yl)pyridin-3-yl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 8.76 (s, 1H), 8.18 (d, <i>J</i> = 1.0 Hz, 1H), 8.04 (s, 1H), 7.83 (s, 1H), 7.39–7.31 (m, 2H), 7.28 (d, <i>J</i> = 6.8 Hz, 1H), 7.23–7.19 (m, 1H), 6.42 (s, 1H), 5.75 (dd, <i>J</i> = 8.8, 5.1 Hz, 1H), 4.20 (td, <i>J</i> = 7.9, 3.7 Hz, 1H), 3.90 (s, 3H), 3.87 (s, 3H), 3.15 (s, 5H), 2.81 (dtd, <i>J</i> = 12.0, 8.1, 3.7 Hz, 4H), 2.27–2.16 (m, 2H), 1.91 (s, 3H); 564.3 [M+H] <sup>+</sup>
25 30 35 40 45 50		6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-(1-methyl-1H-pyrazol-4-yl)-5-((2R,3S)-2-methyl-3-((methylsulfonyl)methyl)azetidin-1-yl)-4-morpholinophenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, DMSO- <i>d</i> <sub>6</sub> ) δ 9.30 (s, 1H), 8.23 (d, <i>J</i> = 0.9 Hz, 1H), 7.72 (s, 1H), 7.40 (s, 1H), 7.29 (d, <i>J</i> = 7.0 Hz, 1H), 7.23–7.19 (m, 2H), 6.87 (s, 1H), 6.81 (d, <i>J</i> = 2.5 Hz, 1H), 6.49 (d, <i>J</i> = 1.0 Hz, 1H), 4.60 (t, <i>J</i> = 7.8 Hz, 1H), 4.22 (d, <i>J</i> = 3.8 Hz, 1H), 3.89 (s, 3H), 2.99 (s, 4H), 2.74–2.70 (m, 2H), 2.68–2.66 (m, 1H), 2.33 (t, <i>J</i> = 1.9 Hz, 1H), 2.26–2.19 (m, 3H), 1.78 (s, 6H), 1.38 (d, <i>J</i> = 6.0 Hz, 3H), 1.24 (s, 3H); 681.3 [M+H] <sup>+</sup>

5 10 15 20		(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N4-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-N2-methylpyrimidin-2,4-diamine	595.3 [M+H] <sup>+</sup>
25 30 35		(R)-4-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyridin-2-amine	565.3 [M+H] <sup>+</sup>
40 45 50 55		(R)-N-(4-(4-(4-allylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-2-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.31 (s, 1H), 7.58 (d, <i>J</i> = 8.7 Hz, 1H), 7.36-7.29 (m, 1H), 7.11-6.98 (m, 2H), 6.79 (s, 1H), 6.59-6.48 (m, 2H), 6.44 (d, <i>J</i> = 1.0 Hz, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 5.30 (d, <i>J</i> = 12.9 Hz, 2H), 4.10 (td, <i>J</i> = 7.9, 4.2 Hz,

		yl)pyrimidin-4-amine	1H), 3.92-3.82 (m, 4H), 3.73 (d, $J = 12.2$ Hz, 2H), 3.32-2.69 (m, 13H), 2.35-2.25 (m, 1H), 2.22-1.63 (m, 6H); 592.4 [M+H] <sup>+</sup>
5 10 15 20 25 30 35	 <p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-ethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>		<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.31 (s, 1H), 7.56 (d, $J = 9.1$ Hz, 1H), 7.36-7.29 (m, 1H), 7.11-6.99 (m, 2H), 6.76 (s, 1H), 6.54 (dq, $J = 5.9, 2.6$ Hz, 2H), 6.43 (d, $J = 1.0$ Hz, 1H), 5.91 (dd, $J = 8.9, 4.7$ Hz, 1H), 4.10 (td, $J = 7.9, 4.2$ Hz, 1H), 3.91-3.82 (m, 4H), 3.71 (d, $J = 12.3$ Hz, 2H), 2.99 (s, 6H), 2.89-2.69 (m, 6H), 2.62-2.49 (m, 1H), 2.30 (dtd, $J = 12.7, 8.2, 4.6$ Hz, 1H), 2.06-1.93 (m, 2H), 1.79-1.66 (m, 2H), 1.44-1.23 (m, 4H); 580.43 [M+H] <sup>+</sup>
40 45 50 55	 <p>(R)-N-(4-(4-(4-cyclobutylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine</p>		<sup>1</sup> H NMR (400 MHz, Methanol- <i>d</i> <sub>4</sub> ) $\delta$ 8.11 (d, $J = 1.0$ Hz, 1H), 7.36-7.30 (m, 2H), 7.16 (dddd, $J = 16.0, 12.9, 8.6, 7.1$ Hz, 2H), 6.71 (d, $J = 2.6$ Hz, 1H), 6.62 (dd, $J = 8.7, 2.5$ Hz, 1H), 6.27 (d, $J = 1.1$ Hz, 1H), 5.79 (dd, $J = 8.8, 4.8$ Hz, 1H), 4.14 (td, $J = 7.9,$

5

10

15

azolidin-2-  
yl)pyrimidin-4-  
amine

4.1 Hz, 1H), 3.92 (q,  $J = 8.0$  Hz, 1H),  
3.85 (s, 3H), 3.81 (d,  $J = 12.3$  Hz,  
2H), 3.02 (p,  $J = 8.2$  Hz, 1H), 2.92-  
2.74 (m, 7H), 2.64 (s, 2H), 2.59-2.51  
(m, 2H), 2.33-2.22 (m, 1H), 2.20-2.09  
(m, 2H), 2.08-1.97 (m, 4H), 1.83-1.64  
(m, 5H); 606.42[M+H]<sup>+</sup>

20

25


30

35

40

45

20  
9







(R)-6-(3-(2,3-  
difluorophenyl)isox  
azolidin-2-yl)-N-  
(2-methoxy-4-(4-(4-  
(oxetan-3-  
yl)piperazin-1-  
yl)piperidin-1-  
yl)phenyl)pyrimidin  
-4-amine



<sup>1</sup>H NMR (400 MHz, Chloroform-*d*)  $\delta$  8.30  
(s, 1H), 7.54 (d,  $J = 8.9$  Hz, 1H),  
7.35-7.31 (m, 1H), 7.11-6.99 (m, 2H),  
6.76 (s, 1H), 6.58-6.52 (m, 2H),  
6.46-6.40 (m, 1H), 5.91 (dd,  $J = 8.8$ ,  
4.7 Hz, 1H), 4.68 (t,  $J = 6.6$  Hz, 2H),  
4.62 (t,  $J = 6.1$  Hz, 2H), 4.09 (td,  
 $J = 7.9$ , 4.2 Hz, 1H), 3.91-3.83 (m,  
4H), 3.72 (d,  $J = 12.2$  Hz, 2H), 3.53  
(t,  $J = 6.5$  Hz, 1H), 2.87-2.63 (m,  
7H), 2.45 (s, 4H), 2.30 (dtd,  $J =$   
12.6, 8.0, 4.7 Hz, 2H), 2.03-1.92 (m,  
2H), 1.80-1.69 (m, 2H); 608.37[M+H]<sup>+</sup>



50

55



5 10 15 20 25		<p>(R)-N-(4-(4-(4-(cyclopropylmethyl)piperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (s, 1H), 7.55 (d, <i>J</i> = 9.1 Hz, 1H), 7.35-7.30 (m, 1H), 7.10-6.99 (m, 2H), 6.80 (s, 1H), 6.57-6.51 (m, 2H), 6.43 (s, 1H), 5.91 (dd, <i>J</i> = 8.9, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.92-3.80 (m, 4H), 3.71 (d, <i>J</i> = 12.0 Hz, 2H), 3.01 (s, 6H), 2.86-2.49 (m, 8H), 2.31 (ddd, <i>J</i> = 15.3, 8.0, 4.2 Hz, 1H), 2.07-1.95 (m, 2H), 1.80-1.68 (m, 2H), 1.14 (s, 1H), 0.68 (s, 2H), 0.30 (s, 2H); 606.33 [M+H] <sup>+</sup>
30 35 40 45 50 55		<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(6-methyl-2,6-diazaspiro[3.3]heptan-2-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.28 (s, 1H), 7.47 (d, <i>J</i> = 8.3 Hz, 1H), 7.32 (t, <i>J</i> = 7.2 Hz, 1H), 7.12-6.99 (m, 2H), 6.93 (s, 1H), 6.57-6.48 (m, 2H), 6.44-6.37 (m, 1H), 5.90 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.09 (td, <i>J</i> = 7.9, 4.1 Hz, 1H), 3.92-3.86 (m, 1H), 3.84 (s, 3H), 3.68-3.64 (m, 2H), 3.07-2.88 (m, 5H), 2.84-2.67 (m, 4H), 2.31 (tt, <i>J</i> = 12.3, 8.0 Hz, 2H), 2.21-2.10 (m, 2H), 2.05 (s, 3H), 1.79 (d, <i>J</i> = 13.0 Hz, 2H), 1.52-1.41 (m, 2H); 578.4


			[M+H] <sup>+</sup>
5			
10			
15	21	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-3-(dimethylamino)pyrrolidin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (s, 1H), 7.57 (d, <i>J</i> = 8.3 Hz, 1H), 7.34-7.30 (m, 1H), 7.12-7.01 (m, 2H), 6.84 (s, 1H), 6.57-6.50 (m, 2H), 6.43 (s, 1H), 5.91 (dd, <i>J</i> = 8.5, 4.5 Hz, 1H), 4.11 (td, <i>J</i> = 7.9, 4.1 Hz, 1H), 3.93-3.83 (m, 4H), 3.64 (d, <i>J</i> = 10.9 Hz, 2H), 3.15 (s, 2H), 2.86-2.74 (m, 4H), 2.63 (s, 6H), 2.30 (ddd, <i>J</i> = 16.3, 7.7, 4.0 Hz, 2H), 2.17-1.99 (m, 4H), 1.97-1.70 (m, 4H); 580.33[M+H] <sup>+</sup>
20	2		
25			
30			
35	21	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-3,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Methanol- <i>d</i> <sub>4</sub> ) δ 8.12 (s, 1H), 7.34-7.11 (m, 4H), 6.74 (s, 1H), 6.67 (s, 1H), 6.16 (s, 1H), 5.83-5.75 (m, 1H), 4.29 (td, <i>J</i> = 7.7, 4.0 Hz, 1H), 4.06 (q, <i>J</i> = 7.8 Hz, 1H), 3.85 (s, 4H), 3.46 (d, <i>J</i> = 15.1 Hz, 1H), 3.32-3.22 (m, 3H), 3.17-3.03 (m, 2H), 3.02-2.66 (m, 8H), 2.55-2.44 (m, 1H), 2.37 (td, <i>J</i> = 13.2, 7.6 Hz, 1H), 2.10-1.99 (m, 2H), 1.82-1.70 (m, 2H), 1.37 (d, <i>J</i> = 6.5 Hz, 3H); 580.38[M+H] <sup>+</sup>
40	3		
45			
50			



5 10 15 20 25 30	<p>21 4</p> 	<p>6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((S)-3,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Methanol-<i>d</i><sub>4</sub>) δ 7.99 (d, <i>J</i> = 1.0 Hz, 1H), 7.24-7.18 (m, 2H), 7.11-6.97 (m, 2H), 6.59 (d, <i>J</i> = 2.5 Hz, 1H), 6.49 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H), 6.15 (d, <i>J</i> = 1.0 Hz, 1H), 5.67 (dd, <i>J</i> = 8.8, 4.8 Hz, 1H), 4.02 (td, <i>J</i> = 7.9, 4.1 Hz, 1H), 3.80 (q, <i>J</i> = 8.0 Hz, 1H), 3.73 (s, 3H), 3.69 (d, <i>J</i> = 12.3 Hz, 2H), 2.98 (t, <i>J</i> = 12.9 Hz, 3H), 2.75 (dtd, <i>J</i> = 12.1, 8.0, 4.1 Hz, 1H), 2.71-2.61 (m, 2H), 2.60-2.38 (m, 7H), 2.21-2.10 (m, 2H), 1.92 (d, <i>J</i> = 12.4 Hz, 2H), 1.58 (q, <i>J</i> = 12.1, 11.4 Hz, 2H), 1.11 (d, <i>J</i> = 6.3 Hz, 3H); 580.38 [M+H]<sup>+</sup></p>
35 40 45 50 55	<p>21 5</p> 	<p>6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((3S,5R)-4-ethyl-3,5-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.31 (s, 1H), 7.55 (d, <i>J</i> = 7.4 Hz, 1H), 7.35-7.30 (m, 1H), 7.10-7.01 (m, 2H), 6.72 (s, 1H), 6.54 (dd, <i>J</i> = 4.6, 2.3 Hz, 2H), 6.44 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.92-3.80 (m, 4H), 3.70 (d, <i>J</i> = 11.9 Hz, 2H), 3.44-3.33 (m, 1H), 3.28-3.20 (m, 1H), 3.11-3.03 (m,</p>



5  10		idin-4-amine	1H), 2.98-2.89 (m, 2H), 2.86-2.77 (m, 2H), 2.74 (t, <i>J</i> = 12.4 Hz, 2H), 2.54-2.43 (m, 1H), 2.37-2.25 (m, 2H), 1.97-1.89 (m, 2H), 1.71-1.64 (m, 2H), 1.61-1.44 (m, 10H); 608.41[M+H] <sup>+</sup>
15  20  25  30  35	21  6  	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((S)-2,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (s, 1H), 7.59 (s, 1H), 7.31 (t, <i>J</i> = 7.1 Hz, 1H), 7.11-7.01 (m, 2H), 6.89 (s, 1H), 6.58-6.50 (m, 2H), 6.42 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.11 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.90 (t, <i>J</i> = 8.0 Hz, 1H), 3.86 (s, 3H), 3.74 (d, <i>J</i> = 6.3 Hz, 2H), 3.46-3.36 (m, 1H), 3.11-2.94 (m, 3H), 2.91-2.65 (m, 6H), 2.36-2.26 (m, 1H), 1.99-1.79 (m, 3H), 1.75-1.46 (m, 6H), 1.31-1.12 (m, 2H); 580.43[M+H] <sup>+</sup>
40  45  50  55	21  7  	6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-2,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.31 (s, 1H), 7.59 (s, 1H), 7.36-7.29 (m, 1H), 7.12-7.01 (m, 2H), 6.75 (s, 1H), 6.58-6.50 (m, 2H), 6.44 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.14-4.05 (m, 1H), 3.89 (q, 1H), 3.86 (s, 3H), 3.73 (d, <i>J</i> = 14.1 Hz, 2H), 3.38-3.31





5		methoxyphenyl)pyrimidin-4-amine	(m, 1H), 3.07-2.90 (m, 3H), 2.86-2.70 (m, 6H), 2.31 (ddt, $J = 16.1, 7.7, 4.2$ Hz, 1H), 1.93-1.81 (m, 3H), 1.73-1.50 (m, 6H), 1.32-1.26 (m, 1H), 1.18-1.12 (m, 1H); 580.38 [M+H] <sup>+</sup>
15 20 25 30 35 40		(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(3-(4-methylpiperazin-1-yl)azetidin-1-yl)phenyl)pyrimidin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.28 (s, 1H), 7.38 (d, $J = 8.4$ Hz, 1H), 7.36-7.30 (m, 1H), 7.12-6.99 (m, 2H), 6.62 (s, 1H), 6.33 (s, 1H), 6.07 (dd, $J = 8.4, 2.4$ Hz, 1H), 6.04 (d, $J = 2.4$ Hz, 1H), 5.90 (dd, $J = 8.9, 4.7$ Hz, 1H), 4.08 (td, $J = 8.0, 4.2$ Hz, 1H), 4.01 (t, $J = 7.0$ Hz, 2H), 3.91-3.84 (m, 1H), 3.82 (s, 3H), 3.72 (dd, $J = 7.1, 5.4$ Hz, 2H), 3.41 (p, $J = 6.1$ Hz, 1H), 2.89-2.55 (m, 9H), 2.46 (s, 3H), 2.29 (dtd, $J = 12.6, 8.1, 4.8$ Hz, 1H); 502.3 [M+H] <sup>+</sup>
45 50 55		6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(6-ethyl-3,6-diazabicyclo[3.1.1]heptan-3-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.31 (d, $J = 3.6$ Hz, 1H), 7.64 (d, $J = 8.6$ Hz, 1H), 7.33 (d, $J = 7.6$ Hz, 1H), 7.12-6.99 (m, 2H), 6.86 (s, 1H), 6.55 (d, $J = 5.0$ Hz, 1H), 6.49 (d, $J = 8.9$ Hz, 1H), 6.44 (s, 1H), 5.94-5.87 (m,

5 10 15 20 25		yl)piperidin-1-yl)- 2- methoxyphenyl)pyrim idin-4-amine	1H), 4.10 (tt, $J = 16.2, 12.4, 5.5$ Hz, 2H), 3.94-3.87 (m, 2H), 3.85 (s, 3H), 3.76-3.53 (m, 4H), 3.44-3.32 (m, 1H), 3.30-3.16 (m, 1H), 2.97-2.77 (m, 4H), 2.74 (d, $J = 11.2$ Hz, 1H), 2.69- 2.53 (m, 1H), 2.48 (d, $J = 12.0$ Hz, 1H), 2.30 (dtd, $J = 20.7, 14.5, 12.9,$ 6.0 Hz, 2H), 2.19 (d, $J = 13.2$ Hz, 1H), 2.09-1.95 (m, 2H), 1.85-1.71 (m, 1H), 1.38 (t, $J = 7.3$ Hz, 3H); 592.4 [M+H] <sup>+</sup>
30 35 40 45 50 55	22 0 	6-((R)-3-(2,3- difluorophenyl)isox azolidin-2-yl)-N- (4-(4-(3-ethyl-3,6- diazabicyclo[3.1.1] heptan-6- yl)piperidin-1-yl)- 2- methoxyphenyl)pyrim idin-4-amine	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.25 (s, 1H), 7.52 (s, 1H), 7.32 (t, $J =$ 10.5 Hz, 1H), 7.07 (d, $J = 6.3$ Hz, 2H), 6.53 (s, 2H), 6.50 (d, $J = 8.3$ Hz, 1H), 6.32 (s, 1H), 5.89-5.84 (m, 1H), 4.30-4.10 (m, 4H), 3.94 (p, $J =$ 8.0 Hz, 3H), 3.86-3.83 (m, 3H), 3.75- 3.73 (m, 1H), 3.35-3.31 (m, 1H), 3.14-3.10 (m, 1H), 2.96-2.80 (m, 6H), 2.68-2.59 (m, 1H), 2.39-2.30 (m, 3H), 2.18 (d, $J = 2.8$ Hz, 1H), 2.05-1.95 (m, 2H), 1.41 (t, $J = 8.5$ Hz, 3H); 592.4 [M+H] <sup>+</sup>




<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p> <p>30</p>	<p>22</p> <p>1</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methyl-1,4-diazepan-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.31 (s, 1H), 7.59 (d, <i>J</i> = 8.6 Hz, 1H), 7.34 - 7.29 (m, 1H), 7.11 - 6.99 (m, 2H), 6.83 (s, 1H), 6.54 (d, <i>J</i> = 7.1 Hz, 2H), 6.43 (s, 1H), 5.91 (dd, <i>J</i> = 8.6, 4.3 Hz, 1H), 4.11 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.93 - 3.83 (m, 4H), 3.72 (d, <i>J</i> = 12.5 Hz, 2H), 3.30 (s, 2H), 3.07 (s, 2H), 2.87 - 2.71 (m, 7H), 2.32 (ddd, <i>J</i> = 15.2, 7.8, 3.9 Hz, 2H), 2.05 - 1.96 (m, 2H), 1.81 - 1.69 (m, 2H), 1.34 - 1.23 (m, 3H), 0.91 - 0.83 (m, 2H) ;580.43[M+H]<sup>+</sup></p>
<p>35</p> <p>40</p> <p>45</p> <p>50</p> <p>55</p>	<p>22</p> <p>2</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-isopropylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.31 (s, 1H), 7.55 (d, <i>J</i> = 8.9 Hz, 1H), 7.36-7.30 (m, 1H), 7.11-6.99 (m, 2H), 6.74 (s, 1H), 6.54 (dq, <i>J</i> = 5.9, 2.6 Hz, 2H), 6.45-6.41 (m, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 7.9, 4.2 Hz, 1H), 3.92-3.83 (m, 4H), 3.71 (d, <i>J</i> = 12.3 Hz, 2H), 3.10-2.71 (m, 9H), 2.55-2.45 (m, 1H), 2.30 (ddt, <i>J</i> = 16.5, 8.2, 4.8 Hz, 1H), 1.97 (s, 2H), 1.91-1.82 (m, 1H), 1.78-1.64</p>



5			(m, 3H), 1.56-1.47 (m, 1H), 1.45-1.23 (m, 6H); 594.44 [M+H] <sup>+</sup>
10 15 20 25		(R)-1-(1-(4-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-3-methoxyphenyl)piperidin-4-yl)-4-methylpiperazin-2-one	580.3 [M+H] <sup>+</sup>
30 35 40 45 50 55		(R)-4-(1-(4-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-3-methoxyphenyl)piperidin-4-yl)-1-methylpiperazin-2-one	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.30 (s, 1H), 7.54 (d, <i>J</i> = 8.4 Hz, 1H), 7.36-7.29 (m, 1H), 7.12-7.00 (m, 2H), 6.85 (s, 1H), 6.59-6.50 (m, 2H), 6.42 (d, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.10 (td, <i>J</i> = 8.4, 7.9, 4.6 Hz, 1H), 3.89 (q, 1H), 3.85 (s, 3H), 3.69 (d, <i>J</i> = 11.2 Hz, 2H), 3.34 (t, <i>J</i> = 5.4 Hz, 2H), 3.31 (s, 2H), 2.97 (s, 3H), 2.78 (ddd, <i>J</i> = 23.8, 11.4, 3.9 Hz, 5H), 2.43 (tt, <i>J</i> = 11.1, 3.7 Hz, 1H), 2.30 (dtd, <i>J</i> = 12.6, 8.1, 4.7

			Hz, 1H), 1.96 (d, $J = 12.5$ Hz, 2H), 1.70 (qd, $J = 11.9, 4.0$ Hz, 2H); 580.33 [M+H] <sup>+</sup>
5 10 15 20 25	 <p>22 5</p>	(R)-1-methyl-5-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)pyridin-2(1H)-one	448.19 [M+H] <sup>+</sup>
30 35 40 45	 <p>22 6</p>	(R)-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)-1-methylpyridin-2(1H)-one	561.33 [M+H] <sup>+</sup>

50



55

5 22 10 7 15		(S)-1-methyl-5-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)pyridin-2(1H)-one	448.19 [M+H] <sup>+</sup>
20 25 22 30 8 35 40		(S)-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)-1-methylpyridin-2(1H)-one	561.33 [M+H] <sup>+</sup>
45 22 9 50 55		N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylbenzo[d]isoxazol-2(3H)-yl)pyrimidin-4-	<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) δ 8.36 (s, 1H), 8.01 (dd, <i>J</i> = 7.2, 1.9 Hz, 1H), 7.39 (p, <i>J</i> = 4.4 Hz, 5H), 6.98-6.75 (m, 9H), 5.64 (s, 1H), 3.22 (d, <i>J</i> = 4.3 Hz, 4H), 2.63 (t, <i>J</i> = 5.0 Hz, 4H), 2.39 (s, 3H); 465.2 [M+H] <sup>+</sup>

		amine	
5			<sup>1</sup> H NMR (400 MHz, Chloroform- <i>d</i> ) $\delta$ 8.38
10		N-(2-methoxy-4-(4-	( <i>s</i> , 1H), 8.01 (dd, <i>J</i> = 7.6, 1.6 Hz,
15		(4-methylpiperazin-	1H), 7.46-7.37 (m, 5H), 6.99-6.86 (m,
20		1-yl)piperidin-1-	4H), 6.79 (dd, <i>J</i> = 7.6, 1.5 Hz, 1H),
25		yl)phenyl)-6-(3-	6.60 ( <i>s</i> , 1H), 6.50 (d, <i>J</i> = 2.5 Hz,
		phenylbenzo[d]isoxa-	1H), 6.39 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H),
		zol-2(3H)-	5.70 ( <i>s</i> , 1H), 3.75 ( <i>s</i> , 3H), 3.71 (d,
		yl)pyrimidin-4-	<i>J</i> = 12.0 Hz, 2H), 2.80-2.64 (m, 6H),
		amine	2.59-2.37 (m, 5H), 2.32 ( <i>s</i> , 3H), 1.98
			(d, <i>J</i> = 12.4 Hz, 2H), 1.73 (tt, <i>J</i> =
			13.0, 6.6 Hz, 2H); 578.3[M+H] <sup>+</sup>
30		( <i>S</i> )-N-(4-(4-	<sup>1</sup> H NMR (400 MHz, MeOD) $\delta$ 8.13 (d, <i>J</i> =
35		methylpiperazin-1-	0.8 Hz, 1H), 7.43-7.25 (m, 7H), 7.07-
40		yl)phenyl)-6-(7-	7.00 (m, 2H), 6.37 (d, <i>J</i> = 0.8 Hz,
45		phenyl-5-oxa-6-	1H), 5.13 ( <i>s</i> , 1H), 4.07-3.97 (m, 2H),
		azaspiro[2.4]heptan-	3.37-3.33 (m, 4H), 3.10-2.99 (m, 4H),
		-6-yl)pyrimidin-4-	2.67 ( <i>s</i> , 3H), 0.85 (ddd, <i>J</i> = 9.9, 5.8,
		amine	4.5 Hz, 1H), 0.72 (dtd, <i>J</i> = 15.7, 9.8,
			6.1 Hz, 2H), 0.38 (dt, <i>J</i> = 10.7, 5.4
			Hz, 1H); 443.25 [M+H] <sup>+</sup>

50



55

5 10 15 20	<p>23 2</p> 	<p>(R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, MeOD) δ 8.13 (d, <i>J</i> = 0.9 Hz, 1H), 7.46-7.21 (m, 7H), 7.08-6.95 (m, 2H), 6.37 (d, <i>J</i> = 0.8 Hz, 1H), 5.13 (s, 1H), 4.09-3.96 (m, 2H), 3.35 (dd, <i>J</i> = 9.6, 4.2 Hz, 4H), 3.14-3.02 (m, 4H), 2.69 (s, 3H), 0.88-0.81 (m, 1H), 0.72 (dtd, <i>J</i> = 15.8, 9.8, 6.1 Hz, 2H), 0.38 (dt, <i>J</i> = 10.7, 5.4 Hz, 1H); 443.21 [M+H]<sup>+</sup></p>
25 30 35 40 45	<p>23 3</p> 	<p>(S)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, MeOD) δ 8.10 (d, <i>J</i> = 0.6 Hz, 1H), 7.37-7.28 (m, 6H), 6.68 (d, <i>J</i> = 2.3 Hz, 1H), 6.58 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H), 6.23 (s, 1H), 5.12 (s, 1H), 4.02-3.97 (m, 2H), 3.85-3.75 (m, 5H), 2.72 (dd, <i>J</i> = 29.9, 19.2 Hz, 10H), 2.49-2.42 (m, 1H), 2.39 (s, 3H), 2.02 (d, <i>J</i> = 12.0 Hz, 2H), 1.69-1.64 (m, 2H), 0.86-0.80 (m, 1H), 0.70 (ddd, <i>J</i> = 14.7, 9.6, 3.9 Hz, 2H), 0.36 (dt, <i>J</i> = 10.7, 5.4 Hz, 1H); 556.30 [M+H]<sup>+</sup></p>

50



55



<p>5</p> <p>10</p> <p>15</p> <p>20</p> <p>25</p>	<p>23</p> <p>4</p> 	<p>(R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, MeOD) δ 8.09 (d, <i>J</i> = 0.9 Hz, 1H), 7.43–7.24 (m, 6H), 6.68 (d, <i>J</i> = 2.5 Hz, 1H), 6.59 (dd, <i>J</i> = 8.7, 2.5 Hz, 1H), 6.23 (d, <i>J</i> = 0.5 Hz, 1H), 5.12 (s, 1H), 4.05–3.94 (m, 2H), 3.85–3.73 (m, 5H), 2.95–2.58 (m, 10H), 2.48 (ddd, <i>J</i> = 11.5, 7.9, 3.7 Hz, 1H), 2.43 (d, <i>J</i> = 9.3 Hz, 3H), 2.08–1.98 (m, 2H), 1.72–1.62 (m, 2H), 0.87–0.80 (m, 1H), 0.70 (dtd, <i>J</i> = 15.7, 9.8, 6.1 Hz, 2H), 0.36 (dt, <i>J</i> = 10.7, 5.4 Hz, 1H); 556.30 [M+H]<sup>+</sup></p>
<p>30</p> <p>35</p> <p>40</p> <p>45</p>	<p>23</p> <p>5</p> 	<p>(R)-6-(3-isopropylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, MeOD) δ 8.04 (s, 1H), 7.23–7.17 (m, 2H), 6.95–6.88 (m, 2H), 6.19 (d, <i>J</i> = 0.4 Hz, 1H), 4.12 (td, <i>J</i> = 7.9, 4.7 Hz, 1H), 3.93 (td, <i>J</i> = 8.1, 3.5 Hz, 1H), 3.59 (dt, <i>J</i> = 15.4, 7.7 Hz, 1H), 2.96–2.90 (m, 4H), 2.56–2.51 (m, 7H), 2.19–2.00 (m, 2H), 1.77 (dq, <i>J</i> = 13.6, 6.8 Hz, 1H), 0.90 (dd, <i>J</i> = 11.4, 6.7 Hz, 6H); 383.30 [M+H]<sup>+</sup></p>

50

55

<p>5</p> <p>10</p> <p>15</p> <p>20</p>	<p>23</p> <p>6</p> 	<p>(S)-6-(3-isopropylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, MeOD) δ 8.16 (s, 1H), 7.36-7.29 (m, 2H), 7.07-7.00 (m, 2H), 6.31 (s, 1H), 4.24 (td, <i>J</i> = 7.9, 4.7 Hz, 1H), 4.05 (td, <i>J</i> = 8.1, 3.4 Hz, 1H), 3.70 (dt, <i>J</i> = 15.6, 7.8 Hz, 1H), 3.09-3.02 (m, 4H), 2.66 (d, <i>J</i> = 7.1 Hz, 7H), 2.31-2.10 (m, 2H), 1.89 (dq, <i>J</i> = 13.6, 6.8 Hz, 1H), 1.01 (dd, <i>J</i> = 11.4, 6.7 Hz, 6H); 383.30 [M+H]<sup>+</sup></p>
<p>25</p> <p>30</p> <p>35</p> <p>40</p> <p>45</p> <p>50</p>	<p>23</p> <p>7</p> 	<p>(R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-(2-(methylsulfonyl)ethyl)piperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine</p>	<p><sup>1</sup>H NMR (400 MHz, Chloroform-<i>d</i>) δ 8.30 (s, 1H), 7.55-7.49 (m, 1H), 7.36-7.28 (m, 1H), 7.10-7.01 (m, 2H), 6.82 (s, 1H), 6.58-6.51 (m, 2H), 6.42 (s, 1H), 5.91 (dd, <i>J</i> = 8.8, 4.7 Hz, 1H), 4.09 (td, <i>J</i> = 7.9, 4.3 Hz, 1H), 3.88 (q, <i>J</i> = 8.0 Hz, 1H), 3.85 (s, 3H), 3.71 (d, <i>J</i> = 12.1 Hz, 2H), 3.15 (t, <i>J</i> = 6.4 Hz, 2H), 3.04 (s, 3H), 2.89 (t, <i>J</i> = 6.5 Hz, 2H), 2.86-2.70 (m, 3H), 2.64 (s, 3H), 2.58 (s, 3H), 2.43-2.36 (m, 1H), 2.34-2.26 (m, 1H), 1.95 (d, <i>J</i> = 12.5 Hz, 3H), 1.69 (ddd, <i>J</i> = 23.8, 11.7, 3.3 Hz, 3H); 658.35 [M+H]<sup>+</sup></p>

55 <Experimental Example 1> Evaluation of proliferation inhibitory activity of EGFR mutant overexpression Ba/F3 cell

[0180] The following experiments were performed to evaluate the inhibitory activity of the compounds according to the present disclosure on Ba/F3 proliferation expressing EGFR C797S, L861Q, G719A, S768I, L718Q, and/or G724S

EP 4 342 895 A1

mutations.

**[0181]** For Ba/F3 cells, RPMI-1640 containing 10% FBS and 5 ng/ml IL-3 (R&D Systems) was used. Transduced Ba/F3 cells were cultured after adding 1 ug/ml puromycin (Invitrogen) to the same medium.

**[0182]** At 24 hours before treatment with the compounds, 3000 to 5000 cells were aliquoted into each well of a white clear bottom 96 well plate (Corning). The compounds were diluted in dimethyl sulfoxide (3 times dilution, 12 concentrations in total), and injected in an amount of 1 µl so that the final concentration was 0.2 nM to 5 µM. For measurement of living cells, after 72 hours of compound treatment, the cells were stored at room temperature for 10 min using CellTiter-Glo luminescent cell-viability reagent (Promega), and then luminescence intensity was measured using a reader (Synergy-Neo, Biotek). Each test was repeated three times. Result values were calculated as the cell growth rate (%) compared to the control. The graphs were created using GraphPad Prism version 8.3.0 program and the GI<sub>50</sub> values were calculated.

**[0183]** Table 2 below shows the evaluation results of the proliferation inhibitory activity of Ba/F3 cells expressing EGFR Del19/C797S (EGFR DC) and EGFR L858R/C797S (EGFR LC) mutations.

[Table 2]

Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)	Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)	Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)
1	A	A	79	A	-	141	A	-
2	B	-	80	B	-	142	A	-
3	A	-	81	B	-	143	A	-
4	A	A	83	A	-	145	A	-
5	A	A	84	A	A	146	A	-
6	B	B	85	A	-	147	A	-
7	B	B	86	A	-	152	A	-
8	A	A	87	A	-	154	A	-
10	B	B	88	A	-	155	A	-
11	B	B	89	B	-	157	A	-
12	B	B	90	A	-	159	B	-
13	B	B	91	A	-	160	B	-
14	B	-	92	A	-	161	A	-
15	A	A	93	A	-	162	A	-
17	B	-	94	A	-	163	A	-
18	B	-	95	A	-	164	A	-
19	B	-	96	A	-	165	A	-
20	A	-	97	A	-	166	A	-
21	A	-	98	A	-	167	A	-
22	A	-	99	A	-	169	B	-
23	B	-	100	B	-	170	B	-
24	B	-	101	A	-	174	B	-
25	A	-	102	A	A	176	B	-
26	A	-	103	A	-	177	B	-
27	B	-	104	A	A	180	B	-
28	B	-	105	A	-	182	A	-
29	A	-	106	A	-	184	B	-
30	A	-	107	A	-	193	B	-

EP 4 342 895 A1

(continued)

Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)	Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)	Example Compound	Ba/F3 (EGFR DC)	Ba/F3 (EGFR LC)
31	A	-	108	A	-	195	A	-
32	A	-	109	A	-	199	A	-
33	A	-	110	A	-	200	A	-
34	A	-	111	B	-	201	A	-
35	A	-	112	B	-	202	A	-
36	B	-	113	A	-	203	A	-
38	B	-	114	A	-	206	A	-
41	B	-	115	A	-	207	A	-
48	A	-	116	A	-	208	A	-
49	A	-	117	A	-	209	A	-
52	A	-	118	A	-	210	A	-
53	A	-	119	A	-	211	A	-
54	A	-	120	A	-	212	A	-
56	A	-	121	A	-	213	A	-
57	A	-	122	A	-	214	A	-
59	A	-	123	A	-	215	A	-
61	A	-	124	A	-	216	A	-
62	A	-	125	A	-	217	A	-
63	A	-	126	A	-	218	A	-
64	A	-	127	A	-	219	A	-
65	A	-	128	A	A	220	A	-
66	A	-	129	A	A	221	A	-
67	A	-	130	B	-	222	A	-
68	A	-	131	B	-	224	A	-
69	A	-	132	A	A	229	B	-
72	B	-	133	B	-	230	B	-
73	B	-	134	A	-	231	B	-
74	A	-	135	A	-	233	A	-
75	A	-	136	A	-	234	B	-
76	A	-	138	A	-	235	A	-
77	A	-	139	A	-	237	A	-
78	A	-	140	A	-			

A:  $GI_{50} < 50$  nM; B:  $50$  nM  $\leq GI_{50} < 500$  nM; C:  $500$  nM  $\leq GI_{50} < 5000$  nM; D:  $5000$  nM  $\leq GI_{50}$ ;

[0184] Table 3 below shows the evaluation results of the proliferation inhibitory activity of Ba/F3 cells such as L861Q, G719A, S768I, L718Q, G724S, and the like, expressing EGFR rare (or uncommon) and drug-resistant mutations.

[Table 3]

Example Compound	Ba/F3 G719A	Ba/F3 L861Q	Ba/F3 S768I	Ba/F3 L718Q	Ba/F3 G724S
<b>4</b>	A	A	B	-	-
<b>34</b>	A	B	B	B	A

A:  $GI_{50} < 50$  nM; B:  $50$  nM  $\leq GI_{50} < 500$  nM; C:  $500$  nM  $\leq GI_{50} < 5000$  nM; D:  $5000$  nM  $\leq GI_{50}$ ;

**[0185]** Table 4 below shows the activity values for EGFR-family mutant enzymes obtained by requesting on external entrusted organization, Reaction Biology (<https://www.reactionbiology.com/>).

[Table 4]

EGFR Mutant Enzyme	Compound <b>34</b> ( $IC_{50}$ ; nM)
EGFR (d746-750)	8.0
EGFR (d746-750/C797A)	3.5
EGFR (d746-750/C797S)	<0.5
EGFR (d746-750/T790M/C797S)	60.5
EGFR (D761Y)	<0.5
EGFR (G719C)	0.5
EGFR (G719D)	<0.5
EGFR (G719S)	2.7
EGFR (L718Q)	49.8
EGFR (L747S)	<0.5
EGFR (L792F)	8.7
EGFR (L792F/L858R)	78.1
EGFR (L861Q)	<0.5
ERBB2/HER2	5.3
ERBB4/HER4	41.7


**[0186]** As shown in Tables 2 to 4 above, it could be appreciated that the Example compounds of the present disclosure exhibited high inhibitory ability against overexpressing cell lines or enzymes including EGFR C797S mutations, rare mutations, and the like.

**[0187]** As described above, although the present disclosure has been described in detail through preferred Preparation Examples, Examples and Experimental Examples, the scope of the present disclosure is not limited to specific Example compounds, and should be interpreted by the appended claims. In addition, those skilled in the art will understand that many modifications and variations can be made without departing from the scope of the present disclosure.

## Claims

1. A compound represented by the following Chemical Formula 1, an optical isomer thereof, or a pharmaceutically acceptable salt thereof:

[Chemical Formula 1]



20 in the Chemical Formula 1,

$X_1$  to  $X_3$  are each independently CH or N;

$R_X$  is -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -NH<sub>2</sub>, -NH(-C<sub>1-6</sub>alkyl), or -N(-C<sub>1-6</sub>alkyl)(-C<sub>1-6</sub>alkyl);

25  $Y$  is -C<sub>1-6</sub>alkyl, -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>hydroaryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl in which at least one H of the -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>hydroaryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkyl-O-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -(C=O)NR<sub>1</sub>R<sub>2</sub>, -(C=O)OR<sub>3</sub>, -NR<sub>4</sub>R<sub>5</sub>, -OR<sub>6</sub>, -halo, =O, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the cycloalkyl, heterocycloalkyl, aryl, or heteroaryl may be substituted with

30 -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, or -halo;

$n$  is 0, 1, 2, 3, or 4;

$R_1$  to  $R_3$  are each independently -H, -C<sub>1-6</sub>alkyl, or cycloalkyl;

$R_4$  and  $R_5$  are each independently -H or -C<sub>1-6</sub>alkyl;

$R_6$  is -H, -C<sub>1-6</sub>alkyl, or phenyl in which at least one H of the phenyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, or halo;

35  $R_{Y1}$  to  $R_{Y5}$  are each independently -H or -C<sub>1-6</sub>alkyl, or  $R_{Y2}$  and  $R_{Y3}$  may be linked to each other to form cycloalkyl or heterocycloalkyl,  $R_{Y4}$  and  $R_{Y5}$  may be linked to each other to form cycloalkyl or heterocycloalkyl, and  $R_{Y3}$  and  $R_{Y4}$  may be linked to each other to form aryl or heteroaryl;

$L$  is -(CH<sub>2</sub>)<sub>m</sub>-, -C(=O)-, or null;

$m$  is 0, 1, 2, 3, or 4;

40 a ring  $Z$  is aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>cyanoalkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -NR<sub>7</sub>R<sub>8</sub>, -OH, -O-C<sub>1-6</sub>alkyl, -O-C<sub>1-6</sub>haloalkyl, -S-C<sub>1-6</sub>alkyl, -S-C<sub>1-6</sub>haloalkyl, -C(=O)-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>haloalkyl, -C(=O)O-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or  $Z_1$ , or two or more substituents of the aryl, heteroaryl, hydroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -O-C<sub>1-6</sub>alkyl, -halo, or  $Z_1$ ;

45  $R_7$  and  $R_8$  are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -C(=O)-C<sub>1-6</sub>alkyl, or -C(=O)-C<sub>1-6</sub>haloalkyl;

50  $Z_1$  is cycloalkyl, heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-C(=O)-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>9</sub>R<sub>10</sub>, -halo, cycloalkyl, or  $Z_2$ ;

55  $R_9$  and  $R_{10}$  are each independently -H or -C<sub>1-6</sub>alkyl;

Z<sub>2</sub> is heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, aryl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>11</sub>R<sub>12</sub>, cycloalkyl, or Z<sub>3</sub>; R<sub>11</sub> and R<sub>12</sub> are each independently -H or -C<sub>1-6</sub>alkyl; and Z<sub>3</sub> is heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl-C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or cycloalkyl.

2. The compound represented by Chemical Formula I, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of claim 1, wherein

X<sub>1</sub> to X<sub>3</sub> are each independently CH or N;

R<sub>x</sub> is -H, -NH<sub>2</sub>, -NH(-C<sub>1-6</sub>alkyl), or -N(-C<sub>1-6</sub>alkyl)(-C<sub>1-6</sub>alkyl);

Y is -C<sub>1-6</sub>alkyl, -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl in which at least one H of the -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -(C=O)NR<sub>1</sub>R<sub>2</sub>, -(C=O)OR<sub>3</sub>, -NR<sub>4</sub>R<sub>5</sub>, -OR<sub>6</sub>, -halo, =O, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the heterocycloalkyl, aryl, or heteroaryl may be substituted with -halo;

n is 0, 1, or 2;

R<sub>1</sub> to R<sub>3</sub> are each independently -H, -C<sub>1-6</sub>alkyl, or cycloalkyl;

R<sub>4</sub> and R<sub>5</sub> are each independently -H or -C<sub>1-6</sub>alkyl;

R<sub>6</sub> is -C<sub>1-6</sub>alkyl or phenyl in which at least one H of the phenyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, or halo;

R<sub>Y1</sub> to R<sub>Y5</sub> are each independently -H or -C<sub>1-6</sub>alkyl, or R<sub>Y2</sub> and R<sub>Y3</sub> may be linked to each other to form cycloalkyl, and R<sub>Y3</sub> and R<sub>Y4</sub> may be linked to each other to form aryl;

L is -(CH<sub>2</sub>)<sub>m</sub>-, -C(=O)-, or null;

m is 0, 1, or 2;

a ring Z is aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>cyanoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -NR<sub>7</sub>R<sub>8</sub>, -O-C<sub>1-6</sub>alkyl, -O-C<sub>1-6</sub>haloalkyl, -S-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>haloalkyl, -C(=O)O-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or Z<sub>1</sub>, or two or more substituents of the aryl, heteroaryl, hydroheteroaryl, cycloalkyl, or heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl or Z<sub>1</sub>;

R<sub>7</sub> and R<sub>8</sub> are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-NH-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -(C=O)-C<sub>1-6</sub>alkyl, or -(C=O)-C<sub>1-6</sub>haloalkyl;

Z<sub>1</sub> is cycloalkyl, heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl in which the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>9</sub>R<sub>10</sub>, -halo, cycloalkyl, or Z<sub>2</sub>;

R<sub>9</sub> and R<sub>10</sub> are each independently -H or -C<sub>1-6</sub>alkyl;

Z<sub>2</sub> is heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -NR<sub>11</sub>R<sub>12</sub>, cycloalkyl, or Z<sub>3</sub>;

R<sub>11</sub> and R<sub>12</sub> are each independently -H or -C<sub>1-6</sub>alkyl; and

Z<sub>3</sub> is heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl in which the heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl contains, in the ring, at least one selected from the group consisting

of N, O, P, P (=O) and S, and at least one H of the heterocycloalkyl, heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or cycloalkyl.

3. The compound represented by Chemical Formula I, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of claim 1, wherein

X<sub>1</sub> is N;  
X<sub>2</sub> and X<sub>3</sub> are each independently CH or N; and  
R<sub>X</sub> is -H, -NH<sub>2</sub>, or -NH(-C<sub>1-6</sub>alkyl).

4. The compound represented by Chemical Formula I, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of claim 1, wherein

Y is -C<sub>1-6</sub>alkyl, -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl in which at least one H of the -(CH<sub>2</sub>)<sub>n</sub>aryl, -(CH<sub>2</sub>)<sub>n</sub>heteroaryl, or -(CH<sub>2</sub>)<sub>n</sub>hydroheteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkynyl, -CN, -(C=O)NH-cycloalkyl, -(C=O)O-C<sub>1-6</sub>alkyl, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -O(C<sub>1-6</sub>alkyl), -O-phenyl, -halo, =O, heterocycloalkyl, aryl, or heteroaryl, wherein at least one H of the heterocycloalkyl, aryl, or heteroaryl may be substituted with -halo;

n is 0 or 1; and

R<sub>Y1</sub> to R<sub>Y5</sub> are each independently -H or -C<sub>1-6</sub>alkyl, or R<sub>Y2</sub> and R<sub>Y3</sub> may be linked to each other to form 3- to 6-membered cycloalkyl, and R<sub>Y3</sub> and R<sub>Y4</sub> may be linked to each other to form phenyl.

5. The compound represented by Chemical Formula I, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of claim 1, wherein

L is -(CH<sub>2</sub>)<sub>m</sub>-, -C(=O)-, or null;  
m is 0 or 1;

a ring Z is aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl in which at least one H of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>cyanoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -CN, -NR<sub>7</sub>R<sub>8</sub>, -O-C<sub>1-6</sub>alkyl, -O-C<sub>1-6</sub>haloalkyl, -S-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>alkyl, -C(=O)-C<sub>1-6</sub>haloalkyl, -C(=O)O-C<sub>1-6</sub>alkyl, -S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, -C(=N-O-C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), =O, -halo, or Z<sub>1</sub>, wherein at least one H of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl or -halo, or two or more substituents of the aryl, heteroaryl, hydroheteroaryl, 3- to 7-membered cycloalkyl, or 5- to 7-membered heterocycloalkyl ring may be linked to each other to form a fused ring or a spiro ring, wherein at least one H of the fused ring or spiro ring may be substituted with -C<sub>1-6</sub>alkyl or Z<sub>1</sub>;

R<sub>7</sub> and R<sub>8</sub> are each independently -H, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -(C=O)-C<sub>1-6</sub>alkyl, or -(C=O)-C<sub>1-6</sub>haloalkyl;

Z<sub>1</sub> is 3- to 7-membered cycloalkyl, 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, P, P(=O) and S, and at least one H of the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, 6- to 10-membered heterospiroalkyl, -C<sub>1-6</sub>alkyl-heterocycloalkyl, -(C=O)-heterocycloalkyl, -NH-heterocycloalkyl, or heteroaryl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>aminoalkyl, -C<sub>1-6</sub>haloalkyl, -C<sub>1-6</sub>alkenyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), -halo, cycloalkyl, or Z<sub>2</sub>;

Z<sub>2</sub> is 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, and S, and at least one H of the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -NH-heterocycloalkyl ring may be substituted with -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>hydroxyalkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -C(=O)-C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkyl-S(=O)<sub>2</sub>-C<sub>1-6</sub>alkyl, =O, -N(C<sub>1-6</sub>alkyl)(C<sub>1-6</sub>alkyl), 3- to 7-membered cycloalkyl, or Z<sub>3</sub>; and

Z<sub>3</sub> is 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl in which the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl contains, in the ring, at least one selected from the group consisting of N, O, and S, and at least one H of the 5- to 7-membered heterocycloalkyl, 6- to 10-membered heterobicycloalkyl, or -C<sub>1-6</sub>alkyl-heterocycloalkyl



ring may be substituted with -C<sub>1-6</sub>alkyl or 3- to 7-membered cycloalkyl.

6. The compound represented by Chemical Formula I, the optical isomer thereof, or the pharmaceutically acceptable salt thereof of claim 1, wherein the compound represented by the Chemical Formula 1 is selected from the group consisting of the following compounds:

- (1) (R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (2) (S)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (3) (S)-6-(3-benzylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (4) (R)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (5) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (6) (R)-N-(4-(4-(4-cyclopropylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (7) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-morpholinophenyl)pyrimidin-4-amine;
- (8) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (9) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-5-yl)-4-morpholinophenyl)pyrimidin-4-amine;
- (10) (R)-1'-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-methoxy-2-(1-methyl-1H-pyrazol-4-yl)phenyl)-N,N-dimethyl-[1,4'-bipiperidin]-4-amine;
- (11) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-5-(methylsulfonyl)phenyl)pyrimidin-4-amine;
- (12) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(methylsulfonyl)-5-(4-morpholinopiperidin-1-yl)phenyl)pyrimidin-4-amine;
- (13) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-(4-methylpiperazin-1-yl)-5-(methylsulfonyl)phenyl)pyrimidin-4-amine;
- (14) (R)-2-(3-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-((4-methylpiperazin-1-yl)methyl)phenyl)-2-methylpropanenitrile;
- (15) (R)-2-(3-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-(4-methylpiperazin-1-yl)phenyl)-2-methylpropanenitrile;
- (16) (R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-(3-phenoxyphenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (17) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (18) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (19) (R)-N<sup>1</sup>-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-N<sup>4</sup>-(2-(dimethylamino)ethyl)-N<sup>4</sup>-methylbenzene-1,4-diamine;
- (20) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (21) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (22) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(dimethylamino)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (23) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-3-(trifluoromethyl)phenyl)pyrimidin-4-amine;
- (24) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-3-(methylsulfonyl)phenyl)pyrimidin-4-amine;
- (25) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (26) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (27) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(3-(trifluoromethyl)phenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (28) (R)-6-(3-(2-fluoro-3-(trifluoromethyl)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (29) (R)-6-(3-(3-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phe-

- nyl)pyrimidin-4-amine;
- (30) (R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 5 (31) (R)-6-(3-(3-chloro-4-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (32) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (33) (R)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 10 (34) (R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (35) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(thiophen-2-yl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (36) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(naphthalen-1-yl)isoxazolidin-2-yl)pyrimidin-4-amine;
- 15 (37) (R)-6-(3-(3-ethynylphenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (38) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(6-methylpyridin-3-yl)isoxazolidin-2-yl)pyrimidin-4-amine;
- 20 (39) (R)-6-(3-(3-([1,2,4]triazolo[1,5-a]pyridin-7-yl)phenyl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (40) (R)-6-(3-(3-fluoro-5-thiomorpholinophenyl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (41) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(3-methoxyphenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- 25 (42) (R)-6-(3-methyl-3-phenylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (43) isopropyl (R)-3-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzoate;
- (44) (R)-N-cyclohexyl-3-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzamide;
- 30 (45) (R)-6-(3-(3',5'-difluoro-[1,1'-biphenyl]-3-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (46) (R)-N,N-dimethyl-7-(2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-7-azaspiro[3.5]nonan-2-amine;
- (47) (R)-N,N-dimethyl-2-(2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-2-azaspiro[3.5]nonan-7-amine;
- 35 (48) (R)-3-(1-methylpiperidin-4-yl)-N-(6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-1,2,3,4,4a,5-hexahydrobenzo[b]pyrazino[1,2-d][1,4]oxazin-8-amine;
- (49) (6aR,8S)-8-(4-cyclopropylpiperazin-1-yl)-2-methoxy-N-(6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-6,6a,7,8,9,10-hexahydrobenzo[b]pyrido[1,2-d][1,4]oxazin-3-amine;
- 40 (50) N-(4-(4-(5-ethylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (51) (R)-N-(2-methoxy-4-(4-((1-methylpiperidin-4-yl)amino)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (52) (R)-N-(2-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 45 (53) (R)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (54) (R)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (55) (R)-N-(2-ethoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 50 (56) N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-2-(((R)-1,1,1-trifluoropropan-2-yl)oxy)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (57) N-(2-methoxy-4-(4-((1R,4R)-5-methyl-2,5-diazabicyclo[2.2.1]heptan-2-yl)piperidin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 55 (58) (R)-N-(2-methoxy-4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (59) N-(4-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;

- (60) (R)-1-(4-(1-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperidin-4-yl)piperazin-1-yl)ethan-1-one;
- (61) (R)-N-(4-(4-(4-cyclopropyl-3,3-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 5 (62) (R)-N-(2-methoxy-4-(4-(pyrrolidin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (63) (R)-N-(4-(4-(4-cyclopropylpiperazin-1-yl)-[1,4'-bipiperidin]-1'-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (64) (R)-N-(4-(2-(dimethylamino)ethoxy)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 10 (65) (R)-N-(4-((2-(dimethylamino)ethyl)thio)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (66) (R)-N-(2-methoxy-4-thiomorpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (67) (R)-N-(2-methoxy-4-(4-methyl-1,4-diazepan-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (68) (R)-N-(4-(4-allylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 15 (69) N-(4-((R)-3-(dimethylamino)pyrrolidin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (70) N-(4-((S)-2-((dimethylamino)methyl)pyrrolidin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (71) (R)-N-(1-(1-(2-(dimethylamino)ethyl)piperidin-4-yl)-1H-pyrazol-4-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 20 (72) (R)-(4-methylpiperazin-1-yl)(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)methanone;
- (73) (R)-6-(3-phenylisoxazolidin-2-yl)-N-(3-(trifluoromethyl)phenyl)pyrimidin-4-amine;
- (74) N-(5-((1S,4S)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-2-fluoropyridin-3-yl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (75) (R)-1-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)pyrrolidin-2-one;
- 25 (76) (R)-2-methyl-5-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)isoindolin-1-one;
- (77) (R)-6-(3-phenylisoxazolidin-2-yl)-N-(4-(piperazin-1-yl)phenyl)pyrimidin-4-amine;
- (78) (R)-N-(6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)-1,2,3,4-tetrahydroisoquinolin-6-amine;
- (79) (R)-N-(5-(4-(dimethylamino)piperidin-1-yl)pyridin-2-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (80) (R)-N-(8-(1-methyl-1H-pyrazol-5-yl)-2,3-dihydrobenzo[b][1,4]dioxin-5-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 30 (81) (R)-1-cyclopropyl-4-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-1,4-azaphosphinane 4-oxide;
- (82) N-(4-((2R,6S)-2,6-dimethylmorpholino)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (83) 6-((R)-3-phenylisoxazolidin-2-yl)-N-((R)-7-(pyrrolidin-1-yl)-6,7,8,9-tetrahydro-5H-benzo[7]annulen-2-yl)pyrimidin-4-amine;
- 35 (84) (R)-N,N-dimethyl-1'-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-[1,4'-bipiperidin]-4-amine;
- (85) (R)-N-(5-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-2-(4-(dimethylamino)-[1,4'-bipiperidin]-1'-yl)-4-methoxyphenyl)propionamide;
- 40 (86) 3-chloro-N-(5-((6-((R)-3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-2-(4-(hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)piperidin-1-yl)-4-methoxyphenyl)propanamide;
- (87) (R)-N-(5-((6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)-[1,4'-bipiperidin]-1'-yl)phenyl)propionamide;
- (88) (R)-7-(5-methoxy-2-methyl-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-N,N-dimethyl-7-azaspiro[3.5]nonan-2-amine;
- 45 (89) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(naphthalen-2-yl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (90) (R)-6-(3-(3,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 50 (91) (R)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (92) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (93) (R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 55 (94) (R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (95) (R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-

- yl)phenyl)pyrimidin-4-amine;
- (96) (R)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (97) N-(4-(4-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (98) (R)-N-(4-(4-(diethylamino)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (99) N-(2-methoxy-4-((R)-2-methyl-4-(1-methylpiperidin-4-yl)piperazin-1-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (100) isopropyl (R)-3-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzoate;
- (101) (R)-N-(2-methoxy-4-(4-(1-methylpiperidin-4-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (102) (R)-N-(3-methoxy-4-(4-(1-methylpiperidin-4-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (103) (R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (104) (R)-6-(3-(2,4-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (105) (R)-6-(3-(4-chloro-2-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (106) (R)-6-(3-(2,5-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (107) (R)-N-(3-methyl-4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (108) (R)-N-(2-methyl-4-morpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (109) (R)-N-(5-ethyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (110) (R)-N-(2-methoxy-4-morpholinophenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (111) (R)-N4-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-2,4-diamine;
- (112) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-4-(3-phenylisoxazolidin-2-yl)-1,3,5-triazin-2-amine;
- (113) (R)-2-methoxy-N4-(1-methylpiperidin-4-yl)-N1-(6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)benzene-1,4-diamine;
- (114) (R)-N-(2-methoxy-5-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (115) (R)-1'-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)-N,N-dimethyl-[1,4'-bipiperidin]-4-amine;
- (116) N-(4-(4-((1R,4R)-5-ethyl-2,5-diazabicyclo[2.2.1]heptan-2-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (117) (R)-N-(3-ethyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (118) (R)-N-(4-(4-(4-ethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (119) (R)-N-(4-(4-(dimethylamino)piperidin-1-yl)-2-methoxyphenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (120) (R)-N-(2-methoxy-4-(4-morpholinopiperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (121) (R)-2-(4-(1-(2-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperidin-4-yl)piperazin-1-yl)ethan-1-ol;
- (122) (R)-1-(4-(4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)piperazin-1-yl)ethan-1-one;
- (123) (R)-N-(3,5-difluoro-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (124) (R)-N-(4-(4-(oxetan-3-yl)piperazin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (125) (R)-6-(3-(3-chloro-2-fluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (126) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (127) (R)-N-(4-(4-(dimethylamino)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;

- (128) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (129) (R)-6-(3-(4-chloro-3-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 5 (130) (R)-6-(3-(3-chloro-2,4-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (131) (R)-6-(3-(3-(dimethylamino)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 10 (132) (R)-N-(5-chloro-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (133) (R)-N-(1-methyl-1H-pyrazol-4-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (134) tert-butyl (R)-7-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)-3,4-dihydroisoquinoline-2(1H)-carboxylate;
- 15 (135) (R)-N-(4-(9-methyl-3,9-diazaspiro[5.5]undecan-3-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (136) N-(4-(6-methyl-3,6-diazabicyclo[3.1.1]heptan-3-yl)phenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (137) (R)-3-(3-methoxy-4-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)oxazolidin-2-one;
- 20 (138) (R)-N-(5-(4-(dimethylamino)piperidin-1-yl)pyridin-2-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (139) (R)-N-(4-(4-methylpiperazin-1-yl)-3-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (140) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(3-fluoro-4-(4-methylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- 25 (141) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(1-methylpiperidin-4-yl)phenyl)pyrimidin-4-amine;
- (142) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-isopropyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (143) (R)-N-(5-cyclopropyl-2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 30 (144) (R)-N-(4-(4-methylpiperazin-1-yl)-2-(3-((methylsulfonyl)methyl)azetidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (145) (R)-1-(5-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)ethan-1-one;
- 35 (146) (R,E)-1-(5-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-4-methoxy-2-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)ethan-1-one O-methyl oxime;
- (147) (R)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-6-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)pyridin-3-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (148) (R)-6-(3-(4-fluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 40 (149) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (150) (R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-5'-methoxyspiro[cyclopropane-1,3'-indoline]-6'-amine;
- 45 (151) (R)-6'-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5'-methoxyspiro[cyclopropane-1,3'-indolin]-2'-one;
- (152) (R)-1-(6-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-methoxy-1'-methylspiro[indolin-3,4'-piperidin]-1-yl)-2,2,2-trifluoroethan-1-one;
- (153) (R)-6-((6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-7-methoxyspiro[benzo[b][1,4]oxazin-2,1'-cyclopropane]-3(4H)-one;
- 50 (154) (R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-5-methoxy-1'-methylspiro[indolin-3,4'-piperidin]-6-amine;
- (155) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine;
- 55 (156) (R)-N-(6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)-3-methoxy-7-(4-methylpiperazin-1-yl)-9H-carbazol-2-amine;
- (157) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine;
- (158) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-methyl-1H-pyrazol-5-yl)-2,3-dihydrobenzofuran-7-

- yl)pyrimidin-4-amine;
- (159) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine;
- (160) (R)-6-(3-(3,5-difluorophenyl)isoxazolidin-2-yl)-N-(4-(1-(1-methylpiperidin-4-yl)-1H-pyrazol-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-4-amine;
- 5 (161) (R)-N-(4-methoxy-2-(4-methylpiperazin-1-yl)-5-((6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)propionamide;
- (162) (R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-methylpiperazin-1-yl)-2,3-dihydrobenzofuran-7-yl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (163) (R)-N-(2-(4-(4-cyclopropylpiperazin-1-yl)piperidin-1-yl)-4-methoxy-5-((6-(3-(thiophene-2-yl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)propionamide;
- 10 (164) (R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (165) (R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-morpholinopiperidin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- 15 (166) (R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-(4-methylpiperazin-1-yl)-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (167) (R)-N-(5-(1-methyl-1H-pyrazol-4-yl)-4-morpholino-2-(2,2,2-trifluoroethoxy)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (168) (S)-6-(3-(5-fluoropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 20 (169) (R)-6-(3-(5-fluoropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (170) (R)-3-fluoro-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzotrile;
- 25 (171) (S)-3-fluoro-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)benzotrile;
- (172) tert-butyl (R)-3-(4-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate;
- (173) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((R)-2-methylisoxazolidin-3-yl)phenyl)pyrimidin-4-amine;
- 30 (174) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((R)-isoxazolidin-3-yl)phenyl)pyrimidin-4-amine;
- (175) tert-butyl (S)-3-(4-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate;
- (176) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((S)-2-methylisoxazolidin-3-yl)phenyl)pyrimidin-4-amine;
- 35 (177) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-((S)-isoxazolidin-3-yl)phenyl)pyrimidin-4-amine;
- (178) (R)-6-(3-(1-methyl-1H-pyrazol-4-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (179) (R)-6-(3-(furan-2-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- 40 (180) (R)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (181) (S)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;
- (182) (R)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (183) (S)-6-(3-(5-chloropyridin-3-yl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 45 (184) (R)-6-(3-(3-(difluoromethyl)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (185) (S)-6-(3-(3-(difluoromethyl)phenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- 50 (186) tert-butyl (R)-3-(3-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate;
- (187) tert-butyl (S)-3-(3-((6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)phenyl)isoxazolidin-2-carboxylate;
- (188) (R)-N-(6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)cyclopropanecarboxamide;
- 55 (189) (R)-N-(6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)benzamide;
- (190) (R)-N-(cyclopentylmethyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (191) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-fluorobenzyl)pyrimidin-4-amine;
- (192) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(1-(methylsulfonyl)piperidin-4-yl)pyrimidin-4-amine;

- (193) (R)-6-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-5-methoxy-2-methylisoindolin-1-one;
- (194) (R)-3-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-1H-indazol-6-carbonitrile;
- (195) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(9-(1-fluoro-2-methylpropan-2-yl)-3,9-diazaspiro[5.5]undecan-3-yl)-2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)phenyl)pyrimidin-4-amine;
- (196) (R)-N-(2,5-dichloro-4-(9-(1-fluoro-2-methylpropan-2-yl)-3,9-diazaspiro[5.5]undecan-3-yl)phenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (197) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methyl-5-(3-((methylsulfonyl)methyl)azetidin-1-yl)-4-(4-morpholinopiperidin-1-yl)phenyl)pyrimidin-4-amine;
- (198) (R)-N-(5-(3-chloro-1-methyl-1H-pyrazol-4-yl)-2-methoxy-4-(4-methylpiperazin-1-yl)phenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (199) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-methylpiperazin-1-yl)-5-vinylphenyl)pyrimidin-4-amine;
- (200) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(5-ethynyl-2-methoxy-4-(4-methylpiperazine-1-yl)phenyl)pyrimidin-4-amine;
- (201) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-methoxy-6-(1-methyl-1H-pyrazol-4-yl)-5-(4-methylpiperazin-1-yl)pyridin-2-yl)pyrimidin-4-amine;
- (202) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-5-(1-methyl-1H-pyrazol-4-yl)-6-(4-methylpiperazin-1-yl)pyridin-3-yl)pyrimidin-4-amine;
- (203) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(3-(1-methyl-1H-pyrazol-4-yl)-5-((2R,3S)-2-methyl-3-((methylsulfonyl)methyl)azetidin-1-yl)-4-morpholinophenyl)pyrimidin-4-amine;
- (204) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-N2-methylpyrimidin-2,4-diamine;
- (205) (R)-4-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)pyridin-2-amine;
- (206) (R)-N-(4-(4-(4-allylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (207) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-ethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (208) (R)-N-(4-(4-(4-cyclobutylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (209) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-(oxetan-3-yl)piperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (210) (R)-N-(4-(4-(4-(cyclopropylmethyl)piperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-amine;
- (211) (R)-N-(2-methoxy-4-(4-(6-methyl-2,6-diazaspiro[3.3]heptan-2-yl)piperidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (212) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-3-(dimethylamino)pyrrolidin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (213) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-3,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (214) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((S)-3,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (215) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((3S,5R)-4-ethyl-3,5-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (216) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((S)-2,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (217) 6-((R)-3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-((R)-2,4-dimethylpiperazin-1-yl)piperidin-1-yl)-2-methoxyphenyl)pyrimidin-4-amine;
- (218) (R)-N-(2-methoxy-4-(3-(4-methylpiperazin-1-yl)azetidin-1-yl)phenyl)-6-(3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (219) N-(4-(4-(6-ethyl-3,6-diazabicyclo[3.1.1]heptan-3-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (220) N-(4-(4-(3-ethyl-3,6-diazabicyclo[3.1.1]heptan-6-yl)piperidin-1-yl)-2-methoxyphenyl)-6-((R)-3-phenylisoxazolidin-2-yl)pyrimidin-4-amine;
- (221) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(4-methyl-1,4-diazepan-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine;
- (222) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(4-(4-(4-isopropylpiperazin-1-yl)piperidin-1-yl)-2-meth-

oxyphenyl)pyrimidin-4-amine;

(223) (R)-1-(1-(4-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-3-methoxyphenyl)piperidin-4-yl)-4-methylpiperazin-2-one;

(224) (R)-4-(1-(4-((6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)pyrimidin-4-yl)amino)-3-methoxyphenyl)piperidin-4-yl)-1-methylpiperazin-2-one;

(225) (R)-1-methyl-5-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)pyridin-2(1H)-one;

(226) (R)-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)-1-methylpyridin-2(1H)-one;

(227) (S)-1-methyl-5-(2-(6-((4-(4-methylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)pyridin-2(1H)-one;

(228) (S)-5-(2-(6-((2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)amino)pyrimidin-4-yl)isoxazolidin-3-yl)-1-methylpyridin-2(1H)-one;

(229) N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(3-phenylbenzo[d]isoxazol-2(3H)-yl)pyrimidin-4-amine;

(230) N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(3-phenylbenzo[d]isoxazol-2(3H)-yl)pyrimidin-4-amine;

(231) (S)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine;

(232) (R)-N-(4-(4-methylpiperazin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine;

(233) (S)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine; and

(234) (R)-N-(2-methoxy-4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-6-(7-phenyl-5-oxa-6-azaspiro[2.4]heptan-6-yl)pyrimidin-4-amine.

(235) (R)-6-(3-isopropylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine;

(236) (S)-6-(3-isopropylisoxazolidin-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)pyrimidin-4-amine; and

(237) (R)-6-(3-(2,3-difluorophenyl)isoxazolidin-2-yl)-N-(2-methoxy-4-(4-(2-(methylsulfonyl)ethyl)piperazin-1-yl)piperidin-1-yl)phenyl)pyrimidin-4-amine.

7. A pharmaceutical composition for the treatment or prevention of cancer, comprising, as an active ingredient, the compound according to any one of claims 1 to 6, the optical isomer thereof, or the pharmaceutically acceptable salt thereof.

8. The pharmaceutical composition of claim 7, wherein the composition inhibits EGFR and/or HER2.

9. The pharmaceutical composition of claim 8, wherein the composition inhibits any one or more selected from the group consisting of

EGFR Del19/C797S, EGFR L858R/C797S, EGFR L861Q, EGFR G719A, EGFR S768I, EGFR L718Q, EGFR G724S, EGFR d746-750, EGFR d746-750/C797A, EGFR d746-750/C797S, EGFR d746-750/T790M/C797S, EGFR D761Y, EGFR G719C, EGFR G719D, EGFR G719S, EGFR L747S, EGFR L792F, EGFR L858R, EGFR L792F/L858R, ERBB2/HER2, and ERBB4/HER4.

10. The pharmaceutical composition of claim 7, wherein the cancer is one or more selected from the group consisting of pseudomyxoma, intrahepatic biliary tract cancer, hepatoblastoma, liver cancer, thyroid cancer, colon cancer, testicular cancer, myelodysplastic syndrome, glioblastoma, oral cancer, labial cancer, mycosis fungoides, acute myeloid leukemia, acute lymphocytic leukemia, basal cell cancer, ovarian epithelial cancer, ovarian germ cell cancer, male breast cancer, brain cancer, pituitary adenoma, multiple myeloma, gallbladder cancer, biliary tract cancer, colorectal cancer, chronic myelogenous leukemia, chronic lymphocytic leukemia, retinoblastoma, choroidal melanoma, ampulla of Vater cancer, bladder cancer, peritoneal cancer, parathyroid cancer, adrenal cancer, sinus cancer, non-small cell lung cancer, tongue cancer, astrocytoma, small cell lung cancer, pediatric brain cancer, pediatric lymphoma, pediatric leukemia, small intestine cancer, meningioma, esophageal cancer, glioma, renal pelvic cancer, kidney cancer, heart cancer, duodenal cancer, malignant soft tissue cancer, malignant bone cancer, malignant lymphoma, malignant mesothelioma, malignant melanoma, eye cancer, vulvar cancer, ureter cancer, urethral cancer, cancer of unknown primary site, gastric lymphoma, gastric cancer, gastric carcinoma, gastrointestinal stromal cancer, Wilms cancer, breast cancer, sarcoma, penile cancer, pharyngeal cancer, gestational choriocarcinoma, cervical cancer, endometrial cancer, uterine sarcoma, prostate cancer, metastatic bone cancer, metastatic brain cancer, mediastinal cancer, rectal cancer, rectal carcinoma, vaginal cancer, spinal cord cancer, acoustic tumor, pancreatic cancer, salivary gland cancer, Kaposi's sarcoma, Paget's disease, tonsil cancer, squamous cell carcinoma, lung



**EP 4 342 895 A1**

adenocarcinoma, lung cancer, lung squamous cell carcinoma, skin cancer, anal cancer, rhabdomyosarcoma, laryngeal cancer, pleural cancer, blood cancer, and thymus cancer.

- 5
11. A method for treating or preventing EGFR- and/or HER2-related diseases, comprising administering to a subject in need thereof a therapeutically effective amount of the compound, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to any one of claims 1 to 10.
- 10
12. Use of the compound, the optical isomer thereof, or the pharmaceutically acceptable salt thereof according to any one of claims 1 to 10 for use in preparation of a medicament to treat or prevent EGFR- and/or HER2-related diseases.

15

20

25

30

35

40

45

50

55

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2022/006994

5

**A. CLASSIFICATION OF SUBJECT MATTER**  
**C07D 413/04**(2006.01); **A61K 31/506**(2006.01); **A61K 31/5377**(2006.01); **A61K 31/541**(2006.01); **A61P 35/00**(2006.01); **C07D 413/14**(2006.01); **C07D 471/04**(2006.01)  
 According to International Patent Classification (IPC) or to both national classification and IPC

10

**B. FIELDS SEARCHED**  
 Minimum documentation searched (classification system followed by classification symbols)  
 C07D 413/04(2006.01); A61K 31/136(2006.01); A61K 31/404(2006.01); A61K 31/506(2006.01); C07D 403/04(2006.01); C07D 413/14(2006.01); C07D 487/04(2006.01); C07D 491/107(2006.01); C07D 495/04(2006.01)

15

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
 Korean utility models and applications for utility models: IPC as above  
 Japanese utility models and applications for utility models: IPC as above  
 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
 eKOMPASS (KIPO internal), STN (Registry, Caplus), Google & keywords: 상피 성장인자 수용체(epidermal growth factor receptor, EGFR), 암(cancer, tumor), 피리미딘(pyrimidine), 트리아진(triazine)

20

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KR 10-2020-0111644 A (VORONOI INC. et al.) 29 September 2020 (2020-09-29) See abstract; claims 1 and 12-15; and table 1.	1-10,12
A	KR 10-2017-0066650 A (YUHAN CORPORATION) 14 June 2017 (2017-06-14) See claims 1-20.	1-10,12
A	WO 2020-235902 A1 (VORONOI CO., LTD. et al.) 26 November 2020 (2020-11-26) See claims 1-14.	1-10,12
A	WO 2019-010295 A1 (CS PHARMATECH LIMITED) 10 January 2019 (2019-01-10) See entire document.	1-10,12
A	US 2017-0313714 A1 (SHANGHAI HANSOH BIOMEDICAL CO., LTD. et al.) 02 November 2017 (2017-11-02) See entire document.	1-10,12

35

Further documents are listed in the continuation of Box C.  See patent family annex.

\* Special categories of cited documents:  
 "A" document defining the general state of the art which is not considered to be of particular relevance  
 "D" document cited by the applicant in the international application  
 "E" earlier application or patent but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed  
 "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 "&" document member of the same patent family

40

45

Date of the actual completion of the international search <b>29 August 2022</b>	Date of mailing of the international search report <b>29 August 2022</b>
--	---

50

Name and mailing address of the ISA/KR <b>Korean Intellectual Property Office Government Complex-Daejeon Building 4, 189 Cheongsaro, Seo-gu, Daejeon 35208</b> Facsimile No. +82-42-481-8578	Authorized officer  Telephone No.
--	---

55

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/KR2022/006994

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: **11**  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claim 11 pertains to a method for treatment of the human body by surgery or therapy, as well as a diagnostic method (PCT Article 17(2)(a)(i) and PCT Rule 39.1(iv)).
2.  Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

INTERNATIONAL SEARCH REPORT  
Information on patent family members

International application No.

PCT/KR2022/006994

Patent document cited in search report	Publication date (day/month/year)	Patent family member(s)	Publication date (day/month/year)
KR 10-2020-0111644 A	29 September 2020	AU 2020-240382 A1	11 November 2021
		CA 3134261 A1	24 September 2020
		EP 3943491 A1	26 January 2022
		JP 2022-522250 A	14 April 2022
		KR 10-2021-0130123 A	29 October 2021
		US 2022-0073505 A1	10 March 2022
		US 2022-0162203 A1	26 May 2022
		WO 2020-190119 A1	24 September 2020
KR 10-2017-0066650 A	14 June 2017	AU 2015-331166 A1	30 March 2017
		AU 2015-331166 B2	26 April 2018
		CN 106795144 A	31 May 2017
		CN 106795144 B	16 June 2020
		CN 111686110 A	22 September 2020
		EP 3207035 A2	23 August 2017
		EP 3207035 B1	20 November 2019
		EP 3604294 A1	05 February 2020
		EP 3604294 B1	05 May 2021
		EP 3929190 A1	29 December 2021
		JP 2017-530999 A	19 October 2017
		JP 2019-163277 A	26 September 2019
		JP 2020-196740 A	10 December 2020
		JP 6524221 B2	05 June 2019
		JP 6754864 B2	16 September 2020
		KR 10-2021-0011068 A	29 January 2021
		KR 10-2208775 B1	28 January 2021
WO 2020-235902 A1	26 November 2020	US 2016-0102076 A1	14 April 2016
		US 9593098 B2	14 March 2017
		WO 2016-060443 A2	21 April 2016
		WO 2016-060443 A3	23 June 2016
		EP 3971190 A1	23 March 2022
		KR 10-2020-0133188 A	26 November 2020
WO 2019-010295 A1	10 January 2019	CA 3068854 A1	10 January 2019
		CN 111093645 A	01 May 2020
		EP 3648753 A1	13 May 2020
		JP 2020-526499 A	31 August 2020
		KR 10-2020-0028966 A	17 March 2020
		US 2020-0131176 A1	30 April 2020
US 2017-0313714 A1	02 November 2017	AU 2015-330506 A1	06 April 2017
		AU 2015-330506 B2	30 January 2020
		CN 106661000 A	10 May 2017
		CN 106661000 B	09 April 2019
		CN 111170999 A	19 May 2020
		CN 111171000 A	19 May 2020
		CN 111187221 A	22 May 2020
		EP 3205650 A1	16 August 2017
		EP 3205650 B1	04 August 2021
		JP 2017-532326 A	02 November 2017
		JP 6691281 B2	28 April 2020
		KR 10-2017-0118681 A	25 October 2017
		US 10259820 B2	16 April 2019

Form PCT/ISA/210 (patent family annex) (July 2019)

INTERNATIONAL SEARCH REPORT  
Information on patent family members

International application No. <b>PCT/KR2022/006994</b>
---

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Patent document cited in search report	Publication date (day/month/year)	Patent family member(s)	Publication date (day/month/year)
		US 10428081 B2	01 October 2019
		US 2019-0100528 A1	04 April 2019
		WO 2016-054987 A1	14 April 2016
		ZA 201701465 B	28 August 2019
.....			

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Non-patent literature cited in the description**

- **NIEDERST MJ et al.** *Clin Cancer Res*, 2015, vol. 17 (21), 3924-3933 **[0004]**
- **THRESS KS et al.** *Nature Medicine*, 2015, vol. 21, 560-562 **[0004]**
- **DI FIORE PP et al.** *Science*, 1987, vol. 237 (481), 178-182 **[0006]**
- **HARDWICK RH et al.** *Eur. J Surg Oncol.*, 1997, vol. 23 (1), 30-35 **[0006]**
- **KORKAYA H. et al.** *Oncogene*, 2008, vol. 27 (47 **[0006]**