



US 20230416700A1

(19) **United States**

(12) **Patent Application Publication**  
**Wang et al.**

(10) **Pub. No.: US 2023/0416700 A1**

(43) **Pub. Date: Dec. 28, 2023**

(54) **COMPOSITIONS USEFUL IN TREATMENT OF ORNITHINE TRANSCARBAMYLASE (OTC) DEFICIENCY**

(60) Provisional application No. 61/950,157, filed on Mar. 9, 2014.

**Publication Classification**

(71) Applicant: **The Trustees of the University of Pennsylvania**, Philadelphia, PA (US)

(51) **Int. Cl.**  
*C12N 9/10* (2006.01)  
*C12N 15/86* (2006.01)  
*A61K 48/00* (2006.01)

(72) Inventors: **Lili Wang**, Phoenixville, PA (US);  
**James M. Wilson**, Philadelphia, PA (US)

(52) **U.S. Cl.**  
CPC .... *C12N 9/1018* (2013.01); *C12Y 201/03003* (2013.01); *C12N 15/86* (2013.01); *A61K 48/0058* (2013.01); *C12N 2750/14032* (2013.01); *C12N 2750/14071* (2013.01); *C12N 2750/14033* (2013.01); *C12N 2710/10343* (2013.01); *C12N 2740/15043* (2013.01); *C12N 2750/14143* (2013.01); *C12N 2800/22* (2013.01); *C12N 2750/14043* (2013.01)

(73) Assignee: **The Trustees of the University of Pennsylvania**, Philadelphia, PA (US)

(21) Appl. No.: **18/339,457**

(22) Filed: **Jun. 22, 2023**

**Related U.S. Application Data**

(63) Continuation of application No. 16/995,572, filed on Aug. 17, 2020, now Pat. No. 11,732,246, which is a continuation of application No. 16/191,709, filed on Nov. 15, 2018, now Pat. No. 10,781,430, which is a continuation of application No. 15/122,853, filed on Aug. 31, 2016, now Pat. No. 10,167,454, filed as application No. PCT/US2015/019513 on Mar. 9, 2015.

(57) **ABSTRACT**

Viral vectors comprising engineered hOTC DNA and RNA sequences are provided which when delivered to a subject in need thereof are useful for treating hyperammonemia, ornithine transcarbamylase deficiency and symptoms associated therewith. Also provided are methods of using hOTC for treatment of liver fibrosis and/or cirrhosis in OTCD patients by administering hOTC.

**Specification includes a Sequence Listing.**

FIG. 1A

1 atgctgttta atctgaggat cctgttaaac aatgcagctt ttagaaatgg tcacaacttc  
61 atggttcgaa attttcggtg tggacaacca ctacaaaata aagtgcagct gaagggccgt  
121 gaccttctca ctctaaaaaa ctttaccgga gaagaaatta aatatatgct atggctatca  
181 gcagatctga aatthaggat aaaacagaaa ggagagtatt tgcctttatt gcaaggaag  
241 tccttaggca tgatthttga gaaaagaagt actcgaacaa gattgtctac agaaacaggc  
301 tttgcacttc tgggaggaca tccttgthtt cttaccacac aagatattca tttgggtgtg  
361 aatgaaagtc tcacggacac ggcccgtgta ttgtctagca tggcagatgc agtattggct  
421 cgagtgtata aacaatcaga tttggacacc ctggctaaag aagcatccat cccaattatc  
481 aatgggctgt cagatthtga ccatcctatc cagatcctgg ctgattacct cacgctccag  
541 gaacactata gctctctgaa aggtcttacc ctcagctgga tcgggggatgg gaacaatatc  
601 ctgcactcca tcatgatgag cgcagcgaaa ttcggaatgc accttcaggc agctactcca  
661 aagggttatg agccggatgc tagtghtaac aagttggcag agcagtatgc caagagaat  
721 ggtaccaagc tghtgctgac aatgatcca ttggaagcag cgcattggagg caatgtatta  
781 attacagaca cttggataag catgggacaa gaagaggaga agaaaaagcg gctccaggct  
841 ttccaagggt accagggttac aatgaagact gctaaagttg ctgcctctga ctggacattt  
901 ttacactgct tgcccagaaa gccagaagaa gtggatgatg aagtctthta ttctctctga  
961 tcactagtgt tcccagaggc agaaaacaga aagtggacaa tcattggctgt catggtgtcc  
1021 ctgctgacag attactcacc tcagctccag aagcctaaat tt

FIG. 1B

Met Leu Phe Asn Leu Arg Ile Leu Leu Asn Asn Ala Ala Phe Arg Asn  
 1 5 10 15

Gly His Asn Phe Met Val Arg Asn Phe Arg Cys Gly Gln Pro Leu Gln  
 20 25 30

Asn Lys Val Gln Leu Lys Gly Arg Asp Leu Leu Thr Leu Lys Asn Phe  
 35 40 45

Thr Gly Glu Glu Ile Lys Tyr Met Leu Trp Leu Ser Ala Asp Leu Lys  
 50 55 60

Phe Arg Ile Lys Gln Lys Gly Glu Tyr Leu Pro Leu Leu Gln Gly Lys  
 65 70 75 80

Ser Leu Gly Met Ile Phe Glu Lys Arg Ser Thr Arg Thr Arg Leu Ser  
 85 90 95

Thr Glu Thr Gly Phe Ala Leu Leu Gly Gly His Pro Cys Phe Leu Thr  
 100 105 110

Thr Gln Asp Ile His Leu Gly Val Asn Glu Ser Leu Thr Asp Thr Ala  
 115 120 125

Arg Val Leu Ser Ser Met Ala Asp Ala Val Leu Ala Arg Val Tyr Lys  
 130 135 140

Gln Ser Asp Leu Asp Thr Leu Ala Lys Glu Ala Ser Ile Pro Ile Ile  
 145 150 155 160

Asn Gly Leu Ser Asp Leu Tyr His Pro Ile Gln Ile Leu Ala Asp Tyr  
 165 170 175

Leu Thr Leu Gln Glu His Tyr Ser Ser Leu Lys Gly Leu Thr Leu Ser  
 180 185 190

FIG. 1C

Trp Ile Gly Asp Gly Asn Asn Ile Leu His Ser Ile Met Met Ser Ala  
 195 200 205

Ala Lys Phe Gly Met His Leu Gln Ala Ala Thr Pro Lys Gly Tyr Glu  
 210 215 220

Pro Asp Ala Ser Val Thr Lys Leu Ala Glu Gln Tyr Ala Lys Glu Asn  
 225 230 235 240

Gly Thr Lys Leu Leu Leu Thr Asn Asp Pro Leu Glu Ala Ala His Gly  
 245 250 255

Gly Asn Val Leu Ile Thr Asp Thr Trp Ile Ser Met Gly Gln Glu Glu  
 260 265 270

Glu Lys Lys Lys Arg Leu Gln Ala Phe Gln Gly Tyr Gln Val Thr Met  
 275 280 285

Lys Thr Ala Lys Val Ala Ala Ser Asp Trp Thr Phe Leu His Cys Leu  
 290 295 300

Pro Arg Lys Pro Glu Glu Val Asp Asp Glu Val Phe Tyr Ser Pro Arg  
 305 310 315 320

Ser Leu Val Phe Pro Glu Ala Glu Asn Arg Lys Trp Thr Ile Met Ala  
 325 330 335

Val Met Val Ser Leu Leu Thr Asp Tyr Ser Pro Gln Leu Gln Lys Pro  
 340 345 350

Lys Phe

## FIG. 2

1 atgctgttca acctgcgaat cctgctgaac aatgccgctt ttcggaacgg gcacaatttc  
61 atggtgagga actttcgtcg eggacagccc ctccagaaca aggtccagct gaagggcagg  
121 gacctgctga ccctgaaaaa ttccacaggg gaggaaatca agtacatgct gtggctgtca  
181 gccgatctga agttccggat caagcagaag ggccaatc tgcctctgct ccagggcaaa  
241 agcctgggga tgatcttcga aaagcgcagt actcggacca gactgtcaac agagactgga  
301 ttcgcactgc tgggaggaca cccatgtttt ctgaccacac aggacattca tctgggagtg  
361 aacgagtccc tgaccgacac agcacgcgtc ctgagctcca tggctgatgc agtgctggct  
421 cgagtctaca aacagtctga cctggatacc ctggccaagg aagcttctat cccaatcatt  
481 aatggcctga gtgacctgta tcaccccatc cagattctgg ccgattacct gaccctccag  
541 gagcattatt ctagtctgaa agggctgaca ctgagctgga ttggggacgg aaacaatc  
601 ctgcactcca ttatgatgag cgccccaag tttggaatgc acctccaggc tgcaaccca  
661 aaaggctacg aacccgatgc ctccgtgaca aagctggcag aacagtatgc caaagagaac  
721 ggactaagc tgctgctgac caatgaccct ctggaggccg ctcacggagg caacgtgctg  
781 atcactgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccaggcc  
841 ttccagggt accaggtgac aatgaaaact gctaaggctc cagccagcga ctggacctt  
901 ctgcattgcc tgcccagaaa gcctgaagag gtggacgatg aggtcttcta ctcaccaga  
961 agcctggtgt ttctgaagc tgagaatagg aagtggacaa tcatggcagt gatggtcagc  
1021 ctgctgactg attattcccc tcagctccag aaaccaaagt tctgataa

FIG. 3

1 ctgcagccgc caccatgctg ttcaacctgc gaatcctgct gaacaacgcc gcttttcgga  
61 acgggcacaa ctttatgggtg aggaactttc gctgcggaca gcccctccag aataaggctc  
121 agctgaaggg cagggacctg ctgacctga aaaatttcac aggggaggaa atcaagtata  
181 tgctgtggct gtcagctgat ctgaagttcc ggatcaagca gaagggcgaa tatctgcctc  
241 tgctccaggg caaaagcctg gggatgatct tcgaaaagcg cagtactcgg accagactgt  
301 caaccgagac tggattcgct ctgctgggag gacacccttg ttttctgacc actcaggaca  
361 ttcacctggg agtgaacgag tccctgacct aactgctcg cgtcctgagc tctatggccg  
421 acgctgtgct agctcgagtc tacaacagc cgcacctgga taccctggcc aaggaagctt  
481 ctatcccaat tattaacggc ctgtcagacc tgtatcacc catccagatt ctggccgatt  
541 acctgacctt ccaggagcac tattctagtc tgaaagggct gacactgagt tggattgggg  
601 acggaaacaa tatcctgcac tctattatga tgtcagccgc caagtttggg atgcacctcc  
661 aggtgcaac cccaaaaggc tacgaacctg atgcctcagt gacaaagctg gctgaacagt  
721 acgcaaaga gaacggcact aagctgctgc tgaccaacga ccctctggag gccgctcag  
781 gaggcaacgt gctgatcacc gatacctgga ttagtatggg acaggaggaa gagaagaaga  
841 agcggctcca ggccttccag ggetaccagg tgacaatgaa aaccgctaag gtcgcagcca  
901 gcgattggac ctttctgcac tgctgcca gaaagcccga agaggtggac gacgaggtct  
961 tctactctcc cagaagcctg gtgtttcccg aagctgagaa taggaagtgg acaattatgg  
1021 cagtgatggc cagcctgctg actgattatt cacctcagct ccagaaacca aagttctgat  
1081 aagcggccgc

FIG. 4

```
1 ctgcagccgc caccatgctg ttcaacctgc gaatcctgct gaacaacgcc gcttttcgga
61 acgggcacaa ctttatgggtg aggaactttc gctgcgga gcccctccag aataaggctc
121 agctgaaggg cagggacctg ctgacctga aaaatttcac aggggaggaa atcaagtata
181 tgctgtggct gtcagctgat ctgaagttcc ggatcaagca gaagggcgaa tatctgcctc
241 tgctccaggg caaaagcctg gggatgatct tcgaaaagcg cagtactcgg accagactgt
301 caaccgagac tggattcgtc ctgctgggag gacacccttg ttttctgacc actcaggaca
361 ttcacctggg agtgaacgag tccctgaccg aactgctcg cgtcctgagc tctatggccg
421 acgctgtgct ggctcgagtc tacaacagct cgcacctgga taccctggcc aaggaagctt
481 ctatcccaat tattaacggc ctgtcagacc tgtatcacc catccagatt ctggccgatt
541 acctgaccct ccaggagcac tattctagtc tgaaagggct gacactgagt tggattgggg
601 acggaaacaa tatcctgcac tctattatga tgtcagccgc caagtttggg atgcacctcc
661 aggctgcaac cccaaaaggc tacgaaccgg atgcctcagt gacaaagctg gctgaacagt
721 acgccaaga gaacggcact aagctgctgc tgaccaacga ccctctggag gccgctcacg
781 gaggcaacgt gctgatcacc gatacctgga ttagtatggg acaggaggaa gagaagaaga
841 agcggtcca ggccttccag ggctaccagg tgacaatgaa aaccgctaag gtcgcagcca
901 gcgattggac ctttctgcac tgctgcca gaaagcccga agaggtggac gacgaggtct
961 tctactctcc cagaagcctg gtgtttcccg aagctgagaa taggaagtgg acaattatgg
1021 cagtgatggt cagcctgctg actgattatt cacctcagct ccagaaacca aagttctgat
1081 aagcgccgc
```

FIG. 5A

```

hOTCLW5 -----ATGCTGTTCAACCTGAGAATCCTGCTGAACAACGCCGCTTCAGAA 46
hOTCLW6 -----ATGCTGTTCAACCTGCGCATCCTGCTGAACAACGCCGCTTCCGCA 46
hOTCco -----ATGCTGTTCAACCTGCGAATCCTGCTGAACAATGCCGCTTTTCGGA 46
LW3 CTGCAGCCGCCACCATGCTGTTCAACCTGCGAATCCTGCTGAACAACGCCGCTTTTCGGA 60
LW4 CTGCAGCCGCCACCATGCTGTTCAACCTGCGAATCCTGCTGAACAACGCCGCTTTTCGGA 60
hOTwt -----ATGCTGTTAATCTGAGGATCCTGTTAAACAATGCAGCTTTTAGAA 46
          ***** ** ** * ***** * ***** ** ** * *

hOTCLW5 ACGGCCACAACCTTCATGGTGAGAACTTCAGATGCGGCCAGCCCCGTCAGAACAAGGTGC 106
hOTCLW6 ACGGCCACAACCTTCATGGTGCGCAACTTCCGCTGCGGCCAGCCCCGTCAGAACAAGGTGC 106
hOTCco ACGGGCACAAATTTTCATGGTGAGGAACCTTTCGCTGCGGACAGCCCCGTCAGAACAAGGTCC 106
LW3 ACGGGCACAACTTTATGGTGAGGAACCTTTCGCTGCGGACAGCCCCGTCAGAATAAGGTCC 120
LW4 ACGGGCACAACTTTATGGTGAGGAACCTTTCGCTGCGGACAGCCCCGTCAGAATAAGGTCC 120
hOTwt ATGGTCACAACCTTCATGGTTCGAAATTTTCGGTGTGACAACCCTACAAAATAAAGTGC 106
          * ** ***** ** ***** * ** ** * ** ** * ** ** * ** ** *

hOTCLW5 AGCTGAAGGGCAGAGACCTGCTGACCCTGAAGAACTTCACCGGCGAGGAGATCAAGTACA 166
hOTCLW6 AGCTGAAGGGCCGCGACCTGCTGACCCTGAAGAACTTCACCGGCGAGGAGATCAAGTACA 166
hOTCco AGCTGAAGGGCAGGGACCTGCTGACCCTGAAAAATTTACAGGGGAGGAAATCAAGTACA 166
LW3 AGCTGAAGGGCAGGGACCTGCTGACCCTGAAAAATTTACAGGGGAGGAAATCAAGTATA 180
LW4 AGCTGAAGGGCAGGGACCTGCTGACCCTGAAAAATTTACAGGGGAGGAAATCAAGTATA 180
hOTwt AGCTGAAGGGCCGTGACCTTCTCACTCTAAAAAACTTTACCGGAGAAGAAATTAATATA 166
          ***** * ***** ** ** ** * ** ** * ** ** * ** ** *

hOTCLW5 TGCTGTGGCTGAGCGCCGACCTGAAGTTCAGAATCAAGCAGAAGGGCGAGTACCTGCCCC 226
hOTCLW6 TGCTGTGGCTGAGCGCCGACCTGAAGTTCGCGCATCAAGCAGAAGGGCGAGTACCTGCCCC 226
hOTCco TGCTGTGGCTGTCAGCCGATCTGAAGTTCGCGATCAAGCAGAAGGGCGAATATCTGCCTC 226
LW3 TGCTGTGGCTGTCAGCTGATCTGAAGTTCGCGATCAAGCAGAAGGGCGAATATCTGCCTC 240
LW4 TGCTGTGGCTGTCAGCTGATCTGAAGTTCGCGATCAAGCAGAAGGGCGAATATCTGCCTC 240
hOTwt TGCTATGGCTATCAGCAGATCTGAAATTTAGGATAAAAACAGAAAGGAGATATTTGCCTT 226
          **** ***** ** ** ***** ** * ** ** ***** ** ** ** *****

hOTCLW5 TGCTGCAGGGCAAGAGCCTGGGCATGATCTTCGAGAAGAGAAGCACCAGAACCAGACTGA 286
hOTCLW6 TGCTGCAGGGCAAGAGCCTGGGCATGATCTTCGAGAAGCGCAGCACC CGCACCCGCCTGA 286
hOTCco TGCTCCAGGGCAAAAGCCTGGGGATGATCTTCGAAAAGCGCAGTACTCGGACCAGACTGT 286
LW3 TGCTCCAGGGCAAAAGCCTGGGGATGATCTTCGAAAAGCGCAGTACTCGGACCAGACTGT 300
LW4 TGCTCCAGGGCAAAAGCCTGGGGATGATCTTCGAAAAGCGCAGTACTCGGACCAGACTGT 300
hOTwt TATTGCAAGGGAAGTCTTAGGCATGATTTTTGAGAAAAGAAGTACTCGAACAAGATTGT 286
          * * ** ** ** * ** ** ***** ** ** ** * ** ** * ** ** *

hOTCLW5 GCACCGAGACCGGCCTGGCCCTGCTGGGCGGCCACCCCTGCTTCCCTGACCACCCAGGACA 346
hOTCLW6 GCACCGAGACCGGCCTGGCCCTGCTGGGCGGCCACCCCTGCTTCCCTGACCACCCAGGACA 346
hOTCco CAACAGAGACTGGATTTCGCACTGCTGGGAGGACACCCATGTTTTCTGACCACACAGGACA 346
LW3 CAACCGAGACTGGATTTCGCTCTGCTGGGAGGACACCCCTTGTTTTCTGACCACTCAGGACA 360
LW4 CAACCGAGACTGGATTTCGCTCTGCTGGGAGGACACCCCTTGTTTTCTGACCACTCAGGACA 360
hOTwt CTACAGAAACAGGCTTTGCACTTCTGGGAGGACATCCTTGTTTTCTTACCACACAAGATA 346
          ** ** ** ** * ** ** ***** ** ** ** * ** ** * ** ** *

hOTCLW5 TTCACCTGGGCGTGAACGAGAGCCTGACCGACACCCGAGAGTGTGAGCAGCATGGCCG 406
hOTCLW6 TTCACCTGGGCGTGAACGAGAGCCTGACCGACACCCGCGTGTGAGCAGCATGGCCG 406
hOTCco TTCATCTGGGAGTGAACGAGTCCCTGACCGACACAGCAGCGTCTGAGCTCCATGGCTG 406
LW3 TTCACCTGGGAGTGAACGAGTCCCTGACCGACACTGCTCGCGTCTGAGCTCTATGGCCG 420
LW4 TTCACCTGGGAGTGAACGAGTCCCTGACCGACACTGCTCGCGTCTGAGCTCTATGGCCG 420
hOTwt TTCATTTGGGTGTGAATGAAAGTCTCACGGACACGGCCCGTGTATTGTCTAGCATGGCAG 406
          * ** ***** ***** ** ** ** ***** ** * ** ** ***** *
    
```



FIG. 5B

```

hOTCLW5      ACGCCGTGCTGGCCAGAGTGTACAAGCAGAGCGACCTGGACACCCCTGGCCAAGGAGGCCA 466
hOTCLW6      ACGCCGTGCTGGCCCGCGTGTACAAGCAGAGCGACCTGGACACCCCTGGCCAAGGAGGCCA 466
hOTCco       ATGCAGTGTGCTGGCTCGAGTCTACAAAAGTCTGACCTGGATAACCCCTGGCCAAGGAAGCTT 466
LW3          ACGCTGTGCTAGCTCGAGTCTACAAAAGTCCGACCTGGATAACCCCTGGCCAAGGAAGCTT 480
LW4          ACGCTGTGCTGGCTCGAGTCTACAAAAGTCCGACCTGGATAACCCCTGGCCAAGGAAGCTT 480
hOTwt        ATGCAGTATTGGCTCGAGTGTATAACAATCAGATTGGACACCCCTGGCTAAAGAAGCAT 466
* * * * *
hOTCLW5      GCATCCCCATCATCAACGGCCTGAGCGACCTGTACCACCCCATCCAGATCCTGGCCGACT 526
hOTCLW6      GCATCCCCATCATCAACGGCCTGAGCGACCTGTACCACCCCATCCAGATCCTGGCCGACT 526
hOTCco       CTATCCCAATCATTAATGGCCTGAGTACCTGTATCACCCCATCCAGATTCTGGCCGATT 526
LW3          CTATCCCAATTATTAACGGCCTGTGAGACCTGTATCACCCCATCCAGATTCTGGCCGATT 540
LW4          CTATCCCAATTATTAACGGCCTGTGAGACCTGTATCACCCCATCCAGATTCTGGCCGATT 540
hOTwt        CCTATCCCAATTATCAATGGCCTGTGAGATTGTACCATCCTATCCAGATCCTGGCTGATT 526
* * * * *
hOTCLW5      ACCTGACCCTGCAGGAGCACTACAGCAGCCTGAAGGGCCTGACCCTGAGCTGGATCGGCC 586
hOTCLW6      ACCTGACCCTGCAGGAGCACTACAGCAGCCTGAAGGGCCTGACCCTGAGCTGGATCGGCC 586
hOTCco       ACCTGACCCTCCAGGAGCATTATCTAGTCTGAAAGGGCTGACACTGAGCTGGATTGGGG 586
LW3          ACCTGACCCTCCAGGAGCACTATTCTAGTCTGAAAGGGCTGACACTGAGTTGGATTGGGG 600
LW4          ACCTGACCCTCCAGGAGCACTATTCTAGTCTGAAAGGGCTGACACTGAGTTGGATTGGGG 600
hOTwt        ACCTCAGCTCCAGGAACACTATAGTCTCTGAAAGGTCTTACCCTCAGCTGGATCGGGG 586
* * * * *
hOTCLW5      ACGGCAACAACATCCTGCACAGCATCATGATGAGCGCCGCCAAGTTCCGCATGCACCTGC 646
hOTCLW6      ACGGCAACAACATCCTGCACAGCATCATGATGAGCGCCGCCAAGTTCCGCATGCACCTGC 646
hOTCco       ACGGAAACAATATCCTGCACCTCATTATGATGAGCGCCGCCAAGTTTGAATGCACCTCC 646
LW3          ACGGAAACAATATCCTGCACCTCATTATGATGATGTCAGCCGCCAAGTTTGAATGCACCTCC 660
LW4          ACGGAAACAATATCCTGCACCTCATTATGATGTCAGCCGCCAAGTTTGAATGCACCTCC 660
hOTwt        ATGGGAACAATATCCTGCACCTCATTATGATGATGAGCGCAGCGAAATTCCGAATGCACCTTC 646
* * * * *
hOTCLW5      AGGCCGCCACCCCAAGGGCTACGAGCCCGACGCCAGCGTGACCAAGCTGGCCGAGCAGT 706
hOTCLW6      AGGCCGCCACCCCAAGGGCTACGAGCCCGACGCCAGCGTGACCAAGCTGGCCGAGCAGT 706
hOTCco       AGGCTGCAACCCCAAAAGGCTACGAAACCCGATGCCTCCGTGACAAAGCTGGCAGAACAGT 706
LW3          AGGCTGCAACCCCAAAAGGCTACGAAACCCGATGCCTCAGTGACAAAGCTGGCTGAACAGT 720
LW4          AGGCTGCAACCCCAAAAGGCTACGAAACCCGATGCCTCAGTGACAAAGCTGGCTGAACAGT 720
hOTwt        AGGCAGCTACTCCAAAGGGTTATGAGCCGGATGCTAGTGTAAACCAAGTTGGCAGAGCAGT 706
* * * * *
hOTCLW5      ACGCCAAGGAGAACGGCACCAAGCTGCTGCTGACCAACGACCCCTGGAGGCCGCCACG 766
hOTCLW6      ACGCCAAGGAGAACGGCACCAAGCTGCTGCTGACCAACGACCCCTGGAGGCCGCCACG 766
hOTCco       ATGCCAAAGAGAACGGCACTAAGCTGCTGCTGACCAATGACCCCTCTGGAGGCCGCTCACG 766
LW3          ACGCCAAGGAGAACGGCACTAAGCTGCTGCTGACCAACGACCCCTCTGGAGGCCGCTCACG 780
LW4          ACGCCAAGGAGAACGGCACTAAGCTGCTGCTGACCAACGACCCCTCTGGAGGCCGCTCACG 780
hOTwt        ATGCCAAAGAGAATGGTACCAAGCTGTTGCTGACAAATGATCCATTGGAAGCAGCGCATG 766
* * * * *
hOTCLW5      GCGGCAACGTGCTGATCACCGACACCTGGATCAGCATGGGCCAGGAGGAGGAGAAGAAGA 826
hOTCLW6      GCGGCAACGTGCTGATCACCGACACCTGGATCAGCATGGGCCAGGAGGAGGAGAAGAAGA 826
hOTCco       GAGGCAACGTGCTGATCACTGATACCTGGATTAGTATGGGACAGGAGGAAGAGAAGAAGA 826
LW3          GAGGCAACGTGCTGATCACCGATACCTGGATTAGTATGGGACAGGAGGAAGAGAAGAAGA 840
LW4          GAGGCAACGTGCTGATCACCGATACCTGGATTAGTATGGGACAGGAGGAAGAGAAGAAGA 840
hOTwt        GAGGCAATGTATTAATTACAGACACTGGATAAGCATGGGACAAGAAGAGGAGAAGAAA 826
* * * * *
    
```

FIG. 5C

```

hOTCLW5      AGAGACTGCAGGCCTTCCAGGGCTACCAGGTGACCATGAAGACCGCCAAGGTGGCCGCCA 886
hOTCLW6      AGCGCCTGCAGGCCTTCCAGGGCTACCAGGTGACCATGAAGACCGCCAAGGTGGCCGCCA 886
hOTCco       AGCGGCTCCAGGCCTTCCAGGGCTACCAGGTGACAATGAAAACCGCTAAGGTTCGCAGCCA 886
LW3          AGCGGCTCCAGGCCTTCCAGGGCTACCAGGTGACAATGAAAACCGCTAAGGTTCGCAGCCA 900
LW4          AGCGGCTCCAGGCCTTCCAGGGCTACCAGGTGACAATGAAAACCGCTAAGGTTCGCAGCCA 900
hOTwt        AGCGGCTCCAGGCTTTCCAAGGTTACCAGGTTACAATGAAGACTGCTAAAGTTGCTGCCT 886
             ** * ** ***** ** ***** ** ***** ** ** ** **

hOTCLW5      GCGACTGGACCTTTCCTGCACTGCCTGCCAGAAAGCCCGAGGAGGTGGACGACGAGGTGT 946
hOTCLW6      GCGACTGGACCTTTCCTGCACTGCCTGCCCGCAAGCCCGAGGAGGTGGACGACGAGGTGT 946
hOTCco       GCGACTGGACCTTTCCTGCATTGCCTGCCAGAAAGCCTGAAGAGGTGGACGATGAGGTCT 946
LW3          GCGATTGGACCTTTCCTGCACTGCCTGCCAGAAAGCCCGAAGAGGTGGACGACGAGGTCT 960
LW4          GCGATTGGACCTTTCCTGCACTGCCTGCCAGAAAGCCCGAAGAGGTGGACGACGAGGTCT 960
hOTwt        CTGACTGGACATTTTTACACTGCTTGCAGAAAGCCAGAAAGTGGATGATGAAGTCT 946
             ** ***** ** * ** ** ***** * ***** ** ** ***** ** ** **

hOTCLW5      TCTACAGCCCCAGAAGCCTGGTGTTCCTCCGAGGCCGAGAACAGAAAGTGGACCATCATGG 1006
hOTCLW6      TCTACAGCCCCCGCAGCCTGGTGTTCCTCCGAGGCCGAGAACCGCAAGTGGACCATCATGG 1006
hOTCco       TCTACTCAGCCAGAAGCCTGGTGTTCCTGAAGCTGAGAATAGGAAGTGGACAATCATGG 1006
LW3          TCTACTCTCCCAGAAGCCTGGTGTTCCTCCGAAAGCTGAGAATAGGAAGTGGACAATTATGG 1020
LW4          TCTACTCTCCCAGAAGCCTGGTGTTCCTCCGAAAGCTGAGAATAGGAAGTGGACAATTATGG 1020
hOTwt        TTTATTCTCCTCGATCACTAGTGTTCCTCCGAGGCCAGAAAACAGAAAGTGGACAATCATGG 1006
             * ** ** * ** ***** ** ** ** **

hOTCLW5      CCGTGATGGTGAGCCTGCTGACCGACTACAGCCCCAGCTGCAGAAGCCCAAGTTCTGA- 1065
hOTCLW6      CCGTGATGGTGAGCCTGCTGACCGACTACAGCCCCAGCTGCAGAAGCCCAAGTTCTGA- 1065
hOTCco       CAGTGATGGTCAGCCTGCTGACTGATTATTCCTCCAGCTCCAGAAACCAAGTTCTGAT 1066
LW3          CAGTGATGGTCAGCCTGCTGACTGATTATTCACCTCAGCTCCAGAAACCAAGTTCTGAT 1080
LW4          CAGTGATGGTCAGCCTGCTGACTGATTATTCACCTCAGCTCCAGAAACCAAGTTCTGAT 1080
hOTwt        CTGTCAATGGTGTCCCTGCTGACAGATTACTCACCTCAGCTCCAGAAGCCTAAATTT--- 1062
             * ** ***** ***** ** ** ** ***** ***** ** ** **

hOTCLW5      -----
hOTCLW6      -----
hOTCco       AA----- 1068
LW3          AAGCGGCCGC 1090
LW4          AAGCGGCCGC 1090
hOTwt        -----
    
```

**COMPOSITIONS USEFUL IN TREATMENT  
OF ORNITHINE TRANSCARBAMYLASE  
(OTC) DEFICIENCY**

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

**[0001]** This application is a continuation of U.S. patent application Ser. No. 16/995,572, filed Aug. 17, 2020, which is a continuation of U.S. patent application Ser. No. 16/191,709, filed Nov. 15, 2018, now U.S. Pat. No. 10,781,430, issued Sep. 22, 2020, which is a continuation of U.S. patent application Ser. No. 15/122,853, filed Aug. 31, 2016, now U.S. Pat. No. 10,167,454, issued Jan. 1, 2019, which is a national stage application under 35 U.S.C. 371 of PCT/US2015/019513, filed on Mar. 9, 2015, now expired, which claims the benefit of U.S. Patent Application No. 61/950,157, filed Mar. 9, 2014, now expired. These applications are incorporated by reference in their entirety.

**[0002]** This invention was made with support under grant numbers HD057247, HL059407 and DK047757 awarded by the National Institutes of Health. The US government has certain rights in the invention.

**INCORPORATION-BY-REFERENCE OF  
MATERIAL SUBMITTED IN ELECTRONIC  
FORM**

**[0003]** Applicant hereby incorporates by reference the Sequence Listing material filed in electronic form herewith. This file is labeled "UPN-14-7037USC3.xml" (created on Jun. 22, 2023, 43,968 bytes).

**BACKGROUND OF THE INVENTION**

**[0004]** Ornithine transcarbamylase (OTC) deficiency accounts for nearly half of all cases of inborn errors of urea synthesis, with a prevalence estimated to be at least 1 in 15,000. Urea cycle defects put patients at risk of life threatening elevation of ammonia that can lead to irreversible cognitive impairment, coma and death. Newborn males with complete deficiency develop hyperammonemic coma within the first 3 days of life, which if untreated, is lethal.

**[0005]** Current therapies for OTC deficiency (OTCD) have numerous challenges. Patients can be managed with a low protein diet in combination with the use of medications that activate alternate nitrogen clearance pathways, but this does not prevent hyperammonemic crises. Despite the use of dialysis and alternate pathway therapy, there is almost a 50% mortality rate in neonates. Liver transplantation can cure OTCD, but donor liver is limiting, the procedure carries significant morbidity and immunosuppressive drugs are necessary for the duration of the subject's life.

**[0006]** Gene therapy of a metabolic disease such as OTCD presents a more challenging model for gene replacement therapy than other conditions. Because the gene acts in a cell-autonomous manner (i.e., it can only influence the cell in which it is expressed), therapeutic effects should be directly correlated with the number of target cells that are transduced, rather than with the net level of expression in liver such as with a secreted protein where high expression per cell can overcome low transduction. Furthermore, there has been at least one published report that hOTCwt mRNA is unstable. [Wang, L., et al, Molecular Genetics and Metabolism, 105 (2012) 203-211].

**[0007]** There have been published reports of using viral vectors to try to treat OTC deficiency. For example, several groups have tried this in murine models of OTC deficiency, using recombinant adenoviruses carrying rat, mouse, or human OTC cDNA. Some measure of successful reconstitution of liver OTC activity and correction of metabolic derangements have been reported in animal models with viruses carrying rat or mouse OTC cDNA. Previous studies using adenoviral vectors have illustrated the difficulties of expressing sufficient levels of active human OTC in OTCD mice.

**[0008]** Therefore, there is a need for other approaches to OTCD therapy.

**SUMMARY OF THE INVENTION**

**[0009]** In one aspect, the invention provides a recombinant viral vector having an expression cassette comprising an engineered nucleic acid sequence encoding human ornithine transcarbamylase (hOTCase) and expression control sequences which direct expression of hOTC in a liver cell, wherein the hOTC nucleic acid sequence is less than 80% identical to the wild-type hOTC sequence over the full-length hOTC of the wild-type sequence (e.g., SEQ ID NO:1), or a fragment thereof which comprises the mature hOTC but lacking at least the native leader sequence, or another intermediate which comprises at least the mature hOTC and expresses a functional hOTCase. Suitably, the engineered sequence has been preferably codon optimized and further improved such that it enhances at least one of transduction, transcription and/or translation of the enzyme.

**[0010]** The nucleic acid sequence may comprise the mature hOTC of SEQ ID NO: 5, or a nucleic acid sequence at least about 96 to about 99% identical thereto or a nucleic acid sequence comprising at least the mature hOTC of SEQ ID NO: 9, or a nucleic acid sequence at least about 96 to about 99% identical thereto, which expresses a functional hOTCase. In one embodiment, the hOTC is the full-length of SEQ ID NO: 5 or a nucleic acid sequence at least about 96 to about 99% identical thereto or a nucleic acid sequence of the full-length of SEQ ID NO: 9, or a nucleic acid sequence at least about 96 to about 99% identical thereto. The hOTC sequence may be that of the corresponding nucleotides of SEQ ID NO: 3, 4, 8 or 9. Encompassed within the scope of the invention are the strands complementary to those in the sequence listing. The viral vector may be selected from an adeno-associated virus (AAV) vector, an adenoviral vector, and a lentiviral vector.

**[0011]** In a further aspect, the invention provides a recombinant adeno-associated virus (rAAV) having an AAV capsid and packaged therein an expression cassette comprising at least one AAV inverted terminal repeat (ITR) sequence, an engineered nucleic acid sequence encoding human ornithine transcarbamylase (hOTCase) and expression control sequences which direct expression of hOTC in a liver cell, said expression control sequences comprising a liver-specific promoter. The engineered hOTC nucleic acid sequence is less than 80% identical to the wild-type hOTC sequence over the mature sequence or full-length hOTC of the wild-type sequence (e.g., SEQ ID NO:1) and expresses a functional hOTCase. The synthetic hOTC nucleic acid sequence comprises at least the mature hOTC of SEQ ID NO: 5 or a nucleic acid sequence at least about 96 to about 99.9% identical thereto or a nucleic acid sequence comprising at

least the mature hOTC of SEQ ID NO: 9 or a nucleic acid sequence at least about 96 to about 99.9% identical thereto.

**[0012]** In still a further aspect, the invention provides a viral vector comprising a hOTC gene encoding a chimeric ornithine transcarbamylase which comprises mature human ornithine transcarbamylase with a heterologous transit peptide, wherein the coding sequence from the mature human ornithine transcarbamylase is selected from the nucleic acid sequence comprising at least the mature hOTC of SEQ ID NO: 3, 4, 5, 8 or 9. Optionally, the full-length sequences of any of these sequences, which include the transit sequence, may be selected. Alternatively, a chimeric OTC gene including a heterologous transit sequence, as described herein, and these mature hOTC may be selected.

**[0013]** In another aspect, a pharmaceutical composition comprising a carrier and an effective amount of a vector as described herein is provided.

**[0014]** Yet another aspect is a viral vector as described herein used in preparing a medicament for delivering ornithine transcarbamylase to a subject in need thereof and/or for treating ornithine transcarbamylase deficiency. In one particularly desirable embodiment, the subject is a human subject. The subject may be homozygous or heterozygous for ornithine transcarbamylase deficiency.

**[0015]** In still another aspect, use of a viral vector comprising a nucleic acid sequence encoding functional human ornithine transcarbamylase in preventing and/or treating fibrosis or ornithine transcarbamylase deficiency (OTCD)-related cirrhosis in a subject for OTCD is provided. In one embodiment, the subject is a human patient. In a further embodiment, the subject is heterozygous and may exhibit late onset of symptoms.

**[0016]** In yet a further aspect, use of a viral vector comprising a nucleic acid sequence encoding functional human ornithine transcarbamylase in preventing and/or treating hepatocellular carcinoma in a subject having OTCD is provided. In one embodiment, the subject is a human patient. In a further embodiment, the subject is heterozygous for OTCD and may exhibit late onset of symptoms.

**[0017]** Other aspects and advantages of the invention will be readily apparent from the following detailed description of the invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0018]** FIG. 1A provides a wild-type hOTC cDNA, which has 324 A, 223 C, 246 G, and 269 T [SEQ ID NO: 1].

**[0019]** FIG. 1B-1C provides the human ornithine transcarbamylase sequence encoded by the sequence of FIG. 1A [SEQ ID NO:2].

**[0020]** FIG. 2 provides an engineered hOTC cDNA, with an altered GC ratio. The base count in the sequence is 283 A, 285 C, 284 G, and 216 T [SEQ ID NO: 3].

**[0021]** FIG. 3 provides an engineered hOTC cDNA termed LW3 (SEQ ID NO: 18). The base count in this sequence is 279 A, 303 C, 288 G, and 220 T. The start codon for the hOTC LW3 open reading frame (ORF) (SEQ ID NO: 4) is preceded by a Kozak sequence in this figure. The coding sequence (SEQ ID NO: 4) for the leader begins at nucleotide 15 (first 96 nucleotides), followed by the coding sequence for the 322 amino acid hOTCase. In this figure (SEQ ID NO: 18), the stop codon is followed by a NotI restriction site (GCGGCCGC) which is a remnant of the vector.

**[0022]** FIG. 4 provides an engineered hOTC cDNA termed LW4 (SEQ ID NO: 17). The base count in this sequence is 278 A, 303 C, 289 G, and 220 T. The coding sequence (SEQ ID NO: 5) for the leader begins at nucleotide 15 (first 96 nucleotides), followed by the coding sequence for the 322 amino acid hOTCase. In this figure (SEQ ID NO: 17), the stop codon is followed by a NotI restriction site (GCGGCCGC) which is a remnant of the vector.

**[0023]** FIGS. 5A-5C provides an alignment of the cDNA sequences of the wild-type hOTC, and five engineered sequences: hOTCco (SEQ ID NO: 3), LW3 (SEQ ID NO: 18), LW4 (SEQ ID NO: 17), LW5 (SEQ ID NO: 8) and LW6 (SEQ ID NO: 9). The aligned sequences contain a Kozak sequence (first 14 nucleotides of LW3 and LW4) and a restriction enzyme site (following termination codon for LW3 and LW4), which are not part of the open reading frames of LW3 (SEQ ID NO: 4) or LW4 (SEQ ID NO: 5).

#### DETAILED DESCRIPTION OF THE INVENTION

**[0024]** An engineered human (h) ornithine transcarbamylase (OTC) cDNA is provided herein, which was designed to maximize translation and improve mRNA stability as compared to the wild-type hOTC DNA and/or mRNA. Also provided herein are engineered hOTC mRNA sequences. These compositions may be used in therapeutic and/or prophylactic methods as described herein. Optionally, these compositions are used in combination other therapies consistent with the standard of care for the conditions for which the subject (e.g., a human subject) has been diagnosed.

**[0025]** For comparison purposes, a wild-type human OTC cDNA sequence is illustrated in FIG. 1A. This sequence encodes the human ornithine transcarbamylase of the amino acid sequence of FIGS. 1A-1C. This same amino acid sequence is encoded by the engineered hOTC genes of FIG. 2A-FIG. 5. The hOTC enzyme, which may be referred to as hOTCase to distinguish from the gene, is expressed from this sequence in the form of a pre-protein having a 32 amino acid leader peptide at its N-terminus (encoded by nt 1-96 of FIG. 1, about amino acids 1 to about 32 of SEQ ID NO: 2) which is cleaved after directing the enzyme to the cellular mitochondria, leaving the 322 amino acid residue "mature" protein (about amino acid 33 to about amino acid 354 of SEQ ID NO: 2). This "so-called mature" hOTCase is a homo-trimeric protein with a 322 amino acid residue sequence in each polypeptide chain. Optionally, as an alternative to the wild-type sequence of SEQ ID NO:2, one may select a sequence which includes one or more of the naturally occurring polymorphic positions that are not involved in disease: F101, L111, W193-194 of SEQ ID NO: 2 (see, e.g., [www.uniprot.org/uniprot/P00480](http://www.uniprot.org/uniprot/P00480)).

**[0026]** Although all of the engineered cDNA sequences are about 77% to about 78% identical to the wt hOTC nucleic acid sequence of FIG. 1A (SEQ ID NO: 1), there are structural differences between these sequences (see alignment in FIG. 5 illustrating same). Particularly, there is about 4% difference in nucleic acid sequences between hOTCco of FIG. 2 (SEQ ID NO: 3) and the hOTCcoLW4 (SEQ ID NO: 5) of FIG. 4. There is only one nt difference between LW3 (FIG. 3, SEQ ID NO: 18) and LW4 (FIG. 4, SEQ ID NO: 17), i.e., 0.094% (1/1062) difference (an A in LW3 is changed to a G in LW4 as shown in FIG. 5).

**[0027]** In one embodiment, a modified hOTC coding sequence is provided which sequence has less than about

80% identity, preferably about 77% identity or less to the full-length wild-type hOTC coding sequence (FIG. 1A, SEQ ID NO: 1), which encodes functional hOTCase. In one embodiment, the modified hOTC coding sequence is characterized by improved stability as compared to wt hOTC following AAV-mediated delivery (e.g., rAAV). Additionally or alternatively, a modified hOTC coding sequence is provided which lacks alternative reading frames for proteins of at least about 9 amino acids in length. Additionally, or alternatively, a modified hOTC coding sequence is provided which has hOTCase expression levels at least about 25-fold, at least about 50-fold, or at least about 100-fold when measured following expression from a viral vector, as compared to the hOTCase wild-type. Additionally, or alternatively, a modified hOTC coding sequence is provided which has hOTCase liver activity which is at least about 10-fold higher, at least about 20-fold higher, or at least about 30-fold higher as compared to the hOTCase wild-type expressed from a viral vector.

**[0028]** In one embodiment, a modified hOTC coding sequence is 96% to 99.9% identical to the sequence encoding the mature enzyme (about nt 99 to about 1068) or full-length of FIG. 4 (hOTCco-LW4, SEQ ID NO: 5), or 96.5% to 99% identical, or about 97%, or about 98% identical to SEQ ID NO: 17 (FIG. 4).

**[0029]** In one embodiment, a modified hOTC coding sequence is 96% to 99.9% identical to the sequence encoding the mature enzyme (about nt 99 to about 1068) of FIG. 3 (hOTCco-LW3, SEQ ID NO: 4), or 96.5% to 99% identical, or about 97%, or about 98% identical to SEQ ID NO: 18 (FIG. 3).

**[0030]** In another embodiment, a modified hOTC coding sequence is 96% to 99.9% identical to the sequence encoding the mature enzyme (about nt 99 to about 1068) or the full-length of FIG. 2 (hOTCco, SEQ ID NO: 3), or 96.5% to 99% identical, or about 97%, or 98% identical to SEQ ID NO: 3 (FIG. 2).

**[0031]** In still another embodiment, a modified hOTC coding sequence has the sequence encoding the mature protein (about nt 99 to about 1068) or the full-length of hOTCco-LW5 [SEQ ID NO: 8] or hOTCco-LW6 [SEQ ID NO:9], or a sequence 96% to 99.9% identical thereto. hOTCco-LW5 and hOTCco-LW6 are about 97% identical to each other, and each is about 78% identical to the wild-type sequence [SEQ ID NO: 1].

**[0032]** The sequences of FIGS. 2-5 are provided as the sense strand of the cDNA sequences. The present invention also encompasses the anti-sense strands corresponding to these cDNA sequences and corresponding RNA, e.g., mRNA, sequences. For example, the engineered mRNA of SEQ ID NO: 10, corresponds to the DNA of SEQ ID NO:4; the engineered RNA of SEQ ID NO: 11, corresponds to the DNA of SEQ ID NO: 5; the engineered RNA of SEQ ID NO: 12, corresponds to the DNA of SEQ ID NO: 8; and the RNA of SEQ ID NO: 13 corresponds to the DNA of SEQ ID NO:9. These RNA sequences, and sequences which are 95% to 99%, or about 97%, or about 98% identical to one or more of these sequences are encompassed within the scope of this invention. Methods for aligning and determining RNA identity are known in the art and include published and publicly available web-based or commercially available databases and services. See, e.g., LocARNA, CARNA, as well as other programs identified elsewhere therein.

**[0033]** In one embodiment of the invention, the mRNA sequence may be delivered using a selected RNA delivery system, examples of which are supplied herein.

**[0034]** Also encompassed herein are fragments, e.g., the sequences encoding the transit peptide (amino acids 1 to about 32), about amino acids 332 to about 354, an intermediate hOTC enzyme, or the mature enzyme, or other fragments as may be desired. Reference may be made to SEQ ID NO:2. See, e.g., Ye et al. 2001, Hum Gene Ther 12: 1035-1046.

**[0035]** In another embodiment, a chimeric OTC is provided in which the N-terminal presequence of wild-type OTC is replaced with a transit sequence from another source which is compatible with the subject's system such that it effectively transports the mature hOTCase encoded by the chimeric OTC gene to the desired organelle. See, e.g., Ye et al. 2001, Hum Gene Ther. 12: 1035-1046. Such transit sequences encode a transit peptide (also termed a signal peptide, targeting signal, or localization signal) which is fused to the coding sequence for the mature hOTC of SEQ ID NO: 1, 3, 4, 5, 8 and/or 9. For example, the wild-type hOTC transit sequence corresponds to about the first 98 nucleotides of SEQ ID NO: 1. To construct a chimeric OTC, these wild-type N-terminal sequences may be removed (about nucleic acids 1 to about nt 96-nt 98) and replaced with a heterologous transit sequence. Suitable transit peptides are preferably, although not necessarily of human origin. Suitable transit peptides may be chosen from proline. [bic.nus.edu.sg/spdb/zhang270.htm](http://bic.nus.edu.sg/spdb/zhang270.htm), which is incorporated by reference herein, or may be determined using a variety of computational programs for determining the transit peptide in a selected protein. Although not limited, such sequences may be from about 15 to about 50 amino acids in length, or about 20 to about 28 amino acids in length, or may be larger or smaller as required.

**[0036]** The term "percent (%) identity", "sequence identity", "percent sequence identity", or "percent identical" in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for correspondence. The length of sequence identity comparison may be over the full-length of the genome, the full-length of a gene coding sequence, or a fragment of at least about 500 to 5000 nucleotides, is desired. However, identity among smaller fragments, e.g. of at least about nine nucleotides, usually at least about 20 to 24 nucleotides, at least about 28 to 32 nucleotides, at least about 36 or more nucleotides, may also be desired.

**[0037]** Percent identity may be readily determined for amino acid sequences over the full-length of a protein, polypeptide, about 32 amino acids, about 330 amino acids, or a peptide fragment thereof or the corresponding nucleic acid sequence coding sequences. A suitable amino acid fragment may be at least about 8 amino acids in length, and may be up to about 700 amino acids. Generally, when referring to "identity", "homology", or "similarity" between two different sequences, "identity", "homology" or "similarity" is determined in reference to "aligned" sequences. "Aligned" sequences or "alignments" refer to multiple nucleic acid sequences or protein (amino acids) sequences, often containing corrections for missing or additional bases or amino acids as compared to a reference sequence.

**[0038]** Alignments are performed using any of a variety of publicly or commercially available Multiple Sequence Alignment Programs. Sequence alignment programs are

available for amino acid sequences, e.g., the “Clustal X”, “MAP”, “PIMA”, “MSA”, “BLOCKMAKER”, “MEME”, and “Match-Box” programs. Generally, any of these programs are used at default settings, although one of skill in the art can alter these settings as needed. Alternatively, one of skill in the art can utilize another algorithm or computer program which provides at least the level of identity or alignment as that provided by the referenced algorithms and programs. See, e.g., J. D. Thomson et al, Nucl. Acids. Res., “A comprehensive comparison of multiple sequence alignments”, 27(13):2682-2690 (1999).

**[0039]** Multiple sequence alignment programs are also available for nucleic acid sequences. Examples of such programs include, “Clustal W”, “CAP Sequence Assembly”, “BLAST”, “MAP”, and “MEME”, which are accessible through Web Servers on the internet. Other sources for such programs are known to those of skill in the art. Alternatively, Vector NTI utilities are also used. There are also a number of algorithms known in the art that can be used to measure nucleotide sequence identity, including those contained in the programs described above. As another example, polynucleotide sequences can be compared using Fasta™, a program in GCG Version 6.1. Fasta™ provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta™ with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference.

**[0040]** In one embodiment, the modified hOTC genes described herein are engineered into a suitable genetic element (vector) useful for generating viral vectors and/or for delivery to a host cell, e.g., naked DNA, phage, transposon, cosmid, episome, etc., which transfers the hOTC sequences carried thereon. The selected vector may be delivered by any suitable method, including transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. The methods used to make such constructs are known to those with skill in nucleic acid manipulation and include genetic engineering, recombinant engineering, and synthetic techniques. See, e.g., Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, NY.

**[0041]** As used herein, an “expression cassette” refers to a nucleic acid molecule which comprises the hOTC sequences, promoter, and may include other regulatory sequences therefor, which cassette may be packaged into the capsid of a viral vector (e.g., a viral particle). Typically, such an expression cassette for generating a viral vector contains the hOTC sequences described herein flanked by packaging signals of the viral genome and other expression control sequences such as those described herein. For example, for an AAV viral vector, the packaging signals are the 5' inverted terminal repeat (ITR) and the 3' ITR.

**[0042]** In one embodiment, the ITR sequences from AAV2, or the deleted version thereof (AITR), are used for convenience and to accelerate regulatory approval. However, ITRs from other AAV sources may be selected. Where the source of the ITRs is from AAV2 and the AAV capsid is from another AAV source, the resulting vector may be termed pseudotyped.

**[0043]** Typically, an expression cassette for an AAV vector comprises an AAV 5' ITR, the hOTC coding sequences and any regulatory sequences, and an AAV 3' ITR. However, other configurations of these elements may be suitable. A shortened version of the 5' ITR, termed AITR, has been described in which the D-sequence and terminal resolution site (trs) are deleted. In other embodiments, the full-length AAV 5' and 3' ITRs are used.

**[0044]** In one embodiment, the construct is a DNA molecule (e.g., a plasmid) useful for generating viral vectors. An illustrative plasmid containing desirable vector elements is illustrated by pAAVsc.TBG.hOTCco-LW4, the sequence of which is SEQ ID NO: 6 and which is incorporated by reference. This illustrative plasmid contains an expression cassette comprising: scITR (nt 5-109 of SEQ ID NO: 6), a TATA signal (nt 851-854 of SEQ ID NO:6), a synthetic hOTC coding sequence (nt 976-2037 of SEQ ID NO: 6), a poly A (nt 2182-2046 on the complement of SEQ ID NO: 6), a scITR (nt 2378-2211 on the complement of SEQ ID NO: 6), and a liver specific (TBG) promoter (nt 4172-4760) of SEQ ID NO: 6). Other expression cassettes may be generated using the other synthetic hOTC coding sequences, and other expression control elements, described herein.

**[0045]** The abbreviation “sc” in this context refers to self-complementary. “Self-complementary AAV” refers to a construct in which a coding region carried by a recombinant AAV nucleic acid sequence has been designed to form an intra-molecular double-stranded DNA template. Upon infection, rather than waiting for cell mediated synthesis of the second strand, the two complementary halves of scAAV will associate to form one double stranded DNA (dsDNA) unit that is ready for immediate replication and transcription. See, e.g., D M McCarty et al, “Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis”, Gene Therapy, (August 2001), Vol 8, Number 16, Pages 1248-1254. Self-complementary AAVs are described in, e.g., U.S. Pat. Nos. 6,596,535; 7,125,717; and 7,456,683, each of which is incorporated herein by reference in its entirety.

**[0046]** The expression cassette typically contains a promoter sequence as part of the expression control sequences, e.g., located between the selected 5' ITR sequence and the hOTC coding sequence. The illustrative plasmid and vector described herein uses the liver-specific promoter thyroxin binding globulin (TBG). Alternatively, other liver-specific promoters may be used [see, e.g., The Liver Specific Gene Promoter Database, Cold Spring Harbor, [rulai.schl.edu/LSPD/](http://rulai.schl.edu/LSPD/), such as, e.g., alpha 1 anti-trypsin (AIAT); human albumin Miyatake et al., J. Virol., 71:5124-32 (1997), humAlb; and hepatitis B virus core promoter, Sandig et al., Gene Ther., 3:1002-9 (1996)]. TTR minimal enhancer/promoter, alpha-antitrypsin promoter, LSP (845 nt)25 (requires intron-less scAAV); or LSP1. Although less desired, other promoters, such as constitutive promoters, regulatable promoters [see, e.g., WO 2011/126808 and WO 2013/04943], or a promoter responsive to physiologic cues may be used in the vectors described herein.

**[0047]** In addition to a promoter, an expression cassette and/or a vector may contain one or more other appropriate transcription initiation, termination, enhancer sequences, efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation effi-

ciency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. Examples of suitable polyA sequences include, e.g., SV40, SV50, bovine growth hormone (bGH), human growth hormone, and synthetic polyAs. Examples of suitable enhancers include, e.g., the alpha fetoprotein enhancer, the TTR minimal promoter/enhancer, LSP (TH-binding globulin promoter/alpha1-microglobulin/bikunin enhancer), amongst others. In one embodiment, the expression cassette comprises one or more expression enhancers. In one embodiment, the expression cassette contains two or more expression enhancers. These enhancers may be the same or may differ from one another. For example, an enhancer may include an Alpha mic/bik enhancer. This enhancer may be present in two copies which are located adjacent to one another. Alternatively, the dual copies of the enhancer may be separated by one or more sequences. In still another embodiment, the expression cassette further contains an intron, e.g., the Promega intron. Other suitable introns include those known in the art, e.g., such as are described in WO 2011/126808. Optionally, one or more sequences may be selected to stabilize mRNA. An example of such a sequence is a modified WPRE sequence, which may be engineered upstream of the polyA sequence and downstream of the coding sequence [see, e.g., M A Zanta-Boussif, et al, *Gene Therapy* (2009) 16: 605-619.

**[0048]** These control sequences are “operably linked” to the hOTC gene sequences. As used herein, the term “operably linked” refers to both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

**[0049]** Recombinant AAV viral vectors are well suited for delivery of the hOTC expression sequences described herein. Such AAV vectors may contain ITRs which are from the same AAV source as the capsid. Alternatively, the AAV ITRs may be from a different AAV source than that which supplies the capsid.

**[0050]** Where pseudotyped AAV is to be produced, the ITRs in the expression are selected from a source which differs from the AAV source of the capsid. For example, AAV2 ITRs may be selected for use with an AAV capsid having a particular efficiency for targeting liver (e.g., hepatocytes). AAV capsids may be selected from AAV8 [U.S. Pat. Nos. 7,790,449; 7,282,199] and rh10 [WO 2003/042397] for the compositions described herein. However, other AAV, including, e.g., AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV9, and others such as, e.g., those described in WO 2003/042397; WO 2005/033321, WO 2006/110689; U.S. Pat. No. 7,588,772 B2, which are incorporated by reference herein] may be used in human subjects.

**[0051]** In one embodiment, a self-complementary AAV is provided. This viral vector may contain a  $\Delta 5'$  ITR and an AAV 3' ITR. In one example, the viral vector is scAAV2/8.TBG.hOTCco. In another example, the viral vector is scAAV2/rh10.TBG.hOTCco. These vectors both contain the 5'  $\Delta$ ITR from AAV2, the liver-specific TBG promoter, an engineered hOTCco coding sequence of the invention, an SV40 polyA, and the 3' AAV2 ITR in an AAV8 capsid [see, e.g., U.S. Pat. No. 8,318,480B2] or AAV rh10 capsid. The sequence may be selected from engineered hOTC of one of SEQ ID NO: 3, 4, 5, 8 or 9. Optionally, the transit sequence

of the engineered hOTC may be substituted with a heterologous transit sequence to provide a chimeric hOTC, which retains the mature hOTCase.

**[0052]** In another embodiment, a single-stranded AAV viral vector is provided. Such a vector may contain a 5' AAV ITR and a 3' ITR. One example is AAV2/8.TBG.hOTCco, which contains the full-length AAV2—5' ITR, the liver-specific TBG promoter, the hOTC coding sequence, a bovine growth hormone polyA, and AAV2

**[0053]** 3' ITR. Another example is AAV2/8.TBG.hOTCco-.WPRE.bGH, which contains the same vector elements, and additionally contains the woodchuck hepatitis virus post-transcriptional regulatory element (WPRE). In other embodiments, WPRE is absent from constructs to be used in vivo. The engineered hOTC sequence (abbreviated herein hOTCco) may be selected from engineered hOTC of one of SEQ ID NO: 3, 4, 5, 8 or 9. Optionally, the transit peptide sequence of the engineered hOTC may be substituted with a heterologous transit sequence to provide a chimeric hOTC, which retains the mature hOTCase.

**[0054]** Still other promoters may be selected, including tissue specific promoters. Methods for generating and isolating AAV viral vectors suitable for delivery to a subject are known in the art. See, e.g. US Published Patent Application No. 2007/0036760 (Feb. 15, 2007), U.S. Pat. Nos. 7,790,449; 7,282,199; WO 2003/042397; WO 2005/033321, WO 2006/110689; and U.S. Pat. No. 7,588,772 B2]. The sequences of AAV8 and methods of generating vectors based on the AAV8 capsid are described in U.S. Pat. No. 7,282,199 B2, U.S. Pat. Nos. 7,790,449, and 8,318,480, which are incorporated herein by reference. In a one system, a producer cell line is transiently transfected with a construct that encodes the transgene flanked by ITRs and a construct (s) that encodes rep and cap. In a second system, a packaging cell line that stably supplies rep and cap is transiently transfected with a construct encoding the transgene flanked by ITRs. In each of these systems, AAV virions are produced in response to infection with helper adenovirus or herpesvirus, requiring the separation of the rAAVs from contaminating virus. More recently, systems have been developed that do not require infection with helper virus to recover the AAV—the required helper functions (i.e., adenovirus E1, E2a, VA, and E4 or herpesvirus UL5, UL8, UL52, and UL29, and herpesvirus polymerase) are also supplied, in trans, by the system. In these newer systems, the helper functions can be supplied by transient transfection of the cells with constructs that encode the required helper functions, or the cells can be engineered to stably contain genes encoding the helper functions, the expression of which can be controlled at the transcriptional or posttranscriptional level. In yet another system, the transgene flanked by ITRs and rep/cap genes are introduced into insect cells by infection with baculovirus-based vectors. For reviews on these production systems, see generally, e.g., Zhang et al., 2009, “Adenovirus-Adeno-associated virus hybrid for large-scale recombinant adeno-associated virus production,” *Human Gene Therapy* 20:922-929, the contents of each of which is incorporated herein by reference in its entirety. Methods of making and using these and other AAV production systems are also described in the following U.S. patents, the contents of each of which is incorporated herein by reference in its entirety: U.S. Pat. Nos. 5,139,941; 5,741,683; 6,057,152;

6,204,059; 6,268,213; 6,491,907; 6,660,514; 6,951,753; 7,094,604; 7,172,893; 7,201,898; 7,229,823; and 7,439,065.

**[0055]** The available space for packaging may be conserved by combining more than one transcription unit into a single construct, thus reducing the amount of required regulatory sequence space. For example, a single promoter may direct expression of a single RNA that encodes two or three or more genes, and translation of the downstream genes are driven by IRES sequences. In another example, a single promoter may direct expression of an RNA that contains, in a single open reading frame (ORF), two or three or more genes separated from one another by sequences encoding a self-cleavage peptide (e.g., T2A) or a protease recognition site (e.g., furin). The ORF thus encodes a single polyprotein, which, either during (in the case of T2A) or after translation, is cleaved into the individual proteins (such as, e.g., transgene and dimerizable transcription factor). It should be noted, however, that although these IRES and polyprotein systems can be used to save AAV packaging space, they can only be used for expression of components that can be driven by the same promoter. In another alternative, the transgene capacity of AAV can be increased by providing AAV ITRs of two genomes that can anneal to form head to tail concatamers.

**[0056]** Optionally, the hOTC genes described herein may be used to generate viral vectors other than rAAV. Such other viral vectors may include any virus suitable for gene therapy may be used, including but not limited to adenovirus; herpes virus; lentivirus; retrovirus; etc. Suitably, where one of these other vectors is generated, it is produced as a replication-defective viral vector.

**[0057]** A “replication-defective virus” or “viral vector” refers to a synthetic or artificial viral particle in which an expression cassette containing a gene of interest is packaged in a viral capsid or envelope, where any viral genomic sequences also packaged within the viral capsid or envelope are replication-deficient; i.e., they cannot generate progeny virions but retain the ability to infect target cells. In one embodiment, the genome of the viral vector does not include genes encoding the enzymes required to replicate (the genome can be engineered to be “gutless”—containing only the transgene of interest flanked by the signals required for amplification and packaging of the artificial genome), but these genes may be supplied during production. Therefore, it is deemed safe for use in gene therapy since replication and infection by progeny virions cannot occur except in the presence of the viral enzyme required for replication. Such replication-defective viruses may be adeno-associated viruses (AAV), adenoviruses, lentiviruses (integrating or non-integrating), or another suitable virus source.

**[0058]** The pharmaceutical compositions described herein are designed for delivery to subjects in need thereof by any suitable route or a combination of different routes. Direct or intrahepatic delivery to the liver is desired and may optionally be performed via intravascular delivery, e.g., via the portal vein, hepatic vein, bile duct, or by transplant. Alternatively, other routes of administration may be selected (e.g., oral, inhalation, intranasal, intratracheal, intraarterial, intraocular, intravenous, intramuscular, and other parenteral routes). The hOTC delivery constructs described herein may be delivered in a single composition or multiple compositions. Optionally, two or more different AAV may be delivered [see, e.g., WO 2011/126808 and WO 2013/049493]. In another embodiment, such multiple viruses may contain

different replication-defective viruses (e.g., AAV, adenovirus, and/or lentivirus). Alternatively, delivery may be mediated by non-viral constructs, e.g., “naked DNA”, “naked plasmid DNA”, RNA, and mRNA; coupled with various delivery compositions and nano particles, including, e.g., micelles, liposomes, cationic lipid—nucleic acid compositions, poly-glycan compositions and other polymers, lipid and/or cholesterol-based—nucleic acid conjugates, and other constructs such as are described herein. See, e.g., X. Su et al, *Mol. Pharmaceutics*, 2011, 8 (3), pp 774-787; web publication: Mar. 21, 2011; WO2013/182683, WO 2010/053572 and WO 2012/170930, both of which are incorporated herein by reference. Such non-viral hOTC delivery constructs may be administered by the routes described previously.

**[0059]** The viral vectors, or non-viral DNA or RNA transfer moieties, can be formulated with a physiologically acceptable carrier for use in gene transfer and gene therapy applications. In the case of AAV viral vectors, quantification of the genome copies (“GC”) may be used as the measure of the dose contained in the formulation. Any method known in the art can be used to determine the genome copy (GC) number of the replication-defective virus compositions of the invention. One method for performing AAV GC number titration is as follows: Purified AAV vector samples are first treated with DNase to eliminate un-encapsidated AAV genome DNA or contaminating plasmid DNA from the production process. The DNase resistant particles are then subjected to heat treatment to release the genome from the capsid. The released genomes are then quantitated by real-time PCR using primer/probe sets targeting specific region of the viral genome (usually poly A signal). The replication-defective virus compositions can be formulated in dosage units to contain an amount of replication-defective virus that is in the range of about  $1.0 \times 10^9$  GC to about  $1.0 \times 10^{15}$  GC (to treat an average subject of 70 kg in body weight), and preferably  $1.0 \times 10^{12}$  GC to  $1.0 \times 10^{14}$  GC for a human patient. Preferably, the dose of replication-defective virus in the formulation is  $1.0 \times 10^9$  GC,  $5.0 \times 10^9$  GC,  $1.0 \times 10^{10}$  GC,  $5.0 \times 10^{10}$  GC,  $1.0 \times 10^{11}$  GC,  $5.0 \times 10^{11}$  GC,  $1.0 \times 10^{12}$  GC,  $5.0 \times 10^{12}$  GC, or  $1.0 \times 10^{13}$  GC,  $5.0 \times 10^{13}$  GC,  $1.0 \times 10^{14}$  GC,  $5.0 \times 10^{14}$  GC, or  $1.0 \times 10^{15}$  GC.

**[0060]** DNA and RNA is generally measured in the nanogram (ng) to microgram ( $\mu$ g) amounts of the nucleic acids. In general, for a treatment in a human preferably dosages of the RNA is the range of 1 ng to 700  $\mu$ g, 1 ng to 500  $\mu$ g, 1 ng to 300  $\mu$ g, 1 ng to 200  $\mu$ g, or 1 ng to 100  $\mu$ g are formulated and administered. Similar dosage amounts of a DNA molecule containing an expression cassette and not delivered to a subject via a viral vector may be utilized for non-viral hOTC DNA delivery constructs.

**[0061]** Production of lentivirus is measured as described herein and expressed as IU per volume (e.g., mL). IU is infectious unit, or alternatively transduction units (TU); IU and TU can be used interchangeably as a quantitative measure of the titer of a viral vector particle preparation. The lentiviral vector is typically non-integrating. The amount of viral particles is at least about  $3 \times 10^6$  IU, and can be at least about  $1 \times 10^7$  IU, at least about  $3 \times 10^7$  IU, at least about  $1 \times 10^8$  IU, at least about  $3 \times 10^8$  IU, at least about  $1 \times 10^9$  IU, or at least about  $3 \times 10^9$  IU.

**[0062]** The above-described recombinant vectors may be delivered to host cells according to published methods. The rAAV, preferably suspended in a physiologically compatible



carrier, may be administered to a human or non-human mammalian patient. Suitable carriers may be readily selected by one of skill in the art in view of the indication for which the transfer virus is directed. For example, one suitable carrier includes saline, which may be formulated with a variety of buffering solutions (e.g., phosphate buffered saline). Other exemplary carriers include sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextran, agar, pectin, peanut oil, sesame oil, and water. The selection of the carrier is not a limitation of the present invention.

**[0063]** Optionally, the compositions of the invention may contain, in addition to the rAAV and carrier(s), other conventional pharmaceutical ingredients, such as preservatives, or chemical stabilizers. Suitable exemplary preservatives include chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl vanillin, glycerin, phenol, and parachlorophenol. Suitable chemical stabilizers include gelatin and albumin.

**[0064]** The viral vectors described herein may be used in preparing a medicament for delivering ornithine transcarbamylase to a subject (e.g., a human patient) in need thereof, supplying functional hOTCase to a subject, and/or for treating ornithine transcarbamylase deficiency. A course of treatment may optionally involve repeat administration of the same viral vector (e.g., an AAV8 vector) or a different viral vector (e.g., an AAV8 and an AAVrh10). Still other combinations may be selected using the viral vectors and non-viral delivery systems described herein.

**[0065]** In another embodiment, the nucleic acid sequences described herein may be delivered via a non-viral route. For example, a hOTC sequence may be via a carrier system for expression or delivery in RNA form (e.g., mRNA) using one of a number of carrier systems which are known in the art. Such carrier systems include those provided by commercial entities, such as PhaseRx' so-called "SMARTT" technology. These systems utilize block copolymers for delivery to a target host cell. See, e.g., US 2011/0286957 entitled, "Multiblock Polymers", published Nov. 24, 2011; US 2011/0281354, published Nov. 17, 2011; EP2620161, published Jul. 31, 2013; and WO 2015/017519, published Feb. 5, 2015. See, also, S. Uchida et al, (February 2013) PLoS ONE 8(2): e56220. Still other methods involve generating and injecting synthetic dsRNAs [see Soutschek et al. Nature (2004) 432 (7014): 173-8; see also Morrissey et al. Hepatol. (2005) 41(6): 1349-56], local administration to the liver has also been demonstrated by injecting double stranded RNA directly into the circulatory system surrounding the liver using renal vein catheterization. [See Hamar et al. PNAS (2004) 101(41): 14883-8.]. Still other systems involve delivery of dsRNA and particularly siRNA using cationic complexes or liposomal formulations [see, e.g., Landen et al. Cancer Biol. Ther. (2006) 5(12); see also Khoury et al. Arthritis Rheumatol. (2006) 54(6): 1867-77. Other RNA delivery technologies are also available, e.g., from Veritas Bio [see, e.g., US 2010/0323001, Dec. 23, 2010, "In vivo delivery of double stranded RNA to a target cell" (cytosolic content including RNAs, e.g., mRNA, expressed siRNA/shRNA/miRNA, as well as injected/introduced siRNA/shRNA/miRNA, or possibly even transfected DNA present in the cytosol packaged within exovesicles and be transported to distal sites such as the liver)]. Still other systems for in vivo delivery of RNA sequences have been described. See, e.g., US 2012/0195917 (Aug. 2, 2012) (5'-cap analogs of RNA to improve stability and increase RNA expression),

WO 2013/143555A1, Oct. 3, 2013, and/or are commercially available (BioNTech, Germany; Valera (Cambridge, MA); Zata Pharmaceuticals).

**[0066]** Thus, in one embodiment, the invention provides an engineered hOTC mRNA of the mature sequence (at least about nt 99-1098) or the full-length of SEQ ID NO: 10, 11, 12, 13, or a sequence having at least 97% to 99% identity thereto, in a composition for delivery of double-stranded or single stranded RNA which results in expression of the mature hOTCase in a target host cell, e.g., a liver cell.

**[0067]** The kinetics of the composition described herein which contain mRNA (delivered directly, as compared to transcribed from a DNA delivery molecule) are particularly well suited for use in subjects in acute crisis, as expression of the hOTCase from the mRNA may be seen within a period of several hours. In order to avoid rapid clearance of the RNA, it is modified as described herein (e.g., using a cap or a modified base), such that its effects may be retained for over 24 hours, over 48 hours, or up to about 3 days (about 72 hours). It may be desirable to co-administer an mRNA directly as described herein and co-administer at the same or substantially the same time, a DNA or viral vector-based hOTC composition as defined herein. Thus, a subject may receive immediate treatment, and at such time as the mRNA-mediated expression begins to wane, the longer-term hOTC expression conferred by a viral vector-mediated expression begins to take effect. Alternatively, a subject may receive a second administration of an mRNA-based composition as defined herein. The mRNA compositions described herein may be used in other therapeutic regimens or methods, including those involving OTCD patients who are not in acute crisis.

**[0068]** The compositions according to the present invention may comprise a pharmaceutically acceptable carrier, such as defined above. In one embodiment, the composition comprises a block copolymer associated with a hOTC polynucleotide as described herein. The block copolymers may form a micelle, such that the micelle comprises a plurality of block copolymer.

**[0069]** Typically, such a composition contains a nucleic acid molecule comprising the mRNA sequence corresponding to the hOTC sequence encoding the mature hOTCase (at least about nt 99 to 1098) or the full-length of any of SEQ ID NO: 10, 11, 12, 13. In addition, this nucleic acid molecule may include the 5' untranslated region (UTR), also known as the leader sequence or leader RNA, and one or more of an optional intron(s), an optional exon(s), an optional a Kozak sequence, an optional WPRE, and a polyA, and the 3' UTR flanking the coding sequences. Suitable leader sequences include those discussed above in connection with the hOTC DNA sequences, which discussion is incorporated by reference herein. Examples of sources of suitable leader sequences, other than the native hOTC leader sequences, or those corresponding to FIG. 2, FIG. 3 or FIG. 4, or FIG. 5 are discussed above. Similarly, sources of suitable introns, polyA, and Kozak sequences are discussed above and are applicable to the delivery of the corresponding RNA sequences discussed in the present paragraph. Further, various modifications to the RNA may be generated, e.g., a modified 5' cap structure may be engineered into the construct in order to avoid rapid clearance of the mRNA in vivo, or for another desired reason. Methods of generating such 5' cap structures is known to those of skill in the art. See, e.g., US 2012/0195917 and WO 2013/143555A1, Oct. 3, 2013.

In addition, modified nucleotides can be used to make mRNA in vitro, like pseudouridine. Also RNA may be dosed repetitively, or subject can be dosed first with mRNA to manage neonatal crises followed up by viral vector-mediated delivery (e.g., AAV) for long term therapy and to prevent fibrosis/cirrhosis and/or hepatocellular carcinoma.

**[0070]** mRNA can be synthesized from the hOTC DNA sequences described herein, using techniques that are well known in the art. For example, Cazenave C, Uhlenbeck O C, RNA template-directed RNA synthesis by T7 RNA polymerase. Proc Natl Acad Sci USA. 1994 Jul. 19; 91(15): 6972-6, describe the use of the T7 RNA polymerase for generating RNA from cDNA or RNA templates. See also, Wichlacz AI, Legiewicz M, Ciesiolka J., Generating in vitro transcripts with homogenous 3' ends using trans-acting antigenomic delta ribozyme, Nucleic Acids Res. 2004 Feb. 18; 32(3):e39; Krieg P A, Melton D A., Functional messenger RNAs are produced by SP6 in vitro transcription of cloned cDNAs, Nucleic Acids Res. 1984 Sep. 25; 12(18): 7057-70; and Rio, D. C., et al. RNA: A Laboratory Manual. Cold Spring Harbor: Cold Spring Harbor Laboratory Press, 2011, 205-220. Each of these references is incorporated herein by reference. In addition, kits and protocols for generating mRNA are available commercially including, without limitation, the Riboprobe® In Vitro Transcription System (Promega Corp.); RiboMAX™ Large Scale RNA Production Systems (Promega Corp.); MAXIscript Kit (Ambion); MEGIscript Kit (Ambion); MessageAmp™ aRNA Kit (Ambion); mMESSAgE mMACHINE® Kits (Ambion); and HiScribe™ T7 High Yield RNA Synthesis Kit (New England Biolabs® Inc.). Custom RNA can also be generated commercially from companies including, without limitation, TriLink Biotechnologies; bioSYNTHESIS; GE Dharmacon; and IBA Lifesciences.

**[0071]** The hOTC DNA sequences described herein can be generated in vitro and synthetically, using techniques well known in the art. For example, the PCR-based accurate synthesis (PAS) of long DNA sequence method may be utilized, as described by Xiong et al, PCR-based accurate synthesis of long DNA sequences, Nature Protocols 1, 791-797 (2006). A method combining the dual asymmetrical PCR and overlap extension PCR methods is described by Young and Dong, Two-step total gene synthesis method, Nucleic Acids Res. 2004; 32(7): e59. See also, Gordeeva et al, J Microbiol Methods. Improved PCR-based gene synthesis method and its application to the *Citrobacter freundii* phytase gene codon modification. 2010 May; 81(2):147-52. Epub 2010 Mar. 10; see, also, the following patents on oligonucleotide synthesis and gene synthesis, Gene Seq. 2012 April; 6(1):10-21; U.S. Pat. Nos. 8,008,005; and 7,985,565. Each of these documents is incorporated herein by reference. In addition, kits and protocols for generating DNA via PCR are available commercially. These include the use of polymerases including, without limitation, Taq polymerase; OneTaq® (New England Biolabs); Q5® High-Fidelity DNA Polymerase (New England Biolabs); and GoTaq® G2 Polymerase (Promega). DNA may also be generated from cells transfected with plasmids containing the hOTC sequences described herein. Kits and protocols are known and commercially available and include, without limitation, QIAGEN plasmid kits; Chargeswitch® Pro Filter Plasmid Kits (Invitrogen); and GenElute™ Plasmid Kits (Sigma Aldrich). Other techniques useful herein include sequence-specific isothermal amplification methods, that

eliminate the need for thermocycling. Instead of heat, these methods typically employ a strand-displacing DNA polymerase, like Bst DNA Polymerase, Large Fragment (New England Biolabs), to separate duplex DNA. DNA may also be generated from RNA molecules through amplification via the use of Reverse Transcriptases (RT), which are RNA-dependent DNA Polymerases. RTs polymerize a strand of DNA that is complimentary to the original RNA template and is referred to as cDNA. This cDNA can then be further amplified through PCR or isothermal methods as outlined above. Custom DNA can also be generated commercially from companies including, without limitation, GenScript; GENEWIZ®; GeneArt® (Life Technologies); and Integrated DNA Technologies.

**[0072]** The term “expression” is used herein in its broadest meaning and comprises the production of RNA or of RNA and protein. With respect to RNA, the term “expression” or “translation” relates in particular to the production of peptides or proteins. Expression may be transient or may be stable.

**[0073]** The term “translation” in the context of the present invention relates to a process at the ribosome, wherein an mRNA strand controls the assembly of an amino acid sequence to generate a protein or a peptide.

**[0074]** According to the present invention, a “therapeutically effective amount” of the hOTC is delivered as described herein to achieve a desired result, i.e., treatment of OTC deficiency or one or more symptoms thereof. As described herein, a desired result includes reducing orotic acid levels, reducing hyperammonemia and/or minimizing or eliminating one or more of the neurophysical complications including developmental delay, learning disabilities, intellectual disability, attention deficit hyperactivity disorder, and executive function deficits. Treatment may include treatment of subjects having severe neonatal-onset disease (males or, more rarely, females), and late-onset (partial) disease in males and females, which may present from infancy to later childhood, adolescence, or adulthood. In certain embodiments, the invention provides a method of treating and/or preventing fibrosis and/or cirrhosis in subjects, particularly those late-onset heterozygous subjects by administering hOTC as described herein. In one embodiment, therapeutic goals for OTC deficiency are to maintain plasma ammonia at less than <80 μmol/L, plasma glutamine <1,000 μmol/L, argininemia 80-150 μmol/L and branched chain amino acids within the normal range. However, other therapeutic endpoints may be selected by the treating physician.

**[0075]** In yet another embodiment, the invention provides a method of rescuing and/or treating a neonatal subject OTCD comprising the step of delivering a hOTC gene to the liver of a newborn subject (e.g., a human patient). This method may utilize any nucleic acid sequence encoding a functional hOTC, whether a synthetic hOTC as described herein or a wild-type hOTC, or a hOTC from another source, or a combination thereof. In one embodiment, neonatal treatment is defined as being administered a hOTC as described herein within 8 hours, the first 12 hours, the first 24 hours, or the first 48 hours of delivery. In another embodiment, particularly for a primate, neonatal delivery is within the period of about 12 hours to about 1 week, 2 weeks, 3 weeks, or about 1 month, or after about 24 hours to about 48 hours. To address dilution due to the rapid turnover of liver cells in a growing mammal (e.g., a non-

human or human primate), neonatal therapy is desirably followed by readministration at about 3 months of age, about 6 months, about 9 months, or about 12 months. Optionally, more than one readministration is permitted. Such readministration may be with the same type of vector, a different viral vector, or via non-viral delivery. In one embodiment, an RNA based delivery system for functional hOTC is used to stabilize a subject (e.g., a human patient) in crisis, followed by delivery of a viral vector mediated delivery of a functional hOTC. In another embodiment, initial therapy involves co-administration of viral and non-viral—mediated hOTC delivery systems. In a further embodiment, the hOTC DNA and RNA constructs may be used alone, or in combination with the standard of care for the patient's diagnosis and condition.

**[0076]** As described in the working examples herein, the inventors have found that heterozygous OTCD subjects, including those with late onset OTCD, have increased fibrosis and/or microvesicular steatosis throughout the liver. Such liver fibrosis and/or microvesicular steatosis can lead to OTCD-related cirrhosis. Thus, in another embodiment, the invention provides methods of preventing liver fibrosis and/or the associated medical condition OTCD-related cirrhosis by delivering to the subject (e.g., a human patient) a hOTC. This aspect of the invention may utilize a viral or non-viral delivery system. The nucleic acid expression cassette may contain a synthetic hOTC DNA or RNA as provided herein, or another suitable sequence which expresses functional hOTC. In one embodiment, a method of treating and/or preventing liver fibrosis, microvesicular steatosis, and/or OTCD-related cirrhosis is provided which involves delivering OTCase to a subject having OTCD. The subject may be a human patient. In one embodiment, the patient is heterozygous and has late onset OTCD. The patient may have been previously untreated for OTCD, or may have received other conventional treatments. At present, there is no existing standard of care for OTCD, but rather symptoms are managed, e.g., through discontinuation of protein intake, compensatory increases in dietary carbohydrates and lipids, hemodialysis for comatose patients with extremely high blood levels; and/or intravenous administration of sodium benzoate, arginine, and/or sodium phenylacetate. The US FDA has approved glycerol phenylbutyrate (Ravicti®) for long-term management of some urea cycle disorders for patients aged 2 years and older; this drug helps rid the body of ammonia and is intended for patients who cannot be managed by a protein-restricted diet or amino acid supplements alone. In one embodiment, treatment of the patient (e.g., a first injection) is initiated prior to the first year of life. In another embodiment, treatment is initiated after the first 1 year, or after the first 2 to 3 years of age, after 5 years of age, after 11 years of age, or at an older age.

**[0077]** In one embodiment, the method of the invention provides for treating and/or reversing liver fibrosis and/or OTCD-related cirrhosis by delivering to the subject a functional OTCase which is encoded by an engineered DNA of SEQ ID NO: 1, 3, 4, 5, 8 or 9, or a chimeric DNA as defined herein. Delivery of the DNA may be mediated by a viral vector containing the engineered DNA in an expression cassette, or by a non-viral delivery system, either of which mediates expression of functional OTCase in the liver cells of the subject. In another embodiment, the subject is administered an engineered RNA of SEQ ID NO: 10, 11, 12 or 13, or a chimeric RNA as defined herein. Delivery of the RNA

may be mediated by a viral vector containing the engineered RNA in an expression cassette, or by a non-viral delivery system, either of which mediates expression of functional OTCase in the liver cells of the subject.

**[0078]** Heterozygous OTCD subjects have an increased risk of developing hepatocellular carcinoma (HCC). See, e.g., J M Wilson et al, *Molecular Genetics and Metabolism* (2012), *Mol Genet Metab.* 2012 February; 105(2): 263-265, Published online 2011 Nov. 7. Thus, in another embodiment, the invention provides methods of preventing treating and/or preventing HCC by delivering to the subject (e.g., a human patient) a hOTC. This aspect of the invention may utilize a viral or non-viral delivery system. The nucleic acid expression cassette may contain a synthetic hOTC DNA or RNA as provided herein, or another suitable sequence which expresses functional hOTC. The patient may have been previously untreated for OTCD, or may have received other conventional treatments, i.e., standard of care. In one embodiment, treatment of the patient (e.g., a first injection) is initiated prior diagnosis with HCC. In another embodiment, treatment of the patient is initiated following HCC diagnosis. Optionally, treatment involves co-administration with sorafenib (commercially available as Nexavar®), or being used in conjunction with chemoembolization, radiation, thermal ablation, percutaneous ethanol injection, targeted therapy (e.g., anti-angiogenesis drugs), hepatic arterial infusion of anti-cancer drugs, immunotherapy, or with surgical options including, e.g., resection, cryosurgery, and liver transplant. When used for treatment of HCC, it may be desirable to select a non-integrating delivery system (e.g., direct RNA delivery, or non-integrating viruses such as adenoviruses or non-integrating lentiviruses) for delivery of a synthetic hOTC DNA or RNA as described herein.

**[0079]** By “functional OTC”, is meant a gene which encodes the wild-type OTCase such as characterized by SEQ ID NO: 2 or another OTCase which provides at least about 50%, at least about 75%, at least about 80%, at least about 90%, or about the same, or greater than 100% of the biological activity level of the wild-type human ornithine transcarbamylase enzyme, which may be characterized by the sequence of SEQ ID NO:2 or a natural variant or polymorph thereof which is not associated with disease. More particularly, as heterozygous patients may have as low an OTCase functional level as about 50% or lower, effective treatment may not require replacement of OTCase activity to levels within the range of “normal” or non-OTCD patients. Similarly, patients having no detectable amounts of OTCase may be rescued by delivering OTCase function to less than 100% activity levels, and may optionally be subject to further treatment subsequently. As described herein, the gene therapy described herein, whether viral or non-viral, may be used in conjunction with other treatments, i.e., the standard of care for the subject's (patient's) diagnosis.

**[0080]** In one embodiment, such a functional OTCase has a sequence which has about 95% or greater identity to the mature protein (i.e., about the last 322 amino acids) or full-length sequence of SEQ ID NO: 2, or about 97% identity or greater, or about 99% or greater to SEQ ID NO: 2 at the amino acid level. Such a functional OTCase may also encompass natural polymorphs which are not associated with any disease (e.g., F101, L111, and/or W1193-194 of SEQ ID NO: 2). Identity may be determined by preparing an alignment of the sequences and through the use of a variety of algorithms and/or computer programs known in the art or

commercially available [e.g., BLAST, ExpASY; ClustalO; FASTA; using, e.g., Needleman-Wunsch algorithm, Smith-Waterman algorithm].

**[0081]** A variety of assays exist for measuring OTC expression and activity levels in vitro. See, e.g., X Ye, et al, 1996 Prolonged metabolic correction in adult ornithine transcarbamylase-deficient mice with adenoviral vectors. *J Biol Chem* 271:3639-3646) or in vivo. For example, OTC enzyme activity can be measured using a liquid chromatography mass spectrometry stable isotope dilution method to detect the formation of citrulline normalized to [1,2,3,4,5-<sup>13</sup>C<sub>5</sub>] citrulline (98% <sup>13</sup>C). The method is adapted from a previously developed assay for detection of N-acetylglutamate synthase activity [Morizono H, et al, Mammalian N-acetylglutamate synthase. *Mol Genet Metab.* 2004; 81(Suppl 1):S4-11.]. Slivers of fresh frozen liver are weighed and briefly homogenized in buffer containing 10 mM HEPES, 0.5% Triton X-100, 2.0 mM EDTA and 0.5 mM DTT. Volume of homogenization buffer is adjusted to obtain 50 mg/ml tissue. Enzyme activity is measured using 250 µg liver tissue in 50 mM Tris-acetate, 4 mM ornithine, 5 mM carbamyl phosphate, pH 8.3. Enzyme activity is initiated with the addition of freshly prepared 50 mM carbamyl phosphate dissolved in 50 mM Tris-acetate pH 8.3, allowed to proceed for 5 minutes at 25° C. and quenched by addition of an equal volume of 5 mM <sup>13</sup>C<sub>5</sub>-citrulline in 30% TCA. Debris is separated by 5 minutes of microcentrifugation, and the supernatants are transferred to vials for mass spectroscopy. Ten µL of sample are injected into an Agilent 1100 series LC-MS under isocratic conditions with a mobile phase of 93% solvent A (1 ml trifluoroacetic acid in 1 L water):7% solvent B (1 ml trifluoroacetic acid in 1 L of 1:9 water/acetonitrile). Peaks corresponding to citrulline [176.1 mass:charge ratio (m/z)] and <sup>13</sup>C<sub>5</sub>-citrulline (181.1 m/z) are quantitated, and their ratios are compared to ratios obtained for a standard curve of citrulline run with each assay. Samples are normalized to either total liver tissue or to protein concentration determined using a Bio-Rad protein assay kit (Bio-Rad, Hercules, CA). Other assays, which do not require liver biopsy, may also be used. One such assay is a plasma amino acid assays in which the ratio of glutamine and citrulline is assessed and if glutamine is high (>800 microliters/liter) and citrulline is low (e.g., single digits), a urea cycle defect is suspected. Plasma ammonia levels can be measured and a concentration of about 100 micromoles per liter is indicative of OTCD. Blood gases can be assessed if a patient is hyperventilating; respiratory alkalosis is frequent in OTCD. Orotic acid in urine, e.g., greater than about 20 micromoles per millimole creatine is indicative of OTCD, as is elevated urinary orotate after allopurinol challenge test. Diagnostic criteria for OTCD have been set forth in Tuchman et al, 2008, Urea Cycle Disorders Consortium (UCDC) of the Rare Disease Clinical Research Network (RDCRN). Tuchman M, et al., Consortium of the Rare Diseases Clinical Research Network. Cross-sectional multicenter study of patients with urea cycle disorders in the United States. *Mol Genet Metab.* 2008; 94:397-402, which is incorporated by reference herein. See, also, [www.ncbi.nlm.nih.gov/books/NBK154378/](http://www.ncbi.nlm.nih.gov/books/NBK154378/), which provides a discussion of the present standard of care for OTCD.

**[0082]** It is to be noted that the term “a” or “an” refers to one or more. As such, the terms “a” (or “an”), “one or more,” and “at least one” are used interchangeably herein.

**[0083]** The words “comprise”, “comprises”, and “comprising” are to be interpreted inclusively rather than exclusively. The words “consist”, “consisting”, and its variants, are to be interpreted exclusively, rather than inclusively. While various embodiments in the specification are presented using “comprising” language, under other circumstances, a related embodiment is also intended to be interpreted and described using “consisting of” or “consisting essentially of” language.

**[0084]** As used herein, the term “about” means a variability of 10% (±10%) from the reference given, unless otherwise specified.

**[0085]** The term “regulation” or variations thereof as used herein refers to the ability of a compound of formula (I) to inhibit one or more components of a biological pathway.

**[0086]** A “subject” is a mammal, e.g., a human, mouse, rat, guinea pig, dog, cat, horse, cow, pig, or non-human primate, such as a monkey, chimpanzee, baboon or gorilla.

**[0087]** As used herein, “disease”, “disorder” and “condition” are used interchangeably, to indicate an abnormal state in a subject.

**[0088]** Unless defined otherwise in this specification, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art and by reference to published texts, which provide one skilled in the art with a general guide to many of the terms used in the present application.

**[0089]** The following examples are illustrative only and are not intended to limit the present invention.

#### Example 1—scAAV Vectors Containing hOTC

**[0090]** pAAVsc.TBG.hOTCwt and pAAVsc.TBG.hOTCco-LW4 were constructed by replacing the mOTC coding sequencing with wild-type (WT) hOTC (hOTCwt) or hOTCcoLW cDNA, respectively, in a plasmid derived from the previously described pAAVsc.TBG.mOTC1.3 with the intron disrupted [Moscioni D, et al, “Long-term correction of ammonia metabolism and prolonged survival in ornithine transcarbamylase-deficient mice following liver-directed treatment with adeno-associated viral vectors”, *Mol Ther.* 2006; 14:25-33; Cunningham S C, et al, “AAV2/8-mediated correction of OTC deficiency is robust in adult but not neonatal Spf(ash) mice”, *Mol Ther.* 2009; 17:1340-1346; Wang L, et al., “Sustained correction of OTC deficiency in spf<sup>ash</sup> mice using optimized self-complementary AAV2/8 vectors”, *Gene Ther.* 2012 April; 19(4):404-10, Epub 2011 Aug. 18].

**[0091]** The scAAV2/8.TBG.hOTCco-LW4 contains an AAV2 3' ITR and a 5' ITR with a deletion in the D-sequence and trs (terminal resolution site), a TBG promoter, the hOTCco-LW4 gene, and a 137 bp SV40 polyA. The two vector preps (AAV2/8sc.TBG.hOTCwt and AAV2/8sc.TBG.hOTCco-LW4) used in the initial comparison study were purified by two rounds of cesium chloride gradient centrifugation, as previously described [Wang L, et al, Systematic evaluation of AAV vectors for liver directed gene transfer in murine models. *Mol Ther.* 2010; 18:118-125]. Vectors used in the rest of the study were produced by a scaled production method based on polyethylenimine (PEI) transfection and purified from supernatant or total lysate by iodixanol gradient centrifugation as described [Lock M, et al, *Hum Gene Ther.* 2010; 21:1259-1271]. Genome titers [genome copies (GC)/ml] of AAV vectors were determined by real-time PCR using primer and probe sets targeting the TBG promoter

(forward primer 5'-AAACTGCCAATCCACTGCTG-3' [SEQ ID NO: 14], reverse primer 5'-CCAT-AGGCAAAGCACCAAGA-3' [SEQ ID NO: 15], probe 6FAM-TTGGCCCAATAGTGAGAACTTTTCCTGC [SEQ ID NO: 16]-TAMRA), and using a linearized plasmid as the standard. The forward primer is located 400 bp downstream of the 5' closed hairpin. Fagone et al [Systemic errors in quantitative polymerase chain reaction titration of self-complementary adeno-associated viral vectors and improved over alternative methods, Hum Gene Ther Methods. 2012 February; 23(1):1-7.] recently reported that the quantitative PCR (Q-PCR) method could significantly underestimate the titer of scAAV vectors, especially when the PCR primers were close to the closed hairpin of the scAAV vector. The titer of one lot of AAV2/8sc.TBG.hOTCco-LW4 vector using a primer and probe set targeting the polyA region (1900 bp downstream of the 5' closed hairpin), and the genome titer was 1.1-fold of the original titer, which was within the intra-assay error of Q-PCR.

**[0092]** OTC protein expression levels and OTC activity were evaluated in the liver of  $spf^{zsh}$  mice 14 days after i.v. injection of  $1 \times 10^{11}$  GC of AAV2/8sc.TBG.hOTCwt or AAV2/8sc.TBG.hOTCco-LW4 vectors. The  $spf^{zsh}$  mice are a model for late onset OTC disease in humans. All animal procedures were performed in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Pennsylvania.  $Spf^{zsh}$  mice were maintained at the Animal Facility of the Translational Research Laboratories at the University of Pennsylvania as described previously. Three to six months old  $spf^{zsh}$  mice and their normal littermates were used in the studies. Vectors were administered by intravenous (i.v.) injection via the tail vein. The extent of gene transfer based on resident vector genomes was not statistically different between the two groups. Urine samples were collected before and at various time points after vector treatment for orotic acid analysis as previously described [Moscioni D, et al, Long-term correction of ammonia metabolism and prolonged survival in ornithine transcarbamylase-deficient mice following liver-directed treatment with adeno-associated viral vectors. Mol Ther. 2006; 14:25-33].

**[0093]** Western blot analysis to detect hOTC expression in liver lysate was performed as previously described [Wang L, et al, 2012, epub 2011]. The primary antibody to detect hOTC was a custom rabbit polyclonal antibody provided by Hiroki Morizono's laboratory at the Children's National Medical Center. Liver lysates (10  $\mu$ g/lane) were also blotted and probed by an anti-tubulin antibody (Abcam, Cambridge, MA). Western analysis demonstrated 100-fold higher expression of hOTC from the hOTCco-LW4 vector as compared to the hOTCwt vector, reaching levels in slight excess of those seen in WT mice.

**[0094]** OTC enzyme activity was measured using a liquid chromatography mass spectrometry stable isotope dilution method to detect the formation of citrulline normalized to [1,2,3,4,5-<sup>13</sup>C<sub>5</sub>] citrulline (98% <sup>13</sup>C). The method is adapted from a previously developed assay for detection of N-acetylglutamate synthase activity [Morizono H, et al, Mammalian N-acetylglutamate synthase. Mol Genet Metab. 2004; 81(Suppl 1):54-11.]. Slivers of fresh frozen liver were weighed and briefly homogenized in buffer containing 10 mM HEPES, 0.5% Triton X-100, 2.0 mM EDTA and 0.5 mM DTT. Volume of homogenization buffer was adjusted to obtain 50 mg/ml tissue. Enzyme activity was measured

using 250  $\mu$ g liver tissue in 50 mM Tris-acetate, 4 mM ornithine, 5 mM carbamyl phosphate, pH 8.3. Enzyme activity was initiated with the addition of freshly prepared 50 mM carbamyl phosphate dissolved in 50 mM Tris-acetate pH 8.3, allowed to proceed for 5 minutes at 25° C. and quenched by addition of an equal volume of 5 mM 13C<sub>5</sub>-citrulline in 30% TCA. Debris was separated by 5 minutes of microcentrifugation, and the supernatants were transferred to vials for mass spectroscopy. Ten  $\mu$ L of sample was injected into an Agilent 1100 series LC-MS under isocratic conditions with a mobile phase of 93% solvent A (1 ml trifluoroacetic acid in 1 L water):77% solvent B (1 ml trifluoroacetic acid in 1 L of 1:9 water/acetonitrile). Peaks corresponding to citrulline [176.1 mass:charge ratio (m/z)] and 13C<sub>5</sub>-citrulline (181.1 m/z) were quantitated, and their ratios were compared to ratios obtained for a standard curve of citrulline run with each assay. Samples were normalized to either total liver tissue or to protein concentration determined using a Bio-Rad protein assay kit (Bio-Rad, Hercules, CA).

**[0095]** The vector carrying engineered hOTC cDNA termed herein LW4 (FIG. 4) was found to improve expressed hOTC protein levels by 100-fold. An assessment of OTC enzyme activity generally correlated with the OTC Western blot experiments although OTC protein was more elevated than OTC enzyme activity when compared to endogenous OTC. When subtracting the background activity levels in the  $spf^{zsh}$  mice, the hOTCco-LW4 resulted in over 33-fold higher activity than the hOTCwt. Sustained and dose-correlated hOTC expression and activity levels were observed in the treated  $spf^{zsh}$  mice. Compared to a previously described murine OTC vector which differed mainly in the cDNA, the vector carrying the hOTCco-LW4 vector was about 10-fold more potent.

**[0096]** The illustrative vector carrying the modified hOTCco-LW4 (FIG. 4) provided high level of transduction, as measured by OTC histological assays, throughout a broad range of doses. Between doses  $1 \times 10^{11}$  GC and  $3 \times 10^9$  GC, transduction efficiency, as measured by histochemical staining, varied between 50-70%. At the lowest dose of  $1 \times 10^9$  GC, 40% of the liver areas were positive by OTC histochemical staining. The lack of a clear dose effect by histochemistry and immunostaining could be due to the fact that codon optimization significantly improved hOTC expression in the transduced hepatocytes. This leads to improved sensitivity to detect transduced cells with low vector genome copies. Transduction could be saturated with high vector doses ( $1 \times 10^{11}$ - $1 \times 10^{10}$  GC), and therefore transduction efficiency measured by in situ detection methods would not discriminate between low and high dose groups in contrast to OTC enzyme activity on liver lysates measured by mass spectrometry.

**[0097]** A further study was performed in which neonatal expression of hOTC was assessed in  $spf^{zsh}$  mice, injected on day 1 of life, using the scAAV2/8.TBG.hOTCco at a dose of  $5 \times 10^{10}$  GC/pup injected via the temporal vein. Robust expression was detected at 24 and 48 hours. Additional studies were performed using doses of  $1 \times 10^{11}$ ,  $3 \times 10^{10}$ , and  $1 \times 10^{10}$ , and assessed for 12 weeks. Over the 16 week period of the study, a reduction in the initial robust expression levels was observed at each of the doses. This is believed to be due to dilution, i.e., a natural result of the proliferation of liver cells in growing animals. Thus, while initial restoration of OTC liver activity is observed following neonatal gene

transfer in  $sp^{fzsh}$  mice, this result is temporary, with OTC activity dropping from about 1000% of wild-type (wt) levels at about 1 week, to about 50% of wt levels at 4 weeks, to about 10% of wt levels at 12 weeks ( $1 \times 10^{11}$  GC level); or about 500% of wt levels at week 1, to about 20% of wt levels, or about 10% of wt levels at week 1 ( $3 \times 10^{10}$  GC dose) or about 200% wt levels at week 1, to about 10% wt levels at week 4 ( $1 \times 10^{10}$  GC dose). In one study, using animals receiving the first injection of  $3 \times 10^{10}$  GC at day 1, animals were injected with a second AAV vector carrying the hOTCco gene (scAAVrh10.hOTCco;  $1.8 \times 10^{10}$  GC) at week 4. As a control, one group of animals received no readministration and one group received only the second vector at 4 weeks. Readministration of the AAV.hOTCco resulted in restoration of liver OTC activity.

**[0098]** Further studies were designed to assess the ability to rescue OTC-KO pups by neonatal gene therapy, both short-term and long-term.

#### Example 2—Production of scAAV Vectors Having Codon Optimized Sequences

**[0099]** A. scAAV8.TBG.hOTC-co

**[0100]** Plasmids containing a codon optimized hOTC sequence of SEQ ID NO: 3, 4, 5, 9 or 10, respectively, are cloned as described in Example 1 by replacing the mOTC coding sequencing with hOTCco in a plasmid derived from the previously described pAAVsc.TBG.mOTC1.3 with the intron. The resulting plasmid pAAVsc.TBG.hOTCco is cloned into an AAV8 capsid [Gao et al, PNAS USA, 2002, 99:11854-11859] using conventional techniques.

**[0101]** B. scAAVrh10.TBG.hOTC-co

**[0102]** Plasmids containing a codon optimized hOTC sequence of SEQ ID NO: 3, 4, 5, 9 or 10, respectively, are cloned as described in Example 1 by replacing the mOTC coding sequencing with hOTCco in a plasmid derived from the previously described pAAVsc.TBG.mOTC1.3 with the intron. The resulting plasmids pAAVsc.TBG.hOTCco are cloned into an AAVrh10 capsid [Gao et al, PNAS USA, 2002, 99:11854-11859] using conventional techniques.

#### Example 3—Production of ssAAV Vectors Having Codon Optimized Sequences

**[0103]** ssAAV2/8.LSP1.hOTC-co

**[0104]** Plasmids containing the codon optimized hOTCco sequences are cloned as described by replacing the mOTC coding sequencing of the pLSP1mOTC plasmid [Cunningham et al, Mol Ther, 2009, 17: 1340-1346] with the corresponding cDNA sequence of SEQ ID NO:3, 4, 5, 9 or 10. The resulting plasmids pAAVsc.LSP1.hOTCco are cloned into AAV8 capsids to form the corresponding ssAAV2/8.LSP1.hOTC-co vectors using techniques described in Example 1.

**[0105]** B. ssAAV2/rh10.LSP1. hOTC-co

**[0106]** Plasmids containing the codon optimized hOTCco sequences are cloned as described by replacing the mOTC coding sequencing of the  $\mu$ LSP1mOTC plasmid [Cunningham et al, Mol Ther, 2009, 17: 1340-1346] with the corresponding cDNA sequence of SEQ ID NO:3, 4, 5, 9 or 10. The resulting plasmids pAAVsc.LSP1.hOTCco are cloned into AAV8 capsids to form the corresponding ssAAV2/8.LSP1.hOTC-co vectors using techniques described in Example 1.

**[0107]** The vectors generated according to Part A or B may be purified by two rounds of cesium chloride gradient centrifugation, buffered-exchanged with PBS, and concentrated using Amicon Ultra 15 centrifugal filter devices-100K (Millipore, Bedford, MA). Genome titer (GC/ml) of AAV vectors can be determined by real-time PCR using a primer/probe set corresponding to the TBG promoter and linearized plasmid standards. Vectors can be subject to additional quality control tests including sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis for vector purity and Limulus ameocyte lysate (LAL) for endotoxin detection (Cambrex Bio Science, Walkersville, MD, USA).

#### Example 4—ssAAV8.TBG.hOTCco in Model of Late Onset of OTCD

**[0108]** An AAV8 vector was generated using the methods described herein. The vector has packaged therein a 5' AAV2 ITR, a TBG promoter, an intron, a hOTCco, a WPRE element, a bovine growth hormone polyA, and a 3' AAV2 ITR. The expression and kinetics of this vector was compared to a self-complementary AAV8 vector with or without the WPRE element. The results show that the single-stranded constructs with the WPRE element outperformed those vectors lacking the WPRE element; at comparable doses both single-stranded vectors (with and without WPRE) were less robust than the self-complementary vector lacking WPRE in the time points measured.

**[0109]** However, the single-stranded vectors may have other desirable properties, e.g., in terms of kinetics, depending upon the age and condition of the patient.

#### Example 5—Production of Adenovirus Vectors Having Codon Optimized Sequences

**[0110]** hOTCco cDNA [SEQ ID NO: 3, 4, 5, 9 and 10] with NotI linkers is cloned downstream of a rat PEPCK promoter to generate pPEPCK-hOTC as described in Mian et al, Molecular Therapy, 2004, 10: 492-499 (2004). This plasmid is digested with AscI, and the resultant PEPCK-hOTCco fragment is inserted into the adenoviral backbone plasmid pC4HSU31 to generate the parental plasmids pC4HSU-PEPCK-hOTCco. Plasmid pWPRE is digested with ClaI to release the WPRE, which is then inserted into the MluI site of pPEPCK-hOTC, to generate pPEPCK-hOTCco-WPRE plasmid with their respective hOTCco sequences. The remaining steps to generate the adenoviral plasmid pC4HSU-PEPCK-hOTCco-WPRE are as previously described. All cloning sites are confirmed by DNA sequence analysis. The identity of recombinant adenoviral plasmids can be confirmed by restriction enzyme digestion with HindIII and BamHI. The adenoviral plasmids are linearized with PmeI before transfection into 293Cre4 cells. Adenoviral vectors are rescued and amplified with 293Cre4 cells and helper virus AdLC8cluc. Suspension 293N3Scre8 cells may be used in the final step of vector production. Purification, quantification by OD260 and viral DNA extraction are performed as described in detail elsewhere [Brunetti-Pierri, N., et al. (2004). Acute toxicity after high-dose systemic injection of helper-dependent adenoviral vectors into nonhuman primates. Hum. Gene Ther. 15: 35-46; Ng, P., Parks, R. J. and Graham, F. L. (2002). Preparation of helper-dependent adenoviral vectors. *Methods Mol. Med.* 69: 371-388].

#### Example 6—Production of hOTCco Lentiviral Vectors

**[0111]** A. Replication-defective lentiviral vectors containing the hOTCco sequences provided herein can be produced by replacing the rat OTC gene sequence insert of the plasmid pLenti-GIII-CMV-GFP-2A-Puro [commercially available from Applied Biological Materials (ABM) Inc.; Canada] with the desired hOTCco sequence [SEQ ID NO: 3, 4, 5, 9 and 10]. The viruses are generated according to manufacturer instructions. The ABM system includes an enhancer deletion in the U3 region of 3'ΔLTR to ensure self-inactivation of the lentiviral vector following transduction and integration into the target cell's genomic DNA; contains minimal lentiviral genes necessary for packaging, replication and transduction (Gag/Pol/Rev), derived from different plasmids all lacking packaging signals; further, none of the Gag, Pol, or Rev genes are incorporated into the packaged viral genome, thus making the mature virus replication-incompetent.

**[0112]** B. Replication-Defective, Non-Integrating hOTC Lentiviral Vectors

**[0113]** A DNA construct containing a liver specific promoter and the hOTCco DNA of SEQ ID NO: 3, 4, 5, 9 and 10 are engineered into lentivirus vectors which are pseudotyped into sindbis virus E2 enveloped produced as described in US2011/0064763, which is incorporated by reference herein. All vectors contain splice donor, packaging signal (psi), a Rev-responsive element (RRE), splice donor, splice acceptor, central poly-purine tract (cPPT). The WPRE element is eliminated in certain viruses.

**[0114]** C. The hOTCco DNA of SEQ ID NO: 3, 4, 5 9 and 10, is cloned into a lentivirus pseudotyped with a vesicular stomatitis virus (VSV) envelope gene, purchased from Invi-vogen (SanDiego, CA) using manufacturer's instructions.

#### Example 7—Production hOTCco RNA Delivery Systems

**[0115]** RNA may be prepared by in vitro transcription from a DNA template or synthesized. The RNA expression cassette is prepared which includes a 5' UTR, an optional intron with splice donor and acceptor sites, an optional Kozak sequence, the hOTC coding sequence provided herein, a polyA, and a 3' UTR using known techniques.

**[0116]** A. A suitable amount of mRNA are incorporated into a lipid-enveloped pH-responsive polymer nanoparticles generated using published techniques. [X. Su et al, Mol. Pharmaceutics, 2011, 8 (3), pp 774-787; web publication: Mar. 21, 2011].

**[0117]** B. Polymeric nanoparticle formulations with 25 kDa branched polyethyleneimine (PEI) are prepared as follows. When PEI is present, it may be branched PEI of a molecular weight ranging from 10 to 40 kDa, e.g., 25 kDa branched PEI (Sigma #408727). Additional exemplary polymers suitable for the present invention include those described in PCT Publication WO2013182683, the contents of which is hereby incorporated by reference. The required amount of mRNA is diluted just before application in water for injection (Braun, Melsungen) to a total volume of 4 ml and added quickly to 4 ml of an aqueous solution of branched PEI 25 kDa using a pipette at an N/P ratio of 10. The solution is mixed by pipetting up and down.

**[0118]** C. For a lipid-based nanoparticle, a lipid formulation is created using expression cassette containing the

hOTCco RNA in a formulation of cK-E12:DOPE:Chol:PEG-DMG2K (relative amounts 50:25:20:5 (mg:mg:mg:mg)) to provide a solution for delivery. The cationic lipid cK-E12 is used (see, e.g., WO 2013/063468), and is combined with dioleoylphosphatidyl-ethanolamine or "DOPE", cholesterol (chol), and polyethylene glycol (PEG) or a PEGylated lipid (PEG-DMG2K) using e formulation methods described in international patent publications WO 2010/053572 and WO 2012/170930, both of which are incorporated herein by reference.

#### Example 8—hOTCco DNA Delivery Systems

**[0119]** A. Naked Plasmid DNA—The hOTCco sequences [SEQ ID NO: 3, 4, or 5] are engineered as naked plasmid DNA constructs which are delivered to a target liver cell (e.g., via intravascular administration) and express the human OTC protein in the target cell.

**[0120]** B. Cationic lipid-DNA complexes—Cationic lipid—DNA complexes are prepared using a suitable amount of an expression cassette containing at least a promoter, an optional intron, an optional Kozak sequences, an hOTCco of SEQ ID NO: 3, 4 or 5, a polyA, and other optional expression control sequences. The promoter may be a liver specific promoter. Alternatively, another non-tissue specific promoter may be selected. For example, a suitable amount of DNA is formulated with a cationic lipid of cK—E12:DOPE:Chol:PEG-DMG2K (relative amounts 50:25:20:5 (mg:mg:mg:mg)) to form a cationic lipid—DNA complex suitable for delivery to a subject. The cationic lipid cK-E12 is used (see, e.g., WO 2013/063468), and is combined with dioleoylphosphatidyl-ethanolamine or "DOPE", cholesterol (chol), and polyethylene glycol (PEG) or a PEGylated lipid (PEG-DMG2K) using e formulation methods described in international patent publications WO 2010/053572 and WO 2012/170930, both of which are incorporated herein by reference.

#### Example 9 Long-term Correction of a Neonatal Lethal Form of OTC Deficiency by Multiple Treatments with AAV Vectors of Different Serotypes

**[0121]** In the current study, the scAAV8.TBG.hOTCcoLW4 prepared as described in Example 1 was used to rescue animals in a mouse model of neonatal (early) onset OTCD. OTC KO mice were generated through the deletion of exons 2-3, and the properties of this mouse characterized in terms of similarity to human patients with null mutations of OTC. In summary, the OTC knockout (KO) model generated in our laboratory through the deletion of exons 2-3 closely mimics the severe neonatal onset form of OTCD in humans. Neonatal male OTC KO pups have elevated plasma ammonia levels due to the absence of OTC expression in the liver, and they inevitably die within 24 hours after birth. Heterozygous females breed normally, have normal plasma ammonia levels, reduced liver OTC enzyme activity, elevated urine orotic acid levels, and in some cases lower body weight compared to wild type (WT) littermates. A single injection of scAAV8-hOTCco vector prepared as described in Example 1 at a dose of 1-3×10<sup>10</sup> GC/pup immediately after birth is able to rescue the OTC KO pups and extend the life to 6 weeks. To achieve long-term correction, a group of 4-week-old OTC-KO mice received a

second vector administration of scAAVrh10-hOTCco vector, which had been prepared as described in Example 1.

**[0122]** Over 30 OTC-KO pups retrieved by Cesarean section have been successfully rescued with gene delivery. The rescued pups had lower body weight than their normal littermates and had a transient phenotype of sparse fur and abnormal skin. Most importantly, their plasma ammonia levels were in the normal range. However, the efficacy cannot be maintained beyond 6 weeks due to loss of vector genome during fast liver proliferation in neonatal stage. A second vector administration of scAAVrh10-hOTCco vector in 4-week-old OTC-KO mice is able to further extend their lives to adulthood. The oldest mice have reached 18 months of age. The long-term rescued mice show close to normal levels of plasma ammonia, although urine orotic acid levels in a subset of these mice were significantly elevated. Sirius red staining on liver samples from heterozygous mice of different ages (6, 12, and 18 months old) showed liver fibrosis in aged (18-month old) OTC-KO heterozygous female mice, similar to a liver sample from a 11-year-old OTCD patient.

#### Example 10—Treatment of Late Onset OTC Deficiency (OTCD)

**[0123]** Two-month old OTC-KO heterozygous mice received a single tail vein injection of a self-complementary AAV8 vector encoding a codon-optimized human OTC gene (SEQ ID NO: 5) at  $1 \times 10^{10}$ ,  $3 \times 10^{10}$ , and  $1 \times 10^{11}$  vector genome copies per mouse. One week following vector treatment, mice in all three vector dose groups had normal urine orotic acid levels which were maintained throughout the study (16 months). Liver samples were harvested from 18 month old treated mice for pathology analysis and compared to age-matched untreated heterozygous mice and WT littermates. All treated mice showed normal liver histology similar to WT, in contrast to the untreated heterozygous animals which had fibrosis throughout the liver. In conclusion, a single injection of AAV8sc-hOTCco vector can prevent liver fibrosis in OTC-KO heterozygous and has great potential for correction of liver fibrosis in OTCD patients.

**[0124]** Gene therapy vectors described herein are capable of rapid, robust and prolonged gene expression even in mice with a complete lack of OTC. Heterozygous females are able to reproduce and deliver hemizygous male offspring, but these pups die within a day of birth if untreated. Untreated old heterozygous female mice show evidence of increased fibrosis and microvesicular steatosis, a finding that appears

similar to observations in human heterozygous patients. A regimen of gene transfer that is able to rescue affected males has been developed and treated males have survived over 72 weeks.

**[0125]** Thus, these data demonstrate that liver-specific gene therapy with hOTC can prevent liver fibrosis. These data correlate with in treatment of heterozygous OTC deficient humans, e.g., subjects having late onset of OTCD.

#### Sequence Listing Free Text

**[0126]** The following information is provided for sequences containing free text under numeric identifier <223>.

SEQ ID NO: (containing free text)	Free text under <223>
3	Engineered hOTC
4	Engineered hOTC
5	Engineered hOTC
6	Plasmid pscAAVTBGhOTCLW
8	Engineered hOTC
9	Engineered hOTC
10	Engineered hOTC RNA
11	Engineered hOTC RNA
12	Engineered hOTC RNA
13	Engineered hOTC RNA
14	PCR forward primer
15	PCR reverse primer
16	Probe
17	Engineered hOTC cDNA termed LW4
18	Engineered hOTC cDNA termed LW3

**[0127]** This application contains a Sequence Listing labeled “14-7037C2US\_ST25.txt” dated Aug. 17, 2020 which is hereby incorporated by reference. The priority U.S. Provisional Patent Application No. 61/950,157, filed Mar. 9, 2014, PCT Application No. PCT/US15/19513, filed on Mar. 9, 2015, U.S. patent application Ser. No. 15/122,853, filed on Aug. 31, 2016, U.S. patent application Ser. No. 16/191,709, filed on Nov. 15, 2018, and all published documents cited in this specification are incorporated herein by reference. Similarly, the SEQ ID NO which are referenced herein and which appear in the appended Sequence Listing are incorporated by reference. While the invention has been described with reference to particular embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

#### SEQUENCE LISTING

```

Sequence total quantity: 18
SEQ ID NO: 1      moltype = DNA  length = 1062
FEATURE          Location/Qualifiers
source           1..1062
                 mol_type = other DNA
                 organism = Homo sapiens
CDS
SEQUENCE: 1
atgctgttta atctgaggat cctgttaaac aatgcagcct ttagaaatgg tcacaacttc 60
atggttcgaa attttcggtg  tggacaacca ctacaaaata aagtgcagct gaagggccgt 120
gaccttctca ctctaaaaaa ctttaccgga gaagaaatta aatatatgct atggctatca 180
gcagatctga aatttaggat aaaacagaaa ggagagtatt tgccctttatt gcaagggaag 240
tccttaggca tgatttttga gaaaagaagt actgaacaa gattgtctac agaaacaggc 300

```



-continued

```

tttgacttc tgggaggaca tccttgttt cttaccacac aagatattca tttgggtgtg 360
aatgaaagtc tcacggacac ggccccgtgta ttgtctagca tggcagatgc agtattggct 420
cgagtgtata aacaatcaga tttggacacc ctggctaaag aagcatccat cccaattatc 480
aatgggctgt cagatttgta ccatcctatc cagatcctgg ctgattacct cacgctccag 540
gaacactata gctctctgaa aggtctttacc ctcagctgga tcggggatgg gaacaatatc 600
ctgactcca tcctgatgag cgcagcgaaa ttcggaatgc accttcaggc agctactcca 660
aagggttatg agccggatgc tagtgaacc aagttggcag agcagtatgc caaagagaat 720
ggtagcaagc tgttctgac aaatgatcca ttggaagcag cgcattggagg caatgtatta 780
attacagaca cttggataag catgggacaa gaagaggaga agaaaaagcg gctccaggct 840
ttccaaggtt accaggttac aatgaagact gctaaagttg ctgcctctga ctggacattt 900
ttacactgct tgcccagaaa gccagaagaa gtggatgatg aagtctttta ttctcctcga 960
tcactagtggt tcccagagcg agaaaaacaga aagttggacaa tcattggctgt catgggtgctc 1020
ctgctgacag attactcacc tcagctccag aagcctaaat tt 1062

```

```

SEQ ID NO: 2          moltype = AA length = 354
FEATURE              Location/Qualifiers
source                1..354
                     mol_type = protein
                     organism = Homo sapiens

```

```

SEQUENCE: 2
MLFNLRILLN NAAFRNGHNF MVRNFRCGQP LQNKVQLKGR DLLTLKNFTG EEIKYMLWLS 60
ADLKFRIRQK GEYLPPLQ GK SLGMIPEKRS TRRLSTETG FALLGGHPCF LTTQDIHLGV 120
NESLTD TARV LSSMADAVLA RVYKQSDLDL LAKEASIIPII NGLSDLYHPI QILADYLTQL 180
EHYSSLKGLT LSWIGDGNNI LHSIMMSAAK FGMHLQAATP KGYEPDASVT KLAEQYAKEN 240
GTKLLLTNDP LEAAHGDNVL ITDTWISMGO EEEKKRRLQA FQGYQVTMKT AKVAASDWTFF 300
LHCLPRKPEE VDDEVFYSR SLVFPBAENR KWTIMAVMVS LLTDYSPQLQ KPKF 354

```

```

SEQ ID NO: 3          moltype = DNA length = 1068
FEATURE              Location/Qualifiers
misc_feature          1..1068
                     note = engineered hOTC
source                1..1068
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 3
atgctgttca acctcgcaat cctgctgaac aatgccgctt ttcggaacgg gcacaatttc 60
atggtgagga actttcgctg cggacagccc ctccagaaca aggtccagct gaagggcagg 120
gacctgctga ccctgaaaaa ttccacaggg gaggaaatca agtacctgct gttgctgtca 180
gccgatctga agttccggat caagcagaag ggcgaatcgc tgcctctgct ccagggcaaa 240
agcctgggga tgatctctga aaagcgcagt actcggacca gactgtcaac agagactgga 300
ttcgactgca tgggaggaca cccatgtttt ctgaccacac aggcatttca tctgggagtg 360
aacgagtcct tgaccgacac agcaccgctc ctgagctcca tggctgatgc agtctggct 420
cgagtctaca aacagtctga cctggatacc ctggccaagg aagcttctat cccaatcatt 480
aatggcctga gtgacctgta tcaccccatc cagattctgg ccgattacct gaccctccag 540
gagcattatt ctagtctgaa agggctgaca ctgagctgga ttggggacgg aaacaatatc 600
ctgactcca ttatgatgag cgcgcccaag ttggaatgc acctccaggc tgcaacccca 660
aaaggctacg aaccggatgc ctccgtgaca aagctggcag aacagtatgc caaagagaac 720
ggcactaacg tgctgctgac caatgacct ctggaggccg ctcacggagg caactgtgct 780
atcactgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccaggcc 840
ttccagggct accaggtgac aatgaaaact gctaaggtcg cagccagcga ctggacctt 900
ctgactgccc tgcccagaaa gccctaagag gtggacgatg aggtcttcta ctaccaccaga 960
agcctgggtg ttctctgagc tgcagaatagg aagttggacaa tcattggcagt gatggtcagc 1020
ctgctgactg attattcccc tcagctccag aaaccaaagt tctgataa 1068

```

```

SEQ ID NO: 4          moltype = DNA length = 1068
FEATURE              Location/Qualifiers
misc_feature          1..1068
                     note = engineered hOTC coding sequence
source                1..1068
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 4
atgctgttca acctcgcaat cctgctgaac aacgccgctt ttcggaacgg gcacaacttt 60
atggtgagga actttcgctg cggacagccc ctccagaata aggtccagct gaagggcagg 120
gacctgctga ccctgaaaaa ttccacaggg gaggaaatca agtatactgct gttgctgtca 180
gctgatctga agttccggat caagcagaag ggcgaatcgc tgcctctgct ccagggcaaa 240
agcctgggga tgatctctga aaagcgcagt actcggacca gactgtcaac cgagactgga 300
ttcgctctgc tgggaggaca cccttgtttt ctgaccactc aggcatttca cctgggagtg 360
aacgagtcct tgaccgacac tgctcgctc ctgagctcta tggccgacgc tgtgctagct 420
cgagtctaca aacagtcgta cctggatacc ctggccaagg aagcttctat cccaattatt 480
aacggcctgt cagacctgta tcaccccatc cagattctgg ccgattacct gaccctccag 540
gagcactatt ctagtctgaa agggctgaca ctgagttgga ttggggacgg aaacaatatc 600
ctgactcta ttatgatgac agccgccaag ttggaatgc acctccaggc tgcaacccca 660
aaaggctacg aaccggatgc ctccgtgaca aagctggctg aacagtatgc caaagagaac 720
ggcactaacg tgctgctgac caacgacct ctggaggccg ctcacggagg caactgtgct 780
atcaccgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccaggcc 840

```

-continued

```

ttccagggt accaggtgac aatgaaaacc gctaaggctg cagccagcga ttggacctt 900
ctgcaactgcc tgcccagaaa gcccgaaagag gtggacgacg aggtcttcta ctctcccaga 960
agcctgggtg ttcccgaagc tgagaatagg aagtggacaa ttatggcagt gatggtcagc 1020
ctgctgactg attattcacc tcagctccag aaaccaaagt tctgataa 1068

```

```

SEQ ID NO: 5          moltype = DNA length = 1068
FEATURE
misc_feature         1..1068
                    note = engineered hOTC
source              1..1068
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 5
atgctgttca acctgccaat cctgctgaa aacgcccgtt ttggaaacgg gcacaacttt 60
atggtgagga accttcgctg cggacagccc ctccagaata aggtccagct gaagggcagg 120
gacctgctga ccttcaaaaa ttccacaggg gaggaatca agtatatgct gtggctgtca 180
gctgatctga agttccggat caagcagaag gccgaatata tgctctgct ccagggcaaa 240
agcctgggga tgatcttoga aaagcgcagt actcggacca gactgtcaac cgagactgga 300
ttcgctctgc tgggaggaca cccttgtttt ctgacocactc aggcattca cctgggagtg 360
aacgagtcctc tgaccgacac tgctcgcgtc ctgagctcta tggccgacgc tgtgctggct 420
cgagtctaca aacagtcoga cctggatacc ctggccaagg aagcttctat cccaattatt 480
aacggcctgt cagacctgta tcaccccatc cagattctgg ccgattacct gaccctccag 540
gagcactatt ctagtctgaa agggctgaca ctgagttgga ttggggacgg aaacaatatc 600
ctgcaactta ttatgatgct agccgccaag ttggaatgc acctccaggc tgcaacccca 660
aaaggctacg aaccgatgc ctccagtaca aagctggctg aacagtagcg caaagagAAC 720
ggcactaacg tgctgctgac caacgacct ctggaggccg ctcacggagg caacgtgctg 780
atcaccgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccagggc 840
ttccagggtc accaggtgac aatgaaaacc gctaaggctg cagccagcga ttggacctt 900
ctgcaactgcc tgcccagaaa gcccgaaagag gtggacgacg aggtcttcta ctctcccaga 960
agcctgggtg ttcccgaagc tgagaatagg aagtggacaa ttatggcagt gatggtcagc 1020
ctgctgactg attattcacc tcagctccag aaaccaaagt tctgataa 1068

```

```

SEQ ID NO: 6          moltype = DNA length = 5195
FEATURE
misc_feature         1..5195
                    note = Plasmid pscAAVTBGhOTCLW
repeat_region       5..109
                    note = ITR
regulatory           851..854
                    regulatory_class = TATA_box
misc_feature         2046..2182
                    note = polyA - ITR, located on complement
repeat_region       2211..2378
                    note = ITR, located on complement
misc_feature         4172..4760
                    note = TBG/promoter
source              1..5195
                    mol_type = other DNA
                    organism = synthetic construct
CDS                 976..2037
                    note = hOTCco-LW4

```

```

SEQUENCE: 6
taggtcgcgc gctcgcctgc tcaactgaggc cgccccgggca aagcccgggc gtcggggcgc 60
ctttggctgc ccggcctcag tgagcgagcg agcgcgcaga gagggagtg agccatgctc 120
taggaagatc aattcaattc acgctgggta cctagaacta tagctagaat tcgcccttaa 180
gctagcaggt taatttttaa aaagcagtc aagttccaag tggcccttgg cagcatttac 240
tctctctggt tgctctgggt aataatctca ggagcacaac cattccagat ccagggtaat 300
tttataaaag cagtcaaaaag tccaagtggc ccttggcagc atttactctc tctggttgc 360
ctggttaata atctcaggag cacaaacatt ccagatccgg cgcgccaggg ctggaagcta 420
cctttgacat cacttctcct gcgaatgcat gtataatttc tacagaacct attagaaagg 480
atcaccacgc ctctgctttt gtacaacttt cccttaaaaa actgccaatt ccaactgctg 540
ttggcccaat agtgagaact ttttctgct gcctcttggg gcttttgctc atggccctta 600
ttctgcctgc tgaagacact ctggccagca tggacttaaa cccctccagc tctgacaatc 660
ctctttctct tttgttttac atgaagggtc tggcagccaa agcaatcact caaagttaa 720
accttatcat ttttgctttt gttcctcttg gcttgggttt tgtacatcag ctttgaaat 780
accatcccag ggttaatgct ggggttaatt tataactaag agtgctctag ttttgcaata 840
caggacatgc tataaaaaatg gaaagatgtt gctttctgag agacagcttt attgcccgtag 900
ttatcacag ttaaattgct aacgcagtc gtgcttctga cacaaacagtc tcgaacttaa 960
gctgcagccg ccaccatgct gttcaacctg cgaatcctgc tgaacaacgc cgcttttcgg 1020
aacgggcaca actttatggt gaggaacttt cgctcggcag agcccctcca gaataagggtc 1080
cagctgaaagg gcagggagct gctgaccctg aaaaatttca caggggagga aatcaagat 1140
atgctgtggc tgctcagctga tctgaaagtc cggatcaagc agaagggcga atatctgcct 1200
ctgctccagg gcaaaagcct ggggatgatc ttcgaaaagc gcagtactcg gaccagactg 1260
tcaaccgaga ctggattcgc tctgctggga ggacaccctt gttttctgac cactcaggac 1320
attcacctgg cagtgaacga gtccctgacc gacactgctc gcgtcctgag ctctatggcc 1380
gacgctgtgc tggctcaggt ctacaaacag tccgacctgg atacctggc caaggaagct 1440

```

-continued

```

tctatcccaa ttattaacgg cctgtcagac ctgtatcacc ccatccagat tctggccgat 1500
tacctgaccc tccaggagca ctattctagt ctgaaagggc tgacactgag ttggattggg 1560
gacggaaaca atatccctgca ctctattatg atgtcagccg ccaagtttgg aatgcacctc 1620
caggctgcaa ccccaaaagg ctacgaaccc gatgcctcag tgacaaagct ggctgaacag 1680
tacgccaaag agaacggcac taagctgctg ctgaccaacg accctctgga ggcgctcac 1740
ggaggcaacg tgctgatcac cgatacctgg attagtagtg gacaggagga agagaagaag 1800
aagcggctcc aggccttcca gggctaccag gtgacaatga aaaccgctaa ggtcgcagcc 1860
agcgattgga cctttctgca ctgcctgccc agaaagcccg aagaggtgga cgcagaggtc 1920
ttctactctc ccagaagcct ggtgtttccc gaagctgaga ataggaagtg gacaattatg 1980
gcagtgatgg tcagcctgct gactgattat tcacctcagc tccagaaacc aaagtctga 2040
taagcggccg ctattttgga aatttgtgat gctattgctt tatttgaac cattaataagc 2100
tgcaataaac aagttaacaa caacaattgc attcatttta tgmttcaggt tcagggggag 2160
gtgtgggagg ttttttaggc atcgataagg atcttctag agcatggcta cgtagataag 2220
tagcatggcg gtttaatcag taactacaag gaaccctag tgatggagt ggccactccc 2280
tctctgcgcg ctgcctcgct cactgagccc gggcgaccaa aggtgcgccc acgcccgggc 2340
ttgcccgggg cggcctcagt gagcgcagca gcgcgcagcc ttaattaacc taattcactg 2400
ttcgtcgttt tacaacgtcg tgaactggaa aaccctggcg ttaccaact taatcgcctt 2460
gcagcacatc cccctttcgc cagctggcgt aatagcgaag aggcccgcac cgcctgcctt 2520
tcccaacagt tgcgcagcct gaatggcgaa tgggacgcgc cctgtagcgg cgcattaagc 2580
gcgggcgggtg ttgtgggttac gcgcagcgtg accgctacac ttgccagcgc cctagcgcctc 2640
ctcctcttcg ctttcttccc ttcccttctc gccacgttcc cggccttccc cgcctcaagt 2700
gtaaatcggg ggcctccctt agggttccga tttagtgtct tacggcact cgaccccaaa 2760
aaacttgatt agggtgatgg ttcaactgat gggccatcgc cctgatagac ggtttttcgc 2820
cctttgacgt ttgagctcac gttctttaat agtggactct tgmtccaac tggaaacaaca 2880
ctcaacccta tctcggctca ttctttgat ttataagga ttttgcgat ttcggctcat 2940
tgggtaaaaa atgagctgat ttaacaaaaa tttaacgcga attttaacaa aatattaacg 3000
cttacaattt aggtggcact tttcggggaa atgtgcgccc aaccctatt tgmtttttt 3060
tctaataaca tccaattatg tatccgctca tgagacaata accctgataa atgctcaat 3120
aatattgaaa aaggaagatg atgagtatc aacatttccc tgtgcctctt attccctttt 3180
ttgcccattt ttgcttctc gtttttgctc acccagaaac gctggtgaaa gtaaaagatg 3240
ctgaagatca gttgggtgca cgaagtgggtt acatcgaact ggatctcaac agcggtaaga 3300
tccttgagag ttttcgcccc gaagaacggt ttccaatgat gagcaacttt aaagtctcgc 3360
tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cgcgcatac 3420
actattctca gaatgactgt gttgactact caccagtcac agaaaagcat cttacggatg 3480
gcatgacagt aagagaatta tgcagtgctg ccataaccat gagtataaac actgcggcca 3540
acttactctc gacaacgatg ggaggaccga aggagctaac cgcttttttg cacacaatgg 3600
gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg 3660
acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg 3720
gcgaactact tactctagct tcccggcaac aatataaga ctggatggag gcggataaa 3780
ttgcaggacc acttctgcgc tcggcccttc cggctggctg gtttattgct gataaatctg 3840
gagccgggtg gcgtgggtct cgcggtatca ttgcagcact ggggcccagat ggtaagccct 3900
cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaatatagc 3960
agatcgctga gataggtgct tcactgatta agcattggta actgtcagac caagttaact 4020
catatatact ttagattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga 4080
tcctttttga taatctcatg accaaaatcc cttaacgtga gttttcgctc cactgagcgt 4140
cagaccocgt agaaaagatc aaaggatctt cttgagatcc ttttttctg cgcgtaatct 4200
gctgcttgca aacaaaaaaa ccaccgctac cagcgggtgt ttgtttgccc gatcaagagc 4260
taccactctt ttttccgaag gtaactggct tcagcagagc gcagatacca aatactgttc 4320
ttctagtgtg gccgtagtta gccaccactc tcaagaactc tgtagcacct cctacatacc 4380
tcgctctgct aactcctgta ccagtggtg ctgcccagtg cgataaagtc tgtcttaccg 4440
ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggtgta acgggggggt 4500
cgtgcacaca gccagcttg gatcgaacga cctacaccga actgagatc ctagcagctg 4560
agctatgaga aagcgcacg ctctcccgaag ggagaaaggg ggacaggtat ccggtaagcg 4620
gcagggtcgg aacaggagag cgcacgaggg agcttccagg gggaaacgccc tggtatcttt 4680
atagtcctgt cgggtttcgc caccctgac ttgagcgtcg atttttgtga tgctcgtcag 4740
ggggcgggag cctatggaaa aacgccagca acgcccctt tttacggttc ctggcctttt 4800
gctggccttt tgctcacatg ttcttctcctg cgttatcccc tgattctgtg gataaccgta 4860
ttaccgcctt tgagttagct gataccgctc gccgcagccc aacgacagcg cgcagcaggt 4920
cagtgagcga ggaagcggaa gagcgcacca tacgcaaac gcctctcccc gcgcgttggc 4980
cgattcattt atgcagctgg caccagaggt ttcccagctg gaaagcgggc agtgagcga 5040
acgcaattaa tgtgagttag ctcaactatt aggcacccca ggctttacac tttatgcttc 5100
cggctcgtat gttgtgtgga attgtgagcg gataacaatt tcacacagga aacagctatg 5160
accatgatta cgccagattt aattaaggcc ttaat 5195

```

```

SEQ ID NO: 7 moltype = AA length = 354
FEATURE Location/Qualifiers
REGION 1..354
note = Synthetic Construct
source 1..354
mol_type = protein
organism = synthetic construct

```

```

SEQUENCE: 7
MLFNLRILLN NAAFRNGHNF MVRNFRCGQP LQNKVQLKGR DLLTLKNFTG EEIKYMLWLS 60
ADLKFRKIQK GEYLPLQGGK SLGMIFEKRS TRTRLSTETG FALLGGHPCF LTTQDIHLGV 120
NESLTDARV LSSMADAVLA RVYKQSDLDL LAKEASIP II NGLSDLYHPI QILADYVTLQ 180
EHYSLKGLT LSWIGDGNNI LHSIMMSAAK FGMHLQAATP KGYEPDASVT KLABQYAKEN 240

```

-continued

---

GTKLLLTNDP	LEAAHGGNVL	ITDTWISMGQ	EEEKKKRLQA	FQGYQVTMKT	AKVAASDWTG	300
LHCLPRKPEE	VDDEVFYSR	SLVFPEAENR	KWTIMAVMVS	LLTDYSPQLQ	KPKF	354

SEQ ID NO: 8                   moltype = DNA   length = 1065  
 FEATURE                    Location/Qualifiers  
 misc\_feature               1..1065  
                             note = engineered hOTC  
 source                     1..1065  
                             mol\_type = other DNA  
                             organism = synthetic construct

SEQUENCE: 8

atgctgttca	acctgagaat	cctgctgaac	aacgcccct	tcagaaacgg	ccacaacttc	60
atggtgagaa	acttcaagatg	cggccagccc	ctgcagaaca	aggtgcagct	gaagggcaga	120
gacctgctga	ccctgaagaa	cttcaccggc	gaggagatca	agtagatgct	gtggctgagc	180
gcccacctga	agttcagaat	caagcagaag	ggcagatacc	tgcccctgct	gcagggcaag	240
agcctgggca	tgatcttoga	gaagagaagc	accagaacca	gactgagcac	cgagaccggc	300
ctggccctgc	tgggcggcca	ccctgcttc	ctgaccaccc	aggacatcca	cctgggctgt	360
aacgagagcc	tgaccgacac	cgccagagtg	ctgagcagca	tggccgacgc	cgtgctggcc	420
agagtgtaca	agcagagcga	cctggacacc	ctggccaagg	aggccagcat	ccccatcatc	480
aacggcctga	gacacctgta	ccacccatc	cagatcctgg	ccgactacct	gaccctgcag	540
gagcactaca	gcagcctgaa	gggcctgacc	ctgagctgga	tggcgacgg	caacaacatc	600
ctgcacagca	tcagatgag	cgccgccaag	ttcggcatgc	acctgcagcg	cgccaccccc	660
aagggctacg	agcccagcgc	cagcgtgacc	aagctggccg	agcagtacgc	caaggagAAC	720
ggcaccAagc	tgtctgtgac	caacgacccc	ctggaggccg	cccacggcgg	caactgtgctg	780
atcaccgaca	ctggatcag	ctgggcccag	gaggaggaga	agaagaagag	actgcaggcc	840
ttccagggct	accaggtgac	catgaagacc	gccaaggtgg	ccgcccagcga	ctggacccttc	900
ctgcaactgcc	tgcccagaaa	gcccagaggag	gtggacgacg	aggtgttcta	cagccccaga	960
agcctgggtg	tccccgaggg	cgagaacaga	aagtggacca	tcattggccgt	gatggtgagc	1020
ctgctgaccg	actacagccc	ccagctgcag	aagcccaagt	tctga		1065

SEQ ID NO: 9                   moltype = DNA   length = 1065  
 FEATURE                    Location/Qualifiers  
 misc\_feature               1..1065  
                             note = engineered hOTC  
 source                     1..1065  
                             mol\_type = other DNA  
                             organism = synthetic construct

SEQUENCE: 9

atgctgttca	acctgagcat	cctgctgaac	aacgcccct	tccgcaacgg	ccacaacttc	60
atggtgagca	acttccgctg	cggccagccc	ctgcagaaca	aggtgcagct	gaagggcgcg	120
gacctgctga	ccctgaagaa	cttcaccggc	gaggagatca	agtagatgct	gtggctgagc	180
gcccacctga	agttccgat	caagcagaag	ggcagatacc	tgcccctgct	gcagggcaag	240
agcctgggca	tgatcttoga	gaagcgaagc	accgcaaccc	gcctgagcac	cgagaccggc	300
ctggccctgc	tgggcggcca	ccctgcttc	ctgaccaccc	aggacatcca	cctgggctgt	360
aacgagagcc	tgaccgacac	cgcccgctg	ctgagcagca	tggccgacgc	cgtgctggcc	420
cgctgttaca	agcagagcga	cctggacacc	ctggccaagg	aggccagcat	ccccatcatc	480
aacggcctga	gacacctgta	ccacccatc	cagatcctgg	ccgactacct	gaccctgcag	540
gagcactaca	gcagcctgaa	gggcctgacc	ctgagctgga	tggcgacgg	caacaacatc	600
ctgcacagca	tcagatgag	cgccgccaag	ttcggcatgc	acctgcagcg	cgccaccccc	660
aagggctacg	agcccagcgc	cagcgtgacc	aagctggccg	agcagtacgc	caaggagAAC	720
ggcaccAagc	tgtctgtgac	caacgacccc	ctggaggccg	cccacggcgg	caactgtgctg	780
atcaccgaca	ctggatcag	ctgggcccag	gaggaggaga	agaagaagcg	cctgcaggcc	840
ttccagggct	accaggtgac	catgaagacc	gccaaggtgg	ccgcccagcga	ctggacccttc	900
ctgcaactgcc	tgcccagcaa	gcccagaggag	gtggacgacg	aggtgttcta	cagcccccgc	960
agcctgggtg	tccccgaggg	cgagaaccca	aagtggacca	tcattggccgt	gatggtgagc	1020
ctgctgaccg	actacagccc	ccagctgcag	aagcccaagt	tctga		1065

SEQ ID NO: 10                  moltype = RNA   length = 1068  
 FEATURE                    Location/Qualifiers  
 misc\_feature               1..1068  
                             note = engineered hOTC RNA sequence  
 source                     1..1068  
                             mol\_type = other RNA  
                             organism = synthetic construct

SEQUENCE: 10

atgctgttca	acctgcaaat	cctgctgaac	aacgcccgtt	ttcggaaacgg	gcacaacttt	60
atggtgagga	actttcctgt	cggacagccc	ctccagaata	aggtccagct	gaagggcagg	120
gacctgctga	ccctgaaaaa	tttcacaggg	gaggaataca	agtagatgct	gtggctgtca	180
gctgatctga	agttccggat	caagcagaag	ggcgaatata	tgctctgct	ccagggcaaa	240
agcctgggga	tgatcttoga	aaagcagcag	actcggacca	gactgtcaac	cgagactgga	300
ttcctctgct	tgaggagaca	ccctgtttt	ctgaccactc	aggacattca	cctgggagtg	360
aacgagtccc	tgaccgacac	tgctcgcgtc	ctgagctcta	tggccgacgc	tgtgctagct	420
cgagtctaca	aacagtcoga	cctggatacc	ctggccaagg	aagcttctat	ccccattatt	480
aacggcctgt	cagacctgta	tcacccatc	cagattctgg	ccgattacct	gaccctccag	540
gagcactatt	ctagctgaa	agggctgaca	ctgagttgga	tggggacgg	aaacaatata	600
ctgcaactcta	ttatgatgct	agccgccaag	tttggaatgc	acctccagcg	tgcaacccca	660

-continued

```

aaaggctacg aaccgatgc ctcagtgaca aagctggctg aacagtacgc caaagagaac 720
ggcactaagc tgctgctgac caacgacct ctggaggccg ctcacggagg caacgtgctg 780
atcaccgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccaggcc 840
ttccagggtc accagggtac aatgaaaacc gctaaggctc cagccagcga ttggacctt 900
ctgcaactgc tgccccgaaa gccccaagag gtggacgacg aggtcttcta ctctcccaga 960
agcctgggtg ttccccgaag tgagaatagg aagtggacaa ttatggcagt gatggtcagc 1020
ctgctgactg attattcacc tcagctccag aaaccaaagt tctgataa 1068

```

```

SEQ ID NO: 11      moltype = RNA length = 1068
FEATURE
misc_feature      1..1068
                  note = engineered hOTC RNA
source            1..1068
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 11
atgctgttca acctgcgaat cctgctgaac aacgccgctt ttccgaaacgg gcacaacttt 60
atggtgagga actttcgctg cggacagccc ctccagaata aggtccagct gaagggcagg 120
gacctgctga cctgaaaaaa ttccacaggg gaggaatca agtatatgct gtggctgtca 180
gctgatctga agttccggat caagcagaag ggcgaatca tgctctgct ccagggcaaa 240
agcctgggga tgatcttoga aaagcgcagt actcggacca gactgtcaac cgagactgga 300
ttcgctctgc tgggaggaca cccttgtttt ctgaccactc aggacattca cctgggagtg 360
aacgagtccc tgaccgacac tgctcgcgtc ctgagctcta tggccgacgc tgtgctggct 420
cgagtctaca aacagtcoga cctggatacc ctggccaagg aagcttctat cccaattatt 480
aacggcctgt gacacctgta tcaccccatc cagattctgg ccgattacct gaccctccag 540
gagcactatt ctagtctgaa agggctgaca ctgagttgga ttggggacgg aaacaatctc 600
ctgcaactta ttatgatgac agccgccaag ttggaatgc acctccaggg tgcaacccca 660
aaaggctacg aaccgatgc ctcagtgaca aagctggctg aacagtacgc caaagagaac 720
ggcactaagc tgctgctgac caacgacct ctggaggccg ctcacggagg caacgtgctg 780
atcaccgata cctggattag tatgggacag gaggaagaga agaagaagcg gctccaggcc 840
ttccagggtc accagggtac aatgaaaacc gctaaggctc cagccagcga ttggacctt 900
ctgcaactgc tgccccgaaa gccccaagag gtggacgacg aggtcttcta ctctcccaga 960
agcctgggtg ttccccgaag tgagaatagg aagtggacaa ttatggcagt gatggtcagc 1020
ctgctgactg attattcacc tcagctccag aaaccaaagt tctgataa 1068

```

```

SEQ ID NO: 12      moltype = RNA length = 1065
FEATURE
misc_feature      1..1065
                  note = engineered hOTC
source            1..1065
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 12
atgctgttca acctgcgcat cctgctgaac aacgccgctt tccgcaacgg ccacaacttc 60
atggtgagca acttccgctg cggccagccc ctgcagaaca aggtgcagct gaagggccgc 120
gacctgctga cctgaaagaa cttcacccggc gaggagatca agtatatgct gtggctgtca 180
gccgacctga agttccgcat caagcagaag ggcgagtacc tgcccctgct gcagggcaag 240
agcctgggca tgatcttoga gaagcgacgc acccgcaacc gcctgagcac cgagaccggc 300
ctggccctgc tggggcgcca ccctgtcttc ctgaccaccc aggacatcca cctgggagtg 360
aacgagagcc tgaccgacac cgcccgcgtg ctgagcagca tggccgacgc cgtgctggcc 420
cgctgtgaca agcagagcga cctggacacc ctggccaagg aggcacacct ccccatcatc 480
aacggcctga gacacctgta ccaccccatc cagatcctgg ccgactacct gaccctgcag 540
gagcactaca gcagcctgaa gggcctgacc ctgagctgga tcggcgacgg caacaacatc 600
ctgacacagca tcatgatgag cgccgccaag ttccgctatgc acctgcaggg cgccaccccc 660
aaaggctacg agcccagcgc cagcctgacc aagctggccg agcagtacgc caaggagaac 720
ggcaccaagc tgctgctgac caacgacccc ctggaggccg cccacggcgg caacgtgctg 780
atcaccgaca cctggatcag catgggcccag gaggaggaga agaagaagcg cctgcaggcc 840
ttccagggtc accagggtac catgaagacc gccaaaggtgg ccgcccagcga ctggacctt 900
ctgcaactgc tgccccgcaa gccccagggag gtggacgacg aggtgttcta cagccccccg 960
agcctgggtg tccccgaggg cgagaaccgc aagtggacaa tcatggccgt gatggtgagc 1020
ctgctgaccg actacagccc ccagctgcag aagcccaaagt tctga 1065

```

```

SEQ ID NO: 13      moltype = RNA length = 1065
FEATURE
misc_feature      1..1065
                  note = engineered hOTC
source            1..1065
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 13
atgctgttca acctgagaat cctgctgaac aacgccgctt tcagaaacgg ccacaacttc 60
atggtgagaa acttcagatg cggccagccc ctgcagaaca aggtgcagct gaagggcaga 120
gacctgctga cctgaaagaa cttcacccggc gaggagatca agtatatgct gtggctgtca 180
gccgacctga agttcagaat caagcagaag ggcgagtacc tgcccctgct gcagggcaag 240
agcctgggca tgatcttoga gaagagaagc accagaacca gactgagcac cgagaccggc 300
ctggccctgc tggggcgcca ccctgtcttc ctgaccaccc aggacatcca cctgggagtg 360

```

-continued

```

aacgagagcc tgaccgacac cgccagagtg ctgagcagca tggccgacgc cgtgctggcc 420
agagtgtaca agcagagcga cctggacacc ctggccaagg aggccagcat ccccatcatc 480
aacggcctga gcgacctgta ccaccccatc cagatcctgg ccgactacct gacctgcag 540
gagcactaca gcagcctgaa gggcctgacc ctgagctgga tcggcgacgg caacaacatc 600
ctgcacagca tcatgatgag cgccgccaag ttcggcatgc acctgcaggc cgccaccccc 660
aagggtctacg agcccagcgc cagcctgacc aagctggccc agcagtagcc caaggagaac 720
ggcacciaagc tgctgtgtac caacgacccc ctggaggccg cccacggcgg caactgtgtg 780
atcaccgaca cctggatcag catgggcccag gaggaggaga agaagaagag actgcaggcc 840
ttccagggtc accagggtgac catgaagacc gccaaaggtg ccgcccagca ctggaccttc 900
ctgcaactgcc tgcccagaaa gcccgaggag gtggacgacg aggtgttcta cagccccaga 960
agcctgggtg tccccagggc cgagaacaga aagtggacca tcatggcctg gatggtgagc 1020
ctgctgaccg actacagccc ccagctgcag aagcccaagt tctga 1065

```

```

SEQ ID NO: 14      moltype = DNA length = 21
FEATURE           Location/Qualifiers
misc_feature      1..21
                  note = PCR forward primer
source            1..21
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 14
aaactgccaa ttccactgct g 21

```

```

SEQ ID NO: 15      moltype = DNA length = 21
FEATURE           Location/Qualifiers
misc_feature      1..21
                  note = PCR reverse primer
source            1..21
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 15
ccataggcaa aagcaccaag a 21

```

```

SEQ ID NO: 16      moltype = DNA length = 29
FEATURE           Location/Qualifiers
misc_feature      1..29
                  note = Probe
source            1..29
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 16
ttggcccaat agtgagaact ttttctgc 29

```

```

SEQ ID NO: 17      moltype = DNA length = 1090
FEATURE           Location/Qualifiers
misc_feature      1..1090
                  note = Engineered hOTC cDNA termed LW4
source            1..1090
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 17
ctgcagccgc caccatgctg ttcaacctgc gaatcctgct gaacaacgcc gcttttcgga 60
acgggcacaa ctttatgggt aggaactttc gctgcggaca gccctccag aataagggtc 120
agctgaaggg cagggacctg ctgacctga aaaatttcac aggggaggaa atcaagtata 180
tgctgtggct gtcagctgat ctgaagtacc ggatcaagca gaaggcgcaa tatctgctc 240
tgctccaggg caaaagcctg gggatgatct tcgaaaagcg cagtactcgg accagactgt 300
caaccgagac tggattcgct ctgctgggag gacaccctgt tttctgacc actcaggaca 360
ttcacctggg agtgaacgag tccctgaccg acactgctcg cgtcctgagc tctatggccg 420
acgctgtgct ggtcctgagtc tacaaacagt ccgacctgga tacctctggc aaggaagctt 480
ctatcccaat tattaacggc ctgtcagacc tgtatcacc catccagatt ctggccgatt 540
acctgaccct ccaggagcac tattctagtc tgaaaaggct gacactgagt tggattgggg 600
acggaacaaa taccctgcac tctattatga tgtcagccgc caagtttggg atgcacctcc 660
aggctgcaac cccaaaaggc tacgaaccgc atgctcagc gacaaagctg gctgaacagt 720
acgcaaaaga gaacggcact aagctgctgc tgaccaacga ccctctggag gccgctcacg 780
gaggcaacgt gctgatcacc gatacctgga ttagtatggg acaggaggaa gagaagaaga 840
agcggctcca ggccttcag ggctaccagg tgacaatgaa aaccgctaa gtcgagccca 900
gagattggac ctttctgcac tgctctgcca gaaagccgca agaggtggac gacgaggtct 960
tctactctcc cagaagcctg gtgtttcccg aagctgagaa taggaagtgg acaattatgg 1020
cagtgatggt cagcctgctg actgattatt cacctcagct ccagaaacca aagttctgat 1080
aagcggccgc 1090

```

```

SEQ ID NO: 18      moltype = DNA length = 1090
FEATURE           Location/Qualifiers
misc_feature      1..1090
                  note = Engineered hOTC cDNA termed LW3
source            1..1090

```

-continued

---

```

mol_type = other DNA
organism = synthetic construct
SEQUENCE: 18
ctgcagccgc caccatgctg ttcaacctgc gaatcctgct gaacaacgcc gcttttcgga 60
acgggcacaa ctttatgggt aggaactttc gctgeggaca gcccctccag aataaggtcc 120
agctgaaggg cagggacctg ctgacctga aaaatttcac aggggaggaa atcaagtata 180
tgctgtggct gtcagctgat ctgaagtcc ggatcaagca gaaggcgaa tatctgctc 240
tgctccaggg caaaagcctg gggatgatct tcgaaaagcg cagtactcgg accagactgt 300
caaccgagac tggattgct ctgctgggag gacaccttg tttctgacc actcaggaca 360
ttcaactggg agtgaacgag tccctgaccg acactgctcg cgtcctgagc tctatggccg 420
acgctgtgct agctcgagtc tacaacagc cggacctgga taccttgccc aaggaagctt 480
ctatcccaat cttaaaggc ctgtcagacc tgtatcacc catccagatt ctggccgatt 540
acctgacctt ccaggagcac tattctagtc tgaaagggct gacctgagt tggattgggg 600
acggaaacaa tatectgcac tetattatga tgtcagccgc caagtttggga atgcacctcc 660
aggctgcaac cccaaaaggc tacgaaccgc atgcctcagt gacaaagctg gctgaacagt 720
acgcaaaaga gaacggcact aagctgctgc tgaccaacga ccctctggag gccgctcacg 780
gaggcaactg gctgatcacc gatacctgga ttagtatggg acaggaggaa gagaagaaga 840
agcggctcca ggccttcag ggtaccagg tgacaatgaa aaccgctaac gtcgcagcca 900
gagattggac ctttctgcac gcctgccc gaaagcccga agaggtggac gacgaggtct 960
tctactctcc cagaagcctg gtgtttccc aagctgagaa taggaagtgg acaattatgg 1020
cagtgatggg cagcctgctg actgattatt cacctcagct ccagaaacca aagttctgat 1080
aagcgccgc 1090

```

---

What is claimed is:

1. A composition comprising a nucleic acid sequence encoding human ornithine transcarbamylase (hOTCase) and expression control sequences which direct expression of hOTC in a liver cell, wherein the hOTC nucleic acid sequence is less than 80% identical to the wild-type hOTC sequence over the mature sequence or full length hOTC of SEQ ID NO:1, and expresses a functional hOTCase, wherein said hOTC nucleic acid sequence is selected from the nucleic acid sequence comprising SEQ ID NO: 5 or a nucleic acid sequence at least about 96 to about 99% identical thereto or a nucleic acid sequence selected from SEQ ID NO: 9, or a nucleic acid sequence at least about 96 to about 99% identical thereto.

2. The recombinant viral vector according to claim 1, wherein the hOTC nucleic acid sequence has the sequence of SEQ ID NO: 4.

3. The recombinant viral vector according to claim 1, wherein the hOTC nucleic acid sequence has the sequence of SEQ ID NO: 3.

4. The recombinant viral vector according to claim 1, wherein the hOTC nucleic acid sequence has the sequence of SEQ ID NO: 8.

5. The recombinant viral vector according to claim 1, wherein the hOTC is a chimeric OTC comprises a heterologous transit sequence substituted for the native transit sequence of SEQ ID NO: 5 or 9.

6. The recombinant viral vector according to claim 1, wherein the viral vector is selected from an adeno-associated virus (AAV) vector, an adenoviral vector, and a lentiviral vector.

7. The recombinant viral vector according to claim 1, wherein the expression control sequences further comprise a liver-specific promoter.

8. The recombinant viral vector according to claim 7, wherein the liver specific promoter is selected from a thyroxin-binding globulin (TBG) promoter or a lymphocyte-specific protein 1 (LSP1) promoter.

9. The recombinant viral vector according to claim 1, wherein the expression cassette further comprises one or more of an intron, a Kozak sequence, a poly A, and a post-transcriptional regulatory elements.

10. The recombinant viral vector of claim 1, wherein the recombinant viral vector is a recombinant AAV vector which comprises an AA V capsid which has packaged therein a nucleic acid sequence which comprises at least one ITR sequence and the synthetic hOTC.

11. The recombinant viral vector of claim 10, wherein the AAV capsid is selected from AAV8, AAV9 and/or AAVrh10.

12. A recombinant adeno-associated virus (rAAV) having an AAV capsid and packaged therein an expression cassette comprising at least one AAV inverted terminal repeat (ITR) sequence, an engineered nucleic acid sequence encoding human at least the mature ornithine transcarbamylase (hOTCase), and expression control sequences which direct expression of the hOTCase in a liver cell, said expression control sequences comprising a liver-specific promoter, wherein the hOTC nucleic acid sequence is less than 80% identical to the wild-type hOTC sequence over at least the mature hOTC of SEQ ID NO: 1 and comprises at least the mature hOTC of SEQ ID NO: 5 or a nucleic acid sequence at least about 96 to about 99.9% identical thereto or at least the mature hOTC of SEQ ID NO: 9 or a nucleic acid sequence at least about 96 to about 99.9% identical thereto.

13. The rAAV according to claim 12, wherein the AAV capsid is selected from AAV8, AAV9, or AAVrh10.

14. The rAAV according to claim 12, wherein the expression cassette further comprises a 5' AAV inverted terminal repeat (ITR) sequence and a 3' ITR sequence.

15. The rAAV according to claim 12, wherein the at least one AAV ITR comprises a 5' ITR in which the D-sequence and the terminal resolution site is deleted.

16. The rAAV according to claim 12, wherein the 5' and 3' ITRs are from AAV2.

17. The rAAV according to claim 12, wherein the synthetic hOTC has the coding sequence of SEQ ID NO: 3.

18. The rAAV according to claim 12, wherein the synthetic hOTC has the coding sequence of SEQ ID NO: 4.

19. A viral vector comprising a hOTC gene encoding a chimeric ornithine transcarbamylase which comprises at least mature human ornithine transcarbamylase with a heterologous transit sequence, wherein the coding sequence

from the mature human ornithine transcarbamylase is selected from that of a nucleic acid sequence of SEQ ID NO: 3, 4, 5, 8 or 9.

20. A pharmaceutical composition comprising a carrier and an effective amount of the vector according to claim 1.

\* \* \* \* \*