



US 20210207093A1

(19) **United States**

(12) **Patent Application Publication**

Jantz et al.

(10) **Pub. No.: US 2021/0207093 A1**

(43) **Pub. Date: Jul. 8, 2021**

(54) **GENETICALLY-MODIFIED CELLS
COMPRISING A MODIFIED HUMAN T
CELL RECEPTOR ALPHA CONSTANT
REGION GENE**

(71) Applicant: **Precision BioSciences, Inc.**, Durham,
NC (US)

(72) Inventors: **Derek Jantz**, Durham, NC (US); **James
Jefferson Smith**, Morrisville, NC (US);
Michael G. Nicholson, Chapel Hill, NC
(US); **Daniel T. MacLeod**, Durham,
NC (US); **Jeyaraj Antony**, Chapel Hill,
NC (US); **Victor Bartsevich**, Durham,
NC (US)

(73) Assignee: **Precision BioSciences, Inc.**, Durham,
NC (US)

(21) Appl. No.: **17/189,243**

(22) Filed: **Mar. 1, 2021**

Related U.S. Application Data

(63) Continuation of application No. 16/150,179, filed on
Oct. 2, 2018, which is a continuation of application
No. 15/964,446, filed on Apr. 27, 2018, now Pat. No.
10,093,900, which is a continuation of application
No. 15/865,089, filed on Jan. 8, 2018, now Pat. No.
9,969,975, which is a continuation of application No.
PCT/US2016/055492, filed on Oct. 5, 2016.

(60) Provisional application No. 62/297,426, filed on Feb.
19, 2016, provisional application No. 62/237,394,
filed on Oct. 5, 2015.

Publication Classification

(51) **Int. Cl.**
C12N 5/0783 (2006.01)
A61K 39/00 (2006.01)
A61K 35/17 (2006.01)
C12N 15/09 (2006.01)
C12N 15/113 (2006.01)
C12N 15/86 (2006.01)
A61K 48/00 (2006.01)
C07K 14/705 (2006.01)
C07K 14/725 (2006.01)
C07K 16/30 (2006.01)
C07K 16/28 (2006.01)


(52) **U.S. Cl.**
CPC *C12N 5/0636* (2013.01); *C07K 2319/01*
(2013.01); *A61K 35/17* (2013.01); *C12N 15/09*
(2013.01); *C12N 15/113* (2013.01); *C12N*
15/86 (2013.01); *A61K 48/0008* (2013.01);
C07K 14/70503 (2013.01); *C07K 14/7051*
(2013.01); *C07K 16/30* (2013.01); *C07K*
16/3061 (2013.01); *C07K 16/2803* (2013.01);
A61K 2039/5156 (2013.01); *A61K 2039/5158*
(2013.01); *C12N 2510/00* (2013.01); *A61K*
2039/505 (2013.01); *C07K 2317/53* (2013.01);
C07K 2319/02 (2013.01); *C07K 2319/03*
(2013.01); *C07K 2319/33* (2013.01); *C07K*
2319/40 (2013.01); *C07K 2319/74* (2013.01);
A61K 39/0011 (2013.01)

(57) **ABSTRACT**

Disclosed herein is a genetically-modified cell comprising in
its genome a modified human T cell receptor alpha constant
region gene, wherein the cell has reduced cell-surface
expression of the endogenous T cell receptor. The present
disclosure further relates to methods for producing such a
genetically-modified cell, and to methods of using such a
cell for treating a disease in a subject.

Specification includes a Sequence Listing.

A.



B.




FIGURE 1

198
 SEQ ID NO: 33 TRC 1-2x.87 EE KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 34 TRC 1-2x.87 QE KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 35 TRC 1-2x.87 EQ KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 36 TRC 1-2x.87 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 37 TRC 1-2x.6 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 38 TRC 1-2x.20 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 39 TRC 1-2x.55 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 40 TRC 1-2x.60 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 41 TRC 1-2x.105 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 42 TRC 1-2x.163 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 43 TRC 1-2x.113...3 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ

271

7
 SEQ ID NO: 44 TRC 1-2x.5 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 45 TRC 1-2x.8 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 46 TRC 1-2x.25 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 47 TRC 1-2x.72 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 48 TRC 1-2x.80 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 49 TRC 1-2x.84 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 50 TRC 1-2x.120 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 51 TRC 1-2x.113...1 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ
 52 TRC 1-2x.113...2 KEFLLYLAGFVDDGGSINACIAPRQSSKFKHRILKIGFAVGGKQTORRWFLDKLVDEIGVGVYDRGSVSEYVLSQ

80

FIGURE 2A

344

272

TRC 1-2x.87 BE IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.87 CE IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.87 EQ IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.87 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.6 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.20 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.55 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.60 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.105 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.163 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.113...3 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

153

81

TRC 1-2x.5 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.8 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.25 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.72 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.80 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.84 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.120 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.113...1 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

TRC 1-2x.113...2 IKPLHNFLTQLQFFLKLKQKQANLVLKIIEQLPSAKESPDKFFLEVC TWVDQIAAALNDSKTRKTTSETVRAVLD

FIGURE 2B

7
 SEQ ID NO: 58 TRC 1-2x.87 EE K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 59 TRC 1-2x.87 QE K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 60 TRC 1-2x.87 EQ K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 61 TRC 1-2x.87 K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 62 TRC 1-2x.6 K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L V P A V Y Q K T Q R R W F L D K L V D E I G V G Y V D A G S V S E Y R L S E
 63 TRC 1-2x.20 K E F L L Y L A G F V D G D G S I F A S I Y P D Q R K E K H S L R L N F S V F Q K T Q R R W F L C K L V D E I G V G Y V D A G S V S E Y R L S E
 64 TRC 1-2x.55 K E F L L Y L A G F V D G D G S I F A S I Y P D Q R K E K H V L R L F S V F Q K T Q R R W F L D K L V D E I G V G Y V D A G S V S E Y R L S E
 65 TRC 1-2x.60 K E F L L Y L A G F V D G D G S I F A S I Y P D Q R K M K F K H V L L H P S V F Q K T Q R R W F L D K L V D E I G V G Y V D G S V S E Y R L S E
 66 TRC 1-2x.105 K E F L L Y L A G F V D G D G S I F A S I Y P D Q R A K E K H S L K L T F S V Y Q K T Q R R W F L D K L V D E I G V G Y V D G S V S E Y R L S E
 67 TRC 1-2x.163 K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L C K L V D E I G V G Y V D S G S V S E Y R L S E
 68 TRC 1-2x.113_3 K E F L L Y L A G F V D G D G S I F A S I Y P D Q R A K E K H V L R L F S V F Q K T Q R R W F L D K L V D E I G V G Y V D A G S V S E Y R L S E

Hypervariable Region 2 (HVR2)

80

198
 SEQ ID NO: 69 TRC 1-2x.5 K E F L L Y L A G F V D G D G S I F A T I I P D Q R A K E K H L K L F S V F Q K T Q R R W F L D K L V D E I G V G Y V D G S V S E Y R L S E
 70 TRC 1-2x.8 K E F L L Y L A G F V D G D G S I F A T I I P D Q R K E K H Q L A L F N V Q K T Q R R W F L D K L V D E I G V G Y V D G S V S E Y R L S E
 71 TRC 1-2x.25 K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 72 TRC 1-2x.72 K E F L L Y L A G F V D G D G S I F A T I I P C R K F K H L K L T F S V F Q K T Q R R W F L D K L V D E I G V G Y V D A G S V S E Y R L S E
 73 TRC 1-2x.80 K E F L L Y L A G F V D G D G S I F A T I I P D Q R M K F K H Q L R L F V H Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 74 TRC 1-2x.84 K E F L L Y L A G F V D G D G S I F A S I Y P H Q R A K E K H F L K L T F A V Y Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 75 TRC 1-2x.120 K E F L L Y L A G F V D G D G S I F A T I I P D Q R M K F K H Q L R L F V H Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 76 TRC 1-2x.113_1 K E F L L Y L A G F V D G D G S I F A T I I P D Q R M K F K H Q L R L F V H Q K T Q R R W F L D K L V D E I G V G Y V D S G S V S E Y R L S E
 77 TRC 1-2x.113_2 K E F L L Y L A G F V D G D G S I F A T I I P D Q R M K F K H Q L R L F A V H Q K T Q R R W F L D K L V D E I G V G Y V D G S V S E Y R L S E

Hypervariable Region 2 (HVR2)

271

FIGURE 3A

153

81

TRC 1-2x.87 EE IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.87 QE IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.87 EQ IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.87 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.6 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.20 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.55 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.60 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.105 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.163 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.113_3 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD

344

272

TRC 1-2x.5 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.8 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.25 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.72 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.80 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.84 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.120 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.113_1 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD
 TRC 1-2x.113_2 IKPLHNFLTQLQPFLLKLLKQKQANLVLKIIIEQLPSAKESPDKFFLEVCTWVDQIAALNDSKFRKKTTSSETVRAVLD

FIGURE 3B

SEQ ID NO: 7 Hypervariable: Region 1 (HVR1) 80

53 TRC 3-4N.3 KEFLLYLACFVDGDSGKATICPEQKFKHMLLDFVVKOKTORRWFLDKLVDEIGVGYVDDGSSVSCYRLSQ

54 TRC 3-4N.19 KEFLLYLACFVDGDSGKATICPEQKFKHMLLDFVVKOKTORRWFLDKLVDEIGVGYVDDGSSVSCYRLSQ

81 153

TRC 3-4x.3 EKPLHNFLLTQLQPFLLKQKQCANLVKIIIEQLPSAKESPDKFLEVCTWVDQIAALNDSKTRKTTTSETVRAVLD

TRC 3-4x.19 EKPLHNFLLTQLQPFLLKQKQCANLVKIIIEQLPSAKESPDKFLEVCTWVDQIAALNDSKTRKTTTSETVRAVLD

FIGURE 4

SEQ ID		Hypervariable Region 2 (HVR2)	271
NO:	198		
78	TRC 3-4x.3	KEFLLYLAGFVDGDSI NACIQPCQVYKFKHQLNIRFTVHOKTORRWFLEKLVDEIGVGYVYDAGSVSYCLSQ	
79	TRC 3-4x.19	KEFLLYLAGFVDGDSI NACIQPCQVYKFKHQLNIRFTVHOKTORRWFLEKLVDEIGVGYVYDAGSVSYCLSQ	
	272		344
	TRC 3-4x.3	IKPLHNFLTIQDLPFLKQKQANLVLKIIEQLPSAKESPDKFLFVCTWVDQIAALNDSKTRKTTSETVRAVLD	
	TRC 3-4x.19	IKPLHNFLTIQDLPFLKQKQANLVLKIIEQLPSAKESPDKFLFVCTWVDQIAALNDSKTRKTTSETVRAVLD	

FIGURE 5

SEQ ID NO: 7 TRC 7-8x.7
KEFLLYLAGFVDGDSI
ACIIFPQDDMKFKHRLQLLRFVYQNTQRRWFLDKLVDEIGVGVDGSSSYRLSE
Hypervariable Region 1 (HVRL1)
80

SEQ ID NO: 198 TRC 7-8x.9
KEFLLYLAGFVDGDSI
ACIIFPQDDMKFKHRLQLLRFVYQNTQRRWFLDKLVDEIGVGVDGSSSYRLSE
Hypervariable Region 1 (HVRL1)
271

FIGURE 6A

153

81 TRC 7-8x.7 IKPLHNFLTQLQPPFLKQKQANLVKIIIEQLPSAKESPDKFLVCTWVDQIAALNDSKTRKTTSETVRAVLD

344

272 TRC 7-8x.9 IKPLHNFLTQLQPPFLKQKQANLVKIIIEQLPSAKESPDKFLVCTWVDQIAALNDSKTRKTTSETVRAVLD
TRC 7-8x.14 IKPLHNFLTQLQPPFLKQKQANLVKIIIEQLPSAKESPDKFLVCTWVDQIAALNDSKTRKTTSETVRAVLD

FIGURE 6B

SEQ ID NO: 198 TRC 7-8x.7 KEFLLYLAGFVDGDSI 271
Hypervariable Region 2 (HVR2)
NASINPCCAKFKHRLILFFVQKTRRWFLLDKLVDEIGVGYVDGGSSYLSQ

SEQ ID NO: 7 TRC 7-8x.9 KEFLLYLAGFVDGDSI 80
Hypervariable Region 2 (HVR2)
NASINPCCAKFKHRLILFFVQKTRRWFLLDKLVDEIGVGYVDGGSSYLSQ

FIGURE 7A

344

272

TRC 7-8x.7 IKPLHNFLTQLOPFLKLLKQKQANLVLKKIIEQLPSAKESPDKFLEVCTWVDQIAALNDSKTRKTTSETVRAVLD

153

81

TRC 7-8x.9 IKPLHNFLTQLOPFLKLLKQKQANLVLKKIIEQLPSAKESPDKFLEVCTWVDQIAALNDSKTRKTTSETVRAVLD
TRC 7-8x.14 IKPLHNFLTQLOPFLKLLKQKQANLVLKKIIEQLPSAKESPDKFLEVCTWVDQIAALNDSKTRKTTSETVRAVLD

FIGURE 7B





FIGURE 8

A.



B.





FIGURE 9

C.



D.





FIGURE 9 (cont.)

E.



F.




FIGURE 9 (cont.)

G.




FIGURE 9 (cont.)




FIGURE 10






FIGURE 11







FIGURE 12

A.



B.




FIGURE 13

SEQ ID NO:	162	233
86	Wild-Type	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAGCAAAATCTGA
87	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAG---CAAAATCTGAC
88	Indel	ATGGACTTCAAAGAGCAACA-----AACAATCTGAC
89	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAG-----AATCTGAC
90	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAG-----TCTGAC
91	Indel	ATGGACTTCAAAGAGCAA-----ACAATCTGAC
92	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAG---ACAATCTGAC
93	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAGCAA-----TCTGAC
94	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAGCAA-----CAAATCTGAC
95	Indel	ATGGACTTCAAAGAGCAACAGTGTCTGTGGCCCTGGAGCAA-----CAAATCTGAC

FIGURE 14




FIGURE 15




FIGURE 16






FIGURE 17

A.



B.



C.




FIGURE 18





FIGURE 19

A.



B.





FIGURE 20

A.



B.




FIGURE 21

A.

SEQ ID NO:		162	233
92	Wild-Type	ATGGACTTCAAGAGCAACAGTGC	GTGGCCCTGGAGCAACAATCTGGAC
102	Indel	ATGGACTTCAAGAGCAACAGTGC	GTGGCCCTGG--CAACAATCTGGAC
103	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
104	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
105	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
106	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
107	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
108	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
109	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS
110	Indel	ATGGACTTCAAGAGCAACAGTGC	TGTTGGATGACTTTGGCATGFS

B.

SEQ ID NO:		181	214
111	Wild-Type	GTGCTG	GTGCTG
112	Insertion	GTGCTG	GTGCTG

Seq1_Site

FIGURE 22




FIGURE 23




FIGURE 24




FIGURE 25




FIGURE 26






FIGURE 27

A.



B.




FIGURE 28

B.





FIGURE 29
CONTINUED




FIGURE 30




FIGURE 31




FIGURE 32




FIGURE 33

B.








FIGURE 33
CONTINUED

A.




FIGURE 34

B.




FIGURE 34
CONTINUED

C.




FIGURE 34
CONTINUED




FIGURE 35

B.





FIGURE 35
CONTINUED




FIGURE 36




FIGURE 37




FIGURE 37
CONTINUED





FIGURE 37
CONTINUED

A.



B.




FIGURE 38




FIGURE 39






FIGURE 39
CONTINUED


M.



N.



O.



P.




FIGURE 39
CONTINUED





FIGURE 40




FIGURE 41




FIGURE 42

B.








FIGURE 42
CONTINUED




FIGURE 43




FIGURE 44




FIGURE 45

**GENETICALLY-MODIFIED CELLS
COMPRISING A MODIFIED HUMAN T
CELL RECEPTOR ALPHA CONSTANT
REGION GENE**

CROSS REFERENCE TO RELATED
APPLICATIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 15/964,446, filed Apr. 27, 2018, which is a continuation of U.S. patent application Ser. No. 15/865,089, filed Jan. 8, 2018, which claims priority to International Patent Application No. PCT/US2016/055492, filed Oct. 5, 2016, which claims priority to U.S. Provisional Application No. 62/297,426, entitled “Genetically-Modified Cells Comprising a Modified Human T Cell Receptor Alpha Constant Region Gene.” filed Feb. 19, 2016, and U.S. Provisional Application No. 62/237,394, entitled “Genetically-Modified Cells Comprising a Modified Human T Cell Receptor Alpha Constant Region Gene,” filed Oct. 5, 2015, the disclosures of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

[0002] The invention relates to the fields of oncology, cancer immunotherapy, molecular biology and recombinant nucleic acid technology. In particular, the invention relates to a genetically-modified cell comprising in its genome a modified human T cell receptor alpha constant region gene, wherein the cell has reduced cell-surface expression of the endogenous T cell receptor. The invention further relates to methods for producing such a genetically-modified cell, and to methods of using such a cell for treating a disease, including cancer, in a subject.

REFERENCE TO A SEQUENCE LISTING
SUBMITTED AS A TEXT FILE VIA EFS-WEB

[0003] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Apr. 25, 2018, is named P109070014US10-SEQ-MJT.txt, and is 264,046 bytes in size.

BACKGROUND OF THE INVENTION

[0004] T cell adoptive immunotherapy is a promising approach for cancer treatment. This strategy utilizes isolated human T cells that have been genetically-modified to enhance their specificity for a specific tumor associated antigen. Genetic modification may involve the expression of a chimeric antigen receptor or an exogenous T cell receptor to graft antigen specificity onto the T cell. By contrast to exogenous T cell receptors, chimeric antigen receptors derive their specificity from the variable domains of a monoclonal antibody. Thus, T cells expressing chimeric antigen receptors (CAR T cells) induce tumor immunoreactivity in a major histocompatibility complex non-restricted manner. To date, T cell adoptive immunotherapy has been utilized as a clinical therapy for a number of cancers, including B cell malignancies (e.g., acute lymphoblastic leukemia (ALL), B cell non-Hodgkin lymphoma (NHL), and chronic lymphocytic leukemia), multiple myeloma, neuroblastoma, glioblastoma, advanced gliomas, ovarian cancer, mesothelioma, melanoma, and pancreatic cancer.

[0005] Despite its potential usefulness as a cancer treatment, adoptive immunotherapy with CAR T cells has been limited, in part, by expression of the endogenous T cell receptor on the cell surface. CAR T cells expressing an endogenous T cell receptor may recognize major and minor histocompatibility antigens following administration to an allogeneic patient, which can lead to the development of graft-versus-host-disease (GVHD). As a result, clinical trials have largely focused on the use of autologous CAR T cells, wherein a patient's T cells are isolated, genetically-modified to incorporate a chimeric antigen receptor, and then re-infused into the same patient. An autologous approach provides immune tolerance to the administered CAR T cells; however, this approach is constrained by both the time and expense necessary to produce patient-specific CAR T cells after a patient's cancer has been diagnosed.

[0006] Thus, it would be advantageous to develop “off the shelf” CAR T cells, prepared using T cells from a third party donor, that have reduced expression of the endogenous T cell receptor and do not initiate GVHD upon administration. Such products could be generated and validated in advance of diagnosis, and could be made available to patients as soon as necessary. Therefore, a need exists for the development of allogeneic CAR T cells that lack an endogenous T cell receptor in order to prevent the occurrence of GVHD.

[0007] Genetic modification of genomic DNA can be performed using site-specific, rare-cutting endonucleases that are engineered to recognize DNA sequences in the locus of interest. Methods for producing engineered, site-specific endonucleases are known in the art. For example, zinc-finger nucleases (ZFNs) can be engineered to recognize and cut pre-determined sites in a genome. ZFNs are chimeric proteins comprising a zinc finger DNA-binding domain fused to the nuclease domain of the FokI restriction enzyme. The zinc finger domain can be redesigned through rational or experimental means to produce a protein that binds to a pre-determined DNA sequence ~18 basepairs in length. By fusing this engineered protein domain to the FokI nuclease, it is possible to target DNA breaks with genome-level specificity. ZFNs have been used extensively to target gene addition, removal, and substitution in a wide range of eukaryotic organisms (reviewed in Durai et al. (2005), *Nucleic Acids Res* 33, 5978). Likewise, TAL-effector nucleases (TALENs) can be generated to cleave specific sites in genomic DNA. Like a ZFN, a TALEN comprises an engineered, site-specific DNA-binding domain fused to the FokI nuclease domain (reviewed in Mak et al. (2013), *Curr Opin Struct Biol.* 23:93-9). In this case, however, the DNA binding domain comprises a tandem array of TAL-effector domains, each of which specifically recognizes a single DNA basepair. A limitation that ZFNs and TALENs have for the practice of the current invention is that they are heterodimeric, so that the production of a single functional nuclease in a cell requires co-expression of two protein monomers.

[0008] Compact TALENs have an alternative endonuclease architecture that avoids the need for dimerization (Beurdeley et al. (2013), *Nat Commun.* 4:1762). A Compact TALEN comprises an engineered, site-specific TAL-effector DNA-binding domain fused to the nuclease domain from the 1-TevI homing endonuclease. Unlike FokI, 1-TevI does not need to dimerize to produce a double-strand DNA break so a Compact TALEN is functional as a monomer.

[0009] Engineered endonucleases based on the CRISPR/Cas9 system are also known in the art (Ran et al. (2013), *Nat Protoc.* 8:2281-2308; Mali et al. (2013), *Nat Methods* 10:957-63). A CRISPR endonuclease comprises two components: (1) a caspase effector nuclease, typically microbial Cas9; and (2) a short “guide RNA” comprising a ~20 nucleotide targeting sequence that directs the nuclease to a location of interest in the genome. By expressing multiple guide RNAs in the same cell, each having a different targeting sequence, it is possible to target DNA breaks simultaneously to multiple sites in the genome. Thus, CRISPR/Cas9 nucleases are suitable for the present invention. The primary drawback of the CRISPR/Cas9 system is its reported high frequency of off-target DNA breaks, which could limit the utility of the system for treating human patients (Fu et al. (2013), *Nat Biotechnol.* 31:822-6).

[0010] Homing endonucleases are a group of naturally-occurring nucleases that recognize 15-40 base-pair cleavage sites commonly found in the genomes of plants and fungi. They are frequently associated with parasitic DNA elements, such as group 1 self-splicing introns and inteins. They naturally promote homologous recombination or gene insertion at specific locations in the host genome by producing a double-stranded break in the chromosome, which recruits the cellular DNA-repair machinery (Stoddard (2006), *Q. Rev. Biophys.* 38: 49-95). Homing endonucleases are commonly grouped into four families: the LAGLIDADG (SEQ ID NO:7) family, the GIY-YIG family, the His-Cys box family and the HNH family. These families are characterized by structural motifs, which affect catalytic activity and recognition sequence. For instance, members of the LAGLIDADG (SEQ ID NO:7) family are characterized by having either one or two copies of the conserved LAGLIDADG (SEQ ID NO:7) motif (see Chevalier et al. (2001), *Nucleic Acids Res.* 29(18): 3757-3774). The LAGLIDADG (SEQ ID NO:7) homing endonucleases with a single copy of the LAGLIDADG (SEQ ID NO:7) motif form homodimers, whereas members with two copies of the LAGLIDADG (SEQ ID NO:7) motif are found as monomers.

[0011] I-CreI (SEQ ID NO: 6) is a member of the LAGLIDADG (SEQ ID NO:7) family of homing endonucleases that recognizes and cuts a 22 basepair recognition sequence in the chloroplast chromosome of the algae *Chlamydomonas reinhardtii*. Genetic selection techniques have been used to modify the wild-type I-CreI cleavage site preference (Sussman et al. (2004), *J. Mol. Biol.* 342: 31-41; Chames et al. (2005), *Nucleic Acids Res.* 33: e178; Seligman et al. (2002), *Nucleic Acids Res.* 30: 3870-9. Arnould et al. (2006), *J. Mol. Biol.* 355: 443-58). More recently, a method of rationally-designing mono-LAGLIDADG (SEQ ID NO:7) homing endonucleases was described that is capable of comprehensively redesigning I-CreI and other homing endonucleases to target widely-divergent DNA sites, including sites in mammalian, yeast, plant, bacterial, and viral genomes (WO 2007/047859).

[0012] As first described in WO 2009/059195, I-CreI and its engineered derivatives are normally dimeric but can be fused into a single polypeptide using a short peptide linker that joins the C-terminus of a first subunit to the N-terminus of a second subunit (Li et al. (2009), *Nucleic Acids Res.* 37:1650-62; Grizot et al. (2009), *Nucleic Acids Res.* 37:5405-19). Thus, a functional “single-chain” meganuclease can be expressed from a single transcript.

[0013] The use of engineered meganucleases for cleaving DNA targets in the human T cell receptor alpha constant region was previously disclosed in International Publication WO 2014/191527. The '527 publication discloses variants of the I-OnuI meganuclease that are engineered to target a recognition sequence (SEQ ID NO:3 of the '527 publication) within exon 1 of the TCR alpha constant region gene. Although the '527 publication discusses that a chimeric antigen receptor can be expressed in TCR knockout cells, the authors do not disclose the insertion of the chimeric antigen receptor coding sequence into the meganuclease cleavage site in the TCR alpha constant region gene.

[0014] The use of other nucleases and mechanisms for disrupting expression of the endogenous TCR have also been disclosed. For example, the use of zinc finger nucleases for disrupting TCR genes in human T cells was described by U.S. Pat. No. 8,956,828 and by U.S. Patent Application Publication No. US2014/0349402. U.S. Publication No. US2014/0301990 describes the use of zinc finger nucleases and transcription-activator like effector nucleases (TALENs), and a CRISPR/Cas system with an engineered single guide RNA for targeting TCR genes in an isolated T cell. U.S. Patent Application Publication No. US2012/0321667 discloses the use of small-hairpin RNAs that target nucleic acids encoding specific TCRs and/or CD3 chains in T cells.

[0015] However, the present invention improves upon the teachings of the prior art. The present inventors are the first to teach genetically-modified cells that comprise an exogenous polynucleotide sequence (e.g., a chimeric antigen receptor or exogenous TCR coding sequence) inserted into the human TCR alpha constant region gene, which simultaneously disrupts expression of the endogenous T cell receptor at the cell surface. Further, the prior art does not teach the meganucleases or the recognition sequences described herein, or their use for producing such genetically-modified cells.

SUMMARY OF THE INVENTION

[0016] The present invention provides a genetically-modified cell comprising in its genome a modified T cell receptor (TCR) alpha constant region gene. Such a cell is a genetically-modified human T cell, or a genetically-modified cell derived from a human T cell. Further, such a cell has reduced cell-surface expression of the endogenous TCR when compared to an unmodified control cell. The present invention also provides a method for producing the genetically-modified cell. The present invention further provides a method of immunotherapy for treating cancer by administering the genetically-modified cell.

[0017] Thus, in one aspect, the invention provides a genetically-modified cell comprising in its genome a modified human TCR alpha constant region gene, wherein the modified human TCR alpha constant region gene comprises from 5' to 3': (a) a 5' region of the human TCR alpha constant region gene; (b) an exogenous polynucleotide; and (c) a 3' region of the human TCR alpha constant region gene. The genetically-modified cell is a genetically-modified human T cell or a genetically-modified cell derived from a human T cell. Further, the genetically-modified cell has reduced cell-surface expression of the endogenous TCR when compared to an unmodified control cell.

[0018] In one embodiment, the exogenous polynucleotide comprises a nucleic acid sequence encoding a chimeric antigen receptor, wherein the chimeric antigen receptor

comprises an extracellular ligand-binding domain and one or more intracellular signaling domains.

[0019] In one such embodiment, the chimeric antigen receptor comprises an extracellular ligand-binding domain having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:112, wherein the extracellular ligand-binding domain binds to CD19.

[0020] In another such embodiment, the chimeric antigen receptor comprises an intracellular cytoplasmic signaling domain having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:113.

[0021] In another such embodiment, the chimeric antigen receptor comprises an intracellular co-stimulatory signaling domain having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:114.

[0022] In another such embodiment, the chimeric antigen receptor further comprises a signal peptide. In some embodiments, the signal peptide can have at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:115.

[0023] In another such embodiment, the chimeric antigen receptor further comprises a hinge domain. In some embodiments, the hinge domain has at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:116.

[0024] In another such embodiment, the chimeric antigen receptor further comprises a transmembrane domain. In some embodiments, the transmembrane domain has at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:117.

[0025] In another such embodiment, the chimeric antigen receptor has at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:111.

[0026] In another embodiment, the exogenous polynucleotide comprises a promoter sequence that drives expression of the exogenous polynucleotide. In one such embodiment, the promoter sequence has at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:118.

[0027] In another embodiment, the nucleic acid sequence of the exogenous polynucleotide has at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:119.

[0028] In another embodiment, the exogenous polynucleotide is inserted into the TCR gene at a position within a recognition sequence comprising SEQ ID NO:3. In one such embodiment, the modified human TCR alpha constant region gene comprises a nucleic acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:120.

[0029] In another embodiment, the exogenous polynucleotide is inserted into the TCR alpha constant region gene at a position within a recognition sequence comprising SEQ ID NO:4. In one such embodiment, the modified human TCR alpha constant region gene comprises a nucleic acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:121.

[0030] In another embodiment, the exogenous polynucleotide is inserted into the TCR alpha constant region gene at a position within a recognition sequence comprising SEQ ID

NO:5. In one such embodiment, the modified human TCR alpha constant region gene comprises a nucleic acid sequence having at least 80%, at least 85%, at least 90%, at least 95%, or up to 100% sequence identity to SEQ ID NO:122.

[0031] In another aspect, the invention provides a pharmaceutical composition comprising a genetically-modified cell, as described herein, and a pharmaceutically acceptable carrier.

[0032] In another aspect, the invention provides a genetically-modified cell, as described herein, for use as a medicament. The invention further provides the use of a genetically-modified cell, as described herein, in the manufacture of a medicament for treating a disease in a subject in need thereof. In one such aspect, the medicament is useful in the treatment of cancer. In some embodiments, the treatment of cancer is immunotherapy.

[0033] In another aspect, the invention provides a method for producing a genetically-modified cell comprising a modified human TCR alpha constant region gene, the method comprising: (a) introducing into a cell: (i) a first nucleic acid sequence encoding an engineered nuclease; or (ii) an engineered nuclease protein; wherein the engineered nuclease produces a cleavage site at a recognition sequence within the human TCR alpha constant region gene; and (b) introducing into the cell a second nucleic acid sequence comprising an exogenous polynucleotide. In such a method, the cell is a human T cell or is derived from a human T cell. Additionally, the sequence of the exogenous polynucleotide is inserted into the human TCR alpha constant region gene at the cleavage site. Further, the genetically-modified cell has reduced cell-surface expression of the endogenous TCR when compared to an unmodified control cell.

[0034] In various embodiments of the method, the first nucleic acid sequence or the engineered nuclease protein can be introduced into the cell prior to introducing the second nucleic acid, or subsequent to introducing the second nucleic acid.

[0035] In one embodiment of the method, the second nucleic acid sequence comprises from 5' to 3': (a) a 5' homology arm that is homologous to the 5' upstream sequence flanking the cleavage site; (b) the exogenous polynucleotide; and (c) a 3' homology arm that is homologous to the 3' downstream sequence flanking the cleavage site. In such an embodiment, the sequence of the exogenous polynucleotide is inserted into the human TCR alpha constant region gene at the cleavage site by homologous recombination.

[0036] In another embodiment of the method, the second nucleic acid lacks substantial homology to the cleavage site, and the sequence of the exogenous polynucleotide is inserted into the human TCR alpha constant region gene by non-homologous end-joining.

[0037] In another embodiment of the method, the exogenous polynucleotide comprises a nucleic acid sequence encoding a chimeric antigen receptor.

[0038] In another embodiment of the method, the exogenous polynucleotide comprises a first promoter sequence that drives expression of the exogenous polynucleotide.

[0039] In another embodiment of the method, the first nucleic acid encoding the engineered nuclease is introduced into the cell using an mRNA. In some embodiments, the mRNA can be a polycistronic mRNA comprising a coding sequence for at least one engineered nuclease described

herein and a coding sequence for at least one additional protein (e.g., a second nuclease). In particular embodiments, a polycistronic mRNA can encode two or more engineered nucleases described herein that target different recognition sequences within the same gene (e.g., the T cell receptor alpha constant region gene). In other embodiments, a polycistronic mRNA can encode an engineered nuclease described herein and a second nuclease that recognizes and cleaves a different recognition sequence within the same gene (e.g., the T cell receptor alpha constant region gene) or, alternatively, recognizes and cleaves a different recognition sequence within another gene of interest in the genome. In such embodiments, genetically-modified cells produced using such polycistronic mRNA can have multiple genes knocked out simultaneously. In additional embodiments, a polycistronic mRNA can encode at least one engineered nuclease described herein and one additional protein that is beneficial to the cell, improves efficiency of insertion of an exogenous sequence of interest into a cleavage site, and/or is beneficial in the treatment of a disease.

[0040] In another embodiment of the method, at least the second nucleic acid sequence is introduced into the cell by contacting the cell with a viral vector comprising the second nucleic acid sequence. In some embodiments, both the first nucleic acid sequence and the second nucleic acid sequence are introduced by contacting the cell with a single viral vector comprising both the first nucleic acid sequence and the second nucleic acid sequence. Alternatively, the cell can be contacted with a first viral vector comprising the first nucleic acid sequence and a second viral vector comprising the second nucleic acid sequence.

[0041] In such an embodiment of the method, wherein the second nucleic acid sequence is introduced by a viral vector, the second nucleic acid can further comprise a second promoter sequence positioned 5' upstream of the 5' homology arm or, alternatively, positioned 3' downstream of the 3' homology arm. In embodiments where the second promoter is positioned 3' downstream of the 3' homology arm, the promoter may be inverted.

[0042] In another particular embodiment of the method, at least the second nucleic acid sequence is introduced into the cell by contacting the cell with a recombinant adeno-associated virus (AAV) vector comprising the second nucleic acid sequence. In some embodiments, both the first nucleic acid sequence and the second nucleic acid sequence are introduced by contacting the cell with a single recombinant AAV comprising both the first nucleic acid sequence and the second nucleic acid sequence. Alternatively, the cell can be contacted with a first recombinant AAV comprising the first nucleic acid sequence and a second recombinant AAV comprising the second nucleic acid sequence.

[0043] In such an embodiment of the method, wherein the second nucleic acid sequence is introduced by a recombinant AAV vector, the second nucleic acid can further comprise a second promoter sequence positioned 5' upstream of the 5' homology arm or, alternatively, positioned 3' downstream of the 3' homology arm. In embodiments where the second promoter is positioned 3' downstream of the 3' homology arm, the promoter may be inverted.

[0044] In another such embodiment of the method, the recombinant AAV vector is a self-complementary AAV vector.

[0045] In another such embodiment of the method, the recombinant AAV vector can have any serotype. In a par-

ticular embodiment of the method, the recombinant AAV vector has a serotype of AAV2. In another particular embodiment of the method, the recombinant AAV vector has a serotype of AAV6.

[0046] In another embodiment of the method, at least the second nucleic acid sequence is introduced into the cell using a single-stranded DNA template.

[0047] In a particular embodiment of the method, the first nucleic acid sequence encoding an engineered nuclease described herein is introduced into the cell by an mRNA, and the second nucleic acid sequence comprising an exogenous polynucleotide is introduced into the cell using a viral vector, preferably a recombinant AAV vector, wherein the cell is a human T cell, and wherein the sequence of interest encodes a chimeric antigen receptor. In such an embodiment, the method produces a genetically-modified T cell comprising a chimeric antigen receptor and reduced cell-surface expression of the endogenous T cell receptor when compared to a control cell.

[0048] In another embodiment of the method, the engineered nuclease is a recombinant meganuclease, a recombinant zinc-finger nuclease (ZFN), a recombinant transcription activator-like effector nuclease (TALEN), a CRISPR/Cas nuclease, or a megaTAL nuclease. In a particular embodiment of the method, the engineered nuclease is a recombinant meganuclease.

[0049] In such an embodiment of the method, the recombinant meganuclease recognizes and cleaves a recognition sequence within residues 93-208 of the human T cell receptor alpha constant region (SEQ ID NO:1). Such a recombinant meganuclease comprises a first subunit and a second subunit, wherein the first subunit binds to a first recognition half-site of the recognition sequence and comprises a first hypervariable (HVR1) region, and wherein the second subunit binds to a second recognition half-site of the recognition sequence and comprises a second hypervariable (HVR2) region.

[0050] In one such embodiment of the method, the recognition sequence comprises SEQ ID NO:3 (i.e., the TRC 1-2 recognition sequence).

[0051] In another such embodiment of the method, the first meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 198-344 of any one of SEQ ID NOs:8-18 or residues 7-153 of any one of SEQ ID NOs:19-27, and the second meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 7-153 of any one of SEQ ID NOs:8-18 or residues 198-344 of any one of SEQ ID NOs:19-27.

[0052] In another such embodiment of the method, the HVR1 region comprises Y at a position corresponding to: (a) position 215 of any one of SEQ ID NOs:8-18; or (b) position 24 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR1 region comprises G at a position corresponding to: (a) position 233 of any one of SEQ ID NOs:8-18; or (b) position 42 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR1 region comprises one or more of Y and G at positions corresponding to (a) positions 215 and 233, respectively, of any one of SEQ ID NOs:8-18; or (b) positions 24 and 42, respectively, of any one of SEQ ID NOs:19-27.

[0053] In another such embodiment of the method, the HVR2 region comprises T at a position corresponding to: (a)

position 26 of any one of SEQ ID NOs:8-18; or (b) position 217 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises F or Y at a position corresponding to: (a) position 28 of any one of SEQ ID NOs:8-18; or (b) position 219 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises F at a position corresponding to: (a) position 38 of any one of SEQ ID NOs:8-18; or (b) position 229 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises S at a position corresponding to: (a) position 44 of any one of SEQ ID NOs:8-18; or (b) position 235 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises F or Y at a position corresponding to: (a) position 46 of any one of SEQ ID NOs:8-18; or (b) position 237 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises one or more of T, F or Y, F, S, and F or Y, and R at positions corresponding to: (a) positions 26, 28, 38, 44, and 46, respectively, of any one of SEQ ID NOs:8-18; or (b) positions 217, 219, 229, 235, and 237, respectively, of any one of SEQ ID NOs:19-27.

[0054] In another such embodiment of the method, the HVR1 region comprises residues 215-270 of any one of SEQ ID NOs:8-18 or residues 24-79 of any one of SEQ ID NOs:19-27. In another such embodiment, the HVR2 region comprises residues 24-79 of any one of SEQ ID NOs:8-18 or residues 215-270 of any one of SEQ ID NOs:19-27.

[0055] In another such embodiment of the method, the first meganuclease subunit comprises residues 198-344 of any one of SEQ ID NOs:8-18 or residues 7-153 of any one of SEQ ID NOs:19-27. In another such embodiment, the second meganuclease subunit comprises residues 7-153 of any one of SEQ ID NOs:8-18 or residues 198-344 of any one of SEQ ID NOs:19-27.

[0056] In another such embodiment of the method, the recombinant meganuclease is a single-chain meganuclease comprising a linker, wherein the linker covalently joins the first subunit and the second subunit.

[0057] In another such embodiment of the method, the recombinant meganuclease comprises the amino acid sequence of any one of SEQ ID NOs:8-27.

[0058] In a further embodiment of the method, the recognition sequence comprises SEQ ID NO:4 (i.e., the TRC 3-4 recognition sequence).

[0059] In one such embodiment of the method, the first meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 7-153 of SEQ ID NO:28 or 29, and the second meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 198-344 of SEQ ID NO:28 or 29.

[0060] In another such embodiment of the method, the HVR1 region comprises Y at a position corresponding to position 24 of SEQ ID NO:28 or 29. In another such embodiment, the HVR1 region comprises T at a position corresponding to position 26 of SEQ ID NO:30 or 31. In another such embodiment, the HVR1 region comprises Y at a position corresponding to position 46 of SEQ ID NO:28 or 29. In another such embodiment, the HVR1 region comprises one or more of Y, T, and Y at positions corresponding to positions 24, 26, and 46, respectively, of SEQ ID NO:28 or 29.

[0061] In another such embodiment of the method, the HVR2 region comprises H at a position corresponding to position 215 of SEQ ID NO:28 or 29. In another such embodiment, the HVR2 region comprises T at a position corresponding to position 266 of SEQ ID NO:28 or 29. In another such embodiment, the HVR2 region comprises C at a position corresponding to position 268 of SEQ ID NO:28 or 29. In another such embodiment, the HVR2 region comprises one or more of H, T, and C at positions corresponding to positions 215, 266, and 268 of SEQ ID NOs:28 or 29.

[0062] In another such embodiment of the method, the HVR1 region comprises residues 24-79 of SEQ ID NO:28 or 29. In another such embodiment, the HVR2 region comprises residues 215-270 of SEQ ID NO:28 or 29.

[0063] In another such embodiment of the method, the first meganuclease subunit comprises residues 7-153 of SEQ ID NO:28 or 29. In another such embodiment, the second meganuclease subunit comprises residues 198-344 of SEQ ID NO:28 or 29.

[0064] In another such embodiment of the method, the recombinant meganuclease is a single-chain meganuclease comprising a linker, wherein the linker covalently joins the first subunit and the second subunit.

[0065] In another such embodiment of the method, the recombinant meganuclease comprises the amino acid sequence of SEQ ID NO:28 or 29.

[0066] In a further embodiment of the method, the recognition sequence comprises SEQ ID NO:5 (i.e., the TRC 7-8 recognition sequence).

[0067] In one such embodiment of the method, the first meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 7-153 of SEQ ID NO:30 or residues 198-344 of SEQ ID NO:31 or 32, and the second meganuclease subunit comprises an amino acid sequence having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to residues 198-344 of SEQ ID NO:30 or residues 7-153 of SEQ ID NO:31 or 32.

[0068] In another such embodiment of the method, the HVR1 region comprises Y at a position corresponding to: (a) position 24 of SEQ ID NO:30; or (b) position 215 of SEQ ID NO:31 or 32.

[0069] In another such embodiment of the method, the HVR2 region comprises Y or W at a position corresponding to: (a) position 215 of SEQ ID NO:30; or (b) position 24 of SEQ ID NO:31 or 32. In another such embodiment, the HVR2 region comprises M, L, or W at a position corresponding to: (a) position 231 of SEQ ID NO:30; or (b) position 40 of SEQ ID NO:31 or 32. In another such embodiment, the HVR2 region comprises Y at a position corresponding to: (a) position 237 of SEQ ID NO:30; or (b) position 46 of SEQ ID NO:31 or 32. In another such embodiment, the HVR2 region comprises one or more of Y or W, M, L, or W, and Y at positions corresponding to: (a) positions 215, 231, and 237, respectively, of SEQ ID NO:30; or (b) positions 24, 40, and 46, respectively, of SEQ ID NO:31 or 32.

[0070] In another such embodiment of the method, the HVR1 region comprises residues 24-79 of SEQ ID NO:30 or residues 215-270 of SEQ ID NO:31 or 32. In another such embodiment, the HVR2 region comprises residues 215-270 of SEQ ID NO:30 or residues 24-79 of SEQ ID NO:31 or 32.

[0071] In another such embodiment of the method, the first meganuclease subunit comprises residues 7-153 of SEQ ID NO:30 or residues 198-344 of SEQ ID NO:31 or 32. In another such embodiment, the second meganuclease subunit comprises residues 198-344 of SEQ ID NO:30 or residues 7-153 of SEQ ID NO:31 or 32.

[0072] In another such embodiment of the method, the recombinant meganuclease is a single-chain meganuclease comprising a linker, wherein the linker covalently joins the first subunit and the second subunit.

[0073] In another such embodiment of the method, the recombinant meganuclease comprises the amino acid sequence of any one of SEQ ID NOs:30-32.

[0074] In another aspect, the invention provides a method of immunotherapy for treating cancer in a subject in need thereof. In some embodiments, the method comprises administering to the subject a pharmaceutical composition comprising a genetically-modified cell, as described herein, and a pharmaceutically acceptable carrier. In some embodiments, the method comprises administering to the subject a pharmaceutical composition comprising a genetically-modified cell produced according to the methods described herein, and a pharmaceutically acceptable carrier.

[0075] In another embodiment of the method, the cancer to be treated is selected from the group consisting of a cancer of B-cell origin, breast cancer, gastric cancer, neuroblastoma, osteosarcoma, lung cancer, melanoma, prostate cancer, colon cancer, renal cell carcinoma, ovarian cancer, rhabdomyo sarcoma, leukemia, and Hodgkin's lymphoma.

[0076] In another embodiment of the method, the cancer of B-cell origin is selected from the group consisting of B-lineage acute lymphoblastic leukemia, B-cell chronic lymphocytic leukemia, and B-cell non-Hodgkin's lymphoma.

[0077] In some embodiments, the CAR comprises an extracellular antigen-binding domain. In some embodiments, the extracellular ligand-binding domain or moiety can be in the form of a single-chain variable fragment (scFv) derived from a monoclonal antibody, which provides specificity for a particular epitope or antigen (e.g., an epitope or antigen preferentially present on the surface of a cell, such as a cancer cell or other disease-causing cell or particle). The scFv can be attached via a linker sequence. The extracellular ligand-binding domain can be specific for any antigen or epitope of interest. In some embodiments, the scFv can be humanized. The extracellular domain of a chimeric antigen receptor can also comprise an autoantigen (see. Payne et al. (2016), *Science* 353(6295): 179-184), which can be recognized by autoantigen-specific B cell receptors on B lymphocytes, thus directing T cells to specifically target and kill autoreactive B lymphocytes in antibody-mediated autoimmune diseases. Such CARs can be referred to as chimeric autoantibody receptors (CAARs), and their use is encompassed by the invention.

[0078] The foregoing and other aspects and embodiments of the present invention can be more fully understood by reference to the following detailed description and claims. Certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. All combinations of the embodiments are specifically embraced by the present invention and are disclosed herein just as if each and every combination was individually and explicitly disclosed. Conversely, various features of the invention,

which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination. All sub-combinations of features listed in the embodiments are also specifically embraced by the present invention and are disclosed herein just as if each and every such sub-combination was individually and explicitly disclosed herein. Embodiments of each aspect of the present invention disclosed herein apply to each other aspect of the invention *mutatis mutandis*.

BRIEF DESCRIPTION OF THE FIGURES

[0079] FIG. 1. TRC recognition sequences in the human TRC alpha constant region gene. A) Each recognition sequence targeted by a recombinant meganuclease of the invention comprises two recognition half-sites. Each recognition half-site comprises 9 base pairs, separated by a 4 base pair central sequence. The TRC 1-2 recognition sequence (SEQ ID NO:3) spans nucleotides 187-208 of the human T cell alpha constant region (SEQ ID NO:1), and comprises two recognition half-sites referred to as TRC1 and TRC2. The TRC 3-4 recognition sequence (SEQ ID NO:4) spans nucleotides 93-114 of the human T cell alpha constant region (SEQ ID NO:1), and comprises two recognition half-sites referred to as TRC3 and TRC4. The TRC 7-8 recognition sequence (SEQ ID NO:5) spans nucleotides 118-139 of the human T cell alpha constant region (SEQ ID NO:1), and comprises two recognition half-sites referred to as TRC7 and TRC8. B) The recombinant meganucleases of the invention comprise two subunits, wherein the first subunit comprising the HVR1 region binds to a first recognition half-site (e.g., TRC1, TRC3, or TRC7) and the second subunit comprising the HVR2 region binds to a second recognition half-site (e.g., TRC2, TRC4, or TRC8). In embodiments where the recombinant meganuclease is a single-chain meganuclease, the first subunit comprising the HVR1 region can be positioned as either the N-terminal or C-terminal subunit. Likewise, the second subunit comprising the HVR2 region can be positioned as either the N-terminal or C-terminal subunit.

[0080] FIG. 2A-B. Amino acid alignment of TRC1-binding subunits. A-B) Some recombinant meganucleases encompassed by the invention comprise one subunit that binds the 9 base pair TRC1 recognition half-site of SEQ ID NO:3. Amino acid sequence alignments are provided for the TRC1-binding subunits (SEQ ID NOs:33-52) of the recombinant meganucleases set forth in SEQ ID NOs:8-27. As shown, the TRC1-binding subunit of SEQ ID NOs:8-18 comprises residues 198-344, whereas the TRC1-binding subunit of SEQ ID NOs:19-27 comprises residues 7-153. Each TRC1-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded, with the most frequent amino acids at each position further highlighted; the most prevalent residues are bolded, whereas the second most prevalent are bolded and italicized. Residues outside of the hypervariable region are identical in each subunit, with the exception of a Q or E residue at position 80 or position 271 (see. U.S. Pat. No. 8,021,867). All TRC1-binding subunits provided in FIG. 2 share at least 90% sequence identity to the TRC1-binding subunit (residues 198-344) of the TRC 1-2x.87 EE meganuclease (SEQ ID NO:33). Residue numbers shown are those of SEQ ID NOs:8-27.

[0081] FIG. 3A-B. Amino acid alignment of TRC2-binding subunits. A-B) Some recombinant meganucleases

encompassed by the invention comprise one subunit that binds the 9 base pair TRC2 recognition half-site of SEQ ID NO:3. Amino acid sequence alignments are provided for the TRC2-binding subunits (SEQ ID NOs:58-77) of the recombinant meganucleases set forth in SEQ ID NOs:8-27. As shown, the TRC2-binding subunit of SEQ ID NOs:8-18 comprises residues 7-153, whereas the TRC2-binding subunit of SEQ ID NOs:19-27 comprises residues 198-344. Each TRC2-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded, with the most frequent amino acids at each position further highlighted; the most prevalent residues are bolded, whereas the second most prevalent are bolded and italicized. Residues outside of the hypervariable region are identical in each subunit, with the exceptions of a Q or E residue at position 80 or position 271 (see, U.S. Pat. No. 8,021,867), and an R residue at position 139 of meganucleases TRC 1-2x.87 EE, TRC 1-2x.87 QE, TRC 1-2x.87 EQ, TRC 1-2x.87, and TRC 1-2x.163 (shaded grey and underlined). All TRC2-binding subunits provided in FIG. 3 share at least 90% sequence identity to the TRC2-binding subunit (residues 7-153) of the TRC 1-2x.87 EE meganuclease (SEQ ID NO:58). Residue numbers shown are those of SEQ ID NOs:8-27.

[0082] FIG. 4. Amino acid alignment of TRC3-binding subunits. Some recombinant meganucleases encompassed by the invention comprise one subunit that binds the 9 base pair TRC3 recognition half-site of SEQ ID NO:4. Amino acid sequence alignments are provided for the TRC3-binding subunits (SEQ ID NOs:53 and 54) of the recombinant meganucleases set forth in SEQ ID NOs:28 and 29. As shown, the TRC3-binding subunit of SEQ ID NOs:28 and 29 comprises residues 7-153. Each TRC3-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded. Residues outside of the hypervariable region are identical in each subunit, with the exceptions of a Q or E residue at position 80 (see, U.S. Pat. No. 8,021,867). The TRC3-binding subunits of the TRC 3-4x.3 and TRC 3-4x.19 meganucleases share 97% sequence identity. Residue numbers shown are those of SEQ ID NOs:28 and 29.

[0083] FIG. 5. Amino acid alignment of TRC4-binding subunits. Some recombinant meganucleases encompassed by the invention comprise one subunit that binds the 9 base pair TRC4 recognition half-site of SEQ ID NO:4. Amino acid sequence alignments are provided for the TRC4-binding subunits (SEQ ID NOs:78 and 79) of the recombinant meganucleases set forth in SEQ ID NOs:28 and 29. As shown, the TRC4-binding subunit of SEQ ID NOs:28 and 29 comprises residues 198-344. Each TRC4-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded. Residues outside of the hypervariable region are identical in each subunit, with the exceptions of a Q or E residue at position 80 (see, U.S. Pat. No. 8,021,867). The TRC4-binding subunits of the TRC 3-4x.3 and TRC 3-4x.19 meganucleases share 97% sequence identity. Residue numbers shown are those of SEQ ID NOs:28 and 29.

[0084] FIG. 6A-B. Amino acid alignment of TRC7-binding subunits. A-B) Some recombinant meganucleases encompassed by the invention comprise one subunit that binds the 9 base pair TRC7 recognition half-site of SEQ ID NO:5. Amino acid sequence alignments are provided for the TRC7-binding subunits (SEQ ID NOs:55-57) of the recom-

binant meganucleases set forth in SEQ ID NOs:30-32. As shown, the TRC7-binding subunit of SEQ ID NO:30 comprises residues 7-153, whereas the TRC7-binding subunit of SEQ ID NOs:31 and 32 comprises residues 198-344. Each TRC7-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded, with the most frequent amino acids at each position further highlighted; the most prevalent residues are bolded, whereas the second most prevalent are bolded and italicized. Residues outside of the hypervariable region are identical in each subunit, with the exception of a Q or E residue at position 80 or position 271 (see, U.S. Pat. No. 8,021,867). All TRC7-binding subunits provided in FIG. 6 share at least 90% sequence identity to the TRC7-binding subunit (residues 7-153) of the TRC 7-8x.7 meganuclease (SEQ ID NO:55). Residue numbers shown are those of SEQ ID NOs:30-32.

[0085] FIG. 7A-B. Amino acid alignment of TRC8-binding subunits. A-B) Some recombinant meganucleases encompassed by the invention comprise one subunit that binds the 9 base pair TRC8 recognition half-site of SEQ ID NO:5. Amino acid sequence alignments are provided for the TRC8-binding subunits (SEQ ID NOs:80-82) of the recombinant meganucleases set forth in SEQ ID NOs:30-32. As shown, the TRC8-binding subunit of SEQ ID NO:30 comprises residues 198-344, whereas the TRC8-binding subunit of SEQ ID NOs:31 and 32 comprises residues 7-153. Each TRC8-binding subunit comprises a 56 amino acid hypervariable region as indicated. Variable residues within the hypervariable region are shaded, with the most frequent amino acids at each position further highlighted; the most prevalent residues are bolded, whereas the second most prevalent are bolded and italicized. Residues outside of the hypervariable region are identical in each subunit, with the exception of a Q or E residue at position 80 or position 271 (see, U.S. Pat. No. 8,021,867). All TRC8-binding subunits provided in FIG. 7 share at least 90% sequence identity to the TRC8-binding subunit (residues 198-344) of the TRC 7-8x.7 meganuclease (SEQ ID NO:80). Residue numbers shown are those of SEQ ID NOs:30-32.

[0086] FIG. 8. Schematic of reporter assay in CHO cells for evaluating recombinant meganucleases targeting recognition sequences found in the T cell receptor alpha constant region (SEQ ID NO:1). For the recombinant meganucleases described herein, a CHO cell line was produced in which a reporter cassette was integrated stably into the genome of the cell. The reporter cassette comprised, in 5' to 3' order: an SV40 Early Promoter, the 5' $\frac{2}{3}$ of the GFP gene; the recognition sequence for an engineered meganuclease of the invention (e.g., the TRC 1-2 recognition sequence, the TRC 3-4 recognition sequence, or the TRC 7-8 recognition sequence); the recognition sequence for the CHO-23/24 meganuclease (WO/2012/167192); and the 3' $\frac{1}{3}$ of the GFP gene. Cells stably transfected with this cassette did not express GFP in the absence of a DNA break-inducing agent. Meganucleases were introduced by transduction of plasmid DNA or mRNA encoding each meganuclease. When a DNA break was induced at either of the meganuclease recognition sequences, the duplicated regions of the GFP gene recombined with one another to produce a functional GFP gene. The percentage of GFP-expressing cells could then be determined by flow cytometry as an indirect measure of the frequency of genome cleavage by the meganucleases.

[0087] FIG. 9. Efficiency of recombinant meganucleases for recognizing and cleaving recognition sequences in the human T cell receptor alpha constant region (SEQ ID NO:1) in a CHO cell reporter assay. Each of the recombinant meganucleases set forth in SEQ ID NOs:8-32 were engineered to target the TRC 1-2 recognition sequence (SEQ ID NO:3), the TRC 3-4 recognition sequence (SEQ ID NO:4), or the TRC 7-8 recognition sequence (SEQ ID NO:5), and were screened for efficacy in the CHO cell reporter assay. The results shown provide the percentage of GFP-expressing cells observed in each assay, which indicates the efficacy of each meganuclease for cleaving a TRC target recognition sequence or the CHO-23/24 recognition sequence. A negative control (RHO 1-2 bs) was further included in each assay. A)-C) Meganucleases targeting the TRC 1-2 recognition sequence. D) Meganucleases targeting the TRC 3-4 recognition sequence. E)-F) Meganucleases targeting the TRC 7-8 recognition sequence. G) Variants of the TRC 1-2x.87 meganuclease, wherein the Q at position 271 is substituted with E (TRC 1-2x.87 QE), the Q at position 80 is substituted with E (TRC 1-2x.87 EQ), or the Q at position 80 and the Q at position 271 are both substituted with E (TRC 1-2x.87 EE).

[0088] FIG. 10. Time course of recombinant meganuclease efficacy in CHO cell reporter assay. The TRC 1-2x.87 QE, TRC 1-2x.87 EQ, and TRC 1-2x.87 EE meganucleases were evaluated in the CHO reporter assay, with the percentage of GFP-expressing cells determined 1, 4, 6, 8, and 12 days after introduction of meganuclease-encoding mRNA into the CHO reporter cells.

[0089] FIG. 11. Analysis of Jurkat cell genomic DNA following transfection with TRC 1-2 meganucleases. At 72 hours post-transfection with mRNA encoding TRC 1-2 meganucleases, genomic DNA was harvested and a T7 endonuclease assay was performed to estimate genetic modification at the endogenous TRC 1-2 recognition sequence.

[0090] FIG. 12. Dose-response of TRC 1-2 meganuclease expression in Jurkat cells on genetic modification at the endogenous TRC 1-2 recognition sequence. Jurkat cells were transfected with either 3 μ g or 1 μ g of a given TRC 1-2 meganuclease mRNA. At 96 hours, genomic DNA was analyzed using a T7 endonuclease assay.

[0091] FIG. 13. Cleavage of TRC 1-2 recognition sequence in human T cells. A) CD3+ T cells were stimulated with anti-CD3 and anti-CD28 antibodies for 3 days, then electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease. Genomic DNA was harvested at 3 days and 7 days post-transfection, and analyzed using a T7 endonuclease assay. B) To determine whether mutations at the endogenous TRC 1-2 recognition sequence were sufficient to eliminate surface expression of the T cell receptor, cells were analyzed by flow cytometry using an anti-CD3 antibody. Control cells (transfected with water) and TRC 1-2x.87 EE-transfected cells were analyzed at day 3 and day 7 post-transfection, and the percentage of CD3-positive and CD3-negative T cells was determined.

[0092] FIG. 14. Nucleic acid sequences of representative deletions that were observed at the TRC 1-2 recognition sequence in human T cells following expression of TRC 1-2 meganucleases.

[0093] FIG. 15. Diagram illustrating sequence elements of recombinant AAV vectors and their use in combination with

an engineered nuclease to insert an exogenous nucleic acid sequence into the endogenous TRC alpha constant region gene.

[0094] FIG. 16. Map of plasmid used to produce the AAV405 vector.

[0095] FIG. 17. Map of plasmid used to produce the AAV406 vector.

[0096] FIG. 18. Determining the timing of meganuclease mRNA transfection and recombinant AAV transduction to enhance AAV transduction efficiency. Human CD3+ T cells were electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease and at 2, 4, or 8 hours post-transfection, cells were transduced with a recombinant AAV vector encoding GFP (GFP-AAV). T cells were analyzed by flow cytometry for GFP expression at 72 hours post-transduction to determine transduction efficiency.

[0097] FIG. 19. Analyzing human T cells for insertion of an exogenous nucleic acid sequence using recombinant AAV vectors. CD3+ T cells transfected with TRC 1-2x.87 EE mRNA and subsequently transduced (2 hours post-transfection) with AAV405 or AAV406. Transduction-only controls were mock transfected (with water) and transduced with either AAV405 or AAV406. Meganuclease-only controls were transfected with TRC 1-2x.87 EE and then mock transduced (with water) at 2 hours post-transfection. Genomic DNA was harvested from T cells and the TRC 1-2 locus was amplified by PCR using primers that recognized sequences beyond the region of homology in the AAV vectors. PCR primers outside of the homology regions only allowed for amplification of the T cell genome, not from the AAV vectors. PCR products were purified and digested with *EagI*. PCR products were then analyzed for cleavage.

[0098] FIG. 20. Characterization of *EagI* insertion into the TRC 1-2 recognition sequence of human T cells using AAV405. A) Undigested PCR product generated from previous experiments was cloned into a pCR-blunt vector. Colony PCR was performed using M13 forward and reverse primers and a portion of PCR products from cells transfected with TRC 1-2x.87 EE and AAV405 was analyzed by gel electrophoresis. Analysis shows a mix of full-length PCR products (approximately 1600 bp), smaller inserts, and empty plasmids (approximately 300 bp). B) In parallel, another portion of PCR products were digested with *EagI* to determine the percent of clones that contain the *EagI* recognition site inserted in the TRC 1-2 recognition sequence. PCR products cleaved with *EagI* generated expected fragments of approximately 700 and 800 bp.

[0099] FIG. 21. Characterization of *EagI* insertion into the TRC 1-2 recognition sequence of human T cells using AAV406. A) Undigested PCR product generated from previous experiments was cloned into a pCR-blunt vector. Colony PCR was performed using M13 forward and reverse primers and a portion of PCR products from cells transfected with TRC 1-2x.87 EE and AAV406 was analyzed by gel electrophoresis. Analysis shows a mix of full-length PCR products (approximately 1600 bp), smaller inserts, and empty plasmids (approximately 300 bp). B) In parallel, another portion of PCR products were digested with *EagI* to determine the percent of clones that contain the *EagI* recognition site inserted in the TRC 1-2 recognition sequence. PCR products cleaved with *EagI* generated expected fragments of approximately 700 and 800 bp.

[0100] FIG. 22. A) Nucleic acid sequences of representative deletions and insertions (i.e., indels) that were observed

at the TRC 1-2 recognition sequence in human T cells following expression of TRC 1-2 meganucleases. B) Nucleic acid sequence of the TRC 1-2 recognition sequence confirming insertion of the exogenous nucleic acid sequence comprising the EagI restriction site.

[0101] FIG. 23. Enhancement of recombinant AAV transduction efficiency. Transduction efficiency was further analyzed by optimizing the timing of meganuclease mRNA transfection and subsequent AAV transduction. Human CD3+ T cells were electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease and subsequently transduced with GFP-AAV immediately after transfection or 2 hours post-transfection. Additionally, non-stimulated resting T cells were transduced with GFP-AAV. Mock transduced cells were also analyzed. At 72 hours post-transduction, cells were analyzed by flow cytometry for GFP expression to determine AAV transduction efficiency.

[0102] FIG. 24. Map of plasmid used to produce the AAV-CAR100 (AAV408) vector.

[0103] FIG. 25. Map of plasmid used to produce the AAV-CAR763 (AAV412) vector.

[0104] FIG. 26. Insertion of chimeric antigen receptor coding sequence at TRC 1-2 recognition site in human T cells. A PCR-based assay was developed to determine whether the AAV412 HDR template was utilized to repair double-strand breaks at the TRC 1-2 recognition sequence.

[0105] FIG. 27. Insertion of chimeric antigen receptor coding sequence at TRC 1-2 recognition site in human T cells. A PCR-based assay was developed to determine whether the AAV408 HDR template was utilized to repair double-strand breaks at the TRC 1-2 recognition sequence. A) PCR products generated using a primer pair that only amplifies a product on the 5' end of the TRC 1-2 recognition sequence locus if the CAR gene has been inserted into that locus. B) PCR products generated using a primer pair that only amplifies a product on the 3' end of the TRC 1-2 recognition sequence locus if the CAR gene has been inserted into that locus.

[0106] FIG. 28. Digital PCR. A) Schematic of a digital PCR assay developed to quantitatively determine insertion efficiency of the chimeric antigen receptor coding sequence into the TRC 1-2 recognition site in human T cells. B) Results of digital PCR on genomic DNA from human T cells electroporated with a TRC 1-2x.87EE meganuclease mRNA and/or increasing amounts of AAV408.

[0107] FIG. 29. Cell-surface expression of CD19 chimeric antigen receptor on human T cells. The expression level of the anti-CD19 chimeric antigen receptor was determined in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV408 as the HDR template. Cell-surface expression was analyzed by flow cytometry. A) Cells that were mock electroporated and mock transduced (MOI-0), and cells that were mock electroporated and transduced with increasing amounts of AAV408. B) Cells that were electroporated with TRC 1-2x.87EE and mock transduced (MOI-0), and cells that were electroporated with TRC 1-2x.87EE and transduced with increasing amounts of AAV408.

[0108] FIG. 30. Map of plasmid used to produce the AAV421 vector.

[0109] FIG. 31. Map of plasmid used to produce the AAV422 vector.

[0110] FIG. 32. Insertion of chimeric antigen receptor coding sequence. PCR methods were used to determine if

the chimeric antigen receptor coding sequence introduced by AAV421 or AAV422 inserted at the TRC 1-2 recognition site cleaved by the TRC 1-2x.87EE meganuclease. A) Analysis of insertion following transduction with AAV421. B) Analysis of insertion following transduction with AAV422.

[0111] FIG. 33. Cell-surface expression of CD19 chimeric antigen receptor on human T cells. The expression level of the anti-CD19 chimeric antigen receptor was determined in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV421 as the HDR template. Cell-surface expression was analyzed by flow cytometry. A) Cells that were mock electroporated and mock transduced (MOI-0), and cells that were mock electroporated and transduced with increasing amounts of AAV421. B) Cells that were electroporated with TRC 1-2x.87EE and mock transduced (MOI-0), and cells that were electroporated with TRC 1-2x.87EE and transduced with increasing amounts of AAV421.

[0112] FIG. 34. Expansion of human T cells expressing a cell-surface chimeric antigen receptor. Methods were determined for preferentially expanding and enriching a CD3⁺/CAR⁺ T cell population following electroporation with mRNA for the TRC 1-2x.87EE meganuclease and transduction with AAV421. A) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL). B) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL), and incubation with mitomycin C-inactivated IM-9 cells. C) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL), and two incubations with mitomycin C-inactivated IM-9 cells.

[0113] FIG. 35. Cell-surface expression of CD19 chimeric antigen receptor on human T cells. The expression level of the anti-CD19 chimeric antigen receptor was determined in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV422 as the HDR template. Cell-surface expression was analyzed by flow cytometry. A) Cells that were mock electroporated and mock transduced (MOI-0), and cells that were mock electroporated and transduced with increasing amounts of AAV422. B) Cells that were electroporated with TRC 1-2x.87EE and mock transduced (MOI-0), and cells that were electroporated with TRC 1-2x.87EE and transduced with increasing amounts of AAV422.

[0114] FIG. 36. Expansion of human T cells expressing a cell-surface chimeric antigen receptor. Methods were determined for preferentially expanding and enriching a CD3⁺/CAR⁺ T cell population following electroporation with mRNA for the TRC 1-2x.87EE meganuclease and transduction with AAV422. A) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL). B) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL), and incubation with mitomycin C-inactivated IM-9 cells. C) Supplementation with IL-7 (10 ng/mL) and IL-15 (10 ng/mL), and two incubations with mitomycin C-inactivated IM-9 cells.

[0115] FIG. 37. Meganuclease knockout efficiency using single-strand AAV. Experiments were conducted to examine the knockout efficiency of two meganucleases in human T cells when simultaneously transduced with a single-stranded AAV vector. A) Cells electroporated with mRNA for TRC 1-2x.87EE and transduced with increasing amounts of the single-stranded AAV412. B) Cells electroporated with mRNA for a meganuclease targeting the beta-2 microglobulin gene and transduced with increasing amounts of the single-stranded AAV412. C) Cells electroporated with

mRNA for TRC 1-2x.87EE and transduced with increasing amounts of the single-stranded AAV422.

[0116] FIG. 38. Functional activity of anti-CD19 CAR T cells. A) IFN-gamma ELISPOT assay, in which either CD19⁺ Raji cells or CD19⁻ U937 cells were the target population. B) Cell killing assay in which luciferase-labeled CD19⁺ Raji cells were the target.

[0117] FIG. 39. Expression of chimeric antigen receptors following transduction with linearized DNA donor templates. These experiments generated plasmids that contain an anti-CD19 CAR gene flanked by homology arms that are homologous to the TRC 1-2 recognition sequence locus. Different promoters were used in some plasmids, and homology arms were either “short” (200 bp on the 5' homology arm and 180 bp on the 3' homology arm) or “long” (985 bp on the 5' homology arm and 763 bp on the 3' homology arm). CAR donor plasmids were linearized at a restriction site in the vector backbone and gel purified. A) Background CD3⁻/CAR⁺ staining. B) Cells electroporated with TRC 1-2x.87EE mRNA alone. C) Cells co-electroporated with TRC 1-2x.87EE mRNA and a long homology arm vector with an EF1 α core promoter with an HTLV enhancer. D) Cells co-electroporated with TRC 1-2x.87EE mRNA and a short homology arm vector with EF1 α core promoter (with no enhancer). E) Cells electroporated with a long homology arm vector with an EF1 α core promoter with an HTLV enhancer in the absence of TRC 1-2x.87EE mRNA. F) Cells electroporated with a short homology arm vector with EF1 α core promoter (with no enhancer) in the absence of TRC 1-2x.87EE mRNA. G) Cells electroporated with a long homology arm construct that contains an MND promoter driving expression of the CAR and an intron in the 5' end of the CAR gene, as well as TRC 1-2x.87EE mRNA. H) Cells electroporated with a long homology arm construct that contains an MND promoter driving expression of the CAR and no intron, as well as TRC 1-2x.87EE mRNA. I) Cells electroporated with a short homology arm plasmid with the MND promoter and no intron, as well as TRC 1-2x.87EE mRNA. J) Cells electroporated with a long homology arm construct that contains an MND promoter driving expression of the CAR and an intron in the 5' end of the CAR gene, but no TRC 1-2x.87EE mRNA. K) Cells electroporated with a long homology arm construct that contains an MND promoter driving expression of the CAR and no intron, but no TRC 1-2x.87EE mRNA. L) Cells electroporated with a short homology arm plasmid with the MND promoter and no intron, but no TRC 1-2x.87EE mRNA. M) Cells electroporated with a short homology arm construct that contained a JeT promoter, as well as TRC 1-2x.87EE mRNA. N) Cells electroporated with a long homology arm construct that contained a CMV promoter, as well as TRC 1-2x.87EE mRNA. O) Cells electroporated with a short homology arm construct that contained a JeT promoter, but no TRC 1-2x.87EE mRNA. P) Cells electroporated with a long homology arm construct that contained a CMV promoter, but no TRC 1-2x.87EE mRNA.

[0118] FIG. 40. PCR analysis to determine whether the chimeric antigen receptor coding region delivered by linearized DNA constructs was inserted into the TRC 1-2 recognition sequence in human T cells.

[0119] FIG. 41. Map of plasmid used to produce the AAV423 vector.

[0120] FIG. 42. Cell-surface expression of CD19 chimeric antigen receptor on human T cells. The expression level of

the anti-CD19 chimeric antigen receptor was determined in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV423 as the HDR template. Cell-surface expression was analyzed by flow cytometry. A) Cells that were mock electroporated and mock transduced (MOI-0), and cells that were mock electroporated and transduced with increasing amounts of AAV423. B) Cells that were electroporated with TRC 1-2x.87EE and mock transduced (MOI-0), and cells that were electroporated with TRC 1-2x.87EE and transduced with increasing amounts of AAV423.

[0121] FIG. 43. Insertion of chimeric antigen receptor coding sequence. PCR methods were used to determine if the chimeric antigen receptor coding sequence introduced by AAV423 inserted at the TRC 1-2 recognition site cleaved by the TRC 1-2x.87EE meganuclease.

[0122] FIG. 44. Phenotype analysis of anti-CD19 CAR T cells. A) Activated T cells were electroporated with TRC 1-2x.87EE mRNA, then transduced with an AAV6 vector comprising an anti-CD19 CAR expression cassette driven by a JeT promoter and flanked by homology arms. Following 5 days of culture with IL-2 (10 ng/mL), cells were analyzed for cell-surface CD3 and anti-CD19 CAR expression by flow cytometry. B) CD3⁻ cells were enriched by depleting CD3⁺ cells using anti-CD3 magnetic beads. Depleted cells were then cultured for 3 days in IL-15 (10 ng/mL) and IL-21 (10 ng/mL) and re-analyzed for cell-surface expression of CD3 and anti-CD19 CAR. C) The purified population of CD3⁻ CD19⁻ CAR T cells was analyzed by flow cytometry to determine the percentage of cells that were CD4⁺ and CD8⁺. D) The purified population of CD3⁻ CD19⁻ CAR T cells was further analyzed by flow cytometry to determine whether they were central memory T cells, transitional memory T cells, or effector memory T cells by staining for CD62L and CD45RO.

[0123] FIG. 45. Raji disseminated lymphoma model. Raji cells stably expressing firefly luciferase (flLuc)⁴⁴ were injected i.v. into 5-6 week old female NSG mice on Day 1, at a dose of 2.0x10⁵ cells per mouse. On Day 4 mice were injected i.v. with PBS or PBS containing gene edited control TCR KO T cells prepared from the same healthy donor PBMC or PBS containing the indicated doses of CAR T cells prepared from the same donor. On the indicated days, live mice were injected i.p. with Luciferin substrate (150 mg/kg in saline), anesthetized, and Luciferase activity measured after 7 minutes using IVIS SpectrumCT® (Perkin Elmer, Waltham, Mass.). Data was analyzed and exported using Living Image software 4.5.1 (Perkin Elmer, Waltham, Mass.). Luminescence signal intensity is represented by radiance in p/sec/cm²/sr.

BRIEF DESCRIPTION OF THE SEQUENCES

[0124] SEQ ID NO: 1 sets forth the nucleotide sequence of the human T cell receptor alpha constant region gene (NCBI Gene ID NO. 28755).

[0125] SEQ ID NO: 2 sets forth the amino acid sequence encoded by the human T cell receptor alpha constant region.

[0126] SEQ ID NO: 3 sets forth the amino acid sequence of the TRC 1-2 recognition sequence.

[0127] SEQ ID NO: 4 sets forth the nucleotide sequence of the TRC 3-4 recognition sequence.

[0128] SEQ ID NO: 5 sets forth the nucleotide sequence of the TRC 7-8 recognition sequence.

- [0129] SEQ ID NO: 6 sets forth the amino acid sequence of I-CreI.
- [0130] SEQ ID NO: 7 sets forth the amino acid sequence of the LAGLIDADG motif.
- [0131] SEQ ID NO: 8 sets forth the amino acid sequence of the TRC 1-2x.87 EE meganuclease.
- [0132] SEQ ID NO: 9 sets forth the amino acid sequence of the TRC 1-2x.87 QE meganuclease.
- [0133] SEQ ID NO: 10 sets forth the amino acid sequence of the TRC 1-2x.87 EQ meganuclease.
- [0134] SEQ ID NO: 11 sets forth the amino acid sequence of the TRC 1-2x.87 meganuclease.
- [0135] SEQ ID NO: 12 sets forth the amino acid sequence of the TRC 1-2x.6 meganuclease.
- [0136] SEQ ID NO: 13 sets forth the amino acid sequence of the TRC 1-2x.20 meganuclease.
- [0137] SEQ ID NO: 14 sets forth the amino acid sequence of the TRC 1-2x.55 meganuclease.
- [0138] SEQ ID NO: 15 sets forth the amino acid sequence of the TRC 1-2x.60 meganuclease.
- [0139] SEQ ID NO: 16 sets forth the amino acid sequence of the TRC 1-2x.105 meganuclease.
- [0140] SEQ ID NO: 17 sets forth the amino acid sequence of the TRC 1-2x.163 meganuclease.
- [0141] SEQ ID NO: 18 sets forth the amino acid sequence of the TRC 1-2x.113_3 meganuclease.
- [0142] SEQ ID NO: 19 sets forth the amino acid sequence of the TRC 1-2x.5 meganuclease.
- [0143] SEQ ID NO: 20 sets forth the amino acid sequence of the TRC 1-2x.8 meganuclease.
- [0144] SEQ ID NO: 21 sets forth the amino acid sequence of the TRC 1-2x.25 meganuclease.
- [0145] SEQ ID NO: 22 sets forth the amino acid sequence of the TRC 1-2x.72 meganuclease.
- [0146] SEQ ID NO: 23 sets forth the amino acid sequence of the TRC 1-2x.80 meganuclease.
- [0147] SEQ ID NO: 24 sets forth the amino acid sequence of the TRC 1-2x.84 meganuclease.
- [0148] SEQ ID NO: 25 sets forth the amino acid sequence of the TRC 1-2x.120 meganuclease.
- [0149] SEQ ID NO: 26 sets forth the amino acid sequence of the TRC 1-2x.113_1 meganuclease.
- [0150] SEQ ID NO: 27 sets forth the amino acid sequence of the TRC 1-2x.113_2 meganuclease.
- [0151] SEQ ID NO: 28 sets forth the amino acid sequence of the TRC 3-4x.3 meganuclease.
- [0152] SEQ ID NO: 29 sets forth the amino acid sequence of the TRC 3-4x.19 meganuclease.
- [0153] SEQ ID NO: 30 sets forth the amino acid sequence of the TRC 7-8x.7 meganuclease.
- [0154] SEQ ID NO: 31 sets forth the amino acid sequence of the TRC 7-8x.9 meganuclease.
- [0155] SEQ ID NO: 32 sets forth the amino acid sequence of the TRC 7-8x.14 meganuclease.
- [0156] SEQ ID NO: 33 sets forth residues 198-344 of the TRC 1-2x.87 EE meganuclease.
- [0157] SEQ ID NO: 34 sets forth residues 198-344 of the TRC 1-2x.87 QE meganuclease.
- [0158] SEQ ID NO: 35 sets forth residues 198-344 of the TRC 1-2x.87 EQ meganuclease.
- [0159] SEQ ID NO: 36 sets forth residues 198-344 of the TRC 1-2x.87 meganuclease.
- [0160] SEQ ID NO: 37 sets forth residues 198-344 of the TRC 1-2x.6 meganuclease.
- [0161] SEQ ID NO: 38 sets forth residues 198-344 of the TRC 1-2x.20 meganuclease.
- [0162] SEQ ID NO: 39 sets forth residues 198-344 of the TRC 1-2x.55 meganuclease.
- [0163] SEQ ID NO: 40 sets forth residues 198-344 of the TRC 1-2x.60 meganuclease.
- [0164] SEQ ID NO: 41 sets forth residues 198-344 of the TRC 1-2x.105 meganuclease.
- [0165] SEQ ID NO: 42 sets forth residues 198-344 of the TRC 1-2x.163 meganuclease.
- [0166] SEQ ID NO: 43 sets forth residues 198-344 of the TRC 1-2x.113_3 meganuclease.
- [0167] SEQ ID NO: 44 sets forth residues 7-153 of the TRC 1-2x.5 meganuclease.
- [0168] SEQ ID NO: 45 sets forth residues 7-153 of the TRC 1-2x.8 meganuclease.
- [0169] SEQ ID NO: 46 sets forth residues 7-153 of the TRC 1-2x.25 meganuclease.
- [0170] SEQ ID NO: 47 sets forth residues 7-153 of the TRC 1-2x.72 meganuclease.
- [0171] SEQ ID NO: 48 sets forth residues 7-153 of the TRC 1-2x.80 meganuclease.
- [0172] SEQ ID NO: 49 sets forth residues 7-153 of the TRC 1-2x.84 meganuclease.
- [0173] SEQ ID NO: 50 sets forth residues 7-153 of the TRC 1-2x.120 meganuclease.
- [0174] SEQ ID NO: 51 sets forth residues 7-153 of the TRC 1-2x.113_1 meganuclease.
- [0175] SEQ ID NO: 52 sets forth residues 7-153 of the TRC 1-2x.113_2 meganuclease.
- [0176] SEQ ID NO: 53 sets forth residues 7-153 of the TRC 3-4x.3 meganuclease.
- [0177] SEQ ID NO: 54 sets forth residues 7-153 of the TRC 3-4x.19 meganuclease.
- [0178] SEQ ID NO: 55 sets forth residues 7-153 of the TRC 7-8x.7 meganuclease.
- [0179] SEQ ID NO: 56 sets forth residues 198-344 of the TRC 7-8x.9 meganuclease.
- [0180] SEQ ID NO: 57 sets forth residues 198-344 of the TRC 7-8x.14 meganuclease.
- [0181] SEQ ID NO: 58 sets forth residues 7-153 of the TRC 1-2x.87 EE meganuclease.
- [0182] SEQ ID NO: 59 sets forth residues 7-153 of the TRC 1-2x.87 QE meganuclease.
- [0183] SEQ ID NO: 60 sets forth residues 7-153 of the TRC 1-2x.87 EQ meganuclease.
- [0184] SEQ ID NO: 61 sets forth residues 7-153 of the TRC 1-2x.87 meganuclease.
- [0185] SEQ ID NO: 62 sets forth residues 7-153 of the TRC 1-2x.6 meganuclease.
- [0186] SEQ ID NO: 63 sets forth residues 7-153 of the TRC 1-2x.20 meganuclease.
- [0187] SEQ ID NO: 64 sets forth residues 7-153 of the TRC 1-2x.55 meganuclease.
- [0188] SEQ ID NO: 65 sets forth residues 7-153 of the TRC 1-2x.60 meganuclease.
- [0189] SEQ ID NO: 66 sets forth residues 7-153 of the TRC 1-2x.105 meganuclease.
- [0190] SEQ ID NO: 67 sets forth residues 7-153 of the TRC 1-2x.163 meganuclease.
- [0191] SEQ ID NO: 68 sets forth residues 7-153 of the TRC 1-2x.113_3 meganuclease.
- [0192] SEQ ID NO: 69 sets forth residues 198-344 of the TRC 1-2x.5 meganuclease.

- [0193] SEQ ID NO: 70 sets forth residues 198-344 of the TRC 1-2x.8 meganuclease.
- [0194] SEQ ID NO: 71 sets forth residues 198-344 of the TRC 1-2x.25 meganuclease.
- [0195] SEQ ID NO: 72 sets forth residues 198-344 of the TRC 1-2x.72 meganuclease.
- [0196] SEQ ID NO: 73 sets forth residues 198-344 of the TRC 1-2x.80 meganuclease.
- [0197] SEQ ID NO: 74 sets forth residues 198-344 of the TRC 1-2x.84 meganuclease.
- [0198] SEQ ID NO: 75 sets forth residues 198-344 of the TRC 1-2x.120 meganuclease.
- [0199] SEQ ID NO: 76 sets forth residues 198-344 of the TRC 1-2x.113_1 meganuclease.
- [0200] SEQ ID NO: 77 sets forth residues 198-344 of the TRC 1-2x.113_2 meganuclease.
- [0201] SEQ ID NO: 78 sets forth residues 198-344 of the TRC 3-4x.3 meganuclease.
- [0202] SEQ ID NO: 79 sets forth residues 198-344 of the TRC 3-4x.19 meganuclease.
- [0203] SEQ ID NO: 80 sets forth residues 198-344 of the TRC 7-8x.7 meganuclease.
- [0204] SEQ ID NO: 81 sets forth residues 7-153 of the TRC 7-8x.9 meganuclease.
- [0205] SEQ ID NO: 82 sets forth residues 7-153 of the TRC 7-8x.14 meganuclease.
- [0206] SEQ ID NO: 83 sets forth the nucleotide sequence of the antisense strand of the TRC 1-2 recognition sequence.
- [0207] SEQ ID NO: 84 sets forth the nucleotide sequence of the antisense strand of the TRC 3-4 recognition sequence.
- [0208] SEQ ID NO: 85 sets forth the nucleotide sequence of the antisense strand of the TRC 7-8 recognition sequence.
- [0209] SEQ ID NO: 86 sets forth nucleotides 162-233 of SEQ ID NO:1.
- [0210] SEQ ID NO: 87 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0211] SEQ ID NO: 88 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0212] SEQ ID NO: 89 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0213] SEQ ID NO: 90 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0214] SEQ ID NO: 91 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0215] SEQ ID NO: 92 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0216] SEQ ID NO: 93 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0217] SEQ ID NO: 94 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising an insertion resulting from cleavage and NHEJ.
- [0218] SEQ ID NO: 95 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising an insertion resulting from cleavage and NHEJ.
- [0219] SEQ ID NO: 96 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0220] SEQ ID NO: 97 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0221] SEQ ID NO: 98 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0222] SEQ ID NO: 99 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0223] SEQ ID NO: 100 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0224] SEQ ID NO: 101 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0225] SEQ ID NO: 102 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0226] SEQ ID NO: 103 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0227] SEQ ID NO: 104 sets forth nucleotides 162-233 of SEQ ID NO:1 comprising a deletion resulting from cleavage and NHEJ.
- [0228] SEQ ID NO: 105 sets forth nucleotides 181-214 of SEQ ID NO:1.
- [0229] SEQ ID NO: 106 sets forth nucleotides 181-214 of SEQ ID NO:1 comprising an exogenous nucleic acid sequence inserted via homologous recombination.
- [0230] SEQ ID NO: 107 sets forth the nucleotide sequence of a plasmid used to generate the AAV405 vector.
- [0231] SEQ ID NO: 108 sets forth the nucleotide sequence of a plasmid used to generate the AAV406 vector.
- [0232] SEQ ID NO: 109 sets forth the nucleotide sequence of a plasmid used to generate the AAV-CAR100 (AAV408) vector.
- [0233] SEQ ID NO: 110 sets forth the nucleotide sequence of a plasmid used to generate the AAV-CAR763 (AAV412) vector.
- [0234] SEQ ID NO: 111 sets forth the amino acid sequence of an anti-CD19 chimeric antigen receptor.
- [0235] SEQ ID NO: 112 sets forth the amino acid sequence of an anti-CD19 extracellular ligand-binding domain.
- [0236] SEQ ID NO: 113 sets forth the amino acid sequence of a chimeric antigen receptor intracellular cytoplasmic signaling domain.
- [0237] SEQ ID NO: 114 sets forth the amino acid sequence of a chimeric antigen receptor intracellular co-stimulatory domain.
- [0238] SEQ ID NO: 115 sets forth the amino acid sequence of a chimeric antigen receptor signal peptide domain.
- [0239] SEQ ID NO: 116 sets forth the amino acid sequence of a chimeric antigen receptor hinge region.
- [0240] SEQ ID NO: 117 sets forth the amino acid sequence of a chimeric antigen receptor transmembrane domain.
- [0241] SEQ ID NO: 118 sets forth the nucleotide sequence of an EF-1 alpha core promoter.
- [0242] SEQ ID NO: 119 sets forth the nucleotide sequence of an exogenous polynucleotide insert.

[0243] SEQ ID NO: 120 sets forth the nucleotide sequence of the human TCR alpha constant region gene comprising an exogenous nucleic acid sequence inserted within the TRC 1-2 recognition sequence.

[0244] SEQ ID NO: 121 sets forth the nucleotide sequence of the human TCR alpha constant region gene comprising an exogenous nucleic acid sequence inserted within the TRC 3-4 recognition sequence.

[0245] SEQ ID NO: 122 sets forth the nucleotide sequence of the human TCR alpha constant region gene comprising an exogenous nucleic acid sequence inserted within the TRC 7-8 recognition sequence.

[0246] SEQ ID NO: 123 sets forth the nucleic acid sequence of a plasmid used to generate the AAV421 vector.

[0247] SEQ ID NO: 124 sets forth the nucleic acid sequence of a plasmid used to generate the AAV422 vector.

[0248] SEQ ID NO: 125 sets forth the nucleic acid sequence of a plasmid used to generate the AAV423 vector.

DETAILED DESCRIPTION OF THE INVENTION

1.1 References and Definitions

[0249] The patent and scientific literature referred to herein establishes knowledge that is available to those of skill in the art. The issued US patents, allowed applications, published foreign applications, and references, including GenBank database sequences, which are cited herein are hereby incorporated by reference to the same extent as if each was specifically and individually indicated to be incorporated by reference.

[0250] The present invention can be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art. For example, features illustrated with respect to one embodiment can be incorporated into other embodiments, and features illustrated with respect to a particular embodiment can be deleted from that embodiment. In addition, numerous variations and additions to the embodiments suggested herein will be apparent to those skilled in the art in light of the instant disclosure, which do not depart from the instant invention.

[0251] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention.

[0252] All publications, patent applications, patents, and other references mentioned herein are incorporated by reference herein in their entirety.

[0253] As used herein, “a,” “an,” or “the” can mean one or more than one. For example, “a” cell can mean a single cell or a multiplicity of cells.

[0254] As used herein, unless specifically indicated otherwise, the word “or” is used in the inclusive sense of “and/or” and not the exclusive sense of “either/or.”

[0255] As used herein, the term “meganuclease” refers to an endonuclease that binds double-stranded DNA at a recognition sequence that is greater than 12 base pairs. Preferably, the recognition sequence for a meganuclease of the

invention is 22 base pairs. A meganuclease can be an endonuclease that is derived from I-CreI, and can refer to an engineered variant of I-CreI that has been modified relative to natural I-CreI with respect to, for example, DNA-binding specificity, DNA cleavage activity, DNA-binding affinity, or dimerization properties. Methods for producing such modified variants of I-CreI are known in the art (e.g., WO 2007/047859). A meganuclease as used herein binds to double-stranded DNA as a heterodimer or as a “single-chain meganuclease” in which a pair of DNA-binding domains are joined into a single polypeptide using a peptide linker. The term “homing endonuclease” is synonymous with the term “meganuclease.” Meganucleases of the invention are substantially non-toxic when expressed in cells, particularly in human T cells, such that cells can be transfected and maintained at 37° C. without observing deleterious effects on cell viability or significant reductions in meganuclease cleavage activity when measured using the methods described herein.

[0256] As used herein, the term “single-chain meganuclease” refers to a polypeptide comprising a pair of nuclease subunits joined by a linker. A single-chain meganuclease has the organization: N-terminal subunit-Linker-C-terminal subunit. The two meganuclease subunits will generally be non-identical in amino acid sequence and will recognize non-identical DNA sequences. Thus, single-chain meganucleases typically cleave pseudo-palindromic or non-palindromic recognition sequences. A single-chain meganuclease may be referred to as a “single-chain heterodimer” or “single-chain heterodimeric meganuclease” although it is not, in fact, dimeric. For clarity, unless otherwise specified, the term “meganuclease” can refer to a dimeric or single-chain meganuclease.

[0257] As used herein, the term “linker” refers to an exogenous peptide sequence used to join two meganuclease subunits into a single polypeptide. A linker may have a sequence that is found in natural proteins, or may be an artificial sequence that is not found in any natural protein. A linker may be flexible and lacking in secondary structure or may have a propensity to form a specific three-dimensional structure under physiological conditions. A linker can include, without limitation, those encompassed by U.S. Pat. No. 8,445,251. In some embodiments, a linker may have an amino acid sequence comprising residues 154-195 of any one of SEQ ID NOs:8-32.

[0258] As used herein, the term “TALEN” refers to an endonuclease comprising a DNA-binding domain comprising 16-22 TAL domain repeats fused to any portion of the FokI nuclease domain.

[0259] As used herein, the term “Compact TALEN” refers to an endonuclease comprising a DNA-binding domain with 16-22 TAL domain repeats fused in any orientation to any catalytically active portion of nuclease domain of the I-TevI homing endonuclease.

[0260] As used herein, the term “CRISPR” refers to a caspase-based endonuclease comprising a caspase, such as Cas9, and a guide RNA that directs DNA cleavage of the caspase by hybridizing to a recognition site in the genomic DNA.

[0261] As used herein, the term “megaTAL” refers to a single-chain nuclease comprising a transcription activator-like effector (TALE) DNA binding domain with an engineered, sequence-specific homing endonuclease.

[0262] As used herein, with respect to a protein, the term “recombinant” means having an altered amino acid sequence as a result of the application of genetic engineering techniques to nucleic acids that encode the protein, and cells or organisms that express the protein. With respect to a nucleic acid, the term “recombinant” means having an altered nucleic acid sequence as a result of the application of genetic engineering techniques. Genetic engineering techniques include, but are not limited to, PCR and DNA cloning technologies; transfection, transformation and other gene transfer technologies; homologous recombination; site-directed mutagenesis; and gene fusion. In accordance with this definition, a protein having an amino acid sequence identical to a naturally-occurring protein, but produced by cloning and expression in a heterologous host, is not considered recombinant.

[0263] As used herein, the term “wild-type” refers to the most common naturally occurring allele (i.e., polynucleotide sequence) in the allele population of the same type of gene, wherein a polypeptide encoded by the wild-type allele has its original functions. The term “wild-type” also refers to a polypeptide encoded by a wild-type allele. Wild-type alleles (i.e., polynucleotides) and polypeptides are distinguishable from mutant or variant alleles and polypeptides, which comprise one or more mutations and/or substitutions relative to the wild-type sequence(s). Whereas a wild-type allele or polypeptide can confer a normal phenotype in an organism, a mutant or variant allele or polypeptide can, in some instances, confer an altered phenotype. Wild-type nucleases are distinguishable from recombinant or non-naturally-occurring nucleases.

[0264] As used herein with respect to recombinant proteins, the term “modification” means any insertion, deletion or substitution of an amino acid residue in the recombinant sequence relative to a reference sequence (e.g., a wild-type or a native sequence).

[0265] As used herein, the term “recognition sequence” refers to a DNA sequence that is bound and cleaved by an endonuclease. In the case of a meganuclease, a recognition sequence comprises a pair of inverted, 9 basepair “half sites” that are separated by four basepairs. In the case of a single-chain meganuclease, the N-terminal domain of the protein contacts a first half-site and the C-terminal domain of the protein contacts a second half-site. Cleavage by a meganuclease produces four basepair 3’ “overhangs”. “Overhangs”, or “sticky ends” are short, single-stranded DNA segments that can be produced by endonuclease cleavage of a double-stranded DNA sequence. In the case of meganucleases and single-chain meganucleases derived from I-CreI, the overhang comprises bases 10-13 of the 22 basepair recognition sequence. In the case of a Compact TALEN, the recognition sequence comprises a first C>NNNGN sequence that is recognized by the I-TevI domain, followed by a non-specific spacer 4-16 basepairs in length, followed by a second sequence 16-22 bp in length that is recognized by the TAL-effector domain (this sequence typically has a 5’ T base). Cleavage by a Compact TALEN produces two basepair 3’ overhangs. In the case of a CRISPR, the recognition sequence is the sequence, typically 16-24 basepairs, to which the guide RNA binds to direct Cas9 cleavage. Cleavage by a CRISPR produced blunt ends.

[0266] As used herein, the term “target site” or “target sequence” refers to a region of the chromosomal DNA of a cell comprising a recognition sequence for a nuclease.

[0267] As used herein, the term “DNA-binding affinity” or “binding affinity” means the tendency of a meganuclease to non-covalently associate with a reference DNA molecule (e.g., a recognition sequence or an arbitrary sequence). Binding affinity is measured by a dissociation constant, K_d . As used herein, a nuclease has “altered” binding affinity if the K_d of the nuclease for a reference recognition sequence is increased or decreased by a statistically significant percent change relative to a reference nuclease.

[0268] As used herein, the term “homologous recombination” or “HR” refers to the natural, cellular process in which a double-stranded DNA-break is repaired using a homologous DNA sequence as the repair template (see, e.g., Cahill et al. (2006), *From. Biosci.* 11:1958-1976). The homologous DNA sequence may be an endogenous chromosomal sequence or an exogenous nucleic acid that was delivered to the cell.

[0269] As used herein, the term “non-homologous end-joining” or “NHEJ” refers to the natural, cellular process in which a double-stranded DNA-break is repaired by the direct joining of two non-homologous DNA segments (see, e.g., Cahill et al. (2006), *Front. Biosci.* 11:1958-1976). DNA repair by non-homologous end-joining is error-prone and frequently results in the untemplated addition or deletion of DNA sequences at the site of repair. In some instances, cleavage at a target recognition sequence results in NHEJ at a target recognition site. Nuclease-induced cleavage of a target site in the coding sequence of a gene followed by DNA repair by NHEJ can introduce mutations into the coding sequence, such as frameshift mutations, that disrupt gene function. Thus, engineered nucleases can be used to effectively knock-out a gene in a population of cells.

[0270] As used herein, a “chimeric antigen receptor” or “CAR” refers to an engineered receptor that confers or grafts specificity for an antigen onto an immune effector cell (e.g., a human T cell). A chimeric antigen receptor typically comprises an extracellular ligand-binding domain or moiety and an intracellular domain that comprises one or more stimulatory domains that transduce the signals necessary for T cell activation. In some embodiments, the extracellular ligand-binding domain or moiety can be in the form of single-chain variable fragments (scFvs) derived from a monoclonal antibody, which provide specificity for a particular epitope or antigen (e.g., an epitope or antigen preferentially present on the surface of a cancer cell or other disease-causing cell or particle). The extracellular ligand-binding domain can be specific for any antigen or epitope of interest. In a particular embodiment, the ligand-binding domain is specific for CD19.

[0271] The extracellular domain of a chimeric antigen receptor can also comprise an autoantigen (see. Payne et al. (2016), *Science* 353 (6295): 179-184), that can be recognized by autoantigen-specific B cell receptors on B lymphocytes, thus directing T cells to specifically target and kill autoreactive B lymphocytes in antibody-mediated autoimmune diseases. Such CARs can be referred to as chimeric autoantibody receptors (CAARs), and their use is encompassed by the invention.

[0272] The scFvs can be attached via a linker sequence. The intracellular stimulatory domain can include one or more cytoplasmic signaling domains that transmit an acti-

vation signal to the immune effector cell following antigen binding. Such cytoplasmic signaling domains can include, without limitation, CD3-zeta. The intracellular stimulatory domain can also include one or more intracellular co-stimulatory domains that transmit a proliferative and/or cell-survival signal after ligand binding. Such intracellular co-stimulatory domains can include, without limitation, a CD28 domain, a 4-1BB domain, an OX40 domain, or a combination thereof. A chimeric antigen receptor can further include additional structural elements, including a transmembrane domain that is attached to the extracellular ligand-binding domain via a hinge or spacer sequence.

[0273] As used herein, an “exogenous T cell receptor” or “exogenous TCR” refers to a TCR whose sequence is introduced into the genome of an immune effector cell (e.g., a human T cell) that may or may not endogenously express the TCR. Expression of an exogenous TCR on an immune effector cell can confer specificity for a specific epitope or antigen (e.g., an epitope or antigen preferentially present on the surface of a cancer cell or other disease-causing cell or particle). Such exogenous T cell receptors can comprise alpha and beta chains or, alternatively, may comprise gamma and delta chains. Exogenous TCRs useful in the invention may have specificity to any antigen or epitope of interest.

[0274] As used herein, the term “reduced expression” refers to any reduction in the expression of the endogenous T cell receptor at the cell surface of a genetically-modified cell when compared to a control cell. The term reduced can also refer to a reduction in the percentage of cells in a population of cells that express an endogenous polypeptide (i.e., an endogenous T cell receptor) at the cell surface when compared to a population of control cells. Such a reduction may be up to 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or up to 100%. Accordingly, the term “reduced” encompasses both a partial knockdown and a complete knockdown of the endogenous T cell receptor.

[0275] As used herein with respect to both amino acid sequences and nucleic acid sequences, the terms “percent identity,” “sequence identity,” “percentage similarity,” “sequence similarity” and the like refer to a measure of the degree of similarity of two sequences based upon an alignment of the sequences that maximizes similarity between aligned amino acid residues or nucleotides, and that is a function of the number of identical or similar residues or nucleotides, the number of total residues or nucleotides, and the presence and length of gaps in the sequence alignment. A variety of algorithms and computer programs are available for determining sequence similarity using standard parameters. As used herein, sequence similarity is measured using the BLASTp program for amino acid sequences and the BLASTn program for nucleic acid sequences, both of which are available through the National Center for Biotechnology Information (www.ncbi.nlm.nih.gov/), and are described in, for example, Altschul et al. (1990), *J. Mol. Biol.* 215:403-410; Gish and States (1993), *Nature Genet.* 3:266-272; Madden et al. (1996), *Meth. Enzymol.* 266:131-141; Altschul et al. (1997), *Nucleic Acids Res.* 25:33 89-3402; Zhang et al. (2000), *J. Comput. Biol.* 7(1-2):203-14. As used herein, percent similarity of two amino acid sequences is the score based upon the following parameters for the BLASTp algorithm: word size=3; gap opening penalty=-11; gap extension penalty=-1; and scoring matrix=BLOSUM62. As used herein, percent similarity of two nucleic acid sequences is the score based upon the following parameters for the

BLASTn algorithm: word size=11; gap opening penalty=-5; gap extension penalty=-2; match reward=1 and mismatch penalty=-3.

[0276] As used herein with respect to modifications of two proteins or amino acid sequences, the term “corresponding to” is used to indicate that a specified modification in the first protein is a substitution of the same amino acid residue as in the modification in the second protein, and that the amino acid position of the modification in the first proteins corresponds to or aligns with the amino acid position of the modification in the second protein when the two proteins are subjected to standard sequence alignments (e.g., using the BLASTp program). Thus, the modification of residue “X” to amino acid “A” in the first protein will correspond to the modification of residue “Y” to amino acid “A” in the second protein if residues X and Y correspond to each other in a sequence alignment, and despite the fact that X and Y may be different numbers.

[0277] As used herein, the term “recognition half-site,” “recognition sequence half-site,” or simply “half-site” means a nucleic acid sequence in a double-stranded DNA molecule that is recognized by a monomer of a homodimeric or heterodimeric meganuclease, or by one subunit of a single-chain meganuclease.

[0278] As used herein, the term “hypervariable region” refers to a localized sequence within a meganuclease monomer or subunit that comprises amino acids with relatively high variability. A hypervariable region can comprise about 50-60 contiguous residues, about 53-57 contiguous residues, or preferably about 56 residues. In some embodiments, the residues of a hypervariable region may correspond to positions 24-79 or positions 215-270 of any one of SEQ ID NOs:8-32. A hypervariable region can comprise one or more residues that contact DNA bases in a recognition sequence and can be modified to alter base preference of the monomer or subunit. A hypervariable region can also comprise one or more residues that bind to the DNA backbone when the meganuclease associates with a double-stranded DNA recognition sequence. Such residues can be modified to alter the binding affinity of the meganuclease for the DNA backbone and the target recognition sequence. In different embodiments of the invention, a hypervariable region may comprise between 1-20 residues that exhibit variability and can be modified to influence base preference and/or DNA-binding affinity. In particular embodiments, a hypervariable region comprises between about 15-18 residues that exhibit variability and can be modified to influence base preference and/or DNA-binding affinity. In some embodiments, variable residues within a hypervariable region correspond to one or more of positions 24, 26, 28, 29, 30, 32, 33, 38, 40, 42, 44, 46, 66, 68, 70, 72, 73, 75, and 77 of any one of SEQ ID NOs:8-32. In other embodiments, variable residues within a hypervariable region correspond to one or more of positions 215, 217, 219, 221, 223, 224, 229, 231, 233, 235, 237, 248, 257, 259, 261, 263, 264, 266, and 268 of any one of SEQ ID NOs:8-32.

[0279] As used herein, the terms “T cell receptor alpha constant region gene” and “TCR alpha constant region gene” are used interchangeably and refer to the human gene identified by NCBI Gen ID NO. 28755 (SEQ ID NO:1).

[0280] The terms “recombinant DNA construct,” “recombinant construct,” “expression cassette,” “expression construct,” “chimeric construct,” “construct,” and “recombinant DNA fragment” are used interchangeably herein and are

single or double-stranded polynucleotides. A recombinant construct comprises an artificial combination of single or double-stranded polynucleotides, including, without limitation, regulatory and coding sequences that are not found together in nature. For example, a recombinant DNA construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source and arranged in a manner different than that found in nature. Such a construct may be used by itself or may be used in conjunction with a vector.

[0281] As used herein, a “vector” or “recombinant DNA vector” may be a construct that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. If a vector is used then the choice of vector is dependent upon the method that will be used to transform host cells as is well known to those skilled in the art. Vectors can include, without limitation, plasmid vectors and recombinant AAV vectors, or any other vector known in that art suitable for delivering a gene encoding a meganuclease of the invention to a target cell. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells comprising any of the isolated nucleotides or nucleic acid sequences of the invention.

[0282] As used herein, a “vector” can also refer to a viral vector. Viral vectors can include, without limitation, retroviral vectors, lentiviral vectors, adenoviral vectors, and adeno-associated viral vectors (AAV).

[0283] As used herein, a “polycistronic” mRNA refers to a single messenger RNA that comprises two or more coding sequences (i.e., cistrons) and encodes more than one protein. A polycistronic mRNA can comprise any element known in the art to allow for the translation of two or more genes from the same mRNA molecule including, but not limited to, an IRES element, a T2A element, a P2A element, an E2A element, and an F2A element.

[0284] As used herein, a “human T cell” or “T cell” refers to a T cell isolated from a human donor. Human T cells, and cells derived therefrom, include isolated T cells that have not been passaged in culture, T cells that have been passaged and maintained under cell culture conditions without immortalization, and T cells that have been immortalized and can be maintained under cell culture conditions indefinitely.

[0285] As used herein, a “control” or “control cell” refers to a cell that provides a reference point for measuring changes in genotype or phenotype of a genetically-modified cell. A control cell may comprise, for example: (a) a wild-type cell, i.e., of the same genotype as the starting material for the genetic alteration that resulted in the genetically-modified cell; (b) a cell of the same genotype as the genetically-modified cell but that has been transformed with a null construct (i.e., with a construct that has no known effect on the trait of interest); or, (c) a cell genetically identical to the genetically-modified cell but that is not exposed to conditions or stimuli or further genetic modifications that would induce expression of altered genotype or phenotype.

[0286] As used herein, the recitation of a numerical range for a variable is intended to convey that the invention may be practiced with the variable equal to any of the values within that range. Thus, for a variable that is inherently discrete, the variable can be equal to any integer value

within the numerical range, including the end-points of the range. Similarly, for a variable that is inherently continuous, the variable can be equal to any real value within the numerical range, including the end-points of the range. As an example, and without limitation, a variable that is described as having values between 0 and 2 can take the values 0, 1 or 2 if the variable is inherently discrete, and can take the values 0.0, 0.1, 0.01, 0.001, or any other real values ≥ 0 and ≤ 2 if the variable is inherently continuous.

2.1 Principle of the Invention

[0287] The present invention is based, in part, on the discovery that engineered nucleases can be utilized to recognize and cleave recognition sequences found within the human TCR alpha constant region gene (SEQ ID NO:1), such that NHEJ at the cleavage site disrupts expression of the TCR alpha chain subunit and, ultimately, expression of the T cell receptor at the cell surface. Moreover, according to the invention, an exogenous polynucleotide sequence is inserted into the TCR alpha constant region gene at the nuclease cleavage site, for example by homologous recombination, such that a sequence of interest is concurrently expressed in the cell. Such exogenous sequences can encode, for example, a chimeric antigen receptor, an exogenous TCR receptor, or any other polypeptide of interest.

[0288] Thus, the present invention allows for both the knockout of the endogenous T cell receptor and the expression of an exogenous nucleic acid sequence (e.g., a chimeric antigen receptor or exogenous TCR) by targeting a single recognition site with a single engineered nuclease. In particular embodiments where a sequence encoding a chimeric antigen receptor is inserted into the TCR alpha constant region gene, the invention provides a simplified method for producing an allogeneic T cell that expresses an antigen-specific CAR and has reduced expression, or complete knockout, of the endogenous TCR. Such cells can exhibit reduced or no induction of graft-versus-host-disease (GVHD) when administered to an allogeneic subject.

2.2 Nucleases for Recognizing and Cleaving Recognition Sequences within the T Cell Receptor Alpha Constant Region Gene

[0289] It is known in the art that it is possible to use a site-specific nuclease to make a DNA break in the genome of a living cell, and that such a DNA break can result in permanent modification of the genome via mutagenic NHEJ repair or via homologous recombination with a transgenic DNA sequence. NHEJ can produce mutagenesis at the cleavage site, resulting in inactivation of the allele. NHEJ-associated mutagenesis may inactivate an allele via generation of early stop codons, frameshift mutations producing aberrant non-functional proteins, or could trigger mechanisms such as nonsense-mediated mRNA decay. The use of nucleases to induce mutagenesis via NHEJ can be used to target a specific mutation or a sequence present in a wild-type allele. The use of nucleases to induce a double-strand break in a target locus is known to stimulate homologous recombination, particularly of transgenic DNA sequences flanked by sequences that are homologous to the genomic target. In this manner, exogenous nucleic acid sequences can be inserted into a target locus. Such exogenous nucleic acids can encode, for example, a chimeric antigen receptor, an exogenous TCR, or any sequence or polypeptide of interest.

[0290] In different embodiments, a variety of different types of nuclease are useful for practicing the invention. In one embodiment, the invention can be practiced using recombinant meganucleases. In another embodiment, the invention can be practiced using a CRISPR nuclease or CRISPR Nickase. Methods for making CRISPRs and CRISPR Nickases that recognize pre-determined DNA sites are known in the art, for example Ran, et al. (2013) *Nat Protoc.* 8:2281-308. In another embodiment, the invention can be practiced using TALENs or Compact TALENs. Methods for making TALE domains that bind to pre-determined DNA sites are known in the art, for example Reyon et al. (2012) *Nat Biotechnol.* 30:460-5. In a further embodiment, the invention can be practiced using megaTALs.

[0291] In preferred embodiments, the nucleases used to practice the invention are single-chain meganucleases. A single-chain meganuclease comprises an N-terminal subunit and a C-terminal subunit joined by a linker peptide. Each of the two domains recognizes half of the recognition sequence (i.e., a recognition half-site) and the site of DNA cleavage is at the middle of the recognition sequence near the interface of the two subunits. DNA strand breaks are offset by four base pairs such that DNA cleavage by a meganuclease generates a pair of four base pair, 3' single-strand overhangs.

[0292] In some examples, recombinant meganucleases of the invention have been engineered to recognize and cleave the TRC 1-2 recognition sequence (SEQ ID NO:3). Such recombinant meganucleases are collectively referred to herein as "TRC 1-2 meganucleases." Exemplary TRC 1-2 meganucleases are provided in SEQ ID NOs:8-27.

[0293] In additional examples, recombinant meganucleases of the invention have been engineered to recognize and cleave the TRC 3-4 recognition sequence (SEQ ID NO:4).

Such recombinant meganucleases are collectively referred to herein as "TRC 3-4 meganucleases." Exemplary TRC 3-4 meganucleases are provided in SEQ ID NOs:28 and 29.

[0294] In further examples, recombinant meganucleases of the invention have been engineered to recognize and cleave the TRC 7-8 recognition sequence (SEQ ID NO:5). Such recombinant meganucleases are collectively referred to herein as "TRC 7-8 meganucleases." Exemplary TRC 7-8 meganucleases are provided in SEQ ID NOs:30-32.

[0295] Recombinant meganucleases of the invention comprise a first subunit, comprising a first hypervariable (HVR1) region, and a second subunit, comprising a second hypervariable (HVR2) region. Further, the first subunit binds to a first recognition half-site in the recognition sequence (e.g., the TRC1, TRC3, or TRC7 half-site), and the second subunit binds to a second recognition half-site in the recognition sequence (e.g., the TRC2, TRC4, or TRC8 half-site). In embodiments where the recombinant meganuclease is a single-chain meganuclease, the first and second subunits can be oriented such that the first subunit, which comprises the HVR1 region and binds the first half-site, is positioned as the N-terminal subunit, and the second subunit, which comprises the HVR2 region and binds the second half-site, is positioned as the C-terminal subunit. In alternative embodiments, the first and second subunits can be oriented such that the first subunit, which comprises the HVR1 region and binds the first half-site, is positioned as the C-terminal subunit, and the second subunit, which comprises the HVR2 region and binds the second half-site, is positioned as the N-terminal subunit. Exemplary TRC 1-2 meganucleases of the invention are provided in Table 1. Exemplary TRC 3-4 meganucleases of the invention are provided in Table 2. Exemplary TRC 7-8 meganucleases of the invention are provided in Table 3.

TABLE 1

Exemplary recombinant meganucleases engineered to recognize and cleave the TRC 1-2 recognition sequence (SEQ ID NO: 3)							
Meganuclease	AA SEQ ID	TRC1 Subunit Residues	TRC1 Subunit SEQ ID	*TRC1 Subunit %	TRC2 Subunit Residues	TRC2 Subunit SEQ ID	*TRC2 Subunit %
TRC 1-2x.87 EE	8	198-344	33	100	7-153	58	100
TRC 1-2x.87 QE	9	198-344	34	100	7-153	59	99.3
TRC 1-2x.87 EQ	10	198-344	35	99.3	7-153	60	100
TRC 1-2x.87	11	198-344	36	99.3	7-153	61	99.3
TRC 1-2x.6	12	198-344	37	99.3	7-153	62	94.6
TRC 1-2x.20	13	198-344	38	99.3	7-153	63	91.2
TRC 1-2x.55	14	198-344	39	95.9	7-153	64	91.8
TRC 1-2x.60	15	198-344	40	91.8	7-153	65	91.2
TRC 1-2x.105	16	198-344	41	95.2	7-153	66	95.2
TRC 1-2x.163	17	198-344	42	99.3	7-153	67	99.3
TRC 1-2x.113_3	18	198-344	43	99.3	7-153	68	91.2
TRC 1-2x.5	19	7-153	44	99.3	198-344	69	93.2
TRC 1-2x.8	20	7-153	45	92.5	198-344	70	92.5
TRC 1-2x.25	21	7-153	46	99.3	198-344	71	98.6
TRC 1-2x.72	22	7-153	47	99.3	198-344	72	92.5
TRC 1-2x.80	23	7-153	48	99.3	198-344	73	92.5
TRC 1-2x.84	24	7-153	49	95.2	198-344	74	98.6
TRC 1-2x.120	25	7-153	50	99.3	198-344	75	92.5
TRC 1-2x.113_1	26	7-153	51	100	198-344	76	92.5
TRC 1-2x.113_2	27	7-153	52	99.3	198-344	77	92.5

**"TRC1 Subunit %" and "TRC2 Subunit %" represent the amino acid sequence identity between the TRC1-binding and TRC2-binding subunit regions of each meganuclease and the TRC1-binding and TRC2-binding subunit regions, respectively, of the TRC 1-2x.87 EE meganuclease.

TABLE 2

Exemplary recombinant meganucleases engineered to recognize and cleave the TRC 3-4 recognition sequence (SEQ ID NO: 4)							
Meganuclease	AA SEQ ID	TRC3 Subunit Residues	TRC3 Subunit SEQ ID	*TRC3 Subunit %	TRC4 Subunit Residues	TRC4 Subtotal SEQ ID	TRC4 Subunit %
TRC 3-4x.3	28	7-153	53	100	198-344	78	100
TRC 3-4x.19	29	7-153	54	96.6	198-344	79	96.6

**“TRC3 Subunit %” and “TRC4 Subunit %” represent the amino acid sequence identity between the TRC3-binding and TRC4-binding subunit regions of each meganuclease and the TRC3-binding and TRC4-binding subunit regions, respectively, of the TRC 3-4x.3 meganuclease.

TABLE 3

Exemplary recombinant meganucleases engineered to recognize and cleave the TRC 7-8 recognition sequence (SEQ ID NO: 5)							
Meganuclease	AA SEQ ID	TRC7 Subunit Residues	TRC7 Subunit SEQ ID	*TRC7 Subunit %	TRC8 Subunit Residues	TRC8 Subunit SEQ ID	TRC8 Subunit %
TRC 7-8x.7	30	7-153	55	100	198-344	80	100
TRC 7-8x.9	31	198-344	56	97.3	7-153	81	91.2
TRC 7-8x.14	32	198-344	57	97.9	7-153	82	90.5

**“TRC7 Subunit %” and “TRC8 Subunit %” represent the amino acid sequence identity between the TRC7-binding and TRC8-binding subunit regions of each meganuclease and the TRC7-binding and TRC8-binding subunit regions, respectively, of the TRC 7-8x.7 meganuclease.

2.3 Methods for Producing Genetically-Modified Cells

[0296] The invention provides methods for producing genetically-modified cells using engineered nucleases that recognize and cleave recognition sequences found within the human TCR alpha constant region gene (SEQ ID NO:1). Cleavage at such recognition sequences can allow for NHEJ at the cleavage site and disrupted expression of the human T cell receptor alpha chain subunit, leading to reduced expression and/or function of the T cell receptor at the cell surface. Additionally, cleavage at such recognition sequences can further allow for homologous recombination of exogenous nucleic acid sequences directly into the TCR alpha constant region gene.

[0297] Engineered nucleases of the invention can be delivered into a cell in the form of protein or, preferably, as a nucleic acid encoding the engineered nuclease. Such nucleic acid can be DNA (e.g., circular or linearized plasmid DNA or PCR products) or RNA. For embodiments in which the engineered nuclease coding sequence is delivered in DNA form, it should be operably linked to a promoter to facilitate transcription of the meganuclease gene. Mammalian promoters suitable for the invention include constitutive promoters such as the cytomegalovirus early (CMV) promoter (Thomsen et al. (1984). *Proc Natl Acad Sci USA*. 81(3):659-63) or the SV40 early promoter (Benoist and Chambon (1981). *Nature*. 290(5804):304-10) as well as inducible promoters such as the tetracycline-inducible promoter (Dingermann et al. (1992), *Mol Cell Biol*. 12(9):4038-45).

[0298] In some embodiments, mRNA encoding the engineered nuclease is delivered to the cell because this reduces the likelihood that the gene encoding the engineered nuclease will integrate into the genome of the cell. Such mRNA encoding an engineered nuclease can be produced using methods known in the art such as in vitro transcription. In

some embodiments, the mRNA is capped using 7-methyl-guanosine. In some embodiments, the mRNA may be polyadenylated.

[0299] In particular embodiments, an mRNA encoding an engineered nuclease of the invention can be a polycistronic mRNA encoding two or more nucleases that are simultaneously expressed in the cell. A polycistronic mRNA can encode two or more nucleases of the invention that target different recognition sequences in the same target gene. Alternatively, a polycistronic mRNA can encode at least one nuclease described herein and at least one additional nuclease targeting a separate recognition sequence positioned in the same gene, or targeting a second recognition sequence positioned in a second gene such that cleavage sites are produced in both genes. A polycistronic mRNA can comprise any element known in the art to allow for the translation of two or more genes (i.e., cistrons) from the same mRNA molecule including, but not limited to, an IRES element, a T2A element, a P2A element, an E2A element, and an F2A element.

[0300] Purified nuclease proteins can be delivered into cells to cleave genomic DNA, which allows for homologous recombination or non-homologous end-joining at the cleavage site with a sequence of interest, by a variety of different mechanisms known in the art.

[0301] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are coupled to a cell penetrating peptide or targeting ligand to facilitate cellular uptake. Examples of cell penetrating peptides known in the art include poly-arginine (Jearawiriyapaisarn, et al. (2008) *Mol Ther*. 16:1624-9). TAT peptide from the HIV virus (Hudecz et al. (2005), *Med. Res. Rev.* 25: 679-736), MPG (Simeoni, et al. (2003) *Nucleic Acids Res.* 31:2717-2724). Pep-1 (Deshayes et al. (2004) *Biochemistry* 43: 7698-7706, and HSV-1 VP-22 (Deshayes et al. (2005) *Cell Mol Life Sci.* 62:1839-49. In an alternative embodiment, engineered nucleases, or DNA/mRNA encoding engi-

neered nucleases, are coupled covalently or non-covalently to an antibody that recognizes a specific cell-surface receptor expressed on target cells such that the nuclease protein/DNA/mRNA binds to and is internalized by the target cells. Alternatively, engineered nuclease protein/DNA/mRNA can be coupled covalently or non-covalently to the natural ligand (or a portion of the natural ligand) for such a cell-surface receptor. (McCall, et al. (2014) *Tissue Barriers*. 2(4): e944449; Dinda, et al. (2013) *Curr Pharm Biotechnol*. 14:1264-74; Kang, et al. (2014) *Curr Pharm Biotechnol*. 15(3):220-30; Qian et al. (2014) *Expert Opin Drug Metab Toxicol*. 10(11):1491-508).

[0302] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are coupled covalently or, preferably, non-covalently to a nanoparticle or encapsulated within such a nanoparticle using methods known in the art (Sharma, et al. (2014) *Biomed Res Int*. 2014). A nanoparticle is a nanoscale delivery system whose length scale is <1 μm , preferably <100 nm. Such nanoparticles may be designed using a core composed of metal, lipid, polymer, or biological macromolecule, and multiple copies of the recombinant meganuclease proteins, mRNA, or DNA can be attached to or encapsulated with the nanoparticle core. This increases the copy number of the protein/mRNA/DNA that is delivered to each cell and, so, increases the intracellular expression of each engineered nuclease to maximize the likelihood that the target recognition sequences will be cut. The surface of such nanoparticles may be further modified with polymers or lipids (e.g., chitosan, cationic polymers, or cationic lipids) to form a core-shell nanoparticle whose surface confers additional functionalities to enhance cellular delivery and uptake of the payload (Jian et al. (2012) *Biomaterials*. 33(30): 7621-30). Nanoparticles may additionally be advantageously coupled to targeting molecules to direct the nanoparticle to the appropriate cell type and/or increase the likelihood of cellular uptake. Examples of such targeting molecules include antibodies specific for cell-surface receptors and the natural ligands (or portions of the natural ligands) for cell surface receptors.

[0303] In some embodiments, the engineered nucleases or DNA/mRNA encoding the engineered nucleases, are encapsulated within liposomes or complexed using cationic lipids (see, e.g., Lipofectamine™, Life Technologies Corp., Carlsbad, Calif.; Zuris et al. (2015) *Nat Biotechnol*. 33: 73-80; Mishra et al. (2011) *J Drug Deliv*. 2011:863734). The liposome and lipoplex formulations can protect the payload from degradation, and facilitate cellular uptake and delivery efficiency through fusion with and/or disruption of the cellular membranes of the cells.

[0304] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are encapsulated within polymeric scaffolds (e.g., PLGA) or complexed using cationic polymers (e.g., PEI, PLL) (Tamboli et al. (2011) *Ther Deliv*. 2(4): 523-536).

[0305] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are combined with amphiphilic molecules that self-assemble into micelles (Tong et al. (2007) *J Gene Med*. 9(11): 956-66). Polymeric micelles may include a micellar shell formed with a hydrophilic polymer (e.g., polyethyleneglycol) that can prevent aggregation, mask charge interactions, and reduce nonspecific interactions outside of the cell.

[0306] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are formulated into an emulsion or a nanoemulsion (i.e., having an average particle diameter of <1 nm) for delivery to the cell. The term “emulsion” refers to, without limitation, any oil-in-water, water-in-oil, water-in-oil-in-water, or oil-in-water-in-oil dispersions or droplets, including lipid structures that can form as a result of hydrophobic forces that drive apolar residues (e.g., long hydrocarbon chains) away from water and polar head groups toward water, when a water immiscible phase is mixed with an aqueous phase. These other lipid structures include, but are not limited to, unilamellar, paucilamellar, and multilamellar lipid vesicles, micelles, and lamellar phases. Emulsions are composed of an aqueous phase and a lipophilic phase (typically containing an oil and an organic solvent). Emulsions also frequently contain one or more surfactants. Nanoemulsion formulations are well known, e.g., as described in US Patent Application Nos. 2002/0045667 and 2004/0043041, and U.S. Pat. Nos. 6,015,832, 6,506,803, 6,635,676, and 6,559,189, each of which is incorporated herein by reference in its entirety.

[0307] In some embodiments, engineered nuclease proteins, or DNA/mRNA encoding engineered nucleases, are covalently attached to, or non-covalently associated with, multifunctional polymer conjugates. DNA dendrimers, and polymeric dendrimers (Mastorakos et al. (2015) *Nanoscale*. 7(9): 3845-56; Cheng et al. (2008) *J Pharm Sci*. 97(1): 123-43). The dendrimer generation can control the payload capacity and size, and can provide a high payload capacity. Moreover, display of multiple surface groups can be leveraged to improve stability and reduce nonspecific interactions.

[0308] In some embodiments, genes encoding an engineered nuclease are introduced into a cell using a viral vector. Such vectors are known in the art and include lentiviral vectors, adenoviral vectors, and adeno-associated virus (AAV) vectors (reviewed in Vannucci, et al. (2013) *New Microbiol*. 36:1-22). Recombinant AAV vectors useful in the invention can have any serotype that allows for transduction of the virus into the cell and insertion of the nuclease gene into the cell genome. In particular embodiments, recombinant AAV vectors have a serotype of AAV2 or AAV6. Recombinant AAV vectors can also be self-complementary such that they do not require second-strand DNA synthesis in the host cell (McCarty, et al. (2001) *Gene Ther*. 8:1248-54).

[0309] If the engineered nuclease genes are delivered in DNA form (e.g. plasmid) and/or via a viral vector (e.g. AAV) they must be operably linked to a promoter. In some embodiments, this can be a viral promoter such as endogenous promoters from the viral vector (e.g. the LTR of a lentiviral vector) or the well-known cytomegalovirus- or SV40 virus-early promoters. In a preferred embodiment, nuclease genes are operably linked to a promoter that drives gene expression preferentially in the target cell (e.g., a human T cell).

[0310] The invention further provides for the introduction of an exogenous nucleic acid into the cell, such that the exogenous nucleic acid sequence is inserted into the TRC alpha constant region gene at a nuclease cleavage site. In some embodiments, the exogenous nucleic acid comprises a 5' homology arm and a 3' homology arm to promote recombination of the nucleic acid sequence into the cell genome at the nuclease cleavage site.

[0311] Exogenous nucleic acids of the invention may be introduced into the cell by any of the means previously discussed. In a particular embodiment, exogenous nucleic acids are introduced by way of a viral vector, such as a lentivirus, retrovirus, adenovirus, or preferably a recombinant AAV vector. Recombinant AAV vectors useful for introducing an exogenous nucleic acid can have any serotype that allows for transduction of the virus into the cell and insertion of the exogenous nucleic acid sequence into the cell genome. In particular embodiments, the recombinant AAV vectors have a serotype of AAV2 or AAV6. The recombinant AAV vectors can also be self-complementary such that they do not require second-strand DNA synthesis in the host cell.

[0312] In another particular embodiment, an exogenous nucleic acid can be introduced into the cell using a single-stranded DNA template. The single-stranded DNA can comprise the exogenous nucleic acid and, in preferred embodiments, can comprise 5' and 3' homology arms to promote insertion of the nucleic acid sequence into the nuclease cleavage site by homologous recombination. The single-stranded DNA can further comprise a 5' AAV inverted terminal repeat (ITR) sequence 5' upstream of the 5' homology arm, and a 3' AAV ITR sequence 3' downstream of the 3' homology arm.

[0313] In another particular embodiment, genes encoding an endonuclease of the invention and/or an exogenous nucleic acid sequence of the invention can be introduced into the cell by transfection with a linearized DNA template. In some examples, a plasmid DNA encoding an endonuclease and/or an exogenous nucleic acid sequence can be digested by one or more restriction enzymes such that the circular plasmid DNA is linearized prior to transfection into the cell.

[0314] When delivered to a cell, an exogenous nucleic acid of the invention can be operably linked to any promoter suitable for expression of the encoded polypeptide in the cell, including those mammalian promoters and inducible promoters previously discussed. An exogenous nucleic acid of the invention can also be operably linked to a synthetic promoter. Synthetic promoters can include, without limitation, the JeT promoter (WO 2002/012514).

[0315] In examples where the genetically-modified cells of the invention are human T cells, or cells derived therefrom, such cells may require activation prior to introduction of a meganuclease and/or an exogenous nucleic acid sequence. For example, T cells can be contacted with anti-CD3 and anti-CD28 antibodies that are soluble or conjugated to a support (i.e., beads) for a period of time sufficient to activate the cells.

[0316] Genetically-modified cells of the invention can be further modified to express one or more inducible suicide genes, the induction of which provokes cell death and allows for selective destruction of the cells in vitro or in vivo. In some examples, a suicide gene can encode a cytotoxic polypeptide, a polypeptide that has the ability to convert a non-toxic pro-drug into a cytotoxic drug, and/or a polypeptide that activates a cytotoxic gene pathway within the cell. That is, a suicide gene is a nucleic acid that encodes a product that causes cell death by itself or in the presence of other compounds. A representative example of such a suicide gene is one that encodes thymidine kinase of herpes simplex virus. Additional examples are genes that encode thymidine kinase of varicella zoster virus and the bacterial

gene cytosine deaminase that can convert 5-fluorocytosine to the highly toxic compound 5-fluorouracil. Suicide genes also include as non-limiting examples genes that encode caspase-9, caspase-8, or cytosine deaminase. In some examples, caspase-9 can be activated using a specific chemical inducer of dimerization (CID). A suicide gene can also encode a polypeptide that is expressed at the surface of the cell that makes the cells sensitive to therapeutic and/or cytotoxic monoclonal antibodies. In further examples, a suicide gene can encode recombinant antigenic polypeptide comprising an antigenic motif recognized by the anti-CD20 mAb Rituximab and an epitope that allows for selection of cells expressing the suicide gene. See, for example, the RQR8 polypeptide described in WO2013153391, which comprises two Rituximab-binding epitopes and a QBEndO-binding epitope. For such a gene, Rituximab can be administered to a subject to induce cell depletion when needed.

2.4 Pharmaceutical Compositions

[0317] In some embodiments, the invention provides a pharmaceutical composition comprising a genetically-modified cell of the invention, or a population of genetically-modified cells of the invention, and a pharmaceutical carrier. Such pharmaceutical compositions can be prepared in accordance with known techniques. See, e.g., Remington, *The Science And Practice of Pharmacy* (21st ed. 2005). In the manufacture of a pharmaceutical formulation according to the invention, cells are typically admixed with a pharmaceutically acceptable carrier and the resulting composition is administered to a subject. The carrier must, of course, be acceptable in the sense of being compatible with any other ingredients in the formulation and must not be deleterious to the subject. In some embodiments, pharmaceutical compositions of the invention can further comprise one or more additional agents useful in the treatment of a disease in the subject. In additional embodiments, where the genetically-modified cell is a genetically-modified human T cell (or a cell derived therefrom), pharmaceutical compositions of the invention can further include biological molecules, such as cytokines (e.g., IL-2, IL-7, IL-15, and/or IL-21), which promote in vivo cell proliferation and engraftment. Pharmaceutical compositions comprising genetically-modified cells of the invention can be administered in the same composition as an additional agent or biological molecule or, alternatively, can be co-administered in separate compositions.

[0318] Pharmaceutical compositions of the invention can be useful for treating any disease state that can be targeted by T cell adoptive immunotherapy. In a particular embodiment, the pharmaceutical compositions of the invention are useful in the treatment of cancer. Such cancers can include, without limitation, carcinoma, lymphoma, sarcoma, blastomas, leukemia, cancers of B-cell origin, breast cancer, gastric cancer, neuroblastoma, osteosarcoma, lung cancer, melanoma, prostate cancer, colon cancer, renal cell carcinoma, ovarian cancer, rhabdomyo sarcoma, leukemia, and Hodgkin's lymphoma. In certain embodiments, cancers of B-cell origin include, without limitation, B-lineage acute lymphoblastic leukemia, B-cell chronic lymphocytic leukemia, and B-cell non-Hodgkin's lymphoma.

2.5 Methods for Producing Recombinant AAV Vectors

[0319] In some embodiments, the invention provides recombinant AAV vectors for use in the methods of the

invention. Recombinant AAV vectors are typically produced in mammalian cell lines such as HEK-293. Because the viral cap and rep genes are removed from the vector to prevent its self-replication to make room for the therapeutic gene(s) to be delivered (e.g. the endonuclease gene), it is necessary to provide these in trans in the packaging cell line. In addition, it is necessary to provide the “helper” (e.g. adenoviral) components necessary to support replication (Cots D, Bosch A, Chillon M (2013) *Curr. Gene Ther.* 13(5): 370-81). Frequently, recombinant AAV vector are produced using a triple-transfection in which a cell line is transfected with a first plasmid encoding the “helper” components, a second plasmid comprising the cap and rep genes, and a third plasmid comprising the viral ITRs containing the intervening DNA sequence to be packaged into the virus. Viral particles comprising a genome (ITRs and intervening gene (s) of interest) encased in a capsid are then isolated from cells by freeze-thaw cycles, sonication, detergent, or other means known in the art. Particles are then purified using cesium-chloride density gradient centrifugation or affinity chromatography and subsequently delivered to the gene(s) of interest to cells, tissues, or an organism such as a human patient.

[0320] Because recombinant AAV particles are typically produced (manufactured) in cells, precautions must be taken in practicing the current invention to ensure that the site-specific endonuclease is NOT expressed in the packaging cells. Because the viral genomes of the invention comprise a recognition sequence for the endonuclease, any endonuclease expressed in the packaging cell line will be capable of cleaving the viral genome before it can be packaged into viral particles. This will result in reduced packaging efficiency and/or the packaging of fragmented genomes. Several approaches can be used to prevent endonuclease expression in the packaging cells, including:

[0321] 1. The endonuclease can be placed under the control of a tissue-specific promoter that is not active in the packaging cells. For example, if a viral vector is developed for delivery of (an) endonuclease gene(s) to muscle tissue, a muscle-specific promoter can be used. Examples of muscle-specific promoters include C5-12 (Liu, et al. (2004) *Hum Gene Ther.* 15:783-92), the muscle-specific creatine kinase (MCK) promoter (Yuasa, et al. (2002) *Gene Ther.* 9:1576-88), or the smooth muscle 22 (SM22) promoter (Haase, et al. (2013) *BMC Biotechnol.* 13:49-54). Examples of CNS (neuron)-specific promoters include the NSE, Synapsin, and MSCP2 promoters (Lentz, et al. (2012) *Neurobiol Dis.* 48:179-88). Examples of liver-specific promoters include albumin promoters (such as Palb), human α 1-antitrypsin (such as PalAT), and hemopexin (such as Phpx) (Kramer, M G et al., (2003) *Mol. Therapy* 7:375-85). Examples of eye-specific promoters include opsin, and corneal epithelium-specific K12 promoters (Martin K R G, Klein R L, and Quigley H A (2002) *Methods* (28): 267-75) (Tong Y, et al., (2007) *J Gene Med.* 9:956-66). These promoters, or other tissue-specific promoters known in the art, are not highly-active in HEK-293 cells and, thus, will not be expected to yield significant levels of endonuclease gene expression in packaging cells when incorporated into viral vectors of the present invention. Similarly, the viral vectors of the present invention contemplate the use of other cell lines with the use of incompatible tissue

specific promoters (i.e., the well-known HeLa cell line (human epithelial cell) and using the liver-specific hemopexin promoter). Other examples of tissue specific promoters include: synovial sarcoma PDZD4 (cerebellum), C6 (liver), ASB5 (muscle), PPP1R12B (heart), SLC5A12 (kidney), cholesterol regulation APOM (liver), ADPRHL1 (heart), and monogenic malformation syndromes TP73L (muscle). (Jacox E, et al., (2010) *PLoS One* v. 5(8):e12274).

[0322] 2. Alternatively, the vector can be packaged in cells from a different species in which the endonuclease is not likely to be expressed. For example, viral particles can be produced in microbial, insect, or plant cells using mammalian promoters, such as the well-known cytomegalovirus- or SV40 virus-early promoters, which are not active in the non-mammalian packaging cells. In a preferred embodiment, viral particles are produced in insect cells using the baculovirus system as described by Gao, et al. (Gao, H., et al. (2007) *J. Biotechnol.* 131(2):138-43). An endonuclease under the control of a mammalian promoter is unlikely to be expressed in these cells (Airenne, K J, et al. (2013) *Mol. Ther.* 21(4):739-49). Moreover, insect cells utilize different mRNA splicing motifs than mammalian cells. Thus, it is possible to incorporate a mammalian intron, such as the human growth hormone (HGH) intron or the SV40 large T antigen intron, into the coding sequence of an endonuclease. Because these introns are not spliced efficiently from pre-mRNA transcripts in insect cells, insect cells will not express a functional endonuclease and will package the full-length genome. In contrast, mammalian cells to which the resulting recombinant AAV particles are delivered will properly splice the pre-mRNA and will express functional endonuclease protein. Haifeng Chen has reported the use of the HGH and SV40 large T antigen introns to attenuate expression of the toxic proteins barnase and diphtheria toxin fragment A in insect packaging cells, enabling the production of recombinant AAV vectors carrying these toxin genes (Chen. H (2012) *Mol Ther Nucleic Acids.* 1(11): e57).

[0323] 3. The endonuclease gene can be operably linked to an inducible promoter such that a small-molecule inducer is required for endonuclease expression. Examples of inducible promoters include the Tet-On system (Clontech; Chen H., et al., (2015) *BMC Biotechnol.* 15(1):4) and the RheoSwitch system (Intron; Sowa G., et al., (2011) *Spine*, 36(10): E623-8). Both systems, as well as similar systems known in the art, rely on ligand-inducible transcription factors (variants of the Tet Repressor and Ecdysone receptor, respectively) that activate transcription in response to a small-molecule activator (Doxycycline or Ecdysone, respectively). Practicing the current invention using such ligand-inducible transcription activators includes: 1) placing the endonuclease gene under the control of a promoter that responds to the corresponding transcription factor, the endonuclease gene having (a) binding site(s) for the transcription factor; and 2) including the gene encoding the transcription factor in the packaged viral genome. The latter step is necessary because the endonuclease will not be expressed in the target cells or tissues following recombinant AAV delivery if the transcription activator is not also provided to the

same cells. The transcription activator then induces endonuclease gene expression only in cells or tissues that are treated with the cognate small-molecule activator. This approach is advantageous because it enables endonuclease gene expression to be regulated in a spatio-temporal manner by selecting when and to which tissues the small-molecule inducer is delivered. However, the requirement to include the inducer in the viral genome, which has significantly limited carrying capacity, creates a drawback to this approach.

[0324] 4. In another preferred embodiment, recombinant AAV particles are produced in a mammalian cell line that expresses a transcription repressor that prevents expression of the endonuclease. Transcription repressors are known in the art and include the Tet-Repressor, the Lac-Repressor, the Cro repressor, and the Lambda-repressor. Many nuclear hormone receptors such as the ecdysone receptor also act as transcription repressors in the absence of their cognate hormone ligand. To practice the current invention, packaging cells are transfected/transduced with a vector encoding a transcription repressor and the endonuclease gene in the viral genome (packaging vector) is operably linked to a promoter that is modified to comprise binding sites for the repressor such that the repressor silences the promoter. The gene encoding the transcription repressor can be placed in a variety of positions. It can be encoded on a separate vector; it can be incorporated into the packaging vector outside of the ITR sequences; it can be incorporated into the cap/rep vector or the adenoviral helper vector; or, most preferably, it can be stably integrated into the genome of the packaging cell such that it is expressed constitutively. Methods to modify common mammalian promoters to incorporate transcription repressor sites are known in the art. For example, Chang and Roninson modified the strong, constitutive CMV and RSV promoters to comprise operators for the Lac repressor and showed that gene expression from the modified promoters was greatly attenuated in cells expressing the repressor (Chang B D, and Roninson I B (1996) *Gene* 183:137-42). The use of a non-human transcription repressor ensures that transcription of the endonuclease gene will be repressed only in the packaging cells expressing the repressor and not in target cells or tissues transduced with the resulting recombinant AAV vector.

2.6 Engineered Nuclease Variants

[0325] Embodiments of the invention encompass the engineered nucleases, and particularly the recombinant meganucleases, described herein, and variants thereof. Further embodiments of the invention encompass isolated polynucleotides comprising a nucleic acid sequence encoding the recombinant meganucleases described herein, and variants of such polynucleotides.

[0326] As used herein, “variants” is intended to mean substantially similar sequences. A “variant” polypeptide is intended to mean a polypeptide derived from the “native” polypeptide by deletion or addition of one or more amino

acids at one or more internal sites in the native protein and/or substitution of one or more amino acids at one or more sites in the native polypeptide. As used herein, a “native” polynucleotide or polypeptide comprises a parental sequence from which variants are derived. Variant polypeptides encompassed by the embodiments are biologically active. That is, they continue to possess the desired biological activity of the native protein; i.e., the ability to recognize and cleave recognition sequences found in the human T cell receptor alpha constant region (SEQ ID NO:1), including, for example, the TRC 1-2 recognition sequence (SEQ ID NO:3), the TRC 3-4 recognition sequence (SEQ ID NO:4), and the TRC 7-8 recognition sequence (SEQ ID NO:5). Such variants may result, for example, from human manipulation. Biologically active variants of a native polypeptide of the embodiments (e.g., SEQ ID NOs:8-32), or biologically active variants of the recognition half-site binding subunits described herein (e.g., SEQ ID NOs:33-82), will have at least about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99%, sequence identity to the amino acid sequence of the native polypeptide or native subunit, as determined by sequence alignment programs and parameters described elsewhere herein. A biologically active variant of a polypeptide or subunit of the embodiments may differ from that polypeptide or subunit by as few as about 1-40 amino acid residues, as few as about 1-20, as few as about 1-10, as few as about 5, as few as 4, 3, 2, or even 1 amino acid residue.

[0327] The polypeptides of the embodiments may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants can be prepared by mutations in the DNA. Methods for mutagenesis and polynucleotide alterations are well known in the art. See, for example, Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel et al. (1987) *Methods in Enzymol.* 154:367-382; U.S. Pat. No. 4,873,192; Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff et al. (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be optimal.

[0328] A substantial number of amino acid modifications to the DNA recognition domain of the wild-type I-CreI meganuclease have previously been identified (e.g., U.S. Pat. No. 8,021,867) which, singly or in combination, result in recombinant meganucleases with specificities altered at individual bases within the DNA recognition sequence half-site, such that the resulting rationally-designed meganucleases have half-site specificities different from the wild-type enzyme. Table 4 provides potential substitutions that can be made in a recombinant meganuclease monomer or subunit to enhance specificity based on the base present at each half-site position (-1 through -9) of a recognition half-site.

TABLE 4

Posn.	Favored Sense-Strand Base										
	A	C	G	T	A/T	A/C	A/G	C/T	G/T	A/G/T	A/C/C/T
-1	Y75	R70*	K70	Q70*				T46*			G70
	L75*	H75*	E70*	C70							A70
	C75*	R75*	E75*	L70							S70
	Y139*	H46*	E46*	Y75*							G46*
	C46*	K46*	D46*	Q75*							
	A46*	R46*		H75*							
-2	Q70	E70	H70	Q44*	C44*						
	T44*	D70	D44*								
	A44*	K44*	E44*								
	V44*	R44*									
	I44*										
	L44*										
-3	Q68	E68	R68	M68		H68		Y68	K68		
	C24*	F68		C68							
	I24*	K24*		L68							
-4	A26*	E77	R77					S77			S26*
	Q77	K26*	E26*					Q26*			
-5		E42	R42			K28*	C28*				M66
							Q42				K66
-6	Q40	E40	R40	C40	A40						S40
	C28*	R28*		I40	A79						S28*
				V40	A28*						
				C79	H28*						
				I79							
				V79							
-7	N30*	E38	K38	I38			C38				H38
	Q38	K30*	R38	L38							N38
		R30*	E30*								Q30*
-8	F33	E33	F33	L33		R32*	R33				
	Y33	D33	H33	V33							
				I33							
-9		E32	R32	L32				D32			S32
			K32	V32				I32			N32
				A32							H32
				C32							Q32
											T32

[0329] For polynucleotides, a “variant” comprises a deletion and/or addition of one or more nucleotides at one or more sites within the native polynucleotide. One of skill in the art will recognize that variants of the nucleic acids of the embodiments will be constructed such that the open reading frame is maintained. For polynucleotides, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the polypeptides of the embodiments. Variant polynucleotides include synthetically derived polynucleotides, such as those generated, for example, by using site-directed mutagenesis but which still encode a recombinant meganuclease of the embodiments. Generally, variants of a particular polynucleotide of the embodiments will have at least about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or more sequence identity to that particular polynucleotide as determined by sequence alignment programs and parameter described elsewhere herein. Variants of a particular polynucleotide of the embodiments (i.e., the reference poly-

nucleotide) can also be evaluated by comparison of the percent sequence identity between the polypeptide encoded by a variant polynucleotide and the polypeptide encoded by the reference polynucleotide.

[0330] The deletions, insertions, and substitutions of the protein sequences encompassed herein are not expected to produce radical changes in the characteristics of the polypeptide. However, when it is difficult to predict the exact effect of the substitution, deletion, or insertion in advance of doing so, one skilled in the art will appreciate that the effect will be evaluated by screening the polypeptide for its ability to preferentially recognize and cleave recognition sequences found within the human T cell receptor alpha constant region gene (SEQ ID NO:1).

EXAMPLES

[0331] This invention is further illustrated by the following examples, which should not be construed as limiting. Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and proce-

dures described herein. Such equivalents are intended to be encompassed in the scope of the claims that follow the examples below.

Example 1

Characterization of Meganucleases that Recognize and Cleave TRC Recognition Sequences

[0332] 1. Meganucleases that Recognize and Cleave the TRC 1-2 Recognition Sequence

[0333] Recombinant meganucleases (SEQ ID NOs:8-27), collectively referred to herein as “TRC 1-2 meganucleases,” were engineered to recognize and cleave the TRC 1-2 recognition sequence (SEQ ID NO:3), which is present in the human T cell receptor alpha constant region. Each TRC 1-2 recombinant meganuclease comprises an N-terminal nuclease-localization signal derived from SV40, a first meganuclease subunit, a linker sequence, and a second meganuclease subunit. A first subunit in each TRC 1-2 meganuclease binds to the TRC recognition half-site of SEQ ID NO:3, while a second subunit binds to the TRC2 recognition half-site (see, FIG. 1A).

[0334] As illustrated in FIGS. 2 and 3, TRC1-binding subunits and TRC2-binding subunits each comprise a 56 base pair hypervariable region, referred to as HVR1 and HVR2, respectively. TRC1-binding subunits are identical outside of the HVR1 region except at position 80 or position 271 (comprising a Q or E residue), and are highly conserved within the HVR1 region. Similarly, TRC2-binding subunits are also identical outside of the HVR2 region except at position 80 or position 271 (comprising a Q or E residue), and at position 139 of meganucleases TRC 1-2x.87 EE. TRC 1-2x.87 QE. TRC 1-2x.87 EQ. TRC 1-2x.87, and TRC 1-2x.163 which comprise an R residue (shaded grey and underlined). Like the HVR1 region, the HVR2 region is also highly conserved.

[0335] The TRC1-binding regions of SEQ ID NOs:8-27 are illustrated in FIG. 2 and are provided as SEQ ID NOs:33-52, respectively. Each of SEQ ID NOs:33-52 share at least 90% sequence identity to SEQ ID NO:33, which is the TRC1-binding region of the meganuclease TRC 1-2x.87 EE (SEQ ID NO:8). TRC2-binding regions of SEQ ID NOs:8-27 are illustrated in FIG. 3 and are provided as SEQ ID NOs:58-77, respectively. Each of SEQ ID NOs:58-77 share at least 90% sequence identity to SEQ ID NO:58, which is the TRC2-binding region of the meganuclease TRC 1-2x.87 EE (SEQ ID NO:8).

2. Meganucleases that Recognize and Cleave the TRC 3-4 Recognition Sequence

[0336] Recombinant meganucleases (SEQ ID NOs:28 and 29), collectively referred to herein as “TRC 3-4 meganucleases,” were engineered to recognize and cleave the TRC 3-4 recognition sequence (SEQ ID NO:4), which is present in the human T cell receptor alpha constant region. Each TRC 3-4 recombinant meganuclease comprises an N-terminal nuclease-localization signal derived from SV40, a first meganuclease subunit, a linker sequence, and a second meganuclease subunit. A first subunit in each TRC 3-4 meganuclease binds to the TRC3 recognition half-site of SEQ ID NO:4, while a second subunit binds to the TRC4 recognition half-site (see, FIG. 1A).

[0337] As illustrated in FIGS. 4 and 5, TRC3-binding subunits and TRC4-binding subunits each comprise a 56 base pair hypervariable region, referred to as HVR1 and

HVR2, respectively. TRC3-binding subunits are identical outside of the HVR1 region except at position 80 or position 271 (comprising a Q or E residue), and are highly conserved within the HVR1 region. Similarly, TRC4-binding subunits are also identical outside of the HVR2 region except at position 80 or position 271 (comprising a Q or E residue), and are highly conserved within the HVR2 region.

[0338] The TRC3-binding regions of SEQ ID NOs:28 and 29 are illustrated in FIG. 4 and are provided as SEQ ID NOs:53 and 54, respectively. SEQ ID NOs:53 and 54 share 96.6% sequence identity. TRC4-binding regions of SEQ ID NOs:28 and 29 are illustrated in FIG. 5 and are provided as SEQ ID NOs:78 and 79, respectively. SEQ ID NOs:78 and 79 also share 96.6% sequence identity.

3. Meganucleases that Recognize and Cleave the TRC 7-8 Recognition Sequence

[0339] Recombinant meganucleases (SEQ ID NOs:30-32), collectively referred to herein as “TRC 7-8 meganucleases,” were engineered to recognize and cleave the TRC 7-8 recognition sequence (SEQ ID NO:5), which is present in the human T cell receptor alpha constant region. Each TRC 7-8 recombinant meganuclease comprises an N-terminal nuclease-localization signal derived from SV40, a first meganuclease subunit, a linker sequence, and a second meganuclease subunit. A first subunit in each TRC 7-8 meganuclease binds to the TRC7 recognition half-site of SEQ ID NO:5, while a second subunit binds to the TRC8 recognition half-site (see, FIG. 1A).

[0340] As illustrated in FIGS. 6 and 7, TRC7-binding subunits and TRC8-binding subunits each comprise a 56 base pair hypervariable region, referred to as HVR1 and HVR2, respectively. TRC7-binding subunits are identical outside of the HVR1 region except at position 80 or position 271 (comprising a Q or E residue), and are highly conserved within the HVR1 region. Similarly, TRC8-binding subunits are also identical outside of the HVR2 region except at position 80 or position 271 (comprising a Q or E residue), and are highly conserved within the HVR2 region.

[0341] The TRC7-binding regions of SEQ ID NOs:30-32 are illustrated in FIG. 6 and are provided as SEQ ID NOs:55-57, respectively. Each of SEQ ID NOs:55-57 share at least 90% sequence identity to SEQ ID NO:55, which is the TRC7-binding region of the meganuclease TRC 7-8x.7 (SEQ ID NO:30). TRC8-binding regions of SEQ ID NOs:30-32 are illustrated in FIG. 7 and are provided as SEQ ID NOs:80-82, respectively. Each of SEQ ID NOs:80-82 share at least 90% sequence identity to SEQ ID NO:80, which is the TRC8-binding region of the meganuclease TRC 7-8x.7 (SEQ ID NO:30).

4. Cleavage of Human T Cell Receptor Alpha Constant Region Recognition Sequences in a CHO Cell Reporter Assay

[0342] To determine whether TRC 1-2, TRC 3-4, and TRC 7-8 meganucleases could recognize and cleave their respective recognition sequences (SEQ ID NOs:3, 4, and 5, respectively), each recombinant meganuclease was evaluated using the CHO cell reporter assay previously described (see, WO/2012/167192 and FIG. 8). To perform the assays, CHO cell reporter lines were produced which carried a non-functional Green Fluorescent Protein (GFP) gene expression cassette integrated into the genome of the cells. The GFP gene in each cell line was interrupted by a pair of recognition sequences such that intracellular cleavage of either recog-

niton sequence by a meganuclease would stimulate a homologous recombination event resulting in a functional GFP gene.

[0343] In CHO reporter cell lines developed for this study, one recognition sequence inserted into the GFP gene was the TRC 1-2 recognition sequence (SEQ ID NO:3), the TRC 3-4 recognition sequence (SEQ ID NO:4), or the TRC 7-8 recognition sequence (SEQ ID NO:5). The second recognition sequence inserted into the GFP gene was a CHO-23/24 recognition sequence, which is recognized and cleaved by a control meganuclease called “CHO-23/24”. CHO reporter cells comprising the TRC 1-2 recognition sequence and the CHO-23/24 recognition sequence are referred to herein as “TRC 1-2 cells.” CHO reporter cells comprising the TRC 3-4 recognition sequence and the CHO-23/24 recognition sequence are referred to herein as “TRC 3-4 cells.” CHO reporter cells comprising the TRC 7-8 recognition sequence and the CHO-23/24 recognition sequence are referred to herein as “TRC 7-8 cells.”

[0344] CHO reporter cells were transfected with plasmid DNA encoding their corresponding recombinant meganucleases (e.g., TRC 1-2 cells were transfected with plasmid DNA encoding TRC 1-2 meganucleases) or encoding the CHO-23134 meganuclease. In each assay, 4×10^5 CHO reporter cells were transfected with 50 ng of plasmid DNA in a 96-well plate using Lipofectamine 2000 (ThermoFisher) according to the manufacturer’s instructions. At 48 hours post-transfection, cells were evaluated by flow cytometry to determine the percentage of GFP-positive cells compared to an untransfected negative control (TRC 1-2bs). As shown in FIG. 9, all TRC 1-2, TRC 3-4, and TRC 7-8 meganucleases were found to produce GFP-positive cells in cell lines comprising their corresponding recognition sequence at frequencies significantly exceeding the negative control.

[0345] The efficacy of the TRC 1-2x.87 QE, TRC 1-2x.87 EQ, and TRC 1-2x.87 EE meganucleases was also determined in a time-dependent manner. In this study, TRC 1-2 cells (1×10^6) were electroporated with 1×10^6 copies of meganuclease mRNA per cell using a BioRad Gene Pulser Xcell according to the manufacturer’s instructions. At 1, 4, 6, 8, and 12 days post-transfection, cells were evaluated by flow cytometry to determine the percentage of GFP-positive cells. As shown in FIG. 10, each TRC 1-2 meganuclease exhibited high efficiency at 2 days post-transfection, with greater than 50% GFP-positive cells observed. This effect persisted over the 12 day period, with no evidence of cell toxicity observed.

5. Conclusions

[0346] These studies demonstrated that TRC 1-2 meganucleases, TRC 3-4 meganucleases, and TRC 7-8 meganucleases encompassed by the invention can efficiently target and cleave their respective recognition sequences in cells.

Example 2

Cleavage of TRC Recognition Sequences in T Cells and Suppression of Cell-Surface T Cell Receptor Expression

1. Cleavage of the TRC 1-2 Recognition Sequence in Jurkat Cells

[0347] This study demonstrated that TRC 1-2 meganucleases encompassed by the invention could cleave the TRC

1-2 recognition sequence in Jurkat cells (an immortalized human T lymphocyte cell line), 1×10^6 Jurkat cells were electroporated with 8×10^6 copies of a given TRC 1-2 meganuclease mRNA per cell using a BioRad Gene Pulser Xcell according to the manufacturer’s instructions. At 72 hours post-transfection, genomic DNA (gDNA) was harvested from cells and a T7 endonuclease I (T7E) assay was performed to estimate genetic modification at the endogenous TRC 1-2 recognition sequence (FIG. 11). In the T7E assay, the TRC 1-2 locus is amplified by PCR using primers that flank the TRC 1-2 recognition sequence. If there are indels (random insertions or deletions) within the TRC 1-2 locus, the resulting PCR product will consist of a mix of wild-type alleles and mutant alleles. The PCR product is denatured and allowed to slowly reanneal. Slow reannealing allows for the formation of heteroduplexes consisting of wild-type and mutant alleles, resulting in mismatched bases and/or bulges. The T7E1 enzyme cleaves at mismatch sites, resulting in cleavage products that can be visualized by gel electrophoresis. FIG. 11 clearly demonstrates that thirteen different versions of the TRC 1-2 meganucleases generated positive results in the T7E1 assay, indicating effective generation of indels at the endogenous TRC 1-2 recognition sequence.

[0348] To further examine the cleavage properties of TRC 1-2 meganucleases, a dose-response experiment was performed in Jurkat cells, 1×10^6 Jurkat cells were electroporated with either 3 μ g or 1 μ g of a given TRC 1-2 meganuclease mRNA per cell using a BioRad Gene Pulser Xcell according to the manufacturer’s instructions. At 96-hours post-transfection, gDNA was harvested and the T7E1 assay was performed as described above. As seen in FIG. 12, fifteen different TRC 1-2 meganucleases showed cleavage at the endogenous TRC 1-2 recognition site, including three different versions of the TRC 1-2x.87 meganuclease. TRC 1-2x.87 EE worked especially well, generating a strong signal in the T7E1 assay with little to no toxicity in Jurkat cells.

2. Cleavage of TRC 1-2 Recognition Sequence in Human T Cells

[0349] This study demonstrated that TRC 1-2 meganucleases encompassed by the invention could cleave the TRC 1-2 recognition sequence in human T cells obtained from a donor. CD3+ T cells were stimulated with anti-CD3 and anti-CD28 antibodies for 3 days, then electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease using the Amaxa 4D-Nucleofector (Lonza) according to the manufacturer’s instructions. At 3 days and 7 days post-transfection, gDNA was harvested and the T7E assay was performed as described above. FIG. 13A demonstrates that TRC 1-2x.87 EE effectively introduced mutations in the endogenous TRC 1-2 recognition sequence in human T cells, indicating that the meganuclease recognized and cleaved the TRC 1-2 recognition sequence. The intensity of cleavage products does not appear to change between day 3 and day 7 post-transfection, suggesting little or no toxicity due to the TRC 1-2x.87 EE meganuclease. To determine whether the mutations at the endogenous TRC 1-2 recognition sequence were sufficient to eliminate surface expression of the T cell receptor, cells were analyzed by flow cytometry using an anti-CD3 antibody. FIG. 13B shows that approximately 50% of transfected T cells stained negative for CD3, indicating knockout of the T cell receptor. The CD3 negative population did not change significantly between day 3 and day 7

post-transfection, further indicating little or no toxicity associated with the TRC 1-2x.87 EE meganuclease, or the loss of T cell receptor expression.

[0350] To verify that loss of CD3 expression was due to mutations in the TRC 1-2 recognition site, gDNA was harvested from transfected T cells and the TRC 1-2 recognition site locus was amplified by PCR. PCR products were cloned into the pCR-blunt vector using the Zero Blunt PCR cloning kit (Thermo Fisher) according to the manufacturer's instructions. Individual colonies were picked and mini-prepped plasmids were sequenced. FIG. 14 shows sequences of several representative deletions that were observed at the TRC 1-2 recognition sequence. The observed sequences are typical of deletions resulting from the non-homologous end joining repair of DNA double-strand breaks generated by endonucleases.

[0351] In addition to TRC 1-2x.87 EE, other TRC 1-2 meganucleases were able to knockout the T cell receptor in human T cells, including TRC 1-2x.55, and TRC 1-2x.72, albeit to a lesser extent than knockout previously observed for TRC 1-2x.87 EE (Tables 5 and 6). TRC 1-2x.72 Q47E carries a mutation in the active site of the meganuclease (amino acid 47) and serves as a negative control.

TABLE 5

Meganuclease	% CD3 ⁺ Cells	
	Day 3	Day 6
TRC 1-2x.72 Q47E	0.38	1.1
TRC 1-2x.55	3.11	10.84

TABLE 6

Meganuclease	% CD3 ⁺ Cells	
	Day 3	Day 5
TRC 1-2x.72 Q47E	0.29	0.4
TRC 1-2x.72	2.09	4.19

3. Conclusions

[0352] These studies demonstrated that TRC 1-2 meganucleases encompassed by the invention can recognize and cleave the TRC 1-2 recognition sequence in both Jurkat cells (an immortalized T lymphocyte cell line) and in T cells obtained from a human donor. Further, these studies demonstrated that NHEJ occurs at the meganuclease cleavage site, as evidenced by the appearance of indels. Moreover, TRC 1-2 meganucleases were shown to reduce cell-surface expression of the T cell receptor on human T cells obtained from a donor.

Example 3

Recombinant AAV Vectors for Introducing Exogenous Nucleic Acids into Human T Cells

1. Recombinant AAV Vectors

[0353] In the present study, two recombinant AAV vectors (referred to as AAV405 and AAV406) were designed to introduce an exogenous nucleic acid sequence, comprising an EagI restriction site, into the genome of human T cells at

the TRC 1-2 recognition sequence via homologous recombination. Each recombinant AAV vector was prepared using a triple-transfection protocol, wherein a cell line is transfected with a first plasmid encoding "helper" components (e.g., adenoviral) necessary to support replication, a second plasmid comprising the cap and rep genes, and a third plasmid comprising the viral inverted terminal repeats (ITRs) containing the intervening DNA sequence to be packaged into the virus (e.g., the exogenous nucleic acid sequence) (see, Cots D. Bosch A. Chillon M (2013) *Curr. Gene Ther.* 13(5): 370-81). FIG. 15 illustrates the general approach for using recombinant AAV vectors to introduce an exogenous nucleic acid sequence into the cell genome at the nuclease cleavage site.

[0354] AAV405 was prepared using the plasmid illustrated in FIG. 16 (SEQ ID NO:107). As shown, the AAV405 plasmid generally comprises sequences for a 5' ITR, a CMV enhancer and promoter sequence, a 5' homology arm, a nucleic acid sequence comprising the EagI restriction site, an SV40 poly(A) signal sequence, a 3' homology arm, and a 3' ITR. AAV406 was prepared using the plasmid illustrated in FIG. 17 (SEQ ID NO:108). As shown, the AAV406 plasmid comprises similar sequences to those of AAV405, but lacks the CMV enhancer and promoter sequences upstream of the 5' homology arm. The present AAV studies further included the use of an AAV vector encoding GFP (GFP-AAV), which was incorporated as a positive control for AAV transduction efficiency.

2. Introducing Exogenous Nucleic Acid Sequences into the TRC 1-2 Recognition Sequence

[0355] To test whether AAV templates would be suitable for homology directed repair (HDR) following generation of a double-strand break with TRC 1-2 meganucleases, a series of experiments were performed using human T cells. In the first experiment, the timing of electroporation with TRC 1-2 RNA and transduction with recombinant AAV vectors was determined. Human CD3⁺ T cells were stimulated with anti-CD3 and anti-CD28 antibodies for 3 days, then electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease (1 µg) using the Amaxa 4D-Nucleofector (Lonza) according to the manufacturer's instructions. At either 2, 4, or 8 hours post-transfection, cells were transduced with GFP-AAV (1e⁵ viral genomes per cell). Cells were analyzed by flow cytometry for GFP expression at 72 hours post-transduction. As shown in FIG. 18, the highest transduction efficiency was observed when cells were transduced at 2 hours post-transfection (88% GFP-positive cells). Transduction efficiency decreased significantly as the time between transfection and transduction increased, with 78% GFP-positive cells at 4 hours and 65% GFP-positive cells at 8 hours.

[0356] Having determined that efficient viral transduction occurred when cells were transduced 2 hours post-transfection, the AAV405 and AAV406 vectors were used as HDR templates in human T cells. CD3⁺ T cells were stimulated and transfected with 1 µg TRC 1-2x.87 EE mRNA as described above. At 2 hours post-transfection, cells were either transduced with AAV405 or AAV406 (1e⁵ viral genomes per cell). As transduction-only controls, cells were mock transfected (with water) and transduced with either AAV405 or AAV406 (1e⁵ viral genomes per cell). For a meganuclease-only control, cells were transfected with TRC 1-2x.87 EE and then mock transduced (with water) at 2 hours post-transfection.

[0357] To determine whether the AAV vectors served as HDR templates, gDNA was harvested from cells and the TRC 1-2 locus was amplified by PCR using primers that recognized sequences beyond the region of homology in the AAV vectors. PCR primers outside of the homology regions only allowed for amplification of the T cell genome, not from the AAV vectors. PCR products were purified and digested with *EagI*. FIG. 19 shows cleavage of the PCR products amplified from cells that were transfected with TRC 1-2x.87 EE and transduced with either AAV vector (see arrows), indicating insertion of the *EagI* site into the TRC 1-2 recognition sequence. The PCR products from all of the control cell populations are not cleaved by *EagI*, demonstrating that the insertion of the *EagI* site requires creation of a DNA double-strand break by a TRC 1-2 meganuclease.

[0358] To further define the insertion of the *EagI* site into human T cells, individual products from the bulk PCR product were examined. Undigested PCR product generated from the above experiment was cloned into the pCR-blunt vector using the Zero Blunt PCR cloning kit (Thermo Fisher) according to the manufacturer's instructions. Colony PCR was performed using M13 forward and reverse primers (pCR blunt contains M13 forward and reverse priming sites flanking the insert) and a portion of PCR products from cells transfected with TRC 1-2x.87 EE and either AAV405 or AAV406 were analyzed by gel electrophoresis (FIGS. 20A and 21A, respectively). In both cases, there are a mix of full-length PCR products (approximately 1600 bp), smaller inserts, and some empty plasmids (approximately 300 bp). In this assay, bands smaller than full-length but larger than empty plasmids are often times sequences containing large deletions within the TRC 1-2 recognition sequence. In parallel, another portion of PCR products were digested with *EagI* to determine the percent of clones that contain the *EagI* recognition site inserted into the TRC 1-2 recognition sequence. FIGS. 20B and 21B show that several PCR products were cleaved with *EagI* (e.g., FIG. 20B, second row, 6 lanes from the left), generating the expected fragments of approximately 700 and 800 bp. These gels allow for the estimation of *EagI* insertion to be approximately 25% and 6% for AAV405 and AAV406, respectively (adjusted for empty vectors).

[0359] To confirm observations from gel electrophoresis of uncut PCR products and digest with *EagI*, the remaining portion of each PCR product was sequenced. FIG. 22A shows sequences of several representative deletions and insertions that were observed at the TRC 1-2 recognition sequence. These sequences are typical of sequences resulting from the non-homologous end joining repair of DNA double-strand breaks generated by endonucleases. All PCR products that were cleaved with *EagI* contained an *EagI* site inserted into the TRC 1-2 recognition sequence (FIG. 22B).

3. Enhanced AAV Transduction Efficiency

[0360] In light of the observation that AAV transduction was more efficient when it was carried out 2 hours post-transfection than when it was carried out later, an experiment was performed to optimize the timing of transfection and transduction. Human CD3+ T cells were stimulated with anti-CD3 and anti-CD28 antibodies for 3 days, then electroporated with the TRC 1-2x.87 EE meganuclease (1 μ g) using the Amaxa 4D-Nucleofector (Lonza) according to the manufacturer's instructions. Immediately after transfection or 2 hours post-transfection, cells were transduced with

GFP-AAV ($1e^5$ viral genomes per cell). Additionally, non-stimulated cells were transduced with GFP-AAV ($1e^5$ viral genomes per cell). At 72 hours post-transduction, cells were analyzed by flow cytometry for GFP expression. FIG. 23 shows that GFP-AAV transduction performed 2 hour post-transfection resulted in 90% GFP-positive cells, but that transduction immediately after transfection resulted in 98% GFP-positive cells. Resting T cells appeared refractive to AAV transduction, with approximately 0% GFP-positive cells. Non-transduced cells also showed approximately 0% GFP-positive cells.

4. Summary

[0361] These studies demonstrate that AAV vectors can be used in conjunction with recombinant meganucleases to incorporate an exogenous nucleic acid sequence into a cleavage site in the TCR alpha constant region via homologous recombination.

Example 4

Recombinant AAV Vectors for Introducing Exogenous Nucleic Acids Encoding a Chimeric Antigen Receptor in Human T Cells

1. Recombinant AAV Vectors

[0362] In the present study, two recombinant AAV vectors (referred to as AAV-CAR 100 and AAV-CAR763) were designed to introduce an exogenous nucleic acid sequence, encoding a chimeric antigen receptor, into the genome of human T cells at the TRC 1-2 recognition sequence via homologous recombination. Each recombinant AAV vector was prepared using the triple-transfection protocol described previously.

[0363] AAV-CAR100 (also referred to herein as AAV408) was prepared using the plasmid illustrated in FIG. 24 (SEQ ID NO:109). As shown, the AAV-CAR100 (AAV408) is designed for producing a self-complementary AAV vector, and generally comprises sequences for a 5' ITR, a 5' homology arm, a nucleic acid sequence encoding an anti-CD19 chimeric antigen receptor, an SV40 poly(A) signal sequence, a 3' homology arm, and a 3' ITR. AAV-CAR763 (also referred to herein as AAV412) was prepared using the plasmid illustrated in FIG. 25 (SEQ ID NO:110). As shown, the AAV-CAR763 (AAV412) plasmid generally comprises the same sequences as AAV-CAR100 (AAV408), but is designed for producing a single-stranded AAV vector. Because a single-stranded AAV vector can accommodate a larger payload, the 5' homology arm and the 3' homology arm are longer in AAV-CAR763 (AAV412) than in AAV-CAR100 (AAV408). The present AAV studies will further include the use of an AAV vector encoding GFP (GFP-AAV), which will be incorporated as a positive control for AAV transduction efficiency.

2. Introducing a Chimeric Antigen Receptor Sequence into the TRC 1-2 Recognition Sequence

[0364] Studies will be conducted to determine the efficiency of using recombinant AAV vectors to insert a chimeric antigen receptor sequence into the TCR alpha constant region gene while, simultaneously, knocking out cell-surface expression of the endogenous TCR receptor.

[0365] To confirm transduction efficiency, human CD3+ T cells will be obtained and stimulated with anti-CD3 and anti-CD28 antibodies for 3 days, then electroporated with

mRNA encoding the TRC 1-2x.87 EE meganuclease (1 µg) using the Amaxa 4D-Nucleofector (Lonza) according to the manufacturer's instructions. Cells will be transduced with GFP-AAV (1e⁵ viral genomes per cell) immediately after transfection as described above. Cells will be analyzed by flow cytometry for GFP expression at 72 hours post-transduction to determine transduction efficiency.

[0366] AAV-CAR100 (AAV408) and AAV-CAR763 (AAV412) vectors will then be used as HDR templates in human T cells for the insertion of the anti-CD19 chimeric antigen receptor sequence. Human CD3+ T cells will be stimulated and transfected with 1 µg TRC 1-2x.87 EE mRNA as described above. Cells will then be transduced with AAV-CAR100 (AAV408) or AAV-CAR763 (AAV412) (1e⁵ viral genomes per cell) either immediately after transfection or within 0-8 hours of transfection. As transduction-only controls, cells will be mock transfected (with water) and transduced with either AAV-CAR100 (AAV408) or AAV-CAR763 (AAV412) (1e⁵ viral genomes per cell). For a meganuclease-only control, cells will be transfected with mRNA encoding TRC 1-2x.87 EE and then mock transduced (with water) immediately post-transfection.

[0367] Insertion of the chimeric antigen receptor sequence will be confirmed by sequencing of the cleavage site in the TCR alpha constant region gene. Cell-surface expression of the chimeric antigen receptor will be confirmed by flow cytometry, using an anti-Fab or anti-CD19 antibody. Knock-out of the endogenous T cell receptor at the cell surface will be determined by flow cytometry as previously described.

Example 5

Insertion and Expression of Chimeric Antigen Receptor

[0368] 1. Insertion of Chimeric Antigen Receptor Sequence into the TRC 1-2 Recognition Sequence

[0369] In the present study, we test whether AAV can provide HDR templates that can be used to insert a chimeric antigen receptor sequence into the TCR alpha constant region gene and, simultaneously, knock out cell-surface expression of the endogenous TCR receptor. In the first experiment, human CD3+ T cells (1e⁶ cells) were stimulated and electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease (2 µg) as described above, then immediately transduced with AAV412 (1e⁵ viral genomes/cell). As controls, cells were mock electroporated, then transduced with AAV412 or electroporated with mRNA encoding TRC 1-2x.87EE, then mock transduced. An additional control of mock electroporated, mock transduced cells was included.

[0370] A PCR-based assay was developed to determine whether the AAV HDR template was utilized to repair double-strand breaks at the TRC 1-2 recognition sequence. Three sets of primer pairs were used for PCR analysis. The first set was designed to amplify a region within the homology arms of AAV412. Since this first primer set (referred to as "Inside homolog arms/CAR region" in Table 7) lies within the homology region, it will either amplify the unmodified TRC 1-2 recognition sequence locus of the genome (349 bp), the AAV412 vector input (2603 bp), or the TRC 1-2 recognition sequence into which the CAR gene has been inserted (2603 bp). The second primer set (referred to as "Outside 5' homology arm" in Table 7) includes one primer that anneals within the CAR region of the AAV412 HDR template, one primer that anneals in the human genome,

outside of the 5' homology arm of the AAV412 HDR template and will amplify an 1872 bp fragment only if the CAR gene was successfully inserted into the TRC 1-2 recognition sequence. The third primer set (referred to as "Outside 3' homology arm" in Table 7) includes one primer that anneals within the CAR region of the AAV412 HDR template, and one primer that anneals in the human genome, outside of the 3' homology arm of the AAV412 HDR template. Similarly to the second primer set, the third primer set will amplify an 1107 bp fragment only if the CAR gene was successfully inserted into the TRC 1-2 recognition sequence. Taken together, PCR products from all three primer sets will indicate whether the CAR sequence is present in cells (primer set 1), and whether it has been inserted into the TRC 1-2 recognition sequence (primer sets 2 and 3).

[0371] On day 4 post-transduction cells were analyzed using the PCR primer pairs described above. Briefly, approximately 3,000 cells were harvested, pelleted, and lysed and PCR was performed to determine whether the CAR gene was inserted into the TRC 1-2 recognition sequence. PCR products were resolved on an agarose gel, shown in FIG. 26 (lane descriptions can be found in Table 7). Lanes 1-3 are PCR products from the sample that was electroporated with mRNA encoding TRC 1-2x.87EE and mock transduced.

[0372] As expected, the first primer pair ("Inside homolog arms/CAR region") amplified the unmodified TRC 1-2 recognition sequence locus, generating a 349 bp band shown in lane 1. Lanes 2 and 3 correspond to primer pairs that only generate a product if the CAR gene has been inserted into the TRC 1-2 recognition sequence, and do not show products. Lanes 7-9 represent samples that were mock electroporated and mock transduced and show the same bands as the TRC 1-2x.87EE mRNA only control described above. Lanes 4-6 show PCR products from the sample that was electroporated with TRC 1-2x.87EE mRNA and transduced with AAV412. Lane 4 shows two bands generated by the first primer pair ("Inside homolog arms/CAR region"), indicating amplification of the unmodified TRC 1-2 recognition sequence locus of the genome (349 bp) and the AAV412 vector input (2603 bp) or the TRC 1-2 recognition sequence into which the CAR gene has been inserted (2603 bp). Lanes 5 and 6 show products generated by the primer pairs that only amplify products if the CAR nucleic acid sequence has been inserted into the TRC 1-2x.87EE recognition site. Both bands are the predicted size (1872 and 1107 bp, respectively). Lanes 10-12 represent the sample that was mock electroporated and transduced with AAV412. Lane 10 shows two bands generated by the first primer pair ("Inside homolog arms/CAR region"), indicating amplification of the unmodified TRC 1-2 recognition sequence locus of the genome (349 bp) and the AAV412 vector input (2603 bp). Lanes 11 and 12 correspond to primer pairs that only generate a product if the CAR gene has been inserted into the TRC 1-2 recognition sequence, and do not show products. The absence of bands in lanes 11 and 12 (which include primers outside of the homology arm) indicates that the 2603 bp band in lane 10 was generated from amplification of the AAV412 input.

[0373] Taken together, the PCR analysis clearly demonstrates that CAR genes are introduced into the TRC 1-2x.87EE recognition site when both TRC 1-2x.87EE mRNA and AAV412 are present in cells. Thus, we conclude that

AAV412 serves to produce suitable HDR templates that can be used to insert a CAR gene into the TRC 1-2x.87EE recognition sequence.

[0374] In a second experiment, human CD3+ T cells were stimulated and electroporated with mRNA encoding the TRC 1-2x.87 EE meganuclease as described above, then immediately transduced with increasing amounts of AAV408 (0 μ L, 3.125 μ L, 6.25 μ L, 12.5 μ L, or 25 μ L, which corresponds to approximately 0, 3.125e³, 6.250e³, 1.25e⁴ and 2.5e⁴ viral genomes/cell). As controls, cells were mock electroporated, then transduced with increasing amounts of AAV408. Additional controls included cells that were mock electroporated and mock transduced, as well as cells that were electroporated with TRC 1-2x.87EE mRNA then mock transduced. On day 4 post-transduction, cells were harvested and analyzed as described above, but only using the primer pairs that amplified a product only if the CAR gene has been inserted into the TRC 1-2 recognition sequence. PCR products were resolved on agarose gels, shown in FIG. 27. FIG. 27A shows the PCR products generated using the primer pair described above (“Outside 5’ homology arm”) which only amplifies a product on the 5’ end of the TRC 1-2 recognition sequence locus if the CAR gene has been inserted into that locus. FIG. 27B shows the PCR products generated using the primer pair described above (“Outside 3’ homology arm”) which only amplifies a product on the 3’ end of the TRC 1-2 recognition sequence locus if the CAR gene has been inserted into that locus. Lane descriptions can be found in Table 8. Lanes 1-5 in both FIGS. 27A and 27B represent the samples that were either mock electroporated or mock electroporated then mock transduced. No PCR products are visible in mock electroporated cells, indicating the HDR templates produced by AAV408 are unable to insert the CAR gene into the TRC 1-2 recognition sequence in the absence of TRC 1-2x.87EE mRNA. Lane 6 represents the sample that was electroporated with TRC 1-2x.87EE mRNA and mock transduced. No PCR products are visible, indicating that the CAR gene had not been inserted into the TRC 1-2 recognition sequence. Lanes 7-10 represent samples that were electroporated with TRC 1-2x.87EE mRNA and transduced with increasing amounts of AAV408. The appropriately sized bands for each PCR are evident, indicating that AAV408 can produce HDR donors for repair of the TRC 1-2 recognition sequence, resulting in insertion of the CAR gene.

TABLE 7-continued

Sample	Nucleofection	Virus (100k MOI)	PCR	Product Size
8	Mock (Water)	—	Outside 5’ homology arm	1872 bp
9	Mock (Water)	—	Outside 3’ homology arm	1107 bp
10	Mock (Water)	AAV412	Inside homolog arms/Genomic = 349 bp CAR region +CD19 = 2603 bp	
11	Mock (Water)	AAV412	Outside 5’ homology arm	1872 bp
12	Mock (Water)	AAV412	Outside 3’ homology arm	1107 bp

TABLE 8

Sample	Nucleofection	Virus (AAV408) μ L
1	Mock (Water)	0
2	Mock (Water)	1125
3	Mock (Water)	6.75
4	Mock (Water)	12.5
5	Mock (Water)	25
6	TRC1-2x87EE	0
7	TRC1-2x87EE	3.125
8	TRC1-2x87EE	62.5
9	TRC1-2x87EE	12.5
10	TRC1-2x87EE	25

[0375] The PCR-based assays described above are useful in determining whether the CAR gene had been inserted into the TRC 1-2 recognition sequence, but do not give information on efficiency. To determine the efficiency of CAR insertion, we developed a digital PCR-based assay (schematic shown in FIG. 28A). In this assay, two primer sets are used. The first set amplifies an irrelevant gene sequence and serves a reference sequence to control for template number. The second set consists of one primer that anneals within the CAR gene and one primer that anneals outside of the 3’ homology arm, such that a product is only amplified if the CAR gene has been inserted into the TRC 1-2 recognition sequence. A VIC-labeled probe anneals within the amplicon generated from the first primer set and FAM-labeled probe anneals within the amplicon generated by the second set of primers. By dividing the number of amplicons detected by the FAM-labeled probe to the number of reference sequence amplicons detected by the VIC-labeled probe, it is possible to accurately quantitate the percent of TRC 1-2 recognition sequence loci that were modified by insertion of the CAR gene.

[0376] FIG. 28B shows the results of the digital PCR assay for samples that were either mock electroporated then transduced, electroporated with TRC 1-2x.87EE mRNA then mock transduced, or electroporated with TRC 1-2x.87EE mRNA then transduced with increasing amounts of AAV408. Digital PCR was performed using genomic DNA isolated from cells approximately 1 week post-transduction. Consistent with the observations from the PCR described in FIG. 27, both control samples (transduction only or electroporation only) were found to have 0% CAR gene inserted into the TRC 1-2x.87EE recognition sequence. Samples that were electroporated with mRNA encoding TRC 1-2x.87EE then transduced with increasing amounts of AAV408 were found to have between approximately 1.5% and 7%. The assay was performed on two different instruments (labeled

TABLE 7

Sample	Nucleofection	Virus (100k MOI)	PCR	Product Size
1	TRC1-2x87EE	—	Inside homolog arms/Genomic = 349 bp CAR region +CD19 = 2603 bp	
2	TRC1-2x87EE	—	Outside 5’ homology arm	1872 bp
3	TRC1-2x87EE	—	Outside 3’ homology arm	1107 bp
4	TRC1-2x87EE	AAV412	Inside homolog arms/Genomic = 349 bp CAR region +CD19 = 2603 bp	
5	TRC1-2x87EE	AAV412	Outside 5’ homology arm	1872 bp
6	TRC1-2x87EE	AAV412	Outside 3’ homology arm	1107 bp
7	Mock (Water)	—	Inside homolog arms/Genomic = 349 bp CAR region +CD19 = 2603 bp	

QX200 and QS3D) and showed remarkable agreement, demonstrating the sensitivity and precision of this digital PCR-based assay.

2. Expression of Anti-CD19 Chimeric Antigen Receptor on T Cells

[0377] In addition to determining whether CAR insertion occurred at the molecular level, we sought to determine the expression level of the anti-CD19 chimeric antigen receptor in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV408 as the HDR template. Additionally, we examined the efficiency in which insertion of the CAR into the TRC 1-2x.87EE recognition sequence resulted in knockout of the T cell receptor. Samples described above and analyzed in FIGS. 27 and 28 were also analyzed for CAR and CD3 expression by flow cytometry. Approximately 4 days post-transduction, cells were labeled with antibodies that recognize the anti-CD19 CAR (anti-Fab-Alexa647) or CD3 (CD3-BB515) and analyzed by flow cytometry. FIG. 29A shows flow cytometry plots, with anti-CAR labeling shown on the Y axis and anti-CD3 labeling shown on the X axis. Cells that were mock electroporated and mock transduced (MOI-0) were overwhelmingly CD3⁺/CAR⁻ (the lower right quadrant, 98.7%). Cells that were mock electroporated then transduced with increasing amounts of AAV408 looked essentially identical to the control cells, with the CD3⁺/CAR⁻ populations at 98.8%, 99.99%, and 99.1%. Thus we conclude that the AAV408 virus alone is not driving detectable levels of CAR expression, nor is it capable of disrupting expression of the T cell receptor.

[0378] FIG. 29B shows flow cytometry plots for samples that were either electroporated with mRNA encoding TRC 1-2x.87EE then mock transduced or cells that were electroporated with TRC 1-2x.87EE then transduced with increasing amounts of AAV408. Cells that were electroporated then mock transduced show 47.1% CD3⁻ cells, indicating efficient knockout of the T cell receptor complex. Background labeling with the anti-CD19 CAR was very low, with 0.6% in the CD3⁻ population and 0.78% in the CD3⁺ population. Samples that were electroporated with mRNA encoding TRC 1-2x.87EE then transduced with increasing amounts of AAV408 showed CAR labeling in the CD3⁻ population, ranging from 2.09% to 5.9%. There was also a slight increase in CAR labeling in the CD3⁺ population, ranging from 1.08% to 1.91%. We did not determine the cause of the increase in CAR⁺ cells in the CD3⁺ population, although it is possible that the CAR was inserted into the non-expressed T cell receptor allele (only one allele of the T cell receptor alpha chain is expressed and incorporated into the T cell receptor complex).

[0379] These data correlated well with the quantitative digital PCR-based assay described above. For example, at the highest MOI of AAV408 (2.5e⁴ viral genomes/cell), the digital PCR assay showed approximately 6% CAR insertion, and the flow cytometry assay showed 5.9% CAR⁺/CD3⁻ cells. If one takes into account the CAR⁺/CD3⁺ population, the data are still quite comparable, with the flow cytometry assay showing approximately 7.8% CAR⁺ compared to 6% by digital PCR.

Example 6

Characterization of Additional AAV Vectors

[0380] 1. Insertion of a Chimeric Antigen Receptor Sequence into the TRC 1-2 Recognition Sequence

[0381] Having shown that AAV vectors could provide suitable HDR templates to insert CAR genes into the TRC 1-2x.87EE recognition sequence, we sought to optimize the configuration of the AAV vector. We generated a vector that could be used to produce self-complementary AAV genomes that included the CAR gene expression cassette driven by a JeT promoter, flanked by short regions of homology to the TRC 1-2 recognition sequence locus and AAV ITRs. This vector is referred to as AAV421 (FIG. 30; SEQ ID NO:123). Short homology arms were necessary due to limited packaging capacity of self-complementary AAV. Additionally, we generated a vector that could be used to produce single-strand AAV genomes that includes the CAR gene expression cassette driven by a CMV promoter, flanked by long homology arms and AAV ITRs. This vector is referred to as AAV422 (FIG. 31; SEQ ID NO:124). Since single-strand AAV genomes have a larger cargo capacity, we were able to utilize longer homology arms than in the self-complementary vector.

[0382] To test whether AAV421 and AAV422 were useful to target insertion of the CAR gene into the TRC 1-2 recognition sequence, several experiments similar to those described above were carried out in human CD3⁺ T cells. In a first experiment, human CD3⁻ T cells (1e⁶ cells) were either mock electroporated then transduced with increasing amounts of AAV421 or 422, or electroporated with TRC 1-2x.87EE mRNA (2 µg) then transduced with increasing amounts of AAV421 or AAV422. AAV422 MOIs were significantly higher than AAV421 in this experiment than in the experiments described above (approximate MOIs were 1.25e⁴, 2.5e⁴, 5e⁴ and less viral genomes/cell) because earlier experiments with AAV408 suggested that higher MOIs would result in more efficient CAR insertion. The AAV421 virus stock was not concentrated enough to allow for titers significantly higher than in the experiments described earlier. As controls, cells were electroporated (mock or with TRC 1-2x.87EE mRNA) then mock transduced. As an additional component to this experiment, a “large scale” condition was performed, in which 10e⁶ cells (10 times more than a typical experiment) were electroporated with TRC 1-2x.87EE mRNA then transduced with AAV422 (2.5e⁴ viral genomes/cell). Lastly, we also tested a second virus stock of AAV421 to compare to the primary virus stock.

[0383] Insertion of the CAR was determined by PCR as described above, using primer pairs that only amplify products if the CAR gene has been inserted into the TRC 1-2x.87EE recognition sequence. PCR was resolved by agarose gel, shown in FIGS. 32A and 32B (lane descriptions can be found in Tables 9 and 10). Sample 1 in FIG. 32A was mock electroporated then mock transduced, and samples 2-5 were mock electroporated then transduced with AAV421. The gel shows that none of these samples generated PCR products, indicating that AAV421, in the absence of TRC 1-2x.87EE mRNA, is unable to drive insertion of the CAR gene into the TRC 1-2 recognition sequence. Additionally, the control sample that was electroporated with TRC 1-2x.87EE mRNA then mock transduced (sample 6), did not show any PCR products. Samples 7-10 in FIG. 32A were electroporated with TRC 1-2x.87EE mRNA, then trans-

duced with increasing amounts of AAV421. The gel shows PCR bands for products extending beyond both the 5' and 3' homology arm (the two bands under each sample number), demonstrating integration of the CAR gene into the TRC 1-2 recognition sequence. Lastly in FIG. 32A, lanes 11 and 12 represent samples that were electroporated with TRC 1-2x.87EE mRNA then transduced with AAV422, either starting with $1e^6$ or $10e^6$ cells/sample, respectively. The presence of both PCR bands (larger in the first set, because different primer was used to account for a longer homology arm) indicate successful insertion of the CAR gene into the TRC 1-2 recognition sequence.

[0384] Sample 1 in FIG. 32B was mock electroporated then mock transduced, and samples 2-5 were mock electroporated then transduced with increasing amounts of AAV422 (Table 10). The gel shows that none of these samples generated PCR products, indicating that AAV422, in the absence of TRC 1-2x.87EE mRNA, is unable to drive insertion of the CAR gene into the TRC 1-2 recognition sequence. Samples 7-10 in FIG. 32B were electroporated with TRC 1-2x.87EE mRNA, then transduced with increasing amounts of AAV422. The gel shows PCR bands for products extending beyond both the 5' and 3' homology arm, demonstrating integration of the CAR gene into the TRC 1-2 recognition sequence. Lastly, sample 11 represents the sample that was electroporated with TRC 1-2x.87EE mRNA then transduced with a AAV421 from a different virus stock than samples shown in FIG. 32A. The presence of bands indicate insertion of the CAR gene into the TRC 1-2 recognition sequence and confirms reproducibility between different virus stocks. Taken together, FIG. 32 clearly demonstrates that both AAV421 and AAV422 are capable of generating HDR templates suitable for inserting the CAR gene into the TRC 1-2 recognition sequence.

TABLE 9

Sample	Nucleofection	AAV Virus	μ l AAV	MOI (approximate)
1	Mock (Water)	421	0	0
2	Mock (Water)	421	3.125	3906
3	Mock (Water)	421	6.25	7813
4	Mock (Water)	421	12.5	15625
5	Mock (Water)	421	25	31250
6	TRC1-2x87EE	421	0	0
7	TRC1-2x87EE	421	3.125	3906
8	TRC1-2x87EE	421	6.25	7813
9	TRC1-2x87EE	421	12.5	15625
10	TRC1-2x87EE	421	25	31250
11	TRC1-2x87EE	422	6.25	25000
12	TRC1-2x87EE	422	62.5	25000

Large Scale

TABLE 10

Sample	Nucleofection	AAV Virus	μ l AAV	MOI (approximate)
1	Mock (Water)	422	0	0
2	Mock (Water)	422	3.125	12500

TABLE 10-continued

Sample	Nucleofection	AAV Virus	μ l AAV	MOI (approximate)
3	Mock (Water)	422	6.25	25000
4	Mock (Water)	422	12.5	50000
5	Mock (Water)	422	25	100000
6	TRC1-2x87EE	422	0	0
7	TRC1-2x87EE	422	3.125	12500
8	TRC1-2x87EE	422	6.25	25000
9	TRC1-2x87EE	422	12.5	50000
10	TRC1-2x87EE	422	25	100000
11	TRC1-2x87EE	421B	25	10000

2. Expression of Anti-CD19 Chimeric Antigen Receptor on T Cells Using AAV421

[0385] Here, we sought to determine the expression level of the anti-CD19 chimeric antigen receptor in cells that had the CAR gene inserted into the TRC 1-2 recognition sequence using AAV421. Samples described above and analyzed in FIG. 32A were also analyzed for CAR and CD3 expression by flow cytometry. Approximately 4 days post-transduction, cells were labeled with antibodies that recognize the anti-CD19 CAR or CD3 and analyzed by flow cytometry. FIG. 33A shows flow cytometry plots for cells that were mock electroporated and transduced with AAV421, along with control cells that were mock electroporated and mock transduced. Cells that were mock electroporated and mock transduced (MOI-0) were overwhelmingly $CD3^+/CAR^-$ (the lower right quadrant, 98.8%). Cells that were mock electroporated then transduced with increasing amounts of AAV421 looked essentially identical to the control cells, with the $CD3^+/CAR^-$ populations at 98.8%, 98.6%, 98.8% and 97.9%. Thus, we conclude that the AAV421 virus alone is not driving detectable levels of CAR expression, nor is it capable of disrupting expression of the T cell receptor.

[0386] FIG. 33B shows flow cytometry plots for samples that were either electroporated with TRC 1-2x.87EE mRNA then mock transduced or cells that were electroporated with TRC 1-2x.87EE then transduced with increasing amounts of AAV421. Cells that were electroporated then mock transduced show 56.7% $CD3^-$ cells, indicating efficient knockout of the T cell receptor complex. Background labeling with the anti-CD19 CAR was very low, with 0.48% in the $CD3^-$ population and 0.36% in the $CD3^+$ population. Samples that were electroporated with mRNA encoding TRC 1-2x.87EE then transduced with increasing amounts of AAV412 showed significant amounts of CAR labeling in the $CD3^-$ population, ranging from 4.99% to 13.4%. There was also a slight increase in CAR labeling in the $CD3^-$ population, ranging from 1.27% to 3.95%. As mentioned above, it is possible that the CAR gene was inserted into the non-expressed T cell receptor allele. Also in contrast to experiments with AAV408, the CAR^+ population was much better defined, with a higher mean fluorescence intensity, suggesting that the JeT promoter drives higher expression than the eF1 α core promoter.

[0387] While evaluating insertion of the CAR gene using AAV421 in conjunction with TRC 1-x.87EE, we sought to determine a method that would allow us to preferentially expand and enrich the CD3⁻/CAR⁺ population. From the experiment described above and shown in FIG. 33, we used cells that were electroporated with TRC 1-2x.87EE mRNA (2 µg) then transduced with AAV421 (3.13e⁴ viral genomes/cell). Control samples were mock electroporated and mock transduced, mock electroporated and transduced with AAV421, or electroporated with TRC 1-2x.87EE and mock transduced taken from the experiment described above and shown in FIG. 33. As a control enrichment and expansion process, these cells were incubated for 6 days in complete growth medium supplemented with IL-7 and IL-15 (both at 10 ng/mL). Cells were then labeled with antibodies against the anti-CD19 CAR and CD3 and analyzed by flow cytometry (FIG. 34A). Cells that were mock electroporated and mock transduced showed low levels of background staining in the CD3⁻/CAR⁺ quadrant (0.13%). The CD3⁻/CAR⁺ population was essentially the same in samples that were either mock electroporated then transduced with AAV or electroporated with TRC 1-2x.87EE mRNA then mock transduced (0.16% and 0.55%, respectively). Cells that were electroporated with TRC 1-2x.87EE mRNA and mock transduced had a CD3⁻/CAR⁻ population of 53.2%, very close to the amount stained in the first part of this experiment shown in FIG. 33B (56.7%). Cells that were electroporated with TRC 1-2x.87EE and transduced with AAV showed 12.6% CD3⁻/CAR⁺ cells, almost identical to the original labeling of these cells shown in FIG. 33 (13.4%), demonstrating that mixture of IL-7 and IL-15 is insufficient to enrich or expand the specific CD3⁻/CAR⁺ cell population.

[0388] We next sought to enrich for the CD3⁻/CAR⁺ population in an antigen-specific manner by incubating the 4 samples described above with IM-9 cells, which present CD19 on the cell surface. IM-9 cells were inactivated by pre-treatment with mitomycin C and incubated with samples at a 1:1 ratio for 6 days in the presence of IL-7 and IL-15 (10 ng/mL). Cells were then labeled with antibodies against CD3 and the anti-CD19 CAR and analyzed by flow cytometry (FIG. 34B). Cells that were mock electroporated and mock transduced showed low levels of background staining in the CD3⁻/CAR⁺ quadrant (0.2%). The CD3⁻/CAR⁺ population was the same in samples that were mock electroporated then transduced with AAV (0.2%) and slightly higher in cells that were electroporated with TRC 1-2x.87EE and mock transduced (1.24%). The increase in CD3⁻/CAR⁺ cells the TRC 1-2x.87EE alone control is considered background since no CAR nucleic acid was ever introduced into the system. Cells that were electroporated with TRC 1-2x.87EE mRNA and mock transduced had a CD3⁻/CAR⁻ population of 42.5%, which is significantly lower than they were prior to expansion (56.7%, FIG. 33) suggesting that CD⁺ cells may have a growth advantage in this system. However, cells that were electroporated with TRC 1-2x.87EE and transduced with AAV showed 49.9% CD3⁻/CAR⁺ cells, a dramatic increase compared to the original labeling of these cells shown in FIG. 33 (13.4%), demonstrating that incubation of this sample with IM-9 cells in the presence of IL-7 and IL-15 is quite effective in enriching and expanding the CD3⁻/CAR⁺ population. The CD3⁻/CAR⁺ population was also expanded under these conditions, with the mock electroporated/AAV transduced sample and the TRC 1-2x.87EE

electroporated/AV transduced sample showing 2.53% and 15.3% CD3⁻/CAR⁺, respectively.

[0389] In the cells that were electroporated with TRC 1-2x.87EE then transduced with AAV421, 24.2% of the CD3⁻ population was CAR⁺ prior to expansion (FIG. 33B). After incubation in medium supplemented with IL-7 and IL-15, that 25.3% of the CD3⁻ cells were CAR⁺ (FIG. 34A) indicating that the ratio of gene knock-in to gene-knockout was unchanged. However, after incubation with IM-9 cells in addition to IL-7 and IL-15, over 80% (80.35%, FIG. 34B) of the CD3⁻ cells were CAR⁺, demonstrating that incubation with IM-9 cells resulted in antigen-specific enrichment.

[0390] Since mitomycin C inactivates cells very potently and IM-9 cells were not persisting long in the mixed culture, we reasoned that a second infusion of IM-9 cells might further increase enrichment of CD3⁻/CAR⁺ cells. Some of the cells described above and shown in FIG. 34B would mixed with fresh IM-9 cells (pre-treated with mitomycin C) in medium containing IL-7 and IL-15 and were incubated another 6 days. Cells were then stained for CD3 and anti-CD19 CAR and analyzed by flow cytometry (FIG. 34C). The percentage of CD3⁻/CAR⁺ cells in any of the control samples were essentially unchanged compared to the first round of enrichment on IM-9 cells.

[0391] However, the cells that were electroporated with TRC 1-2x.87EE and transduced with AAV421 showed a significant enrichment of the CD3⁻/CAR⁺ population, increasing from 49.9% (after the first round if incubation with IM-9 cells, FIG. 34B) to 65.7% (FIG. 34C). Importantly, 93.75% of the CD3⁻ population was CAR⁺, indicating further antigen-specific expansion.

3. Expression of Anti-CD19 Chimeric Antigen Receptor on T Cells Using AAV422

[0392] We also examined expression of the anti-CD19 CAR from cells in which AAV422 was used to provide the HDR template (described above, PCR results shown in FIG. 32B). Approximately 4 days post-transduction, cells were labeled with antibodies that recognize the anti-CD19 CAR or CD3 and analyzed by flow cytometry. FIG. 35A shows flow cytometry plots for cells that were mock electroporated and transduced with increasing amounts of AAV422, along with control cells that were mock electroporated and mock transduced. Cells that were mock electroporated and mock transduced (MOI-0) were overwhelmingly CD3⁺/CAR⁻ (the lower right quadrant, 98.8%). Cells that were mock electroporated then transduced with increasing amounts of AAV422 looked essentially identical to the control cells, with the CD3⁺/CAR⁻ populations at 98.6%, 98.6%, 98.9% and 98.4%. Thus, the AAV422 vector alone is not driving detectable levels of CAR expression, nor is it capable of disrupting expression of the T cell receptor.

[0393] FIG. 35B shows flow cytometry plots for samples that were either electroporated with TRC 1-2x.87EE mRNA then mock transduced or cells that were electroporated with TRC 1-2x.87EE then transduced with increasing amounts of AAV422. Cells that were electroporated then mock transduced show 59.3% CD3⁻ cells, indicating efficient knockout of the T cell receptor complex. Background labeling with the anti-CD19 CAR was very low, with 1.47% in the CD3⁻ population and 0.52% in the CD3⁻ population. Samples that were electroporated with mRNA encoding TRC 1-2x.87EE then transduced with increasing amounts of AAV422 showed significant amounts of CAR labeling in the CD3⁻

population, ranging from 14.7% to 20.3%. There was also a slight increase in CAR labeling in the CD3⁺ population, ranging from 2.3% to 2.7%.

[0394] Surprisingly, we observed a noticeable increase in T cell receptor knockout efficiency in the presence of AAV422. Overall CD3 knockout efficiency with increasing AAV422 was 71.6%, 74.9%, 77.8% and 74.4% compared to 59.3% in the TRC 1-2x.87EE electroporation alone. In contrast, overall CD3 knockout efficiency with increasing AAV421 was 56.99%, 56.62%, 57.4% and 55.4% compared to 57.18% in the TRC 1-2x.87EE electroporation alone (FIG. 33B). Thus, it appears that electroporation with TRC 1-2x.87EE in the presence of single-stranded AAV genomes, but not self-complementary AAV genomes, results in an increase in the overall knockout efficiency of the TRC 1-2x.87EE nuclease. Because of this increase, the percent of CD3⁻ cells that are CAR⁺ is not significantly different between cells transduced with AAV421 and AAV422 despite the higher numbers of CD3⁻/CAR⁺ cells. The highest percent of CD3⁻ cells that were CAR⁺ using AAV421 was 24.18% (MOI=3.13e⁴ viral genomes/cell) compared to 26.48% with AAV422 (MOI=1e⁵ viral genomes/cell). This observation is particularly interesting considering the large difference in MOI between AAV421 and AAV422.

[0395] The concept of utilizing IM-9 cells to specifically enrich for CD3⁻/CAR⁺ cells was tested using cells from this experiment. Again, rather than testing the entire panel, we only attempted enrichment of either cells mock electroporated then transduced with AAV422 or electroporated with TRC 1-2x.87EE then transduced with AAV422 (2.5e⁴ viral genomes/cell) in a new experiment. FIG. 36A shows flow cytometry plots at approximately day 4 post-transduction. Mock electroporated/transduced cells showed background staining of CD3⁻/CAR⁺ cells at 0.13%. In comparison, cells electroporated with TRC 1-2x.87EE the transduced with AAV422 showed 4.44% CD3⁻/CAR⁺ cells. Cells were incubated with IM-9 cells (pre-treated with mitomycin) in the presence of IL-7 and IL-15 for 6 days as described above, then analyzed by flow cytometry. FIG. 36B shows that incubation with IM-9 cells dramatically increased the CD3⁻/CAR⁺ population in AAV422 transduced cells to 35.8%. The CAR⁺ cells make up 45.2% of the total CD3⁻ population, compared to 6.69% prior to enrichment (FIG. 36A). As above, we also further enriched by a second addition of IM-9 cells (FIG. 36C). Two rounds of incubation with IM-9 cells resulted in 65.1% CD3⁻/CAR⁺ cells. The CAR⁺ cells make up 78.25% of the total CD3⁻ population, indicating significant, antigen-dependent enrichment of CD3⁻/CAR⁺ cells.

[0396] These data, in conjunction with the data presented above, clearly demonstrate that cells that have had an anti-CD19 CAR gene inserted into the TRC 1-2 recognition sequence can be successfully enriched by incubation with IM-9 cells in the presence of IL-7 and IL-15, and can result in a CD3⁻ population that is over 90% CAR⁺ in as little as 12 days of culture.

4. Increased Knockout Efficiency Observed when Using Single-Strand AAV Vectors

[0397] In the present study, we followed up on the observation that single-stranded AAV vectors increased knockout efficiency of the TRC 1-2x.87EE nuclease. In a first experiment, cells were electroporated with TRC 1-2x.87EE (2 µg) and either mock transduced or transduced with increasing amounts of AAV412 (6.25e⁴, 1.25e⁴, 2.5e⁴ or e⁴ viral genomes/cell). On day 4 post-transduction, cells were

labeled with an antibody against CD3 and analyzed by flow cytometry (FIG. 37A). In the mock transduced cells, 20.7% are CD3⁻ compared to 21.6%, 23.7%, 25.5% and 25% with increasing AAV412, indicating that TRC 1-2x.87EE knockout efficiency is up to 23% higher in the presence of AAV412 (25.5% compared to 20.7%).

[0398] To determine whether this increase in knockout efficiency was nuclease specific, in an additional experiment, cells were electroporated with mRNA (2 µg) encoding a nuclease targeting the P2-microglobulin gene and either mock transduced or transduced with increasing amounts of AAV412. Cells were stained for β2-microglobulin on day 4 post-transduction and analyzed by flow cytometry (FIG. 37B). In the mock transduced cells, P2-microglobulin knockout efficiency was 64.5% and increased in the transduced cells to 68.6%, 70.7%, 77.2% and 82.5% with increasing amounts of AAV412, demonstrating an increase in knockout efficiency of up to 27.9% (82.5% compared to 64.5%).

[0399] In a parallel experiment, cells were electroporated with TRC 1-2x.87EE mRNA and either mock transduced or transduced with AAV422 (using the same MOIs as AAV412). Cells were labeled with an antibody against CD3 and cells were analyzed by flow cytometry (FIG. 37C). The mock transduced cells showed 62.2% T cell receptor knockout, and with increasing amounts of AAV, the T cell receptor knockout frequency increased to 72.6%, 75.5%, 78.3% and 75.1%. Here, the presence of AAV422 increases the knockout efficiency of TRC 1-2x.87EE by up to 25.8% (78.3% compared to 62.2%). It is striking that the increase in percent knockout efficiency is almost identical between these three experiments, using two different nucleases and two different AAV vectors. Taken together, these data strongly indicate that transduction of cells with single strand AAV vectors increase the knockout efficiency of our nucleases, irrespective of nuclease or AAV cargo.

5. Activity of T Cells Expressing Anti-CD19 Chimeric Antigen Receptor

[0400] The above experiments clearly demonstrate the generation of CAR T cells by electroporating cells with TRC 1-2x.87EE mRNA, then immediately transducing cells with AAV421, and that these cells can be enriched for a CD3⁻/CAR⁺ population by co-culture with CD19 expressing IM-9 cells. We next examined the activity of these CAR T cells against target cells. In the first experiment, the cells described above and shown in FIG. 34C were used in an IFN-gamma ELISPOT assay, in which either CD19⁺ Raji cells or CD19⁻ U937 cells were the target population. As shown in FIG. 38A, when anti-CD19 CAR T cells were incubated with U937 cells, they did not secrete IFN-gamma regardless of the target:effector ratio. Incubating CAR T cells with Raji cells, however, resulted in high levels of IFN-gamma secretion, in a dose-dependent manner, indicating that secretion of IFN-gamma is antigen-specific.

[0401] These CAR T cells were also used in a cell killing assay in which luciferase-labeled Raji cells were the target. Briefly, CAR T cells were incubated with luciferase-labeled Raji cells at a ratio of 10:1. At several time points, cells were washed and lysed to measure luciferase activity as a measure of how many cells remained. Control cells showed luciferase activity greater than 5500 arbitrary units (FIG. 38B). Co-incubation for 2, 3, 4 and 5 hours resulted in a decrease in luciferase activity to 4598, 3292, 2750 and 1932 arbitrary

units, respectively. Thus, within 5 hours of co-incubation, luciferase activity was reduced approximately 65%, indicating strong cytolytic activity of the CAR T cells.

[0402] Taken together, these data demonstrate that anti-CD19 CAR T cells generated according to the methods described herein are effective at killing CD19⁺ cells.

Example 7

Linearized Plasmid DNA

[0403] 1. Expression of Chimeric Antigen Receptor from Linearized Plasmid DNA

[0404] Since HDR templates produced by AAV are linear DNA molecules, we hypothesized that linear DNA from any source may be a suitable HDR template for inserting a CAR gene into the TRC 1-2 recognition sequence. To test this, we generated several plasmids that contain an anti-CD19 CAR gene flanked by homology arms that are homologous to the TRC 1-2 recognition sequence locus. Different promoters were used in some plasmids, and homology arms were either “short” (200 bp on the 5' homology arm and 180 bp on the 3' homology arm) to mimic the self-complimentary AAV vectors, or “long” (985 bp on the 5' homology arm and 763 bp on the 3' homology arm) to mimic the single strand AAV vectors. Plasmids with short homology arms are labeled “pDS” and those with long homology arms are labeled “pDI.” Additionally, some plasmid contained an intron upstream of the CAR gene.

[0405] The CAR donor plasmids were linearized at a restriction site in the vector backbone and gel purified. Human CD3⁺ T cells were either electroporated with the linearized CAR donor plasmid alone (varying amounts between 500 ng and 1000 ng, depending on the concentration of the purified linearized plasmid), or co-electroporated with TRC 1-2.87EE mRNA (2 µg). As controls, cells were either mock electroporated or electroporated with TRC 1-2x.87EE alone. The graphs in FIG. 39 are labelled with descriptions for all electroporations. Approximately 4 days post-electroporation, cells were labelled with antibodies against CD3 and the anti-CD19 CAR and analyzed by flow cytometry (FIG. 39). FIG. 39A shows background CD3⁺/CAR⁺ staining of 0.15%. It should be noted that the background CD3⁺/CAR⁺ staining was unusually high at 4.31%. FIG. 39B shows cells that were electroporated with TRC 1-2x.87EE mRNA alone, demonstrating 60.8% CD3 knock-out. FIGS. 39C and 39D represent samples that were co-electroporated with TRC 1-2x.87EE mRNA and either the long homology arm vector with an EF1α core promoter with an HTLV enhancer or the short homology arm vector with EF1α core promoter (with no enhancer). Interestingly, the linearized CAR donor with the EF1α core promoter alone generated a CD3⁺/CAR⁺ population of 2.38%, while the vector harboring the EF1α core promoter with the HTLV enhancer did not generate a significant percentage of CD3⁺/CAR⁺ cells. Cells that were electroporated with these two vectors in the absence of TRC 1-2x.87EE mRNA showed no significant increase in the CD3⁺/CAR⁺ population (FIGS. 39E and 39F). The increase in the CD3⁺/CAR⁺ population with the EF1α core promoter vector in the presence of TRC 1-2x.87EE suggested that a linearized plasmid could serve as an HDR template to repair double strand breaks at the TRC 1-2 recognition sequence.

[0406] FIGS. 39G and 39H show two long homology arm constructs that both contain an MND promoter driving

expression of the CAR. One of these constructs, shown in FIG. 39G, also contains an intron in the 5' end of the CAR gene. Surprisingly, the long homology arm plasmid with an MND promoter and intron showed significant CAR expression (FIG. 39G, 4.14% CD3⁺/CAR⁺) while the intron-less construct (FIG. 39H) did not show detectable CAR expression when co-electroporated with TRC 1-2x.87EE mRNA. A short homology arm plasmid with the MND promoter, but with no intron, was also tested with TRC 1-2x.87EE mRNA and did not demonstrate any CAR expression (FIG. 39I). None of the MND promoter-containing constructs generated any CAR⁺ cells in the absence of TRC 1-2x.87EE mRNA (FIGS. 39J, 39K, and 39L).

[0407] Lastly in this experiment, we tested a short homology arm construct that contained a JeT promoter driving expression of the CAR and a “long” homology arm construct with a CMV promoter driving expression of the CAR. Alone, neither of these linearized plasmids resulted in significant CAR⁺ cells (FIGS. 39O and 39P). When cells were co-electroporated with TRC 1-2x.87EE mRNA, the JeT containing construct showed 2.69% CD3⁺-CAR⁺ cells and the CMV containing construct yielded 2.7% CD3⁺/CAR⁺ cells.

[0408] The flow plots shown in FIG. 39 clearly demonstrate that linearized plasmid DNA that encodes the CAR, flanked by homology arms, can serve as HDR templates to repair DNA breaks caused by TRC 1-2x.87EE, resulting in insertion of the CAR nucleic acid. It is clear that promoter strength plays a significant role in expression of the CAR, and some promoters drive more efficient expression when there is an intron in the gene.

[0409] To confirm that insertion of the CAR using linearized DNA constructs was specific to the TRC 1-2 recognition sequence locus, we analyzed cells as described above using primers that sat within the CAR and outside of the homology arms (FIG. 40, Table 11). Samples 1 and 2 are PCR products from cells that were either mock electroporated or electroporated with only mRNA encoding TRC 1-2x.87EE. Consistent with results shown above, no PCR bands are present indicating the lack of CAR gene in the TRC 1-2 recognition site. Samples 3, 4 and 5 are from cells that were co-electroporated with TRC 1-2x.87EE and a linearized CAR homology plasmid (samples names in FIG. 40). Each sample shows two PCR bands of the predicted size indicating insertion of the CAR gene expression cassette into the TRC 1-2 recognition site. Samples 6, 7, and 8 are from cells that were electroporated with the same linearized CAR homology plasmids as samples 3, 4, and 5 but without TRC 1-2x.87EE mRNA. As expected, no PCR bands are present. Samples 9 and 10 are PCR products from cells that were either mock electroporated or electroporated with only mRNA encoding TRC 1-2x.87EE and show no PCR bands. Samples 11, 12, 13 and 14 are from cells that were co-electroporated with TRC 1-2x.87EE and a linearized CAR homology plasmid (samples names in FIG. 40). Each sample shows two PCR bands of the predicted size indicating insertion of the CAR gene into the TRC 1-2 recognition site. Samples 15, 16, 17, and 18 are from cells that were electroporated with the same linearized CAR homology plasmids as samples 11, 12, 13, and 14 but without TRC 1-2x.87EE mRNA. As expected, no PCR bands are present.

[0410] FIGS. 39 and 40 clearly demonstrate that co-electroporating human CD3⁺ T cells with mRNA encoding

TRC 1-2x.87EE and a linearized CAR homology plasmid is an effective method to insert the CAR gene into the TRC 1-2 recognition sequence.

TABLE 11

Sample	Nucleofection	Linearized plasmid
1	Mock (Water)	—
2	TRC1-2x87EE	—
3	TRC1-2x87EE	pDS EF1- α Core
4	TRC1-2x87EE	pDS 200 MND NC
5	TRC1-2x87EE	pDS 200 JET NC
6	Mock (Water)	pDS EF1- α Core
7	Mock (Water)	pDS 200 MND NC
8	Mock (Water)	pDS 200 JET NC
9	Mock (Water)	—
10	TRC1-2x87EE	—
11	TRC1-2x87EE	pDI EF1- α NC
12	TRC1-2x87EE	pDI MND intron NC
13	TRC1-2x87EE	pDI MND NC
14	TRC1-2x87EE	pDI CMV 985 NC 763
15	Mock (Water)	pDI EF1- α NC
16	Mock (Water)	pDI MND intron NC
17	Mock (Water)	pDI MND NC
18	Mock (Water)	pDI CMV 985 NC 763
19	Mock (Water)	—
20	TRC1-2x87EE	—
21	TRC1-2x87EE	pDS MCS
22	Mock (Water)	pDS MCS

Example 8

Characterization of Additional AAV Vectors

[0411] 1. Use of AAV with JeT Promoter and Long Homology Arms

[0412] Collectively, the data shown above indicate that vectors utilizing the JeT promoter drive high, consistent expression of the CAR and that longer homology arms may increase gene insertion efficiency. We designed and generated the vector shown in FIG. 41 (SEQ ID NO:125), which was used to make single-strand AAV with long homology arms, and a JeT promoter driving expression of the anti-CD19 CAR (referred to herein as AAV423). Human CD3⁺ T cells were electroporated with mRNA encoding TRC 1-2x.87EE and transduced with increasing amounts of AAV423. Since data shown above suggested that higher MOIs may result in increased insertion efficiency, we used titers ranging from 1.875×10^4 to 1.5×10^5 . As controls, cells were either electroporated with mRNA encoding TRC 1-2x.87EE then mock transduced or mock electroporated then transduced with increasing amounts of AAV423. On day 6 post-transduction, cells were labeled with antibodies recognizing CD3 or the anti-CD19 CAR and analyzed by flow cytometry. As shown in FIG. 42, cells that were mock electroporated then transduced with increasing amounts of AAV423 are overwhelmingly CD3⁺/CAR⁻ (ranging from 96.6% to 98.5%). Cells that were electroporated with mRNA encoding TRC 1-2x.87EE and mock transduced were 39% CD3⁻ indicating efficient knockout of the T cell receptor. In these cells, background CAR staining was very low (around 2%). Cells that were electroporated with mRNA encoding TRC 1-2x.87EE then transduced with

increasing amounts of AAV423 showed dramatic CAR staining in conjunction with CD3 knockout. CD3⁻/CAR⁺ populations ranged from 21.6% to 22.7%, while CD3⁺/CAR⁺ populations were around 2%. As described above, the presence of single-strand AAV increased the overall gene modification efficiency at the TRC 1-2 recognition site, with total CD3⁻ populations increasing from 41.44% in the control cells to 57.6%, 59.2%, 58.7%, and 56.1% in cells that were electroporated then transduced with increasing amounts of AV423. The percent of CD3⁻ cells that were CAR⁺ ranged from 37.5% to 39.9% indicating a dramatic increase in insertion efficiency compared to data described above.

[0413] To confirm that insertion of the CAR using AAV423 was specific to the TRC 1-2 recognition sequence locus, we analyzed cells as described above using primers that sat within the CAR and outside of the homology arms (FIG. 43, Table 12).

TABLE 12

Sample	Nucleofection	AAV (μ l)	MOI
1	Mock (Water)	—	—
2	Mock (Water)	—	—
3	Mock (Water)	pDI JET Prep A (3.125)	18750
4	Mock (Water)	pDI JET Prep A (6.25)	37500
5	Mock (Water)	pDI JET Prep A (12.5)	7500
6	Mock (Water)	pDI JET Prep A (25)	150000
7	TRC1-2x87EE	—	—
8	TRC1-2x87EE	pDI JET Prep A (3.125)	18750
9	TRC1-2x87EE	pDI JET Prep A (6.25)	37500
10	TRC1-2x87EE	pDI JET Prep A (12.5)	7500
11	TRC1-2x87EE	pDI JET Prep A (25)	150000

[0414] Samples 1 and 2 are PCR products from cells that were mock electroporated. Consistent with results shown above, no PCR bands are present indicating the lack of CAR gene in the TRC 1-2 recognition site. Samples 3-6 are from cells that were mock electroporated then transduced with increasing amounts of AAV423. Consistent with results above, there are no PCR bands present. Sample 7 is from cells electroporated with mRNA encoding TRC 1-2x.87EE then mock transduced, and shows no PCR bands. Samples 8-11 are from cells electroporated with mRNA encoding TRC 1-2x.87EE then transduced with increasing amounts of AAV423, and show the PCR bands expected if the CAR is inserted into the TRC 1-2 recognition sequence.

[0415] Given the ability of AAV423 to insert the CAR sequence into the TRC 1-2 recognition site following cleavage, it is further envisioned that the AAV423 plasmid (FIG. 41) could be linearized by digestion with a and delivered to the cell by digestion with one or more restriction enzymes, such that the T cells could be transfected with a linearized DNA template which could integrate into the TRC 1-2 recognition site and encode an anti-CD19 CAR.

Example 9

In Vivo Efficacy of Anti-CD19 TCR-Negative CAR T Cells

1. Murine Model of Disseminated B Cell Lymphoma

[0416] The efficacy of the gene-edited anti-CD19 CAR T cells was evaluated in a murine model of disseminated B cell lymphoma. Activated T cells were electroporated with TRC

1-2x.87 EE mRNA as described above, then transduced with an AAV6 vector comprising an anti-CD19 CAR expression cassette driven by a JeT promoter and flanked by homology arms. Following 5 days of culture with IL-2 (10 ng/mL), cells were analyzed for cell-surface CD3 and anti-CD19 CAR expression by flow cytometry as previously described (FIG. 44A). CD3⁻ cells were enriched by depleting CD3⁺ cells using anti-CD3 magnetic beads. Depleted cells were then cultured for 3 days in IL-15 (10 ng/mL) and IL-21 (10 ng/mL) and re-analyzed for cell-surface expression of CD3 and anti-CD19 CAR (FIG. 44B). Isolation of the CD3⁻ population was quite efficient, yielding 99.9% purity as measured by flow cytometry following depletion of CD3⁺ cells (FIG. 44B). The purified CD3⁻ population comprised 56% CD4⁺ and 44% CD8⁺ cells (FIG. 44C), and had primarily central memory/transitional memory phenotypes, determined by staining for CD62L and CD45RO (FIG. 44D).

[0417] Studies utilizing the Raji disseminated lymphoma model were conducted by Charles River Laboratories International Inc. (Morrisville, N.C., USA). CD19⁺ Raji cells stably expressing firefly luciferase (fLuc)⁴⁴ were injected i.v. into 5-6 week old female NSG mice on Day 1, at a dose of 2.0×10⁵ cells per mouse. On Day 4 mice were injected i.v. with PBS or PBS containing gene edited control TCR KO T cells prepared from the same healthy donor PBMC or PBS containing the indicated doses of CAR T cells prepared from the same donor. On the indicated days, live mice were

injected i.p. with Luciferin substrate (150 mg/kg in saline), anesthetized, and Luciferase activity measured after 7 minutes using IVIS SpectrumCT (Perkin Elmer, Waltham, Mass.). Data was analyzed and exported using Living Image software 4.5.1 (Perkin Elmer, Waltham, Mass.). Luminescence signal intensity is represented by radiance in p/sec/cm²/sr.

2. Results

[0418] As shown in FIG. 45, growth of CD19⁺ Raji cells was evident in all mice at low levels by day 8, and increased significantly in untreated and TCR⁻ control groups by day 11. In control groups, significant tumor growth was observed by day 15, and by day 18 or 19 all control groups were euthanized. In contrast, all groups of mice treated with anti-CD19 CAR T cells showed no evidence of tumor growth by day 11 and, with the exception of a single mouse in the low dose group, remained tumor-free through day 29 of the study. Tumor re-growth was observed in three mice in the low dose cohort around day 36. One of the three died at day 42, though imaging revealed only low levels of tumor in this animal, so it is unlikely that death was tumor-related.

3. Conclusions

[0419] These results provide clear evidence for in vivo clearance of CD19⁺ tumor cells by gene-edited CD3⁻ CAR T cells and support further preclinical development of this platform for allogeneic CAR T cell therapy.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 125

<210> SEQ ID NO 1

<211> LENGTH: 4627

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaatacc agtgacaagt      60
ctgtctgcct attcaccgat tttgattctc aaacaaatgt gtcacaaagt aaggattctg    120
atgtgtatat cacagacaaa actgtgctag acatgaggtc tatggacttc aagagcaaca    180
gtgctgtggc ctggagcaac aaatctgact ttgcatgtgc aaacgccttc aacaacagca    240
ttattccaga agacaccttc tccccagcc caggtaaggg cagctttggt gccttcgcag    300
gctgtttcct tgcttcagga atggccaggt tctgcccaga gctctggtea atgatgtcta    360
aaactcctct gattgtgggt ctggccctta tccattgcca ccaaaaccct ctttttacta    420
agaaacagtg agccttgctc tggcagtgca gagaatgaca cgggaaaaaa gcagatgaag    480
agaaggtggc aggagagggc acgtggccca gcctcagtct ctccaactga gttcctgcct    540
gcctgccttt gctcagactg tttgccctt actgctcttc taggcctcat tctaagcccc    600
ttctccaagt tgctctcctt tatttctccc tgtctgcca aaaatcttcc ccagctcact    660
aagtcagtct cagcagtgca ctcatcaacc caccaatcac tgattgtgcc ggcacatgaa    720
tgcaccaggt gttgaagtgg aggaattaa aagtcagatg aggggtgtgc ccagaggaag    780
caccattcta gttgggggag cccatctgtc agctgggaaa agtccaaata acttcagatt    840
ggaatgtgtt ttaactcagg gttgagaaaa cagctacctt caggacaaaa gtcaggggaag    900

```

-continued

ggctctctga	agaaatgcta	cttgaagata	ccagccctac	caagggcagg	gagaggacc	960
tatagaggcc	tgggacagga	gctcaatgag	aaaggagaag	agcagcaggc	atgagttgaa	1020
tgaaggaggc	agggccgggt	cacagggcct	tctaggccat	gagagggtag	acagtattct	1080
aaggacgcca	gaaagctggt	gatcggttc	aagcagggga	gggacaccta	atttgccttt	1140
cttttttttt	tttttttttt	tttttttttt	tgagatggag	ttttgctctt	gttgcccagg	1200
ctggagtgca	atggtgcatc	ttggctcact	gcaacctccg	cctcccagg	tcaagtgatt	1260
ctcctgcctc	agcctccoga	gtagctgaga	ttacaggcac	ccgccaccat	gcctggctaa	1320
ttttttgtat	tttttagtaga	gacaggggtt	cactatgttg	gccaggetgg	tctcgaactc	1380
ctgacctcag	gtgatccacc	cgcttcagcc	tcccaaagtg	ctgggattac	aggcgtgagc	1440
caccacaccc	ggcctgcttt	tcttaagat	caatctgagt	gctgtacgga	gagtgggttg	1500
taagccaaga	gtagaagcag	aaagggagca	gttcgagcag	agagatgatg	gaggcctggg	1560
caggtgggtg	gcagggagg	aaccaacacc	atcaggttt	caaaggtaga	accatgcagg	1620
gatgagaaa	caaagagggg	atcaaggaag	gcagctggat	tttggcctga	gcagctgagt	1680
caatgatagt	gccgtttact	aagaagaaac	caaggaaaaa	atttgggggtg	cagggatcaa	1740
aactttttgg	aacatagtaa	agtacgtgtt	tatactcttt	atggcccttg	tcactatgta	1800
tgccctcctg	cctccattgg	actctagaat	gaagccaggc	aagagcaggg	tctatgtgtg	1860
atggcacatg	tggccagggt	catgcaacat	gtactttgta	caaacagtgt	atattgagta	1920
aatagaaaatg	gtgtccagga	gccgaggtat	cggtcctgcc	agggccaggg	gctctcccta	1980
gcaggtgctc	atatgctgta	agttccctcc	agatctctcc	acaaggaggc	atggaaaaggc	2040
tgtagtgtt	cacctgcccc	agaactagga	ggtctgggg	gggagagtca	gcctgctctg	2100
gatgctgaaa	gaatgtctgt	ttttcctttt	agaaagtccc	tgatgatgca	agctggctga	2160
gaaaagcttt	gaaacaggtg	agacaggggt	ctagcctggg	tttgccacagg	attgcggaag	2220
tgatgaaccc	gcaataaccc	tgcttgatg	agggagtggt	aagaaattag	tagatgtggg	2280
aatgaatgat	gaggaatgga	aacagcgggt	caagacctgc	ccagagctgg	gtgggtctc	2340
tcttgaatcc	ctctcaccat	ctctgacttt	ccattctaag	cactttgagg	atgagtttct	2400
agcttcaata	gaccaaggac	tctctcctag	gcctctgtat	tcctttcaac	agctccactg	2460
tcaagagagc	cagagagagc	ttctgggtgg	cccagctgtg	aaatttctga	gtcccttagg	2520
gatagcccta	aacgaaccag	atcatcctga	ggacagccaa	gaggttttgc	cttctttcaa	2580
gacaagcaac	agtactcaca	taggctgtgg	gcaatggtcc	tgtctctcaa	gaatccctg	2640
ccactcctca	cacccaccct	gggcccata	tcaattccat	ttgagttggt	cttattgagt	2700
catccttctc	gtggtagcgg	aactcactaa	ggggcccata	tggacccgag	gtattgtgat	2760
gataaattct	gagcacctac	cccatcccc	gaagggctca	gaaataaaat	aagagccaag	2820
tctagtctgt	gtttcctgtc	ttgaaacaca	atactgttgg	ccctggaaga	atgcacagaa	2880
tctgtttgta	aggggatatg	cacagaagct	gcaagggaca	ggaggtgcag	gagctgcagg	2940
cctccccac	ccagcctgct	ctgccttggg	gaaaaccgtg	ggtgtgtcct	gcaggccatg	3000
caggcctggg	acatgcaagc	ccataaccgc	tgtggcctct	tggttttaca	gatacgaacc	3060
taaaacttca	aaacctgtca	gtgattgggt	tccgaatcct	cctcctgaaa	gtggccgggt	3120
ttaatctgct	catgacgtg	cggtctgtgt	ccagctgagg	tgaggggct	tgaagctggg	3180

-continued

```

agtggggttt agggacgagg gtctctgggt gcctcctaag ctctgagagc aaacctccct 3240
gcagggtctt gcttttaagt ccaaagcctg agcccaccaa actctcctac ttcttcctgt 3300
tacaaattcc tcttgtgcaa taataatgac ctgaaacgct gtaaaatatic ctcatctcag 3360
ccgcctcagt tgcactcttc cccatgagg taggaagaac agttgtttag aaacgaagaa 3420
actgaggccc cacagctaata gaggaggaga agagagacac ttgtgtacac cacatgcctt 3480
gtgttgact tctctcaccg tgtaacctcc tcatgtcctc tctcccagc acggctctct 3540
tagctcagta gaaagaagac attacactca tattacacc ccaatcctggc tagagtctcc 3600
gcacctcct cccccagggt cccagctgt cttgtgaca actgcactct gttccatcac 3660
catcaaaaaa aaactccagg ctgggtgagg gggctcacac ctgtaatccc agcactttgg 3720
gaggcagagg caggaggagc acaggagctg gagaccagcc tgggcaaac agggagacc 3780
cgctctaca aaaagtgaaa aaattaacca ggtgtggtgc tgcacacctg tagtcccagc 3840
tacttaagag gctgagatgg gaggatcgt tgagccctgg aatgttgagg ctacaatgag 3900
ctgtgattgc gtcactgcac tccagcctgg aagacaaagc aagatcctgt ctcaaataat 3960
aaaaaaaaata agaactccag ggtacatttg ctctagaac tctaccacat agccccaaac 4020
agagccatca ccacacac cctaacagtc ctgggtcttc ctgagtgtcc agcctgactt 4080
ctgttcttcc tattccaga tctgcaagat tgtaagacag cctgtgtctc ctgctctt 4140
cctctgcatt gccctcttc tccctctcca aacagaggga actctcctac cccaaggag 4200
gtgaaagctg ctaccacctc tgtgcccccc cggcaatgcc accaactgga tccatccga 4260
atztatgatt aagattgctg aagagctgcc aaacactgct gccacccct ctgttccctt 4320
attgtgctt gtcactgct gacattcac gcagaggcaa ggctgtgca gcctccctg 4380
gctgtgcaca ttcctctctg ctcccagag actgctccg ccaccccaca gatgatggat 4440
cttcagtggt ttctctggg ctctaggtcc tgcagaatgt tgtgaggggt ttatttttt 4500
ttaatagtgt tcaaaagaa atacatagta ttctctctc caagacgtgg ggggaaatta 4560
tctcattatc gagccctgc tatgtgtgt atctgggct gttgtatgtc ctgctgccga 4620
tgcttc 4627

```

```

<210> SEQ ID NO 2
<211> LENGTH: 142
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Pro Asn Ile Gln Asn Pro Asp Pro Ala Val Tyr Gln Leu Arg Asp Ser
1 5 10 15
Lys Ser Ser Asp Lys Ser Val Cys Leu Phe Thr Asp Phe Asp Ser Gln
20 25 30
Thr Asn Val Ser Gln Ser Lys Asp Ser Asp Val Tyr Ile Thr Asp Lys
35 40 45
Thr Val Leu Asp Met Arg Ser Met Asp Phe Lys Ser Asn Ser Ala Val
50 55 60
Ala Trp Ser Asn Lys Ser Asp Phe Ala Cys Ala Asn Ala Phe Asn Asn
65 70 75 80
Ser Ile Ile Pro Glu Asp Thr Phe Phe Pro Ser Pro Glu Ser Ser Cys
85 90 95

```

-continued

```

Asp Val Lys Leu Val Glu Lys Ser Phe Glu Thr Asp Thr Asn Leu Asn
      100                             105                             110

Phe Gln Asn Leu Ser Val Ile Gly Phe Arg Ile Leu Leu Leu Lys Val
      115                             120                             125

Ala Gly Phe Asn Leu Leu Met Thr Leu Arg Leu Trp Ser Ser
      130                             135                             140

```

```

<210> SEQ ID NO 3
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 3

```

```

tggcctggag caacaaatct ga 22

```

```

<210> SEQ ID NO 4
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 4

```

```

acaaatgtgt cacaaagtaa gg 22

```

```

<210> SEQ ID NO 5
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 5

```

```

ctgatgtgta taccacagac aa 22

```

```

<210> SEQ ID NO 6
<211> LENGTH: 163
<212> TYPE: PRT
<213> ORGANISM: Chlamydomonas reinhardtii

```

```

<400> SEQUENCE: 6

```

```

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
  1      5      10      15

Val Asp Gly Asp Gly Ser Ile Ile Ala Gln Ile Lys Pro Asn Gln Ser
      20      25      30

Tyr Lys Phe Lys His Gln Leu Ser Leu Ala Phe Gln Val Thr Gln Lys
      35      40      45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
      50      55      60

Gly Tyr Val Arg Asp Arg Gly Ser Val Ser Asp Tyr Ile Leu Ser Glu
      65      70      75      80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
      85      90      95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Trp Arg Leu
      100      105      110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
      115      120      125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
      130      135      140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys

```

-continued

145	150	155	160
Ser Ser Pro			
<210> SEQ ID NO 7			
<211> LENGTH: 9			
<212> TYPE: PRT			
<213> ORGANISM: Chlamydomonas reinhardtii			
<400> SEQUENCE: 7			
Leu	Ala	Gly	Leu Ile Asp Ala Asp Gly
1		5	
<210> SEQ ID NO 8			
<211> LENGTH: 354			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthesized			
<400> SEQUENCE: 8			
Met	Asn	Thr	Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
1		5	10 15
Val	Asp	Gly	Asp Gly Ser Ile Phe Ala Ser Ile Tyr Pro His Gln Arg
	20		25 30
Ala	Lys	Phe	Lys His Phe Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys
	35		40 45
Thr	Gln	Arg	Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
	50		55 60
Gly	Tyr	Val	Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu
65		70	75 80
Ile	Lys	Pro	Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
	85		90 95
Leu	Lys	Gln	Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
	100		105 110
Pro	Ser	Ala	Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
	115		120 125
Val	Asp	Gln	Ile Ala Ala Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr
	130		135 140
Ser	Glu	Thr	Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
145		150	155 160
Gly	Leu	Ser	Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
	165		170 175
Ser	Pro	Gly	Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
	180		185 190
Gly	Thr	Gly	Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
	195		200 205
Asp	Gly	Asp	Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
210		215	220
Lys	Phe	Lys	His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
225		230	235 240
Gln	Arg	Arg	Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
	245		250 255
Tyr	Val	Tyr	Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile
	260		265 270

-continued

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350

Ser Pro

<210> SEQ ID NO 9
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 9

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15

Val Asp Gly Asp Gly Ser Ile Phe Ala Ser Ile Tyr Pro His Gln Arg
 20 25 30

Ala Lys Phe Lys His Phe Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys
 35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60

Gly Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr
 130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205

Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
 210 215 220

Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255

Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile

-continued

260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

 <210> SEQ ID NO 10
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

 <400> SEQUENCE: 10
 Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Phe Ala Ser Ile Tyr Pro His Gln Arg
 20 25 30
 Ala Lys Phe Lys His Phe Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
 210 215 220
 Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255

-continued

Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 11
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 11

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Phe Ala Ser Ile Tyr Pro His Gln Arg
 20 25 30
 Ala Lys Phe Lys His Phe Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
 210 215 220
 Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240

-continued

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
245 250 255

Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
340 345 350

Ser Pro

<210> SEQ ID NO 12

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 12

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
1 5 10 15

Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Tyr Pro His Gln Arg
20 25 30

Ala Lys Phe Lys His Leu Leu Lys Leu Val Phe Ala Val His Gln Lys
35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
50 55 60

Gly Tyr Val Tyr Asp Ala Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
195 200 205

Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
210 215 220

Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr

-continued

```

225                230                235                240
Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
      245                250                255
Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
      260                265                270
Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
      275                280                285
Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
      290                295                300
Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
      305                310                315                320
Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
      325                330                335
Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
      340                345                350

Ser Pro

<210> SEQ ID NO 13
<211> LENGTH: 354
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 13
Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1      5      10      15
Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Tyr Pro Asp Gln Arg
 20     25     30
Thr Lys Phe Lys His Gly Leu Arg Leu Asn Phe Ser Val Phe Gln Lys
 35     40     45
Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50     55     60
Gly Tyr Val Phe Asp Ala Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65     70     75     80
Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85     90     95
Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
100    105    110
Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
115    120    125
Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
130    135    140
Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
145    150    155    160
Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
165    170    175
Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
180    185    190
Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
195    200    205
Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
210    215    220

```

-continued

Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 14
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 14

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile His Pro Asp Gln Arg
 20 25 30
 Ser Lys Phe Lys His Tyr Leu Arg Leu Phe Phe Ser Val Phe Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Ala Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205

-continued

Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Ala Pro Cys Gln Arg Ala
 210 215 220

Lys Phe Lys His Arg Leu Lys Leu Gly Phe Thr Val Gly Gln Lys Thr
 225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255

His Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile
 260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350

Ser Pro

<210> SEQ ID NO 15
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 15

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15

Val Asp Gly Asp Gly Ser Ile Phe Ala Gln Ile Lys Pro Asp Gln Lys
 20 25 30

Met Lys Phe Lys His Tyr Leu Ser Leu His Phe Ser Val Phe Gln Lys
 35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60

Gly Tyr Val Tyr Asp Gly Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val

-continued

195					200					205					
Asp	Gly	Asp	Gly	Ser	Ile	Tyr	Ala	Gln	Ile	Lys	Pro	Gln	Gln	Arg	Ala
210						215					220				
Lys	Phe	Lys	His	Arg	Leu	Leu	Leu	Ala	Phe	Thr	Val	Ser	Gln	Lys	Thr
225						230					235				240
Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val	Gly
				245					250					255	
Tyr	Val	Ile	Asp	Arg	Gly	Gly	Val	Ser	Glu	Tyr	Ile	Leu	Ser	Glu	Ile
			260						265					270	
Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu
		275					280					285			
Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro
	290					295					300				
Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp	Val
305						310					315				320
Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr	Ser
				325					330					335	
Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Ser	Glu	Lys	Lys	Lys	Ser
			340						345					350	

Ser Pro

<210> SEQ ID NO 16

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 16

Met	Asn	Thr	Lys	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe
1				5					10					15	
Val	Asp	Gly	Asp	Gly	Ser	Ile	Phe	Ala	Ser	Ile	Asn	Pro	Asp	Gln	Arg
		20						25					30		
Ala	Lys	Phe	Lys	His	Ser	Leu	Lys	Leu	Thr	Phe	Ser	Val	Tyr	Gln	Lys
		35					40					45			
Thr	Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val
50						55					60				
Gly	Tyr	Val	Tyr	Asp	Thr	Gly	Ser	Val	Ser	Glu	Tyr	Arg	Leu	Ser	Gln
65				70							75				80
Ile	Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys
				85					90					95	
Leu	Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu
		100					105						110		
Pro	Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp
		115					120					125			
Val	Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr
	130					135					140				
Ser	Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Pro	Gly	Ser	Val	Gly
145						150					155				160
Gly	Leu	Ser	Pro	Ser	Gln	Ala	Ser	Ser	Ala	Ala	Ser	Ser	Ala	Ser	Ser
				165					170					175	
Ser	Pro	Gly	Ser	Gly	Ile	Ser	Glu	Ala	Leu	Arg	Ala	Gly	Ala	Gly	Ser
		180						185						190	

-continued

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Tyr Ala Ser Ile Arg Pro Ser Gln Arg Ser
 210 215 220
 Lys Phe Lys His Lys Leu Gly Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 17
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 17

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Phe Ala Ser Ile Tyr Pro His Gln Arg
 20 25 30
 Ala Lys Phe Lys His Phe Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175

-continued

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205

Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
 210 215 220

Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
 225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255

Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
 260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350

Ser Pro

<210> SEQ ID NO 18
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 18

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15

Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Asp Gln Arg
 20 25 30

Ala Lys Phe Lys His Tyr Leu Arg Leu Gln Phe Ser Val Phe Gln Lys
 35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60

Gly Tyr Val Phe Asp Ala Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
 65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser

-continued

```

165           170           175
Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
180           185           190
Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
195           200           205
Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser
210           215           220
Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr
225           230           235           240
Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
245           250           255
Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile
260           265           270
Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
275           280           285
Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
290           295           300
Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
305           310           315           320
Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
325           330           335
Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
340           345           350

```

Ser Pro

```

<210> SEQ ID NO 19
<211> LENGTH: 354
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

<400> SEQUENCE: 19

```

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
1           5           10           15
Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly
20           25           30
Ser Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys
35           40           45
Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
50           55           60
Gly Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln
65           70           75           80
Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
85           90           95
Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
100          105          110
Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
115          120          125
Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
130          135          140
Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
145          150          155          160

```

-continued

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Tyr Pro Asp Gln Arg Ala
 210 215 220
 Lys Phe Lys His Ala Leu Lys Leu Ile Phe Ser Val Phe Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Gly Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 20
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 20

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Arg Pro Ala Gln Arg
 20 25 30
 Ala Lys Phe Lys His Arg Leu Val Leu Gly Phe Glu Val Gly Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Gly Gly Ser Val Ser Lys Tyr Arg Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140

-continued

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Ala Pro Asp Gln Arg Pro
 210 215 220
 Lys Phe Lys His Gln Leu Arg Leu Ile Phe Asn Val Cys Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Thr Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 21
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 21

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly
 20 25 30
 Ser Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr

-continued

130					135					140					
Ser	Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Pro	Gly	Ser	Val	Gly
145				150					155					160	
Gly	Leu	Ser	Pro	Ser	Gln	Ala	Ser	Ser	Ala	Ala	Ser	Ser	Ala	Ser	Ser
				165					170					175	
Ser	Pro	Gly	Ser	Gly	Ile	Ser	Glu	Ala	Leu	Arg	Ala	Gly	Ala	Gly	Ser
			180					185					190		
Gly	Thr	Gly	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe	Val
		195					200					205			
Asp	Gly	Asp	Gly	Ser	Ile	Phe	Ala	Ser	Ile	Tyr	Pro	His	Gln	Arg	Ala
210						215					220				
Lys	Phe	Lys	His	Phe	Leu	Lys	Leu	Thr	Phe	Ala	Val	Tyr	Gln	Lys	Thr
225					230					235					240
Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val	Gly
				245					250					255	
Tyr	Val	Tyr	Asp	Ser	Gly	Ser	Val	Ser	Glu	Tyr	Arg	Leu	Ser	Gln	Ile
			260					265						270	
Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu
		275					280					285			
Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro
290					295					300					
Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp	Val
305					310					315					320
Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr	Ser
				325					330					335	
Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Ser	Glu	Lys	Lys	Lys	Ser
			340					345						350	

Ser Pro

<210> SEQ ID NO 22
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 22

Met	Asn	Thr	Lys	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe
1				5					10					15	
Val	Asp	Gly	Asp	Gly	Ser	Ile	Tyr	Ala	Cys	Ile	Ala	Pro	Arg	Gln	Gly
		20						25					30		
Ser	Lys	Phe	Lys	His	Arg	Leu	Lys	Leu	Gly	Phe	Ala	Val	Gly	Gln	Lys
		35					40					45			
Thr	Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val
		50				55					60				
Gly	Tyr	Val	Tyr	Asp	Arg	Gly	Ser	Val	Ser	Glu	Tyr	Val	Leu	Ser	Gln
65						70					75				80
Ile	Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys
				85					90					95	
Leu	Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu
			100						105					110	
Pro	Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp
			115					120						125	

-continued

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Val Pro Glu Gln Arg Ser
 210 215 220
 Lys Phe Lys His Tyr Leu Lys Leu Thr Phe Ser Val Phe Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Arg Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Ala Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 23
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 23

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly
 20 25 30
 Ser Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

-continued

```

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
   115                               120                               125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
   130                               135                               140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
  145                               150                               155                               160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
   165                               170                               175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
   180                               185                               190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
   195                               200                               205

Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met
  210                               215                               220

Lys Phe Lys His Gln Leu Arg Leu His Phe Cys Val His Gln Lys Thr
  225                               230                               235                               240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
   245                               250                               255

Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile
   260                               265                               270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
   275                               280                               285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
   290                               295                               300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
  305                               310                               315                               320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
   325                               330                               335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
   340                               345                               350

Ser Pro

```

<210> SEQ ID NO 24

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 24

```

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
  1                               5                               10                               15

Val Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Ala Pro Cys Gln Arg
   20                               25                               30

Ala Lys Phe Lys His Arg Leu Lys Leu Gly Phe Thr Val Gly Gln Lys
   35                               40                               45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
   50                               55                               60

Gly His Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Gln
   65                               70                               75                               80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
   85                               90                               95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu

```

-continued

100					105					110					
Pro	Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp
		115					120					125			
Val	Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr
	130					135					140				
Ser	Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Pro	Gly	Ser	Val	Gly
145						150					155				160
Gly	Leu	Ser	Pro	Ser	Gln	Ala	Ser	Ser	Ala	Ala	Ser	Ser	Ala	Ser	Ser
				165					170					175	
Ser	Pro	Gly	Ser	Gly	Ile	Ser	Glu	Ala	Leu	Arg	Ala	Gly	Ala	Gly	Ser
				180					185					190	
Gly	Thr	Gly	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe	Val
				195					200					205	
Asp	Gly	Asp	Gly	Ser	Ile	Phe	Ala	Ser	Ile	Tyr	Pro	His	Gln	Arg	Ala
210						215					220				
Lys	Phe	Lys	His	Phe	Leu	Lys	Leu	Thr	Phe	Ala	Val	Tyr	Gln	Lys	Thr
225						230					235				240
Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val	Gly
				245					250					255	
Tyr	Val	Tyr	Asp	Ser	Gly	Ser	Val	Ser	Glu	Tyr	Arg	Leu	Ser	Gln	Ile
				260					265					270	
Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu
				275					280					285	
Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro
290						295					300				
Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp	Val
305						310					315				320
Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr	Ser
				325					330					335	
Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Ser	Glu	Lys	Lys	Lys	Ser
				340					345					350	

Ser Pro

<210> SEQ ID NO 25

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 25

Met	Asn	Thr	Lys	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe
1				5					10					15	
Val	Asp	Gly	Asp	Gly	Ser	Ile	Tyr	Ala	Cys	Ile	Ala	Pro	Arg	Gln	Gly
				20					25					30	
Ser	Lys	Phe	Lys	His	Arg	Leu	Lys	Leu	Gly	Phe	Ala	Val	Gly	Gln	Lys
				35					40					45	
Thr	Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val
				50					55					60	
Gly	Tyr	Val	Tyr	Asp	Arg	Gly	Ser	Val	Ser	Glu	Tyr	Val	Leu	Ser	Gln
65						70					75			80	
Ile	Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys
				85					90					95	

-continued

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met
 210 215 220
 Lys Phe Lys His Gln Leu Arg Leu His Phe Cys Val His Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 26
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 26

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly
 20 25 30
 Ser Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys
 35 40 45
 Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60
 Gly Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Glu
 65 70 75 80

-continued

```

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
      85
Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
      100
Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
      115
Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
      130
Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
      145
Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
      165
Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
      180
Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
      195
Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met
      210
Lys Phe Lys His Gln Leu Arg Leu His Phe Cys Val His Gln Lys Thr
      225
Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
      245
Tyr Val Tyr Asp Ser Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile
      260
Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
      275
Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
      290
Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
      305
Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
      325
Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
      340
Ser Pro

```

<210> SEQ ID NO 27

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 27

```

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
  1
Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Ala Pro Gly Gln Gly
      20
Ser Lys Phe Lys His Arg Leu Lys Leu Gly Phe Ala Val Gly Gln Lys
      35
Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
      50
Gly Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Val Leu Ser Glu

```

-continued

65	70	75	80
Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys 85 90 95			
Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu 100 105 110			
Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp 115 120 125			
Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr 130 135 140			
Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly 145 150 155 160			
Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser 165 170 175			
Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser 180 185 190			
Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val 195 200 205			
Asp Gly Asp Gly Ser Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met 210 215 220			
Lys Phe Lys His Gln Leu Arg Leu Gly Phe Ala Val His Gln Lys Thr 225 230 235 240			
Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly 245 250 255			
Tyr Val Tyr Asp Arg Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile 260 265 270			
Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu 275 280 285			
Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro 290 295 300			
Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val 305 310 315 320			
Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser 325 330 335			
Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser 340 345 350			

Ser Pro

<210> SEQ ID NO 28

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 28

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe 1 5 10 15
Val Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Cys Pro Cys Gln Thr 20 25 30
Leu Lys Phe Lys His Tyr Leu Thr Leu Ser Phe Ser Val Tyr Gln Lys 35 40 45
Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val 50 55 60

-continued

Gly Tyr Val Tyr Asp Gln Gly Ser Val Ser Cys Tyr Arg Leu Ser Gln
 65 70 75 80
 Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95
 Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110
 Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125
 Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140
 Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160
 Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175
 Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190
 Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205
 Asp Gly Asp Gly Ser Ile His Ala Cys Ile Gln Pro Gln Gln Asp Val
 210 215 220
 Lys Phe Lys His Gln Leu His Leu Arg Phe Thr Val His Gln Lys Thr
 225 230 235 240
 Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255
 Tyr Val Tyr Asp Ala Gly Ser Val Ser Thr Tyr Cys Leu Ser Gln Ile
 260 265 270
 Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285
 Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300
 Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320
 Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335
 Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350
 Ser Pro

<210> SEQ ID NO 29
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 29

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15
 Val Asp Gly Asp Gly Ser Ile Tyr Ala Thr Ile Cys Pro Asp Gln Ala
 20 25 30
 Leu Lys Phe Lys His Tyr Leu Ser Leu Thr Phe Ala Val Tyr Gln Lys
 35 40 45

-continued

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60

Gly Tyr Val Tyr Asp Gln Gly Ser Val Ser Cys Tyr Arg Leu Ser Gln
 65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205

Asp Gly Asp Gly Ser Ile His Ala Cys Ile Gln Pro Met Gln Ser Met
 210 215 220

Lys Phe Lys His Tyr Leu His Leu Arg Phe Thr Val His Gln Lys Thr
 225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Thr Gly Val Gly
 245 250 255

Tyr Val Tyr Asp Ala Gly Ser Val Ser Thr Tyr Cys Leu Ser Gln Ile
 260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350

Ser Pro

<210> SEQ ID NO 30
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 30

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
 1 5 10 15

Val Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp
 20 25 30

Met Lys Phe Lys His Arg Leu Gln Leu Arg Phe Cys Val Thr Gln Lys

-continued

	35					40					45								
Thr	Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val				
	50					55					60								
Gly	Tyr	Val	Gln	Asp	Cys	Gly	Ser	Val	Ser	Glu	Tyr	Arg	Leu	Ser	Glu				
65				70						75					80				
Ile	Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys				
				85					90					95					
Leu	Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu				
			100						105					110					
Pro	Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp				
							120						125						
Val	Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr				
	130					135						140							
Ser	Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Pro	Gly	Ser	Val	Gly				
145					150					155					160				
Gly	Leu	Ser	Pro	Ser	Gln	Ala	Ser	Ser	Ala	Ala	Ser	Ser	Ala	Ser	Ser				
				165					170						175				
Ser	Pro	Gly	Ser	Gly	Ile	Ser	Glu	Ala	Leu	Arg	Ala	Gly	Ala	Gly	Ser				
			180					185						190					
Gly	Thr	Gly	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe	Val				
						200								205					
Asp	Gly	Asp	Gly	Ser	Ile	Val	Ala	Ser	Ile	Lys	Pro	Gln	Gln	Val	Ala				
	210					215					220								
Lys	Phe	Lys	His	Arg	Leu	Met	Leu	Glu	Phe	Tyr	Val	Tyr	Gln	Lys	Thr				
225					230					235					240				
Gln	Arg	Arg	Trp	Phe	Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val	Gly				
				245						250					255				
Tyr	Val	Tyr	Asp	Leu	Gly	Gly	Ala	Ser	Arg	Tyr	Val	Leu	Ser	Gln	Ile				
			260						265					270					
Lys	Pro	Leu	His	Asn	Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu				
		275					280							285					
Lys	Gln	Lys	Gln	Ala	Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro				
	290					295					300								
Ser	Ala	Lys	Glu	Ser	Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp	Val				
305					310					315					320				
Asp	Gln	Ile	Ala	Ala	Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr	Ser				
			325						330						335				
Glu	Thr	Val	Arg	Ala	Val	Leu	Asp	Ser	Leu	Ser	Glu	Lys	Lys	Lys	Ser				
			340						345						350				

Ser Pro

<210> SEQ ID NO 31
 <211> LENGTH: 354
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized
 <400> SEQUENCE: 31

Met	Asn	Thr	Lys	Tyr	Asn	Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe				
1				5					10					15					
Val	Asp	Gly	Asp	Gly	Ser	Ile	Tyr	Ala	Cys	Ile	Lys	Pro	Asp	Gln	Ala				
			20					25						30					

-continued

Ala Lys Phe Lys His Arg Leu Leu Leu Glu Phe Thr Val Cys Gln Lys
35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
50 55 60

Gly Tyr Val Val Asp Gln Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln
65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
195 200 205

Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp Met
210 215 220

Lys Phe Lys His Arg Leu Gln Leu Arg Phe Cys Val Thr Gln Lys Thr
225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
245 250 255

Tyr Val Gln Asp His Gly Gly Ala Ser Glu Tyr Arg Leu Ser Glu Ile
260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
340 345 350

Ser Pro

<210> SEQ ID NO 32

<211> LENGTH: 354

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 32

Met Asn Thr Lys Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe
1 5 10 15

-continued

Val Asp Gly Asp Gly Ser Ile Trp Ala Ser Ile Arg Pro Thr Gln Leu
 20 25 30

Ala Lys Phe Lys His Ala Leu Trp Leu Gly Phe Ala Val Tyr Gln Lys
 35 40 45

Thr Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val
 50 55 60

Gly Tyr Val Tyr Asp Ser Gly Ser Val Ser Lys Tyr Thr Leu Ser Glu
 65 70 75 80

Ile Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys
 85 90 95

Leu Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu
 100 105 110

Pro Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp
 115 120 125

Val Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr
 130 135 140

Ser Glu Thr Val Arg Ala Val Leu Asp Ser Leu Pro Gly Ser Val Gly
 145 150 155 160

Gly Leu Ser Pro Ser Gln Ala Ser Ser Ala Ala Ser Ser Ala Ser Ser
 165 170 175

Ser Pro Gly Ser Gly Ile Ser Glu Ala Leu Arg Ala Gly Ala Gly Ser
 180 185 190

Gly Thr Gly Tyr Asn Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val
 195 200 205

Asp Gly Asp Gly Ser Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp Met
 210 215 220

Lys Phe Lys His Arg Leu Gln Leu Arg Phe Cys Val Thr Gln Lys Thr
 225 230 235 240

Gln Arg Arg Trp Phe Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly
 245 250 255

Tyr Val Gln Asp Lys Gly Ser Ala Ser Glu Tyr Arg Leu Ser Glu Ile
 260 265 270

Lys Pro Leu His Asn Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu
 275 280 285

Lys Gln Lys Gln Ala Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro
 290 295 300

Ser Ala Lys Glu Ser Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val
 305 310 315 320

Asp Gln Ile Ala Ala Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser
 325 330 335

Glu Thr Val Arg Ala Val Leu Asp Ser Leu Ser Glu Lys Lys Lys Ser
 340 345 350

Ser Pro

<210> SEQ ID NO 33
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized
 <400> SEQUENCE: 33

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser

-continued

1	5	10	15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg	20	25	30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg	50	55	60
Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 34
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 34

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser	5	10	15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg	20	25	30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg	50	55	60
Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 35
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

-continued

<400> SEQUENCE: 35

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
 20 25 30
 Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 36

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 36

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
 20 25 30
 Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 37

<211> LENGTH: 147

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 37

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1           5           10           15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
20           25           30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35           40           45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50           55           60
Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65           70           75           80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85           90           95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100          105          110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115          120          125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130          135          140

Val Leu Asp
145

```

```

<210> SEQ ID NO 38
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 38

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1           5           10           15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
20           25           30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35           40           45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50           55           60
Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65           70           75           80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85           90           95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100          105          110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115          120          125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130          135          140

Val Leu Asp
145

```

-continued

<210> SEQ ID NO 39
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 39

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Thr Ile Ala Pro Cys Gln Arg Ala Lys Phe Lys His Arg
 20 25 30
 Leu Lys Leu Gly Phe Thr Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly His Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 40
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 40

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Gln Ile Lys Pro Gln Gln Arg Ala Lys Phe Lys His Arg
 20 25 30
 Leu Leu Leu Ala Phe Thr Val Ser Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Ile Asp Arg
 50 55 60
 Gly Gly Val Ser Glu Tyr Ile Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala

-continued

130 135 140

Val Leu Asp
145

<210> SEQ ID NO 41
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 41

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Tyr Ala Ser Ile Arg Pro Ser Gln Arg Ser Lys Phe Lys His Lys
20 25 30

Leu Gly Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130 135 140

Val Leu Asp
145

<210> SEQ ID NO 42
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 42

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
20 25 30

Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100 105 110

-continued

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130 135 140

Val Leu Asp
145

<210> SEQ ID NO 43
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 43

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
20 25 30

Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130 135 140

Val Leu Asp
145

<210> SEQ ID NO 44
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 44

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
20 25 30

Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

-continued

```

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100                      105                      110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115                      120                      125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130                      135                      140
Val Leu Asp
145

```

```

<210> SEQ ID NO 45
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 45

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1      5                      10                      15
Ile Tyr Ala Thr Ile Arg Pro Ala Gln Arg Ala Lys Phe Lys His Arg
      20                      25                      30
Leu Val Leu Gly Phe Glu Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
      35                      40                      45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Gly
50     55                      60
Gly Ser Val Ser Lys Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
65     70                      75                      80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
      85                      90                      95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100                     105                     110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115                     120                     125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130                     135                     140
Val Leu Asp
145

```

```

<210> SEQ ID NO 46
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 46

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1      5                      10                      15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
      20                      25                      30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
      35                      40                      45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
50     55                      60
Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn

```

-continued

```

65              70              75              80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
      85              90              95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100             105             110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115             120             125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130             135             140
Val Leu Asp
145

```

```

<210> SEQ ID NO 47
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 47

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1              5              10              15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
      20              25              30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
      35              40              45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
      50              55              60
Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
      65              70              75              80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
      85              90              95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100             105             110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115             120             125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130             135             140
Val Leu Asp
145

```

```

<210> SEQ ID NO 48
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 48

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1              5              10              15
Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
      20              25              30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
      35              40              45

```


-continued

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

Val Leu Asp
 145

<210> SEQ ID NO 49
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 49

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

Ile Tyr Ala Thr Ile Ala Pro Cys Gln Arg Ala Lys Phe Lys His Arg
 20 25 30

Leu Lys Leu Gly Phe Thr Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly His Val Tyr Asp Arg
 50 55 60

Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

Val Leu Asp
 145

<210> SEQ ID NO 50
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 50

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
 20 25 30

-continued

Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Val Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 51
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 51

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Cys Ile Ala Pro Arg Gln Gly Ser Lys Phe Lys His Arg
 20 25 30
 Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 52
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 52

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser

-continued

1	5	10	15
Ile Tyr Ala Cys Ile Ala Pro Gly Gln Gly Ser Lys Phe Lys His Arg	20	25	30
Leu Lys Leu Gly Phe Ala Val Gly Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg	50	55	60
Gly Ser Val Ser Glu Tyr Val Leu Ser Glu Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 53
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 53

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser	5	10	15
Ile Tyr Ala Thr Ile Cys Pro Cys Gln Thr Leu Lys Phe Lys His Tyr	20	25	30
Leu Thr Leu Ser Phe Ser Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Gln	50	55	60
Gly Ser Val Ser Cys Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 54
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

-continued

<400> SEQUENCE: 54

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Thr Ile Cys Pro Asp Gln Ala Leu Lys Phe Lys His Tyr
 20 25 30
 Leu Ser Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Gln
 50 55 60
 Gly Ser Val Ser Cys Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 55

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 55

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp Met Lys Phe Lys His Arg
 20 25 30
 Leu Gln Leu Arg Phe Cys Val Thr Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Gln Asp Cys
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 56

<211> LENGTH: 147

-continued

<212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

 <400> SEQUENCE: 56

 Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

 Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp Met Lys Phe Lys His Arg
 20 25 30

 Leu Gln Leu Arg Phe Cys Val Thr Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Gln Asp His
 50 55 60

 Gly Gly Ala Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80

 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

 Val Leu Asp
 145

<210> SEQ ID NO 57
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

 <400> SEQUENCE: 57

 Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

 Ile Tyr Ala Cys Ile Thr Pro Gln Gln Asp Met Lys Phe Lys His Arg
 20 25 30

 Leu Gln Leu Arg Phe Cys Val Thr Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Gln Asp Lys
 50 55 60

 Gly Ser Ala Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80

 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

 Val Leu Asp
 145

-continued

<210> SEQ ID NO 58
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 58

```
Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1           5           10           15
Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
20          25          30
Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
35          40          45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
50          55          60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
65          70          75          80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85          90          95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100         105        110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115        120        125
Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130        135        140
Val Leu Asp
145
```

<210> SEQ ID NO 59
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 59

```
Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1           5           10           15
Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
20          25          30
Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
35          40          45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
50          55          60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
65          70          75          80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85          90          95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100         105        110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115        120        125
Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
```

-continued

130 135 140

Val Leu Asp
145

<210> SEQ ID NO 60
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 60

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
20 25 30

Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
50 55 60

Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
115 120 125

Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
130 135 140

Val Leu Asp
145

<210> SEQ ID NO 61
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 61

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1 5 10 15

Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
20 25 30

Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
50 55 60

Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
100 105 110

-continued

```

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
    100                               105                110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
    115                               120                125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
    130                               135                140
Val Leu Asp
145

```

```

<210> SEQ ID NO 64
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 64

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1      5      10      15
Ile Phe Ala Thr Ile His Pro Asp Gln Arg Ser Lys Phe Lys His Tyr
 20     25     30
Leu Arg Leu Phe Phe Ser Val Phe Gln Lys Thr Gln Arg Arg Trp Phe
 35     40     45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ala
 50     55     60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65     70     75     80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85     90     95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100    105    110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115    120    125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130    135    140
Val Leu Asp
145

```

```

<210> SEQ ID NO 65
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 65

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1      5      10      15
Ile Phe Ala Gln Ile Lys Pro Asp Gln Lys Met Lys Phe Lys His Tyr
 20     25     30
Leu Ser Leu His Phe Ser Val Phe Gln Lys Thr Gln Arg Arg Trp Phe
 35     40     45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Gly
 50     55     60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn

```

-continued

```

65              70              75              80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
      85              90              95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100             105             110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115             120             125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130             135             140
Val Leu Asp
145

```

```

<210> SEQ ID NO 66
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 66

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1          5          10          15
Ile Phe Ala Ser Ile Asn Pro Asp Gln Arg Ala Lys Phe Lys His Ser
      20          25          30
Leu Lys Leu Thr Phe Ser Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
      35          40          45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Thr
      50          55          60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
      65          70          75          80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
      85          90          95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100             105             110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115             120             125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130             135             140
Val Leu Asp
145

```

```

<210> SEQ ID NO 67
<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 67

```

```

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
1          5          10          15
Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
      20          25          30
Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
      35          40          45

```

-continued

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
 50 55 60

Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

Leu Asn Asp Ser Arg Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

Val Leu Asp
 145

<210> SEQ ID NO 68
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 68

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

Ile Tyr Ala Cys Ile Ala Pro Asp Gln Arg Ala Lys Phe Lys His Tyr
 20 25 30

Leu Arg Leu Gln Phe Ser Val Phe Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Phe Asp Ala
 50 55 60

Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

Val Leu Asp
 145

<210> SEQ ID NO 69
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 69

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

Ile Tyr Ala Thr Ile Tyr Pro Asp Gln Arg Ala Lys Phe Lys His Ala
 20 25 30

-continued

Leu Lys Leu Ile Phe Ser Val Phe Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Gly
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 70
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 70

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Phe Ala Thr Ile Ala Pro Asp Gln Arg Pro Lys Phe Lys His Gln
 20 25 30
 Leu Arg Leu Ile Phe Asn Val Cys Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Thr
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Glu Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 71
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 71

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser

-continued

1	5	10	15
Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe	20	25	30
Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser	50	55	60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 72
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 72

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser	5	10	15
Ile Phe Ala Thr Ile Val Pro Glu Gln Arg Ser Lys Phe Lys His Tyr	20	25	30
Leu Lys Leu Thr Phe Ser Val Phe Gln Lys Thr Gln Arg Arg Trp Phe	35	40	45
Leu Asp Arg Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ala	50	55	60
Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn	65	70	80
Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala	85	90	95
Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser	100	105	110
Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala	115	120	125
Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala	130	135	140
Val Leu Asp			
145			

<210> SEQ ID NO 73
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

-continued

<400> SEQUENCE: 73

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met Lys Phe Lys His Gln
 20 25 30
 Leu Arg Leu His Phe Cys Val His Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 74

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 74

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Phe Ala Ser Ile Tyr Pro His Gln Arg Ala Lys Phe Lys His Phe
 20 25 30
 Leu Lys Leu Thr Phe Ala Val Tyr Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 75

<211> LENGTH: 147

-continued

<212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

 <400> SEQUENCE: 75

 Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

 Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met Lys Phe Lys His Gln
 20 25 30

 Leu Arg Leu His Phe Cys Val His Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
 50 55 60

 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

 Val Leu Asp
 145

<210> SEQ ID NO 76
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

 <400> SEQUENCE: 76

 Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15

 Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met Lys Phe Lys His Gln
 20 25 30

 Leu Arg Leu His Phe Cys Val His Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45

 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ser
 50 55 60

 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80

 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95

 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110

 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125

 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140

 Val Leu Asp
 145

-continued

<210> SEQ ID NO 77
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 77

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile Phe Ala Thr Ile Phe Pro Asp Gln Arg Met Lys Phe Lys His Gln
 20 25 30
 Leu Arg Leu Gly Phe Ala Val His Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Arg
 50 55 60
 Gly Ser Val Ser Glu Tyr Arg Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
 130 135 140
 Val Leu Asp
 145

<210> SEQ ID NO 78
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 78

Lys Glu Phe Leu Leu Tyr Leu Ala Gly Phe Val Asp Gly Asp Gly Ser
 1 5 10 15
 Ile His Ala Cys Ile Gln Pro Gln Gln Asp Val Lys Phe Lys His Gln
 20 25 30
 Leu His Leu Arg Phe Thr Val His Gln Lys Thr Gln Arg Arg Trp Phe
 35 40 45
 Leu Asp Lys Leu Val Asp Glu Ile Gly Val Gly Tyr Val Tyr Asp Ala
 50 55 60
 Gly Ser Val Ser Thr Tyr Cys Leu Ser Gln Ile Lys Pro Leu His Asn
 65 70 75 80
 Phe Leu Thr Gln Leu Gln Pro Phe Leu Lys Leu Lys Gln Lys Gln Ala
 85 90 95
 Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
 100 105 110
 Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
 115 120 125
 Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala

-continued

130	135	140
-----	-----	-----

Val Leu Asp
145

<210> SEQ ID NO 79
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 79

Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe	Val	Asp	Gly	Asp	Gly	Ser
1			5						10					15	
Ile	His	Ala	Cys	Ile	Gln	Pro	Met	Gln	Ser	Met	Lys	Phe	Lys	His	Tyr
			20					25					30		
Leu	His	Leu	Arg	Phe	Thr	Val	His	Gln	Lys	Thr	Gln	Arg	Arg	Trp	Phe
		35					40					45			
Leu	Asp	Lys	Leu	Val	Asp	Glu	Thr	Gly	Val	Gly	Tyr	Val	Tyr	Asp	Ala
	50					55					60				
Gly	Ser	Val	Ser	Thr	Tyr	Cys	Leu	Ser	Gln	Ile	Lys	Pro	Leu	His	Asn
65					70					75					80
Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu	Lys	Gln	Lys	Gln	Ala
				85					90					95	
Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro	Ser	Ala	Lys	Glu	Ser
			100					105						110	
Pro	Asp	Lys	Phe	Leu	Glu	Val	Cys	Thr	Trp	Val	Asp	Gln	Ile	Ala	Ala
		115					120					125			
Leu	Asn	Asp	Ser	Lys	Thr	Arg	Lys	Thr	Thr	Ser	Glu	Thr	Val	Arg	Ala
	130					135						140			

Val Leu Asp
145

<210> SEQ ID NO 80
 <211> LENGTH: 147
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 80

Lys	Glu	Phe	Leu	Leu	Tyr	Leu	Ala	Gly	Phe	Val	Asp	Gly	Asp	Gly	Ser
1			5						10					15	
Ile	Val	Ala	Ser	Ile	Lys	Pro	Gln	Gln	Val	Ala	Lys	Phe	Lys	His	Arg
			20					25					30		
Leu	Met	Leu	Glu	Phe	Tyr	Val	Tyr	Gln	Lys	Thr	Gln	Arg	Arg	Trp	Phe
		35					40					45			
Leu	Asp	Lys	Leu	Val	Asp	Glu	Ile	Gly	Val	Gly	Tyr	Val	Tyr	Asp	Leu
	50					55					60				
Gly	Gly	Ala	Ser	Arg	Tyr	Val	Leu	Ser	Gln	Ile	Lys	Pro	Leu	His	Asn
65					70					75					80
Phe	Leu	Thr	Gln	Leu	Gln	Pro	Phe	Leu	Lys	Leu	Lys	Gln	Lys	Gln	Ala
				85					90					95	
Asn	Leu	Val	Leu	Lys	Ile	Ile	Glu	Gln	Leu	Pro	Ser	Ala	Lys	Glu	Ser
			100					105						110	

-continued

```

Asn Leu Val Leu Lys Ile Ile Glu Gln Leu Pro Ser Ala Lys Glu Ser
      100              105              110

Pro Asp Lys Phe Leu Glu Val Cys Thr Trp Val Asp Gln Ile Ala Ala
      115              120              125

Leu Asn Asp Ser Lys Thr Arg Lys Thr Thr Ser Glu Thr Val Arg Ala
      130              135              140

Val Leu Asp
145

<210> SEQ ID NO 83
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

accggacctc gttgtttaga ct                               22

<210> SEQ ID NO 84
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

tgtttacaca gtgttcatt cc                               22

<210> SEQ ID NO 85
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

gactacacat atagtgtctg tt                               22

<210> SEQ ID NO 86
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

atggacttca agagcaacag tgctgtggcc tggagcaaca aatctgactt tgcattgtgca 60
aacgccttca ac                                          72

<210> SEQ ID NO 87
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 87

atggacttca agagcaacag tgctgtggcc tggagcaaat ctgactttgc atgtgcaaac 60
gccttcaac                                          69

<210> SEQ ID NO 88
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

-continued

<400> SEQUENCE: 88

atggacttca agagcaacaa acaaatctga ctttgcattg gcaaacgcct tcaac 55

<210> SEQ ID NO 89

<211> LENGTH: 67

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 89

atggacttca agagcaacag tgctgtggcc tggagaatct gactttgcat gtgcaaacgc 60

cttcaac 67

<210> SEQ ID NO 90

<211> LENGTH: 65

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 90

atggacttca agagcaacag tgctgtggcc tggagtctga ctttgcattg gcaaacgcct 60

tcaac 65

<210> SEQ ID NO 91

<211> LENGTH: 52

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 91

atggacttca agagcaacaa aatctgactt tgcattgtgca aacgccttca ac 52

<210> SEQ ID NO 92

<211> LENGTH: 70

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 92

atggacttca agagcaacag tgctgtggcc tggagacaaa tctgactttg catgtgcaaa 60

cgcccttcaac 70

<210> SEQ ID NO 93

<211> LENGTH: 68

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 93

atggacttca agagcaacag tgctgtggcc tggagcaatc tgactttgca tgtgcaaacg 60

ccttcaac 68

<210> SEQ ID NO 94

<211> LENGTH: 77

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 94

atggacttca agagcaacag tgctgtggcc tggagcaacg caacaaatct gactttgcat 60

gtgcaaacgc cttcaac 77

<210> SEQ ID NO 95
<211> LENGTH: 76
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 95

atggacttca agagcaacag tgctgtggcc tggagcaaag aacaaatctg actttgcatg 60

tgcaaacgcc ttcaac 76

<210> SEQ ID NO 96
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 96

atggacttca agagcaacag tgctgtggcc tggcaacaaa tctgactttg catgtgcaaa 60

cgctttcaac 70

<210> SEQ ID NO 97
<211> LENGTH: 62
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 97

atggacttca agagcaacag tgctgtggca aatctgactt tgcattgtgca aacgccttca 60

ac 62

<210> SEQ ID NO 98
<211> LENGTH: 68
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 98

atggacttca agagcaacag tgctgtggcc tggacaaatc tgactttgca tgtgcaaacg 60

ccttcaac 68

<210> SEQ ID NO 99
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 99

-continued

atggacttca agagcaacag tgctgtggcc tggagcaaat ctgactttgc atgtgcaaac 60

gccttcaac 69

<210> SEQ ID NO 100
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 100

atggacttca agagcaacag tgctgtggcc tggagcaatg tgcaaacgcc ttcaac 56

<210> SEQ ID NO 101
<211> LENGTH: 69
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 101

atggacttca agagcaacag tgctgtggcc tggaaacaaat ctgactttgc atgtgcaaac 60

gccttcaac 69

<210> SEQ ID NO 102
<211> LENGTH: 71
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 102

atggacttca agagcaacag tgctgtggcc tggagcacaa atctgacttt gcatgtgcaa 60

acgccttcaa c 71

<210> SEQ ID NO 103
<211> LENGTH: 56
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 103

atggacttca agagcaacag tgctgtggcc tggagcaatg tgcaaacgcc ttcaac 56

<210> SEQ ID NO 104
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 104

atcaaatctg actttgcatg tgcaaacgcc ttcaac 36

<210> SEQ ID NO 105
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

-continued

 gtgctgtggc ctggagcaac aaatctgact ttgc 34

<210> SEQ ID NO 106
 <211> LENGTH: 62
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 106

gtgctgtggc ctggagcaag aattcatgcg gccgcaatct agagcaacaa atctgacttt 60

gc 62

<210> SEQ ID NO 107
 <211> LENGTH: 6053
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 107

cagcagctgg cgtaatagcg aagaggcccg caccgatcgc ccttccaac agttgcgag 60

cctgaatggc gaattgaatt ccagacgatt gagcgtcaaa atgtaggat ttccatgagc 120

gtttttctcg ttgcaatggc tggcggtaat attgttctgg atattaccag caaggccgat 180

agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgcgaca 240

acggttaatt tgcgtgatgg acagactctt ttactcggtg gcctcactga ttataaaaac 300

acttctcagg attctggcgt accggttctg tctaaaaatcc ctttaatcgg cctcctgttt 360

agctcccgcg ctgattctaa cgaggaaagc acgttatacg tgctcgtcaa agcaaccata 420

gtacgcgccc tgtagcggcg cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac 480

cgctacactt gccagcgcgc tagcgcgcgc tcctttcgtt ttcttccctt cctttctcgc 540

cacgttcgcc ggctttcccc gtcaagctct aaatcggggg ctccctttag ggttccgatt 600

tagtgcttta cggcacctcg accccaaaaa acttgattag ggtgatggtt cacgtagtgg 660

gccatcgccc tgatagacgg tttttcgcgc tttgacgttg gagtccacgt tctttaatag 720

tggactcttg ttccaaaactg gaacaacact caaccctatc tcggtctatt cttttgattt 780

ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt aacaaaaatt 840

taacgcgaat ttttaaaaaa tattaacggt tacaatttaa atatttgctt atacaatctt 900

cctgtttttg gggcttttct gattatcaac cggggtacat atgattgaca tgctagtttt 960

acgattaccg ttcacgcgcc tgcgcgctcg ctgcctcact gaggccgccc gggcaaaagc 1020

cgggcgctcg gcgaccttg gtcgcccggc ctcaagtgagc gagcgcgagc gcagagaggg 1080

agtgaattc acgcgtggat cttaatagta atcaattacg gggtcattag ttcatagcc 1140

atatatggag ttccgcgcta cataacttac ggtaaatggc ccgcctggct gaccgcccaa 1200

cgacccccgc ccattgacgt caataatgac gtatgttccc atagtaacgc caatagggac 1260

ttccattga cgtaaatggg tggagtattt acggtaaact gcccacttgg cagtacatca 1320

agtgtatcat atgccaagtc cgcctctat tgacgtcaat gacggtaaat ggcccgcctg 1380

gcattatgcc cagtacatga ccttacggga ctttctact tggcagtaca tctacgtatt 1440

agtcacgct attaccatgg tgatgcggtt ttggcagtac accaatgggc gtggatagcg 1500

-continued

gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga gtttgtttg 1560
gcacaaaaat caacgggact ttccaaaatg tcgtaataac cccgcccgt tgacgcaaat 1620
gggcggtagg cgtgtacggt gggagggtcta tataagcaga gctcgttttag tgaaccgtca 1680
gatcactaga agctttctg gacaccccct catctgactt ttttaattcct ccacttcaac 1740
acctggtgca ttcattgtgccc ggcacaaatca gtgattggtg ggtaaatgag tgactgcgtg 1800
agactgactt agtgagctgg gaaagatttt ttggcagaca gggagaaata aggagaggca 1860
acttgagaaa ggggcttaga atgaggccta gaagagcagt aaggggcaaa cagtctgagc 1920
aaaggcaggc aggcaggaac tcagttggag agactgaggc tgggcccagt gccctctcct 1980
gccaccttct cttcatctgc ttttttccc gtcattcttc tggactgcca gaacaaggct 2040
cactgtttct tagtaaaaag agggtttttg tggcaatgga taaggccgag accaccaatc 2100
agaggagttt tagacatcat tgaccagagc tctgggcaga acctggccat tctgaagca 2160
aggaaacagc ctgcgaaggc accaaagctg ccttacctg ggctggggaa gaaggtgtct 2220
tctggaataa tgctgttgtt gaaggcgtt gcacatgcaa agtcagattt gttgctctag 2280
attgcggcgc catgaattct tgctccaggc cacagcactg ttgctcttga agtccataga 2340
cctcatgtct agcacagttt tgtctgtgat atacacatca gaatccttac tttgtgacac 2400
atattgttga gaataaaaaat cgtgaaatag gcagacagac ttgtcactgg atttagagtc 2460
tctcagctgg tacacggcag ggtcagggtt ctggatatct gtgggacaag aggatcaggg 2520
ttaggacatg atctcatttc cctccttgcc ccaaccagg ctggagtcca gatgccagtg 2580
atggacaagg gcggggctct gtggggctgg caagtcacgg tctcatgctt tatacgggaa 2640
atagcatctt agaaaccagc tgctcgtgat ggactgggac tcagggacag gcacaagcta 2700
tcaatcctgg ccaagaggcc atgatttcag tgaacgttca cggccaggcc tggcctgcca 2760
ctcaagggaaa cctgaaatgc agggctactt aataaactg cttattcttt tatttaaatg 2820
gatcttcttc aaaaccccag caatataact ctggcagagt aaaggcaggc atgggaaaaa 2880
ggcccagcaa agcaaaactgt acatcttga atctggagtg gtctcccga cttaggctgg 2940
gcattagcag aatgggaggt ttatggtatg ttggcattaa gttgggaaat ctatcacatt 3000
accaggagat tgctctctca ttgatagagg ttttgaacta taaatcagaa cacctgcgtc 3060
taagccccag cgcaattgtt gttgttaact tgtttattgc agcttataat ggttacaat 3120
aaagcaatag catcacaaat ttcacaaata aagcattttt ttcactgcat tctagttgtg 3180
gtttgtccaa actcatcaat gtatcttaag gcgggaattg atctaggaac ccctagtgat 3240
ggagttggcc actccctctc tgcgcgctcg ctgcctcact gaggccgccc gggcaagcc 3300
cgggcgctcg gcgaccttg gtcgcccggc ctcaagtgagc gagcgagcgc gcagagaggg 3360
agtggccaac ccccccccc cccccccggc gattctcttg tttgctccag actctcaggc 3420
aatgacctga tagcctttgt agagacctct caaaaatagc taccctctcc ggcatgaatt 3480
tatcagctag aacgggtgaa tatcatattg atggtgattt gactgtctcc ggcctttctc 3540
accggtttga atctttacct acacattact caggcattgc atttaaaata tatgagggtt 3600
ctaaaaattt ttatccttgc gttgaaataa aggtctctcc cgcaaaagta ttacagggtc 3660
ataatgtttt tggtaacaac gatttagctt tatgctctga ggctttattg cttaattttg 3720
ctaattcttt gccttgctg tatgatttat tggatgttg aattcctgat gcggtatfff 3780

-continued

ctccttacgc atctgtgogg tatttcacac cgcataatggt gcaactctcag tacaatctgc 3840
tctgatgocg catagttaag ccagccccga caccgcgcaa caccogctga cgcgocctga 3900
cgggcttgtc tgctcccggc atccgcttac agacaagctg tgaccgtctc cgggagctgc 3960
atgtgtcaga ggttttcacc gtcacaccg aaacgcgcga gacgaaaggg cctcgtgata 4020
cgcctatfff tataggtaa tgcacatgata ataatggttt cttagacgtc aggtggcact 4080
tttcggggaa atgtgocgog aacccctatt tgtttatfff tctaaataca ttcaaatatg 4140
tatccgctca tgagacaata accctgataa atgcttcaat aatattgaaa aaggaagagt 4200
atgagtattc aacatttcog tgcgcocctt attccctfff ttgcggcatt ttgccttctt 4260
gtttttgctc acccagaaac gctggtgaaa gtaaaagatg ctgaagatca gttgggtgca 4320
cgagtgggtt acatcgaact ggatctcaac agcggtaaga tccttgagag ttttcgcccc 4380
gaagaacgtt ttccaatgat gagcactfff aaagtctctc tatgtggcgc ggtattatcc 4440
cgtattgacg ccgggcaaga gcaactcogt cgcgcacatc actattctca gaatgacttg 4500
gttgagtact caccagtcac agaaaagcat cttacggatg gcatgacagt aagagaatta 4560
tgcagtgtcg ccataacat gagtgataac actgcggcca acttacttct gacaacgatc 4620
ggaggaccga aggagctaac cgcttttttg cacaacatgg gggatcatgt aactcgcctt 4680
gatcgttggg aaccggagct gaatgaagcc ataccaaaag acgagcgtga caccacgatg 4740
cctgtagcaa tggcaacaac gttgcgaaa ctattaactg gcgaactact tactctagct 4800
tcccggcaac aattaataga ctggatggag gcgataaag ttgcaggacc acttctgcgc 4860
tcggcccttc cggtggctg gtttattgct gataaatctg gagccggtga gcgtgggtct 4920
cgcggtatca ttgcagcact ggggccagat ggtaagccct cccgtatcgt agttatctac 4980
acgacgggga gtcaggcaac tatggatgaa cgaaatagac agatcogtga gataggtgcc 5040
tcaactgatta agcattggta actgtcagac caagtttact catatatact ttagattgat 5100
ttaaaacttc attttaatt taaaaggatc taggtgaaga tcctttttga taatctcatg 5160
acaaaaatcc cttaacgtga gttttcgttc cactgagcgt cagaccccg agaaaagatc 5220
aaaggatctt cttgagatcc tttttttctg cgcgtaatct gctgcttgca aacaaaaaaa 5280
ccaccgctac cagcgggtgt ttgtttgocg gatcaagagc taccaactct tttccgaag 5340
gtaactggct tcagcagagc gcagatacca aatactgtcc ttctagtga gccgtagtta 5400
ggccaccact tcaagaactc ttagcaccg cctacatacc tcgctctgct aatcctgtta 5460
ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg ggttgactc aagacgatag 5520
ttaccggata agcgcagcgc gtcgggctga acggggggtt cgtgcacaca gccagcttg 5580
gagcgaacga cctacaccga actgagatac ctacagcgtg agctatgaga aagcgcacg 5640
cttcccgaag ggagaaaggc ggacaggat ccggtaaagc gcagggtcgg aacaggagag 5700
cgcacgaggg agcttccagg gggaaaagcc tggtatcttt atagtcctgt cgggtttcgc 5760
cacctctgac ttgagcgtcg atttttgtga tgctcgtcag gggggcggag cctatggaaa 5820
aacgccagca acgcggcctt tttacggctt cttggccttt gctggccttt tgctcacatg 5880
ttctttctg cgttatcccc tgattctgtg gataaacgta ttaccgcctt tgagtgagct 5940
gataccgctc gccgcagccg aacgaccgag cgcagcaggt cagtgagcga ggaagcggaa 6000
gagcgcacaa tacgcaaac gcctctcccc gcgcgttggc cgattcatta atg 6053

-continued

```

<210> SEQ ID NO 108
<211> LENGTH: 5458
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 108
cagcagctgg cgtaatagcg aagaggcccg caccgatcgc ccttccaac agttgcgag      60
cctgaatggc gaatggaatt ccagacgatt gagcgtcaaa atgtaggat tccatgagc      120
gttttctctg ttgcaatggc tggcggtaat attgttctgg atattaccag caaggccgat      180
agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgcgaca      240
acggttaatt tgcgtgatgg acagactctt ttactcggtg gcctcactga ttataaaaa      300
acttctcagg attctggcgt accgttctct tctaaaatcc cttaaatcgg cctcctgttt      360
agctcccgct ctgattctaa cgaggaaagc acggtatacg tgctcgtcaa agcaaccata      420
gtacgcgccc tgtagcggcg cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac      480
cgctacactt gccagcgcgc tagcgcgcgc tcctttcgtc ttcttcctt cctttctcgc      540
cacgttcgcc ggctttcccc gtcaagctct aaatcggggg ctccccttag ggttccgatt      600
tagtgcttta cggcacctcg acccaaaaaa acttgattag ggtgatgggt cacgtagtgg      660
gccatcgcgc tgatagacgg tttttcgcgc tttgacgttg gagtccacgt tctttaatag      720
tggactcttg ttccaaactg gaacaacact caaccctatc tcggtctatt cttttgattt      780
ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt aacaaaaatt      840
taacgcgaat ttaacaaaaa tattaacggt tacaatttaa atatttgctt atacaatctt      900
cctgtttttg gggcttttct gattatcaac cggggtacat atgattgaca tgctagtttt      960
acgattacgg ttcacgcgcc tgcgcgctcg ctgcctcact gaggccgcgc gggcaaagcc      1020
cgggcgctcg gcgaccttg gtcgcccggc ctcagtgagc gagcgcgcgc gcagagaggg      1080
agtggaattc acgcgtgctt tctgggcaca cccctcatct gactttttaa ttctccact      1140
tcaacacctg gtgcattcat gtgccggcac aatcagtgat tgggtgggta atgagtgact      1200
gcgtgagact gacttagtga gctgggaaag attttttggc agacagggag aaataaggag      1260
aggcaacttg gagaaggggc ttagaatgag gcctagaaga gcagtaaggg gcaaacagtc      1320
tgagcaaagg caggcaggca ggaactcagt tggagagact gaggctgggc cacgtgccct      1380
ctcctgccac cttctcttca tctgcttttt tcccggtgca ttctctggac tgccagaaca      1440
aggctcactg tttcttagta aaaagagggg tttggtggca atggataagg ccgagaccac      1500
caatcagagg agtttttagac atcattgacc agagctctgg gcagaacctg gccattcctg      1560
aagcaaggaa acagcctcgc aaggcaccaa agctgccctt acctgggctg gggagaaggg      1620
tgtcttctgg aataatgctg ttgttgaagg cgtttgcaca tgcaaagtca gatttgttgc      1680
tctagattgc ggccgcatga attcctgctc caggccacag cactggttct cttgaagtcc      1740
atagacctca tgtctagcac agttttgtct gtgatataca catcagaate cttactttgt      1800
gacacatttg tttgagaate aaaatcggtg aataggcaga cagacttgtc actggattta      1860
gagtccttca gctggtacac ggcagggcga gggttctgga tatctgtggg acaagaggat      1920
cagggtagg acatgatctc atttccctct ttgccccaac ccaggtgga gtccagatgc      1980

```

-continued

cagtgatgga caagggcggg gctctgtggg gctggcaagt cacggtctca tgctttatac 2040
gggaaatagc atcttagaaa ccagctgctc gtgatggact gggactcagg gacaggcaca 2100
agctatcaat cttggccaag aggccatgat ttcagtgaac gttcacggcc aggccctggcc 2160
tgccactcaa ggaacctga aatgcagggc tacttaataa tactgcttat tcttttattt 2220
aataggatct tcttcaaac ccagcaata taactctggc agagtaaagg caggcatggg 2280
aaaaaggccc agcaaagcaa actgtacatc ttggaatctg gagtggctc cccaacttag 2340
gctgggcatt agcagaatgg gaggtttatg gtatgtggc attaagttgg gaaatctatc 2400
acattaccag gagattgctc tctcattgat agaggttttg aactataaat cagaacacct 2460
gcgtctaagc cccagcga aatgtttgt taactgttt attgcagctt ataatggta 2520
caaataaagc aatagcatca caaattcac aaataaagca tttttttcac tgcattctag 2580
ttgtggttg tccaaactca tcaatgtatc ttaaggcggg aattgatcta ggaacccta 2640
gtgatggagt tggcaactcc ctctctgcgc gctcgcctgc tcaactgaggc cgcccgggca 2700
aagcccgggc gtcgggcgac ctttgctgc cggcctcag tgagcgcgc agcgcgcaga 2760
gagggagtgg ccaaccccc ccccccccc cggcgattc tctgtttgc tccagactct 2820
caggcaatga cctgatagcc tttgtagaga cctctcaaaa atagctacc tctccggcat 2880
gaatttatca gctagaacgg ttgaatatca tattgatggg gatttgactg tctccggcct 2940
ttctcaccog tttgaatctt tacctacaca ttactcaggc attgcattta aaatatatga 3000
gggttctaaa aatttttatc cttgcgttga aataaaggct tctcccga aagtattaca 3060
gggtcataat gtttttgta caaccgattt agctttatgc tctgaggctt tattgcttaa 3120
ttttgctaat tctttgcctt gcctgtatga tttattggat gttggaattc ctgatcgggt 3180
atctctctc tacgcctctg tgcggtatct cacaccgcat atggtgcaact ctcagtacaa 3240
tctgctctga tgccgcatag ttaagccagc cccgacacc gccaacacc gctgacgcgc 3300
cctgacgggc ttgtctgctc ccggcatccg cttacagaca agctgtgacc gtctccggga 3360
gctgcatgtg tcagaggttt tcaccgtcat caccgaaacg cgcgagacga aaggcctcgc 3420
tgatacgctt atttttatag gttaatgtca tgataataat ggtttcttag acgtcagggtg 3480
gcacttttgc gggaaatgtg cgcggaacc ctattgttt atttttctaa atacattcaa 3540
atatgtatcc gctcatgaga caataaccct gataaatgct tcaataatat tgaaaagga 3600
agagtatgag tattcaacat ttcctgtctc cctttattcc cttttttgcg gcattttgce 3660
ttcctgtttt tgctcaccga gaaacgctgg tgaaagtaaa agatgctgaa gatcagttgg 3720
gtgcacgagt gggttacatc gaactggatc tcaacagcgg taagatcctt gagagttttc 3780
gccccgaaga acgttttoca atgatgagca cttttaaagt tctgctatgt ggcgcgggat 3840
tatcccgat tgacgcggg caagagcaac tcggtcgcgc catacactat tctcagaatg 3900
acttgggtga gtactcacca gtcacagaaa agcatcttac ggatggcatg acagtaagag 3960
aattatgcag tgctgccata accatgagtg ataacactgc ggccaactta cttctgacaa 4020
cgatcggagg accgaaggag ctaaccgctt ttttgcaaa catgggggat catgtaactc 4080
gccttgatcg ttgggaaccg gagctgaatg aagccatacc aaacgacgag cgtgacacca 4140
cgatgcctgt agcaatggca acaacgttgc gcaactatt aactggcgaa ctacttactc 4200
tagcttcccg gcaacaatta atagactgga tggaggcggg taaagttgca ggaccacttc 4260

-continued

```

tgcgctcggc ccttccggct ggctggttta ttgctgataa atctggagcc ggtgagcgtg 4320
ggtctcggcg tatcattgca gcaactgggc cagatggtaa gccctcccgat atcgtagtta 4380
tctacacgac ggggagtcag gcaactatgg atgaacgaaa tagacagatc gctgagatag 4440
gtgcctcact gattaagcat tggttaactgt cagaccaagt ttactcatat atactttaga 4500
ttgatttaaa acttcatttt taatttaaaa ggatctaggt gaagatcctt tttgataatc 4560
tcatgaccaa aatcccttaa cgtgagtttt cgttccactg agcgtcagac cccgtagaaa 4620
agatcaaagg atcttcttga gatccttttt ttctgcgctg aatctgctgc ttgcaaaaaa 4680
aaaaaccacc gctaccagcg gtggtttgtt tgccggatca agagctacca actctttttc 4740
cgaaggtaac tggttccagc agagcgcaga taccaaatc tgctcttcta gtgtagccgt 4800
agttaggcca ccacttcaag aactctgtag caccgcctac atacctcgtc ctgctaatcc 4860
tgttaccagt ggctgctgcc agtggcgata agtcgtgtct taccgggttg gactcaagac 4920
gatagttacc ggataaggcg cagcggtcgg gctgaacggg gggttcgtgc acacagccca 4980
gcttgaggcg aacgacctac accgaactga gatacctaca gcgtgagcta tgagaaagcg 5040
ccacgcttcc cgaagggaga aagcgggaca ggtatccggt aagcggcagg gtcggaacag 5100
gagagcgcac gagggagcct ccagggggaa acgcctggta tctttatagt cctgtcgggt 5160
ttcgccacct ctgacttgag cgtcgatttt tgtgatgctc gtcagggggg cggagcctat 5220
ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc cttttgctgg ctttttgcctc 5280
acatgttctt tcctcgctta tcccctgatt ctgtggataa ccgtattacc gcctttgagt 5340
gagctgatac cgctcgcgcg agccgaacga ccgagcgcag cgagtcagtg agcggaggaag 5400
cggaaagagcg cccaatacgc aaaccgcctc tccccgcgcg ttggccgatt cattaatg 5458

```

<210> SEQ ID NO 109

<211> LENGTH: 6082

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 109

```

cagcagctgg cgtaatagcg aagaggcccg caccgatcgc ctttcccaac agttgcgcag 60
cctgaatggc gaatggaatt ccagacgatt gagcgtcaaa atgtaggtat ttccatgagc 120
gtttttcctg ttgcaatggc tggcggtaat attgttctgg atattaccag caaggccgat 180
agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgcgaca 240
acggttaatt tgctgtatgg acagactcct ttactcggtg gcctcactga ttataaaaac 300
acttctcagg attctggcgt accggttctg tctaaaatcc ctttaacgg cctcctgttt 360
agctcccgcct ctgattctaa cgaggaaagc acgttatacg tgctcgtcaa agcaaccata 420
gtacgcgccc tgtagcggcg cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac 480
cgctacactt gccagcgccc tagcgcgcgc tccttctcgt ttcttccctt cctttctcgc 540
cacgttcgcc ggetttcccc gtcgaagctct aaatcggggg ctccccttag ggttccgatt 600
tagtgcttta cggcacctcg accccaaaaa acttgattag ggtgatgggt cacgtagtgg 660
gccatcgcgc tgatagacgg ttttctgccc tttgacgttg gagtccacgt tctttaatag 720
tggactcttg ttccaaactg gaacaacact caaccctatc tcggtctatt cttttgattt 780

```

-continued

ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt aacaaaaatt	840
taacgcgaat tttacaacaaa tattaacggt tacaatttaa atatttgctt atacaatctt	900
cctgtttttg gggcttttct gattatcaac cgggggtacat atgattgaca tgctagtttt	960
acgattaccg ttcategccc tgcgcgctcg ctcgctcact gaggcgccc gggcaaagcc	1020
cgggcgctcg gcgaccttg gtcgcccggc ctcagtgagc gagcgagcgc gcagagaggg	1080
agtggaattc acgcgtactg gcatctggac tccagcctgg gttggggcaa agagggaaat	1140
gagatcatgt cctaaccctg atcctcttgt cccacagata tccagaaccc tgaccctgcc	1200
gtgtaccagc tgagagactc taaatccagt gacaagtctg tctgctatt caccgatttt	1260
gattctcaaa caaatgtgtc acaagtaag gattctgatg tgatatcac agacaaaact	1320
gtgctagaca tgaggctat ggacttcaag agcaacagtg ctgtggcctg gagcaactag	1380
tgggcagagc gcacatcgcc cacagtccc gagaagttgg ggggaggggt cggcaattga	1440
accggtgctc agagaaggtg gcgcggggta aactgggaaa gtgatgtcgt gtactggctc	1500
cgctttttc cggaggttg gggagaaccg tatataagtg cagttagtgc cgtgaacgtt	1560
cttttctgca acgggtttgc cgccagaaca cagctagcac catggcgtc ccagtgcag	1620
ccttactttt acctctggcg ttattattgc acgcggctcg tcctgacata cagatgactc	1680
agactacctc ttcctatctc gctcttttag gcgaccgagt aacaatatct tgcggggcca	1740
gccaggacat ctcaaaatac ttaaactggt atcagcagaa gccggacgga acagttaagt	1800
tgctcattta ccacacgtcg agattacact caggcgttcc tagccgattt tcgggttccg	1860
gttccggtac ggactacagc ctgacaatca gtaacctga gcaggaggac atcgccacct	1920
acttctgtca gcagggaac acgctcccgt acacattcgg tgggggaact aagctggaga	1980
ttaccggagg cggttggcagc ggtggcggcg gcagcggggg tggcggctcg gaggtcaagt	2040
tacaggagag cggaccgggc ttggtcgcac ctagccagag cctctcagtc acgtgcactg	2100
tgtctggagt cagtctccca gactacgggg tatcatggat acgacagccg cctagaaagg	2160
gcttagagtg gctgggggtt atctggggaa gtgaaaccac atactacaac tcagctctca	2220
agagccgctc caccatcatt aaggacaaca gtaagtcgca ggttttctta aagatgaact	2280
ctctccagac tgacgacacc gctatttact actgcgcgaa gcactactac tacggcggga	2340
gttacgcaat ggactactgg ggtcagggca cttctgtgac cgatccagc actactacc	2400
cagccccaag tcccccaag ccagctccaa cgatagcaag tcagccctta tctcttcgcc	2460
ctgaggcttg caggcccgcg gcggcggcg ccgttcacac gcgaggacta gacttcgct	2520
gcgacatcta catctgggca ccactagccg ggacttgccg agtgttgttg ttgagcttg	2580
taataacgct ctactgcaag cgtgggagaa agaagctctt gtacattttc aagcagccat	2640
tcatcgctcc cgttcagacg actcaggagg aggacggctg ctcgtgccga ttcccggagg	2700
aggaggaggg cggttgcgaa ctcagagtga agttctctcg ctccgcggac gcaccgctt	2760
accagcaggg tcagaaccag ctatacaacg agttaaacct ggggcgcccg gaggagtacg	2820
acgtgttaga caagcgtaga ggtagggacc cggagatggg aggcaagcct cggagaaaga	2880
acccccagga gggcctgtac aacgaactcc agaaggacaa gatggctgag gcgtactcgg	2940
agattggtat gaagggcgag agacgtcgcg gaaagggaca cgacggctta taccaggggc	3000
tttccaccgc gaccaaggac acatacgacg cgctgcacat gcaagcctta ccacctgat	3060

-continued

gataagatac attgatgagt ttggacaaac cacaactaga atgcagtgaa aaaaatgctt	3120
tatttgtgaa atttgtgatg ctattgcttt atttgaacc attataagct gcaataaaca	3180
agttctagag caacaaatct gactttgcat gtgcaaacgc cttcaacaac agcattattc	3240
cagaagacac cttcttcccc agcccaggta agggcagctt tgggtccttc aattgcctct	3300
ctgcgcgctc gctcgctcac tgaggcggcc cgggcaaacg cggggcgctg ggcgaccttt	3360
ggtcgccccg cctcagtgag cgagcgagcg cgcagagagg gagtggccaa ccccgcgat	3420
tctcttgttt gctccagact ctcaggcaat gacctgatag cctttgtacc tgcaggtctc	3480
aaaaatagct accctctccg gcatgaattt atcagctaga acggttgaat atcatattga	3540
tgggtgattg actgtctccg gcctttctca cccgtttgaa tctttaccta cacattactc	3600
aggcattgca tttaaaatat atgagggttc taaaaatttt tatccttgcg ttgaaataaa	3660
ggcttctccc gcaaaagtat tacagggtca taatgttttt ggtacaaccg atttagcttt	3720
atgctctgag gctttattgc ttaattttgc taattctttg ccttgccctg atgatttatt	3780
ggatgttggg attcctgatg cggtatcttc tccttacgca tctgtgcggt atttcacacc	3840
gcatatggtg cactctcagt acaatctgct ctgatgccgc atagttaagc cagccccgac	3900
accgcaccaac acccgctgac gcgcccctgac gggcttgtct gctcccggca tccgcttaca	3960
gacaagctgt gaccgtctcc gggagctgca tgtgtcagag gttttcaccg tcatcaccga	4020
aacgcgcgag acgaaaagggc ctcgtgatac gcctatcttt ataggttaat gtcataataa	4080
taatggtttc tttagactca ggtggcactt ttcggggaaa tgtgcgcgga acccctattt	4140
gtttatcttt ctaataacat tcaaatatgt atccgctcat gagacaataa cctgataaaa	4200
tgcttcaata atattgaaaa aggaagagta tgagtattca acatttccgt gtcgccctta	4260
ttcccttttt tgccgcatct tgccctctctg tttttgctca cccagaaaac ctggtgaaag	4320
taaaagatgc tgaagatcag ttgggtgcac gagggggta catcgaactg gatctcaaca	4380
gcggtaaagt ccttgagagt tttcgccccg aagaacgttt tccaatgatg agcactttta	4440
aagttctgct atgtggcgcg gtattatccc gtattgacgc cgggcaagag caactcggctc	4500
gccgcataca ctattctcag aatgacttgg ttgagtactc accagtcaca gaaaagcatc	4560
ttacggatgg catgacagta agagaattat gcagtgtctc cataaccatg agtgataaca	4620
ctgcccgaac cttactctctg acaacgatcg gaggaccgaa ggagctaacc gcttttttgc	4680
acaacatggg ggatcatgta actcgccttg atcgttggga accggagctg aatgaagcca	4740
tacaaaacga cgagcgtgac accacgatgc ctgtagcaat ggcaacaacg ttgocgaaac	4800
tattaactgg cgaactactt actctagctt cccggcaaca attaatagac tggatggagg	4860
cggataaagt tgcaggacca cttctgctct cggcccttcc ggctggctgg tttattgctg	4920
ataaatctgg agccggtgag cgtgggtctc gcggtatcat tgcagcactg gggccagatg	4980
gtaagccctc ccgtagctga gttatctaca cgacggggag tcaggcaact atggatgaac	5040
gaaatagaca gatcgtgag ataggtgcct cactgattaa gcattggtaa ctgtcagacc	5100
aagtttactc atatatactt tagattgatt taaaacttca tttttaattt aaaaggatct	5160
aggtgaagat cctttttgat aatctcatga ccaaaatccc ttaacgtgag ttttcgttcc	5220
actgagcgtc agaccccgta gaaaagatca aaggatcttc ttgagatcct tttttctgctc	5280
gcgtaactctg ctgcttgcaa acaaaaaaac caccgctacc agcggtggtt tgtttgocgg	5340

-continued

```

atcaagagct accaactcct ttccgaagg taactggctt cagcagagcg cagataccaa 5400
atactgtcct tctagtgtag ccgtagttag gccaccactt caagaactct gtagcaccgc 5460
ctacatacct cgctctgcta atcctgttac cagtggctgc tgccagtggc gataagtcgt 5520
gtcttaccgg gttggactca agacgatagt taccggataa ggccagcggg tcgggctgaa 5580
cgggggggtc gtgcacacag cccagcttgg agcgaacgac ctacaccgaa ctgagatacc 5640
tacagcgtga gctatgagaa agcggcacgc ttccgaagg gagaaggcg gacaggtatc 5700
cggtaagcgg cagggtcgga acaggagagc gcacgagggg gcttccaggg ggaacgcct 5760
ggtatcttta tagtcctgtc gggtttccg acctctgact tgagcgtcga tttttgtgat 5820
gctcgtcagg gggggcggagc ctatggaaaa acgccagcaa cgccgctttt ttacggttcc 5880
tggccttttg ctggcctttt gctcacatgt tctttctgc gttatccctt gattctgtgg 5940
ataaccgat taccgccttt gagtgagctg ataccgctcg ccgcagccga acgaccgagc 6000
gcagcgagtc agtgagcgag gaagcggaag agcgcccaat acgcaaacgg cctctccccg 6060
cgcgttggcc gattcattaa tg 6082

```

```

<210> SEQ ID NO 110
<211> LENGTH: 7887
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

```

```

<400> SEQUENCE: 110

```

```

cagcagctgg cgtaatagcg aagaggccc caccgatcgc ccttcccaac agttgcgcag 60
cctgaatggc gaatggaatt ccagacgatt gagcgtcaaa atgtaggtat ttccatgagc 120
gtttttcctg ttgcaatggc tggcggtaat attgttctgg atattaccag caaggccgat 180
agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgcgaca 240
acggttaatt tgcgtgatgg acagactcct ttactcgggtg gcctcactga ttataaaaac 300
acttctcagg attctggcgt accgttctctg tctaaaatcc cttaaatcgg cctcctgttt 360
agctcccgct ctgattctaa cgaggaaagc acggtatacg tgctcgtcaa agcaaccata 420
gtacgcgccc tgtagcggcg cattaagcgc ggccgggtgtg gtggttacgc gcagcgtgac 480
cgctacactt gccagcggcc tagcggcccgc tcctttcgtc ttcttccctt cctttctcgc 540
cacgttcgcc ggttttcccc gtaagctct aaatcggggg ctccctttag ggttccgatt 600
tagtgcttta cggcacctcg accccaaaaa acttgattag ggtgatggtt cacgtagtgg 660
gccatcgccc tgatagacgg tttttcgccc tttgacgttg gagtccacgt tctttaatag 720
tggactcttg ttccaaaactg gaacaacact caaccctatc tcggtctatt cttttgattt 780
ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt aacaaaaatt 840
taacgcgaat tttacaaaa tattaacgct tacaatttaa atatttgctt atacaatcct 900
cctgtttttg gggcttttct gattatcaac cggggtacat atgattgaca tgctagtttt 960
acggcgcgcc gggttggcca ctccctctct gcgcgctcgc tcgctcactg aggccggggc 1020
accaaaggtc gcccgacgcc cgggctttgc cggggcggcc tcagtgagcg agegagcgcg 1080
cagagagggg gtggccaact ccatcactag gggttcctac gcgtagatct catattctgg 1140
cagggtcagt ggctccaact aacatttgtt tggtaactta cagtttatta aatagatggt 1200

```

-continued

tatatggaga agctctcatt tctttctcag aagagcctgg cttaggaaggt ggatgaggca 1260
ccatattcat tttgcagggtg aaattcctga gatgtaagga gctgctgtga cttgctcaag 1320
gccttataac aagtaaaccg tagcgtggtg gcttagacgc aggtggtctg atttatagtt 1380
caaaacctct atcaatgaga gagcaatctc ctggtaatgt gatagatttc ccaacttaat 1440
gccaacatac cataaacctc ccattctgct aatgcccagc ctaagttggg gagaccactc 1500
cagattccaa gatgtacagt ttgctttgct gggccttttt cccatgcctg cctttactct 1560
gccagagtta tattgctggg gttttgaaga agatcctatt aaataaaaga ataagcagta 1620
ttattaagta gccctgcatt tcaggtttcc ttgagtggca ggcagggcct ggccgtgaaac 1680
gttactgaa atcatggcct cttggccaag attgatagct tgtgcctgtc cctgagtcce 1740
agtccatcac gagecagctgg tttctaagat gctattttcc gtataaagca tgagaccgtg 1800
acttgccagc cccacagagc cccgccttg tccatcactg gcatctggac tccagcctgg 1860
gttggggcaa agagggaaat gagatcatgt cctaaccctg atcctctgt cccacagata 1920
tccagaaccc tgaccctgcc gtgtaccagc tgagagactc taaatccagt gacaagtctg 1980
tctgcctatt caccgatttt gattctcaaa caaatgtgtc acaaagtaag gattctgatg 2040
tgtatatcac agacaaaact gtgctagaca tgaggtctat ggacttcaag agcaacagtg 2100
ctgtggcctg gagcaactag tggatctgct atcgtctccg tgcccgtcag tgggcagagc 2160
gcacatcgcc cacagtcctc gagaagtgg ggggaggggt cggcaattga acgggtgcct 2220
agagaaggtg ggcgggggta aactgggaaa gtgatgtcgt gtactggctc cgcctttttc 2280
ccgaggggtg gggagaaccg tatataagt cagtagtctc cgtgaacgtt cttttctgca 2340
acgggtttgc cgcagaaca cagctgaagc ttcgaggggc tcgcatctct ccttcaagcg 2400
cccgcctccc tacctgaggc cgcctccac gccggttgag tcgctgtctg cgcctccc 2460
cctgtggtgc ctctgaact gcgtccgctg tctaggtaag tttaaagctc aggtcgagac 2520
cgggcctttg tccggcctc ccttgagcc tacctagact cagccgctc tccacgctt 2580
gcctgacct gcttgcctca ctctacgtct ttgtttcgtt ttctgtctg cgcggttaca 2640
gatccaagct gtgaccggcg cctactctag agctagcga gtcagtgtct ctgacacaac 2700
agtctcgaac ttaactagca ccatggcct cccagtgaca gccttacttt tacctctggc 2760
gttattattg cacgcggctc gtctgacat acagatgact cagactacct cttccctatc 2820
tgcttcttta ggcgaccgag taacaatctc ttgcccggcc agccaggaca tctcaaaata 2880
cttaactgg tatcagcaga agccggacgg aacagttaag ttgctcattt accacacgctc 2940
gagattacac tcaggcgctc ctgaccgatt ttcgggttcc gggtccggta cggactacag 3000
cctgacaatc agtaaccttg agcaggagga catcgccacc tacttctgct agcagggcaa 3060
cacgctccc tacacattcg gtgggggaa taagctggag attacggag gcggtggcag 3120
cgggtggcgc ggcagcggg gtggcggctc ggaggtcaag ttacaggaga gcggaccggg 3180
cttggctgca cctagccaga gcctctcagt cacgtgcact gtgtctggag tcagtctccc 3240
agactacggg gtatcatgga tacgacagcc gcctagaaag ggcttagagt ggctgggggt 3300
tatctgggga agtgaaacca catactaaa ctcaactctc aagagccgctc tcaccatcat 3360
taaggacaac agtaagtgc aggttttctt aaagatgaac tctctccaga ctgacgacac 3420
cgctatttac tactgcgoga agcactacta ctacggcggg agttacgcaa tggactactg 3480

-continued

gggtcagggc	acttctgtga	ccgtatccag	cactactacc	ccagcccac	gtccccccac	3540
gccagctcca	acgatagcaa	gtcagccctt	atctcttcgc	cctgaggctt	gcaggcccgc	3600
ggcggggcgc	gccgttcaca	cgcgaggact	agacttcgcc	tgcgacatct	acatctgggc	3660
accactagcc	gggacttgcg	gagtgttgtt	gttgagcttg	gtaataacgc	tctactgcaa	3720
gcgtgggaga	aagaagctct	tgtacathtt	caagcagcca	ttcatgcgtc	ccgttcagac	3780
gactcaggag	gaggacggct	gctcgtgccg	attcccggag	gaggaggagg	gcggttgcga	3840
actcagagtg	aagttctctc	gctccgcgga	cgcacccgct	taccagcagg	gtcagaacca	3900
gctatacaac	gagttaaac	tggggcgccg	ggaggagtac	gacgtgttag	acaagcgtag	3960
aggtagggac	ccggagatgg	gaggcaagcc	tccgagaaa	aacccccagg	agggcctgta	4020
caacgaactc	cagaaggaca	agatggctga	ggcgtactcg	gagattggta	tgaagggcga	4080
gagacgtcgc	ggaaggggac	acgacggctt	ataccagggg	ctttccaccg	cgaccaagga	4140
cacatacgac	gcgctgcaca	tgcaagcctt	accacctcga	tgaggtacca	gcggccgctt	4200
cgagcagaca	tgataagata	cattgatgag	tttgacaaa	ccacaactag	aatgcagtga	4260
aaaaaatgct	ttatttgtga	aatttgtgat	gctattgctt	tatttgaac	cattataagc	4320
tgcaataaac	aagttaacaa	caacaattcg	aatttaaatc	ggatccgcaa	caaatctgac	4380
tttgcattgtg	caaacgcctt	caacaacagc	attattccag	aagacacctt	cttccccagc	4440
ccaggtaagg	gcagctttgg	tgccctcgca	ggctgtttcc	ttgcttcagg	aatggccagg	4500
ttctgcccag	agctctggtc	aatgatgtct	aaaactcctc	tgattggtgg	tctcggcctt	4560
atccattgoc	accaaaaacc	tctttttact	aagaaacagt	gagccttgtt	ctggcagtc	4620
agagaatgac	acgggaaaaa	agcagatgaa	gagaagggtg	caggagaggg	cacgtggccc	4680
agcctcagtc	tctccaactg	agttcctgcc	tgccctgcctt	tgctcagact	gtttgccect	4740
tactgctctt	ctaggcctca	ttetaagccc	cttctccaag	ttgctctctc	ttatttctcc	4800
ctgtctgcca	aaaaatcttt	cccagctcac	taagtcagtc	tcacgcagtc	actcattaac	4860
ccaccaatca	ctgattgtgc	cggcacatga	atgcaccagg	tggtgaagtg	gaggaattaa	4920
aaagtccagat	gaggggtgtg	cccagaggaa	gcaccattct	agttggggga	gcccactctgt	4980
cagctgggaa	aagtccaaat	aacttcagat	tggaaatgtg	tttaactcag	ggttgagaaa	5040
acagccacct	tcaggacaaa	agtcagggaa	gggctctctg	aagaaatgct	acttgaagat	5100
accagcccta	ccaagggcag	ggagaggacc	aattgatgga	gttggccact	ccctctctgc	5160
gcgctcgctc	gctcactgag	gccgccggg	caaagcccgg	gcgtcgggcg	acctttggtc	5220
gccccgcctc	agtgagcgag	cgagcgcgca	gagagggagt	ggccaacggc	gcgcctgcag	5280
gtctcaaaaa	tagctaccct	ctccggcatg	aatttatcag	ctagaacggt	tgaatatcat	5340
attgatggtg	atttgactgt	ctccggcctt	tctcaccgct	ttgaatcttt	acctacacat	5400
tactcaggoa	ttgcatttaa	aatatatgag	ggttctaaaa	atthttatcc	ttgogttgaa	5460
ataaaggctt	ctcccgcaaa	agtattacag	ggtcataatg	tttttggtac	aacogattta	5520
gctttatgct	ctgaggcttt	attgcttaat	tttgetaatt	ctttgccttg	cctgtatgat	5580
ttattggatg	ttggaattcc	tgatgctgta	ttttctcctt	acgcactctg	gcggatttcc	5640
acaccgcata	tgggtgcactc	tcagtacaat	ctgctctgat	gccgcatagt	taagccagcc	5700
ccgacaccog	ccaacaccog	ctgacgcgcc	ctgacgggct	tgtctgctcc	cggeatccgc	5760

-continued

ttacagacaa gctgtgaccg tctccgggag ctgcatgtgt cagaggtttt cacogtcac 5820
accgaaacgc gcgagacgaa agggcctcgt gatacgccta tttttatagg ttaatgtcat 5880
gataataatg gtttcttaga cgtcaggtgg cacttttcgg ggaaatgtgc gcggaacccc 5940
tatttgttta tttttctaaa tacattcaaa tatgtatccg ctcatgagac aataaccctg 6000
ataaatgctt caataatatt gaaaaaggaa gagtatgagt attcaacatt tccgtgtcgc 6060
ccttattccc ttttttgcgg cattttgcct tectgttttt gctcaccag aaacogctgg 6120
gaaagtaaaa gatgctgaag atcagttggg tgcacgagtg ggttacatcg aactggatct 6180
caacagcggg aagatccttg agagttttcg ccccgaaaga cgttttccaa tgatgagcac 6240
ttttaaagtt ctgcatgtg gcgcggtatt atcccgatt gacgocggg aagagcaact 6300
cggtcgccc atacactatt ctcaaatga cttggttgag tactcaccag tcacagaaaa 6360
gcatcttaog gatggcatga cagtaagaga attatgcagt gctgccataa ccatgagtga 6420
taacactgog gccaaactac tctgacaac gatcggagga ccgaaggagc taaccgcttt 6480
tttgacaaac atgggggatc atgtaactcg cettgatcgt tgggaaccgg agctgaatga 6540
agccatacca aacgacgagc gtgacaccac gatgcctgta gcaatggcaa caacgttgcg 6600
caactatta actggcgaac tacttactct agcttcccgg caacaattaa tagactggat 6660
ggagcgggat aaagttgcag gaccattct gcgctcggcc cttccggctg gctggtttat 6720
tgctgataaa tctggagccg gtgagcgtgg gtctcgcggt atcattgcag cactggggcc 6780
agatggtaag ccctcccga tcgtagttat ctacacgacg gggagtcagg caactatgga 6840
tgaacgaaat agacagatcg ctgagatagg tgcctcactg attaagcatt ggtaactgtc 6900
agaccaagtt tactcatata tactttagat tgatttaaaa cttcattttt aatttaaaag 6960
gatctaggty aagatccttt ttgataatct catgacaaa atccctaac gtgagttttc 7020
gttccactga gcgtcagacc ccgtagaaaa gatcaaaagg tcttcttgag atcctttttt 7080
tctgcgcgta atctgctgct tgcaaaaaaaa aaaaccaccg ctaccagcgg tggtttgttt 7140
gccgatcaa gagctaccaa ctctttttcc gaaggttaact ggcttcagca gagcgcagat 7200
accaaatact gtccttctag tgtagccgta gttaggccac cacttcaaga actctgtagc 7260
accgcctaca tacctcgtc tgctaactct gttaccagtg gctgotgcca gtggcgataa 7320
gtcgtgtctt accgggttgg actcaagacg atagttaccg gataaggcgc agcggctcggg 7380
ctgaacgggg ggttcgtgca cacagcccag cttggagcga acgacctaca ccgaactgag 7440
atacctacag cgtgagctat gagaaagcgc cacgcttccc gaaggagaa agcgggacag 7500
gtatccgta agcggcaggg tcggaacagg agagcgcagc agggagcttc cagggggaaa 7560
cgctcgtat ctttatagtc ctgtcgggtt tcgccacctc tgacttgagc gtcgattttt 7620
gtgatgctcg tcaggggggc ggagcctatg gaaaaacgcc agcaacgcgg cctttttacg 7680
gttctcggcc ttttgctggc cttttgctca catgttcttt cctgogttat cccctgattc 7740
tgtggataac cgtattaccg cctttgagtg agctgatacc gctcgcgca gccgaacgac 7800
cgagcgcagc gagtcagtg gcgaggaagc ggaagagcgc ccaatacgc aaccgcctct 7860
ccccgcgct tggccgattc attaatg 7887

<210> SEQ ID NO 111

<211> LENGTH: 486

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 111

Met Ala Leu Pro Val Thr Ala Leu Leu Leu Pro Leu Ala Leu Leu Leu
 1           5           10           15

His Ala Ala Arg Pro Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu
 20           25           30

Ser Ala Ser Leu Gly Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln
 35           40           45

Asp Ile Ser Lys Tyr Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr
 50           55           60

Val Lys Leu Leu Ile Tyr His Thr Ser Arg Leu His Ser Gly Val Pro
 65           70           75           80

Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile
 85           90           95

Ser Asn Leu Glu Gln Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly
 100          105          110

Asn Thr Leu Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Thr
 115          120          125

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu
 130          135          140

Val Lys Leu Gln Glu Ser Gly Pro Gly Leu Val Ala Pro Ser Gln Ser
 145          150          155          160

Leu Ser Val Thr Cys Thr Val Ser Gly Val Ser Leu Pro Asp Tyr Gly
 165          170          175

Val Ser Trp Ile Arg Gln Pro Pro Arg Lys Gly Leu Glu Trp Leu Gly
 180          185          190

Val Ile Trp Gly Ser Glu Thr Thr Tyr Tyr Asn Ser Ala Leu Lys Ser
 195          200          205

Arg Leu Thr Ile Ile Lys Asp Asn Ser Lys Ser Gln Val Phe Leu Lys
 210          215          220

Met Asn Ser Leu Gln Thr Asp Asp Thr Ala Ile Tyr Tyr Cys Ala Lys
 225          230          235          240

His Tyr Tyr Tyr Gly Gly Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly
 245          250          255

Thr Ser Val Thr Val Ser Ser Thr Thr Thr Pro Ala Pro Arg Pro Pro
 260          265          270

Thr Pro Ala Pro Thr Ile Ala Ser Gln Pro Leu Ser Leu Arg Pro Glu
 275          280          285

Ala Cys Arg Pro Ala Ala Gly Gly Ala Val His Thr Arg Gly Leu Asp
 290          295          300

Phe Ala Cys Asp Ile Tyr Ile Trp Ala Pro Leu Ala Gly Thr Cys Gly
 305          310          315          320

Val Leu Leu Leu Ser Leu Val Ile Thr Leu Tyr Cys Lys Arg Gly Arg
 325          330          335

Lys Lys Leu Leu Tyr Ile Phe Lys Gln Pro Phe Met Arg Pro Val Gln
 340          345          350

Thr Thr Gln Glu Glu Asp Gly Cys Ser Cys Arg Phe Pro Glu Glu Glu
 355          360          365

```

-continued

Glu Gly Gly Cys Glu Leu Arg Val Lys Phe Ser Arg Ser Ala Asp Ala
 370 375 380

Pro Ala Tyr Gln Gln Gly Gln Asn Gln Leu Tyr Asn Glu Leu Asn Leu
 385 390 395 400

Gly Arg Arg Glu Glu Tyr Asp Val Leu Asp Lys Arg Arg Gly Arg Asp
 405 410 415

Pro Glu Met Gly Gly Lys Pro Arg Arg Lys Asn Pro Gln Glu Gly Leu
 420 425 430

Tyr Asn Glu Leu Gln Lys Asp Lys Met Ala Glu Ala Tyr Ser Glu Ile
 435 440 445

Gly Met Lys Gly Glu Arg Arg Arg Gly Lys Gly His Asp Gly Leu Tyr
 450 455 460

Gln Gly Leu Ser Thr Ala Thr Lys Asp Thr Tyr Asp Ala Leu His Met
 465 470 475 480

Gln Ala Leu Pro Pro Arg
 485

<210> SEQ ID NO 112
 <211> LENGTH: 242
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 112

Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
 1 5 10 15

Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Ser Lys Tyr
 20 25 30

Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
 35 40 45

Tyr His Thr Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Gln
 65 70 75 80

Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro Tyr
 85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Thr Gly Gly Gly Ser
 100 105 110

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Val Lys Leu Gln Glu
 115 120 125

Ser Gly Pro Gly Leu Val Ala Pro Ser Gln Ser Leu Ser Val Thr Cys
 130 135 140

Thr Val Ser Gly Val Ser Leu Pro Asp Tyr Gly Val Ser Trp Ile Arg
 145 150 155 160

Gln Pro Pro Arg Lys Gly Leu Glu Trp Leu Gly Val Ile Trp Gly Ser
 165 170 175

Glu Thr Thr Tyr Tyr Asn Ser Ala Leu Lys Ser Arg Leu Thr Ile Ile
 180 185 190

Lys Asp Asn Ser Lys Ser Gln Val Phe Leu Lys Met Asn Ser Leu Gln
 195 200 205

Thr Asp Asp Thr Ala Ile Tyr Tyr Cys Ala Lys His Tyr Tyr Tyr Gly
 210 215 220

-continued

Gly Ser Tyr Ala Met Asp Tyr Trp Gly Gln Gly Thr Ser Val Thr Val
225 230 235 240

Ser Ser

<210> SEQ ID NO 113
 <211> LENGTH: 112
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 113

Arg Val Lys Phe Ser Arg Ser Ala Asp Ala Pro Ala Tyr Gln Gln Gly
1 5 10 15
 Gln Asn Gln Leu Tyr Asn Glu Leu Asn Leu Gly Arg Arg Glu Glu Tyr
20 25 30
 Asp Val Leu Asp Lys Arg Arg Gly Arg Asp Pro Glu Met Gly Gly Lys
35 40 45
 Pro Arg Arg Lys Asn Pro Gln Glu Gly Leu Tyr Asn Glu Leu Gln Lys
50 55 60
 Asp Lys Met Ala Glu Ala Tyr Ser Glu Ile Gly Met Lys Gly Glu Arg
65 70 75 80
 Arg Arg Gly Lys Gly His Asp Gly Leu Tyr Gln Gly Leu Ser Thr Ala
85 90 95
 Thr Lys Asp Thr Tyr Asp Ala Leu His Met Gln Ala Leu Pro Pro Arg
100 105 110

<210> SEQ ID NO 114
 <211> LENGTH: 42
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 114

Lys Arg Gly Arg Lys Lys Leu Leu Tyr Ile Phe Lys Gln Pro Phe Met
1 5 10 15
 Arg Pro Val Gln Thr Thr Gln Glu Glu Asp Gly Cys Ser Cys Arg Phe
20 25 30
 Pro Glu Glu Glu Glu Gly Gly Cys Glu Leu
35 40

<210> SEQ ID NO 115
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 115

Met Ala Leu Pro Val Thr Ala Leu Leu Leu Pro Leu Ala Leu Leu Leu
1 5 10 15
 His Ala Ala Arg Pro
20

<210> SEQ ID NO 116
 <211> LENGTH: 45
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 116

```

Thr Thr Thr Pro Ala Pro Arg Pro Pro Thr Pro Ala Pro Thr Ile Ala
1           5           10           15
Ser Gln Pro Leu Ser Leu Arg Pro Glu Ala Cys Arg Pro Ala Ala Gly
          20           25           30
Gly Ala Val His Thr Arg Gly Leu Asp Phe Ala Cys Asp
          35           40           45

```

<210> SEQ ID NO 117

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 117

```

Ile Tyr Ile Trp Ala Pro Leu Ala Gly Thr Cys Gly Val Leu Leu Leu
1           5           10           15
Ser Leu Val Ile Thr Leu Tyr Cys
          20

```

<210> SEQ ID NO 118

<211> LENGTH: 493

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 118

```

gggcagagcg cacatcgccc acagtccccg agaagttggg gggagggggtc ggcaattgaa      60
cgggtgccta gagaaggtgg cgcggggtaa actgggaaag tgatgctgtg tactggtccc      120
gcctttttcc cgagggtggg ggagaaccgt atataagtgc agtagtcgcc gtgaacgttc      180
tttttcgcaa cgggtttgcc gccagaacac agctgaagct tcgaggggct cgcattcttc      240
cttcacgcgc cgcgccccct acctgaggcc gccatccacg ccggttgagt cgcgttctgc      300
cgctccccgc ctgtggtgcc tctgaactg cgtccgcegt ctaggtaagt ttaaagctca      360
ggtcgagacc gggcctttgt ccggcgctcc cttggagcct acctagactc agccggctct      420
ccacgctttg cctgaccctg cttgctcaac tctacgtctt tgtttcgttt tctggtctgc      480
gccgttacag atc                                                                493

```

<210> SEQ ID NO 119

<211> LENGTH: 2184

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 119

```

gggcagagcg cacatcgccc acagtccccg agaagttggg gggagggggtc ggcaattgaa      60
cgggtgccta gagaaggtgg cgcggggtaa actgggaaag tgatgctgtg tactggtccc      120
gcctttttcc cgagggtggg ggagaaccgt atataagtgc agtagtcgcc gtgaacgttc      180
tttttcgcaa cgggtttgcc gccagaacac agctgaagct tcgaggggct cgcattcttc      240

```

-continued

```

cttcacgcgc cgcgcgcct acctgaggcc gccatccacg ccggttgagt cgcgttctgc 300
cgcctcccgc ctgtggtgcc tctgaactg cgtccgcctg ctaggtaagt ttaaagctca 360
ggtcgagacc gggcctttgt cggcgctcc cttggagcct acctagactc agcggctct 420
ccacgctttg cctgaccctg cttgctcaac tctacgtctt tgtttcgttt tctgttctgc 480
gccgttacag atccaagctg tgaccggcgc ctactctaga gctagcgcag tcagtgett 540
tgacacaaca gtctcgaact taactagcac catggcgctc ccagtgcagc ccttactttt 600
acctctggcg ttattattgc acgcggtctg tcttgacata cagatgactc agactactc 660
ttccctatct gcttctttag gcgaccgagt aacaatatct tgccgggcca gccaggacat 720
ctcaaaatac ttaaactggt atcagcagaa gccggacgga acagttaagt tgctcattta 780
ccacacgtcg agattacact caggcgttcc tagccgattt tcgggttccg gttccggtac 840
ggactacagc ctgacaatca gtaacctga gcaggaggac atcgccacct acttctgtca 900
gcagggcaac acgtcccggt acacattcgg tgggggaaact aagctggaga ttaccggagg 960
cgggtggcagc ggtggcggcg gcagcggggg tggcggctcg gaggtcaagt tacaggagag 1020
cggaccgggc ttggtcgcac ctgaccagag cctctcagtc acgtgcactg tgtctggagt 1080
cagttcccga gactacgggg tatcatggat acgacagcgg cctagaaaagg gcttagagt 1140
gctgggggtt atctggggaa gtgaaaccac atactacaac tcagctctca agagccgcct 1200
caccatcatt aaggacaaca gtaagtcgca ggttttctta aagatgaact ctctccagac 1260
tgacgacacc gctatttact actgcgcgaa gcactactac tacggcggga gttacgcaat 1320
ggactactgg ggtcagggca cttctgtgac cgtatccagc actactacc cagcccacg 1380
tccccccacg ccagctccaa cgatagcaag tcagccctta tctcttcgcc ctgaggcttg 1440
caggcccgcg gcggcgcgcg ccgttcacac gcgaggacta gacttcgctt gcgacatcta 1500
catctgggca ccactagccg ggacttgccg agtgttgttg ttgagcttgg taataacgct 1560
ctactgcaag cgtgggagaa agaagctctt gtacatttcc aagcagccat tcatgctcc 1620
cgttcagaag actcaggagg aggacggctg ctcgtgccga ttcccggagg aggaggagg 1680
cggttgcgaa ctcagagtga agttctctcg ctccgaggac gcacccgctt accagcagg 1740
tcagaaccag ctatacaacg agttaaacct gggcgccggg gaggagtacg acgtgttaga 1800
caagcgtaga ggtagggacc cggagatggg aggcaagcct cggagaaaaga acccccagga 1860
gggcctgtac aacgaactcc agaaggacaa gatggctgag gcgtactcgg agattggtat 1920
gaaggcgag agacgtcgcg gaaagggaca gcacggctta taccaggggc ttccaccgc 1980
gaccaaggac acatacgagc cgtgcacat gcaagcctta ccacctcgat gaggtaccag 2040
cggccgcttc gagcagacat gataagatac attgatgagt ttggacaaac cacaactaga 2100
atgcagtgaa aaaaatgctt tatttgtgaa atttgtgatg ctattgcttt atttgtaacc 2160
attataagct gcaataaaca agtt 2184

```

<210> SEQ ID NO 120

<211> LENGTH: 6811

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 120

-continued

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt	60
ctgtctgcct attcaccgat tttgattctc aaacaaatgt gtcacaaagt aaggattctg	120
atgtgtatat cacagacaaa actgtgctag acatgaggtc tatggacttc aagagcaaca	180
gtgctgtggc ctggagcaag ggcagagcgc acatcgccca cagtccccga gaagttgggg	240
ggaggggtcg gcaattgaac gggtgccctag agaaggtggc gcggggtaaa ctgggaaagt	300
gatgtcgtgt actggctcgc cctttttccc gaggggtggg gagaaccgta tataagtgca	360
gtagtgcgcg tgaacgttct ttttcgcaac gggtttgccg ccagaacaca gctgaagctt	420
cgaggggctc gcactctctc ttcacgcgcc cgccgcccta cctgaggccg ccatccacgc	480
cggttgagtc gcgttctgcc gctcccgcgc tgtggtgcct cctgaactgc gtcgcgcgtc	540
taggtaagtt taaagctcag gtcgagaccg ggcctttgtc cggcgctccc ttggagccta	600
cctagactca gccggctctc cacgctttgc ctgaccctgc ttgctcaact ctacgtcttt	660
gtttcgtttt ctgttctgcg ccgttacaga tccaagctgt gaccggcgcc tactctagag	720
ctagcgcagt cagtgttctt gacacaacag tctcgaactt aactagcacc atggcgctcc	780
cagtgcagc cttactttta cctctggcgt tattattgca cgcggtcgt cctgacatac	840
agatgactca gactacctct tccctatctg cttcttttag cgaccgagta acaatatctt	900
gccgggccag ccaggacatc tcaaaatact taaactggta tcagcagaag ccggacggaa	960
cagttaagtt gctcatttac cacacgtcga gattacactc aggcgttccct agccgatttt	1020
cggttcccg ttcgggtacg gactacagcc tgacaatcag taaccttgag caggaggaca	1080
tgcacaccta cttctgtcag cagggcaaca cgctcccgtc cacattcggg gggggaacta	1140
agctggagat tacccggaggc ggtggcagcg gtggcggcgg cagcgggggt ggcggctcgg	1200
aggtcaagtt acaggagagc ggaccgggct tggtcgcacc tagccagagc ctctcagtca	1260
cgtgcactgt gtctggagtc agtctcccag actacggggt atcatggata cgacagccgc	1320
ctagaaaggg cttagagtgg ctggggggtta tctggggaag tgaaccaca tactacaact	1380
cagctctcaa gagccgcctc accatcatta aggacaacag taagtgcag gttttcttaa	1440
agatgaactc tctccagact gacgacaccg ctatttacta ctgcgcgaag cactactact	1500
acggcgggag ttacgcaatg gactactggg gtcagggcac ttctgtgacc gtatccagca	1560
ctactacccc agcccccagc cccccacgc cagctccaac gatagcaagt cagcccttat	1620
ctcttcgccc tgaggcttgc aggcccgcg cgggcggcgc cgttcacacg cgaggactag	1680
acttcgcctg cgacatctac atctgggcac cactagccgg gacttgcgga gtgttgtgt	1740
tgagcttggg aataacgctc tactgcaagc gtgggagaaa gaagctcttg tacattttca	1800
agcagccatt catgctccc gttcagacga ctcaggagga ggaccgctgc tcgtgccgat	1860
tcccggagga ggaggagggc ggttgcgcaac tcagagtga gttctctcgc tcccgggacg	1920
caccgcctta ccagcagggc cagaaccagc tataacaaga gttaaacctg gggcgccggg	1980
aggagtacga cgtgttagac aagcgtagag gtagggaccc ggagatggga ggcaagcctc	2040
ggagaaagaa cccccaggag ggcctgtaca acgaaactca gaaggacaag atggctgagg	2100
cgtactcgga gattggtatg aagggcgaga gacgtcgcgg aaaggacac gacggcttat	2160
accaggggct ttccaccgcg accaaggaca catacgagc gctgcacatg caagccttac	2220
cacctcgatg aggtaccagc ggcgcttcg agcagacatg ataagataca ttgatgagtt	2280

-continued

tggacaaacc	acaactagaa	tgcaagtgaaa	aaaatgcttt	atgtgtgaaa	ttgtgatgc	2340
tattgcttta	tttgaacca	ttataagctg	caataaacia	gttcaaatct	gactttgcat	2400
gtgcaaacgc	cttcaacaac	agcattatct	cagaagacac	cttcttcccc	agcccaggta	2460
agggcagctt	tggtgccttc	gcaggctgtt	tccttgcctc	aggaatggcc	aggttctgcc	2520
cagagctctg	gtcaatgatg	tctaaaactc	ctctgattgg	tggtctcggc	cttatccatt	2580
gccacaaaa	ccctcttttt	actaagaaac	agtgagcctt	gttctggcag	tccagagaat	2640
gacacgggaa	aaaagcagat	gaagagaagg	tggcaggaga	gggcacgtgg	cccagcctca	2700
gtctctccaa	ctgagtctct	gcctgcctgc	ctttgctcag	actgtttgcc	ccttactgct	2760
cttctaggcc	tcattctaa	cccctctctc	aagttgcctc	tccttatttc	tccttctctg	2820
ccaaaaaatc	tttcccagct	cactaagtca	gtctcacgca	gtcactcatt	aaccacccaa	2880
tcactgattg	tgccggcaca	tgaatgcacc	aggtgttgaa	gtggaggaa	taaaaagtca	2940
gatgaggggt	gtgcccagag	gaagcaccat	tctagtggg	ggagcccatc	tgtcagctgg	3000
gaaaagtcca	aataacttca	gattggaatg	tgttttaact	cagggttgag	aaaacagcta	3060
ccttcaggac	aaaagtcagg	gaagggctct	ctgaagaaat	gctacttgaa	gataccagcc	3120
ctaccaaggg	cagggagagg	accctataga	ggcctgggac	aggagctcaa	tgagaaagga	3180
gaagagcagc	aggcatgagt	tgaatgaagg	aggcagggcc	gggtcacagg	gccttctagg	3240
ccatgagagg	gtagacagta	ttctaaggac	gccagaaagc	tggtgatcgg	cttcaagcag	3300
gggagggaca	cctaatttgc	ttttcttttt	tttttttttt	tttttttttt	tttttgagat	3360
ggagttttgc	tcttgttgcc	caggctggag	tgcaatggtg	catcttggct	actgcaacc	3420
tccgcctccc	aggttcaagt	gattctcctg	cctcagcctc	ccgagtagct	gagattacag	3480
gcacccgcc	ccatgcctgg	ctaatttttt	gtatttttag	tagagacagg	gtttcactat	3540
gttgccagg	ctggtctoga	actcctgacc	tcaggtgatc	caccogcttc	agcctcccaa	3600
agtgtcggga	ttacagcgct	gagccaccac	accggcctg	cttttcttaa	agatcaatct	3660
gagtgcctgta	cgagagtggt	gttgtaagcc	aagagtagaa	gcagaaaggg	agcagttgca	3720
gcagagagat	gatggaggcc	tgggcagggt	ggtggcaggg	aggtaaccaa	caccattcag	3780
gtttcaaagg	tagaacctatg	cagggatgag	aaagcaaaga	gggatcaag	gaaggcagct	3840
ggattttggc	ctgagcagct	gagtcaatga	tagtgccgtt	tactaagaag	aaaccaagga	3900
aaaaatttgg	ggtgcaggga	tcaaaacttt	ttggaacata	tgaaagtacg	tgtttatact	3960
ctttatggcc	cttgtcacta	tgtatgcctc	gctgcctcca	ttggactcta	gaatgaagcc	4020
aggcaagagc	agggtctatg	tgtgatggca	catgtggcca	gggtcatgca	acatgtactt	4080
tgtacaaaca	gtgtatattg	agtaaataga	aatggtgtcc	aggagccgag	gtatcggctc	4140
tgccagggcc	aggggctctc	cctagcaggt	gctcatatgc	tgtaagttcc	ctccagatct	4200
ctccacaagg	aggcatggaa	aggctgtagt	tgttcacctg	cccaagaact	aggaggtctg	4260
gggtgggaga	gtcagcctgc	tctggatgct	gaaagaatgt	ctgtttttcc	ttttagaaa	4320
ttcctgtgat	gtcaagctgg	tcgagaaaag	ctttgaaaca	ggtaagacag	gggtctagcc	4380
tgggtttgca	caggattgctg	gaagtgatga	accgcgaata	accctgcctg	gatgagggag	4440
tgggaagaaa	ttagtagatg	tgggaatgaa	tgatgaggaa	tggaaacagc	ggttcaagac	4500
ctgccagag	ctgggtgggg	tctctcctga	atccctctca	ccatctctga	ctttccattc	4560

-continued

taagcacttt gaggatgagt ttctagcttc aatagaccaa ggactctctc ctaggcctct	4620
gtattccttt caacagctcc actgtcaaga gagccagaga gagcttctgg gtggcccagc	4680
tgtgaaatth ctgagtcocct tagggatagc cctaaacgaa ccagatcctc ctgaggacag	4740
ccaagaggtt ttgccttctt tcaagacaag caacagtact cacataggct gtgggcaatg	4800
gtcctgtctc tcaagaatcc cctgccactc ctccacccca cctgggccc atattcattt	4860
ccatttgagt tgttcttatt gagtcctct tctctgtgta gcggaactca ctaaggggcc	4920
catctggaoc cgaggtattg tgatgataaa ttctgagcac ctacccctc cccagaaggg	4980
ctcagaaata aaataagagc caagtctagt cgggtgttcc tgtcttgaaa cacaatactg	5040
ttggccctgg aagaatgcac agaactctgt tgtaagggga tatgcacaga agctgcaagg	5100
gacaggaggt gcaggagctg caggcctccc ccaccagcc tgctctgctc tggggaaaac	5160
cgtgggtgtg tctctcagcc catgcaggcc tgggacatgc aagcccataa ccgctgtggc	5220
ctcttggttt tacagatacg aacctaaact ttcaaaacct gtcagtgatt gggttccgaa	5280
tctcctcct gaaagtggcc gggtttaact tgctcatgac gctcggctg tgggtccagct	5340
gaggtgaggg gccttgaagc tgggagtggg gtttagggac gcgggtctct ggggtgcacc	5400
taagctctga gagcaaacct cctgcaggg tcttgctttt aagtccaaag cctgagccca	5460
ccaaactctc ctacttcttc ctgttcaaaa ttctctctgt gcaataataa tggcctgaaa	5520
cgctgtaaaa tatectcatt tcagccgctc cagttgcact tctcccctat gaggtaggaa	5580
gaacagttgt ttgaaaaaga aaaaactgag gcccacagc taatgagtgg aggaagagag	5640
acacttgtgt acaccacatg ccttgtgtg tacttctctc accgtgtaac ctctcatgt	5700
cctctctccc cagtaacgct ctcttagctc agtagaaaga agacattaca ctcatattac	5760
acccaatcc tggetagagt ctccgacccc tctccccca gggccccag tegtcttget	5820
gacaactgca tctgtttcca tcaccatcaa aaaaaactc caggctgggt gcgggggctc	5880
acacctgtaa tcccagcact ttgggaggca gaggcaggag gagcacagga gctggagacc	5940
agcctgggca acacagggag accccgctc tacaaaaagt gaaaaatta accaggtgtg	6000
gtgctgcaca cctgtagtcc cagctactta agaggctgag atgggaggat cgcttgagcc	6060
ctggaatggt gaggtacaaa tgagctgtga ttgctcact gcactccagc ctggaagaca	6120
aagcaagatc ctgtctcaaa taataaaaa aataagaact ccagggtaca tttgctccta	6180
gaactctacc acatagcccc aaacagagcc atcaccatca catccctaac agtctgggt	6240
cttctcaggt gtccagctg acttctgttc tctctcttc cagatctgca agattgtaag	6300
acagcctgtg ctccctcgtc ccttctctg cattgccctt cttctccctc tccaaacaga	6360
gggaactctc ctacccccaa ggaggtgaaa gctgctacca cctctgtgcc cccccgcaa	6420
tgccaccaac tggatcttac ccgaatttat gattaagatt gctgaagagc tgccaaacac	6480
tgctgccacc ccctctgttc ccttattgct gcttgcact gcctgacatt caeggcagag	6540
gcaaggctgc tgcagcctcc cctggtgtg cacattccct cctgctcccc agagactgcc	6600
tccgcatcc cacagatgat ggatctcag tgggttctct tgggctctag gtcctgcaga	6660
atggtgtgag gggtttattt ttttttaata gtgttcataa agaaatacat agtattctc	6720
ttctcaagac gtgggggaa attatctcat tatcgaggcc ctgctatgct gtgtatctgg	6780
gcgtgttgta tgcctgctg ccgatgcctt c	6811

-continued

<210> SEQ ID NO 121
<211> LENGTH: 6811
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 121

atatacagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt	60
ctgtctgcct attcaccgat tttgattctc aaacaaatgt gtcacgggca gagcgacat	120
cgcccacagt ccccgagaag ttggggggag gggtcggcaa ttgaacgggt gcctagagaa	180
ggtagggcgg ggtaaacctgg gaaagtgatg tegtgtactg gctccgcctt tttcccgagg	240
gtgggggaga accgtatata agtgcagtag tcgccgtgaa cgttcttttt cgcaacgggt	300
ttgccgccag aacacagctg aagccttcgag gggctcgcct ctctcctca cgcgcccgcc	360
gccctacctg aggcgcgcat ccacgcccgt tgagtgcgct tctgccgctt cccgctgtg	420
gtgcctcctg aactgcgtcc gccgtctagg taagttaaa gctcaggtcg agaccgggcc	480
tttgtccggc gctcccttgg agcctacctt gactcagccg gctctccacg ctttgctga	540
ccctgcttgc tcaactctac gtctttgttt cgttttctgt tctgcccgct tacagatcca	600
agctgtgacc ggcgcctact ctagagctag cgcagtcagt gcttctgaca caacagtctc	660
gaacttaact agcaccatgg cgtcctcagc gacagcctta cttttacctc tggcgttatt	720
attgcacgag gctcgtcctg acatacagat gactcagact acctcttccc tatctgcttc	780
tttaggcgac cgagtaacaa tatcttgccg ggcacgcccag gacatctcaa aatactaaa	840
ctggtatcag cagaagccgg acggaacagt taagttgctc atttaccaca cgtcagagatt	900
aacctcaggg gttcctagcc gattttcggg ttccgggtcc ggtacggact acagcctgac	960
aatcagtaac cttgagcagg aggacatcgc cacctacttc tgtcagcagg gcaacacgct	1020
cccgtacaca ttccggtggg gaactaagct ggagattacc ggaggcgggtg gcagcgggtg	1080
cggcggcagc ggggggtggc gctcggaggt caagttacag gagagcggac cgggcttgggt	1140
cgcacctagc cagagcctct cagtcactg cactgtgtct ggagtcagtc tcccagacta	1200
cggggatca tggatacagc agccgcctag aaaggcctta gagtggctgg gggttatctg	1260
gggaagttaa accacatact acaactcagc tctcaagagc cgcctacca tcattaagga	1320
caacagtaag tcgcaggttt tcttaagat gaactctctc cagactgacg acaccgctat	1380
ttactactgc gcgaagcact actactacgg cgggagttac gcaatggact actggggcca	1440
gggcacttct gtgaccgtat ccagcactac taccgccgcc ccacgtcccc ccacgccagc	1500
tccaacgata gcaagtcagc ccttatctct tcgccctgag gcttgacggc ccgcgccggg	1560
cggcgcgctt cacacgcgag gactagactt cgcctgcgac atctacatct gggcaccact	1620
agccgggact tgcggagtgt tgttgttgag cttgtaata acgctctact gcaagcgtgg	1680
gagaaagaag ctcttgtaca ttttaagca gccattcatg cgtcccgttc agacgactca	1740
ggaggaggac ggctgctcgt gccgattccc ggaggaggag gagggcgggt gcaactcag	1800
agtgaagttc tctcgtccg cggacgcacc cgcttaccag cagggtcaga accagctata	1860
caacgagtta aacctggggc gccgggagga gtacgacgtg ttagacaagc gtagaggtag	1920
ggacccggag atgggaggca agcctcggag aaagaacccc caggagggcc tgtacaacga	1980

-continued

actccagaag gacaagatgg ctgaggcgta ctcgagagatt ggtatgaagg gcgagagacg	2040
tgcgcgaaag ggacacgacg gcttatacca ggggctttcc accgcgacca aggacacata	2100
cgacgcgctg cacatgcaag ccttaccacc tcgatgaggt accagcgccc gcttcgagca	2160
gacatgataa gatacattga tgagtttga caaaccacaa ctagaatgca gtgaaaaaaa	2220
tgctttattt gtgaaatttg tgatgctatt gctttatttg taaccattat aagctgcaat	2280
aaacaagtta aagtaaggat tctgatgtgt atateacaga caaaactgtg ctagacatga	2340
ggtctatgga cttcaagagc aacagtgtg tggcctggag caacaaatct gactttgcat	2400
gtgcaaacgc cttcaacaac agcattatc cagaagacac cttcttcccc agcccaggta	2460
agggcagctt tggtgcttc gcaggctgtt tccttgcttc aggaatggcc aggttctgcc	2520
cagagctctg gtcaatgatg tctaaaactc ctctgattgg tggctctggc cttatccatt	2580
gccacccaaa ccctcttttt actaagaaac agtgagcctt gttctggcag tccagagaat	2640
gacacgggaa aaaagcagat gaagagaagg tggcaggaga gggcacgtgg cccagctca	2700
gtctctcaa ctgagttcct gctgcctgc ctttgctcag actgtttgcc ccttactgct	2760
cttctaggcc tcattctaag ccccttctcc aagttgcctc tccttatttc tcctgtctg	2820
ccaaaaaate tttcccagct cactaagtca gtctcacgca gtcactcatt aaccaccaa	2880
tcactgattg tgccggcaca tgaatgcacc aggtgtttaa gtggaggaat taaaagtca	2940
gatgaggggt gtgccagag gaagcaccat tctagttggg ggagcccac tgtcagctgg	3000
gaaaagtcca aataacttca gattggaatg tgttttaact cagggttgag aaaacagcta	3060
ccttcaggac aaaagtcagg gaaggctct ctgaagaaat gctacttgaa gataccagcc	3120
ctaccaaggg cagggagagg accctataga ggcctgggac aggagctcaa tgagaaagga	3180
gaagagcagc aggcgatgag tgaatgaagg aggcagggcc gggtcacagg gccttctagg	3240
ccatgagagg gtagacagta ttctaaggac gccagaaagc tgttgatcgg cttcaagcag	3300
gggagggaca cctaatttgc tttcttttt ttttttttt ttttttttt tttttgagat	3360
ggagttttgc tcttgttgcc caggctggag tgcaatggtg catcttggct cactgcaacc	3420
tccgcctccc aggttcaagt gattctctg cctcagcctc ccgagtagct gagattacag	3480
gcacccgccca ccatgcctgg ctaatttttt gtatttttag tagagacagg gtttcactat	3540
gttgccagg ctggtctoga actcctgacc tcaggtgatc caccgcttc agcctcccaa	3600
agtctgsgga ttacaggcgt gagccaccac acccggcctg ctttcttaa agatcaatct	3660
gagtctgta cggagagtg gttgtaagcc aagagtagaa gcagaaaggg agcagttgca	3720
gcagagagat gatggaggcc tgggcagggt ggtggcaggg aggtaacca caccattcag	3780
gtttcaaagg tagaacatg cagggatgag aaagcaaga ggggatcaag gaaggcagct	3840
ggattttggc ctgagcagct gagtcaatga tagtgccgtt tactaagaag aaaccaagga	3900
aaaaatttgg ggtgcaggga tcaaaacttt ttggaacata tgaaagtacg tgtttatact	3960
ctttatggcc cttgtcacta tgtatgctc gctgcctcca ttggactcta gaatgaagcc	4020
aggcaagagc agggctctat tgtgatggca catgtggcca gggctatgca acatgtactt	4080
tgtacaaaca gtgtatattg agtaaataga aatggtgtcc aggagccgag gtatcggctc	4140
tgccagggcc aggggctctc cctagcaggt gctcatatgc tgtaagtcc ctccagatct	4200
ctccacaagg aggcattgaa aggctgtagt tgttcacctg cccaagaact aggaggtctg	4260

-continued

gggtgggaga gtcagcctgc tctggatgct gaaagaatgt ctgtttttcc ttttagaaaag	4320
ttcctgtgat gtcagctgg tcgagaaaag ctttgaaaca ggtaagacag gggcttagcc	4380
tgggtttgca caggattgcg gaagtgatga acccgcaata accctgcctg gatgaggag	4440
tgggaagaaa ttagtagatg tgggaatgaa tgatgaggaa tggaaacagc ggttcaagac	4500
ctgccagag ctgggtggg tctctctga atccctctca ccatctctga ctttccattc	4560
taagcacttt gaggatgagt tcttagcttc aatagacca ggactctctc ctaggcctct	4620
gtattccttt caacagctcc actgtcaaga gagccagaga gagcttctgg gtggcccagc	4680
tgtgaaatct ctgagtcct tagggatagc cctaaacgaa ccagatcatc ctgaggacag	4740
ccaagaggtt ttgcttctt tcaagacaag caacagtact cacataggct gtgggcaatg	4800
gtcctgtctc tcaagaatcc cctgccactc ctcacacca ccctgggccc atattcattt	4860
ccatttgagt tgttcttatt gagtcctct tctctgtgta gcggaactca ctaaggggcc	4920
catctggacc cgaggattg tgatgataaa ttctgagcac ctacccctc cccagaaggg	4980
ctcagaaata aaataagagc caagtctagt cgggtttcc tgtcttgaaa cacaactctg	5040
ttggccctgg aagaatgcac agaactctgt tgtaagggga tatgcacaga agctgcaagg	5100
gacaggaggt gcaggagctg caggcctccc ccaccagcc tgetctgct tggggaaaac	5160
cgtgggtgtg tcctgcaggc catgcaggcc tgggacatgc aagcccataa ccgctgtggc	5220
ctcttggttt tacagatagc aacctaaact tcaaaaact gtcagtatt gggttccgaa	5280
tcctcctct gaaagtggcc gggtttaac tgetcatgac gctgaggctg tggctcagct	5340
gaggtgaggg gccttgaagc tgggagtggg gtttagggac gcgggtctct ggggtcatcc	5400
taagctctga gagcaaacct ccctgcaggg tcttgccttt aagtccaaag cctgagccca	5460
ccaaactctc ctactctctc ctgttataaa ttcctctgt gcaataataa tggcctgaaa	5520
cgctgtaaaa taccctcatt tcagccgct cagttgact tctcccctat gaggtaggaa	5580
gaacagttgt ttgaaaacga agaaactgag gcccacagc taatgagtgg aggaagagag	5640
acactgtgt acaccacatg ccttgtgtg tacttctctc accgtgtaac ctctcatgt	5700
cctctctccc cagtacggct ctcttagctc agtagaaga agacattaca ctcatattac	5760
acccaatcc tggctagagt ctccgaccc tcctcccaca gggtocccag tegtcttct	5820
gacaactgca tcctgttoca tcaccatcaa aaaaaaactc caggctgggt gcgggggctc	5880
acacctgtaa tcccagcact ttgggaggca gaggcaggag gagcacagga gctggagacc	5940
agcctgggca acacaggag accccgctc tacaaaaagt gaaaaatta accaggtgtg	6000
gtgctgcaca cctgtagtcc cagctactta agaggctgag atgggaggat cgtttgagcc	6060
ctggaatgtt gaggtacaa tgagctgtga ttgcgtcact gactccagc ctggaagaca	6120
aagcaagatc ctgtctcaaa taataaaaaa aataagaact ccagggtaca tttgctccta	6180
gaactctacc acatagcccc aaacagagcc atcaccatca catecctaac agtctgggt	6240
cttctcagt gtccagcctg acttctgttc ttcctcattc cagatctgca agattgtaag	6300
acagcctgtg ctccctcgt ccttctctg cattgcccct cttctccctc tccaaacaga	6360
gggaactctc ctaccccaa ggagtgaaa gctgctacca cctctgtgcc cccccgcaa	6420
tgccaccaac tggatctac ccgaatttat gattaagatt gctgaagagc tgccaaacac	6480
tgetgccacc ccctctgttc ccttattgct gctgtcact gcctgacatt cacggcagag	6540

-continued

```

gcaaggctgc tgcagcctcc cctggetgtg cacattccct cctgctcccc agagactgcc 6600
tccgccatcc cacagatgat ggatcttcag tgggttctct tgggctctag gtctgcaga 6660
atggtgtgag gggtttattt tttttaata gtgttcataa agaaatacat agtattcttc 6720
ttctcaagac gtggggggaa attatctcat tatcgaggcc ctgctatgct gtgtatctgg 6780
gcgtggtgta tgtctgctg ccgatgcct c 6811

```

<210> SEQ ID NO 122

<211> LENGTH: 6811

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 122

```

atatccagaa ccctgaccct gccgtgtacc agctgagaga ctctaaatcc agtgacaagt 60
ctgtctgcct attcaccgat tttgattctc aaacaaatgt gtcacaaagt aaggattctg 120
atgtgtatat gggcagagcg cacatcgccc acagtccccg agaagttggg gggaggggtc 180
ggcaattgaa cgggtgccta gagaaggtgg cgcggggtaa actgggaaag tgatgtcgtg 240
tactggctcc gcctttttcc cgagggtggg ggagaaccgt atataagtgc agtagtcgcc 300
gtgaacgttc tttttcgcaa cgggtttgcc gccagaacac agctgaagct tcgaggggct 360
cgcctctctc cttcacgcgc ccgccgccct acctgaggcc gccatccacg ccggttgagt 420
cgcgttctgc cgctcccgcc ctgtggtgcc tcttgaactg cgtccgccgt ctaggtaagt 480
ttaaagctca ggtcgagacc gggcctttgt ccggcgcctc cttggagcct acctagactc 540
agccggctct ccacgctttg cctgaccctg cttgctcaac tctacgtctt tgtttcgttt 600
tctgttctgc gccgttacag atccaagctg tgaccggcgc ctactctaga gctagcgcag 660
tcagtgtctc tgacacaaca gtctcgaact taactagcac catggcgctc ccagtgcag 720
ccttactttt acctctggcg ttattattgc acgcggtctg tcttgacata cagatgactc 780
agactacctc ttcctatctc gcttctttag gcgaccgagt aacaatatct tgccgggcca 840
gccaggacat ctcaaaatac ttaaactggt atcagcagaa gccggacgga acagttaagt 900
tgctcattta ccacacgtcg agattacact caggcggtcc tagccgattt tcgggttccg 960
gttcgggtac ggactacagc ctgacaatca gtaacctga gcaggaggac atcgccacct 1020
acttctgtca gcagggcaac acgctcccgt acacattcgg tgggggaact aagctggaga 1080
ttaccggagg cgggtggcagc ggtggcggcg gcagcggggg tggcggctcg gaggtcaagt 1140
tacaggagag cggaccgggc ttggtcgcac ctagccagag cctctcagtc acgtgcactg 1200
tgtctggagt cagtctccca gactacgggg tatcatggat acgacagccg cctagaaagg 1260
gcttagagtg gctggggggt atctggggaa gtgaaaccac atactacaac tcagctctca 1320
agagccgcct caccatcatt aaggacaaca gtaagtcgca ggttttctta aagatgaact 1380
ctctccagac tgacagacc gctatttact actgcgcgaa gcaactactac tacggcggga 1440
gttacgcaat ggactactgg ggtcagggca cttctgtgac cgtatccagc actactacc 1500
cagccccacg tccccccacg ccagctccaa cgatagcaag tcagccctta tctcttcgcc 1560
ctgaggcttg caggcccgcg gcgggcggcg ccgttcacac gcgaggacta gacttcgcct 1620
gcgacateta catctgggca ccactagccg ggacttgagg agtggtgttg ttgagcttg 1680

```

-continued

taataacgct	ctactgcaag	cgtgggagaa	agaagctctt	gtacattttc	aagcagccat	1740
tcatgcgctc	cgttcagacg	actcaggagg	aggacggctg	ctcgtgccga	ttcccggagg	1800
aggaggaggg	cggttgcgaa	ctcagagtga	agttctctcg	ctccgcgga	gcacccgctt	1860
accagcaggg	tcagaaccag	ctatacaacg	agttaaacct	ggggcgccgg	gaggagtacg	1920
acgtgttaga	caagcgtaga	ggtagggacc	cggagatggg	aggcaagcct	cggagaaaga	1980
acccccagga	gggectgtac	aacgaactcc	agaaggacaa	gatggctgag	gcgtactcgg	2040
agattggtat	gaagggcgag	agacgtcgcg	gaaagggaca	cgacggctta	taccaggggc	2100
ttccaccgc	gaccaaggac	acatacgacg	cgctgcacat	gcaagcctta	ccacctcgat	2160
gaggtaccag	cggccgcttc	gagcagacat	gataagatac	attgatgagt	ttggacaaac	2220
cacaactaga	atgcagtgaa	aaaaatgctt	tatttgtaa	atttgatgag	ctattgcttt	2280
attingtaacc	attataagct	gcaataaaca	agttcacaga	caaaactgtg	ctagacatga	2340
ggtctatgga	cttcaagagc	aacagtgtcg	tggcctggag	caacaaatct	gactttgcat	2400
gtgcaaacgc	cttcaacaac	agcattattc	cagaagacac	cttctcccc	agcccaggta	2460
agggcagctt	tgggtgcctc	gcaggctgtt	tccttgcttc	aggaatggcc	aggttctgcc	2520
cagagctctg	gtcaatgatg	tctaaaactc	ctctgattgg	tggctctcggc	cttatccatt	2580
gccacaaaa	ccctcttttt	actaagaaac	agtgagcctt	gttctggcag	tccagagaat	2640
gacacgggaa	aaaagcagat	gaagagaagg	tggcaggaga	gggcacgtgg	cccagcctca	2700
gtctctccaa	ctgagttcct	gctgcctgc	ctttgctcag	actgtttgcc	ccttactgct	2760
cttctaggcc	tcattctaag	cccctctctc	aagttgcctc	tccttatttc	tcctgtctg	2820
ccaaaaaatc	tttcccagct	cactaagtca	gtctcacgca	gtcactcatt	aaccaccaa	2880
tactctgattg	tgcgggcaca	tgaatgcacc	aggtgttgaa	gtggagggaat	taaaaagtca	2940
gatgaggggt	gtgcccagag	gaagcaccat	tctagttggg	ggagcccatc	tgtoagctgg	3000
gaaaagtcca	aataacttca	gattggaatg	tgttttaact	cagggttgag	aaaacagcta	3060
ccttcaggac	aaaagtccag	gaagggctct	ctgaagaaat	gctacttgaa	gataccagcc	3120
ctaccaaggg	cagggagagg	accctataga	ggcctgggac	aggagctcaa	tgagaaagga	3180
gaagagcagc	aggcatgagt	tgaatgaagg	aggcagggcc	gggtcacagg	gccttctag	3240
ccatgagagg	gtagacagta	ttctaaggac	gccagaaagc	tgttgatcgg	cttcaagcag	3300
gggagggaca	cctaatttgc	ttttcttttt	tttttttttt	tttttttttt	tttttgagat	3360
ggagttttgc	tcttgttgcc	caggctggag	tgcaatggtg	catcttgget	cactgcaacc	3420
tccgcctccc	aggttcaagt	gattctctcg	cctcagcctc	ccgagtagct	gagattacag	3480
gcacccgcca	ccatgcctgg	ctaatttttt	gtatttttag	tagagacagg	gtttcactat	3540
gttgccagg	ctggtctoga	actcctgacc	tcaggtgatc	caccgcttc	agcctcccaa	3600
agtgtcggga	ttacaggcgt	gagccaccac	accggcctg	cttttcttaa	agatcaatct	3660
gagtgcctgta	cggagagtgg	gttgtaagcc	aagagtagaa	gcagaaaggg	agcagttgca	3720
gcagagagat	gatggaggcc	tgggcaggg	ggtggcaggg	aggtaaccaa	caccattcag	3780
gtttcaaagg	tagaacctag	cagggatgag	aaagcaaga	gggatcaag	gaaggcagct	3840
ggattttggc	ctgagcagct	gagtcaatga	tagtgccgtt	tactaagaag	aaaccaagga	3900
aaaaatttgg	ggtgcagggga	tcaaaacttt	ttggaacata	tgaaagtacg	tgtttatact	3960

-continued

ctttatggcc	ctgtcacta	tgtatgctc	gctgcctcca	ttggactcta	gaatgaagcc	4020
aggcaagagc	agggtctatg	tgtgatggca	catgtggcca	gggtcatgca	acatgtactt	4080
tgtacaaaca	gtgtatatg	agtaaataga	aatgggtgoc	aggagccgag	gtatcggtcc	4140
tgccagggcc	aggggctctc	cctagcaggt	gctcatatgc	tgtaagttcc	ctccagatct	4200
ctccacaagg	aggcatggaa	aggctgtagt	tgttcacctg	cccaagaact	aggaggtctg	4260
gggtgggaga	gtcagcctgc	tctggatgct	gaaagaatgt	ctgtttttcc	ttttagaaag	4320
ttcctgtgat	gtcaagctgg	tcgagaaaag	ctttgaaaca	ggtaagacag	gggtctagcc	4380
tgggtttgca	caggattgcg	gaagtgatga	acccgcaata	accctgcctg	gatgaggggag	4440
tgggaagaaa	ttagtagatg	tgggaatgaa	tgatgaggaa	tggaaacagc	ggttcaagac	4500
ctgccagag	ctgggtgggg	tctctcctga	atccctctca	ccatctctga	ctttccattc	4560
taagcacttt	gaggatgagt	ttctagcttc	aatagaccaa	ggactctctc	ctaggcctct	4620
gtattccttt	caacagctcc	actgtcaaga	gagccagaga	gagcttctgg	gtggcccagc	4680
tgtgaaattt	ctgagtcctt	tagggatagc	cctaaacgaa	ccagatcatc	ctgaggacag	4740
ccaagagggt	ttgccttctt	tcaagacaag	caacagtact	cacataggct	gtgggcaatg	4800
gtcctgtctc	tcaagaatcc	cctgccactc	ctcacacca	ccctgggccc	atattcattt	4860
ccatttgagt	tgttcttatt	gagtcacctt	tctctgtgta	gcggaactca	ctaaggggcc	4920
catctggacc	cgaggatatt	tgatgataaa	ttctgagcac	ctaccccatc	cccagaaggg	4980
ctcagaaata	aaataagagc	caagtctagt	cggtgtttcc	tgtcttgaaa	cacaatactg	5040
ttggccctgg	aagaatgcac	agaatctggt	tgtaagggga	tatgcacaga	agctgcaagg	5100
gacaggaggt	gcaggagctg	caggcctccc	ccaccacagc	tgctctgcct	tggggaaaac	5160
cgtgggtgtg	tcctgcaggc	catgcaggcc	tgggacatgc	aagcccataa	ccgctgtggc	5220
ctcttggttt	tacagatagc	aacctaaact	ttcaaaacct	gtcagtgatt	gggttccgaa	5280
tcctcctcct	gaaagtggcc	gggtttaatc	tgctcatgac	gctgcggctg	tggtccagct	5340
gagggtgaggg	gccttgaagc	tgggagtggg	gtttagggac	gcggtctctc	gggtgcatcc	5400
taagctctga	gagcaaacct	ccctgcaggg	tcttgctttt	aagtccaaag	cctgagccca	5460
ccaaactctc	ctactctctc	ctgttacaaa	ttcctctgtg	gcaataataa	tggcctgaaa	5520
cgctgtaaaa	tatcctcatt	tcagccgctt	cagttgcact	tctcccctat	gaggtaggaa	5580
gaacagttgt	ttagaaaacga	agaaactgag	gccccacagc	taatgagtg	aggaagagag	5640
acacttggtg	acaccacatg	ccttggtgtg	tactctctc	accgtgtaac	ctcctcatgt	5700
cctctctccc	cagtacggct	ctcttagctc	agtagaaaga	agacattaca	ctcatattac	5760
accccaatcc	tggctagagt	ctccgcaccc	tcctccccc	gggtcccag	tcgtcttget	5820
gacaactgca	tcctgttcca	tcaccatcaa	aaaaaaaa	caggctgggt	gcgggggctc	5880
acacctgtaa	tcccagcact	tggggaggca	gaggcaggag	gagcacagga	gctggagacc	5940
agcctgggca	acacagggag	accccgcctc	tacaaaaagt	gaaaaaatta	accaggtgtg	6000
gtgctgcaca	cctgtagtcc	cagctactta	agaggtgag	atgggaggat	cgcttgagcc	6060
ctggaatggt	gaggctacaa	tgagctgtga	ttgcgtcact	gcactccagc	ctggaagaca	6120
aagcaagatc	ctgtctcaaa	taataaaaa	aataagaact	ccaggttaca	tttgctccta	6180
gaactctacc	acatagcccc	aaacagagcc	atcaccatca	catccctaac	agtcctgggt	6240

-continued

```

cttctcagc gtcacgctg acttctgtc ttcctcattc cagatctgca agattgtaag 6300
acagcctgtg ctccctcgct ccttctcttg cattgcccct cttctccctc tccaaacaga 6360
gggaactctc ctacccccaa ggaggtgaaa gctgctacca cctctgtgcc cccccggcaa 6420
tgccaccaac tggatcctac ccgaatttat gattaagatt gctgaagagc tgccaaacac 6480
tgctgccacc ccctctgttc ccttattgct gcttgtcact gcctgacatt cacggcagag 6540
gcaaggctgc tgcagcctcc cctggctgtg cacattccct cctgctcccc agagactgcc 6600
tccgccatcc cacagatgat ggatctcag tgggttctct tgggctctag gtcctgcaga 6660
atggtgtgag gggtttattt ttttttaata gtgttcataa agaaatacat agtattcttc 6720
ttctcaagac gtggggggaa attatctcat taccgaggcc ctgctatgct gtgtatctgg 6780
gcgtgttgta tgcctctgtg ccgatgcctt c 6811

```

<210> SEQ ID NO 123

<211> LENGTH: 6040

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 123

```

cagcagctgg cgtaatagcg aagaggcccg caccgatcgc cttcccaac agttgctgag 60
cctgaatggc gaatggaatt ccagacgatt gagcgtcaaa atgtaggtat ttccatgagc 120
gtttttctctg ttgcaatggc tggcggtaat attgttcttg atattaccag caaggccgat 180
agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgctgaca 240
acggttaatt tgcgtgatgg acagactctt ttactcggtg gcctcactga ttataaaaac 300
acttctcagg attctggcgt accgttctctg tctaaaaatcc ctttaactgg cctcctgttt 360
agctcccgcct ctgattctaa cgaggaaagc acgttatacg tgctcgtcaa agcaaccata 420
gtacgcgccc tgtagcggcg cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac 480
cgctacactt gccagcgcct tagcgcgcctg tcccttctctg ttcttccctt cctttctctg 540
cacgttcgcc ggctttcccc gtcaagctct aaatcggggg ctcctttag ggttccgatt 600
tagtgcttta cggcacctcg accccaaaaa acttgattag ggtgatggtt cacgtagtgg 660
gccatcgcct tgatagacgg tttttcgcct tttgacgttg gagtccacgt tctttaatag 720
tggactcttg tccaaaactg gaacaacact caaccctatc tcggtctatt cttttgattt 780
ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt acaaaaaatt 840
taacgcgaat ttttaaaaaa tattaacgct tacaatttaa atatttgctt atacaatctt 900
cctgtttttg gggcttttct gattatcaac cggggtacat atgattgaca tgctagtttt 960
acgattaccg ttcacgcctc tgcgcgctcg ctcgctcact gaggccgccc gggcaaaagc 1020
cgggcgctcg gcgacctttg gtcgcccggc ctcagtgagc gacgagaggg 1080
agtgaatcc acgctgata tccagaacct tgaccctgcc gtgtaccagc tgagagactc 1140
taaatccagt gacaagtctg tctgcctatt caccgatttt gattctcaaa caaatgtgtc 1200
acaaagtaag gattctgatg tgcatatcac agacaaaact gtgctagaca tgaggtctat 1260
ggacttcaag agcaacagtg ctgtggcctg gagcaactag tgggaggag tagggcggag 1320
ccaatcagcg tgcgcgcttc cgaaggttgc cttttatggc tgggaggaga atgggagggtg 1380

```

-continued

aacgccgatg	attatataag	gacgcgccgg	gtgtggcaca	gctagttccg	tcgcagccgg	1440
gatttgggtc	gcggttcttg	tttgttccgg	aaagccacca	tggcgctccc	agtgacagcc	1500
ttacttttac	ctctggcggt	attattgcac	gcggtctgtc	ctgacataca	gatgactcag	1560
actacctctt	ccctatctgc	ttctttaggc	gaccgagtaa	caatatcttg	ccgggccagc	1620
caggacatct	caaaataactt	aaactgggat	cagcagaagc	cggacggaac	agttaagttg	1680
ctcatttacc	acacgtcgag	attacactca	ggcgttccta	gccgatttcc	gggttccggt	1740
tccggtacgg	actacagcct	gacaatcagt	aacctgagc	aggaggacat	cgccacctac	1800
ttctgtcagc	agggcaacac	gctcccgtac	acattcggtg	ggggaactaa	gctggagatt	1860
accggaggcg	gtggcagcgg	tggcgccggc	agcgggggtg	gcggtctcga	ggtcaagtta	1920
caggagagcg	gaccgggctt	ggtcgcacct	agccagagcc	tctcagtcac	gtgcaactgtg	1980
tctggagtca	gtctcccaga	ctacggggta	tcatggatac	gacagccgcc	tagaaagggc	2040
ttagagtggc	tgggggttat	ctggggaagt	gaaaccacat	actacaactc	agctctcaag	2100
agccgcctca	ccatcattaa	ggacaacagt	aagtcgcagg	ttttcttaa	gatgaactct	2160
ctccagactg	acgacaccgc	tatttactac	tgcgcgaagc	actactacta	cggcgggagt	2220
tacgcaatgg	actactgggg	tcagggcact	tctgtgacgg	tatccagcac	tactacccca	2280
gccccacgtc	ccccacgccc	agctccaacg	atagcaagtc	agcccttacc	tcttcgcctt	2340
gaggcttgca	ggcccgcggc	ggcgccggcc	gttcacacgc	gaggactaga	cttcgcctgc	2400
gacatctaca	tctgggcacc	actagccggg	acttgcggag	tggtgtgtgt	gagcttggtg	2460
ataacgctct	actgcaagcg	tgggagaaag	aagctctgtt	acattttcaa	gcagccattc	2520
atgcttccc	ttcagacgac	tcaggaggag	gacggctgct	cggtccgatt	cccggaggag	2580
gaggaggggc	gttgcgaact	cagagtgaag	ttctctcgct	ccgcggacgc	accgccttac	2640
cagcagggtc	agaaccagct	atacaacgag	ttaaacctgg	ggcgcgggga	ggagtaacgac	2700
gtgttagaca	agcgtagagg	tagggaccgg	gagatgggag	gcaagcctcg	gagaaagaac	2760
ccccaggagg	gcctgtacaa	cgaactccag	aaggacaaga	tggctgaggg	gtactcggag	2820
attggtatga	agggcgagag	acgtccggga	aaggacacg	acggcttata	ccaggggctt	2880
tccaccgcga	ccaaggacac	atacgcgcgg	ctgcacatgc	aagccttacc	acctcgatga	2940
taagatacat	tgatgagttt	ggacaaaacca	caactagaat	gcagtgaaaa	aaatgcttta	3000
tttgtgaaat	ttgtgatgct	attgctttat	ttgtaacct	tataagctgc	aataaacaag	3060
ttctagagca	acaaatctga	ctttgcatgt	gcaaacgcct	tcaacaacag	cattattcca	3120
gaagacacct	tcttccccag	cccaggtaag	ggcagctttg	gtgccttcgc	aggctgtttc	3180
cttgcttcag	gaatggccag	gttctgccc	gagctctggt	caatgatgtc	taaaactcct	3240
ctgattgcaa	ttgcctctct	gcgcgctcgc	tcgctcaactg	aggccgcgcc	ggcaaagccc	3300
gggcgtcggg	cgacctttgg	tcgcccggcc	tcagtgagcg	agcgagcgcg	cagagagggg	3360
gtggccaacc	ccggcgatcc	tcttgtttgc	tccagactct	caggcaatga	cctgatagcc	3420
tttgtacctg	caggtctcaa	aaatagctac	cctctccggc	atgaatttat	cagctagaac	3480
ggttgaatat	catattgatg	gtgatttgac	tgtctccggc	ctttctcacc	cgtttgaatc	3540
tttaacctaca	cattactcag	gcattgcatt	taaaatatat	gagggttcta	aaaattttta	3600
tccttgcggt	gaaataaagg	cttctcccgc	aaaagtatta	cagggtcata	atgtttttgg	3660

-continued

tacaaccgat	ttagctttat	gctctgagge	tttatgtctt	aattttgcta	attctttgcc	3720
ttgcctgtat	gatttattgg	atgttggaat	tctgatgceg	gtattttctc	cttacgcac	3780
tgtgcggat	ttcacaccgc	atatggtgca	ctctcagtac	aatctgctct	gatgccgcat	3840
agttaagcca	gccccgacac	ccgccaacac	ccgctgacgc	gccctgacgg	gcttgtctgc	3900
tcccggcatc	cgcttacaga	caagctgtga	ccgtctccgg	gagctgcatg	tgctcagaggt	3960
tttcaccgtc	atcacccaaa	cgcgcgagac	gaaagggcct	cgtgatacgc	ctatttttat	4020
aggtaaatgt	catgataata	atggtttctt	agacgtcagg	tggcactttt	cggggaaatg	4080
tgcgcggaac	ccctatttgt	ttatttttct	aaatacattc	aaatatgtat	ccgctcatga	4140
gacaataaacc	ctgataaatg	cttcaataat	attgaaaaag	gaagagtatg	agtattcaac	4200
atctccgtgt	cgcccttatt	cccttttttg	cggcattttg	ccttctgttt	tttgcctacc	4260
cagaaacgct	ggtgaaagta	aaagatgctg	aagatcagtt	gggtgcacga	gtgggttaca	4320
tcgaactgga	tctcaacagc	ggttaagatcc	ttgagagttt	tcgccccgaa	gaacgttttc	4380
caatgatgag	cacttttaaa	gttctgctat	gtggcgcggt	attatcccgt	attgacgccg	4440
ggcaagagca	actcggctgc	cgcatacact	attctcagaa	tgacttggtt	gagtaactcac	4500
cagtccacaga	aaagcatcct	acggatggca	tgacagtaag	agaattatgc	agtgtgtcca	4560
taacatgag	tgataaacact	cgcggccaact	tacttctgac	aacgatcggg	ggaccgaagg	4620
agctaacccg	ttttttgac	aacatggggg	atcatgtaac	tcgccttgat	cgttgggaa	4680
cggagctgaa	tgaagccata	ccaaacgacg	agcgtgacac	cacgatgctt	gtagcaatgg	4740
caacaacggt	cgcgaaacta	ttaactggcg	aactacttac	tctagcttcc	cgcaacaat	4800
taatagactg	gatggaggcg	gataaagttg	caggaccact	tctgcgctcg	gcccttccgg	4860
ctggtgggtt	tattgctgat	aaatctggag	ccggtgagcg	tgggtctcgc	ggtatcattg	4920
cagcactggg	gccagatggt	aagccctccc	gtatcgtagt	tatctacacg	acggggagtc	4980
aggcaactat	ggatgaacga	aatagacaga	tcgctgagat	aggtgcctca	ctgattaagc	5040
attggttaact	gtcagaccaa	gtttactcat	atatacttta	gattgattta	aaacttcatt	5100
tttaatttaa	aaggatctag	gtgaagatcc	tttttgataa	tctcatgacc	aaaatccctt	5160
aacgtgagtt	ttcgttccac	tgagcgtcag	accccgtaga	aaagatcaaa	ggatcttctt	5220
gagatccctt	ttttctgcgc	gtaatctgct	gcttgcaaac	aaaaaaaa	ccgctaccag	5280
cggtgggttg	tttgccgat	caagagctac	caactctttt	tccgaaggta	actggcttca	5340
gcagagcgca	gataccaaat	actgtccttc	tagtgtagcc	gtagttaggc	caccacttca	5400
agaactctgt	agcaccgctt	acatacctcg	ctctgcta	cctgttacca	gtggctgctg	5460
ccagtggcga	taagtctgtt	cttaccgggt	tggactcaag	acgatagtta	ccggataagg	5520
cgcagcggtc	gggctgaacg	gggggttcgt	gcacacagcc	cagcttggag	cgaacgacct	5580
acaccgaact	gagataccta	cagcgtgagc	tatgagaaag	cgccacgctt	cccgaaggga	5640
gaaaaggcga	caggatcccg	gtaagcggca	gggtcggaac	aggagagcgc	acgaggggagc	5700
ttccaggggg	aaacgcctgg	tatctttata	gtcctgtcgc	gtttcgccac	ctctgacttg	5760
agcgtcgatt	tttgtgatgc	tcgtcagggg	ggcggagcct	atggaaaaac	gccagcaacg	5820
cggccttttt	acggttctcg	gccttttgct	ggccttttgc	tcacatgttc	tttctcgcgt	5880
tatcccctga	ttctgtggat	aaccgtatta	ccgccttga	gtgagctgat	accgctcgcc	5940

-continued

```
gcagccgaac gaccgagcgc agcgagtcag tgagcggagga agcgggaagag cgcccaatac 6000
gcaaaccgcc tctcccgcgc cgttgggcca ttcattaatg 6040
```

```
<210> SEQ ID NO 124
<211> LENGTH: 8342
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthesized
```

```
<400> SEQUENCE: 124
```

```
cagcagctgg cgtaatagcg aagaggcccg caccgatcgc ccttccaac agttgcgag 60
cctgaatggc gaatggaatt ccagacgatt gagcgtcaaa atgtaggtat ttccatgagc 120
gtttttctcg ttgcaatggc tggcggtaat attgttctgg atattaccag caaggccgat 180
agtttgagtt cttctactca ggcaagtgat gttattacta atcaaagaag tattgcgaca 240
acggttaatt tgcgtgatgg acagactctt ttactcggtg gcctcaactga ttataaaaa 300
acttctcagg attctggcgt accggttctg tctaaaatcc cttaaatcgg cctcctgttt 360
agctcccgcct ctgattctaa cgaggaaagc acggtatacg tgctcgtcaa agcaaccata 420
gtacgcgccc tgtagcggcg cattaagcgc ggcgggtgtg gtggttacgc gcagcgtgac 480
cgctacactt gccagcgcgc tagcgcgcgc tcctttcgtc ttcttccctt cctttctcgc 540
cacgttcgcc ggctttcccc gtcaagctct aaatcggggg ctccccttag ggttccgatt 600
tagtgcttta cggcacctcg accccaaaaa acttgattag ggtgatgggt cacgtagtgg 660
gccatcgcgc tgatagacgg tttttcgcgc ttgacggtg gagtccacgt tctttaatag 720
tggactcttg ttccaaactg gaacaacact caacctatc tcggtctatt cttttgattt 780
ataagggatt ttgccgattt cggcctattg gttaaaaaat gagctgattt aacaaaaatt 840
taacgcgaat tttaacaaaa tattaacggt tacaatttaa atatttgctt atacaatctt 900
cctgtttttg gggcttttct gattatcaac cggggtacat atgattgaca tgctagtttt 960
acggcgcgccc ggggtggcca ctcccctctc gcgcgctcgc tcgctcactg aggcggggcg 1020
accaaaggtc gcccgacgccc cgggctttgc ccgggcccgc tcagtgagcg agcgagcgcg 1080
cagagaggga gtggccaact ccatacactag gggttcctac gcgtagatct catattctgg 1140
cagggtcagt ggtcccaact aacatttgtt tggtaactta cagtttatta aatagatggt 1200
tatatggaga agctctcatt tctttctcag aagagcctgg ctaggaaggt ggatgaggca 1260
ccatattcat tttgcaggtg aaattcctga gatgtaagga gctgctgtga cttgctcaag 1320
gccttataac gagtaaacgg tagcgtggg gcttagacgc aggtgttctg atttatagtt 1380
caaacctct atcaatgaga gagcaatctc ctggaatgt gatagattc ccaacttaat 1440
gccaacatac cataaacctc ccattctgct aatgccagc ctaagttggg gagaccactc 1500
cagattccaa gatgtacagt ttgctttgct gggccttttt cccatgectg cctttactct 1560
gccagagtta tattgctggg gttttgaaga agatcctatt aaataaaaga ataagcagta 1620
ttattaagta gccctgcatt tcaggtttcc ttgagtggca ggccaggcct ggccgtgaac 1680
gttactgaa atcatggcct cttggccaag attgatagct tgtgctgtc cctgagtccc 1740
agtccatcac gagcagctgg tttctaagat gctatttccc gtataaagca tgagaccgtg 1800
acttccagc cccacagagc cccgccttg tccatcactg gcatctggac tccagcctgg 1860
```

-continued

gttggggcaa	agagggaaat	gagatcatgt	cctaaccctg	atcctcttgt	cccacagata	1920
tccagaaccc	tgaccctgcc	gtgtaccagc	tgagagactc	taaatccagt	gacaagtctg	1980
tctgcctatt	caccgatttt	gattctcaaa	caaatgtgtc	acaaagtaag	gattctgatg	2040
tgtatatac	agacaaaact	gtgctagaca	tgaggtctat	ggacttcaag	agcaacagtg	2100
ctgtggcctg	gagcaactag	tcaatattgg	ccattagcca	tattattcat	tggttatata	2160
gcataaatca	atattggcta	ttggccattg	catacgttgt	atctatatca	taatatgtac	2220
atztatattg	gctcatgtcc	aatatgaccg	ccatgttggc	attgattatt	gaccagttat	2280
taatagtaat	caattacggg	gtcattagtt	catagcccat	atatggagtt	ccgcggtaca	2340
taacttacgg	taaatggccc	gcttggtgta	ccgcccacg	acccccgcc	attgacgtca	2400
ataatgacgt	atgttcccat	agtaacgcca	atagggactt	tccattgacg	tcaatgggtg	2460
gagtatttac	ggtaaaactg	ccacttggca	gtacatcaag	tgtatcataa	tccaagtccg	2520
ccccctattg	acgtcaatga	cggtaaatgg	cccgcctggc	attatgcccc	gtacatgacc	2580
ttacgggact	ttcctacttg	gcagtaacac	tacgtattag	tcatcgctat	taccatgatg	2640
atgdcggttt	ggcagtacac	caatgggctg	ggatagcggg	ttgactcacc	gggatttcca	2700
agtctccacc	ccattgacgt	caatgggagt	ttgttttggc	acaaaaatca	acgggacttt	2760
ccaaaatgtc	gtaataaccc	cgccccgttg	acgcaaatgg	gcggtaggcg	tgtacgggtg	2820
gaggtctata	taagcagagc	tcgttttagtg	aaccgtcaga	tactagaag	ctttattgcg	2880
gtagtttata	acagttaaat	tgtaacgcca	gtcagtgtct	ctgacacaac	agtctcgaac	2940
ttaagctgca	gaagttggtc	gtgaggcaact	gggcaggtaa	gtatcaaggt	tacaagacag	3000
gtttaaggag	accaatagaa	actgggcttg	tcgagacaga	gaagactctt	gcgtttctga	3060
taggcaccta	ttggtcttac	tgacatccac	tttgcccttc	tctccacagg	tgtccactcc	3120
cagttcaatt	acagctctta	aggctagagt	acttaatacg	actcactata	ggccaccatg	3180
gcgctcccag	tgacagcctt	acttttacct	ctggcggtat	tattgcacgc	ggctcgtcct	3240
gacatacaga	tgactcagac	tacctcttcc	ctatctgctt	ctttaggcga	ccgagtaaca	3300
atatcttgcc	gggccagcca	ggacatctca	aaatacttaa	actggtatca	gcagaagccg	3360
gacggaacag	ttaagttgct	catttaccac	acgtcgagat	tacactcagg	cgttcctagc	3420
cgattttcgg	gttccggctc	cgttacggac	tacagcctga	caatcagtaa	ccttgagcag	3480
gaggacatcg	ccacctactt	ctgtcagcag	ggcaacacgc	tcccgtacac	attcgggtggg	3540
ggaactaagc	tggagattac	cggaggcggg	ggcagcgggtg	gcggcgccag	cgggggtggc	3600
ggctcggagg	tcaagttaca	ggagagcggg	ccgggcttgg	tcgcacctag	ccagagcctc	3660
tcagtcacgt	gcactgtgtc	tggagtcaag	ctcccagact	acggggtatc	atggatacga	3720
cagccgccta	gaaagggcct	agagtggtcg	ggggttatct	ggggaagtga	aaccacatac	3780
tacaactcag	ctctcaagag	ccgcctcacc	atcattaagg	acaacagtaa	gtcgcaggtt	3840
ttcttaaaga	tgaactctct	ccagactgac	gacaccgcta	tttactactg	cgcgaagcac	3900
tactactacg	gcgggagtta	cgcaatggac	tactggggtc	agggcacttc	tgtgaccgta	3960
tccagcacta	ctaccccagc	cccacgtccc	cccacgccag	ctccaacgat	agcaagtccg	4020
cccttatctc	ttcgcctgta	ggcttgcaag	cccgcggcgg	gcggcgccgt	tcacacgcga	4080
ggactagact	tcgcctgcga	catctacatc	tgggcaccac	tagccgggac	ttgcggagtg	4140

-continued

ttgttgttga gcttggtaat aacgctctac tgcaagcgtg ggagaaagaa gctcttgtaac	4200
atthtcaagc agccattcat gcgtcccggt cagacgactc aggaggagga cggctgctcg	4260
tgccgattcc cggaggagga ggaggcggt tgcaactca gagtgaagt ctctcgctcc	4320
gchgacgcac ccgcttacca gcagggtcag aaccagctat acaacgagtt aaacctgggg	4380
cgccgggagg agtacgacgt gttagacaag cgtagaggta gggacccgga gatgggaggc	4440
aagctcggga gaaagaacct ccaggagggc ctgtacaacg aactccagaa ggacaagatg	4500
gctgaggcgt actcggagat tggtatgaag ggcgagagac gtcgcgaaa gggacacgac	4560
ggcttatacc aggggcttcc caccgcgacc aaggacacat acgacgcgct gcacatgcaa	4620
gccttaccac ctcgatgagg taccagcggc cgcttcgagc agacatgata agatacattg	4680
atgagtttgg acaaaccaca actagaatgc agtgaaaaaa atgctttatt tgtgaaattt	4740
gtgatgctat tgctttatth gtaaccatta taagctgcaa taaacaagtt aacaacaaca	4800
attcgaattt aaatcggatc cgcaacaaat ctgactttgc atgtgcaaac gccttcaaca	4860
acagcattat tccagaagac accttcttcc ccagcccagg taagggcagc tttggtgcct	4920
tcgcaggctg tttccttgcct tcaggaaatgg ccaggttctg ccagagctc tggccaatga	4980
tgtctaaaaa tcctctgatt ggtggtctcg gccttatcca ttgccaccaa aacctcttt	5040
ttactaagaa acagtgagcc ttgttctggc agtccagaga atgacacggg aaaaaagcag	5100
atgaagagaa ggtggcagga gagggcacgt ggcccagcct cagtctctcc aactgagttc	5160
ctgctgcct gcctttgctc agactgtttg ccccttactg ctcttctagg cctcattcta	5220
agcccttct ccaagttgcc tctccttatt tctccctgct tgccaaaaaa tctttcccag	5280
ctcactaagt cagtctcagc cagtccactca ttaaccacc aatcactgat tgtgccggca	5340
catgaatgca ccagggttg aagtgaggga attaaaaagt cagatgaggg gtgtgccag	5400
aggaagcacc attctagtth ggggagocca tctgtcagct gggaaaagtc caaataactt	5460
cagattgcaa tgtgttttaa ctcagggttg agaaaacagc caccttcagg acaaaagtca	5520
gggaagggct ctctgaagaa atgctacttg aagataccag ccctaccaag ggcagggaga	5580
ggaccaattg atggagttgg ccactccctc tctgcgcgct cgctcgtca ctgaggccgc	5640
ccgggcaaa cccgggctc gggcgacctt tggtcgccc gcctcagtg gcgagcgagc	5700
gcgagagag gtagtgcca acggcgccgc tgcaagctc aaaaatagct acctctccg	5760
gcatgaattt atcagctaga acggttgaat atcatattga tggtgattth actgtctccg	5820
gcctttctca cccgtttgaa tctttacctc cacattactc aggcattgca tttaaaat	5880
atgagggctc taaaaattht tatccttgcg ttgaaataaa ggcttctccc gcaaaagtat	5940
tacagggca taatgttht ggtacaaccg atthtagctth atgctctgag gctthattgc	6000
ttaatthtgc taatcttht ccttgcctgt atgatttht ggatgthtga attcctgatg	6060
cggtatthtc tcttaagca tctgtgctgt atthacacc gcatatggtg cactctcagt	6120
acaatctgct ctgatgccc atagthtaagc cagccccgac acccgccaac acccgctgac	6180
gcccctgac gggcttctc gctcccggca tccgcttaca gacaagctgt gaccgtctcc	6240
gggagctgca tgtgtcagag gthttcaccg tcatcaccga aacgcgcgag acgaaagggc	6300
ctcgtgatac gcctattht atagthtaat gtcatagata taatgthtcc ttagacgtca	6360
ggtggcactt ttcgggaaa tgtgcgcgga acctattht gthtthttht ctaaatat	6420

-continued

tcaaatatgt atccgctcat gagacaataa ccttgataaa tgcttcaata atattgaaaa	6480
aggaagagta tgagtattca acatttccgt gtcgcocta tcccttttt tgcggcattt	6540
tccttctctg tttttgtca ccagaaacg ctggtgaaag taaaagatgc tgaagatcag	6600
ttgggtgcac gagtgggtta catcgaactg gatctcaaca gcggtaaagat ccttgagagt	6660
tttcgccccg aagaacgttt tccaatgatg agcactttta aagttctgct atgtggcgcg	6720
gtattatccc gtattgacgc cgggcaagag caactcggtc gccgcataca ctattctcag	6780
aatgacttgg ttgagtactc accagtcaca gaaaagcadc ttacggatgg catgacagta	6840
agagaattat gcagtgtgac cataaccatg agtgataaca ctgcggccaa cttacttctg	6900
acaacgatcg gaggaccgaa ggagctaacg gcttttttgc acaacatggg ggatcatgta	6960
actcgccttg atcgttggga accggagctg aatgaagcca taccaaaacga cgagcgtgac	7020
accacgatgc ctgtagcaat ggcaacaacg ttgcgcaaac tattaactgg cgaactactt	7080
actctagctt cccggcaaca attaatagac tggatggagg cggataaagt tgcaggacca	7140
cttctgcgct cggcccttcc ggctggctgg tttattgctg ataaatctgg agccggtgag	7200
cgtgggtctc gcggtatcat tgcagcactg gggccagatg gtaagccctc ccgtatcgta	7260
gttatctaca cgacggggag tcaggcaact atggatgaac gaaatagaca gatcgtgag	7320
ataggtgcct cactgattaa gcattggtaa ctgtcagacc aagtttactc atatatactt	7380
tagattgatt taaaaactca tttttaattt aaaaggatct aggtgaagat cctttttgat	7440
aatctcatga ccaaaatccc ttaacgtgag ttttcgttcc actgagcgtc agaccccgta	7500
gaaaagatca aaggatcttc ttgagatcct tttttctgac gcgtaatctg ctgcttcaa	7560
acaaaaaac caccgctacc agcgggtggt tgtttgcccg atcaagagct accaactctt	7620
tttcggaagg taactggctt cagcagagcg cagataccaa atactgtcct tctagtgtag	7680
ccgtagttag gccaccactt caagaactct gtacaccgc ctacatacct cgctctgcta	7740
atcctgttac cagtggctgc tgcagtggtc gataagtcgt gtcttaccgg gttggactca	7800
agacgatagt taccggataa ggcgcagcgg tcgggctgaa cgggggggttc gtgcacacag	7860
cccagcttgg agcgaacgac ctacaccgaa ctgagatacc tacagcgtga gctatgagaa	7920
agcgcaccgc ttcccgaagg gagaaaggcg gacaggtatc cggtaaagcgg cagggtcggg	7980
acaggagagc gcacgagggg gcttccaggg ggaaacgcct ggtatcttta tagtctgtc	8040
gggtttcgcc acctctgact tgagcgtcga tttttgtgat gctcgtcagg ggggcccagc	8100
ctatggaaaa acgccagcaa cgcggccttt ttacggttcc tggccttttg ctggcctttt	8160
gctcacatgt tctttctgac gttatcccct gattctgtgg ataaccgtat taccgccttt	8220
gagtgagctg ataccgctcg ccgcagccga acgaccgagc gcagcagatc agtgagcag	8280
gaagcgggag agcgcaccaat acgcaaaccc cctctccccg cgcgttggcc gattcattaa	8340
tg	8342

<210> SEQ ID NO 125

<211> LENGTH: 7464

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthesized

<400> SEQUENCE: 125

-continued

cagcagctgg	cgtaatagcg	aagaggcccg	caccgatcgc	ccttcccaac	agttgcgcag	60
cctgaatggc	gaatggaatt	ccagacgatt	gagcgtcaaa	atgtaggtat	ttccatgagc	120
gtttttctcg	ttgcaatggc	tggcggtaat	attgttctgg	atattaccag	caaggccgat	180
agtttgagtt	cttctactca	ggcaagtgat	gttattacta	atcaaagaag	tattgcgaca	240
acggttaatt	tgcgtgatgg	acagactctt	ttactcggtg	gcctcactga	ttataaaaac	300
acttctcagg	attctggcgt	accgttctcg	tctaaaatcc	ctttaatcgg	cctcctgttt	360
agctcccgcg	ctgattctaa	cgaggaaagc	acggtatacg	tgctcgtcaa	agcaaccata	420
gtacgcgccc	tgtagcggcg	cattaagcgc	ggcgggtgtg	gtggttacgc	gcagcgtgac	480
cgctacactt	gccagcgcgc	tagcgcgcgc	tcctttcgtc	ttcttccctt	cctttctcgc	540
cacgttcgoc	ggctttcccc	gtcaagctct	aaatcggggg	ctccctttag	ggttccgatt	600
tagtgcttta	cggcacctcg	acccccaaaa	acttgattag	ggtgatggtt	cacgtagtgg	660
gccatcgcgc	tgatagaagg	tttttcgccc	tttgacgttg	gagtccacgt	tctttaatag	720
tggactcttg	ttccaaactg	gaacaacact	caaccctatc	tcggtctatt	cttttgattt	780
ataagggatt	ttgccgattt	cgcctatttg	gttaaaaaat	gagctgattt	aacaaaaatt	840
taacgcgaat	tttaacaaaa	tattaacggt	tacaatttaa	atatttgctt	atacaatctt	900
cctgtttttg	gggcttttct	gattatcaac	cggggtacat	atgattgaca	tgctagtttt	960
acggcgcgcc	gggttgccca	ctccctctct	gcgcgctcgc	tcgctcactg	aggccggggc	1020
accaaaggtc	gcccgacgcc	cgggctttgc	ccgggcggcc	tcagtgagcg	agcgcgagcg	1080
cagagagggg	gtggccaact	ccatcactag	gggttccctac	gcgtagatct	catattctgg	1140
cagggtcagt	ggctccaact	aacatttgtt	tggactttta	cagtttatta	aatagatggt	1200
tatatggaga	agctctcatt	tctttctcag	aagagcctgg	ctaggaaggt	ggatgaggca	1260
ccatattcat	tttgacggtg	aaattcctga	gatgtaagga	gctgctgtga	cttgctcaag	1320
gccttatact	gagtaaacgg	tagcgcctgg	gcttagacgc	aggtgttctg	atttatagtt	1380
caaaaacctct	atcaatgaga	gagcaatctc	ctggtaatgt	gatagatttc	ccaacttaat	1440
gccaacatac	cataaacctc	ccattctgct	aatgcccagc	ctaagttggg	gagaccactc	1500
cagattccaa	gatgtacagt	ttgctttgct	gggccttttt	cccatgcctg	cctttactct	1560
gccagagtta	tattgctggg	gttttgaaga	agatcctatt	aaataaaaga	ataagcagta	1620
ttattaagta	gccctgcatt	tcaggtttcc	ttgagtggca	ggccaggcct	ggcgcgtaac	1680
gttactgaa	atcatggcct	cttggccaag	attgatagct	tgtgcctgtc	cctgagtcct	1740
agtccatcac	gagcagctgg	tttctaagat	gctatttccc	gtataaaagca	tgagaccgtg	1800
acttgccagc	cccacagagc	cccgccttg	tccatcactg	gcatctggac	tccagcctgg	1860
gttggggcaa	agagggaaat	gagatcatgt	cctaaccctg	atcctcttgt	cccacagata	1920
tccagaaccc	tgaccctgcc	gtgtaccagc	tgagagactc	taaatccagt	gacaagtctg	1980
tctgcctatt	caccgatttt	gattctcaaa	caaatgtgtc	acaaagtaag	gattctgatg	2040
tgtatatac	agacaaaact	gtgctagaca	tgaggtctat	ggacttcaag	agcaacagtg	2100
ctgtggcctg	gagcaactag	tgggcggagt	tagggcggag	ccaatcagcg	tgcgccgttc	2160
cgaaagtgc	cttttatggc	tgggcggaga	atgggcggtg	aacgcgatg	attatataag	2220
gacgcgcgg	gtgtggcaca	gctagtccg	tcgcagccgg	gatttgggtc	gcggttcttg	2280

-continued

tttgttccgg aaagccacca tggcgctccc agtgacagcc ttacttttac ctctggcggt 2340
attattgcac gcggtcgtc ctgacataca gatgactcag actacctctt ccctatctgc 2400
ttctttaggc gaccgagtaa caatatcttg ccgggccagc caggacatct caaaatactt 2460
aaactggtat cagcagaagc cggacggaac agttaagttg ctcatattacc acacgtcgag 2520
attacactca ggcgttecta gccgattttc gggttccggt tccggtaacgg actacagcct 2580
gacaatcagt aaccttgagc aggaggacat cgccacctac ttctgtcagc agggcaacac 2640
gctcccgtac acattcggtg ggggaactaa gctggagatt accggaggcg gtggcagcgg 2700
tggcgccggc agcgggggtg gcggtccgga ggtcaagtta caggagagcg gaccgggctt 2760
ggtcgcacct agccagagcc tctcagtcac gtgcactgtg tctggagtca gtctcccaga 2820
ctacgggta tcatggatac gacagccgcc tagaaagggc ttagagtggc tgggggttat 2880
ctggggaagt gaaaccacat actacaactc agctctcaag agccgcctca ccatcattaa 2940
ggacaacagt aagtcgcagg ttttcttaaa gatgaactct ctccagactg acgacaccgc 3000
tatttactac tgcgcgaagc actactacta cggcgggagt tacgcaatgg actactgggg 3060
tcagggcact tctgtgaccg tatccagcac tactacccca gccccacgct cccccacgcc 3120
agctccaacg atagcaagtc agcccttate tcttcgccct gaggcttga gggccgaggc 3180
ggcgccgcc gttcacacgc gaggactaga cttcgcctgc gacatctaca tctgggcacc 3240
actagccggg acttgcggag tgttgttgtt gagcttggtg ataacgctct actgcaagcg 3300
tgggagaaa aagctcttgt acattttcaa gcagccattc atgctgccc ttcagacgac 3360
tcaggaggag gacggctgct cgtgccgatt cccggaggag gaggagggcg gttgcgaact 3420
cagagtgaag ttctctcgct ccgcgacgc acccgcttac cagcagggtc agaaccagct 3480
atacaacgag ttaaacctgg ggcgcgggga ggagtacgac gtgttagaca agcgtagagg 3540
tagggaccog gagatgggag gcaagcctcg gagaaagaac cccagggagg gcctgtacaa 3600
cgaactccag aaggacaaga tggctgaggc gtactcggag attggtatga agggcgagag 3660
acgtcgcgga aagggacacg acggcttata ccaggggctt tccaccgca ccaaggacac 3720
atacgacgog ctgcacatgc aagccttacc acctcgatga ggtaccagcg gccgcttca 3780
gcagacatga taagatacat tgatgagttt ggacaaacca caactagaat gcagtgaaaa 3840
aaatgcttta tttgtgaaat ttgtgatgct attgctttat ttgtaacct tataagctgc 3900
aataaacaag ttaacaacaa caattcgaat ttaaatcgga tccgcaacaa atctgacttt 3960
gcatgtgcaa acgcttcaa caacagcatt attccagaag acaccttctt ccccagccca 4020
ggtaagggca gctttgtgoc cttcgcaggc tgtttccttg cttcaggaat ggccagggtc 4080
tgcccagagc tctggtaaat gatgtctaaa actcctctga ttggtggtct cgcccttatc 4140
cattgccacc aaaacctct ttttactaag aaacagtgag ccttgttctg gcagtccaga 4200
gaatgacacg ggaaaaaagc agatgaagag aaggtggcag gagagggcac gtggcccagc 4260
ctcagtctct ccaactgagt tctgocctgc ctgcctttgc tcagactgtt tgccccttac 4320
tgctcttcta ggcctcattc taagccctt ctccaagttg cctctcctta tttctccctg 4380
tctgcaaaa aatctttccc agctcactaa gtcagtctca cgcagtcact cattaacca 4440
ccaatcactg attgtgocgg cacatgaatg caccaggtgt tgaagtggag gaattaa 4500
gtcagatgag ggggtgcccc agaggaagca ccattctagt tgggggagcc catctgtcag 4560

-continued

ctgggaaaag	tccaaaataac	ttcagattgg	aatgtgtttt	aactcagggt	tgagaaaaca	4620
gccaccttca	ggacaaaagt	caggaaggg	ctctctgaag	aatgctact	tgaagatacc	4680
agccctacca	aggcgagga	gaggaccaat	tgatggagt	ggcactccc	tctctgcgcg	4740
ctcgtctcgt	cactgaggcc	gcccgggcaa	agccgggcg	tcgggagacc	tttggtcgcc	4800
cggcctcagt	gagcgagcga	gcgcgcagag	agggagtggc	caacggcgcg	cctgcaggtc	4860
tcaaaaatag	ctaccctctc	cggcatgaat	ttatcagcta	gaacggttga	atatcatatt	4920
gatggtgatt	tgactgtctc	cggcctttct	cacccttttg	aatctttacc	tacacattac	4980
tcaggcattg	catttaaaat	atatgagggt	tctaaaaatt	tttatccttg	cgttgaaata	5040
aaggcttctc	ccgcaaaagt	attacagggt	cataatgttt	ttggtacaac	cgatttagct	5100
ttatgctctg	aggctttatt	gcttaatttt	gctaattctt	tgccttgctc	gtatgattta	5160
ttggatgttg	gaattctctga	tgcggtattt	tctccttacg	catctgtgcg	gtatttcaca	5220
ccgcatatgg	tgcactctca	gtacaatctg	ctctgatgcc	gcatagttaa	gccagccccg	5280
acacccgcca	acaccgctg	acgcgccttg	acgggcttgt	ctgctccgg	catccgctta	5340
cagacaagct	gtgaccgtct	ccgggagctg	catgtgtcag	aggttttcac	cgtcacacc	5400
gaaaacgcgcg	agacgaaagg	gctctgtgat	acgcctattt	ttataggtta	atgtcatgat	5460
aataatggtt	tcttagacgt	caggtggcac	ttttcgggga	aatgtgcgcg	gaaccctat	5520
ttgtttattt	ttctaatac	attcaaatat	gtatccgctc	atgagacaat	aaccctgata	5580
aatgttcoa	taatattgaa	aaaggaagag	tatgagtatt	caacatttcc	gtgtcgcct	5640
tattcccttt	tttgcgcat	tttgccttcc	tgtttttgc	caccagaaa	cgctggtgaa	5700
agtaaaagat	gctgaagatc	agttgggtgc	acgagtgggt	tacatcgaa	tgatctcaa	5760
cagcggtaa	atccttgaga	gttttcgccc	cgaagaacct	tttccaatga	tgagcacttt	5820
taaagtctg	ctatgtggcg	cggtattatc	ccgtattgac	gccgggcaag	agcaactcgg	5880
tcgcccata	cactattctc	agaatgactt	ggttgagtac	tcaccagtca	cagaaaagca	5940
tcttacggat	ggcatgacag	taagagaatt	atgcagtgtc	gccataacca	tgagtgataa	6000
cactgccc	aacttacttc	tgacaacgat	cggaggaccg	aaggagctaa	ccgctttttt	6060
gcacaacatg	gggatcatg	taactcgcct	tgatcgttgg	gaaccggagc	tgaatgaagc	6120
cataccaaac	gacgagcgtg	acaccaogat	gcctgtagca	atggcaacaa	cgttgcgcaa	6180
actattaact	ggcgaactac	ttactctagc	ttcccggcaa	caattaatag	actggatgga	6240
ggcgataaaa	gttgcaggac	cacttctcgc	ctcggccctt	ccggctggct	ggtttattgc	6300
tgataaatct	ggagccgggtg	agcgtgggtc	tcgcggtatc	attgcagcac	tggggcccaga	6360
tggtaaagccc	tcccgtatcg	tagttatcta	cacgacgggg	agtcaggcaa	ctatggatga	6420
acgaaaataga	cagatcgtc	agatagggtc	ctcactgatt	aagcattgg	aactgtcaga	6480
ccaagtttac	tcatatatac	tttagattga	tttaaaactt	catttttaat	ttaaaaggat	6540
ctaggtgaag	atcctttttg	ataatctcat	gacaaaatc	ccttaacgtg	agttttcgtt	6600
ccactgagcg	tcagacccc	tagaaaagat	caaaggatct	tcttgagatc	ctttttttct	6660
gcgcgtaatc	tgctgcttgc	aaacaaaaa	accaccgcta	ccagcgggtg	ttgttttgc	6720
ggatcaagag	ctaccaactc	ttttccgaa	ggtaactggc	ttcagcagag	cgagatacc	6780
aaatactgtc	cttctagtgt	agccgtagtt	aggccaccac	ttcaagaact	ctgtagcacc	6840

-continued

gctacatac ctcgctctgc taatcctgtt accagtggt gctgccagtg gcgataagtc	6900
gtgtcttacc ggggttgact caagacgata gttaccggat aaggcgcagc ggtcgggctg	6960
aacggggggt tcgtgcacac agcccagctt ggagcgaacg acctacaccg aactgagata	7020
cctacagcgt gagctatgag aaagcgcac gcttcccga gggagaaagg cggacaggtta	7080
tccggtaagc ggcagggtcg gaacaggaga gcgcacgagg gagcttccag ggggaaacgc	7140
ctggtatctt tatagtcctg tcgggtttcg ccacctctga cttgagcgtc gatttttgtg	7200
atgctcgtca gggggcgga gcctatgaa aaacgccagc aacgcggcct ttttacggtt	7260
cctggccttt tgctggcctt ttgctccat gttcttctc gcgttatccc ctgattctgt	7320
ggataaccgt attaccgctt ttgagtgagc tgataccgct cgccgcagcc gaacgaccga	7380
gcgcagcgag tcagtgagcg aggaagcgga agagcgccca atacgaaac cgctctccc	7440
cgcgcttgg cggattcatt aatg	7464

1-69. (canceled)

70. A method for producing a genetically-modified human T cell comprising a modified human TCR alpha constant region gene, said method comprising:

(a) introducing into a human T cell an mRNA comprising a first nucleic acid sequence encoding an engineered nuclease,

wherein said engineered nuclease produces a cleavage site at a recognition sequence within said human TCR alpha constant region gene; and

(b) introducing into said human T cell a second nucleic acid sequence comprising an exogenous polynucleotide encoding a chimeric antigen receptor comprising an extracellular ligand-binding domain, a transmembrane domain, and one or more intracellular signaling domains, or encoding an exogenous T cell receptor, wherein said second nucleic acid sequence is introduced by contacting said human T cell with a recombinant adeno-associated virus (AAV) vector comprising said second nucleic acid sequence;

wherein said second nucleic acid sequence comprises, from 5' to 3':

(i) a 5' homology arm that is homologous to the 5' upstream sequence flanking said cleavage site;

(ii) said exogenous polynucleotide; and

(iii) a 3' homology arm that is homologous to the 3' downstream sequence flanking said cleavage site;

wherein the sequence of said exogenous polynucleotide is inserted into said human TCR alpha constant region gene at said cleavage site by homologous recombination,

and further wherein said genetically-modified human T cell does not express an endogenous TCR on the cell surface.

71. The method of claim **70**, wherein said recombinant AAV vector is a single-strand AAV vector.

72. The method of claim **70**, wherein said recombinant AAV vector has a serotype of AAV6.

73. The method of claim **70**, wherein said recombinant AAV vector has a serotype of AAV2.

74. The method of claim **70**, wherein said engineered nuclease is a recombinant meganuclease, a recombinant zinc-finger nuclease (ZFN), a recombinant transcription

activator-like effector nuclease (TALEN), a CRISPR/Cas nuclease, or a megaTAL nuclease.

75. The method of claim **74**, wherein said engineered nuclease is an engineered meganuclease.

76. The method of claim **75**, wherein said recognition sequence within said human TCR alpha constant region gene consists of SEQ ID NO: 3.

77. The method of claim **76**, wherein said exogenous polynucleotide is inserted between nucleotides 13 and 14 of SEQ ID NO: 3.

78. The method of claim **70**, wherein said exogenous polynucleotide comprises a promoter that drives expression of said chimeric antigen receptor or said exogenous T cell receptor.

79. The method of claim **70**, wherein said recombinant AAV vector is a single-strand AAV vector, and wherein said recombinant AAV vector has a serotype of AAV6 or AAV2.

80. The method of claim **79**, wherein said engineered nuclease is a recombinant meganuclease, a recombinant zinc-finger nuclease (ZFN), a recombinant transcription activator-like effector nuclease (TALEN), a CRISPR/Cas nuclease, or a megaTAL nuclease.

81. The method of claim **80**, wherein said engineered nuclease is an engineered meganuclease.

82. The method of claim **81**, wherein said recognition sequence within said human TCR alpha constant region gene consists of SEQ ID NO: 3.

83. The method of claim **82**, wherein said exogenous polynucleotide is inserted between nucleotides 13 and 14 of SEQ ID NO: 3.

84. The method of claim **70**, wherein said recombinant AAV vector is a single-strand AAV vector, and wherein said recombinant AAV vector has a serotype of AAV6 or AAV2, and wherein said exogenous polynucleotide comprises a promoter that drives expression of said chimeric antigen receptor or said exogenous T cell receptor.

85. The method of claim **84**, wherein said engineered nuclease is a recombinant meganuclease, a recombinant zinc-finger nuclease (ZFN), a recombinant transcription activator-like effector nuclease (TALEN), a CRISPR/Cas nuclease, or a megaTAL nuclease.

86. The method of claim **85**, wherein said engineered nuclease is an engineered meganuclease.

87. The method of claim **86**, wherein said recognition sequence within said human TCR alpha constant region gene consists of SEQ ID NO: 3.

88. The method of claim **87**, wherein said exogenous polynucleotide is inserted between nucleotides 13 and 14 of SEQ ID NO: 3.

* * * * *