



US 20180237830A1

(19) **United States**

(12) **Patent Application Publication**

JI et al.

(10) **Pub. No.: US 2018/0237830 A1**

(43) **Pub. Date: Aug. 23, 2018**

(54) **COMPOSITIONS AND METHODS FOR DETECTION OF MYCOBACTERIUM AVIUM PARATUBERCULOSIS**

(60) Provisional application No. 61/619,196, filed on Apr. 2, 2012.

(71) Applicant: **Life Technologies Corporation**, Carlsbad, CA (US)

Publication Classification

(51) **Int. Cl.**
C12Q 1/689 (2006.01)
C12Q 1/6806 (2006.01)

(72) Inventors: **Yongmei (Alexis) JI**, Chicago, IL (US); **Pius BRZOSKA**, Woodside, CA (US); **Angela BURRELL**, Austin, TX (US); **Craig CUMMINGS**, Pacifica, CA (US); **Catherine O'CONNELL**, Austin, TX (US)

(52) **U.S. Cl.**
CPC *C12Q 1/689* (2013.01); *C12Q 2600/112* (2013.01); *C12Q 2600/16* (2013.01); *C12Q 1/6806* (2013.01)

(21) Appl. No.: **15/891,454**

(57) **ABSTRACT**

(22) Filed: **Feb. 8, 2018**

Disclosed are compositions, assays, methods, diagnostic methods, kits and diagnostic kits for the specific and differential detection of *Mycobacterium avium* subsp. *paratuberculosis* from samples including veterinary samples, clinical samples, food samples, forensic sample, an environmental sample (e.g., soil, dirt, garbage, sewage, air, or water), including food processing and manufacturing surfaces, or a biological sample.

Related U.S. Application Data

(60) Division of application No. 14/978,243, filed on Dec. 22, 2015, now Pat. No. 9,926,609, which is a continuation of application No. 13/785,409, filed on Mar. 5, 2013, now Pat. No. 9,255,299.

Specification includes a Sequence Listing.

COMPOSITIONS AND METHODS FOR DETECTION OF MYCOBACTERIUM AVIUM PARATUBERCULOSIS

CROSS-REFERENCE(S) TO RELATED APPLICATION

[0001] This application is a divisional of U.S. application Ser. No. 14/978,243 filed Dec. 22, 2015, which is a continuation of U.S. application Ser. No. 13/785,409 filed Mar. 5, 2013 (now U.S. Pat. No. 9,255,299), which claims the benefit under 35 U.S.C. § 119(e) to U.S. Provisional Application No. 61/619,196 filed Apr. 2, 2012. The entire contents of the aforementioned applications are incorporated herein.

FIELD

[0002] The present teachings relate to compositions, methods and kits for specific detection, diagnosis and differentiation of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) for differentially detecting MAP from other non-MAP microorganisms including other *Mycobacterium avium* subspecies (such as *M. avium* subsp. *hominissuis* (MAH), *Mycobacterium avium* subsp. *avium* (MAA), and *Mycobacterium avium* subsp. *silvaticum* (MAS)) in a variety of mammalian host species.

SEQUENCE LISTINGS

[0003] The instant application contains a Sequence Listing which has been submitted via EFS and is hereby incorporated by reference in its entirety.

BACKGROUND

[0004] *Mycobacterium avium* subsp. *paratuberculosis* (MAP) is a bacterium that causes Johne's Disease (chronic granulomatous enteritis of the small intestine) in livestock. Johne's disease results in decreased milk production, fetal loss, diarrhea and early death resulting in substantial economic loss to the livestock and dairy industry.

[0005] There is a need for assays for the rapid, sensitive and specific detection of infectious pathogens for differential identification of MAP from other bacteria and *Mycobacterium avium* subspecies. Such assays are sought for diagnostic identification of animals infected by MAP.

SUMMARY

[0006] The present disclosure, in some embodiments, describes compositions, methods and kits for specific and/or differential detection and diagnosis of MAP from other *Mycobacterium avium* subspecies and/or other non-MAP organisms. Differentially detecting MAP microorganisms from non-MAP organisms in a variety of mammalian host species provides elimination of false positives and superior diagnostic tests which may include one-step diagnosis.

[0007] In some embodiments, the present disclosure describes nucleic acid target sequences that are unique to MAP and not present in other *Mycobacterium avium* subspecies. These unique nucleic acid sequences comprise regions of MAP genomes including non-coding regions, coding regions, genes, alleles and variants thereof and/or portions and/or fragments and/or complements thereof. In some embodiments, a nucleic acid target sequence unique to MAP comprises the sequence of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO:

5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences. In some embodiments, fragments comprise fragments having at least 10, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100, contiguous nucleotides, including all values in-between at least 10 and at least 100 contiguous nucleotides (such as 11, 12, . . . 21, 27 . . . 29 etc.), of any of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5. Sequences having 80%-99% sequence identity may include nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and include all values in-between 80% and 99% not explicitly disclosed.

[0008] In some embodiments, compositions comprising isolated nucleic acid sequences of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences are described. In some embodiments, isolated nucleic acid fragments comprise fragments having at least 10, at least 20, at least 25, or at least 30 contiguous nucleotides of any of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5. Isolated nucleic acids having 80%-99% sequence identity include nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and include all values in-between 80% and 99% not explicitly disclosed.

[0009] Some embodiments describe oligonucleotide primers for use in a nucleic acid amplification method (such as but not limited to PCR) for the detection of target nucleic acid sequences that are unique to MAP. Oligonucleotide primers of the present disclosure comprise at least two primers, having at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting primer sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 49, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0010] In some embodiments, primer sets are described having at least 1 or at least 2 primer sets, wherein each primer set has at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences

having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. In some embodiments, primer sets may be nested primers. In some embodiments, primer sets or primers may be degenerate primers.

[0011] Some embodiments describe oligonucleotide probe sequences for use in detection of target nucleic acid sequences and/or amplified target sequences that are unique to MAP. Oligonucleotide probes of the present disclosure are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting probe sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0012] In some embodiments, isolated nucleic acid sequence compositions of the disclosure, including primers, probes and polynucleotides/oligonucleotides, can further comprise one or more label, such as, but not limited to, a dye, a radioactive isotope, a chemiluminescent label, a fluorescent moiety, a bioluminescent label an enzyme, and combinations thereof.

[0013] The present disclosure, in some embodiments, describes methods (assays) utilizing molecular methods such as nucleic acid sequence specific amplification and detection that offer significant improvements in speed, sensitivity and specificity over traditional microbiological methods. Embodiments relate to design and development of molecular detection assays comprising identification of one or more target nucleic acid sequence that is present in a MAP organism to be detected and absent or divergent in organisms not to be detected. Some embodiments further relate to designing primers, including designing degenerate primers, that can bind to and amplify one or more target nucleic acid sequences encoding for a MAP-specific target nucleic acid and/or a complement and/or a fragment thereof, and using the designed primers to amplify and detect such target nucleic acid sequences.

[0014] In some embodiments, methods of detecting in a sample the presence of a MAP microorganism are disclosed. In some embodiments, methods of detecting the presence of a MAP strain are described.

[0015] The specification also discloses methods for detection of a MAP organism from a sample and methods to exclude the presence of MAP organism in a sample, wherein the detection of at least one nucleic acid sequence that is specific for a MAP organism is indicative of the presence of a MAP organism and the absence of detection of any nucleic acid sequence unique to a MAP organism is indicative of the absence of a MAP organism in the sample.

[0016] In some embodiments, methods of detection of MAP comprise detection of one or more target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto.

[0017] Accordingly, a method of the disclosure, in some embodiments, can comprise detecting, in a sample, at least

one (or more) nucleic acid sequence(s) having at least 10 to at least 25 contiguous nucleic acids of one (or more) MAP specific nucleic acid targets and/or complementary sequences thereof, wherein detection of at least one nucleic acid sequence indicates the presence of an MAP organism in the sample. Non-limiting examples of MAP specific nucleic acids that can be detected include SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences (see also Table 6). Methods of detection can also comprise identification steps and can further comprise steps of sample preparation. Such embodiments are described in detail in sections below.

[0018] One embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of an MAP-specific target nucleic acid and/or a fragment or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof by contacting nucleic acids present in the sample with at least one primer set, having one forward primer and one reverse primer that can hybridize to and amplify the MAP-specific nucleic acid and/or a fragment and/or a complement thereof, under conditions suitable for amplification, and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid amplified by the primers confirms the presence of a MAP organism in a sample. In some embodiments, more than one primer can be used to amplify one or more amplification products.

[0019] Non-limiting exemplary primer pairs comprise a primer pair such as SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; complements thereof and sequences having about 90% identity to the foregoing sequences, wherein one of the two primers of each primer set is a forward primer and the other is a reverse primer.

[0020] Accordingly, an example embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of a MAP-specific nucleic acid and/or a fragment and/or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof contacting nucleic acids present in the sample with at least one primer set, each primer set having one forward primer and one reverse primer, comprising the at least one primer set selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49, wherein the contacting is performed under conditions suitable for an

nucleic acid amplification reaction; and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in a sample.

[0021] In one embodiment a method for detection of a MAP organism from a sample comprises: detecting the presence of one or more MAP specific nucleic acids and/or a fragment or a complement thereof comprising: contacting nucleic acids present in a sample with a multiplex of primer sets each primer set having one forward primer and one reverse primer, comprising a first primer set, a second primer set, and optionally a third (a fourth etc.) primer sets, under conditions optimal for an amplification reaction to obtain one or more amplified nucleic acids; and detecting the one or more amplified nucleic acids, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in the sample.

[0022] Some embodiments describe a method for detection of a MAP organism from a sample comprising: detecting the presence of one or more MAP-specific nucleic acids including detecting a first MAP specific nucleic acid and/or a fragment or a complement thereof comprising: a) amplifying from a sample a first MAP specific nucleic acid and/or a fragment or a complement thereof by contacting nucleic acids present in the sample with at least a first primer set, having one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid and/or a fragment or a complement thereof; and b) amplifying simultaneously from the same sample a second MAP specific nucleic acid and/or a fragment or a complement thereof by simultaneously contacting nucleic acids present in the sample with at least a second primer set, having one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid and/or a fragment or a complement thereof, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction; and detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) and/or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample. In some embodiments a first and a second amplification product can be detected to indicate the presence of a MAP organism in the sample. Furthermore, the process can be optionally repeated to detect a third, a fourth and additional MAP specific nucleic acid molecules.

[0023] In other embodiments, not detecting any amplified product using one or more methods described above can be used to exclude the presence of a MAP organism in a sample.

[0024] Amplification reactions can comprise a PCR amplification, an end-point determination, a quantitative amplification, a real-time PCR such as a SYBR® Green Assay or a TaqMan® Assay.

[0025] Detection can be performed by a variety of methods, such as but not limited to, a nucleic acid amplification reaction such as described in the paragraph above. Detection in some embodiments can be performed by hybridization using probes specific to amplified nucleic acid sequences encoding a MAP specific target nucleic acid sequence. Combinations of amplification and hybridization can be used for detection according to some embodiments. Example probe sequences of the disclosure that can be used for detecting in a method of the present disclosure are SEQ

ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0026] In some embodiments, probes can be used to detect and/or to identify a nucleic acid sequence amplified described in the methods above. For example, a probe comprising SEQ ID NO:8 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 6 and SEQ ID NO 7; a probe having SEQ ID NO:11 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 8 and SEQ ID NO 9; a probe having SEQ ID NO 14 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 12 and SEQ ID NO: 13; a probe comprising SEQ ID NO:17 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 15 and SEQ ID NO 16; a probe having SEQ ID NO:20 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 18 and SEQ ID NO 19; a probe having SEQ ID NO 23 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 21 and SEQ ID NO:22; a probe comprising SEQ ID NO:26 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 24 and SEQ ID NO 25; a probe having SEQ ID NO:29 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 27 and SEQ ID NO 28; a probe having SEQ ID NO 32 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 30 and SEQ ID NO:31; a probe comprising SEQ ID NO:35 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 33 and SEQ ID NO 34; a probe having SEQ ID NO:38 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 36 and SEQ ID NO 37; a probe having SEQ ID NO 41 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 39 and SEQ ID NO:40; a probe comprising SEQ ID NO:44 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 42 and SEQ ID NO 43; a probe having SEQ ID NO:47 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 45 and SEQ ID NO 46; and a probe having SEQ ID NO 50 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 48 and SEQ ID NO:49.

[0027] In one embodiment, disclosed is an assay for the detection of a MAP organism in a sample comprising a) hybridizing a first pair (or set) of PCR primers selected from a row in the Table 3 described as forward primers and reverse primers (selected from primer sets described in the paragraph above), and complements thereof to at least a first MAP target polynucleotide sequence and/or fragment thereof; b) amplifying the at least first target MAP polynucleotide sequences; and c) detecting the at least first and the at least second amplified target polynucleotide sequence products; wherein the detection of the at least first amplified target polynucleotide sequence product and the detection of the at least second amplified target polynucleotide sequence product is indicative of the presence of a MAP organism in the sample. The method can also comprise hybridizing a

second pair of PCR primers selected from another row in Table 3 described as forward primers and reverse primers, and complements thereof to at least a second target MAP polynucleotide sequence and/or fragment thereof.

[0028] In further embodiments, the detection can comprise using hybridization with one or more probes. Probes that can be used are described in Table 3 where different probes specific to different amplified MAP target sequences. Primer-probe combinations are outlined in the Table 3 and also set forth in the sections above (for example, amplification product amplified using primers SEQ ID NO:6 and SEQ ID NO:7 can be detected using probe SEQ ID NO:8).

[0029] In some embodiments, hybridization can comprise at least a first probe and a second probe, the first probe further comprising a first label and the second probe further comprising a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAMTM dye and the second probe is labeled with VIC[®] dye.

[0030] In some embodiments, any detection method described above can further comprise preparing a sample for PCR amplification (prior to hybridizing), for example, but not limited to (1) bacterial enrichment, and/or (2) separation of bacterial cells from the sample, and/or (3) cell lysis, and/or (4) nucleic acid extraction. In some embodiments, a detection method can comprise DNA isolation from a sample.

[0031] Samples may include without limitation, veterinary samples, animal-derived samples, clinical samples, food/beverage samples, water samples, and environmental sample. Veterinary samples may be derived from animals such as but not limited to all ruminants, cattle, sheep, bison, deer, rabbits, foxes and birds including poultry birds, other domesticated and wild birds.

[0032] Methods can include multiplex assays such as polymerase chain reactions, wherein hybridizing and amplifying of the first pair of polynucleotide primers occurs in a first vessel and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a second vessel, or hybridizing and amplifying of the first pair of polynucleotide primers and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a single vessel, the detection is a real-time assay, the real-time assay is a SYBR[®] Green dye assay or a TaqMan[®] assay. Methods can also comprise using additional primers such as a third primer pair and a fourth primer pair and so on.

[0033] A method of the disclosure can further comprise providing a first probe and a second probe (and additional probes such as a third probe and a fourth probe and so on), wherein the first and second probes are different from each other, the first probe operable to identify the first amplified target polynucleotide sequence and the second probe operable to identify the second amplified target nucleotide sequence, the first probe further comprises a first label and the second probe further comprises a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAMTM dye and the second probe is labeled with VIC[®] dye; and hybridizing the first and second probes to the PCR amplified fragments to detect the presence of the first amplified target

polynucleotide sequence and the second amplified target polynucleotide sequence from the sample.

[0034] Some embodiments describe kits suitable for identifying the presence of a MAP organism. Such a kit can comprise at least one set of oligonucleotide primers for use in a PCR process for the detection of a MAP target nucleic acid sequence.

[0035] A first probe can further comprise a first label and a second probe further comprise a second label, both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, and the first probe is labeled with FAMTM dye and the second probe is labeled with VIC[®] dye. Kits can without limitation contain other buffers, molecular bio reagents, and one or more container means for kit components.

[0036] In the following description, certain aspects and embodiments will become evident. It should be understood that a given embodiment need not have all aspects and features described herein. It should be understood that these aspects and embodiments are merely exemplary and explanatory and are not restrictive of the invention. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION

[0037] The present disclosure, in some embodiments, describes compositions, kits and methods for detection of MAP. Some embodiments describe specific detection of MAP from a sample even when non-MAP *Mycobacterium avium* organisms are present in the sample.

[0038] MAP is a bacterium that infects livestock and several herd animals and causes Johne's Disease which is characterized by chronic granulomatous enteritis of the small intestine. In livestock especially dairy animals such as cows, sheep and bison, Johne's disease results in decreased milk production, fetal loss, diarrhea and early death resulting in substantial economic loss to the livestock industry. Existing assays that are used to detect the presence of MAP in samples obtained from animals and to diagnose Johne's disease are based on the detection of insertion sequences present in MAP. MAP is also believed to be linked to human Crohn's disease.

[0039] *Mycobacterium avium* complex (MAC) consists of multiple *Mycobacterium avium* subspecies that can be found as environmental contaminants in soil and water (Covert et al., 1999; Falkinham et al., 2001; Le Dantec et al., 2002; Santos et al., 2005; Hilborn et al., 2008), as well as infectious agents for animals such as pigs, cattle, sheep and birds (C. Cayrou et al 2010). Organisms classified in the MAC complex include *Mycobacterium* subsp. *avium*, *Mycobacterium avium* subsp *paratuberculosis*, *Mycobacterium avium* subsp *silvaticum*, and *Mycobacterium* subsp *hominissuis*. Symptoms of infection vary by host species and MAC organism. Common methods used for identifying MAC subspecies include DNA sequencing, particularly of the 16S and *rpoB* genes, and RFLP (restriction fragment length polymorphism) analysis (C. Cayrou et al 2010).

[0040] In some cases, certain nucleic acid insertion sequences have been associated with some *Mycobacterium* subspecies and this has been used to detect these organisms. For example, insertion sequences known as ISMAP02 and

IS900 are detected in existing PCR-based MAP detection assays which rely on detecting or determining the presence of the ISMAP02 or IS900 target nucleic acid sequences in a sample.

[0041] Next-generation sequencing, using an Ion Torrent Personal Genome Machine (PGM) was performed herein to get complete genome sequences for 16 *Mycobacterium avium* subspecies. Sequences were obtained for one *M. avium* subsp. *silvaticum* sample; four *M. avium* subsp. *avium* samples; two *M. avium* subsp. *hominissuis* samples; three *M. avium* samples with an unknown subtype; and six *M. avium* subsp. *paratuberculosis* samples. The sequenced genomes were analyzed to look for the presence of all previously known insertion sequences as well as for new highly specific single-gene targets. A partial ISMAP02 sequence was found in all organisms indicating that ISMAP02 is not an optimal sequence to use for a MAP-specific assay. Results were also confirmed with qPCR testing using the TaqMan® MAP reagents. Results of this analysis are presented in Table 1.

target region to specifically amplify only the unique MAP-specific target sequences against both inclusion (organism to be detected, i.e., MAP organisms) and exclusion genomes (organisms not to be detected, non-MAP organisms).

[0044] Exemplary embodiments of nucleic acid target sequence unique to MAP identified herein include nucleic acids having SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences.

[0045] Genome coordinates for SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, are described in Table 2 below in relation to the MAP genome described in GenBank accession number NC_002944. For example, nucleic acid target sequence unique to MAP comprising the sequence of SEQ ID. NO: 1 corresponds to a 1,863 bp nucleic acid sequence starting from position 238,936 and continuing through position

TABLE 1

Sample Number	Pathogen	Host	1:100X Sample Dilution		ISMAP02 Sequencing Results
			MAP CT 1	MAP CT 2	
ATCC 49884	<i>M. silvaticum</i>	Wood pigeon	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240)
10-5581	<i>M. avium</i> subsp. <i>Avium</i>	Asian Elephant	16.94	16.96	ISMAP02 positive, full sequence
10-9275	<i>M. avium</i> subsp. <i>Avium</i>	Red Tailed Hawk	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240)
11-4751	<i>M. avium</i> subsp. <i>Avium</i>	Ruddy Duck	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240 and 360-519)
05-4293	<i>M. avium</i>	Avian broadbill from Dallas Zoo	Undetermined	37.45	parital ISMAP02 sequence (position 1-240)
10-4249	<i>M. avium</i> subsp. <i>hominissuis</i>	White lipped deer	14.78	14.76	ISMAP02 positive, full sequence
10-5606	<i>M. avium</i> subsp. <i>hominissuis</i>	Pig	30.24	30.95	parital ISMAP02 sequence (position 1-240 and 360-519)
10-5560	Unknown MAC subtype	Pig	15.49	15.48	ISMAP02 positive, full sequence
09-5983	Unknown MAC subtype	Dog	16.43	16.43	ISMAP02 positive, full sequence
11-0986	Unknown MAC subtype	deer	16.22	16.23	ISMAP02 positive, full sequence
10-4404	<i>M. avium</i> subsp. <i>paratuberculosis</i>	bison	14.38	14.40	ISMAP02 positive, full sequence
10-5975	<i>M. avium</i> subsp. <i>paratuberculosis</i>	bison	14.74	14.79	ISMAP02 positive, full sequence
10-5864	<i>M. avium</i> subsp. <i>paratuberculosis</i>	cattle	14.44	14.45	SMAP02 positive, full sequence
10-8425	<i>M. avium</i> subsp. <i>paratuberculosis</i>	cattle	14.74	14.70	ISMAP02 positive, full sequence
08-8281	<i>M. avium</i> subsp. <i>paratuberculosis</i>	sheep	14.82	14.82	ISMAP02 positive, full sequence
11-1786	<i>M. avium</i> subsp. <i>paratuberculosis</i>	sheep	14.76	14.75	ISMAP02 positive, full sequence

[0042] Other insertion sequences, including IS901 were also found to be non-specific for MAP. The IS901 sequence was present in the *M. avium* subsp. *silvaticum* sequence, and two of the *M. avium* subsp. *avium* sequences (isolates 11-4751 and 10-9275).

[0043] In one embodiment of the current teachings, bio-informatic and direct DNA sequencing comparisons of several *Mycobacterium avium* species were conducted to identify MAP-specific target nucleic acids sequences. Alignment of these sequences using custom algorithms identified several MAP-specific target regions to which primer pairs of the disclosure were designed for each identified MAP-specific

240,798 of the MAP genome described as NC_002944. SEQ ID. NOS: 1-5 are described here and in the Sequence Listing below.

[0046] GenBank accession number NC_002944 can be found at http://www.ncbi.nlm.nih.gov/nuccore/nc_002944 referenced as “*Mycobacterium avium* subsp. *paratuberculosis* K-10, complete genome” 16 Sep. 2011, (1. Li, L., Bannantine, J. P., Zhang, Q., Amonsin, A., May, B. J., Alt, D., Banerji, N., Kanjilal, S. and Kapur, V. “The complete genome sequence of *Mycobacterium avium* subspecies *paratuberculosis*,” Proc. Natl. Acad. Sci. U.S.A., 102 (35), 12344-12349 (2005); 2. CONSR™ NCBI Genome Project

Direct Submission, Submitted (11 Sep. 2004) National Center for Biotechnology Information, NIH, Bethesda, Md. 20894, USA; 3. Li, L., Bannantine, J., Zhang, Q., Amonsin, A., Alt, D. and Kapur, V., Direct Submission, Submitted (5 Sep. 2003) Biomedical Genomics Center, University of Minnesota, 1971 Commonwealth Ave., St. Paul, Minn. 55108, USA).

TABLE 2

MAP genome coordinates (NC_002944)				
Assay ID Number	Signature left	Signature right	Signature length	SEQ ID. NO:
61801, 61802, 61803, 61805	238936	240798	1863	SEQ ID. NO: 1
61808	3088062	3089612	1551	SEQ ID. NO: 2
61813, 61815, 61816, 61817	4168103	4169949	1847	SEQ ID. NO: 3
61819, 61822	4169965	4174680	4716	SEQ ID. NO: 4
61823, 61825, 61826, 61829	4194685	4197076	2392	SEQ ID. NO: 5

[0047] In some embodiments, compositions of the disclosure comprise isolated nucleic acids having SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences. Sequences having 80%-99% sequence identity include isolated nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5 and/or fragments thereof and/or complementary sequences thereof and include all values in-between 80% and 99% not explicitly disclosed. Fragments include oligonucleotides or polynucleotides having at least 10 contiguous nucleotide sequences, or at least 20 contiguous nucleotide sequences, or at least 25 contiguous nucleotide sequences, or at least 30 contiguous nucleotide sequences in any part of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5 and/or fragments thereof and/or complementary sequences thereof and sequences having at least 90% identity to the foregoing sequences.

[0048] Compositions of the disclosure comprise probe and/or primer sequences that are specific to hybridize to (and amplify in the case of primers or detect in the case of probes) one or more target nucleic acid sequences that are unique to a MAP organism. A target nucleic acid sequence unique to a MAP organism may include a gene, a non-coding region, an allele or a complement thereof that is present in a MAP organism but absent from a non-MAP organism. In some embodiments, a target nucleic acid sequence unique to a MAP organism is absent from other *Mycobacterium avium* subspecies which are otherwise very similar or closely related. Target nucleic acid sequences as described in this disclosure can comprise portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto and/or from about 70% to about 90% identity to a target nucleic acid sequence unique to a MAP organism and can include a gene, a non-coding region, an allele or a complement thereof that is present in a MAP

organism but absent from a non-MAP organism. Some example probe and primer sequences are described in Table 3 in sections ahead.

[0049] Several programs for designing primers such as Primer3 (Steve Rozen and Helen J. Skaletsky (2000) "Primer3" on the World Wide Web for general users and for biologist programmers as published in: Krawetz S, Misener S (eds) Bioinformatics Methods and Protocols: Methods in Molecular Biology. Humana Press, Totowa, N.J., pp 365-386), Primer Express® software (Applied Biosystems), and OLIGO 7 (Wojciech Rychlik (2007). "OLIGO 7 Primer Analysis Software". *Methods Mol. Biol.* 402: 35-60) and variations thereof can be used for primer designing. Custom algorithms were used in the present disclosure in conjunction with the new uniquely identified targets to arrive at primer and probe compositions described herein. The presently designed PCR primers and probes are useful in assays such as real-time PCR assays to specifically and unambiguously detect MAP organisms with great sensitivity.

[0050] In some embodiments, exemplary non-limiting compositions comprising primer sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence of SEQ ID. NO: 6, SEQ ID. NO: 7, SEQ ID. NO: 9, SEQ ID. NO: 10, SEQ ID. NO: 12, SEQ ID. NO: 13, SEQ ID. NO: 15, SEQ ID. NO: 16, SEQ ID. NO: 18, SEQ ID. NO: 19, SEQ ID. NO: 21, SEQ ID. NO: 22, SEQ ID. NO: 24, SEQ ID. NO: 25, SEQ ID. NO: 27, SEQ ID. NO: 28, SEQ ID. NO: 30, SEQ ID. NO: 31, SEQ ID. NO: 33, SEQ ID. NO: 34, SEQ ID. NO: 36, SEQ ID. NO: 37, SEQ ID. NO: 39, SEQ ID. NO: 40, SEQ ID. NO: 42, SEQ ID. NO: 43, SEQ ID. NO: 45, SEQ ID. NO: 46, SEQ ID. NO: 48, SEQ ID. NO: 49, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0051] In some embodiments, primer sets are described having at least 1 or at least 2 primer sets, wherein each primer set has at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. In some embodiments, primer sets can be nested primers. In some embodiments, primer sets or primers can be degenerate primers. In some embodiments the present disclosure describes designing multiplex primers that are suitable for multiplex PCR assays.

[0052] Some embodiments describe oligonucleotide probe sequences for use in detection of target nucleic acid sequences and/or amplified target sequences that are unique to MAP. Oligonucleotide probes of the present disclosure are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting probe sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID. NO: 8, SEQ ID. NO: 11, SEQ ID. NO: 14, SEQ ID. NO: 17, SEQ ID. NO: 20, SEQ ID. NO: 23, SEQ ID. NO: 26, SEQ ID. NO: 29, SEQ ID. NO: 32, SEQ ID. NO: 35, SEQ ID. NO: 38, SEQ ID. NO: 41, SEQ ID. NO: 44, SEQ ID. NO: 47, and SEQ ID. NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0053] Isolated nucleic acid sequence compositions of the disclosure, including primers and probes according to the disclosure, can further comprise one or more label, such as, but not limited to, a dye, a radioactive isotope, a chemiluminescent label, a fluorescent moiety, a bioluminescent label an enzyme, and combinations thereof.

[0054] The specification also discloses methods for detection and/or diagnosis of a MAP organism from a sample, wherein the detection of at least one unique or target nucleic acid sequence that is expressed in a MAP organism is indicative of the presence of a MAP organism and the absence of detection of any nucleic acid sequence unique to MAP organism is indicative of the absence of a MAP organism in the sample. Accordingly, some methods of the disclosure exclude the presence of a MAP organism in a sample absent of detection of any nucleic acid sequence unique to MAP organism in the sample. In some embodiments, methods of detecting and/or diagnostically determining in a sample the presence of a MAP microorganisms in the presence of other *Mycobacterium avium* organisms of different subspecies and subtypes are disclosed.

[0055] Diagnosis and diagnostic methods of the disclosure can include detecting the presence of a MAP organism to diagnose a MAP organism induced disease or condition. Any animal or human disease or condition caused by a MAP organism is contemplated to be diagnosable by the detection methods disclosed herein that can detect the presence of a MAP organism as defined herein. Non limiting examples include diagnosis of Johnes's disease and Crohn's disease.

[0056] A method of detection and/or diagnosis of MAP comprises detection of one or more target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. Accordingly, a method of the disclosure, in some embodiments, can comprise detecting, in a sample, at least one (or more) nucleic acid sequence(s) having at least 10 to at least 25 nucleic acids of one (or more) MAP specific nucleic acid targets and/or complementary sequences thereof, wherein detection of at least one nucleic acid sequence indicates the presence of an MAP organism in the sample. Non-limiting examples of MAP specific nucleic acids that can be detected include SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO:5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences.

[0057] Methods of detection and/or diagnosis of MAP in a sample can also comprise identification steps and can further comprise steps of sample preparation. Preparing a sample for PCR amplification (prior to hybridizing with primers), can, in some embodiments, comprise one or more steps such as, but not limited to (1) bacterial enrichment, and/or (2) separation of bacterial cells from the sample (isolation of DNA from sample), and/or (3) cell lysis, and/or (4) nucleic acid (NA) extraction.

[0058] Detection of MAP organisms by the use of methods described herein, in some embodiments, is by an amplification reaction such as a polymerase chain reaction for rapid detection. In general methods of the disclosure include comparing for presence of an MAP organism using suitable controls, for example, typically, an internal positive control can be used in a PCR reaction which will have a detectable

signal/positive result and a suitable negative control that will have no detectable signal. Amplification reactions can comprise one or more of the following: a PCR amplification, an end-point determination, a quantitative amplification, a real-time PCR such as a SYBR® Green Assay and/or a TaqMan® Assay.

[0059] Methods can also comprise detecting at least one amplified nucleic acid by hybridization, mass spectrometry, nanostring, microfluidics, chemiluminescence, enzyme technologies and combinations thereof.

[0060] In one embodiment, a method for detection of a MAP organism from a sample comprises: detecting the presence of an MAP-specific target nucleic acid and/or a fragment or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof by contacting nucleic acids present in the sample with at least one primer set, having one forward primer and one reverse primer that can hybridize to and amplify the MAP-specific nucleic acid and/or a fragment and/or a complement thereof, under conditions suitable for amplification, and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid amplified by the primers confirms the presence of a MAP organism in a sample.

[0061] In some embodiments, more than one primers can be used to amplify one or more amplification products. Non-limiting exemplary primer pairs comprise a primer pair such as SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; complements thereof and sequences having about 90% identity to the foregoing sequences, wherein one of the two primers of each primer set is a forward primer and the other is a reverse primer.

[0062] One example embodiment method for detection (and/or diagnosis) of a MAP organism from a sample comprises: detecting the presence of a MAP-specific nucleic acid and/or a fragment and/or a complement thereof comprising: amplifying a MAP-specific nucleic acid and/or a fragment and/or a complement thereof contacting nucleic acids present in the sample with at least one primer set, each primer set having one forward primer and one reverse primer, comprising the at least one primer set selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49, wherein the contacting is performed under conditions suitable for an nucleic acid amplification reaction; and detecting an amplified nucleic acid, wherein detecting an

amplified nucleic acid using the primers confirms the presence of a MAP organism in a sample.

[0063] In one embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of one or more MAP specific nucleic acids and/or a fragment or a complement thereof comprising: contacting nucleic acids present in a sample with a multiplex of primer sets each primer set having one forward primer and one reverse primer, comprising a first primer set, a second primer set, and optionally a third (a fourth etc.) primer sets, under conditions optimal for an amplification reaction to obtain one or more amplified nucleic acids; and detecting the one or more amplified nucleic acids, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in the sample.

[0064] Some embodiments describe a method for detection of a MAP organism from a sample comprising: detecting the presence of one or more MAP-specific nucleic acids including detecting a first MAP specific nucleic acid and/or a fragment or a complement thereof comprising: a) amplifying from a sample a first MAP specific nucleic acid and/or a fragment or a complement thereof by contacting nucleic acids present in the sample with at least a first primer set, having one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid and/or a fragment or a complement thereof; and b) amplifying simultaneously from the same sample a second MAP specific nucleic acid and/or a fragment or a complement thereof by simultaneously contacting nucleic acids present in the sample with at least a second primer set, having one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid and/or a fragment or a complement thereof, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction; and c) detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) and/or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample. In some embodiments a first and a second amplification product are detected to indicate the presence of a MAP organism in the sample. Furthermore, the method can be optionally repeated to detect a third, a fourth and additional MAP specific nucleic acid molecules.

[0065] In other embodiments, not detecting any amplified product using one or more methods described above can be used to exclude the presence of a MAP organism in a sample.

[0066] Methods of the disclosure can also use other detection methods in addition to nucleic acid amplification reactions described in the paragraphs above. Detection in some embodiments can be performed by hybridization using probes specific to MAP specific target nucleic acid sequence. Accordingly, probes specific to detect one or more target MAP nucleic acids can be used in a sample, under conditions to promote sequence specific hybridization, to detect presence of a MAP nucleic acid in the sample. In some embodiments, hybridization comprises using at least a first probe and a second probe, the first probe further comprising a first label and the second probe further comprising a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhod-

amine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye.

[0067] In some embodiments, combinations of amplification and hybridization can be used for detection and amplified nucleic acid sequences encoding a MAP specific target nucleic acid sequence can be detected as MAP-specific amplification products using MAP specific probes. Some non-limiting example probe sequences of the disclosure that can be used for detecting in a method of the present disclosure are described in SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

[0068] In one embodiment, a method for the detection of a MAP organism in a sample comprises: a) hybridizing a first pair (or set) of PCR primers selected from a row in the Table 3 described as forward primers and reverse primers (selected from primer sets described in the paragraph above), and complements thereof to at least a first MAP target polynucleotide sequence and/or fragment thereof; b) amplifying the at least first target MAP polynucleotide sequences; and c) detecting the at least first and the at least second amplified target polynucleotide sequence products; wherein the detection of the at least first amplified target polynucleotide sequence product and the detection of the at least second amplified target polynucleotide sequence product is indicative of the presence of a MAP organism in the sample. The method can also comprise hybridizing a second pair of PCR primers selected from another row in Table 3 described as forward primers and reverse primers, and complements thereof to at least a second target MAP polynucleotide sequence and/or fragment thereof.

[0069] In further embodiments, the detection can comprise using hybridization with one or more probes. Some example probes are described in Table 3 where different probes specific to different amplified MAP target sequences are described. Primer-probe combinations are also outlined in the Table 3. For example, in some non-limiting embodiments, a probe comprising SEQ ID NO:8 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 6 and SEQ ID NO 7; a probe having SEQ ID NO:11 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 8 and SEQ ID NO 9; a probe having SEQ ID NO 14 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 12 and SEQ ID NO: 13; a probe comprising SEQ ID NO:17 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 15 and SEQ ID NO 16; a probe having SEQ ID NO:20 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 18 and SEQ ID NO 19; a probe having SEQ ID NO 23 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 21 and SEQ ID NO:22; a probe comprising SEQ ID NO:26 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 24 and SEQ ID NO 25; a probe having SEQ ID NO:29 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 27 and SEQ ID NO 28; a probe having SEQ ID NO 32 can be used

to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 30 and SEQ ID NO:31; a probe comprising SEQ ID NO:35 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 33 and SEQ ID NO 34; a probe having SEQ ID NO:38 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 36 and SEQ ID NO 37; a probe having SEQ ID NO 41 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 39 and SEQ ID NO:40; a probe comprising SEQ ID NO:44 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 42 and SEQ ID NO 43; a probe having SEQ ID NO:47 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 45 and SEQ ID NO 46; and a probe having SEQ ID NO 50 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 48 and SEQ ID NO:49.

[0070] In some embodiments of the present methods, one assay alone may not be definitive for detecting a MAP organism due to genomic similarity between the genomic regions of other non-MAP organisms. Yet, when two (or more) assays such as but not limited to the assays shown in Table 3 are used either in parallel or as a multiplex assay, e.g., in a real-time TaqMan® assay, for example, where each probe in each of the two (or more) assays has a different label for distinguishing results on a real-time PCR instrument, e.g., a 7500 Fast Real-Time PCR System (Applied Biosystems), a positive result from such an assay is indicative of the presence of a MAP organism. Such dual or multiplex (more than 2 assay sets) assay approach can be used to detect and distinguish MAP organism. Some embodiments describe detecting at least two (or more) of MAP-specific target nucleic acid target regions as positive identification of a MAP organism.

[0071] Methods of the disclosure include multiplex assays such as polymerase chain reactions, wherein hybridizing and amplifying of the first pair of polynucleotide primers occurs in a first vessel and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a second vessel, or hybridizing and amplifying of the first pair of polynucleotide primers and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a single vessel, the detection is a real-time assay, the real-time assay is a SYBR® Green dye assay or a TaqMan® assay. Methods also comprise using additional primers such as a third primer pair and a fourth primer pair and so on.

[0072] A method of the disclosure can further comprise providing a first probe and a second probe (and additional probes such as a third probe and a fourth probe and so on), wherein the first and second probes are different from each other, the first probe operable to identify the first amplified target polynucleotide sequence and the second probe operable to identify the second amplified target nucleotide sequence, the first probe further comprises a first label and the second probe further comprises a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye; and hybridizing the first and second probes to the PCR amplified fragments to detect the presence of the first amplified target

polynucleotide sequence and the second amplified target polynucleotide sequence from the sample.

[0073] Compositions and methods of the present disclosure are ideally suited for the preparation of a kit suitable for identifying the presence of a MAP organism, such as for example a diagnostic kit. Such a kit can comprise at least one set of oligonucleotide primers for use in a PCR process for the amplification and detection of a MAP-specific target nucleic acid sequence. Some kits of the disclosure comprise at least two sets of oligonucleotide primers for simultaneous use in a multiplex PCR process for the amplification and detection of MAP-specific target nucleic acid sequences. Kits can additionally comprise one or more reagents such as but are not limited to, buffers, nucleotide triphosphates, DNA polymerases, intercalating dye, primers, probes, salt, and instructions for the use of the kit.

[0074] In some embodiments, kit primers can be labeled. A kit comprising multiple pairs of primers can have primer pairs each labeled with different labels that may be detectable separately. Probes comprised in kits of the disclosure can be labeled. If a kit comprises multiple probes each probe can be labeled with a different label to allow detection of different target nucleic acids and/or amplification products that are the targets of each different probe.

[0075] An example kit for the detection and/or diagnosis of a MAP organism comprises: at least one pair of forward and reverse PCR primers having primer pairs selected from primer pairs described in Table 3 (i.e., at least one primer set selected from: a first primer set having SEQ ID NO: 6 and SEQ ID NO 7; and/or a second primer set having SEQ ID NO: 9 and SEQ ID NO: 10; and/or a third primer set having SEQ ID NO: 12 and SEQ ID NO:13; and/or a fourth primer set having SEQ ID NO: 15 and SEQ ID NO 16; and/or a fifth primer set having SEQ ID NO: 18 and SEQ ID NO: 19; and/or a sixth primer set having SEQ ID NO: 21 and SEQ ID NO: 22; and/or a seventh primer set having SEQ ID NO: 24 and SEQ ID NO: 25; and/or an eighth primer set having SEQ ID NO: 27 and SEQ ID NO: 28; and/or a ninth primer set having SEQ ID NO: 30 and SEQ ID NO: 31; and/or a tenth primer set having SEQ ID NO: 33 and SEQ ID NO: 34; and/or a eleventh primer set having SEQ ID NO: 36 and SEQ ID NO: 37; and/or a twelfth primer set having SEQ ID NO: 39 and SEQ ID NO: 40; and/or a thirteenth primer set having SEQ ID NO: 42 and SEQ ID NO: 43; and/or a fourteenth primer set having SEQ ID NO: 45 and SEQ ID NO: 46; and/or a fifteenth primer set having SEQ ID NO: 48 and SEQ ID NO: 49, or sequences comprising at least 90% nucleic acid sequence identity thereof, or labeled derivatives thereof). In some embodiments, a kit can also have at least one probe selected from probes described Table 3 (i.e., probe sequences of the disclosure having SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences, each probe corresponding to the first primer set, second primer set, the third primer set, etc. as described above, respectively, selected based on which primer set is selected).

[0076] Another example and kit for the detection and/or diagnosis of a MAP organism comprise: at least two pairs of forward and reverse PCR primers (two primer pairs)

selected from primer pairs described in Table 3; and optionally at least two probes selected from probes described Table 3.

[0077] A kit of the disclosure can further comprise one or more components such as but not limited to: at least one enzyme, dNTPs, at least one buffer, at least one salt, at least one control nucleic acid sample, loading solution for preparation of the amplified material for electrophoresis, genomic DNA as a template control, a size marker to insure that materials migrate as anticipated in a separation medium, and an instruction protocol and manual to educate a user and limit error in use. Components of kits can be individually and/or in various combinations comprised in one or a plurality of suitable container means.

[0078] It is within the scope of these teachings to provide test kits for use in manual applications or test kits for use with automated sample preparation, reaction set-up, detectors or analyzers. In some embodiments, a kit amplification product may be further analyzed by methods such as but not limited to electrophoresis, hybridization, mass spectrometry, nanostring, microfluidics, chemiluminescence and/or enzyme technologies.

[0079] For the purposes of interpreting of this specification, the following definitions may apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. In the event that any definition set forth below conflicts with the usage of that word in any other document, including any document incorporated herein by reference, the definition set forth below shall always control for purposes of interpreting this specification and its associated claims unless a contrary meaning is clearly intended (for example in the document where the term is originally used). It is noted that, as used in this specification and the appended claims, the singular forms “a,” “an,” and “the,” include plural referents unless expressly and unequivocally limited to one referent. The use of “or” means “and/or” unless stated otherwise. The use of “comprise,” “comprises,” “comprising,” “include,” “includes,” and “including” are interchangeable and not intended to be limiting. Furthermore, where the description of one or more embodiments uses the term “comprising,” those skilled in the art would understand that, in some specific instances, the embodiment or embodiments can be alternatively described using the language “consisting essentially of” and/or “consisting of.”

[0080] As used herein, the phrase “nucleic acid,” “oligonucleotide,” and polynucleotide(s)” are interchangeable and not intended to be limiting.

[0081] As used herein, the phrase “stringent hybridization conditions” refers to hybridization conditions which can take place under a number of pH, salt and temperature conditions. The pH can vary from 6 to 9, preferably 6.8 to 8.5. The salt concentration can vary from 0.15 M sodium to 0.9 M sodium, and other cations can be used as long as the ionic strength is equivalent to that specified for sodium. The temperature of the hybridization reaction can vary from 30° C. to 80° C., preferably from 45° C. to 70° C. Additionally, other compounds can be added to a hybridization reaction to promote specific hybridization at lower temperatures, such as at or approaching room temperature. Among the compounds contemplated for lowering the temperature requirements is formamide. Thus, a polynucleotide is typically “substantially complementary” to a second polynucleotide if hybridization occurs between the polynucleotide and the

second polynucleotide. As used herein, “specific hybridization” refers to hybridization between two polynucleotides under stringent hybridization conditions.

[0082] As used herein, the term “polynucleotide” refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxynucleotides, or peptide nucleic acids (PNA), and includes both double- and single-stranded RNA, DNA, and PNA. A polynucleotide may include nucleotide sequences having different functions, including, for instance, coding regions, and non-coding regions such as regulatory regions. A polynucleotide can be obtained directly from a natural source, or can be prepared with the aid of recombinant, enzymatic, or chemical techniques. A polynucleotide can be linear or circular in topology. A polynucleotide can be, for example, a portion of a vector, such as an expression or cloning vector, or a fragment. An “oligonucleotide” refers to a polynucleotide of the present invention, typically a primer and/or a probe.

[0083] As used herein a “target-specific polynucleotide” refers to a polynucleotide having a target-binding segment that is perfectly or substantially complementary to a target sequence, such that the polynucleotide binds specifically to an intended target without significant binding to non-target sequences under sufficiently stringent hybridization conditions. The target-specific polynucleotide can be e.g., a primer or probe and the subject of hybridization with its complementary target sequence.

[0084] The term “target sequence”, “target signature sequence” “target nucleic acid”, “target” or “target polynucleotide sequence” refers to a nucleic acid present in a MAP organism that is not present in other non-MAP organisms and is unique to MAP. The target sequence can be a polynucleotide sequence that is the subject of hybridization with a complementary polynucleotide, e.g. a primer or probe. The target sequence can be composed of DNA, RNA, an analog thereof, and including combinations thereof. The target sequence may be known or not known, in terms of its actual sequence and its amplification can be desired. The target sequence may or may not be of biological significance. Typically, though not always, it is the significance of the target sequence which is being studied in a particular experiment. As non-limiting examples, target sequences may include regions of genomic DNA, regions of genomic DNA which are believed to contain one or more polymorphic sites, DNA encoding or believed to encode genes or portions of genes of known or unknown function, DNA encoding or believed to encode proteins or portions of proteins of known or unknown function, DNA encoding or believed to encode regulatory regions such as promoter sequences, splicing signals, polyadenylation signals, etc.

[0085] As used herein an “amplified target polynucleotide sequence product” refers to the resulting amplicon from an amplification reaction such as a polymerase chain reaction. The resulting amplicon product arises from hybridization of complementary primers to a target polynucleotide sequence under suitable hybridization conditions and the repeating in a cyclic manner the polymerase chain reaction as catalyzed by DNA polymerase for DNA amplification or RNA polymerase for RNA amplification.

[0086] As used herein, the “polymerase chain reaction” or PCR is an amplification of nucleic acid consisting of an initial denaturation step which separates the strands of a double stranded nucleic acid sample, followed by repetition of (i) an annealing step, which allows amplification primers

to anneal specifically to positions flanking a target sequence; (ii) an extension step which extends the primers in a 5' to 3' direction thereby forming an amplicon polynucleotide complementary to the target sequence, and (iii) a denaturation step which causes the separation of the amplicon from the target sequence (Mullis et al., eds, *The Polymerase Chain Reaction*, BirkHauser, Boston, Mass. (1994). Each of the above steps may be conducted at a different temperature, preferably using an automated thermocycler (Applied Biosystems LLC, a division of Life Technologies Corporation, Foster City, Calif.). If desired, RNA samples can be converted to DNA/RNA heteroduplexes or to duplex cDNA by methods known to one of skill in the art.

[0087] As used herein, “amplifying” and “amplification” refers to a broad range of techniques for increasing polynucleotide sequences, either linearly or exponentially. Exemplary amplification techniques include, but are not limited to, PCR or any other method employing a primer extension step. Other nonlimiting examples of amplification include, but are not limited to, ligase detection reaction (LDR) and ligase chain reaction (LCR). Amplification methods may comprise thermal-cycling or may be performed isothermally. In various embodiments, the term “amplification product” includes products from any number of cycles of amplification reactions.

[0088] In certain embodiments, amplification methods comprise at least one cycle of amplification, for example, but not limited to, the sequential procedures of: hybridizing primers to primer-specific portions of target sequence or amplification products from any number of cycles of an amplification reaction; synthesizing a strand of nucleotides in a template-dependent manner using a polymerase; and denaturing the newly-formed nucleic acid duplex to separate the strands. The cycle may or may not be repeated.

[0089] Descriptions of certain amplification techniques can be found, among other places, in H. Ehrlich et al., *Science*, 252:1643-50 (1991), M. Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press, New York, N.Y. (1990), R. Favis et al., *Nature Biotechnology* 18:561-64 (2000), and H. F. Rabenau et al., *Infection* 28:97-102 (2000); Sambrook and Russell, *Molecular Cloning*, Third Edition, Cold Spring Harbor Press (2000) (hereinafter “Sambrook and Russell”), Ausubel et al., *Current Protocols in Molecular Biology* (1993) including supplements through September 2005, John Wiley & Sons (hereinafter “Ausubel et al.”).

[0090] The term “label” refers to any moiety which can be attached to a molecule and: (i) provides a detectable signal; (ii) interacts with a second label to modify the detectable signal provided by the second label, e.g. FRET; (iii) stabilizes hybridization, i.e. duplex formation; or (iv) provides a capture moiety, i.e. affinity, antibody/antigen, ionic complexation. Labeling can be accomplished using any one of a large number of known techniques employing known labels, linkages, linking groups, reagents, reaction conditions, and analysis and purification methods. Labels include light-emitting compounds which generate a detectable signal by fluorescence, chemiluminescence, or bioluminescence (Kricka, L. in *Nonisotopic DNA Probe Techniques* (1992), Academic Press, San Diego, pp. 3-28). Another class of labels are hybridization-stabilizing moieties which serve to enhance, stabilize, or influence hybridization of duplexes, e.g. intercalators, minor-groove binders, and cross-linking functional groups (Blackburn, G. and Gait, M. Eds. “DNA

and RNA structure” in *Nucleic Acids in Chemistry and Biology*, 2.sup.nd Edition, (1996) Oxford University Press, pp. 15-81). Yet another class of labels effect the separation or immobilization of a molecule by specific or non-specific capture, for example biotin, digoxigenin, and other haptens (Andrus, A. “Chemical methods for 5' nonisotopic labeling of PCR probes and primers” (1995) in *PCR 2: A Practical Approach*, Oxford University Press, Oxford, pp. 39-54).

[0091] The terms “annealing” and “hybridization” are used interchangeably and mean the base-pairing interaction of one nucleic acid with another nucleic acid that results in formation of a duplex or other higher-ordered structure. The primary interaction is base specific, i.e. A/T and G/C, by Watson/Crick and Hoogsteen-type hydrogen bonding.

[0092] The term “end-point analysis” refers to a method where data collection occurs only when a reaction is substantially complete.

[0093] The term “real-time analysis” refers to periodic monitoring during PCR. Certain systems such as the ABI 7700 Sequence Detection System (Applied Biosystems, Foster City, Calif.) conduct monitoring during each thermal cycle at a pre-determined or user-defined point. Real-time analysis of PCR with FRET probes measures fluorescent dye signal changes from cycle-to-cycle, preferably minus any internal control signals.

[0094] The term “quenching” refers to a decrease in fluorescence of a first moiety (reporter dye) caused by a second moiety (quencher) regardless of the mechanism.

[0095] A “primer,” as used herein, is an oligonucleotide that is complementary to a portion of target polynucleotide and, after hybridization to the target polynucleotide, may serve as a starting-point for an amplification reaction and the synthesis of an amplification product. Primers include, but are not limited to, spanning primers. A “primer pair” refers to two primers that can be used together for an amplification reaction. A “PCR primer” refers to a primer in a set of at least two primers that are capable of exponentially amplifying a target nucleic acid sequence in the polymerase chain reaction.

[0096] The term “probe” comprises a polynucleotide that comprises a specific portion designed to hybridize in a sequence-specific manner with a complementary region of a specific nucleic acid sequence, e.g., a target nucleic acid sequence. In certain embodiments, the specific portion of the probe may be specific for a particular sequence, or alternatively, may be degenerate, e.g., specific for a set of sequences. In certain embodiments, the probe is labeled. The probe can be an oligonucleotide that is complementary to at least a portion of an amplification product formed using two primers.

[0097] The terms “complement” and “complementary” as used herein, refer to the ability of two single stranded polynucleotides (for instance, a primer and a target polynucleotide) to base pair with each other, where an adenine on one strand of a polynucleotide will base pair to a thymine or uracil on a strand of a second polynucleotide and a cytosine on one strand of a polynucleotide will base pair to a guanine on a strand of a second polynucleotide. Two polynucleotides are complementary to each other when a nucleotide sequence in one polynucleotide can base pair with a nucleotide sequence in a second polynucleotide. For instance, 5'-ATGC and 5'-GCAT are complementary.

[0098] A “label” refers to a moiety attached (covalently or non-covalently), or capable of being attached, to an oligo-

nucleotide, which provides or is capable of providing information about the oligonucleotide (e.g., descriptive or identifying information about the oligonucleotide) or another polynucleotide with which the labeled oligonucleotide interacts (e.g., hybridizes). Labels can be used to provide a detectable (and optionally quantifiable) signal. Labels can also be used to attach an oligonucleotide to a surface.

[0099] A “fluorophore” is a moiety that can emit light of a particular wavelength following absorbance of light of shorter wavelength. The wavelength of the light emitted by a particular fluorophore, is characteristic of that fluorophore. Thus, a particular fluorophore can be detected by detecting light of an appropriate wavelength following excitation of the fluorophore with light of shorter wavelength.

[0100] The term “quencher” as used herein refers to a moiety that absorbs energy emitted from a fluorophore, or otherwise interferes with the ability of the fluorescent dye to emit light. A quencher can re-emit the energy absorbed from a fluorophore in a signal characteristic for that quencher, and thus a quencher can also act as a fluorophore (a fluorescent quencher). This phenomenon is generally known as fluorescent resonance energy transfer (FRET). Alternatively, a quencher can dissipate the energy absorbed from a fluorophore as heat (a non-fluorescent quencher).

[0101] As used herein the term “sample” refers to a starting material suspected of harboring a particular microorganism or group of microorganisms. A “contaminated sample” refers to a sample harboring a pathogenic microbe (such as MAP) thereby comprising nucleic acid material from the pathogenic microbe. Examples of samples include, but are not limited to, veterinary samples (samples obtained from animals suspected of harboring or being infected by a microorganism—these may include any body fluid or tissue sample), food samples (including but not limited to samples from food intended for human or animal consumption such as processed foods, raw food material, produce (e.g., fruit and vegetables), legumes, meats (from livestock animals and/or game animals), fish, sea food, nuts, beverages, drinks, fermentation broths, and/or a selectively enriched food matrix comprising any of the above listed foods), water samples, environmental samples (e.g., soil samples, dirt samples, garbage samples, sewage samples, industrial effluent samples, air samples, or water samples from a variety of water bodies such as lakes, rivers, ponds etc., samples obtained from animal pens/barns/farms), air samples (from the environment or from a room or a building), forensic samples, agricultural samples, pharmaceutical samples, biopharmaceutical samples, samples from food processing and manufacturing surfaces, and/or biological samples. A “biological sample” refers to a sample obtained from eukaryotic or prokaryotic sources. Examples of eukaryotic sources include mammals, such as a human, a ruminant a cow, a bison, a sheep, a horse, a pig, a chicken, a turkey, a livestock animal, a fish, a crab, a crustacean, a rabbit, a game animal, a murine animal such as rat or mouse, etc. A biological sample may include tissue samples, cell samples, blood, serum, plasma, pus, cerebrospinal fluid, bone marrow, urine, feces, saliva, milk, mucus, or other materials from a human or a livestock animal. A biological sample can be, for instance, in the form of a single cell, plurality of cells, in the form of a tissue, or in the form of a fluid.

[0102] A sample may be tested directly, or may be prepared or processed in some manner prior to testing. For example, a sample may be processed to enrich any contami-

nating microbe and may be further processed to separate and/or lyse microbial cells contained therein. Lysed microbial cells from a sample may be additionally processed or prepared to separate, isolate and/or extract genetic material from the microbe for analysis to detect and/or identify the contaminating microbe. Analysis of a sample may include one or more molecular methods. For example, according to some exemplary embodiments of the present disclosure, a sample may be subject to nucleic acid amplification (for example by PCR) using appropriate oligonucleotide primers that are specific to one or more microbe nucleic acid sequences that the sample is suspected of being contaminated with. Amplification products may then be further subject to testing with specific probes (or reporter probes) to allow detection of microbial nucleic acid sequences that have been amplified from the sample. In some embodiments, if a microbial nucleic acid sequence is amplified from a sample, further analysis may be performed on the amplification product to further identify, quantify and analyze the detected microbe (determine parameters such as but not limited to the microbial strain, pathogenicity, quantity etc.).

[0103] As used herein “preparing” or “preparing a sample” or “processing” or processing a sample” refers to one or more of the following steps to achieve extraction and separation of a nucleic acid from a sample: (1) bacterial enrichment, (2) separation of bacterial cells from the sample, (3) cell lysis, and (4) nucleic acid extraction and/or purification (e.g., DNA extraction, total DNA extraction, genomic DNA extraction, RNA extraction). Embodiments of the nucleic acid extracted include, but are not limited to, DNA, RNA, mRNA and miRNA.

[0104] As used herein, “presence” refers to the existence (and therefore to the detection) of a reaction, a product of a method or a process (including but not limited to, an amplification product resulting from an amplification reaction), or to the “presence” and “detection” of an organism such as a pathogenic organism or a particular strain or species of an organism.

[0105] As used herein, “detecting” or “detection” refers to the disclosure or revelation of the presence or absence in a sample of a target polynucleotide sequence or amplified target polynucleotide sequence product. The detecting can be by end point, real-time, enzymatic, and by resolving the amplification product on a gel and determining whether the expected amplification product is present, or other methods known to one of skill in the art.

[0106] The presence or absence of an amplified product can be determined or its amount measured. Detecting an amplified product can be conducted by standard methods well known in the art and used routinely. The detecting may occur, for instance, after multiple amplification cycles have been run (typically referred to an end-point analysis), or during each amplification cycle (typically referred to as real-time). Detecting an amplification product after multiple amplification cycles have been run is easily accomplished by, for instance, resolving the amplification product on a gel and determining whether the expected amplification product is present. In order to facilitate real-time detection or quantification of the amplification products, one or more of the primers and/or probes used in the amplification reaction can be labeled, and various formats are available for generating a detectable signal that indicates an amplification product is present. For example, a convenient label is typically a label that is fluorescent, which may be used in various formats

including, but are not limited to, the use of donor fluorophore labels, acceptor fluorophore labels, fluorophores, quenchers, and combinations thereof. Assays using these various formats may include the use of one or more primers that are labeled (for instance, scorpions primers, amplifluor primers), one or more probes that are labeled (for instance, adjacent probes, TaqMan® probes, light-up probes, molecular beacons), or a combination thereof. The skilled person will understand that in addition to these known formats, new types of formats are routinely disclosed. The present invention is not limited by the type of method or the types of probes and/or primers used to detect an amplified product. Using appropriate labels (for example, different fluorophores) it is possible to combine (multiplex) the results of several different primer pairs (and, optionally, probes if they are present) in a single reaction. As an alternative to detection using a labeled primer and/or probe, an amplification product can be detected using a polynucleotide binding dye such as a fluorescent DNA binding dye. Examples include, for instance, SYBR® Green dye or SYBR® Gold dye (Molecular Probes). Upon interaction with the double-stranded amplification product, such polynucleotide binding dyes emit a fluorescence signal after excitation with light at a suitable wavelength. A polynucleotide binding dye such as a polynucleotide intercalating dye also can be used.

[0107] As used herein a “MAP organism” or “MAP micro-organism” refers to an organism of the subspecies *Mycobacterium avium* subspecies *paratuberculosis* (MAP).

[0108] As used herein, a “non-MAP organism” is an organism of the *Mycobacterium avium* Complex (MAC) with the exception of *Mycobacterium avium* subsp. *paratuberculosis*. Non-MAP organisms include *Mycobacterium avium* subspecies *avium* (MAA), *Mycobacterium avium* subsp. *hominissuis* (MAH) and *Mycobacterium avium* subsp. *silvaticum* (MAS), as well as MAC organisms of indeterminate subspecies.

[0109] As used herein, a “MAP-specific polynucleotide” refers to a nucleic acid sequence that is able to specifically hybridize to a MAP specific target nucleic acid sequence and/or to a portion and/or complement thereof, under suitable hybridization conditions and which does not hybridize with other nucleic acid sequences that do not encode for a MAP-specific nucleic acid sequence, portions thereof or complements thereof. In some embodiments, a “MAP-specific polynucleotide” of the disclosure may be a probe or primer sequence specific to MAP. It is well within the ability of one skilled in the art, using the present teachings, to determine suitable hybridization conditions based on probe length, G+C content, and the degree of stringency required for a particular application.

[0110] It is expected that minor sequence variations in MAP-specific target nucleotide sequences associated with nucleotide additions, deletions and mutations, whether naturally occurring or introduced in vitro, would not interfere with the usefulness of primer and probe sequences disclosed herein in the detection of MAP organisms, as would be understood by one of skill in the art. Therefore, the scope of the present invention as claimed is intended to encompass minor variations in the sequences of the MAP specific target nucleic acid sequences described here and the MAP specific nucleotides, such as, exemplary primer sets and probe sequences set forth here, and all sequences disclosed also comprise sequences having at least a 90% sequence homology to these sequences.

[0111] A probe may be RNA or DNA. Depending on the detection means employed, the probe may be unlabeled, radiolabeled, chemiluminescent labeled, enzyme labeled, or labeled with a dye. The probe may be hybridized with a sample in solution or immobilized on a solid support such as nitrocellulose, a microarray or a nylon membrane, or the probe may be immobilized on a solid support, such as a silicon chip or a microarray.

[0112] Conditions that “allow” an event to occur or conditions that are “suitable” for an event to occur, such as hybridization, strand extension, and the like, or “suitable” conditions are conditions that do not prevent such events from occurring. Thus, these conditions permit, enhance, facilitate, and/or are conducive to the event. Such conditions, known in the art and described herein, may depend upon, for example, the nature of the nucleotide sequence, temperature, and buffer conditions. These conditions may also depend on what event is desired, such as hybridization, cleavage, or strand extension. An “isolated” polynucleotide refers to a polynucleotide that has been removed from its natural environment. A “purified” polynucleotide is one that is at least about 60% free, preferably at least about 75% free, and most preferably at least about 90% free from other components with which they are naturally associated.

[0113] The words “preferred” and “preferably” refer to embodiments of the invention that may afford certain benefits, under certain circumstances. However, other embodiments may also be preferred, under the same or other circumstances. Furthermore, the recitation of one or more preferred embodiments does not imply that other embodiments are not useful, and is not intended to exclude other embodiments from the scope of the invention.

[0114] The terms “comprises” and variations thereof do not have a limiting meaning where these terms appear in the description and claims. Unless otherwise specified, “a,” “an,” “the,” and “at least one” are used interchangeably and mean one or more than one.

[0115] Also herein, the recitations of numerical ranges by endpoints include all numbers subsumed within that range (e.g., 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, 5, etc.). The term “and/or” means one or all of the listed elements or a combination of any two or more of the listed elements.

[0116] There are many known methods of amplifying nucleic acid sequences including e.g., PCR. See, e.g., PCR Technology: Principles and Applications for DNA Amplification (ed. H. A. Erlich, Freeman Press, NY, N.Y., 1992); PCR Protocols: A Guide to Methods and Applications (eds. Innis, et al., Academic Press, San Diego, Calif., 1990); Mattila et al., Nucleic Acids Res. 19, 4967 (1991); Eckert et al., PCR Methods and Applications 1, 17 (1991); PCR (eds. McPherson et al., IRL Press, Oxford); and U.S. Pat. Nos. 4,683,202, 4,683,195, 4,800,159, 4,965,188 and 5,333,675 each of which is incorporated herein by reference in their entireties for all purposes.

[0117] Nucleic acid amplification techniques are traditionally classified according to the temperature requirements of the amplification process. Isothermal amplifications are conducted at a constant temperature, in contrast to amplifications that require cycling between high and low temperatures. Examples of isothermal amplification techniques are: Strand Displacement Amplification (SDA; Walker et al., 1992, Proc. Natl. Acad. Sci. USA 89:392-396; Walker et al., 1992, Nuc. Acids. Res. 20:1691-1696; and EP 0 497 272, all of which are incorporated herein by reference), self-sus-

tained sequence replication (3SR; Guatelli et al., 1990, Proc. Natl. Acad. Sci. USA 87:1874 1878), the Q β replicase system (Lizardi et al., 1988, BioTechnology 6:1197 1202), and the techniques disclosed in WO 90/10064 and WO 91/03573.

[0118] Examples of techniques that require temperature cycling are: polymerase chain reaction (PCR; Saiki et al., 1985, Science 230:1350 1354), ligase chain reaction (LCR; Wu et al., 1989, Genomics 4:560 569; Barringer et al., 1990, Gene 89:117 122; Barany, 1991, Proc. Natl. Acad. Sci. USA 88:189 193), transcription-based amplification (Kwoh et al., 1989, Proc. Natl. Acad. Sci. USA 86:1173 1177) and restriction amplification (U.S. Pat. No. 5,102,784).

[0119] Other exemplary techniques include Nucleic Acid Sequence-Based Amplification (“NASBA”; see U.S. Pat. No. 5,130,238), Q β replicase system (see Lizardi et al., BioTechnology 6:1197 (1988)), and Rolling Circle Amplification (see Lizardi et al., Nat Genet 19:225 232 (1998)). The amplification primers of the present invention may be used to carry out, for example, but not limited to, PCR, SDA or tSDA. Any of the amplification techniques and methods disclosed herein can be used to practice the claimed invention as would be understood by one of ordinary skill in the art.

[0120] PCR is an extremely powerful technique for amplifying specific polynucleotide sequences, including genomic DNA, single-stranded cDNA, and mRNA among others. Various methods of conducting PCR amplification and primer design and construction for PCR amplification will be known to those of skill in the art. Generally, in PCR a double-stranded DNA to be amplified is denatured by heating the sample. New DNA synthesis is then primed by hybridizing primers to the target sequence in the presence of DNA polymerase and excess dNTPs. In subsequent cycles, the primers hybridize to the newly synthesized DNA to produce discreet products with the primer sequences at either end. The products accumulate exponentially with each successive round of amplification.

[0121] The DNA polymerase used in PCR is often a thermostable polymerase. This allows the enzyme to continue functioning after repeated cycles of heating necessary to denature the double-stranded DNA. Polymerases that are useful for PCR include, for example, Taq DNA polymerase, Tth DNA polymerase, Tfi DNA polymerase, Tma DNA polymerase, Tli DNA polymerase, and Pfu DNA polymerase. There are many commercially available modified forms of these enzymes including: AmpliTaq® and AmpliTaq Gold® both available from Applied Biosystems. Many are available with or without a 3- to 5' proofreading exo-

nuclease activity. See, for example, Vent® and Vent®. (exo-) available from New England Biolabs.

[0122] Other suitable amplification methods include the ligase chain reaction (LCR) (e.g., Wu and Wallace, Genomics 4, 560 (1989) and Landegren et al., Science 241, 1077 (1988)), transcription amplification (Kwoh et al., Proc. Natl. Acad. Sci. USA 86, 1173 (1989)), and self-sustained sequence replication (Guatelli et al., Proc. Nat. Acad. Sci. USA, 87, 1874 (1990)) and nucleic acid based sequence amplification (NABSA). (See, U.S. Pat. Nos. 5,409,818, 5,554,517, and 6,063,603). The latter two amplification methods include isothermal reactions based on isothermal transcription, which produce both single-stranded RNA (ss-RNA) and double-stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

[0123] Those having ordinary skill in the art, in light of this specification, will understand that many modifications, alternatives, and equivalents of the embodiments described above are possible. All such modifications, alternatives, and equivalents are intended to be encompassed herein.

EXAMPLES

[0124] The following procedures are representative examples of embodiments according to the disclosure that may be employed for the detection of a MAP organism. These examples are not intended to be limiting to the scope of the claims and/or the disclosure in any way.

Example 1: Compositions & Methods to Detect MAP Organisms

[0125] The present example describes exemplary assays designed to detect MAP organism using probe and primer sequences designed as described herein. The described assays can be used in diagnostic methods and kits to detect a MAP organism in a sample, such as for example a sample obtained from an animal suspected of being infected with a MAP organism, with no cross-reactivity to a non-MAP or other MAC organism.

[0126] A set of real-time qPCR primers and probes that can specifically amplify and detect the presence of *Mycobacterium avium* subsp. *paratuberculosis* without cross-reactivity with other subspecies in the *Mycobacterium avium* complex including *M. avium hominissuis*, and *M. avium avium* are described in Table 3 below.

[0127] Table 3 lists a set of TaqMan® primer/probe sets, where primer sequences comprise sets/pairs of primers, each primer set including a forward and a reverse primer, and a corresponding probe sequence, that have been shown to be specific to the amplification of MAP with a CT cutoff <38.

TABLE 3

Assay	Coordinate on NC_002944.2						
	ID No	Forward	Reverse	Probe	Left	Right	Locus
61801	CGGTGCGCGGTAG GA SEQ ID NO: 6	CGACCTGGTCTGC GTCATC SEQ ID NO: 7	CCGCGAGGACCTG CTGACGC SEQ ID NO: 8	239559	239635	MAP0225c	Hypothetical protein
61802	GCGATCCGGCGAA ACG SEQ ID NO: 9	GACGAGTTAGACG CACAGATTCT SEQ ID NO: 10	ACGCCCTGCAGTT AGCTCCCCG SEQ ID NO: 11	239855	239922	MAP0225c	Hypothetical protein

TABLE 3-continued

Assay	Coordinate on NC_002944.2						
	ID No	Forward	Reverse	Probe	Left	Right	Locus
61803	ATCGACCGGCTGG TGTAC SEQ ID NO: 12	GCGAGCAGCACGT TGAAC SEQ ID NO: 13	TCGACGCGATGAC CGCGCTCG SEQ ID NO: 14	240155	240255	MAP0226	Hypothetical protein
61805	CAGACACCTGCAG AGCTACTAC SEQ ID NO: 15	AGCATGTCGGCGA CGTT SEQ ID NO: 16	CAAGCCCGACTAC GCCGCCCT SEQ ID NO: 17	240712	240777	MAP0226	Hypothetical protein
61808	TCGCGGTGCCGAT CAG SEQ ID NO: 18	GTTTGGATGCCCG TGACATTC SEQ ID NO: 19	CCGCGCTCCCGA AATGCTC SEQ ID NO: 20	3088984	3089043	MAP2756c	Large hypothetical protein
61813	CGGCGAACCCGTG ACA SEQ ID NO: 21	CACAGGTGCTCCC ATTGAGT SEQ ID NO: 22	CCGCGCCAGCGT GTAGGC SEQ ID NO: 23	4168688	4168782	MAP3740	Large hypothetical protein
61815	GCGCAGCTTCACC ATCGA SEQ ID NO: 24	GTGCGAAGGCTTC GAAGAC SEQ ID NO: 25	AACAGGCTGCCGC CACGCT SEQ ID NO: 26	4169324	4169424	MAP3740	Large hypothetical protein
61816	GCCACGACGGTGA CATCA SEQ ID NO: 27	CCAGGGTGCTGGT GAACA SEQ ID NO: 28	CCGCGTTGCGGCC AGCCCG SEQ ID NO: 29	4169626	4169706	MAP3740	Large hypothetical protein
61817	GGAATCGATCTCG GTGGCTTTT SEQ ID NO: 30	CCACCGCCGGAGC AA SEQ ID NO: 31	CCGCGTGGCGCCG CA SEQ ID NO: 32	4169849	4169946	MAP3740	Large hypothetical protein
61819	GCCCGCAGCGTAG GT SEQ ID NO: 33	CCAGATCTCGGCA AGCGT SEQ ID NO: 34	ACGACCCCGGCAC AGCC SEQ ID NO: 35	4170784	417072	MAP3740	Large hypothetical protein
61822	CGGACGGAGAGGT GGAC SEQ ID NO: 36	GGGCTCCGGCACC A SEQ ID NO: 37	TTGCCGACCCGCA GACCTG SEQ ID NO: 38	4174593	4174666	MAP3742	Large hypothetical protein
61823	GGTAACCAGCAAC CCGATCTG SEQ ID NO: 39	CCCGAGGTGAACC ACTTTGA SEQ ID NO: 40	TCGGTTGATGATT TCACCCAGGACG C SEQ ID NO: 41	4194693	4194779	MAP3756c	Hypothetical protein
61825	GGCGTCCGCCGAA CCA SEQ ID NO: 42	GCAACATTGGCGA TCGACATC SEQ ID NO: 43	CAGCGGAGCATCA CCGACCCCGG SEQ ID NO: 44	4195499	4195561	MAP3757c	Hypothetical protein
61826	GGCTCGGGCCTGA TAGC SEQ ID NO: 45	GCGGCCTGAACTC CTATGATC SEQ ID NO: 46	CCGCGTTCAGTGC CGTGGCC SEQ ID NO: 47	4195904	4195980	MAP3757c	Hypothetical protein
61829	GCAATACTCGTGA ACCGTAAACTG SEQ ID NO: 48	GTGGCGCGGGTGT TG SEQ ID NO: 49	CCCTGGCGGGCAT TTCCTCTACG SEQ ID NO: 50	4196988	4197054	MAP3758c- MAP3759c	Intergenic

*Assay ID describes an assay comprising using Forward Primers and Reverse Primers described in one row and in some embodiments Probe described in the same row.

TABLE 5-continued

Assay ID No.	Sample Info							
	Dog Intermediate	deer Intermediate	bison	bison	cattle	cattle	sheep	sheep
	sp	sp	MAP	MAP	MAP	MAP	MAP	MAP
	MAC 09-5983	MAC 11-0986	4404	5975	5864	8425	8281	1786
61801	40.00	40.00	27.12	28.50	28.47	29.31	30.57	27.03
61802	40.00	40.00	20.10	21.17	20.30	21.15	22.04	20.68
61803	40.00	40.00	26.01	27.56	26.35	27.30	28.06	25.11
61805	40.00	40.00	19.73	20.12	19.60	20.69	21.49	20.09
61808	40.00	40.00	18.33	19.69	19.05	19.57	20.04	19.53
61813	40.00	40.00	29.65	30.04	29.86	30.88	31.10	29.91
61815	40.00	40.00	19.78	21.45	21.36	22.16	22.14	21.03
61816	40.00	40.00	21.03	22.30	21.21	23.19	22.74	21.11
61817	40.00	40.00	32.05	33.05	31.55	33.60	34.49	30.34
61819	40.00	40.00	26.62	25.74	28.12	27.34	28.55	27.02
61822	40.00	40.00	19.59	20.26	20.00	21.08	20.75	20.04
61823	40.00	40.00	19.31	20.06	19.83	20.33	20.27	20.28
61825	40.00	40.00	18.21	19.23	19.26	19.63	19.57	19.20
61826	40.00	40.00	20.73	21.02	22.59	23.12	22.45	21.58
61829	40.00	40.00	19.12	19.91	19.78	20.25	20.22	19.56

[0133] The above experiment demonstrates the feasibility of using any of the assays listed below as a specific and highly sensitive detection method for *Mycobacterium avium* subsp. *paratuberculosis*. In some embodiments, the disclosure provides a set of highly-specific TaqMan® assays for the detection of *Mycobacterium avium* subsp. *paratuberculosis*.

[0134] One or more advantages of the methods of the present disclosure are described in the following section. Each individual assay can independently serve as a highly-accurate diagnostic assay for MAP detection. Each assay has the advantage of targeting single-copy gene in the MAP genome instead of genomic insertion sequences that are utilized in other MAP detection assays. Insertion sequences have the capacity to mobilize between genomes and insert into new species, making them suboptimal for a species-specific assay.

[0135] The currently described assays and methods are able to detect MAP in multiple host species including sheep, cattle, and bison quickly and accurately with no cross-reactivity between very closely related *M. avium* subspecies making it ideal for veterinary diagnostics.

[0136] While the foregoing specification teaches the principles of the present invention, with examples provided for

the purpose of illustration, it will be appreciated by one skilled in the art from reading this disclosure that various changes in form and detail can be made without departing from the spirit and scope of the invention. These methods are not limited to any particular type of sample or nucleic acid contained therein for e.g., total genome DNA, RNA, cDNA and the like may be analyzed using some or all of the methods disclosed in this disclosure. This disclosure provides powerful tools for analysis of complex nucleic acid samples. From experiment design to detection of MAP microbes, the above disclosure provides for fast, efficient and inexpensive methods for detection of pathogenic organisms.

[0137] All publications and patent applications cited above are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent application were specifically and individually indicated to be so incorporated by reference. Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

TABLE 6

SEQ ID NOs: 1-50
SEQ ID NO: 1 CGGGTGC GGATTGATCAGCCGCCCATGGGCCACCAGCGATCCGTGTTGCTTGAGCCGCTGGGTGCGGACGTTGACCTCGT AGCCGAGCACCCCGTCGAGCTGGCCGAGTTGCTCGCGCAGGTAGCGGTACAGGTCGTCCGGAATCTCGGCAGATGGCCACG GCCATCAGGTTTTTGGATCCGCTTGTGGCGCCGGCGAAAGCAATCTGCGGATGGGCCGCGATACGTTCCGCCAGCCGTGTG CACGTCGCGAGGCGCAACCCGTGAGCCACAGCATGGCGCTGAGGCGATGGCCGAGCCGTTCCGGTAGTACGTCGACGTCGT AGACCAGGGCGCCGAGGCTCCAGCGCCGCGATCCGCTGCTCACC CGCCACCCGACCAACCCGGTGC CGCGGCCAGT TGGACTGCGGAGTGGCGCGTCGTCCGCGAGCCCGCGAGCAGCGGGCGGCTTTCGGCGGTGGCCGGGACCCGTTGAGT TCCCGACTCCGGCTGGGCCCGCGGCCACGATCCGCTGCGCTTGCCGGTCCGAGAGCGCGCTGCCGTAGCCCGTCCAGG GCGGTTTACCGGGTCGCCAACAATGCGATCAGATCGATGCTGATGTCGGTGACCGCGGGCGGTGC CGGTTAGGAGC TGCGTCAGCAGGTCCTCGGGTTCGCCAGCGCGCACGGATGACGCAGACCAGGTCGGTCCACCCCGCCAGCACATT GGCGTGGGACAGTCGGGGCGTCGCCACAGCGCTCGGCCAGCCGATATGCGGTCGGCGCCGACGGATGCGGCCACA CCAGTCCGCGTCGCCATCGGCCACCAGGTTCTCCAGCCGACGACGCGCACCCCGTCCGCGCGCAGCCGTTGATC CGCCGGGCGACCGTCTGCTCGGTGGCGCCACCACGCTGGCGATCCGGCGAAACGAGACCCGGGGAGCTAACTGCAGGGC GTGCAGAACTGTCGCTAACTCGTCGTCATCAGGAAAAGTATCCGCGTGGACAGGTCCAAGTGAGGATTTTCAGGCCA ATTTGGTCCGGCGAGCGCAATTTCAACGCTCCGCTGTCAGCTGGAAGTCCGGCTAAGGAGCGCTCACATGCACATC

TABLE 6-continued

SEQ ID NOs: 1-50

GAAATCGGACTGCCAGCCATATCGCCACGTGCGGGGGGGCTGACCGTCGAGTGGGCCCGCCGGGCGAACACCGCGG
CTTTGCGCGGCTGGCCGCGATCGACCGGCTGGTGTACCCGAGCCTCGACGCGATGACCGCGCTCGCGGTGGCCGCGGGCG
CGACCACCGGATCGGGTGAAGTTCACAGTGTCTGCTCGCTCCGCTGTACCCCGCGGTGTGCTGGCCAGCAGGTCATC
ACCCTCGCGGAGGCGTCCGGCGGCCGCTGCGCCTGGGGCTGGGCGTGGGAGCCGCTGCGACGACTACACCGCCGTGCG
CGTCGACTATCGGCGGCGCGGGCCGCTCCGACGAGACGCGCGCGCTGCTGCTGCGCGCGTGTGAGGCGAGGTGGTGA
CGGCGACACGCGCTGTGCCCGCGCGGTCGGATCCCGATCCTGTTCCGTTGGAGCGGGCGGACGCGACGATCCCGCGC
GCCGTGACCGCTCGGTGACGGCTGGACCAGCGGGGGCGCTGCGCGATTACGCCAACAGTCCGTTGTTCCGGAGCGGGTCCG
GGCGCATGGGCGCGCGGGCCGCTCCGACGCGCGTGGCTGACGCGGCGGCTCAACTTCGCTTCCGCGATGACGAGG
CCGTGCGCGCTGGCCGACGACACTGACGAGCTACTACGGCTTCAAGCCGACTACGCGCCCTCAACGTCGCGACATG
CTCACACCGCCGAGGAGGCGGC

SEQ ID NO: 2

CCCGCACCGCGGTCCTGCCCGGACAAATGCCCGTGGCTTCCGCGGCTGCCGCGTGCAGCGGCCTTCGGCGGCGAC
TTCGGCGTCCGCGGCGCCAGCACCTCCGCTTCACTGCCATGGCCGCGGGCCCATGTTGCCGATCGCCGTGCTGGCTG
TGGTGGCCCTGGGTCFCGACGCTGAACAGCTTGTCTTCAACATGCCACGACCCGTCGCGATCGGTGACAGGATCGTCGAG
GCGATGTTGTAGCCCTTGATCGCCAACCATGCCGTGCCATTGACCCACACCGGACGGTACCGCATCCAGCAACGGCTT
GTGCTCGGTGAGCCAAATGGCCGACTCCCAAGCACCCCAACGACTTCAATCATCGCTGGCAGAAAGTCTGGCCGATCT
CGATCGGGCTGCGCTCAACGCCCCCTTGAAGTCTTCAATCTTGGCGTGTAGGTCTCTCGGATTCGGTGAATCCCTTG
GTATTGCCCGCACCGTCCGCTGCGCTGACTGACCTCCTTGACCGCGGACGCGGCTTGGGGCCGTTCTCCCGACCAA
CTGCAATAACCGTCCGACAGATTGCTTGGCCCGGACCATGAGGGCCAGCGCTTGGATCATCGACTGGACATCACCCCTGC
CGGACTTGAGCTGATGTTGAAGCTGTCAATGCTTTTCTTGGTTCACCACTGCTCGACAAGGTTGGCTTCCTCAACG
CTCAAGCCCGCGCGTCTTGGGAACTGCTTGTACGCTCAGCATTGTTCTGGATCGCCGTGCGACCGCCCTGCGCCTG
CGCGGCGAGCTTCGAGAACATCTCATTAGCGCTGTCGGCCACCTGCTTGTCTTGAACAGGCGTCCAGCGTGACCAATCC
CGACGCGGCCAAGCTTCGACTGGATCGTGTATCAACAGGTTCCGCGTGCATGAGCGCGCTCCCGAAATGCTCC
TGAATGTACGGGATCAAAACCGAGCTGCTCCATCTCCTGGGACATTTGGTGGGTTGGCTTGGCCAGCGACGTGATCGC
GTGCGCTCATATTTCTGGGTGGCTGATCAGCAGCCATACCCGACTGAGTGAAGCATCGCCAAAGGACATACAGGTCATTGA
CGCTTTCACCAACTCCGCGCGGACCGGCTCGACCTTGTGCAACGCCCGCGAAGTCTTGCAGGAAACCTTCGACAAAC
CCCACGGCCACATCAATTCGAGGTGACGTTGGCGGCTGCTCGACCGGATGTGGTAGTCTGTCATCGTGGTCTGAC
CGCGCTGATCTGCTCAGCCAGGTCGCGCCCTTCTCGGACGCAAGCTCGCTGAAAGCTCAACACTTGTATCGCATCAC
TGCCCGATAGCGGCTTCTCGACACCCAGCGCAGCTCATCAGCTTTGAGGCGCATAACCGACACGCTCGACAGT
TTCAGCACGCCATCGCTGATCGCTTTCAGATTGGCCGGGACTCCTCGGCCACCGTGTGACGCTCTGCAACGACTGCTG
GAAATTCGCCCGCATCAGTCGCCGAGACA

SEQ ID NO: 3

ACATCGGGTGGAGTGGGGGAGATCGAGCACGCGCTGCGCGGCCACCCGCTGGTGCAGCGCGGAGGTTGGTCCCATCC
ACAATGCACTGCGCTGGGCGCGGGATCGTCTGTAACCGGACGCGCGCAGAGCAATTTAGAGACTCCACACCCGGCGCG
CTGCGCGCCATCTCGCGCTCCGACTCCCGCAGTACATGATCCCCAAGGTGTTGCTCTCATGCCCCGAGCTACCGCTCAC
CGCCAAACGGCAAGTTCGAGCAAAATCGCGCGCGCTCGGAGCGGCGCGCGGGCACCCAGCGCTCGACACGT
CATCCACCTCACTGTGTCGAGCGGCTGGTCCGCGAGGTCGTTCCGATGTGTCGGCGCGCGGATCACCGGCGCGGAG
GACAAATTTCTCGCCAGGGCGGCGATAGCTGCGCGCCACCGAAGCGGTGCGCCGACTGACGCGCAGGGGAGTGGCCGG
AGCGGAGGTGGGCCAGCTGCTCAGCCACGAGACGCTTGGGCAAGTTCAGCGCGGCGTGTGCTCGCCGACCCGGCATCCG
AGGCATCCGAGTCCGCGCGCGATGTCGCGCAACCGTACACCGGGCGAGGGTTCCCGCTCACCCGGTTCGAGCAGGCC
TACACGCTGGGCGCGGCGGACTCAATGGGAGCACCTGTGACCAACGATTTCCGCGTGGTGTGCGCCGCGCGCCGA
GCTGCGCGGTATAGACTGGATCGGTTTGGCCGTGGTACCCAGATGCTCGAGCAATTCCGCGATGCTGCGGTGCGCGC
TGGACGCGGACACCCCAACGGTGCAGGTCGACGCGGGCGGTCGCGCTCATGACTTGAATATACAGAGCAGCCCC
GACCCTGTTACTGCGCGCATGGCGGCGCCCGTTCGATCCGATTCGGTTCGGTGTGATCCAGTCTTCGACCCGTCGAG
GTACCCCGTACGTCGCTGCTGATCAGCTATCTGGGCTCGATGCCCGCAGCTGTCCACCGTCTCACACGATCA
TCGCGAATACAGTCCGACCCCGGCGCGGAGTCCGACCCGCGCGGCTTTCGCGCGGTTGCTTCGAAAGC
GCTTGGGGCGAAACGATGTCGACAAACAGCTTTCGCGCCCTCCGCTGCTGCGCTGACGACCCAGCGACGTCAGCCCTT
CGAGCGGTCACCTTCGCGCGCGCGAGCTTACCATCGAAGAACAGGCTGCGCCAGCGTGCAGTGCAGCAGCCGACAC
TCGCGCTCACCCACCGCGCTGCTTTCGAAGCTTTCGACATGCTGCGTGGCGTGCATCGCGCGCGTTCAGCGATTCGCG
GTGACAGTCCCAAGTCTACCGTCCGACTACGCCCCGCGGACCGCGAGGTTGGGCAACTTCACCCGCTGGCGCT
GTGCGAGTTCGACTACCGCGCCGTCGAGACCGGATCTGCGGAAGCGGTTGCGCGCGCAGCGGGAAGTGTGGCGCGCGG
TGAGCCACGAGGTCATCACCGGCGGGTGGCCGCAACCGGACCGCGGGTGGTACCCCGTGGTGTACACAGCACCC
CTGGGCTCACCCATCAGGACCGCAGCGGGTGAACCAAGTGGGACATTGACCCAGACCCGGGCGTCTGGCTGGACTG
CCAGCCGAGGACGAGGTCGCCGAATTCGATAGCTGGGACATAGCTACCAATGTGGTTGCCCGGAATCGATCTCGG
TGGCTTTTCCGATTCGAGGAGGCGGTGCGCGCCACCGCGGGCAAGCCGAGCGCGGGCACGGCCGTTGCTCCGCGG
GTGGGCG

SEQ ID NO: 4

AGTGGGCGAGCGCGTGCATCGCCCGCGCTGCGCCACTGCCACCCGAGCAGGTCGTCGCGCAATACCATGCTGGTG
CGGCGTGGGAGGCACTGCGATACGTCCTCCAGGATACCGCGCTTCGAGCTCGAACGGGCGGCTCGCCGGTGGCCGG
CATGCTCACCGGAGCCGTGTCACCCAGACGCTGATCGGAGACCCGACGCTGACGCGCGAGGCGCTGCTGCTGCGTGAAG
ACCGCATCGGATGGCCCTCGACGACCTGGCCGGGGCCATCTTCGGGACGCGCGCACGCTGGGGCGGCGGCTTCGCGTC
GTGGAAGTCCGCTCCCGGACCGGCTGATCACCGAACGGTTCGACGAGTGGTGGGCGTGGTGGTGGAGAACTTTGTG
CCTGGAGCCAAACCGAGCTCGCCGGAATCGCCGCGGGCGGCGCTTCCCGGCCCGACCCGCGCACGCTCGACGACCCCG
ACGCGCGTCCGGCTCGACGTTGATCTGCTGTTGGTCTGCTGATCAGTTCGCGCACCGCCGACAGGCGGCTCTCGAGGCG
ATCACCCGTCGAGAGCGGTTGGCTGTTGGATGGTGGAGAAATTCGAGGCCACCCAGCGACGCTCATCAGCGCGCGCTG
TCTCGACCCCGGCTGCTCGGTCGATTCGAAGACGCTGCTCCGCGGATCGGTGGTGGGCGCTCATCGCCGACCCAG
GTTGGCGACCGACGACATGATCCAGGACCGGACCGGCTCAGCTCATCGCGCACCGCCGACAGCCCGGATGCGCG
ACACCGCGGCGGACAGCGCCGCGACAGGTCAGGTCGCGCGGCTGTCGCGCGCTGTCGCGGACCGGACCGGACCG
GGTGGTGGCCAGCTTTCGCGAGATTCGCGAGCTCATCTCGCCATTCACACACCCGCGCTCGACGACGACTTCTTCTGTC
TCGGCGGTGACAGCTCGTTCGCGACCCGGGCTACGCGGACTTTCGGGCGCCGCTTTCGGCCAACTCGCTTCTGTCGAC
CTGTTCAACCACTCGACCTCGTTCGAGCTCGCGCACGCGCGGCGCGCACCGCGCGGAAAGTGTGCGTGGCGGCTGA
GTCGACCCGGGCGGACCCAGCAGCCGACCGATTCGCGCTCACCGTCTGACGAAACGCTATCGGCGCGGCGAGAA

TABLE 6-continued

SEQ ID NOs: 1-50

GCGCGTTGATCCTCGGCGGCGTTCGCGCGCACTGCTACTTCGAGTTCGAGCTCGCGGACTTCGACCCGGCCGAGATTCGAT
 TCGGCGCAGCCAACTCGTAGCACGCCACGCCGACTGCGCACCCAGGTGTCACCGCGGGCACCAGCCGGCCCTCCTC
 GGGTAGGTTCGCGCGTTCGAGCACACCGCCCGATCGAGCCCGTTCGAGAGACACGACGACGTCGAGCCGCGATGCGCG
 ACCAGATCATCGACTTGCAGCGCCCGCCGGGCATCGACTTCGGGGTGCAAACCCGCGCGACGGGCGCACCGTTCGTCGGC
 ATCAGCATGGACAACCATGCTCGACGGCGCCAGCATGATGATCGCCCTGTCGAACCTCGATACCTCTATCGCGCGGA
 AACCGTTCACCAATTGCGCGCGCTGGAACGCTCTTCGCGCACTACGTGTGGAACACCACCGGAGCTGCTGCCCGACGCC
 ACGAGGGGTGCTGCGCGCGTGGCGCCAGCGAGACTATTGGCGCGCACGCCGCCATCTTTGCCCGCGCGCCGAAA
 TTGGCCGACATGCTACTGCTGTTTCGAGATCGAGGAGCCGAGGTTGCAACGGGCAACCGCGACATTCGCCCGTTCGACTG
 GTCGCGAGTAACCGCATCGTGCCTGCGCGAGGGCGTCAACGTCGCGTATTTCGTCGCGCAACTATGACCGGGTGTGT
 CTCGGTGGTTCGGGACCAGCACTTCTGCATCAACGTCACGCTATTTCGACCCGCGACCCCGATGTCGTTGGGATCGAAAAC
 GTCGTCGGAGATTTCACTTCCCTGTTGTTGGAGTCCCGAGTTCGATGAGCCCGCTCGATCTGGGAGAGCGTTCGCGC
 TCTGCAGCGCAATTGATGACCGACTGCGCCACCGCGCGCGGACCGCGTGTGGTTCGAACCGCAACTGTCGCGTTTC
 ACGCAACCCGACCGCGCGCTGTTTCCGTCGCTTCAACAGCGGACTGGGCTTGTTCGACGCTCGGCTCGGCGCGCG
 GTCGCGTTTCGCCGAACCGGTATTTCGCCCTTCGACACACCGCAGACCGGTGTTGGACTTCAGGTGTGGGAAAGCGCGGG
 GCGCTGAGGCTGTCGTTGGACTTCGTCAGTCAGCGGTGTCGCCCGCCACCGCGCACTCAGTTCGAGTTCGTTGGTGG
 ACGGCATCACCGGTTCGCGTCCAGCAGCCCGCGCATCGAACACAAGTTGGGCGAGGGGGCATCAATGACGAGCTCCTG
 CAACGTGTTTCGAGGATCTGCGCGTCCGCCCTGGGTGAGCCGAGGTCGAACCTCACGACAATTTCTTCAGCTCGCGCG
 CGATTCGTCAGCGGACCAAGGTGGTTCGACAGATCGGCGGTGAGCTGTCAGCCTCGGCCACCTTCGACTGCTGTTTCG
 CCAATCCGGTTCGCGGACTTCGCCCGCAAAATCGCCGACACTGACAACCGCGGCAACCCGACTGACCGTTGAGGAG
 GCGATGTTATGACCCGCGCCGAGCTCGTCGACCACTGCGGGGTACCGGCTTCAACTGTGGGCGACGCGTGAATTTG
 CGCTACCGAGCACCAACAGTCTCACCAGCGACTGAAAGTCAACTGGCGCGGTCAAAACGGACGTCGATACCTCCT
 GCTGGCGGAGAGACGACCTGCTGCGCGCGCCGAGGACCGGTTTCGAGCCGTTTCGCTCACGACGTCGAAGCCGAT
 ATCTGGTTCGGGCGCACGTCGGCGTTTCAGTGGGCGGGTAGGCTGCCAGGCTACGCCGAGTTCGCGGTTCGACACACC
 GTGGCAACACCGAGCGCCGAGCAATATCGGAGGGCGTGGCGCAAGGTTGCCAGCCGACGACATGTTGCGCTGCGTCTG
 TCATCCCGAGGGGTATCAGGTGATATGCCCCGACGTCGCCGACGACGGGCTGGTTCATCAGTGTACACCGTTCGAA
 ACGTTGCGCGGCTACGTCGCGGGTCAAGCAATCTGCGTAAACCGGATATCCGCTGGGCGAAGCGCGATGATGAC
 CTGGTGCATCAGATGCGCCCTGACGACACCGTGGTCCATCTGTCGTTGACTGTCGTCGCGGACTTCGTCAGCATCTC
 CATCTGATGACCGACTTTCAGCAGTGTGCTTTCGACCCGAATGCGACTTTCGCGCCGTCGATTTTCAGCTTCGCGACT
 ACCTGCTGAATCTCGCTCGCGAGCGAAGCTCGGCCCGGTTAGTCCCGCGGAAACCGGATCTCGCTACTGGCGGGAT
 CGGCTCGATCAGCTGCGCTCACCACTGTCGTTACCGGTGCTGCCGACGACTAGACCGCTCAGCGTCGCGGACCGCAC
 GACCTTTTTCAGCGGCTTCGATGCGCTGTCGGCCGAGCCTTTCGAGGTGCTCAGCCGCGCGCGGACGACCGGAGCGA
 CCGTCAACGTCGCGCATTCGACCGGATTCAGCCGGGCGATCGCACGCTATGGCGATCGCGACATTTCTGCTGACCCCTG
 ACGACCATGAGACCGCCACGCGTTCACTCCCGCGTTCGGCGAGTGGTTCGGGATTTCAACCGCACCGGCTGCTCGAAGT
 CGATGTCGCGGACAGCGCACCTTCGCCGAACGTCGACCGCGTTCGGTATGCTCTCTTCGACGACATGGACATTTCGA
 CCACCGCGCGCTCAACGTCGCGCGGCTTCGGGCGAGCGGACGACGACGAGGTGAGCAGACGCCGCTGCTGTTCCAC
 TCGACACTCGGCGCACGACCCGAATCGACAGCGCGCAACATCTTTGTTGACCCGATTCAGGTTCGCGGCTTCGAGCA
 GATCCTCAGGTGCTGCTGGACTGTCAGGTTCGCGAAATCGACGGCATGCTTCAGGTCGACTGGGACACCCGCGATCAGG
 CCGTCCCGCGGAGTTCGACCGCGGTTTCGCGACTTCGCGCACGCCCTGGATCTGCTCAGCACCGGCGCTCGGCC
 TGGCACCGGCGCTATTGCGCGCCAGCCCGGAGACACACCGGTTCGAGGGGCGACGACACACCATGAACCGGCGCT
 GATCCATACCGGGTTCTTCGCGCAAGTTCGTTGACACCGGATGCGTCGCTCCGACCGGTGATCGGGCCACACGTC
 ACGCCGAATTGCTCGCTGCGCGCACCGCGTGGCGCACGCTGGCGCGACGGGGTGGCGCCGCTGACTACGTCGGT
 ATCCGGTTGCGCGAGGGCCCGCCAGATCGCGGCTCTCCTGGGCGCATGCTGCGCCGGGCGCTATGTCGCTGGA
 CGTCGGCTGGCCACTCACCGCTCGACAGATCGCGGCCAATGCTCGCTGGCAGCTGTCGAGCCGAGCGGAGGAGG
 TGGACCGGCTGCTTCGCCAGCCGCAACTGTTGTCACCGCGCGCGGCGGTTGGTCCGAGGCCCAACAGCGAGGTGCT

SEQ ID NO: 5

CCGACTCCGGTAACACGAAACCCGATCTGGGCTCGGTTGATGATTTACCCCGAGGACGCAAGGGCGCTCCAACCCTCAAA
 GTGGTTCACTTCGGCTGATGTTGGGACAACACGGGCGCGCATCGTGGCCTTGTCAAAAACCGGTTTCGTAATGTT
 CGTATCAAAAGATCAGGTCGGCCCCATACATTCGGCTTGGATGACCGCGGACGCCAGGTCCGGTAGTTCGGGGTCCCG
 GCGGCTGGATCTGAACACCAACGGAGGAGATCAATTGCCCGCACCCGGCTGTCGATCCCGGAGGGTTGAGTCCGGTGG
 GTGATCAGTTCGAGGATGATTGACTGGGATCTGCGCGTGTAGCCATGTGCCGGTGGCAAAGAACCAGCGCCGATTTGGG
 GGACCCGTGCTCGAGCAGCAGGTAGTAGTTCGCTGATCGGGAATACGGTGTGCTGTAGCCCGTTCGATGAGCAGCGTCTCAG
 CGCAGGGCCGGTCCAGGATACCTTGGGCTACAAGGGACAGCTTCGGCGGCTAGTCCACCCATTGCTCGAAGGTGTCGCGC
 CCAAATGATAGGGCAGGGTTCGCGCCAGTTCGAAGGGTACTCGCTGATTCGGGCTGCTCGATCCTACTCGGCGTCAAA
 GCCATGGTTCGATGACGCGCCTTGGCTGACCAACAGGGCAGGGAGTTCGGTGGGTTGGGCGATTTTGGCCGCCAGT
 AGCCACCGGTACTACCAACCCCAATAGCCCAACCGCGCGGTCGAGTTTCGGGTTCGCGCGGATCCACTCCAAACCGGGG
 GTGAACATTCGCTCGGCGTCTCCGAAACAGCAGCGGAGCATCACCGACCCGGGATGTCGATCGCCAAATGTTGCGAC
 GTTCGCGGCGAGCACCTGATCGGTGTCATGCTTCTTGAAGGTGTCGATCCCGCCGAGACGAGCAATACAGGCAGCC
 GGTCCGTCGCTTCGGGACACGAGATGGGCGATGATGCGGTACCCCTCCCGGGCGCGCCGGCGAAGCGGATCTCGACA
 CGCTCGATGGGATGTCGAAGTACCGGATCGCGCAGCAACATTTCTGCGACTTCGCGCTAGGCGCTTCTTACCGGCG
 CGAATTCATCGTTCGGTTCGGGCGGCTCAGGTAGTAGCCATAGGCGCGCAAAATGTTTCTGTCGCGCACACTGGTTCG
 CAATCTGTTCCGGCTTTCGCGCTCGGGCTGATAGCGTGGCGCCACCGGACTGAACCGGCGCGCCAGGCGTCCGATCA
 TAGGAGTTTCAGGCGCTCAACCGCCCTGCACATCATCGCGAGCGTGTACTGAAACCGGATAAATTCGCCCGTGGCCGA
 GACCGCCTAGCGGCTAGCTCGCTCACCCTGCGGTCGCTGTCGCTTACACCGGCAAGTTATGACCGTTCGCGGCTGCA
 GATATTGGAGAAAACCGACATCTCACCGCTGCGTGGGCGAGCCGCAACCGGAAAGTTCTGTTGGGAGCCGCTCCAGCCAGT
 GCCAGGCATTCGCTGATCAGTGGGACTGATCGGCGTAAACCGCTTCGATCGCTAGATCGGCGCCACCGGATTCGGTGTATC
 ACCGCGATGGATCAAAATCCAGCAGCCGCTGGAACCGGAAAATTCGGCCAAAGCCGCTTGGGGCGAAGCCGACCGCGGCT
 GCAACCGGCAAGCTGCGCTCGCTCAACCCAAACAGAGGAGGCGACATCCGAAACAGAGCGGGCGGGTGGACACGCG
 AGCATCTCGATTGAGTTCGCGCACTGGCTGATCGAACACCGGATCGATCTTGACCAATACCGCGCCAGCACCGCGCCAA
 TAATCGACCCGTTGCCGAAAAGCGTTGGCTCCAGCACATCTCAGCCAATCCGACGCGGATGTCGCCGAACCCGAGT

TABLE 6-continued

SEQ ID NOs: 1-50

CAACCCGGATCTGGGTATCTTAACTCGCTGACAGGATGGCCAGCACCCGCGGCGAGCACCCGGTCGTAACCGCAGA
CCGATCATCGCACCTTCGGTGCCCGCACATTGGTCGTAAGGTTGTGGCCGGACCAGAAACCATGACTGCGCCCTCGGC
GGTGACGAACAGATCGAGCGAACCGTCCGGCATCACCCGAGCGCCACCAGCGGTATTGAGGCGCACCCAGCCACAGT
CCACATACGGCGCGAGCGCACCCAAAGGTCGACACTCCAATACCCGACCATCACGCTCACATTGAGCCAGGTCCTACGT
CGCTCCGCAACAAATTGGGCCGCAATCCTTCGAAGCGCCCGGTCAGCGCAGACTCCAGGGCGCAATACTCGTGAACCG
TAAACTGCGGCGTAGAGAAATGCCCGCAGGGCGCAACCCCGCCACTTCGGCGAGAAAGCGTGACAGG

SEQ ID NO: 6:
CGGTGCGCGTAGGA

SEQ ID NO: 7
CGACTGGTCTGCGTCATC

SEQ ID NO: 8
CCGCGAGGACCTGCTGACGC

SEQ ID NO: 9
GCGATCCGGCGAAACG

SEQ ID NO: 10
GACGAGTTAGACGCACAGATTCT

SEQ ID NO: 11
ACGCCCTGCAGTTAGCTCCCCG

SEQ ID NO: 12
ATCGACCGGCTGGTGTAC

SEQ ID NO: 13
GCGAGCAGCAGTTGAAC

SEQ ID NO: 14
TCGACGCGATGACCCGCTCG

SEQ ID NO: 15
CAGACACCTGCAGAGCTACTAC

SEQ ID NO: 16
AGCATGTGCGGACGTT

SEQ ID NO: 17
CAAGCCGACTACGCCCCCT

SEQ ID NO: 18
TCGCGGTGCCGATCAG

SEQ ID NO: 19
GTTTGGATGCCCGTGACATTC

SEQ ID NO: 20
CCGCGCTCCCCGAAATGCTC

SEQ ID NO: 21
CGCGGAACCCGTGACA

SEQ ID NO: 22
CACAGGTGCTCCATTGAGT

SEQ ID NO: 23
CCGCGCCAGCGTGTAGGC

SEQ ID NO: 24
GCGCAGCTTACCATCGA

SEQ ID NO: 25
GTGCGAAGGCTTCGAAGAC

SEQ ID NO: 26
AACAGGCTGCCGCCACGCT

SEQ ID NO: 27
GCCACGACGGTGACATCA

SEQ ID NO: 28
CCAGGGTGCTGGTGAACA

TABLE 6-continued

SEQ ID NOs: 1-50
SEQ ID NO: 29 CCGCGTTGCGGCCAGCCCG
SEQ ID NO: 30 GGAATCGATCTCGGTGGCTTTT
SEQ ID NO: 31 CCACCGCCGGAGCAA
SEQ ID NO: 32 CCGCGTGGCGCCGCA
SEQ ID NO: 33 GCCGCGACGGTAGGT
SEQ ID NO: 34 CCAGATCTCGGCAAGCGT
SEQ ID NO: 35 ACGACGCCGGCACAGCC
SEQ ID NO: 36 CGGACGGAGAGGTGGAC
SEQ ID NO: 37 GGGCTCCGGCACCA
SEQ ID NO: 38 TGCCGACCCGCAGACCTG
SEQ ID NO: 39 GGTAACCAGCAACCCGATCTG
SEQ ID NO: 40 CCCGAGGTGAACCACTTTGA
SEQ ID NO: 41 TCGGTTGATGATTTACCCAGGACGC
SEQ ID NO: 42 GGCGTCCTCCGAACCA
SEQ ID NO: 43 GCAACATTGGCGATCGACATC
SEQ ID NO: 44 CAGCGGAGCATCACCGACCCCGG
SEQ ID NO: 45 GGCTCGGGCCTGATAGC
SEQ ID NO: 46 GCGGCCTGAACTCCTATGATC
SEQ ID NO: 47 CCGCGTTCAGTGCCGTGGCC
SEQ ID NO: 48 GCAATACTCGTGAACCGTAAACTG
SEQ ID NO: 49 GTGGCGGGGTGTTG
SEQ ID NO: 50 CCCTGGCGGGCATTTCCTCTACG

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 50

<210> SEQ ID NO 1

<211> LENGTH: 1863

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 1

```

cgggtgcgga ttgatcagcc gcccatgggc caccagcgat ccgtggtgct tgagccgctg      60
ggtgcggaag ttgacctcgt agccgagcac cccgtcgagc tggccgagtt gctcgcgcag      120
gtagcggtag aggtcgtcgg aatctcggca gatggccacg gccatcaggt ttttggatcc      180
gcttgtagcg ccggcgaaag caatctgcgg atgggcccgg atacggtcgc cgaccgtgtg      240
cacgtcgcga ggcgcaaccg tgagccacag catggcgctg aggcgatggc cgagccgttc      300
gggtagtagc tcgacgtcgt agaccagggc gcccgaggcc tccagcgccg cgatccgcct      360
gctcaccgcg gccaccgacc aaccggtgcg cgcggccagt tgggactgcg gactgcggcc      420
gtcgtcggcg agcgcgcgca gcagcggcg gtcttcggcg gtgggcccgg accggtgagt      480
tcccgaactc ggctgggccc gcgcggccac gatccgctgc gcttgccggt ccgagagcgc      540
gctgcccgtg cccgtccagg gcgcggtcac cgggtcgcgg aacgaatgca gcatcagatc      600
gatgctgatg tcggtgaccg cggcgggtgc cggtaggagc tgcgtcagca ggtcctcgcg      660
ggtgtcgcgc agcggcgcac ggatgacgca gaccaggctg gtccaccccg ccagcacatt      720
ggcgtgggac acgtcggggc gtcgcaccag cgcgtcggcc agccgcgata tcgcggtccgg      780
cgccgcacgg atgcggcaca cccagtccgc gtcgcccacg gccacccggt tctccagccc      840
gacgacgcgc accaccccgt cgcggcgccg ccggtgatac cgccgggcca ccgtctgctc      900
ggtggcgccc accacgctgg cgatccggcg aaacgagacc cggggagcta actgcagggc      960
gtgcagaatc tgtgctgcta actcgtcggc catcaggaaa agtatccgcg tggacaggtc     1020
caagtgagga tttcaggcca atttggctgc ggcgagcgca atttcaacgc ctccgctgtc     1080
agcctggaag ctccggcctaa ggagcgctca catgcacatc gaaatcggac tgcccagcca     1140
tatcggccac gtcggggggc ggctgaccgt cgagtgggcc cgcggggccg aacaccgcgg     1200
ctttgcccgc ctggccgcga tcgaccggct ggtgtaccgc agcctcgacg cgatgaccgc     1260
gctcgcggtg gcccgggggc cgaccaccgg catcggggtg acgttcaacg tgctgctcgc     1320
tccgctgtac ccccggtgtc tgctggccaa gcaggtcacc accctcgcgg aggcgtccgg     1380
cggccggctg cgccctggggc tgggctcggg gagccgctgc gacgactaca ccgccgtcgg     1440
cgtcgactat cggcgggcgcg ggcgcattct cgacgagacg gcggcgctgc tcggtgccgc     1500
gtgtgaggcc gaggtggtga ccggcgacca gccgctgtgc ccggcgccgg tcgggatccc     1560
gatcctgttc ggtggacggg cggacgcgac gatccgcccg gccgtgaccg tcggtgacgg     1620
ctggaccgcc ggggcgctgc gcgattacgc caaccagtcg gtgttcgccc agcgggtccg     1680
ggcggcatgg gcgcgcgggg gccgctccgg acggcctggt ctgcaggcca gcgtcaactt     1740
cgccttcggc gatgacgagg ccgtcgcccg tggcgcgaga cacctgcaga gctactacgg     1800
cttcaagccc gactacgccc cctcaacgt cgcggacatg ctcaccacgc cgcaggaggg     1860
ggc                                                                                   1863

```

<210> SEQ ID NO 2

-continued

```

<211> LENGTH: 1551
<212> TYPE: DNA
<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 2
cccgcaccgc cggtcctgcc gccggacaat gccgcggttg cttcgccggc tgccgcctg      60
acgcggcctt cggcggcgac ttcggcgctc cggcgggcca gcaacctggc ttcacctgcc      120
atggccgcgg ggcccattgt gcgcatcgcc gtgctggctg tggtgacctg ggtctcgacg      180
ctgaacagct tgtccttcaa catgcccaaga cccgtcccga tcggtgacag gatcgctgag      240
gcatgtttgt agcccttgat cgccaacctat gccgtgccca ttgcacctac accgacggtc      300
accgcatcca gcaacggcct gtgctcggtg agccaatgcg ccgactcccc aagcacccca      360
acgacttcag tcatcgctgg cagaaagtcc tggccgatct cgatgcgggc tgcgctcaac      420
gcccccttga agtcttcat cttggcgctg taggtctcct gggattcggg gaatcccttg      480
gtattgcccc caccgtggc ctcgcctga ctgacctct tgaccgcggc ageggetttg      540
ggcccgttct ccccgaccaa ctgcaatacc gtccgcagat tgtcttgccc gccgacctg      600
agggccagcg cttggatcat cgactggaca tcaccttgc cggacttgag ctgattgtt      660
aagcctgtca atgcttttcc ttggttccac cactgctcga caagggtggc ttcctcaacg      720
ctcaagccgc cgcggtcctt gcggaactgc ttgtaagcct cagcattgtt ctggatcgcc      780
gtcgcgaccc cctgcgctg cgcgggcagc ttcgagaaca tctcattagc gctgtcggcc      840
acctgcttgc tcttgaacca ggcgctccagc gtgacctacc ccgacggggc aagcttcgac      900
tggatcgtgt catacaacag gttcgcggtg ccgatcaggc cgcgctcccc gaaatgctcc      960
tgaatgtcac gggcatccaa accgagctgt cccatctcct gggacatttg ctgggttggc      1020
ttggccagcg acgtgatgc gtgcgtcata ttctgggtgg cctgatcagc acccataccc      1080
gactgagtga gcatcgccaa cgacgcatac aggtcattga cgctttcacc aactccggcc      1140
gcgaccggct cgacgttggt caacgcccc gcgaagtctt gcagcgaaac cttcgacaac      1200
ccccaggcca cattcaattt cgaggtgacg ttggcggcct gctcgaccgg gatgtggtag      1260
tcgtgcatcg tggctgtgac cgcgctgatc gtctcgccca ggtccgcgcc ctcttcggac      1320
gcaagctgcy ctgaagcct caacaccttg atcgcatcac tgccccgata gccggccttc      1380
tcgacaccca gcgcagcgtc catcagcttt tgaggcgcac aaccgaccac gctcgacagt      1440
ttcagcacgc catcgctgat cgccttcaga ttggccggcg actcctcggc caccgctgtc      1500
agcctctgca acgactgctg gaaattcccc gccgcatcag tcgccgagac a          1551

```

```

<210> SEQ ID NO 3
<211> LENGTH: 1847
<212> TYPE: DNA
<213> ORGANISM: Mycobacterium avium

```

```

<400> SEQUENCE: 3
acatcgggtg gagtgcgggg agatcgagca cgcgctgcgc ggccaccgc tggtcgcgc      60
cgcgacggtg gtccccatcc acaactgcac tgcgctgggc gccgggatcg tcgtaaccgg      120
cagcggcgca gagcaatttg acgactccac acccggcgcg ctgcgcgccc atctcgcctg      180
ccgactcccc cagtacatga tccccaaagt gttcgtctca tgccccgagc tacgctcac      240
cgccaacggc aaggtcgacc ggggcaaaat cgcggcgcgc ctggaagcgg ccgcgcgggc      300

```


-continued

```

accccagccg ctcgacacgt catccaccct cactgtggtc gagcggctgg tcgcccaggt 360
ctggtccgat gtgctgggcg cgcgatcac cggccgcgag gacaatttct tcgcccaggg 420
cggcgatagc ctgcgcgccca ccgaagcggg cgcgcgactg acgcgcaggg gagtgccgg 480
agcggaggtg ggccagctgc tcagccacca gacgcttggg cagttcagcg cggcgtgtgt 540
gctcgcggac ccggcatccg aggcattccga gtcggcggcc gatgtcggcg aaccctgac 600
accggggcag gggttccgcg tcaccgggtt gcagcaggcc tacacgctgg gcgcggccgg 660
actcaatggg agcacctgtg caccaacgta ttccgctgtg gtgctggccg ccgcgcccga 720
gtctgccggg atagacctgg atcggtttgc ccgtgtggtc accagatgcg tcgacgaatt 780
cgcgatgctg cggtgccgcg tggacgccga caccacccaa cgggtgcagg tcgacgccgg 840
gccggtgccc gtccatgacc ttgatataca agacgacccc gacctgttac tgcggcgcat 900
ggcggccgcc ccgttcgacg cgcattcggg tccggtgatc cagtgtctcg caccgtcgag 960
gtcaccctgt cacgtcggtc tgtgatcag ctatctgggc ctgatgccc gcagcctgtc 1020
caccgtcgtc accacgatca tcgccgaata ccagtcgcaa ccccgccgcg gccaggtcga 1080
cccgaaccgc gcggtctctg cccggttgcg ctccgaaagc gcttggggcg aaaacgatgt 1140
cgacaacagc gttgccggcc ctccgctgct gccgctgca caccagcgac gtgaccctt 1200
cgagcgggtc accttcgcgc gccgcagctt caccatcga gaacaggctg ccgccacgct 1260
gcgtgagcac gccgcacacc tcggcgtcac ccccaccgcg ctggtctctg aagccttcgc 1320
acatgcgctg gcgtcgatcg gcgccgtca gcgattcgcg gtgacagtcc ccaagtctga 1380
ccgtcccgac tacgcccccg cggaccgcga ggtgttgggc aacttcacc gcctggcgct 1440
gtgcgaggtc gactacggcg ccgtgagacc gggatctgcc gaagcggttg ccgcggcgca 1500
gcggaactg tggcgcgcgg tgagccacga cggtgacatc accggcgggc tggccgcaac 1560
gcggaccgcg ggtggctacc ccgtggtgtt caccagcacc ctggggctca cccatcagga 1620
cgccagcggg ctgaccaacg tcgggacatt gaccagacc ccgggcgtct ggctggactg 1680
ccagaccgag gacgaggtcg ccggaattcg tatgagctgg gacatagcta ccaatgtggt 1740
tgcccgga tcatctcgg tggcttttc ccgattcag gaggcgggtc ggcgccacgc 1800
ggggcaagcc gagccgcgg gcacggccgt tgcctcggcg gtgggcg 1847

```

<210> SEQ ID NO 4

<211> LENGTH: 4716

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 4

```

agtgggcgag cgcggtgatc gccccgcgcg tcgccactg ccgaccgcg caggtgctgc 60
cgcaatacac catgctgggt cggcgtggg aggcactgcg atacgtccc tcaggatagc 120
ccgcttccga cgtcgaacgg gcggctcgcg ggttggccgg catcgtcacc ggagccgtgt 180
caccaccgac gctgatcgga gaccgcgacg tgacgcccga ggcgctgctg ctgctgacg 240
accgatgctg gatggccctc gacgacctg ccggggccat ctccgggca cgcgcgacgc 300
tggggcggcg gcttcgcgct gtggaagtgc gctcccgac cggcctgatc accgaacggg 360
tgaccgagct ggtgggcgtg gtggtcgagg aatacttgtg cctggagcca aaccgcagc 420
tcgccggaat cgcgcgggg cggcgttcc cggccccgac ccgccacgtc gacgaccgc 480

```

-continued

acgcggcgtc	gggcgtcgac	gtggtgatct	gctgtgggtc	gctgcatcag	ctgcccgcag	540
cggaggcggt	cctcgaggcg	atcacctgt	ccgacgcagg	ttggctgtgg	atggtggaga	600
attccgaggg	cacccaagcg	acgctcatca	gcgcgccgt	tctcgacccc	ggcctgctcg	660
cgccgatc	gaagacgctg	cgtccggcgg	atcgggtggg	gcggctcacc	gccgaccacg	720
gttgccgacc	gacgcacatg	atccaggacg	gacccggcct	cacgctcacc	gcgcaccgcc	780
ccgacaagcc	cgccatgccg	acaccgccgg	ccgaacagcg	ccgcgacggg	aggtggctcg	840
gaccggctgt	gccggcgctg	tcgctgccga	ccgacgccac	gggtggggcc	acgcttgccc	900
agatctggca	gcgtcatctc	gccattccaa	cacccggcgt	cgacgcagac	ttcttctctg	960
tcggcggtga	cagcctcgtc	gcgaccgggg	tctacgccga	ccttcgggcc	gccggttctg	1020
gccaaactcgc	tttcgctcgac	ctgttcaacc	actcgacgct	cggtgagctc	gcggcacacg	1080
ccggcccgcg	caccggcccg	gaagtgtcgg	tggcggtga	gtcgaccggg	ggcggaaccc	1140
acgacccgaa	ccgattcccg	ctcacctcg	tcgagaacgc	gtatcgggcc	ggcgagagaag	1200
gcgcgttgat	cctcggcggc	gtcccgccg	actgctactt	cgagtctgag	ctcgcggact	1260
tcgaccggcc	gagattcgat	tcggccgcac	gccaaactcg	agcacgccac	gccggactgc	1320
gcaccaacggg	gtcacccggg	ggcaaccgac	cgccctctc	gggtgaggtc	gccgctgctg	1380
acaccgcgcc	gatcgagccc	gtcgtgcgag	accacgcagc	cggtcgagcc	gcgatgcgcg	1440
accagatcat	cgacttgacg	gcccccgg	gcacgactt	cggtgtgcaa	accgcggcgg	1500
acggggcgac	cgctgtcggc	atcagcatgg	acaacaccat	gctcgacggc	gccagcatga	1560
tgatcgccct	gtccgaaact	gatcacctct	atcgcggcga	aaccggtgac	caattgccgc	1620
cgctggaaaac	gtctttcggc	cactacgtgt	ggaaccaccc	ggagctgctg	cccgcgcgg	1680
acgaggcggt	gctcccggg	ctggccgcca	gccgagacta	ttggcgcgca	cgccctgccat	1740
ctttgccgcc	ggcggcga	ttggccgaca	tgtcaactgct	gttcgagatc	gaggagccga	1800
ggttcgaacg	ggcaaccggc	accattcccg	ccgtcgactg	gtcgcaggta	acgcgatcgt	1860
gccgtgccga	ggcgctcacc	gtcgcgtcat	ttctgctcgc	caactatgca	cggtgtctgt	1920
ctcgtgtgtc	ggggaccgac	cacttctgca	tcaacgtcac	gctattcgac	cgcgaccccg	1980
atgtcgtggg	gatcgaaaac	gtcgtcggag	atttcaactc	cctggtgttg	ttggagtgcc	2040
gagtcgatga	gcccgctcgg	atctggggaga	gcgtgcgcgc	tctgcagcgg	caattgatga	2100
ccgacctgcc	gcaccgcggc	gcggacgcgg	tgtggtgca	acgcgaactg	ctgcggtttc	2160
acggcaaccc	gacggcggc	ctgtttcccg	tcgtcttacc	cagcggactg	ggccttgctg	2220
acgcctcggc	tcggggcgcg	gtccggttcg	ccgaaccggg	attcggccgc	tcgcagacac	2280
cgcagacggg	gttggacttc	caggtgtggg	aaagcgcggg	ggcgtgaaag	ctgtcgtggg	2340
acttcgtcag	tcaggcgggt	tcgcccggca	ccgcgcgcac	tcagctcgag	tcgttggtgg	2400
acggcatcac	cggtgtgcc	acaccgcagc	gccgcacga	acacaagtgg	ggcgaggggg	2460
catccaatga	cgagctcctg	caacgtgttt	cgaggatctg	cgctccgc	ctgggtcagc	2520
cgagggtcga	acctcacgac	aatttcttcc	agctcggcgg	cgattcggtc	agcgcgacca	2580
aggtggctga	acagatcggc	cgtgagctgt	cagcctcggc	cacccttcca	ctgctgttcg	2640
ccaatccggg	gatcggcgac	ttccggcca	aaatcgcca	cactgacaac	gccgacgaac	2700
ccgacctgac	cgttgaggag	ggcatgttat	gaccgcggcc	gagctcgtcg	accacctgcg	2760

-continued

```

gggtatcggc gtccaactgt gggccgacgg tgagaatttg cgctaccgag caccgcaaca 2820
agtcctcacc gcggaacctga aagctcaact ggcggcggtc aaaacggacg tgattaccct 2880
gctggcggaa gagacgaccc tgtgcgcgcc gccgcaggac cggttcgagc cgtttccgct 2940
caccgacgtg caagccgcat atctggtcgg gcgcacgtcg gcgtttcagt ggggccccgg 3000
aggctgccac ggctacgcgc agttccggtt cgaccacacc gtggcaaacac cgagcgccga 3060
gcaatatacg gaggcgtggc gcaaggttgc cgaccgccac gacatgttgc gctgctcgt 3120
tcatcccagc gggatcaggc tgatatgcc cgacgtgccc gacgacgggc tggatcatca 3180
tcagtgtcac acggtcgaag acgttgcccg cgtacgtgcc ggggtcacgg aacatctgcg 3240
taaccggata tatecgctgg gcgaagcgcc gatgtatgac ctggtgatca cgatgggccc 3300
tgacgacacc gtggtccatc tgtccgttga cctgctgac gccgacttcg tcagcatctc 3360
catcctgatg accgactttc agcagtgctt gcttgacccc gaatgcgacc ttgcgcccgt 3420
cgatttcagc ttccgcgact acctgtgaa tctcgctcgc gagcgaagct cggcccgccg 3480
tagtgcccgc cgggaacgcg atctcgcta ctggcgggat cggctcgatc agctgcccgc 3540
accactgtcg ttaccggtgc tgcccacga ctgacgcgc tcagcgtcgc ggacggcacc 3600
gaccttttca cggcgcttga tgccgctgtc ggcgcgagct ttcgaggtgc tcagccggcg 3660
cgcgccagag cacggagcga ccgtcaacgt cgccatcgtg acggcattca gccgggccc 3720
cgcacgctat ggcgatcgcg accatttcc tctgaccctg acgaccatgg accgccacgc 3780
gttcaactcc gccgtcgggc agttggtcgg ggatttcacc ggcaccagcg tgctcgaagt 3840
cgatgtccgc ggacagcgca ccttcgccga actgctgac ggcgtcggtg atcgtctctt 3900
cgacgacatg gaccattoga ccaccggcgg cgtcaacgtc gcccgctcc tggggcagcg 3960
cgacgacgac cgaggtgagc agacgcccgt cgtgttcacc tcgacactcg gcgccacgac 4020
ccgaatcgac agcggcgcaa catctttgtt gcaccgatt caaggtcgcg gtctgagcca 4080
gactcctcag gtgctgctgg actgtcaggt cgccgaaatc gacggcatgc ttgaggtcaa 4140
ctgggacacc cgcgatcagg ccgtgcccgc cgaggtcctc gaccgcgctg tcgccgactt 4200
ccgccacgcc ctggatctgc tcagcaccga cgcctcgccc tggcaccggc cgctattgcc 4260
ggcccagccc cccgagacca caccggtcga ggggccacgc acacaccatg aaccggcgct 4320
gatccatacc gggttcctgc gcaacgtgct ggtgacaccg gatgcccgtc ccatccgcca 4380
cggatgacgg gccaccacgt acgccgaatt gctcgtgccc gcgaccgcgg tggccgacac 4440
gctggccgcg acgggggtgc ggcgcgctga ctacgtcggg atccgggtgc cgcagggccc 4500
cgcccagatc gcggctctcc tgggcgcatt gctggcccgc gccgctatg tgccgttggc 4560
cgtcggctgg cccactcacc gcgtcgacca gatcgcgccc caatgctcgc tggcagcgct 4620
gtgcgagcgc gacggagagg tggaccggct gcttgcggac ccgcagacct ggtaaccgcg 4680
cgcgccgggtg gtgcccggagc cccacagcga ggtgct 4716

```

<210> SEQ ID NO 5

<211> LENGTH: 2392

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 5

```

ccgactccgg taaccagcaa cccgatctgg gctcgggtga tgatttcacc ccaggacgca 60

```

-continued

agggcgctcc aaccctcaaa gtggttcacc tcgggctgat gtggggacaa cacggggccg	120
gcatcgggtgg ccttgtcaaa aaccggttcg tgaaaatggt cgtatccaaa gatcagggtcg	180
gccccatac attcggcttg gatgaccgcg gcaecgccagg tccggtagtt cgggggtccc	240
gccggctgga tctgaacacc aacggaggag atcattgccc gccacccggc tgtcgatccc	300
gcgaggggttg cagtcgggtg gtgatccagt cgaggatgat tgactgggta tctgcccgtg	360
agcccatgtg cccgggtgca aagaaccgcg ccgatttggg ggaccctgfc tcgagcagca	420
ggtagtagtc gctgatcggg aatacgggtg cgtgtagccc gttgatgagc agcgtctcag	480
cgaggggccc gtccaggata ccttggcgta caagggacag ctccggcgcg tagtccaccc	540
attgctcgaa ggtgtcgcgc ccaaatgcat aggcgagggg ctccggccagc tcgaaggggt	600
actcgcctga ttgggcctgc tcgatccact cggcgtcaaa gccatggtcg atgcagccgc	660
cttggctgac cacacaggcg agggagttcc ggtgggtgtg ggcgattttg gccgcccagt	720
agccaccggt actaccaccc caatagocca caccgcgccg gtcgagttcg ggtcggccgg	780
cgatccactc caacacgggg gtgaacattc gctcggcgtc ctccgaacca gccagcggag	840
catcaccgac cccggggatg tcgatcgcca atggtgcgac gttccgggcg agcacctgat	900
cggtgtgcat gtcttctctg aaggtgtcga tcccggccga gacgagcaat acaggcagcc	960
ggtccgctgc cttcgggaca cgcagatggg cgatgatgcg gtcaccctcc ccgggcccgc	1020
cggcgaacgg gatctcgaca cgtcctgatg ggatgtcgaa gtaccgggat gcgcgcagca	1080
acatttctg cgacttcggg taggcgctct tcttaccggc cgaattcctc gtcgggttagc	1140
gggcatgctg gtagtagcca taggcgcgca aatagttttc gtgtgccaca ctggtgtcgc	1200
caatctgttc ggcctctctg gctcgggctt gatagcgtgc ggccacggca ctgaaccgcg	1260
ccgcccaggc gtcgcgatca taggagttca ggccgctcaa cacgccctgc acatcatcgg	1320
cgagcgtgta ctgaaacgga taaattcccg ccgtggccga gaccgccata gcggccatag	1380
ctcggtcacc ggtcgcgtgt gtcgcttcc acggccaagt tatgaccgct cgcggctgca	1440
gatattggag aaaacggaca tctcacgctg gccgtgggcg agccgcaccg ggaagttctg	1500
tgggagccgc tccagccagt gccaggcatt cgctgatcag gtgggactga tcggcgtaac	1560
ccgcctcgat cgctagatcg gccaccgga ttcggtgatc accggcatgg atcaaatcca	1620
gcagccgctg gaaccggaaa atccggccaa gccgcttggg gccgaagccg acccggcgt	1680
cgaaccgacg gcgaagctgc cgctcgtca acccaacaga ggaggcgaca tccgaaacag	1740
agcgggcccg gtggacacgc agcatctcga ttgagtgcgc cactggctga tcgaccacgg	1800
gatcgatctt gaccacatac cgcgccagca ccgcccga taatgcgacc cgttgccgaa	1860
aagcgttggc ctccagcaca tctcagcca atccggacgc ggatgtcccg aacaccgagt	1920
caaccggat ctgggtatct cttaactcgc tgacaggatg gccacgacc gcggcggcag	1980
caccggctcg taaccgcaga ccgatcatcg caccttcggg gcccgccatc tggctgtaaa	2040
aggtgtggc cggaccagaa accatgactg cgccctcggc ggtgacgaac agatcgacgc	2100
aaccgtccgg catcaccgcg agcgcgccac ccgcggtatt cgagcgcacc cagccacagt	2160
ccacatacgg cgcgagcgca cccaaaggtc gacactccaa ataccgacc atcacgctca	2220
cattcagcca ggtcctacgt cgctccgcaa caaatgggc cgcaatcctt cgaagccgcc	2280
cggtcagcgc agactcccag gcggcaatac tcgtgaaccg taaactgcgg cgtagaggaa	2340

-continued

atgcccgcca gggcgcaaca cccgcgccac ttcggcgaga aagcgtgaca gg 2392

<210> SEQ ID NO 6
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 6

cggtgcgcgg tagga 15

<210> SEQ ID NO 7
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 7

cgacctggtc tgcgtcatc 19

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 8

ccgcgaggac ctgctgacgc 20

<210> SEQ ID NO 9
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 9

gcgatccggc gaaacg 16

<210> SEQ ID NO 10
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 10

gacgagttag acgcacagat tct 23

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

-continued

<400> SEQUENCE: 11

acgccctgca gttagctccc cg 22

<210> SEQ ID NO 12

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 12

atcgaccggc tgggtgac 18

<210> SEQ ID NO 13

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 13

gcgagcagca cgttgaac 18

<210> SEQ ID NO 14

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 14

tcgacgcgat gaccgcgctc g 21

<210> SEQ ID NO 15

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 15

cagacacctg cagagctact ac 22

<210> SEQ ID NO 16

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 16

agcatgtcgg cgacgtt 17

<210> SEQ ID NO 17

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

probe

<400> SEQUENCE: 17

caagcccgac tacgcccgc t 21

<210> SEQ ID NO 18
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 18

tcgcggtgcc gatcag 16

<210> SEQ ID NO 19
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 19

gtttggatgc ccgtgacatt c 21

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 20

ccgcgctccc cgaaatgctc 20

<210> SEQ ID NO 21
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 21

cggcgaacc gtgaca 16

<210> SEQ ID NO 22
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 22

cacagtgct cccattgagt 20

<210> SEQ ID NO 23
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 23

ccgcgcccag cgtgtaggc 19

<210> SEQ ID NO 24
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 24

gcgcagcttc accatcga 18

<210> SEQ ID NO 25
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 25

gtgcgaaggc ttcgaagac 19

<210> SEQ ID NO 26
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 26

aacaggctgc cgccacgct 19

<210> SEQ ID NO 27
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 27

gccacgacgg tgacatca 18

<210> SEQ ID NO 28
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 28

ccagggtgct ggtgaaca 18

<210> SEQ ID NO 29
<211> LENGTH: 19

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 29

ccgcggttgcg gccagcccg 19

<210> SEQ ID NO 30
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 30

ggaatcgatc tcggtggctt tt 22

<210> SEQ ID NO 31
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 31

ccaccgccgg agcaa 15

<210> SEQ ID NO 32
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 32

ccgcgtggcg ccgca 15

<210> SEQ ID NO 33
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 33

gccgcgacgg taggt 15

<210> SEQ ID NO 34
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 34

ccagatctcg gcaagcgt 18

-continued

<210> SEQ ID NO 35
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 35

acgacgccgg cacagcc 17

<210> SEQ ID NO 36
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 36

cggacggaga ggtggac 17

<210> SEQ ID NO 37
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 37

gggctccggc acca 14

<210> SEQ ID NO 38
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 38

ttgccgacc gcagacctg 19

<210> SEQ ID NO 39
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 39

ggtaaccagc aaccgatct g 21

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 40

cccgaggtga accacttga 20

-continued

<210> SEQ ID NO 41
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 41

tcggttgatg atttcacccc aggacgc 27

<210> SEQ ID NO 42
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 42

ggcgtcctcc gaacca 16

<210> SEQ ID NO 43
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 43

gcaacattgg cgatcgacat c 21

<210> SEQ ID NO 44
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
probe

<400> SEQUENCE: 44

cagcggagca tcaccgaccc cgg 23

<210> SEQ ID NO 45
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 45

ggctcgggcc tgatagc 17

<210> SEQ ID NO 46
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 46

-continued

```

geggcctgaa ctctatgat c                               21

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 47

ccgcgttcag tgccgtggcc                               20

<210> SEQ ID NO 48
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 48

gcaatactcg tgaaccgtaa actg                           24

<210> SEQ ID NO 49
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      primer

<400> SEQUENCE: 49

gtggcgcggg tgttg                                    15

<210> SEQ ID NO 50
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      probe

<400> SEQUENCE: 50

ccctggcggg catttcctct acg                           23

```

What is claimed is:

1. An isolated nucleic acid sequence having the nucleic acid sequence of SEQ ID NO:1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 or SEQ ID NO:5, fragments thereof, complements thereof, sequences having at least 90% nucleic acid sequence identity thereto and labeled derivatives thereof.

2. A set of oligonucleotide primers comprising: at least one forward primer and at least one reverse primer, that are operable to hybridize to a MAP-specific target nucleotide sequence that is present in a MAP microorganism and is absent from a non-MAP organism, the pair of primers operable for simultaneous use in a PCR process for the detection of a *Mycobacterium avium* subsp. *paratuberculosis* (MAP) microorganism.

3. The set of oligonucleotide primers of claim 2, wherein the MAP-specific target nucleic acid sequence comprises

nucleic acid sequence of SEQ ID NO:1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 or SEQ ID NO:5, fragments thereof, complements thereof and sequences having at least 90% nucleic acid sequence identity to the foregoing sequences.

4. The set of oligonucleotide primers of claim 2, wherein the at least one forward primer and the at least one reverse primer are selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO:

48 and SEQ ID NO: 49; complements thereof and primers having about 90% identity to the foregoing sequences.

5. A method for differentially detecting MAP from other non-MAP microorganisms comprising: detecting, in a sample, at least one MAP-specific target nucleic acid, wherein detection of the at least one MAP-specific target nucleic acid sequence indicates the presence of an MAP organism in the sample.

6. The method of claim 5, wherein the at least one MAP-specific target nucleic acid sequence comprises a nucleic acid having SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 or SEQ ID NO:5, fragments thereof, complements thereof and sequences having at least 90% nucleic acid sequence identity to the foregoing sequences, wherein fragments thereof comprise a nucleic acid sequence having at least 10 to at least 25 contiguous nucleic acids of SEQ ID NO:1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 or SEQ ID NO:5.

7. The method of claim 5, wherein the detecting comprises:

amplifying MAP-specific target nucleic acids from the sample comprising:

obtaining nucleic acids from the sample;

contacting the nucleic acids from the sample with at least one primer set, having one forward primer and one reverse primer that can hybridize to and amplify at least one MAP-specific target nucleic acid, under conditions suitable for amplification; and

detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid amplified by the primer set confirms the presence of a MAP organism in a sample.

8. The method of claim 7, wherein the at least one primer set comprises at least one forward primer and the at least one reverse primer selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; complements thereof and primers having about 90% identity to the foregoing sequences.

9. The method of claim 7, wherein the step of detecting an amplified nucleic acid comprises using a probe that can hybridize to and reveal the presence of the amplified nucleic acid.

10. The method of claim 9, wherein the probe is selected from SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof, sequences having about 90% identity to the foregoing sequences and labeled derivatives thereof.

11. The method of claim 10, wherein the probe having SEQ ID NO:8 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 6 and SEQ ID NO 7; the probe having SEQ ID NO:11 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 8 and SEQ ID NO 9; the probe having SEQ ID

NO 14 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 12 and SEQ ID NO: 13; the probe comprising SEQ ID NO:17 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 15 and SEQ ID NO 16; the probe having SEQ ID NO:20 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 18 and SEQ ID NO 19; the probe having SEQ ID NO 23 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 21 and SEQ ID NO:22; the probe comprising SEQ ID NO:26 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 24 and SEQ ID NO 25; the probe having SEQ ID NO:29 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 27 and SEQ ID NO 28; the probe having SEQ ID NO 32 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 30 and SEQ ID NO:31; the probe comprising SEQ ID NO:35 is used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 33 and SEQ ID NO 34; the probe having SEQ ID NO:38 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 36 and SEQ ID NO 37; the probe having SEQ ID NO 41 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 39 and SEQ ID NO:40; the probe comprising SEQ ID NO:44 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 42 and SEQ ID NO 43; the probe having SEQ ID NO:47 is used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 45 and SEQ ID NO 46; or the probe having SEQ ID NO 50 is used to detect an amplified nucleic acid amplified using primer set having SEQ ID NO: 48 and SEQ ID NO:49.

12. The method of claim 7, further comprising using at least two primer sets, each primer set having one forward primer and one reverse primer, that can hybridize to and amplify the MAP-specific target nucleic acid, under conditions suitable for amplification; and detecting at least two amplified nucleic acids, wherein detecting at least two amplified nucleic acids amplified by the at least primer sets confirms the presence of a MAP organism in a sample and excludes the presence of a non-MAP organism in the sample.

13. The method of claim 5, wherein the sample is an veterinary sample, an animal sample, a food sample, an agricultural sample, a produce sample, a clinical sample, an environmental sample, a biological sample, a water sample or an air sample.

14. The method of claim 13, wherein the animal sample is obtained from a cow, a sheep, a bison, a ruminant, a deer, a rabbit, a fox a bird.

15. The method of claim 5 further comprising:

detecting the presence of at least two MAP-specific nucleic acids comprising:

- a) amplifying from the sample a first MAP specific nucleic acid or a fragment or a complement thereof by contacting nucleic acids present in the sample with at least a first primer set, having one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid or a fragment or a complement thereof; and
- b) amplifying simultaneously from the same sample a second MAP specific nucleic acid and/or a fragment or a complement thereof by simultaneously contact-

ing nucleic acids present in the sample with at least a second primer set, having one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid or a fragment or a complement thereof, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction; and

- c) detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample.

16. The method of claim **15**, further comprising detecting the first and the second amplified nucleic acids amplified by the amplification reactions of steps a) and b).

17. The method of claim **5** performed using a kit for the detection of a MAP organism, the kit comprising:

at least one primer set having one forward primer and one reverse primer that can hybridize to and amplify a MAP-specific target nucleic acid, under conditions suitable for amplification; and

optionally at least one probe, the probe operable to hybridize to the amplified MAP-specific target nucleic acid amplified by the at least one primer set or to a MAP-specific target;

one or more components selected from a group consisting of: at least one enzyme, dNTPs, at least one buffer, at least one salt, at least one control nucleic acid sample; and an instruction protocol.

* * * * *