



US 20240091160A1

(19) **United States**

(12) **Patent Application Publication**
Murphy et al.

(10) **Pub. No.: US 2024/0091160 A1**

(43) **Pub. Date: Mar. 21, 2024**

(54) **PHARMACEUTICAL COMPOSITION
CONTAINING ACETAMINOPHEN AND
IBUPROFEN**

continuation of application No. 16/287,836, filed on
Feb. 27, 2019, now Pat. No. 11,197,830.

Publication Classification

(71) Applicant: **AFT Pharmaceuticals Limited,**
Takapuna (NZ)

(51) **Int. Cl.**
A61K 9/28 (2006.01)
A61K 9/00 (2006.01)
A61K 31/136 (2006.01)
A61K 31/192 (2006.01)

(72) Inventors: **Maura Murphy**, King of Prussia, PA
(US); **Matt Callahan**, Haverford, PA
(US)

(52) **U.S. Cl.**
CPC *A61K 9/2893* (2013.01); *A61K 9/0053*
(2013.01); *A61K 31/136* (2013.01); *A61K*
31/192 (2013.01)

(21) Appl. No.: **18/518,202**

(22) Filed: **Nov. 22, 2023**

Related U.S. Application Data

(57) **ABSTRACT**

(63) Continuation of application No. 18/087,608, filed on
Dec. 22, 2022, now Pat. No. 11,872,317, which is a
continuation of application No. 17/548,877, filed on
Dec. 13, 2021, now Pat. No. 11,534,407, which is a

Solid oral dosage forms containing 325 mg of acetamino-
phen and 97.5 mg of ibuprofen or 500 mg of acetaminophen
and 150 mg of ibuprofen, wherein the ibuprofen has a [D50]
between 1 and 9 μ m, are described.

**PHARMACEUTICAL COMPOSITION
CONTAINING ACETAMINOPHEN AND
IBUPROFEN**

RELATED APPLICATIONS

[0001] This application is a continuation and claims priority to U.S. application Ser. No. 18/087,608, filed on Dec. 22, 2022, which is a continuation and claims priority to U.S. application Ser. No. 17/548,877, filed on Dec. 13, 2021, (U.S. Pat. No. 11,534,407) which is a continuation and claims priority to U.S. application Ser. No. 16/287,836, filed on Feb. 27, 2019, (U.S. Pat. No. 11,197,830) the entire contents of which are hereby incorporated by reference.

BACKGROUND

[0002] For various non-steroidal anti-inflammatory drugs (NSAIDs) there is evidence that high, early plasma concentrations result in better early pain relief, better overall pain relief, longer lasting pain relief and lower rates of remediation. (Moore et al. (2015) *British Journal of Clinical Pharmacology* 80:381). For ibuprofen, more rapid absorption can result in earlier and higher maximum plasma concentrations and this results in earlier onset of analgesia and better overall and longer lasting analgesia in dental pain models. The effect of fast acting formulations can be significant. In one study 200 mg of a fast acting formulation, a ibuprofen salt produced the same or better analgesia as 400 mg ibuprofen acid, and with a reduced requirement for additional analgesic use (Moore et al. (2014) *Pain* 155:14). It is possible to alter the pharmacokinetics of ibuprofen by reducing the particle size to below 1000 nm (WO 2010/121328)

SUMMARY

[0003] Described herein is a solid oral unit dosage pharmaceutical composition containing acetaminophen (325 mg) and ibuprofen (97.5 mg) having a higher maximum plasma concentration for ibuprofen compared to Maxigesic (325). Also described is a solid oral unit dosage pharmaceutical composition containing acetaminophen (500 mg) and ibuprofen (150 mg). These dosage forms are referred to as Rapid Maxigesic 325 and Rapid Maxigesic 500, respectively. This disclosure features these pharmaceutical compositions as well as methods for producing and using such compositions.

[0004] The solid oral dosage forms described herein include ibuprofen (free acid) that is particularly rapidly dissolving in vitro. The impact of increased in vitro dissolution rates on pharmacokinetic parameters is highly variable. In the case of the present dosage forms, the rapidly dissolving ibuprofen results in a product with meaningfully higher C_{max} for ibuprofen. Significantly, the ibuprofen in the present dosage forms has a particle size that is larger than 1,000 nm.

[0005] Described herein is a solid oral dosage form comprising 325 mg of acetaminophen and 97.5 mg of ibuprofen or 500 mg of acetaminophen and 150 mg of ibuprofen, wherein the ibuprofen has a [D50] between 1 and 9 μ m.

[0006] In various embodiments: the ibuprofen has a [D10] between 1 and 3 μ m; the ibuprofen has a [D90] between 3 and 16 μ m; the ibuprofen has a [D50] between 2 and 8 μ m, a [D10] between 1 and 3 μ m and a [D90] between 4 and 16 μ m; the dissolution rate of ibuprofen in the solid oral dosage

form is such that, when tested using USP Apparatus II (paddles) set to rotation speed of 50 rpm in 900 mL of 50 mM pH 5.8 phosphate buffer at 37° C., wherein 80% or more (e.g., at least 85% to 95% or 95% to 100%) of the ibuprofen dissolves in 15 minutes or less; the dosage form is a tablet (e.g., a coated tablet); the [D90] to [D50] ratio is between 4:1 and 1.5:1; the [D90] to [D50] ratio is between 3:1 and 1.5:1; the [D50] to [D10] ratio is between 4:1 and 1.5:1; and the [D50] to [D10] ratio is between 3:1 and 1.5:1.

[0007] In some cases, the dosage form is prepared by a process comprising jet milling a composition comprising 21-23% wt/wt ibuprofen and 73-75% wt/wt acetaminophen. In some cases, the composition subjected to jet milling further comprises a surfactant (e.g., sodium lauryl sulfate).

[0008] Also described is a method for treating pain (e.g., mild to moderate acute pain) comprising administering a dose of 1, 2 or 3 units of the solid oral dosage form comprising 325 mg of acetaminophen and 97.5 mg of ibuprofen. In some cases, the administration is 1, 2, 3 or 4 times daily.

[0009] Also described is a method for treating pain (e.g., mild to moderate acute pain) comprising administering a dose of 1 or 2 units of the solid oral dosage form comprising 500 mg of acetaminophen and 150 mg of ibuprofen. In some cases, the administration is 1, 2, 3 or 4 times daily.

[0010] Also described is a method for preparing a pharmaceutical composition, comprising jet milling a composition comprising acetaminophen, ibuprofen and a surfactant (e.g. sodium lauryl sulfate), wherein the ratio of acetaminophen to ibuprofen is 3:1 (w/w), under conditions and for a time to reduce the particle size of the ibuprofen to a median particle size, on a volume average basis between 2 and 8 μ m.

[0011] Also describes is a method for preparing a pharmaceutical composition, comprising jet milling a composition comprising acetaminophen, ibuprofen and a surfactant (e.g. sodium lauryl sulfate), wherein the ratio of acetaminophen to ibuprofen is 3:1 (w/w), under conditions and for a time to achieve a median bulk particle size between 4 and 15 μ m.

[0012] In some cases: the [D90] of the ibuprofen in Rapid Maxigesic 325 and Rapid Maxigesic 500 is greater than 2 μ m (e.g., greater than 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 μ m) and less than one of: 18 μ m, 17 μ m, 16 μ m; 14 μ m, and 13 μ m (preferably 3-17 μ m, 3-16 μ m or 4-15 μ m); the [D50] of the ibuprofen greater than 1 μ m, 1.5 μ m, 2 μ m, 2.5 μ m, 3 μ m, 3.5 μ m, 4 μ m, 4.5 μ m, 5 μ m, 5.5 μ m, 6 μ m), but less than 12 μ m, 10 μ m, 9 μ m, or 8 μ m (preferably 1-11 μ m, 1-10 μ m, 2-10 μ m, 2-9 μ m); the [D10] of the ibuprofen is less than 7 μ m (e.g., 6.5, 6, 5.5, 5, 4.5, 4, 3.5, 3, 2.5, 2, or 1.5 μ m), but greater than 100 nm; the ratio of [D90] to [D50] is between 4:1 and 2:1; and the ratio of [D50] to [D10] is between 3:1 and 1.5:1 (all measurements on a volume average basis).

[0013] Also described herein is a method for treating pain, e.g., mild to moderate acute pain, comprising administering to a patient in need thereof a therapeutically effective dose (acetaminophen 975 mg+ibuprofen 292.5 mg) of the Rapid Maxigesic 325 unit dosage form (acetaminophen 325 mg+ibuprofen 97.5 mg/unit dose) up to 4 times a day (acetaminophen 3900 mg+ibuprofen 1170 mg/day).

[0014] Also described herein is a method for treating pain, e.g., mild to moderate acute pain, comprising administering to a patient in need thereof a therapeutically effective dose (acetaminophen 1000 mg+ibuprofen 300 mg) of the Rapid Maxigesic 500 unit dosage form (acetaminophen 500

mg+ibuprofen 150 mg/unit dose) up to 4 times a day (acetaminophen 4000 mg+ibuprofen 1200 mg/day).

[0015] Additional dosage regimes for Rapid Maxigesic 325 include: 1 unit dose given 1, 2, 3 or 4 times daily; 2 unit doses given 1, 2, 3 or 4 times daily; and 3 unit doses given 1, 2, 3 or 4 times daily.

[0016] Additional dosage regimes for Rapid Maxigesic 500 include: 1 unit dose given 1, 2, 3 or 4 times daily and 2 unit doses given 1, 2, 3 or 4 times daily.

[0017] The dissolution rate of the ibuprofen in a coated tablet containing 97.5 mg of ibuprofen is such that when tested in 900 ml of pH 5.8 phosphate buffer (50 mM) using USP Apparatus II at 50 rpm and 37° C., is such that at least 75%, 80% or 85% ibuprofen dissolves in 15 min or less (e.g., 14 min or less, 13 min or less, e.g., 85% can dissolve in 15 min). For example, at least 85% can dissolve in 12-19 minutes.

[0018] The dissolution rate of the ibuprofen in a coated tablet containing 150 mg of ibuprofen is such that when tested in 900 ml of pH 5.8 phosphate buffer (50 mM) using USP Apparatus II at 50 rpm and 37° C., is such that at least 75%, 80% or 85% ibuprofen dissolves in 15 min or less (e.g., 14 min or less, 13 min or less, e.g., 85% can dissolve in 15 min). For example, at least 85% can dissolve in 12-19 minutes.

[0019] The dosage form can include various excipients. For example, the dosage form can include one or more of: a diluent, lubricant, disintegrant, binder and wetting agent. For example, the dosage form can include one or more of: magnesium stearate, povidone, lactose, microcrystalline cellulose pregelatinized starch, hypromellose, sodium starch glycolate, sodium starch fumarate, sodium lauryl sulfate, and croscarmellose sodium. The tablet can be uncoated or, preferably, coated with a suitable coating agent.

[0020] The dry milling apparatus used is preferably a jet mill (e.g., a spiral jet mill).

[0021] In another aspect, the disclosure comprises a method for manufacturing a pharmaceutical composition as described herein comprising the step of combining a composition comprising ibuprofen and acetaminophen prepared by a method described herein or a composition as described herein, together with one of a diluent, lubricant, excipient, disintegrant, and wetting agent, to produce a pharmaceutically acceptable solid dosage form.

[0022] The disclosure described herein may include one or more ranges of values (e.g. size, concentration etc.). A range of values will be understood to include all values within the range, including the values defining the range, and values adjacent to the range that lead to the same or substantially the same outcome as the values immediately adjacent to that value which defines the boundary to the range.

[0023] Throughout this specification, unless the context requires otherwise, the word “comprise” or variations, such as “comprises” or “comprising” will be understood to imply the inclusion of a stated integer, or group of integers, but not the exclusion of any other integers or group of integers. It is also noted that in this disclosure, and particularly in the claims and/or paragraphs, terms such as “comprises”, “comprised”, “comprising” and the like can have the meaning attributed to it in US Patent law; e.g., they can mean “includes”, “included”, “including”, and the like.

[0024] “Therapeutically effective amount” as used herein with respect to methods of treatment and in particular drug dosage, shall mean that dosage that provides the specific pharmacological response for which the drug is adminis-

tered in a significant number of subjects in need of such treatment. It is emphasized that “therapeutically effective amount,” administered to a particular subject in a particular instance will not always be effective in treating the diseases described herein, even though such dosage is deemed a “therapeutically effective amount” by those skilled in the art. It is to be further understood that drug dosages are, in particular instances, measured as oral dosages, or with reference to drug concentrations as measured in blood.

[0025] Those skilled in the art will appreciate that the disclosure described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the disclosure includes all such variations and modifications. The disclosure also includes all of the steps, features, compositions and materials referred to or indicated in the specification, individually or collectively and any and all combinations or any two or more of the steps or features.

[0026] The present disclosure is not to be limited in scope by the specific embodiments described herein, which are intended for the purpose of exemplification only. Functionally equivalent products, compositions and methods are clearly within the scope of the disclosure as described herein.

[0027] Other aspects and advantages of the disclosure will become apparent to those skilled in the art from a review of the ensuing description.

DETAILED DESCRIPTION

[0028] Described herein are various rapid release, solid oral dosage forms containing 325 mg acetaminophen and 97.5 mg of ibuprofen with higher maximum plasma concentration of ibuprofen compared to a dosage form referred to as Maxigesic 325. Also described below are various rapid release, solid oral dosage forms containing 500 mg acetaminophen and 150 mg of ibuprofen with higher plasma concentration of ibuprofen compared to a dosage form referred to a Maxigesic 500.

[0029] Maxigesic 325 tablets contain acetaminophen (325 mg) and ibuprofen (97.5 mg) and Maxigesic 500 tablets contain acetaminophen (500 mg) and ibuprofen (150 mg). Both are given in single doses of up to three tablets for the lower strength and up to two tablets for the higher strength. The amount of acetaminophen delivered in a single dose is similar in both cases being either up to three tablets of Maxigesic 325: acetaminophen 975 mg and ibuprofen 292.5 mg or two tablets Maxigesic 500: acetaminophen 1000 mg and ibuprofen 300 mg.

Particle Size

[0030] For measurements made using a laser diffraction the term “median particle size” is defined as the median particle diameter as determined on an equivalent spherical particle volume basis. Where the term median is used, it is understood to describe the particle size that divides the population in half such that 50% of the population on a volume basis is greater than or less than this size. The median particle size is written as: $[D_{50}]$ or $D_{[50]}$ or $[D50]$, $D50$, $D(0.50)$ or $D[0.5]$ or similar. As used herein $[D_{50}]$ or $D_{[50]}$ or $[D50]$, $D50$, $D(0.50)$ or $D[0.5]$ or similar shall be taken to mean median particle size.

[0031] The term “ D_x of the particle size distribution” refers to the x th percentile of the distribution on an equiva-

lent spherical particle volume basis; thus, D90 refers to the 90th percentile, D95 refers to the 95th percentile, and so forth. Taking D90 as an example this can often be written as, [D₉₀] or D_[90] or [D90], D(0.90) or D[0.9] or similar. With respect to the median particle size and Dx an upper case D or lowercase d are interchangeable and have the same meaning.

[0032] The term “D(3,2)” is referred to as the area-weighted mean size or the Sauter diameter; the term “D(4, 3)” is referred to as the volume-weighted mean size. Detailed descriptions of how these values are calculated are known in the art and can be found in, for example, ISO 9276-2:2014(E).

[0033] For many of the materials subject to the methods of this disclosure the particle size can be easily measured. Where the active material has poor water solubility and the matrix it is milled in has good water solubility the powder can simply be dispersed in an aqueous solvent. In this scenario the matrix dissolves leaving the active material dispersed in the solvent. This size of the particles in the suspension can then be measured by laser light diffraction.

Medicaments

[0034] The medicaments of the present disclosure may include the pharmaceutically acceptable material, optionally together with a grinding matrix or at least a portion of the grinding matrix, combined with one or more pharmaceutically acceptable carriers, as well as other agents commonly used in the preparation of pharmaceutically acceptable compositions.

[0035] As used herein “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible.

[0036] Pharmaceutical acceptable carriers according to the disclosure may include one or more of the following examples:

[0037] (1) surfactants and polymers including, but not limited to polyethylene glycol (PEG), polyvinylpyrrolidone (PVP), sodium lauryl sulfate, polyvinylalcohol, crospovidone, polyvinylpyrrolidone-polyvinylacrylate copolymer, cellulose derivatives, hydroxypropylmethyl cellulose, hydroxypropyl cellulose, carboxymethyl-ethyl cellulose, hydroxypropylmethyl cellulose phthalate, polyacrylates and polymethacrylates, urea, sugars, polyols, and their polymers, emulsifiers, sugar gum, starch, organic acids and their salts, vinyl pyrrolidone and vinyl acetate

[0038] (2) binding agents such as various celluloses and cross-linked polyvinylpyrrolidone, microcrystalline cellulose, povidone; and or

[0039] (3) filling agents such as lactose monohydrate, lactose anhydrous, microcrystalline cellulose and various starches; and or

[0040] (4) lubricating agents such as colloidal silicon dioxide, talc, stearic acid, magnesium stearate, calcium stearate, sodium stearyl fumarate; and or

[0041] (5) sweeteners such as any natural or artificial sweetener including sucrose, xylitol, sodium saccharin, cyclamate, aspartame, and acesulfame K; and or

[0042] (6) flavoring agents; and or

[0043] (7) preservatives such as potassium sorbate, methylparaben, propylparaben, benzoic acid and its

salts, other esters of parahydroxybenzoic acid such as butylparaben, alcohols such as ethyl or benzyl alcohol, phenolic chemicals such as phenol, or quaternary compounds such as benzalkonium chloride; and or

[0044] (8) buffers; and or

[0045] (9) Diluents such as pharmaceutically acceptable inert fillers, such as microcrystalline cellulose, lactose, dibasic calcium phosphate, saccharides, and/or mixtures of any of the foregoing; and or

[0046] (10) wetting agents such as docusate sodium, sodium lauryl sulfate, phospholipids, poloxamer, polysorbate 80, sorbitan esters, tricaprilyn, glyceryl monooleate, myristyl alcohol and mixtures thereof; and or

[0047] (11) disintegrants; such as croscarmellose sodium, crospovidone, sodium starch glycolate, and or

[0048] (12) effervescent agents such as effervescent couples such as an organic acid (e.g., citric, tartaric, malic, fumaric, adipic, succinic, and alginic acids and anhydrides and acid salts), or a carbonate (e.g. sodium carbonate, potassium carbonate, magnesium carbonate, sodium glycine carbonate, L-lysine carbonate, and arginine carbonate) or bicarbonate (e.g. sodium bicarbonate or potassium bicarbonate); and or

[0049] (13) other pharmaceutically acceptable excipients.

Example 1: Attritor Milling and Jet Milling Improve Dissolution of Ibuprofen

[0050] Attritor milling of an active pharmaceutical ingredient (API), whether dry or wet, most often takes place in the presence of a milling matrix, i.e., one or more non-active ingredients that can assist in particle size reduction of the API. For example, in a dry milling process, the API can be milled in the presence of lactose or another suitable excipient. Such milling can be used to reduce the median particle size to less than 1,000 nm. Due to the high drug load desired for the present dosage form, the acetaminophen was used as the milling matrix for the attritor milling studies described below.

[0051] Ibuprofen, acetaminophen and sodium lauryl sulfate (SLS) were milled in an attritor mill using the conditions described in Table 1.

TABLE 1

Milling Conditions for Attritor Mill			
Ingredient	Formula	Processing parameters	
	% w/w	Attritor Mill	0.5 gallon, 1S
Ibuprofen	22.5%	Media: powder ratio	35:1
Acetaminophen (APAP)	75.5%	Milling speed	500 rpm
Sodium Lauryl Sulfate (SLS)	2.0%	Milling time	40 min
Total	100%	yield	51.8%

[0052] The attritor milling (Table 1) of a mixture of acetaminophen, ibuprofen and sodium lauryl sulfate at a 200 g scale was successful in that the median particle size of the ibuprofen was reduced to the submicron range (Table 2). Moreover, the dissolution rate of the ibuprofen in the milled

material, either attritor milled or jet milled, was faster than that of the unmilled API (Table 3). However, the attritor milling yield was relatively low due to caking of the milled material.

[0053] Co-jet milling was investigated as an alternative approach that might increase yield and improve scale. The formula in Table 1 was processed in a 4" jet mill with 100 psig pressure. As shown in Table 2, the ibuprofen within the co-jet milled formulation (acetaminophen/ibuprofen/SLS) was larger in size compared to the ibuprofen produced by attritor milling.

TABLE 2

Particle size of ibuprofen			
	Dv10 (μm)	Dv50 (μm)	Dv90 (μm)
Ibuprofen (starting material)	24.2	51.3	94.5
Attritor-milled intermediate	0.282	0.617	2.05
Co-Jet-milled intermediate	1.63	2.73	4.44

[0054] Both milled formulations had substantially faster ibuprofen dissolution compared to the unmilled formulation, demonstrating that particle size reduction improved the ibuprofen dissolution rate. However, while the attritor milled material had much smaller particle size for ibuprofen than the co-jet milled material, the dissolution rates of ibuprofen for both milled materials (Table 3) were similar. This suggests that the co-jet milled material could overcome the dissolution rate limited solubility of ibuprofen, despite the fact that the median particle size was greater than 1,000 nm.

TABLE 3

Ibuprofen dissolution rate in acetaminophen/ ibuprofen powder formulations						
Time (min)	Prior to milling		Attritor milled		Co-Jet-milled	
	% ibuprofen Dissolved	% RSD n = 3	% ibuprofen Dissolved	% RSD n = 3	% ibuprofen Dissolved	% RSD n = 3
2	34.4	4.4	82.5	5.5	96.0	1.3
5	49.2	6.6	94.9	1.7	103.0	0.4
10	62.3	7.4	100.1	0.5	103.5	0.4
15	68.3	9.6	103.5	0.8	104.7	1.2
30	73.1	8.8	103.1	0.2	105.6	0.3
45	75.8	6.7	103.3	0.9	104.0	0.3
Infinity	91.6	2.1	104.7	1.9	102.9	1.2

Methods

Particle Size

[0055] Particle size was measured by first dispersing the sample in 0.1% HPC, 0.01N HCl and sonicating for 1 minute with a probe sonicator. The sonicated samples were added to the laser light diffractor wet sample unit, filled with 0.1% HPC in 0.01N HCl, stirred for 5 minutes and then the sample was measured.

Dissolution Rate

[0056] The dissolution of selected powder intermediate formulations was tested using the USP Apparatus II, with 900 mL of 50 mM pH 5.8 media at 37° C. and a 50 rpm

paddle speed. The powder was added to the top of the media-filled vessels, and samples were assayed by HPLC to obtain the results.

Example 2: Co-Jet Milling Provides Improved Dissolution for Ibuprofen

[0057] The jet milling results suggested that it could provide a means for preparing a dosage form containing a high drug load of both ibuprofen and acetaminophen. Jet milling was further investigated by manufacturing a powder formulation using two different jet milling methods, both of which employed the formula in Table 4. In the first process, ibuprofen was jet milled alone and then blended with the acetaminophen, SLS and hydroxypropyl cellulose. In the second process, all four ingredients were blended and then co-jet milled.

TABLE 4

Formulation for comparisons of jet milling and co-jet milling Formula	
Ingredient	% w/w
Ibuprofen	22%
Acetaminophen (APAP)	74%
Sodium Lauryl Sulfate (SLS)	2%
Hydroxypropyl cellulose (HPC)	2%
Total	100%

[0058] The particle size of jet-milled ibuprofen, whether jet-milled alone or co-jet milled with acetaminophen, SLS and HPC was similar, as shown in Table 5.

TABLE 5

Particle size of jet milled and co-jet milled ibuprofen			
	Dv10 (μm)	Dv50 (μm)	Dv90 (μm)
Jet milled ibuprofen	1.46	2.53	3.97
Co-Jet-milled ibuprofen, APAP, SLS, HPC	1.31	2.51	4.32

[0059] It was unexpectedly found that when ibuprofen was size reduced by co-jet milling with acetaminophen, SLS and hydroxypropyl cellulose, the dissolution rate was improved relative to ibuprofen that was size reduced by jet milling in the absence of acetaminophen, SLS and hydroxypropyl cellulose despite the fact that the two methods yielded similarly-sized ibuprofen particles. Table 6 presents data on the dissolution rate of 1) jet-milled ibuprofen that was blended with acetaminophen, SLS and hydroxypropyl cellulose after jet milling; and 2) ibuprofen that was co-jet milled with acetaminophen, SLS and hydroxypropyl cellulose.

[0060] It can be seen from the dissolution data in Table 6 that the ibuprofen co-jet milled with acetaminophen, SLS and HPC fully dissolved within 2 minutes whereas the ibuprofen milled in the absence of acetaminophen, SLS and HPC took 10 minutes to exceed 85% dissolved. Hence, the presence of acetaminophen, SLS and HPC in the milling improve the dissolution rate of similarly sized ibuprofen.

TABLE 6

Dissolution data for jet milled and co-jet milled ibuprofen				
Time (min)	Jet-milled ibuprofen subsequently blended with acetaminophen, SLS, hydroxypropyl cellulose		Ibuprofen co-jet milled with acetaminophen, SLS, and hydroxypropyl cellulose	
	Avg % ibuprofen dissolved	% RSD (n = 3)	Avg % ibuprofen dissolved	% RSD (n = 3)
2	80.0	13.9	100.0	0.6
5	83.1	14.3	104.5	0.4
10	86.8	11.4	104.6	0.6
15	89.6	9.6	104.8	0.7
30	94.2	6.5	104.8	0.7
45	98.4	3.8	104.6	0.6

[0061] The particle size and dissolution rate methods were the same as those described in Example 1.

Example 3: Preparation of Examples of Rapid Maxigesic 325 Tablets and Rapid Maxigesic 500 Tablets

[0062] Acetaminophen (75.41% w/w), ibuprofen (22.62% w/w) and sodium lauryl sulfate (1.97% w/w) were blended together, and the blend was milled using a spiral jet mill to achieve a target bulk particle size of Dv50 <4.5 micron and Dv90 <14 microns. The milled blend (“Drug Product Intermediate” or DPI) was then blended with excipients and wet granulated using a high shear mixer and fluid bed dryer. The granulation was screened/milled, blended with lubricant and compressed into tablets. The tablets were then film coated. Table 7 shows the formulation of the milled intermediate, and Tables 8 and 9 show the tablet formulations.

TABLE 7

Example of Drug Product Intermediate for Rapid Maxigesic 325 and 500	
Ingredient	% w/w
Acetaminophen, USP	75.41
Ibuprofen, USP	22.62
Sodium Lauryl Sulfate, USP	1.97
	100.00

TABLE 8

Example of Tablet Formulation for Rapid Maxigesic (325) Rapid Maxigesic 325 Tablet (325 mg acetaminophen, 97.5 mg ibuprofen)		
Intragranular	% w/w	mg/tablet
Drug Product Intermediate	74.75	431.0
75.41% APAP		
22.62% Ibuprofen		
1.97% SLS		
Microcrystalline Cellulose (Avicel PH101)	9.00	51.9
Lactose Monohydrate (Lactose 310)	11.74	67.7
Croscarmellose Sodium (Ac-di-Sol)	2.00	11.5

TABLE 8-continued

Example of Tablet Formulation for Rapid Maxigesic (325) Rapid Maxigesic 325 Tablet (325 mg acetaminophen, 97.5 mg ibuprofen)		
Intragranular	% w/w	mg/tablet
Povidone (K30)	2.00	11.5
Purified Water	qs	—
extragranular		
Magnesium stearate	0.50	2.9
Total tablet core	100.00	576.6
Film coating		
Opadry II 57U18539, White	2.25	13.0
(hypromellose, titanium dioxide, polydextrose, talc, maltodextrin, medium chain triglycerides)		
Total		593.8 mg

TABLE 9

Example of Tablet Formulation for Rapid Maxigesic (500) Rapid Maxigesic 500 Tablet (500 mg acetaminophen, 150 mg ibuprofen)		
Intragranular	% w/w	mg/tablet
Drug Product Intermediate	74.75	663.0
75.41% APAP		
22.62% Ibuprofen		
1.97% SLS		
Microcrystalline Cellulose (Avicel PH101)	9.00	79.8
Lactose Monohydrate (Lactose 310)	11.74	103.8
Croscarmellose Sodium (Ac-di-Sol)	2.00	17.7
Povidone (K30)	2.00	17.7
Purified Water	qs	
extragranular		
Magnesium stearate	0.50	4.4
Total tablet core	100.00	887.0
Film coating		
Opadry II 57U18539, White	2.00	17.7
(hypromellose, titanium dioxide, polydextrose, talc, maltodextrin, medium chain triglycerides)		
Total		904.7

Particle Size

[0063] Several lots of DPI were analyzed to determine the particle size of ibuprofen within the co-jet milled intermediate. All testing was conducted using laser light diffraction. To measure ibuprofen particle size the DPI was dispersed in an aqueous media in which the acetaminophen and SLS are soluble (0.1% HPC in 0.01N HCl), leaving only the poorly soluble ibuprofen suspended for particle size testing. Specifically, the powder was directly added to the sample chamber and stirred for 5 minutes, then the sample measurement was taken. The results of this analysis are in Table 10.

TABLE 10

Particle size data for Rapid Maxigesic DPI			
Lot	Ibuprofen		
	D ₁₀ (µm)	D ₅₀ (µm)	D ₉₀ (µm)
10-1	2.15	4.19	8.22
10-2	1.67	3.03	5.48
10-3	1.72	3.45	6.95
10-4	2.55	5.26	10.6
10-5	2.35	4.84	9.97
10-6	2.97	6.89	14.3
10-7	3.15	7.30	14.7

Dissolution Rate

[0064] The dissolution of selected tablet formulations was tested using the USP Apparatus II, with 900 mL of 50 mM pH 5.8 media at 37° C. and a 50 rpm paddle speed. One tablet was added to each media-filled vessel, and samples were assayed by HPLC to obtain the results.

[0065] The film coating is present to aid in swallowing, mask taste, and for pharmaceutical elegance. Film coatings typically take a few minutes to dissolve, creating a small lag time in the dissolution. To facilitate comparison, both film coated and uncoated core tablets were studied. The tablets in these lots were prepared using the DPIs in Table 11. The results of this analysis are depicted in Table 11.

TABLE 11

time	Dissolution of Rapid Maxigesic 325 Ibuprofen dissolution in pH 5.8 buffer									
	LOT 12-A (coated)		LOT 12-B (coated)		LOT 12-C (uncoated)		LOT 12-D (uncoated)		LOT 12-E (uncoated)	
	avg %	% RSD n = 6	avg %	% RSD n = 6	avg %	% RSD n = 6	avg %	% RSD n = 6	avg %	% RSD n = 6
5	54	18.1	28	28.2	47	10.2	50	9.4	63	7.6
10	94	5.4	78	7.8	80	4.6	87	2.2	95	2.3
15	98	2.4	93	2.5	94	0.3	96	0.9	101	0.9
20	100	1.6	96	1.4	95	1	98	1.6	102	0.7
30	100	1.6	97	0.8	96	0.3	97	1.6	102	0.8
45	101	0.8	98	0.5	96	0.6	97	1.9	102	0.6

[0066] In a separate study, the dissolution rate of ibuprofen in two different lots of Rapid Maxigesic 325 tablets (prepared as described above) was compared to that of Maxigesic Tablets 325 (measured in pH 5.8 phosphate buffer, as described above). The results are presented in Table 12. A similar comparison was performed for Rapid Maxigesic 500 and Maxigesic 500 tablets. The results are shown in Table 13.

TABLE 12

Dissolution of Rapid Maxigesic 325 and Maxigesic 325 Tablets in pH 5.8 Phosphate Buffer						
Time (min)	Rapid Maxigesic 325 (Lot X)		Rapid Maxigesic 325 (Lot Y)		Maxigesic 325	
	% Ibuprofen dissolved	% RSD (n = 12)	% Ibuprofen dissolved	% RSD (n = 12)	% Ibuprofen dissolved	% RSD (n = 12)
5	57	13.7	81	8.1	42	12.2
10	91	3.0	95	1.6	59	5.6

TABLE 12-continued

Time (min)	Dissolution of Rapid Maxigesic 325 and Maxigesic 325 Tablets in pH 5.8 Phosphate Buffer					
	Rapid Maxigesic 325 (Lot X)		Rapid Maxigesic 325 (Lot Y)		Maxigesic 325	
	% Ibuprofen dissolved	% RSD (n = 12)	% Ibuprofen dissolved	% RSD (n = 12)	% Ibuprofen dissolved	% RSD (n = 12)
15	95	2.9	98	1.7	69	5.5
20	97	1.3	99	1.3	77	4.3
30	98	1.2	100	1.0	84	4.5
45	99	1.0	100	1.0	89	3.1
F2	34		28		Reference	

TABLE 13

Time (min)	Dissolution of Maxigesic 500 and Rapid Maxigesic 500 Tablets in pH 5.8 Phosphate Buffer			
	Rapid Maxigesic 500 (Lot Z)		Maxigesic 500	
	% Ibuprofen dissolved	% RSD (n = 6)	% Ibuprofen dissolved	% RSD (n = 6)
5	80.8	8.9	42.6	14.5
10	95.6	3.1	65.2	11.3

TABLE 13-continued

Time (min)	Dissolution of Maxigesic 500 and Rapid Maxigesic 500 Tablets in pH 5.8 Phosphate Buffer			
	Rapid Maxigesic 500 (Lot Z)		Maxigesic 500	
	% Ibuprofen dissolved	% RSD (n = 6)	% Ibuprofen dissolved	% RSD (n = 6)
15	98.1	2.2	72.2	6.9
20	99.7	1.5	77.1	3.3
30	100.8	1.3	83.1	2.7
45	100.8	1.5	88.1	2.2
F2	34		Reference	

[0067] From the results describe above, it can be seen that it is possible to prepare solid oral dosages forms with greatly improved dissolution of ibuprofen, despite the ibuprofen having a median particle size (on a volume average basis) that is larger than 1 micron.

Example 4: Pharmacokinetic Analysis of Rapid Maxigesic 325 in Comparison to Maxigesic 325

[0068] A clinical study was undertaken in which the C max for ibuprofen and the C max for acetaminophen was assessed in both the fasting and fed states.

[0069] This data was tested using individual 90% CI, two-tailed, unpaired t-tests. Comparisons were made between Maxigesic Rapid 325 and Maxigesic 325 in either fed or fasting conditions. This allows for a significant difference to be detected at 10% or $P \leq 0.1$. The study protocol already used a 90% CI to calculate the mean differences in the mean log transformed data of acetaminophen and ibuprofen.

[0070] For acetaminophen in the fasting state there was no statistically significant difference in C max between the formulations Table 13.

TABLE 13

Acetaminophen fasting Cmax Maxigesic 325 and Rapid Maxigesic 325	
Difference between means	-459.1 ± 1665
90% confidence interval	-3243 to 2325
Percentage difference between means	-2.99%

[0071] For ibuprofen in the fasting state there was a statistically significant difference in C max between the formulations (Table 14)

TABLE 14

Ibuprofen fasting Cmax Maxigesic 325 and Rapid Maxigesic 325	
Difference between means	-2727 ± 1443
90% confidence interval	-5138 to -315.6
Percentage difference between means	11.93%

[0072] This is an important advantage of the present dosage forms since dosing on an empty stomach is advised in order to achieve fast onset of pain relief.

[0073] For acetaminophen in the fed state there was no statistically significant difference in C max between the formulations (Table 15).

TABLE 15

Acetaminophen fed Cmax Maxigesic 325 and Rapid Maxigesic 325	
Difference between means	-813.1 ± 996
90% confidence interval	-2478 to 851.7
Percentage difference between means	7.82%

[0074] For ibuprofen in the fed state there was no statistically significant difference in C max between the formulations, although a significant increase was again observed (Table 16)

TABLE 16

Ibuprofen fed Cmax Maxigesic 325 and Rapid Maxigesic 325	
Difference between means	2344 ± 1362
90% confidence interval	67.02 to 4622
Percentage difference between means	12.68%

[0075] To summarize, in both studies, fed and fasting, the C max of ibuprofen in Maxigesic Rapid (325) was significantly higher than Maxigesic (325). The 90% confidence intervals did not overlap with 0 which shows a significant difference between the means. There was a difference of >11% between the ibuprofen means, consistent with the Maxigesic Rapid formulation delivering higher C max drug concentrations of ibuprofen.

Example 5: Pharmacokinetic Analysis of Maxigesic 325

[0076] A pharmacokinetic analysis of the Maxigesic 325 formulation described above was conducted under fed and fasting conditions. The results of this study are presented in Tables 17-20.

TABLE 17

Ibuprofen PK Values for Maxigesic Rapid 325 in Fasting condition	
Pharmacokinetic Parameter	(N = 30)
C_{max} (ng/ml)	25579.823 ± 5996.24
$AUC_{0 \rightarrow t}$ (ng · h/ml)	91888.7 ± 24015.73
$AUC_{0 \rightarrow \infty}$ (ng · h/ml)	95621.9 ± 28252.68
t_{max} (h) *	1.25 (0.25-3.00)
K_{el} (1/h)	0.3076 ± 0.06
$t_{1/2el}$ (h)	2.35 ± 0.57
$AUC_{0 \rightarrow t}/AUC_{0 \rightarrow \infty}$ %	96.82 ± 3.22

TABLE 18

Ibuprofen PK Values for Maxigesic Rapid 325 in Fed Condition	
Pharmacokinetic Parameter	(N = 30)
C_{max} (ng/ml)	20834.673 ± 5506.39
$AUC_{0 \rightarrow t}$ (ng · h/ml)	71286.1 ± 16546.13
$AUC_{0 \rightarrow \infty}$ (ng · h/ml)	73449.6 ± 17660.44
t_{max} (h) *	1.25 (0.75-6.00)
K_{el} (1/h)	0.3222 ± 0.05
$t_{1/2el}$ (h)	2.19 ± 0.31
$AUC_{0 \rightarrow t}/AUC_{0 \rightarrow \infty}$ %	97.25 ± 1.42

TABLE 19

Acetaminophen PK Values for Maxigesic Rapid 325 in Fasting Condition	
Pharmacokinetic Parameter	® (N = 29)
C_{max} (ng/ml)	14877.280 ± 5969.64
$AUC_{0 \rightarrow t}$ (ng · h/ml)	44637.6 ± 11784.46
$AUC_{0 \rightarrow \infty}$ (ng · h/ml)	47437.2 ± 13053.06
t_{max} (h) *	0.75 (0.25-2.00)
K_{el} (1/h)	0.2233 ± 0.05
$t_{1/2el}$ (h)	3.25 ± 0.75
$AUC_{0 \rightarrow t}/AUC_{0 \rightarrow \infty}$ %	94.43 ± 2.54

TABLE 20

Acetaminophen PK Values for Maxigesic Rapid 325 in Fed Condition	
Pharmacokinetic Parameter	(N = 30)
C_{max} (ng/ml)	11214.182 ± 4084.46
$AUC_{0 \rightarrow t}$ (ng · h/ml)	39826.6 ± 11211.91
$AUC_{0 \rightarrow \infty}$ (ng · h/ml)	42381.0 ± 12086.45
t_{max} (h) *	1.25 (0.50-6.00)
K_{el} (1/h)	0.2303 ± 0.03
$t_{1/2el}$ (h)	3.05 ± 0.37
$AUC_{0 \rightarrow t} / AUC_{0 \rightarrow \infty}$ %	94.16 ± 2.30

Other Embodiments

[0077] All publications, patents, and patent applications mentioned in this specification are incorporated herein by reference to the same extent as if each independent publi-

cation or patent application was specifically and individually indicated to be incorporated by reference.

[0078] While the disclosure has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the disclosure following, in general, the principles of the disclosure and including such departures from the disclosure that come within known or customary practice within the art to which the disclosure pertains and may be applied to the essential features hereinbefore set forth, and follows in the scope of the claims.

[0079] Other embodiments are within the claims.

What is claimed is:

1. A solid oral dosage form comprising 325 mg of acetaminophen and 97.5 mg of ibuprofen or 500 mg of acetaminophen and 150 mg of ibuprofen, wherein the ibuprofen has a [D50] between 1 and 9 μ m.

* * * * *