



US 20150320021A1

(19) **United States**

(12) **Patent Application Publication**  
**Wang et al.**

(10) **Pub. No.: US 2015/0320021 A1**

(43) **Pub. Date: Nov. 12, 2015**

(54) **HUMANIZED IL-4 AND IL-4RA ANIMALS**

2227/105 (2013.01); *A01K 2267/0393*  
(2013.01)

(71) Applicant: **REGENERON**  
**PHARMACEUTICALS, INC.**,  
Tarrytown, NY (US)

(57) **ABSTRACT**

(72) Inventors: **Li-Hsien Wang**, Somers, NY (US);  
**Yingzi Xue**, Ardsley, NY (US); **Andrew**  
**Murphy**, Croton-on-Hudson, NY (US);  
**Sean Stevens**, San Diego, CA (US)

Non-human animals comprising a human or humanized IL-4 and/or IL-4R $\alpha$  nucleic acid sequence are provided. Non-human animals that comprise a replacement of the endogenous IL-4 gene and/or IL-4R $\alpha$  gene with a human IL-4 gene and/or IL-4R $\alpha$  gene in whole or in part, and methods for making and using the non-human animals, are described. Non-human animals comprising a human or humanized IL-4 gene under control of non-human IL-4 regulatory elements is also provided, including non-human animals that have a replacement of non-human IL-4-encoding sequence with human IL-4-encoding sequence at an endogenous non-human IL-4 locus. Non-human animals comprising a human or humanized IL-4R $\alpha$  gene under control of non-human IL-4R $\alpha$  regulatory elements is also provided, including non-human animals that have a replacement of non-human IL-4R $\alpha$ -encoding sequence with human or humanized IL-4R $\alpha$ -encoding sequence at an endogenous non-human C IL-4R $\alpha$  locus. Non-human animals comprising human or humanized IL-4 gene and/or IL-4R $\alpha$  sequences, wherein the non-human animals are rodents, e.g., mice or rats, are provided.

(21) Appl. No.: **14/706,319**

(22) Filed: **May 7, 2015**

**Related U.S. Application Data**

(60) Provisional application No. 61/989,757, filed on May 7, 2014.

**Publication Classification**

(51) **Int. Cl.**  
**A01K 67/027** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **A01K 67/0278** (2013.01); *A01K 2207/15*  
(2013.01); *A01K 2217/072* (2013.01); *A01K*




Figure 1

### Humanization of IL4 locus




Figure 2A




Figure 2B




Figure 3




Figure 4




Figure 5




Figure 6





Figure 7



Experimental Design




1. IL-4<sup>hchb</sup>/IL-4R<sup>hchb</sup> – PBS; n=4
2. IL-4<sup>hchb</sup>/IL-4R<sup>hchb</sup> – HDM; n=7
3. IL-4<sup>hchb</sup>/IL-4R<sup>hchb</sup> – HDM; REGN129 treatment – 25 mg/kg; n=7
4. IL-4<sup>hchb</sup>/IL-4R<sup>hchb</sup> – HDM; REGN668 treatment – 50 mg/kg; n=7

**Endpoint readouts**

- Collagen deposition in lungs
- serum IgE, HDM-specific IgG1
- BALF cellularity
- lung mucus and infiltrates

**Figure 8**

HDM-14



- A. IL-4<sup>hu/hu</sup>/IL-4R<sup>hu/hu</sup> – Saline+PBS (n=7)
- C. IL-4<sup>hu/hu</sup>/IL-4R<sup>hu/hu</sup> – HDM+REGN668 – dosing at 25mg/kg twice a week, beginning on week 7 (n=7)
- D. IL-4<sup>hu/hu</sup>/IL-4R<sup>hu/hu</sup> – HDM+REGN98 (Isotype Ctrl) – dosing at 25mg/kg twice a week, beginning on week 7 (n=7)

Figure 9




Figure 10A




Figure 10B




Figure 10C

**HUMANIZED IL-4 AND IL-4RA ANIMALS****CROSS REFERENCE TO RELATED APPLICATIONS**

**[0001]** This application claims the benefit of priority to U.S. Provisional Application No. 61/989,757 filed May 7, 2014, the entire contents of which are incorporated herein by reference.

**INCORPORATION BY REFERENCE OF SEQUENCE LISTING**

**[0002]** The Sequence Listing in an ASCII text file, named 31260\_SEQ.txt of 16 KB, created on May 7, 2015, and submitted to the United States Patent and Trademark Office via EFS-Web, is incorporated herein by reference.

**FIELD OF INVENTION**

**[0003]** Non-human animals are disclosed herein which comprise nucleic acid sequences encoding an IL-4 and/or an IL-4R $\alpha$  protein that comprise a human sequence. Transgenic non-human animals are also disclosed herein which comprise an IL-4 and/or an IL-4R $\alpha$  gene that is human in whole or in part. Non-human animals that express human or humanized IL-4 and/or IL-4R $\alpha$  proteins are also disclosed. In addition, methods are disclosed for making and using non-human animals comprising human or humanized IL-4 and/or IL-4R $\alpha$  nucleic acid sequences.

**BACKGROUND**

**[0004]** IL-4 and IL-4R $\alpha$  are therapeutic targets for treatment of a variety of human diseases, disorders and conditions that are associated with abnormal type-2 T helper (Th2) cells. The evaluation of pharmacokinetics (PK) and pharmacodynamics (PD) of therapeutic molecules that specifically target human IL-4 or human IL-4R $\alpha$  proteins are routinely performed in non-human animals, e.g., rodents, e.g., mice or rats. However, the PD of such therapeutic molecules cannot properly be determined in certain non-human animals because these therapeutic molecules do not target the endogenous IL-4 or IL-4R $\alpha$  proteins.

**[0005]** Moreover, the evaluation of therapeutic efficacy of human-specific IL-4 and IL-4R $\alpha$  protein antagonists using various non-human animal models of diseases associated with abnormal Th2 cells is problematic in non-human animals in which such species-specific antagonists do not interact with the endogenous IL-4 or IL-4R $\alpha$  proteins.

**[0006]** Accordingly, there is a need for non-human animals, e.g., rodents, e.g., murine animals, e.g., mice or rats, in which the IL-4 and/or IL-4R $\alpha$  genes of the non-human animal are humanized in whole or in part or replaced (e.g., at the endogenous non-human loci) with human IL-4 and/or IL-4R $\alpha$  genes comprising sequences encoding human or humanized IL-4 and/or IL-4R $\alpha$  proteins, respectively.

**[0007]** There is also a need for non-human animals comprising IL-4 and/or IL-4R $\alpha$  genes (e.g., humanized, or human) in which the IL-4 and/or IL-4R genes are under control of non-human regulatory elements (e.g., endogenous regulatory elements).

**[0008]** There is also a need for humanized non-human animals that express human or humanized IL-4 protein in blood, plasma or serum at a concentration similar to that of IL-4 protein present in blood, plasma or serum of an age-matched non-human animal that expresses functional IL-4 protein, but

does not comprise the human or humanized IL-4 genes, and/or express human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, at a level similar to that of IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, of an age-matched non-human animal that expresses functional IL-4R $\alpha$  protein, but does not comprise the human or humanized IL-4R $\alpha$  gene.

**SUMMARY**

**[0009]** Non-human animals comprising nucleic acid sequences encoding an IL-4 and/or an IL-4R $\alpha$  protein that comprise a human sequence are provided.

**[0010]** Transgenic non-human animals comprising an IL-4 and/or an IL-4R $\alpha$  gene that is human in whole or in part are provided.

**[0011]** Non-human animals that express human or humanized IL-4 and/or IL-4R $\alpha$  proteins are provided.

**[0012]** Non-human animals having a replacement (in whole or in part) of endogenous non-human animal IL-4 and/or IL-4R $\alpha$  genes are provided.

**[0013]** Non-human animals comprising an IL-4 and/or an IL-4R $\alpha$  humanization (in whole or in part) at an endogenous non-human IL-4 and/or IL-4R $\alpha$  loci are provided.

**[0014]** Non-human animals are provided that have a human or humanized IL-4 gene, wherein the non-human animals do not express endogenous IL-4 protein, and wherein the non-human animals express human or humanized IL-4 protein in blood, plasma or serum at a concentration similar to that of IL-4 protein present in blood, plasma or serum of an age-matched non-human animal that expresses functional endogenous IL-4 protein, but does not comprise the human or humanized IL-4 gene.

**[0015]** Non-human animals are provided that have a human or humanized IL-4R $\alpha$  gene, wherein the non-human animals do not express endogenous IL-4R $\alpha$  protein, and express human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, at a level similar to that of IL-4R $\alpha$  protein present on immune cells, e.g., B and T cells, of an age-matched non-human animal that expresses functional endogenous IL-4R $\alpha$  protein, but does not comprise the human or humanized IL-4R $\alpha$  gene.

**[0016]** In one aspect, non-human animals comprising a human or humanized IL-4 and/or IL-4R $\alpha$  nucleic acid sequence are provided.

**[0017]** In one aspect, genetically modified non-human animals are provided that comprise a replacement at an endogenous IL-4 and/or IL-4R $\alpha$  locus of a gene encoding an endogenous IL-4 and/or IL-4R $\alpha$  with a gene encoding a human or humanized IL-4 and/or IL-4R $\alpha$  protein. Rodents, e.g., mice or rats, are provided that comprise a replacement of an endogenous IL-4 gene, at an endogenous rodent IL-4 locus, with a human IL-4 gene, and/or comprise a replacement of an endogenous IL-4R $\alpha$  gene, at an endogenous rodent IL-4R $\alpha$  locus, with a human IL-4R $\alpha$  gene. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0018]** In one aspect, genetically modified rodents, e.g., mice or rats, are provided comprising a humanization of an endogenous rodent IL-4 gene, wherein the humanization comprises a replacement at an endogenous rodent IL-4 locus of a rodent nucleic acid comprising at least one exon of a rodent IL-4 gene with a nucleic acid sequence comprising at least one exon of a human IL-4 gene to form a modified IL-4

gene, wherein expression of the modified IL-4 gene is under control of rodent regulatory elements at the endogenous rodent IL-4 locus.

**[0019]** In one embodiment, the rodent is a mouse or a rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0020]** In one embodiment, the modified IL-4 gene encodes human or humanized IL-4 protein and comprises exon 1 starting from the ATG initiation codon through exon 4 of the human IL-4 gene.

**[0021]** In one embodiment, the rodent is a mouse that is incapable of expressing a mouse IL-4 protein.

**[0022]** In one embodiment, the rodent is a mouse that expresses a mouse IL-4R $\alpha$  protein encoded by an endogenous mouse IL-4R $\alpha$  gene.

**[0023]** In one embodiment, the rodent is mouse that expresses a human or humanized IL-4R $\alpha$  protein.

**[0024]** In one embodiment, the humanized IL-4R $\alpha$  protein comprises the ectodomain of a human IL-4R $\alpha$  protein.

**[0025]** In one embodiment, the humanized IL-4R $\alpha$  protein comprises the transmembrane domain and cytoplasmic domain of a mouse IL-4R $\alpha$  protein.

**[0026]** In one embodiment, the rodent is a mouse that comprises a replacement at an endogenous mouse IL-4R $\alpha$  locus of a mouse nucleic acid comprising at least one exon of a mouse IL-4R $\alpha$  gene with a nucleic acid sequence comprising at least one exon of a human IL-4R $\alpha$  gene to form a modified IL-4R $\alpha$  gene, wherein expression of the modified IL-4R $\alpha$  gene is under control of mouse regulatory elements at the endogenous mouse IL-4R $\alpha$  locus.

**[0027]** In one embodiment, the rodent is a mouse, wherein a contiguous genomic fragment of mouse IL-4 sequence comprising exon 1 starting from the ATG initiation codon through exon 4 of mouse IL-4 is replaced with a contiguous genomic fragment of human IL-4 sequence comprising exon 1 starting from the ATG initiation codon through exon 4 of human IL-4.

**[0028]** In one embodiment, expression of the modified IL-4R $\alpha$  gene encoding the human or humanized IL-4R $\alpha$  protein is under control of mouse regulatory elements at the endogenous mouse IL-4R $\alpha$  locus.

**[0029]** In one aspect, genetically modified rodents, e.g., mice or rats, are provided comprising a humanization of an endogenous rodent IL-4R $\alpha$  gene, wherein the humanization comprises a replacement at an endogenous rodent IL-4R $\alpha$  locus of a rodent nucleic acid comprising an exon of a rodent IL-4R $\alpha$  gene with a nucleic acid sequence encoding at least one exon of a human IL-4R $\alpha$  gene to form a modified (i.e., humanized) IL-4R $\alpha$  gene, wherein expression of the modified, humanized IL-4R $\alpha$  gene is under control of rodent regulatory elements at the endogenous rodent IL-4R $\alpha$  locus.

**[0030]** In one embodiment, the rodent is a mouse or a rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0031]** In one embodiment, the modified IL-4R $\alpha$  gene encodes human or humanized IL-4R $\alpha$  protein and comprises exon 1 starting from the ATG initiation codon through exon 5 of the human IL-4R $\alpha$  gene.

**[0032]** In one embodiment, the rodent is a mouse that is incapable of expressing a mouse IL-4R $\alpha$  protein.

**[0033]** In one embodiment, the rodent is a mouse that expresses a mouse IL-4 protein encoded by an endogenous mouse IL-4 gene.

**[0034]** In one embodiment, the rodent is mouse that expresses a human or humanized IL-4 protein.

**[0035]** In one embodiment, the rodent is a mouse that comprises a replacement at an endogenous mouse IL-4 locus of a mouse nucleic acid comprising an exon of a mouse IL-4 gene with a nucleic acid sequence encoding at least one exon of a human IL-4 gene to form a modified IL-4 gene, wherein expression of the modified IL-4 gene is under control of mouse regulatory elements at the endogenous mouse IL-4 locus.

**[0036]** In one embodiment, the rodent is a mouse, and wherein a contiguous genomic fragment of mouse IL-4R $\alpha$  sequence comprising exon 1 starting from the ATG initiation codon through exon 5 of IL-4R $\alpha$  is replaced with a contiguous genomic fragment of human IL-4R $\alpha$  sequence comprising exon 1 starting from the ATG initiation codon through exon 5 of human IL-4R $\alpha$ .

**[0037]** In one aspect, genetically modified rodents, e.g., a mouse or rat, are provided that express a human or humanized IL-4 protein, wherein the rodent that expresses a human or humanized IL-4 protein comprises a normal immune system, i.e., the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of the rodent expressing human or humanized IL-4 protein are similar to the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of a rodent that expresses functional endogenous IL-4 protein. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0038]** In one aspect, genetically modified rodents, e.g., a mouse or rat, are provided that express IL-4 protein from a human or humanized IL-4 gene, wherein the rodent expresses human or humanized IL-4 protein in its serum. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0039]** In one embodiment, the serum of the rodent that expresses a human or humanized IL-4 protein has approximately the same level of IL-4 protein as a rodent that expresses a functional, endogenous IL-4 protein, e.g., a wild-type mouse or rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0040]** In one embodiment, the mouse expresses human or humanized IL-4 protein in serum at a concentration of at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190% or 200% of the level of IL-4 protein present in the serum of an age-matched mouse that expresses functional endogenous IL-4 protein, but does not comprise a replacement of an endogenous IL-4 gene, at an endogenous mouse IL-4 locus, with a human IL-4 gene.

**[0041]** In one embodiment, the mouse expresses human or humanized IL-4 protein in serum at a concentration of between about 10% and about 200%, between about 20% and about 150%, or between about 30% and about 100% of the level of mouse IL-4 protein present in the serum of an age-matched mouse that expresses functional endogenous IL-4 protein, but does not comprise a replacement of an endogenous IL-4 gene, at an endogenous mouse IL-4 locus, with a human IL-4 gene.

**[0042]** In one aspect, genetically modified rodents, e.g., a mouse or rat, are provided that express a human or humanized IL-4R $\alpha$  protein, wherein the rodent expresses a human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, at a level similar to that of IL-4R $\alpha$  protein present on immune cells, e.g., B and T cells, of an age-matched rodent

that expresses functional endogenous IL-4R $\alpha$  protein. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0043]** In one aspect, genetically modified rodents, e.g., a mouse or rat, are provided that express IL-4R $\alpha$  protein from a human IL-4R $\alpha$  gene, wherein the rodent expresses human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0044]** In one embodiment, the immune cells, e.g., B and T cells, of the rodent that expresses a human or humanized IL-4R $\alpha$  protein have approximately the same level of IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, of a rodent that expresses a functional, endogenous IL-4R $\alpha$  protein, e.g., a wild-type mouse or rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0045]** In one embodiment, the mouse expresses human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, at an amount of at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190% or 200% of the amount of IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, of an age-matched mouse that expresses functional endogenous IL-4R $\alpha$  protein, but does not comprise a replacement of an endogenous IL-4R $\alpha$  gene, at an endogenous mouse IL-4R $\alpha$  locus, with a human IL-4R $\alpha$  gene.

**[0046]** In one embodiment, the mouse expresses human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells, at an amount of between about 10% and about 200%, between about 20% and about 150%, or between about 30% and about 100% of the amount of mouse IL-4R $\alpha$  protein present on immune cells, e.g., B and T cells, of an age-matched mouse that expresses functional endogenous IL-4R $\alpha$  protein, but does not comprise a replacement of an endogenous IL-4R $\alpha$  gene, at an endogenous mouse IL-4R $\alpha$  locus, with a human IL-4R $\alpha$  gene.

**[0047]** In one aspect, a genetically modified rodent is provided, comprising a humanized IL-4R $\alpha$  gene comprising a replacement of rodent IL-4R $\alpha$  ectodomain-encoding sequence with human IL-4R $\alpha$  ectodomain-coding sequence, wherein the humanized IL-4R $\alpha$  gene comprises a rodent IL-4R $\alpha$  transmembrane sequence and a rodent IL-4R $\alpha$  cytoplasmic sequence, wherein the humanized IL-4R $\alpha$  gene is under control of endogenous rodent IL-4R $\alpha$  regulatory elements at the endogenous IL-4R $\alpha$  locus, and wherein the rodent further comprises a humanized IL-4 gene encoding a human or humanized IL-4 protein, wherein the humanized IL-4 gene is under control of endogenous rodent IL-4 regulatory elements at the endogenous IL-4 locus.

**[0048]** In one embodiment, the rodent is a mouse or a rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0049]** In one embodiment, the mouse is incapable of expressing a mouse IL-4 protein and incapable of expressing a mouse IL-4R $\alpha$  protein.

**[0050]** In one embodiment, the rodent regulatory elements or sequences at the endogenous rodent IL-4 locus and/or rodent IL-4R $\alpha$  locus are from a mouse or a rat.

**[0051]** In one embodiment, the rodent regulatory elements or sequences are endogenous rodent regulatory elements or sequences at the rodent IL-4 locus and/or rodent IL-4R $\alpha$  locus.

**[0052]** In one aspect, a non-human animal, e.g., a rodent, e.g., a mouse or rat, is provided that expresses human or

humanized IL-4 and/or IL-4R $\alpha$  proteins, wherein the non-human animal expresses human or humanized IL-4 and/or IL-4R $\alpha$  proteins from an endogenous non-human IL-4R $\alpha$  locus and/or an endogenous non-human IL-4R $\alpha$  locus. In an embodiment, the non-human animal is a rodent. In an embodiment, the rodent is a mouse. In an embodiment, the rodent is a rat.

**[0053]** In one aspect, a genetically modified mouse is provided that expresses human or humanized IL-4 protein from an endogenous mouse IL-4 locus, wherein the endogenous mouse IL-4 gene has been replaced, in whole or in part, with a human IL-4 gene.

**[0054]** In one embodiment, a contiguous mouse genomic nucleic acid of about 6.3 kb at an endogenous mouse IL-4 locus, including exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4, is deleted and replaced with about 8.8 kb of a human IL-4 nucleic acid sequence comprising exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 of the human IL-4 gene. In a specific embodiment, the human IL-4 nucleic acid sequence replacing the mouse genomic nucleic acid comprises exon 1 starting from the ATG initiation codon through exon 4 and a portion of the 3' region downstream of exon 4 of the human IL-4 gene of human BAC RP11-17K19. In a specific embodiment, the modified IL-4 gene comprises mouse IL-4 5' regulatory elements and human IL-4 exon 1 starting from the ATG initiation codon through exon 4, i.e., the IL-4 protein coding sequences.

**[0055]** In one aspect, a genetically modified mouse is provided that comprises a nucleotide sequence encoding a human or humanized IL-4 protein, wherein the nucleotide sequence encoding the human or humanized IL-4 protein replaces, in whole or in part, an endogenous nucleotide sequence encoding an endogenous mouse IL-4 protein.

**[0056]** In one aspect, a genetically modified mouse is provided that expresses human or humanized IL-4R $\alpha$  protein from an endogenous mouse IL-4R $\alpha$  locus, wherein the endogenous mouse IL-4R $\alpha$  gene has been replaced, in whole or in part, with a human IL-4R $\alpha$  gene.

**[0057]** In one embodiment, a contiguous mouse genomic nucleic acid of about 7.1 kb at the endogenous mouse IL-4R $\alpha$  locus, including exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5, is deleted and replaced with about 15.6 kb of human IL-4R $\alpha$  nucleic acid sequence comprising exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 of the human IL-4R $\alpha$  gene. In a specific embodiment, the human IL-4 $\alpha$  nucleic acid replacing the mouse genomic nucleic acid comprises exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 of the human IL-4 $\alpha$  gene of human BAC RP11-16E24. In a specific embodiment, the human IL-4R $\alpha$  nucleic acid replacing the mouse genomic nucleic acid comprises the entire human IL-4R $\alpha$  ectodomain coding sequence.

**[0058]** In one aspect, a method is provided for making a humanized IL-4 rodent, comprising replacing a rodent IL-4 gene sequence encoding rodent IL-4 protein with a human IL-4 nucleic acid sequence comprising one or more exons of the human IL-4 gene sequence to form a modified, humanized IL-4 gene encoding human or humanized IL-4 protein, wherein the replacement is at an endogenous rodent IL-4 locus and the humanized IL-4 gene sequence comprising one



or more exons of the human IL-4 gene sequence and encoding human or humanized IL-4 protein is operably linked to rodent regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous rodent IL-4 locus.

**[0059]** In one embodiment, the rodent is a mouse or a rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0060]** In one embodiment, the rodent regulatory elements or sequences are derived from a mouse. In one embodiment, the rodent regulatory elements or sequences are derived from a rat.

**[0061]** In one embodiment, the rodent regulatory elements or sequences are endogenous rodent regulatory elements or sequences at the rodent IL-4 locus. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0062]** In one embodiment, the human IL-4 nucleic acid sequence replacing the rodent IL-4 gene sequence comprises at least one exon of the human IL-4 gene sequence. In other embodiments, the human IL-4 nucleic acid sequence replacing the rodent IL-4 gene sequence comprises at least 2 or at least 3 exons of the human IL-4 gene sequence. In one embodiment, the human IL-4 nucleic acid sequence replacing the rodent IL-4 gene sequence comprises all 4 exons of the human IL-4 gene sequence. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0063]** In one embodiment, the human IL-4 nucleic acid sequence replacing the rodent IL-4 gene sequence encodes a protein that is about 85%, 90%, 95%, 96%, 97%, 98%, or about 99% identical to a human IL-4 (e.g., the human IL-4 protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000589.3 s).

**[0064]** In one embodiment, the replacement is at an endogenous rodent IL-4 locus and the humanized IL-4 gene sequence comprising one or more exons of the human IL-4 gene sequence and encoding human or humanized IL-4 protein is operably linked to endogenous rodent regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous rodent IL-4 locus.

**[0065]** In one aspect, a method is provided for making a humanized IL-4 mouse, comprising replacing a mouse IL-4 gene sequence encoding mouse IL-4 protein with a human IL-4 gene sequence to form a modified, humanized IL-4 gene encoding human or humanized IL-4 protein.

**[0066]** In one embodiment, the replacement is at an endogenous mouse IL-4 locus, and the resulting humanized IL-4 gene encoding human or humanized IL-4 protein is operably linked to mouse regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous mouse IL-4 locus.

**[0067]** In one embodiment, the replacement is at an endogenous mouse IL-4 locus, and the humanized IL-4 gene encoding human or humanized IL-4 protein is operably linked to endogenous mouse regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous mouse IL-4 locus.

**[0068]** In one aspect, a method is provided for making a humanized IL-4R $\alpha$  rodent, comprising replacing a rodent IL-4R $\alpha$  gene sequence encoding rodent IL-4R $\alpha$  protein with a human IL-4R $\alpha$  nucleic acid sequence comprising one or more exons of the human IL-4R $\alpha$  gene sequence to form a modified, humanized IL-4R $\alpha$  gene encoding human or humanized IL-4R $\alpha$  protein, wherein the replacement is at an endogenous rodent IL-4R $\alpha$  locus and the humanized IL-4R $\alpha$  gene sequence comprising one or more exons of the human

IL-4R $\alpha$  gene sequence and encoding human or humanized IL-4R $\alpha$  protein is operably linked to rodent regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous rodent IL-4R $\alpha$  locus.

**[0069]** In one embodiment, the rodent is a mouse or a rat. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0070]** In one embodiment, the rodent regulatory elements or sequences are derived from a mouse. In one embodiment, the rodent regulatory elements or sequences are derived from a rat.

**[0071]** In one embodiment, the rodent regulatory elements or sequences are endogenous rodent regulatory elements or sequences at the rodent IL-4R $\alpha$  locus. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0072]** In one embodiment, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises at least one exon of the human IL-4R $\alpha$  gene sequence. In other embodiments, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises at least 2, 3, 4, 5, 6, 7, or 8 exons of the human IL-4R $\alpha$  gene sequence. In one embodiment, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises all 9 exons of the human IL-4R $\alpha$  gene sequence. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0073]** In one embodiment, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence encodes a protein that is about 85%, 90%, 95%, 96%, 97%, 98%, or about 99% identical to a human IL-4R $\alpha$  (e.g., the human IL-4R $\alpha$  protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000418.3).

**[0074]** In one embodiment, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises at least one exon of the human IL-4R $\alpha$  gene sequence encoding the ectodomain of the human IL-4R $\alpha$  protein. In other embodiments, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises at least 2, 3, or 4 exons of the human IL-4R $\alpha$  gene sequence encoding the ectodomain of the human IL-4R $\alpha$  protein. In one embodiment, the human IL-4R $\alpha$  nucleic acid sequence replacing the rodent IL-4R $\alpha$  gene sequence comprises all 5 exons of the human IL-4R $\alpha$  gene sequence encoding the ectodomain of the human IL-4R $\alpha$  protein. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0075]** In one embodiment, the human or humanized IL-4R $\alpha$  gene sequence replacing the rodent IL-4R $\alpha$  gene sequence encodes an ectodomain of the IL-4R $\alpha$  protein that is about 85%, 90%, 95%, 96%, 97%, 98%, or about 99% identical to the ectodomain of a human IL-4R $\alpha$  protein (e.g., the human IL-4R $\alpha$  protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000418.3).

**[0076]** In one embodiment, the replacement is at an endogenous rodent IL-4R $\alpha$  locus and the humanized IL-4R $\alpha$  gene sequence comprising one or more exons of the human IL-4R $\alpha$  gene sequence and encoding human or humanized IL-4R $\alpha$  protein is operably linked endogenous rodent regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous rodent IL-4R $\alpha$  locus.

**[0077]** In one aspect, a method is provided for making a humanized IL-4R $\alpha$  mouse, comprising replacing a mouse IL-4R $\alpha$  gene sequence encoding mouse IL-4R $\alpha$  protein with a human IL-4R $\alpha$  nucleic acid sequence to form a humanized IL-4R $\alpha$  gene encoding human or humanized IL-4R $\alpha$  protein.

**[0078]** In one embodiment, the replacement is at an endogenous mouse IL-4R $\alpha$  locus, and the humanized IL-4R $\alpha$  gene encoding human or humanized IL-4R $\alpha$  protein is operably linked to mouse regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous mouse IL-4R $\alpha$  locus.

**[0079]** In one embodiment, the replacement is at an endogenous mouse IL-4R $\alpha$  locus, and the humanized IL-4R $\alpha$  gene encoding human or humanized IL-4R $\alpha$  protein is operably linked to endogenous mouse regulatory elements or sequences (e.g., 5' and/or 3' regulatory elements) at the endogenous mouse IL-4R $\alpha$  locus.

**[0080]** In various aspects, the genetically modified non-human animals, e.g., rodents, e.g., mice or rats, described herein comprise the genetic modifications in their germline.

**[0081]** In one aspect, a non-human animal, e.g., rodent, e.g., a mouse or rat, embryo comprising a genetic modification as described herein is provided.

**[0082]** In one aspect, a non-human animal, e.g., rodent, e.g., a mouse or rat, host embryo is provided that comprises a donor cell that comprises a genetic modification as described herein.

**[0083]** In one aspect, a pluripotent or totipotent non-human animal, e.g., rodent, e.g., mouse or rat, cell comprising a genetic modification as described herein is provided. In one embodiment, the cell is a rodent cell. In one embodiment, the cell is a mouse cell. In one embodiment, the cell is a rodent ES cell. In one embodiment, the cell is a mouse ES cell.

**[0084]** In one aspect, a non-human animal, e.g., rodent, e.g., mouse or rat, egg is provided, wherein the non-human animal egg comprises an ectopic non-human animal chromosome, wherein the ectopic non-human animal chromosome comprises a genetic modification as described herein. In one embodiment, the non-human animal is a rodent. In one embodiment, the rodent is a mouse. In one embodiment, the rodent is a rat.

**[0085]** In one aspect, the mouse embryo, egg, or cell that is genetically modified to comprise a human IL-4 gene or human IL-4R $\alpha$  gene is of a mouse that is of a C57BL strain selected from C57BL/A, C57BL/An, C57BL/GrFa, C57BL/KaLwN, C57BL/6, C57BL/6J, C57BL/6ByJ, C57BL/6NJ, C57BL/10, C57BL/10ScSn, C57BL/10Cr, and C57BL/Ola. In another embodiment, the mouse is a 129 strain selected from the group consisting of a strain that is 129P1, 129P2, 129P3, 129X1, 129S1 (e.g., 129S1/SV, 129S1/SvIm), 129S2, 129S4, 129S5, 129S9/SvEvH, 129S6 (129/SvEvTac), 129S7, 129S8, 129T1, 129T2 (see, e.g., Festing et al. (1999) Revised nomenclature for strain 129 mice, Mammalian Genome 10:836, see also, Auerbach et al (2000) Establishment and Chimera Analysis of 129/SvEv- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines). In a specific embodiment, the genetically modified mouse is a mix of an aforementioned 129 strain and an aforementioned C57BL/6 strain. In another specific embodiment, the mouse is a mix of aforementioned 129 strains, or a mix of aforementioned BL/6 strains. In a specific embodiment, the 129 strain of the mix is a 129S6 (129/SvEvTac) strain. In another embodiment, the mouse is a BALB strain, e.g., BALB/c strain. In yet another embodiment, the mouse is a mix of a BALB strain and another aforementioned strain. In one embodiment, the mouse is Swiss or Swiss Webster mouse.

**[0086]** In various aspects, the non-human animals comprising a human or humanized IL-4R and/or IL-4 nucleic acid sequence are selected from mammals and birds. In one

embodiment, the non-human animals are mammals. In one embodiment, the mammals are murine.

**[0087]** In one aspect, a method of screening for a human-specific IL-4 or IL-4R $\alpha$  antagonist is provided. The method is useful for identifying therapeutic candidates and evaluating therapeutic efficacy. The method comprises administering an agent to a genetically modified rodent that is doubly humanized for IL-4 and IL-4R $\alpha$  as described herein, determining an effect of the agent on a biological function mediated by the IL-4/IL-4R $\alpha$  signaling pathway, and identifying the agent as a human-specific IL-4 or IL-4R $\alpha$  antagonist if it antagonizes the function mediated by the IL-4/IL-4R $\alpha$  signaling pathway in the genetically modified rodent.

**[0088]** In one embodiment, the agent comprises an immunoglobulin variable domain that binds IL-4 or IL-4R $\alpha$ . In one embodiment, the agent specifically binds human IL-4 or IL-4R $\alpha$ , but not rodent IL-4 or IL-4R $\alpha$ . In one embodiment, the agent is an antibody. In a specific embodiment, the agent is an antibody that specifically binds human IL-4R $\alpha$ , but not rodent IL-4R $\alpha$ .

**[0089]** In one embodiment, the screening method utilizes a doubly humanized mouse that expresses a human IL-4 protein, and a humanized IL-4R $\alpha$  protein, wherein the humanized IL-4R $\alpha$  protein includes the ectodomain of a human IL-4R $\alpha$  protein, linked to the transmembrane and cytoplasmic domains of the endogenous mouse IL-4R $\alpha$  protein, and wherein the mouse does not express murine IL-4 or murine IL-4R $\alpha$ .

**[0090]** In some embodiments, the method of screening includes the steps of inducing in a doubly humanized rodent as described herein a disease associated with IL-4/IL-4R $\alpha$  signaling, administering an agent to the rodent, determining whether the agent ameliorates the disease, and identifying the agent as a human-specific IL-4 or IL-4R $\alpha$  antagonist suitable for treating the disease if the agent ameliorates the disease.

**[0091]** In some embodiments, the disease associated with IL-4/IL-4R $\alpha$  signaling is airway inflammation, which can be induced in a rodent by intranasal administration of an allergen (e.g., house dust mite extract) in one or more doses for a period of time. The effect of an agent can be determined by measuring whether the extent of airway inflammation (reflected by e.g., mucus accumulation, infiltrating cells in bronchoalveolar lavage fluid, and/or levels of total circulating IgE), is reduced as a result of the administration of the agent.

**[0092]** In some embodiments, the disease associated with IL-4/IL-4R $\alpha$  signaling is skin inflammation or atopic dermatitis, which can be induced in a rodent by creating skin injury and exposing the injured skin to an allergen (e.g., bacterial toxin or house dust mite extract) in one or more doses for a period of time. The effect of an agent can be determined by measuring whether skin inflammation is reduced as a result of the administration of the agent.

**[0093]** In a further aspect, a triply humanized non-human animal whose IL-4, IL-4R $\alpha$ , and IL-33 genes have been humanized as described herein is used to evaluate the pharmacodynamics (PD) and therapeutic efficacy of a compound or a combination of compounds.

**[0094]** Each of the aspects and embodiments described herein are capable of being used together, unless excluded either explicitly or clearly from the context of the embodiment or aspect.

## BRIEF DESCRIPTION OF THE DRAWINGS

**[0095]** FIG. 1 provides an illustration, not to scale, of the receptors for IL-4 and IL-13 signal transduction and the mechanism of action of dupilumab, a neutralizing fully human monoclonal antibody that binds specifically to the human IL-4 receptor  $\alpha$  chain (IL-4R $\alpha$ ).

**[0096]** FIGS. 2A-2B provide an illustration, not to scale, of the strategies for humanization of the IL-4 (I14) and IL-4R $\alpha$  (I14r $\alpha$ ) loci. (2A) The mouse IL-4 gene (top) spanning the coding region from exon 1 starting at the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 are deleted and replaced by the coding region from exon 1 starting at the ATG codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 of the human IL-4 gene (bottom) along with a floxed hygro selection cassette loxP, as indicated. (2B) The mouse IL-4R $\alpha$  gene (top) spanning the coding region from exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 are deleted and replaced by the coding region from exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 of the human IL-4R $\alpha$  gene (bottom) and a floxed neo selection cassette, as indicated.

**[0097]** FIG. 3 shows the expression of humanized IL-4R $\alpha$  protein on B and T cells from doubly humanized IL-4/IL-4R $\alpha$  (I14<sup>hu/hu</sup>/I14r<sup>hu/hu</sup>) mice.

**[0098]** FIG. 4 shows the IL-4 and IL-13 ligand specificities and receptor functionalities using primary cells derived from humanized IL-4R $\alpha$  (I14r<sup>hu/hu</sup>) mice.

**[0099]** FIG. 5 shows IL-4 dependent IgE production in vivo in wild-type, but not humanized IL-4R $\alpha$  (I14r<sup>hu/hu</sup>) mice.

**[0100]** FIG. 6 shows that dose-dependent IL-4 induction of mIgE production ex vivo in mouse B cells (left panel) is blocked by dupilumab, in a dose-dependent manner (right panel).

**[0101]** FIG. 7 shows that dupilumab, in a dose-dependent manner, prevents IL-25 induced lung pathologies in vivo in humanized IL-4R $\alpha$  (I14r<sup>hu/hu</sup>) mice.

**[0102]** FIG. 8 shows the experimental design for assessing the therapeutic efficacy of dupilumab in a house dust mite extract (HDM) induced pulmonary inflammation model using doubly humanized IL-4 and IL-4R $\alpha$  (IL-4<sup>hu/hu</sup>/IL-4R<sup>hu/hu</sup>) mice. "REGN668" refers to a human monoclonal antibody directed to human IL-4R $\alpha$ , also known as dupilumab. "REGN129" refers to a mouse sol IL-13R $\alpha$ 2-Fc, which is a fusion protein between the ectodomain of mouse IL-13R $\alpha$ 2 and Fc.

**[0103]** FIG. 9 shows the experimental design for assessing the therapeutic efficacy of dupilumab in an HDM induced pulmonary inflammation model using doubly humanized IL-4 and IL-4R $\alpha$  (IL-4<sup>hu/hu</sup>/IL-4R<sup>hu/hu</sup>) mice and an isotype control antibody.

**[0104]** FIGS. 10A-10C illustrate the strategies for humanization of the mouse IL-33 locus. FIG. 10A illustrates that the mouse IL-33 gene (top) spanning the coding region from exon 2 starting at the ATG initiation codon through the stop codon in exon 8 is deleted and replaced by the coding region from exon 2 starting at the ATG codon through exon 8 (including the 3' untranslated region) of the human IL-33 gene (bottom). FIG. 10B shows the humanized IL-33 allele in mouse ES cell clone MAID 7060, which contains a loxP neomycin selection cassette. FIG. 10C shows the humanized IL-33 allele in mouse ES cell clone MAID 7061, in which the

neomycin selection cassette has been deleted, with loxP and cloning sites (77 bp) remaining downstream of the human IL-33 sequence, and mouse 3' UTR retained downstream of the loxP site.

## DETAILED DESCRIPTION

IL-4 and IL-4R $\alpha$  as Therapeutic Targets

**[0105]** Allergic disorders are a spectrum of diseases that are occurring at an increasing rate, especially in developed countries. Atopic dermatitis, asthma, and allergic rhinitis are the most common inflammatory conditions among patients with allergies; these patients often suffer the onset of multiple clinical symptoms. The pathogenesis of allergy is linked to abnormal immune responses against exogenous antigens (see Mueller et al. (2002) Structure, binding, and antagonists in the IL-4/IL-13 receptor system, *Biochim Biophys Acta* 1592: 237-250).

**[0106]** Over-production of antigen-specific IgE is an essential component to trigger allergic inflammation. Abnormal type-2 T helper cell (Th2) polarization contributes to the increased IgE responses.

**[0107]** Interleukin-4 (IL-4) and interleukin-13 (IL-13), originally identified from activated T cells, are major Th2 cytokines that play central roles in initiating and sustaining the immune and inflammatory reactions in allergies.

**[0108]** IL-4 and IL-13 signaling are mediated by two distinct receptor complexes with a shared subunit, IL-4 receptor alpha (IL-4R $\alpha$ ), which may contribute to the overlapping biological responses between these two cytokines. See FIG. 1.

**[0109]** Receptors for Interleukin-4/13 Signal Transduction and the Mechanism of Action of Dupilumab.

**[0110]** IL-4R $\alpha$  forms two distinct heterodimeric receptor complexes to mediate the biological functions of IL-4 and IL-13 in a tissue- and response-specific manner. The type I receptor comprised of IL-4R $\alpha$  and common cytokine receptor gamma chain ( $\gamma$ C) is unique for IL-4. Type II receptor formed between IL-4R $\alpha$  and IL-13R $\alpha$ 1 is the primary receptor for IL-13, but is also functional for IL-4. In addition, IL-13 will bind to a second high affinity receptor, IL-13R $\alpha$ 2, which is generally recognized as a decoy receptor or with a possible, pro-fibrotic effect in the full-length form.

**[0111]** Dupilumab is an antagonistic monoclonal antibody against human IL-4R $\alpha$  that inhibits induced biological activities from IL-4 and IL-13. Dupilumab blocks IL-4 signal transduction by preventing its binding to receptor subunits, whereas the inhibitory effect on IL-13 signaling is likely mediated through interfering with the dimeric receptor interaction.

**[0112]** Dupilumab, a fully-human monoclonal antibody directed against the shared IL-4R $\alpha$  subunit, was developed at Regeneron Pharmaceuticals, Inc. using VelocImmune® mice. Dupilumab is undergoing clinical trials for the treatment of moderate-to-severe asthma and for the treatment of moderate-to-severe atopic dermatitis.

**[0113]** Evaluating the potency of dupilumab in murine models presents multiple challenges: (a) dupilumab does not recognize the cognate mouse IL-4 receptor; and (b) there is a lack of functional interaction between mouse IL-4 protein and human IL-4 receptor.

#### IL-4 Gene and Protein

**[0114]** The IL-4 gene encodes a secreted IL-4 protein, which plays an important role in the activation of B cells, as well as other cell types (see FIG. 1).

**[0115]** Human IL-4.

**[0116]** NCBI Gene ID: 3565; Primary source: HGNC: 6014; RefSeq transcript: NM\_000589.3; UniProt ID: P05112; Genomic assembly: GRCh37; Location: chr5:132,009,743-132,018,576+ strand.

**[0117]** The human IL-4 gene is located on chromosome 5, at 5q31.1. The human IL-4 gene has 4 exons and encodes a precursor polypeptide of 153 amino acids in length, including a 24 amino acid signal peptide, and a 129 amino acid mature IL-4 protein.

**[0118]** Mouse IL-4.

**[0119]** NCBI Gene ID: 16189; Primary source: MGI: 96556; RefSeq transcript: NM\_021283.2; UniProt ID: P07750; Genomic assembly: GRCm38; Location: chr11:53,612,350-53,618,606- strand.

**[0120]** The mouse IL-4 gene is located on chromosome 11, at 11 31.97 cM. The mouse IL-4 gene has 4 exons and encodes a precursor polypeptide of 140 amino acids in length, including a 20 amino acid signal peptide, and a 120 amino acid mature IL-4 protein.

#### IL-4R $\alpha$ Gene and Protein

**[0121]** The IL-4R $\alpha$  gene encodes the transmembrane receptor IL-4R $\alpha$  protein, which is expressed primarily on B and T cells, is a receptor for the IL-4 and IL-13 proteins (see FIG. 1).

**[0122]** Human IL-4R $\alpha$ .

**[0123]** NCBI Gene ID: 3566; Primary source: MGI:6015; RefSeq transcript: NM\_000418.3; UniProt ID: P24394; Genomic assembly: GRCh37; Location: chr16:27,351,525-27,367,111+ strand.

**[0124]** The human IL-4R $\alpha$  gene is located on chromosome 16 at 16p12.1-p11.2. The human IL-4R $\alpha$  gene has 9 coding exons and encodes a precursor polypeptide of 825 amino acids, including a 25 amino acid signal peptide, and an 800 amino acid mature IL-4R $\alpha$  protein, with the first 207 amino acid residues of the mature protein constituting the extracellular domain. The extracellular domain (i.e., ectodomain) of the human IL-4R $\alpha$  protein is encoded by coding exons 1 through 5 of the human IL-4R $\alpha$  gene.

**[0125]** Mouse IL-4R $\alpha$ .

**[0126]** NCBI Gene ID: 16190; Primary source: MGI: 105367; RefSeq transcript: NM\_001008700.3; UniProt ID: P16382; Genomic assembly: GRCm38; Location: chr11:125,565,655-125,572,745+ strand.

**[0127]** The mouse IL-4R $\alpha$  gene is located on chromosome 7 at 7 68.94 cM. The mouse IL-4R $\alpha$  gene has 9 coding exons and encodes a precursor polypeptide of 810 amino acids, including a 25 amino acid signal peptide, and a 785 amino acid mature IL-4R $\alpha$  protein, with the first 208 amino acid residues of the mature protein constituting the extracellular domain. The extracellular domain (i.e., ectodomain) of the mouse IL-4R $\alpha$  protein is encoded by coding exons 1 through 5 of the mouse IL-4R $\alpha$  gene.

#### Species Specificity of IL-4 and IL-4R $\alpha$ Proteins

**[0128]** As shown herein, mouse, but not human, IL-4 is functional in wild-type mice, and, conversely, human, but not mouse, IL-4 is functional in humanized IL-4R $\alpha$  (IL4ra<sup>hu/hu</sup>)

mice. (See also, e.g., Andrews et al. (2001) Reconstitution of a functional human type II IL-4/IL-13 receptor in mouse B cells: demonstration of species specificity, J Immunol. 166: 1716-1722).

#### Species Specificity of Human IL-4 and IL-4R $\alpha$ Inhibitors

**[0129]** Candidate therapeutic molecules that target the IL-4 or IL-4R $\alpha$  proteins are typically evaluated for pharmacokinetics (PK) and pharmacodynamics (PD) in non-human animals, e.g., rodents, e.g., mice or rats. Such therapeutic molecules are also tested for in vivo therapeutic efficacy in non-human animal, e.g., rodent, e.g., mouse or rat, models of human diseases, disorders and conditions associated with abnormal Th2 cells.

**[0130]** However, therapeutic molecules that are specific for the human IL-4 or IL-4R $\alpha$  proteins, e.g., human-specific IL-4 or IL-4R $\alpha$  inhibitors, cannot be adequately evaluated for PD or in vivo therapeutic efficacy in rodents, in particular mice, because the targets of these therapeutic molecules are missing. This problem is not overcome using transgenic non-human animals, e.g., rodents, e.g., mice or rats, expressing human IL-4 or IL-4R $\alpha$  proteins because of the above-mentioned species specificity of IL-4 protein.

**[0131]** Accordingly, in various embodiments, to assess the PD and in vivo therapeutic efficacy of a human-specific IL-4 or IL-4R $\alpha$  protein antagonist or inhibitor in non-human animals, e.g., rodents, e.g., mice or rats, it is desirable to replace the endogenous IL-4 and/or IL-4R $\alpha$  proteins with human IL-4 and/or IL-4R $\alpha$  proteins.

**[0132]** Further, in various embodiments, in order to avoid potential problems of over- or under-expression of the human IL-4 and/or IL-4R $\alpha$  proteins, it is desirable to insert the human IL-4 and/or IL-4R $\alpha$  genes into the genome of the non-human animals, e.g., rodents, e.g., mice or rats, at the endogenous IL-4 and/or IL-4R $\alpha$  gene loci, and to express the human IL-4 and/or IL-4R $\alpha$  proteins in non-human animals, e.g., rodents, e.g., mice or rats, under the control, at least in part, of the endogenous IL-4 and/or IL-4R $\alpha$  regulatory elements.

#### Genetically Modified Non-Human Animals

**[0133]** Genetically modified non-human animals are provided herein whose endogenous IL-4 gene and/or IL-4R $\alpha$  gene has been replaced in whole or in part, at an endogenous IL-4 locus and/or the IL-4R $\alpha$  locus, with a human IL-4 nucleic acid and/or human IL-4R $\alpha$  nucleic acid to form a modified IL-4 gene and/or modified IL-4R $\alpha$  gene which encodes a human or humanized IL-4 and/or human or humanized IL-4R $\alpha$  protein.

**[0134]** The phrase “non-human animal” as used herein refers to any vertebrate organism that is not a human. In some embodiments, the non-human animal is a mammal. In specific embodiments, the non-human animal is a rodent such as a rat or a mouse.

**[0135]** In one aspect, genetically modified rodents, e.g., mice or rats, are provided whose endogenous rodent IL-4 gene has been replaced in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid.

**[0136]** The replacement involves a replacement of at least one exon, i.e., one or more exons, of a rodent IL-4 gene with a human nucleic acid comprising at least one exon of a human IL-4 gene. In some embodiments, a contiguous rodent genomic fragment which includes exon 1 starting from the

ATG initiation codon through exon 4 of a rodent IL-4 gene has been replaced with a contiguous human genomic fragment including exon 1 starting from the ATG initiation codon through exon 4 of a human IL-4 gene. In a specific embodiment, the rodent is a mouse, and a contiguous mouse genomic fragment of about 6.3 kb at an endogenous mouse IL-4 locus, including exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4, is deleted and replaced with about 8.8 kb of a human IL-4 nucleic acid sequence comprising exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 of the human IL-4 gene.

**[0137]** In some embodiments, the replacement results in a modified, humanized IL-4 gene at an endogenous IL-4 gene locus, wherein the expression of the modified IL-4 gene is under control of the endogenous regulatory elements at the endogenous IL-4 locus. The term "regulatory elements" as used herein refer to transcriptional regulatory sequences, including both 5' transcriptional regulatory sequences such as promoter, enhancer, and suppressor elements, and 3' transcriptional regulatory sequences such as a transcriptional termination sequence. In some embodiments, the expression of a modified IL-4 gene is under control of the endogenous 5' regulatory elements. In other embodiments, the expression of a modified IL-4 gene is under control of the endogenous 3' regulatory elements. In certain embodiments, the expression of a modified IL-4 gene is under control of the endogenous 5' and 3' regulatory elements.

**[0138]** The modified, humanized IL-4 gene formed at an endogenous IL-4 locus encodes a human or humanized IL-4 protein. The term "humanized" refer to nucleic acids or proteins which include portions or sequences of a gene or protein found in a non-human animal (e.g., a rodent such as mouse or rat), and also include portions or sequences that differ from those found in a non-human animal but instead correspond to (identical with) portions or sequences of the counterpart human gene or protein. The modified, humanized IL-4 gene can encode an IL-4 protein that is at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical, or is 100% identical, with a human IL-4 protein (e.g., the human IL-4 protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000589.3).

**[0139]** A genetically modified rodent having a replacement of an endogenous rodent IL-4 gene in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid, can be homozygous or heterozygous with respect to the replacement. In some embodiments, the genetically modified rodent is heterozygous with respect to the replacement, i.e., only one of the two copies of the endogenous rodent IL-4 gene has been replaced with a human IL-4 nucleic acid. In other embodiments, the genetically modified rodent is homozygous with respect to the replacement, i.e., both copies of the endogenous rodent IL-4 gene have been replaced with a human IL-4 nucleic acid.

**[0140]** The genetically modified rodent expresses a human or humanized IL-4 protein in the serum. In some embodiments, the genetically modified rodent does not express endogenous rodent IL-4 protein. In one embodiment, the serum of the rodent that expresses a human or humanized IL-4 protein has approximately the same level of IL-4 protein as a rodent that expresses a functional, endogenous IL-4 protein, e.g., a wild-type rodent (e.g., a rodent that expresses

functional endogenous IL-4 protein, but does not comprise a replacement of an endogenous IL-4 gene in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid). By "approximately the same level" it is meant a level that falls within 25%, 20%, 15%, 10%, 5% or less in either direction (i.e., greater than or less than) of the level in a wild-type rodent. In other embodiments, the rodent expresses a human or humanized IL-4 protein in serum at a concentration of at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190% or 200% of the level of IL-4 protein present in the serum of an age-matched rodent that expresses functional endogenous IL-4 protein, but does not comprise a replacement of an endogenous IL-4 gene in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid.

**[0141]** In some embodiments, a genetically modified rodent having a replacement of an endogenous rodent IL-4 gene in whole or in part with a human IL-4 nucleic acid and expressing a human or humanized IL-4 protein in the serum has a normal immune system, i.e., the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of the rodent expressing human or humanized IL-4 protein are similar to the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of a rodent that expresses functional endogenous IL-4 protein and does not have a replacement of an endogenous rodent IL-4 gene in whole or in part with a human IL-4 nucleic acid.

**[0142]** In additional embodiments, a genetically modified rodent having a replacement of an endogenous rodent IL-4 gene in whole or in part with a human IL-4 nucleic acid and expressing a human or humanized IL-4 protein, also includes a replacement of the endogenous rodent IL-4R $\alpha$  gene in whole or in part, at an endogenous IL-4 R $\alpha$  locus, with a human IL-4R $\alpha$  nucleic acid, and as a result, also expresses a human or humanized IL-4R $\alpha$  protein.

**[0143]** In another aspect, genetically modified rodents, e.g., mice or rats, are provided whose endogenous rodent IL-4R $\alpha$  gene has been replaced in whole or in part, at an endogenous IL-4 R $\alpha$  locus, with a human IL-4R $\alpha$  nucleic acid.

**[0144]** The replacement involves replacement of at least one exon, i.e., one or more exons, of a rodent IL-4R $\alpha$  gene with a human nucleic acid comprising at least one exon of a human IL-4R $\alpha$  gene. In some embodiments, the replacement involves replacement of at least one of the exons of a rodent IL-4R $\alpha$  gene encoding the rodent ectodomain with at least one of the exons of human IL-4R $\alpha$  gene encoding the human ectodomain. In some embodiments, the replacement involves replacement with a human nucleic acid comprising at least 2, 3 or 4 of the 5 exons encoding the ectodomain of a human IL-4R $\alpha$  gene. In other embodiments, a contiguous rodent genomic fragment which includes exon 1 starting from the ATG initiation codon through exon 5 of a rodent IL-4R $\alpha$  gene has been replaced with a genomic fragment including exon 1 starting from the ATG initiation codon through exon 5 of a human IL-4R $\alpha$  gene. In a specific embodiment, the rodent is a mouse, and a contiguous mouse genomic fragment of about 7.1 kb at an endogenous mouse IL-4R $\alpha$  locus, including exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5, is deleted and replaced with about 15.6 kb of a human IL-4R $\alpha$  nucleic acid sequence comprising exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 of the human IL-4R $\alpha$  gene.

**[0145]** In some embodiments, the replacement results in a modified, humanized IL-4R $\alpha$  gene at an endogenous IL-4R $\alpha$

gene locus, wherein the expression of the modified IL-4R $\alpha$  gene is under control of the endogenous regulatory elements at the endogenous IL-4R $\alpha$  locus.

**[0146]** The modified, humanized IL-4R $\alpha$  gene formed at an endogenous IL-4R $\alpha$  locus encodes a human or humanized IL-4R $\alpha$  protein. In some embodiments, the modified, humanized IL-4R $\alpha$  gene encodes an IL-4R $\alpha$  protein that is at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical, or is 100% identical, with a human IL-4R $\alpha$  protein (e.g., the human IL-4 protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000418.3). In other embodiments, the modified IL-4R $\alpha$  gene encodes a humanized IL-4R $\alpha$  protein which comprises an ectodomain that is at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical, or is 100% identical, with the ectodomain of a human IL-4R $\alpha$  protein (e.g., the human IL-4 protein encoded by the nucleic acid set forth in GenBank Accession No. NM\_000418.3). In specific embodiments, the transmembrane and cytoplasmic domains of the humanized IL-4R $\alpha$  protein are identical with the transmembrane and cytoplasmic domains of the endogenous rodent IL-4R $\alpha$  protein.

**[0147]** A genetically modified rodent having a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part, at an endogenous IL-4R $\alpha$  locus, with a human IL-4R $\alpha$  nucleic acid, can be homozygous or heterozygous with respect to the replacement. In some embodiments, the genetically modified rodent is heterozygous with respect to the replacement, i.e., only one of the two copies of the endogenous rodent IL-4R $\alpha$  gene has been replaced with a human IL-4R $\alpha$  nucleic acid. In other embodiments, a genetically modified rodent is homozygous with respect to the replacement, i.e., both copies of the endogenous rodent IL-4R $\alpha$  gene have been replaced with a human IL-4R $\alpha$  nucleic acid.

**[0148]** The genetically modified rodent disclosed herein expresses a human or humanized IL-4R $\alpha$  protein on immune cells, e.g., B and T cells. In some embodiments, the genetically modified rodent does not express endogenous rodent IL-4R $\alpha$  protein. In one embodiment, the immune cells of the rodent that expresses a human or humanized IL-4R $\alpha$  protein have approximately the same level of IL-4R $\alpha$  protein on immune cells as a rodent that expresses a functional, endogenous IL-4R $\alpha$  protein on immune cells of a wild-type rodent that expresses a functional, endogenous IL-4R $\alpha$  protein and does not express the human or humanized IL-4R $\alpha$  protein. In other embodiments, the rodent expresses human or humanized IL-4R $\alpha$  protein on immune cells at an amount of at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 110%, 120%, 130%, 140%, 150%, 160%, 170%, 180%, 190% or 200% of the amount of IL-4R $\alpha$  protein present on immune cells of an age-matched rodent that expresses functional endogenous IL-4R $\alpha$  protein, but does not comprise a replacement of an endogenous IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid.

**[0149]** In some embodiments, a genetically modified rodent having a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid and expressing a human or humanized IL-4R $\alpha$  protein has a normal immune system, i.e., the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of the rodent expressing human or humanized IL-4R $\alpha$  protein are similar to the number of immune cells, e.g., B and T cells, in the blood, plasma or serum of a wild-type rodent (e.g., a rodent that expresses functional endogenous IL-4R $\alpha$  protein and

does not have a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid).

**[0150]** In some embodiments, a genetically modified rodent having a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid and expressing a human or humanized IL-4R $\alpha$  protein is capable of and functional in mediating IL-4 dependent signaling and IL-13 dependent signaling. For example, a humanized IL-4R $\alpha$  protein having the ectodomain of a human IL-4R $\alpha$  protein, expressed on immune cells of a genetically modified rodent, interacts with human IL-4 and mediates human IL-4 dependent signaling via forming Type I receptor (see FIG. 1). Such humanized IL-4R $\alpha$  protein having the ectodomain of a human IL-4R $\alpha$  protein also interacts with human and mouse IL-13, and mediates IL-13 dependent signaling via forming Type II receptor (see FIG. 1). The functionality of a humanized IL-4R $\alpha$  protein expressed in a genetically modified rodent can be evaluated in various assays known in the art, including those specifically described in the examples hereinbelow, such as an assay that measures IL-4 induced IgE class switching using primary B cells derived from a genetically modified rodent.

**[0151]** In additional embodiments, a genetically modified rodent having a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid and expressing a human or humanized IL-4R $\alpha$  protein, also includes a replacement of the endogenous rodent IL-4 gene in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid, and as a result, also expresses a human or humanized IL-4.

**[0152]** In a further aspect, doubly humanized rodents, e.g., mice or rats, are provided whose endogenous rodent IL-4 gene has been replaced in whole or in part, at an endogenous IL-4 locus, with a human IL-4 nucleic acid, and whose endogenous rodent IL-4R $\alpha$  gene has also been replaced in whole or in part, at an endogenous IL-4 R $\alpha$  locus, with a human IL-4R $\alpha$  nucleic acid. Such doubly humanized rodents can be homozygous or heterozygous with respect to each humanization replacement. In a specific embodiment, the doubly humanized rodent is homozygous with respect to both humanized IL-4 and humanized IL-4R $\alpha$ .

**[0153]** The genetic modification to an endogenous rodent IL-4 gene in a doubly humanized rodent includes those modifications or replacements described hereinabove for a genetically modified rodent having a replacement of an endogenous rodent IL-4 gene in whole or in part with a human IL-4 nucleic acid. Similarly, the genetic modification to an endogenous rodent IL-4R $\alpha$  gene in a doubly humanized rodent includes those modifications or replacements described hereinabove for a genetically modified rodent having a replacement of an endogenous rodent IL-4R $\alpha$  gene in whole or in part with a human IL-4R $\alpha$  nucleic acid. Thus, the features disclosed hereinabove with respect to humanization of the rodent IL-4 gene and with respect to humanization of the rodent gene, respectively, are incorporated herein specifically for a doubly humanized rodent.

**[0154]** In specific embodiments, a doubly humanized rodent, e.g., a mouse or rat, is provided that expresses a human IL-4 protein and a humanized IL-4R $\alpha$  protein, wherein the humanized IL-4R $\alpha$  protein includes the ectodomain of a human IL-4R $\alpha$  protein and includes the transmembrane and cytoplasmic domains of the rodent's endogenous IL-4R $\alpha$  protein. In particular embodiments, the expression of the human IL-4 protein and the humanized

IL-4R $\alpha$  protein are under control of the endogenous rodent regulatory sequences at the endogenous rodent IL-4 locus and rodent IL-4R $\alpha$  locus, respectively.

**[0155]** In some embodiments, a doubly humanized rodent has a normal immune system (i.e., the number of immune cells is approximately the same as a wild-type rodent), has approximately the same level of IL-4 protein in the serum, and expresses approximately the same amount of IL-4R $\alpha$  protein on immune cells, as a wild-type rodent, a wild-type rodent being a rodent that expresses functional, endogenous IL-4 protein and IL-4R $\alpha$  protein and does not express human or humanized IL-4 protein or IL-4R $\alpha$  protein.

**[0156]** In particular embodiments, a doubly humanized rodent exhibits a functional IL-4 signaling pathway. By “functional IL-4 signaling pathway” it is meant that both a human or humanized IL-4 protein, and a human or humanized IL-4R $\alpha$  protein, are expressed in a doubly humanized rodent and interact with each other in the doubly humanized rodent so as to effectively mediate downstream signal transduction and carry out the biological activities of a normal IL-4 signaling pathway. The biological activities of a normal IL-4 signaling pathway are described hereinabove and illustrated in FIG. 1, including those mediated through Type I receptor such as initiation and maintenance of the Th2 differentiation, activation and growth of B cells, class switching to IgE and IgG4, and those mediated through Type II receptor signaling such as Goblet cell hyperplasia, sub-epithelial fibrosis, and tissue remodeling. For example, a functional IL-4 signaling pathway in a doubly humanized rodent is reflected by an inflammatory phenotype characterized by, e.g., increased IgE in circulation, airway inflammation and/or eosinophilic infiltrating cells in response to a house dust mite challenge, which phenotype is also observed in wild-type rodents without the double humanization.

#### Methods of Making a Genetically Modified Non-Human Animal

**[0157]** A genetically modified non-human animal such as a rodent can be made using methods known in the art. For example, a targeting vector can be made that contains a human nucleic acid (such as a human IL-4 or IL-4R $\alpha$  gene, in whole or in part), flanked by non-human animal homologous upstream and downstream regions. The targeting construct can also contain a drug selection cassette (e.g., a floxed hygromycin selection cassette, which can be subsequently removed by a transient Cre-expression vector), which is positioned 3' to the human nucleic acid. The targeting vector can be introduced into the genome of a non-human animal cell, e.g., an embryonic stem (ES) cell (such as a mouse ES cell) by electroporation, for example. Correctly targeted ES cell clones can then be introduced into an early stage embryo (e.g., 8-cell stage mouse embryo). Non-human animals fully derived from correctly targeted ES cells are identified based on, for example, allele analysis. For non-human animals where suitable genetically modifiable ES cells are not readily available, other methods can be employed to make a non-human animal comprising genetic modifications as described herein. Such methods include, e.g., modifying a non-ES cell genome (e.g., a fibroblast or an induced pluripotent cell) and employing nuclear transfer to transfer the modified genome to a suitable cell, e.g., an oocyte, and gestating the modified cell (e.g., the modified oocyte) in a non-human animal under suitable conditions to form an embryo.

#### Methods of Employing a Genetically Modified Non-Human Animal

**[0158]** In one aspect, genetically modified IL-4 and/or IL-4R $\alpha$  non-human animals disclosed herein are used for evaluating the pharmacodynamics (PD) and therapeutic efficacy of human-specific IL-4 and/or IL-4R $\alpha$  antagonists, e.g., neutralizing anti-IL-4 and/or anti-IL-4R $\alpha$  antibodies (e.g., dupilumab) in various disease models, as further illustrated in the examples below.

**[0159]** In some embodiments, the present invention provides a method of screening for human-specific IL-4 or IL-4R $\alpha$  antagonists using a doubly humanized IL-4 and IL-4R $\alpha$  mice disclosed herein.

**[0160]** By “IL-4 or IL-4R $\alpha$  antagonists” it is meant molecules (e.g., antibodies) that block, suppress or inhibit one or more biological functions mediated by IL-4 or IL-4R $\alpha$ . “Human-specific IL-4 or IL-4R $\alpha$  antagonists” refer to antagonists that are specific to the human IL-4 or IL-4R $\alpha$ , and substantially do not act on rodent IL-4 or IL-4R $\alpha$ .

**[0161]** In specific embodiments, the method of screening utilizes a doubly humanized mouse that expresses a human IL-4 protein, and a humanized IL-4R $\alpha$  protein, wherein the humanized IL-4R $\alpha$  protein includes the ectodomain of a human IL-4R $\alpha$  protein, linked to the transmembrane and cytoplasmic domains of the endogenous mouse IL-4R $\alpha$  protein, and wherein the mouse does not express mouse IL-4 or mouse IL-4R $\alpha$ .

**[0162]** In some embodiments, the method of screening for a human-specific IL-4 or IL-4R $\alpha$  antagonist comprises administering an agent to a genetically modified rodent that is doubly humanized for IL-4 and IL-4R $\alpha$  as described herein, determining an effect of the agent on a biological function mediated by the IL-4/IL-4R $\alpha$  signaling pathway, and identifying the agent as a human-specific IL-4 or IL-4R $\alpha$  antagonist if it antagonizes the function mediated by the IL-4/IL-4R $\alpha$  signaling pathway in the genetically modified rodent.

**[0163]** In one embodiment, the agent comprises an immunoglobulin variable domain that binds IL-4 or IL-4R $\alpha$ . In one embodiment, the agent specifically binds human IL-4 or IL-4R $\alpha$ , but not rodent IL-4 or IL-4R $\alpha$ . In one embodiment, the agent is an antibody. In a specific embodiment, the agent is an antibody that specifically binds human IL-4R $\alpha$ , but not rodent IL-4R $\alpha$ .

**[0164]** In one embodiment, the method of screening utilizes a doubly humanized mouse that expresses a human IL-4 protein, and a humanized IL-4R $\alpha$  protein, wherein the humanized IL-4R $\alpha$  protein includes the ectodomain of a human IL-4R $\alpha$  protein, linked to the transmembrane and cytoplasmic domains of the endogenous mouse IL-4R $\alpha$  protein, and wherein the mouse does not express murine IL-4 or murine IL-4R $\alpha$ .

**[0165]** In some embodiments, the method of screening includes the steps of inducing in a doubly humanized rodent as described herein a disease associated with IL-4/IL-4R $\alpha$  signaling, administering an agent to the rodent, determining whether the agent ameliorates the disease, and identifying the agent as a human-specific IL-4 or IL-4R $\alpha$  antagonist suitable for treating the disease if the agent ameliorates the disease.

**[0166]** By “disease associated with IL-4/IL-4R $\alpha$  signaling” it is meant a disease in which the biological function mediated by IL-4/IL-4R $\alpha$  signaling is implicated. Examples of diseases associated with IL-4/IL-4R $\alpha$  signaling include, e.g., inflammatory diseases or disorders, such as asthma, atopic dermatitis, chronic obstructive pulmonary disease

(COPD) (which may result at least in part from cigarette smoke), inflammatory bowel disease, multiple sclerosis, arthritis, allergic rhinitis, eosinophilic esophagitis and psoriasis. Asthma can be eosinophilic or non-eosinophilic asthma, and steroid sensitive or steroid resistant asthma.

**[0167]** In some embodiments, the disease associated with IL-4/IL-4R $\alpha$  signaling is airway inflammation, which can be induced in a rodent by intranasal administration of an allergen (e.g., house dust mite extract) in one or more doses for a period of time. The effect of an agent can be determined by measuring whether the extent of airway inflammation (reflected by e.g., mucus accumulation, eosinophilic infiltrating cells in bronchoalveolar lavage fluid, levels of total circulating IgE, and/or alteration in expression profile measurable by microarray expression analysis) is reduced as a result of the administration of the agent. The allergen used for inducing airway inflammation and the agent being tested can be administered simultaneously or at different times. In some embodiments, the allergen is given to the rodent in one or more doses, and the agent being tested is administered to the rodent after at least one dose of the allergen has been given to the rodent.

**[0168]** In some embodiments, the disease associated with IL-4/IL-4R $\alpha$  signaling is skin inflammation or atopic dermatitis, which can be induced in a rodent by creating skin injury and exposing the injured skin to an allergen (e.g., bacterial toxin or house dust mite extract) in one or more doses for a period of time. The effect of an agent can be determined by measuring whether skin inflammation is reduced as a result of administration of the agent.

**[0169]** In a further aspect, triply humanized non-human animals, i.e., non-human animals whose IL-4, IL-4R $\alpha$ , and IL-33 genes have been humanized, are used to evaluate the pharmacodynamics (PD) and therapeutic efficacy of candidate compounds such as, e.g., human-specific IL-4 and/or IL-4R $\alpha$  antagonists, and human-specific IL-33 antagonists.

**[0170]** By "IL-33 antagonists" it is meant molecules (e.g., antibodies) that block, suppress or inhibit one or more biological functions or signaling mediated by IL-33. "Human-specific IL-33 antagonists" refer to antagonists that are specific to the human IL-33, and substantially do not act on rodent IL-33. IL-33 is known to stimulate signal transduction through ST2 and IL-1 RAcP, which is diminished in the presence of an antagonist, such as an IL-33 antibody. Inhibition of IL-33 signal transduction through ST2 and IL-1 RAcP can be determined by assaying for IL-33 signal transduction in an in vitro or in vivo assay, such as those described in US Published Application 2014/0271658 A1, the entire contents of which are incorporated herein by reference. For example, an assay such as that described in US Published Application 2014/0271658 A1 can be used to assess the effect of an antibody to IL-33 on lung inflammation in allergen-sensitized animals that are homozygous for expression of human IL-33. An IL-33 antibody that is effective as an IL-33 antagonist should demonstrate a trend towards reduction in inflammatory cells in the lung, as well as a trend towards reduction in cytokines such as IL-4 and IL-5.

**[0171]** In specific embodiments, a triply humanized non-human animal is used herein to evaluate candidate compounds, wherein the triply humanized animal is a triply humanized mouse that expresses a human IL-4 protein, a humanized IL-4R $\alpha$  protein which includes the ectodomain of a human IL-4R $\alpha$  protein linked to the transmembrane and cytoplasmic domains of a mouse IL-4R $\alpha$  protein, and a

human IL-33 protein, wherein the mouse does not express mouse IL-4, mouse IL-4R $\alpha$  or mouse IL-33.

**[0172]** In some embodiments, a triply humanized non-human animal is used to evaluate the pharmacodynamics (PD) and therapeutic efficacy of a candidate compound, such as, e.g., a human-specific IL-4 and/or IL-4R $\alpha$  antagonist, or a human-specific IL-33 antagonist. For example, a human-specific IL-4 antibody, a human-specific IL-4R $\alpha$  antibody, and a human-specific IL-33 antibody, can be tested individually in a triply humanized animal (such as a rodent, e.g., mouse or rat), and their PD profiles and therapeutic efficacies can be evaluated and compared.

**[0173]** In other embodiments, a triply humanized non-human animal is used to evaluate the efficacy of a combination of compounds, e.g., a combination of a human specific IL-4 and/or IL-4R $\alpha$  antagonist antibody, with a human-specific IL-33 antagonist antibody, as compared to the efficacy of the compounds when used individually to determine, for example, whether the combination of compounds exhibits a synergistic therapeutic effect. In specific embodiments, a combination of a human specific IL-4 antibody and a human-specific IL-33 antibody are tested in a triply humanized non-human animal. In other specific embodiments, a combination of a human specific IL-4R $\alpha$  antibody and a human-specific IL-33 antibody are tested in a triply humanized non-human animal.

**[0174]** To evaluate a candidate compound or a combination of compounds, a disease associated with the IL-4/IL-4R $\alpha$  signaling and the IL-33 signaling can be induced in the triply humanized animal. Examples of diseases associated with the IL-4/IL-4R $\alpha$  signaling and the IL-33 signaling include, e.g., inflammatory diseases or disorders, such as asthma, atopic dermatitis, chronic obstructive pulmonary disease (COPD) (which may result at least in part from cigarette smoke), inflammatory bowel disease, multiple sclerosis, arthritis, allergic rhinitis, eosinophilic esophagitis and psoriasis. Asthma can be eosinophilic or non-eosinophilic asthma, and steroid sensitive or steroid resistant asthma. The effect of a compound or a combination of compounds can be assessed similarly to an IL-4/IL-4R $\alpha$  doubly humanized animal as described hereinabove.

**[0175]** The present invention is further illustrated by the following, non-limiting examples.

#### Example 1

##### Replacement of the Endogenous Mouse IL-4 Gene with a Human IL-4 Gene

**[0176]** The 8.8 kb human IL-4 gene containing the coding portion of exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 of the human IL-4 gene replaced 6.3 kb of the murine IL-4 gene locus spanning the coding portion of exon 1 starting from the ATG initiation codon through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4. See FIG. 2A.

**[0177]** A targeting construct for replacing the mouse with the human IL-4 gene in a single targeting step was constructed using VelociGene<sup>®</sup> genetic engineering technology (see Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech*, 21(6):652-659). Mouse and human IL-4 DNA were obtained from bacterial artificial chromo-



some (BAC) clones bMQ-41A12 and RP11-17K19, respectively. Briefly, an SbfI linearized targeting construct generated by gap repair cloning containing mouse IL-4 upstream and downstream homology arms flanking a 8.8 kb human IL-4 sequence extending from the ATG codon in exon 1 through exon 4 (including the 3' untranslated region) and a portion of the 3' region downstream of exon 4 (genomic coordinates: GRCh37: chr5:132,009,743-132,018,576 (+ strand)) and a floxed hygro selection cassette, was electroporated into F1H4 mouse embryonic stem (ES) cells (C57BL/6x129 F1 hybrid). Correctly targeted ES cells (MAID 879) were further electroporated with a transient Cre-expressing vector to remove the drug selection cassette. Targeted ES cell clones without drug cassette (MAID 1553) were introduced into an 8-cell stage SW mouse embryo by the VelociMouse® method (see, U.S. Pat. Nos. 7,294,754, 7,576,259, 7,659,442, and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses, Nature Biotech. 25(1):91-99). VelociMice® (F0 mice fully derived from the donor ES cell) bearing the humanized IL-4 gene were identified by genotyping for loss of mouse allele and gain of human allele using a modification of allele assay (see, Valenzuela et al. (2003)).

**[0178]** Correctly targeted ES cell clones were identified by a loss-of-native-allele (LONA) assay (Valenzuela et al. 2003) in which the number of copies of the native, unmodified IL-4 gene were determined by two TaqMan™ quantitative polymerase chain reactions (qPCRs) specific for sequences in the mouse IL-4 gene that were targeted for deletion. The qPCR assays comprised the following primer-probe sets (written 5' to 3'): upstream forward primer, CATGCACGGA GATG-GATGTG (SEQ ID NO:1); upstream reverse primer, GAC-CCCTCAG GTCCACTTAC C (SEQ ID NO:2); upstream probe, FAM-AACGTCCTCA CAGCAACGA-MGB (SEQ ID NO:3); downstream forward primer, GTGCCAGGT GTGCTCATG (SEQ ID NO:4); downstream reverse primer, CGCCTGCCTC CTCACCTTAT C (SEQ ID NO:5); downstream probe, FAM-ATCTGCTTCA CCATCCACT-MGB (SEQ ID NO:6); in which FAM refers to the 5-carboxyfluorescein fluorescent probe and BHQ refers to the fluorescence quencher of the black hole quencher type (Biosearch Technologies). DNA purified from ES cell clones that have taken up the targeting vector and incorporated in their genomes was combined with TaqMan™ Gene Expression Master Mix (Life Technologies) according to the manufacturer's suggestions in a 384-well PCR plate (MicroAmp™ Optical 384-Well Reaction Plate, Life Technologies) and cycled in an Applied Biosystems Prism 7900HT, which collects fluorescence data during the course of the PCRs and determines a threshold cycle (Ct), the fractional PCR cycle at which the accumulated fluorescence reaches a pre-set threshold. The upstream and downstream IL-4-specific qPCRs and two qPCRs for non-targeted reference genes were run for each DNA sample. The differences in the Ct values ( $\Delta Ct$ ) between each IL-4-specific qPCR and each reference gene qPCR were calculated, and then the difference between each  $\Delta Ct$  and the median  $\Delta Ct$  for all samples assayed was calculated to obtain  $\Delta\Delta Ct$  values for each sample. The copy number of the IL-4 gene in each sample was calculated from the following formula: copy number =  $2 \times 2^{-\Delta\Delta Ct}$ . A correctly targeted clone, having lost one of its native copies, will have a IL-4 gene copy number equal to one. Confirmation that the human IL-4 gene sequence replaced the deleted mouse IL-4 gene sequence in

the humanized allele was confirmed by a TaqMan™ qPCR assay that comprises the following primer-probe sets (written 5' to 3'): human forward primer, GCCTGGACCA AGACTCTGT (SEQ ID NO:7); human reverse primer, ACCGTGGGAC GGCTTCTTAC (SEQ ID NO:8); human upstream probe, FAM-CACCGAGTTG ACCGTAACAG ACATC-BHQ (SEQ ID NO:9). Confirmation that the hygro selection cassette was inserted with the human IL-4 gene sequence in the humanized allele was confirmed by a TaqMan™ qPCR assay that comprises the following primer-probe sets (written 5' to 3'): hygro forward primer, TGCG-GCCGAT CTTAGCC (SEQ ID NO:10); hygro reverse primer, TTGACCGATT CCTTGCGG (SEQ ID NO:11); hygro probe, FAM-ACGAGCGGGT TCGGCCATT C-BHQ (SEQ ID NO:12).

**[0179]** The same LONA assay was used to assay DNA purified from tail biopsies for mice derived from the targeted ES cells to determine their IL-4 genotypes and confirm that the humanized IL-4 allele had transmitted through the germline. Two pups heterozygous for the replacement are bred to generate a mouse that is homozygous for the replacement of the endogenous mouse IL-4 gene by the human IL-4 gene. Pups that are homozygous for the replacement are used for phenotyping.

**[0180]** The upstream junction of the murine IL-4 locus and the sequence containing the human IL-4 gene is designed to be within 5'-TGCTGATTGG CCCAGAATAA CTGACAACTCT GGTGTAATAA AATTTTCCAA TGTAACCTCA TTTTCCCTTG GTTTCAGCAA CTTA-CTACTCT ATATATAGAG AGACCTCTGC CAGCAT-TGCA TTGTTAGCAT CTCTTGATAA ACTTAATTGT CTCTCGTCAC TGACGGCACA GAGCTATTG(A TGGGTCTCAC CTCCCAACTG CTCCCCCTC TGT-TCTTCTCT GCTAGCATGT GCCGGCAACT TTGTC-CACGG ACACAAGTGC GATATCACCCT TACAGGAGAT CATCAAAACT TTGAACAGCC TCACAGACA GMG) GTGAGT ACCATCTGG CACCATCTCT CCAGATGTTT TGGTGATGCT CTCAGTATTT CTAGGCATGA AAACGTTAAC AGCTGCTAGA GAAGTTGGAA CTG-GTGGTTG GTGGCAGTCC AGGGCACACA GCGAG-GCTTC TCCCCTGC (SEQ ID NO:13), wherein the human IL-4 sequences are italicized and the IL-4 coding sequences are bracketed. The downstream junction of the sequence containing the human IL-4 gene and the floxed hygro selection cassette is designed to be within 5'-TGTTTATTTT GCAG (AATTCC TGTCCCTGTGA AGGAAGCCAA CCAGAG-TACG TTGGAAAAC TCTTGAAAAG GCTAAAGACG ATCATGAGAG AGAAATATC AAAGTGTTCG AGCT-GA)ATAT TTAAATTTAT GAGTTTTTGA TAGCTTTATT TTTTAAAGTAT TTATATATTT ATAACCTCATC ATAAAATAM GTATATATAG AATCTAACAG CAATG-GCATT TAAATGATTG GCTATGTTTA CTTGACAAAT GAAATATGG TTTGCAACTT TTAGGGAAT CAATT-TAGTT TACCMGAGA CTATAMTGC TATGGGAGCA AAACAGGAAA GACCACTTCC CCCTCGAGGG GTTC-CCTCTC GAGTTAGGGA CATAACACAC AAGATAATTA AAGAACACAA GGCCATACAA GAT-GCGGCCG CACCGGTATA ACTTCGTATA AGGTATC-CTA TACGAAGTTA TATGCATGGC CTCCCGCCCG GGTTTTGGCG CCTCCCGCG GCGCCCCCT CTT-CACGGCG AGCGCTGCCA CGTCAGACGA AGGGCG-CAGC GAGCGTCTCTG ATCCT (SEQ ID NO:14), wherein the human IL-4 sequences are italicized and the IL-4 coding sequences are bracketed. The downstream junction of the

sequence of the floxed hygro selection cassette and the murine IL-4 locus is designed to be within 5'-TGC-CAAGTTC TAATTCCATC AGACCTCGAC CTGCAGC-CGG CGCGCCATAA CTTCGTATAA GGTATCCTAT ACGAAGTTAT CTCGAGAGGA GTTCCCACCC TTCT-CAAGAG CATAATGCGC AGATCATTA A GGGACA-GATG CAGGCTGGGG AGACGGTTC A GCAGTTAGGA GTACCTGTTG CTCTTCCAGA GGACCCAGGT TCAAT-TCCCG GCACTCACAT AGCAGCTTAA AACAATAACT CAAGTTCCTGG GGGAGCTGAT GCTCTCCTCT GGC-CTCCTGT GGAGGTACAC AGACCACATG CCTGTAG-GCA AGACACCCAC ACACATAAAA ACAAATAAAA ATAAGGATAG AAAGGCCAGG GGGATGAATC CAGAGGTAGA AGAAAACCTTA TTCCCTGGAA TTGTCCTCTG ACTCCCTCC CAAAACCTCT AACACGCAT (SEQ ID NO:15), wherein the hygro cassette sequences are italicized.

#### Example 2

##### Replacement of the Endogenous Mouse IL-4R $\alpha$ Ectodomain Gene Sequence with a Human IL-4R $\alpha$ Ectodomain Gene Sequence

**[0181]** The 15.6 kb human IL-4R $\alpha$  gene containing exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5 of the human IL-4R $\alpha$  gene replaced 7.1 kb of the murine IL-4R $\alpha$  gene locus spanning coding exon 1 starting from the ATG initiation codon through exon 5 and a portion of intron 5. Mouse exons 6 through 9 were retained; only exons 1 through 5 (i.e., the ectodomain) were humanized. See FIG. 2B.

**[0182]** A targeting construct for replacing the mouse with the human IL-4R $\alpha$  gene in a single targeting step was constructed using VelociGene<sup>®</sup> genetic engineering technology (see Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech*, 21(6):652-659). Mouse and human IL-4R $\alpha$  DNA were obtained from bacterial artificial chromosome (BAC) clones RP23-136G14 and RP11-166E24, respectively. Briefly, a NotI linearized targeting construct generated by gap repair cloning containing mouse IL-4R $\alpha$  gene upstream and downstream homology arms flanking a 15.6 kb human IL-4R $\alpha$  sequence extending from the ATG codon in exon 1 through exon 5 and a portion of intron 5 (genomic coordinates: GRCh37: chr16:27,351,525-27,367,111 (+ strand)) and a floxed neo selection cassette, was electroporated into F1H4 mouse embryonic stem (ES) cells (C57BL/6 $\times$ 129 F1 hybrid). Correctly targeted ES cells (MAID 803) were further electroporated with a transient Cre-expressing vector to remove the drug selection cassette. Targeted ES cell clones without drug cassette (MAID 1444) were introduced into an 8-cell stage SW mouse embryo by the VelociMouse<sup>®</sup> method (see, U.S. Pat. Nos. 7,294,754, 7,576,259, 7,659,442, and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses, *Nature Biotech*. 25(1):91-99). VelociMice<sup>®</sup> (F0 mice fully derived from the donor ES cell) bearing the humanized IL-4R $\alpha$  gene were identified by genotyping for loss of mouse allele and gain of human allele using a modification of allele assay (see, Valenzuela et al. (2003)).

**[0183]** Correctly targeted ES cell clones were identified by a loss-of-native-allele (LONA) assay (Valenzuela et al. 2003) in which the number of copies of the native, unmodified

IL-4R $\alpha$  gene were determined by two TaqMan<sup>™</sup> quantitative polymerase chain reactions (qPCRs) specific for sequences in the mouse IL-4R $\alpha$  gene that were targeted for deletion. The qPCR assays comprised the following primer-probe sets (written 5' to 3'): upstream forward primer, CCGCTGGCAT GTGTATTGTG (SEQ ID NO:16); upstream reverse primer, TGAGTGTGGG ACCCTCAAGA G (SEQ ID NO:17); upstream probe, FAM-TGACCCAAGC CCTACATGGC CACT-BHQ (SEQ ID NO:18); downstream forward primer, TGAGGAGAGC TCACGGGAAT C (SEQ ID NO:19); downstream reverse primer, ACCCATCTCC TGCGTTTCTG (SEQ ID NO:20); downstream probe, FAM-TTGACACGCC AGCTACACTG CTCCA-BHQ (SEQ ID NO:21); in which FAM refers to the 5-carboxyfluorescein fluorescent probe and BHQ refers to the fluorescence quencher of the black hole quencher type (Biosearch Technologies). DNA purified from ES cell clones that have taken up the targeting vector and incorporated in their genomes was combined with TaqMan<sup>™</sup> Gene Expression Master Mix (Life Technologies) according to the manufacturer's suggestions in a 384-well PCR plate (MicroAmp<sup>™</sup> Optical 384-Well Reaction Plate, Life Technologies) and cycled in an Applied Biosystems Prism 7900HT, which collects fluorescence data during the course of the PCRs and determines a threshold cycle (Ct), the fractional PCR cycle at which the accumulated fluorescence reaches a pre-set threshold. The upstream and downstream IL-4R $\alpha$ -specific qPCRs and two qPCRs for non-targeted reference genes were run for each DNA sample. The differences in the Ct values ( $\Delta$ Ct) between each IL-4R $\alpha$ -specific qPCR and each reference gene qPCR were calculated, and then the difference between each  $\Delta$ Ct and the median  $\Delta$ Ct for all samples assayed was calculated to obtain  $\Delta\Delta$ Ct values for each sample. The copy number of the IL-4R $\alpha$  gene in each sample was calculated from the following formula: copy number =  $2 \times 2^{-\Delta\Delta Ct}$ . A correctly targeted clone, having lost one of its native copies, will have a IL-4R $\alpha$  gene copy number equal to one. Confirmation that the human IL-4R $\alpha$  gene sequence replaced the deleted mouse IL-4R $\alpha$  gene sequence in the humanized allele was confirmed by a TaqMan<sup>™</sup> qPCR assay that comprises the following primer-probe sets (written 5' to 3'): human forward primer, ACCT-GCGTCT CCGACTACAT G (SEQ ID NO:22); human reverse primer, GAGCTCGGTG CTGCAATTG (SEQ ID NO:23); human probe, FAM-TGGGACCATT CATCTTC-CAC TCGCA-BHQ (SEQ ID NO:24). Confirmation that the neo selection cassette was inserted with the human IL-4R $\alpha$  gene sequence in the humanized allele was confirmed by a TaqMan<sup>™</sup> qPCR assay that comprises the following primer-probe sets (written 5' to 3'): neo forward primer, GGTG-GAGAGG CTATTCCGGC (SEQ ID NO:25); neo reverse primer, GAACACGGCG GCATCAG (SEQ ID NO:26); neo probe, FAM-TGGGCACAAC AGACAATCGG CTG-BHQ (SEQ ID NO:27).

**[0184]** The same LONA assay was used to assay DNA purified from tail biopsies for mice derived from the targeted ES cells to determine their IL-4R $\alpha$  genotypes and confirm that the humanized IL-4R $\alpha$  allele had transmitted through the germline. Two pups heterozygous for the replacement are bred to generate a mouse that is homozygous for the replacement of the endogenous mouse IL-4R $\alpha$  gene by the human IL-4R $\alpha$  gene. Pups that are homozygous for the replacement are used for phenotyping.

**[0185]** The upstream junction of the murine IL-4R $\alpha$  locus and the sequence containing the human IL-4R $\alpha$  gene is

designed to be within 5'-TGGGGGAGGG AGGCCATGAC ACAAATGAAA TGGACCCCGCTGACCCAGGA TCAG-CATCTG CCCACTCTTC TTTCTGCAGG CACCTTTTGT GTCCCCA(ATG GGGTGGCTTT GCTCTGGGCT CCT-GTTCCT GTGAGCTGCC TGGTCCTGCT GCAGGTG-GCA AGCTCTG)GTA AGTCACCACT TCTCAATCAT TCATTTGTTG GCTATTAATG GCGTGCCAGG GTCCT-GCAGT ATGTCACCTG GCC (SEQ ID NO:28), wherein the human IL-4R $\alpha$  sequences are italicized and the IL-4R $\alpha$  coding sequences are underlined. The downstream junction of the sequence containing the human IL-4R $\alpha$  gene and the floxed neo selection cassette is designed to be within 5'-GT-CAGATCGT GGAGGGTCTC GGACGAGGG TCCTGAC-CCT GGGTCTCCAG TCCTGGGAAG TGGAGCCAG GCTGTACCAT GGCTGACCTC AGCTCATGGC Tcccggctc gataactata acggtctctaa ggtagcgact cgagataact tcg-tataatg tatgtctatac gaagtatat gcatggctc cgcgcccggg tttggcgct cccggggcg cccccctct cactggcgagc gctg (SEQ ID NO:29), wherein the human IL-4R $\alpha$  sequences are italicized and the cassette sequences are in lower case. The downstream junction of the sequence of the floxed neo selection cassette and the murine IL-4R $\alpha$  locus is designed to be within 5'-tattgtttg ccaagtcta attccatcag acctcgacct gcagccccta gataactcg tataat-gtat gctatacga gttatctag gttggagctc TCTGTAGCCA GGTAACCAAG GGTCCCAGGG GAACCCCAAG TGTGGACGCG GACTGCACAT GACACAGGGC GGC-CTCCCCA TTCATGACTG TTTTCTCCT TGCAG (ACTTC CAGCTGCCCC TGATACAGCG CCTTCCACTG GGGGTCACCA TCTCCTGCCT CTGCATCCCG TTGTTTTGCC TGTTCTGTTA CTTCAGCATT ACCAA) GTGAG TTCCTGCTTT GGCTGGTGTIC TCTGGCTGGC CCTTCAGCAG TGCTCTCAGA GGTCACAGTC ATTGT-GCTGG CTGAGAAAAG (SEQ ID NO:30), wherein the mouse IL-4R $\alpha$  coding sequences are bracketed, and the neo cassette sequences are in lower case.

### Example 3

#### Generation of Doubly Humanized IL-4/IL-4R $\alpha$ Mice

**[0186]** The doubly humanized IL-4/IL-4R $\alpha$  (Il4<sup>hu/hu</sup>/Il4ra<sup>hu/hu</sup>) mice were generated as follows. ES cell clone MAID 803, comprising the humanized IL-4R $\alpha$  gene and floxed neo cassette, was electroporated with a Cre expression vector to remove the floxed neo cassette to generate ES cell clone MAID 1444, comprising the humanized IL-4R $\alpha$  gene without a drug selection cassette (see Example 2). The same targeting construct that was used to generate ES cell clone MAID 879, comprising the humanized IL-4 gene and floxed hydro cassette (see Example 1), was electroporated into ES cell clone MAID 1444 to generate 879 Het/1444 Het (Il4+/hu/Il4ra+/hu) ES cells, which were subsequently electroporated with a Cre expression vector to remove the floxed hydro cassette to generate an ES cell clone (MAID 1553/1444) comprising humanized IL-4 and IL-4R $\alpha$  genes. The ES cell clone MAID 1553/1444 without drug cassette was introduced into an 8-cell stage SW mouse embryo to generate doubly humanized IL-4/IL-4R $\alpha$  mice.

### Example 4

#### Efficacy Evaluation of Dupilumab, a Fully Human IL-4R $\alpha$ mAb, in Mice with Human IL-4 and IL4R $\alpha$ Gene Replacements

#### Methods

**[0187]** Genetically engineered mice were created using VelociGene® technology to replace both mouse full-length

IL-4 locus with 8.8 kb of human IL-4 genomic sequences (see Example 1 and FIG. 2A) and the extracellular domain (i.e., ectodomain) of mouse IL-4R $\alpha$  (CD124) gene with a 15.6 kb fragment of the corresponding human IL-4R $\alpha$  genomic DNA (see Example 2 and FIG. 2B).

**[0188]** Mice with a homozygous humanized IL-4R $\alpha$  gene were validated for expression and function of the human gene. To determine the expression of human IL-4R $\alpha$  by humanized mice, splenocytes from wild-type and humanized mice were collected and processed for fluorescent-labeled antibody sorting (FACS) analysis with fluorescent-labeled antibodies against mouse CD3, mouse CD19, human CD124, and mouse CD124. (See, e.g., Blaeser et al. (2003) Targeted inactivation of the IL-4 receptor  $\alpha$  chain I4R motif promotes allergic airway inflammation, *J Exp Med* 198(8):1189-1200).

**[0189]** To demonstrate the ligand specificities and receptor functionalities, primary cells derived from humanized IL-4R $\alpha$  mice were used. Bone marrow-derived macrophages were cultured using femoral bone marrow cells from wild-type and humanized IL-4R $\alpha$  mice in DMEM containing 10% fetal bovine serum plus 20% L-cell conditioned medium for 7 days.

**[0190]** Cells were then treated individually with 20 ng/ml of mouse IL-4, mouse IL-13, human IL-4, human IL-13, or vehicle diluted in culture medium for 20 hours. Quadruplicate samples from each condition were harvested for gene expression analysis.

**[0191]** Total RNA from these samples was extracted and amplified into cRNA by incorporating Cy3-CTP. Cy3 labeled cRNA from each sample was then hybridized to a custom Agilent array comprising of 43,538 60-mer oligos covering mouse transcriptomes. Data were extracted from scanned array images using Agilent Feature Extraction Software 9.5.

**[0192]** Differentially expressed genes between experimental groups were identified using Student's t-test ( $p < 0.05$ , fold change 1.5). An expression cluster of these genes was generated using the Pearson correlation clustering algorithm from GeneSpring GX7.3.

**[0193]** The neutralizing effect of dupilumab against IL-4 was evaluated using an in vitro IgE class-switching assay with primary B cells isolated from humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice.

**[0194]** Wild-type (WT) and humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice received a high volume (hydrodynamic) driven gene delivery of naked plasmid DNA solution for the expression of mouse IL-25 in vivo. (See, e.g., Liu et al. (1999) Hydrodynamics-based transfection in animals by systemic administration of plasmid DNA, *Gene Therapy* 6:1258-1266.) Peripheral blood was collected 8 days later to measure serum murine IgE (mIgE) levels using a commercial ELISA kit (R & D systems, MN).

**[0195]** Purified primary B cells from humanized mouse splenocytes were activated with bacterial LPS and mixed with increasing amounts of recombinant human IL-4 in a 7 day culture to induce immunoglobulin class switching. For the antibody blockade experiment, purified B cells were incubated with increasing doses of dupilumab for 30 minutes before adding 0.167 nM recombinant human IL-4 and cultured for 7 days. IgE production in the absence of IL-4 or with isotype control mAb is shown in ( $\diamond$ ) and ( $\Delta$ ), respectively. The murine IgE levels in the culture supernatants were measured using a commercial ELISA kit. (See, e.g., Moon et al. (1989) Regulation of IgG1 and IgE synthesis by interleukin 4 in mouse B cells, *Scand J Immunol* 30:355-361.)

**[0196]** Interleukin-25 (IL-25) is a cytokine produced by Th2 cells whose main activities are mediated through the production of IL-4 and IL-13 to induce tissue specific pathologies, such as increased pulmonary mucus production and goblet cell hyperplasia. (See Fort et al. (2001) IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies in vivo, *Immunity* 15(6):985-995.)

**[0197]** Lack of IL-13 protects animals from IL-25 induced pathologies in target organ. Therefore, an IL-25 driven pulmonary inflammation model was used to assess the pharmacodynamic (PD) properties of dupilumab in vivo in mice comprising humanized IL-4 and/or IL-4R $\alpha$  genes.

**[0198]** The PD responses of dupilumab on type II receptors were characterized using an IL-25-induced inflammation method by measuring pulmonary mucus accumulation in the humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice.

**[0199]** On Day 0, WT and humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice received the hydrodynamic delivery of mouse IL-25 expression vector and followed by an injection of dupilumab or isotype control mAb at the indicated doses. Additional doses of antibodies were administered every other day for a total of 4 doses. On day 8, lung tissues were collected from euthanized mice and processed lung sections were stained by periodic acid-Schiff before blinded scoring for pathological changes.

## Results

**[0200]** The humanized IL-4R $\alpha$  mice were characterized to show: (a) expression of human IL-4R $\alpha$  on primary cells from doubly humanized IL-4/IL-4R $\alpha$  (Il4<sup>hu/hu</sup>/Il4ra<sup>hu/hu</sup>) mice (see FIG. 3); (b) the change of IL-4 ligand specificities in humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice (see FIG. 4); and (c) the functionality of the IL-13 pathway in the humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice (see FIG. 4).

**[0201]** As shown in FIG. 3, in which labeled profiles of IL-4R $\alpha$  (CD124) on gated B and T cell populations are shown and the distribution of corresponding unstained cell population is shown in the shaded area, wild-type and humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice express similar amounts of IL-4R $\alpha$  protein on the surface of B (CD19<sup>+</sup>, CD3<sup>-</sup>) and T (CD19<sup>-</sup>, CD3<sup>+</sup>) cells.

**[0202]** As shown in FIG. 4 (left side), wild-type (Il4ra<sup>+/+</sup>) mice respond to mouse, but not human, IL-4, and respond to both mouse and human IL-13. As shown in FIG. 4 (right side), humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice respond to human, but not mouse, IL-4, and respond to both mouse and human IL-13.

**[0203]** This data shows that IL-4, but not IL-13, displays species-specificity in wild-type and humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice.

**[0204]** As shown in FIG. 5, the role of IL-4 as the major factor mediating IgE class switching was supported by the lack of elevated levels of circulating IgE after mouse IL-25 gene delivery in humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice.

**[0205]** The dupilumab monoclonal antibody was investigated in vitro and in vivo.

**[0206]** As shown in FIG. 6, dupilumab prevents human IL-4 induced IgE production in humanized IL-4R $\alpha$  (Il4ra<sup>hu/hu</sup>) mice-derived primary B cell cultures.

**[0207]** As shown in FIG. 7, dupilumab dose dependently reduced IL-25 induced pulmonary pathologies at 10 mg/kg and above (25 mg/kg reduced mucus pathology

## CONCLUSIONS

**[0208]** The results demonstrate the pharmacological activity of dupilumab, a fully human anti-human IL-4R $\alpha$  monoclonal antibody, in a genetically modified mouse model with cytokine-induced inflammation.

**[0209]** Generation of genetically modified mice with human IL-4 and/or IL-4R $\alpha$  gene replacements provides a powerful tool to evaluate function of gene orthologs and in vivo efficacy of antibody candidates with limited cross-species reactivity.

### Example 5

#### House Dust Mite Extract (HDM) Induced Pulmonary Inflammation Model

**[0210]** Chronic airway inflammation in doubly humanized IL-4 and IL-4R $\alpha$  mice is induced by intranasal challenge of house dust mite (HDM) extract (Greer laboratories). In brief, mice were first sensitized by intranasal instillation of HDM suspension (20  $\mu$ l at the concentration of 2.5  $\mu$ g/ml) for 10 days. After a two-week interval of resolution, mice were re-challenged with intranasal HDM application 3 times per week between week 5 and 12. The treatment of dupilumab (anti-IL4R $\alpha$  antibody) was started from the 7th week at the frequency of twice weekly by subcutaneous injections until the end of experiment at week 12. Tissue samples were collected for further analyses. The experimental design is depicted in FIG. 8.

#### Demonstrating the Therapeutic Efficacy of Dupilumab in the HDM Induced Airway Inflammation Model Using Doubly Humanized IL-4 and IL-4R $\alpha$ Mice

**[0211]** Airway disease was induced in doubly humanized IL-4 and IL-4R $\alpha$  (Il4<sup>hu/hu</sup>/Il4ra<sup>hu/hu</sup>) mice using the protocol described above. The histological analysis of lung tissue showed that intranasal HDM instillation caused increased production of mucus in the airway. Treatment of dupilumab reduced the mucus accumulation in the HDM challenged mice. Analysis of the infiltrating cells in bronchoalveolar lavage fluid (BALF) indicates that the eosinophil counts were increased by the HDM instillation and were reduced by the treatment of dupilumab. The total circulating IgE was elevated by the treatment of HDM in the humanized mice, suggesting a competent IL-4 signaling pathway. Use of dupilumab was capable of reducing the level of IgE. In contrast, a comparator molecule, IL13R2 $\alpha$ -Fc, which antagonizes IL-13 only without interfering the IL-4 signal transduction, had comparable activities in reducing mucus accumulation and preventing eosinophil infiltration. Nonetheless, a differential effect was detected in the circulating IgE level between dupilumab and the IL-13 antagonist, IL13R2 $\alpha$ -Fc. Blockade of IL-13 pathway alone was insufficient to reduce the HDM induced IgE level; whereas dupilumab reduced the production of IgE, a main pathogenic mediator of allergy, by blocking both the IL-4 and IL-13 pathways.

**[0212]** In a separate set of experiments, airway disease was induced in doubly humanized IL-4 and IL-4R $\alpha$  (Il4<sup>hu/hu</sup>/Il4ra<sup>hu/hu</sup>) mice using the same protocol described above, except that a different control was used. An isotype control antibody of the same IgG isotype as dupilumab was used in these experiments. The experimental design is depicted in FIG. 9. mRNA was purified from total RNA using Dynabeads mRNA kit (Life Tech) and strand specific RNA-Seq library

was prepared from mRNA using Scriptseq RNA Library Prep kit (Illumina). The library was sequenced using HiSeq 2000 (Illumina) at read length of 33 bp and gene expression levels were extracted from the raw reads using Clcbio (Qiagen) RNA-Seq workflow. Differentially expressed genes between experimental groups were identified using Student's t-test ( $p < 0.05$ , fold change 1.5). An expression cluster of these genes was generated using the Pearson correlation clustering algorithm from GeneSpring GX7.3. HDM was found to induce alteration of pulmonary gene expression in the doubly humanized IL-4 and IL-4R $\alpha$  mice and such alteration was blocked by dupilumab. Serum samples were collected from euthanized mice at the end of the treatment period. The serum murine IgE levels were measured using a commercial ELISA kit (R & D systems). Statistical analysis was performed using ordinary one-way ANOVA method.

#### Example 6

##### Antigen Induced Cutaneous Inflammation Model

**[0213]** Chronic skin inflammation in doubly humanized IL-4 and IL-4R $\alpha$  mice can be induced by the following procedure. The back hair of humanized mice is shaved with electric clipper and then stripped with adhesive tape to create minor injuries and break skin barrier. A gauze patch soaked with a solution of allergen (such as ovalbumin plus bacterial toxin or house dust mite extract) is attached to the skin for one week followed by two weeks of resolution period. The procedure is repeated three times for a total of 7 weeks to induce atopic dermatitis like skin lesions. The treated mice will have increased IgE levels, pruritis, thickening of the epidermis, typical symptoms of atopic dermatitis.

#### Example 7

##### Characterizing PK Profiles of Anti-Human IL-4R $\alpha$ Antibodies in Mice Expressing Humanized IL-4R $\alpha$

**[0214]** This Example describes experiments conducted to evaluate the PK profiles of REGN 668 (human monoclonal antibody directed to human IL-4R $\alpha$ , also known as "dupilumab") and control antibody REGN646 (monkey surrogate, anti-mfIL-4R non-binding control antibody).

**[0215]** The mice used in these experiments were MAID 1444 (homozygous for humanized IL-4R $\alpha$ , or "IL-4R $\alpha$  HumIn", in which the IL-4R $\alpha$  ectodomain is human and the transmembrane and cytoplasmic regions are mouse) and strain-matched (75% C57BL/6/25%129Sv) wild-type ("WT") mice of 20-23 weeks. The study group included a total of 40 mice, male and female, with a cohort size per drug/per dose of 5 homozygous and 5 strain-matched WT. The antibodies (in PBS buffer) were given to mice via subcutaneous injection at 10 mg/kg. Blood samples were taken for analysis on the day of the injection (time point "0" or day 0), at 6 hr post injection, and on day 1, day 3, day 7, day 10, day 14, day 21, and day 30, respectively.

**[0216]** The circulating drug (i.e., REGN668 or REGN646) levels were determined by total human antibody analysis using an ELISA immunoassay. Briefly, a goat anti-human IgG polyclonal antibody (Jackson ImmunoResearch, #109-005-098) was coated onto 96-well plates to capture the tested human antibodies in the sera, and then plate-bound antibodies were detected using a goat anti-human IgG polyclonal antibody conjugated with horseradish peroxidase (Jackson ImmunoResearch, #109-035-098) and TMB substrate (BD

Pharmingen). The serum samples were in six-dose serial dilutions per sample ranging from 1:100-1:243,000 and reference standards of the respective antibodies were in 12-dose serial dilutions. Drug antibody concentrations in the sera were calculated based on the reference standard curve generated using Graphpad Prism software.

**[0217]** The half-life of REGN 668 was found to be shortened in IL-4R $\alpha$  HumIn mice as compared to wild-type mice with only mouse IL-4R $\alpha$  protein. This difference in PK profiles could be explained by the target mediated interaction and clearance between monoclonal antibodies and human IL-4R $\alpha$  receptor. Therefore, mice expressing human or humanized IL-4R $\alpha$  provide suitable simulation to characterize the PK properties of anti-human IL-4R $\alpha$  antibodies (e.g., dupilumab) in a preclinical mouse model.

Uses for Humanized IL-4 and/or IL-4R $\alpha$  Mice

**[0218]** Humanized IL-4 and/or IL-4R $\alpha$  are useful to evaluate the pharmacodynamics (PD) of human-specific IL-4 and/or IL-4R $\alpha$  antagonists, e.g., neutralizing anti-IL-4 and/or anti-IL-4R $\alpha$  antibodies, e.g., dupilumab.

**[0219]** Pharmacokinetics (PK) and PD assays in humanized IL-4 and/or IL-4R $\alpha$  mice are performed according to standard procedures known in the art.

**[0220]** Humanized IL-4 and/or IL-4R $\alpha$  mice are useful to test the in vivo therapeutic efficacy of human-specific IL-4 and/or IL-4R $\alpha$  antagonists, e.g., neutralizing anti-IL-4 and/or IL-4R $\alpha$  antibodies, e.g., dupilumab, in a variety of disease models known in the art, e.g., as shown hereinabove.

#### Example 8

##### Replacement of the Endogenous Mouse IL-33 Gene with a Human IL-33 Gene

**[0221]** The mouse IL-33 gene (NCBI Gene ID: 77125, Primary source: MGI:1924375; RefSeq transcript: NM\_001164724.1; UniProt ID: Q8BVZ5; Genomic assembly: NCBI37/mm9; Location: chr19:29,999,604-30,035,205+ strand) has 8 exons and encodes a protein of 266 amino acids (GenBank Accession No. NP\_001158196.1).

**[0222]** The human IL-33 gene (NCBI Gene ID: 90865, Primary source: HGNC:16028; RefSeq transcript: NM\_033439.3; UniProt ID: 095760; Genomic assembly: GRCh37/hg19; Location: chr9:6,215,149-6,257,983+ strand) also has 8 exons and encodes a protein of 270 amino acids (GenBank Accession No. NP\_254274.1).

**[0223]** A 16333 bp human genomic segment containing exon 2 starting from the ATG initiation codon through exon 8 (including the 3' untranslated region) of the human IL-33 gene replaced 9381 bp of the mouse IL-33 gene locus spanning exon 2 starting from the ATG initiation codon through the coding portion of exon 8 including the stop codon. See FIG. 10A.

**[0224]** A targeting construct for replacing the mouse IL-33 gene with a human IL-33 genomic segment in a single targeting step was constructed using VelociGene® genetic engineering technology (see Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, Nature Biotech, 21(6): 652-659), similar to the procedure described in Example 1 above for replacing the mouse IL-4 gene with a human IL-4 genomic segment, except that mouse and human IL-33 DNA were obtained from bacterial artificial chromosome (BAC)

clones bMQ-350I18 and CTD-3015M15, respectively, and that the targeting vector contained a loxP neomycin selection cassette (FIG. 10B).

**[0225]** Correctly targeted ES cell clones (MAID 7060) were identified by a loss-of-native-allele (LONA) assay (Valenzuela et al. 2003) in which the number of copies of the native, unmodified IL-33 gene were determined by two TaqMan™ quantitative polymerase chain reactions (qPCRs) specific for sequences in the mouse IL-33 gene that were targeted for deletion. The qPCR assays comprised the following primer-probe sets (written 5' to 3'):

upstream ("mTU"):  
forward primer,  
(SEQ ID NO: 31)  
TTGGACTAGTAACAAGAAGGGTAGCA;  
reverse primer,  
(SEQ ID NO: 32)  
CCTTTCCCATCACCCCTCTAACTT;  
probe (MGB),  
(SEQ ID NO: 33)  
AGCTCTGGTGACAGA;  
downstream ("mTD"):  
forward primer,  
(SEQ ID NO: 34)  
TCTCTGCCAAGCTGCTTATCC;  
reverse primer,  
(SEQ ID NO: 35)  
GGCTGCATGGAAGAGGTGAA;  
probe (MGB),  
(SEQ ID NO: 36)  
CTCTCCACAAATCG.

**[0226]** Confirmation that the human IL-33 gene sequence replaced the mouse IL-33 gene sequence in the humanized allele was confirmed by a TaqMan™ qPCR assay that comprises the following primer-probe sets (written 5' to 3'):

upstream ("hTU")  
forward primer,  
(SEQ ID NO: 37)  
CAGGCAGGAATAGCTGAGATAATCT;  
reverse primer,  
(SEQ ID NO: 38)  
TGTGGAGCAAAAAGTGGTTGAT;  
probe (MGB),  
(SEQ ID NO: 39)  
CCTGTGAATAGTGATAAAC;  
downstream ("hTD"):  
forward primer,  
(SEQ ID NO: 40)  
CAGTTCCAACGATAGGCTCAA;  
reverse primer,  
(SEQ ID NO: 41)  
ATAATCTGTGAAGCATCTGGTCTTC;  
probe (MGB),  
(SEQ ID NO: 42)  
CTAGAGCTGCTAGTAAAA.

**[0227]** The upstream junction of the murine IL-33 locus and the sequence containing the human IL-33 gene (shown as "I" in FIG. 10B) is designed to be within 5'-ATAGCCAAGG TTGCTTCTGA TGACTTCAGG TCCATATAGT TGGAT-

TAATG TTATATTTCA ATCCCACAGA AACCTGAAAA  
ATGAAGCCTA AAATGAAGTA TTCAACCAAC  
AAAATTTCCA CAGCAAAGTG GAAGAACACA  
GCAAGCAAAG CCTTGTGTTT-3' (SEQ ID NO: 43),  
wherein the human IL-33 sequence is italicized and the human start codon ATG is underlined. The downstream junction of the sequence containing the human IL-33 genomic sequence and the loxP neomycin selection cassette (shown as "II" in FIG. 10B) is designed to be within

(SEQ ID NO: 44)  
5'- TTTATATTAT TGAATAAAGT ATATTTTCCA AATGTATGTG  
AGACTATAAT GATTTTATCA TATGATGACT CAATATTCTG/  
CTCGAGATAA CTTCGTATAA TGTATGCTAT ACGAAGTTAT  
ATGCATGGCC TCCGCGCCGG GTTTTGGCGC CTCCCAGGGG -3',

wherein the human IL-33 sequence is italicized and the junction is indicated by the "/" symbol, and the lox P site is underlined. The downstream junction of the sequence of the loxP neo selection cassette and the murine IL-33 locus (shown as "III" in FIG. 10C) is designed to be within

(SEQ ID NO: 45)  
5'- AGCCCCTAGA TAACCTCGTA TAATGTATGC TATACGAAGT  
TATGCTAGTA ACTATAACGG TCCTAAGGTA GCGAGCTAGC/  
CGCCTGTGCG TTCTGGGTTG AATGACTTAA TGCTTCCAAC  
TGAAGAAAAG GTAACAGAGA GAAAGAAAAGC CATTCTTGCC -3',

wherein the junction is shown by the "/" symbol, and the loxP site is underlined.

**[0228]** Correctly targeted ES cells (MAID 7060) were further electroporated with a transient Cre-expressing vector to remove the drug selection cassette and obtain ES cell clones without drug cassette (MAID 7061). The upstream junction in these MAID 7061 ES cells (shown as "I" in FIG. 18C) is the same as in MAID 7060 ES cells. The downstream junction (shown as "II" in FIG. 18C) is designed to be within

(SEQ ID NO: 46)  
5'- TTTATATTAT TGAATAAAGT ATATTTTCCA AATGTATGTG  
AGACTATAAT GATTTTATCA TATGATGACT CAATATTCTG/  
CTCGAGATAA CTTCGTATAA TGTATGCTAT ACGAAGTTAT  
GCTAGTAACT ATACCGTCC TAAGGTAGCG AGCTAGC/  
CGCCTGTGCG TTCTGGGTTG AATGACTTAA TGCTTCCAAC  
TGAAGAAAAG GTAACAGAGA GAAAGAAAAGC CATTCTTGCC -3',

wherein the 3' human IL-33 sequence is italicized before the first "/" symbol, and the mouse IL-33 3' sequence is italicized after the second "/" symbol, and the loxP site is underlined.

**[0229]** Correctly targeted ES cells (MAID 7060 or MAID 7061) were introduced into an 8-cell stage SW mouse embryo by the VelociMouse® method (see, U.S. Pat. Nos. 7,294,754, 7,576,259, 7,659,442, and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses, Nature Biotech. 25(1):91-99). VelociMice® (F0 mice fully derived from the donor ES cell) bearing the humanized

IL-33 gene were identified by genotyping for loss of mouse allele and gain of human allele using a modification of allele assay (see, Valenzuela et al. (2003)). The same LONA assay was used to assay DNA purified from tail biopsies for mice derived from the targeted ES cells to determine their IL-33

genotypes and confirm that the humanized IL-33 allele had transmitted through the germline. Two pups heterozygous for the replacement were bred to generate a mouse that is homozygous for the replacement of the endogenous mouse IL-33 gene by the human IL-33 gene.

---

 SEQUENCE LISTING
 

---

<160> NUMBER OF SEQ ID NOS: 46

<210> SEQ ID NO 1  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 1

catgcacgga gatggatgtg 20

<210> SEQ ID NO 2  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 2

gacccctcag gtccacttac c 21

<210> SEQ ID NO 3  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
 fluorescent probe  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (19)..(19)  
 <223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
 black hole quencher type

<400> SEQUENCE: 3

aacgtcctca cagcaacga 19

<210> SEQ ID NO 4  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 4

gtgcccaggt gtgctcatg 19

<210> SEQ ID NO 5  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 5

---

-continued

---

cgctgcctc ctcactttat c 21

<210> SEQ ID NO 6  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (19)..(19)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
black hole quencher type

<400> SEQUENCE: 6

atctgcttca ccatccact 19

<210> SEQ ID NO 7  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 7

gcttggacca agactctgt 19

<210> SEQ ID NO 8  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 8

accgtgggac ggcttcttac 20

<210> SEQ ID NO 9  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (19)..(19)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
black hole quencher type

<400> SEQUENCE: 9

caccgagttg accgtaacag acatc 25

<210> SEQ ID NO 10  
<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence



-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 10

tgcggccgat cttagcc 17

<210> SEQ ID NO 11  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 11

ttgaccgatt ccttgccg 18

<210> SEQ ID NO 12  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
 fluorescent probe  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (21)..(21)  
 <223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
 black hole quencher type

<400> SEQUENCE: 12

acgagcgggt tcggcccatt c 21

<210> SEQ ID NO 13  
 <211> LENGTH: 458  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Nucleic acid sequence containing the upstream  
 junction of Murine IL4 Locus and Human IL4 sequence

<400> SEQUENCE: 13

tgctgattgg cccagaataa ctgacaatct ggtgtaataa aatthtccaa tgtaaaactca 60

ttttcccttg gtttcagcaa cttaactct atatatagag agacctctgc cagcattgca 120

ttggttagcat ctcttgataa acttaattgt ctctcgtcac tgacggcaca gagctattga 180

tgggtctcac ctcccaactg ctccccctc tgttcttct gctagcatgt gccggcaact 240

ttgtccacgg acacaagtgc gatatcacct tacaggagat catcaaaact ttgaacagcc 300

tcacagagca gaaggtgagt acctatctgg caccatctct ccagatgttc tggatgatgct 360

ctcagtattt ctaggcatga aaacgttaac agctgctaga gaagttggaa ctggtgggtg 420

gtggcagtc agggcacaca gcgaggcttc tccccctgc 458

<210> SEQ ID NO 14  
 <211> LENGTH: 565  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Nucleic acid sequence containing the downstream  
 junction of Human IL4 sequence and floxed hygo selection cassett

-continued

&lt;400&gt; SEQUENCE: 14

```

tgtttatttt gcagaattcc tgcctctgga aggaagccaa ccagagtacg ttggaaaact    60
tcttggaag gctaaagacg atcatgagag agaaatattc aaagtgttcg agctgaatat    120
tttaatttat gagtttttga tagctttatt ttttaagtat ttatatattt ataactcacc    180
ataaaataaa gtatatatag aatctaacag caatggcatt taatgtattg gctatgttta    240
cttgacaaat gaaattatgg tttgcaactt ttagggaat caatttagtt taccaagaga    300
ctataaatgc tatgggagca aaacaggaaa gaccacttcc ccctcgaggg gttccctctc    360
gagttagga cataacacac aagataatta aagaacacaa ggccatacaa gatgcgggccg    420
caccgggata acttcgtata aggtatoccta tacgaagtta tatgcatggc ctccgcccgg    480
ggttttggcg cctcccggcg gcgccccct cctcacggcg agcgctgcca cgtcagacga    540
agggcgcagc gagcgtcctg atcct                                          565

```

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 438

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Nucleic acid sequence containing the downstream junction of floxed hygo selection casset and Murine IL-4 locus

&lt;400&gt; SEQUENCE: 15

```

tgccaagtcc taattccacc agacctcgac ctgcagccgg cgcgccataa ctctgtataa    60
ggtatcctat acgaagtat ctccagagga gttcccaccc ttctcaagag cataatgcgc    120
agatcattaa gggacagatg caggctgggg agacgggtca gcagttagga gtacctgttg    180
ctcttccaga ggaccagggt tcaattcccg gcaactcacat agcagcttaa aacaataact    240
caagttctgg gggagctgat gctctcctct gccctcctgt ggaggtacac agaccacatg    300
cctgtaggca agacaccac acacataaaa acaaaataaa ataaggatag aaaggccagg    360
gggatgaatc cagaggtaga agaaaactta ttccctggaa ttgtcctctg actcccctcc    420
caaaacctct aacacgca                                          438

```

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 20

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide

&lt;400&gt; SEQUENCE: 16

```

ccgctggcat gtgtattgtg                                          20

```

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide

&lt;400&gt; SEQUENCE: 17

```

tgagtgtggg accctcaaga g                                          21

```

&lt;210&gt; SEQ ID NO 18

---

-continued

---

<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (24)..(24)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
black hole quencher type

<400> SEQUENCE: 18

tgaccaagc cctacatggc cact 24

<210> SEQ ID NO 19  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 19

tgaggagagc tcacgggaat c 21

<210> SEQ ID NO 20  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 20

accatctcc tgcgtttctg 20

<210> SEQ ID NO 21  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (25)..(25)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
black hole quencher type

<400> SEQUENCE: 21

ttgacacgcc agctacactg ctcca 25

<210> SEQ ID NO 22  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 22

---

-continued

---

acctgcgtct ccgactacat g 21

<210> SEQ ID NO 23  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 23

gagctcggtg ctgcaattg 19

<210> SEQ ID NO 24  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (25)..(25)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the  
black hole quencher type

<400> SEQUENCE: 24

tgggaccatt catcttccac tcgca 25

<210> SEQ ID NO 25  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 25

ggtggagagg ctattcggc 19

<210> SEQ ID NO 26  
<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 26

gaacacggcg gcatcag 17

<210> SEQ ID NO 27  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: Labeled with FAM: 5-carboxyfluorescein  
fluorescent probe  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (23)..(23)  
<223> OTHER INFORMATION: Labeled with BHQ: fluorescence quencher of the

-continued

---

black hole quencher type

<400> SEQUENCE: 27

tgggcacaac agacaatcgg ctg 23

<210> SEQ ID NO 28  
 <211> LENGTH: 243  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Portion of Murine IL-4Ralpha locus and sequence containing IL-4Ralpha gene

<400> SEQUENCE: 28

tgggggaggg aggccatgac acaaatgaaa tggaccccgc tgaccaggga tcagcatctg 60  
 cccactcttc tttctgcagg caccttttgt gtccccaatg ggggtgcttt gctctgggct 120  
 cctgttcctc gtgagctgcc tggctctgct gcaggtggca agctctggta agtcaccact 180  
 tctcaatcat tcatttgggt gctattaatg gcgtgccagg gtccctgcagt atgtcacctg 240  
 gcc 243

<210> SEQ ID NO 29  
 <211> LENGTH: 243  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Nucleic acid sequence containing the downstream junction of Human IL-4Ralpha sequence and floxed neo selection cassett

<400> SEQUENCE: 29

gtcagatcgt ggagggcttc ggacgagggt cctgacctg ggtctccagt cctgggaagt 60  
 ggagcccagg ctgtaccatg gctgacctca gctcatggct cccgggctcg ataactataa 120  
 cggtcctaag gtatcgactc gagataactt cgtataatgt atgctatacg aagttatatg 180  
 catggcctcc gcgcccgggt ttggcgctc ccgcccggcg cccctcctc acggcgagcg 240  
 ctg 243

<210> SEQ ID NO 30  
 <211> LENGTH: 390  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Portion of the sequence of the floxed neo selection cassett and the Murine IL-4Ralpha locus

<400> SEQUENCE: 30

tattgttttg ccaagttcta attccatcag acctcgacct gcagccccta gataacttcg 60  
 tataatgtat gctatacga gttatcctag gttggagctc tctgtagcca ggtaaccaag 120  
 ggtcccaggg gaacccccag tgtggacgcg gactgcacat gacacagggc ggcctcccca 180  
 ttcattgactg tttttctcct tgcagacttc cagctgcccc tgatacagcg ccttccactg 240  
 ggggtcacca tctcctgect ctgcatcccg ttgttttgcc tgttctgtta cttcagcatt 300  
 accaagttag ttcctgcttt ggctgggtgc tctggctggc ccttcagcag tgctctcaga 360  
 ggtcacagtc attgtgctgg ctgagaaaag 390

<210> SEQ ID NO 31

---

-continued

<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 31

ttggactagt aacaagaagg gtagca 26

<210> SEQ ID NO 32  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 32

cctttcccat cacctctaa ctt 23

<210> SEQ ID NO 33  
<211> LENGTH: 16  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 33

agctctggtg gacaga 16

<210> SEQ ID NO 34  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 34

tctctgccaa gctgcttacc c 21

<210> SEQ ID NO 35  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 35

ggctgcatgg aagagtgaa 20

<210> SEQ ID NO 36  
<211> LENGTH: 14  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 36

ctctccacaa atcg 14

<210> SEQ ID NO 37  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 37

caggcaggaa tagctgagat aatct 25

<210> SEQ ID NO 38  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 38

tgtggagcaa aaagtggttg at 22

<210> SEQ ID NO 39  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 39

cctgtgaata gtgataaac 19

<210> SEQ ID NO 40  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 40

cagttccaaa cgataggctc aa 22

<210> SEQ ID NO 41  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 41

ataattctgt gaagcatctg gtcttc 26

<210> SEQ ID NO 42  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 42

ctagagctgc tagtaaaa 18

<210> SEQ ID NO 43  
<211> LENGTH: 160  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 43

-continued

---

```

atagccaagg ttgcttctga tgacttcagg tccatatagt tggattaatg ttatatttca    60
atccccacaga aacctgaaaa atgaagccta aatgaagta ttcaaccaac aaaatttcca    120
cagcaaaagtg gaagaacaca gcaagcaaag ccttgtgttt                            160

```

```

<210> SEQ ID NO 44
<211> LENGTH: 160
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

```

```

<400> SEQUENCE: 44

```

```

tttatattat tgaataaagt atattttcca aatgtatgtg agactataat gattttatca    60
tatgatgact caatattctg ctcgagataa ctctgtataa tgtatgctat acgaagttaa    120
atgcatggcc tccgcgcggg gttttggcgc ctcccgcggg                            160

```

```

<210> SEQ ID NO 45
<211> LENGTH: 160
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

```

```

<400> SEQUENCE: 45

```

```

agcccctaga taacttcgta taatgtatgc tatacgaagt tatgctagta actataacgg    60
tcctaaggta gcgagctagc cgctgtgctg ttctgggttg aatgacttaa tgettccaac    120
tgaagaaagg gtaacagaga gaaagaaagc cattcttggc                            160

```

```

<210> SEQ ID NO 46
<211> LENGTH: 237
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

```

```

<400> SEQUENCE: 46

```

```

tttatattat tgaataaagt atattttcca aatgtatgtg agactataat gattttatca    60
tatgatgact caatattctg ctcgagataa ctctgtataa tgtatgctat acgaagttaa    120
gctagtaact ataacggctc taaggtagcg agctagccgc ctgtgcttgc tgggttgaat    180
gacttaatgc ttccaactga agaaagggta acagagagaa agaaagccat tcttggc      237

```

---

1. A rodent, comprising a humanization of an endogenous rodent IL-4 gene, wherein the humanization comprises a replacement at the endogenous rodent IL-4 locus of a rodent nucleic acid comprising an exon of an IL-4 gene with a nucleic acid sequence comprising at least one exon of a human IL-4 gene to form a modified IL-4 gene, wherein expression of the modified IL-4 gene is under control of rodent regulatory elements at the endogenous rodent IL-4 locus.

2. The rodent of claim 1, wherein the rodent is a mouse or a rat.

3. The rodent of claim 1, wherein the modified IL-4 gene encodes a human or humanized IL-4 protein, and comprises exon 1 starting from the ATG initiation codon through exon 4 of the human IL-4 gene.

4. The rodent of claim 1, wherein the rodent is a mouse that is incapable of expressing a mouse IL-4 protein.

5. The rodent of claim 1, wherein the rodent is a mouse that expresses a mouse IL-4R $\alpha$  protein encoded by an endogenous mouse IL-4R $\alpha$  gene.

6. The rodent of claim 1, wherein the rodent is a mouse that expresses a human or humanized IL-4R $\alpha$  protein.

7. The rodent of claim 6, wherein the humanized IL-4R $\alpha$  protein comprises the ectodomain of the human IL-4R $\alpha$  protein.

8. The rodent of claim 7, wherein the humanized IL-4R $\alpha$  protein comprises the transmembrane domain and cytoplasmic domain of the mouse IL-4R $\alpha$  protein.

9. The rodent of claim 6, wherein the mouse comprises a replacement at an endogenous mouse IL-4R $\alpha$  locus of a mouse nucleic acid comprising an exon of a mouse IL-4R $\alpha$



gene with a nucleic acid sequence encoding at least one exon of a human IL-4R $\alpha$  gene to form a modified IL-4R $\alpha$  gene, wherein expression of the modified IL-4R $\alpha$  gene is under control of mouse regulatory elements at the endogenous mouse IL-4R $\alpha$  locus.

**10.** The rodent of claim **1**, wherein the rodent is a mouse, and wherein a contiguous genomic fragment of mouse IL-4 sequence comprising exon 1 starting from the ATG initiation codon through exon 4 of mouse IL-4 is replaced with a contiguous genomic fragment of human IL-4 sequence comprising exon 1 starting from the ATG initiation codon through exon 4 of human IL-4.

**11-44.** (canceled)

**45.** A rodent, comprising a humanized IL-4R $\alpha$  gene and a humanized IL-4 gene, wherein the humanized IL-4R $\alpha$  gene results from a replacement of a rodent IL-4R $\alpha$  ectodomain-coding sequence with a human IL-4R $\alpha$  ectodomain-coding sequence, and the humanized IL-4R $\alpha$  gene comprises a rodent IL-4R $\alpha$  transmembrane domain coding sequence and a rodent IL-4R $\alpha$  cytoplasmic domain coding sequence, and wherein the humanized IL-4R $\alpha$  gene is under control of endogenous rodent IL-4R $\alpha$  regulatory elements at the endogenous rodent IL-4R $\alpha$  locus; and wherein the humanized IL-4 gene encodes a human or humanized IL-4 protein, and the humanized IL-4 gene is under control of endogenous rodent IL-4 regulatory elements at the endogenous rodent IL-4 locus.

**46.** The rodent of claim **45**, wherein the rodent is a mouse or a rat.

**47.** The rodent of claim **45**, wherein the rodent is a mouse that is incapable of expressing a mouse IL-4 protein and incapable of expressing a mouse IL-4R $\alpha$  protein.

**48.** The rodent of claim **45**, wherein the rodent regulatory elements or sequences at the endogenous rodent IL-4 locus and/or endogenous rodent IL-4R $\alpha$  locus are from a mouse or a rat.

**49.** The rodent of claim **45**, wherein the rodent regulatory elements or sequences are endogenous rodent regulatory elements or sequences at the rodent IL-4 locus and/or rodent IL-4R $\alpha$  locus.

**50.** A method of screening for human-specific IL-4 or IL-4R $\alpha$  antagonist, comprising:

providing a doubly humanized IL-4 and IL-4R $\alpha$  mouse, inducing pulmonary inflammation in said mouse, administering an agent to said mouse, determining whether pulmonary inflammation is reduced by the agent, and identifying the agent as a human specific IL-4 or IL-4R $\alpha$  antagonist based on its ability to reduce pulmonary inflammation.

**51.** The method of claim **50**, wherein the extent of pulmonary inflammation is determined by measuring mucus accumulation in the airway, eosinophilic infiltrating cells in bronchoalveolar lavage fluid, and/or total circulating IgE.

**52.** A method of screening for human-specific IL-4 or IL-4R $\alpha$  antagonist, comprising:

providing a doubly humanized IL-4 and IL-4R $\alpha$  mouse, inducing skin inflammation in said mouse, administering an agent to said mouse, determining whether skin inflammation is reduced by the agent, and identifying the agent as a human specific IL-4 or IL-4R $\alpha$  antagonist based on its ability to reduce skin inflammation.

**53.** The method of claim **50**, wherein the humanized IL-4R $\alpha$  gene results from a replacement of a murine IL-4R $\alpha$  ectodomain-coding sequence with a human IL-4R $\alpha$  ectodomain-coding sequence, and the humanized IL-4R $\alpha$  gene is under control of endogenous murine IL-4R $\alpha$  regulatory elements at the endogenous murine IL-4R $\alpha$  locus; and wherein the humanized IL-4 gene encodes a human or humanized IL-4 protein, and the humanized IL-4 gene is under control of endogenous rodent IL-4 regulatory elements at the endogenous rodent IL-4 locus.

**54.** The method of claim **52**, wherein the humanized IL-4R $\alpha$  gene results from a replacement of a murine IL-4R $\alpha$  ectodomain-coding sequence with a human IL-4R $\alpha$  ectodomain-coding sequence, and the humanized IL-4R $\alpha$  gene is under control of endogenous murine IL-4R $\alpha$  regulatory elements at the endogenous murine IL-4R $\alpha$  locus; and wherein the humanized IL-4 gene encodes a human or humanized IL-4 protein, and the humanized IL-4 gene is under control of endogenous rodent IL-4 regulatory elements at the endogenous rodent IL-4 locus.

\* \* \* \* \*