




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 31/40, C07D 209/56</p>	<p>A1</p>	<p>(11) International Publication Number: WO 93/00087 (43) International Publication Date: 7 January 1993 (07.01.93)</p>
<p>(21) International Application Number: PCT/US92/04624 (22) International Filing Date: 5 June 1992 (05.06.92) (30) Priority data: 723,394 28 June 1991 (28.06.91) US (71) Applicant: CYTOPHARM, INC. [US/US]; 1330 O'Brien Drive, Menlo Park, CA 94025 (US). (72) Inventors: VOGEL, Emanuel ; Raschdorffstrasse 7, D-5000 Köln 41 (DE). BENNINGHAUS, Thomas ; Schwalbacher Strasse 12, D-5000 Köln 51 (DE). RICHERT, Clemens ; Grüner Weg 36, D-4409 Havixbeck (DE). MULLER, Martin ; Lenaustrasse 11, D-5000 Köln 30 (DE). CROSS, Alexander, Dennis ; 286 Park Lane, Atherton, CA 94025 (US).</p>		<p>(74) Agents: WALKER, William, B. et al.; Oblon, Spivak, McClelland, Maier & Neustadt, Fourth Floor, 1755 South Jefferson Davis Highway, Arlington, VA 22202 (US). (81) Designated States: AU, BB, BG, BR, CA, CS, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, PL, RO, RU, SD, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: PORPHYCENE COMPOUNDS FOR PHOTODYNAMIC THERAPY



(57) Abstract

A porphycene having structure (I) wherein each R¹ is, independently, (a) -(CH₂)_n-X, where n = 1-4, X is OR² and R² is C₁₋₆ alkyl, aralkyl or aryl; CN; OH; OSO₂R²; NH₂; NHR²; NR²₂; SH; SR²; S(O)₁₋₂R²; COOH; CO₂R²; C(O)NH₂; C(O)NHR²; C(O)NR²₂; halogen; or CHO; (b) -(CH₂)_mCH=CH₂ where m is 0-2; or (c) -(CH₂)_n-O-G where G is a mono- or oligosaccharide; (d) -(CH₂)_{2n}-X, where X is an amino acid, oligopeptide covalently bonded by an ether-, ester- or amine-bond or -Y-(CH₂)_n-porphycene² (porphycene² being a compound of the same structure and Y is a direct bond; -O-; or -CH=CH₂); or (e) where one, two or three of the substituents R¹ are C₁₋₆ alkyl or aryl and the remaining substituents are as above under (a)-(d), and salts and metal complexes thereof. The porphycene compounds and pharmaceutical compositions containing the compounds are useful in photodynamic therapy treatment of tumors and psoriasis.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NI	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

Porphycene Compounds for Photodynamic Therapy

Technical Field

The invention relates to novel porphycene compounds
5 and pharmaceutical compositions containing these
compounds which are useful for therapeutic treatment.

Background Art

During the past few years there has developed a
widespread recognition that modern, though sophisticated,
10 cancer diagnosis and treatments have served neither to
reduce overall the number of cases of reported cancers in
the U.S.A. nor, save the notable cases, the death rate.
This is a disheartening result for the billions of
dollars invested in conquering the disease. Moreover,
15 surgery, radiotherapy and chemotherapy are all associated
with major debilitating side effects such as trauma,
severe immunosuppression or toxicity which are not easily
surmounted by patients already compromised by ill-health.

Early work in the 1970's, followed by rapidly
20 expanding studies in the 1980's, has shown that
photodynamic therapy (PDT) offers a viable, less toxic
and generally less painful avenue to treatment of cancer.
Not all cancers are candidates for PDT. However,
neoplasms of hollow organs and skin, including multifocal
25 carcinoma in situ, sometimes inoperable, and with no good
track record for treatment by established therapeutic
procedures, appear to be targets for PDT.

In photodynamic therapy, porphyrinoid dyes are
administered to a patient and localize in neoplastic
30 tissues (Lipson et al, J. Thoracic Cardiovascular

Surgery, 1961, 42:623-629). Irradiation of the porphyrinoid dye with light at a wavelength which corresponds to an absorption band of the dye results in destruction of the neoplastic tissue. See also Kessel, 5 D., "Methods in Porphyrin Photosensitization", Plenum Press, New York, 1985; Gomer, C. J., "Photodynamic Therapy", Pergamon Press, Oxford, 1987 and Doiron, D. R. and Gomer, C. J., "Porphyrin Localization and Treatment of Tumors", Liss, New York, 1984. The use of a fiber 10 optic laser light source is described in U.S. 4,957,481.

Dougherty et al (Cancer Res., 1978, 38:2628; Photochem. Photobiol, 1987, 45:879) pioneered the field with infusion of photoactivatable dyes, followed by appropriate long wavelength radiation of the tumors (600+ 15 nm) to generate a lethal shortlived species of oxygen which destroyed the neoplastic cells. Early experiments utilized a mixture termed hematoporphyrin derivative (HPD). See also Lipson et al, J.N.C.I., 1961, 26:1; Dougherty et al, J.N.C.I., 1975, 55:115; Diamond et al, 20 Lancet, 1972(II), 1175; D. Dolphin, "The Porphyrins", vol. I, Academic Press, New York, 1978; and D. Kessel, Photochem. PhotoBiol., 1984, 39:851. The deficiencies of HPD, especially prolonged phototoxicity caused by 25 retained HPD components in human skin led to its displacement by a purified fraction termed dihematoporphyrin ether (DHE) which, although yielding improvements over HPD, nevertheless still suffered certain practical limitations. Relatively weak absorption in the wavelength range above 600 nm, 30 retention in dermal cells (potentially leading to phototoxicity), only modest or low selectivity for tumor cells versus other cell types in vital organs, the inability to use available, modern, inexpensive diode lasers, and uncertain chemical constitution of the

-3-

mixtures are all known negative features of DHE and HPD. The great majority of the earlier PDT agents studied have been derived from natural sources (porphyrins, chlorins, purpurins, etc.) or from known chemicals originating in the dyestuffs industry (e.g., cyanine dyes). For more recent PDT agents derived from natural sources see U.S. 4,961,920 and U.S. 4,861,876.

In animal and cell culture experiments one observes, following PDT, depending on the incubation time, damage to the vasculature, cell membranes, mitochondria and specific enzymes. When absorbed in tumor cells, an increased selectivity can be obtained by injecting the porphyrinoid sensitizers enclosed in liposomes (Ricchelli and Jori, Photochem. Photobiol., 1986, 44:151). Porphyrinoid dyes can be transported in the blood with the aid of lipoproteins such as low-density lipoprotein (Jori et al, Cancer Lett., 1984, 24:291).

PDT has been used to treat bladder, bronchial, bone marrow and skin tumors (Dougherty, Photochem. Photobiol., 1987, 45:879, Sieber et al, Leukemia Res., 1987, 11:43) as well as severe psoriasis (Diezel et al, Dermatol. Monatsschr., 1980, 166:793; Emtenstam et al, Lancet, 1989 (I), 1231). Treatment of viruses in transfused blood has also been reported (Matthews et al, Transfusion, 1988, 28:81; Sieber et al, Semin. Hematol., 1989, 26:35).

As the deficiencies of earlier PDT agents have become apparent, it also becomes possible to define activity parameters for improved chemically pure photoactivatable dyes for PDT therapy, available by chemical synthesis. Moreover, the products of synthesis lend themselves more readily to further chemical structural manipulation than do the naturally occurring

starting materials which can be expensive and bear chemically sensitive constituents. The synthesis of novel porphycene macrocycles embracing four pyrrole rings has been described by Vogel and coworkers. Alkylated porphycenes have also been prepared (R = Me, Et, n-Pr, n-octyl, phenyl) and the photochemical properties determined. The suitability of these compounds for PDT was noted and confirmed in animal studies (Guardiano et al, Cancer Letters, 1989, 44, 1).

10 Synthetic efforts have focused on porphyrinoid compounds which are highly absorptive in the longer wavelength range of about 600-1200 nm, where the transparency of tissue is higher. Compounds such as purpurines (Morgan et al, J. Org. Chem., 1986, 51:1347; 15 Morgan et al, Cancer Res., 1987, 47:496; Morgan et al, J. Med. Chem., 1989, 32:904; Hooper et al, Photochem. Photobiol., 1988, 48:579), naphthocyanin silicon complexes (Firey et al, J. Am. Chem. Soc., 1988, 110:7626), chlorins (Robert et al, J.N.C.I., 1988, 20 80:330; Kessel, Cancer Res., 1986, 46:2248), bacteriochlorins (Beams et al, Photochem. Photobiol., 1987, 46:639) and substituted phenylporphyrins (Kreimer-Birnbaum, Semin. Hematol., 1989, 26:157) have been prepared and tested *in vivo*. Additional PDT agents are 25 described in EP 276,121.

Pyrrole-containing ring systems larger than porphycene have also been prepared and evaluated as photosensitizers. Sessler et al have prepared and studied texaphyrin (J. Am. Chem. Soc., 1988, 110:5586) 30 and Woodward et al and Johnson et al have prepared and investigated the sapphyrin ring system. Additionally, the platyrin system has been studied by LeGoff (Tetrahedron, Lett., 1978, 4225; J. Org. Chem., 1987,

710) and vinylogous porphyrins have been studied by Franck (Angew. Chem., 1986, 98:1107; Angew. Chem. Int. Ed. Eng., 1986, 25:1100; Angew. Chem., 1988, 100:1203; Angew. Chem. Int. Ed. Eng., 1988, 27:1170).

5 A need continues to exist, therefore, for new compounds for use in PDT therapy, which compounds are easily available, have low intrinsic toxicity, are efficient photosensitizers for singlet oxygen production, have selective uptake in rapidly proliferating cells, are
10 rapidly or at least moderately rapidly degraded and eliminated from the tissues after administration and which are available as chemically pure and stable compounds easily subject to synthetic modification. The compound should penetrate tissue quickly, especially if
15 used for topical application.

Disclosure of the Invention

Accordingly, one object of the present invention is to provide new and effective compounds for use in photodynamic therapy whose properties and characteristics
20 approach the ideal characteristics of PDT dyes listed above.

This and other objects which will become apparent from the following specification have now been achieved with the compounds of the present invention. The present
25 compounds have utility as PDT dyes for use in cancer therapy and dermatological diseases, blood purification (elimination of viruses and bacteria, e.g., CMV, HIV), etc.

Brief Description of the Drawings

A more complete appreciation of the invention and many of the attendant advantages thereof will be readily obtained as the same becomes better understood by
5 reference to the following detailed description when considered in connection with the accompanying drawing, wherein:

Figure 1 illustrates a synthesis of the tetrakis(alkoxyalkyl)porphycene compounds of the present
10 invention, specifically tetrakis(methoxyethyl)porphycene.


Figure 2 illustrates the preparation of mono-, bis-, tris- and tetrakis(bromoalkyl)porphycenatonickel compounds which, following demetallation, provide
compounds of the present invention.

15 Figure 3 illustrates the preparation of mono-, bis-, tris and tetrakis(hydroxyalkyl)porphycene compounds of the present invention.

Figure 4 illustrates the preparation of porphycene carboxylic acid compounds of the present invention.

20 Best Mode for Carrying Out the Invention

The porphycene compounds of the present invention have the structure shown below:



SUBSTITUTE SHEET

-7-

where each R^1 in the 2, 7, 12 and 17-positions of the porphyr- cene structure is, independently of one another, (a) the group $-(CH_2)_n-X$, where $n=1-4$, X is OR^2 and R^2 is C_{1-6} alkyl, aralkyl or aryl; or X is CN ; OH ; $OCOR^2$;
5 OSO_2R^2 ; NH_2 ; NHR^2 ; NR_2^2 ; SH ; SR^2 ; $S(O)_{1-2}R^2$; $COOH$; $C(O)NH_2$,
 $C(O)NHR^2$, $C(O)NR_2^2$, CO_2R^2 , halogen or CHO ; (b) the group
 $-(CH_2)_m-CH=CH_2$ where $m = 0-2$ or (c) the group $-(CH_2)_n-O-G$
where G is a mono- or oligosaccharide covalently bonded
to the porphycene or (d) the group $-(CH_2)_n-X$ where X is an
10 amino acid, oligopeptide covalently bonded by an ether-,
ester- or amine-bond or $Y(CH_2)_n$ -porphycene² (porphycene²
being a compound of the same structure and Y is a direct
bond; $-O-$; or $-CH=CH-$) or (e) where one, two or three of
the substituents R^1 are C_{1-6} alkyl or aryl and the
15 remaining substituents are as described above under (a)-
(d), salts and metal complexes thereof. For all four
substituents $R^1 =$ alkyl, see U.S. 4,913,907.

Suitable aryl groups include C_{6-20} carbocyclic aryl
groups, optionally substituted with one or more C_{1-6} alkyl
20 groups. Examples include phenyl, naphthyl, indanyl etc.
Suitable aralkyl groups are the aryl groups defined above
bonded to a C_{1-6} alkylene group. Examples include benzyl,
phenylethyl, phenylpropyl, phenylbutyl, etc.


Each amino acid may have either the D or L form.
25 Preferred oligopeptides will have 2-6 amino acid
residues, more preferably 2-3 residues. It is convenient
to use the 20 naturally occurring amino acids.

The oligosaccharide covalently bonded to the
porphycene may have 2-6, preferably 2-3 saccharide units.
30 Both pentose and hexose saccharides may be used,
including, but not limited to, glucose, mannose,
galactose and fructose.


-8-

Preferred compounds are those in which at least one R^1 is $-(CH_2)_n-X$ where X is OR^2 , OH, $OCOR^2$, OSO_2R^2 or halogen (I, Br, Cl), and compounds in which R^1 is $-(CH_2)_m-CH=CH_2$. Especially preferred are compounds in which at least one R^1 is $-(CH_2)_2-X$ where X is OCH_3 , OH, bromine and compounds in which R^1 is $-(CH_2)_n-O-G$, where G is a pentose or hexose.

The compounds of the present invention may be prepared by coupling appropriately substituted dialdehydes to form the porphycene ring structure as is shown below.



Porphycenes with two different substituents in the 2,7- and 12,17-positions are readily obtained by mixed coupling of two different dialdehydes as is shown below for the case of 2,7-bis(methoxyethyl)-12,17-di-n-propylporphycene.



The coupling of the dialdehyde components may be accomplished with a McMurry reaction utilizing low valency titanium in a non-reactive solvent. See for example McMurry and Fleming, *J. Am. Chem. Soc.*, 1974, 96:4708; McMurry, *Chem. Rev.*, 1989, 89:1513; Lenoir, *Synthesis*, 1989, 883; Mukaiyama et al, *Chem. Lett.*, 1973, 1041. Typically, reductive coupling is carried out using a titanium amalgam prepared by reacting titanium tetrachloride with an additional metal such as zinc, copper, aluminum and salts thereof. The reductive coupling reaction is carried out in a non-reactive solvent such as a hydrocarbon or ether solvent. Typical solvents include alkanes having from 5-10 carbon atoms such as pentane, hexane, heptane, etc., as well as ether solvents such as diethyl ether, crown ethers, glyme and

tetrahydrofuran (THF). If desired, the coupling reaction can be heated to facilitate the reaction, i.e., the reaction can be conducted under reflux conditions. The dialdehyde components to gain the bis(alkoxyalkyl)-
5 dialkylporphycenes, are on one hand those described below and, on the other hand, well known dialdehydes, preferred in the present invention, such as 5,5'-diformyl-4,4'-di-
n-propyl-bipyrrole (e.g., E.Vogel et al; Angew. Chem. Int. Ed. Engl. 26 (1987)928).

10 Synthesis of compounds in which R^1 is $-(CH_2)_n-X$ where X is OR^2 represent a convenient starting synthesis for the compounds of the present invention. With an appropriate cyano ether, it is possible to synthesize the dialdehyde coupling components which ultimately provide
15 the porphycene ring structure. Beginning with cyano ethers having the structure $R^2O-(CH_2)_n-CN$ where R^2 is C_{1-6} alkyl and n is 1-4, one can systematically construct the dialdehyde coupling components required to prepare the porphycene derivatives in which R^1 is $-(CH_2)_n-OR^2$. Figure
20 1 illustrates a synthesis of tetrakis(methoxyethyl)-porphycene starting from the cyano ether in which R^2 is CH_3 and $n = 2$. The remaining porphycene compounds in which R^1 is $-(CH_2)_n-OR^2$ can be synthesized by an analogous synthetic procedure using the appropriate cyano ether.
25 The cyano ethers may be purchased commercially or are available by cyano-dehalogenation displacement reactions on the corresponding halo-ether (Friedrich and Wallenfels, in Rappoport, "The Chemistry of the Cyano Group", pages 77-86, InterScience, New York, 1970). For
30 example, tetrakis(ethoxyethyl)porphycene, tetrakis(propoxyethyl)porphycene, tetrakis(butoxyethyl)-porphycene, tetrakis(pentoxylethyl)porphycene and tetrakis(hexyloxyethyl)porphycene can be prepared from starting cyanoethers in which $n = 2$ and R^2 is straight

-11-

chain or branched ethyl, propyl, butyl, pentyl and hexyl. Similarly, tetrakis(C₁₋₆ alkoxyethyl)porphycene, tetrakis(C₁₋₆ alkoxypropyl)porphycene, tetrakis(C₁₋₆ alkoxybutyl)porphycene, tetrakis(C₁₋₆ alkoxy

5 alkoxyethyl)porphycene and tetrakis(C₁₋₆ alkoxyhexyl)porphycene compounds can be prepared from the corresponding cyano ethers where R² is C₁₋₆ alkyl and n is 1, 3, 4, 5 or 6. A particularly preferred compound is tetrakis(methoxyethyl)porphycene. The bromoester which

10 is condensed with the cyano ether in the first step shown in Figure 1, is preferably a lower alkyl (C₁₋₆) bromo ester, more preferably a methyl or ethyl ester.

Reaction of the tetrakis(alkoxyalkyl)porphycene as its nickel or zinc complex, preferably the nickel

15 complex, with about 0.7 moles of BBr₃/B(OH)₃ results in conversion of a single alkoxyalkyl group into the corresponding bromoalkyl group. Reaction of about 1.2 moles of BBr₃/B(OH)₃ results in the corresponding bis(bromoalkyl)porphycenatonickel complex, and reaction

20 with 2 moles of BBr₃/B(OH)₃ gives the tris-product. The use of higher molar amounts of BBr₃/B(OH)₃ provides the tetrakis(bromoalkyl)porphycene compounds. The same products are obtained if solid boric acid is added to a solution of the substrate porphycene and boron tribromide

25 is added subsequently.

The reaction is generally conducted at low temperatures, for example about -40 to about -120°C, preferably about -78°C, and the reaction mixture then warmed to room temperature. The reaction is generally

30 conducted in an inert organic solvent which can be readily removed by distillation or vacuum evaporation. Preferred solvents are halogenated hydrocarbons and a particularly preferred solvent is dichloromethane.

Purification of the desired brominated porphycenatonickel complex is accomplished by conventional extraction and purification procedures. Preparation of the mono-, bis-, tris- and tetrakis(bromoalkyl)porphycenatonickel complex
5 of a preferred embodiment of the present invention is shown in Figure 2.

Metal complexes, preferably smaller metal complexes such as Ni^{+2} , Zn^{+2} , etc., of the porphycene compounds of the present invention can be easily prepared by the
10 addition of metal salts, e.g., metal acetates, to the porphycene compounds in acid medium, such as glacial acetic acid. Demetallation occurs when the metal complex is reacted with concentrated sulfuric acid at room temperature with stirring. Hydrogen ions displace the
15 metal atom during the demetallation reaction (Buchler, J.W. in Smith, K.M. (Ed): "Porphyrins and Metalloporphyrins", Elsevier, Amsterdam, 1975; Buchler, J.W. in Dolphin, D. (Ed), "The Porphyrins," Vol. I, Academic Press, New York, 1978; Dorrough et al, J. Am.
20 Chem. Soc., 1951, 73:4315).

Reaction of the tetrakis(alkoxyalkyl)porphycene compounds with BBr_3 may also yield hydroxyalkyl porphycenes. By controlling the amount of boron tribromide reagent, it is possible to optimize the
25 production of 1, 2, 3 or 4 hydroxyalkyl substituents (R^1 is $-(CH_2)_n-OH$), leaving the remaining alkoxyalkyl substituents intact. The synthesis of hydroxyalkyl substituents on the porphycene is possible if boric acid is omitted and the substrate is the uncomplexed
30 porphycene.

The remaining compounds of the present invention are available synthetically by further reactions on any of

-13-

the mono-, bis-, tris- or tetrakis(bromoalkyl)- or (hydroxyalkyl)porphycene compounds. Olefinic groups (R^1 is $-(CH_2)_m-CH=CH_2$), such as vinyl groups, can be introduced into the porphycene compound by reaction of the appropriate bromoalkyl porphycene under dehydrohalogenation conditions. Loss of HBr results in the synthesis of mono-, bis-, tris- and tetrakis(vinyl)porphycene compounds from the corresponding mono-, bis-, tris- and tetrakis(bromoethyl)porphycene compounds ($n=2$). Analogously, a carbon-carbon double bond can be introduced into substituent R^1 by eliminating HBr from a substituent having the formula $-(CH_2)_n-Br$, where $n = 3$ or 4. The dehydrohalogenation can be conducted under any suitable dehydrohalogenation conditions, generally through the addition of a base such as an alkoxide, amine or hydroxide base. For example, t-butoxide, diazobicycloundecene (DBU) or aqueous sodium hydroxide/pyridine may be used to dehydrohalogenate the bromoalkyl porphycene and thereby introduce a carbon-carbon double bond. Obviously, other dehydrohalogenation conditions not specifically disclosed are suitable for the dehydrohalogenation reaction so long as dehydrohalogenation is effected. A particularly preferred dehydrohalogenating reagent is DBU.

Through dehydrohalogenation, porphycene compounds in which R^1 is $-(CH_2)_m-CH=CH_2$ where m is 0, 1 or 2 can be prepared. The dehydrohalogenation reaction can be used to prepare porphycene compounds in which 1, 2, 3 or 4 of the R^1 substituents contain a carbon-carbon double bond by dehydrohalogenating the corresponding mono-, bis-, tris- or tetrakis(bromoalkyl)porphycene.

Additional porphycene compounds are available by

displacing the bromine atom or better the methanesulfonate group, which is prepared in high yield from the hydroxy compound, in R^1 groups having the formula $-(CH_2)_n-Br$ or $-(CH_2)_n-OSO_2CH_3$ respectively. For example, nucleophilic displacement of a bromide ion by cyanide, chloride, iodide, ammonia, primary amines (NH_2R^2), secondary amines (HNR^2_2), H_2S and thiolate ions ($-SR^2$) provides porphycene compounds where R^1 is $-(CH_2)_n-X$ and X is $I, Cl, CN, NH_2, NHR^2, NR^2_2, SH$ and SR^2 .

10 Conventional oxidation of the SR^2 group with known oxidizing agents yields, the $S(O)_{1-2}R^2$ groups (Oae, in Oae, "The Chemistry of Sulfur", Plenum, New York, 1977). These aliphatic nucleophilic substitution reactions are well known in the art of organic synthesis (March,

15 "Advanced Organic Chemistry, Reactions, Mechanisms, and Structure", 3rd Ed., John Wiley & Sons, New York, 1985 and references cited therein).

Hydrolysis of the cyano porphycene compounds (R^1 is $-(CH_2)_n-CN$) with alcoholic hydrogen chloride provides the

20 corresponding carboxylic esters in which R^1 is $-(CH_2)_n-COOR^2$. These are readily converted to the corresponding carboxylic acids, which can also be prepared by conventional hydrolysis methods but in low yield. Hydrolysis of the porphycene ester with alkali or

25 alkaline earth metal hydroxide solutions yields the corresponding alkali and alkaline earth metal carboxylate salts by well known conventional hydrolysis reactions for example.

Carboxylic acid-containing porphycene compounds can

30 further be converted to the acid halide by the conventional addition of thionyl bromide, chloride, PBr_3 or PCl_3 . The acid halide derivative is then readily converted to the corresponding amide by the addition of

ammonia, a primary amine or secondary amine to provide porphycene derivatives in which R^1 is $-(CH_2)_n-C(O)NH_2$, $-(CH_2)_n-C(O)NHR^2$, and $-(CH_2)_n-C(O)NR_2^2$. The acid halide derivative may also be converted to the corresponding ester (R^1 is $-(CH_2)_n-CO_2R^2$) by reaction with an alcohol (HOR²). Esters may also be prepared by esterification of the carboxylic acid porphycene compounds using known esterification reactions. Porphycene derivatives in which R^1 is $-(CH_2)_n-CHO$ can be readily prepared by reducing the carboxylic acid to the aldehyde using known reagents such as $(iso-Bu)_2AlH$ or by oxidizing the alcohol (R^1 is $-(CH_2)_{n-1}-CH_2OH$) using a weak oxidizing agent such as the chromium trioxide/pyridine, MnO_2 or other suitable reagents (see March, loc. cit.).

To improve water solubility of the porphycene compounds, a saccharide may be covalently attached to the porphycene via an ether linkage to provide porphycene compounds in which R^1 is $-(CH_2)_n-O-G$, where G is a monosaccharide. Preferred monosaccharides are pentose and hexose monosaccharides such as glucose, galactose, mannose, xylose, fructose, etc. The monosaccharide derivatives can be prepared by known chemistry, for example by reacting a porphycene containing at least one R^1 group having the formula $-(CH_2)_n-OH$ with a OH-protected bromosaccharide, e.g., tetraacetyl bromoglucose. See the examples below and Fulling et al, Angew. Chem., 1989, 101:1550.

The monosaccharide-containing porphycene glycosides confer an increased level of hydrophilicity to the porphycene compounds which facilitates the preparation of aqueous solutions for pharmaceutical use. The glycosides have adequate solubility for preparation of aqueous, aqueous-alcoholic (preferably ethanolic) and aqueous

-16-

dimethylsulfoxide (DMSO) solutions which may be directly topically applied for PDT therapy. Topical application of these aqueous solutions is particularly effective for treating psoriatic lesions. To increase water solubility
5 the porphycene compounds may also be covalently attached to an amino acid or peptide by reaction of the substituent side chain, $-(\text{CH}_2)_n\text{-X}$ (X = OH or halogen, for example) or $-(\text{CH}_2)_n\text{-CH=CH}_2$ with an alcohol/phenol-, a carboxyl-, an amine or a thiol-group of the amino acid or
10 peptide.

The present invention includes porphycene metal complexes, for example zinc, nickel, magnesium and tin complexes, of the porphycene compounds described above. The invention also includes pharmaceutically acceptable
15 acid and base addition salts of the porphycene compounds which may be prepared by the known addition of acids such as HCl, HBr, H_2SO_4 , H_3PO_4 , malic acid, tartaric acid, maleic acid, fumaric acid, etc. Base addition salts are prepared by the addition of alkali and alkaline earth
20 metal salts such as sodium, potassium, calcium and magnesium carbonates, bicarbonates, sulfates, phosphates, etc. as well as by addition of ammonia, amines, preferably primary, secondary and tertiary C_{1-6} alkyl amines, amino acids, etc. Any conventional acid or base
25 addition salt which is pharmaceutically acceptable is considered to be within the scope of the present invention.

The porphycene compounds may also be covalently bonded to a stationary phase, such as beads, plates,
30 fibers, etc. by reaction of the substituent side chain, $-(\text{CH}_2)_n\text{-X}$ (X = OH or halogen, for example) or $-(\text{CH}_2)_m\text{-CH=CH}_2$ with a reactive group in the stationary support by addition or displacement reactions. When

-17-

attached to a stationary solid phase, the porphycene compounds may be used to purify and decontaminate fluids passed over the solid phase. For example, blood for use in transfusions or from a patient undergoing

5 plasmaphoresis procedures can be passed through a column containing resin beads having bound thereto the porphycene compounds of the present invention. Irradiation of the porphycene dye bound to the solid support at the absorption maximum wavelength for the

10 particular porphycene dye results in the generation of singlet oxygen species which are lethal to viral, retroviral and bacterial contaminants in the blood or biological fluid. Addition of small amounts of the porphycene compounds into blood, followed by irradiation

15 and subsequent transfusion into the patient undergoing therapy is also possible. The porphycene dyes are metabolized *in vivo* and excreted from the patient.

The porphycene compounds of the present invention may be formulated as therapeutic formulations for

20 administration to patients in need of photodynamic therapy.

THERAPEUTIC FORMULATIONS

Therapeutic compositions containing the compounds of the present invention include liposome or microvesicle

25 preparations, dispersions, solutions for parenteral injection, etc. and including topical dermatological preparations.

Parenteral Solutions

The photoactivatable porphycene dyes generally are

30 used with additional solvents and adjuvants to prepare

-18-

solutions suitable for intravenous injection. A number of solvents and co-solvents that are miscible with water and suitable surfactants can be used to achieve solutions for parenteral use. The most important solvents in this group are ethanol, polyethylene glycols of the liquid series and propylene glycol. A more comprehensive listing includes acetone, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide ethanol, glycerin, polyethylene glycol 300, and 400, propylene glycol, sorbitol, polyoxyethylene sorbitan fatty acid esters such as laureate, palmitate, stearate, and oleate, polyoxyethylated vegetable oil, sorbitan monopalmitate, 2-pyrrolidone; n-methyl-2-pyrrolidine; n-ethyl-1-pyrrolidine; tetrahydrofurfuryl alcohol, tween 80 and dimethyl isosorbide. Dimethyl isosorbide (ARLASOLVE® DMI, ICI Specialty Chemicals) has the advantage of being both water- and oil-soluble. Additionally, dimethyl isosorbide may be readily gelled with a gelling agent to produce gel formulations with, for example, 4% KLUCEL® (Hercules).

Other additives may be necessary to enhance or maintain chemical stability and physiological suitability. Examples are antioxidants, chelating agents, inert gases, buffers and isotonicifiers.

Examples of antioxidants and typical concentration ranges include acetone sodium bisulfite (0.1-0.8%), ascorbic acid (0.05-1.0%), monothioglycerol (0.1-1.0%), potassium metabisulfite (0.05-0.1%), propyl gallate (0.02%), sodium bisulfite (0.01-1.0%), sodium formaldehyde sulfoxylate (0.03-0.1%), sodium metabisulfite (0.02-0.25%), sodium sulfite (0.01-0.1%), sodium thioglycolate (0.05-0.1%).

-19-

Examples of chelating/complexing agents and typical concentration ranges include edetate sodium (0.005-0.1%), edetate calcium disodium (0.005%-0.01%), gentisic acid ethanolamide (1.0%-2.0%), niacinamide (1.0%-2.5%), sodium
5 citrate (0.01%-2.5%), citric acid (0.001%-1.0%).

Examples of inert gases are nitrogen and carbon dioxide.

Buffers are used primarily to stabilize a solution against the chemical degradation that might occur if the
10 pH changed appreciably. Buffer systems employed normally have as low a buffer capacity as feasible in order to not disturb significantly the body buffer systems when injected. The buffer range and effect of the buffer on activity must be evaluated. Appropriate adjustment is
15 useful to provide the optimum conditions for pH dependent partition into the target malignant tissues or lesion area.

Examples of such buffer systems include the following acids: acetic, adipic, ascorbic, benzoic,
20 citric, glycine, lactic, tartaric, hydrochloric, phosphoric, sulfuric, carbonic and bicarbonic; and their corresponding salts such as: potassium, sodium, magnesium, calcium and diethanolamine salts.

Osmoticity is of great importance and hypotonic
25 solutions usually have their tonicity adjusted by the addition of salts such as sodium chloride, potassium chloride, magnesium chloride and calcium chloride and sugars such as dextrose, lactose, mannitol and sorbitol.

When the solution will be dispensed from multiple
30 dose containers, antimicrobial agents in bacteriostatic

or fungistatic concentrations must be added. Among the compounds and concentrations most frequently employed are phenylmercuric acid (0.002-0.01%), thimerosal (0.01%), benzethonium chloride (0.01%), benzalkonium chloride
 5 (0.01%), phenol or cresol (0.5%), chlorbutanol (0.5%), benzyl alcohol (2.0%), methyl p-hydroxybenzoate (0.18%), and propyl p-hydroxybenzoate (0.02%).

After the solution of the porphycene with its solvents and additives has been compounded, the solution
 10 is generally filtered to remove particulate matter above 2 μ m in size and a further step eliminating particulate matter down to 0.2 μ m can eliminate microorganisms and accomplish cold sterilization. The solution is filled under aseptic conditions. The final solution can be
 15 additionally sterilized in its final container by thermal methods such as autoclaving or non-thermal methods such as ionizing radiation. The process of freeze drying (lyophilization) can be employed to avoid adverse thermal and oxidative decomposition and provide enhanced
 20 stability and improved solubility.

The following formula provides an example of the utilization of various solvents and additives such as have been heretofore mentioned in the creation of a suitable parenteral solution of the porphycene. The
 25 formula is by way of example only and is not limiting to this invention. Suitable combinations and variations are obvious to those skilled in the art.

Formula example for tetrakis(methoxyethyl)porphycene

(TMEP)

30		<u>Grams</u>
	TMEP	0.1
	Tetrahydrofurfurylalcohol	40.0

-21-

Polysorbate 20	1.0
Sodium chloride	0.9
Citric acid buffer	0.1
water* enough to make 100 ml	

- 5 *water may be water for injection, bacteriostatic water for injection or sterile water for injection.

Method of Preparation

1. Dissolve porphycene in tetrahydrofurfuryl alcohol and polysorbate 20, using heat and stirring as
10 needed.
2. Dissolve sodium chloride and citrate buffer in water.*
3. Add the water solution slowly with stirring and heat as necessary to the solution.
- 15 4. Sterile fill using aseptic conditions and use terminal sterilization as needed.

This solution is suitable for a broad dosage range such as 0.1-10 mg/kg and preferably 0.2-5.0 mg/kg and may be infused as such or added to suitable large volume
20 parenteral solutions such as dextrose, saline, ringers solutions for slower intravenous administration. Suitable solutions are described, for example, in REMINGTON'S PHARMACEUTICAL SCIENCES, 15th Ed., Easton: Mack Publishing Co. incorporated herein by reference.

Topical Formulations

The porphycene compounds of the present invention may be formulated for topical application in penetrating solvents or in the form of a lotion, cream, ointment or
5 gel containing a sufficient amount of the porphycene compound to be effective for PDT therapy.

Suitable penetrating solvents are solvents for the porphycene compound which will enhance percutaneous penetration of the porphycene compound. Solvents which
10 have this property include dimethyl sulfoxide, dimethyl acetamide, dimethylformamide, 1-methyl-2-pyrrolidone, diisopropyladipate, diethyltoluamide and to a lesser extent propylene glycol. Additional solvents include
15 substituted azacycloalkan-2-ones having from 5 to 7 carbons in the cycloalkyl group such as 1-dodecylazacycloheptan-2-one (AZONE) and other azacycloalkan-2-ones such as described in U.S. 3,989,816 incorporated herein by reference.

Also included are N-bis-azocyclopentan-2-onyl
20 alkanes described in U.S. 3,989,815 (hereby incorporated by reference), 1-substituted azacyclopentan-2-ones described in U.S. 3,991,203 (hereby incorporated by reference) and water-soluble tertiary amine oxides described in U.S. 4,411,893 (hereby incorporated by
25 reference).

The topical formulations contain a sufficient amount of the porphycene compound to be effective in PDT therapy. Generally, concentrations in the range of 0.001 to 5 wt.%, preferably from about 1 to 5 wt.%, may be
30 used. Typical lotion and cream formulations are shown below.

-23-

LOTION

	<u>Parts by Weight</u>	<u>Ingredient</u>
	5	polyoxyethylene-40-sterate
	3	sorbitan monostearate
5	12	*mixture of lanolin, mineral oil and lanolin alcohol
	6	cetyl alcohol
	20	soybean oil
10	53.7	water
	0.2	methyl paraben
	0.1	propyl paraben

*AMERCOL BL (Amerchol Corp. Edison, N.J.)

15

CREAM

	<u>Parts by Weight</u>	<u>Ingredient</u>
	3	polyoxyethylene-40-sterate
	2.5	sorbitan monostearate
20	10	*mixture of lanolin, mineral oil and lanolin alcohol
	10	cetyl alcohol
	1	soybean oil
	73.2	water
25	0.2	methyl paraben
	0.1	propyl paraben

*AMERCOL BL (Amerchol Corp. Edison, N.J.)

Additional topical formulations which may be used in conjunction with the porphycene compounds of the present invention are disclosed in U.S. 3,592,930 and U.S. 4,017,615 (hereby incorporated by reference).

5 Topical formulations may be prepared in gel form by combining the porphycene with a solvent such as diethyltoluamide (DEET) or diisopropyl adipate (DIPA) and adding a gelling agent. A preferred gelling agent is fumed silica (CAB-O-SIL®, Cabot Corp., Tuscola, IL), and
10 particularly grade M-5. The gelling agent is generally used in amounts of about 5-12 wt% to obtain a gel with the desired viscosity. Obviously, gels containing more or less gelling agent will have slightly higher or lower viscosity. One skilled in the art can readily obtain the
15 desired gel viscosity by adjusting the concentration of gelling agent. Additives, such as cosolvents and/or surfactants, frequently improve the gel properties and may be added as desired. Suitable cosolvents/surfactants include propylene glycol and glycerine. The additives
20 may be incorporated into the gel by mechanically mixing the additives into a mixture of solvent and gelling agent.

Liposome or Microvesicle Preparations

Liposomes and methods of preparing liposomes are
25 known and are described for example in U.S. 4,452,747 and U.S. 4,448,765 incorporated herein by reference. Liposomes are microvesicles which encapsulate a liquid within lipid or polymeric membranes. The porphycene compounds of the present invention may be incorporated
30 into liposome microvesicles and used in this form for both topical and parenteral application. Topical and parenteral liposome preparations are known in the art.

-25-

Sonified unilamellar liposomes made from certain unsaturated lipids are known stable carriers for some of the porphycenes of the invention.

U.S. 4,837,028 discloses injectable liposome
5 formulations having enhanced circulation time. The liposomes have a size of about 0.08-0.5 microns, contain at least 50 mole % of a membrane rigidifying component such as sphingomyelin and further contain about 5-15 mole % ganglioside G_{M1}. Liposome preparations for
10 encapsulating sparingly soluble pharmaceutical compounds are disclosed in U.S. 4,721,612. The specification of these U.S. patents is incorporated herein by reference.

After administration of a therapeutically effective amount of one or more of the porphycene compounds in the
15 pharmaceutical composition or preparation, to a patient having a treatable condition such as a solid tumor (cancer) or psoriasis, for example, the patients affected body area is exposed to a therapeutically sufficient amount of light having an appropriate wavelength for
20 absorption by the particular porphycene compound used. Suitable wavelengths are generally from about 600 to about 900 nm, preferably from about 600 to about 650 nm, more preferably 620-650 nm. Irradiation of the accumulated porphycene generates singlet oxygen which is
25 thought to be the actual lethal species responsible for destruction of the neoplastic cells.

Photodynamic therapy using the porphycene compounds of the present invention has a number of advantages. The porphycene compound itself is minimally toxic in the
30 unexcited state. Each porphycene molecule can be repeatedly photoactivated and lead each time to cell-lethal events, that is, the generation of singlet

-26-

molecular oxygen. The half-life of singlet molecular oxygen is approximately four microseconds in water at room temperature. The target cell is therefore affected without the opportunity for migration of the lethal
5 singlet molecular oxygen to neighboring healthy tissue cells. Preferably, the singlet oxygen molecules rupture chemical bonds in the target cell wall or mitochondria resulting in destruction of the target cell. Destruction of target cell tissue commences promptly upon irradiation
10 of the porphycene compounds. Indirect target cell death can also result from destruction of the tumor vascular system with concomitant restriction of oxygen supply.

Photodynamic therapy using the compounds of the present invention is therefore selective and minimally
15 toxic to healthy tissue. Singlet oxygen molecules produced which do not react rapidly decay to harmless ground state oxygen molecules.

A variety of phototherapy and irradiation methodologies are known to those skilled in the art and
20 can be used with the novel porphycene compounds of the present invention. The time and duration of therapy and repetition of the irradiation treatment can be selected by the therapist (physician or radiologist) according to known photodynamic therapy criteria. The dosage of the
25 porphycene compound may be varied according to the size and location of the target tissues which are to be destroyed and the method of administration. Generally, the dosage will be in the range of 0.05-10 mg of porphycene compound per kilogram of body weight, more
30 preferably in the range of 0.1-5.0 mg/kg.

Irradiation generally takes place not less than one hour nor more than four days after parenteral

-27-

administration of the porphycene compound. Usually, phototherapy is begun approximately 3 hours to 48 hours after administration of the photodynamic therapy agent. With topically administered dye, radiation may commence
5 as soon as 10 minutes after dye application for treatment of psoriasis, genital warts, bacterial infections, etc. Exposure to non-therapeutic light sources should be avoided immediately following phototherapy to minimize light toxicity. Appropriate draping of the patient can
10 be used to limit the area affected by phototherapy.

Light sources which are appropriate for use are well known in the art and may vary from white light sources with appropriate filters to lasers. As noted above, preferred wavelengths are from 600 to 950 nm, preferably
15 from about 600 to about 800 nm. The total amount of light which is applied to the affected area will vary with the method used and the location of the tumor or topical lesion. Generally, the amount of light is in the range of about 50 to 1000 J-cm² preferably in the range
20 of 100 to 350 J-cm².

Having generally described this invention, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only and are not intended to be limiting
25 unless otherwise specified. Procedures which are constructively reduced to practice herein are described in the present tense and procedures which have been carried out in the laboratory are set forth in the past tense.

-28-

EXAMPLESZinc/copper pair

A solution of 20g (0.1 mol) $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ in 400 mL of glacial acetic acid was treated rapidly with 196 g (3
5 mol) zinc powder with vigorous stirring. Within 15 seconds the solution discolored and was subsequently stirred for 30 seconds. The solution was then filtered under inert gas and washed in succession twice with acetone, toluene, and absolute THF. The Zn/Cu pair thus
10 produced can be used without drying.

5-Methoxy-3-oxopentanoic acid alkylester (1) (see Figure 1)

68 g (0.8 mol) of 3-methoxypropionitrile were dissolved in 1000 mL THF freshly distilled over LiAlH_4
15 and treated with 150.3 g (2.2 mol) Zn/Cu pair under inert gas. With moderate reflux 267.2 g (1.6 mol) of 2-bromoacetic acid ethyl ester were added drop-by-drop to the vigorously stirred suspension within 75 minutes. After the reaction had started, the heat was turned off
20 and the slight reflux adjusted via the drop velocity. To complete the reaction, the solution was stirred for 30 minutes at the boiling temperature of THF. Then, 420 mL of 10% HCl were added drop-by-drop within 30 minutes at 15°C and stirred for another 30 minutes. The solution
25 was filtered from the non-converted zinc and the filtrate was extracted three times with 300 mL of CHCl_3 each. The organic phases were washed four times with water and once with 5% NaHCO_3 solution and the solvent was removed following drying over MgSO_4 under vacuum. The obtained
30 reddish liquid was distilled under oil pump vacuum. The fraction at 85°C/0.5 torr yielded 106.8 g (0.62 mol) of

5-methoxy-3-oxopentanic acid ethyl ester as a colorless, fruity smelling oil with a refractive index $n_D^{22}=1.434$. The yield was to 77.5% based on 3-methoxypropionitrile.

Elementary analysis: calculated: C 55.16% H 8.10%
 5 found: C 54.82% H 7.92%

If 2-bromoacetic acid methyl ester is added as the α -bromoester, 97.2 g (0.61 mol) of 5-methoxy-3-oxopentanic acid methyl ester having a boiling point of 62 - 63°C/0.12 torr are obtained with
 10 the same method. The refractive index of the colorless oil, which was produced in a yield of 76% based on 3-methylpropionitrile, is $n_D^{21}=1.435$.

$^1\text{H-NMR}$ spectrum of (1a) (CDCl_3 , 80 MHz), δ (ppm) =
 2.62 (triplet, 2H) $\text{MeOCH}_2\text{CH}_2$; 3.16 (singlet, 3H) CH_3OCH_2 ;
 15 3.35 (singlet, 2H) COCH_2CO ; 3.48 (triplet, 2H) MeOCH_2 ;
 3.56 (singlet, 3H) CO_2CH_3 ; signals at 4.9 (singlet) and 2.3 ppm (triplet) through the enol form present in the equilibrium.

Mass spectrum of (1a), EI, 70 eV: $m/z = 160$ (M^+ , 3%); 129 ($\text{M-O}^+\text{Me}$, 12%); 101 ($\text{M-CO}_2\text{Me}^+$, 40%); 100 ($\text{M-MeOCH}_2\text{-CH}_3^+$, 33%); 87 ($\text{M-CH}_2\text{CO}_2\text{Me}^+$, 85%).

$^1\text{H-NMR}$ spectrum of (1b), CDCl_3 , 80 MHz, δ (ppm) =
 1.06 (triplet, 3H) CH_2CH_3 ; 2.57 (triplet, 2H) $\text{MeOCH}_2\text{CH}_2$;
 3.10 (singlet, 3H) CH_3OCH_2 ; 3.28 (singlet, 2H) COCH_2CO ;
 25 3.43 (triplet, 2H) MeOCH_2 ; 3.97 (quartet, 2H) CH_2CH_3 .

Mass spectrum of (1b), EI, 70 eV: $m/z = 174$ (M^+ , 1%); 143 ($\text{M-O}^+\text{Me}$, 12%); 100 ($\text{M-MeOCH}_2\text{-CH}_2\text{CH}_3^+$, 40%); 87 ($\text{M-CH}_2\text{CO}_2\text{Et}^+$, 85%).

-30-

IR spectrum of (1b), film: ν (cm^{-1}) = 2987, 2941, 2903 (CH), 1743, 1724 (C=O), 1118 (C-O, ether)

2,4-Bis(alkoxycarbonyl)-3-methoxyethyl-5-methylpyrroles
(2)

5 While stirring and ice cooling, a cold saturated aqueous solution of 42 g (0.6 mol) sodium nitrite was added so slowly drop-by-drop to a solution of 104.4 g (0.6 mol) methoxy-pentanoic acid ethyl ester in 450 mL of glacial acetic acid, that no noticeable NO_x development
10 was detected. The solution was stirred for two hours and treated with 83.2 g (0.64 mol) acetoacetic acid ethyl ester. Then while stirring vigorously, a mixture of 80 g (1.22 mol) of zinc powder and 100 g (1.22 mol) sodium acetate was added in portions within 40 minutes. The
15 inner temperature ranged from 80 to 85°C, a state that was reached by cooling occasionally. Following completion of the addition, the solution was stirred another 30 minutes and then filtered from the nonreacted zinc. The warm solution was poured into 4 L of water and
20 left standing overnight to complete the precipitation. The raw product was filtered off, washed with a little ice cold ether, dried over P_4O_{10} under vacuum for one day, and recrystallized from ethanol/water. Thus, 113.2 g (0.4 mol) 2,4-bis(ethoxycarbonyl)-3-methoxyethyl-
25 5-methylpyrrole (2b) was obtained as colorless needles with a melting point of 86°C. Yield: 65% based on (1b). For further reaction of (2b) it was not necessary to recrystallize the raw product from ethanol/water.

Elementary analysis: calculated: C 59.16% H 7.47% N 4.94%
30 found: C 59.24% H 7.33% N 4.99%

The workup to isolate the corresponding pyrrol-

-31-

dimethyl ester (2a) was the same as in the case of the application of 96 g (0.6 mol) 5-methoxy-3-oxopentanoic acid methyl ester (1a) and 74.2 g (0.64 mol) acetoacetic acid methyl ester. However, the mother liquor of the first crystallization was extracted once with CHCl_3 and the solid obtained following neutralization and drying over MgSO_4 and evaporation of the solvent was combined with the quantity of substance that precipitated first. In this case recrystallization from methanol/water for esterification was avoided. The result was 96.4 g (0.38 mol) colorless needles of 2,4-bis(methoxycarbonyl)3-methoxyethyl-5-methylpyrrole (2a), which melt at 118°C . Yield: 63%.

Elementary analysis: calculated: C: 56.46 H: 6.71 N: 5.49
found: C: 56.42 H: 6.96 N: 5.45

$^1\text{H-NMR}$ spectrum of (2a), CDCl_3 , 300 MHz, δ (ppm) = 2.45 (singlet, 3H) $\alpha\text{-CH}_3$; 3.33 (singlet, 3H) CH_3OCH_2 ; 3.33 (triplet, 2H) $\text{MeOCH}_2\text{CH}_2$; 3.50 (triplet, 2H) MeOCH_2 ; 3.77 (singlet, 3H) $\beta\text{-CO}_2\text{CH}_3$; 3.80 (singlet, 3H) $\alpha\text{-CO}_2\text{CH}_3$; 9.98 (wide singlet) NH .

$^{13}\text{C-NMR}$ spectrum of (2a), CDCl_3 , 75.5 MHz, δ (ppm) = 14.01 ($\alpha\text{-CH}_3$); 25.57 ($\text{MeOCH}_2\text{CH}_2$); 50.66 ($\beta\text{-CO}_2\text{CH}_3$); 51.36 ($\alpha\text{-CO}_2\text{CH}_3$); 58.1 (H_3COCH_2); 72.78 ($\text{MeOCH}_2\text{CH}_2$); 112.88 (C-4); 118.21 (C-2); 130.80 (C-3); 139.86 (C-5) 161.94 ($\beta\text{-CO}_2\text{CH}_3$); 165.44 ($\alpha\text{-CO}_2\text{CH}_3$).

Mass spectrum of (2a), EI, 70 eV: m/z = 255 (M^+ , 12%); 223 (M-MeOH^+ , 77%); 201 (M-MeOCH_2^+ , 47%); 178 (M-MeOH-MeOCH_2^+ , 100%).

IR spectrum of (2a), KBr pellet: ν (cm^{-1}) = 3306, (NH); 2962, 2943, 2868 (CH), 1711 (C=O).

-32-

UV/VIS spectrum of (2a) CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 227
(1.4×10^4); 270 (1.6×10^4).

$^1\text{H-NMR}$ spectrum of (2b), CDCl_3 , 80 MHz, δ (ppm) =
1.33 (2 triplets, 6H) CH_2CH_3 ; 2.48 (singlet, 3H) $\alpha\text{-CH}_3$;
5 3.35 (singlet, 3H) CH_3OCH_2 ; 3.45 (A_2B_2 system, 4H)
 $\text{MeOCH}_2\text{CH}_2$; 4.27 (quartet, 2H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 4.32 (quartet,
2H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 9.5 (wide singlet): NH.

IR spectrum of (2b) KBr pellet: ν (cm^{-1}) 3289, (NH);
2989, 2932, 2891 (CH); 1703, 1663 (C=O); 1438 (CH).

10 Mass spectrum of (2b) EI, 70 eV: m/z = 283 (M^+ ,
13%); 251 (M-MeOCH_2^+ , 15%); 164 ($\text{M-MeOCH}_2\text{-CH}_3\text{-EtOH}^+$, 18%).

UV/VIS spectrum of (2b), CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 226
(1.3×10^4); 271 (1.6×10^4).

15 3,5-Bis(alkoxycarbonyl)-2-carboxy-4-methoxyethylpyrroles
(3)

At one time 48 g (0.3 mol) of distilled bromine were
added to a solution of 85 g (0.3 mol)
 α -methylpyrroldiethyl ester in 280 mL glacial acetic acid
and 55 mL of acetic anhydride at 0°C . Subsequently 128.5
20 g (1.05 mol) freshly distilled sulfuryl chloride were
added drop-by-drop at this temperature in the dark over a
period of two hours. Since the reaction solution becomes
viscous in the interim, the use of a KPG® stirrer is
recommended. The red solution obtained was then stirred
25 at 0°C for two hours. Then, 190 mL of water were added
dropwise within 30 minutes, and the inner temperature
rose to 70°C . After another 30 minutes of stirring, the
mixture precipitated out by pouring in 2.5 L of water.
It was left standing overnight and was filtered the next

-33-

day. If the result was an oily raw product, it was decanted. At this stage the mixture was suspended in 2 L of 70°C warm water and treated in portions with solid NaHCO₃ until the generation of gas was no longer
5 observed. It was filtered and slowly neutralized with semiconcentrated HCl while stirring vigorously. The mixture was left standing at 0°C for eight hours in order to complete the precipitation and the resulting fine crystalline solid was filtered off. The pyrrolcarboxylic
10 acid (3b) was recrystallized from ethanol/water and dried under vacuum for two days. 47 g (0.15 mol) of colorless needles having a solid point of 136°C were obtained. The yield of (3b) was 50%. For further reaction of (2b) it was not necessary to recrystallize the raw product from
15 ethanol/water.

With the same method, 76.5 g (0.3 mol) methylpyrroldi-methyl ester (2a) yield, following recrystallization from MeOH/water, 42.8 g (0.15 mol) of
3,5-bis(methoxycarbonyl)-2-carboxy-4-methoxyethylpyrrole
20 (3a). The colorless needles have a melting point of 142°C. The yield was 50%.

¹H-NMR spectrum of (3a), CDCl₃, 300 MHz, δ (ppm) =
3.36 (singlet, 3H) CH₃OCH₂; 3.45 (A₂B₂ system, 4H)
MeOCH₂CH₂; 3.91 (singlet, 3H) β-CO₂CH₂CH₃; 4.02 (singlet,
25 3H) α-CO₂CH₃; 10.30 (wide singlet) NH.

IR spectrum of (3a), KBr pellet: ν (cm⁻¹) 3247 (NH); 2958, 2900 (CH), 2616 (OH); 1722, 1632 (C=O); 1441 (CH).

Mass spectrum of (3a), EI, 70 eV: m/z = 285 (M+, 28%); 270 (M-CH₃+, 11%); 253 (M-MeOH+, 85%); 222
30 (M-MeOH-MeO+, 100%); 208 (M-MeOCH₂-MeOH+, 75%).

-34-

$^1\text{H-NMR}$ spectrum of (3b), CDCl_3 , 80 MHz, δ (ppm) = 1.38 (triplet, 3H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 1.45 (triplet, 3H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 3.33 (singlet, 3H) CH_3OCH_2 ; 3.46 (A_2B_2 system, 4H) $\text{MeOCH}_2\text{CH}_2$; 4.40 (quartet, 2H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 4.50 (quartet, 2H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 10.25 (wide singlet) NH ; 14.64 (wide singlet) CO_2H .

IR spectrum of (3b), KBr pellet: ν (cm^{-1}) 3258, (NH); 2980, 2934, 2881 (CH), 2568 (OH); 1737, 1693 (C=O); 1116 (C-O, ether).

10 Mass spectrum of (3b), EI, 70 eV: $m/z = 313$ (M^+ , 34%); 281 (M-MeOH^+ , 41%); 252 ($[\text{M-EtOH-CH}_3]^+$, 28%); 222 ($\text{M-MeOCH}_2\text{-EtOH}^+$, 48%).

UV/VIS spectrum of (3b), CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 237 (2.7×10^4); 257 (9.2×10^3); 288 (8.0×10^3).

15 3,5-Bis(alkoxycarbonyl)-2-iodo-4-methoxyethylpyrroles (4)

While stirring, 47 g (0.15 mol) bis(ethoxycarbonyl) pyrrole carboxylic acid were suspended in 350 mL of water and treated in portions with 40.9 g (0.48 mol) of sodium hydrogen carbonate at 75°C . At this stage a solution of 20 38.1 g (0.15 mol) I_2 and 49.8 g (0.3 mol) KI in 280 ml of water were added dropwise to this clear solution within two hours. The reaction solution foamed due to the resulting CO_2 . The product (4b) began to precipitate after some time. Following completion of the addition, 25 the temperature was maintained for another 30 minutes while stirring and then the still warm reaction solution was poured on 700 g of ice. The precipitate was filtered off, washed with water and ice cold pentane and recrystallized from ethanol so that colorless needles 30 were obtained. The 53.3 g (0.135 mol) of the title

-35-

compound that were obtained following drying under vacuum exhibit a melting point of 132°C. Yield: 90%.

Elementary analysis: calculated: C 39.51% H 4.59% N 3.54%
found: C 39.36% H 4.54% N 3.45%

5 When 42.8 g (0.15 mol) bis(methoxycarbonyl) pyrrolcarboxylic acid (3a) were used, 49.5 g (0.135 mol) iodopyrroledimethyl ester (4a) in 90% yield was obtained following recrystallization from methanol with the same procedure. The melting point of the colorless needles
10 was 114°C.

$^1\text{H-NMR}$ spectrum of (4a), CDCl_3 , 80 MHz, δ (ppm) = 3.35 (singlet, 3H) H_3COCH_2 ; 3.45 (A_2B_2 system, 4H) $\text{MeOCH}_2\text{CH}_2$; 3.84 (singlet, 3H) $\beta\text{-CO}_2\text{CH}_3$; 3.88 (singlet, 3H) $\alpha\text{-CO}_2\text{CH}_3$; 9.54 (wide singlet) NH .

15 Mass spectrum of (4a), EI, 70 eV: m/z = 367 (M^+ , 28%); 355 (M-MeOH^+ , 100%); 322 (M-MeOCH_2^+ , 35%); 290 ($\text{M-MeOCH}_2\text{-MeOH}^+$, 59%).

IR spectrum of (4a), KBr pellet: ν (cm^{-1}) = 3263 (NH); 2955, 2902 (CH), 1676 (C=O); 1440 (CH); 1120 (C-O ,
20 ether).

UV/VIS spectrum of (4a) MeOH, CH_2Cl_2 , λ_{max} [nm] (ϵ) = 222 (2.0×10^4); 266 (1.8×10^4).

$^1\text{H-NMR}$ spectrum of (4b), CDCl_3 , 80 MHz, δ (ppm) = 1.36 (triplet, 3H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 1.37 (triplet, 3H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$;
25 3.34 (singlet, 3H) CH_3OCH_2 ; 3.46 (A_2B_2 system, 4H) $\text{MeOCH}_2\text{CH}_2$; 4.33 (quartet, 2H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 4.36 (quartet, 2H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 9.5 (wide singlet) NH .

-36-

Mass spectrum of (4b), EI, 70 eV: $m/z = 395$ (M+, 30%); 363 (M-MeOH+, 100%); 322 (M-CO₂Et+, 31%); 276 (M-EtOH-CH₂H₅-MeOCH₂+, 42%).

IR spectrum of (4b), KBr pellet: ν (cm⁻¹) 3140 (NH);
5 2989, 2830 (CH), 1707 (C=O); 1105 (C-O, ether).

UV/VIS spectrum of (4b), CH₂Cl₂, λ_{max} [nm] (ϵ) = 227 (1.4 x 10⁴); 267 (1.8 x 10⁴).

3,3',5,5'-Tetrakis(alkoxycarbonyl)-4,4'-bis(methoxyethyl)
2,2'-bipyrroles (5)

10 45 g (0.7 mol) copper bronze were added to a solution of 49.1 g (0.12 mol) of iodopyrrolediethyl ester in 250 mL of absolute dimethylformamide and the mixture was stirred at room temperature for twenty hours. In so doing, the suspension assumed a greenish-brown color.
15 The raw product was subsequently precipitated by slowly adding 1.2 L of water and filtering off over Celite® together with the non-converted copper. The filter residue was extracted with 400 mL of hot chloroform. The extract was washed briefly with 20% HNO₃, twice with
20 water and once with 5% NaHCO₃ solution. After drying over MgSO₄ the solvent was evaporated and the remaining solid was digested with 50 mL of cold hexane and subsequently dried under oil pump vacuum. Following recrystallization from ethanol, 39.9 g (74.4 mmol) of
25 bipyrroltetraethyl ester (5b) were obtained in the form of matted needles having a melting point of 150°C.
Yield: 67%

Elementary analysis: calculated: C 58.20% H 6.76% N 5.22%
found: C 58.05% H 6.72% N 5.25%

-37-

Under analogous condition, 45.5 g (94.8 mmol) of bipyrrrole (5a) was recovered from 44 g (0.12 mol) of the iodopyrroledimethyl ester (4a) following recrystallization from methanol. The melting point of the colorless, matted needles was at 221°C. Yield: 79%.

$^1\text{H-NMR}$ spectrum of (5a), CDCl_3 , 80 MHz, δ (ppm) = 3.38 (singlet, 6H) H_3COCH_2 ; 3.50 (A_2B_2 system, 8H) $\text{MeOCH}_2\text{CH}_2$; 3.96 (singlet, 6H) $\beta\text{-CO}_2\text{CH}_3$; 3.98 (singlet, 6H) $\alpha\text{-CO}_2\text{CH}_3$; 13.95 (wide singlet) NH .

10 Mass spectrum of (5a), FAB, 75 eV: m/z = 480 (M^+ , 76%); 449 (M-MeO^+ , 40%); 417 (M-MeOH-MeO^+ , 24%); 403 ($\text{M-MeOCH}_2\text{-MeOH}^+$, 12%); 307 (M-4MeOH-MeOCH_2^+ , 33%).

IR spectrum, KBr pellet, ν (cm^{-1}) = 2953, 2985 (CH), 1711 (C=O), 1451, 1311.

15 UV/VIS spectrum of (5a), CHCl_3 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 250 (2.5×10^4); 288 (1.5×10^4); 346 (2.2×10^4).

IR spectrum of (34b), KBr pellet: ν (cm^{-1}) = 2988, 2893 (CH), 1716 (C=O), 1479, 1383 (CH); 1117 (C-O, ether).

20 $^1\text{H-NMR}$ spectrum of (5b), CDCl_3 , 80 MHz, δ (ppm) = 1.41 (triplet, 3H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 1.42 (triplet, 3H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 3.34 (singlet, 6H) H_3COCH_2 ; 3.49 (A_2B_2 system, 8H) $\text{MeOCH}_2\text{CH}_2$; 4.39 (quartet, 4H) $\beta\text{-CO}_2\text{CH}_2\text{CH}_3$; 4.43 (quartet, 4H) $\alpha\text{-CO}_2\text{CH}_2\text{CH}_3$; 14.10 (wide singlet) NH .

25 Mass spectrum of (5b) EI, 75 eV: m/z = 536 (M^+ , 100%); 504 (M-MeOH^+ , 3%); 490 (M-MeO-CH^+ , 11%); 445 ($\text{M-MeOCH}_2\text{-EtOH}^+$, 26%).

-38-

UV/VIS spectrum of (5b), CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 228 (1.9×10^4); 249 (2.6×10^4); 289 (1.5×10^4); 347 (2.3×10^4).

3,3',5,5'-Tetracarboxy-4,4'-bis(methoxyethyl)-2,2'-
5 bipyrrole (6)

A solution of 26.8 g (50 mmol) of bipyrrole tetraethyl ester (5b) or 24 g (50 mmol) of tetramethyl ester (5a) in 1350 mL of methanol were treated with a solution of 30 g (0.75 mol) NaOH in 540 mL of water and
10 heated with reflux for 40 hours. Following distillation of methanol, the mixture was diluted with water to 4 L and slowly neutralized with diluted hydrochloric acid while stirring. The mixture was stirred at the neutral point for fifteen minutes at room temperature and then
15 precipitated at pH 2 with the further addition of acid. The mixture was left standing at 0°C for six hours and the precipitate was filtered, washed with water and dried for five days over P_4O_{10} under vacuum. 1.5 g (46 mmol) of
20 that, following recrystallization from DMSO/water yielded a microcrystalline, colorless substance, which decomposed at 248°C.

Elementary analysis: calculated: C 50.95% H 4.75% N 6.60%
found: C 50.94% H 4.87% N 6.47%

25 $^1\text{H-NMR}$ spectrum of (6), DMSO-d_6 , 300 MHz, δ (ppm) = 3.23 (singlet, 6H) H_3COCH_2 ; 3.40 (A_2B_2 system, 8H) $\text{MeOCH}_2\text{CH}_2$; 12.70 (very wide singlet) CO_2H ; 13.15 (wide singlet) NH .

$^{13}\text{C-NMR}$ spectrum of (6), DMSO-d_6 , 75.5 MHz, δ (ppm)
30 = 25.27 ($\text{MeOCH}_2\text{CH}_2$); 57.61 (H_3COCH_2); 72.50 ($\text{MeOCH}_2\text{CH}_2$);

-39-

114.83, 120.52, 129.61, 132.68 (C-2, 3, 4, 5): 161.78, 167.22 (CO₂H).

UV/VIS spectrum of (6), DMSO, λ_{\max} [nm] (ϵ) = 258 (1.5 x 10⁴); 277 (1.5 x 10⁴); 343 (1.3 x 10⁴).

5 Mass spectrum of (6), EI, 70 eV: m/z = 292 (M-3CO₂⁺, 10%); 248 (M-4CO₂⁺, 36%); 203 (M-4CO₂MeOCH₂⁺, 26%); 171 (M-4CO₂-MeOCH₂-MeOH⁺, 100%).

IR spectrum of (6), KBr pellet, ν (cm⁻¹) = 2922 (CH), 1633 (C=O), 1554, 1187, 756.

10 4,4'-Bis(methoxyethyl)-2,2'-bipyrrole (7)

8.5 g (20 mmol) of bipyrrole tetracarboxylic acid (6) were decarboxylated in a sublimator at 230°C (bath temperature)/ 0.1 - 0.2 torr. The product (7) separated as a colorless, amorphous or pale green crystalline
15 solid. The bipyrrole (7) decomposed slowly above 75°C and melts when rapidly heated at 98°C. Weighing yielded 4.7 g (18.8 mmol) of the title compound in a 94% yield.

The bipyrrole was sensitive to acid and oxygen and was stored under protective gas at low temperature.

20 ¹H-NMR spectrum of (7), acetone-d₆, 300 MHz, δ (ppm) = 2.64 (triplet, 4H) MeOCH₂CH₂; 3.27 (singlet, 6H) H₃COCH₂; 3.46 (triplet, 4H) MeOCH₂CH₂; 6.11 (multiplet, 2H) H at C-3; 6.2 (multiplet, 2H) H at C-5; 9.79 (wide singlet) NH.

25 Mass spectrum of (7), EI, 70 eV: m/z = 248 (M⁺, 82%); 203 (M-MeOCH₂⁺, 28%); 171 (M-MeOCH₂-MeOH⁺, 100%); 157 (M-MeOCH₂-MeOCH₃⁺, 33%).

-40-

IR spectrum (7), KBr pellet, ν (cm^{-1}) = 3284 (NH); 3123, 3089, 2949, 2852 (CH) 1568, 1185, 1082, 946, 802.

UV/VIS spectrum of (7), CH_3CN , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 343 (8.0×10^2); 284 (1.8×10^4).

5 5,5'-Diformyl-4,4'-bis(methoxyethyl)-2,2-bipyrrole (8)

6.14 g (40 mmol) of distilled phosphoryl chloride was added dropwise to a solution of 2.48 g (10 mmol) bipyrrole (7) in 50 mL of absolute dimethylformamide under protective gas at 0°C within 30 minutes. The
10 mixture was heated to 60°C for one hour and subsequently poured into a solution of 60 g sodium acetate in 480 mL of water. The mixture was stirred for one hour at 85°C , wherein the dialdehyde (8) precipitated out in yellow
15 flakes. The solution was cooled to 0°C , the precipitate filtered off and washed with cold water. After drying over P_4O_{10} under vacuum, 2.58 g (8.5 mmol) of the title compound which was recrystallized from tetrahydrofuran, were obtained. The title compound decomposes at 186°C without melting. The yield was 85%.

20 Elementary analysis: calculated: C: 63.14 H: 6.62 N: 9.21
found: C: 63.07 H: 6.54 N: 9.15

$^1\text{H-NMR}$ spectrum of (8), CDCl_3 , 300 MHz, δ (ppm) =
3.03 (triplet, 4H) $\text{MeOCH}_2\text{CH}_2$; 3.37 (singlet, 6H) H_3COCH_2 ;
3.62 (triplet, 4H) $\text{MeOCH}_2\text{CH}_2$; 6.54 (doublet, 2H) H at C-3;
25 9.7 (singlet, 2H) CHO ; 12.27 (wide singlet) NH .

$^{13}\text{C-NMR}$ spectrum of (8), 75.5 MHz, δ (ppm) = 26.03 ($\text{MeOCH}_2\text{CH}_2$); 58.74 (H_3COCH_2); 72.91 ($\text{MeOCH}_2\text{CH}_2$); 111.79, 130.60, 131.57, 136.10 (C-2, 3, 4, 5), 177.71 (C-5¹).

-41-

Mass spectrum of (8), EI, 75 eV: $m/z = 304$ (M+, 100%); 272 (M-MeOH+, 9%); 227 (M-MeOCH₂-MeOH+, 19%); 199 (M-MeOCH₂-MeOH-CO+, 19%).

IR spectrum of (8), KBr pellet: ν (cm⁻¹) = 3266 (NH); 2925, 2868 (CH), 1648 (C=O), 1116 (C-O, ether) 1607, 809.

UV/VIS spectrum of (8), CH₂Cl₂ λ_{max} [nm] (ϵ) = 235 (9.6 x 10³); 271 (1.5 x 10⁴); 381 (3.6 x 10⁴).

2,7,12,17-Tetrakis(methoxyethyl)porphycene (9)

10 Under protective gas over 10 minutes, 16.5 ml (0.15 mol) TiCl₄ were added dropwise to a suspension of 19.6 g (0.3 mol) zinc powder and 1.95 g (9.5 mmol) CuCl in 800 mL of THF freshly distilled over LiAlH₄. Subsequently the mixture was heated with reflux for three hours. At
15 this stage a solution of 1.83 g (6 mmol) dialdehyde (8) in 600 mL of absolute THF was added dropwise within 20 minutes to the black McMurry reagent thus prepared while stirring vigorously. The reaction can be followed by means of thin layer chromatography (silica gel/CH₂Cl₂).
20 The reaction was stirred for ten minutes at the boiling temperature of the tetrahydrofuran and then cooled to 0°C. At this temperature 300 ml of 6% NH₃ solution were added dropwise over one hour. The reaction mixture was treated with 600 mL of dichloromethane and filtered off
25 over Celite® after fifteen minutes of stirring. The residue was extracted with another 200 mL of CH₂Cl₂ and the combined organic phases were washed three times with 300 mL of water. Following drying over MgSO₄, the solvent was evaporated under vacuum and the residue was
30 chromatographed with CH₂Cl₂ on Al₂O₃ (Brockmann, II-III activity, column 5 x 10 cm). The first blue, red

-42-

fluorescing fraction was collected and rechromatographed with CH_2Cl_2 /ethylacetate/methanol (100:20:1) on silica gel (column 4 x 40 cm) in order to separate any possibly existing small quantities of degraded porphycenes as a
5 small blue fraction, which was eluted prior to the title compound. Following evaporation of the solvent and crystallization from CH_2Cl_2 /MeOH, (9) was obtained in the form of long, violet, metallicly glistening needles having a melting point of 172°C. A 25% yield of 406 mg
10 (0.75 mmol) of the porphycene tetraether (9) was obtained.

Elementary analysis: calculated C 70.83% H 7.06% N 10.32%
found C 70.71% H 6.93% N 10.40%

$^1\text{H-NMR}$ spectrum of (9), CDCl_3 , 300 MHz, δ (ppm) =
15 3.11 (wide singlet) NH; 3.60 (singlet, 12H); 4.31
(singlet, 16H) $\text{MeOCH}_2\text{CH}_2$; 9.34 (singlet, 4H) H at C-3, 6,
13, 16; 9.71 (singlet, 4H) H at C-9, 10, 19, 20.

$^{13}\text{C-NMR}$ spectrum of (9), CDCl_3 , 75.5 MHz, -60°C , δ
(ppm) = 28.57 (C- $\text{MeOCH}_2\text{CH}_2$); 59.24 ($\text{H}_3\text{C}\text{COCH}_2$); 73.54
20 ($\text{MeOCH}_2\text{CH}_2$); 110.41 (C-9, 10, 19, 20); 123.37 (C-3, 6, 13,
16); 133.54 (C-4, 5, 14, 15); 140.28, 142.80 (C-1, 2, 7,
8, 11, 12, 17, 18).

Mass spectrum of (9) EI, 75 eV: m/z = 542 (M+,
81%); 497 (M- MeOCH_2 +, 100%); 452 (M-2 MeOCH_2 +, 20%); 407
25 (M-3 MeOCH_2 +, 24%); 375 (M-3 MeOCH_2 - MeOH +, 3%).

IR spectrum; CsI pellet, ν (cm^{-1}) 2920, 2867 (CH),
1113 (C-O, ether), 1459, 1017, 816.

UV/VIS spectrum of (9) CH_2Cl_2 , λ_{max} [nm] (ϵ) = 370
(1.4×10^5); 563 (3.5×10^4); 602 (3.3×10^4); 634 ($4.7 \times$

10⁴).

¹H-NMR spectrum of 2-methyl-7,12,17-tri(methoxyethyl)-porphycene, CDCl₃, 80 MHz, δ (ppm) = 3.12 (wide singlet) NH; 3.6 (singlet, 12H) H₃COCH₂ and H₃C at C-2); 4.36 (singlet, 12H) MeOCH₂CH₂; 9.25 (singlet, 1H), 9.34 (3 singlets, 3H) H at C-3, 6, 13, 16; 9.68, 9.71 (2 singlets, 4H) H at C-9, 10, 19, 20.

Mass spectrum of 2-methyl-7,12,17-tri(methoxyethyl)-porphycene, EI, 75 eV: m/z = 498 (M⁺, 78%); 453 (M-MeOCH₂⁺, 100%); 408 (M-2MeOCH₂⁺, 26%).

9-Acetoxy-2,7,12,17-tetrakis(methoxyethyl)porphycene

A solution of 542 mg (1 mmol) 2,7,12,17-tetrakis(methoxyethyl)porphycene in 75 ml CH₂CL₂ freshly distilled over LiAlH₄, and 75 ml glacial acetic acid was combined with 179 mg (0.75 mmol) PbO₂ and stirred for 20 min. at room temperature. The reaction mixture was then poured into 500 ml water and extracted with 250 ml dichloromethane. After washing the organic phase once with concentrated aqueous sodium hydrogen carbonate (150 ml) and twice with water (200 ml), the organic layer was evaporated under vacuum. The residue was chromatographed with dichloromethane/ethylacetate (1:1) on silica gel (column 40 X 5 cm). The first eluted compound consisted of unchanged tetrakis(methoxyethyl)-porphycene (195 mg). Following evaporation of the solvent and crystallization the second, largest fraction from methanol/water, the title compound 9-acetoxy-tetrakis(methoxyethyl)porphycene was obtained in the form of small, violet needles having a melting point of 107°C. Yield: 153 mg, 40%

Mass spectrum, EI, 70 eV, m/z = 600 (M⁺, 87%); 558

-44-

([M-CH₂CO]⁺, 100%); 557 ([M-CH₃CO]⁺, 22%); 513 ([M-MeOCH₂-CH₂CO]⁺, 36%).

UV/VIS spectrum, CH₂Cl₂, λ_{max}[nm] (ε) = 370 (1.39 x 10⁵); 382 (9.1 x 10⁴); 562 (2.9 x 10⁴); 604 (3.2 x 10⁴);
5 633 (3.2 x 10⁴); 640 (3.0 x 10⁴).

IR spectrum, CsI pellet, ν (cm⁻¹) 2924, 2872 (C-H), 1758 (C=O), 1115 (C-O), 1458, 1074, 1002, 816.

Solubility of the title compound in selected solvents:

10 Excellent: ethyl acetate, THF, CHCl₃, CH₂Cl₂;

Good: DMSO, DEET, DMF, Toluene, acetone;

Moderate: EtOH, MeOH, i-propanol;

Insoluble: hexane, water.

2,7-Bis(methoxyethyl)-12,17-di-n-propylporphycene

15 A stirred suspension under protective gas of 17.4 g (260 mmol) zinc powder and 1.7 g (16.8 mmol) anhydrous CuCl in 700 ml of THF freshly distilled over LiAlH₄ was mixed in 30 minutes dropwise with 14.2 ml (130 mmol) TiCl₄. Subsequently the mixture was heated under reflux
20 for three hours. At this stage, a solution of 1.36 g (5 mmol) 5,5'-diformyl-4,4'-di-n-propyl-2,2'-bipyrrole and 1.52 g (5 mmol) 5,5'-diformyl-4,4'-bis(methoxyethyl)-2,2'-bipyrrole in 300 ml THF, was added, with stirring,
25 within 45 minutes to the black McMurry reagent. The reaction was stirred for two minutes at the boiling temperature of the tetrahydrofuran and then cooled to

-45-

0°C. At this temperature 250 ml of 10% NH₃ solution were added dropwise over one hour. The reaction mixture was treated with 500 ml THF, stirred under oxygen for 90 minutes and filtered off. The residue was extracted with 5 200 ml of CH₂Cl₂ and the combined organic layers were evaporated under vacuum. The residue, which contained black polymers, was dissolved as thoroughly as possible in dichloromethane and filtered over Celite. The black residue was ground up and extracted with further 10 dichloromethane. The combined organic phases were evaporated under vacuum and the dark blue residue was taken up into CH₂Cl₂ and placed on an alumina column. (Brockmann, II-III activity, column 10 x 5 cm). All the blue, red fluorescing fractions were collected and 15 rechromatographed with CH₂Cl₂/ethyl acetate (100:4) on silica gel (column 25 x 5 cm). The first eluted compound consisted of tetra-n-propylporphycene, followed by the main compound bis(methoxyethyl)-di-n-propylporphycene. The third fraction was the 20 tetrakis(methoxyethyl)porphycene, eluted with ethyl acetate. After dividing those three fractions, it was necessary to further remove small quantities of minor compounds by recrystallization. The first fraction with tetra-n-propylporphycene was recrystallized after 25 evaporation of the solvent from dichloromethane/hexane.

By evaporation of the solvent and recrystallization of the second fraction from CH₂Cl₂/MeOH, the title compound bis(methoxyethyl)-di-n-propylporphycene was obtained in the form of long, violet, metallicly 30 glistening needles having a melting point of 142°C. The porphycene-tetraether (third fraction) was purified by recrystallization from CH₂Cl₂/MeOH.

Total yield of three porphycenes: 22% (560 mg; 1.10 mmol)

-46-

- 2,7-bis(methoxyethyl)-12,17-di-n-propylporphycene: 11% (280 mg; 0.55 mmol)
- 2,7,12,17-tetrakis(methoxyethyl)-porphycene: 5.5% (146 mg; 0.27 mmol)
- 5 2,7,12,17-tetra-n-propylporphycene: 5.5% (134 mg; 0.28 mmol)

$^1\text{H-NMR}$ spectrum of bis(methoxyethyl)-di-n-propylporphycene, CDCl_3 , 300 MHz, δ (ppm) = 1.34 (triplet, 6H) $-\text{CH}_2\text{CH}_2\text{CH}_3$; 2.41 (multiplet, 4H) $-\text{CH}_2\text{CH}_2\text{CH}_3$; 3.14 (wide singlet, 2H) NH ; 3.59 (singlet, 6H) $-\text{CH}_2\text{CH}_2\text{OCH}_3$; 4.00 (triplet, 4H) $-\text{CH}_2\text{CH}_2\text{CH}_3$; 4.31 (singlet, 8H) $-\text{CH}_2\text{CH}_2\text{OCH}_3$; 9.27 (singlet, 2H) H-13, H-16; 9.34 (singlet, 2H) H-3, H-6; 9.71 (singlet, 4H) H-9, H-10, H-19, H-20.

$^{13}\text{C-NMR}$ spectrum, CDCl_3 , 75.5 MHz, δ (ppm) = 14.51 ($-\text{CH}_2\text{CH}_2\text{CH}_3$); 25.22 ($-\text{CH}_2\text{CH}_2\text{CH}_3$); 28.89 ($-\text{CH}_2\text{CH}_2\text{OCH}_3$); 30.45 ($-\text{CH}_2\text{CH}_2\text{CH}_3$); 58.99 ($-\text{CH}_2\text{CH}_2\text{OCH}_3$); 74.09 ($-\text{CH}_2\text{CH}_2\text{OCH}_3$); 110.52 (C-9, C-10, C-19, C-20); 122.88 (C-13, C-16); 123.42 (C-6); 134.16 (C-4, C-5); 134.22 (C-14, C-15); 140.65 (C-2, C-7); 143.34 (C-11, C-18); 143.82 (C-1, C-8); 145.01 (C-12, C-17).

Mass spectrum, EI, 70 eV, m/z = 510 (M^+ , 100%); 481 ($[\text{M}-\text{C}_2\text{H}_5]^+$, 13%); 466 ($[\text{M}-\text{C}_3\text{H}_8]^+$, 31%); 466 ($[\text{M}-\text{MeOCH}_2]^+$, 88%); 420 ($[\text{M}-2\text{MeOCH}_2]^+$, 32%); 255 (M^{2+} , 4%).

UV/VIS spectrum, CH_2Cl_2 , λ max[nm] (ϵ) = 370 (142000); 382 (100000); 562 (36000); 601 (34000); 633 (49000).

IR spectrum, CsI pellet, ν (cm^{-1}) 2955, 2927, 2868 (C-H); 1116 (C-O); 1458, 1214, 1083, 1018, 816.

2,7,12,17-Tetrakis(methoxyethyl)porphycenatonickel (II)

A suspension of 380 mg (0.7 mmol) tetrakis(methoxyethyl)porphycene in 120 mL of acetic acid was heated with reflux with 1.7 g (7 mmol) Ni(OAc)₂·4H₂O for five hours. The reaction can be followed by means of thin layer chromatography (CH₂Cl₂/ethyl acetate (4:1), silica gel). In so doing, the nickel complex appears as a less mobile, blue fraction without fluorescence. Following complete complexing, the mixture was poured into 600 mL of water and extracted three times with 150 mL CHCl₃. The combined organic phases were washed twice with water and a saturated NaHCO₃ solution and again twice with water, then dried over MgSO₄ and the solvent evaporated under vacuum. 398 mg (0.67 mmol) bluish-violet, matted needles of the title compound having a melting point of 182°C was obtained from the residue through recrystallization from CH₂Cl₂/MeOH. The yield was 95%.

If the raw product of (9) obtained following the first chromatography is added, a mixture of partially degraded porphycenes can be separated in the form of its nickel complex even at this stage by means of chromatography on silica gel (column 4 x 40 cm) with CH₂Cl₂/ethyl acetate (3:1).

¹H-NMR spectrum of the title compound, CDCl₃, 300 MHz, δ (ppm) = 3.58 singlet, (12H) H₃COCH₂; 4.20 (A₂B₂ system, 16H) MeOCH₂CH₂; 8.82 (singlet, 4H) H at C-3, 6, 13, 16; 9.28 (singlet, 4H) H at C-9, 10, 19, 20.

Mass spectrum, DEI, 75 eV; m/z = 599 (M⁺, 15%); 554 (M-MeOCH₂⁺, 8%).

-48-

IR spectrum, CsI pellet, ν (cm^{-1}) 2887, 2870, 2810 (CH), 1117 (C-O, ether), 1489, 974, 839, 812.

UV/VIS spectrum CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 265 (3.1×10^4); 387 (1.27×10^5); 603 (6.3×10^4).

5 Bis-, tris-, and tetrakis(bromoethyl)porphycenes

120 mg (0.2 mmol) tetrakis(methoxyethyl)-porphycenatonickel (II) were dissolved under protective gas in 150 mL of CH_2Cl_2 freshly distilled over LiAlH_4 and treated with 100 mg (1.6 mmol) boric acid. The reaction was cooled to -78°C and 100 mg (75 μL , 0.8 mmol) BBr_3 were added at one time under cooling and in the dark. Cooling the boron tribromide to -20°C facilitates its handling. The reaction was thawed over ten hours and then 100 mL of 5% NaHCO_3 solution were added at 0°C within 30 minutes while stirring vigorously. Another 200 mL of CH_2Cl_2 are added and the organic phase was separated out. This was washed twice, each time with a saturated NaHCO_3 solution and water and dried over MgSO_4 . The solvent was evaporated under vacuum and the residue was dried at 0.1 torr.

The blue solid obtained was treated with 20 mL of 98% H_2SO_4 and stirred for 30 minutes. The resulting blue, red fluorescing solution was poured into 1.2 L of deionized water and extracted four times with 150 mL of CHCl_3 . The combined organic phases were washed with water and a 5% sodium hydrogen carbonate solution and the solvent was removed under vacuum. The residue was chromatographed twice with chloroform on silica gel (column 2 x 30 cm). From the first blue fraction of the second chromatography, 29 mg (0.04 mmol) of 2,7,12,17 tetrakis(bromoethyl)porphycene were obtained by means of

-49-

crystallization from CHCl_3 in the form of small, violet needles, which did not exhibit a melting point until 300°C . The yield was 20%.

A similar procedure can be applied to the three
5 isomeric bis(bromoethyl)bis(methoxyethyl)porphycenes and 2,7,12-tris(bromoethyl)-17-(methoxyethyl)porphycene. In the case of bis(bromoethyl)bis(methoxyethyl)porphycenes, 30 mg (0.48 mmol) boric acid and 60 mg (0.24 mmol) BBr_3 were used for 120 mg (0.2 mmol) of the nickel complex.
10 To isolate the tris(bromoethyl) compound, 50 mg (0.8 mmol) B(OH)_3 and 100 mg (0.4 mmol) BBr_3 were added to the same quantity of the nickel complex. The yield was 25% and 30%, respectively. The tris(bromoethyl)porphycene begins to decompose at 250°C , becoming black, without
15 exhibiting a melting point until 300°C when heated rapidly.

Mass spectrum of 2,7,12,17-tetrakis(bromoethyl)-porphycenatonickel (II), EI, 75 eV: $m/z = 794/96$ (M^+ , 3%); $704/06$ ($\text{M}-\text{Br}^+$, 58%); $467/69$ ($\text{M}-4\text{Br}^+$, 30%).

20 ^1H -NMR spectrum of 2,7,12,17-tetrakis(bromoethyl)porphycene (TBEP), D_2SO_4 , 80 MHz, δ (ppm) = 4.32 and 4.53 (2 triplets, 16H) BrCH_2CH_2 ; 10.16 (singlet, 4H) H at 9, 10, 19, 20.

IR spectrum of (TBEP), CsI pellet, ν (cm^{-1}) = 2958,
25 2920 (CH), 1261, 1046, 799.

UV/VIS spectrum of (TBEP), CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 369 (1.02×10^5); 383 (7.9×10^4); 564 (2.5×10^4); 605 (2.9×10^4); 637 (3.2×10^4 .)

^1H -NMR spectrum of bis(bromoethyl)bis(methoxyethyl)-

porphycene (mixture of isomers), CDCl_3 , 80 MHz, δ (ppm) =
3.20 (wide singlet) NH ; 3.61 (singlet, 6H) H_3COCH_2 ; 4.33
(singlet, 8H) $\text{MeOCH}_2\text{CH}_2$; 4.26, 4.50, 4.59 (3 multiplets,
8H) BrCH_2CH_2 ; 9.37 (multiplet, 4H) H at C-3, 6, 13, 16;
5 9.76, 9.66 (3 multiplets, 4H) H at C-9, 10, 19, 20.

$^1\text{H-NMR}$ spectrum of 2,7,12-tris(bromoethyl)-17-
methoxyethylporphycene, CDCl_3 , 80 MHz, δ (ppm) = 3.11
(wide singlet) NH ; 3.61 (singlet, 3H) H_3COCH_2 ; 4.19, 4.58
(2 multiplets, 12H) BrCH_2CH_2 ; 4.32 (singlet, 4H)
10 $\text{MeOCH}_2\text{CH}_2$; 9.34, 9.35 (2 singlets, 4H) H at C-3, 6, 13,
16; 9.62, 9.67, 9.68 (3 singlets, 4H) H at C-9, 10, 19,
20.

Mass spectrum, EI, 75 eV; m/z = 497 ($\text{M}-2\text{Br}-\text{MeOH}+$,
2%); 94/96 ($\text{CH}_3\text{Br}+$, 100%).

15 IR spectrum, CsI pellet, ν (cm^{-1}) 2962, 2925 (CH),
1140 (C-O, ether), .963, 816.

2,7,12,17-Tetravinylporphycene

At room temperature while stirring in the dark, 243
mg (248 μL , 1.6 mmol) DBU were added all at once to a
20 solution of 14.8 mg (20 μmol)
tetrakis(bromoethyl)porphycene in 50 mL of absolute THF.
After thirty minutes, the agitator was turned off and the
mixture left to stand at room temperature for two days.
Subsequently the greenish-blue solution was treated with
25 80 mL of CH_2Cl_2 and poured into 80 mL of 0.5% HCl. The
organic phase was shaken out and washed with water and a
2% NaHCO_3 solution, dried over Na_2SO_4 and the solvent was
evaporated under vacuum. The residue was chromatographed
with CH_2Cl_2 on aluminum oxide (Brockmann, II-III activity
30 stage, column 2 x 10 cm) and the first green fraction was

-51-

collected. The solvent was slowly evaporated under a slight vacuum and the remaining fine crystalline solid was washed with 3 mL of ice cooled, absolute pentane. Following drying under a high vacuum, 4.1 mg (10 μ mol) 5 tetravinylporphycene were obtained. The title compound decomposed even with traces of oxygen in the presence of light. The yield was 50%.

All reagents and solvents were purged of oxygen by introducing inert gas and all purification steps were 10 carried out under subdued light, as was the reaction.

$^1\text{H-NMR}$ spectrum of, CD_2Cl_2 , 300 MHz, δ (ppm) = 3.64 (wide singlet) NH; 6.01 (doublet of doublets, 4H) $\text{CH}=\underline{\text{CH}}_2$ -cis $^3\text{J}_{\text{cis}}=11.0$ Hz; 6.68 (doublet of doublets, 4H) $\text{CH}=\underline{\text{CH}}_2$ -trans $^3\text{J}_{\text{cis}}=17.4$ Hz; 8.31 (doublet of doublets, 4H) 15 $\underline{\text{CH}}=\text{CH}_2$; 9.68 (singlet, 4H) H at C-3, 6, 13, 16; 9.86 (singlet, 4H) H at C-9, 10, 19, 20.

UV/VIS spectrum, CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 383 (3.2×10^4); 597 (1.1×10^4); 642 (1.5×10^4); 676 (1.3×10^4).

20 2-Bromoethyl-7,12,17-tris(methoxyethyl)porphyceneato-nickel(II)

A solution of 419 mg (0.7 mmol) tetrakis(methoxyethyl)-porphycenatonickel complex in 250 mL of CH_2Cl_2 freshly distilled over LiAlH_4 was treated 25 with 62 mg (1 mmol) boric acid and 126 mg (48 μL , 0.5 mmol) BBr_3 were added all at once under protective gas at -78°C . The mixture was left to thaw over eight hours and 50 mL of 5% NaHCO_3 solution were added dropwise over 10 minutes at 0°C . The organic phase was separated off and 30 washed sequentially with a saturated NaHCO_3 solution and

-52-

water. Following drying over MgSO_4 , the solvent was evaporated under vacuum and purified by means of chromatography on silica gel (column 4 x 40 cm, CH_2Cl_2 /ethyl acetate/methanol (80:20:1)). The first three
5 fractions contained various ether-split nickel porphycenes; the 2-bromoethyl-7,12,17-tris(methoxyethyl)porphycenato-nickel (II) was eluted as the fourth fraction. Following removal of the solvent and recrystallization from CH_2Cl_2 /methanol there resulted
10 159 mg (0.245 mmol, yield 35%, reaction yield 70%) of bluish-violet crystals having a melting point of 186-189°C. The fifth blue band yielded after evaporation of the solvent and crystallization 209 mg (0.35 mmol) of the educt, 2,7,12,17-tetrakis(methoxyethyl)porphycenato-
15 nickel (II).

$^1\text{H-NMR}$ spectrum, CDCl_3 , 300 MHz, δ (ppm) = 3.58, 3.59 (2 singlets, 9H) H_3COCH_2 ; 4.06, 4.10 (2 multiplets, 16 H) $\text{MeOCH}_2\text{CH}_2$ and BrCH_2CH_2 ; 8.34, 8.45 (2 singlets, 2H) H at C-3, 6; 8.55, 8.57 (2 singlets, 2H) H at C-13, 16; 8.70
20 (A_2B_2 system, 2H) H at C-19, 20; 8.95 (singlet, 2H) H at C-9, 10.

$^{13}\text{C-NMR}$ spectrum, CDCl_3 , 75.5 MHz, δ (ppm) = 29.1 ($\text{CH}_2\text{CH}_2\text{OMe}$; 32.3, 32.4 ($\text{CH}_2\text{CH}_2\text{Br}$); 59.0 (CH_2OCH_3); 73.7 ($\text{CH}_2\text{CH}_2\text{OMe}$); 106.6, 106.8, 107.0, 107.2, (C-9, 10, 19,
25 20); 119.1, 119.4, 119.5, 119.6 (C-3, 6, 13, 16); 145.1, 144.1, 143.8, 142.6, 146.6 (C-4, 5, 14, 15, 1, 2, 7, 8, 11, 12, 17, 18).

IR spectrum, CsI pellet, ν (cm^{-1}) = 2920, 2870 (CH) 1116 (C-O, ether), 1618, 1560, 1488, 974, 811.

30 Mass spectrum, EI, 75 eV: m/z = 648 (M^+ , 45%); 603 (M-MeOCH_2^+ , 54%); 432 ($\text{M-3MeOCH}_2\text{-Br}^+$, 18%); 418

-53-

(M-3MeOCH₂-CH₂Br⁺, 21%); 94 (CH₃Br⁺, 35%); 80 (HBr⁺, 100%).

UV/VIS spectrum, CH₂Cl₂, λ_{max}[nm] (ε) = 264 (3.2 x 10⁴); 387 (1.28 x 10⁵); 605 (6.5 x 10⁴).

5 2-Bromoethyl-7,12,17-tris(methoxyethyl)porphycene

To demetallate, 97.2 mg (0.15 mmol) of the above-described bromoethylporphycenatonickel complex were suspended in 12 mL of concentrated sulfuric acid and stirred for ten minutes until a clear solution was
10 produced. The reaction solution was poured in 1 L of water and extracted three times with 150 mL of CHCl₃. The combined organic phases were washed three times with water and once with a 1% NaHCO₃ solution and dried over
15 MgSO₄. Following evaporation of the solvent under vacuum and recrystallization from CH₂Cl₂/MeOH, 80 mg (0.135 mmol) of the title compound were obtained in the form of long, violet, metallicly glistening needles with a melting point of 164°C. Yield: 90 percent.

Elementary analysis: calc. C 62.94% H 5.96% N 9.47% Br 13.51%
20 found C 62.87% H 5.87% N 9.55% Br 13.56%

¹H-NMR spectrum, CF₃CO₂D, 300 MHz, δ (ppm) = 3.83, 3.84 (2 singlets, 9H) H₃COCH₂; 4.27 (triplet, 2 H) BrCH₂CH₂; 4.56, 4.66 (2 multiplets, 14H) MeOCH₂CH₂ and BrCH₂CH₂; 10.03 (multiplet 4H) H at C-3, 6, 13, 16; 10.54
25 (multiplet, 4H) H at C-9, 10, 19, 20.

¹³C-NMR spectrum, CDCl₃, 75.5 MHz, δ (ppm) = 28.85 (MeOCH₂CH₂); 32.2, 32.9 (BrCH₂CH₂); 59.0 (H₃COCH₃); 73.9 (MeOCH₂CH₂); 110.5, 110.8, 110.9, 111.1, (C-9, 10, 19, 20); 123.5, 123.6, 123.9, 124.0 (C-3, 6, 13, 16); 133.5,

-54-

134.1, 134.2, 135.0 (C-4, 5, 14, 15); 140.58, 140.61,
140.9, 141.0, 142.9, 143.0, 144.5, (C-1, 2, 7, 8, 11, 12,
17, 18).

IR spectrum, CsI pellet, ν (cm^{-1}) = 2908, 2860 (CH)
5 1116 (C-O, ether), 1558, 1208, 937, 814.

Mass spectrum, EI, 75 eV: m/z = 592 (M^+ , 8%); 547
($M-\text{MeOCH}_2^+$, 2%); 511 ($M-\text{Br}^+$, 1%).

UV/VIS spectrum, CH_2Cl_2 , $\lambda_{\text{max}}[\text{nm}]$ (ϵ) = 243
(1.3×10^4); 370 (1.33×10^5); 832 (9.5×10^4); 564
10 (3.4×10^4); 546 (3.4×10^4); 603 (3.2×10^4); 635
(4.5×10^4).

2-Vinyl-7,12,17-tris(methoxyethyl)porphycene

All of the reagents and solvents used in the
following procedure were purged of oxygen by introducing
15 inert gas and all of the purification steps were carried
out, as in the reaction itself, under subdued light.

71 mg (0.12 mmol) of 2-bromoethyl-7,12,17-
tris(methoxyethyl)porphycene were dissolved in 50 mL of
absolute THF and treated with 1.8 g (1.86 ml, 12 mmol)
20 DBU under protective gas. The reaction solution was
stirred at 40°C for 90 minutes in the dark and
subsequently treated with 150 mL of absolute
dichloromethane. The mixture was poured into 100 mL of
5% hydrochloric acid, neutralized by shaking with a 2%
25 NaHCO_3 solution and then washed with water. The organic
phase obtained was separated off and evaporated under a
slight vacuum. The residue was chromatographed with
 CH_2Cl_2 on aluminum oxide (Brockmann, activity stage
II-III, column 3 x 10 cm). Following crystallization

-55-

from dichloro-methane and drying under oil pump vacuum, 52.6 mg (0.103 mmol) of the title compound were obtained from the single mobile blue fraction as needle-shaped crystals having a melting point of 119-120°C (with decomposition). Yield: 86%.

¹H-NMR spectrum, CDCl₃, 300 MHz, δ (ppm) = 3.05 (wide singlet) NH; 3.59, 3.95, 3.40 (3 singlets, 9H) H₃COCH₂; 4.28 (multiplet, 12H) MeOCH₂CH₂; 5.95 (doublet of doublets, 1H) CH=CH₂ cis; 6.60 (doublet of doublets, 1H) CH=CH₂ trans; 8.25 (doublet of doublets) CH=CH₂; 9.25, 9.26 (2 singlets, 3H) H at C-3, 6, 13, 16; 9.53 (singlet, 1H) H at C-3; 9.60 (AB system, 2H) H at C-9, 10; 9.69 (AB system, 2H) H at C-19, 20.

Mass spectrum, FAB, 75 eV; m/z = 510 (M⁺, 100%); 465 (M-MeOCH₂⁺, 23%).

IR spectrum, CsI pellet: ν (cm⁻¹) 2968, 2922, 2870 (CH), 1116 (C-O, ether), 1727, 1557, 1460, 1440, 801.

UV/VIS spectrum, CH₂Cl₂, λ_{max}[nm] (ε) = 373 (9.8 x 10⁴); 572 (2.6 x 10⁴); 613 (2.8 x 10⁴); 643 (3.3 x 10⁴).

Cleavage of 2,7,12,17-tetrakis(methoxyethyl)porphycene with BBr₃

The following procedure was carried out in a totally dry and oxygen free atmosphere with especially dried equipment and solvents.

70 μL (0.74 mmol) freshly distilled boron tribromide in 70 mL of dichloromethane were added during 60 minutes to a vigorously stirred solution of 542 mg (1 mmol)

-56-

tetrakis(methoxyethyl)porphycene in 150 mL of the same solvent at -30°C . Over the next 16 hours, the mixture was allowed to reach room temperature and 25 mL of a 8% aqueous solution of sodium hydrogen carbonate were added in one portion. The precipitate was filtrated off by the aid of Celite®, washed with sodium hydrogen carbonate solution and with water. The collected solids were extracted with methanol. The dichloromethane phase of the filtrate was extracted once with sodium hydrogen carbonate solution, twice with water and then combined with the methanol extract. The raw product mixture was preadsorbed by evaporating the organic phase on addition of 15g of alumina, and fractionated on a column of silica gel (2 x 60 cm) with dichloromethane/ethyl acetate/ethanol (100:10:1). After a forerun of 300-360 mg (50-65% depending on the extent of remaining water in solvent and equipment) unchanged starting material (tetrakis(methoxyethyl)porphycene), the 2-hydroxyethyl-7,12,17-tris(methoxyethyl)porphycene was eluted, which was then crystallized from toluene/hexane (1:1) to afford 17-18% (93 mg) of dark violet cubes melting at 131°C . By the use of chloroform/methanol (8:1) the isomeric bis(hydroxyethyl)bis(methoxyethyl)porphycenes as well as the tris(hydroxyethyl)mono(methoxyethyl)porphycene were eluted, which could be obtained in 10-12% (53 mg) and 4% (20 mg) yield, respectively.

2-Hydroxyethyl-7,12,17-tris(methoxyethyl)porphycene

Mass spectrum: (75 eV) m/z = 528 (80%; $\text{M}^{\cdot+}$); 497 (10%; $[\text{M}-\text{HOCH}_2]^+$); 483 (100%; $[\text{M}-\text{CH}_3\text{OCH}_2]^+$).

IR spectrum: (CsI): ν (cm^{-1}) 3462 (OH); 2921; 2869; 1561; 1460; 1405; 1215; 1115 (ether); 1017; 968; 884; 815.

-57-

UV/VIS spectrum: (dichloromethane) λ_{\max} [nm] (ϵ) =
370 (134000); 563 (33600); 602 (31200); 635 (43900).

2,12-Bis(hydroxyethyl)-7,17-bis(methoxyethyl)porphycene
(3rd fraction)

5 Mass spectrum: (75 eV) $m/z=514$ (90%; M^+); 483 (40%;
[M-HOCH₂·]⁺); 469 (100%; [M-CH₃OCH₂·]⁺).

¹H-NMR spectrum: (80 MHz, DMSO-d₆) δ (ppm) = 9.92
(singlet 4 H) H-9, H-10, H-19, H-20; 9.67 (singlet 4 H)
H-3, H-6, H-13, H-16; 5.12 (triplet 2H)-OH; 4.31
10 (multiplet 8 H) HOCH₂CH₂; 4.30 (singlet 8 H) H₃COCH₂CH₂;
3.50 (singlet 6 H) H₃CO-; 3.09 (wide singlet 2 H) NH.

IR spectrum: (CsI) ν (cm⁻¹) = 3424 (OH); 2912 (CH);
1870; 1560; 1459; 1415; 1388; 1216; 1108; 1064; 1013;
974; 889; 815; 625; 522.

15 UV/VIS spectrum: (DMSO) λ_{\max} [nm] = 251 (10000); 309
(12300); 371 (130000); 383 (92800); 530 (5600) (sh); 554
(22600) (sh); 562 (31900); 601 (32100); 632 (44400).

2,7-Bis(hydroxyethyl)-12,17-bis(methoxyethyl)porphycene
and 2,17-Bis(hydroxyethyl)-7,12-
20 bis(methoxyethyl)porphycene (4th fraction)

Mass spectrum: (75 eV) $m/z = 514$ (100%; M^+), 483
(40%; [M-HOCH₂·]⁺); 469 (100%; [M-CH₃OCH₂·]⁺).

¹H-NMR spectrum: (80 MHz, DMSO-d₆) δ (ppm) = 9.94
(singlet; 4 H) H-9, H-10, H-19, H-20; 9.68 (singlet; 4 H)
25 H-3, H-6, H-13, H-16; 5.73 (triplet; 2 H) -OH; 4.32
(multiplet; 8 H) HOCH₂CH₂; 4.31 (singlet; 8 H) H₃COCH₂CH₂;
3.50 (singlet; 6 H) -OCH₃; 3.09 (wide singlet; 2H) NH.

-58-

IR spectrum: (CsI) ν (cm^{-1}) = 2877 (CH); 1851; 1654; 1560; 1459; 1380; 1215; 1116; 1050; 1007; 887; 815; 626, 522.

UV/VIS spectrum: (DMSO) λ_{max} [nm] (ϵ) = 262 (8000);
5 310 (13000); 370 (136700); 383 (99000); 529 (5800) (sh);
551 (20800) (sh); 562 (34200); 601 (34600); 632 (48000).

2,7,12-Tris(hydroxyethyl)-17-methoxyethylporphycene

Mass spectrum: (75 eV) m/z = 500 (100%; M^+); 469
(70%; $[M-\text{HOCH}_2\cdot]^+$); 455 (75%; $[M-\text{CH}_3\text{OCH}_2\cdot]^+$).

10 IR spectrum: (CsI) ν (cm^{-1}) = 3331 (OH); 2870 (CH);
1870; 1560; 1460; 1371; 1220; 1046; 1008; 967; 886; 816;
626; 518.

UV/VIS spectrum: (DMSO) λ_{max} [nm] (ϵ) = 370 (134000);
382 (97000); 563 (33600); 602 (31200); 635 (43900).

15 2,7,12,17-Tetrakis(acetoxyethyl)porphycene

At -30°C and in an atmosphere of argon a solution of
108.4 mg (0.2 mmol) tetrakis(methoxyethyl)porphycene in
100 ml freshly dried dichloromethane was treated with 0.2
ml (2.1 mmol) pure boron tribromide. The solution was
20 stirred for 16 hours, quenched with 10 ml of dilute
ammonia and the precipitate was separated, washed with
dichloromethane, water and ether and dried in vacuo. The
crude 2,7,12,17-tetrakis(hydroxyethyl)porphycene was
dissolved in 25 ml pyridine and 5 ml acetic anhydride
25 stirred for 16 hours, evaporated and washed with water.
The residue was chromatographed with
dichloromethane/acetone (40:1) on silica gel (2 x 15 cm).
The second blue fraction contained 66 mg (50%) of the

-59-

title compound, which after recrystallization from benzene yielded violet needles melting at 188-189°C.

Mass spectrum: (75 eV) $m/z = 654$ (100%; M^+); 594 (35%; $[M-CH_3CO_2H]^+$); 581 (15%; $[M-CH_3CO_2CH_2\cdot]^+$); 534 (35%; $[M-2CH_3CO_2H]^+$).

1H -NMR spectrum: (80 MHz, $CDCl_3$) δ (ppm) = 9.67 (singlet; 4 H) H-9, H-10, H-19, H-20; 9.29 (singlet; 4H) H-3, H-6, H-13, H-16; 4.99 (AX-System $^3J(H-3a, H-3b) = 7.0$ Hz; 8 H) H-2a, H-7a, H-12a, H-17a; 4.34 (AX-System; 8H) H-2b, H-7b, H-12b, H-17b; 3.06 (wide singlet; 2H) NH; 2.14 (singlet, 12 H)-OAc.

IR spectrum: (CsI) ν (cm^{-1}) = 2955 (CH); 1733 (C=O); 1465; 1373; 1238; 1027 (C-O-ester); 965; 887; 806; 608; 517; 478.

UV/VIS spectrum: (dichloromethane) λ_{max} [nm] (ϵ) = 242 (14000); 369 (144000); 381 (103000); 564 (37000); 603 (35000); 635 (50000).

2,7,12,17-Tetrakis(hydroxyethyl)porphycene

32.7 mg (50 μ mol) Tetrakis(acetoxyethyl)porphycene were dissolved in 20 ml of dry tetrahydrofuran and 27 mg (0.5 mmol) sodium methoxide in 1 ml absolute methanol was added. The solution was allowed to stand without agitation while the pure tetrakis(hydroxyethyl)porphycene crystallized as small violet needles, which did not melt below 350°C (yield: 22 mg = 90%).

Mass spectrum: (75 eV) $m/z = 486$ (50%; M^+); 455 (80%; $[M-HOCH_2\cdot]^+$); 425 (30%; $[M-HOCH_2\cdot-OCH_2]^+$); 393 (25%; $[M-3HOCH_2\cdot]^+$).

-60-

IR spectrum: (CsI) ν (cm^{-1}) = 3328 (OH); 2895; 1463; 1372; 1220; 1043 (C-O); 880; 824; 396.

UV/VIS spectrum: (DMSO) λ_{max} [nm] (ϵ) = 372 (92400); 385 (83000); 570 (26200); 611 (20400); 647 (28300).

5 2-Chloroethyl-7,12,17-tris(methoxyethyl)porphycene

39.6 mg (75 μmol) 2-Hydroxyethyl-7,12,17-tris(methoxyethyl)porphycene were dissolved in 10 ml dimethylformamide (free from water and amine) and at 0°C 2 ml (27.5 mmol) of purified thionylchloride were added quickly. The solution warmed up and was directly poured into 100 ml dilute aqueous ammonia. The ammonia was extracted three times with trichloromethane, and the combined organic phases were then washed five times with water. After evaporation of the solvents the crude product was chromatographed on silica gel (2 x 80 cm) with trichloromethane/ethyl acetate (20:1; third blue fraction) and repeatedly recrystallized from dichloromethane/hexane and washed with ether and hexane. This procedure gave 4.1 mg (10%) chloroethylporphycene as tiny violet needles.

Mass spectrum: (70 eV) m/z = 546/548 (45%; M^+); 512 (90%; $[M+H-\text{Cl}]^+$); 501/503 (55%; $[M-\text{CH}_3\text{OCH}_2]^+$); 482 (15%; $[M-\text{CH}_3\text{O}-\text{Cl}]^+$); 467 (100%; $[M+H-\text{CH}_3\text{OCH}_2-\text{Cl}]^+$).

$^1\text{H-NMR}$ spectrum: (80 MHz, CDCl_3) δ (ppm) = 9.69 (singlet; 2 H) H-9, H-10; 9.63 (A_2B_2 -System; 2 H) H-19, H-20; 9.30 (singlet; 4 H) H-3, H-6, H-13, H-16; 4.42 (A_2B_2 -System; 4 H) H-2a, H-2b; 4.30 (singlet; 12 H) H-7a, H-7b, H-12a, H-12b, H-17a, H-17b; 3.60 (singlet; 12H)- OCH_3 ; 3.04 (broad singlet; 2 H)- NH .

-61-

IR spectrum: (CsI) ν (cm^{-1}) = 2984 (CH); 2903 (CH); 2808 (CH); 1121 (CO); 1846; 1560; 1458; 1420; 1391; 1263; 1214; 1066; 1019; 999; 969; 885; 819; 768; 692; 652; 553; 520; 497.

5 Methanesulfonate of 2-hydroxyethyl-7,12,17-
tris(methoxyethyl)porphycene

To a well-stirred solution of 52.8 mg (0.1 mmol) 2-hydroxyethyl-7,12,17-tris(methoxyethyl)porphycene in 100 ml dichloromethane and 9 ml pyridine were added during
10 one hour 8.1 ml (1.05 mmol) methanesulfonylchloride. After another hour the solution was poured onto 200 ml ice chilled water, the organic phase was separated and the aqueous phase was extracted twice with
15 dichloromethane. The combined organic phases were washed five times with water, the solvents were evaporated and the resulting blue oil was chromatographed thrice with trichloromethane/ethyl acetate (20:1) on silicagel (2 x 20cm). The content of the blue fractions were
20 recrystallized from dichloromethane/petrol to yield 55 mg (91%) of violet at 145°C melting crystals.

Mass spectrum: (75 eV) = m/z = 606 (2%; M^+); 562 (2%; $[M-CH_3OCH_2]^+$); 512 (10%; $[M-CH_2SO_2OCH_2]^+$); 467 (13%; $[M-CH_2SO_2O-CH_3OCH_2]^+$); 96 (80%; $CH_2SO_2O^+$); 79 (100%; $CH_3SO_2^+$).

25 IR spectrum: (CsI) ν (cm^{-1}) = 2978 (CH); 2869 (CH); 1357 (S=O); 1176 (S-O); 1114 (C-O); 1852; 1560; 1460; 1390; 1215; 1069; 1019; 973; 886; 816; 728; 649; 626; 556; 528; 508.

UV/VIS spectrum: (dichloromethane) λ_{max} [nm] (ϵ) =
30 242 (13400); 310 (12300); 369 (141000); 381 (104000); 532

-62-

(6200); 563 (35700); 603 (34100); 635 (48400).

2,3,4,6-Tetra-O-acetyl- β -D-galactopyranoside of 2-hydroxyethyl-7,12,17-tris(methoxyethyl)-porphycene

1 g of dry, freshly prepared silver carbonate and 1
5 g anhydrous sodium sulfate were added to a solution of
42.2 mg (0.08 mmol) 2-hydroxyethyl-7,12,17-
tris(methoxyethyl)porphycene and 658 mg 2,3,4,6-tetra-O-
acetyl- α -D-galactopyranosyl bromide in 50 ml of anhydrous
dichloromethane, and the mixture was stirred in the dark
10 with rigorous protection from moisture until thin layer
chromatography (tlc) (silica gel: dichlorometh-
ane/methanol, 20:1) indicated the disappearance of educt
(about 3 days). The reaction mixture was filtered
through a pad of Celite, and the inorganic solids were
15 washed with dichloromethane.

The filtrate and washings were combined, washed with
water, dried (sodium sulfate) and evaporated to give a
syrup which was fractionated by repeated
recrystallization from ethanol and finally ether/pentane.
20 This procedure yielded 6.9 mg (91%) of violet cubes
melting at 95°C.

Mass spectrum: (75 eV) m/z = 858 (100%; $M^{+\cdot}$); 814
(45%; $[M-CH_2CO]^{+\cdot}$); 511 (10%; $[M-C_6H_5O(OAc)_4O]^{+\cdot}$); 497
(10%; $[M-C_6H_5O(OAc)_4OCH_2]^{+\cdot}$); 375 (5%; $[M-C_6H_5O(OAc)_4O-$
25 $3CH_3OCH_2]^{+\cdot}$); 43 (80%; $CH_3CO^{+\cdot}$).

IR spectrum: (CsI) ν (cm^{-1}) = 2932 (CH); 1752;
(C=O); 1560; 1460; 1371; 1227; 1114; 1057; 966; 888; 814;
602; 536.

UV/VIS spectrum: (dichloromethane) λ_{max} [nm] (ϵ) =

-63-

242 (13100); 309 (12300); 369 (140000); 382 (100000); 532 (6000) (sh); 563 (35100); 602 (33500); 634 (47500).

β -D-Galactopyranoside of 2-hydroxyethyl-7,12,17-tris(methoxyethyl)porphycene

5 A solution of 17.2 mg (0.02 mmol) of the tetra-O-acetyl-compound (the mother liquors of the previous reaction could also be used) in 10 ml absolute tetrahydrofuran was treated with methanolic 0.05 M sodium methoxide (40 ml). The solution was kept for 2 hours at
10 room temperature, and 10 ml brine was added as well as 20 ml water. The organic phase was separated, the aqueous phase was extracted twice with 20 ml tetrahydrofuran and the combined extracts were washed thrice with dilute aqueous sodium chloride, dried (sodium
15 sulfate) and evaporated. The resulting residue was recrystallized from methanol to yield, after washing with water and cold ether small violet needles. m.p. = 124°C.

IR spectrum: (CsI) ν (cm⁻¹) = 3417 (OH); 2926 (CH); 1116 (C-O); 1561; 1461; 1383; 1218; 1018; 966; 888; 816;
20 626; 529.

UV/VIS spectrum: (ethanol) λ_{\max} [nm] (ϵ) = 373 (125800); 384 (103900) (sh); 572 (33500); 613 (37400); 645 (44700).

2-Cyanoethyl-7,12,17-tris(methoxyethyl)porphycene

25 202 mg (0.33 mmol) methanesulfonate or 197 mg (0.33 mmol) monobromoethyl porphycene were dissolved in 120 ml dry dimethylsulfoxide, and sodium cyanide (491 mg/10 mmol, dried for 16 hours at 110°C) was added. The suspension was stirred in the dark under an atmosphere of

-64-

argon until no more educt could be detected (tlc, silica gel: dichloromethane/acetone, 40:1, 3 hours). The dimethylsulfoxide was distilled off at an maximum bath temperature of 50°C, and the residue was taken up in 300 ml trichloromethane and washed with three 100 ml portions of water. After drying and evaporating the solvent the resulting product was fractionated on a column (3 x 30 cm) of silica gel eluting with dichloromethane/ethyl acetate (15:1). When the bromo compound is used as the starting material, protection from light is advisable.

The second blue fraction afforded after recrystallization from benzene 135 mg (75%) in the case of the methanesulfonate as starting material, and 104 mg (58%) in the case of reacting with the bromide, of blue violet needles of the nitrile; m.p. 166-167°C.

Mass spectrum: (70 eV) $m/z = 537$ (60%, M^+); 492 (100%, $[M-CH_3OH]^+$); 452 (4%, $[M-CH_3OCH_2]^+$); 447 (10%, $[M-2CH_3OCH_2]^+$); 407 (20%, $[M-NCCH_2-2CH_3OCH_2]^+$); 402 (15%, $[M-3CH_3OCH_2]^+$); 362 (10%, $[M-NCCH_2-CH_3OCH_2]^+$).

IR spectrum: (CsI) ν (cm^{-1}) = 2920; 2900 (CH); 2243 (CN); 1846; 1215; 1119 (C-O).

UV/VIS spectrum: (dichloromethane) λ_{max} [nm] (ϵ) = 370 (115000); 382 (86700); 564 (29000); 603 (28700); 635 (40400).

25 2,7,12-Tris(methoxyethyl)-17-methoxypropionylporphycene

A suspension of 53.7 mg (0.1 mmol) cyanoethylporphycene in 15 ml anhydrous methanol was saturated at 0°C with dry hydrogen chloride, so that the porphycene was completely dissolved and then stirred for

-65-

18 hours under protection from light. The now green solution was added to 100 g ice, neutralized with aqueous 5N sodium hydroxide and the precipitate was extracted with 200 ml dichloromethane. The solution was washed with water, dried, the solvent was evaporated and the product was purified by chromatography with 8:1 dichloromethane/acetone (1.5 x 10 cm) followed by recrystallization from benzene/hexane to afford 46.1 mg (81%) of violet needles, which melted at 137-139°C.

10 Mass spectrum: (70 eV) $m/z = 570$ (80%, M^+); 525 (100%, $[M-CH_3OCH_2]^+$); 480 (20%, $[M-2CH_3OCH_2]^+$).

IR spectrum: (CsI) ν (cm^{-1}) = 2870 (CH); 1849; 1736 (C=O); 1562; 1458; 1389; 1361; 1213; 1116 (C-O-ester); 1067 (C-O-ether); 1018; 998; 968; 882; 812; 710; 546; 15 364.

UV/VIS spectrum: (dichloromethane) λ_{max} [nm] (ϵ) = 370 (146000); 382 (105000); 563 (37200); 602 (35500); 634 (50700).

2-Carboxyethyl-7,12-17-tris(methoxyethyl)porphycene

20 A solution of 45.6 mg (80 μ mol), 2,7,12-tris(methoxyethyl)-17-methoxypropionylporphycene in 15 ml tetrahydrofuran was treated with 10 ml aqueous potassium hydroxide (2N) and stirred for 40 hours in the dark. The precipitate was separated from the nearly colorless solution, washed successively with water and 25 dichloromethane and taken up in a mixture of dichloromethane and 1N hydrochloric acid. The dichloromethane phase was twice washed with 0.5N hydrochloric acid, dried over magnesium sulfate, and 30 evaporated to give 29 mg (72%) of a microcrystalline powder.

Mass spectrum: (75 eV) $m/z = 556$ (60%; M^+); 511 (100%, $[M-CHO_2]^+$; $[M-CH_3OCH_2]^+$); 466 (25%, $[M-2CH_3OCH_2]^+$; $[M-CHO_2-CH_3OCH_2]^+$); 421 (25%, $[M-CO_2H-2CH_3OCH_2]^+$); 407 (10%, $[M-CH_2CO_2H-2CH_3OCH_2]^+$); 375 (10%,
5 $[M-CO_2H-3CH_3OCH_2]^+$).

1H -NMR spectrum: (80 MHz, CF_3COOD) δ (ppm) = 10.61 (wide singlet, 4 H) C-9, C-10, C-19, C-20; 10.09 (multiplet, 4 H) C-3, C-13, C-16; 4.67 (multiplet; 16 H) C-2a, C-2b, C-7a, C-7b, C-12a, C-12b, C-17a, C-17b; 3.91
10 (singlet, 9H) $-OCH_3$.


IR spectrum: (CsI) ν (cm^{-1}) = 3444 (OH); 2922 (C-H); 1710 (C=O); 1562; 1459; 1208; 1117 (C-O); 1016; 964; 883; 812.

UV/VIS spectrum: (dichloromethane/4%
15 trifluoroacetic acid) λ_{max} [nm] (ϵ) = 370 (120000); 382 (90000); 562 (30000); 602 (28000); 634 (41000).

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within
20 the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

Claims

1. A porphycene having the structure



wherein each R^1 is, independently,

- (a) $-(CH_2)_n-X$, where $n=1-4$, X is OR^2 and R^2 is C_{1-6} alkyl, aralkyl or aryl; CN ; OH ; $OCOR^2$; OSO_2R^2 ; NH_2 ; NHR^2 ; NR_2^2 ; SH ; SR^2 ; $S(O)_{1-2}R^2$; $COOH$; CO_2R^2 ; $C(O)NH_2$; $C(O)NHR^2$; $C(O)NR_2^2$; halogen; or CHO ;
- (b) $-(CH_2)_mCH=CH_2$ where m is 0-2; or
- (c) $-(CH_2)_n-O-G$ where G is a mono- or oligosaccharide;
- (d) $-(CH_2)_n-X$ where X is an amino acid, oligopeptide covalently bonded by an ether-, ester- or amine-bond or where X is $-Y-(CH_2)_n$ -porphycene², where porphycene² is a compound of the same said structure and Y is a direct bond; $-O-$; or $-CH=CH-$; or
- (e) where one, two or three of the substituents R^1 are C_{1-6} alkyl or aryl and the remaining substituents are as above under (a)-(d), and salts and metal complexes thereof.

2. The porphycene of Claim 1, wherein R^1 is $-(CH_2)_n-$ X and X is OR^2 .
3. The porphycene of Claim 1, wherein R^1 is $-(CH_2)_mCH=CH_2$.
- 5 4. The porphycene of Claim 1, wherein X is CN, COOH, CO_2R^2 or CHO.
5. The porphycene of Claim 1, wherein X is OH.
6. The porphycene of Claim 1, wherein X is NH_2 , NHR^2 , NR_2^2 , $C(O)NH_2$, $C(O)NHR^2$ or $C(O)NR_2^2$.
- 10 7. The porphycene of Claim 1, wherein X is halogen.
8. The porphycene of Claim 7, wherein said halogen is bromine.
9. The porphycene of Claim 1, wherein X is SH, SR^2 , OSO_2R^2 , or $S(O)_{1-2}R^2$.
- 15 10. The porphycene of Claim 1, wherein R^1 is O-G and G is a hexose or pentose.
11. The porphycene of Claim 2, wherein R^2 is methyl.
12. The porphycene of Claim 3, wherein $m = 0$.
- 20 13. The porphycene of Claim 1, wherein at least one R^1 is $-(CH_2)_n-OH$, $-(CH_2)_m-CH = CH_2$ or $-(CH_2)_n-O-G$.
14. The porphycene of Claim 13, wherein one R^1 is $-(CH_2)_n-OH$, $-(CH_2)_m-CH = CH_2$ or $-O-G$.

15. A pharmaceutical composition comprising the porphycene of Claim 1 and a pharmaceutically acceptable carrier or diluent.

16. The pharmaceutical composition of Claim 15,
5 wherein said composition is a solution of said porphycene in water, water-alcohol or dimethyl sulfoxide.

17. The pharmaceutical composition of Claim 15, wherein R^1 is $-(CH_2)_n-X$ and X is OR^2 .

18. The pharmaceutical composition of Claim 15,
10 wherein R^1 is $-(CH_2)_mCH=CH_2$.

19. The pharmaceutical composition of Claim 15, wherein R^1 is $-(CH_2)_n-X$ and X is halogen.

20. A method of photodynamic therapy, comprising the steps of:

15 administering to a mammal in need thereof, an effective amount of the porphycene of Claim 1, and

irradiating said mammal with light at a wavelength in the absorption spectrum of said porphycene.

21. The method of Claim 20, wherein said
20 administering is topical administration.

22. The method of Claim 20, wherein said administering is enteral, parenteral or intramuscular administration.

23. A method of purifying or decontaminating a
25 fluid, comprising

-70-

contacting said fluid with the porphycene of Claim
1, and

irradiating said fluid in contact with the
porphycene with light at a wavelength in the absorption
5 spectrum of said porphycene.

24. The method of Claim 23, wherein said fluid is
blood.

25. The method of Claim 23, wherein said porphycene
is covalently bonded to a stationary support

10 26. The compound 9-acetoxy-2,7,12,17-
tetrakis(methoxyethyl)porphycene.

27. A method of photodynamic therapy, comprising
the steps of:

administering to a mammal in need thereof, an
15 effective amount of the porphycene of Claim 26, and

irradiating said mammal with light at a wavelength
in the absorption spectrum of said porphycene.




FIG. 1

SUBSTITUTE SHEET

2/4





FIG. 2



4 / 4




FIG. 4

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/04624

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 31/40; C07D 209/56
US CL :540/465, 472; 514/410; 604/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 540/465, 472; 514/410; 604/20; 424/9

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,E See entire document.	US, A, 5,124,449 (FRANCK ET AL) 23 JUNE 1992	1-27
A,E	US, A, 5,132,101 (VOGEL ET AL) 21 JULY 1992 See entire document.	1-27

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search
10 OCTOBER 1992

Date of mailing of the international search report
24 NOV 1992

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT

Authorized officer

