



US 20180344840A1

(19) **United States**

(12) **Patent Application Publication**

Weiner et al.

(10) **Pub. No.: US 2018/0344840 A1**

(43) **Pub. Date: Dec. 6, 2018**

(54) **FILOVIRUS CONSENSUS ANTIGENS, NUCLEIC ACID CONSTRUCTS AND VACCINES MADE THEREFROM, AND METHODS OF USING SAME**

(71) Applicant: **The Trustees of the University of Pennsylvania**, Philadelphia, PA (US)

(72) Inventors: **David B. Weiner**, Merion, PA (US); **Devon Shedlock**, Philadelphia, PA (US)

(21) Appl. No.: **16/050,142**

(22) Filed: **Jul. 31, 2018**

Related U.S. Application Data

(60) Continuation of application No. 15/431,203, filed on Feb. 13, 2017, now Pat. No. 10,034,930, which is a division of application No. 14/391,952, filed on Oct. 10, 2014, now Pat. No. 9,597,388, filed as application No. PCT/US13/36413 on Apr. 12, 2013.

(60) Provisional application No. 61/623,428, filed on Apr. 12, 2012.

Publication Classification

(51) **Int. Cl.**
A61K 39/12 (2006.01)
C07K 14/005 (2006.01)
A61K 38/20 (2006.01)
A61N 1/32 (2006.01)
C12N 7/00 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
 CPC *A61K 39/12* (2013.01); *A61K 2039/572* (2013.01); *A61K 38/208* (2013.01); *A61K 38/2086* (2013.01); *A61N 1/327* (2013.01); *A61K 38/20* (2013.01); *C12N 7/00* (2013.01); *A61K 2039/57* (2013.01); *C07K 2319/02* (2013.01); *A61K 2039/545* (2013.01); *A61K 2039/53* (2013.01); *A61K 2039/70* (2013.01); *C12N 2760/14271* (2013.01); *C12N 2760/14122* (2013.01); *C12N 2760/14134* (2013.01); *C12N 2760/14171* (2013.01); *C12N 2760/14222* (2013.01); *C12N 2760/14234* (2013.01); *C07K 14/005* (2013.01)

(57) **ABSTRACT**

Nucleic acid molecules and compositions comprising one or more nucleic acid sequences that encode a consensus filovirus immunogen including a consensus Marburgvirus filovirus glycoprotein MARV GP immunogen, a consensus Ebolavirus Sudan filovirus glycoprotein SEBOV GP immunogen and a consensus Ebolavirus Zaire glycoprotein ZEBOV GP immunogen are disclosed. The coding sequences optionally include operable linked coding sequence that encode a signal peptide. Immunomodulatory methods and methods of inducing an immune response against filovirus, particularly Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire are disclosed. Method of preventing filovirus infection, particularly infection by Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire and methods of treating individuals infected with filovirus infection, particularly infection by Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire are disclosed. Consensus filovirus proteins are disclosed.

Specification includes a Sequence Listing.




Figure 1A




Figure 1B




Figure 1C




Figure 2A




Figure 2B




Figure 2C




Figure 2D




Figure 2E




Figure 2F




Figure 2G




Figure 2H




Figure 3A




Figure 3B




Figure 3C




Figure 4A




Figure 4B




Figure 4C




Figure 4D




Figure 5A




Figure 5B




Figure 5C




Figure 5D




Figure 6




Figure 7A




Figure 7B




Figure 8A




Figure 8B

**FILOVIRUS CONSENSUS ANTIGENS,
NUCLEIC ACID CONSTRUCTS AND
VACCINES MADE THEREFROM, AND
METHODS OF USING SAME**

FIELD OF THE INVENTION

[0001] The present invention relates to vaccines for inducing immune responses and preventing filovirus infection and/or treating individuals infected with filovirus, particularly infection by Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire. The present invention relates to consensus filovirus proteins, particularly Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire filovirus proteins and nucleic acid molecules which encode the same.

BACKGROUND OF THE INVENTION

[0002] The Filoviridae are non-segmented, single stranded RNA viruses which contain two divergent genera, Marburgvirus (MARV) and Ebolavirus (EBOV). Members from each can cause severe and highly lethal hemorrhagic fever disease to which there is no cure or licensed vaccine (Bradfute S. B., et al. (2011) Filovirus vaccines. *Hum Vaccin* 7: 701-711; Falzarano D., et al. (2011) Progress in filovirus vaccine development: evaluating the potential for clinical use. *Expert Rev Vaccines* 10: 63-77; Fields B. N., et al. (2007) Fields' virology. Philadelphia: Lippincott Williams & Wilkins 2 v. (xix, 3091, 1-3086 p.); Richardson J. S., et al. (2009) Enhanced protection against Ebola virus mediated by an improved adenovirus-based vaccine. *PLoS One* 4: e5308; and Towner J. S., et al. (2006) Marburgvirus genomics and association with a large hemorrhagic fever outbreak in Angola. *J Virol* 80: 6497-6516).

[0003] Due to lethality rates of up to 90% they have been described as "one of the most virulent viral diseases known to man" by the World Health Organization. The US Centers for Disease Control and Prevention has classified them as 'Category A Bioterrorism Agents' due in part to their potential threat to national security if weaponized (Burki T. K. (2011) USA focuses on Ebola vaccine but research gaps remain. *Lancet* 378: 389). These 'high priority' agents could in theory be easily transmitted, result in high mortality rates, cause major public health impact and panic, and require special action for public health preparedness (CDC (2011) Bioterrorism Agents/Diseases. Atlanta: Centers for Disease Control and Prevention).

[0004] The haemorrhagic fever diseases are acute infectious with no carrier state, although they are easily transmissible among humans and nonhuman primates by direct contact with contaminated bodily fluids, blood, and tissue (Feldmann H., et al. (2003) Ebola virus: from discovery to vaccine. *Nat Rev Immunol* 3: 677-685). During outbreak situations, reuse of medical equipment, health care facilities with limited resources, and untimely application of prevention measures escalate transmission of the disease, allowing amplification of infections in medical settings.

[0005] Since the natural reservoirs of these zoonotic pathogens are likely to be African bats and pigs (Kobinger G. P., et al. (2011) Replication, pathogenicity, shedding, and transmission of Zaire ebolavirus in pigs. *J Infect Dis* 204: 200-208), the latter possibly being more of an amplifying host, the manner in which the virus first appears at the start of an outbreak is thought to occur through human contact with an infected animal. Unpredictable endemic surfacing in

the Philippines, potentially Europe, and primarily Africa of this disease further constitutes a major public health concern (Outbreak news. (2009) Ebola Reston in pigs and humans, Philippines. *Wkly Epidemiol Rec* 84: 49-50).

[0006] Experiments have been performed to determine the capacity of the vaccine for inducing protective efficacy and broad CTL including experiments in rodent preclinical studies. (Fenimore P W, et al. (2012). Designing and testing broadly-protective filoviral vaccines optimized for cytotoxic T-lymphocyte epitope coverage. *PLoS ONE* 7: e44769; Hensley L E, et al. (2010). Demonstration of cross-protective vaccine immunity against an emerging pathogenic Ebolavirus Species. *PLoS Pathog* 6: e1000904; Zahn R, et al (2012). Ad35 and ad26 vaccine vectors induce potent and cross-reactive antibody and T-cell responses to multiple filovirus species. *PLoS ONE* 7: e44115; Geisbert T W, Feldmann H (2011). Recombinant vesicular stomatitis virus-based vaccines against Ebola and Marburg virus infections. *J Infect Dis* 204 Suppl 3: S1075-1081; and Grant-Klein R J, Van Deusen N M, Badger C V, Hannaman D, Dupuy L C, Schmaljohn C S (2012). A multiagent filovirus DNA vaccine delivered by intramuscular electroporation completely protects mice from ebola and Marburg virus challenge. *Hum Vaccin Immunother* 8; Grant-Klein R J, Altamura L A, Schmaljohn C S (2011). Progress in recombinant DNA-derived vaccines for Lassa virus and filoviruses. *Virus Res* 162: 148-161).

[0007] Vaccine-induced adaptive immune responses have been described in numerous preclinical animal models (Blaney J E, et al. (2011). Inactivated or live-attenuated bivalent vaccines that confer protection against rabies and Ebola viruses. *J Virol* 85: 10605-10616; Dowling W, et al. (2007). Influences of glycosylation on antigenicity, immunogenicity, and protective efficacy of ebola virus GP DNA vaccines. *J Virol* 81: 1821-1837; Jones S M, et al. (2005). Live attenuated recombinant vaccine protects nonhuman primates against Ebola and Marburg viruses. *Nat Med* 11: 786-790; Kalina W V, Warfield K L, Olinger G G, Bavari S (2009). Discovery of common marburgvirus protective epitopes in a BALB/c mouse model. *Virol J* 6: 132; Kobinger G P, et al. (2006). Chimpanzee adenovirus vaccine protects against Zaire Ebola virus. *Virology* 346: 394-401; Olinger G G, et al. (2005). Protective cytotoxic T-cell responses induced by Venezuelan equine encephalitis virus replicons expressing Ebola virus proteins. *J Virol* 79: 14189-14196; Rao M, Bray M, Alving C R, Jahrling P, Matyas G R (2002). Induction of immune responses in mice and monkeys to Ebola virus after immunization with liposome-encapsulated irradiated Ebola virus: protection in mice requires CD4(+) T cells. *J Virol* 76: 9176-9185; Rao M, Matyas G R, Grieder F, Anderson K, Jahrling P B, Alving C R (1999). Cytotoxic T lymphocytes to Ebola Zaire virus are induced in mice by immunization with liposomes containing lipid A. *Vaccine* 17: 2991-2998; Richardson J S, et al. (2009) Enhanced protection against Ebola virus mediated by an improved adenovirus-based vaccine. *PLoS One* 4: e5308; Vanderzanden L, et al (1998). DNA vaccines expressing either the GP or NP genes of Ebola virus protect mice from lethal challenge. *Virology* 246: 134-144; Warfield K L, et al. (2005). Induction of humoral and CD8+ T cell responses are required for protection against lethal Ebola virus infection. *J Immunol* 175: 1184-1191; Jones S M, et al. (2007). Assessment of a vesicular stomatitis virus-based vaccine by use of the mouse model of Ebola virus hemorrhagic fever. *J*

Infect Dis 196 Suppl2: S404-412 Grant-Klein R J, Van Deusen N M, Badger C V, Hannaman D, Dupuy L C, Schmaljohn C S (2012). A multiagent filovirus DNA vaccine delivered by intramuscular electroporation completely protects mice from ebola and Marburg virus challenge. *Hum Vaccin Immunother* 8.; Geisbert T W, et al. (2010). Vector choice determines immunogenicity and potency of genetic vaccines against Angola Marburg virus in nonhuman primates. *J Virol* 84: 10386-10394.) Viral vaccines have shown promise and include mainly the recombinant adenoviruses and vesicular stomatitis viruses. Non-infectious strategies such as recombinant DNA and Ag-coupled virus-like particle (VLP) vaccines have also demonstrated levels of pre-clinical efficacy and are generally considered to be safer than virus-based platforms. Virus-specific Abs, when applied passively, can be protective when applied either before or immediately after infection (Gupta M, Mahanty S, Bray M, Ahmed R, Rollin P E (2001). Passive transfer of antibodies protects immunocompetent and immunodeficient mice against lethal Ebola virus infection without complete inhibition of viral replication. *J Virol* 75: 4649-4654; Marzi A, et al. (2012). Protective efficacy of neutralizing monoclonal antibodies in a nonhuman primate model of Ebola hemorrhagic fever. *PLoS ONE* 7: e36192; Parren P W, Geisbert T W, Maruyama T, Jahrling P B, Burton D R (2002). Pre- and postexposure prophylaxis of Ebola virus infection in an animal model by passive transfer of a neutralizing human antibody. *J Virol* 76: 6408-6412; Qiu X, et al. (2012). Ebola GP-Specific Monoclonal Antibodies Protect Mice and Guinea Pigs from Lethal Ebola Virus Infection. *PLoS Negl Trop Dis* 6: e1575; Wilson J A, et al. (2000). Epitopes involved in antibody-mediated protection from Ebola virus. *Science* 287: 1664-1666; Sullivan N J, et al. (2011). CD8(+) cellular immunity mediates rAd5 vaccine protection against Ebola virus infection of nonhuman primates. *Nat Med* 17: 1128-1131; Bradfute S B, Warfield K L, Bavari S (2008). Functional CD8+ T cell responses in lethal Ebola virus infection. *J Immunol* 180: 4058-4066; Warfield K L, Olinger G G (2011). Protective role of cytotoxic T lymphocytes in filovirus hemorrhagic fever. *J Biomed Biotechnol* 2011: 984241). T cells have also been shown to provide protection based on studies performed in knockout mice, depletion studies in NHPs, and murine adoptive transfer studies where efficacy was greatly associated with the lytic function of adoptively-transferred CD8+ T cells. However, little detailed analysis of this response as driven by a protective vaccine has been reported.

[0008] Countermeasure development will ultimately require an improved understanding of protective immune correlates and how they are modulated during infection. This proves difficult when infected individuals who succumb to filoviral disease fail to mount an early immune response. These fast-moving hemorrhagic fever diseases result in immune dysregulation, as demonstrated by the lack of a virus-specific Ab response and a great reduction in gross T cell numbers, leading to uncontrolled viral replication and multi-organ infection and failure. Conversely, survivors of Ebola virus (EBOV) disease exhibit an early and transient IgM response, which is quickly followed by increasing levels of virus-specific IgG and CTL. These observations suggest that humoral and cell-mediated immune responses play a role in conferring protection against disease. These data are also supported by numerous preclinical efficacy studies demonstrating the contribution of vaccine-induced

adaptive immunity to protection against lethal challenge. However, mounting evidence has demonstrated a critical role for T cells in providing protection where efficacy was greatly associated with the functional phenotype of CD8+ T cells. While these recent studies highlight the importance of T cells in providing protection, their precise contributions remain uncharacterized and controversial. Furthermore, little detailed analysis of this response driven by a protective vaccine has been reported.

SUMMARY OF THE INVENTION

[0009] A composition comprising a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, and a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen is provided. The amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1, an amino acid sequence that is homologous to SEQ ID NO:1, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:1. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:1. Fragments of SEQ ID NO:1 or fragments of amino acid sequences that are homologous to SEQ ID NO:1 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2, an amino acid sequence that is homologous to SEQ ID NO:2, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:2. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:2. Fragments of SEQ ID NO:2 or fragments of amino acid sequences that are homologous to SEQ ID NO:2 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen may be SEQ ID NO:3 (MARV ANG), a fragment of SEQ ID NO:3, an amino acid sequence that is homologous to SEQ ID NO:3, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:3. Amino acid sequences that are homologous to SEQ ID NO:3 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:3. Fragments of SEQ ID NO:3 or fragments of amino acid sequences that are homologous to SEQ ID NO:3 are typically 600 or more, 637 or more, or 670 or more amino acids. The amino acid sequence may optionally comprise a leader sequences such as the IgE leader.

[0010] A composition comprising a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a Marburg marburgvirus first consensus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a Marburg marburgvirus second consensus envelope glycoprotein immunogen, and a nucleic acid sequence that encodes a Marburg marburgvirus third consensus envelope glycoprotein immunogen is also provided. The amino acid sequence

of the consensus Zaire ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1, an amino acid sequence that is homologous to SEQ ID NO:1, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:1. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:1. Fragments of SEQ ID NO:1 or fragments of amino acid sequences that are homologous to SEQ ID NO:1 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2, an amino acid sequence that is homologous to SEQ ID NO:2, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:2. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:2. Fragments of SEQ ID NO:2 or fragments of amino acid sequences that are homologous to SEQ ID NO:2 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence of the Marburg marburgvirus first consensus envelope glycoprotein immunogen may be SEQ ID NO:4 (MARV RAV), a fragment of SEQ ID NO:4, an amino acid sequence that is homologous to SEQ ID NO:4, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:4. Amino acid sequences that are homologous to SEQ ID NO:4 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:4. Fragments of SEQ ID NO:4 or fragments of amino acid sequences that are homologous to SEQ ID NO:4 are typically 600 or more, 637 or more, or 670 or more amino acids. The amino acid sequence of the Marburg marburgvirus second consensus envelope glycoprotein immunogen may be SEQ ID NO:5 (MARV OZO), a fragment of SEQ ID NO:5, an amino acid sequence that is homologous to SEQ ID NO:5, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:5. Amino acid sequences that are homologous to SEQ ID NO:5 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:4. Fragments of SEQ ID NO:5 or fragments of amino acid sequences that are homologous to SEQ ID NO:5 are typically 600 or more, 637 or more, or 670 or more amino acids. The amino acid sequence of the Marburg marburgvirus third consensus envelope glycoprotein immunogen may be SEQ ID NO:6 (MARV MUS), a fragment of SEQ ID NO:6, an amino acid sequence that is homologous to SEQ ID NO:6, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:6. Amino acid sequences that are homologous to SEQ ID NO:6 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:6. Fragments of SEQ ID NO:6 or fragments of amino acid sequences that are homologous to SEQ ID NO:6 are typically 600 or more, 637 or more, or 670 or more amino acids. The amino acid sequence may optionally comprise a leader sequences such as the IgE leader. In some embodiments, the composition further comprises a nucleic acid sequence that encodes the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen. The amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen may be SEQ ID NO:3 (MARV ANG), a fragment of SEQ ID NO:3,

an amino acid sequence that is homologous to SEQ ID NO:3, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:3. Amino acid sequences that are homologous to SEQ ID NO:3 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:3. Fragments of SEQ ID NO:3 or fragments of amino acid sequences that are homologous to SEQ ID NO:3 are typically 600 or more, 637 or more, or 670 or more amino acids. The amino acid sequence may optionally comprise a leader sequences such as the IgE leader.

[0011] Also provided is a composition comprising a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, and a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen. The amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1, an amino acid sequence that is homologous to SEQ ID NO:1, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:1. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:1. Fragments of SEQ ID NO:1 or fragments of amino acid sequences that are homologous to SEQ ID NO:1 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2, an amino acid sequence that is homologous to SEQ ID NO:2, or a fragment of an amino acid sequence that is homologous to SEQ ID NO:2. Amino acid sequences that are homologous to SEQ ID NO:1 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:2. Fragments of SEQ ID NO:2 or fragments of amino acid sequences that are homologous to SEQ ID NO:2 are typically 600 or more, 630 or more, or 660 or more amino acids. The amino acid sequence may optionally comprise a leader sequences such as the IgE leader.

[0012] A composition comprising a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, and a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen is provided. The nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:64, a fragment of SEQ ID NO:64, a nucleic acid sequence that is homologous to SEQ ID NO:64, or a fragment of a nucleotide sequence that is homologous to SEQ ID NO:64. Nucleic acid sequences that are homologous to SEQ ID NO:64 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:64. Fragments of SEQ ID NO:64 or fragments of amino acid sequences that are homologous to SEQ ID NO:64 typically encode 600 or more, 630 or more, or 660 or more amino acids of the consensus Zaire ebolavirus envelope glycoprotein immunogen encoded by SEQ ID NO:64. The nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen may be SEQ ID NO:65, a fragment of SEQ ID NO:65, a nucleic acid sequence that is homologous to SEQ ID NO:65, or a fragment of a nucleotide sequence

that is homologous to SEQ ID NO:65. Nucleic acid sequences that are homologous to SEQ ID NO:65 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:65. Fragments of SEQ ID NO:65 or fragments of amino acid sequences that are homologous to SEQ ID NO:65 typically encode 600 or more, 630 or more, or 660 or more amino acids of the consensus Sudan ebolavirus envelope glycoprotein immunogen encoded by SEQ ID NO:65. The nucleic acid sequence that encodes Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen may be SEQ ID NO:66, a fragment of SEQ ID NO:66, a nucleic acid sequence that is homologous to SEQ ID NO:66, or a fragment of a nucleotide sequence that is homologous to SEQ ID NO:66. Nucleic acid sequences that are homologous to SEQ ID NO:66 are typically 95% or more, 96% or more, 97% or more, 99% or more, or 99% or more, homologous to SEQ ID NO:66. Fragments of SEQ ID NO:66 or fragments of amino acid sequences that are homologous to SEQ ID NO:66 typically encode 600 or more, 630 or more, or 670 or more amino acids of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen encoded by SEQ ID NO:66. The nucleic acid sequences may optionally include sequences that encode leader sequences such as the IgE leader linked to the sequences encoding the immunogens.

[0013] Each of the different nucleic acid sequences may be on a single nucleic acid molecule, may each be on a separate nucleic acid molecules or various permutations. Nucleic acid molecules may be plasmids.

[0014] The composition may be formulated for delivery to an individual using electroporation.

[0015] The composition may further comprise nucleic acid sequences that encode one or more proteins selected from the group consisting of: IL-12, IL-15 and IL-28.

[0016] The composition may be used in methods of inducing an immune response against a filovirus. The filovirus may be selected from the group consisting of: Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire.

[0017] Methods of treating an individual who has been diagnosed with filovirus comprising administering a therapeutically effective amount of the composition to an individual are provided. The filovirus may be selected from the group consisting of: Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire.

[0018] Method of preventing filovirus infection in an individual are provided. The methods comprise administering a prophylactically effective amount of the composition to an individual. The filovirus may be selected from the group consisting of: Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire.

[0019] Compositions comprising two or more proteins selected from the group consisting of: a consensus Zaire ebolavirus envelope glycoprotein immunogen, a consensus Sudan ebolavirus envelope glycoprotein immunogen, a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, a first consensus Marburg marburgvirus envelope glycoprotein immunogen, a second consensus Marburg marburgvirus envelope glycoprotein immunogen and a third consensus Marburg marburgvirus envelope glycoprotein immunogen are provided.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIGS. 1A-1C refer to the polyvalent-vaccine construction strategy and expression experiments in Example 1. FIG. 1A shows phylogenetic trees for MGP (top), SGP (lower right), and ZGP (lower left). Significant support values are indicated (*) as verified by bootstrap analysis. A consensus strategy was adopted for the ZGP and SGP immunogens (CON VACCINE). Scale bars signify distance of amino acids per site and analyses were conducted using MEGA version 5 software. GP transgenes were commercially synthesized, genetically optimized, and subcloned into modified pVAX1 mammalian expression vectors. Antigen expression was analyzed following transfection of HEK 293T cells by Western immunoblotting and FACS. Western immunoblotting results are shown in FIG. 1B and FACS in FIG. 1C. For a comparative control, rVSV expressing MGP, SGP, or ZGP was run concurrently with each GP sample and species-specific anti-GP1 mAbs were used for detection. Size is indicated (kDa). For FACS, transfected cells were indirectly stained with mouse-derived GP-specific serum reagents followed by extensive washing and goat anti-mouse IgG and MHC class I. Western immunoblotting and FACS experiments were repeated at least three times with similar results. Significance for unrooted phylogenetic trees was determined by maximum-likelihood method and verified by bootstrap analysis and significant support values ($\geq 80\%$; 1,000 bootstrap replicates) were determined by MEGA version 5 software.

[0021] FIGS. 2A-2H show results from experiments in Example 1 in which complete protection against MARV and ZEBOV challenge was observed. Animal survival data is shown in FIG. 2A and FIG. 2E. FIG. 2A shows trivalent vaccinated animals survived post MARV challenge while control animals all died by day 10. FIG. 2E shows trivalent vaccinated animals survived post ZEBOV challenge while control animals all died by day 7. Data for % change in body weight for vaccinated and control animal are displayed in FIG. 2B for vaccinated challenged with MARV. The y axis indicates change in body weight as shown in FIG. 2F. The light solid line is for Trivalent vaccinated animals. The light dashed line is for TriAVE, the average results of the Trivalent vaccinated animals. The dark solid line is for control animals. The dark dashed line is for Control AVE, the average result for the control animals. The light solid lines and light dashed lines remain steady on the graph in the days post challenge showing no significant weight loss among vaccinated animals. The dark solid lines and dark dashed lines decline on the graph from days 0-9 post challenge ending with the dagger denote animals that succumbed to disease by day 10. Data for % change in body weight for vaccinated and control animal are displayed in FIG. 2F for vaccinated challenged with ZEBOV. The y axis shows change in body weight as a percent. The light solid line is for Trivalent vaccinated animals. The light dashed line is for TriAVE, the average results of the Trivalent vaccinated animals. The dark solid line is for control animals. The dark dashed line is for Control AVE, the average result for the control animals. The light solid lines and light dashed lines remain steady on the graph in the days post challenge showing no significant weight loss among vaccinated animals. The dark solid lines and dark dashed lines decline on the graph from days 0-6 post challenge ending with the dagger denote animals that succumbed to disease before day 8. (n=3 for gpMARV and n=6 for gpZEBOV). Binding Abs

(FIG. 2C and FIG. 2G) and NAbs (FIG. 2D and FIG. 2H) were measured in serum from vaccinated animals before (Pre) and after the first (1x) and second (2x) immunizations. Analysis was conducted on pooled serum (FIG. 2H). * $p < 0.1$; *** $p < 0.001$; **** $p < 0.0001$.

[0022] FIGS. 3A-3C show results from Example 1 demonstrating induction of neutralizing Abs. B cell responses were assessed in mice ($n=5$ /group) 20 days following each of two vaccinations, spaced three weeks between injections with 40 μg of E-DNA vaccination. FIG. 3A shows serum GP-specific IgG responses from vaccinated (solid lines) mice or pre-bled (dotted lines) mice were measured by ELISA. The data is summarized in FIG. 3B. All responses from pEBOS- and pEBOZ-immunized animals were measured against sucrose-purified ZGP since SGP was not available for this study. IgG responses from pMARV-immunized mice were measured against MARV-Ozolin GP or with negative control sucrose-purified Nipah G protein. Neutralization activity of serum samples was measured against ZEBOV-EGFP, SUDV-Boniface and MARV-Angola in a BSL-4 facility and NAb titers are shown in FIG. 3C. NAbs against SUDV-Boniface were assayed based on cytopathic effect (CPE) on CV-1 cells and those against MARV-Angola were assayed using an immunofluorescent assay. Averages are shown in FIG. 3B and FIG. 3C and error bars represent SEM. Group analyses were completed by matched, two-tailed, unpaired t test. Experiments were repeated at least two times with similar results and * $p < 0.1$; ** $p < 0.01$; *** $p < 0.001$.

[0023] FIGS. 4A-4D shows refer to broad T cells responses generated by vaccination. In FIG. 4A H-2^b (light bars) and H-2^d (dark bars) mice ($n=5$ /group) were immunized twice with either pMARV, pEBOS or pEBOZ DNA, and IFN γ responses were measured by IFN γ ELISPOT assay. Splenocytes harvested 8 days after the second immunization were incubated in the presence of individual GP peptides (15-mers overlapping by 9 amino acids) and results are shown in stacked bar graphs. Epitope-containing peptides were identified (≥ 10 AVE spots AND $\geq 80\%$ response rate), confirmed by flow cytometry and characterized in the population of total activated IFN γ + and CD44+CD4+ and/or CD8+ T cells (Tables 1-6), and peptide numbers of positive inducers are indicated above the bars. Peptides containing CD4+ epitopes alone, CD8+ epitopes alone (*), and dual CD4+ and CD8+ epitopes (***) are numbered. Putative shared and/or partial epitopes were explored for contiguous positive peptide responses (Tables 1-6). FIG. 4B shows amino acid similarity plots comparing GP sequences from MARV, SUDV, and ZEBOV viruses displayed in FIG. 1A. FIG. 4C is a diagram showing putative domains within the ZEBOV GP (GenBank #VGP_EBOZM). SP, signal peptide; RB, receptor binding; MUC, mucin-like region; FC, furin cleavage site; TM, transmembrane region. In FIG. 4D, total subdominant (darker shade) and immunodominant (lighter shade) T cell epitopic responses are displayed as a percentage of the total IFN γ response generated by each vaccine. Experiments were repeated at least two times with similar results.

[0024] FIGS. 5A-5D show data from experiment assessing protective 'single-dose' vaccination induced neutralizing Abs and CTL. H-2^k mice ($n=10$ /group) were vaccinated once i.m. with pEBOZ E-DNA and then challenged 28 days later with 1,000 LD₅₀ of mZEBOV in a BSL-4 facility. Mice were weighed daily and monitored for disease progression.

Animal survival data in FIG. 5A. Vaccinated animals survived challenge while control animals died by day 7. FIG. 5B shows data for % change in body weight in challenged animals. Data from immunized animals is shown as a solid light line; the average data for immunized animals is shown as a dashed light line. Data from control animals is shown as a solid dark line; the average data for control animals is shown as a dashed dark line. The light solid lines and light dashed lines remain steady within the range of about 85%-120% on the graph in the days post challenge showing no significant weight loss among vaccinated animals. The dark solid lines and dark dashed lines decline on the graph from days 0-6 post challenge ending with the dagger denote animals that succumbed to disease by day 7. NAbs measured prior to challenge; the data shown in FIG. 5C. T cell responses after a single pEBOZ immunization as measured by FACS are summarized as AVE % of total CD44+/IFN γ + CD4+(dark) or CD8+(light) cells in FIG. 5D. T_H1-type effector markers were assessed (TNF and T-bet) and data for CD44+/IFN γ +CD4+ and CD8+ T cells were compared with total T cell data which was as follows: For Total Cells: TNF 2.9 \pm 0.8, Tbet 13.0 \pm 1.1. For CD4+/CD44+/IFN γ + Cells: TNF 61.4 \pm 3.1, Tbet 72.6 \pm 2.0. For CD8+/CD44+/IFN γ + Cells: TNF 33.0 \pm 3.3, Tbet 992.1 \pm 1.4 (* $p < 0.1$; *** $p < 0.001$; **** $p < 0.0001$). Group analyses were completed by matched, two-tailed, unpaired t test and survival curves were analyzed by log-rank (Mantel-Cox) test. Experiments were performed twice with similar results and error bars represent SEM.

[0025] FIG. 6 shows a GP-specific T cell gating disclosed in Example 1.

[0026] FIGS. 7A and 7B show that vaccination experiments in Example 1 generated robust T cells.

[0027] FIGS. 8A and 8B show T cell induction by 'single-dose' vaccination disclosed in Example 1.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0028] In one aspect of the invention, it is desired that the consensus antigen provides for improved transcription and translation, including having one or more of the following: low GC content leader sequence to increase transcription; mRNA stability and codon optimization; eliminating to the extent possible cis-acting sequence motifs (i.e., internal TATA-boxes).

[0029] In some aspects of the invention, it is desired to generate a consensus antigen that generates a broad immune response across multiple strains, including having one or more of the following: incorporate all available full-length sequences; computer generate sequences that utilize the most commonly occurring amino acid at each position; and increase cross-reactivity between strains.

[0030] Diversity among the Filoviridae is relatively high. Intensive efforts have been aimed at developing a universal and broadly-reactive filovirus vaccine that would ideally provide protection against multiple species responsible for the highest human case-fatality rates. However, this proves difficult due to the relative high level of diversity among the Filovirida. The EBOV are currently classified into five distinct species, Zaire ebolavirus (ZEBOV), Sudan ebolavirus (SUDV), Reston ebolavirus (RESTV), Bundibugyo ebolavirus (BDBV) and Taï Forest ebolavirus (TAFV; formerly Cote d'Ivoire ebolavirus), the first two responsible for the highest lethality rates and the most likely candidates for

weaponization. Diversity is lower among the Marburg viruses (MARV) of which can also be up to 90% lethal. Currently, there is only one classified species, Marburg marburgvirus (formerly Lake Victoria marburgvirus), although a recent amendment proposes that it contain two viruses including the Ravn virus (RAVV). Adding to the complexity for polyvalent-vaccine development, the MARV and EBOV are highly divergent, in which there exists about 67% divergence at the nucleotide level. Furthermore, phylogenetic diversity among the filoviral GP is also very high (82% overall). These allude to the potential of the filoviruses to evolve, as demonstrated by the recent emergence of BDBV in 2007. Therefore, due to relative divergence among the Filoviridae, we hypothesized that development of an effective polyvalent-filovirus vaccine will likely require a cocktail of immunogenic components.

[0031] A synthetic polyvalent-filovirus DNA vaccine against Marburg marburgvirus (MARV), Zaire ebolavirus (ZEBOV), and Sudan ebolavirus (SUDV) was developed. The novel polyvalent-filovirus vaccine comprised by three DNA plasmids encoding the envelope glycoprotein (GP) genes of Marburg marburgvirus (MARV), Sudan ebolavirus (SUDV) or Zaire ebolavirus (ZEBOV), adopting the multi-agent approach. As a filoviral vaccine candidate, an enhanced DNA (DNA)-based platform exhibits many advantages given recent advances in genetic optimization and delivery techniques (Bagarazzi M L, et al. (2012). Immunotherapy Against HPV16/18 Generates Potent TH1 and Cytotoxic Cellular Immune Responses. *Sci Transl Med* 4: 155ra138; Kee S T, Gehl J, W. LE (2011). *Clinical Aspects of Electroporation*, Springer, New York, N.Y.; Hirao L A, et al. (2011). Multivalent smallpox DNA vaccine delivered by intradermal electroporation drives protective immunity in nonhuman primates against lethal monkeypox challenge. *J Infect Dis* 203: 95-102). As such, each GP was genetically-optimized, subcloned into modified mammalian expression vectors, and then delivered using in vivo electroporation (EP).

[0032] Preclinical efficacy studies were performed in guinea pigs and mice using rodent-adapted viruses, while murine T cell responses were extensively analyzed using a novel modified assay described herein. T cell responses were extensively analyzed including the use of a novel method for epitope identification and characterization described herein. This model provides an important preclinical tool for studying protective immune correlates that could be applied to existing platforms.

[0033] Vaccination in preclinical rodent studies was highly potent, elicited robust neutralizing antibodies (NAbs) and CTL expressing T_H1 -type markers, and completely protected against MARV and ZEBOV challenge. Comprehensive T cell analysis as extensively analyzed using a novel modified assay described herein (Shedlock D J, et al. (2012). Vaccination with synthetic constructs expressing cytomegalovirus immunogens is highly T cell immunogenic in mice. *Hum Vaccin Immunother* 8: 1668-1681) revealed cytotoxic T lymphocytes of great magnitude, epitopic breadth, and T_H1 -type marker expression. In total, 52 novel T cell epitopes from two different mouse genetic backgrounds were identified (19 of 20 MARV epitopes, 15 of 16 SUDV, and 18 of 22 ZEBOV) and occurred primarily in highly conserved regions of their respective glycoproteins (GPs). These data represent the most comprehensive report of preclinical glycoprotein epitopes to date.

[0034] In developing a strategy to provide protection against multiple species responsible for the highest human case-fatality rates, we focused on MARV, SUDV, and ZEBOV. Due to their relative divergence, we hypothesized that development of a polyvalent-filovirus vaccine would require a cocktail of components that can be quickly and easily adapted in response to future outbreak strains and/or species. While overall diversity among the EBOV is about 33%, amino acid identity increases substantially when SUDV and ZEBOV are analyzed separately (~94% identity within each species). Therefore, as shown in FIG. 1A, a two component strategy for coverage of the most lethal EBOV, one plasmid GP vaccine for SUDV and another for ZEBOV was designed. Since GP diversity among each species was relatively low (5.6% for SUDV and 7.1% for ZEBOV), consensus immunogens were developed increase inter-species coverage, a strategy shown previously to enhance protection among divergent strains of influenza and HIV. These GP sequences were consensus for all reported outbreak sequences (GenBank) as determined by alignment using Vector NTI software (Invitrogen, CA, USA). Non-consensus residues, 4 amino acids each in SUDV (95, 203, 261, and 472) and ZEBOV (314, 377, 430, and 440), were weighted towards Gulu and Mbomo/Mbanza, respectively. Gulu was chosen since it was responsible for the highest human case-fatality rate of any Filoviridae outbreak (n=425), while Mbomo/Mbanza was chosen since they were the most recent and lethal outbreaks with published sequence data. The consensus GP for SUDV (SUDV CON VACCINE) and ZEBOV (ZEBOV CON VACCINE) were phylogenetically intermediary their parentally aligned strains.

[0035] Identification of proteins in FIG. 1A are as follows: MARV Durba (05DRC99) '99: ABE27085; Uganda (OIUga07) '07: ACT79229; Durba (07DRC99) '99: ABE27078; Ozolin '75: VGP_MABVO; Musoke '80: VGP_MABVM; Popp '67: VGP_MABVP; Leiden '08: AEW11937; Angola '05: VGP_MABVA; Ravn '87: VGP_MABVR; Durba (09DRC99) '99: ABE27092; Uganda (02Uga07) '07: ACT79201. SUDV: Boniface '76: VGP_EBOSB; Maleo '79: VGP_EBOSM; Yambio '04: ABY75325; Gulu '00: VGP_EBOSU. ZEBOV: Booue '96: AAL25818; Mayibout '96: AEK25495; Mekouka '94: AAC57989, VGP_EBOG4; Kikwit '95: VGP_EBOZ5; Yambuku (Ekron) '76: VGP_EBOEC; Yambuku (Mayinga) '76: VGP_EBOZM; Kasai '08: AER59712; Kassai '07: AER59718; Etoumbi '05: ABW34742; Mbomo/Mbandza '03: ABW34743.

[0036] A sequence listing provided herewith contains a list of 66 sequences including the following

[0037] SEQ ID NO:1 is the amino acid sequence of ZEBOV CON, which is a consensus Zaire ebolavirus envelope glycoprotein immunogen.

[0038] SEQ ID NO:2 is the amino acid sequence of SUDV CON, which is a consensus Sudan ebolavirus envelope glycoprotein immunogen.

[0039] SEQ ID NO:3 is the amino acid sequence of MARV or MARV ANG, which the amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein and a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen.

[0040] SEQ ID NO:4 is the amino acid sequence of MARV CON1, which is the first consensus Marburg marburgvirus envelope glycoprotein immunogen.

[0041] SEQ ID NO:5 is the amino acid sequence of MARV CON2, which is the second consensus Marburg marburgvirus envelope glycoprotein immunogen.

[0042] SEQ ID NO:6 is the amino acid sequence of MARV CON3, which is the third consensus Marburg marburgvirus envelope glycoprotein immunogen.

[0043] SEQ ID NOS:7-25 are peptides derived from MARV ANG.

[0044] SEQ ID NO:26-41 are peptides derived from SUDV CON.

[0045] SEQ ID NO: 42-62 are peptides derived from ZEBOV CON.

[0046] SEQ ID NO:63 is the sequence of the IgE signal peptide: MDWTWILFLVAAATRVHS.

[0047] SEQ ID NO:64 is the nucleotide sequence insert in plasmid pEBOZ which encodes consensus Zaire ebolavirus envelope glycoprotein immunogen.

[0048] SEQ ID NO:65 is the nucleotide sequence insert in plasmid pEBOS which encodes consensus Sudan ebolavirus envelope glycoprotein immunogen.

[0049] SEQ ID NO:66 is the nucleotide sequence insert in plasmid pMARZ ANG which encodes the Marburg marburgvirus Angola 2005 envelope glycoprotein.

[0050] In some embodiments, the strategy employs coding sequences for three filovirus immunogens: MARV, SUDV, and ZEBOV. MARV immunogen is the glycoprotein of the Angola 2005 isolate. For SUDV and ZEBOV, consensus glycoprotein sequences were designed.

[0051] In some embodiments, the strategy employs coding sequences for five filovirus immunogens. Three MARV immunogens are provided. Consensus glycoprotein Ozolin, Musoke, or Ravn derived from three clusters, were designed. These three MARV immunogens are targets for immune responses together the SUDV and ZEBOV consensus glycoprotein sequences that were designed.

[0052] In some embodiments, the strategy employs coding sequences for six filovirus immunogens. Four MARV immunogens are provided: three consensus glycoproteins derived from three clusters were designed. These three MARV immunogens are targets for immune responses together the he SUDV and ZEBOV consensus glycoprotein sequences that were designed and the MARV immunogen is the glycoprotein of the Angola 2005 isolate.

[0053] As a candidate for filoviral vaccines, DNA vaccines exhibit a multitude of advantages including rapid and inexpensive up-scale production, stability at room temperature, and ease of transport, all of which further enhance this platform from an economic and geographic perspective. Due to the synthetic nature of the plasmids, Ag sequences can be quickly and easily modified in response to newly emergent species and/or expanded to include additional vaccine components and/or regimen for rapid response during outbreak settings. For example, the MARV strategies herein can be easily expanded for greater coverage by the co-administration of additional plasmids encoding consensus MARV GP (MGP) immunogens for other phylogenetic clusters.

[0054] While ‘first-generation’ DNA vaccines were poorly immunogenic, recent technological advances have dramatically improved their immunogenicity in clinical trials. Optimization of plasmid DNA vectors and their encoded Ag genes have led to increases in in vivo immunogenicity. Cellular uptake and subsequent Ag expression are substantially amplified when highly-concentrated plasmid vaccine formulations are administered with in vivo electroporation,

a technology that uses brief square-wave electric pulses within the vaccination site to drive plasmids into transiently permeabilized cells. In theory, a cocktail of DNA plasmids could be assembled for directing a highly-specialized immune response against any number of variable Ags. Immunity can be further directed by co-delivery with plasmid molecular adjuvants encoding species-specific cytokine genes as well as ‘consensus-engineering’ of the Ag amino acid sequences to help bias vaccine-induced immunity towards particular strains. This strategy has been shown to enhance protection among divergent strains of influenza virus and HIV. Due in parts to these technological advancements, immunization regimens including these DNA vaccines are highly versatile and extremely customizable.

1. Definitions

[0055] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise.

[0056] For recitation of numeric ranges herein, each intervening number there between with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the numbers 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, and 7.0 are explicitly contemplated.

[0057] a. Adjuvant

[0058] “Adjuvant” as used herein may mean any molecule added to the DNA plasmid vaccines described herein to enhance antigenicity of the one or more consensus filovirus immunogens encoded by the DNA plasmids and encoding nucleic acid sequences described hereinafter.

[0059] b. Antibody

[0060] “Antibody” may mean an antibody of classes IgG, IgM, IgA, IgD or IgE, or fragments, fragments or derivatives thereof, including Fab, F(ab')₂, Fd, and single chain antibodies, diabodies, bispecific antibodies, bifunctional antibodies and derivatives thereof. The antibody may be an antibody isolated from the serum sample of mammal, a polyclonal antibody, affinity purified antibody, or mixtures thereof which exhibits sufficient binding specificity to a desired epitope or a sequence derived therefrom.

[0061] c. Coding Sequence

[0062] “Coding sequence” or “encoding nucleic acid” as used herein may mean refers to the nucleic acid (RNA or DNA molecule) that comprise a nucleotide sequence which encodes a protein. The coding sequence may further include initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of an individual or mammal to whom the nucleic acid is administered. In some embodiments, the coding sequence may optionally further comprise a start codon that encodes an N terminal methionine or a signal peptide such as an IgE or IgG signal peptide.

[0063] d. Complement

[0064] “Complement” or “complementary” as used herein may mean a nucleic acid may mean Watson-Crick (e.g., A-T/U and C-G) or Hoogsteen base pairing between nucleotides or nucleotide analogs of nucleic acid molecules.

[0065] e. Consensus or Consensus Sequence

[0066] “Consensus” or “consensus sequence” as used herein may mean a synthetic nucleic acid sequence, or corresponding polypeptide sequence, constructed based on analysis of an alignment of multiple subtypes of a particular filovirus antigen, that can be used to induce broad immunity against multiple subtypes or serotypes of a particular filovirus antigen.

[0067] Consensus Zaire ebolavirus envelope glycoprotein immunogen refers to SEQ ID NO:1, fragments of SEQ ID NO:1, variants of SEQ ID NO:1 and fragment of variants of SEQ ID NO:1. (ZEBOV or ZEBOV CON or ZEBOV CON VACCINE). Plasmids comprising coding sequences of SEQ ID NO:1 may be referred to as pZEBOV or pEBOZ. Coding sequences for consensus Zaire ebolavirus envelope glycoprotein immunogen include SEQ ID NO:64, fragments of SEQ ID NO:64, variants of SEQ ID NO:64 and fragment of variants of SEQ ID NO:64. Plasmid pEBOZ comprises SEQ ID NO:64.

[0068] Consensus Sudan ebolavirus envelope glycoprotein immunogen refers to SEQ ID NO:2, fragments of SEQ ID NO:2, variants of SEQ ID NO:2 and fragment of variants of SEQ ID NO:2. (SUDV or SUDV CON or SUDV CON VACCINE) Plasmids comprising coding sequences of SEQ ID NO:2 may be referred to as pSUDV or pEBOS. Coding sequences for consensus Sudan ebolavirus envelope glycoprotein immunogen include SEQ ID NO:65, fragments of SEQ ID NO:65, variants of SEQ ID NO:65 and fragment of variants of SEQ ID NO:65. Plasmid pEBOS comprises SEQ ID NO:65.

[0069] Marburg marburgvirus Angola 2005 envelope glycoprotein is not a consensus but a protein sequence derived from an isolate. It has sequence SEQ ID NO:3. Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen refers to SEQ ID NO:3, fragments of SEQ ID NO:3, variants of SEQ ID NO:3 and fragment of variants of SEQ ID NO:3. (MARV or MARV ANG or MARV ANG or MARV ANG VACCINE) Plasmids comprising coding sequences of SEQ ID NO:3 may be referred to as pMARV or pMARV-ANG. Coding sequences for Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen include SEQ ID NO:66, fragments of SEQ ID NO:66, variants of SEQ ID NO:66 and fragment of variants of SEQ ID NO:66. Plasmid pMARV ANG comprises SEQ ID NO:66.

[0070] The first consensus Marburg marburgvirus envelope glycoprotein immunogen refers to SEQ ID NO:4, fragments of SEQ ID NO:4, variants of SEQ ID NO:4 and fragment of variants of SEQ ID NO:4. SEQ ID NO:4 is a Marburg marburgvirus consensus sequence from the Ravn cluster consensus (Ravn, Durba (09DRC99) and Uganda (02Uga07Y)). (MARV CON1 or MARV-RAV CON or MARV-RAV CON VACCINE) Plasmids comprising coding sequences of SEQ ID NO:4 may be referred to as pMARV-RAV.

[0071] The second consensus Marburg marburgvirus envelope glycoprotein immunogen refers to SEQ ID NO:5, fragments of SEQ ID NO:5, variants of SEQ ID NO:5 and fragment of variants of SEQ ID NO:5. SEQ ID NO:5 is a Marburg marburgvirus consensus sequence from the Ozolin cluster consensus (Ozolin, Uganda (01Uga07), and Durba (05 and 07DRC99)). (MARV CON2 or MARV-OZO CON

or MARV-OZO CON VACCINE) Plasmids comprising coding sequences of SEQ ID NO:5 may be referred to as pMARV-OZO.

[0072] The third consensus Marburg marburgvirus envelope glycoprotein immunogen refers to SEQ ID NO:6, fragments of SEQ ID NO:6, variants of SEQ ID NO:6 and fragment of variants of SEQ ID NO:6. SEQ ID NO:6 is a Marburg marburgvirus consensus sequence from the Musoke cluster consensus (Musoke, Popp, and Leiden). (MARV CON1 or MARV-MUS CON or MARV-MUS CON VACCINE) Plasmids comprising coding sequences of SEQ ID NO:6 may be referred to as pMARV-MUS.

[0073] f. Constant Current

[0074] “Constant current” as used herein to define a current that is received or experienced by a tissue, or cells defining said tissue, over the duration of an electrical pulse delivered to same tissue. The electrical pulse is delivered from the electroporation devices described herein. This current remains at a constant amperage in said tissue over the life of an electrical pulse because the electroporation device provided herein has a feedback element, preferably having instantaneous feedback. The feedback element can measure the resistance of the tissue (or cells) throughout the duration of the pulse and cause the electroporation device to alter its electrical energy output (e.g., increase voltage) so current in same tissue remains constant throughout the electrical pulse (on the order of microseconds), and from pulse to pulse. In some embodiments, the feedback element comprises a controller.

[0075] g. Current Feedback or Feedback

[0076] “Current feedback” or “feedback” as used herein may be used interchangeably and may mean the active response of the provided electroporation devices, which comprises measuring the current in tissue between electrodes and altering the energy output delivered by the EP device accordingly in order to maintain the current at a constant level. This constant level is preset by a user prior to initiation of a pulse sequence or electrical treatment. The feedback may be accomplished by the electroporation component, e.g., controller, of the electroporation device, as the electrical circuit therein is able to continuously monitor the current in tissue between electrodes and compare that monitored current (or current within tissue) to a preset current and continuously make energy-output adjustments to maintain the monitored current at preset levels. The feedback loop may be instantaneous as it is an analog closed-loop feedback.

[0077] h. Decentralized Current

[0078] “Decentralized current” as used herein may mean the pattern of electrical currents delivered from the various needle electrode arrays of the electroporation devices described herein, wherein the patterns minimize, or preferably eliminate, the occurrence of electroporation related heat stress on any area of tissue being electroporated.

[0079] i. Electroporation

[0080] “Electroporation,” “electro-permeabilization,” or “electro-kinetic enhancement” (“EP”) as used interchangeably herein may refer to the use of a transmembrane electric field pulse to induce microscopic pathways (pores) in a bio-membrane; their presence allows biomolecules such as plasmids, oligonucleotides, siRNA, drugs, ions, and water to pass from one side of the cellular membrane to the other.

[0081] j. Feedback Mechanism

[0082] “Feedback mechanism” as used herein may refer to a process performed by either software or hardware (or firmware), which process receives and compares the impedance of the desired tissue (before, during, and/or after the delivery of pulse of energy) with a present value, preferably current, and adjusts the pulse of energy delivered to achieve the preset value. A feedback mechanism may be performed by an analog closed loop circuit.

[0083] k. Fragment

[0084] “Fragment” may mean a polypeptide fragment of a filovirus immunogen that is capable of eliciting an immune response in a mammal against filovirus by recognizing the particular filovirus antigen. The filovirus envelope glycoprotein immunogen may optionally include a signal peptides and/or a methionine at position 1, proteins 98% or more homologous to the consensus sequences set forth herein, proteins 99% or more homologous to the consensus sequences set forth herein, and proteins 100% identical to the consensus sequences set forth herein, in each case with or without signal peptides and/or a methionine at position 1. A fragment may or may not for example comprise a fragment of a filovirus immunogen linked to a signal peptide such as an immunoglobulin signal peptide for example IgE signal peptide or IgG signal peptide.

[0085] Fragments of any of ZEBOV CON, SUDV CON, MARV ANG, MARV-RAV CON, MARV-OZO CON or MARV-MUS CON or variants thereof, in each case with or without signal peptides and/or a methionine at position 1, may comprise 20% or more, 25% or more, 30% or more, 35% or more, 40% or more, 45% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 91% or more, 92% or more, 93% or more, 94% or more, 95% or more, 96% or more, 97% or more, 98% or more, 99% or more percent of the length of the particular full length ZEBOV CON, SUDV CON, MARV ANG, MARV-RAV CON, MARV-OZO CON or MARV-MUS CON or variants thereof. Fragments refer to fragments polypeptides 100% identical to the sequences ZEBOV CON, SUDV CON, MARV ANG, MARV-RAV CON, MARV-OZO CON or MARV-MUS CON, in each case with or without signal peptides and/or a methionine at position 1. Fragments also refer to fragments of variants, i.e. polypeptides that 95% or more, 98% or more, or 99% or more homologous to the sequences ZEBOV CON, SUDV CON, MARV ANG, MARV-RAV CON, MARV-OZO CON or MARV-MUS CON, in each case with or without signal peptides and/or a methionine at position 1. The fragment may comprise a fragment of a polypeptide that is 98% or more homologous, 99% or more homologous, or 100% identical to the filovirus immunogens set forth in SEQ ID NOs: 1-6 and additionally comprise a signal peptide such as an immunoglobulin signal peptide which is not included when calculating percent homology. In some embodiments, a fragment of SEQ ID NOs: 1-6 linked to a signal peptide such as an immunoglobulin signal peptide for example IgE signal peptide or IgG signal peptide. The fragment may comprise fragments of SEQ ID NOs: 1-6 including the N terminal methionine. Fragments also refer to fragments of a polypeptide that is 95% or more, 98% or more, or 99% or more homologous to the sequence disclosed in SEQ ID NOs: 1-6. If a signal peptide is present it is not included when calculating percent homology.

[0086] In some embodiments, fragments of SEQ ID NOs: 1-6 or variants thereof may comprise 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670 or more contiguous amino acids of any of SEQ ID NOs: 1-6 or variants thereof. In some embodiments, fragments of SEQ ID NOs: 1-6 or variants thereof may comprise 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670 or less contiguous amino acids of any of SEQ ID NOs: 1-6 or variants thereof.

[0087] “Fragment” may also mean a fragment of a nucleic acid sequence that encodes a filovirus immunogen, the nucleic acid fragment encoding a fragment of filovirus immunogen that is capable of eliciting an immune response in a mammal against filovirus by recognizing the particular filovirus antigen. Fragments of nucleic acid fragment encoding a filovirus immunogen or variants thereof, in each case with or without signal peptides and/or a methionine at position 1, may comprise 20% or more, 25% or more, 30% or more, 35% or more, 40% or more, 45% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 91% or more, 92% or more, 93% or more, 94% or more, 95% or more, 96% or more, 97% or more, 98% or more, 99% or more percent of the length of the particular full length nucleic acid sequence that encodes a filovirus immunogen or variants thereof. The fragment may comprise a fragment of nucleotide sequence that encodes polypeptide that is 98% or more homologous, 99% or more homologous, or 100% identical to the filovirus immunogens set forth in SEQ ID NOs: 1-6 and additionally comprise a signal peptide such as an immunoglobulin signal peptide which is not included when calculating percent homology. In some embodiments, fragment of nucleotide sequence that encodes a fragment of SEQ ID NOs: 1-6 linked to a signal peptide such as an immunoglobulin signal peptide for example IgE signal peptide or IgG signal peptide. Coding sequences of a signal peptide is present it is not included when calculating percent homology. In some embodiments, fragment of nucleotide sequence that encodes fragments of SEQ ID NOs: 1-6 or variants thereof may comprise sequences that encode 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670 or more contiguous amino acids of any of SEQ ID NOs: 1-6 or variants thereof. In some embodiments, fragment of nucleotide sequence that encodes fragments of SEQ ID NOs: 1-6 or variants thereof may sequences that encode comprise 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670, 675 or less contiguous amino acids of any of SEQ ID NOs: 1-6 or variants thereof.

[0088] In some embodiments, fragments are fragments of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66. Fragments of SEQ ID NO:64, SEQ ID NO:65 or SEQ ID NO:66, in each case with or without signal peptides and/or a methionine at position 1, may comprise 20% or more, 25% or more, 30% or more, 35% or more, 40% or more, 45% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 91% or more, 92% or more, 93% or more, 94% or more, 95% or more, 96% or more, 97% or more, 98% or more, 99% or more percent of the length of the particular full length nucleic acid sequence that encodes a filovirus immunogen or variants thereof. The fragment of

SEQ ID NO:64, SEQ ID NO:64, or SEQ ID NO:66 may comprise a fragment of nucleotide sequence that encodes polypeptide that is 98% or more homologous, 99% or more homologous, or 100% identical to the filovirus immunogens encoded by of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 and additionally comprise a signal peptide such as an immunoglobulin signal peptide which is not included when calculating percent homology. Fragments of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 may comprises sequences that encode 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670 or more contiguous amino acids of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 or variants thereof. Fragments of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 may comprises sequences that encode 15, 20, 30, 40, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 610, 620, 630, 640, 650, 660, 670, 675 or less contiguous amino acids of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 or variants thereof.

[0089] l. Identical

[0090] “Identical” or “identity” as used herein in the context of two or more nucleic acids or polypeptide sequences, may mean that the sequences have a specified percentage of residues that are the same over a specified region. The percentage may be calculated by optimally aligning the two sequences, comparing the two sequences over the specified region, determining the number of positions at which the identical residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the specified region, and multiplying the result by 100 to yield the percentage of sequence identity. In cases where the two sequences are of different lengths or the alignment produces one or more staggered ends and the specified region of comparison includes only a single sequence, the residues of single sequence are included in the denominator but not the numerator of the calculation. When comparing DNA and RNA, thymine (T) and uracil (U) may be considered equivalent. Identity may be performed manually or by using a computer sequence algorithm such as BLAST or BLAST 2.0.

[0091] m. Impedance

[0092] “Impedance” as used herein may be used when discussing the feedback mechanism and can be converted to a current value according to Ohm’s law, thus enabling comparisons with the preset current.

[0093] n. Immune Response

[0094] “Immune response” as used herein may mean the activation of a host’s immune system, e.g., that of a mammal, in response to the introduction of one or more filovirus consensus antigen via the provided DNA plasmid vaccines. The immune response can be in the form of a cellular or humoral response, or both.

[0095] o. Nucleic Acid

[0096] “Nucleic acid” or “oligonucleotide” or “polynucleotide” as used herein may mean at least two nucleotides covalently linked together. The depiction of a single strand also defines the sequence of the complementary strand. Thus, a nucleic acid also encompasses the complementary strand of a depicted single strand. Many variants of a nucleic acid may be used for the same purpose as a given nucleic acid. Thus, a nucleic acid also encompasses substantially identical nucleic acids and complements thereof. A single strand provides a probe that may hybridize to a target

sequence under stringent hybridization conditions. Thus, a nucleic acid also encompasses a probe that hybridizes under stringent hybridization conditions.

[0097] Nucleic acids may be single stranded or double stranded, or may contain portions of both double stranded and single stranded sequence. The nucleic acid may be DNA, both genomic and cDNA, RNA, or a hybrid, where the nucleic acid may contain combinations of deoxyribo- and ribo-nucleotides, and combinations of bases including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine and isoguanine. Nucleic acids may be obtained by chemical synthesis methods or by recombinant methods.

[0098] p. Operably Linked

[0099] “Operably linked” as used herein may mean that expression of a gene is under the control of a promoter with which it is spatially connected. A promoter may be positioned 5’ (upstream) or 3’ (downstream) of a gene under its control. The distance between the promoter and a gene may be approximately the same as the distance between that promoter and the gene it controls in the gene from which the promoter is derived. As is known in the art, variation in this distance may be accommodated without loss of promoter function.

[0100] q. Promoter

[0101] “Promoter” as used herein may mean a synthetic or naturally-derived molecule which is capable of conferring, activating or enhancing expression of a nucleic acid in a cell. A promoter may comprise one or more specific transcriptional regulatory sequences to further enhance expression and/or to alter the spatial expression and/or temporal expression of same. A promoter may also comprise distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A promoter may be derived from sources including viral, bacterial, fungal, plants, insects, and animals. A promoter may regulate the expression of a gene component constitutively, or differentially with respect to cell, the tissue or organ in which expression occurs or, with respect to the developmental stage at which expression occurs, or in response to external stimuli such as physiological stresses, pathogens, metal ions, or inducing agents. Representative examples of promoters include the bacteriophage T7 promoter, bacteriophage T3 promoter, SP6 promoter, lac operator-promoter, tac promoter, SV40 late promoter, SV40 early promoter, RSV-LTR promoter, CMV IE promoter, SV40 early promoter or SV40 late promoter and the CMV IE promoter.

[0102] r. Stringent Hybridization Conditions

[0103] “Stringent hybridization conditions” as used herein may mean conditions under which a first nucleic acid sequence (e.g., probe) will hybridize to a second nucleic acid sequence (e.g., target), such as in a complex mixture of nucleic acids. Stringent conditions are sequence-dependent and will be different in different circumstances. Stringent conditions may be selected to be about 5-10° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m may be the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Stringent conditions may be those in which the salt concentration is less than about 1.0 M

sodium ion, such as about 0.01-1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., about 10-50 nucleotides) and at least about 60° C. for long probes (e.g., greater than about 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal may be at least 2 to 10 times background hybridization. Exemplary stringent hybridization conditions include the following: 50% formamide, 5×SSC, and 1% SDS, incubating at 42° C., or, 5×SSC, 1% SDS, incubating at 65° C., with wash in 0.2×SSC, and 0.1% SDS at 65° C.

[0104] s. Substantially Complementary

[0105] “Substantially complementary” as used herein may mean that a first sequence is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical to the complement of a second sequence over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more nucleotides or amino acids, or that the two sequences hybridize under stringent hybridization conditions.

[0106] t. Substantially Identical

[0107] “Substantially identical” as used herein may mean that a first and second sequence are at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% identical over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more nucleotides or amino acids, or with respect to nucleic acids, if the first sequence is substantially complementary to the complement of the second sequence.

[0108] u. Variant

[0109] “Variant” used herein with respect to a nucleic acid may mean (i) a portion or fragment of a referenced nucleotide sequence; (ii) the complement of a referenced nucleotide sequence or portion thereof; (iii) a nucleic acid that is substantially identical to a referenced nucleic acid or the complement thereof; or (iv) a nucleic acid that hybridizes under stringent conditions to the referenced nucleic acid, complement thereof, or a sequences substantially identical thereto.

[0110] “Variant” with respect to a peptide or polypeptide that differs in amino acid sequence by the insertion, deletion, or conservative substitution of amino acids, but retain at least one biological activity. Variant may also mean a protein with an amino acid sequence that is substantially identical to a referenced protein with an amino acid sequence that retains at least one biological activity. A conservative substitution of an amino acid, i.e., replacing an amino acid with a different amino acid of similar properties (e.g., hydrophilicity, degree and distribution of charged regions) is recognized in the art as typically involving a minor change. These minor changes can be identified, in part, by considering the hydrophobic index of amino acids, as understood in the art. Kyte et al., *J. Mol. Biol.* 157:105-132 (1982). The hydrophobic index of an amino acid is based on a consideration of its hydrophobicity and charge. It is known in the art that amino acids of similar hydrophobic indexes can be substituted and still retain protein function. In one aspect, amino acids having hydrophobic indexes of ± 2 are substituted. The hydrophilicity of amino acids can also be used to reveal substitutions that would result in proteins retaining biological function. A consideration of the hydrophilicity of amino acids in the context of a peptide permits calculation of the

greatest local average hydrophilicity of that peptide, a useful measure that has been reported to correlate well with antigenicity and immunogenicity. U.S. Pat. No. 4,554,101, incorporated fully herein by reference. Substitution of amino acids having similar hydrophilicity values can result in peptides retaining biological activity, for example immunogenicity, as is understood in the art. Substitutions may be performed with amino acids having hydrophilicity values within ± 2 of each other. Both the hydrophobicity index and the hydrophilicity value of amino acids are influenced by the particular side chain of that amino acid. Consistent with that observation, amino acid substitutions that are compatible with biological function are understood to depend on the relative similarity of the amino acids, and particularly the side chains of those amino acids, as revealed by the hydrophobicity, hydrophilicity, charge, size, and other properties. Variants are preferably homologous to SEQ ID NO:1-6 by 95% or more, 96% or more, 97% or more, 98% or more or 99% or more.

[0111] “Variant” with respect to a nucleic acid sequence that encodes the same specific amino acid sequence differs in nucleotide sequence by use of different codons. Variants of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 that encode the same amino acid sequence as those encoded by SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 may be any degree of homology, preferably 80% or more, 85% or more, 90% or more, 95% or more, 96% or more, 97% or more, 98% or more or 99% or more. Variant may also be variants of SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 that encode protein which are variants of the proteins encoded by SEQ ID NO:64, SEQ ID NO:65, or SEQ ID NO:66 with an amino acid sequence that is substantially identical to a referenced protein with an amino acid sequence that retains at least one biological activity, typically the amino acid sequences are homologous by 95% or more, 96% or more, 97% or more, 98% or more or 99% or more.

[0112] v. Vector

[0113] “Vector” used herein may mean a nucleic acid sequence containing an origin of replication. A vector may be a plasmid, bacteriophage, bacterial artificial chromosome or yeast artificial chromosome. A vector may be a DNA or RNA vector. A vector may be either a self-replicating extrachromosomal vector or a vector which integrates into a host genome.

2. Proteins

[0114] Provided herein are filovirus immunogens that can be used to induce broad immunity against multiple subtypes or serotypes of a particular filovirus antigen. Consensus filovirus antigens may include consensus amino acid sequences of Marburgvirus filovirus glycoprotein MARV RAV immunogen, consensus amino acid sequences of Marburgvirus filovirus glycoprotein MARV OZO immunogen, consensus amino acid sequences of Marburgvirus filovirus glycoprotein MARV MUS immunogen, isolate amino acid sequences of Marburgvirus filovirus glycoprotein MARV ANG immunogen, consensus amino acid sequences of Zaire ebolavirus glycoprotein ZEBOV immunogen and consensus amino acid sequences of Sudan ebolavirus glycoprotein SUDV immunogen, respectively. In some embodiments, the immunogens may comprise a signal peptide from a different protein such as an immunoglobulin protein, for example an IgE signal peptide or an IgG signal peptide.

[0115] The amino acid sequence for immunogens include SEQ ID NO:1-6, variants thereof and fragments of SEQ ID NO:1-6 and variants thereof, optionally including a signal peptide such as for example an IgE or IgG signal peptide.

3. Coding Sequences Encoding Proteins

[0116] Coding sequences encoding the proteins set forth herein may be generated using routine methods. Composition comprising a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen are provided and a nucleic acid sequence that encodes a first consensus Marburg marburgvirus envelope glycoprotein immunogen, a nucleic acid sequence that encodes a second consensus Marburg marburgvirus envelope glycoprotein immunogen, and a nucleic acid sequence that encodes a third consensus Marburg marburgvirus envelope glycoprotein immunogen can be generated based upon the amino acid sequences disclosed.

[0117] Nucleic acid sequence may encodes a full length consensus Zaire ebolavirus envelope glycoprotein immunogen, a full length consensus Sudan ebolavirus envelope glycoprotein immunogen, a full length Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, a full length first consensus Marburg marburgvirus envelope glycoprotein immunogen, a full length second consensus Marburg marburgvirus envelope glycoprotein immunogen, or a full length third consensus Marburg marburgvirus envelope glycoprotein immunogen. Nucleic acid sequences may comprise a sequence that encodes SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. Nucleic acid sequence may comprise SEQ ID NO:64, SEQ ID NO:65 or SEQ ID NO:66. Nucleic acid sequence may optionally comprise coding sequences that encode a signal peptide such as for example an IgE or IgG signal peptide.

[0118] Nucleic acid sequence may encode a fragment of a full length consensus Zaire ebolavirus envelope glycoprotein immunogen, a fragment of a full length consensus Sudan ebolavirus envelope glycoprotein immunogen, a fragment of a full length Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, a fragment of a full length first consensus Marburg marburgvirus envelope glycoprotein immunogen, a fragment of a full length second consensus Marburg marburgvirus envelope glycoprotein immunogen, or a fragment of a full length third consensus Marburg marburgvirus envelope glycoprotein immunogen. Nucleic acid sequence may comprise a sequence that encodes a fragment of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. Nucleic acid sequence may comprise a fragment of SEQ ID NO:64, SEQ ID NO:65 or SEQ ID NO:66. Fragment sizes are disclosed herein as set forth in section entitled "Fragments". Nucleic acid sequence may optionally comprise coding sequences that encode a signal peptide such as for example an IgE or IgG signal peptide.

[0119] Nucleic acid sequences may encode a protein homologous to a full length consensus Zaire ebolavirus envelope glycoprotein immunogen, a protein homologous to a full length consensus Sudan ebolavirus envelope glycoprotein immunogen, a protein homologous to a full length

Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, a protein homologous to a full length first consensus Marburg marburgvirus envelope glycoprotein immunogen, a protein homologous to a full length second consensus Marburg marburgvirus envelope glycoprotein immunogen, or a protein homologous to a full length third consensus Marburg marburgvirus envelope glycoprotein immunogen. Nucleic acid sequence may comprise a sequence that encodes a protein homologous to SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. Nucleic acid sequence may be homologous to SEQ ID NO:64, SEQ ID NO:65 or SEQ ID NO:66. Degrees of homology are discussed herein such as in the section referring to Variants. Nucleic acid sequence may optionally comprise coding sequences that encode a signal peptide such as for example an IgE or IgG signal peptide.

[0120] Nucleic acid sequence may encode a protein homologous to fragment of a full length consensus Zaire ebolavirus envelope glycoprotein immunogen, a protein homologous to a fragment of a full length consensus Sudan ebolavirus envelope glycoprotein immunogen, a protein homologous to a fragment of a full length Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, a protein homologous to a fragment of a full length first consensus Marburg marburgvirus envelope glycoprotein immunogen, a protein homologous to a fragment of a full length second consensus Marburg marburgvirus envelope glycoprotein immunogen, or a protein homologous to a fragment of a full length third consensus Marburg marburgvirus envelope glycoprotein immunogen. Nucleic acid sequence may comprise a sequence that encodes a protein homologous to a fragment of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5 or SEQ ID NO:6. Nucleic acid sequence may comprise a fragment homologous to SEQ ID NO:64, SEQ ID NO:65 or SEQ ID NO:66. Degrees of homology are discussed herein such as in the section referring to Variants. Nucleic acid sequence may optionally comprise coding sequences that encode a signal peptide such as for example an IgE or IgG signal peptide.

[0121] SEQ ID NO:64 is the nucleotide sequence insert in plasmid pEBOZ which encodes consensus Zaire ebolavirus envelope glycoprotein immunogen.

[0122] SEQ ID NO:65 is the nucleotide sequence insert in plasmid pEBOS which encodes consensus Sudan ebolavirus envelope glycoprotein immunogen.

[0123] SEQ ID NO:66 is the nucleotide sequence insert in plasmid pMARZ ANG which encodes the Marburg marburgvirus Angola 2005 envelope glycoprotein.

4. Plasmid

[0124] Plasmid may comprise a nucleic acid sequence that encodes one or more of the various immunogens disclosed above including coding sequences that encode synthetic, consensus antigen capable of eliciting an immune response against filoproteins.

[0125] A single plasmid may contain coding sequence for a single filoprotein immunogen, coding sequence for two filoprotein immunogens, coding sequence for three filoprotein immunogens, coding sequence for four filoprotein immunogens, coding sequence for five filoprotein immunogens or coding sequence for six filoprotein immunogens. A single plasmid may contain a coding sequence for a single

filoprotein immunogen which can be formulated together. In some embodiments, a plasmid may comprise coding sequence that encodes IL-12, IL-15 and/or IL-28.

[0126] The plasmid may further comprise an initiation codon, which may be upstream of the coding sequence, and a stop codon, which may be downstream of the coding sequence. The initiation and termination codon may be in frame with the coding sequence.

[0127] The plasmid may also comprise a promoter that is operably linked to the coding sequence. The promoter operably linked to the coding sequence may be a promoter from simian virus 40 (SV40), a mouse mammary tumor virus (MMTV) promoter, a human immunodeficiency virus (HIV) promoter such as the bovine immunodeficiency virus (BIV) long terminal repeat (LTR) promoter, a Moloney virus promoter, an avian leukosis virus (ALV) promoter, a cytomegalovirus (CMV) promoter such as the CMV immediate early promoter, Epstein Barr virus (EBV) promoter, or a Rous sarcoma virus (RSV) promoter. The promoter may also be a promoter from a human gene such as human actin, human myosin, human hemoglobin, human muscle creatine, or human metallothionein. The promoter may also be a tissue specific promoter, such as a muscle or skin specific promoter, natural or synthetic. Examples of such promoters are described in US patent application publication no. US20040175727, the contents of which are incorporated herein in its entirety.

[0128] The plasmid may also comprise a polyadenylation signal, which may be downstream of the coding sequence. The polyadenylation signal may be a SV40 polyadenylation signal, LTR polyadenylation signal, bovine growth hormone (bGH) polyadenylation signal, human growth hormone (hGH) polyadenylation signal, or human β -globin polyadenylation signal. The SV40 polyadenylation signal may be a polyadenylation signal from a pCEP4 plasmid (Invitrogen, San Diego, Calif.).

[0129] The plasmid may also comprise an enhancer upstream of the coding sequence. The enhancer may be human actin, human myosin, human hemoglobin, human muscle creatine or a viral enhancer such as one from CMV, FMDV, RSV or EBV. Polynucleotide function enhancers are described in U.S. Pat. Nos. 5,593,972, 5,962,428, and WO94/016737, the contents of each are fully incorporated by reference.

[0130] The plasmid may also comprise a mammalian origin of replication in order to maintain the plasmid extrachromosomally and produce multiple copies of the plasmid in a cell. The plasmid may be pVAX1, pCEP4 or pREP4 from Invitrogen (San Diego, Calif.), which may comprise the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region, which may produce high copy episomal replication without integration. The backbone of the plasmid may be pAV0242. The plasmid may be a replication defective adenovirus type 5 (Ad5) plasmid.

[0131] The plasmid may also comprise a regulatory sequence, which may be well suited for gene expression in a cell into which the plasmid is administered. The coding sequence may comprise a codon that may allow more efficient transcription of the coding sequence in the host cell.

[0132] The coding sequence may also comprise an Ig leader sequence. The leader sequence may be 5' of the coding sequence. The consensus antigens encoded by this

sequence may comprise an N-terminal Ig leader followed by a consensus antigen protein. The N-terminal Ig leader may be IgE or IgG.

[0133] The plasmid may be pSE420 (Invitrogen, San Diego, Calif.), which may be used for protein production in *Escherichia coli* (*E. coli*). The plasmid may also be pYES2 (Invitrogen, San Diego, Calif.), which may be used for protein production in *Saccharomyces cerevisiae* strains of yeast. The plasmid may also be of the MAXBAC™ complete baculovirus expression system (Invitrogen, San Diego, Calif.), which may be used for protein production in insect cells. The plasmid may also be pcDNA I or pcDNA3 (Invitrogen, San Diego, Calif.), which may be used for protein production in mammalian cells such as Chinese hamster ovary (CHO) cells.

5. Compositions

[0134] Compositions are provided which comprise nucleic acid molecules. The compositions may comprise a plurality of copies of a single nucleic acid molecule such a single plasmid, a plurality of copies of two or more different nucleic acid molecules such as two or more different plasmids. For example a composition may comprise plurality of two, three, four, five, six, seven, eight, nine or ten or more different nucleic acid molecules. Such compositions may comprise plurality of two, three, four, five, six, or more different plasmids.

[0135] Compositions may comprise nucleic acid molecules, such as plasmids, that collectively contain coding sequence for a single filoprotein immunogen selected from the group consisting of one or more a consensus Zaire ebolavirus envelope glycoprotein immunogen, a consensus Sudan ebolavirus envelope glycoprotein immunogen, the Marburg marburgvirus Angola 2005 envelope glycoprotein, the first consensus Marburg marburgvirus envelope glycoprotein immunogen, the second consensus Marburg marburgvirus envelope glycoprotein immunogen and the third consensus Marburg marburgvirus envelope glycoprotein immunogen.

[0136] Composition comprise nucleic acid sequence that encode the combination of: a consensus Zaire ebolavirus envelope glycoprotein immunogen and a consensus Sudan ebolavirus envelope glycoprotein immunogen; or a consensus Zaire ebolavirus envelope glycoprotein immunogen, a consensus Sudan ebolavirus envelope glycoprotein immunogen and the Marburg marburgvirus Angola 2005 envelope glycoprotein; or a consensus Zaire ebolavirus envelope glycoprotein immunogen, a consensus Sudan ebolavirus envelope glycoprotein immunogen, the first consensus Marburg marburgvirus envelope glycoprotein immunogen, the second consensus Marburg marburgvirus envelope glycoprotein immunogen and the third consensus Marburg marburgvirus envelope glycoprotein immunogen; or a consensus Zaire ebolavirus envelope glycoprotein immunogen, a consensus Sudan ebolavirus envelope glycoprotein immunogen, the Marburg marburgvirus Angola 2005 envelope glycoprotein, the first consensus Marburg marburgvirus envelope glycoprotein immunogen, the second consensus Marburg marburgvirus envelope glycoprotein immunogen and the third consensus Marburg marburgvirus envelope glycoprotein immunogen. Each coding sequence for each filoprotein immunogens is preferably included on a separate plasmid. Accordingly, compositions that comprise nucleic acid sequence that encode a consensus Zaire ebolavirus

receptor, Flt, Apo-1, p55, WSL-1, DR3, TRAMP, Apo-3, AIR, LARD, NGRF, DR4, DR5, KILLER, TRAIL-R2, TRICK2, DR6, Caspase ICE, Fos, c-jun, Sp-1, Ap-1, Ap-2, p38, p65Rel, MyD88, IRAK, TRAF6, Ikb, Inactive NIK, SAPK, SAP-1, JNK, interferon response genes, NFkB, Bax, TRAIL, TRAIL_{rec}, TRAIL_{rec}DRC5, TRAIL-R3, TRAIL-R4, RANK, RANK LIGAND, Ox40, Ox40 LIGAND, NKG2D, MICA, MICB, NKG2A, NKG2B, NKG2C, NKG2E, NKG2F, TAP1, TAP2 and functional fragments thereof or a combination thereof. In some embodiments adjuvant may be one or more proteins and/or nucleic acid molecules that encode proteins selected from the group consisting of: CCL-20, IL-12, IL-15, IL-28, CTACK, TECK, MEC or RANTES. Examples of IL-12 constructs and sequences are disclosed in PCT application no. PCT/US1997/019502 and corresponding U.S. application Ser. No. 08/956,865, and U.S. Provisional Application Ser. No. 61/569,600 filed Dec. 12, 2011, which are each incorporated herein by reference. Examples of IL-15 constructs and sequences are disclosed in PCT application no. PCT/US04/18962 and corresponding U.S. application Ser. No. 10/560,650, and in PCT application no. PCT/US07/00886 and corresponding U.S. application Ser. No. 12/160,766, and in PCT application no. PCT/US10/048827, which are each incorporated herein by reference. Examples of IL-28 constructs and sequences are disclosed in PCT application no. PCT/US09/039648 and corresponding U.S. application Ser. No. 12/936,192, which are each incorporated herein by reference. Examples of RANTES and other constructs and sequences are disclosed in PCT application no. PCT/US1999/004332 and corresponding U.S. application Ser. No. 09/622,452, which are each incorporated herein by reference. Other examples of RANTES constructs and sequences are disclosed in PCT application no. PCT/US11/024098, which is incorporated herein by reference. Examples of RANTES and other constructs and sequences are disclosed in PCT application no. PCT/US1999/004332 and corresponding U.S. application Ser. No. 09/622,452, which are each incorporated herein by reference. Other examples of RANTES constructs and sequences are disclosed in PCT application no. PCT/US11/024098, which is incorporated herein by reference. Examples of chemokines CTACK, TECK and MEC constructs and sequences are disclosed in PCT application no. PCT/US2005/042231 and corresponding U.S. application Ser. No. 11/719,646, which are each incorporated herein by reference. Examples of OX40 and other immunomodulators are disclosed in U.S. application Ser. No. 10/560,653, which is incorporated herein by reference. Examples of DR5 and other immunomodulators are disclosed in U.S. application Ser. No. 09/622,452, which is incorporated herein by reference.

[0142] The vaccine may further comprise a genetic vaccine facilitator agent as described in U.S. Ser. No. 021,579 filed Apr. 1, 1994, which is fully incorporated by reference.

[0143] The vaccine may comprise the consensus antigens and plasmids at quantities of from about 1 nanogram to 100 milligrams; about 1 microgram to about 10 milligrams; or preferably about 0.1 microgram to about 10 milligrams; or more preferably about 1 milligram to about 2 milligram. In some preferred embodiments, pharmaceutical compositions according to the present invention comprise about 5 nanogram to about 1000 micrograms of DNA. In some preferred embodiments, the pharmaceutical compositions contain about 10 nanograms to about 800 micrograms of DNA. In

some preferred embodiments, the pharmaceutical compositions contain about 0.1 to about 500 micrograms of DNA. In some preferred embodiments, the pharmaceutical compositions contain about 1 to about 350 micrograms of DNA. In some preferred embodiments, the pharmaceutical compositions contain about 25 to about 250 micrograms, from about 100 to about 200 microgram, from about 1 nanogram to 100 milligrams; from about 1 microgram to about 10 milligrams; from about 0.1 microgram to about 10 milligrams; from about 1 milligram to about 2 milligram, from about 5 nanogram to about 1000 micrograms, from about 10 nanograms to about 800 micrograms, from about 0.1 to about 500 micrograms, from about 1 to about 350 micrograms, from about 25 to about 250 micrograms, from about 100 to about 200 microgram of the consensus antigen or plasmid thereof. **[0144]** The vaccine may be formulated according to the mode of administration to be used. An injectable vaccine pharmaceutical composition may be sterile, pyrogen free and particulate free. An isotonic formulation or solution may be used. Additives for isotonicity may include sodium chloride, dextrose, mannitol, sorbitol, and lactose. The vaccine may comprise a vasoconstriction agent. The isotonic solutions may include phosphate buffered saline. Vaccine may further comprise stabilizers including gelatin and albumin. The stabilizing may allow the formulation to be stable at room or ambient temperature for extended periods of time such as LGS or polycations or polyanions to the vaccine formulation.

7. Methods of Delivery the Vaccine

[0145] Provided herein is a method for delivering the vaccine for providing genetic constructs and proteins of the consensus antigen which comprise epitopes that make them particular effective against immunogens of filovirus, particularly Marburgvirus, Ebolavirus Sudan and/or Ebolavirus Zaire, against which an immune response can be induced. The method of delivering the vaccine or vaccination may be provided to induce a therapeutic and prophylactic immune response. The vaccination process may generate in the mammal an immune response against filovirus, particularly Marburgvirus, Ebolavirus Sudan and/or Ebolavirus Zaire. The vaccine may be delivered to an individual to modulate the activity of the mammal's immune system and enhance the immune response. The delivery of the vaccine may be the transfection of the consensus antigen as a nucleic acid molecule that is expressed in the cell and delivered to the surface of the cell upon which the immune system recognized and induces a cellular, humoral, or cellular and humoral response. The delivery of the vaccine may be used to induce or elicit and immune response in mammals against filovirus, particularly Marburgvirus, Ebolavirus Sudan and/or Ebolavirus Zaire by administering to the mammals the vaccine as discussed above.

[0146] Upon delivery of the vaccine and plasmid into the cells of the mammal, the transfected cells will express and secrete consensus antigens for each of the plasmids injected from the vaccine. These proteins will be recognized as foreign by the immune system and antibodies will be made against them. These antibodies will be maintained by the immune system and allow for an effective response to subsequent infections by filovirus, particularly Marburgvirus, Ebolavirus Sudan and/or Ebolavirus Zaire.

[0147] The vaccine may be administered to a mammal to elicit an immune response in a mammal. The mammal may

be human, primate, non-human primate, cow, cattle, sheep, goat, antelope, bison, water buffalo, bison, bovids, deer, hedgehogs, elephants, llama, alpaca, mice, rats, and chicken.

[0148] a. Combination Treatments

[0149] The vaccine may be administered in combination with other proteins and/or genes encoding CCL20, α -interferon, γ -interferon, platelet derived growth factor (PDGF), TNF α , TNF β , GM-CSF, epidermal growth factor (EGF), cutaneous T cell-attracting chemokine (CTACK), epithelial thymus-expressed chemokine (TECK), mucosae-associated epithelial chemokine (MEC), IL-12, IL-15 including IL-15 having the signal sequence deleted and optionally including the different signal peptide such as the IgE signal peptide, MHC, CD80, CD86, IL-28, IL-1, IL-2, IL-4, IL-5, IL-6, IL-10, IL-18, MCP-1, MIP-1 α , MIP-1 β , IL-8, RANTES, L-selectin, P-selectin, E-selectin, CD34, GlyCAM-1, MadCAM-1, LFA-1, VLA-1, Mac-1, p150.95, PECAM, ICAM-1, ICAM-2, ICAM-3, CD2, LFA-3, M-CSF, G-CSF, mutant forms of IL-18, CD40, CD40L, vascular growth factor, fibroblast growth factor, IL-7, nerve growth factor, vascular endothelial growth factor, Fas, TNF receptor, Flt, Apo-1, p55, WSL-1, DR3, TRAMP, Apo-3, AIR, LARD, NGRF, DR4, DR5, KILLER, TRAIL-R2, TRICK2, DR6, Caspase ICE, Fos, c-jun, Sp-1, Ap-1, Ap-2, p38, p65Rel, MyD88, IRAK, TRAF6, I κ B, Inactive NIK, SAP K, SAP-1, JNK, interferon response genes, NF κ B, Bax, TRAIL, TRAILrec, TRAILrecDRC5, TRAIL-R3, TRAIL-R4, RANK, RANK LIGAND, Ox40, Ox40 LIGAND, NKG2D, MICA, MICB, NKG2A, NKG2B, NKG2C, NKG2E, NKG2F, TAP1, TAP2 and functional fragments thereof or combinations thereof. In some embodiments, the vaccine is administered in combination with one or more of the following nucleic acid molecules and/or proteins: nucleic acid molecules selected from the group consisting of nucleic acid molecules comprising coding sequence that encode one or more of CCL20, IL-12, IL-15, IL-28, CTACK, TECK, MEC and RANTES or functional fragments thereof, and proteins selected from the group consisting of: CCL20, IL-12 protein, IL-15 protein, IL-28 protein, CTACK protein, TECK protein, MEC protein or RANTES protein or functional fragments thereof.

[0150] The vaccine may be administered by different routes including orally, parenterally, sublingually, transdermally, rectally, transmucosally, topically, via inhalation, via buccal administration, intrapleurally, intravenous, intraarterial, intraperitoneal, subcutaneous, intramuscular, intranasal, intrathecal, and intraarticular or combinations thereof. For veterinary use, the composition may be administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal. The vaccine may be administered by traditional syringes, needleless injection devices, “microprojectile bombardment gone guns”, or other physical methods such as electroporation (“EP”), “hydrodynamic method”, or ultrasound.

[0151] The plasmid of the vaccine may be delivered to the mammal by several well-known technologies including DNA injection (also referred to as DNA vaccination) with and without in vivo electroporation, liposome mediated, nanoparticle facilitated, recombinant vectors such as recombinant adenovirus, recombinant adenovirus associated virus and recombinant vaccinia. The consensus antigen may be delivered via DNA injection and along with in vivo electroporation.

[0152] b. Electroporation

[0153] Administration of the vaccine via electroporation of the plasmids of the vaccine may be accomplished using electroporation devices that can be configured to deliver to a desired tissue of a mammal a pulse of energy effective to cause reversible pores to form in cell membranes, and preferable the pulse of energy is a constant current similar to a preset current input by a user. The electroporation device may comprise an electroporation component and an electrode assembly or handle assembly. The electroporation component may include and incorporate one or more of the various elements of the electroporation devices, including: controller, current waveform generator, impedance tester, waveform logger, input element, status reporting element, communication port, memory component, power source, and power switch. The electroporation may be accomplished using an in vivo electroporation device, for example CELLECTRA EP system (VGX Pharmaceuticals, Blue Bell, Pa.) or Elgen electroporator (Genetronics, San Diego, Calif.) to facilitate transfection of cells by the plasmid.

[0154] The electroporation component may function as one element of the electroporation devices, and the other elements are separate elements (or components) in communication with the electroporation component. The electroporation component may function as more than one element of the electroporation devices, which may be in communication with still other elements of the electroporation devices separate from the electroporation component. The elements of the electroporation devices existing as parts of one electromechanical or mechanical device may not be limited as the elements can function as one device or as separate elements in communication with one another. The electroporation component may be capable of delivering the pulse of energy that produces the constant current in the desired tissue, and includes a feedback mechanism. The electrode assembly may include an electrode array having a plurality of electrodes in a spatial arrangement, wherein the electrode assembly receives the pulse of energy from the electroporation component and delivers same to the desired tissue through the electrodes. At least one of the plurality of electrodes is neutral during delivery of the pulse of energy and measures impedance in the desired tissue and communicates the impedance to the electroporation component. The feedback mechanism may receive the measured impedance and can adjust the pulse of energy delivered by the electroporation component to maintain the constant current.

[0155] A plurality of electrodes may deliver the pulse of energy in a decentralized pattern. The plurality of electrodes may deliver the pulse of energy in the decentralized pattern through the control of the electrodes under a programmed sequence, and the programmed sequence is input by a user to the electroporation component. The programmed sequence may comprise a plurality of pulses delivered in sequence, wherein each pulse of the plurality of pulses is delivered by at least two active electrodes with one neutral electrode that measures impedance, and wherein a subsequent pulse of the plurality of pulses is delivered by a different one of at least two active electrodes with one neutral electrode that measures impedance.

[0156] The feedback mechanism may be performed by either hardware or software. The feedback mechanism may be performed by an analog closed-loop circuit. The feedback occurs every 50 μ s, 20 μ s, 10 μ s or 1 μ s, but is preferably a real-time feedback or instantaneous (i.e., substantially

instantaneous as determined by available techniques for determining response time). The neutral electrode may measure the impedance in the desired tissue and communicates the impedance to the feedback mechanism, and the feedback mechanism responds to the impedance and adjusts the pulse of energy to maintain the constant current at a value similar to the preset current. The feedback mechanism may maintain the constant current continuously and instantaneously during the delivery of the pulse of energy.

[0157] Examples of electroporation devices and electroporation methods that may facilitate delivery of the DNA vaccines of the present invention, include those described in U.S. Pat. No. 7,245,963 by Draghia-Akli, et al., U.S. Patent Pub. 2005/0052630 submitted by Smith, et al., the contents of which are hereby incorporated by reference in their entirety. Other electroporation devices and electroporation methods that may be used for facilitating delivery of the DNA vaccines include those provided in co-pending and co-owned U.S. patent application Ser. No. 11/874,072, filed Oct. 17, 2007, which claims the benefit under 35 USC 119(e) to U.S. Provisional Application Ser. No. 60/852,149, filed Oct. 17, 2006, and 60/978,982, filed Oct. 10, 2007, all of which are hereby incorporated in their entirety.

[0158] U.S. Pat. No. 7,245,963 by Draghia-Akli, et al. describes modular electrode systems and their use for facilitating the introduction of a biomolecule into cells of a selected tissue in a body or plant. The modular electrode systems may comprise a plurality of needle electrodes; a hypodermic needle; an electrical connector that provides a conductive link from a programmable constant-current pulse controller to the plurality of needle electrodes; and a power source. An operator can grasp the plurality of needle electrodes that are mounted on a support structure and firmly insert them into the selected tissue in a body or plant. The biomolecules are then delivered via the hypodermic needle into the selected tissue. The programmable constant-current pulse controller is activated and constant-current electrical pulse is applied to the plurality of needle electrodes. The applied constant-current electrical pulse facilitates the introduction of the biomolecule into the cell between the plurality of electrodes. The entire content of U.S. Pat. No. 7,245,963 is hereby incorporated by reference.

[0159] U.S. Patent Pub. 2005/0052630 submitted by Smith, et al. describes an electroporation device which may be used to effectively facilitate the introduction of a biomolecule into cells of a selected tissue in a body or plant. The electroporation device comprises an electro-kinetic device (“EKD device”) whose operation is specified by software or firmware. The EKD device produces a series of programmable constant-current pulse patterns between electrodes in an array based on user control and input of the pulse parameters, and allows the storage and acquisition of current waveform data. The electroporation device also comprises a replaceable electrode disk having an array of needle electrodes, a central injection channel for an injection needle, and a removable guide disk. The entire content of U.S. Patent Pub. 2005/0052630 is hereby incorporated by reference.

[0160] The electrode arrays and methods described in U.S. Pat. No. 7,245,963 and U.S. Patent Pub. 2005/0052630 may be adapted for deep penetration into not only tissues such as muscle, but also other tissues or organs. Because of the configuration of the electrode array, the injection needle (to deliver the biomolecule of choice) is also inserted com-

pletely into the target organ, and the injection is administered perpendicular to the target issue, in the area that is pre-delineated by the electrodes. The electrodes described in U.S. Pat. No. 7,245,963 and U.S. Patent Pub. 2005/005263 are preferably 20 mm long and 21 gauge.

[0161] Additionally, contemplated in some embodiments that incorporate electroporation devices and uses thereof, there are electroporation devices that are those described in the following patents: U.S. Pat. No. 5,273,525 issued Dec. 28, 1993, U.S. Pat. No. 6,110,161 issued Aug. 29, 2000, U.S. Pat. No. 6,261,281 issued Jul. 17, 2001, and U.S. Pat. No. 6,958,060 issued Oct. 25, 2005, and U.S. Pat. No. 6,939,862 issued Sep. 6, 2005. Furthermore, patents covering subject matter provided in U.S. Pat. No. 6,697,669 issued Feb. 24, 2004, which concerns delivery of DNA using any of a variety of devices, and U.S. Pat. No. 7,328,064 issued Feb. 5, 2008, drawn to method of injecting DNA are contemplated herein. The above-patents are incorporated by reference in their entirety.

[0162] c. Method of Preparing DNA Plasmids

[0163] Provided herein is methods for preparing the DNA plasmids that comprise the DNA vaccines discussed herein. The DNA plasmids, after the final subcloning step into the mammalian expression plasmid, can be used to inoculate a cell culture in a large scale fermentation tank, using known methods in the art.

[0164] The DNA plasmids for use with the EP devices of the present invention can be formulated or manufactured using a combination of known devices and techniques, but preferably they are manufactured using an optimized plasmid manufacturing technique that is described in a licensed, co-pending U.S. provisional application U.S. Ser. No. 60/939,792, which was filed on May 23, 2007. In some examples, the DNA plasmids used in these studies can be formulated at concentrations greater than or equal to 10 mg/mL. The manufacturing techniques also include or incorporate various devices and protocols that are commonly known to those of ordinary skill in the art, in addition to those described in U.S. Ser. No. 60/939,792, including those described in a licensed patent, U.S. Pat. No. 7,238,522, which issued on Jul. 3, 2007. The above-referenced application and patent, U.S. Ser. No. 60/939,792 and U.S. Pat. No. 7,238,522, respectively, are hereby incorporated in their entirety.

EXAMPLES

[0165] The present invention is further illustrated in the following Example. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. Thus, various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

Examples 1

Methods

Plasmid Vaccine Construction

[0166] The pMARV, pEBOS, and pEBOZ plasmid DNA constructs encode full-length GP proteins. An amino acid consensus strategy was used for the pEBOS and pEBOZ, while a type-matched sequence from the 2005 Angola outbreak strain was used (GenBank #VGP_MABVR) for pMARV (Towner J S, et al. (2006). Marburgvirus genomics and association with a large hemorrhagic fever outbreak in Angola. *J Virol* 80: 6497-6516). Consensus sequences were determined by alignment of the prevailing ZEBOV and SUDV GP amino acid sequences and generating a consensus for each. Each vaccine GP gene was genetically optimized for expression in humans (including codon- and RNA-optimization for enhancing protein expression (GenScript, Piscataway, N.J.)), synthesized commercially, and then sub-cloned (GenScript, Piscataway, N.J.) into modified pVAX1 mammalian expression vectors (Invitrogen, Carlsbad, Calif.) under the control of the cytomegalovirus immediate-early (CMV) promoter; modifications include 2A>C, 3C>T, 4T>G, 241C>G, 1,942C>T, 2,876A>-, 3,277C>T, and 3,753G>C. Phylogenetic analysis was performed by multiple-alignment with ClustalW using MEGA version 5 software. Alternatively, GP diversity among the MARV was much higher (~70% identity) in comparison, so a consensus strategy was not adopted. For coverage of MARV, we chose to utilize the MGP sequence from the 2005 outbreak in Angola (GenBank #VGP_MABVR) since it was solely responsible for the largest and deadliest MARV outbreak to date. This sequence was greater than 10% divergent from either of its closest cluster of relative strains including Musoke, Popp and Leiden (10.6% divergence), or Uganda (OIUga07), Durba (05DRC99 and 07DRC99) and Ozolin (10.3% divergence). Altogether, a three-plasmid strategy formed the foundation for our novel trivalent polyvalent-filovirus vaccine strategy.

Transfections and Immunoblotting

[0167] Human Embryonic Kidney (HEK) 293T cells were cultured, transfected, and harvested. Briefly, cells were grown in DMEM with 10% FBS, 1% Pen-strep, sodium pyruvate, and L-glutamine. Cells were cultured in 150 mm Corning dishes and grown to 70% confluence overnight in a 37° incubator with 5% CO₂. Dishes were transfected with 10-25 µg of Filoviridae pDNA using either a Calphos™ Mammalian Transfection Kit protocol (Clontech) or Lipofectamine™ 2000 reagent (Invitrogen) per the manufacturer's protocol and then incubated for 24-48 h. Cells were harvested with ice cold PBS, centrifuged and washed, and then pelleted for Western immunoblot or FACS analysis. Standard Western blotting was used and GP-specific MAbs for GP1 detection were generated. Data from Western immunoblotting experiments is shown in FIG. 1B. Data from FACS analysis is shown in FIG. 1C.

Animals, Vaccinations, and Challenge

[0168] Adult female C57BL/6 (H-2^b), BALB/cJ (H-2^d), and B10.Br (H-2^k) mice were purchased from The Jackson Laboratory (Bar Harbor, Me.) while Hartley guinea pigs were from Charles River (Wilmington, Mass.). All animal

experimentation was conducted following UPenn IACUC and School of Medicine Animal Facility, or NML Institutional Animal Care Committee of the PHAC and the Canadian Council on Animal Care guidelines for housing and care of laboratory animals and performed in accordance with recommendations in the Guide for the Care and Use of Laboratory Animals of NIH after pertinent review and approval by the abovementioned institutions. UPenn and NML comply with NIH policy on animal welfare, the Animal Welfare Act, and all other applicable federal, state and local laws.

[0169] Mice were immunized i.m. by needle injection with 40 µg of plasmid resuspended in water, while guinea pigs were immunized i.d., with 200 µg of each into three separate vaccination sites. Vaccinations were immediately followed by EP at the same site. Briefly, a three-pronged CELLECTRA® adaptive constant current Minimally Invasive Device was inserted approximately 2 mm i.d. (Inovio Pharmaceuticals, Inc., Blue Bell, Pa.). Square-wave pulses were delivered through a triangular 3-electrode array consisting of 26-gauge solid stainless steel electrodes and two constant-current pulses of 0.1 Amps were delivered for 52 msec/pulse separated by a 1 sec delay.

[0170] For lethal challenge studies, challenges were limited to rodent-adapted ZEBOV and MARV. Guinea pigs were challenged 28 days after the final vaccination by i.p. injection with 1,000 LD₅₀ of guinea pig-adapted ZEBOV (21.3 FFU/animal) (Richardson J S, Abou M C, Tran K N, Kumar A, Sahai B M, Kobinger G P (2011). Impact of systemic or mucosal immunity to adenovirus on ad-based Ebola virus vaccine efficacy in guinea pigs. *J Infect Dis* 204 Suppl 3: S1032-1042) or 1,000 LD₅₀ MARV-Angola (681 TCID50/animal). Briefly, the guinea-pig adapted MARV was made by the serial passage of wild-type MARV-Angola in outbred adult female Hartley guinea pigs. Seven days after inoculation, the animals were euthanized and livers were harvested and homogenized. This homogenate was then injected i.p. into naive adult guinea pigs and the process repeated until animals lost weight, gloss of hair, and succumbed to infection similar to EBOV adaptation in guinea pigs. For mouse lethal challenge studies (Kobinger G P, et al. (2006). Chimpanzee adenovirus vaccine protects against Zaire Ebola virus. *Virology* 346: 394-401), mice were injected i.p. with 200 µl of a 1,000 LD₅₀ (10 FFU/animal) of mouse-adapted ZEBOV. All animals were weighed daily and monitored for disease progression using an approved score sheet for at least 18 days for mice and 22 days for guinea pigs. All infectious work was performed in a 'Bio-safety Level 4' (BSL4) facility at NML, PHAC.

ELISA and Neutralization Assays

[0171] Antibody (Ab) titers were determined using 96-well ELISA plates coated with either sucrose-purified MARV Ozolin GP or ZGP, or with negative control sucrose-purified Nipah G protein at a concentration of 1:2,000. Briefly, the plates were then incubated for 18 h at 4° C., washed with PBS and 0.1% Tween-20, and 100 µl/sample of the sera were tested in triplicate (at dilutions 1:100, 1:400, 1:1,600, and 1:6,400 in PBS with 5% skim milk and 0.5% Tween-20). Following an incubation at 37° C. for 1 h in a moist container, the plates were washed and then 100 µl of goat anti-mouse IgG-conjugated HRP antibody (Cedarlane) was added (1:2,000 dilution) and incubated for another 37° C. for 1 h in a moist container. After a wash, 100 µl of the

ABST (2,2'-azino-bis(3-ethylbenthiiazoline-6-sulphonic acid) and peroxidase substrate (Cedarlane) was added to visualize Ab binding. Again in a moist container, the plate was incubated for 30 min at 37° C. and then later read at 405 nm. Positive binding results were characterized by being >3 SD when subtracting the positive control from the negative control serum.

[0172] The ZEBOV neutralization assay was performed. Briefly, Sera collected from immunized mice and guinea pigs were inactivated at 56° C. for 45 minutes and serial dilutions of each sample (1:20, 1:40, etc . . . , for mice and 1:50 for guinea pigs, in 50 µl of DMEM) was mixed with equal volume of ZEBOV expressing the EGFP reporter gene (ZEBOV-EGFP) (100 transducing units/well, according to EGFP expression) and incubated at 37° C. for 90 minutes. The mixture was then transferred onto sub-confluent VeroE6 cells in 96-well flat-bottomed plates and incubated for 5-10 minutes at RT. Control wells were infected with equal amounts of the ZEBOV-EGFP virus without addition of serum or with non-immune serum. 100 µl of DMEM supplemented with 20% FBS was then added to each well and plates were incubated at 37° C. in 5% CO₂ for 48 h.

[0173] Alternatively, neutralization of MARV-Angola 368 was assessed using an immunofluorescent assay. A primary rabbit anti-MARV Ab and secondary goat anti-rabbit IgG FITC-conjugated Ab was used for detection. Neutralizing Abs (NAbs) against SUDV Boniface were assayed based on cytopathic effect (CPE) on CV-1 cells. Cells were incubated with equal parts of immunized sera and SUDV Boniface for 10 days before subsequently fixed with 10% buffered formalin for 24 hours and examined under a light microscope. EGFP and FITC positive cells were counted in each well and sample dilutions showing >50% reduction in the number of green cells compared to controls scored positive for NAb. Alternatively, NAbs against SUDV-Boniface were assayed based on cytopathic effect (CPE) on CV-1 cells. All infectious work was performed in the BSL4 laboratory of NML, PHAC.

Splenocyte Isolation

[0174] Mice were sacrificed 8-11 days following the final immunization and the spleens were harvested. Briefly, spleens were placed in RPMI 1640 medium (Mediatech Inc., Manassas, Va.) supplemented with 10% FBS, 1× Anti-anti (Invitrogen), and 1× β-ME (Invitrogen). Splenocytes were isolated by mechanical disruption of the spleen using a Stomacher machine (Seward Laboratory Systems Inc., Bohemia, N.Y.), and the resulting product was filtered using a 40 µm cell strainer (BD Falcon). The cells were then treated for 5 min with ACK lysis buffer (Lonza, Switzerland) for lysis of RBCs, washed in PBS, and then resuspended in RPMI medium for use in ELISPOT or FACS assay.

ELISPOT Assays

[0175] Standard IFN γ ELISPOT assay was performed. Briefly, 96-well plates (Millipore, Billerica, Mass.) were coated with anti-mouse IFN- γ capture antibody and incubated for 24 h at 4° C. (R&D Systems, Minneapolis, Minn.). The following day, plates were washed with PBS and then blocked for 2 h with blocking buffer (1% BSA and 5% sucrose in PBS). Splenocytes (1-2×10⁵ cells/well) were plated in triplicate and stimulated overnight at 37° C. in 5% CO₂ and in the presence of either RPMI 1640 (negative

control), Con A (positive control), or GP peptides either individually (15-mers overlapping by 9 amino acids and spanning the lengths of their respective GP) or whole pooled (2.5 µg/ml final). After 18-24 h of stimulation, the plates were washed in PBS and then incubated for 24 h at 4° C. with biotinylated anti-mouse IFN- γ mAb (R&D Systems, Minneapolis, Minn.). Next, the plates were washed again in PBS, and streptavidin-alkaline phosphatase (MabTech, Sweden) was added to each well and incubated for 2 h at RT. Lastly, the plates were washed again in PBS and then BCIP/NBT Plus substrate (MabTech) was added to each well for 5-30 min for spot development. As soon as the development process was complete upon visual inspection, the plate were rinsed with distilled water and then dried overnight at RT. Spots were enumerated using an automated ELISPOT reader (Cellular Technology Ltd., Shaker Heights, Ohio).

[0176] For comprehensive analysis of T cell breadth, standard IFN γ ELISPOT was modified herein as previously described in Shedlock D J, et al. (2012). SUPRA. Identification and measurement of subdominant and immunodominant T cell epitopes were assessed by stimulating splenocytes with individual peptides as opposed to whole or matrix peptide pools; the traditional practice of pooling peptides for the sake of sample preservation, such as the use of matrix array pools, results in a reduction of assay sensitivity since total functional responses in pools containing multiple epitope-displaying peptides will effectively lower assay resolution, i.e. 'drown-out' those of lower magnitude. Thus, modified ELISPOT was performed with individual peptides (15-mers overlapping by 9 amino acids; 2.5 µg/ml final) spanning each GP immunogen. Peptides containing T cell epitopes were identified (10 AVE IFN γ + spots AND 80% animal response rate; summarized in Tables 1-6) and then later confirmed functionally and phenotypically by FACS. No shared or partial epitopes were identified, nor did FACS data or web-based epitope prediction software suggest the presence of a CD4+ or CD8+ T cell epitope that was preserved within consecutive peptides. Here, possible shared/partial T cell epitopes were addressed for all instances of contiguous peptide responses as identified by modified ELISPOT assay. Cells were stimulated individually with each of the contiguous peptides, as well as paired in combination for direct comparison, and were defined as 'shared/partial' if the combined response was not greater than either of the two individual responses. Also, it must be noted, that the epitopic response presented herein may not have been completely comprehensive since the '15-mer overlapping by 9 amino acids' algorithm for generating peptides is biased towards complete coverage of CD8 T cell epitopes which may underestimate CD4 T cell responses due to the nature of class II-restricted epitopes being longer than 15 amino acids. Lastly, amino acid similarity plots were generated using Vector NTI software and the results are shown in FIG. 4B.

Flow Cytometry

[0177] Splenocytes were added to a 96-well plate (1×10⁶ cells/well) and stimulated for 5-6 h with either individual peptides or 'Minimal Peptide Pools' (2.5 µg/ml final). Individual peptides stimulation was used for functional confirmation of all peptides identified by modified ELISPOT (Tables 1-6) as well as phenotypic characterization. Splenocytes and transfected 293 Ts were first pre-stained with

LIVE/DEAD® Fixable Violet Dead Cell Stain Kit (Invitrogen). For splenocytes, cells were surface-stained for CD19 (V450; clone 1D3), CD4 (PE-Cy7; clone RM4-5), CD8a (APC-Cy7; clone 53-6.7) and CD44 (PE-Cy5; clone IM7) (BD Biosciences, San Jose, Calif.), washed three times in PBS+1% FBS, permeabilized with BD Cytotfix/Cytoperm™ kit, and then stained intracellularly with IFN γ (APC; clone XMG1.2), TNF (FITC; clone MP6-XT22), CD3 (PE-cy5.5; clone 145-2C11), and T-bet (PE; clone 4B10) (eBioscience, San Diego, Calif.). GP expression in transfected 293T cells was assessed 24 h post-transfection. Indirect staining was performed following a 30 min incubation at 4° C. in PBS+1% FBS containing the indicated mouse-derived GP-specific polyclonal serum reagent (1:200 dilution), each produced by pooling serum from H-2^b mice immunized three times with their respective DNA vaccine or pVAX1 empty vector control. Cells were then stained with FITC-conjugated goat anti-mouse IgG (BioLegend, San Diego, Calif.), washed extensively, and then stained for MHC class I (HLA-ABC; PE-Cy7; clone G46-2.6; BD). All cells were fixed in 1% paraformaldehyde. All data was collected using a LSRII flow cytometer (BD) and analyzed using FlowJo software (Tree Star, Ashland, Oreg.). Splenocytes were gated for activated IFN γ -producing T cells that were CD3+ CD44+, CD4+ or CD8+, and negative for the B cell marker CD19 and viability dye.

[0178] FIG. 6 shows a GP-specific T cell gating. Functional and phenotypic analysis for peptides containing T cell epitopes as identified by ELISPOT was performed by FACS gating of total lymphocytes, live (LD) CD3+ cells that were negative for CD 19 and LIVE-DEAD (dump channel), singlets (excludes cell doublets), CD4+ and CD8+ cells, activated cells (CD44+), and peptide-specific IFN γ -producing T cells.

Statistical Analysis

[0179] Significance for unrooted phylogenetic trees was determined by maximum-likelihood method and verified by bootstrap analysis and significant support values (80%; 1,000 bootstrap replicates) were determined by MEGA version 5 software. Group analyses were completed by matched, two-tailed, unpaired t test and survival curves were analyzed by log-rank (Mantel-Cox) test. All values are mean \pm SEM and statistical analyses were performed by GraphPad Prism (La Jolla, Calif.).

Results

Vaccine Construction and Expression

[0180] Phylogenetic analysis revealed relative conservation among the EBOV GPs (94.4% for SUDV and 92.9% for ZEBOV), whereas the MARV GP (MGP) were more divergent (~70% conserved). Thus, a consensus strategy, as determined by alignment of the prevailing ZEBOV and SUDV GP amino acid sequences, was adopted for the EBOV GPs, while a type-matched strategy was used for MARV employing the 2005 Angola outbreak sequence which was solely responsible for the largest and deadliest MARV outbreak. Each GP transgene was genetically optimized, synthesized commercially, and then subcloned into modified pVAX1 mammalian expression vector. Altogether, a three-plasmid strategy formed the foundation for our novel polyvalent-filovirus vaccine strategy.

[0181] HEK 293T cells were transfected separately with each plasmid and GP expression was assessed by Western immunoblotting and FACS. A ~130 kDa protein was observed for each in cell lysates harvested 48 h post-transfection using species-specific anti-GP1 mAbs for detection. Results are shown in FIG. 1B. For a comparative control, recombinant vesicular stomatitis viruses (rVSV) expressing the respective GPs were loaded in concurrent lanes. Next, GP expression on the cell surface was analyzed 24 h post-transfection by indirect staining with GP-specific or control polyclonal serum by FACS. Results are shown in FIG. 1C. Cell surface expression was detected for all vaccine plasmids while little non-specific binding was observed; control serum did not react with GP-transfected cells nor did the positive sera with pVAX1-transfected cells (data shown for pEBOZ). As expected for the EBOV GPs, cell surface expression sterically occluded recognition of surface MHC class I, as well as β 1-integrin (Francica J R, Varela-Rohena A, Medvec A, Plesa G, Riley J L, Bates P (2010). Steric shielding of surface epitopes and impaired immune recognition induced by the ebola virus glycoprotein. *PLoS Pathog* 6: e1001098).

Complete Protection Against MARV and ZEBOV Challenge

[0182] To determine protective efficacy, we employed the guinea pig preclinical challenge model. Preclinical immunogenicity and efficacy studies were performed herein using the guinea pig and mouse models. The guinea pig preclinical model has been extensively used as a screening and ‘proof-of-concept’ tool for filoviral vaccine development. Although primary isolates of MARV and EBOV cause non-fatal illness in guinea pigs, a small number of passages in this host results in selection of variants able to cause fatal disease with pathological features similar to those seen in filovirus-infected primates. Similarly, mice have also been widely used for filoviral vaccine development, however, unlike the guinea pig model, immunodetection reagents for assessing immunity and T cell responses are extensively available. Infection with a murine-adapted ZEBOV (mZEBOV) results in disease characterized by high levels of virus in target organs and pathologic changes in livers and spleens akin to those found in EBOV-infected primates.

[0183] Guinea pigs (n=24) were immunized i.d. two times with 200 μ g of each plasmid (pEBOZ, pEBOS and pMARV) into three separate vaccination sites or with pVAX1 empty vector control (n=9), and then boosted with the same vaccines one month later. Animals were challenged 28 days following the second immunization with 1,000 LD₅₀ of a guinea pig-adapted MARV-Angola (gpMARV) (n=9) or ZEBOV (gpZEBOV) (n=15) in a BSL-4 facility, and then observed and weighed daily. Results are shown in FIGS. 2A-2H. Vaccinated animals were completely protected while control-vaccinated animals succumbed to gpMARV by 10 days post-challenge (n=3; P=0.0052) or to gpZEBOV by day 7 post-challenge (n=6; P=0.0008) (FIG. 2A and FIG. 2E). Additionally, vaccinated animals were protected from weight loss (FIG. 2B and FIG. 2F; P<0.0001). It is likely that vaccine-induced Abs may have contributed to protection since GP-specific Abs in pooled serum exhibited a significant increase in binding (FIG. 2C and FIG. 2G) and neutralization (FIG. 2D and FIG. 2H) titers. Experiments were performed in a BSL-4 facility and repeated twice with similar results and error bars in Figures A-2H represent

SEM. Group analyses were completed by matched, two-tailed, unpaired t test and survival curves were analyzed by log-rank (Mantel-Cox) test.

Plasmid Vaccines were Highly Immunogenic

[0184] To better characterize immune correlates as driven by the protective DNA vaccine (plasmids pEBOZ, pEBOS and pMARV, also referred to as trivalent DNA vaccine), we next employed the mouse model which has been widely used as a screening and 'proof-of-concept' tool for filoviral vaccine development and in which extensive immunodetection reagents are available. First, B cell responses were assessed in H-2^d mice (n=5/group) 20 days following each of two vaccinations, three weeks between injections with 40 µg of respective monovalent DNA vaccine. Data from these experiments is shown in FIGS. 3A-3C. While little GP-specific IgG was observed in pre-bleed control samples, as shown in FIG. 3A and FIG. 3B, a significant increase was detected in all animals following vaccination. Since purified SGP was not available, purified ZGP was used as a surrogate. IgG in SUDV-vaccinated mice bound ZGP, demonstrating the ability for vaccine-induced Ab generation as well as its capability for cross-species recognition. Additionally, seroconversion occurred in 100% of vaccinated animals after only one immunization, after which responses were significantly increased by homologous boost; AVE reciprocal endpoint dilution titres were boosted 22.1-fold in pMARV-immunized mice, and 3.4-fold and 8.6-fold in pEBOS- and pEBOZ-vaccinated animals, respectively. Samples were next assayed for neutralization of ZEBOV, SUDV-Boniface, and MARV-Angola in a BSL-4 facility. The results of the neutralization assay are shown in FIG. 3C. Significant increases in NAb titres were detected following vaccination in all animals.

[0185] Mice from two different genetic backgrounds (H-2^d and H-2^b; n=5/group) were immunized with 40 µg of respective plasmids pEBOZ, pEBOS and pMARV, homologous boosted after two weeks, and then sacrificed 8 days later for T cell analysis. Results from a novel modified ELISPOT assay to assess the comprehensive vaccine-induced T cell response, in which splenocytes were stimulated using individual peptides as opposed to matrix pools are shown in FIG. 4A. DNA vaccination induced robust IFNγ+ responses that recognized a diversity of T-cell epitopes (Tables 1-6). All positive epitope-comprising peptides were subsequently gated (See FIG. 6), confirmed, and further characterized by FACS. This modified ELISPOT approach proved extremely sensitive since background responses from control wells were low (7.2±0.2 IFNγ-producing SFC/10⁶ splenocytes in H-2^b and 9.2±0.5 in H-2^d mice). Results as shown in FIG. 4A revealed that vaccination with pMARV induced 9 measurable epitopes in H-2^b mice and 11 in H-2^d, pEBOS induced 9 and 8, and pEBOZ generated 10 and 12, in these respective strains. While five of nine (55.6%) of the epitopes from pMARV-immunized H-2^b mice were CD8+, they accounted for about 57.3% of the total MGP-specific IFNγ+ response as measured by both ELISPOT and FACS confirmation and phenotypic analysis. Similarly, only 33% and 38% of confirmed epitopes were CD8-restricted in pEBOS-immunized H-2^b and H-2^d mice, respectively. However these epitopes comprised roughly 50-90% of the total response; CD8+ T cell responses were estimated to be approximately 56% in both mouse strains while FACS estimates were 51% and 90% in H-2^b and H-2^d mice, respectively. Total CD8+ responses were lower in pEBOZ-

vaccinated animals and measured between 33% and 57% (33% for both strains by ELISPOT and 6% and 57% for H-2^b and H-2^d mice, respectively, by FACS).

[0186] A single immunodominant epitope was detected in both mouse strains receiving pEBOS where an immunodominant epitope was loosely defined as generating an IFNγ response at least two-fold over the highest subdominant epitope; pMARV induced four H-2^b-restricted immunodominant CD8+ epitopes within peptides MGP₂₅₋₃₉ (#5), MGP₆₇₋₈₁ (#12), MGP₁₈₁₋₁₉₅ (#31) and MGP₃₈₅₋₃₉₉ (#65), and an H-2^d-restricted CD4+ epitope in MGP₁₅₁₋₁₇₁ (#27). Four of these epitopes occurred within highly conserved regions of MARV GP1, including three of which were located within the putative receptor binding domain, while only one occurred within the variable mucin-like region (MGP₃₈₅₋₃₉₉ (#65)) as shown in FIG. 4B and FIG. 4C. pEBOS stimulated CD8+ epitopes occurring in SUDV GP (SGP)₁₉₋₃₃ (#4) and SGP₂₄₁₋₂₅₅ (#41) in H-2^b and H-2^d mice, respectively, both in highly conserved regions of GP1. However, pEBOZ immunization revealed three immunodominant epitopes in H-2^d mice (a CD8-restricted epitope located in the ZEBOV GP receptor binding domain (GP)₁₃₉₋₁₅₃ (#24), and two CD4-restricted epitopes ZGP₁₇₅₋₁₈₉ (#30) and ZGP₃₉₁₋₄₀₅ (#66)), occurring within the receptor binding domain and the mucin-like region, respectively. Only one immunodominant epitope was defined in H-2^b mice which contained both a CD4+ and a CD8+ epitope (#89) and occurred in a highly conserved region of GP2. Overall, diverse epitope hierarchies were consistent and reproducible in each vaccine group. Furthermore, as shown in FIG. 4D, the subdominant response comprised a significant proportion of the total response; the total AVE subdominant response as measured by the modified ELISPOT assay was approximately 12%, 62%, and 74% in pMARV-, pEBOS- and pEBOZ-immunized H-2^b mice, respectively, while responses in H-2^d mice were 47%, 50% and 34%, respectively.

[0187] Lastly, total GP-specific T cell responses were measured by FACS using stimulation with minimal peptide pools containing only confirmed epitope-comprising peptides identified. Robust responses were detected in each of the vaccinated animals and were, in a majority of cases, comprised by both activated CD4+ and CD8+ T cells. Responses were GP-specific, since little IFNγ production was observed with a control peptide (h-Clip), and correlated well with ELISPOT data. The only instance where immunization did not induce remarkable CTL as measured by FACS was in H-2^d mice vaccinated with pMARV in which no epitope identified by ELISPOT was confirmed to be CD8-restricted. Altogether, these data show that each of the vaccine plasmids was highly immunogenic in mice and yielded robust GP-specific T cell responses recognizing a diverse array of T cell epitopes including immunodominant epitopes within highly conserved regions of the GP. Furthermore, the highly diverse subdominant T cell response characterized herein might have otherwise been overlooked using traditional matrix array peptide pools for epitope identification.

[0188] T cell responses were measured for reactivity against minimal peptide pools comprised by all positively identified peptides for each respective GP by FACS. FIG. 7A shows DNA vaccine-induced T cell responses are shown from a representative animal and IFNγ-producing CD4+ (right) and CD8+(left) cells are gated. FACS plots are

shown. Incubation with h-CLIP peptide served as a negative control (Control). FIG. 7B shows results of gated cells in FIG. 7A are summarized as average % of total CD44+/IFN γ +CD4+ or CD8+ cells and error bars represent SEM. Experiments were repeated at least two times with similar results.

'Single-Dose' Protection in Mice

[0189] Vaccine efficacy against ZEBOV challenge was next assessed in the preclinical murine model. Mice were vaccinated only once due to strong NAb induction and protection data observed. Mice (H-2^k; n=10/group) were immunized with 40 μ g of the pEBOZ DNA and protection was evaluated 28 days later by challenge with 1,000 LD₅₀ of mouse-adapted ZEBOV (mZEBOV) in a BSL4 facility. While all control animals succumbed to infection by day 7 post-challenge, FIG. 5A shows DNA-vaccinated mice were completely protected (P=0.0002). In addition, as shown in FIG. 5B, control mice exhibited progressive loss of body weight until death (P<0.0001).

[0190] To better understand the mechanisms of DNA-induced protection in a 'single-dose' model, we next assessed NAb and T cell generation. NAb were assessed 25 days post-vaccination, 3 days prior to challenge, and, as shown in FIG. 5C, a significant (P<0.0001) increase was detected in all vaccinated animals (n=10/group); reciprocal endpoint dilution titers ranged from 19 to 42, 27.3 \pm 2.5.

[0191] We next evaluated the generation of ZGP-specific T cells and increased the scope of our analysis to compare responses in mice immunized with either the pEBOZ alone, or in a trivalent formulation IFN- γ production (n=5) was assessed 11 days later by FACS using whole ZGP peptide pools; the data is shown in FIG. 5D. IFN γ -producing T cells were detected in all animals and were specific for ZGP peptides since stimulation with a control peptide did not induce cytokine production. Immunization with either the monovalent or trivalent formulation induced robust IFN γ T cell responses that, when compared, were not significantly different P=0.0920).

[0192] Since CTL may be important in eliminating virus-infected cells (Warfield K L, et al. (2005). Induction of humoral and CD8+ T cell responses are required for protection against lethal Ebola virus infection. *J Immunol* 175: 1184-1191; Kalina W V, Warfield K L, Olinger G G, Bavari S (2009). Discovery of common marburgvirus protective epitopes in a BALB/c mouse model. *Viol J* 6: 132; Olinger G G, et al. (2005). Protective cytotoxic T-cell responses induced by venezuelan equine encephalitis virus replicons expressing Ebola virus proteins. *J Virol* 79: 14189-14196; Sullivan N J, et al. (2011). CD8(+) cellular immunity mediates rAd5 vaccine protection against Ebola virus infection of nonhuman primates. *Nat Med* 17: 1128-1131; and Geisbert T W, et al. (2010). Vector choice determines immunogenicity and potency of genetic vaccines against Angola Marburg virus in nonhuman primates. *J Virol* 84: 10386-10394), production of an additional effector cytokine, TNF, as well as a developmental restriction factor, T-box transcription factor TBX21 (T-bet), known to correlate with T_H1-type CTL immunity and cytotoxicity were measured and the results were as follows. For Total Cells: TNF 2.9 \pm 0.8, Tbet 13.0 \pm 1.1. For CD4+/CD44+/IFN γ + Cells: TNF 61.4 \pm 3.1, Tbet 72.6 \pm 2.0. For CD8+/CD44+/IFN γ + Cells: TNF 33.0 \pm 3.3, Tbet 992.1 \pm 1.4 (*p<0.1; ***p<0.001; ****p<0.0001). We found that ~61% and ~33% of activated

CD4+ and CD8+ T cells, respectively, also produced TNF in addition to IFN γ . Furthermore, a majority of IFN γ -producing T cells expressed high levels of T-bet; about 73% and 92% of CD8+ and CD4+ T cells, respectively, were CD44+ and produced IFN γ following ZGP peptide stimulation.

[0193] FIGS. 8A and 8B show T cell induction by 'single-dose' vaccination. T cell responses in H-2^k mice after a single pEBOZ immunization or a single trivalent vaccination, comprised by the three vaccine plasmids in separate sites, as measured by FACS are shown (a) and summarized (b) as AVE % of total CD44+/IFN γ +CD4+(purple) or CD8+(orange) cells. Pseudocolor FACS plots are from a representative animal and IFN γ -producing CD4+(right) and CD8+(left) cells are gated. Incubation with h-CLIP peptide served as a negative control (Control). Experiments were performed twice with similar results, error bars represent SEM; ns, no significance.

Discussion

[0194] We report development and evaluation of a polyvalent-filoviral vaccine in preclinical rodent immunogenicity and efficacy studies. Complete protection against challenge with gpMARV and gpZEBOV was observed following two DNA vaccine doses in guinea pigs, as well as with a 'single-dose' DNA vaccine in mice against mZEBOV. To date, genetic vaccination of guinea pigs has included either injection of naked DNA (Sullivan N J, Sanchez A, Rollin P E, Yang Z Y, Nabel G J (2000). Development of a preventive vaccine for Ebola virus infection in primates. *Nature* 408: 605-609) or DNA delivered by gene gun (Dowling W, et al. (2006). The influences of glycosylation on the antigenicity, immunogenicity, and protective efficacy of Ebola virus GP DNA vaccines. *J Virol* 81: 1821-1837; Vanderzanden L, et al. (1998). DNA vaccines expressing either the GP or NP genes of Ebola virus protect mice from lethal challenge. *Virology* 246: 134-144; and Riemenschneider J, et al (2003). Comparison of individual and combination DNA vaccines for *B. anthracis*, Ebola virus, Marburg virus and Venezuelan equine encephalitis virus. *Vaccine* 21: 4071-4080), however, either method required at least three vaccinations to achieve complete protection. Improved protection herein may be due to the induction of robust Abs since a single DNA vaccination generated GP-specific IgG binder titers that were comparable in magnitude to titers in protected animals following gene gun administration; DNA vaccination induced 3.85 and 2.18 log₁₀ ZGP and MGP-specific Ab titers, respectively, after a single administration versus 2.7 and 3.0 after three gene gun vaccinations. For comparison with an alternative 'single-dose' protective strategy in guinea pigs, an Ag-coupled virus-like particle (VLP) platform generated Ab titers that were only slightly higher than observed following DNA vaccination (Swenson D L, Warfield K L, Negley D L, Schmaljohn A, Aman M J, Bavari S (2005). Virus-like particles exhibit potential as a pan-filovirus vaccine for both Ebola and Marburg viral infections. *Vaccine* 23: 3033-3042). Furthermore, a recombinant adenovirus (rAd) approach induced ZGP-specific NAb titers that were lower than those from a single DNA vaccination (53 reciprocal endpoint dilution titer verses 88 herein) (Kobinger G P, et al. (2006). Chimpanzee adenovirus vaccine protects against Zaire Ebola virus. *Virology* 346: 394-401). Vaccination with rVSV (Jones S M, et al (2007). Assessment of a vesicular stomatitis virus-based vaccine by use of the mouse model of Ebola virus hemorrhagic fever. *J*

Infect Dis 196 Suppl 2: S404-412) generated ZGP-specific Ab titers that were similar to the current platform. Altogether, these data demonstrate that DNA vaccination was capable of inducing binding and neutralizing Abs that were comparable to non-replicating viral platforms and that these data may help, in part, to explain strong guinea pig survival data herein.

[0195] The generation of NABs by protective DNA vaccination may have benefitted by transgene-expressed mature GP structures. In vitro transfection studies confirmed that the vaccine-encoded GP were highly expressed, post-translationally cleaved (FIG. 1B), transported to the cell surface, and sterically occluded the immunodetection of cell surface molecules (FIG. 1C). Therefore, it was highly likely that the vaccine immunogens formed herein matured into heterotrimeric spikes that would otherwise be functional upon virion assembly during infection. This may be important for the generation and display of virologically-relevant neutralizing determinants which would be subsequently critical for the induction of conformation-dependent Nabs (Dowling W, et al. (2007). Influences of glycosylation on antigenicity, immunogenicity, and protective efficacy of ebola virus GP DNA vaccines. *J Virol* 81: 1821-1837; Shedlock D J, Bailey M A, Popernack P M, Cunningham J M, Burton D R, Sullivan N J (2010). Antibody-mediated neutralization of Ebola virus can occur by two distinct mechanisms. *Virology* 401: 228-235). Thus, in this regard, the expression of native anchored structures may be superior to soluble derivatives in the capacity for generating NABs (Sullivan N J, et al. (2006). Immune protection of nonhuman primates against Ebola virus with single low-dose adenovirus vectors encoding modified GPs. *PLoS Med* 3: e177; Xu L, et al. (1998). Immunization for Ebola virus infection. *Nat Med* 4: 37-42).

[0196] To better characterize T cells responses as driven by a protective vaccine, we performed immunogenicity and efficacy studies in mice and determined 'single-dose' complete protection against mZEBOV with DNA vaccination (FIGS. 5A-5D). To date, the most effective platforms conferring complete protection in this model are VLP, either with (Warfield K L, et al. (2005). Induction of humoral and CD8+ T cell responses are required for protection against lethal Ebola virus infection. *J Immunol* 175: 1184-1191; Warfield K L, Swenson D L, Olinger G G, Kalina W V, Aman M J, Bavari S (2007). Ebola virus-like particle-based vaccine protects nonhuman primates against lethal Ebola virus challenge. *J Infect Dis* 196 Suppl 2: S430-437) or without (Sun Y, et al. (2009). Protection against lethal challenge by Ebola virus-like particles produced in insect cells. *Virology* 383: 12-21) adjuvant, rAd vaccination ((Kobinger G P, et al. (2006) SUPRA; Choi J H, et al. (2012). A single sublingual dose of an adenovirus-based vaccine protects against lethal Ebola challenge in mice and guinea pigs. *Mol Pharm* 9: 156-167; Richardson J S, et al. (2009). Enhanced protection against Ebola virus mediated by an improved adenovirus-based vaccine. *PLoS One* 4: e5308), or rRABV vaccination (Blaney J E, et al. (2011). Inactivated or live-attenuated bivalent vaccines that confer protection against rabies and Ebola viruses. *J Virol* 85: 10605-10616). However, characterization of T cell responses were severely limited in these studies and were restricted to splenocyte stimulation with either two (Warfield K L, (2007), SUPRA) or one (Warfield K L, et al. (2005) SUPRA) peptides previously described to contain ZGP T cell epitopes (Warfield K L, et al. (2005) SUPRA; Olinger G G, et al. (2005)

SUPRA; Kobinger G P, et al. (2006), SUPRA; Sun Y, et al. (2009). Choi, J H, et al. (2012). Herein, we report induction of robust and broad CTL by protective vaccination as extensively analyzed by a novel modified T cell assay (FIG. 4A and Tables 1-6). In total, 52 novel T cell epitopes were identified including numerous immunodominant epitopes occurring primarily in highly conserved regions of GP. Of the 22 total ZGP epitopes identified, only 4 have been previously reported. Moreover, only one of the 20 MGP (Kalina W V, Warfield K L, Olinger G G, Bavari S (2009). Discovery of common marburgvirus protective epitopes in a BALB/c mouse model. *Virology* 6: 132) and one of 16 SGP epitopes were previously described. As such, this the most comprehensive report of preclinical GP epitopes to date, describing GP epitopes from multiple filoviruses in two different mouse genetic backgrounds.

[0197] Another novel finding resulting from these analyses was the assessment of the vaccine-induced subdominant T cell responses, which we show comprised a significant percentage of the total T cell response, widely ranging between 12%-74% (FIG. 4D). This may be particularly important since subdominant responses can significantly contribute to protection. Thus, it may prove informative in the future to determine the specific contributions of the subdominant and immunodominant epitopic T cell responses to protection. Notably, these responses may have otherwise been overlooked using traditional matrix array peptide pools for epitope identification. As such, limited epitope detection in previous studies may have been directly related to lower levels of vaccine-induced immunity, the use of less sensitive standard assays, and/or the use of peptide arrangements and/or algorithms favoring detection of immunodominant CD8+ epitopes.

[0198] Although immune correlates of protection against the filoviruses remain controversial, data generated by this highly immunogenic approach provides a unique opportunity with which to study T cell immunity as driven by a protective vaccine. DNA vaccination herein induced strong ZGP-specific T cells, a large part of which were characterized by T_H1-type multifunctional CTL expressing high levels of T-bet, also shown to correlate with T cell cytotoxicity in humans. It is clear that previous stand-alone DNA vaccine platforms capable of generating mainly humoral immune responses and cellular immunity skewed towards CD4+ T cells may likely benefit from in vivo EP delivery which has been recently demonstrated to induce potent CD8+ T cells in NHPs and the clinic. Thus, data herein are consistent with this approach as a stand-alone or prime-boost modality in NHP immunogenicity and efficacy studies. This approach offers an attractive vaccination strategy that can be quickly and inexpensively modified and/or produced for rapid response during Filoviridae bio-threat situations and outbreaks. In addition, this model approach provides an important tool for studying protective immune correlates against filoviral disease and could be applied to existing platforms to guide future strategies.

Example 2

[0199] A trivalent vaccine is provided which comprises three plasmids. The first plasmid comprises a nucleic acid sequence that encodes a Zaire ebolavirus consensus immunogen which is based upon ZEBOV CON, SEQ ID NO:1, modified to include an IgE signal peptide at the N terminus of the Zaire ebolavirus consensus immunogen. The second

plasmid comprises a nucleic acid sequence that encodes a Sudan ebolavirus consensus immunogen which is based upon SUDV CON, SEQ ID NO:2, modified to include an IgE signal peptide at the N terminus of the Sudan ebolavirus consensus immunogen. The third plasmid comprises a nucleic acid sequence that encodes a Marburg marburgvirus Angola (MARV immunogen which is based upon MARV ANG, SEQ ID NO:3, modified to include an IgE signal peptide at the N terminus of the Marburg marburgvirus Angola immunogen.

Example 3

[0200] A five plasmid vaccine is provided. The first plasmid comprises a nucleic acid sequence that encodes a Zaire ebolavirus consensus immunogen which is ZEBOV CON, SEQ ID NO:1. The second plasmid comprises a nucleic acid sequence that encodes a Sudan ebolavirus consensus immunogen which is SUDV CON, SEQ ID NO:2. The third plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:4, a Marburg marburgvirus—Ravn cluster consensus (MARV-RAV CON) using Marburg marburgvirus Ravn, Durba (09DRC99) and Uganda (02Uga07Y). The fourth plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:5, a Marburg marburgvirus—Ozolin cluster consensus (MARV-OZO CON) using Ozolin, Uganda (01Uga07), and Durba (05 and 07DRC99). The fifth plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:6, a Marburg marburgvirus—Musoke cluster consensus (MARV-MUS CON) using (Musoke, Popp, and Leiden).

Example 4

[0201] A five plasmid vaccine is provided. The first plasmid comprises a nucleic acid sequence that encodes a Zaire ebolavirus consensus immunogen which is based upon ZEBOV CON, SEQ ID NO:1, modified to include an IgE signal peptide at the N terminus of the Zaire ebolavirus consensus immunogen. The second plasmid comprises a nucleic acid sequence that encodes a Sudan ebolavirus consensus immunogen which is based upon SUDV CON, SEQ ID NO:2, modified to include an IgE signal peptide at the N terminus of the Sudan ebolavirus consensus immunogen. The third plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Rav consensus based upon SEQ ID NO:4, a Marburg marburgvirus—Ravn cluster consensus (MARV-RAV CON) using Marburg marburgvirus Ravn Durba (09DRC99) and Uganda (02Uga07Y) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Rav immunogen. The fourth plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Ozo consensus based upon SEQ ID NO:5, a Marburg marburgvirus—Ozolin cluster consensus (MARV-OZO CON) using Ozolin, Uganda (01Uga07), and Durba (05 and 07DRC99) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Ozo immunogen. The fifth plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Mus consensus based upon SEQ ID NO:6, a Marburg marburgvirus—Musoke cluster consensus (MARV-MUS CON) using (Musoke, Popp, and Leiden) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Mus immunogen.

Example 5

[0202] A six plasmid vaccine is provided. The first plasmid comprises a nucleic acid sequence that encodes a Zaire ebolavirus consensus immunogen which is ZEBOV CON, SEQ ID NO:1. The second plasmid comprises a nucleic acid sequence that encodes a Sudan ebolavirus consensus immunogen which is SUDV CON, SEQ ID NO:2. The third plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:4, a Marburg marburgvirus—Ravn cluster consensus (MARV-RAV CON) using Marburg marburgvirus Ravn, Durba (09DRC99) and Uganda (02Uga07Y). The fourth plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:5, a Marburg marburgvirus—Ozolin cluster consensus (MARV-OZO CON) using Ozolin, Uganda (01Uga07), and Durba (05 and 07DRC99). The fifth plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:6, a Marburg marburgvirus—Musoke cluster consensus (MARV-MUS CON) using (Musoke, Popp, and Leiden). The sixth plasmid comprises a nucleic acid sequence that encodes SEQ ID NO:3, a Marburg marburgvirus Angola 2005 isolate glycoproteins immunogen.

Example 6

[0203] A five plasmid vaccine is provided. The first plasmid comprises a nucleic acid sequence that encodes a Zaire ebolavirus consensus immunogen which is based upon ZEBOV CON, SEQ ID NO:1, modified to include an IgE signal peptide at the N terminus of the Zaire ebolavirus consensus immunogen. The second plasmid comprises a nucleic acid sequence that encodes a Sudan ebolavirus consensus immunogen which is based upon SUDV CON, SEQ ID NO:2, modified to include an IgE signal peptide at the N terminus of the Sudan ebolavirus consensus immunogen. The third plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Rav consensus based upon SEQ ID NO:4, a Marburg marburgvirus—Ravn cluster consensus (MARV-RAV CON) using Marburg marburgvirus Ravn Durba (09DRC99) and Uganda (02Uga07Y) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Rav immunogen. The fourth plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Ozo consensus based upon SEQ ID NO:5, a Marburg marburgvirus—Ozolin cluster consensus (MARV-OZO CON) using Ozolin, Uganda (01Uga07), and Durba (05 and 07DRC99) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Ozo immunogen. The fifth plasmid comprises a nucleic acid sequence that encodes Marburg marburgvirus Mus consensus based upon SEQ ID NO:6, a Marburg marburgvirus—Musoke cluster consensus (MARV-MUS CON) using (Musoke, Popp, and Leiden) and modified to include an IgE signal peptide at the N terminus of the consensus Marburg marburgvirus—Mus immunogen. The sixth plasmid comprises a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 isolate glycoproteins immunogen which is based upon MARV ANG, SEQ ID NO:3, modified to include an IgE signal peptide at the N terminus of the Marburg marburgvirus Angola immunogen.

TABLE 1

Plasmid Vaccine pMARV GP sequence MARV ANG							
Peptide Number	Sequence	SEQ		ELISPOT		FACS	
		ID NO:	Position	H-2	AVE	±SEM	T cell restr.
3	IQGVKTLPILEIASN	7	13-27	d	62	34	4+
5	ASNIQPQNVDVSVCSG	8	25-39	b	743	186	8+
12	SKRWAFRAGVPPKNV	9	67-81	b	694	204	4+
27	GKVFTEGNIAAMIVN	10	157-171	d	602	75	4+
28	GNIAAMIVNKTVHKM	11	163-177	b/d	126	28	8+
	GNIAAMIVNKTVHKM	12		d	30	10	4+
29	IVNKTVHKMIFSRQG	13	169-183	d	92	17	4+
30	HKMIFSRQGQYRHM	14	175-189	d	31	10	4+
31	RQGQGYRHMNLTSTN	15	181-195	b	674	112	8+
32	RHMNLTSTNKYWTSS	16	187-201	b	44	16	8+
65	LPTENPTTAKSTNST	17	385-399	b/d	398/16	107/2	4+
71	PNSTAQHLVYFRKR	18	421-435	d	29	6	4+
72	HLVYFRKRKNILWRE	19	427-441	d	145	18	4+
89	GLSWIPFPGGIEGL	20	529-543	b	26	8	4+
92	GLIKNQNNLVCLRR	21	547-561	d	29	10	4+
93	NNLVCLRRRLANQTA	22	553-567	d	34	13	4+
97	TTEERTFSLINRHAI	23	577-591	b	46	18	8+
99	HAIDFLLARWGGTCK	24	589-603	d	63	12	4+
101	TCKVLGPDCICIGIED	25	601-615	b	97	37	4+

"Epitope-containing peptides were identified by IFN γ ELISPOT (≥ 10 SFC/ 10^6 splenocytes AND $\geq 80\%$ response rate) and then confirmed by FACS ($\geq 3-5 \times 10^4$ CD3+ cells were acquired). Responses for each were further characterized by FACS (expression of CD4 and/or CD8 by CD3+/CD44+/IFN γ + cells). Predicted CD8+ epitopes are underlined (best consensus % rank by IEDB) and previously-described epitopes are referenced. Immunodominant epitopes are displayed (*).

TABLE 2

Plasmid Vaccine pEBOS GP sequence SUDV CON							
Peptide Number	Sequence	SEQ		ELISPOT		FACS	
		ID NO:	Position	H-2	AVE	±SEM	T cell restr.
4	FFVWVILFQKAFSM	26	19-33	b	310	139	8+
15	RWGFRSGVPPKVVSYS	27	85-99	b	108	59	4+
19	YNLEIKKPDGSECLP	28	109-123	b	55	25	4+
24	HKAQGTGPCPGDYAF	29	139-153	d	13	3	8+
27	GAFFLYDRLASTVIY	30	157-171	d	29	9	8+
30	NPAEGVIAFLILAKP	31	175-189	d	31	6	4+
36	SYIATSYLEYEIEINF	32	211-225	b	60	16	4+

TABLE 2-continued

		Plasmid Vaccine pEBOS					
		GP sequence SUDV CON					
Peptide		SEQ ID	ELISPOT				FACS T cell
Number	Sequence	NO:	Position	H-2	AVE	±SEM	restr.
41	FVLLDRPHTPQFLFQ	33	241-255	d	338	55	8+
78	NITTAVKTVLPQEST	34	463-477	b/d	28/105	12/18	4+
82	TGILGSLGLRKRSR	35	487-501	d	82	14	4+
83	LGLRKRSRQVNTRA	36	493-507	d	69	12	4+
89	IAWIPYFGPGAEGYI	37	529-543	b	123	40	8+/4+
97	TELRTYTILNRKAID	38	577-591	d	12	5	4+
101	CRILGPDCCIEPHDW	39	601-615	b	80	41	4+
105	QIIHDFIDNPLPNQD	40	625-639	b	28	23	4+
110	GIGITGIIIIAIIALL	41	655-669	b	27	19	8+

"Epitope-containing peptides were identified by IFN γ ELISPOT (≥ 10 SFC/10 6 splenocytes AND $\geq 80\%$ response rate) and then confirmed by FACS ($\geq 3 \times 10^4$ CD3+ cells were acquired). Responses for each were further characterized by FACS (expression of CD4 and/or CD8 by CD3+/CD44+/IFN γ + cells). Predicted CD8+ epitopes are underlined (best consensus % rank by IEDB) and previously-described epitopes are referenced. Immunodominant epitopes are displayed (*).

TABLE 3

		Plasmid Vaccine pEBOZ					
		GP sequence ZEBOV CON					
Peptide		SEQ ID	ELISPOT				FACS T cell
Number	Sequence	NO:	Position	H-2	AVE	±SEM	restr.
6	FSIPLGVIHNS [*] TLQV	42	31-45	d	78	31	8+
15	RWGFRSGVPPKVVNY	43	85-99	b	44	12	4+
19	YNLEIKKPDGSECLP	44	109-123	b	29	12	4+
24	HKVSGTGPCAGDFAF	45	139-153	d	484	85	8+
27	GAFFLYDRLASTVIY	46	157-171	d	72	18	8+
30	TFAEGVVAFLILPQA	47	175-189	d	581	85	4+
32	PQAKKDFSSHPLRE	48	187-201	b	18	6	4+
33	FFSSHPLREPVNATE	49	193-207	b	21	8	4+
40	EVDNLTYYVQLESRFT	50	235-249	d	32	17	4+
41	YVQLESRFTPQFL [*] LQ	51	241-255	d	97	23	4+
48	TTIGEWAFWETKKNL	52	283-297	d	219	70	4+
49	AFWETKKNL [*] TRKIRS	53	289-303	d	32	15	4+
50	KNL [*] TRKIRSEELSFT	54	295-309	d	105	37	4+
60	SQGREAAVSHLTTLA	55	355-369	b	16	7	4+
65	DNSTHNTPVYKLDIS	56	385-399	d	29	18	4+
66	TPVYKLDISEATQVE	57	391-405	d	371	118	4+
71	PPATTAAGPPKAENT	58	421-435	b	21	8	4+

TABLE 3-continued

		Plasmid Vaccine pEBOZ GP sequence ZEBOV CON					
Peptide		SEQ ID	ELISPOT			FACS T cell	
Number	Sequence	NO:	Position	H-2	AVE	±SEM	restr.
84	TRREAIVNAQPKCNP	59	499-513	b	12	5	8+
89	LAWIPYFGPAEAGIY	60	529-543	b	93	8	8+/4+
97	TELRTFSILNRKAID	61	577-591	b/d	14/82	4/42	8+
101	CHILGPDCCIEPHDW	62	601-615	b	96	62	4+

"Epitope-containing peptides were identified by IFN γ ELISPOT (≥ 10 SFC/10⁶ splenocytes AND $\geq 80\%$ response rate) and then confirmed by FACS ($\geq 3-5 \times 10^4$ CD3+ cells were acquired). Responses for each were further characterized by FACS (expression of CD4 and/or CD8 by CD3+/CD44+/IFN γ + cells). Predicted CD8+ epitopes are underlined (best consensus % rank by IEDB) and previously-described epitopes are referenced. Immunodominant epitopes are displayed (*).

TABLE 4

		Plasmid Vaccine pMARV GP sequence MARV ANG								
Peptide		SEQ ID	Best con % rank (IEDB)						Previously defined (80%)	
Number	Sequence	NO:	CD8+ (≥ 0.5)			CD4+ (<25)			Blast; Allele	
			D ^b	K ^b	D ^d	L ^d	I-A ^b	I-A ^d	I-E ^d	
3	IQGVKTLPILEIASN	7						12.1		
5	ASNIQPQNVDSDVCSG	8	0.4							
12	SKRWAFRAGVPPKNV	9					0.8			
27	GKVFTEGNIAAMIVN	10					12.9			
28	GNIAAMIVNKTVHKM	11	0.2					3.9		
	GNIAAMIVNKTVHKM	12	0.2					3.9		
29	IVNKTVHKMIFSRQG	13						17.2		
30	HKMIFSRQGGYRHM	14								
31	RQGGYRHMNLTST N	15		0.1				23.9		
32	RHMNLTSTNKYWTSS	16								H-2 ^d class I
65	LPTENPTAKSTNST	17					24.0			
71	PNSTAQHLVYFRRKR	18							7.5	
72	HLVYFRRKRNLWRE	19			0.3				8.3	
89	GLSWIPFFGPGIEGL	20					7.0			
92	GLIKNQNLLVCLRR	21								
93	NNLVCLRRLLANQTA	22						13.3		
97	TTEERTFSLINRHAI	23	0.1		0.4					
99	HAIDFLLARWGGTCK	24							21.8	
101	TCKVLGPDCCIGIED	25			0.4					

"Epitope-containing peptides were identified by IFN γ ELISPOT (≥ 10 SFC/10⁶ splenocytes AND $\geq 80\%$ response rate) and then confirmed by FACS ($\geq 3-5 \times 10^4$ CD3+ cells were acquired). Responses for each were further characterized by FACS (expression of CD4 and/or CD8 by CD3+/CD44+/IFN γ + cells). Predicted CD8+ epitopes are underlined (best consensus % rank by IEDB) and previously-described epitopes are referenced. Immunodominant epitopes are displayed (*).

TABLE 6-continued

		Plasmid Vaccine pEBOZ GP sequence ZEBOV CON									
		SEQ	Best con % rank (IEDB)								Previously
Peptide	ID	CD8+ (≤ 0.5)				CD4+ (<25)				defined (80%	
Number	Sequence	NO:	D ^b	K ^b	D ^d	K ^d	L ^d	I-A ^b	I-A ^d	I-E ^d	Blast; Allele
48	TTIGEWAFWETKKNL	52								12.9	
49	AFWETKKNLTRKIRS	53								22.9	
50	KNLTRKIRSEELSFT	54							22.7		
60	SQGREAAVSHLTTLA	55	0.3					23.1	3.9		
65	DNSTHNTVPYKLDIS	56									
66	TPVYKLDISEATQVE	57						22.6	5.5		
71	PPATTAAGPPKAENT	58						2.1			
84	TRREAIVNAQPKCNP	59	0.3					14.6	7.9		H-2 ^b class I
89	LAWIPYFGPAAEGII	60			0.1			0.8			H-2 ^k class I
97	TELRTFSILNRKAID	61	0.1							22.2	
101	CHILGPDCCIEPHDW	62									

"Epitope-containing peptides were identified by IFN γ ELISPOT (≥ 10 SFC/10⁶ splenocytes AND $\geq 80\%$ response rate) and then confirmed by FACS ($\geq 3 \times 10^4$ CD3+ cells were acquired). Responses for each were further characterized by FACS (expression of CD4 and/or CD8 by CD3+/CD44+/IFN γ + cells). Predicted CD8+ epitopes are underlined (best consensus % rank by IEDB) and previously-described epitopes are referenced. Immunodominant epitopes are displayed (*).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 66

<210> SEQ ID NO 1

<211> LENGTH: 676

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic consensus Zaire ebolavirus glycoprotein

<400> SEQUENCE: 1

Met Gly Val Thr Gly Ile Leu Gln Leu Pro Arg Asp Arg Phe Lys Arg
1 5 10 15

Thr Ser Phe Phe Leu Trp Val Ile Ile Leu Phe Gln Arg Thr Phe Ser
20 25 30

Ile Pro Leu Gly Val Ile His Asn Ser Thr Leu Gln Val Ser Asp Val
35 40 45

Asp Lys Leu Val Cys Arg Asp Lys Leu Ser Ser Thr Asn Gln Leu Arg
50 55 60

Ser Val Gly Leu Asn Leu Glu Gly Asn Gly Val Ala Thr Asp Val Pro
65 70 75 80

Ser Ala Thr Lys Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val
85 90 95

Val Asn Tyr Glu Ala Gly Glu Trp Ala Glu Asn Cys Tyr Asn Leu Glu
100 105 110

Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro Ala Ala Pro Asp Gly
115 120 125

-continued

Ile	Arg	Gly	Phe	Pro	Arg	Cys	Arg	Tyr	Val	His	Lys	Val	Ser	Gly	Thr
130						135					140				
Gly	Pro	Cys	Ala	Gly	Asp	Phe	Ala	Phe	His	Lys	Glu	Gly	Ala	Phe	Phe
145					150					155					160
Leu	Tyr	Asp	Arg	Leu	Ala	Ser	Thr	Val	Ile	Tyr	Arg	Gly	Thr	Thr	Phe
				165					170						175
Ala	Glu	Gly	Val	Val	Ala	Phe	Leu	Ile	Leu	Pro	Gln	Ala	Lys	Lys	Asp
			180					185					190		
Phe	Phe	Ser	Ser	His	Pro	Leu	Arg	Glu	Pro	Val	Asn	Ala	Thr	Glu	Asp
		195					200					205			
Pro	Ser	Ser	Gly	Tyr	Tyr	Ser	Thr	Thr	Ile	Arg	Tyr	Gln	Ala	Thr	Gly
	210					215					220				
Phe	Gly	Thr	Asn	Glu	Thr	Glu	Tyr	Leu	Phe	Glu	Val	Asp	Asn	Leu	Thr
225					230					235					240
Tyr	Val	Gln	Leu	Glu	Ser	Arg	Phe	Thr	Pro	Gln	Phe	Leu	Leu	Gln	Leu
				245					250						255
Asn	Glu	Thr	Ile	Tyr	Thr	Ser	Gly	Lys	Arg	Ser	Asn	Thr	Thr	Gly	Lys
			260					265						270	
Leu	Ile	Trp	Lys	Val	Asn	Pro	Glu	Ile	Asp	Thr	Thr	Ile	Gly	Glu	Trp
		275					280					285			
Ala	Phe	Trp	Glu	Thr	Lys	Lys	Asn	Leu	Thr	Arg	Lys	Ile	Arg	Ser	Glu
	290					295					300				
Glu	Leu	Ser	Phe	Thr	Ala	Val	Ser	Asn	Arg	Ala	Lys	Asn	Ile	Ser	Gly
305					310					315					320
Gln	Ser	Pro	Ala	Arg	Thr	Ser	Ser	Asp	Pro	Gly	Thr	Asn	Thr	Thr	Thr
				325					330						335
Glu	Asp	His	Lys	Ile	Met	Ala	Ser	Glu	Asn	Ser	Ser	Ala	Met	Val	Gln
			340					345					350		
Val	His	Ser	Gln	Gly	Arg	Glu	Ala	Ala	Val	Ser	His	Leu	Thr	Thr	Leu
		355					360					365			
Ala	Thr	Ile	Ser	Thr	Ser	Pro	Gln	Ser	Pro	Thr	Thr	Lys	Pro	Gly	Pro
	370					375						380			
Asp	Asn	Ser	Thr	His	Asn	Thr	Pro	Val	Tyr	Lys	Leu	Asp	Ile	Ser	Glu
385					390					395					400
Ala	Thr	Gln	Val	Glu	Gln	His	His	Arg	Arg	Thr	Asp	Asn	Asp	Ser	Thr
				405					410						415
Ala	Ser	Asp	Thr	Pro	Pro	Ala	Thr	Thr	Ala	Ala	Gly	Pro	Pro	Lys	Ala
			420					425					430		
Glu	Asn	Thr	Asn	Thr	Ser	Lys	Ser	Thr	Asp	Leu	Leu	Asp	Pro	Ala	Thr
		435				440						445			
Thr	Thr	Ser	Pro	Gln	Asn	His	Ser	Glu	Thr	Ala	Gly	Asn	Asn	Asn	Thr
	450					455						460			
His	His	Gln	Asp	Thr	Gly	Glu	Glu	Ser	Ala	Ser	Ser	Gly	Lys	Leu	Gly
465					470					475					480
Leu	Ile	Thr	Asn	Thr	Ile	Ala	Gly	Val	Ala	Gly	Leu	Ile	Thr	Gly	Gly
				485					490						495
Arg	Arg	Thr	Arg	Arg	Glu	Ala	Ile	Val	Asn	Ala	Gln	Pro	Lys	Cys	Asn
			500						505					510	
Pro	Asn	Leu	His	Tyr	Trp	Thr	Thr	Gln	Asp	Glu	Gly	Ala	Ala	Ile	Gly
		515					520					525			

-continued

Leu Ala Trp Ile Pro Tyr Phe Gly Pro Ala Ala Glu Gly Ile Tyr Thr
 530 535 540
 Glu Gly Leu Met His Asn Gln Asp Gly Leu Ile Cys Gly Leu Arg Gln
 545 550 555 560
 Leu Ala Asn Glu Thr Thr Gln Ala Leu Gln Leu Phe Leu Arg Ala Thr
 565 570 575
 Thr Glu Leu Arg Thr Phe Ser Ile Leu Asn Arg Lys Ala Ile Asp Phe
 580 585 590
 Leu Leu Gln Arg Trp Gly Gly Thr Cys His Ile Leu Gly Pro Asp Cys
 595 600 605
 Cys Ile Glu Pro His Asp Trp Thr Lys Asn Ile Thr Asp Lys Ile Asp
 610 615 620
 Gln Ile Ile His Asp Phe Val Asp Lys Thr Leu Pro Asp Gln Gly Asp
 625 630 635 640
 Asn Asp Asn Trp Trp Thr Gly Trp Arg Gln Trp Ile Pro Ala Gly Ile
 645 650 655
 Gly Val Thr Gly Val Ile Ile Ala Val Ile Ala Leu Phe Cys Ile Cys
 660 665 670
 Lys Phe Val Phe
 675

<210> SEQ ID NO 2
 <211> LENGTH: 676
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic consensus sequence Sudan
 ebolavirus glycoprotein

<400> SEQUENCE: 2

Met Glu Gly Leu Ser Leu Leu Gln Leu Pro Arg Asp Lys Phe Arg Lys
 1 5 10 15
 Ser Ser Phe Phe Val Trp Val Ile Ile Leu Phe Gln Lys Ala Phe Ser
 20 25 30
 Met Pro Leu Gly Val Val Thr Asn Ser Thr Leu Glu Val Thr Glu Ile
 35 40 45
 Asp Gln Leu Val Cys Lys Asp His Leu Ala Ser Thr Asp Gln Leu Lys
 50 55 60
 Ser Val Gly Leu Asn Leu Glu Gly Ser Gly Val Ser Thr Asp Ile Pro
 65 70 75 80
 Ser Ala Thr Lys Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val
 85 90 95
 Val Ser Tyr Glu Ala Gly Glu Trp Ala Glu Asn Cys Tyr Asn Leu Glu
 100 105 110
 Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro Pro Pro Asp Gly
 115 120 125
 Val Arg Gly Phe Pro Arg Cys Arg Tyr Val His Lys Ala Gln Gly Thr
 130 135 140
 Gly Pro Cys Pro Gly Asp Tyr Ala Phe His Lys Asp Gly Ala Phe Phe
 145 150 155 160
 Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr Arg Gly Val Asn Phe
 165 170 175
 Ala Glu Gly Val Ile Ala Phe Leu Ile Leu Ala Lys Pro Lys Glu Thr
 180 185 190

-continued

Phe Leu Gln Ser Pro Pro Ile Arg Glu Ala Val Asn Tyr Thr Glu Asn
195 200 205
Thr Ser Ser Tyr Tyr Ala Thr Ser Tyr Leu Glu Tyr Glu Ile Glu Asn
210 215 220
Phe Gly Ala Gln His Ser Thr Thr Leu Phe Lys Ile Asn Asn Asn Thr
225 230 235 240
Phe Val Leu Leu Asp Arg Pro His Thr Pro Gln Phe Leu Phe Gln Leu
245 250 255
Asn Asp Thr Ile His Leu His Gln Gln Leu Ser Asn Thr Thr Gly Lys
260 265 270
Leu Ile Trp Thr Leu Asp Ala Asn Ile Asn Ala Asp Ile Gly Glu Trp
275 280 285
Ala Phe Trp Glu Asn Lys Lys Asn Leu Ser Glu Gln Leu Arg Gly Glu
290 295 300
Glu Leu Ser Phe Glu Thr Leu Ser Leu Asn Glu Thr Glu Asp Asp Asp
305 310 315 320
Ala Thr Ser Ser Arg Thr Thr Lys Gly Arg Ile Ser Asp Arg Ala Thr
325 330 335
Arg Lys Tyr Ser Asp Leu Val Pro Lys Asp Ser Pro Gly Met Val Ser
340 345 350
Leu His Val Pro Glu Gly Glu Thr Thr Leu Pro Ser Gln Asn Ser Thr
355 360 365
Glu Gly Arg Arg Val Asp Val Asn Thr Gln Glu Thr Ile Thr Glu Thr
370 375 380
Thr Ala Thr Ile Ile Gly Thr Asn Gly Asn Asn Met Gln Ile Ser Thr
385 390 395 400
Ile Gly Thr Gly Leu Ser Ser Ser Gln Ile Leu Ser Ser Ser Pro Thr
405 410 415
Met Ala Pro Ser Pro Glu Thr Gln Thr Ser Thr Thr Tyr Thr Pro Lys
420 425 430
Leu Pro Val Met Thr Thr Glu Glu Pro Thr Thr Pro Pro Arg Asn Ser
435 440 445
Pro Gly Ser Thr Thr Glu Ala Pro Thr Leu Thr Thr Pro Glu Asn Ile
450 455 460
Thr Thr Ala Val Lys Thr Val Leu Pro Gln Glu Ser Thr Ser Asn Gly
465 470 475 480
Leu Ile Thr Ser Thr Val Thr Gly Ile Leu Gly Ser Leu Gly Leu Arg
485 490 495
Lys Arg Ser Arg Arg Gln Val Asn Thr Arg Ala Thr Gly Lys Cys Asn
500 505 510
Pro Asn Leu His Tyr Trp Thr Ala Gln Glu Gln His Asn Ala Ala Gly
515 520 525
Ile Ala Trp Ile Pro Tyr Phe Gly Pro Gly Ala Glu Gly Ile Tyr Thr
530 535 540
Glu Gly Leu Met His Asn Gln Asn Ala Leu Val Cys Gly Leu Arg Gln
545 550 555 560
Leu Ala Asn Glu Thr Thr Gln Ala Leu Gln Leu Phe Leu Arg Ala Thr
565 570 575
Thr Glu Leu Arg Thr Tyr Thr Ile Leu Asn Arg Lys Ala Ile Asp Phe
580 585 590

-continued

Leu Leu Arg Arg Trp Gly Gly Thr Cys Arg Ile Leu Gly Pro Asp Cys
 595 600 605

 Cys Ile Glu Pro His Asp Trp Thr Lys Asn Ile Thr Asp Lys Ile Asn
 610 615 620

 Gln Ile Ile His Asp Phe Ile Asp Asn Pro Leu Pro Asn Gln Asp Asn
 625 630 635 640

 Asp Asp Asn Trp Trp Thr Gly Trp Arg Gln Trp Ile Pro Ala Gly Ile
 645 650 655

 Gly Ile Thr Gly Ile Ile Ile Ala Ile Ile Ala Leu Leu Cys Val Cys
 660 665 670

 Lys Leu Leu Cys
 675

<210> SEQ ID NO 3
 <211> LENGTH: 681
 <212> TYPE: PRT
 <213> ORGANISM: Marburg virus

<400> SEQUENCE: 3

Met Lys Thr Thr Cys Leu Leu Ile Ser Leu Ile Leu Ile Gln Gly Val
 1 5 10 15

 Lys Thr Leu Pro Ile Leu Glu Ile Ala Ser Asn Ile Gln Pro Gln Asn
 20 25 30

 Val Asp Ser Val Cys Ser Gly Thr Leu Gln Lys Thr Glu Asp Val His
 35 40 45

 Leu Met Gly Phe Thr Leu Ser Gly Gln Lys Val Ala Asp Ser Pro Leu
 50 55 60

 Glu Ala Ser Lys Arg Trp Ala Phe Arg Ala Gly Val Pro Pro Lys Asn
 65 70 75 80

 Val Glu Tyr Thr Glu Gly Glu Glu Ala Lys Thr Cys Tyr Asn Ile Ser
 85 90 95

 Val Thr Asp Pro Ser Gly Lys Ser Leu Leu Leu Asp Pro Pro Thr Asn
 100 105 110

 Ile Arg Asp Tyr Pro Lys Cys Lys Thr Ile His His Ile Gln Gly Gln
 115 120 125

 Asn Pro His Ala Gln Gly Ile Ala Leu His Leu Trp Gly Ala Phe Phe
 130 135 140

 Leu Tyr Asp Arg Ile Ala Ser Thr Thr Met Tyr Arg Gly Lys Val Phe
 145 150 155 160

 Thr Glu Gly Asn Ile Ala Ala Met Ile Val Asn Lys Thr Val His Lys
 165 170 175

 Met Ile Phe Ser Arg Gln Gly Gln Gly Tyr Arg His Met Asn Leu Thr
 180 185 190

 Ser Thr Asn Lys Tyr Trp Thr Ser Ser Asn Gly Thr Gln Thr Asn Asp
 195 200 205

 Thr Gly Cys Phe Gly Thr Leu Gln Glu Tyr Asn Ser Thr Lys Asn Gln
 210 215 220

 Thr Cys Ala Pro Ser Lys Lys Pro Leu Pro Leu Pro Thr Ala His Pro
 225 230 235 240

 Glu Val Lys Leu Thr Ser Thr Ser Thr Asp Ala Thr Lys Leu Asn Thr
 245 250 255

 Thr Asp Pro Asn Ser Asp Asp Glu Asp Leu Thr Thr Ser Gly Ser Gly
 260 265 270

-continued

Cys Arg Ile Phe Thr Lys Tyr Ile Gly
675 680

<210> SEQ ID NO 4
 <211> LENGTH: 681
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic consensus sequence 1 of Marburgvirus glycoprotein

<400> SEQUENCE: 4

Met Lys Thr Ile Tyr Phe Leu Ile Ser Leu Ile Leu Ile Gln Ser Ile
1 5 10 15
 Lys Thr Leu Pro Val Leu Glu Ile Ala Ser Asn Ser Gln Pro Gln Asp
20 25 30
 Val Asp Ser Val Cys Ser Gly Thr Leu Gln Lys Thr Glu Asp Val His
35 40 45
 Leu Met Gly Phe Thr Leu Ser Gly Gln Lys Val Ala Asp Ser Pro Leu
50 55 60
 Glu Ala Ser Lys Arg Trp Ala Phe Arg Thr Gly Val Pro Pro Lys Asn
65 70 75 80
 Val Glu Tyr Thr Glu Gly Glu Glu Ala Lys Thr Cys Tyr Asn Ile Ser
85 90 95
 Val Thr Asp Pro Ser Gly Lys Ser Leu Leu Leu Asp Pro Pro Ser Asn
100 105 110
 Ile Arg Asp Tyr Pro Lys Cys Lys Thr Val His His Ile Gln Gly Gln
115 120 125
 Asn Pro His Ala Gln Gly Ile Ala Leu His Leu Trp Gly Ala Phe Phe
130 135 140
 Leu Tyr Asp Arg Val Ala Ser Thr Thr Met Tyr Arg Gly Lys Val Phe
145 150 155 160
 Thr Glu Gly Asn Ile Ala Ala Met Ile Val Asn Lys Thr Val His Arg
165 170 175
 Met Ile Phe Ser Arg Gln Gly Gln Gly Tyr Arg His Met Asn Leu Thr
180 185 190
 Ser Thr Asn Lys Tyr Trp Thr Ser Ser Asn Glu Thr Arg Arg Asn Asp
195 200 205
 Thr Gly Cys Phe Gly Ile Leu Gln Glu Tyr Asn Ser Thr Asn Asn Gln
210 215 220
 Thr Cys Ser Pro Ser Leu Lys Pro Pro Ser Leu Pro Thr Val Thr Pro
225 230 235 240
 Ser Ile His Ser Thr Asn Thr Gln Ile Asn Thr Ala Lys Ser Gly Thr
245 250 255
 Met Asn Pro Ser Ser Asp Asp Glu Asp Leu Met Ile Ser Gly Ser Gly
260 265 270
 Ser Gly Glu Gln Gly Pro His Thr Thr Leu Asn Val Val Thr Glu Gln
275 280 285
 Lys Gln Ser Ser Thr Ile Leu Ser Thr Pro Ser Leu His Pro Ser Thr
290 295 300
 Ser Gln His Glu Gln Asn Ser Thr Asn Pro Ser Arg His Ala Val Thr
305 310 315 320
 Glu His Asn Gly Thr Asp Pro Thr Thr Gln Pro Ala Thr Leu Leu Asn
325 330 335

-continued

```

Asn Thr Asn Thr Thr Pro Thr Tyr Asn Thr Leu Lys Tyr Asn Leu Ser
      340                               345                               350

Thr Pro Ser Pro Pro Thr Arg Asn Ile Thr Asn Asn Asp Thr Gln Arg
      355                               360                               365

Glu Leu Ala Glu Ser Glu Gln Thr Asn Ala Gln Leu Asn Thr Thr Leu
      370                               375                               380

Asp Pro Thr Glu Asn Pro Thr Thr Ala Gln Asp Thr Asn Ser Thr Thr
      385                               390                               395                               400

Asn Ile Ile Met Thr Thr Ser Asp Ile Thr Ser Lys His Pro Thr Asn
      405                               410                               415

Ser Ser Pro Asp Ser Ser Pro Thr Thr Arg Pro Pro Ile Tyr Phe Arg
      420                               425                               430

Lys Lys Arg Ser Ile Phe Trp Lys Glu Gly Asp Ile Phe Pro Phe Leu
      435                               440                               445

Asp Gly Leu Ile Asn Thr Glu Ile Asp Phe Asp Pro Ile Pro Asn Thr
      450                               455                               460

Glu Thr Ile Phe Asp Glu Ser Pro Ser Phe Asn Thr Ser Thr Asn Glu
      465                               470                               475                               480

Glu Gln His Thr Pro Pro Asn Ile Ser Leu Thr Phe Ser Tyr Phe Pro
      485                               490                               495

Asp Lys Asn Gly Asp Thr Ala Tyr Ser Gly Glu Asn Glu Asn Asp Cys
      500                               505                               510

Asp Ala Glu Leu Arg Ile Trp Ser Val Gln Glu Asp Asp Leu Ala Ala
      515                               520                               525

Gly Leu Ser Trp Ile Pro Phe Phe Gly Pro Gly Ile Glu Gly Leu Tyr
      530                               535                               540

Thr Ala Gly Leu Ile Lys Asn Gln Asn Asn Leu Val Cys Arg Leu Arg
      545                               550                               555                               560

Arg Leu Ala Asn Gln Thr Ala Lys Ser Leu Glu Leu Leu Leu Arg Val
      565                               570                               575

Thr Thr Glu Glu Arg Thr Phe Ser Leu Ile Asn Arg His Ala Ile Asp
      580                               585                               590

Phe Leu Leu Thr Arg Trp Gly Gly Thr Cys Lys Val Leu Gly Pro Asp
      595                               600                               605

Cys Cys Ile Gly Ile Glu Asp Leu Ser Lys Asn Ile Ser Glu Gln Ile
      610                               615                               620

Asp Lys Ile Arg Lys Asp Glu Gln Lys Glu Glu Thr Gly Trp Gly Leu
      625                               630                               635                               640

Gly Gly Lys Trp Trp Thr Ser Asp Trp Gly Val Leu Thr Asn Leu Gly
      645                               650                               655

Ile Leu Leu Leu Leu Ser Ile Ala Val Leu Ile Ala Leu Ser Cys Ile
      660                               665                               670

Cys Arg Ile Phe Thr Lys Tyr Ile Gly
      675                               680

```

<210> SEQ ID NO 5

<211> LENGTH: 681

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic consensus sequence 2 of Marburgvirus glycoprotein

-continued

<400> SEQUENCE: 5

```

Met Arg Thr Thr Cys Phe Phe Ile Ser Leu Ile Leu Ile Gln Gly Ile
1          5          10          15
Lys Thr Leu Pro Ile Leu Glu Ile Ala Ser Asn Asp Gln Pro Gln Asn
20          25          30
Val Asp Ser Val Cys Ser Gly Thr Leu Gln Lys Thr Glu Asp Val His
35          40          45
Leu Met Gly Phe Thr Leu Ser Gly Gln Lys Val Ala Asp Ser Pro Leu
50          55          60
Glu Ala Ser Lys Arg Trp Ala Phe Arg Thr Gly Val Pro Pro Lys Asn
65          70          75          80
Val Glu Tyr Thr Glu Gly Glu Glu Ala Lys Thr Cys Tyr Asn Ile Ser
85          90          95
Val Thr Asp Pro Ser Gly Lys Ser Leu Leu Leu Asp Pro Pro Thr Asn
100         105         110
Val Arg Asp Tyr Pro Lys Cys Lys Thr Ile His His Ile Gln Gly Gln
115         120         125
Asn Pro His Ala Gln Gly Ile Ala Leu His Leu Trp Gly Ala Phe Phe
130         135         140
Leu Tyr Asp Arg Ile Ala Ser Thr Thr Met Tyr Arg Gly Lys Val Phe
145         150         155         160
Thr Glu Gly Asn Ile Ala Ala Met Ile Val Asn Lys Thr Val His Lys
165         170         175
Met Ile Phe Ser Arg Gln Gly Gln Gly Tyr Arg His Met Asn Leu Thr
180         185         190
Ser Thr Asn Lys Tyr Trp Thr Ser Ser Asn Gly Thr Gln Thr Asn Asp
195         200         205
Thr Gly Cys Phe Gly Thr Leu Gln Glu Tyr Asn Ser Thr Lys Asn Gln
210         215         220
Thr Cys Ala Pro Ser Lys Thr Pro Pro Pro Pro Thr Ala Arg Pro
225         230         235         240
Glu Ile Lys Pro Thr Ser Thr Pro Thr Asp Ala Thr Arg Leu Asn Thr
245         250         255
Thr Asn Pro Asn Ser Asp Asp Glu Asp Leu Thr Thr Ser Gly Ser Gly
260         265         270
Ser Gly Glu Gln Glu Pro Tyr Thr Thr Ser Asp Ala Val Thr Lys Gln
275         280         285
Gly Leu Ser Ser Thr Met Pro Pro Thr Pro Ser Pro Gln Pro Gly Thr
290         295         300
Pro Gln Gln Gly Gly Asn Asn Thr Asn His Ser Gln Asp Ala Ala Thr
305         310         315         320
Glu Leu Asp Asn Thr Asn Thr Thr Ala Gln Pro Pro Thr Pro Ser His
325         330         335
Asn Thr Thr Thr Ile Ser Thr Asn Asn Thr Ser Lys His Asn Leu Ser
340         345         350
Thr Leu Ser Glu Pro Pro Gln Asn Thr Thr Asn Pro Asn Thr Gln Ser
355         360         365
Met Ala Thr Glu Asn Glu Lys Thr Ser Ala Pro Pro Lys Thr Thr Leu
370         375         380
Pro Pro Thr Glu Ser Pro Thr Thr Glu Lys Ser Thr Asn Asn Thr Lys
385         390         395         400

```


-continued

50					55					60					
Glu	Ala	Ser	Lys	Arg	Trp	Ala	Phe	Arg	Thr	Gly	Val	Pro	Pro	Lys	Asn
65				70						75					80
Val	Glu	Tyr	Thr	Glu	Gly	Glu	Glu	Ala	Lys	Thr	Cys	Tyr	Asn	Ile	Ser
			85						90					95	
Val	Thr	Asp	Pro	Ser	Gly	Lys	Ser	Leu	Leu	Leu	Asp	Pro	Pro	Thr	Asn
			100					105						110	
Ile	Arg	Asp	Tyr	Pro	Lys	Cys	Lys	Thr	Ile	His	His	Ile	Gln	Gly	Gln
		115					120					125			
Asn	Pro	His	Ala	Gln	Gly	Ile	Ala	Leu	His	Leu	Trp	Gly	Ala	Phe	Phe
	130					135					140				
Leu	Tyr	Asp	Arg	Ile	Ala	Ser	Thr	Thr	Met	Tyr	Arg	Gly	Arg	Val	Phe
145				150						155					160
Thr	Glu	Gly	Asn	Ile	Ala	Ala	Met	Ile	Val	Asn	Lys	Thr	Val	His	Lys
			165					170						175	
Met	Ile	Phe	Ser	Arg	Gln	Gly	Gln	Gly	Tyr	Arg	His	Met	Asn	Leu	Thr
		180						185					190		
Ser	Thr	Asn	Lys	Tyr	Trp	Thr	Ser	Asn	Asn	Gly	Thr	Gln	Thr	Asn	Asp
		195					200					205			
Thr	Gly	Cys	Phe	Gly	Ala	Leu	Gln	Glu	Tyr	Asn	Ser	Thr	Lys	Asn	Gln
	210					215					220				
Thr	Cys	Ala	Pro	Ser	Lys	Ile	Pro	Ser	Pro	Leu	Pro	Thr	Ala	Arg	Pro
225					230					235					240
Glu	Ile	Lys	Pro	Thr	Ser	Thr	Pro	Thr	Asp	Ala	Thr	Lys	Leu	Asn	Thr
			245						250					255	
Thr	Asp	Pro	Asn	Ser	Asp	Asp	Glu	Asp	Leu	Ala	Thr	Ser	Gly	Ser	Gly
			260					265					270		
Ser	Gly	Glu	Gln	Glu	Pro	His	Thr	Thr	Ser	Asp	Ala	Val	Thr	Lys	Gln
		275					280					285			
Gly	Leu	Ser	Ser	Thr	Met	Pro	Pro	Thr	Pro	Ser	Pro	Gln	Pro	Ser	Thr
	290					295					300				
Pro	Gln	Gln	Glu	Gly	Asn	Asn	Thr	Asp	His	Ser	Gln	Asp	Ala	Val	Thr
305					310					315					320
Glu	Pro	Asn	Lys	Thr	Asn	Thr	Thr	Ala	Gln	Pro	Ser	Met	Pro	Pro	His
			325						330					335	
Asn	Thr	Thr	Ala	Ile	Ser	Thr	Asn	Asn	Thr	Ser	Lys	His	Asn	Phe	Ser
			340						345				350		
Thr	Leu	Ser	Ala	Pro	Leu	Gln	Asn	Thr	Thr	Asn	Tyr	Asp	Thr	Gln	Ser
		355					360					365			
Thr	Ala	Thr	Glu	Asn	Glu	Gln	Thr	Ser	Ala	Pro	Ser	Lys	Thr	Thr	Leu
	370					375					380				
Pro	Pro	Thr	Gly	Asn	Leu	Thr	Thr	Ala	Lys	Ser	Thr	Asn	Asn	Thr	Lys
385					390					395					400
Gly	Pro	Thr	Thr	Thr	Ala	Pro	Asn	Met	Thr	Asn	Gly	His	Leu	Thr	Ser
			405						410					415	
Pro	Ser	Pro	Thr	Pro	Asn	Pro	Thr	Thr	Gln	His	Leu	Val	Tyr	Phe	Arg
			420						425				430		
Lys	Lys	Arg	Ser	Ile	Leu	Trp	Arg	Glu	Gly	Asp	Met	Phe	Pro	Phe	Leu
		435						440				445			
Asp	Gly	Leu	Ile	Asn	Ala	Pro	Ile	Asp	Phe	Asp	Pro	Val	Pro	Asn	Thr
	450					455					460				

-continued

Lys Thr Ile Phe Asp Glu Ser Ser Ser Ser Gly Ala Ser Ala Glu Glu
 465 470 475 480

Asp Gln His Ala Ser Pro Asn Ile Ser Leu Thr Leu Ser Tyr Phe Pro
 485 490 495

Asn Ile Asn Glu Asn Thr Ala Tyr Ser Gly Glu Asn Glu Asn Asp Cys
 500 505 510

Asp Ala Glu Leu Arg Ile Trp Ser Val Gln Glu Asp Asp Leu Ala Ala
 515 520 525

Gly Leu Ser Trp Ile Pro Phe Phe Gly Pro Gly Ile Glu Gly Leu Tyr
 530 535 540

Thr Ala Gly Leu Ile Lys Asn Gln Asn Asn Leu Val Cys Arg Leu Arg
 545 550 555 560

Arg Leu Ala Asn Gln Thr Ala Lys Ser Leu Glu Leu Leu Leu Arg Val
 565 570 575

Thr Thr Glu Glu Arg Thr Phe Ser Leu Ile Asn Arg His Ala Ile Asp
 580 585 590

Phe Leu Leu Thr Arg Trp Gly Gly Thr Cys Lys Val Leu Gly Pro Asp
 595 600 605

Cys Cys Ile Gly Ile Glu Asp Leu Ser Arg Asn Ile Ser Glu Gln Ile
 610 615 620

Asp Gln Ile Lys Lys Asp Glu Gln Lys Glu Gly Thr Gly Trp Gly Leu
 625 630 635 640

Gly Gly Lys Trp Trp Thr Ser Asp Trp Gly Val Leu Thr Asn Leu Gly
 645 650 655

Ile Leu Leu Leu Leu Ser Ile Ala Val Leu Ile Ala Leu Ser Cys Ile
 660 665 670

Cys Arg Ile Phe Thr Lys Tyr Ile Gly
 675 680

<210> SEQ ID NO 7
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 1

<400> SEQUENCE: 7

Ile Gln Gly Val Lys Thr Leu Pro Ile Leu Glu Ile Ala Ser Asn
 1 5 10 15

<210> SEQ ID NO 8
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 2

<400> SEQUENCE: 8

Ala Ser Asn Ile Gln Pro Gln Asn Val Asp Ser Val Cys Ser Gly
 1 5 10 15

<210> SEQ ID NO 9
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 3

-continued

<400> SEQUENCE: 9

Ser Lys Arg Trp Ala Phe Arg Ala Gly Val Pro Pro Lys Asn Val
1 5 10 15

<210> SEQ ID NO 10

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Marburg virus epitope fragment 4

<400> SEQUENCE: 10

Gly Lys Val Phe Thr Glu Gly Asn Ile Ala Ala Met Ile Val Asn
1 5 10 15

<210> SEQ ID NO 11

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Marburg virus epitope fragment 5

<400> SEQUENCE: 11

Gly Asn Ile Ala Ala Met Ile Val Asn Lys Thr Val His Lys Met
1 5 10 15

<210> SEQ ID NO 12

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Marburg virus epitope fragment 6

<400> SEQUENCE: 12

Gly Asn Ile Ala Ala Met Ile Val Asn Lys Thr Val His Lys Met
1 5 10 15

<210> SEQ ID NO 13

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Marburg virus epitope fragment 7

<400> SEQUENCE: 13

Ile Val Asn Lys Thr Val His Lys Met Ile Phe Ser Arg Gln Gly
1 5 10 15

<210> SEQ ID NO 14

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Marburg virus epitope fragment 8

<400> SEQUENCE: 14

His Lys Met Ile Phe Ser Arg Gln Gly Gln Gly Tyr Arg His Met
1 5 10 15

<210> SEQ ID NO 15

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 9

<400> SEQUENCE: 15

Arg Gln Gly Gln Gly Tyr Arg His Met Asn Leu Thr Ser Thr Asn
 1 5 10 15

<210> SEQ ID NO 16
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 10

<400> SEQUENCE: 16

Arg His Met Asn Leu Thr Ser Thr Asn Lys Tyr Trp Thr Ser Ser
 1 5 10 15

<210> SEQ ID NO 17
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 11

<400> SEQUENCE: 17

Leu Pro Thr Glu Asn Pro Thr Thr Ala Lys Ser Thr Asn Ser Thr
 1 5 10 15

<210> SEQ ID NO 18
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 12

<400> SEQUENCE: 18

Pro Asn Ser Thr Ala Gln His Leu Val Tyr Phe Arg Arg Lys Arg
 1 5 10 15

<210> SEQ ID NO 19
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 13

<400> SEQUENCE: 19

His Leu Val Tyr Phe Arg Arg Lys Arg Asn Ile Leu Trp Arg Glu
 1 5 10 15

<210> SEQ ID NO 20
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 14

<400> SEQUENCE: 20

Gly Leu Ser Trp Ile Pro Phe Phe Gly Pro Gly Ile Glu Gly Leu
 1 5 10 15

<210> SEQ ID NO 21
 <211> LENGTH: 15

-continued

<212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 15
 <400> SEQUENCE: 21

Gly Leu Ile Lys Asn Gln Asn Asn Leu Val Cys Arg Leu Arg Arg
 1 5 10 15

<210> SEQ ID NO 22
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 16
 <400> SEQUENCE: 22

Asn Asn Leu Val Cys Arg Leu Arg Arg Leu Ala Asn Gln Thr Ala
 1 5 10 15

<210> SEQ ID NO 23
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 17
 <400> SEQUENCE: 23

Thr Thr Glu Glu Arg Thr Phe Ser Leu Ile Asn Arg His Ala Ile
 1 5 10 15

<210> SEQ ID NO 24
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 18
 <400> SEQUENCE: 24

His Ala Ile Asp Phe Leu Leu Ala Arg Trp Gly Gly Thr Cys Lys
 1 5 10 15

<210> SEQ ID NO 25
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Marburg virus epitope fragment 19
 <400> SEQUENCE: 25

Thr Cys Lys Val Leu Gly Pro Asp Cys Cys Ile Gly Ile Glu Asp
 1 5 10 15

<210> SEQ ID NO 26
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 1
 <400> SEQUENCE: 26

Phe Phe Val Trp Val Ile Ile Leu Phe Gln Lys Ala Phe Ser Met
 1 5 10 15

-continued

<210> SEQ ID NO 27
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 2

<400> SEQUENCE: 27

Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val Val Ser Tyr
1 5 10 15

<210> SEQ ID NO 28
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 3

<400> SEQUENCE: 28

Tyr Asn Leu Glu Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro
1 5 10 15

<210> SEQ ID NO 29
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 4

<400> SEQUENCE: 29

His Lys Ala Gln Gly Thr Gly Pro Cys Pro Gly Asp Tyr Ala Phe
1 5 10 15

<210> SEQ ID NO 30
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 5

<400> SEQUENCE: 30

Gly Ala Phe Phe Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr
1 5 10 15

<210> SEQ ID NO 31
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 6

<400> SEQUENCE: 31

Asn Phe Ala Glu Gly Val Ile Ala Phe Leu Ile Leu Ala Lys Pro
1 5 10 15

<210> SEQ ID NO 32
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 7

<400> SEQUENCE: 32

Ser Tyr Tyr Ala Thr Ser Tyr Leu Glu Tyr Glu Ile Glu Asn Phe
1 5 10 15

-continued

<210> SEQ ID NO 33
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 8

<400> SEQUENCE: 33

Phe Val Leu Leu Asp Arg Pro His Thr Pro Gln Phe Leu Phe Gln
1 5 10 15

<210> SEQ ID NO 34
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 9

<400> SEQUENCE: 34

Asn Ile Thr Thr Ala Val Lys Thr Val Leu Pro Gln Glu Ser Thr
1 5 10 15

<210> SEQ ID NO 35
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 10

<400> SEQUENCE: 35

Thr Gly Ile Leu Gly Ser Leu Gly Leu Arg Lys Arg Ser Arg Arg
1 5 10 15

<210> SEQ ID NO 36
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 11

<400> SEQUENCE: 36

Leu Gly Leu Arg Lys Arg Ser Arg Arg Gln Val Asn Thr Arg Ala
1 5 10 15

<210> SEQ ID NO 37
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 12

<400> SEQUENCE: 37

Ile Ala Trp Ile Pro Tyr Phe Gly Pro Gly Ala Glu Gly Ile Tyr
1 5 10 15

<210> SEQ ID NO 38
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 13

<400> SEQUENCE: 38

-continued

Thr Glu Leu Arg Thr Tyr Thr Ile Leu Asn Arg Lys Ala Ile Asp
1 5 10 15

<210> SEQ ID NO 39
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 14

<400> SEQUENCE: 39

Cys Arg Ile Leu Gly Pro Asp Cys Cys Ile Glu Pro His Asp Trp
1 5 10 15

<210> SEQ ID NO 40
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 15

<400> SEQUENCE: 40

Gln Ile Ile His Asp Phe Ile Asp Asn Pro Leu Pro Asn Gln Asp
1 5 10 15

<210> SEQ ID NO 41
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Sudan ebolavirus peptide fragment 16

<400> SEQUENCE: 41

Gly Ile Gly Ile Thr Gly Ile Ile Ile Ala Ile Ile Ala Leu Leu
1 5 10 15

<210> SEQ ID NO 42
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 1

<400> SEQUENCE: 42

Phe Ser Ile Pro Leu Gly Val Ile His Asn Ser Thr Leu Gln Val
1 5 10 15

<210> SEQ ID NO 43
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 2

<400> SEQUENCE: 43

Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val Val Asn Tyr
1 5 10 15

<210> SEQ ID NO 44
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 3

-continued

<400> SEQUENCE: 44

Tyr Asn Leu Glu Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro
1 5 10 15

<210> SEQ ID NO 45

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 4

<400> SEQUENCE: 45

Gly Ala Phe Phe Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr
1 5 10 15

<210> SEQ ID NO 46

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 5

<400> SEQUENCE: 46

Gly Ala Phe Phe Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr
1 5 10 15

<210> SEQ ID NO 47

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 6

<400> SEQUENCE: 47

Thr Phe Ala Glu Gly Val Val Ala Phe Leu Ile Leu Pro Gln Ala
1 5 10 15

<210> SEQ ID NO 48

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 7

<400> SEQUENCE: 48

Pro Gln Ala Lys Lys Asp Phe Phe Ser Ser His Pro Leu Arg Glu
1 5 10 15

<210> SEQ ID NO 49

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 8

<400> SEQUENCE: 49

Phe Phe Ser Ser His Pro Leu Arg Glu Pro Val Asn Ala Thr Glu
1 5 10 15

<210> SEQ ID NO 50

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 9

<400> SEQUENCE: 50

Glu Val Asp Asn Leu Thr Tyr Val Gln Leu Glu Ser Arg Phe Thr
 1 5 10 15

<210> SEQ ID NO 51

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 10

<400> SEQUENCE: 51

Tyr Val Gln Leu Glu Ser Arg Phe Thr Pro Gln Phe Leu Leu Gln
 1 5 10 15

<210> SEQ ID NO 52

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 11

<400> SEQUENCE: 52

Thr Thr Ile Gly Glu Trp Ala Phe Trp Glu Thr Lys Lys Asn Leu
 1 5 10 15

<210> SEQ ID NO 53

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 12

<400> SEQUENCE: 53

Ala Phe Trp Glu Thr Lys Lys Asn Leu Thr Arg Lys Ile Arg Ser
 1 5 10 15

<210> SEQ ID NO 54

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 13

<400> SEQUENCE: 54

Lys Asn Leu Thr Arg Lys Ile Arg Ser Glu Glu Leu Ser Phe Thr
 1 5 10 15

<210> SEQ ID NO 55

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 14

<400> SEQUENCE: 55

Ser Gln Gly Arg Glu Ala Ala Val Ser His Leu Thr Thr Leu Ala
 1 5 10 15

<210> SEQ ID NO 56

<211> LENGTH: 15

<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 15

<400> SEQUENCE: 56

Asp Asn Ser Thr His Asn Thr Pro Val Tyr Lys Leu Asp Ile Ser
 1 5 10 15

<210> SEQ ID NO 57
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 16

<400> SEQUENCE: 57

Thr Pro Val Tyr Lys Leu Asp Ile Ser Glu Ala Thr Gln Val Glu
 1 5 10 15

<210> SEQ ID NO 58
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 17

<400> SEQUENCE: 58

Pro Pro Ala Thr Thr Ala Ala Gly Pro Pro Lys Ala Glu Asn Thr
 1 5 10 15

<210> SEQ ID NO 59
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 18

<400> SEQUENCE: 59

Thr Arg Arg Glu Ala Ile Val Asn Ala Gln Pro Lys Cys Asn Pro
 1 5 10 15

<210> SEQ ID NO 60
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 19

<400> SEQUENCE: 60

Leu Ala Trp Ile Pro Tyr Phe Gly Pro Ala Ala Glu Gly Ile Tyr
 1 5 10 15

<210> SEQ ID NO 61
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 20

<400> SEQUENCE: 61

Thr Glu Leu Arg Thr Phe Ser Ile Leu Asn Arg Lys Ala Ile Asp
 1 5 10 15

<210> SEQ ID NO 62

-continued

<211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Zaire ebolavirus consensus fragment 21

<400> SEQUENCE: 62

Cys His Ile Leu Gly Pro Asp Cys Cys Ile Glu Pro His Asp Trp
 1 5 10 15

<210> SEQ ID NO 63
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Human

<400> SEQUENCE: 63

Cys His Ile Leu Gly Pro Asp Cys Cys Ile Glu Pro His Asp Trp
 1 5 10 15

<210> SEQ ID NO 64
 <211> LENGTH: 2034
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic Zaire ebolavirus glycoprotein
 consensus

<400> SEQUENCE: 64

atgggggtca ctgggattct gcagctgcct agagatcgct tcaagcgaac ctctttcttt 60
 ctgtgggtca tcattctgtt ccagaggact tttagtatcc ctctgggcgt cattcacaat 120
 tctaccctgc aggtgagtga cgtcgataag ctgggtgtgc gggacaaact gagctccacc 180
 aaccagctga gatctgtcgg cctgaatctg gaggggaacg gagtggctac cgatgtccca 240
 agtgcaacaa agagatgggg gtttcgctca ggagtgcccc ctaaagtggg caattacgag 300
 gccggggaat gggctgagaa ttgctataac ctggaaatca agaaaccoga cggatcagag 360
 tgtctgccag ccgctccoga tgggattcgc ggattcccta gatgcagata cgtgcacaag 420
 gtcagcggca ccgggccatg tgcaggagac ttcgcctttc ataaagaagg cgccttcttt 480
 ctgtacgata gactggcttc caccgtgatc tataggggga ccacattcgc cgagggagtg 540
 gtcgcttttc tgattctgcc tcaggccaag aaagacttct tttctagtca tcctctgctg 600
 gaaccagtga acgctaccga ggaccccgac agcggctact attccactac catcagatac 660
 caggccacag gattcggcac caatgagaca gaatacctgt ttgaagtgga caacctgaca 720
 tatgtccagc tggagtctag gttcactccc cagtttctgc tgcagctgaa tgaaactatc 780
 tataccagtg gcaagcgtc aaatacaact gggagctgga tttggaaagt gaaccctgag 840
 atcgatacca caattggcga atgggccttt tgggagacca agaaaaacct gacacggaag 900
 atcagaagcg aggaactgtc cttcaccgca gtgagtaata gggccaaaaa catttcaggc 960
 cagagcccag cacgaacttc ctctgacccc gggaccaata ctaccacaga agatcacaag 1020
 atcatggcca gcgagaacag ttcagctatg gtgcaggctc actcccaggg aaggagggca 1080
 gccgtgtctc atctgactac cctggccaca atctctacta gtccccagag ccccacaact 1140
 aagcccgggc ctgacaatag caccataac acacctgtgt acaaactgga tatctccgaa 1200
 gccacccagg tcgagcagca ccacgggaga acagacaatg attccactgc atctgacacc 1260
 cctccagcaa ccacagctgc aggaccccc aaggctgaga atactaacac cagcaaaagc 1320

-continued

accgacctgc tggacccgc aactaccaca taccacaga accacagcga gacagccggg	1380
aacaataaca ctcacatca ggacaccgga gaggaatccg ccagctccgg caagctgggg	1440
ctgatcacia atactattgc tggagtggca ggactgatca caggcgggag gcgaactcga	1500
cgagaagcta ttgtgaacgc acagcccaaa tgcaatccta acctgcacta ttggactacc	1560
caggacgagg gagcagctat cggactggca tggattccat actttgggcc cgcagccgaa	1620
ggaatctata ccgagggcct gatgcataat caggatggac tgatctgtgg cctgcggcag	1680
ctggctaacg aaacaactca ggcactgcag ctgttctctc gagctaccac agagctgcgg	1740
accttagca tcctgaatcg caaggcaatt gacttctctc tgcagcgatg gggaggcaca	1800
tgccacatcc tgggaccaga ctgctgtatt gagcctcatg attggacaaa gaacatcact	1860
gacaaaattg atcagatcat tcacgacttc gtggataaaa cactgccaga tcagggggac	1920
aatgataact ggtggactgg atggagacag tggattcccg ccggcattgg cgtcacccgc	1980
gtcattattg ccgtcattgc tctgttctgt atttgtaagt tcgtgttctg ataa	2034

<210> SEQ ID NO 65

<211> LENGTH: 2034

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic Sudan ebolavirus glycoprotein consensus

<400> SEQUENCE: 65

atggagggac tgtcactgct gcagctgcct agagataagt tcaggaaaag ctcttcttt	60
gtgtgggtca tcattctgtt ccagaaggcc ttttcaatgc ccctgggctg ggtoactaat	120
agcaccctgg aagtacaga gatcgatcag ctggctgtga aggaccctt ggettaact	180
gatcagctga aaagcgtggg gctgaacctg gagggatcag gcgtcagcac tgatattcct	240
tctgcaacca agagatgggg atttcgcagc ggagtgcctc taaagtggg ctctacgaa	300
gcaggggagt gggccgaaaa ttgctataac ctggagatca agaaaccaga tggcagcgaa	360
tgtctgccac cccctccaga cggggtgcgc ggattcccca gatgcagata cgtccacaag	420
gcccagggga ccggaccttg tccaggagac tatgccttcc ataaagtgg cgcttcttt	480
ctgtacgacc gcctggctag tacagtgatc tatcgaggcg tcaatttcgc cgagggcgtg	540
atcgcttttc tgattctggc aaagccaaaa gaaaccttcc tgcagagccc tcccattagg	600
gaggccgtga actacacaga aaacacttct agttactacg ctacatccta cctggagtat	660
gaaatcgaga actttggcgc tcagcactct accacactgt tcaagattaa caataacacc	720
tttgtgctgc tggatcgcct tcatacacca cagttctctg ttcagctgaa cgacactatc	780
cacctgcatc agcagctgag caatactacc ggaaaaactga tttggacact ggacgctaat	840
atcaacgcag atattggcga gtgggccttc tgggaaaata agaaaaacct gtcgagcag	900
ctgcggggag aggaactgag ctttgaacaa ctgtccctga atgaaactga ggacgatgac	960
gccacctcaa gccgaacaac taagggcccg atctctgatc gggctaccag aaagtacagt	1020
gatctggtgc caaaagactc tcccggcatg gtgagtctgc acgtccctga aggggagacc	1080
acactgccat cccagaactc tactgagggc cggagagtgg acgtcaatac ccaggagact	1140
atcaccgaaa ctaccgcaac aatcattggc actaacggga ataacatgca gatcagcacc	1200

-continued

```

attggcacag ggctgtcctc tagtcagatt ctgtcaagct cccctacat ggccccctcc 1260
cctgagacac agacttctac aacttataca cccaagctgc ctgtgatgac cacagaggaa 1320
cccactaccc caccagaaa cagtectggg tcaacaactg aggcacccac cctgaccaca 1380
cctgaaaata tcaactaccg cgtgaaaaca gtccctgcctc aggagtctac tagtaacgga 1440
ctgatcacca gcacagtgac tggaaattctg ggcagtctgg ggctgcgcaa gcgatcaagg 1500
cgccaagtga atactcgggc taccggcaaa tgcaatccaa acctgcacta ctggaccgca 1560
caggagcagc ataacgcgcg tgggatcgcg tggattcctt acttcggacc aggcgcagag 1620
gggatctata ccgaaggact gatgcataat cagaacgccc tgggtgtgtgg cctgagacag 1680
ctggcaaatg agacaactca ggcctcgcag ctgttctga gagcaaccac agaactgagg 1740
acctatacaa tcctgaaccg gaaggccatt gatcttctgc tgcgacgatg gggcgggacc 1800
tgcagaatcc tgggaccaga ctgctgtatt gagccccacg attggaccaa gaacatcaca 1860
gacaagatca accagatcat tcatgatttc atcgacaacc cactgcccac tcaggacaac 1920
gatgacaatt ggtggaccgg atggcgacag tggattcccg caggaattgg aatcacccgga 1980
attattattg ccattattgc tctgctgtgt gtctgtaagc tgctgtgttg ataa 2034

```

<210> SEQ ID NO 66

<211> LENGTH: 2049

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic Marburgvirus glycoprotein consensus sequence

<400> SEQUENCE: 66

```

atgaaaacca cttgtctgct gatctcactg attctgattc agggcgtcaa aacactgccc 60
attctgaaa ttgcctctaa catccagcca cagaacgtgg actccgtctg ttctgggacc 120
ctgcagaaga cagaggatgt gcacctgatg ggcttcaccc tgagcgggca gaaggtcgca 180
gactcaccoc tggaaagccag caaacgatgg gcatttcggg ccggagtgcc ccctaagaac 240
gtcgagtaca ccgaaggcga ggaagccaaa acatgctata atatctccgt gactgatcct 300
agtggcaagt cactgctgct ggaccacccc accaacatta gggattaccc taagtgtaaa 360
acaatccacc atattcaggg ccagaatcca cacgctcagg ggatgcact gcatctgtgg 420
ggagccttct ttctgtacga caggattgct agcaccacaa tgatcgcgg gaaagtgttc 480
accgagggaa acatcgccgc tatgattgtg aataagacag tccacaaaat gatcttttct 540
cgccagggcc aggggtaccg acatatgaac ctgaccagta caaataagta ttggaccagc 600
tccaacggca ctcagaccaa tgacactggg tgcttcggaa ccctgcagga gtacaacagt 660
actaaaaatc agacctgtgc tccatcaaag aaaccactgc cactgcctac cgcacaccca 720
gaggtgaagc tgacaagtac ttcaaccgac gccacaaaac tgaacactac cgaccccaat 780
agtgcagatg aagatctgac aactagcgga tccggtctct gggagcagga accttatacc 840
acatccgatg cagccaccaa gcagggcctg tctagtacaa tgctccaac tccatctccc 900
cagcctagta ctccccagca gggcgggaac aataccaacc attcccaggg cgtggtcaca 960
gagccagggg agactaacac taccgccag ccctctatgc cccctcaca tacaactacc 1020
atctccacca acaatacatc taaacataac ctgagcacac cttccgtgcc aatccagaac 1080
gtactaact acaactca gtctaccgca cccgagaatg aacagacttc tgcccctagt 1140

```


-continued

aagacaactc	tgtgtccac	cgagaaccct	accacagcca	agtcaacaaa	tagcactaaa	1200
ccccacta	ccacagtgc	aaactactacc	aataagtaca	gtacctcacc	aagccccacc	1260
cctaactcca	cagcacagca	cctgggtctat	ttccggagaa	aaagaaatat	cctgtggagg	1320
gagggcgaca	tgttcccttt	tctggatggg	ctgatcaacg	ctccaattga	cttcgatcca	1380
gtgccaata	caaagactat	ctttgacgaa	tcaagctcct	ctggcgctc	tgctgaggaa	1440
gatcagcacg	cctcacccaa	cattagcctg	acactgtcct	actttcctaa	agtgaacgag	1500
aatactgcc	atagcgggga	gaacgaaaat	gactcggatg	ctgagctcgg	gatctggagc	1560
gtccaggaag	acgatctggc	tgcaggactg	tcttggatcc	cattctttgg	accggcatt	1620
gagggactgt	ataccgcgg	cctgattaag	aaccagaaca	acctgggtg	cagactgagg	1680
cgctggcca	atcagaccgc	taaatcactg	gaactgtgc	tggggctac	aactgaggaa	1740
agaacattca	gcctgatcaa	ccgacatgct	attgactttc	tgctggcacg	ctggggaggc	1800
acctgcaagg	tgctgggacc	agactgctgt	atcggcattg	aggatctgtc	tcgcaatc	1860
agtgaacaga	tcgaccagat	taagaagat	gagcagaagg	aaggaaccgg	atggggactg	1920
ggcggcaagt	ggtggaccag	cgattggggc	gtgctgacaa	acctgggaat	cctgctgctg	1980
ctgtccatcg	ccgtcctgat	tgctctgtcc	tgtatttctc	ggattttcac	taagtatatt	2040
gggtgataa						2049

1-3. (canceled)

4. A composition comprising: a first plasmid that comprises a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, or a fragment thereof; a second plasmid that comprises a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, or a fragment thereof; and one or more additional plasmid that comprises a nucleic acid sequence that encodes one or more Marburg marburgvirus envelope glycoprotein immunogen, or a fragment thereof.

5-7. (canceled)

8. The composition of claim **4** comprising: a first plasmid that comprises a nucleic acid sequence that encodes SEQ ID NO:1 or SEQ ID NO:1 linked to an IgE signal peptide; a second plasmid that comprises a nucleic acid sequence that encodes SEQ ID NO:2 or SEQ ID NO:2 linked to an IgE signal peptide; and a third plasmid that comprises a nucleic acid sequence that encodes SEQ ID NO:3 or SEQ ID NO:3 linked to an IgE signal peptide.

9. The composition of claim **4** comprising:

i) a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:1 that comprises 600 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:1 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino

acid sequence that is 95% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:1 that comprises 600 or more amino acids linked to an IgE signal peptide;

ii) a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:2 that comprises 600 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:2 that comprises 600 or more amino acids linked to an IgE signal peptide;

iii) a nucleic acid sequence that encodes a first consensus Marburg marburgvirus envelope glycoprotein immunogen, the amino acid sequence of the first consensus Marburg marburgvirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:4 (MARV CON1), a fragment of SEQ ID NO:4 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:4, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:4 that comprises 600 or more amino acids, SEQ ID NO:4 (MARV CON1)

- linked to an IgE signal peptide, a fragment of SEQ ID NO:4 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:4 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:4 that comprises 600 or more amino acids linked to an IgE signal peptide;
- iv) a nucleic acid sequence that encodes a second consensus Marburg marburgvirus envelope glycoprotein immunogen, the amino acid sequence of the second consensus Marburg marburgvirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:5 (MARV CON2), a fragment of SEQ ID NO:5 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:5, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:5 that comprises 600 or more amino acids, SEQ ID NO:5 (MARV CON2) linked to an IgE signal peptide, a fragment of SEQ ID NO:5 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:5 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:5 that comprises 600 or more amino acids linked to an IgE signal peptide; and
- v) a nucleic acid sequence that encodes a third consensus Marburg marburgvirus envelope glycoprotein immunogen, the amino acid sequence of the third consensus Marburg marburgvirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:6 (MARV CON3), a fragment of SEQ ID NO:6 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:6, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:6 that comprises 600 or more amino acids, SEQ ID NO:6 (MARV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:6 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:6 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:6 that comprises 600 or more amino acids linked to an IgE signal peptide.
10. (canceled)
11. The composition of claim 9 further comprising
- vi) a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, the amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:3 (MARV), a fragment of SEQ ID NO:3 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:3, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:3 that comprises 600 or more amino acids, SEQ ID NO:3 (MARV) linked to an IgE signal peptide, a fragment of SEQ ID NO:3 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:3 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:3 that comprises 600 or more amino acids linked to an IgE signal peptide.
12. (canceled)
13. The composition of claim 9 wherein:
- i) the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:1 that comprises 630 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:1 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:1 that comprises 630 or more amino acids linked to an IgE signal peptide;
- ii) the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein is selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:2 that comprises 630 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:2 that comprises 630 or more amino acids linked to an IgE signal peptide;
- iii) the amino acid sequence of the first consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:4 (MARV CON1), a fragment of SEQ ID NO:4 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:4, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:4 that comprises 630 or more amino acids, SEQ ID NO:4 (MARV CON1) linked to an IgE signal peptide, a fragment of SEQ ID NO:4 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:4 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:4 that comprises 630 or more amino acids linked to an IgE signal peptide;
- iv) the amino acid sequence of the second consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:5 (MARV CON2), a fragment of SEQ ID NO:5 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:5, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:5 that comprises 630 or more amino acids, SEQ ID NO:5 (MARV CON2) linked to an IgE signal peptide, a fragment of SEQ ID

NO:5 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:5 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:5 that comprises 630 or more amino acids linked to an IgE signal peptide; and

- v) the amino acid sequence of the third consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:6 (MARV CON3), a fragment of SEQ ID NO:6 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:6, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:6 that comprises 630 or more amino acids, SEQ ID NO:6 (MARV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:6 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:6 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:6 that comprises 630 or more amino acids linked to an IgE signal peptide.

14. The composition of claim **13** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that comprises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; and a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen.

15. The composition of claim **13** further comprising

- vi) a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, the amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:3 (MARV), a fragment of SEQ ID NO:3 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:3, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:3 that comprises 630 or more amino acids, SEQ ID NO:3 (MARV) linked to an IgE signal peptide, a fragment of SEQ ID NO:3 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:3 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:3 that comprises 630 or more amino acids linked to an IgE signal peptide.

16. The composition of claim **15** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that com-

prises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen, and a sixth plasmid that comprises a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen.

17. The composition of claim **9** wherein:

- i) the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:1 that comprises 660 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:1 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:1 that comprises 660 or more amino acids linked to an IgE signal peptide;
- ii) the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein is selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:2 that comprises 660 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:2 that comprises 660 or more amino acids linked to an IgE signal peptide;
- iii) the amino acid sequence of the first consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:4 (MARV CON1), a fragment of SEQ ID NO:4 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:4, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:4 that comprises 660 or more amino acids, SEQ ID NO:4 (MARV CON1) linked to an IgE signal peptide, a fragment of SEQ ID NO:4 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:4 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:4 that comprises 660 or more amino acids linked to an IgE signal peptide;
- iv) the amino acid sequence of the second consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:5 (MARV CON2), a fragment of SEQ ID NO:5 that

comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:5, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:5 that comprises 660 or more amino acids, SEQ ID NO:5 (MARV CON2) linked to an IgE signal peptide, a fragment of SEQ ID NO:5 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:5 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:5 that comprises 660 or more amino acids linked to an IgE signal peptide; and

- v) the amino acid sequence of the third consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:6 (MARV CON3), a fragment of SEQ ID NO:6 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:6, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:6 that comprises 660 or more amino acids, SEQ ID NO:6 (MARV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:6 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:6 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:6 that comprises 660 or more amino acids linked to an IgE signal peptide.

18. The composition of claim **17** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that comprises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; and a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen.

19. The composition of claim **17** further comprising

- vi) a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, the amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:3 (MARV), a fragment of SEQ ID NO:3 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:3, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:3 that comprises 660 or more amino acids, SEQ ID NO:3 (MARV) linked to an IgE signal peptide, a fragment of SEQ ID NO:3 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:3 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:3 that comprises 660 or more amino acids linked to an IgE signal peptide.

20. The composition of claim **19** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that comprises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen, and a sixth plasmid that comprises a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen.

21. The composition of claim **9** wherein:

- i) the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON) and SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide;
- ii) the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein is selected from the group consisting of: SEQ ID NO:2 (SUDV CON) and SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide;
- iii) the amino acid sequence of the first consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:4 (MARV CON1) and SEQ ID NO:4 (MARV CON1) linked to an IgE signal peptide;
- iv) the amino acid sequence of the second consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:5 (MARV CON2) and SEQ ID NO:5 (MARV CON2) linked to an IgE signal peptide; and
- v) the amino acid sequence of the third consensus Marburg marburgvirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:6 (MARV CON3) and SEQ ID NO:6 (MARV CON) linked to an IgE signal peptide.

22. The composition of claim **21** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that comprises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; and a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen.

23. The composition of claim **21** further comprising

- vi) a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen, the amino acid sequence of the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:3 (MARV) and SEQ ID NO:3 (MARV) linked to an IgE signal peptide.

24. The composition of claim **21** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; a third plasmid that comprises a nucleic acid sequence that encodes the first consensus Marburg marburgvirus envelope glycoprotein immunogen; a fourth plasmid that comprises a nucleic acid sequence that encodes the second consensus Marburg marburgvirus envelope glycoprotein immunogen; a fifth plasmid that comprises a nucleic acid sequence that encodes the third consensus Marburg marburgvirus envelope glycoprotein immunogen, and a sixth plasmid that comprises a nucleic acid sequence that encodes a Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen.

25. A composition comprising:

a nucleic acid sequence that encodes a consensus Zaire ebolavirus envelope glycoprotein immunogen, the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:1 that comprises 600 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:1 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:1 that comprises 600 or more amino acids linked to an IgE signal peptide; and

a nucleic acid sequence that encodes a consensus Sudan ebolavirus envelope glycoprotein immunogen, the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein immunogen selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 600 or more amino acids, an amino acid sequence that is 95% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:2 that comprises 600 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 600 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 95% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 95% homologous to SEQ ID NO:2 that comprises 600 or more amino acids linked to an IgE signal peptide.

26. (canceled)

27. The composition of claim **25** wherein:

the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:1 that comprises 630 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide,

a fragment of SEQ ID NO:1 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:1 that comprises 630 or more amino acids linked to an IgE signal peptide; and

the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein is selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 630 or more amino acids, an amino acid sequence that is 98% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:2 that comprises 630 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 630 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 98% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 98% homologous to SEQ ID NO:2 that comprises 630 or more amino acids linked to an IgE signal peptide.

28. The composition of claim **27** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen; and a third plasmid that comprises a nucleic acid sequence that encodes the Marburg marburgvirus Angola 2005 envelope glycoprotein immunogen.

29. The composition of claim **25** wherein:

the amino acid sequence of the consensus Zaire ebolavirus envelope glycoprotein immunogen is selected from the group consisting of: SEQ ID NO:1 (ZEBOV CON), a fragment of SEQ ID NO:1 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:1, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:1 that comprises 660 or more amino acids; SEQ ID NO:1 (ZEBOV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:1 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:1 linked to an IgE signal peptide, and a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:1 that comprises 660 or more amino acids linked to an IgE signal peptide; and

the amino acid sequence of the consensus Sudan ebolavirus envelope glycoprotein is selected from the group consisting of: SEQ ID NO:2 (SUDV CON), a fragment of SEQ ID NO:2 that comprises 660 or more amino acids, an amino acid sequence that is 99% homologous to SEQ ID NO:2, a fragment of an amino acid sequence that is 99% homologous to SEQ ID NO:2 that comprises 660 or more amino acids, SEQ ID NO:2 (SUDV CON) linked to an IgE signal peptide, a fragment of SEQ ID NO:2 that comprises 660 or more amino acids linked to an IgE signal peptide, an amino acid sequence that is 99% homologous to SEQ ID NO:2 linked to an IgE signal peptide, and a fragment of an amino acid

sequence that is 99% homologous to SEQ ID NO:2 that comprises 660 or more amino acids linked to an IgE signal peptide.

30. The composition of claim **29** comprising: a first plasmid that comprises a nucleic acid sequence that encodes the consensus Zaire ebolavirus envelope glycoprotein immunogen; and a second plasmid that comprises a nucleic acid sequence that encodes the consensus Sudan ebolavirus envelope glycoprotein immunogen.

31. The composition of claim **25** wherein: the consensus Zaire ebolavirus envelope glycoprotein immunogen has an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:1 linked to an IgE signal peptide; and the consensus Sudan ebolavirus envelope glycoprotein immunogen has an amino acid sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:2 linked to an IgE signal peptide.

32. The composition of claim **31** comprising: a first plasmid that comprises a nucleic acid sequence that encodes SEQ ID NO:1 or SEQ ID NO:1 linked to an IgE signal peptide; and a second plasmid that comprises a nucleic acid sequence that encodes SEQ ID NO:2 or SEQ ID NO:2 linked to an IgE signal peptide.

33-36. (canceled)

37. A method of treating an individual who has been diagnosed with filovirus comprising administering a therapeutically effective amount of the composition of claim **4** to an individual.

38. A method of treating an individual who has been diagnosed with filovirus selected from the group consisting of: Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire, the method comprising administering a therapeutically effective amount of the composition of claim **4** to an individual.

39. A method of preventing filovirus infection in an individual comprising administering a prophylactically effective amount of the composition of claim **4** to an individual.

40. A method of preventing filovirus infection in an individual, wherein the filovirus is selected from the group consisting of: Marburgvirus, Ebolavirus Sudan and Ebolavirus Zaire, the method comprising administering a prophylactically effective amount of the composition of claim **4** to an individual.

41. (canceled)

* * * * *