

(19)



(11)

EP 4 385 511 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

19.06.2024 Bulletin 2024/25

(51) International Patent Classification (IPC):

A61K 31/519^(2006.01)(21) Application number: **24173596.8**

(52) Cooperative Patent Classification (CPC):

**A61P 19/02; A61P 29/00; A61P 35/00;
C07D 491/147; Y02P 20/55**(22) Date of filing: **30.04.2020**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**• **WANG, Laibao****Shanghai, 201417 (CN)**• **LIU, Haihui****Shanghai, 201417 (CN)**(30) Priority: **06.06.2019 CN 201910490711**(74) Representative: **Mewburn Ellis LLP****Aurora Building****Counterslip****Bristol BS1 6BX (GB)**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

20817954.9 / 3 981 398(71) Applicant: **Hangzhou HighlightlI Pharmaceutical Co., Ltd.****Hangzhou, Zhejiang 310018 (CN)**Remarks:

• This application was filed on 30-04-2024 as a divisional application to the application mentioned under INID code 62.

• Claims filed after the date of filing of the application / after the date of receipt of the divisional application (Rule 68(4) EPC).

(72) Inventors:

- **LIANG, Congxin**
Jupiter, 33458 (US)

(54) **METHOD FOR SYNTHESIZING FUROIMIDAZOPYRIDINE COMPOUND, POLYMORPHIC SUBSTANCE AND POLYMORPHIC SUBSTANCE OF SALT**

(57) A method for synthesizing a compound 2-[(2R,5S)-5-[2-methylfuro[3,2-b]imidazo[4,5-d]pyridin-1-yl]tetrahydropyran-2-yl]acetonitrile as a selective JAK1/TYK2 kinase inhibitor. The compound is prepared by taking 7-chloro-6-nitrofuro[3,2-b]pyridine as the starting material, and by nucleophilic substitution, palladium on carbon reduction and cyclization reactions. The

present synthesis method has mild reaction conditions, high product yield and high purity, and is suitable for industrial production. A crystal form of the compound, crystal forms of the salts thereof and preparation methods thereof. The crystal form of the compound and the crystal forms of the salts thereof have good physical and chemical properties and are suitable for drug development.




Figure 1

Description

TECHNICAL FIELD

5 **[0001]** The present invention relates to the field of drug substance synthesis, specifically to the synthesis method of the compound 2-[(2*R*,5*S*)-5-[2-methylfuro[3,2-*b*]imidazo[4,5-*d*]pyridin-1-yl] tetrahydropyran-2-yl] acetonitrile (hereinafter referred to as compound I or a compound of formula I) as a selective JAK1/TYK2 kinase inhibitor. The present invention also relates to the crystal forms of compound I and its salts and their preparation methods. In addition, the present invention also relates to a pharmaceutical composition and pharmaceutical formulation comprising the crystal form of
10 compound I and/or crystal form of its salts as well as use of crystal forms of compound I and its salts in treating JAK1/TYK2-related diseases and conditions.


BACKGROUND

15 **[0002]** Protein kinases represent a family of proteins that play an important role in modulating multiple cell processes and maintaining cell functions. These kinases at least include: non-receptor tyrosine kinase, such as Janus kinase family (JAK1, JAK2, JAK3 and TYK2); receptor tyrosine kinase, such as platelet-derived growth factor receptors (PDGFR); and serine/threonine kinase, such as b-RAF.

[0003] Janus kinase family includes 4 known family members: JAK1, JAK2, JAK3 and tyrosine kinase 2 (TYK2). These
20 cytoplasmic tyrosine kinases are related to membrane cytokine receptor (such as common γ -chain receptor and glycoprotein 130 (gp130) transmembrane protein) (Murray, J. Immunol. 178 (5): 2623 - 2629, 2007). Almost 40 cytokine receptors transmit signals by the combination of these four JAK family members and their 7 downstream substrates: signal transduction activator of transcription (STAT) family members (Ghoreschi et al., Immunol Rev. 228 (1): 273 - 287, 2009). Cytokine that binds to its receptor activates JAK by trans and/or autophosphorylation. In turn, the activated JAK
25 family kinase phosphorylates a cytokine receptor, generates binding sites for proteins (such as STAT factor and other regulator) containing Src homology 2 (SH2), and JAK phosphorylation then activates them. The activated STAT enters the cell nucleus, starts to promote the expression of survival factors, cytokines, chemokines and molecules of white blood cell transport (Schindler et al., J. Biol. Chem. 282(28):20059-20063, 2007). JAK activation also causes cell proliferation by pathways mediated by phosphoinositide-3-kinase (PI3K) and protein kinase B.

[0004] JAK3 and JAK1 are components of common γ -chain cytokine receptor compound, and blocking any one of the
30 two can inhibit signal transduction of inflammatory cytokines (interleukin (IL)-2, 4, 7, 9, 15 and 21) (Ghoreschi et al., Immunol. Rev. 228 (1): 273 - 287, 2009). In contrast, other pathologically related cytokines (such as IL-6) only depend on JAK1. Therefore, JAK1 blocking inhibits signal transduction of many proinflammatory cytokines (Guschin et al, EMBO J. 14 (7): 1421 - 1429, 1995). Clinical efficacy of IL-6 receptor neutralizing antibody - tocilizumab in rheumatoid arthritis (RA) has been observed (Maini et al, Arthritis Rheum. 54(9):2817-2829, 2006).

[0005] International patent application WO2018067422A1 discloses 1*H*-furo[3,2-*b*]imidazo[4,5-*d*]pyridine derivatives as selective JAK1 kinase inhibitors and their preparation method, including compound I and its preparation method. The synthesis route is as follows:
35




[0006] Biological tests indicate that compound I is a potent and selective JAK1 inhibitor, demonstrates selective inhibition of IL-6-induced STAT3 phosphorylation and does not show selective inhibition of thrombopoietin-induced STAT3 phosphorylation. However, international patent application WO2018067422A1 doesn't disclose the biological activities of TYK2. In addition, the disclosed preparation method of compound I involves high temperature, produces too many impurities, and has low yield, hereby is not suitable for large scale production. Therefore, it is necessary to develop a preparation method of compound I with milder reaction conditions, higher product yield, higher purity and is suitable for large scale/industrial production.

[0007] Currently, there is no report of the crystal form of compound I and its salts. Comprehensive and systematic polymorph screening and the selection of a crystal form that is most suitable for development are one of the indispensable and important research contents. Accordingly, it is necessary to further screen the crystal form of compound I and its salts, develop a crystal form with good stability, low hygroscopicity and is suitable for large scale production and provides more and better choices for subsequent developments of drugs.

SUMMARY

[0008] The objective of the present invention is to provide a method for preparing a compound of formula I (that is, compound I) with mild reaction conditions, high product yield and purity and is suitable for industrial production. The synthesis route of the method is as follows:



the method comprises the following steps:

step 1:

adding ethanol, a compound of formula IV, a compound of formula V and DIPEA to a reaction container, starting stirring;

heating to raise the temperature to 65 - 90 °C, maintaining the temperature and stirring overnight;

terminating the reaction and lowering the temperature of the system to 15 - 30 °C;

adding water to the system dropwise and keeping stirring;

filtering and washing the filter cake;

drying the filter cake to obtain a compound of formula III;

step 2:

EP 4 385 511 A2

adding tetrahydrofuran, the compound of formula III obtained in step 1 and palladium on carbon to a reaction container;

5 purging the system with nitrogen and then hydrogen;

maintaining the temperature between 20 - 35 °C and stirring for 16 - 120 hours under 0.1 - 1.0 MPa hydrogen pressure;

10 after the reaction is completed, filtering the reaction liquid and washing the filter cake;

combining the filtrate and concentrating to obtain a compound of formula II concentrate;

step 3:

15 adding the compound of formula II concentrate or the compound of formula II obtained in step 2, and trimethyl orthoacetate and tetrahydrofuran to a reaction container; heating the material system in the reaction container until tetrahydrofuran reflux; adding pyridine hydrochloride to the reaction container, reacting the obtained material system for 4 - 20 hours at a temperature between 50 - 90 °C, separating and purifying to obtain a compound of formula I.

20 **[0009]** In some of the embodiments of the above step 1:

the volume mass ratio (mL/g) of ethanol to the compound of formula IV is between 5:1 and 20:1, preferably 10:1;

25 the molar ratio of the compound of formula IV, the compound of formula V and DIPEA is 1:1-1.1:2-3, preferably 1:1.01:2.2;

after starting stirring, under nitrogen protection, heating to raise the temperature to 65 - 90°C, preferably 70 - 90 °C, more preferably 70 - 80 °C, maintaining the temperature and stirring for 5 - 16 hours, preferably 10 - 16 hours;

30 after terminating the reaction, lowering the temperature of the system to 15 - 25 °C;

the volume mass ratio (mL/g) of the water added to the system to the compound of formula IV is between 10:1 and 20:1, preferably 15:1;

35 after adding water to the system, stirring for 2 - 6 hours, preferably 4 hours, at a temperature between 0 - 30 °C, preferably 5 - 15 °C, more preferably 5 - 10 °C;

the filter cake is washed with ethanol aqueous solution, the volume ratio (mL/mL) of ethanol to water in the ethanol aqueous solution is between 1:1 and 1:2, preferably 1:1.5 - 1:2; the

40 volume mass ratio (mL/g) of the ethanol aqueous solution to the compound of formula IV is between 2:1 and 10:1, preferably 2:1 - 5:1, more preferably 2:1 - 3:1;

drying the filter cake under vacuum or with an air blower at a temperature between 45 - 55°C, preferably 50 °C.

45 **[0010]** In some of the embodiments of the above step 2:

the volume mass ratio (mL/g) of tetrahydrofuran to the compound of formula III is between 10:1 and 70:1, preferably 20:1 - 70:1;

50 the palladium on carbon is 5% Pd/C, 50% water wet, the mass ratio (g/g) of the palladium on carbon to the compound of formula III is between 0.15:1 and 0.16:1, preferably 0.15:1;

maintaining the temperature between 25 - 35 °C and stirring for 24 - 96 hours under 0.5 - 1.0 MPa hydrogen pressure;

55 the compound of formula II concentrate obtained by combining the filtrate and concentrating is a compound of formula II in tetrahydrofuran, wherein the volume mass ratio (mL/g) of the tetrahydrofuran for washing to the compound of formula II is between 2:1 and 4:1, preferably 2:1 - 3:1 (the mass of the compound of formula II calculated according to a 100% yield of step 2); preferably, exchanging the compound of formula II in tetrahydrofuran with

EP 4 385 511 A2

ethanol to obtain a compound of formula II in ethanol, wherein the volume mass ratio (mL/g) of ethanol to the compound of formula II is between 2:1 and 5:1, preferably 2:1 - 4:1, more preferably 2:1 - 3:1 (the mass of the compound of formula II calculated according to a 100% yield of step 2).

5 **[0011]** In some of the embodiments of the above step 3, the volume mass ratio (mL:mg) of tetrahydrofuran to the compound of formula II in the compound of formula II concentrate is between 1.5:1 and 5.0:1; or in some of the embodiments of the above step 3, the volume mass ratio (mL:mg) of tetrahydrofuran to the compound of formula II is between 1.5:1 and 5.0:1.

10 **[0012]** In some of the embodiments of the above step 3, exchanging the compound of formula II concentrate with toluene, tetrahydrofuran or methyl tertiary-butyl ether for subsequent steps; in some of the embodiments, the volume mass ratio (mL:mg) of toluene, tetrahydrofuran or methyl tertiary-butyl ether used for exchanging to the compound of formula II concentrate is between 2.0:1 and 4.0:1;

15 in some of the embodiments of the above step 3, the molar ratio of the compound of formula II in the compound of formula II concentrate to trimethyl orthoacetate is between 3.0:1 and 3.5:1; or in some of the embodiments of the above step 3, the molar ratio of the compound of formula II to trimethyl orthoacetate is between 3.0:1 and 3.5:1;

20 in some of the embodiments of the above step 3, the molar ratio of the compound of formula II in the compound of formula II concentrate to pyridine hydrochloride is between 0.2:1 and 0.3:1; or in some of the embodiments of the above step 3, the molar ratio of the compound of formula II to pyridine hydrochloride is between 0.2:1 and 0.3:1;

25 in some of the embodiments of the above step 3, after adding the compound of formula II concentrate or the compound of formula II, and trimethyl orthoacetate and the solvent to the reaction container, under nitrogen protection, heating the material system in the reaction container until the solvent reflux;


after adding pyridine hydrochloride to the reactor, under nitrogen protection, reacting the material system for 4 - 20 hours, preferably 5 - 15 hours at a temperature between 50 - 90 °C, preferably 65 - 75 °C;

30 in some of the embodiments of the above step 3, after the reaction is completed, purifying the product with a solvent selected from the group consisting of water, methanol, ethanol, methyl tertiary-butyl ether and any combinations thereof.

35 **[0013]** In some of the embodiments of the above step 3, separating and purifying the compound of formula I obtained by column chromatography, wherein the eluent is a mixed solution of ethyl acetate and n-heptane ($V_{EA}:V_{n\text{-heptane}} = 1:1 - 1:0$, mL/mL);

in some of the embodiments of the above step 3, the obtained compound of formula I is dried under vacuum or with an air blower between 50 - 55 °C.

40 **[0014]** Another objective of the present invention is to provide a crystal form of a compound of formula I, which is named as crystal form 1 of a compound of formula I hereinafter.



50
I

55 **[0015]** Crystal form 1 of a compound of formula I has an X-ray powder diffraction pattern showing characteristic peaks at 2theta (2θ) angles of 13.4° ± 0.2°, 17.6° ± 0.2° and 21.9° ± 0.2°.

[0016] In some embodiments, the X-ray powder diffraction pattern of crystal form 1 of a compound of formula I shows characteristic peaks at 2theta angles of 9.0° ± 0.2°, 13.4° ± 0.2°, 17.6° ± 0.2°, 18.1° ± 0.2°, 21.9° ± 0.2° and 27.3° ± 0.2°.

[0017] In some embodiments, the X-ray powder diffraction pattern of crystal form 1 of a compound of formula I shows

EP 4 385 511 A2

characteristic peaks at 2θ angles of $9.0^\circ \pm 0.2^\circ$, $10.4^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $21.9^\circ \pm 0.2^\circ$, $24.1^\circ \pm 0.2^\circ$ and $27.3^\circ \pm 0.2^\circ$.

[0018] Non-restrictively, the X-ray powder diffraction data of crystal form 1 of a compound of formula I of the invention are shown in Table 1.

Table 1

Diffraction angle 2θ ($^\circ$)	d (Å)	Relative diffraction intensity I% (based on peak height)
9.0	9.84	25.0
10.4	8.51	20.9
13.4	6.60	63.7
17.6	5.02	37.5
18.1	4.91	27.3
18.7	4.74	20.8
21.9	4.05	100.0
24.1	3.69	16.4
27.3	3.26	25.8

[0019] Non-restrictively, the X-ray powder diffraction (XRPD) pattern of crystal form 1 of a compound of formula I of the invention is shown in Figure 1.

[0020] Non-restrictively, the differential scanning calorimetry (DSC) thermogram of crystal form 1 of a compound of formula I of the invention is shown in Figure 2. The DSC thermogram shows that the initial melting point of crystal form 1 of a compound of formula I of the invention is 173.38°C .

[0021] Non-restrictively, the thermogravimetric analysis (TGA) thermogram of crystal form 1 of a compound of formula I of the invention is shown in Figure 3. The TGA thermogram shows that there is only a 0.42% weight loss of crystal form 1 of a compound of formula I of the invention from 25°C to 162°C . Crystal form 1 of a compound of formula I doesn't contain crystal water or solvent.

[0022] Non-restrictively, the dynamic vapour sorption (DVS) isotherm plot of crystal form 1 of a compound of formula I of the invention is shown in Figure 4. The DVS isotherm plot shows a 13.86% weight gain of crystal form 1 of a compound of formula I of the invention by moisture absorption from 0%RH to 95%RH, indicating that the sample is hygroscopic. The moisture absorption curve during desorption exhibited hysteresis, combining with XRPD pattern of the sample before and after the DVS test (see Figure 5 for the XRPD pattern after the test), show that after moisture absorption of crystal form 1 of a compound of formula I, the crystal form changes.

[0023] The present invention provides a preparation method of crystal form 1 of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I crude product with methanol, stirring between $40 - 60^\circ\text{C}$ for 0.5 - 2 hours, cooling to $5 - 15^\circ\text{C}$, stirring for 15 minutes - 1 hour, filtering, washing the filter cake with MTBE, and drying to obtain crystal form 1 of a compound of formula I.

[0024] In some embodiments, after dissolving the compound of formula I crude product with methanol, stirring at 50°C for 1 hour, cooling to 10°C , stirring for 0.5 hour, filtering, washing the filter cake with MTBE, and drying the filter cake under vacuum at 50°C for 16 hours to obtain crystal form 1 of a compound of formula I.

[0025] In some embodiments, the volume ratio of methanol to MTBE is between 3:1 - 2:1, preferably 8:3;

[0026] In some embodiments, after dissolving the compound of formula I crude product with methanol, adding a silicon-based metal eliminator and an activated carbon to the system.\

[0027] Another objective of the present invention is to provide crystal forms of a compound of formula I, specifically, a crystal form of a hydrochloride, a crystal form of a sulfate, a crystal form of a phosphate, a crystal form of a mesylate, a crystal form of a hydrobromide, a crystal form of a fumarate, a crystal form of a benzene sulfonate, a crystal form of a citrate, a crystal form of a L-(+)-tartrate (which is named as tartrate for short in the present application) of a compound of formula I, they are named as crystal form A of a hydrochloride, crystal form B of a hydrochloride, crystal form C of a hydrochloride, crystal form D of a sulfate, crystal form E of a phosphate, crystal form F of a phosphate, crystal form G of a mesylate, crystal form H of a hydrobromide, crystal form J of a hydrobromide, crystal form K of a hydrobromide, crystal form L of a fumarate, crystal form M of a benzene sulfonate, crystal form N of a citrate, crystal form O of a tartrate of a compound of formula I respectively hereinafter.

[0028] Crystal form A of a hydrochloride of a compound of formula I of the present invention has an X-ray powder

EP 4 385 511 A2

diffraction pattern showing characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$ and $20.9^\circ \pm 0.2^\circ$.

[0029] In some embodiments, the X-ray powder diffraction pattern of crystal form A of a hydrochloride of a compound of formula I shows characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $20.9^\circ \pm 0.2^\circ$, $23.5^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

[0030] In some embodiments, the X-ray powder diffraction pattern of crystal form A of a hydrochloride of a compound of formula I shows characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $10.6^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $14.0^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $20.9^\circ \pm 0.2^\circ$, $23.5^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

[0031] Non-restrictively, the X-ray powder diffraction data of crystal form A of a hydrochloride of a compound of formula I of the invention are shown in Table 2.

Table 2

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
7.3	12.12	91.5
10.6	8.30	30.4
12.1	7.29	91.2
12.8	6.92	31.1
14.0	6.32	22.5
18.7	4.73	41.0
20.9	4.24	100.0
23.5	3.79	40.9
24.0	3.70	42.3

[0032] Non-restrictively, the XRPD pattern of crystal form A of a hydrochloride of a compound of formula I of the invention is shown in Figure 6.

[0033] Non-restrictively, the DSC thermogram of crystal form A of a hydrochloride of a compound of formula I of the invention is shown in Figure 7.

[0034] The present invention provides a preparation method of the crystal form A of a hydrochloride of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding hydrochloric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form A of a hydrochloride of a compound of formula I.

[0035] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and then is dissolved in acetone;

[0036] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 50 mg/mL, preferably 20 mg/mL;

[0037] In some embodiments, the concentration of the hydrochloric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

[0038] In some embodiments, after adding the hydrochloric acid in acetone, keeping stirring at a room temperature for 4 - 48 hours, preferably 24 hours;

[0039] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0040] Crystal form B of a hydrochloride of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$ and $22.6^\circ \pm 0.2^\circ$.

[0041] In some embodiments, the X-ray powder diffraction pattern of the crystal form B of a hydrochloride of a compound of formula I shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $10.2^\circ \pm 0.2^\circ$, $11.5^\circ \pm 0.2^\circ$, $18.0^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $25.9^\circ \pm 0.2^\circ$.

[0042] In some embodiments, the X-ray powder diffraction pattern of the crystal form B of a hydrochloride of a compound of formula I shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $10.2^\circ \pm 0.2^\circ$, $11.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $14.5^\circ \pm 0.2^\circ$, $18.0^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $25.9^\circ \pm 0.2^\circ$.

[0043] Non-restrictively, the X-ray powder diffraction data of crystal form B of a hydrochloride of a compound of formula I of the invention are shown in Table 3:

EP 4 385 511 A2

Table 3

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
7.2	12.26	100.0
10.2	8.62	41.7
11.5	7.72	41.5
14.1	6.29	35.3
14.5	6.09	25.6
18.0	4.93	48.9
20.0	4.44	69.9
22.6	3.93	62.3
25.9	3.43	41.8

[0044] Non-restrictively, the XRPD pattern of crystal form B of a hydrochloride of a compound of formula I of the invention is shown in Figure 8.

[0045] Non-restrictively, the DSC thermogram of crystal form B of a hydrochloride of a compound of formula I of the invention is shown in Figure 9.

[0046] The present invention provides a preparation method of crystal form B of a hydrochloride of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding hydrochloric acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form B of a hydrochloride of a compound of formula I.

[0047] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate;

[0048] In some embodiments, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL;

[0049] In some embodiments, the concentration of the hydrochloric acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL;

[0050] In some embodiments, after adding the hydrochloric acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

[0051] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0052] Crystal form C of a hydrochloride of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of 10.7° ± 0.2°, 21.5° ± 0.2° and 24.3° ± 0.2°.

[0053] In some embodiments, the X-ray powder diffraction pattern of the crystal form C of a hydrochloride of a compound of formula I shows characteristic peaks at 2θ angles of 5.3° ± 0.2°, 10.7° ± 0.2°, 21.5° ± 0.2°, 24.3° ± 0.2°, and 30.4° ± 0.2°.

[0054] Non-restrictively, the X-ray powder diffraction data of crystal form C of a hydrochloride of a compound of formula I of the invention are shown in Table 4:

Table 4

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
5.3	16.5427	10.1
10.7	8.2756	100.0
21.5	4.1379	52.3
24.3	3.6532	20.1
30.4	2.9351	15.8

[0055] Non-restrictively, the XRPD pattern of crystal form C of a hydrochloride of a compound of formula I of the invention is shown in Figure 10.

[0056] Non-restrictively, the DSC thermogram of crystal form C of a hydrochloride of a compound of formula I of the invention is shown in Figure 11.

EP 4 385 511 A2

[0057] Conducting recrystallization or crystal transformation of crystal form A of a hydrochloride of a compound of formula I with a solvent to obtain crystal form C of a hydrochloride of a compound of formula I, wherein the solvent is selected from the group consisting of methanol, acetonitrile, n-heptane, methyl ethyl ketone and any combinations thereof.

[0058] In some embodiments, mixing the solvent with crystal form A of a hydrochloride of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form C of a hydrochloride of a compound of formula I.

[0059] In some embodiments, adding the solvent to a container with crystal form A of a hydrochloride of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form C of a hydrochloride of a compound of formula I.

[0060] In some embodiments, the duration of the stirring is 4 - 48 hours, preferably 24 hours.

[0061] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0062] Crystal form D of a sulfate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $6.0^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$ and $25.2^\circ \pm 0.2^\circ$.

[0063] In some embodiments, the X-ray powder diffraction pattern of the crystal form D of a sulfate of a compound of formula I shows characteristic peaks at 2theta angles of $6.0^\circ \pm 0.2^\circ$, $12.3^\circ \pm 0.2^\circ$, $17.5^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $25.2^\circ \pm 0.2^\circ$.

[0064] Non-restrictively, the X-ray powder diffraction data of crystal form D of a sulfate of a compound of formula I of the invention are shown in Table 5:

Table 5

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
6.0	14.59	100.0
12.3	7.19	12.0
17.5	5.06	9.0
22.8	3.90	36.3
25.0	3.53	17.1

[0065] Non-restrictively, the XRPD pattern of crystal form D of a sulfate of a compound of formula I of the invention is shown in Figure 12.

[0066] Non-restrictively, the DSC thermogram of crystal form D of a sulfate of a compound of formula I of the invention is shown in Figure 13.

[0067] The present invention provides a preparation method of crystal form D of a sulfate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding sulfuric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form D of a sulfate of a compound of formula I.

[0068] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone.

[0069] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL.

[0070] In some embodiments, the concentration of the sulfuric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL.

[0071] In some embodiments, after adding sulfuric acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours.

[0072] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0073] Crystal form E of a phosphate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $6.2^\circ \pm 0.2^\circ$, $15.5^\circ \pm 0.2^\circ$, $17.4^\circ \pm 0.2^\circ$ and $24.6^\circ \pm 0.2^\circ$.

[0074] Non-restrictively, the X-ray powder diffraction data of crystal form E of a phosphate of a compound of formula I of the invention are shown in Table 6:

Table 6

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
6.2	14.24	100.0

EP 4 385 511 A2

(continued)

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
15.5	5.70	22.4
17.4	5.09	16.9
24.6	3.62	16.1

[0075] Non-restrictively, the XRPD pattern of crystal form E of a phosphate of a compound of formula I of the invention is shown in Figure 14.

[0076] Non-restrictively, the DSC thermogram of crystal form E of a phosphate of a compound of formula I of the invention is shown in Figure 15.

[0077] The present invention provides a preparation method of crystal form E of a phosphate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding phosphoric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form E of a phosphate of a compound of formula I.

[0078] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone.

[0079] In some embodiments, the molar ratio of the compound of formula I to phosphoric acid is between 1:1.0 - 1:1.5.

[0080] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL.

[0081] In some embodiments, the concentration of the phosphoric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL.

[0082] In some embodiments, after adding the phosphoric acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours.

[0083] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0084] In some embodiments, conducting recrystallization or crystal transformation of crystal form E of a phosphate of a compound of formula I with a solvent, and the product is still crystal form E of a phosphate of a compound of formula I, wherein the solvent is selected from the group consisting of methanol, acetonitrile, n-heptane, methyl ethyl ketone and any combinations thereof.

[0085] In some embodiments, during recrystallization or crystal transformation, mixing the solvent and crystal form E of a phosphate of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying.

[0086] In some embodiments, during recrystallization or crystal transformation, the duration of the stirring is between 4 - 48 hours, preferably stirring overnight;

[0087] In some embodiments, during recrystallization or crystal transformation, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0088] Crystal form F of a phosphate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$ and $22.6^\circ \pm 0.2^\circ$.

[0089] In some embodiments, the X-ray powder diffraction pattern of the crystal form F of a phosphate of a compound of formula I shows characteristic peaks at 2θ angles of $11.6^\circ \pm 0.2^\circ$, $14.8^\circ \pm 0.2^\circ$, $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $26.6^\circ \pm 0.2^\circ$.

[0090] In some embodiments, the X-ray powder diffraction pattern of the crystal form F of a phosphate of a compound of formula I shows characteristic peaks at 2θ angles of $11.1^\circ \pm 0.2^\circ$, $11.6^\circ \pm 0.2^\circ$, $14.8^\circ \pm 0.2^\circ$, $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$, $21.2^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $26.6^\circ \pm 0.2^\circ$.

[0091] Non-restrictively, the X-ray powder diffraction data of crystal form F of a phosphate of a compound of formula I of the invention are shown in Table 7:

Table 7

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
11.1	7.97	15.9
11.6	7.65	35.0
14.8	5.97	32.4
16.6	5.33	69.7

EP 4 385 511 A2

(continued)

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
17.2	5.16	49.1
21.2	4.19	25.3
22.6	3.93	100.0
26.6	3.34	31.9

[0092] Non-restrictively, the XRPD pattern of crystal form F of a phosphate of a compound of formula I of the invention is shown in Figure 16.

[0093] Non-restrictively, the DSC thermogram of crystal form F of a phosphate of a compound of formula I of the invention is shown in Figure 17. The DSC thermogram shows that the initial melting point of crystal form F of a phosphate of a compound of formula I of the invention is 198.78 °C.

[0094] Non-restrictively, the DVS isotherm plot of crystal form F of a phosphate of a compound of formula I of the invention is shown in Figure 18. The DVS isotherm plot shows a 6.5% weight gain of crystal form F of a phosphate of a compound of formula I of the invention by moisture absorption from 0%RH to 95%RH. At a humidity of 85%RH, crystal form F of a phosphate of a compound of formula I reaches a weight gain of 0.72%; at 70%RH, crystal form F of a phosphate of a compound of formula I reaches a weight gain of 1.95%. After the moisture absorption, crystal form F of a phosphate of a compound of formula I doesn't change (see Figure 19 for the XRPD pattern after the moisture absorption).

[0095] The present invention provides a preparation method of crystal form F of a phosphate of a compound of formula I, specifically, the method is described as follows:

dissolving crystal form E of a phosphate of a compound of formula I with a first solvent to obtain crystal form E of a phosphate of a compound of formula I in the first solvent, adding an anti-solvent, stirring, collecting the solid, and drying to obtain crystal form F of a phosphate of a compound of formula I; or

in some embodiments, the first solvent is a solvent that can dissolve crystal form E of a phosphate of a compound of formula I, preferably methanol; the anti-solvent is a solvent that is difficult to dissolve crystal form E of a phosphate of a compound of formula I, preferably isopropyl acetate.

[0096] In some embodiments, the first solvent is added in an amount that can completely dissolve crystal form E of a phosphate of a compound of formula I.

[0097] In some embodiments, the anti-solvent is used to dilute crystal form E of a phosphate of a compound of formula I in the first solvent at 5 - 15 folds, preferably 10 folds.

[0098] In some embodiments, after dissolving crystal form E of a phosphate of a compound of formula I with the first solvent, adding a small amount of seed crystals of crystal form F of a phosphate of a compound of formula I until the system is slightly turbid, then adding the anti-solvent.

[0099] In some embodiments, after adding the anti-solvent, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours.

[0100] In some embodiments, collecting the solid by centrifugation and drying under vacuum between 30 - 60 °C, preferably 50 °C.

[0101] Crystal form G of a mesylate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of 8.6° ± 0.2°, 19.9° ± 0.2° and 24.9° ± 0.2°.

[0102] In some embodiments, the X-ray powder diffraction pattern of the crystal form G of a mesylate of a compound of formula I shows characteristic peaks at 2θ angles of 8.6° ± 0.2°, 18.1° ± 0.2°, 18.6° ± 0.2°, 19.9° ± 0.2°, 24.0° ± 0.2° and 24.9° ± 0.2°.

[0103] Non-restrictively, the X-ray powder diffraction data of crystal form G of a mesylate of a compound of formula I of the invention are shown in Table 8.

Table 8

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
8.6	10.32	100.0
18.1	4.88	56.9

EP 4 385 511 A2

(continued)

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
18.6	4.76	62.7
19.9	4.456	87.1
24.0	3.71	23.0
25.0	3.57	76.1

[0104] Non-restrictively, the XRPD pattern of a crystal form G of a mesylate of a compound of formula I of the invention is shown in Figure 20.

[0105] Non-restrictively, the DSC thermogram of crystal form G of a mesylate of a compound of formula I of the invention is shown in Figure 21. The DSC thermogram shows that the initial melting point of crystal form G of a mesylate of a compound of formula I of the invention is 218.78 °C.

[0106] Non-restrictively, the DVS isotherm plot of crystal form G of a mesylate of a compound of formula I of the invention is shown in Figure 22.

[0107] The present invention provides a preparation method of crystal form G of a mesylate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding methanesulfonic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form G of a mesylate of a compound of formula I.

[0108] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone.

[0109] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL.

[0110] In some embodiments, the concentration of the methanesulfonic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL.

[0111] In some embodiments, after adding the methanesulfonic acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours.

[0112] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0113] Crystal form H of a hydrobromide of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of 7.2° ± 0.2°, 20.7° ± 0.2° and 24.0° ± 0.2°.

[0114] In some embodiments, the X-ray powder diffraction pattern of the crystal form H of a hydrobromide of a compound of formula I shows characteristic peaks at 2θ angles of 7.2° ± 0.2°, 17.9° ± 0.2°, 18.8° ± 0.2°, 20.7° ± 0.2° and 24.0° ± 0.2°.

[0115] In some embodiments, the X-ray powder diffraction pattern of the crystal form H of a hydrobromide of a compound of formula I shows characteristic peaks at 2θ angles of 7.2° ± 0.2°, 11.9° ± 0.2°, 17.0° ± 0.2°, 17.9° ± 0.2°, 18.8° ± 0.2°, 20.7° ± 0.2°, 24.0° ± 0.2° and 27.5° ± 0.2°.

[0116] Non-restrictively, the X-ray powder diffraction data of crystal form H of a hydrobromide of a compound of formula I of the invention are shown in Table 9.

Table 9

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
7.2	12.34	100.0
11.9	7.40	22.1
17.0	5.21	29.1
17.9	4.94	50.0
18.8	4.73	45.7
20.7	4.28	73.1
24.0	3.70	57.3
27.5	3.24	18.9

[0117] Non-restrictively, the XRPD pattern of crystal form H of a hydrobromide of a compound of formula I of the invention is shown in Figure 23.

[0118] Non-restrictively, the DSC thermogram of crystal form H of a hydrobromide of a compound of formula I of the invention is shown in Figure 24.

[0119] The present invention provides a preparation method of crystal form H of a hydrobromide of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding hydrobromic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form H of a hydrobromide of a compound of formula I.

[0120] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone.

[0121] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL.

[0122] In some embodiments, the concentration of the hydrobromic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL.

[0123] In some embodiments, after adding the hydrobromic acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hour.

[0124] In some embodiments, collecting the solid by centrifugation and drying under vacuum overnight at 30 - 60 °C.

[0125] In some embodiments, conducting recrystallization or crystal transformation of crystal form H of a hydrobromide of a compound of formula I with a solvent, and the product is still crystal form H of a hydrobromide, wherein the solvent is selected from the group consisting of acetonitrile, methyl ethyl ketone and any combinations thereof.

[0126] In some embodiments, the recrystallization or crystal transformation comprises the following steps:

mixing one or both of acetonitrile and methyl ethyl ketone with crystal form H of a hydrobromide of a compound of formula I, to prepare suspension, stirring at room temperature, centrifuging, collecting the solid and drying;

preferably, adding one or both of acetonitrile and methyl ethyl ketone to a container with crystal form H of a hydrobromide of a compound of formula I, to prepare suspension, stirring at room temperature, centrifuging, collecting the solid and drying.

[0127] Crystal form J of a hydrobromide of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2 θ angles of $6.2^\circ \pm 0.2^\circ$ and $15.0^\circ \pm 0.2^\circ$.

[0128] Non-restrictively, the X-ray powder diffraction data of crystal form J of a hydrobromide of a compound of formula I of the invention are shown in Table 10.

Table 10

Diffraction angle 2 θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
6.2	14.23	51.2
15.0	5.88	43.0

[0129] Non-restrictively, the XRPD pattern of crystal form J of a hydrobromide of a compound of formula I of the invention is shown in Figure 25.

[0130] The present invention provides a preparation method of crystal form J of a hydrobromide of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding hydrobromic acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form J of a hydrobromide of a compound of formula I.

[0131] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate.

[0132] In some embodiments, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL.

[0133] In some embodiments, the concentration of the hydrobromic acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL.

[0134] In some embodiments, after adding the hydrobromic acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

[0135] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0136] Crystal form K of a hydrobromide of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $17.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$ and $24.2^\circ \pm 0.2^\circ$.

[0137] The X-ray powder diffraction pattern of the crystal form K of a hydrobromide of a compound of formula I shows characteristic peaks at 2theta angles of $17.1^\circ \pm 0.2^\circ$, $20.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $24.2^\circ \pm 0.2^\circ$ and $28.8^\circ \pm 0.2^\circ$.

[0138] The X-ray powder diffraction pattern of the crystal form K of a hydrobromide of a compound of formula I shows characteristic peaks at 2theta angles of $9.5^\circ \pm 0.2^\circ$, $17.1^\circ \pm 0.2^\circ$, $20.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $24.2^\circ \pm 0.2^\circ$, $27.7^\circ \pm 0.2^\circ$ and $28.8^\circ \pm 0.2^\circ$.

[0139] Non-restrictively, the X-ray powder diffraction data of crystal form K of a hydrobromide of a compound of formula I of the invention are shown in Table 11:

Table 11

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
9.5	9.26	23.3
17.1	5.19	55.6
20.1	4.40	33.9
22.0	4.04	100.0
22.6	3.94	31.1
23.6	3.76	30.6
24.2	3.67	36.7
27.7	3.21	19.3
28.8	3.10	32.9

[0140] Non-restrictively, the XRPD pattern of crystal form K of a hydrobromide of a compound of formula I of the invention is shown in Figure 26.

[0141] Non-restrictively, the DSC thermogram of crystal form K of a hydrobromide of a compound of formula I of the invention is shown in Figure 27.

[0142] Non-restrictively, the DVS isotherm plot of crystal form K of a hydrobromide of a compound of formula I of the invention is shown in Figure 28. The DVS isotherm plot shows a 11.84% weight gain of crystal form K of a hydrobromide of a compound of formula I of the invention by moisture absorption from 0%RH to 95%RH, indicating that the sample is hygroscopic. The moisture absorption curve during desorption exhibited hysteresis, combining with the XRPD pattern of the sample before and after the DVS test (see Figure 29 for the XRPD pattern after the test), showing that the crystal form of crystal form K of a hydrobromide of a compound of formula I changed after moisture absorption.

[0143] The present invention provides a preparation method of the crystal form K of a hydrobromide of a compound of formula I, specifically, the method is described as follows:

conducting recrystallization or crystal transformation of crystal form H of a hydrobromide of a compound of formula I with n-heptane, to obtain crystal form K of a hydrobromide of a compound of formula I.

[0144] In some embodiments, mixing n-heptane with crystal form H of a hydrobromide of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form K of a hydrobromide of a compound of formula I.

[0145] In some embodiments, adding n-heptane to a container with crystal form H of a hydrobromide of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form K of a hydrobromide of a compound of formula I.

[0146] In some embodiments, the duration of the stirring is 4 - 48 hours, preferably 24 hours;

[0147] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0148] Crystal form L of a fumarate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

[0149] The X-ray powder diffraction pattern of the crystal form L of a fumarate of a compound of formula I shows characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $15.7^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

[0150] The X-ray powder diffraction pattern of the crystal form L of a fumarate of a compound of formula I shows characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $15.7^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $23.2^\circ \pm 0.2^\circ$, $23.8^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

[0151] Non-restrictively, the X-ray powder diffraction data of crystal form L of a fumarate of a compound of formula I

of the invention are shown in Table 12.

Table 12

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
6.1	14.56	100.0
13.4	6.60	12.6
15.7	5.63	15.8
16.3	5.43	68.1
22.6	3.93	11.4
23.2	3.84	11.3
23.8	3.74	9.6
26.4	3.37	37.6

[0152] Non-restrictively, the XRPD pattern of crystal form L of a fumarate of a compound of formula I of the invention is shown in Figure 30.

[0153] Non-restrictively, the DSC thermogram of crystal form L of a fumarate of a compound of formula I of the invention is shown in Figure 31.

[0154] The present invention provides a preparation method of crystal form L of a fumarate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with a solvent to obtain a compound of formula I in the solvent, and adding fumaric acid in ethanol to the compound of formula I in the solvent under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form L of a fumarate of a compound of formula I.

[0155] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in a solvent, wherein the solvent is selected from the group consisting of ethyl acetate, acetone and any combinations thereof.

[0156] In some embodiments, the concentration of the compound of formula I in the solvent is 10 - 30 mg/mL, preferably 20 mg/mL.

[0157] In some embodiments, the concentration of the fumaric acid in ethanol is 15 - 35 mg/mL, preferably 25 mg/mL.

[0158] In some embodiments, after adding the fumaric acid in ethanol, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours.

[0159] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0160] Crystal form M of a benzene sulfonate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of $7.5^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

[0161] The X-ray powder diffraction pattern of the crystal form M of a benzene sulfonate of a compound of formula I shows characteristic peaks at 2θ angles of $7.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $15.2^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $22.4^\circ \pm 0.2^\circ$, $23.0^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

[0162] The X-ray powder diffraction pattern of the crystal form M of a benzene sulfonate of a compound of formula I shows characteristic peaks at 2θ angles of $7.5^\circ \pm 0.2^\circ$, $12.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $15.2^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $22.4^\circ \pm 0.2^\circ$, $23.0^\circ \pm 0.2^\circ$, $24.6^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

[0163] Non-restrictively, the X-ray powder diffraction data of crystal form M of a benzene sulfonate of a compound of formula I of the invention are shown in Table 13.

Table 13

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
7.5	11.72	100.0
12.5	7.08	16.5
14.1	6.28	19.4
15.2	5.82	19.0
18.5	4.80	48.6
22.4	3.96	39.3

EP 4 385 511 A2

(continued)

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
23.0	3.87	22.7
24.6	3.62	17.6
25.2	3.54	45.3
29.8	2.99	51.3

[0164] Non-restrictively, the XRPD pattern of crystal form M of a benzene sulfonate of a compound of formula I of the invention is shown in Figure 32.

[0165] Non-restrictively, the DSC thermogram of crystal form M of a benzene sulfonate of a compound of formula I of the invention is shown in Figure 33. The DSC thermogram shows that the initial melting point of crystal form M of a benzene sulfonate of a compound of formula I of the invention is 198.73 °C.

[0166] Non-restrictively, the DVS isotherm plot of crystal form M of a benzene sulfonate of a compound of formula I of the invention is shown in Figure 34. The DVS isotherm plot shows a 4.6% weight gain of crystal form M of a benzene sulfonate of a compound of formula I of the invention by moisture absorption from 0%RH to 95%RH. At a humidity of 85%RH, crystal form M of a benzene sulfonate of a compound of formula I reaches a weight gain of 0.54%; at 70%RH, crystal form M of a benzene sulfonate of a compound of formula I reaches a weight gain of 0.97%. After moisture absorption, the crystal form of crystal form M of a benzene sulfonate of a compound of formula I doesn't change (see Figure 35 for the XRPD pattern after moisture absorption).

[0167] The present invention provides a preparation method of crystal form M of a benzene sulfonate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding benzenesulfonic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form M of a benzene sulfonate of a compound of formula I.

[0168] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone.

[0169] In some embodiments, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;

[0170] In some embodiments, the concentration of the benzenesulfonic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

[0171] In some embodiments, after adding the benzenesulfonic acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

[0172] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0173] Crystal form N of a citrate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2θ angles of 15.8° ± 0.2°, 17.0° ± 0.2° and 21.1° ± 0.2°.

[0174] Non-restrictively, the X-ray powder diffraction data of crystal form N of a citrate of a compound of formula I of the invention are shown in Table 14.

Table 14

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
4.8	18.35	4.3
15.8	5.62	81.4
17.0	5.21	17.7
21.1	4.21	15.5

[0175] Non-restrictively, the XRPD pattern of crystal form N of a citrate of a compound of formula I of the invention is shown in Figure 36.

[0176] Non-restrictively, the DSC thermogram of crystal form N of a citrate of a compound of formula I of the invention is shown in Figure 37.

[0177] The present invention provides a preparation method of the crystal form N of a citrate of a compound of formula I, specifically, the method is described as follows:

dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding

EP 4 385 511 A2

citric acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form N of a citrate of a compound of formula I.

[0178] In some embodiments, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate;

[0179] In some embodiments, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL;

[0180] In some embodiments, the concentration of the citric acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL;

[0181] In some embodiments, after adding the citric acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

[0182] In some embodiments, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

[0183] Crystal form O of a tartrate of a compound of formula I of the present invention has an X-ray powder diffraction pattern showing characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $26.1^\circ \pm 0.2^\circ$ and $26.9^\circ \pm 0.2^\circ$.

[0184] In some embodiments, the X-ray powder diffraction pattern of the crystal form O of a tartrate of a compound of formula I shows characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $12.5^\circ \pm 0.2^\circ$, $15.1^\circ \pm 0.2^\circ$, $26.1^\circ \pm 0.2^\circ$, $26.9^\circ \pm 0.2^\circ$ and $27.5^\circ \pm 0.2^\circ$.

[0185] In some embodiments, the X-ray powder diffraction pattern of the crystal form O of a tartrate of a compound of formula I shows characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $11.4^\circ \pm 0.2^\circ$, $12.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $14.4^\circ \pm 0.2^\circ$, $15.1^\circ \pm 0.2^\circ$, $26.1^\circ \pm 0.2^\circ$, $26.9^\circ \pm 0.2^\circ$ and $27.5^\circ \pm 0.2^\circ$.

[0186] Non-restrictively, the X-ray powder diffraction data of crystal form O of a tartrate of a compound of formula I of the invention are shown in Table 15:

Table 15

Diffraction angle 2θ (°)	d (Å)	Relative diffraction intensity I% (based on peak height)
6.3	14.00	100.0
11.4	7.73	11.0
12.5	7.05	18.3
14.1	6.26	10.9
14.4	6.14	6.2
15.1	5.86	12.6
26.1	3.41	38.5
26.9	3.32	24.2
27.5	3.24	11.7

[0187] Non-restrictively, the XRPD pattern of crystal form O of a tartrate of a compound of formula I of the invention is shown in Figure 38.

[0188] Non-restrictively, the DSC thermogram of crystal form O of a tartrate of a compound of formula I of the invention is shown in Figure 39. The DSC thermogram shows that the initial melting point of crystal form O of a tartrate of a compound of formula I of the invention is 218.80 °C.

[0189] Non-restrictively, the TGA thermogram of crystal form O of a tartrate of a compound of formula I of the invention is shown in Figure 40. The TGA thermogram shows that there is only a 0.05% weight loss of crystal form O of a tartrate of a compound of formula I of the invention from 26°C to 120°C, indicating crystal form O of a tartrate of a compound of formula I doesn't contain crystal water or solvent.

[0190] Non-restrictively, the DVS isotherm plot of crystal form O of a tartrate of a compound of formula I of the invention is shown in Figure 41. The DVS isotherm plot shows a 6.85% weight gain of crystal form O of a tartrate of a compound of formula I of the invention by moisture absorption from 0%RH to 95%RH. At a humidity of 80%RH, crystal form O of a tartrate of a compound of formula I reaches a 1.80% weight gain. After moisture absorption, the crystal form of crystal form O of a tartrate of a compound of formula I doesn't change (see Figure 42 for the XRPD pattern after moisture absorption).

[0191] The present invention provides a preparation method of crystal form O of a tartrate of a compound of formula I, specifically, the method is described as follows:

mixing a compound of formula I with a first solvent, dissolving the compound until the solution is clear and obtaining a compound of formula I in the first solvent; mixing tartaric acid with a second solvent, dissolving the compound until the solution is clear, and obtaining tartaric acid in the second solvent; adding the tartaric acid in the second solvent to the

EP 4 385 511 A2

compound of formula I in the first solvent under stirring, conducting controlled-rate cooling under stirring, collecting the solid, drying and obtaining crystal form O of a tartrate of a compound of formula I.

[0192] In some embodiments, the first solvent and the second solvent are selected from the group consisting of acetone, ethyl acetate and any combinations thereof.

[0193] In some embodiments, the molar ratio of the compound of formula I to tartaric acid is 1: (0.5 - 1.5), preferably 1: (0.5 - 0.7), more preferably 1: (0.55 - 0.6).

[0194] In some embodiments, in the crystal form O of a tartrate of a compound of formula I, the molar ratio of the compound of formula I to tartaric acid is 2: 1.

[0195] In some embodiments, the concentration of the compound of formula I in acetone is 15 - 70 mg/mL, preferably 40 - 60 mg/mL, more preferably preferably 50 mg/mL.

[0196] In some embodiments, the concentration of the tartaric acid in acetone is 5 - 35 mg/mL, preferably 10 - 25 mg/mL, more preferably preferably 15 mg/mL.

[0197] In some embodiments, mixing the compound of formula I with acetone, raising the temperature to 40 - 60 °C, preferably 50 - 55 °C, to dissolve the compound of formula I until the solution is clear.

[0198] In some embodiments, mixing the tartaric acid with acetone, raising the temperature to 40 - 60 °C, preferably 50 - 55 °C, to dissolve the tartaric acid until the solution is clear.

[0199] In some embodiments, adding the tartaric acid in acetone at 40 - 60 °C, preferably 45 - 55 °C, to the compound of formula I in acetone.

[0200] In some embodiments, the controlled-rate cooling is realized by the following steps:

stirring the system for 0.5 - 3 hours, preferably 1 - 2 hours, at room temperature between 35 - 60 °C, preferably 40 - 60 °C;

keeping cooling the system to 15 - 35 °C, maintaining the temperature, stirring for 0.5 - 3 hours, preferably 1 - 2 hours;

keeping cooling the system to 5 - 15 °C, such as 5 - 10 °C, and maintaining the temperature, stirring for 0.5 - 3 hours, preferably 1 - 2 hours.

[0201] In the present invention, the controlled-rate cooling can gradually cool the system in steps, and maintain a specific temperature range for a certain duration.

[0202] In some embodiments, during the controlled-rate cooling, after stirring the system for 0.5 - 3 hours, preferably 1 - 2 hours, at room temperature between 35 - 60 °C, preferably 40 - 60 °C, concentrating the system to one third to two thirds of the original volume, preferably half of the original volume.

[0203] In some embodiments, during the controlled-rate cooling, after keeping cooling the system to 15 - 35 °C, maintaining the temperature and stirring for 0.5 - 3 hours, preferably 1 - 2 hours, concentrating the system to one third to two thirds of the original volume, preferably half of the original volume.

[0204] In some embodiments, the purity of the compound of formula I is more than 90%, preferably more than 95%, more preferably more than 99%.

[0205] In some embodiments, drying the collected solid between 40 - 60 °C under reduced pressure or with an air blower for 5 - 48 hours, preferably 16 - 28 hours.

[0206] The present invention also provides a pharmaceutical composition comprising crystal form 1 of a compound of formula I, crystal form A of a hydrochloride of a compound of formula I, crystal form B of a hydrochloride of a compound of formula I, crystal form C of a hydrochloride of a compound of formula I, crystal form D of a sulfate of a compound of formula I, crystal form E of a phosphate of a compound of formula I, crystal form F of a phosphate of a compound of formula I, crystal form G of a mesylate of a compound of formula I, crystal form H of a hydrobromide of a compound of formula I, crystal form J of a hydrobromide of a compound of formula I, crystal form K of a hydrobromide of a compound of formula I, crystal form L of a fumarate of a compound of formula I, crystal form M of a benzene sulfonate of a compound of formula I, crystal form N of a citrate crystal of a compound of formula I and/or crystal form O of a tartrate of a compound of formula I.

[0207] The present invention also provides a pharmaceutical formulation comprising crystal form 1 of a compound of formula I, crystal form A of a hydrochloride of a compound of formula I, crystal form B of a hydrochloride of a compound of formula I, crystal form C of a hydrochloride of a compound of formula I, crystal form D of a sulfate of a compound of formula I, crystal form E of a phosphate of a compound of formula I, crystal form F of a phosphate of a compound of formula I, crystal form G of a mesylate of a compound of formula I, crystal form H of a hydrobromide of a compound of formula I, crystal form J of a hydrobromide of a compound of formula I, crystal form K of a hydrobromide of a compound of formula I, crystal form L of a fumarate of a compound of formula I, crystal form M of a benzene sulfonate of a compound of formula I, crystal form N of a citrate of a compound of formula I and/or crystal form O of a tartrate of a compound of formula I.

5 [0208] Use of crystal form 1 of a compound of formula I, crystal form A of a hydrochloride of a compound of formula I, crystal form B of a hydrochloride of a compound of formula I, crystal form C of a hydrochloride of a compound of formula I, crystal form D of a sulfate of a compound of formula I, crystal form E of a phosphate of a compound of formula I, crystal form F of a phosphate of a compound of formula I, crystal form G of a mesylate of a compound of formula I, crystal form H of a hydrobromide of a compound of formula I, crystal form J of a hydrobromide of a compound of formula I, crystal form K of a hydrobromide of a compound of formula I, crystal form L of a fumarate of a compound of formula I, crystal form M of a benzene sulfonate of a compound of formula I, crystal form N of a citrate of a compound of formula I and/or crystal form O of a tartrate of a compound of formula I in preparing medicaments for treating JAK1/TYK2-related diseases or conditions, wherein the diseases or conditions can be autoimmune diseases or disorders, such as rheumatoid arthritis or inflammatory diseases or disorders, and cancers or tumor proliferative diseases or disorders.

10 [0209] In the present invention, unless otherwise specified, the involved temperatures refer to the internal temperatures of the reaction system.

15 [0210] In terms of the melting point, a person skilled in the art can understand that in a DSC test, there are a certain range of changes in the actually measured initial melting point due to the influences of measuring instrument, heating rate and crystalline shape, etc.; generally, these changes are within ± 5 °C.

BRIEF DESCRIPTION OF THE DRAWING

20 [0211]

Figure 1 is the XRPD pattern of crystal form 1 of a compound of formula I of the invention.

Figure 2 is the DSC thermogram of crystal form 1 of a compound of formula I of the invention.

25 Figure 3 is the TGA thermogram of crystal form 1 of a compound of formula I of the invention.

Figure 4 is the DVS isotherm plot of crystal form 1 of a compound of formula I of the invention.

30 Figure 5 is the XRPD overlay pattern of crystal form 1 of a compound of formula I of the invention before and after the DVS test.

Figure 6 is the XRPD pattern of crystal form A of a hydrochloride of a compound of formula I of the invention.

35 Figure 7 is the DSC thermogram of crystal form A of a hydrochloride of a compound of formula I of the invention.

Figure 8 is the XRPD pattern of crystal form B of a hydrochloride of a compound of formula I of the invention.

Figure 9 is DSC thermogram of crystal form B of a hydrochloride of a compound of formula I of the invention.

40 Figure 10 is the XRPD pattern of crystal form C of a hydrochloride of a compound of formula I of the invention.

Figure 11 is the DSC thermogram of crystal form C of a hydrochloride of a compound of formula I of the invention.

45 Figure 12 is the XRPD pattern of crystal form D of a sulfate of a compound of formula I of the invention.

Figure 13 is the DSC thermogram of crystal form D of a sulfate of a compound of formula I of the invention.

Figure 14 is the XRPD pattern of crystal form E of a phosphate of a compound of formula I of the invention.

50 Figure 15 is the DSC thermogram of crystal form E of a phosphate of a compound of formula I of the invention.

Figure 16 is the XRPD pattern of crystal form F of a phosphate of a compound of formula I of the invention.

55 Figure 17 is the DSC thermogram of crystal form F of a phosphate of a compound of formula I of the invention.

Figure 18 is the DVS isotherm plot of crystal form F of a phosphate of a compound of formula I of the invention.

Figure 19 is the XRPD overlay pattern of crystal form F of a phosphate of a compound of formula I of the invention

before and after the DVS test.

Figure 20 is the XRPD pattern of crystal form G of a mesylate of a compound of formula I of the invention.

5 Figure 21 is the DSC thermogram of crystal form G of a mesylate of a compound of formula I of the invention.

Figure 22 is the DVS isotherm plot of crystal form G of a mesylate of a compound of formula I of the invention.

10 Figure 23 is the XRPD pattern of crystal form H of a hydrobromide of a compound of formula I of the invention.

Figure 24 is the DSC thermogram of crystal form H of a hydrobromide of a compound of formula I of the invention.

Figure 25 is the XRPD pattern of crystal form J of a hydrobromide of a compound of formula I of the invention.

15 Figure 26 is the XRPD pattern of crystal form K of a hydrobromide of a compound of formula I of the invention.

Figure 27 is the DSC thermogram of crystal form K of a hydrobromide of a compound of formula I of the invention.

20 Figure 28 is the DVS isotherm plot of crystal form K of a hydrobromide of a compound of formula I of the invention.

Figure 29 is the XRPD overlay pattern of crystal form K of a hydrobromide of a compound of formula I of the invention before and after the DVS test.

25 Figure 30 is the XRPD pattern of crystal form L of a fumarate of a compound of formula I of the invention.

Figure 31 is the DSC thermogram of crystal form L of a fumarate of a compound of formula I of the invention.

Figure 32 is the XRPD pattern of crystal form M of a benzene sulfonate of a compound of formula I of the invention.

30 Figure 33 is the DSC thermogram of crystal form M of a benzene sulfonate of a compound of formula I of the invention.

Figure 34 is the DVS isotherm plot of crystal form M of a benzene sulfonate of a compound of formula I of the invention.

35 Figure 35 is the XRPD overlay pattern of crystal form M of a benzene sulfonate of a compound of formula I of the invention before and after the DVS test.

Figure 36 is the XRPD pattern of crystal form N of a citrate of a compound of formula I of the invention.

40 Figure 37 is the DSC thermogram of crystal form N of a citrate of a compound of formula I of the invention.

Figure 38 is the XRPD pattern of crystal form O of a tartrate of a compound of formula I of the invention.

Figure 39 is the DSC thermogram of crystal form O of a tartrate of a compound of formula I of the invention.

45 Figure 40 is the TGA thermogram of crystal form O of a tartrate of a compound of formula I of the invention.

Figure 41 is the DVS isotherm plot of crystal form O of a tartrate of a compound of formula I of the invention.

50 Figure 42 is the XRPD overlay pattern of crystal form O of a tartrate of a compound of formula I of the invention before and after the DVS test.

Figure 43 is the ¹H NMR spectrum of crystal form 1 of a compound of formula I of the invention.

55 Figure 44 is the ¹H NMR spectrum of crystal form O of a tartrate of a compound of formula I of the invention.

Figure 45 is the XRPD overlay pattern of crystal form F of a phosphate of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

EP 4 385 511 A2

Figure 46 is the XRPD overlay pattern of crystal form O of a tartrate of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

Figure 47 is the DSC overlay thermogram of crystal form F of a phosphate of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

Figure 48 is the DSC overlay thermogram of crystal form O of a tartrate of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

Figure 49 is the XRPD overlay pattern of crystal form 1 of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

Figure 50 is the DSC overlay thermogram of crystal form 1 of a compound of formula I of the invention after being placed at a high temperature and under accelerated conditions for 2 weeks.

EMBODIMENTS

[0212] The following embodiments further explain the invention, but don't constitute a restriction or limitation to the scope of the invention.

No.	Instrument	Model	Manufacturer	Test methods
1	High performance liquid chromatograph (UPLC)	Agilent 1200, DAD	Agilent	Instrument: Agilent 1200 DAD HPLC System or Similar configuration Chromatographic column: Waters XBridge Shield RP18 4.6×150mm, 3.5 μm Mobile phase: A: 0.05% phosphoric acid aqueous solution; B: Acetonitrile
2	Liquid chromatography-mass spectrometry (LC-MS)	Agilent 1200 HPLC/6100 SQ System	Agilent	Instrument: Agilent 1200 HPLC/6100 SQ System Chromatographic column: Agilent XDB-C18, 4.6mmx50mm, 1.8μm Mobile phase: A: 0.05% TFA in water; B: 0.05% TFA in acetonitrile
3	Nuclear magnetic resonance spectroscopy ¹ HNMR	AVANCE III 400MHz	BRUKER	Ultrashield-Plus Digital NMR Spectroscopy Experiment: N PROTON 1H experiment (default parameters)
4	X-ray powder diffractometer (XRPD)	D8 Advance	BRUKER	Light source is CuK. X-ray intensity is 40 KV/40 mA. Scanning mode is Theta-theta. Scanning angle range is 4 - 40°. Step length is 0.05°. Scanning speed is 0.5 seconds/step.
5	Differential scanning calorimeter (DSC)	Q1000	TA	Weighing 2 - 4 mg of sample and placing the sample into a unsealed aluminum pan, allowing the sample to reach the equilibrium in nitrogen flow (50 mL/min) at 25 °C, and raising the temperature from 25 °C to 300 °C at 10 °C/min.
6	Thermogravimetric analyzer (TGA)	Q500	TA	Weighing 10 - 20 mg of sample and placing the sample into a platinum sample pan, allowing the sample in nitrogen flow (60 mL/min) and the balance in nitrogen flow (40 mL/min) to reach equilibrium at 25 °C, and raising the temperature from 25 °C to 300 °C at 10 °C/min.

EP 4 385 511 A2

(continued)

No.	Instrument	Model	Manufacturer	Test methods
7	Dynamic vapor sorption analyzer (DVS)	Advantage 1	Surface Measurement System	Weighing about 10 mg of the sample, setting the temperature as 25 °C, drying for 60 minutes at a humidity of 0%RH, and determining the moisture absorption characteristics of the samples when the humidity changes from 0%RH to 95%RH, and the dehumidification characteristics of the samples when the humidity changes from 95%RH to 0%RH. The humidity change step is 5%RH. When the mass change rate dm/dt is less than 0.002%, it is considered as the scale balance. The mass change rate less than 0.01%/min within 5 minutes is the balance criterion in the test and the maximum equilibration time is 2 hours.
8	Ion chromatography	ICS-2000 + AS40 Automated sampler	DIONEX	Chromatographic column: IonPac® AS11-HC 4*250 mm; Column temperature: 30 °C; Eluent: 10 mM KOH aqueous solution Flow rate: 1.00 mL/min Suppressor: DionexAERS 500 4 mm Electric current of suppressor: 25 mA Run time: 15 min

[0213] Information on raw materials and reagents used in the invention is as follows:

Material	Purity/grade	Lot No.	Manufacturer
Compound of formula V	≥98.0%	KM 1009-1804001	ZINNOVA
Compound of formula IV	≥98.0%	KM 1008-1804001	ZINNOVA
N,N-Diisopropylethylamine (DIPEA)	AR	KM261A-1801002	Shanghai Qiao Chemical Science Co., Ltd.
Ethanol	/	160321047B	Nanjing Chemical Reagent Co., Ltd.
Methanol	AR	P1176856	GENERAL-REAGENT
Tetrahydrofuran (THF)	AR	P1167158	GENERAL-REAGENT
Dichloromethane (DCM)	AR	P1216848	GENERAL-REAGENT
Ethyl acetate (EA)	AR	P1080359	GENERAL-REAGENT
Acetone	AR	P1160778	GENERAL-REAGENT
Acetonitrile	HPLC	6308IX20	Anhui Fulltime Specialized Solvent & Reagent Co., Ltd.
N-hexane	AR	P1196621	GENERAL-REAGENT
Purified water	Milli-Q	Prepared on the same day	Milli-Q
Methyl tert-butyl ether (MTBE)	AR	P1135054	GENERAL-REAGENT
Isopropanol	HPLC	6553IU13	Anhui Fulltime Specialized Solvent & Reagent Co., Ltd.
Palladium on carbon (Pd/C)	AR	KM416A-1603001	Shaanxi Rock New Materials Co., Ltd.
Trimethyl orthoacetate	98%+	KM1013-1805001	Shanghai Titan Scientific Co., Ltd.


(continued)

	Material	Purity/grade	Lot No.	Manufacturer
5	Pyridine hydrochloride	98%+	KM616-1703001	Jiangsu Heng An Chemical Industry Co., Ltd.
	Hydrobromic acid	45%	P1337848	Adamas-beta
	Tartaric acid (L)	99%+	P1311486	Adamas-beta
10	Benzenesulfonic acid monohydrate	98%+	P1257168	Adamas-beta
	Hydrochloric acid	36-38%	P1246465	GENERAL-REAGENT
	Sulfuric acid	95-98%	20140301	Wuxi Jiani Chemistry Co., Ltd.
15	Methanesulfonic acid	>98.0%	P1133997	GENERAL-REAGENT
	Fumaric acid	99%	LU80M51	J&KCHEMICAL
	Citric acid	Pharmaceutical-grade	160105001C	Nanjing Chemical Reagent Co., Ltd.
20	P-toluenesulfonic acid	>99.5%	20101208	Shanghai Lingfeng Chemical Reagent Co., Ltd.
	Phosphoric acid	85%	H2160	Honeywell
25	Silicon based metal eliminator (Thiol silica gel)	Sulphur content \geq 3.2%	-	Shanghai Chiral Chemistry Co., Ltd.
	Methanolic potassium hydroxide VS	AR	0.1 mol/L	-
30	Potassium hydrogen phthalate	AR	\geq 99.5%	-

Examples

Preparation of a compound of formula III

[0214]



Example 1 Preparation of a compound of formula III

[0215] Ethanol (4 mL), a compound of formula IV (0.20 g, 1.0 eq), a compound of formula V (0.18 g, 1.0 eq), and DIPEA (0.39 g, 3.0 eq) were added to a 25 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to reflux (70 - 80 °C), stirred overnight at the reflux temperature; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (4 mL) was added to the system dropwise, the system was stirred for 2 hours at room temperature (15 - 20 °C); filtered, the filter cake was washed with ethanol aqueous solution (2 mL, V/V, 1:1), the filter cake was dried under vacuum at 45 - 50 °C for 16 hours; about 0.21 g of yellow solids were obtained, with a LC-MS purity of 96.4% (214 nm) and a yield of 69%.

EP 4 385 511 A2

[0216] MS-ESI: [M +1]⁺: 303.1

[0217] ¹H NMR(400MHz, CDCl₃): 9.238 (s, 1H), 8.400 (d, 1H), 7.968 (d, 1H), 6.987 (d, 1H), 4.537 - 4.613 (m, 1H), 4.305 - 4.350 (m, 1H), 3.661 - 3.722 (m, 1H), 3.313 - 3.366 (m, 1H), 2.590 - 2.699 (m, 2H), 2.407 - 2.454 (m, 1H), 1.815 - 2.035 (m, 1H), 1.688 - 1.806 (m, 2H).

5

Example 2 Preparation of a compound of formula III

[0218] Ethanol (120 mL, 20V), a compound of formula IV (6.0 g, 1.0 eq), a compound of formula V (5.4 g, 1.01 eq), and DIPEA (11.7 g, 3.0 eq) were added to a 250 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C (internal temperature) and stirred with the temperature maintained for 8 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (120 mL, 20V) was added to the system dropwise, the system was stirred for 2 hours at room temperature (10 - 15 °C); filtered, the filter cake was washed with ethanol aqueous solution (30 mL, 1:1), the filter cake was dried under vacuum at 50 °C for 16 hours; about 7.7 g of yellow solids were obtained in total, with an HPLC purity of 95.5% and a yield of 84.3%.

10

15

[0219] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 3 Preparation of a compound of formula III

[0220] Ethanol (5 mL, 10V), a compound of formula IV (0.50 g, 1.0 eq), a compound of formula V (0.45 g, 1.01 eq), and DIPEA (0.98 g, 3.0 eq) were added to a 25 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C, and was allowed to reflux and was stirred for 5 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (5 mL, 10V) was added to the system dropwise, the system was stirred for 2 hours at room temperature (10 - 15 °C); filtered, the filter cake was washed with ethanol aqueous solution (1:1) (1.5 mL, 3V), the filter cake was dried under vacuum at 50 °C for 16 hours; about 0.54 g of brown solids were obtained in total, with an HPLC purity of 95.4% and a yield of 71%.

20

25

[0221] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 4 Preparation of a compound of formula III

[0222] Ethanol (5 mL, 10V), a compound of formula IV (0.50 g, 1.0 eq), a compound of formula V (0.45 g, 1.01 eq), and DIPEA (0.72 g, 2.2 eq) were added to a 25 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C, and was allowed to reflux and was stirred for 5 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (7.5 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1) (1.5 mL, 3V), the filter cake was dried under vacuum at 50 °C for 16 hours; about 0.57 g of brown solids were obtained in total, with an HPLC purity of 91.4% and a yield of 75%.

30

35

[0223] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 5 Preparation of a compound of formula III

[0224] Ethanol (50 mL, 10V), a compound of formula IV (5.0 g, 1.0 eq), a compound of formula V (4.5 g, 1.01 eq), and DIPEA (7.2 g, 2.2 eq) were added to a 250 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C, and was allowed to reflux and was stirred for 5 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (75 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1, 15 mL), the filter cake was dried under vacuum at 50 °C for 16 hours; about 6.6 g of yellow solids were obtained in total, with an HPLC purity of 94.2% and a yield of 86.7%.

40

45

[0225] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 6 Preparation of a compound of formula III

[0226] Ethanol (180 mL, 10V), a compound of formula IV (17.8 g, 1.0 eq), a compound of formula V (16.0 g, 1.01 eq), and DIPEA (25.7 g, 2.2 eq) were added to a 500 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C, and was allowed to reflux and was stirred for 5 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (270 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred

55

EP 4 385 511 A2

for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (ethanol:water=1:1.5, v/v, 40 mL), the filter cake was dried under vacuum at 50 °C for 16 hours; about 23.0 g of brown solids were obtained in total, with an HPLC purity of 95.3% and a yield of 85.2%.

[0227] The MS-ESI and ¹H NMR data are consistent with example 1.

5

Example 7 Preparation of a compound of formula III

[0228] Ethanol (1000 mL, 10V), a compound of formula IV (100 g, 1.0 eq), a compound of formula V (89.9 g, 1.01 eq), and DIPEA (143.2 g, 2.2 eq) were added to a 3000 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 85 - 90 °C (internal temperature: about 75°C), and was allowed to reflux and was stirred for 10 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (1500 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1.5, v/v, 200 mL), the filter cake was dried under vacuum at 50 °C for 16 hours; about 130 g of reddish brown solids were obtained in total, with an HPLC purity of 94.2% and a yield of 85.5%.

10

15

[0229] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 8 Preparation of a compound of formula III

[0230] Ethanol (2000 mL, 10V), a compound of formula IV (200 g, 1.0 eq), a compound of formula V (179.7 g, 1.01 eq), and DIPEA (286.4 g, 2.2 eq) were added to a 5000 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 70 - 80 °C (internal temperature: about 65 - 70°C), and was allowed to reflux and was stirred for 16 hours; the system was cooled to room temperature (15 - 20 °C), solids were precipitated during cooling; water (3000 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1.5, v/v, 400 mL), the filter cake was dried with an air blower at 50 °C for 16 hours; about 251 g of reddish brown solids were obtained in total, with an HPLC purity of 93.4%, a content of 94.7% and a content yield of 78.1%.

20

25

[0231] The MS-ESI and ¹H NMR data are consistent with example 1.

30

Example 9 Preparation of a compound of formula III

[0232] Ethanol (5000 mL, 10V), a compound of formula IV (500 g, 1.0 eq), a compound of formula V (450 g, 1.01 eq), and DIPEA (723 g, 2.2 eq) were added to a 20000 mL three-necked flask, and were stirred; under nitrogen protection, the system was heated to 80 - 90 °C (internal temperature: about 70 - 80°C), and was allowed to reflux and was stirred for 16 hours; the system was cooled to room temperature (25 - 30 °C), solids were precipitated during cooling; water (7500 mL, 15V) was added to the system dropwise, the system was stirred for 1 hour at room temperature (25 - 30 °C); the system was cooled to 10 - 15 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1.5, v/v, 1000 mL), the filter cake was dried in an oven under vacuum at 50 - 55 °C for 24 hours; about 623 g of products were obtained in total, with an HPLC purity of 93.7%, ethanol residue of 0.5%, a content of 93.1%, and a content yield of 76.2%.

35

40

[0233] The MS-ESI and ¹H NMR data are consistent with example 1.

Example 10 Preparation of a compound of formula III

[0234] Ethanol (100 mL, 10V), a compound of formula IV (10.0 g, 1.0 eq), a compound of formula V (9.0 g, 1.01 eq), and DIPEA (14.3 g, 2.2 eq) were added to a 500 mL three-necked flask, and were stirred; the system was heated to 70 - 80 °C, and was allowed to reflux and was stirred for 16 hours; the system was cooled to room temperature (20 - 30 °C), solids were precipitated during cooling; water (150 mL, 15V) was added to the system, the system was stirred for 2 hours at room temperature (20 - 30 °C); the system was cooled to 5 - 10 °C and was stirred for 2 hours; filtered, the filter cake was washed with ethanol aqueous solution (1:1.5, v/v, 25 mL), the filter cake was dried in an oven under vacuum at 50 - 55 °C for 16 hours; about 13.7 g of products were obtained in total, with an HPLC purity of 93.7%, and a yield of 90%.

45

50

[0235] The MS-ESI and ¹H NMR data are consistent with example 1.

55

Example 11 Preparation of a compound of formula III

[0236] Ethanol (17 Kg, 10V), a compound of formula IV (2.2 Kg, 1.0 eq), a compound of formula V (1.98 Kg, 1.01 eq), and DIPEA (3.19 Kg, 2.2 eq) were added to a R0462 reactor, and were stirred; under nitrogen protection, the system


EP 4 385 511 A2

was heated to 75 - 80 °C (internal temperature, about 70 - 80 °C), and was stirred for 16 hours; the system was cooled to room temperature (15 - 25°C), solids were precipitated during cooling; water (33 Kg, 15V) was added to the system dropwise, the system was stirred for 2 hours at room temperature (10 - 15 °C); the system was cooled to 5 - 10 °C and was stirred for 4 hours; filtered, the filter cake was washed with ethanol aqueous solution (ethanol:water=1:2, v/v, 6.2 Kg), the filter cake was dried at a jacket temperature between 45 - 55 °C and under vacuum ≤ -0.08 MPa for 16 hours; about 2.64 g of brown solids were obtained in total, with an HPLC purity of 94.0%, a content of 93.4% and a content yield of 79.04%.

[0237] The MS-ESI and ^1H NMR data are consistent with example 1.

Preparation of a compound of formula II

[0238]



Example 12 Preparation of a compound of formula II

[0239] A compound of formula III (5.0 g), THF (50 mL, 10 V) and palladium on carbon (0.75 g, 10% Pd/C, 50% wet) were added to a 100 mL stainless steel autoclave sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the pressure of the system was increased to 0.50 MPa with hydrogen, then the system was heated to 25 - 35 °C and stirred 24 hours with the temperature maintained; the reaction liquid was filtered with diatomite, the filter cake was washed with THF (20 mL), and the filtrate was concentrated to dryness to obtain 4.2 g of brown solids, with an HPLC purity of 94.9% and a yield of 93.3%.

[0240] MS-ESI: $[\text{M} + 1]^+$: 273.1

[0241] ^1H NMR (400MHz, CDCl_3): 7.988 (s, 1H), 7.688 (d, 1H), 6.805 (d, 1H), 4.190 - 4.338 (m, 3H), 3.584 - 3.648 (m, 1H), 3.147 - 3.206 (t, 1H), 2.594 - 2.651 (d, 2H), 2.318 - 2.364 (m, 1H), 1.917 - 1.974 (m, 1H), 1.633 - 1.738 (m, 1H), 1.456 - 1.525 (m, 1H).

Example 13 Preparation of a compound of formula II

[0242] A compound of formula III (120.0 g), THF (2400 mL, 20 V) and palladium on carbon (18 g, 10% Pd/C, 50% wet) were added to a 5000 mL stainless steel autoclave sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the pressure of the system was increased to 0.50 MPa with hydrogen, then the system was heated to 25 - 35 °C and stirred 24 hours with the temperature maintained; the reaction liquid was filtered with diatomite, the filter cake was washed with THF (600 mL) (until TLC almost did not show fluorescence), and the filtrate was concentrated to obtain 130 g of black semi-oily solids, with an HPLC purity of 91.7% and a yield of 120.26%.

[0243] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 14 Preparation of a compound of formula II

[0244] A compound of formula III (100.0 g), THF (2000 mL, 20 V) and palladium on carbon (15.0 g, 10% Pd/C, 50% wet) were added to a 5 L stainless steel autoclave sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the pressure of the system was increased to 0.5 - 1.0 MPa with hydrogen, the temperature of the jacket was set to 30 °C, the system was stirred for 16 hours with the temperature maintained, the reaction liquid was filtered with diatomite, the filter cake was washed with THF (1000 mL), 3877 g of a compound of formula II in THF was obtained in total.

[0245] Post-treatment 1: the above filtrate (1820 g, about 40 g of a compound of formula II calculated according to a 100% yield) was concentrated to (2 - 3 V, 80 - 120 mL) with a rotary evaporator, the system was exchanged with ethanol (150 mL \times 2) to (2 - 3 V, 80 - 120 mL); 78 g of a compound of formula II in ethanol was obtained, with a content of

EP 4 385 511 A2

47.25%, and a content yield of 92.14%.

[0246] Post-treatment 2: the above filtrate (450 g, about 10 g of a compound of formula II calculated according to a 100% yield) was concentrated to dryness with a rotary evaporator; 10.5 g of brownish red solids were obtained.

[0247] Post-treatment 3: the above filtrate (450 g, about 10 g of a compound of formula II after calculation) was added to a flask, concentrated to about 30 - 40 mL (3 - 4 V) with a rotary evaporator; the concentrated residue was exchanged with ethanol (50 mL \times 2) to about 30 - 40 mL (3 - 4 V); black oily concentrated residues were obtained, the concentrated residues were directly fed to the next step of reaction.

[0248] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 15 Preparation of a compound of formula II

[0249] THF (240 mL, 20 V), a compound of formula III (12.0 g), and palladium on carbon (1.8 g, 5% Pd/C, 50% wet) were added to a 5000 mL three-necked flask sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the system was stirred for 48 hours with the temperature maintained at room temperature (25 - 30 $^{\circ}\text{C}$) and under hydrogen pressure (about 0.1 MPa); the filter liquid was filtered, the filter cake was washed with THF (60 mL); the combined filtrate was concentrated with a rotary evaporator to 20 - 30 mL, the system was exchanged with ethanol (60 mL \times 2) to 20 - 30 mL; 24g of a compound of formula II in ethanol was obtained, which is directly used for the next step of reaction.

[0250] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 16 Preparation of a compound of formula II

[0251] THF (1500 mL, 15 V), a compound of formula III (100 g), and palladium on carbon (15 g, 5% Pd/C, 50% wet) were added to a 5000 mL three-necked flask sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the system was stirred for 48 hours with the temperature maintained at room temperature (20 - 25 $^{\circ}\text{C}$) and under hydrogen pressure (about 0.1 MPa); the filter liquid was filtered, the filter cake was washed with THF (200 mL); the combined filtrate was concentrated with a rotary evaporator to 200 - 300 mL, 185.6g of a compound of formula II in THF was obtained, with an HPLC purity of 94.2%, a content of 43.2% and a content yield of 94.0%.

[0252] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 17 Preparation of a compound of formula II

[0253] THF (12400 mL, 20 V), a compound of formula III (620 g), and palladium on carbon (93 g, 5% Pd/C, 50% wet) were added to a 20000 mL three-necked flask sequentially; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the system was stirred for 48 hours with the temperature maintained at room temperature (30 - 35 $^{\circ}\text{C}$) and under hydrogen pressure (about 0.1 MPa); the filter liquid was filtered with diatomite (200g), the filter cake was washed with THF (1200 mL); the combined filtrate was concentrated with a rotary evaporator to 1200 - 1800 mL, 1664 g of a compound of formula II in THF was obtained, with an HPLC purity of 93.8%, a content of 34.57% and a content yield of 110.6%.

[0254] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 18 Preparation of a compound of formula II

[0255] THF (140 mL, 70 V), a compound of formula III (2.0 g), and palladium on carbon (0.3 g, 5% Pd/C, 50% wet) were added to a 250 mL autoclave; the autoclave was covered with a cap and the nut was tightened; the system was purged with nitrogen 3 times, then with hydrogen 3 times; the autoclave was charged with hydrogen to about a pressure of 0.50 ± 0.05 MPa, the inlet valve was closed; the stirring apparatus was started at a rotating speed of 500 r/min; the hydrogen pressure of the autoclave was maintained at 0.5 ± 0.05 MPa between 25 - 35 $^{\circ}\text{C}$, the system was stirred for reacting for 96 hours, the reaction liquid was filtered with diatomite (10 g), the filter cake was washed with THF (60 mL); the combined filtrate was concentrated with a rotary evaporator to dryness to obtain 1.8 g of semi-oily solids with an HPLC purity of 91.2%, and a yield of 99.9%.

[0256] The MS-ESI and ^1H NMR data are consistent with example 12.

Example 19 Preparation of a compound of formula II

[0257] THF (167 Kg, 70 V), a compound of formula III (2.64 g), and palladium on carbon (0.4 Kg, 5% Pd/C, 50% wet) were added to a 500 L autoclave; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the autoclave was charged with hydrogen to about a pressure of 0.50 ± 0.05 MPa, the inlet valve was closed; the stirring apparatus

was started; the hydrogen pressure of the autoclave was maintained at 0.5 ± 0.05 MPa between 25 - 35 °C, the system was stirred for reacting for 120 hours, the system was filtered under a pressure, the filter cake was washed with THF (13 Kg); the combined filtrate was distilled under reduced pressure (to 2 V - 3 V) to obtain 11Kg of a compound of formula II in THF, with an HPLC purity of 90.7%, a content of 18.5% and a content yield of 91.9%.

[0258] The MS-ESI and ^1H NMR data are consistent with example 12.


Example 20 Preparation of a compound of formula II

[0259] THF (60 mL, 12 V), a compound of formula III (5.0 g), and palladium on carbon (0.75 g, 5% Pd/C, 50% wet) were added to a 100 mL stainless steel autoclave; the system was purged with nitrogen 5 times, then with hydrogen 5 times; the pressure of the system was increased to 0.5 - 1.0 MPa with hydrogen, the temperature of the jacket was set to 30 °C, the system was stirred for 42 hours with the temperature maintained; after the reaction was completed, the reaction liquid was filtered with diatomite, the filter cake was washed with THF (100 mL); 197.8 g of a compound of formula II in THF was obtained in total; the solution was concentrated with a rotary evaporator to (2 - 3 V, 10 - 15 mL); the system was exchanged with ethanol (25 mL \times 2) to (2 - 3 V, 10 - 15mL); the obtained compound of formula II in ethanol was directly used for the next step of reaction.

[0260] The MS-ESI and ^1H NMR data are consistent with example 12.

Preparation of a compound of formula I

[0261]



Preparation of a compound of formula I **Example 21** Preparation of a compound of formula I A compound of formula II (5 g, 1.0 eq), trimethyl orthoacetate (6.6 g, 3.0 eq) and THF (125 mL) were added to a 250 mL three-necked flask; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (210 mg, 0.1 eq) was added to the three-necked flask; under nitrogen protection, the system was heated to 75 ± 5 °C (internal temperature 60 - 63 °C) and reacted for 8 hours. HPLC monitoring showed the compound of formula II was completely transformed. The purity of the compound of formula I in the residual reaction liquid is 93.1%.

[0262] The system was cooled to room temperature, the reaction liquid in the system was concentrated with a rotary evaporator until there was basically no fraction flowing out; water (50 mL) was added to the system, the pH value of the system was adjusted to 9 - 10 with a 4M sodium hydroxide solution; the system was extracted with ethyl acetate (50 mL), filtered, the filter cake was washed with ethyl acetate (10 mL) to obtain 3.5 g of a compound of formula I filter cake wet product. LC-MS test showed the purity of the compound of formula I was 99.22%; the filtrate obtained by filtration and washing was separated, the aqueous phase was extracted with ethyl acetate (50 mL), the organic phases were combined, and concentrated to dryness to obtain 2.8 g of a compound of formula I crude product. LC-MS test showed the purity of the compound of formula I was 95.08%.

[0263] MS-ESI: $[\text{M}+1]^+$: 297.0 ^1H NMR(400MHz, DMSO): 8.78 (s, 1H), 8.32 (d, 1H), 7.25 (d, 1H), 4.60 (m, 1H), 4.10-4.13 (t, 2H), 3.91 (m, 1H), 2.93-2.98 (m, 1H), 2.80-2.86 (m, 1H), 2.84 (s, 3H), 2.50 (m, 1H), 2.16-2.19 (m, 1H), 1.99-2.02 (m, 1H), 1.69-1.77 (m, 1H).

Example 22 Preparation of a compound of formula I

[0264] A compound of formula II (54 g, 1.0 eq), trimethyl orthoacetate (71.5 g, 3.0 eq) and THF (1.35 L) were added to a 3 L three-necked flask; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (2.27 g, 0.1 eq) was added to the three-necked flask; under nitrogen protection, the system was heated to 75 ± 5 °C (internal temperature 60 - 63 °C) and reacted for 8 hours. HPLC monitoring showed 1.5% of the compound of formula II was left in the system; the system was cooled to room temperature, and trimethyl orthoacetate (11.9 g, 0.5 eq) and pyridine hydrochloride (1.14 g, 0.05 eq) was supplemented to the system; under nitrogen protection, the system was heated to

EP 4 385 511 A2

75 ± 5 °C (internal temperature 60 - 63 °C) and reacted for 4 hours. HPLC monitoring showed 0.4% of the compound of formula II was left in the system, the purity of the compound of formula II in the reaction liquid of the system was 91.6%.

[0265] The system was cooled to room temperature, the reaction liquid in the system was concentrated with a rotary evaporator until there was basically no fraction flowing out; water (540 mL, 10V) was added to the system, the pH value of the system was adjusted to 9 - 10 with a 4M sodium hydroxide solution; the system was filtered, the filter cake was washed with water (270 mL) and then MTBE (270 mL), the obtained filter cake was dried under vacuum at 50 °C for 16 hours to obtain 56 g of a compound of formula II crude product, with an HPLC purity of 96.32%; the obtained crude product was dissolved in 600 mL of methanol until the solution was clear, a silicon based metal eliminator (43 g) and an activate carbon (5.4 g) were added to the system, the mixture was heated to reflux and the temperature was maintained for 1 hour (internal temperature 50 °C); the system was cooled to room temperature, filtered with diatomite, washed with methanol (15 mL) until the filtrate does not show fluorescence; the methanol solution was concentrated to dryness and the dropping speed of the distillate became slower, MTBE (540 mL) was added to the obtained solids, the system was heated to 50 °C and allowed for reflux and was then stirred for 1 hour until the solids were completely dissolved; the system was cooled to 10 - 15 °C and was stirred for 1 hour, filtered, the filter cake was washed with cold MTBE (100 mL); the obtained filter cake was dried under vacuum at 50 °C for 16 hour to obtain 28.0 g of a compound of formula I with an HPLC purity of 98.8%.

[0266] The compound of formula I was recovered from the mother liquid;

[0267] The filtrate obtained in the previous step was concentrated to dryness to obtain about 23 g of light yellow solids; MTBE (230 mL) was added thereto, the temperature was raised to 50 °C, and the system was allowed to reflux for 10 minutes; methanol was added to the system in divided doses, until the total amount of the methanol added was about 30 mL and the material was basically dissolved and the solution was clear; the system was cooled to 10 - 15 °C and was stirred for 1 hour; filtered, the filter cake was washed with cold MTBE (50 mL); the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 8.7 g of off-white solids of a compound of formula I, with an HPLC purity of 97.8%.

Further purification:

[0268] The compound of formula I (11.7 g) with a purity of 97.8% obtained in the previous step and MTBE (60 mL) were added to a reaction flask, the system was triturated at room temperature for 4 hours, filtered, the filter cake was washed with MTBE (20 mL), and the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 10.8 g of off-white compound of formula I, with an HPLC purity of 98.1%.

[0269] The MS-ESI and ¹H NMR data of the above compound of formula I product are consistent with example 21.

Example 23 Preparation of a compound of formula I

[0270] A compound of formula II in THF (45 g, including about 1 g of the compound of formula II, 1 eq) that was not subjected to a post-treatment and was prepared in example 14 was added to a flask, and the solution was concentrated by a rotary evaporator to 3 mL. Toluene (5 mL) was added to the flask, was then subjected to rotary evaporation to 3 mL, the step was repeated twice to obtain black oily concentrated residues.

[0271] 1.0 g of black oily concentrated residues obtained in the previous step trimethyl orthoacetate (1.32 g, 3.0 eq) and THF (25 mL) were added to a 100 mL three-necked flask; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (0.08 g, 0.2 eq) was added to the three-necked flask; under nitrogen protection, the system was heated to 65 - 70 °C (internal temperature) and reacted for 5 hours; sampled and tested. HPLC monitoring showed 0.48% of the compound of formula II in the reaction liquid was left, and the purity of the compound of formula I was 90.10%; allowed the system to reflux and reacted for 5 hours, sampled and tested. HPLC showed the compound of formula II in the reaction liquid was completely transformed, and the purity of the compound of formula I was 91.79%.

Example 24 Preparation of a compound of formula I

[0272] The compound of formula II in THF (450 g, including about 10.0 g of the compound of formula II, 1 eq) not subjected to a post-treatment and was prepared in example 14 was added to a flask, the solution was concentrated by a rotary evaporator to 20 - 30 mL. Toluene (50 mL) was added to the flask, the system was subjected to rotary evaporation to 20-30 mL, the step was repeated twice to obtain black oily concentrated residues, and the residues were dissolved in THF (20 mL, 2 V) until the solution was clear.

[0273] The THF solution obtained in the previous step, trimethyl orthoacetate (13.2 g, 3.0 eq) and THF (230 mL) were added to a 500 mL three-necked flask; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (0.8 g, 0.2 eq) was added to the three-necked flask; under nitrogen protection, the system was heated to 65 - 70 °C (internal temperature) and reacted for 10 hours; sampled and tested. HPLC monitoring showed 0.7% of the compound of formula II in the reaction liquid was left, and the purity of the compound of formula I was 90.1%.

EP 4 385 511 A2

[0274] The system was cooled to room temperature, water (20 mL, 2 V) was added; the system was concentrated by a rotary evaporator until there was basically no fraction flowing out; water (100 mL, 10 V) was supplemented to the system; the pH value of the system was adjusted to 9 with saturated potassium carbonate solution; the system was filtered, the filter cake was washed with water (50 mL, 5 V) and MTBE (50 mL, 5 V) sequentially; the washed filter cake was dried under vacuum at 50 °C for 16 hours to obtain 9.2 g of earthy yellow compound of formula I crude product, with an HPLC purity of 89.7% and a crude product yield of 84.5%.

[0275] Purification of a compound of formula I:

The compound of formula I crude product (5.0 g) with a purity of 89.7% obtained in the previous step and ethanol (50 mL) were added to a flask, the system was stirred at room temperature for 20 minutes, until the materials were basically dissolved and the solution was clear; silica gel (5.0 g, 1X) was added to the system, the system was concentrated by a rotary evaporator to dryness for later use; the solid crude product obtained by the rotary evaporation was allowed to pass through a silica gel column (40 g, 8X), and the column was eluted with a mixed solution of ethyl acetate and petroleum ether ($V_{EA}: V_{PE} = 2:1$); through a TLC test, the fractions containing the compound of formula I were collected and concentrated to dryness to obtain 0.5 g of the compound of formula I with a purity of 96.5% and 3.2 g of light yellow compound of formula I with an HPLC purity of 99.3%.

[0276] The compound of formula I with an HPLC purity of 99.3% and MTBE (30 mL) were added to a flask, the system was heated to reflux, and allowed the system to reflux for 1 hour; the system was cooled to 5 - 10 °C, the temperature was maintained and the system was stirred for 1 hour, filtered, the filter cake was washed with MTBE (5 mL), and the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 2.8 g of light yellow to off-white solids of the compound of formula I, with an HPLC purity of 99.8%, without impurities with the content that was > 0.1%. The total yield of the second and third steps was 47.3%.

Example 25 Preparation of a compound of formula I

[0277] The compound of formula II in THF (130 g, including about 56 g of the compound of formula II, 1 eq) not subjected to a post-treatment and was prepared in example 14 was added to a flask, the solution was exchanged by toluene (280 mL × 2) and was concentrated to 120 - 130 mL, then was exchanged with THF (280 mL) and was concentrated to 120 - 180 mL, THF (500 mL) was added until the solid was dissolved and the solution was clear, the obtained solution was transferred to a 2L three-necked flask, THF (700 mL) and trimethyl orthoacetate (74.0 g, 3.0 eq) were added to the system; under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 10 hours; sampled and tested. HPLC monitoring showed 0.1% of the compound of formula II in the reaction liquid was left, and the purity of the compound of formula I was 93.2%.

[0278] The above reaction liquid was cooled to room temperature, some of the reaction liquid (corresponding to the amount that containing 55g of the compound of formula II before the reaction) was taken, water (110 mL, 2 V) was added; the system was concentrated by a rotary evaporator to 110 - 160 mL (2 - 3 V); water (400 mL, 7 V) was added to the concentrated residue slowly; the system was stirred for 30 minutes at room temperature, water (440 mL, 8V) was then added, the system was stirred for 30 minutes at room temperature (25 - 30 °C); the pH of the system was adjusted to 8 - 9 with 50% potassium carbonate solution (1.5g (the mass of all the solution)); the system was stirred for 30 minutes at room temperature (25 - 30 °C); the temperature of the system was cooled to 10 - 15°C, and the system was stirred for 2 hours under 10 - 15°C, and was subjected to suction filtration, the filter cake was dried at 50 °C for 24 hours to obtain 55g earthy yellow compound of formula I with a purity of 96.6%, a content of 87.53%, and a content yield of a crude product of 80.4%.

[0279] Purification of a compound of formula I:

The compound of formula I crude product (55 g) with an HPLC purity of 96.6% obtained in the previous step, silica gel (110g, 2X) and ethanol (500 mL) were added to a flask; the system was heated to 50 °C, and was stirred for 30 minutes at 50 °C; the system was concentrated by a rotary evaporator until there was basically no fraction flowing out, then was exchanged with n-heptane (200 mL) to dryness, silica gel was mixed with the system, the system was eluted by a column packed with silica gel (550 g, 10X); the eluent was a mixed solution of ethyl acetate and n-heptane ($V_{EA}: V_{n\text{-heptane}} = 1:1$ to pure EA); through a TLC test, compound of formula I component A and cross component B were collected.

[0280] The component A was concentrated by a rotary evaporator to about 100 mL, the concentrated residue was exchanged with about 200 mL of methanol twice, then was exchanged with MTBE (about 200 mL) twice, then about 300 mL of MTBE was added thereto; the system was heated to reflux and allowed the system to reflux for 1 hour, then the system was cooled to room temperature (25 - 30 °C), and was stirred at room temperature (1 hour); the system was cooled to 5 - 10 °C, stirred between 5 - 10 °C for 2 hours, filtered, the filter cake was washed with MTBE (30 mL); the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 37.6 g of light yellow solids of compound of formula I, with an HPLC purity of 99.85%, without impurities with a content that was > 0.1%.

[0281] The component B was concentrated by a rotary evaporator to dryness, triturated with MTBE (50 mL) for 1 hour, the system was cooled to 5 - 10 °C, stirred between 5 - 10 °C for 2 hours, filtered, the filter cake was washed with MTBE

EP 4 385 511 A2

(10 mL); the washed filter cake was dried under vacuum at 50 °C for 16 hours to obtain 6 g of light yellow solids of the compound of formula I, with an HPLC purity of 99.35%, and with 2 impurities with a content that was > 0.1%.

[0282] The above compound of formula I (4.8 g) with a purity of 99.35%, MTBE (50 mL) and ethanol (5 mL) were added to a flask, the system was heated to 55 - 60 °C, after allowing the system to reflux for 0.5 hours, the system was cooled to room temperature (25 - 30 °C), stirred at room temperature for 1 hour; the system was cooled to 5 - 10 °C, stirred between 5 - 10 °C for 2 hours, filtered, the filter cake was washed with MTBE (10 mL); the washed filter cake was dried under vacuum at 50 - 55 °C for 16 hours to obtain 4.0 g of the compound of formula I, with a purity of 99.75%, without impurities with a content that was > 0.1%.

Example 26 Preparation of a compound of formula I

[0283] The compound of formula II in THF (14.5 g, containing about 5.0 g of the compound of formula II, 1 eq) not subject to post-treatment and was prepared in example 14 was added to a flask, the solution was exchanged with toluene (25 mL × 2) and concentrated to about 10 - 15 mL, then was exchanged with THF (25 mL) and concentrated to about 10 - 15 mL, then THF (115 mL) was added until the solid was dissolved and the solution was clear, the obtained solution was transferred to a 500 L three-necked flask; trimethyl orthoacetate (6.6 g, 3.0 eq) was added to the system; under nitrogen protection, the system was heated until reflux; pyridine hydrochloride (0.42 g, 0.2 eq) was added to a three-necked flask; under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 15 hours; sampled and tested. HPLC showed 4.0% of the compound of formula II in the reaction liquid was left; trimethyl orthoacetate (0.5 g) and pyridine hydrochloride (0.1 g) were supplemented to the system, allowed the system to reflux between 70 - 75 °C (internal temperature) for 5 hours; sampled and tested. HPLC showed 0.05% of the compound of formula II in the reaction liquid was left and the purity of the compound of formula I was 92.80%.

Example 27 Preparation of a compound of formula I

[0284] The compound of formula II in THF (817 g, containing about 282.6 g of the compound of formula II, 1 eq) not subject to post-treatment and was prepared in example 14 was added to a flask, the solution was exchanged with toluene (about 1400 mL × 2) and concentrated to about 550 - 850 mL, then was exchanged with THF (about 1400 mL × 2) and concentrated to about 550-850 mL, THF (about 6500 mL) was added until the solid was dissolved and the solution was clear, the solution was transferred to a 10 L three-necked flask; trimethyl orthoacetate (374.0 g, 3.0 eq) was added to the system; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (24.0 g, 0.2 eq) was added to a three-necked flask; under nitrogen protection, the system was heated to 70 -75 °C (internal temperature) and reacted for 12 hours; sampled and tested. HPLC showed 4.1% of the compound of formula II in the reaction liquid was left; the purity of the compound of formula I was 85.5%; trimethyl orthoacetate (22 g) and pyridine hydrochloride (1.4 g) were added to the reaction system; under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 5 hours; sampled and tested. HPLC showed 0.7% of the compound of formula II in the reaction liquid was left and the purity of the compound of formula I was 91.4%.

[0285] The reaction liquid obtained in the previous step was cooled to room temperature, 570 g of water was added thereto, the system was concentrated by a rotary evaporator to 600 - 900 mL (2 - 3 V), the concentrated residue was transferred to a 10 L flask, 2000 g of water (about 7V) was added thereto slowly, the system was stirred at room temperature for 1 hour, 2300 g of water (about 8 V) was then added, the system was stirred at room temperature (25 - 30 °C) for 1 hour, the pH value of the system was adjusted to 8 - 9 with 50% potassium carbonate solution (8.5 g); the system was stirred at room temperature (25 - 30 °C) for 1 hour, the system was cooled to 10 - 15 °C, and was stirred between 10 - 15 °C for 2 hours, filtered, the filter cake was washed with water (500 g); the washed filter cake was dried at 50 °C for 72 hours, sampled and the water content was tested. The water content tested with Karl Fischer method was 3.2%. 252 g of earthy yellow compound of formula I crude product was obtained, with an HPLC purity of 97.5%, a content of 89.8%, and a content yield of the crude product was 72.3%.

[0286] Purification of a compound of formula I:

252 g of the compound of formula I crude product obtained in the previous step and ethanol (1004 g, ~1000 mL) were added to a flask; the system was heated to 50 - 60 °C, stirred between 50 - 60 °C for 30 minutes, until the material was basically dissolved and the solution was clear; the system was divided into two equal proportions, silica gel (252 g) was added to each proportion, each proportion was concentrated by a rotary evaporator until there was basically no fraction flowing out; each proportion was exchanged with n-heptane (272 g, ~ 400 mL) until there was basically no fraction flowing out for later use; a silica gel column was compacted and filled with silica gel (3000 g, 200 - 300 meshes) and n-heptane (5.4 kg, ~ 8 L), the crude product obtained in the previous step was separated by column chromatography, and eluted with a mixture of n-heptane and ethyl acetate (1:1 V/V, 15.5 kg, ~ 20 L; 1:2 V/V, 28.5 kg, ~ 35 L; 1:5 V/V, 25.5 kg, ~ 25 L) and pure ethyl acetate (62 kg, 70 L); through TLC test, compound of formula I component A and cross component B were collected.

EP 4 385 511 A2

[0287] The component A was concentrated by a rotary evaporator until there was basically no fraction flowing out; the concentrated residue was transferred to a 2000 mL flask, exchanged with MTBE (370 g, ~ 500 mL) until there was basically no fraction flowing out; MTBE (1330 g, ~ 1800 mL) was added to the concentrated residue; the system was heated to reflux and allowed the system to reflux for 1 hour; the system was cooled to room temperature (25 - 30 °C), stirred at room temperature (1 hour); cooled to 5 - 10 °C, stirred between 5 - 10 °C for 2 hours; filtered, the filter cake was washed with MTBE (75 g, ~ 100 mL); the HPLC purity of the washed filter cake was 99.9%; the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 190 g of the compound of formula I, with an HPLC purity of 99.9%, and a water content by a KF test of 0.07%.

[0288] The component B was concentrated by a rotary evaporator to dryness; the obtained solid was transferred to a 500 mL single-necked flask; the system was exchanged with MTBE (85 g, ~ 120 mL) until there was basically no fraction flowing out; MTBE (200 g, ~ 300 mL) and methanol (23 g, ~ 30 mL) were added to the concentrated residue; the system was heated to reflux and allowed the system to reflux for 1 hour; the system was cooled to room temperature (25 - 30 °C), stirred at room temperature (1 hour); cooled to 5 - 10 °C, stirred between 5 - 10 °C for 1 hour; filtered, the filter cake was washed with MTBE (22 g, ~ 30 mL); the washed filter cake was dried under vacuum at 50 °C for 16 hours to obtain 20 g of light yellow solid of the compound of formula I, with an HPLC purity of 99.6%.

Example 28 Preparation of a compound of formula I

[0289] The compound of formula II in THF (27.0 g, containing about 5.0 g of the compound of formula II, 1 eq) prepared in example 19 was added to a flask, the solution was exchanged with toluene (25 mL×2) and was concentrated to about 10 - 15 mL; then was exchanged with THF (25 mL) and was concentrated to about 10 - 15 mL; THF (115 mL) was added until the solid was dissolved and the solution was clear, and the solution was transferred to a 500 mL three-necked flask; trimethyl orthoacetate (6.6 g, 3.0 eq) was added to the system; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (0.42 g, 0.2 eq) was added to a three-necked flask; under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 12 hours; sampled and tested. HPLC showed 0.19% of the compound of formula II in the reaction liquid was left, and the purity of the compound of formula I was 93.3%.

Example 29 Preparation of a compound of formula I

[0290] The compound of formula II in THF (5.40 g, containing about 1.0 g of the compound of formula II, 1 eq) prepared in example 19 was transferred to a flask, the solution was exchanged with toluene (4.3 kg×2) and concentrated to about 2 L, then was exchanged with THF (4.3 kg) and concentrated to about 2 L, after THF (19 kg) was added, the system was transferred into a 50 L reactor; trimethyl orthoacetate (1.32 kg, 3.0 eq) was added to the system; under nitrogen protection, the system was heated to reflux; pyridine hydrochloride (85.0 g, 0.2 eq) was added to a three-necked flask; under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 12 hours; sampled and tested. HPLC showed 4.22% of the compound of formula II in the reaction liquid was left; the purity of the compound of formula I was 80.30%, trimethyl orthoacetate (80 g) and pyridine hydrochloride (5 g) were supplemented to the reaction system, under nitrogen protection, the system was heated to 70 - 75 °C (internal temperature) and reacted for 5 hours; sampled and tested. HPLC showed 0.56% of the compound of formula II in the reaction liquid was left and the purity of the compound of formula I was 92.48%.

[0291] The reaction liquid of the previous step was cooled to 25°C, 2.0 kg of water was added; a distillation under reduced pressure was conducted at $45 \pm 5^\circ\text{C}$ to 2 L volume, 2 L of water was added to the rotary evaporation flask, the material liquid in the flask was transferred to a reactor, 5 kg of water was added slowly, stirred at 25°C for 1 hour; 5 kg of water was added to the reactor, stirred at 25°C for 1 hour; 32 g of 50% potassium carbonate solution was added to the reactor dropwise, the pH of the system was adjusted to 8 - 9, the system was stirred at 25 °C for 1 hour; the material liquid in the reactor was cooled to 10 - 15 °C, stirred for 2 hours; the system was filtered, the filter cake was washed with 2 kg of water, and was then dried under vacuum at 45 - 55 °C for 48 hours to obtain 0.91 kg of the compound of formula I, with a water content of 0.2% tested with a KF method, an HPLC purity of 95.71%, a content of 85.39% and a content yield of 71.4%.

[0292] Purification of a compound of formula I:

The compound of formula I (0.91 kg) with an HPLC purity of 95.71% obtained in the previous step and ethanol (3.6 kg) were added to a 20 L rotary evaporation flask; the system was heated to 50 - 60 °C, stirred between 50 - 60 °C for 30 minutes, the material was dissolved and the solution was basically clear; silica gel (1.82 kg) was added to the above rotary evaporation flask, the system was concentrated under reduced pressure between 50 - 60 °C to dry powder; n-heptane (1.82 kg) was added to the above rotary evaporation flask, the system was concentrated and exchanged between 40 - 50 °C to dry powder; an appropriate cleaned column was prepared, silica gel (11 kg, 200 - 300 meshes) was added to the column, the column was compacted with nitrogen; n-heptane (27 kg) was added to the column and the column was compacted with nitrogen; the silica gel-like compound of formula I concentrated to dryness in the previous step was

EP 4 385 511 A2

added to the column, the column was eluted with n-heptane (18 kg), a mixture of n-heptane/ethyl acetate (1:1 V/V, 72 kg; 1:2 V/V, 158 kg; 1:5 V/V, 142 kg), a mixture of n-heptane/ethyl acetate (1:5 V/V, 70 kg), a mixture of n-heptane/ethyl acetate (1:5 V/V, 175 kg), and ethyl acetate (205 kg) sequentially, a TLC test was conducted, and a compound of formula I component A and cross component B were collected.

5 **[0293]** The component A was added to a 50 L reactor, the system was concentrated under vacuum at 40 - 50 °C to the minimum stirring volume (~ 6 L); MTBE (5.0 kg×5) was added to the reactor, the system was concentrated and exchanged 5 times; MTBE (1.2 kg) was added to the reactor, the system was heated to reflux (50 - 60 °C), and allowed the system to reflux for 1 hour with the temperature maintained; the system was cooled to 20 - 30 °C, and was stirred (1 hour) with the temperature maintained; the system was cooled to 5 - 10 °C, stirred between 5 - 10 °C for 2 hours; 10 filtered, the filter cake was washed with MTBE (0.25 kg); 0.62 kg of the compound of formula I was obtained, with an HPLC purity of 100.0%; the filter cake was dried at 50 °C for 16 hours to obtain 0.55 kg of the compound of formula I, with a water content of 0.04% tested by a KF method and a Pd residue < 2 ppm.

15 **[0294]** The component B was added to the reaction flask, the system was concentrated under vacuum in a 45 - 50 °C water bath to the minimum stirring volume (about 1 L); MTBE (1.5 kg ×2) was added to the reaction flask, the system was heated in a 45 - 50 °C water bath, distilled under vacuum to the minimum stirring volume, exchanged twice; MTBE (1.5 kg) and absolute ethanol (0.14 kg) were added to the reaction flask, the system was heated to 50 - 60 °C, stirred for 1 hour; the material liquid in the reaction flask was cooled to 20 - 25 °C, the system was stirred for 1 hour with the temperature maintained; cooled, the temperature of the material liquid in the reaction flask was lowered to 6 - 10 °C, the temperature was maintained and the system was stirred for 2 hours; filtered, the filter cake was washed with MTBE 20 (0.24 kg); the filter cake, MTBE (1.5 kg)/absolute ethanol (0.14 kg) were added to the reaction flask; the system was stirred, heated to raise the temperature to 50 - 60 °C, stirred for 1 hour with the temperature maintained; cooled, the temperature of the material liquid in the reaction flask was lowered to 20 - 25 °C, the temperature was maintained and the system was stirred for 1 hour; cooled, the temperature of the material liquid in the reaction flask was lowered to 6 - 10 °C, the temperature was maintained, and the system was stirred for 2 hours; filtered, the filter cake was washed 25 with MTBE (0.24 kg); 115 g of wet products were obtained, with an HPLC purity of 99.8%, and a maximum individual impurity content of 0.09%; the wet products were dried between 45 - 55 °C, under a vacuum degree ≤ - 0.080 MPa for 16 hours; 0.10 kg of the compound of formula I was obtained, with a water content of 0.08% tested with a KF method, an HPLC purity of 99.8%, a maximum individual impurity content of 0.09%, and a Pd residue ≤ 2 ppm.

30 **[0295]** Unless otherwise specified, the compound of formula I finally purified and prepared in example 27 is taken as the starting material in the following examples.

Preparation of crystal form 1 of a compound of formula I

Example 30

35 **[0296]** The wet product and the crude product of the filter cake of the compound of formula I obtained in example 21 were combined, dissolved with methanol (40 mL); a silicon based metal eliminator (4.0 g) and an activated carbon (1.0 g) were added to the methanol solution, the system was heated to 50 °C and stirred for 1 hour; the system was cooled to 10 ± 5 °C and was stirred at the temperature for 0.5 hour; filtered, the filter cake was washed with MTBE (15 mL); 40 the filter cake was dried under vacuum at 50 °C for 16 hours to obtain 2.5 g of off-white solid of the compound of formula I, with an HPLC purity of 98.4%. Upon testing, the solid was crystal form 1 of a compound of formula I. See Figures 1 - 4 for the XRPD pattern, the DSC thermogram, the TGA thermogram and the DVS isotherm plot.

Preparation of crystal form A of a hydrochloride of a compound of formula I

Example 31

45 **[0297]** About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to 50 prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.73 mL of hydrochloric acid in acetone (the concentration of the hydrochloric acid in acetone was 25 mg/mL) was slowly added dropwise, white precipitates were produced, bottle cap was covered tightly at room temperature and the system was stirred for 1 day, the suspension was then centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain compound of formula I hydrochloride solid. Upon testing, the solid 55 was crystal form A of a hydrochloride of a compound of formula I. See Figures 6 and 7 for the XRPD pattern and the DSC thermogram.

Preparation of crystal form B of a hydrochloride of a compound of formula I**Example 32**

5 [0298] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of ethyl acetate was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in ethyl acetate. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.73 mL of hydrochloric acid in ethyl acetate (the concentration of the hydrochloric acid in ethyl acetate was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was then centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I hydrochloride solid. Upon testing, the solid was crystal form B of a hydrochloride of a compound of formula I. See Figures 8 and 9 for the XRPD pattern and the DSC thermogram.

Preparation of a compound of formula I hydrochloride crystal form C**Example 33**

20 [0299] 5 mg of crystal form A of a hydrochloride of a compound of formula I prepared in example 31 was weighed and placed into a small bottle, a suitable amount of methanol was added, magnetic stirring of the sample suspension obtained was conducted at room temperature overnight, the system was centrifuged to separate the solid and liquid, the solid was collected, and dried under vacuum overnight at 40 °C to obtain a compound of formula I hydrochloride solid. Upon testing, the solid was crystal form C of a hydrochloride of a compound of formula I. See Figures 10 and 11 for the XRPD pattern and the DSC thermogram.

Examples 34 - 36

30 [0300] The crystallization method that was the same as that in example 33 was adopted. The solvent was changed to acetonitrile, n-heptane and methyl ethyl ketone to prepare crystal form C of a hydrochloride of a compound of formula I. Upon testing, the XRPD pattern of the solid compounds prepared in examples 35 - 37 are consistent with Figure 10.

Preparation of crystal form D of a sulfate of a compound of formula I**Example 37**

35 [0301] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.77 mL of sulfuric acid in acetone (the concentration of the sulfuric acid in acetone was 25 mg/mL), white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was then centrifuged, the collected solid was dried under vacuum at 40 °C overnight to obtain compound of formula I sulfate solid. Upon testing, the solid was crystal form D of a sulfate of a compound of formula I. See Figures 12 and 13 for the XRPD pattern and the DSC thermogram.

Preparation of crystal form E of a phosphate of a compound of formula I**Example 38**

50 [0302] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.86 mL of phosphoric acid in acetone (the concentration of the phosphoric acid in acetone was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was then centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain compound of formula I phosphate solid. Upon testing, the solid was crystal form E of a phosphate of a compound of formula I. See Figures 14 and 15 for the XRPD pattern and the DSC thermogram.

55 [0303] The salt-forming percentage test of crystal form E of a phosphate of a compound of formula I was conducted with ion chromatography:

EP 4 385 511 A2

[0304] About 0.5 g of crystal form E of a phosphate of a compound of formula I was weighed and placed into a liquid phase small bottle, completely dissolved with 1 mL of water, then was used as a sample solution. The bulk (1000 ppm) of phosphoric acid standard solution was diluted at 10 and 20 folds with water, to obtain 100 ppm and 50 ppm standard sample solution respectively.

[0305] Ion chromatography tests were conducted on the the sample solution and the standard solution, respectively, see Table 16 for the test method. A standard curve was plotted with the peak area of the countra-ion on the ion chromatogram corresponding to the concentration of the countra-ion in the standard sample solution, the concentration of the countra-ion in each sample was calculated using an external standard method, the content of the countra-ion in crystal form E of a phosphate of a compound of formula I was calculated, to determine the salt-forming percentage of a compound of formula I and the corresponding countra-ion in crystal form E of a phosphate of a compound of formula I.

Table 16 Ion chromatography determination method

Instrument model	ICS-2000 + AS40 Automated sampler
Chromatographic column	IonPac®AS11-HC 4×250mm
Column temperature	30°C
Eluent	10Mm KOH solution
Flow rate	1.00 mL/min
Suppressor	Dionex AERS 500 4mm
Electric current of suppressor	25 mA
Run time	15 min

[0306] See Table 17 for the countra-ion content in crystal form E of a phosphate of a compound of formula I. In the 0.5 g of crystal form E of a phosphate of a compound of formula I, the compound of formula I and phosphoric acid were fed at a molar ratio of 1:1.1 and reacted. The actually measured content of phosphate anion in the obtained phosphate was 25.6%, which was basically consistent with the theoretical content of 24.9%. The salt-forming molar ratio was 1:1.04 (the compound of formula I: phosphoric acid).

Table 17 Countra-ion content determination results in crystal form E of a phosphate of a compound of formula I

Sample name	Countra-ion at different salt-forming percentage		Measured countra-ion content %
	Theoretical content %		
Crystal form E of a phosphate of a ompound of formula I	Free base: phosphoric acid = 1:1 salt formation	24.9%	25.6%

Example 39

[0307] 5 mg of crystal form E of a phosphate of a compound of formula I prepared in example 38 was weighed and placed into a small bottle, a suitable amount of methanol was added, magnetic stirring of the sample suspension was conducted at room temperature overnight, the system was centrifuged and the solid and liquid were separated, the solids were collected, and dried under vacuum overnight at 40 °C to obtain a compound of formula I phosphate solid. Upon testing, the solid was still crystal form E of a phosphate of a compound of formula I. Its XRPD pattern is consistent with Figure 14.

Examples 40 - 42

[0308] The method that was the same as that in example 39 was adopted. The solvent was changed to acetonitrile, n-heptane and methyl ethyl ketone to prepare crystal form E of a phosphate of a compound of formula I. Upon testing, the XRPD pattern of the solid compounds prepared in examples 40 - 42 were consistent with Figure 14.

Preparation of crystal form F of a phosphate of a compound of formula I**Example 43**

5 [0309] About 500 mg of a compound of formula I was weighed and placed into a small bottle, 20 mL of acetone was added, the system was subjected to ultrasonication and heating until the sample was completely dissolved to obtain 25 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic heating stirrer, magnetic stirring was conducted and 8.57 mL of phosphoric acid in acetone (the concentration of the phosphoric acid in acetone was 25 mg/mL) was slowly added dropwise overnight. The suspension was subjected to suction filtration, the solids were dried under vacuum at 50 °C, and the solid were collected and placed into a 100 mL glass bottle. Methanol was slowly added dropwise, magnetic stirring was conducted at room temperature, until the solution was clear, and then the solution was diluted with anti-solvent acetic acid isopropyl ester at 10 folds. The system was stirred overnight and the suspension was filtered, the solids were vacuum dried at 50 °C, and the solids were collected to obtain compound of formula I phosphate solids. Upon testing, the solid was crystal form F of a phosphate of a compound of formula I. See Figures 16 - 19 for the XRPD pattern, the DSC thermogram, and DVS isotherm plot and the XRPD pattern after the DVS test.

Preparation of crystal form G of a mesylate of a compound of formula I**Example 44**

20 [0310] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.73 mL of methanesulfonic acid in acetone (the concentration of the methanesulfonic acid in acetone was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I mesylate solid. Upon testing, the solid was crystal form G of a mesylate of a compound of formula I. See Figures 20, 21 and 22 for the XRPD pattern, the DSC thermogram and the DVS isotherm plot.

Preparation of crystal form H of a hydrobromide of a compound of formula I**Example 45**

35 [0311] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 1.32 mL of hydrobromic acid in acetone (the concentration of the hydrobromic acid in acetone was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I hydrobromide solid. Upon testing, the solid was crystal form H of a hydrobromide of a compound of formula I. See Figures 23 and 24 for the XRPD pattern and the DSC isotherm plot.

Example 46

45 [0312] 5 mg of crystal form H of a hydrobromide of a compound of formula I prepared in example 46 was weighed and placed into a small bottle, a suitable amount of acetonitrile was added, magnetic stirring of the sample suspension was conducted at room temperature overnight, the system was centrifuged and the solid and liquid were separated, the solid was collected, and dried under vacuum overnight at 40 °C to obtain a compound of formula I hydrobromide solid. Upon testing, the solid was still crystal form H of a hydrobromide of a compound of formula I. Its XRPD pattern is consistent with Figure 23.

Example 47

55 [0313] The method that was the same as that in example 46 was adopted. The solvent was changed to methyl ethyl ketone to prepare crystal form H of a hydrobromide of a compound of formula I. Upon testing, the XRPD pattern of the solid compound prepared in example 48 is consistent with Figure 23.

Preparation of crystal form J of a hydrobromide of a compound of formula I**Example 48**

5 [0314] About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of ethyl acetate was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in ethyl acetate. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 1.32 mL of hydrobromic acid in ethyl acetate (the concentration of the hydrobromic acid in ethyl acetate was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I hydrobromide solid. Upon testing, the solid was crystal form J of a hydrobromide of a compound of formula I. See Figure 25 for the XRPD pattern.

Preparation of crystal form K of a hydrobromide of a compound of formula I**Example 49**

15 [0315] 5 mg of crystal form H of a hydrobromide of a compound of formula I prepared in example 46 was weighed and placed into a small bottle, a suitable amount of n-heptane was added, magnetic stirring of the sample suspension was conducted at room temperature overnight, the system was centrifuged and the solid and liquid were separated, the solid was collected, and dried overnight under vacuum at 40 °C to obtain a compound of formula I hydrobromide solid. Upon testing, the solid was crystal form K of a hydrobromide of a compound of formula I. See Figures 26 - 29 for the XRPD pattern, the DSC thermogram, and the DVS isotherm plot and the XRPD pattern after the DVS test.

Preparation of crystal form L of a fumarate of a compound of formula I**Example 50**

25 [0316] About 50 mg of a compound of formula I was weight and placed into a small bottle, 2.5 mL of ethyl acetate was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in ethyl acetate. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 0.87 mL of fumaric acid in ethanol (the concentration of the fumaric acid in ethanol was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I fumarate solid. Upon testing, the solid was crystal form L of a fumarate of a compound of formula I. See Figures 30 and 31 for the XRPD pattern and the DSC thermogram.

Example 51

30 [0317] The crystallization method that was the same as that in example 50 was adopted. Ethyl acetate was changed to acetone to prepare crystal form L of a fumarate of a compound of formula I. Upon testing, the XRPD pattern of the solid compound prepared in example 51 is consistent with Figure 30.

Preparation of crystal form M of a benzene sulfonate of a compound of formula I**Example 52**

45 [0318] About 50 mg of a compound of formula I wa weighed and placed into a small bottle, 2.5 mL of acetone was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 1.34 mL of benzenesulfonic acid in acetone (the concentration of the benzenesulfonic acid in acetone was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I benzene sulfonate solid. Upon testing, the solid was crystal form M of a benzene sulfonate of a compound of formula I. See Figures 32 - 35 for the XRPD pattern, the DSC thermogram, and the DVS isotherm plot and the XRPD pattern after the DVS test.

Preparation of crystal form N of a citrate of a compound of formula I**Example 53**

5 **[0319]** About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of ethyl acetate was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in ethyl acetate. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 1.58 mL of citric acid in ethyl acetate (the concentration of the citric acid in ethyl acetate was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuged, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I citrate solid. Upon testing, the solid was crystal form N of a citrate of a compound of formula I. See Figures 36 and 37 for the XRPD pattern and the DSC thermogram.

Preparation of crystal form O of a tartrate of a compound of formula I**Example 54**

20 **[0320]** About 50 mg of a compound of formula I was weighed and placed into a small bottle, 2.5 mL of ethyl acetate was added, the system was subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 20 mg/mL a compound of formula I in ethyl acetate. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 1.12 mL of L-(+)-tartaric acid in ethyl acetate (the concentration of the L-(+)-tartaric acid in ethyl acetate was 25 mg/mL) was slowly added dropwise, white precipitates were produced, the bottle cap was covered tightly at room temperature, the system was stirred for 1 day, the suspension was centrifuge, and the collected solid was dried under vacuum at 40 °C overnight to obtain a compound of formula I L-(+)-tartrate solid. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. See Figures 38 - 42 for the XRPD pattern, the DSC thermogram, the TGA thermogram and the DVS isotherm plot and the XRPD pattern after the DVS test.

[0321] The salt-forming percentage test of crystal form O of a tartrate of a compound of formula I was conducted with a ¹H NMR method:

30 About 5 mg of a compound of formula I and crystal form O of a tartrate of a compound of formula I were weighed and placed into a magnetic tube respectively, the samples were dissolved with 0.6 mL of DMSO d6 until each solution was clear, and each the sample solution was scanned with Bruker AVANCE 400MHz nuclear magnetic resonance spectrometer using the general method to collect the ¹H-NMR data of the samples.

35 **[0322]** The NMR spectrums showed that the ¹H NMR of crystal form O of a tartrate of a compound of formula I contained one more hydrogen than that of the compound of formula I. Because tartaric acid is a diacid. It can be seen that the salt-forming molar ratio of the free base to tartaric acid is 2:1. See Figures 43 and 44 for the specific information.

[0323] The salt-forming percentage test of crystal form O of a tartrate of a compound of formula I was conducted with a chemical titration:

40 A nonaqueous titration instrument was adopted, and methanolic potassium hydroxide VS was adopted to titrate the sample. The content of tartaric acid in the sample was calculated with the formula according to the concentration of the titrant and the volume consumed of the titrant.

Table 18 Blank titration condition

System	Tiamo 2.2 light
Electrode	pH-electrode filled with 1 M LiCl/EtOH, Metrohm, No. 6.0299.010
Titrator	Metrohm 809 Titrando
Titrant	0.1 mol/ L methanolic potassium hydroxide VS
Titration volume	1 mL

Table 19 Sample titration condition

System	Tiamo 2.2 light
Electrode	pH-electrode filled with 1 M LiCl/EtOH, Metrohm, No. 6.0299.010

EP 4 385 511 A2

(continued)

System	Tiamo 2.2 light
Titrator	Metrohm 809 Titrande
Titrant	0.1 mol/ L methanolic potassium hydroxide VS
Titration volume	1 mL
Sample concentration	1.25 mg/mL

[0324] 80 mL of methanol solution was taken and placed into a titration cup, titration was conducted according to the blank titration condition in triplicate. 160 mg of standard potassium acid phthalate that has been dried to a constant weight at 105 °C was weighed accurately, 50 ml of methanol solution was added, the system was subjected to ultrasonication to dissolve, and was then transferred to the titration cup, and titration was conducted according to the sample titration condition in triplicate. 100 mg of crystal form O of a tartrate of a compound of formula I was weighed accurately, 80 mL of methanol solution was added, the system was subjected to ultrasonication to dissolve and was then transferred to the titration cup, and titration was conducted according to the sample titration condition in triplicate.

[0325] The titer (T) of the titrant was calculated according to the following formula:

$$T \text{ (mol/L)} = \frac{W}{M \times (V - V_0)} \times 1000$$

W: Weighting of standard potassium hydrogen phthalate (g)

V: The volume of methanolic potassium hydroxide VS consumed for titrating standard potassium acid phthalate solution (mL)

V₀: The volume of methanolic potassium hydroxide VS consumed for titrating blank solution (mL)

M: The molecular weight of standard potassium hydrogen phthalate

[0326] The tartaric acid content in the sample was calculated according to the following formula:

$$\text{Assay (\%)} = \frac{T \times (V - V_0)}{1000 \times W \times 2} \times 100\%$$

T: The titer of a calibrated methanolic potassium hydroxide VS (mol/L)

V: The volume of methanolic potassium hydroxide VS consumed for titrating sample solution (mL)

V₀: The volume of methanolic potassium hydroxide VS consumed for titrating blank solution (mL)

M: The molecular weight of tartaric acid

W: Sample weighting (g)

[0327] The final titration test results indicated that the content of tartaric acid in crystal form O of a tartrate of a compound of formula I was 19.2 w/w% and 21.1 w/w%, respectively, which was consistent with the theoretical value 20.2 w/w% when the molar ratio of the free base: tartaric acid was 2:1.

Example 55

[0328] About 500 mg of a compound of formula I was weighed and placed into a small bottle, 20 mL of acetone was added, the system as subjected to ultrasonication and heating until the compound was completely dissolved, to prepare 25 mg/mL a compound of formula I in acetone. The sample bottle was placed on a magnetic stirring plate, magnetic stirring was conducted and 11.2 mL of L-(+)-tartaric acid in acetone (the concentration of the L-(+)-tartaric acid in acetone

EP 4 385 511 A2

was 25 mg/mL) was slowly added dropwise, the system was stirred overnight, filtered, and the solid was dried under vacuum at 50 °C to obtain a compound of formula I L-(+)-tartrate solid. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

5 Example 56

[0329] 2.0 g of a compound of formula I with an HPLC purity of 99.9% prepared by the purification of component A in example 27 and 40 mL of acetone (20 V) were added to flask 1#, the system was stirred to dissolve until the solution was clear; 0.61 g of L-(+)-tartaric acid and 40 mL of acetone (20 V) were added to flask 2#, the system was stirred to dissolve until the solution was clear; the solution in flask 2# was added to flask 1# within 2 - 3 minutes; the temperature of the system was raised to 50 - 60 °C and the system was stirred for 2 hours; the system was cooled to room temperature; and was concentrated to about 40 mL; the system was stirred at room temperature between 25 - 30 °C for 1 hour; the system was cooled to 5 - 10 °C, stirred between 5 - 10 °C for 1 hour; filtered, and the filter cake was dried with an air blower between 50 - 55 °C for 16 hours to obtain 2.4 g of product, with an HPLC purity of 99.6% and a yield of 95.6%. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

Example 57

[0330] 2.0 g of a compound of formula I with an HPLC purity of 99.9% prepared by the purification of component A in example 27 and 40 mL of acetone (20 V) were added to flask 1#, the temperature was raised to 50 - 55 °C, the system was stirred to dissolve until the solution was clear; 0.61 g of L-(+)-tartaric acid and 40 mL of acetone (20 V) were added to flask 2#, the temperature was raised to 50 - 55 °C, the system was stirred until the solution was clear the solution in flask 2# was added to flask 1# within 2 - 3 minutes; the system was stirred between 45 - 50 °C for 2 hours; the system was concentrated under vacuum to about 40 mL at 45 - 50 °C; cooled to 20 - 25 °C, stirred for 1 hour; cooled to 5 - 10 °C, stirred between 5 - 10 °C for 1 hour; filtered, and the filter cake was dried with an air blower between 50 - 55 °C for 16 hours to obtain 2.4 g of product, with an HPLC purity of 99.8% and a yield of 95.62%. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

30 Example 58

[0331] 36.0 g of a compound of formula I with an HPLC purity of 99.85% prepared by the purification of component A in example 26 and 720 mL of acetone (20 V) were added to flask 1#, the temperature was raised to 50 - 55 °C, the system was stirred to dissolve until the solution was clear; 11.0 g of L-(+)-tartaric acid and 720 mL of acetone (20 V) were added to flask 2#, the temperature was raised to 50 - 55 °C, the system was stirred to dissolve until the solution was clear; the solution in flask 2# was added to flask 1# between 45 - 55 °C within 2 - 3 minutes; the system was stirred between 45 - 50 °C for 2 hours; concentrated under vacuum to about 720 mL; cooled to 5 - 10 °C, stirred for 1 hour between 5 - 10 °C; filtered, and the filter cake was dried with an air blower between 50 - 55 °C for 16 hours to obtain 43.6 g of product, with an HPLC purity of 99.96% and a yield of 96.6%. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

Example 59

[0332] 189.6 g of a compound of formula I with an HPLC purity of 99.9% prepared by the purification of component A in example 28 and 3792 mL of acetone (20 V) were added to flask 1#, the temperature was raised to 50 - 55 °C, the system was stirred to dissolve until the solution was clear; 57.6 g of L-(+)-tartaric acid and 3792 mL of acetone (20 V) were added to flask 2#, the temperature was raised to 50 - 55 °C, the system was stirred to dissolve until the solution was clear; the solution in flask 2# was added to flask 1# between 45-55°C; the system was stirred between 45 - 50 °C for 2 hours; concentrated under vacuum to about 3800 mL (about 20 V); cooled to 17 - 21 °C, stirred for 1 hour; cooled to 5 - 10 °C, stirred between 5 - 10 °C for 1 hour; filtered, and the filter cake was dried with an air blower between 50 - 55 °C for 28 hours to obtain 223.2 g of product, with an HPLC purity of 99.98% and a yield of 94.0%. Upon testing, the solid was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

Example 60

[0333] 0.65 kg of a compound of formula I with an HPLC purity of 100.0% prepared by the purification of component A in example 30 and 7.7 kg of acetone were added to a rotary flask, the flask was rotated in a 40 - 50 °C water bath for 1 hour to make the solution clear; the clear solution was transferred to PT1 (reactor 1), compressed with nitrogen through

EP 4 385 511 A2

a pipeline filter to the reactor in the purification area; 2.6 kg of acetone was added to the rotary flask, the flask was washed and then transferred to PT1, compressed to the reactor through a pipeline filter; the volume of the reaction liquid in the reactor in the purification room was calibrated to 13.0 L; 7.7 kg of acetone and 0.198 kg of L-tartaric acid were added to the rotary flask, the rotary flask was rotated in a 40 - 50 °C water bath for 1 hour to make the solution clear; the clear solution was transferred to PT1, compressed with nitrogen through a pipeline filter to PT2 (reactor 2); 2.6 kg of acetone was added to the rotary flask, the flask was washed and the system was transferred to PT1, compressed with nitrogen through a pipeline filter to PT2; the temperature of the material liquid in the reactor was heated to 40 - 50 °C, the materials in PT2 was added to the reactor within 1 hour; the temperature of the materials in the reactor was controlled between 40 - 50 °C, the system was stirred and reacted for 4 hours; cooled, the temperature in reactor was lowered to 25 - 35 °C; the system was distilled under vacuum, the vacuum was controlled $\leq - 0.080$ MPa, the material liquid in the reactor was distilled under vacuum to the calibration volume of 13.0 L; the temperature of the material liquid in the reactor was cooled to 15 - 25 °C, and the system was stirred at this temperature for 1 hour; the temperature in the reactor was cooled to 0 - 10 °C; the temperature in the reactor was controlled between 0 - 10 °C and the system was stirred for 1 hour; the system was subjected to suction filtration, the filter cake was washed with acetone (2.5 kg); samples were taken and analyzed. The HPLC purity of the compound of formula I tartrate in the filter cake was 99.98%; the maximum individual impurity: 0.02%; the filter cake was dried for 24 hours at 45 - 55 °C, under vacuum $\leq - 0.080$ MPa; samples were taken and analyzed, acetone residue ≤ 5000 ppm; the oven was cooled to 15 - 25 °C; dried and 0.74 kg of product was obtained, samples were taken and analyzed. The HPLC purity of the compound of formula I tartrate: 99.95%; the maximum individual impurity: 0.03%; the residual solvents comply with the requirements. Upon further testing, the product was crystal form O of a tartrate of a compound of formula I. Its XRPD pattern is consistent with Figure 38.

Experimental section

Experimental example 1: Solubility test

[0334] 4 proportions of each of crystal form 1 of a compound of formula I, crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I with suitable amounts were weighed and each placed into a 4 mL transparent glass bottle, 1 mL of water, simulated gastric fluid (SGF), fasted-state simulated intestinal fluid (FaSSIF) and fed-state simulated intestinal fluid (FeSSIF) were added respectively to obtain sample suspension and the suspension was transferred to a shaker quickly (37 °C, 200 rpm) and was shaken. The samples were observed 5 minutes later, a quantity of sample or medium was supplemented to obtain mild suspension, and sampling was conducted at 30 minutes, 2 hour, 4 hours and 24 hours, respectively. The samples were centrifuged for 10 minutes at 12000 rpm to obtain the supernatant, the supernatant was diluted appropriately and was subjected to high performance liquid chromatography. See Table 20 for chromatographic conditions.

Table 20 High performance liquid chromatographic conditions of the solubility test

Instrument	Agilent 1200 DAD HPLC system
Chromatographic column	Waters XBridge Shield RP18 4.6×150mm, 3.5 μm
Mobile phase	A: 0.1% trifluoroacetic acid solution B: 0.1% trifluoroacetic acid in acetonitrile
Column temperature	30°C
Detector	DAD
Detection wavelength	230 nm
Injection volume	5 μL
Column flow rate	1.0 mL/min
Run time	15 min
Collection time	15 min

EP 4 385 511 A2

(continued)

Elution procedure	Time (min)	A (%)	B (%)
	0.0	95	5
	7.0	65	35
	10.0	5	95
	10.1	95	5
15	95	5	

[0335] The sample concentration was calculated with an external standard method. The test results are shown in Table 21.

Table 21 Results of solubilities of crystal form 1 of a compound of formula I, crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I in water, SGF, FaSSIF, and FeSSIF at different time points

Sample	Medium	Solubility (based on a compound of formula I, mg/mL)			
		30 min	2 h	4 h	24 h
Crystal form 1 of a compound of formula I	Water	1.35	1.73	1.49	1.43
	Simulated gastric fluid (SGF)	16.40	16.96	15.17	16.41
	Fasted-state simulated intestinal fluid (FaSSIF)	2.59	1.86	1.81	1.56
	Fed-state simulated intestinal fluid (FeSSIF)	2.91	2.20	2.61	3.29
Crystal form F of a phosphate of a compound of formula I	Water	35.73	35.14	36.32	39.53
	Simulated gastric fluid (SGF)	155.84	176.85	148.76	174.35
	Fasted-state simulated intestinal fluid (FaSSIF)	57.23	79.34	85.21	93.28
	Fed-state simulated intestinal fluid (FeSSIF)	4.22	3.89	3.56	3.64
Crystal form O of a tartrate of a compound of formula I	Water	12.53	12.11	12.13	12.97
	Simulated gastric fluid (SGF)	31.18	29.85	33.21	30.22
	Fasted-state simulated intestinal fluid (FaSSIF)	9.95	10.03	10.24	11.36
	Fed-state simulated intestinal fluid (FeSSIF)	4.28	4.31	4.41	3.98

[0336] The results showed that crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I can significantly improve the solubility of the compound in water, SGF, and FaSSIF. The solubilities of crystal form F of a phosphate of a compound of formula I in water, SGF, FaSSIF and FeSSIF at 24-

EP 4 385 511 A2

hour time point were 27, 10, 60 and 1 time that of the compound of formula I, respectively; the solubilities of crystal form O of a tartrate of a compound of formula I in water, SGF, FaSSIF and FeSSIF at 24-hour time point were 9, 2, 7 and 1 time that of the compound of formula I, respectively.

5 Experimental example 2: Stability test 1

[0337] About 1 mg of crystal form 1 of a compound of formula I, crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I samples were weighed and each was placed into a 20 mL transparent glass bottle respectively, and each sample was placed to a stability chamber in accelerated conditions (40°C/75%RH, open) and at a high temperature (60°C, sealed). For the open samples, the bottle cap was removed and the bottle neck was covered with an aluminium-foil paper stabbed with pinholes to avoid cross contamination; for the closed samples, the bottled were covered and sealed tightly. Samples were taken at weeks 1 and 2, respectively, the samples were diluted with a diluent (methanol/water (1/1) (v/v)), the liquid phase was injected according to the chromatographic conditions in Table 22 and the sample purities were determined.

Table 22 High performance liquid chromatographic conditions in the solid state stability test

Instrument	Agilent 1200 DAD HPLC system		
Chromatographic column	WATERS Xbridge C 18 250 × 4.6mm 5 μm		
Mobile phase	A: 20 mM dipotassium phosphate solution; B: methanol		
Column temperature	40°C		
Detector	DAD		
Detection wavelength	220 nm		
Injection volume	5.0 μL		
Column flow rate	1.0 mL/min		
Run time	35 min		
Post run time	42 min		
Sample concentration	0.5mg/mL		
Elution procedure	Time (min)	A (%)	B (%)
	0	95	5
	30	25	75
	35	25	75

[0338] The sample purities were calculated with an area normalization method. The test results are shown in Table 23.

Table 23 Results of short-term solid state stabilities of crystal form 1 of a compound of formula I, crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I

Sample	Stability study (purity, Area%)						
	0 day	60°C			40°C/75%RH		
		1 week	2 weeks	Crystal form	1 week	2 weeks	Crystal form
Crystal form 1 of a compound of formula I	99.80	99.81	99.83	Change	99.82	99.82	Change
Crystal form F of a phosphate of a compound of formula I	99.34	99.32	99.16	Unchange	99.33	99.35	Unchange
Crystal form O of a tartrate of a compound of formula I	99.22	99.34	99.36	Unchange	99.31	99.36	Unchange

[0339] The results indicated that the appearances of crystal form 1 of a compound of formula I, crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I didn't change within 2 weeks and they were off-white powder. There were no significant differences in the purities and no obvious increased in impurities, indicating good chemical stabilities within 2 weeks. The XRPD and DSC tests (Figures 45 - 50) indicated that there were no significant differences in the crystal forms and the initial melting points of crystal form F of a phosphate of a compound of formula I and crystal form O of a tartrate of a compound of formula I samples compared with those on Day 0, showing good physical stability of the compound of formula I phosphate and tartrate within 2 weeks at a high temperature (60 °C) and in accelerated conditions (40°C/75%RH). The crystal form of crystal form 1 of a compound of formula I changed at a high temperature (60 °C) and in accelerated conditions (40°C/75%RH). On the basis of the characterization results of the free base before and after the DVS test, it can be seen that the stability of the crystal form of the free base was poor.

Experimental example 3: Stability test 2

[0340] Crystal form O of a tartrate of a compound of formula I tartrate crystal form O was tested with a suspension balance method, a heating-fast/slow cooling crystallization method, an anti-solvent method and a solution volatilization and crystallization method to investigate whether crystal transformation occurred in different solvents and test conditions, so as to further verify its thermodynamic stability.

1. Slow solution volatilization method (EVA)

[0341] 5 proportions (10 mg/proportion) of crystal form O of a tartrate of a compound of formula I prepared in example 55 were weighed and each was placed into a sample bottle, a suitable amount of tetrahydrofuran, ethanol, methanol, acetone and isopropanol (see Table 24 for the specific amounts) were added respectively, the bottled were subjected to ultrasonication to dissolve the samples, the obtained solution was filtered to a new sample bottle with a 0.45 μm nylon membrane, the sample bottle was opened and placed in a fuming cupboard, the solvent was evaporated naturally at room temperature (about 20 - 25 °C), the precipitated solid was collected. Upon testing, the obtained solid in 5 tests was crystal form O of a tartrate of a compound of formula I and their XRPD patterns are consistent with Figure 38.

Table 24 Test conditions and results summary of the slow solution volatilization method

Solvent	Total volume of solvent (mL)	Crystal form
Tetrahydrofuran	1.8	Crystal form O
Ethanol	1.5	Crystal form O
Methanol	0.6	Crystal form O
Acetone	3.6	Crystal form O
Isopropanol	4.8	Crystal form O

2 Suspension balance method (Slurry)

[0342] 18 proportions of crystal form O of a tartrate of a compound of formula I samples, each with a suitable amount were weighed, then a certain amount of tetrahydrofuran, ethanol, ethyl acetate, n-heptane, toluene, methyl tertiary-butyl ether, isopropanol, methanol and acetone (see Table 25 for the specific amounts) was added respectively to obtain two proportions of suspension sample in each solvent system, the samples were stored at room temperature and at a high temperature (50 °C) and were slurried. The sample bottles (wrapped with a tin foil paper to protect from light) in a room temperature system were placed on Labquaker rotator for rotating 360°; the samples in a high temperature system were placed in a 50 °C thermostatic shaking incubator and were slurried, part of the suspension samples were taken on days 3, 7 and 14, respectively and were centrifuged, solid residue was collected, the solvents were volatilized at room temperature (20 - 25 °C) to dryness, the solid was collected. Upon testing, the obtained solid was crystal form O of a tartrate of a compound of formula I and their XRPD pattern are consistent with Figure 38.

EP 4 385 511 A2

Table 25 Test conditions and results summary of the suspension balance method

	Mass of crystal form O of a tartrate of a compound of formula I (mg)	Solvent	Total volume of solvent (mL)	Crystal form
5	34.5	Tetrahydrofuran	1	Crystal form O
	28.1	Ethanol	1	Crystal form O
10	23.1	Ethyl acetate	1	Crystal form O
	22.4	N-heptane	1	Crystal form O
15	20.0	Toluene	1	Crystal form O
	20.3	Methyl tert-butyl ether	1	Crystal form O
20	30.1	Isopropanol	1	Crystal form O
	50.6	Methanol	0.5	Crystal form O
25	28.9	Acetone	1	Crystal form O

3. Anti-solvent method (Anti-solvent)

30 [0343] 21 proportions of crystal form O of a tartrate of a compound of formula I samples were weighed, a certain volume of good solvents in Table 26 were added sequentially, the systems were subjected ultrasonication to dissolve, the obtained solutions were filtered with a 0.45 μm nylon membrane to a new sample bottle, different anti-solvents were slowly added dropwise to each sample bottle under magnetic stirring, the solvent system wherein solid was precipitated was centrifuged, the solid was collected, the solvent was volatilized at room temperature (20 - 25 °C) to dryness; the solvent system wherein no solids were precipitated was stirred for 48 hours. If no solid was precipitated, the system was stirred with the bottle opened until solid was precipitated. Upon testing, the obtained solid was crystal form O of a tartrate of a compound of formula I and their XRPD patterns are consistent with Figure 38.

Table 26 Test conditions and results summary of the anti-solvent method

Solvent			Product crystal form
Anti-solvent	Good solvent	Volume ratio (mL: mL) (Anti-solvent/good solvent)	Crystal form O
N-heptane	Tetrahydrofuran	10/2.2	
	Ethanol	8.5/1.7	
	Isopropanol	16.8/4.2	
	Methanol	5/0.5	
	Acetone	8/4	

55

(continued)

Solvent			Product crystal form
Anti-solvent	Good solvent	Volume ratio (mL: mL) (Anti-solvent/good solvent)	Crystal form O
Toluene	Tetrahydrofuran	20/2.2	Crystal form O
	Ethanol	8.5/1.8	
	Isopropanol	25.2/4.2	
	Methanol	5/0.5	
	Acetone	24/4	
Methyl tert-butyl ether	Tetrahydrofuran	20/2.2	Crystal form O
	Ethanol	17/1.8	
	Isopropanol	25.2/4.2	
	Methanol	5/0.5	
	Acetone	24/4	
Ethyl acetate	Tetrahydrofuran	20/2.3	Crystal form O
	Ethanol	17/1.7	
	Isopropanol	25.2/4.2	
	Methanol	5/0.5	
	Acetone	24/4	

5. Solution heating - fast cooling method (HFC)

[0344] 5 proportions (about 20 mg/proportion) of crystal form O of a tartrate of a compound of formula I samples were weighed and placed into sample bottles, a suitable amount of tetrahydrofuran, acetone, ethanol, isopropanol and methanol (see Table 27 for the specific amounts) was added respectively, the sample bottles were placed on a magnetic heating stirrer, heated to dissolve in a water bath at about 50°C at 200 rpm. The temperature was maintained for 15 minutes, the solution was filtered with a 0.45 μm membrane when hot and was transferred to new sample bottles, the bottled were immediately transferred to a -20 °C refrigerator overnight, the solvent systems wherein solid was precipitated were centrifuged and the solid was collected, the solvent was volatilized at room temperature (20 - 25 °C) to dryness; the solvent systems wherein no solid was precipitated were placed to a -20 °C refrigerator until a large amount of solids precipitated. In the tetrahydrofuran and acetone systems, solid didn't precipitate all the time. Upon testing, the obtained solid obtained from ethanol, isopropanol and methanol systems was crystal form O of a tartrate of a compound of formula I and their XRPD spectra were consistent with Figure 38.

Table 27 Test conditions and results summary of the solution heating-fast cooling method

Solvent	Volume of solvent (mL)	Crystal form
Tetrahydrofuran	1.5	/
Acetone	2.0	/
Ethanol	0.8	Crystal form O
Isopropanol	2.0	Crystal form O
Methanol	0.3	Crystal form O

6. Solution heating-slow cooling method (HSC)

[0345] 5 proportions (about 20 mg/proportion) of crystal form O of a tartrate of a compound of formula I samples were

EP 4 385 511 A2

weighed and placed in sample bottles, a suitable amount of tetrahydrofuran, acetone, ethanol, isopropanol and methanol (see Table 28 for the specific amounts) was added respectively, the sample bottles were placed on a magnetic heating stirrer, heated to dissolve in a water bath at about 50°C at 200 rpm. The temperature was maintained for 15 minutes, the solution was filtered with a 0.45 µm membrane when hot and transferred to new sample bottles, slowly cooled to room temperature at 6 °C/h overnight, the sample bottles were placed in a refrigerator (2 - 8 °C), the solvent systems wherein solid precipitated were centrifuged and the solid was collected, the solvent was volatilized at room temperature (20 - 25 °C) to dryness; the solvent systems wherein no solid precipitated were placed in a - 20 °C refrigerator until a large amount of solids precipitated. In the tetrahydrofuran, acetone and methanol systems, solids didn't precipitate all the time. Upon testing, the obtained solid obtained from the ethanol and isopropanol systems was crystal form O of a tartrate of a compound of formula I and their XRPD pattern were consistent with Figure 38.

Table 28 Test conditions and results summary of the solution heating-fast cooling method

Solvent	Volume of solvent (ml)	Crystal form
Tetrahydrofuran	1.5	/
Methanol	0.3	/
Acetone	2.0	/
Ethanol	0.8	Crystal form O
Isopropanol	2.0	Crystal form O

[0346] The above test results indicated that after processing crystal form O of a tartrate of a compound of formula I with different methods including suspension balance, heating-fast/slow cooling crystallization, anti-solvent and solution volatilization and crystallization methods, the products were still the single crystal form O. Crystal form O of a tartrate of a compound of formula I is a thermally stable preponderant crystal form.

Test 4: TYK2 biochemical test

A suitable amount of a compound of formula I was weighed for TYK2 biochemical test.

[0347] The test was conducted by Reaction Biology Corp, Malvern, PA (Anastassiadis et al. Nat Biotechnol. 2011; 29(11): 1039-45). The steps are briefly described as follows.

Reagents:

[0348] Basic reaction buffer: 20 mM Hepes (pH 7.5), 10 mM MgCl₂, 1 mM EGTA, 0.02% Brij35, 0.02 mg/ml BSA, 0.1 mM Na₃VO₄, 2 mM DTT and 1% DMSO. The required cofactors were added to each kinase reaction.

Reaction steps:

[0349]

1. Preparing the designated substrate in the newly prepared basic reaction buffer;
2. Transferring the required cofactor to the above matrix solution;
3. Transferring the designated kinase to the substrate solution and mixing well slightly;
4. Transferring a compound of formula I in DMSO to a kinase reaction mixture with Acoustic technique (Echo550; nanoliter range), culturing for 20 minutes at room temperature;
5. Introducing ³³P-ATP (specific activity: 10 µCi/µl) to the reaction mixture to trigger a reaction;
6. Culturing at room temperature and conducting a kinase reaction for 2 hours;
7. Plotting the reaction on P81 ion exchange paper;


8. Testing the kinase activity with a filter binding assay.

[0350] The test results indicated that a compound of formula I was also a potent TYK2 inhibitor and its IC_{50} was less than 10 nM.

[0351] A person skilled in the art can understand and make some modifications or changes to the invention under the instruction of the present description. These modifications and changes should be in the scope specified in the claims of the invention.

[0352] The following clauses form part of the present disclosure.

1. A method for preparing a compound of formula I, wherein the synthesis route of the method is as follows:



the method comprises the following steps:

step 1:

adding ethanol, a compound of formula IV, a compound of formula V and DIPEA to a reaction container, starting stirring;

heating to raise the temperature to 65 - 90 °C, maintaining the temperature and stirring overnight;

terminating the reaction and lowering the temperature of the system to 15 - 30 °C;

adding water to the system dropwise and keeping stirring;

filtrering and washing the filter cake;

drying the filter cake to obtain a compound of formula III;

step 2:

adding tetrahydrofuran, the compound of formula III obtained in step 1 and palladium on carbon to a reaction container;

EP 4 385 511 A2

purging the system with nitrogen and then hydrogen;

maintaining the temperature between 20 - 35 °C and stirring for 16 - 120 hours under 0.1 - 1.0 MPa hydrogen pressure;

after the reaction is completed, filtering the reaction liquid and washing the filter cake;

combining the filtrate and concentrating to obtain a compound of formula II concentrate;

step 3:

adding the compound of formula II concentrate or the compound of formula II obtained in step 2, and trimethyl orthoacetate and tetrahydrofuran to a reaction container; heating the material system in the reaction container until tetrahydrofuran reflux; adding pyridine hydrochloride to the reaction container, reacting the obtained material system for 4 - 20 hours at a temperature between 50 - 90 °C, separating and purifying to obtain a compound of formula I.

2. The method according to clause 1, wherein in the above step 1:

the volume mass ratio (mL/g) of ethanol to the compound of formula IV is between 5:1 and 20:1, preferably 10:1;

the molar ratio of the compound of formula IV, the compound of formula V and DIPEA is 1:1-1.1:2-3, preferably 1:1.01:2.2;

after starting stirring, under nitrogen protection, heating to raise the temperature to 65 - 90°C, preferably 70 - 90 °C, more preferably 70 - 80 °C, maintaining the temperature and stirring for 5 - 16 hours, preferably 10 - 16 hours;

after terminating the reaction, lowering the temperature of the system to 15 - 25 °C;

the volume mass ratio (mL/g) of the water added to the system to the compound of formula IV is between 10:1 and 20:1, preferably 15:1;

after adding water to the system, stirring for 2 - 6 hours, preferably 4 hours, at a temperature between 0 - 30 °C, preferably 5 - 15 °C, more preferably 5 - 10 °C;

the filter cake is washed with ethanol aqueous solution, the volume ratio (mL/mL) of ethanol to water in the ethanol aqueous solution is between 1:1 and 1:2, preferably 1:1.5 - 1:2; the volume mass ratio (mL/g) of the ethanol aqueous solution to the compound of formula IV is between 2:1 and 10:1, preferably 2:1 - 5:1, more preferably 2:1 - 3:1;

drying the filter cake under vacuum or with an air blower at a temperature between 45 - 55°C, preferably 50 °C.

3. The method according to clause 1 or 2, wherein in the above step 2:

the volume mass ratio (mL/g) of tetrahydrofuran to the compound of formula III is between 10:1 and 70:1, preferably 20:1 - 70:1;

the palladium on carbon is 5% Pd/C, 50% water wet, the mass ratio of the palladium on carbon to the compound of formula III is between 0.15:1 and 0.16:1, preferably 0.15:1;

maintaining the temperature between 25 - 35 °C and stirring for 24 - 96 hours under 0.5 - 1.0 MPa hydrogen pressure;

the compound of formula II concentrate obtained by combining the filtrate and concentrating is a compound of formula II in tetrahydrofuran, wherein the volume mass ratio of tetrahydrofuran to the compound of formula II is between 2:1 and 4:1, preferably 2:1 - 3:1;

preferably, exchanging the compound of formula II in tetrahydrofuran with ethanol to obtain a compound of

EP 4 385 511 A2

formula II in ethanol, wherein the volume mass ratio of ethanol to the compound of formula II is between 2:1 and 5:1, preferably 2:1 - 4:1, more preferably 2:1 - 3:1.

4. The method according to any one of clauses 1-3, wherein in the above step 3:

the volume mass ratio (mL:mg) of tetrahydrofuran to the compound of formula II in the compound of formula II concentrate is between 1.5:1 and 5.0:1; or the volume mass ratio (mL:mg) of tetrahydrofuran to the compound of formula II is between 1.5:1 and 5.0:1;

preferably, exchanging the compound of formula II concentrate with toluene, tetrahydrofuran or methyl tertiary-butyl ether for subsequent steps; preferably, the volume mass ratio (mL:mg) of toluene, tetrahydrofuran or methyl tertiary-butyl ether used for exchanging to the compound of formula II concentrate is between 2.0:1 and 4.0:1;

preferably, the molar ratio of the compound of formula II in the compound of formula II concentrate to trimethyl orthoacetate is between 3.0:1 and 3.5:1; or the molar ratio of the compound of formula II to trimethyl orthoacetate is between 3.0:1 and 3.5:1;

preferably, the molar ratio of the compound of formula II in the compound of formula II concentrate to pyridine hydrochloride is between 0.2:1 and 0.3:1; or the molar ratio of the compound of formula II to pyridine hydrochloride is between 0.2:1 and 0.3:1;

preferably, after adding the compound of formula II concentrate or the compound of formula II, and trimethyl orthoacetate and the solvent to the reaction container, under nitrogen protection, heating the material system in the reaction container until the solvent reflux;


after adding pyridine hydrochloride to the reactor, under nitrogen protection, reacting the material system for 4 - 20 hours, preferably 5-15 hours at a temperature between 50 - 90 °C, preferably 65 - 75 °C;

preferably, after the reaction is completed, purifying the product with a solvent selected from the group consisting of water, methanol, ethanol, methyl tertiary-butyl ether and any combinations thereof;

preferably, separating and purifying the compound of formula I obtained by column chromatography, wherein the eluent is a mixed solution of ethyl acetate and n-heptane ($V_{EA}: V_{n\text{-heptane}} = 1:1 - 1:0$, mL/mL);

preferably, the obtained compound of formula I is dried under vacuum or with an air blower between 50 - 55 °C.

5. Crystal form 1 of a compound of formula I



I

wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$ and $21.9^\circ \pm 0.2^\circ$.

6. The crystal form 1 of a compound of formula I according to clause 5, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $9.0^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $21.9^\circ \pm 0.2^\circ$ and $27.3^\circ \pm 0.2^\circ$.

EP 4 385 511 A2

7. The crystal form 1 of a compound of formula I according to clause 5 or 6, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $9.0^\circ \pm 0.2^\circ$, $10.4^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $21.9^\circ \pm 0.2^\circ$, $24.1^\circ \pm 0.2^\circ$ and $27.3^\circ \pm 0.2^\circ$.

8. A preparation method of the crystal form 1 of a compound of formula I according to any of clauses 5-7, comprising: dissolving a compound of formula I crude product with methanol, stirring between 40 - 60 °C for 0.5 - 2 hours, cooling to 5 - 15 °C, stirring for 15 minutes - 1 hour, filtering, washing the filter cake with MTBE, and drying to obtain crystal form 1 of a compound of formula I.


9. The preparation method according to clause 8, comprising:

after dissolving the compound of formula I crude product with methanol, stirring at 50 °C for 1 hour, cooling to 10 °C, stirring for 0.5 hour, filtering, washing the filter cake with MTBE, and drying the filter cake under vacuum at 50 °C for 16 hours to obtain crystal form 1 of a compound of formula I;

preferably, the volume ratio of methanol to MTBE is between 3:1 - 2:1, preferably 8:3;

preferably, after dissolving the compound of formula I crude product with methanol, adding a silicon-based metal eliminator and an activated carbon to the system.

10. Crystal form A of a hydrochloride of a compound of formula I



wherein, the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$ and $20.9^\circ \pm 0.2^\circ$.

11. The crystal form A of a hydrochloride of a compound of formula I according to clause 10, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $20.9^\circ \pm 0.2^\circ$, $23.5^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

12. The crystal form A of a hydrochloride of a compound of formula I according to clause 10 or 11, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.3^\circ \pm 0.2^\circ$, $10.6^\circ \pm 0.2^\circ$, $12.1^\circ \pm 0.2^\circ$, $12.8^\circ \pm 0.2^\circ$, $14.0^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $20.9^\circ \pm 0.2^\circ$, $23.5^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

13. A preparation method of the crystal form A of a hydrochloride of a compound of formula I according to any one of clauses 10 - 12, comprising:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding hydrochloric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form A of a hydrochloride of a compound of formula I.

14. The preparation method according to clause 13, wherein the compound of formula I is subjected to ultrasonication, heating and then is dissolved in acetone;

preferably, the concentration of the compound of formula I in acetone is 10 - 50 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the hydrochloric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

EP 4 385 511 A2


preferably, after adding the hydrochloric acid in acetone, keeping stirring at a room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

5

15. Crystal form B of a hydrochloride of a compound of formula I

10



15

I

20

wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$ and $22.6^\circ \pm 0.2^\circ$.

25

16. The crystal form B of a hydrochloride of a compound of formula I according to clause 15, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $10.2^\circ \pm 0.2^\circ$, $11.5^\circ \pm 0.2^\circ$, $18.0^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $25.9^\circ \pm 0.2^\circ$.

30

17. The crystal form B of a hydrochloride of a compound of formula I according to clause 15 or 16, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $10.2^\circ \pm 0.2^\circ$, $11.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $14.5^\circ \pm 0.2^\circ$, $18.0^\circ \pm 0.2^\circ$, $20.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $25.9^\circ \pm 0.2^\circ$.

35

18. A preparation method of the crystal form B of a hydrochloride of a compound of formula I according to any one of clauses 15 - 17, comprising:

dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding hydrochloric acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form B of a hydrochloride of a compound of formula I.

40

19. The preparation method according to clause 18, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate;

preferably, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the hydrochloric acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL;

45


preferably, after adding the hydrochloric acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

50

20. Crystal form C of a hydrochloride of a compound of formula I

55



I

wherein the X-ray power diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $10.7^\circ \pm 0.2^\circ$, $21.5^\circ \pm 0.2^\circ$ and $24.3^\circ \pm 0.2^\circ$.

21. The crystal form C of a hydrchloride of a compound of formula I according to clause 20, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $5.3^\circ \pm 0.2^\circ$, $10.7^\circ \pm 0.2^\circ$, $21.5^\circ \pm 0.2^\circ$, $24.3^\circ \pm 0.2^\circ$, and $30.4^\circ \pm 0.2^\circ$.

22. A preparation method of the crystal form C of a hydrchloride of a compound of formula I according to clause 20 or 21, comprising:
conducting recrystallization or crystal transformation of crystal form A of a hydrchloride of a compound of formula I with a solvent to obtain crystal form C of a hydrchloride of a compound of formula I, wherein the solvent is selected from of the group consisting of methanol, acetonitrile, n-heptane, methyl ethyl ketone and any combinations thereof.


23. The preparation method according to clause 22, wherein, mixing the solvent with crystal form A of a hydrchloride of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form C of a hydrchloride of a compound of formula I;

preferably, adding the solvent to a container with crystal form A of a hydrchloride of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form C of a hydrchloride of a compound of formula I;

preferably, the duration of the stirring is 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

24. Crystal form D of a sulfate of a compound of formula I



I

wherein the X-ray power diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.0^\circ \pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$ and $25.2^\circ \pm 0.2^\circ$.

25. The crystal form D of a sulfate of a compound of formula I according to clause 24, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.0^\circ \pm 0.2^\circ$, $12.3^\circ \pm 0.2^\circ$, 17.5°

EP 4 385 511 A2

$\pm 0.2^\circ$, $22.8^\circ \pm 0.2^\circ$, and $25.2^\circ \pm 0.2^\circ$.

26. A preparation method of crystal form D of a sulfate of a compound of formula I according to clause 24 or 25, comprising:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding sulfuric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form D of a sulfate of a compound of formula I.

27. The preparation method according to clause 26, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone;


preferably, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the sulfuric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding sulfuric acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

28. Crystal form E of a phosphate of a compound of formula I



I

wherein the X-ray power diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.2^\circ \pm 0.2^\circ$, $15.5^\circ \pm 0.2^\circ$, $17.4^\circ \pm 0.2^\circ$ and $24.6^\circ \pm 0.2^\circ$.

29. The preparation method of crystal form E of a phosphate of a compound of formula I according to clause 28, comprising:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding phosphoric acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form E of a phosphate of a compound of formula I.

30. The preparation method according to clause 29, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone;

preferably, the molar ratio of the compound of formula I to phosphoric acid is between 1:1.0 - 1: 1.5;

preferably, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the phosphoric acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding the phosphoric acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

EP 4 385 511 A2


31. The preparation method according to clause 29 or 30, wherein, conducting recrystallization or crystal transformation of crystal form E of a phosphate of a compound of formula I with a solvent, and the product is still crystal form E of a phosphate of a compound of formula I, wherein the solvent is selected from the group consisting of methanol, acetonitrile, n-heptane, methyl ethyl ketone and any combinations thereof;

preferably, during recrystallization or crystal transformation, mixing the solvent and crystal form E of a phosphate of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying;

preferably, during recrystallization or crystal transformation, the duration of the stirring is between 4 - 48 hours, preferably stirring overnight;

preferably, during recrystallization or crystal transformation, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

32. Crystal form F of a phosphate of a compound of formula I



I

wherein the X-ray power diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$ and $22.6^\circ \pm 0.2^\circ$.

33. The crystal form F of a phosphate of a compound of formula I according to clause 32, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $11.6^\circ \pm 0.2^\circ$, $14.8^\circ \pm 0.2^\circ$, $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $26.6^\circ \pm 0.2^\circ$.

34. The crystal form F of a phosphate of a compound of formula I according to clause 32 or 33, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $11.1^\circ \pm 0.2^\circ$, $11.6^\circ \pm 0.2^\circ$, $14.8^\circ \pm 0.2^\circ$, $16.6^\circ \pm 0.2^\circ$, $17.2^\circ \pm 0.2^\circ$, $21.2^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$ and $26.6^\circ \pm 0.2^\circ$.

35. A preparation method of the crystal form F of a phosphate of a compound of formula I in any one of clauses 32 - 34, comprising:

dissolving crystal form E of a phosphate of a compound of formula I with a first solvent to obtain crystal form E of a phosphate of a compound of formula I in the first solvent, adding an anti-solvent, stirring, collecting the solid, and drying to obtain crystal form F of a phosphate of a compound of formula I;

preferably, the first solvent is a solvent that can dissolve crystal form E of a phosphate of a compound of formula I, preferably methanol; the anti-solvent is a solvent that is difficult to dissolve crystal form E of a phosphate of a compound of formula I, preferably isopropyl acetate;

preferably, the first solvent is added in an amount that can completely dissolve crystal form E of a phosphate of a compound of formula I;

preferably, the anti-solvent is used to dilute crystal form E of a phosphate of a compound of formula I in the first solvent at 5 - 15 folds, preferably 10 folds;

preferably, after dissolving crystal form E of a phosphate of a compound of formula I with


EP 4 385 511 A2

the first solvent, adding a small amount of seed crystals of crystal form F of a phosphate of a compound of formula I until the system is slightly turbid, then adding the anti-solvent;

preferably, after adding the anti-solvent, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying under vacuum between 30 - 60 °C, preferably 50 °C.

36. Crystal form G of a mesylate of a compound of formula I



I

wherein the X-ray powder diffraction pattern shows characteristic peaks at 2theta angles of $8.6^\circ \pm 0.2^\circ$, $19.9^\circ \pm 0.2^\circ$ and $24.9^\circ \pm 0.2^\circ$.

37. The crystal form G of a mesylate of a compound of formula I according to clause 36, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $8.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $18.6^\circ \pm 0.2^\circ$, $19.9^\circ \pm 0.2^\circ$, $24.0^\circ \pm 0.2^\circ$ and $24.9^\circ \pm 0.2^\circ$.

38. A preparation method of the crystal form G of a mesylate of a compound of formula I according to clause 36 or 37, comprising:

dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding methanesulfonic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form G of a mesylate of a compound of formula I.

39. The preparation method according to clause 38, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone;


preferably, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the methanesulfonic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding the methanesulfonic acid in acetone, keeping stirring at room temperature for 4-48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

40. Crystal form H of a hydrobromide of a compound of formula I



I

wherein, the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $20.7^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

41. The crystal form H of a hydrobromide of a compound of formula I according to clause 40, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $17.9^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $20.7^\circ \pm 0.2^\circ$ and $24.0^\circ \pm 0.2^\circ$.

42. The crystal form H of a hydrobromide of a compound of formula I according to clause 40 or 41, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.2^\circ \pm 0.2^\circ$, $11.9^\circ \pm 0.2^\circ$, $17.0^\circ \pm 0.2^\circ$, $17.9^\circ \pm 0.2^\circ$, $18.8^\circ \pm 0.2^\circ$, $20.7^\circ \pm 0.2^\circ$, $24.0^\circ \pm 0.2^\circ$ and $27.5^\circ \pm 0.2^\circ$.

43. The preparation method according to any one of clauses 40 - 42, comprising:
dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding hydrobromic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form H of a hydrobromide of a compound of formula I.

44. The preparation method according to clause 43, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone;

preferably, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the hydrobromic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding the hydrobromic acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying under vacuum overnight at 30 - 60 °C.


45. The preparation method according to clause 43 or 44, wherein, conducting recrystallization or crystal transformation of crystal form H of a hydrobromide of a compound of formula I with a solvent, and the product is still crystal form H of a hydrobromide, wherein the solvent is selecte from the group consisting of acetonitrile, methyl ethyl ketone and any combinations thereof;

preferably, the recrystallization or crystal transformation comprises the following steps:

mixing one or both of acetonitrile and methyl ethyl ketone with crystal form H of a hydrobromide of a compound of formula I, to prepare suspension, stirring at room temperature, centrifuging, collecting the solid and drying;

more preferably, adding one or both of acetonitrile and methyl ethyl ketone to a container with crystal form H of a hydrobromide of a compound of formula I, to prepare suspension, stirring at room temperature, centrifuging, collecting the solid and drying.

46. Crystal form J of a hydrobromide of a compound of formula I



I

15 wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.2^\circ \pm 0.2^\circ$ and $15.0^\circ \pm 0.2^\circ$.

20 47. The preparation method according to clause 46, comprising:
dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding hydrobromic acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring,
20 collecting the solid, and drying to obtain crystal form J of a hydrobromide of a compound of formula I.

25 48. The preparation method according to clause 47, wherein, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate;


25 preferably, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the hydrobromic acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL;

30 preferably, after adding the hydrobromic acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

35 49. Crystal form K of a hydrobromide of a compound of formula I



I

50 wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $17.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$ and $24.2^\circ \pm 0.2^\circ$.

55 50. The crystal form K of a hydrobromide of a compound of formula I according to clause 49, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $17.1^\circ \pm 0.2^\circ$, $20.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $24.2^\circ \pm 0.2^\circ$ and $28.8^\circ \pm 0.2^\circ$.

51. The crystal form K of a hydrobromide of a compound of formula I according to clause 49 or 50, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $9.5^\circ \pm 0.2^\circ$, $17.1^\circ \pm 0.2^\circ$, $20.1^\circ \pm 0.2^\circ$, $22.0^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $24.2^\circ \pm 0.2^\circ$, $27.7^\circ \pm 0.2^\circ$ and $28.8^\circ \pm 0.2^\circ$.

EP 4 385 511 A2

52. A preparation method of the crystal form K of a hydrobromide of a compound of formula I according to any one of clauses 49 - 51, comprising:
conducting recrystallization or crystal transformation of crystal form H of a hydrobromide of a compound of formula I with n-heptane, to obtain crystal form K of a hydrobromide of a compound of formula I.


53. The preparation method according to clause 52, comprising: mixing n-heptane with crystal form H of a hydrobromide of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form K of a hydrobromide of a compound of formula I;

preferably, adding n-heptane to a container with crystal form H of a hydrobromide of a compound of formula I to prepare suspension, stirring at room temperature, collecting the solid, and drying to obtain crystal form K of a hydrobromide of a compound of formula I;

preferably, the duration of the stirring is 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

54. Crystal form L of a fumarate of a compound of formula I



I

Wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

55. The crystal form L of a fumarate of a compound of formula I according to clause 54, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $15.7^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

56. The crystal form L of a fumarate of a compound of formula I according to clause 54 or 55, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.1^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $15.7^\circ \pm 0.2^\circ$, $16.3^\circ \pm 0.2^\circ$, $22.6^\circ \pm 0.2^\circ$, $23.2^\circ \pm 0.2^\circ$, $23.8^\circ \pm 0.2^\circ$ and $26.4^\circ \pm 0.2^\circ$.

57. A preparation method of the crystal form L of a fumarate of a compound of formula I according to any one of clauses 54 - 56, comprising:
dissolving a compound of formula I with a solvent to obtain a compound of formula I in the solvent, and adding fumaric acid in ethanol to the compound of formula I in the solvent under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form L of a fumarate of a compound of formula I.

58. The preparation method according to clause 57, wherein, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in a solvent, wherein the solvent is selected from the group consisting of ethyl acetate, acetone and any combinations thereof;

preferably, the concentration of the compound of formula I in the solvent is 10 - 30 mg/mL, preferably 20 mg/mL;


preferably, the concentration of the fumaric acid in ethanol is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding the fumaric acid in ethanol, keeping stirring at room temperature for 4 - 48 hours,

preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

5 59. Crystal form M of a benzene sulfonate of a compound of formula I



I

20 wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.5^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

25 60. The crystal form M of a benzene sulfonate of a compound of formula I according to clause 59, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $15.2^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $22.4^\circ \pm 0.2^\circ$, $23.0^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

30 61. The crystal form M of a benzene sulfonate of a compound of formula I according to clause 59 or 60, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $7.5^\circ \pm 0.2^\circ$, $12.5^\circ \pm 0.2^\circ$, $14.1^\circ \pm 0.2^\circ$, $15.2^\circ \pm 0.2^\circ$, $18.5^\circ \pm 0.2^\circ$, $22.4^\circ \pm 0.2^\circ$, $23.0^\circ \pm 0.2^\circ$, $24.6^\circ \pm 0.2^\circ$, $25.2^\circ \pm 0.2^\circ$ and $29.8^\circ \pm 0.2^\circ$.

35 62. A preparation method of the crystal form M of a benzene sulfonate of a compound of formula I according to any one of clauses 59 - 61, comprising:
dissolving a compound of formula I with acetone to obtain a compound of formula I in acetone, and adding benzenesulfonic acid in acetone to the compound of formula I in acetone under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form M of a benzene sulfonate of a compound of formula I.

40 63. The preparation method according to clause 62, wherein, the compound of formula I is subjected to ultrasonication, heating and is then dissolved in acetone;

preferably, the concentration of the compound of formula I in acetone is 10 - 30 mg/mL, preferably 20 mg/mL;


preferably, the concentration of the benzenesulfonic acid in acetone is 15 - 35 mg/mL, preferably 25 mg/mL;

45 preferably, after adding the benzenesulfonic acid in acetone, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

50 64. Crystal form N of a citrate of a compound of formula I

55



I

wherein, the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $15.8^\circ \pm 0.2^\circ$, $17.0^\circ \pm 0.2^\circ$ and $21.1^\circ \pm 0.2^\circ$.

65. A preparation method of the crystal form N of a citrate of a compound of formula I according to clause 64, comprising:

dissolving a compound of formula I with ethyl acetate to obtain a compound of formula I in ethyl acetate, and adding citric acid in ethyl acetate to the compound of formula I in ethyl acetate under stirring, keeping stirring, collecting the solid, and drying to obtain crystal form N of a citrate of a compound of formula I.

66. The preparation method according to clause 65, wherein the compound of formula I is subjected to ultrasonication, heating and is then dissolved in ethyl acetate;


preferably, the concentration of the compound of formula I in ethyl acetate is 10 - 30 mg/mL, preferably 20 mg/mL;

preferably, the concentration of the citric acid in ethyl acetate is 15 - 35 mg/mL, preferably 25 mg/mL;

preferably, after adding the citric acid in ethyl acetate, keeping stirring at room temperature for 4 - 48 hours, preferably 24 hours;

preferably, collecting the solid by centrifugation and drying overnight under vacuum at 30 - 60 °C.

67. Crystal form O of a tartrate of a compound of formula I



I

wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $26.1^\circ \pm 0.2^\circ$ and $26.9^\circ \pm 0.2^\circ$.

68. The crystal form O of a tartrate of a compound of formula I according to clause 67, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $12.5^\circ \pm 0.2^\circ$, $15.1^\circ \pm 0.2^\circ$, $26.1^\circ \pm 0.2^\circ$, $26.9^\circ \pm 0.2^\circ$ and $27.5^\circ \pm 0.2^\circ$.

69. The crystal form O of a tartrate of a compound of formula I according to clause 67 or 68, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $6.3^\circ \pm 0.2^\circ$, $11.4^\circ \pm$

EP 4 385 511 A2

0.2°, 12.5° ± 0.2°, 14.1° ± 0.2°, 14.4° ± 0.2°, 15.1° ± 0.2°, 26.1° ± 0.2°, 26.9° ± 0.2° and 27.5° ± 0.2°.

70. A preparation method of crystal form O of a tartrate of a compound of formula I according to any one of clauses 67 - 69, comprising:

5 mixing a compound of formula I with a first solvent, dissolving the compound until the solution is clear and obtaining a compound of formula I in the first solvent; mixing tartaric acid with a second solvent, dissolving the compound until the solution is clear, and obtaining tartaric acid in the second solvent; adding the tartaric acid in the second solvent to the compound of formula I in the first solvent under stirring, conducting controlled-rate cooling under stirring, collecting the solid, drying and obtaining crystal form O of a tartrate of a compound of formula I.

10 71. The preparation method according to clause 70, wherein the first solvent and the second solvent are selected from the group consisting of acetone, ethyl acetate and any combinations thereof;

15 preferably, the molar ratio of the compound of formula I to tartaric acid is 1: (0.5 - 1.5), preferably 1: (0.5 - 0.7), more preferably 1: (0.55 - 0.6);

preferably, in the crystal form O of a tartrate of a compound of formula I, the molar ratio of the compound of formula I to tartaric acid is 2:1;

20 preferably, the concentration of the compound of formula I in acetone is 15 - 70 mg/mL, preferably 40 - 60 mg/mL, more preferably preferably 50 mg/mL;

25 preferably, the concentration of the tartaric acid in acetone is 5 - 35 mg/mL, preferably 10 - 25 mg/mL, more preferably preferably 15 mg/mL;

preferably, mixing the compound of formula I with acetone, raising the temperature to 40 - 60 °C, preferably 50 - 55 °C, to dissolve the compound of formula I until the solution is clear;

30 preferably, mixing the tartaric acid with acetone, raising the temperature to 40 - 60 °C, preferably 50 - 55 °C, to dissolve the tartaric acid until the solution is clear;

preferably, adding the tartaric acid in acetone at 40 - 60 °C, preferably 45 - 55 °C, to the compound of formula I in acetone;

35 preferably, drying the collected solid between 40 - 60 °C under reduced pressure or with an air blower for 5 - 48 hours, preferably 16 - 28 hours.

72. The preparation method according to clause 70 or 71, wherein the controlled-rate cooling is realized by the following steps:

40 1) stirring the system for 0.5 - 3 hours, preferably 1 - 2 hours, at room temperature between 35 - 60 °C, preferably 40 - 60 °C;

45 2) keeping cooling the system to 15 - 35 °C, maintaining the temperature, stirring for 0.5 - 3 hours, preferably 1 - 2 hours;

3) keeping cooling the system to 5 - 15 °C, maintaining the temperature, stirring for 0.5-3 hours, preferably 1 - 2 hours;

50 73. The preparation method according to clause 72, wherein, in step 1), after stirring the system for 0.5 - 3 hours, preferably 1 - 2 hours, at room temperature between 35 - 60 °C, preferably 40 - 60 °C, concentrating the system to one third to two thirds of the original volume, preferably half of the original volume;

55 preferably, in step 2), after keeping cooling the system to 15 - 35 °C, maintaining the temperature and stirring for 0.5 - 3 hours, preferably 1 - 2 hours, concentrating the system to one third to two thirds of the original volume, preferably half of the original volume.

74. The preparation method according to any one of clauses 70 - 73, wherein the purity of the compound of formula I is more than 90%, preferably more than 95%, more preferably preferably more than 99%.


5 75. A pharmaceutical composition comprising the crystal form 1 of a compound of formula I according to any one of clauses 5 - 7, the crystal form A of a hydrochloride of a compound of formula I according to any one of clauses 10 - 12, the crystal form B of a hydrochloride of a compound of formula I according to any one of clauses 15 - 17, the crystal form C of a hydrochloride of a compound of formula I according to clause 20 or 21, the crystal form D of a sulfate of a compound of formula I according to clause 24 or 25, the crystal for E of a phosphate of a compound of formula I according to clause 28, the crystal form F of a phosphate of a compound of formula I according to any one of clauses 32 - 34, the crystal form G of a mesylate of a compound of formula I according to clause 36 or 37, the crystal form H of a hydrobromide of a compound of formula I according to any one of clauses 40 - 42, the crystal form J of a hydrobromide of a compound of formula I according to clause 46, the crystal form K of a hydrobromide of a compound of formula I according to any one of clauses 49 - 51, the crystal form L of a fumarate of a compound of formula I according to any one of clauses 54 - 56, the crystal form M of a benzene sulfonate of a compound of formula I according to any one of clauses 59 - 61, the crystal form N of a citrate crystal of a compound of formula I according to clause 64 and/or the crystal form O of a tartrate of a compound of formula I according to any one of clauses 67 - 69.

15 76. A pharmaceutical formulation comprising the crystal form 1 of a compound of formula I according to any one of clauses 5 - 7, the crystal form A of a hydrochloride of a compound of formula I according to any one of clauses 10 - 12, the crystal form B of a hydrochloride of a compound of formula I according to any one of clauses 15 - 17, the crystal form C of a hydrochloride of a compound of formula I according to clause 20 or 21, the crystal form D of a sulfate of a compound of formula I according to clause 24 or 25, the crystal form E of a phosphate of a compound of formula I according to clause 28, the crystal form F of a phosphate of a compound of formula I according to any one of clauses 32 - 34, the crystal form G of a mesylate of a compound of formula I according to clause 36 or 37, the crystal form H of a hydrobromide of a compound of formula I according to any one of clauses 40 - 42, the crystal form J of a hydrobromide of a compound of formula I according to clause 46, the crystal form K of a hydrobromide of a compound of formula I according to any one of clauses 49 - 51, the crystal form L of a fumarate of a compound of formula I according to any one of clauses 54 - 56, the crystal form M of a benzene sulfonate of a compound of formula I according to any one of clauses 59 - 61, the crystal form N of a citrate of a compound of formula I according to clause 64 and/or the crystal form O of a tartrate of a compound of formula I according to any one of clauses 67 - 69.

20 77. Use of the crystal form 1 of a compound of formula I according to any one of clauses 5 - 7, the crystal form A of a hydrochloride of a compound of formula I according to any one of clauses 10 - 12, the crystal form B of a hydrochloride of a compound of formula I according to any one of clauses 15 - 17, the crystal form C of a hydrochloride of a compound of formula I according to clause 20 or 21, the crystal form D of a sulfate of a compound of formula I according to clause 24 or 25, the crystal form E of a phosphate of a compound of formula I according to clause 28, the crystal form F of a phosphate of a compound of formula I according to any one of clauses 32 - 34, the crystal form G of a mesylate of a compound of formula I according to clause 36 or 37, the crystal form H of a hydrobromide of a compound of formula I according to any one of clauses 40 - 42, the crystal form J of a hydrobromide of a compound of formula I according to clause 46, the crystal form K of a hydrobromide of a compound of formula I according to any one of clauses 49 - 51, the crystal form L of a fumarate of a compound of formula I according to any one of clauses 54 - 56, the crystal form M of a benzene sulfonate of a compound of formula I according to any one of clauses 59 - 61, the crystal form N of a citrate of a compound of formula I according to clause 64 and/or the crystal form O of a tartrate of a compound of formula I according to any one of clauses 67 - 69 in preparing medicaments for treating JAK1/TYK2-related diseases or conditions; preferably, the diseases or conditions are autoimmune diseases or disorders, more preferably preferably rheumatoid arthritis or inflammatory diseases or disorders, and cancers or tumor proliferative diseases or disorders.

Claims

50 1. Crystal form 1 of a compound of formula I



I

wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$ and $21.9^\circ \pm 0.2^\circ$.

2. The crystal form 1 of a compound of formula I according to claim 1, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $9.0^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $21.9^\circ \pm 0.2^\circ$ and $27.3^\circ \pm 0.2^\circ$.
3. The crystal form 1 of a compound of formula I according to claim 1 or 2, wherein the X-ray powder diffraction pattern of the crystal form shows characteristic peaks at 2theta angles of $9.0^\circ \pm 0.2^\circ$, $10.4^\circ \pm 0.2^\circ$, $13.4^\circ \pm 0.2^\circ$, $17.6^\circ \pm 0.2^\circ$, $18.1^\circ \pm 0.2^\circ$, $18.7^\circ \pm 0.2^\circ$, $21.9^\circ \pm 0.2^\circ$, $24.1^\circ \pm 0.2^\circ$ and $27.3^\circ \pm 0.2^\circ$.
4. A preparation method of the crystal form 1 of a compound of formula I according to any of claims 1 to 3, comprising dissolving a compound of formula I crude product with methanol, stirring between 40 - 60 °C for 0.5 - 2 hours, cooling to 5 - 15 °C, stirring for 15 minutes - 1 hour, filtering, washing the filter cake with MTBE, and drying to obtain crystal form 1 of a compound of formula I.
5. The preparation method according to claim 4 comprising: after dissolving the compound of formula I crude product with methanol, stirring at 50 °C for 1 hour, cooling to 10 °C, stirring for 0.5 hour, filtering, washing the filter cake with MTBE, and drying the filter cake under vacuum at 50 °C for 16 hours to obtain crystal form 1 of a compound of formula I.
6. The preparation method according to claim 5 wherein the volume ratio of methanol to MTBE is between 3:1 - 2:1, preferably 8:3.
7. The preparation method according to claim 6 wherein, after dissolving the compound of formula I crude product with methanol, adding a silicon-based metal eliminator and an activated carbon to the system.
8. A pharmaceutical composition comprising the crystal form 1 of a compound of formula I according to any one of claims 1 to 3.
9. A pharmaceutical formulation comprising the crystal form 1 of a compound of formula I according to any one of claims 1 to 3.
10. The crystal form 1 of a compound of formula I according to any one of claims 1 to 3 for treating JAK1/TYK2-related diseases or conditions.
11. The crystal form 1 of a compound of formula I for use according to claim 10, wherein the diseases or conditions are autoimmune diseases or disorders.
12. The crystal form 1 of a compound of formula I for use according to claim 11 wherein the diseases or conditions are rheumatoid arthritis or inflammatory diseases or disorders, and cancers or tumor proliferative diseases or disorders.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29




Figure 30




Figure 31




Figure 32




Figure 33




Figure 34




Figure 35




Figure 36




Figure 37




Figure 38




Figure 39




Figure 40




Figure 41




Figure 42




Figure 43




Figure 44




Figure 45




Figure 46




Figure 47




Figure 48




Figure 49




Figure 50

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2018067422 A1 [0005] [0006]

Non-patent literature cited in the description

- **MURRAY**. *J. Immunol.*, 2007, vol. 178 (5), 2623-2629 [0003]
- **GHOESCHI et al.** *Immunol Rev.*, 2009, vol. 228 (1), 273-287 [0003]
- **SCHINDLER et al.** *J. Biol. Chem.*, 2007, vol. 282 (28), 20059-20063 [0003]
- **GHOESCHI et al.** *Immunol. Rev.*, 2009, vol. 228 (1), 273-287 [0004]
- **GUSCHIN et al.** *EMBO J.*, 1995, vol. 14 (7), 1421-1429 [0004]
- **MAINI et al.** *Arthritis Rheum.*, 2006, vol. 54 (9), 2817-2829 [0004]
- **ANASTASSIADIS et al.** *Nat Biotechnol.*, 2011, vol. 29 (11), 1039-45 [0347]