

(12) United States Plant Patent

Dümmen

(10) **Patent No.:**

US PP20.852 P2

(45) **Date of Patent:**

Mar. 16, 2010

(54) CALIBRACHOA PLANT NAMED 'DUEALTIOR'

(50) Latin Name: Calibrachoa sp. Varietal Denomination: **Duealtior**

Inventor: Tobias Dümmen, Rheinberg (DE)

Assignee: Capital Green Investments Ltd., Grand

Cayman (KY)

Subject to any disclaimer, the term of this (*) Notice:

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

Appl. No.: 12/313,688

(22) Filed: Nov. 21, 2008

(51) Int. Cl.

A01H 5/00 (2006.01)

(52) U.S. Cl. Plt./413

(58) Field of Classification Search Plt./413 See application file for complete search history.

(56)References Cited

OTHER PUBLICATIONS

Upov-rom Plant Variety Database, GTI Jouve Retrieval Software, Citation for Calibrachoa 'Duealtior', one page.*

* cited by examiner

Primary Examiner—June Hwu (74) Attorney, Agent, or Firm—C. A. Whealy

ABSTRACT

A new and distinct cultivar of Calibrachoa plant named 'Duealtior', characterized by its upright and outwardly spreading to trailing and decumbent plant habit; numerous large orange-colored flowers with red-colored centers; and good garden performance.

1 Drawing Sheet

1

Botanical designation: Calibrachoa sp. Cultivar denomination: 'Duealtior'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of Calibrachoa plant, botanically known as Calibrachoa sp. and hereinafter referred to by the name 'Duealtior'.

The new Calibrachoa plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Ger- 10 many. The objective of the breeding program is to create new Calibrachoa cultivars with large attractive flowers.

The new Calibrachoa originated from a cross-pollination made by the Inventor in August, 2005 in Rheinberg, Germany of a proprietary selection of Calibrachoa sp. identified as 15 code number F-13-0104, not patented, as the female, or seed, parent with a proprietary selection of Calibrachoa sp. identified as code number A-04-0015-006, not patented, as the male, or pollen, parent. The new Calibrachoa was discovered and selected by the Inventor as a single flowering plant from 20 within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2007.

Asexual reproduction of the new Calibrachoa plant by vegetative terminal cuttings in a controlled greenhouse envi-25 ronment in Rheinberg, Germany since May, 2007, has shown that the unique features of this new Calibrachoa plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new Calibrachoa have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment and cultural practices such as temperature and light intensity with- 35 out, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Duealtior'.

2

These characteristics in combination distinguish 'Duealtior' as a new and distinct cultivar of Calibrachoa:

- 1. Upright and outwardly spreading to trailing and decumbent plant habit.
- 2. Numerous large orange-colored flowers with red-colored centers.
- 3. Good garden performance.

30

The new Calibrachoa can be compared to plants of the female parent selection. Plants of the new Calibrachoa differ primarily from plants of the female parent selection in the following characteristics:

- 1. Plants of the new Calibrachoa are more compact than plants of the female parent selection.
- 2. Plants of the new Calibrachoa have more intense orange-colored flowers than plants of the female parent

The new Calibrachoa can be compared to plants of the male parent selection. Plants of the new Calibrachoa differ primarily from plants of the male parent selection in flower color as plants of the male parent selection have red-colored

Plants of the new Calibrachoa can be compared to plants of Calibrachoa 'KLECA06124', disclosed in U.S. Plant Pat. No. 19,476. Plants of the new Calibrachoa differ from plants of 'KLECA06124' in the following characteristics:

- 1. Plants of the new Calibrachoa have longer internodes than plants of 'KLECA06124'.
 - 2. Plants of the new Calibrachoa have larger flowers than plants of 'KLECA06124'.
- 3. Plants of the new Calibrachoa have darker orange-colored flowers than plants of 'KLECA06124'.
- 4. Plants of the new Calibrachoa have larger sepals than plants of 'KLECA06124'.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new Calibrachoa, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly 20

45

3 4

from the color values cited in the detailed botanical description which accurately describe the colors of the new Calibra-

The photograph is a top perspective view of a typical flowering lateral branch of 'Duealtior'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in Rheinberg, Germany, under commercial practice during the spring in a glass-covered greenhouse with day and night temperatures averaging 18° C. and light levels averaging 4,500 lux. Rooted young plants were pinched one time about three weeks after planting had been growing for 16 weeks when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are

Botanical classification: Calibrachoa sp. 'Duealtior'. Parentage:

Female, or seed, parent.—Proprietary selection of Calibrachoa sp. identified as code number F-13-0104, not

Male, or pollen, parent.—Proprietary selection of Cali-25 brachoa sp. identified as code number A-04-0015-006, not patented.

Propagation:

Type.—By vegetative cuttings.

Time to initiate roots, summer.—About seven days at 20°

Time to initiate roots, winter.—About ten days at 20° C. Time to produce a rooted young plant, summer.—About three weeks at temperatures of 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures of 20° C.

Root description.—Fine, fibrous; white in color. Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Upright and outwardly spreading to trailing and decumbent plant habit. 40 Freely branching habit with about 10 to 15 lateral branches; pinching enhances branching; vigorous growth habit.

Plant height.—About 11.5 cm.

Plant diameter.—About 22 cm.

Lateral branch description:

Length.—About 11 cm.

Diameter.—About 1.5 mm. Internode length.—About 1.4 cm.

Strength.—Strong.

Aspect.—Initially upright to outwardly spreading to 50 trailing and decumbent.

Texture.—Pubescent.

Color.—Close to 144B.

Foliage description:

Arrangement.—Before flowering, alternate, simple; 55 after flowering, opposite, simple.

Length.—About 3.6 cm.

Width.—About 1.5 cm.

Shape.—Oblanceolate.

Apex.—Rounded.

Base.—Cuneate. Margin.—Entire.

Texture, upper and lower surfaces.—Pubescent.

Venation pattern.—Pinnate; arcuate.

surface: Close to 137C; venation, close to 144C.

Developing and fully expanded leaves, lower surface: Close to 138B to 138C; venation, close to 144C. Petiole: Length: About 2 mm. Diameter: About 1 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: Close to 144B.

Flower description:

Flower arrangement and habit.—Single salverform flowers arising from leaf axils. Freely flowering habit with usually about 40 open flowers and flower buds per plant. Flowers face upright or outwardly.

Fragrance.—None detected.

Natural flowering season.—Plants of the new Calibrachoa initiate and develop flowers about four to ten weeks after planting. Plants flower continuously from the spring until the autumn in Germany.

Flower longevity.—Individual flowers last about seven to ten days on the plant; flowers not persistent.

Flower diameter.—About 3.8 cm.

Flower length (height).—About 2.4 cm.

Flower throat diameter.—About 9 mm.

Flower tube length.—About 1.6 cm.

Flower tube diameter, base.—About 2 mm.

Flower bud.—Shape: Elongated oblong. Length: About 2.2 cm. Diameter: About 5 mm. Color: Close to 145C tinted with close to 22A

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal length from throat: About 1.5 cm. Petal lobe width: About 1.9 cm. Petal shape: Roughly spatulate. Petal apex: Shallowly cordate. Petal margin: Entire. Petal texture, upper and lower surfaces: Smooth, glabrous. Throat texture: Smooth, glabrous. Tube texture: Smooth, glabrous. Color: Petal, when opening and fully opened, upper surface: Close to 26A to 26B; towards the base, close to 44A; venation, close to 59A to 59B; color does not fade with development. Petal, when opening and fully opened, lower surface: Close to 26B; venation, close to 59A. Throat: Close to 7A; venation, close to 59A. Tube: Close to 151C; venation, close to 59A.

Calvx.—Arrangement: One star-shaped calvx tube with five sepals fused at the base. Sepal length: About 1.3 cm. Sepal width: About 2.6 mm. Sepal shape: Lanceolate. Sepal apex: Acute. Sepal margin: Entire. Sepal texture, upper and lower surfaces: Pubescent. Color, upper surface: Close to 137C. Color, lower surface: Close to 138A.

Peduncles.—Length: About 2.1 cm. Diameter: About 1 mm. Angle: About 45° from stem axis. Strength: Moderately strong. Texture: Pubescent. Color: Close to 144B.

Reproductive organs.—Stamens: Quantity/arrangement: Five per flower. Filament length: About 8 mm. Filament color: Close to 145C. Anther shape: Ellipsoidal. Anther length: About 1.5 mm. Anther color: Close to 7B. Pollen amount: Scarce. Pollen color: Close to 13C. Pistils: Quantity: One per flower. Pistil length: About 1.2 cm. Style length: About 9 mm. Style color: Close to 144D. Stigma shape: Ellipsoidal. Stigma color: Close to 144B. Ovary color: Close to 144D. Seed/fruit: Seed and fruit development have not been observed on plants of the new Calibrachoa.

Color.—Developing and fully expanded leaves, upper 65 Garden performance: Plants of the new Calibrachoa have been observed to have good garden performance and tol 5

6

erate wind, rain and temperatures ranging from about 2° C. to about 38° C.

Pathogen/pest resistance: Plants of the new *Calibrachoa* have not been observed to be resistant to pathogens and pests common to Calibrachoa.

It is claimed:
1. A new and distinct *Calibrachoa* plant named 'Duealtior' as illustrated and described.

