



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A01N 57/00, 57/10, 57/18, 57/26, 59/26, C05B 7/00, C05G 3/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 98/38863</p> <p>(43) International Publication Date: 11 September 1998 (11.09.98)</p>
<p>(21) International Application Number: PCT/US98/03459</p> <p>(22) International Filing Date: 24 February 1998 (24.02.98)</p> <p>(30) Priority Data: 08/812,865 6 March 1997 (06.03.97) US</p> <p>(71)(72) Applicant and Inventor: TAYLOR, John [US/US]; 1420 Lemon Street, Deland, FL 32720 (US).</p> <p>(74) Agent: SCHOONOVER, Donald, R.; Lathrop & Gage, LC, Suite 2500, 2345 Grand Boulevard, Kansas City, MO 64108 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: PLANT FERTILIZER COMPOSITIONS CONTAINING PHOSPHONATE AND PHOSPHATE SALTS, AND DERIVATIVES THEREOF</p>		
<p>(57) Abstract</p> <p>A fertilizer composition for plants containing phosphonate and phosphate salts, and derivatives thereof is disclosed. The composition provides a single product which may be employed to stimulate the growth response in plants.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MX	Mexico	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Plant Fertilizer Compositions Containing
Phosphonate and Phosphate Salts, and Derivatives Thereof

5 Cross-Reference to Related Application

The present invention is a continuation-in-part of co-pending Serial Number 08/705,594 filed on August 30, 1996.

Background of the Invention


10 The present invention is broadly concerned with fungicidal compositions, and methods of use, which provide improved efficacy in controlling parasitic fungi in plants. More particularly, the compositions and methods of use of the invention include fungicidally effective amounts of both phosphate, preferably in the form of either mono, di, tri or dipotassium phosphate (KH_2PO_4 , K_2HPO_4) and phosphonate, preferably in the form of either mono or dipotassium phosphonate (KH_2PO_3 , K_2HPO_3), in aqueous
15 solution.

 According to another aspect of the present invention, the compositions and methods of use of the invention include growth response effective amounts of both phosphate, preferably in the form of either mono or dipotassium phosphate (KH_2PO_4 , K_2HPO_4) and phosphonate, preferably in the form of either mono or dipotassium
20 phosphonate (KH_2PO_3 , K_2HPO_3), in aqueous solution.

 Phosphorus is an essential major element in plant nutrition because it governs the energy producing reactions, including those that are oxidative, and photophosphorylative and the production of adenosine diphosphate (ADP) and adenosine triphosphate (ATP).

5 Energy-rich phosphate bonds of ADP and ATP provide the energy for many of the physiological reactions that occur in plants.

The element phosphorous appears in two general forms that concern the present invention — phosphonate and phosphate. The term “phosphonate,” sometimes also referred to as “phosphite,” means the salts (organic or inorganic) of either phosphonic acid or phosphorous acid. Phosphonic and phosphorous acids have the formula H_3PO_3 and a molecular weight of 82.00. Their structures from the International Union of Pure and Applied Chemistry are shown below:



20 The term “phosphate” means the salts (organic or inorganic) of phosphoric acid having the formula H_3PO_4 , molecular weight of 98, and has the following structure:



30 In the past, various phosphonate compounds have been proposed as useful in fungicidal and fertilizer compositions for application to plants. See, e.g., U.S. Patent

5 Nos. 4,075,324 and 4,119,724 to Thizy, describing phosphorous acid, its inorganic and
organic salts, as a plant fungicide; U.S. Patent No. 4,139,616 to Dueret, describing
fungicidal compositions based on phosphorous acid esters and salts thereof; U.S. Patent
No. 4,542,023 to Lacroix et al., describing organophosphorus derivatives as possessing
systemic and contact fungistatic and fungicidal activity; U.S. Patent Nos. 4,698,334 and
10 4,806,445 and 5,169,646 to Horriere et al., describing fungicidal compositions based on
alkyl phosphonates; U.S. Patent Nos. 4,935,410 and 5,070,083 to Barlet, describing
fungicidal aluminum tris-alkyl-phosphonate compositions; and U.S. Patent No. 5,514,200
to Lovatt, describing formulations of phosphorous-containing acid fertilizer for plants.
(The teachings of the proceeding U.S. Patents are hereby incorporated by reference.) The
15 above references disclose phosphonate compositions have been found to be effective for
protecting plants, and particularly grape vines, citrus and fruit trees and tropical plants,
against fungal attack.

Once assimilated, phosphonates have been shown to enhance the phytoimmune
system. The phosphonate induced stimulation of the phytoimmune system is triggered
20 by the induction of ethylene production, followed by a rapid accumulation of
phytoalexins at the site of infection. Phosphonates have also been shown to have
detrimental effect on the growth of Phycomycetes. See, Pegg, K.G. and deBoer, R.F.,
"Proceedings of the Phosponic (Phosphorous) Acid Work Shop, "*Australiasian Plant
Pathology*, Vol. 19 (4) 1990.

25 In accordance with this development of the present invention, however, it has
recently been discovered that phosphonates exacerbate the non-target Ascomycete fungi,
and other fungi producing an eiphytotic outbreak of much greater magnitude than the

5 infections occurring without phosphonate treatment. This phenomenon is known as pathological acerbation.

In the past, phosphates were not viewed as a solution to pathological acerbation of Ascomycete fungal infections. This is because phosphates are viewed primarily as a fertilizer with only limited, or even detrimental, fungicidal properties. For example, U.S. Patent 5,514,200 teaches that phosphate fertilizers inhibit beneficial symbiosis between 10 plant roots and mycorrhizal fungi, and further promote bacterial and fungal growth in the rhizosphere, including the growth of pathogenic fungi and other small soil-borne organisms. (Col. 2, lines 18-28). Phosphates have also been considered to be a competitive inhibitor for phosphonate assimilation, thus inhibiting the ability of 15 phosphonates to protect against fungus attack. See, Pegg, K.G. and deBoer, R.F., "Proceedings of the Phosphonic (Phosphorous) Acid Work Shop, *Australasian Plant Pathology*, Vol. 19 (4), pp. 117 and 144, 1990. Yet further, phosphonates and phosphates were believed to be "biological strangers," with the presence of phosphonates or its esters, exerting little or no influence on enzyme reactions involving phosphates. 20 Robertson, H.E. and Boyer, P.D., "The Biological Inactivity of Glucose 6 — phosphonate, Inorganic Phosphites and Other Phosphites," *Archives of Biochemistry and Biophysics*, 62 pp. 380-395 (1956).

Accordingly, the requirements for a successful phosphonate-based fungicide depend on the elimination of the phosphonate-induced pathological acerbation of 25 Ascomycete fungal infections.

5 Moreover, the prior art teaching that phosphates and phosphonates are "biological strangers" is relevant to the lack of teaching of use of a composition of both for any agricultural use.

Summary of the Invention

10 The present invention addresses the problems outlined above, and provides an improved anti-fungicidal composition for plants that contains, as active ingredients, fungicidally effective amounts of both phosphonates and phosphates. According to the present invention, it has been discovered that the application to a plant of the inventive phosphonate/phosphate composition substantially eliminates pathological acerbatation of Ascomycete fungi, while at the same time, provides a means to control Phycomycetes and Ascomycetes and other fungi with a single product.

15 Thus, an object of present invention is to provide a fungicidal composition for protection for plants against fungal infection, especially Phycomycetes and Ascomycetes.

20 Another object of the invention is to provide such anti-fungal protection with a single product that upon application does not cause pathological acerbatation of Ascomycetes infections.

 A further object of the invention is to provide a method of treating plants and to provide anti-fungal protection in plants against attack by Phycomycetes, Ascomycete and other fungi and bacteria.

25 A yet further object of the invention is to provide an anti-fungal composition for treating plants that is environmentally safe, inexpensive to use and has low mammalian toxicity.

5 These and other objects of the invention are attained by the invention disclosed
below. According to the invention, anti-fungicidal compositions for the protection of
plants, especially citrus and fruit trees and vines against fungus attack, preferably contain
as an active material a fungicidally effective amount of at least a first salt formula
selected from the group consisting of KH_2PO_3 , K_2HPO_3 , and K_3PO_3 , and at least a
10 second salt selected from the group consisting of KH_2PO_4 , K_2HPO_4 and K_3PO_4 , in
a mixture with an agriculturally acceptable carrier.

The composition preferably comprises an aqueous solution wherein each salt is
present in solution from about 20 millimole to about 5% vol./vol..

15 According to another aspect of the invention, the amount of the first salt is one
part by weight and the amount of the second salt is between 0.001 and 1,000 parts by
weight.

Phosphonate salts useful in the practice of the invention also include those organic
and inorganic salts taught by U.S. Patent Nos. 4,075,324 and 4,119,724 to Thizy et al.,
(see, e.g., col. 1, ln. 51-69 through col. 2, ln. 1-4).

20 The present invention, in another aspect, also has been found to be effective as
a growth stimulator or fertilizer for plants, and addresses the problems in the art with
regard to finding effective fertilizers. Thus, the present invention provides a means for
applying a single product to plants which is both an effective fungicide and an effective
fertilizer.

25 It is therefore an object of this aspect of the invention to present a composition
and method of use which functions as a fertilizer for plants.

5 A further object of this aspect of the invention is to function as a fertilizer which provides substantial growth response results.

Another object of this aspect of the invention is to function as a fertilizer which is ecologically and human compatible, is economical, and is efficient.

10 These and other objects, features and advantages of the invention will become more apparent with reference to the accompanying specification and claims.

A Detailed Description of the Preferred Embodiments.

The following examples set for preferred concentrations and techniques for formulation thereof, as well as methods of application and use and test results demonstrating the efficacy of the inventive concentration in protecting plants against attack by Ascomycete and Phycomycete fungi. It is to be understood, however, that these
15 Examples are presented by way of illustration only and nothing therein shall be taken as a limitation upon the overall scope of the invention. The specific components tested in the Examples were prepared and applied as follows.

Method of Preparation

20 A. Potassium Phosphonate (KH_2PO_3) Aqueous Solution. H_3PO_3 is produced by the hydrolysis of phosphorus trichloride according to the reaction: $\text{PCl}_3 + 3\text{H}_2\text{O} > \text{H}_3\text{PO}_3 + 3\text{HCl}$. The HCl is removed by stripping under reduced pressure and the phosphonic acid (H_3PO_3) is sold as a 70% acid solution.

The phosphonic acid is then neutralized in aqueous solution by potassium
25 hydroxide according to the reaction: $\text{H}_3\text{PO}_3 + \text{KOH} > \text{KH}_2\text{PO}_3 + \text{H}_2\text{O}$ to about pH 6.5 and to produce a 0-22-20 liquid weighing 11.15 lbs./gal. This solution is commercially

5 available and is sold under the trademark "Phos-Might" by Foliar Nutrients, Inc., Cairo, GA 31728.

B. Potassium Phosphate (K₂HPO₄) Aqueous Solution. Mono potassium phosphate (0-51.5-34) is reacted with 45% potassium hydroxide in aqueous solution to produce dipotassium phosphate by the following reaction: This solution is commercially
10 available and is sold under trademark "K-Phos" by Foliar Nutrients, Inc., Cairo, GA 31724

$\text{KH}_2\text{PO}_4 + \text{KOH} > \text{K}_2\text{HPO}_4 + \text{H}_2\text{O}$ with a product density of 1.394 at 20 C and a solution pH of 7.6 producing a 0-18-20 analysis.

C. KH₂PO₃/K₂HPO₄ Solution. Varying amounts of each compound
15 (K₂HPO₃;KH₂PO₃; K₂HPO₄; or KH₂PO₄) in aqueous solution are combined at rates ranging from 20 millimole to 5% vol./vol., depending on crop host and the pathogen complex and level of infection.

Method of Application

The preferable method of application is foliar, either by ground or aerial
20 equipment, but is not limited to that method alone. Injection or soil applications, for example, could also be efficacious depending on specific crops and pathogens.

The inventive compositions have utility on fruit crops, agronomic crops, ornamentals, trees, grasses, vegetables, grains, and floricultural crops, as well as, some aquatic crops including water cress.

25 The fungicidal properties of the compounds according to the invention are various, but are particularly interesting in the case described in the following examples:

5 Examples 1-5 exemplify a portion of these applications. In Examples 1-4, fungicidal treatments were applied to ornamentals, citrus and vegetables. In each of Examples 1-4, treatments were applied as a one gallon solution by a back pack sprayer, maintained at about 60 psi, in sufficient quantities of water to achieve thorough coverage.

10 All treatments were applied to the appropriate number of experimental units assigned in a randomized complete block (RCB) design replicated four times. Example #5 was a pre-harvest fungicidal spray evaluation on "Nova" tangelos.

 As used in the examples, "percent infection" means percent of leaves that exhibit fungus lesions. "Phytotoxicity" means number crop injury to leaves that fell off the plant following the application of the solution.

15 The Ascomycete fungi were the subjects of testing in Examples 1 through 5 and are all members of the Ascomycete family.

EXAMPLE NO. 1

Dogwood/Powdery Mildew Fungicide Trial Trail Ridge Nursery, Keystone Hts., Florida

20 The compounds according to the invention are studied for the effect on the powdery mildew (*Oidium Spp.*), which is responsible for foliar infections in dogwood.

 Potassium phosphonate and potassium phosphate solutions were applied in compositions as set forth in Table 1 below. An average of 5 six-inch pots per experimental unit with 4 repetitions in a randomized complete block ("RCB") design
25 were considered. The solutions were applied to Dogwood (*Cornis Florida Var. "Weaver"*). The application of the compositions were made in April 25, 1996, May 8,

5 1996, and May 23, 1996. The effects of the various compositions were rated on May 30,
1996.

10 Table 1 shows that when a potassium phosphonate solution (containing no appreciable amount of phosphate) is applied to dogwood, pathological acerbation of the Ascomycete fungus occurs in 100 percent of the dogwood leaves, in contrast to the control, which exhibited only 30 percent infection. The inventive composition that combine potassium phosphate and potassium phosphonate showed a complete elimination of the pathological acerbation phenomenon, and in fact, dropped the amount of infection by about 20 percent.

TABLE 1

15	<u>TREATMENT</u>	<u>RATE/100 gal.</u>	<u>*%INFECTION</u>	<u>PHYTOTOXICITY</u>
	1) POTASSIUM PHOS- PHONATE SOLUTION	1%	100	0
	2) POTASSIUM PHOS- PHATE SOLUTION	1%	0	0
20	3) POTASSIUM PHOS- PHONATE SOLUTION	1%	10	0
	+ POTASSIUM PHOS- PHATE SOLUTION	+ 1%		0
25	4) POTASSIUM PHOS- PHATE SOLUTION	2%	0	0
	5) CONTROL	-	30	0

5

EXAMPLE NO. 2

Shumard Oak/Powdery Mildew Fungicide Trial
Trail Ridge Nursery, Keystone Hts., Florida

10

In Example No. 2, the inventive compositions were tested for effectiveness against the powdery mildew (*Phyllactinia corylea*) that infects shumard oak (*Quercus shumardii*). In Example No. 2, an average of ten 3-gallon plots per experimental unit with 4 repetitions in a randomized complete block design were examined.

15

When only potassium phosphonate solution was applied, the percentage of infection increased to 40 percent, compared to a 20 percent infection in the control, indicating a pathological acerbation of the Ascomycete fungi. When the inventive composition including potassium phosphonate and potassium phosphate was applied, the infection was completely eliminated.

TABLE 2

<u>TREATMENT</u>	<u>RATE/100 gal.</u>	<u>*%INFECTION</u>	<u>PHYTOTOXICITY</u>
20 1) POTASSIUM PHOS- PHONATE SOLUTION	1%	40	0
2) POTASSIUM PHOS- PHATE SOLUTION	1%	0	0
25 3) POTASSIUM PHOS- PHONATE SOLUTION +	1% +	0	0
POTASSIUM PHOS- PHATE SOLUTION	1%		
4) POTASSIUM PHOS- PHATE SOLUTION	2%	0	0
30 5) CONTROL	-	20	0

5

EXAMPLE NO. 3

Citrus Alternaria Fungicide Trial
Kerr Center, Vero Beach, Florida

In Example No. 3, the inventive composition was tested for effectiveness in citrus trees (*Alternaria citri*) that were infected with the fungi.

10

As used in Table No. 3 below, “% Y.L.D.” means percent young leaf drop, and “% Y.F.D.” means percent young fruit drop. %Y.L.D. and %Y.F.D. were determined by examining 4 terminals/tree with 4 repetitions in a randomized complete block design. The date on which the compositions were applied were March 21, 1996. The compositions were rated on April 2, 1996. When only potassium phosphonate solutions were applied, %Y.L.D. and %Y.F.D. increased from 6.2 and 1.5 to 23.9 and 37.9, respectively, indicating pathological acerbation of the Ascomycete fungi. Application of the inventive composition not only eliminated the acerbation, but showed an improvement in Y.L.D. and Y.F.D.

15

TABLE 3

20

<u>TREATMENT</u>	<u>RATE/100 gal.</u>	<u>%Y.L.D.</u>	<u>% Y . F . D .</u>	
<u>PHYTOTOXICITY</u>				
1) POTASSIUM PHOSPHATE SOLUTION	0.5%	5.2	3.8	0
2) POTASSIUM PHOSPHATE SOLUTION	1 %	2.5	2.5	0
3) POTASSIUM PHOSPHATE SOLUTION	2 %	1.5	1.0	0
4) POTASSIUM PHOSPHONATE SOLUTION	1 %	23.9	37.5	0

25

5	5) POTASSIUM PHOS- PHATE SOLUTION + POTASSIUM PHOS- PHONATE SOLUTION	0.5% + 0.5%	2.2	3.7	0
10	6) CONTROL	-	6.2	1.5	0

EXAMPLE NO. 4

Carrot/*Alternaria* Fungicide Trial

In Example No. 4, carrot plots were inoculated with *Alternaria dauci* fungi. The plot sizes were single rows X 25 feet X 4 repetitions in a RCB design. The dates that compositions were applied were February 2, 9, 15, 22, and March 8, 14, 22, and 28, 1996. The fungi infections were rated on April 5 and April 23, 1996. The second rating was 25 days after the last fungicide application. The test took place at Sanford, Florida.

Application of the inventive solutions including potassium phosphonate and potassium phosphate improved the extent of infection from that otherwise occurring in the control carrots. Although the solution making use of only potassium phosphonate caused a slight improvement from the infection occurring in the control, it was substantially less than that associated with the inventive composition.

Also shown in Example No. 4 is a comparison of the effectiveness of the inventive composition to other fungicides, including Cu-Alexin and Fe-Alexin. This test demonstrates that the inventive composition is as effective as other well known fungicides, but without being as environmentally toxic.

5

TABLE 4

	<u>TREATMENT</u>	<u>RATE/100 GAL.</u>	<u>AVG%INFECTION</u>	
			<u>4/5/96</u>	<u>4/23/96</u>
	1) POTASSIUM PHOSPHATE SOLUTION	1%	6.9	8.2
10	2) POTASSIUM PHOSPHONATE SOLUTION	1%	18.7	28.8
	3) POTASSIUM PHOSPHATE SOLUTION	0.5%	8.9	10.7
15	+ POTASSIUM PHOSPHONATE SOLUTION	+ 0.5%		
	4) Cu-ALEXIN	0.2 lb ai	8.8	11.6
	5) Fe-ALEXIN	0.2 lb ai	12.7	12.9
	6) CONTROL	-	23.0	34.8

20

EXAMPLE NO. 5

Citrus Preharvest Spray Trials
Kerr Center, Vero Beach, Florida

25

In Example No. 5, twenty-two fruits, each from five nova tangelo trees, were randomly selected and marked. The fruits were scuffed with an emery board and then sprayed with the compositions set forth below in Table 5, with five treatments on the same day. Seven days later, the fruit was picked and stored for two weeks in sealed white plastic bags, then examined for penicillium mold.

30

The degree of infection was rated using following scores: 0 = no infection, 1 = light infection (less than 30% coverage), 2 = severe infection (100% coverage). The degree of infection is the mean of scores from 22 fruit.

5

TABLE 5

<u>TREATMENT</u>	<u>RATE</u>	<u>DEGREE OF INFECTION</u>	<u>#ROTTEN FRUIT</u>	<u>PHYTOTOXICITY</u>
1) POTASSIUM PHOSPHONATE SOLUTION	1%	1	9	0
2) POTASSIUM PHOSPHONATE SOLUTION	2%	0.91	12	0
3) POTASSIUM PHOSPHATE SOLUTION	1%	0	0	0
4) POTASSIUM PHOSPHATE SOLUTION + POTASSIUM PHOSPHONATE SOLUTION	0.5% + 0.5%	0.27	3	0
5) CONTROL	-	1.41	12	0

10

15

20


25

The above Examples demonstrate that the inventive compositions are useful in protecting plants against attack by Phycomycete , Ascomycete and other fungi and bacteria with the application of one solution.


30

It will also be appreciated that compositions for controlling Phycomycete and Ascomycete fungi diseases in plants may also contain phosphate and phosphonate compounds comprising a fungicidally effective amount of at least a first salt having the

5 following formula:



And a second salt having the following formula:



20 Where R₁ is selected from group consisting of H, K, an alkyl radical containing from 1 to 4 carbon atoms, halogen-substituted alkyl or nitro-substituted alkyl radical, an alkenyl, halogen-substituted alkenyl, alkynyl, halogen-substituted alkynyl, alkoxy-substituted alkyl radical, ammonium substituted by alkyl and hydroxy alkyl radicals;

R₂ and R₃ are selected from a group consisting of H and K;

25 Me is selected from a group consisting of K, alkaline earth metal cations, aluminum atom; ammonium cation; and

n is a whole number from 1 to 3, equal to the valence of Me.

It will be appreciated that foliar applications of the inventive compositions will be effective as a common agricultural practice to control root pathogens caused by *Phytophthora*, *Phythium*, and foliar infections caused by *Plasmopara*.

5 It will also be appreciated that the inventive compositions will have biocidal and
arthropod pest control activity, and also have fertilizer effects, in plants. (See e.g., U.S.
Pat. Nos. 5,206,228, 5,133,891 and 5,514,200).

According to another aspect of the invention, compositions including both
potassium phosphate and potassium phosphonate have been found to produce substantial
10 growth response in certain plants. Table 6 below is illustrative:

TABLE 6

FOLIAR NUTRITIONAL EVALUATIONS OF K-PHOS AND PHOS-MIGHT ON
RHAPHIOLEPIS INDICA

15	TREATMENT	RATE	No. of Shoots/Rep.				AVG
			A	B	C	D	
	1) POTASSIUM PHOSPHONATE SOLUTION	1%	22	13	20	15	17.5
20	2) POTASSIUM PHOSPHATE SOLUTION	1%	28	23	25	32	27.0
	3) POTASSIUM PHOSPHONATE SOLUTION	1/2%	23	26	24	8	20.3
25	+ POTASSIUM PHOSPHATE SOLUTION	+ 1/2%					
30	4) POTASSIUM PHOSPHATE SOLUTION	2%	24	38	32	24	29.5
	5) CONTROL-	--	4	1	7	3	3.8

5 Experimental units = 3 x 3 gallon containers x 4 repetitions with treatments assigned in a randomized complete block design. Application dates for treatments A-D were approximately one, two, four, and three weeks apart respectively (October 31, 1996; November 7, 1996; November 20, 1996, December 19, 1996, and January 8, 1997 respectively; rated January 14, 1997).

10 As can be seen by Table 6, all four nutritional treatments provided growth responses greater than the untreated control, and K-PHOS at 1% and 2% were superior to PHOS-MIGHT at 1% and the combination of 1/2% each of PHOS-MIGHT + K-PHOS. The combination, however, produced substantially higher growth responses than the control, in fact, on a level similar to the individual treatments of K-PHOS or PHOS-MIGHT. The higher growth responses, compared to the control, can be expected when
15 the inventive composition of Potassium Phosphate salt and Potassium phosphonate salt aqueous solution wherein each said salt is present in solution from about .25% vol./vol. to about 3.0% vol./vol., and preferably between .5% vol./vol. and 2.0% vol./vol. It is theorized that the unexpected growth responses are due to the slow conversion of PO_3 in
20 the inventive composition to PO_4 after application. This composition further provides the added benefit of being fungicidal as well.

The disclosures in all references cited herein are incorporated by reference.

What is Claimed is:

1 1. A fertilizer composition for stimulating growth in plants comprising:
2 a growth stimulating effective amount of at least a first salt selected from a group
3 consisting of K_2HPO_3 , KH_2PO_3 and K_3PO_3 ; and a second salt selected from a group
4 consisting of K_2HPO_4 , KH_2PO_4 and K_3PO_4 .

1 2. The composition according to claim 1 wherein said composition
2 comprises an aqueous solution, wherein each said first and second salt being present in
3 solution from about .25% vol./vol. to about 5% vol./vol.


1 3. The composition according to claim 1 wherein the amount of said first
2 said salt is one part by weight and the amount of said second salt is between 0.001 and
3 1,000 parts by weight.

1 4. A method of stimulating growth in plants comprising:
2 applying to the plants in growth stimulating effective amounts at least a first salt selected
3 from a group consisting of K_2HPO_3 , KH_2PO_3 and K_3PO_3 ; and a second salt selected
4 from a group consisting of K_2HPO_4 , KH_2PO_4 and K_3PO_4 .


1 5. The method according to claim 4 wherein said composition comprises an
2 aqueous solution, wherein each said first and second salt being present in solution from
3 about .25% vol./vol. to about 5% vol./vol.

1 6. The method according to claim 4 wherein the amount of said first said salt
2 is one part by weight and the amount of said second salt is between 0.001 and 1,000 parts
3 by weight.

1 7. A composition for fertilizing comprising:
2 a growth stimulating effective amount of at least a first salt having the following
3 formula:



9 and a second salt having the following formula:



17 where R₁ is selected from a group consisting of H, K, an alkyl radical containing
18 from 1 to 4 carbon atoms, halogen-substituted alkyl or nitro-substituted alkyl radical, an
19 alkenyl, halogen-substituted alkenyl, alkynyl, halogen-substituted alkynyl, alkoxy-
20 substituted alkyl radical, ammonium substituted by alkyl or hydroxy alkyl radicals;

21 R₂ and R₃ are selected from a group consisting of H and K;

22 Me is selected from a group consisting of K, alkaline earth metal cations, or
23 aluminum atom; ammonium cation; and

24 n is a whole number from 1 to 3, equal to the valence of Me.

1 8. The composition according to claim 7 wherein said composition
2 comprises an aqueous solution, wherein each said first and second salt being present in
3 solution from about .25% vol./vol. to about 5% vol./vol.

1 9. The composition according to claim 7 wherein the amount of said first
2 said salt is one part by weight and the amount of said second salt is between 0.001 and
3 1,000 parts by weight.

4 10. The composition according to claim 1 which is fungicidal.

1 11. A fungicidal/fertilizer composition comprising from about .25% vol./vol.
2 to about 5% vol./vol. of a phosphite salt and a phosphate salt mixture.

1 12. A composition according to claim 10 wherein the phosphite salt and the
2 phosphate salt each comprise about 1/2% by weight of the composition.

1 13. A composition according to claim 10 which is aqueous.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/03459

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A01N 57/00, 57/10, 57/18, 57/26, 59/26; C05B 7/00; C05G 3/00
US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 71/34; 424/601, 605; 504/101; 514/76, 114, 131, 134, 135, 141, 143, 148

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,139,616 A (DUCRET et al.) 13 February 1979, see entire document.	1-13
Y	US 5,395,418 A (VETANOVETZ et al.) 07 March 1995, see entire document.	1-13
Y	US 5,169,646 A (HORRIERE et al.) 08 December 1992, see entire document.	1-13
Y	US 5,514,200 A (LOVATT) 07 May 1996, see entire document.	1-13
Y	US 4,075,324 A (THIZY et al.) 21 February 1978, see entire document.	1-13
Y	US 4,849,219 A (STAUB et al.) 18 July 1989, see entire document.	1-13



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

16 APRIL 1998

Date of mailing of the international search report

16 JUL 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ALLEN J. ROBINSON

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/03459

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,395,418 A (VETANOVETZ et al.) 07 March 1995, see entire document.	1-13
Y	US 5,070,083 A (BARLET) 03 December 1991, see entire document.	1-13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/03459

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

71/34; 424/601, 605; 504/101; 514/76, 114, 131, 134, 135, 141, 143, 148