



US007354754B2

(12) **United States Patent**
Mitani et al.

(10) **Patent No.:** **US 7,354,754 B2**
(45) **Date of Patent:** **Apr. 8, 2008**

(54) **MICROORGANISM SENSITIVE TO
LYSOZYME**

JP 61-280273 12/1986
JP 07-255484 10/1995

(75) Inventors: **Yasuo Mitani**, Sapporo (JP); **Nobutaka Nakashima**, Sapporo (JP); **Tomohiro Tamura**, Sapporo (JP)

(73) Assignee: **National Institute of Advanced Industrial Science and Technology**, Tokyo (JP)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 296 days.

(21) Appl. No.: **10/524,630**

(22) PCT Filed: **Aug. 14, 2003**

(86) PCT No.: **PCT/JP03/10342**

§ 371 (c)(1),
(2), (4) Date: **Feb. 16, 2005**

(87) PCT Pub. No.: **WO2004/018651**

PCT Pub. Date: **Mar. 4, 2004**

(65) **Prior Publication Data**

US 2006/0166312 A1 Jul. 27, 2006

(30) **Foreign Application Priority Data**

Aug. 20, 2002 (JP) 2002-239554

(51) **Int. Cl.**
C12N 1/22 (2006.01)

(52) **U.S. Cl.** **435/252**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

EP 0 064 680 A2 11/1982

OTHER PUBLICATIONS

Hashimoto et al., Site-directed mutagenesis for cysteine residues of cobalt-containing nitrile hydratase, *J Inorg Biochem.* Jul. 2002 vol. 91, No. 1 pp. 70-77.*

Arensdorf et al., Chemostat approach for the directed evolution of biodesulfurization gain-of-function mutants, *Environ Microbiol.* Feb. 2002, vol. 68, No. 2, pp. 691-698.*

Bailey et al., "Emulsification of Crude Oil by *Rhodococcus erythropolis* Strain ST-2 via a Cell-Surface, Lysozyme-Sensitive Glycoprotein," *System. Appl. Microbiol.*, 1997, vol. 20, pp. 545-548.

Inoue et al., "Isolation and Characterization of Lysozyme-Sensitive Mutants of *Staphylococcus aureus*," *Journal of Bacteriology*, Dec. 1980, pp. 1186-1189.

Hirasawa et al., "A Mutation in the *Corynebacterium glutamicum ItsA* Gene Causes Susceptibility to Lysozyme, Temperature-Sensitive Growth, and L-Glutamate Production," *Journal of Bacteriology*, May 2000, pp. 2696-2701.

* cited by examiner

Primary Examiner—Robert B. Mondesi

(74) *Attorney, Agent, or Firm*—Foley & Lardner LLP

(57) **ABSTRACT**

A microorganism of the genus *Rhodococcus* is provided which has a higher sensitivity to lysozyme at a low concentration than a wild-type strain, which can easily cause cell lysis, and from which a recombinant protein expressed therein is easily recovered. More specifically, a mutant microorganism of the genus *Rhodococcus* having a higher sensitivity to lysozyme than a wild-type microorganism of the genus *Rhodococcus*.

1 Claim, 5 Drawing Sheets
(2 of 5 Drawing Sheet(s) Filed in Color)

Fig. 1




Fig. 2

Rhodococcus erythropolis L-65




Fig. 3

Rhodococcus erythropolis L-88





Fig. 4

Rhodococcus erythropolis JCM3201



1

MICROORGANISM SENSITIVE TO LYSOZYME

TECHNICAL FIELD

The present invention relates to a microorganism of the genus *Rhodococcus* suitable for production of a recombinant protein, and more specifically, to a mutant strain more sensitive to lysozyme at a low concentration than a wild-type strain and capable of easily causing cell lysis. Use of the mutant strain makes it easier to extract and recover an expressed protein.

BACKGROUND ART

As a technique for expressing a recombinant protein in a microbial host, an expression system using *Escherichia coli* (*E. coli*) as a host has been generally and widely used. This is because *E. coli* is extremely easy to handle in a laboratory. More specifically, *E. coli* is confirmed as a safe microbial host and proliferates at a high rate, and its molecular biological operations in a laboratory are well established. On the other hand, development of host microorganisms having usefulness and advantages over *E. coli* in view of recombinant protein expression has progressed.

Microorganisms of the genus *Rhodococcus* are not pathogenic, except a few, and easily cultured in an ordinary laboratory. In addition to such essential conditions, they have the function as microbial catalysts, which is considered to be extremely useful from an industrial point of view. For these reasons, recently, various molecular biological techniques have been developed by use of such microorganisms. For example, in an attempt to add a further useful function to the microorganisms, techniques involving gene recombination has been developed. As a result, a shuttle vector was established which can replicate autonomously both in *E. coli* and in a microorganism of the genus *Rhodococcus* (R, De Mot et al., *Microbiology* 143, 3137-3147, (1997)). Furthermore, there is a report that a transposable transposon is present in a microorganism of the genus *Rhodococcus* (I, Nagy et al., *J. Bacteriol.* 179, 4635-4638 (1997)). Thus, it is expected to improve the microorganism in function, for example, by destroying the gene or integrating an exogenous gene into the chromosome.

In an attempt to further improve a microbial catalytic action based on such a molecular biological establishment, development of a vector for expressing a recombinant protein has been underway (JP Patent Publication (Kokai) No. 10-248578 A (1998)).

A microorganism of the genus *Rhodococcus*, namely, *Rhodococcus erythropolis* is not only useful as a microbial catalyst but also advantageous in that it can grow under a low temperature condition of 4° C. For this reason, it is expected that *Rhodococcus erythropolis* may produce a recombinant protein or the like in a temperature range where *E. coli* could not be used. Development of an inducible expression vector has been underway for such a purpose (the application already filed by Tamura, on Aug. 12, (2002)).

However, the cell wall of a microorganism of the genus *Rhodococcus* is particular and rigid in structure compared to those of other gram-positive bacteria. Therefore, extraction of a cellular content from the microorganism is complicated and difficult compared to the case of *E. coli*. More specifically, a microorganism of the genus *Rhodococcus* has an extremely strong resistance to a cell-wall lytic enzyme used generally for microbial cell lysis, such as lysozyme.

2

Examples of a cell lysis method include a method of exposing cell wall to a high-concentration antibiotic, such as penicillin, for a predetermined time to weaken the cell wall and then being subjected to cell lysis with lysozyme, and a method of applying ultrasonic treatment to bacterial cells for a long time to physically destroy them. However, these methods are complicated in process, it is difficult to treat a large amount of cells, and specimens are not likely to be treated uniformly. These problems are significant in view of industrial use. The effectiveness of an antibiotic such as penicillin is brought by inhibiting a de-novo synthesis of a cell wall and therefore the cell wall completed in synthesis is not affected by such an antibiotic. Therefore, the effect of such an antibiotic is considered to be low in low-temperature conditions where rapid growth is not expected.

It is known that the cell wall structure of a microorganism of the genus *Rhodococcus* is commonly seen in bacteria of the genus *Corynebacterium* (C. E. Barry III et al., *Prog. Lipid Res.* 37, 143-179 (1988)) and an invention similar to the present invention has been made in view of an object of facilitating a molecular biological operation such as transformation (JP Patent Publication (Kokoku) No. 01-003475 B (1989), T. Hirasawa et al., *J. Bacteriol.* 182, 2696-2701 (2000)).

DISCLOSURE OF THE INVENTION

The present invention is directed to providing a microorganism of the genus *Rhodococcus* improved in sensitivity to lysozyme and capable of being lysed with lysozyme at a low concentration, the microorganism allowing recovery of the protein by treatment of the microorganism with lysozyme after an exogenous gene is integrated to the microorganism and allowed to express. Furthermore, the present invention provides a method of producing an exogenous protein by use of the microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme.

The present inventors conducted studies with a view toward to attain an expression system for a recombinant protein by use of a microorganism of the genus *Rhodococcus* by overcoming a difficulty in extracting a cellular content. As a result, they found a novel microorganism of the genus *Rhodococcus* more sensitive to lysozyme at an extremely low concentration than a wild type strain. More specifically, mutation was induced in a wild type strain to obtain a mutant that cannot grow in a medium containing lysozyme. Mutation is usually induced by a chemical mutagen such as nitrosoguanidine or irradiation with radioactive rays. However, taking safety and convenience into consideration, ultraviolet ray irradiation is employed in the present invention. Furthermore, the present inventors found that cell lysis can be performed only by lysozyme treatment without pretreatment with penicillin or the like, and that a cellular content such as a recombinant protein accumulated in the cells can be extracted in a much easier manner than a conventional method. Based on these findings, the present invention was completed.

More specifically, the present invention includes

(1) A mutant of a microorganism of the genus *Rhodococcus* having a higher sensitivity to lysozyme than a wild-type microorganism of the genus *Rhodococcus*.

(2) The microorganism of the genus *Rhodococcus* according to item (1), in which the microorganism of the genus *Rhodococcus* is *Rhodococcus erythropolis*.

(3) The microorganism of the genus *Rhodococcus* according to item (2), in which the *Rhodococcus erythropolis* is *Rhodococcus erythropolis* strain L-65 (deposited on Jun. 12,

2002, originally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan) under Accession No. FERM BP-8443) or *Rhodococcus erythropolis* strain L-88 (deposited on Jun. 12, 2002, originally at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan, Accession No. FERM BP-8444).

(4) A method of producing a protein comprising transforming a mutant of a microorganism of the genus *Rhodococcus* having a higher sensitivity to lysozyme than a wild-type microorganism of the genus *Rhodococcus* by a gene encoding an exogenous protein; expressing the gene; and treating the microorganism of the genus *Rhodococcus* with lysozyme, thereby extracting and recovering the protein.

(5) The method of producing a protein according to item (4), in which the microorganism of the genus *Rhodococcus* is *Rhodococcus erythropolis*.

(6) The method of producing a protein according to item (5), in which the *Rhodococcus erythropolis* is *Rhodococcus erythropolis* strain L-65 (Accession No. FERM BP-8443) or *Rhodococcus erythropolis* strain L-88 (Accession No. FERM BP-8444).

Now, the present invention will be explained in detail.

A microorganism of the genus *Rhodococcus* according to the present invention is a mutant microorganism of the genus *Rhodococcus*, which has a higher sensitivity to lysozyme than a wild-type microorganism of the genus *Rhodococcus*. The microorganism of the genus *Rhodococcus* is not limited to a specific species and includes *Rhodococcus erythropolis*, *Rhodococcus fascians*, and *Rhodococcus opacus*. The wild-type microorganism of the genus *Rhodococcus* refers to a microorganism belonging to the genus *Rhodococcus* and having no genetic mutation, for example, *Rhodococcus erythropolis* strain JCM 3201. More specifically, the microorganism of the genus *Rhodococcus* according to the present invention having a higher sensitivity to lysozyme is a mutant derived from a wild-type microorganism of the genus *Rhodococcus* as a parent strain, and having an increased sensitivity to lysozyme compared to the parent strain. The phrase "having an increased sensitivity to lysozyme" means that cell lysis may occur at a low lysozyme concentration. If cell growth is inhibited when lysozyme is added to a medium where a microorganism is cultured, it is said that the microorganism has a sensitivity to lysozyme. The sensitivity to lysozyme can be expressed by a minimum lysozyme concentration capable of inhibiting the growth of a microorganism (a minimum growth inhibitory lysozyme concentration). A source providing lysozyme is not limited, for example, egg-white lysozyme may be used. The minimum growth inhibitory lysozyme concentration is obtained as follows: for example, a microorganism of the genus *Rhodococcus* is prepared in a liquid medium at a density of 1×10^5 to 1×10^6 cells/10 μ l. Lysozyme is serially diluted from a concentration of several hundreds μ g/ml to several μ g/ml. Each of the serial dilutions is added to 10 μ l of the liquid medium prepared above. The microorganism is cultured for several days. The lysozyme concentration that inhibits the growth of a microorganism of the genus *Rhodococcus* represents the minimum growth inhibitory lysozyme concentration. Alternatively, the degree of sensitivity to lysozyme can be determined by adding lysozyme to a predetermined concentration of a microorganism of the genus *Rhodococcus*, and monitoring a change in absorbance

while culturing. In this case, a strain non-sensitive to lysozyme continues to grow without causing cell lysis by lysozyme, and thus absorbance increases with time, whereas a strain sensitive to lysozyme causes cell lysis by lysozyme, and absorbance decreases rapidly.

The minimum growth inhibitory lysozyme concentration of a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention is preferably 50 μ g/ml or less, more preferably, 25 μ g/ml or less, and most preferably 13 μ g/ml or less. This is equal to or less than $1/8$, preferably $1/16$, and particularly preferably, $1/30$ of the minimum growth inhibitory lysozyme concentration of a wild type, that is, a parent strain.

A microorganism of the genus *Rhodococcus* generally has a high resistance to lysozyme, so that the microorganism cannot be lysed with lysozyme alone. Therefore, an antibiotic such as penicillin must be used at a high concentration to inhibit the cell wall synthesis of the microorganism during growth to weaken the cell wall, and then lysozyme is applied to the microorganism. However, the organism of the genus *Rhodococcus* according to the present invention can be lysed with lysozyme alone.

The organism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention can be obtained by treating a wild-type microorganism of the genus *Rhodococcus* such as *Rhodococcus erythropolis* strain JCM 3201 with a chemical mutagen or a physical mutagen, culturing it in an agar medium, transferring the colonies thus grown onto a medium containing lysozyme and a medium not containing lysozyme, culturing both, and selecting a bacterial cell not grown in the medium containing lysozyme. The sensitivity to lysozyme can be determined by the aforementioned sensitivity test. Examples of such chemical mutagens include alkylation agents such as N-methyl-N'-nitro-N-nitrosoguanidine and mustard gas, non-alkylation agents such as hydrazine and nitrite, DNA nucleotide analogs such as 5-bromo uracil, 2-aminopterin, and DNA intercalators, such as acrydine orange. Examples of such physical mutagens include ultraviolet rays, X-rays, γ -rays, and neutron beam. A method of treating a microorganism with a mutagen, the concentration of a chemical mutagen to be used, and the intensity of a physical mutagen to be used may be appropriately selected in accordance with a known method.

Examples of such a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention include *Rhodococcus erythropolis* strain L-65 (Accession No. FERM BP-8443) and *Rhodococcus erythropolis* strain L-88 (Accession No. FERM BP-8444).

In a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention, its sensitivity to lysozyme is higher than that of a wild-type strain, that is, a parent strain; however, the sensitivity to at least one member of antibiotics such as ampicillin, kanamycin, chloramphenicol, tetracycline, and thiostreptone is equal to that of the wild strain and thus has no significant difference. Even if the sensitivity differs between them, the difference from the wild type is not so large as that from lysozyme sensitivity. More specifically, a microorganism of the genus *Rhodococcus* having a high sensitivity to the lysozyme according to the present invention has a resistant gene to a certain antibiotic integrated as a selection marker. Therefore, when the microorganism of the present invention is transformed by an expression vector having an exogenous gene integrated therein and a transformant is selected based on the selection marker, non-transformed

microorganism of the genus *Rhodococcus* cannot grow since the sensitivity to the antibiotic is not lowered and thus only transformant can be selected. In this respect, as long as a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention has a sensitivity to the antibiotic to be used for selection even if the sensitivity to other antibiotics is low, it can be used.

A recombinant protein can be efficiently obtained by using a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention. To explain more specifically, the protein can be obtained by transforming a microorganism of the genus *Rhodococcus* having a high sensitivity to lysozyme according to the present invention with a gene encoding an exogenous protein derived from other species, culturing the transformed *Rhodococcus* microorganism in the conditions where the gene can be expressed, thereby expressing the exogenous protein, treating the microorganism having the expressed protein therein with lysozyme, thereby extracting the protein, and purifying and recovering the protein from the extract solution. The transformation of a microorganism of the genus *Rhodococcus* of the present invention may be performed in accordance with a known method. At this time, a transformation efficiency of a microorganism of the genus *Rhodococcus* with a reduced sensitivity to lysozyme is equivalent to that of a wild type strain, that is, a parent strain, even if there is a difference between them, the difference is not so significant. In some cases, the transformation efficiency is more or less lowered by the effect of introduction of mutation; however, there is no case where the efficiency of expressing and producing an exogenous protein is significantly reduced.

Transformation can be performed by using a known expression vector for a microorganism of the genus *Rhodococcus* or by expression vector pHN170 constructed by the present inventors such that the expression of the vector can be induced by thioestreptone.

A microorganism of the genus *Rhodococcus* transformed by integrating an exogenous gene therein is cultured. After the exogenous gene is expressed, the cells of the microorganism are collected by centrifugation, or the like, suspended in a buffer solution, such as a phosphate buffer, having lysozyme dissolved therein, and incubated at a temperature near an optimal temperature of lysozyme for several tens to several hours. The cells of the microorganism are lysed by the action of lysozyme and the expressed protein is extracted into the buffer solution. The extracted protein is purified by a known protein purification method to obtain the protein. The concentration of lysozyme to be used in cell lysis is 0.1 mg/ml to 10 mg/ml, preferably about 1 mg/ml. Purification can be performed by use of any separation and purification method. For example, ammonium sulfate precipitation, gel filtration, ion exchange chromatography, and affinity chromatography may be used singly or in an appropriate combination. In the case where an expression product is present in the form of a fusion protein with GST, His tag, it may be purified by use of the nature of a peptide or a protein that is fused to a desired protein. To explain more specifically, since GST has an affinity for glutathione, the desired protein can be efficiently purified by affinity chromatography by use of a column having a carrier to which glutathione is attached.

The specification includes the specification and/or contents of the drawings of JP Patent Application No. 2002-239554 based on which the present application claims the priority.

BRIEF DESCRIPTION OF THE DRAWINGS

The application contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

FIG. 1 shows photographs of LB agar mediums onto which serially diluted culture solutions are spotted for comparing their growth;

FIG. 2 is a graph showing a growth curve of *Rhodococcus erythropolis* strain L-65;

FIG. 3 is a graph showing a growth curve of *Rhodococcus erythropolis* strain L-88;

FIG. 4 is a graph showing a growth curve of *Rhodococcus erythropolis* strain JCM 3201; and

FIG. 5 shows SDS polyacrylamide electrophoresis of the cases where a PIP protein was expressed by *Rhodococcus erythropolis* strain L-65, L-88 and JCM3201.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention will be now explained with reference to Examples, which should not be construed as limiting the present invention.

EXAMPLE 1

Production of Lysozyme-sensitive Bacterial Strain

Rhodococcus erythropolis strain JCM 3201 was cultured in LB medium (1% Difco Bacto Tryptone, 0.5% Difco Yeast Extract, and 1% sodium chloride) with shaking at 30° C. The LB medium was taken in the middle of the logarithmic growth period and appropriately diluted. The dilution was applied onto an LB medium plate containing 1.5% agar at a density of about 5×10^3 bacterial cells per plate, and the application surface was irradiated with 254 nm ultraviolet ray by means of an ultraviolet-ray irradiation apparatus (manufactured by Atto, power: 4 W) placed at a distance of 15 cm from the application surface for 20 seconds. The medium irradiated with the ultraviolet ray was cultured stand still at 30° C. for 2 days to obtain about 5×10^2 colonies per plate. The colonies were scraped by a cocktail stick and inoculated onto a 96-well plate filled with about 150 μ l of LB medium. After the colonies were sufficiently suspended, a part of the suspension was inoculated onto a 96-well plate filled with 150 μ l of LB medium containing lysozyme derived from egg-white in a concentration of 50 μ g/ml (manufactured by Sigma, hereinafter simply referred to as "lysozyme"). The couple of plates thus obtained were cultured stand still at 30° C. for 2 days. As a result, a mutant strain capable of growing only in lysozyme-free LB medium was obtained as a lysozyme sensitive strain. Examples of such a novel lysozyme sensitive microorganism according to the present invention include *Rhodococcus erythropolis* strain L-65 and *Rhodococcus erythropolis* strain L-88, which was originally deposited on Jun. 12, 2002 under Accession Nos. FERM BP-8443 and FERM BP-8444, respectively at the International Patent Organism Depository of the National Institute of Advanced Industrial Science and Technology (Tsukuba Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki, Japan). A request for transferring the microorganisms from the original deposition to the international deposition based on the Budapest Treaty was made and accepted as of Jul. 28, 2003. The bacterial strain was inoculated in LB medium and cultured with shaking at 30° C. A part of the culture solution was taken in the middle of the logarithmic

growth period and diluted with fresh LB medium so as to contain 1×10^5 , 1×10^4 , 1×10^3 , 1×10^2 , and 1×10 cells in 10 μ l of the LB medium. The diluted culture solutions thus prepared were spotted onto each of LB agar mediums containing lysozyme in concentrations of 50, 25, 12.5 and 6.3 μ g/ml. After the mediums were cultured at 30° C. for 2 days, the presence or absence of bacterial cells grown on mediums was checked. In this manner, the minimum growth inhibition concentration was determined (Table 1 and FIG. 1). As shown in the figure, the culture solutions of bacterial strains JCM3201, L-65, and L-88 were dropped onto an LB agar medium containing no lysozyme and an LB agar medium containing lysozyme (12.5 μ g/ml) and then subjected to culturing. The bacterial strains JCM3201, L-65, and L-88 cells were dropped respectively onto the upper stage, the middle stage, and the lower stage of the LB agar medium and cultured as shown in FIG. 1. The numbers of bacterial cells contained in culture solutions were 1×10^5 , 1×10^4 , 1×10^3 , 1×10^2 , and 1×10 cells in this order from the left.

TABLE 1

Bacterial strain	Deposition No.	Minimum growth-inhibiting lysozyme concentration (μ g/ml)
<i>Rhodococcus erythropolis</i> strain L-65	FERM BP-8443	12.5
<i>Rhodococcus erythropolis</i> strain L-88	FERM BP-8444	12.5
<i>Rhodococcus erythropolis</i> strain JCM 3201	ATCC25544	>400

EXAMPLE 2

Turbidity Change of *Rhodococcus erythropolis* Strain L-65 Culture Solution by Addition of Lysozyme

To 100 ml of LB medium, *Rhodococcus erythropolis* strain L-65 was inoculated and cultured with shaking at 30° C. The absorbance (OD₆₀₀) of the culture solution was measured at an absorption wavelength of 600 nm every hour from the beginning of the logarithmic growth period. When OD₆₀₀ reached about 0.2, the volume of the culture solution was divided into halves. To one of them, lysozyme was added to a final concentration of 12.5 μ g/ml. No lysozyme was added to the other. While both solutions were further cultured continuously, the absorbance was measured. The results are shown in FIG. 2. The growth profiles of *Rhodococcus erythropolis* strain L-65 culture solutions with Lysozyme (12.5 μ g/ml) and without lysozyme were shown by absorbance at 600 nm. When OD₆₀₀ reached about 0.2, lysozyme was added (Indicated by the arrow in the figure). When lysozyme was added, a sharp decrease in absorbance was observed. This is considered because bacterial cell lysis was caused by lysozyme.

EXAMPLE 3

Turbidity Change of *Rhodococcus erythropolis* Strain L-88 Culture Solution by Addition of lysozyme

The same operation as in Example 2 was performed by use of *Rhodococcus erythropolis* strain L-88. Absorbance was measured and the results are shown in FIG. 3. The growth profiles of culture solutions of *Rhodococcus erythropolis* strain L-88 with lysozyme (12.5 μ g/ml) and without lysozyme were shown by absorbance at 600 nm. When OD₆₀₀ reached about 0.2, lysozyme was added (Indicated

the arrow in the figure). When lysozyme was added, a sharp decrease in absorbance was observed. This is considered because bacterial cell lysis was caused by lysozyme.

COMPARATIVE EXAMPLE 1

Turbidity Change of *Rhodococcus erythropolis* Strain JCM 3201 Culture Solution by Addition of Lysozyme

The same operation as in Example 2 was performed by use of *Rhodococcus erythropolis* strain JCM 3201. Absorbance was measured and the results are shown in FIG. 4. The growth profiles of culture solutions of *Rhodococcus erythropolis* strain JCM 3201 with lysozyme (12.5 μ g/ml) and without lysozyme were shown by absorbance at 600 nm. When OD₆₀₀ reached about 0.2, lysozyme was added (Indicated by the arrow in the figure). Regardless of the presence or absence of lysozyme, the same tendency of growth was observed.

EXAMPLE 4

Sensitivity of Lysozyme Sensitive Bacterial Strain to Ampicillin

The sensitivity of *Rhodococcus erythropolis* strain L-65 and L-88 to ampicillin was determined in the same manner as in Example 1. To explain more specifically, the bacterial cells of each strain were inoculated in LB medium and cultured with shaking at 30° C. A part of the culture solution was taken in the middle of logarithmic growth period and diluted with fresh LB medium so as to contain 1×10^5 , 1×10^4 , 1×10^3 , 1×10^2 , and 1×10 cells in 10 μ l of the LB medium. The diluted culture solutions thus prepared were dropped onto each of LB agar mediums containing lysozyme in concentrations of 15, 10, 1, and 0.1 μ g/ml. After the mediums were cultured at 30° C. for 2 days and the presence or absence of bacterial cells grown on the mediums was checked. In this manner, the minimum growth inhibition concentration was determined (Table 2). Similarly, the wild type and the mutant strain were compared for sensitivity to kanamycin, chloramphenicol, tetracycline, and thiostreptone; however, no significant difference was observed between them.

TABLE 2

Bacterial strain	Deposition No.	Minimum growth-inhibiting ampicillin concentration (μ g/ml)
<i>Rhodococcus erythropolis</i> strain L-65	FERM BP-8443	1
<i>Rhodococcus erythropolis</i> strain L-88	FERM BP-8444	1
<i>Rhodococcus erythropolis</i> strain JCM 3201	ATCC25544	15

EXAMPLE 5

Transformation Efficiency of Lysozyme Sensitive Strain

The transformation of *Rhodococcus erythropolis* was performed by an electroporation method. The method will be described in detail below. *Rhodococcus erythropolis* strain JCM 3201, L-65, and L-88 were cultured in 100 ml of LB medium with shaking at 30° C. until they reach their logarithmic growth periods. The culture solutions were cooled on ice for 30 minutes, and centrifugally separated to recover cells. To the recovered cells, 100 ml of ice-cooled sterilized water was added, stirred well, and again centrifu-

gally separated to recover cells. To the recovered cells, 100 ml of an ice-cooled 10% glycerin solution was added, stirred well, and centrifugally separated to recover cells. The cells were washed again with the ice-cooled 10% glycerin solution and suspended in 5 ml of an ice-cold 10% glycerin solution. Then, 400 μ l of the resultant cells were mixed with plasmid DNA (pHN144; Nakashima and Tamura; the full length sequence is represented by SEQ ID No: 1) capable of self-replicating in *Rhodococcus erythropolis*. The mixture solution was transferred to an electroporation cuvette (0.2 cm gap cuvette manufactured by Bio-Rad) and applied with an electric pulse by a gene introduction apparatus, gene pulser II, (manufactured by Bio-Rad) at an electric field of 12.5 kV/cm in strength, a capacitance of 25 μ F (the pulse controller), and an external resistance of 400 Ω . The mixture of cells and DNA treated with the electric pulse was mixed with 1 ml of LB medium and cultured at 30° C. for 4 hours. Thereafter, cells were collected and applied onto LB agar medium containing thioestreptone in a concentration of 10 μ g/ml and cultured at 30° C. for 3 days to obtain transformants for each case. The transformation efficiency (the number of colonies formed) per 1 μ g of DNA is shown in Table 3.

TABLE 3

Bacterial strain	Deposition No.	Transformation efficiency rate
<i>Rhodococcus erythropolis</i> strain L-65	FERM BP-8443	2.6×10^5
<i>Rhodococcus erythropolis</i> strain L-88	FERM BP-8444	2.5×10^5
<i>Rhodococcus erythropolis</i> strain JCM 3201	ATCC25544	4.0×10^5

EXAMPLE 6

Extraction of Recombinant Protein Produced by *Rhodococcus erythropolis* Strain L-65

A plasmid (pHN170, Nakashima and Tamura: the full length sequence is represented by SEQ ID No: 2) was constructed such that it could self-replicate in a bacterial cell of *Rhodococcus erythropolis* and could be induced by thioestreptone to express a proline iminopeptidase (hereinafter referred to as "PIP") protein (T. Tamura et al., FEBS Lett. 398, 101-105 (1996)) having a 6 \times histidine tag at the C terminal. This plasmid was introduced into *Rhodococcus erythropolis* strain L-65 by an electroporation method. Transformants were screened on LB agar medium containing tetracycline (20 μ g/ml). The transformants were inoculated on 4 ml of LB medium containing tetracycline (8 μ g/ml) and cultured with shaking at 30° C. until the absorbance at an absorption wavelength of 600 nm reached 0.8. The entire culture solution was added to 40 ml of LB medium containing thioestreptone (1 μ g/ml) and cultured with rotation in a vanned flask for 16 hours. After PIP protein was induced to express, bacterial cells were centrifugally collected at 1,500 \times g for 15 minutes. After the cells thus collected was suspended in 4 ml of a 50 mM phosphate buffer (pH8.0) containing 300 mM salt, lysozyme was added so as to obtain a final concentration of 1 mg/ml. The resultant solution was incubated at 37° C. for one hour, cooled on ice, and centrifuged at 10,000 \times g for 15 minutes to separate the supernatant (s) and the precipitate (p). An aliquot of 1 ml was taken from the obtained supernatant (s) and placed in another microcentrifuge tube. To the microcentrifuge tube, 50 μ l of Ni-NTA Superflow (manufactured

by QIAGEN) was added, which had been previously equilibrated with a 50 mM phosphate buffer (pH 8.0) containing 300 mM salt. The resultant mixture was incubated at 4° C. for one hour while turning it upside down, washed three times with 1 ml of a 50 mM phosphate buffer (pH 6.0) containing 300 mM salt and 10% glycerin, and thereafter, eluted with 50 μ l of a 50 mM phosphate buffer (pH 6.0) containing 500 mM EDTA, 300 mM salt and 10% glycerin to obtain 6 \times histidine-fused PIP protein. An aliquot (10 μ l) was taken from the protein thus obtained and subjected to SDS polyacrylamide gel electrophoresis. As a result, a clear band was detected near the molecular weight (34.3 KDa), which was predicted from the amino acid sequence of the 6 \times histidine-fused PIP protein (FIG. 5). On the other hand, the precipitate (p) was resuspended in 1 ml of a 100 mM sodium dihydrogen phosphate -10 mM tris chloride buffer (pH 8.0) containing 8M urea, allowed to stand alone for 30 minutes, and subjected to centrifugation at 10,000 \times g for 15 minutes. The resultant supernatant was transferred to a new microcentrifuge tube, 50 μ l of Ni-NTA Superflow, which had been previously equilibrated with a 100 mM sodium dihydrogen phosphate -10 mM tris chloride buffer (pH 8.0) containing 8M urea, was added to the microcentrifuge tube, and incubated at room temperature for one hour while turning upside down. After the microcentrifuge tube was washed three times with 1 ml of a 100 mM sodium dihydrogen phosphate -10 mM tris chloride buffer (pH 6.3) containing 8M urea, it was subjected to elution with 50 μ l of a 100 mM sodium dihydrogen phosphate -10 mM tris chloride buffer (pH 8.0) containing 500 mM EDTA and 8M urea to obtain 6 \times histidine-fused PIP protein. An aliquot (10 μ l) was taken from the protein thus obtained and subjected to SDS polyacrylamide gel electrophoresis (FIG. 5). Reference symbol M represents a molecular marker and molecular weight is shown at the left side of the figure to indicate an approximate molecular weight of each band. Band patterns of individual lanes are shown in FIG. 5 as follows.

Lane 1 (JCM 3201,s) shows an electrophoretic pattern of a sample obtained from the supernatant in the case where PIP is expressed by bacterial strain JCM 3201. Since cell lysis rarely takes place in a buffer solution under nondenaturation condition (containing no urea), a desired band (indicated by the arrow) is not detected. Lane 2 (JSM3201, p) shows an electrophoretic pattern of a sample obtained from the precipitate in the case where PIP is expressed by bacterial strain JSM 3201. Lysis takes place slightly in a buffer solution under denaturation conditions (containing urea) and thus a thin band of interest is detected.

Lane 3 (JCM3201+amp,s) shows an electrophoretic pattern of a sample obtained from the supernatant in the case where PIP is expressed by bacterial strain JCM 3201 where the sample is treated with ampicillin for 2 hours before cell collection. Since the sensitivity to lysozyme is increased by the treatment with ampicillin, cell lysis takes place even in nondenaturation conditions. As a result, a desired band can be clearly confirmed.

Lane 4 (JSM3201+amp, p) shows an electrophoretic pattern of a sample obtained from the precipitate in the case where PIP is expressed by bacterial strain JCM 3201 where the sample is treated with ampicillin for 2 hours before cell collection. It is considered that even though the cells previously treated with ampicillin do not cause cell lysis in a buffer under the nondenaturation conditions, they lyse in a buffer under denaturation conditions to give a detectable desired band.

Lane 5 (L-65, s) shows an electrophoretic pattern of a sample obtained from the supernatant in the case where PIP

11

is expressed by bacterial strain L-65. Cells were lysed completely with lysozyme treatment to give a detectable desired band.

Lane 6 (L-65, p) shows an electrophoretic pattern of a sample obtained from the precipitate in the case where PIP is expressed by bacterial strain L-65. Since the precipitate is considered to contain only residual cells after lysis, the desired band is not detected.

Lane 7 (L-88, s) shows the electrophoretic pattern of a sample obtained from the supernatant in the case where PIP is expressed by bacterial strain L-88. The same phenomenon as in the case of bacterial strain L-65 is considered.

Lane 8 (L-88, p) shows the electrophoretic pattern of a sample obtained from the precipitate in the case where PIP is expressed by bacterial strain L-88. The same phenomenon as in the case of strain L-65 is considered.

As for the antibiotic used in the aforementioned operation, a required amount of a solution containing 5 mg of tetracycline dissolved in 1 ml of 50 wt % ethanol or 10 mg of thiostrepton dissolved in 1 ml of dimethylsulfoxide was used.

EXAMPLE 7

Extraction of Recombinant Protein Produced by *Rhodococcus erythropolis* Strain L-88

The same operation as in Example 6 was performed except that *Rhodococcus erythropolis* strain L-88 was used in place of *Rhodococcus erythropolis* strain L-65 (FIG. 5).

COMPARATIVE EXAMPLE 2

Extraction of Recombinant Protein Produced by *Rhodococcus erythropolis* Strain JCM 3201

The same operation as in Example 6 was performed except that *Rhodococcus erythropolis* strain JCM 3201 was used in place of *Rhodococcus erythropolis* strain L-65 (FIG. 5).

12

COMPARATIVE EXAMPLE 3

Extraction of Recombinant Protein Produced by *Rhodococcus erythropolis* Strain JCM 3201

A transformant was prepared in the same manner as in Example 6 except that *Rhodococcus erythropolis* strain JCM 3201 was used in place of *Rhodococcus erythropolis* strain L-65. The expression of PIP protein was induced by thiostrepton. Two hours before cell collection, 480 µl of an aqueous solution of ampicillin (50 mg/ml) was added (a final concentration of 600 µg/ml) and subjected to cell collection. Thereafter, the same operations as in Example 6 were performed and the obtained sample was subjected to electrophoresis (FIG. 5).

All publications, patents and patent applications cited herein are incorporated herein in its entirety by reference.

INDUSTRIAL APPLICABILITY

As shown in Examples, a microorganism of the genus *Rhodococcus* of the present invention has an increased sensitivity to lysozyme compared to a wild type strain. The transformation efficiency of the microorganism of the present invention is not significantly changed from that of the wild-type strain. It is therefore possible to efficiently transform a microorganism of the genus *Rhodococcus* of the present invention by a gene encoding the exogenous protein, express the exogenous protein, cause cell lysis with lysozyme, and extract and recover the protein easily.

Sequence listing free text

SEQ ID No. 1: Plasmid pHN 144

SEQ ID No. 2: Plasmid pHN 170

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1

<211> LENGTH: 5108

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Plasmid pHN144

<400> SEQUENCE: 1

```

gagctcgacc ggcggggtcc cggacgggga agagcgggga gctttgccag agagcgacga      60
cttccccttg cgttggtgat tgcgggtcag ggcagccatc cgccatcgtc gcgtagggtg      120
tcacacccca ggaatcgctt cactgaacac agcagccggt aggacgacca tgactgagtt      180
ggacaccatc gcaaatccgt ccgatcccgc ggtgcagcgg atcatcgatg tcaccaagcc      240
gtcacgatcc aacataaaga caacgttgat cgaggacgtc gagcccctca tgcacagcat      300
cgcgccgggg gtggagtcca tcgaggctca cggcagcgac agcagtcctt ttccatctga      360
gttgctggat ctgtgcgggc ggcagaacat accggtccgc ctcatcgact cctcgatcgt      420
caaccagttg ttcaaggggg agcggaaggg caagacattc ggcacgccc gcgtccctcg      480

```

-continued

ccccgccagg	ttcggcgata	tcgcgagccg	gcgtggggac	gtcgtcgttc	tcgacggggt	540
gaagatcgtc	gggaacatcg	gcgcgatagt	acgcacgtcg	ctcgcgctcg	gagcgtcggg	600
gatcatcctg	gtggacagtg	acatcaccag	catcgcggac	cggcgtctcc	aaagggccag	660
ccgaggttac	gtcttctccc	ttcccgtcgt	tctctccggt	cgcgaggagg	ccatcgcctt	720
cattcgggac	agcggtatgc	agctgatgac	gctcaaggcg	gatggcgaca	tttccgtgaa	780
ggaactcggg	gacaatccgg	atcggctggc	cttgcgttcc	ggcagcgaaa	aggggtgggc	840
ttccgacctg	ttcgaggagg	cgtcttccgc	ctcggtttcc	atccccatga	tgagccagac	900
cgagtctctc	aacgtttccg	ttccctcgg	aatcgcgctg	cacgagagga	tcgacaggaa	960
tctcgcggcc	aaccgataag	cgcctctggt	cctcggacgc	tcggttctct	gacctcgatt	1020
cgtcagtgat	gatcacctca	cacggcagcg	atcaccactg	acatatacgag	gtcaacggtc	1080
gtggtccggg	cgggcactcc	tcgaaggcgc	ggccgacgcc	cttgaacgac	tcgatgactc	1140
tagaggatcc	ccgggtaccg	agctcgtcag	gtggcacttt	tcggggaaat	gtgcgcggaa	1200
ccccatattg	tttatttttc	taaatacatt	caaataatga	tccgctcatg	agacaataac	1260
cctgataaat	gcttcaataa	tattgaaaaa	ggaagagtat	gagtattcaa	catttccgtg	1320
tcgcccttat	tccctttttt	gcggcatttt	gccttcctgt	ttttgctcac	ccagaaacgc	1380
tggtgaaagt	aaaagatgct	gaagatcagt	tgggtgcacg	agtgggttac	atcgaactgg	1440
atctcaacag	cggtaagatc	cttgagagtt	ttcgcctcga	agaacgtttt	ccaatgatga	1500
gcacttttaa	agttctgcta	tgtggcgcgg	tattatcccg	tattgacgcc	gggcaagagc	1560
aactcggtcg	ccgcatacac	tattctcaga	atgacttggg	tgagtactca	ccagtcacag	1620
aaaagcatct	tacggatggc	atgacagtaa	gagaattatg	cagtgtctgcc	ataaccatga	1680
gtgataaac	tcgggccaac	ttactttctg	caacgatcgg	aggaccgaag	gagctaaccg	1740
cttttttgca	caacatgggg	gatcatgtaa	ctcgccttga	tcgttgggaa	ccggagctga	1800
atgaagccat	accaaacgac	gagcgtgaca	ccacgatgcc	tgtagcaatg	gcaacaacgt	1860
tcgcaaaact	attaactggc	gaactactta	ctctagcttc	ccggcaacaa	ttaatagact	1920
ggatggaggc	ggataaagtt	gcaggaccac	ttctgcgctc	ggcccttccg	gctggctggt	1980
ttattgctga	taaactctga	gccggtgagc	gtgggtctcg	cggtatcatt	gcagcactgg	2040
ggccagatgg	taagccctcc	cgtatcgtag	ttatctacac	gacggggagt	caggcaacta	2100
tgatgaacg	aaatagacag	atcgtgaga	taggtgcctc	actgattaag	catttgtaac	2160
tgtcagacca	agtttactca	tatatacttt	agattgattt	aaaacttcat	ttttaattta	2220
aaaggatcta	ggtgaagatc	ctttttgata	atctcatgac	caaaatccct	taacgtgagt	2280
ttcgtttcca	ctgagcgtca	gacccogtag	aaaagatcaa	aggatcttct	tgagatcctt	2340
ttttctcgcg	cgtaatctgc	tgcttgcaaa	caaaaaaac	accgctacca	gcgggtggtt	2400
gtttgccgga	tcaagagcta	ccaactcttt	ttccgaaggt	aactggcttc	agcagagcgc	2460
agataccaaa	tactgttctt	ctagtgtagc	cgtagttagg	ccaccacttc	aagaactctg	2520
tagcaccgcc	tacatacctc	gctctgctaa	tcctgttacc	agtggtgctg	gccagtggcg	2580
ataagtctgt	tcttaccggg	ttggaactca	gacgatagtt	accggataag	gcgcagcggg	2640
cgggctgaac	ggggggttcc	tgcacacagc	ccagcttggg	gcgaacgacc	tacaccgaac	2700
tgagatacct	acagcgtgag	ctatgagaaa	gcgccacgct	tcccgaaggg	agaaaggcgg	2760
acaggtatcc	ggtaacgggc	agggctggaa	caggagagcg	cacgaggggg	cttccagggg	2820
gaaacgctcg	gtatctttat	agtctctgct	ggtttcgcca	cctctgactt	gagcgtcgat	2880

-continued

ttttgtgatg	ctcgtcaggg	gggcgagacc	tatggaaaaa	cgccagcaac	gcggcctttt	2940
tacggttcct	ggccttttgc	tggccttttg	ctcacatggt	ctttcctgcg	ttatccccctg	3000
attctgtgga	taacctattt	accgcctttg	agtgagctga	taccgctcgc	cgcagccgaa	3060
cgaccgagcg	cagcgagtca	gtgagcgagg	aagcggaaga	gcccacaata	cgaaaaccgc	3120
ctctccccgc	gcggtggccg	attcattaat	gcagctggca	cgactagtgtg	tacacccgag	3180
aagctcccag	cgtcctcctg	ggccgcgata	ctcgaccacc	acgcacgcac	accgcactaa	3240
cgattcggcc	ggcgctcgat	tggccggcg	ctcgattcgg	cggcgctcgc	attcggcccg	3300
cgctcgatc	ggccggcgct	cgattcggcc	gagcagaaga	gtgaacaacc	accgaccacg	3360
cttccgctct	gcgcgccgta	cccgaacctac	ctcccgcagc	tcgaagcagc	tcccgggagt	3420
accgccttac	tcaccgcctt	gtgctcacca	tccaccgacg	caaagcccaa	cccgagcaca	3480
cctcttgca	caaggtgccc	accgtggctt	tccgctcgca	gggttccaga	agaatcga	3540
cgatccagcg	cggcaaggtt	caaaaagcag	gggttggtgg	ggaggaggtt	ttggggggtg	3600
tgcgcgggat	acctgatatg	gctttgtttt	gcgtagtcga	ataattttcc	atatagcctc	3660
ggcgctcgg	actcgaatag	ttgatgtggg	cgggcacagt	tgcccatga	aatccgcaac	3720
ggggggcggtg	ctgagcgatc	ggcaatgggc	ggatgctggtg	ttgcttcgcg	accggccgctt	3780
cgcgacgaac	aacctccaac	gaggtcagta	cggatgagc	cgcgacgacg	cattggcaat	3840
gcggtacgct	gagcattcac	cgcacgcggt	gctcggatct	atcgtcatcg	actgcatca	3900
cgttgacgcc	gcgatgctcg	cattcagca	accatccgac	catccggcgc	cgaactgggt	3960
tgcaaatcgc	ccgtccggcc	gcgcacacat	cggatggtgg	ctcggcccca	accacgtgtg	4020
ccgcaccgac	agcggccgac	tgacgccact	gcgctacgcc	caccgcatcg	aaaccggcct	4080
caagatcagc	gtcggcgccg	atttcgcgta	tggcgggcaa	ctgacaaaa	accgattca	4140
ccccgattgg	gagacgatct	acggcccgcc	caccccgctac	acattgcccgc	agctggccac	4200
catccacaca	ccccggcaga	tgccgctcgc	gcccgatcgg	gccgtgggcc	tgggcccga	4260
cgtcaccatg	ttcagcgcca	cccggcgatg	ggcatacccg	cagtgggtggc	aacaccgaaa	4320
cggaaaccggc	cgcgactggg	accatctcgt	cctgcagcac	tgccacgccc	tcaaacccga	4380
gttcacgaca	ccactgcctg	tcaccgaagt	acgcgccacc	gcgcaatcca	tctccaaatg	4440
gatctggcgc	aatttcaccg	aagaacagta	cgcgcccga	caagcgcac	tcgggtcaaaa	4500
aggcggcaag	gcaacgacac	tcgccaaca	agaagcctgc	cgaacaatg	caagaaagta	4560
cgaagaacat	acgatgagag	aggcgattat	ctgatgggcg	gagcaaaaa	tccggtgccc	4620
cgaagatga	cggcagcagc	agcagccgaa	aaattcgggtg	cctccactcg	cacaatccaa	4680
cgttggtttg	ctgagccgcg	tgacgattac	ctcggccggtg	cgaagctcgc	ccgtgacaaa	4740
gctgtcagc	tgccgaagca	ggggttgaag	taccgggaaa	tcgccgaagc	gatggaactc	4800
tcgaccggga	tcgtcggccc	attactgcac	gacgcccgca	ggcacggcga	gatttcagcg	4860
gaggatctgt	cggcgtaac	aagtcagcgg	ggtgtcgggt	tccggccggc	gctcggcaact	4920
cggaccggcc	ggcgatggtg	gttctgcctc	tggcgagcgc	tcagctaccg	ccgaaggcct	4980
gtcatcgacc	ggcttcgact	gaagtatgag	caacgtcaca	gcctgtgatt	ggatgatccg	5040
ctcagctcgc	accgctacct	gttcagctgc	cgcctcctgg	gcagtgagca	cggccaactc	5100
tcgttcaa						5108

-continued

```

<211> LENGTH: 8971
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Plasmid
      pHN170

<400> SEQUENCE: 2
gagctcgacc gcgcggtgcc cggacgggga agagcgggga gctttgccag agagcgacga      60
cttcccccttg cgttggtgat tgccggtcag ggcagccatc cgccatcgtc gcgtaggggtg    120
tcacacccca ggaatcgcgt cactgaacac agcagccggg aggacgacca tgactgagtt      180
ggacaccatc gcaaatccgt ccgatcccgc ggtgcagcgg atcatcgatg tcaccaagcc      240
gtcacgatcc aacataaaga caacgttgat cgaggacgtc gagcccctca tgcacagcat      300
cgcggccggg gtggagttea tcgaggtcta cggcagcgac agcagtcctt ttccatctga      360
gttgctggat ctgtgcgggc ggcagaacat accggtccgc ctcatcgact cctcgatcgt      420
caaccagttg ttcaaggggg agcgggaagg caagacattc ggcacgccc gcgtccctcg      480
cccggccagg ttcggcgata tcgcgagccg gcgtggggac gtcgtcgttc tcgacggggt      540
gaagatcgtc gggaacatcg gcgcatagtg acgcacgtcg ctcgcgctcg gagcgtcggg      600
gatcatcctg gtggacagtg acatcaccag catcgcggac cggcgtctcc aaagggccag      660
cggaggttac gtcttctccc ttcccgtcgt tctctccggg cgcgaggagg ccatcgctt      720
cattcgggac agcggtatgc agctgatgac gctcaaggcg gatggcgaca tttccgtgaa      780
ggaactcggg gacaatccgg atcggctggc cttgctgttc ggcagcgaaa aggggtgggcc      840
ttccgacctg ttcgaggagg cgtcttccgc ctcggtttcc atccccatga tgagccagac      900
cgagtctctc aacgtttccg tttccctcgg aatcgcgctg cacgagagga tcgacaggaa      960
tctcgcggcc aaccgataag cgcctctgtt cctcggacgc tcggttctc gacctcgatt    1020
cgtcagtgat gatcacctca cacggcagcg atcaccactg acatatacgag gtcacacggtc    1080
gtggtccggg cgggcactcc tcgaaggcgc ggcgcagccc cttgaacgac tcgatgactc    1140
tagagtaacg ggctactccg tttaacggac cccgttctca cgctttaggc ttgaccccgg    1200
agcctgcatg gggcattccg ccgtgaaccc ggtggaatgc ccccggcacc cgggctttcc    1260
agcaaagatc acctggcgcg gatgagtaa gcgtaacagaa cactccaca ggaggaccgt    1320
cgagatgaaa tctaacaatg cgctcatcgt catcctcggc accgtcaacc tggatgetgt    1380
aggcataggc ttggttatgc cggtaactgc gggcctcttg cgggatatcg tccattccga    1440
cagcatcgcc agtcactatg gcgtgctgct agcgtatat gcgttgatgc aatttctatg    1500
cgcacccggt ctcggagcac tgtccgaccg ctttggccgc cgcccagtcc tgctcgttc    1560
gctacttgga gccactatcg actacggcat catggcgacc acaccctcc tgtggattct    1620
ctaocccgga cgcactgtgg ccggcatcac cggcgccaca ggtgcggttg ctggcgctta    1680
tatcgcgcag atcaccgatg ggaagatcg ggctcgccac ttcgggctca tgagcgcttg    1740
tttcggcgtg ggtatggtgg caggccccgt ggcgggggga ctgttgggcg ccattctctt    1800
gcatgcacca ttcttgcgg cggcggtgct caacggcctc aacctactac tgggctgctt    1860
cctaatgcag gagtcgcata agggagagcg tcgtccgatg cccttgagag ccttcaaccc    1920
agtcagctcc ttccggtggg cgcggggcat gactatcgtc gccgcactta tgactgtctt    1980
ctttatcatg caactcgtag gacaggtgcc ggcagcgtc tgggtcattt tcggcgagga    2040
ccgctttcgc tggagcgcga cgatgatcgg cctgtcgtt gcggtattcg gaatcttga    2100

```

-continued

cgccctcget caagccttcg tcaactggtcc cgccacaaa cgtttcggcg agaagcagge	2160
cattatcgcc ggcattggcgg ccgacgcgct gggctacgtc ttgctggcgt tcgacgcg	2220
aggctggatg gccttcccca ttatgattct tctcgcttcc gggggcatcg ggatgcccgc	2280
ggtgcagccc atgctgtcca ggcaggtaga tgacgacat cagggacagc ttcaaggatc	2340
gctcgcggct cttaccagcc taacttcgat cattggaccg ctgatcgtca cggcgattta	2400
tgccgcctcg gcgagacat ggaacgggtt ggcattggatt gtaggcgccc ccctatacct	2460
tgtctgcctc cccgcgttgc gtcgcgggtc atggagccgg gccacctcga cctgaatgga	2520
agccggcggc acctcgttaa cggattcacc actccaagaa ttggagcaa tcaattcttg	2580
cggagaactg tgaatcgcga aaccaacct tggcagaaca tatccatcgc gtcgcacctc	2640
tccagcagcc gcacgcggcg catctcgggc agcgttgggt cctggccacg ggtgcgcaac	2700
tagaattgat ctctcgcacc gccaatggg catctgagaa tcctctcgt ttctcgcacg	2760
caacgtactt gcaacgttgc aactcctagt gttgtgaatc acacccacc ggggggtggg	2820
attgcagtca ccgatttggg ggggtcggcc aggaagatca cgtttacata ggagcttgca	2880
atgagctact ccgtgggaca ggtggccggc ttcgccggag tgacggtgcg cagcctgcac	2940
cactacgacg acatcggcct gctcgtaccg agcagcgcga gccacgcggg ccaccggcgc	3000
tacagcgacg ccgacctcga ccggctgcag cagatcctgt tctaccggga gctgggcttc	3060
ccgctcgcag aggtcgcgc cctgctgcac gaccgcggcg cggaccgcg cgcgcacctg	3120
cgccgcagc acgagctgct gtcgcggcgg atcgggaaac tgcagaagat gggggcggcc	3180
gtggagcagg cgatggaggc acgcagcatg ggaatcaacc tcaccccgga ggagaagttc	3240
gaggtcttcg gcgacttcga ccccgaccag tacgaggagg aggtccggga acgctggggg	3300
aacaccgacg cctaccgcca gtcgaaggag aagaccgct cgtacaccaa ggaggactgg	3360
cagcgcacc aggacgagge cgacgagctc acccgcgct tcgtcgccct gatggacgcg	3420
ggtgagcccg ccgactccga gggggcgatg gacgcgcggc aggaccaccg gcagggeatc	3480
gcccgaacc actacgactg cgggtacgag atgcacacct gcctgggcca gatgtacgtg	3540
tccgacgaac gtttcacgcg aaacatcgac gccgccaagc cgggctcgc cgcctacatg	3600
cgcgacgcga tcctcgcгаа cgcctcggc cacacccct gagcgggtgt cgtggcccgg	3660
gtctcccgc cggctcacc ccaaggctca ctcccggcc acgaccaccg ccgtcccgtc	3720
cgcgacacc tcggtgccc cgtccgcgc ctccgtcagc tcgaaacgga agatcccgg	3780
gtaccgagct cgtcaggtgg cacttttcgg ggaatgtgc gcggaacccc tatttgttta	3840
tttttctaaa tacattcaaa tatgtatcgg ctcatgagac aataacctg ataaatgctt	3900
caataatatt gaaaaaggaa gactatgagt attcaacatt tccgtgtcgc ccttattccc	3960
tttttgcgg cattttgcct tcctgtttt gctcaccag aaacgctggg gaaagtaaaa	4020
gatgctgaag atcagttggg tgacagagtg ggttacatcg aactggatct caacagcggg	4080
aagatccttg agagttttcg cccgaagaa cgttttcaa tgatgagcac ttttaaagtt	4140
ctgctatgtg gcgcggtatt atcccgtatt gacgcgggc aagagcaact cggtcgccc	4200
atacactatt ctcaaatga cttggttgag tactcaccag tcacagaaaa gcactttacg	4260
gatggcatga cagtaagaga attatgcagt gctgcataa ccatgagtga taacctgcg	4320
gccaaacttac ttctgacaac gatcggagga ccgaaggagc taaccgcttt ttgacacaac	4380
atgggggatc atgtaactcg ccttgatcgt tgggaaccgg agctgaatga agccatacca	4440
aacgacgagc gtgacaccac gatgcctgta gcaatggcaa caacgttgcg caaactatta	4500

-continued

actggcgaac	tacttactct	agcttcccgg	caacaattaa	tagactggat	ggaggcggat	4560
aaagttgcag	gaccacttct	gcgctcggcc	cttccggctg	gctggtttat	tgctgataaa	4620
tctggagccg	gtgagcgtgg	gtctcgcggt	atcattgcag	cactggggcc	agatggtaag	4680
ccctcccgta	tcgtagttat	ctacacgacg	gggagtcagg	caactatgga	tgaacgaaat	4740
agacagatcg	ctgagatagg	tgccctcactg	attaagcatt	ggtaactgtc	agaccaagtt	4800
tactcatata	tacttttagat	tgatttaaaa	cttcattttt	aatttaaaag	gatctagggtg	4860
aaagatccttt	ttgataatct	catgacccaa	atcccttaac	gtgagttttc	gttccactga	4920
gcgtcagacc	ccgtagaaaa	gatcaaaagga	tcttcttgag	atcctttttt	tctgcgcgta	4980
atctgctgct	tgcaaaaaa	aaaaccaccg	ctaccagcgg	tggtttgttt	gccggatcaa	5040
gagctaccaa	ctctttttcc	gaaggtaact	ggcttcagca	gagcgcagat	accaaatact	5100
gttcttctag	tgtagccgta	gttaggccac	cacttcaaga	actctgtagc	accgcctaca	5160
tacctcgctc	tgctaactct	gttaccagtg	gctgctgcca	gtggcgataa	gtcgtgtcct	5220
accgggttgg	actcaagacg	atagttaccg	gataaggcgc	agcggtcggg	ctgaacgggg	5280
ggttcgtgca	cacagcccag	cttgagcga	acgacctaca	ccgaactgag	atacctacag	5340
cgtgagctat	gagaaagcgc	cacgcttccc	gaaggggagaa	aggcggacag	gtatccggtg	5400
agcggcaggg	tcggaacagg	agagcgcacg	agggagcttc	cagggggaaa	cgctcggtat	5460
ctttatagtc	ctgtcggggt	tcgccacctc	tgacttgagc	gtcgattttt	gtgatgctcg	5520
tcaggggggc	ggagcctatg	gaaaaacgcc	agcaacgcgg	cctttttacg	gttctctggc	5580
ttttgctggc	cttttctca	catgttcttt	cctgcgttat	ccctgattc	tgtggataac	5640
cgtattaccg	cctttgagtg	agctgatacc	gctcgcgcga	gccgaacgac	cgagcgcagc	5700
gagtcagtga	gcgaggaagc	ggaagagcgc	ccaatacga	aaccgcctct	ccccgcgcgt	5760
tgcccgattc	attaatgcag	ctggcacgac	tagagtcccg	ctgaggcggc	gtagcaggtc	5820
agccgcccc	gcggtggtca	ccaaccgggg	tggaacggcg	ccggtatcgg	gtgtgtccgt	5880
ggcgcctcatt	ccaacctccg	tgtgtttgtg	caggtttcgc	gtgttgacgt	ccctcgcacc	5940
ggcaccgcga	gcgaggggct	cacgggtgcc	ggtgggtcga	ctagtttatt	aatgatgatg	6000
atgatgatgc	aggtgtttca	ggatgaaatc	cgaaagcaac	ttgttgtatc	cttcacgatc	6060
ctcccacatc	gtgaggtgcg	agcaatccct	gaagacgtga	agtccgaac	cagctatttt	6120
ttcatgtatg	actctggcca	cgtttgccgt	gacctcatcg	tattcgcca	ccgttataag	6180
gggtggggatc	tttattgcag	atattttgtc	cgtgatatcc	cagtccttta	tcgtgccgggt	6240
tatggtgaac	tcattcgggc	cgttcattat	cctgtatacg	tttcgccttt	ccgcgtattc	6300
tagtgatttg	agtacctcgg	gcggccaatc	ctctgatctc	agcagatgct	gatggtaaaa	6360
gtagttcacg	gcctcctgat	attctggatt	ctcgtaagat	ccagatgaac	cgtatttttt	6420
aatggcatct	ctgtactttg	ccgggagctc	gtcaatgagc	ctgttcatct	ccttcaccgt	6480
cagaggggact	gaagataagc	ctccggatag	gatgagccct	ttcagatgat	cctgggtactt	6540
gactgcgtat	gccagcgcga	gcgctccacc	atatgatgac	cccatcaaaa	ataccttctc	6600
gttgccgaac	agctttgate	ttagggcctc	tgccctctcc	acaccatagt	caattgtgaa	6660
tttagactga	tcgggttctc	cggatctacc	gcataccaaac	tgatcgtaga	atagaaccgt	6720
tatccccttc	ttggtcatat	ccctgagaga	aagcaggtaa	tcgtgggaca	tgccccggcc	6780
cccgtgcatg	gtcattagct	ttgctttctc	ctcaggggct	ttgcacagct	tgtaaataat	6840

-continued

ataaattccg ttacaccttg cgtagttttc tatgcattcc tgateccatgg ccgctccctt	6900
ctctgacgcc gtccacgctg cctcctcacg tgacgtgagg tgcaagcccg gacgttccgc	6960
gtgccacgcc gtgagccgcc gcgtgccctc ggctccctca gcccgggcgg ccgtgggagc	7020
ccgctcgtat atgtacaccc gagaagctcc cagcgtctctc ctggggccgg ataactcgacc	7080
accacgcacg cacaccgcac taacgattcg gccggcgctc gattcggccc gcgctcgatt	7140
cggccggcgc tcgattcggc cggcgctcga ttcggccggc gctcgattcg gccgagcaga	7200
agagtgaaca accaccgacc acgcttccgc tctgcgcgcc gtacccgacc tacctcccgc	7260
agctcgaagc agctcccggg agtacccgcg tactcaccgg cctgtgctca ccatccaccg	7320
acgcaaaacc caaccggagc acacctcttg caccaagggtg ccgaccgtgg ctttccgctc	7380
gcagggttcc agaagaaatc gaacgatcca gcgcggcaag gttcaaaaag caggggttgg	7440
tggggaggag gttttggggg gtgtcgcggg gatacctgat atggctttgt tttgcgtagt	7500
cgaataattt tccatatagc ctccggcggc cggactcgaa tagttgatgt gggcgggac	7560
agttgccccg tgaatccgc aacggggggc gtgctgagcg atcggcaatg ggcggatgcg	7620
gtgttgcttc cgcaccggcc gttcgcgacg aacaacctcc aacgaggtca gtaccggatg	7680
agccgcgacg acgcatggcc aatgcggtag gtcgagcatt caccgcacgc gttgctcggg	7740
tctatcgtca tcgactcgca tcacgttgac gcccgatgc gcgcattcga gcaaccatcc	7800
gaccatccgg cgcgaactg ggttgcaaaa tcgccgtccg gccgcgcaca catcgatgg	7860
tggctcggcc ccaaccacgt gtgccgcacc gacagcggcc gactgacgcc actgcccctac	7920
gcccaccgca tcgaaaccgg cctcaagatc agcgtcggcg gcgatttcgc gtatggcggg	7980
caactgacca aaaaccgat tcacccgat tgggagacga tctacggccc ggcacccccg	8040
tacacattgc ggcagctggc caccatccac acacccggc agatgcccgcg tcggcccgat	8100
cgggcccgtg gcctggggcc caacgtcacc atgttcgacg ccacccggcg atgggcatac	8160
ccgcagtggt ggcaacaccg aaacggaaac gcccgcgact gggaccatct cgtcctgcag	8220
cactgcacag ccgtcaaac caggttcacg acaccactgc cgttcaccga agtacgcgcc	8280
accgcgcaat ccatctcaa atggatctgg cgcaatttca ccgaagaaca gtaccgagcc	8340
cgacaagcgc atctcggta aaaaggcggc aaggcaacga cactcgcaa acaagaagcc	8400
gtccgaaaca atgcaagaaa gtacgacgaa catacgatgc gagaggcgat tatctgatgg	8460
gcccagccaa aatccgggtg cgcgaaaga tgacggcagc agcagcagcc gaaaaattcg	8520
gtgcctccac tcgcacaatc caacgcttgt ttgctgagcc gcgtgacgat tacctcggcc	8580
gtgcgaaagc tcgccgtgac aaagctgtcg agctgcggaa gcaggggttg aagtaccggg	8640
aaatcgccga agcgatggaa ctctcgaccg ggatcgtcgg ccgattactg cacgacgccc	8700
gcaggcacgg cgagatttca gcggaggatc tgctggcgta accaagtcag cgggttctcg	8760
ggttccggcc ggccgtcggc actcggaccg gccggcgat ggtgttctgc ctctggcgca	8820
gcgtcagcta ccgccgaagg cctgtcatcg accggcttcg actgaagtat gagcaacgctc	8880
acagcctgtg attgatgat ccgctcacgc tcgaccgcta cctgttcagc tgccgcccgc	8940
tgggcatgag caacggccaa ctctcgttca a	8971

The invention claimed is:

1. An isolated microorganism of the genus *Rhodococcus* that is *Rhodococcus erythropolis* strain L-65 (Accession No.

FERM BP-8443) or *Rhodococcus erythropolis* strain L-88 (Accession No. FERM BP-8444).

* * * * *