



US 20230066720A1

(19) **United States**(12) **Patent Application Publication**
PERNI et al.(10) **Pub. No.: US 2023/0066720 A1**(43) **Pub. Date: Mar. 2, 2023**(54) **NOVEL PRODRUGS AND CONJUGATES OF DIMETHYLTRYPTAMINE***C07D 403/06* (2006.01)*C07D 401/06* (2006.01)(71) Applicant: **ATAI Life Sciences AG**, Berlin (DE)(52) **U.S. Cl.**CPC *C07D 209/16* (2013.01); *C07D 209/08* (2013.01); *C07D 209/12* (2013.01); *C07F 9/5728* (2013.01); *C07D 209/14* (2013.01); *C07D 403/06* (2013.01); *C07D 401/06* (2013.01)(72) Inventors: **Robert B. PERNI**, Marlborough, MA (US); **Glenn SHORT**, Scituate, MA (US); **Srinivasan G. RAO**, Encinitas, CA (US); **Tanweer A. KHAN**, Bridgewater, NJ (US)

(57)


ABSTRACT

Provided herein are compounds of Formula (I), or pharmaceutically acceptable salt thereof, wherein R₁, R₂, R₃, R₄, R₅, R₈, R₉, X, Y, Z, W, and m are defined herein. Also provided herein are pharmaceutical compositions comprising a compound of Formula (I) or pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.


(21) Appl. No.: **17/889,013**(22) Filed: **Aug. 16, 2022****Related U.S. Application Data**

(63) Continuation of application No. 17/836,984, filed on Jun. 9, 2022.

(60) Provisional application No. 63/229,879, filed on Aug. 5, 2021, provisional application No. 63/208,874, filed on Jun. 9, 2021.

Publication Classification(51) **Int. Cl.***C07D 209/16* (2006.01)*C07D 209/08* (2006.01)*C07D 209/12* (2006.01)*C07F 9/572* (2006.01)*C07D 209/14* (2006.01)

(I)









FIG. 1A

D



PH-VLS-02-23-0



PH-VLS-02-23-10

FIG. 1B




FIG. 1C




FIG. 2A

PO Dose	Mean plasma Concentration (ng/mL)	
	Time(h)	VLS-02-23-0
		VLS-02-23-10
	0.25	NA
	0.5	NA
	1	NA
	3	NA
	5	NA
	8	NA




FIG. 2B

NOVEL PRODRUGS AND CONJUGATES OF DIMETHYLTRYPTAMINE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present Application is a continuation of U.S. application Ser. No. 17/836,984 filed Jun. 9, 2022, which claims priority to U.S. Provisional Patent Application No. 63/208,874 filed Jun. 9, 2021, and U.S. Provisional Application No. 63/229,879 filed Aug. 5, 2021, the contents of which are hereby incorporated by reference in their entirety.

BACKGROUND OF THE INVENTION


[0002] N,N-dimethyltryptamine (DMT) is a naturally occurring compound found in numerous plant species and botanical preparations, such as the hallucinogenic infusion known as ayahuasca, and classified as a classic serotonergic psychedelic that induces intense modifications in perception, emotion, and cognition in humans. At higher doses, DMT has a rapid onset, intense psychedelic effects, and a relatively short duration of action with an estimated half-life of less than fifteen minutes. Like other hallucinogens in the tryptamine family, DMT binds to serotonin receptors to produce euphoria and psychedelic effects. Unfortunately, DMT is metabolically unstable and is readily converted by monoamine oxidases (MAO's) to indoleacetic acid and N-oxidation metabolites resulting in poor oral bioavailability.

[0003] Serotonergic psychedelics have also demonstrated promising antidepressant, anxiolytic, and anti-addictive properties.

[0004] There remains a need for improved prodrugs of tryptamines, such as N,N-dimethyltryptamine (DMT).

SUMMARY OF THE INVENTION

[0005] In one aspect, the present disclosure provides a compound of Formula (I):




[0006] or a pharmaceutically acceptable salt thereof;

[0007] wherein,

[0008] R_1 and R_2 are independently hydrogen, alkyl, cycloalkyl, —or R_1 and R_2 together with the atoms to which they are attached form a heterocycle;

[0009] X, Y and Z are independently hydrogen, halogen, —O—, —S—, —NR_A—, or —(P=O)—OR_A (—OR'), wherein R_A is hydrogen or alkyl;

[0010] R' , R_3 , R_4 , R_5 and R_6 are independently absent, hydrogen, alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0011] R_7 and R_8 are independently hydrogen or alkyl;

[0012] R_9 is hydrogen, alkyl, or halogen, preferably a bromo


[0013] L and M are independently alkyl or aryl;

[0014] W is oxygen, sulfur or NR₆;


[0015] n is an integer from 2-7; and

[0016] m is an integer from 1-4,


[0017] wherein at least one of R' , R_3 , R_4 , R_5 and R_6 is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0018] In some embodiments, the compound of Formula (I) is a compound of the Formula (IV):




[0019] wherein R_{10} is (C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0020] A is a pharmaceutically acceptable anion and


[0021] wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_8 , R_9 , X, Y, Z, W, and m are defined herein.

[0022] In some embodiments, the compound of Formula (I) is a compound of the formula:



[0023] wherein $R_1, R_{10}, R_3, R_4, R_5, R_5, R_8, R_9, X, Y, Z,$ W, and m are defined herein.

[0024] In some embodiments, the present disclosure provides a compound of Formula (III):



[0025] or a pharmaceutically acceptable salt thereof;

[0026] wherein,


[0027] R_1 and R_2 are independently hydrogen, alkyl, cycloalkyl or R_1 and R_2 together with the atoms to which they are attached form a heterocycle;

[0028] X, Y and Z are independently hydrogen, halogen, $-O-$, $-S-$, $-NR_A-$, or $(P=O)-OR_A(-OR')$,

[0029] wherein R_A is hydrogen or alkyl,

[0030] $R', R_3, R_4,$ and R_5 are independently absent, hydrogen, alkyl, $-(C=O)-alkyl,$ $-(C=O)CHR_7(NH_2),$ $-(C=O)(CH_2)_n(NH_2),$ $-(C=O)CH(NH_2)(CH_2)_n(NH_2),$ $-(C=O)CH(NH_2)$

[0031] $(CH_2)_n(NH)(C=NH)-NH_2,$ $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0032] R_7 and R_8 are independently hydrogen or alkyl;

[0033] R_9 is hydrogen, alkyl, or halogen;


[0034] L and M are independently alkyl or aryl;

[0035] n is an integer from 2-7; and


[0036] m is an integer from 1-4,

[0037] wherein at least one of $R', R_3, R_4,$ and R_5 is $-(C=O)-alkyl,$ $-(C=O)CHR_7(NR_2),$ $-(C=O)$


$(CH_2)_n(NH_2),$ $(NH_2),$ $-(C=O)CH((NH_2)(CH_2)_n(NH_2),$ $-(C=O)(CH_2)_m(NH)(C=NH)-NH_2$ or



[0038] In some embodiments, the compound of Formula (V) is a compound of the formula:



[0039] wherein R_{10} is $-(C=O)-,$ $-(C=O)CHR_7(NH_2),$ $-(C=O)(CH_2)_n(NH_2),$ $-(C=O)CH(NH_2)(CH_2)_n(NH_2),$ $-(C=O)CH(NH_2)(CH_2)_n(NH)(C=NH)-NH_2,$ $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0040] A is a pharmaceutically acceptable anion,

[0041] and wherein $R_1, R_2, R_3, R_4, R_5, R_8, R_9, X, Y, Z,$ and m are defined herein.

[0042] In some embodiments, the present disclosure provides a pharmaceutical composition comprising a compound of Formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.

[0043] In some embodiments, the present disclosure provides a pharmaceutical composition comprising a compound of Formula (II) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.

BRIEF DESCRIPTION OF THE DRAWINGS

[0044] FIGS. 1A-1C show PK studies of N-phosphonoxyethyl produg 6-1-2 (VLS-02-23-0) and metabolite (VLS-02-023-10) following intravenous (1 mg/kg) and oral administration (30 mg/kg) to male CD1 mice (in plasma).

[0045] FIGS. 2A-2B show PK studies of N-phosphonoxyethyl produg 6-1-2 (VLS-02-23-0) and metabolite

(OU LS-02-023-10) following intravenous (1 mg/kg) and oral administration (30 mg/kg) to male CD1 mice (in the brain).

DETAILED DESCRIPTION

[0046] Throughout this disclosure, various patents, patent applications and publications are referenced. The disclosures of these patents, patent applications and publications in their entireties are incorporated into this disclosure by reference for all purposes in order to more fully describe the state of the art as known to those skilled therein as of the date of this disclosure. This disclosure will govern in the instance that there is any inconsistency between the patents, patent applications and publications cited and this disclosure.

Definitions

[0047] For convenience, certain terms employed in the specification, examples and claims are collected here. Unless defined otherwise, all technical and scientific terms used in this disclosure have the same meanings as commonly understood by one of ordinary skill in the art to which this disclosure belongs.

[0048] The term “about” when immediately preceding a numerical value means a range (e.g., plus or minus 10% of that value). For example, “about 50” can mean 45 to 55, “about 25,000” can mean 22,500 to 27,500, etc., unless the context of the disclosure indicates otherwise, or is inconsistent with such an interpretation. For example in a list of numerical values such as “about 49, about 50, about 55, . . .”, “about 50” means a range extending to less than half the interval(s) between the preceding and subsequent values, e.g., more than 49.5 to less than 50.5. Furthermore, the phrases “less than about” a value or “greater than about” a value should be understood in view of the definition of the term “about” provided herein. Similarly, the term “about” when preceding a series of numerical values or a range of values (e.g., “about 10, 20, 30” or “about 10-30”) refers, respectively to all values in the series, or the endpoints of the range.

[0049] The terms “administer,” “administering” or “administration” as used herein refer to administering a compound or pharmaceutically acceptable salt of the compound or a composition or formulation comprising the compound or pharmaceutically acceptable salt of the compound to a patient.

[0050] The term “pharmaceutically acceptable salts” includes both acid and base addition salts. Pharmaceutically acceptable salts include those obtained by reacting the active compound functioning as a base, with an inorganic or organic acid to form a salt, for example, salts of hydrochloric acid, sulfuric acid, phosphoric acid, methanesulfonic acid, camphorsulfonic acid, oxalic acid, maleic acid, succinic acid, citric acid, formic acid, hydrobromic acid, benzoic acid, tartaric acid, fumaric acid, salicylic acid, mandelic acid, carbonic acid, etc. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, i.e., salts containing pharmaceutically acceptable anions, including but not limited to malate, oxalate, chloride, bromide, iodide, nitrate, acetate, tartrate, oleate, fumarate, formate, benzoate, glutamate, methanesulfonate, benzenesulfonate, and p-toluenesulfonate salts. Base addition salts include but are not limited to, ethylenediamine,

N-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris-(hydroxymethyl)-aminomethane, tetramethylammonium hydroxide, triethylamine, dibenzylamine, ephenamine, dehydroabietylamine, N-ethylpiperidine, benzylamine, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, ethylamine, basic amino acids, e.g., lysine and arginine dicyclohexylamine and the like. Examples of metal salts include lithium, sodium, potassium, magnesium, calcium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methylammonium, dimethylammonium, trimethylammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like. Examples of organic bases include lysine, arginine, guanidine, diethanolamine, choline and the like. Those skilled in the art will further recognize that acid addition salts may be prepared by reaction of the compounds with the appropriate inorganic or organic acid via any of a number of known methods.

[0051] The term “treating” as used herein with regard to a patient, refers to improving at least one symptom of the patient’s disorder. In some embodiments, treating can be improving, or at least partially ameliorating a disorder or one or more symptoms of a disorder.

[0052] The term “preventing” as used herein with regard to a patient or subject, refers to preventing the onset of disease development if none had occurred, preventing the disease or disorder from occurring in a subject or a patient that may be predisposed to the disorder or disease but has not yet been diagnosed as having the disorder or disease, and/or preventing further disease/disorder development if already present.

[0053] The term “therapeutically effective” applied to dose or amount refers to that quantity of a compound or pharmaceutical formulation that is sufficient to result in a desired clinical benefit after administration to a patient in need thereof.

[0054] When a range of values is listed, it is intended to encompass each value and sub-range within the range. For example, “C₁-C₆ alkyl” is intended to encompass C₁, C₂, C₃, C₄, C₅, C₆, C₁₋₆, C₁₋₅, C₁₋₄, C₁₋₃, C₁₋₂, C₂₋₆, C₂₋₅, C₂₋₄, C₂₋₃, C₃₋₆, C₃₋₅, C₃₋₄, C₄₋₆, C₄₋₅, and C₅₋₆ alkyl.

[0055] “Alkyl” or “alkyl group” refers to a fully saturated, straight or branched hydrocarbon chain having from one to twelve carbon atoms, and which is attached to the rest of the molecule by a single bond. Alkyls comprising any number of carbon atoms from 1 to 12 are included. An alkyl comprising up to 12 carbon atoms is a C₁-C₁₂ alkyl, an alkyl comprising up to 10 carbon atoms is a C₁-C₁₀ alkyl, an alkyl comprising up to 6 carbon atoms is a C₁-C₆ alkyl and an alkyl comprising up to 5 carbon atoms is a C₁-C₅ alkyl. A C₁-C₅ alkyl includes C₅ alkyls, C₄ alkyls, C₃ alkyls, C₂ alkyls and C₁ alkyl (i.e., methyl). A C₁-C₆ alkyl includes all moieties described above for C₁-C₅ alkyls but also includes C₆ alkyls. A C₁-C₁₀ alkyl includes all moieties described above for C₁-C₅ alkyls and C₁-C₆ alkyls, but also includes C₇, C₈, C₉ and C₁₀ alkyls. Similarly, a C₁-C₁₂ alkyl includes all the foregoing moieties, but also includes C₁₁ and C₁₂ alkyls. Non-limiting examples of C₁-C₁₂ alkyl include methyl, ethyl, n-propyl, i-propyl, sec-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, n-pentyl, n-amyl, n-hexyl, n-hep-

tyl, n-octyl, n-nonyl, n-decyl, n-undecyl, and n-dodecyl. Unless stated otherwise specifically in the specification, an alkyl group can be optionally substituted.

[0056] “Aryl” refers to a hydrocarbon ring system comprising hydrogen, 6 to 18 carbon atoms and at least one aromatic ring, and which is attached to the rest of the molecule by a single bond. For purposes of this disclosure, the aryl can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused or bridged ring systems. Aryls include, but are not limited to, aryls derived from acenaphthylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, as-indacene, s-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and triphenylene. Unless stated otherwise specifically in the specification, the “aryl” can be optionally substituted.

[0057] “Cycloalkyl” refers to a stable non-aromatic monocyclic or polycyclic fully saturated hydrocarbon consisting solely of carbon and hydrogen atoms, which can include fused, bridged, or spirocyclic ring systems, having from three to twenty carbon atoms (e.g., having from three to ten carbon atoms) and which is attached to the rest of the molecule by a single bond. Monocyclic cycloalkyls include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyls include, for example, adamantyl, norbornyl, decalyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, a cycloalkyl group can be optionally substituted.

[0058] “Heterocyclyl,” “heterocyclic ring” or “heterocycle” refers to a stable saturated, unsaturated, or aromatic 3- to 20-membered ring which consists of two to nineteen carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, and which is attached to the rest of the molecule by a single bond. Heterocyclyl or heterocyclic rings include heteroaryls, heterocyclalalkyls, heterocyclalalkenyls, and heterocyclalalkynyls. Unless stated otherwise specifically in the specification, the heterocyclyl can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused, bridged, or spirocyclic ring systems; and the nitrogen, carbon or sulfur atoms in the heterocyclyl can be optionally oxidized; the nitrogen atom can be optionally quaternized; and the heterocyclyl can be partially or fully saturated. Examples of such heterocyclyl include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, and 1,1-dioxo-thiomorpholinyl. Unless stated otherwise specifically in the specification, a heterocyclyl group can be optionally substituted.

[0059] “Heteroaryl” refers to a 5- to 20-membered ring system comprising hydrogen atoms, one to nineteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, at least one aromatic ring, and which is attached to the rest of the molecule by a single bond. For purposes of this disclosure, the heteroaryl can be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which can include fused or bridged

ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl can be optionally oxidized; the nitrogen atom can be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzothiazolyl, benzindolyl, benzodioxolyl, benzofuranlyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranlyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranlyl, benzopyranonyl, benzofuranlyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl, dibenzofuranlyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinazoliny, quinoxaliny, quinolinyl, quinuclidinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiazidazolyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (i.e. thienyl). Unless stated otherwise specifically in the specification, a heteroaryl group can be optionally substituted.

[0060] The term “substituted” used herein means any of the groups described herein (e.g., alkyl, alkenyl, alkynyl, alkoxy, aryl, cycloalkyl, cycloalkenyl, haloalkyl, heterocyclyl, and/or heteroaryl) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atoms such as, but not limited to: a halogen atom such as F, Cl, Br, and I; an oxygen atom in groups such as hydroxyl groups, alkoxy groups, and ester groups; a sulfur atom in groups such as thiol groups, thioalkyl groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, alkylamines, dialkylamines, arylamines, alkylaryl amines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as trialkylsilyl groups, dialkylarylsilyl groups, alkyl diarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. “Substituted” also means any of the above groups in which one or more hydrogen atoms are replaced by a higher-order bond (e.g., a double- or triple-bond) to a heteroatom such as oxygen in oxo, carbonyl, carboxyl, and ester groups; and nitrogen in groups such as imines, oximes, hydrazones, and nitriles. For example, “substituted” includes any of the above groups in which one or more hydrogen atoms are replaced with $-\text{NR}_g\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{NR}_g\text{C}(=\text{O})\text{OR}_h$, $-\text{NR}_g\text{SO}_2\text{R}_h$, $-\text{OC}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{OR}_g$, $-\text{SR}_g$, $-\text{SOR}_g$, $-\text{SO}_2\text{R}_g$, $-\text{OSO}_2\text{R}_g$, $-\text{SO}_2\text{NR}_g$, $-\text{NSO}_2\text{R}_g$, and $-\text{SO}_2\text{NR}_g\text{R}_h$. “Substituted” also means any of the above groups in which one or more hydrogen atoms are replaced with $-\text{C}(=\text{O})\text{R}_g$, $-\text{C}(=\text{O})\text{OR}_g$, $-\text{C}(=\text{O})\text{NR}_g\text{R}_h$, $-\text{CH}_2\text{SO}_2\text{R}_g$, $-\text{CH}_2\text{SO}_2\text{NR}_g\text{R}_h$. In the foregoing, R_g and R_h are the same or different and independently hydrogen, alkyl, alkenyl, alkynyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, haloalkyl, haloalkenyl, haloalkynyl, heterocyclyl, N-heterocyclyl, heterocyclalalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl. “Substituted” further means any of the above groups in which one or more hydrogen atoms are replaced by a bond to an amino, cyano, hydroxyl, imino, nitro, oxo, thioxo, halo, alkyl, alkenyl, alkynyl, alkoxy, alkylamino, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl,

haloalkyl, haloalkenyl, haloalkynyl, heterocyclyl, N-heterocyclyl, heterocyclylalkyl, heteroaryl, N-heteroaryl and/or heteroarylalkyl group. In addition, each of the foregoing substituents can also be optionally substituted with one or more of the above substituents.


Compounds

[0061] The present disclosure provides compounds that are prodrugs of tryptamines and relates scaffolds, for example N,N-dimethyltryptamine (DMT) as well as pharmaceutical compositions thereof.

[0062] DMT is metabolically unstable and is readily converted by monoamine oxidases (MAO's) to indoleacetic acid and N-oxidation metabolites resulting in poor oral bioavailability. In some embodiments, the compounds of the present disclosure allow for the controlled release of DMT. Advantages of compounds of the present disclosure may include increased metabolic stability, increased absorption, decreased maximal plasma concentrations of parent drug DMT over time, and less frequent dosing. In some embodiments, compounds of the present disclosure prevent or inhibit N-oxidation to promote oral bioavailability and increased exposure.

[0063] In some embodiments, compounds of the present disclosure comprise an enacarbil moiety. Without being bound by theory such compounds may for example, increase bioavailability by virtue of active transport of the prodrug in the small intestine by high-capacity nutrient transporters, including mono-carboxylate transporter-1 (MCT-1). DMT-enacarbil prodrugs may also reduce abuse potential by preventing absorption by snorting or insufflation.

[0064] In some embodiments, the present disclosure provides a compound of Formula (I):




[0065] or a pharmaceutically acceptable salt thereof;

[0066] wherein,

[0067] R_1 and R_2 are independently hydrogen, alkyl, cycloalkyl, —or R_1 and R_2 together with the atoms to which they are attached form a heterocycle;

[0068] X, Y and Z are independently hydrogen, halogen, —O—, —S—, —NR_A—, or —(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl;

[0069] R', R₃, R₄, R₅ and R₆ are independently absent, hydrogen, alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂) (CH₂)_n(NH)(C=NH)—NH₂, or —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0070] R₇ and R₈ are independently hydrogen or alkyl;

[0071] R₉ is hydrogen, alkyl, or halogen;


[0072] L and M are independently alkyl or aryl;

[0073] W is oxygen, sulfur or NR₆;


[0074] n is an integer from 2-7; and

[0075] m is an integer from 1-4,

[0076] wherein at least one of R', R₃, R₄, R₅ and R₆ is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), or —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0077] In some embodiments, provided herein is a compound of Formula (II)



[0078] or a pharmaceutically acceptable salt thereof;

[0079] wherein,

[0080] R₁ and R₂ are independently hydrogen, alkyl, cycloalkyl, —or R₁ and R₂ together with the atoms to which they are attached form a heterocycle;

[0081] X, Y and Z are independently hydrogen, halogen, —O—, —S—, —NR_A—, or (P=O)—OR_A(—OR'),

[0082] wherein R_A is hydrogen or alkyl;

[0083] R', R₃, R₄, R₅ and R₆ are independently absent, hydrogen, alkyl, —(C=O)-alkyl, (C=O)CHR₇, —(C=O)(CH₂)_n(NH₂), —(C)CH(NH₂)(CH₃), —(C=O)CH(NH₂), (CH₂)_n(NH)(C=NH), or —(C=O)(CH₂)_n(NH)(C=NH)—NH₂;

[0084] R₇ and R₈ are independently hydrogen or alkyl;

[0085] R₉ is hydrogen, alkyl, or halogen;


[0086] L and M are independently alkyl or aryl;

[0087] W is oxygen, sulfur or NR₆;

[0088] n is an integer from 2-7; and


[0089] m is an integer from 1-4,

[0090] wherein at least one of R', R₃, R₄, and R₅ and R₆ is $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0091] In some embodiments, R₉ is a halogen. In some embodiments, R₉ is bromo.

[0092] In some embodiments, provided herein is a compound of Formula (III):




[0093] or a pharmaceutically acceptable salt thereof;

[0094] wherein.

[0095] R₁ and R₂ are independently hydrogen, alkyl, cycloalkyl, —or R₁ and R₂ together with the atoms to which they are attached form a heterocycle;

[0096] X, Y and Z are independently hydrogen, halogen, —O—, —S—, —NR_A—, —O(P=O)—OR_A(—OR'), or —(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl,

[0097] R', R₃, R₄, and R₅ are independently absent, hydrogen, alkyl, $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH)(C=NH)-NH_2$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0098] R₇ and R₈ are independently hydrogen or alkyl;


[0099] R₉ is hydrogen, alkyl, or halogen;

[0100] L and M are independently alkyl or aryl;

[0101] n is an integer from 2-7; and

[0102] m is an integer from 1-4,

[0103] wherein at least one of R', R₃, R₄, and R₅ is $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0104] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are independently hydrogen, alkyl, or cycloalkyl, —or R₁ and R₂ together with the atoms to which they are attached form a heterocycle.

[0105] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are independently hydrogen, alkyl, or cycloalkyl.

[0106] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ together with the atoms to which they are attached form a heterocycle.

[0107] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are independently hydrogen or C₁-C₆ alkyl.

[0108] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are independently C₁-C₆ alkyl.

[0109] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are independently methyl, ethyl, propyl, and isopropyl.

[0110] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₁ and R₂ are methyl.

[0111] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), X, Y and Z are independently hydrogen, halogen, —O—, —S—, —NR_A—, —O(P=O)—OR_A(—OR'), or —(P=O)—OR_A(—OR').

[0112] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), X, Y and Z are H.

[0113] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), two of X, Y and Z are hydrogen.

[0114] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), two of X, Y and Z are hydrogen and the non-hydrogen X, Y or Z is —O—, —S—, —NR_A—, —O(P=O)—OR_A(—OR'), or —(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0115] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), X and Y are hydrogen; and Z is —O—, —S—, —NR_A—, —O(P=O)—OR_A(—OR'), or —(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0116] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), X and Z are hydrogen; and Y is —O—, —S—, —NR_A—, —O(P=O)—OR_A(—OR'), or —(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0117] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), Y and Z are

hydrogen; and X is —O—, —S—, —NR₄—, —O(P=O)—OR₄(—OR'), or —(P=O)—OR₄(—OR'), wherein R₄ is hydrogen or alkyl.

[0118] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), at least one of X, Y, or Z is —O(P=O)—OR₄(—OR').

[0119] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), X is —O(P=O)—OR₄(—OR') and R₃ is absent.


[0120] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), Y is —O(P=O)—OR₄(—OR') and R₄ is absent.

[0121] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), Z is —O(P=O)—OR₄(—OR') and R₅ is absent.


[0122] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R' is hydrogen or alkyl.

[0123] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₄ is hydrogen or alkyl.


[0124] In some embodiments of the compounds of Formula (I) or Formula (II), R', R₃, R₄, R₅ and R₆ are independently absent, hydrogen alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0125] wherein at least one of R', R₃, R₄, R₅ and R₆ is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0126] In some embodiments of the compounds of Formula (III), R', R₃, R₄, and R₅ are independently absent, hydrogen, alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, or —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



wherein at least one of R', R₃, R₄, and R₅ is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0127] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R' is absent, hydrogen, alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, or —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0128] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R' is hydrogen or alkyl.

[0129] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₃ is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0130] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R₄ is: —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or




[0131] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_5 is:


$-(C=O)$ -alkyl, $-CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0132] In some embodiments of the compounds of Formula (I), R_6 is: $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0133] In some embodiments of the compounds of Formula (I), X, Y and Z are H; W is NR_6 ; and R_6 is $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or




[0134] In some embodiments of the compounds of Formula (I), at least one of R' , R_3 , R_4 , R_5 and R_6 is $-(C=O)(CH_2)_n(NH_2)$ and n is 3.

[0135] In some embodiments of the compounds of Formula (I), at least one of R' , R_3 , R_4 , R_5 and R_6 is $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, and n is 4.

[0136] In some embodiments of the compounds of Formula (I), at least one of R' , R_3 , R_4 , R_5 and R_6 is $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ and n is 3.

[0137] In some embodiments of the compounds of Formula (I), at least one of R' , R_3 , R_4 , R_5 and R_6 is $-(C=O)CH(NH)(CH_2)_n(C=NH)-NH_2$ and n is 2.

[0138] In some embodiments of the compounds of Formula (I), at least one of R' , R_3 , R_4 , R_5 and R_6 is




[0139] In some embodiments of the compounds of Formula (III), at least one of R' , R_3 , R_4 , and R_5 is $-(C=O)(CH_2)_n(NH_2)$ and n is 3.

[0140] In some embodiments of the compounds of Formula (III), at least one of R' , R_3 , R_4 , and R_5 is $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$ and n is 4.

[0141] In some embodiments of the compounds of Formula (III), at least one of R' , R_3 , R_4 , and R_5 is $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ and n is 3.

[0142] In some embodiments of Formula (III), at least one of R' , R_3 , R_4 , and R_5 is $-(C=O)CH(NH_2)(CH_2)_n(NH)(C=NH)-NH_2$ and n is 2.

[0143] In some embodiments of Formula (III), at least one of R' , R_3 , R_4 , and R_5 is



[0144] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), L and M are independently alkyl or aryl.

[0145] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), L is alkyl and M is alkyl or aryl.

[0146] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), is methyl or isopropyl and M is alkyl or aryl.

[0147] In some embodiments of the compounds of Formula (I) or Formula (II), W is oxygen, sulfur or NR_6 ;

[0148] In some embodiments of the compounds of Formula (I) or Formula (II), W is NR_6 and R_6 is hydrogen or alkyl.

[0149] In some embodiments of the compounds of Formula (I) or Formula (II), R_7 , R_8 , and R_9 are independently hydrogen or alkyl.

[0150] In some embodiments of the compounds of Formula (III), R_8 , and R_9 are independently hydrogen or alkyl.

[0151] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_8 is hydrogen.

[0152] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_8 is alkyl.

[0153] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is C_1 - C_6 alkyl.

[0154] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is hydrogen, halogen, or alkyl.

[0155] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is hydrogen.

[0156] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is alkyl.

[0157] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is C_1 - C_6 alkyl.

[0158] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is halogen. In some embodiments, R_9 is $-F$, $-Cl$, $-Br$, or $-I$. In some embodiments, R_9 is $-Cl$, $-Br$, or $-I$. In some embodiments, R_9 is $-Cl$ or $-Br$. In some embodiments, R_9 is $-F$. In some embodiments, R_9 is $-Cl$. In some embodiments, R_9 is $-Br$. In some embodiments, R_9 is 4.

[0159] In some embodiments of the compounds of Formula (I) or Formula (II), R_7 and R_8 are independently hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.

[0160] In some embodiments of the compounds of Formula (III), R_8 is hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.

[0161] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), R_9 is hydrogen, halogen, or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.


[0162] In some embodiments of the compounds of Formula (I) or Formula (II), R_7 , R_8 and R_9 are independently hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.

[0163] In some embodiments of the compounds of Formula (III), R_8 and R_9 are independently hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.


[0164] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), n is an integer from 2-7 (i.e. 2, 3, 4, 5, 6, or 7). In some embodiments, n is 2. In some embodiments, n is 3. In some embodiments, n is 4. In some embodiments, n is 5. In some embodiments, n is 6. In some embodiments, n is 7.

[0165] In some embodiments of the compounds of Formula (I), Formula (II), or Formula (III), m is an integer from 1-4 (i.e. 1, 2, 3, or 4). In some embodiments, m is 1. In some embodiments, m is 2. In some embodiments, m is 3. In some embodiments, m is 4.

[0166] In some embodiments of the compounds of Formula (I), the compound is of Formula (IV):




[0167] wherein R_{10} is $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or




[0168] A is a pharmaceutically acceptable anion,

[0169] and wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_8 , R_9 , X , Y , Z , W , and m are defined herein.

[0170] In some embodiments, the compound of Formula (I) is a compound of Formula (V):




[0171] wherein R_{10} is $-(C=O)$ -alkyl, $-(C=O)CHR_7(NH_2)$, $-(C=O)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH_2)$, $-(C=O)CH(NH_2)(CH_2)_n(NH)(C=NH)-NH_2$, $(C=O)(CH_2)_n(NH)(C=NH)-NH_2$ or



[0172] A is a pharmaceutically acceptable anion,

[0173] and wherein R_1 , R_2 , R_3 , R_4 , R_5 , R_8 , R_9 , Y , Z , and W are defined herein.

[0174] In some embodiments, the present disclosure provides compounds of Formula (VI):




[0175] wherein R_1 , R_2 , R_3 , R_4 , and R_5 are as defined herein,

[0176] X is carbon or nitrogen;

[0177] W is carbon, oxygen, or nitrogen; and


[0178] Z is hydrogen, deuterium, or oxygen.

[0179] In some embodiments, the compound of Formula (VII) is:




(VII)


[0180] In some embodiments, the compound of Formula (VI) or (VII) is:




1-2-1




1-2-2



1-2-3




1-2-4




1-2-5


-continued




1-2-6




1-2-7




1-2-8




1-2-9



1-2-10




1-2-11




1-2-12

-continued




1-2-13



1-2-14

[0181] In some embodiments, the present disclosure provides compounds of Formula (VII):



(VIII)


wherein R_1 , R_2 , R_3 , R_4 , R_5 , and R_9 , are as defined herein,

[0182] R_{11} and R_{12} are independently a pharmaceutically acceptable salt, hydrogen, deuterium, alkyl, substituted alkyl, cycloalkyl, —or R_{11} and R_{12} together with the atoms to which they are attached form an alkyl, cycloalkyl, heterocycles and are ether, hemiaminal ketal and amino acid.

[0183] In some embodiments of the compounds of Formula (VIII), R_{11} and R_{12} together with the atoms to which they are attached form a heterocycle.

[0184] In some embodiments of the compounds of Formula (VIII), the pharmaceutically acceptable salt is Na^+ , K^+ , or NH_4^+ .

[0185] In some embodiments, provided herein is a compound of Formula (IX):




(IX)

or a pharmaceutically acceptable salt thereof; wherein,

R_1 and R_2 are each independently hydrogen, alkyl, —(C=O)—alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH

(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n, (C=NH)—NH₂, —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



X, Y and Z are each independently hydrogen, halogen, —O—, —S—, —NR_A—, or —O(P=O)—OR_A(—OR'), wherein R' and R_A is hydrogen or alkyl;

R_3 , R_4 , R_5 and R_6 are each independently absent, hydrogen, or alkyl;

R_7 and R_8 are each independently hydrogen or alkyl;

R_9 is hydrogen, alkyl, or halogen;


L and M are each independently alkyl or aryl;

W is oxygen, sulfur or NR₆;

n is an integer from 2-7; and

m is an integer from 1-4,

wherein at least one of R_1 and R_2 is —(C=O)—alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0186] In some embodiments of the compounds of Formula (IX), X, Y and Z are each independently hydrogen, halogen, —O—, —S—, —NR_A—, or —O(P=O)—OR_A(—OR').

[0187] In some embodiments of the compounds of Formula (IX), X, Y and Z are hydrogen.

[0188] In some embodiments of the compounds of Formula (IX), two of X, Y and Z are hydrogen.

[0189] In some embodiments of the compounds of Formula (IX), two of X, Y and Z are hydrogen and the non-hydrogen X, Y or Z is —O—, —S—, NR_A—, or —O(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0190] In some embodiments of the compounds of Formula (IX), X and Y are hydrogen; and Z is —O—, —S—, —NR_A—, or —O(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0191] In some embodiments of the compounds of Formula (IX), X and Z are hydrogen; and Y is —O—, —S—, —NR_A—, or —O(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0192] In some embodiments of the compounds of Formula (IX), Y and Z are hydrogen; and X is —O—, —S—, —NR_A—, or —O(P=O)—OR_A(—OR'), wherein R_A is hydrogen or alkyl.

[0193] In some embodiments of the compounds of Formula (IX), at least one of X, Y, or Z is $-\text{O}(\text{P}=\text{O})-\text{OR}'_4$ ($-\text{OR}'$).

[0194] In some embodiments of the compounds of Formula (IX), X is $-\text{O}(\text{P}=\text{O})-\text{OR}'_4$ ($-\text{OR}'$) and R_3 is absent.


[0195] In some embodiments of the compounds of Formula (IX), Y is $-\text{O}(\text{P}=\text{O})-\text{OR}'_4$ ($-\text{OR}'$) and R_4 is absent.

[0196] In some embodiments of the compounds of Formula (IX), Z is $-\text{O}(\text{P}=\text{O})-\text{OR}'_4$ ($-\text{OR}'$) and R_5 is absent.


[0197] In some embodiments of the compounds of Formula (IX), R' is hydrogen or alkyl.

[0198] In some embodiments of the compounds of Formula (IX), R'_4 is hydrogen or alkyl.


[0199] In some embodiments of the compounds of Formula (IX), R_1 and R_2 are each independently absent, hydrogen, alkyl, $-(\text{C}=\text{O})$ -alkyl, $-(\text{C}=\text{O})\text{CHR}_7(\text{NH}_2)$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$, $(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ or



wherein at least one of R_1 and R_2 is $-(\text{C}=\text{O})$ -alkyl, $-(\text{C}=\text{O})\text{CHR}_7(\text{NH}_2)$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ or




[0200] In some embodiments of the compounds of Formula (IX), R_1 is absent, hydrogen, alkyl, $-(\text{C}=\text{O})$ -alkyl, $-(\text{C}=\text{O})\text{CHR}_7(\text{NH}_2)$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ or



[0201] In some embodiments of the compounds of Formula (IX), R^1 is hydrogen or alkyl.

[0202] In some embodiments of the compounds of Formula (IX), R_2 is absent, hydrogen, alkyl, $-(\text{C}=\text{O})$ -alkyl, $-(\text{C}=\text{O})\text{CHR}_7(\text{NH}_2)$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})$

$\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH}_2)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ or



[0203] In some embodiments of the compounds of Formula (IX), R_2 is hydrogen or alkyl.

[0204] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH}_2)$ and n is 3.


[0205] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH}_2)$ and n is 4.

[0206] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ and n is 3.

[0207] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})\text{CH}(\text{NH})(\text{CH}_2)_n(\text{C}=\text{NH})-\text{NH}_2$ and n is 2.

[0208] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})$ -alkyl.

[0209] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is



[0210] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ and n is 4.

[0211] In some embodiments of the compounds of Formula (IX), at least one of R_1 and R_2 is $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})-\text{NH}_2$ and n is 3.

[0212] In some embodiments of the compounds of Formula (IX), L and M are each independently alkyl or aryl.

[0213] In some embodiments of the compounds of Formula (IX), L is alkyl and M is alkyl or aryl.

[0214] In some embodiments of the compounds of Formula (IX), L is methyl or isopropyl and M is alkyl or aryl.

[0215] In some embodiments, R_3 is hydrogen.

[0216] In some embodiments, R_3 is alkyl.

[0217] In some embodiments, R_3 is C_1 - C_6 alkyl.

[0218] In some embodiments, R_4 is absent.

[0219] In some embodiments, R_4 is hydrogen.

[0220] In some embodiments, R_4 is alkyl.

[0221] In some embodiments, R_4 is C_1 - C_6 alkyl.


[0222] In some embodiments, R_1 is absent.

[0223] In some embodiments, R_5 is hydrogen.

[0224] In some embodiments, R_5 is alkyl.

[0225] In some embodiments, R_5 is C_1 - C_6 alkyl.

- [0226] In some embodiments, R_5 is absent.
- [0227] In some embodiments of the compounds of Formula (IX), W is oxygen, sulfur or NR_6 .
- [0228] In some embodiments of the compounds of Formula (IX), W is NR_6 and R_6 is hydrogen or alkyl.
- [0229] In some embodiments of the compounds of Formula (IX), R_7 , R_8 , and R_9 are each independently hydrogen or alkyl.
- [0230] In some embodiments of the compounds of Formula (IX), R_8 is hydrogen.
- [0231] In some embodiments of the compounds of Formula (IX), R_8 is alkyl.
- [0232] In some embodiments of the compounds of Formula (IX), R_8 is C_1 - C_6 alkyl.
- [0233] In some embodiments of the compounds of Formula (IX), R_9 is hydrogen, halogen, or alkyl.
- [0234] In some embodiments of the compounds of Formula (IX), R_9 is hydrogen.
- [0235] In some embodiments of the compounds of Formula (IX), R_9 is alkyl.
- [0236] In some embodiments of the compounds of Formula (IX), R_9 is C_1 - C_6 alkyl.
- [0237] In some embodiments of the compounds of Formula (IX), R_9 is halogen. In some embodiments, R_9 is —F, —Cl, —Br, or in some embodiments, R_9 is —Cl, —Br, or —I. In some embodiments, R_9 is —Cl or —Br. In some embodiments, R_9 is —F. In some embodiments, R_9 is —Cl. In some embodiments, R_9 is —Br. In some embodiments, R_9 is
- [0238] In some embodiments of the compounds of Formula (IX), R_7 and R_8 are each independently hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.
- [0239] In some embodiments of the compounds of Formula (IX), R_9 is hydrogen, halogen, or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.
- [0240] In some embodiments of the compounds of Formula (IX), R_7 , R_8 and R_9 are each independently hydrogen or C_1 - C_8 alkyl optionally substituted with 1-3 groups independently selected from hydroxyl, thiol and halogen.
- [0241] In some embodiments of the compounds of Formula (IX), n is an integer from 2-7 (i.e. 2, 3, 4, 5, 6, or 7). In some embodiments n is 2. In some embodiments, n is 3. In some embodiments, n is 4. In some embodiments, n is 5. In some embodiments, n is 6. In some embodiments, n is 7.
- [0242] In some embodiments of the compounds of Formula (IX), m is an integer from 1-4 (i.e. 1, 2, 3, or 4). In some embodiments, m is 1. In some embodiments, m is 2. In some embodiments, M is 3. In some embodiments, m is 4.
- [0243] In some embodiments, provided herein is a compound of Formula (X):
wherein




(X)


R_1 and R_2 together with the atoms to which they are attached form a heterocycle;

X and Y are each independently hydrogen, halogen, —O—, —S—, — NR_4 —, or —O(P=O)— OR_4 (— OR'), wherein R_4 is hydrogen or alkyl; and

R' , R_3 and R_4 are each independently absent, hydrogen, alkyl, —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(MH₂), —(C=O)CH(NH₂)(CH₂)_n(NH)(C=NH)—NH₂, —(C=O)(CH₂)_n(C=NH)—NH₂ or



wherein at least one of R' , R_3 , and R_4 is —(C=O)-alkyl, —(C=O)CHR₇(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)(CH₂)_n(NH₂), —(C=O)CH(NH₂)(CH₂)_n(NH), —(C=O)(CH₂)_n(NH)(C=NH)—NH₂ or



[0244] In some embodiments, provided herein is one or more compounds selected from Table 1.

[0245] In some embodiments, provided herein is one or more pharmaceutically acceptable salts of a compound selected from Table 1.

TABLE 1

Compounds	
No.	Structure
1-1	

TABLE 1-continued





Compounds	
No.	Structure
1-2	
1-3	
1-4	
1-5	

TABLE 1-continued









Compounds	
No.	Structure
1-6	
1-7	
1-8	
1-9	

TABLE 1-continued

Compounds	
No.	Structure
1-10	
1-11	
1-12	
1-13	

[0246] In some embodiments, provided herein is one or more compounds selected from Table 2.

[0247] In some embodiments, provided her is one or more pharmaceutically acceptable salts of a compound selected from Table 2.

TABLE 2






Compounds	
No.	Structure
2-1	
2-2	
2-3	
2-4	


TABLE 2-continued

Compounds	
No.	Structure


2-5



2-6



2-7



2-8






TABLE 2-continued

Compounds	
No.	Structure


2-9




2-10




2-11



2-12



2-13



[0248] In some embodiments of the compounds of Table 2, X and Y are hydrogen and R₃ and R₄ are absent.

[0249] In some embodiments of the compounds of Table 2, X is hydrogen, R is absent, Y is O, and R₄ is hydrogen.

[0250] In some embodiments of the compounds of Table 2, X is hydrogen, R₃ is absent, Y is O, and R₄ is Me.

[0251] In some embodiments of the compounds of Table 2, X is hydrogen, R₃ is absent, Y is O, and R₄ is Ac.

[0252] In some embodiments of the compounds of Table 2, X is O, R₃ is hydrogen, Y is hydrogen, and R₄ is absent.

[0253] In some embodiments of the compounds of Table 2, X is O, R₃ is Me, Y is hydrogen, and R₄ is absent.

[0254] In some embodiments of the compounds of Table 2, X is O, R₃ is Ac, Y is hydrogen, and R₄ is absent. In some embodiments, provided herein is one or more compounds selected from Table 3.

[0255] In some embodiments, provided herein is one or more pharmaceutically acceptable salts of a compound selected from Table 3.

TABLE 3

Compounds	
No.	Structure
3-1	
3-2	
3-3	
3-4	

wherein X, Y, R₃ and R₄ have the definitions provided herein.

[0256] In some embodiments of the compounds of Table 3, R₃ is absent, X is hydrogen, R₄ is absent, and Y is hydrogen.

[0257] In some embodiments of the compounds of Table 3, R₃ is absent, X is hydrogen, R₄ is hydrogen, and Y is —O—.

[0258] In some embodiments of the compounds of Table 3, R₃ is absent, X is hydrogen, R₄ is alkyl, and Y is —O—.

[0259] In some embodiments of the compounds of Table 3, R₃ is hydrogen, X is —O—, R₄ is absent hydrogen, and Y is hydrogen.

[0260] In some embodiments of the compounds of Table 3, R₃ is alkyl, X is —O—, R₄ is absent hydrogen, and Y is hydrogen.

[0261] In some embodiments of the compounds of Table 3, R₃ is absent, X is hydrogen, R₄ is absent, Y is —O(P=O)(OR₄)(OR'), R₄ is hydrogen or alkyl, and R' is hydrogen or alkyl.

[0262] In some embodiments of the compounds of Table 3, R₃ is absent, X is hydrogen, R₄ is absent, and Y is —O(P=O)(OH)(OH).

[0263] In some embodiments of the compounds of Table 3, R₃ is absent, X is —O(P=O)(OR₄)(OR'), R₄ is hydrogen or alkyl, R' is hydrogen or alkyl, R₄ is absent, and Y is hydrogen.

[0264] In some embodiments of the compounds of Table 3, R₃ is absent, X is —O(P=O)(OH)(OH), R₄ is absent, and Y is hydrogen.

[0265] In some embodiments, provided herein is one or more compounds selected from Table 4.

[0266] In some embodiments, provided herein is one or more pharmaceutically acceptable salts of a compound selected from Table 4.

TABLE 4

Compounds	
1-2-1	
1-2-2	
1-2-3	

TABLE 4-continued






Compounds	
1-2-4	 1-2-4
1-2-5	 1-2-5
1-2-6	 1-2-6
1-2-7	 1-2-7
1-2-8	 1-2-8

TABLE 4-continued








Compounds	
1-2-9	 1-2-9
1-2-10	 1-2-10
1-2-11	 1-2-11
1-2-12	 1-2-12
1-2-13	 1-2-13
1-2-14	 1-2-14

TABLE 4-continued


Compounds

2-1-2




2-1-2

3-1-2




3-1-2

3-1-3




3-1-3

3-1-4



3-1-4

4-1-1




4-1-1

TABLE 4-continued


Compounds

4-1-2




4-1-2

4-1-3




4-1-3

4-1-4



4-1-4

5-1-1



5-1-1

TABLE 4-continued







Compounds	
5-1-2	 5-1-2
5-1-3	 5-1-3
5-1-4	 5-1-4
6-1-1	 6-1-1
6-1-2	 6-1-2

TABLE 4-continued

Compounds	
6-1-3	 6-1-3

Compositions

[0267] In some embodiments of the present disclosure, a pharmaceutical composition comprises a therapeutically effective amount of one or more compounds of the present disclosure (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.

[0268] The pharmaceutically acceptable excipients and adjuvants are added to the composition or formulation for a variety of purposes. In some embodiments, a pharmaceutical composition comprising one or more compounds disclosed herein, or a pharmaceutically acceptable salt thereof, further comprise a pharmaceutically acceptable carrier. In some embodiments, a pharmaceutically acceptable carrier includes a pharmaceutically acceptable excipient, binder, and/or diluent. In some embodiments, suitable pharmaceutically acceptable carriers include, but are not limited to, inert solid fillers or diluents and sterile aqueous or organic solutions. In some embodiments, suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylase, magnesium stearate, talc, silicic acid, viscous paraffin, and the like.

[0269] For the purposes of this disclosure, the compounds of the present disclosure can be formulated for administration by a variety of means including orally, parenterally, by inhalation spray, topically, or rectally in formulations containing pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used here includes subcutaneous, intravenous, intramuscular, and intraarterial injections with a variety of infusion techniques. Intraarterial and intravenous injection as used herein includes administration through catheters.

Methods

[0270] In one aspect, the present disclosure provides methods of treating or preventing neurological disorders in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject.

[0271] In some embodiments, the neurological disorder is a mood disorder. In some embodiments, the mood disorder is clinical depression, postnatal depression or postpartum depression, perinatal depression, atypical depression, melancholic depression, psychotic major depression, catonic depression, seasonal affective disorder, dysthymia, double depression, depressive personality disorder, recurrent brief depression, major depressive disorder, minor depressive disorder, bipolar disorder or manic depressive disorder, depression caused by chronic medical conditions, treatment-resistant depression, refractory depression, suicidality, suicidal ideation, or suicidal behavior. In some embodiments, the method described herein provides therapeutic effect to a subject suffering from depression (e.g., moderate or severe depression). In some embodiments, the mood disorder is associated with neuroendocrine diseases and disorders, neurodegenerative diseases and disorders (e.g., epilepsy), movement disorders, tremor (e.g., Parkinson's Disease), or women's health disorders or conditions. In some embodiments, the mood disorder is depression. In some embodiments, the mood disorder is treatment-resistant depression or major depressive disorder. In some embodiments, the mood disorder is major depressive disorder. In some embodiments, the mood disorder is treatment-resistant depression.

[0272] In some embodiments, the present disclosure provides methods of treating or preventing PTSD, mood disorders, general anxiety disorder, addictive disorders, and/or drug dependence in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject.

[0273] In some embodiments, the present disclosure provides methods of treating or preventing PTSD in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject.

[0274] In some embodiments, the methods include treating PTSD through induction and maintenance therapy by administering a compound disclosed herein, a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof. In some embodiments, the compounds of the present disclosure are used for induction and maintenance therapy to treat PTSD with an improved safety profile when compared to treatment with the entactogenic, oneirophrenic or psychedelic compound (e.g., dimethyltryptamine or related compound, psilocybin, or MDMA) alone.

[0275] In some embodiments, the present disclosure provides methods of treating or preventing behavioral or mood disorders in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition

thereof to the subject, In some embodiments, the behavioral or mood disorder includes anxiety, such as social anxiety in autistic subjects (e.g., autistic adults) and anxiety related to life-threatening illnesses. In some embodiments, the behavioral or mood disorder includes stress (where moderation thereof is measured, e.g., by effects on amygdala responses). In some embodiments, the anxiety disorder is panic disorder, obsessive-compulsive disorder, and/or general anxiety disorder. In some embodiments, the subject suffers from a lack of motivation, attention, lack of accuracy in memory recall, speed of response, perseveration, and/or cognitive engagement. Further examples include depression (e.g., MDD or TRD), attention disorders, disorders of executive function and/or cognitive engagement, obsessive compulsive disorder, bipolar disorder, panic disorder, phobia, schizophrenia, psychopathy, antisocial personality disorder and/or neurocognitive disorders.

[0276] In some embodiments, the present disclosure provides methods for treating an addictive disorder in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject. In some embodiments, the addictive disorder is alcohol abuse, substance abuse, smoking, obesity, or mixtures thereof. In some embodiments, the disorder is an eating disorder (e.g., anorexia nervosa, bulimia, nervosa, binge eating disorder, etc.) or an auditory disorder.

[0277] In some embodiments, the present disclosure provides methods for treating an impulsive disorder in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject. In some embodiments, the impulsive disorder is attention deficit hyperactivity disorder (ADHD), attention deficit disorder (ADD), Tourette's syndrome, autism, or combinations thereof.

[0278] In some embodiments, the present disclosure provides methods for treating a compulsive disorder in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof to the subject. In some embodiments, the compulsive disorder is obsessive compulsive disorder (OCD), gambling, aberrant sexual behavior, or combinations thereof.

[0279] In some embodiments, the present disclosure provides methods for treating a personality disorder in a subject in need thereof, the methods comprising administering a therapeutically effective amount of a compound disclosed herein (e.g., a compound of Formulas (I)-(X) or Compounds of Tables 1-4), a pharmaceutically acceptable salt thereof, or a pharmaceutical composition thereof. In some embodi-

ments, the personality disorder is conduct disorder, antisocial personality, aggressive behavior, or combinations thereof to the subject.


EXAMPLES

[0280] Compounds of the present disclosure can be synthesized using the following exemplary methods or other methods that are known to those skilled in the art.

[0281] General reaction conditions are provided, and reaction products can be purified by known methods including silica gel chromatography using various organic solvents such as hexane, dichloromethane, ethyl acetate, methanol and the like or preparative reverse phase high pressure liquid chromatography.

[0282] Preparation of compounds can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Greene and Wuts, *Protective Groups in Organic Synthesis*, 44th, Ed., Wiley & Sons, 2006, as well as in Jerry March, *Advanced Organic Chemistry*, 4th edition, John Wiley & Sons, publisher, New York, 1992 which are incorporated herein by reference in their entirety.


Scheme 1: Representative synthesis of compounds of the disclosure



$n = 1-3$, $R = \text{H, NH}_2$, $R_{11} = \text{H, C} = \text{NH(NH}_2\text{)}$

[0283] As shown in Scheme 1, 2-(1H-indol-3-yl)-N,N-dimethylethan-1-amine can be coupled with a carboxylic acid of Formula (A) where PG_1 is a protecting group (such as Boc) under amide coupling conditions (e.g., in the presence of 1,1'-carbonyldiimidazole (CDI), dimethylaminopyridine (DMAP) and a strong base, such as sodium hydride), followed by deprotection of the protecting group PG_1 (e.g., using TFA) to provide compounds of Formula (I-a).


Scheme 2: Representative synthesis of compounds of the disclosure



$n = 1-3$, $R = \text{H, NH}_2$, $R_{11} = \text{H, C} = \text{NH(NH}_2\text{)}$

[0284] As shown in Scheme 2, 2-(1H-indol-3-yl)-N-methylethan-1-amine can be reacted with a carboxylic acid of Formula (A) under amide coupling conditions (e.g., in the presence of CDI, DMAP and a base, such as a tertiary amine base like triethylamine) where PG_1 is a protecting group (such as Boc) followed by deprotection of the protecting group PG_1 (e.g., using TFA) to provide compounds of Formula (I-b).


Scheme 3: Representative synthesis of compounds of the disclosure



[0285] As shown in Scheme 3, 3-(2-(dimethylamino)ethyl)-1H-indol-4-ol can be converted into 3-(2-(dimethylamino)ethyl)-1H-indol-4-yl dihydrogen phosphate in the presence of POCl_3 , 3-(2-(dimethylamino)ethyl)-1H-indol-4-yl dihydrogen phosphate can be reacted with a carboxylic acid of Formula (B) in the presence of CDI and a tertiary amine such as triethylamine to form a compound of Formula (Ic). Compounds of Formula (Ic) can be deprotected e.g., with TFA to provide compounds of Formula (I-d).

1. Synthesis of 1-2

[0286]




To a solution of tryptophol (6 g, 37.27 mmol, 1.0 equiv) in dimethyl sulfoxide (60 mL) was added 2-Iodoxybenzoic acid (12.5 g, 44.72 mmol, 1.2 equiv) at 40° C. under nitrogen atmosphere. The mixture solution was stirred for 2 h at 40° C. The reacting solution was diluted with 200 mL of dichloromethane, washed with 2×200 mL saturated sodium bicarbonate solution and 2×200 mL, saturated aqueous sodium chloride respectively. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated

under vacuum. The residue was applied onto a silica gel column with petroleum ether/ethyl acetate (3:1). 4 g (67%) of 1-2 was obtained as a yellow oil.

2. Synthesis of 1-2-1

[0287]




1-2-1

[0288] To a solution of 1-2 (100 mg, 0.63 mmol, 1.00 equiv) and 1-isopropylpiperazine (241.5 mg, 1.89 mmol, 3.00 equiv) in dichloromethane (1 mL) were added triethylamine (190.9 mg, 1.89 mmol, 3.00 equiv) and the solution of SnCl_4 in dichloromethane (1N, 0.32 mL, 0.5 equiv) at 25° C. The resulting mixture was stirred for 1 h at 25° C. under nitrogen atmosphere. Then sodium cyanoborohydride (58.6 mg, 0.95 mmol, 1.50 equiv) was added in portions. The mixture was stirred for 2 h at 25° C. The reaction was diluted with 20 mL of dichloromethane. The organic layer was

washed with 2x20 mL saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The crude product was purified by Prep-HPLC for 2 times with the following condition: Column: XTfridge Prep OBD C₁₈ Column, 30*150 mm, 5 μm; Mobile Phase A: Water (10 mmol/L NH₄HCO₃+0.1% NH₃.H₂O), Mobile Phase B: ?CCN; Flow rate: 60 milmin; Gradient: 22% B to 50% B in 9 min, 50% B; Wave Length: 254 nm; 4.6 mg (3%) of 1-2-1 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 272.25. ¹H NMR (300 MHz, DMSO-d₆) δ 10.76 (s, 1H), 7.51-7.49 (m, 1H), 7.34-7.32 (m, 1H), 7.14-6.97 (m, 3H), 2.83 (s, 2H), 2.58-2.46 (m, 11H), 0.97 (s, 6H).

3. Synthesis of 1-2-2

[0289]




1-2-2

[0290] In a similar manner described for 1-2-1, compound 1-2-2 was prepared from 100 mg of 1-2 and 189 mg of 1-methylpiperazine. 5.1 mg (3%) of 1-2-2 was obtained as a brown oil. MS m/z [M+H]⁺ (ESI): 244.10. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=7.5 Hz, 1H), 7.32 (d, J=8.1 Hz, 1H), 7.13 (s, 1H), 7.05 (t, 7.5 Hz, 1H), 6.95 (t, J=7.4 Hz, 1H), 2.85-2.79 (m, 2H), 2.58-2.53 (m, 2H), 2.47-2.33 (m, 8H), 2.15 (s, 3H).

4. Synthesis of 1-2-3

[0291]




1-2-3

[0292] In a similar manner described for 1-2-1, compound 1-2-3 was prepared from 100 mg of 1-2 and 164.4 mg of morpholine. 25.5 mg (17%) of 1-2-3 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 231.05. ¹H NMR (300 MHz, DMSO-d₆) δ 10.77 (s, 1H), 7.50 (d, J=7.8 Hz, 1H), 7.32 (d, J=7.8 Hz, 1H), 7.15 (s, 1H), 7.07-6.94 (m, 2H), 3.60 (t, J=4.2 Hz, 4H), 2.84 (t, J=7.8 Hz, 2H), 2.54 (t, J=7.8 Hz, 2H), 2.47-2.45 (m, 4H).

Synthesis of 1-2-4

[0293]




1-2-4

[0294] In a similar manner described for 1-2-1, compound 1-2-4 was prepared from 100 mg of 1-2 and 168.2 mg of piperidine. 3.4 mg (2%) of 1-2-4 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 229.20. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=5.1 Hz, 1H), 7.32 (d, J=6.0 Hz, 1H), 7.12 (s, 1H), 7.04-6.95 (m, 2H), 2.82 (s, 2H), 2.55 (s, 2H), 2.42 (s, 4H), 1.51-1.40 (m, 6H).

Synthesis of 1-2-5

[0295]




1-2-5

[0296] In a similar manner described for 1-2-1, compound 1-2-5 was prepared from 100 mg of 1-2 and 190.9 mg of 4-Hydroxypiperidine, 18.9 mg (12%) of 1-2-5 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 245.10. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=7.8 Hz, 1H), 7.32 (d, J=7.8 Hz, 1H), 7.13-6.93 (m, 3H), 4.53 (s, 1H), 3.49-3.38 (m, 1H), 2.87-2.73 (m, 4H), 2.59-2.52 (m, 2H), 2.12-2.01 (m, 2H), 1.75-1.70 (m, 2H), 1.45-1.33 (m, 2H).

7. Synthesis of 1-2-6

[0297]




1-2-6

[0298] To a solution of methyl 4-piperidineacetate (296.7 mg, 1.89 mmol, 3.00 equiv) and acetic acid (0.1 mL) in methanol (1 mL) were added sodium cyanoborohydride (58.6 mg, 0.95 mmol, 1.50 equiv) at 0° C. Then 1-2 (100 mg,

0.63 mmol, 1.00 equiv) was added to the mixture dropwise at 0° C. The mixture was stirred at 25° C. for 2 h. The reaction was diluted with 20 mL of dichloromethane. The organic layer was washed with 2x20 mL saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following condition: Column: (Bridge Prep OBD C₁₈ Column, 30*150 mm, 5 μm; Mobile Phase A: Water (10 mmol/L NH₄HCO₃+0.1%N₃·H₂O), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 30% B to 58% B in 9 min, 58% B; Wavelength: 254 nm. 27.1 mg (14%) of 1-2-6 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 301.05. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=7.8 Hz, 1H), 7.32 (d, J=8.1 Hz, 1H), 7.12 (d, J=1.8 Hz, 1H), 7.07-7.02 (m, 1H), 6.97-6.93 (m, 1H), 3.59 (s, 3H), 2.96-2.92 (m, 2H), 2.84-2.79 (m, 2H), 2.57-2.51 (m, 2H), 2.24 (d, J=6.6 Hz, 2H), 1.98-1.89 (m, 2H), 1.72-1.60 (m, 3H), 1.29-1.13 (m, 2H).

8. Synthesis of 1-2-7


[0299]



[0300] In a similar manner described for 1-2-1, compound 1-2-7 was prepared from 100 mg of 1-2 and 270.2 mg of piperidin-4-ylacetic acid. 7.8 mg (4%) of 1-2-7 was obtained as a white solid. MS m/z [M+H]⁺ (ESI): 287.10. ¹H NMR (300 MHz, DMSO-d₆) δ 10.76 (s, 1H), 7.50 (d, J=7.5 Hz, 1H), 7.33 (d, J=7.4 Hz, 1H), 7.13-6.95 (m, 3H), 3.02-2.89 (m, 2H), 2.88-2.74 (m, 2H), 2.57-2.53 (m, 2H), 2.11 (s, 2H), (s, 2H), 1.99-1.90 (m, 2H), 1.67-1.63 (m, 3H), 1.26-1.12 (m, 2H).

9. Synthesis of 1-2-8


[0301]



[0302] In a similar manner described for 1-2-7, compound 1-2-8 was prepared from 100 mg of 1-2 and 270.2 mg of methyl isonipecotate. 26.1 mg (14%) of 1-2-8 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 287.05. ¹H NMR (300 MHz, DMSO-d₆) δ 10.76 (s, 1H), 7.49 (d, J=7.8 Hz, 1H), 7.34-7.29 (m, 1H), 7.13 (d, J=2.1 Hz, 1H), 7.07-7.02 (m, 1H), 6.98-6.93 (m, 1H), 3.60 (s, 3H), 2.94-2.89 (m, 2H), 2.85-2.80 (m, 2H), 2.58-2.53 (m, 2H), 2.38-2.26 (m, 1H), 2.09-1.99 (m, 2H), 1.86-1.77 (m, 2H), 1.64-1.51 (m, 2H).

10. Synthesis of 1-2-9


[0303]



[0304] In a similar manner described for 1-2-1, compound 1-2-9 was prepared from 100 mg of 1-2 and 243.8 mg of isonipecotic acid. 4.3 mg (2%) of 1-2-9 was obtained as a white solid. MS m/z [M+H]⁺ (ESI): 273.10. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=7.2 Hz, 1H), 7.31 (d, J=7.5 Hz, 1H), 7.13 (s, 1H), 7.09-6.90 (m, 2H), 2.95-2.78 (m, 4H), 2.58-2.53 (m, 2H), 2.22-1.96 (m, 3H), 1.82-1.77 (m, 2H), 1.61-1.48 (m, 2H).

1. Synthesis of 1-2-10


[0305]



[0306] In a similar manner described for 1-2-1, compound 1-2-10 was prepared from 100 mg of 1-2 and 134.2 mg of pyrrolidine. 8.0 mg (6%) of 1-2-10 was obtained as a brown solid. MS m/z [M+H]⁺ (ESI): 215.10. ¹H NMR (300 MHz, DMSO-d₆) δ 10.76 (s, 1H), 7.49 (d, J=7.8 Hz, 1H), 7.32 (d, J=8.1 Hz, 1H), 7.13 (d, J=2.4 Hz, 1H), 7.07-7.02 (m, 1H), 6.98-6.93 (m, 1H), 2.87-2.82 (m, 2H), 2.69-2.64 (m, 2H), 2.59-2.50 (m, 4H), 1.75-1.64 (m, 4H).

2. Synthesis of 1-2-11


[0307]



[0308] In a similar manner described for 1-2-1, compound 1-2-11 was prepared from 100 mg of 1-2 and 164.4 mg of 3-Pyrrolidinol. 5.5 mg (4%) of 1-2-11 (racemate) was obtained as a white solid. MS m/z [M+H]⁺ (ESI): 231.05. ¹H NMR (300 MHz, DMSO-d₆) δ 10.75 (s, 1H), 7.49 (d, J=7.8 Hz, 1H), 7.32 (d, J=7.8 Hz, 1H), 7.13 (s, 1H), 7.07-7.02 (m, 1H), 6.98-6.93 (m, 1H), 4.67 (s, 1H), 4.19 (s, 1H), 2.84-2.73 (m, 3H), 2.67-2.58 (m, 3H), 2.45-2.34 (m, 2H), 2.01-1.95 (m, 1H), 1.56-1.51 (m, 1H).

3. Synthesis of 1-2-12

[0309]




1-2-12

[0310] In a similar manner described for 1-2-6, compound 1-2-12 was prepared from 100 mg of 1-2 and 243.8 mg of methyl 3-pyrrolidinecarboxylate. 34.1 mg (20%) of 1-2-12 was obtained as a brown solid. MS m/z. $[M+H]^+$ (ESI): 273.05. 1H NMR (300 MHz, DMSO- d_6) δ 10.77 (s, 1H), 7.50 (d, J=7.8 Hz, 1H), 7.32 (d, J=8.1 Hz, 1H), 7.14 (d, J=2.1 Hz, 1H), 7.07-7.02 (m, 1H), 6.98-6.93 (m, 1H), 3.62 (s, 3H), 3.08-2.98 (m, 1H), 2.86-2.79 (m, 3H), 2.72-2.53 (m, 5H), 2.04-1.91 (m, 2H).

4. Synthesis of 1-2-13

[0311]




1-2-13

[0312] In a similar manner described for 1-2-1, compound 1-2-13 was prepared from 100 mg of 1-2 and 217.3 mg of methyl 3-Pyrrolidinecarboxylic acid. 24.0 mg (15%) of 1-2-13 was obtained as a white solid, MS m/z (ESI): 259.05. 1H NMR (300 MHz, DMSO- d_6) δ 10.80 (s, 1H), 7.50 (d, J=6.9 Hz, 1H), 7.32 (d, J=7.5 Hz, 1H), 7.14 (s, 1H), 7.08-6.93 (m, 2H), 2.84-2.55 (m, 9H), 1.95 (s, 2H).

15. Synthesis of 1-2-14

[0313]




1-2-14

[0314] In a similar manner described for 1-2-1, compound 1-2-14 was prepared from 100 mg of 1-2 and 228.7 mg of N-Ethylaniline. 6.3 mg (4%) of 1-2-14 was obtained as a white solid, MS m/z $[M+H]^+$ (ESI): 265.10. NMR (300 MHz, DMSO- d_6) δ 10.84 (s, 1H), 7.55 (d, J=6.6 Hz, 1H), 7.35 (d, J=7.2 Hz, 1H), 7.22-7.15 (m, 3H), 7.11-6.96 (m,

2H), 6.72 (d, J=6.9 Hz, 2H), 6.60-6.54 (m, 1H), 3.55-3.49 (m, 2H), 3.41-3.35 (m, 2H), 2.95-2.90 (m, 2H), 1.11-1.06 (m, 3H).

16. Synthesis of 2-1-2

[0315]




2-1-2

[0316] To a solution of tryptamine (200 mg, 1.25 mmol, 1.0 equiv) in 3 mL of methanol was added sodium cyanohydroxide (196.9 mg, 3.13 mmol, 2.50 equiv), acetic acid (0.15 mL) and formaldehyde (253.8 mg, 3.13 mol, 2.5 equiv, 37% in aq. soln) in methanol (1.5 mL) at 0° C. The mixture solution was stirred for overnight at 25° C. and then concentrated under vacuum. The resulting solution was diluted with 20 mL of saturated sodium bicarbonate and extracted with 3x20 mL of ethyl acetate. The combined organic layer was washed with 2x20 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The resulting mixture was dissolved in 2 mL of acetonitrile, then di-tert-butyl (chloromethyl) phosphate (273.5 mg, 1.06 mmol, 2.0 equiv), Me₃-piperidine (164.5 mg, 1.06 mmol, 2.0 equiv) and sodium iodide (8.0 mg, 0.05 mmol, 0.1 equiv) was added at 25° C. under nitrogen atmosphere. The mixture was stirred for 24 h at 40° C. The resulting solution was diluted with 20 mL of ethyl acetate and washed with 2x10 mL of saturated sodium bicarbonate solution and 10 mL of saturated sodium chloride solution respectively. The combined organic layer was washed with 2x20 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was used in the next step directly without further purification. 120 mg of 2-1-1 an intermediate phosphate was Obtained as a white solid, which was used directly for the next reaction without further characterization.

[0317] To a solution of 2-1-1 (120 mg, 0.29 mmol, 1.0 equiv) in dichloromethane (1.2 mL) with an inert atmosphere of argon was added trifluoroacetic acid (0.6 mL) slowly at room temperature. The resulting solution was stirred for 1 h at room temperature. The resulting solution was concentrated under reduced pressure. The crude product was purified by Flash with the following conditions: Column, C₁₈ silica gel; mobile layer, water and methanol (20% methanol up to 100% in 15 min and hold 100% for 5 min); Detector, 220 nm. The fractions were concentrated under reduced pressure. 28.3 mg (27.6% yield) of 2-1-2 was Obtained as a white solid. MS m/z $[M+H]^+$ (ESI): 299.05. 1H NMR (300 MHz, DMSO- d_6) δ 11.11 (s, 1H), 7.65 (d, J=7.8 Hz, 1H), 7.37 (d, J=8.1 Hz, 1H), 7.26 (s, 1H), 7.11-6.98 (m, 2H), 4.89 (d, J=8.7 Hz, 2H), 3.51-3.45 (m, 2H), 3.18-3.13 (m, 2H), 3.07 (s, 6H). ^{31}P NMR (121 MHz, DMSO- d_6) δ -1.88.

19. Synthesis of 3-1-4

[0318]



[0319] In a 250 mL flask in the ice bath, 12.83 g (0.11 mol, 1.1 equiv) of thionylchloride was added and then 8.91 g (0.10 mol, 1 equiv) of N,N-dimethyl ethanolamine was added dropwise over 0.5 h under stirring. The resulting mixture was stirred for 1 h at 50° C. Then 150 mL of ethanol was added and heated to reflux for 0.5 h. After cooling to room temperature, 3-1-1 (6 g, 42%) was collected by filtration as a light-yellow solid.

21. Synthesis of 3-1-2

[0320] A solution of 3-1-1 (6 g, 41.9 mmol, 1 equiv) in N,N-dimethylformamide (60 mL) was treated with KOH (11.75 g, 209.5 mmol, 5 equiv) for 1 h at 25° C., 7-methoxy-1H-indole (6.17 g, 41.9 mmol, 1 equiv) was added to the mixture at 25° C. The resulting mixture was stirred for 16 h at 50° C. The reaction was diluted with dichloromethane (200 mL), washed with water (3×200 mL). The organic phase was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by Flash with the following condition: Column, C₁₈ silica gel; mobile layer, water (0.05% NH₄HCO₃) and acetonitrile (10% acetonitrile up to 100% in 20 min and hold 100% for 5 min); Detector, 220 nm. The fractions were concentrated under reduced pressure to afford 3-1-2 (5 g, 54%) as a white solid. MS m/z [M+H]⁺ (ESI): 219.14.

22. Synthesis of 3-1-3

[0321] To a solution of 3-1-2 (600 mg, 2.75 inmol, 1 equiv) in dichloromethane (20 mL) was added aluminium chloride (1.83 g, 13.75 mmol, 5 equiv) in portions at 0° C. The mixture was stirred overnight at 25° C. The reaction was quenched with methanol (20 mL), filtered and concentrated under reduced pressure. The crude product was purified by

Flash with the following condition: Column, C₁₈ silica gel; mobile layer, water (0.1% NH₃H₂O) and methanol (10% methanol up to 100% in 20 min and hold 100% for 5 min); Detector, 220 nm. The fractions were concentrated under reduced pressure. This resulted in 3-1-3 (415 mg, 73%) as a brown oil, MS m/z [M+H]⁺ (ESI): 205.13.

23. Synthesis of 3-1-4

[0322] To a solution of 3-1-3 (100 mg, 0.49 mmol, 1 equiv) in dichloromethane (1 mL) was added pentanoic acid (75 mg, 0.74 mmol, 1.5 equiv), 4-dimethylaminopyridine (17.9 mg, 0.15 mmol, 0.3 equiv), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (141.2 mg, 0.74 mmol, 1.5 equiv) at 25° C. The mixture was stirred for 2 h at 25° C. then diluted with 20 mL of dichloromethane, washed with 2×10 mL of saturated sodium bicarbonate solution and 2×10 mL of saturated aqueous sodium chloride solution respectively. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The crude product was purified by Flash with the following condition: Column, C₁₈ silica gel; mobile layer, water (0.05% NH₄HCO) and acetonitrile (20% acetonitrile up to 100% in 20 min and hold 100% for 5 min); Detector, 220 nm. The fractions were concentrated under reduced pressure. This resulted in 3-1-4 (22.6 mg, 16%) as a brown oil. MS m/z [M+H]⁺ (ESI): 289.25. ¹H NMR (300 MHz, Methanol-d₄) δ 7.42-7.38 (m, 1H), 7.16 (s, 1H), 7.01-6.94 (m, 1H), 6.82-6.78 (m, 1H), 6.48-6.44 (m, 1H), 4.33-4.26 (m, 2H), 2.75-2.60 (m, 4H), 2.23 (s, 6H), 1.81-1.69 (m, 2H), 1.53-1.43 (m, 2H), 1.03-0.96 (m, 3H).


24. Synthesis of 4-1-2, 4-1-3, and 4-1-4

[0323] To a solution of tryptamine (400 mg, 2.5 mmol, 1.0 equiv) in 3 ml of methanol was added sodium cyanoboro-

hydride (393.8 mg, 6.26 mmol, 2.50 equiv), acetic acid (0.3 mL) and formaldehyde (507.6 mg, 6.26 mol, 2.5 equiv, 37% in aq. sola) in methanol (4 mL) at 0° C. The mixture solution was stirred for overnight at 25° C. and then concentrated under vacuum. The resulting solution was diluted with 20 mL of saturated sodium bicarbonate and extracted with 3×20 mL of ethyl acetate. The combined organic layer was washed with 2×20 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The resulting mixture dissolved in 3 mL of dichloromethane, then 483.5 mg of triethylamine was added. 1-chloroethyl carbonochloridate (340 mg, 2.39 mmol, 1.5 equiv) dropwise at 0° C. The mixture was stirred at 25° C. for 2 h. The reaction was diluted with dichloromethane (20mL), washed with saturated aqueous sodium chloride (2×20 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. 300 mg (crude) of 4-1-1 (intermediate) was obtained as a yellow solid. It was used for next step directly without further purification.

25. Synthesis of 44-2 (Racemate)

[0324] To a solution of 4-1-1 (300 mg, 1.02 mmol, 1.0 equiv) and triethylamine (2.06 g, 20.40 mmol, 20 equiv) in dimethylformamide (6 mL) was added isobutyric acid (449 mg, 5.10 mmol, 5 equiv) at 25° C. The mixture was stirred for 16 h at 60° C. The reaction was diluted with 50 mL of dichloromethane, washed with 1×50 mL saturated sodium chloride aqueous solution. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by Prep-HPLC for 2 times with the following condition: Column: X Select CSH Fluoro Phenyl, 30*150 mm, 5 μm; Mobile Phase A: Water (0.05% HCl), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 28% B to 58% B in 8 min, 58% B; Wavelength: 254 nm; 9.0 mg of 4-1-2 was obtained as a light brown solid. Combined the other batches, 26 mg of 4-1-2 was obtained. MS m/z [M+H]⁺ (ESI): 347.25. ¹H NMR (300 MHz, DMSO-d₆) δ 8.06 (d, J=8.1 Hz, 1H), 7.75-7.67 (m, 2H), 7.44-7.32 (m, 2H), 7.02-6.97 (m, 1H), 3.40-3.35 (m,



2H), 3.12 -3.07 (m, 2H), 2.84 (s, 6H), 2.66-2.56 (m, 1H), 1.65 (d, 5.4 Hz, 3H), 1.11 (d, J=7.2 Hz, 6H).

26. Synthesis of 4-1-3 (Racemate)


[0325] In a similar manner described for 4-1-2, compound 4-1-3 was prepared from 300 mg of 4-1-1 and 306 mg of acetic acid. 5.0 mg of 4-1-3 was obtained as a light brown solid. Combined the other batches, 40 mg of 4-1-3 was obtained. MS m/z $[M+H]^+$ (ESI): 319.10. 1H NMR (400 MHz, DMSO- d_6) δ 10.16 (s, 1H), 8.08 (d, J=8.0 Hz, 1H), 7.75 (d, J=7.6 Hz, 1H), 7.65 (s, 1H), 7.43-7.32 (m, 2H), 7.01-6.97 (m, 1H), 3.39-3.33 (m, 2H), 3.15-3.11 (m, 2H), 2.83 (s, 6H), 2.10 (s, 3H), 1.63 (d, J=5.6 Hz, 3H).

27. Synthesis of 4-1-4 (Racemate)

[0326] In a similar manner described for 4-1-2, compound 4-1-4 was prepared from 300 mg of 4-1-1 and 520 mg of pivalic acid. The crude product (300 mg) was purified by Prep-HPLC with the following conditions (Column: ?Bridge Shield RP18 OBD Column, 30*150 mm, 5 μ m; Mobile Phase A: Water (10 mmol/L, NH_4HCO_3 +0.1% $NH_3\cdot H_2O$), Mobile Phase B: ACN; Flow rate: 60 mL/min; Gradient: 48% B to 78% B in 9 min, 78% B; Wavelength: 254 nm. 5 mg of 4-1-4 was obtained as oil. MS m/z $[M+H]^+$ (ESI): 361.25. 1H NMR. (300 MHz, DMSO- d_6) δ 8.03 (d, J=8.1 Hz, 1H), 7.65-7.62 (m, 1H), 7.51, (s, 1H), 7.39-7.26 (m, 6.98-6.93 (m, 1H), 2.83-2.78 (m, 2H), 2.57-2.54 (m, 2H), 2.20 (s, 6H), 1.64 (d, J=5.4 Hz, 3H), 1.17 (s, 9H).

28. Synthesis of 5-1-4

[0327]



[0328] To a stirred solution of 4-methoxy-1H-indole (1.0 g, 6.80 mmol, 1.0 equiv) in diethyl ether (20.0 mL) was added oxalyl chloride (0.9 g, 6.80 mmol, 1.0 equiv) dropwise at 0° C. under nitrogen atmosphere. The resulting mixture was stirred for additional 3 h at room temperature. The resulting mixture 5-1-1 was used in the next step directly without further purification.

[0329] To a stirred solution of dimethylamine hydrochloride (1.7 g, 21.04 mmol, 5.0 equiv) and triethylamine (3.4 g, 33.66 mmol, 8.0 equiv) in tetrahydrofuran (10.0 mL) was added 5-14 (crude, obtained at last step) dropwise at 0° C. under nitrogen atmosphere. The mixture was stirred for additional 0.5 h at room temperature. The reaction was quenched by the addition of water (50 mL) at 0° C. The resulting mixture was extracted with ethyl acetate (3x50 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The crude was purified by Flash with the following condition: Column, C_{18} silica gel; mobile layer, water (0.05% NH_4HCO_3) and acetonitrile (10% acetonitrile up to 100% in 15 min and hold 100% for 5 min); Detector, 254 nm. 600 mg (58% yield) of 5-1-2 was obtained as a yellow solid. MS m/z $[M+H]^+$ (ESI): 247.10.






[0330] To a stirred solution of 5-1-2 (600.0 mg, 2.44 mmol, 1.0 equiv) in tetrahydrofuran (10.0 mL) was added Lithium Aluminum Hydride (184.9 mg, 4.87 mmol, 2.0 equiv) in portions at 0° C. The resulting mixture was stirred for 1.5 days at 70° C. The reaction was quenched by the addition of water (20.0 mL) at 0° C. The aqueous layer was extracted with ethyl acetate (3x30 mL). The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The crude was purified by Flash with the following condition: Column, C_{18} silica gel; mobile layer, water (0.1% $NH_3\cdot H_2O$ +10 mmol/L NH_4HCO_3) and acetonitrile (10% acetonitrile up to 100% in 15 min and hold 100% for 5 min); Detector, 254 nm, 350 mg (66% yield) of 5-1-3 was obtained as a yellow solid. MS m/z $[M+H]^+$ (ESF): 219.15.

[0331] To a stirred solution of 5-1-3 (80.0 mg, 0.37 mmol, 1.0 equiv) in dichloromethane (1.0 mL) was added tribromoborane (459.0 mg, 1.83 mmol, 5.0 equiv) dropwise at -78° C. under nitrogen atmosphere. The resulting mixture was stirred for 9 h at room temperature. The mixture was concentrated under vacuum. The residue was dissolved in dichloromethane (30.0 mL), added 50 mg of $KHCO_3$ and

15.0 mL of methanol. After stirring for 30 min, the resulting mixture was concentrated under reduced pressure. The resulting mixture was dissolved in N,N-dimethylformamide (1.0 mL), added entanoic acid (30.0 mg, 0.15 mmol, 1.0 equiv), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (56.3 mg, 0.29 mmol, 2.0 equiv) and 4-dimethylaminopyridine (9.0 mg, 0.07 mmol, 0.5 equiv) at room temperature under nitrogen atmosphere. The resulting mixture was stirred for 16 h at room temperature. The reaction was diluted with ethyl acetate (3×10 mL), washed with water (3×10 mL). The organic phase was dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure. The crude product was purified by Flash with the following condition: Column, C₁₈ silica gel; mobile layer, water (0.05% NH₄HCO₃) and acetonitrile (10% acetonitrile up to 100% in 20 min and hold 100% for 5 min); Detector, 254 nm. This resulted in 5-1-5 (20.0 mg, 47%) as a colorless oil. MS m/z [M+H]⁺ (ESI): 289.25. ¹H NMR (300 MHz, DMSO-d₆) δ 11.03 (s, 1H), 7.23 (d, J=8.1 Hz, 1H), 7.15 (d, J=2.4 Hz, 1H), 7.06-7.01 (m, 1H), 6.66-6.63 (m, 1H), 2.78-2.65 (m, 4H), 2.47-2.43 (m, 2H), 2.20 (s, 6H), 1.73-1.63 (m, 2H), 1.48-1.35 (m, 2H), 0.94 (t, J=7.3 Hz, 3H).

Improved Prodrugs of Tryptamines, Such as N,N-Dimethyltryptamine (DMT).

[0332] Prodrug 3-1-4, 4-1-2 and 4-1-4 was supposed to be stable in formulation and should release the parent compound in vivo.



[0333] Preliminary studies demonstrated the stability of prodrug (Tables 5-9). While solution stability has been shown for prodrug, the plasma stability data clearly showed release of parent drug in mouse plasma instantly. The following prodrugs were tested.

[0334] Prodrugs 3-1-4, 4-1-2 and 4-1-4 Stability Results in
PBS pH 6.5 With and Without Pepsin

TABLE 5

Detection of VLS-03-1 Parent in PBS pH 6.5, SGF with and without pepsin								
Peak Area of MRM: 205.0/159.8								
Compound	Analyte	Incubation	VLS-03-1 standard	0 min	30 min	60 min	120 min	240 min
VLS-03-1	VLS-03-1 Parent	PBS (pH 6.5)	4.49E+04	5.04E+04	4.71E+04	4.71E+04	4.86E+04	4.75E+04
VLS-03-1	VLS-03-1 Parent	SGF with pepsin	4.22E+04	5.72E+04	5.37E+04	5.01E+04	5.48E+04	5.27E+04
VLS-03-1	VLS-03-1 Parent	SGF without pepsin	3.75E+04	4.53E+04	4.75E+04	4.51E+04	4.66E+04	4.57E+04

TABLE 6

Detection of VLS-03-10 Parent in PBS pH 6.5, SGF with and without pepsin								
Peak Area of MRM: 188.7/144.0								
Compound	Analyte	Incubation	VLS-03-10 standard	0 min	30 min	60 min	120 min	240 min
VLS-03-10	VLS-03-10 Parent	PBS (pH 6.5)	4.34E+03	5.32E+03	4.68E+03	5.00E+03	4.72E+03	4.95E+03
VLS-03-10	VLS-03-10 Parent	SGF with pepsin	4.05E+03	5.13E+03	5.10E+03	5.25E+03	5.32E+03	5.28E+03
VLS-03-10	VLS-03-10 Parent	SGF without pepsin	3.46E+03	5.29E+03	4.98E+03	4.66E+03	4.93E+03	4.61E+03

TABLE 7

Detection of VLS-03-15 Parent in PBS pH 6.5, SGF with and without pepsin								
Peak Area of MRM: 205.0/159.8								
Compound	Analyte	Incubation	VLS-03-15 standard	0 min	30 min	60 min	120 min	240 min
VLS-03-15	VLS-03-15 Parent	PBS (pH 6.5)	6.74E+05	6.56E+05	6.55E+05	6.28E+05	5.86E+05	5.51E+05
VLS-03-15	VLS-03-15 Parent	SGF with pepsin	6.26E+05	6.75E+05	7.21E+05	7.26E+05	7.14E+05	7.15E+05
VLS-03-15	VLS-03-15 Parent	SGF without pepsin	6.10E+05	7.05E+05	6.94E+05	6.55E+05	7.09E+05	6.96E+05

Data Summary

[0335]

TABLE 8

Stability results of test compounds in PBS pH 6.5, SGF with and without pepsin							
Compound	Incubation	T _{1/2} (min)	Remaining Percentage (%)				
			0 min	30 min	60 min	120 min	240 min
Chlorambucil	PBS (pH 6.5)	47.25	100.00	64.51	42.26	16.83	2.99
Erythromycin	SGF with pepsin	43.62	100.00	56.55	37.33	14.06	2.14
Erythromycin	SGF without pepsin	39.47	100.00	52.67	28.46	8.27	1.47
VLS-03-1	PBS (pH 6.5)	9999.00	100.00	96.32	99.71	100.03	101.35
VLS-03-1	SGF with pepsin	8950.90	100.00	101.74	90.49	102.16	96.47
VLS-03-1	SIG F without pepsin	9999.00	100.00	106.25	99.72	104.22	103.22
VLS-03-10	PBS (pH 6.5)	19876.96	100.00	92.62	100.03	92.63	97.66
VLS-03-10	SGF with pepsin	1604.11	100.00	96.82	93.09	91.42	89.32
VLS-03-10	SGF without pepsin	7008.36	100.00	99.86	104.95	98.41	98.86
VLS-03-15	PBS (pH 6.5)	922.46	100.00	96.93	89.82	89.44	82.54
VLS-03-15	SGF with pepsin	9999.00	100.00	97.29	107.53	103.18	102.22
VLS-03-15	SGF without pepsin	7839.52	100.00	101.12	103.35	105.28	97.98

If calculated T_{1/2} < 0, then T_{1/2} was reported as 9999 min.

Prodrugs 3-1-4, 4-1-2 and 4-1-4 Stability Results in Mouse Plasma


[0336]

TABLE 9


Stability Results of Test Compounds in Mouse Plasma							
Compound	Species	T _{1/2} (min)	Remaining Percentage (%)				
			0 min	15 min	30 min	60 min	120 min
Propranolol	Mouse	32.60	100.00	85.78	70.74	29.46	8.74
VLS-03-1		N.A.	100.00	0.00	0.00	0.00	0.00
VLS-03-10		N.A.	100.00	0.00	0.00	0.00	0.00
VLS-03-15		N.A.	100.00	0.00	0.00	0.00	0.00

N-Phosphonoxyethyl Prodrug 64-2 (PH-VLS-02-23-0)

[0337]




-continued



[0338] Using methods described in Gunda I. George (*Journal of Medicinal Chemistry*, 1999, Vol. 42, 309) N-Phosphonoxyethyl Prodrug compounds were prepared.



Illustration of the Prodrug Strategy:

[0339]



[0340] 6-1-2 was synthesized, as a novel prodrug strategy for improving the water solubility of tertiary amine-containing drugs. And tertiary amine DMT was converted in two step reaction to a very polar, N-Phosphonoxyethyl 1 Prodrug 6-1-3.

[0341] The prodrug 6-1-2 releases the parent DMT in vivo through a two-step bio-reversion process. The rate determining first step is via an enzymatic process, where prodrug, bio-reversion involves a phosphatase-catalyzed dephosphorylation to give the resultant hydroxymethyl quaternary ammonium intermediate and inorganic phosphate. The second step involves conversion of hydroxymethyl quaternary ammonium intermediate chemically to DMT and formaldehyde at physiological pH.



6-1-2 was synthesized, from commercially available in 3 simple steps.

[0342] Chemical stability of prodrug 6-1-2 (VLS-02-23) was evaluated and the results enumerated in Tables 10-11. PK studies of prodrug 6-1-2 and metabolite (VLS-02-23-10) in mice were also conducted and both plasma exposure and brain exposure were analyzed as shown in FIGS. 1A-1C and FIGS. 2A-2B, respectively.

Chemical Stability of N-Phosphonoxyethyl Prodrug 6-1-2 (NTS-02-23)

Data Summary

[0343]

TABLE 10

Stability results of test compounds in PBS pH 6.5, SGF with and without pepsin							
Compound	Incubation	T _{1/2} (min)	Remaining Percentage (%)				
			0 min	30 min	60 min	120 min	240 min
Chlorambucil	PBS (pH 6.5)	43.17	100.00	61.12	43.17	15.12	2.15
Erythromycin	SGF with pepsin	40.26	100.00	56.55	34.88	11.22	1.60
Erythromycin	SGF without pepsin	37.36	100.00	59.45	33.60	10.27	1.20
VLS-02-23	PBS (pH 6.5)	9999.00	100.00	104.71	106.37	107.29	117.35
VLS-02-23	SGF with pepsin	6227.35	100.00	96.70	106.43	102.36	96.82
VLS-02-23	SIG F without pepsin	9999.00	100.00	98.76	107.08	101.90	108.80

Plasma Stability of N-Phosphonoxyethyl Prodrug 6-1-2 (NTS-02-23)

Data Summary


[0344]

TABLE 11

Stability Results of Test Compounds in Mouse Plasma							
Compound	Species	T _{1/2} (min)	Remaining Percentage (%)				
			0 min	15 min	30 min	60 min	120 min
Propranolol	Mouse	34.09	100.00	80.81	64.04	34.86	9.05
VLS-02-03	Mouse	568.34	100.00	114.24	110.21	109.95	91.16

What is claimed:


1. A compound of Formula:



wherein R_1 and R_2 are independently hydrogen, alkyl, or cycloalkyl; or R_1 and R_2 , together with the atoms to which they are attached form a heterocyclic ring;

X , Y , and Z are independently hydrogen, halogen, $-O-$, $-S-$, $-NR_4-$, or $(P=O)-OR_4(-OR')$, wherein R' and R_4 are independently hydrogen or alkyl;

R_3 , R_4 , and R_5 are independently absent, hydrogen, alkyl, $-(C=O)-alkyl$, $-(C=O)CHR_7$, $-(C=O)(CH_2)_n$ (NH_2), $-(C)CH(NH_2)(CH_3)$, $-(C=O)CH(NH_2)$, $-(CH_2)_n(NH)(C=NH)$, $-(C=O)(CH_2)_n(NH)(C=NH)-NH_2$, or



L and M are independently alkyl or aryl;

R_7 is hydrogen or alkyl;


R_9 is hydrogen, alkyl, or halogen;

R_{11} is a pharmaceutically acceptable cation, hydrogen, deuterium, alkyl, or cycloalkyl; and


n is an integer from 2-7.

2. The compound of claim 1, wherein the pharmaceutically acceptable cation is Na^+ , K^+ , or NH_4^+ .


3. The compound of claim 1, of the formula:




4. The compound of claim 1, of the formula:




5. The compound of claim 1, of the formula:



6. The compound of claim 1, of the formula:



7. A compound of Formula (VIII):




wherein R_1 and R_2 are independently hydrogen, alkyl, or cycloalkyl; or R_1 and R_2 , together with the atoms to which they are attached form a heterocyclic ring;

X , Y , and Z are independently hydrogen, halogen, $-O-$, $-S-$, $-NR_4-$, or $(P=O)-OR_4(-OR')$, wherein R' and R_4 are independently hydrogen or alkyl;

R_3 , R_4 , and R_5 are independently absent, hydrogen, alkyl, $-(C=O)-alkyl$, $-(C=O)CHR_7$, $-(C=O)(CH_2)_n$

(NH_2) , $-(\text{C})\text{CH}(\text{NH}_2)(\text{CH}_3)$, $-(\text{C}=\text{O})\text{CH}(\text{NH}_2)$,
 $-(\text{CH}_2)_n(\text{NH})(\text{C}=\text{NH})$, $-(\text{C}=\text{O})(\text{CH}_2)_n(\text{NH})$
 $(\text{C}=\text{NH})-\text{NH}_2$, or



L and M are independently alkyl or aryl;

R_7 is hydrogen or alkyl;

R_9 is hydrogen, alkyl, or halogen;

R_{11} and R_{17} are independently a pharmaceutically acceptable cation, hydrogen, deuterium, alkyl, cycloalkyl, —or R_{11} and R_{12} together with the atoms to which they are attached form a heterocyclic ring, and

n is an integer from 2-7.

8. The compound of claim 7, wherein the pharmaceutically acceptable cation is Na^+ , K^+ , or NH_4^+ .

* * * * *