




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : C12N 15/31, 15/82, A01H 5/00</p>	A1	<p>(11) International Publication Number: <b>WO 00/09696</b></p> <p>(43) International Publication Date: 24 February 2000 (24.02.00)</p>
<p>(21) International Application Number: PCT/GB99/02652</p> <p>(22) International Filing Date: 12 August 1999 (12.08.99)</p> <p>(30) Priority Data: 9817707.4 13 August 1998 (13.08.98) GB</p> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(71) Applicant (for US only): STEWART, Lesley (Personal representative of STEWART, Gordon, Sidney, Anderson, Bernie) [/GB]; 14 James Avenue, Loughborough LE11 5QL (GB).</p> <p>(72) Inventor: STEWART, Gordon, Sidney, Anderson, Bernie (deceased) (deceased).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): FRAY, Rupert, George [GB/GB]; University of Nottingham, School of Biological Sciences, Division of Plant Science, Sutton Bonington Campus, Loughborough LE12 5RD (GB). THROUP, John, Peter [GB/US]; SmithKline Beecham, Pharmaceuticals R &amp; D, UP1345 South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426 (US). WALLACE, Andrew, David</p>		<p>[GB/GB]; Unilever Research, Colworth Laboratory, Colworth House, Sharnbrook, Bedfordshire MK44 1LQ (GB). GRIERSON, Donald [GB/GB]; University of Nottingham, School of Biological Sciences, Division of Plant Science, Sutton Bonington Campus, Loughborough LE12 5RD (GB).</p> <p>(74) Agents: HUSKISSON, Frank, Mackie et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> With international search report.</p>

(54) Title: EXPRESSION OF BACTERIAL SIGNAL MOLECULES IN PLANTS

## (57) Abstract

The ability of a plant to defend against attack by bacteria, and any virus borne by the bacteria, is enhanced by transforming the plant genome with a gene of bacterial origin which enables the plant to produce a bacterial pheromone, N-acyl-L-homoserine lactone. Such plants also secrete the lactone into the soil enhancing the protective effect of antifungal rhizobacteria.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## EXPRESSION OF BACTERIAL SIGNAL MOLECULES IN PLANTS

This invention relates to the expression of bacterial signal molecules in plants which allows, for example, modulation of the interaction between plants and infecting or symbiotic bacteria.

The ability of bacteria to respond to environmental cues such as nutrient availability, temperature or pH is critical to microbe success. It is apparent that individual bacteria can also sense the density and state of the local bacterial population of which they are members. This sensing ability, referred to as "quorum sensing", allows a bacterial community to synchronise growth and development and, when the minimum population or "quorum" has been achieved, to initiate a concerted population response. Quorum sensing is thus an example of multicellular behaviour in prokaryotes and regulates diverse physiological processes including bioluminescence, swarming, antibiotic biosynthesis, plasmid conjugal transfer and the production of virulence determinants in pathogens.

The signalling pheromones upon which quorum sensing is based have been identified as *N*-acyl-L-homoserine lactones (reviewed by Swift *et.al.* "Quorum sensing: a population-density component in the determination of bacterial phenotype", *Trends in Biochemical Science*, **21**, 214-219 (1996). *N*-acyl-L-homoserine lactones molecules comprise a homoserine lactone moiety (derived from amino acid metabolism, possibly via S-adenosyl methionine) linked to an acyl sidechain (probably derived from fatty acid synthesis). A number of *N*-acyl-L-homoserine lactones with different acyl side chains have been identified in different bacterial systems where they elicit a wide range of quorum-dependent responses such as swarming, pathogenicity, conjugation or production of colour, light or antibiotics.

Several bacterial species produce the same *N*-acyl-L-homoserine lactone, although in some of the species it may regulate a different biological process. For example, the *luxI* gene product of *Photobacterium fischeri* synthesises *N*-(3-oxohexanoyl)-L-homoserine lactone which regulates bioluminescence in a cell density-dependent manner, whilst the *carI* gene product of *Erwinia carotovora* also produces *N*-(3-oxohexanoyl)-L-homoserine lactone which in this bacterium is responsible for the induction of secreted plant cell wall degrading exoenzymes and of the antibiotic carbapenem. The *cviI* gene of *Chromobacterium violaceum* encodes the enzyme for synthesis of *N*-hexanoyl-L-homoserine lactone which is

structurally very similar to the oxohexanoyl analogue and which induces production of the purple pigment violacein. Inactivation of *luxI*, *carI* or *cviI* results in loss of the density dependent bioluminescence, virulence or violacein production respectively. The relevant operons can, however, be induced by the addition of an exogenous supply of the *N*-acyl-L-homoserine lactone to the mutant bacteria.

*CarI* mutants of *Erwinia carotovora* appear to be completely avirulent when tested on tobacco. They can neither macerate plant tissue nor multiply *in planta* because they lack pectin lyase, pectate lyase, polygalacturonase, cellulase and protease. It is pertinent to ask how the expression of these exoenzymes only at high cell density in the wild-type cells may contribute to the success of *Erwinia* as a plant pathogen. It has been suggested that under aerobic conditions, a successful *E. carotovora* infection requires a relatively high inoculum ( $10^6$  -  $10^7$  c.f.u.) and the progression of the disease is then a competition between bacterial multiplication and development of plant resistance. Thus, the production of macerating enzymes at low cell densities would not give rise to a successful infection, but would result in the induction of the local and systemic plant defence response, which in turn would hamper subsequent infections. Such resistance to *E. carotovora* infection is seen when the plant defence response is artificially induced by the application of salicylic acid.

While not wishing to be bound by any theory as to the manner in which the invention proposed herein operates, the following explanation of the naturally occurring phenomenon of quorum sensing is offered. Using *Photobacterium fischeri* as a convenient example, the expression of two regulatory genes, *luxI* and *luxR*, is necessary for the expression of the genes necessary for bioluminescence. Expression of *luxI* leads to production of the pheromone *N*-(3-hydroxy)hexanoyl-L-homoserine lactone, the mechanisms by which the lactone is synthesised being largely irrelevant to this discussion. A complex of the pheromone with the protein produced by the *luxR* gene gives a phenotypic response, in the case of *P. fischeri*, bioluminescence. At low population density of bacteria, *luxI* and *luxR* are transcribed at low level and there is insufficient accumulation of the pheromone (*N*-acyl-L-homoserine lactone) to elicit *luxI*-dependent transcription of the operon responsible for visible bioluminescence. It has been suggested that in the absence of sufficient pheromone, and/or a chaperonin known as GroESL, *luxR* is unstable and sensitive to degradation. As the population grows, however, the concentration of the pheromone increases gradually. At a critical level of the pheromone, which represents a critical population density, a complex between *luxR* and the pheromone is thought to bind to a palindromic sequence within the *luxI*

operator thereby activating increased transcription of the operon necessary for increased production of the pheromone and for bioluminescence.

The present invention seeks to provide a method and means of manipulating plant/microbe interactions.

5           According to the present invention there is provided a method of protecting a plant against bacterial infection and/or virus infection transmitted by bacteria, comprising introducing into the genome of the plant by transformation the ability to synthesise a *N*-acyl-L-homoserine lactone.

10           Further according to the invention there is provided a method of protecting a plant against bacterial infection and/or virus infection transmitted by bacteria, comprising introducing into the genome of the plant by transformation the ability to synthesise an analogue of *N*-acyl-L-homoserine lactone.

15           The invention also provides a method of enhancing interaction between an antifungal rhizobacterium and a plant comprising introducing into the genome of the plant by transformation the ability to synthesise the *N*-acyl-L-homoserine lactone naturally produced by the rhizobacterium.


The invention also provides a recombinant plant genome containing a gene construct for *in planta* expression of an *N*-acyl-L-homoserine lactone.

Preferably expression of introduced genes is targeted to plant chloroplasts.

20           The gene specifying the *N*-acyl-L-homoserine lactone may be selected from the group consisting of, the *yenI* gene of *Yersinia enterocolitica*; the *cviI* gene of *Chromobacterium violaceum*; the *luxI* gene of *Photobacterium fischeri*; the *carI* gene of *Erwinia carotovora*; the *traI* gene of *Agrobacterium tumefaciens* and the *lasI* and *vsml* genes of *Pseudomonas aeruginosa*.

25           Examples of suitable sources of DNAs specifying *N*-acyl-L-homoserine lactones and the acyl group involved are as follows:

Table 1

**N-acyl-L-homoserine lactone**

Bacterium	Signal Generator	Response Regulator	N-acyl-group R
<i>Yersinia enterocolitica</i>	<i>yenI</i>	<i>yenR</i>	3-oxohexanoyl
<i>Chromobacterium violaceum</i>	<i>cviI</i>	<i>cviR</i>	3-hexanoyl
<i>Photobacterium fischeri</i>	<i>luxI</i>	<i>luxR</i>	3-oxohexanoyl
<i>Erwinia carotovora</i>	<i>carI</i>	<i>carR</i>	3-oxohexanoyl
<i>Agrobacterium tumefaciens</i>	<i>traI</i>	<i>traR</i>	3-oxo-octanoyl
<i>Pseudomonas aeruginosa</i>	<i>lasI</i>	<i>lasR</i>	3-oxo-dodecanoyl
<i>Pseudomonas aeruginosa</i>	<i>vsmI</i>	<i>vsmR</i>	butanoyl

5 These examples in Table 1 are quoted in Swift *et al.*, *Trends in Biochemical Science*, **21**, 214-219 (1996).

Table 2 below gives further examples along with references and the appropriate GenBank Accession Numbers.

Table 2

Organism	Signal generator	Response Regulator	Signal Molecule	GenBank Accession number	References
<i>Aeromonas hydrophila</i>	<i>AhyL</i>	<i>AhyR</i>	unknown	X89469	
<i>Agrobacterium tumefaciens</i>	<i>TraI</i>	<i>TraR</i>	N-(3-oxo)-octanoyl-L-homoserine Lactone (OOHL)	L17024, L22207	Fuqua <i>et al.</i> , 1994; Hwang <i>et al.</i> 1995
<i>Chromobacterium violaceum</i>	<i>CviI</i>	<i>CviR</i>	N-hexanoyl-L-homoserine lactone (OHL)		Winson, <i>et al.</i> , (1994)
<i>Enterobacter</i>	<i>EagI</i>	unknown	N-(3-	x74300	Swift <i>et al.</i> , 1993

<i>agglomerans</i>			oxo)hexanoyl-L-homoserine lactone (OHHL)		
<i>Erwinia carotovora subsp carotova</i>	<i>Carl</i>	<i>CarR</i>	OHHL	U17224, X72891, X74299, X80475	McGowan et al., 1995
<i>Erwinia stewartii</i>	<i>EsaI</i>	<i>EsaR</i>	OHHL	L32183, L32184	Beck von Bodman and Farrand, 1995
<i>Escherichia coli</i>	unknown	<i>SdiA</i>	unknown	Xo3691	Sitnikov et al 1995
<i>Photobacterium fischeri</i>	<i>LuxI</i>	<i>LuxR</i>	OHHL,OOHL	M19039, M96844, M25752	Meignhen, 1994; Devine et al, 1988
<i>Pseudomonas aeruginosa</i>	<i>LasI</i>	<i>LasR</i>	<i>N</i> -(3-oxo)-dodecanoyl-L-homoserine lactone (OdDHL)	M59425	Winson et al 1995;.
	<i>VsmI</i>	<i>vsmR</i>	<i>N</i> -butanoyl-L-homoserine lactone (BHL), HHL	L08962, U11811, U15644	Winson et al., 1995; Latifi et al 1995; Ochsner and Reiser, 1995.
<i>Pseudomonas aureofaciens</i>	<i>PhzI</i>	<i>PhzR</i>	unknown	L32729, L33724	Wood and Piersen, 1996
<i>Rhizobium leguminosarum</i>	unknown	<i>RhiR</i>	<i>N</i> -(3-hydroxy)-tetradecanoyl-L-homoserine lactone (HtDeHL)	M98835	Fuqua et al., 1994; Gray et al., 1996.
<i>Serratia liquefaciens</i>	<i>SwrI</i>	unknown	BHL	U2823	
<i>Aeromonas hydrophila</i>	<i>ahyI</i>	<i>ahyR</i>	BHL		Swift et al., 1997
<i>Aeromonas salmonicida</i>	<i>Asal</i>	unknown	BHL, <i>N</i> -hexanoyl-L-homoserine lactone		Swift et al., 1997
<i>Vibrio</i>	<i>vanI</i>	<i>vanR</i>	<i>N</i> -(3-oxo-		Milton et al.,

<i>anguillarum</i>			decanoyl)-L-homoserine lactone (ODHL)		1997
<i>Vibrio harveyi</i>	<i>LuxLM</i>	<i>LuxN</i>	<i>N</i> -(3-hydroxy)-butanoyl-L-homoserine lactone (HBHL)	L13940	Meighen, 1994; Bassler <i>et al.</i> , 1994.
<i>Yersinia enterocolitica</i>	<i>YenI</i>	<i>YenR</i>	OHHL,HHL	X76082	Throup <i>et al.</i> , 1996.

- References:
- Bassler et.al. *Molecular Microbiology*, **12**, 403-412 (1994)
- Beck et.al. *J.Bacteriol.*, **177**, 5000-5008 (1995)
- Devine et.al. *Biochemistry*, **27**,837-842 (1988)
- Fuqua et.al. *J.Bacteriol.* **176**, 269-275 (1994)
- Gray et.al. *J.Bacteriol.* **178**, 372-376 (1996)
- Hwang et.al. *J.Biotech.* **177**, 449-458 (1995)
- Latifi et.al. *Molecular Microbiology*, **17**, 333-343(1995)
- McGowan et.al.*Microbiology*, **141**, 541-550 (1995)
- Meighen *Ann. Rev. Genet.* **28**,117-139(1994)
- Milton et.al. *J.Bacteriol.* **179**, 3004-3012 (1994)
- Ochsner and Reiser *Proc.Natl.Acad.Sci.USA*, **92**, 6424-6428 (1995)
- Sitnikov et.al. *Molecular Microbiology*, **17**, 801-812 (1995)
- Swift et.al. *Molecular Microbiology*, **10**, 511-520 (1993)
- Swift et.al. *J.Bacteriol.* **179**, 5271-5281 (1997)
- Throup et.al. *Molecular Microbiology*, **17**, 345-356 (1996)
- Winson et.al. *Proc.Natl.Acad.Sci USA*, **92**, 9427-9431 (1995)
- Wood and Piersen, *Gene* **168**, 49-53 (1996)

Our invention is founded on our reasoning that if the inoculating bacteria were to encounter levels of *N*-acyl-L-homoserine lactone that gave a false indication of the local population size, the course of the ensuing infection would be drastically reduced.

A second aspect of the invention concerns engineering the plant to take advantage of the potential protective effect of antifungal rhizobacteria. There exist in the rhizosphere certain bacteria which are capable of attacking potential pathogenic fungal microorganisms which are also present in the soil, perhaps the best known of which are certain strains of *Pseudomonas fluorescens* and *P. aureofaciens*. But the population of such antifungal bacterial strains in the soil will normally be low and their antifungal activity dependent on the quorum sensing phenomenon to be activated. By imparting to the plant the ability to produce the activator molecule, the *N*-acyl-L-homoserine lactone, appropriate to the antifungal bacteria the antifungal activity may be initiated at low colony size providing earlier than normal protection of the plant against the pathogenic fungi. The rhizosphere-



expressed genes of the *rhiABC* operon of the symbiotic nitrogen-fixing bacterium *Rhizobium leguminosarum*, for example, are regulated by an AHL with a C14 side chain containing hydroxylated carbon in the 3 position and a single carbon-carbon double bond.

Transgenic plants producing an AHL signal molecule enhance the establishment of an antifungal environment on the rhizosphere. This phenomenon would also enable the use of disarmed bacterial strains to be used as crop protection biocontrol agents in conjunction with the AHL-producing transgenic plants.

The invention will now be described in the following Examples

The ability of AHLs to induce changes in neighbouring bacteria was tested in four ways;

(1) the ability of the AHLs to diffuse out of intact leaves was demonstrated by placing intact transgenic leaves on agar and subsequently removing it before overlaying with *C.violaceum* CV026 (see Example 4 below) and the outline of the whole leaf could be seen showing that the AHL diffused out of the leaf surface and not just the cut stem;

(2) being interested in whether the AHLs were only produced in the chloroplasts or whether they could be found in other tissues such as roots, the ability of the AHLs to diffuse from the roots was demonstrated in that AHLs in the vicinity of the roots were able to induce bioluminescence in a recombinant *E.coli* strain carrying an AHL-inducible operon: this showed that the root plastids are competent to synthesise the AHLs are, alternatively, that the AHLs synthesised in green tissue can be transported to the roots but in either case the roots were clearly capable of signalling to nearby bacteria.

(3) AHL-producing plant tissue is capable of restoring *G.graminis* growth-inhibiting activity to the disarmed *P.aureofaciens* 30-84 *phzI*- strain (see Example 9 below)

(4) *Erwinia carotovora carI (exl)* mutants, which have greatly reduced virulence in their natural host plants were shown to infect transgenic tobacco plants which are producing AHLs (see Example 10 below).

Figure 1 herewith shows the components of the constructs pBDHEYI and pBDHERBYI described in the Examples.

### **Example 1**

#### **Preparation of pBDHEYI**

pBDHEYI was constructed by fusing the alfalfa mosaic virus (AMV) translation enhancer sequence from pBI526 (Datla et.al., *Plant Science* **94**, 139-149 (1993)) to the *yenI*

coding sequence from *Yersinia enterocolitica*. The *yenI* sequence had previously been amplified by PCR to create an *NcoI* site overlapping the translation initiation sequence. This changed the second amino acid from leucine to valine but did not affect the ability of the encoded enzyme to synthesise *N*-acyl-L-homoserine lactones in a bacterial assay. The AMV/*yenI* fusion was cloned on a *BglIII/BamHI* fragment into the *BamHI* of pDH51 (Pietrzak *et al.*, *Nucl. Acids Res.* **14**, 5857-5868(1986)) to give pDHEYI. An *EcoRI* fragment of pDHEYI was cloned into the *EcoRI* site of pBIN19 (Bevan, *Nucl. Acids Res.* **12**, 8711-8721 (1984)) to give pBDHEYI.

### Example 2

#### 10 **Preparation of pBDHERBYI**

pBDHERBYI was constructed by fusing the petunia SSU611 ribulose biphosphate carboxylase small subunit (*rbcS*) chloroplast targeting sequence (Dean *et al.* *Mol. Gen. Genet.*, 206, 465-474 (1987)) to the AMV translation enhancer sequence of pBI526. An *NcoI* site was engineered to overlap the initiating ATG codon of *rbcS*. An *SphI* site was engineered to overlap the initiating ATG codon of *yenI* and the *yenI* coding sequence cloned into the *SphI* site of the SSU611 fragment. This site spans the cleavage site of the encoded chloroplast transit peptide. The AMV/*rbcS/yeI* fusion was cloned on a *BglIII/BamHI* fragment into the *BamHI* site of pDH51 to give pDHERBYI. An *EcoRI* fragment from pDHERBYI was cloned into the *EcoRI* site of pBIN19 to give pBDHERBYI.

20 The rationale for producing pBDHERBYI and believing that it would be active in chloroplasts was as follows: in *E.coli* homoserine lactone is not produced by mutants of the threonine biosynthetic pathway that are blocked prior to homoserine synthesis but is produced by those mutants when supplied with an exogenous source of homoserine. However, *TraI*, the *N*-acyl-L-homoserine lactone biosynthetic enzyme in *Agrobacterium tumefaciens*, has been found to utilise *S*-adenosylmethionine and not homoserine as a substrate *in vitro*. There is also evidence for the acyl moiety being derived from fatty acid biosynthetic intermediates. In plants the enzymes of the threonine biosynthetic pathway are located in the chloroplast and this organelle is also active in fatty acid metabolism. Therefore the chloroplasts may be expected to contain the necessary precursors for *N*-acyl-L-homoserine lactone synthesis by *yenI* and more closely approximate to the environment in which *yenI* is normally active than would be the cytoplasm.

### Example 3

#### **Generation of Transgenic Plants**

Construct pBDHEYI for Example 1 and pBDHERBYI from Example 2 were transferred to the *Agrobacterium tumefaciens* strain LBA 4404 and used to transform tobacco leaf discs according to standard protocol (Draper et.al., pages 69-160, In Plant Genetic Transformation and gene expression: a laboratory manual; Draper et.al. (Eds) Blackwell Scientific Publications, London (1988)) .

The transgenic status of the resulting kanamycin positive explants was confirmed by Southern analysis (data not given)

#### **Example 4**

##### **Complementation of Violacein Production**

Leaf segments of the transgenic plants produced in Example 3 were tested for their ability to synthesise *N*(3-oxohexanoyl)-L-homoserine lactone or a related analogue.

A transgenic tobacco leaf was placed in an agar plate overnight. The leaf was then removed and the *cviI* mutant of *Chromobacterium violaceum* spread over the plate. Violacein production by the bacteria could be seen where the *N*-(3-oxohexanoyl)-L-homoserine lactone had diffused out of the leaf and into the agar.

Two leaf segments tested positive as indicated by the ability of a diffusible product to complement *C.violaceum*, inducing the production of the purple pigment violacein by the bacteria.

#### **Example 5**

##### **Complementation of *carI***

Construct pBDHERBYI (Example 2) was transferred to the *Agrobacterium tumefaciens* strain LBA 4404 and transformed into tobacco. Leaf segments were tested for their ability to synthesise *N*(3-oxohexanoyl)-L-homoserine lactone or a related analogue.

An untransformed control and a transgenic BDHERBYI tobacco leaf were inoculated with *Erwinia carotovora* mutant for *carI*. The bacteria were applied at a high culture density (OD600 of 2.5) in a volume of 10 µl to a small wound site made with a hypodermic needle. A second BDHERBYI leaf was mock inoculated with bacterial culture medium alone.

The leaves were inspected after four days. The untransformed control and the mock inoculated leaf remained substantially unchanged. The sample inoculated with *E.carotovora* displayed advanced disease symptoms demonstrating that the pathogen can perceive and respond to the *N*-acyl-L-homoserine lactone being made by the transgenic plant.

#### **Example 6**

##### **Complementation of *luxI***

Following a similar protocol as described above, the *luxR* *N*-acyl-L-homoserine lactone response regulator and the *lux* operon (minus *luxI*) of *Pseudomonas fischeri* was inserted into *E.coli*. When transgenic tobacco carrying the BDHERBYI construct was challenged with the *E.coli*, bioluminescence was induced in the bacteria demonstrating that the *luxR* gene was able to respond to the *N*-acyl-L-homoserine lactone produced by the plant.

Twenty-nine tobacco plants that were independently transformed with either BDHERBYI or BDHEYI were challenged with *C.violaceum* mutant for *cviI* (Example 3) and *E.coli* carrying an *N*-acyl-L-homoserine lactone-inducible *lux* operon. Table I summarises the results.

TABLE 1					
Number of plants					
		Positive reaction		Negative reaction	
Construct	<i>cviI</i>	<i>luxI</i>	<i>cviI</i>	<i>luxI</i>	Total
BDHERBYI	8	8	5	5	13
BDHEYI	0	0	16	16	16

10

### Example 7

#### **Extraction and TLC analysis of AHLs**

For thin-layer chromatographic analysis, transgenic plant extracts were made by grinding two grams of plant tissue to a fine powder in liquid nitrogen and mixing the frozen powder with 200ml of warm distilled water. After five minutes, solid matter was filtered off and the filtrate extracted with an equal volume of ethyl acetate. The ethyl acetate layer was then dried over anhydrous magnesium sulphate, filtered and evaporated to dryness. The residue was taken up in 500  $\mu$ l of acetonitrile and 20  $\mu$ l of this was applied to a C18 reverse phase TLC plate (Merck). A similar extract from an untransformed control plant was also spotted on to the plate. *N*-hexanoyl-L-homoserine lactone (HHL) ( $1 \times 10^{-8}$ g) and *N*-(3-oxohexanoyl)-L homoserine lactone (OHHL) ( $5 \times 10^{-7}$ g) were applied as standards and the chromatogram developed with methanol/water (60:40 vol/vol) as running solvent (Shaw P.D. *et.al.* Proc.Natl.Acad.Sci. USA **94**: 6036-6041 (1997)). After drying, AHLs were located on the TLC plate by overlaying *C.violaceum* strain CV026 in top agar as described by McClean *et.al.* (Microbiology-UK, 143: 3703-3711 (1997)). After 16 hours growth at 28°C the presence of AHLs was indicated by localised violacein production.

25

Two different molecules with  $R_f$  values identical to the synthetic HHL and OHHL standards were observed.

### **Example 8**

#### **HPLC and LC-MS Analyses**

5 For HPLC and LC-MS analyses, transgenic plant extracts were made by grinding the issue in ethyl acetate. The supernatant was taken and the plant residue re-extracted with ethyl acetate, the supernatants pooled and the process repeated until the plant residue was white/brown n colour and free of chlorophyll. The ethyl acetate layer was separated from a small plant-derived aqueous layer and dried over anhydrous magnesium sulphate, filtered,  
10 and evaporated to dryness. The residue was resuspended in 500  $\mu$ l of methanol, this was brought to 60% methanol with sterile distilled water and placed at -20°C overnight to precipitate out the majority of the chlorophyll. After pelleting any solid matter by centrifugation in a bench-top microfuge, the AHL-containing supernatant was partitioned against 10 volumes of ethyla acetate and the organic phase evaporated to dryness. The  
15 residue was taken up into 500 $\mu$ l of acetonitrile. For both LC-MS and HPLC analyses linear gradients of acetonitrile in water were run (20-100% over 32 minutes) as described by Camara *et.al.* In Methods in Microbiology: Bacterial Pathogenesis Vol. 27: 319-330, Williams *et.al.* (Eds) (1998). OHHL and HHL eluted at 9 minutes and 13.5 minutes respectively.

20 The presence of HHL and OHHL, detected in TLC analysis, were confirmed.

### **Example 9**

#### **Assay for restoration of activity to *P.aureofaciens* mutant**

Leaf material from transgenic and non-transformed control plants were placed in wells cut in a potato dextrose agar plate (Oxoid). *P.aureofaciens* strain 80-84I (*phzI*<sup>-</sup>) was inoculated adjacent to  
25 the wells and the plates incubated for 24 hours at 22°C. The *G.graminis* var. tritici was then introduced on the opposite side of the plate and the whole incubated for a further four days.

The antifungal activity of the *P.aureofaciens phzI*<sup>-</sup> strain against the *G.graminis* was found to have been restored.

### **Example 10**

#### **Assay for restoration of virulence to *Erwinia carotovora* avirulent mutant**

30 Untransformed and control BDHERBYI tobacco leaves were inoculated with the avirulent *E.carotovora* mutant PNP22 (Bainton *et.al.*, Biochem.Journal, **288**: 997-1004

(1992) and also Jones *et.al.*, The EMBO Journal, 12: 2477-2482 (1993)) The bacteria were applied at high culture density (OD<sub>600</sub>=2.5) in a volume of 10µl to a small wound site made with a hypodermic needle.

Normally these *Erwinia* mutants are avirulent in the tobacco system, in which they  
5 can neither macerate plant tissue nor multiply *in planta* because they are defective in the production of plant cell-wall-degrading enzymes pectin lyase, pectate lyase, polygalacturonase, cellulase and protease. The regulated expression of plant cell wall-degrading enzymes only at high density in wild-type bacteria may contribute to the success of *Erwinia* as a plant pathogen. Under aerobic conditions, *E.carotovora* infection only  
10 occurs when the bacteria has reached sufficiently high population density such that disease progression depends on competition between bacterial multiplication and the plant host defences. Thus the production of macerating enzymes at low cell densities would not give rise to a successful infection, but would result in the premature induction of the local and systemic plant defence response, which in turn would hamper subsequent infection. Thus, if  
15 the infecting pathogen were to encounter AHL levels that gave a false indication of the local bacterial population size the course of the ensuing infection will be substantially reduced as the plant is able to mount a successful defence to a weak attack.

Other modifications of the present invention will be apparent to those skilled in the art without departing from the scope of the invention.

## CLAIMS

5

1. A method of protecting a plant against bacterial infection and/or virus infection transmitted by bacteria, comprising introducing into the genome of the plant by transformation the ability to synthesise a *N*-acyl-L-homoserine lactone.

10

2. A method of protecting a plant against bacterial infection and/or virus infection transmitted by bacteria, comprising introducing into the genome of the plant by transformation the ability to synthesise an analogue of *N*-acyl-L-homoserine lactone capable of competing with the *N*-acyl-L-homoserine lactone secreted by infecting bacteria for *N*-acyl-L-homoserine lactone receptor sites therein.

15

3. A method of enhancing interaction between a rhizobacterium and a plant comprising introducing into the genome of the plant by transformation the ability to synthesise the *N*-acyl-L-homoserine lactone naturally produced by the rhizobacterium.

20

4. A method as claimed in any of claims 1 to 3 in which the gene expressing the *N*-acyl-L-homoserine lactone is selected from the group consisting of, the *yenI* gene of *Yersinia enterocolitica*; the *cviI* gene of *Chromobacterium violaceum*; the *luxI* gene of *Photobacterium fischeri*; the *carI* gene of *Erwinia carotovora*; the *traI* gene of *Agrobacterium tumefaciens* and the *lasI* and *vsmI* genes of *Pseudomonas aeruginosa*.

25



5. A recombinant plant genome containing a gene construct for *in planta* expression of an *N*-acyl-L-homoserine lactone and/or the response regulator thereof.

30

6. A genome as claimed in claim 5 in which expression of the said *N*-acyl-L-homoserine lactone is targeted to plant chloroplasts.

1/1

Figure 1



**pBDHERBYI**



**INTERNATIONAL SEARCH REPORT**

International Application No  
PCT/GB 99/02652

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C12N15/31 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07K C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category <sup>o</sup>	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 195 48 301 C (DUERING KLAUS DR) 27 February 1997 (1997-02-27) the whole document ---	1-6
A	WO 97 48719 A (BECKERMAN JANNA L ; TEXAS A & M UNIVERSITY SYST (US); ZHANG LEI (US)) 24 December 1997 (1997-12-24) the whole document ---	1-6
A	WO 96 29392 A (UNISEARCH LTD ; KJELLEBERG STAFFAN (AU); STEINBERG PETER (AU); NYS) 26 September 1996 (1996-09-26) the whole document ---	2
	-/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

<sup>o</sup> Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

2 November 1999

17/11/1999

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 99/02652

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GRAY K. ET AL.: "Cell-to cell signaling in the symbiotic nitrogen-fixing bacterium <i>Rhizobium leguminosarum</i> : autoinduction of a stationary phase and rhizosphere-expressed genes" JOURNAL OF BACTERIOLOGY, vol. 178, no. 2, 1996, pages 372-376, XP002084285 the whole document ----	3
A	ROSEMEYER ET AL: "luxI- and luxR-homologous genes of <i>Rhizobium etli</i> CNPAF512 contribute to synthesis of autoinducer molecules and nodulation of <i>Phaseolus vulgaris</i> " JOURNAL OF BACTERIOLOGY, vol. 180, no. 4, 1 February 1998 (1998-02-01), pages 815-821, XP002084284 ISSN: 0021-9193 ----	3
A	THROUP J. ET AL.: MOLECULAR MICROBIOLOGY, vol. 17, 1996, pages 345-56, XP002121181 cited in the application the whole document ----	1-6
A	ROBSON N D ET AL: "Bacterial N-acyl-homoserine-lactone-dependent signalling and its potential biotechnological applications" TRENDS IN BIOTECHNOLOGY, vol. 15, no. 11, 1 November 1997 (1997-11-01), page 458-464 XP004092668 ISSN: 0167-7799 see the whole document; esp. p.461 r. col. ----	1-6
A	SWIFT S ET AL: "Quorum sensing: a population-density component in the determination of bacterial phenotype" TIBS TRENDS IN BIOCHEMICAL SCIENCES, vol. 21, no. 6, 1 June 1996 (1996-06-01), page 214-219 XP004050894 ISSN: 0968-0004 cited in the application the whole document -----	1-6

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 99/02652

Patent document cited in search report	C	Publication date	Patent family member(s)	Publication date
DE 19548301	C	27-02-1997	NONE	
WO 9748719	A	24-12-1997	AU 3570997 A	07-01-1998
WO 9629392	A	26-09-1996	AU 708962 B	19-08-1999
			AU 4999696 A	08-10-1996
			BR 9607661 A	16-06-1998
			CA 2215797 A	26-09-1996
			CN 1185173 A	17-06-1998
			EP 0815201 A	07-01-1998
			JP 11502108 T	23-02-1999
			NZ 303630 A	26-01-1998