



US006121416A

United States Patent [19]

[11] Patent Number: 6,121,416

Clark et al.

[45] Date of Patent: Sep. 19, 2000

- [54] **INSULIN-LIKE GROWTH FACTOR AGONIST MOLECULES**
- [75] Inventors: **Ross G. Clark**, Pacifica; **Henry B. Lowman**, El Granada, both of Calif.; **Iain C. A. F. Robinson**, St. Albans, United Kingdom
- [73] Assignee: **Genentech, Inc.**, South San Francisco, Calif.
- [21] Appl. No.: **08/825,852**
- [22] Filed: **Apr. 4, 1997**
- [51] **Int. Cl.**⁷ **C07K 14/00**; C07K 14/65; A61K 38/04; A61K 38/16
- [52] **U.S. Cl.** **530/326**; 530/300; 530/303; 530/321; 530/324; 530/325; 530/327; 530/328; 514/12; 514/13; 514/14; 514/15
- [58] **Field of Search** 530/300, 303, 530/321, 324, 325, 326, 327, 328; 514/12, 13, 14, 15

- WO 89/09792 10/1989 WIPO .
- WO 91/03253 3/1991 WIPO .
- WO 92/11865 7/1992 WIPO .
- WO 92/15679 9/1992 WIPO .
- WO 93/08826 5/1993 WIPO .
- WO 93/20836 10/1993 WIPO .
- WO 93/23071 11/1993 WIPO .
- WO 93/25219 12/1993 WIPO .
- WO 94/04569 3/1994 WIPO .
- WO 94/16722 8/1994 WIPO .
- WO 94/16723 8/1994 WIPO .
- WO 95/07697 3/1995 WIPO .
- WO 95/17422 6/1995 WIPO .
- WO 95/17423 6/1995 WIPO .
- WO 96/01124 1/1996 WIPO .
- WO 96/15148 5/1996 WIPO .
- WO 96/33216 10/1996 WIPO .
- WO 96/37514 11/1996 WIPO .
- WO 96/40189 12/1996 WIPO .
- WO 96/40750 12/1996 WIPO .
- WO 97/37010 10/1997 WIPO .
- WO 97/39032 10/1997 WIPO .

OTHER PUBLICATIONS

- Humbel E.R. Insulin-like growth factors I and II. *European J. Biochem.* 190:445-462 (1990).
- Rudinger, J. "Characteristics of the amino acids as components of a peptide hormone sequence", in *Peptide Hormones*, University Park Press, Jun. 1976.
- Alberts et al., *Molecular Biology of the Cell*, 1994, Garland Publishing, Inc., New York, NY.
- Bowie et al. Deciphering the message in protein sequences: tolerance to amino acid substitutions. *Science*, (Mar. 16, 1990) 247 (4948) 1306-10.
- Brandt et al. Role of natriuretic peptide clearance receptor in vivo control of C-type natriuretic peptide. *Am. J. Physiol.*, (Jul. 1995) 269 (1 Pt 2) H326-31.
- Baxter et al. Structural determinants for binary and ternary complex formation between insulin-like growth factor-I (IGF-I) and IGF binding protein-3. *J. Biol. Chem.*, (Jan. 5, 1992) (1) 60-5.
- Cascieri et al. Analysis of the interaction of IGF-I analogs with the IGF-I receptor and IGF binding proteins. *Adv Exp Med Biol* 1993;343-33-40.
- Clemmons et al. Competition for binding to insulin-like growth factor (IGF) binding protein-2, 3, 4, and 5 by the IGFs and IGF analogs. *Endocrinology* Aug. 1992;131(2):890-895.
- Heding et al. Biosensor measurement of the binding of insulin-like growth factors I and II and their analogues to the insulin-like growth factor-binding protein-3. *J Biol Chem* Jun. 14 1996;271(24):13948-13952.

(List continued on next page.)

[56] **References Cited**

U.S. PATENT DOCUMENTS

- 4,411,890 10/1983 Momany 424/177
- 4,876,242 10/1989 Applebaum 514/3
- 4,988,675 1/1991 Froesch et al. 514/4
- 5,068,224 11/1991 Fryklund et al. 514/21
- 5,077,276 12/1991 Ballard et al. 514/12
- 5,093,317 3/1992 Lewis et al. 514/12
- 5,106,832 4/1992 Froesch et al. 514/3
- 5,126,324 6/1992 Clark et al. 514/12
- 5,164,370 11/1992 Ballard et al. 514/12
- 5,187,151 2/1993 Clark et al. 514/3
- 5,202,119 4/1993 Clark et al. 424/88
- 5,206,235 4/1993 Fisher et al. 514/213
- 5,210,017 5/1993 Carlsson et al. 435/7.8
- 5,223,409 6/1993 Ladner et al. 435/69.7
- 5,273,961 12/1993 Clark 514/8
- 5,374,620 12/1994 Clark et al. 514/12
- 5,466,670 11/1995 Dunger et al. 514/12
- 5,470,828 11/1995 Ballard et al. 514/12
- 5,534,617 7/1996 Cunningham et al. 530/399
- 5,569,648 10/1996 Lewis et al. 514/12
- 5,593,844 1/1997 Carlsson et al. 435/7.1
- 5,597,797 1/1997 Clark 514/12
- 5,622,932 4/1997 DiMarchi et al. 514/12
- 5,652,214 7/1997 Lewis et al. 514/12
- 5,703,045 12/1997 Lewis et al. 514/12
- 5,714,460 2/1998 Gluckman et al. 514/3
- 5,776,897 7/1998 Lewis et al. 514/12

FOREIGN PATENT DOCUMENTS

- 128733 12/1984 European Pat. Off. .
- 135094 3/1985 European Pat. Off. .
- 230869 8/1987 European Pat. Off. .
- 288451 10/1988 European Pat. Off. .
- 294021 12/1988 European Pat. Off. .
- 369943 5/1990 European Pat. Off. .
- 375438 6/1990 European Pat. Off. .
- 379338 7/1990 European Pat. Off. .
- 681842 11/1995 European Pat. Off. .
- 742228 11/1996 European Pat. Off. .
- WO 89/05822 6/1989 WIPO .
- WO 89/08667 9/1989 WIPO .
- WO 89/09268 10/1989 WIPO .

Primary Examiner—Elizabeth Kemmerer
Assistant Examiner—David S. Romeo
Attorney, Agent, or Firm—Janet E. Hasak

[57] **ABSTRACT**

Compounds are provided that prevent the interaction of an IGF with any one of its binding proteins and not to a human IGF receptor. These IGF agonist compounds, which include peptides, are useful to increase serum and tissue levels of active IGFs in a mammal

9 Claims, 44 Drawing Sheets

OTHER PUBLICATIONS

- Maack et al. Physiological role of silent receptors of atrial natriuretic factor. *Science*, Oct. 30, 1987, vol. 238, pp. 675–678.
- Bach and Rechler, “Insulin-like Growth Factor Binding Proteins” *Diabetes Reviews* 3:38–61 (1995).
- Ballard et al., “Does IGF-I Ever Act Through the Insulin Receptors” *The Insulin-like Growth Factors and Their Regulatory Proteins*, Baxter, eds., Amsterdam: Elsevier pp. 131–138 (1994).
- Bar et al., “Tissue localization of perfused endothelial cell IGF binding protein is markedly altered by association with IGF-I” *Endocrinology* 127:3243–3245 (1990).
- Bass et al., “Hormone Phage: An Enrichment Method for Variant Proteins with Altered Binding Properties” *Proteins: Structure, Function, and Genetics* 8(4):309–314 (1990).
- Baxter, “Physiological Roles of IGF Binding Proteins” *Modern Concepts of Insulin-like Growth Factors*, Spencer, eds., Elsevier, New York pp. 371–380 (1991).
- Baxter, “The somatomedins: insulin-like growth factors” *Advances in Clinical Chemistry* 25:49–115 (1986).
- Baxter & Martin, “Binding Proteins for Insulin-Like Growth Factors in Adult Rat Serum. Comparison With Other Human and Rat Binding Proteins” *Biochem. & Biophys. Res. Comm.* 147(1):408–415 (1987).
- Baxter, Robert C., “The Insulin-Like Growth Factors and Their Binding Proteins” *Comp. Biochem. Physiol.* 91B(2):229–235 (1988).
- Bayne et al., “The C region of human insulin-like growth factor (IGF) I is required for high affinity binding to the type 1 IGF receptor” *Journal of Biological Chemistry* 264a:11004–11008 (1988).
- Bayne et al., “The roles of tyrosines 24, 31, and 60 in the high affinity binding of insulin-like growth factor-I to the type I insulin-like growth factor receptor” *Journal of Biological Chemistry* 265:15648–15652 (1990).
- Bayne et al., “Structural analogs of human insulin-like growth factor I with reduced affinity for serum binding proteins and the type 2 insulin-like growth factor receptor” *Journal of Biological Chemistry* 263:6233–6239 (1988).
- Binkert et al., “Cloning, sequence analysis and expression of a cDNA encoding a novel insulin-like growth factor binding protein (IGFBP-2)” *EMBO Journal* 8:2497–2502 (1989).
- Binoux, M., “Donnees recentes sur les somatomedines” *Annales d'Endocrinologie* 41:157–191 (1980) English abstract only.
- Bondy, C., “Clinical uses of insulin-like growth factor I” *Annals of Internal Medicine* 120:593–601 (1994).
- Bowers, C. Y., “GH Releasing Peptides—Structure and Kinetics” *J. Pediatr. Endocrinology* 6(1):21–31 (1993).
- Brewer et al., “Cloning, Characterization, and Expression of a Human Insulin-Like Growth Factor Binding Protein” *Biochem. & Biophys. Res. Comm.* 152(3):1289–1297 (1988).
- Brinkman et al., “Isolation and characterization of a cDNA encoding the low molecular weight insulin-like growth factor binding protein (IBP-1)” *The EMBO J.* 7:2417–2423 (1988).
- Carlsson et al., “Growth Hormone and Growth in Diabetic Rats: Effects of Insulin and Insulin-Like Growth Factor-I Infusions” *J. Endocrinol.* 122:661–670 (1989).
- Cascieri et al., “Mutants of human insulin-like growth factor I with reduced affinity for the type 1 insulin-like growth factor receptor” *Biochemistry* 27:3229–3233 (1988).
- Cascieri et al., “Structural analogs of human insulin-like growth factor (IGF) I with altered affinity for type 2 IGF receptors” *Journal of Biological Chemistry* 264:2199–2202 (1989).
- Charlton et al., “Growth hormone-deficient dwarfism in the rat: a new mutation” *J. of Endocrinology* 119:51–58 (1988).
- Cheetham et al., “The Effects of Recombinant Human Insulin-like Growth Factor I on Growth Hormone Secretion in Adolescents With Insulin Dependent Diabetes Mellitus” *Clin. Endocrinol.* 40:515–522 (1994).
- Cheetham et al., “The Effects of Recombinant Insulin-like Growth Factor I Administration on Growth Hormone Levels and Insulin Requirements in Adolescents With Type 1 (Insulin-dependent) Diabetes Mellitus” *Diabetologia* 36:678–681 (1993).
- Chen et al., “Recombinant human IGF-I infusion results in transient improvement in nitrogen balance: evidence for IGF-I autoregulation” *US Endocrine Meeting* (Abstract 1596) pp. 449 (1993).
- Clark et al., “Growth-Responses to Patterned GH Delivery” *Endocrine* 3:717–723 (1995).
- Clark et al., “Insulin-Like Growth Factor and Growth Hormone (GH) Have Distinct and Over-Lapping Effects in GH-Deficient Rats” *Endocrine* 3:297–304 (1995).
- Clemmons and Van Wyk, “Somatomedin: physiological control and effects on cell proliferation” *Handbook Exp. Pharmacol.* 57:161–208 (1981).
- Clemmons et al., “Discrete Alterations of the Insulin-like Growth Factor I Molecule Which Alter Its Affinity for Insulin-like Growth Factor-binding Proteins Result in Changes in Bioactivity” *Journal of Biological Chemistry* 265(21):12210–12216 (1990).
- Cwirla et al., “Peptides on phage: a vast library of peptides for identifying ligands” *Proc. Natl. Acad. Sci. USA* 87(16):6378–6382 (1990).
- Devlin et al., “Random peptide libraries: a source of specific protein binding molecules” *Science* 249:404–406 (1990).
- Duerr et al., “Insulin-like growth factor-1 enhances ventricular hypertrophy and function during the onset of experimental cardiac failure” *J. Clin. Invest.* 95:619–627 (1995).
- Elahi et al., “Hemodynamic and metabolic responses to human insulin-like growth factor I (IGF-I) in men” *Modern Concepts of Insulin-Like Growth Factors*, Spencer, EM, ed., New York:Elsevier Science Publ. Co. pp. 219–224 (1991).
- Fielder et al., “Differential long-term effects of insulin-like growth factor-I (IGF-I) growth hormone (GH), and IGF-I plus GH on body growth and IGF binding proteins in hypophysectomized rats” *Endocrinology* 137:1913–1920 (1996).
- Franklin et al., “Insulin-Like Growth Factor I Preserves Renal Function Postoperatively” *Am. J. Physiol.* 272:F257–F259 (1997).
- Froesch et al., “Metabolic and Therapeutic Effects of Insulin-Like Growth Factor I” *Horm. Res.* 42:66–71 (1994).
- Furnsinn et al., “Insulin-Like Growth Factor-I Inhibits Insulin and Amylin Secretion in Conscious Rats” *Endocrinology* 135(5):2144–2149 (1994).
- Geysen et al., “A priori delineation of a peptide which mimics a discontinuous antigenic determinant” *Molecular Immunology* 23(7):709–715 (1986).
- Ghazzi et al., “Cardiac and glycemic benefits of troglitazone treatment in NIDDM. The Troglitazone Study Group” *Diabetes* 46:433–439 (1997).

- Giebel et al., "Screening of cyclic peptide phage libraries identifies ligands that bind streptavidin with high affinities" *Biochemistry* 34:15430–15435 (1995).
- Guler et al., "Effects of Insulin-like Growth Factor I in Man" *Acta Paediatr. Scand.* 367:52–54 (Suppl. 1990).
- Guler et al., "Effects of recombinant insulin-like growth factor I on insulin secretion and renal function in normal human subjects" *Proc. Natl. Acad. Sci. USA* 86:2868–2872 (Apr. 1989).
- Guler et al., "Insulin-like growth factor I increases glomerular filtration rate and renal plasma flow in man" *Acta Endocrinologica* 121:101–106 (1989).
- Guler et al., "Recombinant human insulin-like growth factor 1 stimulates growth and has distinct effects on organ size in hypophysectomized rats" *Proc. Natl. Acad. Sci. USA* 85:4889–4893 (1988).
- Guler et al., "Short-term metabolic effects of recombinant human insulin-like growth factor I in healthy adults" *New England J. of Medicine* 317(3):137–140 (1987).
- Hammerman and Miller, "The growth hormone insulin-like factor axis in kidney revisited" *Am. J. Physiol.* 265:F1–F14 (1993).
- Hammerman and Miller, "Therapeutic use of growth factors in renal failure" *J. Am. Soc. Nephrol.* 5:1–11 (1994).
- Hartman et al., "A low dose euglycemic infusion of recombinant human insulin-like growth factor I rapidly suppresses fasting-enhanced pulsatile growth hormone secretion in humans" *J. Clin. Invest.* 91:2453–2462 (1993).
- Hasegawa et al., "The free form of insulin-like growth factor I increases in circulation during normal human pregnancy" *J. Clin. Endocrinol. Metabol.* 80:3284–3286 (1995).
- Hirschberg et al., "Effects of insulin-like growth factor I on renal function in normal men" *Kidney International* 43:387–397 (1993).
- Hizuka et al., "Measurement of free form of insulin-like growth factor I in human plasma" *Growth Regulation* 1:51–55 (1991).
- Jabri et al., "Adverse effects of recombinant human insulin-like growth factor I in obese insulin-resistant type II diabetic patients" *Diabetes* 43:369–374 (1994).
- Johnson et al., "Underexpression of beta cell high Km glucose transporters in noninsulin-dependent diabetes" *Science* 250:546–549 (1990).
- Jones et al., "Insulin-Like Growth Factors and Their Binding Proteins: Biological Actions" *Endocr. Rev.* 16:3–34 (1995).
- Juul et al., "Serum concentrations of free and total insulin-like growth factor-I, IGF binding proteins -1 and -3 and IGFBP-3 protease activity in boys with normal or precocious puberty" *Clin. Endocrinology* 44:515–523 (1996).
- Kay et al., "An M13 phage library displaying random 38-amino-acid peptides as a source of novel sequences with affinity to selected targets" *Gene* 128:59–65 (1993).
- Kerr et al., "Effect of Insulin-like Growth Factor 1 on the Responses to and Recognition of Hypoglycemia" *American Diabetes Association (ADA), San Antonio, Texas, Jun. 20–23, 1992* (abstract #225), 52nd Annual Meeting edition pp. 60A.
- Kerr et al., "Effect of Insulin-like Growth Factor-1 on the Responses to and Recognition of Hypoglycemia in Humans" *J. Clin. Invest.* 91:141–147 (1993).
- Kupfer et al., "Enhancement of the anabolic effects of growth hormone and insulin-like growth factor I by use of both agents simultaneously" *J. Clin. Invest.* 91:391–396 (1993).
- Kuzuya et al., "Trial of insulinlike growth factor I therapy for patients with extreme insulin resistance syndromes" *Diabetes* 42:696–705 (1993).
- Lassale et al., "ESM-1 is a novel human endothelial cell-specific molecule expressed in lung and regulated by cytokines" *Journal of Biological Chemistry* 271:20458–20464 (1996).
- Leahy et al., "Insulin-Like Growth Factor-I at Physiological Concentrations is a Potent Inhibitor of Insulin Secretion" *Endocrinology* 126(3):1593–1598 (1990).
- Lee et al., "Insulin-Like Growth Factor (IGF) Binding Protein Complementary Deoxyribonucleic Acid from Human HEP G2 Hepatoma Cells: Predicted Protein Sequence Suggests an IGF Binding Domain Different from Those of the IGF-I and IGF-II Receptors" *Mol. Endocrinol.* 2(5):404–411 (1988).
- Lee et al., "Regulation and function of insulin-like growth factor-binding protein-1" *Proc. Soc. Exp. Biol. & Med.* 204:4–29 (1993).
- Leung et al., "Growth hormone receptor and serum binding protein: purification, cloning and expression" *Nature* 330:537–543 (1987).
- Lewitt and Baxter, "Insulin-like growth factor-binding protein-1: a role in glucose counterregulation?" *Mol. Cell. Endocrinology* 79(1–3):C147–152 (1991).
- Lewitt et al., "Bioavailability of insulin-like growth factors (IGFs) in rats determined by the molecular distribution of human IGF-binding protein-3" *Endocrinology* 133:1797–1802 (1993).
- Lewitt et al., "Insulin-like Growth Factor-binding Protein-1 Modulates Blood Glucose Levels" *Endocrinology* 129(4):2254–2265 (1991).
- Lieberman et al., "Anabolic effects of recombinant insulin-like growth factor I in AIDS-associated cachexia" *US Endocrine Meeting* (Abstract 1664) pp. 466 (1993).
- Lieberman et al., "Anabolic effects of recombinant insulin-like growth factor-I in cachectic patients with the acquired immunodeficiency syndrome" *J. Clin. Endocrinol. and Metab.* 78(2):404–410 (1994).
- Lieberman et al., "Effects of recombinant human insulin-like growth factor-I (rhIGF-I) on total and free IGF-I concentrations, IGF-binding proteins, and glycemic response in humans" *J. Clin. Endocrinol. and Metab.* 75(1):30–36 (1992).
- Livnah et al., "Functional mimicry of a protein hormone by a peptide agonist: the EPO receptor complex at 2.8 Å" *Science* 273:464–471 (1996).
- Martin & Baxter, "Insulin-like Growth Factor-binding Protein from Human Plasma" *Journal of Biological Chemistry* 261(10):8754–8760 (1996).
- McLafferty et al., "M13 bacteriophage displaying disulfide-constrained microproteins" *Gene* 128:29–36 (1993).
- Miller et al., "Effects of IGF-I on renal function in end-stage chronic renal failure" *Kidney Int.* 46:201–207 (1994).
- Morrow et al., "Recombinant Human (rh) IGF-1 Reverses Hyperglycemia and Improves Insulin Sensitivity in Severe Insulin Resistance" *Diabetes-53rd Annual Meeting, Jun. 12–15, 1993* (abstract No. 269) 42:83A (Suppl. 1 1993).
- O'Neil et al., "Identification of novel peptide antagonists for GPIIb/IIIa from a conformationally constrained phage peptide library" *Proteins* 14:509–515 (1992).
- O'Shea and Layish, "Growth hormone and the kidney: a case presentation and review of the literature" *J. Am. Soc. Nephrol.* 3:157–161 (1992).

- O'Shea et al., "Effects of IGF-I on renal function in patients with chronic renal failure" *Am. J. Physiol.* 264:F917-F922 (1993).
- Oh et al., "Characterization of the affinities of insulin-like growth factor (IGF)-binding proteins 1-4 for IGF-I, IGF-II, IGF-I/insulin hybrid, and IGF-I analogs" *Endocrinology* 132:1337-1344 (1993).
- Oh et al., "Synthesis and characterization of insulin-like growth factor-binding protein (IGFBP)-7. Recombinant human mac25 protein specifically binds IGF-I and -II" *Journal of Biological Chemistry* 271:30322-30325 (1996).
- Oldenburg et al., "Peptide ligands for a sugar-binding protein isolated from a random peptide library" *Proc. Natl. Acad. Sci.* 89:5393-5397 (1992).
- Quin et al., "Acute Response to Recombinant Insulin-like Growth Factor I in a Patient with Mendenhall's Syndrome" *New England J. of Medicine* 323:1425-1426 (1990).
- Rinderknecht and Humbel, "The amino acid sequence of human insulin-like growth factor I and its structural homology with proinsulin" *Journal of Biological Chemistry* 253(8):2769-2776 (1978).
- Rinderknecht and Humbel, "Polypeptides with nonsuppressible insulin-like and cell-growth promoting activities in human serum: isolation, chemical characterization, and some biological properties of forms I and II" *Proc. Natl. Acad. Sci USA* 73(7):2365-2369 (1976).
- Ross et al., "The Role of Insulin, Growth Hormone and IGF-I as Anabolic Agents in the Critically Ill" *Intensive Care Med.* 19(2):S54-57 (Suppl. 1993).
- Saad et al., "Low-doses of Inulin-like Growth Factor-I Improve Insulin Sensitivity" *Diabetologia* (Abstract) 37:A40 (Supp. 1 1994).
- Schalch et al., "Short-Term Effects of Recombinant Human insulin-Like Growth Factor I on Metabolic Control of Patients with Type II Diabetes Mellitus" *J. of Clinical Endocrinology & Metabolism* 77(6):1563-1568 (1993).
- Schalch et al., "Short-term metabolic effects of recombinant human insulin-like growth factor I Control of Patients with Type II Diabetes Mellitus" *J. of Clinical Endocrinology & Metabolism* 77(6):1563-1568 (1993).
- Schalch et al., "Short-term metabolic effects of recombinant human insulin-like growth factor I (rhIGF-I) in type II diabetes mellitus" *Modern Concepts of Insulin-Like Growth Factors*, Spencer, ed., New York:Elsevier Science Publ. Co. pp. 705-713 (1991).
- Schoen et al., "Growth Hormone Secretagogues" *Annual Reports in Medicinal Chemistry: Section IV-Endocrinology & Metabolic Diseases*, William K. Hagmann, Chapter 19, vol. 28:177-186 (1993).
- Schoenle et al., "Recombinant human insulin-like growth factor I(rhIGF I) reduces hyperglycaemia in patients with extreme insulin resistance" *Diabetologia* 34:675-679 (1991).
- Scott et al., "Searching for peptide ligands with an epitope library" *Science* 249:386-390 (1990).
- Sherwin et al., "Metabolic Effects of Insulin-like Growth Factor I in Normal Humans" *Horm. Res.* 41:97-101 (Suppl. 2 1994).
- Skottner et al., "Growth responses in a mutant dwarf rat to human growth hormone and recombinant human insulin-like growth factor I" *Endocrinology* 124(5):2519-2526 (1989).
- Smith, "Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface" *Science* 228(4705):1315-1317 (1985).
- Stern et al., "Insulin resistance and pancreatic insulin release in the" *Proc. Soc. Exp. Biol. Med.* 139:66-69 (1972).
- Suikkari et al., "Insulin regulates the serum levels of low molecular weight insulin-like growth factor-binding protein" *J. Clin. Endocrinology Metabol.* 66:266-272 (1988).
- Swisshelm et al., "Enhanced expression of an insulin growth factor-like binding protein (mac25) in senescent human mammary epithelial cells and induced expression with retinoic acid" *Proc. Natl. Acad. Sci.* 92:4472-4476 (1995).
- Takano et al., "Effects of sc Administration of Recombinant Human Insulin-like Growth Factor I (IGF-I) on Normal Human Subjects" *Endocrinol. Japan* 37(2):309-317 (1990).
- Tanner et al., "Comparative rapidity of response of height, limb muscle and limb fat to treatment with human growth hormone in patients with and without growth hormone deficiency" *Acta Endocrinologica* 84:681-696 (1977).
- Underwood et al., "Regulation of somatomedin-c/insulin-like growth factor I by nutrients" *Hormone Res.* 24:166-176 (1986).
- Usala et al., "Brief report: treatment of insulin-resistant diabetic ketoacidosis with insulin-like growth factor I in an adolescent with insulin-dependent diabetes" *New England J. of Medicine* 327(12):853-857 (1992).
- Uthne et al., "Effects of human somatomedin preparations on membrane transport and protein synthesis in the isolated rat diaphragm" *J. Clin. Endocrinol. Metab.* 39(3):548-554 (1974).
- Van Wyk et al., "The somatomedins: a family of insulinlike hormones under growth hormone control" *Recent Prog. Horm. Res.* 30:259-318 (1974).
- Vlachopapadopoulou et al., "Metabolic and Clinical Response to Recombinant Human Insulin-like Growth Factor I in Myotonic Dystrophy—A Clinical Research Center Study" *J. Clin. Endo. Metab.* 80(12):3715-3723 (1995).
- Wells and Lowman, "Rapid Evolution of Peptide and Protein Binding Properties in Vitro" *Curr. Opin. Struct. Biol.* 2:597-604 (1992).
- Wilton et al., "Treatment with recombinant human insulin-like growth factor I of children with growth hormone receptor deficiency (Laron syndrome)" *Acta Paediatr Suppl* 383:137-141 (1992).
- Wood et al., "Cloning and expression of the growth hormone-dependent insulin-like growth factor-binding protein" *Molecular Endocrinology* 2:1176-1185 (1988).
- Wood et al., "Crystal structure analysis of deamino-oxytocin: conformational flexibility and receptor binding" *Science* 232:633-636 (1986).
- Wrighton et al., "Small peptides as potent mimetics of the protein hormone erythropoietin" *Science* 273:458-464 (1996).
- Yamauchi et al., "Purification and molecular cloning of prostacyclin-stimulating factor from serum-free conditioned medium of human diploid fibroblast cells" *Biochemical Journal* 303(Part 2):591-598 (1994).
- Zenobi et al., "Effects of insulin-like growth factor-I on glucose tolerance, insulin levels, and insulin secretion" *J. Clin. Invest.* 89:1908-1913 (1992).
- Zenobi et al., "Insulin-like growth factor-I improves glucose lipid metabolism in type 2 diabetes mellitus" *J. Clin. Invest.* 90:2234-2241 (1992).
- Hampton et al., "Purification and Characterization of an Insulin-like Growth Factor II Variant from Human Plasma" *Journal of Biological Chemistry* 264(32):19155-19160 (Nov. 15, 1989).

Peterkofsky et al., "Elevated Activity of Low Molecular Weight Insulin-Like Growth Factor-Binding Proteins in Sera of Vitamin C-Deficient and Fasted Guinea Pigs" *Endocrinology* 128(4):1769-1779 (1991).

Kletzien et al., "Enhancement of adipocyte differentiation by an insulin-sensitizing agent" *Molecular Pharmacology* 41(2):393-398 (Feb. 1992).

Loddick et al., "Displacement of insulin-like growth factors from their binding proteins as a potential treatment for stroke" *Proc. Natl. Acad. Sci. USA* 95(4):1894-1898 (Feb. 17, 1998).

Lowman et al., "Molecular mimics of insulin-like growth factor 1 (IGF-1) for inhibiting IGF-1: IGF-binding protein interactions" *Biochemistry* 37(25):8870-8878 (Jun. 23, 1998).

TCACGTAAAA AGGGTATCTA GAATTATGAT GATTACTCTG CGCAAACTTC CTCTGGCGGT TGCCGTGCA GCGGGCGTAA TGCTCTCTCA GGCCATGGCC
 AGTGCATTTT TCCCATAGAT CTTAATACTA CTAATGAGAC GCGTTTGAAG GAGACCGCCA ACGGCAGCGT CGCCCGCATT ACAGACGAGT CCGGTACCGG
 MetMe tileThrLeu ArgLysLeuP rLeuAlaVa lAlaValAla AlaGlyValM etSerAlaG1 nAlaMetAla
 ^Start of lamB signal sequence

GGTCCCGAAA CTCTGTGCGG TGCTGAACTG GTTACCGCTC TGCAGTTCGT ATGTGGTGAT CGAGGCTTCC TGTTCAACAA ACCGACTGGG GCTGGATCCT
 CCAGGGCTTT GAGACACGCC ACGACTTGAC CAACTGCGAG ACGTCAAGCA TACACACTA GCTCCGAAGG ACAAGTTGTT TGGCTGACCC CGACCTAGGA
 GlyProGluT hrLeuCysG1 YAlaGluLeu ValAspAlaL euGlnPheVa lCysGlyAsp ArgGlyPheL euPheAsnLy sProThrGly AlaGlySerSer
 ^Start of IGF-I (Y24L, Y31A)

CCTCTCGTCG TGCTCCCGAG ACTGGTATTG TTGACGAAAG CTGCTTTCGT TCTTGGGACC TGCGTCGTCT GGAAATGTAT TGCGCTCCCC TGAACCCCGC
 GGAGAGCAGC ACGAGGGGTC TGACCATAAC AACTGCTTAC GACGAAAGCA AGAACGCTGG ACGCAGCAGA CCTTTACATA ACGCGAGGGG ACTTTGGGCG
 SerArgAr gAlaProGln ThrGlyIlev alAspGlucy sCysPheArg SerCysAspL euArgArgLe uGluMetTyr CysAlaProL euLysProAla

TAAATCTGCT TAGAAGCTCC TAACGCTCGG TTGCCCGCCGG GCGTTTTTTA TTGTTAACTC ATGTTTGACA GCTTATCATC GATAAGCTTT AATGCGGTAG
 ATTTAGACGA ATCTTCGAGG ATTGCGAGCC AACGGCGGCC CGCAAAAAAT AACCAATTGAG TACAAAACCTGT CGAATAGTAG CTATTCGAAA TTACGCCCATC
 LysSerAla Am*

Nucleotide and Amino Acid Sequence of the LamB Signal Sequence and IGF-I (Y24L, Y31A)

FIG. 1




FIG. 2

plasmid IGfMI
length: 5115 (circular)

```

1  GAAATCAACT TCTCCATACT TTGGATAAGG AAATACAGAC ATGAAAAAATC TCATTGCTGA GTTGTATATT AAGCTTGCCC AAAAAAGAAG AGAGTCGAAT
   CTTAAGTTGA AGAGGTATGA AACCTATTCC TTTATGTCTG TACTTTTTAG AGTAACGACT CAACRAATAA TTCCGAACGGG TTTTCTTCTT TCTCAGCTTA

101 GAACGTGTG CGCAGGTAGA AGCTTTGGAG AATTATCGTCA CTGCAATGCT TCGCAATATG GCGRAAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG
   CTTGACACAC GCGTCCATCT TCGAAACCTC TAATAGCAGT GACGTTACGA AGCGTTATAC CCGGTTTTAC TGGTTGTGCG CAACATACTA GTCCATCTCC

201 GGGCGCTGTA CGAGGTAAAG CCCGATGCCA GCATTCCTGA CGACCATACG GAGCTGCTGC GCGATTACGT AAAGAAGTTA TTGAAGCATC CTCGTCAATA
   CCCGCGACAT GCTCCATTTT CCGTACCGT CGTRAGACT GCTGCTATGC CTCGACGACG CGCTAATGCA TTTCTTCAAT AACTTCGTAG GAGCAGTCAT

301 AAAAGTAAAT CTTTTCAACA GCTGTCAATA AGTTGTACAG GCCGAGACTT ATAGTCGCTT TGTTTTTAAT TTTTAAATGTA TTTGTAACATA GTACGCAAGT
   TTTTCAATTA GAAAAGTTGT CGACAGTATT TCAACAGTGC CCGCTCTGAA TATCAGCGAA AAAAAATAA AAAATTACAT AAACATTGAT CATCGGTTCA

401 TCACGTAAAA AGGGTATCTA GAATTATGAT GATTACTCTG CGCAAACTTC CTCTGGCGGT TGCCGTGCGA GCGGCGGTAA TGTCTGCTCA GCCCATGGCC
   AGTGCATTTT TCCCATAGAT CTTAATACTA CTRATGAGAC GCGTTTGAAG GAGACCGCCA ACGGCAGCST CGCCCGGANT ACAGACGAGT CCGGTACCCGG
   MetMe tileThrLeu ArgLysLeup roleuAlaVal AlaAlaGlyValM etSerAlaGl nAlaMetAla

1  501 GGTCCCGAAA CTCTGTCCAG TGTGAACTG GTTACCGTC TGCAAGTCTG ATGTGGTGAT CGAGGCTTCC TGTTCACAA ACCGACTGGG GCTGGATCCT
   CCAGGGCTTT GAGACACGCC ACGACTTGAC CAACTGCGAG ACGTCAAGCA TACACCACATA GCTCCGAAGG ACAAGTTGTT TGGCTGACCC CGACCTAGGA

26 GlyProGluT hrLeuCysGl yAlaGluLeu ValAspAlaL euGlnPheVa lCysGlyAsp ArgGlyPheL euPheAsnly sProThrGly AlaGlySerSer

601 CCTCTCGTCG TGTCCCCAG ACTGGTATTG TTGACGAATG CTGCTTTCGT TCTTTCGACC TCGTCTGCTT GGAATGTAT TCGCTCCCC TGAACCCCGC
   GGAGAGCAGC ACGAGGGTC TGACCATAAC AACTGCTTAC GACGAAAGCA AGAACGCTGG ACGCAGCAGA CCTTTACATA ACCGAGGGG ACTTTGGCGG

60 SerArgAr gAlaProGln ThrGlyIleV alAspGluCy sCysPheArg SerCysAspL euArgArgLe uGluMetTyr CysAlaProL euLysProAla

701 TAAATCTGCT TAGAAGCTCC TAACGCTCGG TTGCCGCGCG GCGTTTTTTA TTGTTAACTC ATGTTTGACA GCTTATCATC GATAAGCTTT AATGCGGTAG
   ATTTAGACGA ATCTTCGAGG ATTGCGAGC AACGGCGGCC CGCAAAAAAT AACAAATTGAG TACAAACTGT CGAATAGTAG CTATTTCGAAA TTACGCCATC

93 LysSerAla Am*

801 TTTATCACAG TTAATTTGCT AACCCAGTCA GGCACCGTGT ATGAAATCTA ACAATCGCT CACGTCATC CTCGGCACCG TCACCCCTGA TGCTGTAGGC
   AAATAGTGTG AATTTAACGA TTGCGTCAGT CCGTGGCACA TACTTTAGAT TGTTACCCTG TGTACAGTAG GAGCCGTGGC AGTGGGACCT ACGACATCCG

901 ATAGGCTTGG TTAGCCGGT ACTGCCGGC CTCTTGCGGG ATATCGTCCA TTCCGACAGC ATCCGACGTC ACTATGGCGT GCTGTAGCG CTATATGCGT
   TATCCGAACC AATACGGCCA TGACGGCCCG GAGAACGCC TATAGCAGGT AAGGCTTCC TAGSGTCCAG TGATACCCGA CGACGATCCG GATATACGCA

1001 TGATGCAATT TCTATGGCA CCCGTTCTCG GAGCACTGTC CGACCGCTTT GGCCGCCGCC CAGTCTGCT CGCTTCGCTA CTTGGAGCCA CTATCGACTA
   ACTACGTTAA AGATACGCGT GGGCAAGAGC CTCGTGACAG GCTGGCGAAA CCGCGCGCGG GTCAGGACGA GCGAAGCGAT GAACCTCGGT GATAGCTGAT
    
```

FIG. 3A

1101 CGGATCATG GGCACCAC CCGTCCGTG GATCCTCTAC GCGGACGCA TCGTGCCGG CATCACCCGG GCCACAGTG CGGTTGCTGG CGCTATATC
 GCGCTACTAC CGCTGGTGTG GGCAGGAC CTAGAGATG CGCCCTGCGT AGCACCCGCC GTAGTGGCCG CCGTGTCCAC GCCAACGACC GCGGATATAG
 1201 GCCGACATCA CCGATGGGA AGATCGGCT CGCCACTTCG GGCATGAG CGCTTGTTC GCGTGGGTA TGGTGGCAGG CCCCCTGGCC GGGGGACTGT
 CCGCTGTAGT GGTACTCCCT TCTAGCCCGA CCGGTGAAG CCGAGTACTC GCGAACAAAG CCGCACCCAT ACCACCGTCC GGGGCACCG CCCCCTGACA
 1301 TGGGGCCAT CTCCTTGCAT GCACCATCC TTGGGCGGC GGTGCTCAAC GGCCTCAACC TACTACTGG CTGCTTCCTA ATGCAGGAGT CGCATAAAGG
 ACCCGCGTA GAGGAACGTA CGTGGTAAAG AACGCGCCG CCACAGTTG CCGGAGTTG ATGATGACCC GACGAAGGAT TACGTCCCTCA GCCTATTCCC
 1401 AGAGCTCGA CCGATGCCCT TGAGAGCCTT CAACCAGTC AGCTCTTCC GGTGGGCGG GGCATGACT ATCGTCCCG CACTTATGAC TGTCTTCTTT
 TCTCGCAGCT GGCTACGGGA ACTCTCGGAA GTTGGTCTAG TCGAGGAAG CCACCCGCG CCGTACTGA TAGCAGCGGC GTGAATACTG ACAGAAGAAA
 1501 ATCATGCAAC TCGTAGGACA GGTGCCGGA CCGCTCTGG GCAATTTCCG CGAGGACCG TCAATTTCCG TTTCCGCTGA GCAGCAGAT GATCGGCTG TCGCTTGC
 TAGTACGTTG AGCATCCTGT CCACGGCCGT CCGAGACCC AGTAAAGCC GCTCCTGGG AAAGCGACT CCGCTGTCTA CTAGCCGGAC AGCGAACGCC
 1601 TATTCGGAAT CTTGACGCC CTCGCTCAG CCTTCGTCAC TGGTCCCGC ACCAAACGTT TCGGCGAAA GCAGGCCATT ATCGCCGGA TGGCGGCCGA
 ATAAGCCCTA GAACGTCCGG GAGCGAGTTC GGAAGCAGTG ACCAGGCGG TGGTTTGCAA AGCCGCTCTT CGTCCGGTAA TAGCGCCGT ACCGCGCGT
 1701 CCGGCTGGC TACGCTTGC TGGCGTTCC GACGCGAGC TGGATGGCT TCCCATTTAT GATTTCTTC GCITCCCGG GCATCGGGAT GCCCGGTTG
 GCGGACCCG ATGCAGAAC ACCGCAAGC CTGCGCTCC ACCTACCAG GAGGTAATA CTAAGAAGAG CGAAGGCCG CTAGCCCTA CCGGCGAAC
 1801 CAGGCCATG TGTCCAGGA GTFAGATGAC GACCATCAG GACAGTTCA AGGATCGTC CCGGCTCTTA CCAGCTAAC TTCGATCACT GGACCCTGA
 GTCCGGTACG ACAGGTCCT CCACTACTG CTGGTAGTCC CTGTGAGT TCCTAGCAG CCGCGGAAAT GGTCCGATTG AAGCTAGTGA CCTGGCAGT
 1901 TCGTACCGC GATTTATGC GCCTCGGGA GCACATGGA CCACATGGA CCGGTTGGA TGGATGTAG GCGCCGCCCT ATACTTGTG TGCCTCCCCG CGTTGCGTGC
 AGCAGTGCCG CTAATAACG CCGAGCCGCT CGTGTACCCT GCCCAACCCT ACCTAACATC CCGCGCGGTA TATGGAACAG ACGGAGGGG GCAACCGCAGC
 2001 CCGTGCATGG AGCCGGGCA CCTCGACCTG AATGGAAGC GCGGACCT CGGTAAACGGA TTCACCCTC CAAGRAATGG AGCCAATCAA TTCTTGGCGA
 GCCACGTACC TCGGCCCGT GAGCTGGAC TTACCTCCG CCGCCGTTGA CCGATTCCTT CCGTAAACG AAGTGGTGA GTTCTTAACC TCGGTAGTT AAGAACGCT
 2101 GAACTGTGAA TCGCAAAAC AACCTTGGC AGAACATATC CATCGCTCC GCCATCTCA GCAGCCGAC CCGGCGCATC TCGGCGAGG TTGGTCTGTG
 CTTGACACTT ACGCGTTGG TTGGAAACG TCTTGTATG GTAGCGCAG CCGTAGAGT CCGTGGCGTG CGCCGCGTAG AGCCCGTCC AACCCAGGAC
 2201 GCCACGGGTG CGCATGATG TGTCTCTGC GTTGAAGCC CCGCTAGCT GCGGGGTTG CCTTACTGTT TAGCAATG AATCACCGAT ACGGAGCGA
 CCGTGCCCCAC GCGTACTAG ACGAGGACAG CAACTCCITG GCCGATCCGA CCGCCCAAC GGAATGACCA ATCGTCTTAC TTAGTGGCTA TCGGCTCGT
 2301 ACGTGAAGC ACTGCTGCTG CAAAACGCTT GCGACCTGAG CAACAACATG AATGGTCTT GGTTCCTGG TTTCCGTAAG TCTGGAACG CCGAAGTCA
 TGCACCTCGC TGACGACGAC GTTTTGCAGA CCGTGGACTC GTTGTGTATC TTACCAGAAG CCAAAGGCAC AAAGCATTTC AGACCTTTG GCCTTCACTC
 2401 CGCCCTGCAC CATTATGTT CCGATCTGCA TCGCAGGATG CTGCTGGCTA CCCTGTGGAA CACTACATC TGTATTAAACG AAGCGTGGC ATTGACCTG
 CCGGACGCTG GTAAATACAG GCCTAGACGT AGCTCCTAC GACGACCTT GGGACACCTT GTGGATGAG ACATATTTG TTCGCGACC TAACCTGGAC

FIG. 3B

2501 AGTGATTTT CTCTGGTCCC GCCGATCCA TACCGCCAGT TGTTTACCCT CACAACGTTT CAGTACCCG GCATGTTTCA CATCAGTAAC CCGTATCGTG
TCACTAAAA GAGACCAGG ATGGCGGTCA ACAATGGGA GTGTGCAAG GCATTTGGCC CGTACAAGTA GTAGTCATTG GGCATAGCAC

2601 AGCATCCTCT CTGTTTCAT CCGTATCAAT ACCCCATGA ACAGAAATTC CCCCTTACAC GGAGGCATCA AGTGACCAAA CAGGAAAAA CCGCCCTTAA
TCGTAGGAGA GAGCAAAGTA GCCATAGTAA TGGGGTACT TGCTTTAAG GGGGAATGT CCTCCGTAGT TCACTGGTTT GTCCTTTTT GCGGGGAATT

2701 CATGCCCGC TTTATCAGAA GCCAGACAT AACGGTTCTG GAGAAACTCA ACGAGCTGGA CGCGGATGAA CAGGCAGACA TCTGTGAATC GCITTCACGAC
GTACCCGGCG AATAGTCTT CCGTCTGTAA TTGCGAAGAC CTCITTAGT TGCTCGACCT GCGCCTACTT GTCCGTCTGT AGACACTTAG CGAAGTGCTG

2801 CACGGTGAT AGCTTTACCG CAGTGCCTC CAGCGTTTCG TGAAAACCTC TGACACATGC AGCTCCCGGA GACGGTCACA GCITTGTCTGT
GTGGACTAC TCGAAATGGC GTCGACGGAG CCGCAAAGC CACTACTGCC ACTTTTGGAG ACTGTGTAGC TCGAGGGCCT CTGCCAGTGT CGAACAGACA

2901 AAGCGGATG CCGGAGCAGA CAAGCCGTC AGGGCGGTC AGCGGTGTT GCGGGTGTG GGGCGGCAG CATGACCCAG TCACGTAGCG ATAGCGGAGT
TTCGCCTAGC GCCCTCGTCT GTTCGGGCAG TCCCGGCAG TCCCCACAA CCGCCACAG CCCCAGGTC GTACTGGTC AGTGCATCGC TATCGCCCTCA

3001 GTATCTGGC TTAACATGC GGCATCAGG CAGATTGTAC TGAGAGTGCA CCATATGCGG TGTGAAATAC CGCACAGATG CGTAAGGAGA AAATACCGCA
CATATGACCG AATTGATACG CCGTAGTCTC GTCTAACATG ACTCTCACGT GGATACGCC AACTTTATG GCGTGTCTAC GCATTCCTCT TTTATGGCGT

3101 TCAGCGCTC TTCGGTTC TCGCTCACTG ACTCGTGGC CTCCGTCTGT CCGTGGCGG GAGCGTATC AGCTCACTCA AAGCGGTAA TACGGTTATC
AGTCCCGGAG AAGCGAAGG AGCGAGTGAC TGAGCGACGC GAGCAGCAA GCGACGCGC CTCGCCATAG TCGAGTGAGT TTCGCCATT ATGCCAATAG

3201 CACAGATCA GGGATTAAC GAGGAAGAA CATGTGAGCA AAAGCCAGC AAAGGCCAG GAACCGTAAA AAGSCCGGT TGCTGGCGT TTTCCATAGG
GTGTCTTAGT CCCCTATTGC GTCCCTTCTT GTACACTCGT TTTCCGGTGC TTTTCCGGTC CTTGGCATT TTTCCGGCGCA ACGACCCGAA AAAGGTATCC

3301 CTCGCCCCC CTGACGAGCA TCACAAAAAT CGACGCTCAA GTCAGAGGTG GCGAAACCCG ACAGGACTAT AAAGATACCA GCGGTTTCCC CCTGGAAGCT
GAGGCGGGGG GACTGCTCCT AGTGTTTTTA GCTGCGAGTT CAGICTCCAC CGCTTTGGGC TGTCTGATA TTTCTATGGT CCGCAAAGGG GGACCTTCGA

3401 CCCTCGTGG CTCTCCTGT CCGACCTGC ATACCTTCC GCCTTTCTCC CTTCCGGGAA GCGTGGCGCT TCTCATAGCT CACGCTGTAG
GGAGCCACC GAGAGGACAA GGCTGGACC GCGAATGGCC TATGGACAGG CCGAAGAGG GAAGCCCTTC GCACCCGAA AGAGTATCGA GTGCGACATC

3501 GTATCTCAG TCGGTGTAG TCGTTCGCTC CAAGCTGGC TGTGTGCAGC AACCCCCCGT TCAGCCCGAC CCGTGCCTT TATCCGGTAA CTATCGTCTT
CATAGATCA AGCCACATCC AGCAAGCGAG GTTCGACCCG ACACACGTG TTGGGGGCA AGTCGGGCTG CCGACCCGGA ATAGGCCATT GATAGCAGAA

3601 GAGTCCACC CCGTAAGACA CGACTTATCG CCACTGGCAG CAGCCACTGG TAACAGGATT AGCAGAGCGA GGTATGTAGG CCGTGTACA GAGTTCCTGA
CTCAGGTTGG GCCATTCTGT G.TGAATAGC GGTGACCGTC GTCGGIAGC ATTGTCTTAA TCGTCTCGCT CCATACATCC GCCACGATG CTCAGAAGACT

3701 AGTGGTGGC TAACTACGGC TACACTAGAA GGRACGATTT TGGTATCTGC GCTCTGCTGA AGCCAGTTAC CTTCCGAAAA AGAGTTGGTA GCTCTTGATC
TCACCACCCG ATTGATGCCG ATGTGATCTT CCTGTCATAA ACCATAGAGC CGAGACGACT TCGGTCAATG GAAGCCTTTT TCTCAACCAT CGAGAACTAG

FIG. 3C

3801 CGGCCAACA ACCACGCTG GTAGCGGTGG TTTTTTTGTT TCGAAGCAGC AGATTACGGC CAGAAAAA GGATCTCAAG AAGATCCCTT GATCTTTTCT
 GCCGTTTGT TGGTGGCGAC CATCGCCACC AAAAACAACAA ACGTTCGTGC ICTAATGCGC GTCITTTTTT TTCTAGGAAA CTAGAAAAA

3901 ACGGGGTCTG ACGCTCAGTG GAACGAAAAC TCACGTTAAG GGATTTTGGT CATGAGATTA TCAAAAAGGA TCTTCACTTA GATCCTTTTA AATTAATAAT
 TGCCCCAGAC TCGGAGTCAC CTTGCTTTTG AGTGCAATTC CCTAAAAACCA GTACTCTAAT AGTTTTTCTT AGAAGTGGAT CTAGGAAAAT TTAATTTTTA

4001 GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT TACCATGCT TAATCAGTGA GGCACCTATC TCACGGATCT GTCTATTTCCG
 CTTCAAAT TAGTTAGATT TCATATATAC TCATTTGAAC CAGACTGTCA ATGGTTACGA ATTAGTCACT CCGTGGATAG AGTCGCTAGA CAGATAAAGC

4101 TTCATCCATA GTTGCTGAC TCCCCGTGCT AGCATAAATC AGCAATAAAC CAGCCAGCCG GAAGGGCCGA GCGCAGAAGT GGTCTTGCAA CTTTATCCCG TCTATTAATT
 AAGTAGGIAT CAACGGACTG AGGGCAGCA CATCTATTGA TGCTATGCC TCCCGAATGG TAGACCGGGG TCACGACGTT ACTATGGCGC TCTGGGTGCG

4201 TCACCGGCTC CAGATTTATC AGCAATAAAC CAGCCAGCCG GAAGGGCCGA GCGCAGAAGT GGTCTTGCAA CTTTATCCCG TCTATTAATT
 AGTGGCCGAG GTCATAATAG TCGTTATTTG GTCGGTCCGC CTTCCCGGCT CCGCTTCTCA CCAGGACGTT GAAATAGGCG GAGGTAGGTC AGATAATTAA

4301 GTTGCCGGGA AGCTAGAGTA AGTAGTTCCG CAGTTAATAG TTTGCGCAAC GTTGTGCCA TTGCTGCAGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT
 CAACGGCCCT TCGATCTCAT TCATCAAGCG GTCAATTATC AAACGCGTTG CAACAACGGT AACGACGTTCC GTFACACCAC AGTGGGAGCA GCAAACCATA

4401 GGCTTCATTC AGCTCCGGTT CCCAACGATC AAGCCGAGTT ACATGATCCC CCATGTTGT CAAAAAGCG GTTAGCTCCT TCGGTCTCCT GATCGTTGTC
 CCGAAGTAAG TCGAGGCCAA GGGTGTCTAG TTCCGCTCAA TGTACTAGG GGTACAACAC GTTTTTCCG CAATCCAGGA AGCCAGGAGG CTAGCAACAG

4501 AGAAGTAAGT TGGCCGCAGT GTTATCACTC ATGGTTATGG CAGCACTGCA TAAATCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTTCT GTGACTGGTG
 TCTTCATTCA ACCGGCTCA CAATAGTGAG TACCAATACC GTCGTGACGT ATTAAGAGAA TGCACAGTACG GTAGGCATTC TACGAAAAGA CACTGACCCAC

4601 AGTACTCAAC CAAGTCATTC TGAGAAATAG GTATGCGGCG ACCGAGTTGC TCTTGCCCGG CGTCAACACG GGATAATACC GCGCCACATA GCAGAACTTT
 TCATGAGTTG GTTCAGTAAG ACTCTTATCA CATA CGCCCGC TGGCTCAACG AGAACGGCC GCAGTTGTGC CCTATTATGG CCGGTTGTAT CGTCTTGAAA

4701 AAAAGTCTC ATCATTGGAA AACGTTCTTC GGGCGAAAA CTCTCAAGGA TCTTACCCTT AGAATGGCGA CAACCTTAGG TCAAGCTACA TTGGGTGAGC ACGTGGGTTG
 TTTTCACGAG TAGTACCTT TTGCAAGAAG CCCCCTTTT GAGAGTTCTT GAGAGTTCTT AGAATGGCGA CAACCTTAGG TCAAGCTACA TTGGGTGAGC ACGTGGGTTG

4801 TGATCTCAG CATCTTTTAC TTTCAACCAG GTTTCTGGT GAGCAAAAAC AGRAAGGCAA AATGCCGCAA AAGGGGAAAT AAGGGCGACA CGGAAATGTT
 ACTAGRAGTC GTAGAAAATG AAAGTGGTCC CAAAGACCCA CTCGTTTTTG TCCCTCCGTT TTACGGCGTT TTTTCCCTTA TTCCCGCTGT GCCTTTACAA

4901 GAATACTCAT ACTCTTCTT TTTCAAVATT APTGAAGCAT TTATCAGGGT TATTGTCTCA TGAGCGGATA CATAATTTGA TGTATTTAGA AAAATAAACA
 CTTATGAGTA TGAGAAGGAA AAAGTTATAA TAACTTCGTA AATFATCCCA AATACAGAGT ACTCGCCTAT GTATAAACIT ACATAAATCT TTTTATTTGT

5001 AATAGGGGTT CCGCGCACAT TTCCCCGAAA AGTGCCACCT GACGTCTAAG AAACCTTAT TATCATGACA TTAACCTATA AAAATAGGCG TATCACGAGG
 TTATCCCCAA GGCGCGTGTG AAGGGCTTT TCACGGTGGG CTGCAGATTC TTTTGGTAATA ATAGTACTGT AATTTGGATAT TTTTATCCCG ATAGTCTCC

5101 CCCTTTCTGC TTCAA GGGARAGCAG AAGTT

FIG. 3D




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8A




FIG. 8B

—□— Control —●— IGF-Mutant




FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B




FIG. 16




FIG. 17A




FIG. 17B




FIG. 18




FIG. 19A




FIG. 19B




FIG. 20A




FIG. 20B




FIG. 21A




FIG. 21B




FIG. 22A




FIG. 22B

- | | | | |
|---|----------------------------|---|-----------------------------|
| ● | IGF-1
(150 µg, tid) | □ | IGF Mutant
(150 µg, tid) |
| ○ | IGF Mutant
(50 µg, tid) | ▲ | Excipient
Control |




FIG. 23A




FIG. 23B

- IGF-1 (150 µg, tid)
- ▨ IGF-Mutant (150 µg, tid)
- ▩ IGF-Mutant (50 µg, tid)
- Control

plasmid t4.g8
length: 5140 (circular)

```

1  GAATTCAACT TCTCCATACT TTGGATAAGG AAATACAGAC ATGAAAATC TCAITGCTGA GTGTATATT AACCTTGCCC AAAAAGAAGA AGAGTCGAAT
   CTTAAGTTGA AGAGTATGA AACCTATTCC TTTATGCTCG TACTTTTTAG AGTAACGACT CAACAATAAA TTCCGAACGGG TTTTCTTCT TCTCAGCTTA
101 GAACCTGTGTG CGCAGGTAGA ACCTTTGGAG ATTATCGTCA CTGCAATGCT TCGCAATATG GCGAAAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG
   CTTGACACAC GCGTCCATCT TCGAAACCCTC TAAATAGCAGT GACGTTACGA AGCGTTATAC CCGCTTTTAC TGGTTGTCCG CAACTAACA GTCCATCTCC
201 GGGCGGTGTA CGAGGTAAG CCCGATGCCA GCATTCCTGA CGACGATACG GAGCTGCTGC GCGATTACGT AAAGAAGTTA TTGAAGCATC CTCGTGAGTA
   CCCCGGACAT GCTCCGATTC GGGCTACGGT CGTAAGGACT GCTGCTAIGC CTCGACGACG CGCTAATGCA TTTTCTCAAT AACCTCGTAG GAGCAGTCAT
301 AAAAGTTAAT CTTTCAACA GCTGTCATAA AGTTGTACG GCCGAGACTT ATAGTCGCTT TGTTTTTATT TTTTAAATGTA TTTGTAAC TAACGCAAGT
   TTTTCAATA GAAAAGTTGT CGACAGTATT TCAACAGTGC CGGCTCTGAA TATCAGCGAA ACAAAAATAA AAAATTACAT AAACATTGAT CATGCGTTCA
401 TCACGTTAAA AGGGTATCTA GAGGTTGAGG TGATTTTATG AAAAAAATA TCGCATTTCT TCTTGCACTT ATGTTCTGTTT TTTTCTATTG TACAAATGCC
   AGTGCATTTT TCCCATAGAT CTCCAACCTCC ACTAAAATAC TTTTTCTTAT AGCGTAAGA AGAACGTAGA TACAAGCAAA AAAGATAACG ATGTTTACGG
501 TATGCATCTG GTACCGCCAT GGCCTGATCCG AACCGTTTCC GCGGTAAAGA TCTGGCAGGT TCACCAGGTG GAGGATCCGG AGGAGGCGCC GAGGTTGACG
   ATACGTAGAC CATGGCGGTA CCGACTAGGC TTGGCAAAGG CGCCATTTCT AGACCGTCCA AGTGGTCCAC CTCTTAGGCC TCCTCCCGG CTCCCACTGC
1   serg lyThrAlaPhe tAlaAspPro AsnArgPhea rgGlyLysAs pLeuAlaGly SerProGlyG lyGlySerG1 yGlyGlyAla GluGlyAspAsp
601 ATCCCGCAAA AGCGGCCCTTT AACCTCCTGC AAGCTCAGC GACCGAATAT ATCGGTTATG CGTGGGCGAT GGTGTGTTGTC ATTGTCCGGC CAACTATCGG
   TAGGGCGTTT TCGCCGGAAA TTGAGGACG TTCCGAGTCCG CTGGCTTATA TAGCCAATAC GCACCCGCTA CCAACAACAG TAACACCGCG GTTGATAGCC
33  ProAlaLy salaAlaPhe AsnSerLeuG lnAlaSerAl aThrGluTyI IleGlyTyra laTrpAlaMe tValValVal IleValGlyA laThrIleGly
701 TATCAAGCTG TTTAAGAAAT TCACCTCGAA AGCAAGCTGA TAAACCGATA CAATTAAGG CTCCTTTTGG AGCCTTTTTT TTTGGAGATT TTCAAACGTTGA
   ATAGTTCGAC AAATTCCTTA AGTGGAGCTT TCGTTCCGACT ATTTGGCTAT GTTAATTTCC GAGGAAAACC TCGGAAAAAA AAACCTCTAA AAGTTGCACT
66  IleLysLeu PheLysLysP heThrSerLy salaSer
801 AAAAATATT ATTTCGCAATT CCTTTAGTTG TTCTCACTCC GCTGAAACTG TTGAAAGTTG TTTAGCAAAA CCCCATACAG AAAATTCATT
   TTTTAAATAA TAAGCGTTAA GAAATCAAC AAGGAAAGAT AAGAGTGAGG CGACTTTGAC AACTTCAAC AAATCGTTTT GGGGTATGTC TTTTAAATAA
901 TACTAACGTC TGGAAAAGC ACAAAACTTT AGATCGTTAC GCTAATATG AGGTTGTCT GTGGAATGCT ACAGCGTTG TAGTTGTAC TGGTGACGAA
   ATGATTCGAC ACCTTCTGC TGTATTGAAA TCTAGCAATG CGATTGATAC TCCCAACAGA CACTTACGA TGTCCGCAAC ATCAACAATG ACCACTGTT
1001 ACTCACTGTC TAGCTAGAGT GCGGTCCGG CTGGTTCCGG TGATTTTGTAT TATGAAAAGA TGCCAAACGC TAATAAGGGG GCTATGACCG AAAATGCCGA
   TGAGTCACAG ATCGATCTCA CCGCCACCGA GACCAAGGCC ACTAAAAC TAACATTTCT ACCGTTTGGC ATTATTCCCG CGATACTGGC TTTTACGGCT

```

FIG. 24A

1101 TGAAAACGGG CTACAGTCTG ACGCTAAAG CAAACTTGAT TCTGTGCGTA CTGATTACGG TGCTGCTATC GATGTTTCA TTGGTGACGF TTCCGGCCTT
ACTTTTGGCG GATGTCAGAC TGCGATTTC GTTGAACATA AGACAGCGAT GACTAATGCC ACGACGATAG CTACCAAAGT AACCACTGCA AAGCCCGGAA

1201 GCTAATGGTA ATGGTGCTAC TGGTGATTTT GCTGGCTCTA ATTCCCRAAT GGCTCAAAGTC GGTGACGGTG ATAAATCACC TTTAATGAAT AATTTCCGTC
CGATTACCAT TACCACGATG ACCACTAAA CGACCGAGAT TAAGGGTTTA CCGAGTTTCCG CCACTGCCAC TATTAAGTGG AAATTACTTA TTAAGGCAG

1301 AATATTTACC TTCCCTCCCT CAATCGGTTG AATGTCGCC CTTTGTCTTT AGCGTGGTA AACCATATGA ATTTTCTATT GATTGTGACA AAATAAACTT
TTATAAATGG AAGGGAGGGA GTTAGCCAAC TTAGACGGG AAAACAGAAA TCGGACCAT TTGGTATACT TAAAAGATAA CTAACACTGT TTTATTTGAA

1401 ATTCCGGTGT GTCTTTGCGT TCTTTTATA TGTTCGCCACC TTTATGTATG TATTTTCTAC GTTTGCTAAC ATACTGCGTA ATAAGGAGTC TTAATCATGC
TAAGGCACCA CAGAAACGCA AAGAAATAT ACAACGGTGG AAATACATAC ATAAAAGATG CAAACGATTG TATGACCGAT TATTCCTCAG AATTAGTACG

3201 ACTCAAAGC GGTAAATACGG TTAATCCACAG AATCAGGGGA AATCGCAGGA AAGAACATGT GAGCAAAAAG CCACGAAACC GTCAGGAAACC GTAAAAAGGC
TGAGTTTCCG CCAATATGCC AATAGGTGTC TTAGTCCCTT ATTTGGTCCCT TTTCTGTACA CTCGTTTTCC GGTCCGTTTC CCGTCCCTGG CATTTTTCCG

3301 CGCGTTGCTG GCGTTTTC ATAGGCTCCG CCCCCTGAC GAGCATACA AAAATCGAG CTCAAAGTCAG AGGTGGCGAA ACCCGACAGG ACTATAAAGA
GGCAACGAC CGCAAAAAGG TATCCGAGGC GGGGGACTG CTCGTAGTGT TTTTAGCTGC GAGTTCAGTC TCCACCCTTT TGGGCTGTCC TGATATTTCT

3401 TACCAGGCGT TTCCCCTGG AAGCTCCCTC GTGCGTCTC CTGTTCCGAC CCGTCCGCTT ACCGGATACC TGTCGCCCTT TCTCCCTTCG GGAAGCGTGG
ATGGTCCGCA AAGGGGACC TTCAGGGAG CACGCGAGAG GACAAGGCTG GGACGGCGAA TGGCCTATGG ACAGGGCGAA AGAGGAAAGC CTTTCGCCACC

3501 CGCTTTCTCA TAGTCAAGC TGTAGGTATC TCAAGTCCGT GTAGTCCGTT CGTCCAAGC TGGGCTGTGT GCACGAAACC CCCGTTACAG CCGACCGCTG
GCGAAAGAGT ATCGAGTCCG ACATCCATAG AGTCAAGCCA CATCCAGCAA GCGAGGTTCC ACCCGACACA CGTGTCTGGG GGGCAAAGTC GGCTGGCGAC

3601 CGCCTTATCC GGTAACTATC GTCTTGAGTC CAACCCGGTA AGACACGACT TATCGCCACT GGCAGCAGCC ACTGGTAAACA GGATTAGCAG AGCGAGGTAT
GCGGAATAGG CCATTGATAG CAGAACTCAG GTTGGGCCAT TCTGTGCTGA ATAGCGGTGA CCGTCCGTCGG TGACCAATGT CCTAATCGTC TCGCTCCATA

3701 GTAGCGGGT CTACAGAGTT CTTGAAGTGG TGGCCTAATC ACGGCTACAC TAGAAGGACA GTATTGGTA TCTGCGCTCT GCTGAAGCCA GTTACCTTCG
CATCCGCCAC GATGCTCAA GAACCTCACC ACCGGATTGA TGCCGATGTC ATCTTCCCTGT CATAAAACCAT AGACGGGAGA CGACTTCGGT CAATGGAAGC

3801 GAAAAGAGT TGGTAGTCT TGATCCGGCA AACAAACCAC CGCTGGTAGC GGTGGTTTTT TTGTTTGCAA GCAGCAGATT ACGCGCAGAA AAAAAGGATC
CTTTTCTCA ACCATCGAGA ACTAGGCCGT TTGTTTGGTG GCGACCATCG CCACCAAAA AACAAAACGTT CGTCCGCTAA TCGCGTCTT TTTTCCCTAG

3901 TCAAGAAGAT CCTTTGATCT TTTCTACGGG GTCTGACGCT CAGTGGAAAG AAAACTCAGG TTAAGGGATT TTGGTCATGA GATTATCAA AAGGATCTTC
AGTTCTTCTA GGAACACTAGA AAAGATGCC CAGACTGCCA GTCACCTTGC TTTTGAGTGC AATTCCTTAA AACCACTACT CTAATAGTTT TTCTTAGAAG

4001 ACCTAGATCC TTTTAAATTA AAAATGAAGT TTTAAATCAA TCTAAAGTAT ATATGAGTAA ACTTGGTCTG ACAGTTACCA ATGCTTAATC AGTGAGGCAC
TGGATCTAGG AAAATTTAAT TTTTACTTCA AAATTTAGTT AGATTTTATA TATACTCAT TGAACCCAGAC TGTCAATGGT TAGGAATTAG TCACTCCGTC

FIG. 24B

4101 CTATCTCAGC GATCTGTCTA TTTCTGTTTCAT CCATAGTTGC CTGACTCCCC CTGCGTGTAGA TAACTACGAT ACGGGAGGGC TTACCATCTG GCCCCAGTGC
 GATAGAGTCC CTAGACAGAT AAAGCAAGTA GGTATCRAAG GACTGAGGG CAGCACATCT ATTGATGCTA TGCCCTCCCG AATGGTAGAC CGGGGTCACG
 4201 TGCAATGATA CCGGAGACC CACGTCACC GGTCCAGAT TTATCAGCAA TAAACCAGCC AGCCGGAAGG GCCGAGCGCA GAAGTGGTCC TGCAACTTTA
 ACGTFACTAT GCGGCTCTGG GTCCGAGTGG CCGAGGTCIA AATAGTCGTI ATTTGGTCCG TCGSCCTTCC CCGCTCCGCT CTTACCAGG ACGTTGAAAT
 4301 TCCGCCCTCCA TCCAGTCTAT TAAITGTTGC CCGGAAGCTA GAGTAAGTAG TTCGCCAGTT AATAGTTTGC GCRAACGTTGT TGCCATTGCT GCAGGCATCG
 AGGCGGAGGT AGGTCAGATA ATTAACAACG GCCCTTCGAT CTCATTCATC AAGCGGTCAA TTATCAAAACG CGTTGCAACA ACGGTAAACA CGTCCGTAGC
 4401 TGGTGTACG CTCGTCGTTT GGTATGGCTT CATTAGCTC CCGTTCCTCAA CGATCAAGGC GAGTTACATG ATCCCCCATG TTGTGCAAAA AAGCGGTTAG
 ACCACAGTGC GAGCAGCAAA CCATACCGAA GTAAGTCGAG GCCAAGGTT GCTAGTCCG CTCRAATGTAC TAGGGGTAC AACACGTTTT TTCGCCAATC
 4501 CTCCTTCGGT CCTCCGATCG TTGTGAGAG TAAGTTGGCC GCAGTGTAT CACTCATGGT TATGGCAGCA CTGCATAATT CTCTTACTGT CATGCCATCC
 GAGGAAGCCA GGAGGCTAGC AACAGTCTTC ATTCAACCGG CGTCACAATA GTGAGTACCA ATACCCGTCG GACGTATTAA GAGAATGACA GTACGGTAGG
 4601 GTAAGATGCT TTTCTGTGAC TGGTGAGTAC ACCACTCAIG AGTTGGTTCA GTAAGACTCT TATCACATAC GCGCGACCGA GTTGCTCTTG CCGCGGTCA ACACGGGATA
 CATTCACGA AAAGACACTG ACCACTCAIG AGTTGGTTCA GTAAGACTCT TATCACATAC GCGCGTGGCT CAACGAGAAC GCGCGCAGT TGTGCCCTAT
 4701 ATACCGGCC ACATAGCAGA ACTTTAAAAG TGCTCATCAT TGGA AAAACGT TCTTCGGGGC GAAAACTCTC AAGGATCTTA CCGCTGTTGA GATCCAGTTC
 TATGGCGCG TGTATCGTCT TGAAATTTTC ACGAGTAGTA ACCTTTGTCA AGAAGCCCCG CTTTTGAGAG TTCCCTAGAAAT GCGCACAACT CTAGTCAAG
 4801 GATGTAACC ACTCGTGAC CCAACTGATC TTCAGCATCT TTTACTTTCA CCAGCGTTTC TGGGTGAGCA AAAACAGGAA GGC AAAATGC CGCAAAAAG
 CTACATTTGG TGAGCACCGT GGTGACTAG AAGTCGTAGA AAAAGAAAAGT GGTCCGAAAAG ACCCACTCGT TTTTGTCCCT CCGTTTTTACG GCGTTTTTTC
 4901 GGAATRAGG CGACACGGAA ATGTTGAATA CTCATACTCT TCCTTTTTCA ATATTATGA AGCATTATC AGGGTTATTG TCTCATGAGC GGATACATAT
 CCTTATTTCC GCTGTGCCIT TACAACCTTAT GAGTATGAGA AGGAAAAGT TATAATAACT TCCGTAATAAG TCCCAATAAC AGAGTACTCG CCTATGTATA
 5001 TTGAATGTAT TTAGAAAAAT AAACAATAAG GGGTTCCGGC CACATTTCCC CGAAAAGTGC CACCTGACGT CTAAGAAAAC ATTAATATCA TGACATTAAC
 AACTTACATA AATCTTTTTA TTTGTTTTAT CCCAAGGCGC GTGTAAAGGG GCTTTTCCAG GTGGACTGCA GATTCITTTGG TAAATAATAGT ACTGTAATTG
 5101 CTATAAAAAT AGGGGTATCA CGAGGCCCTT TCGTCTTCAA
 GATATTTTTA TCCGCATAGT GCTCCGGGAA AGCAGAAAGT

FIG. 24C




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37A




FIG. 37B




FIG. 37C




FIG. 37D




FIG. 38




FIG. 39




FIG. 40A




FIG. 40B




FIG. 41




FIG. 42





FIG. 43

Effect of IGF-I Treatment on IGFBP-3

(Mean \pm SE)



Treatment Visits (Week)

FIG. 44

INSULIN-LIKE GROWTH FACTOR AGONIST MOLECULES

BACKGROUND OF THE INVENTION

1. Field of Invention

This invention relates to molecules useful as agonists of the insulin-like growth factors (IGFs). More particularly, these molecules prevent the interaction of an IGF with one or more of its IGF binding proteins. Such molecules can be used, for example, in any methods where the IGFs are used, for example, in treating hyperglycemic, obesity-related, neurological, cardiac, renal, immunologic, and anabolic disorders.

2. Description of Background and Related Art

There is a large body of literature on the actions and activities of IGFs (IGF-I, IGF-II, and IGF variants). Human IGF-I is a 7649-dalton polypeptide with a pI of 8.4 (Rinderknecht and Humbel, *Proc. Natl. Acad. Sci. USA*, 73: 2365 (1976); Rinderknecht and Humbel, *J. Biol. Chem.*, 253: 2769 (1978)) belonging to a family of somatomedins with insulin-like and mitogenic biological activities that modulate the action of growth hormone (GH). Van Wyk et al., *Recent Prog. Horm. Res.*, 30: 259 (1974); Binoux, *Ann. Endocrinol.*, 41: 157 (1980); Clemmons and Van Wyk, *Handbook Exp. D Pharmacol.*, 57: 161 (1981); Baxter, *Adv. Clin. Chem.*, 25: 49 (1986); U.S. Pat. No. 4,988,675; WO 91/03253; WO 93/23071.

Like GH, IGF-I is a potent anabolic protein. See Tanner et al., *Acta Endocrinol.*, 84: 681-696 (1977); Uthne et al., *J. Clin. Endocrinol. Metab.*, 39: 548-554 (1974). See also Ross et al., *Intensive Care Med.*, 19 Suppl. 2: S54-57 (1993), which is a review of the role of insulin, GH, and IGF-I as anabolic agents in the critically ill. IGF-I has hypoglycemic effects similar to those of insulin, but also promotes positive nitrogen balance. Underwood et al., *Hormone Res.*, 24: 166 (1986); Guler et al., *N. Engl. J. Med.*, 317: 137 (1987). Due to this range of activities, IGF-I is being tested in humans for such widely disparate uses as wound healing, treatment of diabetes, reversal of whole body catabolic states, treatment of heart conditions such as congestive heart failure, and treatment of neurological disorders. Guler et al., *Proc. Natl. Acad. Sci. USA*, 85: 4889-4893 (1988); Duerr et al., *J. Clin. Invest.*, 95: 619-627 (1995); and *Science*, 264: 772-774 (1994).

U.S. Pat. Nos. 5,273,961; 5,466,670; 5,126,324; 5,187,151; 5,202,119; 5,374,620; 5,106,832; 4,988,675; 5,106,832; 5,068,224; 5,093,317; 5,569,648; and 4,876,242; WO 92/11865; WO 96/01124; WO 91/03253; WO 93/25219; WO 93/08826; and WO 94/16722 disclose various methods of treating mammals, especially human patients, using IGF-I. In addition, clinical uses of IGF-I are described, for example, in Bondy, *Ann Intern. Med.*, 120: 593-601 (1994).

As one specific use, IGF-I has been found to exert a variety of actions in the kidney. Hammerman and Miller, *Am. J. Physiol.*, 265: F1-F14 (1993). It has been recognized for decades that the increase in kidney size observed in patients with acromegaly is accompanied by a significant enhancement of glomerular filtration rate. O'Shea and Layish, *J. Am. Soc. Nephrol.*, 3: 157-161 (1992). U.S. Pat. No. 5,273,961 discloses a method for prophylactic treatment of mammals at risk for acute renal failure. In humans IGF-I has been shown to preserve renal function post-operatively. Franklin et al., *Am. J. Physiol.*, 272: F257-F259 (1997). Infusion of the peptide in humans with normal renal function increases glomerular filtration rate and renal plasma flow. Guler et al., *Acta Endocrinol.*, 121: 101-106 (1989); Guler

et al., *Proc. Natl. Acad. Sci. USA*, 86: 2868-2872 (1989); Hirschberg et al., *Kidney Int.*, 43: 387-397 (1993); U.S. Pat. No. 5,106,832. Further, humans with moderately reduced renal function respond to short-term (four days) IGF-I administration by increasing their rates of glomerular filtration and renal plasma flow. Hence, IGF-I is a potential therapeutic agent in the setting of chronic renal failure. O'Shea et al., *Am. J. Physiol.*, 264: F917-F922 (1993).

Despite the fact that IGF-I can enhance renal function for those experiencing end-stage chronic renal failure, the enhancements of the glomerular filtration rate and renal plasma flow induced by IGF-I short-term do not persist during long-term administration and incidence of side-effects is high. Miller et al., *Kidney International*, 46: 201-207 (1994).

For complete reviews of the effect of IGF-I on the kidney, see, e.g., Hammerman and Miller, *Am. J. Physiol.*, 265: F1-F14 (1993) and Hammerman and Miller, *J. Am. Soc. Nephrol.*, 5: 1-11 (1994).

As to anabolic indications for IGF-I, in HIV-infected patients treated consecutively with IGF-I, the IGF-I promoted anabolism, but tachyphylaxis developed rapidly in the patients. Lieberman et al., *U.S. Endocrine Meeting*, June 1993 (Abst. 1664); Lieberman et al., *J. Clin. Endo. Metab.*, 78: 404-410 (1994). In patients with severe head injuries, a condition associated with profound hypercatabolism and nitrogen loss, infusion of IGF-I produced only a transient positive nitrogen balance. In the first week the patients experienced a positive nitrogen balance, but during the second week, a negative nitrogen balance developed. Chen et al., *U.S. Endocrine Meeting*, June 1993 (Abst. 1596).

IGF-I has hypoglycemic effects in humans similar to those of insulin when administered by intravenous bolus injection. Underwood et al., *Hormone Research*, 24: 166 (1986). IGF-I is known to exert glucose-lowering effects in both normal (Guler et al., *N. Engl. J. Med.*, supra) and diabetic individuals (Schoenle et al., *Diabetologia*, 34: 675-679 (1991); Zenobi et al., *J. Clin. Invest.*, 90: 2234-2241 (1992); Sherwin et al., *Hormone Research*, 41 (Suppl. 2): 97-101 (1994); Takano et al., *Endocrinol. Japan*, 37: 309-317 (1990); Guler et al., *Acta Paediatr. Scand. (Suppl.)*, 367: 52-54 (1990)), with a time course described as resembling regular insulin. See also Kerr et al., "Effect of Insulin-like Growth Factor 1 on the responses to and recognition of hypoglycemia," American Diabetes Association (ADA), 52nd Annual Meeting, San Antonio, Tex., Jun. 20-23, 1992, which reported an increased hypoglycemia awareness following recombinant human IGF-I (rhIGF-I) administration. In addition, single administration of rhIGF-I reduces overnight GH levels and insulin requirements in adolescents with IDDM. Cheetham et al., *Clin. Endocrinol.*, 40: 515-555 (1994); Cheetham et al., *Diabetologia*, 36: 678-681 (1993).

The administration of rhIGF-I to Type II diabetics, as reported by Schalch et al., *J. Clin. Metab.*, 77: 1563-1568 (1993), demonstrated a fall in both serum insulin as well as a paralleled decrease in C peptide levels. This indicated a reduction in pancreatic insulin secretion after five days of IGF-I treatment. This effect has been independently confirmed by Froesch et al., *Horm. Res.*, 42: 66-71 (1994). In vivo studies in normal rats also illustrate that IGF-I infusion inhibits pancreatic insulin release. Fursinn et al., *Endocrinology*, 135: 2144-2149 (1994). In addition, in pancreas perfusion preparations, IGF-I also suppressed insulin secretion. Leahy et al., *Endocrinology*, 126: 1593-1598 (1990). Despite these clear in vivo inhibitory effects of IGF-I

on insulin secretion in humans and animals, in vitro studies have not yielded such uniform results.

RhIGF-I has the ability to improve insulin sensitivity. For example, rhIGF-I (70 $\mu\text{g}/\text{kg}$ bid) improved insulin sensitivity in non-diabetic, insulin-resistant patients with myotonic dystrophy. Vlachopapadopoulou et al., *J. Clin. Endo. Metab.*, 12: 3715–3723 (1995). Saad et al., *Diabetologia*, 37: Abstract 40 (1994) reported dose-dependent improvements in insulin sensitivity in adults with obesity and impaired glucose tolerance following 15 days of rhIGF-I treatment (25 μg and 100 $\mu\text{g}/\text{kg}$ bid). RhIGF-I also improved insulin sensitivity and glycemic control in some patients with severe type A insulin resistance (Schoenle et al., *Diabetologia*, 34: 675–679 (1991); Morrow et al., *Diabetes*, 42 (Suppl.): 269 (1993) (abstract); Kuzuya et al., *Diabetes*, 42: 696–705 (1993)) and in other patients with non-insulin dependent diabetes mellitus. Schalch et al., “Short-term metabolic effects of recombinant human insulin-like growth factor I (rhIGF-I) in type II diabetes mellitus”, in: Spencer EM, ed., *Modern Concepts of Insulin-like Growth Factors* (New York: Elsevier: 1991) pp. 705–715; Zenobi et al., *J. Clin. Invest.*, 90: 2234–2241 (1993).

A general scheme for the etiology of some clinical phenotypes that give rise to insulin resistance and the possible effects of administration of IGF-I on selected representative subjects is given in several references See, e.g., Elahi et al., “Hemodynamic and metabolic responses to human insulin-like growth factor-i (IGF-I) in men,” in: *Modern Concepts of Insulin-Like Growth Factors*, (Spencer, EM, ed.), Elsevier, N.Y., pp. 219–224 (1991); Quinn et al., *New Engl. J. Med.*, 323: 1425–1426 (1990); Schalch et al., “Short-term metabolic effects of recombinant human insulin-like growth factor 1 (rhIGF-I) in type II diabetes mellitus,” in: *Modern Concepts of Insulin-Like Growth Factors*, (Spencer, EM, ed.), Elsevier, N.Y., pp. 705–714 (1991); Schoenle et al., *Diabetologia*, 34: 675–679 (1991); Usala et al., *N. Eng. J. Med.*, 327: 853–857 (1992); Lieberman et al., *J. Clin. Endo. Metab.*, 75: 30–36 (1992); Zenobi et al., *J. Clin. Invest.*, 90: 2234–2241 (1992); Zenobi et al., *J. Clin. Invest.*, 89: 1908–1913 (1992); Kerr et al., *J. Clin. Invest.*, 91: 141–147 (1993). When IGF-I was used to treat Type II diabetic patients in the clinic at a dose of 120–160 $\mu\text{g}/\text{kg}$ twice daily, the side effects outweighed the benefit of the treatment. Jabri et al., *Diabetes*, 43: 369–374 (1994). See also Wilton, *Acta Paediatr.*, 383: 137–141 (1992) regarding side effects observed upon treatment of patients with IGF-I.

The IGF binding proteins (IGFBPs) are a family of at least six proteins (Jones and Clemmons, *Endocr. Rev.*, 16: 3–34 (1995); Bach and Rechler, *Diabetes Reviews*, 3: 38–61 (1995)), with other related proteins also possibly binding the IGFs. The IGFBPs bind IGF-I and IGF-II with varying affinities and specificities. Jones and Clemmons, supra; Bach and Rechler, supra. For example, IGFBP-3 binds IGF-I and IGF-II with a similar affinity, whereas IGFBP-2 and IGFBP-6 bind IGF-II with a much higher affinity than they bind IGF-I. Bach and Rechler, supra; Oh et al., *Endocrinology*, 132, 1337–1344 (1993).

Unlike most other growth factors, the IGFs are present in high concentrations in the circulation, but only a small fraction of the IGFs is not protein bound. For example, it is generally known that in humans or rodents, less than 1% of the IGFs in blood is in a “free” or unbound form. Juul et al., *Clin. Endocrinol.*, 44: 515–523 (1996); Hizuka et al., *Growth Regulation*, 1: 51–55 (1991); Hasegawa et al., *J. Clin. Endocrinol. Metab.*, 80: 3284–3286 (1995). The overwhelming majority of the IGFs in blood circulate as part of a non-covalently associated ternary complex composed of

IGF-I or IGF-II, IGFBP-3, and a large protein termed the acid-labile subunit (ALS). This complex is composed of equimolar amounts of each of the three components. The ternary complex of an IGF, IGFBP-3, and ALS has a molecular weight of approximately 150,000 daltons, and it has been suggested that the function of this complex in the circulation may be to serve as a reservoir and buffer for IGF-I and IGF-II, preventing rapid changes in free IGF-I or IGF-II.

IGF-I naturally occurs in human body fluids, for example, blood and human cerebral spinal fluid. Although IGF-I is produced in many tissues, most circulating IGF-I is believed to be synthesized in the liver. The IGFBPs are believed to modulate the biological activity of IGF-I (Jones and Clemmons, supra), with IGFBP-1 (Lee et al., *Proc. Soc. Exp. Biol. & Med.*, 204: 4–29 (1993)) being implicated as the primary binding protein involved in glucose metabolism. Baxter, “Physiological roles of IGF binding proteins”, in: Spencer (Ed.), *Modern Concepts of Insulin-like Growth Factors* (Elsevier, N.Y., 1991), pp. 371–380. IGFBP-1 production by the liver is regulated by nutritional status, with insulin directly suppressing its production. Suikkari et al., *J. Clin. Endocrinol. Metab.*, 66: 266–272 (1988).

The function of IGFBP-1 in vivo is poorly understood. The administration of purified human IGFBP-1 to rats has been shown to cause an acute, but small, increase in blood glucose. Lewitt et al., *Endocrinology*, 129: 2254–2256 (1991). The regulation of IGFBP-1 is somewhat better understood. It has been proposed (Lewitt and Baxter, *Mol. Cell Endocrinology*, 79: 147–152 (1991)) that when blood glucose rises and insulin is secreted, IGFBP-1 is suppressed, allowing a slow increase in “free” IGF-I levels that might assist insulin action on glucose transport. Such a scenario places the function of IGFBP-1 as a direct regulator of blood glucose.

The IGF system is also composed of membrane-bound receptors for IGF-I, IGF-II, and insulin. The Type 1 IGF receptor is closely related to the insulin receptor in structure and shares some of its signaling pathways. Jones and Clemmons, supra. The IGF-II receptor is a clearance receptor that appears not to transmit an intracellular signal. Jones and Clemmons, supra. Since IGF-I and IGF-II bind to the Type 1 IGF-I receptor with a much higher affinity than to the insulin receptor, it is most likely that most of the effects of IGF-I and IGF-II are mediated by the Type 1 IGF receptor. Ballard et al., “Does IGF-I ever act through the insulin receptor?”, in Baxter et al. (Eds.), *The Insulin-Like Growth Factors and Their Regulatory Proteins*, (Amsterdam: Elsevier, 1994), pp. 131–138.

There has been much work identifying the domains on IGF-I and IGF-II that bind to the IGFBPs. Bayne et al., *J. Biol. Chem.*, 265: 15648–15652 (1990); U.S. Pat. Nos. 5,077,276; 5,164,370; 5,470,828. For example, it has been discovered that the N-terminal region of IGF-I and IGF-II is critical for binding to the IGFBPs. U.S. Pat. Nos. 5,077,276; 5,164,370; 5,470,828. Thus, the natural IGF-I variant, designated des(1-3)IGF-I, binds poorly to IGFBPs.

A similar amount of research has been devoted to identifying the domains on IGF-I and IGF-II that bind to the Type 1 IGF receptor. Bayne et al., supra; Oh et al., supra. It was found that the tyrosine residues in IGF-I at positions 24, 31, and 60 are crucial to the binding of IGF-I to the Type 1 IGF receptor. Bayne et al., supra. Mutant IGF-I molecules where one or more of these tyrosine residues are substituted showed progressively reduced binding to Type 1 IGF receptors. Bayne et al., supra, also investigated whether such

mutants of IGF-I could bind to the Type 1 IGF receptor and to the IGFBPs. They found that quite different residues on IGF-I and IGF-II are used to bind to the IGFBPs from those used to bind to the Type 1 IGF receptor. It is therefore possible to produce IGF variants that show reduced binding to the IGFBPs, but, because they bind well to the Type 1 IGF receptor, show maintained activity in in vitro activity assays.

Also reported was an IGF variant that binds to IGFBPs but not to IGF receptors and therefore shows reduced activity in in vitro activity assays. Bar et al., *Endocrinology*, 127: 3243-3245 (1990). In this variant, designated (1-27, gly⁴,38-70)-hIGF-I, residues 28-37 of the C region of human IGF-I are replaced by a four-residue glycine bridge. Bar et al. studied the transport of the mutant IGF-I when it was perfused as a complex with IGFBP through the heart in terms of the localization of IGFBPs bound to the mutant IGF or to IGF itself. There were no data supplied by Bar et al. on the localization of the IGF mutant given alone, only data on the localization of the complex of the IGF mutant and IGFBP. Further, Bar et al. provided no data on any biological or efficacy response to the administration of the IGF mutant.

Other truncated IGF-I variants are disclosed. For example, in the patent literature, WO 96/33216 describes a truncated variant having residues 1-69 of authentic IGF-I. EP 742,228 discloses two-chain IGF-I superagonists which are derivatives of the naturally occurring single-chain IGF-I having an abbreviated C domain. The IGF-I analogs are of the formula: BCⁿ,A wherein B is the B domain of IGF-I or a functional analog thereof, C is the C domain of IGF-I or a functional analog thereof, n is the number of amino acids in the C domain and is from about 6 to about 12, and A is the A domain of IGF-I or a functional analog thereof.

Additionally, Cascieri et al., *Biochemistry*, 27: 3229-3233 (1988) discloses four mutants of IGF-I, three of which have reduced affinity to the Type 1 IGF receptor. These mutants are: (Phe²³,Phe²⁴,Tyr²⁵)IGF-I (which is equipotent to human IGF-I in its affinity to the Types 1 and 2 IGF and insulin receptors), (Leu²⁴)IGF-I and (Ser²⁴)IGF-I (which have a lower affinity than IGF-I to the human placental Type 1 IGF receptor, the placental insulin receptor, and the Type 1 IGF receptor of rat and mouse cells), and desoctapeptide (Leu²⁴)IGF-I (in which the loss of aromaticity at position 24 is combined with the deletion of the carboxyl-terminal D region of hIGF-I, which has lower affinity than (Leu²⁴)IGF-I for the Type 1 receptor and higher affinity for the insulin receptor). These four mutants have normal affinities for human serum binding proteins.

Bayne et al., *J. Biol. Chem.*, 263: 6233-6239 (1988) discloses four structural analogs of human IGF-I: a B-chain mutant in which the first 16 amino acids of IGF-I were replaced with the first 17 amino acids of the B-chain of insulin, (Gln³,Ala⁴)IGF-I, (Tyr¹⁵,Leu¹⁶)IGF-I, and (Gln³,Ala⁴,Tyr¹⁵,Leu¹⁶)IGF-I. These studies identify some of the domains of IGF-I that are responsible for maintaining high-affinity binding with the serum binding protein and the Type 2 IGF receptor.

Bayne et al., *J. Biol. Chem.*, 264: 11004-11008 (1988) discloses three structural analogs of IGF-I: (1-62)IGF-I, which lacks the carboxyl-terminal 8-amino-acid D region of IGF-I; (1-27,Gly⁴,38-70)IGF-I, in which residues 28-37 of the C region of IGF-I are replaced by a four-residue glycine bridge; and (1-27,Gly⁴,38-62)IGF-I, with a C region glycine replacement and a D region deletion.

Cascieri et al., *J. Biol. Chem.*, 264: 2199-2202 (1989) discloses three IGF-I analogs in which specific residues in the A region of IGF-I are replaced with the corresponding

residues in the A chain of insulin. The analogs are: (Ile⁴¹, Glu⁴⁵, Gln⁴⁶, Thr⁴⁹, Ser⁵⁰,Ile⁵¹, Ser⁵³, Tyr⁵⁵, Gln⁵⁶)IGF-I, an A chain mutant in which residue 41 is changed from threonine to isoleucine and residues 42-56 of the A region are replaced; (Thr⁴⁹,Ser⁵⁰,Ile⁵¹)IGF-I; and (Tyr⁵⁵, Gln⁵⁶) IGF-I.

Clemmons et al., *J. Biol. Chem.*, 265: 12210-12216 (1990) discloses use of IGF-I analogs that have reduced binding affinity for either the Type 1 IGF receptor or binding proteins to study the ligand specificity of IGFBP-1 and the role of IGFBP-1 in modulating the biological activity of IGF-I.

WO 94/04569 discloses a specific binding molecule, other than a natural IGFBP, that is capable of binding to IGF-I and can enhance the biological activity of IGF-I.

U.S. Pat. Nos. 5,593,844 and 5,210,017 disclose a ligand-mediated immunofunctional binding protein assay method that can be used to quantitate the amount of GH binding protein or IGFBP in a liquid sample by the use of antibodies, where complex formation takes place between one of these binding proteins and the hormone ligand that binds to it.

The direction of research into IGF variants has mostly been to make IGF variants that do not bind to the IGFBPs but show maintained binding to the IGF receptor. The idea behind the study of such molecules is that the major actions of the IGFBPs are proposed to be an inhibition of the activity of the IGFs. Chief among these variants is the natural molecule, des(1-3)IGF-I, which shows selectively reduced affinity for some of the IGF binding proteins, yet a maintained affinity for the IGF receptor. U.S. Pat. Nos. 5,077,276; 5,164,370; 5,470,828, supra.

There is a need in the art for a molecule that acts as an IGF agonist, and also for a molecule that binds to IGF binding proteins with high affinity and specificity for therapeutic or diagnostic purposes.

SUMMARY OF THE INVENTION

This invention relates to a novel method for providing releasing factors which, as part of their actions, prevent binding of an IGF to an IGFBP such as by binding to an IGFBP to agonize the action of IGF. Accordingly, the present invention provides a compound that prevents the interaction of an IGF with any one of its IGFBPs and does not bind to a human IGF receptor, excluding (1-27, gly⁴, 38-70)-hIGF-I, excluding antibodies against an IGFBP that do not bind to a human IGF receptor, excluding antibodies that bind to an IGF, and excluding peptides having the native sequence of human IGF-I with the tyrosine residues at positions 24, 31, and/or 60 replaced or deleted.

Preferably, the compound herein binds to an IGFBP, preferably a serum IGFBP. Also, preferably, the compound reduces plasma insulin secretion, reduces plasma GH, and/or reduces blood glucose levels in a mammal.

In other preferred embodiments, the compound herein is a peptide, especially a peptide having about 10 to about 25 amino acid residues, and/or having a cysteine residue at position 5, 6, 7, or 8 numbered from its N-terminus or having a cysteine residue at position 5, 6, 7, or 8 numbered from its C-terminus, or both such cysteine residues, or a cysteine residue at position 2 numbered from its N-terminus.

In another embodiment, the invention provides a peptide comprising an amino acid sequence selected from the group consisting of the following peptides:

BP3-23	ELDGWVCIKVGEQNL CYLAEG	(SEQ ID NO: 1),
BP3-24	WFKTVCYEWEDVQC YTLEEG	(SEQ ID NO: 2),
BP3-25	RVGAYISCSETECWVEDLLDG	(SEQ ID NO: 3),
BP3-4D3.11 (BP14)	VAWEVCWDRHDQGYICTTDS	(SEQ ID NO: 4),
BP3-4D3.11DEL	AWEVCWDRHQGYICTTDS	(SEQ ID NO: 5),
BP13	CWDRHDQGYICTTDS	(SEQ ID NO: 6),
BP3-4B3.3	EESECFEGPGYVICGLVG	(SEQ ID NO: 7),
BP3-02-OX	DMGVCADGPWVMYVCEWTE	(SEQ ID NO: 8),
BP3-01-OX	SEEVCPVAEAWYLCNMWG	(SEQ ID NO: 9),
BP15	SEEVCPVAEAWYLCN	(SEQ ID NO: 10),
BP16	VCWPVAEAWYLCNMWG	(SEQ ID NO: 11),
BP17	VCWPVAEAWYLCN	(SEQ ID NO: 12),
BP06	TGVDCQCGPVHCVCMDWA	(SEQ ID NO: 13),
BP08	TVANCDCYMPLCLCYDSD	(SEQ ID NO: 14),
bp1-01	CRAGPLQWLCEKYFG	(SEQ ID NO: 15), and
bp1-02	SEVGCRAQPLQWLCEKYFG	(SEQ ID NO: 16).

Also provided herein is a composition comprising one of the compounds or peptides described above in a pharmaceutically acceptable carrier. Preferably, this composition is sterile.

Uses of these compounds and peptides include all uses that liberate or enhance at least one biological activity of exogenous or endogenous IGFs. They can be used in treating or preventing conditions in which an IGF such as IGF-I is useful, as described below.

Additionally provided herein is a method for increasing serum and tissue levels of biologically active IGF in a mammal comprising administering to the mammal an effective amount of a compound that prevents the interaction of an IGF with any one of its IGFs and does not bind a human IGF receptor. Preferably, this compound also reduces plasma insulin secretion, plasma GH secretion, or blood glucose levels in a mammal, and does not directly stimulate the secretion or release of endogenous GH from any species. In other preferred embodiments, this compound binds to an IGFs, such as IGFs-1 and/or IGFs-3, and/or does not bind to a human Type 1 IGF-I receptor. In addition, the mammal is preferably human and the compound is preferably a peptide, more preferably one having about 10 to about 25 amino acid residues. Also preferred is where administering the compound, preferably in an amount effective to produce body weight gain, causes an increase in anabolism in the mammal. Additionally preferred is that glycemic control is effected in the mammal after the compound is administered.

Isolated nucleic acid encoding the compound herein, if it is a peptide, is also provided, and may be used for in vivo or ex vivo gene therapy.

The compound herein can be administered alone or together with another agent such as GH, a GH releasing peptide (GHRP), a GH releasing factor (GHRF), a GH releasing hormone (GHRH), a GH secretagogue, an IGF, an IGF in combination with an IGFs, an IGFs, GH in combination with a GH binding protein (GHBP), insulin, or a hypoglycemic agent (which includes in the definition below an insulin-sensitizing agent such as thiazolidinedione).

In yet another aspect of the invention, a method is provided for effecting glycemic control in a mammal comprising administering to the mammal an effective amount of a compound that prevents the interaction of an IGF with any one of its IGFs and does not bind a human IGF receptor. Preferably, the compound also reduces plasma insulin secretion and blood glucose levels in a mammal and binds an IGFs. Also preferably, the mammal has a hyperglycemic

disorder such as diabetes. This method can additionally comprise administering to the mammal an effective amount of a hypoglycemic agent or insulin.

Also provided is a method for increasing serum and tissue levels of biologically active IGF in a mammal, or a method for increasing anabolism in a mammal, or a method for controlling glycemia in a mammal comprising administering to the mammal an effective amount of the composition containing the compound herein.

In another embodiment, a method is provided for determining appropriate dosing of a compound that prevents the interaction of an IGF with any one of its IGFs and does not bind to a human IGF receptor comprising:

(a) measuring the level of an IGF in a body fluid;

(b) contacting the fluid with the compound herein using single or multiple doses; and

(c) re-measuring the level of an IGF in the fluid, wherein if the fluid IGF level has fallen by an amount sufficient to produce the desired efficacy for which the compound is to be administered, then the dose of the compound is adjustable or adjusted to produce maximal efficacy.

In yet another embodiment, a method is provided for determining the amount of a particular IGFs or the amount of the compound bound to a particular IGFs in a biological fluid so that dosing of the compound can be adjusted appropriately. This method involves:

(a) contacting the fluid with 1) a first antibody attached to a solid-phase carrier, wherein the first antibody is specific for epitopes on the IGFs such that in the presence of antibody the IGF binding sites remain available on the IGFs for binding to the compound, thereby forming a complex between the first antibody and the IGFs; and 2) the above-identified compound for a period of time sufficient to saturate all available IGF binding sites on the IGFs, thereby forming a saturated complex;

(b) contacting the saturated complex with a detectably labeled second antibody which is specific for epitopes on the compound which are available for binding when the compound is bound to the IGFs; and

(c) quantitatively analyzing the amount of the labeled second antibody bound as a measure of the IGFs in the biological fluid, and therefore as a measure of the amount of the compound bound.

Also contemplated herein is a kit comprising a container containing a pharmaceutical composition containing the compound herein and instructions directing the user to utilize the composition. This kit may optionally further comprise a container containing a GH, a GHRP, a GHRF, a

GHRH, a GH secretagogue, an IGF, an IGF complexed to an IGFBP, an IGFBP, a GH complexed with a GHBP, insulin, or a hypoglycemic agent.

Also included herein is a method for predicting the relative affinity for binding to a ligand of a peptide that competes with a polypeptide for binding to the ligand, which peptide is derived from a phage-displayed library, which method comprises incubating a phagemid clone corresponding to the peptide with the polypeptide in the presence of the ligand, serially diluting the phage, and measuring the degree to which binding of the phagemid clone to the ligand is inhibited by the peptide, wherein a phagemid clone that is inhibited only at low phage concentrations has a higher affinity for the ligand than a phagemid clone that is inhibited at both high and low phage concentrations.

In another embodiment herein, a method for directing endogenous IGF either away from, or towards, a particular site in a mammal comprises administering to the mammal an effective amount of the compound herein that is specific for an IGFBP that is either prevalent at, or absent from, the site.

A further embodiment is a method for detecting endogenous or exogenous IGF bound to an IGF binding protein or the amount of a compound that binds to an IGF binding protein and does not bind to a human IGF receptor bound to an IGF binding protein or detecting the level of unbound IGF in a biological fluid comprising:

(a) contacting the fluid with 1) a means for detecting the compound attached to a solid-phase carrier, wherein the means is specific for the compound such that in the presence of the compound the IGF binding sites remain available on the compound for binding to the IGF binding protein, thereby forming a complex between the means and the IGF binding protein; and 2) the compound for a period of time sufficient to saturate all available IGF binding sites on the IGF binding protein, thereby forming a saturated complex;

(b) contacting the saturated complex with a detectably labeled second means which is specific for the IGF binding protein when the compound is bound to the IGF binding protein; and

(c) quantitatively analyzing the amount of the labeled means bound as a measure of the IGFBP in the biological fluid, and therefore as a measure of the amount of bound compound and IGF binding protein, bound IGF and IGF binding protein, or active IGF present in the fluid.

There has been much debate as to the role of the IGFbps in the action of an IGF. The activity of the IGFs in various situations has been shown to be either inhibited, enhanced, or unaffected by the presence of the IGFbps. Jones and Clemmons, supra; Bach and Rechler, supra. It has been unclear if the presence of IGFbps is obligatory for some actions of the IGFs. For some actions it was thought possible that it was necessary for the IGFs to be bound to the IGFbps, or that it was necessary for the IGFbps to be present if IGF-I were to be fully active. Before the present studies it was therefore unclear as to what would be the net biological effect in vivo of administering molecules that prevent the interaction of an IGF with any one of its IGFbps.

The compounds herein are superior to IGF mutants such as des(1-3)IGF-I, since the latter have short half-lives and effects, whereas the compounds herein have longer half lives and effects, and, if they bind to IGFbps, this binding avoids normal renal filtration which would otherwise eliminate short peptides and other small molecules rapidly. Further, administering the compound herein together with exogenous GH or GH secretagogues would have the advantage of minimizing diabetogenic effects of such GH and secreta-

gogues. Yet another advantage of the compounds herein is that there is a ceiling of the effects of the IGF agonist compound herein. That is, it cannot exert more effects than the maximum capacity of IGFbps to carry IGFs, unlike IGF-I, which can have unwanted side effects if used in large concentrations over its maximum efficacy.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts the nucleotide sequence (SEQ ID NO:17) and translated amino acid sequence (SEQ ID NO:18) of the lamB signal connected at the 5' end of DNA encoding an IGF-I mutant designated herein as Y24L,Y31A, or (Leu²⁴, Ala³¹)hIGF-I, or IGF-M, where the Tyr at position 24 is changed to Leu and the Tyr at position 31 is changed to Ala.

FIG. 2 depicts the construction of the plasmid pIGFMI from the vector fragment of p131TGF, from a 120-bp and a 190-bp fragment of pBKIGF-2B, and from a synthetic piece of DNA.

FIGS. 3A-3D depicts the full nucleotide sequence of pIGFMI (SEQ ID NO:19).

FIG. 4 is a standard curve for receptor phosphorylation when IGF-I (squares) or (Leu²⁴, Ala³¹)hIGF-I (circles) is added to MCF-7 cells in a KIRA phosphorylation assay.

FIG. 5 shows thymidine incorporation into mouse 3T3 cells using IGF-I or (Leu²⁴, Ala³¹)hIGF-I.

FIG. 6 shows the binding affinity of (Leu²⁴, Ala³¹)hIGF-I to IGFBP-1.

FIG. 7 shows the binding affinity of (Leu²⁴, Ala³¹)hIGF-I to IGFBP-3.

FIGS. 8A and 8B depict the responses in plasma glucose (FIG. 8A) and plasma insulin (FIG. 8B) of normal rats treated, in a first study, with (Leu²⁴, Ala³¹)hIGF-I (solid circles) or a control (PBS) (open squares), expressed as a percentage of the values in the pre-treatment blood samples which were averaged and set at 100%.

FIGS. 9A and 9B depict the responses in plasma glucose (FIG. 9A) and plasma insulin (FIG. 9B) of normal rats treated, in a second study, with (Leu²⁴, Ala³¹)hIGF-I (solid triangles), or IGF-I (solid circles), or a control (open squares), expressed as a percentage of the values in the pre-treatment blood samples which were averaged and set at 100%.

FIGS. 10A and 10B show the percentage change in plasma insulin from baseline (set as 100%) (FIG. 10A) and the percentage change in plasma glucose from baseline (set as 100%) (FIG. 10B) of diabetic rats treated with (Leu²⁴, Ala³¹)hIGF-I, IGF-I, or a control.

FIGS. 11A and 11B show the percentage change in plasma insulin from baseline (set as 100%) (FIG. 11A) and the percentage change in plasma glucose from baseline (set as 100%) (FIG. 11B) of diabetic (ZDF) rats treated, in a second study, with (Leu²⁴, Ala³¹)hIGF-I, IGF-I, or a control.

FIG. 12 shows the final body weights of rats treated with (Leu²⁴, Ala³¹)hIGF-I at two doses, with GH, with a combination of GH and (Leu²⁴, Ala³¹)hIGF-I, or with a control over seven days.

FIGS. 13A and 13B show the weight of the spleen in the study described above for FIG. 12, with FIG. 13A showing absolute spleen weight and FIG. 13B showing spleen weight expressed as a percentage of body weight.

FIGS. 14A and 14B show the weight of the thymus in the study described above for FIG. 12, with FIG. 14A showing absolute thymus weight and FIG. 14B showing thymus weight expressed as a percentage of body weight.

FIGS. 15A and 15B show the weight of the heart in the study described above for FIG. 12, with FIG. 15A showing absolute heart weight and FIG. 15B showing heart weight expressed as a percentage of body weight.

FIG. 16 shows the epiphyseal plate widths from the five treatment groups in the study described for FIG. 12.

FIGS. 17A and 17B show the amount of rat IGF-I (FIG. 17A) and the amount of total IGF-I (FIG. 17B) in the blood in the five treatment groups in the study described for FIG. 12.

FIG. 18 depicts the final body weights over seven days of dwarf rats treated with placebo (open squares), (Leu²⁴, Ala³¹)hIGF-I (open triangles), IGF-I (open circles), a combination of (Leu²⁴, Ala³¹)hIGF-I and IGF-I (solid circles), GH (half open/half solid squares), and a combination of (Leu²⁴, Ala³¹)hIGF-I and GH (solid squares).

FIGS. 19A and 19B show the weight of the spleen in the study described above for FIG. 18, with FIG. 19A showing absolute spleen weight and FIG. 19B showing spleen weight expressed as a percentage of body weight.

FIGS. 20A and 20B show the weight of the kidney in the study described above for FIG. 18, with FIG. 20A showing absolute kidney weight and FIG. 20B showing kidney weight expressed as a percentage of body weight.

FIGS. 21A and 21B show the amount of rat IGF-I (FIG. 21A) and the amount of total IGF-I (FIG. 21B) in the blood in the six treatment groups in the study described for FIG. 18.

FIGS. 22A and 22B depict body weight gain (FIG. 22A) and percent increase in blood glucose levels (FIG. 22B) over a longer time period of diabetic rats treated with excipient control (solid triangles), with (Leu²⁴, Ala³¹)hIGF-I at 150 μ g, tid (three times daily) (open squares), with (Leu²⁴, Ala³¹)hIGF-I at 50 μ g, tid (open circles), and with IGF-I at 150 μ g, tid (solid circles).

FIGS. 23A and 23B show the increase in blood glucose levels (FIG. 23A) and the changes in insulin levels (FIG. 23B) in the study described for FIG. 22. The control is indicated by black bars, the mutant at 150 μ g by dark stippled bars, the mutant at 50 μ g by light stippled bars, and the IGF-I by open bars.

FIGS. 24A–C depicts the partial DNA sequence (SEQ ID NO: 20) of plasmid pt4.g8 used as a template to construct a phage library. Also shown is the amino acid sequence (SEQ ID NO:21) of an antibody-recognizable (gD-tag) peptide fused to g8p of bacteriophage M13.

FIG. 25 shows gene-8 naive phage library enrichments with a selection using four library pools each and the targets IGF-I, IGFBP-1, and IGFBP-3.

FIG. 26 shows an IGF-I blocking assay using g8-phage peptides from IGFBP-3 selections, where the phage titration is with 100 nM IGF-I. In the Figure, the open circles are peptide 4A3.1, the open triangles are peptide 4B3.4, the open squares are peptide 4C3.2, the solid circles are peptide 4D3.3, the solid triangles are peptide 4D3.4, and the solid squares are peptide 4D3.5.

FIG. 27 shows an IGF-I blocking assay using g8-phage peptides from IGFBP-3 selections, where the phage titration is without IGF-I. The designations for the peptides are the same as those described above for FIG. 26.

FIG. 28 shows an IGF-I blocking assay using a g8-phage peptide from an IGFBP-1 selection (peptide bp1-01), where the phage titration is with 1 μ M IGF-I.

FIG. 29 shows an IGF-I blocking assay using g8-phage peptides from IGFBP-3 selections, where the peptides

(4C3.2, 4D3.8, 4D3.9, 4D3.11, and 4D3.12) are from a NEUTRAVIDIN™/DTT selection. The solid bars are with 100 μ M IGF-I and the open bars are without IGF-I.

FIG. 30 shows an IGF-I blocking assay using g8-phage peptides from IGFBP-3 selections where the peptides (indicated on the x axis) are from direct-coat/HCl selection. The solid bars are with 100 μ M IGF-I and the open bars are without IGF-I.

FIG. 31 depicts a competition assay of IGFBP-3 inhibition by a peptide binding to IGFBP-3 (designated BP3-01) using a BIAcore™ surface-plasmon-resonance device to measure free binding protein. The circles indicate 800 response units (RU) of IGF-I and the squares indicate 400 RU of immobilized IGF-I.

FIG. 32 depicts a competition assay of IGFBP-3 inhibition by a peptide binding to IGFBP-3 (designated BP3-02) using a BIAcore™ surface-plasmon-resonance device to measure free binding protein. The circles indicate 800 RU of IGF-I and the squares indicate 400 RU of immobilized IGF-I.

FIG. 33 shows inhibition of biotinylated IGFBP-1 binding to IGF-I on plates by three peptides that bind to IGFBP-1 or IGFBP-3 but do not bind to the Type 1 IGF receptor (bp1-01: solid circles, bp1-02: open circles, and bp3-01-ox: open triangles).

FIG. 34 shows inhibition of biotinylated IGFBP-3 binding to IGF-I on plates by two peptides that bind to IGFBP-1 but not to the Type 1 IGF receptor (bp1-01 (bp3): solid circles, and bp1-02 (bp3): open circles).

FIG. 35 shows a radiolabeled IGF-I plate assay of the ability of two peptides that bind to IGFBP-3 but not to the Type 1 IGF receptor (bp3-01-ox: circles, and bp3-02-ox: squares) to inhibit IGFBP-3.

FIG. 36 shows a radiolabeled IGF-I plate assay of the ability of the two IGFBP-3 binding peptides described for FIG. 35 to inhibit IGFBP-1 (symbols are the same).

FIG. 37 depicts KIRA assays of IGF-I activity using three peptides (bp1-01: squares, bp1-02: circles, and bp03-ox: triangles).

FIG. 37A depicts the peptides alone, FIG. 37B depicts the peptides plus IGF-I plus IGFBP-1, FIG. 37C depicts the peptides plus IGF-I, and FIG. 37D depicts the peptides plus IGF-I plus IGFBP-3.

FIG. 38 depicts an IGF-II competition assay of IGFBP-3 inhibition by four peptides, designated bp3-01-ox (open squares), BP14 (open circles), BP15 (closed circles), and BP17 (closed squares), using a BIAcore™ surface-plasmon-resonance device to measure free binding protein. Each peptide was tested using 20 nM IGFBP-3 and approximately 1500 RU of immobilized IGF-II.

FIG. 39 depicts a one-dimensional ¹H NMR spectrum of a peptide binding to IGFBP-1 (bp1-01) in H₂O solution at 30° C.

FIGS. 40A and 40B disclose a three-dimensional model of the structure of the peptide bp1-01 in solution. These are stereoviews of a representative structure of bp1-01 from the ensemble of structures calculated using restraints derived from NMR data. The backbone fold is depicted as a ribbon, and all side-chain heavy atoms are shown; each non-glycine residue is labeled. The two views differ by approximately 90°. The relatively flat hydrophobic surface (on the left in FIG. 40A, and towards the viewer in FIG. 40B) is involved in self association, and may also be involved in IGFBP-1 binding.

FIG. 41 is a chromatogram demonstrating the ability of the mutant (Leu²⁴, Ala³¹)hIGF-I to displace ¹²⁵I-IGF-I from

endogenous IGF-BPs present in serum from normal humans. Data are expressed as cpm per fraction (n=1). In the figure, the solid lines with circles are the mutant at time 0 of incubation, the long dashed lines with circles are the mutant at time 0.5 hr of incubation, the shorter dashed lines with diamonds are the mutant at time 3 hr of incubation, and the shortest dotted lines with squares are the mutant at time 16 hr of incubation.

FIG. 42 shows the concentrations of IGF-I in the blood of type II diabetic patients treated with placebo (p) or 10 (1), 20 (2), 40 (4), or 80 (8) $\mu\text{g}/\text{kg}$ of rhIGF-I for 12 weeks by twice daily subcutaneous injection.

FIG. 43 shows the concentrations of IGF-II in the blood of the patients treated as described for FIG. 42.

FIG. 44 shows the concentrations of IGF-BP-3 in the blood of the patients treated as described for FIG. 42.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

A. Definitions

As used herein, "mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic, and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, sheep, pigs, cows, etc. The preferred mammal herein is a human. The term "non-adult" refers to mammals that are from perinatal age (such as low-birth-weight infants) up to the age of puberty, the latter being those that have not yet reached full growth potential.

As used herein, "IGF" refers to native insulin-like growth factor-I and native insulin-like growth factor-II as well as natural variants thereof such as brain IGF, otherwise known as des(1-3)IGF-I.

As used herein, "IGF-I" refers to insulin-like growth factor-I from any species, including bovine, ovine, porcine, equine, and human, preferably human, and, if referring to exogenous administration, from any source, whether natural, synthetic, or recombinant. Human native-sequence, mature IGF-I, more preferably without a N-terminal methionine, is prepared, e.g., by the process described in EP 230,869 published Aug. 5, 1987; EP 128,733 published Dec. 19, 1984; or EP 288,451 published Oct. 26, 1988. More preferably, this native-sequence IGF-I is recombinantly produced and is available from Genentech, Inc., South San Francisco, Calif. for clinical investigations.

As used herein, "IGF-II" refers to insulin-like growth factor-II from any species, including bovine, ovine, porcine, equine, and human, preferably human, and, if referring to exogenous administration, from any source, whether natural, synthetic, or recombinant. It may be prepared by the method described in, e.g., EP 128,733, supra.

An "IGFBP" or an "IGF binding protein" refers to a protein or polypeptide normally associated with or bound or complexed to IGF-I or IGF-II, whether or not it is circulatory (i.e., in serum or tissue). Such binding proteins do not include receptors. This definition includes IGFBP-1, IGFBP-2, IGFBP-3, IGFBP-4, IGFBP-5, IGFBP-6, Mac 25 (IGFBP-7), and prostacyclin-stimulating factor (PSF) or endothelial cell-specific molecule (ESM-1), as well as other proteins with high homology to IGFBPs. Mac 25 is described, for example, in Swisshelm et al., *Proc. Natl. Acad. Sci. USA*, 92: 4472-4476 (1995) and Oh et al., *J. Biol. Chem.*, 271: 30322-30325 (1996). PSF is described in Yamauchi et al., *Biochemical Journal*, 303: 591-598 (1994). ESM-1 is described in Lassalle et al., *J. Biol. Chem.*, 271: 20458-20464 (1996). For other identified IGFBPs, see, e.g., EP 375,438 published Jun. 27, 1990; EP 369,943 published

May 23, 1990; WO 89/09268 published Oct. 5, 1989; Wood et al., *Molecular Endocrinology*, 2: 1176-1185 (1988); Brinkman et al., *The EMBO J.*, 7: 2417-2423 (1988); Lee et al., *Mol. Endocrinol.*, 2: 404-411 (1988); Brewer et al., *BBRC*, 152: 1289-1297 (1988); EP 294,021 published Dec. 7, 1988; Baxter et al., *BBRC*, 147: 408-415 (1987); Leung et al., *Nature*, 330: 537-543 (1987); Martin et al., *J. Biol. Chem.*, 261: 8754-8760 (1986); Baxter et al., *Comp. Biochem. Physiol.*, 91B: 229-235 (1988); WO 89/08667 published Sep. 21, 1989; WO 89/09792 published Oct. 19, 1989; and Binkert et al., *EMBO J.*, 8: 2497-2502 (1989).

The term "body fluid" refers to a biological sample of liquid from a mammal, preferably from a human. Such fluids include aqueous fluids such as serum, plasma, lymph fluid, synovial fluid, follicular fluid, seminal fluid, amniotic fluid, milk, whole blood, urine, cerebrospinal fluid, saliva, sputum, tears, perspiration, mucus, tissue culture medium, tissue extracts, and cellular extracts.

As used herein, "human IGF receptor" refers to any receptor for an IGF found in humans and includes the Type 1 and Type 2 IGF receptors in humans to which both human IGF-I and IGF-II bind, such as the placental Type 1 IGF-I receptor, etc.

A compound that "prevents the interaction of an IGF with any one of its IGFBPs" refers to a molecule that increases serum and tissue levels of biologically active IGF, no matter how this increase occurs. For instance, the compound may partially or completely displace active IGF from a complex in which the IGF is bound to one or more of its IGFBPs. The compound under this definition may bind to an IGFBP, and possibly thereby act to displace an endogenous IGF formerly bound to the IGFBP, or it may bind to an IGF itself at a site remote from that involved in receptor interactions so as to inhibit or prevent the interaction of the IGF with one or more of its IGFBPs, but not inhibit or prevent the interaction of the IGF with any of its receptors. Further, while the compound will occupy the IGFBPs, the effect on the ternary complex will depend on whether the binary complexes can form ternary ones. IGF agonist compounds that can form complexes with ALS will replace IGFs but not affect the concentration of IGFBP-3 or of ternary complexes. IGF agonist compounds that cannot form complexes with ALS will occupy IGFBP-3, and the amount of ALS/IGFBP-3/IGF complex will be reduced. This may differ between full-length IGF mutants which may form ternary complexes and small peptides, which might not. With respect to the structure of one exemplary peptide herein and its interaction with an IGFBP, see Example 9 below.

A compound that "binds to IGF binding protein" refers to a compound that binds an IGFBP to at least some degree, whether with high affinity or not.

A compound that "does not bind to a human IGF receptor" does not bind at all to any such receptor, or binds to such receptor with an affinity more than about 200-fold less than wild-type human IGF-I (hIGF-I) or wild-type human IGF-II (hIGF-II) binds to such receptor. Preferably, the compound binds to such receptor with an affinity of more than about 250-fold less than wild-type hIGF-I or hIGF-II binds to the same receptor or does not bind at all. Such a compound is additionally defined as one that does not phosphorylate the human Type 1 IGF receptor and does not stimulate the mouse IGF-I receptor as measured by thymidine uptake into mouse 3T3 cells using the KIRA and mouse 3T3 assays of Example 1, i.e., the compound acts like (Leu²⁴,Ala³¹)hIGF-I or binds the receptor even less than this mutant in these assays. Further, IGF-II could be construed as an IGF-I agonist, and IGF-I could be construed as an IGF-II agonist.

However, both IGF-I and IGF-II bind to the Type 1 IGF receptor, and thus are both receptor-active molecules and not within the scope of the compounds as defined herein.

A "disorder" is any condition that would benefit from treatment with an IGF, including but not limited to, for example, lung diseases, hyperglycemic disorders as set forth below, renal disorders, such as acute and chronic renal insufficiency, end-stage chronic renal failure, glomerulonephritis, interstitial nephritis, pyelonephritis, glomerulosclerosis, e.g., Kimmelstiel-Wilson in diabetic patients and kidney failure after kidney transplantation, obesity, GH-insufficiency, Turner's syndrome, Laron's syndrome, short stature, undesirable symptoms associated with aging such as obesity and increased fat mass-to-lean ratios, immunological disorders such as immunodeficiencies including decreased CD4 counts and decreased immune tolerance or chemotherapy-induced tissue damage, bone marrow transplantation, diseases or insufficiencies of cardiac structure or function such as heart disfunctions and congestive heart failure, neuronal, neurological, or neuromuscular disorders, e.g., peripheral neuropathy, multiple sclerosis, muscular dystrophy, or myotonic dystrophy, and catabolic states associated with wasting caused by any condition, including, e.g., trauma or wounding or infection such as with a bacterium or human virus such as HIV, wounds, skin disorders, gut structure and function that need restoration, and so forth. The disorder being treated may be a combination of two or more of the above disorders. The preferred disorders targeted for treatment herein are diabetes and obesity, heart disfunctions, kidney disorders, neurological disorders, whole body growth disorders, and immunological disorders.

As used herein, the term "hyperglycemic disorders" refers to all forms of diabetes and disorders resulting from insulin resistance, such as Type I and Type II diabetes, as well as severe insulin resistance, hyperinsulinemia, and hyperlipidemia, e.g., obese subjects, and insulin-resistant diabetes, such as Mendenhall's Syndrome, Werner Syndrome, leprechaunism, lipoatrophic diabetes, and other lipotrophies. The preferred hyperglycemic disorder is diabetes, especially Type I and Type II diabetes. "Diabetes" itself refers to a progressive disease of carbohydrate metabolism involving inadequate production or utilization of insulin and is characterized by hyperglycemia and glycosuria.

As used herein, the term "treating" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disorder as well as those prone to having the disorder or diagnosed with the disorder or those in which the disorder is to be prevented. Consecutive treatment or administration refers to treatment on at least a daily basis without interruption in treatment by one or more days. Intermittent treatment or administration, or treatment or administration in an intermittent fashion, refers to treatment that is not consecutive, but rather cyclic in nature. The treatment regime herein can be either consecutive or intermittent.

As used herein, the term "hypoglycemic agent" refers to compounds that are useful for regulating glucose metabolism, preferably oral agents. More preferred herein for human use are insulin and the sulfonylurea class of oral hypoglycemic agents, which cause the secretion of insulin by the pancreas. Examples include glyburide, glipizide, and gliclazide. In addition, agents that enhance insulin sensitivity or are insulin sensitizing, such as biguanides (including metformin and phenformin) and thiazolidenediones such as REZULIN™ (troglitazone) brand insulin-sensitizing agent, and other compounds that bind to the PPAR γ nuclear receptor, are within this definition, and also are preferred.

As used herein, "insulin" refers to any form of insulin from any species, and whether natively or synthetically or recombinantly derived. Preferably it is NPH insulin.

As used herein, "active" or "biologically active" IGF in the context of changing serum and tissue levels of endogenous IGF refers to IGF that binds to its receptor or otherwise causes a biological activity to occur, such as those biological activities of endogenous or exogenous IGF referred to above.

A "molecule" or "compound" that prevents interaction of an IGF with any one of its IGFbps and does not bind a human IGF receptor includes molecules with high oral bioavailability, exemplified by GH secretagogues, organic chemical molecules modeled after the 3-dimensional model given herein, and peptides as defined below. Such compounds are also referred to herein as "IGF agonists" or "IGF agonist compounds".

"Peptides" include molecules having at least two amino acids and include polypeptides having at least about 50 amino acids. Preferably, the peptides have about 10 to about 25 amino acids, more preferably about 12-25, and most preferably about 15-25 amino acids. The definition includes peptide derivatives, their salts, or optical isomers.

"Growth hormone releasing peptides or factors" ("GHRP" or "GHRF") are described below, as are secretagogues. A "growth hormone releasing hormone" ("GHRH") can be any hormone that releases GH from the cells or tissue. "Growth hormone in combination with a growth hormone binding protein" ("GH" plus "GHBP") means a GH complexed with or otherwise associated with one of its binding proteins. Similarly, "IGF in combination with an IGF binding protein" ("IGF" plus "IGFBP") refers to an IGF complexed with or otherwise associated with one of its IGFbps.

B. Modes for Carrying Out the Invention

The invention herein relates to a compound or molecule that prevents interaction of an IGF with one or more of its IGFbps and does not bind to a human IGF receptor, excluding (1-27, gly⁴, 38-70)-hIGF-I, excluding antibodies against (that bind to) an IGFbp, excluding antibodies against (that bind to) an IGF, and excluding peptides having the native sequence of human IGF-I with the tyrosine residues at positions 24, 31, and/or 60 replaced or deleted. Preferably, the compound binds to an IGFbp and/or to an IGF, especially to IGFbp-1, or to IGFbp-3, or to both IGFbp-1 and IGFbp-2, or to IGF-I, or to IGF-II, or to both IGF-I and IGF-II, or to IGFbp-1 or -3 and IGF-I or -II. Also preferably, the compound is a peptide. More preferably, it is a peptide that binds to an IGFbp, more preferably to a serum IGFbp. Also, preferably, the compound reduces plasma insulin secretion, reduces plasma GH, or reduces blood glucose levels in a mammal. More preferably, the peptide has about 10 to about 25 amino acid residues and/or has a cysteine residue at position 5, 6, 7, or 8 numbered from its N-terminus or a cysteine residue at position 5, 6, 7, or 8 numbered from its C-terminus, or both such cysteine residues, or a cysteine residue at position 2 numbered from its N-terminus. Preferably, the peptide described above has about 12 to 25, more preferably 15 to 25, amino acid residues. More preferably, the peptide has a tryptophan residue at position 1, 2, 3, 4, or 5 numbered from its N-terminus. Still more preferably, the peptide additionally has a valine, serine, or glutamine residue N-terminal to the N-terminal cysteine residue. Even more preferably, the peptide is such that the residue N-terminal to the N-terminal cysteine residue is a valine residue. Still more preferably, the peptide has a glycine-proline or valine-alanine sequence of residues beginning at a position that is three residues

C-terminal to the N-terminal cysteine. This peptide further preferably has a tryptophan residue within three residues C-terminal to the glycine-proline or valine-alanine sequence of residues. This peptide more preferably has a leucine or valine residue within two residues N-terminal to the C-terminal cysteine residue. This peptide even more preferably has a tryptophan residue at position 2 or 3 numbered from its C-terminus.

Also preferred is a peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:1-16. Preferably, this peptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16. More preferably, this peptide comprises an amino acid sequence that is SEQ ID NO:4, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:15, or SEQ ID NO:16. Still more preferably, this peptide comprises an amino acid sequence that is SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:15, or SEQ ID NO:16. Most preferably, this peptide comprises an amino acid sequence that is SEQ ID NO:15 or SEQ ID NO:16.

The compound preferably excludes any IGF-I analogs of the formula: BCⁿ, A

wherein B is the B domain of IGF-I or a functional analog thereof, C is the C domain of IGF-I or a functional analog thereof, n is the number of amino acids in the C domain and is from about 6 to about 12, and A is the A domain of IGF-I or a functional analog thereof. The compounds herein specifically exclude (Leu^{2,4})IGF-I and (Ser^{2,4})IGF-I. The compounds also exclude a B-chain human IGF-I mutant in which the first 16 amino acids of IGF-I are replaced with the first 17 amino acids of the B-chain of insulin; (Gln³,Ala⁴)IGF-I; (Tyr¹⁵,Leu¹⁶)IGF-I; (Gln³,Ala⁴,Tyr¹⁵,Leu¹⁶)IGF-I; (1-62)IGF-I, which lacks the carboxyl-terminal 8-amino-acid D region of IGF-I; (1-27,Gly⁴,38-62)IGF-I, with a C region glycine replacement and a D region deletion; (Ile⁴¹,Glu⁴⁵,Gln⁴⁶,Thr⁴⁹,Ser⁵⁰,Ile⁵¹,Ser⁵³,Tyr⁵⁵,Gln⁵⁶)IGF-I, an A chain mutant in which residue 41 is changed from threonine to isoleucine and residues 42-56 of the A region are replaced; (Thr⁴⁹,Ser⁵⁰,Ile⁵¹)IGF-I; and (Tyr⁵⁵,Gln⁵⁶)IGF-I.

The compounds herein that are not peptides may be made by chemical synthesis or other appropriate methods in the art for making non-peptidyl molecules with high oral bioavailability (such as GH secretagogues) and other orally active organic molecules.

The peptides of this invention can be made by chemical synthesis or by employing recombinant technology. These methods are known in the art. Chemical synthesis, especially solid phase synthesis, is preferred for short (e. g., less than 50 residues) peptides or those containing unnatural or unusual amino acids such as D-Tyr, Ornithine, amino adipic acid, and the like. Recombinant procedures are preferred for longer polypeptides. When recombinant procedures are selected, a synthetic gene may be constructed de novo or a natural gene may be mutated by, for example, cassette mutagenesis. Set forth below are exemplary general recombinant procedures.

From a purified IGF and its amino acid sequence, for example, an IGF agonist that is a peptidyl mutant of an IGF may be produced using recombinant DNA techniques. These techniques contemplate, in simplified form, taking the gene, either natural or synthetic, encoding the peptide; inserting it into an appropriate vector; inserting the vector into an appropriate host cell; culturing the host cell to cause expres-

sion of the gene; and recovering or isolating the peptide produced thereby. Preferably, the recovered peptide is then purified to a suitable degree.

Somewhat more particularly, the DNA sequence encoding a peptidyl IGF agonist is cloned and manipulated so that it may be expressed in a convenient host. DNA encoding parent polypeptides can be obtained from a genomic library, from cDNA derived from mRNA from cells expressing the peptide, or by synthetically constructing the DNA sequence (Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2d ed.), Cold Spring Harbor Laboratory, N.Y., 1989).

The parent DNA is then inserted into an appropriate plasmid or vector which is used to transform a host cell. In general, plasmid vectors containing replication and control sequences which are derived from species compatible with the host cell are used in connection with those hosts. The vector ordinarily carries a replication site, as well as sequences which encode proteins or peptides that are capable of providing phenotypic selection in transformed cells.

For example, *E. coli* may be transformed using pBR322, a plasmid derived from an *E. coli* species. Mandel et al., *J. Mol. Biol.* 53: 154 (1970). Plasmid pBR322 contains genes for ampicillin and tetracycline resistance, and thus provides easy means for selection. Other vectors include different features such as different promoters, which are often important in expression. For example, plasmids pKK223-3, pDR720, and pPL-lambda represent expression vectors with the tac, trp, or PL promoters that are currently available (Pharmacia Biotechnology).

A preferred vector is pB0475. This vector contains origins of replication for phage and *E. coli* that allow it to be shuttled between such hosts, thereby facilitating both mutagenesis and expression. Cunningham et al., *Science*, 243: 1330-1336 (1989); U.S. Pat. No. 5,580,723. Other preferred vectors are pRIT5 and PR1T2T (Pharmacia Biotechnology). These vectors contain appropriate promoters followed by the Z domain of protein A, allowing genes inserted into the vectors to be expressed as fusion proteins.

Other preferred vectors can be constructed using standard techniques by combining the relevant traits of the vectors described above. Relevant traits include the promoter, the ribosome binding site, the decorsin or ornatin gene or gene fusion (the Z domain of protein A and decorsin or ornatin and its linker), the antibiotic resistance markers, and the appropriate origins of replication.

The host cell may be prokaryotic or eukaryotic. Prokaryotes are preferred for cloning and expressing DNA sequences to produce parent IGF-I polypeptide, segment-substituted peptides, residue-substituted peptides, and peptide variants. For example, *E. coli* K12 strain 294 (ATCC No. 31446) may be used as well as *E. coli* B, *E. coli* X1776 (ATCC No. 31537), and *E. coli* c600 and c600hfl, *E. coli* W3110 (F-, gamma-, prototrophic/ATCC No. 27325), bacilli such as *Bacillus subtilis*, and other enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcesans*, and various *Pseudomonas* species. The preferred prokaryote is *E. coli* W3110 (ATCC 27325). When expressed by prokaryotes the peptides typically contain an N-terminal methionine or a formyl methionine and are not glycosylated. In the case of fusion proteins, the N-terminal methionine or formyl methionine resides on the amino terminus of the fusion protein or the signal sequence of the fusion protein. These examples are, of course, intended to be illustrative rather than limiting.

In addition to prokaryotes, eukaryotic organisms, such as yeast cultures, or cells derived from multicellular organisms

may be used. In principle, any such cell culture is workable. However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a reproducible procedure. *Tissue Culture*, Academic Press, Kruse and Patterson, editors (1973). Examples of such useful host cell lines are VERO and HeLa cells, Chinese Hamster Ovary (CHO) cell lines, W138, 293, BHK, COS-7, and MDCK cell lines.

A variation on the above procedures contemplates the use of gene fusions, wherein the gene encoding the desired peptide is associated, in the vector, with a gene encoding another protein or a fragment of another protein. This results in the desired peptide being produced by the host cell as a fusion with another protein or peptide. The "other" protein or peptide is often a protein or peptide which can be secreted by the cell, making it possible to isolate and purify the desired peptide from the culture medium and eliminating the necessity of destroying the host cells which arises when the desired peptide remains inside the cell. Alternatively, the fusion protein can be expressed intracellularly. It is useful to use fusion proteins that are highly expressed.

The use of gene fusions, though not essential, can facilitate the expression of heterologous peptides in *E. coli* as well as the subsequent purification of those gene products. Harris, in *Genetic Engineering*, Williamson, R., Ed. (Academic Press, London, Vol. 4, 1983), p. 127; Ljungquist et al., *Eur. J. Biochem.*, 186: 557-561 (1989) and Ljungquist et al., *Eur. J. Biochem.*, 186: 563-569 (1989). Protein A fusions are often used because the binding of protein A, or more specifically the Z domain of protein A, to IgG provides an "affinity handle" for the purification of the fused protein. It has also been shown that many heterologous proteins are degraded when expressed directly in *E. coli*, but are stable when expressed as fusion proteins. Marston, *Biochem J.*, 240: 1 (1986).

Fusion proteins can be cleaved using chemicals, such as cyanogen bromide, which cleaves at a methionine, or hydroxylamine, which cleaves between an Asn and Gly residue. Using standard recombinant DNA methodology, the nucleotide base pairs encoding these amino acids may be inserted just prior to the 5' end of the gene encoding the desired peptide.

Alternatively, one can employ proteolytic cleavage of fusion protein. Carter, in *Protein Purification: From Molecular Mechanisms to Large-Scale Processes*, Ladisch et al., eds. (American Chemical Society Symposium Series No. 427, 1990), Ch 13, pages 181-193.

Proteases such as Factor Xa, thrombin, and subtilisin or its mutants, and a number of others have been successfully used to cleave fusion proteins. Typically, a peptide linker that is amenable to cleavage by the protease used is inserted between the "other" protein (e.g., the Z domain of protein A) and the desired peptide. Using recombinant DNA methodology, the nucleotide base pairs encoding the linker are inserted between the genes or gene fragments coding for the other proteins. Proteolytic cleavage of the partially purified fusion protein containing the correct linker can then be carried out on either the native fusion protein, or the reduced or denatured fusion protein.

The peptide may or may not be properly folded when expressed as a fusion protein. Also, the specific peptide linker containing the cleavage site may or may not be accessible to the protease. These factors determine whether the fusion protein must be denatured and refolded, and if so, whether these procedures are employed before or after cleavage.

When denaturing and refolding are needed, typically the peptide is treated with a chaotrope, such as guanidine HCl,

and is then treated with a redox buffer, containing, for example, reduced and oxidized dithiothreitol or glutathione at the appropriate ratios, pH, and temperature, such that the peptide is refolded to its native structure.

When peptides are not prepared using recombinant DNA technology, they are preferably prepared using solid-phase synthesis, such as that generally described by Merrifield, *J. Am. Chem. Soc.*, 85: 2149 (1963), although other equivalent chemical syntheses known in the art are employable. Solid-phase synthesis is initiated from the C-terminus of the peptide by coupling a protected α -amino acid to a suitable resin. Such a starting material can be prepared by attaching an α -amino-protected amino acid by an ester linkage to a chloromethylated resin or a hydroxymethyl resin, or by an amide bond to a BHA resin or MBHA resin. The preparation of the hydroxymethyl resin is described by Bodansky et al., *Chem. Ind. (London)*, 38: 1597-1598 (1966). Chloromethylated resins are commercially available from BioRad Laboratories, Richmond, Calif. and from Lab. Systems, Inc. The preparation of such a resin is described by Stewart et al., "Solid Phase Peptide Synthesis" (Freeman & Co., San Francisco 1969), Chapter 1, pp. 1-6. BHA and MBHA resin supports are commercially available and are generally used only when the desired polypeptide being synthesized has an unsubstituted amide at the C-terminus.

The amino acids are coupled to the peptide chain using techniques well known in the art for the formation of peptide bonds. One method involves converting the amino acid to a derivative that will render the carboxyl group more susceptible to reaction with the free N-terminal amino group of the peptide fragment. For example, the amino acid can be converted to a mixed anhydride by reaction of a protected amino acid with ethylchloroformate, phenyl chloroformate, sec-butyl chloroformate, isobutyl chloroformate, pivaloyl chloride or like acid chlorides. Alternatively, the amino acid can be converted to an active ester such as a 2,4,5-trichlorophenyl ester, a pentachlorophenyl ester, a pentafluorophenyl ester, a p-nitrophenyl ester, a N-hydroxysuccinimide ester, or an ester formed from 1-hydroxybenzotriazole.

Another coupling method involves use of a suitable coupling agent such as N,N'-dicyclohexylcarbodiimide or N,N'-diisopropylcarbodiimide. Other appropriate coupling agents, apparent to those skilled in the art, are disclosed in E. Gross & J. Meienhofer, *The Peptides: Analysis, Structure, Biology*, Vol. I: Major Methods of Peptide Bond Formation (Academic Press, New York, 1979).

It should be recognized that the α -amino group of each amino acid employed in the peptide synthesis must be protected during the coupling reaction to prevent side reactions involving their active α -amino function. It should also be recognized that certain amino acids contain reactive side-chain functional groups (e.g., sulfhydryl, amino, carboxyl, and hydroxyl) and that such functional groups must also be protected with suitable protecting groups to prevent a chemical reaction from occurring at that site during both the initial and subsequent coupling steps. Suitable protecting groups, known in the art, are described in Gross and Meienhofer, *The Peptides: Analysis, Structure, Biology*, Vol.3: "Protection of Functional Groups in Peptide Synthesis" (Academic Press, New York, 1981).

In the selection of a particular side-chain protecting group to be used in synthesizing the peptides, the following general rules are followed. An α -amino protecting group (a) must render the α -amino function inert under the conditions employed in the coupling reaction, (b) must be readily removable after the coupling reaction under conditions that

will not remove side-chain protecting groups and will not alter the structure of the peptide fragment, and (c) must eliminate the possibility of racemization upon activation immediately prior to coupling. A side-chain protecting group (a) must render the side chain functional group inert under the conditions employed in the coupling reaction, (b) must be stable under the conditions employed in removing the α -amino protecting group, and (c) must be readily removable upon completion of the desired amino acid peptide under reaction conditions that will not alter the structure of the peptide chain.

It will be apparent to those skilled in the art that the protecting groups known to be useful for peptide synthesis will vary in reactivity with the agents employed for their removal. For example, certain protecting groups such as triphenylmethyl and 2-(p-biphenyl)isopropylloxycarbonyl are very labile and can be cleaved under mild acid conditions. Other protecting groups, such as t-butyloxycarbonyl (BOC), t-amylloxycarbonyl, adamantylloxycarbonyl, and p-methoxybenzyloxycarbonyl are less labile and require moderately strong acids, such as trifluoroacetic, hydrochloric, or boron trifluoride in acetic acid, for their removal. Still other protecting groups, such as benzyloxycarbonyl (CBZ or Z), halobenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, cycloalkyloxycarbonyl, and isopropylloxycarbonyl, are even less labile and require stronger acids, such as hydrogen fluoride, hydrogen bromide, or boron trifluoroacetate in trifluoroacetic acid, for their removal. Among the classes of useful amino acid protecting groups are included:

(1) for an α -amino group, (a) aromatic urethane-type protecting groups, such as fluorenylmethylloxycarbonyl (FMOC) CBZ, and substituted CBZ, such as, e.g., p-chlorobenzyloxycarbonyl, p-6-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, and p-methoxybenzyloxycarbonyl, o-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 2,6-dichlorobenzyloxycarbonyl, and the like; (b) aliphatic urethane-type protecting groups, such as BOC, t-amylloxycarbonyl, isopropylloxycarbonyl, 2-(p-biphenyl)isopropylloxycarbonyl, allyloxycarbonyl, and the like; (c) cycloalkyl urethane-type protecting groups, such as cyclopentylloxycarbonyl, adamantylloxycarbonyl, and cyclohexylloxycarbonyl; and d) allyloxycarbonyl. The preferred α -amino protecting groups are BOC or FMOC.

(2) for the side chain amino group present in Lys, protection may be by any of the groups mentioned above in (1) such as BOC, p-chlorobenzyloxycarbonyl, etc.

(3) for the guanidino group of Arg, protection may be by nitro, tosyl, CBZ, adamantylloxycarbonyl, 2,2,5,7,8-pentamethylchroman-6-sulfonyl or 2,3,6-trimethyl-4-methoxyphenylsulfonyl, or BOC.

(4) for the hydroxyl group of Ser, Thr, or Tyr, protection may be, for example, by C1-C4 alkyl, such as t-butyl; benzyl (BZL); or substituted BZL, such as p-methoxybenzyl, p-nitrobenzyl, p-chlorobenzyloxy, o-chlorobenzyloxy, and 2,6-dichlorobenzyloxy.

(5) for the carboxyl group of Asp or Glu, protection may be, for example, by esterification using groups such as BZL, t-butyl, cyclohexyl, cyclopentyl, and the like.

(6) for the imidazole nitrogen of His, the tosyl moiety is suitably employed.

(7) for the phenolic hydroxyl group of Tyr, a protecting group such as tetrahydropyranyl, tert-butyl, trityl, BZL, chlorobenzyloxy, 4-bromobenzyloxy, or 2,6-dichlorobenzyloxy is suitably employed. The preferred protecting group is 2,6-dichlorobenzyloxy.

(8) for the side chain amino group of Asn or Gln, xanthyl (Xan) is preferably employed.

(9) for Met, the amino acid is preferably left unprotected.

(10) for the thio group of Cys, p-methoxybenzyl is typically employed.

The C-terminal amino acid, e.g., Lys, is protected at the N-amino position by an appropriately selected protecting group, in the case of Lys, BOC. The BOC-Lys-OH can be first coupled to the benzyldiamine or chloromethylated resin according to the procedure set forth in Horiki et al., *Chemistry Letters*, 165-168 (1978) or using isopropylcarbodiimide at about 25° C. for 2 hours with stirring. Following the coupling of the BOC-protected amino acid to the resin support, the α -amino protecting group is removed, as by using trifluoroacetic acid (TFA) in methylene chloride or TFA alone. The deprotection is carried out at a temperature between about 0° C. and room temperature. Other standard cleaving reagents, such as HCl in dioxane, and conditions for removal of specific α -amino protecting groups are described in the literature.

After removal of the α -amino protecting group, the remaining α -amino and side-chain protected amino acids are coupled stepwise within the desired order. As an alternative to adding each amino acid separately in the synthesis, some may be coupled to one another prior to addition to the solid-phase synthesizer. The selection of an appropriate coupling reagent is within the skill of the art. Particularly suitable as a coupling reagent is N,N'-dicyclohexyl carbodiimide or diisopropylcarbodiimide.

Each protected amino acid or amino acid sequence is introduced into the solid-phase reactor in excess, and the coupling is suitably carried out in a medium of dimethylformamide (DMF) or CH₂Cl₂ or mixtures thereof. If incomplete coupling occurs, the coupling procedure is repeated before removal of the N-amino protecting group prior to the coupling of the next amino acid. The success of the coupling reaction at each stage of the synthesis may be monitored. A preferred method of monitoring the synthesis is by the ninhydrin reaction, as described by Kaiser et al., *Anal. Biochem.*, 34: 595 (1970). The coupling reactions can be performed automatically using well-known methods, for example, a BIOSEARCH 9500™ peptide synthesizer.

Upon completion of the desired peptide sequence, the protected peptide must be cleaved from the resin support, and all protecting groups must be removed. The cleavage reaction and removal of the protecting groups is suitably accomplished simultaneously or stepwise. When the resin support is a chloromethylated polystyrene resin, the bond anchoring the peptide to the resin is an ester linkage formed between the free carboxyl group of the C-terminal residue and one of the many chloromethyl groups present on the resin matrix. It will be appreciated that the anchoring bond can be cleaved by reagents that are known to be capable of breaking an ester linkage and of penetrating the resin matrix.

One especially convenient method is by treatment with liquid anhydrous hydrogen fluoride. This reagent not only will cleave the peptide from the resin but also will remove all protecting groups. Hence, use of this reagent will directly afford the fully deprotected peptide. When the chloromethylated resin is used, hydrogen fluoride treatment results in the formation of the free peptide acids. When the benzyldiamine resin is used, hydrogen fluoride treatment results directly in the free peptide amines. Reaction with hydrogen fluoride in the presence of anisole and dimethylsulfide at 0°C for one hour will simultaneously remove the side-chain protecting groups and release the peptide from the resin.

When it is desired to cleave the peptide without removing protecting groups, the protected peptide-resin can undergo

methanolysis to yield the protected peptide in which the C-terminal carboxyl group is methylated. The methyl ester is then hydrolyzed under mild alkaline conditions to give the free C-terminal carboxyl group. The protecting groups on the peptide chain then are removed by treatment with a strong acid, such as liquid hydrogen fluoride. A particularly useful technique for methanolysis is that of Moore et al., *Peptides, Proc. Fifth Amer. Pept. Symp.*, M. Goodman and J. Meienhofer, Eds., (John Wiley, N.Y., 1977), p. 518–521, in which the protected peptide-resin is treated with methanol and potassium cyanide in the presence of crown ether.

Another method for cleaving the protected peptide from the resin when the chloromethylated resin is employed is by ammonolysis or by treatment with hydrazine. If desired, the resulting C-terminal amide or hydrazide can be hydrolyzed to the free C-terminal carboxyl moiety, and the protecting groups can be removed conventionally.

It will also be recognized that the protecting group present on the N-terminal α -amino group may be removed preferentially either before or after the protected peptide is cleaved from the support.

Purification of the polypeptides of the invention is typically achieved using conventional procedures such as preparative HPLC (including reversed-phase HPLC) or other known chromatographic techniques such as gel permeation, ion exchange, partition chromatography, affinity chromatography (including monoclonal antibody columns), or counter-current distribution.

The peptides of this invention may be stabilized by polymerization. This may be accomplished by crosslinking monomer chains with polyfunctional crosslinking agents, either directly or indirectly, through multi-functional polymers. Ordinarily, two substantially identical polypeptides are crosslinked at their C- or N-termini using a bifunctional crosslinking agent. The agent is used to crosslink the terminal amino and/or carboxyl groups. Generally, both terminal carboxyl groups or both terminal amino groups are crosslinked to one another, although by selection of the appropriate crosslinking agent the alpha amino group of one polypeptide is crosslinked to the terminal carboxyl group of the other polypeptide. Preferably, the polypeptides are substituted at their C-termini with cysteine. Under conditions well known in the art a disulfide bond can be formed between the terminal cysteines, thereby crosslinking the polypeptide chains. For example, disulfide bridges are conveniently formed by metal-catalyzed oxidation of the free cysteines or by nucleophilic substitution of a suitably modified cysteine residue. Selection of the crosslinking agent will depend upon the identities of the reactive side chains of the amino acids present in the polypeptides. For example, disulfide crosslinking would not be preferred if cysteine was present in the polypeptide at additional sites other than the C-terminus. Also within the scope hereof are peptides crosslinked with methylene bridges.

Suitable crosslinking sites on the peptides, aside from the N-terminal amino and C-terminal carboxyl groups, include epsilon amino groups found on lysine residues, as well as amino, imino, carboxyl, sulfhydryl, and hydroxyl groups located on the side chains of internal residues of the peptides or residues introduced into flanking sequences. Crosslinking through externally added crosslinking agents is suitably achieved, e.g., using any of a number of reagents familiar to those skilled in the art, for example, via carbodiimide treatment of the polypeptide. Other examples of suitable multi-functional (ordinarily bifunctional) crosslinking agents are found in the literature.

The peptides of this invention also may be conformationally stabilized by cyclization. The peptides ordinarily are

cyclized by covalently bonding the N- and C-terminal domains of one peptide to the corresponding domain of another peptide of this invention so as to form cyclo-oligomers containing two or more iterated peptide sequences, each internal peptide having substantially the same sequence. Further, cyclized peptides (whether cyclo-oligomers or cyclo-monomers) are crosslinked to form 1–3 cyclic structures having from 2 to 6 peptides comprised therein. The peptides preferably are not covalently bonded through α -amino and main-chain carboxyl groups (head to tail), but rather are crosslinked through the side chains of residues located in the N- and C-terminal domains. The linking sites thus generally will be between the side chains of the residues.

Many suitable methods per se are known for preparing mono- or poly-cyclized peptides as contemplated herein. Lys/Asp cyclization has been accomplished using Na-Boc-amino acids on solid-phase support with Fmoc/9-fluorenylmethyl (OFm) side-chain protection for Lys/Asp; the process is completed by piperidine treatment followed by cyclization.

Glu and Lys side chains also have been crosslinked in preparing cyclic or bicyclic peptides: the peptide is synthesized by solid-phase chemistry on a p-methylbenzhydrylamine resin. The peptide is cleaved from the resin and deprotected. The cyclic peptide is formed using diphenylphosphorylazide in diluted methylformamide. For an alternative procedure, see Schiller et al., *Peptide Protein Res.*, 25: 171–177 (1985). See also U.S. Pat. No. 4,547,489.

Disulfide crosslinked or cyclized peptides are generated by conventional methods. The method of Pelton et al. (*J. Med. Chem.*, 29: 2370–2375 (1986)) is suitable, except that a greater proportion of cyclo-oligomers are produced by conducting the reaction in more concentrated solutions than the dilute reaction mixture described by Pelton et al., for the production of cyclo-monomers. The same chemistry is useful for synthesis of dimers or cyclo-oligomers or cyclo-monomers. Also useful are thiomethylene bridges. Lebl and Hruby, *Tetrahedron Letters*, 25: 2067–2068 (1984). See also Cody et al., *J. Med. Chem.*, 28: 583 (1985).

The desired cyclic or polymeric peptides are purified by gel filtration followed by reversed-phase high-pressure liquid chromatography or other conventional procedures. The peptides are sterile filtered and formulated into conventional pharmacologically acceptable vehicles.

The starting materials required for the processes described herein are known in the literature or can be prepared using known methods and known starting materials.

If in the peptides being created carbon atoms bonded to four nonidentical substituents are asymmetric, then the compounds may exist as diastereoisomers, enantiomers, or mixtures thereof. The syntheses described above may employ racemates, enantiomers, or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the asymmetric carbon atoms, when present, may be in one of two configurations (R or S) and both are within the scope of the present invention.

The compounds described in this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Examples of such salts include ammonium, metal salts like sodium, potassium, calcium,

and magnesium; salts with organic bases like dicyclohexylamine, N-methyl-D-glucamine, and the like; and salts with amino acids like arginine or lysine. Salts with inorganic and organic acids may be likewise prepared, for example, using hydrochloric, hydrobromic, sulfuric, phosphoric, trifluoroacetic, methanesulfonic, malic, maleic, fumaric acids, and the like. Non-toxic and physiologically compatible salts are particularly useful, although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. Examples include reaction of the free-acid or free-base form of the peptide with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble; or in a solvent like water after which the solvent is removed by evaporation, distillation, or freeze drying. Alternatively, the free-acid or -base form of the product may be passed over an ion-exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

Certain specific schemes that may be appropriate for chemical synthesis of the peptides herein are shown in WO 96/15148 published May 23, 1996, the disclosure of which is incorporated herein by reference.

The compounds of this invention are shown to prevent the interaction of an IGF with one or more of its binding proteins and thereby agonize IGF action. It is known to those skilled in the art that there are many uses for IGFs. Therefore, administration of the compounds of this invention for purposes of agonizing an IGF action can have the same effects or uses as administration of an exogenous IGF itself. These uses of IGF include the following, which may be additional to or the same as the disorders as defined above: increasing whole body, bone, and muscle growth rate in normal and hypopituitary animals; protection of body weight and nitrogen loss during catabolic states (such as fasting, nitrogen restriction, elevated corticosteroid levels, and/or diabetes); kidney regeneration; treating peripheral and central nervous system (CNS) degenerative disorders and promoting neuroprotection or repair following CNS damage or injury; treating hypoxia; promotion of wound healing; cardiac regeneration; reversal of cancer cachexia; inhibition of angiogenesis; regeneration of the gastrointestinal tract; stimulation of mammary function; counteracting IGF-I-dependent actions of GH such as metabolic stress, age-related decreases in GH activity, and adult GH deficiency; treating maturity-onset diabetes; and/or treating a specific IGF deficiency.

Additional and specific disorders for which the compounds herein are useful include growth disorders such as GH-resistant short stature, GH-insensitivity syndrome, osteoporosis, and catabolic states; disorders where treatment requires regeneration of tissues or cells, for example, peripheral nerves and supporting cells, central nervous system cells including nerves and glia, and other cells such as oligodendrocytes, muscle, skin, and bone; heart disorders, e.g., heart ischemia, cardiac myopathy, and congestive heart disorders; hyperglycemic disorders such as insulin-dependent and non-insulin-dependent diabetes mellitus and extreme insulin resistance; and renal disorders such as renal failure. These also include stimulation of an anabolic response in elderly humans, prevention of catabolic side effects of glucocorticoids, treatment of osteoporosis, stimulation of the immune system, reduction of obesity, acceleration of wound healing, acceleration of bone fracture repair, treatment of growth retardation, treatment of renal failure or

insufficiency resulting in growth retardation, treatment of physiological short stature, including growth-hormone-deficient children, treating short stature associated with chronic illness, treatment of obesity and growth retardation associated with Prader-Willi syndrome and Turner's syndrome, acceleration of the recovery and reduction in the hospitalization of burn patients, treatment of interuterine growth retardation, skeletal dysplasia, hypercortisolism, and Cushing's syndrome, induction of pulsatile growth hormone release, replacement of growth hormone in stressed patients, treatment of osteochondrodysplasias, Noonans syndrome, schizophrenia, depression, peripheral neuropathy, ALS, depression, Alzheimer's disease, diseases of demyelination, multiple sclerosis, and delayed wound healing, stimulation of the immune system, treatment of psychosocial deprivation, treatment of pulmonary dysfunction and ventilator dependency, attenuation of protein catabolic response after a major operation, reduction of cachexia and protein loss due to chronic illness such as cancer or AIDS, treatment of hyperinsulinemia including Type II and Type I diabetes, adjuvant treatment for ovulation induction, stimulation of thymic development and prevention of the age-related decline of thymic function, treatment of immunosuppressed patients, treatment of bone-marrow transplanted patients, improvement in muscle strength, mobility, diseases of muscle function, muscular dystrophy, maintenance of skin thickness, and metabolic homeostasis, enhancement of renal function and homeostasis including acute and chronic renal failure, stimulation of osteoblasts, bone remodeling, and cartilage growth, stimulation of the immune system, and growth promotion in livestock. Various IGF-I uses are found, for example, in WO 94/04569; WO 96/33216; and Bondy, *Ann Intern. Med.*, 120: 593-601 (1994). All these are included in the definition of "disorder."

In one example, the compounds can be administered to commercially important mammals such as swine, cattle, sheep, and the like to accelerate and increase their rate and extent of growth and the efficiency of their conversion of feed into body tissue. The compounds can be administered *in vivo* to adults and children to stimulate IGF action.

The compounds of this invention may be administered to the mammal by any suitable technique, including oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, or subcutaneous injection or infusion, or implant), nasal, pulmonary, vaginal, rectal, sublingual, or topical routes of administration, and can be formulated in dosage forms appropriate for each route of administration. The specific route of administration will depend, e.g., on the medical history of the patient, including any perceived or anticipated side effects using the compound, the type of compound being administered, and the particular disorder to be corrected. Most preferably, the administration is orally or by continuous infusion (using, e.g., slow-release devices or minipumps such as osmotic pumps or skin patches), or by injection (using, e.g., intravenous or subcutaneous means).

The compound to be used in the therapy will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the compound), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amounts" of the compound for purposes herein are thus determined by such considerations and must be amounts that result in bioavailability of the drugs to the mammal and the desired effect.

If a small-molecule antagonist is used as an IGF agonist, it may have cyclical effects and require, for efficacy, an administration regimen appropriate thereto, the variable concentration of IGFBP-1 in blood being an example. Jones and Clemmons, *supra*. For a peptide, a preferred administration is a chronic administration of about two times per day for 4–8 weeks to reproduce the effects of IGF-I. Although injection is preferred, chronic infusion may also be employed using an infusion device for continuous subcutaneous (SC) infusions. A small peptide may be administered orally. An intravenous bag solution may also be employed. The key factor in selecting an appropriate dose for diabetes is the result obtained, as measured by decreases in blood glucose so as to approximate the normal range, or by other criteria for measuring treatment of diabetes as are deemed appropriate by the medical practitioner.

As a general proposition, the total pharmaceutically effective amount of the IGF agonist compound administered parenterally per dose will be in a range that can be measured by a dose-response curve. For example, IGFs bound to IGFBPs or in the blood can be measured in body fluids of the mammal to be treated to determine the dosing. Alternatively, one can administer increasing amounts of the IGF agonist compound to the patient and check the serum levels of the patient for IGF-I and IGF-II. The amount of IGF agonist to be employed can be calculated on a molar basis based on these serum levels of IGF-I and IGF-II. See the examples below on displacement of IGF-I tracer from IGFBPs present in human serum.

Specifically, one method for determining appropriate dosing of the compound entails measuring IGF levels in a biological fluid such as a body or blood fluid. Measuring such levels can be done by any means, including RIA and ELISA. After measuring IGF levels, the fluid is contacted with the compound using single or multiple doses. After this contacting step, the IGF levels are re-measured in the fluid. If the fluid IGF levels have fallen by an amount sufficient to produce the desired efficacy for which the molecule is to be administered, then the dose of the molecule can be adjusted to produce maximal efficacy. This method may be carried out *in vitro* or *in vivo*. Preferably, this method is carried out *in vivo*, i.e., after the fluid is extracted from a mammal and the IGF levels measured, the compound herein is administered to the mammal using single or multiple doses (that is, the contacting step is achieved by administration to a mammal) and then the IGF levels are remeasured from fluid extracted from the mammal.

Another method for determining the amount of a particular IGFBP or the amount of the compound bound to a particular IGFBP in a biological fluid so that dosing of the compound can be adjusted appropriately involves:

(a) contacting the fluid with 1) a first antibody attached to a solid-phase carrier, wherein the first antibody is specific for epitopes on the IGFBP such that in the presence of antibody the IGF binding sites remain available on the IGFBP for binding to the compound, thereby forming a complex between the first antibody and the IGFBP; and 2) the compound for a period of time sufficient to saturate all available IGF binding sites on the IGFBP, thereby forming a saturated complex;

(b) contacting the saturated complex with a detectably labeled second antibody which is specific for epitopes on the compound which are available for binding when the compound is bound to the IGFBP; and

(c) quantitatively analyzing the amount of the labeled second antibody bound as a measure of the IGFBP in the biological fluid, and therefore as a measure of the amount of

the compound bound. This technique can be expanded to include a diagnostic use whereby the compound is administered to a mammal to displace an IGF from a specific IGFBP for which the compound has affinity, such as IGFBP-1 or IGFBP-3, and measuring the amount that is displaced.

The quantitative technique mentioned above using antibodies, called the ligand-mediated immunofunctional method (LIFA), is described for determining the amount of IGFBP by contact with IGF in U.S. Pat. No. 5,593,844, and for determining the amount of GHBP by contact with GH in U.S. Pat. No. 5,210,017. The disclosures of these patents are incorporated herein by reference regarding antibodies and other materials and conditions that can be used in the assay.

Another method for determining dosing is to use antibodies to the IGF agonist or another detection method for the IGF agonist in the LIFA format. This would allow detection of endogenous or exogenous IGFs bound to IGFBP and the amount of IGF agonist bound to the IGFBP.

Another method for determining dosing would be to measure the level of “free” or active IGF in blood. For some uses the level of “free” IGF would be a suitable marker of efficacy and effective doses or dosing.

For example, one method is described for detecting endogenous or exogenous IGF bound to an IGF binding protein or the amount of a compound that binds to an IGF binding protein and does not bind to a human IGF receptor bound to an IGF binding protein or detecting the level of unbound IGF in a biological fluid. This method comprises:

(a) contacting the fluid with 1) a means for detecting the compound that is specific for the compound (such as a first antibody specific for epitopes on the compound) attached to a solid-phase carrier, such that in the presence of the compound the IGF binding sites remain available on the compound for binding to the IGF binding protein, thereby forming a complex between the means and the IGF binding protein; and 2) the compound for a period of time sufficient to saturate all available IGF binding sites on the IGF binding protein, thereby forming a saturated complex;

(b) contacting the saturated complex with a detectably labeled second means which is specific for one or more sites on the IGF binding protein (such as a second antibody specific for epitopes on the IGFBP) which are available for binding when the compound is bound to the IGF binding protein; and

(c) quantitatively analyzing the amount of the labeled means bound as a measure of the IGFBP in the biological fluid, and therefore as a measure of the amount of bound compound and IGF binding protein, bound IGF and IGF binding protein, or active IGF present in the fluid.

Given the above methods for determining dosages, and assuming dosing shares at least some of the characteristics demonstrated in Example 11 for IGF-I, in general, the amount of IGF agonist compound that may be employed can be estimated. An orally active small IGF agonist would have a molecular weight of approximately 500 daltons, compared to 7500 daltons for IGF-I and IGF-II. Assuming the IGF agonist is 16-fold less able to bind to IGFBPs than IGF-I or IGF-II, then equal weights of IGF-I or IGF-II and these molecules could be equally effective, so that doses from about 10 $\mu\text{g}/\text{kg}/\text{day}$ to 200 $\mu\text{g}/\text{kg}/\text{day}$ might be used, based on kg of patient body weight, although, as noted above, this will be subject to a great deal of therapeutic discretion.

A further method is provided to estimate the distribution of IGFs on specific IGFBPs, e.g., on IGFBP-1 or IGFBP-3 using the LIFA format.

The compound is suitably administered by a sustained-release system. Suitable examples of sustained-release com-

positions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include poly(lactides) (U.S. Pat. No. 3,773,919; EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., *Biololymers*, 22, 547-556 (1983), poly(2-hydroxyethyl methacrylate) (Langer et al., *J. Biomed. Mater. Res.*, 15: 167-277 (1981); Langer, *Chem. Tech.*, 12: 98-105 (1982)), ethylene vinyl acetate (Langer et al., *supra*), or poly-D-(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include a liposomally entrapped compound. Liposomes containing the compound are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. U.S.A.*, 82: 3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci. U.S.A.*, 77: 4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appln. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (from or about 200 to 800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the most efficacious therapy.

PEGylated peptides having a longer life can also be employed, based on, e.g., the conjugate technology described in WO 95/32003 published November 30, 1995.

For parenteral administration, in one embodiment, the compound is formulated generally by mixing each at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically, or parenterally, acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the compound uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, a buffered solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; glycine; amino acids such as glutamic acid, aspartic acid, histidine, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose, trehalose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; non-ionic surfactants such as polysorbates, poloxamers, or polyethylene glycol (PEG); and/or neutral salts, e.g., NaCl, KCl, MgCl₂, CaCl₂, etc.

The compound is typically formulated in such vehicles at a pH of from or about 4.5 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of salts of the compound. The final preparation may be a stable liquid or lyophilized solid.

Typical formulations of the peptide or oral secretagogues as pharmaceutical compositions are discussed below. About 0.5 to 500 mg of the compound or mixture of compounds, as the free-acid or -base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules, and the like are a binder such as acacia, corn starch, or gelatin; an excipient such as microcrystalline cellulose; a disintegrating agent like corn starch or alginic acid; a lubricant such as magnesium stearate; a sweetening agent such as sucrose or lactose; a flavoring agent such as peppermint, wintergreen, or cherry. When the dosage form is a capsule, in addition to the above materials, it may also contain a liquid carrier such as a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. A syrup or elixir may contain the active compound, a sweetener such as sucrose, preservatives like propyl paraben, a coloring agent, and a flavoring agent such as cherry. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as water or naturally occurring vegetable oil like sesame, peanut, or cottonseed oil or a synthetic fatty vehicle like ethyl oleate or the like may be desired. Buffers, preservatives, antioxidants, and the like can be incorporated according to accepted pharmaceutical practice.

The compound to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The compound ordinarily will be stored in unit or multi-dose containers, for example, sealed ampules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-mL vials are filled with 5 mL of sterile-filtered 1% (w/v) aqueous solution of compound, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized compound using bacteriostatic Water-for-Injection.

Combination therapy with the IGF agonist compound herein and one or more other appropriate reagents that increase total IGF in the blood or enhance the effect of the IGF agonist is also part of this invention. These reagents generally allow the IGF agonist compound herein to release the generated IGF, and include growth-promoting agents.

Growth-promoting agents for this purpose include, but are not limited to, GH secretagogues that promote the release of endogenous GH in mammals to increase concentrations of the IGF in the blood. Examples include TRH, diethylstilbestrol, theophylline, enkephalins, E series prostaglandins, peptides of the VIP-secretin-glucagon-GRF family, and other GH secretagogues such as GHRP-6, GHRP-1 as described in U.S. Pat. No. 4,411,890, and benzo-fused lactams such as those disclosed in U.S. Pat. No. 5,206,235. See also, e.g., WO 96/15148 published May 23, 1996. Other growth-promoting agents include GHRPs, GHRFs, GH, and their analogs. For example, GHRPs are described in WO 95/17422 and WO 95/17423 both pub-

lished Jun. 29, 1995; Bowers, *J. Pediatr. Endocrinol.*, 6: 21–31 (1993); and Schoen et al., *Annual Reports in Medicinal Chemistry*, 28: 177–186 (1993). GHRFs and their analogs are described, for example, in WO 96/37514 published Nov. 28, 1996.

Additionally, GHRH, any of the IGFFBPs, long-acting GH, GH plus GHBP, insulin, or a hypoglycemic agent can be employed in conjunction with the IGF agonist compound herein for this purpose. In addition, IGF-I or IGF-II or an IGF with an IGFBP such as IGF-I complexed to IGFBP-3 can also be employed with the IGF agonist compound herein. For example, pharmaceutical compositions containing IGF-I and IGFBP in a carrier as described in WO 94/16723 published Aug. 4, 1994 can be used in conjunction with the compound. The entities can be administered sequentially or simultaneously with the IGF agonist compound. In addition, other means of manipulating IGF status, such as regimens of diet or exercise, are also considered to be combination treatments as part of this invention.

If insulin is also administered, it can be any formulation of insulin, but is preferably NPH insulin, and the dose of NPH insulin is from or about 5 to 50 units/injection (i.e., from or about 0.2 to 2 mg) twice a day subcutaneously. For a combination of insulin and the compound, the ratio of NPH insulin to compound in this formulation by weight is generally from or about 10:1 to 1:50, preferably from or about 1:1 to 1:20, more preferably from or about 1:1 to 1:10, still more preferably, from or about 1:1 to 1:5, and most preferably from or about 1:1 to 1:3.

Furthermore, the formulation is suitably administered along with an effective amount of a hypoglycemic agent such as a sulfonylurea. The hypoglycemic agent is administered to the mammal by any suitable technique including parenterally, intranasally, orally, or by any other effective route. Most preferably, the administration is by the oral route. For example, MICRONASE™ tablets (glyburide) marketed by Upjohn in 1.25-, 2.5-, and 5-mg tablet concentrations are suitable for oral administration. The usual maintenance dose for Type II diabetics, placed on this therapy, is generally in the range of from or about 1.25 to 20 mg per day, which may be given as a single dose or divided throughout the day as deemed appropriate. *Physician's Desk Reference*, 2563–2565 (1995). Other examples of glyburide-based tablets available for prescription include GLYNASE™ brand drug (Upjohn) and DIABETA™ brand drug (Hoechst-Roussel). GLUCOTROL™ (Pratt) is the trademark for a glipizide (1-cyclohexyl-3-(p-(2-(5-methylpyrazine carboxamide)ethyl)phenyl)sulfonyl)urea) tablet available in both 5- and 10-mg strengths and is also prescribed to Type II diabetics who require hypoglycemic therapy following dietary control or to patients who have ceased to respond to other sulfonylureas. *Physician's Desk Reference*, 1902–1903 (1995). Other hypoglycemic agents than sulfonylureas, such as the biguanides (e.g., metformin and phenformin) or thiazolidinediones (e.g., troglitazone), or other drugs affecting insulin action may also be employed. If a thiazolidinedione is employed with the compound, it is used at the same level as currently used or at somewhat lower levels, which can be adjusted for effects seen with the compound alone or together with the dione. The typical dose of troglitazone (REZULIN™) employed by itself is about 100–1000 mg per day, more preferably 200–800 mg/day, and this range is applicable herein. See, for example, Ghazzi et al., *Diabetes*, 46: 433–439 (1997). Other thiazolidinediones that are stronger insulin-sensitizing agents than troglitazone would be employed in lower doses.

Another aspect of this invention is a composition comprising an IGF and a thiazolidinedione, or a combination of

an IGF, a thiazolidinedione, and a compound of this invention. Additionally, a method for effecting glycemic control is provided by administering to a mammal in need thereof an effective amount of an IGF and a thiazolidinedione, or an effective amount of an IGF, a thiazolidinedione, and the compound of this invention. The active agents may be administered to the mammal sequentially or together, whether in the same formulation or concurrently. Effective amounts are determined by the practitioner as described above and would generally mean an amount the same as or less than the amount of IGF that is used to treat the condition in question (for example, from about 10 to about 250 $\mu\text{g}/\text{kg}/\text{day}$ of IGF-I for diabetes) and an amount of dione that is known to be useful to treat the condition in question, or if the three are used, the amount of compound using the dosages as determined above.

In addition, the invention contemplates using gene therapy for treating a mammal, using nucleic acid encoding the IGF agonist compound, if it is a peptide. Generally, gene therapy is used to increase (or overexpress) IGF levels in the mammal. Nucleic acids which encode the IGF agonist peptide can be used for this purpose. Once the amino acid sequence is known, one can generate several nucleic acid molecules using the degeneracy of the genetic code, and select which to use for gene therapy.

There are two major approaches to getting the nucleic acid (optionally contained in a vector) into the patient's cells for purposes of gene therapy: *in vivo* and *ex vivo*. For *in vivo* delivery, the nucleic acid is injected directly into the patient, usually at the site where the IGF agonist compound is required. For *ex vivo* treatment, the patient's cells are removed, the nucleic acid is introduced into these isolated cells, and the modified cells are administered to the patient either directly or, for example, encapsulated within porous membranes which are implanted into the patient. See, e.g. U.S. Pat. Nos. 4,892,538 and 5,283,187.

There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells *in vitro*, or *in vivo* in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells *in vitro* include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. A commonly used vector for *ex vivo* delivery of the gene is a retrovirus.

The currently preferred *in vivo* nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and lipid-based systems (useful lipids for lipid-mediated transfer of the gene are DOTMA, DOPE and DC-Chol, for example). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell-surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g., capsid proteins or fragments thereof to facilitate uptake, e.g., capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.*, 262: 4429–4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA*, 87: 3410–3414 (1990). For review of the currently known gene marking and gene therapy protocols, see Ander-

son et al., *Science*, 256: 808–813 (1992). See also WO 93/25673 and the references cited therein.

Kits are also contemplated for this invention. A typical kit would comprise a container, preferably a vial, for the IGF agonist compound formulation comprising IGF agonist compound in a pharmaceutically acceptable buffer and instructions, such as a product insert or label, directing the user to utilize the pharmaceutical formulation. The kit optionally includes a container, preferably a vial, for a GH, a GHRP, a GHRH, a GH secretagogue, an IGF, an IGF complexed to an IGFBP, an IGFBP, a GH complexed with a GHP, insulin, or a hypoglycemic agent.

Also provided is a method for predicting the relative affinity for binding to a ligand of a peptide that competes with a polypeptide for binding to the ligand, which peptide is derived from a phage-displayed library, which method comprises incubating a phagemid clone corresponding to the peptide with the polypeptide in the presence of the ligand, serially diluting the phage, and measuring the degree to which binding of the phagemid clone to the ligand is inhibited by the peptide, wherein a phagemid clone that is inhibited only at low phage concentrations has a higher affinity for the ligand than a phagemid clone that is inhibited at both high and low phage concentrations. Details are provided in Example 7 below. Preferably, the ligand is an IGFBP such as IGFBP-1 or IGFBP-3 and the polypeptide is an IGF.

In another embodiment herein, a method is provided for directing endogenous IGF either away from, or towards, a particular site in a mammal comprising administering to the mammal an effective amount of the compound herein that is specific for an IGFBP that is either prevalent at, or absent from, the site. "Sites" for this purpose include specific tissues or organs such as the heart, or such as the brain via brain-specific IGFbps. Prevalence at the site indicates that the IGFBP in question is located at the site and constitutes a substantial or biologically important portion of the IGFBP at the site. This indication follows from the specificity for IGFBP-1 versus IGFBP-3 of the compounds demonstrated herein.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention. All literature and patent citations mentioned herein are expressly incorporated by reference.

EXAMPLES

To discover the effect of molecules that bind to the IGFbps but not to IGF receptors, a mutant human IGF-I described by Bayne et al., *J. Biol. Chem.*, supra, was produced by recombinant DNA technology in *E. coli*. Specifically, the plasmid pIGFMI was designed for the production of an IGF-I mutant with amino acid changes at residues 24 and 31 (Y24L, Y31A), also designated (Leu²⁴, Ala³¹)hIGF-I or IGF-M in these examples. The plasmid was constructed from a basic backbone of pBR322 (Sutcliffe, *Cold Spring Harb. Symp. Quant. Biol.*, 43: 77–90 (1978)) as described for the construction of phGH1. Chang et al., *Gene*, 55: 189–196 (1987). The transcriptional and translational sequences required for the expression of the gene were provided by the alkaline phosphatase promoter and the trp Shine-Dalgarno sequence. Chang et al., supra. Additionally, the lambda transcriptional termination sequence is located downstream of the gene. Scholtissek and Grosse, *Nucleic Acids Res.*, 15: 3185 (1987). Secretion of the protein from the cytoplasm to the periplasmic space is directed by the lamB signal sequence. Clement and Hofnung, *Cell*, 27:

507–514 (1981). The nucleotide and amino acid sequences for the lamB signal sequence and the IGF-I mutant (Y24L, Y31A) are given in FIG. 1.

The vector fragment for the construction of pIGFMI was isolated by digestion of p131TGF with XbaI and ClaI. The vector p131TGF contains the sequences for the alkaline phosphatase promoter, the ampicillin and tetracycline resistance markers, and the trp Shine-Dalgarno sequence. The lamB signal sequence and amino acids 1–15 of IGF-I were provided by digestion of pBKIGF-IIB (U.S. Pat. No. 5,487,980) with XbaI and PstI. This fragment of approximately 120 bp was isolated and pre-ligated to the following strands of synthetic DNA encoding the amino acid changes Y24L and Y31A (codons underlined):

5'-G TTC GTA TGT GGT GAT CGA GGC TTC
CTG TTC AAC AAA CCG ACT GGG GCT G

3'-ACGT C AAG CAT ACA CCA CTA GCT CCG
AAG GAC AAG TTG TTT GGC TGA CCC CGA
CCTAG

(SEQ ID NOS:22 and 23, respectively)

The remaining IGF-I coding sequence and the lambda transcriptional terminator were provided by isolating the approximately 190-bp BamHI-to-ClaI fragment of pBKIGF-IIB. These fragments were then ligated together to construct pIGFMI as illustrated in FIG. 2. The full nucleotide sequence of pIGFMI is shown in FIGS. 3A–3D. Bayne et al., *J. Biol. Chem.*, supra, described the properties of IGF-I mutants in terms of their binding to IGF binding proteins or to IGF or insulin receptors. This work showed that the tyrosine residues at positions 24, 31, and 60 on human IGF-I are important for binding to the Type 1 IGF-I receptor. The mutant (Leu²⁴, Ala³¹)hIGF-I, where two of these tyrosine residues are mutated, has a half maximal inhibition of ligand binding of 2,000 nM compared to the affinity of wild-type hIGF-I of 8.7 nM. This indicates a relative reduction in affinity for the Type 1 IGF receptor (derived from placental membranes) by about 250-fold of (Leu²⁴, Ala³¹)hIGF-I compared to wild-type hIGF-I.

The functional effect of mutations in IGF-I on activity in vitro was also studied by Bayne et al., *J. Biol. Chem.*, supra.

The assay system used, the incorporation of 3H thymidine into L7 murine fibroblasts, shows a good correlation between the ability of IGF mutants to bind to the Type 1 receptor and their ability to stimulate DNA synthesis. This is reflected in the reduction in activity of (Leu²⁴, Ala³¹)hIGF-I by more than 200-fold in this in vitro activity assay. However, the binding of (Leu²⁴, Ala³¹)hIGF-I to human serum IGFbps is similar to that seen for wild-type IGF-I. Bayne et al., *J. Biol. Chem.*, supra. Therefore, in the following examples, (Leu²⁴, Ala³¹)hIGF-I was chosen to be tested in animals. This molecule was chosen because the mutant contains only two mutations, because Type 1 IGF receptor binding is reduced more than 200-fold, and because binding to the IGFbps is largely maintained.

Example 1

In vitro Activity of (Leu²⁴, Ala³¹)hIGF-I

To test the direct activity of the IGF-I mutant on the IGF receptor, two different assays were employed. A third assay was used to determine if the mutant could displace IGF from IGFbps in a competitive environment.

Assay 1: KIRA for Phosphorylation of the Human Type 1 Receptor

This assay is a direct activity assay for the human Type 1 receptor. When a receptor in the tyrosine kinase family, such

as the Type 1 IGF receptor, is activated, it is phosphorylated on tyrosine residues. In this assay cells containing the Type 1 IGF receptor are activated *in vitro*, then disrupted, and antibodies against the receptor are used to precipitate the IGF receptor. Next, an anti-phosphotyrosine antibody is used to assay the amount of Type 1 IGF receptor that is phosphorylated. If a fixed number of cells is used, then the amount of receptor that is phosphorylated is a direct measure of the activity of a molecule on the Type 1 IGF receptor.

A KIRA for the human Type 1 IGF-I receptor was developed using human MCF-7 cells. This cell line was derived originally from a human breast cancer tumor and is available from the ATCC. The growth of these cells is known to respond to the addition of IGF-I. When IGF-I is added to these cells, a dose-related increase in the phosphorylation of the IGF-I receptor occurs (FIG. 4). The addition of the IGF mutant did not cause any change in receptor phosphorylation (FIG. 4). There was no indication of phosphorylation even at very high concentrations of the mutant IGF.

Specifically, the assay is as follows:

Cells

MCF-7, an adherent cell line isolated from a human breast adenocarcinoma, was purchased from American Type Culture Collection (ATCC-HTB 22; American Type Culture Collection, Rockville, Md.). MCF-7 cells have been shown to express measurable levels of surface IGF-IR by FACS analysis. For culture passage, the cells were cultured in 150 cm² tissue-culture flasks (Corning Inc, Corning, N.Y.), 1.5 × 10⁶/flask for 4–7 days. For the assay, cells were detached from the tissue-culture flasks with PBS/5mM EDTA, quantified, and cultured in flat-bottom microtiter plates (Falcon 3072, Becton Dickinson Labware, Lincoln Park, N.J.), 2 × 10⁵ per well, overnight at 37° C. in 5% CO₂.

Media

Cells were grown in F12/DMEM 50:50 prepared in the media facility of Genentech, Inc. (obtained from Gibco as a custom formulation, Gibco/BRL, Life Technologies, Grand Island, N.Y.). The medium was supplemented with 10% FBS (HyClone, Logan, Utah), 25 mM HEPES (Gibco), and 2 mM L-glutamine (Gibco).

KIRA-ELISA

MCF7 cells (2 × 10⁵) in 100 μl medium were added to each well in a flat-bottom, 96-well culture plate and cultured overnight at 37° C. in 5% CO₂. The following morning the well supernatants were decanted, and the plates were lightly blotted on a paper towel. Stimulation media (F12/DMEM 50:50 with 25 mM HEPES and 2.0% BSA) containing either experimental samples or the recombinant hIGF-I standards were then added to each well. The cells were stimulated at 37° C. for 30 min., the well supernatants were decanted, and the plates were once again lightly blotted on a paper towel. To lyse the cells and solubilize the receptors, 100 μl of lysis buffer were added to each well. Lysis buffer consisted of 150 mM NaCl containing 50 mM HEPES (Gibco), 0.5% Triton-X 100 (Gibco), 0.01% thimerosol, 30 KIU/ml aprotinin (ICN Biochemicals, Aurora, Ohio), 1 mM 4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride (AEBSF; ICN Biochemicals), and 2 mM sodium orthovanadate. The plate was then agitated gently on a plate shaker (Bellico Instruments, Vineland, N.J.) for 60 min. at room temperature.

While the cells were being solubilized, an ELISA microtiter plate (Nunc Maxisorp, Inter Med, Denmark) coated overnight at 4° C. with the polyclonal anti-IGF-IR (antibodies to human Type 1 IGF-I receptor, catalogue 3B7, Santa Cruz Biotech, 5.0 μg/ml in PBS, 100 μl/well) was decanted, blotted on a paper towel and blocked with 150

μl/well of Block Buffer (PBS containing 0.5% BSA (Intergen Company, Purchase, N.Y.) and 0.01% thimerosol) for 60 min. at room temperature with gentle agitation. After 60 minutes, the anti-IGF-IR-coated plate was washed six times with wash buffer (PBS containing 0.05% Tween-20 and 0.01% thimerosol) using an automated plate washer (ScanWasher 300, Skatron Instruments, Inc, Sterling, Va.).

The lysate containing solubilized IGF-IR from the cell-culture microtiter well was transferred (85 μl/well) to anti-IGF-IR-coated and -blocked ELISA wells and incubated for 2 h at room temperature with gentle agitation. The unbound receptor was removed by washing with wash buffer, and 100 μl of biotinylated antibody 4G10 (anti-phosphotyrosine) diluted to 0.1 μg/ml in dilution buffer (PBS containing 0.5% BSA, 0.05% Tween-20, 5 mM EDTA, and 0.01% thimerosol) was added to each well. After incubation for 2 h at room temperature the plate was washed and 100 μl of HRP-conjugated Dextran-streptavidin (Amdex Laboratories) diluted 1:50,000 in dilution buffer was added to each well. The plate was incubated for 30 minutes at room temperature with gentle agitation. The free avidin-conjugate was washed away and 100 μl of freshly prepared substrate solution (tetramethyl benzidine; TMB, 2-component substrate kit; Kirkegard and Perry, Gaithersburg, Md.) was added to each well. The reaction was allowed to proceed for 10 minutes, after which the color development was stopped by the addition at 100 μl/well of 1.0 M H₃PO₄. The absorbance at 450 nm was read with a reference wavelength of 650 nm (Abs₄₅₀), using a vmax plate reader (Molecular Devices, Palo Alto, Calif.) controlled with a MACINTOSH CENTRIS 650™ computer (Apple Computers, Cupertino, Calif.) and SOFTMAX™ software (Molecular Devices).

The standard curve (FIG. 4) was generated by stimulating MCF7 cells with 300, 100, 33.3, 11.1, 3.7, 1.2, 0.4, or 0 ng/ml IGF-I as a reference standard (Genentech, Inc., lot 1189-2 or equivalent). Sample concentrations were obtained by interpolation of their absorbance on the standard curve and are expressed in terms of IGF-I ng/ml activity. The mutant did not phosphorylate the receptor in this assay.

Assay 2: Increase in Cell Number of Mouse 3T3 Cells

The activity of the IGF mutant was also measured in a bioassay for the activity of IGF-I on DNA synthesis and cell replication. In this assay mouse 3T3 cells are cultured, and the IGF mutant or IGF-I is added, followed by tritiated ³H-thymidine. The amount of ³H-thymidine incorporated is a measure of the replication of DNA and thus of cell replication. IGF-I increased thymidine incorporation in a dose-related manner (FIG. 5). The IGF mutant showed no activity in this assay even when added at very high concentrations.

Assay 3: In vitro Binding of (Leu²⁴, Ala³¹)hIGF-I to IGF-BPs

The mutant (Leu²⁴, Ala³¹)hIGF-I was also tested for its ability to displace radio-labeled rhIGF-I off the IGF-BPs in a competitive binding assay. This assay involves coating recombinant human (rh) IGF-BP-1 or rhIGFBP-3 (160 ng/ml) onto 96-well plates overnight, blocking the plates for one hour with 0.5% BSA, adding rhIGF-I (25014 0.08 ng/ml) or (Leu²⁴, Ala³¹)hIGF-I for 1 hour, then adding 20,000 cpm of ¹²⁵I-IGF-I and incubating for 2 hours before washing and counting. FIG. 6 shows that the mutant (Leu²⁴, Ala³¹)hIGF-I binds to recombinant IGF-BP-1 with an affinity only several fold different from that of wild-type hIGF-I. FIG. 7 shows that the mutant (Leu²⁴, Ala³¹)hIGF-I also binds to recombinant IGF-BP-3 with an affinity only several fold different from that of wild-type hIGF-I.

Conclusion

In two *in vitro* assays the mutant (Leu²⁴, Ala³¹)hIGF-I showed little or no direct activity on the Type 1 IGF-

receptor. Firstly, it did not activate the human IGF-I receptor, as measured by the phosphorylation of the receptor in the KIRA assay. Secondly, it did not stimulate the mouse IGF-I receptor directly, as measured by thymidine uptake into 3T3 cells. Therefore, on the basis of the lack of its activation of the IGF receptor, the IGF mutant would also be expected to be inactive in vivo. However, the mutant (Leu²⁴, Ala³¹)hIGF-I did show significant binding to IGFBP-1 and IGFBP-3 in vitro. This in vitro data provided the basis for testing the mutant (Leu²⁴, Ala³¹)hIGF-I in vivo.

Example 2

In vivo Activity of (Leu²⁴, Ala³¹)hIGF-I Introduction

The mutant (Leu²⁴, Ala³¹)hIGF-I tested in Example 1 in vitro was tested in vivo. Even though the mutant (Leu²⁴, Ala³¹)hIGF-I was inactive in vitro, it was hypothesized that molecules of this class (molecules that are inactive directly on receptors, but capable of binding to IGFbps) would show some activity in vivo. In the first studies (Leu²⁴, Ala³¹)hIGF-I was given by IV injection to conscious rats and effects on glycemic control were determined.

Methods

Seven-week-old male Wistar rats (240–250 g, Charles Rivers Laboratories, Hollister, Calif.) were anesthetized with KETAMINE/XYLAZINE™ anesthesia and the right jugular vein was cannulated with a silicone rubber cannula that was developed for chronic blood sampling. Clark et al., *J. Endocrinol.*, 111: 27–35 (1986). Following a two- to three-day recovery period, two basal blood samples were taken at –10 and –5 minutes, the test substances administered to the rats by IV injection, and then blood samples collected after 5, 10, 20, 30, 45, 60, and 120 minutes, and the plasma was immediately separated by centrifugation. The glucose and insulin concentrations were subsequently determined either by a coupled hexokinase procedure using a Chem 1A serum chemistry analyzer or, in the later case, a rat insulin RIA kit (Linco Research, Inc., St. Charles, Mo.).

Statistical comparisons were made by an analysis of variance (ANOVA) with a Duncan's Multiple Range test. A p value of <0.05 was considered as being statistically significant. All data are represented as the mean ± SEM.

Results

Study One

Two treatment groups, with five rats per group, were dosed IV with either (Leu²⁴, Ala³¹)hIGF-I (100 µg) or PBS.

The responses of plasma glucose and plasma insulin to the treatments are shown in FIGS. 8A and 8B, respectively. The data are expressed as a percentage of the values in the pre-treatment blood samples which were averaged and set at 100%. One IV injection of (Leu²⁴, Ala³¹)hIGF-I (100 µg) caused plasma insulin levels to be immediately (after 5 minutes), dramatically, and significantly (after 10 mins, P<0.001 vs. control) decreased to 25% of that of the control group, and remained depressed for 60 minutes. This fall in plasma insulin was accompanied by a brief but statistically significant (after 10 mins, P<0.05 vs. control) fall in blood glucose.

Study Two

The rats in three treatment groups, with 4–5 rats per group, were dosed IV with either (Leu²⁴, Ala³¹)hIGF-I (100/µg) or rhIGF-I (100/µg) and a control group was given PBS.

FIGS. 9A and 9B show the responses of plasma glucose (FIG. 9A) and plasma insulin (FIG. 9B) expressed as a percentage of the values in the pre-treatment blood samples which were averaged and set at 100%. Plasma insulin levels

decreased significantly (after 10 mins, p<0.001 vs. control), and remained depressed for 120 minutes in response to the IV injection of rhIGF-I. In addition, there was a brief but statistically significant fall in blood glucose (after 10 mins, p<0.01 vs. control). These are the expected responses to the injection of rhIGF-I. When (Leu²⁴, Ala³¹)hIGF-I was injected, there was a fall in insulin concentrations of a similar magnitude and degree of statistical significance (after 10 mins, p<0.001 vs. control) to the fall in insulin in the rats given wild-type rhIGF-I. The insulin levels after (Leu²⁴, Ala³¹)hIGF-I administration remained significantly suppressed to the end of the study (after 120 min., p<0.05 vs. control). There was also a small but statistically significant fall in blood glucose (after 10 min., p<0.05 vs. control).

Conclusion

In two separate experiments, (Leu²⁴, Ala³¹)hIGF-I, a molecule that binds to IGFbps with high affinity but binds poorly to IGF receptors, shows IGF-like activity on glycemic response variables when given by IV injection in a normal rat. This activity could not have been predicted from data in the literature. The IGF mutant showed suppressive effects on insulin secretion that were of a similar magnitude to the effect of wild-type rhIGF-I. The size of this effect, and its similarity to that seen following administering rhIGF-I itself, were surprising. The fall in glucose levels was not as large as that seen following the administration of rhIGF-I. This order of responses, a larger effect on insulin secretion than on blood glucose levels, follows from the sensitivity of these responses to rhIGF-I. A suppression of insulin secretion is seen at much lower doses of rhIGF-I than are needed to lower blood glucose levels. Furnsinn et al., *Endocrinology*, 135: 2144–2149 (1994).

Example 3

Injections of (Leu²⁴, Ala³¹)hIGF-I Into Diabetic Rats

Introduction

In Example 2 the IGF-I mutant reduced insulin secretion and lowered blood glucose in normal non-diabetic rats. It was unclear if, in the diabetic state, manipulating the endogenous IGF system would lead to similar changes in glycemic control. Therefore, an animal model of Type II diabetes was also chosen to test the glycemic activity of the IGF-I mutant. The animal chosen, the Zucker Rat, is a well-known model of obesity associated with diabetes in the rat. Stern et al., *Proc. Soc. Exp. Biol. Med.*, 139: 66–69 (1972). A strain of these rats, the Zucker Diabetic Fatty strain (ZDF), is a good model of Type II diabetes, since they become obese and insulin resistant at an early age, with progressive b-cell failure and then frank diabetes. Johnson et al., *Science*, 250: 546–549 (1990). It has been previously shown in ZDF rats that IGF-I administration can in the long term slow the onset and severity of Type II diabetes (WO 96/15148 published May 23, 1996). Therefore, the ZDF rat provides a relevant and sensitive animal model of Type II diabetes to examine the effects of IGF agonist compounds on glycemic control.

In the present example intravenous bolus injections of the IGF mutant were given and effects on the blood glucose and insulin concentrations were studied.

Methods

Eighteen 7-week-old male Zucker Diabetic Fatty rats (250–300 g, Genetic Models Inc.) were anesthetized using KETAMINET (62.5 mg/kg)/ROMPUN XYLAZINE™ (12.5 mg/kg) anesthesia. The right jugular vein was cannulated using a silicone rubber catheter and the rats were allowed to recover.

Study One

Two days after surgery the rats (5–6/group) were divided into three treatment groups and given IV via the jugular catheter 200 μ l of:

- 1) the PBS vehicle;
- 2) IGF mutant (100 μ g); or
- 3) rhIGF-I (100 μ g).

The dose of rhIGF-I was chosen as a dose that might cause a small fall in blood glucose levels and in blood insulin levels. The same dose of IGF mutant was given for comparison.

Study Two

Three days later, using the same rats the study was repeated. The rats that were dosed with IGF-I in the initial study received the IGF mutant in the repeat study and vice versa. Control animals received PBS in both experiments.

Measurements
Two blood samples were collected from each rat via the jugular cannula prior to IV dosing with the hormones and then blood samples were collected after 10, 30, 60, and 120 minutes. The plasma was immediately separated by centrifugation. The glucose concentration was measured by a coupled hexokinase procedure using a Chem 1A serum chemistry analyzer. Insulin concentration was measured using a rat insulin RIA kit (Linco Research, Inc., St. Charles, Mo.).

Statistical comparisons for each time point were made by an ANOVA with a Duncan's Multiple Range test. A p value of <0.05 was considered as being statistically significant. All data are represented as the mean \pm SEM, with five or six animals per treatment group.

Results

The results from Study One are shown in FIGS. 10A and 10B and from Study Two in FIGS. 11A and 11B. In each Figure the top panel (A) shows the percentage change in plasma insulin from baseline (set as 100%) and the bottom panel (B) the percentage change in plasma glucose from baseline (set as 100%).

In Study One the IV injection of rhIGF-I caused an immediate, moderate, and statistically significant ($p<0.01$ vs. control, excipient-treated rats) fall in blood glucose (FIG. 10B). In comparison, there was a small rise in blood glucose in the excipient-treated rats. Blood glucose then rebounded to the level in control animals. A smaller, but statistically significant ($P<0.05$ vs. control, excipient-treated rats) fall in blood glucose was also seen after 10 minutes in animals treated with the IGF mutant. Blood glucose then rebounded to control levels. After 120 minutes blood glucose drifted to lower levels in the animals treated with both rhIGF-I and the IGF mutant. This is suggestive of a long-term effect of the IGF mutant on blood glucose.

These changes in blood glucose were accompanied by changes in the concentrations of insulin in the blood. In the control rats blood insulin levels were high and the levels were maintained throughout the two hours of sampling. The injection of rhIGF-I caused a statistically significant ($p<0.01$ vs. control, excipient-treated rats), large, and maintained fall in plasma insulin. Injection of the IGF mutant also significantly ($p<0.01$, vs. control, excipient-treated rats) suppressed blood insulin concentrations. In this group insulin slowly returned to control levels two hours after the injection.

In Study Two (FIGS. 11A and 11B) similar results were obtained to those found in Study One and shown in FIGS. 10A and 10B. In Study Two an IV injection of rhIGF-I caused an immediate, moderate, and statistically significant ($p<0.01$ vs. control, excipient-treated rats) fall in blood glucose after 10 minutes. Blood glucose then rebounded to

the level in control animals after 30 minutes. In this study a similar and statistically significant ($p<0.05$ vs. control, excipient-treated rats) fall in blood glucose was seen after 10 minutes in animals treated with the IGF mutant. Blood glucose then rebounded to control levels, then drifted to lower levels. Once again, this is suggestive of a long-term effect of the IGF mutant on blood glucose.

The changes in blood glucose were accompanied by changes in the insulin concentrations in the blood. In the controls blood insulin levels were high and were maintained throughout the two hours of sampling. The injection of rhIGF-I caused a statistically significant ($p<0.01$ vs. control excipient-treated rats), large, and maintained fall in plasma insulin. Injection of the IGF mutant caused a similar and significant fall in blood insulin concentrations ($p<0.01$, vs. control, excipient-treated rats). In both groups insulin slowly returned toward control levels, but after two hours did not appear to have fully returned to pre-injection levels or the levels in control rats.

These experiments show that the acute administration of an IGF agonist 1) reduces insulin levels and 2) reduces blood glucose levels. In the present study only one injection was given, and the delivery was intravenous. Since one injection of the IGF mutant showed efficacy on glycemic parameters, it would be most likely that multiple injections would show similar acute glycemic effects leading to beneficial long-term cumulative effects. It would also be expected that injections given subcutaneously would have a similar effect to the intravenous injections used here, but may show a different time course, due to their slower absorption from the subcutaneous injection site.

In humans, infusions of rhIGF-I inhibit insulin release at much lower doses than those needed to reduce blood glucose (Hartman et al., *J. Clin. Invest.*, 91: 2453–2462 (1993)), and such low-dose euglycemic infusions of rhIGF-I also rapidly suppress fasting-enhanced pulsatile GH secretion. Therefore, a fall in insulin is expected to be a sensitive marker of the release of IGF-I after the administration of a molecule that prevents the interaction of an IGF with one of its IGF-BPs, assuming that the IGF agonist increases the bioavailability of IGF-I, causing endogenous active IGF-I levels to rise.

In the examples shown herein, using normal animals and diabetic animals, it can be deduced that sufficient IGF-I is activated to induce significant falls in blood insulin and blood glucose concentrations.

Example 4

Long-term Administration of (Leu²⁴, Ala³¹)hIGF-I to Hypox Rats

Introduction

The previous set of studies addressed the acute effect of administering molecules that preferentially bind to the IGF-BPs rather than to IGE receptors. The short-term effect of the administration of these molecules is now shown to cause an IGF-like agonist effect, as shown by a fall in blood glucose and insulin levels.

An important issue that then arises is the longer-term effects of administration of the IGF agonist. If the mechanism of these activities is a simple displacement of IGF, then it is possible that with time, with long-term exposure, with continual exposure, or with high-dose exposure, the response to the IGF agonist will show tachyphylaxis and the acute response will diminish. If the mechanism of action is other than simple displacement of IGF, then the long-term effect in animals of an IGF agonist is even less certain. In addition, the short-term studies on glucose regulation do not

show the longer-term effects of IGF-I, for example, the anabolic effects, the differentiative effects, the mitotic effects, and the effects on organ function. It was therefore important to administer an IGF agonist long term to animals.

The first model chosen was the hypophysectomized rat. The hypophysectomized rat is very sensitive to the effects of both GH and IGF-I. Guler et al., *Proc. Natl. Acad. Sci. USA*, (1988), supra; Clark et al., *Endocrine*, 3: 297–304 (1995). However, as endogenous IGF-I levels are very low in hypophysectomized rats (perhaps only 10% of normal), it was possible that the IGF mutant might not show activity by itself. Therefore, a group of hypophysectomized rats were also given recombinant human GH. GH treatment in hypophysectomized rats raises the levels of IGF-I in blood (Guler et al., *Proc. Natl. Acad. Sci. USA*, (1988), supra; Clark et al., *Endocrine*, supra), and it was reasonable to assume that this increased blood level of IGF-I might be activated by the co-administration of the IGF mutant.

Methods

Young female SD rats were hypophysectomized at Taconic Laboratories and delivered several days later. The rats were then weighed frequently and those whose weight increased or decreased by more than 7 grams between 2 and 3 weeks following hypophysectomy were excluded from the study. Twenty-five rats were grouped randomly into five groups based on their body weight, and then randomly assigned to cages, with four rats per cage. The animals were allowed ad libitum access to food and water and housed in a room controlled for temperature and lighting.

Experimental Groups

- 1) Excipient pump, excipient injections;
- 2) (Leu²⁴, Ala³¹)hIGF-I (10 μ g/day, by SC minipump), excipient injections;
- 3) (Leu²⁴, Ala³¹)hIGF-I (50 μ g/day, by SC minipump), excipient injections;
- 4) Recombinant human GH (NUTROPIN™ brand from Genentech, Inc., 20 μ g/day, by SC injection, two injections each of 10 μ g/day), excipient minipump; or
- 5) (Leu²⁴, Ala³¹)hIGF-I (50 μ g/day, by SC minipump) plus recombinant human GH (20 μ g/day, by SC injection, two injections each of 10 μ g/day).

The rats were dosed for one week.

Hormones

The mutant (Leu²⁴,Ala³¹)hIGF-I was administered by osmotic minipumps (ALZET₂₀₀₁™, Alza, Palo Alto, Calif.), which were placed in a subcutaneous tunnel in the dorsal neck region of the rats while they were anesthetized with KETAMINET™/XYLAZINE™ anesthesia. The pumps were filled with solutions so that the calculated daily dose was 50 μ g or 10 μ g, assuming that 24 μ l of solution was delivered each day for one week.

Recombinant human GH was administered for one week by twice daily SC injections of 10 μ g in 100 μ l volume.

Animals not receiving injections of rhGH received injections of vehicle and animals which were not administered IGF-I had a saline-filled pump implanted.

Measurements

Body weights were measured daily in the morning. At the end of the week of dosing the animals were anesthetized via CO₂ and exsanguinated via cardiac puncture. The remaining blood was allowed to clot and serum was separated and stored for further analysis.

Spleen, thymus, heart, liver, kidney, and perirenal fat were removed and weighed. The tibia was removed and fixed in neutral-buffered formalin for histological measurement of epiphyseal plate width.

Serum chemistries were measured on an automated Chem I analyzer. Serum insulin concentration was measured using

an RIA Kit supplied by Linco, Inc. To measure the IGF-I concentrations the serum was extracted using acid ethanol and the supernatant diluted to neutrality and assayed for IGF-I by RIA. The concentration of rat IGF-I in serum extract was measured by RIA using a kit from Diagnostic Systems Laboratories, Inc. This rat IGF-I assay does not measure human IGF-I. Human IGF-I concentration was measured by RIA at Genentech, Inc.

Tibia were sectioned longitudinally, stained with Toluene Blue, and mounted on microscope slides. The tibial epiphyseal plate width was measured using an ocular micrometer attached to a microscope.

Data were analyzed statistically by analysis of variance using one factor and two factor ANOVA.

Results and Discussion

Body Weight

FIG. 12 shows the final body weight gains of the animals at the end of the study. There was a significant increase in body weight as a result of hGH injections. This is the expected response in these GH-deficient hypophysectomized rats, which, when treated with excipients, failed to gain weight. By itself, the IGF agonist showed only a small amount of activity at 50 μ g per day. At 10 μ g per day the IGF agonist group had a mean weight gain that was numerically smaller than that of the control group. However, when the IGF agonist was given along with hGH, the IGF agonist showed an enhancement of the activity of hGH on whole body growth. This is shown in the final body weights in FIG. 12.

Organ Growth

The organ weight data is illustrated for spleen (FIGS. 13A and 13B), thymus (FIGS. 14A and 14B), and heart (FIGS. 15A and 15B). In each Figure the absolute organ weight is shown in A, the top panel, and the organ weight expressed as a percentage of body weight is shown in B, the bottom panel.

Absolute spleen weight (FIG. 13A) was increased by both doses of the IGF mutant alone, as it was by hGH. The response to hGH was greater than that to the IGF agonist. However, the combination of hGH plus IGF agonist doubled spleen size with evidence of a synergistic effect. When the spleen size was corrected for the growth of the whole body (by expressing the data as a percent of whole body weight), the IGF agonist again showed evidence of activity when given alone. However, now the synergistic effect of the combination of hGH and IGF mutant was more obvious, as there was a very large response to combination treatment.

Absolute thymus weight (FIG. 14A) was also increased by the high-dose IGF agonist alone. The responses to hGH and to the IGF displacer were almost equal. The combination of hGH plus IGF agonist also increased thymus weight. When the thymus size was corrected for the growth of the whole body (by expressing the data as a percent of whole body weight), the IGF agonist again showed evidence of activity when given alone, with the response again being similar to that caused by hGH. The combination of hGH and IGF mutant also increased relative thymic weight.

In the present Example the IGF agonist at both 10- and 50- μ g doses caused a significant increase in the absolute and the relative size of the heart (FIGS. 15A and 15B). In comparison, treatment with hGH slightly increased absolute heart weight and caused a fall in relative heart weight. Treatment with the IGF mutant in combination with hGH greatly increased the absolute weight of the heart and reversed the GH-induced decline in relative heart weight.

Bone Growth

The epiphyseal plate widths from the five treatment groups in this study are shown in FIG. 16. The width of the

epiphysis in the control hypophysectomized rats (about 200 microns) is similar to that reported in the literature in similar rats. Clark et al., *Endocrine*, 3: 717–723 (1995). When given by itself, the (Leu²⁴,Ala³¹)hIGF-I at 50 μ g/day induced a significant increase in bone growth (P<0.01 vs. control) 5
Low-dose (Leu²⁴,Ala³¹)hIGF-I did not increase bone growth. In comparison, the injections of rhGH induced a very large increase in bone growth, doubling plate width to around 400 μ m. The combination of rhGH and (Leu²⁴,Ala³¹) 10
hIGF-I did not cause a further increase in bone growth. However, an epiphyseal plate width of 400 μ m is near maximal, making it unlikely that a further increase could occur with combination dosing at the doses used. Since there was a growth response to (Leu²⁴,Ala³¹)hIGF-I given alone, it would be expected that at lower doses of hGH a greater 15
effect of the combination would be seen than for each agent given alone.

Serum IGF-I Levels

The levels of IGF-I in the blood were measured in two assays: one assay measured the amount of rat IGF-I in the blood, another the amount of human IGF-I in the blood. FIG. 20
17 shows the amount of rat IGF-I (FIG. 17A) and the amount of total IGF-I (FIG. 17B) in the blood in the five treatment groups. A remarkable aspect of these data (FIG. 17A) is that the blood levels of receptor-active IGF-I (endogenous rat IGF-I) are lower in the animals given the IGF agonist, yet 25
these lower blood IGF-I levels were associated with marked IGF-like responses in various organs and tissues. Clearly, it is counter-intuitive to observe evidence of increased activity of a hormone when the blood levels of the hormone are decreased.

The assay for total IGF-I (FIG. 17B) measures both rat and human IGF-I. In this assay there was a rise in total IGF-I in the rats given 50 μ g of IGF-I agonist, probably due to the IGF-I agonist binding to binding proteins and therefore being present in the blood. At the lower dose of 10 μ g of IGF 30
agonist, there was not a rise in total IGF in the blood. The total IGF in the blood did rise (p<0.05 vs. hGH alone) in the rats given the combination of GH and the IGF agonist, in contrast to the level of rat IGF-I, which tended to fall in this treatment group. This indicates that when GH generates 40
IGF-I and IGFBPs, there is spare binding capacity, and that this capacity has been in part filled by the IGF agonist.

Conclusion

It has been previously shown that the combination of GH and IGF-I, when administered together to animals, including humans (U.S. Pat. No. 5,126,324; Kupfer et al., *J. Clin. Invest.*, 91: 391–396 (1993)), shows greater activity than either agent alone. However, it was unclear if administering an IGF agonist would result in activation of the IGF axis and enhance the effect of GH administration. The present experiment shows that administering compounds that bind tightly to IGFBPs, but bind poorly to IGF receptors, can enhance the activity of GH. 50

It has also been reported that the administration of IGF-I to animals induces a different pattern of growth to that induced by GH. In particular, IGF-I administration causes the spleen, thymus, and kidney to show marked overgrowth. Guler et al., *Proc. Natl. Acad. Sci. USA*, 85: 4889–4893 (1988); Skottner et al., *Endocrinology*, 124: 2519–2526 (1989); Clark et al., *Endocrine*, 3: 297–304 (1995). 60
However, since the whole body size of the rats treated with IGF mutant alone showed only a small response, it might be expected that there would also be little change in the size of the organs of the treated animals. This was not the case; it is shown in this Example that the IGF mutant has very large 65
“IGF-like” effects on some of the organs known to be sensitive to IGF-I, i.e., the spleen and thymus.

A very surprising finding in this study was the effect of the treatments on the size of the heart. In previous studies in hypophysectomized or GH-deficient rats, it had been shown that the administration of IGF-I had a very small or no effect on cardiac size. Guler et al., *Proc. Natl. Acad. Sci. USA*, 85: 4889–4893 (1988); Skottner et al., *Endocrinology*, 124: 2519–2526 (1989); Clark et al., *Endocrine*, 3: 297–304 (1995).

In view of the efficacy of the IGF mutant as an agonist of IGF-I as shown above, the present invention is expected to have application in the treatment of a large group of disorders associated with, or characterized by, a lack of active IGF-I in the bloodstream. Representative of such disorders are diabetes, obesity, anabolic disorders, immunologic disorders, cardiac disorders, and renal disorders, as well as others noted above.

Example 5

Long-term Administration of (Leu²⁴,Ala³¹)hIGF-I to Dwarf Rats

Introduction

The previous Example addressed the long-term effect in hypophysectomized rats of administering molecules that preferentially bind to the IGFBPs rather than to IGF receptors, represented by (Leu²⁴,Ala³¹)hIGF-I. Hypophysectomized rats have very low serum IGF-I levels and very low levels of IGFBPs because they lack pituitary hormones. Fielder et al., *Endocrinology*, 137: 1913–1920 (1996). However, despite these low levels of endogenous proteins, the IGF mutant showed remarkable activity. The next model chosen to test the activity of the IGF mutant was the 30
GH-deficient dwarf (dw/dw) rat (Charlton et al., *J. Endocr.*, 119: 51–58 (1988)), which shows a growth response to both GH and IGF-I. Skottner et al., *Endocrinology*, 124: 2519–2526 (1989); Clark et al., *Endocrine* 3:717–727 (1995). The dw/dw rat is not totally GH-deficient, as is the hypophysectomized rat, and thus has higher levels of serum IGF-I and the IGFBPs. Therefore, the dose of the IGF mutant given to the dw/dw rats was increased, since the blood of the dw/dw rats was expected to contain more IGF binding capacity and more IGF. 40

In the hypophysectomized rat (Example 4) GH was given to produce greater amounts of IGF and IGFBP, and the IGF mutant tested in the presence of this exogenous hormone. In the present Example the addition of exogenous GH was repeated, as was the addition of exogenous IGF-I, to discover the activity of the IGF mutant in the presence of these exogenously administered hormones. The aim of the study was to discover the effects of administering a molecule that binds well to the IGFBPs but binds poorly to IGF receptors. 50

Methods

Animals

Young female dwarf rats (11–12 wk of age, 115–140 g) were bred by homozygous mating (Charles Rivers Laboratories) and delivered to the Genentech Animal House where they were housed five per cage on polystyrene chips and fed a standard animal chow and water ad libitum, and kept in a room of constant humidity and with controlled temperature and lighting. The animals were weighed, and based on uniformity of body weight, 37 of the animals were selected and randomized into treatment groups and cages to give six treatment groups of equal initial body weights (approximately 120 g). 65

Experimental Groups

The study consisted of six groups of rats with six or seven rats per group.

1)	Excipient control (Excipient pumps)	Excipient injections
2)	IGF mutant (IGF-M) (120 μ g/day by pump)	Excipient injections
3)	IGF-I (120 μ g/day by pump)	Excipient injections
4)	IGF-M (120 μ g/day) plus IGF-I (120 μ g/day)	Excipient injections
5)	Excipient pumps	hGH injections (50 μ g/d)
6)	IGF-M (120 μ g/day by pump)	hGH injections (50 μ g/d)

The rats were dosed for eight days.

Hormones

The mutant (Leu²⁴, Ala³¹)hIGF-I and native-sequence recombinant human IGF-I were administered by osmotic minipumps (ALZET₂₀₀₁TM, Alza, Palo Alto, Calif.), which were placed in a subcutaneous tunnel in the dorsal neck region while the rats were anesthetized with KETAMINE/XYLAZINETM anesthesia. The pumps were filled with solutions so that the calculated daily dose was 120 μ g per rat per day (1 mg/kg/day), assuming that the pump delivered as per the manufacturer's description (24 μ l of solution per day).

Recombinant human GH (50 μ g/day) was administered for eight days by twice-daily sc injection, with each injection being of 25 μ g in a volume of 100 μ l.

Animals not receiving injections of rhGH received injections of vehicle and animals which were not administered IGF-M had a saline-filled pump implanted.

Measurements:

Body weights were measured daily in the morning. After 8 days the animals were anesthetized using CO₂ and exsanguinated by cardiac puncture. The remaining blood was allowed to clot and serum was separated and stored for further analysis.

Spleen, thymus, heart, liver, kidney, and perirenal fat were removed and weighed. The tibia was removed and fixed in neutral-buffered formalin for histological measurement of epiphyseal plate width.

Serum chemistries were measured using a TECHNICON CHEMIPLUSTM analyzer. Serum insulin concentration was measured using an RIA kit supplied by Linco, Inc. To measure the IGF-I concentrations the serum was extracted using acid ethanol and the supernatant diluted to neutrality and assayed for IGF-I by RIA. The concentration of rat IGF-I in serum extract was measured by RIA using a kit from Diagnostic Systems Laboratories, Inc. Human IGF-I concentration was measured by RIA at Genentech, Inc.

Tibia were sectioned longitudinally, stained with TOLUENE BLUETM stain, and mounted on microscope slides. The tibial epiphyseal plate width was measured using an ocular micrometer attached to a microscope.

Data were analyzed statistically by analysis of variance using one-factor ANOVA followed by Duncan's Range Test. Results and Discussion

Body Weight

FIG. 18 shows the effect on body weight gain of the treatments over the 8 days of the study. On day 7 of the study it was decided to continue the study for one further day, and a new preparation of hGH was made and injected. The large

weight gain between Days 7 and 8 suggests that this preparation of hGH was different from that used between Days 0 and 7. Therefore, it was believed that the weight gain data for the hGH groups on Day 7 are a better reflection of the overall experiment than those on Day 8. There were significant increases in body weight as a result of infusions of IGF-I or injections of hGH, the expected responses in these GH-deficient dwarf rats. By itself, the IGF agonist showed only a small amount of growth-promoting activity, with the weight gain only approaching statistical significance at day 6 (excipient 1.3 \pm 1.1 g vs. IGF mutant 4.2 \pm 0.6 g, p<0.10). However, when the IGF mutant was given along with IGF-I, there was initially clear weight gain (Day one excipient -0.6 \pm 0.8 g, IGF mutant 0.5 \pm 0.6 g, IGF-I 1.2 \pm 0.4 g, IGF mutant plus IGF-I 3.3 \pm 0.8 g, p<0.05 vs. IGF mutant alone and IGF-I alone), but this response waned with time. In contrast, the difference between hGH alone and hGH plus the IGF mutant increased with time, reaching statistical significance by Day 5 (hGH alone 12.4 \pm 0.9 g vs. hGH plus IGF mutant 15.5 \pm 0.6 g, p<0.05). Therefore, the IGF mutant enhanced the activity of hGH on whole body growth. These effects are shown in the growth curves and the final body weights (FIG. 18).

Organ Growth

The organ weight data are illustrated for spleen (FIG. 19) and kidney (FIG. 20). In each Figure the absolute organ weight is shown in A (the top panel) and the organ weight expressed as a percentage of body weight is shown in B (the bottom panel).

The absolute spleen weight (FIG. 19A) tended to increase in all groups treated with the IGF mutant. For example, the spleen weights in excipient control rats (308 \pm 6 mg) were increased by treatment with the IGF mutant alone to 350 \pm 14 mg (p<0.10 vs. excipient). Combining the IGF-M with either IGF-I or hGH treatments also tended to give larger spleens (FIG. 19A). When the spleen weights are expressed as a percentage of body weight, similar increases in spleen size caused by the IGF mutant were seen (FIG. 19B).

The absolute kidney weight and the relative kidney weight are shown in FIGS. 20A and 20B, respectively. The responses of the kidney to the IGF mutant were similar to those of the spleen. For example, relative kidney weight was increased by treatment with the IGF mutant, especially when it was given in combination with IGF-I (IGF-I alone, 0.85 \pm 0.3% vs IGF-I plus IGF mutant 0.92 \pm 0.04%, p<0.05).

The retroperitoneal fat depot was removed from the rats at sacrifice and weighed. There was a tendency for the IGF mutant to decrease adipose tissue stores (placebo 0.56 \pm 0.09 g, IGF mutant 0.53 \pm 0.10 g, IGF-I 0.48 \pm 0.06 g, and IGF-I plus IGF mutant 0.43 \pm 0.05 g).

Serum Chemistries and Hormones

Table I shows data collected from analyses of the sera collected at sacrifice. There was some effect of the various treatments on the blood glucose levels in the rats, but there were very significant effects on the insulin levels. The diabetogenic effect of GH was shown by the significantly (p<0.05) increased insulin concentrations in GH-treated rats compared to excipient-treated controls. If the rats were co-treated with the IGF mutant, this increase in serum insulin (diabetogenic effect of GH) was prevented (Table I), as the serum insulin level in the co-treatment group was no different from the insulin level in the excipient control group.

TABLE I

Serum Measurements in Dwarf Rats						
Group	Glucose (mg/dl)	Insulin (ng/ml)	BUN (mg/dl)	Creatinine (mg/dl)	AST (U/l)	CK (u/l)
Excipient	198 ± 25	0.82 ± 0.04	19.3 ± 0.6	0.28 ± 0.02	137 ± 9	245 ± 47
IGF-M	174 ± 18	0.88 ± 0.09	17.8 ± 0.4	0.23 ± 0.03	143 ± 8	301 ± 44
IGF-I	190 ± 24	0.54 ± 0.06	16.0 ± 0.7*	0.22 ± 0.05	128 ± 11	225 ± 18
IGF M + IGF-I	193 ± 19	0.69 ± 0.07	13.8 ± 0.8*	0.18 ± 0.02*	101 ± 2*	138 ± 11*
hGH	192 ± 10	1.48 ± 0.29*	15.6 ± 0.7*	0.33 ± 0.02	123 ± 13	184 ± 29
hGH + IGF-M	202 ± 27	1.03 ± 0.20	15.3 ± 0.6*	0.25 ± 0.02	113 ± 9	229 ± 30

Data are Means ± SEM, *p < 0.05 vs excipient control.

The IGF mutant (IGF-M) significantly increased kidney size. Two indications of serum measurements of renal function, the serum creatinine and the blood urea nitrogen (BUN) levels, were reduced by the IGF mutant. Table I shows that the creatinine concentration in blood tended to be reduced by both IGF-I and the IGF mutant when they were given alone; however, these effects were not statistically significant. However, the combination of IGF-I plus IGF mutant significantly ($p < 0.05$) reduced serum creatinine, indicative of an improvement in renal function. A second measure of renal function, the blood urea nitrogen, was also reduced by IGF-I and by IGF-I plus the IGF mutant, with the response to the combination being significantly greater than that to IGF-I alone ($p < 0.05$).

The combination of IGF-I plus IGF mutant decreased the amounts of the enzyme AST (alanine serine transferase) and CK (creatinine kinase) in the blood compared to the excipient controls ($p < 0.05$). AST is a measure of cardiac damage or function and CK is a measure of skeletal muscle function. These data could be interpreted as showing beneficial effects of the IGF mutant on skeletal and cardiac muscle.

Bone Growth

The epiphyseal plate widths were significantly ($p < 0.05$) increased by hGH and by IGF-I treatment, compared to excipient-treated controls. The IGF mutant in all cases increased the mean plate widths compared to the individual treatments, but these increases did not reach statistical significance.

Serum IGF-I Levels

The levels of IGF-I in the blood were measured in two assays (FIGS. 21A and 21B). One assay (FIG. 21A) measured the amount of rat IGF-I, and another assay (FIG. 21B) measured both human and rat IGF-I (total IGF-I) in the blood in the five treatment groups. A remarkable aspect of these data (FIG. 21A) is that the blood levels of receptor-active IGF-I (endogenous rat IGF-I) were lower in the animals given the mutant IGF, yet these lower blood IGF-I levels were associated with marked IGF-like responses in various organs and tissues. Clearly, it is counter-intuitive to observe evidence of increased activity of a hormone when the blood levels of the hormone are decreased.

The assay for total IGF-I (FIG. 21B) measures both rat and human IGF-I. In this assay there was a rise in total IGF-I in the rats given 150 μ g of IGF-I agonist, probably due to the IGF-I agonist binding to binding proteins and therefore being present in the blood. The total IGF level in the blood did rise ($p < 0.05$ vs. IGF-I alone or hGH alone) in the rats given the combination of either IGF-I or GH and the IGF agonist, in contrast to the level of rat IGF-I, which fell in these combination treatment groups. This indicates that when GH generates IGF and IGF-BPs, there is spare binding capacity, and that this capacity has been in part filled by the IGF agonist.

Conclusion

These long-term data in the GH-deficient dwarf rat confirm and extend the long-term data collected in hypophysectomized rats; the IGF mutant produced anabolic and growth-promoting effects. In addition, this experiment confirms that the IGF mutant shows growth-promoting activity when administered with GH. Further, when the IGF mutant was given along with IGF-I to the dwarf rats, the IGF mutant could increase the activity of the administered IGF-I. Therefore, the present experiment shows that the long-term administration of compounds that bind tightly to IGF-BPs, but bind poorly to IGF receptors, can enhance the activity of endogenous IGF-I, exogenously administered GH, and exogenously administered IGF-I.

In this Example several important findings were made which extend the discoveries made in the hypophysectomized rat.

Firstly, the increased size of the kidney in the rats treated with the IGF mutant was accompanied by a fall in the concentration of creatinine and blood urea nitrogen in the blood. These falls in blood metabolites are a hallmark of an increased functioning of the kidney. Additional evidence of a functional effect on cardiac and skeletal muscle was also obtained. Evidence of a decrease in fat mass was obtained using the IGF mutant. This indicates uses for IGF agonists in controlling body composition, especially obesity. See, for example, U.S. Pat. No. 5,597,797 on use of GH and IGF-I to prevent or treat obesity.

Secondly, the administration of GH significantly increased the blood concentrations of insulin, while the co-administration of the IGF mutant prevented this rise in insulin. Therefore, this well-known diabetogenic effect of GH could be reversed by the co-administration of the IGF mutant.

Thirdly, these responses to the IGF mutant occurred in the presence of a fall in the blood concentrations of "active" rat IGF-I.

Therefore, long-term infusions of the IGF mutant showed multiple activities in 1) hypophysectomized rats, which have no detectable GH in their blood and very low IGF-I concentrations, and 2) dwarf rats, which have low GH and low IGF-I levels in their blood.

Example 6

Long-Term Administration of (Leu²⁴,Ala³¹)hIGF-I to Diabetic Rats

In this Example an animal model of diabetes was treated long-term with the IGF mutant. The animal chosen was a Type II diabetic rat (the Zucker Diabetic Fatty (ZDF) rat). This rat has relatively normal pituitary function (in terms of

GH secretion) and relatively normal serum IGF-I concentrations. Therefore, this animal model differs from the two previous rat models, which have a complete lack of GH (hypophysectomized rat) or a clear deficiency of GH (the dw/dw rat) and are both IGF deficient. In addition, in this Example, the mode of administration of the IGF mutant was altered from infusions (used in the two previous Examples) to multiple injections. Furthermore, the dose of the IGF mutant was increased because, as explained above, the ZDF rats are not as IGF deficient as the hypophysectomized or dw/dw rats and therefore have a larger amount of active IGF to be released. The main endpoints used to measure efficacy in the previous Examples were measures of body growth, while in the present Example the major endpoints were blood glucose and insulin concentrations, measures of the diabetic state of the animals.

In the earlier Examples it was shown that the acute intravenous administration of one injection of an IGF agonist 1) reduced insulin levels and 2) reduced blood glucose levels. Since one injection of the IGF agonist showed efficacy on glycemic parameters and long-term infusions showed efficacy (without tachyphylaxis on growth parameters), it was hypothesized that multiple injections would each show acute glycemic effects leading to beneficial long-term cumulative glycemic effects. It was also hypothesized that injections given subcutaneously would show efficacy. Therefore, the long-term study in diabetic rats described below was initiated

Methods and Experimental Groups

Animals

Obese male Zucker Diabetic Fatty (ZDF) rats (6–7 weeks of age, 225–250 g, Genetic Models Inc., Indianapolis, Ind. 46268) were group housed (3/cage) in a room controlled for temperature and lighting and fed the pelleted rat diet specified by the breeders (Purina 5008, 6% fat breeder chow) and tap water ad libitum. After three days acclimation, a blood sample was obtained from the tail vein for measurement of blood glucose and insulin, which was measured by rat-specific RIA (Linco Research Inc., St. Charles, Mo.). Animals were randomized into four treatment groups and into cages to give groups balance so as to have equivalent initial blood glucose levels, insulin levels, and body weights.

The experiment consisted of four groups of rats with seven rats per group:

-
- 1) Excipient injections three times a day,
 - 2) Injections of IGF mutant (IGF-M) (50 μ g three times a day),
 - 3) Injections of IGF-M (150 μ g three times a day), and
 - 4) Injections of rhIGF-I (150 μ g three times a day).
-

The injections were each of 100 μ l given three times a day. Hormones

The mutant (Leu²⁴,Ala³¹)hIGF-I was prepared at two concentrations (1.5 mg/ml and 0.5 mg/ml) so that with three injections per day two doses were given (a total of 450 μ g/d or 150 μ g/d, respectively). rhIGF-I was prepared at one concentration (1.5 mg/ml) so that with three injections per day one dose was given (450 μ g/d). A fourth group of rats were given the excipient. The injections, each of 100- μ l volume, were given at 7 am, 1 pm, and 7 pm for 15 days without disrupting the controlled diurnal lighting of the vivarium room. Body weights were measured each day at 1 pm.

Blood Sampling

Blood samples were collected from non-fasted rats on the evening before the first day of dosing and on days 3, 7, and

10. Blood samples (400 μ l) on days 3, 7, and 10 were taken by bleeding from a tail vein one hour after the mid-day injection. The blood was allowed to clot, and serum was separated and stored before analysis.

A glucose tolerance test was performed on day 14 of the study. Rats received their regular morning injection and then their food was withdrawn. After a 3–5-hour fast a blood sample (400 μ l) was taken via the tail vein. Rats then received 2 g/kg dextrose (50%) by intraperitoneal (ip) injection. A blood sample (200 μ l) was taken via tail vein at 10, 60, and 120 minutes after dextrose administration. Blood was allowed to clot and serum was separated and stored for measurement of glucose and insulin concentrations. After the glucose tolerance test was completed, animals were re-fed.

On day 15, one hour after the morning injection, rats were asphyxiated with carbon dioxide and bled by cardiac puncture. Blood was allowed to clot and serum was separated and frozen. Spleen, thymus, heart, liver, kidney, and perirenal fat were removed, blotted dry, and immediately weighed.

Statistical Analysis:

Statistical comparisons for each time point were made by an analysis of variance with a Duncan's Multiple Range test. A p value of <0.05 was considered as being statistically significant. All data are represented as the mean \pm SEM, with seven animals per treatment group.

Results

The body weight gains plotted against time are shown for the four different treatment groups in FIG. 22A. On day 13 the body weight gains were significantly increased vs. control (control, 93.3 \pm 2.9 g) for the IGF mutant at 50 μ g (100 \pm 2.1 g, p<0.05 vs. control), for the IGF mutant at 150 μ g (105.1 \pm 2.3 g, p<0.01 vs. control), and for IGF-I (114.5 \pm 1.6 g, p<0.0001 vs. control). The dose-related anabolic effect on body weight of the IGF mutant confirms the data in hypophysectomized and dwarf rats in the earlier examples.

In these diabetic animals the reason for the gain in body weight could also be an indirect effect rather than an anabolic effect because the treatments improved the diabetic state of the animals. Carlsson et al., *J. Endocrinol.*, 122: 661 (1989). This interpretation of the data is supported by the illustration in FIG. 22B showing the changes in blood glucose with time in these same animals. Blood glucose was measured before the injections were commenced, and the Figure shows the percentage change from this basal level. FIG. 22B shows that the blood glucose rose steadily in the rats treated with repeated injections of excipient to reach 141 \pm 21% of basal at day 10, while at this time injections of IGF-I (150 μ g three times a day) largely prevented (25 \pm 10%, p<0.001 vs control) this rise in blood glucose. Injections of the IGF mutant at 50 μ g (84 \pm 20%, p<0.05 vs. control) and the IGF mutant at 150 μ g (74 \pm 22%, p<0.05 vs. control) also slowed the rise of blood glucose in these diabetic rats at Day 10. At earlier time points, for example at day 7, the high-dose IGF mutant and IGF-I were equally effective at suppressing the rise in blood glucose.

FIGS. 23A and 23B show the effect of a glucose tolerance test in these diabetic rats on blood glucose and on insulin concentrations. A glucose tolerance test can be viewed as a simulated meal. The rats were bled and then given an intraperitoneal injection of glucose at 2 g/kg of body weight and were then bled again following the injection. FIG. 23A shows the change in blood glucose two hours following the glucose challenge. In the diabetic rats treated with injections of excipient, blood glucose rose substantially (105 \pm 21 mg/dl). Treatment with IGF-I caused blood glucose to rise

much less (39 ± 12 mg/dl, $P < 0.05$ vs control), while treatment with the IGF mutant ($150 \mu\text{g}$, tid) also blunted this rise (59 ± 11 mg/dl, $p < 0.1$ vs. control).

A possible reason for these beneficial effects of treatment with IGF-1 or the IGF mutant is provided by the insulin levels shown in FIG. 23B. In the rats treated with IGF-I more insulin secretion occurred in response to the glucose challenge (control, 8.1 ± 0.6 ng/ml vs. IGF-1, 12.2 ± 0.7 ng/ml, $p < 0.01$) than it did in the rats treated with IGF mutant at $50 \mu\text{g}$ (10.2 ± 0.7 ng/ml, $p < 0.05$ vs. control) or $150 \mu\text{g}$ (10.6 ± 0.6 ng/ml, $p < 0.05$ vs. control). Therefore, long-term treatment with the IGF mutant allowed greater insulin secretion in response to a simulated meal, which was associated with an improved ability to dispose of a glucose load.

Conclusions

This long-term study administering the IGF mutant to diabetic rats shows that blood glucose can be controlled in the long term by manipulating the endogenous IGF system. The body weight gain, perhaps an indirect marker of improved glucose control in diabetic animals, and the direct measures of glucose control (blood glucose and insulin), show the surprising efficacy of this class of receptor-inactive molecules in improving the diabetic state. In the present example multiple subcutaneous daily injections were given for 15 days with no evidence of tachyphylaxis. In the previous examples infusions of this molecule produced maintained anabolic responses. It is therefore reasonable to assume that most routes and patterns of delivery would also show similar efficacy. For example, oral formulations of molecules with long half-lives would be expected to be efficacious as would shorter-half-life molecules delivered by the oral or other routes.

Example 7

Phage-derived Peptides to Bind IGF-I and Binding Proteins

In the next set of examples, common α -amino acids may be described by the standard one- or three-letter amino acid code when referring to intermediates and final products. By common α -amino acids is meant those amino acids incorporated into proteins under mRNA direction. Standard abbreviations are listed in *The Merck Index*, 10th Edition, pp Misc-2–Misc-3. Unless otherwise designated, the common α -amino acids have the natural or "L"-configuration at the alpha-carbon atom. If the code is preceded by a "D", this signifies the opposite enantiomer of the common (α -amino acid. Modified or unusual α -amino acids such as norleucine (Nle) and ornithine (Orn) are designated as described in U.S. Patent and Trademark Office Official Gazette 1114 TMOG, May 15, 1990.

Based upon the results of experiments using the IGF mutant described above, it is predicted that other molecules, such as peptides or small organic molecules, which prevent the interaction of an IGF with an IGFBP, and bind poorly or not at all to the IGF-I receptor, should increase active IGF levels in a subject being treated. In addition, it is possible that another class of molecules, in particular peptidic or small molecules, might bind IGF-I itself at a site remote from that involved in receptor interactions in such a way as to inhibit or prevent the interaction of IGF-I with the IGFbps, but not the interaction of IGF-I with its receptor.

It has been shown that peptides which bind specifically and with measurable affinity to target molecules, such as proteins, can be identified from an initial library of many binding and non-binding peptides through binding selections using bacteriophage coat-protein fusions. Smith, *Science*, 228: 1315 (1985); Scott and Smith, *Science*, 249: 386 (1990); Cwirla et al., *Proc. Natl. Acad. Sci. USA*, 8: 309 (1990); Devlin et al., *Science*, 249: 404 (1990); reviewed by

Wells and Lowman, *Curr. Opin. Struct. Biol.*, 2: 597 (1992); U.S. Pat. No. 5,223,409. In addition, both proteins and peptides displayed on phage can be affinity-enhanced through iterative cycles of mutations, selection, and propagation.

Libraries of peptides differing in sequence at particular residue positions can be constructed using synthetic oligodeoxynucleotides. Peptides are displayed as fusion proteins with a phage coat protein (such as g3p or g8p) on bacteriophage particles, each of which contains a single-stranded DNA genome encoding the particular peptide variant. After cycles of affinity purification, using an immobilized target molecule, individual bacteriophage clones are isolated, and the amino acid sequence of their displayed peptides is deduced from their DNA sequences.

I. Construction of peptide-phage libraries

To identify a set of peptide molecules having the ability to bind to IGF-I or to an IGF binding protein, such as IGFBP-1 or IGFBP-3, several diverse phage libraries of peptides, of length ranging from 18 to 20 residues, were constructed. Peptides of this size were chosen in order to favor the selection of peptides capable of maintaining well-defined structures in solution. Because natural amino acid peptides of this size have a potential sequence diversity of 20^{18} – 20^{20} (i.e., 2.6×10^{23} to 1.0×10^{26}) variants, it is not practical to construct and test all such variants. Instead, certain residues were fixed or constant, which might be expected to allow or promote stable elements of peptide structure such as disulfide bonds or beta-turns, within each peptide.

Structural constraints or frameworks have previously been used for presentation of peptide libraries on phage and for subsequent, successive enhancement of binding affinities through mutation and selection. Such structured frameworks may favor stable binding conformations of peptide segments. By analogy, immunoglobulins provide a stable (and conserved) structural framework for presentation of a diversity of different peptide loops (CDR's, complementarity-determining regions) which can bind different antigens.

Used as a template for library constructions was a plasmid, pt4.g8 (DNA sequence shown in FIGS. 24A–C) expressing an antibody-recognizable (gD-tag) peptide fused to g8p of bacteriophage M13. This plasmid contains single-stranded and double-stranded origins of DNA replication. The phoA promoter and STII secretion-signal sequences are upstream of the gD peptide (underlined below), which is followed by a "linker" peptide (double underlined below), and then the g8p of bacteriophage M13:

SGTAMADPNRFRGKDLAGSPGGGSGG-
GAEGDDPAKAAAFNSLQASATEYI-
GYAWAMVVVIVGATI GIKLFKKFTSKAS (SEQ ID NO:24)

Several random-sequence peptide libraries (Table II) were constructed using single-stranded template-directed mutagenesis (Kunkel et al., *Methods. Enzymol.*, 204:125 (1991)), with the oligonucleotides described below.

TABLE II

Large Naive Libraries for g8 Display

Library	Oligo no.	Peptide motif	SEQ ID NO:
A	HL-300	SGTACX ₂ GPX ₁ CSLAGSP	25
B	HL-301	X ₄ CX ₂ GPX ₁ CX ₄	26
C	HL-302	X ₂₀	27
D	HL-303	X ₇ CX ₄ CX ₇	28
D	HL-304	X ₇ CX ₅ CX ₆	29
65 D	HL-305	X ₆ CX ₆ CX ₆	30
D	HL-306	X ₆ CX ₇ CX ₅	31

TABLE II-continued

Large Naive Libraries for g8 Display			
Library	Oligo no.	Peptide motif	
D	HL-307	X ₅ CX ₆ CX ₅	SEQ ID NO: 32
D	HL-308	X ₅ CX ₉ CX ₄	SEQ ID NO: 33
D	HL-309	X ₄ CX ₁₀ CX ₄	SEQ ID NO: 34

A. Beta-turn sequence motif

An example of a peptide of known three-dimensional structure is given by Wrighton et al., who selected a peptide agonist for the erythropoietin receptor (EPO-R) by phage display. Wrighton et al., *Science*, 273: 458 (1996). The peptide GGTYSCHFGLPTWVCKPQGG (SEQ ID NO:35) (having a disulfide bond joining the two Cys residues) forms a dimer of two beta hairpins, in the crystallized complex with EPO-R. Livnah et al., *Science*, 273: 464 (1996). Although the structure of the unbound form of this peptide in solution has not been reported, the beta-turn structure formed by this peptide in complex with EPO-R suggested that similar structures might be formed by peptides of the form CX₂GPX₄C (SEQ ID NO:36).

As one type of structured peptide library, a portion of the gD peptide was replaced with the motif CX₂GPX₄C (SEQ ID NO:37), leaving the upstream and downstream ("flanking") residues unchanged from that of the starting plasmid. Thus, this library was designed to display on phage particles the peptide SGTACX₂GPX₄CSLAGSP (SEQ ID NO:38), where X represents any of the 20 natural L-amino acids, fused to the linker and g8p described above. This library was constructed using the oligonucleotide HL-300: 5'-GCC TAT GCA TCT GGT ACC GCC TGC NNS NNS GGT CCT NNS NNS NNS TGT TCT CTG GCA GGT TCA CCA G-3' (SEQ ID NO:39), where N indicates a mixture of the nucleotides A, G, C, and T, and S represents a mixture of the nucleotides G and C.

An additional library was constructed to allow for further interactions within the peptide and/or with the target proteins by randomizing the flanking sequences as well. This library was constructed with the form X₄CX₂GPX₄CX₄ (SEQ ID NO:40) by using oligonucleotide HL-301: 5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS TGC NNS NNS GGT CCT NNS NNS NNS TGT NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:41).

B. Disulfide-loop motifs

Because many additional peptide conformations might be productive for binding to a given target protein, it was desirable to test other types of peptide sequence motifs in phage-displayed libraries. For example, a single disulfide bond within a small peptide may favor stable structures which allow for relatively higher-affinity binding than in unconstrained structures. Geysen et al., *Mol. Immunol.*, 23: 709 (1986); Wood et al., *Science*, 232: 633 (1986); Oldenburg et al., *Proc. Natl. Acad. Sci. USA*, 89: 5393 (1992); O'Neil et al., *Proteins*, 14: 509 (1992); McLafferty et al., *Gene*, 128: 29 (1993); Giebel et al., *Biochem.*, 34: 15430 (1995). Several peptide-phage libraries were therefore constructed, of the form X_mCX_nCX_k (SEQ ID NO:42), where m=4, n=10, and k=4, or where m=5, n=8-9, and k=4-5, or m=6, n=6-7, and k=5-6, or m=7, n=4-5, and k=6-7 (SEQ ID NOS:28 to 34). In these peptides, a disulfide bond is predicted to form a stabilizing constraint for peptide conformation.

These peptide libraries (see Table II) were constructed as X₇CX₄CX₇ (SEQ ID NO:43), using oligonucleotide HL-303:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:44);

5 X₇CX₅CX₆ (SEQ ID NO:45), using oligonucleotide HL-304:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:46);

10 X₆CX₆CX₆ (SEQ ID NO:47), using oligonucleotide HL-305:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:48);

15 X₆CX₇CX₅ (SEQ ID NO:49), using oligonucleotide HL-306:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:50);

20 X₅CX₈CX₅ (SEQ ID NO:51), using oligonucleotide HL-307:

25 5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:52);

30 X₅CX₉CX₄ (SEQ ID NO:53), using oligonucleotide HL-308:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:54); and

35 X₄CX₁₀CX₄ (SEQ ID NO:55), using oligonucleotide HL-309:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS TGC NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:56).

C. Unconstrained peptides

Unconstrained libraries (i.e., having no fixed residues within the peptide) have also yielded specific binding molecules. Scott and Smith, *Science*, supra; Cwirla et al., *Proc. Natl. Acad. Sci. USA*, supra; Devlin et al., *Science*, supra;

45 Kay et al., *Gene*, 128: 59 (1993). Such libraries may yield structured peptides, nevertheless, since noncovalent interactions may still induce structure in the bound and/or unbound forms. An unconstrained peptide library, of the form X₂₀ (SEQ ID NO:57), was constructed using oligonucleotide HL-302:

5'-GCT ACA AAT GCC TAT GCA NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS NNS GGT GGA GGA TCC GGA GGA G-3' (SEQ ID NO:58).

55 II. Polyvalent (q8) phage binding selections

The products of random mutagenesis reactions were transformed into XL1-BLUE™ *E. coli* cells (Stratagene) by electroporation and amplified by growing 15-16 h with M13K07 (Vieira and Messing, *Methods Enzymol.*, 153: 3-11 (1987)) or VCSM13 helper phage (Stratagene Corp.). Based upon plating of the initial transformations, the number of transformants per library was approximately 1.8×10⁸ for library HL-300, 7.9×10⁸ for HL-301, 5.0×10⁸ for HL-302, 5.3×10⁸ for HL-303, 5.6×10⁸ for HL-304, 5.0×10⁸ for HL-305, 6.3×10⁸ for HL-306, 4.5×10⁸ for HL-307, 1.9×10⁸ for HL-308, and 2.1×10⁸ for HL-309.

IGFBP-1, IGFBP-3, and IGF-I were biotinylated with a 1.5:1 molar ratio of a cleavable biotin reagent, EZ-LINK™ NHS-SS-Biotin (Pierce), to protein, using the manufacturer's instructions.

The initial selection of peptides for binding to IGFBP-1, IGFBP-3, or IGF-I was carried out using phage pools of approximately 10^{10} phage/ml (100 μ l total volume). MAXISORP™ 96-well plastic plates (Nunc) were coated with a solution of 2 μ g/ml of NEUTRAVIDIN™ brand avidin (Pierce) in 50 mM sodium carbonate buffer, pH 9.6, overnight at 4° C. The NEUTRAVIDIN™ solution was then removed, and the plates were incubated with a blocking solution of 5 g/l of bovine serum albumin, or 5 g/l of ovalbumin, or 5 g/l of instant milk in 50 mM sodium carbonate buffer, for 1–2 h at room temperature. The blocking solution was then removed, and a solution of biotinylated target protein was added. After 1–2 h at room temperature, the target solution was removed, and the plates were washed ten times with PBS/TWEEN™ surfactant (0.05% TWEEN-20™ in PBS buffer).

Phage from the libraries described above were pooled as follows: pool A consisted of HL-300 phage, pool B of HL-301 phage, pool C of HL-302 phage, and pool D of phage from the HL-303, HL-304, HL-305, HL-306, HL-307, HL-308, and HL-309 libraries. Phage were added in PBS/TWEEN™/albumin/biotin (PBS/TWEEN™ buffer with 1 μ M biotin, 5 g/l bovine serum albumin, or ovalbumin) to wells coated with each target, and with control wells that were coated with NEUTRAVIDIN™ or with albumin, but not biotinylated target. The phage were allowed to bind 5–15 h at room temperature. The plates were then washed ten times with PBS/TWEEN™ buffer.

Phage remaining bound to the plates were eluted by incubating with 50 mM DTT for 1–2 h at room temperature. The eluted phage were transfected into *E. coli* cells and allowed to grow overnight at 37° C. to amplify the phage.

The second and third cycles of binding selection were carried out as above, except that streptavidin (0.1 mg/ml) was included in the phage cocktails along with biotin. An aliquot was taken from each target-coated and control well incubated with each library, and serial dilutions of the diluted phage were performed to measure specific binding to target. The diluted phage were then transfected into *E. coli* cells and plated for colony counting.

The fourth round of binding selection was carried out on MAXISORP™ plates directly coated with 2 μ g/ml of each target protein, or with albumin only. The results of phage-binding selections in cycles 2–4 are shown in FIG. 25.

The same initial phage libraries (A, B, C, D) were also used for binding selections to directly-coated IGFBP-3. In this case, MAXISORP™ 96-well plastic plates (Nunc) were coated with a solution of 2 μ g/ml of IGFBP-3 in 50 mM sodium carbonate buffer, pH 9.6, overnight at 4° C. The target solution was then removed, and the plates were incubated with a blocking solution of 5 g/L of bovine serum albumin, for 1–2 h at room temperature. Phage were incubated with the plates as above, and non-binding phage washed away. The phage remaining bound were eluted by incubating with 20 mM HCl for 10 min at room temperature. Thereafter, the acid-eluted phage were neutralized with one-fifth volume of 1 M Tris-HCl, pH 8.0. Phage were transfected for colony counting as described above.

III. Screening of polyvalent phage clones (IGF-blocking phage assay)

Peptide-phage clones were isolated by mixing phage pools with *E. coli* cells, and plating onto antibiotic-containing media.

Colonies were isolated and grown with helper phage (as above) to obtain single-stranded DNA for sequencing. Peptide sequences selected for binding IGFBP-3, IGFBP-1, or IGF-I were deduced from the DNA sequences of phagemid clones. A number of such clones are represented by the peptide sequences in Tables III-V, respectively.

TABLE III

Peptide sequences from g8 display, IGFBP-3 selection		
Name	Peptide sequence	
4A3.1	SGTACYGGPEWCCSLAGSP	(SEQ ID NO: 59)
4A3.3	SGTACYGGPEWCCSLAGSP	(SEQ ID NO: 60)
4A3.4	SGTACYGGPEWCCSLAGSP	(SEQ ID NO: 61)
4B3.1	DLAICAEGPEIWCCEETS	(SEQ ID NO: 62)
4B3.2	DFWICLSGPGWEECLEWW	(SEQ ID NO: 63)
4B3.3	EESECFEGPGYVICGLVG	(SEQ ID NO: 7)
4B3.4	DMGVCADGPPWYVCEWTE	(SEQ ID NO: 8)
4B3.5	DMGVCADGPPWYVCEWTE	(SEQ ID NO: 8)
4C3.1	GSAGQGMTEEWAWIWEWKE	(SEQ ID NO: 64)
4C3.2	ELDGWVCIKVGEQNLCLYAE	(SEQ ID NO: 65)
4C3.4	ELDGWVCIKVGEQNLCLYAE	(SEQ ID NO: 66)
4C3.4	ELDGWVCIKVGEQNLCLYAE	(SEQ ID NO: 67)
4C3.5	ELDGWVCIKVGEQNLCLYAE	(SEQ ID NO: 68)
4D3.1	AIGGWCFIELDSLWCEEQIG	(SEQ ID NO: 69)
4D3.2	SEVVECVQVWENLVCSVEHR	(SEQ ID NO: 70)
4D3.3	SEVVECVQVWENLVCSVEHR	(SEQ ID NO: 71)
4D3.4	RVGAYISCSSETECWVEDLLD	(SEQ ID NO: 72)
4D3.5	WFKTVCYEWDEVEQCYLEE	(SEQ ID NO: 73)
4D3.6	SEVVECVQVWENLVCSVEHR	(SEQ ID NO: 74)
4D3.7	RLEEQCVENYEPSCSFTAN	(SEQ ID NO: 75)
4D3.8	SEVVECVQVWENLVCSVEHR	(SEQ ID NO: 76)
4D3.9	ETVANCDYEMDLCLCYGSDR	(SEQ ID NO: 77)
4D3.10	YHPISCDMHYYLICDETIVN	(SEQ ID NO: 78)
4D3.11	VAVEVCWDRHDGGYICTTDS	(SEQ ID NO: 4)
4D3.12	AEWAECWIAGDQLLCVQKDN	(SEQ ID NO: 79)
23A3.1	EPWLCQYYEAMLYLCWEEG	(SEQ ID NO: 80)
23A3.2	AEGMVGWGTGGWYNLDEL	(SEQ ID NO: 81)
23A3.3	SGGAIYVPEQFIAMAVGK	(SEQ ID NO: 82)
23A3.4	EPWLCQYYEAMLYLCWEEG	(SEQ ID NO: 83)
23A3.5	SGGAIYMPVEQFIAMAVGK	(SEQ ID NO: 84)
23B3.1	TGVDCQCGPVHCVCMDWA	(SEQ ID NO: 13)
23B3.2	EVLLCSDGPQLYLCELYA	(SEQ ID NO: 85)
23B3.4	SGVECVWGPQWGFVVEEY	(SEQ ID NO: 86)
23B3.5	DKEVCYLGPETWLFCWWP	(SEQ ID NO: 87)
23B3.6	EVLLCSDGPQLYLCELYA	(SEQ ID NO: 88)
23B3.7	GDVEICIEGPGWELCVWAD	(SEQ ID NO: 89)
23D3.1	FGWWSQCPTWVDVYVCNFEE	(SEQ ID NO: 90)
23D3.2	AMWVCVSDWETVEECIQYMY	(SEQ ID NO: 91)
23D3.3	AMWVCVSDWETVEECIQYMY	(SEQ ID NO: 92)
23D3.4	AMWVCVSDWETVEECIQYMY	(SEQ ID NO: 93)
23D3.5	AMWVCVSDWETVEECIQYMY	(SEQ ID NO: 94)
23D3.6	TNWFVFCESGHQDICWLAAE	(SEQ ID NO: 95)

TABLE IV

Peptide sequences from g8 display, IGFBP-1 selection			
Clone	Peptide sequence	Library	Frequency
HL-1	SEVGCRAGPLQLCEKYF (SEQ ID NO: 96)	A	6/6
HL-14	SEVGCRAGPLQLCEKYF (SEQ ID NO: 97)	B	5/6
HL-13	KDPVCGEGPLMRICERLFG (SEQ ID NO: 98)	B	1/6
HL-31	EVDGRWWIVETFLAKWDHMAG (SEQ ID NO: 99)	C	6/6

TABLE V

Peptide sequences from g8 display, IGF-I selection			
Clone	Peptide sequence	Library	Frequency
HL-8	WVMECGAGPWPEGCTFML (SEQ ID NO: 100)	B	5/6
HL-26	RKTSOGRGQEMCWETGGCS (SEQ ID NO: 101)	C	1/6
HL-25	SWERGELTYMKLCEYMRLQQ (SEQ ID NO: 102)	C	4/6
HL-30	EHGRANCLITPEAGKLARVT (SEQ ID NO: 103)	C	1/6

Such peptide-phage clones could represent specific target-binding peptides which either do or do not block ligand (IGF-I to IGFBP) binding, or any of a number of non-binding or background members of the selected pool. To distinguish among these possibilities, phage clones were tested for the ability to bind to IGFBP-1 or IGFBP-3 in the presence and absence of IGF-I.

IGFBP-1 or IGFBP-3 was coated directly onto MAXISORP™ plates as above. Phage from clonal cultures were mixed with IGF-I (100 nM final concentration), and incubated with the immobilized IGFBP for 1 hour at room temperature. The plates were then washed ten times, as above, and a solution of rabbit anti-phage antibody mixed with a goat-anti-rabbit conjugate of horseradish peroxidase was added. After an incubation of 1 hour at room temperature, the plates were developed with a chromogenic substrate, o-phenylenediamine (Sigma). The reaction was stopped with addition of ½ volume of 2.5 M H₂SO₄. Optical density at 490 nm was measured on a spectrophotometric plate reader.

Titration of several IGFBP-3-selected peptide-phage clones showed all were inhibited by IGF-I for binding to IGFBP-3 at some phage concentration (FIGS. 26 and 27). These peptides are thus likely to occupy an overlapping site with the IGF-binding epitope on IGFBP-3. Titration of an IGFBP-1-selected peptide-phage clone HL-1 showed it was inhibited by IGF-I for binding to IGFBP-1 at some phage concentration (FIG. 28). Additional peptide-phage clones were screened similarly, at a low concentration of phage, with and without IGF-I.

FIG. 29 shows the results of a blocking assay of several phagemid clones derived from three rounds of DTT elution, followed by one round of HCl elution, as described above. In each case, the phagemid clone was grown from a single colony overnight at 37° C. in a culture volume of 5 ml. The phage particles were precipitated and resuspended in 0.5 ml of PBS buffer. A 50-fold dilution of each phage solution was made into PBS/TWEEN™ buffer, and the phage were incubated with or without 100 nM IGF-I on an IGFBP-3-coated MAXISORP™ plate. As shown in FIG. 29, most clones were >40% inhibited for binding to IGFBP-3 at these phage concentrations, although clone 4D3.11 was only 5% inhibited under these conditions.

FIG. 30 shows the results of a blocking assay of several phagemid clones derived from three rounds of HCl elution, as described above. In each case, the phagemid clone was grown from a single colony overnight at 37° C. in a culture volume of 5 ml.

The phage particles were prepared as described above. In this case, as shown in FIG. 30, most clones were >80% inhibited for binding to IGFBP-3 at these phage concentrations, although clones 23A3.3 and 23A3.5 were only about 20% inhibited under these conditions.

The variation in the degree to which phage binding is blocked by a constant concentration of IGF-I, as a function

of phage dilution (FIGS. 26, 28), or as a function of peptide displayed (FIGS. 29–30), is of interest because, without being limited to any one theory, it may be predictive of (1) the degree of overlap between IGF-I- and peptide-binding epitopes on the IGFBP-3 molecule, and/or (2) the relative affinity of IGF-I versus phage-displayed peptide for binding to IGFBP-3. Since all peptide-phage clones tested here showed some degree of inhibition with IGF-I, it is likely that the epitope for peptide binding on IGFBP-3 for each lies within an area occupied by bound IGF-I. Peptide assays (see below) support this conclusion (i.e., case 1). On the other hand, without being limited to any one theory, it is possible that some peptide epitopes could be simply within an area for which binding of the phage particle displaying such peptides is sterically excluded by bound IGF-I.

The dependence of inhibition upon phage concentration, and the differences among phage clones (FIG. 26) may reflect case 2. In particular, phage clones whose binding to an IGFBP-3-coated plate was inhibited only at low phage concentrations (e.g., 4D3.3, 4B3.4, corresponding to peptides BP3-01-ox and BP3-02-ox, respectively) appear to yield higher-affinity peptides (see below) for IGFBP-3 than do those phage clones whose binding to an IGFBP-3-coated plate was inhibited both at high and at low phage concentrations (e.g., 4C3.2, 4D3.5, corresponding to peptides BP3-23 and BP3-24, respectively).

Thus, this type of phage-titration blocking assay may be generally useful as a means to predict the relative affinities and inhibitory potencies of peptides derived from phage-displayed libraries.

IV. Monovalent (g3) display of IGFBP-3-binding peptides

Affinity maturation of a peptide or protein sequence by successive rounds of random mutagenesis, selection, and propagation can be efficiently accomplished when the copy number of displayed peptides or proteins is limited. Bass et al., *Proteins*, 8: 309 (1990). Such an affinity maturation process is illustrated by the affinity maturation of hGH. U.S. Pat. No. 5,534,617. In this case, the copy number of displayed hGH was limited by fusing the displayed protein to g3, rather than to g8 of bacteriophage particles, restricting the expression level of hGH, and using a helper phage to supply wild-type g3p for phagemid packaging and propagation.

To select for higher-affinity peptide variants from pools of phage-displaying peptides on g8p, peptide cDNAs from two round-4 g8 library pools, 4B and 4D, were transferred to a g3 vector for monovalent phage display. Binding selections were carried out for three rounds, as described above, with acid elution of binding phage.

Peptide sequences obtained after three rounds of selections are shown in Table VI. Two clones, 4B3.3 and 4D3.11, dominated the selected pools, and were seen in the earlier, g8 phage selections. A third clone, 3Ai.2, represents a new peptide sequence that was not identified from g8 display. In phage-ELISA competition assays, the apparent affinity of the g3-4B3.3 and g3-4D3.11 clones was <100 nM; however, the corresponding peptides showed much weaker inhibition (see below).

TABLE VI

Peptide sequences from g3 display, IGFBP-3 selection			
Clone	Peptide sequence	Library	Frequency
65 3Ai.1 = 4B3.3	EESECFEGPGYVICLVG (SEQ ID NO: 7)	4B	6/10

TABLE VI-continued

Peptide sequences from g3 display, IGFBP-3 selection			
Clone	Peptide sequence	Library	Frequency
3Ai.2	VEDECWMGPDWAVCWWTWG (SEQ ID NO: 104)	4B	4/10
3Bi.1 = 4D3.11	VAWEVCWDRHDQGYICTTDS (SEQ ID NO: 4)	4D	10/10

It is anticipated that affinity improvements can be obtained by iteratively mutating, selecting, and propagating peptide-phage libraries, as described for hGH. See, e.g., U.S. Pat. No. 5,534,617.

V. Peptide assays

Peptides were synthesized corresponding to a number of phage-derived sequences. In cases where two Cys residues were found in the peptide sequence, the disulfide (oxidized or "ox" suffix) monomeric form of the peptide was prepared and purified. In cases where four Cys residues were found, the {1-4, 2-3}disulfide form was prepared and purified.

The ability of these peptides to bind IGFBP-1 or IGFBP-3 and block IGF-I binding was tested in one or more of the following assays.

A. BIAcore™ competition assay (for IGFBP-3 binders)

IGF-I was immobilized on a dextran chip for inhibition assays using a BIAcore™ 2000 surface-plasmon-resonance device (BIAcore, Inc., Piscataway, N.J.) to measure free binding protein. IGF-I was biotinylated as described above, and injected over a chip to which streptavidin had been coupled (BIAcore, Inc.) to give 400 to 800 RU (response units) of immobilized IGF-I. The IGF-I showed no detectable dissociation over the time course of each experiment. Serial dilutions of peptide were mixed with a constant concentration (40 nM) of IGFBP-3. After incubation for >1 hour at room temperature, an aliquot of 20 μ L was injected at a flow rate of 20 μ L/min over the IGF-I chip. Following the injection, a response reading was taken to measure the relative amount of IGFBP-3 bound to the IGF-I.

The results (FIGS. 31-32) show a dose-response curve for each peptide's inhibition of IGFBP-3 binding to the chip. In particular, the most effective inhibitors of IGFBP-3 binding tested were peptides BP3-01-ox (corresponding to phage clone 4D3.3), and a truncated form of this peptide, BP15 (see Table VII). In that table, a disulfide bond is formed between the two Cys residues of each 2-Cys-containing peptide. For peptides containing four cysteines, the two Cys* residues form a disulfide bond and the remaining two form a second disulfide bond. These peptides showed IC₅₀'s of 2 μ M and 0.75 μ M, respectively. Other peptides such as BP3-4D3.11 (phage clone 4D3.11 from g8 display and 3Bi.1 from g3 display) showed inhibition with IC₅₀'s of <10 μ M.

IGFBP-1 did not show binding to IGF-I immobilized in this manner.

TABLE VII

Inhibition of IGF-I binding to IGFBP-3 by synthetic peptides		
Peptide	Sequence	BIAcore™ Assay IC ₅₀ (μ M)
BP3-23	ELDGWVCIKVGQNLCYLAEG-nh2 (SEQ ID NO: 1)	220
BP3-24	WFKTVCYEWEDVQCYYTLEEG-nh2 (SEQ ID NO: 2)	100-300

TABLE VII-continued

Inhibition of IGF-I binding to IGFBP-3 by synthetic peptides		
Peptide	Sequence	BIAcore™ Assay IC ₅₀ (μ M)
5 BP3-25	RVGAYISCSSETECWVEDLLDG-nh2 (SEQ ID NO: 3)	>1000
10 BP3-4D3.11	VAWEVCWDRHDQGYICTTDS (SEQ ID NO: 4)	<10
BP3-4D3.11DEL	AWEVCWDRHDQGYICTTDS (SEQ ID NO: 5)	80
BP13	CWDRHDQGYICTTDS (SEQ ID NO: 6)	>1000
15 BP3-4B3.3	EESECFEGPGYVICGLVG (SEQ ID NO: 7)	80
BP3-02-OX	DMGVCADGPWVWVCEWTE (SEQ ID NO: 8)	12
BP3-01-OX	SEEVWCWPVAEWYLCNMWG (SEQ ID NO: 9)	2
20 BP15	SEEVWCWPVAEWYLCN (SEQ ID NO: 10)	0.75
BP16	VCWVPAEWYLCNMWG (SEQ ID NO: 11)	30
BP17	VCWVPAEWYLCN (SEQ ID NO: 12)	9
BP06	TGVDCQC*GPVHC*VCMDWA (SEQ ID NO: 13)	5
25 BP08	TVANCDC*YMPLC*LCYDSD (SEQ ID NO: 14)	15

where nh2 means that the peptide has been blocked with an amide and where the C* indicates a cysteine that has been linked to another cysteine in the peptide.

B. Plate assays

1. Biotin-BP assay (for IGFBP-1 binders)

For inhibition of IGFBP-1 binding, IGFBP-1 (as well as IGFBP-3 as a control) were biotinylated as described above. MAXISORP™ plates were coated and blocked as described above, using 2 μ g/mL IGF-I. In a separate plate, serial dilutions of peptide were premixed with a constant concentration (20 nM) of biotinylated IGFBP-1. After 20 min. at room temperature, the mixture was added to the IGF-I plate. The peptide/IGFBP mix was then removed and the plate was washed ten times with PBS buffer containing 0.05% TWEEN-20™ surfactant. Bound biotin-IGFBP-1 was then detected using streptavidin-conjugated horseradish peroxidase, and a chromogenic substrate.

The results of assays of IGFBP-1 inhibition (FIGS. 33 and 34) show that peptides bp1-01 and bp1-02 (Table VIII) inhibited IGFBP-1 binding with IC₅₀'s of about 0.18 and 0.05 μ M, respectively. In contrast, the IGFBP-3 selectant BP3-01-ox showed little or no inhibition of IGFBP-1 binding. Conversely, bp1-01 and bp1-02 showed no inhibition of biotinylated IGFBP-3 binding in this assay. A bp1-01 variant in which the Cys residues were changed to serine showed an IC₅₀ of greater than 10 μ M.

TABLE VIII

Inhibition of IGF-I binding to IGFBP-1 by synthetic peptides		
Peptide	Sequence	ELISA IC ₅₀ (μ M)
60 bp1-01	CRAGPLQWLCEKYFG-nh2 (SEQ ID NO: 15)	0.18
bp1-02	SEVGCRAGPLQWLCEKYFG-nh2 (SEQ ID NO: 16)	0.05

2. Radiolabeled IGF assay (for IGFBP-3 binders)

As an additional assay of peptide activity, several peptides were tested in an assay using ¹²⁵I-labeled IGF-I to measure

inhibition of IGFBP binding, as described in Example 1 (Assay 3). Serial dilutions of peptide were added to an IGFBP-1 or an IGFBP-3 plate. Thereafter, ^{125}I -labeled IGF-I was added and the plates were incubated for 2 hours. The plates were then washed and counted to determine the amount of bound IGF-I.

FIG. 35 shows the inhibition of two IGFBP-3-selected peptides, bp3-01-ox and bp3-02-ox, for IGF-I binding to an IGFBP-3 plate. In contrast, these peptides did not inhibit IGF-I binding to an IGFBP-1-coated plate (FIG. 36).

C. In vitro activation (KIRA)

The ability of several synthetic peptides to block IGF-I binding to IGFBPs and release functional IGF-I was tested in a KIRA assay of IGF-I activity, as described in Example 1. Cells were treated with IGF-I alone, peptide alone, peptide plus IGF-I, IGF-I plus binding protein (IGFBP-1 or IGFBP-3), or IGF-I plus binding protein (IGFBP-1 or IGFBP-3) and peptide.

The results (FIG. 37) show that the peptides alone had no activity. Furthermore, when mixed with IGF-I, the peptides did not significantly alter IGF-I activity. The peptide BP3-01-ox also showed no significant effect on IGF-I activity when mixed with IGF-I plus IGFBP-1 or IGFBP-3. However, both peptides bp1-01 and bp1-02 appeared to block the inhibition of IGF-I activity by IGFBP-1, with IC₅₀'s of 397 nM and 187 nM, respectively. Neither of these peptides showed activity with IGF-I and IGFBP-3.

In addition, the IGFBP-3 peptide BP15 incubated with IGFBP-3:IGF-1 (at a 3:1 ratio) caused an activation of the KIRA assay, where blocking of IGFBP-3 occurred with an IC₅₀ of 95 μM for BP15. It is believed that this would drop to 30 μM with a 1:1 ratio.

D. BIAcore™ competition assay (for IGFBP-3 binders)

IGF-II was immobilized on a dextran chip for inhibition assays using a BIAcore™ 2000 surface-plasmon-resonance device (BIAcore, Inc., Piscataway, N.J.) to measure free binding protein. IGF-II was biotinylated as described above, and injected over a chip to which streptavidin had been coupled (BIAcore, Inc.) to give approximately 1500 RU of immobilized IGF-II. The IGF-II showed no detectable dissociation over the time course of each experiment. Serial dilutions of peptide were mixed with a constant concentration (20 nM) of IGFBP-3. After incubation for ≥ 1 hour at room temperature, an aliquot of 20 μl was injected at a flow rate of 20 $\mu\text{l}/\text{min}$ over the IGF-II chip. Following the injection, a response reading was taken to measure the relative amount of IGFBP-3 bound to the IGF-II.

The results (e.g., see FIG. 38) show a dose-response curve for each peptide's inhibition of IGFBP-3 binding to IGF-II. Peptides bp3-01-ox, BP14, BP15, and BP17 showed IC₅₀'s of 0.92 μM , 1.0 μM , 0.78 μM , and 5.1 μM , respectively. Thus, these peptides inhibit the binding of IGFBP-3 both to IGF-I and to IGF-II.

Example 8

NMR Spectra of Peptide bp1-01

^1H NMR data were collected on peptide bp1-01 in H₂O solution at 30° C. and pH 5.2 at a concentration of 6.7 millimolar. For generation of the one-dimensional NMR spectrum of FIG. 39, a total of 5.0 mg of purified peptide was dissolved in 440 μl H₂O containing 7% (v/v) $^2\text{H}_2\text{O}$ for the spectrometers field frequency lock. The pH of the sample was adjusted to 5.2 by the addition of 3 μl of 1N NaOH. The spectrum consisting of 32 transients was collected on a BRUKER AMX-500™ spectrometer (^1H frequency of 500.13 MHz) equipped with a 5-mm triple-axis, pulsed-field

gradient probe. The chemical shift scale is in parts per million, referenced to the water resonance at 4.75 p.p.m. The water resonance was suppressed by the excitation sculpting method. Hwang and Shaka, *J. Magn. Reson.*, 112A: 275–279 (1995).

In addition to the one-dimensional spectrum of FIG. 39, two-dimensional double-quantum-filtered correlation spectroscopy (2QF-COSY), total correlation spectra (TOCSY), and rotating-frame Overhauser effect spectra (ROESY) were collected. The experiments were recorded as described by Cavanagh et al. in "Protein NMR Spectroscopy, Principles and Practice" (Academic Press, San Diego: ISBN 0-12-164490-1, 1995), except that pulsed-field gradients were used for coherence selection in the 2QF-COSY (van Zijl et al., *J. Magn. Reson.*, 113A: 265–270 (1995)), and excitation sculpting was used to suppress the water resonance in the TOCSY and ROESY experiments (Hwang and Shaka, supra). After lyophilization and dissolution of the peptide in $^2\text{H}_2\text{O}$, a 2-D ROESY spectrum (Cavanagh et al., supra) and a COSY spectrum with a 35° C. mixing pulse (Cavanagh et al., supra) were acquired. Complete ^1H resonance assignments were obtained from these data by standard methods. Wuethrich, in "NMR of proteins and nucleic acids", (John Wiley & Sons, New York: ISBN 0-471-82893-9, 1986).

Evidence of a well-defined three-dimensional structure for bp1-01 was obtained from the following:

- 1) The resonance positions in the one-dimensional spectrum (FIG. 39) are significantly different from those expected in an unstructured peptide. For example, the amide protons span a range of 8.73–6.64 p.p.m. (unstructured peptides are in the range 8.50–8.00 p.p.m.); and the methyl groups are at lower chemical shift (0.82 p.p.m. for Leu6, and 0.66, 0.59 p.p.m. for Leu9) than for an unstructured molecule (0.90–0.85 p.p.m.).
 - 2) Scalar coupling constants between amide and alpha protons (obtained from the 2QF-COSY spectrum) are distinct from the averaged values observed in unstructured peptides. The values less than 5.5 Hz for Gln7, Trp8, Cys10, Glu11, and Lys12 are indicative of a helix spanning these residues. The values greater than 8.0 Hz observed for Leu6, Tyr13, and Phe14 are indicative of an extended conformation in these regions. Scalar coupling constants were also measured between alpha and beta protons in the COSY-35 spectrum. These data indicate that the side chains of Cys1, Gln7, Trp8, Cys10, Tyr13, and Phe14 have fixed chi-1 angles, i.e., these side chains do not sample the range of chi-1 rotamers that are populated in unstructured peptides.
 - 3) Peaks in the ROESY spectra indicate that there are many proton—proton contacts (<5 Å) between residues that are not adjacent in the primary sequence. These can only occur if the peptide folds up into a well-defined structure. Contacts between residues at position i and i+3 in the primary sequence are prevalent between Leu6 and Tyr13, consistent with the presence of a helix in this region. Many contacts are observed between the three aromatic side chains (Trp8, Tyr13, and Phe14) and the leucine methyl groups (Leu6 and Leu9), indicating the presence of a mini-hydrophobic core along one face of the helix.
- The NMR data were used to derive restraints that could be used to determine a three-dimensional model of the bp1-01 structure. Dihedral angle restraints were derived from the amide-alpha and alpha-beta scalar coupling constants via an appropriate Karplus relationship. Karplus, *J. Phys. Chem.*, 30: 11–15 (1959). Distance restraints were introduced between protons which exhibited a through-space interaction in the ROESY spectrum; the size of the upper bound,

and corrections to the upper bound because of peak overlap or resonance degeneracy were as described by Skelton et al., *Biochemistry*, 33: 13581–92 (1994). These restraints were used to generate a family of structures using the program DGII (Havel, *Prog. Biophys. Mol. Biol.*, 56: 43–78 (1991)), which were subsequently refined by restrained molecular dynamics with the program Discover (MSI, San Diego) using the AMBER all atom force field. Weiner et al., *J. Comput. Chem.*, 7: 230–252 (1986). The resulting structures converged to a single global fold (mean root-mean-squared deviation from the mean structure of 0.37 Å for N, C-alpha, and carbonyl carbons of residues 4–14). The best 25 structures (least violation of the input restraints) agreed with the input data very well (no distance restraint violations greater than 0.1 Å and no dihedral angle violations greater than 1.4°), and had good covalent geometry as judged by the program PROCHECKTM. Laskowski et al., *J. Appl. Cryst.*, 26: 283–291 (1993).

A representative member of the ensemble is shown in FIGS. 40A and 40B. According to the Kabsch and Sander secondary structure algorithm within the PROCHECKTM program, bp1-01 contains a reverse turn centered at Pro5-Leu6 (Type I) and an alpha helix from Gln7 to Lys12; Leu6 and Tyr13 are extensions of the main helix. The residues Cys1, Arg2, Ala3, and Gly15 are not well defined by the NMR restraints and may be more flexible in solution than residues Gly4 to Phe14. FIGS. 40A and 40B indicate that the aromatic rings of Trp8, Tyr13, and Phe13 pack onto the aliphatic portions of Leu6, Leu9, and Lys12 to form a relatively flat, hydrophobic face on one side of the peptide. This packing is accomplished by the periodicity of the helix (bringing Leu6, Leu9, and Tyr13 together), the chi-1 angle of Trp8 (the chi-1 of +60 degrees, relatively unusual for this amino acid, allows Trp8 to pack onto Leu9), and the extended nature of the peptide backbone of Phe14 (allowing the side chain of Phe14 to fold back on to Leu9).

Although bp1-01 is prone to self-association under some conditions, the three-dimensional structure observed for bp1-01 is not stabilized by such intermolecular interactions.

The hydrophobic packing within the helix undoubtedly contributes to the stability of the fold. The disulfide bond between Cys1 and Cys10 is also important for structure and function. Peptides in which both cystine residues are replaced by serine have much reduced affinity for IGFBP-1 (see Example 7) and exhibit no evidence of helical or other stable structure in one- and two-dimensional NMR spectra, suggesting that the structure displayed by bp1-01 is important for its function. Without being limited to any one theory, one possibility is that the reverse turn (Pro5-Leu6) and disulfide bond help to initiate the first turn of helix (Gln7-Cys10), which is then propagated by the hydrophobic side chain-side chain contacts involving residues in the second turn of helix (Glu11-Phe14).

The structure of bp1-01 shown in FIG. 40 suggests several possibilities for the way in which the peptide is able to bind to IGFBP-1, thereby enhancing levels of active IGF-I.

- 1) The flat, hydrophobic surface formed by the side chains of Leu6, Leu9, Trp8, Tyr13, and Phe14 in the bp1-01 monomer may pack onto an exposed hydrophobic region of IGFBP-1; in this case the helix is the key structural feature recognized by IGFBP-1;
- 2) BP-1 may recognize some part of bp1-01 other than the hydrophobic face of the helix, e.g., the five N-terminal residues preceding the helix, or a combination of these residues and the face of the helix away from hydrophobic side chains. In this case, the helix may still be important as a scaffold to set up a geometry appropriate for IGFBP-1 binding elsewhere in the peptide.
- 3) bp1-01 may bind to IGFBP-1 in an aggregated form. Given the likelihood of self-association of bp1-01 via the hydrophobic face of the helix (Leu6, Trp8, Leu9, Tyr13, and Phe14), some other region of the peptide would be involved in IGFBP-1 contact. In this case the helices and the contacts between them would act as a scaffold to present the IGFBP-1 binding region.

Below are the coordinates for the representative structure that is shown in FIG. 39 (standard protein database format).

REMARK	BP1-01 IGF-I-BINDING PROTEIN BINDING PEPTIDE								
REMARK	REPRESENTATIVE STRUCTURE FROM ENSEMBLE CALCULATED								
REMARK	WITH NMR-DERIVED RESTRAINTS								
ATOM	1	N	CYS	1	-5.022	-1.559	0.892	1.00	0.00
ATOM	2	CA	CYS	1	-4.328	-1.133	-0.335	1.00	0.00
ATOM	3	C	CYS	1	-3.837	0.310	-0.220	1.00	0.00
ATOM	4	O	CYS	1	-2.646	0.544	-0.024	1.00	0.00
ATOM	5	CB	CYS	1	-5.220	-1.340	-1.567	1.00	0.00
ATOM	6	SG	CYS	1	-4.498	-0.877	-3.166	1.00	0.00
ATOM	7	1H	CYS	1	-5.842	-0.987	1.034	1.00	0.00
ATOM	8	2H	CYS	1	-5.296	-2.527	0.807	1.00	0.00
ATOM	9	3H	CYS	1	-4.400	-1.455	1.682	1.00	0.00
ATOM	10	HA	CYS	1	-3.456	-1.777	-0.448	1.00	0.00
ATOM	11	1HB	CYS	1	-5.484	-2.396	-1.625	1.00	0.00
ATOM	12	2HB	CYS	1	-6.143	-0.773	-1.446	1.00	0.00
ATOM	13	1LG	CYS	1	-4.970	-1.181	-3.550	1.00	0.00
ATOM	14	2LG	CYS	1	-4.093	-0.385	-2.924	1.00	0.00
ATOM	15	N	ARG	2	-4.756	1.275	-0.336	1.00	0.00
ATOM	16	CA	ARG	2	-4.410	2.685	-0.348	1.00	0.00
ATOM	17	C	ARG	2	-3.778	3.012	-1.703	1.00	0.00
ATOM	18	O	ARG	2	-2.559	3.112	-1.836	1.00	0.00
ATOM	19	CB	ARG	2	-3.519	2.999	0.862	1.00	0.00
ATOM	20	CG	ARG	2	-3.052	4.460	0.950	1.00	0.00
ATOM	21	CD	ARG	2	-4.211	5.459	0.892	1.00	0.00
ATOM	22	NE	ARG	2	-3.742	6.823	1.164	1.00	0.00
ATOM	23	CZ	ARG	2	-4.455	7.940	0.945	1.00	0.00
ATOM	24	NH1	ARG	2	-5.701	7.877	0.455	1.00	0.00
ATOM	25	NH2	ARG	2	-3.914	9.134	1.222	1.00	0.00
ATOM	26	H	ARG	2	-5.723	1.024	-0.475	1.00	0.00

-continued

ATOM	27	HA	ARG	2	-5.340	3.246	-0.248	1.00	0.00
ATOM	28	1HB	ARG	2	-4.084	2.770	1.767	1.00	0.00
ATOM	29	2HB	ARG	2	-2.647	2.346	0.839	1.00	0.00
ATOM	30	1HG	ARG	2	-2.523	4.585	1.896	1.00	0.00
ATOM	31	2HG	ARG	2	-2.356	4.680	0.142	1.00	0.00
ATOM	32	1HD	ARG	2	-4.657	5.430	-0.103	1.00	0.00
ATOM	33	2HD	ARG	2	-4.962	5.189	1.635	1.00	0.00
ATOM	34	HE	ARG	2	-2.813	6.913	1.550	1.00	0.00
ATOM	35	2HH1	ARG	2	-6.227	8.724	0.294	1.00	0.00
ATOM	36	1HH1	ARG	2	-6.122	6.981	0.254	1.00	0.00
ATOM	37	1HH2	ARG	2	-2.979	9.192	1.599	1.00	0.00
ATOM	38	2HH2	ARG	2	-4.442	9.979	1.061	1.00	0.00
ATOM	39	N	ALA	3	-4.640	3.150	-2.718	1.00	0.00
ATOM	40	CA	ALA	3	-4.259	3.352	-4.106	1.00	0.00
ATOM	41	C	ALA	3	-3.843	4.805	-4.352	1.00	0.00
ATOM	42	O	ALA	3	-4.546	5.555	-5.027	1.00	0.00
ATOM	43	CB	ALA	3	-5.430	2.931	-5.001	1.00	0.00
ATOM	44	H	ALA	3	-5.626	3.058	-2.522	1.00	0.00
ATOM	45	HA	ALA	3	-3.412	2.706	-4.345	1.00	0.00
ATOM	46	3HB	ALA	3	-5.643	1.872	-4.851	1.00	0.00
ATOM	47	1HB	ALA	3	-6.320	3.511	-4.753	1.00	0.00
ATOM	48	2HB	ALA	3	-5.182	3.096	-6.049	1.00	0.00
ATOM	49	N	GLY	4	-2.681	5.191	-3.818	1.00	0.00
ATOM	50	CA	GLY	4	-2.069	6.481	-4.087	1.00	0.00
ATOM	51	C	GLY	4	-0.636	6.488	-3.558	1.00	0.00
ATOM	52	O	GLY	4	0.272	6.072	-4.276	1.00	0.00
ATOM	53	H	GLY	4	-2.169	4.528	-3.249	1.00	0.00
ATOM	54	1HA	GLY	4	-2.035	6.637	-5.166	1.00	0.00
ATOM	55	2HA	GLY	4	-2.666	7.280	-3.648	1.00	0.00
ATOM	56	N	PRO	5	-0.410	6.939	-2.314	1.00	0.00
ATOM	57	CA	PRO	5	0.917	7.007	-1.721	1.00	0.00
ATOM	58	C	PRO	5	1.505	5.610	-1.509	1.00	0.00
ATOM	59	O	PRO	5	2.698	5.413	-1.731	1.00	0.00
ATOM	60	CB	PRO	5	0.739	7.773	-0.407	1.00	0.00
ATOM	61	CG	PRO	5	-0.715	7.501	-0.026	1.00	0.00
ATOM	62	CD	PRO	5	-1.419	7.411	-1.380	1.00	0.00
ATOM	63	HA	PRO	5	1.586	7.574	-2.370	1.00	0.00
ATOM	64	1HB	PRO	5	1.439	7.458	0.368	1.00	0.00
ATOM	65	2HB	PRO	5	0.857	8.840	-0.600	1.00	0.00
ATOM	66	1HG	PRO	5	-0.781	6.542	0.486	1.00	0.00
ATOM	67	2HG	PRO	5	-1.126	8.290	0.603	1.00	0.00
ATOM	68	1HD	PRO	5	-2.270	6.732	-1.322	1.00	0.00
ATOM	69	2HD	PRO	5	-1.752	8.405	-1.683	1.00	0.00
ATOM	70	N	LEU	6	0.670	4.646	-1.098	1.00	0.00
ATOM	71	CA	LEU	6	1.043	3.246	-0.926	1.00	0.00
ATOM	72	C	LEU	6	0.520	2.418	-2.110	1.00	0.00
ATOM	73	O	LEU	6	0.206	1.241	-1.944	1.00	0.00
ATOM	74	CB	LEU	6	0.497	2.733	0.422	1.00	0.00
ATOM	75	CG	LEU	6	1.343	3.106	1.651	1.00	0.00
ATOM	76	CD1	LEU	6	1.406	4.613	1.920	1.00	0.00
ATOM	77	CD2	LEU	6	0.748	2.411	2.883	1.00	0.00
ATOM	78	H	LEU	6	-0.298	4.883	-0.937	1.00	0.00
ATOM	79	HA	LEU	6	2.127	3.131	-0.913	1.00	0.00
ATOM	80	1HB	LEU	6	-0.520	3.095	0.556	1.00	0.00
ATOM	81	2HB	LEU	6	0.465	1.645	0.409	1.00	0.00
ATOM	82	HG	LEU	6	2.359	2.738	1.507	1.00	0.00
ATOM	83	2HD1	LEU	6	1.891	4.795	2.879	1.00	0.00
ATOM	84	3HD1	LEU	6	1.994	5.109	1.150	1.00	0.00
ATOM	85	1HD1	LEU	6	0.400	5.032	1.947	1.00	0.00
ATOM	86	2HD2	LEU	6	1.368	2.612	3.757	1.00	0.00
ATOM	87	3HD2	LEU	6	-0.261	2.780	3.069	1.00	0.00
ATOM	88	1HD2	LEU	6	0.707	1.332	2.728	1.00	0.00
ATOM	89	N	GLN	7	0.441	3.001	-3.316	1.00	0.00
ATOM	90	CA	GLN	7	0.009	2.262	-4.497	1.00	0.00
ATOM	91	C	GLN	7	1.046	1.194	-4.848	1.00	0.00
ATOM	92	O	GLN	7	0.696	0.030	-5.022	1.00	0.00
ATOM	93	CB	GLN	7	-0.227	3.214	-5.677	1.00	0.00
ATOM	94	CG	GLN	7	-0.648	2.494	-6.968	1.00	0.00
ATOM	95	CD	GLN	7	-1.990	1.773	-6.841	1.00	0.00
ATOM	96	OE1	GLN	7	-3.015	2.295	-7.270	1.00	0.00
ATOM	97	NE2	GLN	7	-1.991	0.566	-6.271	1.00	0.00
ATOM	98	H	GLN	7	0.702	3.971	-3.428	1.00	0.00
ATOM	99	HA	GLN	7	-0.936	1.778	-4.245	1.00	0.00
ATOM	100	1HB	GLN	7	-1.007	3.925	-5.415	1.00	0.00
ATOM	101	2HB	GLN	7	0.690	3.769	-5.878	1.00	0.00
ATOM	102	1HG	GLN	7	-0.742	3.247	-7.752	1.00	0.00
ATOM	103	2HG	GLN	7	0.120	1.787	-7.282	1.00	0.00
ATOM	104	1HE2	GLN	7	-2.855	0.049	-6.189	1.00	0.00
ATOM	105	2HE2	GLN	7	-1.129	0.171	-5.919	1.00	0.00

-continued

ATOM	106	N	TRP	8	2.319	1.596	-4.940	1.00	0.00
ATOM	107	CA	TRP	8	3.449	0.700	-5.154	1.00	0.00
ATOM	108	C	TRP	8	3.434	-0.449	-4.141	1.00	0.00
ATOM	109	O	TRP	8	3.652	-1.602	-4.505	1.00	0.00
ATOM	110	CB	TRP	8	4.756	1.500	-5.048	1.00	0.00
ATOM	111	CG	TRP	8	5.026	2.107	-3.702	1.00	0.00
ATOM	112	CD1	TRP	8	4.543	3.290	-3.263	1.00	0.00
ATOM	113	CD2	TRP	8	5.753	1.533	-2.573	1.00	0.00
ATOM	114	NE1	TRP	8	4.896	3.483	-1.944	1.00	0.00
ATOM	115	CE2	TRP	8	5.643	2.425	-1.466	1.00	0.00
ATOM	116	CE3	TRP	8	6.450	0.324	-2.354	1.00	0.00
ATOM	117	CZ2	TRP	8	6.203	2.134	-0.213	1.00	0.00
ATOM	118	CZ3	TRP	8	7.061	0.049	-1.117	1.00	0.00
ATOM	119	CH2	TRP	8	6.933	0.948	-0.045	1.00	0.00
ATOM	120	H	TRP	8	2.521	2.575	-4.803	1.00	0.00
ATOM	121	HA	TRP	8	3.378	0.286	-6.161	1.00	0.00
ATOM	122	1HB	TRP	8	5.585	0.835	-5.293	1.00	0.00
ATOM	123	2HB	TRP	8	4.739	2.297	-5.793	1.00	0.00
ATOM	124	HD1	TRP	8	3.930	3.966	-3.841	1.00	0.00
ATOM	125	HE1	TRP	8	4.624	4.279	-1.384	1.00	0.00
ATOM	126	HE3	TRP	8	6.504	-0.411	-3.142	1.00	0.00
ATOM	127	HZ2	TRP	8	6.053	2.799	0.624	1.00	0.00
ATOM	128	HZ3	TRP	8	7.623	-0.862	-0.986	1.00	0.00
ATOM	129	HH2	TRP	8	7.382	0.719	0.911	1.00	0.00
ATOM	130	N	LEU	9	3.164	-0.117	-2.872	1.00	0.00
ATOM	131	CA	LEU	9	3.155	-1.044	-1.753	1.00	0.00
ATOM	132	C	LEU	9	2.024	-2.054	-1.956	1.00	0.00
ATOM	133	O	LEU	9	2.239	-3.260	-1.862	1.00	0.00
ATOM	134	CB	LEU	9	3.025	-0.233	-0.448	1.00	0.00
ATOM	135	CG	LEU	9	3.830	-0.813	0.726	1.00	0.00
ATOM	136	CD1	LEU	9	3.898	0.197	1.877	1.00	0.00
ATOM	137	CD2	LEU	9	3.197	-2.091	1.264	1.00	0.00
ATOM	138	H	LEU	9	2.975	0.853	-2.669	1.00	0.00
ATOM	139	HA	LEU	9	4.112	-1.567	-1.754	1.00	0.00
ATOM	140	1HB	LEU	9	3.418	0.768	-0.625	1.00	0.00
ATOM	141	2HB	LEU	9	1.977	-0.133	-0.166	1.00	0.00
ATOM	142	HG	LEU	9	4.846	-1.031	0.399	1.00	0.00
ATOM	143	3HD1	LEU	9	4.523	-0.200	2.678	1.00	0.00
ATOM	144	1HD1	LEU	9	4.326	1.137	1.535	1.00	0.00
ATOM	145	2HD1	LEU	9	2.899	0.384	2.267	1.00	0.00
ATOM	146	2HD2	LEU	9	3.808	-2.482	2.077	1.00	0.00
ATOM	147	3HD2	LEU	9	2.197	-1.867	1.630	1.00	0.00
ATOM	148	1HD2	LEU	9	3.135	-2.835	0.479	1.00	0.00
ATOM	149	N	CYS	10	0.830	-1.551	-2.288	1.00	0.00
ATOM	150	CA	CYS	10	-0.347	-2.351	-2.581	1.00	0.00
ATOM	151	C	CYS	10	-0.090	-3.354	-3.706	1.00	0.00
ATOM	152	O	CYS	10	-0.370	-4.536	-3.533	1.00	0.00
ATOM	153	CB	CYS	10	-1.528	-1.430	-2.894	1.00	0.00
ATOM	154	SG	CYS	10	-3.024	-2.270	-3.462	1.00	0.00
ATOM	155	H	CYS	10	0.733	-0.547	-2.354	1.00	0.00
ATOM	156	HA	CYS	10	-0.598	-2.913	-1.684	1.00	0.00
ATOM	157	1HB	CYS	10	-1.764	-0.865	-1.995	1.00	0.00
ATOM	158	2HB	CYS	10	-1.251	-0.720	-3.670	1.00	0.00
ATOM	159	1LG	CYS	10	-2.931	-2.146	-4.124	1.00	0.00
ATOM	160	2LG	CYS	10	-3.134	-2.620	-2.887	1.00	0.00
ATOM	161	N	GLU	11	0.449	-2.907	-4.846	1.00	0.00
ATOM	162	CA	GLU	11	0.729	-3.800	-5.964	1.00	0.00
ATOM	163	C	GLU	11	1.792	-4.838	-5.593	1.00	0.00
ATOM	164	O	GLU	11	1.609	-6.024	-5.859	1.00	0.00
ATOM	165	CB	GLU	11	1.168	-3.009	-7.206	1.00	0.00
ATOM	166	CG	GLU	11	0.102	-2.037	-7.727	1.00	0.00
ATOM	167	CD	GLU	11	-1.243	-2.716	-7.973	1.00	0.00
ATOM	168	OE1	GLU	11	-1.333	-3.451	-8.980	1.00	0.00
ATOM	169	OE2	GLU	11	-2.154	-2.493	-7.146	1.00	0.00
ATOM	170	H	GLU	11	0.669	-1.924	-4.942	1.00	0.00
ATOM	171	HA	GLU	11	-0.182	-4.348	-6.204	1.00	0.00
ATOM	172	1HB	GLU	11	2.075	-2.446	-6.979	1.00	0.00
ATOM	173	2HB	GLU	11	1.396	-3.721	-8.001	1.00	0.00
ATOM	174	1HG	GLU	11	-0.034	-1.231	-7.014	1.00	0.00
ATOM	175	2HG	GLU	11	0.452	-1.601	-8.663	1.00	0.00
ATOM	176	N	LYS	12	2.909	-4.388	-5.010	1.00	0.00
ATOM	177	CA	LYS	12	4.080	-5.221	-4.779	1.00	0.00
ATOM	178	C	LYS	12	3.824	-6.277	-3.701	1.00	0.00
ATOM	179	O	LYS	12	4.069	-7.459	-3.936	1.00	0.00
ATOM	180	CB	LYS	12	5.280	-4.325	-4.444	1.00	0.00
ATOM	181	CG	LYS	12	6.590	-5.123	-4.400	1.00	0.00
ATOM	182	CD	LYS	12	7.820	-4.206	-4.385	1.00	0.00
ATOM	183	CE	LYS	12	7.961	-3.450	-3.061	1.00	0.00
ATOM	184	NZ	LYS	12	9.120	-2.542	-3.084	1.00	0.00

-continued

ATOM	185	H	LYS	12	2.989	-3.401	-4.803	1.00	0.00
ATOM	186	HA	LYS	12	4.306	-5.736	-5.715	1.00	0.00
ATOM	187	1HB	LYS	12	5.366	-3.568	-5.225	1.00	0.00
ATOM	188	2HB	LYS	12	5.107	-3.827	-3.490	1.00	0.00
ATOM	189	1HG	LYS	12	6.604	-5.776	-3.526	1.00	0.00
ATOM	190	2HG	LYS	12	6.650	-5.743	-5.296	1.00	0.00
ATOM	191	1HD	LYS	12	8.711	-4.820	-4.529	1.00	0.00
ATOM	192	2HD	LYS	12	7.754	-3.496	-5.210	1.00	0.00
ATOM	193	1HE	LYS	12	7.068	-2.857	-2.869	1.00	0.00
ATOM	194	2HE	LYS	12	8.098	-4.170	-2.255	1.00	0.00
ATOM	195	2HZ	LYS	12	9.000	-1.861	-3.820	1.00	0.00
ATOM	196	3HZ	LYS	12	9.192	-2.066	-2.196	1.00	0.00
ATOM	197	1HZ	LYS	12	9.962	-3.073	-3.250	1.00	0.00
ATOM	198	N	TYR	13	3.345	-5.856	-2.524	1.00	0.00
ATOM	199	CA	TYR	13	3.118	-6.740	-1.388	1.00	0.00
ATOM	200	C	TYR	13	1.704	-7.312	-1.433	1.00	0.00
ATOM	201	O	TYR	13	1.537	-8.530	-1.445	1.00	0.00
ATOM	202	CB	TYR	13	3.347	-5.993	-0.069	1.00	0.00
ATOM	203	CG	TYR	13	4.787	-5.600	0.198	1.00	0.00
ATOM	204	CD1	TYR	13	5.300	-4.391	-0.305	1.00	0.00
ATOM	205	CD2	TYR	13	5.596	-6.412	1.016	1.00	0.00
ATOM	206	CE1	TYR	13	6.582	-3.958	0.070	1.00	0.00
ATOM	207	CE2	TYR	13	6.896	-6.000	1.354	1.00	0.00
ATOM	208	CZ	TYR	13	7.385	-4.766	0.892	1.00	0.00
ATOM	209	OH	TYR	13	8.639	-4.357	1.241	1.00	0.00
ATOM	210	H	TYR	13	3.140	-4.873	-2.402	1.00	0.00
ATOM	211	HA	TYR	13	3.826	-7.571	-1.417	1.00	0.00
ATOM	212	1HB	TYR	13	2.720	-5.102	-0.042	1.00	0.00
ATOM	213	2HB	TYR	13	3.018	-6.640	0.745	1.00	0.00
ATOM	214	HD1	TYR	13	4.707	-3.781	-0.968	1.00	0.00
ATOM	215	HD2	TYR	13	5.218	-7.350	1.396	1.00	0.00
ATOM	216	HE1	TYR	13	6.948	-3.005	-0.278	1.00	0.00
ATOM	217	HE2	TYR	13	7.514	-6.626	1.982	1.00	0.00
ATOM	218	HH	TYR	13	8.872	-3.497	0.883	1.00	0.00
ATOM	219	N	PHE	14	0.685	-6.443	-1.441	1.00	0.00
ATOM	220	CA	PHE	14	-0.707	-6.851	-1.269	1.00	0.00
ATOM	221	C	PHE	14	-1.342	-7.174	-2.628	1.00	0.00
ATOM	222	O	PHE	14	-2.470	-6.768	-2.906	1.00	0.00
ATOM	223	CB	PHE	14	-1.488	-5.752	-0.525	1.00	0.00
ATOM	224	CG	PHE	14	-0.808	-5.116	0.680	1.00	0.00
ATOM	225	CD1	PHE	14	0.044	-5.862	1.519	1.00	0.00
ATOM	226	CD2	PHE	14	-1.039	-3.756	0.964	1.00	0.00
ATOM	227	CE1	PHE	14	0.722	-5.231	2.578	1.00	0.00
ATOM	228	CE2	PHE	14	-0.370	-3.129	2.027	1.00	0.00
ATOM	229	CZ	PHE	14	0.523	-3.862	2.826	1.00	0.00
ATOM	230	H	PHE	14	0.882	-5.453	-1.475	1.00	0.00
ATOM	231	HA	PHE	14	-0.751	-7.755	-0.659	1.00	0.00
ATOM	232	1HB	PHE	14	-1.708	-4.956	-1.237	1.00	0.00
ATOM	233	2HB	PHE	14	-2.441	-6.169	-0.195	1.00	0.00
ATOM	234	HD1	PHE	14	0.196	-6.918	1.352	1.00	0.00
ATOM	235	HD2	PHE	14	-1.733	-3.185	0.367	1.00	0.00
ATOM	236	HE1	PHE	14	1.396	-5.800	3.201	1.00	0.00
ATOM	237	HE2	PHE	14	-0.539	-2.080	2.228	1.00	0.00
ATOM	238	HZ	PHE	14	1.053	-3.370	3.629	1.00	0.00
ATOM	239	N	GLY	15	-0.607	-7.896	-3.480	1.00	0.00
ATOM	240	CA	GLY	15	-0.995	-8.181	-4.849	1.00	0.00
ATOM	241	C	GLY	15	0.182	-8.794	-5.604	1.00	0.00
ATOM	242	O	GLY	15	1.159	-9.233	-4.999	1.00	0.00
ATOM	243	H	GLY	15	0.302	-8.231	-3.186	1.00	0.00
ATOM	244	1HA	GLY	15	-1.832	-8.881	-4.853	1.00	0.00
ATOM	245	2HA	GLY	15	-1.295	-7.254	-5.340	1.00	0.00
ATOM	246	N	NH2	16	0.095	-8.822	-6.935	1.00	0.00
ATOM	247	1HN	NH2	16	-0.720	-8.445	-7.397	1.00	0.00
ATOM	248	2HN	NH2	16	0.854	-9.209	-7.476	1.00	0.00

Example 9

Displacement of IGF-I from IGFBPs using (Leu²⁴, Ala³¹)hIGF

This example shows that the IGF mutant (Leu²⁴,Ala³¹)hIGF used in the above animal studies can displace IGF-I tracer from IGFBPs present in human serum.

To determine if the IGF mutant could displace IGF-I from endogenous IGFBPs present in human serum, the following experiment was performed. Radiolabeled rhIGF-I (¹²⁵I-IGF-

I) was incubated with normal human serum samples at a concentration of 33 ng/ml for 16 hr at 4° C. to allow for the formation of ¹²⁵I-IGF-I:IGFBP complexes. Following complex formation, the IGF mutant (Leu²⁴,Ala³¹)hIGF (40 μg/ml) was added to the incubations for 0, 0.5, 3, or 16 hr. The samples (n=3) were then analyzed by FPLC size-exclusion chromatography (FPLC-SEC) to determine the relative size and amounts of the ¹²⁵I-IGF-I:IGFBP complexes.

FIG. 41 is a chromatogram demonstrating the ability of the IGF mutant to displace ¹²⁵I-IGF-I from endogenous

IGFBPs present in serum from normal humans. Data are expressed as cpm per fraction (n=1). This experiment was repeated three times, was quantitated, and is presented in Table IX.

TABLE IX

Quantitation of FPLC-SEC Data Demonstrating Ability of IGF Mutant to Displace ^{125}I -IGF-I From Endogenous IGFBPs in Normal Human Serum*			
	~150 kDa	~44 kDa	Free (~7 kDa)
IGF Mutant (0 hr)	30.3 ± 1.4	60.5 ± 0.8	9.3 ± 2.3
IGF Mutant (0.5 hr)	24.4 ± 0.8	37.3 ± 2.9	38.2 ± 3.7
IGF Mutant (3 hr)	10.3 ± 1.1	27.8 ± 1.5	61.9 ± 2.6
IGF Mutant (24 hr)	5.5 ± 0.1	12.6 ± 0.4	81.9 ± 0.5

*Data (mean ± SD, n = 3) are expressed as percent of total radioactivity, calculated by dividing the amount of radioactivity in each molecular weight range by total radioactivity in all three molecular weight ranges.

After incubation of normal human serum with ^{125}I -IGF-I for 16 hr, three peaks of radioactivity were observed following FPLC-SEC (FIG. 42). These peaks likely correspond to the following 125I-IGF-I complexes: ~150 kDa (125I-IGF-I:IGFBP-3:ALS); ~44 kDa (125I-IGF-I:IGFBPs 1-4); ~7 kDa (free ^{125}I -IGF-I). As can be seen in FIG. 42, addition of the IGF mutant to the serum resulted in a time-dependent decrease in ^{125}I -IGF-I associated with the ^{125}I -IGF-I:IGFBP complexes and an increase in the amount of free ^{125}I -IGF-I. These results are outlined in the above table. The ^{125}I -IGF-I was more readily displaced from the ~44 kDa than the ~150 kDa ^{125}I -IGF-I:IGFBP complex, suggesting that IGF-I bound to the lower molecular weight IGFBPs is in a more bioavailable form. These data clearly indicate that the IGF mutant has the ability to displace IGF-I from endogenous IGFBPs present in normal human serum and therefore is likely to be active in vivo in humans.

In conclusion, this evidence shows that one would expect the molecules of the nature of the mutant to demonstrate the activities in humans that have been shown in rats.

Example 10

Displacement of IGF-I from IGFBPs using BP15

This Example tests an IGFBP-3-specific peptide, BP15, for its ability to block the binding of ^{125}I -IGF-I in human serum. Human serum was incubated with ^{125}I -IGF-I ± the peptide and the amount of tracer bound to IGFBPs via size-exclusion chromatography was measured. Addition of the peptide resulted in an approximately 42% decrease in ^{125}I -IGF-I associated with the 150-KD IGF/IGFBP-3/ALS complex and a 59% increase in the amount of free ^{125}I -IGF-I. The peptide did not decrease ^{125}I -IGF-I binding to the 44-KD IGFBPs (in fact, it slightly increased it), indicating that the peptide only competes with IGF-I for binding to IGFBP-3.

These results indicate that the analog (at 0.2 mM) can compete with IGF-I for binding to IGFBP-3 in human serum.

Example 11

Treatment of Humans with Human IGF-I

This example shows the principle of how an exogenously administered compound that binds to one or more of the

IGFBPs acts to displace endogenous IGFs and how to dose an IGF agonist for use in humans. In this study human Type II diabetics were administered recombinant human IGF-I or placebo by twice-daily injection at four doses (10, 20, 40, or 80 $\mu\text{g}/\text{kg}$) for 12 weeks. Blood samples were drawn, before, every two weeks during, and after (EP) the 12 weeks of treatment. The concentrations of IGF-I, IGF-II, and IGFBP-3 were measured in all the samples, with the exception of IGF-II not being measured in the samples taken from the patients treated with 10 $\mu\text{g}/\text{day}$ of IGF-I.

FIG. 43 shows the concentrations of IGF-I in the blood of the patients. The unexpected finding was the "plateau" effect of administering 40 and 80 μg of IGF-I; the same total blood concentration of IGF-I was reached with these two doses. FIG. 44 shows the concentrations of IGF-II in the blood of the patients. In contrast to the rising levels of IGF-I, the levels of IGF-II fell in almost a mirror-image pattern to the rise in IGF-I concentrations. As with the plateauing of the rising IGF-I concentrations, the falling IGF-II concentrations also reached a plateau.

FIG. 45 shows the concentrations of IGFBP-3 in the blood of the patients. In contrast to the clear changes in the patterns of IGF-I and IGF-II in the blood, the concentrations of IGFBP-3 showed no statistically significant or clear pattern of change.

Inspection of FIGS. 43 and 44 reveals that the total IGF concentrations (IGF-I plus IGF-II) showed little change with treatment. This was because the rise in the concentrations of IGF-I closely matched the fall in the concentrations of IGF-II. Inspection of all three Figures shows that the dose-related changes in the concentrations of IGF-I and IGF-II in the blood of the patients were not accompanied by a reduced IGFBP-3 binding protein capacity (IGFBP-3 is the major binding protein in blood).

The obvious explanation for the fall in the concentration of IGF-II, and the plateauing of IGF-I and IGF-II concentrations, is that there is a finite amount of IGF binding protein capacity, and in this experiment the doses of IGF-I used caused a dose-related displacement of IGF-II from the binding proteins.

It is a logical extension of the observations in this Example to expect that any molecule with the ability to enhance levels of active IGF would show activities similar to those shown for IGF-I in this Example. In addition, from the doses of IGF-I used and the concentrations of IGFBP and IGF-I and IGF-II demonstrated, it is simple to calculate how much of an IGF agonist should be given to increase levels of active endogenous IGF. The molar size relative to IGF-I, the affinity of the IGF agonist for the IGFBP, and its bioavailability would be other variables taken into account to arrive at doses that increased active IGF in a human.

The present invention has of necessity been discussed herein by reference to certain specific methods and materials. It is to be understood that the discussion of these specific methods and materials in no way constitutes any limitation on the scope of the present invention, which extends to any and all alternative materials and methods suitable for accomplishing the objectives of the present invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 79

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Glu Leu Asp Gly Trp Val Cys Ile Lys Val Gly Glu Gln Asn Leu
 1 5 10 15

Cys Tyr Leu Ala Glu Gly
 20 21

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Trp Phe Lys Thr Val Cys Tyr Glu Trp Glu Asp Glu Val Gln Cys
 1 5 10 15

Tyr Thr Leu Glu Glu Gly
 20 21

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Arg Val Gly Ala Tyr Ile Ser Cys Ser Glu Thr Glu Cys Trp Val
 1 5 10 15

Glu Asp Leu Leu Asp Gly
 20 21

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Val Ala Trp Glu Val Cys Trp Asp Arg His Asp Gln Gly Tyr Ile
 1 5 10 15

Cys Thr Thr Asp Ser
 20

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Ala Trp Glu Val Cys Trp Asp Arg His Gln Gly Tyr Ile Cys Thr
 1 5 10 15

Thr Asp Ser
 18

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Cys Trp Asp Arg His Asp Gln Gly Tyr Ile Cys Thr Thr Asp Ser
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Glu Glu Ser Glu Cys Phe Glu Gly Pro Gly Tyr Val Ile Cys Gly
 1 5 10 15

Leu Val Gly
 18

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Asp Met Gly Val Cys Ala Asp Gly Pro Trp Met Tyr Val Cys Glu
 1 5 10 15

Trp Thr Glu
 18

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ser Glu Glu Val Cys Trp Pro Val Ala Glu Trp Tyr Leu Cys Asn
 1 5 10 15

Met Trp Gly
 18

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

-continued

Ser	Glu	Glu	Val	Cys	Trp	Pro	Val	Ala	Glu	Trp	Tyr	Leu	Cys	Asn
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Val	Cys	Trp	Pro	Val	Ala	Glu	Trp	Tyr	Leu	Cys	Asn	Met	Trp	Gly
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 12 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Val	Cys	Trp	Pro	Val	Ala	Glu	Trp	Tyr	Leu	Cys	Asn
1				5					10		12

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Thr	Gly	Val	Asp	Cys	Gln	Cys	Gly	Pro	Val	His	Cys	Val	Cys	Met
1				5					10					15

Asp	Trp	Ala
		18

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Thr	Val	Ala	Asn	Cys	Asp	Cys	Tyr	Met	Pro	Leu	Cys	Leu	Cys	Tyr
1				5					10					15

Asp	Ser	Asp
		18

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Cys	Arg	Ala	Gly	Pro	Leu	Gln	Trp	Leu	Cys	Glu	Lys	Tyr	Phe	Gly
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids

-continued

(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ser Glu Val Gly Cys Arg Ala Gly Pro Leu Gln Trp Leu Cys Glu
1 5 10 15
Lys Tyr Phe Gly
19

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 400 base pairs
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TCACGTAAAA AGGGTATCTA GAATTATGAT GATTACTCTG CGCAAATTC	50
CTCTGGCGGT TGCCGTCGCA GCGGGCGTAA TGTCTGCTCA GGCCATGGCC	100
GGTCCCGAAA CTCTGTGCGG TGCTGAACTG GTTGACGCTC TGCAGTTCGT	150
ATGTGGTGAT CGAGGCTTCC TGTTCACAAA ACCGACTGGG GCTGGATCCT	200
CCTCTCGTCG TGCTCCCCAG ACTGGTATTG TTGACGAATG CTGCTTTCGT	250
TCTTGCGACC TGCGTCGTCT GAAAATGTAT TGCCTCCCC TGAAACCCGC	300
TAAATCTGCT TAGAAGCTCC TAACGCTCGG TTGCCGCCGG GCGTTTTTTA	350
TTGTTAACTC ATGTTTGACA GCTTATCATC GATAAGCTTT AATGCGGTAG	400

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 95 amino acids
(B) TYPE: Amino Acid
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Met Met Ile Thr Leu Arg Lys Leu Pro Leu Ala Val Ala Val Ala
1 5 10 15
Ala Gly Val Met Ser Ala Gln Ala Met Ala Gly Pro Glu Thr Leu
20 25 30
Cys Gly Ala Glu Leu Val Asp Ala Leu Gln Phe Val Cys Gly Asp
35 40 45
Arg Gly Phe Leu Phe Asn Lys Pro Thr Gly Ala Gly Ser Ser Ser
50 55 60
Arg Arg Ala Pro Gln Thr Gly Ile Val Asp Glu Cys Cys Phe Arg
65 70 75
Ser Cys Asp Leu Arg Arg Leu Glu Met Tyr Cys Ala Pro Leu Lys
80 85 90
Pro Ala Lys Ser Ala
95

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5115 base pairs
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Double
(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

-continued

GAATTCAACT TCTCCATACT TTGGATAAGG AAATACAGAC ATGAAAAATC	50
TCATTGCTGA GTTGTATTAT AAGCTTGCCC AAAAAGAAGA AGAGTCGAAT	100
GAACTGTGTG CGCAGGTAGA AGCTTTGGAG ATTATCGTCA CTGCAATGCT	150
TCGCAATATG GCGCAAAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG	200
GGGCGCTGTA CGAGGTAAAG CCCGATGCCA GCATTCTGTA CGACGATACG	250
GAGCTGTGTC GCGATTACGT AAAGAAGTTA TTGAAGCATC CTCGTCAGTA	300
AAAAGTTAAT CTTTTCAACA GCTGTCATAA AGTTGTCACG GCCGAGACTT	350
ATAGTCGCTT TGTTTTTATT TTTTAATGTA TTTGTAAC TAACGCAAGT	400
TCACGTAAAA AGGGTATCTA GAATATATGAT GATTACTCTG CGCAAAC TTC	450
CTCTGGCGGT TGCCGTCGCA GCGGGCGTAA TGTCTGCTCA GGCCATGGCC	500
GGTCCCGAAA CTCTGTGCGG TGCTGAACTG GTTGACGCTC TGCAGTTCGT	550
ATGTGGTGAT CGAGGCTTCC TGTTCACAAA ACCGACTGGG GCTGGATCCT	600
CCTCTCGTCG TGCTCCCCAG ACTGGTATTG TTGACGAATG CTGCTTTCGT	650
TCTTGCGACC TGCGTCTGCT GAAATGTAT TGCGCTCCCC TGAACCCGC	700
TAAATCTGCT TAGAAGCTCC TAACGCTCGG TTGCCGCCGG GCGTTTTTTA	750
TTGTAACTC ATGTTTGACA GCTTATCATC GATAAGCTTT AATGCGGTAG	800
TTTATCACAG TTAATTTGCT AACGAGTCA GGCACCGTGT ATGAAATCTA	850
ACAATGCGCT CATCGTCATC CTCGGCACCG TCACCCTGGA TGCTGTAGGC	900
ATAGGCTTGG TTATGCCGGT ACTGCCGGGC CTCTTGCGGG ATATCGTCCA	950
TTCCGACAGC ATCGCCAGTC ACTATGGCGT GCTGCTAGCG CTATATGCGT	1000
TGATGCAATT TCTATGCGCA CCCGTTCTCG GAGCACTGTC CGACCGCTTT	1050
GGCCGCCGCC CAGTCCTGCT CGCTTCGCTA CTTGGAGCCA CTATCGACTA	1100
CGCGATCATG GCGACCACAC CCGTCTCTGT GATCCTCTAC GCCGGACGCA	1150
TCGTGGCCGG CATCACCGGC GCCACAGGTG CGGTTGCTGG CGCCTATATC	1200
GCCGACATCA CCGATGGGGA AGATCGGGCT CGCCACTTTC GGCTCATGAG	1250
CGCTTGTTTC GCGTGGGTA TGGTGGCAGG CCCCGTGGCC GGGGACTGT	1300
TGGCGCCAT CTCCTTGCAT GCACCATTC TTGCGGCCGG GGTGCTCAAC	1350
GGCCTCAACC TACTACTGGG CTGCTTCCTA ATGCAGGAGT CGCATAAGGG	1400
AGAGCGTCGA CCGATGCCCT TGAGAGCCTT CAACCCAGTC AGCTCCTTCC	1450
GGTGGGCGCG GGGCATGACT ATCGTCGCCG CACTTATGAC TGTCTTCTTT	1500
ATCATGCAAC TCGTAGGACA GGTGCCGGCA GCGCTCTGGG TCATTTTCGG	1550
CGAGGACCGC TTTCGCTGGA GCGCGACGAT GATCGGCCTG TCGCTTGGCG	1600
TATTCCGAAT CTTGCACGCC CTCGCTCAAG CCTTCGTAC TGGTCCCGCC	1650
ACCAAACGTT TCGGCAGAA GCAGGCCATT ATCGCCGGCA TGGCGGCCGA	1700
CGCGCTGGG TACGTCTTGC TGGCGTTTCG GACGCGAGG TGGATGGCCT	1750
TCCCCATTAT GATTCTTCTC GCTTCCGGCG GCATCGGGAT GCCCGCGTTG	1800
CAGGCCATGC TGTCCAGGCA GGTAGATGAC GACCATCAGG GACAGCTTCA	1850
AGGATCGCTC GCGGCTCTTA CCAGCCTAAC TTCGATCACT GGACCGCTGA	1900
TCGTCACGGC GATTTATGCC GCCTCGGCGA GCACATGGAA CGGGTTGGCA	1950

-continued

TGGATTGTAG	GCGCCGCCCT	ATACCTTGTC	TGCCTCCCCG	CGTTGCGTCG	2000
CGGTGCATGG	AGCCGGGCCA	CCTCGACCTG	AATGGAAGCC	GGCGGCACCT	2050
CGCTAACGGA	TTCACCACTC	CAAGAATTGG	AGCCAATCAA	TTCTTGCGGA	2100
GAAGTGTGAA	TGCGCAAACC	AACCCCTTGGC	AGAACATATC	CATCGCGTCC	2150
GCCATCTCCA	GCAGCCGCAC	GCGGCGCATC	TCGGGCAGCG	TTGGGTCCCTG	2200
GCCACGGGTG	CGCATGATCG	TGCTCCTGTC	GTTGAGGACC	CGGCTAGGCT	2250
GGCGGGTTG	CCTTACTGGT	TAGCAGAATG	AATCACCATG	ACGCGAGCGA	2300
ACGTGAAGCG	ACTGCTGCTG	CAAAACGTCT	GCGACCTGAG	CAACAACATG	2350
AATGGTCTTC	GGTTTCCGTG	TTTCGTAAAG	TCTGAAAACG	CGGAAGTCAG	2400
CGCCCTGCAC	CATTATGTTT	CGGATCTGCA	TCGCAGGATG	CTGCTGGCTA	2450
CCCTGTGGAA	CACCTACATC	TGTATTAACG	AAGCGCTGGC	ATTGACCCCTG	2500
AGTGATTTTT	CTCTGGTCCC	GCCGCATCCA	TACCGCCAGT	TGTTTACCCT	2550
CACAACGTTT	CAGTAACCGG	GCATGTTTAT	CATCAGTAAC	CCGTATCGTG	2600
AGCATCTCTT	CTCGTTTCAT	CGGTATCATT	ACCCCCATGA	ACAGAAATTC	2650
CCCCTTACAC	GGAGGCATCA	AGTGACCAAA	CAGGAAAAAA	CCGCCCTTAA	2700
CATGGCCCGC	TTTATCAGAA	GCCAGACATT	AACGCTTCTG	GAGAAACTCA	2750
ACGAGCTGGA	CGCGGATGAA	CAGGCAGACA	TCTGTGAATC	GCTTCACGAC	2800
CACGCTGATG	AGCTTTACCG	CAGCTGCCTC	GCGCGTTTCG	GTGATGACGG	2850
TGAAAACCTC	TGACACATGC	AGTCCCCTGA	GACGGTCACA	GCTTGTCTGT	2900
AAGCGGATGC	CGGGAGCAGA	CAAGCCCCTC	AGGGCGCGTC	AGCGGGTGT	2950
GGCGGGTGTG	GGGGCGCAGC	CATGACCCAG	TCACGTAGCG	ATAGCGGAGT	3000
GTATACTGGC	TTAACTATGC	GGCATCAGAG	CAGATTGTAC	TGAGAGTGCA	3050
CCATATGCGG	TGTGAAATAC	CGCACAGATG	CGTAAGGAGA	AAATACCGCA	3100
TCAGGCGCTC	TTCCGCTTCC	TCGCTCACTG	ACTCGCTGCG	CTCGGTGCTT	3150
CGGCTGCGGC	GAGCGGTATC	AGCTCACTCA	AAGCGGTAA	TACGGTTATC	3200
CACAGAATCA	GGGGATAACG	CAGGAAAGAA	CATGTGAGCA	AAAGGCCAGC	3250
AAAAGGCCAG	GAACCGTAAA	AAGGCCGCGT	TGCTGGCGTT	TTTCCATAGG	3300
CTCCGCCCCC	CTGACGAGCA	TCACAAAAAT	CGACGCTCAA	GTCAGAGGTG	3350
GCGAAACCCG	ACAGGACTAT	AAAGATACCA	GGCGTTTCCC	CCTGGAAGCT	3400
CCCTCGTGCG	CTCTCCTGTT	CCGACCCCTG	CGCTTACCGG	ATACCTGTCC	3450
GCCTTTCTCC	CTTCGGGAAG	CGTGGCGCTT	TCTCATAGCT	CACGCTGTAG	3500
GTATCTCAGT	TCGGTGTAGG	TCGTTGCTC	CAAGCTGGGC	TGTGTGCACG	3550
AACCCCCCGT	TCAGCCCGAC	CGCTGCGCCT	TATCCGGTAA	CTATCGTCTT	3600
GAGTCCAACC	CGGTAAGACA	CGACTTATCG	CCACTGGCAG	CAGCCACTGG	3650
TAACAGGATT	AGCAGAGCGA	GGTATGTAGG	CGGTGCTACA	GAGTTCTTGA	3700
AGTGGTGGCC	TAACCTACGGC	TACTACTAGAA	GGACAGTATT	TGGTATCTGC	3750
GCTCTGTGTA	AGCCAGTTAC	CTTCGGAAAA	AGAGTTGGTA	GCTCTTGATC	3800
CGGCAAACAA	ACCACCGCTG	GTAGCGGTGG	TTTTTTTTGTT	TGCAAGCAGC	3850
AGATTACGCG	CAGAAAAAAA	GGATCTCAAG	AAGATCCTTT	GATCTTTTCT	3900
ACGGGGTCTG	ACGCTCAGTG	GAACGAAAAA	TCACGTTAAG	GGATTTGGT	3950

-continued

CATGAGATTA TCAAAAAGGA TCTTCACCTA GATCCTTTTA AATTA AAAAT	4000
GAAGTTTTAA ATCAATCTAA AGTATATATG AGTAAACTTG GTCTGACAGT	4050
TACCAATGCT TAATCAGTGA GGCACCTATC TCAGCGATCT GTCTATTTTCG	4100
TTCATCCATA GTTGCCTGAC TCCCCGTCGT GTAGATAACT ACGATACGGG	4150
AGGGCTTACC ATCTGGCCCC AGTGCTGCAA TGATACCGCG AGACCCACGC	4200
TCACCGGCTC CAGATTTATC AGCAATAAAC CAGCCAGCCG GAAGGGCCGA	4250
GCGCAGAAGT GGTCCCTGCAA CTTTATCCGC CTCCATCCAG TCTATTAATT	4300
GTTGCCGGGA AGCTAGAGTA AGTAGTTTCG CAGTTAATAG TTTGCGCAAC	4350
GTTGTTGCCA TTGCTGCAGG CATCGTGGTG TCACGCTCGT CGTTTGGTAT	4400
GGCTTCATTC AGCTCCGGTT CCCAACGATC AAGGCGAGTT ACATGATCCC	4450
CCATGTTGTG CAAAAAGCG GTTAGCTCCT TCGGTCCTCC GATCGTTGTC	4500
AGAAGTAAGT TGGCCGCGT GTTATCACTC ATGGTTATGG CAGCACTGCA	4550
TAATTCTCTT ACTGTCATGC CATCCGTAAG ATGCTTTTCT GTGACTGGTG	4600
AGTACTCAAC CAAGTCATTC TGAGAATAGT GTATGCGGCG ACCGAGTTGC	4650
TCTTGCCCGG CGTCAACACG GGATAATACC GCGCCACATA GCAGAACTTT	4700
AAAAGTGCTC ATCATTGGAA AACGTTCTTC GGGGCGAAAA CTCTCAAGGA	4750
TCTTACCGCT GTTGTAGATCC AGTTCGATGT AACCCACTCG TGCACCCAAC	4800
TGATCTTCAG CATCTTTTAC TTTACCAGC GTTCTGGGT GAGCAAAAAC	4850
AGGAAGGCAA AATGCCGCAA AAAAGGGAAT AAGGCGGACA CGGAAATGTT	4900
GAATACTCAT ACTCTTCCTT TTTCAATATT ATTGAAGCAT TTATCAGGGT	4950
TATTGTCTCA TGAGCGGATA CATATTTGAA TGTATTAGA AAAATAAACA	5000
AATAGGGGTT CCGCGCACAT TTCCCGAAA AGTGCCACCT GACGTCTAAG	5050
AAACCATTAT TATCATGACA TTAACCTATA AAAATAGGCG TATCACGAGG	5100
CCCTTTCGTC TTCAA	5115

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5140 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GAATTCAACT TCCTCACTT TTGGATAAGG AAATACAGAC ATGAAAAATC	50
TCATTGCTGA GTTGTATTTT AAGCTTGCCC AAAAAGAAGA AGAGTCGAAT	100
GAACTGTGTG CGCAGGTAGA AGCTTTGGAG ATTATCGTCA CTGCAATGCT	150
TCGCAATATG GCGCAAAATG ACCAACAGCG GTTGATTGAT CAGGTAGAGG	200
GGGCGCTGTA CGAGGTAAAG CCCGATGCCA GCATTCCTGA CGACGATACG	250
GAGCTGCTGC GCGATTACGT AAAGAAGTTA TTGAAGCATC CTCGTCAGTA	300
AAAAGTTAAT CTTTTCAACA GCTGTCATAA AGTGTGCACG GCCGAGACTT	350
ATAGTCGCTT TTTTTTATT TTTTAATGTA TTTGTAAC TAACGCAAGT	400
TCACGTAAAA AGGGTATCTA GAGGTTGAGG TGATTTTATG AAAAAGAATA	450
TCGCATTTCT TCTTGCACT ATGTTCTGTT TTTCTATTGC TACAAATGCC	500

-continued

TATGCATCTG	GTACCGCCAT	GGCTGATCCG	AACCGTTTCC	GCGGTAAAGA	550
TCTGGCAGGT	TCACCAGGTG	GAGGATCCGG	AGGAGGCGCC	GAGGGTGACG	600
ATCCCGCAA	AGCGGCCTTT	AACTCCCTGC	AAGCCTCAGC	GACCGAATAT	650
ATCGGTTATG	CGTGGGCGAT	GGTTGTTGTC	ATTGTCGGCG	CAACTATCGG	700
TATCAAGCTG	TTTAAGAAAT	TCACCTCGAA	AGCAAGCTGA	TAAACCGATA	750
CAATTAAGG	CTCCTTTTGG	AGCCTTTTTT	TTTGGAGATT	TTCAACGTGA	800
AAAAATTATT	ATTCGCAATT	CCTTTAGTTG	TTCCTTCTA	TTCTCACTCC	850
GCTGAAACTG	TTGAAAGTTG	TTTAGCAAAA	CCCATACAG	AAAATTCATT	900
TACTAACGTC	TGGAAAGACG	ACAAAACTTT	AGATCGTTAC	GCTAACTATG	950
AGGGTTGTCT	GTGGAATGCT	ACAGGCGTTG	TAGTTTGATC	TGGTGACGAA	1000
ACTCAGTGTC	TAGCTAGAGT	GGCGGTGGCT	CTGGTCCGG	TGATTTTGAT	1050
TATGAAAAGA	TGGCAAACGC	TAATAAGGGG	GCTATGACCG	AAAATGCCGA	1100
TGAAAACGCG	CTACAGTCTG	ACGCTAAAGG	CAAACCTGAT	TCTGTCGCTA	1150
CTGATTACGG	TGCTGCTATC	GATGGTTTCA	TTGGTGACGT	TTCCGGCCTT	1200
GCTAATGGTA	ATGGTGCTAC	TGGTGATTTT	GCTGGCTCTA	ATTCCCAAAT	1250
GGCTCAAGTC	GGTGACGGTG	ATAATTCACC	TTTAATGAAT	AATTTCCGTC	1300
AATATTTACC	TTCCCTCCCT	CAATCGGTTG	AATGTCGCC	TTTTGTCTTT	1350
AGCGCTGGTA	AACCATATGA	ATTTTCTATT	GATGTCGACA	AAATAAACTT	1400
ATTCGGTGGT	GTCTTTGCGT	TTCTTTTATA	TGTTGCCACC	TTTATGTATG	1450
TATTTTCTAC	GTTTGCTAAC	ATACTGCCTA	ATAAGGAGTC	TTAATCATGC	1500
CAGTCTTTT	GGCTAGCGCC	GCCCTATACC	TTGTCTGCCT	CCCCGCGTTG	1550
CGTCGCGGTG	CATGGAGCCG	GGCCACCTCG	ACCTGAATGG	AAGCCGGCGG	1600
CACCTCGCTA	ACGGATTAC	CACTCCAAGA	ATTGGAGCCA	ATCAATTCCTT	1650
GCGGAGAACT	GTGAATGCGC	AAACCAACCC	TTGGCAGAAC	ATATCCATCG	1700
CGTCCGCCAT	CTCCAGCAGC	CGCACGCGGC	GCATCTCGGG	CAGCGTTGGG	1750
TCCTGGCCAC	GGGTGCGCAT	GATCGTGCTC	CTGTCGTTGA	GGACCCGGCT	1800
AGGCTGGCGG	GGTTGCCTTA	CTGGTTAGCA	GAATGAATCA	CCGATACGCG	1850
AGCGAACGTG	AAGCGACTGC	TGCTGCAAAA	CGTCTGCGAC	CTGAGCAACA	1900
ACATGAATGG	TCITTCGGTTT	CCGTGTTTCC	TAAAGTCTGG	AAACGCGGAA	1950
GTCAGCGCCC	TGCACCATTA	TGTTCCGGAT	CTGCATCGCA	GGATGCTGCT	2000
GGCTACCCTG	TGGAACACCT	ACATCTGTAT	TAACGAAGCG	CTGGCATTGA	2050
CCCTGAGTGA	TTTTTCTCTG	GTCCCGCCG	ATCCATACCG	CCAGTTGTFT	2100
ACCCTCACAA	CGTTCCAGTA	ACCGGGCATG	TTCATCATCA	GTAACCCGTA	2150
TCGTGAGCAT	CCTCTCTCGT	TTCATCGGTA	TCATTACCCC	CATGAACAGA	2200
AATTCCCCTT	TACACGGAGG	CATCAAGTGA	CCAAACAGGA	AAAAACCGCC	2250
CTTAACATGG	CCCGCTTTAT	CAGAAGCCAG	ACATTAACGC	TTCTGGAGAA	2300
ACTCAACGAG	CTGGACGCGG	ATGAACAGGC	AGACATCTGT	GAATCGCTTC	2350
ACGACCACGC	TGATGAGCTT	TACCGCAGGA	TCCGAAATT	GTAAACGTTA	2400
ATATTTTGTT	AAAATTCGCG	TTAAATTTTT	GTAAATCAG	CTCATTTTTT	2450

-continued

AACCAATAGG	CCGAAATCGG	CAAAATCCCT	TATAAATCAA	AAGAATAGAC	2500
CGAGATAGGG	TTGAGTGTG	TTCCAGTTTG	GAACAAGAGT	CCACTATTAA	2550
AGAACGTGGA	CTCCAACGTC	AAAGGGCGAA	AAACCGTCTA	TCAGGGCTAT	2600
GGCCCACTAC	GTGAACCATC	ACCCTAATCA	AGTTTTTTGG	GGTCGAGGTG	2650
CCGTAAAGCA	CTAAATCGGA	ACCCTAAAGG	GAGCCCCCGA	TTTAGAGCTT	2700
GACGGGGAAA	GCCGGCGAAC	GTGGCGAGAA	AGGAAGGGAA	GAAAGCGAAA	2750
GGAGCGGGCG	CTAGGGCGCT	GGCAAGTGTA	GCGGTCACGC	TGCGCGTAAC	2800
CACCACACCC	GCCGCGCTTA	ATGCGCCGCT	ACAGGGCGCG	TCCGGATCCT	2850
GCCTCGCGCG	TTTCGGTGAT	GACGGTGAAA	ACCTCTGACA	CATGCAGCTC	2900
CCGAGACGG	TCACAGCTTG	TCTGTAAGCG	GATGCCGGGA	GCAGACAAGC	2950
CCGTCAGGGC	GCGTCAGCGG	GTGTTGGCGG	GTGTCGGGGC	GCAGCCATGA	3000
CCCAGTCACG	TAGCGATAGC	GGAGTGTATA	CTGGCTTAAC	TATGCGGCAT	3050
CAGAGCAGAT	TGTACTGAGA	GTGCACCATA	TGCGGTGTGA	AATACCGCAC	3100
AGATGCGTAA	GGAGAAAATA	CCGCATCAGG	CGCTCTCCG	CTCCTCGCT	3150
CACTGACTCG	CTGCGCTCGG	TCGTTCCGGCT	GCGGCGAGCG	GTATCAGCTC	3200
ACTCAAAGGC	GGTAATACGG	TTATCCACAG	AATCAGGGGA	TAACGCAGGA	3250
AAGAACATGT	GAGCAAAAAG	CCAGCAAAAG	GCCAGGAACC	GTAAAAAGGC	3300
CGCGTTGCTG	GCGTTTTTCC	ATAGGCTCCG	CCCCCCTGAC	GAGCATCACA	3350
AAAATCGACG	CTCAAGTCAG	AGGTGGCGAA	ACCCGACAGG	ACTATAAAGA	3400
TACCAGGCGT	TTCCCCCTGG	AAGCTCCCTC	GTGCGCTCTC	CTGTTCCGAC	3450
CCTGCCGCTT	ACCGGATACC	TGTCCGCCTT	TCTCCCTTCG	GGAAGCGTGG	3500
CGCTTTCTCA	TAGCTCACGC	TGTAGGTATC	TCAGTTCCGT	GTAGGTCGTT	3550
CGCTCCAAGC	TGGGCTGTGT	GCACGAACCC	CCCGTTCAGC	CCGACCGCTG	3600
CGCCTTATCC	GGTAACTATC	GTCTTGAGTC	CAACCCGGTA	AGACACGACT	3650
TATCGCCACT	GGCAGCAGCC	ACTGGTAACA	GGATTAGCAG	AGCGAGGTAT	3700
GTAGGCGGTG	CTACAGAGTT	CTTGAAGTGG	TGGCCTAACT	ACGGCTACAC	3750
TAGAAGGACA	GTATTTGGTA	TCTGCGCTCT	GCTGAAGCCA	GTTACCTTCG	3800
GAAAAAGAGT	TGGTAGCTCT	TGATCCGGCA	AACAACCAC	CGCTGGTAGC	3850
GGTGGTTTTT	TTGTTTGCAA	GCAGCAGATT	ACGCGCAGAA	AAAAAGGATC	3900
TCAAGAAGAT	CCTTTGATCT	TTTCTACGGG	GTCTGACGCT	CAGTGGAAAG	3950
AAAATCAGC	TTAAGGGATT	TTGGTCATGA	GATTATCAA	AAGGATCTTC	4000
ACCTAGATCC	TTTTAAATTA	AAAATGAAGT	TTTAAATCAA	TCTAAAGTAT	4050
ATATGAGTAA	ACTTGGTCTG	ACAGTTACCA	ATGCTTAATC	AGTGAGGCAC	4100
CTATCTCAGC	GATCTGTCTA	TTTCGTTTAT	CCATAGTTGC	CTGACTCCCC	4150
GTCGTGTAGA	TAACTACGAT	ACGGGAGGGC	TTACCATCTG	GCCCCAGTGC	4200
TGCAATGATA	CCGCGAGACC	CACGCTCACC	GGCTCCAGAT	TTATCAGCAA	4250
TAAACCAGCC	AGCCGGAAGG	GCCGAGCGCA	GAAGTGGTCC	TGCAACTTTA	4300
TCCGCCTCCA	TCCAGTCTAT	TAATTTGTTG	CGGGAAGCTA	GAGTAAGTAG	4350
TTCCGCAGTT	AATAGTTTGC	GCAACGTTGT	TGCCATTGCT	GCAGGCATCG	4400
TGGTGTACAG	CTCGTCGTTT	GGTATGGCTT	CATTCAGCTC	CGGTTCCCAA	4450

-continued

CGATCAAGGC GAGTTACATG ATCCCCCATG TTGTGCAAAA AAGCGGTTAG	4500
CTCCTTCGGT CCTCCGATCG TTGTGCAAG TAAGTTGGCC GCAGTGTAT	4550
CATCATGGT TATGGCAGCA CTGCATAATT CTCTTACTGT CATGCCATCC	4600
GTAAGATGCT TTTCTGTGAC TGGTGAGTAC TCAACCAAGT CATTCTGAGA	4650
ATAGTGTATG CGGCGACCGA GTTGCTCTTG CCCGGCGTCA ACACGGGATA	4700
ATACCGCGCC ACATAGCAGA ACTTTAAAAG TGCTCATCAT TGGAAAACGT	4750
TCTTCGGGGC GAAAACCTC AAGGATCTTA CCGCTGTGA GATCCAGTTC	4800
GATGTAACCC ACTCGTGCAC CCAACTGATC TTCAGCATCT TTTACTTTCA	4850
CCAGCGTTTC TGGGTGAGCA AAAACAGGAA GGCAAAATGC CGCAAAAAG	4900
GGAATAAGGG CGACACGGAA ATGTTGAATA CTCATACTCT TCCTTTTTCA	4950
ATATTATGA AGCATTATC AGGGTTATTG TCTCATGAGC GGATACATAT	5000
TTGAATGTAT TTAGAAAAAT AAACAAATAG GGGTTCGCG CACATTTCCC	5050
CGAAAAGTGC CACCTGACGT CTAAGAAACC ATTATTATCA TGACATTAAC	5100
CTATAAAAAT AGGCGTATCA CGAGGCCCTT TCGTCTTCAA	5140

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 77 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ser	Gly	Thr	Ala	Met	Ala	Asp	Pro	Asn	Arg	Phe	Arg	Gly	Lys	Asp
1				5					10					15
Leu	Ala	Gly	Ser	Pro	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Ala	Glu	Gly
				20					25					30
Asp	Asp	Pro	Ala	Lys	Ala	Ala	Phe	Asn	Ser	Leu	Gln	Ala	Ser	Ala
				35					40					45
Thr	Glu	Tyr	Ile	Gly	Tyr	Ala	Trp	Ala	Met	Val	Val	Val	Ile	Val
				50					55					60
Gly	Ala	Thr	Ile	Gly	Ile	Lys	Leu	Phe	Lys	Lys	Phe	Thr	Ser	Lys
				65					70					75
Ala	Ser													
														77

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 50 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GTTCGTATGT GGTGATCGAG GCTTCCTGTT CAACAAACCG ACTGGGGCTG	50
--	----

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 58 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

-continued

(ii) MOLECULE TYPE: Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GATCCAGCCC CAGTCGGTTT GTTGAACAGG AAGCCTCGAT CACCACATAC 50

GAACTGCA 58

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Ser Gly Thr Ala Cys Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Cys Ser
1 5 10 15Leu Ala Gly Ser Pro
20

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Xaa Xaa Xaa Xaa Cys Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Cys Xaa
1 5 10 15Xaa Xaa Xaa
18

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15Xaa Xaa Xaa Xaa Xaa
20

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Cys Xaa Xaa
1 5 10 15Xaa Xaa Xaa Xaa Xaa
20

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Cys
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15

Cys Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid

-continued

(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15
 Cys Xaa Xaa Xaa Xaa
 20

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

Gly Gly Thr Tyr Ser Cys His Phe Gly Pro Leu Thr Trp Val Cys
 1 5 10 15
 Lys Pro Gln Gly Gly
 20

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Cys Xaa Xaa Gly Pro Xaa Xaa Xaa Xaa Cys
 1 5 10

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 70 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GCCTATGCAT CTGGTACCGC CTGCNNSNNS GGCCTNNSN NSNNSNNSTG 50
 TTCTCTGGCA GGTTCACCAG 70

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 91 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Nucleic Acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

GCTACAAATG CCTATGCANN SNNSNNSNNS TGCNNSNNSG GTCCTNNSNN 50
 SNNSNNSSTGT NNSNNSNNSN NSGGTGGAGG ATCCGGAGGA G 91

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 97 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSNNSNNSST GCNNSNNSNN	50
SNNSTGCNNS NNSNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG	97
(2) INFORMATION FOR SEQ ID NO:39:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 97 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Single	
(D) TOPOLOGY: Linear	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSNNSNNSST GCNNSNNSNN	50
SNNNSNNSSTGC NNSNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG	97
(2) INFORMATION FOR SEQ ID NO:40:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 97 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Single	
(D) TOPOLOGY: Linear	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSNNSSTGCN NSNNSNNSNN	50
SNNNSNNSSTGC NNSNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG	97
(2) INFORMATION FOR SEQ ID NO:41:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 97 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Single	
(D) TOPOLOGY: Linear	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSNNSSTGCN NSNNSNNSNN	50
SNNNSNNSNS TGCNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG	97
(2) INFORMATION FOR SEQ ID NO:42:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 97 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Single	
(D) TOPOLOGY: Linear	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSTGCNNSN NSNNSNNSNN	50
SNNNSNNSNS TGCNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG	97
(2) INFORMATION FOR SEQ ID NO:43:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 97 base pairs	
(B) TYPE: Nucleic Acid	
(C) STRANDEDNESS: Single	
(D) TOPOLOGY: Linear	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:	
GCTACAAATG CCTATGCANN SNNNSNNSNS NNSTGCNNSN NSNNSNNSNN	50

-continued

SNNSNNSNNS NNSTGCNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG 97

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 97 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GCTACAAATG CCTATGCANN SNNSNNSNNS TGCNNSNNSN NSNNSNNSNN 50

SNNSNNSNNS NNSTGCNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG 97

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 97 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GCTACAAATG CCTATGCANN SNNSNNSNNS NNSNNSNNSN NSNNSNNSNN 50

SNNSNNSNNS NNSNNSNNSN NSNNSNNSGG TGGAGGATCC GGAGGAG 97

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

Ser Gly Thr Ala Cys Tyr Gly Gly Pro Glu Trp Trp Cys Cys Ser
 1 5 10 15

Leu Ala Gly Ser Pro
 20

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Asp Leu Ala Ile Cys Ala Glu Gly Pro Glu Ile Trp Val Cys Glu
 1 5 10 15

Glu Thr Ser
 18

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Asp Phe Trp Ile Cys Leu Ser Gly Pro Gly Trp Glu Glu Cys Leu
 1 5 10 15

Glu Trp Trp

-continued

18

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

Gly Ser Ala Gly Gln Gly Met Thr Glu Glu Trp Ala Trp Ile Trp
 1 5 10 15
 Glu Trp Trp Lys Glu
 20

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

Glu Leu Asp Gly Trp Val Cys Ile Lys Val Gly Glu Gln Asn Leu
 1 5 10 15
 Cys Tyr Leu Ala Glu
 20

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

Ala Ile Gly Gly Trp Cys Phe Ile Glu Leu Asp Ser Leu Trp Cys
 1 5 10 15
 Glu Glu Gln Ile Gly
 20

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Ser Glu Asp Val Glu Cys Trp Gln Val Trp Glu Asn Leu Val Cys
 1 5 10 15
 Ser Val Glu His Arg
 20

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Ser Glu Glu Val Cys Trp Pro Val Ala Glu Trp Tyr Leu Cys Asn
 1 5 10 15

-continued

Met Trp Gly Arg
19

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

Arg Val Gly Ala Tyr Ile Ser Cys Ser Glu Thr Glu Cys Trp Val
1 5 10 15

Glu Asp Leu Leu Asp
20

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Trp Phe Lys Thr Val Cys Tyr Glu Trp Glu Asp Glu Val Gln Cys
1 5 10 15

Tyr Thr Leu Glu Glu
20

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

Arg Leu Glu Glu Gln Cys Val Glu Val Asn Tyr Glu Pro Ser Cys
1 5 10 15

Ser Phe Thr Ala Asn
20

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Ser Glu Glu Val Cys Trp Pro Val Ala Glu Trp Tyr Leu Cys Asn
1 5 10 15

Ile Leu Gly Pro
19

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

Glu Thr Val Ala Asn Cys Asp Cys Tyr Met Asp Leu Cys Leu Cys
1 5 10 15

-continued

Tyr Gly Ser Asp Arg
20

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Tyr His Pro Ile Ser Cys Met Asp His Tyr Tyr Leu Ile Ile Cys
1 5 10 15

Asp Glu Thr Val Asn
20

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

Ala Glu Trp Ala Glu Cys Trp Ile Ala Gly Asp Gln Leu Leu Cys
1 5 10 15

Val Gly Lys Asp Asn
20

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Glu Pro Trp Leu Cys Gln Tyr Tyr Glu Ala Ala Met Leu Tyr Leu
1 5 10 15

Cys Trp Glu Glu Gly
20

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Ala Glu Glu Gly Met Val Trp Gly Trp Thr Gly Gly Trp Tyr Asn
1 5 10 15

Leu Asp Glu Leu Cys
20

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Ser Gly Gly Ala Ile Tyr Trp Pro Val Glu Gln Phe Ile Ala Phe

-continued

1	5	10	15
---	---	----	----

Met Ala Val Gly Lys
20

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

Ser Gly Gly Ala Ile Tyr Met Pro Val Glu Gln Phe Ile Ala Phe			
1	5	10	15

Met Ala Val Gly Lys
20

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Glu Val Leu Leu Cys Ser Asp Gly Pro Gln Leu Tyr Leu Cys Glu			
1	5	10	15

Leu Tyr Ala
18

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Ser Gly Val Glu Cys Val Trp Gly Pro Gln Trp Gly Phe Cys Val			
1	5	10	15

Glu Glu Tyr
18

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Asp Lys Glu Val Cys Tyr Leu Gly Pro Glu Thr Trp Leu Cys Phe			
1	5	10	15

Trp Trp Pro
18

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

-continued

Gly Asp Val Glu Cys Ile Glu Gly Pro Trp Gly Glu Leu Cys Val
 1 5 10 15

Trp Ala Asp
 18

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

Phe Gly Gly Trp Ser Cys Gln Pro Thr Trp Val Asp Val Tyr Val
 1 5 10 15

Cys Asn Phe Glu Glu
 20

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

Ala Met Trp Val Cys Val Ser Asp Trp Glu Thr Val Glu Glu Cys
 1 5 10 15

Ile Gln Tyr Met Tyr
 20

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

Thr Asn Trp Phe Phe Val Cys Glu Ser Gly His Gln Asp Ile Cys
 1 5 10 15

Trp Leu Ala Glu Glu
 20

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

Ser Glu Val Gly Cys Arg Ala Gly Pro Leu Gln Leu Cys Glu Lys
 1 5 10 15

Tyr Phe
 17

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

-continued

Lys Asp Pro Val Cys Gly Glu Gly Pro Leu Met Arg Ile Cys Glu
 1 5 10 15

Arg Leu Phe Gly
 19

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

Glu Val Asp Gly Arg Trp Trp Ile Val Glu Thr Phe Leu Ala Lys
 1 5 10 15

Trp Asp His Met Ala Gly
 20 21

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

Trp Val Met Glu Cys Gly Ala Gly Pro Trp Pro Glu Gly Cys Thr
 1 5 10 15

Phe Met Leu
 18

(2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

Arg Lys Thr Ser Gln Gly Arg Gly Gln Glu Met Cys Trp Glu Thr
 1 5 10 15

Gly Gly Cys Ser
 19

(2) INFORMATION FOR SEQ ID NO:77:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

Ser Trp Glu Arg Gly Glu Leu Thr Tyr Met Lys Leu Cys Glu Tyr
 1 5 10 15

Met Arg Leu Gln Gln
 20

(2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

