



US007563879B2

(12) **United States Patent**
Palli

(10) **Patent No.:** **US 7,563,879 B2**
(45) **Date of Patent:** **Jul. 21, 2009**

(54) **LEAFHOPPER ECDYSONE RECEPTOR
NUCLEIC ACIDS, POLYPEPTIDES, AND USES
THEREOF**

(75) Inventor: **Subba Reddy Palli**, Lansdale, PA (US)
(73) Assignee: **Intrexon Corporation**, Blacksburg, VA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 566 days.

(21) Appl. No.: **10/490,971**

(22) PCT Filed: **Feb. 20, 2002**

(86) PCT No.: **PCT/US02/05026**

§ 371 (c)(1),
(2), (4) Date: **Mar. 25, 2004**

(87) PCT Pub. No.: **WO03/027289**

PCT Pub. Date: **Apr. 3, 2003**

(65) **Prior Publication Data**

US 2004/0197861 A1 Oct. 7, 2004

Related U.S. Application Data

(60) Provisional application No. 60/325,096, filed on Sep. 26, 2001.

(51) **Int. Cl.**

C07H 21/02 (2006.01)
C07H 21/04 (2006.01)
C07K 1/00 (2006.01)
C12N 5/00 (2006.01)
C12P 21/06 (2006.01)

(52) **U.S. Cl.** **536/23.1; 536/23.5; 530/350; 435/4; 435/320.1; 435/69.1; 435/70.1; 435/325; 435/252.3; 435/254.11; 435/419; 435/254.2**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,631,211 A 12/1986 Houghten
4,859,609 A 8/1989 Dull et al.
4,954,655 A 9/1990 Kelly
4,981,784 A 1/1991 Evans et al.
4,985,461 A 1/1991 Hsu et al.
5,010,175 A 4/1991 Rutter et al.
5,071,773 A 12/1991 Evans et al.
5,117,057 A 5/1992 Hsu et al.
5,171,671 A 12/1992 Evans et al.
5,225,443 A 7/1993 Murphy et al.
5,283,173 A 2/1994 Fields et al.
5,378,726 A 1/1995 Yanagi et al.
5,424,333 A 6/1995 Wing
5,429,952 A 7/1995 Garner et al.
5,459,127 A 10/1995 Felgner et al.
5,514,578 A 5/1996 Hogness et al.
5,530,028 A 6/1996 Lidert et al.
5,599,904 A 2/1997 Evans et al.
5,641,652 A 6/1997 Oro et al.
5,668,175 A 9/1997 Evans et al.

5,688,691 A 11/1997 Oro et al.
5,710,004 A 1/1998 Evans et al.
5,723,329 A 3/1998 Mangelsdorf et al.
5,880,333 A 3/1999 Goff et al.
5,919,667 A 7/1999 Gage et al.
5,989,863 A 11/1999 Tang
6,013,836 A 1/2000 Hsu et al.
6,025,483 A 2/2000 Yanofsky
6,096,787 A 8/2000 Evans et al.
6,117,639 A 9/2000 Germann
6,147,282 A 11/2000 Goff et al.
6,245,531 B1 6/2001 Hogness et al.
6,265,173 B1 6/2001 Evans et al.
6,281,330 B1 8/2001 Evans et al.
6,300,488 B1 10/2001 Gage et al.
6,333,318 B1 12/2001 Evans et al.
6,379,945 B1* 4/2002 Jepson et al. 435/243
6,458,926 B1 10/2002 Evans et al.
6,504,082 B1 1/2003 Albertsen et al.
6,635,429 B1 10/2003 Leid et al.
6,723,531 B2 4/2004 Evans et al.
6,756,491 B2 6/2004 Evans et al.
6,875,569 B2 4/2005 Gage et al.
6,939,711 B2 9/2005 Goff et al.
6,958,236 B2* 10/2005 Pascal et al. 435/320.1
7,038,022 B1 5/2006 Evans et al.
7,045,315 B2 5/2006 Evans et al.
7,057,015 B1 6/2006 Gage et al.
7,091,038 B2 8/2006 Palli et al.
7,119,077 B1 4/2007 Evans et al.
2002/0110861 A1 8/2002 Dhadialla et al.
2002/0119521 A1 8/2002 Palli et al.
2004/0033600 A1 2/2004 Palli et al.
2004/0096942 A1 5/2004 Kapitskaya et al.
2004/0197861 A1 10/2004 Palli
2004/0235097 A1 11/2004 Zhang et al.
2005/0266457 A1 12/2005 Palli et al.
2006/0100416 A1 5/2006 Palli et al.
2007/0161086 A1 7/2007 Palli et al.
2008/0064097 A1 3/2008 Palli

FOREIGN PATENT DOCUMENTS

CN 1313276 A 9/2001

(Continued)

OTHER PUBLICATIONS

Michael Ashburner et al., "Temporal Control of Puffing Activity in Polytene Chromosomes", Cold Spring Harbor Symp. Quant. Biol., 1974, 38:655-662.

(Continued)

Primary Examiner—Michael Pak
Assistant Examiner—Nirmal S Basi
(74) *Attorney, Agent, or Firm*—Sterne, Kessler, Goldstein & Fox P.L.L.C

(57) **ABSTRACT**

The present invention relates to a novel isolated leafhopper ecdysone receptor polypeptide. The invention also relates to an isolated nucleic acid encoding the leafhopper ecdysone receptor polypeptide, to vectors comprising them and to their uses, in particular in methods for modulating gene expression in an ecdysone receptor-based gene expression modulation system and methods for identifying molecules that modulate leafhopper ecdysone receptor activity.

FOREIGN PATENT DOCUMENTS

EP	0 461 809	B1	12/1991
EP	461809	A1	12/1991
EP	0 798 378	B1	10/1997
EP	0 965 644	B1	12/1999
EP	1 266 015	B1	12/2002
WO	WO 8912690	A1	12/1989
WO	WO 9200252	A1	1/1992
WO	WO 9428028		12/1994
WO	WO 9518863	A1	7/1995
WO	WO 9521931	A1	8/1995
WO	WO 96/37609		5/1996
WO	WO 9617823	A1	6/1996
WO	WO 9625508	A1	8/1996
WO	WO 9627673		9/1996
WO	WO 97/35985	A1	10/1997
WO	WO 9738117	A1	10/1997
WO	WO 98/33162	A1	7/1998
WO	WO 9902683	A1	1/1999
WO	WO 99/10510	A2	3/1999
WO	WO 99/10510	A3	3/1999
WO	WO 99/27365	A1	6/1999
WO	WO 99/36520	A1	7/1999
WO	WO 9936520	A1	7/1999
WO	WO 99/51777	A2	10/1999
WO	WO 9958155	A1	11/1999
WO	WO 00/71743	A1	11/2000
WO	WO 01/02436		1/2001
WO	WO 01/36447		5/2001
WO	WO 01/62780		8/2001
WO	WO 0170816	A2	9/2001
WO	WO 02029075	A2	4/2002
WO	WO 02066612	A2	8/2002
WO	WO 02066613	A2	8/2002
WO	WO 02066614	A2	8/2002
WO	WO 02066615	A2	8/2002
WO	WO 03/105849	A1	12/2003
WO	WO 2004/005478	A2	1/2004
WO	WO 2004/072254	A2	8/2004
WO	WO 2004/078924	A2	9/2004
WO	WO 2005/017126	A2	2/2005
WO	WO 2005/108617	A2	11/2005
WO	WO 05108617	A2	11/2005
WO	WO 2006/083253	A1	8/2006

OTHER PUBLICATIONS

- Tarlochan S. Dhadialla, et al., "New Insecticides with Ecdysteroidal and Juvenile Hormone Activity", *Annu. Rev. Entomol.*, 1988, 45:545-569.
- Michael R. Koelle et al., "The *Drosophila* EcR Gene Encodes and Ecdysone Receptor, a New Member of the Steroid Receptor Superfamily", *Cell*, 1991, 67: 59-77.
- Ravi Kothapalli et al., "Cloning and Developmental Expression of the Ecdysone Receptor Gene From the Spruce Budworm, *Choristoneura fumiferana*", *Developmental Genetics*, 1995, 17: 319-30.
- Jean-Francois Mouillet et al., "Cloning of two putative ecdysteroid receptor isoforms from *Tenebrio molitor* and their developmental expression in the epidermis during metamorphosis", *Eur. J. Biochem.*, 1997, 248: 856-863.
- Haruhiko Fujiwara et al., "Cloning of an Ecdysone Receptor Homolog from *Manduca sexta* and the Developmental Profile of Its mRNA in Wings", *Insect Biochem. Molec. Biol.*, 1995, 25, 845-856.
- Alberto Martinez et al., "Transcriptional activation of the cloned *Heliothis virescens* (*Lepidoptera*) ecdysone receptor (HvEcR) by MuristeroneA", *Insect Biochemistry and Molecular Biology*, 1999, 29: 915-930.
- Markus O. Imhof et al., Cloning of a *Chironomus tentans* cDNA Encoding a Protein (cEcRH) Homologous to the *Drosophila melanogaster* Ecdysteroid Receptor (dEcR). *Insect Biochem. Molec. Biol.*, 1993, 23: 115-124.
- Luc Swevers et al., "The Silkmoth Homolog of the *Drosophila* Ecdysone Receptor (B1 Isoform): Cloning and Analysis of Expression During Follicular Cell Differentiation", *Insect Biochem. Molec. Biol.*, 1995, 25:857-856.
- Wen-Long Cho et al., "Mosquito Ecdysteroid Receptor: Analysis of the cDNA and Expression During Vitellogenesis", *Insect Biochem. Molec. Biol.*, 1995, 25: 19-27.
- Gary N. Hannan et al., "Cloning and Characterization of LcEcR: A Functional Ecdysone Receptor from the Sheep Blowfly *Lucilia cuprina*", *Insect Biochem. Molec. Biol.*, 1997, 27: 479-488.
- Meletis Verras et al., "Cloning and characterization of CcEcR: An ecdysone receptor homolog from the Mediterranean fruit fly *Ceratitis capitata*", *Eur. J. Biochem.*, 1999, 265: 798-808.
- David S. Saleh et al., "Cloning and characterization of an ecdysone receptor cDNA from *Locusta migratoria*", *Molecular and Cellular Endocrinology*, 1998, 143: 91-99.
- Arthur C. -K. Chung et al., "Cloning of crustacean ecdysteroid receptor and retinoid-X receptor gene homologs and elevation of retinoid-X receptor mRNA by retinoic acid", *Molecular and Cellular Endocrinology*, 1998, 139: 209-277.
- Xiaoping Guo et al., "Isolation of a Functional Ecdysteroid Receptor Homologue from the Ixodid Tick *Amblyomma americanum* (L.)", *Insect Biochem. Molec. Biol.* 1997, 27: 945-962.
- Ronald M. Evans, "The Steroid and Thyroid Hormone Receptor Superfamily", *Science*, 1988, 240: 889-895.
- Lynn M. Riddiford et al., "Ecdysone Receptors and Their Biological Actions", *Vitamins and Hormones*, 2000, 60:1-73.
- Karen S. Christopherson et al., "Ecdysteroid-dependent regulation of genes in mammalian cells by a *Drosophila* ecdysone receptor and chimeric transactivators", *Proc. Natl. Acad. Sci. U.S.A.*, 1992, 89:6314-6318.
- David NO et al., "Ecdysone-inducible gene expression in mammalian cells and transgenic mice", *Proc. Natl. Acad. Sci.*, 1996, 93: 3346-3351.
- Steven T. Suhr et al., High level transactivation by a modified Bombyx ecdysone receptor in mammalian cells without exogenous retinoid X receptor, *Proc. Natl. Acad. Sci. U.S.A.*, 1998, 95:7999-8004.
- Ulrike Heberlein et al., "Characterization of *Drosophila* Transcription Factors That Activate the Tandem Promoters of the Alcohol Dehydrogenase Gene", *Cell*, 1985, 41: 965-977.
- James M. Wilson et al., "Hepatocyte-directed Gene Transfer in Vivo Leads to Transient Improvement of Hypercholesterolemia in Low Density Lipoprotein Receptor-deficient Rabbits", *J. Biol. Chem.*, 1992, 267:963-967.
- Michaels S. Neuberger et al., "Recombinant antibodies possessing novel effector functions", *Nature*, 1984, 312: 604-608.
- Christophe Antoniewski et al., "The Ecdysone Response Enhancer of the Fbp1 Gene of *Drosophila melanogaster* Is a Direct Target for the EcR/USP Nuclear Receptor", *Molecular and Cellular Biology*, 1994, 14, 4465-4474.
- Donald A. Morrison et al., "Isolation of Transformation-Deficient *Streptococcus pneumoniae* Mutants Defective in Control of competence, Using Insertion-Duplication Mutagenesis with the Erythromycin Resistance Determinant of pAMBeta1", *J. Bacteriol.* 1984, 159:870.
- D. Metzger et al., "The human oestrogen receptor functions in yeast", *Nature*, 1988, 334:31-36.
- Paul J. Godowski et al., "Signal Transduction and Transcriptional Regulation by Glucocorticoid Receptor-LexA Fusion Proteins", *Science*, 1988, 241: 812-816.
- Pier Paolo D'Avino et al., "The moulting hormone ecdysone is able to recognize target elements composed of direct repeats", *Molecular and Cellular Endocrinology*, 113: 1-9. 1995.
- Lucy Cherbas et al., "Identification of ecdysone response elements by analysis of the *Drosophila* Eip28/29 gene", *Genes & Development*, 1991, 5:120-131.
- U.S. Appl. No. 11/837,834, filed Aug. 13, 2007.
- U.S. Appl. No. 11/837,848, filed Aug. 13, 2007.
- U.S. Appl. No. 11/852,194, filed Sep. 7, 2007.
- Adrianov, V.G., et al., "4-Aminofurazan-3-Hydroxamic Halides," *Chem. Hetero. Comp.* 28:5815585, Springer New York (1992).

- Andrianov, V.G., et al., "4-Amino- Δ^2 , 4-Oxadiazolines," *Chem. Hetero. Comp.* 22:216-218, Springer New York (1991).
- Belshaw, P.J., et al., "Controlling protein association and subcellular localization with a synthetic ligand that induces heterodimerization of proteins," *Proc. Natl. Acad. Sci.* 93:4604-4607, National Academy of Science (1996).
- Belshaw, P.J., et al., "Rational Design of Orthogonal Receptor - Ligand Combinations," *Angew. Chem. Int. Ed. Engl.* 34:2129-2132, Wiley-VCH (1995).
- Bonneton, F., et al., "Rapid Divergence of the Ecdysone Receptor in Diptera and Lepidoptera Suggests Coevolution Between ECR and USP-RXR," *Mol. Biol. Evol.* 20:541-553, Oxford University Press (Apr. 2003).
- Brennan, J.D., "Preparation and Entrapment of Fluorescently Labeled Proteins for the Development of Reagentless Optical Biosensors," *J. Fluoresc.* 9:295-312, Springer (1999).
- Cao, S., et al., "*N*-*tert*-Butyl-*N*'-aroyl-*N*-(alkoxycarbonylmethyl)-*N*-aroylhydrazines, a novel nonsteroidal ecdysone agonist: syntheses, insecticidal activity, conformational, and crystal structure analysis," *Can. J. Chem.* 79:272-278, NRC Research Press (Mar. 2001).
- Carlson, G.R., et al., "The chemical and biological properties of methoxyfenozide, a new insecticidal ecdysteroid agonist," *Pest Manag. Sci.* 57:115-119, Society of Chemical Industry (Jan. 2001).
- Doyle, D.F., et al., "Engineering Orthogonal Ligand - Receptor Pairs from "Near Drugs", " *J. Am. Chem. Soc.* 123:11367-11371, American Chemical Society (Oct. 2001).
- Egea, P.F., et al., "Effects of Ligand Binding on the Association Properties and Conformation in Solution of Retinoic Acid Receptors RXR and RAR," *J. Mol. Biol.* 307:557-576, Academic Press (Mr. 2001).
- Fields, S. and Song, O., "A novel genetic system to detect protein-protein interactions," *Nature* 340:245-246, Nature Publishing Group (1989).
- Filmus, J., et al., "Synergistic induction of promoters containing metal- and glucocorticoid-responsive elements," *Nucl. Acids Res.* 20:2755-2760, Current Science (1992).
- Glass, C.K., et al., "Nuclear receptor coactivators," *Curr. Opin. Cell Biol.* 9:222-232, Current Science (1997).
- Hayward, D.C., et al., "The structure of the USP/RXR of *Xenos pecki* indicates that Strepsiptera are not closely related to Diptera," *Dev. Genes Evol.* 215:213-219, Springer Berlin / Heidelberg (Jan. 2005).
- Holt, J.R. et al., "Functional Expression of Exogenous Proteins in Mammalian Sensory Hair Cells Infected With Adenoviral Vectors," *J. Neurophysiol.* 81:1881-1888, American Physiological Society (1999).
- Hoppe, U.C. et al., "Adenovirus-Mediated Inducible Gene Expression in Vivo by Hybrid Ecdysone Receptor," *Mol. Ther.* 1:159-164, Academic Press (2000).
- Horwitz, K.B., et al., "Nuclear Receptor Coactivators and Corepressors," *MOL. Endocrinol.* 10:1167-1177, Endocrine Society (1996).
- Kakizawa, T., et al., "Ligand-dependent Heterodimerization of Thyroid Hormone Receptor and Retinoid X Receptor," *J. Biol. Chem.* 272:23799-23804, American Society for Biochemistry and Molecular Biology (1997).
- Kim, J-S., et al., "Design of TATA box-binding protein/zinc finger fusions for targeted regulation of gene expression," *Proc. Natl. Acad. Sci. U.S.A.* 94:3616-3620, National Academy of Science (1997).
- Kirken, R.A. et al., "Two Discrete Regions of Interleukin-2 (IL2) Receptor β Independently Mediate IL2 Activation of a PD98059/Rapamycin/Wortmannin-insensitive Stat5/a/b Serine Kinase," *J. Biol. Chem.* 272:15459-15465, American Society for Biochemistry and Molecular Biology (1997).
- Leid, M., et al., "Purification, cloning, and RXR identity of the HeLa cell factor with which RAR or TR heterodimerizes to bind target sequences efficiently," *Cell* 68:377-395, Cell Press (1992).
- Leonhardt, S.A., et al., "Agonist and Antagonists Induce Homodimerization and Mixed Ligand Heterodimerization of Human Progesterone Receptors in Vivo by a Mammalian Two-Hybrid Assay," *Mol. Endocrinol.* 12:1914-1930, The Endocrine Society (1998).
- Licitra, E.J., et al., "A three-hybrid systems for detecting small ligand-protein receptor interactions," *Proc. Natl. Sci. U.S.A.* 93:12817-12821, National Academy of Science (1996).
- Martinez, A., et al., "Creation of ecdysone receptor chimeras in plants for controlled regulation of gene expression," *Mol. Gen. Genetic* 261:564-552, Springer-Verlag (1999).
- Moradpour, D., et al., "Independent Regulation of Two Separate Gene Activities in a Continuous Human Cell Line," *Biol. Chem.* 379:1189-1191, Walter De Gruyter (1998).
- Nakagawa, Y., et al., "Quantitative structure-activity studies of insect growth regulators: XIX. Effects of substituents on the aromatic moiety of dibenzoylhydrazines on larvicidal activity against the beet armyworm *Spodoptera eximia*," *Pest. Manag. Sci.* 58:131-138, Wiley (2001).
- O'Brien, R.M., et al., "Structural and functional analysis of the human phospho *enol* pyruvate carboxykinase gene promoter," *Biochim. Biophys. Acta* 1264:284-288, Elsevier Science (1995).
- Peet, D.J., et al., "Engineering novel specificities for ligand-activated transcription in the nuclear hormone receptor RXR," *Chem. Biol.* 5:13-21, Current Biology (1998).
- Perera, S.C., et al., "An Analysis Of Ecdysone Receptor Domains Required for Heterodimerization With Ultraspiracle," *Arch. Insect Biochem. Physiol.* 41:61-70, Wiley-Liss (1999).
- Perera, S.C., et al., "Studies on two ecdysone receptor isoforms of the spruce budworm, *Choristoneura fumiferana*," *Mol. Cell. Endocrinol.* 152:73-84, Elsevier Science (1999).
- Pierce, A.C. and Jorgensen, W.L., "Computational Binding Studies of Orthogonal Cyclosporin-Cyclophilin Pairs," *Angew. Chem. Int. Ed. Engl.* 36:1466-1469, Wiley-VCH Verlag (1997).
- Shea, C., et al., "An rxr/usp homolog from the parasitic nematode, *Dirioflaria immitis*," *Gene* 324:171-182, Elsevier/North-Holland (Nov. 2003).
- Spencer, D.M., et al., "Controlling Signal Transduction with Synthetic Ligands," *Science* 262:1019-1024, Association for the Advancement of Science (1993).
- Trisyone, A., and Chippendale, M.G., "Effect of the Nonsteroidal Ecdysone Agonist, Methoxyfenozide and Tebufenozide, on the European Corn Borer (*Lepidoptera: Pyralidae*)," *J. Econ. Entomol.* 90:1486-1492, Entomological Society of America (1997).
- Wing, K.D., "RH 5849, a Nonsteroidal Ecdysone Agonist: Effects on *Drosophila* Cell Line," *Science* 241:467-469, Association for the Advancement of Science (1988).
- Wipf, P., et al., "Combinatorial synthesis and Biological Evaluation of Library of Small-Molecule Ser/Thr-Protein Phosphatase Inhibitors," *Bioorg. Med. Chem.* 5:165-177, Elsevier Science (1997).
- Wurm, F.M., et al., "Inducible overproduction of the mouse c-myc protein in mammalian cells," *Proc. Natl. Acad. Sci. U.S.A.* 83:5414-5418, National Academy of Science (1986).
- Yao, T-P., et al., "Drosophila ultraspiracle Modulates Ecdysone Receptor Function via Heterodimer Formation," *Cell* 71:63-72, Cell Press (1992).
- Yao, T-P., et al., "Functional ecdysone receptor is the product of *EcR* and *Ultraspiracle* gene," *Nature* 366:476-479, Nature Publishing Group (1993).

* cited by examiner




Figure 1




Figure 2

1

LEAFHOPPER ECDYSONE RECEPTOR NUCLEIC ACIDS, POLYPEPTIDES, AND USES THEREOF

This application is the U.S. national phase of International Application No. PCT/US02/05026, filed Feb. 20, 2002, which claims the benefit of U.S. Provisional Application No. 60/325,096, filed Sep. 26, 2001.

FIELD OF THE INVENTION

This invention relates to the field of biotechnology. Specifically, this invention relates to isolated nucleic acids, vectors comprising them, and polypeptides encoded by them, and to their use in the field of gene expression and insecticide discovery. More specifically, this invention relates to a novel nucleic acid encoding an ecdysone receptor polypeptide from the homopteran green leafhopper (*Nephotetix cincticeps*, "NcEcR") and its use in methods of modulating the expression of a gene within a host cell using NcEcR, and in methods of identifying molecules that modulate the activity of the NcEcR.

BACKGROUND OF THE INVENTION

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties. However, the citation of any reference herein should not be construed as an admission that such reference is available as "Prior Art" to the instant application.

Cultivated agriculture has greatly increased efficiency of food production in the world. However, various insect pests have found it advantageous to seek out and exploit cultivated sources of food to their own advantage. These insect pests typically develop by a temporal sequence of events which are characteristic of their order. Many insects initially develop in a caterpillar or maggot-like larval form. Thereafter, they undergo a significant metamorphosis from which an adult emerges having characteristic anatomical features. Anatomic similarity is a reflection of developmental, physiological and biochemical similarities shared by these creatures. In particular, the principles of the insect ecdysteroid-hormone receptors and development, as described by Ashburner et al. (Cold Spring Harbor Symp. Quant. Biol. 38: 655-662, 1974), likely would be shared by many different types of insects.

To prevent or reduce the destruction of cultivated crops by insects, organic molecules with pesticidal properties are used commonly in attempts to eliminate or reduce the insect populations. However, the ecological side effects of these pesticides, due in part to their broad activity and lack of specificity, and in part, to the fact that some of these pesticides are not easily biodegradable, significantly affect populations of both insect and other species of animals. Some of these organisms may be advantageous from an ecological or other perspective. Furthermore, as the insect populations evolve in directions to minimize the effects of the applied pesticides, the amounts of pesticides applied are often elevated so high as to cause significant effects on other animals, including humans, which are affected directly or indirectly by the application of the pesticides. Thus, an important need exists for both highly specific pesticides or highly active pesticides which have biological effects only on the species of animals targeted by the pesticides, and are biodegradable. Novel insect hormones which, like the ecdysteroids, act by complexing with insect members of the steroid receptor superfamily to control insect development, are likely candidates for pesticides with these desirable properties.

2

Growth, molting, and development in insects are regulated by the ecdysone steroid hormone (molting hormone) and the juvenile hormones (Dhadialla, et al., 1998, Annu. Rev. Entomol. 43: 545-569). The molecular target for ecdysone in insects consists of at least ecdysone receptor (EcR) and ultraspiracle protein (USP). EcR is a member of the nuclear steroid receptor super family that is characterized by signature DNA and ligand binding domains, and an activation domain (Koelle et al. 1991, Cell, 67: 59-77). EcR receptors are responsive to a number of steroidal compounds such as ponasterone A and muristerone A. Recently, non-steroidal compounds with ecdysteroid agonist activity have been described, including the commercially available insecticides tebufenozide and methoxyfenozide (see International Patent Application No. PCT/EP96/00686 and U.S. Pat. No. 5,530, 028). Both analogs have exceptional safety profiles to other organisms.

Polynucleotides encoding ecdysone receptors have been cloned from a variety of insect species, including Dipterans (see U.S. Pat. Nos. 5,514,578 and 6,245,531 B1), Lepidopterans, Orthopterans, Hemipterans, and one Homopteran Aphid, all from the class Arthropod. In particular, EcRs have been cloned from spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"; Kothapalli et al., 1995 Dev Genet. 17: 319-30), a yellow meal worm *Tenebrio molitor* EcR ("TmEcR"; Mouillet et al., 1997, Eur. J. biochem. 248: 856-863), a tobacco hornworm *Manduca sexta* EcR ("MsEcR"; Fujiwara et al., 1995, Insect Biochem. Molec. Biol. 25, 845-856), a tobacco budworm *Heliothis virescens* EcR ("HvEcR"; Martinez et al., 1999, *Insect Biochem Mol Biol.* 29: 915-30), a golmidge *Chironomus tentans* EcR ("CtEcR"; Imhof et al., 1993, Insect Biochem. Molec. Biol. 23: 115-124), a silkworm *Bombyx mori* EcR ("BmEcR"; Swevers et al., 1995, Insect Biochem. Molec. Biol. 25: 857-866), a squinting bush brown *Bicyclus anynana* EcR ("BanEcR"), a buckeye *Junonia coenia* EcR ("JcEcR"), a fruit fly *Drosophila melanogaster* EcR ("DmEcR"; Koelle et al., 1991, Cell 67: 59-77), a yellow fever mosquito *Aedes aegypti* EcR ("AaEcR"; Cho et al., 1995, Insect Biochem. Molec. Biol. 25: 19-27), a blowfly *Lucilia capitata* ("LcEcR"), a sheep blowfly *Lucilia cuprina* EcR ("LucEcR"; Hannan and Hill, 1997, Insect Biochem. Molec. Biol. 27: 479-488), a blowfly *Calliphora vicina* EcR ("CvEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"; Verras et al., 1999, *Eur J Biochem.* 265: 798-808), a locust *Locusta migratoria* EcR ("LmEcR"; Saleh et al., 1998, *Mol Cell Endocrinol.* 143: 91-9), an aphid *Myzus persicae* EcR ("MpEcR"; International Patent Application Publication WO99/36520), a fiddler crab *Celuca pugilator* EcR ("CpEcR"; Chung et al., 1998, *Mol Cell Endocrinol* 139: 209-27), and an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"; Guo et al., 1997, Insect Biochem. Molec. Biol. 27: 945-962). The nucleotide and/or amino acid sequences of these ecdysone receptors have been determined and are publicly available.

The ecdysone receptor complex typically includes proteins that are members of the nuclear receptor superfamily wherein all members are generally characterized by the presence of an amino-terminal transactivation domain, a DNA binding domain ("DBD"), and a ligand binding domain ("LBD") separated from the DBD by a hinge region. As used herein, the term "DNA binding domain" comprises a minimal polypeptide sequence of a DNA binding protein, up to the entire length of a DNA binding protein, so long as the DNA binding domain functions to associate with a particular response element. Members of the nuclear receptor superfamily are also characterized by the presence of four or five domains: A/B, C, D, E, and in some members F (see U.S. Pat.

No. 4,981,784 and Evans, *Science* 240:889-895 (1988)). The “A/B” domain corresponds to the transactivation domain, “C” corresponds to the DNA binding domain, “D” corresponds to the hinge region, and “E” corresponds to the ligand binding domain. Some members of the family may also have another transactivation domain on the carboxy-terminal side of the LBD corresponding to “F”.

The DBD is characterized by the presence of two cysteine zinc fingers between which are two amino acid motifs, the P-box and the D-box, which confer specificity for ecdysone response elements. These domains may be either native, modified, or chimeras of different domains of heterologous receptor proteins. The EcR receptor, like a subset of the steroid receptor family, also possesses less well-defined regions responsible for heterodimerization properties. Because the domains of nuclear receptors are modular in nature, the LBD, DBD, and transactivation domains may be interchanged.

Gene switch systems are known that incorporate components from the ecdysone receptor complex. However, in these known systems, whenever EcR is used it is associated with native or modified DNA binding domains and transactivation domains on the same molecule. USP or RXR are typically used as silent partners. Applicants have previously shown that when DNA binding domains and transactivation domains are on the same molecule the background activity in the absence of ligand is high and that such activity is dramatically reduced when DNA binding domains and transactivation domains are on different molecules, that is, on each of two partners of a heterodimeric or homodimeric complex (see PCT/US01/09050).

The insect ecdysone receptor (EcR) heterodimerizes with Ultraspiracle (USP), the insect homologue of the mammalian RXR, and binds ecdysteroids and ecdysone receptor response elements and activates transcription of ecdysone responsive genes (Riddiford et al. 2000, *Vitam Horm*, 60:1-73). The EcR/USP/ligand complexes play important roles during insect development and reproduction. The EcR is a member of the steroid hormone receptor superfamily and has five modular domains, A/B (transactivation), C (DNA binding, heterodimerization), D (Hinge, heterodimerization), E (ligand binding, heterodimerization and transactivation and in some cases, F (transactivation), domains. Some of these domains such as A/B, C and E retain their function when they are fused to other proteins.

Recently, ecdysone receptor based gene expression systems have been developed. Tightly regulated inducible gene expression systems or “gene switches” are useful for various applications such as gene therapy, large scale production of proteins in cells, cell based high throughput screening assays, functional genomics and regulation of traits in transgenic plants and animals. U.S. Pat. No. 6,265,173 B1 discloses that various members of the steroid/thyroid superfamily of receptors can combine with *Drosophila melanogaster* ultraspiracle receptor (USP) or fragments thereof comprising at least the dimerization domain of USP for use in a gene expression system. U.S. Pat. No. 5,880,333 discloses a *Drosophila melanogaster* EcR and ultraspiracle (USP) heterodimer system used in plants in which the transactivation domain and the DNA binding domain are positioned on two different hybrid proteins.

The first version of an EcR-based gene switch used *Drosophila melanogaster* EcR (DmEcR) and *Mus musculus* RXR (MmRXR) and showed that these receptors in the presence of steroid, ponasteroneA, transactivate reporter genes in mam-

1996, *Proc. Natl. Acad. Sci.* 93: 3346-3351). Later, Suhr et al. (1998, *Proc. Natl. Acad. Sci. U.S.A.* 95: 7999-8004) showed that non-steroidal ecdysone agonist, tefubenozone, induced high level of transactivation of reporter genes in mammalian cells through *Bombyx mori* EcR (BmEcR) in the absence of exogenous heterodimer partner.

International Patent Applications Nos. PCT/US97/05330 (WO 97/38117) and PCT/US99/08381 (WO99/58155) disclose methods for modulating the expression of an exogenous gene in which a DNA construct comprising the exogenous gene and an ecdysone response element is activated by a second DNA construct comprising an ecdysone receptor that, in the presence of a ligand therefor, and optionally in the presence of a receptor capable of acting as a silent partner, binds to the ecdysone response element to induce gene expression. The ecdysone receptor of choice was isolated from *Drosophila melanogaster*. Typically, such systems require the presence of the silent partner, preferably retinoid X receptor (RXR), in order to provide optimum activation. In mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and regulates expression of target genes in a ligand dependent manner. International Patent Application No. PCT/US98/14215 (WO 99/02683) discloses that the ecdysone receptor isolated from the silk moth *Bombyx mori* is functional in mammalian systems without the need for an exogenous dimer partner.

Unfortunately, these USP-based systems are constitutive in animal cells and therefore, are not effective for regulating reporter gene expression. Drawbacks of the above described EcR-based gene regulation systems include a considerable background activity in the absence of ligands and non-applicability of these systems for use in both plants and animals (see U.S. Pat. No. 5,880,333).

Recently, an improved ecdysone receptor-based inducible gene expression system has been developed in which the transactivation and DNA binding domains are separated from each other by placing them on two different proteins results in greatly reduced background activity in the absence of a ligand and significantly increased activity over background in the presence of a ligand (pending application PCT/US01/09050, incorporated herein in its entirety by reference). This two-hybrid system is a significantly improved inducible gene expression modulation system compared to the two systems disclosed in applications PCT/US97/05330 and PCT/US98/14215. The two-hybrid system exploits the ability of a pair of interacting proteins to bring the transcription activation domain into a more favorable position relative to the DNA binding domain such that when the DNA binding domain binds to the DNA binding site on the gene, the transactivation domain more effectively activates the promoter (see, for example, U.S. Pat. No. 5,283,173). Briefly, the two-hybrid gene expression system comprises two gene expression cassettes; the first encoding a DNA binding domain fused to a nuclear receptor polypeptide, and the second encoding a transactivation domain fused to a different nuclear receptor polypeptide. In the presence of ligand, the interaction of the first polypeptide with the second polypeptide effectively tethers the DNA binding domain to the transactivation domain. Since the DNA binding and transactivation domains reside on two different molecules, the background activity in the absence of ligand is greatly reduced.

A two-hybrid system also provides improved sensitivity to non-steroidal ligands for example, diacylhydrazines, when compared to steroidal ligands for example, ponasterone A (“PonA”) or muristerone A (“MurA”). That is, when compared to steroids, the non-steroidal ligands provide higher activity at a lower concentration. In addition, since transacti-

vation based on EcR gene switches is often cell-line dependent, it is easier to tailor switching systems to obtain maximum transactivation capability for each application. Furthermore, the two-hybrid system avoids some side effects due to overexpression of RXR that often occur when unmodified RXR is used as a switching partner. In a preferred two-hybrid system, native DNA binding and transactivation domains of EcR or RXR are eliminated and as a result, these hybrid molecules have less chance of interacting with other steroid hormone receptors present in the cell resulting in reduced side effects.

Applicants have now obtained and determined the full length coding sequence of an additional homopteran EcR polynucleotide for use in methods of modulating gene expression in a host cell and methods of identifying molecules that modulate activity of leafhopper EcR. As described herein, Applicants' invention provides novel leafhopper ecdysone receptor polypeptides and novel polynucleotides encoding these polypeptides that are useful as components of gene expression systems for highly specific regulation of recombinant proteins in host cells or in methods for identifying new molecules which may act as agonists or antagonists of a homopteran insect ecdysone receptor.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1: Transactivation of reporter genes through VP16/NcEcR-CDE construct transfected into L57 cells or CBW cells along with 5xEcRElacZ and pFREcRE by 20E or RH-123709. The numbers on top of the bars indicate fold increase over DMSO levels.

FIG. 2: Transactivation of reporter genes through GAL4/NcEcR-DE construct transfected into NIH3T3 cells along with VP16/CfUSP-EF, VP16/DmUSP-EF, VP16/MmRXR α -EF, VP16/MmRXR α /LmUSP-EF chimera, VP16/AmaRXR1-EF, or VP16/AmaRX2-EF, and pFRLuc by PonA or GSTTM-E. The numbers on top of the bars indicate the maximum fold induction for that group.

DETAILED DESCRIPTION OF THE INVENTION

The present invention advantageously provides an isolated polynucleotide encoding a novel leafhopper ecdysone receptor polypeptide. The polynucleotides and polypeptides of the present invention are useful in methods to regulate gene expression of a polypeptide of interest in a host cell and in identifying new molecules that modulate activity of a leafhopper EcR.

The various aspects of the invention will be set forth in greater detail in the following sections, directed to the nucleic acids, polypeptides, vectors, antibodies, compositions, and methods of use of the invention. This organization into various sections is intended to facilitate understanding of the invention, and is in no way intended to be limiting thereof.

Definitions

The following defined terms are used throughout the present specification, and should be helpful in understanding the scope and practice of the present invention.

In a specific embodiment, the term "about" or "approximately" means within 20%, preferably within 10%, more preferably within 5%, and even more preferably within 1% of a given value or range.

The term "substantially free" means that a composition comprising "A" (where "A" is a single protein, DNA molecule, vector, recombinant host cell, etc.) is substantially free of "B" (where "B" comprises one or more contaminating

proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight. It is also preferred that a composition, which is substantially free of contamination, contain only a single molecular weight species having the activity or characteristic of the species of interest.

The term "isolated" for the purposes of the present invention designates a biological material (nucleic acid or protein) that has been removed from its original environment (the environment in which it is naturally present). For example, a polynucleotide present in the natural state in a plant or an animal is not isolated, however the same polynucleotide separated from the adjacent nucleic acids in which it is naturally present, is considered "isolated". The term "purified" does not require the material to be present in a form exhibiting absolute purity, exclusive of the presence of other compounds. It is rather a relative definition.

A polynucleotide is in the "purified" state after purification of the starting material or of the natural material by at least one order of magnitude, preferably 2 or 3 and preferably 4 or 5 orders of magnitude.

As used herein, the term "substantially pure" describes a polypeptide or other material which has been separated from its native contaminants. Typically, a monomeric polypeptide is substantially pure when at least about 60 to 75% of a sample exhibits a single polypeptide backbone. Minor variants or chemical modifications typically share the same polypeptide sequence. Usually a substantially pure polypeptide will comprise over about 85 to 90% of a polypeptide sample, and preferably will be over about 99% pure. Normally, purity is measured on a polyacrylamide gel, with homogeneity determined by staining. Alternatively, for certain purposes high resolution will be necessary and HPLC or a similar means for purification will be used. For most purposes, a simple chromatography column or polyacrylamide gel will be used to determine purity.

The term "substantially free of naturally-associated host cell components" describes a polypeptide or other material which is separated from the native contaminants which accompany it in its natural host cell state. Thus, a polypeptide which is chemically synthesized or synthesized in a cellular system different from the host cell from which it naturally originates will be free from its naturally-associated host cell components.

The terms "nucleic acid" or "polynucleotide" are used interchangeably herein to refer to a polymeric compound comprised of covalently linked subunits called nucleotides. Nucleic acid includes polyribonucleic acid (RNA) and polydeoxyribonucleic acid (DNA), both of which may be single-stranded or double-stranded. DNA includes but is not limited to cDNA, genomic DNA, plasmids DNA, synthetic DNA, and semi-synthetic DNA. DNA may be linear, circular, or supercoiled.

A "nucleic acid molecule" refers to the phosphate ester polymeric form of ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it

to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear or circular DNA molecules (e.g., restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the non-transcribed strand of DNA (i.e., the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

The term "fragment" will be understood to mean a nucleotide sequence of reduced length relative to the reference nucleic acid and comprising, over the common portion, a nucleotide sequence identical to the reference nucleic acid. Such a nucleic acid fragment according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. Such fragments comprise, or alternatively consist of, oligonucleotides ranging in length from at least 6-1100 consecutive nucleotides of a nucleic acid according to the invention.

As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA.

A "gene" refers to an assembly of nucleotides that encode a polypeptide, and includes cDNA and genomic DNA nucleic acids. "Gene" also refers to a nucleic acid fragment that expresses a specific protein or polypeptide, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and/or coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. A chimeric gene may comprise coding sequences derived from different sources and/or regulatory sequences derived from different sources. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene or "heterologous" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

"Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

The term "genome" includes chromosomal as well as mitochondrial, chloroplast and viral DNA or RNA.

A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength (see Sambrook et al., 1989 *infra*). Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly

Chapter 11 and Table 11.1 therein (entirely incorporated herein by reference). The conditions of temperature and ionic strength determine the "stringency" of the hybridization.

Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a T_m of 55°, can be used, e.g., 5×SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5×SSC, 0.5% SDS). Moderate stringency hybridization conditions correspond to a higher T_m , e.g., 40% formamide, with 5× or 6×SSC. High stringency hybridization conditions correspond to the highest T_m , e.g., 50% formamide, 5× or 6×SSC.

Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The term "complementary" is used to describe the relationship between nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA, adenosine is complementary to thymine and cytosine is complementary to guanine. Accordingly, the instant invention also includes isolated nucleic acid fragments that are complementary to the complete sequences as disclosed or used herein as well as those substantially similar nucleic acid sequences.

In a specific embodiment of the invention, polynucleotides are detected by employing hybridization conditions comprising a hybridization step at T_m of 55° C., and utilizing conditions as set forth above. In a preferred embodiment, the T_m is 60° C.; in a more preferred embodiment, the T_m is 63° C.; in an even more preferred embodiment, the T_m is 65° C.

Post-hybridization washes also determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6×SSC, 0.5% SDS at room temperature for 15 minutes (min), then repeated with 2×SSC, 0.5% SDS at 45° C. for 30 minutes, and then repeated twice with 0.2×SSC, 0.5% SDS at 50° C. for 30 minutes. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2×SSC, 0.5% SDS was increased to 60° C. Another preferred set of highly stringent conditions uses two final washes in 0.1×SSC, 0.1% SDS at 65° C.

The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., *supra*, 9.50-0.5 1). For hybridization with shorter nucleic acids, i.e., oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., *supra*, 11.7-11.8).

Selectivity of hybridization exists when hybridization occurs which is more selective than total lack of specificity. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14/25 nucleotides, preferably at least about 65%, more preferably at least about 75%, and most preferably at least about

90%. See, Kanehisa, M. (1984), *Nucleic Acids Res.* 12:203-213, which is incorporated herein by reference. Stringent hybridization conditions will typically include salt concentrations of less than about 1 M, more usually less than about 500 mM and preferably less than about 200 mM. Temperature conditions will typically be greater than 20 degrees Celsius, more usually greater than about 30 degrees Celsius and preferably in excess of about 37 degrees Celsius. As other factors may significantly affect the stringency of hybridization, including, among others, base composition and size of the complementary strands, presence of organic solvents and extent of base mismatching, the combination of parameters is more important than the absolute measure of any one.

In a specific embodiment of the invention, polynucleotides of the invention are detected by employing hybridization conditions comprising a hybridization step in less than 500 mM salt and at least 37 degrees Celsius, and a washing step in 2×SSPE at least 63 degrees Celsius. In a preferred embodiment, the hybridization conditions comprise less than 200 mM salt and at least 37 degrees Celsius for the hybridization step. In a more preferred embodiment, the hybridization conditions comprise 2×SSPE and 63 degrees Celsius for both the hybridization and washing steps.

In one embodiment, the length for a hybridizable nucleic acid is at least about 10 nucleotides. Preferable a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably at least about 20 nucleotides; and even more preferably the length is at least 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

The term “probe” refers to a single-stranded nucleic acid molecule that can base pair with a complementary single stranded target nucleic acid to form a double-stranded molecule.

As used herein, the term “oligonucleotide” refers to a nucleic acid, generally of at least 18 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, a plasmid DNA or an mRNA molecule. Oligonucleotides can be labeled, e.g., with ³²P-nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. A labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid. Oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of a nucleic acid, or to detect the presence of a nucleic acid. An oligonucleotide can also be used to form a triple helix with a DNA molecule. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer. Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

A “primer” is an oligonucleotide that hybridizes to a target nucleic acid sequence to create a double stranded nucleic acid region that can serve as an initiation point for DNA synthesis under suitable conditions. Such primers may be used in a polymerase chain reaction.

“Polymerase chain reaction” is abbreviated PCR and means an in vitro method for enzymatically amplifying specific nucleic acid sequences. PCR involves a repetitive series of temperature cycles with each cycle comprising three stages: denaturation of the template nucleic acid to separate the strands of the target molecule, annealing a single stranded PCR oligonucleotide primer to the template nucleic acid, and extension of the annealed primer(s) by DNA polymerase. PCR provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative condi-

tions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

“Reverse transcription-polymerase chain reaction” is abbreviated RT-PCR and means an in vitro method for enzymatically producing a target cDNA molecule or molecules from an RNA molecule or molecules, followed by enzymatic amplification of a specific nucleic acid sequence or sequences within the target cDNA molecule or molecules as described above. RT-PCR also provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

A DNA “coding sequence” is a double-stranded DNA sequence that is transcribed and translated into a polypeptide in a cell in vitro or in vivo when placed under the control of appropriate regulatory sequences. “Suitable regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, polyadenylation recognition sequences, RNA processing site, effector binding site and stem-loop structure. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from mRNA, genomic DNA sequences, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

“Open reading frame” is abbreviated ORF and means a length of nucleic acid sequence, either DNA, cDNA or RNA, that comprises a translation start signal or initiation codon, such as an ATG or AUG, and a termination codon and can be potentially translated into a polypeptide sequence.

The term “head-to-head” is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-head orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 5' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds away from the 5' end of the other polynucleotide. The term “head-to-head” may be abbreviated (5')-to-(5') and may also be indicated by the symbols (←→) or (3'←5'→3').

The term “tail-to-tail” is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a tail-to-tail orientation when the 3' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds toward the other polynucleotide. The term “tail-to-tail” may be abbreviated (3')-to-(3') and may also be indicated by the symbols (→←) or (5'→3'3'←5').

The term “head-to-tail” is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-tail orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds in the same direction as that

of the other polynucleotide. The term "head-to-tail" may be abbreviated (5')-to-(3') and may also be indicated by the symbols ($\rightarrow\rightarrow$) or ($5'\rightarrow 3'5'\rightarrow 3'$).

The term "downstream" refers to a nucleotide sequence that is located 3' to reference nucleotide sequence. In particular, downstream nucleotide sequences generally relate to sequences that follow the starting point of transcription. For example, the translation initiation codon of a gene is located downstream of the start site of transcription.

The term "upstream" refers to a nucleotide sequence that is located 5' to reference nucleotide sequence. In particular, upstream nucleotide sequences generally relate to sequences that are located on the 5' side of a coding sequence or starting point of transcription. For example, most promoters are located upstream of the start site of transcription.

The terms "restriction endonuclease" and "restriction enzyme" refer to an enzyme that binds and cuts within a specific nucleotide sequence within double stranded DNA.

"Homologous recombination" refers to the insertion of a foreign DNA sequence into another DNA molecule, e.g., insertion of a vector in a chromosome. Preferably, the vector targets a specific chromosomal site for homologous recombination. For specific homologous recombination, the vector will contain sufficiently long regions of homology to sequences of the chromosome to allow complementary binding and incorporation of the vector into the chromosome. Longer regions of homology, and greater degrees of sequence similarity, may increase the efficiency of homologous recombination.

Several methods known in the art may be used to propagate a polynucleotide according to the invention. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As described herein, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (e.g., lambda), and plasmid and cosmid DNA vectors, to name but a few.

A "vector" is any means for the cloning of and/or transfer of a nucleic acid into a host cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (e.g., plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, i.e., capable of replication under its own control. The term "vector" includes both viral and nonviral means for introducing the nucleic acid into a cell *in vitro*, *ex vivo* or *in vivo*. A large number of vectors known in the art may be used to manipulate nucleic acids, incorporate response elements and promoters into genes, etc. Possible vectors include, for example, plasmids or modified viruses including, for example bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives, or the Bluescript vector. For example, the insertion of the DNA fragments corresponding to response elements and promoters into a suitable vector can be accomplished by ligating the appropriate DNA fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells that have incorporated the marker into the cellular genome. Such markers allow identification and/or selection of host cells that incorporate and express the proteins encoded by the marker.

Viral vectors, and particularly retroviral vectors, have been used in a wide variety of gene delivery applications in cells, as well as living animal subjects. Viral vectors that can be used include but are not limited to retrovirus, adeno-associated virus, pox, baculovirus, vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors. Non-viral vectors include plasmids, liposomes, electrically charged lipids (cytofectins), DNA-protein complexes, and biopolymers. In addition to a nucleic acid, a vector may also comprise one or more regulatory regions, and/or selectable markers useful in selecting, measuring, and monitoring nucleic acid transfer results (transfer to which tissues, duration of expression, etc.).

The term "plasmid" refers to an extra chromosomal element often carrying a gene that is not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome integrating sequences, phage or nucleotide sequences, linear, circular, or supercoiled, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

A "cloning vector" is a "replicon", which is a unit length of a nucleic acid, preferably DNA, that replicates sequentially and which comprises an origin of replication, such as a plasmid, phage or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. Cloning vectors may be capable of replication in one cell type and expression in another ("shuttle vector").

Vectors may be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), particle bombardment, use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, *J. Biol. Chem.* 267:963-967; Wu and Wu, 1988, *J. Biol. Chem.* 263:14621-14624; and Hartmut et al., Canadian Patent Application No. 2,012,311, filed Mar. 15, 1990).

A polynucleotide according to the invention can also be introduced *in vivo* by lipofection. For the past decade, there has been increasing use of liposomes for encapsulation and transfection of nucleic acids *in vitro*. Synthetic cationic lipids designed to limit the difficulties and dangers encountered with liposome-mediated transfection can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner et al., 1987, *Proc. Natl. Acad. Sci. U.S.A.* 84: 7413; Mackey, et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85: 8027-8031; and Ulmer et al., 1993, *Science* 259: 1745-1748). The use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, 1989, *Science* 337:387-388). Particularly useful lipid compounds and compositions for transfer of nucleic acids are described in International Patent Publications WO95/18863 and WO96/17823, and in U.S. Pat. No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. It is clear that directing transfection to particular cell types would be particularly preferred in a tissue with cellular heterogeneity, such as pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting (Mackey, et al., 1988, *supra*). Targeted

peptides, e.g., hormones or neurotransmitters, and proteins such as antibodies, or non-peptide molecules could be coupled to liposomes chemically.

Other molecules are also useful for facilitating transfection of a nucleic acid in vivo, such as a cationic oligopeptide (e.g., WO95/21931), peptides derived from DNA binding proteins (e.g., WO96/25508), or a cationic polymer (e.g., WO95/21931).

It is also possible to introduce a vector in vivo as a naked DNA plasmid (see U.S. Pat. Nos. 5,693,622, 5,589,466 and 5,580,859). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., 1992, *Hum. Gene Ther.* 3:147-154; and Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432).

The term "transfection" means the uptake of exogenous or heterologous RNA or DNA by a cell. A cell has been "transfected" by exogenous or heterologous RNA or DNA when such RNA or DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous RNA or DNA when the transfected RNA or DNA effects a phenotypic change. The transforming RNA or DNA can be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or "recombinant" or "transformed" organisms.

The term "genetic region" will refer to a region of a nucleic acid molecule or a nucleotide sequence that comprises a gene encoding a polypeptide.

In addition, the recombinant vector comprising a polynucleotide according to the invention may include one or more origins for replication in the cellular hosts in which their amplification or their expression is sought, markers or selectable markers.

The term "selectable marker" means an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene's effect, i.e., resistance to an antibiotic, resistance to a herbicide, colorimetric markers, enzymes, fluorescent markers, and the like, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic acid of interest. Examples of selectable marker genes known and used in the art include: genes providing resistance to ampicillin, streptomycin, gentamycin, kanamycin, hygromycin, bialaphos herbicide, sulfonamide, and the like; and genes that are used as phenotypic markers, i.e., anthocyanin regulatory genes, isopentenyl transferase gene, and the like. Selectable marker genes may also be considered reporter genes.

The term "reporter gene" means a nucleic acid encoding an identifying factor that is able to be identified based upon the reporter gene's effect, wherein the effect is used to track the inheritance of a nucleic acid of interest, to identify a cell or organism that has inherited the nucleic acid of interest, and/or to measure gene expression induction or transcription. Examples of reporter genes known and used in the art include: luciferase (Luc), green fluorescent protein (GFP), chloramphenicol acetyltransferase (CAT), β -galactosidase (LacZ), β -glucuronidase (Gus), and the like.

"Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in

the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters that cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". Promoters that cause a gene to be expressed in a specific cell type are commonly referred to as "cell-specific promoters" or "tissue-specific promoters". Promoters that cause a gene to be expressed at a specific stage of development or cell differentiation are commonly referred to as "developmentally-specific promoters" or "cell differentiation-specific promoters". Promoters that are induced and cause a gene to be expressed following exposure or treatment of the cell with an agent, biological molecule, chemical, ligand, light, or the like that induces the promoter are commonly referred to as "inducible promoters" or "regulatable promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if the coding sequence contains introns) and translated into the protein encoded by the coding sequence.

"Transcriptional and translational control sequences" are DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

The term "response element" means one or more cis-acting DNA elements which confer responsiveness on a promoter mediated through interaction with the DNA-binding domains of the first chimeric gene. This DNA element may be either palindromic (perfect or imperfect) in its sequence or composed of sequence motifs or half sites separated by a variable number of nucleotides. The half sites can be similar or identical and arranged as either direct or inverted repeats or as a single half site or multimers of adjacent half sites in tandem. The response element may comprise a minimal promoter isolated from different organisms depending upon the nature of the cell or organism into which the response element will be incorporated. The DNA binding domain of the first hybrid protein binds, in the presence or absence of a ligand, to the DNA sequence of a response element to initiate or suppress transcription of downstream gene(s) under the regulation of this response element. Examples of DNA sequences for response elements of the natural ecdysone receptor include: RRGG/TTCANTGAC/ACY (see Cherbas L., et al., (1991), *Genes Dev.* 5: 120-131); AGGTCAN_(n)AGGTC, where N_(n) can be one or more spacer nucleotides (see D'Avino P.P., et al., (1995), *Mol. Cell. Endocrinol.* 113: 1-9);

and GGGTTGAATGAATT (see Antoniewski C., et. al., (1994), Mol. Cell Biol. 14: 4465-4474).

The term “operably linked” refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term “expression”, as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from a nucleic acid or polynucleotide. Expression may also refer to translation of mRNA into a protein or polypeptide.

The terms “cassette”, “expression cassette” and “gene expression cassette” refer to a segment of DNA that can be inserted into a nucleic acid or polynucleotide at specific restriction sites or by homologous recombination. The segment of DNA comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation. “Transformation cassette” refers to a specific vector comprising a polynucleotide that encodes a polypeptide of interest and having elements in addition to the polynucleotide that facilitate transformation of a particular host cell. Cassettes, expression cassettes, gene expression cassettes and transformation cassettes of the invention may also comprise elements that allow for enhanced expression of a polynucleotide encoding a polypeptide of interest in a host cell. These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

For purposes of this invention, the term “gene switch” refers to the combination of a response element associated with a promoter, and an EcR-based gene expression system which, in the presence of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated.

The terms “modulate” and “modulates” mean to induce, reduce or inhibit nucleic acid or gene expression, resulting in the respective induction, reduction or inhibition of protein or polypeptide production.

The plasmids or vectors according to the invention may further comprise at least one promoter suitable for driving expression of a gene in a host cell. The term “expression vector” means a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, i.e., the inserted nucleic acid sequence, is usually placed under the control of control elements such as a promoter, a minimal promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to: viral promoters, bacterial promoters, animal promoters, mammalian promoters, synthetic promoters, constitutive promoters, tissue specific promoter, developmental specific promoters, inducible promoters, light regulated promoters; CYC1, HIS3, GAL1, GAL4, GAL10, ADH1, PGK, PHO5, GAPDH, ADC1, TRP1, URA3, LEU2, ENO, TPI, alkaline phosphatase promoters (useful for expression in *Saccharomyces*); AOX1 promoter (useful for expression in *Pichia*); b-lactamase, lac, ara, tet, trp, IP_L , IP_R , T7, tac, and trc

promoters (useful for expression in *Escherichia coli*); light regulated-, seed specific-, pollen specific-, ovary specific-, pathogenesis or disease related-, cauliflower mosaic virus 35S, CMV 35S minimal, cassava vein mosaic virus (Cs-VMV), chlorophyll a/b binding protein, ribulose 1,5-bisphosphate carboxylase, shoot-specific, root specific, chitinase, stress inducible, rice tungro bacilliform virus, plant super-promoter, potato leucine aminopeptidase, nitrate reductase, mannopine synthase, nopaline synthase, ubiquitin, zein protein, and anthocyanin promoters (useful for expression in plant cells); animal and mammalian promoters known in the art include, but are not limited to, the SV40 early (SV40e) promoter region, the promoter contained in the 3' long terminal repeat (LTR) of Rous sarcoma virus (RSV), the promoters of the E1A or major late promoter (MLP) genes of adenoviruses (Ad), the cytomegalovirus (CMV) early promoter, the herpes simplex virus (HSV) thymidine kinase (TK) promoter, a baculovirus IE1 promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, an albumin promoter, the regulatory sequences of the mouse metallothionein-L promoter and transcriptional control regions, the ubiquitous promoters (HPRT, vimentin, α -actin, tubulin and the like), the promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like), the promoters of therapeutic genes (of the MDR, CFTR or factor VIII type, and the like), pathogenesis or disease related-promoters, and promoters that exhibit tissue specificity and have been utilized in transgenic animals, such as the elastase I gene control region which is active in pancreatic acinar cells; insulin gene control region active in pancreatic beta cells, immunoglobulin gene control region active in lymphoid cells, mouse mammary tumor virus control region active in testicular, breast, lymphoid and mast cells; albumin gene, Apo AI and Apo AII control regions active in liver, alpha-fetoprotein gene control region active in liver, alpha 1-antitrypsin gene control region active in the liver, beta-globin gene control region active in myeloid cells, myelin basic protein gene control region active in oligodendrocyte cells in the brain, myosin light chain-2 gene control region active in skeletal muscle, and gonadotropic releasing hormone gene control region active in the hypothalamus, pyruvate kinase promoter, villin promoter, promoter of the fatty acid binding intestinal protein, promoter of the smooth muscle cell α -actin, and the like. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

Enhancers that may be used in embodiments of the invention include but are not limited to: an SV40 enhancer, a cytomegalovirus (CMV) enhancer, an elongation factor 1 (EF1) enhancer, yeast enhancers, viral gene enhancers, and the like.

Termination control regions, i.e., terminator or polyadenylation sequences, may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary, however, it is most preferred if included. In a preferred embodiment of the invention, the termination control region may be comprise or be derived from a synthetic sequence, synthetic polyadenylation signal, an SV40 late polyadenylation signal, an SV40 polyadenylation signal, a bovine growth hormone (BGH) polyadenylation signal, viral terminator sequences, or the like.

The terms “3' non-coding sequences” or “3' untranslated region (UTR)” refer to DNA sequences located downstream (3') of a coding sequence and may comprise polyadenylation [poly(A)] recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually


characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.

"Regulatory region" means a nucleic acid sequence that regulates the expression of a second nucleic acid sequence. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region) or may include sequences of a different origin that are responsible for expressing different proteins or even synthetic proteins (a heterologous region). In particular, the sequences can be sequences of prokaryotic, eukaryotic, or viral genes or derived sequences that stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, promoters, enhancers, transcriptional termination sequences, and signal sequences which direct the polypeptide into the secretory pathways of the target cell.

A regulatory region from a "heterologous source" is a regulatory region that is not naturally associated with the expressed nucleic acid. Included among the heterologous regulatory regions are regulatory regions from a different species, regulatory regions from a different gene, hybrid regulatory sequences, and regulatory sequences which do not occur in nature, but which are designed by one having ordinary skill in the art.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that is not translated yet has an effect on cellular processes.

A "polypeptide" is a polymeric compound comprised of covalently linked amino acid residues. Amino acids have the following general structure:



Amino acids are classified into seven groups on the basis of the side chain R: (1) aliphatic side chains, (2) side chains containing a hydroxylic (OH) group, (3) side chains containing sulfur atoms, (4) side chains containing an acidic or amide group, (5) side chains containing a basic group, (6) side chains containing an aromatic ring, and (7) proline, an imino acid in which the side chain is fused to the amino group. A polypeptide of the invention preferably comprises at least about 14 amino acids.

A "protein" is a polypeptide that performs a structural or functional role in a living cell.

An "isolated polypeptide" or "isolated protein" is a polypeptide or protein that is substantially free of those compounds that are normally associated therewith in its natural state (e.g., other proteins or polypeptides, nucleic acids, carbohydrates, lipids). "Isolated" is not meant to exclude artificial or synthetic mixtures with other compounds, or the presence of impurities which do not interfere with biological activity, and which may be present, for example, due to incomplete purification, addition of stabilizers, or compounding into a pharmaceutically acceptable preparation.

A "fragment" of a polypeptide according to the invention will be understood to mean a polypeptide whose amino acid sequence is shorter than that of the reference polypeptide and which comprises, over the entire portion with these reference polypeptides, an identical amino acid sequence. Such fragments may, where appropriate, be included in a larger polypeptide of which they are a part. Such fragments of a polypeptide according to the invention may have a length of at least 2-300 amino acids.

A "heterologous protein" refers to a protein not naturally produced in the cell.

A "mature protein" refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

The term "signal peptide" refers to an amino terminal polypeptide preceding the secreted mature protein. The signal peptide is cleaved from and is therefore not present in the mature protein. Signal peptides have the function of directing and translocating secreted proteins across cell membranes. Signal peptide is also referred to as signal protein.

A "signal sequence" is included at the beginning of the coding sequence of a protein to be expressed on the surface of a cell. This sequence encodes a signal peptide, N-terminal to the mature polypeptide, that directs the host cell to translocate the polypeptide. The term "translocation signal sequence" is used herein to refer to this sort of signal sequence. Translocation signal sequences can be found associated with a variety of proteins native to eukaryotes and prokaryotes, and are often functional in both types of organisms.

The term "homology" refers to the percent of identity between two polynucleotide or two polypeptide moieties. The correspondence between the sequence from one moiety to another can be determined by techniques known to the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules by aligning the sequence information and using readily available computer programs. Alternatively, homology can be determined by hybridization of polynucleotides under conditions that form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s) and size determination of the digested fragments.

As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (e.g., the immunoglobulin superfamily) and homologous proteins from different species (e.g., myosin light chain, etc.) (Reeck et al., 1987, Cell 50: 667.). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity. However, in common usage and in the instant application, the term "homologous," when modified

with an adverb such as “highly,” may refer to sequence similarity and not a common evolutionary origin.

Accordingly, the term “sequence similarity” in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that may or may not share a common evolutionary origin (see Reeck et al., 1987, *Cell* 50: 667).

In a specific embodiment, two DNA sequences are “substantially homologous” or “substantially similar” when at least about 50% (preferably at least about 75%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., 1989, *supra*.

As used herein, “substantially similar” refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence. “Substantially similar” also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. “Substantially similar” also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotide bases that do not substantially affect the functional properties of the resulting transcript. It is therefore understood that the invention encompasses more than the specific exemplary sequences. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

Moreover, the skilled artisan recognizes that substantially similar sequences encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1×SSC, 0.1% SDS, 65° C. and washed with 2×SSC, 0.1% SDS followed by 0.1×SSC, 0.1% SDS), with the sequences exemplified herein. Substantially similar nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 70% identical to the DNA sequence of the nucleic acid fragments reported herein. Preferred substantially nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 80% identical to the DNA sequence of the nucleic acid fragments reported herein. More preferred nucleic acid fragments are at least 90% identical to the DNA sequence of the nucleic acid fragments reported herein. Even more preferred are nucleic acid fragments that are at least 95% identical to the DNA sequence of the nucleic acid fragments reported herein.

Two amino acid sequences are “substantially homologous” or “substantially similar” when greater than about 40% of the amino acids are identical, or greater than 60% are similar (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, Version 7, Madison, Wis.) pileup program.

The term “corresponding to” is used herein to refer to similar or homologous sequences, whether the exact position is identical or different from the molecule to which the similarity or homology is measured. A nucleic acid or amino acid

sequence alignment may include spaces. Thus, the term “corresponding to” refers to the sequence similarity, and not the numbering of the amino acid residues or nucleotide bases.

A “substantial portion” of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to putatively identify that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993), *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., in situ hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a “substantial portion” of a nucleotide sequence comprises enough of the sequence to specifically identify and/or isolate a nucleic acid fragment comprising the sequence.

The term “percent identity”, as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, “identity” also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. “Identity” and “similarity” can be readily calculated by known methods, including but not limited to those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects* (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991). Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity calculations may be performed using the Megalign program of the LASER-GENE bioinformatics computing suite (DNASTAR Inc., Madison, Wis.). Multiple alignment of the sequences may be performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*, 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method may be selected: KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

The term “sequence analysis software” refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. “Sequence analysis software” may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, Wis., BLASTP, BLASTN, BLASTX (Altschul et al., *J. Mol. Biol.* 215: 403-410 (1990), and DNASTAR

(DNASTAR, Inc. 1228 S. Park St. Madison, Wis. 53715 USA). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters which originally load with the software when first initialized.

"Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments that are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled in vitro. Manual chemical synthesis of DNA may be accomplished using well-established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

Polynucleotides Encoding Leafhopper Ecdysone Receptor Polypeptides

The present invention provides novel polynucleotides encoding a leafhopper ecdysone receptor polypeptide of the invention, including a full-length leafhopper ecdysone receptor protein, and any leafhopper ecdysone receptor-specific fragments thereof.

In accordance with specific embodiments of the present invention, nucleic acid sequences encoding portions of a novel ecdysone receptor polypeptide have been elucidated and characterized. Specifically, polynucleotides encoding a homopteran ecdysone receptor from green leafhopper (NcEcR) have been characterized. The full-length encoding sequence has been determined and is presented herein as SEQ ID NO: 1. In addition, domains within this polynucleotide encoding the full-length NcEcR polypeptide have been defined and are presented herein as described in Table 1.

TABLE 1

Nucleotide and amino acid sequences corresponding to various domains and helices of leafhopper ecdysone receptor ("NcEcR").		
Full Length NcEcR or NcEcR Domains	Nucleotides of SEQ ID NO: 1	Amino Acids of SEQ ID NO: 2
A/BCDE (Full length)	1-1109	1-370
A/B	1-15	1-5
C	16-213	6-71
D	214-447	72-149
E	448-1109	150-370
CDE	16-1109	6-370
DE	214-1109	72-370
Helices 1-12	418-1109	140-370

Thus, a first subject of the invention relates to an isolated polynucleotide encoding a novel ecdysone receptor polypeptide. More specifically, the invention relates to an isolated polynucleotide encoding a leafhopper ecdysone receptor polypeptide. In a specific embodiment, the isolated polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 1-15 of SEQ ID NO: 1, nucleotides 16-213 of SEQ ID NO: 1, nucle-

otides 214-447 of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1. In another specific embodiment, the isolated polynucleotide comprises a nucleic acid sequence as depicted in SEQ ID NO: 1. In another specific embodiment, the isolated polynucleotide further comprises a region permitting expression of the polypeptide in a host cell.

The present invention also relates to an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 1-5 of SEQ ID NO: 2, amino acids 6-71 of SEQ ID NO: 2, amino acids 72-149 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. In a specific embodiment, the isolated polynucleotide encodes a leafhopper ecdysone receptor polypeptide comprising an amino acid sequence as depicted in SEQ ID NO: 2.

The present invention provides novel isolated polynucleotides encoding leafhopper ecdysone receptor polypeptides. Having elucidated the sequence and structure of this ecdysone receptor, an isolated polynucleotide encoding a leafhopper receptor polypeptide comprising a ligand-binding domain may be used individually or in combination to screen for new ligands that bind this ligand binding domain. Thus, for example, an ecdysone receptor polypeptide according to the invention may be used to control expression of reporter genes for which sensitive assays exist. The ligand binding domain may serve as a reagent for screening new molecules, useful as either agonists or antagonists of the leafhopper ecdysone receptor. Either new classes of molecules may be screened, or selected modifications from known ligands may be used. These new ligands find use as highly specific and highly active, naturally occurring pesticides. Thus, the present invention provides for screening for new ligand molecules.

The polynucleotides of the present invention also provide probes for screening for homologous nucleic acid sequences, both in *Nephotetix* and other genetic sources. This screening allows isolation of homologous genes from both vertebrates and invertebrates.

Accordingly, any leafhopper cell potentially can serve as the nucleic acid source for the molecular cloning of a leafhopper ecdysone receptor polynucleotide. The polynucleotide may be obtained by standard procedures known in the art from cloned DNA (e.g. a DNA "library"), and preferably is obtained from a cDNA library prepared from tissues with high level expression of the protein, by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (See, for example, Sambrook et al., 1989, *supra*; Glover, D. M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived from cDNA will not contain intron sequences. Whatever the source, the polynucleotide should be molecularly cloned into a suitable vector for propagation of the polynucleotide.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the desired leafhopper ecdysone receptor polynucleotide may be accomplished in a number of ways. For example, DNA fragments may be screened by nucleic acid hybridization to a labeled probe (Benton and Davis, 1977, *Science* 196: 180; Grunstein and Hogness, 1975, *Proc. Natl. Acad. Sci. U.S.A.* 72: 3961). Those DNA fragments with substantial homology to the

probe will hybridize. As noted above, the greater the degree of homology, the more stringent hybridization conditions can be used.

Further selection can be carried out on the basis of the properties of the polynucleotide, e.g., if the polynucleotide encodes a polypeptide having the isoelectric, electrophoretic, amino acid composition, or partial amino acid sequence of the leafhopper ecdysone receptor polypeptide as disclosed herein. Thus, the presence of the polynucleotide may be detected by assays based on the physical, chemical, or immunological properties of its expressed product. For example, cDNA clones, or DNA clones which hybrid-select the proper mRNAs, can be selected which produce a polypeptide that, e.g., has similar or identical electrophoretic migration, isoelectric focusing or non-equilibrium pH gel electrophoresis behavior, proteolytic digestion maps, or antigenic properties as known for a leafhopper ecdysone receptor polypeptide. In a specific embodiment, the expressed polypeptide is recognized by a polyclonal antibody that is generated against an epitope specific for a leafhopper ecdysone receptor polypeptide.

Due to the degeneracy of nucleotide coding sequences, other polynucleotides that encode substantially the same amino acid sequence as a leafhopper ecdysone receptor polynucleotide disclosed herein, including an amino acid sequence that contains a single amino acid variant, may be used in the practice of the present invention. These include but are not limited to allelic genes, homologous genes from other species, and nucleotide sequences comprising all or portions of leafhopper ecdysone receptor polynucleotides that are altered by the substitution of different codons that encode the same amino acid residue within the sequence, thus producing a silent change. Likewise, the leafhopper ecdysone receptor derivatives of the invention include, but are not limited to, those comprising, as a primary amino acid sequence, all or part of the amino acid sequence of a leafhopper ecdysone receptor polypeptide including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a conservative amino acid substitution. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations can be produced by various methods known in the art (see Sambrook et al., 1989, *infra*) and are not expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

The present invention also relates to an isolated leafhopper ecdysone receptor polypeptide encoded by a polynucleotide according to the invention.

Leafhopper Ecdysone Receptor Polypeptides

The present invention provides novel isolated leafhopper ecdysone receptor polypeptides, including a full-length leafhopper ecdysone receptor protein, and any leafhopper ecdysone receptor-specific polypeptide fragments thereof.

Thus, the invention relates to an isolated ecdysone receptor polypeptide. More specifically, the invention relates to an isolated leafhopper ecdysone receptor polypeptide. In a specific embodiment, the isolated ecdysone receptor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 1-5 of SEQ ID NO: 2, amino acids 6-71 of SEQ ID NO: 2, amino acids 72-149 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. In another specific embodiment, the isolated ecdysone receptor polypeptide comprises an amino acid sequence as depicted in SEQ ID NO: 2.

In another specific embodiment, the isolated ecdysone receptor polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 1-15 of SEQ ID NO: 1, nucleotides 16-213 of SEQ ID NO: 1, nucleotides 214-447 of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1. In another specific embodiment, the isolated ecdysone receptor polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence as depicted in SEQ ID NO: 1.

One of skill in the art is able to produce other polynucleotides to encode the polypeptides of the invention, by making use of the present invention and the degeneracy or non-universality of the genetic code as described herein.

Additional embodiments of the present invention include an ecdysone receptor polypeptide according to the invention, wherein the ecdysone receptor polypeptide is substantially free of naturally associated cell components. Such polypeptides will typically be either full-length proteins, functional fragments, or fusion proteins comprising segments from an ecdysone receptor polypeptide of the present invention fused to a heterologous, or normally non-contiguous, protein domain. Preferably, the ecdysone receptor polypeptide comprises a transactivation domain, a DNA binding domain, a ligand binding domain, a hinge region, or a heterodimerization domain. More preferably, the ecdysone receptor polypeptide comprises a ligand binding domain that is capable of binding to a ligand selected from the group consisting of a steroid ligand and a non-steroid ligand. As desired, the ecdysone receptor polypeptide may be fused to a second polypeptide to generate a hybrid polypeptide. Preferably, the second polypeptide is a heterologous polypeptide from the steroid hormone nuclear receptor superfamily.

Besides substantially full-length polypeptides, the present invention provides for biologically active fragments of the polypeptides. Significant biological activities include transactivation activity, ligand binding, DNA binding, heterodimerization activity, immunological activity and other biological activities characteristic of steroid receptor superfamily members. Immunological activities include both immunogenic function in a target immune system, as well as sharing of immunological epitopes for binding, serving as either a competitor or substitute antigen for an ecdysone receptor epitope.

For example, transactivation, ligand binding, or DNA-binding domains may be "swapped" between different new fusion polypeptides or fragments. Thus, novel hybrid polypeptides exhibiting new combinations of specificities result from the functional linkage of transactivation, ligand-binding specificities, or DNA-binding domains. This is extremely useful in the design of inducible expression systems.

For immunological purposes, immunogens may be produced that tandemly repeat polypeptide segments, thereby producing highly antigenic proteins. Alternatively, such polypeptides will serve as highly efficient competitors for specific binding. Production of antibodies to NcEcR is described below.

The present invention also provides for other polypeptides comprising fragments of NcEcR. Thus, fusion polypeptides between the NcEcR segments and other homologous or heterologous proteins are provided. Homologous polypeptides may be fusions between different steroid receptor superfamily members, resulting in, for instance, a hybrid protein exhibiting ligand specificity of one member and DNA-binding specificity of another. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with another domain of a receptor, e.g., a DNA-binding domain, so that the presence or location of a desired ligand may be easily determined. See, e.g., Dull et al., U.S. Pat. No. 4,859,609, which is hereby incorporated herein by reference. Other typical gene fusion partners include "zinc finger" segment swapping between DNA-binding proteins, bacterial beta-galactosidase, trpE Protein A, beta-lactamase, alpha amylase, alcohol dehydrogenase and yeast alpha mating factor. See, e.g., Godowski et al. (1988), *Science* 241: 812-816.

Thus, the present invention also provides an isolated polypeptide selected from the group consisting of a) an isolated polypeptide comprising a transactivation domain, a DNA-binding domain, and a leafhopper ecdysone receptor ligand binding domain; b) an isolated polypeptide comprising a DNA-binding domain and a leafhopper ecdysone receptor ligand binding domain; and c) an isolated polypeptide comprising a transactivation domain and a leafhopper ecdysone receptor ligand binding domain. Preferably, the leafhopper ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. In another preferred embodiment, the leafhopper ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1.

The present invention also provides an isolated hybrid polypeptide selected from the group consisting of a) an isolated hybrid polypeptide comprising a transactivation domain, a DNA-binding domain, and a leafhopper ecdysone receptor ligand binding domain; b) an isolated hybrid polypeptide comprising a DNA-binding domain and a leafhopper ecdysone receptor ligand binding domain; and c) an isolated hybrid polypeptide comprising a transactivation domain and a leafhopper ecdysone receptor ligand binding domain. Preferably, the leafhopper ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. In another preferred embodiment, the leafhopper ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides

16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1.

The present invention also relates to compositions comprising an isolated polypeptide according to the invention.

Compositions

The present invention also relates to compositions comprising the isolated polynucleotides or polypeptides according to the invention. Such compositions may comprise a leafhopper ecdysone receptor polypeptide or a polynucleotide encoding a leafhopper ecdysone receptor polypeptide, as defined above, and an acceptable carrier or vehicle. The compositions of the invention are particularly suitable for formulation of biological material for use in a gene expression modulation system or a ligand-screening assay according to the invention. Thus, in a preferred embodiment, the composition comprises a polynucleotide encoding a leafhopper ecdysone receptor polypeptide. In another preferred embodiment, the composition comprises a leafhopper ecdysone receptor polypeptide according to the invention.

The phrase "acceptable" refers to molecular entities and compositions that are physiologically tolerable to the cell or organism when administered. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the composition is administered. Such carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Examples of acceptable carriers are saline, buffered saline, isotonic saline (e.g., monosodium or disodium phosphate, sodium, potassium, calcium or magnesium chloride, or mixtures of such salts), Ringier's solution, dextrose, water, sterile water, glycerol, ethanol, and combinations thereof. 1,3-butanediol and sterile fixed oils are conveniently employed as solvents or suspending media. Any bland fixed oil can be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid also find use in the preparation of injectables. Water or aqueous solution saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E. W. Martin. Pharmaceutical compositions of the invention may be formulated for the purpose of topical, oral, parenteral, intranasal, intravenous, intramuscular, subcutaneous, intraocular, and the like, administration.

Preferably, the compositions comprise an acceptable vehicle for an injectable formulation. This vehicle can be, in particular, a sterile, isotonic saline solution (monosodium or disodium phosphate, sodium, potassium, calcium or magnesium chloride, and the like, or mixtures of such salts), or dry, in particular lyophilized, compositions which, on addition, as appropriate, of sterilized water or of physiological saline, enable injectable solutions to be formed. The preferred sterile injectable preparations can be a solution or suspension in a nontoxic parenterally acceptable solvent or diluent.

In yet another embodiment, a composition comprising a leafhopper ecdysone receptor polypeptide, or polynucleotide encoding the polypeptide, can be delivered in a controlled release system. For example, the polynucleotide or polypeptide may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. Other controlled release systems are discussed in the review by Langer [*Science* 249: 1527-1533 (1990)].

Expression of Leafhopper Ecdysone Receptor Polypeptides

With the sequence of the receptor polypeptides and the polynucleotides encoding them, large quantities of leafhop-

per ecdysone receptor polypeptides may be prepared. By the appropriate expression of vectors in cells, high efficiency production may be achieved. Thereafter, standard purification methods may be used, such as ammonium sulfate precipitations, column chromatography, electrophoresis, centrifugation, crystallization and others. See various volumes of Methods in Enzymology for techniques typically used for protein purification. Alternatively, in some embodiments high efficiency of production is unnecessary, but the presence of a known inducing protein within a carefully engineered expression system is quite valuable. For instance, a combination of: (1) a ligand-responsive enhancer or response element operably linked to (2) a desired gene sequence with (3) the corresponding leafhopper ecdysone receptor polypeptide together in an expression system provides a specifically inducible expression system. Typically, the expression system will be a cell, but an in vitro expression system may also be constructed.

A polynucleotide encoding a leafhopper ecdysone receptor, or fragment, derivative or analog thereof, or a functionally active derivative, including a chimeric protein, thereof, can be inserted into an appropriate expression vector, i.e., a vector which comprises the necessary elements for the transcription and translation of the inserted protein-coding sequence. Thus, the polynucleotide of the invention is operationally linked with a transcriptional control sequence in an expression vector of the invention. Both cDNA and genomic sequences can be cloned and expressed under control of such regulatory sequences. An expression vector also preferably includes a replication origin.

The isolated polynucleotides of the invention may be inserted into any appropriate cloning vector. A large number of vector-host systems known in the art may be used. Possible vectors include, but are not limited to, plasmids or modified viruses, but the vector system must be compatible with the host cell used. Examples of vectors include, but are not limited to, *Escherichia coli*, bacteriophages such as lambda derivatives, or plasmids such as pBR322 derivatives or pUC plasmid derivatives, e.g. pGEX vectors, pmal-c, pFLAG, etc. The insertion into a cloning vector can, for example, be accomplished by ligating the polynucleotide into a cloning vector that has complementary cohesive termini. However, if the complementary restriction sites used to fragment the polynucleotide are not present in the cloning vector, the ends of the polynucleotide molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. Preferably, the cloned gene is contained on a shuttle vector plasmid, which provides for expansion in a cloning cell, e.g., *E. coli*, and purification for subsequent insertion into an appropriate expression cell line, if such is desired. For example, a shuttle vector, which is a vector that can replicate in more than one type of organism, can be prepared for replication in both *E. coli* and *Saccharomyces cerevisiae* by linking sequences from an *E. coli* plasmid with sequences from the yeast 2 μ plasmid.

In addition, the present invention relates to an expression vector comprising a polynucleotide according to the invention, operatively linked to a transcription regulatory element. Preferably, the polynucleotide is operatively linked with an expression control sequence permitting expression of the nuclear receptor ligand binding domain in an expression competent host cell. The expression control sequence may comprise a promoter that is functional in the host cell in which expression is desired. The vector may be a plasmid DNA

molecule or a viral vector. Preferred viral vectors include retrovirus, adenovirus, adeno-associated virus, herpes virus, and vaccinia virus. The invention further relates to a replication defective recombinant virus comprising in its genome, a polynucleotide according to the invention. Thus, the present invention also relates to an isolated host cell comprising such an expression vector, wherein the transcription regulatory element is operative in the host cell.

The desired genes will be inserted into any of a wide selection of expression vectors. The selection of an appropriate vector and cell line depends upon the constraints of the desired product. Typical expression vectors are described in Sambrook et al. (1989). Suitable cell lines may be selected from a depository, such as the ATCC. See, ATCC Catalogue of Cell Lines and Hybridomas (6th ed.) (1988); ATCC Cell Lines, Viruses, and Antisera, each of which is hereby incorporated herein by reference. The vectors are introduced to the desired cells by standard transformation or transfection procedures as described, for instance, in Sambrook et al. (1989).

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation are described generally, for example, in Sambrook et al. (1989), Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, which are incorporated herein by reference. Techniques for synthesis of polypeptides are described, for example, in Merrifield, J. Amer. Chem. Soc. 85: 2149-2156 (1963).

The nucleotide sequences used to produce fusion proteins of the present invention may be derived from natural or synthetic sequences. Many natural gene sequences are obtainable from various cDNA or from genomic libraries using appropriate probes. See, GenBank™, National Institutes of Health. Typical probes for leafhopper ecdysone receptors may be selected from the sequences of Table 1 in accordance with standard procedures. Suitable synthetic DNA fragments may be prepared by the phosphoramidite method described by Beaucage and Carruthers, Tetra. Letts. 22: 1859-1862 (1981). A double stranded fragment may then be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

The natural or synthetic polynucleotide fragments encoding a desired leafhopper ecdysone receptor polypeptide fragment will be incorporated into nucleic acid constructs capable of introduction to and expression in an in vitro cell culture. Usually the nucleic acid constructs will be suitable for replication in a unicellular host, such as yeast or bacteria, but may also be intended for introduction to, with and without and integration within the genome, cultured mammalian or plant or other eukaryotic cell lines. Nucleic acid constructs prepared for introduction into bacteria or yeast will typically include a replication system recognized by the host, the intended DNA fragment encoding the desired receptor polypeptide, transcription and translational initiation regulatory sequences operably linked to the polypeptide encoding segment and transcriptional and translational termination regulatory sequences operably linked to the polypeptide encoding segment. The transcriptional regulatory sequences will typically include a heterologous enhancer, response element, or promoter which is recognized by the host. The selection of an appropriate promoter will depend upon the host, but promoters such as the trp, lac and phage promoters, tRNA promoters and glycolytic enzyme promoters are known. See, Sambrook et al. (1989). Conveniently available expression vectors which include the replication system and transcrip-

tional and translational regulatory sequences together with the insertion site for the steroid receptor DNA sequence may be employed. Examples of workable combinations of cell lines and expression vectors are described in Sambrook et al. (1989); see also, Metzger et al. (1988), Nature 334: 31-36.

Once a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (e.g., lambda), and plasmid and cosmid DNA vectors, to name but a few.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. Expression in yeast can produce a biologically active product. Expression in eukaryotic cells can increase the likelihood of "native" folding. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, leafhopper ecdysone receptor activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent.

Vectors are introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), particle bombardment, use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992, J. Biol. Chem. 267: 963-967; Wu and Wu, 1988, J. Biol. Chem. 263: 14621-14624; Hartmut et al., Canadian Patent Application No. 2,012,311, filed Mar. 15, 1990).

Soluble forms of the protein can be obtained by collecting culture fluid, or solubilizing inclusion bodies, e.g., by treatment with detergent, and if desired sonication or other mechanical processes, as described above. The solubilized or soluble protein can be isolated using various techniques, such as polyacrylamide gel electrophoresis (PAGE), isoelectric focusing, 2-dimensional gel electrophoresis, chromatography (e.g., ion exchange, affinity, immunoaffinity, and sizing column chromatography), centrifugation, differential solubility, immunoprecipitation, or by any other standard technique for the purification of proteins.

Vectors and Gene Expression Cassettes Comprising a Leafhopper Ecdysone Receptor Polynucleotide

Thus, the present invention also relates to a vector comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide according to the invention. The present invention also provides a gene expression cassette comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide according to the invention. The polynucleotides of the invention, where appropriate incorporated in vectors or gene expression cassettes, and the compositions comprising them, are useful for regulating gene expression in an ecdysone receptor-based gene expression system. They may be used for the transfer and expression of genes in vitro or in vivo in any type of cell or tissue. The transformation can, moreover, be targeted (transfer to a particular tissue can, in

particular, be determined by the choice of a vector, and expression by the choice of a particular promoter). The polynucleotides and vectors of the invention are advantageously used for the production in vivo and intracellularly, of polypeptides of interest.

The polynucleotides encoding the leafhopper ecdysone receptor polypeptides of the invention will typically be used in a plasmid vector. Preferably, an expression control sequence is operably linked to the leafhopper ecdysone receptor polynucleotide coding sequence for expression of the leafhopper ecdysone receptor polypeptide. The expression control sequence may be any enhancer, response element, or promoter system in vectors capable of transforming or transfecting a host cell. Once the vector has been incorporated into the appropriate host, the host, depending on the use, will be maintained under conditions suitable for high-level expression of the polynucleotides.

Polynucleotides will normally be expressed in hosts after the sequences have been operably linked to (i.e., positioned to ensure the functioning of) an expression control sequence. These expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. Commonly, expression vectors will contain selection markers, e.g., tetracycline or neomycin, to permit detection of those cells transformed with the desired DNA sequences (see, e.g., U.S. Pat. No. 4,704,362, which is incorporated herein by reference).

Escherichia coli is one prokaryotic host useful for cloning the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species.

Other eukaryotic cells may be used, including yeast cells, insect tissue culture cells, avian cells or the like. Preferably, mammalian tissue cell culture will be used to produce the polypeptides of the present invention (see, Winnacker, From Genes to Clones, VCH Publishers, N.Y. (1987), which is incorporated herein by reference). Yeast and mammalian cells are preferred cells in which to use leafhopper ecdysone receptor-based inducible gene expression systems because they naturally lack the molecules which confer responsiveness to the ligands for ecdysone receptor.

Expression vectors may also include expression control sequences, such as an origin of replication, a promoter, an enhancer, a response element, and necessary processing information sites, such as ribosome-binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferably, the enhancers or promoters will be those naturally associated with genes encoding the steroid receptors, although it will be understood that in many cases others will be equally or more appropriate. Other preferred expression control sequences are enhancers or promoters derived from viruses, such as SV40, Adenovirus, Bovine Papilloma Virus, and the like.

The vectors comprising the polynucleotides of the present invention can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment may be used for other cellular hosts. (See, generally, Sambrook et al. (1989), Molecular Cloning: A Laboratory Manual (2d ed.), Cold Spring Harbor Press, which is incorporated herein by reference). The term "transformed cell" is meant to also include the progeny of a transformed cell.

The necessary transcriptional and translational signals can be provided on a recombinant expression vector, or they may be supplied by the native gene encoding leafhopper ecdysone

receptor and/or its flanking regions. Potential host-vector systems include but are not limited to mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); microorganisms such as yeast containing yeast vectors; or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA.

The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used.

A recombinant leafhopper ecdysone receptor protein of the invention, or functional fragment, derivative, chimeric construct, or analog thereof, may be expressed chromosomally, after integration of the coding sequence by recombination. In this regard, any of a number of amplification systems may be used to achieve high levels of stable gene expression (See Sambrook et al., 1989, supra).

The cell into which the recombinant vector comprising the polynucleotide encoding leafhopper ecdysone receptor is cultured in an appropriate cell culture medium under conditions that provide for expression of leafhopper ecdysone receptor by the cell. Any of the methods previously described for the insertion of DNA fragments into a cloning vector may be used to construct expression vectors containing a gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include in vitro recombinant DNA and synthetic techniques and in vivo recombination (genetic recombination).

A polynucleotide encoding a leafhopper ecdysone receptor polypeptide may be operably linked and controlled by any regulatory region, i.e., promoter/enhancer element known in the art, but these regulatory elements must be functional in the host cell selected for expression. The regulatory regions may comprise a promoter region for functional transcription in the host cell, as well as a region situated 3' of the gene of interest, and which specifies a signal for termination of transcription and a polyadenylation site. All these elements constitute an expression cassette.

Expression vectors comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the invention can be identified by five general approaches: (a) PCR amplification of the desired plasmid DNA or specific mRNA, (b) nucleic acid hybridization, (c) presence or absence of selection marker gene functions, (d) analyses with appropriate restriction endonucleases, and (e) expression of inserted sequences. In the first approach, the nucleic acids can be amplified by PCR to provide for detection of the amplified product. In the second approach, the presence of a foreign gene inserted in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted marker gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "selection marker" gene functions (e.g., β -galactosidase activity, thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of foreign genes in the vector. In another example, if the nucleic acid encoding a leafhopper ecdysone receptor polypeptide is inserted within the "selection marker" gene sequence of the vector, recombinants comprising the leafhopper ecdysone receptor nucleic acid insert can be identified by the absence of the gene function. In the fourth approach, recombinant expression vectors are identified by digestion with appropriate restriction enzymes. In the fifth approach, recombinant expression vectors can be identified by assaying for the activity, biochemi-

cal, or immunological characteristics of the gene product expressed by the recombinant, provided that the expressed protein assumes a functionally active conformation.

A wide variety of host/expression vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include but are not limited to derivatives of SV40 and known bacterial plasmids, e.g., *E. coli* plasmids col E1, pCR1, pBR322, pMal-C2, pET, pGEX (Smith et al., 1988, Gene 67: 31-40), pMB9 and their derivatives, plasmids such as RP4; phage DNAs, e.g., the numerous derivatives of phage λ , e.g., NM989, and other phage DNA, e.g., M13 and filamentous single stranded phage DNA; yeast plasmids such as the 2 μ plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian cells; vectors derived from combinations of plasmids and phage DNAs, such as plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

The present invention also provides a gene expression cassette that is capable of being expressed in a host cell, wherein the gene expression cassette comprises a polynucleotide that encodes a leafhopper ecdysone receptor polypeptide according to the invention. Thus, Applicants' invention also provides novel gene expression cassettes useful in an ecdysone receptor-based gene expression system

In a specific embodiment, the gene expression cassette that is capable of being expressed in a host cell comprises a polynucleotide that encodes a polypeptide selected from the group consisting of a) a polypeptide comprising a transactivation domain, a DNA-binding domain, and a leafhopper ecdysone receptor ligand binding domain; b) a polypeptide comprising a DNA-binding domain and a leafhopper ecdysone receptor ligand binding domain; and c) a polypeptide comprising a transactivation domain and a leafhopper ecdysone receptor ligand binding domain.

In another specific embodiment, the present invention provides a gene expression cassette that is capable of being expressed in a host cell, wherein the gene expression cassette comprises a polynucleotide that encodes a hybrid polypeptide selected from the group consisting of a) a hybrid polypeptide comprising a transactivation domain, a DNA-binding domain, and a leafhopper ecdysone receptor ligand binding domain; b) a hybrid polypeptide comprising a DNA-binding domain and a leafhopper ecdysone receptor ligand binding domain; and c) a hybrid polypeptide comprising a transactivation domain and a leafhopper ecdysone receptor ligand binding domain. A hybrid polypeptide according to the invention comprises at least two polypeptide fragments, wherein each polypeptide fragment is from a different source, i.e., a different polypeptide, a different nuclear receptor, a different species, etc. The hybrid polypeptide according to the invention may comprise at least two polypeptide domains, wherein each polypeptide domain is from a different source.

Preferably, the leafhopper ecdysone receptor ligand binding domain is from a green leafhopper *Nephotetix cincticeps* EcR ("NcEcR").

In a specific embodiment, the leafhopper ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1.

In a specific embodiment, the leafhopper ecdysone receptor ligand binding domain comprises an amino acid sequence

selected from the group consisting of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2.

The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, such as an EcR DBD, or a bacterial LacZ DBD.

The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- κ B AD, a BP64 AD, a B42 acidic activation domain (B42AD), a p65 transactivation domain (p65AD), or an analog, combination, or modification thereof. In a specific embodiment, the AD is a synthetic or chimeric AD, or is obtained from an EcR, a glucocorticoid receptor, VP16, GAL4, NF- κ B, or B42 acidic activation domain AD.

In a specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising either a) a DNA-binding domain, or b) a transactivation domain; and an NcEcR ligand binding domain according to the invention.

The present invention also provides a gene expression cassette comprising: i) a response element comprising a domain recognized by a polypeptide comprising a DNA binding domain; ii) a promoter that is activated by a polypeptide comprising a transactivation domain; and iii) a gene whose expression is to be modulated.

The response element ("RE") may be any response element with a known DNA binding domain, or an analog, combination, or modification thereof. A single RE may be employed or multiple REs, either multiple copies of the same RE or two or more different REs, may be used in the present invention. In a specific embodiment, the RE is an RE from GAL4 ("GAL4RE"), LexA, a steroid/thyroid hormone nuclear receptor RE, such as an ecdysone response element (EcRE), or a synthetic RE that recognizes a synthetic DNA binding domain.

A steroid/thyroid hormone nuclear receptor DNA binding domain, activation domain or response element according to the invention may be obtained from a steroid/thyroid hormone nuclear receptor selected from the group consisting of ecdysone receptor (EcR), ubiquitous receptor (UR), Orphan receptor 1 (OR-1), steroid hormone nuclear receptor 1 (NER-1), RXR interacting protein-15 (RIP-15), liver x receptor β (LXR β), steroid hormone receptor like protein (RLD-1), liver x receptor (LXR), liver x receptor α (LXR α), farnesoid x receptor (FXR), receptor interacting protein 14 (RIP-14), farnesol receptor (HRR-1), thyroid hormone receptor α (TR α), thyroid receptor 1 (c-erbA-1), thyroid hormone receptor β (TR β), retinoic acid receptor α (RAR α), retinoic acid receptor β (RAR β , HAP), retinoic acid receptor γ (RAR γ), retinoic acid receptor gamma-like (RARD), peroxisome proliferator-activated receptor α (PPAR α), peroxisome proliferator-activated receptor β (PPAR β), peroxisome proliferator-activated receptor δ (PPAR δ , NUC-1), peroxisome proliferator-activator related receptor (FFAR), peroxisome proliferator-activated receptor γ (PPAR γ), orphan receptor encoded by non-encoding strand of thyroid hormone receptor α (REVERB α), v-erb A related receptor (EAR-1), v-erb related receptor (EAR-1A), γ), orphan receptor encoded by non-encoding strand of thyroid hormone receptor β (REVERB β), v-erb related receptor (EAR-1 β), orphan nuclear receptor BD73 (BD73), rev-erbA-related receptor (RVR),

zinc finger protein 126 (HZF2), ecdysone-inducible protein E75 (E75), ecdysone-inducible protein E78 (E78), *Drosophila* receptor 78 (DR-78), retinoid-related orphan receptor α (ROR α), retinoid Z receptor α (RZR α), retinoid related orphan receptor β (ROR β), retinoid Z receptor β (RZR β), retinoid-related orphan receptor γ (ROR γ), retinoid Z receptor γ (RZR γ), retinoid-related orphan receptor (TOR), hormone receptor 3 (HR-3), *Drosophila* hormone receptor 3 (DHR-3), *Manduca* hormone receptor 3 (MHR-3), *Gallaria* hormone receptor 3 (GHR-3), *C. elegans* nuclear receptor 3 (CNR-3), *Choristoneura* hormone receptor 3 (CHR-3), *C. elegans* nuclear receptor 14 (CNR-14), vitamin D receptor (VDR), orphan nuclear receptor (ONR-1), pregnane X receptor (PXR), steroid and xenobiotic receptor (SXR), benzoate X receptor (BXR), nuclear receptor (MB-67), constitutive androstane receptor 1 (CAR-1), constitutive androstane receptor α (CAR α), constitutive androstane receptor 2 (CAR-2), constitutive androstane receptor β (CAR β), *Drosophila* hormone receptor 96 (DHR-96), nuclear hormone receptor 1 (NHR-1), hepatocyte nuclear factor 4 (HNF-4), hepatocyte nuclear factor 4G (HNF-4G), hepatocyte nuclear factor 4B (HNF-4B), hepatocyte nuclear factor 4D (HNF-4D, DHNF-4), retinoid X receptor α (RXR α), retinoid X receptor β (RXR β), H-2 region II binding protein (H-2RIIBP), nuclear receptor co-regulator-1 (RCoR-1), retinoid X receptor γ (RXR γ), Ultraspiracle (USP), 2C1 nuclear receptor, chorion factor 1 (CF-1), testicular receptor 2 (TR-2), testicular receptor 2-11 (TR2-11), testicular receptor 4 (TR4), TAK-1, *Drosophila* hormone receptor (DHR78), Tailless (TLL), tailless homolog (TLX), XTLL, chicken ovalbumin upstream promoter transcription factor I (COUP-TFI), chicken ovalbumin upstream promoter transcription factor A (COUP-TFA), EAR-3, SVP-44, chicken ovalbumin upstream promoter transcription factor II (COUP-TFII), chicken ovalbumin upstream promoter transcription factor B (COUP-TFB), ARP-1, SVP40, SVP, chicken ovalbumin upstream promoter transcription factor III (COUP-TFIII), chicken ovalbumin upstream promoter transcription factor G (COUP-TFG), SVP-46, EAR-2, estrogen receptor α (ER α), estrogen receptor β (ER β), estrogen related receptor 1 (ERR1), estrogen related receptor α (ERR α), estrogen related receptor 2 (ERR2), estrogen related receptor β (ERR β), glucocorticoid receptor (GR), mineralocorticoid receptor (MR), progesterone receptor (PR), androgen receptor (AR), nerve growth factor induced gene B (NGFI-B), nuclear receptor similar to Nur-77 (TRS), N10, Orphan receptor (NUR-77), Human early response gene (NAK-1), Nurr related factor 1 (NURR-1), a human immediate-early response gene (NOT), regenerating liver nuclear receptor 1 (RNR-1), hematopoietic zinc finger 3 (HZF-3), Nur related protein-1 (TINOR), Nuclear orphan receptor 1 (NOR-1), NOR1 related receptor (MINOR), *Drosophila* hormone receptor 38 (DHR-38), *C. elegans* nuclear receptor 8 (CNR-8), C48D5, steroidogenic factor 1 (SF1), endozep-like peptide (ELP), fushi tarazu factor 1 (FTZ-F1), adrenal 4 binding protein (AD4BP), liver receptor homolog (LRH-1), Ftz-F1-related orphan receptor A (xFFrA), Ftz-F1-related orphan receptor B (xFFrB), nuclear receptor related to LRH-1 (FFLR), nuclear receptor related to LRH-1 (PHR), fetoprotein transcription factor (FTF), germ cell nuclear factor (GCNFM), retinoid receptor-related testis-associated receptor (RTR), knirps (KNI), knirps related (KNRL), Embryonic gonad (EGON), *Drosophila* gene for ligand dependent nuclear receptor (EAGLE), nuclear receptor similar to trithorax (ODR7), Trithorax, dosage sensitive sex reversal adrenal hypoplasia congenita critical region

chromosome X gene (DAX-1), adrenal hypoplasia congenita and hypogonadotropic hypogonadism (AHCH), and short heterodimer partner (SHP).

For purposes of this invention, nuclear receptors and leafhopper ecdysone receptors also include synthetic and chimeric nuclear receptors and leafhopper ecdysone receptors and their homologs.

Antibodies to Leafhopper Ecdysone Receptor

According to the invention, a leafhopper ecdysone receptor polypeptide produced recombinantly or by chemical synthesis, and fragments or other derivatives or analogs thereof, including fusion proteins, may be used as an antigen or immunogen to generate antibodies. Preferably, the antibodies specifically bind homopteran ecdysone receptor polypeptides, but do not bind other ecdysone receptor polypeptides. More preferably, the antibodies specifically bind a leafhopper ecdysone receptor polypeptide, but do not bind other ecdysone receptor polypeptides.

The present invention also relates to antigenic peptides and antibodies thereto. More particularly, the invention relates to antigenic peptides comprising a fragment of a leafhopper ecdysone receptor polypeptide according to the invention, wherein the fragment has a property selected from the group consisting of:

(a) it is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of nucleotides 1-15 of SEQ ID NO: 1, nucleotides 16-213 of SEQ ID NO: 1, nucleotides 214-447 of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1;

(b) it comprises an amino acid sequence selected from the group consisting of amino acids 1-5 of SEQ ID NO: 2, amino acids 6-71 of SEQ ID NO: 2, amino acids 72-149 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2; and

(c) it specifically binds to an antibody generated against an epitope within a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 1-5 of SEQ ID NO: 2, amino acids 6-71 of SEQ ID NO: 2, amino acids 72-149 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2.

In another embodiment, the invention relates to an antibody which specifically binds an antigenic peptide comprising a fragment of a leafhopper ecdysone receptor polypeptide according to the invention as described above. The antibody may be polyclonal or monoclonal and may be produced by *in vitro* or *in vivo* techniques.

The antibodies of the invention possess specificity for binding to particular homopteran ecdysone receptors. Thus, reagents for determining qualitative or quantitative presence of these or homologous polypeptides may be produced. Alternatively, these antibodies may be used to separate or purify receptor polypeptides.

For production of polyclonal antibodies, an appropriate target immune system is selected, typically a mouse or rabbit. The substantially purified antigen is presented to the immune system in a fashion determined by methods appropriate for the animal and other parameters well known to immunologists. Typical sites for injection are in the footpads, intramuscularly, intraperitoneally, or intradermally. Of course, another species may be substituted for a mouse or rabbit.

An immunological response is usually assayed with an immunoassay. Normally such immunoassays involve some purification of a source of antigen, for example, produced by the same cells and in the same fashion as the antigen was produced. The immunoassay may be a radioimmunoassay, an enzyme-linked assay (ELISA), a fluorescent assay, or any of many other choices, most of which are functionally equivalent but may exhibit advantages under specific conditions.

Monoclonal antibodies with high affinities are typically made by standard procedures as described, e.g., in Harlow and Lane (1988), *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory; or Goding (1986), *Monoclonal Antibodies: Principles and Practice* (2d ed) Academic Press, New York, which are hereby incorporated herein by reference. Briefly, appropriate animals will be selected and the desired immunization protocol followed. After the appropriate period of time, the spleens of such animals are excised and individual spleen cells fused, typically, to immortalized myeloma cells under appropriate selection conditions. Thereafter, the cells are clonally separated and the supernatants of each clone are tested for their production of an appropriate antibody specific for the desired region of the antigen.

Other suitable techniques involve *in vitro* exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse et al., (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246: 1275-1281, hereby incorporated herein by reference.

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radio-nuclides, enzymes, substrates, cofactors, inhibitors, fluorescence, chemiluminescence, magnetic particles and the like. Patents, teaching the use of such labels include U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see Cabilly, U.S. Pat. No. 4,816,567.

A molecule is "antigenic" when it is capable of specifically interacting with an antigen recognition molecule of the immune system, such as an immunoglobulin (antibody) or T cell antigen receptor. An antigenic polypeptide contains at least about 5, and preferably at least about 10 amino acids. An antigenic portion of a molecule can be that portion that is immunodominant for antibody or T cell receptor recognition, or it can be a portion used to generate an antibody to the molecule by conjugating the antigenic portion to a carrier molecule for immunization. A molecule that is antigenic need not be itself immunogenic, i.e., capable of eliciting an immune response without a carrier.

Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. The anti-leafhopper ecdysone receptor antibodies of the invention may be cross-reactive, e.g., they may recognize leafhopper ecdysone receptor from different species. Polyclonal antibodies have greater likelihood of cross reactivity. Alternatively, an antibody of the invention may be specific for a single form of leafhopper ecdysone receptor, such as leafhopper ecdysone receptor. Preferably, such an antibody is specific for leafhopper ecdysone receptor.

Various procedures known in the art may be used for the production of polyclonal antibodies. For the production of antibody, various host animals can be immunized by injection

with the leafhopper ecdysone receptor polypeptide, or a derivative (e.g., fragment or fusion protein) thereof, including but not limited to rabbits, mice, rats, sheep, goats, etc. In one embodiment, the leafhopper ecdysone receptor polypeptide or fragment thereof can be conjugated to an immunogenic carrier, e.g., bovine serum albumin (BSA) or keyhole limpet hemocyanin (KLH). Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (*bacille Calmette-Guerin*) and *Corynebacterium parvum*.

For preparation of monoclonal antibodies directed toward the leafhopper ecdysone receptor polypeptide, or fragment, analog, or derivative thereof, any technique that provides for the production of antibody molecules by continuous cell lines in culture may be used. These include but are not limited to the hybridoma technique originally developed by Kohler and Milstein [*Nature* 256:495-497 (1975)], as well as the trioma technique, the human B-cell hybridoma technique [Kozbor et al., *Immunology Today* 4: 72 (1983); Cote et al., *Proc. Natl. Acad. Sci. U.S.A.* 80: 2026-2030 (1983)], and the EBV-hybridoma technique to produce human monoclonal antibodies [Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)]. In an additional embodiment of the invention, monoclonal antibodies can be produced in germ-free animals [International Patent Publication No. WO 89/12690, published 28 Dec. 1989]. In fact, according to the invention, techniques developed for the production of "chimeric antibodies" [Morrison et al., *J. Bacteriol.* 159: 870 (1984); Neuberger et al., *Nature* 312: 604-608 (1984); Takeda et al., *Nature* 314: 452-454 (1985)] by splicing the genes from a mouse antibody molecule specific for a leafhopper ecdysone receptor polypeptide together with genes from a human antibody molecule of appropriate biological activity can be used; such antibodies are within the scope of this invention. Such human or humanized chimeric antibodies are preferred for use in therapy of human diseases or disorders (described *infra*), since the human or humanized antibodies are much less likely than xenogenic antibodies to induce an immune response, in particular an allergic response, themselves.

According to the invention, techniques described for the production of single chain Fv (scFv) antibodies [U.S. Pat. Nos. 5,476,786 and 5,132,405 to Huston; U.S. Pat. No. 4,946,778] can be adapted to produce leafhopper ecdysone receptor polypeptide-specific single chain antibodies. An additional embodiment of the invention utilizes the techniques described for the construction of Fab expression libraries [Huse et al., *Science* 246: 1275-1281 (1989)] to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for a leafhopper ecdysone receptor polypeptide, or its derivatives, or analogs.

Antibody fragments which contain the idiotype of the antibody molecule can be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, and the Fab fragments which can be generated by treating the antibody molecule with papain and a reducing agent.

In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art,

e.g., radioimmunoassay, ELISA (enzyme-linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitin reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention. For example, to select antibodies which recognize a specific epitope of a leafhopper ecdysone receptor polypeptide, one may assay generated hybridomas for a product which binds to a leafhopper ecdysone receptor polypeptide fragment containing such epitope. For selection of an antibody specific to a leafhopper ecdysone receptor polypeptide from a particular species of animal, one can select on the basis of positive binding with leafhopper ecdysone receptor polypeptide expressed by or isolated from cells of that species of animal.

The foregoing antibodies can be used in methods known in the art relating to the localization and activity of the leafhopper ecdysone receptor polypeptide, e.g., for western blotting, imaging leafhopper ecdysone receptor polypeptide in situ, measuring levels thereof in appropriate physiological samples, etc. using any of the detection techniques mentioned above or known in the art.

In a specific embodiment, antibodies that agonize or antagonize the activity of leafhopper ecdysone receptor polypeptide can be generated. Such antibodies can be tested using the assays described *infra* for identifying ligands. In particular, such antibodies can be scFv antibodies expressed intracellularly.

40 Uses of Novel Polynucleotides and Polypeptides of the Invention

The present invention further provides a number of uses for the leafhopper ecdysone receptor polynucleotides of the present invention and their encoded polypeptides.

45 The leafhopper ecdysone receptor polypeptides of the present invention have a variety of utilities. For example, the polynucleotides and polypeptides of the invention are useful in methods of modulating gene expression in an ecdysone receptor-based gene expression system. Also included are methods for identifying and selecting ligands specific for binding to a ligand binding domain of a polypeptide of the invention, methods for identifying and selecting compounds exhibiting specific binding to the ligand binding domain and methods for modulating insect physiology or development (e.g., killing).

Methods of Modulating Gene Expression

As presented herein, Applicants' novel polynucleotides and polypeptides are useful in an ecdysone receptor-based gene expression system to provide a regulatable gene expression system in both prokaryotic and eukaryotic host cells. Thus, the present invention also relates to the use of the novel leafhopper ecdysone receptor polynucleotides and polypeptides of the present invention in an ecdysone receptor-based gene expression system, and methods of modulating the expression of a gene within a host cell using such an ecdysone receptor-based gene expression system.

This gene expression system may be a “single switch”-based gene expression system in which the transactivation domain, DNA-binding domain and ligand binding domain are on one encoded polypeptide. Alternatively, the gene expression modulation system may be a “dual switch”- or “two-hybrid”-based gene expression modulation system in which the transactivation domain and DNA-binding domain are located on two different encoded polypeptides. Applicants’ have demonstrated for the first time that leafhopper ecdysone receptor polynucleotides and polypeptides of the invention can be used as a component of an ecdysone receptor-based inducible gene expression system to modify gene expression in a host cell.

In particular, the present invention relates to a gene expression modulation system comprising at least one gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide that encodes a leafhopper ecdysone receptor polypeptide. Preferably, the leafhopper ecdysone receptor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 1-5 of SEQ ID NO: 2, amino acids 6-71 of SEQ ID NO: 2, amino acids 72-149 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. More preferably, the leafhopper ecdysone receptor polypeptide comprises amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, or amino acids 140-370 of SEQ ID NO: 2.

In a specific embodiment, the gene expression modulation system comprises a gene expression cassette comprising a polynucleotide that encodes a polypeptide comprising a transactivation domain, a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and a leafhopper ecdysone receptor ligand binding domain (referred to herein as “NcEcR LBD”). The gene expression modulation system may further comprise a second gene expression cassette comprising: i) a response element recognized by the DNA-binding domain of the encoded polypeptide of the first gene expression cassette; ii) a promoter that is activated by the transactivation domain of the encoded polypeptide of the first gene expression cassette; and iii) a gene whose expression is to be modulated.

In another specific embodiment, the gene expression modulation system comprises a gene expression cassette comprising a) a polynucleotide that encodes a polypeptide comprising a transactivation domain, a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and an NcEcR LBD, and b) a second nuclear receptor ligand binding domain selected from the group consisting of a vertebrate retinoid X receptor ligand binding domain, an invertebrate retinoid X receptor ligand binding domain, an ultraspiracle protein ligand binding domain, and a chimeric ligand binding domain comprising two polypeptide fragments, wherein the first polypeptide fragment is from a vertebrate retinoid X receptor ligand binding domain, an invertebrate retinoid X receptor ligand binding domain, or an ultraspiracle protein ligand binding domain, and the second polypeptide fragment is from a different vertebrate retinoid X receptor ligand binding domain, invertebrate retinoid X receptor ligand binding domain, or ultraspiracle protein ligand binding domain. The gene expression modulation system may further comprise a second gene expression cassette comprising: i) a response element recognized by the DNA-binding domain of the encoded polypeptide of the first gene expression cassette; ii) a promoter that is activated by the transactivation domain of

the encoded polypeptide of the first gene expression cassette; and iii) a gene whose expression is to be modulated.

In another specific embodiment, the gene expression modulation system comprises a first gene expression cassette comprising a polynucleotide that encodes a first polypeptide comprising a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated and a nuclear receptor ligand binding domain, and a second gene expression cassette comprising a polynucleotide that encodes a second polypeptide comprising a transactivation domain and a nuclear receptor ligand binding domain, wherein one of the nuclear receptor ligand binding domains is an NcEcR LBD. In a preferred embodiment, the first polypeptide is substantially free of a transactivation domain and the second polypeptide is substantially free of a DNA binding domain. For purposes of the invention, “substantially free of a DNA binding domain” means that the protein in question does not contain a sufficient sequence of the domain in question to provide activation or binding activity. The gene expression modulation system may further comprise a third gene expression cassette comprising: i) a response element recognized by the DNA-binding domain of the first polypeptide of the first gene expression cassette; ii) a promoter that is activated by the transactivation domain of the second polypeptide of the second gene expression cassette; and iii) a gene whose expression is to be modulated.

Wherein when only one nuclear receptor ligand binding domain is an NcEcR LBD, the other nuclear receptor ligand binding domain may be from any other nuclear receptor that forms a dimer with the NcEcR LBD. For example, the other nuclear receptor ligand binding domain (“partner”) may be from another ecdysone receptor, a vertebrate retinoid X receptor (RXR), an invertebrate RXR, an ultraspiracle protein (USP), or a chimeric nuclear receptor comprising at least two different nuclear receptor ligand binding domain polypeptide fragments selected from the group consisting of a vertebrate RXR, an invertebrate RXR, and a USP (see co-pending applications PCT/US01/09050, U.S. 60/294,814, and U.S. 60/294,819, incorporated herein by reference in their entirety). The “partner” nuclear receptor ligand binding domain may further comprise a truncation mutation, a deletion mutation, a substitution mutation, or another modification.

In a specific embodiment, the leafhopper ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2. In another embodiment, the leafhopper ecdysone receptor ligand binding domain is encoded by a polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1.

In a specific embodiment, the gene whose expression is to be modulated is a homologous gene with respect to the host cell. In another specific embodiment, the gene whose expression is to be modulated is a heterologous gene with respect to the host cell.

The ligands for use in the methods of modulating gene expression are described below, when combined with the ligand binding domain of the nuclear receptor(s), which in turn are bound to the response element linked to a gene, provide the means for external temporal regulation of expression of the gene. The binding mechanism or the order in which the various components of this invention bind to each other, that is, for example, ligand to ligand binding domain,

41

DNA-binding domain to response element, transactivation domain to promoter, etc., is not critical.

Thus, Applicants' invention is useful in methods of modulating gene expression in a host cell using a leafhopper ecdysone receptor according to the invention. Specifically, Applicants' invention provides a method of modulating the expression of a gene in a host cell comprising the steps of: a) introducing into the host cell an ecdysone receptor-based gene expression modulation system comprising a leafhopper ecdysone receptor according to the invention; and b) introducing into the host cell a ligand; wherein the gene to be modulated is a component of a gene expression cassette comprising: i) a response element comprising a domain recognized by the DNA binding domain of the gene expression system; ii) a promoter that is activated by the transactivation domain of the gene expression system; and iii) a gene whose expression is to be modulated, whereby upon introduction of the ligand into the host cell, expression of the gene is modulated.

The invention also provides a method of modulating the expression of a gene in a host cell comprising the steps of: a) introducing into the host cell an ecdysone receptor-based gene expression modulation system comprising a leafhopper ecdysone receptor according to the invention; b) introducing into the host cell a gene expression cassette, wherein the gene expression cassette comprises i) a response element comprising a domain recognized by the DNA binding domain from the gene expression system; ii) a promoter that is activated by the transactivation domain of the gene expression system; and iii) a gene whose expression is to be modulated; and c) introducing into the host cell a ligand; whereby upon introduction of the ligand into the host cell, expression of the gene is modulated.

Genes of interest for expression in a host cell using Applicants' methods may be endogenous genes or heterologous genes. Nucleic acid or amino acid sequence information for a desired gene or protein can be located in one of many public access databases, for example, GENBANK, EMBL, Swiss-Prot, and PIR, or in many biology related journal publications. Thus, those skilled in the art have access to nucleic acid sequence information for virtually all known genes. Such information can then be used to construct the desired constructs for the insertion of the gene of interest within the gene expression cassettes used in the methods described herein.

Examples of genes of interest for expression in a host cell using these methods include, but are not limited to: antigens produced in plants as vaccines, enzymes like alpha-amylase, phytase, glucanase, xylase and xylanase, genes for resistance against insects, nematodes, fungi, bacteria, viruses, and abiotic stresses, nutraceuticals, pharmaceuticals, vitamins, genes for modifying amino acid content, herbicide resistance, cold, drought, and heat tolerance, industrial products, oils, protein, carbohydrates, antioxidants, male sterile plants, flowers, fuels, other output traits, genes encoding therapeutically desirable polypeptides or products that may be used to treat a condition, a disease, a disorder, a dysfunction, a genetic defect, such as monoclonal antibodies, enzymes, proteases, cytokines, interferons, insulin, erythropoietin, clotting factors, other blood factors or components, viral vectors for gene therapy, virus for vaccines, targets for drug discovery, functional genomics, and proteomics analyses and applications, and the like.


The term "ligand" is meant herein to refer to a molecule that binds the domain described here as the "ligand binding domain." Also, a ligand for a leafhopper ecdysone receptor is a ligand which serves either as the natural ligand to which the

42

ecdysone receptor binds, or a functional analogue which may serve as an agonist or antagonist.

Acceptable ligands are any that modulate expression of the gene when binding of the DNA binding domain of the gene expression system according to the invention to the response element in the presence of the ligand results in activation or suppression of expression of the genes. Preferred ligands include an ecdysteroid, such as ecdysone, 20-hydroxyecdysone, ponasterone A, muristerone A, and the like, 9-cis-retinoic acid, synthetic analogs of retinoic acid, N,N'-diacylhydrazines such as those disclosed in U.S. Pat. Nos. 6,013,836; 5,117,057; 5,530,028; and 5,378,726; dibenzoylalkyl cyanohydrazines such as those disclosed in European Application No. 461,809; N-alkyl-N,N'-diarylhydrazines such as those disclosed in U.S. Pat. No. 5,225,443; N-acyl-N-alkylcarbonylhydrazines such as those disclosed in European Application No. 234,994; N-aryl-N-alkyl-N'-arylhydrazines such as those described in U.S. Pat. No. 4,985,461; each of which is incorporated herein by reference and other similar materials including 3,5-di-tert-butyl-4-hydroxy-N-isobutyl-benzamide, 8-O-acetylharpagide, and the like.

In a preferred embodiment, the ligand for use in the method of modulating expression of gene is a compound of the formula:



wherein:

E is a (C₄-C₆)alkyl containing a tertiary carbon or a cyano (C₃-C₅)alkyl containing a tertiary carbon;

R¹ is H, Me, Et, i-Pr, F, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, C^oCH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, SCN, or SCHF₂;

R² is H, Me, Et, n-Pr, i-Pr, formyl, CF₃, CHF₂, CHCl₂, CH₂F, CH₂Cl, CH₂OH, CH₂OMe, CH₂CN, CN, C^oCH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe₂, NEt₂, SMe, SEt, SOCF₃, OCF₂CF₂H, COEt, cyclopropyl, CF₂CF₃, CH=CHCN, allyl, azido, OCF₃, OCHF₂, O-i-Pr, SCN, SCHF₂, SOMe, NH—CN, or joined with R³ and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R³ is H, Et, or joined with R² and the phenyl carbons to which R² and R³ are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R⁴, R⁵, and R⁶ are independently H, Me, Et, F, Cl, Br, formyl, CF₃, CHF₂, CHCl₂CH₂F, CH₂Cl, CH₂OH, CN, C^oCH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

In another preferred embodiment, the ligand for use in the method of modulating expression of gene is an ecdysone, 20-hydroxyecdysone, ponasterone A, or muristerone A.

In another preferred embodiment, a second ligand may be used in addition to the first ligand discussed above in the method of modulating expression of a gene. Preferably, this second ligand is 9-cis-retinoic acid or a synthetic analog of retinoic acid.

In another preferred embodiment, the ligand for use in the method of modulating expression of gene is an ecdysone, 20-hydroxyecdysone, ponasterone A, or muristerone A.

In another preferred embodiment, a second ligand may be used in addition to the first ligand discussed above in the method of modulating expression of a gene. Preferably, this second ligand is 9-cis-retinoic acid or a synthetic analog of retinoic acid.

Screening Assays

Identification and isolation of a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the invention provides for expression of leafhopper ecdysone receptor in quantities greater than can be isolated from natural sources, or in indicator cells that are specially engineered to indicate the activity of leafhopper ecdysone receptor expressed after transfection or transformation of the cells. Accordingly, in addition to rational design of agonists and antagonists based on the structure of leafhopper ecdysone receptor polypeptide, the present invention contemplates an alternative method for identifying specific ligands of leafhopper ecdysone receptor using various screening assays known in the art.

Thus, the present invention also relates to methods of screening for a compound that induces or represses transactivation of a leafhopper ecdysone receptor polypeptide in a cell by contacting a leafhopper ecdysone receptor polypeptide with a candidate molecule and detecting reporter gene activity in the presence of the ligand. Candidate compounds may be either agonists or antagonists of the leafhopper ecdysone receptor polypeptide. In a preferred embodiment, the leafhopper ecdysone receptor polypeptide is expressed from a polynucleotide in the cell and the transactivation activity (i.e., expression or repression of a reporter gene) or compound binding activity is measured.

In a specific embodiment, the present invention relates to methods of screening for molecules that stimulate or inhibit leafhopper ecdysone receptor activity in a cell by contacting a leafhopper ecdysone receptor polypeptide with a candidate molecule and detecting leafhopper ecdysone receptor activity in the presence of the molecule. Candidate molecules may be either agonists or antagonists of leafhopper ecdysone receptor. In a preferred embodiment, the leafhopper ecdysone receptor is expressed from a polynucleotide in the cell and the leafhopper ecdysone receptor activity measured is by induction of expression or transactivation of a reporter gene. Induction of reporter gene expression can be measured as described herein.

Thus, one aspect of the present invention is a method for selecting molecules or ligands that modulate the activity of a leafhopper ecdysone receptor polypeptide. In a specific embodiment, the present invention provides a method for identifying a ligand specific for binding to a ligand binding domain of a leafhopper ecdysone receptor comprising

- (a) combining (i) a hybrid polypeptide comprising a leafhopper ecdysone receptor ligand binding domain and a DNA binding domain from a steroid hormone nuclear receptor superfamily; and (ii) a polynucleotide encoding a second polypeptide, wherein the polynucleotide is operably linked to a transcriptional control element that is responsive to the DNA binding domain of the hybrid polypeptide;
- (b) exposing the hybrid polypeptide and the polynucleotide of (a) to a compound;
- (c) determining ligand activity of the compound of (b) by determining induction of expression of the second polypeptide; and
- (d) identifying the compound that results in the induction of expression of the second polypeptide.

The present invention is also useful to search for orthogonal ligands and orthogonal receptor-based gene expression systems such as those described in co-pending U.S. application Ser. No. 60/237,446, which is incorporated herein by reference in its entirety.

The ligand binding domain ("LBD") of the ecdysone receptor, specifically binds steroid and non-steroidal agonist ligands, thereby providing a means to screen for new mol-

ecules possessing the property of binding with high affinity to the ligand binding domain. Thus, the ligand binding domain of a leafhopper ecdysone receptor polypeptide may be used as a reagent to develop a binding assay. On one level, the LBD can be used as an affinity reagent for a batch or in a column selective process, to selectively retain ligands which bind. Alternatively, a functional assay is preferred for its greater sensitivity to ligand binding. By using a reporter molecule for binding, either through a direct assay for binding, or through an expression or other functional linkage between binding and another function, an assay for binding may be developed. For example, by operable linkage of an easily assayable reporter gene to a controlling element responsive to binding by an ecdysone receptor, and where ligand binding is functionally linked to protein induction, an extremely sensitive assay for the presence of a ligand or of a receptor results. Such a construct is useful for assaying the presence of 20-hydroxyecdysone is described below. This construct is useful for screening for agonists or antagonists of homopteran ecdysone receptors, in particular, leafhopper ecdysone receptors.

As presented herein, a leafhopper ecdysone receptor can transactivate gene expression of an ecdysone receptor-based gene expression modulation system. Therefore, agonists of leafhopper ecdysone receptor that enhance its ability to transactivate gene expression will be expected to improve its activity in an ecdysone receptor-based gene expression modulation system. Inhibitors (antagonists) of leafhopper ecdysone receptor activity are useful to reduce its ability to transactivate an ecdysone receptor-based gene expression modulation system.

Any screening technique known in the art can be used to screen for leafhopper ecdysone receptor agonists or antagonists. For example, a suitable cell line expressing both leafhopper ecdysone receptor and an ecdysone receptor-based gene expression modulation system can be transfected with a nucleic acid encoding a marker gene, such as β -galactosidase. Cells are then exposed to a test solution comprising an agonist or antagonist, and then stained for β -galactosidase activity. The presence of more β -gal positive cells relative to control cells not exposed to the test solution is an indication of the presence of a leafhopper ecdysone receptor agonist in the test solution. Conversely, the presence of less β -gal positive cells relative to control cells not exposed to the test solution is an indication of the presence of a leafhopper ecdysone receptor antagonist in the test solution.

The present invention contemplates screens for small molecule ligands or ligand analogs and mimics, as well as screens for natural ligands that bind to and agonize or antagonize leafhopper ecdysone receptor in vivo. For example, natural products libraries can be screened using assays of the invention for molecules that agonize or antagonize leafhopper ecdysone receptor activity.

Knowledge of the primary sequence of leafhopper ecdysone receptor, and the similarity of that sequence with proteins of known function, can provide an initial clue as the inhibitors or antagonists of the protein. Identification and screening of antagonists is further facilitated by determining structural features of the protein, e.g., using X-ray crystallography, neutron diffraction, nuclear magnetic resonance spectrometry, and other techniques for structure determination. These techniques provide for the rational design or identification of agonists and antagonists.

Another approach uses recombinant bacteriophage to produce large libraries. Using the "phage method" [Scott and Smith, 1990, *Science* 249: 386-390 (1990); Cwirla, et al., *Proc. Natl. Acad. Sci.*, 87: 6378-6382 (1990); Devlin et al., *Science*, 249: 404-406 (1990)], very large libraries can be

constructed (10^6 - 10^8 chemical entities). A second approach uses primarily chemical methods, of which the Geysen method [Geysen et al., *Molecular Immunology* 23: 709-715 (1986); Geysen et al. *J. Immunologic Method* 102: 259-274 (1987)] and the method of Fodor et al. [*Science* 251: 767-773 (1991)] are examples. Furka et al. [*14th International Congress of Biochemistry, Volume 5*, Abstract FR:013 (1988); Furka, *Int. J. Peptide Protein Res.* 37: 487-493 (1991)], Houghton [U.S. Pat. No. 4,631,211, issued December 1986] and Rutter et al. [U.S. Pat. No. 5,010,175, issued Apr. 23, 1991] describe methods to produce a mixture of peptides that can be tested as agonists or antagonists.

In another aspect, synthetic libraries [Needels et al., *Proc. Natl. Acad. Sci. USA* 90: 10700-4 (1993); Ohlmeyer et al., *Proc. Natl. Acad. Sci. USA* 90: 10922-10926 (1993); Lam et al., International Patent Publication No. WO 92/00252; Kocis et al., International Patent Publication No. WO 9428028, each of which is incorporated herein by reference in its entirety], and the like can be used to screen for leafhopper ecdysone receptor ligands according to the present invention.

The screening can be performed with recombinant cells that express the leafhopper ecdysone receptor, or alternatively, using purified protein, e.g., produced recombinantly, as described above. For example, labeled, soluble leafhopper ecdysone receptor can be used to screen libraries, as described in the foregoing references.

In one embodiment, leafhopper ecdysone receptor may be directly labeled. In another embodiment, a labeled secondary reagent may be used to detect binding of a leafhopper ecdysone receptor to a molecule of interest, e.g., a molecule attached to a solid phase support. Binding may be detected by in situ formation of a chromophore by an enzyme label. Suitable enzymes include, but are not limited to, alkaline phosphatase and horseradish peroxidase. In a further embodiment, a two-color assay, using two chromogenic substrates with two enzyme labels on different acceptor molecules of interest, may be used. Cross-reactive and singly reactive ligands may be identified with a two-color assay.

Other labels for use in the invention include colored latex beads, magnetic beads, fluorescent labels (e.g., fluorescein isothiocyanate (FITC), phycoerythrin (PE), Texas red (TR), rhodamine, free or chelated lanthanide series salts, especially Eu^{3+} , to name a few fluorophores), chemiluminescent molecules, radio-isotopes, or magnetic resonance imaging labels. Two-color assays may be performed with two or more colored latex beads, or fluorophores that emit at different wavelengths. Labeled may be detected visually or by mechanical/optical means. Mechanical/optical means include fluorescence activated sorting, i.e., analogous to FACS, and micromanipulator removal means.

As exemplified herein, the level of a leafhopper ecdysone receptor polypeptide can be evaluated by metabolic labeling of the proteins. As the metabolic labeling occurs during in vitro incubation of the tissue biopsy in the presence of culture medium supplemented with [^{35}S]-methionine, the level of each of the markers detected may be affected by the in vitro conditions. In addition to metabolic (or biosynthetic) labeling with [^{35}S]-methionine, the invention further contemplates labeling with [^{14}C]-amino acids and [^3H]-amino acids (with the tritium substituted at non-labile positions). Thus, a sample or library of compounds can be directly analyzed after labeling of the proteins therein, e.g., by colorimetric staining using silver, gold, coomassie blue, or amido-schwartz, to mention a few techniques; isotopic labeling, e.g., with [^{32}P]-orthophosphate, [^{125}I], [^{131}I]; fluorescent or chemiluminescent tags; and immunological detection with labeled antibody or specific binding partner of a marker.

Modulating Insect Physiology or Development

The isolation of a leafhopper ecdysone receptor provides for isolation or screening of new ligands for receptor binding. Some of these will interfere with, or disrupt, normal insect development. It may sometimes be important to either accelerate or decelerate insect development, for instance, in preparing sterile adults for release. Alternatively, in certain circumstances, a delay or change in the timing of development may be lethal or may dramatically modify the ability of an insect to affect an agricultural crop. Thus, naturally occurring, biodegradable and highly active molecules to disrupt the timing of insect development will result.

The present invention provides a means for disrupting insect development where new ligand agonists or antagonists are discovered. These compounds are prime candidates as agonists or antagonists to interfere with the normal insect development. By application of new analogues of ligands for a leafhopper ecdysone receptor, it is possible to modify the normal temporal sequence of developmental events. For example, accelerating insect development will minimize generation time. This may be very important in circumstances where large numbers of insects are desired finally, for instance, in producing sterile males. Alternatively, it may be useful to slow development in a pest infestation, such that the insects reach destructive stages of development only after commercial crops may have passed sensitive stages. In another commercial application, ligands discovered by methods provided by the present invention may be used to artificially maintain insects in a specific developmental stage. The development of larvae may also be accelerated to reach a particular developmental stage in their life cycle earlier than naturally.

Other analogues of ligands for a leafhopper ecdysone receptor may be selected which, upon application, may be completely disruptive of normal development, leading to a lethal result and pest control. Indeed, there may be new ligands for a leafhopper ecdysone receptor which may be species specific or may exhibit a particularly useful spectrum of effectiveness. The greater specificity of the ligands will allow avoidance of use of non-specific pesticides possessing undesired deleterious ecological side effects. Furthermore, compounds having structures closely analogous to natural compounds may be susceptible to natural mechanisms of biological degradation.

Thus, the present invention also provides a method for identifying and selecting compounds exhibiting specific binding to the ligand binding domain to modulate insect physiology or development (e.g., killing) comprising the steps of screening compounds for binding to a homopteran ecdysone receptor, selecting compounds exhibiting said binding and administering the ligand to a homopteran insect. In a specific embodiment, a method for modulating insect physiology or development comprises the steps of screening compounds for binding to a leafhopper ecdysone receptor, selecting compounds exhibiting said binding and administering the ligand to a leafhopper.

Polypeptide Production

A purified leafhopper ecdysone receptor polypeptide of the invention is also useful in a method for determining the structural and biosynthetic aspects of the purified leafhopper ecdysone receptor polypeptide. Structural studies of interactions of the ligand-binding domains with selected ligands may be performed by various methods. The preferred method for structural determination is X-ray crystallography but may include various other forms of spectroscopy or chromatography. See, e.g., Connolly, M. L., *J. Appl. Crystall.* 16: 548

(1983); and Connolly, M. L., Science 221: 709 (1983), which are hereby incorporated herein by reference. For example, the structure of the interaction between ligand and ligand-binding domain may be determined to high resolution.

Having provided for the substantially pure polypeptides, biologically active fragments thereof and recombinant polynucleotides encoding them, the present invention also provides cells comprising each of them. By appropriate introduction techniques well known in the field, cells comprising them may be produced. See, e.g., Sambrook et al. (1989).

Host Cells and Non-Human Organisms

Another aspect of the present invention involves cells comprising an isolated polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the present invention. In a specific embodiment, the invention relates to an isolated host cell comprising a vector comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the present invention. The present invention also relates to an isolated host cell comprising an expression vector according to the invention. In another specific embodiment, the invention relates to an isolated host cell comprising a gene expression cassette comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the present invention. In another specific embodiment, the invention relates to an isolated host cell transfected with a gene expression modulation system comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the present invention. In another specific embodiment, the invention also provides an isolated host cell comprising an ecdysone receptor-based gene expression system comprising a leafhopper ecdysone receptor polypeptide according to the invention. In another specific embodiment, the invention relates to an isolated host cell comprising a leafhopper ecdysone receptor polypeptide of the present invention. In still another embodiment, the invention relates to a method for producing a leafhopper ecdysone receptor polypeptide, wherein the method comprises culturing an isolated host cell comprising a polynucleotide encoding a leafhopper ecdysone receptor polypeptide of the present invention in culture medium under conditions permitting expression of the polynucleotide encoding the leafhopper ecdysone receptor polypeptide, and isolating the leafhopper ecdysone receptor polypeptide from the culture.

As described above, the polypeptides of the present invention and the polynucleotides encoding them may be used to modulate gene expression in a host cell. Expression in transgenic host cells may be useful for the expression of various genes of interest. Applicants' invention provides for modulation of gene expression in prokaryotic and eukaryotic host cells. Expression in transgenic host cells is useful for the expression of various polypeptides of interest including but not limited to antigens produced in plants as vaccines, enzymes like alpha-amylase, phytase, glucanase, and xylanase, genes for resistance against insects, nematodes, fungi, bacteria, viruses, and abiotic stresses, antigens, nutraceuticals, pharmaceuticals, vitamins, genes for modifying amino acid content, herbicide resistance, cold, drought, and heat tolerance, industrial products, oils, protein, carbohydrates, antioxidants, male sterile plants, flowers, fuels, other output traits, therapeutic polypeptides, pathway intermediates; for the modulation of pathways already existing in the host for the synthesis of new products heretofore not possible using the host; cell based assays; functional genomics assays, biotherapeutic protein production, proteomics assays, and the like. Additionally the gene products may be useful for conferring higher growth yields of the host or for enabling an alternative growth mode to be utilized.

In a specific embodiment, the isolated host cell is a prokaryotic host cell or a eukaryotic host cell. In another specific embodiment, the isolated host cell is an invertebrate host cell or a vertebrate host cell. Preferably, the isolated host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a nematode cell, an insect cell, a fish cell, a plant cell, an avian cell, an animal cell, and a mammalian cell. More preferably, the isolated host cell is a yeast cell, a nematode cell, an insect cell, a plant cell, a zebrafish cell, a chicken cell, a hamster cell, a mouse cell, a rat cell, a rabbit cell, a cat cell, a dog cell, a bovine cell, a goat cell, a cow cell, a pig cell, a horse cell, a sheep cell, a simian cell, a monkey cell, a chimpanzee cell, or a human cell.

Examples of preferred host cells include, but are not limited to, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, or bacterial species such as those in the genera *Synechocystis*, *Synechococcus*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Rhodococcus*, *Streptomyces*, *Escherichia*, *Pseudomonas*, *Methylobacter*, *Methylobacter*, *Alcaligenes*, *Synechocystis*, *Anabaena*, *Thiobacillus*, *Methanobacterium* and *Klebsiella*; plant species selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat; animal; and mammalian host cells.

In a specific embodiment, the isolated host cell is a yeast cell selected from the group consisting of a *Saccharomyces*, a *Pichia*, and a *Candida* host cell.

In another specific embodiment, the isolated host cell is a *Caenorhabditis elegans* nematode cell.

In another specific embodiment, the isolated host cell is a plant cell selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat cell.

In another specific embodiment, the isolated host cell is a zebrafish cell.

In another specific embodiment, the isolated host cell is a chicken cell.

In another specific embodiment, the isolated host cell is a mammalian cell selected from the group consisting of a hamster cell, a mouse cell, a rat cell, a rabbit cell, a cat cell, a dog cell, a bovine cell, a goat cell, a cow cell, a pig cell, a horse cell, a sheep cell, a monkey cell, a chimpanzee cell, and a human cell.

Host cell transformation is well known in the art and may be achieved by a variety of methods including but not limited to electroporation, viral infection, plasmid/vector transfection, non-viral vector mediated transfection, *Agrobacterium*-mediated transformation, particle bombardment, and the like. Expression of desired gene products involves culturing the transformed host cells under suitable conditions and inducing expression of the transformed gene. Culture conditions and gene expression protocols in prokaryotic and eukaryotic cells are well known in the art (see General Methods section of Examples). Cells may be harvested and the gene products isolated according to protocols specific for the gene product.

In addition, a host cell may be chosen that modulates the expression of the transfected polynucleotide, or modifies and

processes the polypeptide product in a specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification [e.g., glycosylation, cleavage (e.g., of signal sequence)] of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce a non-glycosylated core protein product. However, a polypeptide expressed in bacteria may not be properly folded. Expression in yeast can produce a glycosylated product. Expression in eukaryotic cells can increase the likelihood of "native" glycosylation and folding of a heterologous protein. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, the polypeptide's activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent.

Applicants' invention also relates to a non-human organism comprising an isolated host cell according to the invention. In a specific embodiment, the non-human organism is a prokaryotic organism or a eukaryotic organism. In another specific embodiment, the non-human organism is an invertebrate organism or a vertebrate organism.

Preferably, the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a nematode, an insect, a fish, a plant, a bird, an animal, and a mammal. More preferably, the non-human organism is a yeast, a nematode, an insect, a plant, a zebrafish, a chicken, a hamster, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a cow, a pig, a horse, a sheep, a simian, a monkey, or a chimpanzee.

In a specific embodiment, the non-human organism is a yeast selected from the group consisting of *Saccharomyces*, *Pichia*, and *Candida*.

In another specific embodiment, the non-human organism is a *Caenorhabditis elegans* nematode.

In another specific embodiment, the non-human organism is a plant selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat.

In another specific embodiment, the non-human organism is a *Mus musculus* mouse.

Measuring Gene Expression/Transcription

One useful measurement of the methods of modulating gene expression using the novel polynucleotides, polypeptides, vectors, and/or gene expression cassettes of the present invention is that of the transcriptional state of the cell including the identities and abundances of RNA, preferably mRNA species. Such measurements are conveniently conducted by measuring cDNA abundances by any of several existing gene expression technologies.

Nucleic acid array technology is a useful technique for determining differential mRNA expression. Such technology includes, for example, oligonucleotide chips and DNA microarrays. These techniques rely on DNA fragments or oligonucleotides which correspond to different genes or cDNAs which are immobilized on a solid support and hybridized to probes prepared from total mRNA pools extracted from cells, tissues, or whole organisms and converted to cDNA. Oligonucleotide chips are arrays of oligonucleotides synthesized on a substrate using photolithographic tech-

niques. Chips have been produced which can analyze for up to 1700 genes. DNA microarrays are arrays of DNA samples, typically PCR products, that are robotically printed onto a microscope slide. Each gene is analyzed by a full or partial-length target DNA sequence. Microarrays with up to 10,000 genes are now routinely prepared commercially. The primary difference between these two techniques is that oligonucleotide chips typically utilize 25-mer oligonucleotides which allow fractionation of short DNA molecules whereas the larger DNA targets of microarrays, approximately 1000 base pairs, may provide more sensitivity in fractionating complex DNA mixtures.

Another useful measurement of Applicants' methods of the invention is that of determining the translation state of the cell by measuring the abundances of the constituent protein species present in the cell using processes well known in the art.

Where identification of genes associated with various physiological functions is desired, an assay may be employed in which changes in such functions as cell growth, apoptosis, senescence, differentiation, adhesion, binding to a specific molecule, binding to another cell, cellular organization, organogenesis, intracellular transport, transport facilitation, energy conversion, metabolism, myogenesis, neurogenesis, and/or hematopoiesis is measured.

In addition, selectable marker or reporter gene expression may be used to measure gene expression modulation using Applicants' invention.

Other methods to detect the products of gene expression are well known in the art and include Southern blots (DNA detection), dot or slot blots (DNA, RNA), northern blots (RNA), RT-PCR (RNA), western blots (polypeptide detection), and ELISA (polypeptide) analyses. Although less preferred, labeled proteins can be used to detect a particular nucleic acid sequence to which it hybridizes.

In some cases it is necessary to amplify the amount of a nucleic acid sequence. This may be carried out using one or more of a number of suitable methods including, for example, polymerase chain reaction ("PCR"), ligase chain reaction ("LCR"), strand displacement amplification ("SDA"), transcription-based amplification, and the like. PCR is carried out in accordance with known techniques in which, for example, a nucleic acid sample is treated in the presence of a heat stable DNA polymerase, under hybridizing conditions, with one pair of oligonucleotide primers, with one primer hybridizing to one strand (template) of the specific sequence to be detected. The primers are sufficiently complementary to each template strand of the specific sequence to hybridize therewith. An extension product of each primer is synthesized and is complementary to the nucleic acid template strand to which it hybridized. The extension product synthesized from each primer can also serve as a template for further synthesis of extension products using the same primers. Following a sufficient number of rounds of synthesis of extension products, the sample may be analyzed as described above to assess whether the sequence or sequences to be detected are present.

The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention.

EXAMPLES

General Molecular Biology Techniques

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g.,

Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (herein "Sambrook et al., 1989"); *DNA Cloning: A Practical Approach*, Volumes I and II (D. N. Glover ed. 1985); *Oligonucleotide Synthesis* (M. J. Gait ed. 1984); *Nucleic Acid Hybridization* [B. D. Hames & S. J. Higgins eds. (1985)]; *Transcription And Translation* [B. D. Hames & S. J. Higgins, eds. (1984)]; *Animal Cell Culture* [R. I. Freshney, ed. (1986)]; *Immobilized Cells And Enzymes* [IRL Press, (1986)]; B. Perbal, *A Practical Guide To Molecular Cloning* (1984); F. M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994).

Conventional cloning vehicles include pBR322 and pUC type plasmids and phages of the M13 series. These may be obtained commercially (Bethesda Research Laboratories).

For ligation, DNA fragments may be separated according to their size by agarose or acrylamide gel electrophoresis, extracted with phenol or with a phenol/chloroform mixture, precipitated with ethanol and then incubated in the presence of phage T4 DNA ligase (Biolabs) according to the supplier's recommendations.

The filling in of 5' protruding ends may be performed with the Klenow fragment of *E. coli* DNA polymerase I (Biolabs) according to the supplier's specifications. The destruction of 3' protruding ends is performed in the presence of phage T4 DNA polymerase (Biolabs) used according to the manufacturer's recommendations. The destruction of 5' protruding ends is performed by a controlled treatment with S1 nuclease.

Mutagenesis directed in vitro by synthetic oligodeoxynucleotides may be performed according to the method developed by Taylor et al. [*Nucleic Acids Res.* 13 (1985) 8749-8764] using the kit distributed by Amersham.

The enzymatic amplification of DNA fragments by PCR [Polymerase-catalyzed Chain Reaction, Saiki R. K. et al., *Science* 230 (1985) 1350-1354; Mullis K. B. and Faloona F. A., *Meth. Enzym.* 155 (1987) 335-350] technique may be performed using a "DNA thermal cycler" (Perkin Elmer Cetus) according to the manufacturer's specifications.

Verification of nucleotide sequences may be performed by the method developed by Sanger et al. [*Proc. Natl. Acad. Sci. USA* 74: 5463-5467 (1977)] using the kit distributed by Amersham.

Plasmid DNAs may be purified by the Qiagen Plasmid Purification System according to the manufacturer's instruction.

Example 1

This Example describes the cloning of full-length cDNA encoding green leafhopper *Nephotetix cincticeps* ecdysone receptor polypeptide. To isolate the full length coding sequence of this previously unknown leafhopper ecdysone receptor isoform (herein named "NcEcR"), Applicants used degenerate oligonucleotides (see Table 2) designed based on conserved regions at the end of A/B and E domains to amplify and clone NcEcR from cDNA prepared from RNA obtained from green leafhopper cells.

Polymerase chain reaction (PCR) amplification was performed using the Precision Taq polymerase kit (Stratagene) using the reaction conditions and cycling parameters as follows. PCR was performed using 1× reaction buffer (supplied by manufacturer), 50 ng of dsDNA template, 125 ng of forward primer (Primer 1), 125 ng of reverse complementary primer (Primer 2), and 1 μl of dNTP mix (supplied by manufacturer) in a final reaction volume of 50 μL. The cycling parameters used consisted of one cycle of denaturing at 95° C. for 30 seconds, followed by 35 cycles of denaturing at 94° C. for 1 minute, annealing at 50° C. for 1 minute, and extending at 72° C. for 2 minutes, followed by a final elongation cycle at 72° C. for 2 minutes.

The resulting PCR products were cloned into PCRII vector (Invitrogen), sequenced and the full-length coding sequence determined. The polynucleotide sequence encoding the full length NcEcR is presented as SEQ ID NO: 1. The deduced amino acid sequence of the full-length NcEcR is presented herein as SEQ ID NO: 2 and showed high similarity with the deduced amino acid sequence of other EcRs. As shown in Table 3, NcEcR showed highest similarity with locust, whitefly and mealworm EcRs. It is also interesting that NcEcR showed higher similarity with Crab and Tick EcRs when compared to lepidopteran and dipteran insect EcRs.

Applicants have aligned the polypeptide sequences of the NcEcR and EcRs from other species and identified the percent amino acid similarity as compared to the full-length NcEcR. The results are presented in Table 3.

TABLE 3

Comparison of leafhopper EcR (NcEcR) with other EcRs.	
EcR	Percent Amino Acid Similarity with NcEcR
LmEcR	78.3
BaEcR	75.1
TmEcR	75.9
AmaEcR	63.1
CpEcR	65.6
AaEcR	62.1
DmEcRB	61.0
LcEcR	62.3
CeEcR	61.2
CfEcRB	58.0
BmEcR	55.3
MsEcR	59.1
CtEcR	58.5
FXR	35.0
LXR	37.4
TR	28.7
VDR	25.7

Example 2

This Example describes the construction of leafhopper ecdysone receptor gene expression cassettes and their use in an ecdysone receptor-based gene expression modulation system. The results presented herein demonstrate that a leafhop-

Primer and SEQ ID NO: Primer Nucleic Acid Sequence

Primer 1 (SEQ ID NO: 3) 5'-CAA/G GAA/G GAA/G CTG/C TGT/C CTG/C GT-3'

Primer 2 (SEQ ID NO: 4) 5'-TCC CAA/G ATC/T TCT/C TCG/C AGG/A AA-3'

per ecdysone receptor is functional in an ecdysone receptor-based gene expression modulation system in both insect and mammalian cells.

A) Insect Cells:

Briefly, the NcEcR CDE domains (amino acids 6-370 of SEQ ID NO: 2) were fused to a VP16 transactivation domain (SEQ ID NO: 5) as follows. A construct was prepared by fusing a polynucleotide (nucleotides 16-1109 of SEQ ID NO: 1) encoding an NcEcR-CDE polypeptide to a polynucleotide (SEQ ID NO: 6) encoding a VP16 activation domain at the NH2 terminal end. This VP16NcEcR fusion was then cloned under the control of baculovirus IE1 promoter (SEQ ID NO: 7). The VP16NcEcR gene expression cassette was transfected into L57 cell line (*Drosophila* cell line that lacks endogenous EcR) and a cotton boll weevil ("CBW"; *Anthonomus grandis*) BRL-AG-2 cell line (generously provided by USDA, ARS, Bioscience Research Laboratory, Fargo, N. Dak.) along with a reporter construct EcRELaCZ that comprises a 6×ECRE response element (1×EcRE is shown in SEQ ID NO: 8), an ADH distal promoter (see Heberlein et al., 1985, Cell 41: 965-977 and Koelle et al. 1991, Cell 67: 59-77) and a LacZ reporter gene (SEQ ID NO: 9). The reporter gene activity was quantified in the presence of a series of concentrations of 20-hydroxyecdysone (20E) and N'-tert-butyl-N'-(3,5-dimethylbenzoyl)-3-methoxy-2-ethylbenzohydrazide (RH-123709) ligands.

Ligands: The steroid ligand 20-hydroxyecdysone (20E) was purchased from Sigma Chemical Company. The non-steroidal ligand N'-tert-butyl-N'-(3,5-dimethylbenzoyl)-3-methoxy-2-ethylbenzohydrazide (RH-123709) is a synthetic stable ecdysteroid ligand synthesized at Rohm and Haas Company. The ligands were dissolved in DMSO and the final concentration of DMSO was maintained at 0.1% in both controls and treatments.

Transfections: DNAs corresponding to the gene constructs described above were transfected into L57 or the CBW cells as follows. L57 cells were grown in HyQ-CCM3 medium (Hyclone labs) and transfected with lipofectomine (Life Technologies). The CBW cells were grown in Ex-Cell 401 (JRH Sciences) and transfected with Celfectin (Invitrogen). Standard methods for culture and maintenance of the cells were followed. Cells were harvested when they reached 50% confluency and plated in 6-, 12- or 24-well plates at 125,000, 50,000, or 25,000 cells, respectively, in 2.5, 1.0, or 0.5 ml of growth medium, respectively. The next day, the cells were rinsed with growth medium and transfected for four hours. For 12-well plates, 4 µl of the appropriate transfection reagent was mixed with 100 µl of growth medium. One µg of reporter construct and 0.25 µg of each receptor construct of the receptor pair to be analyzed were added to the transfection mix. A second reporter construct was added [pTKRL (Promega), 0.1 µg/transfection mix] that comprises a Renilla luciferase gene operably linked and placed under the control of a thymidine kinase (TK) constitutive promoter and was used for normalization. The contents of the transfection mix were mixed in a vortex mixer and let stand at room temperature for 30 minutes. At the end of incubation, the transfection mix was added to the cells maintained in 400 µl growth medium. The cells were maintained at 37° C. and 5% CO₂ for four hours. At the end of incubation, 500 µl of growth medium and either dimethylsulfoxide (DMSO; control) or a DMSO solution of 0, 0.001, 0.01, 0.1, 1.0, 10, or 100 µM steroidal ligand or non-steroidal ligand was added and the cells were maintained at 37° C. and 5% CO₂ for 48 hours. The cells were harvested and reporter activity was assayed. The same procedure was followed for 6 and

24 well plates as well except all the reagents were doubled for 6 well plates and reduced to half for 24-well plates.

Reporter Assays: Cells were harvested 48 hours after adding ligand. 125 µl of passive lysis buffer part of Dual-luciferase™ reporter assay system from Promega Corporation) were added to each well of the 24-well plate. The plates were placed on a rotary shaker for 15 minutes. Twenty µl of lysate were assayed. Luciferase activity was measured using Dual-luciferase™ reporter assay system from Promega Corporation following the manufacturer's instructions. β-Galactosidase was measured using Galacto-Star™ assay kit from TROPIX following the manufacturer's instructions. All luciferase and β-galactosidase activities were normalized using Renilla luciferase as a standard. Fold activities were calculated by dividing normalized relative light units ("RLU") in ligand treated cells with normalized RLU in DMSO treated cells (untreated control).

As shown in FIG. 1, the NcEcR construct was able to transactivate reporter gene activity in a dose-dependent manner with both ligands tested in the CBW cells. However, in L57 cells there was very little transactivation in the presence of either ligand. Applicants' previous studies have shown that both CfEcR and DmEcR cause good transactivation in L57 but NcEcR was a poor transactivator in these cells.

B) Mammalian Cells:

Briefly, the NcEcR DE domains (amino acids 72-370 of SEQ ID NO: 2) were fused to a GAL4 DNA binding domain (SEQ ID NO: 10) as follows. A construct was prepared by fusing a polynucleotide (nucleotides 214-1109 of SEQ ID NO: 1) encoding an NcEcR-DE polypeptide to a polynucleotide (SEQ ID NO: 11) encoding a GAL4 DNA-binding domain at the NH2 terminal end. This GAL4/NcEcR fusion was then cloned under the control of a cytomegalovirus (CMV) promoter (SEQ ID NO: 12). In addition, a polynucleotide encoding the EF domains of seven RXR/USPs from a moth *Choristoneura fumiferana* ultraspiracle protein ("CfUSP", SEQ ID NO: 13), a fruit fly *Drosophila melanogaster* ultraspiracle protein ("DmUSP", SEQ ID NO: 14), a locust *Locusta migratoria* ultraspiracle protein (LmUSP; SEQ ID NO: 15), a mouse *Mus musculus* retinoid X receptor isoform α (MmRXRα; SEQ ID NO: 16), a chimeric RXR/USP between MmRXRα and LmUSP (Chimera; SEQ ID NO: 17), a tick *Amblyomma americanum* retinoid X receptor homolog 1 (AmaRXR1; SEQ ID NO: 18), and a tick *Amblyomma americanum* retinoid X receptor homolog 2 (AmaRXR2; SEQ ID NO: 19) were each fused to a polynucleotide (SEQ ID NO: 6) encoding a VP16 activation domain.

The GAL4/NcEcR gene expression cassette was transfected into NIH3T3 cells (ATCC) along with each of the seven VP16RXR/USP gene expression cassettes and a reporter construct pFRLuc that comprises a 5×GAL4RE (1×GAL4RE is shown in SEQ ID NO: 20), a synthetic TATAA (SEQ ID NO: 21) and a luciferase reporter gene (SEQ ID NO: 22) as described above except the cells were cultured in growth media comprising 10% fetal bovine serum (FBS), Superfect™ (Qiagen Inc.) was used as the transfection reagent, and at the end of incubation/transfection, 500 µl of growth medium containing 20% FBS was added to the cells.

The receptor combinations were compared for their ability to transactivate pFRLuc in NIH3T3 cells in the presence of 0, 0.2, 1.0, or 10 µM steroid ligand ponasteroneA (PonA; Invitrogen) or 0, 0.04, 0.2, 1.0, or 10 µM GST™-E. The non-steroidal ligand N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine (GST™-E) is a

synthetic stable ecdysteroid ligand synthesized at Rohm and Haas Company. The ligand was dissolved in DMSO and the final concentration of DMSO was maintained at 0.1% in both controls and treatments. The results are presented in FIG. 2 and the numbers on the top of each bar show the maximum fold induction for that group.

As shown in FIG. 2, NcEcR in combination with MmRXR α , AmaRXR1, or AmaRXR2 supports induction of reporter activity. NcEcR in combination with CfUSP, DmUSP, Chimera or LmUSP produced lower fold induction mainly due to increased background activity of the reporter in the absence of ligand.

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 1109

<212> TYPE: DNA

<213> ORGANISM: *Nephotetix cincticeps*

<400> SEQUENCE: 1

```

caggaggagc tctgctgtt gtgcggagac cgagcgtcgg gataccacta caacgctctc      60
acctgcgaag gatcaaggg cttctttcgg aggagtatca ccaaaaacgc agtgtaccag      120
tccaaatacg gcaccaattg tgaatatagac atgtatatgc ggcgcaagtg ccaggagtgc      180
cgactcaaga agtgctcag tgtagggatg aggccagaat gtgtagtacc tgagtatcaa      240
tgtgcccgtaa aaaggaaaga gaaaaagct caaaaggaca aagataaacc tgtctcttca      300
accaatggct cgctgaaat gagaatagac caggacaacc gttgtgtggt gttgcagagt      360
gaagacaaca ggtacaactc gactacgcc agtttcggag tcaaacccct cagtccagaa      420
caagaggagc tcctccacag gctcgtctac ttccagaacg agtacgaaca ccctgccgag      480
gaggatctca agcggatcga gaacctcccc tgtgacgacg atgacctgtg tgatgttcgc      540
tacaacacac ttacggagat cacaatactc acagtccagc tcctcgtgga gtttgcgaaa      600
aaactgectg gtttcgacaa actactgaga gaggaccaga tcgtgttctc caaggcgtgt      660
tcgagcgagg tgatgatgct gcggatggcg cggaggtacg acgtccagac agactcgatc      720
ctgttcgcca acaaccagcc gtacacgoga gactcgtaca cgatggcagg cgtgggggaa      780
gtcatcgaag atctgctcgc gttcggccga ctcatgtgct ccatgaaggt ggacaatgcc      840
gagtatgctc tgctcacggc catcgtcacc ttctccgagc ggccgaacct ggccgaagga      900
tggaaggttg agaagatcca ggagatctac ctggaggcgc tcaagtccta cgtggacaac      960
cgagtgaaac ctgcagctcc gaccatcttc gccaaactgc tctccttct caccgagctg     1020
cgaacactcg gcaaccagaa ctccgagatg tgcttctcgt taaactacgc aaccgcaaac     1080
atgccaccgt tcctcgaaga aatctggga                                     1109
  
```

<210> SEQ ID NO 2

<211> LENGTH: 370

<212> TYPE: PRT

<213> ORGANISM: *Nephotetix cincticeps*

<400> SEQUENCE: 2

```

Gln Glu Glu Leu Cys Leu Leu Cys Gly Asp Arg Ala Ser Gly Tyr His
1           5           10           15
Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser
  
```

-continued

20					25					30					
Ile	Thr	Lys	Asn	Ala	Val	Tyr	Gln	Ser	Lys	Tyr	Gly	Thr	Asn	Cys	Glu
		35					40					45			
Ile	Asp	Met	Tyr	Met	Arg	Arg	Lys	Cys	Gln	Glu	Cys	Arg	Leu	Lys	Lys
	50					55					60				
Cys	Leu	Ser	Val	Gly	Met	Arg	Pro	Glu	Cys	Val	Val	Pro	Glu	Tyr	Gln
65						70					75				80
Cys	Ala	Val	Lys	Arg	Lys	Glu	Lys	Lys	Ala	Gln	Lys	Asp	Lys	Asp	Lys
				85					90					95	
Pro	Val	Ser	Ser	Thr	Asn	Gly	Ser	Pro	Glu	Met	Arg	Ile	Asp	Gln	Asp
			100					105					110		
Asn	Arg	Cys	Val	Val	Leu	Gln	Ser	Glu	Asp	Asn	Arg	Tyr	Asn	Ser	Ser
		115					120					125			
Thr	Pro	Ser	Phe	Gly	Val	Lys	Pro	Leu	Ser	Pro	Glu	Gln	Glu	Glu	Leu
	130					135					140				
Ile	His	Arg	Leu	Val	Tyr	Phe	Gln	Asn	Glu	Tyr	Glu	His	Pro	Ala	Glu
145						150					155				160
Glu	Asp	Leu	Lys	Arg	Ile	Glu	Asn	Leu	Pro	Cys	Asp	Asp	Asp	Asp	Pro
				165					170					175	
Cys	Asp	Val	Arg	Tyr	Lys	His	Ile	Thr	Glu	Ile	Thr	Ile	Leu	Thr	Val
			180					185					190		
Gln	Leu	Ile	Val	Glu	Phe	Ala	Lys	Lys	Leu	Pro	Gly	Phe	Asp	Lys	Leu
		195					200					205			
Leu	Arg	Glu	Asp	Gln	Ile	Val	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val
	210					215					220				
Met	Met	Leu	Arg	Met	Ala	Arg	Arg	Tyr	Asp	Val	Gln	Thr	Asp	Ser	Ile
225						230					235				240
Leu	Phe	Ala	Asn	Asn	Gln	Pro	Tyr	Thr	Arg	Glu	Ser	Tyr	Thr	Met	Ala
				245					250					255	
Gly	Val	Gly	Glu	Val	Ile	Glu	Asp	Leu	Leu	Arg	Phe	Gly	Arg	Leu	Met
			260					265					270		
Cys	Ser	Met	Lys	Val	Asp	Asn	Ala	Glu	Tyr	Ala	Leu	Leu	Thr	Ala	Ile
		275					280						285		
Val	Ile	Phe	Ser	Glu	Arg	Pro	Asn	Leu	Ala	Glu	Gly	Trp	Lys	Val	Glu
	290					295					300				
Lys	Ile	Gln	Glu	Ile	Tyr	Leu	Glu	Ala	Leu	Lys	Ser	Tyr	Val	Asp	Asn
305						310					315				320
Arg	Val	Lys	Pro	Arg	Ser	Pro	Thr	Ile	Phe	Ala	Lys	Leu	Leu	Ser	Val
				325					330					335	
Leu	Thr	Glu	Leu	Arg	Thr	Leu	Gly	Asn	Gln	Asn	Ser	Glu	Met	Cys	Phe
			340					345					350		
Ser	Leu	Asn	Tyr	Ala	Thr	Ala	Asn	Met	Pro	Pro	Phe	Leu	Glu	Glu	Ile
		355					360					365			
Trp	Asp														
	370														

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Degenerate ECR PCR primer
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (3)..(3)
 <223> OTHER INFORMATION: n = a or g

-continued

```

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: n = a or g
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: n = a or g
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: n = g or c
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: n = t or c
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: n = g or c

```

```

<400> SEQUENCE: 3

```

```

canganganc tntgnctngt

```

20

```

<210> SEQ ID NO 4
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Degenerate EcR PCR primer
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: n = a or g
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: n = c or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: n = t or c
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: n = g or c
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: n = g or a

```

```

<400> SEQUENCE: 4

```

```

tcccanatnt cntcnagnaa

```

20

```

<210> SEQ ID NO 5
<211> LENGTH: 90
<212> TYPE: PRT
<213> ORGANISM: herpes simplex virus 7

```

```

<400> SEQUENCE: 5

```

```

Met Gly Pro Lys Lys Lys Arg Lys Val Ala Pro Pro Thr Asp Val Ser
 1             5             10             15

Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His
           20           25           30

Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp
 35           40           45

Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala
 50           55           60

Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu
 65           70           75           80

```

-continued

Gly Ile Asp Glu Tyr Gly Gly Glu Phe Pro
85 90

<210> SEQ ID NO 6

<211> LENGTH: 681

<212> TYPE: DNA

<213> ORGANISM: herpes simplex virus 7

<400> SEQUENCE: 6

```
atgggcccta aaaagaagaa gcgtaaggtc aaagcgtaa cggccaggct tgaattaatt    60
ccgggcggaa tgaagcggtt aacggccagg caacaagagg tgtttgatct catccgtgat    120
cacatcagcc agacagggat gccgccgacg cgtgcggaaa tcgcgcagcg tttggggttc    180
cgttccccaa acgcggttga agaacatctg aaggcgctgg cacgcaaagg cgttattgaa    240
attgtttccg gcgcatcacg cgggattcgt ctgttgacgg aagaggaaga agggttgccc    300
ctggtaggtc gtgtggctgc cgggtaacca cttctggcgc aacagcatat tgaaggtcac    360
tatcaggtcg atccttcctt attcaagccg aatgctgatt tcctgctgcg cgtcagccgg    420
atgtcgatga aagataticg cattatggat ggtgacttgc tggcagtgca taaaactcag    480
gatgtacgta acggtcaggc cgttgtcgca cgtattgatg acgaagtac cgtaaagcgc    540
ctgaaaaaac agggcaataa agtcgaaactg ttgccagaaa atagcgagtt taaaccaatt    600
gtcgtagatc ttcgtcagca gagcttcacc attgaagggc tggcggttgg ggttattcgc    660
aacggcgact ggctggaatt c                                681
```

<210> SEQ ID NO 7

<211> LENGTH: 703

<212> TYPE: DNA

<213> ORGANISM: Baculovirus

<400> SEQUENCE: 7

```
catatgtatc ccgggccagt tgcacaacac tattatcgat ttgcagttcg ggacataaat    60
gtttaaatat atcgatgtct ttgtgatgcg cgcgacattt ttgtaggtta ttgataaaat    120
gaacggatac gttgcccgcac attatcatta aatccttggc gtagaatttg tcgggtccat    180
tgtccgtgtg cgetagcatg cccgtaaccg acctcgtact tttggcttca aaggttttgc    240
gcacagacaa aatgtgccac acttgcagct ctgcatgtgt gcgcggtacc acaaatccca    300
acggcgagct gtacttgttg tatgcaaata aatctcgata agggcgcggc gcgcgaatgc    360
agctgatcac gtacgctcct cgtgttccgt tcaaggacgg tgttatcgac ctcagattaa    420
tgtttatcgg ccgactgttt tcgatatccg tcaccaaacg cgtttttgca ttaacattgt    480
atgtcggcgg atgttctata tctaatttga ataaataaac gataaccgcg ttggttttag    540
agggcataat aaaagaaata ttgttatcgt gttcgccatt agggcagtat aaattgacgt    600
tcatgttggg tattgtttca gttgcaagtt gacactggcg gcgacaagat cgtgaacaac    660
caagtgacta tagaattcac tcgaggctag cataagatct aag                                703
```

<210> SEQ ID NO 8

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Drosophila melanogaster

<400> SEQUENCE: 8

```
gagacaaggg ttcaatgcac ttgtccaatg                                30
```

-continued

```

<210> SEQ ID NO 9
<211> LENGTH: 3157
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 9
atgggggggtt ctcatcatca tcatcatcat ggtatggcta gcatgactgg tggacagcaa      60
atgggtcggg atctgtacga cgatgacgat aaggtaaccta aggatcagct tggagttagt      120
cccgtcgttt tacaacgtcg tgactgggaa aaccctggcg ttaccaact taatcgccct      180
gcagcacatc cccctttcgc cagctggcgt aatagcgaag aggcccgcac cgatcgccct      240
tcccaacagt tgcgcagcct gaatggcgaa tggcgctttg cctggtttcc ggcaccagaa      300
gcggtgcccg aaagctggct ggagtgcgat ctctctgagg ccgatactgt cgtcgtcccc      360
tcaaactggc agatgcacgg ttacgatcgc cccatctaca ccaacgtaac ctatcccatt      420
acggtcaate cgcctttgtt tcccacggag aatccgacgg gttgttactc gctcacattt      480
aatgttgatg aaagctggct acaggaaggc cagacgcgaa ttattttga tggcgtaaac      540
tcggcgtttc atctgtggtg caacggggcg tgggtcggtt acggccagga cagtcgtttg      600
ccgtctgaat ttgacctgag cgcattttta cgcgccggag aaaaccgcct cgcggtgatg      660
gtgctgcggt ggagtgcacg cagttatctg gaagatcagg atatgtggcg gatgagcggc      720
attttcgctg acgtctcgtt gctgcataaa ccgactacac aatcagcga tttccatggt      780
gccactcgct ttaatgatga tttcagccgc gctgtactgg aggctgaagt tcagatgtgc      840
ggcgagttag gtgactacct acgggtaaca gtttctttat ggcagggtag aacgcaggtc      900
gccagcggca ccgcgccttt cggcggtgaa attatcgatg agcgtggtgg ttatgcccgat      960
cgcgtcacac tacgtctgaa cgtcgaaaaa ccgaaactgt ggagcgcgca aatcccgaat     1020
ctctatcgtg cgggtggtga actgcacacc gccgacggca cgctgattga agcagaagcc     1080
tgcgatgtcg gtttcgcgca ggtgcccatt gaaaatggtc tgctgctgct gaacggcaag     1140
ccgttgctga ttcgagggct taaccgtcac gagcatcatc ctctgcatgg tcaggtcatg     1200
gatgagcaga cgatggtgca ggatatacctg ctgatgaagc agaacaactt taacgcccgtg     1260
cgctgttcgc attatccgaa ccatccgctg tggtagacgc tgtgcgaccg ctacggcctg     1320
tatgtggtgg atgaagccaa tattgaaacc cacggcatgg tgccaatgaa tcgtctgacc     1380
gatgatccgc gctggctacc ggcgatgagc gaacgcgtaa cgcgaatggt gcagcgcgat     1440
cgtaatcacc cgagtgtgat catctggtcg ctggggaatg aatcaggcca cggcgcataat     1500
cacgacgcgc tgtatcgtg gatcaaatct gtcgatcctt cccgcccggg gcagtatgaa     1560
ggcggcggag ccgacaccac ggccaccgat attatttgcc cgatgtacgc gcgcgtggat     1620
gaagaccagc ctttcccggc tgtgcogaaa tgggccatca aaaaatggct ttcgctacct     1680
ggagagacgc gcccgtgat cctttgcgaa tacgcccacg cgatgggtaa cagtcttggc     1740
ggtttcgcta aatactggca ggcgtttcgt cagtatcccc gtttacaggg cggcttcgtc     1800
tgggactggg tggatcagtc gctgattaaa tatgatgaaa acggcaaccc gtggtcggct     1860
tacggcgggt attttgcgca tacgcogaac gatcgcagc tctgtatgaa cggctctggtc     1920
tttgccgacc gcacgcgca tccagcgtg acggaagcaa aacaccagca gcagtttttc     1980
cagttccggt tatccgggca aaccatcgaa gtgaccagcg aatacctggt ccgtcatagc     2040
gataacgagc tcctgcactg gatggtggcg ctggatgta agccgctggc aagcggtgaa     2100
gtgcctctgg atgtcgtcc acaaggtaaa cagttgattg aactgcctga actaccgcag     2160

```

-continued

```

ccggagagcg cggggaact ctggtcaca gtacgcgtag tgcaaccgaa cgcgaccgca 2220
tggtcagaag cggggcacat cagcgcctgg cagcagtggc gtctggcgga aaacctcagt 2280
gtgacgctcc ccgcccgcgc ccacgccatc ccgcatctga ccaccagcga aatggatttt 2340
tgcctcgagc tgggtaataa gcggttgcaa ttaaccgcc agtcaggctt tctttcacag 2400
atgtggattg gcgataaaaa acaactgctg acgcccgtgc gcgatcagtt caccctgca 2460
ccgctggata acgacattgg cgtaagtga gcgaccgca ttgaccctaa cgectgggtc 2520
gaacgctgga agggcgcggg ccattaccag gccgaagcag cgttggtgca gtgcacggca 2580
gatacacttg ctgatgcggt gctgattacg accgctcacg cgtggcagca tcaggggaaa 2640
accttattta tcagccggaa aacctaccgg attgatgta gtggtcaaat ggcgattacc 2700
gttgatgttg aagtggcgag cgataaccg catccggcgc ggattggcct gaactgccag 2760
ctggcgcagg tagcagagcg ggtaactgg ctcggattag ggccgcaaga aactatccc 2820
gaccgcctta ctgcccctg ttttgaccgc tgggatctgc cattgtcaga catgtatacc 2880
ccgtacgtct tcccagcga aaacggctg cgctgcggga cgcgcgaatt gaattatggc 2940
ccacaccagt ggcgcgcgca cttccagtc aacatcagcc gctacagtca acagcaactg 3000
atggaacca gccatgccca tctgtgcac gcggaagaag gcacatggct gaatatcgac 3060
ggtttccata tggggattgg tggcgacgac tcttgagcc cgctcagtc ggcggaatta 3120
cagctgagcg ccggtcgcta ccattaccag ttggtct 3157

```

<210> SEQ ID NO 10

<211> LENGTH: 147

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 10

```

Met Lys Leu Leu Ser Ser Ile Glu Gln Ala Cys Asp Ile Cys Arg Leu
1           5           10          15
Lys Lys Leu Lys Cys Ser Lys Glu Lys Pro Lys Cys Ala Lys Cys Leu
20          25          30
Lys Asn Asn Trp Glu Cys Arg Tyr Ser Pro Lys Thr Lys Arg Ser Pro
35          40          45
Leu Thr Arg Ala His Leu Thr Glu Val Glu Ser Arg Leu Glu Arg Leu
50          55          60
Glu Gln Leu Phe Leu Leu Ile Phe Pro Arg Glu Asp Leu Asp Met Ile
65          70          75          80
Leu Lys Met Asp Ser Leu Gln Asp Ile Lys Ala Leu Leu Thr Gly Leu
85          90          95
Phe Val Gln Asp Asn Val Asn Lys Asp Ala Val Thr Asp Arg Leu Ala
100         105         110
Ser Val Glu Thr Asp Met Pro Leu Thr Leu Arg Gln His Arg Ile Ser
115        120        125
Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys Gly Gln Arg Gln Leu
130        135        140
Thr Val Ser
145

```

<210> SEQ ID NO 11

<211> LENGTH: 441

<212> TYPE: DNA

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 11

-continued

```

atgaagctac tgtcttctat cgaacaagca tgcgatattt gccgacttaa aaagctcaag    60
tgctccaaag aaaaaccgaa gtgcgccaaag tgtctgaaga acaactggga gtgtcgctac    120
tctcccaaaa caaaaggctc tccgctgact agggcacatc tgacagaagt ggaatcaagg    180
ctagaaagac tggaacagct atttctactg atttttcttc gagaagacct tgacatgatt    240
ttgaaaatgg attctttaca ggatataaaa gcattgtaa caggattatt tgtacaagat    300
aatgtgaata aagatgccgt cacagataga ttggcttcag tggagactga tatgcctcta    360
acattgagac agcatagaat aagtgcgaca tcatcatcgg aagagagtag taacaaaggt    420
caaagacagt tgactgtatc g                                     441

```

```

<210> SEQ ID NO 12
<211> LENGTH: 750
<212> TYPE: DNA
<213> ORGANISM: Cytomegalovirus

```

```

<400> SEQUENCE: 12

```

```

tcaatattgg ccattagcca tattattcat tggttatata gcataaatca atattggcta    60
ttggccattg catacgttgt atctatatca taatatgtac atttatattg gctcatgtcc    120
aatatgaccg ccatgttggc attgattatt gactagttat taatagtaat caattacggg    180
gtcattagtt catagcccat atatggagtt ccgcgttaca taacttacgg taaatggccc    240
gcctggctga ccgcccaacg acccccggcc attgacgtca ataatgacgt atgttcccat    300
agtaacgccca atagggactt tccattgacg tcaatgggtg gagtatttac ggtaaactgc    360
ccacttgcca gtacatcaag tgtatcatat gccaaagtcg cccctattg acgtcaatga    420
cggtaaatgg cccgcctggc attatgccca gtacatgacc ttacgggact ttcctacttg    480
gcagtacatc tacgtattag tcatcgtat taccatgggt atgcggtttt ggcagtacac    540
caatgggcgt ggatagcggg ttgactcacg gggatttcca agtctccacc ccattgacgt    600
caatgggagt ttgttttggc accaaaatca acgggacttt caaaatgtc gtaacaactg    660
cgatcgcctg ccccgctgac gcaaatgggc ggtaggcgtg tacgggtggga ggtctatata    720
agcagagctc gtttagtgaa ccgtcagatc                                     750

```

```

<210> SEQ ID NO 13
<211> LENGTH: 798
<212> TYPE: DNA
<213> ORGANISM: Choristoneura fumiferana

```

```

<400> SEQUENCE: 13

```

```

tcggtgcagg taagcgatga gctgtcaatc gagcgcctaa cggagatgga gtctttgggtg    60
gcagatccca gcgaggagtt ccagttcctc cgcgtggggc ctgacagcaa cgtgcctcca    120
cgttaccgcg cgcctgtctc ctccctctgc caaataggca acaagcaaat agcggcgttg    180
gtggatggg cgcgcgacat cctcatttc gggcagctgg agctggacga tcaagtggta    240
ctcatcaagg cctcctggaa tgagctgcta ctcttcgcca tcgctggcg ctetatggag    300
tatttggaa atgagagggg gaacggggac ggaacgcgga gcaccactca gccacaactg    360
atgtgtctca tgcttggcat gacgttgcac cgcaactcgg cgcagcaggc gggcgtgggc    420
gccatcttcg accgcgtgct gtccgagctc agtctgaaga tgcgcacctt gcgcatggac    480
caggccgagt acgtcgcgct caaagccatc gtgctgctca accctgatgt gaaaggactg    540
aagaatcggc aagaagttag cgttttgcca gaaaaaatgt tctcttgctt ggacgactac    600

```

-continued

```

tgccggcggg cgcaagcaa cgaggaaggg cggtttgctg ccttgctgct gcggtgcca 660
gctctccgct caatctcgct caagagcttc gaacacctct acttcttcca cctcgtggcc 720
gaaggctcca tcagcggata catacagagag gcgctccgaa accacgcgcc tccgatcgac 780
gtcaatgcca tgatgtaa 798

```

```

<210> SEQ ID NO 14
<211> LENGTH: 799
<212> TYPE: DNA
<213> ORGANISM: Drosophila melanogaster

```

```

<400> SEQUENCE: 14

```

```

catagaggcc gagcagcgag cggagaccca atgcggcgat cgtgcaactga cgttcctgcg 60
cgttggtccc tattccacag tccagccgga ctacaagggt gccgtgtcgg cctcgtgcca 120
agtggccaac aaacagctct tccagatggt cgaatacgcg cgcgatgatgc cgcactttgc 180
ccaggtgccg ctggacgacc aggtgattct gctgaaagcc gcttggatcg agctgctcat 240
tgcgaacgtg gcttggtgca gcatcgttct gctggatgac ggcgggtgccg gcggcggggg 300
cggtggaacta ggccacgatg gctcctttga gcgacgatca ccgggccttc agccccagca 360
gctgttcctc aaccagagct tctcgtacca tcgcaacagt gcgatcaaag ccgggtgtgc 420
agccatcttc gaccgcatac tgtcggagct gagtgtaaag atgaagcggc tgaatctcga 480
ccgacgcgag ctgtcctgct tgaagccat catactgtac aaccgggaca tacgcgggat 540
caagagccgg gcggagatcg agatgtgccg cgagaaggty tacgcttgcc tggacgagca 600
ctgccgctg gaacatccgg gcgacgatgg acgctttgcy caactgctgc tgcgtctgcc 660
cgctttgcyg tcgatecagcc tgaagtgccg ggatcacctg ttctcttcc gcattaccag 720
cgaccggccg ctggaggagc tctttctcga gcagctggag gcgccgcgc caaccggcct 780
ggcggatgaaa ctggagtag 799

```

```

<210> SEQ ID NO 15
<211> LENGTH: 635
<212> TYPE: DNA
<213> ORGANISM: Locusta migratoria

```

```

<400> SEQUENCE: 15

```

```

tgcatacaga catgctggtt gaacgcatac ttgaagctga aaaacgagtg gagtgcaaag 60
cagaaaaacca agtggaaat gagctggtgg agtgggctaa acacatcccg cacttcacat 120
ccctacctct ggaggaccag gttctcctcc tcagagcagg ttggaatgaa ctgctaattg 180
cagcattttc acatgatctc gtagatggtt aagatggcat agtacttgcc actgggtctca 240
cagtgcatcg aaattctgcc catcaagctg gagtggcac aatatttgac agagttttga 300
cagaactggt agcaaagatg agagaaatga aaatggataa aactgaactt ggctgcttgc 360
gatctggtat tcttttcaat ccagaggatg ggggtttgaa atccgccag gaagttgaac 420
ttctacgtga aaaagtatat gccgctttgg aagaatatac tagaacaaca catcccgatg 480
aaccaggaag atttgcaaaa cttttgcttc gtctgccttc tttacgttcc ataggcetta 540
agtgtttga gcatttggtt ttctttcgcc ttattggaga tgttccaatt gatagcttcc 600
tgatggagat gcttgaatca ccttctgatt cataa 635

```

```

<210> SEQ ID NO 16
<211> LENGTH: 714
<212> TYPE: DNA
<213> ORGANISM: Mus musculus

```


-continued

<400> SEQUENCE: 16

```

gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag    60
actgagacat acgtggaggc aaacatgggg ctgaacccca gctcaccaaa tgaccctggt    120
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg    180
atccacactc tttctgagct gcccctagac gaccaggtea tcctgctacg ggcaggctgg    240
aacgagctgc tgatgcctc cttctcccac cgctccatag ctgtgaaaga tgggattctc    300
ctggccaccg gcctgcacgt acaccggaac agcgctcaca gtgctggggg gggcgccatc    360
ttgacagggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg    420
gagctgggct gcctgcgagc cattgtcctg ttcaaccctg actctaaggg gctctcaaac    480
cctgctgagg tggaggcgtt gagggagaag gtgtatgctg cactagaagc gtactgcaaa    540
cacaagtacc ctgagcagcc gggcaggttt gccaaactgc tgctccgctc gcctgcactg    600
cgttccatcg ggctcaagtg cctggagcac ctgttcttct tcaagctcat cggggacacg    660
cccacgaca cttctctcat ggagatgctg gaggcaccac atcaagccac ctag          714

```

<210> SEQ ID NO 17

<211> LENGTH: 711

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chimeric MmRXRalpha/LmUSP-EF

<400> SEQUENCE: 17

```

gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag    60
actgagacat acgtggaggc aaacatgggg ctgaacccca gctcaccaaa tgaccctggt    120
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg    180
atccacactc tttctgagct gcccctagac gaccaggtea tcctgctacg ggcaggctgg    240
aacgagctgc tgatgcctc cttctcccac cgctccatag ctgtgaaaga tgggattctc    300
ctggccaccg gcctgcacgt acaccggaac agcgctcaca gtgctggggg gggcgccatc    360
ttgacagggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagact    420
gaacttggtc gcttgcgatc tgttattctt ttcaatccag aggtgagggg tttgaaatcc    480
gcccaggaag ttgaacttct acgtgaaaaa gtatatgccg ctttgaaga atatactaga    540
acaacacate ccgatgaacc aggaagattt gcaaaaacttt tgcttcgtct gccttcttta    600
cgttccatag gccttaagtg tttggagcat ttgtttttct ttcgccttat tggagatggt    660
ccaattgata cgttctgatg ggagatgctt gaatcacctt ctgattcata a          711

```

<210> SEQ ID NO 18

<211> LENGTH: 687

<212> TYPE: DNA

<213> ORGANISM: Amblyomma americanum

<400> SEQUENCE: 18

```

cctcctgaga tgctctgga gcgcatactg gaggcagagc tgcgggttga gtcacagacg    60
gggaccctct cggaaagcgc acagcagcag gatccagtga gcagcatctg ccaagctgca    120
gaccgacagc tgcaccagct agttcaatgg gccaaagcaca ttccacattt tgaagagctt    180
ccccttgagg accgcatggt gttgctcaag gctggctgga acgagctgct cattgtgctt    240
ttctcccacc gttctgttga cgtgcgtgat ggcattgtgc tcgctacagg tcttgtggtg    300

```

-continued

```

cagcggcata gtgctcatgg ggctggcggt ggggccatat ttgataggg tctcaactgaa 360
ctggtagcaa agatgcgtga gatgaagatg gaccgcactg agcttgatg cctgcttgc 420
gtggtacttt ttaatcctga ggccaagggg ctgcgacact gcccaagtgg aggccctgag 480
ggagaaaagt tatctgcctt ggaagagcac tgccggcagc agtaccaga ccagcctggg 540
cgctttgcca agctgctgct gcggttgcca gctctgcgca gtattggcct caagtgcctc 600
gaacatctct ttttcttcaa gctcatcggg gacacgcca tcgacaactt tcttcttcc 660
atgctggagg cccctctga cccctaa 687

```

```

<210> SEQ ID NO 19
<211> LENGTH: 693
<212> TYPE: DNA
<213> ORGANISM: Amblyomma americanum

```

```

<400> SEQUENCE: 19

```

```

tctccggaca tgccactcga acgcattctc gaagccgaga tgccgctcga gcagccggca 60
ccgtccgttt tggcgcagac ggccgcactc ggccgcgacc ccgtcaacag catgtgccag 120
gctgccccgc cacttcacga gctcgtacag tgggcccggc gaattccgca cttcgaagag 180
cttcccateg aggatcgcac cgcctgctc aaagccggct ggaacgaact gcttattgce 240
gccttttgc accgttctgt ggcggtgcgc gacggcatcg ttctggccac cgggctggtg 300
gtgcagcggc acagcgcaca cggcgcaggc gttggcgaca tcttcgaccg cgtactagcc 360
gagctggtgg ccaagatgcg cgacatgaag atggacaaaa cggagctcgg ctgcctgcgc 420
gccgtggtgc tcttcaatcc agacgccaag ggtctccgaa acgccaccag agtagaggcg 480
ctccgcgaga aggtgtatgc ggcgctggag gagcactgcc gtcggcacca cccggaccaa 540
ccgggtcgct tgggcaagct gctgctgcgg ctgcctgect tgccgagcat cgggctcaaa 600
tgctcgcgac atctgttctt cttcaagctc atcggagaca ctcccataga cagcttctctg 660
ctcaacatgc tggaggcacc ggcagacccc tag 693

```

```

<210> SEQ ID NO 20
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 20

```

```

cggagtactg tcctccg 17

```

```

<210> SEQ ID NO 21
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic TATAA

```

```

<400> SEQUENCE: 21

```

```

tagagggtat ataatggatc cccgggtacc 30

```

```

<210> SEQ ID NO 22
<211> LENGTH: 1705
<212> TYPE: DNA
<213> ORGANISM: Photinus pyralis

```

```

<400> SEQUENCE: 22

```

```

atggaagacg ccaaaaaacat aaagaaaggc ccggcgccat tctatcctct agaggatgga 60
accgctggag agcaactgca taaggctatg aagagatacg ccctggttcc tggacaact 120

```

-continued

gcttttacag atgcacatat cgaggatgaac atcacgtacg cggaataactt cgaaatgtcc	180
gttcgggttg cagaagctat gaaacgatat gggctgaata caaatcacag aatcgctgta	240
tgcagtgaaa actctcttca attctttatg cgggtggttg gcgcgttatt tatcggagtt	300
gcagttgcgc ccgcgaacga ctttataat gaacgtgaat tgctcaacag tatgaacatt	360
tcgcagccta ccgtagtggt tgtttccaaa aaggggttgc aaaaaatmtt gaacgtgcaa	420
aaaaattac caataatcca gaaaattatt atcatggatt ctaaacgga ttaccaggga	480
tttcagtcga tgtacacggt cgtcacatct catctacctc ccggttttaa tgaatacgat	540
tttgaccag agtcctttga tcgtgacaaa acaattgcac tgataatgaa ttctctgga	600
tctactgggt tacctaaggg tgtggccctt ccgcatagaa ctgcctgcgt cagattctcg	660
catgccagag atcctatttt tggcaatcaa atcattccgg aactgcatg ttaagtgtt	720
gttcatttc atcacggttt tggaatggtt actacactcg gatattgat atgtggattt	780
cgagtcgtct taatgtatag atttgaagaa gagctgtttt tacgatccct tcaggattac	840
aaaaattcaa gtgcgttgct agtaccaacc ctattttcat tcttcgcaa aagcactctg	900
attgacaaat acgatttacc taatttacac gaaattgctt ctgggggcgc acctcttctg	960
aaagaagtcg gggaagcggg tgcaaaacgc ttccatcttc cagggatagc acaaggatat	1020
gggctcactg agactacatc agctattctg attacaccgg agggggatga taaaccgggc	1080
gcggtcggtg aagttgttcc attttttgaa gcgaaggttg tggatctgga tacccggaaa	1140
acgctgggcg ttaatcagag aggccaatta tgtgtcagag gacctatgat tatgtccggt	1200
tatgtaaaac atccggaagc gaccaacgcc ttgattgaca aggatggatg gctacattct	1260
ggagacatag ctactggga cgaagacgaa cacttcttca tagttgaccg cttgaagtct	1320
ttaattaaat acaaaggata tcaggtggcc cccgctgaat tggaatcgat attgttacia	1380
caccccaaac tcttcgacgc gggcgtggca ggtcttcccg acgatgacgc cggatgaactt	1440
cccgcgcccg ttgtgtttt ggagcacgga aagacgatga cggaaaaaga gatcgtggat	1500
tacgtcgcca gtcaagtaac aaccgcgaaa aagttgcgcg gaggagtgtg gtttgggac	1560
gaagtaccga aaggtcttac cggaaaactc gacgcaagaa aaatcagaga gatcctcata	1620
aaggccaaga agggcggaaa gtccaaattg taaaatgtaa ctgtattcag cgatgacgaa	1680
attcttagct attgtaatac tctag	1705

I claim:

1. An isolated polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, nucleotides 448-1109 of SEQ ID NO: 1, nucleotides 16-1109 of SEQ ID NO: 1, nucleotides 214-1109 of SEQ ID NO: 1, and nucleotides 418-1109 of SEQ ID NO: 1.

2. An isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370 of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2.

3. A vector comprising the isolated polynucleotide according to claims 1 or 2.

4. The vector according to claim 3, wherein the isolated polynucleotide is operatively linked to an expression control sequence that permits expression of the isolated polynucleotide in an expression competent host cell.

5. The vector according to claim 4, wherein the expression control sequence comprises a promoter that is functional in a mammalian cell.

6. The vector according to claim 4, wherein the vector is selected from the group consisting of a retroviral vector, plasmid, and a viral vector.

7. The vector according to claim 6, wherein the vector is a plasmid.

8. An isolated host cell transfected with the vector according to claim 3.

9. The isolated host cell according to claim 8, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a nematode cell, an insect cell, a fish cell, a plant cell, an avian cell, an animal cell, and a mammalian cell.

10. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 2, amino acids 150-370 of SEQ ID NO: 2, amino acids 6-370

77

of SEQ ID NO: 2, amino acids 72-370 of SEQ ID NO: 2, and amino acids 140-370 of SEQ ID NO: 2.

11. A method for producing a leafhopper ecdysone receptor polypeptide comprising:

a) culturing the host cell of claim 8 in culture medium under conditions permitting expression of a leafhopper ecdysone receptor polypeptide; and

b) isolating the leafhopper ecdysone receptor polypeptide from the culture medium.

12. A composition comprising the isolated polynucleotide according to claim 1 or 2 and a carrier.

13. A composition comprising the vector according to claim 3 and a carrier.

78

14. A composition comprising the isolated polypeptide according to claim 10 and a carrier.

15. A method of screening for molecules that modulate leafhopper ecdysone receptor activity in a cell, the method comprising:

a) contacting a cell comprising the isolated polypeptide according to claim 10 with a candidate molecule; and

b) detecting leafhopper ecdysone receptor activity in the presence of the molecule.

16. The method according to claim 15, wherein the molecule is an agonist of leafhopper ecdysone receptor.

17. The method according to claim 15, wherein the molecule is an antagonist of leafhopper ecdysone receptor.

* * * * *