



US 20130101619A1

(19) **United States**

(12) **Patent Application Publication**  
**Cook et al.**

(10) **Pub. No.: US 2013/0101619 A1**

(43) **Pub. Date: Apr. 25, 2013**

(54) **RECOMBINANT NON-PATHOGENIC  
MAREK'S DISEASE VIRUS CONSTRUCTS  
ENCODING INFECTIOUS  
LARYNGOTRACHEITIS VIRUS AND  
NEWCASTLE DISEASE VIRUS ANTIGENS**

**Publication Classification**

(71) Applicant: **INTERVET INC., (US)**

(72) Inventors: **Stephanie Cook, Omaha, NE (US);  
Mohamad Morsey, Omaha, NE (US);  
Gary Petersen, Omaha, NE (US);  
Paulus Jacobus Antonius  
Sondermeijer, Boxmeer (NL)**

(51) <b>Int. Cl.</b>	
<i>C12N 7/01</i>	(2006.01)
<i>A61P 31/14</i>	(2006.01)
<i>A61P 37/04</i>	(2006.01)
<i>A61P 31/22</i>	(2006.01)
<i>C12N 15/62</i>	(2006.01)
<i>A61K 39/295</i>	(2006.01)
(52) <b>U.S. Cl.</b>	
USPC .....	<b>424/199.1; 435/235.1; 536/23.4</b>

(21) Appl. No.: **13/655,858**

(57) **ABSTRACT**

(22) Filed: **Oct. 19, 2012**

**Related U.S. Application Data**

(60) Provisional application No. 61/549,844, filed on Oct. 21, 2011.

The present invention discloses novel recombinant multivalent non-pathogenic Marek's Disease virus constructs that encode and express both Infectious Laryngotracheitis Virus and Newcastle Disease virus protein antigens, and methods of their use in poultry vaccines.

FIGURE 1





FIGURE 2



**RECOMBINANT NON-PATHOGENIC  
MAREK'S DISEASE VIRUS CONSTRUCTS  
ENCODING INFECTIOUS  
LARYNGOTRACHEITIS VIRUS AND  
NEWCASTLE DISEASE VIRUS ANTIGENS**

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

**[0001]** This application claims priority under 35 U.S.C. §119(e) of provisional application U.S. Ser. No. 61/549,844 filed Oct. 21, 2011, the contents of which are hereby incorporated by reference in their entireties.

**FIELD OF THE INVENTION**

**[0002]** The present invention relates to novel recombinant multivalent non-pathogenic Marek's Disease virus constructs encoding and expressing Infectious Laryngotracheitis Virus and Newcastle Disease virus protein antigens, and methods of their use in poultry vaccines.

**BACKGROUND OF THE INVENTION**

**[0003]** Pathogenic poultry viruses are not only debilitating to chickens, but they also are costly to chicken breeders because most of the resulting diseases are contagious and the poultry industry relies heavily on confined, large-scale breeding facilities. Vaccinating young chicks is often the only viable means to combat these viruses. Although attenuated or killed poultry viral vaccines remain important in the market place, in recent years significant resources have been expended on developing vaccines containing recombinant viral constructs which express pathogenic viral protein antigens. Furthermore, substantial efforts have been made to construct stable and efficacious multivalent recombinant non-pathogenic Marek's Disease virus (rMDV<sub>np</sub>) vectors that express foreign genes from multiple viral pathogens. Such multivalent vaccines would serve to minimize the number of injections given to the chicks and thereby, reduce discomfort and stress on the vaccinated chick, as well as significantly reduce costs in labor and materials. Vaccinating with such single multivalent constructs also would be preferable to alternative multivalent rMDV<sub>np</sub> vaccines that contain multiple recombinant monovalent rMDV<sub>np</sub> constructs, because these alternative vaccines have, at least to date, resulted in protection against only a single viral pathogen. The failure of such alternative vaccines is presumably due to one of the monovalent rMDV<sub>np</sub> constructs overgrowing the other monovalent rMDV<sub>np</sub> constructs thereby, preventing these other monovalent rMDV<sub>np</sub> constructs from inducing a significant immune response. In any case, despite substantial efforts in the past to construct stable and efficacious multivalent recombinant rMDV<sub>np</sub> vectors that express foreign genes from multiple viral pathogens heretofore, such efforts have proved unsuccessful.

**[0004]** One poultry virus disease that can be controlled through vaccination is Marek's disease. Marek's disease is a pathogenic disease that adversely affects chickens, worldwide. Marek's disease occurs predominantly in young chickens between 2 and 5 months of age. Clinical signs include: progressive paralysis of one or more of the extremities, incoordination due to paralysis of legs, drooping of the limb due to wing involvement, and a lowered head position due to

involvement of the neck muscles. In acute cases, severe depression may result. Bursal and thymic atrophy may also develop.

**[0005]** The etiological agent for Marek's disease is Marek's disease virus serotype 1 (MDV1), a cell-associated virus having a double-stranded DNA genome. MDV1 is a lymphotropic avian alphaherpesvirus that both: (i) infects B cells, which can result in cytolysis, and (ii) latently infects T cells, which can induce T-cell lymphoma. Closely related to the virulent MDV1 strain, Marek's disease virus serotype 2 (MDV2), previously known as Gallid herpes virus 3, is a naturally attenuated MDV strain that has been shown to have little to no pathogenicity in chickens [Petherbridge et al., *J. Virological Methods* 158:11-17 (2009)]. SB-1 is a specific MDV2 strain that has been shown to be useful in vaccines against MDV1 [see e.g., Murthy and Calnek, *Infection and Immunity* 26(2) 547-553 (1979)].

**[0006]** Another closely related alphaherpesvirus, Marek's disease virus serotype 3 (MDV3), more widely known as herpesvirus of turkeys (HVT), is a nonpathogenic virus of domestic turkeys [see e.g., Kingham et al., *J. of General Virology* 82:1123-1135 (2001)]. Two commonly used strains of HVT are the PB1 strain and the FC126 strain. Whereas, HVT is also nonpathogenic in chickens, it does induce a long-lasting protective immune response in chickens against MDV1. Accordingly, HVT has been used in poultry vaccines against virulent MDV1 for many years, generally in combination with SB-1, which is more viraemic than HVT, but considered less safe. Alternatively, when flocks are challenged with particularly virulent MDV1 strains, HVT can be combined with the Rispen's vaccine. The Rispen's vaccine is an isolate that originated from a mildly virulent MDV1 strain that was subsequently further weakened by cell passaging. The Rispen's strain however, retains some virulence towards highly susceptible lines of chickens.

**[0007]** The sequence of the complete genome of HVT has been disclosed [Afonso et al., *J. Virology* 75(2):971-978 (2001)], and as most alphaherpesviruses, HVT possesses a significant number of potential nonessential insertion sites [see e.g., U.S. Pat. No. 5,187,087; U.S. Pat. No. 5,830,745; U.S. Pat. No. 5,834,305; U.S. Pat. No. 5,853,733; U.S. Pat. No. 5,928,648; U.S. Pat. No. 5,961,982; U.S. Pat. No. 6,121,043; U.S. Pat. No. 6,299,882 B1]. HVT also has been shown to be amenable to genetic modification and thus, has been used as a recombinant vector for many years [WO 87/04463]. Accordingly, recombinant HVT vectors have been reported to express foreign genes that encode antigens from e.g., Newcastle Disease Virus (NDV), [Sondermeijer et al., *Vaccine*, 11:349-358 (1993); Reddy et al., *Vaccine*, 14:469-477 (1996)], Infectious Bursal Disease Virus (IBDV), [Darteil et al., *Virology*, 211:481-490 (1995); Tsukamoto et al., *J. of Virology* 76(11):5637-5645 (2002)], and Infectious Laryngotracheitis Virus (ILTV) [Johnson et al., *Avian Disease*, 54(4):1251-1259 (2010); WO 92/03554; U.S. Pat. No. 6,875,856]. The entire genomic sequence of MDV2 is also known [see, GenBank acc. nr: AB049735.1, and Petherbridge et al., supra]. The genomic organization of the MDV2 is very similar to that of HVT, with the US region in particular, being identical to that of HVT [see, Kingham et al., supra].

**[0008]** In addition a recombinant chimeric virus, known as the novel avian herpesvirus (NAHV), has been constructed in which specific regions of the HVT genome have been replaced by the corresponding regions of the MDV1 genome. The NAHV also has been used to express foreign genes that

encode antigens from other poultry viruses [U.S. Pat. No. 5,965,138; U.S. Pat. No. 6,913,751].

**[0009]** Like MDV, infectious laryngotracheitis virus (ILTV) is an alphaherpesvirus that adversely affects chickens, worldwide [Fuchs et al., *Veterinary Research* 38:261-279 (2007)]. ILTV causes acute respiratory disease in chickens, which is characterized by respiratory depression, gasping, and expectoration of bloody exudate. Viral replication is limited to cells of the respiratory tract, where in the trachea the infection gives rise to tissue erosion and hemorrhage.

**[0010]** Newcastle disease is another highly contagious and debilitating disease of chickens. The etiological agent for Newcastle disease is the Newcastle disease virus (NDV). NDV belongs to the order of the Mononegavirales and is in the family of Paramyxoviridae. Newcastle disease viruses have a non-segmented, negative sense, single-stranded RNA genome. NDV has been grouped into three distinct pathotypes according to their virulence. Infection of poultry by the non-pathogenic lentogenic strains of NDV is essentially asymptomatic. In direct contrast, the mesogenic (medium pathogenic) and velogenic (highly pathogenic) NDV strains cause extensive disease that can be fatal. Most types of NDV infect the respiratory system and/or the nervous system, and can result in gasping and torticollis.

**[0011]** Infectious bursal disease virus (IBDV), also called Gumboro disease virus, is the causative agent of infectious bursal disease. IBDV causes an acute, highly-contagious, viral infection of a chicken's lymphoid tissue, with its primary target being the bird's essential immunological organ: the bursa of Fabricius. The morbidity rate in susceptible flocks is high, with rapid weight loss and moderate to high mortality rates. Chicks that recover from the disease may have immune deficiencies because of destruction of (or parts of) the bursa of Fabricius. This makes them particularly vulnerable to secondary infections.

**[0012]** IBDV is a member of the Birnaviridae family. The viruses in this family have a genome consisting of two segments (A and B) of double-stranded RNA. Two serotypes of IBDV exist, serotype 1 and 2, which can be differentiated by virus neutralization (VN) tests. Serotype 1 viruses have been shown to be pathogenic to chickens, while serotype 2 viruses cause only sub-acute disease in turkeys. Historically, IBDV serotype 1 viruses consisted of only one type that is now known as "classic" IBD virus. More recently, so-called "variant" IBDV strains have emerged. Classic and variant strains of IBDV can be identified and distinguished by a virus neutralisation test using a panel of monoclonal antibodies, or by RT-PCR [Wu et al., *Avian Diseases*, 51:515-526 (2007)]. Well-known classic IBDV strains include, D78, Faragher 52/70, and STC, whereas 89/03 is a well-known variant strain. Many live or inactivated IBDV vaccines are commercially available, e.g. a live vaccine such as NOBILIS® Gumboro D78 (MSD Animal Health).

**[0013]** As indicated above, because HVT can act as both an antigen that provides significant protection against Marek's Disease and as a recombinant vector, it is presently used as a platform vector for such multivalent vaccines as Innovax®-ILT (sold by Merck Animal Health), which protects against ILTV; and Innovax®-ND-SB (sold by Merck Animal Health) and Vectormune® HVT-NDV (sold by Ceva), both of which protect against NDV. Notably, however, heretofore, no multivalent vaccine comprising a recombinant HVT encoding antigens from more than one pathogen has been shown to be stable and efficacious, even though such vaccines had been

suggested more than fifteen years ago [see e.g., U.S. Pat. No. 5,965,138]. Indeed, Innovax®-ILT contains the only recombinant HVT that comprises two foreign genes, i.e., ILTV gD and ILTV gI, which has proved to be safe, effective, and stable. However, these two foreign genes are from the same pathogen and moreover, they naturally overlap and need to be co-expressed in order to allow proper immunization against ILTV.

**[0014]** Accordingly, despite the clear advantages of stable, multivalent, recombinant MDV<sub>np</sub> constructs that can efficaciously express foreign antigens from two or more different pathogens, and the substantial efforts to design them, heretofore, none have been forthcoming. Therefore, there is a clear need to overcome the collective industry failure, by constructing novel, stable, recombinant MDV<sub>np</sub> vectors that can be used in multivalent vaccines as the sole active to protect against two or more different non-MDV1 poultry virus pathogens.

**[0015]** The citation of any reference herein should not be construed as an admission that such reference is available as "prior art" to the instant application.

#### SUMMARY OF THE INVENTION

**[0016]** Accordingly, the present invention provides a novel, stable, and efficacious multivalent recombinant nonpathogenic Marek's Disease virus (rMDV<sub>np</sub>) for use as a vector to express foreign genes from multiple viral pathogens. In particular embodiments, the rMDV<sub>np</sub> is a recombinant herpesvirus of turkeys (rHVT). In alternative embodiments, the rMDV<sub>np</sub> is a recombinant Marek's disease virus serotype 2 (rMDV2). An rMDV<sub>np</sub>, e.g., an rHVT or an rMDV2, can be used in vaccines against pathogenic poultry viruses.

**[0017]** In particular embodiments, an rMDV<sub>np</sub> comprises a first nucleic acid inserted in a first nonessential site in the rMDV<sub>np</sub> genome and a second nucleic acid inserted in a second nonessential site in the rMDV<sub>np</sub> genome. The first nucleic acid comprises both a nucleotide sequence that encodes an Infectious Laryngotracheitis Virus (ILTV) gD protein and a nucleotide sequence that encodes an Infectious Laryngotracheitis Virus (ILTV) gI protein. The second nucleic acid comprises a nucleotide sequence that encodes a Newcastle Disease Virus (NDV) F protein. In specific embodiments of this type, the first nucleic acid comprises the nucleotide sequence of SEQ ID NO: 16 and the second nucleic acid comprises the nucleotide sequence of SEQ ID NO: 15. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0018]** In certain embodiments, the first nonessential site of the rMDV<sub>np</sub> is the US2 site, while the second nonessential site is a nonessential site of the rMDV<sub>np</sub> other than the US2 site. In related embodiments, the first nonessential site of the rMDV<sub>np</sub> is the US2 site and the second nonessential site of the rMDV<sub>np</sub> is the UL7/8 site. In yet other embodiments, the first nonessential site of the rMDV<sub>np</sub> is the US2 site and the second nonessential site of the rMDV<sub>np</sub> is the US10 site. In still other embodiments, the first nonessential site of the rMDV<sub>np</sub> is the US2 site and the second nonessential site of the rMDV<sub>np</sub> is the UL 54.5 site. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0019]** In other embodiments, the first nonessential site and the second nonessential site of the rMDV<sub>np</sub> are the same. In specific embodiments of this type, the first nucleic acid and the second nucleic acid are actually constructed as part of the same DNA molecule, which is inserted into a nonessential

site of the rMDV<sub>np</sub>. Such a DNA molecule can be an expression cassette that encodes an Infectious Laryngotracheitis Virus (ILTV) gD protein, an Infectious Laryngotracheitis Virus (ILTV) gI protein, and a Newcastle Disease Virus (NDV) F protein. In particular embodiments of this type, the DNA molecule comprises the nucleotide sequence of SEQ ID NO: 17. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0020]** Accordingly, in particular embodiments, the first nonessential site and the second nonessential site of the rMDV<sub>np</sub> are the US2 site. In other embodiments, the first nonessential site and the second nonessential site of the rMDV<sub>np</sub> are the UL54.5 site. In yet other embodiments, the first nonessential site and the second nonessential site of the rMDV<sub>np</sub> are the UL7/8 site. In still other embodiments, the first nonessential site and the second nonessential site of the rMDV<sub>np</sub> are the US10 site. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0021]** The nucleotide sequences encoding the ILTV gD protein, the ILTV gI protein, and the NDV F protein can be operatively under the control of exogenous promoters, i.e., promoters that are not naturally found in the MDV<sub>np</sub>. In certain embodiments, these three nucleotide sequences are operatively under the control of different promoters, i.e., the nucleotide sequence encoding the ILTV gD protein is operatively under the control of a first promoter, the nucleotide sequence encoding the ILTV gI protein is operatively under the control of a second promoter, and the nucleotide sequence encoding the NDV F protein is operatively under the control of a third promoter, with the first promoter, the second promoter, and the third promoter all being different. In particular embodiments, the promoter for the nucleotide sequence encoding the ILTV gD protein is the endogenous ILTV gD promoter. In certain embodiments, the promoter for the nucleotide sequence encoding the ILTV gI protein is the endogenous ILTV gI promoter. In particular embodiments of this type, the promoter for the nucleotide sequence encoding the ILTV gD protein is the endogenous ILTV gD promoter and the promoter for the nucleotide sequence encoding the ILTV gI protein is the endogenous ILTV gI promoter. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0022]** In certain embodiments, at least one of the promoters operably linked to a nucleotide sequence encoding the ILTV gD protein, the ILTV gI protein, or the NDV F protein is the human cytomegalovirus immediate early (hCMV IE) promoter. In particular embodiments of this type, the promoter for the nucleotide sequence encoding the NDV F protein is the hCMV IE promoter. In specific embodiments, at least one of the promoters operably linked to a nucleotide sequence encoding the ILTV gD protein, the ILTV gI protein or the NDV F protein is the pseudorabies virus (PRV) gpX promoter. In related embodiments, at least one of the promoters operably linked to a nucleotide sequence encoding the ILTV gD protein, the ILTV gI protein or the NDV F protein is the chicken beta-actin gene promoter. In specific embodiments, the promoter for the nucleotide sequence encoding the NDV F protein is the hCMV IE promoter, the promoter for the nucleotide sequence encoding the ILTV gD protein is the endogenous ILTV gD promoter, and the promoter for the nucleotide sequence encoding the ILTV gI protein is the endogenous ILTV gI promoter.

**[0023]** In certain embodiments, an rMDV<sub>np</sub> of the present invention that includes insertions of nucleotide sequences

encoding the ILTV gD protein, the ILTV gI protein, and the NDV F protein also includes one or more exogenous transcription terminator sequences. In specific embodiments of this type, a transcription terminator sequence is downstream from the nucleotide sequence encoding the NDV F protein. In particular embodiments, the nucleotide sequences encoding the ILTV gD protein and the ILTV gI protein share one transcription terminator sequence and the nucleotide sequence encoding the NDV F protein has another. In particular embodiments, at least one of the transcription terminator sequences comprises a synthetic polyadenylation sequence. In related embodiments at least one of the transcription terminator sequences comprises a Herpes Simplex Virus thymidine kinase (HSV TK) polyadenylation sequence. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0024]** The present invention also provides a recombinant nucleic acid comprising in 5' to 3' direction in the following order (i) an Infectious Laryngotracheitis Virus (ILTV) gD promoter, (ii) a coding sequence for the ILTV gD protein, (iii) an ILTV gI promoter, (iv) a coding sequence for the ILTV gI protein, (v) a human cytomegalovirus immediate early (hCMV IE) promoter, (vi) a coding sequence for the NDV F protein, and (viii) a transcription terminator sequence. In a particular embodiment of this type, the recombinant nucleic acid comprises the nucleotide sequence of SEQ ID NO: 17.

**[0025]** The present invention further provides an rMDV<sub>np</sub> in which a recombinant nucleic acid of the present invention has been inserted into a nonessential insertion site of the rMDV<sub>np</sub>. In certain embodiments of this type, the rMDV<sub>np</sub> includes an insert in a nonessential site that comprises a recombinant nucleic acid comprising in 5' to 3' direction in the following order (i) an Infectious Laryngotracheitis Virus (ILTV) gD promoter, (ii) a coding sequence for the ILTV gD protein, (iii) an ILTV gI promoter, (iv) a coding sequence for the ILTV gI protein, (v) a human cytomegalovirus immediate early (hCMV IE) promoter, (vi) a coding sequence for the NDV F protein, and (vii) a transcription terminator sequence. In specific embodiments, intervening nucleotide sequences, such as linkers, spacer sequences, and/or extraneous coding sequences, can also be included, see Example 1 below. In a particular embodiment, the rHVT comprises the nucleotide sequence of SEQ ID NO: 17 inserted into a nonessential site. In particular embodiments of these types, the nonessential site is the US2 site. In other such embodiments, the nonessential site is the UL54.5 site. In still other such embodiments, the nonessential site is the UL7/8 site. In yet other such embodiments, the nonessential site is the US10 site. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0026]** The present invention also provides methods of making an rMDV<sub>np</sub> of the present invention. In certain embodiments, a heterologous nucleic acid is constructed that comprises a nucleotide sequence that encodes an ILTV gD protein, a nucleotide sequence that encodes an ILTV gI protein, and a nucleotide sequence that encodes an NDV F protein. The heterologous nucleic acid is then inserted into a nonessential site of an rMDV<sub>np</sub> of the present invention. In certain embodiments, the heterologous nucleic acid is an expression cassette. In particular embodiments of this type, the expression cassette comprises the nucleotide sequence of SEQ ID NO: 17. In other embodiments, a first heterologous nucleic acid is constructed that comprises a nucleotide sequence that encodes an ILTV gD protein and a nucleotide

sequence that encodes an ILTV gI protein; and a second heterologous nucleic acid is constructed that comprises a nucleotide sequence that encodes an NDV F protein. The first heterologous nucleic acid is inserted into a US2 site of an rMDV<sub>np</sub> and the second heterologous nucleic acid is inserted into an alternative nonessential site of the rMDV<sub>np</sub>. In certain embodiments, such heterologous nucleic acids are expression cassettes. In particular embodiments of this type, the first heterologous nucleic acid comprises the nucleotide sequence of SEQ ID NO: 16, and the second heterologous nucleic acid comprises the nucleotide sequence of SEQ ID NO: 15. In specific embodiments, the method of making an rMDV<sub>np</sub> is a method of making an rHVT. In alternative embodiments, the method of making an rMDV<sub>np</sub> is a method of making an rMDV2.

**[0027]** The present invention further provides immunogenic compositions and/or vaccines that comprise any rMDV<sub>np</sub> of the present invention. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2. In addition, the present invention provides methods for aiding in the protection of poultry against a disease caused by ILTV and/or NDV and/or MDV1 by administering such a vaccine and/or immunogenic composition of the present invention. In specific embodiments, such methods aid in the protection of a chicken. In particular embodiments of this type, a vaccine of the present invention is administered subcutaneously. In other embodiments, a vaccine of the present invention is administered in ovo.

**[0028]** Accordingly in one aspect, the present invention provides stable, safe, and efficacious immunogenic compositions and/or vaccines that comprise an rMDV<sub>np</sub> of the present invention. The present invention also provides immunogenic compositions and/or vaccines that comprise any rMDV<sub>np</sub> of the present invention that is further combined with an additional NDV, ILTV, and/or MDV antigen to improve and expand the immunogenicity provided. In addition, the present invention also provides immunogenic compositions and/or vaccines that comprise any rMDV<sub>np</sub> of the present invention that is further combined with an antigen for a pathogen other than MDV, ILTV, or NDV. In a particular embodiment of this type, the antigen is an Infectious Bursal Disease Virus (IBDV) antigen. In a more particular embodiment the IBDV antigen is a mild live IBDV. In certain embodiments the mild live IBDV is a variant IBDV. The present invention also provides methods for aiding in the protection of poultry against a disease caused by ILTV and/or NDV and/or MDV1 and/or IBDV by administering such a vaccine and/or immunogenic composition to the poultry (e.g., chicken). In particular embodiments of this type, a vaccine of the present invention is administered subcutaneously. In other embodiments, a vaccine of the present invention is administered in ovo.

**[0029]** In certain embodiments the immunogenic compositions and/or vaccines of the present invention comprise an rHVT that comprises as an insertion into its US2 site of a recombinant nucleic acid comprising 5' to 3': (i) an Infectious Laryngotracheitis Virus (ILTV) gD promoter; (ii) a coding sequence for the ILTV gD protein; (iii) an ILTV gI promoter; (iv) a coding sequence for the ILTV gI protein; (v) a human cytomegalovirus immediate early (hCMV IE) promoter; (vi) a coding sequence for the Newcastle Disease Virus fusion protein (NDV F); and (vii) a transcription terminator sequence. In particular embodiments of this type the immunogenic compositions and/or vaccines further comprise a mild live infectious bursal disease virus (IBDV). In certain

embodiments the mild live IBDV is a variant IBDV. In more particular embodiments, the IBDV is 89/03. In even more particular embodiments of this type, the recombinant nucleic acid has the nucleotide sequence of SEQ ID NO: 17.

**[0030]** The present invention further provides immunogenic compositions and/or vaccines that comprise any rMDV<sub>np</sub> of the present invention combined with an additional NDV, ILTV, and/or MDV antigen, and a pathogen other than MDV, ILTV, or NDV.

**[0031]** These and other aspects of the present invention will be better appreciated by reference to the following Figures and the Detailed Description.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0032]** FIG. 1 is a schematic drawing of the HVT (FC126) genome, consisting of a unique long (UL) region, and a unique short (US) region, each denoted by straight lines, and flanked by repeat regions, denoted as boxes. Below the genome schematic, is a bar indicating the location of BamHI restriction enzyme digestion fragments, relative to their genome position, and the lettering nomenclature associated with each fragment. (The largest fragment was given the letter "A", the next largest given the letter "B", and so forth and so on). The positions of each cloned subgenomic fragment (and their designation) used to reconstruct either HVT (FC126) or the rHVT/NDV/ILT viruses are indicated below the BamHI restriction map. The asterisk (\*) indicates the position of the insertion sites: UL7/UL8 in 1196-05.1; UL54.5 in 1332-29.4; US2 in 1332-47.A2 or 1317-15.1-1.

**[0033]** FIG. 2 is a schematic drawing of six different recombinant HVTs, which depict the genes inserted into the HVT backbone and the site of their insertion. Innovax-LT is an rHVT that includes an expression cassette encoding the ILTV gD and ILTV gI genes inserted in the UL54.5 site of the rHVT. Innovax-ND is an rHVT that includes an expression cassette encoding the NDV fusion gene inserted in the US10 site of the rHVT. 1348-34C is an rHVT that includes both an expression cassette encoding the ILTV gD and ILTV gI genes inserted in the UL54.5 site of the rHVT, and an expression cassette encoding the NDV fusion gene inserted in the US10 site of the rHVT. 1332-62E is an rHVT that includes an expression cassette that encodes the ILTV gD, the ILTV gI, and the NDV fusion genes inserted in the US2 site of the rHVT. 1317-46 is an rHVT that includes both an expression cassette encoding the ILTV gD and ILTV gI genes inserted in the US2 site, and an expression cassette encoding the NDV fusion gene inserted between UL7 and UL8 (i.e., the UL7/8 site) of the rHVT. 1332-70B is an rHVT that includes an expression cassette that encodes the ILTV gD, the ILTV gI, and the NDV fusion genes inserted in the UL54.5 site of the rHVT.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0034]** The present invention overcomes the prior industry failure to be able to construct rMDV<sub>np</sub> vectors that both contain foreign antigens and can protect against two or more different poultry virus pathogens by providing unique recombinant MDV<sub>np</sub> vectors that encode and express antigens from ILTV and NDV, and that protect against Marek's disease, Newcastle disease, and Infectious Laryngotracheitis virus. In particular embodiments, an rMDV<sub>np</sub> of the present invention encodes and expresses foreign antigens from only ILTV and NDV, and can aid in the protection against Marek's disease,

Newcastle disease, and Infectious Laryngotracheitis virus. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0035]** Prior to the present invention, an HVT vector already had been constructed containing an NDV gene inserted into the US10 region. This HVT-NDV vector was shown to be stable and to express sufficient levels of the corresponding NDV gene product, the NDV F protein, to protect vaccinated chickens against a virulent NDV challenge. In addition, an HVT vector already had been constructed containing a pair of ILTV genes inserted in the HVT UL54.5 region. This HVT-ILTV vector was shown to be stable and to express sufficient levels of the corresponding ILTV gene products, the ILTV gI and gD proteins, to protect vaccinated chickens against a virulent ILTV challenge virus.

**[0036]** Accordingly, a multivalent HVT construct to protect against both NDV and ILTV was designed based on the successful constructs above, i.e., inserting the NDV-F gene in the US10 site and inserting the ILTV gD and gI genes in UL54.5 site [see, 1348-34C in FIG. 2]. Unexpectedly however, following the passaging of this construct in tissue culture the recombinant virus lost its ability to express the ILTVgD, ILTVgI, and NDV F proteins. This proved to be true with a number of duplicate recombinant rHVT constructs. Indeed, these recombinant viruses were unstable and unsuitable for further development as vaccines. These findings demonstrate that the design of a single multivalent rHVT vector that can stably express both the NDV F protein and the ILTVgD and ILTVgI proteins is not a simple process that can be extrapolated from existing information. Indeed, if such stable and efficacious multivalent rHVT vectors were possible at all, their design needed to be premised on an unpredictable set of complex interactions minimally involving the relationship between the insertion sites used and the foreign genes to be inserted. Heretofore, such design of rHVT constructs was not readily predictable from the known art.

**[0037]** The present invention therefore, provides recombinant rMDV<sub>np</sub> vectors in which two genes from ILTV and one gene from NDV have been inserted. In a particular embodiment of the present invention all three genes were inserted in the US2 region of the HVT genome. Upon vaccination of a chicken or a chicken egg with this rHVT, the cells of the immunized host expressed the proteins encoded by the inserted genes. Furthermore, the NDV and ILTV proteins expressed by the rHVT stimulated an immune response that protected the vaccinated chicken against the disease caused by NDV and ILTV. Accordingly, such rMDV<sub>np</sub> vectors can be used to provide protection against both NDV and ILTV infections. Previously, two separate rHVT vectors were necessary to protect against these two viruses, namely one for protection against ILTV and the other for protection against NDV.

**[0038]** The present invention therefore, is advantageous over current methods because it provides simultaneous protection against ILTV and NDV by inoculation of poultry and/or poultry eggs with only a single recombinant MDV<sub>np</sub>. In particular, this allows for additional vaccines to be administered via the in ovo route, because there is a limit on how much volume can be injected into an egg, and further saves on manufacturing costs because only one rather than two vectors is needed. Moreover, this can allow an additional antigen to be included in the vaccine such as a live IBDV, e.g., strain 89/03.

**[0039]** Moreover, the present invention further includes embodiments that comprise different rMDV<sub>np</sub> constructs in

the same vaccine and/or immunogenic compositions. In certain embodiments of this type, the vaccine and/or immunogenic composition comprise both an rMDV2 and an rHVT, each of which encode one or more foreign antigens. Indeed, unlike the combination of two rHVTs, which inevitably lead to one construct significantly overgrowing the other, combining an rHVT with an rMDV2 leads to no such significant overgrowth. Therefore, in specific embodiments, a vaccine of the present invention comprises an rHVT that encodes an ILTVgD protein, an ILTVgI protein, and an NDV F protein with an rMDV2 that encodes yet another poultry viral antigen.

**[0040]** In order to more fully appreciate the instant invention, the following definitions are provided.

**[0041]** The use of singular terms for convenience in description is in no way intended to be so limiting. Thus, for example, reference to a composition comprising “a polypeptide” includes reference to one or more of such polypeptides.

**[0042]** As used herein a “nonpathogenic Marek’s Disease Virus” or “MDV<sub>np</sub>” or “npMDV” is a virus in the MDV family that shows little to no pathogenicity in poultry. The term “MDV<sub>np</sub>” includes naturally occurring MDVs that have been passaged or otherwise similarly manipulated, but does not include viral constructs in which a specific region of the genome of one MDV serotype is replaced by the corresponding region of a different MDV serotype to form a chimeric virus, such as the novel avian herpesvirus (NAHV). In certain embodiments, the MDV<sub>np</sub> is an HVT. In other embodiments, the MDV<sub>np</sub> is an MDV2. In particular embodiments of this type, the MDV2 is SB1.

**[0043]** As used herein, an MDV<sub>np</sub> that has been genetically modified to encode a heterologous nucleotide sequence (e.g., a foreign gene) is defined as a “recombinant MDV<sub>np</sub>” or “rMDV<sub>np</sub>”.

**[0044]** As used herein, a “nonessential site” is a site in the MDV<sub>np</sub> genome in which an insertion of a heterologous nucleotide sequence into that site does not prevent the MDV<sub>np</sub> from replicating in a host cell. Nonessential sites are generally identified by the gene in which they reside, e.g., the US2 site, or a region between two genes, e.g., the UL7/8 site.

**[0045]** As used herein the term “poultry” can include chickens, turkeys, ducks, geese, quail, and pheasants.

**[0046]** As used herein, a “vaccine” is a composition that is suitable for application to an animal (including, in certain embodiments, humans, while in other embodiments being specifically not for humans) comprising one or more antigens typically combined with a pharmaceutically acceptable carrier such as a liquid containing water, which upon administration to the animal induces an immune response strong enough to minimally aid in the protection from a clinical disease arising from an infection with a wild-type microorganism, i.e., strong enough for aiding in the prevention of the clinical disease, and/or preventing, ameliorating or curing the clinical disease.

**[0047]** As used herein, a “multivalent vaccine” is a vaccine that comprises two or more different antigens. In a particular embodiment of this type, the multivalent vaccine stimulates the immune system of the recipient against two or more different pathogens.

**[0048]** As used herein, the term “aids in the protection” does not require complete protection from any indication of infection. For example, “aids in the protection” can mean that the protection is sufficient such that, after challenge, symptoms of the underlying infection are at least reduced, and/or



that one or more of the underlying cellular, physiological, or biochemical causes or mechanisms causing the symptoms are reduced and/or eliminated. It is understood that “reduced,” as used in this context, means relative to the state of the infection, including the molecular state of the infection, not just the physiological state of the infection.

**[0049]** As used herein, an “adjuvant” is a substance that is able to favor or amplify the cascade of immunological events, ultimately leading to a better immunological response, i.e., the integrated bodily response to an antigen. An adjuvant is in general not required for the immunological response to occur, but favors or amplifies this response.

**[0050]** As used herein, the term “pharmaceutically acceptable” is used adjectivally to mean that the modified noun is appropriate for use in a pharmaceutical product. When it is used, for example, to describe an excipient in a pharmaceutical vaccine, it characterizes the excipient as being compatible with the other ingredients of the composition and not disadvantageously deleterious to the intended recipient.

**[0051]** As used herein, “systemic administration” is administration into the circulatory system of the body (comprising the cardiovascular and lymphatic system), thus affecting the body as a whole rather than a specific locus such as the gastro-intestinal tract (via e.g., oral or rectal administration) and the respiratory system (via e.g., intranasal administration). Systemic administration can be performed e.g., by administering into muscle tissue (intramuscular), into the dermis (intra-dermal or transdermal), underneath the skin (subcutaneous), underneath the mucosa (submucosal), in the veins (intravenous) etc.

**[0052]** As used herein the term “parenteral administration” includes subcutaneous injections, submucosal injections, intravenous injections, intramuscular injections, intradermal injections, and infusion.

**[0053]** The term “approximately” is used interchangeably with the term “about” and signifies that a value is within twenty-five percent of the indicated value i.e., a peptide containing “approximately” 100 amino acid residues can contain between 75 and 125 amino acid residues.

**[0054]** As used herein, the term, “polypeptide” is used interchangeably with the terms “protein” and “peptide” and denotes a polymer comprising two or more amino acids connected by peptide bonds. The term “polypeptide” as used herein includes a significant fragment or segment, and encompasses a stretch of amino acid residues of at least about 8 amino acids, generally at least about 12 amino acids, typically at least about 16 amino acids, preferably at least about 20 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids, e.g., 35, 40, 45, 50, etc. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at residues 1, 2, 3, etc., and ending at, e.g., 155, 154, 153, etc., in all practical combinations.

**[0055]** Optionally, a polypeptide may lack certain amino acid residues that are encoded by a gene or by an mRNA. For example, a gene or mRNA molecule may encode a sequence of amino acid residues on the N-terminus of a polypeptide (i.e., a signal sequence) that is cleaved from, and therefore, may not be part of the final protein.

**[0056]** As used herein the term “antigenic fragment” in regard to a particular protein (e.g., a protein antigen) is a fragment of that protein (including large fragments that are missing as little as a single amino acid from the full-length protein) that is antigenic, i.e., capable of specifically interact-

ing with an antigen recognition molecule of the immune system, such as an immunoglobulin (antibody) or T cell antigen receptor. For example, an antigenic fragment of an NDV fusion protein, is a fragment of that fusion protein that is antigenic. Preferably, an antigenic fragment of the present invention is immunodominant for antibody and/or T cell receptor recognition.

**[0057]** As used herein an amino acid sequence is 100% “homologous” to a second amino acid sequence if the two amino acid sequences are identical, and/or differ only by neutral or conservative substitutions as defined below. Accordingly, an amino acid sequence is about 80% “homologous” to a second amino acid sequence if about 80% of the two amino acid sequences are identical, and/or differ only by neutral or conservative substitutions.

**[0058]** Functionally equivalent amino acid residues often can be substituted for residues within the sequence resulting in a conservative amino acid substitution. Such alterations define the term “a conservative substitution” as used herein. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutions for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

**[0059]** Particularly preferred conservative substitutions are: Lys for Arg and vice versa such that a positive charge may be maintained; Glu for Asp and vice versa such that a negative charge may be maintained; Ser for Thr such that a free—OH can be maintained; and Gln for Asn such that a free NH<sub>2</sub> can be maintained. The amino acids also can be placed in the following similarity groups: (1) proline, alanine, glycine, serine, and threonine; (2) glutamine, asparagine, glutamic acid, and aspartic acid; (3) histidine, lysine, and arginine; (4) cysteine; (5) valine, leucine, isoleucine, methionine; and (6) phenylalanine, tyrosine, and tryptophan.

**[0060]** In a related embodiment, two highly homologous DNA sequences can be identified by their own homology, or the homology of the amino acids they encode. Such comparison of the sequences can be performed using standard software available in sequence data banks. In a particular embodiment two highly homologous DNA sequences encode amino acid sequences having about 80% identity, more preferably about 90% identity and even more preferably about 95% identity. More particularly, two highly homologous amino acid sequences have about 80% identity, even more preferably about 90% identity and even more preferably about 95% identity.

**[0061]** As used herein, protein and DNA sequence percent identity can be determined using software such as MacVector v9, commercially available from Accelrys (Burlington, Mass.) and the Clustal W algorithm with the alignment default parameters, and default parameters for identity. See,

e.g., Thompson, et al., 1994. *Nucleic Acids Res.* 22:4673-4680. ClustalW is freely downloadable for Dos, Macintosh and Unix platforms from, e.g., EMBL, the European Bioinformatics Institute. The present download link is found at <http://www.ebi.ac.uk/clustalw/>. These and other available programs can also be used to determine sequence similarity using the same or analogous default parameters.

**[0062]** As used herein the terms “polynucleotide”, or a “nucleic acid” or a “nucleic acid molecule” are used interchangeably and denote a molecule comprising nucleotides including, but is not limited to, RNA, cDNA, genomic DNA and even synthetic DNA sequences. The terms are also contemplated to encompass nucleic acid molecules that include any of the art-known base analogs of DNA and RNA.

**[0063]** A nucleic acid “coding sequence” or a “sequence encoding” a particular protein or peptide, is a nucleotide sequence which is transcribed and translated into a polypeptide in vitro or in vivo when placed under the control of appropriate regulatory elements.

**[0064]** The boundaries of the coding sequence are determined by a start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., avian) DNA, and even synthetic DNA sequences. A transcription termination sequence can be located 3' to the coding sequence.

**[0065]** “Operably linked” refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, control elements operably linked to a coding sequence are capable of effecting the expression of the coding sequence. The control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter and the coding sequence and the promoter can still be considered “operably linked” to the coding sequence.

**[0066]** As used herein, the term “transcription terminator sequence” is used interchangeably with the term “polyadenylation regulatory element” and is a sequence that is generally downstream from a DNA coding region and that may be required for the complete termination of the transcription of that DNA coding sequence.

**[0067]** As used herein an “expression cassette” is a recombinant nucleic acid that minimally comprises a promoter and a heterologous coding sequence operably linked to that promoter. In many such embodiments, the expression cassette further comprises a transcription terminator sequence. Accordingly, the insertion of an expression cassette into a nonessential site of the rMDV<sub>np</sub> genome can lead to the expression of the heterologous coding sequence by the rMDV<sub>np</sub>. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0068]** A “heterologous nucleotide sequence” as used herein is a nucleotide sequence that is added to a nucleotide sequence of the present invention by recombinant methods to form a nucleic acid that is not naturally formed in nature. In specific embodiments, a “heterologous nucleotide sequence” of the present invention can encode a protein antigen such as the NDV F protein, the ILTV gI protein, or the ILTV gD protein. Heterologous nucleotide sequences can also encode fusion (e.g., chimeric) proteins. In addition, a heterologous nucleotide sequence can encode peptides and/or proteins that

contain regulatory and/or structural properties. In other such embodiments, a heterologous nucleotide sequence can encode a protein or peptide that functions as a means of detecting the protein or peptide encoded by the nucleotide sequence of the present invention after the recombinant nucleic acid is expressed. In still another embodiment, the heterologous nucleotide sequence can function as a means of detecting a nucleotide sequence of the present invention. A heterologous nucleotide sequence can comprise non-coding sequences including restriction sites, regulatory sites, promoters and the like.

**[0069]** Insertion of a nucleic acid encoding an antigen of the present invention into a rMDV<sub>np</sub> vector is easily accomplished when the termini of both the nucleic acid and the vector comprise compatible restriction sites. If this cannot be done, it may be necessary to modify the termini of the nucleotide sequence and/or vector by digesting back single-stranded nucleic acid overhangs (e.g., DNA overhangs) generated by restriction endonuclease cleavage to produce blunt ends, or to achieve the same result by filling in the single-stranded termini with an appropriate polymerase. Alternatively, desired sites may be produced, e.g., by ligating nucleotide sequences (linkers) onto the termini. Such linkers may comprise specific oligonucleotide sequences that define desired restriction sites. Restriction sites can also be generated through the use of the polymerase chain reaction (PCR). [See, e.g., Saiki et al., *Science* 239:487-491 (1988)]. The cleaved vector and the DNA fragments may also be modified, if required, by homopolymeric tailing.

#### Protein Antigens and Nucleic Acids Encoding the Protein Antigens

**[0070]** The ILTV gD gene appears to encode a glycoprotein of 434 amino acids in length having a molecular weight of 48,477 daltons, although others have suggested that a downstream start codon, which leads to an ILTV gD protein comprising only 377 amino acid residues, is the actual start codon [Wild et al., *Virus Genes* 12:104-116 (1996)]. The ILTV gI gene encodes a glycoprotein of 362 amino acids in length having a molecular weight of 39,753 daltons [U.S. Pat. No. 6,875,856, hereby incorporated by reference]. Nucleic acids encoding natural and/or laboratory derived variants of the ILTV gD and ILTV gI may be substituted for those presently exemplified.

**[0071]** In particular embodiments of the present invention, an rMDV<sub>np</sub> comprises a recombinant nucleic acid (e.g., an expression cassette) that encodes an ILTV gD protein comprising the amino acid sequence of SEQ ID NO: 2 or an antigenic fragment thereof. In related embodiments the rMDV<sub>np</sub> comprises a recombinant nucleic acid that encodes an ILTV gD protein comprising an amino acid sequence that has greater than 90%, and/or greater than 95%, and/or greater than 98%, and/or greater than 99% identity to the amino acid sequence of SEQ ID NO: 2. In particular embodiments, the ILTV gD protein is encoded by the nucleotide sequence of SEQ ID NO: 1. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0072]** In certain embodiments of the present invention, an rMDV<sub>np</sub> comprises a recombinant nucleic acid (e.g., an expression cassette) that encodes an ILTV gI protein comprising the amino acid sequence of SEQ ID NO: 4 or an antigenic fragment thereof. In related embodiments, the rMDV<sub>np</sub> comprises a recombinant nucleic acid that encodes an ILTV gI protein comprising an amino acid sequence that

has greater than 90%, and/or greater than 95%, and/or greater than 98%, and/or greater than 99% identity to the amino acid sequence of SEQ ID NO: 4. In particular embodiments, the ILTV gI protein is encoded by the nucleotide sequence of SEQ ID NO: 3. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

**[0073]** The NDV F protein gene encodes the so-called "fusion" protein. One NDV F protein gene exemplified by the present invention was derived from NDV Clone 30, a common lentogenic NDV vaccine strain. Nucleic acids encoding natural and/or laboratory derived variants of the F protein gene would equally be applicable, either from lentogenic, mesogenic or velogenic NDV, as the F protein gene sequence itself is highly conserved in these different NDV pathotypes. In particular embodiments of the present invention, an rMDV<sub>np</sub> comprises a recombinant nucleic acid (e.g., an expression cassette) that encodes an NDV fusion protein comprising the amino acid sequence of SEQ ID NO: 6 or an antigenic fragment thereof. In related embodiments, the rMDV<sub>np</sub> comprises a recombinant nucleic acid that encodes an NDF F protein comprising an amino acid sequence that has greater than 90%, and/or greater than 95%, and/or greater than 98%, and/or greater than 99% identity to the amino acid sequence of SEQ ID NO: 6. In specific embodiments, the NDV fusion protein is encoded by the nucleotide sequence of SEQ ID NO: 5. In certain embodiments of the present invention, an rMDV<sub>np</sub> comprises a recombinant nucleic acid (e.g., an expression cassette) that encodes an NDV fusion protein comprising the amino acid sequence of SEQ ID NO: 8 or an antigenic fragment thereof. In related embodiments, an rMDV<sub>np</sub> comprises a recombinant nucleic acid that encodes an NDF F protein comprising an amino acid sequence that has greater than 90%, and/or greater than 95%, and/or greater than 98%, and/or greater than 99% identity to the amino acid sequence of SEQ ID NO: 8. In particular embodiments, the NDV fusion protein is encoded by the nucleotide sequence of SEQ ID NO: 7. In specific embodiments, the rMDV<sub>np</sub> is an rHVT. In alternative embodiments, the rMDV<sub>np</sub> is an rMDV2.

#### Promoters and Polyadenylation Regulatory Elements

**[0074]** Many alternative promoters can be used to drive the expression of a heterologous gene encoding a protein antigen or antigenic fragment thereof in an rMDV<sub>np</sub> of the present invention. Examples include the pseudorabies virus (PRV) gpX promoter [see, WO 87/04463], the Rous sarcoma virus LTR promoter, the SV40 early gene promoter, the ILTV gD promoter, the ILTV gI promoter [see e.g., U.S. Pat. No. 6,183,753 B1], the human cytomegalovirus immediate early1 (hCMV IE1) gene promoter [U.S. Pat. No. 5,830,745; U.S. Pat. No. 5,980,906], and the chicken beta-actin gene promoter [EP 1 298 139 B1]. More specific examples, as exemplified herein, include the Towne Strain hCMV IE promoter comprising the nucleotide sequence of SEQ ID NO: 12, a truncated hCMV IE promoter comprising the nucleotide sequence of SEQ ID NO: 11, an ILTV gD promoter comprising the nucleotide sequence of SEQ ID NO: 9, and an ILTV gI promoter comprising the nucleotide sequence of SEQ ID NO: 10.

**[0075]** The inclusion of a polyadenylation regulatory element downstream from a DNA coding region is oftentimes required to terminate the transcription of the coding DNA sequence. Accordingly, many genes comprise a polyadenylation regulatory element at the downstream end of their coding sequence. Many such regulatory elements have been identi-

fied and can be used in an rMDV<sub>np</sub> of the present invention. Specific examples of polyadenylation regulatory elements as exemplified herein, include a synthetic polyadenylation signal comprising the nucleotide sequence of SEQ ID NO: 13, and the HSV thymidine kinase polyadenylation signal comprising the nucleotide sequence of SEQ ID NO: 14.

#### Vaccines and Immunogenic Compositions

**[0076]** The present invention relates to the use of the recombinant MDV<sub>np</sub>, the nucleic acid molecules used to construct the MDV<sub>np</sub>, or the host cells to grow them, or any combination thereof, all according to the present invention for the manufacture of a vaccine for poultry. Accordingly, the present invention provides vaccines and/or immunogenic compositions that include a multivalent recombinant MDV<sub>np</sub> of the present invention. Such vaccines can be used to aid in the prevention and/or prevent Newcastle disease, and/or Marek's disease, and/or maladies associated with ILTV infections. A vaccine according to the present invention can be used for prophylactic and/or for therapeutic treatment, and thus can interfere with the establishment and/or with the progression of an infection and/or its clinical symptoms of disease.

**[0077]** A recombinant MDV<sub>np</sub> of the present invention can be grown by any number of means currently practiced in the field. For example, a recombinant MDV<sub>np</sub> of the present invention can be grown through the use of in vitro cultures of primary chicken cells, see e.g., the Examples below where chicken embryo fibroblast cells (CEFs) were used. The CEFs can be prepared by trypsinization of chicken embryos. The CEFs also can be plated in monolayers and then infected with the MDV<sub>np</sub>. This particular process can be readily scaled up to industrial-sized production.

**[0078]** Therefore, a further aspect of the invention relates to a method for the preparation of the vaccine according to the invention comprising the steps of infecting host cells with a recombinant MDV<sub>np</sub> of the present invention, harvesting the infected host cells, and then admixing the harvested infected host cells with a pharmaceutically acceptable carrier. Suitable methods for infection, culture and harvesting are well known in the art and are described and exemplified herein.

**[0079]** Typically, the infected host cells are harvested while still intact to obtain the recombinant MDV<sub>np</sub> in its cell-associated form. These cells can be taken up in an appropriate carrier composition to provide stabilization for storage and freezing. The infected cells can be filled into glass ampoules, which are sealed, frozen and stored in liquid nitrogen. Accordingly, in certain embodiments of the present invention, the vaccines and/or immunogenic compositions of the present invention are stored frozen and accordingly, comprise a cryopreservative, such as dimethyl sulfoxide (DMSO), to preserve the frozen infected cells.

**[0080]** Alternatively, when the recombinant MDV<sub>np</sub> is a recombinant HVT, it can be isolated from its host cell, for instance through sonication at the end of culturing, and then taken up into a stabilizer, and freeze-dried (lyophilized) for stable storage or otherwise reduced in liquid volume, for storage, and then reconstituted in a liquid diluent before or at the time of administration. Such reconstitution may be achieved using, for example, vaccine-grade water. In certain embodiments, a lyophilized portion of a multivalent vaccine can comprise one or more antigens and the diluent can comprise one or more other antigens.

**[0081]** In particular embodiments a vaccine of the present invention (or a portion thereof) can be in a freeze-dried form, e.g., as tablets and/or spheres that are produced by a method described in WO 2010/125084, hereby incorporated by reference in its entirety. In particular, reference is made to the examples, from page 15, line 28 to page 27, line 9 of WO 2010/125084, describing a method to produce such fast disintegrating tablets/spheres. Such freeze-dried forms can be readily dissolved in a diluent, to enable systemic administration of the vaccine.

**[0082]** Vaccines and immunogenic compositions can, but do not necessarily include, physiologically compatible buffers and saline and the like, as well as pharmaceutically acceptable adjuvants. Adjuvants can be useful for improving the immune response and/or increasing the stability of vaccine preparations. Adjuvants are typically described as non-specific stimulators of the immune system, but also can be useful for targeting specific arms of the immune system. One or more compounds which have this activity may be added to the vaccine. Therefore, particular vaccines of the present invention can further comprise an adjuvant. Examples of chemical compounds that can be used as adjuvants include, but are not limited to aluminum compounds (e.g., aluminum hydroxide), metabolizable and non-metabolizable oils, mineral oils including mannide oleate derivatives in mineral oil solution (e.g., MONTANIDE ISA 70 from Seppic SA, France), and light mineral oils such as DRAKEOL 6VR, block polymers, ISCOM's (immune stimulating complexes), vitamins and minerals (including but not limited to: vitamin E, vitamin A, selenium, and vitamin B12) and CARBOPOL®.

**[0083]** Other suitable adjuvants, which sometimes have been referred to as immune stimulants, include, but are not limited to: cytokines, growth factors, chemokines, supernatants from cell cultures of lymphocytes, monocytes, cells from lymphoid organs, cell preparations and/or extracts from plants, bacteria or parasites (*Staphylococcus aureus* or lipopolysaccharide preparations) or mitogens. Generally, an adjuvant is administered at the same time as an antigen of the present invention. However, adjuvants can also or alternatively be administered within a two-week period prior to the vaccination, and/or for a period of time after vaccination, i.e., so long as the antigen, e.g., a recombinant MDV<sub>np</sub> of the present invention persists in the tissues.

**[0084]** The vaccines and/or immunogenic compositions of the present invention may be administered by any route such as in ovo, by parenteral administration, including intramuscular injection, subcutaneous injection, intravenous injection, intradermal injection, by scarification, by oral administration, or by any combination thereof. Furthermore, the multivalent recombinant MDV<sub>np</sub> of the present invention can be used and/or combined with additional NDV, ILTV, and/or MDV antigens to improve and expand the immunogenicity provided, and/or antigens for other pathogens in order to provide immune protection against such other pathogens. These additional antigens can be either live or killed whole microorganisms, other recombinant vectors, cell homogenates, extracts, proteins, or any other such derivative, provided that they do not negatively interfere with the safety, stability, and efficacy of the vaccine according to the present invention.

**[0085]** The combination of a multivalent recombinant MDV<sub>np</sub> of the present invention with an additional MDV, NDV, and/or ILTV antigen can be advantageous in those

cases in which very virulent field strains of MDV, NDV, or ILTV are prevalent, e.g., in a particular geographic region. In this regard, the combination of a multivalent recombinant MDV<sub>np</sub> of the present invention with an MDV1, MDV2, or HVT includes the Rispens (MDV1) strain, the SB1 (MDV2) strain, the FC-126 (HVT) strain and/or PB1 (HVT) strain. To improve the response against NDV, multivalent recombinant MDV<sub>np</sub> may be combined with an NDV vaccine strain, such as the mild live NDV vaccine strain C2.

**[0086]** Examples of other microorganisms that can be used as antigens together with the multivalent recombinant MDV<sub>np</sub> of the present invention include: (i) viruses such as infectious bronchitis virus, adenovirus, egg drop syndrome virus, infectious bursal disease virus, chicken anaemia virus, avian encephalo-myelitis virus, fowl pox virus, turkey rhinotracheitis virus, duck plague virus (duck viral enteritis), pigeon pox virus, avian leucosis virus, avian pneumovirus, and reovirus, (ii) bacteria, such as *Escherichia coli*, *Salmonella* spec., *Ornitobacterium rhinotracheale*, *Haemophilus paragallinarum*, *Pasteurella multocida*, *Erysipelothrix rhusiopathiae*, *Erysipelas* spec., *Mycoplasma* spec., and *Clostridium* spec., (iii) parasites such as *Eimeria* spec., and (iv) fungi, such as *Aspergillus* spec. In particular embodiments of the present invention, a recombinant MDV<sub>np</sub> of the present invention can be combined with a mild live IBDV vaccine strain such as D78 (cloned intermediate strain), PBG98, Cu-1, ST-12 (an intermediate strain), or 89-03 (a live Delaware variant strain) in a multivalent vaccine. Many of such strains are used in commercial vaccines.

**[0087]** The combination vaccine can be made in a variety of ways including by combining the recombinant MDV<sub>np</sub> of the present invention with preparations of virus, or bacteria, or fungi, or parasites, or host cells, or a mixture of any and/or all of these. In particular embodiments, the components for such a combination vaccine are conveniently produced separately and then combined and filled into the same vaccine container.

**[0088]** As described above, a vaccine according to the invention can be used advantageously to provide safe and effective immune protection in poultry to a multiple diseases, by a single inoculation at very young age or in ovo. Alternatively, as would be apparent to anyone skilled in the art of poultry vaccines the combinations described above also could include vaccination schedules in which the multivalent recombinant MDV<sub>np</sub> of the present invention and the additional antigen are not applied simultaneously; e.g., the recombinant MDV<sub>np</sub> may be applied in ovo, and the NDV C2 and/or the IBDV strain (e.g., 89/03) could be applied at a subsequent time/date.

**[0089]** Accordingly, the vaccines of the present invention can be administered to the avian subject in a single dose or in multiple doses. For example, a vaccine of the present invention may be applied at the day of hatch and/or in ovo at day 16-18 (Embryonation Day) ED. When multiple doses are administered, they may be given either at the same time or sequentially, in a manner and time compatible with the formulation of the vaccine, and in such an amount as will be immunologically effective. Therefore, a vaccine of the present invention may effectively serve as a priming vaccination, which later can be followed and amplified by a booster vaccination of the identical vaccine, or with a different vaccine preparation e.g., a classical inactivated, adjuvanted whole-virus vaccine.

**[0090]** The volume per dose of a vaccine of the present invention can be optimized according to the intended route of

application: in ovo inoculation is commonly applied with a volume between 0.05 and 0.5 ml/egg, and parenteral injection is commonly done with a volume between 0.1 and 1 ml/avian. In any case, optimization of the vaccine dose volume is well within the capabilities of the skilled artisan.

Sequence Table		
SEQ ID NO:	Description	Type
1	ILTV gD Glycoprotein	nucleic acid
2	ILTV gD Glycoprotein	amino acid
3	ILTV gI Glycoprotein	nucleic acid
4	ILTV gI Glycoprotein	amino acid
5	NDV F Protein (Clone 30)	nucleic acid
6	NDV F Protein (Clone 30)	amino acid
7	NDV F Protein (B1 Hitchner)	nucleic acid
8	NDV F Protein (B1 Hitchner)	amino acid
9	ILTV gD promoter	nucleic acid
10	ILTV gI promoter	nucleic acid
11	hCMV IE promoter (Truncated)	nucleic acid
12	hCMV IE promoter (Towne Strain)	nucleic acid
13	synthetic polyadenylation signal	nucleic acid
14	HSV TK polyadenylation signal	nucleic acid
15	IE-NDV F insert	nucleic acid
16	ILTV insert	nucleic acid
17	ILTV/IE-NDV F insert	nucleic acid

**[0091]** The present invention may be better understood by reference to the following non-limiting examples, which are provided as exemplary of the invention. The following examples are presented in order to more fully illustrate embodiments of the invention and should in no way be construed as limiting the broad scope of the invention.

## EXAMPLES

### Example 1

#### Construction of Recombinant HVT/NDV/ILTV Virus Vectors

**[0092]** The ability to generate herpesviruses by cotransfection of cloned overlapping subgenomic fragments was first demonstrated for pseudorabies virus [van Zijl et al., *J. Virology* 62:2191-2195 (1988)]. This procedure subsequently was employed to construct recombinant HVT vectors [see, U.S. Pat. No. 5,853,733, hereby incorporated by reference with respect to the methodology disclosed regarding the construction of recombinant HVT vectors] and was used to construct the recombinant HVT/NDV/ILTV vectors of the present invention. In this method, the entire HVT genome is cloned into bacterial vectors as several large overlapping subgenomic fragments constructed utilizing standard recombinant DNA techniques [Maniatis et al., (1982) *Molecular Cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1982); and Sambrook et al., *Molecular Cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)]. An HVT strain FC126 cosmid library was derived from sheared viral DNA cloned into the cosmid vector, pWE15 (Stratagene, now Agilent Technologies of Santa Clara, Calif.). In addition, several large genomic DNA fragments were isolated by restriction digestion with the enzyme, BamHI, and cloned into either pWE15 or the plasmid vector pSP64 (Promega, Madison Wis.). As described in U.S. Pat. No. 5,853,733, cotransfection of these fragments into chicken embryo fibroblast (CEF) cells results in the regeneration of the HVT genome mediated by homologous recombination

across the overlapping regions of the fragments. If an insertion is engineered directly into one or more of the subgenomic fragments prior to the cotransfection, this procedure results in a high frequency of viruses containing the insertion. Five overlapping subgenomic clones are required to generate FC126 HVT, and served as the basis for creating all HVT/NDV/ILTV recombinant viruses.

**[0093]** Construction of HVT/NDV/ILTV 1332-62.E1:

**[0094]** The cosmid regeneration for HVT/NDV/ILTV 1332-62.E1 was performed essentially as described in U.S. Pat. No. 5,853,733 [e.g. FIG. 8 of U.S. Pat. No. 5,853,733; redrawn, at least in part, in FIG. 1, herein]. To allow integrations into the US region of the FC126 HVT genome, the region covered by the cosmid nr. 378-50 in U.S. Pat. No. 5,853,733, was now provided from three smaller plasmids: pSY640 and 556-60.6, and one transfer plasmid (1332-47.A2), overlapping these two, and containing the ILTV/NDV expression cassettes in the US2 gene locus.

**[0095]** The set of seven linearized constructs: 3 cosmids and 4 plasmids are transfected all together into CEFs, using a standard CaCl<sub>2</sub> transfection protocol and the resulting virus stock was plaque purified one time.

**[0096]** Construction of HVT/NDV/ILTV 1332-70.B1:

**[0097]** The cosmid regeneration for HVT/NDV/ILTV 1332-70.B1 was performed essentially as described in U.S. Pat. No. 5,853,733 [e.g., FIG. 8 of U.S. Pat. No. 5,853,733; redrawn, at least in part, in FIG. 1, herein]. To allow integrations into the UL54.5 region of the FC126 HVT genome, the region covered by the cosmid nr. 407-32.1C1 in U.S. Pat. No. 5,853,733 was now provided from three smaller plasmids: 672-01.A40 and 672-07.C40, and one transfer plasmid (1332-29.4), overlapping these two, and containing the ILTV/NDV expression cassettes in the UL54.5 gene locus.

**[0098]** The set of seven linearized constructs: 4 cosmids and 3 plasmids are transfected all together into CEFs, using a standard CaCl<sub>2</sub> transfection protocol, and the resulting virus stock was plaque purified one time.

**[0099]** Construction of HVT/NDV/ILTV 1317-46.A1-1:

**[0100]** The cosmid regeneration for HVT/NDV/ILTV 1317-46.A1-1 was performed essentially as described in U.S. Pat. No. 5,853,733 [e.g., FIG. 8 of U.S. Pat. No. 5,853,733; redrawn, at least in part, in FIG. 1, herein]. To allow integrations into the US region of the FC126 HVT genome, the region covered by the cosmid nr. 378-50 in U.S. Pat. No. 5,853,733, was now provided from three smaller plasmids: pSY640 and 556-60.6, and one transfer plasmid (1317-15.1-1), overlapping these two, and containing the ILTV expression cassette inserted into the US2 gene locus. Insertion into a second site within the FC126 HVT genome was accomplished by replacing the UL region cosmid nr. 407-32.2C3 in U.S. Pat. No. 5,853,733 with the transfer cosmid (1196-05.1), containing the NDV expression cassette inserted between the HVT UL7 and UL8 genes.

**[0101]** The set of seven constructs: 1 uncut cosmid (1196-05.1), the remaining 2 linearized cosmids, and 4 linearized plasmids were transfected all together into CEFs, using a standard CaCl<sub>2</sub> transfection protocol, and the resulting virus stock was plaque purified one time.

**[0102]** Description of Subgenomic Fragments for Generating FC126 HVT:

**[0103]** Subgenomic Clone 407-32.2C3.

**[0104]** Cosmid 407-32.2C3 contains an approximately 40,170 base pair region of genomic HVT DNA [Left terminus—pos. 39,754; Afonso et al., 2001, supra; Acc.

#AF291866]. This region includes HVT BamHI fragments F', L, P, N1, E, D, and 2,092 base pairs of fragment B.

**[0105]** Subgenomic Clone 172-07.BA2.

**[0106]** Plasmid 172-07.BA2 contains a 25,931 base pair region of genomic HVT DNA. It was constructed by cloning the HVT BamHI B fragment [pos. 37,663 to 63,593; Afonso et al., 2001, supra; Acc. #AF291866] into the plasmid pSP64 (Promega, Madison Wis.).

**[0107]** Subgenomic Clone 407-32.5G6.

**[0108]** Cosmid 407-32.5G6 contains a 39,404 base pair region of genomic HVT DNA [pos. 61,852-101,255; Afonso et al., 2001, supra; Acc. #AF291866]. This region includes HVT BamHI fragments H, C, Q, K1, M, K2, plus 1,742 base pairs of fragment B, and 3,880 base pairs of fragment J.

**[0109]** Subgenomic Clone 407-32.1C1.

**[0110]** Cosmid 407-32.1C1 contains a 37,444 base pair region of genomic HVT DNA [pos. 96,095-133,538; Afonso et al., 2001, supra; Acc. #AF291866]. This region includes HVT BamHI fragments J, G, I, F, O, plus 1,281 base pairs of fragment K2, and 6,691 base pairs of fragment A.

**[0111]** Subgenomic Clone 378-50.

**[0112]** Cosmid 378-50 contains a 28,897 base pair region of genomic HVT DNA [see FIG. 8 of U.S. Pat. No. 5,853,733; redrawn, at least in part, in FIG. 1, herein]. It was constructed by cloning the HVT BamHI A fragment [pos. 126848-155744; Afonso et al., 2001, supra; Acc. #AF291866] into the cosmid pWE15.

**[0113]** Additional Insertion Fragments for Generating HVT/NDV/ILTV 1332-62.E1:

**[0114]** Subgenomic Clone 1332-47.A2.

**[0115]** The insertion plasmid 1332-47.A2 contains a 7311 base pair EcoRI fragment of the HVT unique short region [pos. 136880-144190; Afonso et al., 2001, supra; Acc. #AF291866], cloned into the plasmid pSP64 (Promega, Madison Wis.). Inserted into a unique StuI site within the HVT US2 gene [pos. 140540/140541, Afonso et al., 2001, supra; Acc. #AF291866, between amino acid residues 124 and 125] are 2 elements: a 3563 base pair Sall-HindIII fragment from ILTV, NVSL Challenge Strain, Lot #83-2 [pos. 10532-14094; Wild et al., *Virus Genes* 12:104-116 (1996); Acc.#U28832], encoding the full length genes for glycoprotein D (gD) and glycoprotein I (gI), plus partial coding regions from glycoprotein E (amino acids 1-101), and ORF5 (amino acids 734-985); and an expression cassette consisting of the HCMV IE promoter, the NDV, clone 30 strain, fusion gene (F), followed by a synthetic poly-adenylation signal. The ILTV gD, ILTV gI, and NDV F genes are transcribed in the opposite direction relative to the HVT US2 gene.

**[0116]** Subgenomic Clone pSY640.

**[0117]** Plasmid pSY640 contains an approximately 13,600 base pair region of genomic HVT DNA [pos. 126848-140540; Afonso et al., 2001, supra; Acc. #AF291866] derived from BamHI fragment A. To generate this plasmid the region of DNA located upstream of the US2 gene, beginning at the StuI site located in the US2 gene and continuing to the end of the BamHI A fragment, was cloned into the plasmid pSP64 (Promega, Madison Wis.).

**[0118]** Subgenomic Clone 556-60.6.

**[0119]** Plasmid 556-60.6 contains an approximately 12,500 base pair region of genomic HVT DNA derived from BamHI fragment A [approximate pos. 143300 to pos. 155744, Afonso et al., 2001, supra; Acc. #AF291866]. To generate this plasmid the region of DNA located downstream of the US2 gene beginning at the StuI site located in the US2

gene and continuing to the end of the BamHI A fragment was cloned into the plasmid pSP64 (Promega, Madison Wis.), and then treated with exonuclease to "chewed back" from StuI site—150 bp, and recloned into pBR322 plasmid vector.

**[0120]** Additional Insertion Fragments for Generating HVT/NDV/ILTV 1332-70.B1:

**[0121]** Subgenomic Clone 1332-29.4

**[0122]** Plasmid 1332-29.4 contains a 8,636 base pair region of genomic HVT DNA derived from the unique long region [pos. 109489-118124; Afonso et al., 2001, supra; Acc. #AF291866], cloned into a derivative of plasmid pNEB193 (deleted AatII-PvuII). It is flanked by AscI sites and includes HVT BamHI fragments I, S, plus 1337 base pairs of fragment G and 1177 base pairs of fragment F. Inserted into an XhoI site within the HVT UL54.5 open reading frame [pos. 111240/111241, Afonso et al., 2001, supra; Acc. #AF291866, between amino acid residues 21 and 22] are 2 elements: a 3563 base pair Sall-HindIII fragment from ILTV, NVSL Challenge Strain, Lot #83-2 [pos. 10532-14094; Wild et al. 1996, supra; Acc.#U28832], encoding the full length genes for glycoprotein D (gD) and glycoprotein I (gI), plus partial coding regions from glycoprotein E (amino acids 1-101), and ORF5 (amino acids 734-985); and an expression cassette consisting of the HCMV IE promoter, the NDV, clone 30 strain, fusion gene (F), followed by a synthetic poly-adenylation signal. The ILTV gD, ILTV gI and NDV F genes are transcribed in the opposite direction relative to the HVT UL54.5 gene.

**[0123]** Subgenomic Clone 672-01.A40

**[0124]** Plasmid 672-01.A40 contains a 14,731 base pair region of genomic HVT DNA derived from the unique long region [pos. 96095-110825; Afonso et al., 2001, supra; Acc. #AF291866], cloned into a derivative of plasmid pNEB193. This region includes HVT BamHI fragments G, J and 1281 base pairs of K2.

**[0125]** Subgenomic Clone 672-07.C40

**[0126]** Plasmid 672-07.C40 contains a 12,520 base pair region of genomic HVT DNA derived from the unique long region [pos. 116948-129467; Afonso et al., 2001, supra; Acc. #AF291866], cloned into a derivative of plasmid pNEB193. This region includes HVT BamHI fragments F, O and 2620 base pairs of A.

**[0127]** Additional Insertion Fragments for Generating HVT/NDV/ILTV 1317-46.A1-1:

**[0128]** Subgenomic Clone 1196-05.1.

**[0129]** Cosmid 1196-05.1 contains an approximately 40,170 base pair region of genomic HVT DNA [Left terminus—pos. 39,754; Afonso et al., 2001, supra; Acc. #AF291866] cloned into cosmid pWE15. This region includes HVT BamHI fragments F', L, P, N1, E, D, and 2,092 base pairs of fragment B. In addition an expression cassette encoding the NDV Fusion (F) gene, including the HCMV IE promoter and HSV TK poly-adenylation regulatory elements was inserted into a non-coding region between HVT UL7 and UL8 genes within BamHI fragment E [pos. 20030-20035; Afonso et al., 2001, supra; Acc. #AF291866]. The NDV F gene is transcribed the same direction as HVT UL7.

**[0130]** Subgenomic Clone 1317-15.1-1.

**[0131]** Plasmid 1317-15.1-1 contains a 7311 base pair EcoRI fragment of the HVT unique short region [pos. 136880-144190; Afonso et al., 2001, supra; Acc. #AF291866], cloned into the plasmid pSP64 (Promega, Madison Wis.). In addition, a 3563 base pair Sall-HindIII fragment from ILTV, NVSL Challenge Strain, Lot #83-2

[pos. 10532-14094; Wild et al., 1996, supra; Acc.#U28832], encoding the full length genes for glycoprotein D (gD) and glycoprotein I (gI), plus partial coding regions from glycoprotein E (amino acids 1-101), and ORF5 (amino acids 734-985) were cloned into a unique StuI site within the HVT US2 gene [pos. 140540/140541, Afonso et al., 2001, supra; Acc. #AF291866, between amino acid residues 124 and 125]. The ILTV gD and gI genes are transcribed in the opposite direction relative to the HVT US2 gene.

**[0132]** Subgenomic Clone pSY640.

**[0133]** Plasmid pSY640 contains an approximately 13,600 base pair region of genomic HVT DNA [pos. 126848-140540; Afonso et al., 2001, supra; Acc. #AF291866] derived from BamHI fragment A. To generate this plasmid the region of DNA located upstream of the US2 gene, beginning at the StuI site located in the US2 gene and continuing to the end of the BamHI A fragment, was cloned into the plasmid pSP64 (Promega, Madison Wis.).

**[0134]** Subgenomic Clone 556-60.6.

**[0135]** Plasmid 556-60.6 contains an approximately 12,500 base pair region of genomic HVT DNA derived from BamHI fragment A [approximate pos. 143300 to pos. 155744, Afonso et al., 2001, supra; Acc. #AF291866]. To generate this plasmid the region of DNA located downstream of the US2 gene beginning at the StuI site located in the US2 gene and continuing to the end of the BamHI A fragment was cloned into the plasmid pSP64 (Promega, Madison Wis.), and then treated with exonuclease to "chewed back" from StuI site ~150 bp, and recloned into pBR322 plasmid vector.

**[0136]** Standard CaCl<sub>2</sub> Transfection Protocol:

**[0137]** Secondary CEF's are seeded on 6 well culture plates and incubated at 38° C. with 5% CO<sub>2</sub> for 24 hours and confluent monolayers form. For each well a total amount of 0.25 µg DNA of cosmids and plasmids were mixed in Hepes buffer and 125 mM CaCl<sub>2</sub> was added dropwise until precipitation was imminent. This mixture was added to the CEF cell monolayer, and incubated for 2 to 3 hrs. Supernatant was removed and an overlay of 15% Glycerol was added, and kept on the cells for 1 minute. Then this was removed, washed with PBS, and fresh culture medium was added and cells were incubated for 5 days. Next, cells were harvested by trypsinization and cells from individual plates were each seeded on fresh monolayers of CEF cells in 10 cm plates and incubated until

50-90% CPE was achieved. Next, the amplified transfected cells were harvested by trypsinization, and dilutions of 10<sup>-2</sup> to 10<sup>-4</sup> were plated on 10 cm plates with CEF monolayers and incubated. The following day, the plates were covered with agar, and a number of individual plaques of HVT/NDV/ILTV were isolated and amplified on CEFs.

## Example 2

### Recombinant HVT/ND/ILTV Vaccine Protects Day-Old Chicks Against Infectious Laryngotracheitis Virus Challenge

**[0138]** Two vaccines, one comprising HVT/NDV/ILTV-1332-62E1 and the other, comprising 1332-70B1, were evaluated for efficacy in protecting chickens from an Infectious Laryngotracheitis Virus challenge. HVT/NDV/ILTV-1332-62E1 is an rHVT in which the FC126 HVT backbone comprises the nucleic acid sequence of SEQ ID NO: 17 inserted in the US2 site (see, Example 1 above). HVT/NDV/ILTV-1332-70B1 is an rHVT in which the FC126 HVT backbone comprises the nucleic acid sequence of SEQ ID NO: 17 inserted in the UL54.5 site (see, Example 1 above).

**[0139]** The vaccine preparations for both stocks of virus were prepared from stocks passaged through chicken embryo fibroblast tissue culture cells, at least 8 times, and an additional preparation of 11 tissue culture passages was prepared and tested for 1332-62E1.

**[0140]** The vaccines were administered to newly hatched, specific-antigen free (SPF) chicks by the subcutaneous route. Birds were then challenged at four weeks of age with virulent ILTV challenge virus by the intra-tracheal route and observed for 10 days for the clinical signs of the disease. The incidence of disease in these chicks was compared with controls that either received a commercial recombinant HVT/ILTV vaccine (Innovax®-ILT, from Merck Animal Health) or no vaccine. The Federal Code of Registry (9CFR) requires that at least 80% of the unvaccinated control birds must show clinical signs for a test to be valid, and at least 90% of the vaccinated birds must remain free of clinical signs to be considered to provide satisfactory protection. The results of this study are provided in Table 1 below. Both dual recombinant vaccines provided satisfactory protection against a virulent ILTV challenge.

TABLE 1

Efficacy of Multivalent HVT/NDV/ILTV Vaccine Against a Virulent ILTV Challenge						
Group	Vaccine	Dose*	Clinical Signs**	Mortality**	Clinical and Necropsy Results**	% Protection
1	1332-62.E Pass 8	2170	1/36	1/36	1/36 = 2.8%	97.2%
2	1332-62.E Pass 11	1409	0/36	0/36	0/36 = 0%	100%
3	1332-70.B Pass 8	2483	3/36	2/36	3/36 = 8.3%	91.7%
4	Innovax ®-ILT	2200	0/24	0/24	0/24 = 0%	100%
5a	Challenged Controls	NA	10/10	9/10	10/10 = 100%	0%
5b	Non-challenged Controls	NA	0/10	0/10	0/10	NA

\*Dose is described as plaque forming units (pfu)/0.2 mL dose volume.

\*\*Results are given as the number of positive birds per total number of birds (No. of positive/total).

Example 3

Recombinant HVT/ND/ILTV Vaccine Protects Day-Old Chicks Against Newcastle Disease Virus Challenge

[0141] Day-old specific-antigen free (SPF) chicks, or 19-day old embryos were vaccinated with a recombinant vaccine, HVT/NDV/ILTV-1332-62E1, tissue culture passage level 11, or a commercial recombinant HVT/NDV vaccine (Innovax®-ND, sold by Merck Animal Health) and then challenged at four weeks of age with virulent Newcastle Disease (ND) challenge virus, Texas-GB strain, by the intra-muscular route. Following a 14-day observation period, where birds were scored for clinical signs of Newcastle disease, the incidence of disease in each group of chicks was compared with unvaccinated controls. The Federal Code of Registry (9CFR) requires that at least 80% of the unvaccinated control birds must show clinical signs for a test to be valid, and at least 90% of the vaccinated birds must remain free of clinical signs for a vaccine to be considered to provide satisfactory protection. The results of this study indicate the recombinant HVT/NDV/ILTV 1332-62E1 vaccine provided satisfactory ND protection by both routes of administration.

Laryngotracheitis Virus challenge or Newcastle Disease Virus Challenge. HVT/NDV/ILTV-1317-46.1-1 is an rHVT in which the FC126 HVT backbone comprises the nucleic acid sequence of SEQ ID NO: 16 inserted into the US2 site, and the nucleic acid sequence of SEQ ID NO: 15 inserted into the UL7/8 site, i.e., in between the UL7 and UL8 genes of HVT, (see, Example 1 above). The vaccine preparation was prepared from a stock passaged through chicken embryo fibroblast tissue culture cells 15 times.

[0143] The vaccine was administered to newly hatched, specific-antigen free (SPF) chicks by the subcutaneous route. Birds were then challenged at four weeks of age with virulent Infectious Laryngotracheitis (ILT) challenge virus by the intra-tracheal route and observed for 10 days for the clinical signs of the disease, or challenged with virulent Newcastle Disease virus, Texas-GB strain, by the intra-muscular route and observed for 14 days. The incidence of disease in these chicks was compared with controls that either received a commercial recombinant HVT/ILT vaccine, HVT/ND vaccine, or no vaccine. The Federal Code of Registry (9CFR) requires that at least 80% of the unvaccinated control birds must show clinical signs for a test to be valid, and at least 90%

TABLE 2

Efficacy of Multivalent HVT/NDV/ILTV Vaccine Against a Virulent NDV Challenge							
Group	Vaccine	Dose*	Route	No. birds	Clinical Signs**	Mortality**	% Protection
1a	1332-62.E Pass 11	2160	in ovo	31	0/31 = 0%	0/31 = 0%	100%
1b	1332-62.E Pass 11	2010	SC	31	0/31 = 0%	0/31 = 0%	100%
2a	Innovax ®-ND	2046	in ovo	32	3/32 = 9.4%	2/31 = 6.3%	90.6%
2b	Innovax ®-ND	1872	SC	32	1/32 = 3%	1/32 = 3%	96.9%
3	Marek's diluent	NA	SC	12	12/12 = 100%	12/12 = 100%	0%

\*Dose is described as plaque forming units (pfu)/dose volume (0.2 mL/SC dose, 0.1 mL/in ovo dose).  
 \*\*Results are given as the number of positive birds per total number of birds (No. of positive/total).

Example 4

Recombinant HVT/ND/ILTV Vaccine Protects Day-Old Chicks Against Infectious Laryngotracheitis Virus Challenge and Newcastle Disease Virus Challenge

[0142] A vaccine, HVT/NDV/ILT-1317-46.1-1, was evaluated for efficacy in protecting chickens from either Infectious

of the vaccinated birds must remain free of clinical signs to be considered to provide satisfactory protection. The results of this study are provided in the Table 3 below. The HVT/NDV/ILT vaccine provided satisfactory protection against NDV challenge. Although, in this preliminary study the protection provided by this construct against a virulent ILTV challenge fell just short of the federal requirements, it did provide substantial protection.

TABLE 3

Efficacy of Multivalent HVT/NDV/ILTV Vaccine Against a Virulent NDV and ILTV Challenge						
Treatment Group	Dose*	Results following Challenge				
		ILT		NDV		
		No. Positive/ Total**	% Protection	No. Positive/ Total**	% Protection	% Protection
HVT/NDV/ILT 1317-46 (p15)	1356	20	4/20 = 20%	80%	0/19 = 0%	100%



TABLE 3-continued

Efficacy of Multivalent HVT/NDV/ILTV Vaccine Against a Virulent NDV and ILTV Challenge						
Treatment Group	Dose*	Results following Challenge				
		ILT		NDV		
		No. Positive/ No. Total**	% Protection	No. Positive/ No. Total**	% Protection	% Protection
Innovax ®-ILT	1740	20	1/20 = 5%	95%	—	—
Innovax ®-ND	1836	20	—	—	0/20 = 0%	100%
Placebo	N/A	10	10/10 = 100%	0%	8/8 = 100%	0%

\*Dose is described as plaque forming units (pfu)/0.2 mL dose volume.  
 \*\*Results are given as the number of positive birds (clinical signs & mortality) per total number of birds (No. of positive/total).

Example 5

Recombinant HVT/ND/ILTV in Combination with 89/03 Bursal Disease in a VACCINE AGAINST AN INFECTIOUS BURSAL DISEASE VIRUS

**[0144]** Groups of one-day-old chicks (SPF Leghorn) were inoculated with HVT/NDV/ILTV-1332-62E1 combined with IBDV 89/03 vaccine at the time of use. A separate group of chicks were vaccinated with only the IBDV vaccine at 3.5 log<sub>10</sub> TCID<sub>50</sub> per dose. Chickens were challenged at 4 weeks of age with Variant E IBDV challenge. At 10 days post-challenge, birds were euthanized and examined for body/bursa weights and gross lesions consistent with bursal disease. The results were analyzed for acceptability per the applicable 9CFR 113.331 requirements.

**[0145]** IBDV 89/03 is a licensed product used in the poultry industry to protect flocks against both the classical and variant strains of Infectious Bursal Disease virus. The target dose for IBDV 89/03 vaccine was 3.5 log<sub>10</sub> TCID<sub>50</sub> per 0.2 mL dose. The target dose for HVT/NDV/ILT was 3000 PFU per 0.2 mL dose. To achieve the target doses in the final vaccine diluent volume the HVT/NDV/ILTV-1332-62E1 vaccine was diluted to contain 6000 PFU in 0.2 mL, which is double the target dose. The 89/03 vaccine was diluted to contain 3.8 log<sub>10</sub> TCID<sub>50</sub>, which is double the target dose. For Group 1 the combination vaccine was prepared by combining equal volumes of the HVT/NDV/ILTV-1332-62E1 vaccine and the 89/03 vaccine. For Group 2, which received only the 89/03 vaccine, an equal volume of diluent was added. One day old chickens in each treatment group received 0.2 mL of the respective vaccine or placebo by the subcutaneous (SC) route (see, Table 4).

**[0146]** At hatch, chicks in each of the vaccine treatment groups were tagged with a set of randomized tag numbers assigned using the randomization program of EXCEL. In addition, birds removed from each pen at 7 days post-challenge for histological examination of bursas were randomly determined using the randomization program of EXCEL.

**[0147]** The chickens were challenged at four weeks of age with IBDV-Variant E challenge virus. Each chicken received 0.06 mL containing approximately 10<sup>2.2</sup> EID<sub>50</sub> per dose via the eyedrop route. At seven days post-challenge, 6-9 birds from each group were removed for histological evaluation of individual bursae (see, Table 5). Bursa samples were collected from each challenged chicken using care to collect tissue which had not been crushed or squeezed by the forceps. The tissue sample was placed in an individual container of 10% formalin.

**[0148]** Bursa from each chicken challenged with IBD-Var E virus was recorded as negative or positive for bursal atrophy, gross macroscopic lesions and/or lymphocyte depletion as determined by histological examination. Bursal lesions included macroscopic hemorrhage, edema/exudates, cream/yellow color, striations, or gross atrophy. Bursal atrophy was measured by individually weighing each chicken to the nearest gram. Bursae were individually weighed to the nearest hundredth of a gram. Bursa/body weight ratios were computed for each bird employing the formula, BW ratio: (Bursa Weight ÷ Body Weight) X 1000. A bursa to body weight ratio of more than 2 standard deviations from the challenged control is considered negative for and protective from infectious

TABLE 4

EXPERIMENTAL DESIGN						
Group	No.	Vaccine	Dose HVT-(89/03)	Age	IBDV Variant E	
					# birds	Necropsy
1	45	HVT/NDV/ILT + 89/03	3000- (3.5 log <sub>10</sub> TCID <sub>50</sub> )	4 wks	≧40	10 day post-challenge
2	45	89/03	NA- (3.5 log <sub>10</sub> TCID <sub>50</sub> )	4 wks	≧40	10 day post-challenge
3	45	Placebo challenged controls	—	4 wks	≧40	10 day post-challenge
4	30	Placebo non-challenged controls	—	—	≧25	10 day post-challenge

bursal disease. The results of this study showed that vaccine treatment Groups 1 and 2 were negative for IBD (i.e., not statistically different from the placebo non-challenged control) indicating that both vaccines were efficacious and further demonstrating that there was no interference of the protection provided by the 89/03 strain of the vaccine against the IBDV challenge due to the recombinant HVT/NDV/ILT construct also being present in the multivalent vaccine (see, Table 5).

TABLE 5

Day 7 NECROPSY DATA FOR IBDV VARIANT E CHALLENGE			
Group	No.	Vaccine	Average Bursa BW ratio
1	9	HVT/NDV/ILT + 89/03	5.464
2	9	89/03	5.715

TABLE 5-continued

Day 7 NECROPSY DATA FOR IBDV VARIANT E CHALLENGE			
Group	No.	Vaccine	Average Bursa BW ratio
3	9	placebo challenged controls	1.874 (SD + 0.641)**
4	6	placebo non-challenged controls	5.838

\*\*2 SD from Control is statistically different.

Example 6

Sequences

[0149] The following sequences have been used in the exemplary rHVT constructs. The coding sequences provided below include individual stop codons, which can be readily replaced with alternative stop codons without modifying the properties of the protein antigens that the coding sequences encode.

ILTV gD Glycoprotein, coding sequence

(SEQ ID NO: 1)

```

ATGCACCGTCTCTCATCTCAGACGGCACTCGCGTTACTACGCGAAAGGAGAGGTGCTTAACAACACAT
GGATTGCGGTGAAAAACGGTGTCTGCTCAGGCGCAGCTGATTCACTCTTTTCTGGACTTGTGTGAGGA
TTATGCGGGAGCATATCTGCTTTGTACGCAACGCTATGGACCGCCATTTATTTTTGAGGAATGCTTTT
TGGACTATCGTACTGCTTTCTTCTTCGCTAGCCAGAGCACCAGCGCGCTCACGTACGACTACATTTT
AGGCCGTCGCGGCTCGACGCGCTAACCATACCGGCGGTTGGCCGTATAACAGATACCTCACTAGGG
TATCAAGAGGCTGCGACGTTGTGAGCTCAACCCGATTTCTAACGTGGACGACATGATATCGGCGGCC
AAAGAAAAGAGAAGGGGGCCCTTTTCGAGGCCTCCGTCGTCGGTCTACGTGATTAAGGGCGACGA
CGGCGAGGACAAGTACTGTCCAATCTATAGAAAAGAGTACAGGGAATGTGGCGACGTACAACCTGCTAT
CTGAATGCGCCGTTCAATCTGCACAGATGTGGCAGTGGACTATGTTCTTAGCACCCCTTGATCGCGA
AATGGCGCGGACTGACTATATTTCTCCCCACTGCTGCGCTCTCTGGCCAATACTTGCTGACCCTGAA
AATCGGGAGATTTGCGCAAACAGCTCTCGTAACTCTAGAAGTTAACGATCGCTGTTTAAAGATCGGGT
CGCAGCTTAACTTTTACCGTCGAAATGCTGGACAACAGAACAGTATCAGACTGGATTTCAAGGCGAA
CACCTTTATCCGATCGCAGACACCAATACACGACACGCGGACGACGTATATCGGGGATACGAAGATAT
TCTGCAGCGCTGGAATAATTTGCTGAGGAAAAAGAATCCTAGCGCGCCAGACCCTCGTCCAGATAGCG
TCCCGCAAGAAATCCCGCTGTAACCAAGAAAGCGGAAGGGCGCACCCCGGACGCGAAGAGCAGCGAA
AAGAAGGCCCTCCAGAAGACTCGGAGGACGACATGCAGGACAGGGCTTCTGGAGAAAATCTGCGCC
CCTCCCCGAAGACGACGAAGTCCCGAGGACACCGAGCAGATGATCCAAACTCGGATCTGACTATT
ACAATGACATGCCCGCGTATCCCGTGGAGGAGACTACTAAAAGTTCTAATGCCGTCTCCATGCC
ATATTCGCGCGTTCGTAGCCTCGCGGTCGCGCTCGTGGGGCTACTGGTTTGGAGCATCGTAAAATG
CGCGCTAGCTAA
    
```

ILTV gD Glycoprotein

(SEQ ID NO: 2)

```

MHRPHLRRHSRYAKGEVLNKHMDCCGKRCCSGAAVFTLFWTCVRIMREHICFVRNAMDRHLFLRNF
WTIVLLSFSASQSTAAVTYDYILGRRALDALTI PAVGPNRYLTRVSRGCDVVELNPI SNVDDMISAA
KEKEKGGPF EASVWVYVIKGGDDGDKYCP IYRKEYRECGDVQLLSECAVQSAQMWAVDYVPSTLVS
NGAGLTIFSP TAAALSGYLLTLKIGRFAQTALVTLEVNDRLKIGSQLNFLPSKWTTEQYQTFQGE
HLYPIADTNTRHADDVYRGYEDILQRWNNLLRKNPSAPDPRPDSVPQEI PAVTKKAEGRTPDAESSE
    
```

- continued

KKAPPEDEEDMQAEASGENPAALPEDEVPEDTEHDDPNSDPDYNDMPAVIPVEETKSSNAVSM

IFAAFVACAVALVGLLVMSIVKCARS

ILTV gI Glycoprotein, coding sequence

(SEQ ID NO: 3)

ATGGCATCGTACTTGGAACTCTGGCTCTCCTTGCCGCGACGCTCGCACCCCTTCGGCGCGATGGGAAT

CGTGATCACTGGAAATCACGTCTCCGCCAGGATTGACGACGATCACATCGTGATCGTCGCGCTCGCC

CCGAAGTACAATTCAACTGCAGCTATTTTCATGCCTGGCCAGAGACCCACAAACCTACTCAGGA

ACCGTCCGCGTCGCGTTTCGGTCTGATATAACAAACAGTGTACACAGGAACCTTAGCGAGGAGCGCTT

TGAAAATTGCACTCATCGATCGTCTTCTGTTTTTGTGGCTGTAAAGTGACCGAGTACACGTTCTCCG

CCTCGAAGACTAACCGGACCTCCACACCCGTTTAAAGCTCACTATACGAAATCCTCGTCCGAACGAC

AGCGGGATGTTCTACGTAATTGTTTCGGCTAGACGACACCAAAGAACCATTGACGTTCTCGCGATCCA

ACTATCGGTGTATCAATTTCGGAACACCGCGGACTCGCGGACTCTATTCCAAGGCTTCGTGTGCA

CCTTCGGATTACCTACCGTCCAACCTGAGGCTATCTCAGGACCGAGGAAAGTTGGCGCAACTGGCAA

GCGTACGTTGCCACGGAGGCCACGACGACCGCGGAGGCGACAACCCGACGCCCCTACTGCAAC

CAGCGCTCCGAACTTGAAGCGGAACACTTTACCTTTCCCTGGCTAGAAAATGGCGTGGATCATTACG

AACCGACACCCGCAACGAAAATTCAAACGTTACTGTCCGTCTCGGGACAATGAGCCCTACGCTAATT

GGGGTAACCGTGGCTGCCGTCGTGAGCGCAACGATCGGCCCTCGTCATTGTAATTTCCATCGTACCAG

AAACATGTGCACCCCGCACGAAAATTAGACACGGTCTCGAAGACGACGAGAAGCTCCCAAATA

GAAGGGAATCGCGAAAATTTGAGCCCATGTTGCGTGCAAAATAAACAGGGGGCTGACCAGGATAGT

GAACTTGTGGAAGTGGTTGCGATGTTAACCCGTCTGCGCTAAGCTCGCCCGACTCAATAAAAATGTG

A

ILTV gI Glycoprotein

(SEQ ID NO: 4)

MASLLGLTALLAATLAPFGAMGIVITGNHVSARIDDDHIVIVAPRPEATIQLQLFFMPGQRPHPKPYSG

TVRVAFRSDITNQCYQELSBERFENCTHRSSSVFVGKVTETPFSASNRLTGPHFPKLTIRNRPND

SGMFYVIVRLDDTKEPIDVFAIQLSVYQFANTAATRGLYSKASCRTPFGLPTVQLEAYLRTEESWRNWQ

AYVATEATTTSAEATTPPVATASASELEAEHFTFPWLENGVDHYEPTPANENSNVTVRLGTMSPTLI

GVTVAAVVSATIGLVIVISIVTRNMCTPHRKLDTVSQDDEERSQTRRESRKFQPMVACEINKGADQDS

ELVELVAIVNPSALSSPDSIKM

NDV F Protein, coding sequence

(SEQ ID NO: 5): Clone 30

ATGGGCCCCAGACCTTCTACCAAGAACCAGTACCTATGATGCTGACTGTCCGAGTCGCGCTGGTACT

GAGTTGCATCTGTCCGCAAACTCCATTGATGGCAGGCCTCTTGCAGCTGCAGGAATTGGTTACAG

GAGACAAAGCCGTCAACATATACACCTCATCCAGACAGGATCAATCATAGTTAAGCTCCTCCGAAT

CTGCCAAAGGATAAGGAGGATGTGCGAAAGCCCCCTGGATGCATACAACAGGACATTGACCCTTT

GCTCACCCCTTGGTACTCTATCCGTAGGATACAAGAGTCTGTGACTACATCTGGAGGGGGAGAC

AGGGCGCCTTATAGGCGCCATTATTGGCGGTGTGGCTCTTGGGGTTGCAACTGCCGCACAAATAACA

GCGGCCGAGCTCTGATACAAGCCAAACAAAATGCTGCCAACATCCTCCGACTTAAAGAGAGCATTGC

CGCAACCAATGAGGCTGTGATGAGGTCACGACGATTATCGCAACTAGCAGTGGCAGTTGGGAAGA

TGCAGCAGTTTGTAAATGACCAATTTAATAAAAACAGCTCAGGAATTAGACTGCATCAAAATGACACAG

CAAGTTGGTGTAGAGCTCAACCTGTACCTAACCGAATGACTACAGTATTCGGACCACAAATCACTTC

ACCTGCTTTAAACAAGCTGACTATTACGGCACTTTACAATCTAGCTGGTGGAAATATGGATTACTTAT

- continued

TGACTAAGTTAGGTGTAGGGAACAATCAACTCAGCTCATTAAATCGGTAGCGGCTTAATCACCGGTAAC  
CCTATTCTATACGACTCACAGACTCAACTCTTGGGTATACAGGTAACCTACCTTCAGTCGGGAAGCT  
AAATAATATGCGTGCCACCTACTTGAAACCTTATCCGTAAGCACAACCAGGGGATTTGCCTCGGCAC  
TTGTCCCAAAGTGGTGACACAGGTCGGTCTGTGATAGAAGAACTTGACACCTCATACTGTATAGAA  
ACTGACTTACATTTATATTGTACAAGAATAGTAACGTTCCCTATGTCCCTGGTATTTATTCCTGCTT  
GAGCGGCAATACGTCGGCCTGTATGTACTCAAAGACCGAAGGCGCACTTACTACACCATACATGACTA  
TCAAAGTTTCAGTCATCGCCAAC TGCAAGATGACAACATGTAGATGTGTAACCCCCGGGTATCATA  
TCGCAAACTATGGAGAAGCCGTCTCTAATAGATAAACAATCATGCAATGTTTTATCCTTAGCGGG  
GATAACTTTAAGGCTCAGTGGGAATTGATGTAACCTATCAGAAGAATATCTCAATACAAGATTCTC  
AAGTAATAATAACAGGCAATCTTGATATCTCAACTGAGCTTGGGAATGTCAACAACCTCGATCAGTAAT  
GCTTTGAATAAGTTAGAGGAAAGCAACAGAAAAC TAGACAAAGTCAATGTCAAACTGACTAGCACATC  
TGCTCTCATTACCTATATCGTGTGACTATCATATCTCTTGTTTTTGGTATACTTAGCCTGATTCTAG  
CATGCTACCTAATGTACAAGCAAAAGGCGCAACAAAAGACCTTATATGGCTTGGGAATAATACTCTA  
GATCAGATGAGAGCCACTACAAAATGTGA

NDV F Protein

(SEQ ID NO: 6): Clone 30

MGPRPSTKNPVPMLLTVRVALVLSICIPANSIDGRPLAAAGIVVTGDKAVNIYSSQTGSIIVKLLPN  
LPKDKKACAKAPLDAYNRTLTLTLLTPLGDSIRRIQESVTTSGGGRQRLIGAIIGGVALGVATAAQIT  
AAAAALIQAQNAANILRLKESIAATNEAVHEVTDGLSQLAVAVGKMQQFVNDQFNKTAQELDCIKIAQ  
QVGVELNLYLTELTVFGPQITSPALNKLTIQALYNLAGNMDYLLTKLGVGNQLSSLIGSGLITGN  
PILYDSQTQLLGIQVTLPSVGLNMRATYLETLSVSTTRGFASALVPKVVTVQGSVIEELDTSYICIE  
TDLHLYCTRIVTFPMSPGIYSCLSGNTSACMYSKTEGALTPYMTIKGSVIANCKMTTCRCVNPPII  
SQNYGEAVSLIDKQSCNVLSLGGITLRLSGEFDVYQKNISIQDSQV IITGNLDISTELGNVNNISIN  
ALNKLEENRKLDKVNVKLTSTALSALITYIVLTIIISLVFGILSLILACYLMIKQKQKTLNLWLGNNTL  
DQMRATTKM

NDV F Protein, coding sequence

(SEQ ID NO: 7): (B1 Hitchner)

ATGGATCGATCCCGGTTGGCGCCCTCCAGGTGCAGGATGGGCTCCAGACCTCTACCAAGAAGCCAGC  
ACCTATGATGCTGACTATCCGGGTCGCGCTGGTACTGAGTTGCATCTGTCCGGCAAACCTCATTGATG  
GCAGGCCTCTTGAGCTGCAGGAATTGTGGTTACAGGAGACAAAGCAGTCAACATATACACCTCATCC  
CAGACAGGATCAATCATAGTTAAGCTCCTCCGAATCTGCCAAGGATAAGGAGGCATGTGCGAAAGC  
CCCTTGGATGCATACAACAGGACATTGACCACTTTGCTCACCCCCCTGGTACTCTATCCGTAGGA  
TACAAGAGTCTGTGACTACATCTGGAGGGGGAGACAGGGGCGCCTTATAGGCGCCATTATTGGCGGT  
GTGGCTCTTGGGTTGCAACTGCCGCACAAATAACAGCGGCGCAGCTCTGATACAAGCCAAACAAA  
TGCTGCCAACATCCTCCGACTTAAAGAGAGCATTGCCGCAACCAATGAGGCTGTGCATGAGGTCAGT  
ACGGATTATCGCAACTAGCAGTGGCAGTTGGGAAGATGCAGCAGTTCGTTAATGACCAATTTAATAAA  
ACAGCTCAGGAATTAGACTGCATCAAATTCACAGCAAGTTGGTGTAGAGCTCAACCTGTACCTAAC  
CGAATCGACTACAGTATTCGGACCACAAATCACTTCACTGCCTTAAACAAGCTGACTATTACGGCAC  
TTTACAATCTAGCTGGTGGGAATATGGATTACTTATTGACTAAGTTAGGTATAGGAAACAATCAACTC  
AGCTCATTAAATCGGTAGCGGCTTAATCACCGGTAACCTATTCTATACGACTCACAGACTCAACTCTT  
GGGTATACAGGTAACCTTACCTTCAGTCGGGAACCTAAATAATATGCGTGCCACCTACTTGGAAACCT

- continued

TATCCGTAAGCACAAACCAGGGGATTTCCTCGGCACCTTGCCAAAAGTGGTGACACGGGTCGGTTCT  
 GTGATAGAAGAACTTGACACCTCATACTGTATAGAACTGACTTAGATTTATTTGTACAAGAATAGT  
 AACGTTCCCTATGTCCCTGGTATTTACTCCTGCTTGAGCGGCAATACATCGGCCTGTATGTACTCAA  
 AGACCGAAGGCGCACTTACTACCCATATATGACTATCAAAGGCTCAGTCATCGCTAACTGCAAGATG  
 ACAACATGTAGATGTGTAACCCCGGGTATCATATCGCAAACATATGGAGAAGCCGTGTCTCTAAT  
 AGATAACAATCATGCAATGTTTTATCCTTAGCGGGATAACTTTAAGGCTCAGTGGGAATTTCGATG  
 TAACTTATCAGAAGAATATCTCAATACAAGATTCTCAAGTAATAATAACAGGCAATCTTGATATCTCA  
 ACTGAGCTTGGGAATGTCAACAACTCGATCAGTAATGCCTTGAATAAGTTAGAGGAAAGCAACAGAAA  
 ACTAGACAAAGTCAATGTCAAACAGCAGCACATCTGCTCTCATTACCTATATCGTTTTGACTATCA  
 TATCTCTGTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAAAGGCGCAA  
 CAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAGATGAGAGCCACTACAAAATGTGA

NDV F Protein

(SEQ ID NO: 8): (B1 Hitchner)

MDRSRLAPSRCRMGRSPSTKNPAPMMLTIRVALVLSICPANSIDGRPLAAAGIVVTGDKAVNIYTSS  
 QTGSIIVKLLPNLPKDKKAKAPLDAYNRTLTLTLLPLGDSIRRIQESVTTSGGGRQRLIGAIIGG  
 VALGVATAAQITAAAALIQAKQNAANILRLKESIAATNEAVHEVTDGLSQLAVAVGKMQQFVNDQFNK  
 TAQELDCIKIAQQVGVELNLYLTESTTVFGPQITSPALNKLTIQALYNLAGGNMDDLTKLIGINNQL  
 SSLIGSLITGNPILYDSQTLQLLGIQVTLPSVGNLNMNRATYLETLVSTTRGFASALVPKVTVRVGS  
 VIEELDTSYCIETDLDLYCTRIVTFPMSPGIYSCLSGNTSACMYSKTEGALTPYMTIKGSVIANCKM  
 TTCRCVNPPIISQNYGEAVSLIDKQSCNVLSLGGITLRLSGEPDVTYQKNISIQDSQVITGNLDIS  
 TELGNVNNSISNALNKLEESNRKLDKVNKLTSTLSALITYIVLTIISLVFGILSLILACYLQKQKQKQ  
 QKTLWLGNNTLDQMRATTKM

ILTV gD Promoter

(SEQ ID NO: 9)

AAACAGCTGTACTACAGAGTAACCGATGGAAGAACATCGGTCCAGCTAATGTGCCTGTGTCGACAGG  
 CCATTCCTCGGAACCTTACTGTCTTTTCGACACGCTCTTATAGCGAGGAAAAAGATATCGGCCAG  
 AGTTATACTTTACCTCTGATCCGCAAACGGCATACTGCACAATAACTCTGCCGTCGGCGTTGTTCCG  
 AGATTCGAATGGAGCCTTAATAATGTTTCACTGCGGAATATTTGACGGCCACGACCGTTGTTTCGA  
 TACCGCTGGCCAAAGTACAGTGTGGAAGAGCAGCGGAGAGCAGCGAGGCGTGGATTCTGGCCGGG  
 GAGGCAATATATACGAATGCACCGTCTCATCTCAGACGGCACTCGCGTTACTACGCGAAAGGAGAGG  
 TGCTTAAACAACACATGGATTGCGGTGGAACCGGTGCTGCTCAGGCGCAGCTGTATTCACTCTTTTC  
 TGGACTTGTGTCAGGATTATGCGGGAGCATATCTGCTTTGTACGCAACGCT

ILTV gI Promoter

(SEQ ID NO: 10)

TGACTATTACAATGACATGCCCGCGTGATCCCGTGGAGGAGACTACTAAAAGTTCTAATGCCGTCT  
 CCATGCCATATTCGCGCGTTCGTAGCCTGCGCGTTCGCGCTCGTGGGGCTACTGGTTGGAGCATC  
 GTAAAATGCGCGGTAGCTAATCGAGCCTAGAATAGGTGGTTTCTTCTACATGCCACGCCTCACGCT  
 CATAATATAAATCACATGGAATAGCATACCAATGCCATTATTGGGACGTCGAAAAGC

hCMV IE Promoter

(SEQ ID NO: 11): (Truncated)

CGCGCCAGGTCAATTCCCTGGCATTATGCCAGTACATGACCTTATGGGACTTTCCCTACTTGGCAGTA  
 CATCTACGTATTAGTCATCGCTATTACCATGGTGTGCGGTTTTGGCAGTACATCAATGGGCGTGGAT  
 AGCGGTTGACTCACGGGGATTTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTGTTTTGGCAC

- continued

CAAAATCAACGGGACTTTCCAAAATGTCGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAGCGT  
GTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCC  
ACGCTGTTTTGACCTCCATA

hCMV IE Promoter

(SEQ ID NO: 12): (Towne Strain)

GTGAATAATAAAATGTGTGTTGTCGAAAATACGCGTTTGAGATTTCTGTCCCGACTAAATTCATGTC  
GCGCGATAGTGGTGGTTTATCGCCGATAGAGATGGCGATATTGGAAAAATCGATTTGAAAAATAGGC  
ATATTGAAAATGTCGCCGATGTGAGTTTCTGTGTAAGTATCGCCATTTTCCAAAAGTTGATTTT  
TGGGCATACGCGATATCTGGCGATACGCTTATATCGTTTACGGGGATGGCGATAGACGCCTTTGGTG  
ACTTGGGCGATTCTGTGTGTCGCAAAATATCGCAGTTTCGATATAGGTGACAGACGATATGAGGCTATA  
TCGCCGATAGAGGCGACATCAAGCTGGCACATGGCCAAATGCATATCGATCTATACATTGAATCAATAT  
TGGCCATTAGCCATATTATTCATTGGTTATATAGCATAAAATCAATATTGGCTATTGGCCATTGCATAC  
GTTGTATCCATATCATAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCCATGTTGACATT  
GATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCAATTAGTTTATAGCCCATATATGGAGTTC  
CGCGTTACATAACTTACGGTAAATGGCCCGCTGGCTGACCGCCCAACGACCCCCGCCATTGACGTC  
AATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCATTGACGTCAATGGGTGGAGTATT  
TACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCATTGACGTC  
AATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTGGCA  
GTACATCTACGTATTAGTCATCGCTATTACCATGGTGTGCGGTTTGGCAGTACATCAATGGGCGTG  
GATAGCGGTTTACTCACGGGATTTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTGTTTTGG  
CACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACCTCCGCCCCATTGACGCAAATGGGCGGTAG  
GCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCC  
ATCCACGCTGTTTTGACCTCCATAGAAGACACCGG

Synthetic Polyadenylation Signal

(SEQ ID NO: 13)

GGAAATCTAGATCCCACGTCACCTATTGTATACTCTATATTATACTCTATGTATACTCTGTAATCCTA  
CTCAATAAACGTGTCACGCCCTGTGAAACCGTACTAAGTCTCCCGTGTCTTCTTATCACCATCAGGTGA  
CATCCTCGCCAGGCTGTCAATCATGCCGTATCGATTCCAGTAGCACCGGCCCCACGCTGACAACCC  
ACTCTTGACGCTTAGCAGCGCCCTCTTAACAAGCCGACCCCCACCAGCGTCGCGGTTACTAACACT  
CCTCTCCCC

HSV TK polyadenylation signal

(SEQ ID NO: 14)

GGGAGATGGGGGAGGCTAACTGAAACACGGGAAGGAGACAATACCGGAAGGAACCCGCGCTATGACGGC  
AATAAAAAGACAGAATAAAACGCACGGGTGTTGGGTCGTTTGTTCATAAACGCGGGTTCGGTCCCAG  
GGCTGGCACTCTGTGCATACCCACCGAGACCCCATTTGGGACCAATACGCCCGCTTCTTCTTTTC  
CCCACCCCAACCCCAAGTTCCGGTGAAGGCCAGGGCTCGCAGCCAACGTGGGGCGGCAAGCCCTG  
CCATAGCCACGGGCCCGTGGGTTAGGACGGGTCCCCATGGGGAATGGTTTATGGTTCGTGGGGG  
TTATTATTTTGGGCGTTGCGTGGGTCAGGTCCACGACTGGACTGAGCAGACAGACCATGGTTTTTG  
GATGGCCTGGGCATGACCCGATGTACTGGCGCAGACGAACACCGGGCTCTGTGGCTGCCAAACAC  
CCCCGACCCCAAAAACCCCGCGGATTTCTGGCGCCCGGACG

-continued

IE-NDV F Cassette Insert (1317-46 virus)  
(SEQ ID NO: 15): (3593 bp)  
TAATTAACCCGGGAAGCTTGCATGCCTGCAGTGAATAATAAAATGTGTGTTTGTCCGAAATACGCGTT  
TGAGATTTCTGTCCCGACTAAATTCATGTCGCGCGATAGTGGTGTATTCGCGGATAGAGATGGCGAT  
ATTGAAAAATCGATATTTGAAAAATATGGCATATTGAAAAATGTCGCGGATGTGAGTTTCTGTGTAAT  
GATATCGCCATTTTTCCAAAAGTTGATTTTTGGGCATACGCGATATCTGGCGATACGCTTATATCGTT  
TACGGGGGATGGCGATAGACGCCCTTGGTGACTTGGCGGATTCGTGTGTCGCAAAATATCGCAGTTTC  
GATATAGGTGACAGACGATATGAGGCTATATCGCCGATAGAGGCGACATCAAGCTGGCACATGGCCAA  
TGCATATCGATCTATACATTTGAATCAATATTGGCCATTAGCCATATTATTCATTGGTTATATAGCATA  
AATCAATATTGGCTATTGGCCATTGCATACGTTGTATCCATATCATAATATGTACATTTATATTGGCT  
CATGTCCAACATTACGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGG  
TCATTAGTTCATAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCTGGCTG  
ACCGCCCAACGACCCCGCCCATTGACGTCATAATGACGTATGTTCCCATAGTAACGCCAATAGGGA  
CTTTCATTGACGTCAATGGGTGGAGTATTACGGTAACTGCCCACTTGGCAGTACATCAAGTGTAT  
CATATGCCAAGTACGCCCCCTATTGACGTCATGACGGTAAATGGCCCGCTGGCATTATGCCCAGTA  
CATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGA  
TGCGGTTTTGGCAGTACATCAATGGGCGTGATAGCGGTTTACTCACGGGGATTTCCAAGTCTCCAC  
CCCATTGACGTCAATGGGAGTTTGTTTTGGCACCAAAATCAACGGGACTTTCAAAATGTGTAACAA  
CTCCGCCCATTTGACGCAAAATGGGCGGTAGGCGTGTACGGTGGGAGTCTATATAAGCAGAGCTCGTT  
TAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGAC  
CATGGATCGATCCCGTTGGCGCCCTCCAGGTGCAGGATGGGCTCCAGACCTTCTACCAAGAACCAG  
CACCTATGATGCTGACTATCCGGTTCGCGCTGGTACTGAGTTGCATCTGTCCGGCAACTCCATTGAT  
GGCAGGCTCTTGACGCTGCAGGAATTGTGTTACAGGAGACAAAGCAGTCAACATATACACCTCATC  
CCAGACAGGATCAATCATAGTTAAGCTCCTCCGAATCTGCCAAAGGATAAGGAGGCATGTGCGAAAG  
CCCCCTTGATGCATACAACAGGACATGACCACCTTGTCTCACCCCTTGGTGACTCTATCCGTAGG  
ATACAAGAGTCTGTGACTACATCTGGAGGGGGAGACAGGGCGCCTTATAGGCGCATTATTGGCGG  
TGTGGCTCTTGGGTTGCAACTGCCGCACAAATAACAGCGGCCGAGCTCTGATACAAGCCAAACAAA  
ATGCTGCCAATCCTCCGACTTAAAGAGAGCATTGCCGCAACCAATGAGGCTGTGCATGAGGTCCT  
GACGGATTATCGCAACTAGCAGTGGCAGTTGGGAAGATGCAGCAGTTCGTTAATGACCAATTTAATAA  
AACAGCTCAGGAATTAGACTGCATCAAAATGACACAGCAAGTTGGTGTAGAGCTCAACCTGTACTTAA  
CCGAATCGACTACAGTATTCGGACCACAAATCACTTCACTGTCCTTAAACAAGCTGACTATTTCAGGCA  
CTTTACAATCTAGCTGGTGGGAATATGGATTACTTATTGACTAAGTTAGGTATAGGGAACAATCAACT  
CAGCTCATTAATCGGTAGCGGCTTAATCACCGTAACCTATTCTATACGACTCACAGACTCAACTCT  
TGGGTATACAGGTAATCTACCTTCAGTCGGGAACCTAAATAATATGCGTGCCACCTACTTGGAACC  
TTATCCGTAAGCACAACCAGGGGATTTGCCTCGGCACTTGTCCTCAAAAGTGGTGACACGGGTCGGTTC  
TGTGATAGAAGAACTTGACACCTCATACTGTATAGAACTGACTTAGATTTATATTGTACAAGAATAG  
TAACGTTCCCTATGTCCCTGGTATTTACTCCTGCTTGAGCGGCAATACATCGGCCTGTATGTACTCA  
AAGACCGAAGGCGCACTTACTACACCATATATGACTATCAAAGGCTCAGTCATCGCTAACTGCAAGAT  
GACAACATGTAGATGTGTAACCCCGGGTATCATATCGCAAACTATGGAGAAGCCGTGTCTCTAA  
TAGATAACAATCATGAATGTTTTATCCTTAGGCGGATAACTTTAAGGCTCAGTGGGAATTTCGAT

- continued

GTAACCTATCAGAAGAATATCTCAATACAAGATTCTCAAGTAATAATAACAGGCAATCTTGATATCTC  
AACTGAGCTTGGGAATGTCAACAACCTCGATCAGTAATGCC TTGAATAAGTTAGAGGAAAAGCAACAGAA  
AACTAGACAAAGTCAATGTCAAACCTGACCAGCACATCTGCTCTCATTACCTATATCGTTTTGACTATC  
ATATCTCTGTTTTTTGGTATACTTAGCCTGATTCTAGCATGCTACCTAATGTACAAGCAAAGGCGCA  
ACAAAAGACCTTATTATGGCTTGGGAATAATACCCTAGATCAGATGAGAGCCACTACAAAATGTGAA  
CACAGATGAGGAACGAAGTTTCCCTAATAGTAATTTGTGTGAAAGTTCTGGTAGTCTGT CAGTTCCG  
AGAGTTAAGAAAAAAAAAAAAACCCCCCCCCCCCCCCCCCCCCCTGGGTACGATCCTCTAGAGTC  
GGGAGATGGGGGAGGCTAACTGAAACACGGAAGGAGACAATACCGGAAGGAACCCGCGCTATGACGGC  
AATAAAAAGACAGAATAAAACGCACGGGTGTTGGGTCGTTTGTTCATAAACCGGGGTTCCGGTCCCAG  
GGCTGGCACTCTGTGCATACCCACCGAGACCCCAATTGGGACCAATACGCCCGCTTTCTTCTTTTC  
CCCACCCCAACCCCAAGTTCGGGTGAAGGCCAGGGCTCGCAGCCAACGTCGGGGCGGCAAGCCCTG  
CCATAGCCACGGGCCCCGTTGGGTAGGGACGGGTCCCCATGGGGAATGGTTTATGGTTCGTGGGG  
TTATTATTTTGGCGTTGCGTGGGTGAGTCCACGACTGGACTGAGCAGACAGACCCATGGTTTTTG  
GATGGCTGGGCATGGACCGCATGTACTGGCGGACACGAACCCGGGCGTCTGTGGCTGCCAAACAC  
CCCCGACCCCAAAAACCCCGCGGGATTCTGGCGCCGGGACGTCGACTTAAT

ILTV insert sequence

(SEQ ID NO: 16)

(3563 bp SalI - HindIII fragment):

gTCGACGGCAGAGTCGAGACGCCCTATTGGACGTCAAAATTGTAGAGGTGAAGTTTTCAACGATG  
GCGAAGTAACGGCGACTTGCCTTTCCACCGTCAAATCTCCCTATAGGGTAGAAACTAATTGGAAAGTA  
GACCTCGTAGATGTAATGGATGAAATTTCTGGGAACAGTCCCGCCGGGTTTTTAAACAGTAATGAGAA  
ATGGCAGAAAACAGCTGTACTACAGAGTAACCGATGGAAGAACATCGGTCCAGCTAATGTGCCTGTCTG  
GCACGAGCCATTCTCCGGAACCTTACTGTCTTTTCGACACGTCTCTTATAGCGAGGGAAAAAGATATC  
GCGCCAGAGTTATACTTTACCTCTGATCCGCAAACGGCATACTGCACAATAACTCTGCCGTCGGCGT  
TGTTCCGAGATTGCAATGGAGCCTTAATAATGTTTCACTGCCGGAATATTTGACGGCCACGACCGTTG  
TTTCGCATACCGCTGGCCAAAGTACAGTGTGGAAGAGCAGCGGAGAGCAGGCGAGGCGTGGATTTCT  
GGCCGGGAGGCAATATATACGAATGCACCGTCCCTCATCTCAGACGGCACTCGCGTTACTACGCGAAA  
GGAGAGTGTCTTAACAAAACATGGATTGCGGTGGAACCGGTGCTGCTCAGGCGCAGCTGTATTAC  
TCTTTCTGGACTTGTGTCAGGATTATGCGGGAGCATATCTGCTTTGTACGCAACGCTATGGACCGCC  
ATTTATTTTTGAGGAATGCTTTTTGGACTATCGTACTGCTTCTTCTTCCTCGTAGCCAGAGCACCGCC  
GCCGTCACGTACGACTACATTTTAGGCCGTCGCGCGCTCGACGCGCTAACCATACCGGCGGTTGGCCC  
GTATAACAGATACCTCACTAGGGTATCAAGAGGCTGCGACGTTGTGAGCTCAACCCGATTTCTAACG  
TGGACGACATGATATCGCGGCCAAAGAAAAGAGAAGGGGGCCCTTTCGAGGCTCCGTCGTCTGG  
TTCTACGTGATTAAGGGCGACGACGGCGAGGACAAGTACTGTCCAATCTATAGAAAAGAGTACAGGGA  
ATGTGGCGACGTACAACCTGCTATCTGAATGCGCCGTTCAATCTGCACAGATGTGGCAGTGGACTATG  
TTCTAGCACCCCTGTATCGCGAAATGGCGGGACTGACTATATTTCCCCCACTGCTGCGCTCTCT  
GGCCAACTACTGTGACCTGAAAATCGGGAGATTGCGCAAACAGCTCTCGTAACTCTAGAAGTTAA  
CGATCGCTGTTTAAAGATCGGGTCGACGTTAACTTTTACCGTCGAAATGCTGGACAACAGAACAGT  
ATCAGACTGGATTTCAAGGCGAACACCTTTATCCGATCGCAGACCAATAACACGACACGGGACGAC  
GTATATCGGGGATACGAAGATATTCGCGCGCTGGAATAATTTGCTGAGGAAAAGAATCCTAGCGC



- continued

GCCAGACCCCTCGTCCAGATAGCGTCCCGCAAGAAATCCCGCTGTAACCAAGAAAGCGGAAGGGCGCA  
CCCCGGACGCAGAAAGCAGCGAAAGAAGGCCCTCCAGAAGACTCGGAGGACGACATGCAGGCAGAG  
GCTTCTGGAGAAATCTGCGCCCTCCCGAAGACGACGAAGTCCCGAGGACACCGAGCAGATGA  
TCCAACTCGGATCCTGACTATTACAATGACATGCCCGCGTGATCCCGGTGGAGGAGACTACTAAAA  
GTTCTAATGCCGTCTCCATGCCATATTCGCGCGGTTTCGTAGCCTGCGCGGTGCGGCTCGTGGGCTA  
CTGGTTTGGAGCATCGTAAATGCGCGGTAGCTAATCGAGCCTAGAATAGGTGGTTTCTTCTACAT  
GCCACGCCTCAGCTCATAATATAAATCACATGGAATAGCATAACCAATGCCTATTTCATGGGACGTTT  
GAAAGCATGGCATCGTACTTGGAACTCTGGCTCTCCTTCCCGCGAGCTCGCACCCCTTCGGCGCGA  
TGGGAATCGTGATCACTGGAATCACGTCTCCGCCAGGATTGACGACGATCACATCGTGATCGTCGCG  
CCTCGCCCCGAAGCTACAATCAACTGCAGCTATTTTTTCATGCCTGGCCAGAGACCCACAAACCTTA  
CTCAGGAACCGTCCGCGTCCGCTTTCGGTCTGATATAACAACCAAGTGTACCAGGAACTTAGCGAGG  
AGCGCTTTGAAAATGCACTCATCGATCGTCTTCTGTTTTGTGCGGTGTAAGTGACCGAGTACACG  
TTCTCCGCTCGAACAGACTAACCGGACCTCCACACCCGTTAAGCTCACTATACGAAATCCTCGTCC  
GAACGACAGCGGGATGTTCTACGTAATGTTTCGGCTAGACGACACCAAGAACCCATTGACGTCTTCG  
CGATCCAATATCGGTATCAATTGCGAACACCGCGGACTCGCGACTCTATTCCAAGGCTTCG  
TGTCGCACCTTCGGATTACCTACCGTCCAACCTTGAGGCCTATCTCAGGACCGAGGAAAGTTGGCGCAA  
CTGGCAAGCGTACGTTGCCACGAGGCGCACGACACCAGCGCCGAGGCGACAACCCCGACGCCGCTCA  
CTGCAACAGCGCCTCCGAACTTGAAGCGGAACACTTACCTTTCCCTGGCTAGAAAATGGCGTGGAT  
CATTACGAACCGACACCCGAAACGAAAATTCAAAAGTACTGTCCGTCTCGGGACAATGAGCCCTAC  
GCTAATTGGGGTAACCGTGGCTGCCGTGAGCGCAACGATCGGCCCTGTCATTGTAATTTCCATCG  
TCACCAGAAACATGTGCACCCCGACCCGAAAATAGACACGGTCTCGCAAGACGACGAAGAAGCTTCC  
CAAAC TAGAAGGGAATCGGAAAATTTGGACCCATGGTTGCGTGCGAAATAAACAAAGGGGGCTGACCA  
GGATAGTGAACCTGTGGAACCTGGTTGCGATTGTTAACCCGTCTGCGCTAAGCTCGCCGACTCAATAA  
AAATGTGATTAAGTCTGAATGTGGCTCTCCAATCATTTCGATTCTCTAATCTCCAATCCTCTCAAAA  
GGGGCAGTATCGGACACGGACTGGGAGGGCGTACACGATAGTTATATGGTACAGCAGAGGCCTCTGA  
ACACTTAGGAGGAGAATTCAGCCGGGAGAGCCCTGTTGAGTAGGCTTGGGAGCATATTGCAGGATG  
AACATGTTAGTAGATTCTCGCTCTTGTCTTGCAGCCCTAATTTGCGACGCGACACGTCCTCTT  
TTTGAAGGCACTCAGGCTGTCTCGGGGAAGATGATCCAGAAACGTTCCGGAAGGGACTGTAATCA  
AATGGACAAAAGTCTGCGGAACGCGTCAAGATGAAGGCGGCCGATGCTGCTCTTCGCCTAATCTAT  
TGCTTTCATGATTTAATTTACGACGAGGAAAGAAAGACTGCCCGCCCGCGGACCCCTGTCTGCAAA  
CCTGGTAATTTTACTAAAGCGGGCGAAagctt

Dual Expression Cassette Insert

(SEQ ID NO: 17): 5920 bp

gTCGACGGCAGAGTCGACAGCGCCCTATTGGACGTCAAAATGTTAGAGGTGAAGTTTCAAACGATG  
GCGAAGTAACGGCGACTTGCCTTTCCACCGTCAAATCTCCCTATAGGGTAGAACTAATTGGAAAGTA  
GACCTCGTAGATGTAATGGATGAAATTTCTGGGAACAGTCCCGCCGGGTTTTTAAACAGTAATGAGAA  
ATGGCAGAAACAGCTGTACTACAGAGTAACCGATGGAAGAACATCGGTCCAGCTAATGTGCTGTCTGT  
GCACGAGCCATTCTCCGGAACCTTACTGTCTTTTCGACACGCTCTTATAGCGAGGGAAAAAGATATC  
GCGCCAGAGTTATACTTTACCTCTGATCCGCAAACGGCATACTGCACAATAACTCTGCCGTCCGGCGT  
TGTTCCGAGATTCGAATGGAGCCTTAATAATGTTTCACTGCCGGAATATTTGACGGCCACGACCGTTG

- continued

TTTCGCATACCGCTGGCCAAAGTACAGTGTGGAAGAGCAGCGCGAGAGCAGGCGAGGCGTGGATTTCT  
GGCCGGGAGGCAATATATACGAATGCACCGTCTCATCTCAGACGGCACTCGCGTTACTACGCGAAA  
GGAGAGGTGCTTAACAAACACATGGATTGCGGTGGAAAACGGTCTGCTCAGGCGCAGCTGTATTAC  
TCTTTTCTGGACTTGTGTGTCAGGATTATGCGGGAGCATATCTGCTTTGTACGCAACGCTATGGACCGCC  
ATTTATTTTTGAGGAATGCTTTTTGGACTATCTGACTGCTTCTTCCCTCGTAGCCAGAGCACCGCC  
GCCGTCACGTACGACTACATTTTAGGCCGTGCGCGCTCGACGCGCTAACCATACCGCGGTTGGCC  
GTATAACAGATACCTCACTAGGGTATCAAGAGGCTGCGACGTTGTGAGCTCAACCCGATTTCTAACG  
TGGACGACATGATATCGCGGCCAAAGAAAAGAGAAGGGGGCCCTTCGAGGCCCTCCGTGCTGG  
TTCTACGTGATTAAGGGCGACGACGGCGAGGACAAGTACTGTCCAATCTATAGAAAAGAGTACAGGGA  
ATGTGGCGACGTACAACCTGCTATCTGAATGCGCCGTTCAATCTGCACAGATGTGGGCAGTGGACTATG  
TTCTAGCACCCTTGTATCGCGAAATGGCGGGACTGACTATATTCTCCCCACTGCTGCGCTCTCT  
GGCCAATACTTGCTGACCCGAAAATCGGGAGATTGCGCAACAGCTCTCGTAACCTAGAAGTTAA  
CGATCGCTGTTTAAAGATCGGGTTCGACGCTTAACTTTTTACCGTCGAAATGCTGGACAACAGAACAGT  
ATCAGACTGGATTTCAAGGCGAACACCTTTATCCGATCGCAGACCAATACACGACACGCGGACGAC  
GTATATCGGGATACGAAGATATCTGCAGCGTGAATAATTTGCTGAGGAAAAGAATCCTAGCGC  
GCCAGACCCCTCGTCCAGATAGCGTCCCGCAAGAAATCCCGCTGTAACCAAGAAAGCGGAAGGGCGCA  
CCCCGGACGCAGAAAGCAGCGAAAAGAGGCCCTCCAGAAGACTCGGAGGACGACATGCAGGCAGAG  
GCTTCTGGAGAAAATCTGCGCCCTCCCGAAGACGACGAAGTCCCGAGGACACCGAGCACGATGA  
TCCAAACTCGGATCCTGACTATTACAATGACATGCCCGCCGTGATCCCGTGGAGGAGACTACTAAAA  
GTTCTAATGCGCTCTCCATGCCATATTCGCGGCGTTCGTAGCCTGCGCGGTCGCGCTCGTGGGCTA  
CTGGTTTGGAGCATCGTAAAATGCGCGTAGCTAATCGAGCCTAGAATAGGTGTTTCTTCTACAT  
GCCACGCTCAGCTCATAATATAAATCACATGGAATAGCATAACCAATGCCATTTTATTTGGACGTTT  
GAAAAGCATGGCATCGTACTTTGGAATCTGGCTCTCCTTGCCGCGACGCTCGCACCCCTTCGCGCGA  
TGGGAATCGTGATCACTGGAATCACGCTCTCCGCCAGGATTGACGACGATCACATCGTGATCGTCGCG  
CCTCGCCCCGAAGCTACAATTAACCTGCAGCTATTTTTCATGCCTGGCCAGAGACCCACAAACCTA  
CTCAGGAACCGTCCGCGTCCGTTTCGGTCTGATATAACAACAGTGTACTACGGAACCTTAGCGAGG  
AGCGCTTTGAAAATGCACTCATCGATCGTCTTCTGTTTTGTGCGGTGTAAGTGACCGAGTACACG  
TTCTCCGCTCGAACAGACTAACCGACCTCCACACCCGTTTAAAGCTCACTATACGAAATCCTCGTCC  
GAACGACAGCGGATGTTCTACGTAATGTTTCGGCTAGACGACACCAAGAACCCATTGACGTCTTCG  
CGATCCAACCTATCGGTGATCAATTCGCGAACCCCGCGACTCGCGGACTCTATTCCAAGGCTTCG  
TGTCGCACCTTCGGATTACCTACCGTCCAACCTTGAGGCTATCTCAGGACCGAGGAAAGTTGGCGCAA  
CTGGCAAGCGTACGTTGCCACGGAGGCCACGACGACCAGCGCCGAGGCGACAACCCGACGCCGCTCA  
CTGCAACAGCGCTCCGAACTTGAAGCGGAACACTTACCTTTCCCTGGCTAGAAAATGGCGTGGAT  
CATTACGAACCGACACCCGAAACGAAAATTCAAACGTTACTGTCCGCTCTCGGGACAATGAGCCCTAC  
GCTAATGGGGTAACCGTGGCTGCCGTCGTGAGCGCAACGATCGGCCCTCGTCATTGTAATTTCCATCG  
TCACCAGAAACATGTGACCCCGCACCGAAAATAGACACGGTCTCGCAAGACGACGAAGAAGCTTCC  
CAAAC TAGAAGGGAATCGGAAAATTTGGACCCATGGTTGCGTGCGAAATAAACAGGGGGCTGACCA  
GGATAGTGAACCTGTGGAACCTGGTTGCGATGTTAACCCGTCTGCGCTAAGCTCGCCGACTCAATAA  
AAATGTGATTAAGTCTGAATGTGGCTCTCCAATCATTTCGATTCTCTAATCTCCAATCCTCTCAAAA

- continued

GGGGCAGTATCGGACACGGACTGGGAGGGCGGTACACGATAGTTATATGGTACAGCAGAGGCCTCTGA  
ACACTTAGGAGGAGAATTACGCCGGGAGAGCCCTGTTGAGTAGGCTTGGGAGCATATTGCAGGATG  
AACATGTTAGTGATAGTTCTCGCCTCTTGTCTTGC CGCCCTAACTTTTGC GACGCGACACGTCCTCTT  
TTTGGAAAGGCACTCAGGCTGTCTCGGGGAAGATGATCCCAGAAAAGTTCCGGAAAGGACTGTAATCA  
AATGGACAAAAGTCTCGCGAACCGTGAAGATGAAGGCGCCGATGTCTGCTCTTCGCCTAACTAT  
TGCTTTCATGATTTAATTTACGACGGAGGAAAGAAAGACTGCCCGCCGCGGACCCCTGTCTGCAAA  
CCTGGTAATTTTACTAAAAGCGCGGCAAGCTTCGCGCCAGGTCAATTCCTGGCATTATGCCAGTA  
CATGACCTTATGGGACTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGA  
TGCGGTTTTGGCAGTACATCAATGGCGTGGATAGCGGTTTTGACTCACGGGGATTTCCAAGTCTCCAC  
CCCATTGACGTCAATGGGAGTTTGTTTTGGCACCAAAATCAACGGGACTTTCAAAATGTCTGTAACAA  
CTCCGCCCATTTGACGCAAAATGGCGGTAGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTT  
AGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGTTGC  
GCCGCCACCATGGGCCCCAGACCTTCTACCAAGAACCAGTACCTATGATGCTGACTGTCCGAGTCGC  
GCTGGTACTGAGTTGCATCTGTCCGGCAACTCCATTTGATGGCAGGCCTCTTGC GGCTGCAGGAATTG  
TGGTTACAGGAGACAAAGCCGTCAACATATACACCTCATCCCAGACAGGATCAATCATAGTTAAGCTC  
CTCCCGAATCTGCCAAGGATAAGGAGGCATGTGCGAAAGCCCCCTTGGATGCATACAACAGGACATT  
GACCACCTTGTCTACCCCCCTTGGTGACTCTATCCGTAGGATACAAGAGTCTGTGACTACATCTGGAG  
GGGGGAGACAGGGCGCCTTATAGCGCCATTATGGCGGTGTGGCTCTTGGGGTTGCAACTGCCGCA  
CAAATAACAGCGGCCGAGCTCTGATACAAGCCAAAACAAAATGTGCCAACATCCTCCGACTTAAAGA  
GAGCATTGCCGAACCAATGAGGCTGTGCATGAGGTCAGTACCGGATTATCGCAACTAGCAGTGGCAG  
TTGGGAAGATGCAGCAGTTTGTAAATGACCAATTTAATAAAACAGCTCAGGAATTAGACTGCATCAA  
ATTGCACAGCAAGTTGGTGTAGAGCTCAACCTGTACCTAACCGAATTGACTACAGTATTCCGACCACA  
AATCACTTCACTGCTTTAAACAAGCTGACTATTTCAGGCACCTTTACAATCTAGCTGGTGGAAATATGG  
ATTACTTATTGACTAAGTTAGGTGTAGGGAACAATCAACTCAGCTCATTAAATCGGTAGCGGCTTAATC  
ACCGGTAACCTATTCTATACGACTCACAGACTCAACTCTTGGGTATACAGGTAACCTTACCTTCAGT  
CGGAAGCTAAATAATATGCGTGCCACCTACTTGGAAACCTTATCCGTAAGCACAAACAGGGGATTTG  
CCTCGGCACCTGTGCCAAAAGTGTGACACAGGTCGGTTCTGTGATAGAAGAACTTGACACCTCATA  
TGATAGAACTGACTTACATTTATATGTACAAGAAATAGTAACGTTCCCTATGTC CCCTGGTATTTA  
TTCTGCTTGAGCGGCAATACGTCGGCCTGTATGTACTCAAAGACCGAAGGCGCACTTACTACCCAT  
ACATGACTATCAAAGTTTCACTCATCGCCAACTGCAAGATGACAACATGTAGATGTGTAACCCCCG  
GGTATCATATCGAAAACATATGGAGAAGCCGTGTCTAATAGATAAAACAATCATGCAATGTTTTATC  
CTTAGCGGGATAACTTTAAGGCTCAGTGGGGAATTCGATGTAACCTTATCAGAAGAATATCTCAATAC  
AAGATTCTCAAGTAATAATAACAGGCAATCTTGATATCTCAACTGAGCTTGGGAATGTCAACAACCTG  
ATCAGTAATGCTTTGAATAAGTTAGAGGAAAGCAACAGAAAAC TAGACAAAGTCAATGTCAAACCTGAC  
TAGCACATCTGCTCTCATTACCTATATCGTGTGACTATCATATCTCTTGTTTTTGGTATACTTAGCC  
TGATTTAGCATGCTACCTAATGTACAAGCAAAAGGCGCAACAAAAGACCTTATTATGGCTTGGGAAT  
AATACTCTAGATCAGATGAGAGCCACTACAAAATGTGAGGATCTCTCGAGGAATTTAGATCCCAG  
TCACTATTGTATACTCTATATTACTCTATGTTATACTCTGTAATCTACTCAATAAACGTGTCCAG  
CCTGTGAAACCGTACTAAGTCTCCCGTGTCTTCTTATCACCATCAGGTGACATCCTCGCCCAGGCTGT

- continued

CAATCATGCCGGTATCGATTCCAGTAGCACCGGCCCCACGCTGACAACCCACTCTTGACGCGTTAGCA  
 GCGCCCTCTTAAACAAGCCGACCCCCACCAGCGTCGCGGTTACTAACACTCTCTCCCGACCTGCAA  
 CTAGT

**[0150]** The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

**[0151]** It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

---

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1

<211> LENGTH: 1305

<212> TYPE: DNA

<213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 1

atgcaccgctc	ctcatctcag	acggcactcg	cgttactacg	cgaaggaga	ggtgcttaac	60
aaacacatgg	attgcggtgg	aaaacggtgc	tgctcaggcg	cagctgtatt	cactcttttc	120
tggacttgty	tcaggattat	gcgggagcat	atctgctttg	tacgcaacgc	tatggaccgc	180
catttatttt	tgaggaatgc	tttttgact	atcgtaactg	tttcttcctt	cgetagccag	240
agcaccgccg	ccgtcacgta	cgactacatt	ttaggccgctc	gcgcgctcga	cgcgctaacc	300
ataccggcgg	ttggccccta	taacagatac	ctcactaggg	tatcaagagg	ctgcgacggt	360
gtcgagctca	acccgatttc	taacgtggac	gacatgatat	cggcggccaa	agaaaaagag	420
aaggggggccc	ctttcgagge	ctccgtcgtc	tggttctacg	tgattaaggg	cgacgacggc	480
gaggacaagt	actgtccaat	ctatagaaaa	gagtacaggg	aatgtggcga	cgtacaactg	540
ctatctgaat	gcgccgttca	atctgcacag	atgtgggcag	tggactatgt	tcctagcacc	600
cttgatcgc	gaaatggcgc	gggactgact	atattctccc	ccactgctgc	gctctctggc	660
caatacttgc	tgacctgaa	aatcgggaga	tttgcgaaa	cagctctcgt	aactctagaa	720
gttaacgate	gctgtttaa	gatcgggtcg	cagcttaact	ttttaccgctc	gaaatgctgg	780
acaacagaac	agtatcagac	tggatttcaa	ggcgaacacc	tttatccgat	cgcagacacc	840
aatacacgac	acgcggacga	cgtatatcgg	ggatacgaag	atattctgca	gcgctggaat	900
aattttgctga	ggaaaaagaa	tcctagcggc	ccagaccctc	gtccagatag	cgtcccga	960
gaaattccc	ctgtaaccaa	gaaagcggaa	gggcgcacc	cggacgcaga	aagcagcgaa	1020
aagaaggccc	ctccagaaga	ctcggaggac	gacatgcagg	cagaggcttc	tgagaaaaat	1080
ctcgcgcc	tccccgaaga	cgacgaagtc	cccagggaca	ccgagcacga	tgatccaaac	1140
tcggatcctg	actattacaa	tgacatgccc	gccgtgatcc	cggtgaggga	gactactaaa	1200
agttctaagt	ccgtctccat	gcccatatc	gcggcgcttc	tagcctgcgc	ggtcgcgctc	1260
gtggggctac	tggtttggag	catcgtaaaa	tgccgcgcta	gctaa		1305

-continued

---

```

<210> SEQ ID NO 2
<211> LENGTH: 434
<212> TYPE: PRT
<213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 2

Met His Arg Pro His Leu Arg Arg His Ser Arg Tyr Tyr Ala Lys Gly
 1          5          10          15
Glu Val Leu Asn Lys His Met Asp Cys Gly Gly Lys Arg Cys Cys Ser
 20          25          30
Gly Ala Ala Val Phe Thr Leu Phe Trp Thr Cys Val Arg Ile Met Arg
 35          40          45
Glu His Ile Cys Phe Val Arg Asn Ala Met Asp Arg His Leu Phe Leu
 50          55          60
Arg Asn Ala Phe Trp Thr Ile Val Leu Leu Ser Ser Phe Ala Ser Gln
 65          70          75          80
Ser Thr Ala Ala Val Thr Tyr Asp Tyr Ile Leu Gly Arg Arg Ala Leu
 85          90          95
Asp Ala Leu Thr Ile Pro Ala Val Gly Pro Tyr Asn Arg Tyr Leu Thr
100          105          110
Arg Val Ser Arg Gly Cys Asp Val Val Glu Leu Asn Pro Ile Ser Asn
115          120          125
Val Asp Asp Met Ile Ser Ala Ala Lys Glu Lys Glu Lys Gly Gly Pro
130          135          140
Phe Glu Ala Ser Val Val Trp Phe Tyr Val Ile Lys Gly Asp Asp Gly
145          150          155          160
Glu Asp Lys Tyr Cys Pro Ile Tyr Arg Lys Glu Tyr Arg Glu Cys Gly
165          170          175
Asp Val Gln Leu Leu Ser Glu Cys Ala Val Gln Ser Ala Gln Met Trp
180          185          190
Ala Val Asp Tyr Val Pro Ser Thr Leu Val Ser Arg Asn Gly Ala Gly
195          200          205
Leu Thr Ile Phe Ser Pro Thr Ala Ala Leu Ser Gly Gln Tyr Leu Leu
210          215          220
Thr Leu Lys Ile Gly Arg Phe Ala Gln Thr Ala Leu Val Thr Leu Glu
225          230          235          240
Val Asn Asp Arg Cys Leu Lys Ile Gly Ser Gln Leu Asn Phe Leu Pro
245          250          255
Ser Lys Cys Trp Thr Thr Glu Gln Tyr Gln Thr Gly Phe Gln Gly Glu
260          265          270
His Leu Tyr Pro Ile Ala Asp Thr Asn Thr Arg His Ala Asp Asp Val
275          280          285
Tyr Arg Gly Tyr Glu Asp Ile Leu Gln Arg Trp Asn Asn Leu Leu Arg
290          295          300
Lys Lys Asn Pro Ser Ala Pro Asp Pro Arg Pro Asp Ser Val Pro Gln
305          310          315          320
Glu Ile Pro Ala Val Thr Lys Lys Ala Glu Gly Arg Thr Pro Asp Ala
325          330          335
Glu Ser Ser Glu Lys Lys Ala Pro Pro Glu Asp Ser Glu Asp Asp Met
340          345          350
Gln Ala Glu Ala Ser Gly Glu Asn Pro Ala Ala Leu Pro Glu Asp Asp
355          360          365

```

-continued

---

Glu Val Pro Glu Asp Thr Glu His Asp Asp Pro Asn Ser Asp Pro Asp  
 370 375 380

Tyr Tyr Asn Asp Met Pro Ala Val Ile Pro Val Glu Glu Thr Thr Lys  
 385 390 395 400

Ser Ser Asn Ala Val Ser Met Pro Ile Phe Ala Ala Phe Val Ala Cys  
 405 410 415

Ala Val Ala Leu Val Gly Leu Leu Val Trp Ser Ile Val Lys Cys Ala  
 420 425 430

Arg Ser

<210> SEQ ID NO 3  
 <211> LENGTH: 1089  
 <212> TYPE: DNA  
 <213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 3

```

atggcatcgc tacttggaac tctggctctc cttgccgcga cgctcgcacc cttcggcgcg      60
atgggaatcg tgatcactgg aatcacgctc tccgccagga ttgacgacga tcacatcgtg      120
atcgtcgcgc ctgcgcccca agctacaatt caactgcagc tatttttcat gcctggccag      180
agaccccaca aaccctactc aggaaccgctc cgcgtcgcgt ttcggctctga tataacaaac      240
cagtgctacc aggaacttag cgaggagcgc tttgaaaatt gcactcatcg atcgtcttct      300
gtttttgtcg gctgtaaagt gaccgagtac acgttctcgc cctcgaacag actaacggga      360
cctccacacc cgtttaagct cactatacga aatcctcgtc cgaacgacag cgggatgttc      420
tacgtaattg ttcggctaga cgacacccaa gaaccattg acgtcttcgc gatccaacta      480
tcgggtgtatc aattcgcgaa caccgcccgc actcgcggac tctattccaa ggettctgtg      540
cgcaccttcg gattacctac cgtccaactt gaggcctatc tcaggaccga ggaaagtgg      600
cgcaactggc aagcgtactg tgccacggag gccacgacga ccagcgcgga ggcgacaacc      660
ccgacgcccc tcaactgcaac cagcgcctcc gaacttgaag cggaacactt tacctttccc      720
tggttagaaa atggcgtgga tcattacgaa cgcacaccgc caaacgaaaa ttcaaactgt      780
actgtccgctc tcgggacaat gagccctacg ctaattgggg taaccgtggc tgccgtcgtg      840
agcgcaacga tcggcctcgt cattgtaatt tccatcgtca ccagaaacat gtgcaccccg      900
caccgaaaaat tagacacggt ctcgcaagac gacgaagaac gttcccaaac tagaagggaa      960
tcgcgaaaaat ttggaccatc ggttgctgtc gaaataaaca agggggctga ccaggatagt     1020
gaacttgtagg aactggttgc gattgttaac ccgtctgcgc taagctcgcc cgactcaata     1080
aaaatgtga                                         1089
    
```

<210> SEQ ID NO 4  
 <211> LENGTH: 362  
 <212> TYPE: PRT  
 <213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 4

Met Ala Ser Leu Leu Gly Thr Leu Ala Leu Leu Ala Ala Thr Leu Ala  
 1 5 10 15

Pro Phe Gly Ala Met Gly Ile Val Ile Thr Gly Asn His Val Ser Ala  
 20 25 30

Arg Ile Asp Asp Asp His Ile Val Ile Val Ala Pro Arg Pro Glu Ala  
 35 40 45

-continued

---

Thr Ile Gln Leu Gln Leu Phe Phe Met Pro Gly Gln Arg Pro His Lys  
 50 55 60

Pro Tyr Ser Gly Thr Val Arg Val Ala Phe Arg Ser Asp Ile Thr Asn  
 65 70 75 80

Gln Cys Tyr Gln Glu Leu Ser Glu Glu Arg Phe Glu Asn Cys Thr His  
 85 90 95

Arg Ser Ser Ser Val Phe Val Gly Cys Lys Val Thr Glu Tyr Thr Phe  
 100 105 110

Ser Ala Ser Asn Arg Leu Thr Gly Pro Pro His Pro Phe Lys Leu Thr  
 115 120 125

Ile Arg Asn Pro Arg Pro Asn Asp Ser Gly Met Phe Tyr Val Ile Val  
 130 135 140

Arg Leu Asp Asp Thr Lys Glu Pro Ile Asp Val Phe Ala Ile Gln Leu  
 145 150 155 160

Ser Val Tyr Gln Phe Ala Asn Thr Ala Ala Thr Arg Gly Leu Tyr Ser  
 165 170 175

Lys Ala Ser Cys Arg Thr Phe Gly Leu Pro Thr Val Gln Leu Glu Ala  
 180 185 190

Tyr Leu Arg Thr Glu Glu Ser Trp Arg Asn Trp Gln Ala Tyr Val Ala  
 195 200 205

Thr Glu Ala Thr Thr Thr Ser Ala Glu Ala Thr Thr Pro Thr Pro Val  
 210 215 220

Thr Ala Thr Ser Ala Ser Glu Leu Glu Ala Glu His Phe Thr Phe Pro  
 225 230 235 240

Trp Leu Glu Asn Gly Val Asp His Tyr Glu Pro Thr Pro Ala Asn Glu  
 245 250 255

Asn Ser Asn Val Thr Val Arg Leu Gly Thr Met Ser Pro Thr Leu Ile  
 260 265 270

Gly Val Thr Val Ala Ala Val Val Ser Ala Thr Ile Gly Leu Val Ile  
 275 280 285

Val Ile Ser Ile Val Thr Arg Asn Met Cys Thr Pro His Arg Lys Leu  
 290 295 300

Asp Thr Val Ser Gln Asp Asp Glu Glu Arg Ser Gln Thr Arg Arg Glu  
 305 310 315 320

Ser Arg Lys Phe Gly Pro Met Val Ala Cys Glu Ile Asn Lys Gly Ala  
 325 330 335

Asp Gln Asp Ser Glu Leu Val Glu Leu Val Ala Ile Val Asn Pro Ser  
 340 345 350

Ala Leu Ser Ser Pro Asp Ser Ile Lys Met  
 355 360

<210> SEQ ID NO 5  
 <211> LENGTH: 1662  
 <212> TYPE: DNA  
 <213> ORGANISM: Newcastle disease virus

<400> SEQUENCE: 5

atgggccccca gaccttctac caagaaccca gtacctatga tgctgactgt ccgagtccgc 60

ctggtagctga gttgcatctg tccggcaaac tccattgatg gcaggcctct tgcgggtgca 120

ggaattgtgg ttacaggaga caaagccgtc aacatataca cctcatccca gacaggatca 180

atcatagtta agctcctccc gaatctgccc aaggataagg aggcattgtgc gaaagcccc 240

ttggatgcat acaacaggac attgaacct ttgctcacc cccttggtga ctctatccgt 300

-continued

---

```

aggatacaag agtctgtgac tacatctgga ggggggagac aggggcgect tatagcgcc 360
attattggcg gtgtggctct tggggttgca actgccgcac aaataacagc ggcgcagct 420
ctgatacaag ccaaacaaaa tgctgccaac atcctccgac ttaaagagag cattgcccga 480
accaatgagg ctgtgcata ggtcactgac ggattatcgc aactagcagt ggcagtggg 540
aagatgcagc agtttgtaa tgaccaattt aataaacag ctcaggaatt agactgcatc 600
aaaaattgac agcaagttgg tgtagagctc aacctgtacc taaccgaatt gactacagta 660
ttcggaccac aatcacttc acctgcttta aacaagctga ctattcaggc actttacaat 720
ctagctggty gaaatatgga ttacttattg actaagttag gtgtaggaa caatcaactc 780
agctcattaa tcggtagcgg cttaatcacc ggtaacceta ttctatacga ctcacagact 840
caactcttgg gtatacaggt aactctacct tcagtcggga agctaaataa tatgcgtgcc 900
acctacttgg aaaccttate cgtaagcaca accaggggat ttgcctcggc acttgtccca 960
aaagtgggta cacaggtcgg ttctgtgata gaagaacttg acctcata ctgtatagaa 1020
actgacttac atttatattg tacaagaata gtaacgttcc ctatgtcccc tggtatattat 1080
tcctgcttga ggggcaatac gtcggcctgt atgtactcaa agaccgaagg cgcacttact 1140
acaccataca tgactatcaa aggttcagtc atcgccaact gcaagatgac aacatgtaga 1200
tgtgtaaacc ccccggtat catatcgcaa aactatggag aagccgtgtc tctaatagat 1260
aaacaatcat gcaatgtttt atccttaggc gggataactt taaggctcag tggggaattc 1320
gatgtaactt atcagaagaa tatctcaata caagattctc aagtaataat aacaggcaat 1380
cttgatatct caactgagct tgggaatgtc aacaactcga tcagtaatgc tttgaataag 1440
ttagaggaaa gcaacagaaa actagacaaa gtcaatgtca aactgactag cacatctgct 1500
ctcattacct atatcgtgtt gactatcata tctctgttt ttggtatact tagcctgatt 1560
ctagcatgct acctaatagta caagcaaaa ggcgaacaaa agaccttatt atggcttggg 1620
aataaactc tagatcagat gagagccact acaaaaatgt ga 1662

```

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 553

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Newcastle disease virus

&lt;400&gt; SEQUENCE: 6

```

Met Gly Pro Arg Pro Ser Thr Lys Asn Pro Val Pro Met Met Leu Thr
1           5           10           15
Val Arg Val Ala Leu Val Leu Ser Cys Ile Cys Pro Ala Asn Ser Ile
20          25          30
Asp Gly Arg Pro Leu Ala Ala Ala Gly Ile Val Val Thr Gly Asp Lys
35          40          45
Ala Val Asn Ile Tyr Thr Ser Ser Gln Thr Gly Ser Ile Ile Val Lys
50          55          60
Leu Leu Pro Asn Leu Pro Lys Asp Lys Glu Ala Cys Ala Lys Ala Pro
65          70          75          80
Leu Asp Ala Tyr Asn Arg Thr Leu Thr Thr Leu Leu Thr Pro Leu Gly
85          90          95
Asp Ser Ile Arg Arg Ile Gln Glu Ser Val Thr Thr Ser Gly Gly Gly
100         105         110
Arg Gln Gly Arg Leu Ile Gly Ala Ile Ile Gly Gly Val Ala Leu Gly

```



-continued

115					120					125						
Val	Ala	Thr	Ala	Ala	Gln	Ile	Thr	Ala	Ala	Ala	Ala	Ala	Leu	Ile	Gln	Ala
130					135					140						
Lys	Gln	Asn	Ala	Ala	Asn	Ile	Leu	Arg	Leu	Lys	Glu	Ser	Ile	Ala	Ala	
145					150					155					160	
Thr	Asn	Glu	Ala	Val	His	Glu	Val	Thr	Asp	Gly	Leu	Ser	Gln	Leu	Ala	
					165					170					175	
Val	Ala	Val	Gly	Lys	Met	Gln	Gln	Phe	Val	Asn	Asp	Gln	Phe	Asn	Lys	
					180					185					190	
Thr	Ala	Gln	Glu	Leu	Asp	Cys	Ile	Lys	Ile	Ala	Gln	Gln	Val	Gly	Val	
					195					200					205	
Glu	Leu	Asn	Leu	Tyr	Leu	Thr	Glu	Leu	Thr	Thr	Val	Phe	Gly	Pro	Gln	
					210					215					220	
Ile	Thr	Ser	Pro	Ala	Leu	Asn	Lys	Leu	Thr	Ile	Gln	Ala	Leu	Tyr	Asn	
					225					230					235	
Leu	Ala	Gly	Gly	Asn	Met	Asp	Tyr	Leu	Leu	Thr	Lys	Leu	Gly	Val	Gly	
					245					250					255	
Asn	Asn	Gln	Leu	Ser	Ser	Leu	Ile	Gly	Ser	Gly	Leu	Ile	Thr	Gly	Asn	
					260					265					270	
Pro	Ile	Leu	Tyr	Asp	Ser	Gln	Thr	Gln	Leu	Leu	Gly	Ile	Gln	Val	Thr	
					275					280					285	
Leu	Pro	Ser	Val	Gly	Lys	Leu	Asn	Asn	Met	Arg	Ala	Thr	Tyr	Leu	Glu	
					290					295					300	
Thr	Leu	Ser	Val	Ser	Thr	Thr	Arg	Gly	Phe	Ala	Ser	Ala	Leu	Val	Pro	
					305					310					315	
Lys	Val	Val	Thr	Gln	Val	Gly	Ser	Val	Ile	Glu	Glu	Leu	Asp	Thr	Ser	
					325					330					335	
Tyr	Cys	Ile	Glu	Thr	Asp	Leu	His	Leu	Tyr	Cys	Thr	Arg	Ile	Val	Thr	
					340					345					350	
Phe	Pro	Met	Ser	Pro	Gly	Ile	Tyr	Ser	Cys	Leu	Ser	Gly	Asn	Thr	Ser	
					355					360					365	
Ala	Cys	Met	Tyr	Ser	Lys	Thr	Glu	Gly	Ala	Leu	Thr	Thr	Pro	Tyr	Met	
					370					375					380	
Thr	Ile	Lys	Gly	Ser	Val	Ile	Ala	Asn	Cys	Lys	Met	Thr	Thr	Cys	Arg	
					385					390					395	
Cys	Val	Asn	Pro	Pro	Gly	Ile	Ile	Ser	Gln	Asn	Tyr	Gly	Glu	Ala	Val	
					405					410					415	
Ser	Leu	Ile	Asp	Lys	Gln	Ser	Cys	Asn	Val	Leu	Ser	Leu	Gly	Gly	Ile	
					420					425					430	
Thr	Leu	Arg	Leu	Ser	Gly	Glu	Phe	Asp	Val	Thr	Tyr	Gln	Lys	Asn	Ile	
					435					440					445	
Ser	Ile	Gln	Asp	Ser	Gln	Val	Ile	Ile	Thr	Gly	Asn	Leu	Asp	Ile	Ser	
					450					455					460	
Thr	Glu	Leu	Gly	Asn	Val	Asn	Asn	Ser	Ile	Ser	Asn	Ala	Leu	Asn	Lys	
					465					470					475	
Leu	Glu	Glu	Ser	Asn	Arg	Lys	Leu	Asp	Lys	Val	Asn	Val	Lys	Leu	Thr	
					485					490					495	
Ser	Thr	Ser	Ala	Leu	Ile	Thr	Tyr	Ile	Val	Leu	Thr	Ile	Ile	Ser	Leu	
					500					505					510	
Val	Phe	Gly	Ile	Leu	Ser	Leu	Ile	Leu	Ala	Cys	Tyr	Leu	Met	Tyr	Lys	
					515					520					525	

-continued

---

Gln Lys Ala Gln Gln Lys Thr Leu Leu Trp Leu Gly Asn Asn Thr Leu  
530 535 540

Asp Gln Met Arg Ala Thr Thr Lys Met  
545 550

<210> SEQ ID NO 7  
<211> LENGTH: 1698  
<212> TYPE: DNA  
<213> ORGANISM: Newcastle disease virus

<400> SEQUENCE: 7

```

atggatcgat cccggttggc gccctccagg tgcaggatgg gctccagacc ttetaccaag    60
aaccacgac ctatgatgct gactatccgg gtcgcgctgg tactgagttg catctgtccc    120
gcaaactcca ttgatggcag gcctcttgca gctgcaggaa ttgtggttac aggagacaaa    180
gcagtcaaca tatacacctc atcccagaca ggatcaatca tagttaagct cctcccgaat    240
ctgcaaaagg ataaggagcg atgtgcgaaa gcccccttgg atgcatacaa caggacattg    300
accactttgc tcacccccct tgggtactct atccgtagga tacaagagtc tgtgactaca    360
tctggagggg ggagacaggg gcgccttata ggcgccatta ttggcgggtg ggctcttggg    420
gttgcaactg ccgcacaaat aacagcggcc gcagctctga tacaagccaa acaaaatgct    480
gccaacatcc tccgacttaa agagagcatt gccgcaacca atgaggctgt gcatgaggtc    540
actgacggat tatcgcaact agcagtgcca gttgggaaga tgcagcagtt cgttaatgac    600
caatttaata aaacagctca ggaatagac tgcatacaaaa tgcacagca agttggtgta    660
gagctcaacc tgtacctaac cgaatcgact acagtattcg gaccacaaat cacttcacct    720
gccttaaaca agctgactat tcaggcactt tacaatctag ctggtgggaa tatggattac    780
ttattgacta agttaggtat agggaacaat caactcagct cattaatcgg tagcggctta    840
atcaccggta accctattct atacgactca cagactcaac tcttgggtat acaggtaact    900
ctaccttcag tcgggaacct aaataatag cgtgccacct acttggaaac cttatccgta    960
agcacaacca ggggatattgc ctcggcactt gtcccaaaag tggtgacacg ggtcggttct   1020
gtgatagaag aacttgacac ctcatactgt atagaaactg acttagattt atattgtaca   1080
agaatagtaa cgttccctat gtcccctggg atttactcct gcttgagcgg caatacatcg   1140
gcctgtatgt actcaaagac cgaaggcgca cttactacac catatatgac tatcaaaggc   1200
tcagtcacg ctaactgcaa gatgacaaca tgtagatgtg taaaccccc gggtatcata   1260
tcgcaaaact atggagaagc cgtgtctcta atagataaac aatcatgcaa tgttttatcc   1320
ttaggcggga taactttaag gctcagtggg gaattcgatg taacttatca gaagaatc   1380
tcaatacaag attctcaagt aataataaca ggcaatcttg atatctcaac tgagcttggg   1440
aatgtcaaca actcgatcag taatgccttg aataagttag aggaaagcaa cagaaaacta   1500
gacaaagtca atgtcaaact gaccagcaca tctgctctca ttacctatat cgttttgact   1560
atcatatctc ttgtttttgg tatacttagc ctgattctag catgctacct aatgtacaag   1620
caaaaggcgc aacaaaagac cttattatgg cttgggaata atacccctaga tcagatgaga   1680
gccactacaa aaatgtga                                     1698

```

<210> SEQ ID NO 8  
<211> LENGTH: 565  
<212> TYPE: PRT  
<213> ORGANISM: Newcastle disease virus

-continued

&lt;400&gt; SEQUENCE: 8

Met Asp Arg Ser Arg Leu Ala Pro Ser Arg Cys Arg Met Gly Ser Arg  
1 5 10 15  
Pro Ser Thr Lys Asn Pro Ala Pro Met Met Leu Thr Ile Arg Val Ala  
20 25 30  
Leu Val Leu Ser Cys Ile Cys Pro Ala Asn Ser Ile Asp Gly Arg Pro  
35 40 45  
Leu Ala Ala Ala Gly Ile Val Val Thr Gly Asp Lys Ala Val Asn Ile  
50 55 60  
Tyr Thr Ser Ser Gln Thr Gly Ser Ile Ile Val Lys Leu Leu Pro Asn  
65 70 75 80  
Leu Pro Lys Asp Lys Glu Ala Cys Ala Lys Ala Pro Leu Asp Ala Tyr  
85 90 95  
Asn Arg Thr Leu Thr Thr Leu Leu Thr Pro Leu Gly Asp Ser Ile Arg  
100 105 110  
Arg Ile Gln Glu Ser Val Thr Thr Ser Gly Gly Gly Arg Gln Gly Arg  
115 120 125  
Leu Ile Gly Ala Ile Ile Gly Gly Val Ala Leu Gly Val Ala Thr Ala  
130 135 140  
Ala Gln Ile Thr Ala Ala Ala Ala Leu Ile Gln Ala Lys Gln Asn Ala  
145 150 155 160  
Ala Asn Ile Leu Arg Leu Lys Glu Ser Ile Ala Ala Thr Asn Glu Ala  
165 170 175  
Val His Glu Val Thr Asp Gly Leu Ser Gln Leu Ala Val Ala Val Gly  
180 185 190  
Lys Met Gln Gln Phe Val Asn Asp Gln Phe Asn Lys Thr Ala Gln Glu  
195 200 205  
Leu Asp Cys Ile Lys Ile Ala Gln Gln Val Gly Val Glu Leu Asn Leu  
210 215 220  
Tyr Leu Thr Glu Ser Thr Thr Val Phe Gly Pro Gln Ile Thr Ser Pro  
225 230 235 240  
Ala Leu Asn Lys Leu Thr Ile Gln Ala Leu Tyr Asn Leu Ala Gly Gly  
245 250 255  
Asn Met Asp Tyr Leu Leu Thr Lys Leu Gly Ile Gly Asn Asn Gln Leu  
260 265 270  
Ser Ser Leu Ile Gly Ser Gly Leu Ile Thr Gly Asn Pro Ile Leu Tyr  
275 280 285  
Asp Ser Gln Thr Gln Leu Leu Gly Ile Gln Val Thr Leu Pro Ser Val  
290 295 300  
Gly Asn Leu Asn Asn Met Arg Ala Thr Tyr Leu Glu Thr Leu Ser Val  
305 310 315 320  
Ser Thr Thr Arg Gly Phe Ala Ser Ala Leu Val Pro Lys Val Val Thr  
325 330 335  
Arg Val Gly Ser Val Ile Glu Glu Leu Asp Thr Ser Tyr Cys Ile Glu  
340 345 350  
Thr Asp Leu Asp Leu Tyr Cys Thr Arg Ile Val Thr Phe Pro Met Ser  
355 360 365  
Pro Gly Ile Tyr Ser Cys Leu Ser Gly Asn Thr Ser Ala Cys Met Tyr  
370 375 380  
Ser Lys Thr Glu Gly Ala Leu Thr Thr Pro Tyr Met Thr Ile Lys Gly  
385 390 395 400

-continued

---

Ser Val Ile Ala Asn Cys Lys Met Thr Thr Cys Arg Cys Val Asn Pro  
405 410 415

Pro Gly Ile Ile Ser Gln Asn Tyr Gly Glu Ala Val Ser Leu Ile Asp  
420 425 430

Lys Gln Ser Cys Asn Val Leu Ser Leu Gly Gly Ile Thr Leu Arg Leu  
435 440 445

Ser Gly Glu Phe Asp Val Thr Tyr Gln Lys Asn Ile Ser Ile Gln Asp  
450 455 460

Ser Gln Val Ile Ile Thr Gly Asn Leu Asp Ile Ser Thr Glu Leu Gly  
465 470 475 480

Asn Val Asn Asn Ser Ile Ser Asn Ala Leu Asn Lys Leu Glu Glu Ser  
485 490 495

Asn Arg Lys Leu Asp Lys Val Asn Val Lys Leu Thr Ser Thr Ser Ala  
500 505 510

Leu Ile Thr Tyr Ile Val Leu Thr Ile Ile Ser Leu Val Phe Gly Ile  
515 520 525

Leu Ser Leu Ile Leu Ala Cys Tyr Leu Met Tyr Lys Gln Lys Ala Gln  
530 535 540

Gln Lys Thr Leu Leu Trp Leu Gly Asn Asn Thr Leu Asp Gln Met Arg  
545 550 555 560

Ala Thr Thr Lys Met  
565

<210> SEQ ID NO 9  
<211> LENGTH: 527  
<212> TYPE: DNA  
<213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 9

```

aaacagctgt actacagagt aaccgatgga agaacatcgg tccagctaata gtgcctgtcg      60
tgcacgagcc attctccgga accttactgt cttttcgaca cgtctcttat agcgagggaa      120
aaagatatcg cgccagagtt atactttacc tctgatccgc aaacggcata ctgcacaata      180
actctgacct cggcgctgtt tccgagatc gaatggagcc ttaataatgt ttcactgccg      240
gaatatttga cggccaacgac cgttgtttcg cataaccgctg gccaaagtac agtgtggaag      300
agcagcgcgga gacgagcgga ggcgtggatt tctggccggg gaggcaatat atacgaatgc      360
accgtcctca tctcagacgg cactcgcggt actacgcgaa aggagagggt cttaacaaac      420
acatggattg cggtggaaaa cgggtctgct caggcgcagc tgtattcact cttttctgga      480
cttgtgtcag gattatgcgg gagcatatct gctttgtacg caacgct                    527

```

<210> SEQ ID NO 10  
<211> LENGTH: 264  
<212> TYPE: DNA  
<213> ORGANISM: Infectious laryngotracheitis virus

<400> SEQUENCE: 10

```

tgactattac aatgacatgc ccgccgtgat cccggtggag gagactacta aaagttctaa      60
tgccgtctcc atgcccatat tcgcggcggt cgtagcctgc gcggtcgcgc tcgtggggct      120
actggtttgg agcatcgtaa aatgcgcgcg tagctaactc agcctagaat aggtggtttc      180
ttcctacatg ccacgcctca cgctcataat ataaatcaca tggaatagca taccaatgcc      240
tattcattgg gacgttcgaa aagc                                          264

```

-continued

---

```

<210> SEQ ID NO 11
<211> LENGTH: 360
<212> TYPE: DNA
<213> ORGANISM: Human cytomegalovirus

<400> SEQUENCE: 11
cgcgccaggt caattccctg gcattatgcc cagtacatga ccttatggga ctttcctact    60
tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcggtt ttggcagtac    120
atcaatgggc gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac    180
gtcaatggga gtttgttttg gcacaaaaat caacgggact ttccaaaatg tcgtaacaac    240
tccgccccat tgacgcaaat gggcggtagc gtgtacggtg ggaggctat ataagcagag    300
ctcgtttagt gaaccgtcag atcgctgga gacgccatcc acgctgtttt gacctccata    360

```

```

<210> SEQ ID NO 12
<211> LENGTH: 1191
<212> TYPE: DNA
<213> ORGANISM: Human cytomegalovirus

<400> SEQUENCE: 12
gtgaataata aaatgtgtgt ttgtccgaaa tacgcgtttg agatttctgt cccgactaaa    60
ttcatgtcgc gcgatagtgg tgtttatcgc cgatagagat ggcgatattg gaaaaatcga    120
tatttgaaaa tatggcatat tgaaaatgtc gccgatgtga gtttctgtgt aactgatatc    180
gccatttttc caaaagttag tttttgggca taccgatat ctggcgatag gcttatatcg    240
tttacggggg atggcgatag acgccttttg tgacttgggc gattctgtgt gtcgcaata    300
tcgcagtttc gatataggtg acagacgata tgaggctata tcgccgatag aggcgacatc    360
aagctggcac atggccaatg catatcgatc tatacattga atcaatattg gccattagcc    420
atattattca ttggttatat agcataaatc aatattggct attggccatt gcatacgttg    480
tatocatatc ataatatgta catttatatt ggctcatgtc caacattacc gccatgttga    540
cattgattat tgactagtta ttaatagtaa tcaattacgg ggtcattagt tcatagccca    600
tatatggagt tccgcgttac ataacttacg gtaaatggcc cgcttggtg accgcccac    660
gacccccgcc cattgacgtc aataatgacg tatgttccca tagtaacgcc aatagggact    720
ttccattgac gtcaatgggt ggagtattta cggtaaacctg cccacttggc agtacatcaa    780
gtgtatcata tgccaagtac gccccctatt gacgtcaatg acggtaaatg gcccgctgg    840
cattatgccc agtacatgac cttatgggac tttcctactt ggcagtacat ctacgtatta    900
gtcatcgcta ttacatggt gatgctggtt tggcagtaca tcaatgggag tggatagcgg    960
tttgactcac ggggatttcc aagtctccac cccattgacg tcaatgggag tttgttttgg    1020
caccaaaatc aacgggactt tccaaaatgt cgtaacaact cgcgccatt gacgcaaatg    1080
ggcggttagc gtgtacggtg ggaggctat ataagcagag ctcgtttagt gaaccgtcag    1140
atcgctgga gacgccatcc acgctgtttt gacctccata gaagacaccg g          1191

```

```

<210> SEQ ID NO 13
<211> LENGTH: 281
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence

```

-continued

&lt;400&gt; SEQUENCE: 13

```

ggaattctag atcccacgtc actattgtat actctatatt atactctatg ttatactctg    60
taatcctact caataaacgt gtcacgctg tgaaaccgta ctaagtctcc cgtgtcttct    120
tatcaccatc aggtgacatc ctgcccagg ctgtcaatca tgccggtatc gattccagta    180
gcaccggccc cacgctgaca acccaactct gcagcgtag cagcgcccct cttaacaagc    240
cgacccccac cagcgtcgcg gttactaaca ctctctccc c                    281

```

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 523

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: herpes simplex virus

&lt;400&gt; SEQUENCE: 14

```

gggagatggg ggaggctaac tgaaacacgg aaggagacaa taccggaagg aaccgcgct    60
atgacggcaa taaaagaca gaataaacg cacgggtgtt gggtcgtttg ttcataaacg    120
cggggttcgg tcccagggtt ggcaactctg cgatacccca ccgagacccc attgggacca    180
atacgccegc gtttcttct tttccccacc ccaaccccca agttcgggtg aaggcccagg    240
gctcgcagcc aacgtcgggg cggcaagccc tgccatagcc acgggcccgc tgggttaggg    300
acggggtccc ccatgggaa tggtttatgg ttcgtggggg ttattatttt gggcgttgcg    360
tggggtcagg tccacgactg gactgagcag acagacccat ggtttttgga tggcctgggc    420
atggaccgca tgtactggcg cgacaagaac accgggcgtc tgtggctgcc aaacaccccc    480
gacccccaaa aaccaccgcg cggatttctg gcgcccgcgg acg                    523

```

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 3593

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Expression cassette insert

&lt;400&gt; SEQUENCE: 15

```

taattaacc gggaagcttg catgcctgca gtgaataata aaatgtgtgt ttgtccgaaa    60
tacgcgtttg agatttctgt cccgactaaa ttcattgtgc gcgatagtgg tgtttatcgc    120
cgatagagat ggcgatattg gaaaaatcga tatttgaaaa tatggcatat tgaaaatgtc    180
gccgatgtga gtttctgtgt aactgatac gccatttttc caaaagtga tttttgggca    240
tacgcgatat ctggcgatac gtttatatcg tttacggggg atggcgatag acgcctttgg    300
tgacttgggc gattctgtgt gtcgcaaata tcgcagtttc gatatagggt acagacgata    360
tgaggctata tcgccgatag aggcgacatc aagctggcac atggccaatg catatcgatc    420
tatacattga atcaatattg gccattagcc atattattca ttggttatat agcataaatc    480
aatattggct atggccatt gcatacgttg tatccatc ataatatgta catttatatt    540
ggctcatgtc caacattacc gccatgttga cattgattat tgactagtta ttaatagtaa    600
tcaattacgg ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg    660
gtaaattggc cgcttgctg accgcccac gacccccgcc cattgacgtc aataatgacg    720
tatgttccca tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta    780
cggtaaaactg cccacttggc agtacatcaa gtgtatcata tgccaagtac gccccctatt    840
gacgtcaatg acggtaaatg gcccgcctgg cattatgccc agtacatgac cttatgggac    900

```

-continued

---

tttctactt ggcagtacat ctacgtatta gtcacgcta ttaccatggt gatgcggttt	960
tggcagtaca tcaatgggcy tggatagcgg tttgactcac ggggatttcc aagtctccac	1020
cccattgacg tcaatgggag tttgttttgg caccaaaatc aacgggactt tccaaaatgt	1080
cgtaacaact ccgccccatt gacgcaaatg ggcggtaggc gtgtacggtg ggaggtctat	1140
ataagcagag ctcgtttagt gaaccgtcag atcgctgga gacgccatcc acgctgtttt	1200
gacctcata gaagacaccg ggaccatgga tcgatcccg ttggcgccct ccaggtgcag	1260
gatgggtcc agaccttcta ccaagaacct agcacctatg atgctgacta tccgggtcgc	1320
gctggtagt agttgcact gtccggcaaa ctccattgat ggcaggctc ttgcagctgc	1380
aggaattgtg gttacaggag acaaagcagt caacatatac acctcatccc agacaggatc	1440
aatcatagtt aagctcctcc cgaatctgcc aaaggataag gaggcattgtg cgaagcccc	1500
cttggatgca tacaacagga cattgaccac tttgctcacc ccccttggtg actctatccg	1560
taggatacaa gagtctgtga ctacatctgg aggggggaga caggggcgcc ttataggcgc	1620
cattattggc ggtgtggctc ttgggggttc aactgccgca caaataacag cggccgcagc	1680
tctgatacaa gccaaaacaa atgctgcca catcctccga cttaagaga gcattgccgc	1740
aaccaatgag gctgtgcatg aggtcactga cggattatcg caactagcag tggcagttgg	1800
gaagatgcag cagttcgtta atgaccaatt taataaaaca gctcaggaat tagactgcat	1860
caaaattgca cagcaagttg gtgtagagct caacctgtac ctaaccgaat cgactacagt	1920
attcggacca caaatcactt cacctgcctt aaacaagctg actattcagg cactttacaa	1980
tctagctggt gggaatatgg attacttatt gactaagtta ggtataggga acaatcaact	2040
cagctcatta atcggtagcg gcttaatcac cggtaaccct attctatagc actcacagac	2100
tcaactcttg ggtatacagg taactctacc ttcagtcggg aacctaaata atatgctggc	2160
cacctacttg gaaaccttat ccgtaagcac aaccagggga tttgcctcgg cacttgtccc	2220
aaaagtggg acacgggtcg gttctgtgat agaagaactt gacacctcat actgtataga	2280
aactgactta gatttatatt gtacaagaat agtaacgttc cctatgtccc ctggatttta	2340
ctctgcttg agcggcaata catcggcctg tatgtactca aagaccgaag gcgcacttac	2400
tacaccatat atgactatca aaggctcagt catcgctaac tgcaagatga caacatgtag	2460
atgtgtaaac cccccgggta tcatacgcga aaactatgga gaagccgtgt ctctaataga	2520
taaacatca tgcaatggtt tacccttagg cgggataact ttaaggctca gtggggaatt	2580
cgatgtaact tatcagaaga atatctcaat acaagattct caagtaataa taacaggcaa	2640
tcttgatate tcaactgagc ttgggaatgt caacaactcg atcagtaatg ccttgaataa	2700
gtagagggaa agcaacagaa aactagacaa agtcaatgac aaactgacca gcacatctgc	2760
tctcattacc tatatcggtt tgactatcat atctctgtt tttggatac ttagcctgat	2820
tctagcatgc tacctaagt acaagcaaaa ggcgcaacaa aagacctat tatggcttgg	2880
gaataatacc ctagatcaga tgagagccac tacaacaaatg tgaacacaga tgaggaacga	2940
aggtttccct aatagtaatt tgtgtgaaag ttctggtagt ctgtcagttc ggagagttaa	3000
gaaaaaaaaa aaaccccccc ccccccccc cccccccct gggtagcagc ctctagagtc	3060
gggagatggg ggaggctaac tgaacacagc aaggagacaa taccggaagg aacccgcgct	3120
atgacggcaa taaaagaca gaataaacg cacgggtgtt gggtcgtttg ttcataaacg	3180

-continued

---

cggggttcgg	tcccagggct	ggcactctgt	cgatacccca	ccgagacccc	attgggacca	3240
atacgcccgc	gtttcttctt	tttcccacc	ccaaccccca	agttcgggtg	aaggcccagg	3300
gctcgcagcc	aacgtcgggg	cggcaagccc	tgccatagcc	acgggcccgc	tgggttaggg	3360
acggggtccc	ccatggggaa	tggtttatgg	ttcgtggggg	ttattatfff	gggcggtgcg	3420
tggggtcagg	tccacgactg	gactgagcag	acagacccat	ggtttttggg	tggcctgggc	3480
atggaccgca	tgtactggcg	cgacacgaac	accgggcgtc	tgtggctgcc	aaacaccccc	3540
gacccccaaa	aaccaccgcg	cggatttctg	gcgccgcggg	acgtcgcactt	aat	3593

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 3569

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Expression cassette insert

&lt;400&gt; SEQUENCE: 16

gtcgcaggca	gagtcgcaga	cgcccctatt	ggacgtcaaa	attgtagagg	tgaagttttc	60
aaacgatggc	gaagtaacgg	cgacttgctg	ttccaccgtc	aaatctccct	atagggtaga	120
aactaattgg	aaagtagacc	tcgtagatgt	aatggatgaa	atttctggga	acagtcccgc	180
cggggttttt	aacagtaatg	agaaatggca	gaaacagctg	tactacagag	taaccgatgg	240
agaacatcgc	gtccagctaa	tgtgcctgtc	gtgcacgagc	cattctccgg	aaccttactg	300
tcttttcgac	acgtctctta	tagcagggga	aaaagatata	gcccagaggt	tatactttac	360
ctctgatccg	caaacggcat	actgcacaat	aactctgccg	tccggcggtg	ttccgagatt	420
cgaatggagc	cttaataatg	tttcaactgcc	ggaatatttg	acggccacga	ccgttgtttc	480
gcataccgct	ggccaaagta	cagtgtagaa	gagcagcgcg	agagcaggcg	aggcgtggat	540
ttctggccgg	ggaggcaata	tatacgaatg	caccgtcctc	atctcagacg	gcactcgcgt	600
tactacgcga	aaggagaggt	gcttaacaaa	cacatggatt	gcggtggaaa	acgggtgctgc	660
tcaggcgcag	ctgtattcac	tcttttctgg	acttggtgca	ggattatgcg	ggagcatatc	720
tgctttgtac	gcaacgctat	ggaccgccat	ttatftttga	ggaatgcttt	ttggactatc	780
gtactgcttt	cttctctcgc	tagccagagc	accgccgcgg	tcacgtacga	ctacatttta	840
ggcgcgcgcg	cgctcgcagc	gctaaccata	ccggcgggtg	gcccgtataa	cagatacttc	900
actaggggat	caagaggctg	cgacgttgct	gagctcaacc	cgatttctaa	cgtggacgac	960
atgatatcgg	cggccaaaga	aaaagagaag	gggggcccct	tcgaggcctc	cgtcgtctgg	1020
ttctacgtga	ttaagggcga	cgacggcgag	gacaagtact	gtccaatcta	tagaaaagag	1080
tacaggggat	gtggcgcagc	acaactgcta	tctgaatgcg	ccgttcaatc	tgcacagatg	1140
tgggcagtg	actatgttcc	tagcaccctt	gtatcgcgaa	atggcgcggg	actgactata	1200
ttctccccca	ctgctcgcgt	ctctggccaa	tacttgctga	ccctgaaaat	cgggagattt	1260
gcgcaaacag	ctctcgtaac	tctagaagtt	aacgatcgtc	gtttaaagat	cgggtcgcag	1320
cttaactttt	taccgtcgaa	atgctggaca	acagaacagt	atcagactgg	atttcaaggc	1380
gaacaccttt	atccgatcgc	agacaccaat	acacgacacg	cggacgacgt	atatcgggga	1440
tacgaagata	ttctgcagcg	ctggaataat	ttgctgagga	aaaagaatcc	tagcgcgcca	1500
gaccctcgtc	cagatagcgt	cccgaagaa	attcccgtcg	taaccaagaa	agcgggaaggg	1560
cgcaccccgg	acgcagaaa	cagcgaaaa	aaggcccctc	cagaagactc	ggaggacgac	1620



-continued

---

```

atgcaggcag aggcttctgg agaaaatcct gccgccctcc ccgaagacga cgaagtcccc 1680
gaggacaccg agcacgatga tccaaactcg gatcctgact attacaatga catgcccgcc 1740
gtgatcccg tggaggagac tactaaaagt tctaatgccg tctccatgcc catattcgcg 1800
gcgtctctag cctgcgcggt cgcgctctgt gggctactgg tttggagcat cgtaaaatgc 1860
gcgcgtagct aatcgcgcct agaataggtg gtttcttctc acatgccacg cctcacgctc 1920
ataatataaa tcacatggaa tagcatacca atgcctattc attgggacgt tcgaaaagca 1980
tggeatcgct acttggaact ctggctctcc ttgccgcgac gctcgcaacc ttcggcgcgga 2040
tgggaatcgt gatcactgga aatcacgtct ccgccaggat tgacgacgat cacatcgtga 2100
tcgtcgcgcc tcgccccgaa gctacaattc aactgcagct atttttcatg cctggccaga 2160
gacccccaaa accctactca ggaacgctcc gcgtcgctt tcggtctgat ataacaacc 2220
agtgtacca ggaacttagc gaggagcgtc ttgaaaattg cactcatcga tcgtcttctg 2280
ttttgtcgg ctgtaaagt accgagtaca cgttctccgc ctcgaaacaga ctaaccggac 2340
ctccacacc gtttaagtc actatacga atcctcgtcc gaacgacagc gggatgttct 2400
acgtaattgt tcggctagac gacaccaaag aaccattga cgtcttcgcg atccaactat 2460
cgggtgatca attcgcgaac accgcgcgga ctgcgggact ctattccaag gcttcgtgtc 2520
gcaccttcgg attacctacc gtccaacttg aggcctatct caggaccgag gaaagtggc 2580
gcaactggca agcgtacgtt gccacggagg ccacgacgac cagcgcgag gcgacaacc 2640
cgacgcccgt cactgcaacc agcgcctccg aactgaagc ggaacacttt acctttccct 2700
ggctagaaaa tggcgtggat cattaacgaac cgacaccgc aaacgaaaat tcaaacgta 2760
ctgtccgtct cgggacaatg agccctacgc taattggggg aaccgtggct gccgtcgtga 2820
gcgcaacgat cggcctcgtc attgtaattt ccacgtcac cagaacatg tgcaccccgc 2880
accgaaaatt agacacggtc tcgcaagacg acgaagaacg tccccaaact agaagggat 2940
cgcgaaaatt tggaccctag gttgcgtgcg aaataaaca gggggctgac caggatagtg 3000
aacttgtgga actggttgcg attgttaacc cgtctgcgct aagctcgcgc gactcaataa 3060
aatgtgatt aagtctgaat gtggctctcc aatcatttcg attctctaat ctcccaatcc 3120
tctcaaaagg gccagtatcg gacacggact gggagggcg tacacgatag ttatatggta 3180
cagcagagge ctctgaacac ttagggaggag aattcagccg gggagagccc ctgttgagta 3240
ggcttgggag catattgcag gatgaacatg ttagtgatag ttctcgcctc ttgtcttgcg 3300
cgcctaactt ttgcgacgcg acacgtctc tttttggaag gcactcagge tgcctcggg 3360
gaagatgate ccagaaacgt tccggaaggg actgtaatca aatggacaaa agtctcggg 3420
aacgcgtgca agatgaagge gccgatgct tgctcttcgc ctaactattg ctttcatgat 3480
ttaatttacg acggaggaaa gaaagactgc ccgcccgcg gaccctgct tcgaaaacctg 3540
gtaattttac taaagcgcgg cgaaagctt 3569

```

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 5921

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Expression cassette insert

&lt;400&gt; SEQUENCE: 17

-continued

---

gtcgacggca	gagtcgcaga	cgcccctatt	ggacgtcaaa	attgtagagg	tgaagttttc	60
aaacgatggc	gaagtaacgg	cgacttgcgt	ttccaccgtc	aaatctccct	atagggtaga	120
aactaattgg	aaagtagacc	tcgtagatgt	aatggatgaa	atttctggga	acagtcccgc	180
cggggttttt	aacagtaatg	agaaaatggca	gaaacagctg	tactacagag	taaccgatgg	240
aagaacatcg	gtccagctaa	tgtgcctgtc	gtgcacgagc	cattctccgg	aaccttactg	300
tcttttcgac	acgtctctta	tagcgaggga	aaaagatata	gcgccagagt	tatactttac	360
ctctgatccg	caaacggcat	actgcacaat	aactctgccg	tccggcgttg	ttccgagatt	420
cgaatggagc	cttaataatg	tttcaactgcc	ggaatatttg	acggccacga	ccgttgtttc	480
gcataccgct	ggccaaagta	cagtgtagaa	gagcagcgcg	agagcaggcg	aggcgtggat	540
ttctggccgg	ggaggcaata	tatacgaatg	caccgtcctc	atctcagacg	gcactcgcgt	600
tactacgcga	aaggagaggt	gcttaacaaa	cacatggatt	gcggtggaaa	acggtgctgc	660
tcaggcgcag	ctgtattcac	tcttttctgg	acttgtgtca	ggattatgcg	ggagcatata	720
tgctttgtac	gcaacgctat	ggaccgccat	ttatttttga	ggaatgcttt	ttggactata	780
gtactgcttt	cttccttcgc	tagccagagc	accgccgccg	tcacgtacga	ctacatttta	840
ggcgcgcg	cgctcagcgc	gctaaccata	ccggcgggtg	gcccgtataa	cagatacttc	900
actagggtat	caagagcgtg	cgacgttgtc	gagctcaacc	cgatttctaa	cgtggacgac	960
atgatatcgg	cggccaaaga	aaaagagaag	gggggccctt	tcgaggcctc	cgtcgtctgg	1020
ttctacgtga	ttaagggcga	cgacggcgag	gacaagtact	gtccaatcta	tagaaaagag	1080
tacagggaat	gtggcgcagc	acaactgcta	tctgaatgcg	ccgttcaatc	tgacagatg	1140
tgggcagtg	actatgttcc	tagcaccctt	gtatcgcgaa	atggcgcggg	actgactata	1200
ttctccccca	ctgctgcgct	ctctggccaa	tacttgcgta	ccctgaaaat	cgggagattt	1260
gcgcaaacag	ctctcgtaac	tctagaagtt	aacgatcgtc	gtttaagat	cgggtcgcag	1320
cttaactttt	taccgtcgaa	atgctggaca	acagaacagt	atcagactgg	atttcaaggc	1380
gaacaccttt	atccgatcgc	agacaccaat	acacgacacg	cggacgacgt	atatcgggga	1440
tacgaagata	ttctgcagcg	ctggaataat	ttgctgagga	aaaagaatcc	tagcgcgcca	1500
gaccctcgtc	cagatagcgt	cccgaagaa	attcccgtcg	taaccaagaa	agcgggaagg	1560
cgcaccccg	acgcagaaag	cagcgaagaa	aaggcccctc	cagaagactc	ggaggacgac	1620
atgcaggcag	aggcttctgg	agaaaatcct	gccgccctcc	ccgaagacga	cgaagtcccc	1680
gaggacaccg	agcacgatga	tccaaactcg	gatcctgact	attacaatga	catgcccgcc	1740
gtgatcccgg	tggaggagac	tactaaaagt	tctaatagctg	tctccatgcc	catattcgcg	1800
gcgttcgtag	cctgcgcggt	cgcgctcgtg	gggctactgg	tttggagcat	cgtaaaatgc	1860
gcgcgtagct	aatcagcctc	agaatagggtg	gtttcttctc	acatgccacg	cctcacgctc	1920
ataatataaa	tcacatggaa	tagcatacca	atgcctattc	attgggacgt	tcgaaaagca	1980
tggcatcgct	acttggaaact	ctggetctcc	ttgcgcgac	gctcgcaccc	ttcggcgcga	2040
tgggaatcgt	gatcactgga	aatcacgtct	ccgccaggat	tgacgacgat	cacatcgtga	2100
tcgtcgcgcc	tcgccccgaa	gctacaatcc	aactgcagct	atttttcatg	cctggccaga	2160
gacccccaaa	accctactca	ggaaccgtcc	gcgtcgcggt	tcggtctgat	ataacaaacc	2220
agtgtacca	ggaacttagc	gaggagcgtc	ttgaaaattg	cactcatcga	tcgtcttctg	2280
ttttgtcgg	ctgtaaagtg	accgagtaca	cgttctccgc	ctcgaacaga	ctaaccggac	2340

---

-continued

---

ctccacaccc gtttaagctc actatacga aacctcgtcc gaacgacagc gggatgttct 2400  
acgtaattgt tcggctagac gacaccaaag aaccattga cgtcttcgcg atccaactat 2460  
cgggtgatca attcgcgaac accgcgcgga ctcgeggact ctattccaag gcttcgtgtc 2520  
gcaacttcgg attacctacc gtccaacttg aggctatct caggaccgag gaaagtggc 2580  
gcaactggca agcgtacgtt gccacggagg ccacgacgac cagcgccgag gcgacaaccc 2640  
cgagcccgt cactgcaacc agcgcctcgg aacttgaagc ggaacacttt acctttccct 2700  
ggctagaaaa tggcgtggat cattaogaac cgacaccgcg aaacgaaaat tcaaacgtta 2760  
ctgtccgtct cgggacaatg agccctacgc taattggggg aaccgtggct gccgtcgtga 2820  
gcgcaacgat cggcctcgtc attgtaattt ccacgtcac cagaacatg tgcacccgc 2880  
accgaaaatt agacacggtc tcgcaagacg acgaagaacg tccccaaact agaagggat 2940  
cgcgaaaatt tggaccatg gttgcgtgcg aaataaaca gggggctgac caggatagtg 3000  
aacttgtgga actggttgcg attgttaacc cgtctgctc aagctcggc gactcaataa 3060  
aatgtgatt aagtctgaat gtggctctcc aatcatttcg attctctaat ctcccaatcc 3120  
tctcaaaagg ggcagtatcg gacacggact gggagggcg tacacgatag ttatatggta 3180  
cagcagaggc ctctgaacac ttaggaggag aattcagccg gggagagccc ctgttgagta 3240  
ggcttgggag catattgcag gatgaacatg ttagtgatag ttctcgcctc ttgtcttgcg 3300  
cgctaactt ttgcgacgcg acacgtcctc tttttggaag gcactcaggc tgcctcggg 3360  
gaagatgatc ccagaaacgt tccggaaggg actgtaatca aatggacaaa agtctcggg 3420  
aacgctgca agatgaagge ggcgatgctc tgctcttcgc ctaactattg ctttcatgat 3480  
ttaatttacg acggaggaaa gaaagactgc ccgcccggg gaccctgctc tgcaaacctg 3540  
gtaattttac taaagcgcgg cgaaagcttc gcgccaggtc aattccctgg cattatgccc 3600  
agtacatgac cttatgggac tttcctactt ggcagtacat ctacgtatta gtcacgcta 3660  
ttaccatggt gatgcggttt tggcagtaca tcaatgggcg tggatagcgg tttgactcac 3720  
ggggatttcc aagtctccac ccattgacg tcaatgggag tttgttttg caccaaaatc 3780  
aacgggactt tccaaaatgt cgtaacaact ccgccccatt gacgcaaatg ggcggtagcg 3840  
tgtacggtag gaggctata taagcagagc tcgtttagt aaccgtcaga tcgcctggag 3900  
acgccatcca cgctgttttg acctccatag aagacaccgg ttgcgccgcc accatgggce 3960  
ccagacctc tacciaaagc ccagtaccta tgatgctgac tgcctgagtc gcctgggtac 4020  
tgagttgcat ctgtccggca aactccattg atggcaggcc tcttgcggct gcaggaattg 4080  
tggttacagg agacaaagcc gtcaacatat acacctcctc ccagacagga tcaatcatag 4140  
ttaagctcct ccgaatctg cccaaggata aggaggcatg tgcgaaagcc ccctggatg 4200  
catacaacag gacattgacc actttgctca cccccctgg tgactctatc cgtaggatac 4260  
aagagtctgt gactacatct ggagggggga gacaggggcy ccttatagge gccattattg 4320  
gcggtgtggc tcttgggggt gcaactgccc cacaataaac agcggccgca gctctgatac 4380  
aagcacaaca aatgctgccc aacatcctcc gacttaaga gagcattgcc gcaaccaatg 4440  
aggctgtgca tgaggtcact gacggattat cgcaactagc agtggcagtt gggagatgc 4500  
agcagttgt taatgacaa ttaataaaa cagctcagga attagactgc atcaaaattg 4560  
cacagcaagt tgggtgtagc ctcaacctgt acctaaccga attgactaca gtattcggac 4620

-continued

---

cacaaatcac	ttcacctgct	ttaaacaagc	tgactattca	ggcactttac	aatctagctg	4680
gtggaaatat	ggattactta	ttgactaagt	taggtgtagg	gaacaatcaa	ctcagctcat	4740
taatcggtag	cggttaate	accgtaacc	ctattctata	cgactcacag	actcaactct	4800
tgggtataca	ggtaactcta	ccttcagtcg	ggaagctaaa	taatatgctg	gccacctact	4860
tggaaacctt	atccgtaagc	acaaccaggg	gatttgcttc	ggcacttgtc	cmetaagtgg	4920
tgacacaggt	cggttctgtg	atagaagaac	tgacacctc	atactgtata	gaaactgact	4980
tacatttata	ttgtacaaga	atagtaacct	tcctatgtc	ccctggtatt	tattcctgct	5040
tgagcggcaa	tacgtcgccc	tgtatgtact	caaagaccga	aggcgcactt	actacaccat	5100
acatgactat	caaaggttca	gtcatcgcca	actgcaagat	gacaacatgt	agatgtgtaa	5160
acccccggg	tatcatatcg	caaaactatg	gagaagccgt	gtctctaata	gataaacaat	5220
catgcaatgt	tttatectta	ggcgggataa	ctttaaggct	cagtggggaa	ttcgatgtaa	5280
cttatcagaa	gaatatctca	atacaagatt	ctcaagtaat	aataacaggc	aatcttgata	5340
tctcaactga	gcttgggaat	gtcaacaact	cgatcagtaa	tgctttgaat	aagttagagg	5400
aaagcaacag	aaaactagac	aaagtcaatg	tcaaaactgac	tagcacatct	gctctcatta	5460
cctatatcgt	gttgactatc	atatctcttg	tttttggtat	acttagcctg	attctagcat	5520
gctaccctaat	gtacaagcaa	aaggcgcaac	aaaagacctt	attatggctt	gggaataata	5580
ctctagatca	gatgagagcc	actacaaaaa	tgtgaggatc	tctcgaggaa	ttctagatcc	5640
cacgtcacta	ttgtatactc	tatatatac	tctatggtat	actctgtaat	cctactcaat	5700
aaacgtgtca	cgctgtgaa	accgtactaa	gtctcccgtg	tcttcttctc	accatcaggt	5760
gacatcctcg	cccaggtgtg	caatcatgcc	ggatcagatt	ccagtagcac	cggccccacg	5820
ctgacaacct	actcttcgag	cgtagcagc	gcccccttta	acaagccgac	ccccaccagc	5880
gtcgcgggta	ctaactctcc	tctccccgac	ctgcaactag	t		5921

---

We claim:

1. A recombinant nonpathogenic Marek's Disease Virus (rMDV<sub>np</sub>) comprising a first nucleic acid inserted in a first nonessential site in the rMDV<sub>np</sub> genome and a second nucleic acid inserted in a second nonessential site in the rMDV<sub>np</sub> genome;

wherein the first nucleic acid comprises both a nucleotide sequence that encodes an Infectious Laryngotracheitis Virus (ILTV) gD protein and a nucleotide sequence that encodes an Infectious Laryngotracheitis Virus (ILTV) gI protein;

wherein the second nucleic acid comprises a nucleotide sequence that encodes a Newcastle Disease Virus fusion protein (NDV F); and

wherein the first nonessential site and the second nonessential site are either the same or only the first nonessential site is the US2 site.

2. The rMDV<sub>np</sub> of claim 1 wherein the first nonessential site and the second nonessential site are the US2 site.

3. The rMDV<sub>np</sub> of claim 1 wherein the first nonessential site and the second nonessential site are the UL54.5 site.

4. The rMDV<sub>np</sub> of claim 1 wherein the first nonessential site is the US2 site and the second nonessential site is the UL7/8 site.

5. The rMDV<sub>np</sub> of claim 1 wherein the nucleotide sequence encoding the ILTV gD protein is operatively under the control

of a first promoter, the nucleotide sequence encoding the ILTV gI protein is operatively under the control of a second promoter, and the nucleotide sequence encoding the NDV F protein is operatively under the control of a third promoter.

6. The rMDV<sub>np</sub> of claim 5 wherein the first promoter, the second promoter, and the third promoter are all different.

7. The rMDV<sub>np</sub> of claim 6 wherein the first promoter is the endogenous ILTV gD promoter and the second promoter is the endogenous ILTV gI promoter.

8. The rMDV<sub>np</sub> of claim 7 wherein the third promoter is the human cytomegalovirus immediate early (hCMV IE) promoter.

9. The rMDV<sub>np</sub> of claim 8 that is a recombinant herpesvirus of turkeys (rHVT).

10. A recombinant nucleic acid comprising in 5' to 3' direction in the following order:

(i) an Infectious Laryngotracheitis Virus (ILTV) gD promoter;

(ii) a coding sequence for the ILTV gD protein;

(iii) an ILTV gI promoter;

(iv) a coding sequence for the ILTV gI protein;

(v) a human cytomegalovirus immediate early (hCMV IE) promoter;

(vi) a coding sequence for the Newcastle Disease Virus fusion protein (NDV F); and

(vii) a transcription terminator sequence.

**11.** The recombinant nucleic acid of claim **10** which comprises the nucleotide sequence of SEQ ID NO: 17.

**12.** A recombinant nonpathogenic Marek's Disease virus (rMDV<sub>np</sub>) comprising the recombinant nucleic acid of claim **11** inserted into a nonessential site.

**13.** The rMDV<sub>np</sub> of claim **12** wherein the nonessential insertion site is selected from the group consisting of US2 and UL54.5.

**14.** The rMDV<sub>np</sub> of claim **13** that is a recombinant herpesvirus of turkeys (rHVT).

**15.** An rMDV<sub>np</sub> comprising the recombinant nucleic acid of claim **10** inserted into a nonessential site.

**16.** The rMDV<sub>np</sub> of claim **15** that is a recombinant herpesvirus of turkeys (rHVT).

**17.** A vaccine comprising the rMDV<sub>np</sub> of claim **1**.

**18.** The vaccine of claim **17** wherein the rMDV<sub>np</sub> is a recombinant herpesvirus of turkeys (rHVT).

**19.** The vaccine of claim **18** that further comprises a mild live infectious bursal disease virus (IBDV).

**20.** The vaccine of claim **19** wherein the mild live IBDV is strain 89/03.

**21.** A vaccine comprising the rHVT of claim **14**.

**22.** The vaccine of claim **21** that further comprises a mild live IBDV.

**23.** The vaccine of claim **22** wherein the mild live IBDV is strain 89/03.

**24.** A method for aiding in the protection of a chicken against ILTV comprising administering the vaccine of claim **23**.

**25.** A method for aiding in the protection of a chicken against ILTV comprising administering the vaccine of claim **17**.

\* \* \* \* \*