



US011455754B2

(12) **United States Patent**
Chui et al.

(10) **Patent No.:** **US 11,455,754 B2**

(45) **Date of Patent:** **Sep. 27, 2022**

(54) **SYSTEM AND METHOD FOR SYNTHESIZING LOW-DIMENSIONAL IMAGE DATA FROM HIGH-DIMENSIONAL IMAGE DATA USING AN OBJECT GRID ENHANCEMENT**

(71) Applicant: **HOLOGIC, INC.**, Marlborough, MA (US)

(72) Inventors: **Haili Chui**, Fremont, CA (US);
Xiaomin Liu, Sunnyvale, CA (US);
Xiangwei Zhang, Fremont, CA (US)

(73) Assignee: **Hologic, Inc.**, Marlborough, MA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/497,766**

(22) PCT Filed: **Mar. 28, 2018**

(86) PCT No.: **PCT/US2018/024912**

§ 371 (c)(1),

(2) Date: **Sep. 25, 2019**

(87) PCT Pub. No.: **WO2018/183549**

PCT Pub. Date: **Oct. 4, 2018**

(65) **Prior Publication Data**

US 2021/0118199 A1 Apr. 22, 2021

Related U.S. Application Data

(60) Provisional application No. 62/479,008, filed on Mar. 30, 2017.

(51) **Int. Cl.**

G06T 11/00 (2006.01)

G06N 20/00 (2019.01)

(Continued)

(52) **U.S. Cl.**

CPC **G06T 11/005** (2013.01); **A61B 6/502** (2013.01); **G06N 20/00** (2019.01); **G06T 7/0012** (2013.01);

(Continued)

(58) **Field of Classification Search**

CPC ... **G06T 11/005**; **G06T 7/0012**; **G06T 15/205**; **G06T 2207/10072**; **G06T 2207/10116**;

(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,502,878 A 3/1970 Stewart
3,863,073 A 1/1975 Wagner

(Continued)

FOREIGN PATENT DOCUMENTS

AU 2014339982 A1 * 5/2016 A61B 6/481
CN 1846622 10/2006

(Continued)

OTHER PUBLICATIONS

M. Ertas, A. Akan, I. Yildirim, A. Dinler and M. Kamasak, "2D versus 3D total variation minimization in digital breast tomosynthesis," 2015 IEEE International Conference on Imaging Systems and Techniques (IST), Macau, 2015, pp. 1-4, doi: 10.1109/IST.2015.7294553. (Year: 2015).*

(Continued)

Primary Examiner — Vu Le


Assistant Examiner — Courtney Joan Nelson

(74) *Attorney, Agent, or Firm* — Merchant & Gould P.C.

(57) **ABSTRACT**

A method for processing breast tissue image data includes processing image data of a patient's breast tissue to generate a high-dimensional grid depicting one or more high-dimensional objects in the patient's breast tissue; determining a probability or confidence of each of the one or more high-dimensional objects depicted in the high-dimensional grid;

(Continued)



and modifying one or more aspects of at least one of the one or more high-dimensional objects based at least in part on its respective determined probability or confidence to thereby generate a lower-dimensional format version of the one or more high-dimensional objects. The method may further include displaying the lower-dimensional format version of the one or more high-dimensional objects in a synthesized image of the patient's breast tissue.

23 Claims, 9 Drawing Sheets

(51) **Int. Cl.**

A61B 6/00 (2006.01)

G06T 7/00 (2017.01)

G06T 15/20 (2011.01)

(52) **U.S. Cl.**

CPC .. **G06T 15/205** (2013.01); **G06T 2207/10072** (2013.01); **G06T 2207/10116** (2013.01); **G06T 2207/30068** (2013.01)

(58) **Field of Classification Search**

CPC **G06T 2207/30068**; **G06N 20/00**; **A61B 6/502**

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

3,971,950	A	7/1976	Evans et al.	5,592,562	A	1/1997	Rooks
4,160,906	A	7/1979	Daniels	5,594,769	A	1/1997	Pellegrino et al.
4,310,766	A	1/1982	Finkenzeller et al.	5,596,200	A	1/1997	Sharma
4,496,557	A	1/1985	Malen et al.	5,598,454	A	1/1997	Franetzki
4,559,641	A	12/1985	Caugant et al.	5,609,152	A	3/1997	Pellegrino et al.
4,706,269	A	11/1987	Reina et al.	5,627,869	A	5/1997	Andrew et al.
4,744,099	A	5/1988	Huettenrauch	5,642,433	A	6/1997	Lee et al.
4,773,086	A	9/1988	Fujita	5,642,441	A	6/1997	Riley et al.
4,773,087	A	9/1988	Plewes	5,647,025	A	7/1997	Frost et al.
4,819,258	A	4/1989	Kleinman et al.	5,657,362	A	8/1997	Giger et al.
4,821,727	A	4/1989	Levene et al.	5,668,889	A	9/1997	Hara
4,907,156	A	6/1990	Doi et al.	5,671,288	A	9/1997	Wilhelm et al.
4,969,174	A	11/1990	Schied	5,712,890	A	1/1998	Spivey
4,989,227	A	1/1991	Tirelli et al.	5,719,952	A	2/1998	Rooks
5,018,176	A	5/1991	Romeas et al.	5,735,264	A	4/1998	Siczek et al.
RE33,634	E	7/1991	Yanaki	5,763,871	A	6/1998	Ortyn et al.
5,029,193	A	7/1991	Saffer	5,769,086	A	6/1998	Ritchart et al.
5,051,904	A	9/1991	Griffith	5,773,832	A	6/1998	Sayed et al.
5,078,142	A	1/1992	Siczek et al.	5,803,912	A	9/1998	Siczek et al.
5,099,846	A	3/1992	Hardy	5,818,898	A	10/1998	Tsukamoto et al.
5,129,911	A	7/1992	Siczek et al.	5,828,722	A	10/1998	Ploetz
5,133,020	A	7/1992	Giger et al.	5,835,079	A	11/1998	Shieh
5,163,075	A	11/1992	Lubinsky	5,841,124	A	11/1998	Ortyn et al.
5,164,976	A	11/1992	Scheid et al.	5,872,828	A	2/1999	Niklason et al.
5,199,056	A	3/1993	Darrah	5,875,258	A	2/1999	Ortyn et al.
5,219,351	A	6/1993	Teubner	5,878,104	A	3/1999	Ploetz
5,240,011	A	8/1993	Assa	5,878,746	A	3/1999	Lemelson et al.
5,279,309	A	1/1994	Taylor et al.	5,896,437	A	4/1999	Ploetz
5,280,427	A	1/1994	Magnusson	5,941,832	A	8/1999	Tumey
5,289,520	A	2/1994	Pellegrino et al.	5,954,650	A	9/1999	Saito
5,343,390	A	8/1994	Doi et al.	5,986,662	A	11/1999	Argiro
5,359,637	A	10/1994	Webbe	6,005,907	A	12/1999	Ploetz
5,365,562	A	11/1994	Toker	6,022,325	A	2/2000	Siczek et al.
5,386,447	A	1/1995	Siczek	6,067,079	A	5/2000	Shieh
5,415,169	A	5/1995	Siczek et al.	6,075,879	A	6/2000	Roehrig et al.
5,426,685	A	6/1995	Pellegrino et al.	6,091,841	A	7/2000	Rogers
5,452,367	A	9/1995	Bick	6,101,236	A	8/2000	Wang et al.
5,491,627	A	2/1996	Zhang et al.	6,102,866	A	8/2000	Nields et al.
5,499,097	A	3/1996	Ortyn et al.	6,137,527	A	10/2000	Abdel-Malek
5,506,877	A	4/1996	Niklason et al.	6,141,398	A	10/2000	He
5,526,394	A	6/1996	Siczek	6,149,301	A	11/2000	Kautzer et al.
5,539,797	A	7/1996	Heidsieck et al.	6,175,117	B1	1/2001	Komardin
5,553,111	A	9/1996	Moore	6,196,715	B1	3/2001	Nambu
				6,215,892	B1	4/2001	Douglass et al.
				6,216,540	B1	4/2001	Nelson
				6,219,059	B1	4/2001	Argiro
				6,256,370	B1	4/2001	Yavus
				6,233,473	B1	5/2001	Sheperd
				6,243,441	B1	6/2001	Zur
				6,245,028	B1	6/2001	Furst et al.
				6,272,207	B1	8/2001	Tang
				6,289,235	B1	9/2001	Webber et al.
				6,292,530	B1	9/2001	Yavus
				6,293,282	B1	9/2001	Lemelson
				6,327,336	B1	12/2001	Gingold et al.
				6,327,377	B1	12/2001	Rutenberg et al.
				6,341,156	B1	1/2002	Baetz
				6,375,352	B1	4/2002	Hewes
				6,389,104	B1	5/2002	Bani-Hashemi et al.
				6,411,836	B1	6/2002	Patel
				6,415,015	B2	7/2002	Nicolas
				6,424,332	B1	7/2002	Powell
				6,442,288	B1	8/2002	Haerer
				6,459,925	B1	10/2002	Nields et al.
				6,463,181	B2	10/2002	Duarte
				6,468,226	B1	10/2002	McIntyre, IV
				6,480,565	B1	11/2002	Ning
				6,501,819	B2	12/2002	Unger et al.
				6,556,655	B1	4/2003	Chichereau
				6,574,304	B1	6/2003	Hsieh
				6,597,762	B1	7/2003	Ferrant
				6,611,575	B1	8/2003	Alyassin et al.
				6,620,111	B2	9/2003	Stephens et al.
				6,626,849	B2	9/2003	Huitema et al.
				6,633,674	B1	10/2003	Barnes
				6,638,235	B2	10/2003	Miller et al.
				6,647,092	B2	11/2003	Eberhard
				6,650,928	B1	11/2003	Gailly
				6,683,934	B1	1/2004	Zhao

(56)	References Cited					
	U.S. PATENT DOCUMENTS					
	6,744,848 B2	6/2004	Stanton		9,592,019 B2	3/2017 Sugiyama
	6,748,044 B2	6/2004	Sabol et al.		9,805,507 B2	10/2017 Chen
	6,751,285 B2	6/2004	Eberhard		9,808,215 B2	11/2017 Ruth et al.
	6,758,824 B1	7/2004	Miller et al.		9,811,758 B2	11/2017 Ren et al.
	6,813,334 B2	11/2004	Koppe		9,901,309 B2	2/2018 DeFreitas et al.
	6,882,700 B2	4/2005	Wang		10,008,184 B2	6/2018 Kreeger et al.
	6,885,724 B2	4/2005	Li		10,010,302 B2	7/2018 Ruth et al.
	6,901,156 B2	5/2005	Giger et al.		10,092,358 B2	10/2018 DeFreitas
	6,912,319 B1	5/2005	Barnes		10,111,631 B2	10/2018 Gkanatsios
	6,940,943 B2	9/2005	Claus		10,242,490 B2	3/2019 Karssemeijer
	6,978,040 B2	12/2005	Berestov		10,335,094 B2	7/2019 DeFreitas
	6,987,331 B2	1/2006	Koeppe		10,357,211 B2	7/2019 Smith
	6,999,554 B2	2/2006	Mertelmeier		10,410,417 B2	9/2019 Chen et al.
	7,022,075 B2	4/2006	Grunwald et al.		10,413,263 B2	9/2019 Ruth et al.
	7,025,725 B2	4/2006	Dione et al.		10,444,960 B2	10/2019 Marshall
	7,030,861 B1	4/2006	Westerman		10,456,213 B2	10/2019 DeFreitas
	7,110,490 B2	9/2006	Eberhard		10,573,276 B2	2/2020 Kreeger et al.
	7,110,502 B2	9/2006	Tsuji		10,575,807 B2	3/2020 Gkanatsios
	7,117,098 B1	10/2006	Dunlay et al.		10,595,954 B2	3/2020 DeFreitas
	7,123,684 B2	10/2006	Jing et al.		10,624,598 B2	4/2020 Chen
	7,127,091 B2	10/2006	OpDeBeek		10,977,863 B2	4/2021 Chen
	7,142,633 B2	11/2006	Eberhard		10,978,026 B2	4/2021 Kreeger
	7,218,766 B2*	5/2007	Eberhard	A61B 6/463 382/131	2001/0038681 A1	11/2001 Stanton et al.
					2001/0038861 A1	11/2001 Hsu et al.
					2002/0012450 A1	1/2002 Tsuji
					2002/0050986 A1	5/2002 Inoue
					2002/0075997 A1	6/2002 Unger et al.
					2002/0113681 A1	8/2002 Byram
					2002/0122533 A1	9/2002 Marie et al.
					2002/0188466 A1	12/2002 Barrette et al.
					2002/0193676 A1	12/2002 Bodicker
					2003/0007598 A1	1/2003 Wang
					2003/0018272 A1	1/2003 Treado et al.
					2003/0026386 A1	2/2003 Tang
					2003/0048260 A1	3/2003 Matusis
					2003/0073895 A1	4/2003 Nields et al.
					2003/0095624 A1	5/2003 Eberhard et al.
					2003/0097055 A1	5/2003 Yanof
					2003/0128893 A1	7/2003 Castorina
					2003/0135115 A1	7/2003 Burdette et al.
					2003/0169847 A1	9/2003 Karellas
					2003/0194050 A1	10/2003 Eberhard
					2003/0194121 A1	10/2003 Eberhard et al.
					2003/0195433 A1	10/2003 Turovskiy
					2003/0210254 A1	11/2003 Doan
					2003/0212327 A1	11/2003 Wang
					2003/0215120 A1	11/2003 Uppaluri
					2004/0008809 A1	1/2004 Webber
					2004/0008900 A1	1/2004 Jabri et al.
					2004/0008901 A1	1/2004 Avinash
					2004/0036680 A1	2/2004 Davis
					2004/0047518 A1	3/2004 Tiana
					2004/0052328 A1	3/2004 Saboi
					2004/0064037 A1	4/2004 Smith
					2004/0066884 A1	4/2004 Claus
					2004/0066904 A1	4/2004 Eberhard et al.
					2004/0070582 A1	4/2004 Smith et al.
					2004/0077938 A1	4/2004 Mark et al.
					2004/0081273 A1	4/2004 Ning
					2004/0094167 A1	5/2004 Brady
					2004/0101095 A1	5/2004 Jing et al.
					2004/0109028 A1	6/2004 Stern et al.
					2004/0109529 A1	6/2004 Eberhard et al.
					2004/0127789 A1	7/2004 Ogawa
					2004/0138569 A1	7/2004 Grunwald
					2004/0171933 A1	9/2004 Stoller et al.
					2004/0171986 A1	9/2004 Tremaglio, Jr. et al.
					2004/0267157 A1	12/2004 Miller et al.
					2005/0047636 A1	3/2005 Gines et al.
					2005/0049521 A1	3/2005 Miller et al.
					2005/0063509 A1	3/2005 DeFreitas et al.
					2005/0078797 A1	4/2005 Danielsson et al.
					2005/0084060 A1	4/2005 Seppi et al.
					2005/0089205 A1	4/2005 Kapur
					2005/0105679 A1	5/2005 Wu et al.
					2005/0107689 A1	5/2005 Sasano
					2005/0111718 A1	5/2005 MacMahon
					2005/0113681 A1	5/2005 DeFreitas et al.
					2005/0113715 A1	5/2005 Schwindt et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2005/0124845	A1	6/2005	Thomadsen et al.	2009/0278812	A1	11/2009	Yasutake
2005/0135555	A1	6/2005	Claus	2009/0296882	A1	12/2009	Gkanatsios et al.
2005/0135664	A1	6/2005	Kaufhold	2009/0304147	A1	12/2009	Jing et al.
2005/0226375	A1	10/2005	Eberhard	2010/0034348	A1	2/2010	Yu
2006/0009693	A1	1/2006	Hanover et al.	2010/0049046	A1	2/2010	Peiffer
2006/0018526	A1	1/2006	Avinash	2010/0054400	A1	3/2010	Ren et al.
2006/0025680	A1	2/2006	Jeune-Iomme	2010/0079405	A1	4/2010	Bernstein
2006/0030784	A1	2/2006	Miller et al.	2010/0086188	A1	4/2010	Ruth et al.
2006/0074288	A1	4/2006	Kelly et al.	2010/0088346	A1	4/2010	Urness et al.
2006/0098855	A1	5/2006	Gkanatsios et al.	2010/0098214	A1	4/2010	Star-Lack et al.
2006/0129062	A1	6/2006	Nicoson et al.	2010/0105879	A1	4/2010	Katayose et al.
2006/0132508	A1	6/2006	Sadikali	2010/0121178	A1	5/2010	Krishnan
2006/0147099	A1	7/2006	Marshall et al.	2010/0131294	A1	5/2010	Venon
2006/0155209	A1	7/2006	Miller et al.	2010/0131482	A1	5/2010	Linthicum et al.
2006/0197753	A1	9/2006	Hotelling	2010/0135558	A1	6/2010	Ruth et al.
2006/0210131	A1*	9/2006	Wheeler, Jr. G06T 11/008 382/128	2010/0152570	A1	6/2010	Navab
2006/0228012	A1	10/2006	Masuzawa	2010/0166267	A1	7/2010	Zhang
2006/0238546	A1	10/2006	Handley	2010/0195882	A1	8/2010	Ren et al.
2006/0257009	A1	11/2006	Wang	2010/0208037	A1	8/2010	Sendai
2006/0269040	A1	11/2006	Mertelmeier	2010/0231522	A1	9/2010	Li
2006/0291618	A1	12/2006	Eberhard et al.	2010/0246909	A1	9/2010	Blum
2007/0019846	A1	1/2007	Bullitt et al.	2010/0259561	A1	10/2010	Forutanpour et al.
2007/0030949	A1	2/2007	Jing et al.	2010/0259645	A1	10/2010	Kaplan
2007/0036265	A1	2/2007	Jing et al.	2010/0260316	A1	10/2010	Stein et al.
2007/0046649	A1	3/2007	Reiner	2010/0280375	A1	11/2010	Zhang
2007/0052700	A1	3/2007	Wheeler et al.	2010/0293500	A1	11/2010	Cragun
2007/0076844	A1	4/2007	Defreitas et al.	2011/0018817	A1	1/2011	Kryze
2007/0114424	A1	5/2007	Danielsson et al.	2011/0019891	A1	1/2011	Puong
2007/0118400	A1	5/2007	Morita et al.	2011/0054944	A1	3/2011	Sandberg et al.
2007/0156451	A1	7/2007	Gering	2011/0069808	A1	3/2011	Defreitas et al.
2007/0223651	A1	9/2007	Wagenaar et al.	2011/0069906	A1	3/2011	Park
2007/0225600	A1	9/2007	Weibrecht et al.	2011/0087132	A1	4/2011	DeFreitas et al.
2007/0236490	A1	10/2007	Castele	2011/0105879	A1	5/2011	Masumoto
2007/0242800	A1	10/2007	Jing et al.	2011/0109650	A1	5/2011	Kreeger
2007/0263765	A1	11/2007	Wu	2011/0110576	A1*	5/2011	Kreeger G16H 50/20 382/132
2007/0274585	A1	11/2007	Zhang et al.	2011/0150447	A1	6/2011	Li
2008/0019581	A1	1/2008	Gkanatsios et al.	2011/0163939	A1	7/2011	Tam et al.
2008/0043905	A1	2/2008	Hassanpourgol	2011/0178389	A1	7/2011	Kumar et al.
2008/0045833	A1	2/2008	DeFreitas et al.	2011/0182402	A1	7/2011	Partain
2008/0101537	A1	5/2008	Sendai	2011/0234630	A1	9/2011	Batman et al.
2008/0114614	A1	5/2008	Mahesh et al.	2011/0237927	A1	9/2011	Brooks et al.
2008/0125643	A1	5/2008	Huisman	2011/0242092	A1	10/2011	Kashiwagi
2008/0130979	A1	6/2008	Ren	2011/0310126	A1	12/2011	Georgiev et al.
2008/0139896	A1	6/2008	Baumgart	2012/0014504	A1	1/2012	Jang
2008/0152086	A1	6/2008	Hall	2012/0014578	A1*	1/2012	Karssemeijer G06T 7/0012 382/131
2008/0165136	A1	7/2008	Christie et al.	2012/0069951	A1	3/2012	Toba
2008/0187095	A1	8/2008	Boone et al.	2012/0131488	A1	5/2012	Karlsson et al.
2008/0198966	A1	8/2008	Hjarn	2012/0133600	A1	5/2012	Marshall
2008/0221479	A1	9/2008	Ritchie	2012/0133601	A1	5/2012	Marshall
2008/0229256	A1	9/2008	Shibaika	2012/0134464	A1	5/2012	Hoernig et al.
2008/0240533	A1	10/2008	Piron et al.	2012/0148151	A1	6/2012	Hamada
2008/0297482	A1	12/2008	Weiss	2012/0189092	A1	7/2012	Jerebko
2009/0003519	A1	1/2009	DeFreitas	2012/0194425	A1	8/2012	Buelow
2009/0005668	A1	1/2009	West et al.	2012/0238870	A1	9/2012	Smith et al.
2009/0010384	A1	1/2009	Jing et al.	2012/0293511	A1	11/2012	Mertelmeier
2009/0034684	A1	2/2009	Bernard	2013/0022165	A1	1/2013	Jang
2009/0037821	A1	2/2009	O'Neal et al.	2013/0044861	A1	2/2013	Muller
2009/0079705	A1	3/2009	Sizelove et al.	2013/0059758	A1*	3/2013	Haick B82Y 15/00 977/773
2009/0080594	A1	3/2009	Brooks et al.	2013/0108138	A1	5/2013	Nakayama
2009/0080602	A1	3/2009	Brooks et al.	2013/0121569	A1	5/2013	Yadav
2009/0080604	A1	3/2009	Shores et al.	2013/0121618	A1	5/2013	Yadav
2009/0080752	A1	3/2009	Ruth	2013/0202168	A1*	8/2013	Jerebko G06T 11/006 382/131
2009/0080765	A1	3/2009	Bernard et al.	2013/0259193	A1	10/2013	Packard
2009/0087067	A1	4/2009	Khorasani	2014/0033126	A1	1/2014	Kreeger
2009/0123052	A1	5/2009	Ruth	2014/0035811	A1	2/2014	Guehring
2009/0129644	A1	5/2009	Daw et al.	2014/0064444	A1	3/2014	Oh
2009/0135997	A1	5/2009	Defreitas et al.	2014/0073913	A1	3/2014	DeFreitas et al.
2009/0138280	A1	5/2009	Morita et al.	2014/0219534	A1	8/2014	Wiemker et al.
2009/0143674	A1	6/2009	Nields	2014/0219548	A1	8/2014	Wels et al.
2009/0167702	A1	7/2009	Nurmi	2014/0327702	A1	11/2014	Kreeger et al.
2009/0171244	A1	7/2009	Ning	2014/0328517	A1	11/2014	Gluncic
2009/0238424	A1	9/2009	Arakita	2015/0052471	A1	2/2015	Chen et al.
2009/0259958	A1	10/2009	Ban	2015/0061582	A1	4/2015	Smith et al.
2009/0268865	A1	10/2009	Ren et al.	2015/0238148	A1	8/2015	Georgescu
				2015/0302146	A1	10/2015	Marshall

(56)

References Cited

U.S. PATENT DOCUMENTS

2015/0309712 A1 10/2015 Marshall
 2015/0317538 A1 11/2015 Ren et al.
 2015/0331995 A1 11/2015 Zhao
 2016/0000399 A1 1/2016 Halmann et al.
 2016/0022364 A1 1/2016 DeFreitas et al.
 2016/0051215 A1 2/2016 Chen
 2016/0078645 A1 3/2016 Abdurahman et al.
 2016/0140749 A1 5/2016 Erhard
 2016/0228034 A1 8/2016 Gluncic
 2016/0235380 A1 8/2016 Smith
 2016/0367210 A1 12/2016 Gkanatsios
 2017/0071562 A1 3/2017 Suzuki
 2017/0262737 A1 9/2017 Rabinovich
 2018/0047211 A1 2/2018 Chen et al.
 2018/0137385 A1 5/2018 Ren
 2018/0144244 A1 5/2018 Masoud
 2018/0256118 A1 9/2018 DeFreitas
 2019/0015173 A1 1/2019 DeFreitas
 2019/0043456 A1 2/2019 Kreeger
 2019/0290221 A1 9/2019 Smith
 2020/0046303 A1 2/2020 DeFreitas
 2020/0093562 A1 3/2020 DeFreitas
 2020/0184262 A1 6/2020 Chui
 2020/0205928 A1 7/2020 DeFreitas
 2020/0253573 A1 8/2020 Gkanatsios
 2020/0345320 A1 11/2020 Chen
 2020/0390404 A1 12/2020 DeFreitas
 2021/0000553 A1 1/2021 St. Pierre
 2021/0100518 A1 4/2021 Chui
 2021/0100626 A1 4/2021 St. Pierre
 2021/0113167 A1 4/2021 Chui
 2022/0005277 A1 1/2022 Chen
 2022/0013089 A1 1/2022 Kreeger
 2022/0192615 A1 6/2022 Chui

FOREIGN PATENT DOCUMENTS

CN 202161328 3/2012
 CN 102429678 5/2012
 CN 107440730 12/2017
 DE 102010009295 8/2011
 DE 102011087127 5/2013
 EP 775467 5/1997
 EP 982001 3/2000
 EP 1428473 6/2004
 EP 2236085 6/2010
 EP 2215600 8/2010
 EP 2301432 3/2011
 EP 2491863 8/2012
 EP 1986548 1/2013
 EP 2656789 10/2013
 EP 2823464 1/2015
 EP 2823765 1/2015
 EP 3060132 4/2019
 JP H09-198490 7/1997
 JP H09-238934 9/1997
 JP 10-33523 2/1998
 JP H10-33523 2/1998
 JP 2000-200340 7/2000
 JP 2002-282248 10/2002
 JP 2003-189179 7/2003
 JP 2003-199737 7/2003
 JP 2003-531516 10/2003
 JP 2006-519634 8/2006
 JP 2006-312026 11/2006
 JP 2007-130487 5/2007
 JP 2007-330334 12/2007
 JP 2007-536968 12/2007
 JP 2008-068032 3/2008
 JP 2009-034503 2/2009
 JP 2009-522005 6/2009
 JP 2009-526618 7/2009
 JP 2009-207545 9/2009
 JP 2010-137004 6/2010
 JP 2011-110175 A 6/2011

JP 2012-501750 1/2012
 JP 2012011255 1/2012
 JP 2012-061196 3/2012
 JP 2013-244211 12/2013
 JP 2014-507250 3/2014
 JP 2014-534042 12/2014
 JP 2015-506794 3/2015
 JP 2015-144632 A 8/2015
 JP 2016-198197 12/2015
 KR 10-2015-0010515 1/2015
 KR 10-2017-0062839 6/2017
 WO 90/05485 5/1990
 WO 93/17620 9/1993
 WO 94/06352 3/1994
 WO 1997/00649 1/1997
 WO 1998/16903 4/1998
 WO 00/51484 9/2000
 WO 2003/020114 3/2003
 WO 2005051197 6/2005
 WO 2005/110230 11/2005
 WO 2005/110230 11/2005
 WO 2005/112767 12/2005
 WO 2005/112767 12/2005
 WO 2006/055830 5/2006
 WO 2006/058160 6/2006
 WO 2007/095330 8/2007
 WO 08/014670 2/2008
 WO 2008047270 4/2008
 WO 2008/054436 5/2008
 WO 2009/026587 2/2009
 WO 2010/028208 3/2010
 WO 2010059920 5/2010
 WO 2011008239 1/2011
 WO 2011/043838 4/2011
 WO 2011065950 6/2011
 WO 2011073864 6/2011
 WO 2011091300 7/2011
 WO 2012/001572 1/2012
 WO 2012/068373 5/2012
 WO 2012063653 5/2012
 WO 2012/112627 8/2012
 WO 2012/122399 9/2012
 WO 2013/001439 1/2013
 WO 2013/035026 3/2013
 WO 2013/078476 5/2013
 WO WO-2013123091 A1 * 8/2013 A61B 6/466
 WO 2014/149554 9/2014
 WO WO2014/207080 12/2014
 WO WO-2014207080 A1 * 12/2014 G06T 15/08
 WO 2015/061582 4/2015
 WO 2015/066650 5/2015
 WO 2015/130916 9/2015
 WO 2016/103094 6/2016
 WO 2016/184746 11/2016
 WO 2018/183548 10/2018
 WO 2018/183550 10/2018
 WO WO2018/183549 10/2018
 WO 2018/236565 12/2018

OTHER PUBLICATIONS

B. E. Caroline and N. Vijayanthi, "Computer aided detection of masses in digital breast tomosynthesis: A review," 2012 International Conference on Emerging Trends in Science, Engineering and Technology (INCOSSET), Tiruchirappalli, 2012, pp. 186-191, doi: 10.1109/INCOSSET.2012.6513903 (Year: 2012).*

International Search Report and Written Opinion dated Jun. 8, 2018 for PCT application No. PCT/US2018/024912, applicant Hologic, Inc., 10 pages.

Non-Final Office Action for U.S. Appl. No. 16/497,767 dated Feb. 19, 2021.

U.S. Appl. No. 16/497,767, Office Action dated Jul. 16, 2021, 26 pages.

PCT International Preliminary Report on Patentability in International Application PCT/US2018/024912, dated Oct. 10, 2019, 7 pages.

(56)

References Cited

OTHER PUBLICATIONS

- “Filtered Back Projection”, (Nygren), published May 8, 2007, URL: <http://web.archive.org/web/19991010131715/http://www.owl.net/~elec539/Projects97/cult/node2.html>, 2 pgs.
- Berg, WA et al., “Combined screening with ultrasound and mammography vs mammography alone in women at elevated risk of breast cancer”, *JAMA* 299:2151-2163, 2008.
- Carton, AK, et al., “Dual-energy contrast-enhanced digital breast tomosynthesis—a feasibility study”, *BR J Radiol.* Apr. 2010;83 (988):344-50.
- Chen, SC, et al., “Initial clinical experience with contrast-enhanced digital breast tomosynthesis”, *Acad Radio.* Feb. 2007 14(2):229-38.
- Diekmann, F., et al., “Digital mammography using iodine-based contrast media: initial clinical experience with dynamic contrast medium enhancement”, *Invest Radiol* 2005; 40:397-404.
- Dromain C., et al., “Contrast enhanced spectral mammography: a multi-reader study”, *RSNA 2010, 96th Scientific Assembly and Scientific Meeting.*
- Dromain, C., et al., “Contrast-enhanced digital mammography”, *Eur J Radiol.* 2009; 69:34-42.
- Freiherr, G., “Breast tomosynthesis trials show promise”, *Diagnostic Imaging—San Francisco* 2005, V27; N4:42-48.
- Giger et al. “Development of a smart workstation for use in mammography”, in *Proceedings of SPIE*, vol. 1445 (1991), p. 101103; 4 pages.
- Giger et al., “An Intelligent Workstation for Computer-aided Diagnosis”, in *RadioGraphics*, May 1993, 13:3 pp. 647-656; 10 pages.
- Hologic, “Lorad StereoLoc II” Operator’s Manual 9-500-0261, Rev. 005, 2004, 78 pgs.
- Hologic, Inc., 510(k) Summary, prepared Nov. 28, 2010, for Affirm Breast Biopsy Guidance System Special 510(k) Premarket Notification, 5 pages.
- Hologic, Inc., 510(k) Summary, prepared Aug. 14, 2012, for Affirm Breast Biopsy Guidance System Special 510(k) Premarket Notification, 5 pages.
- ICRP Publication 60: 1990 Recommendations of the International Commission on Radiological Protection, 12 pages.
- Jochelson, M., et al., “Bilateral Dual Energy contrast-enhanced digital mammography: Initial Experience”, *RSNA 2010, 96th Scientific Assembly and Scientific Meeting*, 1 page.
- Jong, RA, et al., Contrast-enhanced digital mammography: initial clinical experience. *Radiology* 2003; 228:842-850.
- Kopans, et al. Will tomosynthesis replace conventional mammography? *Plenary Session SFN08: RSNA 2005.*
- Lehman, CD, et al. MRI evaluation of the contralateral breast in women with recently diagnosed breast cancer. *N Engl J Med* 2007; 356:1295-1303.
- Lewin, JM, et al., Dual-energy contrast-enhanced digital subtraction mammography: feasibility. *Radiology* 2003; 229:261-268.
- Lindfors, KK, et al., Dedicated breast CT: initial clinical experience. *Radiology* 2008; 246(3): 725-733.
- Niklason, L., et al., Digital tomosynthesis in breast imaging. *Radiology.* Nov. 1997; 205(2):399-406.
- Poplack, SP, et al., Digital breast tomosynthesis: initial experience in 98 women with abnormal digital screening mammography. *AJR Am J Roentgenology* Sep. 2007 189(3):616-23.
- Prionas, ND, et al., Contrast-enhanced dedicated breast CT: initial clinical experience. *Radiology.* Sep. 2010 256(3):714-723.
- Rafferty, E et al., “Assessing Radiologist Performance Using Combined Full-Field Digital Mammography and Breast Tomosynthesis Versus Full-Field Digital Mammography Alone: Results” . . . presented at 2007 Radiological Society of North America meeting, Chicago IL.
- Shrading, Simone et al., “Digital Breast Tomosynthesis-guided Vacuum-assisted Breast Biopsy: Initial Experiences and Comparison with Prone Stereotactic Vacuum-assisted Biopsy”, the Department of Diagnostic and Interventional Radiology, Univ. of Aachen, Germany, published Nov. 12, 2014, 10 pgs.
- Smith, A., “Full field breast tomosynthesis”, *Radiol Manage.* Sep.-Oct. 2005; 27(5):25-31.
- Weidner N, et al., “Tumor angiogenesis and metastasis: correlation in invasive breast carcinoma”, *New England Journal of Medicine* 1991; 324:1-8.
- Weidner, N, “The importance of tumor angiogenesis: the evidence continues to grow”, *AM J Clin Pathol.* Nov. 2004 122(5):696-703.
- “Supersonic to feature Aixplorer Ultimate at ECR”, *AuntiMinnie.com*, 3 pages (Feb. 2018).
- Bushberg, Jerrold et al., “The Essential Physics of Medical Imaging”, 3rd ed., In: “The Essential Physics of Medical Imaging, Third Edition”, Dec. 28, 2011, Lippincott & Wilkins, Philadelphia, PA, USA, XP05579051, pp. 270-272.
- Dromain, Clarisse et al., “Dual-energy contrast-enhanced digital mammography: initial clinical results”, *European Radiology*, Sep. 14, 2010, vol. 21, pp. 565-574.
- Reynolds, April, “Stereotactic Breast Biopsy: A Review”, *Radiologic Technology*, vol. 80, No. 5, Jun. 1, 2009, pp. 447M-464M, XP055790574.
- Fischer Imaging Corp, *Mammotest Plus manual on minimally invasive breast biopsy system*, 2002, 8 pages. (Reference labeled D13 in 01 Opposition).
- Fischer Imaging Corporation, *Installation Manual, MammoTest Family of Breast Biopsy Systems*, 86683G, 86684G, P-55957-IM, Issue 1, Revision 3, Jul. 2005, 98 pages. (Reference labeled D12 in 01 Opposition).
- Fischer Imaging Corporation, *Operator Manual, MammoTest Family of Breast Biopsy Systems*, 86683G, 86684G, P-55956-OM, Issue 1, Revision 6, Sep. 2005, 258 pages. (Reference labeled D11 in 01 Opposition).
- Koehli, Ossi R., “Available Stereotactic Systems for Breast Biopsy”, Renzo Brun del Re (Ed.), *Minimally Invasive Breast Biopsies, Recent Results in Cancer Research* 173:105-113; Springer-Verlag, 2009. (Reference labeled D10 in 01 Opposition).
- E. Shaw de Paredes et al., “Interventional Breast Procedure”, published Sep./Oct. 1998 in *Curr Probl Diagn Radiol*, pp. 138-184. (D15 in oppo).
- Burbank, Fred, “Stereotactic Breast Biopsy: Its History, Its Present, and Its Future”, published in 1996 at the Southeastern Surgical Congress, 24 pages.
- Georgian-Smith, Dianne, et al., “Stereotactic Biopsy of the Breast Using an Upright Unit, a Vacuum-Suction Needle, and a Lateral Arm-Support System”, 2001, at the American Roentgen Ray Society meeting, 8 pages. (Reference labeled D10 in 0020 Opposition).
- Samani, A. et al., “Biomechanical 3-D Finite Element Modeling of the Human Breast Using MRI Data”, 2001, *IEEE Transactions on Medical Imaging*, vol. 20, No. 4, pp. 271-279.
- Yin, H.M., et al., “Image Parser: a tool for finite element generation from three-dimensional medical images”, *BioMedical Engineering Online*. 3:31, pp. 1-9, Oct. 1, 2004.
- Sakic et al., “Mammogram synthesis using a 3D simulation. I. breast tissue model and image acquisition simulation” *Medical Physics*. 29, pp. 2131-2139 (2002).
- Wodajo, Felasfa, MD, “Now Playing: Radiology Images from Your Hospital PACS on your iPad,” Mar. 17, 2010; web site: <http://www.imedicalapps.com/2010/03/now-playing-radiology-images-from-your-hospital-pacs-on-your-ipad/>, accessed on Nov. 3, 2011 (3 pages).
- Pathmanathan et al., “Predicting tumour location by simulating large deformations of the breast using a 3D finite element model and nonlinear elasticity”, *Medical Image Computing and Computer-Assisted Intervention*, pp. 217-224, vol. 3217 (2004).
- Al Sallab et al., “Self Learning Machines Using Deep Networks”, *Soft Computing and Pattern Recognition (SoCPaR)*, 2011 Int’l. Conference of IEEE, Oct. 14, 2011, pp. 21-26.
- Ghiassi, M. et al., “A Dynamic Architecture for Artificial Networks”, *Neurocomputing*, vol. 63, Aug. 20, 2004, pp. 397-413.
- Chan, Heang-Ping et al., “ROC Study of the effect of stereoscopic imaging on assessment of breast lesions,” *Medical Physics*, vol. 32, No. 4, Apr. 2005, 1001-1009.
- Lilja, Mikko, “Fast and accurate voxel projection technique in free-form cone-beam geometry with application to algebraic reconstruction,” *Applies Sciences on Biomedical and Communication Technologies*, 2008, Isabel ’08, first international symposium on, IEEE, Piscataway, NJ, Oct. 25, 2008.

(56)

References Cited

OTHER PUBLICATIONS

Pediconi, "Color-coded automated signal intensity-curve for detection and characterization of breast lesions: Preliminary evaluation of new software for MR-based breast imaging," International Congress Series 1281 (2005) 1081-1086.

eFilm Mobile HD by Merge Healthcare, web site: <http://itunes.apple.com/bw/app/efilm-mobile-hd/id405261243?mt=8>, accessed on Nov. 3, 2011 (2 pages).

eFilm Solutions, eFilm Workstation (tm) 3.4, website: <http://estore.merge.com/na/estore/content.aspx?productID=405>, accessed on Nov. 3, 2011 (2 pages).

Van Schie, Guido, et al., "Mass detection in reconstructed digital breast tomosynthesis volumes with a computer-aided detection system trained on 2D mammograms", *Med. Phys.* 40(4), Apr. 2013, 41902-1-41902-11.

Van Schie, Guido, et al., "Generating Synthetic Mammograms from Reconstructed Tomosynthesis Volumes", *IEEE Transactions on Medical Imaging*, vol. 32, No. 12, Dec. 2013, 2322-2331.

Diekmann, Felix et al., "Thick Slices from Tomosynthesis Data Sets: Phantom Study for the Evaluation of Different Algorithms", *Journal of Digital Imaging*, Springer, vol. 22, No. 5, Oct. 23, 2007, pp. 519-526.

Conner, Peter, "Breast Response to Menopausal Hormone Therapy—Aspects on Proliferation, apoptosis and Mammographic Density", 2007 *Annals of Medicine*, 39;1, 28-41.

Glick, Stephen J., "Breast CT", *Annual Rev. Biomed. Eng.*, 2007, 9;501-26.

Metheany, Kathrine G. et al., "Characterizing anatomical variability in breast CT images", Oct. 2008, *Med. Phys.* 35 (10); 4685-4694.

Dromain, Clarisse, et al., "Evaluation of tumor angiogenesis of breast carcinoma using contrast-enhanced digital mammography", *AJR*: 187, Nov. 2006, 16 pages.

Zhao, Bo, et al., "Imaging performance of an amorphous selenium digital mammography detector in a breast tomosynthesis system", May 2008, *Med. Phys* 35(5); 1978-1987.

Mahesh, Mahadevappa, "AAPM/RSNA Physics Tutorial for Residents—Digital Mammography: An Overview", Nov.-Dec. 2004, vol. 24, No. 6, 1747-1760.

Zhang, Yiheng et al., "A comparative study of limited-angle cone-beam reconstruction methods for breast tomosynthesis", *Med Phys.*, Oct. 2006, 33(10): 3781-3795.

Sechopoulos, et al., "Glandular radiation dose in tomosynthesis of the breast using tungsten targets", *Journal of Applied Clinical Medical Physics*, vol. 8, No. 4, Fall 2008, 161-171.

Wen, Junhai et al., "A study on truncated cone-beam sampling strategies for 3D mammography", 2004, *IEEE*, 3200-3204.

Ijaz, Umer Zeeshan, et al., "Mammography phantom studies using 3D electrical impedance tomography with numerical forward solver", *Frontiers in the Convergence of Bioscience and Information Technologies* 2007, 379-383.

Kao, Tzu-Jen et al., "Regional admittivity spectra with tomosynthesis images for breast cancer detection", *Proc. Of the 29th Annual Int'l. Conf. of the IEEE EMBS*, Aug. 23-26, 2007, 4142-4145.

Varjonen, Mari, "Three-Dimensional Digital Breast Tomosynthesis in the Early Diagnosis and Detection of Breast Cancer", *IWDM 2006, LNCS 4046*, 152-159.

Taghibakhsh, f. et al., "High dynamic range 2-TFT amplified pixel sensor architecture for digital mammography tomosynthesis", *IET Circuits Devices Syst.*, 2007, 1(10), pp. 87-92.

Chan, Heang-Ping et al., "Computer-aided detection system for breast masses on digital tomosynthesis mammograms: Preliminary Experience", *Radiology*, Dec. 2005, 1075-1080.

* cited by examiner




Figure 1




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 4C




Figure 4D




Figure 5




Figure 6

**SYSTEM AND METHOD FOR
SYNTHESIZING LOW-DIMENSIONAL
IMAGE DATA FROM HIGH-DIMENSIONAL
IMAGE DATA USING AN OBJECT GRID
ENHANCEMENT**

RELATED APPLICATIONS DATA

The present application is a National Phase entry under 35 U.S.C § 371 of International Patent Application No. PCT/US2018/024912, having an international filing date of Mar. 28, 2018, which claims the benefit under 35 U.S.C. § 119 to U.S. Provisional Patent Application Ser. No. 62/479,008, filed Mar. 30, 2017, which is incorporated by reference in its entirety into the present application.

FIELD

The presently disclosed inventions relate to systems and methods for processing and displaying breast tissue images, and in particular to representing high-dimensional (e.g., 3D) structures present in breast tissue image data with a high-dimensional object grid, and then reducing the high-dimensional data to a low-dimensional (e.g., 2D) format version that can be incorporated within a synthesized image to be displayed to a medical professional.

BACKGROUND

Mammography has long been used to screen for breast cancer and other abnormalities. Traditionally, mammograms have been formed on x-ray film. More recently, flat panel digital imagers have been introduced that acquire a mammogram in digital form, and thereby facilitate analysis and storage of the acquired image data, and to also provide other benefits. Further, substantial attention and technological development have been dedicated to obtaining three-dimensional images of the breast using methods such as breast tomosynthesis. In contrast to the 2D images generated by legacy mammography systems, breast tomosynthesis systems construct a 3D image volume from a series of 2D projection images, each projection image obtained at a different angular displacement of an x-ray source relative to the image detector as the x-ray source is scanned over the detector. The constructed 3D image volume is typically presented as a plurality of slices of image data, the slices being mathematically reconstructed on planes typically parallel to the imaging detector. The reconstructed tomosynthesis slices reduce or eliminate the problems caused by tissue overlap and structure noise present in single slice, two-dimensional mammography imaging, by permitting a user (e.g., a radiologist or other medical professional) to scroll through the image slices to view only the structures in that slice.

Imaging systems such as tomosynthesis systems have recently been developed for breast cancer screening and diagnosis. In particular, Hologic, Inc. (hologic.com) has developed a fused, multimode mammography/tomosynthesis system that acquires one or both types of mammogram and tomosynthesis images, either while the breast remains immobilized or in different compressions of the breast. Other companies have introduced systems that include tomosynthesis imaging; e.g., which do not include the ability to also acquire a mammogram in the same compression.

Examples of systems and methods that leverage existing medical expertise in order to facilitate, optionally, the tran-

sition to tomosynthesis technology are described in U.S. Pat. No. 7,760,924, which is hereby incorporated by reference in its entirety. In particular, U.S. Pat. No. 7,760,924 describes a method of generating a synthesized 2D image, which may optionally be displayed along with tomosynthesis projection or reconstructed images, in order to assist in screening and diagnosis.

A 2D synthesized image is designed to provide a concise representation of the 3D reconstruction slices, including any clinically important and meaningful information, such as abnormal lesions and normal breast structures, while representing in relevant part a traditional 2D image. There are many different types of lesions and breast structures, which may be defined as different types of image objects having different characteristics. For any given image object visible in the 3D volume data, it is important to maintain and enhance the image characteristics (e.g., micro-calcifications, architectural distortions, etc.) as much as possible onto the 2D synthesized image. Further, when representing multiple identified objects on the 2D synthesized image, the synthesized image may appear crowded and visually confusing. Accordingly, there exists a need for more effectively processing, synthesizing and displaying breast image data.

SUMMARY

In one embodiment of the disclosed inventions, a method for processing breast tissue image data includes obtaining image data of a patient's breast tissue; processing the image data to generate a high-dimensional grid depicting one or more high-dimensional objects in the patient's breast tissue; determining a probability or confidence of each of the one or more high-dimensional objects depicted in the high-dimensional grid; and generating a lower-dimensional format version of the one or more high-dimensional objects for display in a synthesized image of the patient's breast tissue.

These and other aspects and embodiments of the disclosed inventions are described in more detail below, in conjunction with the accompanying figures.

BRIEF DESCRIPTION OF THE FIGURES

The drawings illustrate the design and utility of embodiments of the disclosed inventions, in which similar elements are referred to by common reference numerals. These drawings are not necessarily drawn to scale. In order to better appreciate how the above-recited and other advantages and objects are obtained, a more particular description of the embodiments will be rendered, which are illustrated in the accompanying drawings. These drawings depict only typical embodiments of the disclosed inventions and are not therefore to be considered limiting of its scope.

FIG. 1 is a block diagram illustrating the flow of data through an exemplary breast image acquisition and processing system in accordance with embodiments of the disclosed inventions;

FIG. 2 is a block diagram illustrating the flow of data through a 2D synthesizer employing a 3D object grid and various modules that reduce objects of the grid to a 2D format for display;

FIG. 3 illustrates a first synthesized image formed from the 3D object grid of FIG. 2 without manipulating overlapping objects, and a second synthesized image formed from the same 3D object grid, but with manipulation of overlapping objects;

FIG. 4A-4D illustrate exemplary techniques for combining objects onto one or more 2D synthesized images;

FIG. 5 illustrates an exemplary flow diagram depicting combining objects from a 3D object grid onto a 2D synthesized image; and

FIG. 6 illustrates an exemplary flow diagram depicting generating one or more 2D synthesized images using a 3D object grid.

DETAILED DESCRIPTION OF THE ILLUSTRATED EMBODIMENTS

All numeric values are herein assumed to be modified by the terms “about” or “approximately,” whether or not explicitly indicated, wherein the terms “about” and “approximately” generally refer to a range of numbers that one of skill in the art would consider equivalent to the recited value (i.e., having the same function or result). In some instances, the terms “about” and “approximately” may include numbers that are rounded to the nearest significant figure. The recitation of numerical ranges by endpoints includes all numbers within that range (e.g., 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, and 5).

As used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the content clearly dictates otherwise. As used in this specification and the appended claims, the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise. In describing the depicted embodiments of the disclosed inventions illustrated in the accompanying figures, specific terminology is employed for the sake of clarity and ease of description. However, the disclosure of this patent specification is not intended to be limited to the specific terminology so selected, and it is to be understood that each specific element includes all technical equivalents that operate in a similar manner. It is to be further understood that the various elements and/or features of different illustrative embodiments may be combined with each other and/or substituted for each other wherever possible within the scope of this disclosure and the appended claims.

Various embodiments of the disclosed inventions are described hereinafter with reference to the figures. It should be noted that the figures are not drawn to scale and that elements of similar structures or functions are represented by like reference numerals throughout the figures. It should also be noted that the figures are only intended to facilitate the description of the embodiments. They are not intended as an exhaustive description of the invention or as a limitation on the scope of the disclosed inventions, which is defined only by the appended claims and their equivalents. In addition, an illustrated embodiment of the disclosed inventions needs not have all the aspects or advantages shown. For example, an aspect or an advantage described in conjunction with a particular embodiment of the disclosed inventions is not necessarily limited to that embodiment and can be practiced in any other embodiments even if not so illustrated.

For the following defined terms and abbreviations, these definitions shall be applied throughout this patent specification and the accompanying claims, unless a different definition is given in the claims or elsewhere in this specification:

An “acquired image” refers to an image generated while visualizing a patient’s tissue. Acquired images can be generated by radiation from a radiation source impacting on a radiation detector disposed on opposite sides of a patient’s tissue, as in a conventional mammogram.

A “reconstructed image” refers to an image generated from data derived from a plurality of acquired images. A reconstructed image simulates an acquired image not included in the plurality of acquired images.

A “synthesized image” refers to an artificial image generated from data derived from a plurality of acquired and/or reconstructed images. A synthesized image includes elements (e.g., objects and regions) from the acquired and/or reconstructed images, but does not necessarily correspond to an image that can be acquired during visualization. Synthesized images are constructed analysis tools.

An “Mp” image is a conventional mammogram or contrast enhanced mammogram, which are two-dimensional (2D) projection images of a breast, and encompasses both a digital image as acquired by a flat panel detector or another imaging device, and the image after conventional processing to prepare it for display (e.g., to a health professional), storage (e.g., in the PACS system of a hospital), and/or other use.

A “Tp” image is an image that is similarly two-dimensional (2D), but is acquired at a respective tomosynthesis angle between the breast and the origin of the imaging x-rays (typically the focal spot of an x-ray tube), and encompasses the image as acquired, as well as the image data after being processed for display, storage, and/or other use.

A “Tr” image is a type (or subset) of a reconstructed image that is reconstructed from tomosynthesis projection images Tp, for example, in the manner described in one or more of U.S. Pat. Nos. 7,577,282, 7,606,801, 7,760,924, and 8,571,289, the disclosures of which are fully incorporated by reference herein in their entirety, wherein a Tr image represents a slice of the breast as it would appear in a projection x-ray image of that slice at any desired angle, not only at an angle used for acquiring Tp or Mp images.

An “Ms” image is a type (or subset) of a synthesized image, in particular, a synthesized 2D projection image, which simulates mammography images, such as craniocaudal (CC) or mediolateral oblique (MLO) images, and is constructed using tomosynthesis projection images Tp, tomosynthesis reconstructed images Tr, or a combination thereof. Ms images may be provided for display to a health professional or for storage in the PACS system of a hospital or another institution. Examples of methods that may be used to generate Ms images are described in the above-incorporated U.S. Pat. Nos. 7,760,924 and 8,571,289.

It should be appreciated that Tp, Tr, Ms and Mp image data encompasses information, in whatever form, that is sufficient to describe the respective image for display, further processing, or storage. The respective Mp, Ms, Tp and Tr images are typically provided in digital form prior to being displayed, with each image being defined by information that identifies the properties of each pixel in a two-dimensional array of pixels. The pixel values typically relate to respective measured, estimated, or computed responses to X-rays of corresponding volumes in the breast, i.e., voxels or columns of tissue. In a preferred embodiment, the geometry of the tomosynthesis images (Tr and Tp) and mammography images (Ms and Mp) are matched to a common coordinate system, as described in U.S. Pat. No. 7,702,142. Unless otherwise specified, such coordinate system matching is assumed to be implemented with respect to the embodiments described in the ensuing detailed description of this patent specification.

The terms “generating an image” and “transmitting an image” respectively refer to generating and transmitting

information that is sufficient to describe the image for display. The generated and transmitted information is typically digital information.

In order to ensure that a synthesized 2D image displayed to an end-user (e.g., an Ms image) includes the most clinically relevant information, it is necessary to detect and identify three-dimensional (3D) objects, such as malignant breast mass, tumors, etc., within the breast tissue. In accordance with the inventions disclosed and described herein, this information may be used to create a high-dimensional grid, e.g., a 3D grid, that helps create a more accurate and enhanced rendering of the most important features in the synthesized 2D image. The high-dimensional object grid may then be used to collapse the most clinically-significant information pertaining to the identified objects to a 2D format onto one or more synthesized 2D images. Various data reduction techniques may be applied to the identified 3D objects to ensure that the most clinically-significant objects are emphasized, and less significant objects are omitted and/or de-emphasized. Additionally, or alternatively, data reduction techniques are applied to ensure that significant features of a 3D object are enhanced, while less significant features of the 3D object are de-emphasized, especially when two objects are competing for display and prominence on the one or more 2D synthesized images. Thus, as disclosed and described herein, a 3D object grid is utilized, i.e., as a component of an algorithm, for reducing high-dimensional data (e.g., 3D tomosynthesis image data) to low-dimensional data (e.g. a 2D synthesized image).

FIG. 1 illustrates the flow of data in an exemplary image generation and display system **100**, which incorporates each of synthesized image generation, object identification, and display technology. It should be understood that, while FIG. 1 illustrates a particular embodiment of a flow diagram with certain processes taking place in a particular serial order or in parallel, the claims and various other embodiments described herein are not limited to the performance of the image processing steps in any particular order, unless so specified.

More particularly, the image generation and display system **100** includes an image acquisition system **101** that acquires tomosynthesis image data for generating Tp images of a patient's breasts, using the respective three-dimensional and/or tomosynthesis acquisition methods of any of the currently available systems. If the acquisition system is a combined tomosynthesis/mammography system, Mp images may also be generated. Some dedicated tomosynthesis systems or combined tomosynthesis/mammography systems may be adapted to accept and store legacy mammogram images, (indicated by a dashed line and legend "Mp_{legacy}" in FIG. 1) in a storage device **102**, which is preferably a DICOM-compliant Picture Archiving and Communication System (PACS) storage device. Following acquisition, the tomosynthesis projection images Tp may also be transmitted to the storage device **102** (as shown in FIG. 1). The storage device **102** may further store a library of known 3D objects that may be used to identify significant 3D image patterns to the end-user. In other embodiments, a separate dedicated storage device (not shown) may be used to store the library of known 3D objects with which to identify 3D image patterns or objects.

The Tp images are transmitted from either the acquisition system **101**, or from the storage device **102**, or both, to a computer system configured as a reconstruction engine **103** that reconstructs the Tp images into reconstructed image "slices" Tr, representing breast slices of selected thickness

and at selected orientations, as disclosed in the above-incorporated patents and applications.

Mode filters **107** are disposed between image acquisition and image display. The filters **107** may additionally include customized filters for each type of image (i.e., Tp, Mp, and Tr images) arranged to identify and highlight certain aspects of the respective image types. In this manner, each imaging mode can be tuned or configured in an optimal way for a specific purpose. For example, filters programmed for recognizing objects across various 2D image slices may be applied in order to detect image patterns that may belong to a particular high-dimensional object. The tuning or configuration may be automatic, based on the type of the image, or may be defined by manual input, for example through a user interface coupled to a display. In the illustrated embodiment of FIG. 1, the mode filters **107** are selected to highlight particular characteristics of the images that are best displayed in respective imaging modes, for example, geared towards identifying objects, highlighting masses or calcifications, identifying certain image patterns that may be constructed into a 3D object, or for creating 2D synthesized images (described below). Although FIG. 1 illustrates only one mode filter **107**, it should be appreciated that any number of mode filters may be utilized in order to identify structures of interest in the breast tissue.

The imaging and display system **100** further includes a 2D image synthesizer **104** that operates substantially in parallel with the reconstruction engine **103** for generating 2D synthesized images using a combination of one or more Tp, Mp, and/or Tr images. The 2D image synthesizer **104** consumes a set of input images (e.g., Mp, Tr and/or Tp images), determines a set of most relevant features from each of the input images, and outputs one or more synthesized 2D images. The synthesized 2D image represents a consolidated synthesized image that condenses significant portions of various slices onto one image. This provides an end-user (e.g., medical personnel, radiologist, etc.) with the most clinically-relevant image data in an efficient manner, and reduces time spent on other images that may not have significant data.

One type of relevant image data to highlight in the synthesized 2D images would be relevant objects found across one or more Mp, Tr and/or Tp images. Rather than simply assessing image patterns of interest in each of the 2D image slices, it may be helpful to determine whether any of the 2D image patterns of interest belong to a larger high-dimensional structure, and if so, to combine the identified 2D image patterns into a higher-dimensional structure. This approach has several advantages, but in particular, by identifying high-dimensional structures across various slices/depths of the breast tissue, the end-user may be better informed as to the presence of a potentially significant structure that may not be easily visible in various 2D slices of the breast.

Further, instead of identifying similar image patterns in two 2D slices (that are perhaps adjacent to each other), and determining whether or not to highlight image data from one or both of the 2D slices, identifying both image patterns as belonging to the same high-dimensional structure may allow the system to make a more accurate assessment pertaining to the nature of the structure, and consequently provide significantly more valuable information to the end-user. Also, by identifying the high-dimensional structure, the structure can be more accurately depicted on the synthesized 2D image. Yet another advantage of identifying high-dimensional structures within the various captured 2D slices of the breast tissue relates to identifying a possible size/scope of

the identified higher-dimensional structure. For example, once a structure has been identified, previously unremarkable image patterns that are somewhat proximate to the high-dimensional structure may now be identified as belonging to the same structure. This may provide the end-user with an indication that the high-dimensional structure is increasing in size/scope.

To this end, the 2D image synthesizer **104** generates high-dimensional object grids **120** (e.g., 3D object grids) comprising one or more high-dimensional structures (e.g., 3D objects) present in the patient's breast tissue. Several techniques may be used to construct 3D object grids **120** that identify various objects in the breast tissue. It should be appreciated that this disclosure is not limited to 3D objects and/or structures, and may refer to even higher-dimensional structures, but for simplicity, the remaining disclosure will refer to 3D objects populated in a 3D object grid **120**.

In one or more embodiments, the 3D object grid **120** is in the form of a 3D (volumetric) coordinate space representing a patient's breast mass, and identifies a location, identity, size, scope, and/or other characteristics of any objects or structures found in the breast mass. Examples of such objects or structures include calcifications, spiculated lesions, benign tumors, irregular masses, dense objects, etc.

In one or more embodiments, the end-user (e.g., a medical professional such as a radiologist) can access and interact with the 3D object grid **120**. In other embodiments, the 3D object grid **120** is solely used by the system processor for constructing synthesized 2D images, and the end-user may not be aware of, or have access to, the 3D object grid **120**.

In accordance with the disclosed embodiments, the 2D image synthesizer **104** also includes a data reduction module **122** configured to reduce the high-dimensional data populated in the 3D object grid **120** to a lower-dimensional format suitable for representation in a 2D synthesized image. The data reduction module **122** evaluates the various objects of the 3D object grid **120**, and determines what objects (or what portions of objects) should be enhanced or emphasized in a final 2D synthesized image to be displayed to the end-user. For example, a clinically significant object and a routine background breast tissue object may have regions of overlap, in which case the data reduction module **122** is preferably configured to de-emphasize portions of the background breast tissue in order to highlight the clinically significant object. Further details on various data reduction techniques that may be employed by the data reduction module **122** are described below.

The synthesized 2D images may be viewed at a display system **105**. The reconstruction engine **103** and 2D image synthesizer **104** are preferably connected to a display system **105** via a fast transmission link. The display system **105** may be part of a standard acquisition workstation (e.g., of acquisition system **101**), or of a standard (multi-display) review station (not shown) that is physically remote from the acquisition system **101**. In some embodiments, a display connected via a communication network may be used, for example, a display of a personal computer or of a so-called tablet, smart phone or other hand-held device. In any event, the display **105** of the system is preferably able to display respective Ms, Mp, Tr, and/or Tp images concurrently, e.g., in separate side-by-side monitors of a review workstation, although the invention may still be implemented with a single display monitor, by toggling between images.

Thus, the imaging and display system **100**, which is described as for purposes of illustration and not limitation, is capable of receiving and selectively displaying tomosynthesis projection images Tp, tomosynthesis reconstruction

images Tr, synthesized mammogram images Ms, and/or mammogram (including contrast mammogram) images Mp, or any one or sub combination of these image types. The system **100** employs software to convert (i.e., reconstruct) tomosynthesis images Tp into images Tr, software for synthesizing mammogram images Ms, software for decomposing 3D objects, software for creating feature maps and object maps. An object of interest or feature in a source image may be considered a 'most relevant' feature for inclusion in a 2D synthesized image based upon the application of the object maps along with one or more algorithms and/or heuristics, wherein the algorithms assign numerical values, weights or thresholds, to pixels or regions of the respective source images based upon identified/detected objects and features of interest within the respective region or between features. The objects and features of interest may include, for example, spiculated lesions, calcifications, and the like.

FIG. 2 illustrates the 2D image synthesizer **104** in further detail. As discussed above, various image slices **218** of a tomosynthesis data set (or "stack") **202** (e.g., filtered and/or unfiltered Mp, Tr and/or Tp images of a patient's breast tissue) are input into the 2D image synthesizer **104**, and then processed to determine portions of the images to highlight in a synthesized 2D image that will be displayed on the display **105**. The image slices **218** may be consecutively-captured cross-sections of a patient's breast tissue. Or, the image slices **218** may be cross-sectional images of the patient's breast tissue captured at known intervals. The tomosynthesis image stack **202** comprising the image slices **218** may be forwarded to the 2D image synthesizer **104**, which evaluates each of the source images in order to (1) identify various types of objects (Tr) for possible inclusion in one or more 2D synthesized images, and/or (2) identify respective pixel regions in the images that contain the identified objects.

As shown in the illustrated embodiment, the tomosynthesis stack **202** comprises a plurality of images **218** taken at various depths/cross-sections of the patient's breast tissue. Some of the images **218** in the tomosynthesis stack **202** comprise 2D image patterns. Thus, the tomosynthesis stack **202** comprises a large number of input images containing various image patterns within the images of the stack.

For example, while the tomosynthesis stack **202** may comprise one hundred images **218** captured at various depths/cross sections of the patient's breast tissue, only a few of the images **218** may include any information of significance. Also, it should be noted that the tomosynthesis stack **202** contains 2D image patterns when viewed at differing z-dimension (depth) locations of the otherwise same x, y locations in the image slices **218**, but it may be difficult to determine the 3D structures based only on the various individual images, each representing a finite cross-sectional image of the breast tissue. However, the tomosynthesis stack **202** may be effectively utilized to create the 3D object grid **120**. In any event, for purposes of this patent specification, it is assumed that the 3D object grid **120** is constructed by any means, including but not limited to being created from the tomosynthesis stack **202**.

The 3D object grid **120** may be considered a 3D volumetric coordinate space representing a patient's breast mass. Rather than depicting 2D image patterns at various image slices, the 3D object grid **120** preferably depicts any identified 3D objects in the entire mass (or portion thereof) that represents the patient's breast tissue. The 3D object grid **120** provides fuller detail regarding various objects in the breast mass as compared to the tomosynthesis stack **202**. For example, the 3D object grid **120** may use simulation techniques to infer a shape of the 3D object, even though an

image slice may not have necessarily been reconstructed at every cross-sectional depth covering the respective 3D object.

The 3D object grid **120** may comprise one or more objects, as shown in the illustrated embodiment. It should be appreciated that these objects may be predefined objects that the system has been trained to identify. However, even in healthy breast tissue that does not necessarily comprise any abnormal objects or structures, the target object recognition/enhancement modules may identify a breast background object. For example, all breast linear tissue and density tissue structures can be displayed as the breast background object. In other embodiments, “healthy” objects such as spherical shapes, oval shapes, etc., may simply be identified through one or more target object recognition/enhancement modules **210**. These identified 3D objects may then be displayed on the 2D synthesized image **206**; of course, out of all identified 2D objects, more clinically-significant objects may be prioritized and/or enhanced when displaying the respective object on the 2D synthesized image, as will be discussed in further detail below.

In one or more embodiments, the 2D synthesizer **104** utilizes both the tomosynthesis image stack **202** along with the created 3D object grid **120** in order to merge the relevant features into one or more 2D synthesized images **206**. As shown in the 2D synthesized image, the 3D objects identified in the 3D object grid **120** are collapsed into a 2D format, but provide more detail when compared to individual image slices of the tomosynthesis image stack **202**. Further, although several objects, as shown in the tomosynthesis image stack **202** overlap in the z direction, identifying them as separate 3D objects allows the system to depict both objects clearly and efficiently. Simply utilizing legacy image recognizing techniques on the tomosynthesis image stack **202** may or may not necessarily provide such an accurate synthesized 2D image **206**. To explain, if there is overlap in the z direction of two structures, the two structures are essentially competing with each other for display on the 2D synthesized image **206**. Thus, important aspects of both structures may be compromised. Or, only one of the two structures may be highlighted at all in the 2D synthesized image **206**. Or, in yet another scenario, the 2D synthesized image may depict both structures as one amorphous structure such that an important structure goes entirely undetectable for the end-user.

It will be appreciated that identifying 3D objects as separate objects with predefined types in the 3D object grid **120** allows the system to depict the structures more accurately on the 2D synthesized image **206**, and allows for various objects to be depicted simultaneously, even if there is an overlap of various objects in the coordinate space. Thus, utilizing the 3D object grid **120** has many advantages for producing a more accurate and visually-effective 2D synthesized image **206**.

In one or more embodiments, data from the tomosynthesis stack **202** and the 3D object grid **120** are processed by one or more modules to produce the 2D synthesized image **206**. More particularly, an object combination module **210** may be configured to identify the various objects of the 3D object grid **120**, and determine a most optimal method to collapse all the objects on a 2D plane/format. For example, the object combination module **210** may determine x and y coordinates for the plurality of objects and determine whether there are overlaps between multiple objects to be displayed on the 2D synthesized image **206**. In some embodiments, the object combination module **210** may further be configured to determine which of the identified objects should be dis-

played on the 2D synthesized image **206**. This may be achieved through a training (or a “learning library” type) database **216** that stores an identity of various objects and associated weights of respective objects. The training database becomes more knowledgeable with the processing of each new patent breast image data, as the system derives 3D object models and (subsequently) detection mechanisms from this database, which will grow to include various samples of the same types of objects.

After the 3D objects are detected, then the next step is to utilize this same knowledge in synthesizing the 2D image. Since there may be many different types (or categories) of 3D objects, the weighting mechanism helps to combine the objects in the synthesis/data reduction process. For example, a dense spherical object may be weighed higher than a calcification (weighed 0.95 and 0.6 respectively in the illustrated embodiment), such that the dense spherical object may be enhanced to a greater degree as compared to a calcification. If the weight of an object is close to zero, the object combination module **210** may determine that the object need not be displayed at all, in some embodiments.

In one or more embodiments, an image details synthesis module **212** may be configured to determine what 3D objects or what areas within a 3D object should be emphasized in the 2D synthesized image **206**. For example, if there is an overlap between two objects, the image details synthesis module **212** may emphasize portions of both objects, and de-emphasize other portions of both objects such that both objects are clearly viewable on the 2D synthesized image. By manipulating aspects of both objects, the end-user may be able to identify both objects clearly. It should be appreciated that without this manipulation, both objects may simply be overlaid on top of each other, such that one object may simply be masked out and missed by the end-user.

For example, the 3D object grid **120** may include a calcification area and a spiculated lesion that overlap in the z direction. Without any specially designed image synthesis, a collapsed 2D format of the spiculated lesion and a collapsed 2D format of the calcification would be displayed on top of each other. Assuming the spiculated mass is larger, the spiculated mass may envelop the calcification entirely such that it is not visible to the end-user. Instead, the image details synthesis module **212** may emphasize the outline of the center portion of the spiculated mass, while deemphasizing the middle portion of the spiculated mass such that the calcification area is visible. This image manipulation allows the end-user a clearer picture of significant objects on the 2D synthesized image **206**. FIG. 3, described below, illustrates this system feature in further detail.

In some embodiments, the image details synthesis module **212** may comprise several algorithms and/or heuristics that are programmed with rules to determine what parts of an object to emphasize/de-emphasize based on the object database **216**. For example, each object in the database **216** may correspond to metadata that defines most prominent and least-prominent features of the respective object. This metadata may be used by the various algorithms to determine which objects and/or which parts of objects to emphasize in the 2D synthesized images **206**. By way of another example, a difference in weight between two overlapping objects may be calculated in order to determine whether both objects should be displayed. If the difference in weight is smaller than a predetermined threshold value, both objects may be displayed, but the assigned weight may be used to determine which of the two objects to emphasize over the other. However, if the difference in weight is larger than the

predetermined threshold value, only the object corresponding to the higher weight may be displayed at all. For example, if a dense spherical mass and calcification area are competing for display (difference of 0.35 in weight as per the illustrated embodiment) and the threshold value is set at 0.4, both objects may be displayed, but the spiculated mass (or parts of the spiculated mass) may be highlighted relative to the calcification area. However, if the spiculated mass and a benign semi-spherical mass are overlapping (difference of 0.65 in weight as per the illustrated embodiment), only the dense spherical mass may be displayed at all. Other rules may be defined to allow the system to modify the objects or portions thereof.

As noted above (and in FIG. 1), the 2D image synthesizer 104 further includes a data reduction engine 122 configured to receive the data input from the respective image details synthesis module 212 and object combination module 210, and to reduce any 3D objects identified therein into a low level 2D format that may be inserted into the 2D synthesized image 206. In particular, and as described in further detail herein, the data reduction engine 122 accurately reduces the identified high-dimensional object of the 3D object grid 120 to a 2D format based on input received from the image details synthesis module 212, the database 216 and the object combination module 210.

FIG. 3 depicts an example of how the 3D object grid may be utilized to generate the 2D synthesized images. In the illustrated embodiment, 3D object grid 304 includes at least two 3D objects: a spiculated mass 310, and a calcification area 312. When consulting a database, the object combination module may determine that both objects are important to display in the 2D synthesized image, the spiculated mass 310 being more significant than the calcification. However, since both 3D objects 310 and 312 overlap in the z direction, the images may have to be manipulated such that both objects are still optimally displayed on the 2D synthesized image. In particular, the 2D synthesized image 306 displays a synthesized image that does not use any image manipulation techniques described in this disclosure. As shown in 2D synthesized image 306, both 3D objects 310 and 312 are competing to be displayed, and neither object is displayed very clearly. More specifically, the calcification 312 is barely visible in the 2D synthesized image 306.

By contrast, referring to 2D synthesized image 308, the techniques described with respect to FIG. 2 are utilized in order to determine what parts of the respective 3D object should be emphasized and de-emphasized such that both objects are clearly discernible in the 2D synthesized image. More particularly, although spiculated mass 310 is more significant than the calcification 312, the center portion of the spiculated mass 310 is slightly de-emphasized such that the calcification area is clearly visible. Similarly, it may be determined that the linear lines radiating from the center portion should be emphasized such that the end-user understands a size or scope of the spiculated mass. In light of the modified image corresponding to the spiculated mass 310, the calcification 312 is now visible even though both objects overlap. Thus, as shown in FIG. 3, the 2D synthesized image 308 provides more details about both 3D objects 310 and 312 when compared to 2D synthesized image 306.

FIGS. 4A-4D depict exemplary embodiments of displaying various objects of the 3D object grid, while preserving clinically-significant information. In particular, the collapsing of a 3D object into the 2D synthesized image may be achieved by the respective object combination module,

image synthesis module and data reduction module that work together to display as much clinically-significant information as possible.

FIG. 4A shows an example embodiment of an intra-object combination. Intra-object combination may refer to techniques used to represent a single object (that is captured on multiple Tr image slices 404) onto the 2D synthesized image. More particularly, an identified 3D object may appear in many consecutive Tr image slices as 408a, 408b and 408c. In theory these image patterns compete with each other for representation on the 2D synthesized image. Thus, an intra-object combination requires recognizing that all the images slices belong to the same 3D object, and only showing relevant information pertaining to the 3D object on the 2D synthesized image 406. Notably, as shown in FIG. 4A, the system may determine that all the image patterns 408a, 408b and 408c from the Tr stack 404 belong to the same 3D object, and may collapse them together such that they appear as one object 410 in the 2D synthesized image 406. In one or more embodiments, techniques such as averaging, MIP (maximum intensity projection), filtering, etc. may be used for intra-object combination. Intra-object combination techniques aim to preserve the structure of the 3D object without losing valuable information from any of the image slices, while minimizing competing information from multiple image slices that do not provide valuable information and/or visually confuse the end-user.

FIG. 4B illustrates an example embodiment of an object vs background combination. Object vs. background combination may be important for creating a natural-looking 2D synthesized image. The goal of this technique is to maintain useful information from objects together with meaningful background information representative of breast tissue. In the illustrated embodiment, the Tr stack 414 comprises two Tr image slices. The first image slice comprises a background image pattern 412. The second image slice comprises an object or a portion of an object 411. In collapsing information from the Tr stack 414 to the 2D synthesized image 416, some aspects of both image slices are emphasized while other aspects are de-emphasized. For example, in 2D synthesized image 416, the object 411 is preserved, and the background 412 is also rendered, but the middle portion of the background 412 is de-emphasized.

FIG. 4C illustrates an example embodiment of inter-object combination without overlapping. In the illustrated embodiment, Tr stack 424 comprises two Tr image slices. One Tr image slice comprises object 422, and the other Tr image slice comprises object 421. As shown in the illustrated embodiment, these objects do not overlap in the z direction. Thus, when collapsing both objects onto the 2D synthesized image 426, both objects 421 and 422 are represented clearly at their respective x-y locations.

FIG. 4D illustrates an example embodiment of inter-object combination with overlap. This technique may be performed when two or more objects overlap to some degree. The two objects may be of the same type or of different types. In case of an overlap of objects, a hierarchical approach may be used to determine which object should be given precedence over the other. For example, if a higher weight is assigned to a first object, the first object may be emphasized in the 2D synthesized object, while the second object (or portions of the second object) may be de-emphasized. Or, if both objects are equally, or almost equally, important, both objects may be represented equally even if they are overlapping (and portions of both objects may be emphasized/de-emphasized such that both objects are clear in the synthesized 2D image).

In the illustrated embodiment, Tr image stack **434** comprises two Tr image slices. One Tr image slice comprises object **432**, and the other Tr image slice comprises object **431**. As shown in the illustrated embodiment, these objects overlap in the z direction. Thus, when collapsing the objects onto the 2D synthesized image **436**, both objects **431** and **432** are represented, but are shown to overlap. Depending on weights assigned to both objects, one object may be highlighted while the other is de-emphasized. In the illustrated embodiment, both objects are represented somewhat equally, even though it is clear that they represent two separate objects. In other embodiments (not shown), if object **431** is assigned a higher weight/priority, object **431** may be emphasized in the foreground, while object **432** may be relegated to the background. Similarly, other combination techniques may be utilized to optimally represent clinically-significant information to the end-user.

FIG. **5** depicts an exemplary embodiment of collapsing information from a plurality of Tr images into a 2D synthesized image. In particular, a Tr image stack **504** may be used to create a 3D object grid similar to the 3D grid shown in FIG. **2**. The Tr stack **504** illustrates four distinct objects including two calcification areas **510** and **512**, a spiculated mass **514**, and a spherical mass **516**. As discussed in detail above, identifying these four objects as separate and distinct objects allows the system to accurately depict the objects as a whole on the 2D synthesized image **506**. In the illustrated embodiment, the spiculated mass **514** is shown most prominently, while the calcifications and the spherical mass **516** are not as emphasized. This allows an end-user to easily identify the most clinically significant part of the 2D synthesized image without being overwhelmed with objects that are less-significant.

FIG. **6** is a flow diagram **600** provided to illustrate exemplary steps that may be performed in an image merge process carried out in accordance with one embodiment of the disclosed inventions. At step **602**, an image data set is acquired. The image data set may be acquired by a tomosynthesis acquisition system, a combination tomosynthesis/mammography system, or by retrieving pre-existing image data from a storage device, whether locally or remotely located relative to an image display device. At step **604**, a 3D object grid may be constructed by identifying various objects that are present in a 3D coordinate space representative of a patient's breast tissue. At step **606**, the objects of the 3D object grid are recognized, and a relative weight/priority of each of the objects is determined. As discussed above, in some embodiments, all objects of the 3D object grid may be displayed, with some objects emphasized more than others. In other embodiments, only a subset of the recognized objects may be displayed at all, while less-significant objects are omitted.

For example, it may be determined that one object is much more clinically significant as compared to another. Or, it may be determined that two overlapping objects are equally significant. In this case, algorithms aiming to visually depict both objects optimally may be utilized, rather than highlighting one object over another. At step **608**, based on the relative weight/priority of the objects, the 3D objects may be reduced to a 2D format to create the 2D synthesized image. This reduction process may highlight one object over another, in some embodiments. In other embodiment, the reduction process may highlight an outline of an object while de-emphasizing an interior of the object.

In yet another embodiment, the reduction process may emphasize one or more features that are deemed to be significant, while de-emphasizing less significant aspects of

the same object. For example, in the case of a spiculated lesion, it may be important to display the blood supply lines emanating from the center of the spiculated mass, but the center of the spiculated mass, even if dense may be displayed with less emphasis. Any number of such enhancement techniques may be used in the data reduction process. At step **610**, the synthesized 2D image is displayed to the end-user.

Having described exemplary embodiments, it should be appreciated that the examples described above and depicted in the accompanying figures are only illustrative, and that other embodiments and examples also are encompassed within the scope of the appended claims. For example, while the flow diagrams provided in the accompanying figures are illustrative of exemplary steps; the overall image merge process may be achieved in a variety of manners using other data merge methods known in the art. The system block diagrams are similarly representative only, illustrating functional delineations that are not to be viewed as limiting requirements of the disclosed inventions. It will also be apparent to those skilled in the art that various changes and modifications may be made to the depicted and/or described embodiments (e.g., the dimensions of various parts), without departing from the scope of the disclosed inventions, which is to be defined only by the following claims and their equivalents. The specification and drawings are, accordingly, to be regarded in an illustrative rather than restrictive sense.

What is claimed is:

1. A method for processing breast tissue image data, comprising:

processing an image data set including a plurality of image slices comprising image data of a patient's breast tissue to identify two or more clinically significant high-dimensional breast tissue objects in the breast tissue and to determine a clinically significant object type for each of the two or more clinically significant high-dimensional breast tissue objects;

generating a high-dimensional object grid depicting each of the two or more high-dimensional breast tissue objects in the patient's breast tissue found across one or more respective image slices of the image data set;

determining a confidence or probability of each of the two or more high-dimensional breast tissue objects depicted in the high-dimensional object grid;

generating a synthesized image comprising a lower-dimensional format version of each of the two or more high-dimensional breast tissue objects depicted in the high-dimensional object grid based at least in part upon the confidence or probability of each of the two or more high-dimensional breast tissue objects, wherein the confidence or probability is determined based at least in part on a weight assigned to each of the two or more high-dimensional breast tissue objects, and wherein the weights assigned to the two or more high-dimensional breast tissue objects is based on the respective object type,

where generating the synthesized image comprises:

determining whether the two or more high-dimensional breast tissue objects are likely to overlap in the displayed synthesized image; and

if it is determined that the two or more high-dimensional breast tissue objects are likely to overlap in the synthesized image, modifying at least one feature of at least one of the likely overlapping high-dimensional breast tissue objects such that one or more

15

most clinically significant features of at least one of the likely overlapping high-dimensional breast tissue objects is displayed.

2. The method of claim 1, further comprising generating the image data set including the plurality of image slices that collectively depict the patient's breast tissue, wherein the high-dimensional object grid is generated based at least in part on the plurality of image slices.

3. The method of claim 1, wherein generating the synthesized image comprising the lower-dimensional format version of the two or more high-dimensional breast tissue objects includes modifying one or more aspects of at least one of the two or more high-dimensional breast tissue objects based at least in part on its respective determined confidence or probability.

4. The method of claim 1, wherein the lower-dimensional format version of the two or more high-dimensional breast tissue objects is based at least in part on an intra-object combination.

5. The method of claim 1, wherein the lower-dimensional format version of the two or more high-dimensional breast tissue objects is based at least in part on an inter-object combination.

6. The method of claim 1, wherein the lower-dimensional format version of the two or more high-dimensional breast tissue objects is based at least in part on combining a breast tissue object with a background.

7. The method of claim 1, further comprising displaying the synthesized image comprising the lower-dimensional format version of the two or more high-dimensional breast tissue objects.

8. The method of claim 1, wherein the high-dimensional object grid comprises two or more high-dimensional breast tissue objects that depict normal and/or abnormal breast tissue structures.

9. The method of claim 8, wherein the two or more high-dimensional breast tissue objects comprise a plurality of objects that collectively represent an entire breast parenchymal tissue structure.

10. The method of claim 8, wherein each of the two or more high-dimensional breast tissue objects is associated with a respective set of attributes, each attribute representing a characteristic of the breast tissue structure depicted by the respective high-dimensional breast tissue object.

11. The method of claim 10, wherein the set of attributes collectively represent one or more of a location, a size, a shape, and a morphology of the respective breast tissue structure depicted by the high-dimensional breast tissue object.

12. The method of claim 8, wherein the two or more high-dimensional breast tissue objects include two or more high-dimensional breast tissue objects corresponding to a first object type, and two or more high-dimensional breast tissue objects corresponding to a second object type.

13. The method of claim 12, wherein the first object type corresponds to abnormal breast lesions including microcalcifications and masses, and the second object type corresponds to normal breast structures including nipples, pectoral muscles, and breast parenchymal tissues.

14. The method of claim 13, wherein a weight assigned to objects of the first object type is greater than a weight assigned to objects of the second object type.

15. The method of claim 12, further comprising using a pattern recognition method for each of the first and second object types to determine whether one or both object types are present in the image data set that depicts the patient's breast tissue.

16

16. The method of claim 15, wherein the pattern recognition method utilizes one or more machine learning algorithms.

17. The method of claim 1, further comprising, if it is determined that displaying both the two or more high-dimensional breast tissue objects that are likely to overlap in the synthesized image.

18. The method of claim 1, wherein modifying at least one feature of at least one of the overlapping high-dimensional breast tissue objects comprises modifying the object to emphasizing a first portion of the modified high-dimensional breast tissue object relative to a second portion of the same object.

19. A method for processing breast tissue image data, comprising:

processing an image data set including a plurality of image slices comprising image data of a patient's breast tissue to identify two or more clinically significant high-dimensional breast tissue objects in the breast tissue and to determine a clinically significant object type for each of the two or more clinically significant high-dimensional breast tissue objects;

generating a high-dimensional object grid depicting each of the two or more high-dimensional breast tissue objects in the patient's breast tissue found across one or more respective image slices of the image data set;

determining a confidence or probability of each of the two or more high-dimensional breast tissue objects depicted in the high-dimensional object grid;

generating a synthesized image comprising a lower-dimensional format version of each of the two or more high-dimensional breast tissue objects depicted in the high-dimensional object grid based at least in part upon the confidence or probability of each of the two or more high-dimensional breast tissue objects, wherein the confidence or probability is determined based at least in part on a weight assigned to each of the two or more high-dimensional breast tissue objects, and wherein the weights assigned to the two or more high-dimensional breast tissue objects is based on the respective object type,

where generating the synthesized image comprises:

determining whether two of the two or more high-dimensional breast tissue objects are likely to overlap in the displayed synthesized image; and

if it is determined that the two of the two or more high-dimensional breast tissue objects are likely to overlap in the synthesized image:

determining a difference in assigned weight of each of the two of the two or more high-dimensional breast tissue objects likely to overlap, and

if the determined difference is lower than a threshold value,

modifying one or more aspects of at least one of two of the two or more high-dimensional breast tissue objects, and

displaying both of the two of the two or more high-dimensional breast tissue objects likely to overlap in the synthesized image, wherein the modification pertains to emphasizing at least a portion of the high-dimensional breast tissue object assigned the higher weight relative to the high-dimensional object assigned the lower weight.

20. The method of claim 19, wherein generating the synthesized image comprising the lower-dimensional format version of the respective two or more high-dimensional

breast tissue objects includes modifying one or more aspects of at least one of the two or more high-dimensional breast tissue objects to thereby emphasize high-dimensional breast tissue objects corresponding to the first object type over high-dimensional breast tissue objects corresponding to the 5 second object type.

21. The method of claim **19**, further comprising, if the determined difference is equal to or greater than the threshold value, displaying only one of the two of the two or more high-dimensional breast tissue objects likely to overlap in 10 the synthesized image, the displayed object being the one assigned the higher weight.

22. The method of claim **19**, wherein the first portion comprises an outline portion of the modified high-dimensional breast tissue object, and the second portion comprises 15 a middle portion of the same object.

23. The method of claim **19**, further comprising, if the determined difference is lower than a threshold value, modifying at least one of the two of the two or more high-dimensional breast tissue objects to emphasizing a first 20 portion of the modified high-dimensional breast tissue object relative to a second portion of the same object.

* * * * *