



US009879321B2

(12) **United States Patent**
Bing et al.

(10) **Patent No.:** **US 9,879,321 B2**

(45) **Date of Patent:** ***Jan. 30, 2018**

(54) **CORN EVENT DAS-59122-7 AND METHODS FOR DETECTION THEREOF**

15/8286 (2013.01); *C12Q 1/6895* (2013.01);
C12Q 2600/13 (2013.01)

(71) Applicants: **E. I. DuPont de Nemours and Company**, Wilmington, DE (US); **Pioneer Hi-Bred International, Inc.**, Johnston, IA (US); **Dow Agrosciences LLC**, Indianapolis, IN (US)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

(72) Inventors: **James Wayne Bing**, Ankeny, IA (US); **Robert F. Cressman, Jr.**, Wilmington, DE (US); **Manju Gupta**, Carmel, IN (US); **Salim M. Hakimi**, Sacramento, CA (US); **David Hondred**, Altoona, IA (US); **Todd L. Krone**, Johnston, IA (US); **Mary E. Hartnett Locke**, Mickleton, NJ (US); **Abigail K. Luckring**, West Chester, PA (US); **Sandra E. Meyer**, Des Moines, IA (US); **Daniel Moellenbeck**, Granger, IA (US); **Kenneth Edwin Narva**, Zionsville, IN (US); **Paul D. Olson**, Kalahoe, HI (US); **Craig D. Sanders**, Bear, DE (US); **Jimei Wang**, Johnston, IA (US); **Jian Zhang**, Urbandale, IA (US); **Gan-Yuan Zhong**, Urbandale, IA (US)

5,474,796	A	12/1995	Brennan	
5,541,308	A	7/1996	Hogan et al.	
6,232,063	B1	5/2001	Beckmann et al.	
6,340,593	B1	1/2002	Cardineau et al.	
6,372,480	B1	4/2002	Narva et al.	
6,395,485	B1	5/2002	De Beuckeleer	
6,458,584	B1	10/2002	Mirzabekov et al.	
6,825,400	B2	11/2004	Behr et al.	
7,112,721	B2	9/2006	Fabijanski et al.	
7,323,556	B2	1/2008	Bing et al.	
7,875,430	B2 *	1/2011	Bing	C07K 14/415 435/6.13
7,888,023	B2 *	2/2011	Bing	C07K 14/415 435/6.13
7,897,342	B2 *	3/2011	Bing	C07K 14/415 435/6.13
7,932,033	B2 *	4/2011	Bing	C07K 14/415 435/6.13
8,592,653	B2 *	11/2013	Bing	C07K 14/415 435/6.1
8,952,223	B2 *	2/2015	Bing	C07K 14/415 536/24.3

(73) Assignees: **Pioneer Hi-Bred International, Inc.**, Johnston, IA (US); **E.I. du Pont de Nemours and Company**, Wilmington, DE (US); **Dow Agrosciences LLC**, Indianapolis, IN (US)

(Continued)

FOREIGN PATENT DOCUMENTS

EP	0 857 791	A1	8/1998
EP	1167531	A1	1/2002

(Continued)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 171 days.

OTHER PUBLICATIONS

This patent is subject to a terminal disclaimer.

Hui Ekw et al., "Mini-Review: Strategies for cloning unknown cellular flanking DNA sequences from foreign integrants." *Cell. Mol. Life Sci.*, 1998, 54:1403-1411.

Zimmermann A. et al. "Event Specific Transgene Detection in Bt11 Corn by Quantitative PCR at the Integration Site." *LWT—Food Science and Technology*, 2000, vol. 33, May: 210-216.

An, et al., (1989) "Functional Analysis of the 3' Control Region of the Potato Wound-Inducible Proteinase Inhibitor II Gene", *The Plant Cell*, vol. 1, (pp. 115-122).

Buck, et al., (1999) "Design Strategies and Performance of Custom DNA Sequencing Primers," *BioTechniques*, vol. 27, (pp. 528-536).

Christensen, et al., (1992) "Maize polyubiquitin genes: structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation", *Plant Molecular Biology*, vol. 18, Issue 4, (pp. 675-689).

(Continued)

Primary Examiner — Young J Kim

(74) *Attorney, Agent, or Firm* — Ballard Spahr LLP

(57) **ABSTRACT**

The invention provides DNA compositions that relate to transgenic insect resistant maize plants. Also provided are assays for detecting the presence of the maize DAS-59122-7 event based on the DNA sequence of the recombinant construct inserted into the maize genome and the DNA sequences flanking the insertion site. Kits and conditions useful in conducting the assays are provided.

(21) Appl. No.: **14/577,063**

(22) Filed: **Dec. 19, 2014**

(65) **Prior Publication Data**

US 2015/0167076 A1 Jun. 18, 2015

Related U.S. Application Data

(60) Continuation of application No. 11/938,483, filed on Nov. 12, 2007, now Pat. No. 8,952,223, which is a division of application No. 11/237,222, filed on Sep. 28, 2005, now Pat. No. 7,323,556.

(60) Provisional application No. 60/614,225, filed on Sep. 29, 2004.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
C07K 14/415 (2006.01)
C12N 15/82 (2006.01)

(52) **U.S. Cl.**
CPC *C12Q 1/6876* (2013.01); *C07K 14/415* (2013.01); *C12N 15/8277* (2013.01); *C12N*

4 Claims, 7 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

2002/0013960	A1	1/2002	Behr et al.
2002/0102582	A1	8/2002	Levine
2002/0199215	A1	12/2002	Boets et al.
2003/0198943	A1	10/2003	Remacle et al.
2006/0141495	A1	6/2006	Wu
2008/0166725	A1	7/2008	Bing et al.
2008/0166726	A1	7/2008	Bing et al.
2008/0171334	A1	7/2008	Bing et al.
2008/0182256	A1	7/2008	Bing et al.

FOREIGN PATENT DOCUMENTS

WO	WO-84/02913	A1	8/1984
WO	WO-01/14417	A2	3/2001
WO	WO-02/15701		2/2002
WO	WO-03/052073	A2	6/2003
WO	WO-2004/011601	A2	2/2004

OTHER PUBLICATIONS

Crickmore, et al., (2009) available at: http://www.lifesci.sussex.ac.uk/Home/Neil_Cruckmore/Bt/, (pp. 1-11).

Database EMBL (2003) "PUFTW67TB ZM_0.6_1_0_KB *Zea mays* genomic clone ZMMBTa0732K13, genomic survey sequence", retrieved from EBI accession No. EM_PRO:CG069192, Database accession No. CG069192 (3 pages).

Ellis, et al., (2002) "Novel *Bacillus thuringiensis* Binary Insecticidal Crystal Proteins Active on Western Corn Rootworm, *Diabrotica virgifera* Le Conte", Applied and Environmental Microbiology, vol. 68, No. 3, (pp. 1137-1145).

Hammack, L., et al., (2003) "Larval Sampling and Instar Determination in Field Populations of Northern and Western Corn Rootworm (*Coleoptera: chrysomelidae*)," Journal of Economic Entomology, vol. 96(4), (pp. 1153-1159).

Herman, et al., (2002) "Binary Insecticidal Crystal Protein from *Bacillus thuringiensis*, Strain PS 149B1: Effects of Individual Protein Components and Mixtures in Laboratory Bioassays", Journal of Economic Entomology, vol. 95(3), (pp. 635-639).

Hunst, et al., (2004) "Application for determination of non-regulated status for *B. t.* Cry34/35Abl insect-resistant, glufosinate-tolerant corn: corn line 59122", U.S. Department of Agriculture, available at: <http://www.agbios.com/docroot/decdocs/05-242-046.pdf> (237 pages).

Lowe, et al., (1990) "A computer program for selection of oligonucleotide primers for polymerase chain reactions," Nucleic Acids Research, vol. 18(7), (pp. 1757-1761).

Mitsuhara, et al., (1996) "Efficient promoter cassettes for enhanced expression of foreign genes in dicotyledonous and monocotyledonous plants", Plant Cell Physiol. vol. 37(1), (pp. 49-59).

Moellenbeck, et al., (2001) "Insecticidal proteins from *Bacillus thuringiensis* protect corn from corn rootworms," Nature Biotechnology, vol. 19, (pp. 668-672).

Ricker, (2004) "Biotechnology consultation note to the file BNF No. 000081", U.S. Food and Drug Administration, available at: <http://www.cfsan.fda.gov/~rdb/bnfm081.html> (6 pages).

Sequence Search Reports, OM nucleic-nucleic search, 2009, (pp. 1-6).

Stratagene Catalog, p. 40, 1988.

Whitelaw, et al., (2003, GenBank Accession No. BZ739496).

Wohlleben, et al., (1988) "Nucleotide sequence of the phosphinothricin N-acetyltransferase gene from *Streptomyces viridochromogenes* Tü494 and its expression in *Nicotiana tabacum*", Gene. vol. 70(1), (pp. 25-37).

Zambryski, et al., (1996, GenBank Accession No. TIP37TD2).

International Search Report issued by the International Searching Authority on Jun. 22, 2006 for application PCT/US2005/034947, filed on Sep. 28, 2005 and published as WO 2006/039376 on Apr. 13, 2006 (Applicant—Pioneer Hi-Bred, Int'l, Inc. // Inventor—Bing, et al.) (3 pages).

Written Opinion issued by the International Searching Authority on Mar. 29, 2007 for application PCT/US2005/034947, filed on Sep. 28, 2005 and published as WO 2006/039376 on Apr. 13, 2006 (Applicant—Pioneer Hi-Bred, Int'l, Inc. // Inventor—Bing, et al.) (12 pages).

International Preliminary Report on Patentability issued by the International Searching Authority on Mar. 29, 2007 for application PCT/US2005/034947, filed on Sep. 28, 2005 and published as WO 2006/039376 on Apr. 13, 2006 (Applicant—Pioneer Hi-Bred, Int'l, Inc. // Inventor—Bing, et al.) (13 pages).

* cited by examiner

Fig. 1A

1 CTGAGCGCAC AACAGCGAGT CGCATGGCAC CGGACGACAT GAGCGAGATT
51 TAGATCGGAG GGTGCGGACA TGGGGCAACC TGCGCAGCTA ACGCAGGGAT
101 CCACACGACC ACCAACGAAG CCAAGCCCGG GCACGTCCCC AGGCAGSTTG
151 GGCCCTGGTT CCACCAGCGG ATGCATGCAG TGAAGCGGGG ACGGAGAGAC
201 AAGCCGAGGG CCGGGGTGGG AATGSCCTCC GGGAGGACGA GTGGAGGAGA
251 AGAATCTAGA GGCATCGAGA TTCGAGAAGC CGACGGAGAC AAGATTGCTG
301 TGGGGGGAGA CAAACCGCGG GGCTGAGCGC CGTTGATATG GGATCAGACG
351 GTGTGGATAA AAAAAGTGAC GTTGATAGAA CGTCTGGCCA GTGAAAAAAC
401 AAAACAACCTC CAACAAAATA CTTTAAAAGC TCTTATAACC TAAATGTAGG
451 GGATCAAACA CGTCTCTACA CTATTTAGCA GCGTCTCTA AATGATCCTC
501 TAAATTTAGA GAACGCTACT AGATTCTCTA TATATAGTTT CTCTAAAACA
551 TCTTTTATCC ATTTAAATAC TTTAAATAAC CGGTTTAAACA AAACFAAAAF
601 ATATACAATA CATTGAGAG TATGACAAAT ACGTATGTAT AAAAATAAAA
651 AATAAAATAA TGTATTAGTC TACTTTGAAT CTTCTTTTCT TCATAATATA
701 ATGATGTATA GCTCTCATGT GCGTTGAGAA AAAAGTTAGA GCTFAGCGTT
751 TAATGTGTAG TGACAGTCTT CGACGAAATC TCCCTAATGA GATGAATTAC
801 TGGAGGTCC ATCAGAAAGT CCCCTGAAAA GAGGCATTTA TTTAGTTTAG
851 TCAGCAATTT CTGGGAACAC AAATATTCTT TTGTTATCAC CACTATPAAA
901 AATCTATGST TATAACTTAT AATAACATGA AAAAATAAAT TAGCATCCCA
951 TATATATAAA AACTGAAGGA AGCCATATAT ACTAACATAA GTTAGGAGAA
1001 ACTAAGAAGG TTGTGCAAAG CTTGCCACTGC TCCAAAATAC TGCAAACAAC
1051 CACTCTCCTC TACCAACCAA AGAAACTCAT GTACTCCCTC CGTCTTTTTT
1101 TATTTGTGCG ATTTTAGTTT AAAAATGAAC TAGCAGTCGA CAAATATTCG
1151 AGAACAGATA TAGTATATAC TAACATAACT TAGGAGATAC TAAGAAAGTT
1201 GCGCAGAGCT TTCACTGTTT CAAATTACTG CAAAGCCTCT CCCCTCTGCC
1251 AGTACATCTA CGAGATGTTT CAGTAAACA AAGATTCAGA CAAGTGATGA
1301 GCCACTTCTT GTCATAGATT GTGTGGTCAA CCAACCCATT GATGCCACGG
1351 TTTTGTGTCG TCCATGCTTT TGTATPAAAA CATCAGTTAT GTTTACCATG
1401 TCCGATATGC TCTACATAAT GACAATCAAC TTGGTGTTC TATATTTTAC
1451 AATGTTAGGA ATTTCAATAG CTACGAACAC TTCAATAGAA GTGCCCTTGT
1501 GGGATCACCT TAATGTGTTG TTGATGTAAG GAGAAGAATC TTAATTTACT
1551 CTTGCTAAAT TTGAACTACA CAAAACCACT GCACTGAGGA TTGTCCTAAF
1601 AAATFACTGC TCATACACGT TAGCATCTGT TCAGATACTG AGCTAATCCC
1651 TAGGATTAAT GGATTTGTAA AAGATATGCC CAATCATTCA TTTTACTTAT
1701 TTATTTCTTA GTTATCCACT TGAAGATTA CATACTTTG AAATAAATTT
1751 CTTAGAGTA AAGTGAAAA CAGTTATTTA AATACATTTT AGTTATTTAT
1801 TTTCTTCTTT TTCTAATTTT TTCCTTGTAT TTGAAGCTG AAAAGATAAC
1851 TTTGCCCTTA TACATATTTT ATCTTCTACG TACGCATCTG AACACCTCT
1901 CTTTGTCCCC TGATCGTGCA GCAATTAGTG CTATGAATCG CGTTTAAAGG
1951 CTGCAAAATC AFGGCTGGGG CTTCSTCCTC GAGTCGTCTT GCTGCTCGAT
2001 GTCACCTCGA GTCCCGCACC GACCTCAGTG CTTGTTCTTG TTGGAGCCAC
2051 CTCTCTCGGA CGATCGCCAA AGACGGATAA GGCCGAAGCC GTCACCTCAG
2101 ACCGCGCTCA TCGCGCGTAG CAGACTCCTA CATAGCAGGG CCAGGGTATG
2151 TGCACCTTTG CAAGTTTAGG ATTTGGAACCA GCGACCAGAA TCCACAAGAT
2201 TGGAGCAAAC GACCAAAAAT TCACAAGGAT TGGCGGCTGA CATTGCCAGC
2251 CGCGGATCSC ATGCGGCGGC GCGGCGCGG GCGAGCACGG GAGCGCGGA
2301 CAGTCGAGCT CCATTGGAAC GTAGAAATAC TTAAGGGCAA GGCTCCCAA
2351 TACTTGAAAA AATAGGAAAA AGAAGAAAAAT ACATGAAATG ATATFGAAAT
2401 CAATTGGAAG ATGTTATGAA TCTTGTTTTT GCAAAGCGAA CGATTCAAGT
2451 GGCAAAACFA TGAATCTTTT TGTTTGAAGT CCCAAATATA AAATTTCTC
2501 GTACTCACCA ACATTGGTGC GCACCTGTGA TTGGCTCATA AAAATCTTTG
2551 GAGGGACCGA AGAAAGAGTG AAGGGATAAG CAAGTAAAAG CGCTCAAACA
2601 CTGATAGTTT AAACGTGAAGG CCGGAAACGA CAATCTGATC ATGAGCGGAG

Fig. 1B

2651 AATTAAGGGA GTCACGTTAT GACCCCCGCC GATGACGCGG GACAAGCCGT
 2701 TTTACGTTTG GAACTGACAG AACCGCAACG TTGAAGGAGC CACTCAGCAA
 2751 GCTTACTAGT AGCGCTGTTT AAACGCTCTT CAACTGGAAG AGCGGTACC
 2801 CGGACCGAAG CTTGCATGCC TGCAGTGCAG CGTACCCCGG TCGTGCCCTT
 2851 CTCTAGAGAT AATGAGCATT GCATGTCTAA GTTATAAAAA ATTACCACAT
 2901 ATTTTTTTTT TCACACTTGT TTGAAGTGCA GTTTATCTAT CTTTATACAT
 2951 ATATTTAAAC TTTACTCTAC GAATAATATA ATCTATAGTA CTACAATAAT
 3001 ATCAGTGTTC TAGAGAATCA TATAAATGAA CAGTTAGACA TGGCTAAAAG
 3051 GACAATTGAG TATTTTGACA ACAGGACTCT ACAGTPTTAT CTTTTTAGTG
 3101 TGCATGTGTT CTCCTTTTTT TTTGCAAATA GCTTCACCTA TATAATACTT
 3151 CATCCATTTT ATTAGTACAT CCATTTAGGG TTTAGGGTTA ATGGTTTTTA
 3201 TAGACTAATT TTTTTAGTAC ATCTATTTTA TTCTATTTTA GCCTCTAAAF
 3251 TAAGAAAAC AAAACTCTAT TTTAGTTTTT TTATTTAATA ATTTAGATAT
 3301 AAAATAGAAT AAAATAAAGT GACTAAAAAT TAAACAAATA CCCTTTAAGA
 3351 AATTAAAAAA ACTAAGGAAA CATTTTTCTT GTTTCGAGTA GATAATGCCA
 3401 GCCTGTAAA CGCCGTCGAC GAGTCTAACG GACACCAACC AGCGAACCAG
 3451 CAGCGTCGCG TCGGBCCAAG CGAAGCAGAC GGCACGGCAT CTCTGTGCTT
 3501 GCCTCTGGAC CCTCTCGAG AGTTCCGCTC CACCGTTGGA CTGTCTCCGC
 3551 TCTCCGCATC CAGAAATGCG GTGGCCGAGC GGCAGACGTC AGCCGGCACG
 3601 GCAGGCGGCC TCCTCCTCCT CTCACGGCAC CGGCAGCTAC GGGGATTC
 3651 TTTCCACCG CTCCTTCGCT TTCCCTTCTT CGCCCGCGT AATAAATAGA
 3701 CACCCCTCC ACACCTCTT TCCCCAACCT CGTGTGPTC GGAGCGCACA
 3751 CACACACAAC CAGATCTCCC CCAAATCCAC CCGTCGGCAC CTCCGCTTCA
 3801 AGGTACGCCG CTCGTCTCTC CCCCCCCCC CTCTCTACCT TCTCTAGATC
 3851 GCGGTTCCGG TCCATGGTTA GGGCCCCGTA GTTCTACTTC TGTTCAATGT
 3901 TGTGTTAGAT CCGTGTGTTT GTTAGATCCG TGCTGCTAGC GTTCGTACAC
 3951 GGATCGGACC TGTACGTCAG ACACGTTCTG ATTGCTAACT TGCCAGGFTT
 4001 TCTCTTTGGG GAATCCTGGG ATGGCTCTAG CCGTTCGCA GCCGGGATCG
 4051 ATTTCAATGAT TTTTTTTGTT TCGTTGCATA GGGTTTGGTT TGCCCTTTTC
 4101 CTTTATTTCA ATATATGCCG TGCACTTGTT TGTGGGTCA TCTTTTTCAF
 4151 CTTTTTTTTG TCTTGGTTGT GATGATGTGG TCTGTTGGG CGGTGCTTCT
 4201 AGATCGGAGT AGAATCTGT TTCAAACCTAC CEGGTGGATT TATTAATTTT
 4251 GGATCTGTAT GTGTGTGCCA TACATATTCA TAGTTACGAA TTGAAGATGA
 4301 TGGATGGAAA TATCGATCTA GGATAGGTAT ACATGTTGAT GCGGGTTTTA
 4351 CTGATGCATA TACAGAGATG CTTTTTGTTT CCTEGTGTG GATGATGTGG
 4401 TPTGGTTGGG CGGTGTTCA TTCGTTCTAG ATCGGAGTAG AATACTGTTT
 4451 CAAACTACCT GGTGTAATTA TTAATTTTGG AACTGTATGT GTGTGTCAFA
 4501 CATCTTCATA GTTACGAGTT TAAGATGGAT GGAAATATCG ATGTAGGATA
 4551 GGTATACATG TTGATGFGGG TTTTACTGAT GCATATACAT GATGGCATAF
 4601 GCAGCATCTA TTCATATGCT CTAACCTTGA GTACCTATCT ATTATAATAA
 4651 ACAAGTATGT TTTATAATTA TTTTGATCTT GATATACTTG GATGATGGCA
 4701 TATGCAGCAG CTATATGTGG ATTTTTTTAG CCGTGCCTTC ATACGCTATT
 4751 TATTTGCTTG GTACTGTTTC TTTTGTGAT GCTCACCTTG TTGTTTGGTG
 4801 TFACTTCTGC AGGTGACTC TAGAGGATCC ACACGACACC ATGTCCGCCC
 4851 GCGAGGTGCA CATCGAGTG AACACAAGA CCGGCCACAC CCTCCAGCTG
 4901 GAGGACAAGA CCAAGCTCGA CGGCGGCAGG TGGCCACCT CCCCAGCCA
 4951 CGTGGCCAAC SACCAGATCA AGACCTTCTT GGCCGAATCC AACGGCTTCA
 5001 TGACCGGCAC CGAGGGCAC ATCTACTACT CAATTAATGG CGAGGGCCGAG
 5051 ATCAGCCTCT ACTTCGACAA CCCGTTCCGC GGCTCCAACA AATACGACGG
 5101 CCACTCCAAC AAGTCCCACT ACCACATCAT CACCCAGGGC GGCTCCGGCA
 5151 ACCAGTCCCA CGTGACCTAC ACCATCCAGA CCACCTCTTC CCGCTACGGC
 5201 CACAAGTCTT GAGTCATGAG TCATGAGTCA GTTAACCTAG ACTTGTCCAT
 5251 CTCTGGATT GGCCAACTTA ATTAATGTAT GAAATAAAG GATGCACACA

Fig. 1C

5301 TAGTGACATG CTAATCACTA TAATGTGGGC ATCAAAGTTG TGTGTTATGT
 5351 GTAATFACTA GTTATCTGAA TAAAAGAGAA AGAGATCATC CATATTTCTT
 5401 ATCCTAAATG AATGTCACGT GTCTTTATAA TTCTTTGATG AACCCAGATGC
 5451 ATTTCAATTA CCAAATCCAT ATACATATAA ATATTAATCA TATATAAATTA
 5501 ATATCAAATG GGTTAGCAAA ACAAATCTAG TCTAGGTGTG TTTTGGGAAT
 5551 GCGGCCGCGG ACCGAATTGG GGATCTGCAT GAAAGAAACT GTCGCACTGC
 5601 TGAACCGCAC CTTGTCACTT TCATCGAACA CGACCTGTGC CCAAGATGAC
 5651 GGTGCTGCGG TCTAAGTGAG GCTGAATTGC CTTGGACAGA AGCGGACTCC
 5701 CTACAATTAG TTAGGCCAAA CGGTGCATCC ATGTGTAGCT CCGGGCTCGG
 5751 GCTGTATCGC CATCTGCAAT AGCATCCATG GAGCTCGTTC CATGTAGTTG
 5801 GAGATGAACC AATGATCGGG CGTGTGGACG TATGTTCCCTG TGTACTCCGA
 5851 TAGTAGAGTA CGTGTAGCT CTTTCATGGT GCAAGTGAAA TTTGFTTGG
 5901 TTTAATFACC CCTACGTTAG TTGCGGGACA GGAGACACAT CATGAATTA
 5951 AAGGCGATGA TGTCTCTCTC TGTAATGTTA TTCTTTTGAT GTGATGAATC
 6001 AAAATGTCAT ATAAAACATT TGTTGCTCTT TAGTTAGGCC ATAGCTFAGA
 6051 ACGAAATGCT CGTGTAGCGG GGCTACGAGC CTATGACGCA ATAACACTGG
 6101 TTTGCCGSCC CGGAGTCSCT TGACAAAAAA AAGCATGTTA AGTTTATTTA
 6151 CAATTCAAAA CCTAACATAT TATATTCCTT CAAAGCAGGT TCACGATCAC
 6201 ACCTGTACCT AAAAAAACA TGAAGAATAT ATTACTCCAT TATTATGAGA
 6251 TGAACCACTT GGCAAGAGTG GTAAGCTATA TAAAAAATG AACATTATTA
 6301 CGAGATGTTA TATGCCATTA TATTGATTGG AAGATATATG TTTCTTTCTC
 6351 CCACGGGCAC CTAACGGATA CATGATAAGG CCAAGGCACA TCACGGGAAA
 6401 TTATTCGAAT ACATGTTACG CCCTATTGCC GGAAAAAATA TGCAGGGCAG
 6451 GTGTTGGCCG TAGCCGATTA AGCACTTAAG CTGGAGGTTG CCACACTTGG
 6501 ATGCCAAGCT CTGACCCCTC TAAAACATCG GCGGCTTTGT CCGTATCCGT
 6551 ATCCCTATC CGACATCTAG CTGGCCACAC GACGGGGCTG GGCATGCTGT
 6601 GGATGCCGGG TCGACGTCTA TCGTCAGCCA TCATAGACCA ATCGACCATC
 6651 TGTTATGGAT GCTTGCTAGC TAGACTAGTC AGACATAAAA TTTGGTACT
 6701 TTCTCCCAAC TGGGAGACGG GGACTGATGT GCAGCTGCAC GTGAGCTAAA
 6751 TTTTTCCTA TAAATATGCA TGAAATACTG CATTATCTTG CCACAGCCAC
 6801 TGCCACAGCC AGATAACAAG TGCAGCTGGT AGCAGCCAAC GCATAGCTCT
 6851 GGACTTGTAG CTAGGTAGCC AACC GGATCC ACACGACACC ATGCTCGACA
 6901 CCAACAAGGT GTACGAGATC AGCAACCACG CCAACGGCCT CTACGCCGCC
 6951 ACCTACCTCT CCTCGACGA CTCCGGCGTG TCCCTCATGA ACAAGAACGA
 7001 CGACGACATC GACGACTACA ACCTCAAGTG GTTCTCTTC CCGATCGACG
 7051 ACGCCAGTA CATCATCACC TCCTACGCC CCAACAACCT CAAGGTGTGG
 7101 AACGTGAACA ACGACAAGAT TAATGTGTCA ACCTACTCCT CCACCAACTC
 7151 CATCCAGAAG TGGCAGATCA AGGCCAACGG CTCTCTCTAC GTGATCCAGT
 7201 CCGACAACGG CAAGGTGCTC ACCGCCGGCA CCGGCCAGGC CCTCGGCCCTC
 7251 ATCCGCCCTA CCGACGAGTC CTCCAACAAC CCGAACCCAGC AATGGAACCT
 7301 GACGTCCGTG CAGACCATCC AGCTCCCGCA GAAGCCGATC ATCGACACCA
 7351 AGCTCAAGGA CTACCCGAAG TACTCCCGCA CCGCAACAT CGACAACGGC
 7401 ACCTCCCGC AGCTCATGGG CTGGACCCTC GTGCCGTGCA TCATGGTGAA
 7451 CGACCCGAAC ATCGACAAGA ACACCCAGAT CAAGACCACC CCGTACTACA
 7501 TCCTCAAGAA GTACCAATAC TGGCAGAGGG CCGTGGGCTC CAACGTCCGG
 7551 CTCGCCCGC ACGAGAAGAA GTCCTACACC TACGAGTGGG GCACCCAGAT
 7601 CGACCAGAAG ACCACCATCA TCAACACCTT OGGCTTCCAG ATCAACATCG
 7651 ACAGCGGCAT GAAGTTCGAC ATCCCGGAGG TGGGCGGCGG TACCGACGAG
 7701 ATCAAGACCC AGCTCAACGA GGAGCTCAAG ATCGAGTATT CACATGAGAC
 7751 GAAGATCATG GAGAAGTACC AGGAGCAGTC CGAGATCGAC AACCCGACCG
 7801 ACCAGTCCAT GAACTCCATC GGCTTCTCTA CCATCACCTC CCTGGAGCTC
 7851 TACCGCTACA ACCGCTCCGA GATCCGCATC ATGCAGATCC AGACCTCCGA
 7901 CAACGACACC TACAACGTGA CCTCTTACCC GAACCACCAG CAGGCCCTGC

Fig. 1D

```

7951   TGCTGCTGAC CAACCACCTCC TACGAGGAGG TGGAGGAGAT CACCAACATC
8001   CCGAAGTCCA CCCTCAAGAA GCTCAAGAAG TACTACTTCT GAGTCATGAG
8051   TCATGAGTCA GTF AACCTAG ACTTGTCCAT CTCTCGGATT GGCCAACCTA
8101   ATTAATGTAT GAAATAAAAG GATGCACACA TAGTGACATG CTAATCCTA
8151   TAATGTGGGC ATCAAAGTTC TGTGTTATGT GTAATTACTA GTTATCTGAA
8201   TAAAAGAGAA AGAGATCATC CATATTTCTT ATCCTAAAATG AATGTCACGT
8251   GTCTTTTATAA TTCTTTGATG AACCAGATGC ATTTTCATTAA CCAAATCCAT
8301   ATACATATAA ATATTAATCA TATATAATTA ATATCAATTC GGTTAGCAAA
8351   ACAAATCTAG TCTAGGTGTG TTTTGC GAAT TCCCATGGAG TCAAAGATTC
8401   AAATAGAGGA CCTAACAGAA CTCGCCGTAA AGACTGGCGA ACAGTTCATA
8451   CAGAGTCTCT TACGACTCAA TGACAAGAAG AAAATCTTCG TCAACATGGT
8501   GGAGCACGAC ACGCTTGTCT ACTCCAAAAA TATCAAAGAT ACAGTCTCAG
8551   AAGACCAAAG GCCAATTCAG ACTTTTCAAC AAAGGGTAAT ATCCGGAAAC
8601   CTCTCGGAT TCCATGCCC AGCTATCTGT CACTTTATTG TGAAGATAGT
8651   GGAAAAGGAA GGTGGCTCCT ACAAATGCCA TCATTGCGAT AAAGGAAAGG
8701   CCATCGTTGA AGATGCCTCT GCCGACAGTG GTCCCAAAGA TGGACCCCCA
8751   CCCACGAGGA GCATCGTGA AAAAGAAGAC GTTCCAACCA CGTCTTCAAA
8801   GCAAGTGGAT TGATGTGATA TCTCCACTGA CGTAAGGGAT GACGCACAAT
8851   CCCACTATCC TTCGCAAGAC CCTTCCTCTA TATAAGGAAG TTCATTTTAT
8901   TTGGAGAGGA CAGGGTACCC GGGGATCCAC CATGTCTCCG GAGAGGAGAC
8951   CAGTTGAGAT TAGGCCAGCT ACAGCAGCTG ATATGGCCGC GGTGTTGTGAT
9001   ATCGTTAACC ATTACATTGA GACGTCTACA GTGAACTTTA GGACAGAGCC
9051   ACAAACACCA CAAGAGTGA TTGATGATCT AGAGAGGTTG CAAGATAGAT
9101   ACCCTTGGTT GGTGCTGAG GTTGAGGGTG TGTGGCTGG TATTGCTTAC
9151   GCTGGGCCCT GGAAGGCTAG GAACGCTTAC GATTGGACAG TTGAGAGTAC
9201   TGTTTACGTG TCACATAGGC ATCAAAGGTT GGGCTAGGA TCCACATTTT
9251   ACACACATTT GCTTAAGTCT ATGGAGGGCC AAGTTTTTAA GTCTGTGGTT
9301   GCTGTTATAG GCCTTCCAAA CGATCCATCT GTFAGGTTGC ATGAGGCTTT
9351   GGGATACACA GCGCGGGTA CATTGCGCGC AGCTGGATAC AAGCATGGTG
9401   GATGGCATGA TGTGGTTTT TGGCAAAGGG ATTTTGAGTT GCCAGCTCCT
9451   CCAAGGCCAG TTAGGCCAGT TACCCAGATC TGAGTCGACC TGCAGGCATG
9501   CCGGCTGAAA TCACCAGTCT CTCTCTACAA ATCTATCTCT CTCTATAATA
9551   ATGTGTGAGT AGTTCACAGA TAAGGGAATT AGGGTTCTTA TAGGGTTCTG
9601   CTGATGTGTT GAGCATATAA GAAACCCCTA GTATGTATTT GTATTTGTAA
9651   AAIACCTCTA TCAATAAAAT TTCTAATTC TAAAACCAA ATCCAGGGCG
9701   AGCTCGGTAC CCGGGGATCC TCTAGAGTCG ACCTGCAGGC ATGCCCGCG
9751   ATATCGATGG GCGCGGCGG AAGCTTCGGT CCGGCCATC GTGGCCTCTT
9801   GCTCTTCAGG ATGAAGAGCT ATGTTTAAAC GTGCAAGCGC TCAATTCGCC
9851   CTATAGTGAG TCGTATTACA ATCGTACGCA ATTCAGTACA TTA AAAACGT
9901   CCGCAATGTG TTATTAAGTT GTCTAAGCT CAATTTTTCC CTCTATGGT
9951   CCGTTTTGTT TATCCTCTAA ATTATATAAT CCAGCTTAAA TAAGTTAAGA
10001  GACAAACAAA CAACACAGAT TATTAATAG ATATGTAAT CTAGATACCT
10051  AGATTAIGTA ATCCATAAGT AGAATATCAG GTGCTTATAT AATCTATGAG
10101  CTCGATTATA TAATCTTAAA AGAAAACAAA CAGAGCCCCT ATAAAAGGG
10151  GTCAAGTGA CACTTGGTCA CTCATTTAAT CCCTCCCTCT CCTCTTTTAT
10201  CCTCTTTTTT GGTGTAATCA CCAATAGTGG TGTGCACCTG TGATTGGCTC
10251  GTAAAATTC TTGGACGAT GGAAGAGTGA AGAGATAAGC AAGTCAAAGA
10301  AAAGTAACAA CGAAGCTTCA TCAGCTACAA ATPTTGGCCC AACTGGTTGC
10351  ACCAGCACCA AACTTACGTA TACATGATTA TCTCTGTTTC CCTCATTTCT
10401  AAGAAAAAAA CGGGTTTCAA AACCACCTGC TTTCAGGAGT AAAAAAGAT
10451  AATAATCTGA AACATTCCTT CCACCTTGGC CCTTATTTGG TTACGTTGCA
10501  ATTCACCCCA ATCCACATGT GGATTGAGAT GGATTGCAGT GTAGCTAGAC
10551  AAACCCTTAG GCCCTGTTG CATAGGAATA CACCAGGAAT TATTCCAGCT

```

Fig. 1E

10601 AATCAAAATT TATATAAATG AGAGAAACAA TTCGGATAGG AATTGTTCCA
 10651 GGACTTCATT CTGCAGTAAC CGAACGGCCC CTTAATCCAC CCCAATACAC
 10701 GTGGATTGGA GTGGATPGAG GTACAGCCAA ACAAGGCCA AGTGCAGATC
 10751 AAATAAAATCA CCCGTCATAT TCTTCTACCT ACAAAAACAG CAATAAACAC
 10801 CTGAATGAAG TTCTAATTTG CACAGTGTAG GTAGGATGAA AATAGTTACC
 10851 TCCTCATGGT CAGTAACTCT TGGCACACAA CTTACATGT AATCGATGTA
 10901 CCACTTGGCT CTTGCCTGAA ACCCAATACA TCTTTAGCAT AAGAATAATA
 10951 TTATGATGGC AAGGCATGAT CACCAGCACT CCTTTATTGT TTAGTAAGTC
 11001 TATCACTCCC CAAAACAATF CAAATGAACA GAGATGCATT GCCCCCAATG
 11051 AATTCTATTT CAATTAGCCG GAAAATFCTA CTTCATCAGA AGCATCCAAA
 11101 TTGCCAGCAT CCTACTAGA CTGACCATGA CCAGGCTGCC GCAGATGCCT
 11151 CTTTTCTGT CCTCTCCTCT TTGCCTTGAG PTTCTCTTCA AGATCCCTCA
 11201 CCCCACGTCT CTTATACATC TTAAAAGCTAA CATGTCTCTC CTCCGCCATC
 11251 TTCCTAACCT TCTCAGTAAT CTCAGCAGCA ATCTGACGGT TGTACAACCT
 11301 CTTTCAGCCCC TTCATCAACT TTGCAAATGT GTCAGGCTGT GGCATCAGTC
 11351 CTGCCTCTAG CATGTCTAAG CAATACAGGC AGGCCTCCTT GACATGTTTC
 11401 TTCGCAAACA GTGCATGAAT CCAGATAGTC CATGCACTCA CATTGAGCTC
 11451 ACAGCCPTTG CTCACAATAC ATTTCCAAAC ATCTTPTGCA AGCTCAAGTT
 11501 TCTCATCTCT GACCAACGCA TPGAGGAGGT CCTTCAGCAC CCCATATTGC
 11551 GGTACCACAA AGAGCCCCCT CCCAACCATG TCTTTAAAAT AACTACATGC
 11601 CTCAATCAGC AAACCCTGCC CAACAAGGCC ACTCACCACG ATAGCAAATG
 11651 TATCGAACCAC AGGACTGAGC CCAGCACTTT CCATCTCATT CCACAATGTC
 11701 ATGGCTTGCT TGGTCTCCCC AAGCCTGCAG GCCAACCAGAA TCACCACATT
 11751 GTATATCTTG AGATCTGGTG GACACCGGCA CTCCCGCATC CTCTCCATCA
 11801 GCTCCAAGCA CTCTCAAGC TGCTCCTTCT TCTCGTGTGC TACAAAAGAAA
 11851 CCATGGTACA CGGCAGGTC CACCCGAGG CCATCCCTCG ACATAGCATC
 11901 CAAGAACTCG TACCCCTGGG AT

Fig. 2





Fig. 3



CORN EVENT DAS-59122-7 AND METHODS FOR DETECTION THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 11/938,483 filed Nov. 12, 2007; which is a division of U.S. patent application Ser. No. 11/237,222 filed Sep. 28, 2005, now U.S. Pat. No. 7,323,556, issued Jan. 29, 2008; which claims the benefit of U.S. Provisional Application No. 60/614,225 filed Sep. 29, 2004, the contents of which are herein incorporated by reference in their entirety.

Pursuant to 37 C.F.R. §1.121(b)(1)(i) and (ii), Applicants respectfully request that the Examiner add a paragraph titled "REFERENCE TO SEQUENCE LISTING" following the paragraph titled "CROSS-REFERENCE TO RELATED APPLICATIONS" on page 1 of the application as presented below.

The Sequence Listing submitted Dec. 19, 2014 as a text file named "36446 0047U3 Sequence Listing.txt," created on Dec. 19, 2014, and having a size of 58,694 bytes is hereby incorporated by reference pursuant to 37 C.F.R. §1.52(e)(5).

FIELD OF INVENTION

Embodiments of the present invention relate to the field of plant molecular biology, specifically an embodiment of the invention relates to a DNA construct for conferring insect resistance to a plant. Embodiments of the invention more specifically relate to an insect resistant corn plant DAS-59122-7 and to assays for detecting the presence of corn plant DAS-59122-7 DNA in a sample and compositions thereof.

BACKGROUND OF INVENTION

An embodiment of this invention relates to the insect resistant corn (*Zea mays*) plant DAS-59122-7, also referred to as maize line DAS-59122-7 or maize event DAS-59122-7, and to the DNA plant expression construct of corn plant DAS-59122-7 and the detection of the transgene/flanking insertion region in corn plant DAS-59122-7 and progeny thereof.

Corn is an important crop and is a primary food source in many areas of the world. Damage caused by insect pests is a major factor in the loss of the world's corn crops, despite the use of protective measures such as chemical pesticides. In view of this, insect resistance has been genetically engineered into crops such as corn in order to control insect damage and to reduce the need for traditional chemical pesticides. One group of genes which have been utilized for the production of transgenic insect resistant crops are the delta-endotoxins from *Bacillus thuringiensis* (B.t.). Delta-endotoxins have been successfully expressed in crop plants such as cotton, potatoes, rice, sunflower, as well as corn, and have proven to provide excellent control over insect pests. (Perlak, F. J. et al. (1990) *Bio/Technology* 8, 939-943; Perlak, F. J. et al. (1993) *Plant Mol. Biol.* 22: 313-321; Fujimoto H. et al. (1993) *Bio/Technology* 11: 1151-1155; Tu et al. (2000) *Nature Biotechnology* 18:1101-1104; PCT publication number WO 01/13731; and Bing J W et al. (2000) Efficacy of Cry 1F Transgenic Maize, 14th Biennial International Plant Resistance to Insects Workshop, Fort Collins, Colo.).

The expression of foreign genes in plants is known to be influenced by their location in the plant genome, perhaps due

to chromatin structure (e.g., heterochromatin) or the proximity of transcriptional regulatory elements (e.g., enhancers) close to the integration site (Weising et al., *Ann. Rev. Genet* 22:421-477, 1988). At the same time the presence of the transgene at different locations in the genome will influence the overall phenotype of the plant in different ways. For this reason, it is often necessary to screen a large number of events in order to identify an event characterized by optimal expression of an introduced gene of interest. For example, it has been observed in plants and in other organisms that there may be a wide variation in levels of expression of an introduced gene among events. There may also be differences in spatial or temporal patterns of expression, for example, differences in the relative expression of a transgene in various plant tissues, that may not correspond to the patterns expected from transcriptional regulatory elements present in the introduced gene construct. For this reason, it is common to produce hundreds to thousands of different events and screen those events for a single event that has desired transgene expression levels and patterns for commercial purposes. An event that has desired levels or patterns of transgene expression is useful for introgressing the transgene into other genetic backgrounds by sexual outcrossing using conventional breeding methods. Progeny of such crosses maintain the transgene expression characteristics of the original transformant. This strategy is used to ensure reliable gene expression in a number of varieties that are well adapted to local growing conditions.

It would be advantageous to be able to detect the presence of a particular event in order to determine whether progeny of a sexual cross contain a transgene of interest. In addition, a method for detecting a particular event would be helpful for complying with regulations requiring the pre-market approval and labeling of foods derived from recombinant crop plants, for example, or for use in environmental monitoring, monitoring traits in crops in the field, or monitoring products derived from a crop harvest, as well as for use in ensuring compliance of parties subject to regulatory or contractual terms.

It is possible to detect the presence of a transgene by any nucleic acid detection method known in the art including, but not limited to, the polymerase chain reaction (PCR) or DNA hybridization using nucleic acid probes. These detection methods generally focus on frequently used genetic elements, such as promoters, terminators, marker genes, etc., because for many DNA constructs, the coding region is interchangeable. As a result, such methods may not be useful for discriminating between different events, particularly those produced using the same DNA construct or very similar constructs unless the DNA sequence of the flanking DNA adjacent to the inserted heterologous DNA is known. For example, an event-specific PCR assay is described in U.S. Pat. No. 6,395,485 for the detection of elite event GAT-ZM1. Accordingly, it would be desirable to have a simple and discriminative method for the identification of event DAS-59122-7.

SUMMARY OF INVENTION

Embodiments of this invention relate to methods for producing and selecting an insect resistant monocot crop plant. More specifically, a DNA construct is provided that when expressed in plant cells and plants confers resistance to insects. According to one aspect of the invention, a DNA construct, capable of introduction into and replication in a host cell, is provided that when expressed in plant cells and plants confers insect resistance to the plant cells and plants.

3

The DNA construct is comprised of a DNA molecule named PHI17662A and it includes three (3) transgene expression cassettes. The first expression cassette comprises a DNA molecule which includes the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a B.t. δ -endotoxin identified as Cry34Ab1 (U.S. Pat. Nos. 6,127, 180, 6,624,145 and 6,340,593) operably connected to a DNA molecule comprising a Pin II transcriptional terminator isolated from potato (Gyheung An et al. (1989) *Plant Cell.* 1:115-122). The second transgene expression cassette of the DNA construct comprises a DNA molecule encoding the wheat peroxidase promoter (Hertig et al. (1991) *Plant Mol. Biol.* 16: 171-174) operably connected to a DNA molecule encoding a B.t. δ -endotoxin identified as Cry35Ab1 (U.S. Pat. Nos. 6,083,499, 6,548,291 and 6,340, 593) operably connected to a DNA molecule comprising a Pin II transcriptional terminator isolated from potato (Gyheung An et al. (1989) *Plant Cell.* 1:115-122). The third transgene expression cassette of the DNA construct comprises a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J. T. et al. (1985) *Nature* 313: 810-812; Mitsuhara et al. (1996) *Plant Cell Physiol.* 37: 49-59) operably connected to a DNA molecule encoding a phosphinothricin acetyltransferase (PAT) gene (Wohlleben W. et al. (1988) *Gene* 70:25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhara et al. (1996) *Plant Cell Physiol.* 37:49-59). Plants containing the DNA construct are also provided.

According to another embodiment of the invention, compositions and methods are provided for identifying a novel corn plant designated DAS-59122-7, which methods are based on primers or probes which specifically recognize the 5' and/or 3' flanking sequence of DAS-59122-7. DNA molecules are provided that comprise primer sequences that when utilized in a PCR reaction will produce amplicons unique to the transgenic event DAS-59122-7. These molecules may be selected from the group consisting of:

(SEQ ID NO: 1) 5' -GTGGCTCCTTCAACGTTGCGGTTCTGTC-3' ;

(SEQ ID NO: 2) 5' -CGTGCAAGCGCTCAATTGCGCCCTATAGTG-3' ;

(SEQ ID NO: 3) 5' -AATTGAGCGCTTGACGTTT-3' ;

(SEQ ID NO: 4) 5' -AACACAAGACCGGCCACACCTC-3' ;

(SEQ ID NO: 5) 5' -GAGGTGGTCTGGATGGTGTAGGTCA-3' ;

(SEQ ID NO: 6) 5' -TACAACCTCAAGTGGTTCTCTCCCGA-3' ;

(SEQ ID NO: 7) 5' -GAGGTCTGGATCTGCATGATGCGGA-3' ;

(SEQ ID NO: 8) 5' -AACCCCTTAGTATGTATTGTATT-3' ;

(SEQ ID NO: 9) 5' -CTCCTTCAACGTTGCGGTTCTGTCAG-3' ;

4

-continued

(SEQ ID NO: 10) 5' -TTTTGCAAAGCGAACGATTTCAGATG-3' ;

(SEQ ID NO: 11) 5' -GCGGGACAAGCCGTTTTACGTTT-3' ;

(SEQ ID NO: 12) 5' -GACGGGTGATTTATTTGATCTGCAC-3' ;

(SEQ ID NO: 13) 5' -CATCTGAATCGTTTCGCTTTGCAAAA-3' ;

(SEQ ID NO: 14) 5' -CTACGTTCCAATGGAGCTCGACTGTC-3' ;

(SEQ ID NO: 15) 5' -GGTCAAGTGGACACTTGGTCACTCA-3' ;

(SEQ ID NO: 16) 5' -GAGTGAAGAGATAAGCAAGTCAAAG-3' ;

(SEQ ID NO: 17) 5' -CATGTATACGTAAGTTTGGTGTCTGG-3' ;

(SEQ ID NO: 18) 5' -AATCCACAAGATTGGAGCAAACGAC-3' ;

(SEQ ID NO: 36) 5' -CGTATTACAATCGTACGCAATTCAG-3' ;

(SEQ ID NO: 37) 5' -GGATAAACAAACGGGACCATAGAAG-3' ;

and complements thereof. The corn plant and seed comprising these molecules is an embodiment of this invention. Further, kits utilizing these primer sequences for the identification of the DAS-59122-7 event are provided.

An additional embodiment of the invention relates to the specific flanking sequences of DAS-59122-7 described herein, which can be used to develop specific identification methods for DAS-59122-7 in biological samples. More particularly, the invention relates to the 5' and/or 3' flanking regions of DAS-59122-7, SEQ ID NO: 19, 5' flanking and SEQ ID NO: 20, 3' flanking, respectively, which can be used for the development of specific primers and probes. A further embodiment of the invention relates to identification methods for the presence of DAS-59122-7 in biological samples based on the use of such specific primers or probes.

According to another embodiment of the invention, methods of detecting the presence of DNA corresponding to the corn event DAS-59122-7 in a sample are provided. Such methods comprise: (a) contacting the sample comprising DNA with a DNA primer set, that when used in a nucleic acid amplification reaction with genomic DNA extracted from corn event DAS-59122-7 produces an amplicon that is diagnostic for corn event DAS-59122-7; (b) performing a nucleic acid amplification reaction, thereby producing the amplicon; and (c) detecting the amplicon.

DNA molecules that comprise the novel transgene/flanking insertion region, SEQ ID NO: 21, 5' flanking plus 1000 internal and SEQ ID NO: 22, 3' flanking plus 1000 internal and are homologous or complementary to SEQ ID NO: 21 and SEQ ID NO: 22 are an embodiment of this invention.

DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO: 21 are an embodiment of this invention. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant DAS-59122-7 of SEQ ID NO: 21 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant DAS-59122-7 are included.

5

In addition, DNA sequences that comprise the novel transgene/flanking insertion region, SEQ ID NO: 22 are provided. DNA sequences that comprise a sufficient length of polynucleotides of transgene insert sequence and a sufficient length of polynucleotides of maize genomic and/or flanking sequence from maize plant DAS-59122-7 of SEQ ID NO: 22 that are useful as primer sequences for the production of an amplicon product diagnostic for maize plant DAS-59122-7 are included.

According to another embodiment of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO: 21 or complements thereof, and a similar length of 5' flanking maize DNA sequence of SEQ ID NO: 21 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event DAS-59122-7. Therefore, embodiments of the invention also include the amplicons produced by DNA primers homologous or complementary to SEQ ID NO: 21.

According to another embodiment of the invention, the DNA sequences that comprise at least 11 or more nucleotides of the transgene portion of the DNA sequence of SEQ ID NO: 22 or complements thereof, and a similar length of 3' flanking maize DNA sequence of SEQ ID NO: 22 or complements thereof are useful as DNA primers in DNA amplification methods. The amplicons produced using these primers are diagnostic for maize event DAS-59122-7. Therefore, embodiments of the invention also include the amplicons produced by DNA primers homologous or complementary to SEQ ID NO: 22.

More specifically, a pair of DNA molecules comprising a DNA primer set, wherein the DNA molecules are identified as SEQ ID NO: 18 or complements thereof and SEQ ID NO: 1 or complements thereof; SEQ ID NO: 2 or complements thereof and SEQ ID NO: 17 or complements thereof; SEQ ID NO: 10 or complements thereof and SEQ ID NO: 9 or complements thereof; SEQ ID NO: 8 or complements thereof and SEQ ID NO: 17 or complements thereof; and SEQ ID NO: 36 or complements thereof and SEQ ID NO: 37 or complements thereof are embodiments of the invention.

Further embodiments of the invention include the amplicon comprising the DNA molecules of SEQ ID NO: 18 and SEQ ID NO: 1; the amplicon comprising the DNA molecules of SEQ ID NO: 2 and SEQ ID NO: 17; the amplicon comprising the DNA molecules of SEQ ID NO: 10 and SEQ ID NO: 9; the amplicon comprising the DNA molecules of SEQ ID NO: 8 and SEQ ID NO: 17; and the amplicon comprising the DNA molecules of SEQ ID NO: 36 and SEQ ID NO: 37.

Further embodiments of the invention include the following primers, which are useful in detecting or characterizing event DAS-59122-7: SEQ ID NO: 11 or complements thereof; SEQ ID NO: 5 or complements thereof; SEQ ID NO: 4 or complements thereof; SEQ ID NO: 7 or complements thereof; SEQ ID NO: 6 or complements thereof; SEQ ID NO: 3 or complements thereof; SEQ ID NO: 18 or complements thereof; SEQ ID NO: 14 or complements thereof; SEQ ID NO: 13 or complements thereof; SEQ ID NO: 15 or complements thereof; SEQ ID NO: 17 or complements thereof; SEQ ID NO: 16 or complements thereof; and SEQ ID NO: 12 or complements thereof. Further embodiments also include the amplicons produced by pairing any of the primers listed above.

According to another embodiment of the invention, methods of detecting the presence of a DNA molecule corresponding to the DAS-59122-7 event in a sample, such

6

methods comprising: (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe, molecule that hybridizes under stringent hybridization conditions with DNA extracted from corn event DAS-59122-7 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA. More specifically, a method for detecting the presence of a DNA molecule corresponding to the DAS-59122-7 event in a sample, such methods, consisting of (a) contacting the sample comprising DNA extracted from a corn plant with a DNA probe molecule that consists of sequences that are unique to the event, e.g. junction sequences, wherein said DNA probe molecule hybridizes under stringent hybridization conditions with DNA extracted from corn event DAS-59122-7 and does not hybridize under the stringent hybridization conditions with a control corn plant DNA; (b) subjecting the sample and probe to stringent hybridization conditions; and (c) detecting hybridization of the probe to the DNA.

In addition, a kit and methods for identifying event DAS-59122-7 in a biological sample which detects a DAS-59122-7 specific region within SEQ ID NO: 23 are provided. DNA molecules are provided that comprise at least one junction sequence of DAS-59122-7 selected from the group consisting of SEQ ID NO: 32, 33, 34, and 35 and complements thereof; wherein a junction sequence spans the junction between heterologous DNA inserted into the genome and the DNA from the corn cell flanking the insertion site, i.e. flanking DNA, and is diagnostic for the DAS-59122-7 event.

According to another embodiment of the invention, methods of producing an insect resistant corn plant that comprise the steps of: (a) sexually crossing a first parental corn line comprising the expression cassettes of the invention, which confers resistance to insects, and a second parental corn line that lacks insect resistance, thereby producing a plurality of progeny plants; and (b) selecting a progeny plant that is insect resistant. Such methods may optionally comprise the further step of back-crossing the progeny plant to the second parental corn line to producing a true-breeding corn plant that is insect resistant.

A further embodiment of the invention provides a method of producing a corn plant that is resistant to insects comprising transforming a corn cell with the DNA construct PHI17662A (SEQ ID NO: 24), growing the transformed corn cell into a corn plant, selecting the corn plant that shows resistance to insects, and further growing the corn plant into a fertile corn plant. The fertile corn plant can be self pollinated or crossed with compatible corn varieties to produce insect resistant progeny.

Another embodiment of the invention further relates to a DNA detection kit for identifying maize event DAS-59122-7 in biological samples. The kit comprises a first primer which specifically recognizes the 5' or 3' flanking region of DAS-59122-7, and a second primer which specifically recognizes a sequence within the foreign DNA of DAS-59122-7, or within the flanking DNA, for use in a PCR identification protocol. A further embodiment of the invention relates to a kit for identifying event DAS-59122-7 in biological samples, which kit comprises a specific probe having a sequence which corresponds or is complementary to, a sequence having between 80% and 100% sequence identity with a specific region of event DAS-59122-7. The sequence of the probe corresponds to a specific region comprising part of the 5' or 3' flanking region of event DAS-59122-7.

The methods and kits encompassed by the embodiments of the present invention can be used for different purposes such as, but not limited to the following: to identify event DAS-59122-7 in plants, plant material or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material; additionally or alternatively, the methods and kits can be used to identify transgenic plant material for purposes of segregation between transgenic and non-transgenic material; additionally or alternatively, the methods and kits can be used to determine the quality of plant material comprising maize event DAS-59122-7. The kits may also contain the reagents and materials necessary for the performance of the detection method. A further embodiment of this invention relates to the DAS-59122-7 corn plant or its parts, including, but not limited to, pollen, ovules, vegetative cells, the nuclei of pollen cells, and the nuclei of egg cells of the corn plant DAS-59122-7 and the progeny derived thereof. The corn plant and seed DAS-59122-7 from which the DNA primer molecules provide a specific amplicon product is an embodiment of the invention. The foregoing and other aspects of the invention will become more apparent from the following detailed description and accompanying drawing.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A-1E. DNA sequence (SEQ ID NO: 23) showing the transgenic insert PH17662A, as well as the sequences flanking the transgenic insert. The 5' and 3' border regions, bp 1 to bp 2593 and bp 9937 to bp 11922, respectively, are underlined. Two nucleotide differences (bp 6526 and bp 6562) based on comparison to the transforming plasmid PHP17662 are noted in bold and underlined.

FIG. 2. Schematic diagram of the B.t. Cry34/35Ab1 event DAS-59122-7 insert region is divided into three separate sections; the 5' border region with corn genomic DNA, the intact T-DNA insert, and the 3' border region with corn genomic DNA. The two arrows beneath the diagram of the insert indicate the start and end points of the sequence derived from 5' and 3' genome walking fragments. Other boxes beneath the diagram of the insert represent PCR fragments that were amplified from genomic DNA of event DAS-59122-7 and sequenced to cover the intact T-DNA insert and the 5' and 3' insert/border junction regions.

FIG. 3. Schematic diagram of the B.t. Cry34/35Ab1 event DAS-59122-7 insert region is divided into three separate sections; the 5' border region with corn genomic DNA, the intact T-DNA insert, and the 3' border region with corn genomic DNA. Boxes beneath the diagram of the insert represent PCR fragments located in either the genomic border regions or across the 5' and 3' junction regions of the T-DNA insert with corn genomic DNA that were amplified from genomic DNA from event DAS-59122-7.

DETAILED DESCRIPTION

The following definitions and methods are provided to better define the present invention and to guide those of ordinary skill in the art in the practice of the present invention. Unless otherwise noted, terms are to be understood according to conventional usage by those of ordinary skill in the relevant art. Definitions of common terms in molecular biology may also be found in Rieger et al., *Glossary of Genetics: Classical and Molecular*, 5th edition, Springer-Verlag; New York, 1991; and Lewin, *Genes V*, Oxford University Press; New York, 1994. The nomenclature for DNA bases as set forth at 37 CFR §1.822 is used.

As used herein, the term "comprising" means "including but not limited to".

As used herein, the term "corn" means *Zea mays* or maize and includes all plant varieties that can be bred with corn, including wild maize species.

As used herein, the term "DAS-59122-7 specific" refers to a nucleotide sequence which is suitable for discriminatively identifying event DAS-59122-7 in plants, plant material, or in products such as, but not limited to, food or feed products (fresh or processed) comprising, or derived from plant material.

As used herein, the terms "insect resistant" and "impacting insect pests" refers to effecting changes in insect feeding, growth, and/or behavior at any stage of development, including but not limited to: killing the insect; retarding growth; preventing reproductive capability; inhibiting feeding; and the like.

As used herein, the terms "pesticidal activity" and "insecticidal activity" are used synonymously to refer to activity of an organism or a substance (such as, for example, a protein) that can be measured by numerous parameters including, but not limited to, pest mortality, pest weight loss, pest attraction, pest repellency, and other behavioral and physical changes of a pest after feeding on and/or exposure to the organism or substance for an appropriate length of time. For example "pesticidal proteins" are proteins that display pesticidal activity by themselves or in combination with other proteins.

"Coding sequence" refers to a nucleotide sequence that codes for a specific amino acid sequence. As used herein, the terms "encoding" or "encoded" when used in the context of a specified nucleic acid mean that the nucleic acid comprises the requisite information to guide translation of the nucleotide sequence into a specified protein. The information by which a protein is encoded is specified by the use of codons. A nucleic acid encoding a protein may comprise non-translated sequences (e.g., introns) within translated regions of the nucleic acid or may lack such intervening non-translated sequences (e.g., as in cDNA).

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. "Foreign" refers to material not normally found in the location of interest. Thus "foreign DNA" may comprise both recombinant DNA as well as newly introduced, rearranged DNA of the plant. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure. The site in the plant genome where a recombinant DNA has been inserted may be referred to as the "insertion site" or "target site".

As used herein, "insert DNA" refers to the heterologous DNA within the expression cassettes used to transform the plant material while "flanking DNA" can exist of either

genomic DNA naturally present in an organism such as a plant, or foreign (heterologous) DNA introduced via the transformation process which is extraneous to the original insert DNA molecule, e.g. fragments associated with the transformation event. A “flanking region” or “flanking sequence” as used herein refers to a sequence of at least twenty (20) base pair, preferably at least fifty (50) base pair, and up to five thousand (5000) base pair which is located either immediately upstream of and contiguous with or immediately downstream of and contiguous with the original foreign insert DNA molecule. Transformation procedures leading to random integration of the foreign DNA will result in transformants containing different flanking regions characteristic and unique for each transformant. When recombinant DNA is introduced into a plant through traditional crossing, its flanking regions will generally not be changed. Transformants will also contain unique junctions between a piece of heterologous insert DNA and genomic DNA, or two (2) pieces of genomic DNA, or two (2) pieces of heterologous DNA. A “junction” is a point where two (2) specific DNA fragments join. For example, a junction exists where insert DNA joins flanking DNA. A junction point also exists in a transformed organism where two (2) DNA fragments join together in a manner that is modified from that found in the native organism. “Junction DNA” refers to DNA that comprises a junction point.

As used herein, “heterologous” in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous nucleotide sequence can be from a species different from that from which the nucleotide sequence was derived, or, if from the same species, the promoter is not naturally found operably linked to the nucleotide sequence. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

“Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements are often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence that can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters that cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types

useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamoto and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect numerous parameters including, processing of the primary transcript to mRNA, mRNA stability and/or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The “3' non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

A “protein” or “polypeptide” is a chain of amino acids arranged in a specific order determined by the coding sequence in a polynucleotide encoding the polypeptide.

A DNA construct is an assembly of DNA molecules linked together that provide one or more expression cassettes. The DNA construct may be a plasmid that is enabled for self replication in a bacterial cell and contains various endonuclease enzyme restriction sites that are useful for introducing DNA molecules that provide functional genetic elements, i.e., promoters, introns, leaders, coding sequences, 3' termination regions, among others; or a DNA construct may be a linear assembly of DNA molecules, such as an expression cassette. The expression cassette contained within a DNA construct comprise the necessary genetic elements to provide transcription of a messenger RNA. The expression cassette can be designed to express in prokaryote cells or eukaryotic cells. Expression cassettes of the embodiments of the present invention are designed to express in plant cells.

The DNA molecules of embodiments of the invention are provided in expression cassettes for expression in an organism of interest. The cassette will include 5' and 3' regulatory sequences operably linked to a coding sequence. “Operably linked” means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. Operably linked is intended to indicate a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes or multiple DNA constructs.

The expression cassette will include in the 5' to 3' direction of transcription: a transcriptional and translational initiation region, a coding region, and a transcriptional and translational termination region functional in the organism serving as a host. The transcriptional initiation region (i.e., the promoter) may be native or analogous, or foreign or heterologous to the host organism. Additionally, the pro-

motor may be the natural sequence or alternatively a synthetic sequence. The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation.

It is to be understood that as used herein the term "transgenic" includes any cell, cell line, callus, tissue, plant part, or plant, the genotype of which has been altered by the presence of a heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term "transgenic" as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

A transgenic "event" is produced by transformation of plant cells with a heterologous DNA construct(s), including a nucleic acid expression cassette that comprises a transgene of interest, the regeneration of a population of plants resulting from the insertion of the transgene into the genome of the plant, and selection of a particular plant characterized by insertion into a particular genome location. An event is characterized phenotypically by the expression of the transgene. At the genetic level, an event is part of the genetic makeup of a plant. The term "event" also refers to progeny produced by a sexual outcross between the transformant and another variety that include the heterologous DNA. Even after repeated back-crossing to a recurrent parent, the inserted DNA and flanking DNA from the transformed parent is present in the progeny of the cross at the same chromosomal location. The term "event" also refers to DNA from the original transformant comprising the inserted DNA and flanking sequence immediately adjacent to the inserted DNA that would be expected to be transferred to a progeny that receives inserted DNA including the transgene of interest as the result of a sexual cross of one parental line that includes the inserted DNA (e.g., the original transformant and progeny resulting from selfing) and a parental line that does not contain the inserted DNA.

An insect resistant DAS-59122-7 corn plant can be bred by first sexually crossing a first parental corn plant consisting of a corn plant grown from the transgenic DAS-59122-7 corn plant and progeny thereof derived from transformation with the expression cassettes of the embodiments of the present invention that confers insect resistance, and a second parental corn plant that lacks insect resistance, thereby producing a plurality of first progeny plants; and then selecting a first progeny plant that is resistant to insects; and selfing the first progeny plant, thereby producing a plurality of second progeny plants; and then selecting from the second progeny plants an insect resistant plant. These steps can further include the back-crossing of the first insect resistant progeny plant or the second insect resistant progeny plant to the second parental corn plant or a third parental corn plant, thereby producing a corn plant that is resistant to insects.

As used herein, the term "plant" includes reference to whole plants, plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, and progeny of same. Parts of transgenic plants understood to be within the scope of the invention comprise, for example, plant cells, protoplasts, tissues, callus, embryos as well as flowers, stems, fruits, leaves, and roots originating in transgenic plants or their progeny previously transformed with a DNA molecule of the invention

and therefore consisting at least in part of transgenic cells, are also an embodiment of the present invention.

As used herein, the term "plant cell" includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledonous and dicotyledonous plants.

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or "gene gun" transformation technology (Klein et al. (1987) *Nature* (London) 327:70-73; U.S. Pat. No. 4,945,050, incorporated herein by reference). Additional transformation methods are disclosed below.

Thus, isolated polynucleotides of the invention can be incorporated into recombinant constructs, typically DNA constructs, which are capable of introduction into and replication in a host cell. Such a construct can be a vector that includes a replication system and sequences that are capable of transcription and translation of a polypeptide-encoding sequence in a given host cell. A number of vectors suitable for stable transfection of plant cells or for the establishment of transgenic plants have been described in, e.g., Pouwels et al., (1985; Supp. 1987) *Cloning Vectors: A Laboratory Manual*, Weissbach and Weissbach (1989) *Methods for Plant Molecular Biology*, (Academic Press, New York); and Flevin et al., (1990) *Plant Molecular Biology Manual*, (Kluwer Academic Publishers). Typically, plant expression vectors include, for example, one or more cloned plant genes under the transcriptional control of 5' and 3' regulatory sequences and a dominant selectable marker. Such plant expression vectors also can contain a promoter regulatory region (e.g., a regulatory region controlling inducible or constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

It is also to be understood that two different transgenic plants can also be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated, as is vegetative propagation. Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several references, e.g., Fehr, in *Breeding Methods for Cultivar Development*, Wilcos J. ed., American Society of Agronomy, Madison Wis. (1987).

A "probe" is an isolated nucleic acid to which is attached a conventional detectable label or reporter molecule, e.g., a radioactive isotope, ligand, chemiluminescent agent, or enzyme. Such a probe is complementary to a strand of a target nucleic acid, in the case of the present invention, to a strand of isolated DNA from corn event DAS-59122-7 whether from a corn plant or from a sample that includes DNA from the event. Probes according to the present invention include not only deoxyribonucleic or ribonucleic acids but also polyamides and other probe materials that

bind specifically to a target DNA sequence and can be used to detect the presence of that target DNA sequence.

“Primers” are isolated nucleic acids that are annealed to a complementary target DNA strand by nucleic acid hybridization to form a hybrid between the primer and the target DNA strand, then extended along the target DNA strand by a polymerase, e.g., a DNA polymerase. Primer pairs of the invention refer to their use for amplification of a target nucleic acid sequence, e.g., by the polymerase chain reaction (PCR) or other conventional nucleic-acid amplification methods. “PCR” or “polymerase chain reaction” is a technique used for the amplification of specific DNA segments (see, U.S. Pat. Nos. 4,683,195 and 4,800,159; herein incorporated by reference).

Probes and primers are of sufficient nucleotide length to bind to the target DNA sequence specifically in the hybridization conditions or reaction conditions determined by the operator. This length may be of any length that is of sufficient length to be useful in a detection method of choice. Generally, eleven (11) nucleotides or more in length, eighteen (18) nucleotides or more, and twenty-two (22) nucleotides or more, are used. Such probes and primers hybridize specifically to a target sequence under high stringency hybridization conditions. Probes and primers according to embodiments of the present invention may have complete DNA sequence similarity of contiguous nucleotides with the target sequence, although probes differing from the target DNA sequence and that retain the ability to hybridize to target DNA sequences may be designed by conventional methods. Probes can be used as primers, but are generally designed to bind to the target DNA or RNA and are not used in an amplification process.

Specific primers can be used to amplify an integration fragment to produce an amplicon that can be used as a “specific probe” for identifying event DAS-59122-7 in biological samples. When the probe is hybridized with the nucleic acids of a biological sample under conditions which allow for the binding of the probe to the sample, this binding can be detected and thus allow for an indication of the presence of event DAS-59122-7 in the biological sample. Such identification of a bound probe has been described in the art. In an embodiment of the invention the specific probe is a sequence which, under optimized conditions, hybridizes specifically to a region within the 5' or 3' flanking region of the event and also comprises a part of the foreign DNA contiguous therewith. The specific probe may comprise a sequence of at least 80%, between 80 and 85%, between 85 and 90%, between 90 and 95%, and between 95 and 100% identical (or complementary) to a specific region of the event.

Methods for preparing and using probes and primers are described, for example, in *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. 1989 (hereinafter, “Sambrook et al., 1989”); *Current Protocols in Molecular Biology*, ed. Ausubel et al., Greene Publishing and Wiley-Interscience, New York, 1992 (with periodic updates) (hereinafter, “Ausubel et al., 1992”); and Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press: San Diego, 1990. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as the PCR primer analysis tool in Vector NTI version 6 (Informax Inc., Bethesda Md.); PrimerSelect (DNASTAR Inc., Madison, Wis.); and Primer (Version 0.5©, 1991, Whitehead Institute for Biomedical Research, Cambridge, Mass.).

Additionally, the sequence can be visually scanned and primers manually identified using guidelines known to one of skill in the art.

A “kit” as used herein refers to a set of reagents for the purpose of performing the method embodiments of the invention, more particularly, the identification of the event DAS-59122-7 in biological samples. The kit of the invention can be used, and its components can be specifically adjusted, for purposes of quality control (e.g. purity of seed lots), detection of event DAS-59122-7 in plant material, or material comprising or derived from plant material, such as but not limited to food or feed products. “Plant material” as used herein refers to material which is obtained or derived from a plant.

Primers and probes based on the flanking DNA and insert sequences disclosed herein can be used to confirm (and, if necessary, to correct) the disclosed sequences by conventional methods, e.g., by re-cloning and sequencing such sequences. The nucleic acid probes and primers of the present invention hybridize under stringent conditions to a target DNA sequence. Any conventional nucleic acid hybridization or amplification method can be used to identify the presence of DNA from a transgenic event in a sample. Nucleic acid molecules or fragments thereof are capable of specifically hybridizing to other nucleic acid molecules under certain circumstances. As used herein, two nucleic acid molecules are said to be capable of specifically hybridizing to one another if the two molecules are capable of forming an anti-parallel, double-stranded nucleic acid structure.

A nucleic acid molecule is said to be the “complement” of another nucleic acid molecule if they exhibit complete complementarity. As used herein, molecules are said to exhibit “complete complementarity” when every nucleotide of one of the molecules is complementary to a nucleotide of the other. Two molecules are said to be “minimally complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under at least conventional “low-stringency” conditions. Similarly, the molecules are said to be “complementary” if they can hybridize to one another with sufficient stability to permit them to remain annealed to one another under conventional “high-stringency” conditions. Conventional stringency conditions are described by Sambrook et al., 1989, and by Haymes et al., In: *Nucleic Acid Hybridization, a Practical Approach*, IRL Press, Washington, D.C. (1985), departures from complete complementarity are therefore permissible, as long as such departures do not completely preclude the capacity of the molecules to form a double-stranded structure. In order for a nucleic acid molecule to serve as a primer or probe it need only be sufficiently complementary in sequence to be able to form a stable double-stranded structure under the particular solvent and salt concentrations employed.

In hybridization reactions, specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. The thermal melting point (T_m) is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284; $T_m = 81.5^\circ \text{C} + 16.6 (\log M) + 0.41 (\% \text{GC}) - 0.61 (\% \text{form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution,

and L is the length of the hybrid in base pairs. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with >90% identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the T_m for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4° C. lower than the T_m ; moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10° C. lower than the T_m ; low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20° C. lower than the T_m .

Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel et al., eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.).

As used herein, a substantially homologous sequence is a nucleic acid molecule that will specifically hybridize to the complement of the nucleic acid molecule to which it is being compared under high stringency conditions. Appropriate stringency conditions which promote DNA hybridization, for example, 6× sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2×SSC at 50° C., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of a destabilizing agent such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C. A nucleic acid of the invention may specifically hybridize to one or more of the nucleic acid molecules unique to the DAS-59122-7 event or complements thereof or fragments of either under moderately stringent conditions.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology

algorithm of Smith et al. (1981) *Adv. Appl. Math.* 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, Calif.); the ALIGN program (Version 2.0); the ALIGN PLUS program (version 3.0, copyright 1997); and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 10 (available from Accelrys, 9685 Scranton Road, San Diego, Calif. 92121, USA). Alignments using these programs can be performed using the default parameters.

The CLUSTAL program is well described by Higgins and Sharp, *Gene* 73:237-244 (1988); Higgins and Sharp, *CABIOS* 5:151-153 (1989); Corpet, et al., *Nucleic Acids Research* 16:10881-90 (1988); Huang, et al., *Computer Applications in the Biosciences* 8:155-65 (1992), and Pearson, et al., *Methods in Molecular Biology* 24:307-331 (1994). The ALIGN and the ALIGN PLUS programs are based on the algorithm of Myers and Miller (1988) supra. The BLAST programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) supra. The BLAST family of programs which can be used for database similarity searches includes: BLASTN for nucleotide query sequences against nucleotide database sequences; BLASTX for nucleotide query sequences against protein database sequences; BLASTP for protein query sequences against protein database sequences; TBLASTN for protein query sequences against nucleotide database sequences; and TBLASTX for nucleotide query sequences against nucleotide database sequences. See, *Current Protocols in Molecular Biology*, Chapter 19, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (1995). Alignment may also be performed manually by visual inspection.

To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) supra. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See www.ncbi.nlm.nih.gov.

As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of

the substitution. Sequences that differ by such conservative substitutions are said to have “sequence similarity” or “similarity”. Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, Calif.).

As used herein, “percentage of sequence identity” means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

Regarding the amplification of a target nucleic acid sequence (e.g., by PCR) using a particular amplification primer pair, “stringent conditions” are conditions that permit the primer pair to hybridize only to the target nucleic-acid sequence to which a primer having the corresponding wild-type sequence (or its complement) would bind and preferably to produce a unique amplification product, the amplicon, in a DNA thermal amplification reaction.

The term “specific for (a target sequence)” indicates that a probe or primer hybridizes under stringent hybridization conditions only to the target sequence in a sample comprising the target sequence.

As used herein, “amplified DNA” or “amplicon” refers to the product of nucleic acid amplification of a target nucleic acid sequence that is part of a nucleic acid template. For example, to determine whether a corn plant resulting from a sexual cross contains transgenic event genomic DNA from the corn plant of the invention, DNA extracted from the corn plant tissue sample may be subjected to a nucleic acid amplification method using a DNA primer pair that includes a first primer derived from flanking sequence adjacent to the insertion site of inserted heterologous DNA, and a second primer derived from the inserted heterologous DNA to produce an amplicon that is diagnostic for the presence of the event DNA. Alternatively, the second primer may be derived from the flanking sequence. The amplicon is of a length and has a sequence that is also diagnostic for the event. The amplicon may range in length from the combined length of the primer pairs plus one nucleotide base pair to any length of amplicon producible by a DNA amplification protocol. Alternatively, primer pairs can be derived from flanking sequence on both sides of the inserted DNA so as to produce an amplicon that includes the entire insert nucleotide sequence of the PH117662A expression construct as well as the sequence flanking the transgenic insert, see FIG. 1 (SEQ ID NO: 23), approximately twelve (12) Kb in size. A member of a primer pair derived from the flanking sequence may be located a distance from the inserted DNA sequence, this distance can range from one nucleotide base pair up to the limits of the amplification reaction, or about twenty thousand nucleotide base pairs. The use of the term

“amplicon” specifically excludes primer dimers that may be formed in the DNA thermal amplification reaction.

Nucleic acid amplification can be accomplished by any of the various nucleic acid amplification methods known in the art, including the polymerase chain reaction (PCR). A variety of amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis et al., Academic press, San Diego, 1990. PCR amplification methods have been developed to amplify up to 22 Kb of genomic DNA and up to 42 Kb of bacteriophage DNA (Cheng et al., *Proc. Natl. Acad. Sci. USA* 91:5695-5699, 1994). These methods as well as other methods known in the art of DNA amplification may be used in the practice of the embodiments of the present invention. It is understood that a number of parameters in a specific PCR protocol may need to be adjusted to specific laboratory conditions and may be slightly modified and yet allow for the collection of similar results. These adjustments will be apparent to a person skilled in the art.

The amplicon produced by these methods may be detected by a plurality of techniques, including, but not limited to, Genetic Bit Analysis (Nikiforov, et al. *Nucleic Acid Res.* 22:4167-4175, 1994) where a DNA oligonucleotide is designed which overlaps both the adjacent flanking DNA sequence and the inserted DNA sequence. The oligonucleotide is immobilized in wells of a microwell plate. Following PCR of the region of interest (using one primer in the inserted sequence and one in the adjacent flanking sequence) a single-stranded PCR product can be hybridized to the immobilized oligonucleotide and serve as a template for a single base extension reaction using a DNA polymerase and labeled ddNTPs specific for the expected next base. Readout may be fluorescent or ELISA-based. A signal indicates presence of the insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Another detection method is the Pyrosequencing technique as described by Winge (*Innov. Pharma. Tech.* 00:18-24, 2000). In this method an oligonucleotide is designed that overlaps the adjacent DNA and insert DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted sequence and one in the flanking sequence) and incubated in the presence of a DNA polymerase, ATP, sulfurylase, luciferase, apyrase, adenosine 5' phosphosulfate and luciferin. dNTPs are added individually and the incorporation results in a light signal which is measured. A light signal indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single or multi-base extension.

Fluorescence Polarization as described by Chen et al., (*Genome Res.* 9:492-498, 1999) is also a method that can be used to detect an amplicon of the invention. Using this method an oligonucleotide is designed which overlaps the flanking and inserted DNA junction. The oligonucleotide is hybridized to a single-stranded PCR product from the region of interest (one primer in the inserted DNA and one in the flanking DNA sequence) and incubated in the presence of a DNA polymerase and a fluorescent-labeled ddNTP. Single base extension results in incorporation of the ddNTP. Incorporation can be measured as a change in polarization using a fluorometer. A change in polarization indicates the presence of the transgene insert/flanking sequence due to successful amplification, hybridization, and single base extension.

Taqman® (PE Applied Biosystems, Foster City, Calif.) is described as a method of detecting and quantifying the presence of a DNA sequence and is fully understood in the instructions provided by the manufacturer. Briefly, a FRET oligonucleotide probe is designed which overlaps the flanking and insert DNA junction. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking genomic sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Hybridization of the FRET probe results in cleavage and release of the fluorescent moiety away from the quenching moiety on the FRET probe. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

Molecular Beacons have been described for use in sequence detection as described in Tyangi et al. (*Nature Biotech.* 14:303-308, 1996). Briefly, a FRET oligonucleotide probe is designed that overlaps the flanking and insert DNA junction. The unique structure of the FRET probe results in it containing secondary structure that keeps the fluorescent and quenching moieties in close proximity. The FRET probe and PCR primers (one primer in the insert DNA sequence and one in the flanking sequence) are cycled in the presence of a thermostable polymerase and dNTPs. Following successful PCR amplification, hybridization of the FRET probe to the target sequence results in the removal of the probe secondary structure and spatial separation of the fluorescent and quenching moieties. A fluorescent signal results. A fluorescent signal indicates the presence of the flanking/transgene insert sequence due to successful amplification and hybridization.

A hybridization reaction using a probe specific to a sequence found within the amplicon is yet another method used to detect the amplicon produced by a PCR reaction.

Embodiments of the present invention are further defined in the following Examples. It should be understood that these Examples are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the embodiments of the invention to adapt it to various usages and conditions. Thus, various modifications of the embodiments of the invention, in addition to those shown and described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

The disclosure of each reference set forth herein is incorporated herein by reference in its entirety.

EXAMPLES

Example 1. Transformation of Maize by *Agrobacterium* Transformation and Regeneration of Transgenic Plants Containing the Cry34Ab1 and Cry35Ab1 (Cry34/35Ab1) Genes

A DNA molecule of approximately 7.4 Kb, designated PH117662A (SEQ ID NO: 24), which includes a first transgene expression cassette comprising a DNA molecule which includes the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (Ubi-1) gene (Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689 and Christensen and Quail (1996) *Transgenic Res.* 5:213-218) operably connected to a DNA molecule encoding a B.t. δ -endotoxin identified as Cry34Ab1 (U.S. Pat. Nos. 6,127,180, 6,624,145 and 6,340,593) operably connected to a DNA molecule

comprising a Pin II transcriptional terminator isolated from potato (Gyheung An et al. (1989) *Plant Cell.* 1:115-122). The second transgene expression cassette of the DNA construct comprises a DNA molecule encoding the wheat peroxidase promoter (Hertig et al. (1991) *Plant Mol. Biol.* 16:171-174) operably connected to a DNA molecule encoding a B.t. δ -endotoxin identified as Cry35Ab1 (U.S. Pat. Nos. 6,083,499, 6,548,291 and 6,340,593) operably connected to a DNA molecule comprising a Pin II transcriptional terminator isolated from potato (Gyheung An et al. (1989) *Plant Cell.* 1:115-122). The third transgene expression cassette of the DNA construct comprises a DNA molecule of the cauliflower mosaic virus (CaMV) 35S promoter (Odell J. T. et al. (1985) *Nature* 313:810-812; Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37:49-59) operably connected to a DNA molecule encoding a phosphinothricin acetyltransferase (PAT) gene (Wohlleben W. et al. (1988) *Gene* 70:25-37) operably connected to a DNA molecule comprising a 3' transcriptional terminator from (CaMV) 35S (see Mitsuhashi et al. (1996) *Plant Cell Physiol.* 37: 49-59) was used to transform maize embryo tissue.

B.t. Cry34/35 Ab1 maize plants were obtained by *Agrobacterium* transformation, the method of Zhao was employed (U.S. Pat. No. 5,981,840, and PCT patent publication WO98/32326; the contents of which are hereby incorporated by reference). Briefly, immature embryos were isolated from maize and the embryos contacted with a suspension of *Agrobacterium*, where the bacteria was capable of transferring PH117662 DNA (SEQ ID NO: 24) to at least one cell of at least one of the immature embryos (step 1: the infection step). Specifically, in this step the immature embryos were immersed in an *Agrobacterium* suspension for the initiation of inoculation. The embryos were co-cultured for a time with the *Agrobacterium* (step 2: the co-cultivation step). Specifically, the immature embryos were cultured on solid medium following the infection step. Following this co-cultivation period a "resting" step was provided. In this resting step, the embryos were incubated in the presence of at least one antibiotic known to inhibit the growth of *Agrobacterium* without the addition of a selective agent for plant transformants (step 3: resting step). In particular, the immature embryos are cultured on solid medium with antibiotic, but without a selecting agent, for elimination of *Agrobacterium* and for a resting phase for the infected cells. Next, inoculated embryos were cultured on medium containing a selective agent and growing transformed callus was recovered (step 4: the selection step). Specifically, the immature embryos were cultured on solid medium with a selective agent resulting in the selective growth of transformed cells. The callus was then regenerated into plants (step 5: the regeneration step), and, specifically, calli grown on selective medium were cultured on solid medium to regenerate the plants. Individual embryos were kept physically separate during culture, and the majority of explants died on the selective medium.

Those embryos that survived and produced healthy, glufosinate-resistant callus tissue were assigned unique identification codes representing putative transformation events, and continually transferred to fresh selection medium. Plants were regenerated from tissue derived from each unique event and transferred to the greenhouse. Leaf samples were taken for molecular analysis to verify the presence of the transgene by PCR and to confirm expression of the Cry34/35Ab1 protein by ELISA. Plants were then subjected to a whole plant bioassay using western corn rootworm insects. Positive plants were crossed with inbred lines to obtain seed from the initial transformed plants. A number of lines were

21

evaluated in the field. The DAS-59122-7 event was selected from a population of independent transgenic events based on a superior combination of characteristics, including insect resistance and agronomic performance.

Example 2. Identification of *Bacillus thuringiensis* Cry34/35Ab1 Maize Line DAS-59122-7

Seed from event DAS-59122-7 was evaluated. The T1S2 seed represents transformation into the Hi-II background, followed by a cross with inbred line PH09B and two rounds of self-crossing. All seed were obtained from Pioneer Hi-Bred (Johnston, Iowa). Primary characterization was conducted on plant leaf tissue during the study by confirmation of phosphinothricin acetyltransferase (PAT) activity via herbicide leaf painting and Cry34Ab1 expression using lateral flow devices.

Control substances in this study were defined as unmodified seed representative of the test substance background. Control seeds of Hi-II and PH09B backgrounds were used as negative controls. These unmodified seed do not contain the plant transcription units for the cry34Ab1, cry35Ab1, and pat genes. All seed were obtained from Pioneer Hi-Bred (Johnston, Iowa).

DNA samples from two additional B.t. Cry34/35Ab1 events, event DAS-45214-4 and event DAS-45216-6, were used as negative controls for event specific PCR analysis. The two events were produced through *Agrobacterium* transformation using the same vector used to produce event DAS-59122-7 and therefore contained the plant transcription units for the cry34Ab1, cry35Ab1, and pat genes. However, the insertion sites of the T-DNA in events DAS-45214-4 and DAS-45216-6, including genomic DNA border regions, were different from that in event DAS-59122-7. DNA samples from event DAS-45214-4 and event DAS-45216-6 were isolated and characterized by Southern blot analysis. (Data not shown.)

Corn seed for event DAS-59122-7 and unmodified control seed (Hi-II and PH09B) were planted in growth chambers at the DuPont Experimental Station (Wilmington, Del.) to produce sufficient numbers of plants for DNA analysis. For characterization of event DAS-59122-7, ten (10) T1 S2 seeds were planted. Ten (10) seeds were also planted for each unmodified control line. One (1) seed was planted per pot, and the pot was uniquely identified. Planting and

22

growing conditions were conducive to healthy plant growth including regulated light and water.

Leaf samples were collected for each of the control and event DAS-59122-7 plants. For each sample, sufficient leaf material from above the growing point was collected and placed in a pre-labeled sample bag. The samples were placed on dry ice and were transferred to an ultralow freezer following collection. All samples were maintained frozen until processing. All leaf samples were uniquely labeled with the plant identifier and the date of harvest.

To confirm the expression of the Cry34Ab1 protein in event DAS-59122-7 and the absence of expression in the controls, leaf samples were collected from all event DAS-59122-7 and control plants, and screened for transgenic protein using lateral flow devices specific for Cry34Ab1 (Strategic Diagnostics, Inc., Newark, Del.). Leaf punches were taken from each plant and ground in a phosphate buffered saline solution with Tween 20 to crudely extract the protein. A strip device was dipped into the extract to determine the presence or absence of the Cry34Ab1 protein. The immunoassay results were used to confirm the identity of the test substance plants prior to molecular analysis as shown in Table 1.

To confirm the expression of phosphinothricin acetyltransferase (PAT) in event DAS-59122-7 plants, herbicide leaf painting was conducted. All plants used in this study were leaf painted to confirm plant identity. Plants were assayed prior to the R1 growth stage. Assays were conducted following a standard procedure known in the art for herbicide leaf painting for the identification of PAT-expressing transgenic plants. Specifically, a portion of one leaf of each plant was treated with approximately 2% solution of glufosinate herbicide, Basta® (Bayer CropScience) in water and visually checked for brown or necrotic tissue in the painted leaf area 4-12 days after application. Results for each plant were recorded and used to determine expression of PAT in each test plant as shown in Table 1. As shown in Table 1, of the ten (10) plants tested for event DAS-59122-7 T1S2 generation, six (6) plants expressed both Cry34Ab1 and PAT, while four (4) plants did not express either protein. All unmodified controls tested negative for both CryAb1 and PAT assays (data not shown).

TABLE 1

Cry34Ab1 and PAT Protein Expression and Southern Hybridization
Data for B.t. Cry34/35Ab1 Event DAS-59122-7

Plant ID	Sample ID	Cry34Ab1 and PAT Expression ¹	Southern Blot cry34Ab1 Probe ²	Southern Blot cry35Ab1 Probe ²	Southern Blot pat Probe ²
02-122C 1	DAS59122-7 T1S2 1	positive	+	+	+
02-122C 2	DAS59122-7 T1S2 2	positive	+	+	+
02-122C 3	DAS59122-7 T1S2 3	positive	+	+	+
02-122C 4	DAS59122-7 T1S2 4	negative	-	-	-
02-122C 5	DAS59122-7 T1S2 5	positive	+	+	+
02-122C 6	DAS59122-7 T1S2 6	negative	-	-	-
02-122C 7	DAS59122-7 T1S2 7	positive	+	+	+

TABLE 1-continued

Cry34Ab1 and PAT Protein Expression and Southern Hybridization Data for B.t. Cry34/35Ab1 Event DAS-59122-7					
Plant ID	Sample ID	Cry34Ab1 and PAT Expression ¹	Southern Blot cry34Ab1 Probe ²	Southern Blot cry35Ab1 Probe ²	Southern Blot pat Probe ²
02-122C 8	DAS59122-7 T1S2 8	negative	-	-	-
02-122C 9	DAS59122-7 T1S2 9	negative	-	-	-
02-122C 10	DAS59122-7 T1S2 10	positive	+	+	+

¹Positive Cry34Ab1 expression indicates detection of protein expression as determined by the immunoassay-based lateral flow device specific for Cry34Ab1 protein detection. Negative indicates no detection of the Cry34Ab1 protein. Positive PAT expression indicates plants that were tolerant to the herbicide treatment and negative indicates plants that were sensitive to the herbicide.

²+ indicates hybridization signal on Southern blot; - indicates no hybridization signal on Southern blot. The cry34Ab1 gene probe hybridized to the expected internal T-DNA fragment of 1.915 kb, the cry35Ab1 gene probe hybridized to the expected internal T-DNA fragment of 2.607 kb, and the pat gene probe hybridized to a 3.4 kb border fragment consistent with a single intact T-DNA insertion as determined by Southern blot analysis.

Example 3. Southern Blot Analysis of *Bacillus thuringiensis* Cry34/35Ab1 Maize Line DAS-59122-7

One gram quantities of leaf samples were ground under liquid nitrogen, and the genomic DNA was isolated using DNeasy® Plant Mini Kit (Qiagen, Valencia, Calif.) or using a standard Urea Extraction Buffer procedure. Following extraction, the DNA was visualized on an agarose gel to determine the DNA quality, and was quantified using Pico Green® reagent (Molecular Probes, Inc., Eugene, Oreg.) and spectrofluorometric analysis.

The 1 Kb DNA Ladder (Invitrogen, Carlsbad, Calif.) was used to estimate DNA fragment sizes on agarose gels.

Genomic DNA isolated from event DAS-59122-7 plants was digested with Nco I and electrophoretically separated, transferred to nylon membranes, and hybridized to the cry34Ab1, cry35Ab1 and pat gene probes using standard procedures known in the art. Blots were exposed to X-ray film for one or more time periods to detect hybridizing fragments and to visualize molecular weight standards. Images were then digitally captured by photographing X-ray films and/or by detection with a Lumi-Imager™ instrument (Roche, Indianapolis, Ind.). The sizes of detected bands were documented for each probe. Southern blot analysis was used as a means of verifying the presence of the insertion in the test plants and confirming that all plants from event DAS-59122-7 contained the same insertion as shown in Table 1. (Southern blots not shown.) Southern blot analysis indicated that event DAS-59122-7 contained a single insertion consisting of an intact copy of the T-DNA region from plasmid PHP17662, while the null segregants, as determined by the protein expression analysis did not hybridize to the gene probes. Further, event DAS-59122-7 plants expressing the two proteins exhibited identical hybridization patterns on Southern blots (data not shown). Specifically, the cry34Ab1 gene probe hybridized to the expected internal T-DNA fragment of 1.915 kb, the cry35Ab1 gene probe hybridized to the expected internal T-DNA fragment of 2.607 kb, and the pat gene probe hybridized to a 3.4 kb border fragment consistent with a single intact T-DNA insertion as determined by Southern blot results.

Example 4. T-DNA Insert and Flanking Border Region Sequencing of *Bacillus thuringiensis* Cry34/35Ab1 Maize Line DAS-59122-7

The T-DNA insert and flanking border regions were cloned from B.t. Cry34/35Ab1 event DAS 59122-7 using

PCR based methods as diagramed in FIGS. 2 and 3. Specifically, sequences bordering the 5' and 3' ends of the insert in event DAS-59122-7 were obtained using two genome walking techniques. The first walking method was essentially the method as described for the Universal Genome Walker Kit (BD Biosciences Clontech, Palo Alto, Calif.), and the second method was conducted according to the splinkerette protocol outlined in Devon et al., (1995) *Nucleic Acids Research* 23 (9):1644-1645, with modifications as described by Stover (2001), U. C. Irvine (personal communication).

Briefly, genomic DNA was digested with various restriction enzymes (Dra I, EcoR V, Pvu II, Sma I and Stu I for the Universal Genome Walker method and BamH I, EcoR I, Hind III, and Xba I for the splinkerette method) and then ligated to blunt-end adaptors for the Genome Walker method and to adaptors specific for the restriction enzyme used for the splinkerette method. The adaptors for both genome walking methods were designed to prevent extension of the 3' end of the adaptor during PCR and thus reduce or eliminate nonspecific amplification. The adaptor-ligated genomic DNA fragments were then referred to as genome walker libraries or splinkerette libraries, one library for each restriction enzyme. Libraries were prepared from genomic DNA isolated from three individual T1S2 plants of B.t. Cry34/35Ab1 event DAS-59122-7; plants DAS-59122-7 T1S2 1, DAS-59122-7 T1S2 2 and DAS-59122-7 T1S2 10, and from one Hi-II and one PH09B control plant.

Following construction of the libraries, nested PCR amplifications were completed to amplify the target sequence using Advantage™-GC Genomic PCR kit (BD Biosciences Clontech, Palo Alto, Calif.). The primary PCR amplification used one primer with identity to the adaptor and one gene specific primer. The adaptor primer will not amplify a product in the first cycle of the primary PCR and only products from the gene specific primer will be produced. Annealing and amplification from the adaptor primer only occurs after the complementary strand has been produced from the gene specific primer. Following primary PCR amplification, a secondary nested PCR reaction was performed to increase the specificity of the genomic PCR reactions. The nested primers consisted of gene-specific and adaptor-specific sequences internal to the respective primers used in the primary PCR.

For 5' flanking border sequences, nested PCR was initiated using primers specific to the 5' end of the inserted T-DNA along with primers complementary to the adaptor

sequence ligated onto the digested DNA. Similarly, cloning of the 3' flanking border sequence started with a primer specific for the 3' end of the inserted T-DNA and a primer complementary to the adaptor sequence. DNA sequences internal to the T-DNA Right Border and Left Border sequences within the T-DNA region were used as the starting points for "walking out" to the maize genomic sequence, because they represented unique sequence (not homologous to endogenous maize genomic sequences) from which to anchor the genome walking primers.

The products produced by the nested PCR were analyzed by agarose gel electrophoresis (data not shown). Fragments visible in libraries prepared from one or more of the event DAS-59122-7 DNA samples and absent in libraries prepared from the Hi-II and PH09B genomic DNA samples were identified for further characterization. The identified PCR amplified fragments were separated by preparatory gel electrophoresis, isolated using a QIAquick Gel Extraction Kit (Qiagen), and sent directly for sequencing or cloned into a pGEM-T Easy plasmid vector using the pGEM-T Easy Vector System I (Promega Corp., Madison, Wis.) prior to DNA sequencing. Sequencing reactions were carried out with primers used for the nested PCR amplification or with primers specific for use with the pGEM-T Easy vector. The sequence obtained was used to design additional gene specific primers to continue "walking out" into the unknown maize genomic sequence. Multiple rounds of genome walking were performed until at least 500 bp of border sequence from the ends of the T-DNA insert were obtained.

To ensure validity of the flanking border sequences, additional event-specific PCR amplifications on genomic DNA from event DAS-59122-7 were performed. The amplified fragments were sequenced in order to further extend the region of sequence overlap from the T-DNA insert region into the 5' and 3' bordering genomic DNA. Primers, shown in Table 2, were designed based on the sequence obtained from the genome walking experiments to amplify a fragment spanning the unique junction of the T-DNA with the corn genomic DNA. Primer set 03-O-506/02-O-476 (SEQ ID NO: 10/SEQ ID NO: 9) spanned the 5' junction and amplified a 313 bp fragment (from bp 2427 to bp 2739, see FIG. 1), and primer set 02-O-447/03-O-577 (SEQ ID NO: 8/SEQ ID NO: 17) spanned the 3' junction and amplified a 754 bp fragment (from bp 9623 to bp 10376, see FIG. 1).

TABLE 2

Primer Sequences		Target Sequence Location (bp to bp) ¹
Primer Name	Sequence (5'-3')	
02-O-215 (SEQ ID NO: 1)	GTGGCTCCTTCAACGTTGCGGTTCTGTCTC	2743-2716
02-O-219 (SEQ ID NO: 2)	CGTGCAAGCGTCAATTCCGCCCTATAGTG	9830-9858
02-O-227 (SEQ ID NO: 3)	AATTGAGCGCTTGACGTTT	9846-9827
02-O-370 (SEQ ID NO: 4)	AACAACAAGACCGGCCACACCTC	4871-4894
02-O-371 (SEQ ID NO: 5)	GAGGTGGTCTGGATGGTGTAGGTCA	5187-5163
02-O-372 (SEQ ID NO: 6)	TACAACCTCAAGTGGTTCTCTCCCGA	7017-7044

TABLE 2-continued

Primer Sequences		Target Sequence Location (bp to bp) ¹
Primer Name	Sequence (5'-3')	
02-O-373 (SEQ ID NO: 7)	GAGGTCTGGATCTGCATGATCGCGGA	7897-7873
02-O-447 (SEQ ID NO: 8)	AACCCTTAGTATGTATTTGTATT	9623-9645
02-O-476 (SEQ ID NO: 9)	CTCCTTCAACGTTGCGGTTCTGTCTAG	2739-2714
03-O-506 (SEQ ID NO: 10)	TTTTGCAAAGCGAACGATTCAGATG	2427-2451
03-O-514 (SEQ ID NO: 11)	GCGGGACAAGCCGTTTTACGTTT	2687-2709
03-O-542 (SEQ ID NO: 12)	GACGGGTGATTTATTTGATCTGCAC	10766-10742
03-O-543 (SEQ ID NO: 13)	CATCTGAATCGTTGCTTTGCAAAA	2451-2427
03-O-564 (SEQ ID NO: 14)	CTACGTTCCAATGGAGCTCGACTGTC	2324-2299
03-O-569 (SEQ ID NO: 15)	GGTCAAGTGGACACTTGGTCACTCA	10150-10174
03-O-570 (SEQ ID NO: 16)	GAGTGAAGAGATAAGCAAGTCAAAG	10275-10299
03-O-577 (SEQ ID NO: 17)	CATGTATACGTAAGTTTGGTGTCTGG	10376-10352
03-O-784 (SEQ ID NO: 18)	AATCCACAAGATTGGAGCAAACGAC	2189-2213
67609 (SEQ ID NO: 36)	CGTATTACAATCGTACGCAATTTCAG	9862-9886
69240 (SEQ ID NO: 37)	GGATAAACAAACGGGACCATAGAAG	9941-9965

¹Location in sequence of Event DAS-59122-7 (see FIG. 1). Bases 1-2593 = 5' border, bases 2594-9936 = T-DNA insert, bases 9937-11922 = 3' border.

For verification of the DNA sequence that inserted into the maize genome, PCR was performed to amplify, clone, and sequence the inserted T-DNA from event DAS-59122-7. PCR primer sets, (SEQ ID NO: 11/SEQ ID NO: 5); (SEQ ID NO: 4/SEQ ID NO: 7); and (SEQ ID NO: 6/SEQ ID NO: 3) shown in Table 3 were used to amplify three overlapping fragments labeled 221-1 (SEQ ID NO: 25), 221-2 (SEQ ID NO: 26), and 221-3 (SEQ ID NO: 27) representing sequence from the 5' region of the T-DNA running through to the 3' region of the T-DNA insert from bp 2687 to bp 9846 for event DAS-59122-7 (see FIG. 1). PCR amplicon information is reported in Table 3 and primer sequences are listed in Table 2.

TABLE 3

PCR Primer and Amplicon Descriptions					
PCR Amplicon	Size (bp)	Target Sequence	Forward Primer	Reverse Primer	Location of PCR Amplicon (bp to bp) ¹
221-1 (SEQ ID NO: 25)	2501	T-DNA insert	03-O-514 (SEQ ID NO: 11)	02-O-371 (SEQ ID NO: 5)	2687-5187
221-2 (SEQ ID NO: 26)	3027	T-DNA insert	02-O-370 (SEQ ID NO: 4)	02-O-373 (SEQ ID NO: 7)	4871-7897
221-3 (SEQ ID NO: 27)	2830	T-DNA insert	02-O-372 (SEQ ID NO: 6)	02-O-227 (SEQ ID NO: 3)	7017-9846
O784/O564 (SEQ ID NO: 28)	136	5' genomic border	03-O-784 (SEQ ID NO: 18)	03-O-564 (SEQ ID NO: 14)	2189-2324
O784/O543 (SEQ ID NO: 29)	263	5' genomic border	03-O-784 (SEQ ID NO: 18)	03-O-543 (SEQ ID NO: 13)	2189-2451
O569/O577 (SEQ ID NO: 30)	227	3' genomic border	03-O-569 (SEQ ID NO: 15)	03-O-577 (SEQ ID NO: 17)	10150-10376
O570/O542 (SEQ ID NO: 31)	492	3' genomic border	03-O-570 (SEQ ID NO: 16)	03-O-542 (SEQ ID NO: 12)	10275-10766
O784/O215 (SEQ ID NO: 32)	555	5' junction	03-O-784 (SEQ ID NO: 18)	02-O-215 (SEQ ID NO: 1)	2189-2743
O219/O577 (SEQ ID NO: 33)	547	3' junction	02-O-219 (SEQ ID NO: 2)	03-O-577 (SEQ ID NO: 17)	9830-10376
O506/O476 (SEQ ID NO: 34)	313	5' junction	03-O-506 (SEQ ID NO: 10)	02-O-476 (SEQ ID NO: 9)	2427-2739
O447/O577 (SEQ ID NO: 35)	754	3' junction	02-O-447 (SEQ ID NO: 8)	03-O-577 (SEQ ID NO: 17)	9623-10376
67609/69240 (SEQ ID NO: 38)	104	3' junction	67609 (SEQ ID NO: 36)	69240 (SEQ ID NO: 37)	9862-9965

¹Location in sequence of Event DAS-59122-7 (see FIG. 1). Bases 1-2593 = 5' border, bases 2594-9936 = T-DNA insert, bases 9937-11922 = 3' border.

PCR GC2 Advantage™ Polymerase kit (BD Biosciences Clontech, Inc.) was used according to manufacturer's instructions to amplify the insert fragments (2211 (SEQ ID NO: 25), 2212 (SEQ ID NO: 26), and 2213 (SEQ ID NO: 27)). Briefly, a 50 µL reaction contained 5' and 3' primers at a final concentration of 0.2 µM and 40 ng of genomic DNA. PCR reactions were set up in duplicate using genomic DNA preparation from plants DAS-59122-7 T1S2 1 and DAS-59122-7 T1S2 2. PCR conditions were as follows: initial denaturation at 95° C. for 1 min, followed by 35 cycles of 94/95° C. for 30 sec, 55° C. for 30 sec, and 68° C. for 5 min, with final extension at 68° C. for 6 min. PCR amplification products were visualized under UV light, following electrophoresis through a 1% agarose gel in IX TBE (89 mM Tris-Borate, 2 mM EDTA, pH 8.3) stained with ethidium bromide.

PCR fragments 221-1 (SEQ ID NO: 25), 221-2 (SEQ ID NO: 26), and 221-3 (SEQ ID NO: 27) were purified by excising the fragments from 0.8% agarose gel in 1xTBE stained with ethidium bromide, and purifying the fragment from the agarose using a QIAquick Gel Extraction Kit (Qiagen). PCR fragments were cloned into a pGEM-T Easy plasmid vector using the pGEM-T Easy Vector System I (Promega Corp.). Cloned fragments were verified by mini-preparation of the plasmid DNA (QIAprep Spin Miniprep Kit, Qiagen) and restriction digestion with Not I. Plasmid clones and/or purified PCR insert fragments were then sent for sequencing of the complete insert. Sequencing reactions

were carried with primers designed to be specific for known T-DNA sequences or with primers specific for use with the pGEM-T Easy vector. Sigma-Genosys, Inc. (The Woodlands, Tex.) synthesized all PCR primers, which were used at a final concentration of 0.2-0.4 µM in the PCR reactions.

PCR reactions with genomic DNA isolated from B.t. Cry34/35Ab1 events DAS-59122-7, DAS-45214-4, and DAS-45216-6, and unmodified control lines Hi-II and PH09B were used to confirm (1) the presence of maize genomic DNA in the sequenced border regions of event DAS-59122-7, and (2) event specific amplification across the junctions of the T-DNA insert and genomic DNA borders in event DAS-59122-7.

PCR primers designed to amplify the border sequence flanking the insert in event DAS-59122-7 were used to confirm the presence of those regions in unmodified control lines as well as in event DAS-59122-7. Two (2) sets of primers each, for the 5' and 3' borders (four (4) sets total) were tested. Primer sets 03-O-784/03-O-564 (SEQ ID NO: 18/SEQ ID NO: 14) and 03-O-784/03-O-543 (SEQ ID NO: 18/SEQ ID NO: 13) were used to amplify 136 bp and 263 bp fragments, respectively, from border sequence 5' to the T-DNA insert in event DAS-59122-7 (FIGS. 2 and 3). Similarly, primer sets 03-O-569/03-O-577 (SEQ ID NO: 15/SEQ ID NO: 17) and 03-O-570/03-O-542 (SEQ ID NO: 16/SEQ ID NO: 12) were used to amplify 227 bp and 492 bp fragments, respectively, from the 3' genomic border (FIGS. 2 and 3).

Primers designed to amplify fragments across the junction of the border sequence and T-DNA insert were used to establish event-specific PCR fragments for event DAS-59122-7. One primer set was selected for each of the two junctions. Primer set 03-O-784/02-O-215 (SEQ ID NO: 18/SEQ ID NO: 1) was designed to amplify a 555 bp fragment across the 5' junction, and primer set 02-O-219/03-O-577 (SEQ ID NO: 2/SEQ ID NO: 17) was designed for amplification of a 547 bp fragment at the 3' junction. A set of primers, IVR1 (0197) (SEQ ID NO: 39) 5'-CCGCTG-TATCACAAAGGGCTGGTACC-3' and IVR2(O198) (SEQ ID NO: 40) 5'-GGAGCCCCGTGTAGAGCATGACGATC-3', based on the endogenous maize invertase gene (Hurst et al., (1999) *Molecular Breeding* 5 (6):579-586), was used to generate a 226 bp amplification product as an internal positive control for all maize genomic DNA samples.

All PCR primers were synthesized by Sigma-Genosys, Inc. and used at a final concentration of 0.2-0.4 µM in the PCR reactions. PCR primer sequences are listed in the Table 2. For PCR amplifications, Advantage™-GC 2 PCR kit (BD Biosciences) was used according to manufacturer's instructions. Approximately 10-100 ng of genomic DNA template was used per 50 µL PCR reaction. PCR conditions were as follows: initial template denaturation at 94° C. for 5 min, followed by 35 cycles of 95° C. for 1 minute, 60° C. for 2 minutes, and 72° C. for 3 min, with final extension at 72° C. for 7 min. The PCR amplification products were visualized under UV light following electrophoresis through a 1% agarose gel with IX TBE and ethidium bromide.

Sequence data obtained for the T-DNA insert and border regions of event DAS-59122-7 was reviewed and assembled using Seqman II™ software Version 4.0.5 (DNASTar, Inc., Madison, Wis.). The 5' and 3' border sequences flanking the insert present in event DAS-59122-7 were used for homology searching against the GenBank public databases in order to further characterize the site of insertion in the maize genome. Analysis to identify open reading frames in the junction regions between the flanking borders and T-DNA

insert in event DAS-59122-7 was conducted using Vector NTI 8.0 (InforMax™, Inc., Frederick, Md.).

In total, 11922 bp of sequence from genomic DNA of event DAS-59122-7 was confirmed (see FIG. 1). At the 5' end of the T-DNA insert, 2593 bp of flanking border sequence was identified, and 1986 bp of flanking border sequence was obtained on the 3' end from fragments derived from genome walking experiments. A total of 7160 bp of the T-DNA insert was cloned and sequenced using PCR primer sets designed to amplify three overlapping fragments labeled 221-1 (2501 bp) (SEQ ID NO: 25), 221-2 (3027 bp) (SEQ ID NO: 26), and 221-3 (2830 bp) (SEQ ID NO: 27) representing sequence from the 5' region of the T-DNA running through to the 3' region of the T-DNA insert for event DAS-59122-7 from bp 2687 to bp 9846 (see FIG. 1). The remainder of the T-DNA insert region was sequenced from two PCR fragments, O506/O476 (SEQ ID NO: 10/SEQ ID NO: 9) and O447/O577 (SEQ ID NO: 8/SEQ ID NO: 17) that spanned the 5' and 3' junctions, respectively, of the T-DNA insert with corn genomic DNA. Primers used were designed based on the sequence obtained from the genome walking experiments to amplify a fragment spanning the unique junction of the T-DNA with the corn genomic DNA. Primer set 03-O-506/03-O-476 (SEQ ID NO: 10/SEQ ID NO: 9) spanned the 5' junction and amplified a 313 bp fragment (from bp 2427 to bp 2739) and primer set 03-O-447/03-O-577 (SEQ ID NO: 8/SEQ ID NO: 17) spanned the 3' junction and amplified a 754 bp fragment (from bp 9623 to bp 10376). Combined, a total of 7343 bp of the T-DNA insert in event DAS-59122-7 was cloned and sequenced (from bp 2594 to bp 9936, see FIG. 1) and compared to the sequence of the transforming plasmid, PHP 17662. Two nucleotide differences at bp 6526 and bp 6562 were observed in the non-translated wheat peroxidase promoter region of the T-DNA insert (see FIG. 1). Neither of the observed base changes affected the open reading frame composition of the T-DNA insert. Both the 3' and 5' end regions of the T-DNA insert were found to be intact, except for deletion of the last 22 bp at the 5' end and 25 bp at the 3' end encompassing the Right and Left T-DNA Border regions, respectively. While T-DNA border sequences are known to play a critical role in T-DNA insertion into the genome, this result is not unexpected since insertions are often imperfect, particularly at the Left T-DNA Border (Tinland (1996) *Trends in Plant Science* 1(6): 178-184).

BLAST (Basic Local Alignment Search Tool) analysis of the genomic border regions of event DAS-59122-7 showed limited homology with publicly available sequences (Release 138.0 GenBank, Oct. 25, 2003). Analysis of the 5' border region found two areas with significant homology to maize genomic and EST (Expressed Sequence Tag) sequences. The first area encompasses 179 bp (bp 477 to bp 655 of the border sequence) and displays similarity to several molecular markers, chromosomal sequences, and consensus sequences obtained by alignment of various ESTs. The second area occurs at bp 1080 to bp 1153 (74 bp) of the 5' border sequence, and shows similarity to a number of different maize ESTs and genomic sequences. The 3' border region also had two small non-contiguous regions of similarity to plant DNA sequences. The inner 3' region of 162 bp (bp 9954 to bp 10115) showed similarity to the 3' untranslated end of two genomic *Zea mays* alcohol dehydrogenase (*adh1*) genes as well as to several EST consensus sequences. A smaller region (57 bp) in the middle of the 3' border (bp 10593 to bp 10649) showed similarity to non-coding regions from multiple maize genomic sequences.

Overall, no homologous regions greater than 179 base pairs were identified in either of the genomic border sequences, nor was more than one homologous region from the same known sequence found. Individual accessions displaying similarity to the event DAS-59122-7 border sequences were examined to determine if the insertion in event DAS-59122-7 occurred in a characterized protein coding sequence. None of the regions of similarity occurred within any known protein coding sequences. Local alignment of the entire transformation plasmid sequence, PHP17662, with the event DAS-59122-7 border sequences showed no significant homologies, indicating that the border regions flanking the T-DNA insert did not contain fragments of the transforming plasmid. Therefore, identification and characterization of the genomic sequence flanking the insertion site in event DAS-59122-7 was limited due to the absence of significant regions of homology to known sequences.

The 5' and 3' junction regions between the maize genomic border sequence and the T-DNA insert in event DAS-59122-7 were analyzed for the presence of novel open reading frames. No open reading frames of significant size (>100 amino acids) were identified in the 5' or 3' border junction regions, indicating that no novel open reading frames were generated as a result of the T-DNA insertion. Additionally, the homology searches did not indicate the presence of endogenous maize open reading frames in the border regions that might have been interrupted by the T-DNA insertion in B.t. Cry34/35Ab1 event DAS-59122-7.

Example 5. PCR Primers

PCR Primers DNA event specific primer pairs were used to produce an amplicon diagnostic for DAS-59122-7. These event primer pairs include, but are not limited to, SEQ ID NO: 18 and SEQ ID NO: 1; SEQ ID NO: 2 and SEQ ID NO: 17; SEQ ID NO: 10 and SEQ ID NO: 9; and SEQ ID NO: 8 and SEQ ID NO: 17; and SEQ ID NO: 36 and SEQ ID NO: 37. In addition to these primer pairs, any primer pair derived from SEQ ID NO: 21 and SEQ ID NO: 22 that when used in a DNA amplification reaction produces a DNA amplicon diagnostic for DAS-59122-7 is an embodiment of the present invention. Any modification of these methods that use DNA primers or complements thereof to produce an amplicon DNA molecule diagnostic for DAS-59122-7 is within the ordinary skill of the art. In addition, control primer pairs, which include IVR1(O197)/IVR2(O198) (SEQ ID NO: 39/SEQ ID NO: 40) for amplification of an endogenous corn gene are included as internal standards for the reaction conditions.

The analysis of plant tissue DNA extracts to test for the presence of the DAS-59122-7 event should include a positive tissue DNA extract control (a DNA sample known to contain the transgenic sequences). A successful amplification of the positive control demonstrates that the PCR was run under conditions that allow for the amplification of target sequences. A negative, or wild-type, DNA extract control in which the template DNA provided is either genomic DNA prepared from a non-transgenic plant, or is a non-DAS-59122-7 transgenic plant, should also be included. Additionally a negative control that contains no template corn DNA extract will be a useful gauge of the reagents and conditions used in the PCR protocol.

Additional DNA primer molecules of sufficient length can be selected from SEQ ID NO: 21 and SEQ ID NO: 22 by those skilled in the art of DNA amplification methods, and conditions optimized for the production of an amplicon diagnostic for event DAS-59122-7. The use of these DNA primer sequences with modifications to the methods shown

31

in these Examples are within the scope of the invention. The amplicon wherein at least one DNA primer molecule of sufficient length derived from SEQ ID NO: 21 and SEQ ID NO: 22 that is diagnostic for event DAS-59122-7 is an embodiment of the invention. The amplicon wherein at least one DNA primer of sufficient length derived from any of the genetic elements of PH117662A that is diagnostic for event DAS-59122-7 is an embodiment of the invention. The assay for the DAS-59122-7 amplicon can be performed by using a Stratagene Robocycler, MJ Engine, Perkin-Elmer 9700, or Eppendorf Mastercycler Gradient thermocycler, or by methods and apparatus known to those skilled in the art.

32

Having illustrated and described the principles of the present invention, it should be apparent to persons skilled in the art that the invention can be modified in arrangement and detail without departing from such principles. We claim all modifications that are within the spirit and scope of the appended claims.

All publications and published patent documents cited in this specification are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 40

<210> SEQ ID NO 1
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 1

gtggctcctt caacgttgcg gttctgtc 28

<210> SEQ ID NO 2
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 2

cgtgcaagcg ctcaattcgc cctatagtg 29

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 3

aattgagcgc ttgcacgttt 20

<210> SEQ ID NO 4
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 4

aacaacaaga cggccacac cctc 24

<210> SEQ ID NO 5
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 5

gagtggtctt ggatggtgta ggtca 25

-continued

<210> SEQ ID NO 6
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 6

tacaacctca agtggttcct cttcccga 28

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 7

gaggtctgga tctgcatgat gcgga 25

<210> SEQ ID NO 8
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 8

aacccttagt atgtatttgt att 23

<210> SEQ ID NO 9
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 9

ctccttcaac gttgcggttc tgtcag 26

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 10

ttttgcaaag cgaacgattc agatg 25

<210> SEQ ID NO 11
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

<400> SEQUENCE: 11

gcgggacaag ccgtttttagc ttt 23

<210> SEQ ID NO 12
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 12
gacgggtgat ttatttgatc tgcac 25

<210> SEQ ID NO 13
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 13
catctgaatc gttcgctttg caaaa 25

<210> SEQ ID NO 14
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 14
ctacgttcca atggagctcg actgtc 26

<210> SEQ ID NO 15
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 15
ggtcaagtgg acacttggtc actca 25

<210> SEQ ID NO 16
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 16
gagtgaagag ataagcaagt caaag 25

<210> SEQ ID NO 17
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 17
catgtatacg taagtttggg gctgg 25

<210> SEQ ID NO 18
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer
<400> SEQUENCE: 18
aatccacaag attggagcaa acgac 25

-continued

```

<210> SEQ ID NO 19
<211> LENGTH: 2593
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' flanking sequence of DAS 59122-7.

<400> SEQUENCE: 19
ctgagcgcac aacagcagagt cgcacggcac cggacgacat gagcagagatt tagatcggag    60
ggtgcggaca tggggcaacc tgcgcagcta acgcagggat ccacacgacc accaacgaa    120
ccaagcccgg gcaagtcgcc aggcaggttg ggccctggtt ccaccagcgg atgcatgcag    180
tgaagcgggg acggagagac aagccgaggg cgcgggtggg aatggcgctcc gggaggacga    240
gtggaggaga agaatctaga ggcacgaga ttcgagaagc cgacggagac aagattcgtg    300
tggggggaga caaacccggg ggctgagcgc cgttgatatg ggatcagacg gtgtggataa    360
aaaaagtgac gttgatagaa cgtctggcca gtgaaaaaac aaaacaactc caacaaaata    420
ctttaaaagc tcttataccc taaatgtagg ggatcaaaca cgtctctaca ctatttagca    480
gcgctcctcta aatgatcctc taaatttaga gaacgctact agattctcta tatatagttt    540
ctctaaacga tcttttatcc atttaaac tttaaataac cggtttaaca aaactaaaat    600
atatacaata catttgagag tatgacaaat acgtatgtat aaaaataaaa aataaaaata    660
tgtattagtc tactttgaat cttcttttct tcataatata atgatgtata gctctcatgt    720
gcggtgagaa aaaagttaga gctagacggt taatgtgtag tgacagtctt cgacgaaatc    780
tccctaataga gatgaattac tggaggttcc atcagaaaagt cccctgaaaa gaggcattta    840
tttagtttag tcagcaatth ctgggaacac aaatattctt ttgttatcac cactattaaa    900
aatctatggt tataacttat aataacatga aaaaataatt tagcatccca tatatataaa    960
aactgaagga agccatatat actaacataa gttaggagaa actaagaagg ttgtgcaaag    1020
cttgcaactgc tccaaaatac tgcaaaacaac cactctcctc taccaaccaa agaaactcat    1080
gtactccctc cgttcttttt tatttgtcgc attttagttt aaaaatgaac tagcagtcga    1140
caaatattcg agaacagata tagtatatac taacataact taggagatac taagaaagt    1200
gcgagagact ttcactgttc caaattactg caaagcctct cccctctgcc agtacatcta    1260
cgagatggtt cagttaaaca aagattcaga caagtgatga gccacttctt gtcatagatt    1320
gtgtgggcaa ccaaccatt gatgccacgg tttttgtgca tccatgcttt tgtattaaaa    1380
catcagttat gtttaccatg tccgatatgc tctacataat gacaatcaac ttgggtgtca    1440
ttatatttac aatgttagga atttcaatag ctacgaacac ttcaatagaa gtgcctttgt    1500
gggatcactc taatgtgttg ttgatgtaag gagaagaatc ttaatttact cttgctaaat    1560
ttgaactaca caaaaccact gcactgagga ttgtcctaataaattactgc tcatacacgt    1620
tagcatctgt tcagatactg agctaataccc taggattaaa ggatttgtaa aagatatgcc    1680
caatcattca ttttagttat ttattcttta gttatccact tgaagattta catacattg    1740
aaataaattt cttagaggta aagtgaaaat cagttattta aatacatttt agttatttat    1800
tttctctttt ttctaattt ttcttctgat ttgaagtctg aaaagataac tttgcctta    1860
tacatatttt atcttctacg tacgcatctg aacaacgtct ctttgcctcc tgatcgtgca    1920
gcaattagtg ctatgaatcg cgtttaagcg ctgcaaaatc atggctgggg cttcgtcctc    1980
gagtcgtcct gctgctcgat gtcacctcga gtcccgcacc gacctcagtg cttgttcttg    2040
ttggagccac ctctctcgga cgatcgccaa agacggataa ggccgaagcc gtcacttcag    2100

```

-continued

```

accgcgctca tgcgccgtag cagactccta catagcaggg ccagggatg tggacctttg 2160
caagtttagg attggaacca gcgaccagaa tccacaagat tggagcaaac gacccaaaat 2220
tcacaaggat tggcggtga cattgccagc ggggatcgc atgcggcggc ggcggccggg 2280
gcgagcacgg gagcaggoga cagtcgagct ccattggaac gtagaaatac ttaagggcaa 2340
ggtctccaaa tacttgaaaa aataggaaaa agaagaaaa acatgaaatg atattgaaat 2400
caattggaag atgttatgaa tctgttttt gcaaagcgaa cgattcagat ggcaaaaacta 2460
tgaatctttt tgtttgaagt cccaaatata aaattttctc gtactcacca acattggtgc 2520
gcacctgtga ttggctcata aaaattcttg gagggacgga agaagagtg aagggataag 2580
caagtaaaag cgc 2593

```

```

<210> SEQ ID NO 20
<211> LENGTH: 1986
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: 3' flanking sequence of DAS 59122-7.

```

```

<400> SEQUENCE: 20

```

```

ttcccttcta tggcccgtt tgtttatcct ctaaattata taatccagct taaataagtt 60
aagagacaaa caaacaacac agattattaa atagattatg taatctagat acctagatta 120
tgtaatccat aagtagaata tcaggtgctt atataatcta tgagctcgat tatataatct 180
taaaagaaaa caaacagagc cctataaaaa aggggtcaag tggacacttg gtcactcatt 240
taatccctcc ctctcctctt ttatccctct ttttgggtga ttcaccaata gtggtgtgca 300
cctgtgattg gctcgtaaaa attcctggac ggatggaaga gtgaagagat aagcaagtca 360
aagaaaagta acaacgaagc ttcacagct acaaatcttg gcccaactgg ttgcaccagc 420
accaaactta cgtatacatg attatctctg tttccctcat ttcgaagaaa aaaacgggtt 480
tcaaaacca ctgcttccag gagtaaaaa agataataat ctgaaacatt gcttccacct 540
tggcccttat ttggttaagt tgcaattcac cccaatccac atgtggattg agatggattg 600
cagtgtagct agacaaaacc ttaggccttg tttgcatagg aatacaccag gaattattcc 660
agctaatcaa aatttatata aatgagagaa acaattcga taggaattgt tccaggactt 720
cattctgcag taaccgaagc gcccttaat ccacccaat acacgtggat tggagtggat 780
tgaggtacag ccaacaagc cctaagtga gatcaataa atcacccgct atattcttct 840
acctacaaaa acagcaataa acacctgaat gaagtctaa tttgcacagt gtaggtagga 900
tgaaaatagt tacctctca tggtcagtaa ctcttggcac acaacttcac atgtaatcga 960
tgtaccactt ggctcttgcc tgaacccaa tacatctta gcataagaat aatattatga 1020
tggcaaggca tgatcaccag cactccttta ttgttagta agtctatcac tccccaaaac 1080
aattcaaag aacagagatg cattgcccc aatgaattct atttcaatta gccggaaaat 1140
tctacttcat cagaagcacc caaattgcca gcacccctac tagactgacc atgaccaggc 1200
tgccgcagat gcctcttttt ctgtcctctc ctctttgctc tgagtttctc tcaagatcc 1260
ctacccccac gtctcttata catcttaaag ctaacatgct tctcctcgc catcttctca 1320
accttctcag taatctcagc agcaatctga cggttgtaca acttcttcag ccccttcac 1380
aactttgcaa atgtgtcagg ctgtggcacc agtctgctc cttagcatgct taagcaatac 1440
aggcaggcct ccttgacatg tttctctgca aacagtgcac gaatccagat agtccatgca 1500
ctcacattga gctcacagcc tttgctcaca atacatttcc aaacatcctt tgcaagctca 1560

```

-continued

```

agtttctcat ctctgaccaa cgcattgagg aggtccttca gcaccccata ttgcggtacc 1620
acaaagagcc ccctcccaac catgtcttta aaataactac atgcctcaat cagcaaaccc 1680
tgcccaacaa ggccactcac cacgatagca aatgtatcga ccacaggact gagcccagca 1740
ctttccatct cattccacaa tgtcatggct tgcttggctt cccaagcct gcaggccaac 1800
cgaatcacca cattgtatat cttgagatct ggtggacacc ggcaactccc catcctctcc 1860
atcagctcca agcactctc aagctgctcc ttcttctcgt gtgctacaaa gaaacctatg 1920
tacacggcag cgteccaccg caggccatcc ctcgacatag catccaagaa ctctgacccc 1980
tgggat 1986

```

<210> SEQ ID NO 21

<211> LENGTH: 3594

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence that represents part of the PHI17662A insert as well as flanking sequence 5' to the insert.

<400> SEQUENCE: 21

```

ctgagcgcac aacagcaggt cgcattggcag cggacgacat gagcagagatt tagatcggag 60
ggtgcggaca tggggcaacc tgcgcagcta acgcagggat ccacacgacc accaacgaag 120
ccaagcccgg gcacgtcccc aggcagggtg gcccctggtt ccaccagcgg atgcatgcag 180
tgaagcgggg acggagagac aagccgaggg cgcgggtggg aatggcgtcc gggaggacga 240
gtggaggaga agaattctaga ggcatcgaga ttcgagaagc cgacggagac aagattcgtg 300
tggggggaga caaacccggg ggctgagcgc cgttgatatg ggatcagacg gtgtggataa 360
aaaaagtgc gttgatagaa cgtctggcca gtgaaaaaac aaaacaactc caacaaaata 420
ctttaaaagc tcttataccc taaatgtagg ggatcaaaac cgtctctaca ctatttagca 480
gcgtcctcta aatgatctc taaatttaga gaacgctact agatttctcta tatatagttt 540
ctctaaacga tcttttatcc atttaatac ttttaataac cggtttaaca aaactaaat 600
atatacaata catttgagag tatgacaaat acgtatgtat aaaaataaaa aataaaataa 660
tgtattagtc tactttgaat cttcttttct tcataatata atgatgtata gctctcatgt 720
gcggtgagaa aaaagttaga gctagacggt taatgtgtag tgacagctct cgacgaaatc 780
tccctaata gatgaattac tggaggttcc atcagaaagt cccctgaaaa gaggcattta 840
tttagtttag tcagcaatct ctgggaacac aaatattcct ttgttatcac cactattaaa 900
aatctatggt tataacttat aataacatga aaaaataaatt tagcatccca tatatataaa 960
aactgaagga agccatata actaacataa gttaggagaa actaagaagg ttgtgcaaag 1020
cttgcaactg tccaaaatac tgcaaacac cactctcctc taccaaccaa agaaactcat 1080
gtactccctc cgttcttttt tatttgtgcg attttagttt aaaaatgaac tagcagtcga 1140
caaatattcg agaacagata tagtatatac taacataact taggagatac taagaaagtt 1200
gcgcagagct ttcactgttc caaattactg caaagcctct cccctctgcc agtacatcta 1260
cgagatgttt cagttaaaca aagattcaga caagtgatga gccacttctt gtcatagatt 1320
gtgtgggtcaa ccaaccatt gatgccacgg tttttgtgca tccatgcttt tgtattaaaa 1380
catcagttat gtttaccatg tccgatatgc tctacataat gacaatcaac ttgggtgtca 1440
ttatatttac aatgttagga atttcaatag ctacgaacac ttcaatagaa gtgcctttgt 1500
gggatcacct taatgtgttg ttgatgtaag gagaagaatc ttaatttact cttgctaaat 1560

```

-continued

```

ttgaactaca caaaaccact gcactgagga ttgtcctaataaattactgc tcatacacgt 1620
tagcatctgt tcagatactg agctaataccc taggattaaa ggatttgtaa aagatatgcc 1680
caatcattca ttttagttat ttattttotta gttatocact tgaagattta catacatttg 1740
aaataaattt ctttagagga aagtgaaaat cagttattta aatacatttt agttatttat 1800
tttcttcttt ttctaattt ttcttctgat ttgaagtctg aaaagataac ttgcccctta 1860
tacatatttt atcttctacg tacgcactcg aacaacgtct ctttgtcccc tgatcgtgca 1920
gcaattagtg ctatgaatcg cgtttaagcg ctgcaaaaatc atggctgggg cttcgtctctc 1980
gagtcgtcct gctgctcgat gtcacctcga gtcccgcacc gacctcagtg cttgttcttg 2040
ttggagccac ctctctcgga cgatcgccaa agacggataa ggccgaagcc gtcacttcag 2100
accgcgtca tgcgccgtag cagactccta catagcaggg ccagggtatg tggaccttg 2160
caagtttagg attggaacca gcgaccagaa tccacaagat tggagcaaac gaccaaaaat 2220
tcacaaggat tggcgggtga cattgcccgc gcgggatcgc atgcggcggc ggcggccggg 2280
gcgagcacgg gagcaggcga cagtcgagct ccattggaac gtagaaatac ttaagggcaa 2340
ggtctccaaa tacttgaaaa aatagaaaa agaagaaaat acatgaaatg atattgaaat 2400
caattggaag atgttatgaa tcttgttttt gcaaagcgaa cgattcagat ggcaaaaacta 2460
tgaatctttt tgtttgaagt cccaaatata aaattttctc gtactcacca acattggtgc 2520
gcacctgtga ttggctcata aaaattcttg gagggacgga agaaagagtg aagggataag 2580
caagtaaaag cgctcaaaaca ctgatagttt aaactgaagg cgggaaacga caatctgatc 2640
atgagcggag aattaagggg gtcacgttat gacccccgcc gatgacgcgg gacaagccgt 2700
tttacgtttg gaactgacag aaccgcaacg ttgaaggagc cactcagcaa gcttactagt 2760
agcgtgttt aaacgctctt caactggaag agcggttacc cggaccgaag cttgcatgcc 2820
tgcaatgcag cgtgaccogg tctgcccct ctctagagat aatgagcatt gcatgtctaa 2880
gttataaaaa attaccacat attttttttg tcacacttgt ttgaagtgca gtttatctat 2940
ctttatacat atatttaaac tttactctac gaataatata atctatagta ctacaataat 3000
atcagtgttt tagagaatca tataaatgaa cagttagaca tggctaaag gacaattgag 3060
tattttgaca acaggactct acagttttat ctttttagtg tgcagtgtt ctctttttt 3120
tttgcaata gcttcaccta tataatactt catccatttt attagtacat ccatttaggg 3180
tttaggggta atggttttta tagactaatt ttttttagtac atctatttta ttctatttta 3240
gcctctaaat taagaaaact aaaactctat tttagttttt ttatttaata atttagatat 3300
aaaatagaat aaaataaagt gactaaaaat taacaaaata ccctttaaga aattaaaaaa 3360
actaagggaaa catttttctt gtttcgagta gataatgcca gcctgttaaa cgccgtcgac 3420
gagtctaacg gacaccaacc agcgaaccag cagcgtcgcg tcgggccaag cgaagcagac 3480
ggcacggcat ctctgtgctt gcctctggac cctctctgag agttccgctc caccgttgga 3540
cttgcctcgc tgteggcatc cagaaattgc gtggcggagc ggcagacgtg agcc 3594

```

<210> SEQ ID NO 22

<211> LENGTH: 2987

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Sequence that represents part of the PHI17662A insert as well as flanking sequence 3' to the insert.

<400> SEQUENCE: 22

-continued

ctccggagag gagaccagtt gagattaggc cagctacagc agctgatatg gccgcggttt	60
gtgatatcgt taaccattac attgagacgt ctacagtga ctttaggaca gagccacaaa	120
caccacaaga gtggattgat gatctagaga ggttgcaaga tagataccct tggttggttg	180
ctgagggtga ggggtgtgtg gctggatttg cttacgctgg gccctggaag gctaggaacg	240
cttacgattg gacagttgag agtactgttt acgtgtcaca taggcatcaa aggttgggcc	300
taggatccac attgtacaca catttgctta agtctatgga ggcgcaaggt ttttaagtctg	360
tggttgctgt tataggcctt ccaaacgac catctgttag gttgcatgag gctttgggat	420
acacagcccg ggggtacattg cgcgcagctg gatacaagca tgggtgatgg catgatggtg	480
gtttttggca aagggatttt gagttgccag ctccctccaag gccagttagg ccagttaccc	540
agatctgagt cgacctgcag gcatgcccgc tgaaatcacc agtctctctc tacaaatcta	600
tctctctcta taataatgtg tgagtagtcc ccagataagg gaattagggt tcttataggg	660
tttcgctcat gtgttgagca tataagaaac ccttagtatg tatttgtatt tgtaaaatac	720
ttctatcaat aaaatttcta attcctaaaa ccaaaatcca gggcgagctc ggtaccggg	780
gatcctctag agtcgacctg caggcatgcc cgcggatacc gatgggcccc ggccgaagct	840
tcggtcctgg ccctcgtggc ctcttgctct tcaggatgaa gagctatggt taaacgtgca	900
agcgtcaat tcgccctata gtgagtcgta ttacaatcgt acgcaattca gtacattaaa	960
aacgtccgca atgtgttatt aagttgtcta agcgtcaatt tttccctctc atggtcccgt	1020
ttgtttatcc tctaaattat ataatccagc ttaaataagt taagagacaa acaacaaca	1080
cagattatta aatagattat gtaactctaga tacctagatt atgtaatcca taagtagaat	1140
atcaggtgct tatataatct atgagctcga ttatataatc ttaaaagaaa acaaacagag	1200
cccctataaa aaggggtcaa gtggacactt ggtcactcat ttaatccctc cctctcctct	1260
tttatccctc tttttggtgt attcaccaat agtgggtgtc acctgtgatt ggctcgtaaa	1320
aattcttggc cggatggaag agtgaagaga taagcaagtc aaagaaaagt aacaacgaag	1380
cttcatcagc tacaaatfff ggcccaactg gttgcaccag caccaaaactt acgtatacat	1440
gattatctct gtttccctca tttcgaagaa aaaaacgggt ttcaaaaccc actgctttca	1500
ggagtaaaaa aagataataa tctgaaacat tgcttccacc ttggccctta tttggttacg	1560
tgcaaatca cccaatcca catgtggatt gagatggatt gcagtgtagc tagacaaacc	1620
cttaggcctt gtttgcatag gaatacacca ggaattatcc cagctaatca aaatttatat	1680
aatgagaga aacaattcgg ataggaattg ttccaggact tcattctgca gtaaccgaac	1740
ggccccctaa tccaccccaa tacacgtgga ttggagtgga ttgaggtaca gccaaacaag	1800
gcctaagtgc agatcaaata aatcacccgt catattcttc tacctacaaa aacagcaata	1860
aacacctgaa tgaagttcta atttgcacag tgtaggtagg atgaaaatag ttacctctc	1920
atggtcagta actcttggca cacaacttca catgtaatcg atgtaccact tggctcttgc	1980
ctgaaaccca atacatcttt agcataagaa taatattatg atggcaaggc atgatcacca	2040
gcactccttt attgtttagt aagtctatca ctccccaaaa caattcaaat gaacagagat	2100
gcattgcccc caatgaattc tatttcaatt agccggaaaa ttctacttca tcagaagcat	2160
ccaaattgcc agcatcccta ctagactgac catgaccagg ctgcccgaga tgctctttt	2220
tctgtcctct cctctttgcc ttgagtttct cttcaagatc cctcaccoca cgtctcttat	2280
acatcttaaa gctaacaatgt ctctcctcgg ccattctcct aaccttctca gtaatctcag	2340

-continued

```

cagcaatctg acggttgtag aacttcttca gcccttcat caactttgca aatgtgtcag 2400
getgtggcat cagtctgcc tctagcatgt ctaagcaata caggcaggcc tccttgacat 2460
gtttcttcgc aaacagtga tgaatccaga tagtccatgc actcacattg agctcacagc 2520
ctttgctcac aatacatttc caaacatcct ttgcaagctc aagtttctca tctctgacca 2580
acgcattgag gaggtccttc agcaccocat attgcggtac cacaaagagc cccctcccaa 2640
ccatgtcttt aaaataacta catgcctcaa tcagcaaac ctgcccaaca aggccactca 2700
ccacgatagc aaatgatcg accacaggac tgagcccagc actttccatc tcattccaca 2760
atgtcatggc ttgcttggtc tcccgaagcc tgcaggccaa ccgaatcacc acattgtata 2820
tcttgagatc tgggtgacac cggcactccc gcactctctc catcagctcc aagcactcct 2880
caagctgctc cttcttctcg tgtgctacaa agaaacctg gtacacggca gcgtccacct 2940
gcaggccatc cctcgacata gcatccaaga actcgtacct ctgggat 2987

```

<210> SEQ ID NO 23

<211> LENGTH: 11922

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: The sequence represents the complete sequence of the insert and flanking regions of event DAS 59122-7.

<400> SEQUENCE: 23

```

ctgagcgcac aacagcgagt cgcattggc cggacgacat gagcagatt tagatcggag 60
ggtgctggaca tggggcaacc tgcgcagcta acgcagggat ccacacgacc accaacgaag 120
ccaagcccgg gcactgccc aggcagggtt ggccctggtt caccacgagg atgcatgcag 180
tgaagcgggg acggagagac aagccagggg cgcgggtggg aatggcgtcc gggaggacga 240
gtggaggaga agaattctaga ggcattcaga ttcgagaagc cgacggagac aagattcgtg 300
tggggggaga caaacccggg ggtgagcgc cgttgatatg ggatcagacg gtgtggataa 360
aaaaagtgac gttgatagaa cgtctggcca gtgaaaaaac aaaacaactc caacaaaata 420
ctttaaaagc tcttataccc taaatgtagg ggatcaaaa cgtctctaca ctatttagca 480
gcgtcctcta aatgatcctc taaatttaga gaacgctact agattctcta tatatagttt 540
ctctaaacga tcttttatcc atttaaac tttaaataac cggtttaaca aaactaaaat 600
atatacaata catttgagag tatgacaaat acgtatgtat aaaaataaaa aataaaaata 660
tgtattagtc tactttgaat cttcttttct tcataatata atgatgtata gctctcatgt 720
gcggttgaaa aaaagttaga gctagacgtt taatgtgtag tgacagtctt cgacgaaatc 780
tcccataatga gatgaattac tggaggttcc atcagaaaag cccctgaaaa gaggcattta 840
tttagtttag tcagcaatth ctgggaacac aaatattctt ttgttatcac cactattaaa 900
aatctatggt tataacttat aataacatga aaaaataatt tagcatccca tatatataaa 960
aactgaagga agccatata actaacataa gttaggagaa actaagaagg ttgtgcaaa 1020
cttgactgac tccaaaatac tgcaaacac cactctctc taccaaccaa agaaactcat 1080
gtactcctc cgttcttttt tatttgctgc attttagttt aaaaatgaac tagcagtcga 1140
caaatattcg agaacagata tagtatatac taacataact taggagatac taagaaagt 1200
gcgagagct ttcactgttc caaattactg caaagcctct cccctctgcc agtacatcta 1260
cgagatgttt cagttaaaca aagattcaga caagtgatga gccacttctt gcatagatt 1320
gtgtggtaaa ccaaccatt gatgccagg tttttgtgca tccatgcttt tgtattaaaa 1380

```

-continued

catcagttat gtttaccatg tccgatatgc tctacataat gacaatcaac ttggtgttca	1440
ttatatttac aatgttagga atttcaatag ctacgaacac ttcaatagaa gtgcctttgt	1500
gggatcacct taatgtgttg ttgatgtaag gagaagaatc ttaatttact cttgctaaat	1560
ttgaactaca caaaaccact gcaactgagga ttgtcctaataa aaattactgc tcatacacgt	1620
tagcatctgt tcagatactg agctaataccc taggattaaa ggatttgtaa aagatatgcc	1680
caatcattca ttttagttat ttattttotta gttatccact tgaagattta catacatttg	1740
aaataaattt cttagaggta aagtgaaaat cagttattta aatacatttt agttatttat	1800
tttcttcttt ttcctaattt ttccttgat ttgaagtctg aaaagataac tttgccctta	1860
tacatatttt atcttctacg tacgcactg aacaacgtct ctttgtcccc tgatcgtgca	1920
gcaattagtg ctatgaatcg cgtttaagcg ctgcaaaatc atggctgggg cttcgtcctc	1980
gagtcgtcct gctcctcgat gtcacctcga gtccccgacc gacctcagtg cttgttcttg	2040
ttggagccac ctctctcgga cgatcgccaa agacggataa ggccgaagcc gtcacttcag	2100
accgcgctca tgcgccgtag cagactccta catagcaggg ccagggtatg tggaccttg	2160
caagtttagg attggaacca gcgaccagaa tccacaagat tggagcaaac gaccaaaaat	2220
tcacaaggat tggcggtga cattgccagc ggggatcgc atgcccggc ggcggccggg	2280
gcgagcacgg gagcaggoga cagtcgagct ccattggaac gtagaaatac ttaagggcaa	2340
ggtctccaaa tacttgaaaa aataggaaaa agaagaaaa acatgaaatg atattgaaat	2400
caattggaag atgttatgaa tctgttttt gcaaagcga cgattcagat ggcaaaacta	2460
tgaatctttt tgtttgaagt cccaaatata aaattttctc gtactacca acattggtgc	2520
gcacctgtga ttggctcata aaaattcttg gagggacgga agaaagagtg aagggataag	2580
caagtaaaag cgctcaaaaca ctgatagttt aaactgaagg cgggaaacga caatctgatc	2640
atgagcggag aattaaggga gtcacgttat gacccccgcc gatgacggg gacaagccgt	2700
tttacgtttg gaactgacag aaccgcaacg ttgaaggagc cactcagcaa gcttactagt	2760
agcgtgtttt aaacgctctt caactggaag agcggttacc cggaccgaag cttgcatgcc	2820
tgcagtgcag cgtgaccogg tctgtcccct ctctagagat aatgagcatt gcatgtctaa	2880
gttataaaaa attaccacat attttttttg tcacacttgt ttgaagtgca gtttatctat	2940
ctttatacat atattttaaac tttactctac gaataatata atctatagta ctacaataat	3000
atcagtgttt tagagaatca tataaatgaa cagttagaca tggctaaaag gacaattgag	3060
tattttgaca acaggactct acagttttat ctttttagtg tgcattgttt ctcctttttt	3120
tttgcaata gcttcaccta tataaactt catccatttt attagtacat ccatttaggg	3180
tttagggta atggttttta tagactaatt tttttagtac atctatttta ttctatttta	3240
gcctctaaat taagaaaaact aaaactctat tttagttttt ttatttataa atttagatat	3300
aaaatagaat aaaataaagt gactaaaaat taaacaaata ccctttaaga aattaaaaaa	3360
actaaggaat catttttctt gtttcgagta gataatgcca gcctgttaa cgccgtcgac	3420
gagctaacg gacaccaacc agcgaaccag cagcgtcgcg tcgggccaag cgaagcagac	3480
ggcacggcat ctctgtcgct gcctctggac ccctctcgag agttccgctc cacogttgga	3540
cttgcctcgc tgteggcatc cagaaattgc gtggcggagc ggcagacgtg agccggcacg	3600
gcaggcggcc tcctcctcct ctcacggcac cggcagctac gggggatttc tttcccaccg	3660
ctcctcgcct ttcctcctcct cgcccccgct aataaataga caccctctcc acaccctctt	3720
tcccaacct cgtgtgttc ggagcgcaca cacacacaac cagatctccc ccaaatccac	3780

-continued

cgcgtagc	ctccgcttca	aggtaacg	ctcgtctcc	cccccccc	ctctctacct	3840
tctctagatc	ggcgttccg	tccatggta	gggccggta	gttctacttc	tggtcatgtt	3900
tgtgtagat	ccgtgttgt	gtagatccg	tgtgctagc	gttcgtacac	ggatgcgacc	3960
tgtacgtcag	acacgttctg	attgctaac	tgccagtgtt	tctctttggg	gaatcctggg	4020
atggctctag	ccgttccgca	gacgggatcg	atctcatgat	ttttttgtt	tcgttgcata	4080
gggtttgggtt	tgccttttc	ctttatttca	atatatgccc	tgcacttgtt	tgtcgggtca	4140
tcttttcagt	cttttttttg	tcttggtgt	gatgatgtg	tctggttggg	cggctgttct	4200
agatcggagt	agaattctgt	ttcaaac	ctggtggatt	tattaat	ggatctgat	4260
gtgtgtgcca	tacatattca	tagttacgaa	ttgaagatga	tggatggaaa	tatcgatcta	4320
ggataggtat	acatgttgat	gcgggtttta	ctgatgcata	tacagagatg	ctttttgttc	4380
gcttggtgtg	gatgatgtg	tgtggtggg	cggctgttca	ttcgttctag	atcggagtag	4440
aatactgttt	caaac	ggtgtattta	ttaat	aactgatgt	gtgtgcata	4500
catcttcata	gttacgagtt	taagatggat	ggaaatctg	atgtaggata	ggtatacatg	4560
ttgatgtggg	tttactgat	gcatacat	gatggcatat	gcagcatcta	ttcatatgct	4620
ctaaccttga	gtacctatct	attataataa	acaagtatgt	tttataatta	ttttgatctt	4680
gatatacttg	gatgatggca	tatgcagcag	ctatatgtgg	atttttttag	cctgcttc	4740
atacgtatt	tatttcttg	gtactgttc	ttttgtcgat	gctcaccctg	ttgtttggtg	4800
ttacttctgc	aggctgactc	tagaggatcc	acacgacacc	atgtccgcc	gcgaggtgca	4860
catcgacgtg	aacaacaaga	cggccacac	cctccagctg	gaggacaaga	ccaagctcga	4920
cggcggcagg	tggcgacct	ccccgacca	cgtggccaac	gaccagatca	agacctcgt	4980
ggccgaatcc	aacggttca	tgaccggcac	cgagggcacc	atctactact	caattaatgg	5040
cgagggcag	atcagcctct	acttcgacaa	cccgttcgcc	ggctccaaca	aatacagcgg	5100
ccactccaac	aagctccagt	acgagatcat	caccagggc	ggctccggca	accagtccca	5160
cgtgacctac	accatccaga	ccacctctc	ccgtacggc	cacaagtcc	gagtcagag	5220
tcatgagtca	gttaacctag	acttgcocat	cttctggatt	ggccaactta	attaatgtat	5280
gaaataaaag	gatgcacaca	tagtgacatg	ctaataacta	taatgtggc	atcaaagtgt	5340
tgtgttatgt	gtaactacta	gttatctgaa	taaaagagaa	agagatc	catatctct	5400
atcctaaaatg	aatgtcagct	gtctttataa	ttctttgatg	aaccagatgc	atttcattaa	5460
ccaaatccat	atacatataa	atattaatca	tatataatta	atatcaattg	ggtagcaaa	5520
acaaatctag	tctaggtgtg	ttttgcgaat	gcggccggg	accgaattgg	ggatctgcat	5580
gaaagaaact	gtcgcactgc	tgaaccgcac	cttgtcactt	tcatcgaaca	cgacctgtgc	5640
ccaagatgac	ggtgctcgg	tctaagtgag	gctgaattgc	cttggacaga	agcggactcc	5700
ctacaattag	ttaggccaaa	cgtgcatcc	atgtgtagct	ccgggctcgg	gctgtagcgc	5760
catctgcaat	agcatccatg	gagctcgttc	catgtagttg	gagatgaacc	aatgatcggg	5820
cgtgtggaog	tatgttctctg	tgtactccga	tagtagagta	cgtgttagct	ctttcatggt	5880
gcaagtga	tttgtgttg	tttaattacc	cctacgttag	ttgcgggaca	ggagacacat	5940
catgaattta	aaggcgatga	tgcctctcc	tgtaatgtta	ttctttgat	gtgatgaatc	6000
aaaatgtcat	ataaacatt	tgttgctctt	tagttaggcc	tgatcgtaga	acgaaatgct	6060
cgtgtagcgg	ggctacgagc	ctatgacgca	ataaacactgg	tttgcggcc	cggagtcgct	6120

-continued

tgacaaaaa aagcatgtta agtttattta caattcaaaa cctaacatat tatattccct	6180
caaagcaggt tcacgatcac acctgtacct aaaaaaaca tgaagaatat attactccat	6240
tattatgaga tgaaccactt ggcaagagtg gtaagctata taaaaaatg aacattatta	6300
cgagatgtta tatgccatta tattgattcg aagatatatg tttctttctc ccacgggcac	6360
ctaacggata catgataagg ccaaggcaga tcacgggaaa ttattcgaat acatgttacg	6420
ccctattgcc ggaaaaaaa tgcagggcag gtgttgcccg tagcgttta agcacttaag	6480
ctggaggttg ccacacttgg atgcaagcgt ctgacccttc taaaacatcg gcggctttgt	6540
ccgtatccgt atcccctatc cgacatctag ctggccacac gacgggctg gccagatcgt	6600
ggatgccggg tcgacgtoga tcgtcagcca tcatagacca atcgaccatc tgttatggat	6660
gcttgctagc tagactagtc agacataaaa tttggatact ttctcccaac tgggagacgg	6720
ggactgatgt gcagctgcac gtgagctaaa tttttcccta taaatatgca tgaatactg	6780
cattatcttg ccacagccac tgcacagcc agataacaag tgcagctggt agcagcaac	6840
gcatagctct ggactttag ctaggtagcc aaccggatcc acacgacacc atgctcgaca	6900
ccaacaaggt gtacgagatc agcaaccacg ccaacggcct ctacgccgc acctaectct	6960
ccctcgacga ctccggcgtg tccctcatga acaagaacga cgacgacatc gacgactaca	7020
acctcaagtg gttcctcttc ccgatcgacg acgaccagta catcatcacc tctacgccg	7080
ccaacaactg caaggtgtgg aacgtgaaca acgacaagat taatgtgtca acctactcct	7140
ccaccaactc catccagaag tggcagatca aggccaacgg ctctcctac gtgatccagt	7200
ccgacaacgg caaggtgctc accgccggca ccggccaggc cctcggcctc atccgcctca	7260
ccgacgagtc ctccaacaac ccgaaccagc aatggaacct gacgtccgtg cagaccatcc	7320
agctcccgca gaagccgac atcgacacca agctcaagga ctaccogaag tactccccga	7380
ccggcaacat cgacaacggc acctccccgc agctcatggg ctggaccctc gtgccgtgca	7440
tcatggtgaa cgacccgaac atcgacaaga acaccagat caagaccacc ccgtaactaca	7500
tcctcaagaa gtaccagtac tggcagaggg ccgtgggctc caacgtcggc ctccgcccgc	7560
acgagaagaa gtcctacacc tacgagtggt gcaccgagat cgaccagaag accaccatca	7620
tcaacacctc cggcttcag atcaacatcg acagcggcat gaagttcgac atcccgagg	7680
tgggcccggg tacccgacgag atcaagacc agctcaacga ggagctcaag atcgagtatt	7740
cacatgagac gaagatcatg gagaagtacc aggagcagtc cgagatcgac aaccgaccg	7800
accagtccat gaactccatc ggcttctca ccatcacctc cctggagctc taccgctaca	7860
acggctccga gatccgcatc atcgagatcc agacctccga caacgacacc tacaactgta	7920
cctcctacc gaaccaccag caggccctgc tgctgctgac caaccactcc tacgaggagg	7980
tggaggagat caccaacatc ccgaagtcca cctcaagaa gctcaagaag tactacttct	8040
gagtcagtag tcatgagtca gttaacctag acttgtccat cttctggatt ggccaactta	8100
attaatgtat gaaataaaag gatgcacaca tagtgacatg ctaatcacta taatgtgggc	8160
atcaaagttg tgtgttatgt gtaattacta gttatctgaa taaaagagaa agagatcatc	8220
catatttctt atcctaagt aatgtcacgt gtctttataa ttctttgatg aaccagatgc	8280
atttcattaa ccaaatccat atacatataa atattaatca tatataatta atatcaattg	8340
ggttagcaaa acaaatctag tctaggtgtg ttttgcgaa tcccatggag tcaaagatc	8400
aaatagagga cctaacagaa ctccgccgta agactggcga acagttcata cagagtctct	8460
tacgactcaa tgacaagaag aaaatcttcg tcaacatggt ggagcacgac acgcttctct	8520

-continued

actccaaaa	tatcaaagat	acagtctcag	aagaccaaag	ggcaattgag	acttttcaac	8580
aaagggtaat	atccggaaac	ctcctcggat	tccattgccc	agctatctgt	cactttattg	8640
tgaagatagt	ggaaaaggaa	ggtggctcct	acaaatgcca	tcattgcat	aaaggaaag	8700
ccatcggtga	agatgectct	gccgacagtg	gtcccaaaga	tggaccccca	cccacgagga	8760
gcatcgtgga	aaaagaagac	gttccaacca	cgtcttcaaa	gcaagtggat	tgatgtgata	8820
tctccactga	cgtaagggat	gacgcacaat	cccactatcc	ttcgcaagac	ccttcctcta	8880
tataaggaag	ttcatttcat	ttggagagga	cagggtaccc	gggatccac	catgtctccg	8940
gagaggagac	cagttgagat	taggccagct	acagcagctg	atatggccgc	ggtttgtgat	9000
atcgtaacc	attacattga	gacgtctaca	gtgaacttta	ggacagagcc	acaaacacca	9060
caagagtgga	ttgatgatct	agagaggttg	caagatagat	acccttggtt	ggttgctgag	9120
gttgagggtg	ttgtggctgg	tattgcttac	gctgggccct	ggaaggctag	gaacgcttac	9180
gattggacag	ttgagagtac	tgtttacgtg	tcacatagcc	atcaaagggt	gggcctagga	9240
tccacattgt	acacacattt	gcttaagtct	atggaggcgc	aaggttttaa	gtctgtggtt	9300
gctgttatag	gccttccaaa	cgatccatct	gttaggttgc	atgaggcttt	gggatacaca	9360
gccccgggta	cattgcgcgc	agctggatac	aagcatggtg	gatggcatga	tgttggtttt	9420
tggcaaaagg	atthttgagt	gccagctcct	ccaaggccag	ttaggccagt	taccagatc	9480
tgagtcgacc	tgaggcatg	cccgtgaaa	tcaccagtct	ctctctacaa	atctatctct	9540
ctctataata	atgtgtgagt	agttcccaga	taagggaatt	agggttctta	tagggtttcg	9600
ctcatgtggt	gagcatataa	gaaaccctta	gtatgtattt	gtatttgtaa	aatacttcta	9660
tcaataaaat	ttctaattcc	taaaaccaa	atccagggcg	agctcggtac	ccggggatcc	9720
tctagagtcc	acctgcagcc	atgcccgcgg	atategatgg	gccccggccg	aagcttcggt	9780
ccgggccatc	gtggcctctt	gctcttcagg	atgaagagct	atgtttaaac	gtgcaagcgc	9840
tcaattccgc	ctatagttag	tcgtattaca	atcgtacgca	atcagtagca	ttaaaaacgt	9900
ccgcaatggt	ttattaagtt	gtctaagcgt	caatttttcc	cttctatggt	ccggtttggt	9960
tatctctaa	attatataat	ccagcttaaa	taagttaa	gacaaacaaa	caacacagat	10020
tattaaatag	attatgtaat	ctagatcct	agattatgta	atccataagt	agaatatcag	10080
gtgcttatat	aatctatgag	ctcgattata	taatcttaa	agaaaaacaaa	cagagcccct	10140
ataaaaagg	gtcaagtgga	cacttgggtca	ctcatttaat	ccctccctct	cctcttttat	10200
ccctctttt	ggtgtattca	ccaatagtgg	tgtgcacctg	tgattggctc	gtaaaaattc	10260
ttggacggat	ggaagagtga	agagataaag	aagcaaaaga	aaagtaacaa	cgaagcttca	10320
tcagctacaa	atthttggccc	aactggttgc	accagcacca	aacttacgta	tacatgatta	10380
tctctgtttc	cctcatttcg	aagaaaaaaa	cgggtttcaa	aacccactgc	ttcaggagt	10440
aaaaaaagat	aataatctga	aacattgctt	ccaccttggc	ccttatttgg	ttacgttgca	10500
atccaccca	atccacatgt	ggattgagat	ggattgcagt	gtagctagac	aaaccttag	10560
gcccgtttg	cataggaata	caccaggaat	tattccagct	aatcaaaatt	tatataaatg	10620
agagaaacaa	ttcgatagg	aattgttcca	ggacttcatt	ctgcagtaac	cgaacggccc	10680
cttaatccac	cccaatacac	gtggattgga	gtggattgag	gtacagccaa	acaaggccta	10740
agtgcagatc	aaataaatca	cccgctatat	tcttctacct	acaaaaacag	caataaacac	10800
ctgaatgaag	ttctaatttg	cacagtgtag	gtaggatgaa	aatagttacc	tcctcatggt	10860

-continued

```

cagtaactct tggcacacaa cttcacatgt aatcgatgta ccacttggtct cttgcctgaa 10920
acccaataca tcttttagcat aagaataata ttatgatggc aaggcatgat caccagcact 10980
cctttattgt ttagtaagtc taccactccc caaaacaatt caaatgaaca gagatgcatt 11040
gcccccaatg aattctatct caattagccg gaaaattcta cttcatcaga agcatccaaa 11100
ttgccagcat ccctactaga ctgacatga ccaggctgcc gcagatgctt ctttttctgt 11160
cctctctctt ttgcttgtag tttctcttca agatccctca ccccagctct cttatacatc 11220
ttaaagctaa catgtctctc ctccgcatc ttcctaacct tctcagtaat ctcagcagca 11280
atctgacggt tgtacaactt cttcagcccc ttcataact ttgcaaatgt gtcaggctgt 11340
ggcatcagtc ctgctcttag catgtctaag caatacaggc aggctctctt gacatgtttc 11400
ttcgcaaaac gtgcatgaat ccagatagtc catgactca cattgagctc acagcctttg 11460
ctcacaatac atttccaac atcctttgca agctcaagtt tctcatctct gaccaacgca 11520
ttgaggagggt ccttcagcac cccatattgc ggtaccacaa agagcccctt cccaacctg 11580
tctttaaat aactacatgc ctcaatcagc aaacctgcc caacaaggcc actcaccacg 11640
atagcaaatg tctgaccac aggactgagc ccagcacttt ccatctcatt ccacaatgct 11700
atggcttctt tggtctcccc aagcctgcag gcccaaccgaa tcaccacatt gtatatcttg 11760
agatctggtg gacaccggca ctcccgcac ctctccatca gctccaagca ctctcaagc 11820
tgctcctctt tctcgtgtgc tacaaagaaa ccatggtaca cggcagcgtc caccgcagc 11880
ccatccctcg acatagcatc caagaactcg taccctggg at 11922

```

<210> SEQ ID NO 24

<211> LENGTH: 7390

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: This sequence represents the DNA molecule used to transform maize line DAS59122-7 and represents insert PHI 17662A.

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)...(25)

<223> OTHER INFORMATION: T-DNA right border

<221> NAME/KEY: misc_feature

<222> LOCATION: (7366)...(7390)

<223> OTHER INFORMATION: T-DNA left border

<400> SEQUENCE: 24

```

gtttaccgc caatatatcc tgcacaacac tgatagtta aactgaaggc gggaaacgac 60
aatctgatca tgagcggaga attaaggag tcaagttatg accccgccc atgacgcggg 120
acaagccgtt ttacgtttgg aactgacaga accgcaactg tgaaggagcc actcagcaag 180
cttactagta gcgctgttta aacgctcttc aactggaaga gcggttacc ggaccgaagc 240
ttgcatgctt gcagtgcagc gtgaccgggt cgtgcccctc tctagagata atgagcattg 300
catgtctaag ttataaaaaa ttaccacata tttttttgt cacacttgtt tgaagtgcag 360
tttatctatc tttatacata tatttaaac ttactctacg aataatataa tctatagtac 420
tacaataata tcagtgtttt agagaatcat ataatgaac agttagacat ggtctaaagg 480
acaattgagt attttgacaa caggactcta cagttttatc tttttagtgt gcatgtgttc 540
tccttttttt ttgcaaatag cttcacctat ataatacttc atccatttta ttagtacatc 600
catttagggt ttagggtaa tggtttttat agactaattt ttttagtaca tctattttat 660
tctattttag cctctaaatt aagaaaacta aaactctatt ttagttttt tatttaataa 720
tttagatata aaatagaata aaataaagt actaaaaatt aaacaaatc cctttaagaa 780

```

-continued

atataaaaaa	ctaaggaaac	atattttcttg	tttcgagtag	ataatgccag	cctgttaaac	840
gccgtcgacg	agtctaacgg	acaccaacca	gcgaaccagc	agcgtcgcgt	cgggccaagc	900
gaagcagacg	gcacggcatc	tctgtcgtcg	cctctggacc	cctctcgaga	gttccgctcc	960
accgttgac	ttgctcgcgt	gtcggcatcc	agaaattgcg	tggcggagcg	gcagacgtga	1020
gccggcacgg	caggcggcct	cctcctcctc	tcacggcacc	ggcagctacg	ggggattcct	1080
ttcccaccgc	tccttcgctt	tcctctcctc	gcccgccgta	ataaatagac	accccctcca	1140
cacctctttt	ccccaacctc	gtgttgctcg	gagcgcacac	acacacaacc	agatctcccc	1200
caaatccacc	cgtcggcacc	tcgcctcaa	ggtacgcgc	tcgtcctccc	ccccccccc	1260
tctctacctt	ctctagatcg	gcgttcgggt	ccatggtag	ggcccggtag	ttctacttct	1320
gttcatgttt	gtgtagatc	cgtgtttgtg	ttagatccgt	gctgctagcg	ttcgtacacg	1380
gatgcgacct	gtacgtcaga	cacgttctga	ttgctaactt	gccagtgttt	ctctttgggg	1440
aatcctggga	tggctctagc	cgttccgcag	acgggatcga	tttcatgatt	ttttttgttt	1500
cgttgcatag	ggtttggttt	gcccttttcc	tttatttcaa	tatatgccgt	gcacttgttt	1560
gtcgggtcat	cttttcatgc	ttttttttgt	cttggttggtg	atgatgtggt	ctggttgggc	1620
ggtcgttcta	gatcggagta	gaattctggt	tcaaactacc	tgggtgattt	attaattttg	1680
gatctgtatg	tgtgtgccat	acatattcat	agttacgaat	tgaagatgat	ggatggaaat	1740
atcgatctag	gataggtata	catgttgatg	cgggttttac	tgatgcatat	acagagatgc	1800
tttttgctcg	cttggttggtg	atgatgtggt	gtggttgggc	ggtcggttcat	tcggttctaga	1860
tcggagttaga	atactgtttc	aaactacctg	gtgtatttat	taattttgga	actgtatgtg	1920
tgtgtcatac	atcttcatag	ttacgagttt	aagatggatg	gaaatctcga	tgtaggatag	1980
gtatacatgt	tgatgtgggt	tttactgatg	catatacatg	atggcatatg	cagcatctat	2040
tcataatgctc	taaccttgag	tacctatcta	ttataataaa	caagtatggt	ttataattat	2100
tttgatcttg	atatacttgg	atgatggcat	atgcagcagc	tatatgtgga	tttttttagc	2160
cctgccttca	tacgctatct	atttgccttg	tactgtttct	tttgcgatg	ctcaccctgt	2220
tgtttgggtg	tacttctgca	ggtcgactct	agaggatcca	cacgacacca	tgtccgccc	2280
cgagggtgac	atcgacgtga	acaacaagac	cggccacacc	ctccagctgg	aggacaagac	2340
caagctcgac	ggcggcaggt	ggcgcacctc	cccaccaac	gtggccaacg	accagatcaa	2400
gaccttcgtg	gccgaatcca	acggcttcat	gaccggcacc	gagggcacca	tctactactc	2460
aattaatggc	gaggccgaga	tcagcctcta	cttcgacaac	ccgttcgccg	gctccaacaa	2520
atacgacggc	cactccaaca	agtcaccagta	cgagatcctc	acccagggcg	gctccggcaa	2580
ccagtcctac	gtgacctaca	ccatccagac	cacctcctcc	cgtacggcc	acaagtctcg	2640
agtcatgagt	catgagtcag	ttaacctaga	cttgtccatc	ttctggattg	gccaacttaa	2700
ttaatgtatg	aaataaaagg	atgcacacat	agtgacatgc	taatcactat	aatgtgggca	2760
tcaaagtgtg	gtgttatgtg	taattactag	ttatctgaat	aaaagagaaa	gagatcatcc	2820
atatttctta	tcctaaatga	atgtcacgtg	tctttataat	tctttgatga	accagatgca	2880
tttcattaac	caaatccata	tacataataa	tattaatcat	atataattaa	tatcaattgg	2940
gttagcaaaa	caaatctagt	ctaggtgtgt	tttgccaatg	cggccgcgga	ccgaattggg	3000
gatctgcatg	aaagaaactg	tcgcactgct	gaaccgcacc	ttgtcacttt	catcgaacac	3060
gacctgtgcc	caagatgacg	gtgctgcgggt	ctaagtgagg	ctgaattgcc	ttggacagaa	3120

-continued

gctgactccc	tacaattagt	taggccaac	ggtgcatcca	tgtgtagctc	cgggctcggg	3180
ctgtatcgcc	atctgcaata	gcatccatgg	agctcgttcc	atgtagttagg	agatgaacca	3240
atgatcgggc	gtgtggaagt	atgttcctgt	gtactccgat	agtagagtac	gtgttagctc	3300
tttcatgggt	caagtgaat	ttgtgttgg	ttaattacc	ctacgttagt	tgcgggacag	3360
gagacacatc	atgaatttaa	aggcgatgat	gtcctctcct	gtaatgttat	tcttttgatg	3420
tgatgaatca	aatgtcata	taaaacattt	gttgcctttt	agttaggcct	gatcgtagaa	3480
cgaaatgctc	gtgtagcggg	gctacgagcc	tatgacgcaa	taacactggt	ttgccggccc	3540
ggagtcgctt	gacaaaaaaaa	agcatgttaa	gtttatttac	aattcaaac	ctaacaatatt	3600
atattccctc	aaagcaggtt	cacgatcaca	cctgtaccta	aaaaaaaaacat	gaagaatata	3660
ttactccatt	attatgagat	gaaccacttg	gcaagagtgg	taagctatat	aaaaaaaaatga	3720
acattattac	gagatgttat	atgccattat	attgattcga	agatatatgt	ttctttctcc	3780
cacgggcacc	taacggatac	atgataaggc	caaggcagat	cacgggaaat	tattcgaata	3840
catgttacgc	cctattgccc	gaaaaaaaaat	gcagggcagg	tgttggccgt	agcgatttaa	3900
gcaactaagc	tggaggttgc	cacacttgg	tgcaagcgtc	tgacccttct	aaaaaatcgg	3960
cggtttgtc	cgtatccgta	tcccctatcc	aacatctagc	tggccacacg	acggggctgg	4020
gcagatcgtg	gatgccgggt	cgacgtcgat	cgtcagccat	catagaccaa	tcgaccatct	4080
gttatggatg	cttctagct	agactagtca	gacataaaat	ttggatactt	tctccaact	4140
gggagacggg	gactgatgtg	cagctgcacg	tgagetaaat	ttttccctat	aaatatgcat	4200
gaaatactgc	attatcttgc	cacagccact	gccacagcca	gataacaagt	gcagctggta	4260
gcacgcaacg	catagctctg	gacttgtagc	taggtagcca	accggatcca	cacgacacca	4320
tgctcgacac	caacaagggt	tacgagatca	gcaaccacgc	caacggcctc	tacgcccga	4380
cctacctctc	cctcgacgac	tccggcgtgt	ccctcatgaa	caagaacgac	gacgacatcg	4440
acgactacaa	cctcaagtgg	ttcctcttcc	cgatcgacga	cgaccagtac	atcatcacct	4500
cctacgccgc	caacaactgc	aaggtgtgga	acgtgaacaa	cgacaagatt	aatgtgtcaa	4560
cctactcctc	caccaactcc	atccagaagt	ggcagatcaa	ggccaacggc	tcctcctacg	4620
tgatccagtc	cgacaacggc	aaggtgctca	cgcggcgac	cgccaggcc	ctcggcctca	4680
tccgcctcac	cgacgagtc	tccaacaacc	cgaaccagca	atggaacctg	acgtccgtgc	4740
agaccatcca	gctcccgcag	aagccgatca	tcgacaccaa	gctcaaggac	taccogaagt	4800
actccccgac	cggcaacatc	gacaacggca	cctccccgca	gctcatgggc	tggaccctcg	4860
tgccgtgcat	catggtgaac	gacccgaaca	tcgacaagaa	caccagatc	aagaccacc	4920
cgtactacat	cctcaagaag	taccagtact	ggcagagggc	cgtgggtctc	aacgtcgcgc	4980
tccgcccgca	cgagaagaag	tcttacacct	acgagtgggg	caccgagatc	gaccagaaga	5040
ccaccatcat	caacaccctc	ggcttccaga	tcaacatcga	cagcggcatg	aagttcgaca	5100
tcccggaggt	gggcggcgg	accgacgaga	tcaagaccca	gctcaacgag	gagctcaaga	5160
tcgagtattc	acatgagacg	aagatcatgg	agaagtacca	ggagcagtcc	gagatcgaca	5220
acccgaccga	ccagtccatg	aactccatcg	gcttctctac	catcacctcc	ctggagctct	5280
accgctacaa	cggctccgag	atccgcatca	tcgagatcca	gacctccgac	aacgacacct	5340
acaacgtgac	ctcctaccgc	aaccaccagc	aggcctctgt	gctgctgacc	aaccactcct	5400
acgaggaggt	ggaggagatc	accaacatcc	cgaagtccac	cctcaagaag	ctcaagaagt	5460
actacttctg	agtcatgagt	catgagtcag	ttaacctaga	cttgtccatc	ttctggattg	5520

-continued

```

gccaacttaa ttaatgtatg aaataaaagg atgcacacat agtgacatgc taatcactat 5580
aatgtgggca tcaaagtgtg gtgttatgtg taattactag ttatctgaat aaaagagaaa 5640
gagatcatcc atattttcta tcctaataatga atgtcacgtg tctttataat tctttgatga 5700
accagatgca tttcattaac caaatccata tacatataaa tattaatcat atataattaa 5760
tatcaattgg gttagcaaaa caaatctagt ctagggtgtg tttgogaatt cccatggagt 5820
caaagattca aatagaggac ctaacagAAC tcgccgtaaa gactggcgaa cagttcatac 5880
agagtctctt acgactcaat gacaagaaga aaatcttctg caacatgggtg gagcacgaca 5940
cgcttgctca ctccaaaaat atcaaagata cagtctcaga agaccAAagg gcaattgaga 6000
cttttcaaca aagggttaata tccggaAAC tctctggatt ccattgcca gctatctgtc 6060
actttattgt gaagatagtg gaaaaggaag gtggctccta caaatgccat cattgcgata 6120
aaggaaaggc catcgttgaa gatgcctctg ccgacagtgg tcccaaagat ggacccccac 6180
ccacgaggag catcgtggaa aaagaagacg ttccaaccac gtcttcaaag caagtggatt 6240
gatgtgatat ctccactgac gtaagggatg acgcacaatc ccactatcct tcgcaagacc 6300
cttctctat ataaggaagt tcatttcatt tggagaggac agggtagccg gggatccacc 6360
atgtctccgg agaggagacc agttgagatt aggccagcta cagcagctga tatggccgcg 6420
gtttgtgata tcgttaacca ttacattgag acgtctacag tgaactttag gacagagcca 6480
caaacaccac aagagtggat tgatgatcta gagaggttgc aagatagata cccttggttg 6540
gttgctgagg ttgaggtgtg tgtggctggt attgcttacg ctgggacctg gaaggctagg 6600
aacgcttacg attggacagt tgagagtact gtttacgtg cacaataggca tcaaagggtg 6660
ggcctaggat ccacattgta cacacatttg cttaagtcta tggaggcgca aggttttaag 6720
tctgtggttg ctgttatagg ccttccaaac gatccatctg ttagggtgca tgaggctttg 6780
ggatacacag cccgggttac attgctcgca gctggataca agcatgggtg atggcatgat 6840
gttggttttt ggcaaagga ttttgagttg ccagctcctc caaggccagt taggccagtt 6900
accagatctc gagtcgacct gcaggcatgc ccgctgaaat caccagtctc tctctacaaa 6960
tctatctctc tctataataa tgtgtgagta gttcccagat aagggaaatta gggttcttat 7020
agggttctgc tcatgtgttg agcatataag aaacccttag tatgtatttg tatttgtaaa 7080
atacttctat caataaaatt tctaattcct aaaacaaaa tccagggcga gctcggtagc 7140
cggggatcct ctagagtoga cctgcaggca tgcccgcgga tatcgatggg ccccgccga 7200
agcttcggtc cgggcatcgc tggcctcttg ctcttcagga tgaagagcta tgtttaaagc 7260
tgcaagcgct caattcgccc tatagtgagt cgtattacaa tcgtacgcaa ttcagtacat 7320
taaaaacgtc cgcaatgtgt tattaagttg tctaagcgtc aatttgttta caccacaata 7380
tatcctgcca 7390

```

<210> SEQ ID NO 25

<211> LENGTH: 2501

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Amplicon 22I-1

<400> SEQUENCE: 25

```

gcgggacaag ccgttttacg tttggaactg acagaaccgc aacgttgaag gagccactca 60
gcaagcttac tagtagcgct gtttaaagcgc tcttcaactg gaagagcggt taccgggacc 120

```

-continued

gaagcttgca	tgectgcagt	gcagcgtgac	ccggtcgtgc	ccctctctag	agataatgag	180
cattgcatgt	ctaagttata	aaaattacc	acatattttt	tttgtcacac	ttgtttgaag	240
tcagctttat	ctatctttat	acatatattt	aaactttact	ctacgaataa	tataatctat	300
agtactacaa	taatatacagt	gttttagaga	atcatataaa	tgaacagtta	gacatggctct	360
aaaggacaat	tgagtatttt	gacaacagga	ctctacagtt	ttatcttttt	agtgtgcatg	420
tgttctcctt	tttttttgca	aatagcttca	cctatataat	acttcatcca	ttttattagt	480
acatccattt	agggtttagg	gttaatgggt	tttatagact	aattttttta	gtacatctat	540
tttattctat	tttagcctct	aaattaagaa	aactaaaact	ctatttttagt	ttttttattt	600
aataattttag	atataaaaata	gaataaaaata	aagtgactaa	aaattaaaca	aatacccttt	660
aagaaattaa	aaaaactaag	gaaacatttt	tcttgtttcg	agtagataat	gccagcctgt	720
taaacgccgt	cgacgagtct	aacggacacc	aaccagcgaa	ccagcagcgt	cgctcggggc	780
caagcgaagc	agacggcacg	gcactctctgt	cgctgcctct	ggacccctct	cgagagtcc	840
gctccaccgt	tggacttget	ccgctgtcgg	catccagaaa	ttgcgtggcg	gagcggcaga	900
cgtgagccgg	cacggcaggc	ggcctctctc	tctctcaccg	gcaccggcag	ctacggggga	960
ttcctttccc	accgctcctt	cgttttccct	tctctgcccg	ccgtaataaa	tagacacccc	1020
ctccacaccc	tctttcccca	acctcgtgtt	gttcggagcg	cacacacaca	caaccagatc	1080
tccccaaat	ccaccgctcg	gcacctccgc	ttcaaggtag	gccgctcgtc	ctccccccc	1140
ccccctctct	accttctcta	gatcggcggt	ccggtccatg	gttagggccc	ggtagttcta	1200
cttctgttca	tgtttgtgtt	agatccgtgt	ttgtgttaga	tccgtgctgc	tagcgttctg	1260
acacggatgc	gacctgtacg	tcagacacgt	tctgattgct	aacttgccag	tgtttctctt	1320
tggggaatcc	tgggatggct	ctagccgttc	cgcagacggg	atcgatttca	tgattttttt	1380
tgtttcgttg	catagggttt	ggtttgccct	tttcttttat	ttcaatata	gccgtgcaact	1440
tgtttgcgg	gtcatctttt	catgcttttt	tttgtcttgg	ttgtgatgat	gtggtctggt	1500
tgggcggtcg	ttctagatcg	gagtagaatt	ctgtttcaaa	ctacctgggtg	gatttattaa	1560
ttttggatct	gtatgtgtgt	gccatacata	ttcatagtta	cgaattgaag	atgatggatg	1620
gaaatatacg	tctaggatag	gtatacatgt	tgatgcgggt	tttactgatg	catatacaga	1680
gatgcttttt	gttcgcttgg	ttgtgatgat	gtggtgtggt	tgggcggtcg	ttcattcgtt	1740
ctagatcgga	gtagaatact	gtttcaaact	acctgggtga	tttattaatt	ttggaactgt	1800
atgtgtgtgt	catacatctt	catagttacg	agtttaagat	ggatggaaat	atcgatgtag	1860
gataggtata	catgttgatg	tgggttttac	tgatgcatat	acatgatggc	atatgcagca	1920
tctattcata	tgctctaacc	ttgagtacct	atctattata	ataaacaagt	atgttttata	1980
attattttga	tcttgatata	cttggatgat	ggcatatgca	gcagctatat	gtggattttt	2040
ttagccctgc	cttcatacgc	tatttatttg	cttggtagctg	tttcttttgt	cgatgctcac	2100
cctgttgttt	ggtgttaact	ctgcaggctg	actctagagg	atccacacga	caccatgtcc	2160
gcccgcgagg	tgacatcga	cgtgaacaac	aagaccggcc	acaccctcca	gctggaggac	2220
aagaccaagc	tcgacggcgg	caggtggcgc	acctccccga	ccaacgtggc	caacgaccag	2280
atcaagacct	tcgtggccga	atccaacggc	ttcatgaccg	gcaccgaggg	caccatctac	2340
tactcaatta	atggcgaggc	cgagatcagc	ctctactctg	acaacccgtt	cgcgggtccc	2400
aacaaatacg	acggccactc	caacaagtcc	cagtacgaga	tcatacccca	gggcccgtcc	2460
ggcaaccagt	cccacgtgac	ctacaccatc	cagaccacct	c		2501

-continued

```

<210> SEQ ID NO 26
<211> LENGTH: 3027
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Amplicon 22I-2.

<400> SEQUENCE: 26
aacaacaaga cggccacac cctccagctg gaggacaaga ccaagctcga cggcggcagg      60
tggcgcacct ccccgaccaa cgtggccaac gaccagatca agaccttctg ggccgaatcc      120
aacggcttca tgaccggcac cgagggcacc atctactact caattaatgg cgaggccgag      180
atcagcctct acttcgacaa cccgttcgcc ggctccaaca aatcgcacgg ccaactccaac      240
aagtcccagt acgagatcat caccagggc ggctccggca accagtccca cgtgacctac      300
accatccaga ccacctcctc ccgctacggc cacaagtcct gagtcatgag tcatgagtca      360
gttaacctag acttgccat cttctggatt ggccaactta attaatgtat gaaataaaag      420
gatgcacaca tagtgacatg ctaatcacta taatgtgggc atcaaagttg tgtgttatgt      480
gtaattacta gttatctgaa taaaagagaa agagatcctc catatttctt atcctaattg      540
aatgtcacgt gtctttataa ttctttgatg aaccagatgc atttcattaa ccaaatccat      600
atacatataa atattaatca tatataatta atatcaattg ggtagcaaa acaaatctag      660
tctaggtgtg ttttgcaat gcggccgagg accgaattgg ggatctgcat gaaagaaact      720
gtcgcactgc tgaaccgcac cttgtcactt tcatcgaaca cgacctgtgc ccaagatgac      780
gggtgtcggg tctaagttag gctgaattgc cttggacaga agcggactcc ctacaattag      840
ttaggccaaa cgggtcatcc atgtgtagct ccgggctcgg gctgtatcgc catctgcaat      900
agcatccatg gagctcgttc catgtagttg gagatgaacc aatgatcggg cgtgtggacg      960
tatgttctct gtactccga tagtagagta cgtgttagct ctttcatggt gcaagtgaaa     1020
tttgtgttgg ttaattacc cctacgttag ttgcgggaca ggagacacat catgaattta     1080
aaggcgatga tgcctctcc tghtaatgta ttcttttgat gtgatgaatc aaaatgtcat     1140
ataaaacatt tgttgcctct tagttaggcc tgatcgtaga acgaaatgct cgtgtagcgg     1200
ggctacgagc ctatgacgca ataactcggg ttgcccggcc cggagtcgct tgacaaaaaa     1260
aagcatgtta agtttattta caattcaaaa cctaacatat tatattccct caaagcagg     1320
tcacgatcac acctgtacct aaaaaaaaaa tgaagaatat attactccat tattatgaga     1380
tgaaccactt ggcaagagtg gtaagctata taaaaaatg aacattatta cgagatgtta     1440
tatgccatta tattgattcg aagatatatg tttctttctc ccacgggcac ctaacggata     1500
catgataagg ccaaggcaga tcacgggaaa ttattcgaat acatgttacg ccctattgcc     1560
ggaaaaaaaa tgcagggcag gtgttggccg tagcgattta agcacttaag ctggaggttg     1620
ccacacttgg atgcaagcgt ctgaccttcc taaaacatcg gcggctttgt ccgtatccgt     1680
atcccctatc cgacatctag ctggccacac gacggggctg ggcagatcgt ggatgccggg     1740
tcgacgtcga tcgtcagcca tcatagacca atcgaccatc tgttatggat gcttgcctagc     1800
tagactagtc agacataaaa tttggatact ttctccaac tgggagacgg ggactgatgt     1860
gcagctgcac gtgagctaaa ttttcccta taaatatgca tgaataactg cattatcttg     1920
ccacagccac tgccacagcc agataacaag tgcagctggt agcacgcaac gcatagctct     1980
ggactttagc ctaggtagcc aaccggatcc acacgacacc atgctcgaca ccaacaaggt     2040

```

-continued

gtacgagatc agcaaccacg ccaacggcct ctacgccgcc acctacctct cctcgcagca	2100
ctccggcggtg tccctcatga acaagaacga cgacgacatc gacgactaca acctcaagtg	2160
gttctctctc cccgatcgacg acgaccagta catcatcacc tctacgccg ccaacaactg	2220
caaggtgtgg aacgtgaaca acgacaagat taatgtgtca acctactcct ccaccaactc	2280
catccagaag tggcagatca aggccaacgg ctccctctac gtgatccagt ccgacaacgg	2340
caaggtgtct accgccggca ccggccaggc cctcggcctc atccgcctca ccgacgagtc	2400
ctccaacaac ccgaaccagc aatggaacct gacgtccgtg cagaccatcc agctcccgca	2460
gaagccgac atcgacacca agctcaagga ctaccogaag tactccccga ccggcaacat	2520
cgacaacggc acctccccgc agctcatggg ctggaccctc gtgccgtgca tcatggtgaa	2580
cgaccogaac atcgacaaga acaccagat caagaccacc ccgtactaca tctcaagaa	2640
gtaccagtac tggcagaggg ccgtgggctc caacgtcgcg ctccgccgcg acgagaagaa	2700
gtcctacacc tacgagtggtg gcaccgagat cgaccagaag accaccatca tcaacacct	2760
cgcttccag atcaacatcg acagcggcat gaagtctgac atcccgagg tggcgggcgg	2820
taccgacgag atcaagacc agctcaacga ggagctcaag atcgagtatt cacatgagac	2880
gaagatcatg gagaagtacc aggagcagtc cgagatcgac aaccgaccg accagtccat	2940
gaactccatc ggcttctca ccatcaoctc cctggagctc taccgtaca acggtccga	3000
gatccgcatc atgcagatcc agacctc	3027

<210> SEQ ID NO 27

<211> LENGTH: 2830

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: PCR Amplicon 22I-3.

<400> SEQUENCE: 27

tacaacctca agtggttcct ctccccgatc gacgacgacc agtacatcat cacctctac	60
gccgccaaca actgcaaggt gtggaacgtg aacaacgaca agattaatgt gtcacctac	120
tctccacca actccatcca gaagtggcag atcaaggcca acggtctctc ctacgtgatc	180
cagtccgaca acggcaaggt gctcaccgcc ggcaccggcc aggcctcgg cctcatccgc	240
ctcaccgacg agtctctcaa caaccogaac cagcaatgga acctgacgtc cgtgcagacc	300
atccagctcc cgcagaagcc gatcatcgac accaagctca aggactacc gaagtaactc	360
ccgaccggca acatcgacaa cggcacctcc ccgacgtca tgggctggac cctcgtgccg	420
tgcatcatgg tgaacgaccc gaacatcgac aagaacaccc agatcaagac cccccgtac	480
tacatcctca agaagtacca gtactggcag agggccgtgg gctccaacgt cgcgctccgc	540
ccgcacgaga agaagtctca cacctacgag tggggcaccg agatcgacca gaagaccacc	600
atcatcaaca ccctcggctt ccagatcaac atcgacagcg gcatgaagtt cgacatcccg	660
gaggtgggcg gcggtaccga cgagatcaag acccagctca acgaggagct caagatcgag	720
tattcacatg agacgaagat catggagaag taccaggagc agtccgagat cgacaacccg	780
accgaccagt ccatgaactc catcggcttc ctccacctca cctccctgga gctctaccgc	840
tacaacggct ccgagatccg catcatgcag atccagacct ccgacaacga cacctacaac	900
gtgacctcct acccgaacca ccagcaggcc ctgctgctgc tgaccaacca ctctacgag	960
gaggtggagg agatcaccaa catcccgaag tccaccctca agaagctcaa gaagtactac	1020
ttctgagtca tgaatcatga gtcagttaac ctgacttgt ccatctctg gattggccaa	1080

-continued

```

cttaattaat gtagaaata aaaggatgca cacatagtg catgctaate actataatgt 1140
gggcatcaaa gttgtgtgtt atgtgtaatt actagttatc tgaataaaag agaaagagat 1200
catccatatt tcttatccta aatgaatgac acgtgtcttt ataattcttt gatgaaccag 1260
atgcatttca ttaaccaaat ccatatacat ataaatatta atcatatata attaatatca 1320
attgggtag caaaacaaat ctagtctagg tgtgttttgc gaattcccat ggagtcaaag 1380
attcaaatag aggacctaac agaactcgcc gtaaagactg gcgaacagtt catacagagt 1440
ctcttacgac tcaatgacaa gaagaaaatc ttcgtcaaca tgggtggagca cgacacgctt 1500
gttactcca aaaatatcaa agatacagtc tcagaagacc aaagggcaat tgagactttt 1560
caacaaaggg taatatcgg aaacctctc ggattccatt gccagctat ctgtcacttt 1620
atttgaaga tagtggaaaa ggaaggtgac tctacaaat gccatcattg cgataaagga 1680
aaggccatcg ttgaagatgc ctctgcccac agtgggtcca aagatggacc cccaccacg 1740
aggagcatcg tggaaaaaga agacgttcca accacgtctt caaagcaagt ggattgatgt 1800
gatatctcca ctgacgtaag ggatgacgca caatccact atccttcgca agacccttcc 1860
tctatataag gaagttcatt tcatttgag aggacaggtt acccggggat ccaccatgac 1920
tccggagagg agaccagttg agattaggcc agctacagca gctgatatgg ccgcggtttg 1980
tgatateggt aaccattaca ttgagacgtc tacagtgaac tttaggacag agccacaaac 2040
accacaagag tggattgatg atctagagag gttgcaagat agataccctt ggttgggtgc 2100
tgaggttag ggtgttggg ctggtattgc ttacgctggg ccctggaagg ctaggaacgc 2160
ttacgattgg acagttgaga gtactgttta cgtgtcacat aggcataaaa ggttgggctt 2220
aggatccaca ttgtacacac atttgcttaa gtctatggag gcgcaaggtt ttaagtctgt 2280
ggttgctggt ataggccttc caaacgatcc atctgttagg ttgcatgagg ctttgggata 2340
cacagcccgg ggtacattgc gcgcagctgg atacaagcat ggtggatggc atgatgttg 2400
tttttggcaa agggattttg agttgcccgc tctccaagg ccagttaggc cagttacca 2460
gatctgagtc gacctgcagg catgcccgtt gaaatcacca gtctctctct acaaatctat 2520
ctctctctat aataatgtgt gagtagttcc cagataaggg aattagggtt cttatagggt 2580
ttcgtctatg tgttgagcat ataagaaacc cttagtatgt atttgtattt gtaaaatact 2640
tctatcaata aaatttctaa ttctaaaac caaaatccag ggcgagctcg gtaccggggg 2700
atcctctaga gtcgacctgc aggcagctcc gcggatatcg atgggccccg gccgaagctt 2760
cggtcggggc catcgtggcc tcttgctctt caggatgaag agctatgttt aaacgtgcaa 2820
gcgctcaatt 2830

```

```

<210> SEQ ID NO 28
<211> LENGTH: 136
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Amplicon 0784/0564

```

```

<400> SEQUENCE: 28

```

```

aatccacaag attggagcaa acgacaaaa attcacaagg attggcggct gacattgcca 60
gcgcgggatc gcatgcggcg gcgcgggccg ggcgagcac gggagcaggc gacagtcgag 120
ctccattgga acgtag 136

```

```

<210> SEQ ID NO 29

```

-continued

<211> LENGTH: 263
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Amplicon 0784/0543

<400> SEQUENCE: 29

aatccacaag attggagcaa acgaccaaaa attcacaagg attggcggct gacattgcca	60
gcgcgggatc gcatgcggcg gcggcggccg gggcgagcac gggagcaggc gacagtcgag	120
ctccattgga acgtagaat acttaagggc aaggtctcca aatacttgaa aaaataggaa	180
aaagaagaaa atacatgaaa tgatattgaa atcaattgga agatggtatg aatcttgttt	240
ttgcaaagcg aacgattcag atg	263

<210> SEQ ID NO 30
 <211> LENGTH: 227
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Amplicon 0569/0577

<400> SEQUENCE: 30

ggtcaagtgg acacttggtc actcatttaa tccctccctc tccctcttta tccctctttt	60
tggtgtattc accaatagtg gtgtgcacct gtgattggct cgtaaaaatt cttggacgga	120
tggaagagtg aagagataag caagtcaaag aaaagtaaca acgaagcttc atcagctaca	180
aattttggcc caactggttg caccagcacc aaacttacgt atacatg	227

<210> SEQ ID NO 31
 <211> LENGTH: 492
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Amplicon 0570/0542

<400> SEQUENCE: 31

gagtgaagag ataagcaagt caaagaaaag taacaacgaa gtttcatcag ctacaaattt	60
tggcccaact ggttgcacca gcaccaaact tacgtataca tgattatctc tgtttccctc	120
atctgaaaga aaaaaacggg tttcaaaacc cactgctttc aggagtaaaa aaagataata	180
atctgaaaca ttgcttccac cttggccctt atttggttac gttgcaattc accccaatcc	240
acatgtggat tgagatggat tgcagtgtag ctagacaaac ccttaggccc tgtttgcata	300
ggaatacacc aggaattatt ccagctaatc aaaatttata taaatgagag aaacaattcg	360
gataggaatt gttccaggac ttcattctgc agtaaccgaa cggcccctta atccacccca	420
atacacgtgg attggagtgg attgaggtac agccaaacaa ggcctaagtg cagatcaaat	480
aaatcacccg tc	492

<210> SEQ ID NO 32
 <211> LENGTH: 555
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Amplicon 0784/0215

<400> SEQUENCE: 32

aatccacaag attggagcaa acgaccaaaa attcacaagg attggcggct gacattgcca	60
gcgcgggatc gcatgcggcg gcggcggccg gggcgagcac gggagcaggc gacagtcgag	120
ctccattgga acgtagaat acttaagggc aaggtctcca aatacttgaa aaaataggaa	180

-continued

```

aaagaagaaa atacatgaaa tgatattgaa atcaattgga agatgttatg aatcttgttt 240
ttgcaaagcg aacgattcag atggcaaaac tatgaatctt tttgtttgaa gtcccaaata 300
taaaattttc tcgtactcac caacattggt gcgcacctgt gattggctca taaaaattct 360
tggaggggacg gaagaaagag tgaagggata agcaagtaaa agcgctcaa cactgatagt 420
ttaaactgaa ggcgggaaac gacaatctga tcatgagcgg agaattaagg gagtcacggt 480
atgacccccg ccgatgacgc gggacaagcc gttttacggt tggaactgac agaaccgcaa 540
cgttgaagga gccac 555

```

```

<210> SEQ ID NO 33
<211> LENGTH: 547
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Amplicon O219/0577

```

```

<400> SEQUENCE: 33

```

```

cgtgcaagcg ctcaattgc cctatagtga gtcgtattac aatcgtaagc aattcagtac 60
attaanaaacg tccgcaatgt gttattaagt tgtctaagcg tcaatttttc ccttctatgg 120
tcccgtttgt ttatcctcta aattatataa tccagcttaa ataagttaag agacaaacaa 180
acaacacaga ttattaaata gattatgtaa tctagatacc tagattatgt aatccataag 240
tagaatatca ggtgcttata taatctatga gctcgattat ataacttaa aagaaaacaa 300
acagagcccc tataaaaagg ggtcaagtgg acaactggtc actcatttaa tccctcctc 360
tcctctttta tccctctttt tgggttatc accaatagtg gtgtgcacct gtgattggct 420
cgtaaaaatt cttggacgga tgaagagtg aagagataag caagtcaaag aaaagtaaca 480
acgaagcttc atcagctaca aattttgccc caactgggtg caccagcacc aaacttacgt 540
atacatg 547

```

```

<210> SEQ ID NO 34
<211> LENGTH: 243
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Amplicon O506/0476

```

```

<400> SEQUENCE: 34

```

```

tctcgtactc accaacattg gtgcgcacct gtgattggct cataaaaatt cttggagggg 60
cggaagaaag agtgaagggg taagcaagta aaagcgtca aacctgata gtttaaactg 120
aaggcgggaa acgacaatct gatcatgagc ggagaattaa gggagtcacg ttatgacccc 180
cgccgatgac gcgggacaag ccgttttacg tttggaactg acagaaccgc aacgttgaag 240
gag 243

```

```

<210> SEQ ID NO 35
<211> LENGTH: 754
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Amplicon O447/0577

```

```

<400> SEQUENCE: 35

```

```

aaccttagt atgtatttgt atttgtaaaa tacttctatc aataaaattt ctaattccta 60
aaaccaaact ccagggcgag ctccgtaccg ggggatcctc tagagtcgac ctgcaggcat 120

```

-continued

```

gccccgggat atcgatgggc cccggccgaa gcttcgggcc gggccatcgt ggcctcttgc 180
tcttcaggat gaagagctat gtttaaacgt gcaagcgctc aattcgccct atagtgagtc 240
gtattacaat cgtacgcaat tcagtacatt aaaaacgtcc gcaatgtgtt attaagttgt 300
ctaagcgtea atttttccct tctatgggcc cgtttgttta tcctctaaat tatataatcc 360
agcttaaata agttaagaga caaacaaca acacagatta ttaaatagat tatgtaatct 420
agatacctag attatgtaat ccataagtag aatatcaggt gcttatataa tctatgagct 480
cgattatata atcttaaaag aaaacaaca gagccctat aaaaaggggt caagtggaca 540
cttggtcact catttaatcc ctccctctcc tcttttatcc ctctttttgg tgtattcacc 600
aatagtggtg tgcacctgtg attggctcgt aaaaattctt ggacggatgg aagagtgaag 660
agataagcaa gtcaaagaaa agtaacaacg aagcttcac agctacaaat ttggcccaa 720
ctggttgcac cagcaccaaa cttacgtata catg 754

```

```

<210> SEQ ID NO 36
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

```

```

<400> SEQUENCE: 36

```

```

cgtattacaa tcgtacgcaa ttcag 25

```

```

<210> SEQ ID NO 37
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide primer

```

```

<400> SEQUENCE: 37

```

```

ggataaacia acgggacccat agaag 25

```

```

<210> SEQ ID NO 38
<211> LENGTH: 104
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Amplicon of SEQ ID NOs: 36 and 37

```

```

<400> SEQUENCE: 38

```

```

cgtattacaa tcgtacgcaa ttcagtcac taaaaacgtc cgcaatgtgt tattaagttg 60

```

```

tctaagcgtc aatttttccc ttctatgggc ccgtttgttt atcc 104

```

```

<210> SEQ ID NO 39
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IVR1 (0197) Primer used to generate a 226 bp
amplicon as an internal positive control

```

```

<400> SEQUENCE: 39

```

```

ccgctgtatc acaagggctg gtacc 25

```

```

<210> SEQ ID NO 40
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: IVR2 (0198) Primer used to generate a 226 bp amplicon as an internal positive control

<400> SEQUENCE: 40

ggagcccgtg tagagcatga cgatc

25

What is claimed is:

1. A method of detecting the presence of DNA corresponding to maize DAS-59122-7 event in a sample, the method comprising:

(a) contacting the sample comprising maize DNA with a polynucleotide probe that hybridizes under stringent hybridization conditions with DNA from maize DAS-59122-7 event and does not hybridize under said stringent hybridization conditions with a non-DAS-59122-7 maize plant DNA;

(b) subjecting the sample and probe to stringent hybridization conditions; and

¹⁰ (c) detecting hybridization of the probe to the DNA, wherein detection of hybridization indicates the presence of the maize DAS-59122-7 event.

2. The method of claim 1, wherein the probe hybridizes to SEQ ID NO: 25.

¹⁵ 3. The method of claim 1, wherein the probe hybridizes to SEQ ID NO: 26.

4. The method of claim 1, wherein the probe hybridizes to SEQ ID NO: 27.

* * * * *