



US009580475B2

(12) **United States Patent**
Ross et al.

(10) **Patent No.:** **US 9,580,475 B2**
(45) **Date of Patent:** ***Feb. 28, 2017**

(54) **COMPUTATIONALLY OPTIMIZED
BROADLY REACTIVE ANTIGENS FOR H1N1
INFLUENZA**

2009/0291472 A1 11/2009 Lu et al.
2009/0327170 A1 12/2009 Donati et al.
2010/0041740 A1 2/2010 Wong et al.

(75) Inventors: **Ted M. Ross**, Port St. Lucie, FL (US);
Corey J. Crevar, North Huntingdon,
PA (US)

FOREIGN PATENT DOCUMENTS

(73) Assignee: **University of Pittsburgh—Of the
Commonwealth System of Higher
Education**, Pittsburgh, PA (US)

JP	2004-501647	1/2004
WO	WO 2002/000885	1/2002
WO	WO 2005/020889	3/2005
WO	WO 2007/011904	1/2007
WO	WO 2008/028946	3/2008
WO	WO 2009/073330	6/2009
WO	WO 2010/003225	1/2010
WO	WO 2010/036948	4/2010
WO	WO 2010/036970	4/2010
WO	WO 2010036948	* 4/2010
WO	WO 2010/115133	10/2010
WO	WO 2011/094358	8/2011
WO	WO 2012/036993	3/2012
WO	WO 2013/119683	8/2013
WO	WO 2013/122827	8/2013
WO	WO 2013/148164	10/2013

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 161 days.
This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **14/126,550**

(22) PCT Filed: **Jun. 20, 2012**

(86) PCT No.: **PCT/US2012/043347**
§ 371 (c)(1),
(2), (4) Date: **Dec. 16, 2013**

(87) PCT Pub. No.: **WO2012/177760**
PCT Pub. Date: **Dec. 27, 2012**

(65) **Prior Publication Data**
US 2014/0127248 A1 May 8, 2014

Related U.S. Application Data

(60) Provisional application No. 61/498,800, filed on Jun.
20, 2011.

(51) **Int. Cl.**
C07K 14/005 (2006.01)
A61K 39/145 (2006.01)
A61K 39/12 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
CPC **C07K 14/005** (2013.01); **A61K 39/12**
(2013.01); **A61K 39/145** (2013.01); **A61K**
2039/5258 (2013.01); **A61K 2039/55505**
(2013.01); **C12N 2760/16122** (2013.01); **C12N**
2760/16123 (2013.01); **C12N 2760/16134**
(2013.01)

(58) **Field of Classification Search**
CPC **C12N 2760/16134**; **A61K 39/145**;
A61K 39/12; **C07K 14/005**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,566,454 B2	7/2009	Lu et al.	
9,309,290 B2 *	4/2016	Ross	C07K 14/005
2005/0181459 A1	8/2005	Baker et al.	
2008/0045472 A1	2/2008	Brahmachari et al.	
2009/0074803 A1	3/2009	Sallberg et al.	

OTHER PUBLICATIONS

Sequence alignment of SEQ ID No. 1 with UniProt database access
No. A4U6Y5_9INFA by Ghedin et al on May 15, 2007.*
Sequence alignment of SEQ ID No. 1 with Published Applica-
tions_AA_Main database in U.S. Appl. No. 14/092,371 SEQ ID
No. 5 by Ross et al on Nov. 27, 2012.*
UniProt Accession No. A4U6Y5, May 15, 2007.
Weaver et al., "Protection against Divergent Influenza H1N1 Virus
by a Centralized Influenza Hemagglutinin," *PLoS ONE*
6(3):e18314, Mar. 28, 2011.
Cai et al., "A Computational Framework for Influenza Antigenic
Cartography," *PLoS Comput. Biol.*, vol. 6:e1000949, 2010.
Carter et al., "Complex Patterns of Human Antisera Reactivity to
Novel 2009 H1N1 and Historical H1N1 Influenza Strains," *PLoS*
ONE 7(7):e39435, 2012.
Ekiert et al., "Antibody recognition of a highly conserved influenza
virus epitope: implications for universal prevention and therapy,"
Science 324(5924):246-251, 2009.
GenBank Accession No. ABA55715, Oct. 8, 2005.
GenBank Accession No. ABD60856.1, Mar. 2, 2006.
GenBank Accession No. ABU99095, May 1, 2008.
GenBank Accession No. ABW21677, Mar. 12, 2010.
GenBank Accession No. EU195416, Mar. 12, 2010.
Giles et al., "A Computationally Optimized Hemagglutinin Virus-
Like Particle Elicits Broadly Reactive Antibodies that Protect
Nonhuman Primates from H5N1 Infection," *J Infect Dis*
205(10):1562-1570, 2012.

(Continued)

Primary Examiner — Shanon A Foley

(74) Attorney, Agent, or Firm — Klarquist Sparkman,
LLP

(57)

ABSTRACT

Described herein is the generation of optimized H1N1
influenza HA polypeptides for eliciting a broadly reactive
immune response to H1N1 influenza virus isolates. The
optimized HA polypeptides were developed through a series
of HA protein alignments, and subsequent generation of
consensus sequences, based on selected H1N1 viruses iso-
lated from 1918-2011. Provided herein are optimized H1N1
HA polypeptides, and compositions, fusion proteins and
VLPs comprising the HA polypeptides. Further provided are
codon-optimized nucleic acid sequences encoding the HA
polypeptides. Methods of eliciting an immune response
against influenza virus in a subject are also provided by the
present disclosure.

6 Claims, 7 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

Giles et al., "A Computationally Optimized Broadly Reactive Antigen (COBRA) Based H5N1 VLP Vaccine Elicits Broadly Reactive Antibodies in Mice and Ferrets," *Vaccine*, vol. 29:3043-3054, 2011.

Giles et al., "Computationally Optimized Antigens to Overcome Influenza Viral Diversity," *Expert Review of Vaccines*, vol. 11:267-269, 2012.

Giles et al., "Antibody Breadth and Protective Efficacy are Increased by Vaccination with Computationally Optimized Hemagglutinin but not with Polyvalent Hemagglutinin-Based H5N1 Virus-Like Particle Vaccines," *Clin. Vaccine Immunol.*, vol. 19:128-139, 2012.

Jiang et al., "Enhanced Protective Efficacy of H5 Subtype Avian Influenza DNA Vaccine with Codon Optimized HA Gene in a pCAGGS Plasmid Vector," *Antiviral Res.*, vol. 75:234-241, 2007.

Parida et al., "Computational Analysis of Proteome of H5N1 Avian Influenza Virus to Define T Cell Epitopes with Vaccine Potential," *Vaccine*, vol. 25:7530-7539, 2007.

Somvanshi et al., "Prediction of Epitopes in Hemagglutinin and Neuraminidase Proteins of Influenza A Virus H5N1 Strain: A Clue for Diagnostic and Vaccine Development," *OMICS* vol. 12:61-69, 2008.

Tang et al., "Hemagglutinin Displayed Baculovirus Protects Against Highly Pathogenic Influenza," *Vaccine*, vol. 28:6821-6831, 2010.

Tang et al., Characterization of Duck H5N1 Influenza Viruses with Differing Pathogenicity in Mallard (*Anas platyrhynchos*) Ducks, *Avian Pathol.*, vol. 38:457-467, 2009.

Tao et al., "Virus-Like Particle Vaccine Comprised of the HA, NA, and M1 Proteins of an Avian Isolated H5N1 Influenza Virus Induces

Protective Immunity Against Homologous and Heterologous Strains in Mice," *Viral Immunol.*, vol. 22:273-281, 2009.

Wang et al., "Hemagglutinin (HA) Proteins from H1 and H3 Serotypes of Influenza A Viruses Require Different Antigen Designs for the Induction of Optimal Protective Antibody Responses as Studied by Codon-Optimized HA DNA Vaccines," *J. Virol.*, vol. 80:11628-11637, 2006.

Chen et al., "A consensus-hemagglutinin-based DNA vaccine that protects mice against divergent H5N1 influenza viruses," *Proc Natl Acad Sci USA* 105(36):16538-13543, 2008.

Fenimore et al., "Designing and Testing Broadly-Protective Filoviral Vaccines Optimized for Cytotoxic T-Lymphocyte Epitope Coverage," *PLoS ONE* 7(10):e44769, 2012.

Laddy et al., "Immunogenicity of novel consensus-based DNA vaccines against avian influenza," *Vaccine* 25:2984-2989, 2007.

Prabakaran et al., "Neutralizing Epitopes of Influenza Virus Hemagglutinin: Target for the Development of a Universal Vaccine against H5N1 Lineages," *J Virol* 84(22):11822-11830, 2010.

GenBank Accession No. ABO44123, Mar. 22, 2007.

GenBank Accession No. ABQ44416, May 16, 2007.

GenBank Accession No. ABF47748, May 8, 2006.

Pushko et al., "Recombinant H1N1 Virus-Like Particle Vaccine Elicits Protective Immunity in Ferrets Against the 2009 Pandemic H1N1 Influenza Virus," *Vaccine*, vol. 28:4771-4776, 2010.


Uniprot Accession No. A4JZ83, version 33, deposited May 3, 2011 (2 pages).


Uniprot Accession No. A5GX63, version 32, deposited May 3, 2011 (2 pages).

Uniprot Accession No. B7NZ42, version 20, deposited May 3, 2011 (2 pages).

Zhao et al., "Induction of protection of against divergent H5N1 influenza viruses using a recombinant fusion protein linking influenza M2e to *Onchocerca volvulus* activation associated protein-1 (ASP-1) adjuvant," *Vaccine*, vol. 28:7233-7240, 2010.

* cited by examiner









FIG. 4
H1N1 Method 5 (1918-2011)








FIG. 6

H1 COBRA Method 1 HAIs





1

COMPUTATIONALLY OPTIMIZED BROADLY REACTIVE ANTIGENS FOR H1N1 INFLUENZA

CROSS REFERENCE TO RELATED APPLICATIONS

This application is the U.S. National Stage of International Application No. PCT/US2012/043347, filed Jun. 20, 2012, published in English under PCT Article 21(2), which claims the benefit of U.S. Provisional Application No. 61/498,800, filed Jun. 20, 2011, which is herein incorporated by reference in its entirety.

FIELD

This disclosure concerns optimized influenza hemagglutinin proteins that elicit broadly reactive immune responses to H1N1 virus isolates and their use as vaccines.

BACKGROUND

Influenza virus is a member of Orthomyxoviridae family. There are three subtypes of influenza viruses, designated influenza A, influenza B, and influenza C. The influenza virion contains a segmented negative-sense RNA genome, which encodes the following proteins: hemagglutinin (HA), neuraminidase (NA), matrix (M1), proton ion-channel protein (M2), nucleoprotein (NP), polymerase basic protein 1 (PB1), polymerase basic protein 2 (PB2), polymerase acidic protein (PA), and nonstructural protein 2 (NS2). The HA, NA, M1, and M2 are membrane associated, whereas NP, PB1, PB2, PA, and NS2 are nucleocapsid associated proteins. The M1 protein is the most abundant protein in influenza particles. The HA and NA proteins are envelope glycoproteins, responsible for virus attachment and penetration of the viral particles into the cell, and the sources of the major immunodominant epitopes for virus neutralization and protective immunity. Both HA and NA proteins are considered the most important components for prophylactic influenza vaccines.

Each year, seasonal influenza causes over 300,000 hospitalizations and 36,000 deaths in the U.S. alone (Simonsen et al., *Lancet Infect Dis* 7:658-66, 2007). The emergence of the novel H1N1 influenza virus in 2009 demonstrated how quickly a new influenza pandemic can sweep across the world.

There are currently two influenza vaccine approaches licensed in the United States—the inactivated, split vaccine and the live-attenuated virus vaccine. The inactivated vaccines can efficiently induce humoral immune responses but generally only poor cellular immune responses. Live virus vaccines cannot be administered to immunocompromised or pregnant patients due to their increased risk of infection. Thus, a need exists for a broadly protective influenza virus vaccine.

SUMMARY

Disclosed herein is the generation of optimized H1N1 influenza HA polypeptides for eliciting a broadly reactive immune response to H1N1 influenza virus isolates. The optimized HA polypeptides were developed through a series of HA protein alignments, and subsequent generation of consensus sequences, based on selected H1N1 viruses isolated from 1918-2011.

2

Provided herein are recombinant influenza HA polypeptides having an optimized amino acid sequence for eliciting a broadly reactive immune response against H1N1 influenza. In some embodiments, the HA polypeptide comprises an amino acid sequence at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11. In some embodiments, the amino acid sequence of the polypeptide comprises no more than 5, no more than 6, no more than 7, no more than 8, no more than 9 or no more than 10 amino acid substitutions relative to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11. In some embodiments, the influenza HA polypeptide lacks the N-terminal methionine residue.

Isolated nucleic acid molecules and vectors encoding the recombinant HA polypeptides are also provided by the present disclosure. Further provided are isolated cells comprising such vectors.

Also provided are influenza virus-like particles (VLPs) and fusion proteins comprising the optimized HA polypeptides disclosed herein.

Further provided are compositions that include the optimized influenza HA polypeptides, fusion proteins or VLPs disclosed herein in a pharmaceutically acceptable carrier. Methods of eliciting an immune response against influenza virus in a subject by administering the disclosed compositions, fusion proteins or VLPs is also provided by the present disclosure.

Also provided are methods of immunizing a subject against influenza virus by administering to the subject a composition comprising a VLP that contains an optimized HA polypeptide.

The foregoing and other objects, features, and advantages of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic of the process used to generate an H1N1 HA consensus sequence according to Method 1.

FIG. 2 is a schematic of the process used to generate an H1N1 HA consensus sequence according to Method 2.

FIG. 3 is a schematic of the process used to generate an H1N1 HA consensus sequence according to Method 4.

FIG. 4 is a schematic of the process used to generate an H1N1 HA consensus sequence according to Method 5.

FIG. 5 is a schematic of the process used to generate an H1N1 HA consensus sequence according to Method 6.

FIG. 6 is a graph showing hemagglutination-inhibition (HAI) serum antibody titers induced by COBRA H1N1 infection of mice. HAI serum antibody titers were determined in mice infected with COBRA H1N1 influenza VLPs (COBRA Method 1). Antisera were tested against the seasonal H1N1 influenza virus, A/New Calcdonia/20/1999. The results are represented as the log₂ transformed geometric mean titer (\pm S.E.M.) from antisera collected at weeks 3, 5, 8 and 12 post-infection.

FIG. 7 is Western blot showing expression of COBRA HA protein (Method 1 consensus sequence). COBRA H1N1 HA was translated in vitro and the cell culture lysates were analyzed by SDS-PAGE and Western blot. Lane 1, A/New Calcdonia/20/1999 virus; Lane 2, H1N1 COBRA secreted

HA; Lane 3, H1N1 COBRA 5 µg VLP; Lane 4, COBRA 10 µg VLP. The COBRA HA (lane 2) migrates at its expected molecular weight confirming expression of the synthetic protein.

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one stand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The Sequence Listing is submitted as an ASCII text file, created on Dec. 11, 2013, 67.6 KB, which is incorporated by reference herein. In the accompanying sequence listing:

SEQ ID NOs: 1-11 are the amino acid sequences of optimized H1N1 HA proteins.

SEQ ID NO: 12 is a consensus amino acid sequence of the optimized H1N1 HA proteins.

SEQ ID NO: 13 is the nucleic acid sequence of a codon-optimized H1N1 HA.

DETAILED DESCRIPTION

I. Abbreviations

COBRA: computationally optimized broadly reactive antigen

HA: hemagglutinin

HAI: hemagglutination inhibition

HRP: horseradish peroxidase

M1: matrix protein 1

NA: neuraminidase

PFU: plaque form unit

VLP: virus-like particle

II. Terms and Methods

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

Adjuvant: A substance or vehicle that non-specifically enhances the immune response to an antigen. Adjuvants can include a suspension of minerals (alum, aluminum hydroxide, or phosphate) on which antigen is adsorbed; or water-in-oil emulsion in which antigen solution is emulsified in mineral oil (for example, Freund's incomplete adjuvant), sometimes with the inclusion of killed mycobacteria (Freund's complete adjuvant) to further enhance antigenicity. Immunostimulatory oligonucleotides (such as those including a CpG motif) can also be used as adjuvants (for example, see U.S. Pat. Nos. 6,194,388; 6,207,646; 6,214,806; 6,218,371; 6,239,116; 6,339,068; 6,406,705; and 6,429,199). Adjuvants also include biological molecules, such as costimulatory molecules. Exemplary biological adjuvants

include IL-2, RANTES, GM-CSF, TNF- α , IFN- γ , G-CSF, LFA-3, CD72, B7-1, B7-2, OX-40L and 41 BBL.

Administer: As used herein, administering a composition to a subject means to give, apply or bring the composition into contact with the subject. Administration can be accomplished by any of a number of routes, such as, for example, topical, oral, subcutaneous, intramuscular, intraperitoneal, intravenous, intrathecal and intradermal.

Antibody: An immunoglobulin molecule produced by B lymphoid cells with a specific amino acid sequence. Antibodies are evoked in humans or other animals by a specific antigen (immunogen). Antibodies are characterized by reacting specifically with the antigen in some demonstrable way, antibody and antigen each being defined in terms of the other. "Eliciting an antibody response" refers to the ability of an antigen or other molecule to induce the production of antibodies.

Antigen: A compound, composition, or substance that can stimulate the production of antibodies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. An antigen reacts with the products of specific humoral or cellular immunity, including those induced by heterologous immunogens. In some embodiments of the disclosed compositions and methods, the antigen is an influenza HA protein.

Codon-optimized: A "codon-optimized" nucleic acid refers to a nucleic acid sequence that has been altered such that the codons are optimal for expression in a particular system (such as a particular species or group of species). For example, a nucleic acid sequence can be optimized for expression in mammalian cells. Codon optimization does not alter the amino acid sequence of the encoded protein.

Fusion protein: A protein generated by expression of a nucleic acid sequence engineered from nucleic acid sequences encoding at least a portion of two different (heterologous) proteins. To create a fusion protein, the nucleic acid sequences must be in the same reading frame and contain to internal stop codons. For example, a fusion protein includes an influenza HA fused to a heterologous protein.

Hemagglutinin (HA): An influenza virus surface glycoprotein. HA mediates binding of the virus particle to a host cells and subsequent entry of the virus into the host cell. The nucleotide and amino acid sequences of numerous influenza HA proteins are known in the art and are publically available, such as those deposited with GenBank. HA (along with NA) is one of the two major influenza virus antigenic determinants.

Immune response: A response of a cell of the immune system, such as a B-cell, T-cell, macrophage or polymorphonucleocyte, to a stimulus such as an antigen or vaccine. An immune response can include any cell of the body involved in a host defense response, including for example, an epithelial cell that secretes an interferon or a cytokine. An immune response includes, but is not limited to, an innate immune response or inflammation. As used herein, a protective immune response refers to an immune response that protects a subject from infection (prevents infection or prevents the development of disease associated with infection). Methods of measuring immune responses are well known in the art and include, for example, measuring proliferation and/or activity of lymphocytes (such as B or T cells), secretion of cytokines or chemokines, inflammation, antibody production and the like.

Immunogen: A compound, composition, or substance which is capable, under appropriate conditions, of stimulating an immune response, such as the production of antibodies.

ies or a T-cell response in an animal, including compositions that are injected or absorbed into an animal. As used herein, an “immunogenic composition” is a composition comprising an immunogen (such as an HA polypeptide).

Immunize: To render a subject protected from an infectious disease, such as by vaccination.

Influenza virus: A segmented negative-strand RNA virus that belongs to the Orthomyxoviridae family. There are three types of Influenza viruses, A, B and C. Influenza A viruses infect a wide variety of birds and mammals, including humans, horses, marine mammals, pigs, ferrets, and chickens. In animals, most influenza A viruses cause mild localized infections of the respiratory and intestinal tract. However, highly pathogenic influenza A strains, such as H5N1, cause systemic infections in poultry in which mortality may reach 100%. In 2009, H1N1 influenza was the most common cause of human influenza. A new strain of swine-origin H1N1 emerged in 2009 and was declared pandemic by the World Health Organization. This strain was referred to as “swine flu.” H1N1 influenza A viruses were also responsible for the Spanish flu pandemic in 1918, the Fort Dix outbreak in 1976, and the Russian flu epidemic in 1977-1978.

Isolated: An “isolated” biological component (such as a nucleic acid, protein or virus) has been substantially separated or purified away from other biological components (such as cell debris, or other proteins or nucleic acids). Biological components that have been “isolated” include those components purified by standard purification methods. The term also embraces recombinant nucleic acids, proteins or viruses (or VLPs), as well as chemically synthesized nucleic acids or peptides.

Linker: One or more amino acids that serve as a spacer between two polypeptides of a fusion protein.

Matrix (M1) protein: An influenza virus structural protein found within the viral envelope. M1 is thought to function in assembly and budding.

Neuraminidase (NA): An influenza virus membrane glycoprotein. NA is involved in the destruction of the cellular receptor for the viral HA by cleaving terminal sialic acid residues from carbohydrate moieties on the surfaces of infected cells. NA also cleaves sialic acid residues from viral proteins, preventing aggregation of viruses. NA (along with HA) is one of the two major influenza virus antigenic determinants.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

Optimized influenza HA protein: As used herein, “optimized influenza HA protein” refers to the HA protein consensus sequence generated by sequence alignments of selected H1N1 influenza viruses isolated between 1918 and 2011 (as described in Example 1 below). The nucleotide sequences encoding optimized HA proteins were (or can be) further optimized for expression in mammalian cells via codon-optimization and RNA optimization (such as to increase RNA stability). Optimized influenza HA proteins disclosed herein (and set forth herein as SEQ ID NOs: 1-11) are also referred to as “COBRA” sequences. Optimized HA polypeptides are designed to elicit broadly reactive immune responses in a subject. In the context of the present disclosure, “broadly reactive” means the protein sequence elicits

an immune response in a subject that is sufficient to inhibit, neutralize or prevent infection of a broad range of influenza viruses (such as most or all influenza viruses within a specific subtype). In some instances, the optimized influenza HA protein is capable of eliciting an immune response, such as a protective immune response, against most or all H1N1 influenza virus isolates.

Outbreak: As used herein, an influenza virus “outbreak” refers to a collection of virus isolates from within a single country in a given year.

Pharmaceutically acceptable vehicles: The pharmaceutically acceptable carriers (vehicles) useful in this disclosure are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, Pa., 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of one or more therapeutic compositions, such as one or more influenza vaccines, and additional pharmaceutical agents.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (for example, powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Polypeptide: A polymer in which the monomers are amino acid residues which are joined together through amide bonds. When the amino acids are alpha-amino acids, either the L-optical isomer or the D-optical isomer can be used. The terms “polypeptide” or “protein” as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. The term “polypeptide” is specifically intended to cover naturally occurring proteins, as well as those which are recombinantly or synthetically produced. The term “residue” or “amino acid residue” includes reference to an amino acid that is incorporated into a protein, polypeptide, or peptide.

Conservative amino acid substitutions are those substitutions that, when made, least interfere with the properties of the original protein, that is, the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. Examples of conservative substitutions are shown below.

Original Residue	Conservative Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
His	Asn; Gln
Ile	Leu, Val
Leu	Ile; Val
Lys	Arg; Gln; Glu
Met	Leu; Ile
Phe	Met; Leu; Tyr

-continued

Original Residue	Conservative Substitutions
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp; Phe
Val	Ile; Leu

Conservative substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

The substitutions which in general are expected to produce the greatest changes in protein properties will be non-conservative, for instance changes in which (a) a hydrophilic residue, for example, seryl or threonyl, is substituted for (or by) a hydrophobic residue, for example, leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, for example, lysyl, arginyl, or histadyl, is substituted for (or by) an electronegative residue, for example, glutamyl or aspartyl; or (d) a residue having a bulky side chain, for example, phenylalanine, is substituted for (or by) one not having a side chain, for example, glycine.

Preventing, treating or ameliorating a disease: "Preventing" a disease refers to inhibiting the full development of a disease. "Treating" refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. "Ameliorating" refers to the reduction in the number or severity of signs or symptoms of a disease.

Promoter: A promoter is an array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription. A promoter also optionally includes distal enhancer or repressor elements. A "constitutive promoter" is a promoter that is continuously active and is not subject to regulation by external signals or molecules. In contrast, the activity of an "inducible promoter" is regulated by an external signal or molecule (for example, a transcription factor). In some embodiments herein, the promoter is a CMV promoter.

Purified: The term "purified" does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified peptide, protein, virus, VLP or other active compound is one that is isolated in whole or in part from naturally associated proteins and other contaminants. In certain embodiments, the term "substantially purified" refers to a peptide, protein, virus, VLP or other active compound that has been isolated from a cell, cell culture medium, or other crude preparation and subjected to fractionation to remove various components of the initial preparation, such as proteins, cellular debris, and other components.

Recombinant: A recombinant nucleic acid, protein, virus or VLP is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques.

Sequence identity: The similarity between amino acid or nucleic acid sequences is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are. Homologs or variants of a given gene or protein will possess a relatively high degree of sequence identity when aligned using standard methods.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988; Higgins and Sharp, *Gene* 73:237-244, 1988; Higgins and Sharp, *CABIOS* 5:151-153, 1989; Corpet et al., *Nucleic Acids Research* 16:10881-10890, 1988; and Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988. Altschul et al., *Nature Genet.* 6:119-129, 1994.

The NCBI Basic Local Alignment Search Tool (BLAST™) (Altschul et al., *J. Mol. Biol.* 215:403-410, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, Md.) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx.

Subject: Living multi-cellular vertebrate organisms, a category that includes both human and non-human mammals, such as non-human primates.

Therapeutically effective amount: A quantity of a specified agent sufficient to achieve a desired effect in a subject being treated with that agent. For example, this may be the amount of an influenza virus vaccine useful for eliciting an immune response in a subject and/or for preventing infection or disease caused by influenza virus. Ideally, in the context of the present disclosure, a therapeutically effective amount of an influenza vaccine is an amount sufficient to increase resistance to, prevent, ameliorate, and/or treat infection caused by influenza virus in a subject without causing a substantial cytotoxic effect in the subject. The effective amount of an influenza vaccine useful for increasing resistance to, preventing, ameliorating, and/or treating infection in a subject will be dependent on, for example, the subject being treated, the manner of administration of the therapeutic composition and other factors.

Transformed cell: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Vaccine: A preparation of immunogenic material capable of stimulating an immune response, administered for the prevention, amelioration, or treatment of disease, such as an infectious disease. The immunogenic material may include, for example, attenuated or killed microorganisms (such as attenuated viruses), or antigenic proteins, peptides or DNA derived from them. Vaccines may elicit both prophylactic (preventative) and therapeutic responses. Methods of administration vary according to the vaccine, but may include inoculation, ingestion, inhalation or other forms of administration. Inoculations can be delivered by any of a number of routes, including parenteral, such as intravenous, subcutaneous or intramuscular. Vaccines may be administered with an adjuvant to boost the immune response.

Vector: A vector is a nucleic acid molecule allowing insertion of foreign nucleic acid without disrupting the ability of the vector to replicate and/or integrate in a host cell. A vector can include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. An insertional vector is capable of inserting itself into a host nucleic acid. A vector can also include one or more selectable marker genes and other genetic elements. An expression vector is a vector that contains the necessary regulatory sequences to allow transcription and translation of an inserted gene or genes. In some embodiments of the present disclosure, the vector encodes an influenza HA, NA or M1 protein. In some embodiments, the vector is the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross et al., *Nat. Immunol.* 1(2):102-103, 2000; Green et al., *Vaccine* 20:242-248, 2001).

Virus-like particle (VLP): Virus particles made up of one or more viral structural proteins, but lacking the viral genome. Because VLPs lack a viral genome, they are non-infectious. In addition, VLPs can often be produced by heterologous expression and can be easily purified. Most VLPs comprise at least a viral core protein that drives budding and release of particles from a host cell. One example of such a core protein is influenza M1. In some embodiments herein, an influenza VLP comprises the HA, NA and/or M1 proteins. Influenza VLPs can be produced by transfection of host cells with plasmids encoding the HA and NA proteins, and optionally the M1 protein. After incubation of the transfected cells for an appropriate time to allow for protein expression (such as for approximately 72 hours), VLPs can be isolated from cell culture supernatants. Example 2 provides an exemplary protocol for purifying influenza VLPs from cell supernatants. In this example, VLPs are isolated by low speed centrifugation (to remove cell debris), vacuum filtration and ultracentrifugation through 20% glycerol. Other methods of producing influenza VLPs are known in the art (see, for example, U.S. Patent Application Publication Nos. 2006/0263804; 2008/0031895; 2010/0166769; and 2010/0239610).

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Hence "comprising A or B" means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

III. Overview of Several Embodiments

Disclosed herein is the generation of optimized H1N1 influenza HA polypeptides for eliciting a broadly reactive immune response to H1N1 influenza. The optimized HA polypeptides were developed through a series of HA protein

alignments, and subsequent generation of consensus sequences, based on selected H1N1 viruses isolated from 1918-2011. The methods used to generate the 11 HA consensus sequences are described in Example 1 and FIGS. 1-5. The amino acid sequences of the 11 consensus HA polypeptides are set forth herein as SEQ ID NOs: 1-11. In addition, an amino acid consensus sequence of SEQ ID NOs: 1-11 is provided herein as SEQ ID NO: 12.

Provided herein are recombinant influenza HA polypeptides having an optimized amino acid sequence for eliciting a broadly reactive immune response against H1N1 influenza. In some embodiments, the HA polypeptide comprises an amino acid sequence at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11. In other embodiments, the amino acid sequence of the polypeptide comprises no more than 5, no more than 6, no more than 7, no more than 8, no more than 9 or no more than 10 amino acid substitutions relative to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11.

In particular embodiments, provided is a recombinant influenza HA polypeptide comprising an amino acid sequence at least 99% identical to SEQ ID NO: 1; at least 99.2% identical to SEQ ID NO: 2; at least 99% identical to SEQ ID NO: 3; at least 99% identical to SEQ ID NO: 4; at least 98% or at least 99% identical to SEQ ID NO: 5; at least 99% identical to SEQ ID NO: 6; at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 7; at least 99% identical to SEQ ID NO: 8; comprising SEQ ID NO: 9; at least 99% identical to SEQ ID NO: 10; or at least 99% identical to SEQ ID NO: 11.

In other particular embodiments, the recombinant influenza HA polypeptide comprises an amino acid sequence at least 99% identical to residues 2-566 of SEQ ID NO: 1; at least 99.2% identical to residues 2-566 of SEQ ID NO: 2; at least 99% identical to residues 2-566 of SEQ ID NO: 3; at least 99% identical to residues 2-566 of SEQ ID NO: 4; at least 98% or at least 99% identical to residues 2-566 of SEQ ID NO: 5; at least 99% identical to residues 2-566 of SEQ ID NO: 6; at least 97%, at least 98% or at least 99% identical to residues 2-566 of SEQ ID NO: 7; at least 99% identical to residues 2-566 of SEQ ID NO: 8; comprising residues 2-566 of SEQ ID NO: 9; at least 99% identical to residues 2-566 of SEQ ID NO: 10; or at least 99% identical to residues 2-566 of SEQ ID NO: 11.

In other embodiments, the amino acid sequence of the HA polypeptide comprises (i) no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 1; (ii) no more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 2; (iii) no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 3; (iv) no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 4; (v) no more than 10, no more than 9, no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 5; (vi) no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid

11

substitution(s) relative to SEQ ID NO: 6; (vii) no more than 10, no more than 9, no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 7; (viii) no more than 10, no more than 9, no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 8; (ix) no more than 8, no more than 7, no more than 6, no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 10; or (x) no more than 5, nor more than 4, no more than 3, no more than 2 or no more than 1 amino acid substitution(s) relative to SEQ ID NO: 11.

In some examples, the influenza HA polypeptide comprises or consists of the amino acid sequence of residues 2-566 of SEQ ID NO: 1, residues 2-566 of SEQ ID NO: 2, residues 2-566 of SEQ ID NO: 3, residues 2-566 of SEQ ID NO: 4, residues 2-566 of SEQ ID NO: 5, residues 2-566 of SEQ ID NO: 6, residues 2-566 of SEQ ID NO: 7, residues 2-566 of SEQ ID NO: 8, residues 2-566 of SEQ ID NO: 9, residues 2-566 of SEQ ID NO: 10 or residues 2-566 of SEQ ID NO: 11.

In other examples, the recombinant HA polypeptide comprises or consists of the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11.

In another example, the recombinant HA polypeptide comprises or consists of the amino acid sequence of SEQ ID NO: 12. In yet another example, the recombinant HA polypeptide comprises or consists of the amino acid sequence of residues 2-566 of SEQ ID NO: 12.

Further provided are isolated nucleic acid molecules encoding the recombinant HA polypeptides disclosed herein. In some embodiments, the nucleic acid molecule is codon-optimized for expression in mammalian cells. The nucleic acid molecule is optionally further optimized for RNA stability. In some embodiments, the sequence of the nucleic acid molecule is at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 13. In particular examples, the sequence of the nucleic acid molecule comprises or consists of SEQ ID NO: 13.

Vectors comprising the nucleic acid molecules encoding recombinant HA polypeptides are also provided by the present disclosure. The vector can be any suitable vector for expression of the HA polypeptide, such as a mammalian expression vector. In particular examples, the vector is the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798, herein incorporated by reference; Ross et al., *Nat. Immunol.* 1(2):102-103, 2000; Green et al., *Vaccine* 20:242-248, 2001).

In some examples, the vector includes a promoter operably linked to the nucleic acid sequence encoding the HA polypeptide. In particular examples, the promoter is a CMV promoter.

Also provided are isolated cells comprising the disclosed vectors. In some cases, the cell is any suitable cell type for production and expression of VLPs, such as a mammalian cell.

Further provided are influenza VLPs comprising an optimized HA polypeptide disclosed herein. The influenza VLPs can further include any additional influenza proteins necessary to form the virus particle. In some embodiments, the influenza VLPs further include influenza neuraminidase (NA) protein, influenza matrix (M1) protein, or both.

12

Also provided are influenza VLPs comprising an influenza HA polypeptide disclosed herein, produced by transfecting a host cell with a vector encoding the HA polypeptide, a vector encoding an influenza NA protein and a vector encoding an influenza M1 protein under conditions sufficient to allow for expression of the HA, M1 and NA proteins.

Fusion proteins comprising an optimized influenza HA polypeptide are further provided by the present disclosure.

Also provided herein are compositions comprising an optimized influenza HA protein as disclosed herein, or a fusion protein or VLP comprising the optimized influenza HA protein. In some embodiments, the compositions further comprise a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides).

Further provided is a method of eliciting an immune response to influenza virus in a subject by administering an optimized influenza HA protein, fusion proteins containing an optimized influenza HA, VLPs containing an optimized influenza HA, or compositions thereof, as disclosed herein. In some embodiments, the influenza virus is an H1N1 influenza virus. In some embodiments, the HA protein, HA fusion protein or VLP can be administered using any suitable route of administration, such as, for example, intramuscular. In some embodiments, the HA protein, fusion protein or VLP is administered as a composition further comprising a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides).

Also provided is a method of immunizing a subject against influenza virus by administering to the subject VLPs containing an optimized influenza HA protein disclosed herein, or administering a composition thereof. In some embodiments of the method, the composition further comprises a pharmaceutically acceptable carrier and/or an adjuvant. For example, the adjuvant can be alum, Freund's complete adjuvant, a biological adjuvant or immunostimulatory oligonucleotides (such as CpG oligonucleotides). In some embodiments, the VLPs (or compositions thereof) are administered intramuscularly.

In some embodiments of the methods of eliciting an immune response or immunizing a subject, the subject is administered about 1 to about 25 µg of the VLPs containing an optimized HA protein. In particular examples, the subject is administered about 5 to about 20 µg of the VLPs, or about 10 to about 15 µg of the VLPs. In one specific non-limiting example, the subject is administered about 15 µg of the VLPs. However, one of skill in the art is capable of determining a therapeutically effective amount (for example an amount that provides protection against H1N1 influenza virus infection) of VLPs to administer to a subject.

IV. Optimized H1N1 Influenza HA Polypeptides and Polynucleotides

Provided herein are 11 different optimized H1N1 HA polypeptide sequences. H1N1 HA amino acid sequences were downloaded from the NCBI Influenza Virus Resource database. H1N1 HA proteins from 1134 isolates from 1918-2011 were used for generating consensus sequences. Example 1 describes the methods that were used to generate each consensus sequence (see also FIGS. 1-5).

13

COBRA METHOD 1

(SEQ ID NO: 1)

MKAKLLVLLCAPTATYADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGNCSIAGWILGNPECESLFSKESWSYIVETP
 NSENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 ASCSHNGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQRAIYHTENAYVSVSSHYSRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKQCTPQ
 GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

COBRA METHOD 2

(SEQ ID NO: 2)

MKAKLLVLLCALATDADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGNCSIAGWILGNPECESLFSKESWSYIAETP
 NSENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 AACSHKGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQRAIYHTENAYVSVSSHYNRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKQCTPQ
 GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

COBRA METHOD 3

(SEQ ID NO: 3)

MKAKLLVLLCAPTATYADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGNCSVAGWILGNPECESLISKESWSYIVETP
 NPENGT CYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 ASCSHNGKSSFYRNLLWLTEKNGLYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQRALYHTENAYVSVSSHYSRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKQCTPQ
 GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

14

-continued

COBRA METHOD 4

(SEQ ID NO: 4)

MKAKLLVLLCALATDADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGKCSIAGWILGNPECESLLSKESWSYIAETP
 NSENGTCYPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 ASCSHKGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQRTIYRTENAYVSVSSNYNRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKQCTPQ
 GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

COBRA METHOD 5

(SEQ ID NO: 5)

MKAKLLVLLCTFTADADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGNCSIAGWILGNPECESLLSKESWSYIVETP
 NSENGTCYPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 AACSHAGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQALYQ TENAYVSVSSHYNRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMHECDTKQCTPQ
 GAINSSLPFQNIHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

COBRA METHOD 6

(SEQ ID NO: 6)

MKAKLLVLLCAPTATDADTICIGYHANNSTDVDTVLEKNVTVTHSVNLL
 ED SHNGKLCRLKGIAPLQLGKCSIAGWILGNPECESLLSKESWSYIVETP
 NSENGTCYPGDFADYEELREQLSSVSSFERFEIFPKESSWPNHVTGKGT
 AACSHAGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWG VHHPS
 NIGDQRALYHTENAYVSVSSHYNRRFTPEIAKRPKVRDQEGRINYWT
 LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKQCTPQ
 GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI
 AGFIEGGWTGMIDGWYGYHHQNEQSGSYAADQKSTQNAINGITNKVNSVI
 EKMNTQFTAVGKEFNKLERRMENLNKKVDDGFLDIWTYNAELLVLENER
 TLD FHD SNVKNLYEKVKSQ LKNNAKEIGNGC FEYHKCNNECMESVKNGT
 YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI
 SFWMCSNGSLQCRICI

15

-continued

COBRA METHOD 5 DEGLYCOSYLATED (SEQ ID NO: 7)

MKAKLLVLLCTFTATDADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDSHNGKLCRLKGIAPLQLGNCAIAGWILGNPECESLLSKKSWSYIVETP

NSENGTCTPGDFIDYEELREQLSSVSSFERFEIFPKESSWPNHAVTKGVT

AACSHAGKSSFYRNLLWLTEKNGSYPNLAKSYVNNKGKEVLVLWGVIHHP

NIGDQQALYQTENAYVSVSSHYNRKFTPEIAKRPKVRDQEGRINYWT

LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNAAMHECDTKCQTPQ

GAINSSLPFQNIHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI

AGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVI

EKMNTQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENER

TLDFHDSNVKNLYEKVKSQKLNNAKEIGNGCFEFYHKCNNECMESVKNGT

YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI

SFWMCSNGSLQCRICI

1918-1957 COBRA (SEQ ID NO: 8)

MKARLLVLLCALAATDADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDSHNGKLCRLKGIAPLQLGKCNIAAGWILGNPECESLLSNRSWSYIVETP

NSENGTCTPGDFIDYEELREQLSSVSSFERFEIFPKESSWPKNHTTKGVT

AACSHAGKSSFYRNLLWLTEKNGSYPNLSNSYVNNKGKEVLVLWGVIHHP

NIDDQQTLYQKENAYVSVSSYNNRRFTPEIAERPVRGQAGRMNYYWT

LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMHECDTKCQTPQ

GAINSSLPFQNIHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI

AGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVI

EKMNTQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENER

TLDFHDSNVKNLYEKVKSQKLRNNAKEIGNGCFEFYHKCDNECMESVKNGT

YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI

SFWMCSNGSLQCRICI

1977-2005 COBRA (SEQ ID NO: 9)

MKAKLLVLLCAPTATDADTICIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDSHNGKLCRLKGIAPLQLGNCSIAAGWILGNPECESLFSKKSWSYIAETP

NSENGTCTPGYFADYEELREQLSSVSSFERFEIFPKESSWPNHTVTKGVT

ASCSHGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVLVLWGVIHHP

NIGDQRAIYHTENAYVSVSSHYNNRRFTPEIAKRPKVRDQEGRINYWT

LEPGDTIIFEANGNLIAPWYAFALSRGFGSGIITSNASMDECDKACQTPQ

GAINSSLPFQNVHPVTIGECPKYVRSTKLRMTGLRNIPSIQSRGLFGAI

AGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAINGITNKVNSVI

EKMNTQFTAVGKEFNKLEKRMENLNKKVDDGFLDIWTYNAELLVLENER

TLDFHDSNVKNLYEKVKSQKLNNAKEIGNGCFEFYHKCNNECMESVKNGT

YDYPKYSEESKLNREKIDGVKLESMGVYQILAIYSTVASSLVLLVSLGAI

SFWMCSNGSLQCRICI

16

-continued

HUMAN SWINE COBRA (SEQ ID NO: 10)

MKAILLVLLCTFAATNADTLCIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDKHNGKLCCKLKGIAPLHLGKCNIAAGWLLGNPECELLLTASSWSYIVETS

NSDNGTCTPGDFIDYEELREQLSSVSSFERFEIFPKTSSWPNHETNKGVT

AACPYAGASSFYRNLIWLKKGNSYPKLSKSYVNNKGKEVLVLWGIHHP

TSTDQQSLYQNADAYVFGSSKYNRKFKPEIAARPKVRDQAGRMNYYWT

IEPGDTITFEATGNLVVPRYAFAMNRGSGSGIIISDAPVHDCNTKCQTPK

GAINSSLFPQNIHPVTIGECPKYVKSTKLRMTGLRNIPSIQSRGLFGAI

AGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAIDGITNKVNSVI

EKMNTQFTAVGKEFNHLEKRIENLNKKVDDGFLDIWTYNAELLVLENER

TLDFHDSNVKNLYEKVRSQKLNNAKEIGNGCFEFYHKCDDTCMESVKNGT

YDYPKYSEESKLNREEIDGVKLESTRIYQILAIYSTVASSLVLLVSLGAI

SFWMCSNGSLQCRICI

SWINE COBRA (SEQ ID NO: 11)

MKAILLVLLCTFTAANADTLCIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDRHNGKLCCKLGGIAPLHLGKCNIAAGWLLGNPECELLLTASSWSYIVETS

NSDNGTCTPGDFINYEELREQLSSVSSFERFEIFPKASSWPNHETNRGVT

AACPYAGANSFYRNLIWLKKGNSYPKLSKSYVNNKGKEVLVLWGIHHP

TSTDQQSLYQNADAYVFGSSKYNRKFKPEIAARPKVRGQAGRMNYYWT

LEPGDTITFEATGNLVVPRYAFAMNRGSGSGIIISDAPVHDCNTKCQTPK

GAINSSLFPQNIHPVTIGECPKYVKSTKLRMTGLRNIPSIQSRGLFGAI

AGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAIDGITNKVNSVI

EKMNTQFTAVGKEFNHLEKRIENLNKKVDDGFLDVWTYNAELLVLENER

TLDFHDSNVKNLYEKVRSQKLRNNAKEIGNGCFEFYHKCDDTCMESVKNGT

YDYPKYSEESKLNREEIDGVKLESTRIYQILAIYSTVASSLVLLVSLGAI

SFWMCSNGSLQCRICI

Consensus of SEQ ID NOs: 1-11 (SEQ ID NO: 12)

MKAXLLVLLCXXXAXXADTXCIGYHANNSTDTVDTVLEKNVTVTHSVNLL

EDXHNGKLCXLXGIAPLXLGXCCXAGWXLGNPECEXLXXXXXSWSYIXETX

NXXNGTCTPGXFXXYEELREQLSSVSSFERFEIFPKXSSWPXHXXXGVT

AXCXXGXGXSFYRNLXLXXXXXXYPXLXSYVNNKXKEVLVLWGXXHP

XXXXDQXXXXXXAYVXVSSXYXRFPEIAXRPKVRXQXGRXNY

WTLXEPGDTIXFEAXGNLXXPYAFAXXRGXGSGIIXSAXXXXCXKXQ

TPXGAINXSLFPQNXHPVTIGECPKYVXSTKLRMTGLRNIPSIQSRGLF

GAIAGFIEGGWTGMIDGWYGYHHQNEQSGYAADQKSTQNAIXGITNKVN

SVIEKMNTQFTAVGKEFNXLEXRXENLNKKVDDGFLDXWTYNAELLVLE

NERTLDFHDSNVKNLYEKVXSQKLNNAKEIGNGCFEFYHKCXXXCMESVK

NGTYDYPKYSEESKLNREXIDGVKLESXXXYQILAIYSTVASSLVLLVSL

GAISFWMCSNGSLQCRICI

17

In some embodiments disclosed herein, the HA polypeptides lack the N-terminal methionine residue. Thus, in some examples, provided are HA polypeptides comprising residues 2-566 of any one of SEQ ID NOs: 1-12.

The COBRA amino acid sequences can be reverse translated and optimized for expression in mammalian cells, including codon usage and RNA optimization (GeneArt; Regensburg, Germany). The optimized nucleic acid sequences can be inserted into an appropriate expression vector, such as the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross et al., *Nat. Immunol.* 1(2):102-103, 2000; Green et al., *Vaccine* 20:242-248, 2001).

The optimized nucleic acid sequence of the COBRA HA sequence of Method 1 is (SEQ ID NO: 13):

```

AAGCTTATGAAGGCCAAGCTGCTGGTGTCTGTGCGCCTTCACAGCCA
CCTACGCCGACACCATCTGCATCGGCTACCAAGCAACAACAGCACCGA
CACCGTGATACCGTGCTGGAAGAAAGCTGACCGTGACCCACAGCGTG
AACCTGTGGAAGATAGCCACAACGGCAAGCTGTGCCGGCTGAAGGGA
ATCGCCCTCTGCAGCTGGGCAACTGCTCTATCGCCGGCTGGATTCTGGG
CAACCCGAGTGCAGAGACCTGTTCTAGCAAGAGTCTGGTCTTACATC
GTGGAACCCCAACAGCGAGAACGGCACCTGTTACCCCGGCTACTTCG
CCGACTACGAGGAAGTGGCGGAACAGCTGAGCAGCGTGTCCAGCTTCGA
GAGATTGAGATTTTCCCCAAGAGAGCAGCTGGCCCAACACACCGTG
ACCAAGGCGTGACCGCTCTGCTCCCAATGGCAAGAGCAGCTTCT
ACAGAACTGTGTGGCTGACCGAGAAGAACGGCAGTACCCCAACCT
GAGCAAGAGCTACGTGAACAACAAAGAAAAAGAGGTGCTGGTGTGTG
GGGCGTGACCAACCTAGCAACATCGCGCAGCAGCGGCCATCTACCAC
ACCGAGAATGCCTACGTGTCCGTGGTGTCCAGCCACTACAGCAGCGGT
TCACCCCGAGATCGCCAAGAGGCCAAAGTGGCGGACAGGAAGGCC
GGATCAACTACTACTGGACACTGTGGAACCCGGCGATACCATCATCTTC
GAGGCCAACGGCAACCTGATCGCCCTTGGTACGCTTCGCCCTGAGCA
GAGGCTTTGGCAGCGGCATCATCACCAGCAACGCCAGCATGGACGAGTG
CGACGCCAAGTGCCAGACACCTCAGGGCGCCATCAATAGCAGCTGCC
TTCCAGAAGCTGCACCCCGTGACCATCGCGCAGTGCCCAAAATACGTGC
GGAGCACCAAGCTGCGGATGGTACCGGCCGTGAGAAACATCCCCAGCAT
CCAGAGCAGGGGCTGTTCGGAGCCATTGCCGGCTTTATCGAGGGCGGC
TGGACCGGCATGATCGACGGGTGGTACGGCTATCACCACGAGAACGAGC
AGGGCAGCGGCTACGCCCGCATCAGAAAGTCTACCCAGAACGCCATCAA
CGGCATCACCAACAAGTGAACAGCGTGATCGAGAAGATGAACACCCA
GTTTACCGCGCTGGGCAAGAGTTCAACAAGCTGGAACGGCGGATGGAA
AACCTGAACAAGAAGGTGGACGACGGCTTCTGGACATCTGGACCTACA
ACGCCGAAGTGTGGTGTCTGTGGAACGAGCGGACCTGGACTTCCA
CGACAGCAACGTGAAGAAGCTGTACGAGAAAGTGAAGTCCAGCTGAA
GAACAACGCCAAGAGATCGGCAACGGCTGCTTCGAGTTCTACCACAAG
TGCAACAACGAGTGCATGGAAGCGTGAAGAATGGCACCTACGACTACC

```

18

-continued

```

CCAAGTACAGCGAGGAAAGCAAGCTGAACCCGAGAAAGATCGACGGCG
TGAAGCTGGAATCCATGGGCGTGTACCAGATCCTGGCCATCTATAGCAC
CGTGCCAGCAGCCTGGTGTCTGTGGCGCCATCAGCTTTT
GGATGTGCAGCAACGGCAGCTGCAGTGCCGATCTGTATCGGCAGCAT
CGGATCC

```

V. Influenza

Influenza viruses are segmented negative-strand RNA viruses that belong to the Orthomyxoviridae family. There are three types of Influenza viruses, A, B and C. Influenza A viruses infect a wide variety of birds and mammals, including humans, horses, marine mammals, pigs, ferrets, and chickens. In animals, most influenza A viruses cause mild localized infections of the respiratory and intestinal tract. However, highly pathogenic influenza A strains, such as H5N1, cause systemic infections in poultry in which mortality may reach 100%. Animals infected with influenza A often act as a reservoir for the influenza viruses and certain subtypes have been shown to cross the species barrier to humans.

Influenza A viruses can be classified into subtypes based on allelic variations in antigenic regions of two genes that encode surface glycoproteins, namely, hemagglutinin (HA) and neuraminidase (NA) which are required for viral attachment and cellular release. Currently, sixteen subtypes of HA (H1-H16) and nine NA (N1-N9) antigenic variants are known for influenza A virus. Previously, only three subtypes were known to circulate in humans (H1N1, H1N2, and H3N2). However, in recent years, the pathogenic H5N1 subtype of avian influenza A has been reported to cross the species barrier and infect humans as documented in Hong Kong in 1997 and 2003, leading to the death of several patients.

In animals, most influenza A viruses cause mild localized infections of the respiratory and intestinal tract. However, highly pathogenic influenza A strains, such as H5N1, cause systemic infections in poultry in which mortality may reach 100%. In 2009, H1N1 influenza was the most common cause of human influenza. A new strain of swine-origin H1N1 emerged in 2009 and was declared pandemic by the World Health Organization. This strain was referred to as "swine flu." H1N1 influenza A viruses were also responsible for the Spanish flu pandemic in 1918, the Fort Dix outbreak in 1976, and the Russian flu epidemic in 1977-1978.

The influenza A virus genome encodes nine structural proteins and one nonstructural (NS1) protein with regulatory functions. The influenza virus segmented genome contains eight negative-sense RNA (nsRNA) gene segments (PB2, PB1, PA, NP, M, NS, HA and NA) that encode at least ten polypeptides, including RNA-directed RNA polymerase proteins (PB2, PB1 and PA), nucleoprotein (NP), neuraminidase (NA), hemagglutinin (subunits HA1 and HA2), the matrix proteins (M1 and M2) and the non-structural proteins (NS1 and NS2) (Krug et al., In "The Influenza Viruses," R. M. Krug, ed., Plenum Press, N.Y., 1989, pp. 89-152).

Influenza virus' ability to cause widespread disease is due to its ability to evade the immune system by undergoing antigenic change, which is believed to occur when a host is infected simultaneously with both an animal influenza virus and a human influenza virus. During mutation and reassortment in the host, the virus may incorporate an HA and/or NA

surface protein gene from another virus into its genome, thereby producing a new influenza subtype and evading the immune system.

HA is a viral surface glycoprotein generally comprising approximately 560 amino acids and representing 25% of the total virus protein. It is responsible for adhesion of the viral particle to, and its penetration into, a host cell in the early stages of infection. Cleavage of the virus HA0 precursor into the HA1 and HA2 sub-fragments is a necessary step in order for the virus to infect a cell. Thus, cleavage is required in order to convert new virus particles in a host cell into virions capable of infecting new cells. Cleavage is known to occur during transport of the integral HA0 membrane protein from the endoplasmic reticulum of the infected cell to the plasma membrane. In the course of transport, hemagglutinin undergoes a series of co- and post-translational modifications including proteolytic cleavage of the precursor HA into the amino-terminal fragment HA1 and the carboxy terminal HA2. One of the primary difficulties in growing influenza strains in primary tissue culture or established cell lines arises from the requirement for proteolytic cleavage activation of the influenza hemagglutinin in the host cell.

Although it is known that an uncleaved HA can mediate attachment of the virus to its neuraminic acid-containing receptors on a cell surface, it is not capable of the next step in the infectious cycle, which is fusion. It has been reported that exposure of the hydrophobic amino terminus of HA2 by cleavage is required so that it can be inserted into the target cell, thereby forming a bridge between virus and target cell membrane. This process is followed by fusion of the two membranes and entry of the virus into the target cell.

Proteolytic activation of HA involves cleavage at an arginine residue by a trypsin-like endoprotease, which is often an intracellular enzyme that is calcium dependent and has a neutral pH optimum. Since the activating proteases are cellular enzymes, the infected cell type determines whether the HA is cleaved. The HA of the mammalian influenza viruses and the nonpathogenic avian influenza viruses are susceptible to proteolytic cleavage only in a restricted number of cell types. On the other hand, HA of pathogenic avian viruses among the H5 and H7 subtypes are cleaved by proteases present in a broad range of different host cells. Thus, there are differences in host range resulting from differences in hemagglutinin cleavability which are correlated with the pathogenic properties of the virus.

Neuraminidase (NA) is a second membrane glycoprotein of the influenza viruses. The presence of viral NA has been shown to be important for generating a multi-faceted protective immune response against an infecting virus. For most influenza A viruses, NA is 413 amino acid in length, and is encoded by a gene of 1413 nucleotides. Nine different NA subtypes have been identified in influenza viruses (N1, N2, N3, N4, N5, N6, N7, N8 and N9), all of which have been found among wild birds. NA is involved in the destruction of the cellular receptor for the viral HA by cleaving terminal neuraminic acid (also called sialic acid) residues from carbohydrate moieties on the surfaces of infected cells. NA also cleaves sialic acid residues from viral proteins, preventing aggregation of viruses. Using this mechanism, it is hypothesized that NA facilitates release of viral progeny by preventing newly formed viral particles from accumulating along the cell membrane, as well as by promoting transportation of the virus through the mucus present on the mucosal surface. NA is an important antigenic determinant that is subject to antigenic variation.

In addition to the surface proteins HA and NA, influenza virus comprises six additional internal genes, which give

rise to eight different proteins, including polymerase genes PB1, PB2 and PA, matrix proteins M1 and M2, nucleoprotein (NP), and non-structural proteins NS1 and NS2 (Horimoto et al., *Clin Microbiol Rev.* 14(1):129-149, 2001).

In order to be packaged into progeny virions, viral RNA is transported from the nucleus as a ribonucleoprotein (RNP) complex composed of the three influenza virus polymerase proteins, the nucleoprotein (NP), and the viral RNA, in association with the influenza virus matrix 1 (M1) protein and nuclear export protein (Marsh et al., *J Virol*, 82:2295-2304, 2008). The M1 protein that lies within the envelope is thought to function in assembly and budding. A limited number of M2 proteins are integrated into the virions (Zebedee, *J. Virol.* 62:2762-2772, 1988). They form tetramers having H⁺ ion channel activity, and when activated by the low pH in endosomes, acidify the inside of the virion, facilitating its uncoating (Pinto et al., *Cell* 69:517-528, 1992). Amantadine is an anti-influenza drug that prevents viral infection by interfering with M2 ion channel activity, thus inhibiting virus uncoating.

NS1, a nonstructural protein, has multiple functions, including regulation of splicing and nuclear export of cellular mRNAs as well as stimulation of translation. The major function of NS1 seems to be to counteract the interferon activity of the host, since an NS1 knockout virus was viable although it grew less efficiently than the parent virus in interferon-nondefective cells (Garcia-Sastre, *Virology* 252: 324-330, 1998).

NS2 has been detected in virus particles (Richardson et al., *Arch. Virol.* 116:69-80, 1991; Yasuda et al., *Virology* 196:249-255, 1993). The average number of NS2 proteins in a virus particle was estimated to be 130-200 molecules. An in vitro binding assay shows direct protein-protein contact between M1 and NS2. NS2-M1 complexes have also been detected by immunoprecipitation in virus-infected cell lysates. The NS2 protein is thought to play a role in the export of RNP from the nucleus through interaction with M1 protein (Ward et al., *Arch. Virol.* 140:2067-2073, 1995).

VI. Influenza VLPs and Administration Thereof

Influenza VLPs comprising an optimized HA (such as the HA having the amino acid sequence set forth as any one of SEQ ID NOs: 1-11) are provided herein. The influenza VLPs are generally made up of the HA, NA and M1 proteins. The production of influenza VLPs has been described in the art and is within the abilities of one of ordinary skill in the art. For example, influenza VLPs can be produced by transfection of host cells with plasmids encoding the HA, NA and M1 proteins. After incubation of the transfected cells for an appropriate time to allow for protein expression (such as for approximately 72 hours), VLPs can be isolated from cell culture supernatants. Example 2 below provides an exemplary protocol for purifying influenza VLPs from cell supernatants. In this example, VLPs are isolated by low speed centrifugation (to remove cell debris), vacuum filtration and ultracentrifugation through 20% glycerol.

The influenza VLPs disclosed herein can be used as influenza vaccines to elicit a protective immune response against H1N1 influenza viruses.

Influenza VLPs, or compositions thereof, can be administered to a subject by any of the routes normally used for introducing recombinant virus into a subject. Methods of administration include, but are not limited to, intradermal, intramuscular, intraperitoneal, parenteral, intravenous, subcutaneous, vaginal, rectal, intranasal, inhalation or oral. Parenteral administration, such as subcutaneous, intrave-

nous or intramuscular administration, is generally achieved by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. Administration can be systemic or local.

Influenza VLPs, or compositions thereof, are administered in any suitable manner, such as with pharmaceutically acceptable carriers. Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions of the present disclosure.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

Some of the compositions may potentially be administered as a pharmaceutically acceptable acid- or base-addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

Administration can be accomplished by single or multiple doses. The dose administered to a subject in the context of the present disclosure should be sufficient to induce a beneficial therapeutic response in a subject over time, or to inhibit or prevent H1N1 influenza virus infection. The dose required will vary from subject to subject depending on the species, age, weight and general condition of the subject, the severity of the infection being treated, the particular composition being used and its mode of administration. An appropriate dose can be determined by one of ordinary skill in the art using only routine experimentation.

Provided herein are pharmaceutical compositions which include a therapeutically effective amount of the influenza VLPs alone or in combination with a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The carrier and composition can be sterile, and the formulation suits the mode of administration. The composition can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The composition can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The composition can be formulated as a suppository, with traditional binders

and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, and magnesium carbonate. Any of the common pharmaceutical carriers, such as sterile saline solution or sesame oil, can be used. The medium can also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. Other media that can be used with the compositions and methods provided herein are normal saline and sesame oil.

The influenza VLPs described herein can be administered alone or in combination with other therapeutic agents to enhance antigenicity. For example, the influenza VLPs can be administered with an adjuvant, such as Freund incomplete adjuvant or Freund's complete adjuvant.

Optionally, one or more cytokines, such as IL-2, IL-6, IL-12, RANTES, GM-CSF, TNF- α , or IFN- γ , one or more growth factors, such as GM-CSF or G-CSF; one or more molecules such as OX-40L or 41 BBL, or combinations of these molecules, can be used as biological adjuvants (see, for example, Salgaller et al., 1998, *J. Surg. Oncol.* 68(2): 122-38; Lotze et al., 2000, *Cancer J. Sci. Am.* 6(Suppl 1):S61-6; Cao et al., 1998, *Stem Cells* 16(Suppl 1):251-60; Kuiper et al., 2000, *Adv. Exp. Med. Biol.* 465:381-90). These molecules can be administered systemically (or locally) to the host.

A number of means for inducing cellular responses, both in vitro and in vivo, are known. Lipids have been identified as agents capable of assisting in priming CTL in vivo against various antigens. For example, as described in U.S. Pat. No. 5,662,907, palmitic acid residues can be attached to the alpha and epsilon amino groups of a lysine residue and then linked (for example, via one or more linking residues, such as glycine, glycine-glycine, serine, serine-serine, or the like) to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated in a liposome, or emulsified in an adjuvant. As another example, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine can be used to prime tumor specific CTL when covalently attached to an appropriate peptide (see, Deres et al., *Nature* 342:561, 1989). Further, as the induction of neutralizing antibodies can also be primed with the same molecule conjugated to a peptide which displays an appropriate epitope, two compositions can be combined to elicit both humoral and cell-mediated responses where that is deemed desirable.

Although administration of VLPs containing an optimized HA protein is exemplified herein, one of skill in the art would understand that it is also possible to administer the optimized influenza HA protein itself (in the absence of a viral particle) or as a fusion protein to elicit an immune response in a subject.

The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the disclosure to the particular features or embodiments described.

EXAMPLES

Example 1

Generation of COBRA Sequences for H1N1 Influenza

Influenza A H1N1 HA amino acid sequences were downloaded from the NCBI Influenza Virus Resource database.

H1N1 HA proteins from 1134 isolates from 1918-2011 were used for generating consensus sequences. Eleven different consensus sequences (SEQ ID NOs: 1-11) were generated using the following methods:

1. COBRA Method 1 (1918-2005)

Sequences (456) were organized by date of isolation and eight primary consensus sequences were generated using isolates from 1918-1934 (8), 1935-1947 (13), 1948-1957 (12), 1977-1983 (68), 1984-1986 (9), 1987-1991 (12), 1992-1999 (59) and 2000-2005 (263). Four secondary consensus sequences were generated by grouping the primary consensus sequences by date, as shown in FIG. 1. The final consensus sequence (the third layer consensus; SEQ ID NO: 1) was generated by alignment of the four secondary consensus sequences.

2. COBRA Method 2 (1918-2005)

As with method 1, sequences (456) were organized by date of isolation to generate eight primary consensus sequences. The final consensus sequence (SEQ ID NO: 2) was generated by aligning the eight primary consensus sequences, as illustrated in FIG. 2.

3. COBRA Method 3 (1918-2005)

The consensus sequence (SEQ ID NO: 3) was generated by alignment of 456 H1N1 virus isolates from 1918-2005.

4. COBRA Method 4 (1918-2005)

Sequences (456) were organized by date of isolation and ten primary consensus sequences were generated using isolates from 1918-1933 (6), 1934-1946 (15), 1947-1956 (12), 1957-1977 (8), 1978-1980 (17), 1981-1985 (49), 1986-1990 (14), 1991-1995 (27), 1996-1998 (27) and 1999-2005 (281), as shown in FIG. 3. The final consensus sequence (SEQ ID NO: 4) was generated by alignment of the ten primary consensus sequences.

5. COBRA Method 5 (1918-2011)

Sequences (1134) were organized by date of isolation and 12 primary consensus sequences were generated using isolates from 1918 (1), 1976 (4), 2009-2011 (123), 1933-1934 (8), 1935-1947 (13), 1948-1957 (12), 1977-1983 (68), 1984-1986 (9), 1987-1991 (12), 1992-1999 (27), 2000-2005 (59) and 2006-2008 (798). Four secondary consensus sequences were generated by grouping the primary consensus sequences according to "swine" sequences or by date, as shown in FIG. 4. The final consensus sequence (the third layer consensus; SEQ ID NO: 5) was generated by alignment of the four secondary consensus sequences.

6. COBRA Method 6

Sequences (1134) were organized by date of isolation and 13 primary consensus sequences were generated using isolates from 1918 (1), 1933-1934 (8), 1935-1947 (13), 1948-1957 (12), 1976 (4), 1977-1983 (68), 1984-1986 (9), 1987-1991 (12), 1992-1999 (27), 2000-2005 (59), 2006 (76), 2007-2008 (722) and 2009-2011 (123), as shown in FIG. 5. The final consensus sequence (SEQ ID NO: 6) was generated by alignment of the 13 primary consensus sequences.

7. COBRA Method 5 Deglycosylated

This sequence (SEQ ID NO: 7) was generated by altering the Method 5 consensus sequence (SEQ ID NO: 5) to remove predicted glycosylation sites.

8. 1918-1957 COBRA

This sequence (SEQ ID NO: 8) was generated by alignment of H1N1 isolates from 1918-1957.

9. 1977-2005 COBRA

This sequence (SEQ ID NO: 9) was generated by alignment of H1N1 isolates from 1977-2005.

10. Human Swine COBRA

This sequence (SEQ ID NO: 10) was generated by alignment of H1N1 human swine influenza isolates. This sequence was aligned using consensus sequences from 1918, 1976 and 2009-2011.

11. Swine COBRA

This sequence (SEQ ID NO: 11) was generated by alignment of H1N1 swine influenza isolates. This sequence was aligned using swine sequences from 1930-2010.

The COBRA amino acid sequence generated according to Method 1 was reverse translated and optimized for expression in mammalian cells, including codon usage and RNA optimization (GeneArt; Regensburg, Germany). The optimized nucleic acid sequence is set forth herein as SEQ ID NO: 13. The remaining COBRA sequences will also be reverse translated and optimized for expression in mammalian cells. The optimized nucleic acid sequences will be inserted into the pTR600 expression vector (U.S. Patent Application Publication No. 2002/0106798; Ross et al., *Nat. Immunol.* 1(2):102-103, 2000; Green et al., *Vaccine* 20:242-248, 2001).

Example 2

Preparation of and Immunization with Influenza VLPs

The following methods can be used to produce and characterize influenza VLPs comprising an optimized HA. Exemplary methods for immunization of mice, ferrets and macaques are also described below (see also, Giles and Ross, *Vaccine* 29(16):3043-3054, 2011).

Vaccine Preparation

293T cells are transiently transfected with plasmids expressing M1, NA and an optimized HA, and incubated for 72 hours at 37° C. The M1, NA and HA coding sequences can be codon-optimized for expression in mammalian cells. Supernatants are collected and cell debris is removed by low speed centrifugation followed by vacuum filtration through a 0.22 µm sterile filter. VLPs are purified via ultracentrifugation (100,000xg through 20% glycerol, weight per volume) for 4 hours at 4° C. The pellets are subsequently resuspended in PBS pH 7.2 and stored in single use aliquots at -80° C. until use. Total protein concentration is determined by Micro BCA™ Protein Assay Reagent Kit (Pierce Biotechnology, Rockford, Ill., USA).

Dose Determination

HA specific content can be determined by western blot and densitometry. Purified recombinant COBRA HA and purified VLPs are prepared in standard total protein amounts and are electrophoresed on a 10% SDS-PAGE gel and transferred to a PVDF membrane. The blot is probed with mouse polyclonal antisera from influenza infected mice and the HA-antibody complexes are detected using a goat anti-mouse IgG conjugated to horseradish peroxidase (HRP) (Southern Biotech; Birmingham, Ala., USA). HRP is detected by chemiluminescent substrate (Pierce Biotechnology; Rockford Ill., USA) and exposed to X-ray film (ThermoFisher; Pittsburgh, Pa., USA). Density of bands is determined using ImageJ software (NIH). Density of recombinant HA bands is used to calculate a standard curve and the density of the purified VLPs is interpolated using the results from the recombinant HA.

Mouse Studies

BALB/c mice (*Mus musculus*, females, 6-8 weeks old) can be purchased from Harlan Sprague Dawley (Indianapolis, Ind., USA). Mice are housed in microisolator units and allowed free access to food and water and are cared for

under USDA guidelines for laboratory animals. Mice are vaccinated with one of three doses of purified COBRA HA VLPs (1.5 µg, 0.3 µg or 0.06 µg), based upon HA content from a densitometry assay, via intramuscular injection at week 0 and then boosted with the same dose at week 3. Vaccines at each dose are formulated with alum adjuvant (Imject Alum, Pierce Biotechnology; Rockford, Ill., USA), CpG oligonucleotides, or vehicle alone. Fourteen to twenty-one days after each vaccination, blood is collected from anesthetized mice via the retro-orbital plexus and transferred to a microfuge tube. Tubes are centrifuged and sera is removed and frozen at $-80\pm5^{\circ}\text{C}$. Hemagglutination inhibition (HAI) serum antibody titer for each vaccine group is determined at week 5 using representative reassortant viruses or COBRA HA VLPs.

Three weeks after final vaccination, mice are challenged intranasally with a highly pathogenic H1N1 virus in a volume of 50 µl. After infection, mice are monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores (Toapanta and Ross, *Respiratory Research* 10(1):112, 2009) and death are recorded for each group on each day after inoculation.

Ferret Studies

Fitch ferrets (*Mustela putorius furo*, female, 6-12-months of age), influenza naïve and de-scented, can be purchased from Marshall Farms (Sayre, Pa., USA). Ferrets are pair housed in stainless steel cages (Shor-line, Kansas City, Kans., USA) containing Sani-chips Laboratory Animal Bedding (P. J. Murphy Forest Products, Montville, N.J., USA). Ferrets are provided with Teklad Global Ferret Diet (Harlan Teklad, Madison, Wis., USA) and fresh water ad libitum. The COBRA HA VLPs are diluted in PBS, pH 7.2 to achieve final concentration. Ferrets are vaccinated with one of two doses of purified COBRA VLPs (15 µg, 3 µg), based upon HA content as determined by densitometry assay, via intramuscular injection in the quadriceps muscle in a volume of 0.25 ml at week 0 and then boosted with the same dose at week 3. Vaccines are stored at -80°C prior to use and formulated with alum adjuvant (Imject Alum; Pierce Biotechnology, Rockford, Ill., USA) immediately prior to use. Animals are monitored for adverse events including weight loss, temperature, decrease in activity, nasal discharge, sneezing and diarrhea weekly during the vaccination regimen. Prior to vaccination, animals are confirmed by HAI assay to be seronegative for circulating influenza A and influenza B viruses. Fourteen to twenty-one days after each vaccination, blood is collected from anesthetized ferrets via the anterior vena cava and transferred to a microfuge tube. Tubes are centrifuged and sera is removed and frozen at $-80\pm5^{\circ}\text{C}$. HAI serum antibody titer for each vaccine group is determined at week 5 using representative reassortant viruses or COBRA HA VLPs.

Three weeks after final vaccination, ferrets are challenged intranasally with a highly pathogenic H1N1 virus in a volume of 1 ml. After infection, ferrets are monitored daily for weight loss, disease signs and death for 14 days after infection. Individual body weights, sickness scores, and death are recorded for each group on each day after inoculation. Nasal washes are performed by instilling 3 ml of PBS into the nares of anesthetized ferrets each day for 7 days after inoculation. Washes are collected and stored at -80°C until use.

Primate Immunizations

Cynomolgus macaques (*Macaca fascicularis*, male, 3-5 years old) can be purchased from Harlan Sprague Dawley (Indianapolis, Ind., USA). Macaques are vaccinated with purified COBRA HA VLPs (15 µg), based upon HA content from a densitometry assay, via intramuscular injection at week 0 and then boosted with the same dose at weeks 3 and 6. Vaccines are formulated with alum adjuvant (Imject Alum, Pierce Biotechnology; Rockford, Ill., USA) immediately prior to use. Twenty-one days after each vaccination, blood is collected from anesthetized macaques via the femoral vein and transferred to a serum separator tube. Tubes are allowed to activate clotting followed by centrifugation and sera is removed and frozen at $-80\pm5^{\circ}\text{C}$. End point IgG titers and HAI serum antibody titer for each vaccine group is determined at week 5 using representative reassortant viruses or COBRA HA VLPs.

Three weeks after final vaccination, macaques are challenged by intranasal, intratracheal, and orbital inoculation with a highly pathogenic H1N1 virus in a volume of 1 ml. After infection, macaques are monitored daily for weight loss, disease signs and death for 5 days after infection. Individual body weights, sickness scores and death are recorded for each group on each day after inoculation.

Example 3

Analysis of COBRA H1N1 Method 1 HA

This example describes the finding that inoculation of mice with COBRA H1N1 influenza VLPs induces significant HAI serum antibody titers, and further shows that translation of COBRA H1N1 HA in vitro results in expression of the protein at the expected molecular weight.

HAI Serum Antibody Titers

To evaluate whether influenza virus containing COBRA Method 1 H1N1 HA is capable of eliciting an antibody response in infected animals, VLPs comprising the COBRA Method 1 HA (SEQ ID NO: 1) were generated as described in Example 2. Mice were inoculated with the VLPs and serum was collected at weeks 3, 5, 8 and 12 post-infection for determination of HAI serum antibody titers. Antisera were tested against the seasonal H1N1 influenza virus, A/New Caledonia/20/1999. As shown in FIG. 6, HAI titers were detected starting at 5 weeks post-infection and increased at weeks 8 and 12.

Expression of COBRA HA In Vitro

As described in Example 2, the COBRA HA Method 1 amino acid sequence was reverse translated and optimized for expression in mammalian cells. The optimized nucleic acid sequence (SEQ ID NO: 13) encoding COBRA HA was inserted into the pTR600 expression vector. To assess whether COBRA H1N1 HA is properly expressed in vitro, cells were transfected with the pTR600 expression vector encoding COBRA H1N1 HA. Cell culture lysates were analyzed by SDS-PAGE, followed by Western blot for detection of influenza HA. As shown in FIG. 7, COBRA HA migrated to its expected molecular weight, confirming expression of the synthetic protein in vitro.

In view of the many possible embodiments to which the principles of the disclosed invention may be applied, it should be recognized that the illustrated embodiments are only preferred examples of the invention and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 566

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 1

```

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Phe Thr Ala Thr Tyr
 1          5          10          15

Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20          25          30

Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35          40          45

Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
 50          55          60

Ala Pro Leu Gln Leu Gly Asn Cys Ser Ile Ala Gly Trp Ile Leu Gly
 65          70          75          80

Asn Pro Glu Cys Glu Ser Leu Phe Ser Lys Glu Ser Trp Ser Tyr Ile
 85          90          95

Val Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Tyr Phe
100          105          110

Ala Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115          120          125

Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Thr
130          135          140

Val Thr Lys Gly Val Thr Ala Ser Cys Ser His Asn Gly Lys Ser Ser
145          150          155          160

Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Ser Tyr Pro
165          170          175

Asn Leu Ser Lys Ser Tyr Val Asn Asn Lys Glu Lys Glu Val Leu Val
180          185          190

Leu Trp Gly Val His His Pro Ser Asn Ile Gly Asp Gln Arg Ala Ile
195          200          205

Tyr His Thr Glu Asn Ala Tyr Val Ser Val Val Ser Ser His Tyr Ser
210          215          220

Arg Arg Phe Thr Pro Glu Ile Ala Lys Arg Pro Lys Val Arg Asp Gln
225          230          235          240

Glu Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
245          250          255

Ile Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe
260          265          270

Ala Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Ser
275          280          285

Met Asp Glu Cys Asp Ala Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn
290          295          300

Ser Ser Leu Pro Phe Gln Asn Val His Pro Val Thr Ile Gly Glu Cys
305          310          315          320

Pro Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg
325          330          335

Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
340          345          350

Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr

```

-continued

355					360					365					
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser
370						375					380				
Thr	Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile
385					390					395					400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys
				405					410					415	
Leu	Glu	Arg	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe
			420					425					430		
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
	435						440					445			
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu
	450					455					460				
Lys	Val	Lys	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly
465				470						475					480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asn	Asn	Glu	Cys	Met	Glu	Ser	Val
				485					490					495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu
		500						505					510		
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr
		515					520					525			
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu
	530					535						540			
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu
545					550					555					560
Gln	Cys	Arg	Ile	Cys	Ile										
				565											

<210> SEQ ID NO 2

<211> LENGTH: 566

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 2

Met	Lys	Ala	Lys	Leu	Leu	Val	Leu	Leu	Cys	Ala	Leu	Thr	Ala	Thr	Asp
1				5						10				15	
Ala	Asp	Thr	Ile	Cys	Ile	Gly	Tyr	His	Ala	Asn	Asn	Ser	Thr	Asp	Thr
			20					25					30		
Val	Asp	Thr	Val	Leu	Glu	Lys	Asn	Val	Thr	Val	Thr	His	Ser	Val	Asn
		35					40					45			
Leu	Leu	Glu	Asp	Ser	His	Asn	Gly	Lys	Leu	Cys	Arg	Leu	Lys	Gly	Ile
	50					55					60				
Ala	Pro	Leu	Gln	Leu	Gly	Asn	Cys	Ser	Ile	Ala	Gly	Trp	Ile	Leu	Gly
65					70					75				80	
Asn	Pro	Glu	Cys	Glu	Ser	Leu	Phe	Ser	Lys	Lys	Ser	Trp	Ser	Tyr	Ile
			85					90					95		
Ala	Glu	Thr	Pro	Asn	Ser	Glu	Asn	Gly	Thr	Cys	Tyr	Pro	Gly	Tyr	Phe
		100						105					110		
Ala	Asp	Tyr	Glu	Glu	Leu	Arg	Glu	Gln	Leu	Ser	Ser	Val	Ser	Ser	Phe
		115					120					125			
Glu	Arg	Phe	Glu	Ile	Phe	Pro	Lys	Glu	Ser	Ser	Trp	Pro	Asn	His	Thr
	130					135					140				
Val	Thr	Lys	Gly	Val	Thr	Ala	Ala	Cys	Ser	His	Lys	Gly	Lys	Ser	Ser

145	150								155								160	
Phe	Tyr	Arg	Asn	Leu	Leu	Trp	Leu	Thr	Glu	Lys	Asn	Gly	Ser	Tyr	Pro			
				165					170					175				
Asn	Leu	Ser	Lys	Ser	Tyr	Val	Asn	Asn	Lys	Glu	Lys	Glu	Val	Leu	Val			
				180					185					190				
Leu	Trp	Gly	Val	His	His	Pro	Ser	Asn	Ile	Gly	Asp	Gln	Arg	Ala	Ile			
				195					200					205				
Tyr	His	Thr	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	His	Tyr	Asn			
				210					215					220				
Arg	Arg	Phe	Thr	Pro	Glu	Ile	Ala	Lys	Arg	Pro	Lys	Val	Arg	Asp	Gln			
				225					230					235				
Glu	Gly	Arg	Ile	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr			
				245					250					255				
Ile	Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe			
				260					265					270				
Ala	Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Ser			
				275					280					285				
Met	Asp	Glu	Cys	Asp	Ala	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn			
				290					295					300				
Ser	Ser	Leu	Pro	Phe	Gln	Asn	Val	His	Pro	Val	Thr	Ile	Gly	Glu	Cys			
				305					310					315				
Pro	Lys	Tyr	Val	Arg	Ser	Thr	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg			
				325					330					335				
Asn	Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly			
				340					345					350				
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr			
				355					360					365				
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser			
				370					375					380				
Thr	Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile			
				385					390					395				
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys			
				405					410					415				
Leu	Glu	Lys	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe			
				420					425					430				
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn			
				435					440					445				
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu			
				450					455					460				
Lys	Val	Lys	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly			
				465					470					475				
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asn	Asn	Glu	Cys	Met	Glu	Ser	Val			
				485					490					495				
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu			
				500					505					510				
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr			
				515					520					525				
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu			
				530					535					540				
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu			
				545					550					555				
Gln	Cys	Arg	Ile	Cys	Ile													

-continued

```

<210> SEQ ID NO 3
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 3

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Phe Thr Ala Thr Tyr
1      5      10      15
Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20      25      30
Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35      40      45
Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
50      55      60
Ala Pro Leu Gln Leu Gly Asn Cys Ser Val Ala Gly Trp Ile Leu Gly
65      70      75      80
Asn Pro Glu Cys Glu Ser Leu Ile Ser Lys Glu Ser Trp Ser Tyr Ile
85      90      95
Val Glu Thr Pro Asn Pro Glu Asn Gly Thr Cys Tyr Pro Gly Tyr Phe
100     105     110
Ala Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115     120     125
Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Thr
130     135     140
Val Thr Lys Gly Val Thr Ala Ser Cys Ser His Asn Gly Lys Ser Ser
145     150     155     160
Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Leu Tyr Pro
165     170     175
Asn Leu Ser Lys Ser Tyr Val Asn Asn Lys Glu Lys Glu Val Leu Val
180     185     190
Leu Trp Gly Val His His Pro Ser Asn Ile Gly Asp Gln Arg Ala Leu
195     200     205
Tyr His Thr Glu Asn Ala Tyr Val Ser Val Val Ser Ser His Tyr Ser
210     215     220
Arg Arg Phe Thr Pro Glu Ile Ala Lys Arg Pro Lys Val Arg Asp Gln
225     230     235     240
Glu Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
245     250     255
Ile Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe
260     265     270
Ala Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Ser
275     280     285
Met Asp Glu Cys Asp Ala Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn
290     295     300
Ser Ser Leu Pro Phe Gln Asn Val His Pro Val Thr Ile Gly Glu Cys
305     310     315     320
Pro Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg
325     330     335
Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
340     345     350
Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
355     360     365

```

-continued

His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser
 370 375 380
 Thr Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile
 385 390 395 400
 Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys
 405 410 415
 Leu Glu Arg Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe
 420 425 430
 Leu Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn
 435 440 445
 Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu
 450 455 460
 Lys Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly
 465 470 475 480
 Cys Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val
 485 490 495
 Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 4
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 4

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Leu Thr Ala Thr Asp
 1 5 10 15
 Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20 25 30
 Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35 40 45
 Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
 50 55 60
 Ala Pro Leu Gln Leu Gly Lys Cys Ser Ile Ala Gly Trp Ile Leu Gly
 65 70 75 80
 Asn Pro Glu Cys Glu Ser Leu Leu Ser Lys Lys Ser Trp Ser Tyr Ile
 85 90 95
 Ala Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Tyr Phe
 100 105 110
 Ala Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115 120 125
 Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Asn
 130 135 140
 Val Thr Lys Gly Val Thr Ala Ser Cys Ser His Lys Gly Lys Ser Ser
 145 150 155 160

Phe	Tyr	Arg	Asn	Leu	Leu	Trp	Leu	Thr	Glu	Lys	Asn	Gly	Ser	Tyr	Pro	
				165					170						175	Pro
Asn	Leu	Ser	Lys	Ser	Tyr	Val	Asn	Asn	Lys	Glu	Lys	Glu	Val	Leu	Val	
				180					185						190	
Leu	Trp	Gly	Val	His	His	Pro	Ser	Asn	Ile	Gly	Asp	Gln	Arg	Thr	Ile	
				195					200						205	
Tyr	Arg	Thr	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	Asn	Tyr	Asn	
				210					215						220	
Arg	Arg	Phe	Thr	Pro	Glu	Ile	Ala	Lys	Arg	Pro	Lys	Val	Arg	Asp	Gln	
				225					230						235	240
Glu	Gly	Arg	Ile	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr	
				245					250						255	
Ile	Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe	
				260					265						270	
Ala	Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Ser	
				275					280						285	
Met	Asp	Glu	Cys	Asp	Thr	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn	
				290					295						300	
Ser	Ser	Leu	Pro	Phe	Gln	Asn	Val	His	Pro	Val	Thr	Ile	Gly	Glu	Cys	
				305					310						315	320
Pro	Lys	Tyr	Val	Arg	Ser	Thr	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg	
				325					330						335	
Asn	Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	
				340					345						350	
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr	
				355					360						365	
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser	
				370					375						380	
Thr	Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile	
				385					390						395	400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys	
				405					410						415	
Leu	Glu	Lys	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe	
				420					425						430	
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn	
				435					440						445	
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu	
				450					455						460	
Lys	Val	Lys	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly	
				465					470						475	480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asn	Asn	Glu	Cys	Met	Glu	Ser	Val	
				485					490						495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu	
				500					505						510	
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr	
				515					520						525	
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu	
				530					535						540	
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu	
				545					550						555	560
Gln	Cys	Arg	Ile	Cys	Ile											
				565												

-continued

```

<210> SEQ ID NO 5
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 5

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Thr Phe Thr Ala Thr Asp
 1             5             10             15

Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
      20             25             30

Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
      35             40             45

Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
      50             55             60

Ala Pro Leu Gln Leu Gly Asn Cys Ser Ile Ala Gly Trp Ile Leu Gly
      65             70             75             80

Asn Pro Glu Cys Glu Ser Leu Leu Ser Lys Lys Ser Trp Ser Tyr Ile
      85             90             95

Val Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Asp Phe
      100            105            110

Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
      115            120            125

Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Thr
      130            135            140

Val Thr Lys Gly Val Thr Ala Ala Cys Ser His Ala Gly Lys Ser Ser
      145            150            155            160

Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Ser Tyr Pro
      165            170            175

Asn Leu Ser Lys Ser Tyr Val Asn Asn Lys Gly Lys Glu Val Leu Val
      180            185            190

Leu Trp Gly Val His His Pro Ser Asn Ile Gly Asp Gln Gln Ala Leu
      195            200            205

Tyr Gln Thr Glu Asn Ala Tyr Val Ser Val Val Ser Ser His Tyr Asn
      210            215            220

Arg Lys Phe Thr Pro Glu Ile Ala Lys Arg Pro Lys Val Arg Asp Gln
      225            230            235            240

Glu Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
      245            250            255

Ile Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe
      260            265            270

Ala Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Ser
      275            280            285

Met His Glu Cys Asp Thr Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn
      290            295            300

Ser Ser Leu Pro Phe Gln Asn Ile His Pro Val Thr Ile Gly Glu Cys
      305            310            315            320

Pro Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg
      325            330            335

Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
      340            345            350

Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
      355            360            365

```

-continued

His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser
 370 375 380
 Thr Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile
 385 390 395 400
 Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys
 405 410 415
 Leu Glu Lys Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe
 420 425 430
 Leu Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn
 435 440 445
 Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu
 450 455 460
 Lys Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly
 465 470 475 480
 Cys Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val
 485 490 495
 Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 6
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 6

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Phe Thr Ala Thr Asp
 1 5 10 15
 Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20 25 30
 Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35 40 45
 Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Lys Leu Lys Gly Ile
 50 55 60
 Ala Pro Leu Gln Leu Gly Lys Cys Ser Ile Ala Gly Trp Ile Leu Gly
 65 70 75 80
 Asn Pro Glu Cys Glu Ser Leu Leu Ser Lys Lys Ser Trp Ser Tyr Ile
 85 90 95
 Val Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Asp Phe
 100 105 110
 Ala Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115 120 125
 Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Thr
 130 135 140
 Val Thr Lys Gly Val Thr Ala Ala Cys Ser His Ala Gly Lys Ser Ser
 145 150 155 160

Phe	Tyr	Arg	Asn	Leu	Leu	Trp	Leu	Thr	Glu	Lys	Asn	Gly	Ser	Tyr	Pro	
				165					170					175		
Asn	Leu	Ser	Lys	Ser	Tyr	Val	Asn	Asn	Lys	Glu	Lys	Glu	Val	Leu	Val	
				180					185					190		
Leu	Trp	Gly	Val	His	His	Pro	Ser	Asn	Ile	Gly	Asp	Gln	Arg	Ala	Leu	
				195					200					205		
Tyr	His	Thr	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	His	Tyr	Asn	
				210					215					220		
Arg	Arg	Phe	Thr	Pro	Glu	Ile	Ala	Lys	Arg	Pro	Lys	Val	Arg	Asp	Gln	
				225					230					235		
Glu	Gly	Arg	Ile	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr	
				245					250					255		
Ile	Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe	
				260					265					270		
Ala	Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Ser	
				275					280					285		
Met	Asp	Glu	Cys	Asp	Ala	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn	
				290					295					300		
Ser	Ser	Leu	Pro	Phe	Gln	Asn	Val	His	Pro	Val	Thr	Ile	Gly	Glu	Cys	
				305					310					315		
Pro	Lys	Tyr	Val	Arg	Ser	Thr	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg	
				325					330					335		
Asn	Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly	
				340					345					350		
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr	
				355					360					365		
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser	
				370					375					380		
Thr	Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile	
				385					390					395		
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Lys	
				405					410					415		
Leu	Glu	Arg	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe	
				420					425					430		
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn	
				435					440					445		
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu	
				450					455					460		
Lys	Val	Lys	Ser	Gln	Leu	Lys	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly	
				465					470					475		
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asn	Asn	Glu	Cys	Met	Glu	Ser	Val	
				485					490					495		
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu	
				500					505					510		
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr	
				515					520					525		
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu	
				530					535					540		
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu	
				545					550					555		
Gln	Cys	Arg	Ile	Cys	Ile											
				565												

-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 7

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Thr Phe Thr Ala Thr Asp
 1             5             10             15

Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20             25             30

Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35             40             45

Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
 50             55             60

Ala Pro Leu Gln Leu Gly Asn Cys Ala Ile Ala Gly Trp Ile Leu Gly
 65             70             75             80

Asn Pro Glu Cys Glu Ser Leu Leu Ser Lys Lys Ser Trp Ser Tyr Ile
 85             90             95

Val Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Asp Phe
 100            105            110

Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115            120            125

Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Ala
 130            135            140

Val Thr Lys Gly Val Thr Ala Ala Cys Ser His Ala Gly Lys Ser Ser
 145            150            155            160

Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Ser Tyr Pro
 165            170            175

Asn Leu Ala Lys Ser Tyr Val Asn Asn Lys Gly Lys Glu Val Leu Val
 180            185            190

Leu Trp Gly Val His His Pro Ser Asn Ile Gly Asp Gln Gln Ala Leu
 195            200            205

Tyr Gln Thr Glu Asn Ala Tyr Val Ser Val Val Ser Ser His Tyr Asn
 210            215            220

Arg Lys Phe Thr Pro Glu Ile Ala Lys Arg Pro Lys Val Arg Asp Gln
 225            230            235            240

Glu Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
 245            250            255

Ile Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe
 260            265            270

Ala Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Ala
 275            280            285

Met His Glu Cys Asp Thr Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn
 290            295            300

Ser Ser Leu Pro Phe Gln Asn Ile His Pro Val Thr Ile Gly Glu Cys
 305            310            315            320

Pro Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg
 325            330            335

Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
 340            345            350

Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
 355            360            365

His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser

```

-continued

370	375	380
Thr Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile		
385	390	395 400
Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys		
	405	410 415
Leu Glu Lys Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe		
	420	425 430
Leu Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn		
	435	440 445
Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu		
	450	455 460
Lys Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly		
465	470	475 480
Cys Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val		
	485	490 495
Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu		
	500	505 510
Asn Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr		
	515	520 525
Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu		
	530	535 540
Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu		
545	550	555 560
Gln Cys Arg Ile Cys Ile		
	565	

<210> SEQ ID NO 8
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 8

Met Lys Ala Arg Leu Leu Val Leu Leu Cys Ala Leu Ala Ala Thr Asp
1 5 10 15
Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20 25 30
Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35 40 45
Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
50 55 60
Ala Pro Leu Gln Leu Gly Lys Cys Asn Ile Ala Gly Trp Ile Leu Gly
65 70 75 80
Asn Pro Glu Cys Glu Ser Leu Leu Ser Asn Arg Ser Trp Ser Tyr Ile
85 90 95
Val Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Asp Phe
100 105 110
Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115 120 125
Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Lys His Asn
130 135 140
Thr Thr Lys Gly Val Thr Ala Ala Cys Ser His Ala Gly Lys Ser Ser
145 150 155 160
Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Ser Tyr Pro

-continued

165				170				175							
Asn	Leu	Ser	Asn	Ser	Tyr	Val	Asn	Asn	Lys	Gly	Lys	Glu	Val	Leu	Val
			180								185				190
Leu	Trp	Gly	Val	His	His	Pro	Ser	Asn	Ile	Asp	Asp	Gln	Gln	Thr	Leu
		195					200					205			
Tyr	Gln	Lys	Glu	Asn	Ala	Tyr	Val	Ser	Val	Val	Ser	Ser	Asn	Tyr	Asn
	210					215					220				
Arg	Arg	Phe	Thr	Pro	Glu	Ile	Ala	Glu	Arg	Pro	Lys	Val	Arg	Gly	Gln
	225				230					235					240
Ala	Gly	Arg	Met	Asn	Tyr	Tyr	Trp	Thr	Leu	Leu	Glu	Pro	Gly	Asp	Thr
			245						250					255	
Ile	Ile	Phe	Glu	Ala	Asn	Gly	Asn	Leu	Ile	Ala	Pro	Trp	Tyr	Ala	Phe
			260				265						270		
Ala	Leu	Ser	Arg	Gly	Phe	Gly	Ser	Gly	Ile	Ile	Thr	Ser	Asn	Ala	Ser
		275					280					285			
Met	His	Glu	Cys	Asp	Thr	Lys	Cys	Gln	Thr	Pro	Gln	Gly	Ala	Ile	Asn
	290					295					300				
Ser	Ser	Leu	Pro	Phe	Gln	Asn	Ile	His	Pro	Val	Thr	Ile	Gly	Glu	Cys
	305				310					315					320
Pro	Lys	Tyr	Val	Arg	Ser	Thr	Lys	Leu	Arg	Met	Val	Thr	Gly	Leu	Arg
			325						330					335	
Asn	Ile	Pro	Ser	Ile	Gln	Ser	Arg	Gly	Leu	Phe	Gly	Ala	Ile	Ala	Gly
		340							345				350		
Phe	Ile	Glu	Gly	Gly	Trp	Thr	Gly	Met	Ile	Asp	Gly	Trp	Tyr	Gly	Tyr
		355					360					365			
His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser
	370					375					380				
Thr	Gln	Asn	Ala	Ile	Asn	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile
	385				390					395					400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Asn
			405						410					415	
Leu	Glu	Lys	Arg	Met	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe
			420				425						430		
Leu	Asp	Ile	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
	435					440						445			
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu
	450				455						460				
Lys	Val	Lys	Ser	Gln	Leu	Arg	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly
	465				470					475					480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Asp	Asn	Glu	Cys	Met	Glu	Ser	Val
			485						490					495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu
		500							505					510	
Asn	Arg	Glu	Lys	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Met	Gly	Val	Tyr
		515					520					525			
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu
	530					535					540				
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu
	545				550					555					560
Gln	Cys	Arg	Ile	Cys	Ile										
			565												

-continued

```

<211> LENGTH: 566
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 9

Met Lys Ala Lys Leu Leu Val Leu Leu Cys Ala Phe Thr Ala Thr Asp
1      5      10      15
Ala Asp Thr Ile Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20      25      30
Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35      40      45
Leu Leu Glu Asp Ser His Asn Gly Lys Leu Cys Arg Leu Lys Gly Ile
50      55      60
Ala Pro Leu Gln Leu Gly Asn Cys Ser Ile Ala Gly Trp Ile Leu Gly
65      70      75      80
Asn Pro Glu Cys Glu Ser Leu Phe Ser Lys Lys Ser Trp Ser Tyr Ile
85      90      95
Ala Glu Thr Pro Asn Ser Glu Asn Gly Thr Cys Tyr Pro Gly Tyr Phe
100     105     110
Ala Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115     120     125
Glu Arg Phe Glu Ile Phe Pro Lys Glu Ser Ser Trp Pro Asn His Thr
130     135     140
Val Thr Lys Gly Val Thr Ala Ser Cys Ser His Lys Gly Lys Ser Ser
145     150     155     160
Phe Tyr Arg Asn Leu Leu Trp Leu Thr Glu Lys Asn Gly Ser Tyr Pro
165     170     175
Asn Leu Ser Lys Ser Tyr Val Asn Asn Lys Glu Lys Glu Val Leu Val
180     185     190
Leu Trp Gly Val His His Pro Ser Asn Ile Gly Asp Gln Arg Ala Ile
195     200     205
Tyr His Thr Glu Asn Ala Tyr Val Ser Val Val Ser Ser His Tyr Asn
210     215     220
Arg Arg Phe Thr Pro Glu Ile Ala Lys Arg Pro Lys Val Arg Asp Gln
225     230     235     240
Glu Gly Arg Ile Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
245     250     255
Ile Ile Phe Glu Ala Asn Gly Asn Leu Ile Ala Pro Trp Tyr Ala Phe
260     265     270
Ala Leu Ser Arg Gly Phe Gly Ser Gly Ile Ile Thr Ser Asn Ala Ser
275     280     285
Met Asp Glu Cys Asp Ala Lys Cys Gln Thr Pro Gln Gly Ala Ile Asn
290     295     300
Ser Ser Leu Pro Phe Gln Asn Val His Pro Val Thr Ile Gly Glu Cys
305     310     315     320
Pro Lys Tyr Val Arg Ser Thr Lys Leu Arg Met Val Thr Gly Leu Arg
325     330     335
Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
340     345     350
Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
355     360     365
His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser
370     375     380

```

-continued

Thr Gln Asn Ala Ile Asn Gly Ile Thr Asn Lys Val Asn Ser Val Ile
 385 390 395 400
 Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn Lys
 405 410 415
 Leu Glu Arg Arg Met Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe
 420 425 430
 Leu Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn
 435 440 445
 Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu
 450 455 460
 Lys Val Lys Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly
 465 470 475 480
 Cys Phe Glu Phe Tyr His Lys Cys Asn Asn Glu Cys Met Glu Ser Val
 485 490 495
 Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Glu Lys Ile Asp Gly Val Lys Leu Glu Ser Met Gly Val Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 10
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 10

Met Lys Ala Ile Leu Leu Val Leu Leu Cys Thr Phe Ala Ala Thr Asn
 1 5 10 15
 Ala Asp Thr Leu Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20 25 30
 Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35 40 45
 Leu Leu Glu Asp Lys His Asn Gly Lys Leu Cys Lys Leu Lys Gly Ile
 50 55 60
 Ala Pro Leu His Leu Gly Lys Cys Asn Ile Ala Gly Trp Leu Leu Gly
 65 70 75 80
 Asn Pro Glu Cys Glu Leu Leu Leu Thr Ala Ser Ser Trp Ser Tyr Ile
 85 90 95
 Val Glu Thr Ser Asn Ser Asp Asn Gly Thr Cys Tyr Pro Gly Asp Phe
 100 105 110
 Ile Asp Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115 120 125
 Glu Arg Phe Glu Ile Phe Pro Lys Thr Ser Ser Trp Pro Asn His Glu
 130 135 140
 Thr Asn Lys Gly Val Thr Ala Ala Cys Pro Tyr Ala Gly Ala Ser Ser
 145 150 155 160
 Phe Tyr Arg Asn Leu Ile Trp Leu Val Lys Lys Gly Asn Ser Tyr Pro
 165 170 175

-continued

Lys Leu Ser Lys Ser Tyr Val Asn Asn Lys Gly Lys Glu Val Leu Val
 180 185 190
 Leu Trp Gly Ile His His Pro Pro Thr Ser Thr Asp Gln Gln Ser Leu
 195 200 205
 Tyr Gln Asn Ala Asp Ala Tyr Val Phe Val Gly Ser Ser Lys Tyr Asn
 210 215 220
 Arg Lys Phe Lys Pro Glu Ile Ala Ala Arg Pro Lys Val Arg Asp Gln
 225 230 235 240
 Ala Gly Arg Met Asn Tyr Tyr Trp Thr Leu Ile Glu Pro Gly Asp Thr
 245 250 255
 Ile Thr Phe Glu Ala Thr Gly Asn Leu Val Val Pro Arg Tyr Ala Phe
 260 265 270
 Ala Met Asn Arg Gly Ser Gly Ser Gly Ile Ile Ile Ser Asp Ala Pro
 275 280 285
 Val His Asp Cys Asn Thr Lys Cys Gln Thr Pro Lys Gly Ala Ile Asn
 290 295 300
 Thr Ser Leu Pro Phe Gln Asn Ile His Pro Val Thr Ile Gly Glu Cys
 305 310 315 320
 Pro Lys Tyr Val Lys Ser Thr Lys Leu Arg Met Ala Thr Gly Leu Arg
 325 330 335
 Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
 340 345 350
 Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
 355 360 365
 His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser
 370 375 380
 Thr Gln Asn Ala Ile Asp Gly Ile Thr Asn Lys Val Asn Ser Val Ile
 385 390 395 400
 Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn His
 405 410 415
 Leu Glu Lys Arg Ile Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe
 420 425 430
 Leu Asp Ile Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn
 435 440 445
 Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu
 450 455 460
 Lys Val Arg Ser Gln Leu Lys Asn Asn Ala Lys Glu Ile Gly Asn Gly
 465 470 475 480
 Cys Phe Glu Phe Tyr His Lys Cys Asp Asp Thr Cys Met Glu Ser Val
 485 490 495
 Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Glu Glu Ile Asp Gly Val Lys Leu Glu Ser Thr Arg Ile Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 11

<211> LENGTH: 566

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 11

Met Lys Ala Ile Leu Leu Val Leu Leu Cys Thr Phe Thr Ala Ala Asn
 1             5             10             15
Ala Asp Thr Leu Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
 20             25             30
Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
 35             40             45
Leu Leu Glu Asp Arg His Asn Gly Lys Leu Cys Lys Leu Gly Gly Ile
 50             55             60
Ala Pro Leu His Leu Gly Lys Cys Asn Ile Ala Gly Trp Leu Leu Gly
 65             70             75             80
Asn Pro Glu Cys Glu Leu Leu Leu Thr Ala Ser Ser Trp Ser Tyr Ile
 85             90             95
Val Glu Thr Ser Asn Ser Asp Asn Gly Thr Cys Tyr Pro Gly Asp Phe
 100            105            110
Ile Asn Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
 115            120            125
Glu Arg Phe Glu Ile Phe Pro Lys Ala Ser Ser Trp Pro Asn His Glu
 130            135            140
Thr Asn Arg Gly Val Thr Ala Ala Cys Pro Tyr Ala Gly Ala Asn Ser
 145            150            155            160
Phe Tyr Arg Asn Leu Ile Trp Leu Val Lys Lys Gly Asn Ser Tyr Pro
 165            170            175
Lys Leu Ser Lys Ser Tyr Val Asn Asn Lys Gly Lys Glu Val Leu Val
 180            185            190
Leu Trp Gly Ile His His Pro Pro Thr Ser Thr Asp Gln Gln Ser Leu
 195            200            205
Tyr Gln Asn Ala Asp Ala Tyr Val Phe Val Gly Ser Ser Lys Tyr Asn
 210            215            220
Arg Lys Phe Lys Pro Glu Ile Ala Ala Arg Pro Lys Val Arg Gly Gln
 225            230            235            240
Ala Gly Arg Met Asn Tyr Tyr Trp Thr Leu Leu Glu Pro Gly Asp Thr
 245            250            255
Ile Thr Phe Glu Ala Thr Gly Asn Leu Val Val Pro Arg Tyr Ala Phe
 260            265            270
Ala Met Asn Arg Gly Ser Gly Ser Gly Ile Ile Ile Ser Asp Ala Pro
 275            280            285
Val His Asp Cys Asn Thr Lys Cys Gln Thr Pro Lys Gly Ala Ile Asn
 290            295            300
Thr Ser Leu Pro Phe Gln Asn Ile His Pro Val Thr Ile Gly Glu Cys
 305            310            315            320
Pro Lys Tyr Val Lys Ser Thr Lys Leu Arg Met Ala Thr Gly Leu Arg
 325            330            335
Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
 340            345            350
Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
 355            360            365
His His Gln Asn Glu Gln Gly Ser Gly Tyr Ala Ala Asp Gln Lys Ser
 370            375            380

```

-continued

Thr Gln Asn Ala Ile Asp Gly Ile Thr Asn Lys Val Asn Ser Val Ile
 385 390 395 400
 Glu Lys Met Asn Thr Gln Phe Thr Ala Val Gly Lys Glu Phe Asn His
 405 410 415
 Leu Glu Lys Arg Ile Glu Asn Leu Asn Lys Lys Val Asp Asp Gly Phe
 420 425 430
 Leu Asp Val Trp Thr Tyr Asn Ala Glu Leu Leu Val Leu Leu Glu Asn
 435 440 445
 Glu Arg Thr Leu Asp Phe His Asp Ser Asn Val Lys Asn Leu Tyr Glu
 450 455 460
 Lys Val Arg Ser Gln Leu Arg Asn Asn Ala Lys Glu Ile Gly Asn Gly
 465 470 475 480
 Cys Phe Glu Phe Tyr His Lys Cys Asp Asp Thr Cys Met Glu Ser Val
 485 490 495
 Lys Asn Gly Thr Tyr Asp Tyr Pro Lys Tyr Ser Glu Glu Ser Lys Leu
 500 505 510
 Asn Arg Glu Glu Ile Asp Gly Val Lys Leu Glu Ser Thr Arg Ile Tyr
 515 520 525
 Gln Ile Leu Ala Ile Tyr Ser Thr Val Ala Ser Ser Leu Val Leu Leu
 530 535 540
 Val Ser Leu Gly Ala Ile Ser Phe Trp Met Cys Ser Asn Gly Ser Leu
 545 550 555 560
 Gln Cys Arg Ile Cys Ile
 565

<210> SEQ ID NO 12
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (4)..(4)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (11)..(13)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (15)..(16)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (20)..(20)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (53)..(53)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (60)..(60)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (62)..(62)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (68)..(68)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (71)..(71)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

-continued

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (73)..(74)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (78)..(78)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (86)..(86)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (88)..(91)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (97)..(97)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (100)..(100)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (102)..(103)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (111)..(111)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (113)..(114)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (137)..(137)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (142)..(142)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (144)..(147)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (152)..(152)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (154)..(156)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (158)..(159)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (166)..(166)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (169)..(170)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (172)..(174)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (177)..(177)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (179)..(180)

[illegible]

-continued

<222> LOCATION: (269)..(269)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (274)..(275)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (278)..(278)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (284)..(284)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (286)..(286)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (288)..(291)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (293)..(294)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (300)..(300)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (305)..(305)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (312)..(312)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (325)..(325)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (332)..(332)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (390)..(390)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (416)..(416)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (419)..(419)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (421)..(421)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (435)..(435)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (467)..(467)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (471)..(471)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (489)..(491)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:

-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (516)..(516)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (525)..(527)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 12

Met Lys Ala Xaa Leu Leu Val Leu Leu Cys Xaa Xaa Xaa Ala Xaa Xaa
1          5          10          15

Ala Asp Thr Xaa Cys Ile Gly Tyr His Ala Asn Asn Ser Thr Asp Thr
20          25          30

Val Asp Thr Val Leu Glu Lys Asn Val Thr Val Thr His Ser Val Asn
35          40          45

Leu Leu Glu Asp Xaa His Asn Gly Lys Leu Cys Xaa Leu Xaa Gly Ile
50          55          60

Ala Pro Leu Xaa Leu Gly Xaa Cys Xaa Xaa Ala Gly Trp Xaa Leu Gly
65          70          75          80

Asn Pro Glu Cys Glu Xaa Leu Xaa Xaa Xaa Xaa Ser Trp Ser Tyr Ile
85          90          95

Xaa Glu Thr Xaa Asn Xaa Xaa Asn Gly Thr Cys Tyr Pro Gly Xaa Phe
100         105         110

Xaa Xaa Tyr Glu Glu Leu Arg Glu Gln Leu Ser Ser Val Ser Ser Phe
115         120         125

Glu Arg Phe Glu Ile Phe Pro Lys Xaa Ser Ser Trp Pro Xaa His Xaa
130         135         140

Xaa Xaa Xaa Gly Val Thr Ala Xaa Cys Xaa Xaa Xaa Gly Xaa Xaa Ser
145         150         155         160

Phe Tyr Arg Asn Leu Xaa Trp Leu Xaa Xaa Lys Xaa Xaa Xaa Tyr Pro
165         170         175

Xaa Leu Xaa Xaa Ser Tyr Val Asn Asn Lys Xaa Lys Glu Val Leu Val
180         185         190

Leu Trp Gly Xaa His His Pro Xaa Xaa Xaa Xaa Asp Gln Xaa Xaa Xaa
195         200         205

Tyr Xaa Xaa Xaa Xaa Ala Tyr Val Xaa Val Xaa Ser Ser Xaa Tyr Xaa
210         215         220

Arg Xaa Phe Xaa Pro Glu Ile Ala Xaa Arg Pro Lys Val Arg Xaa Gln
225         230         235         240

Xaa Gly Arg Xaa Asn Tyr Tyr Trp Thr Leu Xaa Glu Pro Gly Asp Thr
245         250         255

Ile Xaa Phe Glu Ala Xaa Gly Asn Leu Xaa Xaa Pro Xaa Tyr Ala Phe
260         265         270

Ala Xaa Xaa Arg Gly Xaa Gly Ser Gly Ile Ile Xaa Ser Xaa Ala Xaa
275         280         285

Xaa Xaa Xaa Cys Xaa Xaa Lys Cys Gln Thr Pro Xaa Gly Ala Ile Asn
290         295         300

Xaa Ser Leu Pro Phe Gln Asn Xaa His Pro Val Thr Ile Gly Glu Cys
305         310         315         320

Pro Lys Tyr Val Xaa Ser Thr Lys Leu Arg Met Xaa Thr Gly Leu Arg
325         330         335

Asn Ile Pro Ser Ile Gln Ser Arg Gly Leu Phe Gly Ala Ile Ala Gly
340         345         350

Phe Ile Glu Gly Gly Trp Thr Gly Met Ile Asp Gly Trp Tyr Gly Tyr
355         360         365

```

-continued

His	His	Gln	Asn	Glu	Gln	Gly	Ser	Gly	Tyr	Ala	Ala	Asp	Gln	Lys	Ser
370						375					380				
Thr	Gln	Asn	Ala	Ile	Xaa	Gly	Ile	Thr	Asn	Lys	Val	Asn	Ser	Val	Ile
385					390					395					400
Glu	Lys	Met	Asn	Thr	Gln	Phe	Thr	Ala	Val	Gly	Lys	Glu	Phe	Asn	Xaa
			405					410						415	
Leu	Glu	Xaa	Arg	Xaa	Glu	Asn	Leu	Asn	Lys	Lys	Val	Asp	Asp	Gly	Phe
		420					425						430		
Leu	Asp	Xaa	Trp	Thr	Tyr	Asn	Ala	Glu	Leu	Leu	Val	Leu	Leu	Glu	Asn
	435					440						445			
Glu	Arg	Thr	Leu	Asp	Phe	His	Asp	Ser	Asn	Val	Lys	Asn	Leu	Tyr	Glu
450					455						460				
Lys	Val	Xaa	Ser	Gln	Leu	Xaa	Asn	Asn	Ala	Lys	Glu	Ile	Gly	Asn	Gly
465				470						475					480
Cys	Phe	Glu	Phe	Tyr	His	Lys	Cys	Xaa	Xaa	Xaa	Cys	Met	Glu	Ser	Val
			485					490						495	
Lys	Asn	Gly	Thr	Tyr	Asp	Tyr	Pro	Lys	Tyr	Ser	Glu	Glu	Ser	Lys	Leu
		500						505					510		
Asn	Arg	Glu	Xaa	Ile	Asp	Gly	Val	Lys	Leu	Glu	Ser	Xaa	Xaa	Xaa	Tyr
		515					520					525			
Gln	Ile	Leu	Ala	Ile	Tyr	Ser	Thr	Val	Ala	Ser	Ser	Leu	Val	Leu	Leu
530					535						540				
Val	Ser	Leu	Gly	Ala	Ile	Ser	Phe	Trp	Met	Cys	Ser	Asn	Gly	Ser	Leu
545					550					555					560
Gln	Cys	Arg	Ile	Cys	Ile										
			565												

<210> SEQ ID NO 13
 <211> LENGTH: 1719
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 13

aagcttatga aggccaaagct gctggtgctg ctgtgcgcct tcacagccac ctacgccgac	60
accatctgca tcggttacca cgccaacaac agcaccgaca ccgtggatac cgtgctggaa	120
aagaacgtga ccgtgaccca cagcgtgaac ctgctggaag atagccacaa cggcaagctg	180
tgccggctga agggaaatgc cctctgcag ctgggcaact gctctatcgc cggttgatt	240
ctgggcaacc ccgagtgcga gagcctgttc agcaaagagt cctggtccta catcgtggaa	300
accccccaaca gcgagaacgg cacctgttac cccggctact tcgccgacta cgaggaaactg	360
cgggaacagc tgagcagcgt gtccagcttc gagagattcg agattttccc caaagagagc	420
agctggccca accacaccgt gaccaaaggc gtgaccgcct cctgctccca caatggcaag	480
agcagcttct acagaaacct gctgtggctg accgagaaga acggcagcta ccccaacctg	540
agcaagagct acgtgaacaa caaagaaaaa gaggtgctgg tgctgtgggg cgtgcaccac	600
cctagcaaca tcggcgacca gcggggccatc taccacaccg agaatgccta cgtgtccgtg	660
gtgtccagcc actacagcag acgggttacc cccgagatcg ccaagaggcc caaagtgcgg	720
gaccaggaag gccggatcaa ctactactgg aactgctgg aaccggcgca taccatcatc	780
ttcgaggcca acggcaacct gatcgccct tggtacgcct tcgccctgag cagaggtttt	840
ggcagcggca tcatcaccag caacgccagc atggacgagt gcgacgcaa gtgccagaca	900

-continued

cctcagggcg ccatcaatag cagcctgccc ttccagaacg tgcaccccggt gaccatcggc	960
gagtgcccca aatacgtgcg gagcaccaag ctgcggaatgg tcaccggcct gagaaacatc	1020
cccagcatcc agagcagggg cctgttcgga gccattgccc gctttatcga gggcggtg	1080
accggcatga tcgacgggtg gtacggctat caccaccaga acgagcaggg cagcggtac	1140
gccgcccgatc agaagtctac ccagaacgcc atcaacggca tcaccaacaa agtgaacagc	1200
gtgatcgaga agatgaacac ccagttcacc gccgtgggca aagagttcaa caagctggaa	1260
cggcggaatgg aaaacctgaa caagaaggtg gacgacggct tcctggacat ctggacctac	1320
aacgccgaac tgctggtgct gctggaaaac gagcggaccc tggacttcca cgacagcaac	1380
gtgaagaacc tgtacgagaa agtgaagtcc cagctgaaga acaacgcaa agagatcggc	1440
aacggctgct tcgagttcta ccacaagtgc aacaacgagt gcatggaaag cgtgaagaat	1500
ggcacctacg actaccccaa gtacagcgag gaaagcaagc tgaaccgcga gaagatcgac	1560
ggcgtgaagc tggaatccat gggcgtgtac cagatcctgg ccatctatag caccgtggcc	1620
agcagcctgg tgctgctggt gtctctgggc gccatcagct tttggatgtg cagcaacggc	1680
agcctgcagt gccgatctg tatcggcagc atcggtacc	1719

The invention claimed is:

1. A recombinant influenza hemagglutinin (HA) polypeptide comprising:

- (i) an amino acid sequence at least 99% identical to SEQ ID NO: 1, wherein the amino acid sequence of the polypeptide comprises no more than 5 amino acid substitutions relative to SEQ ID NO: 1;
- (ii) the amino acid sequence of SEQ ID NO: 2;
- (iii) an amino acid sequence at least 99% identical to SEQ ID NO: 3, wherein the amino acid sequence of the polypeptide comprises no more than 6 amino acid substitutions relative to SEQ ID NO: 3;
- (iv) an amino acid sequence at least 99% identical to SEQ ID NO: 4, wherein the amino acid sequence of the polypeptide comprises no more than 8 amino acid substitutions relative to SEQ ID NO: 4;
- (v) an amino acid sequence at least 98% identical to SEQ ID NO: 5, wherein the amino acid sequence of the polypeptide comprises no more than 10 amino acid substitutions relative to SEQ ID NO: 5;
- (vi) an amino acid sequence at least 99% identical to SEQ ID NO: 6, wherein the amino acid sequence of the polypeptide comprises no more than 8 amino acid substitutions relative to SEQ ID NO: 6;
- (vii) an amino acid sequence at least 97% identical to SEQ ID NO: 7, wherein the amino acid sequence of the polypeptide comprises no more than 10 amino acid substitutions relative to SEQ ID NO: 7;

(viii) an amino acid sequence at least 99% identical to SEQ ID NO: 8, wherein the amino acid sequence of the polypeptide comprises no more than 10 amino acid substitutions relative to SEQ ID NO: 8;

(ix) the amino acid sequence of SEQ ID NO: 9;

(x) an amino acid sequence at least 99% identical to SEQ ID NO: 10, wherein the amino acid sequence of the polypeptide comprises no more than 8 amino acid substitutions relative to SEQ ID NO: 10; or

(xi) an amino acid sequence at least 99% identical to SEQ ID NO: 11, wherein the amino acid sequence of the polypeptide comprises no more than 5 amino acid substitutions relative to SEQ ID NO: 11.

2. The influenza HA polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10 or SEQ ID NO: 11.

3. A composition comprising the influenza HA polypeptide of claim 1, and a pharmaceutically acceptable carrier.

4. A method of eliciting an immune response to influenza virus in a subject, comprising administering the composition of claim 3 to the subject.

5. The method of claim 4, wherein the composition further comprises an adjuvant.

6. The method of claim 4, wherein the composition is administered intramuscularly.

* * * * *