



US 2013021775A1

(19) **United States**

(12) **Patent Application Publication**
WILTON et al.

(10) **Pub. No.: US 2013/021775 A1**

(43) **Pub. Date: Aug. 22, 2013**

(54) **ANTISENSE OLIGONUCLEOTIDES FOR INDUCING EXON SKIPPING AND METHODS OF USE THEREOF**

of application No. 12/837,359, filed on Jul. 15, 2010, now Pat. No. 8,232,384, which is a continuation of application No. 11/570,691, filed on Jan. 15, 2008, now Pat. No. 7,807,816, filed as application No. PCT/AU2005/000943 on Jun. 28, 2005.

(71) Applicants: **Stephen Donald WILTON**, Applecross (AU); **Sue FLETCHER**, Bayswater (AU); **Graham MCCLOREY**, Bayswater (AU)

(30) **Foreign Application Priority Data**

Jun. 28, 2004 (AU) 2004903474

(72) Inventors: **Stephen Donald WILTON**, Applecross (AU); **Sue FLETCHER**, Bayswater (AU); **Graham MCCLOREY**, Bayswater (AU)

Publication Classification

(73) Assignee: **The University of Western Australia**

(51) **Int. Cl.**
C12N 15/113 (2006.01)
(52) **U.S. Cl.**
CPC **C12N 15/113** (2013.01)
USPC **514/44 A; 536/24.5; 435/320.1**

(21) Appl. No.: **13/826,613**

(22) Filed: **Mar. 14, 2013**

(57) **ABSTRACT**

Related U.S. Application Data

(63) Continuation of application No. 13/168,857, filed on Jun. 24, 2011, now abandoned, which is a continuation

An antisense molecule capable of binding to a selected target site to induce exon skipping in the dystrophin gene, as set forth in SEQ ID NO: 1 to 202.

FIGURE 1

bp	Acceptor	ESE	Donor
u <u>a</u> l <u>g</u> c <u>a</u> c <u>u</u> g <u>a</u> g <u>u</u> g <u>a</u> l <u>c</u> c <u>u</u> c <u>u</u> u <u>u</u> c <u>u</u> c <u>g</u> c <u>a</u> g <u>G</u> C <u>G</u> C <u>U</u> A <u>G</u> C <u>U</u> G <u>G</u> A <u>G</u> C <u>A</u> /// <u>CCGUGCAGACUGACGGguc<u>u</u>ca<u>u</u></u>			

SEQ ID NO:213

SEQ ID NO:214




FIGURE 2

H8A(-06+14) | H8A(-06+18)
M 600 300 100 50 20 UT 600 300 100 50 20 UT M




FIGURE 3

H7A(+45+67) H7A(+2+26)
M 600 300 100 50 20 600NM 600 300 100 50 20 600N M



FIGURE 4




FIGURE 5

6A(+69+91)

M 600 300 200 100 50 20 UT




FIGURE 6

H4A(+13+32)

M 600 300 100 50 20 UT Neg M






FIGURE 7



JSR604 - H11A(+75+97)

FIGURE 8B



JSR610 - H12A(+52+75)

FIGURE 8A




FIGURE 9A

FIGURE 9B




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18

H40A(-05+17)




FIGURE 19




FIGURE 20

H46A(+86+115)

600 300 200 100 L2K UT




FIGURE 21




FIGURE 22

**ANTISENSE OLIGONUCLEOTIDES FOR
INDUCING EXON SKIPPING AND METHODS
OF USE THEREOF**

CROSS-REFERENCE TO RELATED
APPLICATIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 13/168,857, filed Jun. 24, 2011, now pending, which application is a continuation of U.S. patent application Ser. No. 12/837,359, filed Jul. 15, 2010, now issued as U.S. Pat. No. 8,232,384, which application is a continuation of U.S. patent application Ser. No. 11/570,691, filed Jan. 15, 2008, now issued as U.S. Pat. No. 7,807,816, which application is a 35 U.S.C. §371 National Phase Application of PCT/AU2005/000943, filed Jun. 28, 2005, which claims priority to Australian Patent Application No. 2004903474, filed Jun. 28, 2004; which applications are each incorporated herein by reference in their entireties.

STATEMENT REGARDING SEQUENCE
LISTING

[0002] The Sequence Listing associated with the application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is SequenceListing.txt. The text file is 61 Kilobytes, was created on Mar. 11, 2013 and is being submitted electronically via EFS-Web.

FIELD OF THE INVENTION

[0003] The present invention relates to novel antisense compounds and compositions suitable for facilitating exon skipping. It also provides methods for inducing exon skipping using the novel antisense compounds as well as therapeutic compositions adapted for use in the methods of the invention.

BACKGROUND ART

[0004] Significant effort is currently being expended researching methods for suppressing or compensating for disease-causing mutations in genes. Antisense technologies are being developed using a range of chemistries to affect gene expression at a variety of different levels (transcription, splicing, stability, translation). Much of that research has focused on the use of antisense compounds to correct or compensate for abnormal or disease-associated genes in a myriad of different conditions.

[0005] Antisense molecules are able to inhibit gene expression with exquisite specificity and because of this many research efforts concerning oligonucleotides as modulators of gene expression have focused on inhibiting the expression of targeted genes such as oncogenes or viral genes. The antisense oligonucleotides are directed either against RNA (sense strand) or against DNA where they form triplex structures inhibiting transcription by RNA polymerase II. To achieve a desired effect in specific gene down-regulation, the oligonucleotides must either promote the decay of the targeted mRNA or block translation of that mRNA, thereby effectively preventing de novo synthesis of the undesirable target protein.

[0006] Such techniques are not useful where the object is to up-regulate production of the native protein or compensate for mutations which induce premature termination of translation such as nonsense or frame-shifting mutations. Further-

more, in cases where a normally functional protein is prematurely terminated because of mutations therein, a means for restoring some functional protein production through antisense technology has been shown to be possible through intervention during the splicing processes (Sierakowska H, et al., (1996) *Proc Natl Acad Sci USA* 93, 12840-12844; Wilton S D, et al., (1999) *Neuromusc Disorders* 9, 330-338; van Deutekom J C et al., (2001) *Human Mol Genet* 10, 1547-1554). In these cases, the defective gene transcript should not be subjected to targeted degradation so the antisense oligonucleotide chemistry should not promote target mRNA decay.

[0007] In a variety of genetic diseases, the effects of mutations on the eventual expression of a gene can be modulated through a process of targeted exon skipping during the splicing process. The splicing process is directed by complex multi-particle machinery that brings adjacent exon-intron junctions in pre-mRNA into close proximity and performs cleavage of phosphodiester bonds at the ends of the introns with their subsequent reformation between exons that are to be spliced together. This complex and highly precise process is mediated by sequence motifs in the pre-mRNA that are relatively short semi-conserved RNA segments to which bind the various nuclear splicing factors that are then involved in the splicing reactions. By changing the way the splicing machinery reads or recognises the motifs involved in pre-mRNA processing, it is possible to create differentially spliced mRNA molecules. It has now been recognised that the majority of human genes are alternatively spliced during normal gene expression, although the mechanisms invoked have not been identified. Using antisense oligonucleotides, it has been shown that errors and deficiencies in a coded mRNA could be bypassed or removed from the mature gene transcripts.

[0008] In nature, the extent of genetic deletion or exon skipping in the splicing process is not fully understood, although many instances have been documented to occur, generally at very low levels (Sherrat T G, et al., (1993) *Am J Hum Genet* 53, 1007-1015). However, it is recognised that if exons associated with disease-causing mutations can be specifically deleted from some genes, a shortened protein product can sometimes be produced that has similar biological properties of the native protein or has sufficient biological activity to ameliorate the disease caused by mutations associated with the target exon (Lu Q L, et al., (2003) *Nature Medicine* 9, 1009-1014; Aartsma-Rus A et al., (2004) *Am J Hum Genet* 74: 83-92).

[0009] This process of targeted exon skipping is likely to be particularly useful in long genes where there are many exons and introns, where there is redundancy in the genetic constitution of the exons or where a protein is able to function without one or more particular exons (e.g. with the dystrophin gene, which consists of 79 exons; or possibly some collagen genes which encode for repeated blocks of sequence or the huge nebulin or titin genes which are comprised of ~80 and over 370 exons, respectively).

[0010] Efforts to redirect gene processing for the treatment of genetic diseases associated with truncations caused by mutations in various genes have focused on the use of antisense oligonucleotides that either: (1) fully or partially overlap with the elements involved in the splicing process; or (2) bind to the pre-mRNA at a position sufficiently close to the element to disrupt the binding and function of the splicing factors that would normally mediate a particular splicing

reaction which occurs at that element (e.g., binds to the pre-mRNA at a position within 3, 6, or 9 nucleotides of the element to be blocked).

[0011] For example, modulation of mutant dystrophin pre-mRNA splicing with antisense oligoribonucleotides has been reported both in vitro and in vivo. In one type of dystrophin mutation reported in Japan, a 52-base pair deletion mutation causes exon 19 to be removed with the flanking introns during the splicing process (Matsuo et al., (1991) *J Clin Invest.*, 87:2127-2131). An in vitro minigene splicing system has been used to show that a 31-mer 2'-O-methyl oligoribonucleotide complementary to the 5' half of the deleted sequence in dystrophin Kobe exon 19 inhibited splicing of wild-type pre-mRNA (Takeshima et al. (1995), *J. Clin. Invest.*, 95, 515-520). The same oligonucleotide was used to induce exon skipping from the native dystrophin gene transcript in human cultured lymphoblastoid cells.

[0012] Dunckley et al., (1997) *Nucleosides & Nucleotides*, 16, 1665-1668 described in vitro constructs for analysis of splicing around exon 23 of mutated dystrophin in the mdx mouse mutant, a model for muscular dystrophy. Plans to analyse these constructs in vitro using 2' modified oligonucleotides targeted to splice sites within and adjacent to mouse dystrophin exon 23 were discussed, though no target sites or sequences were given.

[0013] 2'-O-methyl oligoribonucleotides were subsequently reported to correct dystrophin deficiency in myoblasts from the mdx mouse from this group. An antisense oligonucleotide targeted to the 3' splice site of murine dystrophin intron 22 was reported to cause skipping of the mutant exon as well as several flanking exons and created a novel in-frame dystrophin transcript with a novel internal deletion. This mutated dystrophin was expressed in 1-2% of antisense treated mdx myotubes. Use of other oligonucleotide modifications such as 2'-O-methoxyethyl phosphodiester are described (Dunckley et al. (1998) *Human Mol. Genetics*, 5, 1083-90).

[0014] Thus, antisense molecules may provide a tool in the treatment of genetic disorders such as Duchenne Muscular Dystrophy (DMD). However, attempts to induce exon skipping using antisense molecules have had mixed success. Studies on dystrophin exon 19, where successful skipping of that exon from the dystrophin pre-mRNA was achieved using a variety of antisense molecules directed at the flanking splice sites or motifs within the exon involved in exon definition as described by Errington et al. (2003) *J Gen Med* 5, 518-527".

[0015] In contrast to the apparent ease of exon 19 skipping, the first report of exon 23 skipping in the mdx mouse by Dunckley et al., (1998) is now considered to be reporting only a naturally occurring revertant transcript or artefact rather than any true antisense activity. In addition to not consistently generating transcripts missing exon 23, Dunckley et al., (1998) did not show any time course of induced exon skipping, or even titration of antisense oligonucleotides, to demonstrate dose dependent effects where the levels of exon skipping corresponded with increasing or decreasing amounts of antisense oligonucleotide. Furthermore, this work could not be replicated by other researchers.

[0016] The first example of specific and reproducible exon skipping in the mdx mouse model was reported by Wilton et al., (1999) *Neuromuscular Disorders* 9, 330-338. By directing an antisense molecule to the donor splice site, consistent and efficient exon 23 skipping was induced in the dystrophin mRNA within 6 hours of treatment of the cultured cells.

Wilton et al, (1999), also describe targeting the acceptor region of the mouse dystrophin pre-mRNA with longer antisense oligonucleotides and being unable to repeat the published results of Dunckley et al., (1998). No exon skipping, either 23 alone or multiple removal of several flanking exons, could be reproducibly detected using a selection of antisense oligonucleotides directed at the acceptor splice site of intron 22.

[0017] While the first antisense oligonucleotide directed at the intron 23 donor splice site induced consistent exon skipping in primary cultured myoblasts, this compound was found to be much less efficient in immortalized cell cultures expressing higher levels of dystrophin. However, with refined targeting and antisense oligonucleotide design, the efficiency of specific exon removal was increased by almost an order of magnitude (see Mann C J et al., (2002) *J Gen Med* 4, 644-654).

[0018] Thus, there remains a need to provide antisense oligonucleotides capable of binding to and modifying the splicing of a target nucleotide sequence. Simply directing the antisense oligonucleotides to motifs presumed to be crucial for splicing is no guarantee of the efficacy of that compound in a therapeutic setting.

SUMMARY OF THE INVENTION

[0019] The present invention provides antisense molecule compounds and compositions suitable for binding to RNA motifs involved in the splicing of pre-mRNA that are able to induce specific and efficient exon skipping and a method for their use thereof.

[0020] The choice of target selection plays a crucial role in the efficiency of exon skipping and hence its subsequent application of a potential therapy. Simply designing antisense molecules to target regions of pre-mRNA presumed to be involved in splicing is no guarantee of inducing efficient and specific exon skipping. The most obvious or readily defined targets for splicing intervention are the donor and acceptor splice sites although there are less defined or conserved motifs including exonic splicing enhancers, silencing elements and branch points.

[0021] The acceptor and donor splice sites have consensus sequences of about 16 and 8 bases respectively (see FIG. 1 for schematic representation of motifs and domains involved in exon recognition, intron removal and the splicing process).

[0022] According to a first aspect, the invention provides antisense molecules capable of binding to a selected target to induce exon skipping.

[0023] For example, to induce exon skipping in exons 3 to 8, 10 to 16, 19 to 40, 42 to 44, 46, 47, and 50 to 53 in the Dystrophin gene transcript the antisense molecules are preferably selected from the group listed in Table 1A.

[0024] In a further example, it is possible to combine two or more antisense oligonucleotides of the present invention together to induce multiple exon skipping in exons 19-20, and 53. This is a similar concept to targeting of a single exon. A combination or "cocktail" of antisense oligonucleotides are directed at adjacent exons to induce efficient exon skipping.

[0025] In another example, to induce exon skipping in exons 19-20, 31, 34 and 53 it is possible to improve exon skipping of a single exon by joining together two or more antisense oligonucleotide molecules. This concept is termed by the inventor as a "weasel", an example of a cunningly

designed antisense oligonucleotide. A similar concept has been described in Aartsma-Rus A et al., (2004) *Am J Hum Genet* 74: 83-92).

[0026] According to a second aspect, the present invention provides antisense molecules selected and/or adapted to aid in the prophylactic or therapeutic treatment of a genetic disorder comprising at least an antisense molecule in a form suitable for delivery to a patient.

[0027] According to a third aspect, the invention provides a method for treating a patient suffering from a genetic disease wherein there is a mutation in a gene encoding a particular protein and the affect of the mutation can be abrogated by exon skipping, comprising the steps of: (a) selecting an antisense molecule in accordance with the methods described herein; and (b) administering the molecule to a patient in need of such treatment.

[0028] The invention also addresses the use of purified and isolated antisense oligonucleotides of the invention, for the manufacture of a medicament for treatment of a genetic disease.

[0029] The invention further provides a method of treating a condition characterised by Duchenne muscular dystrophy, which method comprises administering to a patient in need of treatment an effective amount of an appropriately designed antisense oligonucleotide of the invention, relevant to the particular genetic lesion in that patient. Further, the invention provides a method for prophylactically treating a patient to prevent or at least minimise Duchenne muscular dystrophy, comprising the step of: administering to the patient an effective amount of an antisense oligonucleotide or a pharmaceutical composition comprising one or more of these biological molecules.

[0030] The invention also provides kits for treating a genetic disease, which kits comprise at least an antisense oligonucleotide of the present invention, packaged in a suitable container and instructions for its use.

[0031] Other aspects and advantages of the invention will become apparent to those skilled in the art from a review of the ensuing description, which proceeds with reference to the following figures.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] FIG. 1 Schematic representation of motifs and domains involved in exon recognition, intron removal and the splicing process (SEQ ID NOS: 213 and 214).

[0033] FIG. 2. Diagrammatic representation of the concept of antisense oligonucleotide induced exon skipping to bypass disease-causing mutations (not drawn to scale). The hatched box represents an exon carrying a mutation that prevents the translation of the rest of the mRNA into a protein. The solid black bar represents an antisense oligonucleotide that prevents inclusion of that exon in the mature mRNA.

[0034] FIG. 3 Gel electrophoresis showing differing efficiencies of two antisense molecules directed at exon 8 acceptor splice site. The preferred compound [H8A(-06+18)] induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured normal human muscle cells. The less preferred antisense oligonucleotide [H8A(-06+14)] also induces efficient exon skipping, but at much higher concentrations. Other antisense oligonucleotides directed at exon 8 either only induced lower levels of exon skipping or no detectable skipping at all (not shown).

[0035] FIG. 4 Gel electrophoresis showing differing efficiencies of two antisense molecules directed at internal

domains within exon 7, presumably exon splicing enhancers. The preferred compound [H7A(+45+67)] induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells. The less preferred antisense oligonucleotide [H7A(+2+26)] induces only low levels of exon skipping at the higher transfection concentrations. Other antisense oligonucleotides directed at exon 7 either only induced lower levels of exon skipping or no detectable skipping at all (not shown).

[0036] FIG. 5 Gel electrophoresis showing an example of low efficiency exon 6 skipping using two non-preferred antisense molecules directed at human exon 6 donor splice site. Levels of induced exon 6 skipping are either very low [H6D(+04-21)] or almost undetectable [H6D(+18-04)]. These are examples of non-preferred antisense oligonucleotides to demonstrate that antisense oligonucleotide design plays a crucial role in the efficacy of these compounds.

[0037] FIG. 6 Gel electrophoresis showing strong and efficient human exon 6 skipping using an antisense molecules [H6A(+69+91)] directed at an exon 6 internal domain, presumably an exon splicing enhancer. This preferred compound induces consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells.

[0038] FIG. 7 Gel electrophoresis showing strong human exon 4 skipping using an antisense molecule H4A(+13+32) directed at an exon 6 internal domain, presumably an exon splicing enhancer. This preferred compound induces strong and consistent exon skipping at a transfection concentration of 20 nanomolar in cultured human muscle cells,

[0039] FIG. 8 Gel electrophoresis showing (8B) strong human exon 11 skipping using antisense molecule H11A(+75+97) directed at an exon 11 internal domain; and (8B) strong human exon 12 skipping using antisense molecule H12A(+52+75) directed at exon 12 internal domain.

[0040] FIG. 9 Gel electrophoresis showing (9A) strong human exon 15 skipping using antisense molecules H15A(+48+71) and H15A(-12+19) directed at an exon 15 internal domain; and (9B) strong human exon 16 skipping using antisense molecules H16A(-12+19) and H16A(-06+25).

[0041] FIG. 10 Gel electrophoresis showing human exon 19/20 skipping using antisense molecules H20A(+44+71) and H20A(+149+170) directed at an exon 20 and a "cocktail" of antisense oligonucleotides H19A(+35+65, H20A(+44+71) and H20A(+149+170) directed at exons 19/20.

[0042] FIG. 11 Gel electrophoresis showing human exon 19/20 skipping using "weasels" directed at exons 19 and 20.

[0043] FIG. 12 Gel electrophoresis showing exon 22 skipping using antisense molecules H22A(+125+106), H22A(+47+69), H22A(+80+101) and H22D(+13-11) directed at exon 22.

[0044] FIG. 13 Gel electrophoresis showing exon 31 skipping using antisense molecules H31D(+01-25) and H31D(+03-22); and a "cocktail" of antisense molecules directed at exon 31.

[0045] FIG. 14 Gel electrophoresis showing exon 33 skipping using antisense molecules H33A(+30+56) and H33A(+64+88) directed at exon 33.

[0046] FIG. 15 Gel electrophoresis showing exon 35 skipping using antisense molecules H35A(+141+161), H35A(+116+135), and H35A(+24+43) and a "cocktail" of two antisense molecules, directed at exon 35.

[0047] FIG. 16 Gel electrophoresis showing exon 36 skipping using antisense molecules H32A(+49+73) and H36A(+26+50) directed at exon 36.

[0048] FIG. 17 Gel electrophoresis showing exon 37 skipping using antisense molecules H37A(+82+105) and H37A(+134+157) directed at exon 37.

[0049] FIG. 18 Gel electrophoresis showing exon 38 skipping using antisense molecule H38A(+88+112) directed at exon 38.

[0050] FIG. 19 Gel electrophoresis showing exon 40 skipping using antisense molecule H40A(-05+17) directed at exon 40.

[0051] FIG. 20 Gel electrophoresis showing exon 42 skipping using antisense molecule H42A(-04+23) directed at exon 42.

[0052] FIG. 21 Gel electrophoresis showing exon 46 skipping using antisense molecule H46A(+86+115) directed at exon 46

[0053] FIG. 22 Gel electrophoresis showing exon 51, exon 52 and exon 53 skipping using various antisense molecules directed at exons 51, 52 and 53, respectively. A "cocktail" of antisense molecules is also shown directed at exon 53.

BRIEF DESCRIPTION OF THE SEQUENCE LISTINGS

[0054]

TABLE 1A

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".		
SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
1	H8A(-06 + 18)	GAU AGG UGG UAU CAA CAU CUG UAA
2	H8A (-03 + 18)	GAU AGG UGG UAU CAA CAU CUG
3	H8A(-07 + 18)	GAU AGG UGG UAU CAA CAU CUG UAA G
4	H8A(-06 + 14)	GGU GGU AUC AAC AUC UGU AA
5	H8A(-10 + 10)	GUA UCA ACA UCU GUA AGC AC
6	H7A(+45 + 67)	UGC AUG UUC CAG UCG UUG UGU GG
7	H7A(+02 + 26)	CAC UAU UCC AGU CAA AUA GGU CUG G
8	H7D(+15 - 10)	AUU UAC CAA CCU UCA GGA UCG AGU A
9	H7A(-18 + 03)	GGC CUA AAA CAC AUA CAC AUA
10	C6A(-10 + 10)	CAU UUU UGA CCU ACA UGU GG
11	C6A(-14 + 06)	UUU GAC CUA CAU GUG GAA AG
12	C6A(-14 + 12)	UAC AUU UUU GAC CUA CAU GUG GAA AG
13	C6A(-13 + 09)	AUU UUU GAC CUA CAU GGG AAA G
14	CH6A(+69 + 91)	UAC GAG UUG AUU GUC GGA CCC AG
15	C6D(+12 - 13)	GUG GUC UCC UUA CCU AUG ACU GUG G
16	C6D(+06 - 11)	GGU CUC CUU ACC UAU GA
17	H6D(+04 - 21)	UGU CUC AGU AAU CUU CUU ACC UAU
18	H6D(+18 - 04)	UCU UAC CUA UGA CUA UGG AUG AGA
19	H4A(+13 + 32)	GCA UGA ACU CUU GUG GAU CC
20	H4D(+04 - 16)	CCA GGG UAC UAC UUA CAU UA
21	H4D(-24 - 44)	AUC GUG UGU CAC AGC AUC CAG
22	H4A(+11 + 40)	UGU UCA GGG CAU GAA CUC UUG UGG AUC CUU
23	H3A(+30 + 60)	UAG GAG GCG CCU CCC AUC CUG UAG GUC ACU G
24	H3A(+35 + 65)	AGG UCU AGG AGG CGC CUC CCA UCC UGU AGG U
25	H3A(+30 + 54)	GCG CCU CCC AUC CUG UAG GUC ACU G
26	H3D(+46 - 21)	CUU CGA GGA GGU CUA GGA GGC GCC UC

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
27	H3A(+30 + 50)	CUC CCA UCC UGU AGG UCA CUG
28	H3D(+19 - 03)	UAC CAG UUU UUG CCC UGU CAG G
29	H3A(-06 + 20)	UCA AUA UGC UGC UUC CCA AAC UGA AA
30	H3A(+37 + 61)	CUA GGA GGC GCC UCC CAU CCU GUA G
31	H5A(+20 + 50)	UUA UGA UUU CCA UCU ACG AUG UCA GUA CUU C
32	H5D(+25 - 05)	CUU ACC UGC CAG UGG AGG AUU AUA UUC CAA A
33	H5D(+10 - 15)	CAU CAG GAU UCU UAC CUG CCA GUG G
34	H5A(+10 + 34)	CGA UGU CAG UAC UUC CAA UAU UCA C
35	H5D(-04 - 21)	ACC AUU CAU CAG GAU UCU
36	H5D(+16 - 02)	ACC UGC CAG UGG AGG AUU
37	H5A(-07 + 20)	CCA AUA UUC ACU AAA UCA ACC UGU UAA
38	H5D(+18 - 12)	CAG GAU UGU UAC CUG CCA GUG GAG GAU UAU
39	H5A(+05 + 35)	ACG AUG UCA GUA CUU CCA AUA UUC ACU AAA U
40	H5A(+15 + 45)	AUU UCC AUC UAC GAU GUC AGU ACU UCC AAU A
41	H10A(-05 + 16)	CAG GAG CUU CCA AAU GCU GCA
42	H10A(-05 + 24)	CUU GUC UUC AGG AGC UUC CAA AUG CUG CA
43	H10A(+98 + 119)	UCC UCA GCA GAA AGA AGC CAC G
44	H10A(+130 + 149)	UUA GAA AUC UCU CCU UGU GC
45	H10A(-33 - 14)	UAA AUU GGG UGU UAC ACA AU
46	H11D(+26 + 49)	CCC UGA GGC AUU CCC AUC UUG AAU
47	H11D(+11 - 09)	AGG ACU UAC UUG CUU UGU UU
48	H11A(+118 + 140)	CUU GAA UUU AGG AGA UUC AUC UG
49	H11A(+75 + 97)	CAU CUU CUG AUA AUU UUC CUG UU
50	H12A(+52 + 75)	UCU UCU GUU UUU GUU AGC CAG UCA
51	H12A(-10 + 10)	UCU AUG UAA ACU GAA AAU UU
52	H12A(+11 + 30)	UUC UGG AGA UCC AUU AAA AC
53	H13A(+77 + 100)	CAG CAG UUG CGU GAU CUC CAC UAG
54	H13A(+55 + 75)	UUC AUC AAC UAC CAC CAC CAU
55	H13D(+06 - 19)	CUA AGC AAA AUA AUC UGA CCU UAA G
56	H14A(+37 + 64)	CUU GUA AAA GAA CCC AGC GGU CUU CUG U
57	H14A(+14 + 35)	CAU CUA CAG AUG UUU GCC CAU C
58	H14A(+51 + 73)	GAA GGA UGU CUU GUA AAA GAA CC

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
59	H14D(-02 + 18)	ACC UGU UCU UCA GUA AGA CG
60	H14D(+14 - 10)	CAU GAC ACA CCU GUU CUU CAG UAA
61	H14A(+61 + 80)	CAU UUG AGA AGG AUG UCU UG
62	H14A(-12 + 12)	AUC UCC CAA UAC CUG GAG AAG AGA
63	H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA AGA CAU U
64	H15A(+48 + 71)	UCU UUA AAG CCA GUU GUG UGA AUC
65	H15A(+08 + 28)	UUU CUG AAA GCC AUG CAC UAA
66	H15D(+17 - 08)	GUA CAU ACG GCC AGU UUU UGA AGA C
67	H16A(-12 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AAA ACA A
68	H16A(-06 + 25)	UCU UUU CUA GAU CCG CUU UUA AAA CCU GUU A
69	H16A(-06 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU A
70	H16A(+87 + 109)	CCG UCU UCU GGG UCA CUG ACU UA
71	H16A(-07 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AA
72	H16A(-07 + 13)	CCG CUU UUA AAA CCU GUU AA
73	H16A(+12 + 37)	UGG AUU GCU UUU UCU UUU CUA GAU CC
74	H16A(+92 + 116)	CAU GCU UCC GUC UUC UGG GUC ACU G
75	H16A(+45 + 67)	G AUC UUG UUU GAG UGA AUA CAG U
76	H16A(+105 + 126)	GUU AUC CAG CCA UGC UUC CGU C
77	H16D(+05 - 20)	UGA UAA UUG GUA UCA CUA ACC UGU G
78	H16D(+12 - 11)	GUA UCA CUA ACC UGU GCU GUA C
79	H19A(+35 + 53)	CUG CUG GCA UCU UGC AGU U
80	H19A(+35 + 65)	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C
83	H20A(+185 + 203)	UGA UGG GGU GGU GGG UUG G
84	H20A(-08 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA G
85	H20A(+30 + 53)	CCG GCU GUU CAG UUG UUC UGA GGC
86	H20A(-11 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA GAA A
87	H20D(+08 - 20)	GAA GGA GAA GAG AUU CUU ACC UUA CAA A
88	H20A(+44 + 63)	AUU CGA UCC ACC GGC UGU UC
89	H20A(+149 + 168)	CAG CAG UAG UUG UCA UCU GC
90	H21A(-06 + 16)	GCC GGU UGA CUU CAU CCU GUG C

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
91	H21A(+85 + 106)	CUG CAU CCA GGA ACA UGG GUC C
92	H21A(+85 + 108)	GUC UGC AUC CAG GAA CAU GGG UC
93	H21A(+08 + 31)	GUU GAA GAU CUG AUA GCC GGU UGA
94	H21D(+18 - 07)	UAC UUA CUG UCU GUA GCU CUU UCU
95	H22A(+22 + 45)	CAC UCA UGG UCU CCU GAU AGC GCA
96	H22A(+125 + 106)	CUG CAA UUC CCC GAG UCU CUG C
97	H22A(+47 + 69)	ACU GCU GGA CCC AUG UCC UGA UG
98	H22A(+80 + 101)	CUA AGU UGA GGU AUG GAG AGU
99	H22D(+13 - 11)	UAU UCA CAG ACC UGC AAU UCC CC
100	H23A(+34 + 59)	ACA GUG GUG CUG AGA UAG UAU AGG CC
101	H23A(+18 + 39)	UAG GCC ACU UUG UUG CUC UUG C
102	H23A(+72 + 90)	UUC AGA GGG CGC UUU CUU C
103	H24A(+48 + 70)	GGG CAG GCC AUU CCU CCU UCA GA
104	H24A(-02 + 22)	UCU UCA GGG UUU GUA UGU GAU UCU
105	H25A(+9 + 36)	CUG GGC UGA AUU GUC UGA AUA UCA CUG
106	H25A(+131 + 156)	CUG UUG GCA CAU GUG AUC CCA CUG AG
107	H25D(+16 - 08)	GUC UAU ACC UGU UGG CAC AUG UGA
108	H26A(+132 + 156)	UGC UUU CUG UAA UUC AUC UGG AGU U
109	H26A(-07 + 19)	CCU CCU UUC UGG CAU AGA CCU UCC AC
110	H26A(+68 + 92)	UGU GUC AUC CAU UCG UGC AUC UCU G
111	H27A(+82 + 106)	UUA AGG CCU CUU GUG CUA CAG GUG G
112	H27A(-4 + 19)	GGG GCU CUU CUU UAG CUC UCU GA
113	H27D(+19 - 03)	GAC UUC CAA AGU CUU GCA UUU C
114	H28A(-05 + 19)	GCC AAC AUG CCC AAA CUU CCU AAG
115	H28A(+99 + 124)	CAG AGA UUU CCU CAG CUC CGC CAG GA
116	H28D(+16 - 05)	CUU ACA UCU AGC ACC UCA GAG
117	H29A(+57 + 81)	UCC GCC AUC UGU UAG GGU CUG UGC C
118	H29A(+18 + 42)	AUU UGG GUU AUC CUC UGA AUG UCG C
119	H29D(+17 - 05)	CAU ACC UCU UCA UGU AGU UCC C
120	H30A(+122 + 147)	CAU UUG AGC UGC GUC CAC CUU GUC UG
121	H30A(+25 + 50)	UCC UGG GCA GAC UGG AUG CUC UGU UC
122	H30D(+19 - 04)	UUG CCU GGG CUU CCU GAG GCA UU

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
123	H31D(+06 - 18)	UUC UGA AAU AAC AUA UAC CUG UGC
124	H31D(+03 - 22)	UAG UUU CUG AAA UAA CAU AUA CCU G
125	H31A(+05 + 25)	GAC UUG UCA AAU CAG AUU GGA
126	H31D(+04 - 20)	GUU UCU GAA AUA ACA UAU ACC UGU
127	H32D(+04 - 16)	CAC CAG AAA UAC AUA CCA CA
128	H32A(+151 + 170)	CAA UGA UUU AGC UGU GAC UG
129	H32A(+10 + 32)	CGA AAC UUC AUG GAG ACA UCU UG
130	H32A(+49 + 73)	CUU GUA GAC GCU GCU CAA AAU UGG C
131	H33D(+09 - 11)	CAU GCA CAC ACC UUU GCU CC
132	H33A(+53 + 76)	UCU GUA CAA UCU GAC GUC CAG UCU
133	H33A(+30 + 56)	GUC UUU AUC ACC AUU UCC ACU UCA GAC
134	H33A(+64 + 88)	CCG UCU GCU UUU UCU GUA CAA UCU G
135	H34A(+83 + 104)	UCC AUA UCU GUA GCU GCC AGC C
136	H34A(+143 + 165)	CCA GGC AAC UUC AGA AUC CAA AU
137	H34A(-20 + 10)	UUU CUG UUA CCU GAA AAG AAU UAU AAU GAA
138	H34A(+46 + 70)	CAU UCA UUU CCU UUC GCA UCU UAC G
139	H34A(+95 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG
140	H34D(+10 - 20)	UUC AGU GAU AUA GGU UUU ACC UUU CCC CAG
141	H34A(+72 + 96)	CUG UAG CUG CCA GCC AUU CUG UCA AG
142	H35A(+141 + 161)	UCU UCU GCU CGG GAG GUG ACA
143	H35A(+116 + 135)	CCA GUU ACU AUU CAG AAG AC
144	H35A(+24 + 43)	UCU UCA GGU GCA CCU UCU GU
145	H36A(+26 + 50)	UGU GAU GUG GUC CAC AUU CUG GUC A
146	H36A(-02 + 18)	CCA UGU GUU UCU GGU AUU CC
147	H37A(+26 + 50)	CGU GUA GAG UCC ACC UUU GGG CGU A
148	H37A(+82 + 105)	UAC UAA UUU CCU GCA GUG GUC ACC
149	H37A(+134 + 157)	UUC UGU GUG AAA UGG CUG CAA AUC
150	H38A(-01 + 19)	CCU UCA AAG GAA UGG AGG CC
151	H38A(+59 + 83)	UGC UGA AUU UCA GCC UCC AGU GGU U
152	H38A(+88 + 112)	UGA AGU CUU CCU CUU UCA GAU UCA C
153	H39A(+62 + 85)	CUG GCU UUC UCU CAU CUG UGA UUC

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA. Since these 2'-O-methyl antisense oligonucleotides are more RNA-like, U represents uracil. With other antisense chemistries such as peptide nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
154	H39A(+39 + 58)	GUU GUA AGU UGU CUC CUC UU
155	H39A(+102 + 121)	UUG UCU GUA ACA GCU GCU GU
156	H39D(+10 - 10)	GCU CUA AUA CCU UGA GAG CA
157	H40A(-05 + 17)	CUU UGA GAC CUC AAA UCC UGU U
158	H40A(+129 + 153)	CUU UAU UUU CCU UUC AUC UCU GGG C
159	H42A(-04 + 23)	AUC GUU UCU UCA CGG ACA GUG UGC UGG
160	H42A(+86 + 109)	GGG CUU GUG AGA CAU GAG UGA UUU
161	H42D(+19 - 02)	A CCU UCA GAG GAC UCC UCU UGC
162	H43D(+10 - 15)	UAU GUG UUA CCU ACC CUU GUC GGU C
163	H43A(+101 + 120)	GGA GAG AGC UUC CUG UAG CU
164	H43A(+78 + 100)	UCA CCC UUU CCA CAG GCG UUG CA
165	H44A(+85 + 104)	UUU GUG UCU UUC UGA GAA AC
166	H44D(+10 - 10)	AAA GAC UUA CCU UAA GAU AC
167	H44A(-06 + 14)	AUC UGU CAA AUC GCC UGC AG
168	H46D(+16 - 04)	UUA CCU UGA CUU GCU CAA GC
169	H46A(+90 + 109)	UCC AGG UUC AAG UGG GAU AC
170	H47A(+76 + 100)	GCU CUU CUG GGC UUA UGG GAG CAC U
171	H47D(+25 - 02)	ACC UUU AUC CAC UGG AGA UUU GUC UGC
172	H47A(-9 + 12)	UUC CAC CAG UAA CUG AAA CAG
173	H50A(+02 + 30)	CCA CUC AGA GCU CAG AUC UUC UAA CUU CC
174	H50A(+07 + 33)	CUU CCA CUC AGA GCU CAG AUC UUC UAA
175	H50D(+07 - 18)	GGG AUC CAG UAU ACU UAC AGG CUC C
176	H51A(-01 + 25)	ACC AGA GUA ACA GUC UGA GUA GGA GC
177	H51D(+16 - 07)	CUC AUA CCU UCU GCU UGA UGA UC
178	H51A(+111 + 134)	UUC UGU CCA AGC CCG GUU GAA AUC
179	H51A(+61 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA GUU UGG
180	H51A(+66 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA G
181	H51A(+66 + 95)	CUC CAA CAU CAA GGA AGA UGG CAU UUC UAG
182	H51D(+08 - 17)	AUC AUU UUU UCU CAU ACC UUC UGC U
183	H51A/D(+08 - 17) & (-15+)	AUC AUU UUU UCU CAU ACC UUC UGC UAG GAG CUA AAA

TABLE 1A-continued

Description of 2'-O-methyl phosphorothioate antisense oligonucleotides
that have been used to date to study induced exon
skipping during the processing of the dystrophin pre-mRNA. Since
these 2'-O-methyl antisense oligonucleotides are more RNA-like,
U represents uracil. With other antisense chemistries such as peptide
nucleic acids or morpholinos, these U bases may be shown as "T".

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
184	H51A(+175 + 195)	CAC CCA CCA UCA CCC UCU GUG
185	H51A(+199 + 220)	AUC AUC UCG UUG AUA UCC UCA A
186	H52A(-07 + 14)	UCC UGC AUU GUU GCC UGU AAG
187	H52A(+12 + 41)	UCC AAC UGG GGA CGC CUC UGU UCC AAA UCC
188	H52A(+17 + 37)	ACU GGG GAC GCC UCU GUU CCA
189	H52A(+93 + 112)	CCG UAA UGA UUG UUC UAG CC
190	H52D(+05 - 15)	UGU UAA AAA ACU UAC UUC GA
191	H53A(+45 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU G
192	H53A(+39 + 62)	CUG UUG CCU CCG GUU CUG AAG GUG
193	H53A(+39 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU GAA GGU G
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
195	H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C
196	H53A(+150 + 176)	UGU AUA GGG ACC CUC CUU CCA UGA CUC
197	H53D(+20 - 05)	CUA ACC UUG GUU UCU GUG AUU UUC U
198	H53D(+09 - 18)	GGU AUC UUU GAU ACU AAC CUU GGU UUC
199	H53A(-12 + 10)	AUU CUU UCA ACU AGA AUA AAA G
200	H53A(-07 + 18)	GAU UCU GAA UUC UUU CAA CUA GAA U
201	H53A(+07 + 26)	AUC CCA CUG AUU CUG AAU UC
202	H53A(+124 + 145)	UUG GCU CUG GCC UGU CCU AAG A
203	H46A(+86 + 115)	CUC UUU UCC AGG UUC AAG UGG GAU ACU AGC
204	H46A(+107 + 137)	CAA GCU UUU CUU UUA GUU GCU GCU CUU UUC C
205	H46A(-10 + 20)	UAU UCU UUU GUU CUU CUA GCC UGG AGA AAG
206	H46A(+50 + 77)	CUG CUU CCU CCA ACC AUA AAA CAA AUU C
207	H45A(-06 + 20)	CCA AUG CCA UCC UGG AGU UCC UGU AA
208	H45A(+91 + 110)	UCC UGU AGA AUA CUG GCA UC
209	H45A(+125 + 151)	UGC AGA CCU CCU GCC ACC GCA GAU UCA
210	H45D(+16 - 04)	CUA CCU CUU UUU UCU GUC UG
211	H45A(+71 + 90)	UGU UUU UGA GGA UUG CUG AA

TABLE 1B

Description of a cocktail of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU
82	H20A (+147 + 168)	GUU C CAG CAG UAG UUG UCA UCU GCU C
80	H19A(+35 + 65)	GCC UGA GCU GAU CUG CUG GCA UCU
81	H20A(+44 + 63)	UGC
82	H20A (+147 + 168)	AGU U CUG GCA GAA UUC GAU CCA CCG GCU

TABLE 1B-continued

Description of a cocktail of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
		GUU C CAG CAG UAG UUG UCA UCU GCU C
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
195	H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C
196	H53A (+150 + 175)	UGU AUA GGG ACC CUC CUU CCA UGA CUC

TABLE 1C

Description of a "weasel" of 2'-O-methyl phosphorothioate antisense oligonucleotides that have been used to date to study induced exon skipping during the processing of the dystrophin pre-mRNA.

SEQ ID	SEQUENCE	NUCLEOTIDE SEQUENCE (5'-3')
81	H20A(+44 + 71) -	CUG GCA GAA UUC GAU CCA CCG GCU GUU C-
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C
80	H19A(+35 + 65) -	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
88	H20A(+44 + 63) -	-AUU CGA UCC ACC GGC UGU UC-
79	H20A(+149 + 168)	CUG CUG GCA UCU UGC AGU U
80	H19A(+35 + 65) -	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
88	H20A(+44 + 63)	-AUU CGA UCC ACC GGC UGU UC-
80	H19A(+35 + 65) -	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U
79	H20A(+149 + 168)	-CUG CUG GCA UCU UGC AGU U
138	H34A(+46 + 70) -	CAU UCA UUU CCU UUC GCA UCU UAC G-
139	H34A(+94 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG
124	H31D(+03 - 22) - UU-	UAG UUU CUG AAA UAA CAU AUA CCU G- UU-
144	H35A(+24 + 43)	UCU UCA GGU GCA CCU UCU GU
195	H53A(+23 + 47) - AA-	CUG AAG GUG UUC UUG UAC UUC AUC C- AA-
196	H53A(+150 + 175) - AA-	UGU AUA GGG ACC CUC CUU CCA UGA CUC- AA-
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA
-212	Aimed at exons 19/20/20	CAG CAG UAG UUG UCA UCU GCU CAA CUG GCA GAA UUC GAU CCA CCG GCU GUU CAA GCC UGA GCU GAU CUG CUC GCA UCU UGC AGU

DETAILED DESCRIPTION OF THE INVENTION

General

[0055] Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variation and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in the specification, individually or collectively and any and all combinations or any two or more of the steps or features.

[0056] The present invention is not to be limited in scope by the specific embodiments described herein, which are intended for the purpose of exemplification only. Functionally equivalent products, compositions and methods are clearly within the scope of the invention as described herein.

[0057] Sequence identity numbers (SEQ ID NO:) containing nucleotide and amino acid sequence information included in this specification are collected at the end of the description and have been prepared using the programme Patentln Version 3.0. Each nucleotide or amino acid sequence is identified in the sequence listing by the numeric indicator <210> followed by the sequence identifier (e.g. <210>1, <210>2, etc.). The length, type of sequence and source organism for each nucleotide or amino acid sequence are indicated by information provided in the numeric indicator fields <211>, <212> and <213>, respectively. Nucleotide and amino acid sequences referred to in the specification are defined by the information provided in numeric indicator field <400> followed by the sequence identifier (e.g. <400>1, <400>2, etc.).

[0058] An antisense molecules nomenclature system was proposed and published to distinguish between the different antisense molecules (see Mann et al., (2002) *J Gen Med* 4, 644-654). This nomenclature became especially relevant when testing several slightly different antisense molecules, all directed at the same target region, as shown below:

H#A/D(x;y).

[0059] The first letter designates the species (e.g. H: human, M: murine, C: canine) “#” designates target dystrophin exon number.

[0060] “A/D” indicates acceptor or donor splice site at the beginning and end of the exon, respectively.

[0061] (x y) represents the annealing coordinates where “-” or “+” indicate intronic or exonic sequences respectively. As an example, A(-6+18) would indicate the last 6 bases of the intron preceding the target exon and the first 18 bases of the target exon. The closest splice site would be the acceptor so these coordinates would be preceded with an “A”. Describing annealing coordinates at the donor splice site could be D(+2-18) where the last 2 exonic bases and the first 18 intronic bases correspond to the annealing site of the antisense molecule. Entirely exonic annealing coordinates that would be represented by A(+65+85), that is the site between the 65th and 85th nucleotide from the start of that exon.

[0062] The entire disclosures of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference. No admission is made that any of the references constitute prior art or are part of the common general knowledge of those working in the field to which this invention relates.

[0063] As used necessarily herein the term “derived” and “derived from” shall be taken to indicate that a specific integer may be obtained from a particular source albeit not directly from that source.

[0064] Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

[0065] Other definitions for selected terms used herein may be found within the detailed description of the invention and apply throughout. Unless otherwise defined, all other scientific and technical terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which the invention belongs.

DESCRIPTION OF THE PREFERRED EMBODIMENT

[0066] When antisense molecule(s) are targeted to nucleotide sequences involved in splicing in exons within pre-mRNA sequences, normal splicing of the exon may be inhibited causing the splicing machinery to by-pass the entire mutated exon from the mature mRNA. The concept of antisense oligonucleotide induced exon skipping is shown in FIG. 2. In many genes, deletion of an entire exon would lead to the production of a non-functional protein through the loss of important functional domains or the disruption of the reading frame. In some proteins, however, it is possible to shorten the protein by deleting one or more exons, without disrupting the reading frame, from within the protein without seriously altering the biological activity of the protein. Typically, such proteins have a structural role and/or possess functional domains at their ends. The present invention describes antisense molecules capable of binding to specified dystrophin pre-mRNA targets and re-directing processing of that gene.

Antisense Molecules

[0067] According to a first aspect of the invention, there is provided antisense molecules capable of binding to a selected target to induce exon skipping. To induce exon skipping in exons of the Dystrophin gene transcript, the antisense molecules are preferably selected from the group of compounds shown in Table 1A. There is also provided a combination or “cocktail” of two or more antisense oligonucleotides capable of binding to a selected target to induce exon skipping. To induce exon skipping in exons of the Dystrophin gene transcript, the antisense molecules in a “cocktail” are preferably selected from the group of compounds shown in Table 1B. Alternatively, exon skipping may be induced by antisense oligonucleotides joined together “weasels” preferably selected from the group of compounds shown in Table 1C.

[0068] Designing antisense molecules to completely mask consensus splice sites may not necessarily generate any skipping of the targeted exon. Furthermore, the inventors have discovered that size or length of the antisense oligonucleotide itself is not always a primary factor when designing antisense molecules. With some targets such as exon 19, antisense oligonucleotides as short as 12 bases were able to induce exon skipping, albeit not as efficiently as longer (20-31 bases) oligonucleotides. In some other targets, such as murine dystrophin exon 23, antisense oligonucleotides only 17 residues long were able to induce more efficient skipping than another overlapping compound of 25 nucleotides.

[0069] The inventors have also discovered that there does not appear to be any standard motif that can be blocked or masked by antisense molecules to redirect splicing. In some exons, such as mouse dystrophin exon 23, the donor splice site was the most amenable to target to re-direct skipping of that exon. It should be noted that designing and testing a series of exon 23 specific antisense molecules to anneal to overlapping regions of the donor splice site showed considerable variation in the efficacy of induced exon skipping. As reported in Mann et al., (2002) there was a significant variation in the efficiency of bypassing the nonsense mutation depending upon antisense oligonucleotide annealing (“Improved antisense oligonucleotide induced exon skipping in the mdx mouse model of muscular dystrophy”. *J Gen Med* 4: 644-654). Targeting the acceptor site of exon 23 or several internal domains was not found to induce any consistent exon 23 skipping.

[0070] In other exons targeted for removal, masking the donor splice site did not induce any exon skipping. However, by directing antisense molecules to the acceptor splice site (human exon 8 as discussed below), strong and sustained exon skipping was induced. It should be noted that removal of human exon 8 was tightly linked with the co-removal of exon 9. There is no strong sequence homology between the exon 8 antisense oligonucleotides and corresponding regions of exon 9 so it does not appear to be a matter of cross reaction. Rather the splicing of these two exons is inextricably linked. This is not an isolated instance as the same effect is observed in canine cells where targeting exon 8 for removal also resulted in the skipping of exon 9. Targeting exon 23 for removal in the mouse dystrophin pre-mRNA also results in the frequent removal of exon 22 as well. This effect occurs in a dose dependent manner and also indicates close coordinated processing of 2 adjacent exons.

[0071] In other targeted exons, antisense molecules directed at the donor or acceptor splice sites did not induce exon skipping while annealing antisense molecules to intron regions (i.e. exon splicing enhancers within human dystrophin exon 6) was most efficient at inducing exon skipping. Some exons, both mouse and human exon 19 for example, are readily skipped by targeting antisense molecules to a variety of motifs. That is, targeted exon skipping is induced after using antisense oligonucleotides to mask donor and acceptor splice sites or exon splicing enhancers.

[0072] To identify and select antisense oligonucleotides suitable for use in the modulation of exon skipping, a nucleic acid sequence whose function is to be modulated must first be identified. This may be, for example, a gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. Within the context of the present invention, preferred target site(s) are those involved in mRNA splicing (i.e. splice donor sites, splice acceptor sites, or exon splicing enhancer elements). Splicing branch points and exon recognition sequences or splice enhancers are also potential target sites for modulation of mRNA splicing.

[0073] Preferably, the present invention aims to provide antisense molecules capable of binding to a selected target in the dystrophin pre-mRNA to induce efficient and consistent exon skipping. Duchenne muscular dystrophy arises from mutations that preclude the synthesis of a functional dystrophin gene product. These Duchenne muscular dystrophy gene defects are typically nonsense mutations or genomic rearrangements such as deletions, duplications or micro-dele-

tions or insertions that disrupt the reading frame. As the human dystrophin gene is a large and complex gene with the 79 exons being spliced together to generate a mature mRNA with an open reading frame of approximately 11,000 bases, there are many positions where these mutations can occur. Consequently, a comprehensive antisense oligonucleotide based therapy to address many of the different disease-causing mutations in the dystrophin gene will require that many exons can be targeted for removal during the splicing process.

[0074] Within the context of the present invention, preferred target site(s) are those involved in mRNA splicing (i.e. splice donor sites, splice acceptor sites or exon splicing enhancer elements). Splicing branch points and exon recognition sequences or splice enhancers are also potential target sites for modulation of mRNA splicing.

[0075] The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, “specifically hybridisable” and “complementary” are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense molecule need not be 100% complementary to that of its target sequence to be specifically hybridisable. An antisense molecule is specifically hybridisable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

[0076] While the above method may be used to select antisense molecules capable of deleting any exon from within a protein that is capable of being shortened without affecting its biological function, the exon deletion should not lead to a reading frame shift in the shortened transcribed mRNA. Thus, if in a linear sequence of three exons the end of the first exon encodes two of three nucleotides in a codon and the next exon is deleted then the third exon in the linear sequence must start with a single nucleotide that is capable of completing the nucleotide triplet for a codon. If the third exon does not commence with a single nucleotide there will be a reading frame shift that would lead to the generation of truncated or a non-functional protein.

[0077] It will be appreciated that the codon arrangements at the end of exons in structural proteins may not always break at the end of a codon, consequently there may be a need to delete more than one exon from the pre-mRNA to ensure in-frame reading of the mRNA. In such circumstances, a plurality of antisense oligonucleotides may need to be selected by the method of the invention wherein each is directed to a different region responsible for inducing splicing in the exons that are to be deleted.

[0078] The length of an antisense molecule may vary so long as it is capable of binding selectively to the intended location within the pre-mRNA molecule. The length of such sequences can be determined in accordance with selection procedures described herein. Generally, the antisense molecule will be from about 10 nucleotides in length up to about

50 nucleotides in length. It will be appreciated however that any length of nucleotides within this range may be used in the method. Preferably, the length of the antisense molecule is between 17 to 30 nucleotides in length.

[0079] In order to determine which exons can be connected in a dystrophin gene, reference should be made to an exon boundary map. Connection of one exon with another is based on the exons possessing the same number at the 3' border as is present at the 5' border of the exon to which it is being connected. Therefore, if exon 7 were deleted, exon 6 must connect to either exons 12 or 18 to maintain the reading frame. Thus, antisense oligonucleotides would need to be selected which redirected splicing for exons 7 to 11 in the first instance or exons 7 to 17 in the second instance. Another and somewhat simpler approach to restore the reading frame around an exon 7 deletion would be to remove the two flanking exons. Induction of exons 6 and 8 skipping should result in an in-frame transcript with the splicing of exons 5 to 9. In practise however, targeting exon 8 for removal from the pre-mRNA results in the co-removal of exon 9 so the resultant transcript would have exon 5 joined to exon 10. The inclusion or exclusion of exon 9 does not alter the reading frame. Once the antisense molecules to be tested have been identified, they are prepared according to standard techniques known in the art. The most common method for producing antisense molecules is the methylation of the 2' hydroxyribose position and the incorporation of a phosphorothioate backbone produces molecules that superficially resemble RNA but that are much more resistant to nuclease degradation.

[0080] To avoid degradation of pre-mRNA during duplex formation with the antisense molecules, the antisense molecules used in the method may be adapted to minimise or prevent cleavage by endogenous RNase H. This property is highly preferred as the treatment of the RNA with the unmethylated oligonucleotides either intracellularly or in crude extracts that contain RNase H leads to degradation of the pre-mRNA: antisense oligonucleotide duplexes. Any form of modified antisense molecules that is capable of by-passing or not inducing such degradation may be used in the present method. An example of antisense molecules which when duplexed with RNA are not cleaved by cellular RNase H is 2'-O-methyl derivatives. 2'-O-methyl-oligoribonucleotides are very stable in a cellular environment and in animal tissues, and their duplexes with RNA have higher T_m values than their ribo- or deoxyribo-counterparts.

[0081] Antisense molecules that do not activate RNase H can be made in accordance with known techniques (see, e.g., U.S. Pat. No. 5,149,797). Such antisense molecules, which may be deoxyribonucleotide or ribonucleotide sequences, simply contain any structural modification which sterically hinders or prevents binding of RNase H to a duplex molecule containing the oligonucleotide as one member thereof, which structural modification does not substantially hinder or disrupt duplex formation. Because the portions of the oligonucleotide involved in duplex formation are substantially different from those portions involved in RNase H binding thereto, numerous antisense molecules that do not activate RNase H are available. For example, such antisense molecules may be oligonucleotides wherein at least one, or all, of the inter-nucleotide bridging phosphate residues are modified phosphates, such as methyl phosphonates, methyl phosphorothioates, phosphoromorpholidates, phosphoropiperazidates and phosphoramidates. For example, every other one of the internucleotide bridging phosphate residues may be

modified as described. In another non-limiting example, such antisense molecules are molecules wherein at least one, or all, of the nucleotides contain a 2' lower alkyl moiety (e.g., C₁-C₄, linear or branched, saturated or unsaturated alkyl, such as methyl, ethyl, ethenyl, propyl, 1-propenyl, 2-propenyl, and isopropyl). For example, every other one of the nucleotides may be modified as described.

[0082] While antisense oligonucleotides are a preferred form of the antisense molecules, the present invention comprehends other oligomeric antisense molecules, including but not limited to oligonucleotide mimetics such as are described below.

[0083] Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural inter-nucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their inter-nucleoside backbone can also be considered to be oligonucleosides.

[0084] In other preferred oligonucleotide mimetics, both the sugar and the inter-nucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone.

[0085] Modified oligonucleotides may also contain one or more substituted sugar moieties. Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. Certain nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2° C. and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0086] Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates that enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-S-tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., dihexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-oxysterol moiety.

[0087] It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of

the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds that are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this invention, are antisense molecules, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the increased resistance to nuclease degradation, increased cellular uptake, and an additional region for increased binding affinity for the target nucleic acid.

Methods of Manufacturing Antisense Molecules

[0088] The antisense molecules used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, Calif.). One method for synthesising oligonucleotides on a modified solid support is described in U.S. Pat. No. 4,458,066.

[0089] Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates ~ and alkylated derivatives. In one such automated embodiment, diethyl-phosphoramidites are used as starting materials and may be synthesized as described by Beaucage, et al., (1981) *Tetrahedron Letters*, 22:1859-1862.

[0090] The antisense molecules of the invention are synthesised in vitro and do not include antisense compositions of biological origin, or genetic vector constructs designed to direct the in vivo synthesis of antisense molecules. The molecules of the invention may also be mixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption.

Therapeutic Agents

[0091] The present invention also can be used as a prophylactic or therapeutic, which may be utilised for the purpose of treatment of a genetic disease.

[0092] Accordingly, in one embodiment the present invention provides antisense molecules that bind to a selected target in the dystrophin pre-mRNA to induce efficient and consistent exon skipping described herein in a therapeutically effective amount admixed with a pharmaceutically acceptable carrier, diluent, or excipient.

[0093] The phrase "pharmaceutically acceptable" refers to molecular entities and compositions that are physiologically tolerable and do not typically produce an allergic or similarly untoward reaction, such as gastric upset and the like, when administered to a patient. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the compound is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water or

saline solutions and aqueous dextrose and glycerol solutions are preferably employed as carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in Martin, *Remington's Pharmaceutical Sciences*, 18th Ed., Mack Publishing Co., Easton, Pa., (1990).

[0094] In a more specific form of the invention there are provided pharmaceutical compositions comprising therapeutically effective amounts of an antisense molecule together with pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or carriers. Such compositions include diluents of various buffer content (e.g., Tris-HCl, acetate, phosphate), pH and ionic strength and additives such as detergents and solubilizing agents (e.g., Tween 80, Polysorbate 80), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimersol, benzyl alcohol) and bulking substances (e.g., lactose, mannitol). The material may be incorporated into particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Hylauronic acid may also be used. Such compositions may influence the physical state, stability, rate of in vivo release, and rate of in vivo clearance of the present proteins and derivatives. See, e.g., Martin, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton, Pa. 18042) pages 1435-1712 that are herein incorporated by reference. The compositions may be prepared in liquid form, or may be in dried powder, such as lyophilised form.

[0095] It will be appreciated that pharmaceutical compositions provided according to the present invention may be administered by any means known in the art. Preferably, the pharmaceutical compositions for administration are administered by injection, orally, or by the pulmonary, or nasal route. The antisense molecules are more preferably delivered by intravenous, intra-arterial, intraperitoneal, intramuscular, or subcutaneous routes of administration.

Antisense Molecule Based Therapy

[0096] Also addressed by the present invention is the use of antisense molecules of the present invention, for manufacture of a medicament for modulation of a genetic disease.

[0097] The delivery of a therapeutically useful amount of antisense molecules may be achieved by methods previously published. For example, intracellular delivery of the antisense molecule may be via a composition comprising an admixture of the antisense molecule and an effective amount of a block copolymer. An example of this method is described in US patent application US 20040248833.

[0098] Other methods of delivery of antisense molecules to the nucleus are described in Mann C J et al., (2001) [*Antisense-induced exon skipping and the synthesis of dystrophin in the mdx mouse*]. Proc., Natl. Acad. Science, 98(1) 42-47J and in GebSKI et al., (2003). *Human Molecular Genetics*, 12(15): 1801-1811.

[0099] A method for introducing a nucleic acid molecule into a cell by way of an expression vector either as naked DNA or complexed to lipid carriers, is described in U.S. Pat. No. 6,806,084.

[0100] It may be desirable to deliver the antisense molecule in a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes or liposome formulations.

[0101] Liposomes are artificial membrane vesicles which are useful as delivery vehicles *in vitro* and *in vivo*. These formulations may have net cationic, anionic or neutral charge characteristics and are useful characteristics with *in vitro*, *in vivo* and *ex vivo* delivery methods. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, and DNA can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, et al., *Trends Biochem. Sci.*, 6:77, 1981).

[0102] In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the antisense molecule of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., *Biotechniques*, 6:682, 1988).

[0103] The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

[0104] Alternatively, the antisense construct may be combined with other pharmaceutically acceptable carriers or diluents to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may be formulated for parenteral, intramuscular, intravenous, subcutaneous, intraocular, oral or transdermal administration.

[0105] The routes of administration described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and any dosage for any particular animal and condition. Multiple approaches for introducing functional new genetic material into cells, both *in vitro* and *in vivo* have been attempted (Friedmann (1989) *Science*, 244:1275-1280).

[0106] These approaches include integration of the gene to be expressed into modified retroviruses (Friedmann (1989) *supra*; Rosenberg (1991) *Cancer Research* 51(18), suppl.: 5074S-5079S); integration into non-retrovirus vectors (Rosenfeld, et al. (1992) *Cell*, 68:143-155; Rosenfeld, et al. (1991) *Science*, 252:431-434); or delivery of a transgene linked to a heterologous promoter-enhancer element via liposomes (Friedmann (1989), *supra*; Brigham, et al. (1989) *Am. J. Med. Sci.*, 298:278-281; Nabel, et al. (1990) *Science*, 249:1285-1288; Hazinski, et al. (1991) *Am. J. Resp. Cell Molec. Biol.*, 4:206-209; and Wang and Huang (1987) *Proc. Natl. Acad. Sci. (USA)*, 84:7851-7855); coupled to ligand-specific, cation-based transport systems (Wu and Wu (1988) *J. Biol. Chem.*, 263:14621-14624) or the use of naked DNA, expression vectors (Nabel et al. (1990), *supra*); Wolff et al. (1990) *Science*, 247:1465-1468). Direct injection of transgenes into tissue produces only localized expression (Rosenfeld (1992) *supra*); Rosenfeld et al. (1991) *supra*; Brigham et al. (1989) *supra*; Nabel (1990) *supra*; and Hazinski et al. (1991) *supra*). The Brigham et al. group (*Am. J. Med. Sci.* (1989) 298:278-281 and *Clinical Research* (1991) 39 (abstract)) have reported *in vivo* transfection only of lungs of mice following either

intravenous or intratracheal administration of a DNA liposome complex. An example of a review article of human gene therapy procedures is: Anderson, *Science* (1992) 256:808-813.

[0107] The antisense molecules of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such pro-drugs, and other bioequivalents.

[0108] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

[0109] For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, malefic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, (including by nebulizer, intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intra-arterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

[0110] The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Kits of the Invention

[0111] The invention also provides kits for treatment of a patient with a genetic disease which kit comprises at least an antisense molecule, packaged in a suitable container, together with instructions for its use.

[0112] In a preferred embodiment, the kits will contain at least one antisense molecule as shown in Table 1A, or a cocktail of antisense molecules as shown in Table 1B or a "weasel" compound as shown in Table 1C. The kits may also contain peripheral reagents such as buffers, stabilizers, etc.

[0113] Those of ordinary skill in the field should appreciate that applications of the above method has wide application for identifying antisense molecules suitable for use in the treatment of many other diseases.

Examples

[0114] The following Examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these Examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. The references cited herein are expressly incorporated by reference.

[0115] Methods of molecular cloning, immunology and protein chemistry, which are not explicitly described in the following examples, are reported in the literature and are known by those skilled in the art. General texts that described conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art, included, for example: Sambrook et al, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Glover ed., *DNA Cloning: A Practical Approach*, Volumes I and II, MRL Press, Ltd., Oxford, U.K. (1985); and Ausubel, F., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A., Struhl, K. *Current Protocols in Molecular Biology*. Greene Publishing Associates/Wiley Intersciences, New York (2002).

Determining Induced Exon Skipping in Human Muscle Cells

[0116] Attempts by the inventors to develop a rational approach in antisense molecules design were not completely successful as there did not appear to be a consistent trend that could be applied to all exons. As such, the identification of the most effective and therefore most therapeutic antisense molecules compounds has been the result of empirical studies.

[0117] These empirical studies involved the use of computer programs to identify motifs potentially involved in the splicing process. Other computer programs were also used to identify regions of the pre-mRNA which may not have had extensive secondary structure and therefore potential sites for annealing of antisense molecules. Neither of these approaches proved completely reliable in designing antisense oligonucleotides for reliable and efficient induction of exon skipping.

[0118] Annealing sites on the human dystrophin pre-mRNA were selected for examination, initially based upon known or predicted motifs or regions involved in splicing. 2OMe antisense oligonucleotides were designed to be complementary to the target sequences under investigation and were synthesised on an Expedite 8909 Nucleic Acid Synthesiser. Upon completion of synthesis, the oligonucleotides were cleaved from the support column and de-protected in ammonium hydroxide before being desalted. The quality of the oligonucleotide synthesis was monitored by the intensity of the trityl signals upon each deprotection step during the synthesis as detected in the synthesis log. The

concentration of the antisense oligonucleotide was estimated by measuring the absorbance of a diluted aliquot at 260 nm.

[0119] Specified amounts of the antisense molecules were then tested for their ability to induce exon skipping in an in vitro assay, as described below.

[0120] Briefly, normal primary myoblast cultures were prepared from human muscle biopsies obtained after informed consent. The cells were propagated and allowed to differentiate into myotubes using standard culturing techniques. The cells were then transfected with the antisense oligonucleotides by delivery of the oligonucleotides to the cells as cationic lipoplexes, mixtures of antisense molecules or cationic liposome preparations.

[0121] The cells were then allowed to grow for another 24 hours, after which total RNA was extracted and molecular analysis commenced. Reverse transcriptase amplification (RT-PCR) was undertaken to study the targeted regions of the dystrophin pre-mRNA or induced exonic re-arrangements.

[0122] For example, in the testing of an antisense molecule for inducing exon 19 skipping the RT-PCR test scanned several exons to detect involvement of any adjacent exons. For example, when inducing skipping of exon 19, RT-PCR was carried out with primers that amplified across exons 17 and 21. Amplifications of even larger products in this area (i.e. exons 13-26) were also carried out to ensure that there was minimal amplification bias for the shorter induced skipped transcript. Shorter or exon skipped products tend to be amplified more efficiently and may bias the estimated of the normal and induced transcript.

[0123] The sizes of the amplification reaction products were estimated on an agarose gel and compared against appropriate size standards. The final confirmation of identity of these products was carried out by direct DNA sequencing to establish that the correct or expected exon junctions have been maintained.

[0124] Once efficient exon skipping had been induced with one antisense molecule, subsequent overlapping antisense molecules may be synthesized and then evaluated in the assay as described above. Our definition of an efficient antisense molecule is one that induces strong and sustained exon skipping at transfection concentrations in the order of 300 nM or less.

Antisense Oligonucleotides Directed at Exon 8

[0125] Antisense oligonucleotides directed at exon 8 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0126] FIG. 3 shows differing efficiencies of two antisense molecules directed at exon 8 acceptor splice site. H8A(-06+18) [SEQ ID NO:1], which anneals to the last 6 bases of intron 7 and the first 18 bases of exon 8, induces substantial exon 8 and 9 skipping when delivered into cells at a concentration of 20 nM. The shorter antisense molecule, H8A(-06+14) [SEQ ID NO: 4] was only able to induce exon 8 and 9 skipping at 300 nM, a concentration some 15 fold higher than H8A(-06+18), which is the preferred antisense molecule.

[0127] This data shows that some particular antisense molecules induce efficient exon skipping while another antisense molecule, which targets a near-by or overlapping region, can be much less efficient. Titration studies show one compound is able to induce targeted exon skipping at 20 nM while the less efficient antisense molecules only induced exon skipping at concentrations of 300 nM and above. Therefore, we have

shown that targeting of the antisense molecules to motifs involved in the splicing process plays a crucial role in the overall efficacy of that compound.

[0128] Efficacy refers to the ability to induce consistent skipping of a target exon.

[0129] However, sometimes skipping of the target exons is consistently associated with a flanking exon. That is, we have found that the splicing of some exons is tightly linked. For example, in targeting exon 23 in the mouse model of muscular dystrophy with antisense molecules directed at the donor site of that exon, dystrophin transcripts missing exons 22 and 23 are frequently detected. As another example, when using an antisense molecule directed to exon 8 of the human dystrophin gene, all induced transcripts are missing both exons 8 and 9. Dystrophin transcripts missing only exon 8 are not observed.

[0130] Table 2 below discloses antisense molecule sequences that induce exon 8 (and 9) skipping.

TABLE 2

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
1	H8A(-06 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG UAA	Very strong to 20 nM
2	H8A (-03 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG	Very strong skipping to 40 nM
3	H8A(-07 + 18)	5'-GAU AGG UGG UAU CAA CAU CUG UAA G	Strong skipping to 40 nM
4	H8A(-06 + 14)	5'-GGU GGU AUC AAC AUC UGU AA	Skipping to 300 nM
5	H8A(-10 + 10)	5'-GUA UCA ACA UCU GUA AGC AC	Patchy/weak skipping to 100 nm

Antisense Oligonucleotides Directed at Exon 7

[0131] Antisense oligonucleotides directed at exon 7 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0132] FIG. 4 shows the preferred antisense molecule, H7A (+45+67) [SEQ ID NO: 6], and another antisense molecule, H7A(+2+26) [SEQ ID NO: 7], inducing exon 7 skipping. Nested amplification products span exons 3 to 9. Additional products above the induced transcript missing exon 7 arise from amplification from carry-over outer primers from the RT-PCR as well as heteroduplex formation.

[0133] Table 3 below discloses antisense molecule sequences for induced exon 7 skipping.

TABLE 3

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
6	H7A(+45 + 67)	5'-UGC AUG UUC CAG UCG UUG UGU GG	Strong skipping to 20 nM
7	H7A(+02 + 26)	5'-CAC UAU UCC AGU CAA AUA GGU CUG G	Weak skipping at 100 nM
8	H7D(+15 - 10)	5'-AUU UAC CAA CCU UCA GGA UCG AGU A	Weak skipping to 300 nM
9	H7A(-18 + 03)	5'-GGC CUA AAA CAC AUA CAC AUA	Weak skipping to 300 nM

Antisense Oligonucleotides Directed at Exon 6

[0134] Antisense oligonucleotides directed at exon 6 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0135] FIG. 5 shows an example of two non-preferred antisense molecules inducing very low levels of exon 6 skipping in cultured human cells. Targeting this exon for specific removal was first undertaken during a study of the canine model using the oligonucleotides as listed in Table 4, below. Some of the human specific oligonucleotides were also evaluated, as shown in FIG. 5. In this example, both antisense molecules target the donor splice site and only induced low levels of exon 6 skipping. Both H6D(+4-21) [SEQ ID NO: 17] and H6D(+18-4) [SEQ ID NO: 18] would be regarded as non-preferred antisense molecules.

[0136] One antisense oligonucleotide that induced very efficient exon 6 skipping in the canine model, C6A(+69+91) [SEQ ID NO: 14], would anneal perfectly to the corresponding region in human dystrophin exon 6. This compound was evaluated, found to be highly efficient at inducing skipping of that target exon, as shown in FIG. 6 and is regarded as the preferred compound for induced exon 6 skipping. Table 4 below discloses antisense molecule sequences for induced exon 6 skipping.

Antisense Oligonucleotides Directed at Exon 4

[0137] Antisense oligonucleotides directed at exon 4 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0138] FIG. 7 shows an example of a preferred antisense molecule inducing skipping of exon 4 skipping in cultured human cells. In this example, one preferred antisense compound, H4A(+13+32) [SEQ ID NO:19], which targeted a presumed exonic splicing enhancer induced efficient exon skipping at a concentration of 20 nM while other non-preferred antisense oligonucleotides failed to induce even low levels of exon 4 skipping. Another preferred antisense molecule inducing skipping of exon 4 was H4A(+111+40) [SEQ ID NO:22], which induced efficient exon skipping at a concentration of 20 nM.

[0139] Table 5 below discloses antisense molecule sequences for inducing exon 4 skipping.

TABLE 4

SEQ ID	Antisense Oligo name	Sequence	Ability to induce skipping
10	C6A(-10 + 10)	5' CAU UUU UGA CCU ACA UGU GG	No skipping
11	C6A(-14 + 06)	5' UUU GAC CUA CAU GUG GAA AG	No skipping
12	C6A(-14 + 12)	5' UAC AUU UUU GAC CUA CAU GUG GAA AG	No skipping
13	C6A(-13 + 09)	5' AUU UUU GAC CUA CAU GGG AAA G	No skipping
14	CH6A(+69 + 91)	5' UAC GAG UUG AUU GUC GGA CCC AG	Strong skipping to 20 nM
15	C6D(+12 - 13)	5' GUG GUC UCC UUA CCU AUG ACU GUG G	Weak skipping at 300 nM
16	C6D(+06 - 11)	5' GGU CUC CUU ACC UAU GA	No skipping
17	H6D(+04 - 21)	5' UGU CUC AGU AAU CUU CUU ACC UAU	Weak skipping to 50 nM
18	H6D(+18 - 04)	5' UCU UAC CUA UGA CUA UGG AUG AGA	Very weak skipping to 300 nM

TABLE 5

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
19	H4A(+13 + 32)	5' GCA UGA ACU CUU GUG GAU CC	Skipping to 20 nM
22	H4A(+11 + 40)	5' UGU UCA GGG CAU GAA CUC UUG UGG AUC CUU	Skipping to 20 nM
20	H4D(+04 - 16)	5' CCA GGG UAC UAC UUA CAU UA	No skipping
21	H4D(-24 - 44)	5' AUC GUG UGU CAC AGC AUC CAG	No skipping

Antisense Oligonucleotides Directed at Exon 3

[0140] Antisense oligonucleotides directed at exon 3 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0141] H3A(+30+60) [SEQ ID NO:23] induced substantial exon 3 skipping when delivered into cells at a concentration of 20 nM to 600 nM. The antisense molecule, H3A(+35+65) [SEQ ID NO: 24] induced exon skipping at 300 nM.

[0142] Table 6 below discloses antisense molecule sequences that induce exon 3 skipping.

Antisense Oligonucleotides Directed at Exon 5

[0143] Antisense oligonucleotides directed at exon 5 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0144] H5A(+20+50) [SEQ ID NO:31] induces substantial exon 5 skipping when delivered into cells at a concentration of 100 nM. Table 7 below shows other antisense molecules tested. The majority of these antisense molecules were not as effective at exon skipping as H5A(+20+50). However, H5A(+15+45) [SEQ ID NO: 40] was able to induce exon 5 skipping at 300 nM.

[0145] Table 7 below discloses antisense molecule sequences that induce exon 5 skipping.

TABLE 6

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
23	H3A(+30 + 60)	UAG GAG GCG CCU CCC AUC CUG UAG GUC ACU G	Moderate skipping to 20 to 600 nM
24	H3A(+35 + 65)	AGG UCU AGG AGG CGC CUC CCA UCC UGU AGG U	Working to 300 nM
25	H3A(+30 + 54)	GCG CCU CCC AUC CUG UAG GUC ACU G	Moderate 100-600 nM
26	H3D(+46 - 21)	CUU CGA GGA GGU CUA GGA GGC GCC UC	No skipping
27	H3A(+30 + 50)	CUC CCA UCC UGU AGG UCA CUG	Moderate 20-600 nM
28	H3D(+19 - 03)	UAC CAG UUU UUG CCC UGU CAG G	No skipping
29	H3A(-06 + 20)	UCA AUA UGC UGC UGCCA AAC UGA AA	No skipping
30	H3A(+37 + 61)	CUA GGA GGC GCC UCC CAU CCU GUA G	No skipping

TABLE 7

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
31	H5A(+20 + 50)	UUA UGA UUU CCA UCU ACG AUG UCA GUA CUU C	Working to 100 nM

TABLE 7-continued

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
32	H5D(+25 - 05)	CUU ACC UGC CAG UGG AGG AUU AUA UUC CAA A	No skipping
33	H5D(+10 - 15)	CAU CAG GAU UCU UAC CUG CCA GUG G	Inconsistent at 300 nM
34	H5A(+10 + 34)	CGA UGU CAG UAC UUC CAA UAU UCA C	Very weak
35	H5D(-04 - 21)	ACC AUU CAU CAG GAU UCU	No skipping
36	H5D(+16 - 02)	ACC UGC CAG UGG AGG AUU	No skipping
37	H5A(-07 + 20)	CCA AUA UUC ACU AAA UCA ACC UGU UAA	No skipping
38	H5D(+18 - 12)	CAG GAU UCU UAC CUG CCA GUG GAG GAU UAU	No skipping
39	H5A(+05 + 35)	ACG AUG UCA GUA CUU CCA AUA UUC ACU AAA U	No skipping
40	H5A(+15 + 45)	AUU UCC AUC UAC GAU GUC AGU ACU UCC AAU A	Working to 300 nM

Antisense Oligonucleotides Directed at Exon 10

[0146] Antisense oligonucleotides directed at exon 10 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0147] H10A(-05+16) [SEQ ID NO:41] induced substantial exon 10 skipping when delivered into cells. Table 8 below shows other antisense molecules tested. The antisense molecules ability to induce exon skipping was variable. Table 8 below discloses antisense molecule sequences that induce exon 10 skipping.

Antisense Oligonucleotides Directed at Exon 11

[0148] Antisense oligonucleotides directed at exon 11 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0149] FIG. 8B shows an example of H11A(+75+97) [SEQ ID NO:49] antisense molecule inducing exon 11 skipping in cultured human cells. H11A(+75+97) induced substantial exon 11 skipping when delivered into cells at a concentration of 5 nM. Table 9 below shows other antisense molecules tested. The antisense molecules ability to induce exon skipping was observed at 100 nM.

TABLE 8

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
41	H10A(-05 + 16)	CAG GAG CUU CCA AAU GCU GCA	Not tested
42	H10A(-05 + 24)	CUU GUC UUC AGG AGC UUC CAA AUG CUG CA	Not tested
43	H10A(+98 + 119)	UCC UCA GCA GAA AGA AGC CAC G	Not tested
44	H10A(+130 + 149)	UUA GAA AUC UCU CCU UGU GC	No skipping
45	H10A(-33 - 14)	UAA AUU GGG UGU UAC ACA AU	No skipping

TABLE 9

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
46	H11D(+26 + 49)	CCC UGA GGC AUU CCC AUC UUG AAU	Skipping at 100 nM
47	H11D(+11 - 09)	AGG ACU UAC UUG CUU UGU UU	Skipping at 100 nM
48	H11A(+118 + 140)	CUU GAA UUU AGG AGA UUC AUC UG	Skipping at 100 nM
49	H11A(+75 + 97)	CAU CUU CUG AUA AUU UUC CUG UU	Skipping at 100 nM
46	H11D(+26 + 49)	CCC UGA GGC AUU CCC AUC UUG AAU	Skipping at 5 nM

Antisense Oligonucleotides Directed at Exon 12

[0150] Antisense oligonucleotides directed at exon 12 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0151] H12A(+52+75) [SEQ ID NO:50] induced substantial exon 12 skipping when delivered into cells at a concentration of 5 nM, as shown in FIG. 8A. Table 10 below shows other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. The antisense molecules ability to induce exon skipping was variable.

TABLE 10

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
50	H12A(+52 + 75)	UCU UCU GUU UUU GUU AGC CAG UCA	Skipping at 5 nM
51	H12A(-10 + 10)	UCU AUG UAA ACU GAA AAU UU	Skipping at 100 nM
52	H12A(+11 + 30)	UUC UGG AGA UCC AUU AAA AC	No skipping

Antisense Oligonucleotides Directed at Exon 13

[0152] Antisense oligonucleotides directed at exon 13 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0153] H13A(+77+100) [SEQ ID NO:53] induced substantial exon 13 skipping when delivered into cells at a concentration of 5 nM. Table 11 below includes two other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These other antisense molecules were unable to induce exon skipping.

TABLE 11

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
53	H13A(+77 + 100)	CAG CAG UUG CGU GAU CUC CAC UAG	Skipping at 5 nM
54	H13A(+55 + 75)	UUC AUC AAC UAC CAC CAC CAU	No skipping
55	H13D(+06 - 19)	CUA AGC AAA AUA AUC UGA CCU UAA G	No skipping

Antisense Oligonucleotides Directed at Exon 14

[0154] Antisense oligonucleotides directed at exon 14 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0155] H14A(+37+64) [SEQ ID NO:56] induced weak exon 14 skipping when delivered into cells at a concentration of 100 nM. Table 12 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. The other antisense molecules were unable to induce exon skipping at any of the concentrations tested.

TABLE 12

Antisense SEQ ID	Oligonucleotide name	Sequence	Ability to induce skipping
56	H14A(+37 + 64)	CUU GUA AAA GAA CCC AGC GGU CUU CUG U	Skipping at 100 nM
57	H14A(+14 + 35)	CAU CUA CAG AUG UUU GCC CAU C	No skipping
58	H14A(+51 + 73)	GAA GGA UGU CUU GUA AAA GAA CC	No skipping
59	H14D(-02 + 18)	ACC UGU UCU UCA GUA AGA CG	No skipping

TABLE 12-continued

Antisense SEQ ID	Oligonucleotide name	Sequence	Ability to induce skipping
60	H14D(+14 - 10)	CAU GAC ACA CCU GUU CUU CAG UAA	No skipping
61	H14A(+61 + 80)	CAU UUG AGA AGG AUG UCU UG	No skipping
62	H14A(-12 + 12)	AUC UCC CAA UAC CUG GAG AAG AGA	No skipping

Antisense Oligonucleotides Directed at Exon 15

[0156] Antisense oligonucleotides directed at exon 15 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0157] H15A(-12+19) [SEQ ID NO:63] and H15A(+48+71) [SEQ ID NO:64] induced substantial exon 15 skipping when delivered into cells at a concentration of 10 Nm, as shown in FIG. 9A. Table 13 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 Nm. These other antisense molecules were unable to induce exon skipping at any of the concentrations tested.

TABLE 13

Antisense SEQ ID	Oligonucleotide name	Sequence	Ability to induce skipping
63	H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA CAU U	AGASkipping at 5Nm
64	H15A(+48 + 71)	UCU UUA AAG CCA GUU GUG UGA AUC	Skipping at 5 Nm
65	H15A(+08 + 28)	UUU CUG AAA GCC AUG CAC UAA	No skipping
63	H15A(-12 + 19)	GCC AUG CAC UAA AAA GGC ACU GCA CAU U	AGANo skipping
66	H15D(+17 - 08)	GUA CAU ACG GCC AGU UUU UGA AGA C	No skipping

Antisense Oligonucleotides Directed at Exon 16

[0158] Antisense oligonucleotides directed at exon 16 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0159] H16A(-12+19) [SEQ ID NO:67] and H16A(-06+25) [SEQ ID NO:68] induced substantial exon 16 skipping when delivered into cells at a concentration of 10 nM, as shown in FIG. 9B. Table 14 below includes other antisense molecules tested. H16A(-06+19) [SEQ ID NO:69] and H16A(+87+109) [SEQ ID NO:70] were tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These two antisense molecules were able to induce exon skipping at 25 nM and 100 nM, respectively. Additional antisense molecules were tested at 100, 200 and 300 nM and did not result in any exon skipping.

TABLE 14

Antisense SEQ ID name	Oligonucleotide Sequence	Ability to induce skipping
67 H16A(-12 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AAA ACA A	Skipping at 5 nM
68 H16A(-06 + 25)	UCU UUU CUA GAU CCG CUU UUA AAA CCU GUU A	Skipping at 5 nM
69 H16A(-06 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU A	Skipping at 25 nM
70 H16A(+87 + 109)	CCG UCU UCU GGG UCA CUG ACU UA	Skipping at 100 nM
71 H16A(-07 + 19)	CUA GAU CCG CUU UUA AAA CCU GUU AA	No skipping
72 H16A(-07 + 13)	CCG CUU UUA AAA CCU GUU AA	No skipping
73 H16A(+12 + 37)	UGG AUU GCU UUU UCU UUU CUA GAU CC	No skipping
74 H16A(+92 + 116)	CAU GCU UCC GUC UUC UGG GUC ACU G	No skipping
75 H16A(+45 + 67)	G AUC UUG UUU GAG UGA AUA CAG U	No skipping
76 H16A(+105 + 126)	GUU AUC CAG CCA UGC UUC CGU C	No skipping
77 H16D(+05 - 20)	UGA UAA UUG GUA UCA CUA ACC UGU G	No skipping
78 H16D(+12 - 11)	GUA UCA CUA ACC UGU GCU GUA C	No skipping

Antisense Oligonucleotides Directed at Exon 19

[0160] Antisense oligonucleotides directed at exon 19 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0161] H19A(+35+65) [SEQ ID NO:79] induced substantial exon 19 skipping when delivered into cells at a concentration of 10 nM. This antisense molecule also showed very strong exon skipping at concentrations of 25, 50, 100, 300 and 600 nM.

[0162] FIG. 10 illustrates exon 19 and 20 skipping using a “cocktail” of antisense oligonucleotides, as tested using gel electrophoresis. It is interesting to note that it was not easy to induce exon 20 skipping using single antisense oligonucleotides H20A(+44+71) [SEQ ID NO:81] or H20A(+149+170) [SEQ ID NO:82], as illustrated in sections 2 and 3 of the gel shown in FIG. 10. Whereas, a “cocktail” of antisense oligonucleotides was more efficient as can be seen in section 4 of FIG. 10 using a “cocktail” of antisense oligonucleotides H20A(+44+71) and H20A(+149+170). When the cocktail was used to target exon 19, skipping was even stronger (see section 5, FIG. 10).

[0163] FIG. 11 illustrates gel electrophoresis results of exon 19/20 skipping using “weasels” The “weasels” were

effective in skipping exons 19 and 20 at concentrations of 25, 50, 100, 300 and 600 nM. A further “weasel” sequence is shown in the last row of Table 3C. This compound should give good results.

Antisense Oligonucleotides Directed at Exon 20

[0164] Antisense oligonucleotides directed at exon 20 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0165] None of the antisense oligonucleotides tested induced exon 20 skipping when delivered into cells at a concentration of 10, 25, 50, 300 or 600 nM (see Table 15). Antisense molecules H20A(-11+17) [SEQ ID NO:86] and H20D(+08-20) [SEQ ID NO:87] are yet to be tested.

[0166] However, a combination or “cocktail” of H20A(+44+71) [SEQ ID NO: 81] and H20(+149+170) [SEQ ID NO:82] in a ratio of 1:1, exhibited very strong exon skipping at a concentration of 100 nM and 600 nM. Further, a combination of antisense molecules H19A(+35+65) [SEQ ID NO:79], H20A(+44+71) [SEQ ID NO:81] and H20A(+149+170) [SEQ ID NO:82] in a ratio of 2:1:1, induced very strong exon skipping at a concentration ranging from 10 nM to 600 nM.

TABLE 15

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
81	H20A(+44 + 71)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C	No skipping
82	H20A(+147 + 168)	CAG CAG UAG UUG UCA UCU GCU C	No skipping

TABLE 15-continued

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
83	H20A(+185 + 203)	UGA UGG GGU GGU GGG UUG G	No skipping
84	H20A(-08 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA G	No skipping
85	H20A(+30 + 53)	CCG GCU GUU CAG UUG UUC UGA GGC	No skipping
86	H20A(-11 + 17)	AUC UGC AUU AAC ACC CUC UAG AAA GAA A	Not tested yet
87	H20D(+08 - 20)	GAA GGA GAA GAG AUU CUU ACC UUA CAA A	Not tested yet
81 & 82	H20A(+44 + 71) & H20A(+147 + 168)	CUG GCA GAA UUC GAU CCA CCG GCU GUU C CAG CAG UAG UUG UCA UCU GCU C	Very strong skipping
80, 81 & 82	H19A(+35 + 65); H20A(+44 + 71); H20A(+147 + 168)	GCC UGA GCU GAU CUG CUG GCA UCU UGC AGU U; CUG GCA GAA UUC GAU CCA CCG GCU GUU C; CAG CAG UAG UUG UCA UCU GCU C	Very strong skipping

Antisense Oligonucleotides Directed at Exon 21

[0167] Antisense oligonucleotides directed at exon 21 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0168] H21A(+85+108) [SEQ ID NO:92] and H21A(+85+106) [SEQ ID NO:91] induced exon 21 skipping when delivered into cells at a concentration of 50 nM. Table 16 below includes other antisense molecules tested at a concentration range of 5, 25, 50, 100, 200 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping

TABLE 16

SEQ ID	Antisense Oligonucleotide name	Sequence	Ability to induce skipping
90	H21A(-06 + 16)	GCC GGU UGA CUU CAU CCU GUG C	Skips at 600 nM
91	H21A(+85 + 106)	CUG CAU CCA GGA ACA UGG GUC C	Skips at 50 nM
92	H21A(+85 + 108)	GUC UGC AUC CAG GAA CAU GGG UC	Skips at 50 nM
93	H21A(+08 + 31)	GUU GAA GAU CUG AUA GCC GGU UGA	Skips faintly to
94	H21D(+18 - 07)	UAC UUA CUG UCU GUA GCU CUU UCU	No skipping

Antisense Oligonucleotides Directed at Exon 22

[0169] Antisense oligonucleotides directed at exon 22 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0170] FIG. 12 illustrates differing efficiencies of two antisense molecules directed at exon 22 acceptor splice site.

H22A(+125+106) [SEQ ID NO:96] and H22A(+80+101) [SEQ ID NO: 98] induce strong exon 22 skipping from 50 nM to 600 nM concentration.

[0171] H22A(+125+146) [SEQ ID NO:96] and H22A(+80+101) [SEQ ID NO:98] induced exon 22 skipping when delivered into cells at a concentration of 50 nM. Table 17 below shows other antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 17

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
95	H22A(+22 + 45)	CAC UCA UGG UCU CCU GAU AGC GCA	No skipping
96	H22A(+125 + 146)	CUG CAA UUC CCC GAG UCU CUG C	Skipping to 50 nM
97	H22A(+47 + 69)	ACU GCU GGA CCC AUG UCC UGA UG	Skipping to 300 nM
98	H22A(+80 + 101)	CUA AGU UGA GGU AUG GAG AGU	Skipping to 50 nM
99	H22D(+13 - 11)	UAU UCA CAG ACC UGC AAU UCC CC	No skipping

Antisense Oligonucleotides Directed at Exon 23

[0172] Antisense oligonucleotides directed at exon 23 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0173] Table 18 below shows antisense molecules tested at a concentration range of 25, 50, 100, 300 and 600 nM. These antisense molecules showed no ability to induce exon skipping or are yet to be tested.

TABLE 18

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
100	H23A(+34 + 59)	ACA GUG GUG CUG AGA UAG UAU AGG CC	No skipping
101	H23A(+18 + 39)	UAG GCC ACU UUG UUG CUC UUG C	No Skipping
102	H23A(+72 + 90)	UUC AGA GGG CGC UUU CUU C	No Skipping

Antisense Oligonucleotides Directed at Exon 24

[0174] Antisense oligonucleotides directed at exon 24 were prepared using similar methods as described above. Table 19 below outlines the antisense oligonucleotides directed at exon 24 that are yet to be tested for their ability to induce exon 24 skipping.

TABLE 19

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
103	H24A(+48 + 70)	GGG CAG GCC AUU CCU CCU UCA GA	Needs testing
104	H24A(-02 + 22)	UCU UCA GGG UUU GUA UGU GAU UCU	Needs testing

Antisense Oligonucleotides Directed at Exon 25

[0175] Antisense oligonucleotides directed at exon 25 were prepared using similar methods as described above. Table 20 below shows the antisense oligonucleotides directed at exon 25 that are yet to be tested for their ability to induce exon 25 skipping.

TABLE 20

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
105	H25A(+9 + 36)	CUG GGC UGA AUU GUC UGA AUA UCA CUG	Needs testing
106	H25A(+131 + 156)	CUG UUG GCA CAU GUG AUC CCA CUG AG	Needs testing
107	H25D(+16 - 08)	GUC UAU ACC UGU UGG CAC AUG UGA	Needs testing

Antisense Oligonucleotides Directed at Exon 26

[0176] Antisense oligonucleotides directed at exon 26 were prepared using similar methods as described above. Table 21 below outlines the antisense oligonucleotides directed at exon 26 that are yet to be tested for their ability to induce exon 26 skipping.

TABLE 21

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
108	H26A(+132 + 156)	UGC UUU CUG UAA UUC AUC UGG AGU U	Needs testing
109	H26A(-07 + 19)	CCU CCU UUC UGG CAU AGA CCU UCC AC	Needs testing
110	H26A(+68 + 92)	UGU GUC AUC CAU UCG UGC AUC UCU G	Faint skipping at 600 nM

Antisense Oligonucleotides Directed at Exon 27

[0177] Antisense oligonucleotides directed at exon 27 were prepared using similar methods as described above. Table 22 below outlines the antisense oligonucleotides directed at exon 27 that are yet to be tested for their ability to induce exon 27 skipping.

TABLE 22

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
111	H27A(+82 + 106)	UUA AGG CCU CUU GUG CUA CAG GUG G	Needs testing
112	H27A(-4 + 19)	GGG CCU CUU CUU UAG CUC UCU GA	Faint skipping at 600 and 300 nM
113	H27D(+19 - 03)	GAC UUC CAA AGU CUU GCA UUU C	v. strong skipping at 600 and 300 nM

Antisense Oligonucleotides Directed at Exon 28

[0178] Antisense oligonucleotides directed at exon 28 were prepared using similar methods as described above. Table 23 below outlines the antisense oligonucleotides directed at exon 28 that are yet to be tested for their ability to induce exon 28 skipping.

TABLE 23

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
114	H28A(-05 + 19)	GCC AAC AUG CCC AAA CUU CCU AAG	v. strong skipping at 600 and 300 nM
115	H28A(+99 + 124)	CAG AGA UUU CCU CAG CUC CGC CAG GA	Needs testing
116	H28D(+16 - 05)	CUU ACA UCU AGC ACC UCA GAG	v. strong skipping at 600 and 300 nM

Antisense Oligonucleotides Directed at Exon 29

[0179] Antisense oligonucleotides directed at exon 29 were prepared using similar methods as described above. Table 24 below outlines the antisense oligonucleotides directed at exon 29 that are yet to be tested for their ability to induce exon 29 skipping.

TABLE 24

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
117	H29A(+57 + 81)	UCC GCC AUC UGU UAG GGU CUG UGC C	Needs testing
118	H29A(+18 + 42)	AUU UGG GUU AUC CUC UGA AUG UCG C	v. strong skipping at 600 and 300 nM
119	H29D(+17 - 05)	CAU ACC UCU UCA UGU AGU UCC C	v. strong skipping at 600 and 300 nM

Antisense Oligonucleotides Directed at Exon 30

[0180] Antisense oligonucleotides directed at exon 30 were prepared using similar methods as described above. Table 25 below outlines the antisense oligonucleotides directed at exon 30 that are yet to be tested for their ability to induce exon 30 skipping.

TABLE 25

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
120	H30A(+122 + 147)	CAU UUG AGC UGC GUC CAC CUU GUC UG	Needs testing
121	H30A(+25 + 50)	UCC UGG GCA GAC UGG AUG CUC UGU UC	Very strong skipping at 600 and 300 nM.
122	H30D(+19 - 04)	UUG CCU GGG CUU CCU GAG GCA UU	Very strong skipping at 600 and 300 nM.

Antisense Oligonucleotides Directed at Exon 31

[0181] Antisense oligonucleotides directed at exon 31 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0182] FIG. 13 illustrates differing efficiencies of two antisense molecules directed at exon 31 acceptor splice site and a

“cocktail” of exon 31 antisense oligonucleotides at varying concentrations. H31D(+03-22) [SEQ ID NO:124] substantially induced exon 31 skipping when delivered into cells at a concentration of 20 nM. Table 26 below also includes other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 26

SEQ ID name	Antisense oligonucleotide Sequence	Ability to induce skipping
123 H31D(+06 - 18)	UUC UGA AAU AAC AUA UAC CUG UGC	Skipping to 300 nM
124 H31D(+03 - 22)	UAG UUU CUG AAA UAA CAU AUA CCU G	Skipping to 20 nM
125 H31A(+05 + 25)	GAC UUG UCA AAU CAG AUU GGA	No skipping
126 H31D(+04 - 20)	GUU UCU GAA AUA ACA UAU ACC UGU	Skipping to 300 nM

Antisense Oligonucleotides Directed at Exon 32

[0183] Antisense oligonucleotides directed at exon 32 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0184] H32D(+04-16) [SEQ ID NO:127] and H32A(+49+73) [SEQ ID NO:130] induced exon 32 skipping when delivered into cells at a concentration of 300 nM. Table 27 below also shows other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules did not show an ability to induce exon skipping.

TABLE 27

SEQ ID name	Antisense oligonucleotide Sequence	Ability to induce skipping
127H32D(+04 - 16)	CAC CAG AAA UAC AUA CCA CA	Skipping to 300 nM
128H32A(+151 + 170)	CAA UGA UUU AGC UGU GAC UG	No skipping
129H32A(+10 + 32)	CGA AAC UUC AUG GAG ACA UCU UG	No skipping
130H32A(+49 + 73)	CUU GUA GAC GCU GCU CAA AAU UGG C	Skipping to 300 nM

Antisense Oligonucleotides Directed at Exon 33

[0185] Antisense oligonucleotides directed at exon 33 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0186] FIG. 14 shows differing efficiencies of two antisense molecules directed at exon 33 acceptor splice site. H33A(+64+88) [SEQ ID NO:134] substantially induced exon 33 skipping when delivered into cells at a concentration of 10 nM. Table 28 below includes other antisense molecules tested at a concentration of 100, 200 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 28

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
131	H33D(+09 - 11)	CAU GCA CAC ACC UUU GCU CC	No skipping
132	H33A(+53 + 76)	UCU GUA CAA UCU GAC GUC CAG UCU	Skipping to 200 nM
133	H33A(+30 + 56)	GUG UUU AUC ACC AUU UCC ACU UCA GAC	Skipping to 200 nM
134	H33A(+64 + 88)	GCG UCU GCU UUU UCU GUA CAA UCU G	Skipping to 10 nM

Antisense Oligonucleotides Directed at Exon 34

[0187] Antisense oligonucleotides directed at exon 34 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0188] Table 29 below includes antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed a variable ability to induce exon skipping.

TABLE 29

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
135	H34A(+83 + 104)	UCC AUA UCU GUA GCU GGC AGC C	No skipping
136	H34A(+143 + 165)	CCA GGC AAC UUC AGA AUC CAA AU	No skipping
137	H34A(-20 + 10)	UUU CUG UUA CCU GAA AAG AAU UAU AAU GAA	Not tested
138	H34A(+46 + 70)	CAU UCA UUU CCU UUC GCA UCU UAC G	Skipping to 300 nM
139	H34A(+95 + 120)	UGA UCU CUU UGU CAA UUC CAU AUC UG	Skipping to 300 nM
140	H34D(+10 - 20)	UUC AGU GAU AUA GGU UUU ACC UUU CCC CAG	Not tested
141	H34A(+72 + 96)	CUG UAG CUG CCA GCC AUU CUG UCA AG	No skipping

Antisense Oligonucleotides Directed at Exon 35

[0189] Antisense oligonucleotides directed at exon 35 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0190] FIG. 15 shows differing efficiencies of antisense molecules directed at exon 35 acceptor splice site. H35A(+24+43) [SEQ ID NO:144] substantially induced exon 35 skipping when delivered into cells at a concentration of 20 nM. Table 30 below also includes other antisense molecules tested at a concentration of 100 and 300 nM. These antisense molecules showed no ability to induce exon skipping.

TABLE 30

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
142	H35A(+141 + 161)	UCU UCU GCU CGG GAG GUG ACA	Skipping to 20 nM
143	H35A(+116 + 135)	CCA GUU ACU AUU CAG AAG AC	No skipping
144	H35A(+24 + 43)	UCU UCA GGU GCA CCU UCU GU	No skipping

Antisense Oligonucleotides Directed at Exon 36

[0191] Antisense oligonucleotides directed at exon 36 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0192] Antisense molecule H36A(+26+50) [SEQ ID NO:145] induced exon 36 skipping when delivered into cells at a concentration of 300 nM, as shown in FIG. 16.

Antisense Oligonucleotides Directed at Exon 37

[0193] Antisense oligonucleotides directed at exon 37 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0194] FIG. 17 shows differing efficiencies of two antisense molecules directed at exon 37 acceptor splice site. H37A(+82+105) [SEQ ID NO:148] and H37A(+134+157) [SEQ ID NO:149] substantially induced exon 37 skipping when delivered into cells at a concentration of 10 nM. Table 31 below shows the antisense molecules tested.

TABLE 31

SEQ IDname	Antisense oligonucleotide	Sequence	Ability to induce skipping
147	H37A(+26 + 50)	CGU GUA GAG UCC ACC UUU GGG CGU A	No skipping
148	H37A(+82 + 105)	UAC UAA UUU CCU GCA GUG GUC ACC	Skipping to 10 nM
149	H37A(+134 + 157)	UUC UGU GUG AAA UGG CUG CAA AUC	Skipping to 10 nM

Antisense Oligonucleotides Directed at Exon 38

[0195] Antisense oligonucleotides directed at exon 38 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0196] FIG. 18 illustrates antisense molecule H38A(+88+112) [SEQ ID NO:152], directed at exon 38 acceptor splice site. H38A(+88+112) substantially induced exon 38 skipping when delivered into cells at a concentration of 10 nM. Table 32 below shows the antisense molecules tested and their ability to induce exon skipping.

TABLE 32

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
150	H38A(-01 + 19)	CCU UCA AAG GAA UGG AGG CC	No skipping
151	H38A(+59 + 83)	UGC UGA AUU UCA GCC UCC AGU GGU U	Skipping to 10 nM
152	H38A(+88 + 112)	UGA AGU CUU CCU CUU UCA GAU UCA C	Skipping to 10 nM

Antisense Oligonucleotides Directed at Exon 39

[0197] Antisense oligonucleotides directed at exon 39 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0198] H39A(+62+85) [SEQ ID NO:153] induced exon 39 skipping when delivered into cells at a concentration of 100 nM. Table 33 below shows the antisense molecules tested and their ability to induce exon skipping.

TABLE 33

SEQ IDname	Antisense oligonucleotide	Sequence	Ability to induce skipping
153	H39A(+62 + 85)	CUG GCU UUC UCU CAU CUG UGA UUC	Skipping to 100 nM
154	H39A(+39 + 58)	GUU GUA AGU UGU CUC CUC UU	No skipping

TABLE 33-continued

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
155	H39A(+102 + 121)	UUG UCU GUA ACA GCU GCU GU	No skipping
156	H39D(+10 - 10)	GCU CUA AUA CCU UGA GAG CA	Skipping to 300 nM

Antisense Oligonucleotides Directed at Exon 40

[0199] Antisense oligonucleotides directed at exon 40 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0200] FIG. 19 illustrates antisense molecule H40A(-05+17) [SEQ ID NO:157] directed at exon 40 acceptor splice site. H40A(-05+17) and H40A(+129+153) [SEQ ID NO:158] both substantially induced exon 40 skipping when delivered into cells at a concentration of 5 nM.

Antisense Oligonucleotides Directed at Exon 42

[0201] Antisense oligonucleotides directed at exon 42 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0202] FIG. 20 illustrates antisense molecule H42A(-04+23) [SEQ ID NO:159], directed at exon 42 acceptor splice site. H42A(-4+23) and H42D(+19-02) [SEQ ID NO:161] both induced exon 42 skipping when delivered into cells at a concentration of 5 nM. Table 34 below shows the antisense molecules tested and their ability to induce exon 42 skipping.

TABLE 34

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
159	H42A(-4 + 23)	AUC GUU UCU UCA CGG ACA GUG UGG UGC	Skipping to 5 nM
160	H42A(+86 + 109)	GGG CUU GUG AGA CAU GAG UGA UUU	Skipping to 100 nM
161	H42D(+19 - 02)	A CCU UCA GAG GAC UCC UCU UGC	Skipping to 5 nM

Antisense Oligonucleotides Directed at Exon 43

[0203] Antisense oligonucleotides directed at exon 43 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0204] H43A(+101+120) [SEQ ID NO:163] induced exon 43 skipping when delivered into cells at a concentration of 25 nM. Table 35 below includes the antisense molecules tested and their ability to induce exon 43 skipping.

TABLE 35

SEQ ID name	Antisense oligonucleotide	Sequence	Ability to induce skipping
162	H43D(+10 - 15)	UAU GUG UUA CCU ACC CUU GUC GGU C	Skipping to 100 nM
163	H43A(+101 + 120)	GGA GAG AGC UUC CUG UAG CU	Skipping to 25 nM
164	H43A(+78 + 100)	UCA CCC UUU CCA CAG GCG UUG CA	Skipping to 200 nM

Antisense Oligonucleotides Directed at Exon 44

[0205] Antisense oligonucleotides directed at exon 44 were prepared using similar methods as described above. Testing for the ability of these antisense molecules to induce exon 44 skipping is still in progress. The antisense molecules under review are shown as SEQ ID Nos: 165 to 167 in Table 1A.

Antisense Oligonucleotides Directed at Exon 45

[0206] Antisense oligonucleotides directed at exon 45 were prepared using similar methods as described above. Testing for the ability of these antisense molecules to induce exon 45 skipping is still in progress. The antisense molecules under review are shown as SEQ ID Nos: 207 to 211 in Table 1A.

Antisense Oligonucleotides Directed at Exon 46

[0207] Antisense oligonucleotides directed at exon 46 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0208] FIG. 21 illustrates the efficiency of one antisense molecule directed at exon 46 acceptor splice site. Antisense oligonucleotide H46A(+86+115) [SEQ ID NO:203] showed very strong ability to induce exon 46 skipping. Table 36 below includes antisense molecules tested. These antisense molecules showed varying ability to induce exon 46 skipping.

Antisense Oligonucleotides Directed at Exon 47

[0209] Antisense oligonucleotides directed at exon 47 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0210] H47A(+76+100) [SEQ ID NO:170] and H47A(-09+12) [SEQ ID NO:172] both induced exon 47 skipping when delivered into cells at a concentration of 200 nM. H47D(+25-02) [SEQ ID NO: 171] is yet to be prepared and tested.

Antisense Oligonucleotides Directed at Exon 50

[0211] Antisense oligonucleotides directed at exon 50 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0212] Antisense oligonucleotide molecule H50A(+02+30) [SEQ ID NO: 173] was a strong inducer of exon skipping. Further, H50A(+07+33) [SEQ ID NO:174] and H50D(+07-18) [SEQ ID NO:175] both induced exon 50 skipping when delivered into cells at a concentration of 100 nM.

Antisense Oligonucleotides Directed at Exon 51

[0213] Antisense oligonucleotides directed at exon 51 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0214] FIG. 22 illustrates differing efficiencies of two antisense molecules directed at exon 51 acceptor splice site. Antisense oligonucleotide H51A(+66+90) [SEQ ID NO:180] showed the stronger ability to induce exon 51 skipping. Table 37 below includes antisense molecules tested at a concentration range of 25, 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon 51 skipping. The strongest inducers of exon skipping were antisense oligonucleotide H51A(+61+90) [SEQ ID NO: 179] and H51A(+66+95) [SEQ ID NO: 181].

TABLE 36

SEQ ID name	Antisense oligonucleotide Sequence	Ability to induce skipping
168	H46D(+16 - 04) UUA CCU UGA CUU GCU CAA GC	No skipping
169	H46A(+90 + 109) UCC AGG UUC AAG UGG GAU AC	No skipping
203	H46A(+86 + 115) CUC UUU UCC AGG UUC AAG UGG GAU ACU AGC	Good skipping to 100 nM
204	H46A(+107 + 137) CAA GCU UUU CUU UUA GUU GCU GCU CUU UUC C	Good skipping to 100 nM
205	H46A(-10 + 20) UAU UCU UUU GUU CUU CUA GCC UGG AGA AAG	Weak skipping
206	H46A(+50 + 77) CUG CUU CCU CCA ACC AUA AAA CAA AUU C	Weak skipping

TABLE 37

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
176	H51A(-01 + 25)	ACC AGA GUA ACA GUC UGA GUA GGA GC	Faint skipping
177	H51D(+16 - 07)	CUC AUA CCU UCU GCU UGA UGA UC	Skipping at 300 nM
178	H51A(+111 + 134)	UUC UGU CCA AGC CCG GUU GAA AUC	Needs re-testing
179	H51A(+61 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA GUU UGG	Very strong skipping
180	H51A(+66 + 90)	ACA UCA AGG AAG AUG GCA UUU CUA G	skipping
181	H51A(+66 + 95)	CUC CAA CAU CAA GGA AGA UGG CAU UUC UAG	Very strong skipping
182	H51D(+08 - 17)	AUC AUU UUU UCU CAU ACC UUC UGC U	No skipping
183	H51A/D(+08 - 17) & (-15 + ?)	AUC AUU UUU UCU CAU ACC UUC UGC UAG GAG CUA AAA	No skipping
184	H51A(+175 + 195)	CAC CCA CCA UCA GCC UCU GUG	No skipping
185	H51A(+199 + 220)	AUC AUC UCG UUG AUA UCC UCA A	No skipping

Antisense Oligonucleotides Directed at Exon 52

[0215] Antisense oligonucleotides directed at exon 52 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0216] FIG. 22 also shows differing efficiencies of four antisense molecules directed at exon 52 acceptor splice site. The most effective antisense oligonucleotide for inducing exon 52 skipping was H52A(+17+37) [SEQ ID NO:188].

[0217] Table 38 below shows antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon 50 skipping. Antisense molecules H52A(+12+41) [SEQ ID NO:187] and H52A(+17+37) [SEQ ID NO:188] showed the strongest exon 50 skipping at a concentration of 50 nM.

TABLE 38

SEQ ID	Antisense oligonucleotide name	Sequence	Ability to induce skipping
186	H52A(-07 + 14)	UCC UGC AUU GUU GCC UGU AAG	No skipping
187	H52A(+12 + 41)	UCC AAC UGG GGA CGC CUC UGU UCC AAA UCC	Very strong skipping
188	H52A(+17 + 37)	ACU GGG GAC GCC UCU GUU CCA	Skipping to 50 nM
189	H52A(+93 + 112)	CCG UAA UGA UUG UUC UAG CC	No skipping
190	H52D(+05 - 15)	UGU UAA AAA ACU UAC UUC GA	No skipping

Antisense Oligonucleotides Directed at Exon 53

[0218] Antisense oligonucleotides directed at exon 53 were prepared and tested for their ability to induce exon skipping in human muscle cells using similar methods as described above.

[0219] FIG. 22 also shows antisense molecule H53A(+39+69) [SEQ ID NO:193] directed at exon 53 acceptor splice site. This antisense oligonucleotide was able to induce exon 53 skipping at 5, 100, 300 and 600 nM. A “cocktail” of three

exon 53 antisense oligonucleotides: H53A(+23+47) [SEQ ID NO:195], H53A(+150+176) [SEQ ID NO:196] and H53D(+14-07) [SEQ ID NO:194], was also tested, as shown in FIG. 20 and exhibited an ability to induce exon skipping.

[0220] Table 39 below includes other antisense molecules tested at a concentration range of 50, 100, 300 and 600 nM. These antisense molecules showed varying ability to induce exon 53 skipping. Antisense molecule H53A(+39+69) [SEQ ID NO:193] induced the strongest exon 53 skipping.

TABLE 39

SEQ ID name	Antisense oligonucleotide name	Sequence	Ability to induce skipping
191	H53A(+45 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU G	Faint skipping at 50 nM
192	H53A(+39 + 62)	CUG UUG CCU CCG GUU CUG AAG GUG	Faint skipping at 50 nM
193	H53A(+39 + 69)	CAU UCA ACU GUU GCC UCC GGU UCU GAA GGU G	Strong skipping to 50 nM
194	H53D(+14 - 07)	UAC UAA CCU UGG UUU CUG UGA	Very faint skipping to 50 nM
195	H53A(+23 + 47)	CUG AAG GUG UUC UUG UAC UUC AUC C	Very faint skipping to 50 nM
196	H53A(+150 + 176)	UGU AUA GGG ACC CUC CUU CCA UGA CUC	Very faint skipping to 50 nM
197	H53D(+20 - 05)	CUA ACC UUG GUU UCU GUG AUU UUC U	Not made yet
198	H53D(+09 - 18)	GGU AUC UUU GAU ACU AAC CUU GGU UUC	Faint at 600 nM
199	H53A(-12 + 10)	AUU CUU UCA ACU AGA AUA AAA G	No skipping
200	H53A(-07 + 18)	GAU UCU GAA UUG UUU CAA CUA GAA U	No skipping
201	H53A(+07 + 26)	AUC CCA CUG AUU CUG AAU UC	No skipping
202	H53A(+124 + 145)	UUG GCU CUG GCC UGU CCU AAG A	No skipping

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 214

<210> SEQ ID NO 1

<211> LENGTH: 24

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic Human 2'-O-methyl phosphorothioate antisense oligonucleotide

<400> SEQUENCE: 1

gauagguggu aucaacaucu guaa

24

-continued

<210> SEQ ID NO 2
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 2

gauagguggu aucaacaucu g 21

<210> SEQ ID NO 3
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 3

gauagguggu aucaacaucu gaaag 25

<210> SEQ ID NO 4
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 4

ggugguauca acaucuguaa 20

<210> SEQ ID NO 5
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 5

guaucaacau cuguaagcac 20

<210> SEQ ID NO 6
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 6

ugcauguucc agucguugug ugg 23

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 7

cacuaaucca gucaaaauagg ucugg 25

<210> SEQ ID NO 8
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 8

auuuaccaac cuucaggauc gagua 25

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 9

ggccuaaaac acauacacau a 21

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 10

cauuuuugac cuacaugugg 20

<210> SEQ ID NO 11
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 11

uuugaccuac auguggaaag 20

<210> SEQ ID NO 12
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 12

uacauuuuug accuacaugu ggaaag 26

-continued

<210> SEQ ID NO 13
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 13

auuuuugacc uacaugggaa ag 22

<210> SEQ ID NO 14
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 14

uacgaguuga uugucggacc cag 23

<210> SEQ ID NO 15
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 15

guggucuccu uaccuaugac ugugg 25

<210> SEQ ID NO 16
<211> LENGTH: 17
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Canine 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 16

ggucuccuua ccuauga 17

<210> SEQ ID NO 17
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 17

ugucucagua aucuucuuaac cuau 24

<210> SEQ ID NO 18
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 18

ucuuaccuau gacuauggau gaga 24

<210> SEQ ID NO 19
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 19

gcaugaacuc uuguggauc 20

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 20

ccagguacu acuuacaua 20

<210> SEQ ID NO 21
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 21

aucguguc acagcauca g 21

<210> SEQ ID NO 22
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 22

uguucagggc augaacucu guggauccu 30

<210> SEQ ID NO 23
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 23

uaggaggcgc cuccauccu guaggucac g 31

-continued

<210> SEQ ID NO 24
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 24
aggucuagga ggcgccuccc auccguaggu u 31

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 25
gcgccuccca uccguaggu cacug 25

<210> SEQ ID NO 26
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 26
cuucgaggag gucuaggagg cgccuc 26

<210> SEQ ID NO 27
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 27
cucccauccu guaggucacu g 21

<210> SEQ ID NO 28
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 28
uaccaguuuu ugcccuguca gg 22

<210> SEQ ID NO 29
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 29

ucaauaugcu gcuucccaaa cugaaa 26

<210> SEQ ID NO 30
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 30

cuaggaggcg ccucccaucc uguag 25

<210> SEQ ID NO 31
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 31

uuugauuuc caucuacgau gucaguacuu c 31

<210> SEQ ID NO 32
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 32

cuuaccugcc aguggaggau uauauccaa a 31

<210> SEQ ID NO 33
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 33

caucaggauu cuuaccugcc agugg 25

<210> SEQ ID NO 34
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 34

cgaugucagu acuuccaaua uucac 25

-continued

<210> SEQ ID NO 35
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 35
accauucauc aggauucu 18

<210> SEQ ID NO 36
<211> LENGTH: 18
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 36
accugccagu ggaggauu 18

<210> SEQ ID NO 37
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 37
ccaauuuca cuaaaaucaac cuguuaa 27

<210> SEQ ID NO 38
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 38
caggauucuu accugccagu ggaggauuau 30

<210> SEQ ID NO 39
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 39
acgaugucag uacuuccaau auucacuaaa u 31

<210> SEQ ID NO 40
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 40

auuuccaucu acgaugucag uacuuccaau a 31

<210> SEQ ID NO 41
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 41

caggagcuuc caaaugcugc a 21

<210> SEQ ID NO 42
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 42

cuugucuuca ggagcuucca aaucugca 29

<210> SEQ ID NO 43
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 43

uccucagcag aaagaagcca cg 22

<210> SEQ ID NO 44
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 44

uuagaaaucu cuccuugugc 20

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 45

uaaaugggu guacacaau 20

-continued

<210> SEQ ID NO 46
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 46
cccugaggca uucccaucuu gaau 24

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 47
aggacuuacu ugcuuuguuu 20

<210> SEQ ID NO 48
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 48
cuugaaauua ggagauucau cug 23

<210> SEQ ID NO 49
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 49
caucuucuga uaaauuuuccu guu 23

<210> SEQ ID NO 50
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 50
ucuucuguuu uuguuagcca guca 24

<210> SEQ ID NO 51
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 51

ucuauguaaa cugaaaauuu 20

<210> SEQ ID NO 52
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 52

uucuggagau ccauuaaaac 20

<210> SEQ ID NO 53
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 53

cagcaguugc gugaucucca cuag 24

<210> SEQ ID NO 54
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 54

uucaucaacu accaccacca u 21

<210> SEQ ID NO 55
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 55

cuaagcaaaa uaaucugacc uuaag 25

<210> SEQ ID NO 56
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 56

cuuguaaaag aaccagcgg ucuucugu 28

-continued

<210> SEQ ID NO 57
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 57

caucuacaga uguuugccca uc 22

<210> SEQ ID NO 58
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 58

gaaggauguc uuguaaaaga acc 23

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 59

accguucuu caguaagacg 20

<210> SEQ ID NO 60
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 60

caugacacac cuguucuuca guaa 24

<210> SEQ ID NO 61
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 61

cauuugagaa ggauugcuug 20

<210> SEQ ID NO 62
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 62

aucucccaau accuggagaa gaga 24

<210> SEQ ID NO 63
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 63

gccaugcacu aaaaaggcac ugcaagacau u 31

<210> SEQ ID NO 64
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 64

ucuuuaaagc caguugugug aauc 24

<210> SEQ ID NO 65
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 65

uuucugaaag ccaugcacua a 21

<210> SEQ ID NO 66
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 66

guacauacgg ccaguuuuug aagac 25

<210> SEQ ID NO 67
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 67

cuagaucgc uuuuaaaacc uguuaaaca a 31

-continued

<210> SEQ ID NO 68
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 68
ucuuuuuag auccgcuuuu aaaaccguu a 31

<210> SEQ ID NO 69
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 69
cuagauccgc uuuuaaaacc uguua 25

<210> SEQ ID NO 70
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 70
ccgucuucug ggucacugac uua 23

<210> SEQ ID NO 71
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 71
cuagauccgc uuuuaaaacc uguuaa 26

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 72
ccgcuuuuaa aaccguuaa 20

<210> SEQ ID NO 73
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 73

uggauugcuu uuucuuuucu agaucc 26

<210> SEQ ID NO 74
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 74

caugcuuccg ucuucugggu cacug 25

<210> SEQ ID NO 75
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 75

gaucuuguuu gagugaauac agu 23

<210> SEQ ID NO 76
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 76

guuauccagc caugcuuccg uc 22

<210> SEQ ID NO 77
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 77

ugauaaaugg uaucacuaac cugug 25

<210> SEQ ID NO 78
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 78

guaucacuaa ccugugcugu ac 22

-continued

<210> SEQ ID NO 79
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 79

cugcuggcau cuugcaguu 19

<210> SEQ ID NO 80
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 80

gccugagcug aucugcuggc aucuugcagu u 31

<210> SEQ ID NO 81
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 81

cuggcagaau ucgauccacc ggcuguuc 28

<210> SEQ ID NO 82
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 82

cagcaguagu ugucaucugc uc 22

<210> SEQ ID NO 83
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 83

ugauggggug guggguugg 19

<210> SEQ ID NO 84
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 84

aucugcauaa acaccucua gaaag 25

<210> SEQ ID NO 85
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 85

ccggcuguuc aguuguucug aggc 24

<210> SEQ ID NO 86
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 86

aucugcauaa acaccucua gaaagaaa 28

<210> SEQ ID NO 87
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 87

gaaggagaag agauucuuac cuuacaaa 28

<210> SEQ ID NO 88
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 88

auucgaucca ccggcuguuc 20

<210> SEQ ID NO 89
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 89

cagcaguagu ugucaucugc 20

-continued

<210> SEQ ID NO 90
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 90
gccgguugac ucauccugu gc 22

<210> SEQ ID NO 91
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 91
cugcauccag gaacaugggu cc 22

<210> SEQ ID NO 92
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 92
gucugcaucc aggaacaugg guc 23

<210> SEQ ID NO 93
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 93
guugaagauc ugauagccgg uuga 24

<210> SEQ ID NO 94
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 94
uacuuacugu cuguagcucu uuuc 24

<210> SEQ ID NO 95
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 95

cacucauggu cuccugauag cgca 24

<210> SEQ ID NO 96
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 96

cugcaauucc ccgagucucu gc 22

<210> SEQ ID NO 97
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 97

acugcuggac ccauguccug aug 23

<210> SEQ ID NO 98
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 98

cuaaguugag guauggagag u 21

<210> SEQ ID NO 99
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 99

uaauccacaga ccugcaauuc ccc 23

<210> SEQ ID NO 100
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 100

acaguggugc ugagauagua uaggcc 26

-continued

<210> SEQ ID NO 101
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 101

uagggccacu uguugcucu gc 22

<210> SEQ ID NO 102
<211> LENGTH: 19
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 102

uucagagggc gcuuucuuc 19

<210> SEQ ID NO 103
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 103

gggcaggcca uuccuccuuc aga 23

<210> SEQ ID NO 104
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 104

ucuucagggg uuguauuguga uucu 24

<210> SEQ ID NO 105
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 105

cugggcugaa uugucugaau aucacug 27

<210> SEQ ID NO 106
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 106

cuguuggcac augugauccc acugag 26

<210> SEQ ID NO 107
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 107

gucuauaccu guuggcacau guga 24

<210> SEQ ID NO 108
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 108

ugcuuucugu aaucaucug gaguu 25

<210> SEQ ID NO 109
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 109

ccuccuuucu ggcauagacc uuccac 26

<210> SEQ ID NO 110
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 110

ugugucaucc auucgugcau cucug 25

<210> SEQ ID NO 111
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 111

uuaggccuc uugugcuaca ggugg 25

-continued

<210> SEQ ID NO 112
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 112

gggccucuc uuuagcucuc uga 23

<210> SEQ ID NO 113
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 113

gacuuccaaa gucuugcauu uc 22

<210> SEQ ID NO 114
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 114

gccaaacaugc ccaaacuucc uaag 24

<210> SEQ ID NO 115
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 115

cagagauuuc cucagcuccg ccagga 26

<210> SEQ ID NO 116
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 116

cuuacaucua gcaccucaga g 21

<210> SEQ ID NO 117
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 117

uccgccaucu guuagggucu gugcc 25

<210> SEQ ID NO 118
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 118

auuuggguua uccucugaau gugcc 25

<210> SEQ ID NO 119
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 119

cauaccucu cauguaguuc cc 22

<210> SEQ ID NO 120
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 120

cauuugagcu gguccaccu ugucug 26

<210> SEQ ID NO 121
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 121

uccugggcag acuggaugcu cuguuc 26

<210> SEQ ID NO 122
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 122

uugccugggc uuccugaggc auu 23

-continued

<210> SEQ ID NO 123
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 123

uucugaaaua acauauaccu gugc 24

<210> SEQ ID NO 124
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 124

uaguuuucuga aauaacauau accug 25

<210> SEQ ID NO 125
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 125

gacuugucuaa aucagauugg a 21

<210> SEQ ID NO 126
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 126

guuucugaaa uaacauauac cugu 24

<210> SEQ ID NO 127
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 127

caccagaaau acauaccaca 20

<210> SEQ ID NO 128
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 128

caaugauuuu gcugugacug 20

<210> SEQ ID NO 129
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 129

cgaaacuuca uggagacauc uug 23

<210> SEQ ID NO 130
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 130

cuuguagacg cugcucaaaa uuggc 25

<210> SEQ ID NO 131
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 131

caugcacaca ccuuugcucc 20

<210> SEQ ID NO 132
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 132

ucuguacaau cugacgucca gucu 24

<210> SEQ ID NO 133
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 133

gucuuuauca ccauuuccac uucagac 27

-continued

<210> SEQ ID NO 134
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 134

ccgucugcuu uuucuguaca aucug 25

<210> SEQ ID NO 135
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 135

uccauaucug uagcugccag cc 22

<210> SEQ ID NO 136
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 136

ccaggcaacu ucagaaucca aau 23

<210> SEQ ID NO 137
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 137

uuucuguuac cugaaaagaa uuauaagaa 30

<210> SEQ ID NO 138
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 138

cauucuuuc cuuucgcauc uuacg 25

<210> SEQ ID NO 139
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 139

ugaucucuuu gucaauucca uaucug 26

<210> SEQ ID NO 140
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 140

uucagugaua uagguuuuac cuuucccag 30

<210> SEQ ID NO 141
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 141

cuguagcugc cagccauucu gucaag 26

<210> SEQ ID NO 142
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 142

ucuucugcuc gggaggugac a 21

<210> SEQ ID NO 143
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 143

ccaguuacua uucagaagac 20

<210> SEQ ID NO 144
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 144

ucuucaggug caccuucugu 20

-continued

<210> SEQ ID NO 145
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 145

ugugaugugg uccacauucu gguca 25

<210> SEQ ID NO 146
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 146

ccauguguuu cugguauucc 20

<210> SEQ ID NO 147
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 147

cguguagagu ccaccuuugg gcgua 25

<210> SEQ ID NO 148
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 148

uacuaauuuc cugcaguggu cacc 24

<210> SEQ ID NO 149
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 149

uucuguguga aauggcugca aauc 24

<210> SEQ ID NO 150
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 150

ccucaaagg aauggaggcc 20

<210> SEQ ID NO 151
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 151

ugcugaauu cagccuccag ugguu 25

<210> SEQ ID NO 152
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 152

ugaagucuuc cucuuucaga uucac 25

<210> SEQ ID NO 153
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 153

cuggcuuucu cucaucugug auuc 24

<210> SEQ ID NO 154
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense oligonucleotide

<400> SEQUENCE: 154

guuguaagu gucuccuuu 20

<210> SEQ ID NO 155
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 155

uugucuguaa cagcugcugu 20

-continued

<210> SEQ ID NO 156
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 156

gcucuaauac cuugagagca 20

<210> SEQ ID NO 157
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 157

cuugagacc ucaaauccug uu 22

<210> SEQ ID NO 158
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 158

cuuuuuuuuc cuuucaucuc ugggc 25

<210> SEQ ID NO 159
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 159

aucguuucuu cacggacagu gugcugg 27

<210> SEQ ID NO 160
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 160

gggcuuguga gacaugagug auuu 24

<210> SEQ ID NO 161
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 161

accuucagag gacuccucu gc 22

<210> SEQ ID NO 162
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 162

uauguguuac cuacccuugu cgguc 25

<210> SEQ ID NO 163
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 163

ggagagagcu uccuguagcu 20

<210> SEQ ID NO 164
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 164

ucacccuuuc cacaggcguu gca 23

<210> SEQ ID NO 165
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 165

uuugugucu ucugagaaac 20

<210> SEQ ID NO 166
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 166

aaagacuuac cuuaagauac 20

-continued

<210> SEQ ID NO 167
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 167

aucugucaaa ucgccugcag 20

<210> SEQ ID NO 168
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 168

uuaccuugac uugcucaagc 20

<210> SEQ ID NO 169
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 169

uccagguuca agugggauac 20

<210> SEQ ID NO 170
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 170

gcucuucugg gcuaauggga gcacu 25

<210> SEQ ID NO 171
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 171

accuuuaacc acuggagauu ugucugc 27

<210> SEQ ID NO 172
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 172

uuccaccagu aacugaaaca g 21

<210> SEQ ID NO 173
<211> LENGTH: 29
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 173

ccacucagag cucagauuu cuaacuucc 29

<210> SEQ ID NO 174
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 174

cuuccacuca gagcucagau cuucuaa 27

<210> SEQ ID NO 175
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 175

gggauccagu auacuuacag gcucc 25

<210> SEQ ID NO 176
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 176

accagaguua cagucugagu aggagc 26

<210> SEQ ID NO 177
<211> LENGTH: 23
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 177

cucauaccuu cugcuugaug auc 23

-continued

<210> SEQ ID NO 178
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 178

uucuguccaa gcccgguuga aauc 24

<210> SEQ ID NO 179
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 179

acaucaagga agauggcgau ucuaguugg 30

<210> SEQ ID NO 180
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 180

acaucaagga agauggcgau ucuag 25

<210> SEQ ID NO 181
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 181

cuccaacauc aaggaagaug gcauuucug 30

<210> SEQ ID NO 182
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 182

aucuuuuuu cucuaccuu cugcu 25

<210> SEQ ID NO 183
<211> LENGTH: 36
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 183

aucuuuuuu cucauaccuu cugcuaggag cuaaaa 36

<210> SEQ ID NO 184
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 184

caccacccau caccucugu g 21

<210> SEQ ID NO 185
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 185

aucaucucgu ugauaucuc aa 22

<210> SEQ ID NO 186
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 186

uccugcaug uugccuguaa g 21

<210> SEQ ID NO 187
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 187

uccaacuggg gacgccucug uuccaaaacc 30

<210> SEQ ID NO 188
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 188

acuggggacg ccucuguucc a 21

-continued

<210> SEQ ID NO 189
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 189

ccguaaugau uguucuagcc 20

<210> SEQ ID NO 190
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 190

uguuaaaaaa cuuacuugca 20

<210> SEQ ID NO 191
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 191

cauuccaacug uugccuccgg uucug 25

<210> SEQ ID NO 192
<211> LENGTH: 24
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 192

cuguugccuc cgguucugaa ggug 24

<210> SEQ ID NO 193
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 193

cauuccaacug uugccuccgg uucugaaggu g 31

<210> SEQ ID NO 194
<211> LENGTH: 21
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 194

uacuaaccuu gguuucugug a 21

<210> SEQ ID NO 195
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 195

cugaaggugu ucuuguacuu caucc 25

<210> SEQ ID NO 196
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 196

uguauaggga cccuccuucc augacuc 27

<210> SEQ ID NO 197
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 197

cuaaccuugg uuucugugau uuucu 25

<210> SEQ ID NO 198
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 198

gguaucuuug auacuaaccu ugguuuc 27

<210> SEQ ID NO 199
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 199

auucuuuca cuagaauaaa ag 22

-continued

<210> SEQ ID NO 200
<211> LENGTH: 25
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 200

gauucugaau ucuuuaacu agaau 25

<210> SEQ ID NO 201
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 201

aucccacuga uucugaauc 20

<210> SEQ ID NO 202
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 202

uuggcucugg ccuguccuaa ga 22

<210> SEQ ID NO 203
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 203

cucuuuucca gguucaagug ggauacuagc 30

<210> SEQ ID NO 204
<211> LENGTH: 31
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 204

caagcuuuuc uuuuaguugc ugcucuuuuc c 31

<210> SEQ ID NO 205
<211> LENGTH: 30
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 205

uauucuuuug uucuuuagc cuggagaaag 30

<210> SEQ ID NO 206
<211> LENGTH: 28
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 206

cugcuuccuc caaccuaaaa acaaaauuc 28

<210> SEQ ID NO 207
<211> LENGTH: 26
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 207

ccaaugccau ccuggaguuc cuguaa 26

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 208

uccuguagaa uacuggcauc 20

<210> SEQ ID NO 209
<211> LENGTH: 27
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

<400> SEQUENCE: 209

ugcagaccuc cugccaccgc agauuca 27

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Human 2'-O-methyl phosphorothioate antisense
oligonucleotide

-continued

```

<400> SEQUENCE: 210
cuaccucuuu uuucugucug                               20

<210> SEQ ID NO 211
<211> LENGTH: 20
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 211
uguuuuugag gauugcugaa                               20

<210> SEQ ID NO 212
<211> LENGTH: 84
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      Human 2'-O-methyl phosphorothioate antisense
      oligonucleotide

<400> SEQUENCE: 212
cagcaguagu ugucaucugc ucaacuggca gaauucgauc caccggcugu ucaagccuga   60
gcugaucugc ucgcaucuug cagu                                           84

<210> SEQ ID NO 213
<211> LENGTH: 44
<212> TYPE: RNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 213
uaugcacug agugaccucu uucucgcagg cgcuagcugg agca                       44

<210> SEQ ID NO 214
<211> LENGTH: 22
<212> TYPE: RNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 214
ccgugcagac ugacggucuc au                                             22

```

We claim:

1. An isolated antisense oligonucleotide of 20 to 50 nucleotides in length comprising at least 10 consecutive nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1-212, wherein the oligonucleotide specifically hybridizes to a target region in an exon of the human dystrophin gene inducing exon skipping, and wherein uracil bases in the antisense oligonucleotide are optionally thymine bases.

2. The antisense oligonucleotide of claim 1, wherein the exon in human dystrophin is selected from the group consisting of exons 3-8, 10-16, 19-40, 42-47, and 50-53.

3. The antisense oligonucleotide of claim 1, wherein the oligonucleotide does not activate RNase H.

4. The antisense oligonucleotide of claim 1, comprising a non-natural backbone.

5. The antisense oligonucleotide of claim 1, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties.

6. The antisense oligonucleotide of claim 5, wherein the non-natural moieties are morpholinos.

7. The antisense oligonucleotide of claim 1, wherein the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

8. The antisense oligonucleotide of claim 7, wherein the non-natural inter-nucleotide linkages are modified phosphates.

9. The antisense oligonucleotide of claim 1, wherein the sugar moieties of the oligonucleotide backbone are replaced with non-natural moieties and the inter-nucleotide linkages of the oligonucleotide backbone are replaced with non-natural inter-nucleotide linkages.

10. The antisense oligonucleotide of claim **9**, wherein the non-natural moieties are morpholinos and the non-natural internucleotide linkages are modified phosphates, and wherein the modified phosphates are methyl phosphonates, methyl phosphorothioates, phosphoromorpholidates, phosphoropiperazidates, or phosphoroamidates.

11. The antisense oligonucleotide of claim **1**, wherein the oligonucleotide is a peptide nucleic acid.

12. The antisense oligonucleotide of claim **1**, wherein the oligonucleotide is chemically linked to one or more moieties or conjugates that enhance the activity, cellular distribution, or cellular uptake of the antisense oligonucleotide.

13. The antisense oligonucleotide of claim **12**, wherein the oligonucleotide is conjugated to a polyamine.

14. The antisense oligonucleotide of claim **12**, wherein the oligonucleotide is chemically linked to a polyethylene glycol chain.

15. An isolated antisense oligonucleotide of 20 to 50 nucleotides in length comprising at least 10 consecutive nucle-

otides complementary to a target region of an exon in the human dystrophin gene designated as an annealing site selected from the group set forth in Table 1A, wherein the antisense oligonucleotide specifically hybridizes to the annealing site inducing exon skipping.

16. An expression vector comprising the antisense oligonucleotide of claim **1**.

17. The expression vector of claim **16**, wherein the expression vector is a modified retrovirus or a non-retroviral vector.

18. The expression vector of claim **17**, wherein the non-retroviral vector is an adeno-associated virus (AAV).

19. A pharmaceutical composition, comprising an antisense oligonucleotide of claim **1**, and a saline solution that includes a phosphate buffer.

20. A method of treating Duchenne muscular dystrophy, comprising administering to a patient in need thereof an effective amount of a pharmaceutical composition of claim **19**.

* * * * *