



US 20140058056A1

(19) **United States**

(12) **Patent Application Publication**

Burgard et al.

(10) **Pub. No.: US 2014/0058056 A1**

(43) **Pub. Date: Feb. 27, 2014**

(54) **MICROORGANISMS AND METHODS FOR ENHANCING THE AVAILABILITY OF REDUCING EQUIVALENTS IN THE PRESENCE OF METHANOL, AND FOR PRODUCING 1,4-BUTANEDIOL RELATED THERETO**

(71) Applicant: **Genomatica, Inc.**, San Diego, CA (US)

(72) Inventors: **Anthony P. Burgard**, Bellefonte, PA (US); **Robin E. Osterhout**, San Diego, CA (US); **Stephen J. Van Dien**, Encinitas, CA (US); **Cara Ann Tracewell**, Solana Beach, CA (US); **Priti Pharkya**, San Diego, CA (US); **Stefan Andrae**, San Diego, CA (US)

(73) Assignee: **Genomatica, Inc.**, San Diego, CA (US)

(21) Appl. No.: **13/975,678**

(22) Filed: **Aug. 26, 2013**

Related U.S. Application Data

(60) Provisional application No. 61/693,683, filed on Aug. 27, 2012, provisional application No. 61/766,609, filed on Feb. 19, 2013.

Publication Classification

(51) **Int. Cl.**
C12P 7/18 (2006.01)
(52) **U.S. Cl.**
CPC **C12P 7/18** (2013.01)
USPC .. **528/85**; 435/252.3; 435/254.2; 435/254.11;
435/158; 435/157; 435/147; 568/852; 528/425;
528/308.6; 528/272; 528/271; 528/332;
568/617; 548/543

(57) **ABSTRACT**

Provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway that can enhance the availability of reducing equivalents in the presence of methanol. Such reducing equivalents can be used to increase the product yield of organic compounds produced by the microbial organism, such as 1,4-butanediol (BDO). Also provided herein are methods for using such an organism to produce BDO.




Figure 1




Figure 2




Figure 3




Figure 4

**MICROORGANISMS AND METHODS FOR
ENHANCING THE AVAILABILITY OF
REDUCING EQUIVALENTS IN THE
PRESENCE OF METHANOL, AND FOR
PRODUCING 1,4-BUTANEDIOL RELATED
THERE TO**

CROSS-REFERENCE TO RELATED
APPLICATIONS

[0001] This application claims the benefit of U.S. Ser. No. 61/766,609 filed Feb. 19, 2013, and U.S. Ser. No. 61/693,683 filed Aug. 27, 2012, each of which is incorporated herein by reference in its entirety.

1. SUMMARY

[0002] Provided herein are methods generally relating to metabolic and biosynthetic processes and microbial organisms capable of producing organic compounds. Specifically, provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway that can enhance the availability of reducing equivalents in the presence of methanol. Such reducing equivalents can be used to increase the product yield of organic compounds produced by the microbial organism, such as 1,4-butanediol (BDO). Also provided herein are non-naturally occurring microbial organisms and methods thereof to produce optimal yields of BDO.

[0003] In a first aspect, provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol. In certain embodiments, the methanol metabolic pathway comprises one or more enzymes selected from the group consisting of a methanol methyltransferase; a methylenetetrahydrofolate reductase; a methylenetetrahydrofolate dehydrogenase; a methenyltetrahydrofolate cyclohydrolase; a formyltetrahydrofolate deformylase; a formyltetrahydrofolate synthetase; a formate hydrogen lyase; a hydrogenase; a formate dehydrogenase; a methanol dehydrogenase; a formaldehyde activating enzyme; a formaldehyde dehydrogenase; a S-(hydroxymethyl)glutathione synthase; a glutathione-dependent formaldehyde dehydrogenase; and an S-formylglutathione hydrolase. Such organisms advantageously allow for the production of reducing equivalents, which can then be used by the organism for the production of BDO using any one of the BDO pathways provided herein.

[0004] In certain embodiments, the organism further comprises a BDO pathway. In certain embodiments, said organism comprises at least one exogenous nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO. In certain embodiments, the BDO pathway enzyme is selected from the group consisting of a succinyl-CoA transferase or a succinyl-CoA synthetase (or succinyl-CoA ligase); a succinyl-CoA reductase (aldehyde forming); a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase; a 4-hydroxybutyryl-CoA reductase (aldehyde forming); a 1,4-butanediol dehydrogenase; a succinate reductase; a succinyl-CoA reductase (alcohol forming); a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyrate reductase; a 4-hydroxybutyryl-phosphate reductase; and a 4-hydroxybutyryl-CoA reductase (alcohol forming).

[0005] In other embodiments, the organism having a methanol metabolic pathway, either alone or in combination with a BDO pathway, as provided herein, further comprises a formaldehyde assimilation pathway that utilizes formaldehyde, e.g., obtained from the oxidation of methanol, in the formation of intermediates of certain central metabolic pathways that can be used, for example, in the formation of biomass. In some of the embodiments, the formaldehyde assimilation pathway comprises a hexylose-6-phosphate synthase, 6-phospho-3-hexyloisomerase, dihydroxyacetone synthase or dihydroxyacetone kinase. In certain embodiments, provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol dehydrogenase expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol and/or expressed in a sufficient amount to convert methanol to formaldehyde. In some embodiments, the microbial organism further comprises a formaldehyde assimilation pathway. In certain embodiments, the organism further comprises at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme expressed in a sufficient amount to produce an intermediate of glycolysis. In certain embodiments, the formaldehyde assimilation pathway enzyme is selected from the group consisting of a hexylose-6-phosphate synthase, 6-phospho-3-hexyloisomerase, dihydroxyacetone synthase and dihydroxyacetone kinase.

[0006] In some embodiments, the organism further comprises one or more gene disruptions, occurring in one or more endogenous genes encoding protein(s) or enzyme(s) involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂, and/or amino acids by said microbial organism, wherein said one or more gene disruptions confer increased production of BDO in said microbial organism. In some embodiments, one or more endogenous enzymes involved in native production of ethanol, glycerol, acetate, lactate, formate, CO, and/or amino acids by the microbial organism, has attenuated enzyme activity or expression levels. In certain embodiments, the organism comprises from one to twenty-five gene disruptions. In other embodiments, the organism comprises from one to twenty gene disruptions. In some embodiments, the organism comprises from one to fifteen gene disruptions. In other embodiments, the organism comprises from one to ten gene disruptions. In some embodiments, the organism comprises from one to five gene disruptions. In certain embodiments, the organism comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 gene disruptions or more.

[0007] In another aspect, provided herein is a method for producing BDO, comprising culturing any one of the non-naturally occurring microbial organisms comprising a methanol metabolic pathway and an BDO pathway provided herein under conditions and for a sufficient period of time to produce BDO. In certain embodiments, the organism is cultured in a substantially anaerobic culture medium.

2. BRIEF DESCRIPTION OF THE DRAWINGS

[0008] FIG. 1 shows exemplary metabolic pathways enabling the extraction of reducing equivalents from methanol. The enzymatic transformations shown are carried out by the following enzymes: 1A) a methanol methyltransferase, 1B) a methylenetetrahydrofolate reductase, 1C) a methylenetetrahydrofolate dehydrogenase, 1D) a methenyltetrahydro-

folate cyclohydrolase, 1E) a formyltetrahydrofolate deformylase, 1F) a formyltetrahydrofolate synthetase, 1G) a formate hydrogen lyase, 1H) a hydrogenase, 1I) a formate dehydrogenase, 1J) a methanol dehydrogenase, 1K) a formaldehyde activating enzyme, 1L) a formaldehyde dehydrogenase, 1M) a S-(hydroxymethyl)glutathione synthase, 1N) a glutathione-dependent formaldehyde dehydrogenase, and 1O) a S-formylglutathione hydrolase. In certain embodiments, steps K and/or M are spontaneous.

[0009] FIG. 2 shows exemplary BDO pathways, which can be used to increase BDO yields from carbohydrates when reducing equivalents produced by a methanol metabolic pathway provided herein are available. BDO production is carried out by the following enzymes: 2A) a succinyl-CoA transferase or a succinyl-CoA synthetase, 2B) a succinyl-CoA reductase (aldehyde forming), 2C) a 4-hydroxybutyrate dehydrogenase, 2D) a 4-hydroxybutyrate kinase, 2E) a phosphotrans-4-hydroxybutyrylase, 2F) a 4-hydroxybutyryl-CoA reductase (aldehyde forming), 2G) a 1,4-butanediol dehydrogenase, 2H) a succinate reductase, 2I) a succinyl-CoA reductase (alcohol forming), 2J) a 4-hydroxybutyryl-CoA transferase or 4-hydroxybutyryl-CoA synthetase, 2K) a 4-hydroxybutyrate reductase, 2L) a 4-hydroxybutyryl-phosphate reductase, and 2M) a 4-hydroxybutyryl-CoA reductase (alcohol forming).

[0010] FIG. 3 shows an exemplary formaldehyde assimilation pathway. The enzymatic transformations are carried out by the following enzymes: 3A) a hexylose-6-phosphate synthase, and 3B) a 6-phospho-3-hexyloisomerase.

[0011] FIG. 4 shows an exemplary formaldehyde assimilation pathway. The enzymatic transformations are carried out by the following enzymes: 4A) a dihydroxyacetone synthase, and 4B) a dihydroxyacetone kinase.

3. DETAILED DESCRIPTION OF THE INVENTION

3.1 Definitions

[0012] As used herein, the term “non-naturally occurring” when used in reference to a microbial organism or microorganism of the invention is intended to mean that the microbial organism has at least one genetic alteration not normally found in a naturally occurring strain of the referenced species, including wild-type strains of the referenced species. Genetic alterations include, for example, modifications introducing expressible nucleic acids encoding metabolic polypeptides, other nucleic acid additions, nucleic acid deletions and/or other functional disruption of the microbial organism’s genetic material. Such modifications include, for example, coding regions and functional fragments thereof, for heterologous, homologous or both heterologous and homologous polypeptides for the referenced species. Additional modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. Exemplary metabolic polypeptides include enzymes or proteins within a BDO or 4-hydroxybutyrate (4-FIB) biosynthetic pathway.

[0013] A metabolic modification refers to a biochemical reaction that is altered from its naturally occurring state. Therefore, non-naturally occurring microorganisms can have genetic modifications to nucleic acids encoding metabolic polypeptides, or functional fragments thereof. Exemplary metabolic modifications are disclosed herein.

[0014] As used herein, the term “isolated” when used in reference to a microbial organism is intended to mean an organism that is substantially free of at least one component as the referenced microbial organism is found in nature. The term includes a microbial organism that is removed from some or all components as it is found in its natural environment. The term also includes a microbial organism that is removed from some or all components as the microbial organism is found in non-naturally occurring environments. Therefore, an isolated microbial organism is partly or completely separated from other substances as it is found in nature or as it is grown, stored or subsisted in non-naturally occurring environments. Specific examples of isolated microbial organisms include partially pure microbes, substantially pure microbes and microbes cultured in a medium that is non-naturally occurring.

[0015] As used herein, the terms “microbial,” “microbial organism” or “microorganism” are intended to mean any organism that exists as a microscopic cell that is included within the domains of archaea, bacteria or eukarya. Therefore, the term is intended to encompass prokaryotic or eukaryotic cells or organisms having a microscopic size and includes bacteria, archaea and eubacteria of all species as well as eukaryotic microorganisms such as yeast and fungi. The term also includes cell cultures of any species that can be cultured for the production of a biochemical.

[0016] As used herein, the term “CoA” or “coenzyme A” is intended to mean an organic cofactor or prosthetic group (nonprotein portion of an enzyme) whose presence is required for the activity of many enzymes (the apoenzyme) to form an active enzyme system. Coenzyme A functions in certain condensing enzymes, acts in acetyl or other acyl group transfer and in fatty acid synthesis and oxidation, pyruvate oxidation and in other acetylation.

[0017] As used herein, the term “substantially anaerobic” when used in reference to a culture or growth condition is intended to mean that the amount of oxygen is less than about 10% of saturation for dissolved oxygen in liquid media. The term also is intended to include sealed chambers of liquid or solid medium maintained with an atmosphere of less than about 1% oxygen.

[0018] As used herein, the term “gene disruption,” or grammatical equivalents thereof, is intended to mean a genetic alteration that renders the encoded gene product inactive or attenuated. The genetic alteration can be, for example, deletion of the entire gene, deletion of a regulatory sequence required for transcription or translation, deletion of a portion of the gene which results in a truncated gene product, or by any of various mutation strategies that inactivate or attenuate the encoded gene product. One particularly useful method of gene disruption is complete gene deletion because it reduces or eliminates the occurrence of genetic reversions in the non-naturally occurring microorganisms of the invention. The phenotypic effect of a gene disruption can be a null mutation, which can arise from many types of mutations including inactivating point mutations, entire gene deletions, and deletions of chromosomal segments or entire chromosomes. Specific antisense nucleic acid compounds and enzyme inhibitors, such as antibiotics, can also produce null mutant phenotype, therefore being equivalent to gene disruption.

[0019] As used herein, the term “growth-coupled” when used in reference to the production of a biochemical product is intended to mean that the biosynthesis of the referenced biochemical product is produced during the growth phase of

a microorganism. In a particular embodiment, the growth-coupled production can be obligatory, meaning that the biosynthesis of the referenced biochemical is an obligatory product produced during the growth phase of a microorganism. The term “growth-coupled” when used in reference to the consumption of a biochemical is intended to mean that the referenced biochemical is consumed during the growth phase of a microorganism.

[0020] As used herein, the term “attenuate,” or grammatical equivalents thereof, is intended to mean to weaken, reduce or diminish the activity or amount of an enzyme or protein. Attenuation of the activity or amount of an enzyme or protein can mimic complete disruption if the attenuation causes the activity or amount to fall below a critical level required for a given pathway to function. However, the attenuation of the activity or amount of an enzyme or protein that mimics complete disruption for one pathway, can still be sufficient for a separate pathway to continue to function. For example, attenuation of an endogenous enzyme or protein can be sufficient to mimic the complete disruption of the same enzyme or protein for production of a fatty alcohol, fatty aldehyde or fatty acid product of the invention, but the remaining activity or amount of enzyme or protein can still be sufficient to maintain other pathways, such as a pathway that is critical for the host microbial organism to survive, reproduce or grow. Attenuation of an enzyme or protein can also be weakening, reducing or diminishing the activity or amount of the enzyme or protein in an amount that is sufficient to increase yield of a fatty alcohol, fatty aldehyde or fatty acid product of the invention, but does not necessarily mimic complete disruption of the enzyme or protein.

[0021] “Exogenous” as it is used herein is intended to mean that the referenced molecule or the referenced activity is introduced into the host microbial organism. The molecule can be introduced, for example, by introduction of an encoding nucleic acid into the host genetic material such as by integration into a host chromosome or as non-chromosomal genetic material such as a plasmid. Therefore, the term as it is used in reference to expression of an encoding nucleic acid refers to introduction of the encoding nucleic acid in an expressible form into the microbial organism. When used in reference to a biosynthetic activity, the term refers to an activity that is introduced into the host reference organism. The source can be, for example, a homologous or heterologous encoding nucleic acid that expresses the referenced activity following introduction into the host microbial organism. Therefore, the term “endogenous” refers to a referenced molecule or activity that is present in the host. Similarly, the term when used in reference to expression of an encoding nucleic acid refers to expression of an encoding nucleic acid contained within the microbial organism. The term “heterologous” refers to a molecule or activity derived from a source other than the referenced species whereas “homologous” refers to a molecule or activity derived from the host microbial organism. Accordingly, exogenous expression of an encoding nucleic acid of the invention can utilize either or both a heterologous or homologous encoding nucleic acid.

[0022] It is understood that when more than one exogenous nucleic acid is included in a microbial organism that the more than one exogenous nucleic acids refers to the referenced encoding nucleic acid or biosynthetic activity, as discussed above. It is further understood, as disclosed herein, that such more than one exogenous nucleic acids can be introduced into the host microbial organism on separate nucleic acid mol-

ecules, on polycistronic nucleic acid molecules, or a combination thereof, and still be considered as more than one exogenous nucleic acid. For example, as disclosed herein a microbial organism can be engineered to express two or more exogenous nucleic acids encoding a desired pathway enzyme or protein. In the case where two exogenous nucleic acids encoding a desired activity are introduced into a host microbial organism, it is understood that the two exogenous nucleic acids can be introduced as a single nucleic acid, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be considered as two exogenous nucleic acids. Similarly, it is understood that more than two exogenous nucleic acids can be introduced into a host organism in any desired combination, for example, on a single plasmid, on separate plasmids, can be integrated into the host chromosome at a single site or multiple sites, and still be considered as two or more exogenous nucleic acids, for example three exogenous nucleic acids. Thus, the number of referenced exogenous nucleic acids or biosynthetic activities refers to the number of encoding nucleic acids or the number of biosynthetic activities, not the number of separate nucleic acids introduced into the host organism.

[0023] The non-naturally occurring microbial organisms of the invention can contain stable genetic alterations, which refers to microorganisms that can be cultured for greater than five generations without loss of the alteration. Generally, stable genetic alterations include modifications that persist greater than 10 generations, particularly stable modifications will persist more than about 25 generations, and more particularly, stable genetic modifications will be greater than 50 generations, including indefinitely.

[0024] Those skilled in the art will understand that the genetic alterations, including metabolic modifications exemplified herein, are described with reference to a suitable host organism such as *E. coli* and their corresponding metabolic reactions or a suitable source organism for desired genetic material such as genes for a desired metabolic pathway. However, given the complete genome sequencing of a wide variety of organisms and the high level of skill in the area of genomics, those skilled in the art will readily be able to apply the teachings and guidance provided herein to essentially all other organisms. For example, the *E. coli* metabolic alterations exemplified herein can readily be applied to other species by incorporating the same or analogous encoding nucleic acid from species other than the referenced species. Such genetic alterations include, for example, genetic alterations of species homologs, in general, and in particular, orthologs, paralogs or nonorthologous gene displacements.

[0025] An ortholog is a gene or genes that are related by vertical descent and are responsible for substantially the same or identical functions in different organisms. For example, mouse epoxide hydrolase and human epoxide hydrolase can be considered orthologs for the biological function of hydrolysis of epoxides. Genes are related by vertical descent when, for example, they share sequence similarity of sufficient amount to indicate they are homologous, or related by evolution from a common ancestor. Genes can also be considered orthologs if they share three-dimensional structure but not necessarily sequence similarity, of a sufficient amount to indicate that they have evolved from a common ancestor to the extent that the primary sequence similarity is not identifiable. Genes that are orthologous can encode proteins with sequence similarity of about 25% to 100% amino acid

sequence identity. Genes encoding proteins sharing an amino acid similarity less than 25% can also be considered to have arisen by vertical descent if their three-dimensional structure also shows similarities. Members of the serine protease family of enzymes, including tissue plasminogen activator and elastase, are considered to have arisen by vertical descent from a common ancestor.

[0026] Orthologs include genes or their encoded gene products that through, for example, evolution, have diverged in structure or overall activity. For example, where one species encodes a gene product exhibiting two functions and where such functions have been separated into distinct genes in a second species, the three genes and their corresponding products are considered to be orthologs. For the production of a biochemical product, those skilled in the art will understand that the orthologous gene harboring the metabolic activity to be introduced or disrupted is to be chosen for construction of the non-naturally occurring microorganism. An example of orthologs exhibiting separable activities is where distinct activities have been separated into distinct gene products between two or more species or within a single species. A specific example is the separation of elastase proteolysis and plasminogen proteolysis, two types of serine protease activity, into distinct molecules as plasminogen activator and elastase. A second example is the separation of mycoplasma 5'-3' exonuclease and *Drosophila* DNA polymerase III activity. The DNA polymerase from the first species can be considered an ortholog to either or both of the exonuclease or the polymerase from the second species and vice versa.

[0027] In contrast, paralogs are homologs related by, for example, duplication followed by evolutionary divergence and have similar or common, but not identical functions. Paralogs can originate or derive from, for example, the same species or from a different species. For example, microsomal epoxide hydrolase (epoxide hydrolase I) and soluble epoxide hydrolase (epoxide hydrolase II) can be considered paralogs because they represent two distinct enzymes, co-evolved from a common ancestor, that catalyze distinct reactions and have distinct functions in the same species. Paralogs are proteins from the same species with significant sequence similarity to each other suggesting that they are homologous, or related through co-evolution from a common ancestor. Groups of paralogous protein families include HipA homologs, luciferase genes, peptidases, and others.

[0028] A nonorthologous gene displacement is a non-orthologous gene from one species that can substitute for a referenced gene function in a different species. Substitution includes, for example, being able to perform substantially the same or a similar function in the species of origin compared to the referenced function in the different species. Although generally, a nonorthologous gene displacement will be identifiable as structurally related to a known gene encoding the referenced function, less structurally related but functionally similar genes and their corresponding gene products nevertheless will still fall within the meaning of the term as it is used herein. Functional similarity requires, for example, at least some structural similarity in the active site or binding region of a nonorthologous gene product compared to a gene encoding the function sought to be substituted. Therefore, a nonorthologous gene includes, for example, a paralog or an unrelated gene.

[0029] Therefore, in identifying and constructing the non-naturally occurring microbial organisms of the invention having BDO or 4-hydroxybutyrate biosynthetic capability, those

skilled in the art will understand with applying the teaching and guidance provided herein to a particular species that the identification of metabolic modifications can include identification and inclusion or inactivation of orthologs. To the extent that paralogs and/or nonorthologous gene displacements are present in the referenced microorganism that encode an enzyme catalyzing a similar or substantially similar metabolic reaction, those skilled in the art also can utilize these evolutionally related genes.

[0030] Orthologs, paralogs and nonorthologous gene displacements can be determined by methods well known to those skilled in the art. For example, inspection of nucleic acid or amino acid sequences for two polypeptides will reveal sequence identity and similarities between the compared sequences. Based on such similarities, one skilled in the art can determine if the similarity is sufficiently high to indicate the proteins are related through evolution from a common ancestor. Algorithms well known to those skilled in the art, such as Align, BLAST, Clustal W and others compare and determine a raw sequence similarity or identity, and also determine the presence or significance of gaps in the sequence which can be assigned a weight or score. Such algorithms also are known in the art and are similarly applicable for determining nucleotide sequence similarity or identity. Parameters for sufficient similarity to determine relatedness are computed based on well known methods for calculating statistical similarity, or the chance of finding a similar match in a random polypeptide, and the significance of the match determined. A computer comparison of two or more sequences can, if desired, also be optimized visually by those skilled in the art. Related gene products or proteins can be expected to have a high similarity, for example, 25% to 100% sequence identity. Proteins that are unrelated can have an identity which is essentially the same as would be expected to occur by chance, if a database of sufficient size is scanned (about 5%). Sequences between 5% and 24% may or may not represent sufficient homology to conclude that the compared sequences are related. Additional statistical analysis to determine the significance of such matches given the size of the data set can be carried out to determine the relevance of these sequences.

[0031] Exemplary parameters for determining relatedness of two or more sequences using the BLAST algorithm, for example, can be as set forth below. Briefly, amino acid sequence alignments can be performed using BLASTP version 2.0.8 (Jan. 5, 1999) and the following parameters: Matrix: 0 BLOSUM62; gap open: 11; gap extension: 1; x_dropoff: 50; expect: 10.0; wordsize: 3; filter: on. Nucleic acid sequence alignments can be performed using BLASTN version 2.0.6 (Sep. 16, 1998) and the following parameters: Match: 1; mismatch: -2; gap open: 5; gap extension: 2; x_dropoff: 50; expect: 10.0; wordsize: 11; filter: off. Those skilled in the art will know what modifications can be made to the above parameters to either increase or decrease the stringency of the comparison, for example, and determine the relatedness of two or more sequences.

3.2 Microbial Organisms that Utilize Reducing Equivalents Produced by the Metabolism of Methanol

[0032] Provided herein are methanol metabolic pathways engineered to improve the availability of reducing equivalents, which can be used for the production of product molecules. Exemplary product molecules include, without limitation, BDO and/or 4HB, although given the teachings and

guidance provided herein, it will be recognized by one skilled in the art that any product molecule that utilizes reducing equivalents in its production can exhibit enhanced production through the biosynthetic pathways provided herein.

[0033] Methanol is a relatively inexpensive organic feedstock that can be derived from synthesis gas components, CO and H₂, via catalysis. Methanol can be used as a source of reducing equivalents to increase the molar yield of product molecules from carbohydrates.

[0034] BDO is a valuable chemical for the production of high performance polymers, solvents, and fine chemicals. It is the basis for producing other high value chemicals such as tetrahydrofuran (THF) and gamma-butyrolactone (GBL). The value chain is comprised of three main segments including: (1) polymers, (2) THF derivatives, and (3) GBL derivatives. In the case of polymers, BDO is a comonomer for polybutylene terephthalate (PBT) production. PBT is a medium performance engineering thermoplastic used in automotive, electrical, water systems, and small appliance applications. Conversion to THF, and subsequently to polytetramethylene ether glycol (PTMEG), provides an intermediate used to manufacture spandex products such as LYCRA® fibers. PTMEG is also combined with BDO in the production of specialty polyester ethers (COPE). COPEs are high modulus elastomers with excellent mechanical properties and oil/environmental resistance, allowing them to operate at high and low temperature extremes. PTMEG and BDO also make thermoplastic polyurethanes processed on standard thermoplastic extrusion, calendaring, and molding equipment, and are characterized by their outstanding toughness and abrasion resistance. The GBL produced from BDO provides the feedstock for making pyrrolidones, as well as serving the agrochemical market. The pyrrolidones are used as high performance solvents for extraction processes of increasing use, including for example, in the electronics industry and in pharmaceutical production. Accordingly, provided herein is bioderived BDO produced according to the methods described herein and biobased products comprising or obtained using the bioderived BDO. The biobased product can comprise a polymer, THF or a THF derivative, or GBL or a GBL derivative; or the biobased product can comprise a polymer, a plastic, elastic fiber, polyurethane, polyester, polyhydroxyalkanoate, poly-4-hydroxybutyrate, co-polymer of poly-4-hydroxybutyrate, poly(tetramethylene ether) glycol, polyurethane-polyurea copolymer, spandex, elastane, Lycra™, or nylon; or the biobased product can comprise polybutylene terephthalate (PBT) polymer; or the biobased product can comprise a PBT polymer that comprises a resin, a fiber, a bead, a granule, a pellet, a chip, a plastic, a polyester, a thermoplastic polyester, a molded article, an injection-molded article, an injection-molded part, an automotive part, an extrusion resin, an electrical part and a casing, optionally where the biobased product is reinforced or filled, for example glass-filled or mineral-filled; or the biobased product is THF or a THF derivative, and the THF derivative is polytetramethylene ether glycol (PTMEG), a polyester ether (COPE) or a thermoplastic polyurethane or a fiber; or the biobased product comprises GBL or a GBL derivative and the GBL derivative is a pyrrolidone. The biobased product can comprise at least 5%, at least 10%, at least 20%, at least 30%, at least 40% or at least 50% bioderived BDO. The biobased product can comprise a portion of said bioderived BDO as a repeating unit. The biobased product can be a molded product obtained by molding the biobased product.


[0035] BDO is produced by two main petrochemical routes with a few additional routes also in commercial operation. One route involves reacting acetylene with formaldehyde, followed by hydrogenation. More recently, BDO processes involving butane or butadiene oxidation to maleic anhydride, followed by hydrogenation have been introduced. BDO is used almost exclusively as an intermediate to synthesize other chemicals and polymers. Thus, there exists a need for the development of methods for effectively producing commercial quantities of BDO.

[0036] In numerous engineered pathways, realization of maximum product yields based on carbohydrate feedstock is hampered by insufficient reducing equivalents or by loss of reducing equivalents to byproducts. Methanol is a relatively inexpensive organic feedstock that can be used to generate reducing equivalents by employing one or more methanol metabolic enzymes as shown in FIG. 1. The reducing equivalents produced by the metabolism of methanol by one or more of the methanol metabolic pathways can then be used to power the glucose to BDO production pathways, for example, as shown in FIG. 2.


[0037] The product yields per C-mol of substrate of microbial cells synthesizing reduced fermentation products such as BDO and 4-HB are limited by insufficient reducing equivalents in the carbohydrate feedstock. Reducing equivalents, or electrons, can be extracted from methanol using one or more of the enzymes described in FIG. 1. The reducing equivalents are then passed to acceptors such as oxidized ferredoxins, oxidized quinones, oxidized cytochromes, NAD(P)⁺, water, or hydrogen peroxide to form reduced ferredoxin, reduced quinones, reduced cytochromes, NAD(P)H, H₂, or water, respectively. Reduced ferredoxin, reduced quinones and NAD(P)H are particularly useful as they can serve as redox carriers for various Wood-Ljungdahl pathway, reductive TCA cycle, or product pathway enzymes.

[0038] Specific examples of how additional redox availability from methanol can improve the yield of reduced products such as succinate, 4-hydroxybutyrate, and BDO are shown.



[0039] The maximum theoretical yield of BDO via the pathway shown in FIG. 2 supplemented with the reactions of the oxidative TCA cycle (e.g., citrate synthase, aconitase, isocitrate dehydrogenase, alpha-ketoglutarate dehydrogenase) is 1.09 mol/mol.



[0040] When both feedstocks of sugar and methanol are available, the methanol can be utilized to generate reducing equivalents by employing one or more of the enzymes shown in FIG. 1. The reducing equivalents generated from methanol can be utilized to power the glucose to BDO production pathways, e.g., as shown in FIG. 2. Theoretically, all carbons in glucose will be conserved, thus resulting in a maximal theoretical yield to produce BDO from glucose at 2 mol BDO per mol of glucose under either aerobic or anaerobic conditions as shown in FIG. 2:



[0041] In a similar manner, the maximum theoretical yields of succinate and 4-hydroxybutyrate can reach 2 mol/mol glucose using the reactions shown in FIGS. 1 and 2.



[0042] In a first aspect, provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol. In certain embodiments, the methanol metabolic pathway comprises one or more enzymes selected from the group consisting of a methanol methyltransferase; a methylenetetrahydrofolate reductase; a methylenetetrahydrofolate dehydrogenase; a methenyltetrahydrofolate cyclohydrolase; a formyltetrahydrofolate deformylase; a formyltetrahydrofolate synthetase; a formate hydrogen lyase; a hydrogenase; a formate dehydrogenase; a methanol dehydrogenase; a formaldehyde activating enzyme; a formaldehyde dehydrogenase; a S-(hydroxymethyl)glutathione synthase; a glutathione-dependent formaldehyde dehydrogenase; and an S-formylglutathione hydrolase. Such organisms advantageously allow for the production of reducing equivalents, which can then be used by the organism for the production of BDO or 4-HB using any one of the pathways provided herein.

[0043] In certain embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O or any combination of 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, and 1O, thereof, wherein 1A is a methanol methyltransferase; 1B is a methylenetetrahydrofolate reductase; 1C is a methylenetetrahydrofolate dehydrogenase; 1D is a methenyltetrahydrofolate cyclohydrolase; 1E is a formyltetrahydrofolate deformylase; 1F is a formyltetrahydrofolate synthetase; 1G is a formate hydrogen lyase; 1H is a hydrogenase, 1I is a formate dehydrogenase; 1J is a methanol dehydrogenase; 1K is a formaldehyde activating enzyme; 1L is a formaldehyde dehydrogenase; 1M is a S-(hydroxymethyl)glutathione synthase; 1N is glutathione-dependent formaldehyde dehydrogenase; and 1O is S-formylglutathione hydrolase. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0044] In one embodiment, the methanol metabolic pathway comprises 1A. In another embodiment, the methanol metabolic pathway comprises 1B. In another embodiment, the methanol metabolic pathway comprises 1C. In yet another embodiment, the methanol metabolic pathway comprises 1D. In one embodiment, the methanol metabolic pathway comprises 1E. In another embodiment, the methanol metabolic pathway comprises 1F. In another embodiment, the methanol metabolic pathway comprises 1G. In yet another embodiment, the methanol metabolic pathway comprises 1H. In one embodiment, the methanol metabolic pathway comprises 1I. In another embodiment, the methanol metabolic pathway comprises 1J. In another embodiment, the methanol metabolic pathway comprises 1K. In yet another embodiment, the methanol metabolic pathway comprises 1L. In yet another embodiment, the methanol metabolic pathway comprises 1M. In another embodiment, the methanol metabolic pathway comprises 1N. In yet another embodiment, the methanol metabolic pathway comprises 1O. Any combination of two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen or fifteen methanol metabolic pathway enzymes 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, and 1O is also contemplated.

[0045] In some embodiments, the methanol metabolic pathway is a methanol metabolic pathway depicted in FIG. 1.

[0046] In one aspect, provided herein is a non-naturally occurring microbial organism having a methanol metabolic pathway enzyme expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol, wherein said methanol metabolic pathway comprises: (i) 1A and 1B, (ii) 1J; or (iii) 1J and 1K. In one embodiment, the methanol metabolic pathway comprises 1A and 1B. In another embodiment, the methanol metabolic pathway comprises 1J. In one embodiment, the methanol metabolic pathway comprises 1J and 1K. In certain embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, and 1E. In some embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D and 1F. In some embodiments, the methanol metabolic pathway comprises 1J, 1C, 1D and 1E. In one embodiment, the methanol metabolic pathway comprises 1J, 1C, 1D and 1F. In another embodiment, the methanol metabolic pathway comprises 1J and 1L. In yet another embodiment, the methanol metabolic pathway comprises 1J, 1M, 1N and 1O. In certain embodiments, the methanol metabolic pathway comprises 1J, 1N and 1O. In some embodiments, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1E. In one embodiment, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1F. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0047] In certain embodiments, the methanol metabolic pathway comprises 1I. In certain embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1E and 1I. In some embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1F and 1I. In some embodiments, the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1I. In one embodiment, the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1I. In another embodiment, the methanol metabolic pathway comprises 1J, 1L and 1I. In yet another embodiment, the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1I. In certain embodiments, the methanol metabolic pathway comprises 1J, 1N, 1O and 1I. In some embodiments, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1I. In one embodiment, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F and 1I. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0048] In certain embodiments, the methanol metabolic pathway comprises 1G. In certain embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1E and 1G. In some embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1F and 1G. In some embodiments, the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1G. In one embodiment, the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1G. In another embodiment, the methanol metabolic pathway comprises 1J, 1L and 1G. In yet another embodiment, the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1G. In certain embodiments, the methanol metabolic pathway comprises 1J, 1N, 1O and 1G. In some embodiments, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1G. In one embodiment, the

methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F and 1G. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0049] In certain embodiments, the methanol metabolic pathway comprises 1G and 1H. In certain embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1E, 1G and 1H. In some embodiments, the methanol metabolic pathway comprises 1A, 1B, 1C, 1D, 1F, 1G and 1H. In some embodiments, the methanol metabolic pathway comprises 1J, 1C, 1D, 1E, 1G and 1H. In one embodiment, the methanol metabolic pathway comprises 1J, 1C, 1D, 1F, 1G and 1H. In another embodiment, the methanol metabolic pathway comprises 1J, 1L, 1G and 1H. In yet another embodiment, the methanol metabolic pathway comprises 1J, 1M, 1N, 1O, 1G and 1H. In certain embodiments, the methanol metabolic pathway comprises 1J, 1N, 1O, 1G and 1H. In some embodiments, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E, 1G and 1H. In one embodiment, the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F, 1G and 1H. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0050] In certain embodiments, the formation of 5-hydroxymethylglutathione from formaldehyde is spontaneous (see, e.g., FIG. 1, step M). In some embodiments, the formation of 5-hydroxymethylglutathione from formaldehyde is catalyzed by a S-(hydroxymethyl)glutathione synthase (see, e.g., FIG. 1, step M). In certain embodiments, the formation of methylene-THF from formaldehyde is spontaneous (see, e.g., FIG. 1, step K). In certain embodiments, the formation of methylene-THF from formaldehyde is catalyzed by a formaldehyde activating enzyme (see, e.g., FIG. 1, step K).

[0051] In certain embodiments, the organism comprises two, three, four, five, six or seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises two exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises three exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises four exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises five exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises six exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme.

[0052] Any non-naturally occurring eukaryotic organism comprising a methanol metabolic pathway and engineered to comprise a methanol metabolic pathway enzyme, such as those provided herein, can be engineered to further comprise one or more BDO pathway enzymes.

[0053] In certain embodiments, the non-naturally occurring microbial organism further comprises a BDO pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO. In certain embodiments, the BDO pathway enzyme is selected from the group consisting of a succinyl-CoA transferase or a succinyl-CoA syn-

thetase; a succinyl-CoA reductase (aldehyde forming); a 4-hydroxybutyrate dehydrogenase; a 4-hydroxybutyrate kinase; a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming); a 1,4-butanediol dehydrogenase; a succinate reductase; a succinyl-CoA reductase (alcohol forming); a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase; a 4-hydroxybutyrate reductase; a 4-hydroxybutyryl-phosphate reductase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).

[0054] In some embodiments, the non-naturally occurring microbial organisms having a BDO pathway include a set of BDO pathway enzymes.

[0055] Enzymes, genes and methods for engineering pathways from succinate and succinyl-CoA to various products, such as BDO, into a microorganism, are now known in the art (see, e.g., U.S. Publ. No. 2011/0201089). A set of BDO pathway enzymes represents a group of enzymes that can convert succinate to BDO as shown in FIG. 2. The additional reducing equivalents obtained from the methanol metabolic pathways, as disclosed herein, improve the yields of all these products when utilizing carbohydrate-based feedstock. For example, BDO can be produced from succinyl-CoA via previously disclosed pathways (see for example, Burk et al., WO 2008/115840). Exemplary enzymes for the conversion succinyl-CoA to BDO include succinyl-CoA reductase (aldehyde forming) (FIG. 2, Step B), 4-hydroxybutyrate dehydrogenase (FIG. 2, Step C), 4-hydroxybutyrate kinase (FIG. 2, Step D), phosphotrans-4-hydroxybutyrylase (FIG. 2, Step E), 4-hydroxybutyryl-CoA reductase (aldehyde forming) (FIG. 2, Step F), 1,4-butanediol dehydrogenase (FIG. 2, Step G), succinyl-CoA reductase (alcohol forming) (FIG. 1, Step I), 4-hydroxybutyryl-CoA transferase (FIG. 2, Step J), 4-hydroxybutyryl-CoA synthetase (FIG. 2, Step J), 4-hydroxybutyryl-phosphate reductase (FIG. 2, Step L), 4-hydroxybutyrate reductase (FIG. 2, Step K), and 4-hydroxybutyryl-CoA reductase (alcohol forming) (FIG. 2, Step M). Succinate reductase (FIG. 2, Step H) can be additionally useful in converting succinate directly to the BDO pathway intermediate, succinate semialdehyde.

[0056] In another aspect, provided herein is a non-naturally occurring microbial organism, comprising (1) a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol; and (2) a BDO pathway, comprising at least one exogenous nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO. In one embodiment, the at least one exogenous nucleic acid encoding the methanol metabolic pathway enzyme enhances the availability of reducing equivalents in the presence of methanol in a sufficient amount to increase the amount of BDO produced by the non-naturally microbial organism. In some embodiments, the methanol metabolic pathway comprises any of the various combinations of methanol metabolic pathway enzymes described above or elsewhere herein.

[0057] In certain embodiments, (1) the methanol metabolic pathway comprises: 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O or any combination of 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O, thereof, wherein 1A is a methanol methyltransferase; 1B is a methylenetetrahydrofolate reductase; 1C is a methylenetetrahydrofolate dehydrogenase; 1D is a methenyltetrahydrofolate cyclohydrolase; 1E is a formyltetrahydrofolate deformylase;

1F is a formyltetrahydrofolate synthetase; 1G is a formate hydrogen lyase; 1H is a hydrogenase, 1I is a formate dehydrogenase; 1J is a methanol dehydrogenase; 1K is spontaneous or formaldehyde activating enzyme; 1L is a formaldehyde dehydrogenase; 1M is spontaneous or a S-(hydroxymethyl)glutathione synthase; 1N is glutathione-dependent formaldehyde dehydrogenase and 1O is S-formylglutathione hydrolase; and (2) the BDO pathway comprises 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, 2J, 2K, 2L or 2M or any combination of 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, 2J, 2K, 2L or 2M, wherein 2A is a succinyl-CoA transferase or a succinyl-CoA synthetase; 2B is a succinyl-CoA reductase (aldehyde forming); 2C is a 4-hydroxybutyrate dehydrogenase; 2D is a 4-hydroxybutyrate kinase; 2E is a phosphotrans-4-hydroxybutyrylase; 2F is a 4-hydroxybutyryl-CoA reductase (aldehyde forming); 2G is a 1,4-butanediol dehydrogenase; 2H is a succinate reductase; 2I is a succinyl-CoA reductase (alcohol forming); 2J is a 4-hydroxybutyryl-CoA transferase or 4-hydroxybutyryl-CoA synthetase; 2K is a 4-hydroxybutyrate reductase; 2L is a 4-hydroxybutyryl-phosphate reductase; and 2M is a 4-hydroxybutyryl-CoA reductase (alcohol forming). In some embodiments, 2A is a succinyl-CoA transferase. In some embodiments, 2A is a succinyl-CoA synthetase. In some embodiments, 2J is a 4-hydroxybutyryl-CoA transferase. In some embodiments, 2J is a 4-hydroxybutyryl-CoA synthetase. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0058] In one embodiment, the BDO pathway comprises 2A. In another embodiment, the BDO pathway comprises 2B. In an embodiment, the BDO pathway comprises 2C. In another embodiment, the BDO pathway comprises 2D. In one embodiment, the BDO pathway comprises 2E. In yet another embodiment, the BDO pathway comprises 2F. In some embodiments, the BDO pathway comprises 2G. In other embodiments, the BDO pathway comprises 2H. In another embodiment, the BDO pathway comprises 2I. In one embodiment, the BDO pathway comprises 2J. In one embodiment, the BDO pathway comprises 2K. In another embodiment, the BDO pathway comprises 2L. In an embodiment, the BDO pathway comprises 2M. Any combination of two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or thirteen BDO pathway enzymes 2A, 2B, 2C, 2D, 2E, 2F, 2G, 2H, 2I, 2J, 2K, 2L and 2M is also contemplated. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0059] In some embodiments, the methanol metabolic pathway is a methanol metabolic pathway depicted in FIG. 1, and the BDO pathway is a BDO pathway depicted in FIG. 2. In certain embodiments, the formation of 5-hydroxymethylglutathione from formaldehyde is spontaneous (see, e.g., FIG. 1, step M). In some embodiments, the formation of 5-hydroxymethylglutathione from formaldehyde is catalyzed by a S-(hydroxymethyl)glutathione synthase (see, e.g., FIG. 1, step M). In certain embodiments, the formation of methylene-THF from formaldehyde is spontaneous (see, e.g., FIG. 1, step K). In certain embodiments, the formation of methylene-THF from formaldehyde is catalyzed by a formaldehyde activating enzyme (see, e.g., FIG. 1, step K).

[0060] Exemplary sets of 1,4-BDO pathway enzymes to convert succinate to BDO, according to FIG. 2, include 2A, 2B, 2C, 2D, 2E, 2F, and 2G; 2A, 2B, 2C, 2J, 2F, and 2G; 2A, 2B, 2C, 2J, and 2M; 2A, 2B, 2C, 2D, 2E, and 2M; 2A, 2B, 2C, 2K, and 2G; 2A, 2B, 2C, 2D, 2L, and 2G; 2A, 2I, 2D, 2E, 2F, and 2G; 2A, 2I, 2D, 2E, and 2M; 2A, 2I, 2J, 2F, and 2G; 2A, 2I, 2J, and 2M; 2A, 2I, 2K, and 2G; 2A, 2I, 2D, 2L and 2G; 2H, 2C, 2D, 2E, 2F, and 2G; 2H, 2C, 2J, 2F, and 2G; 2H, 2C, 2J, and 2M; 2H, 2C, 2D, 2E, and 2M; 2H, 2C, 2K, and 2G; and 2H, 2C, 2D, 2L, and 2G. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0061] In one embodiment, the BDO pathway comprises 2B, 2C, 2D, 2E, 2F, and 2G. In another embodiment, the BDO pathway comprises 2B, 2C, 2J, 2F, and 2G. In another embodiment, the BDO pathway comprises 2B, 2C, 2J, and 2M. In yet another embodiment, the BDO pathway comprises 2B, 2C, 2D, 2E, and 2M. In one embodiment, the BDO pathway comprises 2B, 2C, 2K, and 2G. In another embodiment, the BDO pathway comprises 2B, 2C, 2D, 2L, and 2G. In another embodiment, the BDO pathway comprises 2I, 2D, 2E, 2F, and 2G. In yet another embodiment, the BDO pathway comprises 2I, 2D, 2E, and 2M. In one embodiment, the BDO pathway comprises 2I, 2J, 2F, and 2G. In another embodiment, the BDO pathway comprises 2I, 2J, and 2M. In yet another embodiment, the BDO pathway comprises 2I, 2K, and 2G. In one embodiment, the BDO pathway comprises 2I, 2D, 2L and 2G. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0062] In certain embodiments, the BDO pathway further comprises 2A. In one embodiment, the BDO pathway comprises 2A, 2B, 2C, 2D, 2E, 2F, and 2G. In another embodiment, the BDO pathway comprises 2A, 2B, 2C, 2J, 2F, and 2G. In another embodiment, the BDO pathway comprises 2A, 2B, 2C, 2J, and 2M. In yet another embodiment, the BDO pathway comprises 2A, 2B, 2C, 2D, 2E, and 2M. In one embodiment, the BDO pathway comprises 2A, 2B, 2C, 2K, and 2G.

[0063] In another embodiment, the BDO pathway comprises 2A, 2B, 2C, 2D, 2L, and 2G. In another embodiment, the BDO pathway comprises 2A, 2I, 2D, 2E, 2F, and 2G. In yet another embodiment, the BDO pathway comprises 2A, 2I, 2D, 2E, and 2M. In one embodiment, the BDO pathway comprises 2A, 2I, 2J, 2F, and 2G. In another embodiment, the BDO pathway comprises 2A, 2I, 2J, and 2M. In yet another embodiment, the BDO pathway comprises 2A, 2I, 2K, and 2G. In one embodiment, the BDO pathway comprises 2A, 2I, 2D, 2L and 2G. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0064] In another embodiment, the BDO pathway comprises 2H, 2C, 2D, 2E, 2F, and 2G. In another embodiment, the BDO pathway comprises 2H, 2C, 2J, 2F, and 2G. In yet another embodiment, the BDO pathway comprises 2H, 2C, 2J, and 2M. In one embodiment, the BDO pathway comprises 2H, 2C, 2D, 2E, and 2M. In another embodiment, the BDO pathway comprises 2H, 2C, 2K, and 2G. In yet another embodiment, the BDO pathway comprises 2H, 2C, 2D, 2L, and 2G. In one embodiment, 2J is a 4-hydroxybutyryl-CoA transferase. In another embodiment, 2J is a 4-hydroxybutyryl-CoA synthetase.

[0065] In one embodiment, (1) the methanol metabolic pathway comprises: (i) 1A and 1B, (ii) 1J; or (iii) 1J and 1K; and (2) the BDO pathway comprises (a) 2B, 2C, 2D, 2E, 2F,

1C, 1D and 1E; 1J, 1K, 1C, 1D and 1F; 1I; 1A, 1B, 1C, 1D, 1E and 1I; 1A, 1B, 1C, 1D, 1F and 1I; 1J, 1C, 1D, 1E and 1I; 1J, 1C, 1D, 1F and 1I; 1J, 1L and 1I; 1J, 1M, 1N, 1O and 1I; 1J, 1N, 1O and 1I; 1J, 1K, 1C, 1D, 1E and 1I; 1J, 1K, 1C, 1D, 1F and 1I; 1G; 1A, 1B, 1C, 1D, 1E and 1G; 1A, 1B, 1C, 1D, 1F and 1G; 1J, 1C, 1D, 1E and 1G; 1J, 1C, 1D, 1F and 1G; 1J, 1L and 1G; 1J, 1M, 1N, 1O and 1G; 1J, 1N, 1O and 1G; 1J, 1K, 1C, 1D, 1E and 1G; 1J, 1K, 1C, 1D, 1F and 1G; 1G and 1H; 1A, 1B, 1C, 1D, 1E, 1G and 1H; 1A, 1B, 1C, 1D, 1F, 1G and 1H; 1J, 1C, 1D, 1E, 1G and 1H; 1J, 1C, 1D, 1F, 1G and 1H; 1J, 1L, 1G and 1H; 1J, 1M, 1N, 1O, 1G and 1H; 1J, 1N, 1O, 1G and 1H; 1J, 1K, 1C, 1D, 1E, 1G and 1H; or 1J, 1K, 1C, 1D, 1F, 1G and 1H; and (2) a BDO pathway. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is a S-(hydroxymethyl)glutathione synthase.

[0100] Any methanol metabolic pathway provided herein can be combined with any BDO pathway provided herein.

[0101] Also provided herein are exemplary pathways, which utilize formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1, step J) in the formation of intermediates of certain central metabolic pathways that can be used for the formation of biomass. One exemplary formaldehyde assimilation pathway that can utilize formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1) is shown in FIG. 3, which involves condensation of formaldehyde and D-ribulose-5-phosphate to form hexylose-6-phosphate (h6p) by hexylose-6-phosphate synthase (FIG. 3, step A). The enzyme can use Mg^{2+} or Mn^{2+} for maximal activity, although other metal ions are useful, and even non-metal-ion-dependent mechanisms are contemplated. H6p is converted into fructose-6-phosphate by 6-phospho-3-hexyloisomerase (FIG. 3, step B). Another exemplary pathway that involves the detoxification and assimilation of formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1) is shown in FIG. 4 and proceeds through dihydroxyacetone. Dihydroxyacetone synthase is a special transketolase that first transfers a glycoaldehyde group from xylulose-5-phosphate to formaldehyde, resulting in the formation of dihydroxyacetone (DHA) and glyceraldehyde-3-phosphate (G3P), which is an intermediate in glycolysis (FIG. 4, step A). The DHA obtained from DHA synthase is then further phosphorylated to form DHA phosphate by a DHA kinase (FIG. 4, step B). DHAP can be assimilated into glycolysis and several other pathways. Rather than converting formaldehyde to formate and on to CO_2 off-gassed, the pathways provided in FIGS. 3 and 4 show that carbon is assimilated, going into the final product.

[0102] Thus, in one embodiment, an organism having a methanol metabolic pathway, either alone or in combination with a BDO pathway, as provided herein, further comprises a formaldehyde assimilation pathway that utilizes formaldehyde, e.g., obtained from the oxidation of methanol, in the formation of intermediates of certain central metabolic pathways that can be used, for example, in the formation of biomass. In some embodiments, the formaldehyde assimilation pathway comprises 3A or 3B, wherein 3A is a hexylose-6-phosphate synthase and 3B is a 6-phospho-3-hexyloisomerase. In other embodiments, the formaldehyde assimilation pathway comprises 4A or 4B, wherein 4A is a dihydroxyacetone synthase and 4B is a dihydroxyacetone kinase.

[0103] In certain embodiments, provided herein is a non-naturally occurring microbial organism having a methanol

metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol dehydrogenase (1J) expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol and/or expressed in a sufficient amount to convert methanol to formaldehyde. In some embodiments, the microbial organism further comprises a formaldehyde assimilation pathway. In certain embodiments, the organism further comprises at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme expressed in a sufficient amount to produce an intermediate of glycolysis and/or a metabolic pathway that can be used, for example, in the formation of biomass. In certain embodiments, the formaldehyde assimilation pathway enzyme is selected from the group consisting of a hexylose-6-phosphate synthase (3A), 6-phospho-3-hexyloisomerase (3B), dihydroxyacetone synthase (4A) and dihydroxyacetone kinase (4B).

[0104] In one aspect, provided herein is a non-naturally occurring microbial organism, comprising (1) a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol and/or expressed in a sufficient amount to convert methanol to formaldehyde; and (2) a formaldehyde assimilation pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme expressed in a sufficient amount to produce an intermediate of glycolysis and/or a metabolic pathway that can be used, for example, in the formation of biomass. In specific embodiments, the methanol metabolic pathway comprises a methanol dehydrogenase (1J). In certain embodiments, the formaldehyde assimilation pathway enzyme is 3A, and the intermediate is a hexylose-6-phosphate, a fructose-6-phosphate, or a combination thereof. In other embodiments, the formaldehyde assimilation pathway enzyme is 3B, and the intermediate is a hexylose-6-phosphate, a fructose-6-phosphate, or a combination thereof. In yet other embodiments, the formaldehyde assimilation pathway enzyme is 3A and 3B, and the intermediate is a hexylose-6-phosphate, a fructose-6-phosphate, or a combination thereof. In some embodiments, the formaldehyde assimilation pathway enzyme is 4A, and the intermediate is a dihydroxyacetone (DHA), a dihydroxyacetone phosphate, or a combination thereof. In other embodiments, the formaldehyde assimilation pathway enzyme is 4B, and the intermediate is a DHA, a dihydroxyacetone phosphate, or a combination thereof. In yet other embodiments, the formaldehyde assimilation pathway enzyme is 4A and 4B, and the intermediate is a DHA, a dihydroxyacetone phosphate, or a combination thereof. In one embodiment, the at least one exogenous nucleic acid encoding the methanol metabolic pathway enzyme, in the presence of methanol, sufficiently enhances the availability of reducing equivalents and sufficiently increases formaldehyde assimilation to increase the production of BDO or other products described herein by the non-naturally microbial organism. In some embodiments, the methanol metabolic pathway comprises any of the various combinations of methanol metabolic pathway enzymes described above or elsewhere herein.

[0105] In certain embodiments, (1) the methanol metabolic pathway comprises: 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O or any combination of 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O, thereof, wherein 1A is a methanol methyltransferase; 1B is a methyl-

enetetrahydrofolate reductase; 1C is a methylenetetrahydrofolate dehydrogenase; 1D is a methylenetetrahydrofolate cyclohydrolase; 1E is a formyltetrahydrofolate deformylase; 1F is a formyltetrahydrofolate synthetase; 1G is a formate hydrogen lyase; 1H is a hydrogenase, 1I is a formate dehydrogenase; 1J is a methanol dehydrogenase; 1K is spontaneous or formaldehyde activating enzyme; 1L is a formaldehyde dehydrogenase; 1M is spontaneous or a S-(hydroxymethyl)glutathione synthase; 1N is glutathione-dependent formaldehyde dehydrogenase and 1O is S-formylglutathione hydrolase; and (2) the formaldehyde assimilation pathway comprises 3A, 3B or a combination thereof, wherein 3A is a hexylose-6-phosphate synthase, and 3B is a 6-phospho-3-hexyloisomerase. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase. In some embodiments, the intermediate is a hexylose-6-phosphate. In other embodiments, the intermediate is a fructose-6-phosphate. In yet other embodiments, the intermediate is a hexylose-6-phosphate and a fructose-6-phosphate.

[0106] In one embodiment, the formaldehyde assimilation pathway comprises 3A. In another embodiment, the formaldehyde assimilation pathway comprises 3B. In one embodiment, the formaldehyde assimilation pathway comprises 3A and 3B.

[0107] In some embodiments, the methanol metabolic pathway is a methanol metabolic pathway depicted in FIG. 1, and a formaldehyde assimilation pathway depicted in FIG. 3. An exemplary set of formaldehyde assimilation pathway enzymes to convert D-ribulose-5-phosphate and formaldehyde to fructose-6-phosphate (via hexylose-6-phosphate) according to FIG. 3 include 3A and 3B.

[0108] In a specific embodiment, (1) the methanol metabolic pathway comprises 1J; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In other embodiments, (1) the methanol metabolic pathway comprises 1J and 1K; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D and 1E; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D and 1F; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In another embodiment, (1) the methanol metabolic pathway comprises 1J and 1L; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N and 1O; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N and 1O; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1E; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1F; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L and 1I; and

(2) the formaldehyde assimilation pathway comprises 3A and 3B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F and 1I; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F and 1G; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L, 1O and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 3A and 3B. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In some embodiments, the intermediate is a hexylose-6-phosphate. In other embodiments, the intermediate is a fructose-6-phosphate. In yet other embodiments, the intermediate is a hexylose-6-phosphate and a fructose-6-phosphate.

[0109] In certain embodiments, (1) the methanol metabolic pathway comprises: 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O or any combination of 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, or 1O, thereof, wherein 1A is a methanol methyltransferase; 1B is a methylenetetrahydrofolate reductase; 1C is a methylenetetrahydro-

folate dehydrogenase; 1D is a methenyltetrahydrofolate cyclohydrolase; 1E is a formyltetrahydrofolate deformylase; 1F is a formyltetrahydrofolate synthetase; 1G is a formate hydrogen lyase; 1H is a hydrogenase, 1I is a formate dehydrogenase; 1J is a methanol dehydrogenase; 1K is spontaneous or formaldehyde activating enzyme; 1L is a formaldehyde dehydrogenase; 1M is spontaneous or a S-(hydroxymethyl)glutathione synthase; 1N is glutathione-dependent formaldehyde dehydrogenase and 1O is S-formylglutathione hydrolase; and (2) the formaldehyde assimilation pathway comprises 4A, 4B or a combination thereof, wherein 4A is a dihydroxyacetone synthase and 4B is a dihydroxyacetone kinase. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In other embodiments, 1M is a S-(hydroxymethyl)glutathione synthase. In some embodiments, the intermediate is a DHA. In other embodiments, the intermediate is a dihydroxyacetone phosphate. In yet other embodiments, the intermediate is a DHA and a dihydroxyacetone phosphate.

[0110] In one embodiment, the formaldehyde assimilation pathway comprises 4A. In another embodiment, the formaldehyde assimilation pathway comprises 4B. In one embodiment, the formaldehyde assimilation pathway comprises 4A and 4B.

[0111] In some embodiments, the methanol metabolic pathway is a methanol metabolic pathway depicted in FIG. 1, and a formaldehyde assimilation pathway depicted in FIG. 4. An exemplary set of formaldehyde assimilation pathway enzymes to convert xyulose-5-phosphate and formaldehyde to dihydroxyacetone-phosphate (via DHA) according to FIG. 4 include 4A and 4B.

[0112] In a specific embodiment, (1) the methanol metabolic pathway comprises 1J; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In other embodiments, (1) the methanol metabolic pathway comprises 1J and 1K; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D and 1E; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D and 1F; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In another embodiment, (1) the methanol metabolic pathway comprises 1J and 1L; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N and 1O; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N and 1O; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1E; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D and 1F; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and

4B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1I; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F and 1G; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1E, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1C, 1D, 1F, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In another embodiment, (1) the methanol metabolic pathway comprises 1J, 1L, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In yet another embodiment, (1) the methanol metabolic pathway comprises 1J, 1M, 1N, 1O, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In certain embodiments, (1) the methanol metabolic pathway comprises 1J, 1N, 1O, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1E, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In one embodiment, (1) the methanol metabolic pathway comprises 1J, 1K, 1C, 1D, 1F, 1G and 1H; and (2) the formaldehyde assimilation pathway comprises 4A and 4B. In some embodiments, 1K is spontaneous. In other embodiments, 1K is a formaldehyde activating enzyme. In some embodiments, 1M is spontaneous. In some embodiments, the intermediate is a DHA. In other embodiments, the intermediate is a dihydroxyacetone phosphate. In yet other embodiments, the intermediate is a DHA and a dihydroxyacetone phosphate.

[0113] Any methanol metabolic pathway provided herein can be combined with any formaldehyde assimilation pathway provided herein. In addition, any methanol metabolic pathway provided herein can be combined with any BDO pathway and any formaldehyde pathway provided herein.

[0114] Also provided herein are methods of producing formaldehyde comprising culturing a non-naturally occur-

ring microbial organism having a methanol metabolic pathway provided herein. In some embodiments, the methanol metabolic pathway comprises 1J. In certain embodiments, the organism is cultured in a substantially anaerobic culture medium. In specific embodiments, the formaldehyde is an intermediate that is consumed (assimilated) in the production of BDO and other products described herein.

[0115] Also provided herein are methods of producing an intermediate of glycolysis and/or a metabolic pathway that can be used, for example, in the formation of biomass, comprising culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and a formaldehyde assimilation pathway, as provided herein, under conditions and for a sufficient period of time to produce the intermediate. In some embodiments, the intermediate is a hexylose-6-phosphate. In other embodiments, the intermediate is a fructose-6-phosphate. In yet other embodiments, the intermediate is a hexylose-6-phosphate and a fructose-6-phosphate. In some embodiments, the intermediate is a DHA. In other embodiments, the intermediate is a dihydroxyacetone phosphate. In yet other embodiments, the intermediate is a DHA and a dihydroxyacetone phosphate. In some embodiments, the methanol metabolic pathway comprises 1J. In certain embodiments, the organism is cultured in a substantially anaerobic culture medium. Such biomass can also be used in methods of producing any of the products, such as the biobased products, provided elsewhere herein.

[0116] In some embodiments, the organism comprises two, three, four, five, six, seven, eight or more exogenous nucleic acids, each encoding a BDO pathway enzyme. In some embodiments, the organism comprises two exogenous nucleic acids, each encoding a BDO pathway enzyme. In some embodiments, the organism comprises three exogenous nucleic acids, each encoding a BDO pathway enzyme. In some embodiments, the organism comprises four exogenous nucleic acids, each encoding a BDO pathway enzyme. In other embodiments, the organism comprises five exogenous nucleic acids, each encoding a BDO pathway enzyme. In some embodiments, the organism comprises six exogenous nucleic acids, each encoding a BDO pathway enzyme. In other embodiments, the organism comprises seven exogenous nucleic acids, each encoding a BDO pathway enzyme. In certain embodiments, the organism comprises two, three, four, five, six or seven exogenous nucleic acids, each encoding a BDO pathway enzyme; and the organism further comprises two, three, four, five, six or seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises two exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises three exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises four exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises five exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises six exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme.

[0117] In some embodiments, the organism comprises two or more exogenous nucleic acids, each encoding a formalde-

hyde assimilation pathway enzyme. In some embodiments, the organism comprises two exogenous nucleic acids, each encoding a formaldehyde assimilation pathway enzyme. In certain embodiments, the organism comprises two exogenous nucleic acids, each encoding a formaldehyde assimilation pathway enzyme; and the organism further comprises two, three, four, five, six or seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises two exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises three exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism comprises further four exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises five exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises six exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme. In certain embodiments, the organism further comprises seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme.

[0118] In some embodiments, the at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme is a heterologous nucleic acid. In other embodiments, the at least one exogenous nucleic acid encoding a BDO pathway enzyme is a heterologous nucleic acid. In other embodiments, the at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme is a heterologous nucleic acid. In certain embodiments, the at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme is a heterologous nucleic acid, and the at least one exogenous nucleic acid encoding a BDO pathway enzyme is a heterologous nucleic acid. In other embodiments, the at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme is a heterologous nucleic acid, and the at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme is a heterologous nucleic acid.

[0119] In certain embodiments, the organism is in a substantially anaerobic culture medium.

[0120] It is understood that any of the pathways disclosed herein, as described in the Examples and exemplified in the figures, including the pathways of FIGS. 1, 2, 3 and 4 can be utilized to generate a non-naturally occurring microbial organism that produces any pathway intermediate or product, as desired. Non-limiting examples of such intermediate or products are 4-HB and BDO. As disclosed herein, such a microbial organism that produces an intermediate can be used in combination with another microbial organism expressing downstream pathway enzymes to produce a desired product. However, it is understood that a non-naturally occurring organism that produces a BDO pathway intermediate can be utilized to produce the intermediate as a desired product (e.g., 4-hydroxybutanal).

[0121] In certain embodiments, a non-naturally occurring microbial organism comprising a methanol metabolic pathway and a BDO pathway provided herein, further comprises one or more gene disruptions. In certain embodiments, the one or more gene disruptions confer increased production of BDO in the organism. In other embodiments, a non-naturally occurring microbial organism comprising a methanol metabolic pathway and a formaldehyde assimilation pathway provided herein, further comprises one or more gene disruptions.

In some embodiments, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂, amino acids, or any combination thereof, by said microbial organism. In one embodiment, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of ethanol. In another embodiment, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of glycerol. In other embodiments, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of acetate. In another embodiment, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of lactate. In one embodiment, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of formate. In another embodiment, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of CO₂. In other embodiments, the gene disruption is in an endogenous gene encoding a protein and/or enzyme involved in native production of amino acids by said microbial organism. In some embodiments, the protein or enzyme is a pyruvate decarboxylase, an ethanol dehydrogenase, a glycerol dehydrogenase, a glycerol-3-phosphatase, a glycerol-3-phosphate dehydrogenase, a lactate dehydrogenase, an acetate kinase, a phosphotransacetylase, a pyruvate oxidase, a pyruvate:quinone oxidoreductase, a pyruvate formate lyase, an alcohol dehydrogenase, a lactate dehydrogenase, a pyruvate dehydrogenase, a pyruvate formate-lyase-2-ketobutyrate formate-lyase, a pyruvate transporter, a monocarboxylate transporter, a NADH dehydrogenase, a cytochrome oxidase, a pyruvate kinase, or any combination thereof. In certain embodiments, the organism comprises from one to twenty-five gene disruptions. In other embodiments, the organism comprises from one to twenty gene disruptions. In some embodiments, the organism comprises from one to fifteen gene disruptions. In other embodiments, the organism comprises from one to ten gene disruptions. In some embodiments, the organism comprises from one to five gene disruptions. In certain embodiments, the organism comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 gene disruptions or more.

[0122] In other embodiments, a non-naturally occurring microbial organism comprising a methanol metabolic pathway and a BDO pathway provided herein, further comprises one or more endogenous proteins or enzymes involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂ and/or amino acids by said microbial organism, wherein said one or more endogenous proteins or enzymes has attenuated protein or enzyme activity and/or expression levels. In some embodiments, a non-naturally occurring microbial organism comprising a methanol metabolic pathway and a formaldehyde assimilation pathway provided herein, further comprises one or more endogenous proteins or enzymes involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂ and/or amino acids by said microbial organism, wherein said one or more endogenous proteins or enzymes has attenuated protein or enzyme activity and/or expression levels. In one embodiment the endogenous protein or enzyme is a pyruvate decarboxylase, an ethanol dehydrogenase, a glycerol dehydrogenase, a glycerol-3-phosphatase, a glycerol-3-phosphate dehydrogenase, a lactate dehydrogenase, an acetate kinase, a phosphotransacetyl-

lase, a pyruvate oxidase, a pyruvate:quinone oxidoreductase, a pyruvate formate lyase, an alcohol dehydrogenase, a lactate dehydrogenase, a pyruvate dehydrogenase, a pyruvate formate-lyase-2-ketobutyrate formate-lyase, a pyruvate transporter, a monocarboxylate transporter, a NADH dehydrogenase, a cytochrome oxidase, a pyruvate kinase, or any combination thereof.

[0123] Each of the non-naturally occurring alterations provided herein result in increased production and an enhanced level of BDO, for example, during the exponential growth phase of the microbial organism, compared to a strain that does not contain such metabolic alterations, under appropriate culture conditions. Appropriate conditions include, for example, those disclosed herein, including conditions such as particular carbon sources or reactant availabilities and/or adaptive evolution.

[0124] Given the teachings and guidance provided herein, those skilled in the art will understand that to introduce a metabolic alteration, such as attenuation of an enzyme, it can be necessary to disrupt the catalytic activity of the one or more enzymes involved in the reaction. Alternatively, a metabolic alteration can include disrupting expression of a regulatory protein or cofactor necessary for enzyme activity or maximal activity. Furthermore, genetic loss of a cofactor necessary for an enzymatic reaction can also have the same effect as a disruption of the gene encoding the enzyme. Disruption can occur by a variety of methods including, for example, deletion of an encoding gene or incorporation of a genetic alteration in one or more of the encoding gene sequences. The encoding genes targeted for disruption can be one, some, or all of the genes encoding enzymes involved in the catalytic activity. For example, where a single enzyme is involved in a targeted catalytic activity, disruption can occur by a genetic alteration that reduces or eliminates the catalytic activity of the encoded gene product. Similarly, where the single enzyme is multimeric, including heteromeric, disruption can occur by a genetic alteration that reduces or destroys the function of one or all subunits of the encoded gene products. Destruction of activity can be accomplished by loss of the binding activity of one or more subunits required to form an active complex, by destruction of the catalytic subunit of the multimeric complex or by both. Other functions of multimeric protein association and activity also can be targeted in order to disrupt a metabolic reaction of the invention. Such other functions are well known to those skilled in the art. Similarly, a target enzyme activity can be reduced or eliminated by disrupting expression of a protein or enzyme that modifies and/or activates the target enzyme, for example, a molecule required to convert an apoenzyme to a holoenzyme. Further, some or all of the functions of a single polypeptide or multimeric complex can be disrupted according to the invention in order to reduce or abolish the catalytic activity of one or more enzymes involved in a reaction or metabolic modification of the invention. Similarly, some or all of enzymes involved in a reaction or metabolic modification of the invention can be disrupted so long as the targeted reaction is reduced or eliminated.

[0125] Given the teachings and guidance provided herein, those skilled in the art also will understand that an enzymatic reaction can be disrupted by reducing or eliminating reactions encoded by a common gene and/or by one or more orthologs of that gene exhibiting similar or substantially the same activity. Reduction of both the common gene and all orthologs can lead to complete abolishment of any catalytic activity of a

targeted reaction. However, disruption of either the common gene or one or more orthologs can lead to a reduction in the catalytic activity of the targeted reaction sufficient to promote coupling of growth to product biosynthesis. Exemplified herein are both the common genes encoding catalytic activities for a variety of metabolic modifications as well as their orthologs. Those skilled in the art will understand that disruption of some or all of the genes encoding an enzyme of a targeted metabolic reaction can be practiced in the methods of the invention and incorporated into the non-naturally occurring microbial organisms of the invention in order to achieve the increased production of fatty alcohol, fatty aldehyde or fatty acid or growth-coupled product production.

[0126] Given the teachings and guidance provided herein, those skilled in the art also will understand that enzymatic activity or expression can be attenuated using well known methods. Reduction of the activity or amount of an enzyme can mimic complete disruption if the reduction causes activity of the enzyme to fall below a critical level that is normally required for the pathway to function. Reduction of enzymatic activity by various techniques rather than disruption can be important for an organism's viability. Methods of reducing enzymatic activity that result in similar or identical effects of a gene disruption include, but are not limited to: reducing gene transcription or translation; destabilizing mRNA, protein or catalytic RNA; and mutating a gene that affects enzyme kinetics. Natural or imposed regulatory controls can also accomplish enzyme attenuation including: promoter replacement; loss or alteration of transcription factors; introduction of inhibitory RNAs or peptides such as siRNA, antisense RNA, RNA or peptide/small-molecule binding aptamers, ribozymes, aptazymes and riboswitches; and addition of drugs and other chemicals that reduce or disrupt enzymatic activity such as gene splicing.

[0127] One of ordinary skill in the art will also recognize that attenuation of an enzyme can be done at various levels. For example, at the gene level, mutations causing a partial or complete null phenotype or epistatic genetic effects that mask the activity of a gene product can be used to attenuate an enzyme. At the gene expression level, methods for attenuation include: coupling transcription to an endogenous or exogenous inducer such as IPTG, then adding low or 0 levels of inducer during the production phase; introducing or modifying positive or negative regulators; modify histone acetylation/deacetylation in region where gene is integrated; introducing a transposition to disrupt a promoter or a regulatory gene; flipping of a transposable element or promoter region; deleting one allele resulting in loss of heterozygosity in a diploid organism; introducing nucleic acids that increase RNA degradation; or in bacteria, for example, introduction of a tmRNA tag, which can lead to RNA degradation and ribosomal stalling. At the translational level, attenuation can include: introducing rare codons to limit translation; introducing RNA interference molecules that block translation; modifying regions outside the coding sequence, such as introducing secondary structure into UTR regions to block translation or reduce efficiency of translation; adding RNAase sites for rapid transcript degradation; introducing antisense RNA oligomers or antisense transcripts; introducing RNA or peptide aptamers, ribozymes, aptazymes, riboswitches; or introducing translational regulatory elements involving RNA structure that can prevent or reduce translation that can be controlled by the presence or absence of small molecules. At the level of enzyme localization and/or longevity, enzyme

attenuation can include: adding a degradation tag for faster protein turnover; or adding a localization tag that results in the enzyme being localized to a compartment where it would not be able to react normally. At the level of post-translational regulation, enzyme attenuation can include: increasing intracellular concentration of known inhibitors; or modifying post-translational modified sites. At the level of enzyme activity, enzyme attenuation can include: adding endogenous or exogenous inhibitor, such as a target-specific drug, to reduce enzyme activity; limiting availability of essential cofactors, such as B12, for an enzyme that require it; chelating a metal ion that is required for activity; or introducing a dominant negative mutation.

[0128] In some embodiments, microaerobic designs can be used based on the growth-coupled formation of the desired product. To examine this, production cones can be constructed for each strategy by first maximizing and, subsequently minimizing the product yields at different rates of biomass formation feasible in the network. If the rightmost boundary of all possible phenotypes of the mutant network is a single point, it implies that there is a unique optimum yield of the product at the maximum biomass formation rate possible in the network. In other cases, the rightmost boundary of the feasible phenotypes is a vertical line, indicating that at the point of maximum biomass the network can make any amount of the product in the calculated range, including the lowest amount at the bottommost point of the vertical line. Such designs are given a low priority.

[0129] Accordingly, also provided herein is a non-naturally occurring microbial organism having metabolic modifications coupling BDO production to growth of the organism, where the metabolic modifications includes disruption of one or more genes selected from the genes encoding proteins and/or enzymes provided herein.

[0130] Each of the strains can be supplemented with additional deletions if it is determined that the strain designs do not sufficiently increase the production of BDO and/or couple the formation of the product with biomass formation. Alternatively, some other enzymes not known to possess significant activity under the growth conditions can become active due to adaptive evolution or random mutagenesis. Such activities can also be knocked out. However, gene deletions provided herein allow the construction of strains exhibiting high-yield production of BDO, including growth-coupled production of BDO.

[0131] In another aspect, provided herein is a method for producing BDO, comprising culturing any one of the non-naturally occurring microbial organisms comprising a methanol metabolic pathway and an BDO pathway provided herein under conditions and for a sufficient period of time to produce BDO. In certain embodiments, the organism is cultured in a substantially anaerobic culture medium.

[0132] Provided herein are methods for producing BDO, comprising culturing an organism provided herein under conditions and for a sufficient period of time to produce BDO. In some embodiments, the method comprises culturing, for a sufficient period of time to produce BDO, a non-naturally occurring microbial organism, comprising (1) a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol; and (2) a BDO pathway, comprising at least one exogenous

nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO.

[0133] In certain embodiments of the methods provided herein, the organism further comprises at least one nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO. In some embodiments, the nucleic acid is an exogenous nucleic acid. In other embodiments, the nucleic acid is an endogenous nucleic acid. In some embodiments, the organism further comprises one or more gene disruptions provided herein that confer increased production of BDO in the organism. In certain embodiments, the one or more gene disruptions occurs in an endogenous gene encoding a protein or enzyme involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂ and/or amino acids by said microbial organism. In other embodiments, the organism further comprises one or more endogenous proteins or enzymes involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂ and/or amino acids by said microbial organism, wherein said one or more endogenous proteins or enzymes has attenuated protein or enzyme activity and/or expression levels. In certain embodiments, the organism is a Crabtree positive, eukaryotic organism, and the organism is cultured in a culture medium comprising glucose. In certain embodiments, the organism comprises from one to twenty-five gene disruptions. In other embodiments, the organism comprises from one to twenty gene disruptions. In some embodiments, the organism comprises from one to fifteen gene disruptions. In other embodiments, the organism comprises from one to ten gene disruptions. In some embodiments, the organism comprises from one to five gene disruptions. In certain embodiments, the organism comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25 gene disruptions or more.

[0134] In an additional embodiment, the invention provides a non-naturally occurring microbial organism having a BDO pathway, formaldehyde assimilation pathway and/or methanol metabolic pathway, wherein the non-naturally occurring microbial organism comprises at least one exogenous nucleic acid encoding an enzyme or protein that converts a substrate to a product. By way of example, in FIG. 1, the substrate of 1J is methanol, and the product is formaldehyde; the substrate of 1L is formaldehyde, and the product is formate; and so forth. One skilled in the art will understand that these are merely exemplary and that any of the substrate-product pairs disclosed herein suitable to produce a desired product and for which an appropriate activity is available for the conversion of the substrate to the product can be readily determined by one skilled in the art based on the teachings herein. Thus, provided herein are non-naturally occurring microbial organisms containing at least one exogenous nucleic acid encoding an enzyme or protein, where the enzyme or protein converts the substrates and products of a methanol metabolic pathway, such as that shown in FIG. 1; a BDO pathway, such as that shown in FIG. 2; and/or a formaldehyde assimilation pathway, such as that shown in FIG. 3 or 4.

[0135] While generally described herein as a microbial organism that contains a BDO pathway, formaldehyde assimilation pathway and/or a methanol metabolic pathway, it is understood that provided herein are also non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding a BDO, formaldehyde assimilation and/or a methanol metabolic pathway enzyme expressed in a sufficient amount to produce an intermediate of a BDO

pathway, formaldehyde assimilation pathway and/or a methanol metabolic pathway intermediate. For example, as disclosed herein, a BDO pathway is exemplified in FIG. 2. Therefore, in addition to a microbial organism containing a BDO pathway that produces BDO, also provided herein is a non-naturally occurring microbial organism comprising at least one exogenous nucleic acid encoding a BDO pathway enzyme, where the microbial organism produces a BDO pathway intermediate.

[0136] In some embodiments, the carbon feedstock and other cellular uptake sources such as phosphate, ammonia, sulfate, chloride and other halogens can be chosen to alter the isotopic distribution of the atoms present in BDO and/or 4-HB or any BDO and/or 4-HB pathway intermediate. The various carbon feedstock and other uptake sources enumerated above will be referred to herein, collectively, as "uptake sources." Uptake sources can provide isotopic enrichment for any atom present in the product BDO and/or 4-HB or BDO and/or 4-HB pathway intermediate, or for side products generated in reactions diverging away from a BDO and/or 4-HB pathway. Isotopic enrichment can be achieved for any target atom including, for example, carbon, hydrogen, oxygen, nitrogen, sulfur, phosphorus, chlorine or other halogens.

[0137] In some embodiments, the uptake sources can be selected to alter the carbon-12, carbon-13, and carbon-14 ratios. In some embodiments, the uptake sources can be selected to alter the oxygen-16, oxygen-17, and oxygen-18 ratios. In some embodiments, the uptake sources can be selected to alter the hydrogen, deuterium, and tritium ratios. In some embodiments, the uptake sources can be selected to alter the nitrogen-14 and nitrogen-15 ratios. In some embodiments, the uptake sources can be selected to alter the sulfur-32, sulfur-33, sulfur-34, and sulfur-35 ratios. In some embodiments, the uptake sources can be selected to alter the phosphorus-31, phosphorus-32, and phosphorus-33 ratios. In some embodiments, the uptake sources can be selected to alter the chlorine-35, chlorine-36, and chlorine-37 ratios.

[0138] In some embodiments, the isotopic ratio of a target atom can be varied to a desired ratio by selecting one or more uptake sources. An uptake source can be derived from a natural source, as found in nature, or from a man-made source, and one skilled in the art can select a natural source, a man-made source, or a combination thereof, to achieve a desired isotopic ratio of a target atom. An example of a man-made uptake source includes, for example, an uptake source that is at least partially derived from a chemical synthetic reaction. Such isotopically enriched uptake sources can be purchased commercially or prepared in the laboratory and/or optionally mixed with a natural source of the uptake source to achieve a desired isotopic ratio. In some embodiments, a target isotopic ratio of an uptake source can be obtained by selecting a desired origin of the uptake source as found in nature. For example, as discussed herein, a natural source can be a biobased derived from or synthesized by a biological organism or a source such as petroleum-based products or the atmosphere. In some such embodiments, a source of carbon, for example, can be selected from a fossil fuel-derived carbon source, which can be relatively depleted of carbon-14, or an environmental or atmospheric carbon source, such as CO₂, which can possess a larger amount of carbon-14 than its petroleum-derived counterpart.

[0139] Isotopic enrichment is readily assessed by mass spectrometry using techniques known in the art such as Stable Isotope Ratio Mass Spectrometry (SIRMS) and Site-Specific

Natural Isotopic Fractionation by Nuclear Magnetic Resonance (SNIF-NMR). Such mass spectral techniques can be integrated with separation techniques such as liquid chromatography (LC) and/or high performance liquid chromatography (HPLC).

[0140] The unstable carbon isotope carbon-14 or radiocarbon makes up for roughly 1 in 10^{12} carbon atoms in the earth's atmosphere and has a half-life of about 5700 years. The stock of carbon is replenished in the upper atmosphere by a nuclear reaction involving cosmic rays and ordinary nitrogen (^{14}N). Fossil fuels contain no carbon-14, as it decayed long ago. Burning of fossil fuels lowers the atmospheric carbon-14 fraction, the so-called "Suess effect".

[0141] Methods of determining the isotopic ratios of atoms in a compound are well known to those skilled in the art. Isotopic enrichment is readily assessed by mass spectrometry using techniques known in the art such as accelerated mass spectrometry (AMS), Stable Isotope Ratio Mass Spectrometry (SIRMS) and Site-Specific Natural Isotopic Fractionation by Nuclear Magnetic Resonance (SNIF-NMR). Such mass spectral techniques can be integrated with separation techniques such as liquid chromatography (LC), high performance liquid chromatography (HPLC) and/or gas chromatography, and the like.

[0142] In the case of carbon, ASTM D6866 was developed in the United States as a standardized analytical method for determining the biobased content of solid, liquid, and gaseous samples using radiocarbon dating by the American Society for Testing and Materials (ASTM) International. The standard is based on the use of radiocarbon dating for the determination of a product's biobased content. ASTM D6866 was first published in 2004, and the current active version of the standard is ASTM D6866-11 (effective Apr. 1, 2011). Radiocarbon dating techniques are well known to those skilled in the art, including those described herein.

[0143] The biobased content of a compound is estimated by the ratio of carbon-14 (^{14}C) to carbon-12 (^{12}C). Specifically, the Fraction Modern (Fm) is computed from the expression: $Fm = (S - B) / (M - B)$, where B, S and M represent the $^{14}\text{C}/^{12}\text{C}$ ratios of the blank, the sample and the modern reference, respectively. Fraction Modern is a measurement of the deviation of the $^{14}\text{C}/^{12}\text{C}$ ratio of a sample from "Modern." Modern is defined as 95% of the radiocarbon concentration (in AD 1950) of National Bureau of Standards (NBS) Oxalic Acid I (i.e., standard reference materials (SRM) 4990b) normalized to $\delta^{13}\text{C}_{VPDB} = -19$ per mil (Olsson, *The use of Oxalic acid as a Standard*, in *Radiocarbon Variations and Absolute Chronology*, Nobel Symposium, 12th Proc., John Wiley & Sons, New York (1970)). Mass spectrometry results, for example, measured by ASM, are calculated using the internationally agreed upon definition of 0.95 times the specific activity of NBS Oxalic Acid I (SRM 4990b) normalized to $\delta^{13}\text{C}_{VPDB} = -19$ per mil. This is equivalent to an absolute (AD 1950) $^{14}\text{C}/^{12}\text{C}$ ratio of $1.176 \pm 0.010 \times 10^{-12}$ (Karlen et al., *Arkiv Geofysik*, 4:465-471 (1968)). The standard calculations take into account the differential uptake of one isotope with respect to another, for example, the preferential uptake in biological systems of C^{12} over C^{13} over C^{14} , and these corrections are reflected as a Fm corrected for δ^{13} .

[0144] An oxalic acid standard (SRM 4990b or HOx 1) was made from a crop of 1955 sugar beet. Although there were 1000 lbs made, this oxalic acid standard is no longer commercially available. The Oxalic Acid II standard (HOx 2; N.I.S.T designation SRM 4990 C) was made from a crop of

1977 French beet molasses. In the early 1980's, a group of 12 laboratories measured the ratios of the two standards. The ratio of the activity of Oxalic acid II to 1 is 1.2933 ± 0.001 (the weighted mean). The isotopic ratio of HOx II is -17.8 per mille. ASTM D6866-11 suggests use of the available Oxalic Acid II standard SRM 4990 C (Hox2) for the modern standard (see discussion of original vs. currently available oxalic acid standards in Mann, *Radiocarbon*, 25(2):519-527 (1983)). A Fm=0% represents the entire lack of carbon-14 atoms in a material, thus indicating a fossil (for example, petroleum based) carbon source. A Fm=100%, after correction for the post-1950 injection of carbon-14 into the atmosphere from nuclear bomb testing, indicates an entirely modern carbon source. As described herein, such a "modern" source includes biobased sources.

[0145] As described in ASTM D6866, the percent modern carbon (pMC) can be greater than 100% because of the continuing but diminishing effects of the 1950s nuclear testing programs, which resulted in a considerable enrichment of carbon-14 in the atmosphere as described in ASTM D6866-11. Because all sample carbon-14 activities are referenced to a "pre-bomb" standard, and because nearly all new biobased products are produced in a post-bomb environment, all pMC values (after correction for isotopic fraction) must be multiplied by 0.95 (as of 2010) to better reflect the true biobased content of the sample. A biobased content that is greater than 103% suggests that either an analytical error has occurred, or that the source of biobased carbon is more than several years old.

[0146] ASTM D6866 quantifies the biobased content relative to the material's total organic content and does not consider the inorganic carbon and other non-carbon containing substances present. For example, a product that is 50% starch-based material and 50% water would be considered to have a Biobased Content=100% (50% organic content that is 100% biobased) based on ASTM D6866. In another example, a product that is 50% starch-based material, 25% petroleum-based, and 25% water would have a Biobased Content=66.7% (75% organic content but only 50% of the product is biobased). In another example, a product that is 50% organic carbon and is a petroleum-based product would be considered to have a Biobased Content=0% (50% organic carbon but from fossil sources). Thus, based on the well known methods and known standards for determining the biobased content of a compound or material, one skilled in the art can readily determine the biobased content and/or prepared downstream products that utilize of the invention having a desired biobased content.

[0147] Applications of carbon-14 dating techniques to quantify bio-based content of materials are known in the art (Currie et al., *Nuclear Instruments and Methods in Physics Research B*, 172:281-287 (2000)). For example, carbon-14 dating has been used to quantify bio-based content in terephthalate-containing materials (Colonna et al., *Green Chemistry*, 13:2543-2548 (2011)). Notably, polypropylene terephthalate (PPT) polymers derived from renewable 1,3-propanediol and petroleum-derived terephthalic acid resulted in Fm values near 30% (i.e., since $^{3/11}$ of the polymeric carbon derives from renewable 1,3-propanediol and $^{4/11}$ from the fossil end member terephthalic acid) (Currie et al., supra, 2000). In contrast, polybutylene terephthalate polymer derived from both renewable BDO and renewable terephthalic acid resulted in bio-based content exceeding 90% (Colonna et al., supra, 2011).

[0148] Accordingly, in some embodiments, the present invention provides BDO and/or 4-HB or a BDO and/or 4-HB pathway intermediate thereof that has a carbon-12, carbon-13, and carbon-14 ratio that reflects an atmospheric carbon, also referred to as environmental carbon, uptake source. For example, in some aspects the BDO and/or 4-HB or a BDO and/or 4-HB intermediate thereof can have an Fm value of at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or as much as 100%. In some such embodiments, the uptake source is CO₂. In some embodiments, the present invention provides BDO and/or 4-HB or a BDO and/or 4-HB intermediate thereof that has a carbon-12, carbon-13, and carbon-14 ratio that reflects petroleum-based carbon uptake source. In this aspect, the BDO and/or 4-HB or a BDO and/or 4-HB intermediate thereof can have an Fm value of less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, less than 50%, less than 45%, less than 40%, less than 35%, less than 30%, less than 25%, less than 20%, less than 15%, less than 10%, less than 5%, less than 2% or less than 1%. In some embodiments, the present invention provides BDO and/or 4-HB or a BDO and/or 4-HB intermediate thereof that has a carbon-12, carbon-13, and carbon-14 ratio that is obtained by a combination of an atmospheric carbon uptake source with a petroleum-based uptake source. Using such a combination of uptake sources is one way by which the carbon-12, carbon-13, and carbon-14 ratio can be varied, and the respective ratios would reflect the proportions of the uptake sources.

[0149] Further, the present invention relates to biologically produced BDO and/or 4-HB or BDO and/or 4-HB intermediate thereof as disclosed herein, and to the products derived therefrom, wherein the BDO and/or 4-HB or a BDO and/or 4-HB intermediate thereof has a carbon-12, carbon-13, and carbon-14 isotope ratio of about the same value as the CO₂ that occurs in the environment. For example, in some aspects the invention provides bioderived BDO and/or 4-HB or a bioderived BDO and/or 4-HB intermediate thereof having a carbon-12 versus carbon-13 versus carbon-14 isotope ratio of about the same value as the CO₂ that occurs in the environment, or any of the other ratios disclosed herein. It is understood, as disclosed herein, that a product can have a carbon-12 versus carbon-13 versus carbon-14 isotope ratio of about the same value as the CO₂ that occurs in the environment, or any of the ratios disclosed herein, wherein the product is generated from bioderived BDO and/or 4-HB or a bioderived BDO and/or 4-HB intermediate thereof as disclosed herein, wherein the bioderived product is chemically modified to generate a final product. Methods of chemically modifying a bioderived product of BDO and/or 4-HB, or an intermediate thereof, to generate a desired product are well known to those skilled in the art, as described herein. The invention further provides plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as poly-4-hydroxybutyrate (P4HB) or co-polymers thereof, poly(tetramethylene ether) glycol (PTMEG) (also referred to as PTMO, polytetramethylene oxide) and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, having a carbon-12 versus carbon-13 versus carbon-14 isotope ratio of about the same value as the CO₂ that occurs in the environment, wherein the plastics, elastic fibers, polyure-

thanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, are generated directly from or in combination with bioderived BDO and/or 4-HB or a bioderived BDO and/or 4-HB intermediate thereof as disclosed herein.

[0150] BDO and/or 4-HB are chemicals used in commercial and industrial applications. Non-limiting examples of such applications include production of plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like. Moreover, BDO and/or 4-HB are also used as a raw material in the production of a wide range of products including plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like. Accordingly, in some embodiments, the invention provides biobased plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, comprising one or more bioderived BDO and/or 4-HB or bioderived BDO and/or 4-HB intermediate thereof produced by a non-naturally occurring microorganism of the invention or produced using a method disclosed herein.

[0151] As used herein, the term “bioderived” means derived from or synthesized by a biological organism and can be considered a renewable resource since it can be generated by a biological organism. Such a biological organism, in particular the microbial organisms of the invention disclosed herein, can utilize feedstock or biomass, such as, sugars or carbohydrates obtained from an agricultural, plant, bacterial, or animal source. Alternatively, the biological organism can utilize atmospheric carbon. As used herein, the term “biobased” means a product as described above that is composed, in whole or in part, of a bioderived compound of the invention. A biobased or bioderived product is in contrast to a petroleum derived product, wherein such a product is derived from or synthesized from petroleum or a petrochemical feedstock.

[0152] In some embodiments, the invention provides plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, comprising bioderived BDO and/or 4-HB or bioderived BDO and/or 4-HB intermediate thereof, wherein the bioderived BDO and/or 4-HB or bioderived BDO and/or 4-HB intermediate thereof includes all or part of the BDO and/or 4-HB or BDO and/or 4-HB intermediate thereof used in the production of plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like. Thus, in some aspects, the invention provides a biobased plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, comprising at least 2%, at least 3%, at least 5%, at

least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98% or 100% bioderived BDO and/or 4-HB or bioderived BDO and/or 4-HB intermediate thereof as disclosed herein. Additionally, in some aspects, the invention provides a bio-based plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, wherein the BDO and/or 4-HB or BDO and/or 4-HB intermediate thereof used in its production is a combination of bioderived and petroleum derived BDO and/or 4-HB or BDO and/or 4-HB intermediate thereof. For example, a bio-based plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, can be produced using 50% bioderived BDO and/or 4-HB and 50% petroleum derived BDO and/or 4-HB or other desired ratios such as 60%/40%, 70%/30%, 80%/20%, 90%/10%, 95%/5%, 100%/0%, 40%/60%, 30%/70%, 20%/80%, 10%/90% of bioderived/petroleum derived precursors, so long as at least a portion of the product comprises a bioderived product produced by the microbial organisms disclosed herein. It is understood that methods for producing plastics, elastic fibers, polyurethanes, polyesters, including polyhydroxyalkanoates such as P4HB or co-polymers thereof, PTMEG and polyurethane-polyurea copolymers, referred to as spandex, elastane or Lycra™, nylons, and the like, using the bioderived BDO and/or 4-HB or bioderived BDO and/or 4-HB intermediate thereof of the invention are well known in the art.

[0153] In one embodiment, the product is a plastic. In one embodiment, the product is an elastic fiber. In one embodiment, the product is a polyurethane. In one embodiment, the product is a polyester. In one embodiment, the product is a polyhydroxyalkanoate. In one embodiment, the product is a poly-4-hydroxybutyrate. In one embodiment, the product is a co-polymer of poly-4-hydroxybutyrate. In one embodiment, the product is a poly(tetramethylene ether) glycol. In one embodiment, the product is a polyurethane-polyurea copolymer. In one embodiment, the product is a spandex. In one embodiment, the product is an elastane. In one embodiment, the product is a Lycra™. In one embodiment, the product is a nylon.

[0154] In some embodiments, provided herein is a culture medium comprising bioderived BDO. In some embodiments, the bioderived BDO is produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO pathway, as provided herein. In certain embodiments, the bioderived BDO has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide uptake source. In one embodiment, the culture medium is separated from a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO pathway.

[0155] In other embodiments, provided herein is a bioderived BDO. In some embodiments, the bioderived BDO is produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO pathway, as provided herein. In certain embodiments, the bioderived BDO has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide

uptake source. In some embodiments, the bioderived BDO has an Fm value of at least 80%, at least 85%, at least 90%, at least 95% or at least 98%. In certain embodiments, the bioderived BDO is a component of culture medium.

[0156] In certain embodiments, provided herein is a composition comprising a bioderived BDO provided herein, for example, a bioderived BDO produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO pathway, as provided herein. In some embodiments, the composition further comprises a compound other than said bioderived BDO. In certain embodiments, the compound other than said bioderived BDO is a trace amount of a cellular portion of a non-naturally occurring microbial organism having a methanol metabolic pathway and a BDO pathway, as provided herein.

[0157] In some embodiments, provided herein is a biobased product comprising a bioderived BDO provided herein. In certain embodiments, the biobased product is a plastic, elastic fiber, polyurethane, polyester, polyhydroxyalkanoate, poly-4-hydroxybutyrate, co-polymer of poly-4-hydroxybutyrate, poly(tetramethylene ether) glycol, polyurethane-polyurea copolymer, spandex, elastane, Lycra™, or nylon. In certain embodiments, the biobased product comprises at least 5% bioderived BDO. In certain embodiments, the biobased product is (i) a polymer, THF or a THF derivative, or GBL or a GBL derivative; (ii) a plastic, elastic fiber, polyurethane, polyester, polyhydroxyalkanoate, poly-4-hydroxybutyrate, co-polymer of poly-4-hydroxybutyrate, poly(tetramethylene ether) glycol, polyurethane-polyurea copolymer, spandex, elastane, Lycra™, or nylon; (iii) a polymer, a resin, a fiber, a bead, a granule, a pellet, a chip, a plastic, a polyester, a thermoplastic polyester, a molded article, an injection-molded article, an injection-molded part, an automotive part, an extrusion resin, an electrical part and a casing; and optionally where the biobased product is reinforced or filled and further where the biobased product is glass-reinforced or -filled or mineral-reinforced or -filled; (iv) a polymer, wherein the polymer comprises polybutylene terephthalate (PBT); (v) a polymer, wherein the polymer comprises PBT and the biobased product is a resin, a fiber, a bead, a granule, a pellet, a chip, a plastic, a polyester, a thermoplastic polyester, a molded article, an injection-molded article, an injection-molded part, an automotive part, an extrusion resin, an electrical part and a casing; and optionally where the biobased product is reinforced or filled and further where the biobased product is glass-reinforced or -filled or mineral-reinforced or -filled; (vi) a THF or a THF derivative, wherein the THF derivative is polytetramethylene ether glycol (PTMEG), a polyester ether (COPE) or a thermoplastic polyurethane; (viii) a THF derivative, wherein the THF derivative comprises a fiber; or (ix) a GBL or a GBL derivative, wherein the GBL derivative is a pyrrolidone. In certain embodiments, the biobased product comprises at least 10% bioderived BDO. In some embodiments, the biobased product comprises at least 20% bioderived BDO. In other embodiments, the biobased product comprises at least 30% bioderived BDO. In some embodiments, the biobased product comprises at least 40% bioderived BDO. In other embodiments, the biobased product comprises at least 50% bioderived BDO. In one embodiment, the biobased product comprises a portion of said bioderived BDO as a repeating unit. In another embodiment, provided herein is a molded product obtained by molding the biobased product provided herein. In other embodiments, provided herein is a process for producing a biobased product

provided herein, comprising chemically reacting said bio-derived—BDO with itself or another compound in a reaction that produces said biobased product. In certain embodiments, provided herein is a polymer comprising or obtained by converting the bio-derived BDO. In other embodiments, provided herein is a method for producing a polymer, comprising chemically or enzymatically converting the bio-derived BDO to the polymer. In yet other embodiments, provided herein is a composition comprising the bio-derived BDO, or a cell lysate or culture supernatant thereof.

[0158] In some embodiments, provided herein is a culture medium comprising bio-derived 4-HB. In some embodiments, the bio-derived 4-HB is produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO and/or 4-HB pathway, as provided herein. In certain embodiments, the bio-derived 4-HB has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide uptake source.

[0159] In one embodiment, the culture medium is separated from a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO and/or 4-HB pathway.

[0160] In other embodiments, provided herein is a bio-derived 4-HB. In some embodiments, the bio-derived 4-HB is produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO and/or 4-HB pathway, as provided herein. In certain embodiments, the bio-derived 4-HB has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide uptake source. In some embodiments, the bio-derived 4-HB has an Fm value of at least 80%, at least 85%, at least 90%, at least 95% or at least 98%. In certain embodiments, the bio-derived 4-HB is a component of culture medium.

[0161] In certain embodiments, provided herein is a composition comprising a bio-derived 4-HB provided herein, for example, a bio-derived 4-HB produced by culturing a non-naturally occurring microbial organism having a methanol metabolic pathway and BDO and/or 4-HB pathway, as provided herein. In some embodiments, the composition further comprises a compound other than said bio-derived 4-HB. In certain embodiments, the compound other than said bio-derived 4-HB is a trace amount of a cellular portion of a non-naturally occurring microbial organism having a methanol metabolic pathway and a BDO and/or 4-HB pathway, as provided herein.

[0162] In some embodiments, provided herein is a biobased product comprising a bio-derived 4-HB provided herein. In certain embodiments, the biobased product is a plastic, elastic fiber, polyurethane, polyester, polyhydroxyalkanoate, poly-4-hydroxybutyrate, co-polymer of poly-4-hydroxybutyrate, poly(tetramethylene ether) glycol, polyurethane-polyurea copolymer, spandex, elastane, Lycra™, or nylon. In certain embodiments, the biobased product comprises at least 5% bio-derived 4-HB. In certain embodiments, the biobased product comprises at least 10% bio-derived 4-HB. In some embodiments, the biobased product comprises at least 20% bio-derived 4-HB. In other embodiments, the biobased product comprises at least 30% bio-derived 4-HB. In some embodiments, the biobased product comprises at least 40% bio-derived 4-HB. In other embodiments, the biobased product comprises at least 50% bio-derived 4-HB. In one embodiment, the biobased product comprises a portion of said bio-derived 4-HB as a repeating unit. In another embodiment, provided herein is a molded product obtained by molding the biobased product provided herein. In other embodiments, provided

herein is a process for producing a biobased product provided herein, comprising chemically reacting said bio-derived 4-HB with itself or another compound in a reaction that produces said biobased product.

[0163] The invention is described herein with general reference to the metabolic reaction, reactant or product thereof, or with specific reference to one or more nucleic acids or genes encoding an enzyme associated with or catalyzing the referenced metabolic reaction, reactant or product. Unless otherwise expressly stated herein, those skilled in the art will understand that reference to a reaction also constitutes reference to the reactants and products of the reaction. Similarly, unless otherwise expressly stated herein, reference to a reactant or product also references the reaction and that reference to any of these metabolic constituents also references the gene or genes encoding the enzymes that catalyze the referenced reaction, reactant or product. Likewise, given the well known fields of metabolic biochemistry, enzymology and genomics, reference herein to a gene or encoding nucleic acid also constitutes a reference to the corresponding encoded enzyme and the reaction it catalyzes, or a protein associated with the reaction, as well as the reactants and products of the reaction.

[0164] The production of 4-HB via biosynthetic modes using the microbial organisms of the invention is particularly useful because it can produce monomeric 4-HB. The non-naturally occurring microbial organisms of the invention and their biosynthesis of 4-HB and BDO family compounds also is particularly useful because the 4-HB product can be (1) secreted; (2) can be devoid of any derivatizations such as Coenzyme A; (3) avoids thermodynamic changes during biosynthesis; (4) allows direct biosynthesis of BDO, and (5) allows for the spontaneous chemical conversion of 4-HB to γ -butyrolactone (GBL) in acidic pH medium. This latter characteristic also is particularly useful for efficient chemical synthesis or biosynthesis of BDO family compounds such as BDO and/or tetrahydrofuran (THF), for example.

[0165] Microbial organisms generally lack the capacity to synthesize 4-HB and therefore any of the compounds disclosed herein to be within the BDO family of compounds or known by those in the art to be within the BDO family of compounds. Moreover, organisms having all of the requisite metabolic enzymatic capabilities are not known to produce 4-HB from the enzymes described and biochemical pathways exemplified herein. Rather, with the possible exception of a few anaerobic microorganisms described further below, the microorganisms having the enzymatic capability to use 4-HB as a substrate to produce, for example, succinate. In contrast, the non-naturally occurring microbial organisms of the invention can generate BDO and/or 4-HB as a product. The biosynthesis of 4-HB in its monomeric form is not only particularly useful in chemical synthesis of BDO family of compounds, it also allows for the further biosynthesis of BDO family compounds and avoids altogether chemical synthesis procedures.

[0166] The non-naturally occurring microbial organisms of the invention that can produce BDO and/or 4-HB are produced by ensuring that a host microbial organism includes functional capabilities for the complete biochemical synthesis of at least one BDO and/or 4-HB biosynthetic pathway of provided herein. Ensuring at least one requisite BDO and/or 4-HB biosynthetic pathway confers BDO and/or 4-HB biosynthesis capability onto the host microbial organism.

[0167] The organisms and methods are described herein with general reference to the metabolic reaction, reactant or

product thereof, or with specific reference to one or more nucleic acids or genes encoding an enzyme associated with or catalyzing, or a protein associated with, the referenced metabolic reaction, reactant or product. Unless otherwise expressly stated herein, those skilled in the art will understand that reference to a reaction also constitutes reference to the reactants and products of the reaction. Similarly, unless otherwise expressly stated herein, reference to a reactant or product also references the reaction, and reference to any of these metabolic constituents also references the gene or genes encoding the enzymes that catalyze or proteins involved in the referenced reaction, reactant or product. Likewise, given the well known fields of metabolic biochemistry, enzymology and genomics, reference herein to a gene or encoding nucleic acid also constitutes a reference to the corresponding encoded enzyme and the reaction it catalyzes or a protein associated with the reaction as well as the reactants and products of the reaction.

[0168] The non-naturally occurring microbial organisms described herein can be produced by introducing expressible nucleic acids encoding one or more of the enzymes or proteins participating in one or more methanol metabolic, formaldehyde assimilation and/or BDO biosynthetic pathways. Depending on the host microbial organism chosen for biosynthesis, nucleic acids for some or all of a particular methanol metabolic, formaldehyde assimilation and/or BDO biosynthetic pathway can be expressed. For example, if a chosen host is deficient in one or more enzymes or proteins for a desired metabolic, assimilation or biosynthetic pathway, then expressible nucleic acids for the deficient enzyme(s) or protein(s) are introduced into the host for subsequent exogenous expression. Alternatively, if the chosen host exhibits endogenous expression of some pathway genes, but is deficient in others, then an encoding nucleic acid is needed for the deficient enzyme(s) or protein(s) to achieve BDO biosynthesis and/or methanol metabolism. Thus, a non-naturally occurring microbial organism described herein can be produced by introducing exogenous enzyme or protein activities to obtain a desired metabolic pathway and/or a desired biosynthetic pathway can be obtained by introducing one or more exogenous enzyme or protein activities that, together with one or more endogenous enzymes or proteins, produces a desired product such as BDO.

[0169] Host microbial organisms can be selected from, and the non-naturally occurring microbial organisms generated in, for example, bacteria, yeast, fungus or any of a variety of other microorganisms applicable to fermentation processes. Exemplary bacteria include species selected from *Escherichia coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*, *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*. Exemplary yeasts or fungi include species selected from *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger*, *Pichia pastoris*, *Rhizopus arrhizus*, *Rhizopus oryzae*, and the like. *E. coli* is a particularly useful host organism since it is a well characterized microbial organism suitable for genetic engineering. Other particularly useful host organisms include yeast such as *Saccharomyces cerevisiae*. It is understood that any suitable microbial host organ-

ism can be used to introduce metabolic and/or genetic modifications to produce a desired product.

[0170] Depending on the BDO biosynthetic, methanol metabolic and/or formaldehyde assimilation pathway constituents of a selected host microbial organism, the non-naturally occurring microbial organisms provided herein will include at least one exogenously expressed BDO, formaldehyde assimilation and/or methanol metabolic pathway-encoding nucleic acid and up to all encoding nucleic acids for one or more BDO biosynthetic pathways, formaldehyde assimilation pathways and/or methanol metabolic pathways. For example, BDO biosynthesis can be established in a host deficient in a pathway enzyme or protein through exogenous expression of the corresponding encoding nucleic acid. In a host deficient in all enzymes or proteins of a BDO pathway, exogenous expression of all enzyme or proteins in the pathway can be included, although it is understood that all enzymes or proteins of a pathway can be expressed even if the host contains at least one of the pathway enzymes or proteins. For example, exogenous expression of all enzymes or proteins in a pathway for production of BDO can be included. The same holds true for the methanol metabolic pathways and formaldehyde assimilation pathways provided herein.

[0171] Given the teachings and guidance provided herein, those skilled in the art will understand that the number of encoding nucleic acids to introduce in an expressible farm will, at least, parallel the BDO, formaldehyde assimilation, and methanol metabolic pathway deficiencies of the selected host microbial organism. Therefore, a non-naturally occurring microbial organism of the invention can have one, two, three, four, five, six, seven, eight, or up to all nucleic acids encoding the enzymes or proteins constituting a methanol metabolic pathway, formaldehyde assimilation and/or BDO biosynthetic pathway disclosed herein. In some embodiments, the non-naturally occurring microbial organisms also can include other genetic modifications that facilitate or optimize BDO biosynthesis, formaldehyde assimilation and/or methanol metabolism or that confer other useful functions onto the host microbial organism. One such other functionality can include, for example, augmentation of the synthesis of one or more of the BDO pathway precursors.

[0172] Generally, a host microbial organism is selected such that it produces the precursor of a BDO pathway, either as a naturally produced molecule or as an engineered product that either provides de novo production of a desired precursor or increased production of a precursor naturally produced by the host microbial organism. A host organism can be engineered to increase production of a precursor, as disclosed herein. In addition, a microbial organism that has been engineered to produce a desired precursor can be used as a host organism and further engineered to express enzymes or proteins of a BDO pathway.

[0173] In some embodiments, a non-naturally occurring microbial organism provided herein is generated from a host that contains the enzymatic capability to synthesize BDO, assimilate formaldehyde and/or metabolize methanol. In this specific embodiment it can be useful to increase the synthesis or accumulation of a BDO pathway product, formaldehyde assimilation pathway product and/or methanol metabolic pathway product (e.g., reducing equivalents and/or formaldehyde) to, for example, drive BDO pathway reactions toward BDO production. Increased synthesis or accumulation can be accomplished by, for example, overexpression of nucleic acids encoding one or more of the above-described BDO,

formaldehyde assimilation and/or methanol metabolic pathway enzymes or proteins. Over expression the enzyme(s) and/or protein(s) of the BDO pathway, formaldehyde assimilation, and/or methanol metabolic pathway can occur, for example, through exogenous expression of the endogenous gene(s), or through exogenous expression of the heterologous gene(s). Therefore, naturally occurring organisms can be readily generated to be non-naturally occurring microbial organisms, for example, producing BDO through overexpression of one, two, three, four, five, six, seven, eight, up to all nucleic acids encoding BDO biosynthetic pathway, and/or methanol metabolic pathway enzymes or proteins. Naturally occurring organisms can also be readily generated to be non-naturally occurring microbial organisms, for example, assimilating formaldehyde, through overexpression of one, two, three, four, five, six, seven, eight, up to all nucleic acids encoding formaldehyde assimilation pathway, and/or methanol metabolic pathway enzymes or proteins. In addition, a non-naturally occurring organism can be generated by mutagenesis of an endogenous gene that results in an increase in activity of an enzyme in the BDO, formaldehyde assimilation and/or methanol metabolic pathway biosynthetic pathway.

[0174] In particularly useful embodiments, exogenous expression of the encoding nucleic acids is employed. Exogenous expression confers the ability to custom tailor the expression and/or regulatory elements to the host and application to achieve a desired expression level that is controlled by the user. However, endogenous expression also can be utilized in other embodiments such as by removing a negative regulatory effector or induction of the gene's promoter when linked to an inducible promoter or other regulatory element. Thus, an endogenous gene having a naturally occurring inducible promoter can be up-regulated by providing the appropriate inducing agent, or the regulatory region of an endogenous gene can be engineered to incorporate an inducible regulatory element, thereby allowing the regulation of increased expression of an endogenous gene at a desired time. Similarly, an inducible promoter can be included as a regulatory element for an exogenous gene introduced into a non-naturally occurring microbial organism.

[0175] It is understood that, in methods provided herein, any of the one or more exogenous nucleic acids can be introduced into a microbial organism to produce a non-naturally occurring microbial organism provided herein. The nucleic acids can be introduced so as to confer, for example, a BDO biosynthetic, formaldehyde assimilation and/or methanol metabolic pathway onto the microbial organism. Alternatively, encoding nucleic acids can be introduced to produce an intermediate microbial organism having the biosynthetic capability to catalyze some of the required reactions to confer BDO biosynthetic, formaldehyde assimilation and/or methanol metabolic capability. For example, a non-naturally occurring microbial organism having a BDO biosynthetic pathway, formaldehyde assimilation pathway and/or methanol metabolic pathway can comprise at least two exogenous nucleic acids encoding desired enzymes or proteins. Thus, it is understood that any combination of two or more enzymes or proteins of a biosynthetic pathway, formaldehyde assimilation pathway and/or metabolic pathway can be included in a non-naturally occurring microbial organism provided herein. Similarly, it is understood that any combination of three or more enzymes or proteins of a biosynthetic pathway, formaldehyde assimilation pathway and/or metabolic pathway can

be included in a non-naturally occurring microbial organism of the invention, as desired, so long as the combination of enzymes and/or proteins of the desired biosynthetic pathway, formaldehyde assimilation pathway and/or metabolic pathway results in production of the corresponding desired product. Similarly, any combination of four or more enzymes or proteins of a biosynthetic pathway, formaldehyde assimilation pathway and/or methanol metabolic pathway as disclosed herein can be included in a non-naturally occurring microbial organism provided herein, as desired, so long as the combination of enzymes and/or proteins of the desired biosynthetic, assimilation and/or metabolic pathway results in production of the corresponding desired product.

[0176] In addition to the metabolism of methanol, assimilation of formaldehyde, and biosynthesis of BDO, as described herein, the non-naturally occurring microbial organisms and methods provided also can be utilized in various combinations with each other and with other microbial organisms and methods well known in the art to achieve product biosynthesis by other routes. For example, one alternative to produce BDO, other than use of the BDO producers is through addition of another microbial organism capable of converting a BDO pathway intermediate to BDO. One such procedure includes, for example, the fermentation of a microbial organism that produces a BDO pathway intermediate. The BDO pathway intermediate can then be used as a substrate for a second microbial organism that converts the BDO pathway intermediate to BDO. The BDO pathway intermediate can be added directly to another culture of the second organism or the original culture of the BDO pathway intermediate producers can be depleted of these microbial organisms by, for example, cell separation, and then subsequent addition of the second organism to the fermentation broth can be utilized to produce the final product without intermediate purification steps.

[0177] In other embodiments, the non-naturally occurring microbial organisms and methods of the invention can be assembled in a wide variety of subpathways to achieve biosynthesis of, for example, BDO. In these embodiments, biosynthetic pathways for a desired product can be segregated into different microbial organisms, and the different microbial organisms can be co-cultured to produce the final product. In such a biosynthetic scheme, the product of one microbial organism is the substrate for a second microbial organism until the final product is synthesized. For example, the biosynthesis of BDO can be accomplished by constructing a microbial organism that contains biosynthetic pathways for conversion of one pathway intermediate to another pathway intermediate or the product. Alternatively, BDO also can be biosynthetically produced from microbial organisms through co-culture or co-fermentation using two organisms in the same vessel, where the first microbial organism produces a BDO intermediate and the second microbial organism converts the intermediate to BDO.

[0178] Given the teachings and guidance provided herein, those skilled in the art will understand that a wide variety of combinations and permutations exist for the non-naturally occurring microbial organisms and methods together with other microbial organisms, with the co-culture of other non-naturally occurring microbial organisms having subpathways and with combinations of other chemical and/or biochemical procedures well known in the art to produce BDO and/or metabolize methanol.

[0179] Sources of encoding nucleic acids for a BDO, formaldehyde assimilation, or methanol metabolic pathway enzyme or protein can include, for example, any species where the encoded gene product is capable of catalyzing the referenced reaction. Such species include both prokaryotic and eukaryotic organisms including, but not limited to, bacteria, including archaea and eubacteria, and eukaryotes, including yeast, plant, insect, animal, and mammal, including human. Exemplary species for such sources include, for example, *Escherichia coli*, *Saccharomyces cerevisiae*, *Saccharomyces kluyveri*, *Candida boidinii*, *Clostridium kluyveri*, *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *Clostridium saccharoperbutylacetonicum*, *Clostridium perfringens*, *Clostridium difficile*, *Clostridium botulinum*, *Clostridium tyrobutyricum*, *Clostridium tetanomorphum*, *Clostridium tetani*, *Clostridium propionicum*, *Clostridium aminobutyricum*, *Clostridium subterminale*, *Clostridium sticklandii*, *Ralstonia eutropha*, *Mycobacterium bovis*, *Mycobacterium tuberculosis*, *Porphyromonas gingivalis*, *Arabidopsis thaliana*, *Thermus thermophilus*, *Pseudomonas* species, including *Pseudomonas aeruginosa*, *Pseudomonas putida*, *Pseudomonas stutzeri*, *Pseudomonas fluorescens*, *Homo sapiens*, *Oryctolagus cuniculus*, *Rhodobacter spaeroioides*, *Thermoanaerobacter brockii*, *Metallosphaera sedula*, *Leuconostoc mesenteroides*, *Chloroflexus aurantiacus*, *Roseiflexus castenholzii*, *Erythrobacter*, *Simmondsia chinensis*, *Acinetobacter* species, including *Acinetobacter calcoaceticus* and *Acinetobacter baylyi*, *Porphyromonas gingivalis*, *Sulfolobus tokodaii*, *Sulfolobus solfataricus*, *Sulfolobus acidocaldarius*, *Bacillus subtilis*, *Bacillus cereus*, *Bacillus megaterium*, *Bacillus brevis*, *Bacillus pumilus*, *Rattus norvegicus*, *Klebsiella pneumonia*, *Klebsiella oxytoca*, *Euglena gracilis*, *Treponema denticola*, *Moorella thermoacetica*, *Thermotoga maritima*, *Halobacterium salinarum*, *Geobacillus stearothermophilus*, *Aeropyrum pernix*, *Sus scrofa*, *Caenorhabditis elegans*, *Corynebacterium glutamicum*, *Acidaminococcus fermentans*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptococcus thermophilus*, *Enterobacter aerogenes*, *Candida*, *Aspergillus terreus*, *Pedicoccus pentosaceus*, *Zymomonas mobilis*, *Acetobacter pasteurians*, *Kluyveromyces lactis*, *Eubacterium barkeri*, *Bacteroides capillosus*, *Anaerotruncus colihominis*, *Natranerobium thermophilum*, *Campylobacter jejuni*, *Haemophilus influenzae*, *Serratia marcescens*, *Citrobacter amalonaticus*, *Myxococcus xanthus*, *Fusobacterium nucleatum*, *Penicillium chrysogenum*, marine gamma proteobacterium, butyrate-producing bacterium, *Nocardia iowensis*, *Nocardia farcinica*, *Streptomyces griseus*, *Schizosaccharomyces pombe*, *Geobacillus thermoglucosidasius*, *Salmonella typhimurium*, *Vibrio cholera*, *Heliobacter pylori*, *Nicotiana tabacum*, *Oryza sativa*, *Haloferax mediterranei*, *Agrobacterium tumefaciens*, *Achromobacter denitrificans*, *Fusobacterium nucleatum*, *Streptomyces clavuligenus*, *Acinetobacter baumannii*, *Mus musculus*, *Lachanea kluyveri*, *Trichomonas vaginalis*, *Trypanosoma brucei*, *Pseudomonas stutzeri*, *Bradyrhizobium japonicum*, *Mesorhizobium loti*, *Bos taurus*, *Nicotiana glutinosa*, *Vibrio vulnificus*, *Selenomonas ruminantium*, *Vibrio parahaemolyticus*, *Archaeoglobus fulgidus*, *Haloarcula marismortui*, *Pyrobaculum aerophilum*, *Mycobacterium smegmatis* MC2 155, *Mycobacterium avium* subsp. *paratuberculosis* K-10, *Mycobacterium marinum* M Tsukamurella paurometabola DSM 20162, *Cyanobium* PCC7001, *Dictyostelium discoideum* AX4, as well as other exemplary species disclosed herein or available as source organisms for corresponding genes. How-

ever, with the complete genome sequence available for now more than 550 species (with more than half of these available on public databases such as the NCBI), including 395 microorganism genomes and a variety of yeast, fungi, plant, and mammalian genomes, the identification of genes encoding the requisite BDO or 4-hydroxybutyrate biosynthetic activity for one or more genes in related or distant species, including for example, homologues, orthologs, paralogs and non-orthologous gene displacements of known genes, and the interchange of genetic alterations between organisms is routine and well known in the art. Accordingly, the metabolic alterations allowing biosynthesis of BDO or 4-HB, metabolism of methanol and/or assimilation of formaldehyde described herein with reference to a particular organism such as *E. coli* can be readily applied to other microorganisms, including prokaryotic and eukaryotic organisms alike. Given the teachings and guidance provided herein, those skilled in the art will know that a metabolic alteration exemplified in one organism can be applied equally to other organisms.

[0180] In some instances, such as when an alternative BDO biosynthetic, formaldehyde assimilation and/or methanol metabolic pathway exists in an unrelated species, BDO biosynthesis, formaldehyde assimilation and/or methanol metabolism can be conferred onto the host species by, for example, exogenous expression of a paralog or paralogs from the unrelated species that catalyzes a similar, yet non-identical metabolic reaction to replace the referenced reaction. Because certain differences among metabolic networks exist between different organisms, those skilled in the art will understand that the actual gene usage between different organisms may differ. However, given the teachings and guidance provided herein, those skilled in the art also will understand that the teachings and methods provided herein can be applied to all microbial organisms using the cognate metabolic alterations to those exemplified herein to construct a microbial organism in a species of interest that will synthesize BDO, assimilate formaldehyde and/or metabolize methanol.

[0181] Methods for constructing and testing the expression levels of a non-naturally occurring BDO-producing host can be performed, for example, by recombinant and detection methods well known in the art. Such methods can be found described in, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Third Ed., Cold Spring Harbor Laboratory, New York (2001); and Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1999).

[0182] Exogenous nucleic acid sequences involved in a pathway for metabolism of methanol, assimilation of formaldehyde and/or production of BDO can be introduced stably or transiently into a host cell using techniques well known in the art including, but not limited to, conjugation, electroporation, chemical transformation, transduction, transfection, and ultrasound transformation. For exogenous expression in *E. coli* or other prokaryotic cells, some nucleic acid sequences in the genes or cDNAs of eukaryotic nucleic acids can encode targeting signals such as an N-terminal mitochondrial or other targeting signal, which can be removed before transformation into prokaryotic host cells, if desired. For example, removal of a mitochondrial leader sequence led to increased expression in *E. coli* (Hoffmeister et al., *J. Biol. Chem.* 280:4329-4338 (2005)). For exogenous expression in yeast or other eukaryotic cells, genes can be expressed in the cytosol without the addition of leader sequence, or can be targeted to mitochondrion or other organelles, or targeted for

secretion, by the addition of a suitable targeting sequence such as a mitochondrial targeting or secretion signal suitable for the host cells. Thus, it is understood that appropriate modifications to a nucleic acid sequence to remove or include a targeting sequence can be incorporated into an exogenous nucleic acid sequence to impart desirable properties. Furthermore, genes can be subjected to codon optimization with techniques well known in the art to achieve optimized expression of the proteins.

[0183] An expression vector or vectors can be constructed to include one or more BDO biosynthetic, formaldehyde assimilation and/or methanol metabolic pathway encoding nucleic acids as exemplified herein operably linked to expression control sequences functional in the host organism. Expression vectors applicable for use in the microbial host organisms provided include, for example, plasmids, phage vectors, viral vectors, episomes and artificial chromosomes, including vectors and selection sequences or markers operable for stable integration into a host chromosome. Additionally, the expression vectors can include one or more selectable marker genes and appropriate expression control sequences. Selectable marker genes also can be included that, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like which are well known in the art. When two or more exogenous encoding nucleic acids are to be co-expressed, both nucleic acids can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector expression, the encoding nucleic acids can be operationally linked to one common expression control sequence or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The transformation of exogenous nucleic acid sequences involved in a metabolic or synthetic pathway can be confirmed using methods well known in the art. Such methods include, for example, nucleic acid analysis such as Northern blots or polymerase chain reaction (PCR) amplification of mRNA, or immunoblotting for expression of gene products, or other suitable analytical methods to test the expression of an introduced nucleic acid sequence or its corresponding gene product. It is understood by those skilled in the art that the exogenous nucleic acid is expressed in a sufficient amount to produce the desired product, and it is further understood that expression levels can be optimized to obtain sufficient expression using methods well known in the art and as disclosed herein.

[0184] Suitable purification and/or assays to test, e.g., for the production of BDO can be performed using well known methods. Suitable replicates such as triplicate cultures can be grown for each engineered strain to be tested. For example, product and byproduct formation in the engineered production host can be monitored. The final product and intermediates, and other organic compounds, can be analyzed by methods such as HPLC (High Performance Liquid Chromatography), GC-MS (Gas Chromatography-Mass Spectroscopy) and LC-MS (Liquid Chromatography-Mass Spectroscopy) or other suitable analytical methods using routine procedures well known in the art. The release of product in the fermentation broth can also be tested with the culture supernatant. Byproducts and residual glucose can be quantified by HPLC using, for example, a refractive index detector for glucose and alcohols, and a UV detector for organic acids

(Lin et al., *Biotechnol. Bioeng.* 90:775-779 (2005)), or other suitable assay and detection methods well known in the art. The individual enzyme or protein activities from the exogenous DNA sequences can also be assayed using methods well known in the art.

[0185] The BDO can be separated from other components in the culture using a variety of methods well known in the art. Such separation methods include, for example, extraction procedures as well as methods that include continuous liquid-liquid extraction, pervaporation, membrane filtration, membrane separation, reverse osmosis, electrodialysis, distillation, crystallization, centrifugation, extractive filtration, ion exchange chromatography, size exclusion chromatography, adsorption chromatography, and ultrafiltration. All of the above methods are well known in the art.

[0186] Any of the non-naturally occurring microbial organisms described herein can be cultured to produce and/or secrete the biosynthetic products, or intermediates thereof. For example, the BDO producers can be cultured for the biosynthetic production of BDO. Accordingly, in some embodiments, the invention provides culture medium having a BDO, formaldehyde assimilation and/or methanol metabolic pathway intermediate described herein. In some aspects, the culture medium can also be separated from the non-naturally occurring microbial organisms provided herein that produced the BDO, formaldehyde assimilation and/or methanol metabolic pathway intermediate. Methods for separating a microbial organism from culture medium are well known in the art. Exemplary methods include filtration, flocculation, precipitation, centrifugation, sedimentation, and the like.

[0187] In certain embodiments, for example, for the production of BDO, the recombinant strains are cultured in a medium with carbon source and other essential nutrients. It is sometimes desirable and can be highly desirable to maintain anaerobic conditions in the fermenter to reduce the cost of the overall process. Such conditions can be obtained, for example, by first sparging the medium with nitrogen and then sealing the flasks with a septum and crimp-cap. For strains where growth is not observed anaerobically, microaerobic or substantially anaerobic conditions can be applied by perforating the septum with a small hole for limited aeration. Exemplary anaerobic conditions have been described previously and are well-known in the art. Exemplary aerobic and anaerobic conditions are described, for example, in U.S. Publ. No. 2009/0047719. Fermentations can be performed in a batch, fed-batch or continuous manner, as disclosed herein. Fermentations can also be conducted in two phases, if desired. The first phase can be aerobic to allow for high growth and therefore high productivity, followed by an anaerobic phase of high BDO yields.

[0188] If desired, the pH of the medium can be maintained at a desired pH, in particular neutral pH, such as a pH of around 7 by addition of a base, such as NaOH or other bases, or acid, as needed to maintain the culture medium at a desirable pH. The growth rate can be determined by measuring optical density using a spectrophotometer (600 nm), and the glucose uptake rate by monitoring carbon source depletion over time.

[0189] The growth medium, can include, for example, any carbohydrate source which can supply a source of carbon to the non-naturally occurring microorganism. Such sources include, for example, sugars such as glucose, xylose, arabinose, galactose, mannose, fructose, sucrose and starch; or

glycerol, alone as the sole source of carbon or in combination with other carbon sources described herein or known in the art. In one embodiment, the carbon source is a sugar. In one embodiment, the carbon source is a sugar-containing biomass. In some embodiments, the sugar is glucose. In one embodiment, the sugar is xylose. In another embodiment, the sugar is arabinose. In one embodiment, the sugar is galactose. In another embodiment, the sugar is fructose. In other embodiments, the sugar is sucrose. In one embodiment, the sugar is starch. In certain embodiments, the carbon source is glycerol. In some embodiments, the carbon source is crude glycerol. In one embodiment, the carbon source is crude glycerol without treatment. In other embodiments, the carbon source is glycerol and glucose. In another embodiment, the carbon source is methanol and glycerol. In one embodiment, the carbon source is carbon dioxide. In one embodiment, the carbon source is formate. In one embodiment, the carbon source is methane. In one embodiment, the carbon source is methanol. In one embodiment, the carbon source is chemo-electro-generated carbon (see, e.g., Liao et al. (2012) *Science* 335:1596). In one embodiment, the chemo-electro-generated carbon is methanol. In one embodiment, the chemo-electro-generated carbon is formate. In one embodiment, the chemo-electro-generated carbon is formate and methanol. In one embodiment, the carbon source is a sugar and methanol. In another embodiment, the carbon source is a sugar and glycerol. In other embodiments, the carbon source is a sugar and crude glycerol. In yet other embodiments, the carbon source is a sugar and crude glycerol without treatment. In one embodiment, the carbon source is a sugar-containing biomass and methanol. In another embodiment, the carbon source is a sugar-containing biomass and glycerol. In other embodiments, the carbon source is a sugar-containing biomass and crude glycerol. In yet other embodiments, the carbon source is a sugar-containing biomass and crude glycerol without treatment. Other sources of carbohydrate include, for example, renewable feedstocks and biomass. Exemplary types of biomasses that can be used as feedstocks in the methods of the invention include cellulosic biomass, hemicellulosic biomass and lignin feedstocks or portions of feedstocks. Such biomass feedstocks contain, for example, carbohydrate substrates useful as carbon sources such as glucose, xylose, arabinose, galactose, mannose, fructose and starch. Given the teachings and guidance provided herein, those skilled in the art will understand that renewable feedstocks and biomass other than those exemplified above also can be used for culturing the microbial organisms provided herein for the production of BDO and other pathway intermediates.

[0190] In one embodiment, the carbon source is glycerol. In certain embodiments, the glycerol carbon source is crude glycerol or crude glycerol without further treatment. In a further embodiment, the carbon source comprises glycerol or crude glycerol, and also sugar or a sugar-containing biomass, such as glucose. In a specific embodiment, the concentration of glycerol in the fermentation broth is maintained by feeding crude glycerol, or a mixture of crude glycerol and sugar (e.g., glucose). In certain embodiments, sugar is provided for sufficient strain growth. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of from 200:1 to 1:200. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of from 100:1 to 1:100. In some embodiments, the sugar (e.g., glucose) is provided at a molar con-

centration ratio of glycerol to sugar of from 100:1 to 5:1. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of from 50:1 to 5:1. In certain embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 100:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 90:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 80:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 70:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 60:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 50:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 40:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 30:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 20:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 10:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 5:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 2:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:1. In certain embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:100. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:90. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:80. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:70. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:60. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:50. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:40. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:30. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:20. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:10. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:5. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of glycerol to sugar of 1:2. In certain embodiments of the ratios provided above, the sugar is a sugar-containing biomass. In certain other embodiments of the ratios provided above, the glycerol is a crude glycerol or a crude glycerol without further treatment. In other embodiments of the ratios provided above, the sugar is a sugar-containing biomass, and the glycerol is a crude glycerol or a crude glycerol without further treatment.

[0191] Crude glycerol can be a by-product produced in the production of biodiesel, and can be used for fermentation without any further treatment. Biodiesel production methods include (1) a chemical method wherein the glycerol-group of vegetable oils or animal oils is substituted by low-carbon

alcohols such as methanol or ethanol to produce a corresponding fatty acid methyl esters or fatty acid ethyl esters by transesterification in the presence of acidic or basic catalysts; (2) a biological method where biological enzymes or cells are used to catalyze transesterification reaction and the corresponding fatty acid methyl esters or fatty acid ethyl esters are produced; and (3) a supercritical method, wherein transesterification reaction is carried out in a supercritical solvent system without any catalysts. The chemical composition of crude glycerol can vary with the process used to produce biodiesel, the transesterification efficiency, recovery efficiency of the biodiesel, other impurities in the feedstock, and whether methanol and catalysts were recovered. For example, the chemical compositions of eleven crude glycerol collected from seven Australian biodiesel producers reported that glycerol content ranged between 38% and 96%, with some samples including more than 14% methanol and 29% ash. In certain embodiments, the crude glycerol comprises from 5% to 99% glycerol. In some embodiments, the crude glycerol comprises from 10% to 90% glycerol. In some embodiments, the crude glycerol comprises from 10% to 80% glycerol. In some embodiments, the crude glycerol comprises from 10% to 70% glycerol. In some embodiments, the crude glycerol comprises from 10% to 60% glycerol. In some embodiments, the crude glycerol comprises from 10% to 50% glycerol. In some embodiments, the crude glycerol comprises from 10% to 40% glycerol. In some embodiments, the crude glycerol comprises from 10% to 30% glycerol. In some embodiments, the crude glycerol comprises from 10% to 20% glycerol. In some embodiments, the crude glycerol comprises from 80% to 90% glycerol. In some embodiments, the crude glycerol comprises from 70% to 90% glycerol. In some embodiments, the crude glycerol comprises from 60% to 90% glycerol. In some embodiments, the crude glycerol comprises from 50% to 90% glycerol. In some embodiments, the crude glycerol comprises from 40% to 90% glycerol. In some embodiments, the crude glycerol comprises from 30% to 90% glycerol. In some embodiments, the crude glycerol comprises from 20% to 90% glycerol. In some embodiments, the crude glycerol comprises from 20% to 40% glycerol. In some embodiments, the crude glycerol comprises from 40% to 60% glycerol. In some embodiments, the crude glycerol comprises from 60% to 80% glycerol. In some embodiments, the crude glycerol comprises from 50% to 70% glycerol. In one embodiment, the glycerol comprises 5% glycerol. In one embodiment, the glycerol comprises 10% glycerol. In one embodiment, the glycerol comprises 15% glycerol. In one embodiment, the glycerol comprises 20% glycerol. In one embodiment, the glycerol comprises 25% glycerol. In one embodiment, the glycerol comprises 30% glycerol. In one embodiment, the glycerol comprises 35% glycerol. In one embodiment, the glycerol comprises 40% glycerol. In one embodiment, the glycerol comprises 45% glycerol. In one embodiment, the glycerol comprises 50% glycerol. In one embodiment, the glycerol comprises 55% glycerol. In one embodiment, the glycerol comprises 60% glycerol. In one embodiment, the glycerol comprises 65% glycerol. In one embodiment, the glycerol comprises 70% glycerol. In one embodiment, the glycerol comprises 75% glycerol. In one embodiment, the glycerol comprises 80% glycerol. In one embodiment, the glycerol comprises 85% glycerol. In one embodiment, the glycerol comprises 90% glycerol. In one embodiment, the glycerol comprises 95% glycerol. In one embodiment, the glycerol comprises 99% glycerol.

[0192] In one embodiment, the carbon source is methanol or formate. In certain embodiments, methanol is used as a carbon source in the formaldehyde assimilation pathways provided herein. In one embodiment, the carbon source is methanol or formate. In other embodiments, formate is used as a carbon source in the formaldehyde assimilation pathways provided herein. In specific embodiments, methanol is used as a carbon source in the methanol metabolic pathways provided herein, either alone or in combination with the product pathways provided herein.

[0193] In one embodiment, the carbon source comprises methanol, and sugar (e.g., glucose) or a sugar-containing biomass. In another embodiment, the carbon source comprises formate, and sugar (e.g., glucose) or a sugar-containing biomass. In one embodiment, the carbon source comprises methanol, formate, and sugar (e.g., glucose) or a sugar-containing biomass. In specific embodiments, the methanol or formate, or both, in the fermentation feed is provided as a mixture with sugar (e.g., glucose) or sugar-comprising biomass. In certain embodiments, sugar is provided for sufficient strain growth.

[0194] In certain embodiments, the carbon source comprises methanol and a sugar (e.g., glucose). In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of from 200:1 to 1:200. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of from 100:1 to 1:100. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of from 100:1 to 5:1. In some embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of from 50:1 to 5:1. In certain embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 100:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 90:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 80:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 70:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 60:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 50:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 40:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 30:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 20:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 10:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 5:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 2:1. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 1:1. In certain embodiments, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 1:100. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 1:90. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol to sugar of 1:80. In one

nol and formate to sugar of 1:80. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:70. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:60. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:50. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:40. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:30. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:20. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:10. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:5. In one embodiment, the sugar (e.g., glucose) is provided at a molar concentration ratio of methanol and formate to sugar of 1:2. In certain embodiments of the ratios provided above, the sugar is a sugar-containing biomass.

[0197] Given the teachings and guidance provided herein, those skilled in the art will understand that a non-naturally occurring microbial organism can be produced that secretes the biosynthesized compounds when grown on a carbon source such as a carbohydrate. Such compounds include, for example, BDO and any of the intermediate metabolites in the BDO pathway. All that is required is to engineer in one or more of the required enzyme or protein activities to achieve biosynthesis of the desired compound or intermediate including, for example, inclusion of some or all of the BDO biosynthetic pathways. Accordingly, provided herein is a non-naturally occurring microbial organism that produces and/or secretes BDO when grown on a carbohydrate or other carbon source and produces and/or secretes any of the intermediate metabolites shown in the BDO pathway when grown on a carbohydrate or other carbon source. The BDO producing microbial organisms provided herein can initiate synthesis from an intermediate. The same holds true for intermediates in the formaldehyde assimilation and methanol metabolic pathways.

[0198] The non-naturally occurring microbial organisms provided herein are constructed using methods well known in the art as exemplified herein to exogenously express at least one nucleic acid encoding a BDO and/or methanol metabolic pathway enzyme or protein in sufficient amounts to produce BDO. It is understood that the microbial organisms are cultured under conditions sufficient to produce BDO. Following the teachings and guidance provided herein, the non-naturally occurring microbial organisms can achieve biosynthesis of BDO, resulting in intracellular concentrations between about 0.1-500 mM or more. Generally, the intracellular concentration of BDO is between about 3-150 mM, particularly between about 5-125 mM and more particularly between about 8-100 mM, including about 10 mM, 20 mM, 50 mM, 80 mM, or more. Intracellular concentrations between and above each of these exemplary ranges also can be achieved from the non-naturally occurring microbial organisms provided herein.

[0199] In some embodiments, culture conditions include anaerobic or substantially anaerobic growth or maintenance conditions. Exemplary anaerobic conditions have been described previously and are well known in the art. Exem-

plary anaerobic conditions for fermentation processes are described herein and are described, for example, in U.S. Publ. No. 2009/0047719. Any of these conditions can be employed with the non-naturally occurring microbial organisms as well as other anaerobic conditions well known in the art. Under such anaerobic or substantially anaerobic conditions, the BDO producers can synthesize BDO at intracellular concentrations of 5-100 mM or more as well as all other concentrations exemplified herein. It is understood that, even though the above description refers to intracellular concentrations, BDO can produce BDO intracellularly and/or secrete the product into the culture medium.

[0200] Exemplary fermentation processes include, but are not limited to, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation; and continuous fermentation and continuous separation. In an exemplary batch fermentation protocol, the production organism is grown in a suitably sized bioreactor sparged with an appropriate gas. Under anaerobic conditions, the culture is sparged with an inert gas or combination of gases, for example, nitrogen, N₂/CO₂ mixture, argon, helium, and the like. As the cells grow and utilize the carbon source, additional carbon source (s) and/or other nutrients are fed into the bioreactor at a rate approximately balancing consumption of the carbon source and/or nutrients. The temperature of the bioreactor is maintained at a desired temperature, generally in the range of 22-37 degrees C., but the temperature can be maintained at a higher or lower temperature depending on the growth characteristics of the production organism and/or desired conditions for the fermentation process. Growth continues for a desired period of time to achieve desired characteristics of the culture in the fermenter, for example, cell density, product concentration, and the like. In a batch fermentation process, the time period for the fermentation is generally in the range of several hours to several days, for example, 8 to 24 hours, or 1, 2, 3, 4 or 5 days, or up to a week, depending on the desired culture conditions. The pH can be controlled or not, as desired, in which case a culture in which pH is not controlled will typically decrease to pH 3-6 by the end of the run. Upon completion of the cultivation period, the fermenter contents can be passed through a cell separation unit, for example, a centrifuge, filtration unit, and the like, to remove cells and cell debris. In the case where the desired product is expressed intracellularly, the cells can be lysed or disrupted enzymatically or chemically prior to or after separation of cells from the fermentation broth, as desired, in order to release additional product. The fermentation broth can be transferred to a product separations unit. Isolation of product occurs by standard separations procedures employed in the art to separate a desired product from dilute aqueous solutions. Such methods include, but are not limited to, liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene or other suitable solvents, including but not limited to diethyl ether, ethyl acetate, tetrahydrofuran (THF), methylene chloride, chloroform, benzene, pentane, hexane, heptane, petroleum ether, methyl tertiary butyl ether (MTBE), dioxane, dimethylformamide (DMF), dimethyl sulfoxide (DMSO), and the like) to provide an organic solution of the product, if appropriate, standard distillation methods, and the like, depending on the chemical characteristics of the product of the fermentation process.

[0201] In an exemplary fully continuous fermentation protocol, the production organism is generally first grown up in batch mode in order to achieve a desired cell density. When

the carbon source and/or other nutrients are exhausted, feed medium of the same composition is supplied continuously at a desired rate, and fermentation liquid is withdrawn at the same rate. Under such conditions, the product concentration in the bioreactor generally remains constant, as well as the cell density. The temperature of the fermenter is maintained at a desired temperature, as discussed above. During the continuous fermentation phase, it is generally desirable to maintain a suitable pH range for optimized production. The pH can be monitored and maintained using routine methods, including the addition of suitable acids or bases to maintain a desired pH range. The bioreactor is operated continuously for extended periods of time, generally at least one week to several weeks and up to one month, or longer, as appropriate and desired. The fermentation liquid and/or culture is monitored periodically, including sampling up to every day, as desired, to assure consistency of product concentration and/or cell density. In continuous mode, fermenter contents are constantly removed as new feed medium is supplied. The exit stream, containing cells, medium, and product, are generally subjected to a continuous product separations procedure, with or without removing cells and cell debris, as desired. Continuous separations methods employed in the art can be used to separate the product from dilute aqueous solutions, including but not limited to continuous liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene or other suitable solvents, including but not limited to diethyl ether, ethyl acetate, tetrahydrofuran (THF), methylene chloride, chloroform, benzene, pentane, hexane, heptane, petroleum ether, methyl tertiary butyl ether (MTBE), dioxane, dimethylformamide (DMF), dimethyl sulfoxide (DMSO), and the like), standard continuous distillation methods, and the like, or other methods well known in the art.

[0202] In addition to the culturing and fermentation conditions disclosed herein, growth condition for achieving biosynthesis of BDO can include the addition of an osmoprotectant to the culturing conditions. In certain embodiments, the non-naturally occurring microbial organisms provided herein can be sustained, cultured or fermented as described herein in the presence of an osmoprotectant. Briefly, an osmoprotectant refers to a compound that acts as an osmolyte and helps a microbial organism as described herein survive osmotic stress. Osmoprotectants include, but are not limited to, betaines, amino acids, and the sugar trehalose. Non-limiting examples of such are glycine betaine, praline betaine, dimethylthetin, dimethylsulfoniopropionate, 3-dimethylsulfonio-2-methylpropionate, pipercolic acid, dimethylsulfonioacetate, choline, L-carnitine and ectoine. In one aspect, the osmoprotectant is glycine betaine. It is understood to one of ordinary skill in the art that the amount and type of osmoprotectant suitable for protecting a microbial organism described herein from osmotic stress will depend on the microbial organism used. The amount of osmoprotectant in the culturing conditions can be, for example, no more than about 0.1 mM, no more than about 0.5 mM, no more than about 1.0 mM, no more than about 1.5 mM, no more than about 2.0 mM, no more than about 2.5 mM, no more than about 3.0 mM, no more than about 5.0 mM, no more than about 7.0 mM, no more than about 10 mM, no more than about 50 mM, no more than about 100 mM or no more than about 500 mM.

[0203] The culture conditions can include, for example, liquid culture procedures as well as fermentation and other large scale culture procedures. As described herein, particu-

larly useful yields of the biosynthetic products of the invention can be obtained under anaerobic or substantially anaerobic culture conditions.

[0204] As described herein, one exemplary growth condition for achieving biosynthesis of BDO, as well as other pathway intermediates, includes anaerobic culture or fermentation conditions. In certain embodiments, the non-naturally occurring microbial organisms provided can be sustained, cultured or fermented under anaerobic or substantially anaerobic conditions. Briefly, anaerobic conditions refer to an environment devoid of oxygen. Substantially anaerobic conditions include, for example, a culture, batch fermentation or continuous fermentation such that the dissolved oxygen concentration in the medium remains between 0 and 10% of saturation. Substantially anaerobic conditions also includes growing or resting cells in liquid medium or on solid agar inside a sealed chamber maintained with an atmosphere of less than 1% oxygen. The percent of oxygen can be maintained by, for example, sparging the culture with an N₂/CO₂ mixture or other suitable non-oxygen gas or gases.

[0205] The culture conditions described herein can be scaled up and grown continuously for manufacturing of BDO. Exemplary growth procedures include, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. All of these processes are well known in the art. Fermentation procedures are particularly useful for the biosynthetic production of commercial quantities of BDO. Generally, and as with non-continuous culture procedures, the continuous and/or near-continuous production of BDO will include culturing a non-naturally occurring BDO producing organism of the invention in sufficient nutrients and medium to sustain and/or nearly sustain growth in an exponential phase. Continuous culture under such conditions can be included, for example, growth for 1 day, 2, 3, 4, 5, 6 or 7 days or more. Additionally, continuous culture can include longer time periods of 1 week, 2, 3, 4 or 5 or more weeks and up to several months. Alternatively, organisms provided can be cultured for hours, if suitable for a particular application. It is to be understood that the continuous and/or near-continuous culture conditions also can include all time intervals in between these exemplary periods. It is further understood that the time of culturing the microbial organism of the invention is for a sufficient period of time to produce a sufficient amount of product for a desired purpose.

[0206] Fermentation procedures are well known in the art. Briefly, fermentation for the biosynthetic production of BDO can be utilized in, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. Examples of batch and continuous fermentation procedures are well known in the art.

[0207] In addition to the above fermentation procedures using the BDO producers for continuous production of substantial quantities of BDO, the BDO producers also can be, for example, simultaneously subjected to chemical synthesis procedures to convert the product to other compounds or the product can be separated from the fermentation culture and sequentially subjected to chemical conversion to convert the product to other compounds, if desired.

[0208] To generate better producers, metabolic modeling can be utilized to optimize growth conditions. Modeling can also be used to design gene knockouts that additionally optimize utilization of the pathway (see, for example, U.S. Publ.

Nos. 2002/0012939, 2003/0224363, 2004/0029149, 2004/0072723, 2003/0059792, 2002/0168654 and 2004/0009466, and U.S. Pat. No. 7,127,379). Modeling analysis allows reliable predictions of the effects on cell growth of shifting the metabolism towards more efficient production of BDO.

[0209] One computational method for identifying and designing metabolic alterations favoring biosynthesis of a desired product is the OptKnock computational framework (Burgard et al., *Biotechnol. Bioeng.* 84:647-657 (2003)). OptKnock is a metabolic modeling and simulation program that suggests gene deletion or disruption strategies that result in genetically stable microorganisms which overproduce the target product. Specifically, the framework examines the complete metabolic and/or biochemical network of a microorganism in order to suggest genetic manipulations that force the desired biochemical to become an obligatory byproduct of cell growth. By coupling biochemical production with cell growth through strategically placed gene deletions or other functional gene disruption, the growth selection pressures imposed on the engineered strains after long periods of time in a bioreactor lead to improvements in performance as a result of the compulsory growth-coupled biochemical production. Lastly, when gene deletions are constructed there is a negligible possibility of the designed strains reverting to their wild-type states because the genes selected by OptKnock are to be completely removed from the genome. Therefore, this computational methodology can be used to either identify alternative pathways that lead to biosynthesis of a desired product or used in connection with the non-naturally occurring microbial organisms for further optimization of biosynthesis of a desired product.

[0210] Briefly, OptKnock is a term used herein to refer to a computational method and system for modeling cellular metabolism. The OptKnock program relates to a framework of models and methods that incorporate particular constraints into flux balance analysis (FBA) models. These constraints include, for example, qualitative kinetic information, qualitative regulatory information, and/or DNA microarray experimental data. OptKnock also computes solutions to various metabolic problems by, for example, tightening the flux boundaries derived through flux balance models and subsequently probing the performance limits of metabolic networks in the presence of gene additions or deletions. OptKnock computational framework allows the construction of model formulations that allow an effective query of the performance limits of metabolic networks and provides methods for solving the resulting mixed-integer linear programming problems. The metabolic modeling and simulation methods referred to herein as OptKnock are described in, for example, U.S. Publ. No. 2002/0168654, International Patent Application No. PCT/US02/00660, and U.S. Publ. No. 2009/0047719.

[0211] Another computational method for identifying and designing metabolic alterations favoring biosynthetic production of a product is a metabolic modeling and simulation system termed SimPheny®. This computational method and system is described in, for example, U.S. Publ. No. 2003/0233218, and International Patent Application No. PCT/US03/18838. SimPheny® is a computational system that can be used to produce a network model in silico and to simulate the flux of mass, energy or charge through the chemical reactions of a biological system to define a solution space that contains any and all possible functionalities of the chemical reactions in the system, thereby determining a range of

allowed activities for the biological system. This approach is referred to as constraints-based modeling because the solution space is defined by constraints such as the known stoichiometry of the included reactions as well as reaction thermodynamic and capacity constraints associated with maximum fluxes through reactions. The space defined by these constraints can be interrogated to determine the phenotypic capabilities and behavior of the biological system or of its biochemical components.

[0212] These computational approaches are consistent with biological realities because biological systems are flexible and can reach the same result in many different ways. Biological systems are designed through evolutionary mechanisms that have been restricted by fundamental constraints that all living systems must face. Therefore, constraints-based modeling strategy embraces these general realities. Further, the ability to continuously impose further restrictions on a network model via the tightening of constraints results in a reduction in the size of the solution space, thereby enhancing the precision with which physiological performance or phenotype can be predicted.

[0213] Given the teachings and guidance provided herein, those skilled in the art will be able to apply various computational frameworks for metabolic modeling and simulation to design and implement biosynthesis of a desired compound in host microbial organisms. Such metabolic modeling and simulation methods include, for example, the computational systems exemplified above as SimPheny® and OptKnock. For illustration of the invention, some methods are described herein with reference to the OptKnock computation framework for modeling and simulation. Those skilled in the art will know how to apply the identification, design and implementation of the metabolic alterations using OptKnock to any of such other metabolic modeling and simulation computational frameworks and methods well known in the art.

[0214] The methods described above will provide one set of metabolic reactions to disrupt. Elimination of each reaction within the set or metabolic modification can result in a desired product as an obligatory product during the growth phase of the organism. Because the reactions are known, a solution to the bilevel OptKnock problem also will provide the associated gene or genes encoding one or more enzymes that catalyze each reaction within the set of reactions. Identification of a set of reactions and their corresponding genes encoding the enzymes participating in each reaction is generally an automated process, accomplished through correlation of the reactions with a reaction database having a relationship between enzymes and encoding genes.

[0215] Once identified, the set of reactions that are to be disrupted in order to achieve production of a desired product are implemented in the target cell or organism by functional disruption of at least one gene encoding each metabolic reaction within the set. One particularly useful means to achieve functional disruption of the reaction set is by deletion of each encoding gene. However, in some instances, it can be beneficial to disrupt the reaction by other genetic aberrations including, for example, mutation, deletion of regulatory regions such as promoters or cis binding sites for regulatory factors, or by truncation of the coding sequence at any of a number of locations. These latter aberrations, resulting in less than total deletion of the gene set can be useful, for example, when rapid assessments of the coupling of a product are desired or when genetic reversion is less likely to occur.

[0216] To identify additional productive solutions to the above described bilevel OptKnock problem which lead to further sets of reactions to disrupt or metabolic modifications that can result in the biosynthesis, including growth-coupled biosynthesis of a desired product, an optimization method, termed integer cuts, can be implemented. This method proceeds by iteratively solving the OptKnock problem exemplified above with the incorporation of an additional constraint referred to as an integer cut at each iteration. Integer cut constraints effectively prevent the solution procedure from choosing the exact same set of reactions identified in any previous iteration that obligatorily couples product biosynthesis to growth. For example, if a previously identified growth-coupled metabolic modification specifies reactions 1, 2, and 3 for disruption, then the following constraint prevents the same reactions from being simultaneously considered in subsequent solutions. The integer cut method is well known in the art and can be found described in, for example, Burgard et al., *Biotechnol. Prog.* 17:791-797 (2001). As with all methods described herein with reference to their use in combination with the OptKnock computational framework for metabolic modeling and simulation, the integer cut method of reducing redundancy in iterative computational analysis also can be applied with other computational frameworks well known in the art including, for example, SimPheny®.

[0217] The methods exemplified herein allow the construction of cells and organisms that biosynthetically produce a desired product, including the obligatory coupling of production of a target biochemical product to growth of the cell or organism engineered to harbor the identified genetic alterations. Therefore, the computational methods described herein allow the identification and implementation of metabolic modifications that are identified by an in silico method selected from OptKnock or SimPheny®. The set of metabolic modifications can include, for example, addition of one or more biosynthetic pathway enzymes and/or functional disruption of one or more metabolic reactions including, for example, disruption by gene deletion.

[0218] As discussed above, the OptKnock methodology was developed on the premise that mutant microbial networks can be evolved towards their computationally predicted maximum-growth phenotypes when subjected to long periods of growth selection. In other words, the approach leverages an organism's ability to self-optimize under selective pressures. The OptKnock framework allows for the exhaustive enumeration of gene deletion combinations that force a coupling between biochemical production and cell growth based on network stoichiometry. The identification of optimal gene/reaction knockouts requires the solution of a bilevel optimization problem that chooses the set of active reactions such that an optimal growth solution for the resulting network overproduces the biochemical of interest (Burgard et al., *Bio-technol. Bioeng.* 84:647-657 (2003)).

[0219] An in silico stoichiometric model of *E. coli* metabolism can be employed to identify essential genes for metabolic pathways as exemplified previously and described in, for example, U.S. Publ. Nos. 2002/0012939, 2003/0224363, 2004/0029149, 2004/0072723, 2003/0059792, 2002/0168654 and 2004/0009466, and in U.S. Pat. No. 7,127,379. As disclosed herein, the OptKnock mathematical framework can be applied to pinpoint gene deletions leading to the growth-coupled production of a desired product. Further, the solution of the bilevel OptKnock problem provides only one set of deletions. To enumerate all meaningful solutions, that

is, all sets of knockouts leading to growth-coupled production formation, an optimization technique, termed integer cuts, can be implemented. This entails iteratively solving the OptKnock problem with the incorporation of an additional constraint referred to as an integer cut at each iteration, as discussed above.

[0220] As disclosed herein, a nucleic acid encoding a desired activity of a BDO pathway, formaldehyde assimilation pathway, and/or methanol metabolic pathway can be introduced into a host organism. In some cases, it can be desirable to modify an activity of a BDO, formaldehyde assimilation, or methanol metabolic pathway enzyme or protein to increase production of BDO, formaldehyde and/or reducing equivalents. For example, known mutations that increase the activity of a protein or enzyme can be introduced into an encoding nucleic acid molecule. Additionally, optimization methods can be applied to increase the activity of an enzyme or protein and/or decrease an inhibitory activity, for example, decrease the activity of a negative regulator.

[0221] One such optimization method is directed evolution. Directed evolution is a powerful approach that involves the introduction of mutations targeted to a specific gene in order to improve and/or alter the properties of an enzyme. Improved and/or altered enzymes can be identified through the development and implementation of sensitive high-throughput screening assays that allow the automated screening of many enzyme variants (for example, $>10^4$). Iterative rounds of mutagenesis and screening typically are performed to afford an enzyme with optimized properties. Computational algorithms that can help to identify areas of the gene for mutagenesis also have been developed and can significantly reduce the number of enzyme variants that need to be generated and screened. Numerous directed evolution technologies have been developed (for reviews, see Hibbert et al., *Biomol. Eng.* 22:11-19 (2005); Huisman and Lalonde, In *Biocatalysis in the pharmaceutical and biotechnology industries* pgs. 717-742 (2007), Patel (ed.), CRC Press; Often and Quax, *Biomol. Eng.* 22:1-9 (2005).; and Sen et al., *Appl Biochem. Biotechnol* 143:212-223 (2007)) to be effective at creating diverse variant libraries, and these methods have been successfully applied to the improvement of a wide range of properties across many enzyme classes. Enzyme characteristics that have been improved and/or altered by directed evolution technologies include, for example: selectivity/specificity, for conversion of non-natural substrates; temperature stability, for robust high temperature processing; pH stability, for bioprocessing under lower or higher pH conditions; substrate or product tolerance, so that high product titers can be achieved; binding (K_m), including broadening substrate binding to include non-natural substrates; inhibition (K_i), to remove inhibition by products, substrates, or key intermediates; activity (kcat), to increase enzymatic reaction rates to achieve desired flux; expression levels, to increase protein yields and overall pathway flux; oxygen stability, for operation of air sensitive enzymes under aerobic conditions; and anaerobic activity, for operation of an aerobic enzyme in the absence of oxygen.

[0222] A number of exemplary methods have been developed for the mutagenesis and diversification of genes to target desired properties of specific enzymes. Such methods are well known to those skilled in the art. Any of these can be used to alter and/or optimize the activity of a BDO pathway enzyme or protein. Such methods include, but are not limited to EpPCR, which introduces random point mutations by

reducing the fidelity of DNA polymerase in PCR reactions (Pritchard et al., *J. Theor. Biol.* 234:497-509 (2005)); Error-prone Rolling Circle Amplification (epRCA), which is similar to epPCR except a whole circular plasmid is used as the template and random 6-mers with exonuclease resistant thiophosphate linkages on the last 2 nucleotides are used to amplify the plasmid followed by transformation into cells in which the plasmid is re-circularized at tandem repeats (Fujii et al., *Nucleic Acids Res.* 32:e145 (2004); and Fujii et al., *Nat. Protoc.* 1:2493-2497 (2006)); DNA or Family Shuffling, which typically involves digestion of two or more variant genes with nucleases such as Dnase I or EndoV to generate a pool of random fragments that are reassembled by cycles of annealing and extension in the presence of DNA polymerase to create a library of chimeric genes (Stemmer, *Proc. Natl. Acad. Sci. U.S.A.* 91:10747-10751 (1994); and Stemmer, *Nature* 370:389-391 (1994)); Staggered Extension (StEP), which entails template priming followed by repeated cycles of 2 step PCR with denaturation and very short duration of annealing/extension (as short as 5 sec) (Zhao et al., *Nat. Biotechnol.* 16:258-261 (1998)); Random Priming Recombination (RPR), in which random sequence primers are used to generate many short DNA fragments complementary to different segments of the template (Shao et al., *Nucleic Acids Res.* 26:681-683 (1998)).

[0223] Additional methods include Heteroduplex Recombination, in which linearized plasmid DNA is used to form heteroduplexes that are repaired by mismatch repair (Volkov et al., *Nucleic Acids Res.* 27:e18 (1999); and Volkov et al., *Methods Enzymol.* 328:456-463 (2000)); Random Chimeragenesis on Transient Templates (RACHITT), which employs Dnase I fragmentation and size fractionation of single stranded DNA (ssDNA) (Coco et al., *Nat. Biotechnol.* 19:354-359 (2001)); Recombined Extension on Truncated templates (RETT), which entails template switching of unidirectionally growing strands from primers in the presence of unidirectional ssDNA fragments used as a pool of templates (Lee et al., *J. Molec. Catalysis* 26:119-129 (2003)); Degenerate Oligonucleotide Gene Shuffling (DOGS), in which degenerate primers are used to control recombination between molecules; (Bergquist and Gibbs, *Methods Mol. Biol.* 352:191-204 (2007); Bergquist et al., *Biomol. Eng.* 22:63-72 (2005); Gibbs et al., *Gene* 271:13-20 (2001)); Incremental Truncation for the Creation of Hybrid Enzymes (ITCHY), which creates a combinatorial library with 1 base pair deletions of a gene or gene fragment of interest (Ostermeier et al., *Proc. Natl. Acad. Sci. U.S.A.* 96:3562-3567 (1999); and Ostermeier et al., *Nat. Biotechnol.* 17:1205-1209 (1999)); Thio-Incremental Truncation for the Creation of Hybrid Enzymes (THIO-ITCHY), which is similar to ITCHY except that phosphothioate dNTPs are used to generate truncations (Lutz et al., *Nucleic Acids Res.* 29:E16 (2001)); SCRATCHY, which combines two methods for recombining genes, ITCHY and DNA shuffling (Lutz et al., *Proc. Natl. Acad. Sci. U.S.A.* 98:11248-11253 (2001)); Random Drift Mutagenesis (RNDM), in which mutations made via epPCR are followed by screening/selection for those retaining usable activity (Bergquist et al., *Biomol. Eng.* 22:63-72 (2005)); Sequence Saturation Mutagenesis (SeSaM), a random mutagenesis method that generates a pool of random length fragments using random incorporation of a phosphothioate

nucleotide and cleavage, which is used as a template to extend in the presence of "universal" bases such as inosine, and replication of an inosine-containing complement gives random base incorporation and, consequently, mutagenesis (Wong et al., *Biotechnol. J.* 3:74-82 (2008); Wong et al., *Nucleic Acids Res.* 32:e26 (2004); and Wong et al., *Anal. Biochem.* 341:187-189 (2005)); Synthetic Shuffling, which uses overlapping oligonucleotides designed to encode "all genetic diversity in targets" and allows a very high diversity for the shuffled progeny (Ness et al., *Nat. Biotechnol.* 20:1251-1255 (2002)); Nucleotide Exchange and Excision Technology Next, which exploits a combination of dUTP incorporation followed by treatment with uracil DNA glycosylase and then piperidine to perform endpoint DNA fragmentation (Muller et al., *Nucleic Acids Res.* 33:e117 (2005)).

[0224] Further methods include Sequence Homology-Independent Protein Recombination (SHIPREC), in which a linker is used to facilitate fusion between two distantly related or unrelated genes, and a range of chimeras is generated between the two genes, resulting in libraries of single-cross-over hybrids (Sieber et al., *Nat. Biotechnol.* 19:456-460 (2001)); Gene Site Saturation Mutagenesis™ (GSSM™), in which the starting materials include a supercoiled double stranded DNA (dsDNA) plasmid containing an insert and two primers which are degenerate at the desired site of mutations (Kretz et al., *Methods Enzymol.* 388:3-11 (2004)); Combinatorial Cassette Mutagenesis (CCM), which involves the use of short oligonucleotide cassettes to replace limited regions with a large number of possible amino acid sequence alterations (Reidhaar-Olson et al. *Methods Enzymol.* 208:564-586 (1991); and Reidhaar-Olson et al. *Science* 241:53-57 (1988)); Combinatorial Multiple Cassette Mutagenesis (CMCM), which is essentially similar to CCM and uses epPCR at high mutation rate to identify hot spots and hot regions and then extension by CMCM to cover a defined region of protein sequence space (Reetz et al., *Angew. Chem. Int. Ed Engl.* 40:3589-3591 (2001)); the Mutator Strains technique, in which conditional is mutator plasmids, utilizing the mutD5 gene, which encodes a mutant subunit of DNA polymerase III, to allow increases of 20 to 4000-X in random and natural mutation frequency during selection and block accumulation of deleterious mutations when selection is not required (Selifonova et al., *Appl. Environ. Microbiol.* 67:3645-3649 (2001)); Low et al., *J. Mol. Biol.* 260:359-3680 (1996)).

[0225] Additional exemplary methods include Look-Through Mutagenesis (LTM), which is a multidimensional mutagenesis method that assesses and optimizes combinatorial mutations of selected amino acids (Rajpal et al., *Proc. Natl. Acad. Sci. U.S.A.* 102:8466-8471 (2005)); Gene Reassembly, which is a DNA shuffling method that can be applied to multiple genes at one time or to create a large library of chimeras (multiple mutations) of a single gene (Tunable GeneReassembly™ (TGR™) Technology supplied by Verenum Corporation), in Silico Protein Design Automation (PDA), which is an optimization algorithm that anchors the structurally defined protein backbone possessing a particular fold, and searches sequence space for amino acid substitutions that can stabilize the fold and overall protein energetics, and generally works most effectively on proteins with known three-dimensional structures (Hayes et al., *Proc. Natl. Acad.*

Sci. USA. 99:15926-15931 (2002)); and Iterative Saturation Mutagenesis (ISM), which involves using knowledge of structure/function to choose a likely site for enzyme improvement, performing saturation mutagenesis at chosen site using a mutagenesis method such as Stratagene QuikChange (Stratagene; San Diego Calif.), screening/selecting for desired properties, and, using improved clone(s), starting over at another site and continue repeating until a desired activity is achieved (Reetz et al., *Nat. Protoc.* 2:891-903 (2007); and Reetz et al., *Angew. Chem. Int. Ed Engl.* 45:7745-7751 (2006)).

[0226] Any of the aforementioned methods for mutagenesis can be used alone or in any combination. Additionally, any one or combination of the directed evolution methods can be used in conjunction with adaptive evolution techniques, as described herein.

[0227] Throughout this application various publications have been referenced. The disclosures of these publications in their entireties, including GenBank and GI number publications, are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains. Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made without departing from the spirit of the invention.

[0228] It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also included within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

4. EXAMPLES

4.1 Example 1

Production of Reducing Equivalents Via a Methanol Metabolic Pathway

[0229] Exemplary methanol metabolic pathways are provided in FIG. 1.

FIG. 1, Step A—Methanol Methyltransferase

[0230] A complex of 3-methyltransferase proteins, denoted MtaA, MtaB, and MtaC, perform the desired methanol methyltransferase activity (Sauer et al., *Eur. J. Biochem.* 243:670-677 (1997); Naidu and Ragsdale, *J. Bacteriol.* 183:3276-3281 (2001); Tallant and Krzycki, *J. Biol. Chem.* 276:4485-4493 (2001); Tallant and Krzycki, *J. Bacteriol.* 179:6902-6911 (1997); Tallant and Krzycki, *J. Bacteriol.* 178:1295-1301 (1996); Ragsdale, S. W., *Crit. Rev. Biochem. Mol. Biol.* 39:165-195 (2004)).

[0231] MtaB is a zinc protein that can catalyze the transfer of a methyl group from methanol to MtaC, a corrinoid protein. Exemplary genes encoding MtaB and MtaC can be found in methanogenic archaea such as *Methanosarcina barkeri* (Maeder et al., *J. Bacteriol.* 188:7922-7931 (2006)) and *Methanosarcina acetivorans* (Galagan et al., *Genome Res.* 12:532-542 (2002)), as well as the acetogen, *Moorella thermoacetica* (Das et al., *Proteins* 67:167-176 (2007)). In

general, the MtaB and MtaC genes are adjacent to one another on the chromosome as their activities are tightly interdependent. The protein sequences of various MtaB and MtaC encoding genes in *M. barkeri*, *M. acetivorans*, and *M. thermoacetica* can be identified by their following GenBank accession numbers.

Protein	GenBank ID	GI number	Organism
MtaB1	YP_304299	73668284	<i>Methanosarcina barkeri</i>
MtaC1	YP_304298	73668283	<i>Methanosarcina barkeri</i>
MtaB2	YP_307082	73671067	<i>Methanosarcina barkeri</i>
MtaC2	YP_307081	73671066	<i>Methanosarcina barkeri</i>
MtaB3	YP_304612	73668597	<i>Methanosarcina barkeri</i>
MtaC3	YP_304611	73668596	<i>Methanosarcina barkeri</i>
MtaB1	NP_615421	20089346	<i>Methanosarcina acetivorans</i>
MtaB1	NP_615422	20089347	<i>Methanosarcina acetivorans</i>
MtaB2	NP_619254	20093179	<i>Methanosarcina acetivorans</i>
MtaC2	NP_619253	20093178	<i>Methanosarcina acetivorans</i>
MtaB3	NP_616549	20090474	<i>Methanosarcina acetivorans</i>
MtaC3	NP_616550	20090475	<i>Methanosarcina acetivorans</i>
MtaB	YP_430066	83590057	<i>Moorella thermoacetica</i>
MtaC	YP_430065	83590056	<i>Moorella thermoacetica</i>
MtaA	YP_430064	83590056	<i>Moorella thermoacetica</i>

[0232] The MtaB1 and MtaC1 genes, YP_304299 and YP_304298, from *M. barkeri* were cloned into *E. coli* and sequenced (Sauer et al., *Eur. Biochem.* 243:670-677 (1997)). The crystal structure of this methanol-cobalamin methyltransferase complex is also available (Hagemeyer et al., *Proc. Natl. Acad. Sci. U.S.A.* 103:18917-18922 (2006)). The MtaB genes, YP_307082 and YP_304612, in *M. barkeri* were identified by sequence homology to YP_304299. In general, homology searches are an effective means of identifying methanol methyltransferases because MtaB encoding genes show little or no similarity to methyltransferases that act on alternative substrates such as trimethylamine, dimethylamine, monomethylamine, or dimethylsulfide. The MtaC genes, YP_307081 and YP_304611 were identified based on their proximity to the MtaB genes and also their homology to YP_304298. The three sets of MtaB and MtaC genes from *M. acetivorans* have been genetically, physiologically, and biochemically characterized (Pritchett and Metcalf, *Mol. Microbiol.* 56:1183-1194 (2005)). Mutant strains lacking two of the sets were able to grow on methanol, whereas a strain lacking all three sets of MtaB and MtaC genes sets could not grow on methanol. This suggests that each set of genes plays a role in methanol utilization. The *M. thermoacetica* MtaB gene was identified based on homology to the methanogenic MtaB genes and also by its adjacent chromosomal proximity to the methanol-induced corrinoid protein, MtaC, which has been crystallized (Zhou et al., *Acta Crystallogr. Sect. F. Struct. Biol. Cryst. Commun.* 61:537-540 (2005)) and further characterized by Northern hybridization and Western Blotting ((Das et al., *Proteins* 67:167-176 (2007)).

[0233] MtaA is zinc protein that catalyzes the transfer of the methyl group from MtaC to either Coenzyme M in methanogens or methyltetrahydrofolate in acetogens. MtaA can also utilize methylcobalamin as the methyl donor. Exemplary genes encoding MtaA can be found in methanogenic archaea such as *Methanosarcina barkeri* (Maeder et al., *J. Bacteriol.* 188:7922-7931 (2006) and *Methanosarcina acetivorans* (Galagan et al., *Genome Res.* 12:532-542 (2002), as well as the acetogen, *Moorella thermoacetica* ((Das et al., *Proteins* 67:167-176 (2007)). In general, MtaA proteins that catalyze the transfer of the methyl group from CH₃-MtaC are difficult to identify bioinformatically as they share similarity to other corrinoid protein methyltransferases and are not oriented adjacent to the MtaB and MtaC genes on the chromosomes. Nevertheless, a number of MtaA encoding genes have been characterized. The protein sequences of these genes in *M. barkeri* and *M. acetivorans* can be identified by the following GenBank accession numbers.

Protein	GenBank ID	GI number	Organism
MtaA	YP_304602	73668587	<i>Methanosarcina barkeri</i>
MtaA1	NP_619241	20093166	<i>Methanosarcina acetivorans</i>
MtaA2	NP_616548	20090473	<i>Methanosarcina acetivorans</i>

[0234] The MtaA gene, YP_304602, from *M. barkeri* was cloned, sequenced, and functionally overexpressed in *E. coli* (Harms and Thauer, *Eur. J. Biochem.* 235:653-659 (1996)). In *M. acetivorans*, MtaA1 is required for growth on methanol, whereas MtaA2 is dispensable even though methane production from methanol is reduced in MtaA2 mutants (Bose et al.,

methyltetrahydrofolate given the similar roles of methyltetrahydrofolate and Coenzyme M in methanogens and acetogens, respectively. The protein sequences of putative MtaA encoding genes from *M. thermoacetica* can be identified by the following GenBank accession numbers.

Protein	GenBank ID	GI number	Organism
MtaA	YP_430937	83590928	<i>Moorella thermoacetica</i>
MtaA	YP_431175	83591166	<i>Moorella thermoacetica</i>
MtaA	YP_430935	83590926	<i>Moorella thermoacetica</i>
MtaA	YP_430064	83590056	<i>Moorella thermoacetica</i>

FIG. 1, Step B—Methylenetetrahydrofolate Reductase

[0236] The conversion of methyl-THF to methylenetetrahydrofolate is catalyzed by methylenetetrahydrofolate reductase. In *M. thermoacetica*, this enzyme is oxygen-sensitive and contains an iron-sulfur cluster (Clark and Ljungdahl, *J. Biol. Chem.* 259:10845-10849 (1984). This enzyme is encoded by metF in *E. coli* (Sheppard et al., *J. Bacteriol.* 181:718-725 (1999) and CHY_1233 in *C. hydrogenoformans* (Wu et al., *PLoS Genet.* 1:e65 (2005)). The *M. thermoacetica* genes, and its *C. hydrogenoformans* counterpart, are located near the CODH/ACS gene cluster, separated by putative hydrogenase and heterodisulfide reductase genes. Some additional gene candidates found bioinformatically are listed below. In *Acetobacterium woodii* metF is coupled to the Rnf complex through RnfC2 (Poelein et al, *PLoS One.* 7:e33439). Homologs of RnfC are found in other organisms by blast search. The Rnf complex is known to be a reversible complex (Fuchs (2011) *Annu. Rev. Microbiol.* 65:631-658).

Protein	GenBank ID	GI number	Organism
Moth_1191	YP_430048.1	83590039	<i>Moorella thermoacetica</i>
Moth_1192	YP_430049.1	83590040	<i>Moorella thermoacetica</i>
metF	NP_418376.1	16131779	<i>Escherichia coli</i>
CHY_1233	YP_360071.1	78044792	<i>Carboxydotherrmus hydrogenoformans</i>
CLJU_c37610	YP_003781889.1	300856905	<i>Clostridium ljungdahlii</i> DSM 13528
DesfrDRAFT_3717	ZP_07335241.1	303248996	<i>Desulfovibrio fructosovorans</i> JJ
CcarbDRAFT_2950	ZP_05392950.1	255526026	<i>Clostridium carboxidivorans</i> P7
Ccel74_010100023124	ZP_07633513.1	307691067	<i>Clostridium cellulovorans</i> 743B
Cphy_3110	YP_001560205.1	160881237	<i>Clostridium phytofermentans</i> ISDg

J. Bacteriol. 190:4017-4026 (2008)). There are multiple additional MtaA homologs in *M. barkeri* and *M. acetivorans* that are as yet uncharacterized, but may also catalyze corrinoid protein methyltransferase activity.

[0235] Putative MtaA encoding genes in *M. thermoacetica* were identified by their sequence similarity to the characterized methanogenic MtaA genes. Specifically, three *M. thermoacetica* genes show high homology (>30% sequence identity) to YP_304602 from *M. barkeri*. Unlike methanogenic MtaA proteins that naturally catalyze the transfer of the methyl group from CH₃-MtaC to Coenzyme M, an *M. thermoacetica* MtaA is likely to transfer the methyl group to

FIG. 1, Steps C and D—Methylenetetrahydrofolate Dehydrogenase, Methenyltetrahydrofolate Cyclohydrolase

[0237] In *M. thermoacetica*, *E. coli*, and *C. hydrogenoformans*, methenyltetrahydrofolate cyclohydrolase and methyltetrahydrofolate dehydrogenase are carried out by the bifunctional gene products of Moth_1516, foLD, and CHY_1878, respectively (Pierce et al., *Environ. Microbiol.* 10:2550-2573 (2008); Wu et al., *PLoS Genet.* 1:e65 (2005); D'Ari and Rabinowitz, *J. Biol. Chem.* 266:23953-23958 (1991)). A homolog exists in *C. carboxidivorans* P7. Several other organisms also encode for this bifunctional protein as tabulated below.

Protein	GenBank ID	GI number	Organism
Moth_1516	YP_430368.1	83590359	<i>Moorella thermoacetica</i>
folD	NP_415062.1	16128513	<i>Escherichia coli</i>
CHY_1878	YP_360698.1	78044829	<i>Carboxydotherrnus hydrogenoformans</i>
CcarbDRAFT_2948	ZP_05392948.1	255526024	<i>Clostridium carboxidivorans</i> P7
folD	ADK16789.1	300437022	<i>Clostridium ljungdahlii</i> DSM 13528
folD-2	NP_951919.1	39995968	<i>Geobacter sulfurreducens</i> PCA
folD	YP_725874.1	113867385	<i>Ralstonia eutropha</i> 1116
folD	NP_348702.1	15895353	<i>Clostridium acetobutylicum</i> ATCC 824
folD	YP_696506.1	110800457	<i>Clostridium perfringens</i>
MGA3_09460	EIJ83438.1	387591119	<i>Bacillus methanolicus</i> MGA3
PB1_14689	ZP_10132349.1	387929672	<i>Bacillus methanolicus</i> PB1

FIG. 1, Step E—Formyltetrahydrofolate Deformylase

[0238] This enzyme catalyzes the hydrolysis of 10-formyltetrahydrofolate (formyl-THF) to THF and for-

teriol. 170:3255-3261 (1988), and CHY_2385 in *C. hydrogenoformans* (Wu et al., *PLoS Genet.* 1:e65 (2005)). Homologs exist in *C. carboxidivorans* P7. This enzyme is found in several other organisms as listed below.

Protein	GenBank ID	GI number	Organism
Moth_0109	YP_428991.1	83588982	<i>Moorella thermoacetica</i>
CHY_2385	YP_361182.1	78045024	<i>Carboxydotherrnus hydrogenoformans</i>
FHS	P13419.1	120562	<i>Clostridium acidurici</i>
CcarbDRAFT_1913	ZP_05391913.1	255524966	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2946	ZP_05392946.1	255526022	<i>Clostridium carboxidivorans</i> P7
Dhaf_0555	ACL18622.1	219536883	<i>Desulfotobacterium hafsiense</i>
fhs	YP_001393842.1	153953077	<i>Clostridium kluyveri</i> DSM 555
the	YP_003781893.1	300856909	<i>Clostridium ljungdahlii</i> DSM 13528
MGA3_08300	EIJ83208.1	387590889	<i>Bacillus methanolicus</i> MGA3
PB1_13509	ZP_10132113.1	387929436	<i>Bacillus methanolicus</i> PB1

mate. In *E. coli*, this enzyme is encoded by *purU* and has been overproduced, purified, and characterized (Nagy, et al., *J. Bacteriol.* 3:1292-1298 (1995)). Homologs exist in *Corynebacterium* sp. U-96 (Suzuki, et al., *Biosci. Biotechnol. Biochem.* 69(5):952-956 (2005)), *Corynebacterium glutamicum* ATCC 14067, *Salmonella enterica*, and several additional organisms.

Protein	GenBank ID	GI number	Organism
purU	AAC74314.1	1787483	<i>Escherichia coli</i> K-12 MG1655
purU	BAD97821.1	63002616	<i>Corynebacterium</i> sp. U-96
purU	EHE84645.1	354511740	<i>Corynebacterium glutamicum</i> ATCC 14067
purU	NP_460715.1	16765100	<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> str. LT2

FIG. 1, Step F—Formyltetrahydrofolate Synthetase

[0239] Formyltetrahydrofolate synthetase ligates formate to tetrahydrofolate at the expense of one ATP. This reaction is catalyzed by the gene product of *Moth_0109* in *M. thermoacetica* (O'Brien et al., *Experientia Suppl.* 26:249-262 (1976); Lovell et al., *Arch. Microbiol.* 149:280-285 (1988); Lovell et al., *Biochemistry* 29:5687-5694 (1990)), *FHS* in *Clostridium acidurici* (Whitehead and Rabinowitz, *J. Bacteriol.* 167:203-209 (1986); Whitehead and Rabinowitz, *J. Bac-*

FIG. 1, Step G—Formate Hydrogen Lyase

[0240] A formate hydrogen lyase enzyme can be employed to convert formate to carbon dioxide and hydrogen. An exemplary formate hydrogen lyase enzyme can be found in *Escherichia coli*. The *E. coli* formate hydrogen lyase consists of hydrogenase 3 and formate dehydrogenase-H (Maeda et al., *Appl Microbiol Biotechnol* 77:879-890 (2007)). It is activated by the gene product of *fhlA*. (Maeda et al., *Appl Microbiol Biotechnol* 77:879-890 (2007)). The addition of the trace elements, selenium, nickel and molybdenum, to a fermentation broth has been shown to enhance formate hydrogen lyase activity (Soini et al., *Microb. Cell Fact.* 7:26 (2008)). Various hydrogenase 3, formate dehydrogenase and transcriptional activator genes are shown below.

Protein	GenBank ID	GI number	Organism
hycA	NP_417205	16130632	<i>Escherichia coli</i> K-12 MG1655
hycB	NP_417204	16130631	<i>Escherichia coli</i> K-12 MG1655
hycC	NP_417203	16130630	<i>Escherichia coli</i> K-12 MG1655
hycD	NP_417202	16130629	<i>Escherichia coli</i> K-12 MG1655
hycE	NP_417201	16130628	<i>Escherichia coli</i> K-12 MG1655
hycF	NP_417200	16130627	<i>Escherichia coli</i> K-12 MG1655
hycG	NP_417199	16130626	<i>Escherichia coli</i> K-12 MG1655
hycH	NP_417198	16130625	<i>Escherichia coli</i> K-12 MG1655
hycI	NP_417197	16130624	<i>Escherichia coli</i> K-12 MG1655
fdhF	NP_418503	16131905	<i>Escherichia coli</i> K-12 MG1655
fhlA	NP_417211	16130638	<i>Escherichia coli</i> K-12 MG1655

[0241] A formate hydrogen lyase enzyme also exists in the hyperthermophilic archaeon, *Thermococcus litoralis* (Takacs et al., *BMC. Microbiol* 8:88 (2008)).

Protein	GenBank ID	GI number	Organism
mhyC	ABW05543	157954626	<i>Thermococcus litoralis</i>
mhyD	ABW05544	157954627	<i>Thermococcus litoralis</i>
mhyE	ABW05545	157954628	<i>Thermococcus litoralis</i>
myhF	ABW05546	157954629	<i>Thermococcus litoralis</i>
myhG	ABW05547	157954630	<i>Thermococcus litoralis</i>
myhH	ABW05548	157954631	<i>Thermococcus litoralis</i>
fdhA	AAB94932	2746736	<i>Thermococcus litoralis</i>
fdhB	AAB94931	157954625	<i>Thermococcus litoralis</i>

[0242] Additional fat late hydrogen lyase systems have been found in *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Rhodospirillum rubrum*, *Methanobacterium formicicum* (Vardar-Schara et al., *Microbial Biotechnology* 1:107-125 (2008)).

FIG. 1, Step H—Hydrogenase

[0243] Hydrogenase enzymes can convert hydrogen gas to protons and transfer electrons to acceptors such as ferredoxins, NAD⁺, or NADP⁺. *Ralstonia eutropha* H16 uses hydrogen as an energy source with oxygen as a terminal electron acceptor. Its membrane-bound uptake [NiFe]-hydrogenase is an “O₂-tolerant” hydrogenase (Cracknell, et al. *Proc Nat Acad Sci*, 106(49) 20681-20686 (2009)) that is periplasmically-oriented and connected to the respiratory chain via a b-type cytochrome (Schink and Schlegel, *Biochim. Biophys. Acta*, 567, 315-324 (1979); Bernhard et al., *Eur. J. Biochem.* 248, 179-186 (1997)). *R. eutropha* also contains an O₂-tolerant soluble hydrogenase encoded by the Hox operon which is cytoplasmic and directly reduces NAD⁺ at the expense of hydrogen (Schneider and Schlegel, *Biochim. Biophys. Acta* 452, 66-80 (1976); Burgdorf, *J. Bact.* 187(9) 3122-3132 (2005)). Soluble hydrogenase enzymes are additionally present in several other organisms including *Geobacter sulfurreducens* (Coppi, *Microbiology* 151, 1239-1254 (2005)), *Synechocystis* str. PCC 6803 (Germer, *J. Biol. Chem.*, 284 (52), 36462-36472 (2009)), and *Thiocapsa roseopersicina* (Rakhely, *Appl. Environ. Microbiol.* 70(2) 722-728 (2004)). The *Synechocystis* enzyme is capable of generating NADPH from hydrogen. Overexpression of both the Hox operon from *Synechocystis* str. PCC 6803 and the accessory genes encoded by the Hyp operon from *Nostoc* sp. PCC 7120 led to increased hydrogenase activity compared to expression of the Hox genes alone (Germer, *J. Biol. Chem.* 284(52), 36462-36472 (2009)).

Protein	GenBank ID	GI Number	Organism
HoxF	NP_942727.1	38637753	<i>Ralstonia eutropha</i> H16
HoxU	NP_942728.1	38637754	<i>Ralstonia eutropha</i> H16
HoxY	NP_942729.1	38637755	<i>Ralstonia eutropha</i> H16
HoxH	NP_942730.1	38637756	<i>Ralstonia eutropha</i> H16
HoxW	NP_942731.1	38637757	<i>Ralstonia eutropha</i> H16
HoxI	NP_942732.1	38637758	<i>Ralstonia eutropha</i> H16
HoxE	NP_953767.1	39997816	<i>Geobacter sulfurreducens</i>
HoxF	NP_953766.1	39997815	<i>Geobacter sulfurreducens</i>
HoxU	NP_953765.1	39997814	<i>Geobacter sulfurreducens</i>
HoxY	NP_953764.1	39997813	<i>Geobacter sulfurreducens</i>

-continued

Protein	GenBank ID	GI Number	Organism
HoxH	NP_953763.1	39997812	<i>Geobacter sulfurreducens</i>
GSU2717	NP_953762.1	39997811	<i>Geobacter sulfurreducens</i>
HoxE	NP_441418.1	16330690	<i>Synechocystis</i> str. PCC 6803
HoxF	NP_441417.1	16330689	<i>Synechocystis</i> str. PCC 6803
Unknown function	NP_441416.1	16330688	<i>Synechocystis</i> str. PCC 6803
HoxU	NP_441415.1	16330687	<i>Synechocystis</i> str. PCC 6803
HoxY	NP_441414.1	16330686	<i>Synechocystis</i> str. PCC 6803
Unknown function	NP_441413.1	16330685	<i>Synechocystis</i> str. PCC 6803
Unknown function	NP_441412.1	16330684	<i>Synechocystis</i> str. PCC 6803
HoxH	NP_441411.1	16330683	<i>Synechocystis</i> str. PCC 6803
HypF	NP_484737.1	17228189	<i>Nostoc</i> sp. PCC 7120
HypC	NP_484738.1	17228190	<i>Nostoc</i> sp. PCC 7120
HypD	NP_484739.1	17228191	<i>Nostoc</i> sp. PCC 7120
Unknown function	NP_484740.1	17228192	<i>Nostoc</i> sp. PCC 7120
HypE	NP_484741.1	17228193	<i>Nostoc</i> sp. PCC 7120
HypA	NP_484742.1	17228194	<i>Nostoc</i> sp. PCC 7120
HypB	NP_484743.1	17228195	<i>Nostoc</i> sp. PCC 7120
Hox1E	AAP50519.1	37787351	<i>Thiocapsa roseopersicina</i>
Hox1F	AAP50520.1	37787352	<i>Thiocapsa roseopersicina</i>
Hox1U	AAP50521.1	37787353	<i>Thiocapsa roseopersicina</i>
Hox1Y	AAP50522.1	37787354	<i>Thiocapsa roseopersicina</i>
Hox1H	AAP50523.1	37787355	<i>Thiocapsa roseopersicina</i>

[0244] The genomes of *E. coli* and other enteric bacteria encode up to four hydrogenase enzymes (Sawers, G., *Antonie Van Leeuwenhoek* 66:57-88 (1994); Sawers et al., *J Bacteriol.* 164:1324-1331 (1985); Sawers and Boxer, *Eur. J Biochem.* 156:265-275 (1986); Sawers et al., *J Bacteriol.* 168:398-404 (1986)). Given the multiplicity of enzyme activities *E. coli* or another host organism can provide sufficient hydrogenase activity to split incoming molecular hydrogen and reduce the corresponding acceptor. Endogenous hydrogen-lyase enzymes of *E. coli* include hydrogenase 3, a membrane-bound enzyme complex using ferredoxin as an acceptor, and hydrogenase 4 that also uses a ferredoxin acceptor. Hydrogenase 3 and 4 are encoded by the hyc and hyf gene clusters, respectively. Hydrogenase activity in *E. coli* is also dependent upon the expression of the hyp genes whose corresponding proteins are involved in the assembly of the hydrogenase complexes (Jacobi et al., *Arch. Microbiol* 158:444-451 (1992); Rangarajan et al., *J Bacteria* 190:1447-1458 (2008)). The *M. thermoacetica* and *Clostridium ljungdahli* hydrogenases are suitable for a host that lacks sufficient endogenous hydrogenase activity. *M. thermoacetica* and *C. ljungdahli* can grow with CO₂ as the exclusive carbon source indicating that reducing equivalents are extracted from H₂ to enable acetyl-CoA synthesis via the Wood-Ljungdahl pathway (Drake, H. L., *J Bacteriol.* 150:702-709 (1982); Drake and Daniel, *Res Microbiol* 155:869-883 (2004); Kellum and Drake, *J Bacteriol.* 160:466-469 (1984)). *M. thermoacetica* has homologs to several hyp, hyc, and hyf genes from *E. coli*. These protein sequences encoded for by these genes are identified by the following GenBank accession numbers. In addition, several gene clusters encoding hydrogenase functionality are present in *M. thermoacetica* and *C. ljungdahli* (see for example US 2012/0003652).

-continued

Protein	GenBank ID	GI Number	Organism
HypA	NP_417206	16130633	<i>Escherichia coli</i>
HypB	NP_417207	16130634	<i>Escherichia coli</i>
HypC	NP_417208	16130635	<i>Escherichia coli</i>
HypD	NP_417209	16130636	<i>Escherichia coli</i>
HypE	NP_417210	226524740	<i>Escherichia coli</i>
HypF	NP_417192	16130619	<i>Escherichia coli</i>
HycA	NP_417205	16130632	<i>Escherichia coli</i>
HycB	NP_417204	16130631	<i>Escherichia coli</i>
HycC	NP_417203	16130630	<i>Escherichia coli</i>
HycD	NP_417202	16130629	<i>Escherichia coli</i>
HycE	NP_417201	16130628	<i>Escherichia coli</i>
HycF	NP_417200	16130627	<i>Escherichia coli</i>
HycG	NP_417199	16130626	<i>Escherichia coli</i>
HycH	NP_417198	16130625	<i>Escherichia coli</i>
HycI	NP_417197	16130624	<i>Escherichia coli</i>
HyfA	NP_416976	90111444	<i>Escherichia coli</i>
HyfB	NP_416977	16130407	<i>Escherichia coli</i>
HyfC	NP_416978	90111445	<i>Escherichia coli</i>
HyfD	NP_416979	16130409	<i>Escherichia coli</i>
HyfE	NP_416980	16130410	<i>Escherichia coli</i>
HyfF	NP_416981	16130411	<i>Escherichia coli</i>
HyfG	NP_416982	16130412	<i>Escherichia coli</i>
HyfH	NP_416983	16130413	<i>Escherichia coli</i>
HyfI	NP_416984	16130414	<i>Escherichia coli</i>
HyfJ	NP_416985	90111446	<i>Escherichia coli</i>
HyfR	NP_416986	90111447	<i>Escherichia coli</i>

[0245] Proteins in *M. thermoacetica* whose genes are homologous to the *E. coli* hydrogenase genes are shown below.

Protein	GenBank ID	GI Number	Organism
Moth_2175	YP_431007	83590998	<i>Moorella thermoacetica</i>
Moth_2176	YP_431008	83590999	<i>Moorella thermoacetica</i>
Moth_2177	YP_431009	83591000	<i>Moorella thermoacetica</i>
Moth_2178	YP_431010	83591001	<i>Moorella thermoacetica</i>
Moth_2179	YP_431011	83591002	<i>Moorella thermoacetica</i>
Moth_2180	YP_431012	83591003	<i>Moorella thermoacetica</i>
Moth_2181	YP_431013	83591004	<i>Moorella thermoacetica</i>
Moth_2182	YP_431014	83591005	<i>Moorella thermoacetica</i>
Moth_2183	YP_431015	83591006	<i>Moorella thermoacetica</i>
Moth_2184	YP_431016	83591007	<i>Moorella thermoacetica</i>
Moth_2185	YP_431017	83591008	<i>Moorella thermoacetica</i>
Moth_2186	YP_431018	83591009	<i>Moorella thermoacetica</i>
Moth_2187	YP_431019	83591010	<i>Moorella thermoacetica</i>
Moth_2188	YP_431020	83591011	<i>Moorella thermoacetica</i>
Moth_2189	YP_431021	83591012	<i>Moorella thermoacetica</i>
Moth_2190	YP_431022	83591013	<i>Moorella thermoacetica</i>
Moth_2191	YP_431023	83591014	<i>Moorella thermoacetica</i>
Moth_2192	YP_431024	83591015	<i>Moorella thermoacetica</i>
Moth_0439	YP_429313	83589304	<i>Moorella thermoacetica</i>
Moth_0440	YP_429314	83589305	<i>Moorella thermoacetica</i>
Moth_0441	YP_429315	83589306	<i>Moorella thermoacetica</i>
Moth_0442	YP_429316	83589307	<i>Moorella thermoacetica</i>
Moth_0809	YP_429670	83589661	<i>Moorella thermoacetica</i>
Moth_0810	YP_429671	83589662	<i>Moorella thermoacetica</i>
Moth_0811	YP_429672	83589663	<i>Moorella thermoacetica</i>
Moth_0812	YP_429673	83589664	<i>Moorella thermoacetica</i>
Moth_0814	YP_429674	83589665	<i>Moorella thermoacetica</i>
Moth_0815	YP_429675	83589666	<i>Moorella thermoacetica</i>
Moth_0816	YP_429676	83589667	<i>Moorella thermoacetica</i>
Moth_1193	YP_430050	83590041	<i>Moorella thermoacetica</i>
Moth_1194	YP_430051	83590042	<i>Moorella thermoacetica</i>
Moth_1195	YP_430052	83590043	<i>Moorella thermoacetica</i>

Protein	GenBank ID	GI Number	Organism
Moth_1196	YP_430053	83590044	<i>Moorella thermoacetica</i>
Moth_1717	YP_430562	83590553	<i>Moorella thermoacetica</i>
Moth_1718	YP_430563	83590554	<i>Moorella thermoacetica</i>
Moth_1719	YP_430564	83590555	<i>Moorella thermoacetica</i>
Moth_1883	YP_430726	83590717	<i>Moorella thermoacetica</i>
Moth_1884	YP_430727	83590718	<i>Moorella thermoacetica</i>
Moth_1885	YP_430728	83590719	<i>Moorella thermoacetica</i>
Moth_1886	YP_430729	83590720	<i>Moorella thermoacetica</i>
Moth_1887	YP_430730	83590721	<i>Moorella thermoacetica</i>
Moth_1888	YP_430731	83590722	<i>Moorella thermoacetica</i>
Moth_1452	YP_430305	83590296	<i>Moorella thermoacetica</i>
Moth_1453	YP_430306	83590297	<i>Moorella thermoacetica</i>
Moth_1454	YP_430307	83590298	<i>Moorella thermoacetica</i>

[0246] Genes encoding hydrogenase enzymes from *C. ljungdahli* are shown below.

Protein	GenBank ID	GI Number	Organism
CLJU_c20290	ADK15091.1	300435324	<i>Clostridium ljungdahli</i>
CLJU_c07030	ADK13773.1	300434006	<i>Clostridium ljungdahli</i>
CLJU_c07040	ADK13774.1	300434007	<i>Clostridium ljungdahli</i>
CLJU_c07050	ADK13775.1	300434008	<i>Clostridium ljungdahli</i>
CLJU_c07060	ADK13776.1	300434009	<i>Clostridium ljungdahli</i>
CLJU_c07070	ADK13777.1	300434010	<i>Clostridium ljungdahli</i>
CLJU_c07080	ADK13778.1	300434011	<i>Clostridium ljungdahli</i>
CLJU_c14730	ADK14541.1	300434774	<i>Clostridium ljungdahli</i>
CLJU_c14720	ADK14540.1	300434773	<i>Clostridium ljungdahli</i>
CLJU_c14710	ADK14539.1	300434772	<i>Clostridium ljungdahli</i>
CLJU_c14700	ADK14538.1	300434771	<i>Clostridium ljungdahli</i>
CLJU_c28670	ADK15915.1	300436148	<i>Clostridium ljungdahli</i>
CLJU_c28660	ADK15914.1	300436147	<i>Clostridium ljungdahli</i>
CLJU_c28650	ADK15913.1	300436146	<i>Clostridium ljungdahli</i>
CLJU_c28640	ADK15912.1	300436145	<i>Clostridium ljungdahli</i>

[0247] In some cases, hydrogenase encoding genes are located adjacent to a CODH. In *Rhodospirillum rubrum*, the encoded CODH/hydrogenase proteins form a membrane-bound enzyme complex that has been indicated to be a site where energy, in the form of a proton gradient, is generated from the conversion of CO and H₂O to CO₂ and H₂ (Fox et al., *J Bacteriol.* 178:6200-6208 (1996)). The CODH-I of *C. hydrogenoformans* and its adjacent genes have been proposed to catalyze a similar functional role based on their similarity to the *R. rubrum* CODH/hydrogenase gene cluster (Wu et al., *PLoS Genet.* 1:e65 (2005)). The *C. hydrogenoformans* CODH-I was also shown to exhibit intense CO oxidation and CO₂ reduction activities when linked to an electrode (Parkin et al., *J Am. Chem. Soc.* 129:10328-10329 (2007)).

Protein	GenBank ID	GI Number	Organism
CooL	AAC45118	1515468	<i>Rhodospirillum rubrum</i>
CooX	AAC45119	1515469	<i>Rhodospirillum rubrum</i>
CooU	AAC45120	1515470	<i>Rhodospirillum rubrum</i>
CooH	AAC45121	1498746	<i>Rhodospirillum rubrum</i>
CooF	AAC45122	1498747	<i>Rhodospirillum rubrum</i>
CODH (CooS)	AAC45123	1498748	<i>Rhodospirillum rubrum</i>
CooC	AAC45124	1498749	<i>Rhodospirillum rubrum</i>
CooT	AAC45125	1498750	<i>Rhodospirillum rubrum</i>
CooJ	AAC45126	1498751	<i>Rhodospirillum rubrum</i>
CODH-I (CooS-I)	YP_360644	78043418	<i>Carboxydotherrmus hydrogenoformans</i>
CooF	YP_360645	78044791	<i>Carboxydotherrmus hydrogenoformans</i>
HypA	YP_360646	78044340	<i>Carboxydotherrmus hydrogenoformans</i>
CooH	YP_360647	78043871	<i>Carboxydotherrmus hydrogenoformans</i>
CooU	YP_360648	78044023	<i>Carboxydotherrmus hydrogenoformans</i>
CooX	YP_360649	78043124	<i>Carboxydotherrmus hydrogenoformans</i>
CooL	YP_360650	78043938	<i>Carboxydotherrmus hydrogenoformans</i>
CooK	YP_360651	78044700	<i>Carboxydotherrmus hydrogenoformans</i>
CooM	YP_360652	78043942	<i>Carboxydotherrmus hydrogenoformans</i>
CooC	YP_360654.1	78043296	<i>Carboxydotherrmus hydrogenoformans</i>
CooA-1	YP_360655.1	78044021	<i>Carboxydotherrmus hydrogenoformans</i>

[0248] Some hydrogenase and CODH enzymes transfer electrons to ferredoxins. Ferredoxins are small acidic proteins containing one or more iron-sulfur clusters that function as intracellular electron carriers with a low reduction potential. Reduced ferredoxins donate electrons to Fe-dependent enzymes such as ferredoxin-NADP⁺ oxidoreductase, pyruvate:ferredoxin oxidoreductase (PFOR) and 2-oxoglutarate:ferredoxin oxidoreductase (OFOR). The *H. thermophilus* gene *fdx1* encodes a [4Fe-4S]-type ferredoxin that is required for the reversible carboxylation of 2-oxoglutarate and pyruvate by OFOR and PFOR, respectively (Yamamoto et al., *Extremophiles* 14:79-85 (2010)). The ferredoxin associated with the *Sulfolobus solfataricus* 2-oxoacid:ferredoxin reductase is a monomeric dicluster [3Fe-4S][4Fe-4S] type ferredoxin (Park et al. 2006). While the gene associated with this

protein has not been fully sequenced, the N-terminal domain shares 93% homology with the *zfx* ferredoxin from *S. acidocaldarius*. The *E. coli* genome encodes a soluble ferredoxin of unknown physiological function, *fdx*. Some evidence indicates that this protein can function in iron-sulfur cluster assembly (Takahashi and Nakamura, 1999). Additional ferredoxin proteins have been characterized in *Helicobacter pylori* (Mukhopadhyay et al. 2003) and *Campylobacter jejuni* (van Vliet et al. 2001). A 2Fe-2S ferredoxin from *Clostridium pasteurianum* has been cloned and expressed in *E. coli* (Fujinaga and Meyer, Biochemical and Biophysical Research Communications, 192(3): (1993)). Acetogenic bacteria such as *Moorella thermoacetica*, *Clostridium carboxidivorans* P7, *Clostridium ljungdahli* and *Rhodospirillum rubrum* are predicted to encode several ferredoxins, listed below.

Protein	GenBank ID	GI Number	Organism
<i>fdx1</i>	BAE02673.1	68163284	<i>Hydrogenobacter thermophilus</i>
M11214.1	AAA83524.1	144806	<i>Clostridium pasteurianum</i>
<i>Zfx</i>	AAV79867.1	68566938	<i>Sulfolobus acidocaldarius</i>
<i>Fdx</i>	AAC75578.1	1788874	<i>Escherichia coli</i>
<i>hp_0277</i>	AAD07340.1	2313367	<i>Helicobacter pylori</i>
<i>fdxA</i>	CAL34484.1	112359698	<i>Campylobacter jejuni</i>
Moth_0061	ABC18400.1	83571848	<i>Moorella thermoacetica</i>
Moth_1200	ABC19514.1	83572962	<i>Moorella thermoacetica</i>
Moth_1888	ABC20188.1	83573636	<i>Moorella thermoacetica</i>
Moth_2112	ABC20404.1	83573852	<i>Moorella thermoacetica</i>
Moth_1037	ABC19351.1	83572799	<i>Moorella thermoacetica</i>
CcarbDRAFT_4383	ZP_05394383.1	255527515	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2958	ZP_05392958.1	255526034	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2281	ZP_05392281.1	255525342	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_5296	ZP_05395295.1	255528511	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_1615	ZP_05391615.1	255524662	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_1304	ZP_05391304.1	255524347	<i>Clostridium carboxidivorans</i> P7
<i>cooF</i>	AAG29808.1	11095245	<i>Carboxydotherrmus hydrogenoformans</i>
<i>fdxN</i>	CAA35699.1	46143	<i>Rhodobacter capsulatus</i>
Rru_A2264	ABC23064.1	83576513	<i>Rhodospirillum rubrum</i>
Rru_A1916	ABC22716.1	83576165	<i>Rhodospirillum rubrum</i>
Rru_A2026	ABC22826.1	83576275	<i>Rhodospirillum rubrum</i>
<i>cooF</i>	AAC45122.1	1498747	<i>Rhodospirillum rubrum</i>
<i>fdxN</i>	AAA26460.1	152605	<i>Rhodospirillum rubrum</i>
Alvin_2884	ADC63789.1	288897953	<i>Allochroamatium vinosum</i> DSM 180
<i>Fdx</i>	YP_002801146.1	226946073	<i>Azotobacter vinelandii</i> DJ
CKL_3790	YP_001397146.1	153956381	<i>Clostridium kluyveri</i> DSM 555

-continued

Protein	GenBank ID	GI Number	Organism
ferI	NP_949965.1	39937689	<i>Rhodopseudomonas palustris</i> CGA009
Fdx	CAA12251.1	3724172	<i>Thauera aromatica</i>
CHY_2405	YP_361202.1	78044690	<i>Carboxydotherrnus hydrogenoformans</i>
Fer	YP_359966.1	78045103	<i>Carboxydotherrnus hydrogenoformans</i>
Fer	AAC83945.1	1146198	<i>Bacillus subtilis</i>
fdx1	NP_249053.1	15595559	<i>Pseudomonas aeruginosa</i> PA01
yfhL	AP_003148.1	89109368	<i>Escherichia coli</i> K-12
CLJU_c00930	ADK13195.1	300433428	<i>Clostridium ljungdahlii</i>
CLJU_c00010	ADK13115.1	300433348	<i>Clostridium ljungdahlii</i>
CLJU_c01820	ADK13272.1	300433505	<i>Clostridium ljungdahlii</i>
CLJU_c17980	ADK14861.1	300435094	<i>Clostridium ljungdahlii</i>
CLJU_c17970	ADK14860.1	300435093	<i>Clostridium ljungdahlii</i>
CLJU_c22510	ADK15311.1	300435544	<i>Clostridium ljungdahlii</i>
CLJU_c26680	ADK15726.1	300435959	<i>Clostridium ljungdahlii</i>
CLJU_c29400	ADK15988.1	300436221	<i>Clostridium ljungdahlii</i>

[0249] Ferredoxin oxidoreductase enzymes transfer electrons from ferredoxins or flavodoxins to NAD(P)H. Two enzymes catalyzing the reversible transfer of electrons from reduced ferredoxins to NAD(P)⁺ are ferredoxin:NAD⁺ oxidoreductase (EC 1.18.1.3) and ferredoxin:NADP⁺ oxidoreductase (FNR, EC 1.18.1.2). Ferredoxin:NADP⁺ oxidoreductase (FNR, EC 1.18.1.2) has a noncovalently bound FAD cofactor that facilitates the reversible transfer of electrons from NADPH to low-potential acceptors such as ferredoxins or flavodoxins (Blaschkowski et al., *Eur. J. Biochem.* 123:563-569 (1982); Fujii et al., 1977). The *Helicobacter pylori* FNR, encoded by HP1164 (fqrB), is coupled to the activity of pyruvate:ferredoxin oxidoreductase (PFOR) resulting in the pyruvate-dependent production of NADPH (St et al. 2007). An analogous enzyme is found in *Campylobacter jejuni* (St Maurice et al., *J. Bacteriol.* 189:4764-4773 (2007)). A ferredoxin:NADP⁺ oxidoreductase enzyme is encoded in the *E. coli* genome by fpr (Bianchi et al. 1993). Ferredoxin:NAD⁺ oxidoreductase utilizes reduced ferre-

doxin to generate NADH from NAD⁺. In several organisms, including *E. coli*, this enzyme is a component of multifunctional dioxygenase enzyme complexes. The ferredoxin:NAD⁺ oxidoreductase of *E. coli*, encoded by hcaD, is a component of the 3-phenylproppionate dioxygenase system involved in aromatic acid utilization (Diaz et al. 1998). NADH:ferredoxin reductase activity was detected in cell extracts of *Hydrogenobacter thermophilus*, although a gene with this activity has not yet been indicated (Yoon et al. 2006). Additional ferredoxin:NAD(P)⁺ oxidoreductases have been annotated in *Clostridium carboxidivorans* P7. The NADH-dependent reduced ferredoxin: NADP oxidoreductase of *C. kluyveri*, encoded by nfnAB, catalyzes the concomitant reduction of ferredoxin and NAD⁺ with two equivalents of NADPH (Wang et al, *J Bacteriol* 192: 5115-5123 (2010)). Finally, the energy-conserving membrane-associated Rnf-type proteins (Seedorf et al, *PNAS* 105:2128-2133 (2008); and Herrmann, *J. Bacteriol* 190:784-791 (2008)) provide a means to generate NADH or NADPH from reduced ferredoxin.

Protein	GenBank ID	GI Number	Organism
fqrB	NP_207955.1	15645778	<i>Helicobacter pylori</i>
fqrB	YP_001482096.1	157414840	<i>Campylobacter jejuni</i>
RPA3954	CAE29395.1	39650872	<i>Rhodopseudomonas palustris</i>
Fpr	BAH29712.1	225320633	<i>Hydrogenobacter thermophilus</i>
yumC	NB_391091.2	255767736	<i>Bacillus subtilis</i>
Fpr	P28861.4	399486	<i>Escherichia coli</i>
hcaD	AAC75595.1	1788892	<i>Escherichia coli</i>
LOC100282643	NP_001149023.1	226497434	<i>Zea mays</i>
NfnA	YP_001393861.1	153953096	<i>Clostridium kluyveri</i>
NfnB	YP_001393862.1	153953097	<i>Clostridium kluyveri</i>
CcarbDRAFT_2639	ZP_05392639.1	255525707	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2638	ZP_05392638.1	255525706	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2636	ZP_05392636.1	255525704	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_5060	ZP_05395060.1	255528241	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_2450	ZP_05392450.1	255525514	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_1084	ZP_05391084.1	255524124	<i>Clostridium carboxidivorans</i> P7
RnfC	EDK33306.1	146346770	<i>Clostridium kluyveri</i>
RnfD	EDK33307.1	146346771	<i>Clostridium kluyveri</i>
RnfG	EDK33308.1	146346772	<i>Clostridium kluyveri</i>
RnfE	EDK33309.1	146346773	<i>Clostridium kluyveri</i>
RnfA	EDK33310.1	146346774	<i>Clostridium kluyveri</i>
RnfB	EDK33311.1	146346775	<i>Clostridium kluyveri</i>
CLJU_c11410 (RnfB)	ADK14209.1	300434442	<i>Clostridium ljungdahlii</i>
CLJU_c11400 (RnfA)	ADK14208.1	300434441	<i>Clostridium ljungdahlii</i>
CLJU_c11390 (RnfE)	ADK14207.1	300434440	<i>Clostridium ljungdahlii</i>
CLJU_c11380 (RnfG)	ADK14206.1	300434439	<i>Clostridium ljungdahlii</i>
CLJU_c11370 (RnfD)	ADK14205.1	300434438	<i>Clostridium ljungdahlii</i>

-continued

Protein	GenBank ID	GI Number	Organism
CLJU_c11360 (RnfC)	ADK14204.1	300434437	<i>Clostridium ljungdahlii</i>
MOTH_1518 (NfnA)	YP_430370.1	83590361	<i>Moorella thermoacetica</i>
MOTH_1517(NfnB)	YP_430369.1	83590360	<i>Moorella thermoacetica</i>
CHY_1992 (NfnA)	YP_360811.1	78045020	<i>Carboxydotherrmus hydrogenoformans</i>
CHY_1993 (NfnB)	YP_360812.1	78044266	<i>Carboxydotherrmus hydrogenoformans</i>
CLJU_c37220 (NfnAB)	YP_003781850.1	300856866	<i>Clostridium ljungdahlii</i>

FIG. 1, Step I—Formate Dehydrogenase

[0250] Formate dehydrogenase (FDH) catalyzes the reversible transfer of electrons from formate to an acceptor.

moacetica ATCC 39073, *Candida boidinii*, *Candida methylica*, and *Saccharomyces cerevisiae* S288c. The soluble formate dehydrogenase from *Ralstonia eutropha* reduces NAD⁺ (fdsG, -B, -A, -C, -D) (Oh and Bowien, 1998)

Protein	GenBank ID	GI Number	Organism
Moth_2312	YP_431142	148283121	<i>Moorella thermoacetica</i>
Moth_2314	YP_431144	83591135	<i>Moorella thermoacetica</i>
Sfum_2703	YP_846816.1	116750129	<i>Syntrophobacter fumaroxidans</i>
Sfum_2704	YP_846817.1	116750130	<i>Syntrophobacter fumaroxidans</i>
Sfum_2705	YP_846818.1	116750131	<i>Syntrophobacter fumaroxidans</i>
Sfum_2706	YP_846819.1	116750132	<i>Syntrophobacter fumaroxidans</i>
CHY_0731	YP_359585.1	78044572	<i>Carboxydotherrmus hydrogenoformans</i>
CHY_0732	YP_359586.1	78044500	<i>Carboxydotherrmus hydrogenoformans</i>
CHY_0733	YP_359587.1	78044647	<i>Carboxydotherrmus hydrogenoformans</i>
CcarbDRAFT_0901	ZP_05390901.1	255523938	<i>Clostridium carboxidivorans</i> P7
CcarbDRAFT_4380	ZP_05394380.1	255527512	<i>Clostridium carboxidivorans</i> P7
fdhA, MGA3_06625	EIJ82879.1	387590560	<i>Bacillus methanolicus</i> MGA3
fdhA, PB1_11719	ZP_10131761.1	387929084	<i>Bacillus methanolicus</i> PB1
fdhD, MGA3_06630	EIJ82880.1	387590561	<i>Bacillus methanolicus</i> MGA3
fdhD, PB1_11724	ZP_10131762.1	387929085	<i>Bacillus methanolicus</i> PB1
fdh	ACF35003.	194220249	<i>Burkholderia stabilis</i>
FDH1	AAC49766.1	2276465	<i>Candida boidinii</i>
fdh	CAA57036.1	1181204	<i>Candida methylica</i>
FDH2	P0CF35.1	294956522	<i>Saccharomyces cerevisiae</i> S288c
FDH1	NP_015033.1	6324964	<i>Saccharomyces cerevisiae</i> S288c
fdsG	YP_725156.1	113866667	<i>Ralstonia eutropha</i>
fdsB	YP_725157.1	113866668	<i>Ralstonia eutropha</i>
fdsA	YP_725158.1	113866669	<i>Ralstonia eutropha</i>
fdsC	YP_725159.1	113866670	<i>Ralstonia eutropha</i>
fdsD	YP_725160.1	113866671	<i>Ralstonia eutropha</i>

Enzymes with FDH activity utilize various electron carriers such as, for example, NADH (EC 1.2.1.2), NADPH (EC 1.2.1.43), quinols (EC 1.1.5.6), cytochromes (EC 1.2.2.3) and hydrogenases (EC 1.1.99.33). FDH enzymes have been characterized from *Moorella thermoacetica* (Andreesen and Ljungdahl, *J Bacteriol* 116:867-873 (1973); Li et al., *J Bacteriol* 92:405-412 (1966); Yamamoto et al., *J Biol Chem* 258:1826-1832 (1983). The loci, Moth 2312 is responsible for encoding the alpha subunit of formate dehydrogenase while the beta subunit is encoded by Moth_2314 (Pierce et al., *Environ Microbiol* (2008)). Another set of genes encoding formate dehydrogenase activity with a propensity for CO₂ reduction is encoded by Sfum_2703 through Sfum_2706 in *Syntrophobacter fumaroxidans* (de Bok et al., *Eur J Biochem* 270:2476-2485 (2003)); Reda et al., *PNAS* 105:10654-10658 (2008)). A similar set of genes presumed to carry out the same function are encoded by CHY_0731, CHY_0732, and CHY_0733 in *C. hydrogenoformans* (Wu et al., *PLoS Genet* 1:e65 (2005)). Formate dehydrogenases are also found many additional organisms including *C. carboxidivorans* P7, *Bacillus methanolicus*, *Burkholderia stabilis*, *Moorella ther-*

FIG. 1, Step J—Methanol Dehydrogenase

[0251] NAD⁺ dependent methanol dehydrogenase enzymes (EC 1.1.1.244) catalyze the conversion of methanol and NAD⁺ to formaldehyde and NADH. An enzyme with this activity was first characterized in *Bacillus methanolicus* (Heggeset, et al., *Applied and Environmental Microbiology*, 78(15):5170-5181 (2012)). This enzyme is zinc and magnesium dependent, and activity of the enzyme is enhanced by the activating enzyme encoded by act (Kloosterman et al, *J Biol Chem* 277:34785-92 (2002)). Additional NAD(P)⁺ dependent enzymes can be identified by sequence homology. Methanol dehydrogenase enzymes utilizing different electron acceptors are also known in the art. Examples include cytochrome dependent enzymes such as mxalF of the methylotroph *Methylobacterium extorquens* (Nunn et al, *Nucl Acid Res* 16:7722 (1988)). Methanol dehydrogenase enzymes of methanotrophs such as *Methylococcus capsulatus* function in a complex with methane monooxygenase (MMO) (Myronova et al. *Biochem* 45:11905-14 (2006)). Methanol can also be oxidized to formaldehyde by alcohol oxidase enzymes such as methanol oxidase (EC 1.1.3.13) of *Candida boidinii* (Sakai et al, *Gene* 114: 67-73 (1992)).

Protein	GenBank ID	GI Number	Organism
mdh, MGA3_17392	EIJ77596.1	387585261	<i>Bacillus methanolicus</i> MGA3
mdh2, MGA3_07340	EIJ83020.1	387590701	<i>Bacillus methanolicus</i> MGA3
mdh3, MGA3_10725	EIJ80770.1	387588449	<i>Bacillus methanolicus</i> MGA3
act, MGA3_09170	EIJ83380.1	387591061	<i>Bacillus methanolicus</i> MGA3
mdh, PB1_17533	ZP_10132907.1	387930234	<i>Bacillus methanolicus</i> PB1
mdh1, PB1_14569	ZP_10132325.1	387929648	<i>Bacillus methanolicus</i> PB1
mdh2, PB1_12584	ZP_10131932.1	387929255	<i>Bacillus methanolicus</i> PB1
act, PB1_14394	ZP_10132290.1	387929613	<i>Bacillus methanolicus</i> PB1
BFZC1_05383	ZP_07048751.1	299535429	<i>Lysinibacillus fusiformis</i>
BFZC1_20163	ZP_07051637.1	299538354	<i>Lysinibacillus fusiformis</i>
Bsph_4187	YP_001699778.1	169829620	<i>Lysinibacillus sphaericus</i>
Bsph_1706	YP_001697432.1	169827274	<i>Lysinibacillus sphaericus</i>
MCA0299	YP_112833.1	53802410	<i>Methylococcus capsulatus</i>
MCA0782	YP_113284.1	53804880	<i>Methylococcus capsulatus</i>
mxaI	YP_002965443.1	240140963	<i>Methylobacterium extorquens</i>
mxaF	YP_002965446.1	240140966	<i>Methylobacterium extorquens</i>
AOD1	AAA34321.1	170820	<i>Candida boidinii</i>

FIG. 1, Step K—Spontaneous or Formaldehyde Activating Enzyme

[0252] The conversion of formaldehyde and THF to methylenetetrahydrofolate can occur spontaneously. It is also possible that the rate of this reaction can be enhanced by a formaldehyde activating enzyme. A formaldehyde activating enzyme (Fae) has been identified in *Methylobacterium extorquens* AM1 which catalyzes the condensation of formaldehyde and tetrahydropterin to methylene tetrahydropterin (Vorholt, et al., *J. Bacteriol.*, 182(23), 6645-6650 (2000)). It is possible that a similar enzyme exists or can be engineered to catalyze the condensation of formaldehyde and tetrahydrofolate to methylenetetrahydrofolate. Homologs exist in several organisms including *Xanthobacter autotrophicus* Py2 and *Hyphomicrobium denitrificans* ATCC 51888.

Protein	GenBank ID	GI Number	Organism
MexAM1_META1p1766	Q9FA38.3	17366061	<i>Methylobacterium extorquens</i> AM1
Xaut_0032	YP_001414948.1	154243990	<i>Xanthobacter autotrophicus</i> Py2
Hden_1474	YP_003755607.1	300022996	<i>Hyphomicrobium denitrificans</i> ATCC 51888

FIG. 1, Step L—Formaldehyde Dehydrogenase

[0253] Oxidation of formaldehyde to formate is catalyzed by formaldehyde dehydrogenase. An NAD⁺ dependent formaldehyde dehydrogenase enzyme is encoded by fdhA of *Pseudomonas putida* (Ito et al, *J Bacteriol* 176: 2483-2491 (1994)). Additional formaldehyde dehydrogenase enzymes include the NAD⁺ and glutathione independent formaldehyde dehydrogenase from *Hyphomicrobium zavarzinii* (Jerome et al, *Appl Microbiol Biotechnol* 77:779-88 (2007)), the glutathione dependent formaldehyde dehydrogenase of *Pichia pastoris* (Sunga et al, *Gene* 330:39-47 (2004)) and the NAD(P)⁺ dependent formaldehyde dehydrogenase of *Methylobacter marinus* (Speer et al, *FEMS Microbiol Lett*, 121(3): 349-55 (1994)).

Protein	GenBank ID	GI Number	Organism
fdhA	P46154.3	1169603	<i>Pseudomonas putida</i>
faoA	CAC85637.1	19912992	<i>Hyphomicrobium zavarzinii</i>
Fld1	CCA39112.1	328352714	<i>Pichia pastoris</i>
fdh	P47734.2	221222447	<i>Methylobacter marinus</i>

[0254] In addition to the formaldehyde dehydrogenase enzymes listed above, alternate enzymes and pathways for converting formaldehyde to formate are known in the art. For example, many organisms employ glutathione-dependent formaldehyde oxidation pathways, in which formaldehyde is converted to formate in three steps via the intermediates S-hydroxymethylglutathione and S-formylglutathione (Vorholt et al, *J Bacteriol* 182:6645-50 (2000)). The enzymes of this

pathway are S-(hydroxymethyl)glutathione synthase (EC 4.4.1.22), glutathione-dependent formaldehyde dehydrogenase (EC 1.1.1.284) and S-formylglutathione hydrolase (EC 3.1.2.12).

FIG. 1, Step M—Spontaneous or S-(hydroxymethyl)glutathione Synthase

[0255] While conversion of formaldehyde to S-hydroxymethylglutathione can occur spontaneously in the presence of glutathione, it has been shown by Goenrich et al (Goenrich, et al., *J Biol Chem* 277(5); 3069-72 (2002)) that an enzyme from *Paracoccus denitrificans* can accelerate this spontaneous condensation reaction. The enzyme catalyzing the conversion of formaldehyde and glutathione was purified and named

glutathione-dependent formaldehyde-activating enzyme (Gfa). The gene encoding it, which was named *gfa*, is located directly upstream of the gene for glutathione-dependent formaldehyde dehydrogenase, which catalyzes the subsequent oxidation of S-hydroxymethylglutathione. Putative proteins with sequence identity to Gfa from *P. denitrificans* are present also in *Rhodobacter sphaeroides*, *Sinorhizobium meliloti*, and *Mesorhizobium loti*.

Protein	GenBank ID	GI Number	Organism
Gfa	Q51669.3	38257308	<i>Paracoccus denitrificans</i>
Gfa	ABP71667.1	145557054	<i>Rhodobacter sphaeroides</i> ATCC 17025
Gfa	Q92WX6.1	38257348	<i>Sinorhizobium meliloti</i> 1021
Gfa	Q98LU4.2	38257349	<i>Mesorhizobium loti</i> MAFF303099

FIG. 1, Step N—Glutathione-Dependent Formaldehyde Dehydrogenase

[0256] Glutathione-dependent formaldehyde dehydrogenase (GS-FDH) belongs to the family of class III alcohol dehydrogenases. Glutathione and formaldehyde combine non-enzymatically to form hydroxymethylglutathione, the true substrate of the GS-FDH catalyzed reaction. The product, S-formylglutathione, is further metabolized to formic acid.

Protein	GenBank ID	GI Number	Organism
<i>frmA</i>	YP_488650.1	388476464	<i>Escherichia coli</i> K-12 MG1655
<i>SFA1</i>	NP_010113.1	6320033	<i>Saccharomyces cerevisiae</i> S288c
<i>flhA</i>	AAC44551.1	1002865	<i>Paracoccus denitrificans</i>
<i>adhI</i>	AAB09774.1	986949	<i>Rhodobacter sphaeroides</i>

FIG. 1, Step O—S-formylglutathione Hydrolase

[0257] S-formylglutathione hydrolase is a glutathione thiol esterase found in bacteria, plants and animals. It catalyzes conversion of S-formylglutathione to formate and glutathione. The *fghA* gene of *P. denitrificans* is located in the same operon with *gfa* and *flhA*, two genes involved in the oxidation of formaldehyde to formate in this organism. In *E. coli*, *FrmB* is encoded in an operon with *FrmR* and *FrmA*, which are proteins involved in the oxidation of formaldehyde. *YeiG* of *E. coli* is a promiscuous serine hydrolase; its highest specific activity is with the substrate S-formylglutathione.

Protein	GenBank ID	GI Number	Organism
<i>frmB</i>	NP_414889.1	16128340	<i>Escherichia coli</i> K-12 MG1655
<i>yeiG</i>	AAC75215.1	1788477	<i>Escherichia coli</i> K-12 MG1655
<i>fghA</i>	AAC44554.1	1002868	<i>Paracoccus denitrificans</i>

4.2 Example II

Enhanced Yield of 1,4 Butanediol from Carbohydrates Using Methanol

[0258] Exemplary methanol metabolic pathways for enhancing the availability of reducing equivalents are provided in FIG. 2.

FIG. 2, Step A—Succinyl-CoA Transferase or Succinyl-CoA Synthetase (or Succinyl-CoA Ligase)

[0259] The conversion of succinate to succinyl-CoA is catalyzed by succinyl-CoA transferase or synthetase (ligase). Succinyl-CoA transferase enzymes include *catI* of *Clostridium kluyveri* and *ygfH* of *E. coli* (Seedorf et al., *Proc. Natl. Acad. Sci USA* 105:2128-2133 (2008); Sohling et al., *J. Bacteriol.* 178:871-880 (1996); Haller et al., *Biochemistry*, 39(16) 4622-4629). Homologs can be found in, for example, *Citrobacter youngae* ATCC 29220, *Salmonella enterica* subsp. *arizonae* serovar, and *Yersinia intermedia* ATCC 29909. Succinyl-CoA:3-oxoacid-CoA transferase employs succinate as the CoA acceptor. This enzyme is encoded by *pcaI* and *pcaJ* in *Pseudomonas putida* (Kaschabek et al., *J. Bacteriol.* 184:207-215 (2002)). Similar enzymes are found in *Acinetobacter* sp. ADP1 (Kowalchuk et al., *Gene* 146:23-30 (1994)), *Streptomyces coelicolor* and *Pseudomonas knackmussii* (formerly sp. B13) (Gobel et al., *J. Bacteriol.* 184:216-223 (2002); Kaschabek et al., *J. Bacteriol.* 184:207-215 (2002)). Other succinyl-CoA:3-oxoacid-CoA transferases have been characterized in *Helicobacter pylori* (Corthesy-Theulaz et al., *J. Biol. Chem.* 272:25659-25667 (1997)), *Bacillus subtilis* (Stols et al., *Protein Expr. Purif.* 53:396-403 (2007)) and *Homo sapiens* (Fukao, T., et al., *Genomics* 68:144-151 (2000); Tanaka, H., et al., *Mol Hum Reprod* 8:16-23 (2002)). Genbank information related to these genes is summarized below.

Protein	GenBank ID	GI Number	Organism
<i>catI</i>	P38946.1	729048	<i>Clostridium kluyveri</i>
<i>ygfH</i>	NP_417395.1	16130821	<i>Escherichia coli</i>
CIT292_04485	ZP_03838384.1	227334728	<i>Citrobacter youngae</i>
SARI_04582	YP_001573497.1	161506385	<i>Salmonella enterica</i>
yinte0001_14430	ZP_04635364.1	238791727	<i>Yersinia intermedia</i>
<i>pcaI</i>	24985644	AAN69545.1	<i>Pseudomonas putida</i>
<i>pcaJ</i>	26990657	NP_746082.1	<i>Pseudomonas putida</i>
<i>pcaI</i>	50084858	YP_046368.1	<i>Acinetobacter</i> sp. ADP1
<i>pcaJ</i>	141776	AAC37147.1	<i>Acinetobacter</i> sp. ADP1
<i>pcaI</i>	21224997	NP_630776.1	<i>Streptomyces coelicolor</i>
<i>pcaJ</i>	21224996	NP_630775.1	<i>Streptomyces coelicolor</i>
<i>catI</i>	75404583	Q8VVF3	<i>Pseudomonas knackmussii</i>
<i>catJ</i>	75404582	Q8VVF2	<i>Pseudomonas knackmussii</i>
HPAG1_0676	108563101	YP_627417	<i>Helicobacter pylori</i>
HPAG1_0677	108563102	YP_627418	<i>Helicobacter pylori</i>
<i>ScoA</i>	16080950	NP_391778	<i>Bacillus subtilis</i>
<i>ScoB</i>	16080949	NP_391777	<i>Bacillus subtilis</i>
OXCT1	NP_000427	4557817	<i>Homo sapiens</i>
OXCT2	NP_071403	11545841	<i>Homo sapiens</i>

[0260] Succinyl-CoA synthetase, also called succinyl-CoA ligase, is encoded by *sucCD* of *E. coli* and *LSC1* and *LSC2* genes of *Saccharomyces cerevisiae*. These enzymes catalyze the formation of succinyl-CoA from succinate with the con-

comitant consumption of one ATP in a reaction which is reversible in vivo (Buck et al., *Biochemistry* 24:6245-6252 (1985)).

Protein	GenBank ID	GI Number	Organism
sucC	NP_415256.1	16128703	<i>Escherichia coli</i>
sucD	AAC73823.1	1786949	<i>Escherichia coli</i>
LSC1	NP_014785	6324716	<i>Saccharomyces cerevisiae</i>
LSC2	NP_011760	6321683	<i>Saccharomyces cerevisiae</i>

FIG. 2, Step B—Succinyl-CoA Reductase (Aldehyde Forming)

[0261] Enzymes with succinyl-CoA reductase activity are encoded by sucD of *Clostridium kluyveri* (Sohling, *J. Bacteriol.* 178:871-880 (1996)) and sucD of *Porphyromonas gingivalis* (Takahashi, *J. Bacteriol.* 182:4704-4710 (2000)). Additional succinyl-CoA reductase enzymes participate in the 3-hydroxypropionate/4-hydroxybutyrate cycle of thermophilic archaea such as *Metallosphaera sedula* (Berg et al., *Science* 318:1782-1786 (2007)) and *Thermoproteus neutrophilus* (Ramos-Vera et al., *J. Bacteriol.* 191:4286-4297 (2009)). The *M. sedula* enzyme, encoded by Msed_0709, is strictly NADPH-dependent and also has malonyl-CoA reductase activity. The *T. neutrophilus* enzyme is active with both NADPH and NADH.

Protein	GenBank ID	GI Number	Organism
MSED_0709	YP_001190808.1	146303492	<i>Metallosphaera sedula</i>
Tneu_0421	ACB39369.1	170934108	<i>Thermoproteus neutrophilus</i>
sucD	P38947.1	172046062	<i>Clostridium kluyveri</i>
sucD	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>

FIG. 2, Step C—4-Hydroxybutyrate Dehydrogenase

[0262] Enzymes exhibiting 4-hydroxybutyrate dehydrogenase activity (EC 1.1.1.61) have been characterized in *Ralstonia eutropha* (Bravo et al., *J. Forensic Sci.* 49:379-387 (2004)), *Clostridium kluyveri* (Wolff and Kenealy, *Protein Expr. Purif.* 6:206-212 (1995)) and *Arabidopsis thaliana* (Breitkreuz et al., *J. Biol. Chem.* 278:41552-41556 (2003)). Other 4-hydroxybutyrate dehydrogenase enzymes are found in *Porphyromonas gingivalis* and gbd of an uncultured bacterium. Accession numbers of these genes are listed in the table below.

Protein	GenBank ID	GI Number	Organism
4hbd	YP_726053.1	113867564	<i>Ralstonia eutropha</i> H16
4hbd	L21902.1	146348486	<i>Clostridium kluyveri</i> DSM 555
4hbd	Q94B07	75249805	<i>Arabidopsis thaliana</i>
4-hBd	NP_904964.1	34540485	<i>Porphyromonas gingivalis</i> W83
gbd	AF148264.1	5916168	Uncultured bacterium

FIG. 2, Step D—Hydroxybutyrate Kinase

[0263] Activation of 4-hydroxybutyrate to 4-hydroxybutyryl-phosphate is catalyzed by 4-hydroxybutyrate kinase.

Phosphotransferase enzymes in the EC class 2.7.2 transform carboxylic acids to phosphonic acids with concurrent hydrolysis of one ATP. Enzymes suitable for catalyzing this reaction include butyrate kinase, acetate kinase, aspartokinase and gamma-glutamyl kinase. Butyrate kinase carries out the reversible conversion of butyryl-phosphate to butyrate during acidogenesis in *C. acetobutylicum* (Cary et al., *Appl. Environ. Microbiol.* 56:1576-1583 (1990)). This enzyme is encoded by either of the two buk gene products (Huang et al., *J. Mol. Microbiol. Biotechnol.* 2:33-38 (2000)). Other butyrate kinase enzymes are found in *C. butyricum*, *C. beijerinckii* and *C. tetanomorphum* (Twarog and Wolfe, *J. Bacteriol.* 86:112-117 (1963)). A related enzyme, isobutyrate kinase from *Thermotoga maritime*, has also been expressed in *E. coli* and crystallized (Diao et al., *Acta Crystallogr. D. Biol. Crystallogr.* 59:1100-1102 (2003); Diao and Hasson, *J. Bacteriol.* 191:2521-2529 (2009)). Aspartokinase catalyzes the ATP-dependent phosphorylation of aspartate and participates in the synthesis of several amino acids. The aspartokinase III enzyme in *E. coli*, encoded by lysC, has a broad substrate range, and the catalytic residues involved in substrate specificity have been elucidated (Deng and Viola, *Arch. Biochem. Biophys.* 335:73-81 (1996)). Two additional kinases in *E. coli* are also good candidates: acetate kinase and gamma-glutamyl kinase. The *E. coli* acetate kinase, encoded by ackA (Skarstedt and Silverstein, *J. Biol. Chem.* 251:6775-6783 (1976)), phosphorylates propionate in addition to acetate (Hesslinger et al., *Mol. Microbiol.* 27:477-492 (1998)). The *E. coli* gamma-glutamyl kinase, encoded by proB (Smith et al., *J. Bacteriol.* 157:545-551 (1984)), phosphorylates the gamma carbonic acid group of glutamate.

Gene	Accession No.	GI No.	Organism
buk1	NP_349675	15896326	<i>Clostridium acetobutylicum</i>
buk2	Q97H1	20137415	<i>Clostridium acetobutylicum</i>
buk2	Q9X278.1	6685256	<i>Thermotoga maritime</i>
lysC	NP_418448.1	16131850	<i>Escherichia coli</i>
ackA	NP_416799.1	16130231	<i>Escherichia coli</i>
proB	NP_414777.1	16128228	<i>Escherichia coli</i>
buk	YP_001307350.1	150015096	<i>Clostridium beijerinckii</i>
buk2	YP_001311072.1	150018818	<i>Clostridium beijerinckii</i>

FIG. 2, Step E—Phosphotrans-4-hydroxybutyrylase

[0264] Phosphotrans-4-hydroxybutyrylase catalyzes the transfer of the 4-hydroxybutyryl group from phosphate to CoA. Acyltransferases suitable for catalyzing this reaction include phosphotransacetylase and phosphotransbutyrylase. The pta gene from *E. coli* encodes an enzyme that can convert acetyl-phosphate into acetyl-CoA (Suzuki, *Biochim. Biophys. Acta* 191:559-569 (1969)). This enzyme can also utilize propionyl-CoA instead of acetyl-CoA (Hesslinger et al., *Mol. Microbiol.* 27:477-492 (1998)). Similarly, the ptb gene from *C. acetobutylicum* encodes an enzyme that can convert butyryl-CoA into butyryl-phosphate (Walter et al., *Gene* 134:107-111 (1993)); Huang et al., *J. Mol. Microbiol. Biotechnol.* 2:33-38 (2000). Additional ptb genes can be found in Clostridial organisms, butyrate-producing bacterium L2-50 (Louis et al., *J. Bacteriol.* 186:2099-2106 (2004)) and *Bacillus megaterium* (Vazquez et al., *Curr. Microbiol.* 42:345-349 (2001)).

Gene	Accession No.	GI No.	Organism
pta	NP_416800.1	16130232	<i>Escherichia coli</i>
ptb	NP_349676	15896327	<i>Clostridium acetobutylicum</i>
ptb	YP_001307349.1	150015095	<i>Clostridium beijerinckii</i>
ptb	AAR19757.1	38425288	butyrate-producing bacterium L2-50
ptb	CAC07932.1	10046659	<i>Bacillus megaterium</i>

FIG. 2, Step F—4-Hydroxybutyryl-CoA Reductase (Aldehyde Forming)

[0265] 4-hydroxybutyryl-CoA reductase catalyzes the reduction of 4-hydroxybutyryl-CoA to its corresponding aldehyde. Several acyl-CoA dehydrogenases are capable of catalyzing this activity. The succinate semialdehyde dehydrogenases (SucD) of *Clostridium kluyveri* and *P. gingivalis* were shown in ref. (WO/2008/115840) to convert 4-hydroxybutyryl-CoA to 4-hydroxybutanal as part of a pathway to produce 1,4-butanediol. Many butyraldehyde dehydrogenases are also active on 4-hydroxybutyraldehyde, including bld of *Clostridium saccharoperbutylacetonicum* and bphG of *Pseudomonas* sp (Powlowski et al., *J. Bacteriol.* 175:377-385 (1993)). Yet another candidate is the ald gene from *Clostridium beijerinckii* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). This gene is very similar to eutE that encodes acetaldehyde dehydrogenase of *Salmonella typhimurium* and *E. coli* (Toth, *Appl. Environ. Microbiol.* 65:4973-4980 (1999)). These and additional proteins with 4-hydroxybutyryl-CoA reductase activity are identified below.

Protein	GenBank ID	GI Number	Organism
sucD	P38947.1	172046062	<i>Clostridium kluyveri</i>
sucD	NP_904963.1	34540484	<i>Porphyromonas gingivalis</i>
bld	AAP42563.1	31075383	<i>Clostridium saccharoperbutylacetonicum</i>
bphG	BAA03892.1	425213	<i>Pseudomonas</i> sp
Ald	AAT66436	49473535	<i>Clostridium beijerinckii</i>
eutE	AAA80209	687645	<i>Salmonella typhimurium</i>
eutE	P77445	2498347	<i>Escherichia coli</i>
ald	YP_001310903.1	150018649	<i>Clostridium beijerinckii</i> NCIMB 8052
Ald	ZP_03778292.1	225569267	<i>Clostridium hylemonae</i> DSM 15053
Ald	ZP_03705305.1	225016072	<i>Clostridium methylpentosum</i> DSM 5476
Ald	ZP_03715465.1	225026273	<i>Eubacterium hallii</i> DSM 3353
Ald	ZP_01962381.1	153809713	<i>Ruminococcus obeum</i> ATCC 29174
Ald	YP_003701164.1	297585384	<i>Bacillus selenitireducens</i> MLS10
Ald	AAP42563.1	31075383	<i>Clostridium saccharoperbutylacetonicum</i> N1-4
Ald	YP_795711.1	116334184	<i>Lactobacillus brevis</i> ATCC 367
Ald	YP_002434126.1	218782808	<i>Desulfatibacillum alkenivorans</i> AK-01
Ald	YP_001558295.1	160879327	<i>Clostridium phytofermentans</i> ISDg
Ald	ZP_02089671.1	160942363	<i>Clostridium bolteae</i> ATCC BAA-613
Ald	ZP_01222600.1	90414628	<i>Photobacterium profundum</i> 3TCK

-continued

Protein	GenBank ID	GI Number	Organism
Ald	YP_001452373.1	157145054	<i>Citrobacter koseri</i> ATCC BAA-895
Ald	NP_460996.1	16765381	<i>Salmonella enterica typhimurium</i>
Ald	YP_003307836.1	269119659	<i>Sebadella termitidis</i> ATCC 33386
Ald	ZP_04969437.1	254302079	<i>Fusobacterium nucleatum</i> subsp. <i>polymorphum</i> ATCC 10953
Ald	YP_002892893.1	237808453	<i>Tolomonas auensis</i> DSM 9187
Ald	YP_426002.1	83592250	<i>Rhodospirillum rubrum</i> ATCC 11170

FIG. 2, Step G—1,4-butanediol dehydrogenase

[0266] 1,4-butanediol dehydrogenase catalyzes the reduction of 4-hydroxybutyraldehyde to 1,4-butanediol. Exemplary genes encoding this activity include alrA of *Acinetobacter* sp. strain M-1 (Tani et al., *Appl. Environ. Microbiol.* 66:5231-5235 (2000)), yqhD and fucO from *E. coli* (Sulzenbacher et al., *J Mol Biol* 342:489-502 (2004)) and bdh I and bdh II from *C. acetobutylicum* (Walter et al, *J. Bacteriol* 174:7149-7158 (1992)). Additional 1,4-butanediol dehydrogenase enzymes are encoded by bdh in *C. saccharoperbutylacetonicum* and Cbei_1722, Cbei_2181 and Cbei_2421 in *C. beijerinckii*. These and other enzymes with 1,4-butanediol activity are listed in the table below.

Protein	GenBank ID	GI Number	Organism
alrA	BAB12273.1	9967138	<i>Acinetobacter</i> sp. strain M-1
ADH2	NP_014032.1	6323961	<i>Saccharomyces cerevisiae</i>
fucO	NP_417279.1	16130706	<i>Escherichia coli</i>
yqhD	NP_417484.1	16130909	<i>Escherichia coli</i>
bdh I	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
bdh II	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
bdh	BAF45463.1	124221917	<i>Clostridium saccharoperbutylacetonicum</i>
Cbei_1722	YP_001308850	150016596	<i>Clostridium beijerinckii</i>
Cbei_2181	YP_001309304	150017050	<i>Clostridium beijerinckii</i>
Cbei_2421	YP_001309535	150017281	<i>Clostridium beijerinckii</i>
14bdh	AAC76047.1	1789386	<i>Escherichia coli</i> K-12 MG1655
14bdh	YP_001309304.1	150017050	<i>Clostridium beijerinckii</i> NCIMB 8052
14bdh	P13604.1	113352	<i>Clostridium saccharobutylicum</i>
14bdh	ZP_03760651.1	225405462	<i>Clostridium asparagiforme</i> DSM 15981
14bdh	ZP_02083621.1	160936248	<i>Clostridium bolteae</i> ATCC BAA-613
14bdh	YP_003845251.1	302876618	<i>Clostridium cellulovorans</i> 743B
14bdh	ZP_03294286.1	210624270	<i>Clostridium hiranonis</i> DSM 13275
14bdh	ZP_03705769.1	225016577	<i>Clostridium methylpentosum</i> DSM 5476
14bdh	YP_003179160.1	257783943	<i>Atopobium parvulum</i> DSM 20469
14bdh	YP_002893476.1	237809036	<i>Tolomonas auensis</i> DSM 9187

-continued

Protein	GenBank ID	GI Number	Organism
14bdh	ZP_05394983.1	255528157	<i>Clostridium carboxidivorans</i> P7

FIG. 2, Step H—Succinate Reductase

[0267] Direct reduction of succinate to succinate semialdehyde is catalyzed by a carboxylic acid reductase. Exemplary enzymes for catalyzing this transformation are described below (see FIG. 2, Step K).

FIG. 2, Step I—Succinyl-CoA Reductase (Alcohol Forming)

[0268] Succinyl-CoA reductase (alcohol forming) enzymes are bifunctional oxidoreductases that convert succinyl-CoA to 4-hydroxybutyrate. 4-Hydroxybutyryl-CoA reductase (alcohol forming) enzymes candidates, described below (FIG. 2, Step M), are also suitable for catalyzing the reduction of succinyl-CoA.

FIG. 2, Step J—4-Hydroxybutyryl-CoA Transferase or 4-Hydroxybutyryl-CoA Synthetase

[0269] Conversion of 4-hydroxybutyrate to 4-hydroxybutyryl-CoA is catalyzed by a CoA transferase or synthetase. 4-Hydroxybutyryl-CoA transferase enzymes include the gene products of cat1, cat2, and cat3 of *Clostridium kluyveri* (Seedorf et al., *Proc. Natl. Acad. Sci. U.S.A.* 105:2128-2133 (2008); Sohling et al., *J. Bacteriol.* 178:871-880 (1996)). Similar CoA transferase activities are also present in *Trichomonas vaginalis*, *Trypanosoma brucei*, *Clostridium aminobutyricum* and *Porphyromonas gingivalis* (Riviere et al., *J. Biol. Chem.* 279:45337-45346 (2004); van Grinsven et al., *J. Biol. Chem.* 283:1411-1418 (2008)).

Protein	GenBank ID	GI Number	Organism
cat1	P38946.1	729048	<i>Clostridium kluyveri</i>
cat2	P38942.2	172046066	<i>Clostridium kluyveri</i>
cat3	EDK35586.1	146349050	<i>Clostridium kluyveri</i>
TVAG_395550	XP_001330176	123975034	<i>Trichomonas vaginalis</i> G3
Tb11.02.0290	XP_828352	71754875	<i>Trypanosoma brucei</i>
cat2	CAB60036.1	6249316	<i>Clostridium aminobutyricum</i>
cat2	NP_906037.1	34541558	<i>Porphyromonas gingivalis</i> W83

[0270] 4HB-CoA synthetase catalyzes the ATP-dependent conversion of 4-hydroxybutyrate to 4-hydroxybutyryl-CoA. AMP-forming 4-HB-CoA synthetase enzymes are found in organisms that assimilate carbon via the dicarboxylate/hydroxybutyrate cycle or the 3-hydroxypropionate/4-hydroxybutyrate cycle. Enzymes with this activity have been charac-

terized in *Thermoproteus neutrophilus* and *Metallosphaera sedula* (Ramos-Vera et al, *J. Bacteriol* 192:5329-40 (2010); Berg et al, *Science* 318:1782-6 (2007)). Others can be inferred by sequence homology. ADP forming CoA synthetases, such as succinyl-CoA synthetase, are also suitable candidates.

Protein	GenBank ID	GI Number	Organism
Tneu_0420	ACB39368.1	170934107	<i>Thermoproteus neutrophilus</i>
Caur_0002	YP_001633649.1	163845605	<i>Chloroflexus aurantiacus</i> J-10-fl
Cagg_3790	YP_002465062	219850629	<i>Chloroflexus aggregans</i> DSM 9485
acs	YP_003431745	288817398	<i>Hydrogenobacter thermophilus</i> TK-6
Pisl_0250	YP_929773.1	119871766	<i>Pyrobaculum islandicum</i> DSM 4184
Msed_1422	ABP95580.1	145702438	<i>Metallosphaera sedula</i>

FIG. 2, Step K—4-Hydroxybutyrate Reductase

[0271] Reduction of 4-hydroxybutyrate to 4-hydroxybutanal is catalyzed by a carboxylic acid reductase (CAR). Such an enzyme is found in *Nocardia iowensis*. Carboxylic acid reductase enzymes catalyze the ATP and NADPH-dependent reduction of carboxylic acids to their corresponding aldehydes (Venkatasubramanian et al., *J. Biol. Chem.* 282:478-485 (2007)). The *Nocardia iowensis* enzyme, encoded by car, was cloned and functionally expressed in *E. coli* (Venkatasubramanian et al., *J. Biol. Chem.* 282:478-485 (2007)). Expression of the npt gene product improved activity of the enzyme via post-transcriptional modification. The npt gene encodes a specific phosphopantetheine transferase (PPTase) that converts the inactive apo-enzyme to the active holo-enzyme. The natural substrate of this enzyme is vanillic acid, and the enzyme exhibits broad acceptance of aromatic and aliphatic substrates including 4-hydroxybutyrate (Venkatasubramanian et al., in *Biocatalysis in the Pharmaceutical and Biotechnology Industries*, ed. R. N. Patel, Chapter 15, pp. 425-440, CRC Press LLC, Boca Raton, Fla. (2006)).

Gene name	GI Number	Gen Bank ID	Organism
Car	40796035	AAR91681.1	<i>Nocardia iowensis</i> (sp. NRRL 5646)
Npt	114848891	ABI83656.1	<i>Nocardia iowensis</i> (sp. NRRL 5646)

[0272] Additional car and npt genes can be identified based on sequence homology.

Gene name	GI Number	GenBank ID	Organism
fadD9	121638475	YP_978699.1	<i>Mycobacterium bovis</i> BCG
BCG_2812c	121638674	YP_978898.1	<i>Mycobacterium bovis</i> BCG
nfa20150	54023983	YP_118225.1	<i>Nocardia farcinica</i> IFM 10152
nfa40540	54026024	YP_120266.1	<i>Nocardia farcinica</i> IFM 10152

-continued

Gene name	GI Number	GenBank ID	Organism
SGR_6790	182440583	YP_001828302.1	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
SGR_665	182434458	YP_001822177.1	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
MSMEG_2956	YP_887275.1	YP_887275.1	<i>Mycobacterium smegmatis</i> MC2 155
MSMEG_5739	YP_889972.1	118469671	<i>Mycobacterium smegmatis</i> MC2 155
MSMEG_2648	YP_886985.1	118471293	<i>Mycobacterium smegmatis</i> MC2 155
MAP1040c	NP_959974.1	41407138	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
MAP2899c	NP_961833.1	41408997	<i>Mycobacterium avium</i> subsp. <i>paratuberculosis</i> K-10
MMAR_2117	YP_001850422.1	183982131	<i>Mycobacterium marinum</i> M
MMAR_2936	YP_001851230.1	183982939	<i>Mycobacterium marinum</i> M
MMAR_1916	YP_001850220.1	183981929	<i>Mycobacterium marinum</i> M
TpauDRAFT_33060	ZP_04027864.1	227980601	<i>Tsukamurella paurometabola</i> DSM 20162
TpauDRAFT_20920	ZP_04026660.1	ZP_04026660.1	<i>Tsukamurella paurometabola</i> DSM 20162
CPCC7001_1320	ZP_05045132.1	254431429	<i>Cyanobium</i> PCC7001
DDBDRAFT_0187729	XP_636931.1	66806417	<i>Dictyostelium discoideum</i> AX4

[0273] An additional CAR enzyme found in *Streptomyces griseus* is encoded by the *grIC* and *grID* genes. This enzyme is believed to convert 3-amino-4-hydroxybenzoic acid to 3-amino-4-hydroxybenzaldehyde as deletion of either *grIC* or *grID* led to accumulation of extracellular 3-acetylamino-4-hydroxybenzoic acid, a shunt product of 3-amino-4-hydroxybenzoic acid metabolism (Suzuki, et al., *J. Antibiot.* 60(6): 380-387 (2007)). Co-expression of *grIC* and *grID* with SGR_665, an enzyme similar in sequence to the *Nocardia iowensis* *npt*, can be beneficial.

Gene name	GI Number	GenBank ID	Organism
<i>grIC</i>	182438036	YP_001825755.1	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350
<i>Grid</i>	182438037	YP_001825756.1	<i>Streptomyces griseus</i> subsp. <i>griseus</i> NBRC 13350

[0274] An enzyme with similar characteristics, alpha-amino acid reductase (AAR, EC 1.2.1.31), participates in lysine biosynthesis pathways in some fungal species. This enzyme naturally reduces alpha-amino acid to alpha-amino acid semialdehyde. The carboxyl group is first activated through the ATP-dependent formation of an adenylate that is then reduced by NAD(P)H to yield the aldehyde and AMP. Like CAR, this enzyme utilizes magnesium and requires activation by a PPTase. Enzyme candidates for AAR and its corresponding PPTase are found in *Saccharomyces cerevisiae* (Morris et al., *Gene* 98:141-145 (1991)), *Candida albicans* (Guo et al., *Mol. Genet. Genomics* 269:271-279 (2003)), and *Schizosaccharomyces pombe* (Ford et al., *Curr. Genet.* 28:131-137 (1995)). The AAR from *S. pombe* exhibited significant activity when expressed in *E. coli* (Guo et al., *Yeast* 21:1279-1288 (2004)). The AAR from *Penicillium chrysogenum* accepts S-carboxymethyl-L-cysteine as an alternate substrate, but did not react with adipate, L-glutamate or diaminopimelate (Hijarrubia et al., *J. Biol. Chem.* 278:8250-8256 (2003)). The gene encoding the *P. chrysogenum* PPTase has not been identified to date.

Gene name	GI Number	GenBank ID	Organism
LYS2	171867	AAA34747.1	<i>Saccharomyces cerevisiae</i>
LYS5	1708896	P50113.1	<i>Saccharomyces cerevisiae</i>
LYS2	2853226	AAC02241.1	<i>Candida albicans</i>
LYS5	28136195	AAO26020.1	<i>Candida albicans</i>
Lys1p	13124791	P40976.3	<i>Schizosaccharomyces pombe</i>
Lys7p	1723561	Q10474.1	<i>Schizosaccharomyces pombe</i>
Lys2	3282044	CAA74300.1	<i>Penicillium chrysogenum</i>

FIG. 2, Step L—4-Hydroxybutyryl-phosphate Reductase

[0275] 4-Hydroxybutyrylphosphate reductase catalyzes the reduction of 4-hydroxybutyrylphosphate to 4-hydroxybutyraldehyde. An enzyme catalyzing this transformation has not been identified to date. However, similar enzymes include phosphate reductases in the EC class 1.2.1. Exemplary phosphonate reductase enzymes include glyceraldehyde-3-phosphate dehydrogenase (EC 1.2.1.12), aspartate-semialdehyde dehydrogenase (EC 1.2.1.11) acetylglutamylphosphate reductase (EC 1.2.1.38) and glutamate-5-semialdehyde dehydrogenase (EC 1.2.1.-). Aspartate semialdehyde dehydrogenase (ASD, EC 1.2.1.11) catalyzes the NADPH-dependent reduction of 4-aspartyl phosphate to aspartate-4-semialdehyde. ASD participates in amino acid biosynthesis and recently has been studied as an antimicrobial target (Hadfield et al., *Biochemistry* 40:14475-14483 (2001)). The *E. coli* ASD structure has been solved (Hadfield et al., *J. Mol. Biol.* 289:991-1002 (1999)) and the enzyme has been shown to accept the alternate substrate beta-3-methylaspartyl phosphate (Shames et al., *J. Biol. Chem.* 259:15331-15339 (1984)). The *Haemophilus influenzae* enzyme has been the subject of enzyme engineering studies to alter substrate binding affinities at the active site (Blanco et al., *Acta Crystallogr. D. Biol. Crystallogr.* 60:1388-1395 (2004); Blanco et al., *Acta Crystallogr. D. Biol. Crystallogr.* 60:1808-1815 (2004)). Other ASD candidates are found in *Mycobacterium tuberculosis* (Shafiani et al., *J. Appl. Microbiol.* 98:832-838 (2005)),

Methanococcus jannaschii (Faehle et al., *J Mol. Biol.* 353:1055-1068 (2005)), and the infectious microorganisms *Vibrio cholera* and *Heliobacter pylori* (Moore et al., *Protein Expr. Purif.* 25:189-194 (2002)). A related enzyme candidate is acetylglutamylphosphate reductase (EC 1.2.1.38), an enzyme that naturally reduces acetylglutamylphosphate to acetylglutamate-5-semialdehyde, found in *S. cerevisiae* (Paulwels et al., *Eur. J. Biochem.* 270:1014-1024 (2003)), *B. subtilis* (O'Reilly et al., *Microbiology* 140 (Pt 5):1023-1025 (1994)), *E. coli* (Parsot et al., *Gene.* 68:275-283 (1988)), and other organisms. Additional phosphate reductase enzymes of *E. coli* include glyceraldehyde 3-phosphate dehydrogenase (gapA (Branlant et al., *Eur. J. Biochem.* 150:61-66 (1985))) and glutamate-5-semialdehyde dehydrogenase (proA (Smith et al., *J. Bacteriol.* 157:545-551 (1984))). Genes encoding glutamate-5-semialdehyde dehydrogenase enzymes from *Salmonella typhimurium* (Mahan et al., *J Bacteriol.* 156:1249-1262 (1983)) and *Campylobacter jejuni* (Louie et al., *Mol. Gen. Genet.* 240:29-35 (1993)) were cloned and expressed in *E. coli*.

Protein	GenBank ID	GI Number	Organism
asd	NP_417891.1	16131307	<i>Escherichia coli</i>
asd	YP_248335.1	68249223	<i>Haemophilus influenzae</i>
asd	AAB49996	1899206	<i>Mycobacterium tuberculosis</i>
VC2036	NP_231670	15642038	<i>Vibrio cholera</i>
asd	YP_002301787.1	210135348	<i>Heliobacter pylori</i>
ARG5,6	NP_010992.1	6320913	<i>Saccharomyces cerevisiae</i>
argC	NP_389001.1	16078184	<i>Bacillus subtilis</i>
argC	NP_418393.1	16131796	<i>Escherichia coli</i>
gapA	P0A9B2.2	71159358	<i>Escherichia coli</i>
proA	NP_414778.1	16128229	<i>Escherichia coli</i>
proA	NP_459319.1	16763704	<i>Salmonella typhimurium</i>
proA	P53000.2	9087222	<i>Campylobacter jejuni</i>

FIG. 2, Step M—4-Hydroxybutyryl-CoA Reductase (Alcohol Forming)

[0276] 4-Hydroxybutyryl-CoA reductase (alcohol forming) enzymes are bifunctional oxidoreductases that convert an 4-hydroxybutyryl-CoA to 1,4-butanediol. Enzymes with this activity include adhE from *E. coli*, adhE2 from *C. acetobutylicum* (Fontaine et al., *J. Bacteriol.* 184:821-830 (2002)) and the *C. acetobutylicum* enzymes encoded by bdh I and bdh II (Walter, et al., *J. Bacteriol.* 174:7149-7158 (1992)). In addition to reducing acetyl-CoA to ethanol, the enzyme encoded by adhE in *Leuconostoc mesenteroides* has been shown to oxidize the branched chain compound isobutyraldehyde to isobutyryl-CoA (Kazahaya et al., *J. Gen. Appl. Microbiol.* 18:43-55 (1972); Koo et al., *Biotechnol Lett*, 27:505-510 (2005)).

Protein	GenBank ID	GI Number	Organism
adhE	NP_415757.1	16129202	<i>Escherichia coli</i>
adhE2	AAK09379.1	12958626	<i>Clostridium acetobutylicum</i>
bdh I	NP_349892.1	15896543	<i>Clostridium acetobutylicum</i>
bdh II	NP_349891.1	15896542	<i>Clostridium acetobutylicum</i>
adhE	AAV66076.1	55818563	<i>Leuconostoc mesenteroides</i>
adhE	NP_781989.1	28211045	<i>Clostridium tetani</i>
adhE	NP_563447.1	18311513	<i>Clostridium perfringens</i>
adhE	YP_001089483.1	126700586	<i>Clostridium difficile</i>

4.3 Example III

Methods of Using Formaldehyde Produced from the Oxidation of Methanol in the Formation of Intermediates of Central Metabolic Pathways for the Formation of Biomass

[0277] Provided herein are exemplary pathways, which utilize formaldehyde produced from the oxidation of methanol (see, e.g., FIG. 1, step J) in the formation of intermediates of certain central metabolic pathways that can be used for the formation of biomass. Exemplary methanol metabolic pathways for enhancing the availability of reducing equivalents, as well as the producing formaldehyde from methanol (step J), are provided in FIG. 1.

[0278] One exemplary pathway that can utilize formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1) is shown in FIG. 3, which involves condensation of formaldehyde and D-ribulose-5-phosphate to form hexylose-6-phosphate (h6p) by hexylose-6-phosphate synthase (FIG. 3, step A). The enzyme can use Mg²⁺ or Mn²⁺ for maximal activity, although other metal ions are useful, and even non-metal-ion-dependent mechanisms are contemplated. H6p is converted into fructose-6-phosphate by 6-phospho-3-hexyloisomerase (FIG. 3, step B).

[0279] Another exemplary pathway that involves the detoxification and assimilation of formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1) is shown in FIG. 4 and proceeds through dihydroxyacetone. Dihydroxyacetone synthase is a special transketolase that first transfers a glycoaldehyde group from xylulose-5-phosphate to formaldehyde, resulting in the formation of dihydroxyacetone (DHA) and glyceraldehyde-3-phosphate (G3P), which is an intermediate in glycolysis (FIG. 4, step A). The DHA obtained from DHA synthase is then further phosphorylated to form DHA phosphate by a DHA kinase (FIG. 4, step B). DHAP can be assimilated into glycolysis and several other pathways.

FIG. 3, Steps A and B—Hexylose-6-phosphate Synthase (Step A) and 6-phospho-3-hexyloisomerase (Step B)

[0280] Both of the hexylose-6-phosphate synthase and 6-phospho-3-hexyloisomerase enzymes are found in several organisms, including methanotrophs and methylotrophs where they have been purified (Kato et al., 2006, *BioSci Biotechnol Biochem.* 70(1):10-21. In addition, these enzymes have been reported in heterotrophs such as *Bacillus subtilis* also where they are reported to be involved in formaldehyde detoxification (Mitsui et al., 2003, *AEM* 69(10):6128-32, Yasueda et al., 1999, *J Bac* 181(23):7154-60. Genes for these two enzymes from the methylotrophic bacterium *Mycobacterium gastri* MB19 have been fused and *E. coli* strains harboring the hps-phi construct showed more efficient utilization of formaldehyde (Orita et al., 2007, *Appl Microbiol Biotechnol.* 76:439-445). In some organisms, these two enzymes naturally exist as a fused version that is bifunctional.

[0281] Exemplary candidate genes for hexylose-6-phosphate synthase are:

Protein	GenBank ID	GI number	Organism
Hps	AAR39392.1	40074227	<i>Bacillus methanolicus</i> MGA3
Hps	EIJ81375.1	387589055	<i>Bacillus methanolicus</i> PB1
RmpA	BAA83096.1	5706381	<i>Methylomonas aminofaciens</i>

-continued

Protein	GenBank ID	GI number	Organism
RmpA	BAA90546.1	6899861	<i>Mycobacterium gastri</i>
YckG	BAA08980.1	1805418	<i>Bacillus subtilis</i>

[0282] Exemplary gene candidates for 6-phospho-3-hexyloisomerase are:

Protein	GenBank ID	GI number	Organism
Phi	AAR39393.1	40074228	<i>Bacillus methanolicus</i> MGA3
Phi	EJ81376.1	387589056	<i>Bacillus methanolicus</i> PB1
Phi	BAA83098.1	5706383	<i>Methylomonas aminofaciens</i>
RmpB	BAA90545.1	6899860	<i>Mycobacterium gastri</i>

[0283] Candidates for enzymes where both of these functions have been fused into a single open reading frame include the following.

Protein	GenBank ID	GI number	Organism
PH1938	NP_143767.1	14591680	<i>Pyrococcus horikoshii</i> OT3
PF0220	NP_577949.1	18976592	<i>Pyrococcus furiosus</i>
TK0475	YP_182888.1	57640410	<i>Thermococcus kodakaraensis</i>
	NP_127388.1	14521911	<i>Pyrococcus abyssi</i>
MCA2738	YP_115138.1	53803128	<i>Methylococcus capsulatus</i>

FIG. 4, Step A—Dihydroxyacetone Synthase

[0284] Another exemplary pathway that involves the detoxification and assimilation of formaldehyde produced from the oxidation of methanol (e.g., as provided in FIG. 1) is shown in FIG. 4 and proceeds through dihydroxyacetone. Dihydroxyacetone synthase is a special transketolase that first transfers a glycoaldehyde group from xylulose-5-phosphate to formaldehyde, resulting in the formation of dihydroxyacetone (DHA) and glyceraldehyde-3-phosphate (G3P), which is an intermediate in glycolysis (FIG. 4, step A). The DHA obtained from DHA synthase is then further phosphorylated to form DHA phosphate by a DHA kinase (FIG. 4, step B). DHAP can be assimilated into glycolysis and several other pathways.

[0285] The dihydroxyacetone synthase enzyme in *Candida boidinii* uses thiamine pyrophosphate and Mg^{2+} as cofactors and is localized in the peroxisome. The enzyme from the methanol-growing carboxydobacterium, *Mycobacter* sp. strain JC1 DSM 3803, was also found to have DHA synthase and kinase activities (Ro et al., 1997, JBac 179(19):6041-7). DHA synthase from this organism also has similar cofactor requirements as the enzyme from *C. boidinii*. The K_m s for formaldehyde and xylulose 5-phosphate were reported to be 1.86 mM and 33.3 microM, respectively. Several other mycobacteria, excluding only *Mycobacterium tuberculosis*, can use methanol as the sole source of carbon and energy and are reported to use dihydroxyacetone synthase (Part et al., 2003, JBac 185(1):142-7.

Protein	GenBank ID	GI number	Organism
DAS1	AAC83349.1	3978466	<i>Candida boidinii</i>
HPODL_2613	EFW95760.1	320581540	<i>Ogataea parapolyomorpha</i> DL-1 (<i>Hansenula polymorpha</i> DL-1)
	AAG12171.2	18497328	<i>Mycobacter</i> sp. strain JC1 DSM 3803

FIG. 4, Step B—Dihydroxyacetone (DHA) Kinase

[0286] DHA obtained from DHA synthase is further phosphorylated to form DHA phosphate by a DHA kinase. DHAP can be assimilated into glycolysis and several other pathways. Dihydroxyacetone kinase has been purified from *Ogataea angusta* to homogeneity (Bystrkh, 1983, Biokhimiia, 48(10):1611-6). The enzyme, which phosphorylates dihydroxyacetone and, to a lesser degree, glyceraldehyde, is a homodimeric protein of 139 kDa. ATP is the preferred phosphate group donor for the enzyme. When ITP, GTP, CTP and UTP are used, the activity drops to about 30%. In several organisms such as *Klebsiella pneumoniae* and *Citrobacter freundii* (Daniel et al., 1995, JBac 177(15):4392-40), DHA is formed as a result of oxidation of glycerol and is converted into DHAP by the kinase DHA kinase of *K. pneumoniae* has been characterized (Jonathan et al, 1984, JBac 160(1):55-60). It is very specific for DHA, with a K_m of 4 μ M, and has two apparent K_m values for ATP, one at 25 to 35 μ M, and the other at 200 to 300 μ M. DHA can also be phosphorylated by glycerol kinases but the DHA kinase from *K. pneumoniae* is different from glycerol kinase in several respects. While both enzymes can phosphorylate dihydroxyacetone, DHA kinase does not phosphorylate glycerol, neither is it inhibited by fructose-1,6-diphosphate. In *Saccharomyces cerevisiae*, DHA kinases (I and II) are involved in rescuing the cells from toxic effects of dihydroxyacetone (Molin et al., 2003, J Biol Chem. 17; 278(3):1415-23).

[0287] In *Escherichia coli*, DHA kinase is composed of the three subunits DhaK, DhaL, and DhaM and it functions similarly to a phosphotransferase system (PTS) in that it utilizes phosphoenolpyruvate as a phosphoryl donor (Gutknecht et al., 2001, EMBO J. 20(10):2480-6). It differs in not being involved in transport. The phosphorylation reaction requires the presence of the EI and HPr proteins of the PTS system. The DhaM subunit is phosphorylated at multiple sites. DhaK contains the substrate binding site (Garcia-Alles et al., 2004, 43(41):13037-45; Siebold et al., 2003, PNAS. 100(14):8188-92). The K_M for dihydroxyacetone for the *E. coli* enzyme has been reported to be 6 μ M. The K subunit is similar to the N-terminal half of ATP-dependent dihydroxyacetone kinase of *Citrobacter freundii* and eukaryotes.

[0288] Exemplary DHA kinase gene candidates for this step are:

Protein	GenBank ID	GI number	Organism
DAK1	P54838.1	1706391	<i>Saccharomyces cerevisiae</i> S288c
DAK2	P43550.1	1169289	<i>Saccharomyces cerevisiae</i> S288c

-continued

Protein	GenBank ID	GI number	Organism
D186_20916	ZP_16280678.1	421847542	<i>Citrobacter freundii</i>
DAK2	ZP_18488498.1	425085405	<i>Klebsiella pneumoniae</i>
DAK	AAC27705.1	3171001	<i>Ogataea angusta</i>
DhaK	NP_415718.6	162135900	<i>Escherichia coli</i>
DhaL	NP_415717.1	16129162	<i>Escherichia coli</i>
DhaM	NP_415716.4	226524708	<i>Escherichia coli</i>

4.4 Example IV

Methods for Handling Anaerobic Cultures

[0289] This example describes methods used in handling anaerobic cultures.

[0290] A. Anaerobic Chamber and Conditions.

[0291] Exemplary anaerobic chambers are available commercially (see, for example, Vacuum Atmospheres Company, Hawthorne Calif.; MBraun, Newburyport Mass.). Conditions included an O₂ concentration of 1 ppm or less and 1 atm pure N₂. In one example, 3 oxygen scrubbers/catalyst regenerators were used, and the chamber included an O₂ electrode (such as Teledyne; City of Industry Calif.). Nearly all items and reagents were cycled four times in the airlock of the chamber prior to opening the inner chamber door. Reagents with a volume >5 mL were sparged with pure N₂ prior to introduction into the chamber. Gloves are changed twice/yr and the catalyst containers were regenerated periodically when the chamber displays increasingly sluggish response to changes in oxygen levels. The chamber's pressure was controlled through one-way valves activated by solenoids. This feature allowed setting the chamber pressure at a level higher than the surroundings to allow transfer of very small tubes through the purge valve.

[0292] The anaerobic chambers achieved levels of O₂ that were consistently very low and were needed for highly oxygen sensitive anaerobic conditions. However, growth and handling of cells does not usually require such precautions. In an alternative anaerobic chamber configuration, platinum or palladium can be used as a catalyst that requires some hydrogen gas in the mix. Instead of using solenoid valves, pressure release can be controlled by a bubbler. Instead of using instrument-based O₂ monitoring, test strips can be used instead.

[0293] B. Anaerobic Microbiology.

[0294] Serum or media bottles are fitted with thick rubber stoppers and aluminum crimps are employed to seal the bottle. Medium, such as Terrific Broth, is made in a conventional manner and dispensed to an appropriately sized serum bottle. The bottles are sparged with nitrogen for ~30 min of moderate bubbling. This removes most of the oxygen from the medium and, after this step, each bottle is capped with a rubber stopper (such as Bellco 20 mm septum stoppers; Bellco, Vineland, N.J.) and crimp-sealed (Bellco 20 mm). Then the bottles of medium are autoclaved using a slow (liquid) exhaust cycle. At least sometimes a needle can be poked through the stopper to provide exhaust during autoclaving; the needle needs to be removed immediately upon removal from the autoclave. The sterile medium has the remaining medium components, for example buffer or antibiotics, added via syringe and needle. Prior to addition of reducing agents, the bottles are equilibrated for 30-60 minutes with nitrogen (or CO depending upon use). A reducing agent such as a 100×150 mM sodium sulfide, 200 mM cysteine-HCl is added. This is made by weighing the sodium sulfide into a dry beaker and the cysteine into a serum bottle, bringing both into the anaerobic chamber, dissolving the sodium sulfide into anaerobic water, then adding this to the cysteine in the serum bottle. The bottle is stoppered immediately as the sodium sulfide solution generates hydrogen sulfide gas upon contact with the cysteine. When injecting into the culture, a syringe filter is used to sterilize the solution. Other components are added through syringe needles, such as B12 (10 μM cyanocobalamin), nickel chloride (NiCl₂, 20 microM final concentration from a 40 mM stock made in anaerobic water in the chamber and sterilized by autoclaving or by using a syringe filter upon injection into the culture), and ferrous ammonium sulfate (final concentration needed is 100 μM—made as 100-1000× stock solution in anaerobic water in the chamber and sterilized by autoclaving or by using a syringe filter upon injection into the culture). To facilitate faster growth under anaerobic conditions, the 1 liter bottles were inoculated with 50 mL of a preculture grown anaerobically. Induction of the pA1-lacO1 promoter in the vectors was performed by addition of isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.2 mM and was carried out for about 3 hrs.

[0295] Large cultures can be grown in larger bottles using continuous gas addition while bubbling. A rubber stopper with a metal bubbler is placed in the bottle after medium addition and sparged with nitrogen for 30 minutes or more prior to setting up the rest of the bottle. Each bottle is put together such that a sterile filter will sterilize the gas bubbled in and the hoses on the bottles are compressible with small C clamps. Medium and cells are stirred with magnetic stir bars. Once all medium components and cells are added, the bottles are incubated in an incubator in room air but with continuous nitrogen sparging into the bottles.

[0296] Throughout this application various publications have been referenced. The disclosures of these publications in their entireties, including GenBank and GI number publications, are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains. Although the invention has been described with reference to the examples and embodiments provided above, it should be understood that various modifications can be made without departing from the spirit of the invention.

What is claimed is:

1. A non-naturally occurring microbial organism having a methanol metabolic pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a methanol metabolic pathway enzyme expressed in a sufficient amount to enhance the availability of reducing equivalents in the presence of methanol, wherein said methanol metabolic pathway comprises:

- (i) a methanol methyltransferase and a methylenetetrahydrofolate reductase;
- (ii) a methanol dehydrogenase; or
- (iii) a methanol dehydrogenase and a formaldehyde activating enzyme.

2. The organism of claim 1, wherein the methanol metabolic pathway comprises a methanol methyltransferase and a methylenetetrahydrofolate reductase.

3. The organism of claim 1, wherein the methanol metabolic pathway comprises a methanol dehydrogenase.

4. The organism of claim 1, wherein the methanol metabolic pathway comprises a methanol dehydrogenase and a formaldehyde activating enzyme.

5. The organism of claim 2, wherein the methanol metabolic pathway comprises a methanol methyltransferase, a methylenetetrahydrofolate reductase, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase, and a formyltetrahydrofolate deformylase.

6. The organism of claim 2, wherein the methanol metabolic pathway comprises a methanol methyltransferase, a methylenetetrahydrofolate reductase, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase and a formyltetrahydrofolate synthetase.

7. The organism of claim 3, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase and a formyltetrahydrofolate deformylase.

8. The organism of claim 3, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase and a formyltetrahydrofolate synthetase.

9. The organism of claim 3, wherein the methanol metabolic pathway comprises a methanol dehydrogenase and a formaldehyde dehydrogenase.

10. The organism of claim 3, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a S-(hydroxymethyl)glutathione synthase, a glutathione-dependent formaldehyde dehydrogenase and a S-formylglutathione hydrolase.

11. The organism of claim 3, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a glutathione-dependent formaldehyde dehydrogenase and a S-formylglutathione hydrolase.

12. The organism of claim 4, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a formaldehyde activating enzyme, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase and a formyltetrahydrofolate deformylase.

13. The organism of claim 4, wherein the methanol metabolic pathway comprises a methanol dehydrogenase, a formaldehyde activating enzyme, a methylenetetrahydrofolate dehydrogenase, a methenyltetrahydrofolate cyclohydrolase and a formyltetrahydrofolate synthetase.

14. The organism of any one of claims 5 to 13, wherein the methanol metabolic pathway further comprises a formate dehydrogenase.

15. The organism of any one of claims 5 to 13, wherein the methanol metabolic pathway further comprises a formate hydrogen lyase.

16. The organism of claim 15, wherein the methanol metabolic pathway further comprises a hydrogenase.

17. The organism of any one of claims 1 to 16, wherein said organism comprises two, three, four, five, six or seven exogenous nucleic acids, each encoding a methanol metabolic pathway enzyme.

18. The organism of any one of claims 1 to 17, further comprising a 1,4-butanediol (BDO) pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a BDO pathway enzyme expressed in a sufficient amount to produce BDO, and wherein the BDO pathway comprises:

(i) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hy-

droxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(ii) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(iii) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

(iv) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

(v) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase;

(vi) a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase, and a 1,4-butanediol dehydrogenase;

(vii) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(viii) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

(ix) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(x) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

(xi) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase;

(xii) a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase and a 1,4-butanediol dehydrogenase;

(xiii) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(xiv) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase;

(xv) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

(xvi) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming);

- (xvii) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase; and
- (xviii) a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase, and a 1,4-butanediol dehydrogenase.
- 19.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 20.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 21.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 22.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 23.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase.
- 24.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (aldehyde forming), a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase, and a 1,4-butanediol dehydrogenase.
- 25.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 26.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 27.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 28.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 29.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase.
- 30.** The organism of claim **18**, wherein the BDO pathway comprises a succinyl-CoA reductase (alcohol forming), a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase and a 1,4-butanediol dehydrogenase.
- 31.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 32.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, a 4-hydroxybutyryl-CoA reductase (aldehyde forming), and a 1,4-butanediol dehydrogenase.
- 33.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyryl-CoA transferase or a 4-hydroxybutyryl-CoA synthetase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 34.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a phosphotrans-4-hydroxybutyrylase, and a 4-hydroxybutyryl-CoA reductase (alcohol forming).
- 35.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate reductase, and a 1,4-butanediol dehydrogenase.
- 36.** The organism of claim **18**, wherein the BDO pathway comprises a succinate reductase, a 4-hydroxybutyrate dehydrogenase, a 4-hydroxybutyrate kinase, a 4-hydroxybutyryl-phosphate reductase, and a 1,4-butanediol dehydrogenase.
- 37.** The organism of any one of claims **18** to **30**, wherein the BDO pathway further comprises a succinyl-CoA transferase.
- 38.** The organism of any one of claims **18** to **30**, wherein the BDO pathway further comprises a succinyl-CoA synthetase.
- 39.** The organism of any one of claims **18** to **38**, wherein the organism comprises four, five, six or seven exogenous nucleic acids, each encoding a BDO pathway enzyme.
- 40.** The organism of any one of claims **1** to **39**, further comprising one or more gene disruptions, wherein said one or more gene disruptions occur in one or more endogenous genes encoding protein(s) or enzyme(s) involved in native production of ethanol, glycerol, acetate, lactate, formate, CO₂, and/or amino acids, by said microbial organism, and wherein said one or more gene disruptions confers increased production of BDO in said microbial organism.
- 41.** The non-naturally occurring microbial organism of any one of claims **1** to **40**, wherein one or more endogenous enzymes involved in: native production of ethanol, glycerol, acetate, lactate, formate, CO₂ and/or amino acids by said microbial organism, has attenuated enzyme activity or expression levels.
- 42.** The organism of any one of claims **1**, **3**, **4** or **7** to **41**, further comprising a formaldehyde assimilation pathway, wherein said organism comprises at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme expressed in a sufficient amount to produce an intermediate of glycolysis and/or a metabolic pathway that can be used in the formation of biomass, and wherein said formaldehyde assimilation pathway comprises a hexylose-6-phosphate synthase and a 6-phospho-3-hexyloisomerase.
- 43.** The organism of claim **42**, wherein the intermediate is a hexylose-6-phosphate, a fructose-6-phosphate, or a combination thereof.

44. The organism of any one of claims **1**, **3**, **4** or **7** to **41**, further comprising a formaldehyde assimilation pathways, wherein said organism comprises at least one exogenous nucleic acid encoding a formaldehyde assimilation pathway enzyme expressed in a sufficient amount to produce an intermediate of glycolysis and/or a metabolic pathway that can be used in the formation of biomass, and wherein said formaldehyde assimilation pathway comprises a dihydroxyacetone and a dihydroxyacetone-phosphate.

45. The organism of claim **44**, wherein the intermediate is a dihydroxyacetone, a dihydroxyacetone phosphate, or a combination thereof.

46. The organism of any one of claims **42** to **45**, wherein the organism comprises two exogenous nucleic acids, each encoding a formaldehyde assimilation pathway enzyme.

47. The organism of any one of claims **1** to **46**, wherein said at least one exogenous nucleic acid is a heterologous nucleic acid.

48. The organism of any one of claims **1** to **47**, wherein said organism is in a substantially anaerobic culture medium.

49. The non-naturally occurring microbial organism of any one of claims **1** to **48**, wherein said microbial organism is a species of bacteria, yeast, or fungus.

50. A method for producing BDO, comprising culturing the organism of any one of claims **18** to **49** under conditions and for a sufficient period of time to produce BDO.

51. A bioderived BDO produced according to the method of claim **50**.

52. A culture medium comprising the bioderived BDO of claim **51**.

53. The culture medium of claim **52**, wherein said bioderived BDO has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide uptake source.

54. The culture medium of claim **52** or **53**, wherein said culture medium is separated from the non-naturally occurring microbial organism having the BDO pathway.

55. The bioderived BDO of claim **51**, wherein said bioderived BDO has a carbon-12, carbon-13 and carbon-14 isotope ratio that reflects an atmospheric carbon dioxide uptake source.

56. The bioderived BDO of claim **51** or **55**, wherein said bioderived BDO has an Fm value of at least 80%, at least 85%, at least 90%, at least 95% or at least 98%.

57. A composition comprising said bioderived BDO of any one of claim **51**, **55** or **56**, and a compound other than said bioderived BDO.

58. The composition of claim **57** wherein said compound other than said bioderived BDO is a trace amount of a cellular portion of a non-naturally occurring microbial organism having a BDO pathway.

59. A biobased product comprising said bioderived BDO of claim **51**, **55** or **56**, wherein said biobased product is

- (i) a polymer, THF or a THF derivative, or GBL or a GBL derivative;
- (ii) a plastic, elastic fiber, polyurethane, polyester, polyhydroxyalkanoate, poly-4-hydroxybutyrate, co-polymer of poly-4-hydroxybutyrate, poly(tetramethylene ether) glycol, polyurethane-polyurea copolymer, spandex, elastane, Lycra™, or nylon;

- (iii) a polymer, a resin, a fiber, a bead, a granule, a pellet, a chip, a plastic, a polyester, a thermoplastic polyester, a molded article, an injection-molded article, an injection-molded part, an automotive part, an extrusion resin, an electrical part and a casing; and optionally where the biobased product is reinforced or filled and further where the biobased product is glass-reinforced or -filled or mineral-reinforced or -filled;

- (iv) a polymer, wherein the polymer comprises polybutylene terephthalate (PBT);

- (v) a polymer, wherein the polymer comprises PBT and the biobased product is a resin, a fiber, a bead, a granule, a pellet, a chip, a plastic, a polyester, a thermoplastic polyester, a molded article, an injection-molded article, an injection-molded part, an automotive part, an extrusion resin, an electrical part and a casing; and optionally where the biobased product is reinforced or filled and further where the biobased product is glass-reinforced or -filled or mineral-reinforced or -filled;

- (vi) a THF or a THF derivative, wherein the THF derivative is polytetramethylene ether glycol (PTMEG), a polyester ether (COPE) or a thermoplastic polyurethane;

- (viii) a THF derivative, wherein the THF derivative comprises a fiber; or

- (ix) a GBL or a GBL derivative, wherein the GBL derivative is a pyrrolidone.

60. The biobased product of claim **59**, comprising at least 5%, at least 10%, at least 20%, at least 30%, at least 40% or at least 50% bioderived BDO.

61. The biobased product of claim **59** or **60**, wherein said biobased product comprises a portion of said bioderived BDO as a repeating unit.

62. A molded product obtained by molding the biobased product of any one of claims **59** to **61**.

63. A process for producing the biobased product of any one of claims **59** to **61** comprising chemically reacting said bioderived BDO with itself or another compound in a reaction that produces said biobased product.

64. A polymer comprising or obtained by converting the bioderived BDO of claim **51**, **55** or **56**.

65. A method for producing a polymer, comprising chemically or enzymatically converting the bioderived BDO of claim **51**, **55** or **56** to the polymer.

66. A composition comprising the bioderived BDO of claim **51**, **55** or **56**, or a cell lysate or culture supernatant thereof.

67. A method of producing formaldehyde, comprising culturing the organism of any one of claims **1**, **3**, **4** or **7** to **49** under conditions and for a sufficient period of time to produce formaldehyde, and optionally wherein the formaldehyde is consumed to provide a reducing equivalent or to incorporate into BDO or target product.

68. A method of producing an intermediate of glycolysis and/or an intermediate of a metabolic pathway that can be used in the formation of biomass, comprising culturing the organism of any one of claims **42** to **49** under conditions and for a sufficient period of time to produce the intermediate, and optionally wherein the intermediate is consumed to provide a reducing equivalent or to incorporate into BDO or target product.

* * * * *