



(19) **United States**

(12) **Patent Application Publication**

**Ashkenazi et al.**

(10) **Pub. No.: US 2003/0064923 A1**

(43) **Pub. Date: Apr. 3, 2003**

(54) **SECRETED AND TRANSMEMBRANE POLYPEPTIDES AND NUCLEIC ACIDS ENCODING THE SAME**

(75) Inventors: **Avi Ashkenazi**, San Mateo, CA (US); **David Botstein**, Belmont, CA (US); **Luc Desnoyers**, San Francisco, CA (US); **Dan L. Eaton**, San Rafael, CA (US); **Napoleone Ferrara**, San Francisco, CA (US); **Ellen Filvaroff**, San Francisco, CA (US); **Sherman Fong**, Alameda, CA (US); **Wei-Qiang Gao**, Palo Alto, CA (US); **Hanspeter Gerber**, San Francisco, CA (US); **Mary E. Gerritsen**, San Mateo, CA (US); **Audrey Goddard**, San Francisco, CA (US); **Paul J. Godowski**, Burlingame, CA (US); **J. Christopher Grimaldi**, San Francisco, CA (US); **Austin L. Gurney**, Belmont, CA (US); **Kenneth J. Hillan**, San Francisco, CA (US); **Ivar J. Kljavin**, Lafayette, CA (US); **Jennie P. Mather**, Millbrae, CA (US); **James Pan**, Belmont, CA (US); **Nicholas F. Paoni**, Belmont, CA (US); **Margaret Ann Roy**, San Francisco, CA (US); **Timothy A. Stewart**, San Francisco, CA (US); **Daniel Tumas**, Orinda, CA (US); **P. Mickey Williams**, Half Moon Bay, CA (US); **William I. Wood**, Hillsborough, CA (US)

Provisional application No. 60/063,486, filed on Oct. 21, 1997. Provisional application No. 60/062,816, filed on Oct. 24, 1997. Provisional application No. 60/062,814, filed on Oct. 24, 1997. Provisional application No. 60/063,127, filed on Oct. 24, 1997. Provisional application No. 60/063,120, filed on Oct. 24, 1997. Provisional application No. 60/063,121, filed on Oct. 24, 1997. Provisional application No. 60/063,045, filed on Oct. 24, 1997. Provisional application No. 60/063,128, filed on Oct. 24, 1997. Provisional application No. 60/063,329, filed on Oct. 27, 1997. Provisional application No. 60/063,327, filed on Oct. 27, 1997. Provisional application No. 60/063,549, filed on Oct. 28, 1997. Provisional application No. 60/063,541, filed on Oct. 28, 1997. Provisional application No. 60/063,550, filed on Oct. 28, 1997. Provisional application No. 60/063,542, filed on Oct. 28, 1997. Provisional application No. 60/063,544, filed on Oct. 28, 1997. Provisional application No. 60/063,564, filed on Oct. 28, 1997. Provisional application No. 60/063,734, filed on Oct. 29, 1997. Provisional application No. 60/063,738, filed on Oct. 29, 1997. Provisional application No. 60/063,704, filed on Oct. 29, 1997. Provisional application No. 60/063,435, filed on Oct. 29, 1997. Provisional application No. 60/064,215, filed on Oct. 29, 1997. Provisional application No. 60/063,735, filed on Oct. 29, 1997. Provisional application No. 60/063,732, filed on Oct. 29, 1997. Provisional application No. 60/064,103, filed on Oct. 31, 1997. Provisional application No. 60/063,870, filed on Oct. 31, 1997. Provisional application No. 60/064,248,

Correspondence Address:  
**KNOBBE, MARTENS, OLSON & BEAR, LLP**  
**2040 MAIN STREET**  
**FOURTEENTH FLOOR**  
**IRVINE, CA 92614 (US)**

(List continued on next page.)

(73) Assignee: **Genentech, Inc.**

(30) **Foreign Application Priority Data**

Sep. 10, 1998 (WO)..... PCT/US98/18824  
Sep. 14, 1998 (WO)..... PCT/US98/19177

(21) Appl. No.: **09/905,348**

(List continued on next page.)

(22) Filed: **Jul. 13, 2001**

**Publication Classification**

**Related U.S. Application Data**

(63) Continuation of application No. 09/665,350, filed on Sep. 18, 2000.

(51) **Int. Cl.<sup>7</sup>** ..... **A61K 38/17**; C12Q 1/68;  
G01N 33/53; C12P 21/06;  
C12N 9/00; C07K 14/435;  
C12N 5/06  
(52) **U.S. Cl.** ..... **514/12**; 435/6; 435/325; 435/320.1;  
435/69.1; 435/7.1; 435/183;  
530/350; 536/23.2

(60) Provisional application No. 60/059,115, filed on Sep. 17, 1997. Provisional application No. 60/059,184, filed on Sep. 17, 1997. Provisional application No. 60/059,122, filed on Sep. 17, 1997. Provisional application No. 60/059,117, filed on Sep. 17, 1997. Provisional application No. 60/059,113, filed on Sep. 17, 1997. Provisional application No. 60/059,121, filed on Sep. 17, 1997. Provisional application No. 60/059,119, filed on Sep. 17, 1997. Provisional application No. 60/059,263, filed on Sep. 18, 1997. Provisional application No. 60/059,266, filed on Sep. 18, 1997. Provisional application No. 60/062,125, filed on Oct. 15, 1997. Provisional application No. 60/062,287, filed on Oct. 17, 1997. Provisional application No. 60/062,285, filed on Oct. 17, 1997. Provisional appli-

(57) **ABSTRACT**

The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

Related U.S. Application Data	(30)	Foreign Application Priority Data
filed on Nov. 3, 1997. Provisional application No. 60/064,809, filed on Nov. 7, 1997. Provisional application No. 60/065,186, filed on Nov. 12, 1997. Provisional application No. 60/065,846, filed on Nov. 17, 1997. Provisional application No. 60/065,693, filed on Nov. 18, 1997. Provisional application No. 60/066,120, filed on Nov. 21, 1997. Provisional application No. 60/066,364, filed on Nov. 21, 1997. Provisional application No. 60/066,772, filed on Nov. 24, 1997. Provisional application No. 60/066,466, filed on Nov. 24, 1997. Provisional application No. 60/066,770, filed on Nov. 24, 1997. Provisional application No. 60/066,511, filed on Nov. 24, 1997. Provisional application No. 60/066,453, filed on Nov. 24, 1997. Provisional application No. 60/066,840, filed on Nov. 25, 1997. Provisional application No. 60/069,425, filed on Dec. 12, 1997. Provisional application No. 60/088,026, filed on Jun. 4, 1998. Provisional application No. 60/099,803, filed on Sep. 10, 1998. Provisional application No. 60/100,262, filed on Sep. 14, 1998. Provisional application No. 60/100,858, filed on Sep. 17, 1998. Provisional application No. 60/104,080, filed on Oct. 13, 1998. Provisional application No. 60/109,304, filed on Nov. 20, 1998. Provisional application No. 60/113,296, filed on Dec. 22, 1998. Provisional application No. 60/143,048, filed on Jul. 7, 1999. Provisional application No. 60/145,698, filed on Jul. 26, 1999. Provisional application No. 60/146,222, filed on Jul. 28, 1999.	Sep. 16, 1998 Sep. 17, 1998 Dec. 1, 1998 Sep. 8, 1999 Sep. 13, 1999 Sep. 15, 1999 Sep. 15, 1999 Oct. 5, 1999 Nov. 29, 1999 Nov. 30, 1999 Dec. 1, 1999 Dec. 2, 1999 Dec. 2, 1999 Dec. 16, 1999 Dec. 20, 1999 Dec. 20, 1999 Jan. 5, 2000 Feb. 11, 2000 Feb. 22, 2000 Feb. 24, 2000 Mar. 2, 2000 Mar. 20, 2000 Mar. 30, 2000 May 22, 2000 Jun. 2, 2000 Jul. 28, 2000 Aug. 24, 2000	(WO)..... PCT/US98/19330 (WO)..... PCT/US98/19437 (WO)..... PCT/US98/25108 (WO)..... PCT/US99/20594 (WO)..... PCT/US99/20944 (WO)..... PCT/US99/21090 (WO)..... PCT/US99/21547 (WO)..... PCT/US99/23089 (WO)..... PCT/US99/28214 (WO)..... PCT/US99/28313 (WO)..... PCT/US99/28301 (WO)..... PCT/US99/28564 (WO)..... PCT/US99/28565 (WO)..... PCT/US99/30095 (WO)..... PCT/US99/30999 (WO)..... PCT/US99/30911 (WO)..... PCT/US00/00219 (WO)..... PCT/US00/03565 (WO)..... PCT/US00/04414 (WO)..... PCT/US00/05004 (WO)..... PCT/US00/05841 (WO)..... PCT/US00/07377 (WO)..... PCT/US00/08439 (WO)..... PCT/US00/14042 (WO)..... PCT/US00/15264 (WO)..... PCT/US00/20710 (WO)..... PCT/US00/23328

**FIGURE 1**

ACTGCACCTCGGTTCTATCGATTGAATTCCCCGGGGATCCTCTAGAGATCCCTCGACCTCGA  
CCCACGCGTCCGGGCCGGAGCAGCACGGCCGAGGACCTGGAGCTCCGGCTGCGTCTTCCCG  
CAGCGCTACCCGCCATGCGCCTGCCGCGCCGGGCCGCGCTGGGGCTCCTGCCGCTTCTGCTG  
CTGCTGCCGCCCGCGCCGGAGGCCCAAGAAGCCGACGCCCTGCCACCGGTGCCGGGGGCT  
GGTGGACAAGTTTAAACCAGGGGATGGTGGACACCGCAAAGAAGAACTTTGGCGCGGGAACA  
CGGCTTGGGAGGAAAAGACGCTGTCCAAGTACGAGTCCAGCGAGATTGCGCTGCTGGAGATC  
CTGGAGGGGCTGTGCGAGAGCAGCGACTTCGAATGCAATCAGATGCTAGAGGGCGCAGGAGGA  
GCACCTGGAGGCCTGGTGGCTGCAGCTGAAGAGCGAATATCCTGACTTATTGAGTGGTTTT  
GTGTGAAGACACTGAAAGTGTGCTGCTCTCCAGGAACCTACGGTCCCAGCTGTCTCGCATGC  
CAGGGCGGATCCCAGAGGCCCTGCAGCGGAATGGCCACTGCAGCGAGATGGGAGCAGACA  
GGGGCAGGGTCTCGCGGTGCCACATGGGGTACCAGGGCCCGCTGTGCACTGACTGCATGG  
ACGGCTACTTCAGCTCGCTCCGGAACGAGACCCACAGCATCTGCACAGCCTGTGACGAGTCC  
TGCAAGACGTGCTCGGGCTGACCAACAGAGACTGCGGCGAGTGTGAAGTGGGCTGGGTGCT  
GGACGAGGGCCCTGTGTGGATGTGGACGAGTGTGCGGCCGAGCCGCCTCCCTGCAGCGCTG  
CGCAGTTCTGTAAGAACGCCAACGGCTCCTACAGTGCAGAGAGTGTGACTCCAGCTGTGTG  
GGCTGCACAGGGGAAAGGCCAGGAACTGTAAAGAGTGTATCTCTGGCTACCGAGGGAGCA  
CGGACAGTGTGCAGATGTGGACGAGTGTCACTAGCAGAAAAAACCCTGTGTGAGGAAAAACG  
AAAACTGCTACAATACTCCAGGGAGCTACGTCTGTGTGTCTCTGACGGCTTCGAAGAAACG  
GAAGATGCCCTGTGTGCCCGCGCAGAGGCTGAAGCCACAGAAGGAGAAAGCCCGACACAGCT  
GCCCTCCCGCGAAGACCTGTAATGTGCCGACTTACCCTTTAAATTTATTGAGAAGGATGTCC  
CGTGAAAATGTGCCCTGAGGATGCCGTCTCCTGCAGTGGACAGCGGCGGGGAGAGGCTGC  
CTGCTCTTAACGGTTGATTCTCATTTGTCCCTTAAACAGCTGCATTTCTTGGTTGTTCTTA  
AACAGACTTGATATTTTGTATACAGTCTTTGTAATAAAATGACCATTGTAGGTAATCAGG  
AGGAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGCCGCGACTCTAGAGTGCACCTGCAGAAGC  
TTGGCCGCATGGCCCAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCA  
TCACAAATTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTGTCCAAATC  
ATCAATGTATCTTATCATGTCTGGATCGGGAATTAATTCGGCGCAGCACCATGGCCTGAAAT  
AACCTCTGAAAGAGGAACTTGGTTAGGTACCTTCTGAGGCGGAAAGAACCAGCTGTGGAATG  
TGTGTCAAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCAGCAGGCAGAAGTATGCAAGCATGC  
ATCTCAATTAGTCAGCAACCCAGTTTT

## FIGURE 2

><subunit 1 of 1, 353 aa, 0 stop  
><MW: 38192, pI: 4.53, NX(S/T): 2  
MRLPRRAALGLLPLLLLLLPPAPEAAKPTPCHRCRGLVDKFNQGMVDTAKKNFGGNTAWEKTLSESEIRL  
LEILEGLCESSDFECNQMLEAQEHLLEAWLQLKSEYPDLFEWFCVKTLKVVCCSPGTYGPDCLACQGGSSQRPCSG  
NGHCSDGSRQGDGSCRCHMGYQGPLCTDCMDGYFSSLRNETHSICTACDESCKTCSGLTNRDCGECEVGVWLDE  
GACVDVDECAAEPPPCSAQAQFCKNANGSYTCEECDSSCVGCTGEGPGNCKECISGYAREHGQCADVDECSLAET  
CVRKNENYNTPGSYVCVCPDGFBEETEDACVPPAEAEATEGESPTQLPSREDL

**Signal peptide:**

amino acids 1-24

**N-glycosylation sites.**

amino acids 190-194 and 251-255

**Glycosaminoglycan attachment sites.**

amino acids 149-153 and 155-159

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 26-30

**Casein kinase II phosphorylation sites.**

amino acids 58-62, 66-70, 86-90, 197-201, 210-214, 255-259, 295-299, 339-343  
and 349-353

**Tyrosine kinase phosphorylation site.**

amino acids 303-310

**N-myristoylation sites.**

amino acids 44-50, 54-60, 55-61, 81-87, 150-156, 158-164, 164-170, 252-258 and  
313-319

**Aspartic acid and asparagine hydroxylation site.**

amino acids 308-320

**EGF-like domain cysteine pattern signature.**

amino acids 166-178

**Leucine zipper pattern.**

amino acids 94-116

**FIGURE 3**

CAGGTCCAACCTGCACCTCGGTTCTATCGATTGAATTCCTCCGGGGATCCTCTAGAGATCCCTC  
GACCTCGACCCACGCGTCCGCCAGGCCGGGAGGCGACGCGCCAGCCGTCTAAACGGGAACA  
GCCCTGGCTGAGGGAGCTGCAGCGCAGCAGAGTATCTGACGGCGCCAGGTTGCGTAGGTGCG  
GCACGAGGAGTTTTCCCGGCAGCGAGGAGGTTCTGAGCAGCATGCCCCGGAGGAGCGCCTTC  
CCTGCCGCCGCGCTCTGGCTCTGGAGCATCCTCCTGTGCCTGCTGGCACTGCGGGCGGAGGC  
CGGGCCGCCGAGGAGGAGAGCCTGTACCTATGGATCGATGCTCACCAGGCAAGAGTACTCA  
TAGGATTTGAAGAAGATATCCTGATTTGTTTCAGAGGGGAAAATGGCACCTTTTACACATGAT  
TTCAGAAAAGCGCAACAGAGAATGCCAGCTATTCCTGTCAATATCCATTCCATGAATTTTAC  
CTGGCAAGCTGCAGGGCAGGCAGAATACTTCTATGAATTCCTGTCTTGGCTCCCTGGATA  
AAGGCATCATGGCAGATCCAACCGTCAATGTCCCTCTGCTGGGAACAGTGCCCTCAAAAGCA  
TCAGTTGTTCAAGTTGGTTTTCCCATGTCCTGGAAAACAGGATGGGGTGGCAGCATTTGAAGT  
GGATGTGATTTGTTATGAATTTCTGAAGGCAACACCATTCTCCAAAACCTCAAAAATGCTATCT  
TCTTTAAAACATGTCAACAAGCTGAGTGCCAGGCGGGTGCCGAAAATGGAGGCTTTTGTAAAT  
GAAAGACGCATCTGCGAGTGTCTGATGGGTTCCACGGACCTCACTGTGAGAAAAGCCCTTTG  
TACCCACAGATGTATGAATGGTGGACTTTGTGTGACTCCTGGTTTCTGCATCTGCCACCTG  
GATTTCTATGGAGTGAACCTGTGACAAAAGCAAACTGCTCAACCACCTGCTTTAATGGAGGGACC  
TGTTTCTACCCTGGAAAATGTATTTGCCCTCCAGGACTAGAGGGAGAGCAGTGTGAAAATCAG  
CAAATGCCACAAACCTGTGCAAAATGGAGGTAATGCATTGGTAAAAGCAAATGTAAGTGT  
CCAAAGGTTACCAGGGAGACCTCTGTTCAAAGCCTGTCTGCGAGCCTGGCTGTGGTGCACAT  
GGAACCTGCCATGAACCCAAACAATGCCAATGTCAAGAAGGTTGGCATGGAAGACACTGCAA  
TAAAAGGTACGAAGCCAGCCTCATAACATGCCCTGAGGCCAGCAGGCCGCCAGCTCAGGCAGC  
ACACGCCTTCACTTAAAAAGGCCGAGGAGCGGGGATCCACCTGAATCCAATTACATCTGG  
TGAACTCCGACATCTGAAACGTTTTAAGTTACACCAAGTTCATAGCCTTTGTAAACCTTTCA  
TGTGTTGAATGTTCAAATAATGTTTATTACACTTAAGAATACTGGCCTGAATTTTATTAGCT  
TCATTATAAATCACTGAGCTGATAATTTACTCTTCTTTTAAAGTTTTCTAAGTACGTCTGTAG  
CATGATGGTATAGATTTTTCTTGTTCAGTGCCTTGGGACAGATTTTATATTATGTCAATTGA  
TCAGGTTAAAATTTTTAGTGTGTAGTTGGCAGATATTTTCAAATTAACAATGCATTTATGGT  
GTCTGGGGGCAGGGGAACATCAGAAAGGTTAAATTTGGGCAAAAATGCGTAAGTACAAGAAT  
TTGCATGGTGCAGTTAATGTTGAAGTTACAGCATTTTCAGATTTTATGTCAGATATTTAGAT  
GTTTGTACATTTTTAAAAAATGCTCTTAATTTTTTAAACTCTCAATACAATATATTTTGACC  
TTACCATTATTCCAGAGATTCAGTATTAATAAAAAAAAAAATTAACACTGTGGTAGTGGCATTT  
AAACAATATAATATATTTCTAAACACAATGAAATAGGGAATATAATGTATGAACTTTTTGCAT  
TGGCTTTGAAGCAATATAATATATTTGTAACAAAACACAGCTTTACCTAATAAACATTTTAT  
ACTGTTTGTATGTATAAAAATAAAGGTGCTGCTTTAGTTTTTTGGAAAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCGACTCTAGAGTCGACCTGCAGAAGCTTGGC  
CGCCATGGCCCAACTTGTTTTATTGCAGCTTATAATG

## FIGURE 4

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA33094
><subunit 1 of 1, 379 aa, 0 stop
><MW: 41528, pI: 7.97, NX(S/T): 2
MARRSAFPAAALWLSILLCLLALRAEAGPPQEEESLYLWIDAHQARVLIGFEEDILIVSEGK
MAPFTHDFRKAQQRMPAIPVNIHSMNFTWQAAGQAEYFYEFSLRSLDKGIMADPTVNVPLL
GTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNTILQTPQNAIFFKTCQQAECPPGC
RNGGFCNERRICECPDGFHGHCEKALCTPRCMNGGLCVTPGFICPPGFYGVNCDKANCST
TCFNGGTCFYPGKICPPGLEGEQCEISKCPQPCRNGGKICGKSKCKCSKGYQGDLCSPVC
EPGCGAHGTCHEPNKCQCQEGWHGRHCNKRYEASLIHALRPAGAQLRQHTPSLKKAEERRDP
PESNYIW
```

**Signal peptide:**

amino acids 1-28

**N-glycosylation site.**

amino acids 88-92, 245-249

**Casein kinase II phosphorylation site.**

amino acids 319-323

**Tyrosine kinase phosphorylation site.**

amino acids 370-378

**N-myristoylation sites.**

amino acids 184-190, 185-191, 189-195, 315-321

**ATP/GTP-binding site motif A (P-loop).**

amino acids 285-293

**EGF-like domain cysteine pattern signature.**

amino acids 198-210, 230-242, 262-274, 294-306, 326-338

**FIGURE 5**

CGGACGCGTGGGCGTCCGGCGGTCCGACAGCCAGGAGGCGGAGGCGCGCGGGCCAGCCTGGG  
CCCCAGCCACACCTTCACCAGGGCCCAGGAGCCACCATGTTGGCGATGTCCACTGGGGCTAC  
TGCTGTGCTGCCGCTGGCTGGCCACTTGGCTCTGGGTGCCCAGCAGGGTCGTGGGCGCCGG  
GAGCTAGCACCGGGTCTGCACCTGCGGGGCATCCGGGACGCGGGAGGCGGTACTGCCAGGA  
GCAGGACCTGTGCTGCCGCGGGCCGTGCCGACGACTGTGCCCTGCCCTACCTGGGCGCCATCT  
GTTACTGTGACCTCTTCTGCAACCGCACGGTCTCCGACTGCTGCCCTGACTTCTGGGACTTC  
TGCCTCGGCGTGCCACCCCTTTTCCCCGATCCAAGGATGTATGCATGGAGGTCGTATCTA  
TCCAGTCTTGGGAACGTAAGGACAACCTGTAACCGTTGCACCTGCCAGGAGAACAGGCAGT  
GGCATGGTGGATCCAGACATGATCAAAGCCATCAACCAGGGCAACTATGGCTGGCAGGCTGG  
GAACCACAGCGCCTTCTGGGGCATGACCCTGGATTGAGGGCATTGCTACCGCCTGGGCACCA  
TCCGCCCATCTTCCCTCGGTATGAACATGCATGAAATTTATACAGTGTGAACCCAGGGGGAG  
GTGCTTCCACAGCCTTCCAGGGCCTCTGAGAAGTGGCCCAACCTGATTTCATGAGCCTCTTGA  
CCAAGGCAACTGTGCAGGCTCCTGGGCCTTCTCCACAGCAGCTGTGGCATCCGATCGTGTCT  
CAATCCATTCTCTGGGACACATGACGCCCTGTCTGTGCCCCAGAACCTGCTGTCTTGTGAC  
ACCCACCAGCAGCAGGGCTGCCGCGGTGGGCGTCTCGATGGTGCCTGGTGTCTTCCGCGTCC  
CCGAGGGGTGGTGTCTGACCACTGCTACCCCTTCTCGGGCCGTGAACGAGACGAGGCTGGCC  
CTGCGCCCCCTGTATGATGCACAGCCGAGCCATGGGTGGGGCAAGCGCCAGGCCACTGCC  
CACTGCCCAACAGCTATGTTAATAACAATGACATCTACCAGGTCCTCTGTCTACCGCCT  
CGGCTCCAACGACAAGGAGATCATGAAGGAGCTGATGGAGAATGGCCCTGTCCAAGCCCTCA  
TGGAGGTGCATGAGGACTTCTTCTTATAACAAGGGAGGCATCTACAGCCACACGCCAGTGAGC  
CTTGGGAGGCCAGAGAGATACCGCCGGCATGGGACCCACTCAGTCAAGATCACAGGATGGGG  
AGAGGAGACGCTGCCAGATGGAAGGACGCTCAAATACTGGACTGCGGCCAACTCCTGGGGCC  
CAGCCTGGGGCGAGAGGGGGCCACTTCCGCATCGTGC GCGGGCTCAATGAGTGGCAGATCGAG  
AGCTTCGTGCTGGGCGTCTGGGGCCGCGTGGGCATGGAGGACATGGGTCACTACTGAGGCTG  
CGGGCACACGCGGGGTCCGGCCTGGGATCCAGGCTAAGGGCCGGCGGAAGAGGCCCAATG  
GGGCGGTGACCCAGCCTCGCCGACAGAGCCCGGGGCGCAGGCGGGCGCCAGGGCGCTAAT  
CCC GGCGGGGTTCCGCTGACGCAGCGCCCCGCTGGGAGCCGCGGGCAGGCGAGACTGGCG  
GAGCCCCAGACCTCCAGTGGGGACGGGGCAGGGCCTGGCCTGGGAAGAGCACAGCTGCAG  
ATCCCAGGCCTCTGGGCGCCCCACTCAAGACTACCAAAGCCAGGACACCTCAAGTCTCCAGC  
CCCAATACCCCAACCCCAATCCCCTATTTCTTTTTTTTTTTTTTTAGACAGGGTCTTGCTCCG  
TTGCCAGGTTGGAGTGCAGTGGCCATCAGGGCTCACTGTAACCTCCGACTCCTGGGTCA  
AGTGACCTCCACCTCAGCCTCTCAAGTAGCTGGGACTACAGGTGCACCACCACCTGGC  
TAATTTTTGTATTTTTTGTAAAGAGGGGGTCTCACTGTGTTGCCAGGCTGGTTTCGAACT  
CCTGGGCTCAAGCGGTCCACCTGCCTCCGCTCCCAAAGTGCTGGGATTGCAGGCATGAGCC  
ACTGCACCCAGCCCTGTATTCTTATCTTCAGATATTTATTTTTCTTTTCACTGTTTTAAAA  
TAAAACCAAAGTATTGATAAAAAAAAAA

## FIGURE 6

></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA33223

><subunit 1 of 1, 164 aa, 1 stop

><MW: 18359, pI: 7.45, NX(S/T): 1

MWRCPLGLLLLLLPLAGHLALGAQQGRGRRELAPGLHLRGIRDAGGRYCQEQLCCRGRADDC  
ALPYLGAICYCDLFCNRTVSDCCPDFWDFCLGVPPFPPIQGCMHGGRIYPVLGTYWDNCNR  
CTCQENRQWHGGSRHDQSHQPGQLWLAGWEPQRLLGHDPG

### **N-glycosylation site.**

amino acids 78-82, 161-165

### **Casein kinase II phosphorylation site.**

amino acids 80-84, 117-121, 126-130, 169-173, 205-209, 296-300,  
411-415

### **N-myristoylation site.**

amino acids 21-27, 39-45, 44-50, 104-110, 160-164, 224-230,  
269-275, 378-384, 442-448

### **Amidation site.**

amino acids 26-30, 318-322

### **Eukaryotic thiol (cysteine) proteases histidine active site.**

amino acids 398-409



**FIGURE 7**

AGGCTCCTTGGCCCTTTTTCACAGCAAGCTTNTGCNATCCCGATTGTTGTCTCAAATCCA  
ATTCTCTTGGGACACATNACGCCTGTCCTTTNGCCCCAGAACCTGCTGTCTTGTACACCCAC  
CAGCAGCAGGGCTGCCGCGNTGGGCGTCTCGATGGTGCCTGGTGGTTCTTGCCTCGCCGAGG  
GNTGGTGTCTGACCACTGCTACCCCTTCTCGGGCCGTGAACGAGACGAGGCTGGCCCTGCGC  
CCCCCTGTATGATGCACAGCCGAGCCATGGGTCCGGGGCAAGCGCCAGGCCACTGCCCACTGC  
CCCAACAGCTATGTTAATAACAATGACATCTACCAGGTCACTCCTGTCTACCGCCTCGGCTC  
CAACGACAAGGAGATCATGAAGGAGCTGATGGAGAATGGCCCTGTCCAAGCCCTCATGGAGG  
TGCATGAGGACTTCTTCCCTATAACAAGGGAGGCATCTACAGCCACACGCCAGTGAGCCTTGGG  
AGGCCAGAGAGATAACCGCCGGCATGGGACCCACTCAG

**FIGURE 8**

GCTGCTTGCCCTGTTGATGGCAGGCTTGGCCCTGCAGCCAGGCACTGCCCTGCTGTGCTACT  
CCTGCAAAGCCCAGGTGAGCAACGAGGACTGCCTGCAGGTGGAGAACTGCACCCAGCTGGGG  
GAGCAGTGCTGGACCGCGGCATCCGCGCAGTTGGCCTCCTGACCGTCATCAGCAAAGGCTG  
CAGCTTGAAC TGCGTGGATGACTCACAGGACTACTACGTGGGCAAGAAGAACATCACGTGCT  
GTGACACCGACTTGTGCAACGCCAGCGGGGCCATGCCCTGCAGCCGGCTGCCGCCATCCTT  
GCGCTGCTCCCTGCACTCGGCCTGCTGCTCTGGGGACCCGGCCAGCTATAGGCTCTGGGGGG  
CCCCGCTGCAGCCCACACTGGGTGTGGTGGCCAGGCCTCTGTGCCACTCCTCACAGACCTG  
GCCCAGTGGGAGCCTGTCTGGTTCTGAGGCACATCCTAACGCAAGTCTGACCATGTATGT  
CTGCACCCCTGTCCCCACCCCTGACCCCTCCCATGGCCCTCTCCAGGACTCCCACCCGGCAGA  
TCAGCTCTAGTGACACAGATCCGCCTGCAGATGGCCCTCCAACCCTCTCTGCTGCTGTTTC  
CATGGCCAGCATCTCCACCCTTAACCCTGTGCTCAGGCACCTCTTCCCCCAGGAAGCCTT  
CCCTGCCACCCCATCTATGACTTGAGCCAGGTCTGGTCCGTGGTGTCCCCGCACCCAGCA  
GGGGACAGGCACTCAGGAGGGCCAGTAAAAGGCTGAGATGAAGTGGACTGAGTAGAACTGGA  
GGACAAGAGTCGACGTGAGTTCCTGGGAGTCTCCAGAGATGGGGCCTGGAGGCCTGGAGGAA  
GGGGCCAGGCCTCACATTCTGTGGGGCTCCCTGAATGGCAGCCTGAGCACAGCGTAGGCCCTT  
AATAAACACCTGTTGGATAAGCCAAAAAAA

**FIGURE 9**

MTHRTTTWARRTSRAVTPTCATPAGEMPCSRLLPPLRCSLHSACCSGDPASYRLWGAPLQPT  
LGVVPQASVPLLTDLAQWEPVLVPEAHPNASLTMYVCTFVPHDPDMALSRTPTTRQISSSDT  
DPPADGPSNPLCCCFFHGPAFSTLNPVLRHLFPQEAFPAHPIYDLSQVWSVVS PAPS R G Q A L R R A Q

**Signal peptide:**

amino acids 1-47

**N-glycosylation site.**

amino acids 31-35, 74-78, 84-88

**Casein kinase II phosphorylation site.**

amino acids 22-26, 76-80

**N-myristoylation site.**

amino acids 56-60

**Amidation site.**

amino acids 70-74

**FIGURE 10**

CCCACGCGTCCGAACCTCTCCAGCGATGGGAGCCGCCCCGCTGCTGCCCAACCTCACTCTGT  
GCTTACAGCTGCTGATTCTCTGCTGTCAAACCTCAGTACGTGAGGGACCAGGGCGCCATGACC  
GACCAGCTGAGCAGGCGGCAGATCCGCGAGTACCAACTCTACAGCAGGACCAGTGGCAAGCA  
CGTGCAGGTACCCGGGCGTCGCATCTCCGCCACCGCCGAGGACGGCAACAAGTTTGCCAAGC  
TCATAGTGGAGACGGACACGTTTGGCAGCCGGGTTTCGCATCAAAGGGGCTGAGAGTGAGAAG  
TACATCTGTATGAACAAGAGGGGCAAGCTCATCGGGAAGCCCAGCGGGAAGAGCAAAGACTG  
CGTGTTCACGGAGATCGTGTGGAGAACTATAACGGCCTTCCAGAACGCCCGGCACGAGG  
GCTGGTTCATGGCCTTACGCGGCAGGGGCGGCCCGCCAGGCTTCCCGCAGCCGCCAGAAC  
CAGCGCGAGGCCCACCTTCATCAAGCGCCTCTACCAAGGCCAGCTGCCCTTCCCCAACCCACGC  
CGAGAAGCAGAAGCAGTTCGAGTTTGTGGGCTCCGCCCCCACCCGCCGACCAAGCGCACAC  
GGCGGCCCCAGCCCTCACGTAGTCTGGGAGGCAGGGGGCAGCAGCCCCTGGGCCGCTCCC  
CACCCCTTTCCCTTCTTAATCCAAGGACTGGGCTGGGGTGGCGGGAGGGGAGCCAGATCCCC  
GAGGGAGGACCCTGAGGGCCGCGAAGCATCCGAGCCCCCAGCTGGGAAGGGGCAGGCCGGTG  
CCCAGGGGCGGCTGGCACAGTGCCCCCTTCCCGGACGGGTGGCAGGCCCTGGAGAGGAACT  
GAGTGTACCCCTGATCTCAGGCCACCAGCCTCTGCCGGCCTCCAGCCGGGCTCCTGAAGCC  
CGCTGAAAGGTCAGCGACTGAAGGCCTTGCAGACAACCGTCTGGAGGTGGCTGTCTCAAAA  
TCTGCTTCTCGGATCTCCCTCAGTCTGCCCCAGCCCCAAACTCCTCCTGGCTAGACTGTA  
GGAAGGGACTTTTGTFTTGTFTTGTFTTTCAGGAAAAAAGAAAGGGAGAGAGAGGAAAAATAG  
AGGGTTGTCCACTCCTCACATTCCACGACCCAGGCCTGCACCCCCACCCCAACTCCCAGCCC  
CGGAATAAAACCATTTTCCTGC

**FIGURE 11**

MGAARLLPNLTLCLQLLILCCQTQYVRDQAMTDQLSRRQIREYQLYSRTSGKHVQVTGRI  
SATAEDGNKFAKLIVETDFTFGSRVRIKGAESEKYICMNRGKLGKPSGKSKDCVFTEIVLE  
NNYTAFQONARHEGWFMATRQGRPRQASRSRQNRQREAHFIKRLYQGQLPFPNHAEKQKQFEF  
VGSAPTRRTKRTRRPQPLT

**Signal peptide:**

amino acids 1-22

**N-glycosylation site.**

amino acids 9-13, 126-130

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 60-64

**Casein kinase II phosphorylation site.**

amino acids 65-69

**Tyrosine kinase phosphorylation site.**

amino acids 39-48, 89-97

**N-myristoylation site.**

amino acids 69-75, 188-194

**Amidation site.**

amino acids 58-62

**HBGF/FGF family signature.**

amino acids 103-128

**FIGURE 12**

ACTTGCCATCACCTGTTGCCAGTGTGGAAAAATTCTCCCTGTTGAATTTTTTGCACATGGAG  
GACAGCAGCAAAGAGGGCAACACAGGCTGATAAGACCAGAGACAGCAGGGAGATTATTTTAC  
CATACGCCCTCAGGACGTTCCCTCTAGCTGGAGTTCTGGACTTCAACAGAACCCCATCCAGT  
CATTTTGATTTTGCTGTTTATTTTTTTTTTCTTTTTCTTTTTCCACCACATTGTATTTTAT  
TTCCGTACTTCAGAAATGGGCCTACAGACCACAAAGTGGCCCAGCCATGGGGCTTTTTTCT  
GAAGTCTTGGCTTATCATTCCCTGGGGCTCTACTCACAGGTGTCCAAACTCCTGGCCTGCC  
CTAGTGTGTGCCGCTGCGACAGGAACTTGTCTACTGTAATGAGCGAAGCTTGACCTCAGTG  
CCTCTTGGGATCCCGAGGGCGTAACCGTACTCTACCTCCACAACAACCAAATTAATAATGC  
TGGATTTCTGCAGAACTGCACAATGTACAGTCGGTGCACACGGTCTACCTGTATGGCAACC  
AACTGGACGAATCCCCATGAACCTTCCCAAGAAATGTCAGAGTCTCCATTTGCAGGAAAAC  
AATATTCAGACCATTTACGGGCTGCTCTTGCCAGCTCTTGAAGCTTGAAGAGCTGCACCT  
GGATGACAACCTCCATATCCACAGTGGGGGTGGAAGACGGGGCCTTCCGGGAGGCTATTAGCC  
TCAAATGTTGTTTTTGTCTAAGAATCACCTGAGCAGTGTGCCTGTTGGGCTTCTGTGGAC  
TTGCAAGAGCTGAGAGTGGATGAAAATCGAATGCTGTATATCCGACATGGCCTTCCAGAA  
TCTCACGAGCTTGGAGCGTCTTATTGTGGACGGGAACCTCCTGACCAACAAGGGTATCGCCG  
AGGGCACCTTCAGCCATCTCACCAAGCTCAAGGAATTTTCAATTGTACGTAATTCGCTGTCC  
CACCTCCTCCCGATCTCCAGGTACGCATCTGATCAGGCTCTATTTGCAGGACAACCAGAT  
AAACCACATTCCTTTGACAGCCTTCTCAAATCTGCGTAAGCTGGAACGGCTGGATATATCCA  
ACAACAACCTCGGGATGCTGACTCAAGGGGTTTTTGATAATCTCTCCAACCTGAAGCAGCTC  
ACTGCTCGGAATAACCCTTGGTTTTGTGACTGCAGTATTAAATGGGTACAGAAATGGCTCAA  
ATATATCCCTTCACTCTCAACGTGCGGGGTTTCATGTGCCAAGGTCTGAACAAGTCCGGG  
GGATGGCCGTGAGGGAATTAATATGAATCTTTTGTCTGTCCACCACGACCCCGGCCTG  
CCTCTCTTACCCCAGCCCCAAGTACAGCTTCTCCGACCACTCAGCCTCCCACCCTCTCTAT  
TCCAAACCCTAGCAGAAGCTACACGCCTCCAACTCCTACCACATCGAAACTTCCCACGATTC  
CTGACTGGGATGGCAGAGAAAGAGTGACCCACCTATTTCTGAACGGATCCAGCTCTCTATC  
CATTTTGTGAATGATACTTCCATTC AAGTCAGCTGGCTCTCTCTTCCACCGTGATGGCATA  
CAAACCTCACATGGGTGAAAATGGGCCACAGTTTAGTAGGGGCATCGTTCAGGAGCGCATAG  
TCAGCGGTGAGAAGCAACACCTGAGCCTGGTTAACTTAGAGCCCCGATCCACCTATCGGATT  
TGTTTAGTGCCACTGGATGCTTTTAACTACCGCGCGGTAGAAGACACCATTTGTTGAGAGC  
CACCACCCATGCCTCCTATCTGAACAACGGCAGCAACACAGCGTCCAGCCATGAGCAGACGA  
CGTCCCACAGCATGGGCTCCCCCTTTCTGCTGGCGGGCTTGATCGGGGGCGCGGTGATATTT  
GTGCTGGTGGTCTTGCTCAGCGTCTTTTGTCTGGCATATGCACAAAAAGGGGCGCTACACCTC  
CCAGAAGTGGAAATAACAACGGGGCCGGCGGAAAGATGATTATTGCGAGGCAGGCACCAAGA  
AGGACAACCTCCATCCTGGAGATGACAGAAACCAGTTTTTCTAGATCGTCTCCTTAAATAACGAT  
CAACTCCTTAAAGGAGATTTGAGACTGCAGCCATTTACACCCCAAATGGGGGCATTAATTA  
CACAGACTGCCATATCCCCAACAAATGCGATACTGCAACAGCAGCGTGCCAGACCTGGAGC  
ACTGCCATACGTTGACAGCCAGAGGCCAGCGTTATCAAGGCGGACAATTAGACTCTTGAGAA  
CACACTCGTGTGTGCACATAAAGACACGCAGATTACATTTGATAAATGTTACACAGATGCAT  
TTGTGCATTTGAATACTCTGTAATTTATACGGTGTACTATATAATGGGATTTAAAAAAGTG  
CTATCTTTTCTATTTCAAGTTAATTACAAACAGTTTTGTAACTCTTGTCTTTTAAATCTT

**FIGURE 13**

MGLQTTKWPSHGAFFLKSWLIISLGLYSQVSKLLACPSVCRCDRNFVYCNERSLTSVPLGIP  
EGVTVLYLHNNQINNAGFPAELHNVQSVHTVYLYGNQLDEFPMNLPKNVRVLHLQENNIQTI  
SRAALAQLLKLLEELHLDNSISTVGVEDGAFREAISLKLFLSKNHLSSVPVGLPVDLQELR  
VDENRIAVISDMAFQNLTSLERLIVDGNLLTNKGIAEGTFSHLTKLKEFSIVRNSLSHPPPD  
LPGTHLIRLYLQDNQINHIPLTAFSNLRKLERLDISNNQLRMLTQGVFDNLSNLKQLTARNN  
PWFCDCSIKWVTEWLKYIPSSLNVRGFMCGPEQVRGMAVRELNMNLLSCPTTTPGLPLFTP  
APSTASPTTQPPTLSIPNPSRSYTPPTPTTTSKLPTIPDWDGRERVTTPPISERIQLSIHFVND  
TSIQVSWLSLFTVMAYKLTWVKMGHSLVGGIVQERIVSGEKQHLSLVNLEPRSTYRICLVPL  
DAFNRYAVEDTICSEATTHASYLNNGSNTASSHEQTTSHSMGSPFLLAGLIGGAVIFVLVVL  
LSVFCWHMHKGRYTSQKWYNRGRRKDDYCEAGTKKDNSILEMTETSQIVSLNNDQLLKG  
DFRLQPIYTPNGGINYTDCHIPNNMRYCNSVDPLEHCHT

**Signal peptide:**

amino acids 1-42

**Transmembrane domain:**

amino acids 542-561

**N-glycosylation site.**

amino acids 202-206, 298-302, 433-437, 521-525, 635-639, 649-653

**Casein kinase II phosphorylation site.**

amino acids 204-208, 407-411, 527-531, 593-597, 598-602, 651-655

**Tyrosine kinase phosphorylation site.**

amino acids 319-328

**N-myristoylation site.**

amino acids 2-8, 60-66, 149-155, 213-219, 220-226, 294-300,  
522-528, 545-551, 633-639

**Amidation site.**

amino acids 581-585

**Leucine zipper pattern.**

amino acids 164-186

**Phospholipase A2 aspartic acid active site.**

amino acids 39-50

**FIGURE 14**

ACTTGGAGCAAGCGGCGGCGGAGACAGAGGCAGAGGCAGAAGCTGGGGCTCCGTCCCTCGCCTCCCACGAGCG  
ATCCCCGAGGAGAGCCGCGGCCCTCGGCGAGGCGAAGAGGCCGACGAGGAAGACCCGGGTGGCTGCGCCCTGCC  
TCGCTTCCAGGCGCCGCGGCTGCAGCCTTGCCCTCTTGCTCGCCTTGAAAATGGAAAAGATGCTCGCAGGCT  
GCTTTTGCTGATCCTCGGACAGATCGTCTCCTCCCTGCCGAGGCCAGGAGCGGTACAGTGGGAGGTCATCT  
CTAGGGGCGACACGCTCGGACCCACCCGCGAGACGGCCCTTCTGGAGAGTTCCTGTGAGAACAGCGGGCAGACC  
TGGTTTTTCATCATTGACAGCTCTCGCAGTGTCAACAACCCATGACTATGCAAAGGTCAAGGAGTTCATCGTGGACA  
TCTTGCAATTCTTGGACATTGGTCTGATGTACCCGAGTGGCCCTGCTCCAATATGGCAGCACTGTCAAGAATG  
AGTTCTCCCTCAAGACCTTCAAGAGGAAGTCCGAGGTGGAGCGTGTGTCAAGAGGATGGCGCATCTGTCCACGG  
GCACCATGACTGGGCTGGCCATCCAGTATGCCCTGAACATCGCATTCTCAGAAGCAGAGGGGGCCCGGCCCTGA  
GGGAGAATGTGCCACGGGTATAATGATCGTGCAGATGGGAGACCTCAGGACTCCGTGGCCGAGGTGGTGTCTA  
AGGCACGGGACACGGGCATCTAATCTTTGCCATTGGTGTGGGCCAGGTAGACTTCAACACCTTGAAGTCCATTG  
GGAGTGAGCCCCATGAGGACCATGTCTTCTGTGGCCAATTTACGCCAGATTGAGACGCTGACCTCCGTGTTC  
AGAAGAAATTGTGCACGGCCACATGTGCAGCACCTGGAGCATAACTGTGCCACTTCTGCATCAACATCCCTG  
GCTCATACTGTGCAGGTGCAAACAAGGCTACATTCTCAACTCGGATCAGACGACTTGCAGAATCCAGGATCTGT  
GTGCCATGGAGGACCACAATGTGAGCAGTCTGTGTGAATGTGCCGGGCTCCTTCTGTGCCAGTGTCTACAGT  
GCTACGCCCTGGCTGAGGATGGGAAGAGGTGTGTGGCTGTGGACTACTGTGCCTCAGAAAACCCGAGTGTGAAC  
ATGAGTGTGTAATGCTGATGGCTCCTACCTTTGCCAGTGCATGAAGGATTTGCTCTTAACCCAGATGAAAAA  
CGTGCACAAGGATCAACTACTGTGCACTGAACAAACCGGGCTGTGAGCATGAGTGCCTCAACATGGAGGAGAGCT  
ACTACTGCCGCTGCCACCGTGGCTACACTTGGACCCCAATGGCAAAACCTGCAGCCGAGTGGAGCACTGTGCAC  
AGCAGGACCATGGCTGTGAGCAGCTGTGTGAACACGGAGGATTCCTTCTGTGAGGAGGATTCCTTCTGTGAGG  
TCATCAACGAGGACCTCAAGACCTGCTCCCGGTTGGATTACTGCCTGTGAGTGAACATGGTTGTGAATFACTCT  
GTGTCAACATGGACAGATCTTTGCTGTGAGTGTCTGAGGGACACGTCTCCGAGCGATGGGAAGACGTGTG  
CAAAATGGACTCTTGTGCTCTGGGGGACACGGTTGTGAACATTCGTGTGTAAGCAGTGAAGATTCGTTTGTGT  
GCCAGTCTTTGAAGGTTATATACTCCGTGAAGATGGAAAAACCTGCAGAAGGAAAGATGTCTGCCAAGCTATAG  
ACCATGGCTGTGAACACATTTGTGTGAACAGTGCAGACTCATAACCGTGCAGTGTCTGGAGGGATTCGGCTCG  
CTGAGGATGGGAAACGCTGCCGAAGGAAGGATGTCTGCAAATCAACCCATGGCTGCCAACACATTTGTGTTA  
ATAATGGGAATTCCTACATCTGCAAATGCTCAGAGGGATTTGTTTAGCTGAGGACGGAAGACGGTGCAAGAAAT  
GCACTGAAGGCCAATGACCTGGTCTTTGTGATCGATGGATCCAAGAGTCTTGGAGAAGAGAATTTTGAGGTCG  
TGAAGCAGTTTGTCACTGGAATATAGATTCCTTGACAAATTCGCCCAAAGCCGCTCGAGTGGGGCTGTCCAGT  
ATTCACACAGGTCACACAGAGTTCACCTGAGAAAACCTCAACTCAGCCAAAGACATGAAAAAGCCGTGGCCC  
ACATGAAATACATGGAAAAGGGCTCTATGACTGGGCTGGCCCTGAAACACATGTTGAGAGAAGTTTTACCCAAAG  
GAGAAGGGCCAGGCCCTTTCCACAAGGTTGCCAGAGCAGCATTGTGTTCAACCGACGGACGGCTCAGGATG  
ACGTCTCCGAGTGGGCCAGTAAAGCCAAGGCCAATGGTATCACTATGTATGCTGTGGGGTAGGAAAAGCCATTG  
AGGAGGAACTACAAGAGATTGCCTCTGAGCCCAAAACAAGCATCTCTTCTATGCCGAAGACTTCAGCACAATGG  
ATGAGATAAGTGA AAAACTCAAGAAAGGCATCTGTGAAGCTCTAGAAGACTCCGATGGAAGACAGGACTCTCCAG  
CAGGGAACTGCCAAAAACGGTCCAACAGCCAACAGAATCTGAGCCAGTACCATAAATATCCAAGACCTACTTT  
CCTGTCTAATTTTGCAGTGCACAACAGATATCTGTTGAAGAAGACAATCTTTTACGGTCTACACAAAAGCTTT  
CCCATTCAACAAAACCTTCAGGAAGCCCTTTGGAAGAAAAACCGATCAATGCAAAATGTGAAAACCTTATAATGT  
TCCAGAACCTTGCAAACGAAGAAGTAAGAAAATTAACACAGCGCTTAGAAGAAAATGACACAGAGAATGGAAGCCC  
TGGAAAATCGCCTGAGATACAGATGAGAGATTAGAATCGCGACACATTTGTAGTCATTGTATCACGGATTACAAT  
GAACGAGTGCAGAGCCCCAAAGCTCAGGCTATTGTTAAATCAATAATGTTGTGAAGTAAAAAATCAGTACTGA  
GAAACCTGGTTGCACAGAACAAAGACAAGAAGTATACACTAATCTGTATAAATTTATCTAGGAAAAAATCCT  
TCAGAATTCTAAGATGAATTTACCAGGTGAGAAATGAATAAGCTATGCAAGGATTTTGTAAATATACTGTGGACAC  
AACTTGTCTTGCCTCATCTGCCTTAGTGTGCAATCTCAATTTGACTATACGATAAAGTTTGCACAGTCTTACTT  
CTGTAGAACACTGGCCATAGGAAATGCTGTTTTTTGTACTGGACTTTACCTTGATATATGATATGGATGTATG  
CATAAAATCATAGACATAATGACTTTGTGGAACAAGTTGGATTTTTTATACAAATATAAATTCACCCTTCAG



## **FIGURE 15**

MEKMLAGCFLLILGQIVLLPAEARERSRGRSISRGRHARTHPTALLESSCENKRADLVFII  
DSSRSVNTHDYAKVKEFIVDILQFLDIGPDVTRVGLLQYGSTVKNEFSLKTFKRRSEVERAV  
KRMRHLSTGMTGLAIQYALNIAFSEAEGARPLRENVPRVIMIVTDGRPODSVAEVAAKARD  
TGILIFAIGVGQVDFNTLKSIGSEPHEDHVFLVANFSQIETLTSVVFQKKLCTAHMCSTLEHN  
CAHFCINIPGSYVCRCKQGYILNSDQTTCRIQDLCAMEDHNCEQLCVNVPGSFVCQCYSGYA  
LAEDGKRCVAVDYCASENHGCEHECVNADGSYLCQHEGFALNPDEKTCRINYCALKNPGC  
EHECVNMEESYCRCHRGYTLDPNGKTC SRVDHCAQQDHGCEQLCLNTEDSFVCQCSEGFLLI  
NEDLKTCSRVDYCLLSDHGCEYSCVNMDRSFACQCPGHEVLRSDGKTC AKLDSCALGDHGCE  
HSCVSSSEDSFVCQCFEGYILREDGKTCRRKDVQCQAI DHGCEHICVN SDDSYTCECLEGFRLA  
EDGKRCRRKDVCKSTHHGCEHICVNNGNSYICKCSEGFVLAEDGRRCKKCTEGPIDLVFVID  
GSKSLGEENFEVVKQFVTGIIDSLTISP KAA RVGLLQYSTQVHTEFTLRNFNSAKDMKKAVA  
HMKYMGKGSMTGLALKHMFERSFTQGE GARPLSTRVPRAAIVFTDGRAQDDVSEWASKAKAN  
GITMYAVGVGKAIEEELQEIASEPTNKHLFYAEDFSTMDEISEKLKKGICEALESDGRQDS  
PAGELPKTVQQPTSESEPTVINIQDLLSCSNFAVQHRYLFEEDNLLRSTQKLSHSTKPSGSPL  
EEKHDQCKCENLIMFQNLANEVVRKLTQRLEEMTQRMEALENRLRYR

**Signal peptide:**

amino acids 1-23

**N-glycosylation site.**

amino acids 221-225

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 115-119, 606-610, 892-896

**Casein kinase II phosphorylation site.**

amino acids 49-53, 118-122, 149-153, 176-180, 223-227, 243-247,  
401-405, 442-446, 501-505, 624-628, 673-677, 706-710, 780-784,  
781-785, 819-823, 866-870

**N-myristoylation site.**

amino acids 133-139, 258-264, 299-305, 340-346, 453-459, 494-500,  
639-645, 690-696, 752-758, 792-798

**Amidation site.**

amino acids 314-318, 560-564, 601-605

**Aspartic acid and asparagine hydroxylation site.**

amino acids 253-265, 294-306, 335-347, 376-388, 417-423, 458-464,  
540-546, 581-587

**FIGURE 16**

GGAGCCGCCCTGGGTGTCAGCGGCTCGGCTCCCGCGCACGCTCCGGCCGTCGCGCAGCCTCG  
GCACCTGCAGGTCCGTGCGTCCCGCGGCTGGCGCCCCTGACTCCGTCCCGGCCAGGGAGGGC  
CATGATTTCCCTCCCGGGGCCCTGGTGACCAACTTGCTGCGGTTTTTGTTCCTGGGGCTGA  
GTGCCCTCGCGCCCCCTCGCGGGCCAGCTGCAACTGCACTTGCCCCCAACCGGTTGCAG  
GCGGTGGAGGGAGGGGAAGTGGTGCTTCCAGCGTGGTACACCTTGCACGGGGAGGTGTCTTC  
ATCCCAGCCATGGGAGGTGCCCTTTGTGATGTGGTTCTTCAAACAGAAAGAAAAGGAGGATC  
AGGTGTTGTCCCTACATCAATGGGTGTCACAACAAGCAAACCTGGAGTATCCTTGGTCTACTCC  
ATGCCCTCCCGAACCTGTCCCTGCGGCTGGAGGGTCTCCAGGAGAAAGACTCTGGCCCCTA  
CAGCTGCTCCGTGAATGTGCAAGACAAAACAAGGCAAATCTAGGGGCCACAGCATCAAAACCT  
TAGAACTCAATGTACTGGTTCCTCCAGCTCCTCCATCCTGCCGTCTCCAGGGTGTGCCCCAT  
GTGGGGGCAAACGTGACCCTGAGCTGCCAGTCTCCAAGGAGTAAGCCCGCTGTCCAATACCA  
GTGGGATCGGCAGCTTCCATCCTTCCAGACTTCTTTGCACCAGCATTAGATGTCATCCGTG  
GGTCTTTAAGCCTCACCAACCTTTCGTCTTCCATGGCTGGAGTCTATGTCTGCAAGGCCAC  
AATGAGGTGGCACTGCCAATGTAATGTGACGCTGGAAGTGAACACAGGGCCCTGGAGCTGC  
AGTGGTTGCTGGAGCTGTTGTGGGTACCCTGGTTGGACTGGGGTTGCTGGCTGGGCTGGTCC  
TCTTGTACCACCGCCGGGGCAAGGCCCTGGAGGAGCCAGCCAATGATATCAAGGAGGATGCC  
ATTGCTCCCGGACCCTGCCCTGGCCCAAGAGCTCAGACACAATCTCCAAGAATGGGACCCT  
TTCCTCTGTACCTCCGCACGAGCCCTCCGGCCACCCCATGGCCCTCCAGGCCTGGTGCAT  
TGACCCCCACGCCAGTCTCTCCAGCCAGGCCCTGCCCTCACCAAGACTGCCCACGACAGAT  
GGGGCCACCCTCAACCAATATCCCCATCCCTGGTGGGGTTTCTTCCCTCTGGCTTGAGCCG  
CATGGGTGCTGTGCCTGTGATGGTGCTGCCAGAGTCAAGCTGGCTCTCTGGTATTGATGAC  
CCCACCACTCATTGGCTAAAGGATTTGGGGTCTCTCCTTCTATAAGGGTACCTCTAGCAC  
AGAGGCCTGAGTCATGGGAAAGAGTCACTCTGACCCTTAGTACTCTGCCCCACCTCTC  
TTTACTGTGGAAAACCATCTCAGTAAGACCTAAGTGTCCAGGAGACAGAAGGAGAAGAGGA  
AGTGGATCTGGAATTGGGAGGAGCCTCCACCCACCCCTGACTCCTCCTTATGAAGCCAGCTG  
CTGAAATTAGCTACTACCAAGAGTGAGGGGCAGAGACTTCCAGTCACTGAGTCTCCAGGC  
CCCCTTGATCTGTACCCACCCCTATCTAACACCACCCTTGGCTCCCACTCCAGCTCCCTGT  
ATTGATATAACCTGTCAGGCTGGCTTGGTTAGGTTTTACTGGGGCAGAGGATAGGGAATCTC  
TTATTAAAACCTAACATGAAATATGTGTTGTTTTTCAATTTGCAAATTTAAATAAAGATACATAA  
TGTTTGTATGAAAAA

**FIGURE 17**

MISLPGPLVTNLLRFLFLGLSALAPPSRAQLQLHLPANRLQAVEGGEVVLPAWYTLHGCVSS  
SQPWEVFPVMWFFKQKEKEDQVLSYINGVTTSTKPGVSLVYSMPNRNLSLRLEGLQEKDSGPY  
SCSVNVQDKQGKSRGHSIKTLELNVLVPPAPPSCRLQGVPHVGANVTLSQSPRSKPAVQYQ  
WDRQLPSFQTFPAPALDVIRGSLSLTNLSSSMAGVYVCKAHNEVGTAQCNTLEVSTGPGAA  
VVAGAVVGTTLVGLGLLAGLVLLYHRRGKALEEPANDIKEDAIAPRTLTPWPKSSDTISKNGTL  
SSVTSARALRPPHGPPRPGALTPTPSLSSQALPSPRLPTTDGAHPQPIISPPIGGVSSSSGLSR  
MGAVPVMVPAQSQAGSLV

**Signal peptide:**

amino acids 1-29

**Transmembrane domain:**

amino acids 245-267

**N-glycosylation site.**

amino acids 108-112, 169-173, 213-217, 236-240, 307-311

**N-myristoylation site.**

amino acids 90-96, 167-173, 220-226, 231-237, 252-258, 256-262,  
262-268, 308-314, 363-369, 364-370

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 164-175

**FIGURE 18**

CGCCACCACTGCGGCCACCGCCAATGAAACGCCTCCCGCTCCTAGTGGTTTTTTTCCACTTTG  
TTGAATTGTTCTTACTATACTCAAATTCACCAAGACACCTTGTCTCCCAAATGCAAATGTGA  
AATACGCAATGGAATTGAAGCCTGCTATTGCAACATGGGATTTTCAGGAAATGGTGTACAA  
TTTGTGAAGATGATAATGAATGTGAAATTTAACTCAGTCCTGTGGCGAAAAATGCTAATTGC  
ACTAACACAGAAAGTATTATTGTATGTGTACCTGGCTTCAGATCCAGCAGTAACCA  
AGACAGGTTTATCACTAATGATGGAACCGTCTGTATAGAAAATGTGAATGCAAACCTGCCATT  
TAGATAATGTCTGTATAGCTGCAAATATTAATAAACTTTAAACAAAAATCAGATCCATAAAA  
GAACCTGTGGCTTTGCTACAAGAAGTCTATAGAAATCTGTGACAGATCTTTCACCAACAGA  
TATAATTACATATATAGAAATATTAGCTGAATCATCTTCATTACTAGGTTACAGAACAACA  
CTATCTCAGCCAAGGACACCCCTTCTAACTCAACTCTTACTGAATTTGTA AAAAACCCTGAAT  
AATTTTGTCAAAGGGATACATTTGTAGTTTGGGACAAGTTATCTGTGAATCATAGGAGAAC  
ACATCTTACAAAACCTCATGCACACTGTTGAACAAGTACTTTAAGGATATCCAGAGCTTCC  
AAAAGACCACAGAGTTTGATACAAATTCACCGGATATAGCTCTCAAAGTTTTCTTTTTGAT  
TCATATAACATGAAACATATTCATCCTCATATGAATATGGATGGAGACTACATAAATATATT  
TCCAAAGAGAAAAGCTGCATATGATTCAAATGGCAATGTTGCAGTTGCATTTTTATATTATA  
AGAGTATTGGTCTTTGCTTTCATCATCTGACAACCTTCTTATTGAAACCTCAAATTTATGAT  
AATCTGAAGAGGAGGAAAGAGTCAATCTTCAGTAATTTTCAGTCTCAATGAGCTCAAACCC  
ACCCACATTATATGAACCTTGAAAAATAACATTTACATTAAGTCATCGAAAGGTCACAGATA  
GGTATAGGAGTCTATGTGCATTTTGGAACTTACCTGATAACCATGAATGGCAGCTGGTCT  
TCAGAGGGCTGTGAGCTGACATACTCAAATGAGACCCACACCTCATGCCGCTGTAATCACCT  
GACACATTTGCAATTTTGTATGTCTCTGGTCTTCCATTGGTATTAAGATTATAATATTC  
TTACAAGGATCACTCAACTAGGAATAATTAATTCAGTATTTGTCTTGCCATATGCATTTTT  
ACCTCTGGTCTTTCAGTGAATTCAAAGCACCAGGACAACAATTCACAAAAATCTTTGCTG  
TAGCTAATTTCTGTGAACCTGTTTTTCTGTGGGATCAATACAAATACTAAATAAGCTCT  
TCTGTTCAATCATTGCGGACTGCTACACTACTTCTTTTTAGCTGCTTTTGCATGGATGTGC  
ATTGAAGGCATACATCTCTATCTCATTGTTGTGGTGTCTATCTACAACAAGGGATTTTTGCA  
CAAGAATTTTATATCTTTGGCTATCTAAGCCAGCCGTGGTAGTTGGATTTTCGGCAGCAC  
TAGGATACAGATATTATGGCACAACCAAAGTATGTTGGCTTAGCACCGAAAACAACCTTATT  
TGGAGTTTTATAGGACCAGCATGCCTAATCATTCTTGTAAATCTCTTGGCTTTTGGAGTCAT  
CATATACAAAAGTTTTTCGTCACACTGCAGGGTTGAAACCAGAAGTTAGTTGCTTTGAGAAC  
TAAGGTCTTGTGCAAGAGGAGCCCTCGCTCTTCTGTTCTCTCGGCACCACCTGGATCTTT  
GGGTTCTCCATGTTGTGCACGCATCAGTGGTTACAGCTTACCTCTTCACAGTCAGCAATGC  
TTTTCCAGGGATGTTTCATTTTTTTTATTCTGTGTGTTTTATCTAGAAAGATTCAAGAAGAT  
ATTACAGATTGTTCAA AAAATGTCCCTGTTGTTTGGATGTTTAAGGTTAAACATAGAGAATG  
GTGGATAATTACAACCTGCACAAAATAAAAAATTC AAGCTGTGGATGACCAATGTATAAAAA  
TGACTCATCAAATATCCAATTATTAACCTAGACAAAAAGTATTTTAAATCAGTTTTTCT  
GTTTATGTATAGGAACTGTAGATAATAAGGTA AAAATATGTATCATATAGATATACTATGT  
TTTTCTATGTGAAATAGTTCTGTCAA AAATAGTATTGCAGATATTTGAAAGTAATGGT  
CTCAGGAGTGATATCACTGCACCCAAGGAAAGATTTTTCTTCTAACACGAGAAGTATATGAA  
TGTCCTGAAGGAAACCACTGGCTTGATTTCTGTGACTCGTGTGCTTTGAAACTAGTCC  
CCTACCCTCGGTAATGAGCTCCATTACAGAAAGTGAACATAAGAGAATGAAGGGCAGA  
ATATCAAACAGTGAAAAGGGAATGATAAGATGTATTTGAAATGAACTGTTTTTTCTGTAGAC  
TAGCTGAGAAATGTTGACATAAAAATAAAGAAATGAAGAAACACATTTTACCATTTTGTGAA  
TTGTTCTGAACTTAAATGTCCACTAAAACAACCTTAGACTTCTGTTTGCTAAATCTGTTCTT  
TTTCTAATATTTCAAAAAAAAAAAAAAAAAAGGTTTACCTCCACAAATTTGAAAAAAAAAAAAAAAA  
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

**FIGURE 19**

MKRLPLLVVFSTLLNCSYTNCTKTPCLPNAKCEIRNGIEACYCNMGFSGNGVTICEDDNEC  
GNLTQSCGENANCTNTEGSYYCMVPGFRSSSNQDRFITNDGTVCIEVNVNANCHLDNVCIAA  
NINKTLTKIRSIKEPVALLQEVYRNSVTDLSPTDIITYIEILAESSLLGYKNNTISAKDTL  
SNSTLTFVKTIVNNFVQRDTFVVDKLSVNHRRTHLTKLMHTVEQATLRISQSFQKTTEFDT  
NSTDIALKVVFFDSYNMKHIHPHMNDGDYINIFPKRKAAYDSNGNVAVAFLYYKSIGPLLS  
SSDNFLLKPQNYDNSEEEERVISSVISVSMSSNPPTLYELEKITFTLSHRKVTDRYRSLCAF  
WNYSPTMNGSWSSEGCELTYSNETHTSRCRNHLTHFAILMSSGSPSIGIKDYNILTRITQLG  
IIISLICLAICIFTFWFFSEIQSTRTTIHKNLCCSLFLAELVFLVGINTNTNKLFCSTIAGL  
LHYFFLAFAWMCIEGIHLYLIVVGVYIYKNGFLHKNFYIFGYLSPAVVVGFSAALGYRYYGT  
TKVCWLSTENFIWFSFIGPACLIILVNLLAFGVIIYKVFRTAGLKPEVSCFENIRSCARGA  
LALLFLLGTTWIFGVLHVHASVVTAYLFTVSNAPQGMFIFLFLCVLSRKIQEEYYRLFKNV  
PCCFGCLR

**Signal peptide:**

amino acids 1-19

**Transmembrane domain:**

amino acids 430-450, 465-486, 499-513, 535-549, 573-593, 619-636,  
648-664

**N-glycosylation site.**

amino acids 15-19, 21-25, 64-68, 74-78, 127-131, 177-181,  
188-192, 249-253, 381-385, 395-399

**Glycosaminoglycan attachment site.**

amino acids 49-53

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 360-364

**Casein kinase II phosphorylation site.**

amino acids 54-58, 68-72, 76-80, 94-98, 135-139, 150-154,  
155-159, 161-165, 181-185, 190-194, 244-248, 310-314, 325-329,  
346-350, 608-612

**Tyrosine kinase phosphorylation site.**

amino acids 36-44, 669-677, 670-678

**N-myristoylation site.**

amino acids 38-44, 50-56, 52-58, 80-86, 382-388, 388-394,  
434-440, 480-486, 521-527

**Aspartic acid and asparagine hydroxylation site.**

amino acids 75-87

**FIGURE 20**

TGGAAACATATCCTCCCTCATATGAATATGGATGGAGACTACATAAATATATTTCCAAAGNG  
AAAAGCCGGCATATGGATTCAAATGGCAATGTTGCAGTTGCATTTTTATATTATAAGAGTAT  
TGGTCCCTTTGCTTTCATCATCTGACAACTTCTTATTGAAACCTCAAATTTATGATAATTCT  
GAAGAGGAGGAAAGAGTCATATCTTCAGTAATTTTCAGTCTCAATGAGCTCAAACCCACCCAC  
ATTATATGAACTTGAAAAATAACATTTACATTAAGTCATCGAAAGGTCACAGATAGGTATA  
GGAGTCTATGTGGCATTFTTGAATACTCACCTGATACCATGAATGGCAGCTGGTCTTCAGAG  
GGCTGTGAGCTGACATACTCAAATGAGACCCACACCTCATGCCGCTGTAATCACCTGACACA  
TTTTGCAATTTTGATGTCTCTGGTCTTCCATTGGTATTAAAGATTATAATATTCTTACAA  
GGATCACTCAACTAGGAATAATTATTTCACTGATTTGTCTTGCCATATGCATTTTTTACCTTC  
TGGTTCTTCAGTGAAATTCAAAGCACCAGGA

**FIGURE 21**

GCTCCCAGCCAAGAACCTCGGGGCCGCTGCGCGGTGGGGAGGAGTTCCCCGAAACCCGGCCG  
CTAAGCGAGGCCTCCTCCTCCCGCAGATCCGAACGGCCTGGGCGGGGTCACCCCGGCTGGGA  
CAAGAAGCCGCCGCTGCCTGCCCCGGGCCCGGGGAGGGGGCTGGGGCTGGGGCCGGAGGCCG  
GGTGTGAGTGGGTGTGTGCGGGGGCGGAGGCTTGATGCAATCCCGATAAGAAATGCTCGGG  
TGTCTTGGGCACCTACCCGTGGGGCCCGTAAGGCGCTACTATATAAGGCTGCCGGCCCCGGAG  
CCGCCGCGCCGTGAGAGCAGGAGCGCTGCGTCCAGGATCTAGGGCCACGACCATCCCAACCC  
GGCACTCACAGCCCCGAGCGCATCCCGGTGCGCGCCAGCCTCCCGCACCCCCATCGCCGG  
AGCTGCGCCGAGAGCCCCAGGGAGGTGCCATGCGGAGCGGGTGTGTGGTGGTCCACGTATGG  
ATCCTGGCCGGCCTCTGGCTGGCCGTGGCCGGGCGCCCCCTCGCCTTCTCGGACGCGGGCC  
CCACGTGCACTACGGCTGGGGCGACCCCATCCGCCTGCGGCACCTGTACACCTCCGGCCCC  
ACGGGCTCTCCAGCTGCTTCCCTGCGCATCCGTGCCGACGGCGTCTGGACTGCGCGCGGGGC  
CAGAGCGCGCACAGTTTGCTGGAGATCAAGGCAGTCGCTCTGCGGACCGTGGCCATCAAGGG  
CGTGACAGCGTGGCGTACCTCTGCATGGGCGCCGACGGCAAGATGCAGGGGCTGCTTCACT  
ACTCGGAGGAAGACTGTGCTTTCGAGGAGGAGATCCGCCAGATGGCTACAAATGTGTACCGA  
TCCGAGAAGCACCGCCTCCCGGTCTCCCTGAGCAGTGCCAAACAGCGGCAGCTGTACAAGAA  
CAGAGGCTTTCTTCCACTCTCTCATTTCCCTGCCCATGCTGCCCATGGTCCCAGAGGAGCCTG  
AGGACCTCAGGGGCCACTTGGAAATCTGACATGTTCTCTTCGCCCTGGAGACCGACAGCATG  
GACCCATTTGGGCTTGTACCCGGACTGGAGGCCGTGAGGAGTCCCAGCTTTGAGAAGTAACT  
GAGACCATGCCCCGGGCCTTTCACCTGCTGCCAGGGGCTGTGGTACCTGCAGCGTGGGGACG  
TGCTTCTACAAGAACAGTCCCTGAGTCCACGTTCTGTTTAGCTTTAGGAAGAAACATCTAGAA  
GTTGTACATATTAGAGTTTTCCATTGGCAGTGCCAGTTTTCTAGCCAATAGACTTGTCTGAT  
CATAACATTGTAAGCCTGTAGCTTGCCCAGCTGCTGCCTGGGCCCCCATCTGCTCCCTCGA  
GGTTGCTGGACAAGCTGCTGCACTGTCTCAGTTCTGCTTGAATACCTCCATCGATGGGGAAC  
TCACTTCCCTTTGGAAAAATTCTTATGTCAAGCTGAAATTCTCTAATTTTTTCTCATCACTTC  
CCCAGGAGCAGCCAGAAGACAGGCAGTAGTTTTAATTTTCAGGAACAGGTGATCCACTCTGTA  
AAACAGCAGGTAATAATTTCACTCAACCCCATGTGGGAATTGATCTATATCTCTACTTCCAGGG  
ACCATTTGCCCTTCCCAAATCCCTCCAGGCCAGAACTGACTGGAGCAGGCATGGCCACCAG  
GCTTCAGGAGTAGGGGAAGCCTGGAGCCCCACTCCAGCCCTGGGACAACCTGAGAATTCCCC  
CTGAGGCCAGTTCTGTGATGGATGCTGTCTGAGAATAACTTGCTGTCCCGGTGTACCTGC  
TTCCATCTCCAGCCCACCAGCCCTCTGCCACCTCACATGCCTCCCCATGGATTGGGGCCT  
CCCAGGCCCCCACCTTATGTCAACCTGCACTTCTTGTTCAAAAATCAGGAAAAGAAAAGAT  
TTGAAGACCCCAAGTCTTGTCAATAACTTGCTGTGTGGAAGCAGCGGGGAAGACCTAGAAC  
CCTTTCCCCAGCACTTGGTTTTTCAACATGATATTTATGAGTAATTTATTTTGATATGTACA  
TCTCTTATTTTCTTACATTATTTATGCCCCAAATTATATTTATGTATGTAAGTGAGGTTTG  
TTTTGTATATTAATAATGGAGTTTGTTTGT

**FIGURE 22**

MRS GCVVHVWILAGLWLAVAGRPLAFSDAGPHVHYGWGDP IRLRHL YTS GPHGLSSCF LRI  
RADGVVDCARGQSAHSLLEIKAV ALRTVAIKGVHSVRYLCMGADGKMQLLQYSEEDCAFEE  
EIRPDGYNVYRSEKHRLPVSLSSAKQRQLYKNRGFLPLSHFLPMLPMVPEEPEDLRGHLESD  
MFSSPLETDSMDPFGLVTGLEAVRSPSF EK

**Signal peptide:**

amino acids 1-22

**Casein kinase II phosphorylation site.**

amino acids 78-82, 116-120, 190-194, 204-208

**N-myristoylation site.**

amino acids 15-21, 54-60, 66-72, 201-207

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 48-59



**FIGURE 23**

CCCAGAAGTTCAAGGGCCCCCGGCCTCCTGCGCTCCTGCCGCCGGGACCCTCGACCTCCTCA  
GAGCAGCCGGCTGCCGCCCGGGAAGATGGCGAGGAGGAGCCGCCACCGCCTCCTCCTGCTG  
CTGCTGCGCTACCTGGTGGTTCGCCCTGGGCTATCATAAGGCCTATGGGTTTTCTGCCCCAAA  
AGACCAACAAGTAGTCACAGCAGTAGAGTACCAAGAGGCTATTTTAGCCTGCAAAAACCCCAA  
AGAAGACTGTTTTCTCCAGATTAGAGTGAAGAAACTGGGTCCGAGTGTCTCCTTTGTCTAC  
TATCAACAGACTCTTCAAGGTGATTTTAAAAATCGAGCTGAGATGATAGATTTCAATATCCG  
GATCAAAAATGTGACAAGAAGTGTGCGGGGAAATATCGTTGTGAAGTTAGTGCCCATCTG  
AGCAAGGCCAAAACCTGGAAGAGGATACAGTCACTCTGGAAGTATTAGTGCTCCAGCAGTT  
CCATCATGTGAAGTACCCTCTTCTGCTCTGAGTGGAACTGTGGTAGAGCTACGATGTCAAGA  
CAAAGAAGGGAAATCCAGCTCCTGAATACACATGGTTTAAGGATGGCATCCGTTTGCTAGAAA  
ATCCCAGACTTGGCTCCCAAAGCACCAACAGCTCATAACAATGAATACAAAAACTGGAAC  
CTGCAATTTAATACTGTTTCCAAACTGGACACTGGAGAATATTCTGTGAAGCCCGCAATTC  
TGTTGGATATCGCAGGTGTCCTGGGAAACGAATGCAAGTAGATGATCTCAACATAAGTGGCA  
TCATAGCAGCCGTAGTAGTTGTGGCCTTAGTGATTTCCGTTTGTGGCCTTGGTGTATGCTAT  
GCTCAGAGGAAAGGCTACTTTTCAAAGAAACCTCCTTCCAGAAGAGTAATTTCTTCATCTAA  
AGCCACGACAATGAGTGAATGTGCACTGGCTCACGCCGTAAATCCAGCACTTTGGAAGG  
CCGCGCGGGCGGATCACGAGGTCAGGAGTTCTAGACCAGTCTGGCCAATATGGTGAAACCC  
CATCTCTACTAAAATACAAAAATTAGCTGGGCATGGTGGCATGTGCCTGCAGTTCCAGCTGC  
TTGGGAGACAGGAGAATCACTTGAACCCGGGAGGCGGAGGTTGCAGTGAGCTGAGATCACGC  
CACTGCAGTCCAGCCTGGGTAACAGAGCAAGATTCCATCTCAAAAAATAAAATAAATA  
AATAAATACTGGTTTTTACCTGTAGAATTCTTACAATAAATATAGCTTGATATTC

## **FIGURE 24**

MARRSRHRLLLLLLRYLVVALGYHKAYGFSAPKDQQVVTAVEYQEAILACKTPKKTVSSRLE  
WKKLGRSVSVFVYYQOTLQGDFKNRAEMIDFNIRIKNVTRSDAGKYRCEVSAPSEQGQONLEED  
TVTLEVLVAPAVPSCCEVPSSALSGTVVELRCQDKEGNPAPEYTWFKDGI RLLLENPRLGSQST  
NSSYTMNTKTGTTLQFNTVSKLDTGEYSCEARN SVGYRRC PGKRMQVDDL NISGIIAAVVVVA  
LVISVCGLGVCYAQRKGYFSKETS FQKSNSSSKAT TMSENVQWLTPVIPALWCAAAGGSRGQEF

**Signal peptide:**

amino acids 1-20

**Transmembrane domain:**

amino acids 130-144, 238-258

**N-glycosylation site.**

amino acids 98-102, 187-191, 236-240, 277-281

**Casein kinase II phosphorylation site.**

amino acids 39-43, 59-63, 100-104, 149-153, 205-209, 284-288

**N-myristoylation site.**

amino acids 182-188, 239-245, 255-261, 257-263, 305-311

**Amidation site.**

amino acids 226-230

**FIGURE 25**

GACATCGGAGGTGGGCTAGCACTGAAACTGCTTTTCAAGACGAGGAAGAGGAGGAGAAAAG  
AAAGAAGAGGAAGATGTTGGGCAACATTTATTTAACATGCTCCACAGCCCGACCCTGGCAT  
CATGCTGCTATTTCCTGCAAATACTGAAGAAGCATGGGATTTAAATATTTTACTTCTAAATAA  
ATGAATTACTCAATCTCCTATGACCATCTATACTACTCCACCTTCAAAAAGTACATCAATA  
TTATATCATTAAGGAAATAGTAACCTTCTCTTCTCCAATATGCATGACATTTTGGACAATG  
CAATTGTGGCCTGGCCTTATTTCACTGAAGAAAACCTTTGTGGTTCTATGGCATTTCATCA  
TTTGACAAATGCAAGCATCTTCCTTATCAATCAGCTCCTATTGAACCTACTAGCACTGACTG  
TGGAATCCTTAAGGGCCCATTCATTTCTGAAGAAGAAAGCTAAGATGAAGGACATGCCACT  
CCGAATTCATGTGCTACTTGGCCTAGCTATCACTACACTAGTACAAGCTGTAGATAAAAAAG  
TGGATTGTCCACGGTTATGTACGTGTGAAATCAGGCCTTGGTTTACACCCAGATCCATTTAT  
ATGGAAGCATCTACAGTGGATTGTAATGATTTAGGTCTTTTAACTTTCCAGCCAGATTGCC  
AGCTAACACACAGATTTCTTCTCTACAGACTAAACAATATTGCAAAAATTGAATACTCCACAG  
ACTTTCCAGTAAACCTTACTGGCCTGGATTTATCTCAAAACAATTTATCTTCAGTCACCAAT  
ATTAATGTAAAAAGATGCCCTCAGCTCCTTTCTGTGTACCTAGAGGAAAACAACTTACTGA  
ACTGCCTGAAAAATGTCTGTCCGAACTGAGCAACTTACAAGAACTCTATATTAATCACAAC  
TGCTTTCTACAATTTACCTGGAGCCTTTATTGGCCTACATAATCTTCTTCGACTTCATCTC  
AATTCAAATAGATTGCAGATGATCAACAGTAAGTGGTTTGTAGTCTTCCAATCTAGAGAT  
TCTGATGATTGGGGAAAAATCCAATTTATCAGAATCAAAGACATGAACCTTAAAGCCTTTATCA  
ATCTTCGCAGCCTGGTTATAGCTGGTATAAACCTCACAGAAATACCAGATAACGCCTTGGTT  
GGACTGGAAAACCTTAGAAAGCATCTCTTTTTACGATAACAGGCCTTATTAAGTACCCCATGT  
TGCTCTTCAAAAAGTTGTAAATCTCAAAATTTTGGATCTAAATAAAAAATCCTATTAATAGAA  
TAGAAGGGGTGATTTTAGCAATATGCTACACTTAAAGAGTTGGGGATAAAATAATATGCCCT  
GAGCTGATTTCCATCGATAGTCTTGCTGTGGATAACCTGCCAGATTTAAGAAAAATAGAAGC  
TACTAACACCCCTAGATTGTCTTACATTCACCCCAATGCATTTTTCAGACTCCCCAAGCTGG  
AATCACTCATGCTGAACAGCAATGCTCTCAGTGCCTGTACCATGGTACCATTGAGTCTCTG  
CCAAACCTCAAGGAAATCAGCATAACAGTAACCCCATCAGGTGTGACTGTGTTCATCCGTTG  
GATGAACATGAACAAAACCAACATTGATTTCATGGAGCCAGATTCACGTGTTTTGCGTGGACC  
CACCTGAATTCAGGTGAGAAATGTTTCGGCAAGTGCATTTTCAAGGACATGATGGAAATTTGT  
CTCCCTCTTATAGCTCCTGAGAGCTTTTCTTCTAATCTAAATGTAGAAGCTGGGAGCTATGT  
TTCCTTTTCACTGTAGAGCTACTGCAGAACCACAGCCTGAAATCTACTGGATAACACCTTCTG  
GTCAAAAACCTTGCCTAATACCTTGACAGACAAGTTCTATGCTCCATTCAGGGAAACACTA  
GATATAAATGGCGTAACTCCCAAAGAAGGGGTTTATATACTTGTATAGCAACTAACCTAGT  
TGGCGCTGACTTGAAGTCTGTTATGATCAAAGTGGATGGATCTTTTCCACAAGATAACAATG  
GCTCTTTGAATATTAATAAGAGATATTCAAGCCAATTCAGTTTTGGTGTCTTGGAAAGCA  
AGTTCTAAAATCTCAAAATCTAGTGTTAAATGGACAGCCTTTGTCAAGACTGAAAATTTCTCA  
TGCTGCGCAAAGTGTCTCGAATACCATCTGATGTCAAGGTATATAATCTTACTCATCTGAATC  
CATCAACTGAGTATAAAATTTGTATTGATATTTCCACCATCTATCAGAAAAACAGAAAAAA  
TGTGTAATGTCAACCACAAAGGTTTGCACCTGATCAAAAAGAGTATGAAAAGAATAATAC  
CACAACTTATGGCCTGTCTTGGAGGCCTTCTGGGATTATTGGTGTGATATGTCTTATCA  
GCTGCCTCTCTCCAGAAATGAACTGTGATGGTGGACACAGCTATGTGAGGAATTACTTACAG  
AAACCAACCTTTGCATTAGGTGAGCTTTATCCTCCTCTGATAAATCTCTGGGAAGCAGGAAA  
AGAAAAAGTACATCACTGAAAGTAAAAGCAACTGTTATAGGTTTACCAACAAATATGTCTT  
AAAACCAACCAAGGAAACCTACTCCAAAATGAAC

## **FIGURE 26**

MKDMPLRIHVLLGLAITTLVQAVDKKVDPCRLCTCEIRPWFTPRSIYMEASTVDCNDLGLLT  
FPARLPANTQIILLQLTNNIAKIEYSTDFPVNLTGLDLSQNNLSSVTNINVKKMPQLLSVYLE  
ENKLTPELPEKCLSELNLQELYINHNLLSTISPGAFI GLHNLLRLHLNSNRLQMINSKWFDA  
LPNLEILMIGENPIIRIKDMNFKPLINLRSLVIAGINL TEIPDNALVLENLESI SFYDNRL  
IKVPHVALQKVVNLKFLDLNKNPINRIRRGDFSNMLHLKELGINNMPELISIDSLAVDNLDP  
LRKIEATNPNRSLYIHPNAFFRLPKLESIMLNSNALSALYHGHTIESLPNLKEISIHSPNIRC  
DCVIRWMMNKTNIRFMEPDSLFCVDPPEFQGNVRQVHFRDMMEICLPLIAPESFPSNLNV  
EAGSYVSFHCRTAEAPQPEIYWITPSGQKLLPNTLTDKIFYVHSEGLDINGVTPKEGGLYTC  
IATNLVGADLKSVMIKVDGSFPQDNNGSLNIKIRDIQANSVLVSWKASSKILKSSVKWTAFV  
KTENSHAAQSARIPSDVKVYNLTHLNPSTEYKICIDIPTIYQKNRKKCVNVTTKGLHPDQKE  
YEKNNTTTLMACLGGLGIIGVICLISCLSPENMCDGGHSYVRNYLQKPTFALGELYPLIN  
LWEAGKEKSTSLKVKATVIGLPTNMS

**Signal sequence:**

amino acids 1-22

**Transmembrane domain:**

amino acids 633-650

**N-glycosylation site.**

amino acids 93-97, 103-107, 223-227, 382-386, 522-526, 579-583,  
608-612, 624-628, 625-629

**Casein kinase II phosphorylation site.**

amino acids 51-55, 95-99, 242-246, 468-472, 487-491

**Tyrosine kinase phosphorylation site.**

amino acids 570-579

**N-myristoylation site.**

amino acids 13-19, 96-102, 158-164, 221-227, 352-358, 437-443,  
491-497, 492-498, 634-640, 702-708

**Cell attachment sequence.**

amino acids 277-280

**FIGURE 27**

GCCCCGGGACTGGCGCAAGGTGCCCAAGCAAGGAAAGAAATAATGAAGAGACACATGTGTTAG  
CTGCAGCCTTTTGAACACGCAAGAAGGAAATCAATAGTGTGGACAGGGCTGGAACCTTTAC  
CACGCTTGTTGGAGTAGATGAGGAATGGGCTCGTGATTATGCTGACATTCCAGCATGAATCT  
GGTAGACCTGTGGTTAACCCGTTCCCTCTCCATGTGTCTCCTCCTACAAAGTTTGTTCCTTA  
TGATACTGTGCTTTTCACTTCTGCCAGTATGTGTCCAAGGGCTGTCTTTGTTCTTCCTCTGGG  
GGTTTAAATGTCACCTGTAGCAATGCAAATCTCAAGGAAATACCTAGAGATCTTCCTCCTGA  
AACAGTCTTACTGTATCTGGACTCCAATCAGATCACATCTATTCCCAATGAAATTTTTAAGG  
ACCTCCATCAACTGAGAGTTCTCAACCTGTCCAAAATGGCATTGAGTTTATCGATGAGCAT  
GCCTTCAAAGGAGTAGCTGAAACCTTGCAGACTCTGGACTTGTCCGACAATCGGATTCAAAG  
TGTGCACAAAATGCCTTCAATAACCTGAAGGCCAGGGCCAGAATTGCCAACAACCCCTGGC  
ACTGCGACTGTACTCTACAGCAAGTTCTGAGGAGCATGGCGTCCAATCATGAGACAGCCCAC  
AACGTGATCTGTAAAACGTCCGTGTTGGATGAACATGCTGGCAGACCATTCTCAATGCTGC  
CAACGACGCTGACCTTTGTAACCTCCCTAAAAAACTACCGATTATGCCATGCTGGTCACCA  
TGTTTGGCTGGTTCACTATGGTGATCTCATATGTGGTATATTTATGTGAGGCAAAATCAGGAG  
GATGCCCCGGAGACACCTCGAATACTTGAAATCCCTGCCAAGCAGGCAGAAGAAAGCAGATGA  
ACCTGATGATATTAGCACTGTGGTATAGTGTCCAAACTGACTGTCAATTGAGAAAGAAAGAAA  
GTAGTTTGGCATTGCAGTAGAAATAAGTGGTTTACTTCTCCCATCCATTGTAAACATTTGAA  
ACTTTGTATTTTTCAGTTTTTTTTTTGAATTATGCCACTGCTGAACTTTTAACAAACACTACAACA  
TAAATAATTTGAGTTTAGGTGATCCACCCCTTAATTGTACCCCGATGGTATATTTCTGAGT  
AAGCTACTATCTGAACATTAGTTAGATCCATCTCACTATTTAATAATGAAATTTATTTTTTT  
AATTTAAAAGCAAATAAAAGCTTAACTTTGAACCATGGGAAAAAAAAAAAAAAAAAAAAAAAAACA

**FIGURE 28**

MNLVDLWLTRSLSMCLLLQS FVLMILCFHSASMC PKGCLCSSSGGLNVTC SNANLKEIPRDL  
PPETVLLYLDSNQITSIPNEIFKDLHQLRV LNLSKNGIEFIDEHAFKGV AETLQTLDLSDNR  
IQSVHKNAFN NLKARARIANNPWHCDCTLQQVLRSMASNHETAHNVIC KTSVLD E HAGRPFL  
NAANDADLCNLPKKT<sup>T</sup>TDYAMLVTMFGWFTMVISYVVYYVRQNQEDARRHLEYL KSLPSRQKK  
ADEPDDISTVV

**Signal sequence:**

amino acids 1-33

**Transmembrane domain:**

amino acids 205-220

**N-glycosylation site.**

amino acids 47-51, 94-98

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 199-203

**Casein kinase II phosphorylation site.**

amino acids 162-166, 175-179

**N-myristoylation site.**

amino acids 37-43, 45-51, 110-116

**FIGURE 29**

ACCGAGCCGAGCGGACCGAAGGCGCGCCGAGATGCAGGTGAGCAAGAGGATGCTGGCGGGG  
GGCGTGAGGAGCATGCCAGCCCCCTCCTGGCCTGCTGGCAGCCCATCCTCCTGCTGGTGCT  
GGGCTCAGTGCTGTGAGGCTCGGCCACGGGCTGCCCGCCCCGCTGCGAGTGCTCCGCCCAGG  
ACCGCGCTGTGCTGTGCCACCGCAAGTGCCTTTGTGGCAGTCCCCGAGGGCATCCCCACCGAG  
ACGCGCTGCTGGACCTAGGCAAGAACCGCATCAAAACGCTCAACCAGGACGAGTTCCGCCAG  
CTCCCCGCACCTGGAGGAGCTGGAGCTCAACGAGAACATCGTGAGCGCCGTTGGAGCCCGCG  
CCTTCAACAACCTCTTCAACCTCCGGACGCTGGGTCTCCGAGCAACCGCTGAAGCTCATC  
CCGCTAGGCGTCTTCACTGGCCTCAGCAACCTGACCAAGCAGGACATCAGCGAGAACAAGAT  
CGTTATCCTACTGGACTACATGTTTCAGGACCTGTACAACCTCAAGTCACTGGAGGTTGGCG  
ACATGACCTCGTCTACATCTCTCACCGCGCTTCCAGCGCCTCAACAGCCTGGAGCAGCTG  
ACGCTGGAGAAATGCAACCTGACCTCCATCCCCACCGAGGCGCTGTCCACCTGCACGGCCT  
CATCGFCTGAGGCTCCGGCACCTCAACATCAATGCCATCCGGGACTACTCCTTCAAGAGGC  
TGTACCGACTCAAGGTCTTGGAGATCTCCCACTGGCCCTACTTGGACACCATGACACCCAAC  
TGCCCTACGGCCTCAACCTGACGTCCCTGTCCATCACACACTGCAATCTGACCGCTGTGCC  
CTACTGGCCGTCGCCACCTAGTCTATCTCCGCTTCTCAACCTCTCCTACAACCCCATCA  
GCACCATTGAGGGCTCCATGTTGCATGAGCTGCTCCGGCTGCAGGAGATCCAGTGGTGGGC  
GGGCAGCTGGCCGTGGTGGAGCCCTATGCCTTCCGCGGCCTCAACTACCTGCGCGTGTCAA  
TGTCTCTGGCAACCAGCTGACCACACTGGAGGAATCAGTCTTCCACTCGGTGGGCAACCTGG  
AGACACTCATCCTGGACTCCAACCGCTGGCCTGGCAGTGTCCGCTCCTGTGGGTGTTCGG  
CGCCGCTGGCGGCTCAACTTCAACCGGCAGCAGCCACGTGCGCCACGCCGAGTTTGTCCA  
GGCAAGGAGTTCAAGGACTTCCCTGATGTGCTACTGCCCAACTACTTACCTGCCCGCGG  
CCCGCATCCGGGACCGCAAGGCCAGCAGGTGTTTGTGGACGAGGGCCACACGTTGAGTTT  
GTGTGCCGGGCCGATGGCGACCCGCCGCCCATCCTCTGGCTCTCACCCGAAAGCACCT  
GGTCTCAGCCAAGAGCAATGGGCGGCTCACAGTCTTCCCTGATGGCAGCTGGAGGTGCGCT  
ACGCCCAGGTACAGGACAACGGCACGTACCTGTGCATCGCGGCAACGCGGGCGGCAACGAC  
TCCATGCCCGCCACCTGCATGTGCGCAGCTACTCGCCCGACTGGCCCCATCAGCCCAACAA  
GACCTTCGCTTTTCACTCCAACCAGCCGGGCGAGGGAGAGGCCAACAGCACCCGCGCCACTG  
TGCTTTCCCTTCGACATCAAGACCTCATCATCGCCACCACCATGGGCTTCACTCTTTTC  
CTGGCGCTCGTCTCTTCTGCTGGTGTGCTGTTTCTCTGGAGCCGGGGCAAGGGCAACAC  
AAAGCACAACATCGAGATCGAGTATGTGCCCCGAAAGTCCGACGCAGGCATCAGTCCGCCG  
ACGCGCCCCGCAAGTTCAACATGAAGATGATATGAGGGCCGGGGCGGGGGCAGGGACCCCCG  
GGCGCCGGGGCAGGGGAAGGGGCTGGTGCACCTGCTCACTCTCCAGTCTTCCACCTC  
CTCCCTACCCTTCTACACAGTTCTCTTTCTCCCTCCCGCCTCCGTCCCCTGCTGCCCCCG  
CCAGCCCTCACACCTGCCCTCCTTCTACCAGGACCTCAGAAGCCCAGACCTGGGGACCCCA  
CCTACACAGGGGCATTGACAGACTGGAGTTGAAAGCCGACGAACCGACACGCGGACAGTCA  
ATAATTCAATAAAAAAGTTACGAACTTCTCTGTAACCTGGGTTTTCAATAATTATGGATTT  
TATGAAAACCTGAAATAATAAAAAAGAGAAAAAACTAAAAAAAAAAAAAAAAAAAAA

**FIGURE 30**

MQVSKRMLAGGVRSMPSPILLACWQPIILLVLSVLSGSATGCPPRCECSAQDRAVLCHRKCF  
VAVPEGIPTETRLLDLGKNRIKTLNQDEFASFPHLEELNENIVSAVEPGAFNNLFLNRTL  
GLRSNRLKLIPLGVFTGLSNLTKQDISENKIVILLDYMFDLYNLKSLEVGDNLDVYISHRA  
FSGLNSLEQLTLEKCNLTSIPTALSHLHGLIVLRLRHLNINAI RDYSFKRLYRLKVLEISH  
WPYLDTMTPNCLYGLNLTSLSITHCNLTAVPYLAVRHLVYLRFLNLSYNPISTIEGSM LHEL  
LRLQEIQLVGGQLAVVEPYAFRGLNYLRVLNVSGNQLTTLLESVFHSGVGNLETLILDSNPLA  
CDCRLLWVFRRRWRLNFNRRQPTCATPEFVQKFKDFPDVLLPNYFTCRRARIRDRKAQQV  
FVDEGHTVQFVCRADGDPPAILWLSPRKHLVSAKSNGLTVFPDGTLEVRYAQVDNGTYL  
CIAANAGGNDSMPAHLHVRSPDWP HQPNKTFAFISNQPGEGEANSTRATVPFPFDIKTLI  
IATTMGFISFLGVVLFCLVLLFLWSRGKGTKHNI EIEYVPRKSDAGISSADAPRKFNMKMI

**Signal sequence:**

amino acids 1-41

**Transmembrane domain:**

amino acids 556-578

**N-glycosylation site.**

amino acids 144-148, 202-206, 264-268, 274-278, 293-297, 341-345,  
492-496, 505-509, 526-530, 542-546

**Casein kinase II phosphorylation site.**

amino acids 49-53, 108-112, 146-150, 300-304, 348-352, 349-353,  
607-611

**Tyrosine kinase phosphorylation site.**

amino acids 590-598

**N-myristoylation site.**

amino acids 10-16, 32-38, 37-43, 113-119, 125-131, 137-143,  
262-268, 320-326, 344-350, 359-365, 493-499, 503-509, 605-611

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 32-43



**FIGURE 31**

CCCACGCGTCCGCACCTCGGCCCGGGCTCCGAAGCGGCTCGGGGGCGCCCTTTTCGGTCAAC  
ATCGTAGTCCACCCCTCCCATCCCCAGCCCCGGGGATTTCAGGCTCGCCAGCGCCCAGCC  
AGGGAGCCGGCCGGGAAGCGCGATGGGGCCCCAGCCGCCTCGCTCCTGCTCCTGCTCCTGC  
TGTTTCGCTGCTGCTGGGCGCCCGGGGCCAACCTCTCCAGGACGACAGCCAGCCCTGG  
ACATCTGATGAAACAGTGGTGGCTGGTGGCACCGTGGTGTCAAGTGCCAAGTGAAAGATCA  
CGAGGACTCATCCCTGCAATGGTCTAACCTGCTCAGCAGACTCTCTACTTTGGGGAGAAGA  
GAGCCCTTCGAGATAATCGAATTCAGCTGGTTACCTCTACGCCCCACGAGCTCAGCATCAGC  
ATCAGCAATGTGGCCCTGGCAGACGAGGGCGAGTACACCTGCTCAATCTTCACTATGCCTGT  
GCCAACTGCCAAGTCCCTCGTCACTGTGCTAGGAATTCACAGAAGCCCATCATCACTGGTT  
ATAAATCTTCATTACGGGAAAAAGACACAGCCACCCTAAACTGTGAGTCTTCTGGGAGCAAG  
CCTGCAGCCCCGGCTCACCTGGAGAAAGGGTGACCAAGAACTCCACGGAGAACCAACCCGCAT  
ACAGGAAGATCCCAATGGTAAAACCTTCACTGTGAGCAGCTCGGTGACATTCCAGGTTACCC  
GGGAGGATGATGGGGCGAGCATCGTGTGCTCTGTGAACCATGAATCTCTAAAGGGAGCTGAC  
AGATCCACCTCTCAACGCATTGAAGTTTTATACACCAACTGCGATGATTAGGCCAGACCC  
TCCCCATCCTCGTGAGGGCCAGAAGCTGTTGCTACACTGTGAGGGTCGCGGCAATCCAGTCC  
CCCAGCAGTACCTATGGGAGAAGGAGGGCAGTGTGCCACCCCTGAAGATGACCCAGGAGAGT  
GCCCTGATCTTCCCTTTCCCTCAACAAGAGTGACAGTGGCACCTACGGCTGCACAGCCACCAG  
CAACATGGGCAGCTACAAGGCCTACTACACCCTCAATGTTAATGACCCAGTCCGGTGGCCCT  
CCTCCTCCAGCACCTACCACGCCATCATCGGTGGGATCGTGGCTTTTCATTGTCTTCTGCTG  
CTCATCATGCTCATCTTCCCTGGCCACTACTTGATCCGGCACAAAGGAACCTACCTGACACA  
TGAGGCAAAAGGCTCCGACGATGCTCCAGACGCGGACACGGCCATCATCAATGCAGAAGGCG  
GGCAGTCAGGAGGGGACGACAAGAAGGAATATTTTCATCTAGAGGCGCCTGCCACTTCCCTGC  
GCCCCCAGGGGCCCTGTGGGGACTGCTGGGGCCGTCACCAACCCGGACTTGTACAGAGCAA  
CCGCAGGGCCGCCCTCCCGCTTGCTCCCAGCCCACCCACCCCTGTACAGAATGTCTGC  
TTTGGGTGCGGTTTTTGTACTCGGTTTTGGAATGGGGAGGGAGGAGGGCGGGGGAGGGGAGGG  
TTGCCCTCAGCCCTTCCGTGGCTTCTCTGCATTTGGGTATTATTATTTTTGTAAACAATCC  
CAAATCAAATCTGTCTCCAGGCTGGAGAGGCAGGAGCCCTGGGGTGAGAAAAGCAAAAAACA  
AACAAAAACA

**FIGURE 32**

MGAPAASLLLLLLLLFACCWAPGGANLSQDDSQPWTSDETVVAGGTVVVKCQVKDHEDSSLQW  
SNPAQQTLYFGEKRALRDNRILQVLTSTPHELISISNVALADEGEYTCSEFTMPVRTAKSLV  
TVLGIPQKPIITGYKSSLREKDTATLNCQSSGSKPAARLTWRKGDQELHGEPTRIQEDPNGK  
TFTVSSSVTFQVTRREDDGASIVCSVNHESLKGADRSTSQRIEVLYTPTAMIRPDPPHPREGQ  
KLLHCEGRGNPVPQQYLWEKEGSVPPLKMTQESALIFPFLNKSDSGTYGCTATSNMGSYKA  
YYTLNVNDPSPVPSSSSTYHAIIGGIVAFIVFLLLIMLIFLGHYLIRHKGTYLTHEAKGSDD  
APDADTAIINAEGGQSGGDDKKEYFI

**Signal sequence:**

amino acids 1-20

**Transmembrane domain:**

amino acids 331-352

**N-glycosylation site.**

amino acids 25-29, 290-294

**Casein kinase II phosphorylation site.**

amino acids 27-31, 35-39, 89-93, 141-145, 199-203, 388-392

**N-myristoylation site.**

amino acids 2-8, 23-29, 156-162, 218-224, 295-301, 298-304,  
306-310, 334-340, 360-364, 385-389, 386-390

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 7-18

**FIGURE 33**

GGGGGTTAGGGGAGGAAGGAATCCACCCCCACCCCCCAAACCCCTTTTCTTCTCCTTTCTG  
 CTTCGGACATTGGAGCACTAAATGAACCTTGAATTGTGTCTGTGGCGAGCAGGATGGTTCGCTG  
 TTACTTTGTGATGAGATCGGGGATGAATTGCTCGCTTTAAAAATGCTGCTTTGGATTCTGTT  
 GCTGGAGACGTCTCTTTGTTTTGCCGCTGGAAACGTTACAGGGGACGTTTGCAAAGAGAAGA  
 TCTGTTCCCTGCAATGAGATAGAAGGGGACCTACACGTAGACTGTGAAAAAAGGGCTTACA  
 AGTCTGCAGCGTTTCACTGCCCGACTTCCAGTTTTACCATTTATTTCTGCATGGCAATTC  
 CCTCACTCGACTTTTCCCTAATGAGTTCGCTAACTTTTATAATGCGGTAGTTTGCACATGG  
 AAAACAATGGCTTGCATGAAATCGTTCCGGGGGCTTTTCTGGGGCTGCAGCTGGTGAAAAGG  
 CTGCACATCAACAACAACAAGATCAAGTCTTTTCGAAAGCAGACTTTTCTGGGGCTGGACGA  
 TCTGGAATATCTCCAGGCTGATTTTAAATTTATTACGAGATATAGACCCGGGGCCTTCCAGG  
 ACTTGAACAAGCTGGAGGTGCTCATTTTAAATGACAATCTCATCAGCACCTACCTGCCAAC  
 GTGTTCCAGTATGTGCCCATCACCCACCTCGACCTCCGGGGTAACAGGCTGAAAACGCTGCC  
 CTATGAGGAGGTCTTGGAGCAAATCCCTGGTATTGCGGAGATCCTGCTAGAGGATAACCTT  
 GGGACTGCACCTGTGATCTGCTCTCCCTGAAAAGAATGGCTGGAAAAATTCCCAAGAAATGCC  
 CTGATCGGCCGAGTGGTCTGCGAAGCCCCCAGACTGCAGGGTAAAGACCTCAATGAAAC  
 CACCAAGACAGGACTTGTGCTCTTTGAAAAACCAGTGGATTCTAGTCTCCCGGCGCCCCCTG  
 CCCAAGAAGAGACCTTTGCTCCTGGACCCCTGCCAACTCCTTTCAAGCAAATGGGCAAGAG  
 GATCATGCCACACCAGGGTCTGCTCCAAACGGAGGTACAAAGATCCAGGCAACTGGCAGAT  
 CAAAATCAGACCCACAGCAGCGATAGCGACGGGTAGCTCCAGGAACAAACCTTAGCTAACA  
 GTTTACCCTGCCCTGGGGCTGCAGCTGCGACCACATCCAGGGTCCGGTTTAAAGATGAAC  
 TGCAACAACAGGAACGTGAGCAGCTTGGCTGATTTGAAGCCCAAGCTCTCTAACGTGCAGGA  
 GCTTTTCTACGAGATAACAAGATCCACAGCATCCGAAAATCGCACTTTGTGGATTACAAGA  
 ACCTCATTCTGTTGGATCTGGGCAACAATAACATCGCTACTGTAGAGAAACAACACTTTCAAG  
 AACTTTTGGACCTCAGGTGGCTATACATGGATAGCAATTACCTGGACACGCTGTCCCGGGA  
 GAAATTCGCGGGGCTGCAAAACCTAGAGTACCTGAACGTGGAGTACAACGCTATCCAGCTCA  
 TCCTCCCGGCACTTTCAATGCCATGCCCAACTGAGGATCCTCATTCTCAACAACAACCTG  
 CTGAGGTCCCTGCCTGTGGACGTGTTTCGCTGGGGTCTCGCTCTCTAAACTCAGCCTGCACAA  
 CAATTACTTCATGTACCTCCCGGTGGCAGGGGTGCTGGACCAGTTAACCTCCATCATCCAGA  
 TAGACCTCCACGGAAACCCCTGGGAGTGCTCCTGCACAATTGTGCTTTCAAGCAGTGGGCA  
 GAACGCTTGGGTTCCGAAGTGCTGATGAGCGACCTCAAGTGTGAGACGCCGGTGAACCTCTT  
 TAGAAAGGATTTTATGCTCCTCTCCAATGACGAGATCTGCCCTCAGCTGTACGCTAGGATCT  
 CGCCACGTTAACTTCGCACAGTAAAAACAGCACTGGGTGGCGGAGACCGGGACGCACTCC  
 AACTCCTACCTAGACACCAGCAGGGTGTCCATCTCGGTGTTGGTCCCGGACTGCTGCTGGT  
 GTTTGTACCTCCGCTTACCGTGGTGGGCATGCTCGTGTATCTTACCTGAGGAACCGAAAGC  
 GGTCCAAGAGACGAGATGCCAACTCCTCCGCGTCCGAGATTAATCCCTACAGACAGTCTGT  
 GACTCTTCTACTGGCACAATGGGCCTTACAACGCAGATGGGGCCACAGAGTGTATGACTG  
 TGGCTCTCACTCGCTCTCAGACTAAGACCCCAACCCCAATAGGGGAGGGCAGAGGGGAGGCG  
 ATACATCCTTCCCCACCGCAGGCACCCCGGGGGCTGGAGGGGCGTGTAACCAAATCCCGCG  
 CCATCAGCCTGGATGGCATAAGTAGATAAATAACTGTGAGCTCGCACAAACGAAAGGGCCT  
 GACCCCTTACTTAGCTCCCTCCTTGAACAAGAGCAGACTGTGGAGAGCTGGGAGAGCGCA  
 GCCAGCTCGCTCTTTGCTGAGAGCCCCCTTTGACAGAAAGCCAGCACGACCTGTGCTGGAAG  
 AACTGACAGTGCCCTCGCCCTCGGCCCGGGGCTGTGGGGTTGGATGCCCGGTTCTATAC  
 ATATATACATATATCCACATCTATATAGAGAGATAGATATCTATTTTTCCCTGTGGATTAG  
 CCCCCTGATGGCTCCCTGTTGGCTACGCAGGGATGGGCAGTTGCACGAAGGCATGAATGTAT  
 TGTAATAAGTAACTTTGACTTCTGAC

## FIGURE 34

MLLWILLLETSLCFAAGNVTGDVCKEIKCSCNEIEGDLHVDCEKKGFTSLQRF TAPT SQFYH  
LFLHGNSLTRLFPNEFANFYNAVSLHMENGLHEIVPGAFLGLQLVKRLHINNNKIKSFRKQ  
TFLGLDDLEYLQADFNLLRDIDPGAQDLNKLEVLILNDNLISTLPANVFQYVPTTHLDLRG  
NRLKTLPEYEEVLEQIPGIAEILLEDNPWDCTCDLLSLKEWLENI PKNALIGRVVCEAPTRLQ  
GKDLNETTEQDLCPLKNRVDSSLPAPPAQEETFAPGPLEPFPFKTNGQEDHATPGSAPNGGTK  
IPGNWQIKIRPTAAIATGSSRNKPLANS LCPGGCSCDHIPGSSGLKMNCNNRRNVSSLADLKP  
KLSNVQELFLRDNKIHSIRKSHFVDYKNLILLDLGNNNIATVENNTFKNLLDLRWLYMDSNY  
LDTLSREKFAGLQNLLEYLNVEYNAIQILILPGTFFNAMPKLRILILNNNLLRSLPVDVFAGVSL  
SKLSLHNNYFMYLPEVAGVLDQLT SIIQIDLHGPNWECSC TIVPFKQWAERLGSEVLMSDLKC  
ETPVNFFRKDFMLLSNDEICPQLYARISPTLTSHSKNSTGLAETGTHSNSYLDTSRVSISVL  
VPGLLLVFVTSFAFTVVGMLVFILRNRKRKR RDANSSASEINSLQTVCDSSYWHNGPYNADG  
AHRVYDCGSHSLSD

**Signal sequence:**

amino acids 1-15

**Transmembrane domain:**

amino acids 618-638

**N-glycosylation site.**

amino acids 18-22, 253-257, 363-367, 416-420, 595-599, 655-659

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 122-126, 646-650

**Casein kinase II phosphorylation site.**

amino acids 30-34, 180-184, 222-226, 256-260, 366-370, 573-577,  
608-612, 657-661, 666-670, 693-697

**N-myristoylation site.**

amino acids 17-23, 67-73, 100-106, 302-308, 328-334, 343-349,  
354-360, 465-471, 493-499, 598-604, 603-609

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 337-348

**FIGURE 35**

AGTCGACTGCGTCCCCTGTACCCGGCGCCAGCTGTGTTCCCTGACCCCAGAATAACTCAGGGC  
TGCACCCGGCCCTGGCAGCGCTCCGCACACATTTCCCTGTGCGGGCCTAAGGGAAACTGTTGGC  
CGCTGGGCCCCGCGGGGGATTCTTGGCAGTTGGGGGTCCGTGCGGAGCGAGGGCGGAGGGG  
AAGGGAGGGGGAACCGGGTTGGGGAAGCCAGCTGTAGAGGGCGGTGACCGCGCTCCAGACAC  
AGCTCTGCGTCTCGAGCGGGACAGATCCAAGTTGGGAGCAGCTCTGCGTGCGGGGCCTCAG  
AGAATGAGGGCCGGCGTTTCGCCCTGTGCCCTCCTCTGGCAGGCGCTCTGGCCCCGGGCCGGGCGG  
CGGCGAACACCCCACTGCCGACCGTGCCTGGCTGCTCGGCCTCGGGGGCCTGCTACAGCCTGC  
ACCACGCTACCATGAAGCGGCAGGCGGCCGAGGAGCCTGCATCCTGCCAGGTGGGGCGCTC  
AGCACCGTGCCTGCGGGCGAGCTGCGCGTGTGCTGCGCTCCTGCGGGCAGGCCCGCTC  
GCCCCGAGGGGGCTCCAAAGACCTGCTGTTCTGGGTGCGACTGGAGCGCAGGCGTTCCCACT  
GCACCCTGGAGAACGAGCCTTTGCGGGGTTTCTCCTGGCTGTCTCCGACCCCGGCGGTCTC  
GAAAGCGACACGCTGCAGTGGGTGGAGGAGCCCCAACGCTCCTGCACCGCGCGGAGATGCGC  
GGTACTCCAGGCCACCGGTGGGGTCCAGCCCGCAGGCTGGAAGGAGATGCGATGCCACCTGC  
GCGCCAACGGCTACCTGTGCAAGTACCAGTTTGGAGTCTTGTGTCTGCGCCGCGCCCCGGG  
GCCGCTCTAACTTGAGCTATCGCGCGCCCTTCCAGCTGCACAGCGCCGCTCTGGACTTCAG  
TCCACCTGGGACCGAGGTGAGTGCCTCTGCCGGGGACAGCTCCCGATCTCAGTTACTTGCA  
TCGCGGACGAAATCGGCGCTCGCTGGGACAACTCTCGGGCGATGTGTTGTGTCCCTGCCCC  
GGGAGGTACCTCCGTGCTGGCAAATGCGCAGAGCTCCCTAACTGCCTAGACGACTTGGGAGG  
CTTTGCCCTGCGAATGTGCTACGGGCTTCGAGCTGGGGAAGGACGGCCGCTCTTGTGTGACCA  
GTGGGGAAGGACAGCCGACCTTTGGGGGACCGGGTGCCACCAGGGCGCCCGCCGGCCACT  
GCAACCAGCCCCGTGCCGAGAGAACATGGCCAATCAGGGTCCGACGAGAAGCTGGGAGAGAC  
ACCCTTGTCCCTGAACAAGACAATTCAGTAACATCTATTCCCTGAGATTCCTCGATGGGGAT  
CACAGAGCACGATGTCTACCCTTCAAATGTCCCTTCAAGCCGAGTCAAAGGCCACTATCACC  
CCATCAGGGAGCGTGATTTCCAAGTTAATTTCTACGACTTCCCTGCCACTCCTCAGGCTTT  
CGACTCCTCCTCTGCCGTGGTCTTCATATTTGTGAGCACAGCAGTAGTAGTGTGGTGATCT  
TGACCATGACAGTACTGGGGCTTGTCAAGCTCTGCTTTACGAAAGCCCCCTTTCCAGCCA  
AGGAAGGAGTCTATGGGCCCGCGGGCCTGGAGAGTGTATCCTGAGCCCGCTGCTTTGGGCTC  
CAGTTCTGCACATTGCACAAACAATGGGCTGAAAGTCGGGGACTGTGATCTGCGGGACAGAG  
CAGAGGGTGCCTTGTGCGGGAGTCCCCCTTTGGCTCTAGTGATGCATAGGGGAAACAGGGGA  
CATGGGCACTCCTGTGAACAGTTTTTCACTTTTTGATGAAACGGGGAACCAAGAGGAACTTAC  
TTGTGTAACCTGACAATTTCTGCAGAAATCCCCCTTCCCTCTAAATTCCTTTACTCCACTGAG  
GAGCTAAATCAGAACTGCACACTCCTTCCCTGATGATAGAGGAAGTGGAAAGTGCCTTTAGGA  
TGGTGATACTGGGGACCGGTTAGTGTGGGGAGAGATATTTTCTTATGTTTATTCCGAGAA  
TTTGGAGAAGTGATTGAACTTTTCAAGACATTTGGAAACAAATAGAACACAATATAATTTACA  
TTAAAAATAATTTCTACAAAATGGAAAGGAAATGTTCTATGTTTTCAGGCTAGGAGTAT  
ATTGGTTGCAAAATCCAGGGAAAAAATAAAAAATAAAAAATTAAGGATTGTTGAT

## **FIGURE 36**

MRPAFALCLLWQALWPGPGGGGEHPTADRAGCSASGACYSLHHATMKRQAAEEACILRGGALS  
TVRAGAE LRAVLALLRAGPGPGGGSKDLLFWVALERRRSHCTLENEPLRGFSWLSSDPGGLE  
SDTLQWVEEPQRSCTARRCAVLQATGGVEPAGWKEMRCHLRANGYLCKYQFEVLC PAPRPGA  
ASNLSYRAPFQLHSAALDFSPPGTEVSALCRGQLPISVTCIADEIGARWDKLSGDVLCPCPG  
RYLRAGKCAELPNCLDDLGGFACECATGFELGKDGRSCVTS GEGQPTLGGTGVPTRRPPATA  
TSPVQPRTWPPIRVDEKLGETPLVPEQDNSVTSIPEI PRWGSQSTMSTLQMSLQAESKATITP  
SGSVISKFNSTTSSATPQAFDSSSAVVFIFVSTAVVVLVILTMTVLGLVKLCFHESPSSQPR  
KESMGPPGLESDPEPAALGSSSAHCTNNGVKVGDCLDRDRAEGALLAESPLGSSDA

**Signal sequence:**

amino acids 1-16

**Transmembrane domain:**

amino acids 399-418

**N-glycosylation site.**

amino acids 189-193, 381-385

**Glycosaminoglycan attachment site.**

amino acids 289-293

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 98-102, 434-438

**Casein kinase II phosphorylation site.**

amino acids 275-279, 288-292, 342-346, 445-449

**N-myristoylation site.**

amino acids 30-36, 35-41, 58-64, 59-65, 121-127, 151-157,  
185-191, 209-215, 267-273, 350-356, 374-380, 453-459, 463-469,  
477-483

**Aspartic acid and asparagine hydroxylation site.**

amino acids 262-274

**FIGURE 37**

CGGACGCGTGGGATTGAGCAGTGGCCTGTGGCTGCCAGAGCAGCTCCTCAGGGGAAACTAAG  
CGTCGAGTCAGACGGCACCATAATCGCCTTTAAAAGTGCCTCCGCCCTGCCGGCCGCGTATC  
CCCCGGCTACCTGGGCCGCCCCGCGCGGTGCGCGCGTGAGAGGGAGCGCGCGGGCAGCCGA  
GCGCCGGTGTGAGCCAGCGCTGCTGCCAGTGTGAGCGGCGGTGTGAGCGCGGTGGGTGCGGA  
GGGGCGTGTGTGCCGGCGCGCGCCGTGGGGTGCAAACCCCGAGCGTCTACGCTGCCATGA  
GGGGCGGAACGCCTGGGCGCCACTCTGCCTGCTGCTGGCTGCCGCCACCCAGCTCTCGCGG  
CAGCAGTCCCCAGAGAGACCTGTTTTACATGTGGTGGCATTCCTTACTGGAGAGTCTGGATT  
TATTGGCAGTGAAGGTTTTCTGGAGTGTACCCTCCAAATAGCAAATGTACTTGGAAAATCA  
CAGTTCCCGAAGGAAAAGTAGTCGTTCTCAATTTCCGATTCATAGACCTCGAGAGTGACAAC  
CTGTGCCGCTATGACTTTGTGGATGTGTACAATGGCCATGCCAATGGCCAGCGCATTGGCCG  
CTTCTGTGGCACTTTCCGGCCTGGAGCCCTTGTGTCCAGTGGCAACAAGATGATGGTGCACA  
TGATTTCTGATGCCAACACAGCTGGCAATGGCTTCATGGCCATGTTCTCCGCTGTGAACCA  
AACGAAAGAGGGGATCAGTATTGTGGAGGACTCCTTGACAGACCTTCCGGCTCTTTAAAAC  
CCCCAACTGGCCAGACCGGGATTACCCTGCAGGAGTCACTTGTGTGGCACATTGTAGCCC  
CAAAGAATCAGCTTATAGAATTAAGTTTGAGAAGTTTGATGTGGAGCGAGATAACTACTGC  
CGATATGATTATGTGGCTGTGTTAATGGCGGGGAAGTCAACGATGCTAGAAGAATTGGAAA  
GTATTGTGGTGTATAGTCCACCTGCGCCAATTGTGTCTGAGAGAAATGAACTTCTTATTCAGT  
TTTTATCAGACTTAAGTTAACTGCAGATGGGTTTATTTGGTCACTACATATTCAGGCCAAAA  
AAACTGCCTACAACACAGAACAGCCTGTCAACAAAAGTGTAGACGGACGGGGACTCTGGAGGGCAATTATT  
GTTCAAGTGACTTTGTATTAGCCGGCACTGTTATCACAAACCATCACTCGCGATGGGAGTTTG  
CACGCCACAGTCTCGATCATCAACATCTACAAAGAGGGAAATTTGGCGATTCAGCAGGCGGG  
CAAGAACATGAGTGCCAGGCTGACTGTGCTGCAAGCAGTGCCCTCTCCTCAGAAGAGGTC  
TAAATTACATTATTATGGGCCAAGTAGGTGAAGATGGGCGAGGCAAAATCATGCCAAACAGC  
TTTATCATGATGTTCAAGACCAAGAATCAGAAGCTCCTGGATGCCTTAAAAAATAAGCAATG  
TTAACAGTGAAGTGTGTCATTTAAGCTGTATTCTGCCATTGCCTTTGAAAGATCTATGTTCT  
TCTCAGTAGAAAAAAAAATACTTATAAAATTACATATTCTGAAAGAGGATTCGAAAGATGG  
GACTGGTTGACTCTTACATGATGGAGGTATGAGGCCTCCGAGATAGCTGAGGGAAGTTCTT  
TGCCCTGCTGTGAGAGGAGCAGCTATCTGATTGGAAACCTGCCGACTTAGTGCGGTGATAGGA  
AGCTAAAAGTGTCAAGCGTTGACAGCTTGAAGCGTTTATTTATACATCTCTGTAAAAGGAT  
ATTTTAGAATTGAGTTGTGTGAAGATGTCAAAAAAGATTTTAGAAGTGCAATATTTATAGT  
GTTATTTGTTTACCTTCAAGCCTTTGCCCTGAGGTGTTACAATCTTGTCTTGCCTTTTCTA  
AATCAATGCTTAATAAAATATTTTTAAAGGAAAAAAAAAAAA

**FIGURE 38**

MRGANAWAPLCLLLAAATQLSRQOSPVPVFTCGGILTGESGFIFGSEGFPGVYPPNSKCTWK  
ITVPEGKVVVNLNFRFIDLESNDLCRYDFVDVYNGHANGQRIGRFCGTFRPGALVSSGNKMMV  
QMISDANTAGNGFMAMFSAAEPNERGDQYCGLLDRPSGSFKTPNWPDRDYPAGVTCVWHIV  
APKNQLIELKFEKFDVERDNYCRYDYVAVFNGGEVNDARRIGKYCGDSPPAPIVSERNELLI  
QFLSDLSLTADGFIGHYIFRPKKLPTTTEQPVTTFPVTTLGLKPTVALCQQKCRRTGTLEGN  
YCSSDFVLAGTVITTTITRDGSLHATVSIINIYKEGNLAIQQAGKNMSARLTVVCKQCPLLR  
GLNYIIMGQVGEDGRGKIMPNSFIMMFKTKNQKLLDALKNKQC

**Signal sequence:**

amino acids 1-23

**N-glycosylation site.**

amino acids 355-359

**Casein kinase II phosphorylation site.**

amino acids 64-68, 142-146, 274-278

**Tyrosine kinase phosphorylation site.**

amino acids 199-208

**N-myristoylation site.**

amino acids 34-40, 35-41, 100-106, 113-119, 218-224, 289-295,  
305-311, 309-315, 320-326, 330-336

**Cell attachment sequence.**

amino acids 149-152



**FIGURE 39**

CGGACGCGTGGGCGGACGCGTGGGCGGCCACGGCGCCCGGGCTGGGGCGGTTCGCTTCTT  
CCTTCTCCGTGGCCTACGAGGGTCCCCAGCCTGGGTAAAGATGGCCCCATGGCCCCGAAGG  
GCCTAGTCCCAGCTGTGCTCTGGGGCCTCAGCCTCTTCTCAACCTCCCAGGACCTATCTGG  
CTCCAGCCCTCTCCACCTCCCCAGTCTTCTCCCCGCTCAGCCCCATCCGTGTATACCTG  
CCGGGACTGGTTGACAGCTTTAACAAGGGCCTGGAGAGAACCATCCGGGACAACTTTGGAG  
GTGAAACACTGCCTGGGAGGAAGAGAATTTGTCCAATACAAAGACAGTGAGACCCGCTG  
GTAGAGGTGCTGGAGGGTGTGTGCAGCAAGTCAGACTTCGAGTGCCACCGCTGCTGGAGCT  
GAGTGAGGAGCTGGTGGAGAGCTGGTGGTTTCAACAAGCAGCAGGAGGCCCGGACCTCTTCC  
AGTGGCTGTGCTCAGATTCCTGAAGCTCTGCTGCCCGCAGGCACCTTCGGGCCCTCCTGC  
CTTCCCTGTCTGGGGGAACAGAGAGGCCCTGCGGTGGCTACGGGCAGTGTGAAGGAGAAGG  
GACACGAGGGGGCAGCGGGCACTGTGACTGCCAAGCCGGCTACGGGGGTGAGGCCTGTGGCC  
AGTGTGGCCTTGCTACTTTGAGGCAGAACGCAACGCCAGCCATCTGGTATGTTTCGGCTTGT  
TTTGGCCCCGTGCCCCGATGCTCAGGACCTGAGGAATCAAACCTGTTTGAATGCAAGAAGGG  
CTGGGCCCTGCATCACCTCAAGTGTGTAGACATTGATGAGTGTGGCACAGAGGGAGCCAACT  
GTGGAGCTGACCAATCTGCGTGAACACTGAGGGCTCCTATGAGTGCCGAGACTGTGCCAAG  
GCCTGCCTAGGCTGCATGGGGCAGGGCCAGGTGCTGTGTAAGAAGTGTAGCCCTGGCTATCA  
GCAGGTGGGCTCCAAGTGTCTCGATGTGGATGAGTGTGAGACAGAGGTGTGTCCGGGAGAGA  
ACAAGCAGTGTGAAAACACCGAGGGCGGTTATCGCTGCATCTGTGCCGAGGGCTACAAGCAG  
ATGGAAGGCATCTGTGTGAAGGAGCAGATCCCAGAGTCAGCAGGCTTCTTCTCAGAGATGAC  
AGAAGACGAGTTGGTGGTGTGTCAGCAGATGTTCTTTGGCATCATCATCTGTGCACTGGCCA  
CGCTGGCTGCTAAGGGCGACTTGGTGTTCACCGCCATCTTCATTGGGGCTGTGGCGGCCATG  
ACTGGCTACTGGTTGTGTCAGAGCGCAGTGACCGTGTGCTGGAGGGCTTCATCAAGGGCAGATA  
ATCGCGGCCACCACCTGTAGGACCTCCTCCCACCCACGCTGCCCCCAGAGCTTGGGCTGCCC  
TCCTGCTGGACACTCAGGACAGCTTGGTTTATTTTTGAGAGTGGGGTAAGCACCCCTACCTG  
CCTTACAGAGCAGCCCAGGTACCCAGGCCCGGGCAGACAAGGCCCTGGGGTAAAAAGTAGC  
CCTGAAGGTGGATACCATGAGCTCTTACCTGGCGGGGACTGGCAGGCTTACAATGTGTGA  
ATTTCAAAGTTTTTCTTAATGGTGGCTGCTAGAGCTTTGGCCCCGCTTAGGATTAGGTG  
GTCCCTCACAGGGGTGGGGCCATCACAGCTCCCTCCTGCCAGCTGCATGCTGCCAGTTCCTGT  
TCTGTGTTACCACATCCCCACACCCATTGCCACTTATTTATTTCATCTCAGGAAATAAAGA  
AAGGTCTTGGAAGTTAAAAAAAAAAAAAAAAAAAAAAAAA

## **FIGURE 40**

MAPWPPKGLVPAVLWGLSLFLNLP GPIIWLQPSPPPQSSPPPQPHPCHTCRGLVDSFNKGLER  
TIRDNFGGGNTAWEEENLSKYKDSETRLVEVLEGVCSKSDFECHRLLLELSEELVESWWFHKQ  
QEAPDLFQWLCSDSLKLCPPAGTFGSPCLPCPGGTERPCGGYGQCEGEGTRGGSGHCDCQAG  
YGGEACGQCGLGYFEAERNASHLVCSACFGPCARCSGPEESNCLQCKKGWALHHLKCV DIDE  
CGTEGANCGADQFCVNTTEGSYECRDCAKACLGC MGAGPGRCKKCSPGYQQVGSKCLDVDECE  
TEVCPGENKQCENTEGGYRCICAEGYKQMEGICVKEQIPESAGFFSEMTEDELVVLQQMFFG  
IIICALATLAAKGDVFTAIFIGAVAAMTGYWLSERSDRVLEGF IKGR

**Signal sequence:**

amino acids 1-29

**Transmembrane domain:**

amino acids 372-395

**N-glycosylation site.**

amino acids 79-83, 205-209

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 290-294

**Casein kinase II phosphorylation site.**

amino acids 63-67, 73-77, 99-103, 101-105, 222-226, 359-263

**N-myristoylation site.**

amino acids 8-14, 51-57, 59-65, 69-75, 70-76, 167-173, 173-179,  
177-183, 188-194, 250-256, 253-259, 267-273, 280-286, 283-289,  
326-332, 372-378, 395-401

**Aspartic acid and asparagine hydroxylation site.**

amino acids 321-333

**EGF-like domain cysteine pattern signature.**

amino acids 181-193

**FIGURE 41**

TGAGACCCTCCTGCAGCCTTCTCAAGGGACAGCCCCACTCTGCCTCTTGCTCCTCCAGGGCA  
GCACCATGCAGCCCCTGTGGCTCTGCTGGGCACTCTGGGTGTTGCCCTGGCCAGCCCCGGG  
GCCGCCCTGACCGGGGAGCAGCTCCTGGGCAGCCTGCTGCGGCAGCTGCAGCTCAAAGAGGT  
GCCACCCTGGACAGGGCCGACATGGAGGAGCTGGTCATCCCCACCCACGTGAGGGCCCAGT  
ACGTGGCCCTGCTGCAGCGCAGCCACGGGGACCGCTCCCGCGAAAGAGGTTTCAGCCAGAGC  
TTCCGAGAGGTGGCCGGCAGGTTCTTGGCGTTGGAGGCCAGCACACACCTGCTGGTGTTCGG  
CATGGAGCAGCGGCTGCCGCCAACAGCGAGCTGGTGCAGGCCGTGCTGCGGCTCTTCCAGG  
AGCCGGTCCCCAAGGCCGCGCTGCACAGGCACGGGGCGGCTGTCCCCGCGCAGCGCCCCGGCC  
CGGGTGACCGTCGAGTGGCTGCGCGTCCGCGACGACGGCTCCAACCGCACCTCCCTCATCGA  
CTCCAGGCTGGTGTCCGTCCACGAGAGCGGCTGGAAGGCCTTCGACGTGACCGAGGCCGTGA  
ACTTCTGGCAGCAGCTGAGCCGGCCCCGGCAGCCGCTGCTGCTACAGGTGTCCGGTGCAGAGG  
GAGCATCTGGGCCCGCTGGCGTCCGGCGCCCCACAAGCTGGTCCGCTTTGCCTCGCAGGGGGC  
GCCAGCCGGGCTTGGGGAGCCCCAGCTGGAGCTGCACACCCTGGACCTTGGGGACTATGGAG  
CTCAGGGCGACTGTGACCCTGAAGCACC AATGACCGAGGGCACCCGCTGCTGCCGCCAGGAG  
ATGTACATTGACCTGCAGGGGATGAAGTGGGCCGAGAACTGGGTGCTGGAGCCCCGGGCTT  
CCTGGCTTATGAGTGTGTGGGCACCTGCCGGCAGCCCCGGAGGCCCTGGCCTTCAAGTGGC  
CGTTTCTGGGGCCTCGACAGTGCATCGCCTCGGAGACTGACTCGCTGCCCATGATCGTCAGC  
ATCAAGGAGGGAGGCAGGACCAGGCCCCAGGTGGTCAGCCTGCCCAACATGAGGGTGCAGAA  
GTGCAGCTGTGCCTCGGATGGTGCCTCGTGCCAAGGAGGCTCCAGCCATAGGCGCCTAGTG  
TAGCCATCGAGGGACTTGACTTGTGTGTGTTTCTGAAGTGTTCGAGGGTACCAGGAGAGCTG  
GCGATGACTGAACTGCTGATGGACAAATGCTCTGTGCTCTCTAGTGAGCCCTGAATTTGCTT  
CCTCTGACAAGTTACCTCACCTAATTTTTGCTTCTCAGGAATGAGAATCTTTGGCCACTGGA  
GAGCCCTTGCTCAGTTTTCTCTATTCTTATTATTCACTGCACTATATTCTAAGCACTTACAT  
GTGGAGATACTGTAACCTGAGGGCAGAAAGCCCANTGTGTGCTATTGTTTACTTGTCTGTGAC  
TGGATCTGGGCTAAAAGTCCCTCCACCACCACTCTGGACCTAAGACCTGGGGTTAAGTGTGGGT  
TGTGCATCCCCAATCCAGATAATAAAGACTTTGTAAAACATGAATAAAAACATTTTATTCT  
AAAA

**FIGURE 42**

MQPLWLCWALWVLPLASPGAALTGEQLLGSLLRQLQLKEVPTLDRADMEELVIPTHVRAQYV  
ALLQRSHGDRSRGKRFSQSFREVAGRFLALEASTHLLVFGMEQRLPPNSELVQAVLRLRFQEP  
VPKAALHRHGRLSPRSARARVTVEWLRVRRDDGSNRTSLIDSRLVSVHESGWKAFDVTEAVNF  
WQLSRPRQPLLLQVSVQREHLGPLASGAHKLVRFASQGAPAGLGEPQLELHTLDLDG DYGAQ  
GDCDPEAPMTEGTRCCRQEMYIDLQGMKWAENWVLEPPGFLAYECVGTCTCRQPPEALAFKWP  
LGPRQCIASETDSLPMIVSIKEGGRTRPQVVSLPNMRVQKSCASD GALVPRRLQP

**Signal sequence:**

amino acids 1-18

**N-glycosylation site.**

amino acids 158-162

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 76-80

**Casein kinase II phosphorylation site.**

amino acids 68-72, 81-85, 161-165, 169-173, 319-323, 329-333

**N-myristoylation site.**

amino acids 19-25, 156-162, 225-231, 260-266, 274-280

**Amidation site.**

amino acids 74-78

**TGF-beta family signature.**

amino acids 282-298

**FIGURE 43**

GTCTGTTCCAGGAGTCCTTCGGCGGCTGTTGTGTCAGTGGCCTGATCGCGATGGGGACAAA  
GGCGCAAGTCGAGAGGAAACTGTTGTGCCCTCTTCATATTGGCGATCCTGTTGTGCTCCCTGG  
CATTGGGCAGTGTTACAGTGCACCTCTTCTGAACCTGAAGTCAGAATTCCTGAGAATAATCCT  
GTGAAGTTGTCCTGTGCCTACTCGGGCTTTTCTTCTCCCCGTGTGGAGTGGAAAGTTTGACCA  
AGGAGACACCACCAGACTCGTTTTGCTATAATAACAAGATCACAGCTTCCATGAGGACCGGG  
TGACCTTCTTGCCAACTGGTATCACCTTCAAGTCCGTGACACGGGAAGACACTGGGACATAC  
ACTTGTATGGTCTCTGAGGAAGGCGGCAACAGCTATGGGGAGGTCAAGGTCAAGCTCATCGT  
GCTTGTGCCTCCATCCAAGCCTACAGTTAACATCCCCTCCTCTGCCACCATTGGGAACCGGG  
CAGTGCTGACATGCTCAGAACAAGATGGTTCCCCACCTTCTGAATACACCTGGTTCAAAGAT  
GGGATAGTGATGCCTACGAATCCCAAAGCACCCGTGCCTTACGCAACTCTTCCATGTCTCT  
GAATCCCACAACAGGAGAGCTGGTCTTTGATCCCCTGTCAGCCTCTGATACTGGAGAATACA  
GCTGTGAGGCACGGAATGGGTATGGGACACCCATGACTTCAAATGCTGTGCGCATGGAAGCT  
GTGGAGCGGAATGTGGGGGTATCGTGGCAGCCGTCTTGTAAACCCTGATTCTCCTGGGAAT  
CTTGGTTTTTGGCATCTGGTTTGCCTATAGCCGAGGCCACTTTGACAGAACAAAGAAAGGGA  
CTTCGAGTAAGAAGGTGATTTACAGCCAGCCTAGTGCCCGAAGTGAAGGAGAATTCAAACAG  
ACCTCGTCATTCTGGTGAGGCCTGGTGGCTCACCGCCTATCATCTGCATTTGCCTTACT  
CAGGTGCTACCGGACTCTGGCCCCTGATGTCTGTAGTTTACAGGATGCCTTATTTGTCTTC  
TACACCCACAGGGCCCCCTACTTCTTCGGATGTGTTTTTAATAATGTCAGCTATGTGCCCC  
ATCCTCCTTCATGCCCTCCCTCCCTTTCTACCCTGCTGAGTGGCCTGGAACCTGTTTTAAA  
GTGTTTTATTCCCATTCTTTGAGGGATCAGGAAGGAATCCTGGGTATGCCATTGACTTCCC  
TTCTAAGTAGACAGCAAAAATGGCGGGGGTTCGAGGAATCTGCACTCAACTGCCACCTGGC  
TGGCAGGGATCTTTGAATAGGTATCTTGAGCTTGGTTCTGGGCTCTTTCCTTGTGTACTION  
GACCAGGGCCAGCTGTTCTAGAGCGGGAATTAGAGGCTAGAGCGGCTGAAATGGTTGTTTGG  
TGATGACACTGGGGTCTTCCATCTCTGGGGCCACTCTCTTCTGTCTTCCCATGGGAAGTG  
CCACTGGGATCCCTCTGCCCTGTCTCCTGAATACAAGCTGACTGACATTGACTGTGTCTGT  
GGAAAATGGGAGCTCTTGTGTGGAGAGCATAGTAAATTTTCAGAGAACTTGAAGCCAAAAG  
GATTTAAAACCGCTGCTCTAAAGAAAAGAAAACCTGGAGGCTGGGCGCAGTGGCTCACGCCTG  
TAATCCCAGAGGCTGAGGCAGGCGGATCACCTGAGGTGGGAGTTCGGGATCAGCCTGACCA  
ACATGGAGAAACCCTACTGGAAATACAAAGTTAGCCAGGCATGGTGGTGCATGCCGTAGTCT  
CCAGCTGCTCAGGAGCCTGGCAACAAGAGCAAACTCCAGCTCAAAAAAAAAAAAAAAAAA

## **FIGURE 44**

MGTKAQVERKLLCLFILAILLCSLALGSVTVHSSEPEVRIPENNPVKLS CAYSGFSSPRVEW  
KFDQGDTTTRLVCYNNKITASYEDRVTFLLPTGITFKSVTREDTGTYYTCMVSEEGGNSYGEVKV  
KLIVLVPPSKPTVNIPISSATIGNRAVLTCSEQDGSPPSEYTWFKDGI VMP TNP KSTRAFSNS  
SYVLNPTTGELVFDPLSASDTGEYSCEARNGYGTPMTSNAVRMEAVERNVGVIVA AVLVTLLI  
LLGILVFGIWFAYS RGHFDRTKKGTSSKKVIYSQPSARSEGEFKQTSSFLV

**Signal sequence:**

amino acids 1-27

**Transmembrane domain:**

amino acids 238-255

**N-glycosylation site.**

amino acids 185-189

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 270-274

**Casein kinase II phosphorylation site.**

amino acids 34-38, 82-86, 100-104, 118-122, 152-156, 154-158,  
193-197, 203-207, 287-291

**N-myristoylation site.**

amino acids 105-111, 116-122, 158-164, 219-225, 237-243, 256-262

**FIGURE 45**

CAGCGCGTGGCCGGCGCCGCTGTGGGGACAGCATGAGCGGCGGTTGGATGGCGCAGGTTGGA  
GCGTGGCGAACAGGGGCTCTGGGCCTGGCGCTGCTGCTGCTGCTCGGCCTCGGACTAGGCCT  
GGAGGCCGCCGCGAGCCCGCTTTCCACCCGACCTCTGCCCAGGCCCGCAGGCCCCAGCTCAG  
GCTCGTGCCACCCACCAAGTTCCAGTGCCGCACCAGTGGCTTATGCGTGCCCTCACCTGG  
CGCTGCGACAGGGACTTGGACTGCAGCGATGGCAGCGATGAGGAGGAGTGCAGGATTGAGCC  
ATGTACCCAGAAAGGGCAATGCCACCGCCCCCTGGCCTCCCCTGCCCTGCACCGGCGTCA  
GTGACTGCTCTGGGGAACTGACAAGAAACTGCGCAACTGCAGCCGCTGGCCTGCCTAGCA  
GGCGAGCTCCGTTGCACGCTGAGCGATGACTGCATTCCACTCACGTGGCGCTGCGACGGCCA  
CCCAGACTGTCCGACTCCAGCGACGAGCTCGGCTGTGGAACCAATGAGATCCTCCCGAAG  
GGGATGCCACAACCATGGGGCCCCCTGTGACCCTGGAGAGTGTACCTCTCTCAGGAATGCC  
ACAACCATGGGGCCCCCTGTGACCCTGGAGAGTGTCCCCTCTGTGCGGAATGCCACATCCTC  
CTCTGCCGGAGACCAGTCTGGAAGCCCAACTGCCTATGGGGTTATTGCAGCTGCTGCGGTGC  
TCAGTGCAAGCCTGGTCAACGCCACCCCTCCTCCTTTTGTCTGGCTCCGAGCCAGGAGCGC  
CTCCGCCCACTGGGGTTACTGGTGGCCATGAAGGAGTCCCTGCTGCTGTGAGAACAGAAGAC  
CTCGCTGCCCTTGAGGACAAGCACTTGCCACCACCGTCACTCAGCCCTGGGCGTAGCCGGACA  
GGAGGAGAGCAGTGATGCGGATGGGTACCCGGGCACACCAGCCCTCAGAGACCTGAGTTCTT  
CTGGCCACGTGGAACCTCGAACCCGAGCTCCTGCAGAAGTGGCCCTGGAGATTGAGGGTCCC  
TGGACACTCCCTATGGAGATCCGGGGAGCTAGGATGGGGAACCTGCCACAGCCAGAACTGAG  
GGGCTGGCCCCAGGCAGCTCCAGGGGGTAGAACGGCCCTGTGCTTAAGACACTCCCTGCTG  
CCCCGTCTGAGGGTGGCGATTAAAGTTGCTTC

**FIGURE 46**

MSGGWMAQVGAWRTGALGLALLLLLGLGLGLEAAASPLSTPTSAQAAGPSSGSCPPTKFQCR  
TSGLCVPLTWRCDRDLDCSDGSDEEEECRIEPTQKGQCPPPPGLPCPCTGVSDCSGGTDKKL  
RNC SRLACLAGE LRCTLSDDCIPLTWRC DGH PDCPDSSDELGCGTNEILPEGDATTMGPPVT  
LESV TSLRNAT TMGPPVTLESVPSVGNATSSSAGDQSGSPTAYGVIAAAVLSASLV TATLL  
LLSWLRAQERLRPLGLLVAMKESLLLSEQKTSLP

**Signal sequence:**

amino acids 1-30

**Transmembrane domain:**

amino acids 230-246

**N-glycosylation site.**

amino acids 126-130, 195-199, 213-217

**Casein kinase II phosphorylation site.**

amino acids 84-88, 140-144, 161-165, 218-222

**N-myristoylation site.**

amino acids 3-9, 10-16, 26-32, 30-36, 112-118, 166-172, 212-218,  
224-230, 230-236, 263-269

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 44-55

**Leucine zipper pattern.**

amino acids 17-39



**FIGURE 47**

CCCACGCGTCCGGTCTCGCTCGCTCGCGCAGCGGGCGGCAGCAGAGGTCGCGCACAGATGCGG  
GTTAGACTGGCGGGGGAGGAGGCGGAGGAGGGAAGGAAGCTGCATGCATGAGACCCACAGA  
CTCTTGCAAGCTGGATGCCCTCTGTGGATGAAAGATGATCATGGAATGAACCCGAGCAATG  
GAGATGGATTTCTAGAGCAGCAGCAGCAGCAGCAACCTCAGTCCCCCAGAGACTCTTG  
GCCGTGATCCTGTGGTTTCAGCTGGCGCTGTGCTTCGGCCCTGCACAGCTCACGGGCGGGTT  
CGATGACCTTCAAGTGTGTGCTGACCCCGCATTTCCCGAGAATGGCTTCAGGACCCCGAGCG  
GAGGGGTTTTCTTTGAAGGCTCTGTAGCCCGATTTCACTGCCAAGACGGATTCAAGCTGAAG  
GGCGCTACAAAGAGACTGTGTTTTGAAGCATTTTAATGGAACCCCTAGGCTGGATCCCAAGTGA  
TAATTCATCTGTGTGCAAGAAGATTGCCGTATCCCTCAAATCGAAGATGCTGAGATTCATA  
ACAAGACATATAGACATGGAGAGAAGCTAATCATCACTTGTTCATGAAGGATTCAAGATCCGG  
TACCCCGACCTACACAATATGGTTTCATTATGTGCGGATGATGGAACGTGGAATAATCTGCC  
CATCTGTCAAGGCTGCCCTGAGACCTCTAGCCTCTTCTAATGGCTATGTAAACATCTCTGAGC  
TCCAGACCTCCTTCCCGGTGGGGACTGTGATCTCCTATCGCTGCTTTCCCGGATTTAACTT  
GATGGGTCTGCGTATCTTGAGTGCCTTACAAAACCTTATCTGGTCGTCCAGCCCACCCCGGTG  
CCTTGCTCTGGAAGCCCAAGTCTGTCCACTACCTCCAATGGTGAGTCACGGAGATTTCTGCT  
GCCACCCGCGGCCCTTGTGAGCGCTACAACCACGGAACGTGGTGGAGTTTACTGCGATCCT  
GGCTACAGCCTCACAGCGACTACAAGTACATCACCTGCCAGTATGGAGAGTGGTTTTCTTC  
TTATCAAGTCTACTGCATCAAATCAGAGCAAACGTGGCCAGCACCCATGAGACCTCCTGA  
CCACGTGGAAGATTGTGGCGTTCACGGCAACCAGTGTGCTGCTGGTGTGCTGCTCGTCATC  
CTGGCCAGGATGTTCCAGACCAAGTTCAAGGCCACTTTCCCCCAGGGGGCTCCCCGGAG  
TTCCAGCAGTGACCCTGACTTTGTGGTGGTAGACGGCGTGCCCGTCATGCTCCCGTCCCTATG  
ACGAAGCTGTGAGTGGCGGCTTGAGTGCCTTAGGCCCCGGGTACATGGCCTCTGTGGCCAG  
GGCTGCCCCTTACCCGTGGACGACCAGAGCCCCCAGCATAACCCGGCTCAGGGGACACGGA  
CACAGGCCAGGGGAGTCAGAAACCTGTGACAGCGTCTCAGGCTCTTCTGAGCTGCTCCAAA  
GTCTGTATTACCTCCAGGTGCCAAGAGAGCACCCACCCTGCTTCGGACAACCCTGACATA  
ATTGCCAGCACGGCAGAGGAGGTGGCATCCACCAGCCCAGGCATCCATCATGCCACTGGGT  
GTTGTTCTTAAGAAACTGATTGATTAAAAAATTTCCCAAAGTGTCTGAAGTGTCTCTTCAA  
ATACATGTTGATCTGTGGAGTTGATTCTTTCTTCTTGGTTTTAGACAAATGTAAACAA  
AGCTCTGATCCTTAAAATTGCTATGCTGATAGAGTGGTGAGGGCTGGAAGCTTGATCAAGTC  
CTGTTTTCTTCTTGACACAGACTGATTTAAAATTTAAAAGNAAAAA

## **FIGURE 48**

MYHGMNPSNGDGFLEQQQQQQPQSPQRLLAVILWFQALCFGPAQLTGGFDDLQVCADPGI  
PENGFRTPSGGVFFEGSVARFHCQDGFKLKGGATKRLCLKHFNGTLGWIPSDNSICVQEDCRI  
PQIEDAEIHNKTYRHGEKLIITCHEGFKIRYPDLHNMVSLCRDDGTWNNLPICQGCLRPLAS  
SNGYVNISELQTSFPVGTVISYRCFPFGFKLDGSAYLECLQNLIWSSSPRCLALEAQVCPLP  
PMVSHGDFVCHPRPCERYNHGTVVEFYCDPGYSLTSDYKYITCQYGEWFPSYQVYCIKSEQT  
WPSTHETLLTTWKIVAFTATSVLLVLLLVILARMFQTKFKAHFPPRGPPRSSSSSDPDFVVVD  
GVPVMLPSYDEAVSGGLSALGPGYMASVGQGCPLVDDQSPPAYPGSGD TDTGPGESETCDS  
VSGSSELLQSLYSPRCQESTHPASDNPDIIASTAEVASTSPGIHHAHWVFLRN

**Signal sequence:**

amino acids 1-41

**Transmembrane domain:**

amino acids 325-344

**N-glycosylation site.**

amino acids 104-108, 134-138, 192-196

**Casein kinase II phosphorylation site.**

amino acids 8-12, 146-150, 252-256, 270-274, 313-317, 362-366,  
364-368, 380-384, 467-471, 468-472

**N-myristoylation site.**

amino acids 4-10, 61-67, 169-175, 203-209, 387-393, 418-424,  
478-484

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 394-405

**FIGURE 49**

CCCACGCGTCCGCTCCGCGCCCTCCCCCGCCTCCCCTGCGGTCCGTCGGTGGCCTAGAGA  
TGCTGCTGCCGCGGTTGCAGTTGTGCGGCACGCCTCTGCCCGCCAGCCCGCTCCACCGCCGT  
AGCGCCCGAGTGTGGGGGGCGCACCCGAGTCGGGCCATGAGGCCGGAACCGCGCTACAGG  
CCGTGCTGCTGGCCGTGCTGCTGGTGGGGCTGCGGGCCGCGACGGGTGCCTGCTGAGTGCC  
TCGGATTTGGACCTCAGAGGAGGGCAGCCAGTCTGCCGGGGAGGGACACAGAGGCCTTGTTA  
TAAAGTCATTTACTTCCATGATACTTCTCGAAGACTGAACTTTGAGGAAGCCAAAGAAGCCT  
GCAGGAGGGATGGAGGCCAGCTAGTCAGCATCGAGTCTGAAGATGAACAGAACTGATAGAA  
AAGTTCATTGAAAACCTCTTGCCATCTGATGGTACTTCTGGATTGGGCTCAGGAGGCGTGA  
GGAGAAACAAAGCAATAGCACAGCCTGCCAGGACCTTTATGCTTGGACTGATGGCAGCATAT  
CACAATTTAGGAACTGGTATGTGGATGAGCCGTCTGCGGCAGCGAGGTCTGCGTGGTCATG  
TACCATCAGCCATCGGCACCCGCTGGCATCGGAGGCCCTACATGTTCCAGTGGAAATGATGA  
CCGGTGCAACATGAAGAACAATTTCAATTTGCAAATATTCTGATGAGAAACCAGCAGTTCCTT  
CTAGAGAAGCTGAAGGTGAGGAAACAGAGCTGACAACACCTGTACTTCCAGAAGAAACACAG  
GAAGAAGATGCCAAAAAACATTTAAAGAAAGTAGAGAAGCTGCCTTGAATCTGGCCTACAT  
CCTAATCCCCAGCATTCCCCTTCTCCTCCTCCTTGTGGTCACCACAGTTGTATGTTGGGTTT  
GGATCTGTAGAAAAAGAAAACGGGAGCAGCCAGACCCTAGCACAAAGAAGCAACACACCATC  
TGGCCCTCTCCTCACCAGGAAACAGCCCGGACCTAGAGGTCTACAATGTCATAAGAAAAACA  
AAGCGAAGCTGACTTAGCTGAGACCCGGCCAGACCTGAAGAATATTTCAATCCGAGTGTGTT  
CGGGAGAAGCCACTCCCGATGACATGTCTTGTGACTATGACAACATGGCTGTGAACCCATCA  
GAAAGTGGGTTTGTGACTCTGGTGGAGCTGGAGAGTGGATTTGTGACCAATGACATTTATGA  
GTTCTCCCCAGACCAAATGGGGAGGAGTAAGGAGTCTGGATGGGTGGAAAATGAAATATATG  
GTTATTAGGACATATAAAAACTGAAACTGACAACAATGGAAAAGAAATGATAAGCAAAATC  
CTCTTATTTTCTATAAGGAAAATACACAGAAGGTCTATGAACAAGCTTAGATCAGGTCCTGT  
GGATGAGCATGTGGTCCCCACGACCTCCTGTTGGACCCCCACGTTTTGGCTGTATCCTTTAT  
CCCAGCCAGTCATCCAGCTCGACCTTATGAGAAGGTACCTTGCCCAGGTCTGGCACATAGTA  
GAGTCTCAATAAATGFCACTTGGTTGGTTGTATCTAACTTTTAAGGGACAGAGCTTTACCTG  
GCAGTGATAAAGATGGGCTGTGGAGCTTGGAAAACCACCTCTGTTTTCTTGCTCTATACAG  
CAGCACATATTATCATAACAGACAGAAAATCCAGAATCTTTTCAAAGCCCACATATGGTAGCACAG  
GTTGGCCTGTGCATCGGCAATTCTCATATCTGTTTTTTCAAAGAATAAAATCAAATAAAGA  
GCAGGAAAAAAAAA

**FIGURE 50**

MRPGTALQAVLLAVLLVGLRAATGRLLSASDLDLRGGQPVCRGGTQRPCYKVIYFHDTSRRL  
NFEEAKEACRRDGGQLVSI ESEDEQKLI EKFIENLLPSDGD F W I G L R R R E E K Q S N S T A C Q D L  
YAWTDGSISQFRN W Y V D E P S C G S E V C V V M Y H Q P S A P A G I G G P Y M F Q W N D D R C N M K N N F I C K Y  
SDEKPAVPSREAEGETELTTPVLPEETQEEDAKKTFKESREAAALNLAYILIPS I P L L L L L V  
VTTVVCWWICRKRKREQDPSTKKQHTIWPSPHQGNSPDLEVYNVIRKQSEADLAETRPDL  
KNISFRVCSGEATPDDMSCDYDNMAVNPSESGFVTLVSVESGFVTNDIYEFSPDQMGRSKES  
GWVENEIYGY

**Signal sequence:**

amino acids 1-21

**Transmembrane domain:**

amino acids 235-254

**N-glycosylation site.**

amino acids 117-121, 312-316

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 296-300

**Casein kinase II phosphorylation site.**

amino acids 28-32, 30-34, 83-87, 100-104, 214-218, 222-226,  
299-303, 306-310, 323-327

**N-myristoylation site.**

amino acids 18-24, 37-43, 76-82, 146-152

**FIGURE 51**

GGGGTCTCCCTCAGGGCCGGGAGGCACAGCGTCCCTGCTTGCTGAAGGGCTGGATGTACGC  
ATCCGCAGGTTCCCGCGGACTTGGGGGCGCCCGCTGAGCCCCGGCGCCCGCAGAAGACTTGT  
GTTTGCCTCCTGCAGCCTCAACCCGGAGGGCAGCGAGGGCTACCACCATGATCACTGGTGT  
GTTCAGCATGCGCTTGTGGACCCAGTGGGCGTCCCTGACCTCGCTGGCGTACTGCCTGCACC  
AGCGGCGGGTGGCCCTGGCCGAGCTGCAGGAGGCCGATGGCCAGTGTCCGGTCGACCGCAGC  
CTGCTGAAGTTGAAAATGGTGCAGGTCGTGTTTTCGACACGGGGCTCGGAGTCCTCTCAAGCC  
GCTCCCGCTGGAGGAGCAGGTAGAGTGAACCCCCAGCTATTAGAGGTCCCACCCCAAATC  
AGTTTGATTACACAGTCACCAATCTAGCTGGTGGTCCGAAACCATATCTCCTTACGACTCT  
CAATACCATGAGACCACCCTGAAGGGGGCATGTTTGTGGGCAGCTGACCAAGGTGGGCAT  
GCAGCAAATGTTTGCCTTGGGAGAGAGACTGAGGAAGAACTATGTGGAAGACATTCCCTTTC  
TTTACCAACCTTCAACCCACAGGAGGTCTTTATTCGTTCCACTAACATTTTTTCGGAATCTG  
GAGTCCACCCGTTGTTTGTGGCTGGGCTTTTTCCAGTGTGAGAAAGAAGGACCCATCATCAT  
CCCACTGATGAAGCAGATTCAGAAGTCTTGTATCCCAACTACCAAAGCTGCTGGAGCCTGA  
GGCAGAGAACCAGAGGCCGGAGGCAGACTGCCTCTTTACAGCCAGGAATCTCAGAGGATTTG  
AAAAAGGTGAAGGACAGGATGGGCATTGACAGTAGTGATAAAGTGGACTTCTTCATCCTCCT  
GGACAACGTGGCTGCCGAGCAGGCACACAACCTCCCAAGCTGCCCCATGCTGAAGAGATTTG  
CACGGATGATCGAACAGAGAGCTGTGGACACATCCTTGTACATACTGCCCAAGGAAGACAGG  
GAAAGTCTTCAGATGGCAGTAGGCCCATTCCTCCACATCCTAGAGAGCAACCTGCTGAAAGC  
CATGGACTCTGCCACTGCCCCCGACAAGATCAGAAAGCTGTATCTCTATGCGGCTCATGATG  
TGACCTTCATACCGCTCTTAATGACCCTGGGGATTTTTGACCACAAATGGCCACCGTTTGT  
GTTGACCTGACCATGGAACCTTACCAGCACCTGGAATCTAAGGAGTGGTTTGTGCAGCTCTA  
TTACCACGGGAAGGAGCAGGTGCCGAGAGGTTGCCCTGATGGGCTCTGCCCGCTGGACATGT  
TCTTGAATGCCATGTGAGTTTATACCTTAAGCCCAGAAAAATACCATGCACTCTGCTCTCAA  
ACTCAGGTGATGGAAGTTGGAATGAAGAGTAACTGATTTATAAAAGCAGGATGTGTTGATT  
TTAAAATAAAGTGCCTTTATACAATG

**FIGURE 52**

MITGVFSMRLWTPVGVLTSLAYCLHQRRVALAELQEADGQCPVDRSLLKLMVQVVFRHGAR  
SPLKPLPLEEQVEWNPQLLEVPPQTQFDYTVTNLAGGPKPYSPYDSQYHETTLKGGMFAGQL  
TKVGMQQMFALGERLRKKNYVEDIPFLSPTFNPQEVFIRSTNIFRNLESTRCLLAGLFQCQKE  
GPIIIHTDEADSEVLYPNYQSCWSLRQRTRGRRQTASLQPGISEDLKKVKDRMGIDSSDKVD  
FFILLDNVAEQAHNLPSCPMLKRFARMIEQRAVDTSLYILPKEDRESLQMAVGPFLHILES  
NLLKAMDSATAPDKIRKLYLYAAHDVTFIPLLMTLGI FDHKWPPFAVDLTMELYQHLESKEW  
FVQLYYHGKEQVPRGCPDGLCPDMLNAMS VYTLSP EKYHALCSQTQVMEVGNEE

**Signal sequence:**

amino acids 1-23

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 218-222

**Casein kinase II phosphorylation site.**

amino acids 87-91, 104-108, 320-324

**Tyrosine kinase phosphorylation site.**

amino acids 280-288

**N-myristoylation site.**

amino acids 15-21, 117-123, 118-124, 179-185, 240-246, 387-393

**Amidation site.**

amino acids 216-220

**Leucine zipper pattern.**

amino acids 10-32

**Histidine acid phosphatases phosphohistidine signature.**

amino acids 50-65

**FIGURE 53**

CTCCTCTTAACATACTTGCAGCTAAAATAAATATTGCTGCTTGGGGACCTCCTTCTAGCCT  
TAAATTTTCAGCTCATCACCTTCACCTGCCTTGGTCAATGGCTCTGCTATTCTCCTTGATCCTT  
GCCATTTGCACCAGACCTGGATTCCTAGCGTCTCCATCTGGAGTGCGGCTGGTGGGGGGCCT  
CCACCGCTGTGAAGGGCGGGTGGAGGTGGAACAGAAAGGCCAGTGGGGCACCGTGTGTGATG  
ACGGCTGGGACATTAAGGACGTGGCTGTGTTGTGCCGGGAGCTGGGCTGTGGAGCTGCCAGC  
GGAACCCCTAGTGGTATTTTGTATGAGCCACCAGCAGAAAAAGAGCAAAAAGGTCCTCATCCA  
ATCAGTCAGTTGCACAGGAACAGAAGATACATTGGCTCAGTGTGAGCAAGAAGAAGTTTATG  
ATTGTTACATGATGAAGATGCTGGGGCATCGTGTGAGAACCAGAGAGCTCTTCTCCTCCA  
GTCCCAGAGGGTGTGAGGCTGGCTGACGGCCCTGGGCATTGCAAGGGACCGGTGGAAGTGAA  
GCACCAGAACCAGTGGTATACCGTGTGCCAGACAGGCTGGAGCCTCCGGGCCGCAAAGGTGG  
TGTGCCGGCAGCTGGGATGTGGGAGGGCTGTACTGACTCAAAAACGCTGCAACAAGCATGCC  
TATGGCCGAAAACCCATCTGGCTGAGCCAGATGTCATGCTCAGGACGAGAAGCAACCCTTCA  
GGATTGCCCTTCTGGGCCTTGGGGGAAGAACACCTGCAACCATGATGAAGACACGTGGGTG  
AATGTGAAGATCCCTTTGACTTGAGACTAGTAGGAGGAGACAACCTCTGCTCTGGGCGACTG  
GAGGTGCTGCACAAGGGCGTATGGGGCTCTGTCTGTGATGACAACCTGGGGAGAAAAGGAGGA  
CCAGGTGGTATGCAAGCAACTGGGCTGTGGGAAGTCCCTCTCTCCCTCCTTCAGAGACCGGA  
AATGCTATGGCCCTGGGGTTGGCCGCATCTGGCTGGATAAATGTTGCTCAGGGGAGGAG  
CAGTCCCTGGAGCAGTGCCAGCACAGATTTTGGGGGTTTTCAGACTGCACCCACCAGGAAGA  
TGTGGCTGTCTGCTCAGTGTAGGTGGGCATCATCTAATCTGTTGAGTGCCTGAATAGAA  
GAAAAACACAGAAGAAGGGAGCATTTACTGTCTACATGACTGCATGGGATGAACACTGATCT  
TCTTCTGCCCTTGGACTGGGACTTATACTTGGTGCCCTGATTCTCAGGCCTTCAGAGTTGG  
ATCAGAACTTACAACATCAGGTCTAGTTCTCAGGCCATCAGACATAGTTTGGAACTACATCA  
CCACCTTTCCTATGTCTCCACATTGCACACAGCAGATTCCCAGCCTCCATAATTGTGTGTAT  
CAACTACTTAAATACATTCTCACACACACACACACACACACACACACACACACACATA  
CACCATTTGTCCTGTTTCTCTGAAGAACTCTGACAAAATACAGATTTTGGTACTGAAAGAGA  
TTCTAGAGGAACCGAATTTAAGGATAAAATTTTCTGAATTGGTTATGGGGTTTCTGAAATTG  
GCTCTATAATCTAATTAGATATAAAATTTCTGGTAACTTTATTTACAATAATAAAGATAGCAC  
TATGTGTTCAA

## **FIGURE 54**

MALLFSLILAICTRPGFLASPSGVRLVGGGLHRCEGRVEVEQKGQWGTVCDDGWDIKDVAVLC  
RELGCCAASGTPSGILYEPPAEKEQKVLIQSVSCTGTEDTLAQCEQEVEVYDCSHDEDAGASC  
ENPESSFSPVPEGVRLADGPGHCKGRVEVKHQWYTVQCQTGWSLRAAKVVCRLGCGRAVL  
TQKRCNKHAYGRKPIWLSQMSCSGREATLQDCPSGPWGKNTCNHDEDTWVECEDPFDLRLVG  
GDNLCSGRLEVLHKGWVGSVCCDDNWGEKEDQVVCKQLGCGKSLSPSFRDRKCYGPGVGRIWL  
DNVRCSGEEQSLEQCQHRFWGFHDCTHQEDVAVICSV

**Signal sequence:**

amino acids 1-15

**Casein kinase II phosphorylation site.**

amino acids 47-51, 97-101, 115-119, 209-213, 214-218, 234-238,  
267-271, 294-298, 316-320, 336-340

**N-myristoylation site.**

amino acids 29-35, 43-49, 66-72, 68-74, 72-78, 98-104, 137-143,  
180-186, 263-269, 286-292

**Amidation site.**

amino acids 196-200

**Speract receptor repeated domain signature.**

amino acids 29-67, 249-287



**FIGURE 55**

ACTGCACTCGGTTCTATCGATTGAATCCCCGGGGATCCTCTAGAGATCCCTCGACCTCGAC  
CCACGCGTCCGCGGACGCGTGGGCGGACGCGTGGGCCGGCTACCAGGAAGAGTCTGCCGAAG  
GTGAAGGCCATGGACTTCATCACCTCCACAGCCATCCTGCCCTGCTGTTGCGCTGCCTGGG  
CGTCTTCGGCCTCTTCCGGCTGCTGCAGTGGGTGCGCGGGAAGGCCTACCTGCGGAATGCTG  
TGGTGGTGATCACAGGCGCCACCTCAGGGCTGGGCAAAGAATGTGCAAAAGTCTTCTATGCT  
GCGGGTGCTAAACTGGTGCTCTGTGGCCGGAATGGTGGGGCCCTAGAAGAGCTCATCAGAGA  
ACTTACCGTCTTCATGCCACCAAGGTGCAGACACACAAGCCTTACTTGGTGACCTTCGACC  
TCACAGACTCTGGGGCCATAGTTGCAGCAGCAGCTGAGATCCTGCAGTGCTTTGGCTATGTC  
GACATACTTGTCAACAATGCTGGGATCAGCTACCGTGGTACCATCATGGACACCACAGTGGA  
TGTGGACAAGAGGGTTCATGGAGACAACTACTTTGGCCCAGTTGCTCTAACGAAAGCACTCC  
TGCCCTCCATGATCAAGAGGAGGCAAGGCCACATTGTCGCCATCAGCAGCATCCAGGGCAAG  
ATGAGCATTCTTTTCGATCAGCATATGCAGCCTCCAAGCACGCAACCCAGGCTTTCTTTGA  
CTGTCTGCGTGCCGAGATGGAACAGTATGAAATGAGGTGACCGTCATCAGCCCCGGCTACA  
TCCACACCAACCTCTCTGTAAATGCCATCACCGGGATGGATCTAGGTATGGAGTTATGGAC  
ACCACCACAGCCCAGGGCCGAAGCCCTGTGGAGGTGGCCCAGGATGTTCTTGCTGCTGTGGG  
GAAGAAGAAGAAAGATGTGATCCTGGCTGACTTACTGCCTTCCTTGGCTGTTTATCTTCGAA  
CTCTGGCTCCTGGGCTCTTCTTCAGCCTCATGGCCTCCAGGGCCAGAAAAGAGCGGAAATCC  
AAGAACTCCTAGTACTCTGACCAGCCAGGGCCAGGGCAGAGAAGCAGCACTCTTAGGCTTGC  
TTACTCTACAAGGGACAGTTGCATTTGTTGAGACTTTAATGGAGATTTGTCTCACAAGTGGG  
AAAGACTGAAGAAACACATCTCGTGCAGATCTGCTGGCAGAGGACAATCAAAAACGACAACA  
AGCTTCTTCCCAGGGTGAGGGGAAACACTTAAGGAATAAATATGGAGCTGGGGTTTAACT  
AAAACTAGAAATAAACATCTCAAACAGTAAAAAAAAAAAAAAAAAGGGCGCCGCGACTCTAG  
AGTCGACCTGCAGAAGCTTGGCCGCCATGGCCCAACTTGTTTATTGCAGCTTATAATGGTTAC

## **FIGURE 56**

MDFITSTAILPLLFGCLGVFGLFRLLQWVRGKAYLRNAVVVITGATSGLGKECAKVFYAAGA  
KLVLCGRNGGALEELIRELTASHATKVQTHKPYLVTFDLTDSGAIVAAAAEILQCFGYVDIL  
VNNAGISYRGTIMDTTVDVVKRVMETNYFGPVALTKALLPSMIKRRQGHIVAIISSIQKMSI  
PFRSAYAASKHATQAFFDCLRAEMEQYEIEVTVISPGYIHTNLSVNAITADGSRYGVMDTTT  
AQGRSPVEVAQDVLAAVGGKKKDVILADLLPSLAVYLRTLAPGLFFSLMASRARKERKSKNS

**Signal sequence:**

amino acids 1-21

**Transmembrane domain:**

amino acids 104-120, 278-292

**N-glycosylation site.**

amino acids 228-232

**Glycosaminoglycan attachment site.**

amino acids 47-51

**Casein kinase II phosphorylation site.**

amino acids 135-139, 139-143, 253-257

**Tyrosine kinase phosphorylation site.**

amino acids 145-153, 146-153

**N-myristoylation site.**

amino acids 44-50, 105-111, 238-244, 242-248, 291-297

**Amidation site.**

amino acids 265-269

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 6-17



**FIGURE 58**

MKFLLDILLLLPLLIIVCSLESFVKLFIPKRRKSVTGEIVLITGAGHGIGRLTAYEFAKLKSK  
LVLWDINKHGLEETAACKCKGLGAKVHTFVVDCSNREDIYSSAKKVKAEIGDVSILVNNAGVV  
YTSDLFATQDPQIEKTFEVNVLAHFWTTKAFLPAMTKNNHGHI VTVASAAGHVSVPELLAYC  
SSKFAAVGFHKTLTDELAALQITGVKTTCLCPNFVNTGFIKNPSTSLGPTLEPEEVNRLMH  
GILTEQKMIFIPSSIAFLTTLERILPERFLAVLKRKISVKFDAVIGYKMKAO

**Signal sequence:**

amino acids 1-19

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 30-34, 283-287

**Casein kinase II phosphorylation site.**

amino acids 52-56, 95-99, 198-202, 267-271

**N-myristoylation site.**

amino acids 43-49, 72-78, 122-128, 210-216

**FIGURE 59**

CCCACGCGTCCGCGGACGCGTGGGTGCGACTAGTTCCTAGATCGCGAGCGGCCGCCGCGGCTC  
AGGGAGGAGCACCGACTGCGCCGCACCCTGAGAGATGGTTGGTGCCATGTGGAAGGTGATTG  
TTTCGCTGGTCCTGTTGATGCCTGGCCCCCTGTGATGGGCTGTTTCGCTCCCTATACAGAAGT  
GTTTCCATGCCACCTAAGGGAGACTCAGGACAGCCATTATTTCTCACCCCTTACATTGAAGC  
TGGGAAGATCCAAAAAGGAAGAGAATTGAGTTTGGTCCGCCCTTTCCAGGACTGAACATGA  
AGAGTTATGCCGGCTTCCCTCACCGTGAATAAGACTTACAACAGCAACCTCTTCTTGTTTC  
TTCCAGCTCAGATACAGCCAGAAGATGCCCCAGTAGTTCCTCTGGCTACAGGGTGGGCCGGG  
AGGTTTCATCCATGTTTGGACTCTTTGTGGAACATGGGCCTTATGTTGTCAAGTAACATGA  
CCTTGCGTGACAGAGACTTCCCCTGGACCACAACGCTCTCCATGCTTTACATTGACAATCCA  
GTGGGCACAGGCTTTCAGTTTTACTGATGATACCCACGGATATGCAGTCAATGAGGACGATGT  
AGCACGGGATTTATACAGTGCACATAATTCAGTTTTTCCAGATATTTCTGAATATAAAAAATA  
ATGACTTTTATGTCACTGGGGAGTCTTATGCAGGAAATATGTGCCAGCCATTGCACACCTC  
ATCCATTCCTCAACCCTGTGAGAGAGGTGAAGATCAACCTGAACCGAATTGCTATTGGAGA  
TGGATATTCTGATCCCGAATCAATTATAGGGGGCTATGCAGAATTCCTGTACCAATTGGCT  
TGTTGGATGAGAAGCAAAAAAGTACTTCCAGAAGCAGTGCCATGAATGCATAGAACACATC  
AGGAAGCAGAAGTGGTTTGGAGCCCTTTGAAATACTGGATAAACTACTAGATGGCGACTTAAC  
AAGTGATCCTTCTTACTTCCAGAATGTTACAGGATGTAGTAATTAATAACTTTTTTGCGGT  
GCACGGAACCTGAGGATCAGCTTTACTATGTGAAATTTTTGTCACTCCAGAGGTGAGACAA  
GCCATCCACGTGGGGAATCAGACTTTTAAATGATGGAACATAGTTGAAAAGTACTTGCAGAGA  
AGATACAGTACAGTCACTTAAGCCATGGTTAACTGAAATCATGAATAATTATAAGGTTCTGA  
TCTACAATGGCCAACCTGGACATCATCGTGGCAGCTGCCCTGACAGAGCGCTCCTTGATGGGC  
ATGGACTGGAAAAGGATCCAGGAATACAAGAAGGCAGAAAAAAAAGTTTGGAAAGATCTTTAA  
ATCTGACAGTGAAGTGGCTGGTTACATCCGGCAAGCGGGTGAATCCATCAGGTAATTATTC  
GAGGTGGAGGACATATTTTACCCTATGACCAGCCTCTGAGAGCTTTTGACATGATTAATCGA  
TTCATTTATGGAAAAGGATGGGATCCTTATGTTGGATAAAACTACCTTCCCAAAAGAGAACAT  
CAGAGGTTTTTCATGCTGAAAAGAAAATCGTAAAAACAGAAAATGTCATAGGAATAAAAAAA  
TTATCTTTTCATATCTGCAAGATTTTTTTTCATCAATAAAAAATTATCCTTGAAACAAGTGAGC  
TTTTGTTTTTGGGGGAGATGTTTACTACAAAATTAACATGAGTACATGAGTAAGAATTACA  
TTATTTAACTTAAAGGATGAAAGGTATGGATGATGTGACACTGAGACAAGATGTATAAAATGA  
AATTTTGGGTCTTGAATAGGAAGTTTTAATTTCTTCTAAGAGTAAGTGAAGAAGTGCAGTTG  
TAACAAACAAAGCTGTAACATCTTTTTCTGCCAATAACAGAAGTTTGGCATGCCGTGAAGGT  
GTTTGGAAATATTATTGGATAAGAATAGCTCAATTATCCCAAATAAATGGATGAAGCTATAA  
TAGTTTTGGGGAAAAGATTCCTCAATGTATAAAGTCTTAGAACAAAAGAATTCTTTGAAATA  
AAAAATATTATATATAAAAGTAAAAAAA

**FIGURE 60**

MVGAMWKVIVSLVLLMPGPCDGLFRSLYRSVSMPPKGDGQPLFLTPYIEAGKIQKGRELSL  
VGPFPGLNMKSYAGFLTVNKTYNLFFWFPPAQIQPEDAPVVLWLQGGPGGSSMFGLFVEH  
GPYVVTSNMTLRDRDFPWTTLTSMLYIDNPVGTGFSFTDDTHGYAVNEDDVARLDLYSALIQF  
FQIFPEYKNNDFYVTGESYAGKYVPAIAHLIHSNPNVREVKINLNGIAIGDGYSDPESIIGG  
YAEFLYQIGLLDEKQKKYFQKQCHECIEHIRKQNWFEAFEILDKLLDGLTSDPSYFQNVGTG  
CSNYYNFLRCTEPEDQLYVVKFSLPEVRQAIHVGNQTFNDGTIVEKYLREDTVQSVKPWLT  
EIMNNYKVLINYGQLDIIVAAALTERSLMGMDWKGSGEYKKAEEKVWKIFKSDSEVAGYIRQ  
AGDFHQVIIRGGGHILPYDQPLRAFDMINRFIYGKGDWPYVG

**Signal sequence:**

amino acids 1-22

**N-glycosylation site.**

amino acids 81-85, 132-136, 307-311, 346-350

**Casein kinase II phosphorylation site.**

amino acids 134-138, 160-164, 240-244, 321-325, 334-338, 348-352,  
353-357, 424-428

**Tyrosine kinase phosphorylation site.**

amino acids 423-432

**N-myristoylation site.**

amino acids 22-28, 110-116, 156-162, 232-238

**Serine carboxypeptidases, serine active site.**

amino acids 200-208

**Crystallins beta and gamma 'Greek key' motif signature.**

amino acids 375-391

**FIGURE 61**

CGAGGGCTTTTCCGGCTCCGGAATGGCACATGTGGGAATCCCAGTCTTGTTGGCTACAACAT  
TTTTCCCTTTCTAACAAAGTTCTAACAGCTGTTCTAACAGCTAGTGATCAGGGGTTCTTCTT  
GCTGGAGAAGAAAGGGCTGAGGGCAGAGCAGGGCACTCTCACTCAGGGTGACCAGCTCCTTG  
CCTCTCTGTGGATAACAGAGCATGAGAAAGTGAAGAGATGCAGCGGAGTGAGGTGATGGAAG  
TCTAAAATAGGAAGGAATTTTGTGTGCAATATCAGACTCTGGGAGCAGTTGACCTGGAGAGC  
CTGGGGGAGGGCCTGCCTAACAAAGCTTTCAAAAACAGGAGCGACTTCCACTGGGCTGGGAT  
AAGACGTGCCGGTAGGATAGGGAAGACTGGGTTTAGTCCTAATATCAAATTGACTGGCTGGG  
TGAACTTCAACAGCCTTTTAACTCTCTGGGAGATGAAAACGATGGCTTAAGGGGCCAGAAA  
TAGAGATGCTTTGTAAAATAAAATTTTAAAAAAGCAAGTATTTTATAGCATAAAGGCTAGA  
GACCAAATAGATAACAGGATTCCCTGAACATTCCTAAGAGGGAGAAAGTATGTTAAAAATA  
GAAAAACCAAATGCAGAAGGAGGAGACTCACAGAGCTAAACCAGGATGGGGACCCTGGGTG  
AGGCCAGCCTCTTTGCTCCTCCCGAAATATTTTTTGGTCTGACCACTCTGCCTTGTGT  
GCAGAATCATGTGAGGGCCAACCGGGGAAGGTGGAGCAGATGAGCACACACAGGAGCCGCT  
CCTCACCGCCGCCCTCTCAGCATGGAACAGAGGCAGCCCTGGCCCCGGGCCCTGGAGGTGG  
ACAGCCGCTCTGTGGTCTGCTCTCAGTGGTCTGGGTGCTGCTGGCCCCCAGCAGCCGGC  
ATGCCTCAGTTCAGCACCTTCCACTCTGAGAATCGTGACTGGACCTTCAACCACTTGACCGT  
CCACCAAGGGAGCGGGGCCGTCTATGTGGGGGCCATCAACCGGGTCTATAAGCTGACAGGCA  
ACCTGACCATCCAGGTGGCTCATAAGACAGGGCCAGAAGAGGACAACAAGTCTCGTTACCCG  
CCCCTCATCGTGACGCCCTGCAGCGAAGTGTCTACCCCTCACCAACAATGTCAACAAGCTGCT  
CATCATTGACTACTCTEAGAACCGCCTGCTGGCCTGTGGGAGCCTCTACCAGGGGGTCTGCA  
AGCTGCTGCGGCTGGATGACCTCTTCATCCTGGTGGAGCCATCCCACAAGAAGGAGCACTAC  
CTGTCCAGTGTCAACAAGACGGGCACCATGTACGGGGTGATTGTGCGCTCTGAGGGTGAGGA  
TGGCAAGCTCTTCATCGGCACGGCTGTGGATGGGAAGCAGGATTACTTCCCAGCCCTGTCCA  
GCCGGAAGCTGCCCCGAGACCCTGAGTCTCAGCCATGCTCGACTATGAGCTACACAGCGAT  
TTTGTCTCCTCTCTCATCAAGATCCCTTTCAGACACCCTGGCCCTGGTCTCCCACTTTGACAT  
CTTCTACATCTACGGCTTTGCTAGTGGGGCTTTGTCTACTTTCTCACTGTCCAGCCCGAGA  
CCCCTGAGGGTGTGGCCATCAACTCCGCTGGAGACCTCTTCTACACCTCACGCATCGTGCGG  
CTCTGCAAGGATGACCCCAAGTTCCACTCATAACGFTGTCCTGCCCCTCGGCTGCACCCGGG  
CGGGGTGGAATACCGCCTCCTGCAGGTGCTTACCTGGCCAAGCCTGGGGACTCACTGGCCC  
AGGCCTTCAATATCACCAAGCCAGGACGATGTACTCTTTGCCATCTTCTCCAAAGGGCAGAAG  
CAGTATCACCAACCCGCGGATGACTCTGCCCTGTGTGCCCTTCCCTATCCGGGCCATCAACTT  
GCAGATCAAGGAGCGCCTGCAGTCTGCTACCAGGGCGAGGGCAACCTGGAGCTCAACTGGC  
TGCTGGGGAAGGACGTCCAGTGCACGAAGGCGCCTGTCCCATCGATGATAACTTCTGTGGA  
CTGGACATCAACCAGCCCCCTGGGAGGCTCAACTCCAGTGGAGGGCCTGACCCTGTACACCAC  
CAGCAGGGACCGCATGACCTCTGTGGCCTCCTACGTTTACAACGGCTACAGCGTGGTTTTTG  
TGGGGACTAAGAGTGGCAAGCTGAAAAAGGTAAGAGTCTATGAGTTTCAAGTCTCCAATGCC  
ATTACCTCTCAGCAAAGAGTCCCTCTTGGAAAGGTAGCTATTGGTGGAGATTTAACTATAG  
GCAACTTTATTTTCTTGGGGAACAAAGGTGGAATGGGGAGGTAAGAAGGGGTTAATTTTGTG  
ACTTAGCTTCTAGCTACTTCTCCAGCCATCAGTCACTGGGTATGTAAGGAATGCAAGCGTA  
TTTTCAATATTTCCCAAACCTTAAAGAAAAAATTTAAGAAGGTACATCTGCAAAAGCAA

**FIGURE 62**

MGTLGQASLFAPPNGYFWSHDHSAFCFAESCEGQPGKVEQMSTHRSRLLTAAPLSMEQRQPWP  
RALEVDSRSVLLSVVWVLLAPPAAGMPQFSTFHSENRDWTFNHLTVHQGTGAVYVGAINRV  
YKLTGNLTIQVAHKTGPEEDNKSRYPLIVQPCSEVLTLTNNVNKLLI IDYSENRLACGSL  
YQGVCKLLRLDDLFILVEPSHKKEHYLSSVNKTGTMYGVIVRSEGEDGKLFIGTAVDGKQDY  
FPTLSSRKLPRDPESSAMLDYELHSDVSSLIKIPSDTLALVSHFDIFYIYGFASGGFVYFL  
TVQPETPEGVAINSAGDLFYTSRIVRLCKDDPKFHSYVSLPFGCTRAGVEYRLLQAAYLAKP  
GDSLAAQAFNITSQDDVLFAlFSKGQKQYHHPDSDALCAFPiRAiNLQIKERLQSCYQEGEN  
LELNWLLGKDVQCTKAPVPIDDNFCGLDINQPLGGSTPVEGLTLYTTSRDRMTSVASYVYNG  
YSVVFVGTGKSGKLLKVRVYEFRCsNAIHLLSKEsLLEGSYWWRFNYRQLYFLGEQR

**Signal sequence:**

amino acids 1-32

**Transmembrane domain:**

amino acids 71-87

**N-glycosylation site.**

amino acids 130-134, 145-149, 217-221, 381-385

**Casein kinase II phosphorylation site.**

amino acids 139-143, 229-233, 240-244, 291-295, 324-328, 383-387,  
384-388, 471-475, 481-485, 530-534

**N-myristoylation site.**

amino acids 220-226, 319-325, 353-359, 460-466, 503-509



**FIGURE 63**

AGGCTCCCGCGCGGGCTGAGTGGGACTGGAGTGGGAACCCGGGTCCCGCGCTTAGAGAACACGCGATGACCA  
CGTGGAGCCTCCGGCGGAGGCCGGCCCGCACGCTGGGACTCCGCTGCTGGTGTCTTGGGCTTCTGGTGCTCC  
GCAGGCTGGACTGGAGCACCCTGGTCCCTCTGCGGCTCCGCCATCGACAGCTGGGGCTGCAGGCCAAGGGCTGGA  
ACTTCATGCTGGAGATTCCACCTTCTGGATCTTCGGGGCTCCATCCACTATTTCCGTGTGCCAGGGAGTACT  
GGAGGGACCGCTGCTGAAGATGAAGGCCTGTGGCTTGAACACCCTCACCACCTATGTTCCGTGGAACCTGCATG  
AGCCAGAAAGAGGCCAAATTTGACTTCTCTGGGAACCTGGACCTGGAGGCCTTCGTCCGTGATGGCCGAGAGATCG  
GGCTGTGGGTGATTCTGCGTCCAGGCCCTACATCTGCAGTGAGATGGACCTCGGGGGCTTGCCAGCTGGCTAC  
TCCAAGACCCTGGCATGAGGCTGAGGACAACCTTACAAGGGCTTCAACGAAGCAGTGGACCTTTATTTTGACCACC  
TGATGTCAGGGTGGTGCCACTCCAGTACAAGCGTGGGGACCTATCATTTGCCGTGCAGGTGGAGAATGAATATG  
GTTCTATAATAAAGACCCCGCATACATGCCCTACGTCAAGAAGGCCTGGAGGACCGTGGCATGTGGAACTGC  
TCCTGACTTCAGACAACAAGGATGGGCTGAGCAAGGGGATTGTCAGGGAGTCTTGGCCACCATCAACTTGCAGT  
CAACACACGAGCTGCAGCTACTGACCACCTTTCTTCAACGTCAGGGGACTCAGCCAAAGATGGTGAATGGAGT  
ACTGGACGGGGTGGTTTACTCGTGGGGAGGCCCTCACAATATCTTGGATTCTTCTGAGGTTTTGAAAACCGTGT  
CTGCCATTGTGGACCGCGGCTCCTCCATCAACCTCTACATGTTCCACGGAGGCCACCAACTTTGGCTTCATGAATG  
GAGCATGCATTTCCATGACTACAAGTCAGATGTCACAGCTATGACTATGATGCTGTGCTGACAGAAGCCGGCG  
ATTACACGGCCAAGTACATGAAGCTTCGAGACTTCTTCCGCTCCATCTCAGGCATCCCTCTCCCTCCCCACCTG  
ACCTTCTTCCAAAGATGCCGTATGAGCCCTTAACGCCAGTCTGTACCTGTCTCTGTGGACGCCCTCAAGTACC  
TGGGGGAGCCAAATCAAGTCTGAAAAGCCCATCAACATGGAGAACCTGCCAGTCAATGGGGGAAATGGACAGTCT  
TCGGGTACATTTCTATGAGACCAGCATCACCTCGTCTGGCATCCTCAGTGGCCACGTGCATGATCGGGGGCAGG  
TGTTTGTGAACACAGTATCCATAGGATTTCTGGACTACAAGACAACGAAGATTGCTGTCCCCCTGATCCAGGGTT  
ACACCGTCTGAGGATCTTGGTGGAGAATCGTGGCGGAGTCAACTATGGGGAGAATATTGATGACCAGCGCAAAG  
GCTTAATGGAAATCTCTATCTGAATGATTCAACCCCTGAAAACTTCAAGATCTATAGCCTGGATATGAAGAAGA  
GCTTCTTTAGAGGTTCCGGCTGGACAATGGNGTTCCCTCCAGAAAACACCCACATTACCTGTCTTCTTCTGG  
GTAGCTTGTCCATCAGCTCCACGCCCTTGTGACACCTTTCTGAAGCTGGAGGGCTGGGAGAAGGGGGTGTATTCA  
TCAATGGCCAGAACCCTGGACGTTACTGGAACATTTGGACCCAGAAAGCGCTTTACCTCCAGGTCCCTGGTTGA  
GCAGCGGAATCAACCAGGTCAFCGTTTTTGGAGAGACGATGGCGGGCCCTGCATTAACAATTCACGGAAACCCCTC  
ACCTGGGCAGGAACAGTACATTAAGTGAAGCGGTGGCACCCCTCCTGCTGGTGCCAGTGGGAGACTGCCGCCCTC  
CTCTTGACCTGAAGCCTGGTGGCTGCTGCCCCACCCCTCACTGCAAAGCATCTCCTTAAGTAGCAACCTCAGGG  
ACTGGGGGCTACAGTCTGCCCTGTCTCAGCTCAAAACCCCTAAGCCTGCAGGGAAAGGTGGGATGGCTCTGGGCC  
TGGCTTGTGATGATGGCTTCTTACAGCCCTGCTCTTGTGCCGAGGCTGTGGGCTGTCTCTAGGGTGGGAGC  
AGCTAATCAGATCGCCAGCCTTTGGCCCTCAGAAAAGTGTGAAACGTGCCCTTGCACCGGACGTCACAGCCC  
TGCGAGCATCTGCTGGACTCAGGCGTGTCTTTGCTGGTTCTTGGGAGGCTTGGCCACATCCCTCATGGCCCAT  
TTTATCCCGAAATCCTGGGTGTGTCAACAGTGTAGAGGGTGGGGAAGGGGTGTCTCACCTGAGCTGACTTTGTT  
CTTCTTCAACACCTTCTGAGCCTTCTTGGGATTTCTGGAAGGAACCTGGCGTGAGAAACATGTGACTTCCCTT  
TCCCTTCCCCTCGCTGCTTCCACAGGGTGACAGGCTGGGCTGGAGAAAACAGAAATCCTCACCTGCGTCTTCC  
CAAGTTAGCAGGTGTCTCTGGTGTTCAGTGAGGAGGACATGTGAGTCTTGGCAGAAGCCATGGCCCATGTCTGCA  
CATCCAGGGAGGAGGACAGAAGGCCAGCTCACATGTGAGTCTTGGCAGAAGCCATGGCCCATGTCTGCACATCC  
AGGGAGGAGGACAGAAGGCCAGCTCACATGTGAGTCTTGGCAGAAGCCATGGCCCATGTCTGCACATCCAGGGA  
GGAGGACAGAAGGCCAGCTCACATGTGAGTCTTGGCAGAAGCCATGGCCCATGTCTGCACATCCAGGGAGGAGG  
ACAGAAGGCCAGCTCAGTGGCCCCGCTCCCCACCCCCACGCCGAACAGCAGGGGGCAGAGCAGCCCTCCTTC  
GAAGTGTCTCAAGTCCGCAATTTGAGCCTTGTCTGGGGCCACGCCAACCTGGCTTGGGCTCACTGTCTGA  
GTTGCAGTAAAGCTATAACCTTGAATCACA

**FIGURE 64**

MTTWSLRRRPPARTLGLLLLLVVLGFLVLRRLDWSTLVPLRLRHRQLGLQAKGWNFMLEDSTFW  
IFGGSIHYFRVPREYWRDRLLKMKACGLNLTITYVPWNLHEPERGKFDGSGNLDLEAFVLM  
AEIGLWVILRPGPYICSEMDLGLPSWLLQDPGMRLRTTYKGFTEAVDLYFDHLMSRVVPLQ  
YKRGGPIIAVQVENEYGSYNKDPAYMPYVKKALEDRGIVELLLTSDNKDGLSKGIVQGVLAT  
INLQSTHELQLLTTFLEFNVQGTQPKMVMMEYWTGWFDGSGPHNILDSSEVLKTVSAIVDAGS  
SINLYMFHGGTNFGFMNGAMHFHDYKSDVTSYDYDAVLTEAGDYTAKYMKLRDFFGSI  
LPPPPDLLPKMPYEPLTPVLYLSLWDALKYLGEPIKSEKPINMENLPVNGGNGQSFGYILYE  
TSITSSGILSGHVHVRGQVFNVTVSIGFLDYKTTKIAVPLIQGYTVLRILVENRGRVNYGEN  
IDDQRKGLIGNLYLNDSPKFNRIYSLDMKKSFFQRFGLDKWXSLETPPTLPAFFLGSLSIS  
STPCDTFLKLEGWEKGVVFINGQNLGRYWNIGPQKTLYLPGPWLSSGINQVIVFEETMAGPA  
LQFTETPHLGRNQYIK

**Signal sequence:**

amino acids 1-27

**Casein kinase II phosphorylation site.**

amino acids 141-118, 253-257, 340-344, 395-399, 540-544, 560-564

**N-myristoylation site.**

amino acids 146-152, 236-242, 240-246, 244-250, 287-293, 309-315,  
320-326, 366-372, 423-429, 425-431, 441-447, 503-509, 580-586

**FIGURE 65**

GGGGACGCGGAGCTGAGAGGCTCCGGGCTAGCTAGGTGTAGGGGTGGACGGGTCCCAGGACC  
CTGGTGAGGGTTCTCTACTTGGCCTTCGGTGGGGGTCAAGACGCAGGCACCTACGCCAAAGG  
GGAGCAAAGCCGGGCTCGGCCCGAGGCCCCAGGACCTCCATCTCCCAATGTTGGAGGAATC  
CGACACGTGACGGTCTGTCCGCCGTCTCAGACTAGAGGAGCGCTGTAACGCCATGGCTCCC  
AAGAAGCTGTCTGCCTTCGTTCCCTGCTGCTGCCGCTCAGCCTGACGCTACTGCTGCCCA  
GGCAGACACTCGGTCGTTTCGTAGTGGATAGGGGTCAAGACCGGTTTCTCTAGACGGGGCCC  
CGTTCGGCTATGTGTCTGGCAGCCTGCACTACTTTTCGGGTACC CGCGGGTGCPTTGGGCCGAC  
CGGCTTTTGAAGATGCGATGGAGCGGCTCAACGCCATACAGTTTTATGTGCCCTGGAACTA  
CCACGAGCCACAGCCTGGGGTCTATAACTTTAATGGCAGCCGGGACCTCATTGCCTTTCTGA  
ATGAGGCAGCTCTAGCGAACCTGTTGGTCACTGAGACCAGGACCTTACATCTGTGCAGAG  
TGGGAGATGGGGGTCTCCCATCCTGGTTGCTTCGAAAACCTGAAATTCATCTAAGAACCTC  
AGATCCAGACTTCTTGC CGCAGTGGACTCCTGGTTC AAGGTCTTGCTGCCCAAGATATATC  
CATGGCTTTATCACAAATGGGGGCAACATCATTAGCATTTCAGGTGGAGAATGAATATGGTAGC  
TACAGAGCCTGTGACTTCAGCTACATGAGGCCTTGGCTGGGCTCTCCGTCGACTGCTAGG  
AGAAAAGATCTTGCTCTTACCACAGATGGGCTGAAGGACTCAAGTGTGGCTCCCTCCGGG  
GACTCTATACCACTGTAGATTTTGGCCCAGCTGACAACATGACCAAAATCTTTACCCCTGCTT  
CGGAAGTATGAACCCCATGGGCCATTGGTAAACTCTGAGTACTACACAGGCTGGCTGGATTA  
CTGGGGCCAGAACTACTCCACACGGTCTGTGTGCTGAGCTGTAACCAAAGGACTAGAGAACATGC  
TCAAGTTGGGAGCCAGTGTGAACATGTACATGTTCCATGGAGGTACCAACTTTGGATATTGG  
AATGGTCCGATAAGAAGGGACGCTTCTTCCGATTACTACCAGCTATGACTATGATGCACC  
TATATCTGAAGCAGGGGACCCACACCTAAGCTTTTTCGCTCTTCGAGATGTCATCAGCAAGT  
TCCAGGAAGTTCCCTTGGGACCTTTACCTCCCCGAGCCCCAAGATGATGCTTGGACCTGTG  
ACTCTGCACCTGGTTGGGCATTTACTGGCTTTCTAGACTTGCTTTGCCCCCGTGGGCCCAT  
TCATTCAATCTTGCCAATGACCTTTGAGGCTGTCAAGCAGGACCATGGCTTCATGTTGTACC  
GAACCTATATGACCCATACCAATTTTGGAGCCAACACCAATTCCTGGGTGCCAAATAATGGAGTC  
CATGACCGTGCCATGTGATGGTGGATGGGGTGTTCAGGGTGTGTGGAGCGAAAATATGAG  
AGACAAACTATTTTGGACGGGAAAACGGGGTCCAAACTGGATATCTTGGTGGAGAACATGG  
GGAGGCTCAGCTTTGGGTCTAACAGCAGTGACTTCAAGGGCCTGTTGAAGCCACCAATTCCTG  
GGGCAAACAATCCTTACCCAGTGGATGATGTTCCCTCTGAAAATTGATAACCTGTGAAGTG  
GTGGTTTCCCCTCCAGTTGCCAAAATGGCCATATCCTCAAGCTCCTTCTGGCCCCACATTCCT  
ACTCCAAAACATTTCCAATTTTAGGCTCAGTTGGGGACACATTTCTATATCTACCTGGATGG  
ACCAAGGGCCAAGTCTGGATCAATGGGTTTAACTTGGGCCGGTACTGGACAAAGCAGGGGCC  
ACAACAGACCCCTTACGTGCCAAGATTCCTGCTGTTTCCCTAGGGGAGCCCTCAACAAAATTA  
CATTGCTGGAAGTAGAAGATGTACCTCTCCAGCCCCAAGTCCAATTTTTGGATAAGCCTATC  
CTCAATAGCACTAGTACTTTGCACAGGACACATATCAATTCCTTTTCAGCTGATACTACTGAG  
TGCTCTGAACCAATGGAGTTAAGTGGGCACTGAAAGGTAGGCCGGGCATGGTGGCTCATGC  
CTGTAATCCCAGCACTTTGGGAGGCTGAGACGGGTGGATTACCTGAGGTCAGGACTTCAAGA  
CCAGCCTGGCCAACATGGTGAAACCCGCTCCTCACTAAAAATACAAAAATFAGCCGGGCGTG  
ATGGTGGGCACCTCTAATCCCAGTACTTGGGAGGCTGAGGGCAGGAGAAATGCTTGAATCC  
AGGAGGCAGAGGTTGCAGTGAGTGGAGGTTGTACCCTGCCTCCAGCCTGGCTGACAGTGA  
GACACTCCATCTCAAAAAAAAAAAAA

**FIGURE 66**

MAPKKLSCLRSLLLPLSLTLLLPQADTRSFVVDGRGHDRFLLDGAPFRYVSGSLHYFRVPRVL  
WADRLMKMRWSGLNAIQFYVPWNYHEPQPGVYNFNNGSRDLIAFLNEAALANLLVILRPGPYI  
CAEWEMGGLPSWLLRKPEIHLRTSDPDFLAAVDSWFKVLLPKIYPWLYHNGGNIISIQVENE  
YGSYRACDFS YMRHLAGLFRALLGEKILLFTTDGPEGLKCGSLRGLYTTVDFGPADNMTKIF  
TLRKYEPHGFLVNSEYYTGWLDYWGQNHSTRSVSAVTKLENMLKLGASVNMVMFHGGTNF  
GYWNGADKKGRFLPITTSYDYDAPISEAGDPTPKLFALRDVISKFQEVPLGPLPPSPKMML  
GPVTLHLVGHLLAFLDLLCPRGPIHSILPMTFEAVKQDHGFMLYRTYMTHTIFEPTPFWVFN  
NGVHDRAYVMVDGVFQGVVERNMRDKLFLTGKLGSKLDILVENMGRLSFGSNSDFKGLLKP  
PILGQTIILTQWMMFPLKIDNLVKWWFPLQLPKWPYPQAPSGPTFYSKTFPILGSGDFTFLYL  
PGWTKGQVWINGFNLGRYWTQGPQOTLYVPRFLLFPRGALNKITLLELEDVPLQPQVQFLD  
KPILNSTSTLHRTHINSL SADTLSASEPMELSGH

**Signal sequence:**

amino acids 1-27

**N-glycosylation site.**

amino acids 97-101, 243-247, 276-280, 486-490, 625-629

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 4-8

**Casein kinase II phosphorylation site.**

amino acids 148-152, 234-238, 327-331, 423-427, 469-473, 550-554,  
603-607, 644-648

**Tyrosine kinase phosphorylation site.**

amino acids 191-198

**N-myristoylation site.**

amino acids 131-137, 176-182, 188-194, 203-209, 223-229, 227-233,  
231-237, 274-280, 296-300, 307-313, 447-453, 484-490

**FIGURE 67**

GCTTTGAACACGTCTGCAAGCCCAAAGTTGAGCATCTGATTGGTTATGAGGTATTTGAGTGC  
ACCCACAATATGGGCTTACATGTTGAAAAAGCTTCTCATCAGTTACATATCCATTATTTGTGT  
TTATGGCTTTATCTGCCTCTACACTCTCTTCTGGTTATTCAGGATACCTTTGAAGGAATATT  
CTTTTCGAAAAAGTCAGAGAAGAGAGCAGTTTTAGTGACATTCCAGATGTCAAAAACGATTTT  
GCGTTCCTTCTTACATGGTAGACCAGTATGACCAGCTATATTTCCAAGCGTTTTGGTGTGTT  
CTTGTGAGAAGTTAGTGAAAATAAACTTAGGGAAATTAGTTTGAACCATGAGTGGACATTTG  
AAAACTCAGGCAGCACATTTACGCAACGCCAGGACAAGCAGGAGTTGCATCTGTTTCATG  
CTGTGCGGGGTGCCCGATGCTGTCTTTGACCTCACAGACCTGGATGTGCTAAAGCTTGAAC  
AATTCCAGAAGCTAAAATTCCTGCTAAGATTTCTCAAATGACTAACCTCCAAGAGCTCCACC  
TCTGCCACTGCCCTGCAAAAGTTGAACAGACTGCTTTTAGCTTTCTTCGCGATCACTTGAGA  
TGCCTTACGTGAAGTTCACTGATGTGGCTGAAATTCCTGCCTGGGTGTATTTGCTCAAAAA  
CCTTCGAGAGTTGTACTTAATAGGCAATTTGAACTCTGAAAAAATAAGATGATAGGACTTG  
AATCTCTCCGAGAGTTGCGGCACCTTAAGATTTCTCCACGTGAAGAGCAATTTGACCAAAGTT  
CCCTCCAACATTACAGATGTGGCTCCACATCTTACAAAGTTAGTCATTACATAATGACGGCAC  
TAAACTCTTGGTACTGAACAGCCTTAAGAAAATGATGAATGTGCTGAGCTGGAACCTCCAGA  
ACTGTGAGCTAGAGAGAATCCACATGCTATTTTTCAGCCTCTCTAATTTACAGGAACCTGGAT  
TTAAAGTCCAATAACATTCGCACAATTTGAGGAAATCATCAGTTTCCAGCATTTAAAACGACT  
GACTTGTTTAAAAATTATGGCATAACAAAATTTGTTACTATTCTCCCTCTATTACCCATGTCA  
AAACTTGGAGTCACTTTATTTCTCTAAACAACAAGCTCGAATCCTTACCAGTGGCAGTATTT  
AGTTTACAGAACTCAGATGCTTAGATGTGAGCTACAACAACATTTCAATGATTTCCAATAGA  
AATAGGATTTGCTTCAGAACCTGCAGCATTTGCATATCACTGGGAACAAAGTGGACATTTCTGC  
CAAAACAATTGTTTAAATGCATAAAGTTGAGGACTTTGAATCTGGGACAGAAGTGCATCACC  
TCACTCCCAGAGAAAGTTGGTTCAGCTCTCCAGCTCACTCAGCTGGAGCTGAAGGGGAACTG  
CTTGGACCGCTGCCAGCCAGCTGGGCCAGTGTGGATGCTCAAGAAAAGCGGGCTTGTG  
TGGAAGATCACCTTTTGTATACCCTGCCACTCGAAGTCAAAGAGGCATTTGAATCAAGACATA  
AATATTCCTTTGCAAATGGGATTTAACTAAGATAATATATGCACAGTGATGTGCAGGAAC  
AACTTCCTAGATTGCAAGTGTTCACGTACAAGTTATTACAAGATAATGCATTTTAGGAGTAG  
ATACATCTTTTAAAAATAAAACAGAGAGGATGCATAGAAGGCTGATAGAAGACATAACTGAAT  
GTTCAATGTTTGTAGGGTTTTAAGTCATTCATTTCCAAATCATTTTTTTTTTTCTTTTGGGG  
AAAGGAAGGAAAAATATAATCACTAATCTTGGTTCTTTTTAAATGTTTGTAACTTGGAT  
GCTGCCGCTACTGAATGTTTACAAATGCTTGCCTGCTAAAGTAAATGATTAAATTTGACATT  
TTCTTACTAAAAA

## FIGURE 68

MAYMLKLLISYISIIICVYGFICLYTLFWLFRIPLKEYSFEKVVREESSFSDIPDVKNDF AFL  
LHMVDQYDQLYSKRFGVFLSEVSENKLR EISLNHEWTFEKL RQHISRNAQDKQELHLFMLS G  
VPDAVFDLTDLDV LKLELIPEAKIPAKISQMTNLQELHLCHCPAKVEQTAFSFLRDHLRCLH  
VKFTDVAEIPAWVYLLKNLRELYLIGNL NSENKMI GLESLREL RHLKILHVKS NLTKVPSN  
ITDVAPHLTKLVIHNDGT KLLVLNSLKKMMNVAE LELQNC ELERIPHAIFSLSNLQELDLKS  
NNIRTIEEIIISFQHLKRLTCLKLWHNKIVTIPPSITHVKNLES LYFSNNKLES LPVAVFSLQ  
KLRCLDVSYNNISMIP IEIGLLQNLQHLHITGNKV DILPKQLFKCIKLR TLNLGQNCITSLP  
EKVGQLSOLTQLELKGNC LDR LPAQLGQCRMLK KSGLVVEDHLFD TLPLEVKEALNQDINIP  
FANGI

**Signal sequence:**

amino acids 1-20

**N-glycosylation site.**

amino acids 241-245, 248-252, 383-387

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 326-330

**Casein kinase II phosphorylation site.**

amino acids 48-52, 133-137, 226-230, 315-319, 432-436, 444-448

**Tyrosine kinase phosphorylation site.**

amino acids 349-355, 375-381

**N-myristoylation site.**

amino acids 78-84, 124-130, 212-218, 392-398

**FIGURE 69**

CCCACGCGTCCGGCCCTTCTCTCGGACTTTGCATTTCCATTCCTTTTCATGACAAACTGACTTTTTTTTATTCT  
TTTTTTCATCTCTGGGCCAGCTTTGGGATCCTAGGCCGCCCTGGGAAGACATTTGTGTTTTACACACATAAGGAT  
CTGTGTTTTGGGTTTTCTTCTCCCTCCCTGACATTGGCATTGCTTAGTGGTGTGTGGGGAGGGAGACCACGTGG  
GCTCAGTGCCTTGCTTGCATTTATCTGCCTAGGTACATCGAAGTCTTTTGCCTCCATACAGTGATTATGCCTGTC  
ATCGCTGGTGGTATCCTGGCGGCCCTTGCTCCTGCTGATAGTTGTGCTGCTCTGTCTTTACTTCAAAATACACAAC  
GCGCTAAAAGCTGCAAGGAACCTGAAGCTGTGGCTGTAAAAAATCACAAACCAGACAAGGTGTGTGGGCCAAG  
AACAGCCAGGCCAAAACCATGCCCAGGAGTCTTGTCTGCCCTGCACTGCTGTGAAGGATATAGAAATGTGTGCC  
AGTTTTGATTCCTGCCACCTTGCTGTTGCGACATAAATGAGGGCCTCTGAGTTAGGAAAGGCTCCCTTCTCAAA  
GCAGAGCCCTGAAGACTTCAATGATGTCAATGAGGCCACCTGTTTGTGATGTGCAGGCACAGAAGAAAGGCACAG  
CTCCCATCAGTTTTCATGGAAAATAACTCAGTGCCTGCTGGGAACCAGCTGCTGGAGATCCTACAGAGAGCTTC  
CACTGGGGCAACCCCTCCAGGAAGGAGTTGGGGAGAGAGAACCCTCACTGTGGGGAATGCTGATAAACCACTCA  
CACAGCTGCTCTATTCTCACACAAAATCACCCCTTGCGTGGCTGGAACCTGACGTTTCCCTGGAGGTGTCCAGAAA  
GCTGATGTAACACAGAGCCTATAAAAGCTGTGGTCTTAAAGGCTGCCAGCGCCCTTGCCAAAATGGAGCTTGTA  
AGAAGGCTCATGCCATTGACCCCTTTAATTTCTCTCTGTTTGGCGGAGCTGACAAATGGCGGAGGCTGAAGGCAAT  
GCAAGCTGCACAGTCACTAGGGGGTGCCAAATATGGCAGAGACCACAAAGCCATGATCCTGCAACTCAATCCC  
AGTGAGAAGCTGCACCTGGACAATAGAAAAGACCAGAAAACAAAAGCATCAGAATTTACTTTTTCTATGTCCAGCTT  
GATCCAGATGGAAGCTGTGAAAGTGAAGCAATTAAGTCTTTGACGGAACTCCAGCAATGGGCCCTGTAGGG  
CAAGTCTGCAGTAAAAACGACTATGTTCTGTATTTGAATCATCATCCAGTACATTGACGTTTTCAATAGTTACT  
GACTCAGCAAGAATTCAAAGAAGTGTCTTTGTCTTCTACTACTTCTCTCTCTCTCAACATCTCTATTCCAAAAGT  
GGCGGTTACCTGGATACCTTGAAGGATCCTTACCAGCCCCAATTACCCAAAGCCGCATCCTGAGCTGGCTTAT  
TGTGTGTGGCACATACAAGTGGAGAAAGATTACAAGATAAAAATAAACTTCAAAGAGATTTTCTAGAAAATAGAC  
AAACAGTGCAAAATTTGATTTTTCTTGCCATCTATGATGGCCCTCCACCAACTCTGGCCTGATGGACAAGTCTGT  
GGCCGTGTGACTCCCACTCGAATCGTCAACAACTCTCTGACTGTGCTGTGTCTACAGATTAAGCCAAATCT  
TACCGGGGATTTTCTGCTTCTTACACTCAATTTATGCAGAAAACATCAACACTACATCTTTAACTGTCTTCT  
GACAGGATGAGAGTTATTATAAGCAAACTCTACCTAGAGGCTTTTAACTTAATGGGAATAACTTGAACATAAAA  
GACCCAACTTGCAGACCAAAATTAACAATGTTGTGGAATTTCTGTCCCTTTAATGGATGTGGTACAATCAGA  
AAGGTAGAAGATCAGTCAATTACTTACACCAATATAATCACCTTTTCTGCATCCTCAACTCTGAAGTGATCACC  
CGTCAAAAACAACTCCAGATTATTGTAAGTGTGAAATGGGACATAATTCTACAGTGGAGATAATATAACATAACA  
GAAGATGATGTAATCAAAATCAAAATGCACTGGGCAAATATAACACCAGCATGGCTTTTTTTGAATCCAATTC  
TTTTAAAAGACTATACTTGAATCACCATAATTATGTGGATTTGAACCAAACTCTTTTTGTTCAAGTTAGTCTGCAC  
ACCTCAGATCCAAATTTGGTGGTGTCTTGTGATACCTGTAGAGCCTCTCCACCTCTGACTTTGCATCTCCAACC  
TAGCACCATAACAGAGTGGATGTAGTTCGAGATGAACTTGTAAAGGTGTATCCCTTATTTGGACACTATGGGAGA  
TTCCAGTTTTAATGCCTTTAAATTTCTGAGAAGTATGAGCTCTGTGTATCTGCAGTGTAAAGTTTTGATATGTGAT  
AGCAGTGACCACAGTCTCGCTGCAATCAAGGTTGTGTCTCCAGAAGCAAACGAGACATTTCTTATATAAATGG  
AAAACAGATTCATCATAGACCCATTCGTCTGAAAAGGGATCGAAGTGCAGTGGCAATTGAGGATTTGAGCAT  
AATGTGGTACTGTAGCGACAATCACAGTGAAGCATTTTGTAAATCAACGGGCAGACTACAAATACCAGAAGCTG  
CAGAACTATTAACTAACAGTCCAACCCCTAAGTGAGACATGTTCTCCAGGATGCCAAAGGAAATGTACCTCGT  
GGCTACACATATTATGAATAAATGAGGAAGGGCCTGAAAGTGACACACAGGCCTGCATGTAAAAAAA

## **FIGURE 70**

MELVRRMLPLTLLILSCLAELTMAEAEAGNASCTVSLGGANMAETHKAMILQLNPSENCTWTI  
ERPENKSIRIIFSYVQLDPDGSCESENIKVFDGTSSNGPLLQVCSKNDYVPVFESSSSTLT  
FQIVTDSARIQRTVFVFFYYFFSPNISIPNCGGYLDTLEGSFTSPNYPKPHELAYCVWHIQV  
EKDYKIKLNFKEIFLEIDKQCKFDLAIYDGPSTNSGLIGQVCGRVTPTFESSNSLTVVLS  
TDYANSYRGFSASYTSIYAENINTTSLTSSDRMRVIIKSYLEAFNSNGNNLQKDPCTCRP  
KLSNVVEFSVPLNGCGTIRKVEDQSITYTNIITFSASSTSEVITRQKQLQIIVKCEMGHNST  
VEIIYITEDDDVIQSQNALGKYNTSMALFESNSFEKTIILESPYYVDLNQTLFVQVSLHTSDPN  
LVVFLDTCRASPTSDFASPTYDLIKSGCSRDETCKVYPLFGHYGRFQFNAFKFLRSMSVYL  
QCKVLI CDSSDHQSRCNQGCVSRSKRDISSYKWKTDSSIIGPIRLKRDRSASGNSGFQHETHA  
EETPNQPFNSVHLFSFMVLALNVVTVATITVRHFVNQRADYKYQKLQNY

**Signal sequence:**

amino acids 1-24

**Transmembrane domain:**

amino acids 571-586

**N-glycosylation site.**

amino acids 29-33, 57-61, 67-71, 148-152, 271-275, 370-374,  
394-398, 419-423

**Casein kinase II phosphorylation site.**

amino acids 22-26, 108-112, 289-293, 348-352, 371-375, 379-383,  
408-412, 463-467, 520-524, 556-560

**Tyrosine kinase phosphorylation site.**

amino acids 172-180, 407-415, 407-416, 519-528

**N-myristoylation site.**

amino acids 28-34, 38-44, 83-89, 95-101, 104-110, 226-232

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 7-18



**FIGURE 71**

GACGGAAGAACAGCGCTCCCCGAGGCCGCGGGAGCCTGCAGAGAGGACAGCCGGCCTGCGCCG  
GGACAATGCGGCCCCCAGGAGCTCCCCAGGCTCGCGTTCCCGTTGCTGCTGTTGCTGTTGCTGC  
TGCTGCCGCGCCCGCCCGTGCCTGCCACAGCGCCACGCGCTTCGACCCCACCTGGGAGTCC  
CTGGACGCCCCGCCAGCTGCCCGCGTGGTTTTGACCAGGCCAAGTTCGGCATCTTCATCCACTG  
GGGAGTGTTTTTCCGTGCCCAGCTTCGGTAGCGAGTGGTTCTGGTGGTATTGGCAAAGGAAA  
AGATACCGAAGTATGTGGAATTTATGAAAGATAATTACCCTCCTAGTTTCAAATATGAAGAT  
TTTGGACCACATTTACAGCAAAATTTTTTAATGCCAACCAAGTGGGCAGATATTTTTCAGGC  
CTCTGGTGCCAAATACATTTGTCTTAACTTCCAACATCATGAAGGCTTTACCTTGTGGGGGT  
CAGAATAATTCGTGGAACCTGGAATGCCATAGATGAGGGGCCAAGAGGGACATTGTCAAGGAA  
CTTGAGGTAGCCATTAGGAACAGAACTGACCTGCGTTTTGGACTGTACTATTCCTTTTTGA  
ATGGTTTTCATCCGCTCTTCCTTGAGGATGAATCCAGTTCATTCCATAAGCGGCAATTTCCAG  
TTTTCAAGACATTGCCAGAGCTCTATGAGTTAGTGAACAACATCAGCCTGAGGTTCTGTGG  
TCGGATGTTGACGGAGGAGCACCGGATCAATACTGGAACAGCACAGGCTTCTTGGCCTGGTT  
ATATAATGAAAGCCCAGTTCCGGGACAGTAGTCACCAATGATCGTTGGGGAGCTGGTAGCA  
TCTGTAAGCATGGTGGCTTCTATACCTGCAGTGATCGTTATAACCCAGGACATCTTTTTGCCA  
CATAAATGGGAAAACATGCATGACAATAGACAACACTGCTCTGGGGCTATAGGAGGGAAGCTGG  
AATCTCTGACTATCTTACAATTTGAAGAATTTGGTGAAGCAACTTGTAGAGACAGTTTCCATGTG  
GAGGAAATCTTTTTGATGAATATTTGGGCCACACTAGATGGCACCATTTCTGTAGTTTTTGAG  
GAGCGACTGAGGCAAGTGGGGTCTGGCTAAAAGTCAATGGAGAAGCTATTTATGAAACCTA  
TACCTGGCGATCCCAGAAATGACACTGTCACCCAGATGTGTGGTACACATCCAAGCCTAAAG  
AAAAATTAGTCTATGCCATTTTTCTTAAATGGCCACATCAGGACAGCTGTTCTTGGCCAT  
CCCAAAGCTATTTCTGGGGGCAACAGAGGTGAAACTACTGGGCCATGGACAGCCACTTAAC TG  
GATTTCTTTGGAGCAAAATGGCATTATGGTAGAACTGCCACAGCTAACCATTATCAGATGC  
CGTGTAATGGGGCTGGGCTCTAGCCCCTAACTAATGTGATCTAAAGTGCAGCAGAGTGGCTG  
ATGCTGCAAGTTATGCTAAGGCTAGGAACATCAGGTGTCTATAATTTGTAGCACATGGAGA  
AAGCAATGTAAACTGGATAAGAAAATTTTGGCAGTTCAGCCCTTTCCCTTTTTCCACTA  
AATTTTTCTTAAATTACCCATGTAACCATTTTAACTCTCCAGTGCACTTTGCCATTAAAGTC  
TCTTCACATTTGATTTGTTTCCATGTGTGACTCAGAGGTGAGAATTTTTTTCACATTATAGTAG  
CAAGGAATTTGGTGGTATTATGGACCGAACTGAAAATTTTATGTTGAAGCCATATCCCCATG  
ATTATATAGTTATGCATCACTTAATATGGGGATATTTTCTGGGAAATGCATTGCTAGTCAAT  
TTTTTTTTGTGCCAACATCATAGAGTGTATTTACAAAATCCTAGATGGCATAGCCTACTACA  
CACCTAATGTGTATGGTATAGACTGTTGCTCCTAGGCTACAGACATATACAGCATGTTACTG  
AATACGTAGGCAATAGTAAACAGTGGTATTTGTATATCGAAACATATGGAAACATAGAGAAG  
GTACAGTAAAAATACGTAAAATAAATGGTGCACCTGTATAGGGCACTTACCACGAATGGAG  
CTTACAGGACTGGAAGTTGCTCTGGGTGAGTCAGTGAGTGAATGTGAAGGCCTAGGACATTA  
TTGAACACTGCCAGACGTTATAAATACTGTATGCTTAGGCTACACTACATTTATAAAAAAAA  
GTTTTTCTTTCTTCAATTATAAATTAACATAAGTGTACTGTAACTTTACAAACGTTTTAATT  
TTTAAAACCTTTTTGGCTCTTTTGTAAATAACACTTAGCTTAAAACATAAACTCATTTGTGCAA  
ATGTAA

**FIGURE 72**

MRPQELPRLAFPLLLLLLLLLLLLLPPPPCPAHSATRFDPTWESLDARQLPAWFDQAKFGIFIHWG  
VFSVPSFGSEWFWWYQKEKI PKYVEFMKDNYPSPFKYEDFGPLFTAKFFNANQWADIFQAS  
GAKYIVLTSKHHEGFTLWGSEYSWNWNAIDEGPKRDIVKELEVAIRNRTDLRFGLYYSLEFEW  
FHPLFLEDESSSFHKRQFPVSKTLPELYELVNNYQPEVLWSDGDGGAPDQYWNSTGFLAWLY  
NESPVRGTVVTNDRWGAGSICKHGGFYTCSDRYNPGHLLPHKWENCMTIDKLSWGYRREAGI  
SDYLTIEELVKQLVETVSCGGNLLMNIGPTLDGTISVVFEERLRQVGSWLKVNGEAIYETYT  
WRSQNDTIVTPDVWYTSKPKEKLVYAI FLKWPTSGQLFLGHPKAILGATEVKLLGHGQPLNWI  
SLEQNGIMVELPQLTIHQMPCKKGWALALTNVI

**Signal sequence:**

amino acids 1-28

**N-glycosylation site.**

amino acids 171-175, 239-243, 377-381

**Casein kinase II phosphorylation site.**

amino acids 32-36, 182-186, 209-213, 227-231, 276-280, 315-319,  
375-375

**Tyrosine kinase phosphorylation site.**

amino acids 361-369, 389-397

**N-myristoylation site.**

amino acids 143-149, 178-184, 255-261, 272-278, 428-434

**Leucine zipper pattern.**

amino acids 410-432

**Alpha-L-fucosidase putative active site.**

amino acids 283-295

**FIGURE 73**

AGCAGGGAAATCCGGATGTCTCGGTTATGAAGTGGAGCAGTGAGTGTGAGCCTCAACATAGT  
 TCCAGAACTCTCCATCCGGACTAGTTATTGAGCATCTGCCTCTCATATCACCAGTGGCCATC  
 TGAGGTGTTTCCCTGGCTCTGAAGGGGTAGGCACGATGGCCAGGTGCTTCAGCCTGGTGTG  
 CTTCTCACTTCCATCTGGACCACGAGGCTCCTGGTCCAAGGCTCTTTGCGTGCAGAAGAGCT  
 TTCCATCCAGGTGTCATGCAGAATTATGGGGATCACCCCTTGTGAGCAAAAAGGCCGAACCAGC  
 AGCTGAATTTACAGAAGCTAAGGAGGCCTGTAGGCTGCTGGGACTAAGTTTGGCCGGCAAG  
 GACCAAGTTGAAACAGCCTTGAAAGCTAGCTTTGAAACTTGCAGCTATGGCTGGGTGGAGA  
 TGGATTTCGTGGTCATCTCTAGGATTAGCCCAAACCCCAAGTGTGGGAAAAATGGGGTGGGTG  
 TCCTGATTTGGAAGGTTCCAGTGCAGCCGACAGTTTGCAGCCTATFTTACAACCTCATCTGAT  
 AACTGGACTAACTCGTGCATTCAGAAATTATACCACCAAAGATCCCATATTTCAACACTCA  
 AACTGCAACACAAACAGAAATTTATTGTCACTGACAGTACCTACTCGGTGGCATCCCCTT  
 ACTCTACAATACCTGCCCTACTACTCTCCTCTGCTCCAGCTTCCACTTCCATTTCCACGG  
 AGAAAAAATFGATTTGTGTACAGAAAGTTTTTATGAAACTAGCACCATGTCTACAGAAAC  
 TGAACCATTTGTTGAAAATAAAGCAGCATTCAAGAATGAAGCTGCTGGGTTTGGAGGTGTC  
 CCACGGCTCTGCTAGTGTCTCTCCTCTTCTTTGGTGTGCAGCTGGTCTTGGATTTTGC  
 TATGTCAAAAGGTATGTGAAGGCCCTCCCTTTTACAAACAAGAATCAGCAGAAGGAAATGAT  
 CGAAACCAAAGTAGTAAAGGAGGAGAAGGCCAATGATAGCAACCCTAATGAGGAATCAAAGA  
 AAAGTATAAAAACCCAGAAGAGTCCAAGAGTCCAAGCAAACTACCGTGCAGTGCCTGGAA  
 GCTGAAGTTTAGATGAGACAGAAATGAGGAGACACACCTGAGGCTGGTTTCTTTTCATGCTCC  
 TTACCCTGCCCCAGCTGGGAAATCAAAGGGCCAAAGAACCAAAAGAAAGTCCACCCTT  
 GGTTCCCTAACTGGAATCAGCTCAGGACTGCCATTGGACTATGGAGTGCACCAAAGAGAATGC  
 CCTTCTCCTTATTGTAACCTGTCTGGATCCTATCCTCCTACCTCCAAAGCTTCCCACGGCC  
 TTTCTAGCCTGGCTATGTCTTAATAATATCCCAGTGGGAGAAAGGAGTTTTCGAAAGTGCAG  
 GGACCTAAAACATCTCATCAGTATCCAGTGGTAAAAAGGCCTCCTGGCTGTCTGAGGCTAGG  
 TGGGTTGAAAGCCAAGGAGTCACTGAGACCAAGGCTTTCTCTACTGATTCCGCAGCTCAGAC  
 CCTTCTTCAGCTCTGAAAGAGAAACAGTATCCACCTGACATGTCTTCTGAGCCCGGTA  
 AGAGCAAAAGAAATGGCAGAAAAGTTTAGCCCTGAAAGCCATGGAGATCTCATAACTTGAG  
 ACCTAATCTCTGTAAAGCTAAAATAAAGAAATAGAACAAGGCTGAGGATACGACAGTACACT  
 GTCAGCAGGGACTGTAAACACAGACAGGGTCAAAGTGTTTTCTCTGAACACATTGAGTTGGA  
 ATCACTGTTTAGAACACACACACTTACTTTTTCTGGTCTCTACCCTGCTGATATTTTCTCT  
 AGGAAATATACTTTTACAAGTAACAAAAATAAAACTCTTATAAAATTTCTATTTTTATCTGA  
 GTTACAGAAATGATTACTAAGGAAGATTACTCAGTAATTTGTTAAAAAGTAATAAAATTCA  
 ACAACATTTGCTGAATAGCTACTATATGTCAAGTGTGTGCAAGGTATTACTCTGTAAT  
 TGAATATTATTCCTCAAAAATTTGCACATAGTAGAACGCTATCTGGGAAGCTATTTTTTTCA  
 GTTTTGGATATTTCTAGCTTATCTACTTCCAACTAATTTTTATTTTTGCTGAGACTAATCTT  
 ATTCATTTTCTCTAATATGGCAACCATTATAACCTTAATTTATTATTAACATACCTAAGAAG  
 TACATTGTTACCTCTATATACCAAAGCACATTTTAAAAGTGCCATTAACAAATGTATCACTA  
 GCCCTCCTTTTCCAACAAGAAGGACTGAGAGATGCAGAAATATTTGTGACAAAAAATTA  
 AGCATTTAGAAAACCTT

**FIGURE 74**

MARCFSLVLLLTISIWTTRLLVQGSRLAEELSIQVSCRIMGITLVSKKANQQQLNFTEAKEACR  
LLGLSLAGKDQVETALKASFETCSYGWVGDGFVVISRISPNPKCGKNGVGVLIWKVPVSRQF  
AAYCYNSSDTWTNSCIPEIITTKDPIFNTQTATQTTEFIVSDSTYSVASPYSTIPAPTTTPP  
APASTSIPRRKKLICVTEVFMETSTMSTETEPFVENKAAFKNEAAGFGGVPTALLVLALLFF  
GAAAGLGFCYVKRYVKAFPFTNKNQOKEMIETKVVKEEKANDSNPNEESKKTDKNPEESKSP  
SKTTVRCLEAEV

**Signal sequence:**

amino acids 1-16

**Transmembrane domain:**

amino acids 235-254

**N-glycosylation site.**

amino acids 53-57, 130-134, 289-293

**Casein kinase II phosphorylation site.**

amino acids 145-149, 214-218

**Tyrosine kinase phosphorylation site.**

amino acids 79-88

**N-myristoylation site.**

amino acids 23-29, 65-71, 234-240, 235-239, 249-255, 253-259

**FIGURE 75**

AGATGGCGGTCTTGGCACCTCTAATTGCTCTCGTGATTCGGTGCCGCGACTTTCACGATGG  
CTCGCCCAACCTTACTACCTTCTGTGCGCCCTGCTCTCTGCTGCCTTCTACTCGTGAGGAA  
ACTGCCGCCGCTCTGCCACGGTCTGCCACCCAACGCGAAGACGGTAACCCGTGTGACTTTG  
ACTGGAGAGAAGTGGAGATCCTGATGTTTCTCAGTGCCATTGTGATGATGAAGAACCGCAGA  
TCCATCACTGTGGAGCAACATATAGGCAACATTTTCATGTTTAGTAAAGTGGCCAACACAAT  
TCTTTTCTTCCGCTTGGATATTCGCATGGGCCACTTTACATCACACTCTGCATAGTGTTC  
TGATGACGTGCAAACCCCCCTATATATGGGCCCTGAGTATATCAAGTACTTCAATGATAAA  
ACCATTGATGAGGAAC TAGAACGGGACAAGAGGGTCACTTGGATTGTGGAGTTCTTTGCCAA  
TTGGTCTAATGACTGCCAATCATTTGCCCTATCTATGCTGACCTCTCCCTTAAATACAAC  
GTACAGGGCTAAATTTTGGGAAGGTGGATGTTGGACGCTATACTGATGTTAGTACGCGGTAC  
AAAGTGAGCACATCACCCCTCACCAAGCAACTCCCTACCCCTGATCCTGTTCCAAGGTGGCAA  
GGAGGCAATGCGGCGGCCACAGATTGACAAGAAAGGACGGGCTGTCTCATGGACCTTCTCTG  
AGGAGAATGTGATCCGAGAAATTAACCTTAAATGAGCTATAACCAGCGGGCCAAGAACTATCA  
AAGGCTGGAGACAATATCCCTGAGGAGCAGCCTGTGGCTTCAACCCCCACCACAGTGTGAGA  
TGGGAAAACAAGAAGGATAAATAAGATCCTCACTTTGGCAGTGCTTCCCTCTCTCTGCAATT  
CCAGGCTCTTTCCATAACCACAAGCCTGAGGCTGCAGCCTTTNATTNATGTTTTCCCTTTGG  
CTGNGACTGGNTGGGGCAGCATGCAGCTTCTGATTTTAAAGAGGCATCTAGGGAATTGTCAG  
GCACCTACAGGAAGGCTGCCATGCTGTGGCCAACTGTTTCACTGGAGCAAGAAAGAGATC  
TCATAGGACCGAGGGGAAATGGTTTTCCCTCCAAGCTTGGGTGAGTGTGTTAACTGCTTATC  
AGCTATTCAGACATCTCCATGGTTTTCTCCATGAAACTCTGTGGTTTTCATCATTCCCTCTTAG  
TTGACCTGCACAGCTTGGTTAGACCTAGATTTAACCCCTAAGGTAAGATGCTGGGGTATAGAA  
CGCTAAGAATTTTCCCCAAGGACTCTTGCTTCCCTAAGCCCTTCTGGCTTCGTTTATGGTC  
TTCATTAAAAGTATAAGCCTAACTTTGTGCTAGTCCCTAAGGAGAAACCTTTAACCAAAAG  
TTTTTATCATTGAAGACAATATTGAACAACCCCTATTTTGTGGGGATTGAGAAGGGGTGAA  
TAGAGGCTTGAGACTTTCCTTTGTGTGGTAGGACTTGGAGGAGAAATCCCCTGGACTTTCAC  
TAACCTCTGACATACTCCCCACCCAGTTGATGGCTTTCGTAATAAAAAGATTGGGATT  
TCCTTTTG

**FIGURE 76**

MAVLAPLIALVYSVPERLSRWLAQPYLLSALLSAAFLLVKRLPPLCHGLPTQREDGNPCDFD  
WREVEILMFLSAIVMMKNRRSITVEQHIGNIFMFSKVANTILFFRLDIRMGLLYITLCIVFL  
MTCKPPLYMGPEYIKYFNDKTIDEELERDKRVTWIVEFFANWSNDCQSFAPIYADLSLKYN  
TGLNFGKVDVGRYTDVSTRYKVSTSP LTKQLPTLLILFQGGKEAMRRPQIDKKGRAVSWTFSE  
ENVIREFNLNELYQRAKKLSKAGDNIPEEQPVASTPTTVSDGENKKDK

**Signal sequence:**

amino acids 1-48

**Transmembrane domain:**

amino acids 111-125

**N-glycosylation site.**

amino acids 165-169, 185-189

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 154-158, 265-269

**Casein kinase II phosphorylation site.**

amino acids 51-55, 145-149, 245-249, 286-290, 288-292

**N-myristoylation site.**

amino acids 188-194, 225-231

**Myb DNA-binding domain repeat signature 1.**

amino acids 244-253



## **FIGURE 78**

MGLLLLVPLLLLPGSYGLPFYNGFYYSNSANDQNLGNHGHGKDLLNGVKLVVETPEETLFTYQ  
GASVILPCRYRYEPALVSPRRVRVKWKKLSENGAPEKDVLVAIGLRHRSFGDYQGRVHLRQD  
KEHDVSLLEIQDLRLEDYGRYRCEVIDGLEDESGLVELELRGVVFPYQSPNGRYQFNFHEGQQ  
VCAEQAAVVASFEOQFRAWEEGLDWCNAGWLQDATVQYPIMLRQPCGGPGLAPGVRSYGPR  
HRRLHRYDVFCFATALKGRVYYLEHPEKLTLTAREACQEDDATIAKVGQLFAAWKFHGLDR  
CDAGWLADGSVRYPVVHHPNCGPPEPGVRSFGFPDQSRLYGVYCYRQH

**Signal sequence:**

amino acids 1-17

**Casein kinase II phosphorylation site.**

amino acids 29-33, 53-57, 111-115, 278-282

**Tyrosine kinase phosphorylation site.**

amino acids 137-145

**N-myristoylation site.**

amino acids 36-42, 184-190, 208-214, 237-243, 297-303, 307-313





**FIGURE 80**

MMWRPSVLLLLLLLLLRHGAQ GKPS PDAGPHGQGRVHQAAPLSDAPHDDAHGNFQYDHEAFLGR  
EVAKEFDQLTPEESQARLGRIVDRMDRAGDGDGWVSLAELRAWIAHTQQRHIRDSVSAAWDT  
YDTRDRGRVGEELRNATYGHYAPGEEFHDVEDAETYKKMLARDERRFRVADQDGDSMATRE  
ELTAFLHPEEFPHMRDIVIAETLEDLDRNKDGYVQVEEYIADLYSAEPGEEEPWVQTERQQ  
FRDFRDLNKDGHLDGSEVGHVWLPPAQDQPLVEANHLLHESDTPDKGRLSKAEILGNWNMFV  
GSQATNYGEDLTRHHDEL

**Signal sequence:**

amino acids 1-20

**N-glycosylation site.**

amino acids 140-144

**Casein kinase II phosphorylation site.**

amino acids 72-76, 98-102, 127-131, 184-188, 208-212, 289-293,  
291-295, 298-302

**N-myristoylation site.**

amino acids 263-269, 311-317

**Endoplasmic reticulum targeting sequence.**

amino acids 325-330

**FIGURE 81**

GGGGCCTTGCCFFCCGCACTCGGGCGCAGCCGGGTGGATCTCGAGCAGGTGCGGAGCCCCGG  
 GCGGCGGGCGCGGGTGCAGGGGATCCCTGACGCCTCTGTCCCTGTTTCTTTGTGCTCCAG  
 CCTGTCTGTGCTGCTTTTGGCGCCCCCGCCTCCCCGCGGTGCGGGGTTGCACACCGATCCTG  
 GGCTTCGCTCGATTTGCGCCGAGGCGCCTCCAGACCTAGAGGGGCGCTGGCCTGGAGCAG  
 CGGGTCGTCTGTCTCTCTCTCTGCGCCGCGCCCGGGGATCCGAAGGGTGCGGGGCTCT  
 GAGGAGGTGACGCGCGGGGCTCCCGCACCTGGCCTTGCCCGCATTCCTCCTCTCTCCAG  
 GTGTGAGCAGCCTATCAGTCACCATGTCCGCGCCTGGATCCCGGCTCTCGGCCTCGGTGTG  
 TGTCTGCTGCTGCTGCCGGGGCCCGGGGCAGCGAGGGAGCCGCTCCCATTTGTATCACATG  
 TTTTACCAGAGGCTTGACATCAGGAAAGAGAAAGCAGATGTCTCTGCCAGGGGGCTGCC  
 CTCTTGAGGAATTTCTGTGTATGGGAACATAGTATATGCTTCTGTATCGAGCATATGTGGG  
 GCTGCTGTCCACAGGGGAGTAATCAGCAACTCAGGGGGACCTGTACGAGTCTATAGCCTACC  
 TGGTTCGAGAAAATTTCTCAGTAGATGCCAATGGCATCCAGTCTCAAATGCTTTCTAGAT  
 GGTCTGCTTCTTTCACAGTAACTAAAGGCAAAAGTAGTACACAGGAGGCCACAGGACAAGCA  
 GTGTCCACAGCACATCCACCAACAGGTAAACGACTAAAGAAAACACCCGAGAAGAAAAGTGG  
 CAATAAAGATTTGTAAGCAGACATTCATTTCTGATTGATGGAAGCTTTAATAATGGGGCAGC  
 GCCGATTTAATTTACAGAAGAATTTTGGTTGGAAAAGTGGCTCTAATGTTGGGAATTTGGAACA  
 GAAGGACCACATGTGGGCCTTGTTCAGCCAGTGAACATCCCAAATAGAAATTTTACTTGAA  
 AAATTTACATCAGCCAAAGATGTTTTGTTGCCATAAAGGAAGTAGGTTTCAGAGGGGGTA  
 ATTTCAATACAGGAAAAGCCTTGAAGCATACTGCTCAGAAAATTTTCACGGTAGATGCTGGA  
 GTAAGAAAAGGGATCCCCAAAGTGGTGGTGGTATTTATTGATGGTTGGCCTTCTGATGACAT  
 CGAGGAAGCAGGCATTTGTGGCCAGAGAGTTTGGTGTCAATGTATTTATAGTTTCTGTGGCCA  
 AGCCTATCCCTGAAGAACTGGGGATGGTTTCAGGATGTCACATTTGTTGACAAGGCTGTCTGT  
 CGGAATAATGGCTTCTTCTTACCACATGCCCAACTGGTTTGGCACCACAAAATACGTA  
 GCCTCTGGTACAGAAGCTGTGCACTCATGAACAAATGATGTGCAGCAAGACCTGTTATAACT  
 CAGTGAACATTGCCTTTCTAATTGATGGCTCCAGCAGTGTGGAGATAGCAATTTCCGCCTC  
 ATGCTTGAATTTGTTTCCAACATAGCCAAGACTTTTGAAATCTCGGACATTTGGTCCAAAGAT  
 AGCTGCTGTACAGTTTACTTATGATCAGCGCACGGAGTTTCTGACTACTATAGCACC  
 AAGAGAATGTCCTAGCTGTATCAGAAACATCCGCTATATGAGTGGTGGAAACAGCTACTGGT  
 GATGCCATTTCTTCACTGTTAGAAATGTGTTTGGCCCTATAAGGGAGAGCCCCAACAAAGAA  
 CTTCTAGTAATTTGTACAGATGGGCAGTCTATGATGATGTCCAAGGCCCTGCAGCTGCTG  
 CACATGATGCAGGAATCACTATCTTCTCTGTTGGTGTGGCTTGGGCACCTCTGGATGACCTG  
 AAAGATATGGCTTCTAAACCGAAGGAGTCTCACGCTTTCTTCAAGAGAGTTTACAGGATT  
 AGAACCAATTTGTTTCTGATGTCATCAGAGGCATTTGTAGAGATTTCTTAGAATCCAGCAAT  
AATGGTAACATTTTGGACAACGTAAAGAAAAAGTACAAGGGGATCCAGTGTGTAATTTGTATT  
 CTCATAATACTGAAATGCTTTAGCATACTAGAATCAGATACAAAATATTAAGTATGTCAAC  
 AGCCATTTAGGCAAATAAGCACTCCTTTAAAGCCGCTGCCTTCTGGTTACAATTTACAGTGT  
 ACTTTGTTAAAAACTGCTGAGGCTTATAATCATGGCTCTTAGAAAATCAGGAAAAGAGGA  
 GATAATGTGGATTTAAACCTTAAGAGTTCTAACCATGCCTACTAAATGTACAGATATGCAAAA  
 TTCCATAGCTCAATAAAGAATCTGATACCTTAGACCAAAAAAAAAA

**FIGURE 82**

MSAAWI PALGLGVCLLLLLPGPAGSEGAAPIAITCFTRGLDIRKEKADVLC PGGCPLEEF SVY  
GNIVYASVSSICGAAVHRGVISNSGGPVRVYSLPGRENYSSVDANGIQSQMLSRWSASFTVT  
K GKSS TQEATGQAVSTAH PPTGKRLKKTPEKKTGNKDKADIAFLIDGSFNIGQRRFNLQKN  
FVGKVALMLGIGTEGPHVGLVQASEHPKIEFYLNFTSAKDVLFAIKEVGFRRGNSNTGKAL  
KHTAQKFFTFVDAGVRKGI PKVVVVFIDGWPSDDIEEAGIVAREFGVNVFIVSVAKPIPEELG  
MVQDVT FVDKAVCRNNGFFSYHMPNWF GTTKYVKPLVQKLC THEQMMCSKTCYNSVNIAFLI  
DGSSSVGDSNFRMLLEFVSNIAKTFEISDIGAKIAAVQFTYDQRTFESFTDYSTKENVLAVI  
RNIRYMSGGTATGDAISFTVRNVFGPIRESPNKNFLVIVTDGQSYDDVQGPAAAAHDAGIT I  
FSVGVAWAPLDDDLKDMASKPKESHAF FTREFTGLEPIVSDVIRGICRDFLESQQ

**Signal sequence:**

amino acids 1-24

**N-glycosylation site.**

amino acids 100-104, 221-225

**Casein kinase II phosphorylation site.**

amino acids 102-106, 129-133, 224-228, 316-320, 377-381, 420-424,  
425-429, 478-482, 528-532

**N-myristoylation site.**

amino acids 10-16, 23-29, 81-87, 135-141, 158-164, 205-211,  
239-245, 240-246, 261-267, 403-409, 442-448, 443-449

**Amidation site.**

amino acids 145-149

**FIGURE 83**

CGCCGCGCTCCCGCACCCGCGGCCCGCCACCGCGCCGCTCCCGCATCTGCACCCGAGCCC  
GGCGGCTCCCGGCGGGAGCGAGCAGATCCAGTCCGGCCCCGAGCGCAACTCGGTCCAGTCCG  
GGCGGCGGCTGCGGGCGCAGAGCGGAGATGCAGCGGCTTGGGGCCACCCTGCTGTGCCTGC  
TGCTGGCGGCGGCGGTCCCCACGGCCCCCGCGCCCGCTCCGACGGCGACCTCGGCTCCAGTC  
AAGCCCGGCCCGGCTCTCAGCTACCCGAGGAGGAGGCCACCCTCAATGAGATGTTCCGCGA  
GGTTGAGGAAGCTGATGGAGGACACGCAGCACAAATTGCGCAGCGCGGTGGAAGAGATGGAGG  
CAGAAGAAGCTGCTGCTAAAGCATCATCAGAAGTGAACCTGGCAAACCTACCTCCCAGCTAT  
CACAAATGAGACCAACACAGACACGAAGGTTGGAAATAATACCATCCATGTGCACCGAGAAAT  
TCACAAGATAACCAACAACAGACTGGACAAATGGTCTTTTTAGAGACAGTTATCACATCTG  
TGGGAGACGAAGAAGGCAGAGAGGAGCCACGAGTGCATCATCGACGAGGACTGTGGGCCAGC  
ATGTACTGCCAGTTTTCAGCTTCCAGTACACCTGCCAGCCATGCCGGGGCCAGAGGATGCT  
CTGCACCCGGGACAGTGTGCTGTGGAGACCAGCTGTGTGTCTGGGGTCACTGCACCAAAA  
TGGCCACCAGGGGAGCAATGGGACCATCTGTGACAACAGAGGGACTGCCAGCCGGGGCTG  
TGCTGTGCCCTCCAGAGAGGCTGCTGTTCCCTGTGTGCACACCCCTGCCCGTGGAGGGCGA  
GCTTTCATGACCCCGCCAGCCGGCTTCTGGACCTCATCACCTGGGAGCTAGAGCCTGATG  
GAGCCTTGGACCGATGCCCTTGTGCCAGTGGCCTCCTCTGCCAGCCCCACAGCCACAGCCTG  
GTGTATGTGTGCAAGCCGACCTTCGTGGGAGCCGTGACCAAGATGGGGAGATCCTGCTGCC  
CAGAGAGGTCCCCGATGAGTATGAAGTTGGCAGCTTCATGGAGGAGGTGCGCCAGGAGCTGG  
AGGACCTGGAGAGGAGCCTGACTGAAGAGATGGCGCTGGGGAGCCTGCCGCTGCCGCCGCT  
GCACTGCTGGGAGGGGAAGAGATTTAGATCTGGACCAGGCTGTGGGTAGATGTGCAATAGAA  
ATAGCTAATTTATTTCCCCAGGTGTGTGCTTTAGGCGTGGGCTGACCAGGCTTCTTCCCTACA  
TCTTCTTCCAGTAAGTTTCCCTCTGGCTTGACAGCATGAGGTGTTGTGCATTTGTTTCAGC  
TCCCCCAGGCTGTTCTCCAGGCTTCCAGTCTGGTGTCTGGGAGAGTCAGGCAGGGTTAAAC  
TGACGGAGCAGTTTCCACCCCTGTCCAGATTATTGGCTGCTTTGCCCTTACCAGTTGGCAG  
ACAGCCGTTTGTCTACATGGCTTTTGATAAATTGTTTGGAGGGAGGAGATGGAAACAATGTGG  
AGTCTCCCTCTGATTGGTTTTGGGAAATGTGGAGAAGAGTGCCTGCTTTGCCAAACATCAA  
CCTGGCAAAAATGCAACAATGAATTTCCACGCAGTTCTTCCATGGGCATAGGTAAGCTG  
TGCTTTCAGCTGTTGCAGATGAAATGTTCTGTTCCACCTGCATTACATGTGTTTATTCATCC  
AGCAGTGTGCTCAGCTCCTACCTCTGTGCCAGGGCAGCATTTCATATCCAAGATCAATTC  
CCTCTCTCAGCACAGCCTGGGGAGGGGTCAATTGTTCTCCTCGTCCATCAGGGATCTCAGAG  
GCTCAGAGACTGCAAGCTGCTTGCCCAAGTCACACAGCTAGTGAAGACCAGAGCAGTTTCAT  
CTGGTTGTGACTTAAGCTCAGTGTCTCTCCACTACCCACACCAGCCTTGGTGGCCACCAA  
AAGTGTCCCCAAAAGGAAGGAGAATGGGATTTTTCTTGAGGCATGCACATCTGGAATTAAG  
GTCAAACATAATCTCACATCCCTCTAAAAGTAACTACTGTTAGGAAACAGCAGTGTCTCAC  
AGTGTGGGGCAGCCGCTCCTTAATGAAGACAATGATATTGACACTGTCCCTCTTTGGCAGT  
TGCATTAGTAATTTGAAAGGTATATGACTGAGCGTAGCATAACAGGTTAACCTGCAGAAACA  
GTACTTAGGTAATTTGAGGGCGAGGATTATAAATGAAATTTGCAAAATCACTTAGCAGCAAC  
TGAAGACAATTTCAACCAGTGGAGAAAATCAAACCAGCAGGGCTGTGTGAAACATGGTT  
GTAATATGCGACTGCCAACACTGAACTCTACGCCACTCCACAAATGATGTTTTTCAGGTGTCA  
TGGACTGTTGCCACCATGTATTATCCAGAGTTCTTAAAGTTTAAAGTTGCACATGATTGTA  
TAAGCATGCTTTTGGAGTTTTAAATTTATGTATAAACATAAGTTGCATTTAGAAATCAAGC  
ATAAATCACTTCAACTGCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

**FIGURE 84**

MQRLGATLLCLLLAAVPTAPAPAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMEDTQ  
HKLRSAVEEMEAEAAAASSEVNLANLPPSYHNETNTDTKVGNNTIHVHREIHKITNNQTG  
QMFVSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQPCRQMLCTRDSECCG  
DQLCVWGHCTKMATRGSNGTICDNQRDCQPLCCAFQRLGFLFPVCTPLPVEGELCHDPASRL  
LDLITWELEPDGALDRPCASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILLPREVPDEYEV  
GSFMEEVRQELEDLERSLTEEMALGEPAAAAAALLGGEEI

**Signal sequence:**

amino acids 1-19

**N-glycosylation site.**

amino acids 96-100, 106-110, 121-125, 204-208

**Casein kinase II phosphorylation site.**

amino acids 46-50, 67-71, 98-102, 135-139, 206-210, 312-316,  
327-331

**N-myristoylation site.**

amino acids 202-208, 217-223

**Amidation site.**

amino acids 140-144

**FIGURE 85**

AAGGAGGCTGGGAGGAAAGAGGTAAGAAAGGTTAGAGAACCTACCTCACATCTCTCTGGGCTCAGAAGGACTCTG  
AAGATAACAATAAATTCAGCCCATCCACTCTCCTTCCCTCCCAAACACACATGTGCATGTACACACACATACA  
CACACATACACCTTCTCTCCTTCACTGAAGACTCACAGTCACTCACTCTGTGAGCAGGTCATAGAAAAGGACAC  
TAAAGCCTTAAGGACAGGCCTGGCCATTACCTCTGCAGCTCCTTTGGCTTGTGAGTCAAAAAACATGGGAGGGG  
CCAGGCACGGTGACTCACACCTGTAATCCCAGCATTTTGGGAGACCCGAGGTGAGCAGATCACTTGAGGTCAGGAG  
TTCGAGACCAGCCTGGCCAACATGGAGAAACCCCATCTCTACTAAAAATACAAAAATAGCCAGGAGTGGTGGC  
AGGTGCCTGTAATCCCAGTACTCAGGTGGCTGAGCCAGGAGAATCGCTTGAATCCAGGAGGCCGAGGATGCAGT  
CAGCTGAGTGCACCCTGCATCCAGCCTGGTGAACAATGAGACTCTGTCTCAAAACAAACAAACACGGGAGGA  
GGGTAGATACTGCTTCTCTGCAACTCCTTAACTCTGCATCCTTCTTCCAGGGCTGCCCTGATGGGGCCTG  
GCAATGACTGAGCAGGCCAGCCCCAGAGGACAAGGAAGAGAAGGCATATTGAGGAGGGCAAGAAGTGCACCCCG  
GTGTAGAATGACTGCCCTGGGAGGGTGGTCCCTTGGGCCCTGGCAGGGTTGCTGACCCTTACCCTGCAAAACACA  
AAGAGCAGGACTCCAGACTCCTTTGTGAATGGTCCCTGACCCTGACCACTGAGGCTTCTCGTGGCCCA  
ACTCTTGCTAGCTTGGGTGGCTGGTGCACCTGCCACTGTGCCCTGGTACCCTGGCATGTTCCCTGCCCCCTCA  
GTGTGCCTGCCAGATCCGGCCCTGGTATACGCCCCGCTCGTCTTACC CGAGGCTACCACTGTGGACTGCAATGA  
CCTATTCTGACGGCAGTCCCCCGGCACTCCCCGAGGCACACAGACCCTGCTCCTGCAGAGCAACAGCATTGT  
CCGTGTGGACCAGAGTGGCTGGGCTACCTGGTCAATCTCACAGAGCTGGACCTGTCAGAGAACTTTTCGGA  
TGCCCGAGACTGTGATTTCCATGCCCTGCCCCAGCTGCTGAGCCTGCACCTAGAGGAGAACCAGCTGACCCGGCT  
GGAGGACCACAGCTTTGCGAGGCTGGCCAGCCTACAGGAACCTATCTCAACCACAACCAGCTTACC CGCATCGC  
CCCCAGGGCCTTTCTGGCTCAGCAACTTGTGCGGCTGCACCTCAACTCAACCTCCTGAGGGCCATTGACAG  
CCGCTGGTTTGAATGCTGCCAACTTGGAGATACTCATGATTGGCGGCAACAAGGTAGATGCCATCCTGGACAT  
GAACCTCCGGCCCTGGCCAACCTGCGTAGCCTGGTGTGACAGGCATGAACCTGCCGGGAGATCTCCGACTATGC  
CCTGGAGGGGCTGCAAGGCTGGAGAGCCTCTCCTTCTATGACAACCAGCTGGCCCGGGTGCACAGGCGGGCACT  
GGAAACAGGTGCCCGGCTCAAGTTCCTAGACCTCAACAAGAACCCTCCAGCGGGTAGGGCCGGGGGACTTTGC  
CAACATGCTGCACCTTAAGGAGCTGGACTGAACAACATGGAGGAGCTGGTCTCCATCGACAAGTTTGGCCTGGT  
GAACCTCCCGAGCTGACCAAGCTGGACATCACAATAACCCACGGCTGTCTTCCATCCACCCCGCGCTTCCA  
CCACTGCCCCAGATGGAGACCTCATGCTCAACAACAACGCTCTCAGTGCTTGCACCCAGAGACCGTGGAGTC  
CCTGCCCAACCTGCAGGAGGTAGTCTCCACGGCAACCCCATCCGCTGTGACTGTGTCACTCCGCTGGGCCAATGC  
CACGGGCACCCGTGTCCGCTTCATCGAGCCGCAATCCACCCCTGTGTGCGGAGCCTCCGACCTCCAGGCCCTCC  
GGTCCGTGAGGTGCCCTTCCGGGAGATGACGGACCACTGTTTGGCCCTCATCTCCCACGAAGCTTCCCCCAAG  
CCTCCAGGTAGCCAGTGGAGAGAGCATGGTGTGCATTGCCGGGCACTGGCCGAACCCGAACCCGAGATCTACTG  
GGTCACTCCAGCTGGCTTCGACTGACACCTGCCCATGACAGGCAGGAGGTACCGGGTGTACCCCGAGGGGACCT  
GGAGCTGCCGAGGGTGACAGCAGAAAGAGGCAGGGCTATACACCTGTGTGGCCAGAACCTGGTGGGGGCTGACAC  
TAAGACGGTTAGTGTGGTTGTGGGCGTGCTCTCCTCCAGCCAGGCAGGACGAAGGACAGGGGCTGGAGCTCCG  
GGTGCAGGAGACCCACCCCTATCACATCCTGCTATCTTGGGTACCCCCACCAACACAGTGTCCACCAACCTCAC  
CTGGTCCAGTGCCTCCTCCCTCCGGGGCCAGGGGGCCACAGCTCTGGCCCGCTGCCCTCGGGGACCCACAGCTA  
CAACATTACCCGCTCCTTCCAGGCCACGGAGTACTGGGCTGCCCTGCAAGTGGCTTTGCTGATGCCACACCCA  
GTTGGCTTGTGATGGGCCAGGACCAAGAGGCCACTTCTTGCCACAGAGCCTTAGGGGATCGTCTGGGCTCAT  
TGCATCCTGGCTCTCCCTGTCTTCTCCTGGCAGCTGGGCTAGCGGCCACCTTGGCACAGGCAACCCAGGAA  
GGGTGTGGTGGGAGGCGGCTCTCCCTCCAGCCTGGGCTTTCTGGGGCTGGAGTGCCTTCTGTCCGGGTTGT  
GTCTGCTCCCTCTGCTTCCCTGGAATCCAGGGAGGAAGCTGCCAGATCCTCAGAAGGGGAGACACTGTTGCC  
ACCATTGTCTCAAAATCTTGAAGCTCAGCCTGTTCTCAGCAGTAGAGAAATCACTAGGACTACTTTTACCAA  
AGAGAAGCAGTCTGGCCAGATGCCCTGCCAGGAAAGGGACATGGACCCACGTCCTTGAGGCTGGCAGCTGGGC  
CAAGACAGATGGGCTTTGTGGCCCTGGGGTGCTTCTGCAGCCTTGAAAAAGTTGCCCTTACCTCCTAGGGTCA  
CCTCTGCTGCCATCTGAGGAACATCTCAAGGAACAGGAGGGACTTTGGCTAGAGCCTCTGCTCCCATCTT  
CTCTCTGCCAGAGGCTCCTGGGCCCTGGCTTGGCTGTCCCTACCTGTGTCCCGGGCTGCACCCCTTCTCTT  
TCTTCTCTGTACAGTCTCAGTTGCTTGTCTTGTGCTTCTGGGCAAGGGCTGAAGGAGGCCACTCCATCTCAC  
CTCGGGGGCTGCCCTCAATGTGGGAGTGACCCAGCCAGATCTGAAGGACATTTGGGAGAGGGATGCCAGGAA  
CGCTCATCTCAGAGCCTGGGCTCGGCATTCGAAGCTGACTTCTATAGGCAATTTGTACCTTTGTGGAGAA  
ATGTGTCACTCCCAACCCGATTCACTCTTCTCCTGTTTTGTAAAAATAAAAAATAAATAACAATAAA  
AAAA

## **FIGURE 86**

MRLLVAPLLLAWVAGATATVPVVPWHVPCPPQCACQIRPWYTPRSSYREATTVDCNDLFLTA  
VPPALPAGTQTLTLLQSNISIVRVDQSELGYLANLTELDSLQNSFSDARDCDFHALPQLLSLHL  
EENQLTRLEDHSFAGLASLQELYLNHNQLYRIAPRAFSGLSNLLRLHLNSNLLRAIDSRWFE  
MLPNLEILMIGGNKVDAILDMNFRPLANLRSLVLAGMNLREISDYALEGLQSLESLSFYDNQ  
LARVPRRALEQVPGLKFLDLNKNPLQRVGPDFANMLHLKELGLNNMEELVSIKDFALVNL  
ELTKLDITNNPRLSFIHPRAFHHLPQMETLMLNNALSALHQQTVESLPLNQEVGLHGPNIR  
CDCVIRWANATGTRVRFIEPOSTLCAEPPDLQRLPVREVPFREMTHCLPLISPRSFPPSLQ  
VASGESMVLHCRALAEPEPEIYWVTPAGLRLTPAHAGRRYRVYPEGTLELRRVTAEAGLYT  
CVAQNLVGADTKTVSVVGRALLQGRDEGQGLELRVQETHPHYILLSWVTPPNTVSTNLTW  
SSASSLRGQGATALARLPRGTHSYNITRLLQATEYWACLQVAFADAHTQLACVWARTKEATS  
CHRALGDRPGLIAILALAVLLLAAGLAAHLGTGQPRKGVGGRRPLPPAWAFWGSAPSVRV  
SAPLVLPWNPGRKLPRSSEGETLLPPLSQNS

**Signal sequence:**

amino acids 1-18

**Transmembrane domain:**

amino acids 629-648

**N-glycosylation site.**

amino acids 94-98, 381-385, 555-559, 583-587

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 485-489

**Casein kinase II phosphorylation site.**

amino acids 46-50, 51-55, 96-100, 104-108, 130-134, 142-146,  
243-247, 313-317, 488-492, 700-704

**Tyrosine kinase phosphorylation site.**

amino acids 532-540

**N-myristoylation site.**

amino acids 15-21, 493-499, 566-572

**Amidation site.**

amino acids 470-474, 660-664, 692-696



**FIGURE 87**

GCAAGCCAAGGGCGTGTGTTGAGAAGGTGAAGAAGTTCGGGACCCATGTGGAGGAGGGGGACATTGTGTACCGCCT  
CTACATGCGGCAGACCATCATCAAGGTGATCAAGTTCATCCTCATCATCTGCTACACCGTCTACTACGTGCACAA  
CATCAAGTTCGACGTGGACTGCACCGTGGACATTGAGAGCCTGACGGGCTACCGCACCTACCGCTGTGCCACCC  
CCTGGCCACACTCTCAAGATCCTGGCGTCCTTCTACATCAGCCTAGTTCATCTTCTACGGCCTCATCTGCATGTA  
CACACTGTGGTGGATGCTACGGCGTCCCTCAAGAAGTACTCGTTTGTAGTCGATCCGTGAGGAGAGCAGCTACAG  
CGACATCCCGACGTCAAGAACGACTTGCCTTCATGCTGCACCTCATTGACCAATACGACCCGCTCTACTCCAA  
GGCTTCGCGCTCTTCTGTGCGAGGTGAGTGAGAACAAGCTGCGGAGCTGAAACCTCAACAACGATGGACGCT  
GGACAAGTCCGGCAGCGGCTCAACAAGAACCGCAGGACAAGCTGGAGCTGCACCTGTTTATGCTCAGTGGCAT  
CCCTGACACTGTGTTGACCTGGTGGAGCTGGAGGTCCTCAAGCTGGAGCTGATCCCCGACGTGACCATCCCCG  
CAGCATTGCCAGCTCACGGCCTCAAGGAGCTGTGGCTTACCACACAGCGGCAAGATTGAAGCGCCTGCGCT  
GGCTTCGTGCGGAGAACCTGCGGGCGCTGCACATCAAGTTCACCGACATCAAGGAGATCCCCCACTCCATCTT  
TAGCCTGAAGACACTGGAGGAGCTGCACCTGACGGGCAACCTGAGCGCGGAGAACAACCGCTACATCGTTCATCGA  
CGGGCTGCGGGAGCTCAAAACGCTCAAGGTGCTGCGGCTCAAGAGCAACCTAAGCAAGCTGCCACAGGTGGTCA  
AGATGTGGGCGTGCACCTGCAGAAGCTGTCCATCAACAATGAGGGCACCAAGCTCATCGTCTCAACAGCCTCAA  
GAAGATGGCGAACCTGACTGAGCTGGAGTGCACCTGCGCTGCGACCTGGAGCGCATCCCCCACTCCATCTT  
CCACAACCTGCAGGAGATTGACCTCAAGGACAACAACCTCAAGACCATCGAGGAGATCATCAGCTTCCAGCACCT  
GCACCGCCTCACCTGCCTTAAGCTGTGTTACAACACATCGCCTACATCCCAATCCAGATCGGCAACCTCACCAA  
CCTGGAGCGCCTCTACTGAACCGCAACAAGATCGAGAAGATCCCCACCCAGCTCTTCTACTGCGCAAGCTGCG  
TACCTGGACCTCAGCCACAACAACCTGACCTTCCCTGCCGACATCGGCTCCTGCAGAACCTCCAGAACCT  
AGCCATCACGGCCAACCGGATCGAGACGCTCCCTCCGGAGCTTCCAGTGCCGGAAGCTGCGGGCCTGCACCT  
GGCAACAACGCTGCTGCAGTCACTCCCTCCAGGTTGGGCGAGCTGACCAACCTGACGCGAGATCGAGCTGCGGG  
CAACCGCTGGAGTGCCTGCCTGTGGAGCTGGGCGAGTGCCACTGCTCAAGCGCAGCGGCTTGGTGGTGGAGGA  
GGACCTGTTCAACACACTGCCACCCGAGGTGAAGGAGCGGCTGTGGAGGGCTGACAAGGAGCAGGCTGAGCGG  
GCCGGCCAGCACAGCAAGCAGCAGACCGCTGCCAGTCTCAGGCCGAGGGGAGGCTAGCTTCTCCAG  
AACTCCCGCAGCAGCCAGGACAGCCTCGCGGCTGGCAGGAGCCTGGGGCCGCTTGTGAGTCAGGCCAGAGCGAGA  
GGACAGTATCTGTGGGGTGGCCCTTTCTCCCTCTGAGACTCACGTCCCCAGGGCAAGTGTGTTGGAGGAG  
AGCAAGTCTCAAGAGCGCAGTATTGGATAATCAGGGTCTCCTCCTGGAGGCCAGCTGCCCCAGGGGCTGAG  
CTGCCACCAGAGGTCCTGGGACCTCACTTTAGTCTTGGTATTTATTTTCTCCATCTCCACCTCCTTCATCC  
AGATAACTTATACATTTCCAAGAAAGTTGAGCCAGATGGAAGGTGTTGAGGGAAAGGTGGGCTGCCTTTTCCC  
TTGTCCTATTTAGCGATGCCGCCGGGCATTTAACACCCACCTGGACTTCAGCAGAGTGGTCCGGGGCGAACCCAG  
CCATGGGACGGTCAACAGCAGTGCAGGCTGGGCTCTGCGGTGCGGTCCACGGGAGAGCAGGCTCCAGCTGGA  
AAGGCCAGGCTGGAGCTTGCTCTTTCAGTTTTTGTGGCAGTTTTAGTTTTTTTTTTTTTTTTTAATCAA  
AAACAATTTTTTTTTAAAAAAGCTTTGAAAATGGATGGTTGGGTATTAAGAAAGAAAAAAGCTTAAAAA  
AAAAGACACTAACGGCCAGTGTGAGTGTGAGTCTCAGGGCAGGGTGGCAGTTTTCCCTGAGCAAAGCAGCCAGACCT  
TGAACGTGTTTTCTTTCCCTGGGCGCAGGGTGCAGGGTGTCTTCCGGATCTGTGTGACCTTGGTCCAGGAGTT  
CTATTTGTTCTTGGGGAGGGAGGTTTTTTTTGTTGTTTTTTGGGTTTTTTTTGGTGTCTGTTTTCTTCTCCTCC  
ATGTGCTTGGCAGGCACTCATTTCTGTGGCTGTGCGCCAGAGGGAATGTTCTGGAGCTGCCAAGGAGGGAGGAG  
ACTCGGTTGGCTAATCCCCGGATGAACGGTGTCCATTGCGACCTCCCTCCTCGTGCCTGCCCTGCCTCTCCA  
CGCACAGTGTAAAGGAGCCAAGAGGAGCCACTTCGCCAGACTTTGTTTCCCACTCCTGCGGCATGGGTGTGT  
CCAGTGCCACCGCTGGCCTCCGCTGCTTCCATCAGCCCTGTGCGCACCTGGTCTTTCATGAAGAGCAGACCTTA  
GAGGCTGTTGCGGAATGGGGAGGTGCGCCCTGGGAGGGCAGGCGTTGGTTCAGCCGGTCCCGTCCCTGGCGC  
CTGGAGTGACACAGCCAGTGGCACCTGGTGGCTGGAAGCCAACCTGCTTTAGATCACTCGGGTCCCCACCTT  
AGAAGGTCCCCGCTTAGATCAATCACGTGGACACTAAGGCACGTTTTAGAGTCTCTTGTCTTAATGATTTATGT  
CCATCCGTCTGTCCGTCCATTTGTGTTTTCTGCGTGTGTCATTGGATATAATCCTCAGAAATAATGCACACTAG  
CCTCTGACAACCATGAAGCAAAAATCCGTTACATGTGGGTCTGAACCTGTAGACTCGGTACAGTATCAAATAAA  
ATCTATAACAGAAAAAAAAAAAAAAAA

**FIGURE 88**

MRQTIKVIKFIILIICTVYVYVHNIKFDVDCTVDIESLTGYRTRYCAHPLATLFKILASFYI  
SLVIFYGLICMYTLWWMRLRRSLKKYSFESIREESSYSDIPDVKNDFAFMLHLIDQYDPLYSK  
RFAVFLSEVSENKLRQLNLNNEWTLDKLRQLTKNAQDKLELHLFMLSIGIPDPTVFDLVELEV  
LKLELIPDVTIPPSIAQLTGLKELWLYHTAAKIEAPALAFLENLRLALHIKFTDIKEIPLWI  
YSLKTLLEELHLTGNSAENNRYIVIDGLRELKRLKVLRLKSNLSKLPQVVTDVGVHLQKLSI  
NNEGTKLIVLNSLKKMANLTELELIRCDLERIPHSIFSLHNLQEQIDLDKNNLKTIEEIIISFQ  
HLHRLTCLKLWYNHAIYIPIQIGNLTNLERLYLNRNKIEKIPTQLFYCRKLRYLDLSHNNLT  
FLPADIGLLQNLQNLAITANRIETLPPPELQCRKLRALHLGNNVLQSLPSRVGELTNLTQIE  
LRGNRLECLPVELGECPLLKRSGLVVEEDLFNTLPPEVKERLWRADKEQA

**Transmembrane domain:**

amino acids 51-75 (type II)

**N-glycosylation site.**

amino acids 262-266, 290-294, 328-332, 396-400, 432-436, 491-495

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 85-89

**Casein kinase II phosphorylation site.**

amino acids 91-95, 97-101, 177-181, 253-257, 330-334, 364-368,  
398-402, 493-497

**N-myristoylation site.**

amino acids 173-179, 261-267, 395-401, 441-447

**FIGURE 89**

GCCTGTTGCTGATGCTGCCGTGCCGTA~~CTTGT~~ATGGAGCTGGCACTGCGGCGCTCTCCCCT  
CCCCGGTGGTTGCTGCTGCTGCCGCTGCTGCTGGGCCTGAACGCAGGAGCTGTCA~~TTGACT~~  
GGCCACAGAGGAGGGCAAGGAAGTATGGGATTATGTGACGGTCCGCAAGGATGCCTACATG  
TTCTGGTGGCTCTATTATGCCACCAACTCCTGCAAGAACTTCTCAGAACTGCCCCCTGGTCAT  
GTGGCTTCAGGGCGGTCCAGGCGGTTCTAGCACTGGATTTGGAACTTTGAGGAAATTGGGC  
CCCTTGACAGTGATCTCAAACCACGGAAAACCACCTGGCTCCAGGCTGCCAGTCTCCTATTT  
GTGATAATCCCCTGGGCACTGGGTTCA~~GTTATGTGAATGGTAGTGGTGCCTATGCCAAGGA~~  
CCTGGCTATGGTGGCTTCAGACATGATGGTTCTCCTGAAGACCTTCTTCAGTTGCCACAAG  
AATTCAGACAGTTCATTCTACATTTTCTCAGAGTCCTATGGAGGAAAAATGGCAGCTGGC  
ATTGGTCTAGAGCTTTATAAGGCCATTCAAGTCAACTTTGCGGGGGT  
TGCCTTGGGTGATTCTGGATCTCCCCTGTTGATTCCGGTGCTCTCCTGGGGACCTTACCTGT  
ACAGCATGTCTTCTCGAAGACAAAGGTCTGGCAGAGGTGTCTAAGGTTGCAGAGCAAGTA  
CTGAATGCCGTAAATAAGGGGCTCTACAGAGAGGCCACAGAGCTGTGGGGGAAAGCAGAAAT  
GATCATTGAACAGAACACAGATGGGGTGA~~ACTTCTATAACATCTTAACTAAAAGCACTCCCA~~  
CGTCTACAATGGAGTCGAGTCTAGAATTCACACAGAGCCACCTAGFTTGTCTTTGTCAGCGC  
CACGTGAGACACCTACAACGAGATGCCTTAAGCCAGCTCATGAATGGCCCCATCAGAAAGAA  
GCTCAA~~AATTATT~~CCTGAGGATCAATCCTGGGGAGGCCAGGCTACCAACGTCTTTGTGAACA  
TGGAGGAGGACTTCATGAAGCCAGTCATTAGCATTGTGGACGAGTTGCTGGAGGCAGGGATC  
AACGTGACGGTGTATAATGGACAGCTGGATCTCATCGTAGATACCATGGGTGAGGAGGCCCTG  
GGTGC~~GGAAACTGAAGTGGCCAGAACTGCCTAAATT~~CAGTCAGCTGAAGTGAAGGCCCTGT  
ACAGTGACCCTAAATCTTTGGAAACATCTGCTTTTGTCAAGTCTACAAGAACCTTGCTTTC  
TACTGGATTCTGAAAGCTGGTTCATATGGTTCTTCTGACCAAGGGGACATGGCTCTGAAGAT  
GATGAGACTGGTGACTCAGCAAGAATAGGATGGATGGGGCTGGAGATGAGCTGGTTTGGCCT  
TGGGGCACAGAGCTGAGCTGAGGCCGCTGAAGCTGTAGGAAGCGCCATTCTTCCCTGTATCT  
AACTGGGGCTGTGATCAAGAAGGTTCTGACCAGCTTCTGCAGAGGATAAAATCATTGTCTCT  
GGAGGCAATTTGGAAATTATTCTGCTTCTTAAAAAACCTAAGATTTTTTAAAAAATTGAT  
TTGTTTTGATCAAAATAAAGGATGATAATAGATATTAA

**FIGURE 90**

MELALRRSPVPRWLLLLLPLLLGLNAGAVIDWPTEEGKEVWDYVTVRKDAYMFWWLYYATNSC  
KNFSELPLVMWLQGGPGGSSTGFGNFEEIGPLDSDLKPRKTTWLQAASLLFVDNPNVGTGFSY  
VNGSGAYAKDLAMVASDMMVLLKTTFFSCHKEFQTVPFYIFSESYGGKMAAGIGLELYKAIQR  
GTIKCNFAGVALGDSWISPVDSVLSWGPYLYSMSLLEDKGLAEVSKVAEQVLNAVNKGLYRE  
ATELWGKAEMII EQNTDGVNFYNI LTKSTPTSTMESLEFTQSHLVCLQRHVRHLQDALS  
QLMNGPIRKKLKIIPEDQSWGQATNVFVNMEEDFMKPVISIVDELLEAGINVTVYNGQLDL  
IVDTMGQEAWVRKWKWPELPKFSQLKWKALYSDPKSLETSFAFKSYKNLAFYWILKAGHMVP  
SDQGDMAKMMRLVTQQE

**Signal sequence:**

amino acids 1-25

**N-glycosylation site.**

amino acids 64-68, 126-130, 362-366

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 101-105

**Casein kinase II phosphorylation site.**

amino acids 204-208, 220-224, 280-284, 284-288, 351-355, 449-453

**N-myristoylation site.**

amino acids 22-28, 76-82, 79-85, 80-86, 119-125, 169-175,  
187-193, 195-201, 331-337, 332-338, 360-366

**FIGURE 91**

GGCCGCGGGAGAGGAGGCCATGGGCGCGCGGGGCGCTGCTGCTGGCGCTGCTGCTGGCTC  
GGGCTGGACTCAGGAAGCCGGAGTCGCAGGAGGCGGGCGCCGTTATCAGGACCATGCGGCCGA  
CGGGTCATCACGTTCGCGCATCGTGGGTGGAGAGGACGCCGAAGTTCGGGCGTTGGCCGTGGCA  
GGGGAGCCTGCGCCTGTGGGATTCCCACGTATGCGGAGTGAGCCTGCTCAGCCACCGCTGGG  
CACTCACGGCGGGCGCACTGCTTTGAAAACCTATAGTGACCTTAGTGATCCCTCCGGGTGGATG  
GTCCAGTTTGGCCAGCTGACTTCCATGCCATCCTTCTGGAGCCTGCAGGCCTACTACACCCG  
TTACTTCGTATCGAATATCTATCTGAGCCCTCGCTACCTGGGGAATTCACCCTATGACATTG  
CCTTGGTGAAGCTGTCTGCACCTGTCACCTACACTAAACACATCCAGCCCATCTGTCTCCAG  
GCCTCCACATTTGAGTTTGGAGAACCGGACAGACTGCTGGGTGACTGGCTGGGGGTACATCAA  
AGAGGATGAGGCACTGCCATCTCCCCACACCCTCCAGGAAGTTCAGGTCGCCATCATAAACA  
ACTCTATGTGCAACCACCTCTTCCCTCAAGTACAGTTTCCGCAAGGACATCTTTGGAGACATG  
GTTTGTGCTGGCAACGCCCAAGGCGGGAAGGATGCCTGCTTCGGTGACTCAGGTGGACCCTT  
GGCCTGTAACAAGAATGGACTGTGGTATCAGATTGGAGTCGTGAGCTGGGGAGTGGGCTGTG  
GTCGGCCCAATCGGCCCGGTGTCTACACCAATATCAGCCACCACTTTGAGTGGATCCAGAAG  
CTGATGGCCCAGAGTGGCATGTCCAGCCAGACCCCTCCTGGCCACTACTCTTTTTCCCTCT  
TCTCTGGGCTCTCCCACTCCTGGGGCCGGTTCAGCCTACCTGAGCCCATGCAGCCTGGGGC  
CACTGCCAAGTCAGGCCCTGGTTCTCTTCTGTCTTGTTTGGTAATAAACACATTCAGTTGA  
TGCCTTGCAGGGCATTCTTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

**FIGURE 92**

MGARGALLLALLLLARAGLRKPEAQEAPLSGPCGRRVITSRIVGGEDAELGRWPWQGSRLRW  
DSHVCVSVLLSHRWALTAAHCFETYSDLSDPSGWMVQFGQLTSMPSFWSLQAYYTRYFVSNI  
YLSPRYLGNSPYDIALVKLSAPVYTKHIQPICLOASTFEFENRTDCWVTGWGYIKEDEALP  
SPHTLQEVQVAIINNSMCNHLFLKYSFRKDI FGDMVCAGNAQGGKDACFGDSGGPLACNKNG  
LWYQIGVSVWGVGCGRPNRPGVYTNISHHFEWIQKLMAQSGMSQPDPSWPLLFFPLLWALPL  
LGPV

**Signal sequence:**

amino acids 1-18

**N-glycosylation site.**

amino acids 167-171, 200-204, 273-277

**Casein kinase II phosphorylation site.**

amino acids 86-90, 134-138, 161-165, 190-194, 291-295

**N-myristoylation site.**

amino acids 2-8, 44-50, 101-107, 225-231, 229-235, 239-245,  
259-265, 269-275

**Amidation site.**

amino acids 33-37

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 252-263,

**Serine proteases, trypsin family, histidine active site.**

amino acids 78-84

**FIGURE 93**

CCCACGCGTCCGCGGACGCGTGGGAAGGGCAGAATGGGACTCCAAGCCTGCCTCCTAGGGCT  
CTTTGCCCTCATCCTCTCTGGCAAATGCAGTTACAGCCCCGGAGCCCCGACCAGCGGAGGACGC  
TGCCCCCAGGCTGGGTGTCCCTGGGCCGTGCGGACCCTGAGGAAGAGCTGAGTCTCACCTTT  
GCCCTGAGACAGCAGAATGTGGAAAGACTCTCGGAGCTGGTGCAGGCTGTGTGGATCCCAG  
CTCTCCTCAATACGGAATAACCTGACCCTAGAGAATGTGGCTGATCTGGTGAGGCCATCCC  
CACTGACCCTCCACACGGTGCAAAAATGGCTCTTGGCAGCCGGAGCCCAGAAGTGCCATTCT  
GTGATCACACAGGACTTCTGACTTGCTGGCTGAGCATCCGACAAGCAGAGCTGCTGCTCCC  
TGGGGCTGAGTTTCATCACTATGTGGGAGGACCTACGGAACCCATGTTGTAAGGTCCCCAC  
ATCCCTACCAGCTTCCACAGGCTTGGCCCCCATGTGGACTTTGTGGGGGGACTGCACCGT  
TTTCCCCAACATCATCCCTGAGGCAACGTCTCTGAGCCGACGGTGACAGGGACTGTAGGCCCT  
GCATCTGGGGGTAACCCCTCTGTGATCCGTAAGCGATACAACTTGACCTACAAGACGTGG  
GCTCTGGCACCAGCAATAACAGCCAAGCCTGTGCCAGTTCTGGAGCAGTATTTCCATGAC  
TCAGACCTGGCTCAGTTCATGCGCCTCTTCGGTGGCAACTTTGCACATCAGGCATCAGTAGC  
CCGTGTGGTTGGACAACAGGGCCGGGGCCGGGGCCGGGATTGAGGCCAGTCTAGATGTGCAGT  
ACCTGATGAGTGTGGTGCCAACATCTCCACCTGGGTCTACAGTAGCCCTGGCCGGCATGAG  
GGACAGGAGCCCTTCTGCAAGTGGCTCATGCTGCTCAGTAATGAGTCAGCCCTGCCACATGT  
GCATACTGTGAGTATGGAGATGATGAGGACTCCCTCAGCAGCGCCTACATCCAGCGGGTCA  
ACACTGAGCTCATGAAGGCTGCCGCTCGGGGTCTCACCTGCTCTTCGCCTCAGGTGACAGT  
GGGGCCGGGTGTTGGTCTGTCTCTGGAAGACACCAGTTCGCCCTACCTTCCCTGCCTCCAG  
CCCCTATGTCACCACAGTGGGAGGCACATCCTTCCAGGAACCTTTCTCATCAAAATGAAA  
TTGTTGACTATATCAGTGGTGGTGGCTTCAGCAATGTGTTCCACGGCCTTCATACCAGGAG  
GAAGCTGTAACGAAGTTCTGAGCTCTAGCCCCACCTGCCACCATCCAGTTACTTCAATGC  
CAGTGGCCGTGCCATACCAGATGTGGCTGCACTTTCTGATGGCTACTGGGTGGTGCAGCAACA  
GAGTGCCCATTCATGGGTGTCCGGAACCTCGGCCTCTACTCCAGTGTFTGGGGGGATCCTA  
TCCTTGATCAATGAGCACAGGATCCTTAGTGGCCGCCCCCTCTGGCTTTCTCAACCCAAG  
GCTCTACCAGCAGCATGGGGCAGGTCTCTTTGATGTAACCCGTGGCTGCCATGAGTCCCTGTC  
TGGATGAAGAGGTAGAGGGCCAGGGTTTCTGCTCTGGTCTGGCTGGGATCCTGTAAACAGGC  
TGGGGAACACCAACTTCCAGCTTTGCTGAAGACTCTACTCAACCCCTGACCTTTTCTATC  
AGGAGAGATGGCTTGTCCCCTGCCCTGAAGCTGGCAGTTCAGTCCCTTATTCTGCCCTGTTG  
GAAGCCCTGCTGAACCCCTCAACTATTGACTGCTGCAGACAGCTTATCTCCCTAACCCCTGAAA  
TGCTGTGAGCTTGACTTGACTCCCAACCCCTACCATGCTCCATCATACTCAGGTCTCCCTACT  
CCTGCCTTAGATTCCTCAATAAGATGCTGTAAGTACTAGCATTTTTTGAATGCCTCTCCCTCCGC  
ATCTCATCTTTCTCTTTTCAATCAGGCTTTTCCAAAGGGTTGTATACAGACTCTGTGCACCTA  
TTTCACTTGATATTCAATCCCAATTCACCTGCAAGGAGACCTCTACTGTCACCGTTTACTCT  
TTCTTACCCTGACATCCAGAAACAATGGCCTCCAGTGCATACTTCTCAATCTTTGCTTTATG  
GCCTTTCCATCATAGTTGCCCACTCCCTCTCCTTACTTAGCTTCCAGGTCTTAACCTCTCTG  
ACTACTCTTGTCTTCTCTCATCAATTTCTGCTTCTTCATGGAATGCTGACCTTCATTGC  
TCCATTTGTAGATTTTTGCTCTTCTCAGTTTACTCATTGTCCCCTGGAACAAATCACTGACA  
TCTACAACCATTACCATCTCACTAAATAAGACTTTCTATCCAATAATGATTGATACCTCAA  
TGTAAAAAA

**FIGURE 94**

MGLQACLLGLFALILSGKCSYSPEPDQRRTLPPGWVSLGRADPEEELSLTFALRQQNVERLS  
ELVQAVSDPSSPQYGKYLTLENVADLVRPSPLTLHTVQKWLLAAGAQQKCHSVITQDFLTCWL  
SIRQAELLLPGAEFHHYVGGPTETHVVRSPHPYQLPQALAPHVDFVGG LHRFPPTSSLRQRP  
EPQVTGTVGLHLGVTPSVIRKRYNLTSQDVGSGTSNNSQACAQFLEQYFHSDLAQFMRLFG  
GNFAHQASVARVVGQQGRGRAGIEASLDVQYLSAGANISTWVYSSPGRHEGQEPFLQWLML  
LSNESALPHVHTVSYGDDSDLSAYIQRVNTELMKAAARGLTLLFASGDSGAGCWSVSGRH  
QFRPTFPASSPYVTTVGGTSFQEPFLITNEIVDYISGGGFSNVFPRPSYQEEAVTKFLSSSP  
HLPPSSYFNASGRAYPDVAALSDGYWVSNRVPWPVSGTSASTPVFGGILSLINEHRILSG  
RPPLGFLNPRLYQQHGAGLFDVTRGCHESCLDEEVEGQGFCSGPGWDPVTGWGTPTSQLC

**Signal sequence:**

amino acids 1-16

**N-glycosylation site.**

amino acids 210-214, 222-226, 286-290, 313-317, 443-447

**Glycosaminoglycan attachment site.**

amino acids 361-365, 408-412, 538-542

**Casein kinase II phosphorylation site.**

amino acids 212-216, 324-328, 392-396, 420-424, 525-529

**N-myristoylation site.**

amino acids 2-8, 107-113, 195-201, 199-205, 217-223, 219-225,  
248-254, 270-276, 284-290, 409-415, 410-416, 473-479, 482-488,  
521-527, 533-539, 549-555



**FIGURE 95**

GCCGCGCGCTCTCTCCCGGCGCCACACCTGTCTGAGCGGCGCAGCGAGCCGCGGCCCGGGC  
GGGCTGCTCGGCGCGGAACAGTGTCTGGCAATGGCAGGGATTCCAGGGCTCCTCTTCCTTCTC  
TTCTTTCTGCTCTGTGCTGTTGGGCAAGTGAGCCCTTACAGTGCCCCCTGGAACCCACTTG  
GCCTGCATACCGCTCCCTGTGCTCTTGCCCCAGTCTACCCTCAATTTAGCCAAGCCAGACT  
TTGGAGCCGAAGCCAAATTAGAAGTATCTTCTTCATGTFGGACCCCAGTGTGATAAGGGA  
CCACTGCCACTTACGAAGAGGCCAAGCAATATCTGTCTTATGAAACGCTCTATGCCAATGG  
CAGCCGCACAGAGACGCAGGTGGGCATCTACATCCTCAGCAGTAGTGGAGATGGGGCCCAAC  
ACCGAGACTCAGGGTCTTCAGGAAAGTCTCGAAGGAAGCGGCAGATTTATGGCTATGACAGC  
AGGTTTCAGCATTTTTGGGAAGGACTTCTGTCTCAACTACCCTTTCTCAACATCAGTGAAGTT  
ATCCACGGGCTGCACCGGCACCCTGGTGGCAGAGAAGCATGTCTTCACAGCTGCCCACTGCA  
TACACGATGGAAAACCTATGTGAAAGGAACCCAGAAGCTTCGAGTGGGCTTCCATAAGCCC  
AAGTTTAAAGATGGTGGTTCGAGGGGCCAACGACTCCACTTCAGCCATGCCCGAGCAGATGAA  
ATTTTCAGTGGATCCGGGTGAAACGCACCCATGTGCCAAGGGTTGGATCAAGGGCAATGCCA  
ATGACATCGGCATGGATTATGATTATGCCCTCCTGGAACCTCAAAAAGCCCCACAAGAGAAAA  
TTTATGAAGATTGGGGTGAGCCCTCTGCTAAGCAGCTGCCAGGGGGCAGAATTCACTTCTC  
TGTTTATGACAATGACCGACCAGGCAATTTGGTGTATCGCTTCTGTGACGTCAAAGACGAGA  
CCTATGACTTGCTCTACCAGCAATGCGATGCCAGCCAGGGGCCAGCGGGTCTGGGGTCTAT  
GTGAGGATGTGGAAGAGACAGCAGCAGAAAGTGGGAGCGAAAAATTATTGGCATTTTTTTCAGG  
GCACCAGTGGGTGGACATGAATGGTTCACACAGGATTTCAACGTGGCTGTGAGAATCACTC  
CTCTCAAATATGCCAGATTTGCTATTTGGATTAAAGGAACTACCTGGATTGTAGGGAGGGG  
TGACCACAGTGTTCCTCCTGGCAGCAATTAAGGGTCTTCATGTTCTTATTTTAGGAGAGGCC  
AAATTGTTTTTTGTCAATTGGCGTGACACCGTGTGTGTGTGTGTGTGTGTGTGTAAGGTGT  
CTTATAATCTTTTACCTATTTCTTACAATTGCAAGATGACTGGCTTTACTATTTGAAAAC TG  
GTTTGTGTATCATATCATATATCATTTAAGCAGTTTGAAGGCATACTTTTGCATAGAAATAA  
AAAAAATACTGATTTGGGGCAATGAGGAATATTTGACAATTAAGTTAATCTTCACGTTTTTG  
CAAACCTTGATTTTTTATTTTCATCTGAACTTGTTCAAAAGATTTATATTAATATTTGGCATA  
CAAGAGATATGAAAAAAAAAAAAAAAAA

**FIGURE 96**

MAGIPGLLFLFFLLCAVGVSPYSAPWKPTWPAYRLPVVLPQSTLNLAKPDFGAEAKLEVS  
SSCGPQCHKGTPLPTYEEAKQYLSYETLYANGSRTETQVGIYILSSSGDGAQHRDSGSSGKS  
RRKRQIYGYDSRFSIFGKDFLLNYPFSTSVKLSSTGCTGTLVAEKHVLTAAHCIHDGKTYVKG  
TQKLRVGFLLKPKFKDGGRGANDSTSAMPEQMKFQWIRVKRTHVPGWIKGNANDIGMDYDYA  
LLELKKPHKRKFMKIGVSPPAKQLPGGRIHFSGYDNDRPGNLVYRFCDVKDETYDLLYQQCD  
AQPASGSGVYVRMWRQOQKWERKIIGIFSGHQWVDMNGSPQDFNVAVRITPLKYAQICYW  
IKGNYLDCREG

**Signal sequence:**

amino acids 1-19

**N-glycosylation site.**

amino acids 93-97, 207-211

**Glycosaminoglycan attachment site.**

amino acids 109-113, 316-320

**Casein kinase II phosphorylation site.**

amino acids 77-81, 95-99, 108-112, 280-284, 351-355

**N-myristoylation site.**

amino acids 159-165, 162-168, 202-208, 205-211, 314-320, 338-344

**Serine proteases, trypsin family, histidine active site.**

amino acids 171-177

**FIGURE 97**

GCATCGCCCTGGGTCTCTCGAGCCTGCTGCCTGCTCCCCCGCCCCACCAGCCATGGTGGTTT  
CTGGAGCGCCCCCAGCCCTGGGTGGGGGCTGTCTCGGCACCTTCACCTCCCTGCTGCTGCTG  
GCGTCGACAGCCATCCTCAATGCGGCCAGGATACCTGTTCCCCCAGCCTGTGGGAAGCCCCA  
GCAGCTGAACCGGTTGTGGGCGGCGAGGACAGCACTGACAGCGAGTGGCCCTGGATCGTGA  
GCATCCAGAAGAATGGGACCCACCACTGCGCAGGTTCTCTGCTCACCAGCCGCTGGGTGATC  
ACTGCTGCCCCTGTTTTCAAGGACAACCTGAACAAACCATACTGTTCTCTGTGCTGCTGGG  
GGCCTGGCAGCTGGGGAACCCCTGGCTCTCGGTCCCAGAAAGGTGGGTGTTGCCCTGGGTGGAGC  
CCCACCCTGTGTATTCTTGAAAGGAAGTGCCTGTGCAGACATTGCCCTGGTGCCTCTCGAG  
CGCTCCATACAGTTCTCAGAGCGGGTCTGCCCATCTGCCTACCTGATGCCTCTATCCACCT  
CCCTCCAAACACCCACTGCTGGATCTCAGGCTGGGGGAGCATCCAAGATGGAGTTCCTTGC  
CCCACCCTCAGACCCTGCAGAAGCTGAAGGTTCCCTATCATCGACTCGGAAGTCTGCAGCCAT  
CTGTACTGGCGGGGAGCAGGACAGGGACCCATCACTGAGGACATGCTGTGTGCCGGCTACTT  
GGAGGGGGAGCGGGATGCTTGTCTGGGCGACTCCGGGGGGCCCCCTCATGTGCCAGGTGGACG  
GCGCCTGGCTGCTGGCCGGCATCATCAGCTGGGGCGAGGGCTGTGCCGAGCGCAACAGGCCC  
GGGGTCTACATCAGCCTCTCTGCGCACCGCTCCTGGGTGGAGAAGATCGTGCAAGGGGTGCA  
GCTCCGCGGGCGCGCTCAGGGGGTGGGGCCCTCAGGGCACCGAGCCAGGGCTCTGGGGCCG  
CCGCGCGCTCTAGGGCGCAGCGGGACGCGGGGCTCGGATCTGAAAGGCGGCCAGATCCACA  
TCTGGATCTGGATCTGCGGCGGCCTCGGGCGGTTTCCCCCGCCGTAAATAGGCTCATCTACC  
TCTACCTCTGGGGGCCCGGACGGCTGCTGCGGAAAGGAAAACCCCTCCCCGACCCGCCCGAC  
GGCCTCAGGCCCCCTCCAAGGCATCAGGCCCCGCCAACGGCTCATGTCCCCGCCCCAC  
GACTTCCGGCCCCGCCCCGGGCCCCAGCGCTTTTGTGTATATAAATGTTAATGATTTTTAT  
AGGTATTTGTAACCCTGCCACATATCTTATTTATTCCTCCAATTTCAATAAATTATTTATT  
CTCCAAAAA

## **FIGURE 98**

></usr/seqdb2/sst/DNA/Dnaseqs.full/ss.DNA43318

><subunit 1 of 1, 317 aa, 1 stop

><MW: 33732, pI: 7.90, NX(S/T): 1

MVVGAPPALGGGCLGTFSTLSTLLASTAILNAARIPVPPACGKQPQLNRVVGGEDSTDSEWP  
WIVSIQKNGTHHCAGSLLTSRWVITAAHCFKDNLNKPYLFSVLLGAWQLGNPGRSRSQKVGVA  
WVEPHPVYSWKEGACADIALVRLERSIQFSERVLPICLPDASIHLPNTHCWISGWGSIQDG  
VPLPHPQTLQKLVPIIDSEVCSHLYWRGAGQGPIEDMLCAGYLEGERDACLGDSGGPLMC  
QVDGAWLLAGIISWEGCAERNRPGVYISLSAHRSWVEKIVQGVQLRGRAQGGGALRAPSQG  
SGAAARS

### **Signal sequence:**

amino acids 1-32

### **N-glycosylation site.**

amino acids 62-66, 96-100, 214-218, 382-386, 409-413, 455-459,  
628-632, 669-673, 845-849, 927-931, 939-943, 956-960

### **Glycosaminoglycan attachment site.**

amino acids 826-830

### **Casein kinase II phosphorylation site.**

amino acids 17-21, 39-43, 120-124, 203-207, 254-258, 264-268,  
314-318, 323-327, 347-351, 464-468, 548-552, 632-636, 649-653,  
671-675, 739-743, 783-787, 803-807, 847-851, 943-947, 958-962,  
1013-1017, 1019-1023, 1021-1025

### **Tyrosine kinase phosphorylation site.**

amino acids 607-615

### **N-myristoylation site.**

amino acids 179-185, 197-203, 320-326, 367-373, 453-459, 528-534,  
612-618, 623-629, 714-720, 873-879

**FIGURE 99**

GACGGCTGGCCACCATGCACGGCTCCTGCAGTTTCCTGATGCTTCTGCTGCCGCTACTGCTA  
CTGCTGGTGGCCACCACAGGCCCGTTGGAGCCCTCACAGATGAGGAGAAACGTTTGATGGT  
GGAGCTGCACAACCTCTACCGGGCCAGGTATCCCCGACGGCCTCAGACATGCTGCACATGA  
GATGGGACGAGGAGCTGGCCGCTTCGCCAAGGCCCTACGCACGGCAGTGCCTGTGGGGCCAC  
AAKAAGGAGCGCGGGCGCCGCGGCGAGAATCTGTTCCGCATCACAGACGAGGGCATGGACGT  
GCCGCTGGCCATGGAGGAGTGGCACCACGAGCGTGAGCACTACAACCTCAGCGCCGCCACCT  
GCAGCCCAGGCCAGATGTGCGGCCACTACACGCAGGTGGTATGGGCCAAGACAGAGAGGATC  
GGCTGTGGTTCCCCTTCTGTGAGAAGCTCCAGGGTGTGAGGAGACCAACATCGAATTA  
GGTGTGCAACTATGAGCCTCCGGGAAACGTGAAGGGGAAACGGCCCTACCAGGAGGGGACTC  
CGTGCTCCCAATGTCCTCTGGCTACCCTGCAAGAATCCCTCTGTGAACCCATCGGAAGC  
CCGGAAGATGCTCAGGATTTGCCCTTACCTGGTAACCTGAGGCCCATCCTTCCGGGCGACTGA  
AGCATCAGACTCTAGGAAAAATGGGTACTCCTTCTTCCCTAGCAACGGGGATTCCGGCTTTCT  
TGGTAACAGAGGTCTCAGGCTCCCTGGCAACCAAGGCTCTGCCTGCTGTGGAAACCCAGGCC  
CCAACCTCCTTAGCAACGAAAGACCCGCCCTCCATGGCAACAGAGGCTCCACCTTGCGTAAC  
AACTGAGGTCCCTTCCATTTTGGCAGCTCACAGCCTGCCCTCCTTGGATGAGGAGCCAGTTA  
CCTTCCCCAAATCGACCCATGTTCTTATCCCCAAATCAGCAGACAAAGTGACAGACAAAACA  
AAAGTGCCCTCTAGGAGCCCAGAGAATCTCTGGACCCCAAGATGTCCCTGACAGGGGCAAG  
GGAATCCTACCCCATGCCAGGAGGAGGCTGAGGCTGAGGCTGAGTTGCCTCCTTCCAGTG  
AGGFCTTGGCCTCAGTTTTTCCAGCCCAGGACAAGCCAGGTGAGCTGCAGGCCACACTGGAC  
CACACGGGGCACACCTCCTCCAAGTCCCTGCCCAATTTCCCCAATACCTCTGCCACCGCTAA  
TGCCACGGGTGGGCGTGCCCTGGCTCTGCAGTCGTCCTTGCCAGGTGCAGAGGGCCCTGACA  
AGCCTAGCGTTGTGTCAGGGCTGAACTCGGGCCCTGGTCATGTGTGGGGCCCTCTCCTGGGA  
CTACTGCTCCTGCCTCCTCTGGTGTGGCTGGAATCTTCTGAATGGGATACCACTCAAAGGG  
TGAAGAGGTCAGCTGTCCCTCCTGTCATCTTCCCCACCCTGTCCCCAGCCCCTAAACAAGATA  
CTTCTTGGTTAAGGCCCTCCGGAAGGGAAAGGCTACGGGGCATGTGCCTCATCACACCATCC  
ATCCTGGAGGCACAAGGCTGGCTGGCTGCGAGCTCAGGAGGCCGCTGAGGACTGCACACC  
GGGCCACACCTCTCCTGCCCTCCTCCTGAGTCTGGGGGTGGGAGGATTTGAGGGAGCT  
CACTGCCTACCTGGCCTGGGGCTGTCTGCCACACAGCATGTGCGCTCTCCCTGAGTGCCTG  
TGTAGCTGGGGATGGGGATTCTTAGGGGCAGATGAAGGACAAGCCCCACTGGAGTGGGGTTTC  
TTTGAGTGGGGGAGGCAGGGACGAGGGAAGGAAAGTAACTCCTGACTCTCCAATAAAAACT  
GTCCAACCTGTGAAA

## **FIGURE 100**

MHGSCSFLMLLLPLLLLLLVATTGPVGAITDEEKRLMVELHNLRYAQVSPITASDMLHMRWDEE  
LAFAKAYARQCVWGHNKERGRGENLFAITDEGMDVPLAMEEWHHEREHYNLSAATCSPGQ  
MCGHYTQVVWAKTERIGCGSHFCEKLOQVEETNIELLVVCNYEPPGNVKGKRPYQEGTPCSQC  
PSGYHCKNSLCEPIGSPEDAQDLPLYLVTEAPSFRAEASDSRKMGTTPSSLATGIPAFVLVTEV  
SGSLATKALPAVETQAPTSLATKDPSPMATEAPPVTEVPSILAAHSLPSLDEEPVTFPKS  
THVPIPKSADKVTDKTKVPSRSPENSLDPKMSLTGARELLPHAQEEAEAEELPPSSEVLAS  
VFPAQDKPGELQATLDHTGHTSSKSLPNFPNTSATANATGGRALALQSSSLPGAEGPDKPSV  
SGLNSGPGHVWGPLLGLLLLLPPLVLAGIF

**Signal sequence:**

amino acids 1-22

**N-glycosylation site.**

amino acids 114-118, 403-407, 409-413

**Glycosaminoglycan attachment site.**

amino acids 439-443

**Casein kinase II phosphorylation site.**

amino acids 29-33, 50-54, 156-160, 195-199, 202-206, 299-303

**N-myristoylation site.**

amino acids 123-129, 143-149, 152-158, 169-175, 180-186, 231-237,  
250-256

**Amidation site.**

amino acids 82-86, 172-176

**Peroxidases proximal heme-ligand signature.**

amino acids 287-298

**Extracellular proteins SCP/Tpx-1/Ag5/PR-1/Sc7 signature 1.**

amino acids 127-138

**Extracellular proteins SCP/Tpx-1/Ag5/PR-1/Sc7 signature 2.**

amino acids 160-172

**FIGURE 101**

GTAACTGAAGTCAGGCTTTTCATTGGGAAGCCCCCTCAACAGAATTCCGTCATTCTCCAAGTTATGTTGGACGT  
 ACTTCTGTTGTTCTCCCTCGCTTGCTTTTTCACATTAGCAGACCCGGACTTAAGTCACAACAGATTATCTTTCAT  
 CAAGGCAAGTTCATGAGCCACCTTCAAAGCCTTCGAGAAGTGAACCTGAACAACAATGAATTGGAGACCATTCC  
 AAATCTGGGACCAGTCTCGGCAAAATATTACACTTCTCTCCTTGGCTGGAAACAGGATTTGTTGAAATACTCCCTGA  
 ACATCTGAAAGAGTTTCAGTCCCTTGAACCTTTGGACCTTAGCAGCAACAATATTTAGAGCTCCAACTGCATT  
 TCCAGCCCTACAGCTCAAATATCTGTATCTCAACAGCAACCGAGTCACATCAATGGAACCTGGGTATTTTGACAA  
 TTTGGCCAAACACTCTCCTTGTGTTAAAGCTGAACAGGAACCGAATCTCAGCTATCCCAACCAGATGTTTAAACT  
 GCCCAACTGCAACATCTCGAATTGAACCGAAACAAGATTAATAATGTAGATGGACTGACATTTCCAAGCCCTGG  
 TGCTCTGAAGTCTCTGAAAATGCAAAGAAATGGAGTAACGAAACTTATGGATGGAGCTTTTTGGGGGCTGAGCAA  
 CATGGAAATTTGCAGCTGGACCATAACAACCTAACAGAGATTACCAAAGGCTGGCTTTACGGCTTGCTGATGCT  
 GCAGGAACCTCATCTCAGCCAAATGCCATCAACAGGATCAGCCCTGATGCCCTGGGAGTTCCTGCCAGAAGCTCAG  
 TGAGCTGGACCTAACTTTCAATCACTTATCAAGTTAGATGATTCAAGCTTCTTGGCCTAAGCTTACTAAATAC  
 ACTGCACATTTGGGAACAACAGAGTCAGCTACATTTGCTGATTTGCTTCCGGGGGCTTTCCAGTTTAAAGACTTT  
 GGATCTGAAGAACATGAAATTTCTGACTATGAAGACATGAATGGTGTCTTCTCTGGGCTTGACAAAACCTGAG  
 GCGACTGATACTCCAAGGAAATCGGATCCGTTCTATTACTAAAAAAGCCTTCACTGGTTTGGATGCATTGGAGCA  
 TCTAGACCTGAGTGACAACGCAATCATGCTTTTACAAGGCAATGCATTTTCAAAATGAAGAACTGCAACAAT  
 GCATTTAAATACATCAAGCCTTTTGTGCGATTGCCAGCTAAAAATGGCTCCACAGTGGGTGGCGGAAACAATTT  
 TCAGAGCTTTGTAATGCGAGTTTGCCTCCTCAGCTGCTAAAAGGAAAGAGCATTTTGTGTTAGCCCA  
 TGGCTTTGTGTGTGATGATTTTCCAAACCCAGATCAGGTTTCAAGCAAGCAACAGTCCGGCAATAAAGGTTT  
 CAATTTGAGTTTTCATCTGCTCAGCTGCCAGCAGCAGTATTCCCAATGACTTTTGTCTGGAAAAAGACAATGA  
 ACTACTGCATGATGCTGAAATGGAAAATTTATGCACACCTCCGGGCCCAAGGTTGGCAGGATGAGGATATACCAC  
 CATCCTTCCGCTGCGCAGGTTGAAATTTGCCAGTGAAGGAAATATCAGTGTGTCATCTCCAATCACTTTGGTTC  
 ATCCTACTCTGTCAAAGCCAAAGCTTACAGTAAATATGCTTCCCTCATTCAACAAGACCCCAATGGATCTCACCAT  
 CCGAGCTGGGGCCATGGCACGCTTGGAGTGTGCTGCTGTGGGGCACCCAGCCCCAGATAGCCTGGCAGAAGGA  
 TGGGGCACAGACTTCCAGCTGCACGGGAGAGACGATGCATGTGATGCCCGAGGATGACGTGTTCTTTATCGT  
 GGATGTGAAGTAGAGGACATTTGGGTATACAGCTGCAGCTCAGAACAGTGCAGGAAGTATTTACGCAAAATGC  
 AACTCTGACTGTCTAGAAAACCATCATTTTTTGGGGCCACTGTTGGACCCGAAGTGAACCAAGGGAGAAACAGC  
 CGTCCCTACAGTGCATGTCTGGAGGAAGCCCTCCCCCTAAACTGAACTGGACCAAAAGATGATAGCCCATTTGGTGGT  
 AACCGAGAGGCATTTTTTGCAGCAGGCAATCAGCTTCTGATTATTTGGACTCAGATGTAGTGTGGGAA  
 ATACACATGTGAGATGTCTAACCCCTTGGCACTGAGAGAGGAAACGTCGCTCAGTGTGATCCCACTCCAAC  
 CTGCGACTCCCTCAGATGACAGCCCATCGTTAGACGATGACGGATGGGCCACTGTGGGTGTGATCATAGC  
 CGTGGTTTGTGTGTTGGGACGCTACTCGTGTGGTGGTCAATCATATACCACACAAGGCGGAGGAATGAAGA  
 TTGCAGCATTACCAACACAGATGAGACCAACTTGCAGCAGATATTCCTAGTTATTTGTCATCTCAGGGAACGTT  
 AGCTGACAGGCAGGATGGGTACGTGCTTTCAGAAAGTGGAGCCACCACAGTTTGTCAATCTTCAGGTGCTGG  
 ATTTTTCTTACCACAACATGACAGTAGTGGGACCTGCCATATTGACAATAGCAGTGAAGCTGATGTGGAGCTGC  
 CACAGATCTGTTTCTTTGTCCGTTTTTGGGATCCACAGGCCCTATGTATTTGAAGGGAAATGTGATGGCTCAGA  
 TCCTTTTGAAACATATACACAGGTGACAGTCTGACCCAAAGAACAGTTTAAATGGACCCTATGAGCCAGTTA  
 CATAAAGAAAAGAGTGTACCCATGTTCTCATCCTTCAGAAGAATCCFGCGAACGGAGCTTCAATATATATC  
 GTGGCCTTACATGTGAGGAAGCTACTTAACTAGTTACTCTCAATGAAGGACCTGGAATGAAAAATCTGTG  
 TCTAAACAAGTCTCTTTAGATTTTGTAGATTTTGTGCAAAATCCAGAGCCAGCGTGGTTCCTCGAGTAATTTTCATGGG  
 TACCTTTGGAAAAGCTCTCAGGAGACCTCACTTAGATGCCATTCAAGCTTTGGACAGCCATCAGATTGTCAGCC  
 AAGAGCCTTTTATTTGAAAGCTCATTTCTCCCAAGACTTGGACTCTGGGTTCAGAGGAGATGGGAAAGAAAGGAC  
 AGATTTTCAGGAAGAAAATCACATTTGTACTTTTAAACAGACTTTAGAAAACCTACAGGACTCCAATTTTCAGTC  
 TTATGACTTGGACACATAGACTGAAATGAGACCAAGGAAAAGCTTAACTACTACCTCAAGTGAACCTTTTATTTA  
 AAAGAGAGAGAACTTTATGTTTTTAAATGGAGTTATGAATTTTAAAAGGATAAAAAATGCTTTATTTATACAGAT  
 GAACCAAAATTAACAAAAGTTATGAAAATTTTATACTGGGAATGATGCTCATATAAGAAATACCTTTTAAACTA  
 TTTTTTAACTTTGTTTTATGCAAAAAGTATCTTACGTAATTAATGATATAAATCATGATTTTATGATTTT  
 TTATAATGCCAGATTTCTTTTATGAAAATGAGTTACTAAGCATTTTAAATAATACCTGCCCTGTACCATTTT  
 TTAATAGAAGTTACTTCAATATATTTTGCACATTATATTTAATAAAATGTGTCAATTTGAA

## **FIGURE 102**

MVDVLLLFSLCLLFHISRPDLSHNRLSFIKASSMSHLQSLREVKLNNELETIPNLGPVSAN  
ITLLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAPPALQLKYLYLNSNRVTSMEP  
GYFDNLANTLLVLKLNRRRISAI PPKMFKLPQLQHLELNRRNKIKNVDGLTFQGLGALKSLKM  
QRNGVTKLMDGAFWGLSNMEI LQLDHNNLTETTKGWLYGLLMLQELHLSQNAINRISPDAWE  
FCQKLSELDLTFNHL SRLDDSSFLGLSLNLT LHI GNNRVSYIADCAFRGLSSSLKTLDLKNN  
ISWTIEDMNGAFSGLDKLRRLILQGNRIRSIKKAF TGLDALEHLDLSDNAIMSLQGNAFSQ  
MKKLQQLHLNLTSSLLCDCQLKWL PQWVAENNFQSFVNASC AHPQLLKGRSIFAVSPDGFVCD  
DFPKPQITVQPETQSAIKGSNLSFICSAASSSDSPMTFAWKKNELLDHAEMENYAHLRAQG  
GEVMEYTTILRLREVEFASEGKYQCVISNHFGSSYSVKAKLTVNMLPSFTKTPMDLTIRAGA  
MARLECAAVGHPAPQIAWQKGGTDFPAARERRMHVMPEDDVFFIVDKIEDIGVYSCTAQN  
SAGSISANATLTVLETPSFLRPLLDRTVTKGETAVLQCIAGGSPPPKLNWTKDDSPLVVTER  
HFFAAGNQLLIIVDSVDVSDAGKYTCEMSNTLGTERRGNVRLSVIPTPTCDSPQMTAPSLDDD  
WATVGVVVI IAVVCCVVGTSLVVVVI IYHTRRRNEDCSITWDETNLPADIPSYLSSQGTLD  
RQDGYVSSSESGSHHQFVTSSGAGFFLPQHDSSGTCHIDNSSEADVEAATDLFLCPFLGSTGP  
MYLKGNVYGSDFPETYHTGCS PDRPTVLMHDHYEPSYIKKKECYPCSHPESESCERSFSNISW  
PSHVRKLLNTSYSHNEGPGMKNLCLNKSSLDFSANPEPASVASSNSFMGTGFKALRRPHLDA  
YSSFGQPSDCQPRAFYLKAHSSPDLDSGSEEDGKERTDFQEEHNICTFKQTLNRYRTPNFQS  
YDLDT

**Signal sequence:**

amino acids 1-19

**Transmembrane domain:**

amino acids 746-765

**N-glycosylation site.**

amino acids 62-66, 96-100, 214-220, 382-386, 409-413, 455-459,  
628-632, 669-673, 845-849, 927-931, 939-943, 956-960

**Glycosaminoglycan attachment site.**

amino acids 826-830

**Casein kinase II phosphorylation site.**

amino acids 17-21, 39-43, 120-124, 203-207, 254-258, 264-268,  
314-318, 323-327, 347-351, 464-468, 548-552, 632-636, 649-653,  
671-675, 739-743, 783-787, 803-807, 847-851, 943-947, 958-962,  
1013-1017, 1019-1023, 1021-1025

**Tyrosine kinase phosphorylation site.**

amino acids 607-615

**N-myristoylation site.**

amino acids 179-185, 197-203, 320-326, 367-373, 453-459, 528-534,  
612-618, 623-629, 714-720, 873-879



**FIGURE 103**

GGGGAGAGGAATTGACCATGTA AAAAGGAGACTTTTTTTTTTGGTGGTGGTGGCTGTGGGTGCCTTGCAAAAATG  
AAGGATGCAGGACGCAGCTTTCTCCTGGAACCGAACGCAATGGATAAACTGATTGTGCAAGAGAGAAGGAAGAAC  
GAAGCTTTTTCTTGTGAGCCCTGGATCTTAACACAAATGTGTATATGTGCACACAGGGAGCATCAAGAATGAAA  
TAAACCAGAGTTAGACCCGCGGGGGTGGTGTGTTCTGACATAAAATAAATAATCTTAAAGCAGCTGTCCCTCC  
CCACCCCAAAAAAAGGATGATTGGAAATGAAGAACCAGGATTCACAAAAGAAAAAGTATGTTTCATTTTTCTC  
TATAAAGGAGAAAGTGAGCCAAGGAGATATTTTTGGAAATGAAAAGTTTGGGGCTTTTTAGTAAAGTAAAGAACT  
GGTGTGGTGTGTTTTCTTTCTTTTTGAATTTCCCAAGAGGAGAGGAAATTAATAATACATCTGCAAAAGAAA  
TTTTAGAGAAAGAAAGTTGACCGCGCAGATTGAGGCATTGATTGGGGGAGAGAAACCAGCAGAGCACAGTTGGA  
TTTTGCTTATGTTGACTAAAATTGACGGATAATGTCAGTTGGATTTTTCTTCATCAACCTCTTTTTTTAAAT  
TTTTATTCTTTTTGGTATCAAGATCATGCGTTTTCTCTTGTCTTAACCACCTGGATTCCATCTGGATGTTGCT  
GTGATCAGTCTGAAATACAACCTGTTTGAATCCAGAAGGACCAACACAGATAAAATATGAATGTTGAACAAGAT  
GACCTTACATCCACAGCAGATAATGATAGGTCTTAGGTTAACAGGGCCCTATTTGACCCCTGCTTGTGGTGTCT  
GCTGGCTCTTCAACTCTTGTGGTGGCTGGTCTGGTGCAGGCTCAGACCTGCCCTTCTGTGTGCTCCTGCAGCAA  
CCAGTTCAGCAAGGTGATTTGTGTTCCGAAAAACCTGCGTGAGGTTCCGGATGGCATCTCCACCAACACAGGCT  
GCTGAACCTCCATGAGAACCAATCCAGATCATCAAAGTGAACAGCTTCAAGCATTGAGGCATTGGAAATCCT  
ACAGTTGAGTAGGAACCATATCAGAACCATTGAAATGGGGCTTCAATGGTCTGGCGAACCTCAACACTCTGGA  
ACTCTTTGACAACTCTTACTTACCATCCCGAATGGAGCTTTTTGTATACTTGTCTAAACTGAAGGAGCTCTGGTT  
GCGAAACAACCCATTGAAAGCATCCCTTCTTATGCTTTTAAACAGAATTCCTTCTTTGCGCCGACTAGACTTAGG  
GGAATGAAAAGACTTTCATACATCTCAGAAGGTGCCCTTTGAAGGTCTGTCCAACCTGAGGTATTTGAACTTGC  
CATGTGCAACCTTCGGGAAATCCCTAACCTCACACCCGCTATAAACTAGATGAGCTGGATCTTCTGGGAATCA  
TTTTATCTGCCATCAGGCCTGGCTCTTTCAGGGTTTGATGCACCTTCAAAAACTGTGGATGATACAGTCCAGAT  
TCAAGTGATTGAACGGAATGCCCTTTGACAACCTTCAGTCACTAGTGGAGATCAACCTGGCACAATAATCTAAC  
ATTACTGCCCTCATGACCTCTTCACTCCCTGTCATCATCTAGAGCGGATACATTTACATCACACCTTGGAACTG  
TAACCTGTGACATACTGTGGCTCAGCTGGTGGATAAAAGACATGGCCCTCGAACACAGCTTGTGTGCCCGGTG  
TAACACTCTCCCAATCTAAAGGGGAGGTACATTGGAGAGCTCGACCAGAATTACTTCACATGCTATGCTCCGGT  
GATTGTGGAGCCCTGCGAGACCTCAATGTCACTGAAGGCATGGCAGCTGAGCTGAAATGTGGGCTCCACATC  
CCTGACATCTGTATCTGGATTACTCCAAATGGAACAGTCATGACACATGGGGCGTACAAAGTGGGATAGCTGT  
GCTCAGTGATGGTACGTTAAATTTCACAATGTAACCTGTGCAAGATACAGGCATGTACACATGTATGGTGGATAA  
TTCCGTTGGGAATACTACTGCTTCAGCCACCCTGAATGTTACTGCAGCAACCCTACTCTTTCTTACTTTTT  
AACCTCACAGTAGAGACTATGGAACCGTCTCAGGATGAGGCACGGACCAAGATAAACAATGTTGGTCCACTCC  
AGTGGTGCAGTGGGAGACCACCAATGTGACCCTCTCTCACACCACAGAGCACAAGGTCGACAGAGAAAACTT  
CACCATCCAGTGA CTGATATAAACAGTGGGATCCAGGAATTTGATGAGGTGATGAAGACTACAAAAATCATCAT  
TGGGTGTTTTGTGGCCATCACACTCATGGCTGCAGTGATGCTGGTCATTTTCTACAAGATGAGGAAGCAGACCA  
TCGGCAAAACCATCACGCCCAACAAGGACTGTTGAAATTTAATGTGGATGATGAGATTACGGGAGACACACC  
CATGGAAGCCACTGCCATGCCTGCTATCGAGCATGAGCACCTAAATCACTATAACTCATACAAATCTCCCTT  
CAACCACACAACAACAGTTAACACAATAAATCAATACACAGTTTCACTGATGAACCGTTATTGATCCGAATGAA  
CTCTAAAGACAATGTACAAGAGACTCAAATCTAAACATTTACAGAGTTACAAAAACAACAATCAAAAAAAA  
GACAGTTTATAAAAATGACACAAATGACTGGGCTAAATCTACTGTTTCAAAAAAGTGTCTTTACAAAAAACAA  
AAAAGAAAAGAAATTTATTTATTAATAAATCTATTTGTGATCTAAAGCAGACAAAA

## **FIGURE 104**

MLNKMTLHPQQIMIGPRFNRALFDPLLVLALLQQLLVVAGLVRAQTCPSVCSCSNQFSKVIC  
VRKNLREVPDGI STNTRLLNLHENQIQI I KVNSFKHLRHLEILQLSRNHIRTIEIGAFNGLA  
NLNTLELFDNRLTTIPNGAFVYLSKLELWLRNNPIESIPSYAFNRIPSLRRLDLGELKRLS  
YISEGAFEGLSNLRYLNLAMCNLREIPNLTPLIKLDELDSLGNHLSAIRPGSFQGLMHLQKL  
WMIQSQIQVIERNAFDNLQSLVEINLAHNNLTLPHDLFTPLHHLERIHLLHNPWNCNDIL  
WLSWWIKDMAPSNTACCARCNTPPNLKGRYIGELDQNYFTCYAPVIVEPPADLNVTEGMAAE  
LKCRASSTLSVSWITPNGTVMTHGAYKVRIAVLSDGTLNFTNVTVQDTGMYTCMVSNVSGN  
TTASATLNVTAATTTFFSYFSTVTVETMEPSQDEARTDNNVGPFPVVDWETTNTVTTSLTPQ  
STRSTEKFTTIPVTDINSIGIPGIDEVMKTKIIIGCFVAITLMAAVMLVIFYKMRKQHRQN  
HHAPTRTVEIINVDDEITGDTMPESHLPMPAIEHEHLNHYSYKSPFNHTTTVNTINSIHSS  
VHEPLLIRMNSKDNVQETQI

**Signal sequence:**

amino acids 1-44

**Transmembrane domain:**

amino acids 523-543

**N-glycosylation site.**

amino acids 278-282, 364-368, 390-394, 412-416, 415-419, 434-438,  
442-446, 488-492, 606-610

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 183-187

**Casein kinase II phosphorylation site.**

amino acids 268-272, 417-421, 465-469, 579-583, 620-624

**N-myristoylation site.**

amino acids 40-46, 73-79, 118-124, 191-197, 228-234, 237-243,  
391-397, 422-428, 433-439, 531-537

**FIGURE 105**

AGCCGACGCTGCTCAAGCTGCAACTCTGTTGCAGTTGGCAGTTCTTTTCGGTTTCCCTCCTGCTGTTTGGGGGCA  
TGAAAGGGCTTCGCCGCGGGAGTAAAGAAGGAATTGACCGGGCAGCGCGAGGGAGGAGCGCGCACGCCGCCG  
GAGGGCGGGCGTGCACCCCTCGGCTGGAAGTTTGTGCCGGGCCCCGAGCGCGCGCCGGCTGGGAGCTTCGGGTAGA  
GACCTAGGCCGCTGGACCGCGATGAGCGCGCCGAGCCTCCGTGCGCGCGCCGGGGTTGGGGCTGCTGCTGTGC  
GCGGTGCTGGGGCGCCTGGCCGGTCCGACAGCGCGGTCGCGGGGAACTCGGGCAGCCCTCTGGGGTAGCCGCC  
GAGCGCCCATGCCCCACTACCTGCCGCTGCCCTCGGGGACCTGCTGGACTGCAGTCTGAAGCGGCTAGCGCGTCTT  
CCCCGAGCCATCCCCTGCTGGGTGCTGCGCTGGACTTAAGTCACAACAGATTATCTTTCATCAAGGCAAGTTCC  
ATGAGCCACCTTCAAAGCCTTCGAGAAGTGAACCTGAACAACAATGAATGGAGACCATTCCAATCTGGGACCA  
GTCTCGCAATATTACACTTCTCTCTGGCTGGAACAAGGATTGTTGAAATACTCCCTGAACATCTGAAAGAG  
TTTTAGTCCCTTGAACCTTGGACCTTAGCAGCAACAATATTTAGAGCTCCAACCTGCATTTCCAGCCCTACAG  
CTCAATATCTGTATCTCAACAGCAACCGAGTCAATCAATGGAACCTGGGTATTTTGACAATTTGGCCAACACA  
CTCCTTGTGTTAAAGCTGAACAGGAACCGAATCTCAGCTATCCCACCAAGATGTTAAACTGCCCAACTGCAA  
CATCTCGAATTTGAACCGAAACAAGATTAAAAATGTAGATTGACTGACATTCCAAGCCTTGGTGTCTGAACTCT  
CTGAAATGCAAAGAAATGGAGTAACGAACTTATGGATGGAGCTTTTGGGGGCTGAGCAACATGGAAATTTG  
CAGCTGGACCATAACACCTAACAGAGATTACCAAGGCTGGCTTTACGGCTTGTGTGCTGCAGGAACCTCAT  
CTCAGCAAAATGCCATCAACAGGATCAGCCCTGATGCTGGGAGTCTGCCAGAAGTCTGATGCTGCAAGTCA  
ACTTTCAACTTATCAAGGTTAGATTCAAGCTTCTTGGCCTAAGCTTACTAAATACACTGCACATTGGG  
AACACAGAGTCACTACATTGCTGATTTGCTTCCGGGGCTTTCAGTTTAAAGACTTTGGATCTGAAGAAC  
AATGAAATTTCTGGACTATTGAAGACATGAATGGTGTCTTCTGGGCTTGACAACTGAGGCGACTGATACTC  
CAAGGAAATCGGATCCGTTCTATTAACAAGGCAATGCAATTTTCAAAATGAAGAACTGCAACAATTTGCAATTA  
GACAACGCAATCATGCTTTTACAGGCAATGCAATTTTCAAAATGAAGAACTGCAACAATTTGCAATTAATA  
TCAAGCCTTTTGTGCGATTGCCAGCTAAAATGGCTCCCACAGTGGGTGGCGGAAACAACTTTCCAGAGCTTTGTA  
AATGCCAGTTGTGCCATCTCAGCTGCTAAAAGGAAGAAGCATTGCTGTTAGCCAGATGGCTTTGTGTGT  
GATGATTTCCCAAACCCAGATCACGGTTGAGCCAGAAACACAGTCCGCAATAAAGGTTCCAATTTGAGTTTC  
ATCTGCTCAGCTGCCAGCAGCAGTGTATCCCAATGACTTTTGGCTGGAAAAAGACAATGAACTACTGCATGAT  
GCTGAAATGGAATATTATGCACACCTCCGGGCCAAGGTGGCGAGGTGATGGAGTATACCACCATCTTCCGCTG  
CGCGAGGTGGAATTTGCCAGTGAAGGGAAATATCAGTGTGTCTCAATCACTTTGGTTTATCCTACTCTGTC  
ATGAAACACCATCATTTTGGCGCCACTGTTGGACCGAACTGTAACCAAGGAGAAACAGCCCTCTACAGTGC  
ATTGCTGGAGGAAGCCCTCCCCCTAACTGAACTGGACCAAGATGATAGCCCATTTGGTGGTAACCGAGAGGCAC  
TTTTTTGCAGCAGGCAATCAGCTTCTGATATTGTTGGACTCAGATGTGATGCTGGGAAATACACATGTGAG  
ATGCTAACACCCCTTGGCACTGAGAGAGGAAACGTCGCCTCAGTGTGATCCCCACTCCAACCTGGCACTCCCT  
CAGATGACAGCCCCATCGTTAGACGATGACGGATGGGCCACTGTGGGTGTGCTGATCATAGCCGTTGCTGT  
GTGGTGGGCACGTCACCTGTTGGGTGGTTCATCATATACCACACAAGGCGGAGGAATGAAGATTGCAGCATTACC  
AACACAGATGAGACCAACTTGCCAGCAGATATCTTAGTTATTTGTCTCATCTCAGGGAACGTTAGCTGACAGGCAG  
GATGGGTACGTTCTTCAAGAAAGTGAAGCCACCACAGTTGTGATCTTCAAGTGTGGATTTTCTTACCA  
CAACATGACAGTAGTGGGACCTGCCATAATTGACAATAGCAGTGAAGCTGATGTGGAAGCTGCCACAGATCTGTT  
CTTTGTCCGTTTTTGGGATCCACAGGCCCTATGTATTTGAAGGAAATGTGTATGGCTCAGATCCTTTTGAACA  
TATCATACAGTTGAGTCTTGCACCAAGAACAGTTTAAATGGACCACTATGAGCCAGTTACATAAAGAAAAG  
GAGTGTACCCATGTTCTCATCTTFCAGAAGAATCTGCGAACCGGAGCTTCAGTAATATATCGTGGCCCTTACAT  
GTGAGGAAGCTACTTAACTAGTTACTCTCACAAATGAAGGACCTGGAATGAAAATCTGTGTCTAAACAAGTCC  
TCTTTAGATTTTGTGCAAAATCCAGAGCCAGCGTGGTTGCCCTCGAGTAATCTTTCATGGGTACCTTTGAAAA  
GCTCTCAGGAGACCTCACTAGATGCCTATTTCAAGCTTTGGACAGCCATCAGATTGTGAGCAAGAGCCTTTTAT  
TTGAAAGCTCATTTCCCGAGACTTTGGACTCTGGGTGAGGAAAGATGGGAAAGAAAGGACAGATTTTCAGGAA  
GAAAATCACATTTGTACCTTTAAACAGACTTTAGAAAACACTACAGGACTCCAATTTTTCAGTCTTATGACTTGGAC  
ACATAGACTGAATGAGACCAAGGAAAAGCTTAACTACTACTCAAGTGAACCTTTTATTTAAAAGAGAGAGAAT  
CTTATGTTTTTAAATGGAGTTTATGAATTTTAAAAGGATAAAAATGCTTTATTTATACAGATGAACCAAAATAC  
AAAAAGTTATGAAAATTTTATACTGGGAATGATGCTCATATAAGAATACCTTTTAAACTATTTTTTAACTTTG  
TTTTATGCAAAAAGTATCTTACGTAATTAATGATATAAATCATGATTTATTTATGATTTTTTATAATGCCAGA  
TTTTCTTTTATGAAAAATGAGTTACTAAAGCATTTTAAATAATACCTGCCTTGTACCATTTTTTAAATAGAAGTT  
ACTTCATTATTTTTGCACATTATTTAATAAATGTGTCAATTTGAAAAAATAAAAAAAAAAAAAAAAAAAAAA

## **FIGURE 106**

MSAPSLRARAAGLGLLLCAVLGRAGRSDSGGRGELGQPSGVAAERPCPTTCRCLGDLLDCSR  
KRLARLPEPLPSWVARLDLSHNRLSFIKASSMSHLQSLREVKLNNELETIPNLGPVSANIT  
LLSLAGNRIVEILPEHLKEFQSLETLDLSSNNISELQTAFFPALQLKYLYLNSNRVTSMEPGY  
FDNLANTLLVLKLNRRNRI SAIPPKMFKLPQLQHLELNRNKIKNVDGLTFQGLGALKSLKMQR  
NGVTKLMGDAFWGLSNMEILQLDHNNTLTIKGWLYGLLMLQELHLSQNAINRISPDAWEFC  
QKLSELDLTFNHL SRLDDSSFLGLSLLNTLHIGNNRVSYIADCAFRGLSSLKTLDLKNNEIS  
WTI EDMNGAFSGLDKLRRLILQGNRIRSITKKAFTGLDALEHLDLSDNAIMSLQGNAFSQMK  
KLQQLHLNNTSSLLCDCQLKWLPOQVAENNFQSFVNASCAPQLLKGRSIFAVSPDGFVCDDF  
PKPQITVQPETQSAIKGSNLSFICSAASSSDSPMTFAWKKNELLHDAEMENYAHLRAQGGE  
VMEYTTILRLREVEFASEGKYQCVISNHFGSSYSVKAKLTVNMLPSFTKTPMDLTI RAGAMA  
RLECAAVGHPAPQIAWQKDGTDPAARERRMHVMPEDDVFFIVDVKIEDIGVYSCTAQN SA  
GSISANATLTVLETSPFLRPLLDRTVTKGETAVLQCIAGGSPPPKLNWTKDSDPLVVTERHF  
FAAGNQLLIIVSDSDVSDAGKYTCEMSNTLGTERGNVRLSVIPTPTCDSPQMTAPSLDDDGWA  
TVGVVIIAVVCCVVGTSLVVWVVIYHTRRRNEDCSITNTDETNPADIPSYLSSQGT LADRQ  
DGYVSSSESGSHHQFVTSSGAGFFLPQHDSSTGCHIDNSSEADVEAATDLFLCPFLGSTGPMY  
LKGNVYGSDFPETYHTGCSPDPRTVLMHDHYEPSYIKKKECYPCHPSEESCERSFSNISWPS  
HVRKLLNTSYSHNEGPGMKNLCLNKSSLDFSANPEPASVASSNSFMGTFGKALRRPHLDAYS  
SFGQPSDCQPRAFYLKAHSSPDLDSGSEEDGKERTDFQEENHICTFKQTLNRYTPNFQSYDLDT

**Signal sequence:**

amino acids 1-27

**Transmembrane domain:**

amino acids 808-828

**N-glycosylation site.**

amino acids 122-126, 156-160, 274-278, 442-446, 469-473, 515-519,  
688-692, 729-733, 905-909, 987-991, 999-1003, 1016-1020

**Glycosaminoglycan attachment site.**

amino acids 886-890

**Casein kinase II phosphorylation site.**

amino acids 99-103, 180-184, 263-267, 314-318, 324-328, 374-378,  
383-387, 407-411, 524-528, 608-612, 692-696, 709-713, 731-735,  
799-803, 843-847, 863-867, 907-911, 1003-1007, 1018-1022,  
1073-1077, 1079-1083, 1081-1085

**Tyrosine kinase phosphorylation site.**

amino acids 667-675

**N-myristoylation site.**

amino acids 14-20, 36-42, 239-245, 257-263, 380-386, 427-433,  
513-519, 588-594, 672-678, 683-687, 774-780, 933-939

**Leucine zipper pattern.**

amino acids 58-80, 65-87

**FIGURE 107**

CAAACTTGCGTCGCGGAGAGCGCCAGCTTGACTTGAAATGGAAGGAGCCCGAGCCCGCGGAGCGCAGCTGAGAC  
TGGGGGAGCGCGTTCGGCCTGTGGGGCGCCGCTCGGCCTGGGGCGCAGCAGGGAAGGGGAAGCTGTGGTCTGCC  
CTGCTCCACGAGGCGCCACTGGTGTGAACCGGGAGAGCCCTGGGTGGTCCCGTCCCCTATCCCTCCTTTATATA  
GAAACCTTCCCACTGGGAAGGCAGCGCGAGGCAGGAGGGCTCATGGTGAGCAAGGAGGCGGCTGATCTGCAG  
GCGCACAGCATTCCGAGTTTACAGATTTTACAGATACCAAATGGAAGCGCAGGAGGCAGAACAGCCTGCCTGGT  
TCCATCAGCCCTGGCGCCAGGCGCATCTGACTCGGCACCCCTGCAGGCACCATGGCCAGAGCCGGTGTGC  
TGCTCCTGCTGTGCTGCCGCGACAGCTGCACCTGGGACCTGTGCTTGCCTGAGGGCCCCAGGATTTGGCCGAA  
GTGGCGCCACAGCCTGAGCCCGAAGAGAACGAATTTGGGGAGGAGGAGCCGGTGTGGTACTGAGCCCTGAGG  
AGCCCGGGCCTGGCCAGCCGGTCACTGCTGCCCGGAGACTGTGCCTGTTCCAGGAGGGCGTCTGTGGACTGTG  
GCGGTATTGACCTGCGTGAAGTTCCCGGGGACCTGCCTGAGCACACCAACCACCTATCTCTGCAGAACAAACAGC  
TGGAAAAGATCTACCTGAGGAGCTCTCCCGGCTGCACCGGTGGAGACACTGAACCTGCAAAACAACCGCCTGA  
CTTCCCGAGGGCTCCAGAGAAGGCGTTGAGCATCTGACCAACCTCAATTACCTGTACTTGGCCAATAACAAGC  
TGACCTTGGCACCCCGCTTCTGCCAAACCGCCTGATCAGTGTGGACTTTGCTGCCAACTATCTCACCAAGATCT  
ATGGGCTCACCTTTGGCCAGAAGCCAACTTGAAGTCTGTGTACTCTGCACAACAACAAGCTGGCAGACGCGGGC  
TGCCGGACAACATGTTCAACGGCTCCAGCAACGTCGAGGTCCTCATCTGTCCAGCAACTTCTCGCCACGCTGC  
CCAAGCAGCTGCCGCTGCCCTGTACAAGCTGCACCTCAAGAACAACAAGCTGGAGAAGATCCCCCGGGGCGCT  
TCAGCGAGCTGAGCAGCCTGCGCGAGCTATACCTGCAGAACAACCTACTGACTGACGAGGGCCTGGACAACGAGA  
CCTTCTGGAAGCTCTCCAGCTTGGAGTACCTGGATCTGTCCAGCAACAACCTGTCTCGGGTCCAGCTGGGCTGC  
CGCGCAGCCTGGTGTGCTGCCTTGGAGAAGAACCCTCCGGAGCGTGGACCGAATGTGCTGACCCCATCC  
GCAGCTGGAGTACCTGTCTGCACAGCAACCAGCTGCGGGAGCAGGGCATCCACCCACTCGCCCTCCAGGGCC  
TCAAGCGGTTGCACACGGTGCACCTGTACAACAACCGCTGGAGCGCTGCCAGTGGCCTGCCTCGCCGCTGC  
GCACCTCATGATCTGCACAACCAGATCACAGGATTGGCCGGAAGACTTTGCCACCACCTACTTCTGGAGG  
AGCTCAACCTCAGCTACAACCGCATCACAGCCACAGGTGCACCGGACGCCTTCCGAAGCTGCGCCTGTGC  
GCTCGCTGGACCTGTGCGGCAACCGGCTGCACACGCTGCCACTTGGGCTGCCTCGAAATGTCCATGTGCTGAAGG  
TCAAGCGAATGAGCTGCCTGCTTGGCACGAGGGCGCTGGCGGCATGGCTCAGCTGCGTGTGACTGTACTCA  
CCAGCAACCGACTGCGCAGCCGAGCCCTGGGCCCCGCTGCCCTGGTGGACCTCGCCCATCTGCAGCTGCTGGACA  
TCGCCGGGAATCAGCTCACAGAGATCCCGAGGGGCTCCCGAGTCACTTGAGTACCTGTACTCTGCAGAACAA  
AGATTAGTGGGTGCCCGCCAAATGCCTTGCATCCACGCCAACCTCAAGGGGATCTTCTCAGGTTTAAACAAGC  
TGGCTGTGGGCTCCGTGGTGGACAGTGCCTTCCGGAGGCTGAAGCACCTGCAGGTCTTGGACATTGAAGGCACT  
TAGAGTTTGGTGACATTTCCAAGGACCGTGGCCGCTTGGGGAAGGAAAAGGAGGAGGAAGAGGAGGAGGAGG  
AGGAAGAGGAAACAAGATAGTGAAGGTGATGCAGATGTGACCTAGGATGATGGACCGCCGGACTCTTTCTGC  
AGCACACGCTGTGTGCTGTGAGCCCCCACTCTGCCGTGCTCACACAGACACCCAGCTGCACACATGAGGCA  
TCCCACATGACACGGGCTGACACAGTCTCATATCCCCACCCCTTCCCACGGCGTGTCCACGGCCAGACACATGC  
ACACACATCACACCTCAAACACCCAGCTCAGCCACACACAACCTACCCTCAAACCACCACAGTCTCTGTACAC  
CCCCACTACCGCTGCCACGCCCTCTGAATCATGCAGGGAAGGGTCTGCCCTGCCCTGGCACACACAGGCACCCA  
TTCCCTCCCCCTGTGACATGTGTATGCGTATGCATACACCACACACACACATGCACAAGTATGTGCGAA  
CAGCCCTCAAAGCCTATGCCACAGACAGCTCTTGCCCCAGCCAGAATCAGCCATAGCAGCTCGCCGTCTGCCCT  
GTCCATCTGTCCGTCCGTTCCCTGGAGAAGACACAAGGGTATCCATGCTCTGTGGCCAGGTGCCCTGCCACCCTCT  
GGAACCTCAAAAAGCTGGCTTTTATTCCTTCCCATCTATGGGGACAGGAGCCTTCAAGACTGCTGGCCTGGCC  
TGGCCACCCTGCTCCTCAGGTGCTGGGCGTCACTCTGCTAAGAGTCCCTCCCTGCCACGCCCTGGCAGGACA  
CAGGCACTTTCAAATGGGCAAGCCAGTGGAGGCAGGATGGGAGAGCCCTGGGTGCTGCTGGGGCCTTGGGG  
CAGGAGTGAAGCAGAGGTGATGGGCTGGCTGAGCCAGGAGGAAGGACCCAGCTGCACCTAGGAGACACCTTT  
GTTCTTCAAGCCTGTGGGGGAAGTTCGGGTGCCTTTATTTTATTCTTTCTAAGGAAAAAATGATAAAAT  
CTCAAAGCTGATTTTCTTGTATAGAAAACTAATATAAAAGCATTATCCCTATCCCTGCAAAAAA

**FIGURE 108**

MEGEEAEQPAWFHQWPRPGASDSAPPAGTMAQSRVLLLLLLLLPPQLHLGPVLAVRAPGFGRS  
GGHLSLSP ENEFAEEEPVLVLSPEEPGPGPAAVSCPRDCACSQEGVVDCCGIDLREFPGDLP  
EHTNHLSQLNNQLEKIYPEELSRHRLETLNLQNNRLTSRGLPEKAFEHLTNLNYLYLANNK  
LTLAPRFLPNALISVDFAAANYLTKIYGLTFGQKPNLRSVYLHNNKLADAGLPDNMFNGSSNV  
EVLILSSNFLRHVPKHLPPALYKHLKNNKLEKIIPGAFSELSSIRELYLQNNYLTDEGLDN  
ETFWKLSLEYLDLSSNNLSRVPAGLPRSLVLLHLEKNAIRSVDANVLTPIRSLEYLLLHSN  
QLREQGIHPLAFQGLKRLHTVHLYNNALERVPSGLPRRVRTLMILHNQITGIGREDFATTYF  
LEELNLSYNRITSPQVHRDAFRKLRLLRSLDLSGNRLHTLPPGLPRNVHVLKVRNELAALA  
RGALAGMAQLRELYLTSNRLRSFALGPRAWVDLAHLQLLDIAGNQLTEIPEGLPESLEYLYL  
QNNKISAVPANAFDSTPNLKGIFLRFNKLAVGSVVDSAFRRLKHLQVLDIEGNLEFGDISKD  
RGR LGKEKEEEEEEEEEEEEEETR

**Signal sequence:**

amino acids 1-48

**N-glycosylation site.**

amino acids 243-247, 310-314, 328-332, 439-443

**Casein kinase II phosphorylation site.**

amino acids 68-72, 84-88, 246-250, 292-296, 317-321, 591-595

**N-myristoylation site.**

amino acids 19-25, 107-113, 213-219, 217-223, 236-242, 335-341,  
477-483, 498-502, 539-545, 548-554

**Leucine zipper pattern.**

amino acids 116-138, 251-273, 258-280, 322-344, 464-486, 471-493,  
535-557

**FIGURE 109**

GGGAGGGGGCTCCGGGCGCCGCGCAGCAGACCTGCTCCGGCCGCGCGCTCGCCGCTGTCTCCGGGAGCGGGCAG  
CAGTAGCCCGGGCGCGGAGGGCTGGGGTTCTCGAGACTCTCAGAGGGGCGCCTCCCATCGGCGCCACCACCC  
CAACCTGTTCTCGCGCGCCACTGCGCTGCGCCCCAGGACCCGCTGCCAACATGGATTTTCTCTGGCGCTGGT  
GCTGTGATCCTCGCTTACCTGCAGGCGGCGCCGAGTTTCAGCGGGAGTGGCCAGGCAAAATAGTGTATCGAT  
TGGCCTATGTCGTTATGGTGGGAGGATTGACTGCTGCTGGGGCTGGGCTCGCCAGTCTTGGGGACAGTGTACGCC  
TGTGTGCCAACCCAGATGCAAAACATGGTGAATGTATCGGGCCAAACAAGTGCAGTGTATCCTGGTTATGCTGG  
AAAAACCTGTAATCAAGATCTAAATGAGTGTGGCCTGAAGCCCCGGCCCTGTAAGCACAGGTGCATGAACACTTA  
CGGCAGCTACAGTGTACTGTCTCAACGGATATATGCTCATGCGGATGGTTCTGCTCAAGTGCCCTGACCTG  
CTCCATGGCAAACTGTCAAGTATGGCTGTGATGTTGTTAAAGGACAAATACGGTGCCAGTGCCCATCCCCGGCCT  
GCACCTGGCTCCTGATGGGAGGACCTGTGTAGATGTTGATGAATGTGCTACAGGAAGAGCCTCCTGCCCTAGATT  
TAGGCAATGTGTCAACACTTTTGGGAGCTACATCTGCAAGTGTCTATAAAGGCTTCGATCTCATGTATATTGGAGG  
CAAATATCAATGTATGACATAGACGAATGCTCACTGGTCAATCATGTCAGCAGCTTTGCTCGATGTTATAA  
CGTACGTGGGCTCTACAAGTGCAAATGTAAGAAGGATACCAGGGTGTGGACTGACTTGTGTGTATATCCCAA  
AGTTATGATTGAACCTTCAGGTCCAATTCATGTACCAAAGGAAATGGTACCAATTTAAAGGGTGACACAGGAAA  
TAATAATTTGGATTCTGATGTTGGAAGTACTTGGTGGCTCCGAAGACACCATAATTCCCTCATATTACCAA  
CAGGCTTACTTCTAAGCCAACAACAAGACCTACACCAAAGCCAAACCAATTCCTACTCCACCACCACCACC  
CTGCCAACAGAGCTCAGAACACTCTACCACCTACAAACCCAGAAAGGCCAACCAATTCCTACTCCACCACCACC  
ACCAGCTGCCAGTACACTCCAGGAGGGATTACAGTTGACAAACAGGGTACAGACAGACCCTCAGAAACCCAGAGG  
AGATGTGTTCACTGTTCTGGTACACAGTTGTAATTTTGACCATGGACTTTGTGGATGGATCAGGGAGAAAGCAA  
TGACTTGCATGGGAACCAATCAGGGACCCAGCAGGTGGACAAATCTGACAGTGTCCGCAGCCAAAGCCCCAGG  
GGGAAAAGCTGCACGCTTGGTGTACTCTCGGCCCGCTCATGCATTCAGGGGACCTGTGCCTGTCAATCAGGCA  
CAAGGTGACGGGGCTGCACTCTGGCACACTCCAGGTGTTTGTGAGAAAACACGGTGCACCGGAGCAGCCCTGTG  
GGGAAGAAATGTTGGCCATGGCTGGAGGCAAACAAGATCACCTTGCAGGGGGCTGACATCAAGAGCGAATCACA  
AAGATGATTAAGGGTTGGAAAAAAGATCTATGATGGAATAAAGGAAGTGGGATTTTGGAGCTGGAGAAG  
AGAAGACTGAGGGGCAACCAATGATGTTTTCAAGTATATGAAGGGTTGGCACAGAGAGGGTGGCGACCAAGCTG  
TTCTCCATATGCACTAAGAATAGAACAGAGGAAACTGGCTTAGACTAGAGTATAAGGGAGCATTTCTTGGCAGG  
GGCCATGTTAGAACTCTCAAAAAAAGAGTGTGAAATCTCAGTATCTCTCTCTCTTCTAAAAAATAGA  
TAAAAATTTGTCTATTTAAGATGGTTAAAGATGTTCTTACCCAGGAAAAGTAAACAAATTAAGAAATTTCCAAA  
AGATGTTTTGATCCTACTAGTAGTATGCAGTGAATACTTTAGAACTAAATAATTTGGACAAGGCTTAATTTAGG  
CATTTCCCTCTTGACCTCCTAATGGAGAGGGATGAAAGGGGAGAGCCACCAATGCTGAGCTCACTGAAATA  
TCTCTCCCTTATGGCAATCCTAGCAGTATTAAGAAAAAAGGAAACTATTTATTTCAAATGAGAGTATGATGGAC  
AGATAATTTAGTATCTCAGTAATGTCTAGTGTGGCGGTGGTTTTCAATGTTTTCTCATGGTAAAGGTATAAGCC  
TTTCATTTGTTCAATGGATGATGTTTCAGATTTTTTTTTTTTTTAAAGAGATCCTTCAAGGAACACAGTTCAGAGAG  
ATTTTTCATCGGGTGCATCTCTCTGCTTCGTGTGTGACAAGTTATCTTGCTGCTGAGAAAAGAGTGCCCTGCCCC  
ACACCGGCAGACCTTTCCTTCACCTCATCAGTATGATTGATTTCTTTATCAATTTGGACTCTCCAGGTTCAC  
AGAACAGTAATATTTTTGAACAATAGGTACAATAGAAGGTCTTCTGTCAATTTAACTGGTAAAGGCAGGGCTGG  
AGGGGAAAATAAATCATTAAGCCTTTGAGTAAACGGCAGAAATATATGGCTGTAGATCCATTTTTAATGGTTTATT  
TCCTTTATGGTCAATAACTGCACAGCTGAAGATGAAAGGGGAAAAATAAATGAAAAATTTACTTTTTGATGCCAA  
TGATACATTTGCATAAACTGATGGAAGAAGTTATCCAAAGTACTGTATAACATCTTGTATTTAATTTAATGTTTT  
CTAAAAATAAAAAATGTTAGTGGTTTTCCAATGGCCTAATAAAAAACAATTTATTTGTAATAAAAAACTGTTAGTAAT

## **FIGURE 110**

MDFLLALVLVSSLYLQAAA EFDGRWPRQIVSSIGLCRYGGRIDCCWGWARQSWGQCQPVCQP  
RCKHGECIGPNKCKCHPGYAGKTCNQDLNECGLKPRPCKHRCMNTYGSYKCYCLNGYMLMPD  
GSCSSALTCSMANCQYGCDVVKGQIRCQCPSPLHLAPDGRTCVDDVDECATGRASCPRFRQC  
VNTFGSYICKCHKGFDLMIYGGKYQCHDIDEC SLGQYQCSSFARCYNVRGSYKCKCKEGYQG  
DGLTCVYIPKVMIEPSGPIHVPKNGTILKGD TGNNNWI PDVGSTWWPPKTPYIPPIITNRP  
TSKPTTRPTPKPTPIPTPPPPPLPTELRTPLPPTT PERPTTGLTTIAPAASTPPGGITVDN  
RVQTD PQKPRGDVFSVLVHSCNFDHGLCGWIREKDN DLHWEPIRDPAGGQYLTVSAAKAPGG  
KAARLVLP LGRMLHSGDLCLSFRHKVTGLHSGTLQVFVRKHGAHGAALWGRNGGHGWROTQI  
TLRGADIKSESQR

**Signal sequence:**

amino acids 1-17

**N-glycosylation site.**

amino acids 273-277

**Casein kinase II phosphorylation site.**

amino acids 166-170, 345-349

**Tyrosine kinase phosphorylation site.**

amino acids 199-206

**N-myristoylation site.**

amino acids 109-115, 125-131, 147-153, 191-197, 221-227, 236-242,  
421-427, 433-439, 462-468, 476-482

**Aspartic acid and asparagine hydroxylation site.**

amino acids 104-116, 186-198, 231-243

**Cell attachment sequence.**

amino acids 382-385

**EGF-like domain cysteine pattern signature.**

amino acids 75-87



**FIGURE 111**

CTTCTTTGAAAAGGATTATCACCTGATCAGGTTCTCTCTGCATTTGCCCTTTAGATTGTGA  
AATGTGGCTCAAGGTCTTCACAACTTTCCTTTCTTTGCAACAGGTGCTTGCTCGGGGCTGA  
AGGTGACAGTGCCATCACACACTGTCCATGGCGTCAGAGGTCAGGCCCTCTACCTACCCGTC  
CACTATGGCTTCCACACTCCAGCATCAGACATCCAGATCATATGGCTATTTGAGAGACCCCA  
CACAATGCCCAAATACTTACTGGGCTCTGTGAATAAGTCTGTGGTTCTGACTTGGAATACC  
AACACAAGTTCACCATGATGCCACCCAATGCATCTCTGCTTATCAACCCACTGCAGTTCCCT  
GATGAAGGCAATTACATCGTGAAGGTCAACATTCAGGGAATGGAACCTATCTGCCAGTCA  
GAAGATACAAGTCACGGTTGATGATCCTGTACAAAGCCAGTGGTGCAGATTCATCCTCCCT  
CTGGGGCTGTGGAGTATGTGGGGAACATGACCCTGACATGCCATGTGGAAGGGGGCACTCGG  
CTAGCTTACCAATGGCTAAAAATGGGAGACCTGTCCACACCAGCTCCACCTACTCCTTTTC  
TCCCCAAAACAATACCCCTTCATATTGCTCCAGTAACCAAGGAAGACATTTGGGAATACAGCT  
GCCTGGTGAGGAACCCTGTCAAGTGAATGGAAAGTGATATCATTATGCCCATCATATATTAT  
GGACCTTATGGACTTCAAGTGAATTTCTGATAAAGGGCTAAAAGTAGGGGAAGTGTTTACTGT  
TGACCTTGAGAGGGCCATCCTATTTGATTGTTCTGCTGATTCTCATCCCCCAACACCTACT  
CCTGGATTAGGAGGACTGACAATACTACATATATCATTAAGCATGGGCCCTCGCTTAGAAGTT  
GCATCTGAGAAAGTAGCCAGAAAGACAATGGACTATGTGTGCTGTGCTTACAACAACATAAC  
CGGCAGGCAAGATGAACTCATTTTCACAGTTATCATCACTTCCGTAGGACTGGAGAAGCTTG  
CACAGAAAGGAAAATCATTGTACCTTTAGCAAGTATAACTGGAATATCACTATTTTTGATT  
ATATCCATGTGTCTTCTTCCCTATGGAAAAATATCAACCCTACAAAGTTATAAAACAGAA  
ACTAGAAGGCAGGCCAGAAACAGAAATACAGGAAAGCTCAAAACATTTTTCAGGCCATGAAGATG  
CTCTGGATGACTTCGGAATAATGAATTTGTTGCTTTTCCAGATGTTCTGGTGTTCAGG  
ATTTCCAAGCAGGTCTGTTCCAGCCTCTGATTGTGTATCGGGCAAGATTTGCACAGTACAGT  
GTATGAAGTTATTCAGCACATCCCTGCCAGCAGCAAGACCATCCAGAGTGAACCTTTTATGG  
GCTAAACAGTACATTCGAGTGAATTTCTGAAGAAACATTTTAAGGAAAAACAGTGGAAAAGT  
ATATTAATCTGGAATCAGTGAAGAAACCAGGACCAACACCTCTTACTCATTATTCCTTTACA  
TGCAGAATAGAGGCATTTATGCAAATGAACTGCAGGTTTTTCAGCATATACACAATGTCTT  
GTGCAACAGAAAAACATGTTGGGGAAATATTCCTCAGTGGAGAGTCGTTCTCATGCTGACGG  
GGAGAACGAAAAGTGACAGGGGTTTTCTCATAAGTTTTGTATGAAATATCTCTACAAAACCTCA  
ATTAGTTCTACTCTACACTTTCACTATCATCAACACTGAGACTATCCTGTCTCACCTACAAA  
TGTGGAAACTTTACATTGTTTCGATTTTTTCAGCAGACTTTGTTTTATTAATTTTTATTAGTG  
TTAAGAATGCTAAATTTATGTTTTCAATTTTTATTTCCAAATTTCTATCTTGTATTTTGTACAA  
CAAAGTAATAAGGATGGTTGTCAAAAAACAAAACCTATGCCCTCTCTTTTTTTTCAATCACC  
AGTAGTATTTTTGAGAAGACTTGTGAACACTTAAGGAAATGACTATTAAGTCTTATTTTTTA  
TTTTTTTTCAAGGAAAGATGGATTCAAATAAATTTATCTGTTTTTGCTTTAAAAA

## **FIGURE 112**

MWLKVFTTFLSFATGACSGLKVTVPSTVHGVGRGQALYLPVHYGFHTPASDIQIIWLFERPH  
TMPKYLLGSVNKSVPDLEYQHKFTMMPPNASLLINPLQFPDEGNYIVKVNIQNGTLSASQ  
KIQVTVDDPVTKPVVQIHPPSGAVEYVGNMTLTCHVEGGTRLAYQWLKNGRPVHTSSTYSFS  
PQNNTLHIAPVTKEDIGNYSCLVRNPVSEMSDIIMPIIYYGPYGLQVNSDKGLKVGEVFTV  
DLGEAILFDCSADSHPPNTYSWIRRTDNTTYIIKHGPRLEVASEKVAQKTMDYVCCAYNNIT  
GRQDETHFTVITTSVGLEKLAQKGSLSPLASITGISLFLIISMCLLFLWKKYQPYKVIKQK  
LEGRPETEYRKAQTFSGHEDALDDFGIYEFVAFPDVSGVSRIPSRVSPASDCVSGQDLHSTV  
YEVIQHIPAQQQDHPE

**Signal sequence:**

amino acids 1-18

**Transmembrane domain:**

amino acids 341-359

**N-glycosylation site.**

amino acids 73-77, 92-96, 117-121, 153-157, 189-193, 204-208,  
276-280, 308-312

**Casein kinase II phosphorylation site.**

amino acids 129-133, 198-202, 214-218, 388-392, 426-430, 433-437

**Tyrosine kinase phosphorylation site.**

amino acids 272-280

**N-myristoylation site.**

amino acids 15-21, 19-25, 118-124, 163-167, 203-209, 231-237,  
239-245

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 7-18

**FIGURE 113**

GCAAGCGGCGAAATGGCGCCCTCCGGGAGTCTTGCAGTTC CCCCTGGCAGTCC TGGTGCTGTT  
GCTTTGGGGTGC TCCCTGGACGCACGGGCGGCGGAGCAACGTTCCGCTCATCACGGGACGAGA  
ACTGGAGAGAACTGCTGGAAGGAGACTGGATGATAGAATTTTATGCCCCGTGGTGCCCTGCT  
TGTCAAATCTTCAACCGGAATGGGAAAGTTTTGCTGAATGGGGAGAAGATCTTGAGGTTAA  
TATTGCGAAAGTAGATGTCACAGAGCAGCCAGGACTGAGTGGACGGTTTATCATAACTGCTC  
TTCCTACTATTTATCATTTGTAAGATGGTGAATTTAGGCGCTATCAGGGTCCAAGGACTAAG  
AAGGACTTCATAAAC TTTATAAGTGATAAAGAGTGGAAAGATATTGAGCCCGTTTCATCATG  
GTTTGGTCCAGGTTCTGTTCTGATGAGTAGTATGTCAGCACTCTTTCAGCTATCTATGTGGA  
TCAGGACGTGCCATAACTACTTTATTGAAGACCTTGGATTGCCAGTGTGGGGATCATATACT  
GTTTTGCTTTAGCAACTCTGTTTTCCGGACTGTTATTAGGACTCTGTATGATATTTGTGGC  
AGATTGCCTTTGTCTTCAA AAAAGGCGCAGACCACAGCCATACCCATACCTTCAA AAAAAT  
TATATCAGAATCTGCACAACCTTTGAAAAAGTGGAGGAGGAA CAAGAGGCGGATGAAGAA  
GATGTTTCAGAAGAAGAAGCTGAAAGTAAAGAAGGAACAAA CAAGACTTTCCACAGAATGC  
CATAAGACAACGCTCTCTGGGTCCATCATTTGGCCACAGATAAATCCTAGT TAAATTTTATAG  
TTATCTTAATATTTATGATTTTGTATAAAAACAGAAGATTGATCATT TTTGTTTGGTTTGAAGTG  
AACTGTGACTTTTTTTGAATATTGCAGGTTTCAGTCTAGATTGTCA TTTAAATTTGAAGAGTCTA  
CATTCAGAACATAAAAGCACTAGGTATACAAGTTTGAATATGATTTAAGCACAGTATGATG  
GTTTAAATAGTCTCTAATTTTGA AAAATCGTGCCAAGCAATAAGATTTATGTATATTTGT  
TTAATAATAACCTATTTCAAGTCTGAGTTTTGAAAATTTACATTTCCCAAGTATTGCATTAT  
TGAGGTATTTAAGAAGATTATTTTAGAGAAAAATATTTCTCATTTGATATAAATTTTTCTCTG  
TTTCACTGTGTGAAAAAAGAAGATATTTCCCATAAATGGGAAGTTTGC CCATTTGTCTCAAG  
AAATGTGTATTTCA GTGACAATTTCTGTGGTCTTTTTAGAGGTATATCCAAAATTTCTTGT  
ATTTTTAGGTTATGCAACTAATAAAAAC TACCTTACATTAATTAAT TACAGTTTTCTACACA  
TGGTAATACAGGATATGCTACTGATTTAGGAAGTTTTTAAGTTCA TGGTATTCTCTTGATTC  
CAACAAAGTTTGATTTTCTCTTGATTTTTCTTACTTACTATGGGTTACATTTTTTATTTTT  
CAAATGGATGATAATTTCTTGAAACATTTTTTATGTTTTAGTAAACAGTATTTTTTTGT  
GTTTCAAAC TGAAGTTTACTGAGAGATCCATCAAATGAA CAATCTGTTGTAATTTAAAAAT  
TTGGCCACTTTTTTTCAGATTTTACATCAATCTTGCTGAACTTCAACTTGAAAATGTTTTTTTT  
TTTTTTTTGGATGTGAAGGTGAACATTCCTGATTTTTGTCTGATGTGAAAAAGCCTTGGTA  
TTTTACATTTTGA AAATTCAAAGAAGCTTAATATAAAAGTTTGCATTTCTACTCAGGAAAAAG  
CATCTTCTTGTATATGCTTAAATGTATTTTTGTCTCATATACAGAAAGTTCTTAATTTGAT  
TTTACAGTCTGTAATGCTTGATGTTTTAAAATAATAACATTTTTTATATTTTTTAAAAGACAA  
ACTTCATATATCCTGTGTTCTTTCTGACTGGTAATATTGTGTGGGATTTACAGGTA AAA  
GTCAGTAGGATGGAACATTTTAGTGTATTTTTACTCCTTAAAGAGCTAGAATACATAGTTTT  
CACCTTAAAAGAAGGGGAAAATCATAAAATACAATGAATCAACTGACCATTACGTAGTAGAC  
AATTTCTGTAATGTCCCTTCTTTCTAGGCTCTGTTGCTGTGTGAATCCATTAGATTTACAG  
TATCGTAATATACAAGTTTTCTTTAAAGCCCTCTCCTTTAGAATTTAAAATATTGTACCATT  
AAAGAGTTTGGATGTGTAAC TTGTGATGCCTTAGAAAAATATCCTAAGCACAAAATAAACCT  
TTCTAACCACTTCATTAAGCTGAAAAA AAAAAAAAAAAAAA

## **FIGURE 114**

MAPSGSLAVPLAVLVLLLWGAPWTHGRRSNVRVITDENWRELLEGDWMIEFYAPWCPACQNL  
QPEWESFAEWGEDLEVNIKVDVTEQPGLSGRFIITALPTIYHCKDGEFRRYQGPRTKKDFI  
NFISDKEWKSIEPVSSWFGPGSVLMSSMSALFQLSMWIRTCHNYFIEDLGLPVWGSYTVFAL  
ATLFSGLLLGLCMIFVADCLCPSKRRRPQYPYPYPSKLLSESAQPLKKVEEQEAEDEEDVSE  
EEAESKEGTNKDFPQNAIRQRSLGPSLATDKS

**Signal sequence:**

amino acids 1-26

**Transmembrane domain:**

amino acids 182-201

**Casein kinase II phosphorylation site.**

amino acids 68-72, 119-123, 128-132, 247-251, 257-261

**Tyrosine kinase phosphorylation site.**

amino acids 107-115

**N-myristoylation site.**

amino acids 20-26, 192-198

**Amidation site.**

amino acids 25-29

**FIGURE 115**

GCGAGTGTCCAGCTGCGGAGACCCGTGATAATTTCGTTAACTAATTCAACAAACGGGACCCTT  
CTGTGTGCCAGAAACCGCAAGCAGTTGCTAACCAGTGGGACAGGCGGATTTGGAAGAGCGGG  
AAGGTCCTGGCCAGAGCAGTGTGACACTTCCCTCTGTGACCATGAAACTCTGGGTGTCTGC  
ATTTGCTGATGGCCTGGTTTGGTGTCTGAGCTGTGTGCAGGCCGAATTTTACCTCTATTG  
GGCAGATGACTGACCTGATTTATGCAGAGAAAGAGCTGGTGCAGTCTCTGAAAGAGTACATC  
CTTGTGGAGGAAGCCAAGCTTTCCAAGATTAAGAGCTGGGCCAACAAAATGGAAGCCTTGAC  
TAGCAAGTCAGCTGCTGATGCTGAGGGCTACCTGGCTCACCTGTGAATCCCTACAAACTGG  
TGAAGCGGCTAAACACAGACTGGCCTGCGCTGGAGGACCTTGTCTCTGCAGGACTCAGCTGCA  
GGTTTTATCGCCAACCTCTCTGTGCAGCGGCAGTTCTTCCCCTGATGAGGACGAGATAGG  
AGCTGCCAAAGCCCTGATGAGACTTCAGGACACATACAGGCTGGACCAGGCACAATTTCCA  
GAGGGGAACCTTCCAGGAACCAAGTACCAGGCAATGCTGAGTGTGGATGACTGCTTTGGGATG  
GGCCGCTCGGCCTACAATGAAGGGGACTATTATCATAACGGTGTGTGGATGGAGCAGGTGCT  
AAAGCAGCTTGATCCCGGGGAGGAGGCCACCACAACCAAGTCACAGGTGCTGGACTACCTCA  
GCTATGCTGTCTTCCAGTTGGGTGATCTGCACCGTGCCCTGGAGCTCACCCGCGCCTGCTC  
TCCCTTGACCCAAAGCCACGAACGAGCTGGAGGGAATCTGCGGTACTTTGAGCAGTTATTGGA  
GGAAGAGAGAGAAAAAACGTTAAACAAATCAGACAGAAGCTGAGCTAGCAACCCAGAAGGCA  
TCTATGAGAGGCCTGTGGACTACCTGCCTGAGAGGGATGTTTACGAGAGCCTCTGTGCTGGG  
GAGGTGTCAAACCTGACACCCCGTAGACAGAAGAGGCTTTTCTGTAGGTACCACCATGGCAA  
CAGGGCCCCACAGCTGCTCATTGCCCCCTTCAAAGAGGAGGACGAGTGGGACAGCCCGCACA  
TCGTCAGGTAACGATGTCTGTCTGATGAGGAAATCGAGAGGATCAAGGAGATCGCAAAA  
CCTAAACTTGCACCGAGCCACCGTTCTGTGATCCCAAGACAGGAGTCTCACTGTCGCCAGCTA  
CCGGGTTTCCAAAAGCTCCTGGCTAGAGGAAGATGATGACCCTGTTGTGGCCCGAGTAAATC  
GTCGGATGCAGCATATCACAGGGTTAACAGTAAAGACTGCAGAATTTGTTACAGGTTGCAAA  
TATGGAGTGGGAGGACAGTATGAACCGCACTTCGACTTCTCTAGGCGACCTTTTGACAGCGG  
CCTCAAAACAGAGGGGAATAGGTTAGCGACGTTTTCTTAACACATGAGTGTGATGATGAGAAGCTG  
GTGGTGCCACCGTCTTCCCTGATCTGGGGGCTGCAATTTGGCCTAAGAAGGGTACAGCTGTG  
TTCTGGTACAACCTCTTGGGAGCGGGGAAGGTGACTACCGAACAAAGACATGCTGCCTGCCC  
TGTGCTTGTGGGCTGCAAGTGGGTCTCCAATAAGTGGTTCATGAACGAGGACAGGAGTTCT  
TGAGACCTTGTGGATCAACAGAAGTTGACTGAACATCCTTTTCTGTCTTCCCCTTCCCTGGTC  
CTTCAGCCCATGTCAACGTGACAGACACCTTTGTATGTTCCCTTTGTATGTTCCATACAGGCT  
GATTTTGGAGAAATGAATGTTTGTCTGGAGCAGAGGGAGACCATACTAGGGCGACTCCTGT  
GTGACTGAAGTCCCAGCCCTTCCATTAGCCTGTGCCATCCCTGGCCCCAAGGCTAGGATCA  
AAGTGGCTGCAGCAGAGTTAGCTGTCTAGCGCCTAGCAAGGTGCCTTTGTACCTCAGGTGTT  
TTAGGTGTGAGATGTTTCAGTGAACCAAAAGTTCTGATACCTTGTTTACATGTTTGTTTTTAT  
GGCATTTCTATCTATTGTGGCTTTACCAAAAAATAAAATGTCCCTACCAGAAAAA

## **FIGURE 116**

MKLWVSALLMAWFGVLSVCVQAEFFTSIGHMTDLIYAEKELVQSLKEYILVEEAKLSKIKSWA  
NKMEALTSKSAADAEGYLAHPVNAYKLVKRLNTDWPALDVLVQDSAAGFIANLSVQRQFFP  
TDEDEIGAAKALMRLQDFTYRLDPGTISRGEPLGTYQAMLSVDDCFGMGRSAYNEGDYYHTV  
LWMEQVLKQLDAGEEATTTKSQVLDYLSYAVFQLGDLHRALELTRRLSLDPSHERAGGNLR  
YFEQLLEEREKTLTNQTEAELATPEGIYERPVDYLPERDVYESLCRGEVGLTPRRQKRLF  
CRYHHGNRAPQLLIAPFKEEEDWDSPHIVRYDVMSEETIERIKEIAKPKLARATVRDPKTF  
VLTVASYRVSKSSWLEEDDDPVVARVNRMMQHITGLTVKTAELLQVANYGVGGQYEPHFDFS  
RRPFDSGLKTEGNRLATFLNYMSDVEAGGATVFPDLGAAIWPKKGTAVFWYNLLRSGECDYR  
TRHAACPVLVGCKWVSNKWFHERGQEFRLPCGSTEVD

**Signal sequence:**

amino acids 1-17

**N-glycosylation site.**

amino acids 115-119, 264-268

**Glycosaminoglycan attachment site.**

amino acids 490-494

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 477-481

**Casein kinase II phosphorylation site.**

amino acids 43-47, 72-76, 125-129, 151-155, 165-169, 266-270,  
346-350, 365-369, 385-389, 457-461, 530-534

**Tyrosine kinase phosphorylation site.**

amino acids 71-80, 489-496

**N-myristoylation site.**

amino acids 14-20, 131-137, 171-177, 446-452

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 8-19

**Leucine zipper pattern.**

amino acids 213-235

**FIGURE 117**

GCAGTATTGAGTTTACTTCTCTCTTTTGGTGGAAAGACAGACCATAATCCAGTGTGAGTGAAATGATTGT  
TTCATTTATTACCGTTTTGGCTGGGGTTAGTTCGACACCTTACAGTTGAAGAGCAGGCAGGAGGAGTTGTGA  
AGACAGGACAATCTTCTGGGGATGCTGGTCTGGAAGCCAGCGGGCTTGTCTGTCTTTGGCCTCATTGACCC  
CAGTTCTCTGGTAAAACTGAAGCCCTACTACTGGCCTGGTGCCCATCAATCCATTGATCCTTGAGGCTGTGCC  
CCTGGGGCACCCACTGGCAGGGCCTACCACCAATGCGACTGAGCTCCTCTGTTGGCTCTGCTGCGGCCAGCGCTTC  
CCTTCATCTTAGGGTGTCTCTGGGGTGACCCCTGAGCCTCCTGCGGGTTTCTGGATCCAGGGGAGGGAGAAG  
ATCCTCTGTGTCGAGGCTGTAGGGGAGCGAGGAGGGCCACAGAATCCAGATTGAGAGCTCGGCTAGACCAAAGTG  
ATGAGACTTCAAACCCCGATTGTCCCTACTACAGGGACCCCAACAGCCCTACAAGAAGGTGCTCAGGACTC  
GGTACATCCAGACAGAGCTGGGCTCCGCTGAGCGGTTGCTGGTGGCTGTCTGACCTCCCGAGCTACACTGTCCA  
CTTTGGCCCTGGCTGTGAACCGTACGGTGGCCCATCACTTCCCTCGGTTACTCTACTTCACTGGGCAGCGGGGG  
CCCGGGCTCCAGCAGGGATGCGAGTGGTGTCTCATGGGGATGAGCGGCCCGCTGGCTCATGTGAGAGCCCTGC  
GCCACTTCCACACACTTTGGGGCCGACTACGACTGGTTCCTCATCATGCAGGATGACACATATGTGCGAGGCC  
CCCGCTGGCAGCCTTGTGGCCACCTCAGCATCAACCAAGACCTGTACTTAGCCCGGCAGAGGAGTTCAATG  
GCGCAGGCGAGCAGGCCCGTACTGTCTATGGGGCTTTGGCTACCTGTTGTACCGAGTCTCCTGCTTCGTCTGC  
GGCCACATCTGGATGGCTGCCGAGGAGACATCTCAGTGCCCTCCTGACGAGTGGCTTGGACCTGCCTCATTG  
ACTCTCTGGGCGTGGCTGTGTCTCACAGCACCAGGGGCGAGTATCGCTCATTTGAACTGGCCAAAATAGGG  
ACCTGAGAAGGAAGGGAGCTCGGCTTTCCTGAGTGCCTTCGCGGTGCACCCCTGTCTCGAAGGTACCCCTCATGT  
ACCGCTCCACAAACGCTTTCAGCGCTCTGGAGTTGGAGCGGGCTTACAGTGAATAGAAACAATGCAGGCTCAGA  
TCCGGAACCTGACCGTGTGACCCCGAAGGGGAGGCAGGGCTGAGCTGGCCCGTGGGCTCCCTGCTCCTTTCA  
CACCACACTCTCGCTTTGAGGTGCTGGGCTGGGACTACTTCACAGAGCAGCACACCTTCTCCTGTGCAGATGGGG  
CTCCCAAGTGCCACTACAGGGGCTAGCAGGGCGGACGTGGGTGATGCGTTGGAGACTGCCCTGGAGCAGCTCA  
ATCGGCGCTATCAGCCCCGCTGCGCTTCCAGAAGCAGCGACTGCTCAACGGCTATCGGCGCTTCGACCCAGC  
GGGGCATGGAGTACACCTGGACCTGCTGTGGAATGTGTGACACAGCGTGGGCACCGCGGGCCCTGGCTCGCA  
GGGTGAGCCTGCTGCGGCCACTGAGCCGGGTGGAATCCTACCTATGCCCTATGTCACTGAGGCCACCCGAGTGC  
AGCTGGTGTGCCACTCCTGGTGTGCTGAAGCTGTGCGAGCCCGGCTTTCCTCGAGGCGTTTGAGCCAAATGTCC  
TGGAGCCACGAGAACATGCATTGCTCACCTGTGTCTGGTCTACGGGCCACGAGAAGGTGGCCGTGGAGCTCCAG  
ACCCATTTCTTGGGGTGAAGGCTGCAGCAGCGGAGTTAGAGCGACGGTACCC'TGGGACGAGGCTGGCCTGGCTCG  
CTGTGCGAGCAGAGGCCCTTCCAGGTGCGACTCATGGACGTGGTCTCGAAGAAGCACCCCTGTGGACACTCTCT  
TCTTCTTACCACCGTGTGGACAAGGCCTGGGCCCGAAGTCCCTCAACCGCTGTGCGATGAATGCCATCTCTGGCT  
GGCAGGCTTCTTTCCAGTCCATTTCCAGGAGTTCAATCCTGCCCTGTCAACCAAGAGATCACCCCAAGGGCC  
CGGGGCTGGCCCTGACCCCCCTCCCTCCTGGTGTGACCCCTCCCGGGGGCTCCTATAGGGGGGAGATTTG  
ACCGGAGGCTTCTGCGGAGGGCTGCTTCTACAACGCTGACTACCTGGCGGCCCGAGCCCGGCTGGCAGGTGAAC  
TGGCAGGCCAGGAAGAGGAGGAAGCCCTGGAGGGGCTGGAGGTGATGGATGTTTTCTCCGGTTCTCAGGGCTCC  
ACCTCTTTCGGGCCGTAGAGCCAGGCTGGTGAGAAGTTCTCCTGCGAGACTGCAGCCACGGCTCAGTGAAG  
AACTTACCACCGCTGCCGCTCAGCAACTGGAGGGCTAGGGGGCCGTGCCAGCTGGCTATGGCTCTCTTTG  
AGCAGGAGCAGGCCAATAGCACTTAGCCCGCTGGGGCCCTAACCTCATACCTTTCTTTGTCTGCCTCAGCC  
CCAGGAAGGCCAAGCAAGATGGTGGACAGATAGAAATTGTTGCTGTATTTTTAAATATGAAAATGTTATTAA  
ACATGTCTTCTGCC

**FIGURE 118**

MRLSSLLALLRPALPLILGLSLGCSLSLLRVSWIQEGEGEDPCVEAVGERGGPQNPDSRARLD  
QSDDFKPRIVPYRDPNKPYPYKVLRLRYIQTELGSRERLLVAVLTSRATLSTLAVAVNRTV  
AHHFPRLLYFTGQRGARAPAGMQVVSNGDERPAWLMSETLRHLHTHFGADYDFFFIMQDDTY  
VQAPRLAALAGHLSINQDLYLGRAEEFIGAGEQARYCHGGFGYLLSRLLLLRPHLDGCRG  
DILSARPDEWLGRCLIDSLGVCVSHQHQQQYRSFELAKNRDPEKEGSSAFLSAFVHPVSE  
GTLMYRLHKRFSALELERAYSEIEQLQAQIRNLTVLTPEGEAGLSWPVGLPAPFTPHSRFEV  
LGWDYFTEQHTFSCADGAPKCPLOQASRADVGDALLETALQLNRRYQPRLLRFQKQRLNNGYR  
RFDPARMEYTLDLLLECVTQRGHRRALARRVSLLRPLSRVEILPMPYVTEATRVQLVLPPL  
VAEAAAAPAFLEAFAANVLEPREHALLTLLLTVYGPREGGRGAPDPFLGVKAAAELERRYPG  
TRLAWLAVRAEAPSQVRLMDVVSKKHPVDTLFFLTWTRPGPEVLNRCRMNAISGWQAFPP  
VHFQEFNPALSPQRSPPGPPGAGDPSPGADPSRGAPIGGRFDRQASAEGCFYNADYLAA  
RARLAGELAQEEEEALEGLEVMDFLRFSGHLHFRAVEPGLVQKFSLRDCSPRLSEELYHR  
CRLSNLEGLGGRAQLAMALFEQEANST

**Signal sequence:**

amino acids 1-15

**Transmembrane domain:**

amino acids 489-507

**N-glycosylation site.**

amino acids 121-125, 342-346

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 319-323, 464-468

**Casein kinase II phosphorylation site.**

amino acids 64-68, 150-154, 322-326, 331-337, 368-372, 385-389,  
399-403, 409-413, 473-477, 729-733, 748-752

**Tyrosine kinase phosphorylation site.**

amino acids 736-743

**N-myristoylation site.**

amino acids 19-25, 23-29, 136-142, 397-403, 441-447, 544-550,  
558-564, 651-657, 657-663, 672-678

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 14-25

**Cell attachment sequence.**

amino acids 247-250



**FIGURE 119**

CGGAGTGGTGCGCCAACGTGAGAGGAAACCCGTGCGCGGCTGCGCTTTCCTGTCCCCAAGCC  
GTTCTAGACGCGGGAAAAATAGCTTTCTGAAAGCAGCTCCTTTTTGAAAGGGTGTGATGCTTGG  
AAGCATTTTCTGTGCTTTTGATCACTATGCTAGGACACATTAGGATTGGTCATGGAAATAGAA  
TGCACCACCATGAGCATCATCACCTACAAGCTCCTAACAAAGAAGATATCTTGAAAAATTTCA  
GAGGATGAGCGCATGGAGCTCAGTAAGAGCTTTCGAGTATACTGTATTATCCTTGTA AAAAC  
CAAAGATGTGAGTCTTTGGGCTGCAGTAAAGGAGACTTGGACCAAACACTGTGACAAAAGCAG  
AGTTCCTCAGTTCTGAAAATGTTAAAGTGTGAGTCAATTAATATGGACACAAATGACATG  
TGGTTAATGATGAGAAAAGCTTACAAATACGCCTTTGATAAGTATAGAGACCAATACAACTG  
GTTCTTCCTTGCACGCCCCACTACGTTTGCTATCATTGAAAACCTAAAGTATTTTTTGTAA  
AAAAGGATCCATCACAGCCTTTCTATCTAGGCCACACTATAAAATCTGGAGACCTTGAATAT  
GTGGGTATGGAAGGAGGAATTGTCTTAAGTGTAGAATCAATGAAAAGACTTAAACAGCCTTCT  
CAATATCCCAGAAAAGTGTCTGAACAGGGAGGGATGATTTGGAAGATATCTGAAGATAAAC  
AGCTAGCAGTTTGCCTGAAAATATGCTGGAGTATTTGCAGAAAATGCAGAAGATGCTGATGGA  
AAAGATGTATTTAATACCAAATCTGTTGGGCTTTCTATTAAAGAGGCAATGACTTATCACCC  
CAACCAGGTAGTAGAAGGCTGTTGTTTCTAGATATGGCTGTTACTTTAATGGACTGACTCCAA  
ATCAGATGCATGTGATGATGTATGGGGTATACCGCCTTAGGGCATTGTTGGGCATATTTTCAAT  
GATGCATTGGTTTTCTTACCTCCAATGGTTCTGACAATGACTGAGAAGTGGTAGAAAAGCG  
TGAATATGATCTTTGTATAGGACGTGTGTTGTCATTATTTGTAGTAGTAACTACATATCCAA  
TACAGCTGTATGTTTCTTTTTCTTTTCTAATTTGGTGGCACTGGTATAACCACACATTAAAG  
TCAGTAGTACATTTTTAAATGAGGGTGGTTTTTTTTCTTTAAAACACATGAACATTGTA AATG  
TGTGAAAAGAAGTGTTTAAGAATAATAATTTTGCAAATAAACTATTAATAAATATTATAT  
GTGATAAATTCTAAATTATGAACATTAGAAATCTGTGGGGCACATATTTTTGCTGATTGGTT  
AAAAAATTTTAAACAGGTCTTTAGCGTTCTAAGATATGCAAATGATATCTCTAGTTGTGAATT  
TGTGATTAAAGTAAAACTTTTAGCTGTGTGTTCCCTTTACTTCTAATACTGATTTATGTTCT  
AAGCCTCCCCAAGTTCCAATGGATTTGCCTTCTCAAATGTACAACCTAAGCAACTAAAGAAA  
ATTAAAGTAAAAGTTGAAAAAT

**FIGURE 120**

MLSESSSFLKGVMLGSIFCALITMLGHIRIGHGNRMHHHEHHHLQAPNKEDILKISEDERME  
LSKSFrvYCIILVKPKDVSLWAAVKETWTKHCDKAEFFSSENVKVFESINMDTNDMWLMMRK  
AYKYAFDKYRDQYNWFFLARPTTFAIIEENLKYFLLKKDPSQPFYLGHTIKSGDLEYVGMEGG  
IVLSVESMKRLNSLLNIPEKCPEQGGMIWKISEDKQLAVCLKYAGVFAENAEDADGKDVFN  
KSVGLSIKEAMTYHPNQVVEGCCSDMAVTFNGLTPNQMHVMMYGVYRLRAFGHIFNDALVFL  
PPNGSDND

**Signal sequence:**

amino acids 1-33

**N-glycosylation site.**

amino acids 121-125, 342-346

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 319-323, 464-468

**Casein kinase II phosphorylation site.**

amino acids 64-132, 150-154, 322-326, 331-335, 368-372, 385-389,  
399-403, 409-413, 473-477, 729-733, 748-752

**Tyrosine kinase phosphorylation site.**

amino acids 736-743

**N-myristoylation site.**

amino acids 19-25, 23-29, 136-142, 397-403, 441-447, 544-550,  
558-564, 651-657, 657-663, 672-672

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 14-25

**Cell attachment sequence.**

amino acids 247-250



**FIGURE 122**

MNSSKSSETQCTERGCFFSSQMFLWTVAGIPILFLSACFITRCVVTFRIFQTCDEKKFQLPEN  
FTELSYNYGSGSVKNCCPLNWEYFQSSCYFFSTDTISWALSLKNCSSAMGAHLVVINSQEEQ  
EFLSYKKPKMREFFIGLSDQVVEGQWQWVDGTPLTKSLSFWDVGEPPNIATLEDCATMRDSS  
NPRQNWNDVTCFLNYFRICEMVGINPLNKGKSL

**Signal sequence:**

amino acids 1-42

**N-glycosylation site.**

amino acids 2-6, 62-66, 107-111

**Casein kinase II phosphorylation site.**

amino acids 51-55, 120-124, 163-167, 175-179, 181-185

**N-myristoylation site.**

amino acids 15-21, 74-80, 155-161

**Prokaryotic membrane lipoprotein lipid attachment site.**

amino acids 27-38

**FIGURE 123**

GGGACTACAAGCCGCGCCGCTGCGCTGGCCCTCAGCAACCCCTCGACATGGCGCTGAGGCGGCCACCGCGAC  
TCCGGCTCTGCGCTCGGCTGCCTGACTTCTTCTCTGCTGCTTTTTCAGGGGCTGCCGTAGAGGGCTGTAAATC  
TCAAATCCAGCAATCGAACCCCACTGGTACAGGAATTTGAAAGTGTGGAACGTCTTGATCATATTACGGATTTCGC  
AGACAAGTGACCCAGGATCGAGTGGAGAAAATCAAGATGAACAAACCACATATGTGTTTTTGACAACAAAA  
TTCAGGGAGACTTGGCGGGTCTGTCAGAAAATCTGGGGAAGACATCCCTGAAGATCTGGAATGTGACACGGAGAG  
ACTCAGCCCTTATCGCTGTGAGGTCTGTGCTCGAAATGACCGCAAGGAAATGTATGAGATTGTGATCGAGTTAA  
CTGTGCAAGTGAAGCCAGTGACCCCTGTCTGTAGAGTGCAGAGGCTGTACCAGTAGGCAAGATGGCAACACTGC  
ACTGCCAGGAGAGTGAGGGCCACCCCGGCTCACTACAGCTGGTATCGCAATGATGTACCACAGCCACGGATT  
CCAGAGCCAATCCGATTTTCGCAATTTCTTTCCACTTAAACTCTGAAACAGGCACCTTTGGTGTTCACCTGCTG  
TTCACAAGGACGACTCTGGGCAGTACTACTGCTTCCAATGACCGAGGCTCAGCCAGGTGTGAGGAGCAGG  
AGATGGAAGTCTATGACCTGAACATTTGGCGGAATTAATGGGGGGTCTGGTTGCTCTGCTGACTGGCCCTGA  
TCACGTTGGGCATCTGCTGTGCATACAGACGTGGCTACTTTCATCAACAATAAACAGGATGGAGAAAGTTACAAGA  
ACCCAGGAAACCCAGATGGAGTTAACTACATCCGCACTGACGAGGAGGGCGACTTCAGACACAAGTCTCGTTG  
TGATCTGAGACCCGCGGTGTGGCTGAGAGCGCACAGAGCGCAGTGCATACCTCTGCTAGAAACTCCGTCAA  
GGCAGCGAGAGCTGATGCACTCGACAGAGCTAGACACTCATTGAAAGCTTTTCGTTTTGGCCAAAGTTGACCA  
CTACTCTTCTACTCTAACAAGCCACATGAATAGAAGAAATTTCCCTCAAGATGGACCCGGTAAATATAACACAA  
GGAAGCGAAATCGGTGCGTTCATGAGTGGGTTCCCTAATCTGTTCTGGCTGATCCCGCATGAGTATTAGG  
GTGATCTTAAAGAGTTTGTCTCACGTAACCGCCGCTGCTGGGCCCTGTAAGCCAGCATGTTCCACTGGTCTG  
CAGCAGCCACGACAGCACCATGTGAGATGGCGAGGTGGCTGGACAGCACAGCAGCGCATCCCGCGGGAAACCA  
GAAAAGGCTTCTACACAGCAGCCCTTACTTTCATCGGCCACAGACACCACCCAGTTTCTTCTTAAAGGCTCTGC  
TGATCGGTGTTGCAAGTTCATTTGTGGAGAAGCTTTTGGATCAGCATTTTGTAAAAACAACAAAATCAGGAAG  
GTAAATTTGGTTGCTGGAAGAGGGATCTTGCTGAGGAACCTGCTTGTCCACAGGGTGTGAGGATTTAAGGAAA  
ACCTTCGCTTAGGCTAAGTCTGAAATGGTACTGAAATATGCTTTTCTATGGGTCTTGTATTTTATAAAATTT  
TACATCTAAATTTTGTCTAAGGATGTATTTGATTAATGAAAAGAAAATTTCTATTTAAACTGTAAATATATTGT  
CATACAATGTTAAATAACCTATTTTTTAAAAAAGTTCAACTTAAGGTAGAAGTTCAAGCTACTAGTGTAAAT  
TGGAAAATATCAATAATTAAGAGTATTTTACCCAAGGAATCCTCTCATGGAAGTTTACTGTGATTTCTTTTCT  
CACACAAGTTTAGCCTTTTTTACAAGGGAACCTCATACTGTCTACACATCAGACCATAGTTGCTTAGGAAACCTT  
TAAAAATCCAGTTAAGCAATGTTGAAATCAGTTTGCATCTCTTCAAAAGAAACCTCTCAGGTTAGCTTTGAACT  
GCCTCTTCTGAGATGACTAGGACAGTCTGTACCCAGAGGCCACCCAGAAGCCCTCAGATGTACATACACAGATG  
CCAGTCAGCTCCTGGGGTGGCCAGGCGCCCGCTCTAGCTCACTGTTGCTCGCTGTCTGCCAGGAGGCCCT  
GCCATCCTTGGGCCCTGGCAGTGGCTGTGTCAGTGGCTTTACTCACGTGGCCCTTGCTTCATCCAGCACAGC  
TCTCAGGTGGGCACTGCAGGGACACTGGTGTCTTCCATGTAGCGTCCCAGCTTTGGGCTCCTGTAACAGACCTCT  
TTTTGGTTATGGATGGCTCAAAAATAGGGCCCCAATGCTATTTTTTTTTTTAAGTTTGTAAATTAATTTGTT  
AAGATTGTCTAAGGCCAAAGGCAATGCGAAATCAAGTCTGTCAAGTACAATAACATTTTTTAAAGAAAATGGAT  
CCCCTGTTCTCTTTGCCACAGAGAAAGCACCCAGAGCCACAGGCTCTGTGCAATTTCAAAACAAACCATGAT  
GGAGTGGCGCCAGTCCAGCCTTTTAAAGAAGCTCAGGTGGAGCAGCCAGGTGAAAGGCTGGCGGGGAGGAAAG  
TGAAACGCCGTAATCAAAGCAGTTTTCTAATTTGACTTTAAATTTTTTCATCCGCGGAGACACTGCTCCCAT  
TGTGGGGGACATTAGCAACATCACTCAGAAGCTGTGTCTTCAAGAGCAGGTGTTCTCAGCCTCACATGCCCT  
GCCGTGCTGGACTCAGGACTGAAGTGTGTAAGCAAGGAGCTGCTGAGAAGGAGCACTCCACTGTGTGCTTGG  
GAATGGCTCTCACTACTCACCTTGTCTTTTCAAGTCTCAGTGTCTTGGGTTTTTATACTTTGACAGCTTTTTTT  
AATTGCATACATGAGACTGTGTTGACTTTTTTTAGTTATGTGAAACACTTTGCCCGAGGCCGCTGGCAGAGGCA  
GGAAATGCTCCAGCAGTGGCTCAGTGTCTCCCTGGTGTCTGTGCATGGCATCCTGGATGCTTAGCATGCAAGTTC  
CCTCCATCATTGCCACCTTGGTAGAGAGGGATGGCTCCCCACCTCAGCGTTGGGGATTCCAGCTCCAGCCTCCT  
TCTTGGTTGTATAGTATAGGGTAGCCTTATTGCCCCCTTCTTATACCTAAAACCTTCTACACTAGTGGCA  
TGGGAACCGGTCTGAAAAGTAGAGAGAAGTGAAGTAGAGTCTGGGAAGTAGCTGCCTATAACTGAGACTAGA  
CGGAAAAGGAATACTCGTGTATTTAAGATATGAATGTGACTCAAGACTCGAGGCCGATACGAGGCTGTGATTCT  
GCCTTTGGATGGATGTTGCTGTACACAGATGCTACAGACTTGTACTAACACACCCGTAATTTGGCATTGTTTAA  
CTCATTTAATAAAGCTTCAAAAAACCA

## FIGURE 124

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA77624
><subunit 1 of 1, 310 aa, 1 stop
><MW: 35020, pI: 7.90, NX(S/T): 3
MALRRPPRLRLCARLPDFFLLLLFRGCLIGAVNLKSSNRTPVVQEFESVELSCIITDSQTS
D
PRIEWKKIQDEQTTYVFFDNKIQGDLAGRAEILGKTSLSKIWNVTRRDSALYRCEVVARNDRK
EIDEIVIELTVQVKPVTTPVCRVPKAVPVGKMATLHCQESEGHPRPHYSWYRNDVPLPTDSRA
NPRFRNSSFHLNSETGTLVFTAVHKDDSGQYYCIASNDAGSARCEEQEMEVDLNIIGGIIGG
VLVVLAVLALITLIGICCAYYRRGYFINNKQDGESYKNPGKPDGVNYIRTDEEGDFRHKSSFVI
```

**Important features of the protein:**

**Signal peptide:**

amino acids 1-30

**Transmembrane domain:**

amino acids 243-263

**N-glycosylation sites.**

amino acids 104-107, 192-195

**cAMP- and cGMP-dependent protein kinase phosphorylation site.**

amino acids 107-110

**Casein kinase II phosphorylation site.**

amino acids 106-109, 296-299

**Tyrosine kinase phosphorylation site.**

amino acids 69-77

**N-myristoylation sites.**

amino acids 26-31, 215-220, 226-231, 243-248, 244-249, 262-267

**SECRETED AND TRANSMEMBRANE  
POLYPEPTIDES AND NUCLEIC ACIDS  
ENCODING THE SAME**

**FIELD OF THE INVENTION**

[0001] The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel polypeptides.

**BACKGROUND OF THE INVENTION**

[0002] Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

[0003] Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.* 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0004] Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and nerve growth factor receptor.

[0005] Membrane-bound proteins and receptor molecules have various industrial applications, including as pharma-

ceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

[0006] Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

[0007] 1. PRO211 and PRO217

[0008] Epidermal growth factor (EGF) is a conventional mitogenic factor that stimulates the proliferation of various types of cells including epithelial cells and fibroblasts. EGF binds to and activates the EGF receptor (EGFR), which initiates intracellular signaling and subsequent effects. The EGFR is expressed in neurons of the cerebral cortex, cerebellum, and hippocampus in addition to other regions of the central nervous system (CNS). In addition, EGF is also expressed in various regions of the CNS. Therefore, EGF acts not only on mitotic cells, but also on postmitotic neurons. In fact, many studies have indicated that EGF has neurotrophic or neuromodulatory effects on various types of neurons in the CNS. For example, EGF acts directly on cultured cerebral cortical and cerebellar neurons, enhancing neurite outgrowth and survival. On the other hand, EGF also acts on other cell types, including septal cholinergic and mesencephalic dopaminergic neurons, indirectly through glial cells. Evidence of the effects of EGF on neurons in the CNS is accumulating, but the mechanisms of action remain essentially unknown. EGF-induced signaling in mitotic cells is better understood than in postmitotic neurons. Studies of cloned pheochromocytoma PC12 cells and cultured cerebral cortical neurons have suggested that the EGF-induced neurotrophic actions are mediated by sustained activation of the EGFR and mitogen-activated protein kinase (MAPK) in response to EGF. The sustained intracellular signaling correlates with the decreased rate of EGFR down-regulation, which might determine the response of neuronal cells to EGF. It is likely that EGF is a multi-potent growth factor that acts upon various types of cells including mitotic cells and postmitotic neurons.

[0009] EGF is produced by the salivary and Brunner's glands of the gastrointestinal system, kidney, pancreas, thyroid gland, pituitary gland, and the nervous system, and is found in body fluids such as saliva, blood, cerebrospinal fluid (CSF), urine, amniotic fluid, prostatic fluid, pancreatic juice, and breast milk, Plata-Salaman, *Peptides* 12: 653-663 (1991).

[0010] EGF is mediated by its membrane specific receptor, which contains an intrinsic tyrosine kinase. Stoscheck et al., *J. Cell Biochem.* 31: 135-152 (1986). EGF is believed to function by binding to the extracellular portion of its receptor which induces a transmembrane signal that activates the intrinsic tyrosine kinase.

[0011] Purification and sequence analysis of the EGF-like domain has revealed the presence of six conserved cysteine residues which cross-bind to create three peptide loops, Savage et al., *J. Biol. Chem.* 248: 7669-7672 (1979). It is

now generally known that several other peptides can react with the EGF receptor which share the same generalized motif  $X_nCX_7CX_{4/5}CX_{10}CXCX_5GX_2CX_n$ , where X represents any non-cysteine amino acid, and n is a variable repeat number. Non isolated peptides having this motif include TGF- $\alpha$ , amphiregulin, schwannoma-derived growth factor (SDGF), heparin-binding EGF-like growth factors and certain virally encoded peptides (e.g., Vaccinia virus, Reiser, *Nature* 313: 801-803 (1985), Shope fibroma virus, Chang et al., *Mol Cell Biol.* 7: 535-540 (1987), Molluscum contagiosum, Porter and Archard, *J. Gen. Virol.* 68: 673-682 (1987), and Myxoma virus, Lemon et al., *J. Virol.* 61: 1271-1275 (1987), Prigent and Lemoine, *Prog. Growth Factor Res.* 4: 1-24 (1992).

[0012] EGF-like domains are not confined to growth factors but have been observed in a variety of cell-surface and extracellular proteins which have interesting properties in cell adhesion, protein-protein interaction and development, Laurence and Gusterson, *Tumor Biol.* 11: 229-261 (1990). These proteins include blood coagulation factors (factors VI, IX, X, XII, protein C, protein S, protein Z, tissue plasminogen activator, urokinase), extracellular matrix components (laminin, cytactin, entactin), cell surface receptors (LDL receptor, thrombomodulin receptor) and immunity-related proteins (complement C1r, uromodulin).

[0013] Even more interesting, the general structure pattern of EGF-like precursors is preserved through lower organisms as well as in mammalian cells. A number of genes with developmental significance have been identified in invertebrates with EGF-like repeats. For example, the notch gene of *Drosophila* encodes 36 tandemly arranged 40 amino acid repeats which show homology to EGF, Wharton et al., *Cell* 43: 557-581 (1985). Hydrophathy plots indicate a putative membrane spanning domain, with the EGF-related sequences being located on the extracellular side of the membrane. Other homeotic genes with EGF-like repeats include Delta, 95F and 5ZD which were identified using probes based on Notch, and the nematode gene Lin-12 which encodes a putative receptor for a developmental signal transmitted between two specified cells.

[0014] Specifically, EGF has been shown to have potential in the preservation and maintenance of gastrointestinal mucosa and the repair of acute and chronic mucosal lesions, Konturek et al., *Eur. J. Gastroenterol Hepatol.* 7 (10), 933-37 (1995), including the treatment of necrotizing enterocolitis, Zollinger-Ellison syndrome, gastrointestinal ulceration gastrointestinal ulcerations and congenital microvillus atrophy, Guglietta and Sullivan, *Eur. J. Gastroenterol Hepatol.* 7(10), 945-50 (1995). Additionally, EGF has been implicated in hair follicle differentiation; du Cros, *J. Invest. Dermatol.* 101 (1 Suppl.), 106S-113S (1993), Hillier, *Clin. Endocrinol.* 33(4), 427-28 (1990); kidney function, Hamm et al., *Semin. Nephrol.* 13 (1): 109-15 (1993), Harris, *Am. J. Kidney Dis.* 17(6): 627-30 (1991); tear fluid, van Setten et al., *Int. Ophthalmol* 15(6); 359-62(1991); vitamin K mediated blood coagulation, Stenflo et al., *Blood* 78(7): 1637-51(1991). EGF is also implicated various skin disease characterized by abnormal keratinocyte differentiation, e.g., psoriasis, epithelial cancers such as squamous cell carcinomas of the lung, epidermoid carcinoma of the vulva and gliomas. King et al., *Am. J. Med. Sci.* 296: 154-158 (1988).

[0015] Of great interest is mounting evidence that genetic alterations in growth factors signaling pathways are closely linked to developmental abnormalities and to chronic diseases including cancer. Aaronson, *Science* 254: 1146-1153 (1991). For example, c-erb-2 (also known as HER-2), a proto-oncogene with close structural similarity to EGF receptor protein, is overexpressed in human breast cancer. King et al., *Science* 229: 974-976 (1985); Gullick, *Hormones and their actions*, Cooke et al., eds, Amsterdam, Elsevier, pp 349-360 (1986).

[0016] We herein describe the identification and characterization of novel polypeptides having homology to EGF, wherein those polypeptides are herein designated PRO211 and PRO217.

[0017] 2. PRO230

[0018] Nephritis is a condition characterized by inflammation of the kidney affecting the structure and normal function of the kidney. This condition can be chronic or acute and is generally caused by infection, degenerative process or vascular disease. In all cases, early detection is desirable so that the patient with nephritis can begin treatment of the condition.

[0019] An approach to detecting nephritis is to determine the antigens associated with nephritis and antibodies thereto. In rabbit, a tubulointerstitial nephritis antigen (TIN-ag) has been reported in Nelson, T. R., et al., *J. Biol. Chem.*, 270(27):16265-70 (July 1995) (GENBANK/U24270). This study reports that the rabbit TIN-ag is a basement membrane glycoprotein having a predicted amino acid sequence which has a carboxyl-terminal region exhibiting 30% homology with human preprocathepsin B, a member of the cysteine proteinase family of proteins. It is also reported that the rabbit TIN-ag has a domain in the amino-terminal region containing an epidermal growth factor-like motif that shares homology with laminin A and S chains, alpha 1 chain of type I collagen, von Willebrand's factor and mucin, indicating structural and functional similarities. Studies have also been conducted in mice. However, it is desirable to identify tubulointerstitial nephritis antigens in humans to aid in the development of early detection methods and treatment of nephritis.

[0020] Proteins which have homology to tubulointerstitial nephritis antigens are of particular interest to the medical and industrial communities. Often, proteins having homology to each other have similar function. It is also of interest when proteins having homology do not have similar functions, indicating that certain structural motifs identify information other than function, such as locality of function. We herein describe the identification and characterization of a novel polypeptide, designated hgerin as PRO230, which has homology to tubulointerstitial nephritis antigens.

[0021] 3. PRO232

[0022] Stem cells are undifferentiated cells capable of (a) proliferation, (b) self maintenance, (c) the production of a large number of differentiated functional progeny, (d) regeneration of tissue after injury and/or (e) a flexibility in the use of these options. Stem cells often express cell surface antigens which are capable of serving as cell specific markers that can be exploited to identify stem cells, thereby providing a means for identifying and isolating specific stem cell populations.



[0023] Having possession of different stem cell populations will allow for a number of important applications. For example, possessing a specific stem cell population will allow for the identification of growth factors and other proteins which are involved in their proliferation and differentiation. In addition, there may be as yet undiscovered proteins which are associated with (1) the early steps of dedication of the stem cell to a particular lineage, (2) prevention of such dedication, and (3) negative control of stem cell proliferation, all of which may be identified if one has possession of the stem cell population. Moreover, stem cells are important and ideal targets for gene therapy where the inserted genes promote the health of the individual into whom the stem cells are transplanted. Finally, stem cells may play important roles in transplantation of organs or tissues, for example liver regeneration and skin grafting.

[0024] Given the importance of stem cells in various different applications, efforts are currently being undertaken by both industry and academia to identify new, native stem cell antigen proteins so as to provide specific cell surface markers for identifying stem cell populations as well as for providing insight into the functional roles played by stem cell antigens in cell proliferation and differentiation. We herein describe the identification and characterization of novel polypeptides having homology to a stem cell antigen, wherein those polypeptides are herein designated as PRO232 polypeptides.

[0025] 4. PRO187

[0026] Growth factors are molecular signals or mediators that enhance cell growth or proliferation, alone or in concert, by binding to specific cell surface receptors. However, there are other cellular reactions than only growth upon expression to growth factors. As a result, growth factors are better characterized as multifunctional and potent cellular regulators. Their biological effects include proliferation, chemotaxis and stimulation of extracellular matrix production. Growth factors can have both stimulatory and inhibitory effects. For example, transforming growth factor (TGF- $\beta$ ) is highly pleiotropic and can stimulate proliferation in some cells, especially connective tissue, while being a potent inhibitor of proliferation in others, such as lymphocytes and epithelial cells.

[0027] The physiological effect of growth stimulation or inhibition by growth factors depends upon the state of development and differentiation of the target tissue. The mechanism of local cellular regulation by classical endocrine molecules involves comprehends autocrine (same cell), juxtacrine (neighbor cell), and paracrine (adjacent cells) pathways. Peptide growth factors are elements of a complex biological language, providing the basis for intercellular communication. They permit cells to convey information between each other, mediate interaction between cells and change gene expression. The effect of these multifunctional and pluripotent factors is dependent on the presence or absence of other peptides.

[0028] FGF-8 is a member of the fibroblast growth factors (FGFs) which are a family of heparin-binding, potent mitogens for both normal diploid fibroblasts and established cell lines, Gospodarowicz et al. (1984), *Proc. Natl. Acad. Sci. USA* 81:6963. The FGF family comprises acidic FGF (FGF-1), basic FGF (FGF-2), INT-2 (FGF-3), K-FGF/HST (FGF-4), FGF-5, FGF-6, KGF (FGF-7), AIGF (FGF-8) among

others. All FGFs have two conserved cysteine residues and share 30-50% sequence homology at the amino acid level. These factors are mitogenic for a wide variety of normal diploid mesoderm-derived and neural crest-derived cells, including granulosa cells, adrenal cortical cells, chondrocytes, myoblasts, corneal and vascular endothelial cells (bovine or human), vascular smooth muscle cells, lens, retina and prostatic epithelial cells, oligodendrocytes, astrocytes, chondrocytes, myoblasts and osteoblasts.

[0029] Fibroblast growth factors can also stimulate a large number of cell types in a non-mitogenic manner. These activities include promotion of cell migration into wound area (chemotaxis), initiation of new blood vessel formation (angiogenesis), modulation of nerve regeneration and survival (neurotrophism), modulation of endocrine functions, and stimulation or suppression of specific cellular protein expression, extracellular matrix production and cell survival. Baird & Bohlen, *Handbook of Exp. Pharmacol.* 95(1): 369-418, Springer, (1990). These properties provide a basis for using fibroblast growth factors in therapeutic approaches to accelerate wound healing, nerve repair, collateral blood vessel formation, and the like. For example, fibroblast growth factors have been suggested to minimize myocardium damage in heart disease and surgery (U.S. Pat. No. 4,378,347).

[0030] FGF-8, also known as androgen-induced growth factor (AIGF), is a 215 amino acid protein which shares 30-40% sequence homology with the other members of the FGF family. FGF-8 has been proposed to be under androgenic regulation and induction in the mouse mammary carcinoma cell line SC3. Tanaka et al., *Proc. Natl. Acad. Sci. USA* 89: 8928-8932 (1992); Sato et al., *J. Steroid Biochem. Molec. Biol.* 47: 91-98 (1993). As a result, FGF-8 may have a local role in the prostate, which is known to be an androgen-responsive organ. FGF-8 can also be oncogenic, as it displays transforming activity when transfected into NIH-3T3 fibroblasts. Kouhara et al., *Oncogene* 9 455-462 (1994). While FGF-8 has been detected in heart, brain, lung, kidney, testis, prostate and ovary, expression was also detected in the absence of exogenous androgens. Schmitt et al., *J. Steroid Biochem. Mol. Biol.* 57 (3-4): 173-78 (1996).

[0031] FGF-8 shares the property with several other FGFs of being expressed at a variety of stages of murine embryogenesis, which supports the theory that the various FGFs have multiple and perhaps coordinated roles in differentiation and embryogenesis. Moreover, FGF-8 has also been identified as a protooncogene that cooperates with Wnt-1 in the process of mammary tumorigenesis (Shackelford et al., *Proc. Natl. Acad. Sci. USA* 90, 740-744 (1993); Heikinheimo et al., *Mech. Dev.* 48: 129-138 (1994)).

[0032] In contrast to the other FGFs, FGF-8 exists as three protein isoforms, as a result of alternative splicing of the primary transcript. Tanaka et al., supra. Normal adult expression of FGF-8 is weak and confined to gonadal tissue, however northern blot analysis has indicated that FGF-8 mRNA is present from day 10 through day 12 of murine gestation, which suggests that FGF-8 is important to normal development. Heikinheimo et al., *Mech Dev.* 48(2): 129-38 (1994). Further in situ hybridization assays between day 8 and 16 of gestation indicated initial expression in the surface ectoderm of the first bronchial arches, the frontonasal process, the forebrain and the midbrain-hindbrain junction. At

days 10-12, FGF-8 was expressed in the surface ectoderm of the forelimb and hindlimb buds, the nasal pits and nasopharynx, the infundibulum and in the telencephalon, diencephalon and metencephalon. Expression continues in the developing hindlimbs through day 13 of gestation, but is undetectable thereafter. The results suggest that FGF-8 has a unique temporal and spatial pattern in embryogenesis and suggests a role for this growth factor in multiple regions of ectodermal differentiation in the post-gastrulation embryo.

[0033] We herein describe the identification of novel polypeptides having homology to FGF-8, wherein those polypeptides are herein designated PRO187 polypeptides.

[0034] 5. PRO265

[0035] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0036] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leucine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0037] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Other studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vouv. Rev. Fr. Hematol.* (Germany), 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome and Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats. Another protein of particular interest which has been reported to have leucine-rich repeats is the SLIT protein which has been reported to be useful in treating neurodegenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanistsakonakos, S. and Rothberg, J. M., WO9210518-A1 by Yale University. Other studies reporting on the biological functions of proteins having leucine-rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-1133 (October 1995) (kidney disease involvement); and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer

Research Foundation (decorin binding to transforming growth factor- $\beta$  involvement for treatment for cancer, wound healing and scarring). Also of particular interest is fibromodulin and its use to prevent or reduce dermal scarring. A study of fibromodulin is found in U.S. Pat. No. 5,654,270 to Ruoslahti, et al.

[0038] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions. Of particular interest are those proteins having leucine rich repeats and homology to known proteins having leucine rich repeats such as fibromodulin, the SLIT protein and platelet glycoprotein V. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound proteins having leucine rich repeats. We herein describe the identification and characterization of novel polypeptides having homology to fibromodulin, herein designated as PRO265 polypeptides.

[0039] 6. PRO219

[0040] Human matrilin-2 polypeptide is a member of the von Willebrand factor type A-like module superfamily. von Willebrand factor is a protein which plays an important role in the maintenance of hemostasis. More specifically, von Willebrand factor is a protein which is known to participate in platelet-vessel wall interactions at the site of vascular injury via its ability to interact and form a complex with Factor VIII. The absence of von Willebrand factor in the blood causes an abnormality with the blood platelets that prevents platelet adhesion to the vascular wall at the site of the vascular injury. The result is the propensity for bruising, nose bleeds, intestinal bleeding, and the like comprising von Willebrand's disease.

[0041] Given the physiological importance of the blood clotting factors, efforts are currently being undertaken by both industry and academia to identify new, native proteins which may be involved in the coagulation process. We herein describe the identification of a novel full-length polypeptide which possesses homology to the human matrilin-2 precursor polypeptide.

[0042] 7. PRO246

[0043] The cell surface protein HCAR is a membrane-bound protein that acts as a receptor for subgroup C of the adenoviruses and subgroup B of the coxsackieviruses. Thus, HCAR may provide a means for mediating viral infection of cells in that the presence of the HCAR receptor on the cellular surface provides a binding site for viral particles, thereby facilitating viral infection.

[0044] In light of the physiological importance of membrane-bound proteins and specifically those which serve as a cell surface receptor for viruses, efforts are currently being undertaken by both industry and academia to identify new, native membrane-bound receptor proteins. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor proteins. We herein describe a novel membrane-bound polypeptide (designated herein as PRO246) having homology to the cell surface protein HCAR and to various tumor antigens including A33 and carcinoembryonic antigen, wherein this polypeptide may be a novel cell surface virus receptor or tumor antigen.

**[0045]** 8. PRO228

**[0046]** There are a number of known seven transmembrane proteins and within this family is a group which includes CD97 and EMR1. CD97 is a seven-span transmembrane receptor which has a cellular ligand, CD55, DAF. Hamann, et al., *J. Exp. Med.* (U.S.), 184(3):1189 (1996). Additionally, CD97 has been reported as being a dedifferentiation marker in human thyroid carcinomas and as associated with inflammation. Aust, et al., *Cancer Res.* (U.S.), 57(9):1798 (1997); Gray, et al., *J. Immunol.* (U.S.), 157(12):5438 (1996). CD97 has also been reported as being related to the secretin receptor superfamily, but unlike known members of that family, CD97 and EMR1 have extended extracellular regions that possess several EGF domains at the N-terminus. Hamann, et al., *Genomics*, 32(1):144 (1996); Harmann, et al., *J. Immunol.*, 155(4):1942 (1995). EMR1 is further described in Lin, et al., *Genomics*, 41(3):301 (1997) and Baud, et al., *Genomics*, 26(2):334 (1995). While CD97 and EMR1 appear to be related to the secretin receptors, a known member of the secretin family of G protein-coupled receptors includes the alpha-latroxin receptor, latrophilin, which has been described as calcium independent and abundant among neuronal tissues. Lelianova, et al., *J. Biol. Chem.*, 272(34), 21504 (1997); Davletov, et al., *J. Biol. Chem.* (U.S.), 271(38):23239 (1996). Both members of the secretin receptor superfamily and non-members which are related to the secretin receptor superfamily, or CRF and calcitonin receptors are of interest. In particular, new members of these families, identified by their homology to known proteins, are of interest.

**[0047]** Efforts are being undertaken by both industry and academia to identify new membrane-bound receptor proteins, particularly transmembrane proteins with EGF repeats and large N-terminuses which may belong to the family of seven-transmembrane proteins of which CD97 and EMR1 are members. We herein describe the identification and characterization of novel polypeptides having homology to CD97 and EMR1, designated herein as PRO228 polypeptides.

**[0048]** 9. PRO533

**[0049]** Growth factors are molecular signals or mediators that enhance cell growth or proliferation, alone or in concert, by binding to specific cell surface receptors. However, there are other cellular reactions than only growth upon expression to growth factors. As a result, growth factors are better characterized as multifunctional and potent cellular regulators. Their biological effects include proliferation, chemotaxis and stimulation of extracellular matrix production. Growth factors can have both stimulatory and inhibitory effects. For example, transforming growth factors (TGF- $\beta$ ) is highly pleiotropic and can stimulate proliferation in some cells, especially connective tissues, while being a potent inhibitor of proliferation in others, such as lymphocytes and epithelial cells.

**[0050]** The physiological effect of growth stimulation or inhibition by growth factors depends upon the state of development and differentiation of the target tissue. The mechanism of local cellular regulation by classical endocrine molecules comprehends autocrine (same cell), juxtacrine (neighbor cell), and paracrine (adjacent cell) pathways. Peptide growth factors are elements of a complex biological language, providing the basis for intercellular communication.

They permit cells to convey information between each other, mediate interaction between cells and change gene expression. The effect of these multifunctional and pluripotent factors is dependent on the presence or absence of other peptides.

**[0051]** Fibroblast growth factors (FGFs) are a family of heparin-binding, potent mitogens for both normal diploid fibroblasts and established cell lines, Godpodarowicz, D. et al. (1984), *Proc. Natl. Acad. Sci. USA* 81: 6983. The FGF family comprises acidic FGF (FGF-1), basic FGF (FGF-2), INT-2 (FGF-3), K-FGF/HST (FGF-4), FGF-5, FGF-6, KGF (FGF-7), AIGF (FGF-8) among others. All FGFs have two conserved cysteine residues and share 30-50% sequence homology at the amino acid level. These factors are mitogenic for a wide variety of normal diploid mesoderm-derived and neural crest-derived cells, inducing granulosa cells, adrenal cortical cells, chondrocytes, myoblasts, corneal and vascular endothelial cells (bovine or human), vascular smooth muscle cells, lens, retina and prostatic epithelial cells, oligodendrocytes, astrocytes, chondrocytes, myoblasts and osteoblasts.

**[0052]** Fibroblast growth factors can also stimulate a large number of cell types in a non-mitogenic manner. These activities include promotion of cell migration into a wound area (chemotaxis), initiation of new blood vessel formation (angiogenesis), modulation of nerve regeneration and survival (neurotrophism), modulation of endocrine functions, and stimulation or suppression of specific cellular protein expression, extracellular matrix production and cell survival. Baird, A. & Bohlen, P., *Handbook of Exp. Pharmacol.* 95(1): 369-418 (1990). These properties provide a basis for using fibroblast growth factors in therapeutic approaches to accelerate wound healing, nerve repair, collateral blood vessel formation, and the like. For example, fibroblast growth factors, have been suggested to minimize myocardium damage in heart disease and surgery (U.S. Pat. No. 4,378,437).

**[0053]** We herein describe the identification and characterization of novel polypeptides having homology to FGF, herein designated PRO533 polypeptides.

**[0054]** 10. PRO245

**[0055]** Some of the most important proteins involved in the above described regulation and modulation of cellular processes are the enzymes which regulate levels of protein phosphorylation in the cell. For example, it is known that the transduction of signals that regulate cell growth and differentiation is regulated at least in part by phosphorylation and dephosphorylation of various cellular proteins. The enzymes that catalyze these processes include the protein kinases, which function to phosphorylate various cellular proteins, and the protein phosphatases, which function to remove phosphate residues from various cellular proteins. The balance of the level of protein phosphorylation in the cell is thus mediated by the relative activities of these two types of enzymes.

**[0056]** Although many protein kinase enzymes have been identified, the physiological role played by many of these catalytic proteins has yet to be elucidated. It is well known, however, that a number of the known protein kinases function to phosphorylate tyrosine residues in proteins, thereby leading to a variety of different effects. Perhaps most impor-

tantly, there has been a great deal of interest in the protein tyrosine kinases since the discovery that many oncogene products and growth factors possess intrinsic protein tyrosine kinase activity. There is, therefore, a desire to identify new members of the protein tyrosine kinase family.

[0057] Given the physiological importance of the protein kinases, efforts are being undertaken by both industry and academia to identify new, native kinase proteins. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel kinase proteins. We herein describe the identification and characterization of novel polypeptides having homology to tyrosine kinase proteins, designated herein as PRO245 polypeptides.

[0058] 11. PRO220, PRO221 and PRO227

[0059] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0060] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leucine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0061] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Others studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vouv. Rev. Fr. Hematol. (Germany)*, 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome and Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats. Another protein of particular interest which has been reported to have leucine-rich repeats is the SLIT protein which has been reported to be useful in treating neurodegenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanistsakonas, S. and Rothberg, J. M., WO9210518-A1 by Yale University. Other studies reporting on the biological functions of proteins having leucine-rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-

1133 (October 1995) (kidney disease involvement); and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer Research Foundation (decorin binding to transforming growth factor $\beta$  involvement for treatment for cancer, wound healing and scarring).

[0062] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions. Of particular interest are those proteins having leucine rich repeats and homology to known proteins having leucine rich repeats such as the SLIT protein and platelet glycoprotein V.

[0063] 12. PRO258

[0064] Immunoglobulins are antibody molecules, the proteins that function both as receptors for antigen on the B-cell membrane and as the secreted products of the plasma cell. Like all antibody molecules, immunoglobulins perform two major functions: they bind specifically to an antigen and they participate in a limited number of biological effector functions. Therefore, new members of the Ig superfamily are always of interest. Molecules which act as receptors by various viruses and those which act to regulate immune function are of particular interest. Also of particular interest are those molecules which have homology to known Ig family members which act as virus receptors or regulate immune function. Thus, molecules having homology to poliovirus receptors, CRTAM and CD166 (a ligand for lymphocyte antigen CD6) are of particular interest.

[0065] Extracellular and membrane-bound proteins play important roles in the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuro-peptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment, usually at a membrane-bound receptor protein.

[0066] We herein describe the identification and characterization of novel polypeptides having homology to CRTAM, designated herein as PRO258 polypeptides.

[0067] 13. PRO266

[0068] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0069] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leucine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor

protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0070] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Others studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vouv. Rev. Fr. Hematol.* (Germany), 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome and Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats. Another protein of particular interest which has been reported to have leucine-rich repeats is the SLIT protein which has been reported to be useful in treating neurodegenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanistsakonias, S. and Rothberg, J. M., WO9210518-A1 by Yale University. Other studies reporting on the biological functions of proteins having leucine-rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-1133 (October 1995) (kidney disease involvement); and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer Research Foundation (decorin binding to transforming growth factors involvement for treatment for cancer, wound healing and scarring).

[0071] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions, neuronal development and adhesion molecules. Of particular interest are those proteins having leucine rich repeats and homology to known proteins having leucine rich repeats such as the SLIT protein. We herein describe novel polypeptides having homology to SLIT, designated herein as PRO266 polypeptides.

[0072] 14. PRO269

[0073] Thrombomodulin binds to and regulates the activity of thrombin. It is important in the control of blood coagulation. Thrombomodulin functions as a natural anticoagulant by accelerating the activation of protein C by thrombin. Soluble thrombomodulin may have therapeutic use as an antithrombotic agent with reduced risk for hemorrhage as compared with heparin. Thrombomodulin is a cell surface trans-membrane glycoprotein, present on endothelial cells and platelets. A smaller, functionally active form of thrombomodulin circulates in the plasma and is also found in urine. (In Haeberli, A., Human Protein Data, VCH Oub., N.Y., 1992). Peptides having homology to thrombomodulin are particularly desirable.

[0074] We herein describe the identification and characterization of novel polypeptides having homology to thrombomodulin, designated herein as PRO269 polypeptides.

[0075] 15. PRO287

[0076] Procollagen C-proteinase enhancer protein binds to and enhances the activity of bone morphogenic protein "BMP1"/procollagen C-proteinase (PCP). It plays a role in extracellular matrix deposition. BMP1 proteins may be used to induce bone and/or cartilage formation and in wound healing and tissue repair. Therefore, procollagen C-proteinase enhancer protein, BMP1 and proteins having homology thereto, are of interest to the scientific and medical communities.

[0077] We herein describe the identification and characterization of novel polypeptides having homology to procollagen C-proteinase enhancer protein precursor and procollagen C-proteinase enhancer protein, designated herein as PRO287 polypeptides.

[0078] 16. PRO214

[0079] Growth factors are molecular signals or mediators that enhances cell growth or proliferation, alone or in concert, by binding to specific cell surface receptors. However, there are other cellular reactions than only growth upon expression to growth factors. As a result, growth factors are better characterized as multifunctional and potent cellular regulators. Their biological effects include proliferation, chemotaxis and stimulation of extracellular matrix production. Growth factors can have both stimulatory and inhibitory effects. For example, transforming growth factor  $\beta$  (TGF- $\beta$ ) is highly pleiotropic and can stimulate proliferation in some cells, especially connective tissue, while being a potent inhibitor of proliferation in others, such as lymphocytes and epithelial cells.

[0080] The physiological effect of growth stimulation or inhibition by growth factors depends upon the state of development and differentiation of the target tissue. The mechanism of local cellular regulation by classical endocrine molecules involves comprehends autocrine (same cell), juxtacrine (neighbor cell), and paracrine (adjacent cells) pathways. Peptide growth factors are elements of a complex biological language, providing the basis for intercellular communication. They permit cells to convey information between each other, mediate interaction between cells and change gene expression. The effect of these multifunctional and pluripotent factors is dependent on the presence or absence of other peptides.

[0081] Epidermal growth factor (EGF) is a conventional mitogenic factor that stimulates the proliferation of various types of cells including epithelial cells and fibroblasts. EGF binds to and activates the EGF receptor (EGFR), which initiates intracellular signaling and subsequent effects. The EGFR is expressed in neurons of the cerebral cortex, cerebellum, and hippocampus in addition to other regions of the central nervous system (CNS). In addition, EGF is also expressed in various regions of the CNS. Therefore, EGF acts not only on mitotic cells, but also on postmitotic neurons. In fact, many studies have indicated that EGF has neurotrophic or neuromodulatory effects on various types of neurons in the CNS. For example, EGF acts directly on cultured cerebral cortical and cerebellar neurons, enhancing neurite outgrowth and survival. On the other hand, EGF also

acts on other cell types, including septal cholinergic and mesencephalic dopaminergic neurons, indirectly through glial cells. Evidence of the effects of EGF on neurons in the CNS is accumulating, but the mechanisms of action remain essentially unknown. EGF-induced signaling in mitotic cells is better understood than in postmitotic neurons. Studies of cloned pheochromocytoma PC12 cells and cultured cerebral cortical neurons have suggested that the EGF-induced neurotrophic actions are mediated by sustained activation of the EGFR and mitogen-activated protein kinase (MAPK) in response to EGF. The sustained intracellular signaling correlates with the decreased rate of EGFR down-regulation, which might determine the response of neuronal cells to EGF. It is likely that EGF is a multi-potent growth factor that acts upon various types of cells including mitotic cells and postmitotic neurons.

**[0082]** EGF is produced by the salivary and Brunner's glands of the gastrointestinal system, kidney, pancreas, thyroid gland, pituitary gland, and the nervous system, and is found in body fluids such as saliva, blood, cerebrospinal fluid (CSF), urine, amniotic fluid, prostatic fluid, pancreatic juice, and breast milk, Plata-Salaman, CR *Peptides* 12: 653-663 (1991).

**[0083]** EGF is mediated by its membrane specific receptor, which contains an intrinsic tyrosine kinase. Stoscheck C M et al., *J. Cell Biochem.* 31: 135-152 (1986). EGF is believed to function by binding to the extracellular portion of its receptor which induces a transmembrane signal that activates the intrinsic tyrosine kinase.

**[0084]** Purification and sequence analysis of the EGF-like domain has revealed the presence of six conserved cysteine residues which cross-bind to create three peptide loops, Savage C R et al., *J. Biol. Chem.* 248: 7669-7672 (1979). It is now generally known that several other peptides can react with the EGF receptor which share the same generalized motif  $X_nCX_7CX_{4/5}CX_{10}CXCX_5GX_2CX_n$ , where X represents any non-cysteine amino acid, and n is a variable repeat number. Non isolated peptides having this motif include TGF- $\alpha$ , amphiregulin, schwannoma-derived growth factor (SDGF), heparin-binding EGF-like growth factors and certain virally encoded peptides (e.g., Vaccinia virus, Reisner A H, *Nature* 313: 801-803 (1985), Shope fibroma virus, Chang W., et al., *Mol Cell Biol.* 7: 535-540 (1987), Molluscum contagiosum, Porter C D & Archard L C, *J. Gen. Virol.* 68: 673-682 (1987), and Myxoma virus, Upton C et al., *J. Virol.* 61: 1271-1275 (1987). Prigent S A & Lemoine N. R., *Prog. Growth Factor Res.* 4: 1-24 (1992).

**[0085]** EGF-like domains are not confined to growth factors but have been observed in a variety of cell-surface and extracellular proteins which have interesting properties in cell adhesion, protein-protein interaction and development, Laurence D J R & Gusterson B A, *Tumor Biol.* 11: 229-261 (1990). These proteins include blood coagulation factors (factors VI, IX, X, XII, protein C, protein S, protein Z, tissue plasminogen activator, urokinase), extracellular matrix components (laminin, cytotoactin, entactin), cell surface receptors (LDL receptor, thrombomodulin receptor) and immunity-related proteins (complement C1r, uromodulin).

**[0086]** Even more interesting, the general structure pattern of EGF-like precursors is preserved through lower organisms as well as in mammalian cells. A number of genes with developmental significance have been identified in invertebrates with EGF-like repeats. For example, the notch gene of *Drosophila* encodes 36 tandemly arranged 40 amino acid repeats which show homology to EGF, Wharton W et al., *Cell* 43: 557-581 (1985). Hydropathy plots indicate a putative membrane spanning domain, with the EGF-related sequences being located on the extracellular side of the membrane. Other homeotic genes with EGF-like repeats include Delta, 95F and 5ZD which were identified using probes based on Notch, and the nematode gene Lin-12 which encodes a putative receptor for a developmental signal transmitted between two specified cells.

**[0087]** Specifically, EGF has been shown to have potential in the preservation and maintenance of gastrointestinal mucosa and the repair of acute and chronic mucosal lesions, Konturek, P C et al., *Eur. J. Gastroenterol Hepatol.* 7 (10), 933-37 (1995), including the treatment of necrotizing enterocolitis, Zollinger-Ellison syndrome, gastrointestinal ulceration, congenital microvillus atrophy, A. Guglietta & P B Sullivan, *Eur. J. Gastroenterol Hepatol.* 7(10), 945-50 (1995). Additionally, EGF has been implicated in hair follicle differentiation; C. L. du Cros, *J. Invest. Dermatol.* 101 (1 Suppl.), 106S-113S (1993), S G Hillier, *Clin. Endocrinol.* 33(4), 427-28 (1990); kidney function, L. L. Hamm et al., *Semin. Nephrol.* 13 (1): 109-15 (1993), R C Harris, *Am. J. Kidney Dis.* 17(6): 627-30 (1991); tear fluid, G B van Setten et al., *Int. Ophthalmol* 15(6): 359-62 (1991); vitamin K mediated blood coagulation, J. Stenflo et al., *Blood* 78(7): 1637-51 (1991). EGF is also implicated various skin disease characterized by abnormal keratinocyte differentiation, e.g., psoriasis, epithelial cancers such as squamous cell carcinomas of the lung, epidermoid carcinoma of the vulva and gliomas. King, L E et al., *Am. J. Med. Sci.* 296: 154-158 (1988).

**[0088]** Of great interest is mounting evidence that genetic alterations in growth factors signaling pathways are closely linked to developmental abnormalities and to chronic diseases including cancer. Aaronson S A, *Science* 254: 1146-1153 (1991). For example, c-erb-2 (also known as HER-2), a proto-oncogene with close structural similarity to EGF receptor protein, is overexpressed in human breast cancer. King et al., *Science* 229: 974-976 (1985); Gullick, W J, *Hormones and their actions*, Cooke B A et al., eds, Amsterdam, Elsevier, pp 349-360 (1986).

**[0089]** 17. PRO317

**[0090]** The TGF- $\beta$  supergene family, or simply TGF- $\beta$  superfamily, a group of secreted proteins, includes a large number of related growth and differentiation factors expressed in virtually all phyla. Superfamily members bind to specific cell surface receptors that activate signal transduction mechanisms to elicit their multifunctional cytokine effects. Kolodziejczyk and Hall, *Biochem. Cell. Biol.*, 74: 299-314 (1996); Attisano and Wrana, *Cytokine Growth Factor Rev.*, 7: 327-339 (1996); and Hill, *Cellular Signaling*, 8: 533-544 (1996).

**[0091]** Members of this family include five distinct forms of TGF- $\beta$  (Sporn and Roberts, in *Peptide Growth Factors and Their Receptors*, Sporn and Roberts, eds. (Springer-Verlag: Berlin, 1990) pp. 419-472), as well as the differentiation factors vg1 (Weeks and Melton, *Cell*, 51: 861-867 (1987)) and DPP-C polypeptide (Padgett et al., *Nature*, 325: 81-84(1987)), the hormones activin and inhibin (Mason et al., *Nature*, 318: 659-663 (1985); Mason et al., *Growth*

*Factors*, 1: 77-88 (1987)), the Mullerian-inhibiting substance (MIS) (Cate et al., *Cell*, 45: 685-698 (1986)), the bone morphogenetic proteins (BMPs) (Wozney et al., *Science*, 242: 1528-1534 (1988); PCT WO 88/00205 published Jan. 14, 1988; U.S. Pat. No. 4,877,864 issued Oct. 31, 1989), the developmentally regulated proteins Vgr-1 (Lyons et al., *Proc. Natl. Acad. Sci. USA*, 86: 4554-4558 (1989)) and Vgr-2 (Jones et al., *Molec. Endocrinol.*, 6: 1961-1968 (1992)), the mouse growth differentiation factor (GDF), such as GDF-3 and GDF-9 (Kingsley, *Genes Dev.*, 8: 133-146 (1994); McPherron and Lee, *J. Biol. Chem.*, 268: 3444-3449 (1993)), the mouse lefty/Stra1 (Meno et al., *Nature*, 381: 151-155 (1996); Bouillet et al., *Dev. Biol.*, 170: 420-433 (1995)), glial cell line-derived neurotrophic factor (GDNF) (Lin et al., *Science*, 260: 1130-1132 (1993)), neurturin (Kotzbauer et al., *Nature*, 384: 467-470 (1996)), and endometrial bleeding-associated factor (EBAF) (Kothapalli et al., *J. Clin. Invest.*, 99: 2342-2350 (1997)). The subset BMP-2A and BMP-2B is approximately 75% homologous in sequence to DPP-C and may represent the mammalian equivalent of that protein.

[0092] The proteins of the TGF- $\beta$  superfamily are disulfide-linked homo- or heterodimers encoded by larger precursor polypeptide chains containing a hydrophobic signal sequence, a long and relatively poorly conserved N-terminal pro region of several hundred amino acids, a cleavage site (usually polybasic), and a shorter and more highly conserved C-terminal region. This C-terminal region corresponds to the processed mature protein and contains approximately 100 amino acids with a characteristic cysteine motif, i.e., the conservation of seven of the nine cysteine residues of TGF- $\beta$  among all known family members. Although the position of the cleavage site between the mature and pro regions varies among the family members, the C-terminus of all of the proteins is in the identical position, ending in the sequence Cys-X-Cys-X, but differing in every case from the TGF- $\beta$  consensus C-terminus of Cys-Lys-Cys-Ser. Sporn and Roberts, 1990, supra.

[0093] There are at least five forms of TGF- $\beta$  currently identified, TGF- $\beta$ 1, TGF- $\beta$ 2, TGF- $\beta$ 3, TGF- $\beta$ 4, and TGF- $\beta$ 5. The activated form of TGF- $\beta$ 1 is a homodimer formed by dimerization of the carboxy-terminal 112 amino acids of a 390 amino acid precursor. Recombinant TGF- $\beta$ 1 has been cloned (Derynck et al., *Nature*, 316:701-705 (1985)) and expressed in Chinese hamster ovary cells (Gentry et al., *Mol. Cell. Biol.*, 7: 3418-3427 (1987)). Additionally, recombinant human TGF- $\beta$ 2 (deMartin et al., *EMBO J.*, 6: 3673 (1987)), as well as human and porcine TGF- $\beta$ 3 (Derynck et al., *EMBO J.*, 7: 3737-3743 (1988); ten Dijke et al., *Proc. Natl. Acad. Sci. USA*, 85: 4715 (1988)) have been cloned. TGF- $\beta$ 2 has a precursor form of 414 amino acids and is also processed to a homodimer from the carboxy-terminal 112 amino acids that shares approximately 70% homology with the active form of TGF- $\beta$ 1 (Marquardt et al., *J. Biol. Chem.*, 262: 12127 (1987)). See also EP 200,341; 169,016; 268,561; and 267,463; U.S. Pat. No. 4,774,322; Cheifetz et al., *Cell*, 48: 409-415 (1987); Jakowlew et al., *Molecular Endocrinol.*, 2: 747-755 (1988); Derynck et al., *J. Biol. Chem.*, 261: 4377-4379 (1986); Sharples et al., *DNA*, 6: 239-244 (1987); Derynck et al., *Nucl. Acids. Res.*, 15: 3188-3189 (1987); Derynck et al., *Nucl. Acids. Res.*, 15: 3187 (1987); Seyedin et al., *J. Biol. Chem.*, 261: 5693-5695 (1986); Madisen et al., *DNA*, 7: 1-8 (1988); and Hanks et al., *Proc. Natl. Acad. Sci. (U.S.A.)*, 85: 79-82 (1988).

[0094] TGF- $\beta$ 4 and TGF- $\beta$ 5 were cloned from a chicken chondrocyte cDNA library (Jakowlew et al., *Molec. Endocrinol.*, 2: 1186-1195 (1988)) and from a frog oocyte cDNA library, respectively.

[0095] The pro region of TGF- $\beta$  associates non-covalently with the mature TGF- $\beta$  dimer (Wakefield et al., *J. Biol. Chem.*, 263: 7646-7654 (1988); Wakefield et al., *Growth Factors*, 1: 203-218 (1989)), and the pro regions are found to be necessary for proper folding and secretion of the active mature dimers of both TGF- $\beta$  and activin (Gray and Mason, *Science*, 247: 1328-1330 (1990)). The association between the mature and pro regions of TGF- $\beta$  masks the biological activity of the mature dimer, resulting in formation of an inactive latent form. Latency is not a constant of the TGF- $\beta$  superfamily, since the presence of the pro region has no effect on activin or inhibin biological activity.

[0096] A unifying feature of the biology of the proteins from the TGF- $\beta$  superfamily is their ability to regulate developmental processes. TGF- $\beta$  has been shown to have numerous regulatory actions on a wide variety of both normal and neoplastic cells. TGF- $\beta$  is multifunctional, as it can either stimulate or inhibit cell proliferation, differentiation, and other critical processes in cell function (Sporn and Roberts, supra).

[0097] One member of the TGF- $\beta$  superfamily, EBAF, is expressed in endometrium only in the late secretory phase and during abnormal endometrial bleeding. Kothapalli et al., *J. Clin. Invest.*, 99: 2342-2350 (1997). Human endometrium is unique in that it is the only tissue in the body that bleeds at regular intervals. In addition, abnormal endometrial bleeding is one of the most common manifestations of gynecological diseases, and is a prime indication for hysterectomy. In situ hybridization showed that the mRNA of EBAF was expressed in the stroma without any significant mRNA expression in the endometrial glands or endothelial cells.

[0098] The predicted protein sequence of EBAF showed a strong homology to the protein encoded by mouse lefty/stra3 of the TGF- $\beta$  superfamily. A motif search revealed that the predicted EBAF protein contains most of the cysteine residues which are conserved among the TGF- $\beta$ -related proteins and which are necessary for the formation of the cysteine knot structure. The EBAF sequence contains an additional cysteine residue, 12 amino acids upstream from the first conserved cysteine residue. The only other family members known to contain an additional cysteine residue are TGF- $\beta$ s, inhibins, and GDF-3. EBAF, similar to LEFTY, GDF-3/Vgr2, and GDF-9, lacks the cysteine residue that is known to form the intermolecular disulfide bond. Therefore, EBAF appears to be an additional member of the TGF- $\beta$  superfamily with an unpaired cysteine residue that may not exist as a dimer. However, hydrophobic contacts between the two monomer subunits may promote dimer formation. Fluorescence in situ hybridization showed that the ebafe gene is located on human chromosome 1 at band q42.1.

[0099] Additional members of the TGF- $\beta$  superfamily, such as those related to EBAF, are being searched for by industry and academics. We herein describe the identification and characterization of novel polypeptides having homology to EBAF, designated herein as PRO317 polypeptides.

**[0100]** 18. PRO301

**[0101]** The widespread occurrence of cancer has prompted the devotion of considerable resources and discovering new treatments of treatment. One particular method involves the creation of tumor or cancer specific monoclonal antibodies (mAbs) which are specific to tumor antigens. Such mAbs, which can distinguish between normal and cancerous cells are useful in the diagnosis, prognosis and treatment of the disease. Particular antigens are known to be associated with neoplastic diseases, such as colorectal cancer.

**[0102]** One particular antigen, the A33 antigen is expressed in more than 90% of primary or metastatic colon cancers as well as normal colon epithelium. Since colon cancer is a widespread disease, early diagnosis and treatment is an important medical goal. Diagnosis and treatment of colon cancer can be implemented using monoclonal antibodies (mAbs) specific therefore having fluorescent, nuclear magnetic or radioactive tags. Radioactive gene, toxins and/or drug tagged mAbs can be used for treatment in situ with minimal patient description. mAbs can also be used to diagnose during the diagnosis and treatment of colon cancers. For example, when the serum levels of the A33 antigen are elevated in a patient, a drop of the levels after surgery would indicate the tumor resection was successful. On the other hand, a subsequent rise in serum A33 antigen levels after surgery would indicate that metastases of the original tumor may have formed or that new primary tumors may have appeared. Such monoclonal antibodies can be used in lieu of, or in conjunction with surgery and/or other chemotherapies. For example, U.S. Pat. No. 4,579,827 and U.S. Ser. No. 424,991 (E.P. 199,141) are directed to therapeutic administration of monoclonal antibodies, the latter of which relates to the application of anti-A33 mAb.

**[0103]** Many cancers of epithelial origin have adenovirus receptors. In fact, adenovirus-derived vectors have been proposed as a means of inserting antisense nucleic acids into tumors (U.S. Pat. No. 5,518,885). Thus, the association of viral receptors with neoplastic tumors is not unexpected.

**[0104]** We herein describe the identification and characterization of novel polypeptides having homology to certain cancer-associated antigens, designated herein as PRO301 polypeptides.

**[0105]** 19. PRO224

**[0106]** Cholesterol uptake can have serious implications on one's health. Cholesterol uptake provides cells with most of the cholesterol they require for membrane synthesis. If this uptake is blocked, cholesterol accumulates in the blood and can contribute to the formation of atherosclerotic plaques in blood vessel walls. Most cholesterol is transported in the blood bound to protein in the form of complexes known as low-density lipoproteins (LDLs). LDLs are endocytosed into cells via LDL receptor proteins. Therefore, LDL receptor proteins, and proteins having homology thereto, are of interest to the scientific and medical communities.

**[0107]** Membrane-bound proteins and receptors can play an important role in the formation, differentiation and maintenance of multicellular organisms. The LDL receptors are an example of membrane-bound proteins which are involved in the synthesis and formation of cell membranes, wherein the health of an individual is affected directly and indirectly by its function. Many membrane-bound proteins act as

receptors such as the LDL receptor. These receptors can function to endocytose substrates or they can function as a receptor for a channel. Other membrane-bound proteins function as signals or antigens.

**[0108]** Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule regulators of the relevant receptor/ligand interaction. In the case of the LDL receptor, it is desirable to find molecules which enhance endocytosis so as to lower blood cholesterol levels and plaque formation. It is also desirable to identify molecules which inhibit endocytosis so that these molecules can be avoided or regulated by individuals having high blood cholesterol. Polypeptides which are homologous to lipoprotein receptors but which do not function as lipoprotein receptors are also of interest in the determination of the function of the fragments which show homology.

**[0109]** The following studies report on previously known low density lipoprotein receptors and related proteins including apolipoproteins: Sawamura, et al., Nippon Chemphar Co, Japan patent application J09098787; Novak, S., et al., *J. Biol. Chem.*, 271:(20)11732-6 (1996); Blaas, D., *J. Virol.*, 69(11)7244-7 (November 1995); Scott, J., *J. Inherit. Metab. Dis.* (UK), 9/Supp. 1 (3-16) (1986); Yamamoto, et al., *Cell*, 39:27-38 (1984); Rebecce, et al., *Neurobiol. Aging*, 15:5117 (1994); Novak, S., et al., *J. Biol. Chemistry*, 271:11732-11736(1996); and Sestavel and Fruchart, *Cell Mol. Biol.*, 40(4):461-81 (June 1994). These publications and others published prior to the filing of this application provide further background to peptides already known in the art.

**[0110]** Efforts are being undertaken by both industry and academia to identify new, native membrane-bound receptor proteins, particularly those having homology to lipoprotein receptors. We herein describe the identification and characterization of novel polypeptides having homology to lipoprotein receptors, designated herein as PRO224 polypeptides.

**[0111]** 20. PRO222

**[0112]** Complement is a group of proteins found in the blood that are important in humoral immunity and inflammation. Complement proteins are sequentially activated by antigen-antibody complexes or by proteolytic enzymes. When activated, complement proteins kill bacteria and other microorganisms, affect vascular permeability, release histamine and attract white blood cells. Complement also enhances phagocytosis when bound to target cells. In order to prevent harm to autologous cells, the complement activation pathway is tightly regulated.

**[0113]** Deficiencies in the regulation of complement activation or in the complement proteins themselves may lead to immune-complex diseases, such as systemic lupus erythematosus, and may result in increased susceptibility to bacterial infection. In all cases, early detection of complement deficiency is desirable so that the patient can begin treatment. Thus, research efforts are currently directed toward identification of soluble and membrane proteins that regulate complement activation.

**[0114]** Proteins known to be important in regulating complement activation in humans include Factor H and



Complement receptor type 1 (CR1). Factor H is a 150 kD soluble serum protein that interacts with complement protein C3b to accelerate the decay of C3 convertase and acts as a cofactor for Factor I-mediated cleavage of complement protein C4b. Complement receptor type 1 is a 190-280 kD membrane bound protein found in mast cells and most blood cells. CR1 interacts with complement proteins C3b, C4b, and iC3b to accelerate dissociation of C3 convertases, acts as a cofactor for Factor I-mediated cleavage of C3b and C4b, and binds immune complexes and promotes their dissolution and phagocytosis.

**[0115]** Proteins which have homology to complement proteins are of particular interest to the medical and industrial communities. Often, proteins having homology to each other have similar function. It is also of interest when proteins having homology do not have similar functions, indicating that certain structural motifs identify information other than function, such as locality of function.

**[0116]** Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound proteins, particularly those having homology to known proteins involved in the complement pathway. Proteins involved in the complement pathway were reviewed in Birmingham D J (1995), *Critical Reviews in Immunology*, 15(2):133-154 and in Abbas A K, et al. (1994) *Cellular and Molecular Immunology*, 2nd Ed. W. B. Saunders Company, Philadelphia, pp 295-315.

**[0117]** We herein describe the identification and characterization of novel polypeptides having homology to complement receptors, designated herein as PRO222 polypeptides.

**[0118]** 21. PRO234

**[0119]** The successful function of many systems within multicellular organisms is dependent on cell-cell interactions. Such interactions are affected by the alignment of particular ligands with particular receptors in a manner which allows for ligand-receptor binding and thus a cell-cell adhesion. While protein-protein interactions in cell recognition have been recognized for some time, only recently has the role of carbohydrates in physiologically relevant recognition been widely considered (see B. K. Brandley et al., *J. Leuk. Biol.* 40: 97 (1986) and N. Sharon et al., *Science* 246: 227 (1989). Oligosaccharides are well positioned to act as recognition novel lectins due to their cell surface location and structural diversity. Many oligosaccharide structures can be created through the differential activities of a smaller number of glycosyltransferases. The diverse structures of oligosaccharides can be generated by transcription of relatively few gene products, which suggests that the oligosaccharides are a plausible mechanism by which is directed a wide range of cell-cell interactions. Examples of differential expression of cell surface carbohydrates and putative carbohydrate binding proteins (lectins) on interacting cells have been described (J. Dodd & T. M. Jessel, *J. Neurosci.* 5: 3278 (1985); L. J. Regan et al., *Proc. Natl. Acad. Sci. USA* 83: 2248 (1986); M. Constantine-Paton et al., *Nature* 324: 459 (1986); and M. Tiemeyer et al., *J. Biol. Chem.* 263: 1671 (1989). One interesting member of the lectin family are selectins.

**[0120]** The migration of leukocytes to sites of acute or chronic inflammation involves adhesive interactions

between these cells and the endothelium. This specific adhesion is the initial event in the cascade that is initiated by inflammatory insults, and it is, therefore, of paramount importance to the regulated defense of the organism.

**[0121]** The types of cell adhesion molecules that are involved in the interaction between leukocytes and the endothelium during an inflammatory response currently stands at four: (1) selectins; (2) (carbohydrate and glycoprotein) ligands for selectins; (3) integrins; and (4) integrin ligands, which are members of the immunoglobulin gene superfamily.

**[0122]** The selectins are cell adhesion molecules that are unified both structurally and functionally. Structurally, selectins are characterized by the inclusion of a domain with homology to a calcium-dependent lectin (C-lectins), an epidermal growth factor (egf)-like domain and several complement binding-like domains, Bevilacqua, M. P. et al., *Science* 243: 1160-1165 (1989); Johnston et al., *Cell* 56: 1033-1044 (1989); Lasky et al., *Cell* 56: 1045-1055 (1989); Siegalman, M. et al., *Science* 243: 1165-1172 (1989); Stoolman, L. M., *Cell* 56: 907-910 (1989). Functionally, selectins share the common property of their ability to mediate cell binding through interactions between their lectin domains and cell surface carbohydrate ligands (Brandley, B, et al., *Cell* 63, 861-863 (1990); Springer, T. and Lasky, L. A., *Nature* 349, 19-197 (1991); Bevilacqua, M. P. and Nelson, R. M., *J. Clin. Invest.* 91 379-387 (1993) and Tedder et al., *J. Exp. Med.* 170: 123-133 (1989).

**[0123]** There are three members identified so far in the selectin family of cell adhesion molecules: L-selectin (also called peripheral lymph node homing receptor (pNHR), LEC-CAM-1, LAM-1, gp90<sup>MEL</sup>, gp100<sup>MEL</sup>, gp110<sup>MEL</sup>, MEL-14 antigen, Leu-8 antigen, TQ-1 antigen, DREG antigen), E-selectin (LEC-CAM-2, LECAM-2, ELAM-1) and P-selectin (LEC-CAM-3, LECAM-3, GMP-140, PAD-GEM).

**[0124]** The identification of the C-lectin domain has led to an intense effort to define carbohydrate binding ligands for proteins containing such domains. E-selectin is believed to recognize the carbohydrate sequence NeuNAc $\alpha$ 2-3Gal $\beta$ 1-4(Fuca $\alpha$ 1-3)GlcNAc (sialyl-Lewis x, or sLe<sup>x</sup>) and related oligosaccharides, Berg et al., *J. Biol. Chem.* 265: 14869-14872 (1991); Lowe et al., *Cell* 63: 475-484 (1990); Phillips et al., *Science* 250: 1130-1132 (1990); Tiemeyer et al., *Proc. Natl. Acad. Sci. USA* 88: 1138-1142 (1991).

**[0125]** L-selectin, which comprises a lectin domain, performs its adhesive function by recognizing carbohydrate-containing ligands on endothelial cells. L-selectin is expressed on the surface of leukocytes, such as lymphocytes, neutrophils, monocytes and eosinophils, and is involved with the trafficking of lymphocytes to peripheral lymphoid tissues (Gallatin et al., *Nature* 303: 30-34 (1983)) and with acute neutrophil-mediated inflammatory responses (Watson, S. R., *Nature* 349: 164-167 (1991)). The amino acid sequence of L-selectin and the encoding nucleic acid sequence are, for example, disclosed in U.S. Pat. No. 5,098,833 issued Mar. 24, 1992.

**[0126]** L-selectin (LECAM-1) is particularly interesting because of its ability to block neutrophil influx (Watson et al., *Nature* 349: 164-167 (1991)). It is expressed in chronic lymphocytic leukemia cells which bind to HEV (Sperini et

al., *Nature* 349: 691-694 (1991). It is also believed that HEV structures at sites of chronic inflammation are associated with the symptoms of diseases such as rheumatoid arthritis, psoriasis and multiple sclerosis.

[0127] E-selectin (ELAM-1), is particularly interesting because of its transient expression on endothelial cells in response to IL-1 or TNF. Bevilacqua et al., *Science* 243: 1160 (1989). The time course of this induced expression (2-8 h) suggests a role for this receptor in initial neutrophil induced extravasation in response to infection and injury. It has further been reported that anti-ELAM-1 antibody blocks the influx of neutrophils in a primate asthma model and thus is beneficial for preventing airway obstruction resulting from the inflammatory response. Gundel et al., *J. Clin. Invest.* 88: 1407 (1991).

[0128] The adhesion of circulating neutrophils to stimulated vascular endothelium is a primary event of the inflammatory response. P-selectin has been reported to recognize the Lewis x structure (Gal $\beta$ 1-4(Fuc $\alpha$ 1-3) GlcNAc), Larsen et al., *Cell* 63: 467-474(1990). Others report that an additional terminal linked sialic acid is required for high affinity binding, Moore et al., *J. Cell. Biol.* 112: 491-499 (1991). P-selectin has been shown to be significant in acute lung injury. Anti-P-selectin antibody has been shown to have strong protective effects in a rodent lung injury model. M. S. Mulligan et al., *J. Clin. Invest.* 90: 1600 (1991).

[0129] We herein describe the identification and characterization of novel polypeptides having homology to lectin proteins, herein designated as PRO234 polypeptides.

[0130] 22. PRO231

[0131] Some of the most important proteins involved in the above described regulation and modulation of cellular processes are the enzymes which regulate levels of protein phosphorylation in the cell. For example, it is known that the transduction of signals that regulate cell growth and differentiation is regulated at least in part by phosphorylation and dephosphorylation of various cellular proteins. The enzymes that catalyze these processes include the protein kinases, which function to phosphorylate various cellular proteins, and the protein phosphatases, which function to remove phosphate residues from various cellular proteins. The balance of the level of protein phosphorylation in the cell is thus mediated by the relative activities of these two types of enzymes.

[0132] Protein phosphatases represent a growing family of enzymes that are found in many diverse forms, including both membrane-bound and soluble forms. While many protein phosphatases have been described, the functions of only a very few are beginning to be understood (Tonks, *Semin. Cell Biol.* 4:373-453 (1993) and Dixon, *Recent Prog. Horm. Res.* 51:405-414 (1996)). However, in general, it appears that many of the protein phosphatases function to modulate the positive or negative signals induced by various protein kinases. Therefore, it is likely that protein phosphatases play critical roles in numerous and diverse cellular processes.

[0133] Given the physiological importance of the protein phosphatases, efforts are being undertaken by both industry and academia to identify new, native phosphatase proteins. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel phosphatase proteins. Examples of

screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0134] We herein describe the identification and characterization of novel polypeptides having homology to acid phosphatases, designated herein as PRO231 polypeptides.

[0135] 23. PRO229

[0136] Scavenger receptors are known to protect IgG molecules from catabolic degradation. Riechmann and Hollinger, *Nature Biotechnology*, 15:617 (1997). In particular, studies of the CH2 and CH3 domains have shown that specific sequences of these domains are important in determining the half-lives of antibodies. Ellerson, et al., *J. Immunol.*, 116: 510 (1976); Yasmeeen, et al., *J. Immunol.* 116: 518 (1976; Pollock, et al., *Eur. J. Immunol.*, 20: 2021 (1990). Scavenger receptor proteins and antibodies thereto are further reported in U.S. Pat. No. 5,510,466 to Krieger, et al. Due to the ability of scavenger receptors to increase the half-life of polypeptides and their involvement in immune function, molecules having homology to scavenger receptors are of importance to the scientific and medical community.

[0137] Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins, particularly those having homology to scavenger receptors. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0138] We herein describe the identification and characterization of novel polypeptides having homology to scavenger receptors, designated herein as PRO229 polypeptides.

[0139] 24. PRO238

[0140] Oxygen free radicals and antioxidants appear to play an important role in the central nervous system after cerebral ischemia and reperfusion. Moreover, cardiac injury, related to ischaemia and reperfusion has been reported to be caused by the action of free radicals. Additionally, studies have reported that the redox state of the cell is a pivotal determinant of the fate of the cells. Furthermore, reactive oxygen species have been reported to be cytotoxic, causing inflammatory disease, including tissue necrosis, organ failure, atherosclerosis, infertility, birth defects, premature aging, mutations and malignancy. Thus, the control of oxidation and reduction is important for a number of reasons including for control and prevention of strokes, heart attacks, oxidative stress and hypertension. In this regard, reductases, and particularly, oxidoreductases, are of interest. Publications further describing this subject matter include Kelsey, et al., *Br. J. Cancer*, 76(7):852-4 (1997); Friedrich and Weiss, *J. Theor. Biol.*, 187(4):529-40 (1997) and Pieuille, et al., *J. Bacteriol.*, 179(18):5684-92 (1997).

[0141] Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins, particularly secreted proteins which have homology to reductase. Many efforts are focused on the screening of mammalian recombinant DNA libraries to

identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637)].

[0142] We herein describe the identification and characterization of novel polypeptides having homology to reductase, designated herein as PRO238 polypeptides.

[0143] 25. PRO233

[0144] Studies have reported that the redox state of the cell is an important determinant of the fate of the cell. Furthermore, reactive oxygen species have been reported to be cytotoxic, causing inflammatory disease, including tissue necrosis, organ failure, atherosclerosis, infertility, birth defects, premature aging, mutations and malignancy. Thus, the control of oxidation and reduction is important for a number of reasons, including the control and prevention of strokes, heart attacks, oxidative stress and hypertension. Oxygen free radicals and antioxidants appear to play an important role in the central nervous system after cerebral ischemia and reperfusion. Moreover, cardiac injury, related to ischaemia and reperfusion has been reported to be caused by the action of free radicals. In this regard, reductases, and particularly, oxidoreductases, are of interest. In addition, the transcription factors, NF-kappa B and AP-1, are known to be regulated by redox state and to affect the expression of a large variety of genes thought to be involved in the pathogenesis of AIDS, cancer, atherosclerosis and diabetic complications. Publications further describing this subject matter include Kelsey, et al., *Br. J. Cancer*, 76(7):852-4 (1997); Friedrich and Weiss, *J. Theor. Biol.*, 187(4):529-40 (1997) and Pieulle, et al., *J. Bacteriol.*, 179(18):5684-92 (1997). Given the physiological importance of redox reactions in vivo, efforts are currently being undertaken to identify new, native proteins which are involved in redox reactions. We describe herein the identification of novel polypeptides which have homology to reductase, designated herein as PRO233 polypeptides.

[0145] 26. PRO223

[0146] The carboxypeptidase family of exopeptidases constitutes a diverse group of enzymes that hydrolyze carboxyl-terminal amide bonds in polypeptides, wherein a large number of mammalian tissues produce these enzymes. Many of the carboxypeptidase enzymes that have been identified to date exhibit rather strong cleavage specificities for certain amino acids in polypeptides. For example, carboxypeptidase enzymes have been identified which prefer lysine, arginine, serine or amino acids with either aromatic or branched aliphatic side chains as substrates at the carboxyl terminus of the polypeptide.

[0147] With regard to the serine carboxypeptidases, such amino acid specific enzymes have been identified from a variety of different mammalian and non-mammalian organisms. The mammalian serine carboxypeptidase enzymes play important roles in many different biological processes including, for example, protein digestion, activation, inactivation, or modulation of peptide hormone activity, and alteration of the physical properties of proteins and enzymes.

[0148] In light of the physiological importance of the serine carboxypeptidases, efforts are being undertaken by

both industry and academia to identify new, native secreted and membrane-bound receptor proteins and specifically novel carboxypeptidases. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. We describe herein novel polypeptides having homology to one or more serine carboxypeptidase polypeptides, designated herein as PRO223 polypeptides.

[0149] 27. PRO235

[0150] Plexin was first identified in *Xenopus* tadpole nervous system as a membrane glycoprotein which was shown to mediate cell adhesion via a homophilic binding mechanism in the presence of calcium ions. Strong evolutionary conservation between *Xenopus*, mouse and human homologs of plexin has been observed. [Kaneyama et al., *Biochem. And Biophys. Res. Comm.* 226: 524-529 (1996)]. Given the physiological importance of cell adhesion mechanisms in vivo, efforts are currently being undertaken to identify new, native proteins which are involved in cell adhesion. We describe herein the identification of a novel polypeptide which has homology to plexin, designated herein as PRO235.

[0151] 28. PRO236 and PRO262

[0152]  $\beta$ -galactosidase is a well known enzymatic protein which functions to hydrolyze  $\beta$ -galactoside molecules.  $\beta$ -galactosidase has been employed for a variety of different applications, both in vitro and in vivo and has proven to be an extremely useful research tool. As such, there is an interest in obtaining novel polypeptides which exhibit homology to the  $\beta$ -galactosidase polypeptide.

[0153] Given the strong interest in obtaining novel polypeptides having homology to  $\beta$ -galactosidase, efforts are currently being undertaken by both industry and academia to identify new, native  $\beta$ -galactosidase homolog proteins. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel  $\beta$ -galactosidase-like proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.* 93:7108-7113 (1996); U.S. Pat. No. 5,536,637)]. We herein describe novel polypeptides having significant homology to the  $\beta$ -galactosidase enzyme, designated herein as PRO236 and PRO262 polypeptides.

[0154] 29. PRO239

[0155] Densin is a glycoprotein which has been isolated from the brain which has all the hallmarks of an adhesion molecule. It is highly concentrated at synaptic sites in the brain and is expressed prominently in dendritic processes in developing neurons. Densin has been characterized as a member of the O-linked sialoglycoproteins. Densin has relevance to medically important processes such as regeneration. Given the physiological importance of synaptic processes and cell adhesion mechanisms in vivo, efforts are currently being undertaken to identify new, native proteins which are involved in synaptic machinery and cell adhesion. We describe herein the identification of novel polypeptides which have homology to densin, designated herein as PRO239 polypeptides.

[0156] 30. PRO257

[0157] Ebnerin is a cell surface protein associated with von Ebner glands in mammals. Efforts are being undertaken by both industry and academia to identify new, native cell surface receptor proteins and specifically those which possess sequence homology to cell surface proteins such as ebnerin. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor proteins. We herein describe the identification of novel polypeptides having significant homology to the von Ebner's gland-associated protein ebnerin, designated herein as PRO257 polypeptides.

[0158] 31. PRO260

[0159] Fucosidases are enzymes that remove fucose residues from fucose containing proteoglycans. In some pathological conditions, such as cancer, rheumatoid arthritis, and diabetes, there is an abnormal fucosylation of serum proteins. Therefore, fucosidases, and proteins having homology to fucosidase, are of importance to the study and abrogation of these conditions. In particular, proteins having homology to the alpha-1-fucosidase precursor are of interest. Fucosidases and fucosidase inhibitors are further described in U.S. Pat. Nos. 5,637,490, 5,382,709, 5,240,707, 5,153,325, 5,100,797, 5,096,909 and 5,017,704. Studies are also reported in Valk, et al., *J. Virol.*, 71(9):6796 (1997), Aktogu, et al., *Monaldi. Arch. Chest Dis. (Italy)*, 52(2):118 (1997) and Focarelli, et al., *Biochem. Biophys. Res. Commun. (U.S.)*, 234(1):54 (1997).

[0160] Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins. Of particular interest are proteins having homology to the alpha-1-fucosidase precursor. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0161] We herein describe the identification and characterization of novel polypeptides having homology to fucosidases, designated herein as PRO260 polypeptides.

[0162] 32. PRO263

[0163] CD44 is a cell surface adhesion molecule involved in cell-cell and cell-matrix interactions. Hyaluronic acid, a component of the extracellular matrix is a major ligand. Other ligands include collagen, fibronectin, laminin, chondroitin sulfate, mucosal addressin, serglycin and osteopontin. CD44 is also important in regulating cell traffic, lymph node homing, transmission of growth signals, and presentation of chemokines and growth factors to traveling cells. CD44 surface proteins are associated with metastatic tumors and CD44 has been used as a marker for HIV infection. Certain splice variants are associated with metastasis and poor prognosis of cancer patients. Therefore, molecules having homology with CD44 are of particular interest, as their homology indicates that they may have functions related to those functions of CD44. CD44 is further described in U.S. Pat. Nos. 5,506,119, 5,504,194 and 5,108,904; Gerberick, et al., *Toxicol. Appl. Pharmacol.*, 146(1):1 (1997); Wittig, et

al., *Immunol. Letters (Netherlands)*, 57(1-3):217 (1997); and Oliveira and Odell, *Oral Oncol. (England)*, 33(4):260 (1997).

[0164] Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins, particularly transmembrane proteins with homology to CD44 antigen. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0165] We herein describe the identification and characterization of novel polypeptides having homology to CD44 antigen, designated herein as PRO263 polypeptides.

[0166] 33. PRO270

[0167] Thioredoxins effect reduction-oxidation (redox) state. Many diseases are potentially related to redox state and reactive oxygen species may play a role in many important biological processes. The transcription factors, NF-kappa B and AP-1, are regulated by redox state and are known to affect the expression of a large variety of genes thought to be involved in the pathogenesis of AIDS, cancer, atherosclerosis and diabetic complications. Such proteins may also play a role in cellular antioxidant defense, and in pathological conditions involving oxidative stress such as stroke and inflammation in addition to having a role in apoptosis. Therefore, thioredoxins, and proteins having homology thereto, are of interest to the scientific and medical communities.

[0168] We herein describe the identification and characterization of novel polypeptides having homology to thioredoxin, designated herein as PRO270 polypeptides.

[0169] 34. PRO271

[0170] The proteoglycan link protein is a protein which is intimately associated with various extracellular matrix proteins and more specifically with proteins such as collagen. For example, one primary component of collagen is a large proteoglycan called aggrecan. This molecule is retained by binding to the glycosaminoglycan hyaluronan through the amino terminal G1 globular domain of the core protein. This binding is stabilized by the proteoglycan link protein which is a protein that is also associated with other tissues containing hyaluronan binding proteoglycans such as versican.

[0171] Link protein has been identified as a potential target for autoimmune antibodies in individuals who suffer from juvenile rheumatoid arthritis (see Guerassimov et al., *J. Rheumatology* 24(5):959-964 (1997)). As such, there is strong interest in identifying novel proteins having homology to link protein. We herein describe the identification and characterization of novel polypeptides having such homology, designated herein as PRO271 polypeptides.

[0172] 35. PRO272

[0173] Reticulocalbin is an endoplasmic reticular protein which may be involved in protein transport and luminal protein processing. Reticulocalbin resides in the lumen of the endoplasmic reticulum, is known to bind calcium, and may be involved in a luminal retention mechanism of the endoplasmic reticulum. It contains six domains of the EF-hand motif associated with high affinity calcium binding. We describe herein the identification and characterization of a

novel polypeptide which has homology to the reticulocalbin protein, designated herein as PRO272.

[0174] 36. PRO294

[0175] Collagen, a naturally occurring protein, finds wide application in industry. Chemically hydrolyzed natural collagen can be denatured and renatured by heating and cooling to produce gelatin, which is used in photographic and medical, among other applications. Collagen has important properties such as the ability to form interchain aggregates having a conformation designated as a triple helix. We herein describe the identification and characterization of a novel polypeptide which has homology to portions of the collagen molecule, designated herein as PRO294.

[0176] 37. PRO295

[0177] The integrins comprise a supergene family of cell-surface glycoprotein receptors that promote cellular adhesion. Each cell has numerous receptors that define its cell adhesive capabilities. Integrins are involved in a wide variety of interaction between cells and other cells or matrix components. The integrins are of particular importance in regulating movement and function of immune system cells. The platelet IIb/IIIa integrin complex is of particular importance in regulating platelet aggregation. A member of the integrin family, integrin  $\beta$ -6, is expressed on epithelial cells and modulates epithelial inflammation. Another integrin, leucocyte-associated antigen-1 (LFA-1) is important in the adhesion of lymphocytes during an immune response. The integrins are expressed as heterodimers of non-covalently associated alpha and beta subunits. Given the physiological importance of cell adhesion mechanisms in vivo, efforts are currently being undertaken to identify new, native proteins which are involved in cell adhesion. We describe herein the identification and characterization of a novel polypeptide which has homology to integrin, designated herein as PRO295.

[0178] 38. PRO293

[0179] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0180] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leucine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0181] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in

pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Others studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vouv. Rev. Fr. Hematol.* (Germany), 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome and Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats. Another protein of particular interest which has been reported to have leucine-rich repeats, is the SLIT protein which has been reported to be useful in treating neurodegenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanistsakonias, S. and Rothberg, J. M., WO921 0518-A1 by Yale University. Other studies reporting on the biological functions of proteins having leucine-rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-1133 (October 1995) (kidney disease involvement); and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer Research Foundation (decorin binding to transforming growth factor $\beta$  involvement for treatment for cancer, wound healing and scarring).

[0182] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions. Of particular interest are those proteins having leucine rich repeats and homology to known neuronal leucine rich repeat proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound proteins having leucine rich repeats. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0183] We describe herein the identification and characterization of a novel polypeptide which has homology to leucine rich repeat proteins, designated herein as PRO293.

[0184] 39. PRO247

[0185] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0186] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leucine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible

for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0187] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Others studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vow. Rev. Fr. Hematol.* (Germany), 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome and Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats. Another protein of particular interest which has been reported to have leucine-rich repeats is the SLIT protein which has been reported to be useful in treating neurodegenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanistsakonas, S. and Rothberg, J. M., WO9210518-A1 by Yale University. Other studies reporting on the biological functions of proteins having leucine-rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-1133 (October 1995) (kidney disease involvement); and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer Research Foundation (decorin binding to transforming growth factor $\beta$  involvement for treatment for cancer, wound healing and scarring).

[0188] Densin is a glycoprotein which has been isolated from the brain which has all the hallmarks of an adhesion molecule. It is highly concentrated at synaptic sites in the brain and is expressed prominently in dendritic processes in developing neurons. Densin has been characterized as a member of the O-linked sialoglycoproteins. Densin has relevance to medically important processes such as regeneration. Given the physiological importance of synaptic processes and cell adhesion mechanisms in vivo, efforts are currently being undertaken to identify new, native proteins which are involved in synaptic machinery and cell adhesion. Densin is further described in Kennedy, M. B, *Trends Neurosci.* (England), 20(6):264 (1997) and Apperson, et al., *J. Neurosci.*, 16(21):6839 (1996).

[0189] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions. Of particular interest are those proteins having leucine rich repeats and homology to known proteins having leucine rich repeats such as KIAA0231 and densin. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound proteins having leucine rich repeats. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0190] We describe herein the identification and characterization of a novel polypeptide which has homology to leucine rich repeat proteins, designated herein as PRO247.

[0191] 40. PRO302, PRO303, PRO304, PRO307 and PRO343

[0192] Proteases are enzymatic proteins which are involved in a large number of very important biological processes in mammalian and non-mammalian organisms. Numerous different protease enzymes from a variety of different mammalian and non-mammalian organisms have been both identified and characterized. The mammalian protease enzymes play important roles in many different biological processes including, for example, protein digestion, activation, inactivation, or modulation of peptide hormone activity, and alteration of the physical properties of proteins and enzymes.

[0193] In light of the important physiological roles played by protease enzymes, efforts are currently being undertaken by both industry and academia to identify new, native protease homologs. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637]. We herein describe the identification of novel polypeptides having homology to various protease enzymes, designated herein as PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides.

[0194] 41. PRO328

[0195] The GLIP protein family has been characterized as comprising zinc-finger proteins which play important roles in embryogenesis. These proteins may function as transcriptional regulatory proteins and are known to be amplified in a subset of human tumors. Glioma pathogenesis protein is structurally related to a group of plant pathogenesis-related proteins. It is highly expressed in glioblastoma. See U.S. Pat. No. 5,582,981 (issued Dec. 10, 1996) and U.S. Pat. No. 5,322,801 (issued Jun. 21, 1996), Ellington, A. D. et al., *Nature*, 346:818 (1990), Grindley, J. C. et al., *Dev. Biol.*, 188(2):337 (1997), Marine, J. C. et al., *Mech. Dev.*, 63(2):211 (1997). The CRISP or cysteine rich secretory protein family are a group of proteins which are also structurally related to a group of plant pathogenesis proteins. [Schwidetzky, U., *Biochem. J.*, 321:325 (1997), Pfisterer, P., *Mol. Cell Biol.*, 16(11):6160 (1996), Kratzschmar, J., *Eur. J. Biochem.*, 236(3):827 (1996)]. We describe herein the identification of a novel polypeptide which has homology to GLIP and CRISP, designated herein as PRO328 polypeptides.

[0196] 42. PRO335, PRO331 and PRO326

[0197] Protein-protein interactions include receptor and antigen complexes and signaling mechanisms. As more is known about the structural and functional mechanisms underlying protein-protein interactions, protein-protein interactions can be more easily manipulated to regulate the particular result of the protein-protein interaction. Thus, the underlying mechanisms of protein-protein interactions are of interest to the scientific and medical community.

[0198] All proteins containing leucine-rich repeats are thought to be involved in protein-protein interactions. Leu-

cine-rich repeats are short sequence motifs present in a number of proteins with diverse functions and cellular locations. The crystal structure of ribonuclease inhibitor protein has revealed that leucine-rich repeats correspond to beta-alpha structural units. These units are arranged so that they form a parallel beta-sheet with one surface exposed to solvent, so that the protein acquires an unusual, nonglobular shape. These two features have been indicated as responsible for the protein-binding functions of proteins containing leucine-rich repeats. See, Kobe and Deisenhofer, *Trends Biochem. Sci.*, 19(10):415-421 (October 1994).

[0199] A study has been reported on leucine-rich proteoglycans which serve as tissue organizers, orienting and ordering collagen fibrils during ontogeny and are involved in pathological processes such as wound healing, tissue repair, and tumor stroma formation. Iozzo, R. V., *Crit. Rev. Biochem. Mol. Biol.*, 32(2):141-174 (1997). Others studies implicating leucine rich proteins in wound healing and tissue repair are De La Salle, C., et al., *Vow. Rev. Fr. Hematol.* (Germany), 37(4):215-222 (1995), reporting mutations in the leucine rich motif in a complex associated with the bleeding disorder Bernard-Soulier syndrome, Chlemetson, K. J., *Thromb. Haemost.* (Germany), 74(1):111-116 (July 1995), reporting that platelets have leucine rich repeats and Ruoslahti, E. I., et al., WO9110727-A by La Jolla Cancer Research Foundation reporting that decorin binding to transforming growth factor $\beta$  has involvement in a treatment for cancer, wound healing and scarring. Related by function to this group of proteins is the insulin like growth factor (IGF), in that it is useful in wound-healing and associated therapies concerned with re-growth of tissue, such as connective tissue, skin and bone; in promoting body growth in humans and animals; and in stimulating other growth-related processes. The acid labile subunit of IGF (ALS) is also of interest in that it increases the half-life of IGF and is part of the IGF complex in vivo.

[0200] Another protein which has been reported to have leucine-rich repeats is the SLIT protein which has been reported to be useful in treating neuro-degenerative diseases such as Alzheimer's disease, nerve damage such as in Parkinson's disease, and for diagnosis of cancer, see, Artavanisakonas, S. and Rothberg, J. M., WO9210518-A1 by Yale University. Of particular interest is LIG-1, a membrane glycoprotein that is expressed specifically in glial cells in the mouse brain, and has leucine rich repeats and immunoglobulin-like domains. Suzuki, et al., *J. Biol. Chem.* (U.S.), 271(37):22522 (1996). Other studies reporting on the biological functions of proteins having leucine rich repeats include: Tayar, N., et al., *Mol. Cell Endocrinol.*, (Ireland), 125(1-2):65-70 (December 1996) (gonadotropin receptor involvement); Miura, Y., et al., *Nippon Rinsho* (Japan), 54(7):1784-1789 (July 1996) (apoptosis involvement); Harris, P. C., et al., *J. Am. Soc. Nephrol.*, 6(4):1125-1133 (October 1995) (kidney disease involvement).

[0201] Efforts are therefore being undertaken by both industry and academia to identify new proteins having leucine rich repeats to better understand protein-protein interactions. Of particular interest are those proteins having leucine rich repeats and homology to known proteins having leucine rich repeats such as LIG-1, ALS and decorin. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound proteins having leu-

cine rich repeats. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0202] We describe herein the identification and characterization of novel polypeptides which have homology to proteins of the leucine rich repeat superfamily, designated herein as PRO335, PRO331 and PRO326 polypeptides.

[0203] 43. PRO332

[0204] Secreted proteins comprising a repeat characterized by an arrangement of conserved leucine residues (leucine-rich repeat motif) have diverse biological roles. Certain proteoglycans, such as biglycan, fibromodulin and decorin, are, for example, characterized by the presence of a leucine-rich repeat of about 24 amino acids [Ruoslahti, *Ann. Rev. Cell. Biol.* 4 229-255 (1988); Oldberg et al., *EMBO J.* 8, 2601-2604 (1989)]. In general, proteoglycans are believed to play a role in regulating extracellular matrix, cartilage or bone function. The proteoglycan decorin binds to collagen type I and II and affects the rate of fibril formation. Fibromodulin also binds collagen and delays fibril formation. Both fibromodulin and decorin inhibit the activity of transforming growth factor beta (TGF- $\beta$ ) (U.S. Pat. No. 5,583, 103 issued Dec. 10, 1996). TGF- $\beta$  is known to play a key role in the induction of extracellular matrix and has been implicated in the development of fibrotic diseases, such as cancer and glomerulonephritis. Accordingly, proteoglycans have been proposed for the treatment of fibrotic cancer, based upon their ability to inhibit TGF- $\beta$ 's growth stimulating activity on the cancer cell. Proteoglycans have also been described as potentially useful in the treatment of other proliferative pathologies, including rheumatoid arthritis, arteriosclerosis, adult respiratory distress syndrome, cirrhosis of the liver, fibrosis of the lungs, post-myocardial infarction, cardiac fibrosis, post-angioplasty restenosis, renal interstitial fibrosis and certain dermal fibrotic conditions, such as keloids and scarring, which might result from burn injuries, other invasive skin injuries, or cosmetic or reconstructive surgery (U.S. Pat. No. 5,654,270, issued Aug. 5, 1997).

[0205] We describe herein the identification and characterization of novel polypeptides which have homology to proteins of the leucine rich repeat superfamily, designated herein as PRO332 polypeptides.

[0206] 44. PRO334

[0207] Microfibril bundles and proteins found in association with these bundles, particularly attachment molecules, are of interest in the field of dermatology, particularly in the study of skin which has been damaged from aging, injuries or the sun. Fibrillin microfibrils define the continuous elastic network of skin, and are present in dermis as microfibril bundles devoid of measurable elastin extending from the dermal-epithelial junction and as components of the thick elastic fibres present in the deep reticular dermis. Moreover, Marfan syndrome has been linked to mutations which interfere with multimerization of fibrillin monomers or other connective tissue elements.

[0208] Fibulin-1 is a modular glycoprotein with amino-terminal anaphalatoxin-like modules followed by nine epidermal growth factor (EGF)-like modules and, depending on alternative splicing, four possible carboxyl termini. Fibu-

lin-2 is a novel extracellular matrix protein frequently found in close association with microfibrils containing either fibronectin or fibrillin. Thus, fibrillin, fibulin, and molecules related thereto are of interest, particularly for the use of preventing skin from being damaged from aging, injuries or the sun, or for restoring skin damaged from same. Moreover, these molecules are generally of interest in the study of connective tissue and attachment molecules and related mechanisms. Fibrillin, fibulin and related molecules are further described in Adams, et al., *J. Mol. Biol.*, 272(2):226-36 (1997); Kielty and Shuttleworth, *Microsc. Res. Tech.*, 38(4):413-27 (1997); and Child, *J. Card. Surg.* 12(2Supp.):131-5 (1997).

[0209] Currently, efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins, particularly secreted proteins which have homology to fibulin and fibrillin. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0210] We herein describe the identification and characterization of novel polypeptides having homology to fibulin and fibrillin, designated herein as PRO334 polypeptides.

[0211] 45. PRO346

[0212] The widespread occurrence of cancer has prompted the devotion of considerable resources and discovering new treatments of treatment. One particular method involves the creation of tumor or cancer specific monoclonal antibodies (mAbs) which are specific to tumor antigens. Such mAbs, which can distinguish between normal and cancerous cells are useful in the diagnosis, prognosis and treatment of the disease. Particular antigens are known to be associated with neoplastic diseases, such as colorectal and breast cancer. Since colon cancer is a widespread disease, early diagnosis and treatment is an important medical goal. Diagnosis and treatment of cancer can be implemented using monoclonal antibodies (mAbs) specific therefore having fluorescent, nuclear magnetic or radioactive tags. Radioactive genes, toxins and/or drug tagged mAbs can be used for treatment in situ with minimal patient description.

[0213] Carcinoembryonic antigen (CEA) is a glycoprotein found in human colon cancer and the digestive organs of a 2-6 month human embryos. CEA is a known human tumor marker and is widely used in the diagnosis of neoplastic diseases, such as colon cancer. For example, when the serum levels of CEA are elevated in a patient, a drop of CEA levels after surgery would indicate the tumor resection was successful. On the other hand, a subsequent rise in serum CEA levels after surgery would indicate that metastases of the original tumor may have formed or that new primary tumors may have appeared. CEA may also be a target for mAb, antisense nucleotides

[0214] 46. PRO268

[0215] Protein disulfide isomerase is an enzymatic protein which is involved in the promotion of correct refolding of proteins through the establishment of correct disulfide bond formation. Protein disulfide isomerase was initially identi-

fied based upon its ability to catalyze the renaturation of reduced denatured RNase (Goldberger et al., *J. Biol. Chem.* 239:1406-1410 (1964) and Epstein et al., *Cold Spring Harbor Symp. Quant. Biol.* 28:439-449 (1963)). Protein disulfide isomerase has been shown to be a resident enzyme of the endoplasmic reticulum which is retained in the endoplasmic reticulum via a -KDEL or -HDEL amino acid sequence at its C-terminus.

[0216] Given the importance of disulfide bond-forming enzymes and their potential uses in a number of different applications, for example in increasing the yield of correct refolding of recombinantly produced proteins, efforts are currently being undertaken by both industry and academia to identify new, native proteins having homology to protein disulfide isomerase. Many of these efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel protein disulfide isomerase homologs. We herein describe a novel polypeptide having homology to protein disulfide isomerase, designated herein as PRO268.

[0217] 47. PRO330

[0218] Prolyl 4-hydroxylase is an enzyme which functions to post-translationally hydroxylate proline residues at the Y position of the amino acid sequence Gly-X-Y, which is a repeating three amino acid sequence found in both collagen and procollagen. Hydroxylation of proline residues at the Y position of the Gly-X-Y amino acid triplet to form 4-hydroxyproline residues at those positions is required before newly synthesized collagen polypeptide chains may fold into their proper three-dimensional triple-helical conformation. If hydroxylation does not occur, synthesized collagen polypeptides remain non-helical, are poorly secreted by cells and cannot assemble into stable functional collagen fibrils. Vuorio et al., *Proc. Natl. Acad. Sci. USA* 89:7467-7470 (1992). Prolyl 4-hydroxylase is comprised of at least two different polypeptide subunits, alpha and beta.

[0219] Efforts are being undertaken by both industry and academia to identify new, native secreted and membrane-bound receptor proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted and membrane-bound receptor proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108-7113 (1996); U.S. Pat. No. 5,536,637]. Based upon these efforts, Applicants have herein identified and describe a novel polypeptide having homology to the alpha subunit of prolyl 4-hydroxylase, designated herein as PRO330.

[0220] 48. PRO339 and PRO310

[0221] Fringe is a protein which specifically blocks serrate-mediated activation of notch in the dorsal compartment of the *Drosophila* wing imaginal disc. Fleming, et al., *Development*, 124(15):2973-81 (1997). Therefore, fringe is of interest for both its role in development as well as its ability to regulate serrate, particularly serrate's signaling abilities. Also of interest are novel polypeptides which may have a role in development and/or the regulation of serrate-like molecules. Of particular interest are novel polypeptides having homology to fringe as identified and described herein, designated herein as PRO339 and PRO310 polypeptides.



[0222] 49. PRO244

[0223] Lectins are a class of proteins comprising a region that binds carbohydrates specifically and non-covalently. Numerous lectins have been identified in higher animals, both membrane-bound and soluble, and have been implicated in a variety of cell-recognition phenomena and tumor metastasis.

[0224] Most lectins can be classified as either C-type (calcium-dependent) or S-type (thiol-dependent).

[0225] Lectins are thought to play a role in regulating cellular events that are initiated at the level of the plasma membrane. For example, plasma membrane associated molecules are involved in the activation of various subsets of lymphoid cells, e.g. T-lymphocytes, and it is known that cell surface molecules are responsible for activation of these cells and consequently their response during an immune reaction.

[0226] A particular group of cell adhesion molecules, selecting, belong in the superfamily of C-type lectins. This group includes L-selectin (peripheral lymph node homing receptor (pnHR), LEC-CAM-1, LAM-1, gp90<sup>MEL</sup>, gp100<sup>MEL</sup>, gp110<sup>MEL</sup>, MEL-14 antigen, Leu-8 antigen, TQ-1 antigen, DREG antigen), E-selectin (LEC-CAM-2, LECAM-2, ELAM-1), and P-selectin (LEC-CAM-3, LECAM-3, GMP-140, PADGEM). The structure of selectins consists of a C-type lectin (carbohydrate binding) domain, an epidermal growth factor-like (EGF-like) motif, and variable numbers of complement regulatory (CR) motifs. Selectins are associated with leukocyte adhesion, e.g. the attachment of neutrophils to venular endothelial cells adjacent to inflammation (E-selectin), or with the trafficking of lymphocytes from blood to secondary lymphoid organs, e.g. lymph nodes and Peyer's patches (L-selectin).

[0227] Another exemplary lectin is the cell-associated macrophage antigen, Mac-2 that is believed to be involved in cell adhesion and immune responses. Macrophages also express a lectin that recognizes Tn Ag, a human carcinoma-associated epitope.

[0228] Another C-type lectin is CD95 (Fas antigen/APO-1) that is an important mediator of immunologically relevant regulated or programmed cell death (apoptosis). "Apoptosis" is a non-necrotic cell death that takes place in metazoan animal cells following activation of an intrinsic cell suicide program. The cloning of Fas antigen is described in PCT publication WO 91/10448, and European patent application EP510691. The mature Fas molecule consists of 319 amino acids of which 157 are extracellular, 17 constitute the transmembrane domain, and 145 are intracellular. Increased levels of Fas expression at T cell surface have been associated with tumor cells and HIV-infected cells. Ligation of CD95 triggers apoptosis in the presence of interleukin-1 (IL-2).

[0229] C-type lectins also include receptors for oxidized low-density lipoprotein (LDL). This suggests a possible role in the pathogenesis of atherosclerosis.

[0230] We herein describe the identification and characterization of novel polypeptides having homology to C-type lectins, designated herein as PRO244 polypeptides.

## SUMMARY OF THE INVENTION

[0231] 1. PRO211 and PRO217

[0232] Applicants have identified cDNA clones that encode novel polypeptides having homology to EGF, designated in the present application as "PRO211" and "PRO217" polypeptides.

[0233] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO211 or PRO217 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding EGF-like homologue PRO211 and PRO217 polypeptides of **FIG. 2** (SEQ ID NO:2) and/or **4** (SEQ ID NO:4) indicated in **FIG. 1** (SEQ ID NO:1) and/or **FIG. 3** (SEQ ID NO:3), respectively, or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0234] In another embodiment, the invention provides isolated PRO211 and PRO217 EGF-like homologue PRO211 and PRO217 polypeptides. In particular, the invention provides isolated native sequence PRO211 and PRO217 EGF-like homologue polypeptides, which in one embodiment, includes an amino acid sequence comprising residues: 1 to 353 of **FIG. 2** (SEQ ID NO:2) or (2) 1 to 379 of **FIG. 4** (SEQ ID NO: 4).

[0235] 2. PRO230

[0236] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO230".

[0237] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO230 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO230 polypeptide having amino acid residues 1 through 467 of **FIG. 6** (SEQ ID NO:12), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0238] In another embodiment, the invention provides isolated PRO230 polypeptide. In particular, the invention provides isolated native sequence PRO230 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 through 467 of **FIG. 6** (SEQ ID NO:12).

[0239] In another embodiment, the invention provides an expressed sequence tag (EST) comprising the nucleotide sequence of SEQ ID NO:13 (**FIG. 7**) which is herein designated as DNA20088.

[0240] 3. PRO232

[0241] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO232".

[0242] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO232 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO232 polypeptide having amino acid residues 1 to 114 of **FIG. 9** (SEQ ID NO:18), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0243] In another embodiment, the invention provides isolated PRO232 polypeptide. In particular, the invention provides isolated native sequence PRO232 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 114 of **FIG. 9** (SEQ ID NO:18).

[0244] 4. PRO187

[0245] Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "PRO187".

[0246] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO187 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO187 polypeptide of **FIG. 11** (SEQ ID NO:23), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides a nucleic acid comprising the coding sequence of **FIG. 10** (SEQ ID NO:22) or its complement. In another aspect, the invention provides a nucleic acid of the full length protein of clone DNA27864-1155, deposited with the ATCC under accession number ATCC 209375, alternatively the coding sequence of clone DNA27864-1155, deposited under accession number ATCC 209375.

[0247] In yet another embodiment, the invention provides isolated PRO187 polypeptide. In particular, the invention provides isolated native sequence PRO187 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 205 of **FIG. 11** (SEQ ID NO:23). Alternatively, the invention provides a polypeptide encoded by the nucleic acid deposited under accession number ATCC 209375.

[0248] 5. PRO265

[0249] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO265".

[0250] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO265 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO265 polypeptide having amino acid residues 1 to 660 of **FIG. 13** (SEQ ID NO:28), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0251] In another embodiment, the invention provides isolated PRO265 polypeptide. In particular, the invention provides isolated native sequence PRO265 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 660 of **FIG. 13** (SEQ ID NO:28). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO265 polypeptide.

[0252] 6. PRO219

[0253] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO219".

[0254] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO219 polypeptide. In one aspect, the isolated nucleic acid

comprises DNA encoding the PRO219 polypeptide having amino acid residues 1 to 915 of **FIG. 15** (SEQ ID NO:34), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0255] In another embodiment, the invention provides isolated PRO219 polypeptide. In particular, the invention provides isolated native sequence PRO219 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 915 of **FIG. 15** (SEQ ID NO:34).

[0256] 7. PRO246

[0257] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO246".

[0258] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO246 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO246 polypeptide having amino acid residues 1 to 390 of **FIG. 17** (SEQ ID NO:39), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0259] In another embodiment, the invention provides isolated PRO246 polypeptide. In particular, the invention provides isolated native sequence PRO246 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 390 of **FIG. 17** (SEQ ID NO:39). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO246 polypeptide.

[0260] 8. PRO228

[0261] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to CD97, EMR1 and latrophilin, wherein the polypeptide is designated in the present application as "PRO228".

[0262] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO228 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO228 polypeptide having amino acid residues 1 to 690 of **FIG. 19** (SEQ ID NO:49), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0263] In another embodiment, the invention provides isolated PRO228 polypeptide. In particular, the invention provides isolated native sequence PRO228 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 690 of **FIG. 19** (SEQ ID NO:49). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO228 polypeptide.

[0264] In another embodiment, the invention provides an expressed sequence tag (EST) comprising the nucleotide sequence of SEQ ID NO:50, designated herein as DNA21951.

[0265] 9. PRO533

[0266] Applicants have identified a cDNA clone (DNA49435-1219) that encodes a novel polypeptide, designated in the present application as PRO533.

[0267] In one embodiment, the invention provides an isolated nucleic acid molecule having at least about 80% sequence identity to (a) a DNA molecule encoding a PRO533 polypeptide comprising the sequence of amino acids 23 to 216 of FIG. 22 (SEQ ID NO:59), or (b) the complement of the DNA molecule of (a). The sequence identity preferably is about 85%, more preferably about 90%, most preferably about 95%. In one aspect, the isolated nucleic acid has at least about 80%, preferably at least about 85%, more preferably at least about 90%, and most preferably at least about 95% sequence identity with a polypeptide having amino acid residues 23 to 216 of FIG. 22 (SEQ ID NO:59). Preferably, the highest degree of sequence identity occurs within the secreted portion (amino acids 23 to 216 of FIG. 22, SEQ ID NO:59). In a further embodiment, the isolated nucleic acid molecule comprises DNA encoding a PRO533 polypeptide having amino acid residues 1 to 216 of FIG. 22 (SEQ ID NO:59), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides a nucleic acid of the full length protein of clone DNA49435-1219, deposited with the ATCC under accession number ATCC 209480.

[0268] In yet another embodiment, the invention provides isolated PRO533 polypeptide. In particular, the invention provides isolated native sequence PRO533 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 23 to 216 of FIG. 22 (SEQ ID NO:59). Native PRO533 polypeptides with or without the native signal sequence (amino acids 1 to 22 in FIG. 22 (SEQ ID NO:59)), and with or without the initiating methionine are specifically included. Alternatively, the invention provides a PRO533 polypeptide encoded by the nucleic acid deposited under accession number ATCC 209480.

[0269] 10. PRO245

[0270] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO245".

[0271] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO245 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO245 polypeptide having amino acid residues 1 to 312 of FIG. 24 (SEQ ID NO:64), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0272] In another embodiment, the invention provides isolated PRO245 polypeptide. In particular, the invention provides isolated native sequence PRO245 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 312 of FIG. 24 (SEQ ID NO:64).

[0273] 11. PRO220, PRO221 and PRO227

[0274] Applicants have identified cDNA clones that each encode novel polypeptides, all having leucine rich repeats. These polypeptides are designated in the present application as PRO220, PRO221 and PRO227.

[0275] In one embodiment, the invention provides isolated nucleic acid molecules comprising DNA respectively encoding PRO220, PRO221 and PRO227, respectively. In one

aspect, provided herein is an isolated nucleic acid comprises DNA encoding the PRO220 polypeptide having amino acid residues 1 through 708 of FIG. 26 (SEQ ID NO:69), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. Also provided herein is an isolated nucleic acid comprises DNA encoding the PRO221 polypeptide having amino acid residues 1 through 259 of FIG. 28 (SEQ ID NO:71), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. Moreover, also provided herein is an isolated nucleic acid comprises DNA encoding the PRO227 polypeptide having amino acid residues 1 through 620 of FIG. 30 (SEQ ID NO:73), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0276] In another embodiment, the invention provides isolated PRO220, PRO221 and PRO227 polypeptides. In particular, provided herein is the isolated native sequence for the PRO220 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 708 of FIG. 26 (SEQ ID NO:69). Additionally provided herein is the isolated native sequence for the PRO221 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 259 of FIG. 28 (SEQ ID NO:71). Moreover, provided herein is the isolated native sequence for the PRO227 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 620 of FIG. 30 (SEQ ID NO:73).

[0277] 12. PRO258

[0278] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to CRTAM and poliovirus receptor precursors, wherein the polypeptide is designated in the present application as "PRO258".

[0279] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO258 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO258 polypeptide having amino acid residues 1 to 398 of FIG. 32 (SEQ ID NO:84), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0280] In another embodiment, the invention provides isolated PRO258 polypeptide. In particular, the invention provides isolated native sequence PRO258 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 398 of FIG. 32 (SEQ ID NO:84). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO258 polypeptide.

[0281] 13. PRO266

[0282] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO266".

[0283] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO266 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO266 polypeptide having

amino acid residues 1 to 696 of **FIG. 34** (SEQ ID NO:91), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0284] In another embodiment, the invention provides isolated PRO266 polypeptide. In particular, the invention provides isolated native sequence PRO266 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 696 of **FIG. 34** (SEQ ID NO:91).

[0285] 14. PRO269

[0286] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as PRO269.

[0287] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO269 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO269 polypeptide having amino acid residues 1 to 490 of **FIG. 36** (SEQ ID NO:96), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0288] In another embodiment, the invention provides isolated PRO269 polypeptide. In particular, the invention provides isolated native sequence PRO269 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 490 of **FIG. 36** (SEQ ID NO:96). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO269 polypeptide.

[0289] 15. PRO287

[0290] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO287".

[0291] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO287 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO287 polypeptide having amino acid residues 1 to 415 of **FIG. 38** (SEQ ID NO:104), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0292] In another embodiment, the invention provides isolated PRO287 polypeptide. In particular, the invention provides isolated native sequence PRO287 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 415 of **FIG. 38** (SEQ ID NO:104).

[0293] 16. PRO214

[0294] Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "PRO214".

[0295] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO214 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO214 polypeptide of **FIG. 40** (SEQ ID NO:109), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides a

nucleic acid comprising the coding sequence of **FIG. 39** (SEQ ID NO:108) or its complement. In another aspect, the invention provides a nucleic acid of the full length protein of clone DNA32286-1191, deposited with ATCC under accession number ATCC 209385.

[0296] In yet another embodiment, the invention provides isolated PRO214 polypeptide. In particular, the invention provides isolated native sequence PRO214 polypeptide, which in one embodiment, includes an amino acid sequence comprising the residues of **FIG. 40** (SEQ ID NO:109). Alternatively, the invention provides a polypeptide encoded by the nucleic acid deposited under accession number ATCC 209385.

[0297] 17. PRO317

[0298] Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "PRO317".

[0299] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding PRO317 polypeptide. In one aspect, the isolated nucleic acid comprises DNA (SEQ ID NO:113) encoding PRO317 polypeptide having amino acid residues 1 to 366 of **FIG. 42**, or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0300] In another embodiment, the invention provides isolated PRO317 polypeptide. In particular, the invention provides isolated native-sequence PRO317 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 366 of **FIG. 42** (SEQ ID NO:114).

[0301] In yet another embodiment, the invention supplies a method of detecting the presence of PRO317 in a sample, the method comprising:

[0302] a) contacting a detectable anti-PRO317 antibody with a sample suspected of containing PRO317; and

[0303] b) detecting binding of the antibody to the sample; wherein the sample is selected from the group consisting of a body fluid, a tissue sample, a cell extract, and a cell culture medium.

[0304] In a still further embodiment a method is provided for determining the presence of PRO317 mRNA in a sample, the method comprising:

[0305] a) contacting a sample suspected of containing PRO317 mRNA with a detectable nucleic acid probe that hybridizes under moderate to stringent conditions to PRO317 mRNA; and

[0306] b) detecting hybridization of the probe to the sample.

[0307] Preferably, in this method the sample is a tissue sample and the detecting step is by in situ hybridization, or the sample is a cell extract and detection is by Northern analysis.

[0308] Further, the invention provides a method for treating a PRO317-associated disorder comprising administering to a mammal an effective amount of the PRO317 polypeptide or a composition thereof containing a carrier, or with an

effective amount of a PRO317 agonist or PRO317 antagonist, such as an antibody which binds specifically to PRO317.

[0309] 18. PRO301

[0310] Applicants have identified a cDNA clone (DNA40628-1216) that encodes a novel polypeptide, designated in the present application as "PRO301".

[0311] In one embodiment, the invention provides an isolated nucleic acid molecule having at least about 80% sequence identity to (a) a DNA molecule encoding a PRO301 polypeptide comprising the sequence of amino acids 28 to 258 of **FIG. 44** (SEQ ID NO:119), or (b) the complement of the DNA molecule of (a). The sequence identity preferably is about 85%, more preferably about 90%, most preferably about 95%. In one aspect, the isolated nucleic acid has at least about 80%, preferably at least about 85%, more preferably at least about 90%, and most preferably at least about 95% sequence identity with a polypeptide having amino acid residues 28 to 258 of **FIG. 44** (SEQ ID NO:119). Preferably, the highest degree of sequence identity occurs within the extracellular domains (amino acids 28 to 258 of **FIG. 44**, SEQ ID NO:119). In a further embodiment, the isolated nucleic acid molecule comprises DNA encoding a PRO301 polypeptide having amino acid residues 28 to 299 of **FIG. 44** (SEQ ID NO:119), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides a nucleic acid of the full length protein of clone DNA40628-1216, deposited with the ATCC under accession number ATCC 209432, alternatively the coding sequence of clone DNA40628-1216, deposited under accession number ATCC 209432.

[0312] In yet another embodiment, the invention provides isolated PRO301 polypeptide. In particular, the invention provides isolated native sequence PRO301 polypeptide, which in one embodiment, includes an amino acid sequence comprising the extracellular domain residues 28 to 258 of **FIG. 44** (SEQ ID NO:119). Native PRO301 polypeptides with or without the native signal sequence (amino acids 1 to 27 in **FIG. 44** (SEQ ID NO:119), and with or without the initiating methionine are specifically included. Additionally, the sequences of the invention may also comprise the transmembrane domain (residues 236 to about 258 in **FIG. 44**; SEQ ID NO:119) and/or the intracellular domain (about residue 259 to 299 in **FIG. 44**; SEQ ID NO:119). Alternatively, the invention provides a PRO301 polypeptide encoded by the nucleic acid deposited under accession number ATCC 209432.

[0313] 19. PRO224

[0314] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO224".

[0315] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO224 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO224 polypeptide having amino acid residues 1 to 282 of **FIG. 46** (SEQ ID NO:127), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0316] In another embodiment, the invention provides isolated PRO224 polypeptide. In particular, the invention provides isolated native sequence PRO224 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 282 of **FIG. 46** (SEQ ID NO:127).

[0317] 20. PRO222

[0318] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO222".

[0319] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO222 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO222 polypeptide having amino acid residues 1 to 490 of **FIG. 48** (SEQ ID NO:132), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0320] In another embodiment, the invention provides isolated PRO222 polypeptide. In particular, the invention provides isolated native sequence PRO222 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 490 of **FIG. 48** (SEQ ID NO:132).

[0321] 21. PRO234

[0322] Applicants have identified a cDNA clone that encodes a novel lectin polypeptide molecule, designated in the present application as "PRO234".

[0323] In one embodiment, the invention provides an isolated nucleic acid encoding a novel lectin comprising DNA encoding a PRO234 polypeptide. In one aspect, the isolated nucleic acid comprises the DNA encoding PRO234 polypeptides having amino acid residues 1 to 382 of **FIG. 50** (SEQ ID NO:137), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides an isolated nucleic acid molecule comprising the nucleotide sequence of **FIG. 49** (SEQ ID NO:136).

[0324] In another embodiment, the invention provides isolated novel PRO234 polypeptides. In particular, the invention provides isolated native sequence PRO234 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 382 of **FIG. 50** (SEQ ID NO:137).

[0325] In yet another embodiment, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences.

[0326] 22. PRO231

[0327] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to a putative acid phosphatase, wherein the polypeptide is designated in the present application as "PRO231".

[0328] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO231 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO231 polypeptide having amino acid residues 1 to 428 of **FIG. 52** (SEQ ID NO:142), or is complementary to such encoding nucleic acid

sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0329] In another embodiment, the invention provides isolated PRO231 polypeptide. In particular, the invention provides isolated native sequence PRO231 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 428 of FIG. 52 (SEQ ID NO:142).

[0330] 23. PRO229

[0331] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to scavenger receptors wherein the polypeptide is designated in the present application as "PRO229".

[0332] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO229 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO229 polypeptide having amino acid residues 1 to 347 of FIG. 54 (SEQ ID NO:148), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0333] In another embodiment, the invention provides isolated PRO229 polypeptide. In particular, the invention provides isolated native sequence PRO229 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 347 of FIG. 54 (SEQ ID NO:148).

[0334] 24. PRO238

[0335] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to reductase, wherein the polypeptide is designated in the present application as "PRO238".

[0336] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO238 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO238 polypeptide having amino acid residues 1 to 310 of FIG. 56 (SEQ ID NO:153), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0337] In another embodiment, the invention provides isolated PRO238 polypeptide. In particular, the invention provides isolated native sequence PRO238 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 310 of FIG. 56 (SEQ ID NO:153).

[0338] 25. PRO233

[0339] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO233".

[0340] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO233 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO233 polypeptide having amino acid residues 1 to 300 of FIG. 58 (SEQ ID NO:159), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0341] In another embodiment, the invention provides isolated PRO233 polypeptide. In particular, the invention provides isolated native sequence PRO233 polypeptide,

which in one embodiment, includes an amino acid sequence comprising residues 1 to 300 of FIG. 58 (SEQ ID NO:159).

[0342] 26. PRO223

[0343] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to serine carboxypeptidase polypeptides, wherein the polypeptide is designated in the present application as "PRO223".

[0344] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO223 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO223 polypeptide having amino acid residues 1 to 476 of FIG. 60 (SEQ ID NO:164), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0345] In another embodiment, the invention provides isolated PRO223 polypeptide. In particular, the invention provides isolated native sequence PRO223 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 476 of FIG. 60 (SEQ ID NO:164).

[0346] 27. PRO235

[0347] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO235".

[0348] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO235 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO235 polypeptide having amino acid residues 1 to 552 of FIG. 62 (SEQ ID NO:170), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0349] In another embodiment, the invention provides isolated PRO235 polypeptide. In particular, the invention provides isolated native sequence PRO235 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 552 of FIG. 62 (SEQ ID NO:170).

[0350] 28. PRO236 and PRO262

[0351] Applicants have identified cDNA clones that encode novel polypeptides having homology to  $\beta$ -galactosidase, wherein those polypeptides are designated in the present application as "PRO236" and "PRO262".

[0352] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO236 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO236 polypeptide having amino acid residues 1 to 636 of FIG. 64 (SEQ ID NO:175), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0353] In another embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO262 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO262 polypeptide having amino acid residues 1 to 654 of FIG. 66 (SEQ ID NO:177), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0354] In another embodiment, the invention provides isolated PRO236 polypeptide. In particular, the invention provides isolated native sequence PRO236 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 636 of FIG. 64 (SEQ ID NO:175).

[0355] In another embodiment, the invention provides isolated PRO262 polypeptide. In particular, the invention provides isolated native sequence PRO262 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 654 of FIG. 66 (SEQ ID NO:177).

[0356] 29. PRO239

[0357] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO239".

[0358] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO239 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO239 polypeptide having amino acid residues 1 to 501 of FIG. 68 (SEQ ID NO:185), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0359] In another embodiment, the invention provides isolated PRO239 polypeptide. In particular, the invention provides isolated native sequence PRO239 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 501 of FIG. 68 (SEQ ID NO:185).

[0360] 30. PRO257

[0361] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO257".

[0362] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO257 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO257 polypeptide having amino acid residues 1 to 607 of FIG. 70 (SEQ ID NO:190), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0363] In another embodiment, the invention provides isolated PRO257 polypeptide. In particular, the invention provides isolated native sequence PRO257 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 607 of FIG. 70 (SEQ ID NO:190). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO257 polypeptide.

[0364] 31. PRO260

[0365] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO260".

[0366] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO260 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO260 polypeptide having amino acid residues 1 to 467 of FIG. 72 (SEQ ID NO:195), or is complementary to such encoding nucleic acid

sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0367] In another embodiment, the invention provides isolated PRO260 polypeptide. In particular, the invention provides isolated native sequence PRO260 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 467 of FIG. 72 (SEQ ID NO:195).

[0368] 32. PRO263

[0369] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to CD44 antigen, wherein the polypeptide is designated in the present application as "PRO263".

[0370] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO263 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO263 polypeptide having amino acid residues 1 to 322 of FIG. 74 (SEQ ID NO:201), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0371] In another embodiment, the invention provides isolated PRO263 polypeptide. In particular, the invention provides isolated native sequence PRO263 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 322 of FIG. 74 (SEQ ID NO:201). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO263 polypeptide.

[0372] 33. PRO270

[0373] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO270".

[0374] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO270 polypeptide. In one aspect, the isolated nucleic acid comprises DNA which includes the sequence encoding the PRO270 polypeptide having amino acid residues 1 to 296 of FIG. 76 (SEQ ID NO:207), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0375] In another embodiment, the invention provides isolated PRO270 polypeptide. In particular, the invention provides isolated native sequence PRO270 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 296 of FIG. 76 (SEQ ID NO:207).

[0376] 34. PRO271

[0377] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to the proteoglycan link protein, wherein the polypeptide is designated in the present application as "PRO271".

[0378] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO271 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO271 polypeptide having amino acid residues 1 to 360 of FIG. 78 (SEQ ID NO:213), or is complementary to such encoding nucleic acid

sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0379] In another embodiment, the invention provides isolated PRO271 polypeptide. In particular, the invention provides isolated native sequence PRO271 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 360 of **FIG. 78** (SEQ ID NO:213).

[0380] 35. PRO272

[0381] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO272".

[0382] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO272 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO272 polypeptide having amino acid residues 1 to 328 of **FIG. 80** (SEQ ID NO:221), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0383] In another embodiment, the invention provides isolated PRO272 polypeptide. In particular, the invention provides isolated native sequence PRO272 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 328 of **FIG. 80** (SEQ ID NO:211).

[0384] 36. PRO294

[0385] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO294".

[0386] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO294 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO294 polypeptide having amino acid residues 1 to 550 of **FIG. 82** (SEQ ID NO:227), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0387] In another embodiment, the invention provides isolated PRO294 polypeptide. In particular, the invention provides isolated native sequence PRO294 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 550 of **FIG. 82** (SEQ ID NO:227).

[0388] 37. PRO295

[0389] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO295".

[0390] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO295 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO295 polypeptide having amino acid residues 1 to 350 of **FIG. 84** (SEQ ID NO:236), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0391] In another embodiment, the invention provides isolated PRO295 polypeptide. In particular, the invention provides isolated native sequence PRO295 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 350 of **FIG. 84** (SEQ ID NO:236).

[0392] 38. PRO293

[0393] Applicants have identified a cDNA clone that encodes a novel human neuronal leucine rich repeat polypeptide, wherein the polypeptide is designated in the present application as "PRO293".

[0394] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO293 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO293 polypeptide having amino acid residues 1 to 713 of **FIG. 86** (SEQ ID NO:245), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0395] In another embodiment, the invention provides isolated PRO293 polypeptide. In particular, the invention provides isolated native sequence PRO293 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 713 of **FIG. 86** (SEQ ID NO:245). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO293 polypeptide.

[0396] 39. PRO247

[0397] Applicants have identified a cDNA clone that encodes a novel polypeptide having leucine rich repeats wherein the polypeptide is designated in the present application as "PRO247".

[0398] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO247 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO247 polypeptide having amino acid residues 1 to 546 of **FIG. 88** (SEQ ID NO:250), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0399] In another embodiment, the invention provides isolated PRO247 polypeptide. In particular, the invention provides isolated native sequence PRO247 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 546 of **FIG. 88** (SEQ ID NO:250). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO247 polypeptide.

[0400] 40. PRO302, PRO303, PRO304, PRO307 and PRO343

[0401] Applicants have identified cDNA clones that encode novel polypeptides having homology to various proteases, wherein those polypeptide are designated in the present application as "PRO302", "PRO303", "PRO304", "PRO307" and "PRO343" polypeptides.

[0402] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO302 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO302 polypeptide having amino acid residues 1 to 452 of **FIG. 90** (SEQ ID NO:255), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0403] In another embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a



PRO303 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO303 polypeptide having amino acid residues 1 to 314 of **FIG. 92** (SEQ ID NO:257), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0404] In yet another embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO304 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO304 polypeptide having amino acid residues 1 to 556 of **FIG. 94** (SEQ ID NO:259), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0405] In another embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO307 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO307 polypeptide having amino acid residues 1 to 383 of **FIG. 96** (SEQ ID NO:261), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0406] In another embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO343 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO343 polypeptide having amino acid residues 1 to 317 of **FIG. 98** (SEQ ID NO:263), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0407] In another embodiment, the invention provides isolated PRO302 polypeptide. In particular, the invention provides isolated native sequence PRO302 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 452 of **FIG. 90** (SEQ ID NO:255).

[0408] In another embodiment, the invention provides isolated PRO303 polypeptide. In particular, the invention provides isolated native sequence PRO303 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 314 of **FIG. 92** (SEQ ID NO:257).

[0409] In another embodiment, the invention provides isolated PRO304 polypeptide. In particular, the invention provides isolated native sequence PRO304 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 556 of **FIG. 94** (SEQ ID NO:259).

[0410] In another embodiment, the invention provides isolated PRO307 polypeptide. In particular, the invention provides isolated native sequence PRO307 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 383 of **FIG. 96** (SEQ ID NO:261).

[0411] In another embodiment, the invention provides isolated PRO343 polypeptide. In particular, the invention provides isolated native sequence PRO343 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 317 of **FIG. 98** (SEQ ID NO:263).

[0412] 41. PRO328

[0413] Applicants have identified a cDNA clone that encodes a novel polypeptide, wherein the polypeptide is designated in the present application as "PRO328".

[0414] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO328 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO328 polypeptide having amino acid residues 1 to 463 of **FIG. 100** (SEQ ID NO:285), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0415] In another embodiment, the invention provides isolated PRO328 polypeptide. In particular, the invention provides isolated native sequence PRO328 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 463 of **FIG. 100** (SEQ ID NO:285). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO306 polypeptide.

[0416] 42. PRO335, PRO331 and PRO326

[0417] Applicants have identified three cDNA clones that respectively encode three novel polypeptides, each having leucine rich repeats and homology to LIG-1 and ALS. These polypeptides are designated in the present application as PRO335, PRO331 and PRO326, respectively.

[0418] In one embodiment, the invention provides three isolated nucleic acid molecules comprising DNA respectively encoding PRO335, PRO331 and PRO326, respectively. In one aspect, herein is provided an isolated nucleic acid comprising DNA encoding the PRO335 polypeptide having amino acid residues 1 through 1059 of **FIG. 102** (SEQ ID NO:290), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. Also provided herein is an isolated nucleic acid comprising DNA encoding the PRO331 polypeptide having amino acid residues 1 through 640 of **FIG. 104** (SEQ ID NO:292), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. Additionally provided herein is an isolated nucleic acid comprising DNA encoding the PRO326 polypeptide having amino acid residues 1 through 1119 of **FIG. 106** (SEQ ID NO:294), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0419] In another embodiment, the invention provides isolated PRO335, PRO331 and PRO326 polypeptides or extracellular domains thereof. In particular, the invention provides isolated native sequence for the PRO335 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 through 1059 of **FIG. 102** (SEQ ID NO:290). Also provided herein is the isolated native sequence for the PRO331 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 through 640 of **FIG. 104** (SEQ ID NO:292). Also provided herein is the isolated native sequence for the PRO326 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 through 1119 of **FIG. 106** (SEQ ID NO:294).

[0420] 43. PRO332

[0421] Applicants have identified a cDNA clone (DNA40982-1235) that encodes a novel polypeptide, designated in the present application as "PRO332."

[0422] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA having at least about 80% sequence identity to (a) a DNA molecule encoding a PRO358 polypeptide comprising the sequence of amino acids 49 to 642 of **FIG. 108** (SEQ ID NO:310), or (b) the complement of the DNA molecule of (a). The sequence identity preferably is about 85%, more preferably about 90%, most preferably about 95%. In one aspect, the isolated nucleic acid has at least about 80%, preferably at least about 85%, more preferably at least about 90%, and most preferably at least about 95% sequence identity with a polypeptide having amino acid residues 1 to 642 of **FIG. 108** (SEQ ID NO:310). Preferably, the highest degree of sequence identity occurs within the leucine-rich repeat domains (amino acids 116 to 624 of **FIG. 108**, SEQ ID NO:310). In a further embodiment, the isolated nucleic acid molecule comprises DNA encoding a PRO332 polypeptide having amino acid residues 49 to 642 of **FIG. 108** (SEQ ID NO:310), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0423] In another embodiment, the invention provides isolated PRO332 polypeptides. In particular, the invention provides isolated native sequence PRO332 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 49 to 624 of **FIG. 108** (SEQ ID NO:310). Native PRO332 polypeptides with or without the native signal sequence (amino acids 1 to 48 in **FIG. 108**, SEQ ID NO:310), and with or without the initiating methionine are specifically included.

[0424] 44. PRO334

[0425] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to fibulin and fibrillin, wherein the polypeptide is designated in the present application as "PRO334".

[0426] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO334 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO334 polypeptide having amino acid residues 1 to 509 of **FIG. 110** (SEQ ID NO:315), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0427] In another embodiment, the invention provides isolated PRO334 polypeptide. In particular, the invention provides isolated native sequence PRO334 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 509 of **FIG. 110** (SEQ ID NO:315).

[0428] 45. PRO346

[0429] Applicants have identified a cDNA clone (DNA44167-1243) that encodes a novel polypeptide, designated in the present application as "PRO346."

[0430] In one embodiment, the invention provides an isolated nucleic acid molecule having at least about 80% sequence identity to (a) a DNA molecule encoding a PRO346 polypeptide comprising the sequence of amino acids 19 to 339 of **FIG. 112** (SEQ ID NO: 320), or (b) the complement of the DNA molecule of (a). The sequence identity preferably is about 85%, more preferably about 90%, most preferably about 95%. In one aspect, the isolated

nucleic acid has at least about 80%, preferably at least about 85%, more preferably at least about 90%, and most preferably at least about 95% sequence identity with a polypeptide having amino acid residues 19 to 339 of **FIG. 112** (SEQ ID NO:320). Preferably, the highest degree of sequence identity occurs within the extracellular domains (amino acids 19 to 339 of **FIG. 112**, SEQ ID NO:320). In alternative embodiments, the polypeptide by which the homology is measured comprises the residues 1-339, 19-360 or 19-450 of **FIG. 112**, SEQ ID NO:320). In a further embodiment, the isolated nucleic acid molecule comprises DNA encoding a PRO346 polypeptide having amino acid residues 19 to 339 of **FIG. 112** (SEQ ID NO:320), alternatively residues 1-339, 19-360 or 19-450 of **FIG. 112** (SEQ ID NO:320) or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the invention provides a nucleic acid of the full length protein of clone DNA44167-1243, deposited with the ATCC under accession number ATCC 209434, alternatively the coding sequence of clone DNA44167-1243, deposited under accession number ATCC 209434.

[0431] In yet another embodiment, the invention provides isolated PRO346 polypeptide. In particular, the invention provides isolated native sequence PRO346 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 19 to 339 of **FIG. 112** (SEQ ID NO:320). Native PRO346 polypeptides with or without the native signal sequence (residues 1 to 18 in **FIG. 112** (SEQ ID NO:320), with or without the initiating methionine, with or without the transmembrane domain (residues 340 to 360) and with or without the intracellular domain (residues 361 to 450) are specifically included. Alternatively, the invention provides a PRO346 polypeptide encoded by the nucleic acid deposited under accession number ATCC 209434.

[0432] 46. PRO268

[0433] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to protein disulfide isomerase, wherein the polypeptide is designated in the present application as "PRO268".

[0434] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO268 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO268 polypeptide having amino acid residues 1 to 280 of **FIG. 114** (SEQ ID NO:325), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0435] In another embodiment, the invention provides isolated PRO268 polypeptide. In particular, the invention provides isolated native sequence PRO268 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 280 of **FIG. 114** (SEQ ID NO:325). An additional embodiment of the present invention is directed to an isolated extracellular domain of a PRO268 polypeptide.

[0436] 47. PRO330

[0437] Applicants have identified a cDNA clone that encodes a novel polypeptide having homology to the alpha subunit of prolyl 4-hydroxylase, wherein the polypeptide is designated in the present application as "PRO330".

[0438] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO330 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO330 polypeptide having amino acid residues 1 to 533 of FIG. 116 (SEQ ID NO:332), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0439] In another embodiment, the invention provides isolated PRO330 polypeptide. In particular, the invention provides isolated native sequence PRO330 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 533 of FIG. 116 (SEQ ID NO:332).

[0440] 48. PRO339 and PRO310

[0441] Applicants have identified two cDNA clones wherein each clone encodes a novel polypeptide having homology to fringe, wherein the polypeptides are designated in the present application as "PRO339" and "PRO310".

[0442] In one embodiment, the invention provides isolated nucleic acid molecules comprising DNA encoding a PRO339 and/or a PRO310 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding the PRO339 polypeptide having amino acid residues 1 to 772 of FIG. 118 (SEQ ID NO:339), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions. In another aspect, the isolated nucleic acid comprises DNA encoding the PRO310 polypeptide having amino acid residues 1 to 318 of FIG. 120 (SEQ ID NO:341), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0443] In another embodiment, the invention provides isolated PRO339 as well as isolated PRO310 polypeptides. In particular, the invention provides isolated native sequence PRO339 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 772 of FIG. 118 (SEQ ID NO:339). The invention further provides isolated native sequence PRO310 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 318 of FIG. 120 (SEQ ID NO:341).

[0444] 49. PRO244

[0445] Applicants have identified a cDNA clone that encodes a novel polypeptide, designated in the present application as "PRO244".

[0446] In one embodiment, the invention provides an isolated nucleic acid molecule comprising DNA encoding PRO244 polypeptide. In one aspect, the isolated nucleic acid comprises DNA encoding PRO244 polypeptide having amino acid residues 1 to 219 of FIG. 122 (SEQ ID NO:377), or is complementary to such encoding nucleic acid sequence, and remains stably bound to it under at least moderate, and optionally, under high stringency conditions.

[0447] In another embodiment, the invention provides isolated PRO244 polypeptide. In particular, the invention provides isolated native sequence PRO244 polypeptide, which in one embodiment, includes an amino acid sequence comprising residues 1 to 219 of FIG. 122 (SEQ ID NO:377).

[0448] 50. Additional Embodiments

[0449] In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

[0450] In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

[0451] In another embodiment, the invention provides an antibody which specifically binds to any of the above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

[0452] In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences, wherein those probes may be derived from any of the above or below described nucleotide sequences.

[0453] In other embodiments, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

[0454] In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0455] In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity,

tity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein or the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0456] In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0457] Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

[0458] Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes or for encoding fragments of a PRO polypeptide that

may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody. Such nucleic acid fragments are usually at least about 20 nucleotides in length, preferably at least about 30 nucleotides in length, more preferably at least about 40 nucleotides in length, yet more preferably at least about 50 nucleotides in length, yet more preferably at least about 60 nucleotides in length, yet more preferably at least about 70 nucleotides in length, yet more preferably at least about 80 nucleotides in length, yet more preferably at least about 90 nucleotides in length, yet more preferably at least about 100 nucleotides in length, yet more preferably at least about 110 nucleotides in length, yet more preferably at least about 120 nucleotides in length, yet more preferably at least about 130 nucleotides in length, yet more preferably at least about 140 nucleotides in length, yet more preferably at least about 150 nucleotides in length, yet more preferably at least about 160 nucleotides in length, yet more preferably at least about 170 nucleotides in length, yet more preferably at least about 180 nucleotides in length, yet more preferably at least about 190 nucleotides in length, yet more preferably at least about 200 nucleotides in length, yet more preferably at least about 250 nucleotides in length, yet more preferably at least about 300 nucleotides in length, yet more preferably at least about 350 nucleotides in length, yet more preferably at least about 400 nucleotides in length, yet more preferably at least about 450 nucleotides in length, yet more preferably at least about 500 nucleotides in length, yet more preferably at least about 600 nucleotides in length, yet more preferably at least about 700 nucleotides in length, yet more preferably at least about 800 nucleotides in length, yet more preferably at least about 900 nucleotides in length and yet more preferably at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

[0459] In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

[0460] In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more

preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein.

[0461] In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% sequence identity, preferably at least about 81% sequence identity, more preferably at least about 82% sequence identity, yet more preferably at least about 83% sequence identity, yet more preferably at least about 84% sequence identity, yet more preferably at least about 85% sequence identity, yet more preferably at least about 86% sequence identity, yet more preferably at least about 87% sequence identity, yet more preferably at least about 88% sequence identity, yet more preferably at least about 89% sequence identity, yet more preferably at least about 90% sequence identity, yet more preferably at least about 91% sequence identity, yet more preferably at least about 92% sequence identity, yet more preferably at least about 93% sequence identity, yet more preferably at least about 94% sequence identity, yet more preferably at least about 95% sequence identity, yet more preferably at least about 96% sequence identity, yet more preferably at least about 97% sequence identity, yet more preferably at least about 98% sequence identity and yet more preferably at least about 99% sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

[0462] In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence scoring at least about 80% positives, preferably at least about 81% positives, more preferably at least about 82% positives, yet more preferably at least about 83% positives, yet more preferably at least about 84% positives, yet more preferably at least about 85% positives, yet more preferably at least about 86% positives, yet more preferably at least about 87% positives, yet more preferably at least about 88% positives, yet more preferably at least about 89% positives, yet more preferably at least about 90% positives, yet more preferably at least about 91% positives, yet more preferably at least about 92% positives, yet more preferably at least about 93% positives, yet more preferably at least about 94% positives, yet more preferably at least about 95% positives, yet more preferably at least about 96% positives, yet more preferably at least about 97% positives, yet more preferably at least about 98% positives and yet more preferably at least about 99% positives when compared with the amino acid sequence of a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein or an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein.

[0463] In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by

a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0464] Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0465] In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

[0466] In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

[0467] In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

[0468] Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0469] FIG. 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO211 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA32292-1131".

[0470] FIG. 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in FIG. 1.

[0471] FIG. 3 shows a nucleotide sequence (SEQ ID NO:3) of a native sequence PRO217 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA33094-1131".

[0472] FIG. 4 shows the amino acid sequence (SEQ ID NO:4) derived from the coding sequence of SEQ ID NO:3 shown in FIG. 3.

[0473] FIG. 5 shows a nucleotide sequence (SEQ ID NO:11) of a native sequence PRO230 cDNA, wherein SEQ ID NO:11 is a clone designated herein as "DNA33223-1136".

- [0474] FIG. 6 shows the amino acid sequence (SEQ ID NO:12) derived from the coding sequence of SEQ ID NO:11 shown in FIG. 5.
- [0475] FIG. 7 shows a nucleotide sequence designated herein as DNA20088 (SEQ ID NO:13).
- [0476] FIG. 8 shows a nucleotide sequence (SEQ ID NO:17) of a native sequence PRO232 cDNA, wherein SEQ ID NO:17 is a clone designated herein as "DNA34435-1140".
- [0477] FIG. 9 shows the amino acid sequence (SEQ ID NO:18) derived from the coding sequence of SEQ ID NO:17 shown in FIG. 8.
- [0478] FIG. 10 shows a nucleotide sequence (SEQ ID NO:22) of a native sequence PRO187 cDNA, wherein SEQ ID NO:22 is a clone designated herein as "DNA27864-1155".
- [0479] FIG. 11 shows the amino acid sequence (SEQ ID NO:23) derived from the coding sequence of SEQ ID NO:22 shown in FIG. 10.
- [0480] FIG. 12 shows a nucleotide sequence (SEQ ID NO:27) of a native sequence PRO265 cDNA, wherein SEQ ID NO:27 is a clone designated herein as "DNA36350-1158".
- [0481] FIG. 13 shows the amino acid sequence (SEQ ID NO:28) derived from the coding sequence of SEQ ID NO:27 shown in FIG. 12.
- [0482] FIG. 14 shows a nucleotide sequence (SEQ ID NO:33) of a native sequence PRO219 cDNA, wherein SEQ ID NO:33 is a clone designated herein as "DNA32290-1164".
- [0483] FIG. 15 shows the amino acid sequence (SEQ ID NO:34) derived from the coding sequence of SEQ ID NO:33 shown in FIG. 14.
- [0484] FIG. 16 shows a nucleotide sequence (SEQ ID NO:38) of a native sequence PRO246 cDNA, wherein SEQ ID NO:38 is a clone designated herein as "DNA35639-1172".
- [0485] FIG. 17 shows the amino acid sequence (SEQ ID NO:39) derived from the coding sequence of SEQ ID NO:38 shown in FIG. 16.
- [0486] FIG. 18 shows a nucleotide sequence (SEQ ID NO:48) of a native sequence PRO228 cDNA, wherein SEQ ID NO:48 is a clone designated herein as "DNA33092-1202".
- [0487] FIG. 19 shows the amino acid sequence (SEQ ID NO:49) derived from the coding sequence of SEQ ID NO:48 shown in FIG. 18.
- [0488] FIG. 20 shows a nucleotide sequence designated herein as DNA21951 (SEQ ID NO:50).
- [0489] FIG. 21 shows a nucleotide sequence (SEQ ID NO:58) of a native sequence PRO533 cDNA, wherein SEQ ID NO:58 is a clone designated herein as "DNA49435-1219".
- [0490] FIG. 22 shows the amino acid sequence (SEQ ID NO:59) derived from the coding sequence of SEQ ID NO:58 shown in FIG. 21.
- [0491] FIG. 23 shows a nucleotide sequence (SEQ ID NO:63) of a native sequence PRO245 cDNA, wherein SEQ ID NO:63 is a clone designated herein as "DNA35638-1141".
- [0492] FIG. 24 shows the amino acid sequence (SEQ ID NO:64) derived from the coding sequence of SEQ ID NO:63 shown in FIG. 23.
- [0493] FIG. 25 shows a nucleotide sequence (SEQ ID NO:68) of a native sequence PRO220 cDNA, wherein SEQ ID NO:68 is a clone designated herein as "DNA32298-1132".
- [0494] FIG. 26 shows the amino acid sequence (SEQ ID NO:69) derived from the coding sequence of SEQ ID NO:68 shown in FIG. 25.
- [0495] FIG. 27 shows a nucleotide sequence (SEQ ID NO:70) of a native sequence PRO221 cDNA, wherein SEQ ID NO:70 is a clone designated herein as "DNA33089-1132".
- [0496] FIG. 28 shows the amino acid sequence (SEQ ID NO:71) derived from the coding sequence of SEQ ID NO:70 shown in FIG. 27.
- [0497] FIG. 29 shows a nucleotide sequence (SEQ ID NO:72) of a native sequence PRO227 cDNA, wherein SEQ ID NO:72 is a clone designated herein as "DNA33786-1132".
- [0498] FIG. 30 shows the amino acid sequence (SEQ ID NO:73) derived from the coding sequence of SEQ ID NO:72 shown in FIG. 29.
- [0499] FIG. 31 shows a nucleotide sequence (SEQ ID NO:83) of a native sequence PRO258 cDNA, wherein SEQ ID NO:83 is a clone designated herein as "DNA35918-1174".
- [0500] FIG. 32 shows the amino acid sequence (SEQ ID NO:84) derived from the coding sequence of SEQ ID NO:83 shown in FIG. 31.
- [0501] FIG. 33 shows a nucleotide sequence (SEQ ID NO:90) of a native sequence PRO266 cDNA, wherein SEQ ID NO:90 is a clone designated herein as "DNA37150-1178".
- [0502] FIG. 34 shows the amino acid sequence (SEQ ID NO:91) derived from the coding sequence of SEQ ID NO:90 shown in FIG. 33.
- [0503] FIG. 35 shows a nucleotide sequence (SEQ ID NO:95) of a native sequence PRO269 cDNA, wherein SEQ ID NO:95 is a clone designated herein as "DNA38260-1180".
- [0504] FIG. 36 shows the amino acid sequence (SEQ ID NO:96) derived from the coding sequence of SEQ ID NO:95 shown in FIG. 35.
- [0505] FIG. 37 shows a nucleotide sequence (SEQ ID NO:103) of a native sequence PRO287 cDNA, wherein SEQ ID NO:103 is a clone designated herein as "DNA39969-1185".
- [0506] FIG. 38 shows the amino acid sequence (SEQ ID NO:104) derived from the coding sequence of SEQ ID NO:103 shown in FIG. 37.

- [0507] FIG. 39 shows a nucleotide sequence (SEQ ID NO:108) of a native sequence PRO214 cDNA, wherein SEQ ID NO:108 is a clone designated herein as "DNA32286-1191".
- [0508] FIG. 40 shows the amino acid sequence (SEQ ID NO:109) derived from the coding sequence of SEQ ID NO:108 shown in FIG. 39.
- [0509] FIG. 41 shows a nucleotide sequence (SEQ ID NO:113) of a native sequence PRO317 cDNA, wherein SEQ ID NO:113 is a clone designated herein as "DNA33461-1199".
- [0510] FIG. 42 shows the amino acid sequence (SEQ ID NO:114) derived from the coding sequence of SEQ ID NO:113 shown in FIG. 41.
- [0511] FIG. 43 shows a nucleotide sequence (SEQ ID NO:118) of a native sequence PRO301 cDNA, wherein SEQ ID NO:118 is a clone designated herein as "DNA40628-1216".
- [0512] FIG. 44 shows the amino acid sequence (SEQ ID NO:119) derived from the coding sequence of SEQ ID NO:118 shown in FIG. 43.
- [0513] FIG. 45 shows a nucleotide sequence (SEQ ID NO:126) of a native sequence PRO224 cDNA, wherein SEQ ID NO:126 is a clone designated herein as "DNA33221-1133".
- [0514] FIG. 46 shows the amino acid sequence (SEQ ID NO:127) derived from the coding sequence of SEQ ID NO:126 shown in FIG. 45.
- [0515] FIG. 47 shows a nucleotide sequence (SEQ ID NO:131) of a native sequence PRO222 cDNA, wherein SEQ ID NO:131 is a clone designated herein as "DNA33107-1135".
- [0516] FIG. 48 shows the amino acid sequence (SEQ ID NO:132) derived from the coding sequence of SEQ ID NO:131 shown in FIG. 47.
- [0517] FIG. 49 shows a nucleotide sequence (SEQ ID NO:136) of a native sequence PRO234 cDNA, wherein SEQ ID NO:136 is a clone designated herein as "DNA35557-1137".
- [0518] FIG. 50 shows the amino acid sequence (SEQ ID NO:137) derived from the coding sequence of SEQ ID NO:136 shown in FIG. 49.
- [0519] FIG. 51 shows a nucleotide sequence (SEQ ID NO:141) of a native sequence PRO231 cDNA, wherein SEQ ID NO:141 is a clone designated herein as "DNA34434-1139".
- [0520] FIG. 52 shows the amino acid sequence (SEQ ID NO:142) derived from the coding sequence of SEQ ID NO:141 shown in FIG. 51.
- [0521] FIG. 53 shows a nucleotide sequence (SEQ ID NO:147) of a native sequence PRO229 cDNA, wherein SEQ ID NO:147 is a clone designated herein as "DNA33100-1159".
- [0522] FIG. 54 shows the amino acid sequence (SEQ ID NO:148) derived from the coding sequence of SEQ ID NO:147 shown in FIG. 53.
- [0523] FIG. 55 shows a nucleotide sequence (SEQ ID NO:152) of a native sequence PRO238 cDNA, wherein SEQ ID NO:152 is a clone designated herein as "DNA35600-1162".
- [0524] FIG. 56 shows the amino acid sequence (SEQ ID NO:153) derived from the coding sequence of SEQ ID NO:152 shown in FIG. 55.
- [0525] FIG. 57 shows a nucleotide sequence (SEQ ID NO:158) of a native sequence PRO233 cDNA, wherein SEQ ID NO:158 is a clone designated herein as "DNA34436-1238".
- [0526] FIG. 58 shows the amino acid sequence (SEQ ID NO:159) derived from the coding sequence of SEQ ID NO:158 shown in FIG. 57.
- [0527] FIG. 59 shows a nucleotide sequence (SEQ ID NO:163) of a native sequence PRO223 cDNA, wherein SEQ ID NO:163 is a clone designated herein as "DNA33206-1165".
- [0528] FIG. 60 shows the amino acid sequence (SEQ ID NO:164) derived from the coding sequence of SEQ ID NO:163 shown in FIG. 59.
- [0529] FIG. 61 shows a nucleotide sequence (SEQ ID NO:169) of a native sequence PRO235 cDNA, wherein SEQ ID NO:169 is a clone designated herein as "DNA35558-1167".
- [0530] FIG. 62 shows the amino acid sequence (SEQ ID NO:170) derived from the coding sequence of SEQ ID NO:169 shown in FIG. 61.
- [0531] FIG. 63 shows a nucleotide sequence (SEQ ID NO:174) of a native sequence PRO236 cDNA, wherein SEQ ID NO:174 is a clone designated herein as "DNA35599-1168".
- [0532] FIG. 64 shows the amino acid sequence (SEQ ID NO:175) derived from the coding sequence of SEQ ID NO:174 shown in FIG. 63.
- [0533] FIG. 65 shows a nucleotide sequence (SEQ ID NO:176) of a native sequence PRO262 cDNA, wherein SEQ ID NO:176 is a clone designated herein as "DNA36992-1168".
- [0534] FIG. 66 shows the amino acid sequence (SEQ ID NO:177) derived from the coding sequence of SEQ ID NO:176 shown in FIG. 65.
- [0535] FIG. 67 shows a nucleotide sequence (SEQ ID NO:184) of a native sequence PRO239 cDNA, wherein SEQ ID NO:184 is a clone designated herein as "DNA34407-1169".
- [0536] FIG. 68 shows the amino acid sequence (SEQ ID NO:185) derived from the coding sequence of SEQ ID NO:184 shown in FIG. 67.
- [0537] FIG. 69 shows a nucleotide sequence (SEQ ID NO:189) of a native sequence PRO257 cDNA, wherein SEQ ID NO:189 is a clone designated herein as "DNA35841-1173".
- [0538] FIG. 70 shows the amino acid sequence (SEQ ID NO:190) derived from the coding sequence of SEQ ID NO:189 shown in FIG. 69.

- [0539] FIG. 71 shows a nucleotide sequence (SEQ ID NO:194) of a native sequence PRO260 cDNA, wherein SEQ ID NO:194 is a clone designated herein as "DNA33470-1175".
- [0540] FIG. 72 shows the amino acid sequence (SEQ ID NO:195) derived from the coding sequence of SEQ ID NO:194 shown in FIG. 71.
- [0541] FIG. 73 shows a nucleotide sequence (SEQ ID NO:200) of a native sequence PRO263 cDNA, wherein SEQ ID NO:200 is a clone designated herein as "DNA34431-1177".
- [0542] FIG. 74 shows the amino acid sequence (SEQ ID NO:201) derived from the coding sequence of SEQ ID NO:200 shown in FIG. 73.
- [0543] FIG. 75 shows a nucleotide sequence (SEQ ID NO:206) of a native sequence PRO270 cDNA, wherein SEQ ID NO:206 is a clone designated herein as "DNA39510-1181".
- [0544] FIG. 76 shows the amino acid sequence (SEQ ID NO:207) derived from the coding sequence of SEQ ID NO:206 shown in FIG. 75.
- [0545] FIG. 77 shows a nucleotide sequence (SEQ ID NO:212) of a native sequence PRO271 cDNA, wherein SEQ ID NO:212 is a clone designated herein as "DNA39423-1182".
- [0546] FIG. 78 shows the amino acid sequence (SEQ ID NO:213) derived from the coding sequence of SEQ ID NO:212 shown in FIG. 77.
- [0547] FIG. 79 shows a nucleotide sequence (SEQ ID NO:220) of a native sequence PRO272 cDNA, wherein SEQ ID NO:220 is a clone designated herein as "DNA40620-1183".
- [0548] FIG. 80 shows the amino acid sequence (SEQ ID NO:221) derived from the coding sequence of SEQ ID NO:220 shown in FIG. 79.
- [0549] FIG. 81 shows a nucleotide sequence (SEQ ID NO:226) of a native sequence PRO294 cDNA, wherein SEQ ID NO:226 is a clone designated herein as "DNA40604-1187".
- [0550] FIG. 82 shows the amino acid sequence (SEQ ID NO:227) derived from the coding sequence of SEQ ID NO:226 shown in FIG. 81.
- [0551] FIG. 83 shows a nucleotide sequence (SEQ ID NO:235) of a native sequence PRO295 cDNA, wherein SEQ ID NO:235 is a clone designated herein as "DNA38268-1188".
- [0552] FIG. 84 shows the amino acid sequence (SEQ ID NO:236) derived from the coding sequence of SEQ ID NO:235 shown in FIG. 83.
- [0553] FIG. 85 shows a nucleotide sequence (SEQ ID NO:244) of a native sequence PRO293 cDNA, wherein SEQ ID NO:244 is a clone designated herein as "DNA37151-1193".
- [0554] FIG. 86 shows the amino acid sequence (SEQ ID NO:245) derived from the coding sequence of SEQ ID NO:244 shown in FIG. 85.
- [0555] FIG. 87 shows a nucleotide sequence (SEQ ID NO:249) of a native sequence PRO247 cDNA, wherein SEQ ID NO:249 is a clone designated herein as "DNA35673-1201".
- [0556] FIG. 88 shows the amino acid sequence (SEQ ID NO:250) derived from the coding sequence of SEQ ID NO:249 shown in FIG. 87.
- [0557] FIG. 89 shows a nucleotide sequence (SEQ ID NO:254) of a native sequence PRO302 cDNA, wherein SEQ ID NO:254 is a clone designated herein as "DNA40370-1217".
- [0558] FIG. 90 shows the amino acid sequence (SEQ ID NO:255) derived from the coding sequence of SEQ ID NO:254 shown in FIG. 89.
- [0559] FIG. 91 shows a nucleotide sequence (SEQ ID NO:256) of a native sequence PRO303 cDNA, wherein SEQ ID NO:256 is a clone designated herein as "DNA42551-1217".
- [0560] FIG. 92 shows the amino acid sequence (SEQ ID NO:257) derived from the coding sequence of SEQ ID NO:256 shown in FIG. 91.
- [0561] FIG. 93 shows a nucleotide sequence (SEQ ID NO:258) of a native sequence PRO304 cDNA, wherein SEQ ID NO:258 is a clone designated herein as "DNA39520-1217".
- [0562] FIG. 94 shows the amino acid sequence (SEQ ID NO:259) derived from the coding sequence of SEQ ID NO:258 shown in FIG. 93.
- [0563] FIG. 95 shows a nucleotide sequence (SEQ ID NO:260) of a native sequence PRO307 cDNA, wherein SEQ ID NO:260 is a clone designated herein as "DNA41225-1217".
- [0564] FIG. 96 shows the amino acid sequence (SEQ ID NO:261) derived from the coding sequence of SEQ ID NO:260 shown in FIG. 95.
- [0565] FIG. 97 shows a nucleotide sequence (SEQ ID NO:262) of a native sequence PRO343 cDNA, wherein SEQ ID NO:262 is a clone designated herein as "DNA43318-1217".
- [0566] FIG. 98 shows the amino acid sequence (SEQ ID NO:263) derived from the coding sequence of SEQ ID NO:262 shown in FIG. 97.
- [0567] FIG. 99 shows a nucleotide sequence (SEQ ID NO:284) of a native sequence PRO328 cDNA, wherein SEQ ID NO:284 is a clone designated herein as "DNA40587-1231".
- [0568] FIG. 100 shows the amino acid sequence (SEQ ID NO:285) derived from the coding sequence of SEQ ID NO:284 shown in FIG. 99.
- [0569] FIG. 101 shows a nucleotide sequence (SEQ ID NO:289) of a native sequence PRO335 cDNA, wherein SEQ ID NO:289 is a clone designated herein as "DNA41388-1234".
- [0570] FIG. 102 shows the amino acid sequence (SEQ ID NO:290) derived from the coding sequence of SEQ ID NO:289 shown in FIG. 101.



[0571] FIG. 103 shows a nucleotide sequence (SEQ ID NO:291) of a native sequence PRO331 cDNA, wherein SEQ ID NO:291 is a clone designated herein as "DNA40981-1234".

[0572] FIG. 104 shows the amino acid sequence (SEQ ID NO:292) derived from the coding sequence of SEQ ID NO:291 shown in FIG. 103.

[0573] FIG. 105 shows a nucleotide sequence (SEQ ID NO:293) of a native sequence PRO326 cDNA, wherein SEQ ID NO:293 is a clone designated herein as "DNA37140-1234".

[0574] FIG. 106 shows the amino acid sequence (SEQ ID NO:294) derived from the coding sequence of SEQ ID NO:293 shown in FIG. 105.

[0575] FIG. 107 shows a nucleotide sequence (SEQ ID NO:309) of a native sequence PRO332 cDNA, wherein SEQ ID NO:309 is a clone designated herein as "DNA40982-1235".

[0576] FIG. 108 shows the amino acid sequence (SEQ ID NO:310) derived from the coding sequence of SEQ ID NO:309 shown in FIG. 107.

[0577] FIG. 109 shows a nucleotide sequence (SEQ ID NO:314) of a native sequence PRO334 cDNA, wherein SEQ ID NO:314 is a clone designated herein as "DNA41379-1236".

[0578] FIG. 110 shows the amino acid sequence (SEQ ID NO:315) derived from the coding sequence of SEQ ID NO:314 shown in FIG. 109.

[0579] FIG. 111 shows a nucleotide sequence (SEQ ID NO:319) of a native sequence PRO346 cDNA, wherein SEQ ID NO:319 is a clone designated herein as "DNA44167-1243".

[0580] FIG. 112 shows the amino acid sequence (SEQ ID NO:320) derived from the coding sequence of SEQ ID NO:319 shown in FIG. 111.

[0581] FIG. 113 shows a nucleotide sequence (SEQ ID NO:324) of a native sequence PRO268 cDNA, wherein SEQ ID NO:324 is a clone designated herein as "DNA39427-1179".

[0582] FIG. 114 shows the amino acid sequence (SEQ ID NO:325) derived from the coding sequence of SEQ ID NO:324 shown in FIG. 113.

[0583] FIG. 115 shows a nucleotide sequence (SEQ ID NO:331) of a native sequence PRO330 cDNA, wherein SEQ ID NO:331 is a clone designated herein as "DNA40603-1232".

[0584] FIG. 116 shows the amino acid sequence (SEQ ID NO:332) derived from the coding sequence of SEQ ID NO:331 shown in FIG. 115.

[0585] FIG. 117 shows a nucleotide sequence (SEQ ID NO:338) of a native sequence PRO339 cDNA, wherein SEQ ID NO:338 is a clone designated herein as "DNA43466-1225".

[0586] FIG. 118 shows the amino acid sequence (SEQ ID NO:339) derived from the coding sequence of SEQ ID NO:338 shown in FIG. 117.

[0587] FIG. 119 shows a nucleotide sequence (SEQ ID NO:340) of a native sequence PRO310 cDNA, wherein SEQ ID NO:340 is a clone designated herein as "DNA43046-1225".

[0588] FIG. 120 shows the amino acid sequence (SEQ ID NO:341) derived from the coding sequence of SEQ ID NO:340 shown in FIG. 119.

[0589] FIG. 121 shows a nucleotide sequence (SEQ ID NO:376) of a native sequence PRO244 cDNA, wherein SEQ ID NO:376 is a clone designated herein as "DNA35668-1171".

[0590] FIG. 122 shows the amino acid sequence (SEQ ID NO:377) derived from the coding sequence of SEQ ID NO:376 shown in FIG. 121.

[0591] FIG. 123 shows a nucleotide sequence (SEQ ID NO:422) of a native sequence PRO1868 cDNA, wherein SEQ ID NO:422 is a clone designated herein as "DNA77624-2515".

[0592] FIG. 124 shows the amino acid sequence (SEQ ID NO:423) derived from the coding sequence of SEQ ID NO:422 shown in FIG. 123.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

##### [0593] I. Definitions

[0594] The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods.

[0595] A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream

from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

[0596] The PRO polypeptide “extracellular domain” or “ECD” refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

[0597] The approximate location of the “signal peptides” of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

[0598] “PRO polypeptide variant” means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, preferably at least about 81% amino acid sequence identity, more preferably at least about 82% amino acid sequence identity, more preferably at least about 83% amino acid sequence identity, more preferably at least about 84% amino acid sequence identity, more preferably at least about 85% amino acid sequence identity, more preferably at least about 86% amino

acid sequence identity, more preferably at least about 87% amino acid sequence identity, more preferably at least about 88% amino acid sequence identity, more preferably at least about 89% amino acid sequence identity, more preferably at least about 90% amino acid sequence identity, more preferably at least about 91% amino acid sequence identity, more preferably at least about 92% amino acid sequence identity, more preferably at least about 93% amino acid sequence identity, more preferably at least about 94% amino acid sequence identity, more preferably at least about 95% amino acid sequence identity, more preferably at least about 96% amino acid sequence identity, more preferably at least about 97% amino acid sequence identity, more preferably at least about 98% amino acid sequence identity and most preferably at least about 99% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, often at least about 20 amino acids in length, more often at least about 30 amino acids in length, more often at least about 40 amino acids in length, more often at least about 50 amino acids in length, more often at least about 60 amino acids in length, more often at least about 70 amino acids in length, more often at least about 80 amino acids in length, more often at least about 90 amino acids in length, more often at least about 100 amino acids in length, more often at least about 150 amino acids in length, more often at least about 200 amino acids in length, more often at least about 300 amino acids in length, or more.

[0599] “Percent (%) amino acid sequence identity” with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating

system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

**[0600]** In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

**[0601]** where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X," "Y" and "Z" each represent different hypothetical amino acid residues.

**[0602]** Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

**[0603]** Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402

(1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

**[0604]** In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

**[0605]** where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

**[0606]** "PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, more preferably at least about 81% nucleic acid sequence identity, more preferably at least about 82% nucleic acid sequence identity, more preferably at least about 83% nucleic acid sequence identity, more preferably at least about 84% nucleic acid sequence identity, more preferably at least about 85% nucleic acid sequence identity, more preferably at least about 86% nucleic acid sequence identity, more preferably at least about 87% nucleic acid sequence identity, more preferably at least about 88% nucleic acid sequence identity, more preferably at least about 89% nucleic acid sequence identity, more preferably at least about 90% nucleic acid sequence identity, more preferably at least about 91% nucleic acid sequence identity, more preferably at least about 92% nucleic acid sequence identity, more preferably at least about 93% nucleic acid sequence identity, more preferably at least about 94% nucleic acid sequence identity, more preferably at least about 95% nucleic acid sequence identity, more preferably at least about 96% nucleic acid sequence identity, more preferably at least about 97% nucleic acid sequence identity, more preferably at least about 98% nucleic acid sequence identity and yet more preferably at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native

sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

[0607] Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, often at least about 60 nucleotides in length, more often at least about 90 nucleotides in length, more often at least about 120 nucleotides in length, more often at least about 150 nucleotides in length, more often at least about 180 nucleotides in length, more often at least about 210 nucleotides in length, more often at least about 240 nucleotides in length, more often at least about 270 nucleotides in length, more often at least about 300 nucleotides in length, more often at least about 450 nucleotides in length, more often at least about 600 nucleotides in length, more often at least about 900 nucleotides in length, or more.

[0608] "Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0609] In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

[0610] where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C

is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

[0611] Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

[0612] Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov>. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0613] In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that

has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

100 times the fraction  $W/Z$

[0614] where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

[0615] In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

[0616] The term "positives", in the context of sequence comparison performed as described above, includes residues in the sequences compared that are not identical but have similar properties (e.g. as a result of conservative substitutions, see Table 6 below). For purposes herein, the % value of positives is determined by dividing (a) the number of amino acid residues scoring a positive value between the PRO polypeptide amino acid sequence of interest having a sequence derived from the native PRO polypeptide sequence and the comparison amino acid sequence of interest (i.e., the amino acid sequence against which the PRO polypeptide sequence is being compared) as determined in the BLO-SUM62 matrix of WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest.

[0617] Unless specifically stated otherwise, the % value of positives is calculated as described in the immediately preceding paragraph. However, in the context of the amino acid sequence identity comparisons performed as described for ALIGN-2 and NCBI-BLAST-2 above, includes amino acid residues in the sequences compared that are not only identical, but also those that have similar properties. Amino acid residues that score a positive value to an amino acid residue of interest are those that are either identical to the amino acid residue of interest or are a preferred substitution (as defined in Table 6 below) of the amino acid residue of interest.

[0618] For amino acid sequence comparisons using ALIGN-2 or NCBI-BLAST2, the % value of positives of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % positives to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction  $X/Y$

[0619] where X is the number of amino acid residues scoring a positive value as defined above by the sequence alignment program ALIGN-2 or NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % positives of A to B will not equal the % positives of B to A.

[0620] "Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide in situ within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[0621] An "isolated" PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

[0622] The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0623] Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0624] The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-PRO monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polypeptopic specificity, single chain anti-PRO antibodies,

and fragments of anti-PRO antibodies (see below). The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

[0625] "Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

[0626] "Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C.

[0627] "Moderately stringent conditions" may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989., and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37-50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

[0628] The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an

antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

[0629] As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

[0630] "Active" or "activity" for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

[0631] The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

[0632] "Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

[0633] "Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activ-

ity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

[0634] "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

[0635] Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[0636] "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

[0637] "Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments; diabodies; linear antibodies (Zapata et al., *Protein Eng.* 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0638] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

[0639] "Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V<sub>H</sub>-V<sub>L</sub> dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0640] The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is

the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0641] The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

[0642] Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

[0643] "Single-chain Fv" or "sFv" antibody fragments comprise the V<sub>H</sub> and V<sub>L</sub> domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V<sub>H</sub> and V<sub>L</sub> domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

[0644] The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V<sub>H</sub>) connected to a light-chain variable domain (V<sub>L</sub>) in the same polypeptide chain (V<sub>H</sub>-V<sub>L</sub>). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

[0645] An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's, natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0646] The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

[0647] By “solid phase” is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Pat. No. 4,275,149.

[0648] A “liposome” is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

[0649] A “small molecule” is defined herein to have a molecular weight below about 500 Daltons.

[0650] “PRO317-associated disorder” refers to a pathological condition or disease wherein PRO317 is over- or

underexpressed. Such disorders include diseases of the female genital tract or of the endometrium of a mammal, including hyperplasia, endometritis, endometriosis, wherein the patient is at risk for infertility due to endometrial factor, endometrioma, and endometrial cancer, especially those diseases involving abnormal bleeding such as a gynecological disease. They also include diseases involving angiogenesis, wherein the angiogenesis results in a pathological condition, such as cancer involving solid tumors (the therapy for the disorder would result in decreased vascularization and a decline in growth and metastasis of a variety of tumors). Alternatively, the angiogenesis may be beneficial, such as for ischemia, especially coronary ischemia. Hence, these disorders include those found in patients whose hearts are functioning but who have a blocked blood supply due to atherosclerotic coronary artery disease, and those with a functioning but underperfused heart, including patients with coronary arterial disease who are not optimal candidates for angioplasty and coronary artery by-pass surgery. The disorders also include diseases involving the kidney or originating from the kidney tissue, such as polycystic kidney disease and chronic and acute renal failure.

TABLE 1

```

/*
 *
 * C—C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop—stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */
int _day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ {2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ {0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ {0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ {0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ {1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ {0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ {_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ {1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ {0, 1, -5, , 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ {1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ {1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ {0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ {0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4}
};
/*
 */
#include <stdio.h>
#include <ctype.h>
#define MAXJMP 16 /* max jumps in a diag */
#define MAXGAP 24 /* don't continue to penalize gaps larger than this */
#define JMPS 1024 /* max jmps in an path */

```



TABLE 1-continued

```

#define MX      4      /* save if there's at least MX-1 bases since last jmp */
#define DMAT    3      /* value of matching bases */
#define DMIS    0      /* penalty for mismatched bases */
#define DINS0   8      /* penalty for a gap */
#define DINS1   1      /* penalty per base */
#define PINS0   8      /* penalty for a gap */
#define PINS1   4      /* penalty per residue */
struct jmp {
    short      n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
};
/* limits seq to 216-1 */
struct diag {
    int      score; /* score at last jmp */
    long     offset; /* offset of prey block */
    short    jmp; /* current jmp index */
    struct jmp *jp; /* list of jmps */
};
struct path {
    int      spc; /* number of leading spaces */
    short    n[JMP]; /* size of jmp (gap) */
    int      x[JMP]; /* loc of jmp (last elem before gap) */
};
char      *ofile; /* output file name */
char      *name[2]; /* seq names: getseqs() */
char      *prog; /* prog name for err msgs */
char      *seq[2]; /* seqs: getseqs() */
int      dmax; /* best diag: nw() */
int      dmax0; /* final diag */
int      dna; /* set if dna: main() */
int      endgaps; /* set if penalizing end gaps */
int      gapx, gapy; /* total gaps in seqs */
int      len0, len1; /* seq lens */
int      ngapx, ngapy; /* total size of gaps */
int      smax; /* max score: nw() */
int      *xbm; /* bitmap for matching */
long     offset; /* current offset in jmp file */
struct    diag *dx; /* holds diagonals */
struct    path *pp[2]; /* holds path for seqs */
char      *calloc(), *malloc(), *index(), *strcpy();
char      *getseq(), *g_calloc();
/* Needleman-Wunsch alignment program
 *
 * usage: prog file1 file2
 * where file1 and file2 are two dna or two protein sequences.
 * The sequences can be in upper- or lower-case and may contain ambiguity
 * Any lines beginning with ';', '>' or '<' are ignored
 * Max file length is 65535 (limited by unsigned short x in the jmp struct)
 * A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
 * Output is in the file "align.out"
 *
 * The program may create a tmp file in /tmp to hold info about traceback.
 * Original version developed under BSD 4.3 on a vax 8650
 */
#include "nw.h"
#include "day.h"
static __dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
static __pbval[26] = {
    1, 2|(1<<('D'-'A'))|(1<<('N'-'A')), 4 ,8, 16, 32, 64
    128, 256, 0xFFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25|(1<<('E'-'A'))|(1<<('Q'-'A'))
};
main(ac, av)
int      ac;
char      *av[];
{
    prog = av[0];
    if (ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';', '>' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
}

```

TABLE 1-continued

```

}
namex[0] = av[1];
namex[1] = av[2];
seqx[0] = getseq(namex[0], &len0);
seqx[1] = getseq(namex[1], &len1);
xbm = (dna)? _dbval : _pbval;
endgaps = 0; /* 1 to penalize endgaps */
ofile = "align.out"; /* output file */
nw(); /* fill in the matrix, get the possible jmps */
readjmps(); /* get the actual jmps */
print(); /* print stats, alignment */
cleanup(0); /* unlink any tmp files */
}
/* do the alignment, return best score: main()
* dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
* pro: PAM 250 values
* When scores are equal, we prefer mismatches to any gap, prefer
* a new gap to extending an ongoing gap, and prefer a gap in seqx
* to a gap in seq y.
*/
nw()
{
char *px, *py; /* seqs and ptrs */
int *ndely, *dely; /* keep track of dely */
int ndelx, delx; /* keep track of delx */
int *tmp; /* for swapping row0, row1 */
int mis; /* score for each type */
int ins0, ins1; /* insertion penalties */
register id; /* diagonal index */
register ij; /* jmp index */
register *col0, *col1; /* score for curr, last row */
register xx, yy; /* index into seqs */
dx = (struct diag *)g_calloc("to get diags", len0+len1+1, sizeof(struct diag));
ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
ins0 = (dna)? DINS0 : PINS0;
ins1 = (dna)? DINS1 : PINS1;
smax = -10000;
if (endgaps) {
for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
col0[yy] = dely[yy] = col0[yy-1] - ins1;
ndely[yy] = yy;
}
col0[0] = 0; /* Waterman Bull Math Biol 84 */
}
else
for (yy = 1; yy <= len1; yy++)
dely[yy] = -ins0;
/* fill in match matrix
*/
for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
/* initialize first entry in col
*/
if (endgaps) {
if (xx == 1)
col1[0] = delx = -(ins0+ins1);
else
col1[0] = delx = col0[0] - ins1;
ndelx = xx;
}
else {
col1[0] = 0;
delx = -ins0;
ndelx = 0;
}
}
}
for (py = seqx[1], yy = 1; yy <= len1; py++, yy++)
mix = col0[yy-1];
if (dna)
mix += (xbm[*pm-'A']&xbm[*py-'A'])? DMAT : DMIS;
else
mix += _day[*px-'A'][*py-'A'];
/* update penalty for del in x seq;
*favor new del over ongoing del

```

TABLE 1-continued

```

* ignore MAXGAP if weighting endgaps
*/
if (endgaps || ndely[yy] < MAXGAP) {
    if (col0[yy] - ins0 >= dely[yy]) {
        dely[yy] = col0[yy] - (ins0+ins1);
        ndely[yy] = 1;
    } else {
        dely[yy] -= ins1;
        ndely[yy]++;
    }
} else {
    if (col0[yy] - (ins0+ins1) >= dely[yy]) {
        dely[yy] = col0[yy] - (ins0+ins1);
        ndely[yy] = 1;
    } else
        ndely[yy]++;
}
/* update penalty for del in y seq;
* favor new del over ongong del
*/
if (endgaps || ndelx < MAXGAP) {
    if (col1[yy-1] - ins0 >= delx) {
        delx = col1[yy-1] - (ins0+ins1);
        ndelx = 1;
    } else {
        delx -= ins1;
        ndelx++;
    }
} else {
    if (col1[yy-1] - (ins0+ins1) >= delx) {
        delx = col1[yy-1] - (ins0+ins1);
        ndelx = 1;
    } else
        ndelx++;
}
/* pick the maximum score; we're favoring
* mis over any del and delx over dely
*/
id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    col1[yy] = mis;
else if (delx >= dely[yy]) {
    col1[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if(++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    col1[yy] = dely[yy];
    ij = dx[id].ijmp;
if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if(xx == len0 && yy < len1) {

```

TABLE 1-continued

```

        /* last col
        */
        if (endgaps)
            col1[yy] -= ins0+ins1*(len1-yy);
        if(col1[yy] > smax) {
            smax = col1[yy];
            dmax = id;
        }
    }
}
if (endgaps && xx < len1)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if(col1[yy-1] > smax) {
    smax = col1[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}
/*
*
* print() - only routine visible outside this module
*
* static:
* getmat() - trace back best path, count matches: print()
* pr_align() -- print alignment of described in array p[]: print()
* dumpblock() -- dump a block of lines with numbers, stars: pr_align()
* nums() -- put out a number line: dumpblock()
* putline() -- put out a line (name, [num], seq, [num]): dumpblock()
* stars() --put a line of stars: dumpblock()
* stripname() -- strip any path and prefix from a seqname
*/
#include "nw.h"
#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */
extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */
print()
{
    int lx, ly, firstgap, lastgap; /* overlap */
    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", nameX[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", nameX[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}
/*

```

print

TABLE 1-continued

```

* trace back the best path, count matches
*/
static
getmat(lx, ly, firstgap, lastgap)                                     getmat
int    lx, ly; /* "core" (minus endgaps) */
int    firstgap, lastgap; /* leading trailing overlap */
{
    int    nm, i0, i1, siz0, siz1;
    char   outx[32];
    double pct;
    register    n0, n1;
    register char *p0, *p1;
    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;
    nm = 0;
    while (*p0 && *p1) {
        if (siz0) {
            p1 ++;
            n1 ++;
            siz0--;
        }
        else if (siz1) {
            p0 ++;
            n0 ++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A']*xbm[*p1-'A'])
                nm++;
            if (n0 ++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1 ++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }
    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < lent)? len0: len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*(double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "<%d match%s in an overlap of %d: %.2f percent similarity\n",
            nm, (nm == 1)? "" : "es", lx, pct);
    fprintf(fx, "<gaps in first sequence: %d", gapx);
    if (gapx) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
    }
    if (dna)
        fprintf(fx,
            "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",

```

TABLE 1-continued

```

lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
else
    fprintf(fx, "<endgaps not penalized\n");
}
static      nm;          /* matches in core -- for checking */
static      lmax;       /* lengths of stripped file names */
static      ij[2];      /* imp index for a path */
static      nc[2];      /* number at start of current line */
static      ni[2];      /* current elem number -- for gapping */
static      siz[2];     /*
static char  *ps[2];     /* ptr to current element */
static char  *po[2];     /* ptr to next output char slot */
static char  out[2][P_LINE]; /* output line */
static char  star[P_LINE]; /* set by stars() */
/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nm;          /* char count */
    int      more;
    register i;
    for(i = 0, lmax = 0; i < 2; i++) {
        nm = stripname(name[x[i]]);
        if (nm > lmax)
            lmax = nm;
        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seq[x[i]];
        po[i] = out[i];
    }
    for (nm = nm = 0, more = 1; more;) {
        for (i = more = 0; i < 2; i++) {
            /*
             * do we have more of this sequence?
             */
            if (!*ps[i])
                continue;
            more++;
            if (pp[i].spc) { /* leading space */
                *po[i]++ = ' ';
                pp[i].spc--;
            }
            else if (siz[i]) { /* in a gap */
                *po[i]++ = '-';
                siz[i]--;
            }
            else { /* we're putting a seq element
             */
                *po[i] = *ps[i];
                if (islower(*ps[i]))
                    *ps[i] = toupper(*ps[i]);
                po[i]++;
                ps[i]++;
                /*
                 * are we at next gap for this seq?
                 */
                if (ni[i] == pp[i].x[ij[i]]) {
                    /*
                     * we need to merge all gaps
                     * at this location
                     */
                    siz[i] = pp[i].n[ij[i]++];
                    while (ni[i] == pp[i].x[ij[i]])
                        siz[i] += pp[i].n[ij[i]++];
                }
                ni[i]++;
            }
        }
        if(++nn == olen || !more && nm) {
            dumpblock();
            for (i = 0; i < 2; i++)
                po[i] = out[i];
            nn = 0;

```

pr\_align

...pr\_align

TABLE 1-continued

```

    }
}
}
/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()
{
    register i;
    for(i = 0; i < 2; i++)
        *po[i] = '\0';

    (void) putc("\n", fx);
    for (i = 0; i < 2; i++) {
        if (*out[i] && (*out[i]) != ' ' || *(po[i] != ' ')) {
            if(i == 0)
                nums(i);
            if(i == 0 && *out[1])
                stars();
            putline(i);
            if (i == 0 && *out[1])
                fprintf(fx, star);
            if(i == 1)
                nums(i);
        }
    }
}
/*
 * put out a number line: dumpblock()
 */
static
nums(ix)
int    ix;          /* index in out[] holding seq line */
{
    char    nline[P_LINE];
    register i, j;
    register char *pn, *px, *py;
    for(pn = nline, i = 0; i < lmax+P_SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if(*py == ' ' || *py == '-')
            *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j /= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = 1;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}
/*
 * put out a line (name, [num], seq. [num]): dumpblock()
 */
static
putline(ix)
int    ix;
{
    int        i;
    register char *px;
    for (px = namex[ix], i = 0; *px && *px != ':'; px++, i++)
        (void) putc(*px, fx);
    for(; i < lmax+P_SPC; i++)
        (void) putc(' ', fx);
}

```

dumpblock

...dumpblock

nums

putline

...putline

TABLE 1-continued

```

/* these count from 1:
 * ni[] is current element (from 1)
 * nc[] is number at start of current line
 */
for (px = out[ix]; *px; px++)
    (void) putc(*px&0x7F, fx);
(void) putc('\n', fx);
}
/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
static
stars()
{
    int            i;
    register char  *p0, *p1, cx, *px;
    if (!*out[0] || (*out[0] == ' ' && *(p0[0] == ' ') ||
        !*out[1] || (*out[1] == ' ' && *(p0[1] == ' ')))
        return;
    px = star;
    for (i = lmax+P_SPC; i--;)
        *px++ = ' ';
    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm ++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '_';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)
char *pn; /* file name (may be path) */
{
    register char *px, *py;
    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq. set dna, len, maxlen
 * g_calloc() -- calloc() with error checkin
 * readjimps() -- get the good jimps, from tmp file if necessary
 * writejimps() -- write a filled array of jimps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>
char *jname = "/tmp/homgXXXXXX"; /* tmp file for jimps */
FILE *fj;
int cleanup(); /* cleanup tmp file */
long lseek();
/*
 * remove any tmp file if we blow
 */
cleanup(i)
int i;
{
    if (fj)

```



TABLE 1-continued

```

        (void) unlink(jname);
        exit(i);
    }
    /*
    * read, return ptr to seq, set dna, len, maxlen
    * skip lines starting with ';', '<', or '>'
    * seq in upper or lower case
    */
    char *
    getseq(file, len)
        char *file;          /* file name */
        int *len;           /* seq len */
    {
        char line[1024], *pseq;
        register char *px, *py;
        int natgc, tlen;
        FILE *fp;
        if ((fp = fopen(file, "r")) == 0) {
            fprintf(stderr, "%s: can't read %s\n", prog, file);
            exit(1);
        }
        tlen = natgc = 0;
        while (fgets(line, 1024, fp)) {
            if (*line == ';' || *line == '<' || *line == '>')
                continue;
            for (px = line; *px != '\n'; px++)
                if (isupper(*px) || islower(*px))
                    tlen++;
        }
        if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
            fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
            exit(1);
        }
        pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

        py = pseq + 4;
        *len = tlen;
        rewind(fp);
        while (fgets(line, 1024, fp)) {
            if (*line == ';' || *line == '<' || *line == '>')
                continue;
            for (px = line; *px != '\n'; px++) {
                if (isupper(*px))
                    *py++ = *px;
                else if (islower(*px))
                    *py++ = toupper(*px);
                if (index("ATGCU", *(py-1)))
                    natgc++;
            }
        }
        *py++ = '\0';
        *py = '/0';
        (void) fclose(fp);
        dna = natgc > (tlen/3);
        return(pseq+4);
    }
    char *
    g__calloc(msg, nx, sz)
        char *msg;          /* program, calling routine */
        int nx, sz;        /* number and size of elements */
    {
        char *px, *calloc();
        if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
            if (*msg) {
                fprintf(stderr, "%s: g__calloc() failed %s (n=%d, sz=%d)\n", prog, msg, nx, sz);
                exit(1);
            }
        }
        return(px);
    }
    /*
    * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
    */
    readjmps()
    {
        int fd = -1;

```

getseq

...getseq

g\_\_calloc

readjmps

TABLE 1-continued

```

int     siz, i0, i1;
register i, j, xx;
if (fj) {
    (void) fclose(fj);
    if ((fd = open(jname, O_RDONLY, 0)) < 0) {
        fprintf(stderr, "%s: can't open() %s\n", prog, jname);
        cleanup(1);
    }
}
for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; ; i++) {
    while (1) {
        for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
            ;

        if (j < 0 && dx[dmax].offset && fj) {
            (void) lseek(fd, dx[dmax].offset, 0);
            (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
            (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
            dx[dmax].ijmp = MAXJMP-1;
        }
        else
            break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;
            /* id = xx - yy + len1 - 1
             */
            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy ++;
            ngapy -= siz;
        }
        /* ignore MAXGAP when doing endgaps */
        siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
        i1++;
    }
    else if (siz > 0) { /* gap in first seq */
        pp[0].n[i0] = siz;
        pp[0].x[i0] = xx;
        gapx ++;
        ngapz += siz;
        /* ignore MAXGAP when doing endgaps */
        siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
        i0++;
    }
}
else
    break;
}
/* reverse the order of jmps
 */
for (j = 0, i0--, j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--, j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}
/*
 * write a filled jmp struct offset of the prey one (if any): nw()

```

TABLE 1-continued

```

*/
writejmps(ix)
int ix;
{
char *mktemp();
if(!fj) {
if (mktemp(jname) < 0) {
fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
cleanup(1);
}
if ((fj = fopen(jname, "w")) == 0) {
fprintf(stderr, "%s: can't write %s\n", prog, jname);
exit(1);
}
}
(void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
(void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}
    
```

[0651]

TABLE 2

PRO	XXXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison	XXXXXXXXYYYYYYYY	(Length = 12 amino acids)
Protein		
% amino acid		
sequence		
identity =		
(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 15 = 33.3%		

[0652]

TABLE 3

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison	XXXXXXXXYYYYZZYZ	(Length = 15 amino acids)
Protein		
% amino acid		
sequence		
identity =		
(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%		

[0653]

TABLE 4

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison	NNNNNNLLLLLLLL	(Length = 16 nucleotides)
DNA		
% nucleic acid		
sequence		
identity =		
(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 14 = 42.9%		

[0654]

TABLE 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison	NNNNLLLV	(Length = 9 nucleotides)
DNA		
% nucleic acid		
sequence		
identity =		
(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 12 = 33.3%		

[0655] II. Compositions and Methods of the Invention

[0656] A. Full-Length PRO Polypeptides

[0657] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

[0658] As disclosed in the Examples below, various cDNA clones have been deposited with the ATCC. The actual nucleotide sequences of those clones can readily be determined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

**[0659]** 1. Full-length PRO211 and PRO217 Polypeptides

**[0660]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO211 and PRO217. In particular, Applicants have identified and isolated cDNA encoding PRO211 and PRO217 polypeptides, as disclosed in further detail in the Examples below. Using BLAST (FastA format) sequence alignment computer programs, Applicants found that cDNA sequences encoding full-length native sequence PRO211 and PRO217 have homologies to known proteins having EGF-like domains. Specifically, the cDNA sequence DNA32292-1131 (**FIG. 1**, SEQ ID NO:1) has certain identity and a Blast score of 209 with PAC6\_RAT and certain identity and a Blast score of 206 with Fibulin-1, isoform c precursor. The cDNA sequence DNA33094-1131 (**FIG. 3**, SEQ ID NO:3) has 36% identity and a Blast score of 336 with eastern newt tenascin, and 37% identity and a Blast score of 331 with human tenascin-X precursor. Accordingly, it is presently believed that PRO211 and PRO217 polypeptides disclosed in the present application are newly identified members of the EGF-like family and possesses properties typical of the EGF-like protein family.

**[0661]** 2. Full-length PRO230 Polypeptides

**[0662]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO230. In particular, Applicants have identified and isolated cDNA encoding a PRO230 polypeptide, as disclosed in further detail in the Examples below. Using known programs such as BLAST and FastA sequence alignment computer programs, Applicants found that a cDNA sequence encoding full-length native sequence PRO230 has 48% amino acid identity with the rabbit tubulointerstitial nephritis antigen precursor. Accordingly, it is presently believed that PRO230 polypeptide disclosed in the present application is a newly identified member of the tubulointerstitial nephritis antigen family and possesses the ability to be recognized by human autoantibodies in certain forms of tubulointerstitial nephritis.

**[0663]** 3. Full-length PRO232 Polypeptides

**[0664]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO232. In particular, Applicants have identified and isolated cDNA encoding a PRO232 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a portion of The full-length native sequence PRO232 (shown in **FIG. 9** and SEQ ID NO:18) has 35% sequence identity with a stem cell surface antigen from Gallus gallus. Accordingly, it is presently believed that the PRO232 polypeptide disclosed in the present application may be a newly identified stem cell antigen.

**[0665]** 4. Full-length PRO187 Polypeptides

**[0666]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO187. In particular, Applicants have identified and isolated cDNA encoding a PRO187 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length

native sequence PRO187 (shown in **FIG. 15**) has 74% amino acid sequence identity and BLAST score of 310 with various androgen-induced growth factors and FGF-8. Accordingly, it is presently believed that PRO187 polypeptide disclosed in the present application is a newly identified member of the FGF-8 protein family and may possess identify activity or property typical of the FGF-8-like protein family.

**[0667]** 5. Full-length PRO265 Polypeptides

**[0668]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO265. In particular, Applicants have identified and isolated cDNA encoding a PRO265 polypeptide, as disclosed in further detail in the Examples below. Using programs such as BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO265 polypeptide have significant homology with the fibromodulin protein and fibromodulin precursor protein. Applicants have also found that the DNA encoding the PRO265 polypeptide has significant homology with platelet glycoprotein V, a member of the leucine rich related protein family involved in skin and wound repair. Accordingly, it is presently believed that PRO265 polypeptide disclosed in the present application is a newly identified member of the leucine rich repeat family and possesses protein binding capabilities, as well as be involved in skin and wound repair as typical of this family.

**[0669]** 6. Full-length PRO219 Polypeptides

**[0670]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO219. In particular, Applicants have identified and isolated cDNA encoding a PRO219 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO219 polypeptide have significant homology with the mouse and human matrilin-2 precursor polypeptides. Accordingly, it is presently believed that PRO219 polypeptide disclosed in the present application is related to the matrilin-2 precursor polypeptide.

**[0671]** 7. Full-length PRO246 Polypeptides

**[0672]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO246. In particular, Applicants have identified and isolated cDNA encoding a PRO246 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a portion of the PRO246 polypeptide has significant homology with the human cell surface protein HCAR. Accordingly, it is presently believed that PRO246 polypeptide disclosed in the present application may be a newly identified membrane-bound virus receptor or tumor cell-specific antigen.

**[0673]** 8. Full-length PRO228 Polypeptides

**[0674]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO228. In particular, Applicants have identified and isolated cDNA encoding a PRO228 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence align-

ment computer programs, Applicants found that various portions of the PRO228 polypeptide have significant homology with the EMR1 protein. Applicants have also found that the DNA encoding the PRO228 polypeptide has significant homology with latrophilin, macrophage-restricted cell surface glycoprotein, B0457.1 and leucocyte antigen CD97 precursor. Accordingly, it is presently believed that PRO228 polypeptide disclosed in the present application is a newly identified member of the seven transmembrane superfamily and possesses characteristics and functional properties typical of this family. In particular, it is believed that PRO228 is a new member of the subgroup within this family to which CD97 and EMR1 belong.

**[0675]** 9. Full-length PRO533 Polypeptides

**[0676]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO533. In particular, Applicants have identified and isolated cDNA encoding a PRO533 polypeptide, as disclosed in further detail in the Examples below. Using BLAST-2 and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO533 (shown in **FIG. 22** and SEQ ID NO:59) has a Blast score of 509 and 53% amino acid sequence identity with fibroblast growth factor (FGF). Accordingly, it is presently believed that PRO533 disclosed in the present application is a newly identified member of the fibroblast growth factor family and may possess activity typical of such polypeptides.

**[0677]** 10. Full-length PRO245 Polypeptides

**[0678]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO245. In particular, Applicants have identified and isolated cDNA encoding a PRO245 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a portion of the amino acid sequence of the PRO245 polypeptide has 60% amino acid identity with the human c-myb protein. Accordingly, it is presently believed that the PRO245 polypeptide disclosed in the present application may be a newly identified member of the transmembrane protein tyrosine kinase family.

**[0679]** 11. Full-length PRO220, PRO221 and PRO227 Polypeptides

**[0680]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO220, PRO221 and PRO227. In particular, Applicants have identified and isolated cDNAs encoding a PRO220, PRO221 and PRO227 polypeptide, respectively, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, PRO220 has amino acid identity with the amino acid sequence of a leucine rich protein wherein the identity is 87%. PRO220 additionally has amino acid identity with the neuronal leucine rich protein wherein the identity is 55%. The neuronal leucine rich protein is further described in Taguchi, et al., *Mol. Brain Res.*, 35:31-40 (1996).

**[0681]** PRO221 has amino acid identity with the SLIT protein precursor, wherein different portions of these two proteins have the respective percent identities of 39%, 38%, 34%, 31%, and 30%.

**[0682]** PRO227 has amino acid identity with the amino acid sequence of platelet glycoprotein V precursor. The same results were obtained for human glycoprotein V. Different portions of these two proteins show the following percent identities of 30%, 28%, 28%, 31%, 35%, 39% and 27%.

**[0683]** Accordingly, it is presently believed that PRO220, PRO221 and PRO227 polypeptides disclosed in the present application are newly identified members of the leucine rich repeat protein superfamily and that each possesses protein-protein binding capabilities typical of the leucine rich repeat protein superfamily. It is also believed that they have capabilities similar to those of SLIT, the leucine rich repeat protein and human glycoprotein V.

**[0684]** 12. Full-length PRO258 Polypeptides

**[0685]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO258. In particular, Applicants have identified and isolated cDNA encoding a PRO258 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO258 polypeptide have significant homology with the CRTAM and poliovirus receptors. Accordingly, it is presently believed that PRO258 polypeptide disclosed in the present application is a newly identified member of the Ig superfamily and possesses virus receptor capabilities or regulates immune function as typical of this family.

**[0686]** 13. Full-length PRO266 Polypeptides

**[0687]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO266. In particular, Applicants have identified and isolated cDNA encoding a PRO266 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO266 polypeptide have significant homology with the SLIT protein from *Drosophila*. Accordingly, it is presently believed that PRO266 polypeptide disclosed in the present application is a newly identified member of the leucine rich repeat family and possesses ligand-ligand binding activity and neuronal development typical of this family. SLIT has been shown to be useful in the study and treatment of Alzheimer's disease, supra, and thus, PRO266 may have involvement in the study and cure of this disease.

**[0688]** 14. Full-length PRO269 Polypeptides

**[0689]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO269. In particular, Applicants have identified and isolated cDNA encoding a PRO269 polypeptide, as disclosed in further detail in the Examples below. Using BLAST, FastA and sequence alignment computer programs, Applicants found that the amino acid sequence encoded by nucleotides 314 to 1783 of the full-length native sequence PRO269 (shown in **FIG. 35** and SEQ ID NO:95) has significant homology to human urinary thrombomodulin and various thrombomodulin analogues respectively, to which it was aligned. Accordingly, it is presently believed that PRO269 polypeptide disclosed in the present application is a newly identified member of the thrombomodulin family.

**[0690]** 15. Full-length PRO287 Polypeptides

**[0691]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO287. In particular, Applicants have identified and isolated cDNA encoding a PRO287 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO287 polypeptide have significant homology with the type 1 procollagen C-proteinase enhancer protein precursor and type 1 procollagen C-proteinase enhancer protein. Accordingly, it is presently believed that PRO287 polypeptide disclosed in the present application is a newly identified member of the C-proteinase enhancer protein family.

**[0692]** 16. Full-length PRO214 Polypeptides

**[0693]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO214. In particular, Applicants have identified and isolated cDNA encoding a PRO214 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO214 polypeptide (shown in **FIG. 40** and SEQ ID NO:109) has 49% amino acid sequence identity with HT protein, a known member of the EGF-family. The comparison resulted in a BLAST score of 920, with 150 matching nucleotides. Accordingly, it is presently believed that the PRO214 polypeptide disclosed in the present application is a newly identified member of the family comprising EGF domains and may possess activities or properties typical of the EGF-domain containing family.

**[0694]** 17. Full-length PRO317 Polypeptides

**[0695]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO317. In particular, cDNA encoding a PRO317 polypeptide has been identified and isolated, as disclosed in further detail in the Examples below. Using BLAST™ and FastA™ sequence alignment computer programs, it was found that a full-length native-sequence PRO317 (shown in **FIG. 42** and SEQ ID NO:114) has 92% amino acid sequence identity with EBAF-1. Further, it is closely aligned with many other members of the TGF-superfamily.

**[0696]** Accordingly, it is presently believed that PRO317 disclosed in the present application is a newly identified member of the TGF-superfamily and may possess properties that are therapeutically useful in conditions of uterine bleeding, etc. Hence, PRO317 may be useful in diagnosing or treating abnormal bleeding involved in gynecological diseases, for example, to avoid or lessen the need for a hysterectomy. PRO317 may also be useful as an agent that affects angiogenesis in general, so PRO317 may be useful in anti-tumor indications, or conversely, in treating coronary ischemic conditions.

**[0697]** Library sources reveal that ESTs used to obtain the consensus DNA for generating PRO317 primers and probes were found in normal tissues (uterus, prostate, colon, and pancreas), in several tumors (colon, brain (twice), pancreas, and mullerian cell), and in a heart with ischemia. PRO317 has shown up in several tissues as well, but it does look to

have a greater concentration in uterus. Hence, PRO317 may have a broader use by the body than EBAF-1. It is contemplated that, at least for some indications, PRO317 may have opposite effects from EBAF-1.

**[0698]** 18. Full-length PRO301 Polypeptides

**[0699]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO301. In particular, Applicants have identified and isolated cDNA encoding a PRO301 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO301 (shown in **FIG. 44** and SEQ ID NO:119) has a Blast score of 246 corresponding to 30% amino acid sequence identity with human A33 antigen precursor. Accordingly, it is presently believed that PRO301 disclosed in the present application is a newly identified member of the A33 antigen protein family and may be expressed in human neoplastic diseases such as colorectal cancer.

**[0700]** 19. Full-length PRO224 Polypeptides

**[0701]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO224. In particular, Applicants have identified and isolated cDNA encoding a PRO224 polypeptide, as disclosed in further detail in the Examples below. Using known programs such as BLAST and FastA sequence alignment computer programs, Applicants found that full-length native PRO224 (**FIG. 46**, SEQ ID NO:127) has amino acid identity with apolipoprotein E receptor 2906 from homo sapiens. The alignments of different portions of these two polypeptides show amino acid identities of 37%, 36%, 30%, 44%, 44% and 28% respectively. Full-length native PRO224 (**FIG. 46**, SEQ ID NO:127) also has amino acid identity with very low-density lipoprotein receptor precursor from gall. The alignments of different portions of these two polypeptides show amino acid identities of 38%, 37%, 42%, 33%, and 37% respectively. Additionally, full-length native PRO224 (**FIG. 46**, SEQ ID NO:127) has amino acid identity with the chicken oocyte receptor P95 from Gallus gallus. The alignments of different portions of these two polypeptides show amino acid identities of 38%, 37%, 42%, 33%, and 37% respectively. Moreover, full-length native PRO224 (**FIG. 46**, SEQ ID NO:127) has amino acid identity with very low density lipoprotein receptor short form precursor from humans. The alignments of different portions of these two polypeptides show amino acid identities of 32%, 38%, 34%, 45%, and 31%, respectively. Accordingly, it is presently believed that PRO224 polypeptide disclosed in the present application is a newly identified member of the low density lipoprotein receptor family and possesses the structural characteristics required to have the functional ability to recognize and endocytose low density lipoproteins typical of the low density lipoprotein receptor family. (The alignments described above used the following scoring parameters: T=7, S+65, S2=36, Matrix: BLOSUM62.)

**[0702]** 20. Full-length PRO222 Polypeptides

**[0703]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO222. In particu-

lar, Applicants have identified and isolated cDNA encoding a PRO222 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a sequence encoding full-length native sequence PRO222 (shown in FIG. 48 and SEQ ID NO:132) has 25-26% amino acid identity with mouse complement factor h precursor, has 27-29% amino acid identity with complement receptor, has 25-47% amino acid identity with mouse complement C3b receptor type 2 long form precursor, has 40% amino acid identity with human hypothetical protein k1aa0247. Accordingly, it is presently believed that PRO222 polypeptide disclosed in the present application is a newly identified member of the complement receptor family and possesses activity typical of the complement receptor family.

[0704] 21. Full-length PRO234 Polypeptides

[0705] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO234. In particular, Applicants have identified and isolated cDNA encoding a PRO234 polypeptide, as disclosed in further detail in the Examples below. Using BLAST (FastA-format) sequence alignment computer programs, Applicants found that a cDNA sequence encoding full-length native sequence PRO234 has 31% identity and Blast score of 134 with E-selectin precursor. Accordingly, it is presently believed that the PRO234 polypeptides disclosed in the present application are newly identified members of the lectin/selectin family and possess activity typical of the lectin/selectin family.

[0706] 22. Full-length PRO231 Polypeptides

[0707] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO231. In particular, Applicants have identified and isolated cDNA encoding a PRO231 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that the full-length native sequence PRO231 polypeptide (shown in FIG. 52 and SEQ ID NO:142) has 30% and 31% amino acid identity with human and rat prostatic acid phosphatase precursor proteins, respectively. Accordingly, it is presently believed that the PRO231 polypeptide disclosed in the present application may be a newly identified member of the acid phosphatase protein family.

[0708] 23. Full-length PRO229 Polypeptides

[0709] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO229. In particular, Applicants have identified and isolated cDNA encoding a PRO229 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO229 polypeptide have significant homology with antigen wc1.1, M130 antigen, T cell surface glycoprotein CD6 and CD6. It also is related to Sp-alpha. Accordingly, it is presently believed that PRO229 polypeptide disclosed in the present application is a newly identified member of the family containing scavenger receptor homology, a sequence motif found in a number of proteins involved in immune function and thus possesses immune function and/or segments which resist degradation, typical of this family.

[0710] 24. Full-length PRO238 Polypeptides

[0711] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO238. In particular, Applicants have identified and isolated cDNA encoding a PRO238 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO238 polypeptide have significant homology with reductases, including oxidoreductase and fatty acyl-CoA reductase. Accordingly, it is presently believed that PRO238 polypeptide disclosed in the present application is a newly identified member of the reductase family and possesses reducing activity typical of the reductase family.

[0712] 25. Full-length PRO233 Polypeptides

[0713] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO233. In particular, Applicants have identified and isolated cDNA encoding a PRO233 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO233 polypeptide have significant homology with the reductase protein. Applicants have also found that the DNA encoding the PRO233 polypeptide has significant homology with proteins from *Caenorhabditis elegans*. Accordingly, it is presently believed that PRO233 polypeptide disclosed in the present application is a newly identified member of the reductase family and possesses the ability to effect the redox state of the cell typical of the reductase family.

[0714] 26. Full-length PRO223 Polypeptides

[0715] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO223. In particular, Applicants have identified and isolated cDNA encoding a PRO223 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that the PRO223 polypeptide has significant homology with various serine carboxypeptidase polypeptides. Accordingly, it is presently believed that PRO223 polypeptide disclosed in the present application is a newly identified serine carboxypeptidase.

[0716] 27. Full-length PRO235 Polypeptides

[0717] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO235. In particular, Applicants have identified and isolated cDNA encoding a PRO235 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO235 polypeptide have significant homology with the various plexin proteins. Accordingly, it is presently believed that PRO235 polypeptide disclosed in the present application is a newly identified member of the plexin family and possesses cell adhesion properties typical of the plexin family.

**[0718]** 28. Full-length PRO236 and PRO262 Polypeptides

**[0719]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO236 and PRO262. In particular, Applicants have identified and isolated cDNA encoding PRO236 and PRO262 polypeptides, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO236 and PRO262 polypeptides have significant homology with various  $\beta$ -galactosidase and  $\beta$ -galactosidase precursor polypeptides. Accordingly, it is presently believed that the PRO236 and PRO262 polypeptides disclosed in the present application are newly identified  $\beta$ -galactosidase homologs.

**[0720]** 29. Full-length PRO239 Polypeptides

**[0721]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO239. In particular, Applicants have identified and isolated cDNA encoding a PRO239 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO239 polypeptide have significant homology with densin proteins. Accordingly, it is presently believed that PRO239 polypeptide disclosed in the present application is a newly identified member of the densin family and possesses cell adhesion and the ability to effect synaptic processes as is typical of the densin family.

**[0722]** 30. Full-length PRO257 Polypeptides

**[0723]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO257. In particular, Applicants have identified and isolated cDNA encoding a PRO257 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO257 polypeptide have significant homology with the ebnerin precursor and ebnerin protein. Accordingly, it is presently believed that PRO257 polypeptide disclosed in the present application is a newly identified protein member which is related to the ebnerin protein.

**[0724]** 31. Full-length PRO260 Polypeptides

**[0725]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO260. In particular, Applicants have identified and isolated cDNA encoding a PRO260 polypeptide, as disclosed in further detail in the Examples below. Using programs such as BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO260 polypeptide have significant homology with the alpha-1-fucosidase precursor. Accordingly, it is presently believed that PRO260 polypeptide disclosed in the present application is a newly identified member of the fucosidase family and possesses enzymatic activity related to fucose residues typical of the fucosidase family.

**[0726]** 32. Full-length PRO263 Polypeptides

**[0727]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO263. In particu-

lar, Applicants have identified and isolated cDNA encoding a PRO263 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO263 polypeptide have significant homology with the CD44 antigen and related proteins. Accordingly, it is presently believed that PRO263 polypeptide disclosed in the present application is a newly identified member of the CD44 antigen family and possesses at least one of the properties associated with these antigens, i.e., cancer and HIV marker, cell-cell or cell-matrix interactions, regulating cell traffic, lymph node homing, transmission of growth signals, and presentation of chemokines and growth factors to traveling cells.

**[0728]** 33. Full-length PRO270 Polypeptides

**[0729]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO270. In particular, Applicants have identified and isolated cDNA encoding a PRO270 polypeptide, as disclosed in further detail in the Examples below. Using BLAST, FastA and sequence alignment computer programs, Applicants found that various portions of the PRO270 polypeptide have significant homology with various thioredoxin proteins. Accordingly, it is presently believed that PRO270 polypeptide disclosed in the present application is a newly identified member of the thioredoxin family and possesses the ability to effect reduction-oxidation (redox) state typical of the thioredoxin family.

**[0730]** 34. Full-length PRO271 Polypeptides

**[0731]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO271. In particular, Applicants have identified and isolated cDNA encoding a PRO271 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that the PRO271 polypeptide has significant homology with various link proteins and precursors thereof. Accordingly, it is presently believed that PRO271 polypeptide disclosed in the present application is a newly identified link protein homolog.

**[0732]** 35. Full-length PRO272 Polypeptides

**[0733]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO272. In particular, Applicants have identified and isolated cDNA encoding a PRO272 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO272 polypeptide have significant homology with the human reticulocalbin protein and its precursors. Applicants have also found that the DNA encoding the PRO272 polypeptide has significant homology with the mouse reticulocalbin precursor protein. Accordingly, it is presently believed that PRO272 polypeptide disclosed in the present application is a newly identified member of the reticulocalbin family and possesses the ability to bind calcium typical of the reticulocalbin family.

**[0734]** 36. Full-length PRO294 Polypeptides

**[0735]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides



referred to in the present application as PRO294. In particular, Applicants have identified and isolated cDNA encoding a PRO294 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO294 polypeptide have significant homology with the various portions of a number of collagen proteins. Accordingly, it is presently believed that PRO294 polypeptide disclosed in the present application is a newly identified member of the collagen family.

**[0736]** 37. Full-length PRO295 Polypeptides

**[0737]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO295. In particular, Applicants have identified and isolated cDNA encoding a PRO295 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO295 polypeptide have significant homology with integrin proteins. Accordingly, it is presently believed that PRO295 polypeptide disclosed in the present application is a newly identified member of the integrin family and possesses cell adhesion typical of the integrin family.

**[0738]** 38. Full-length PRO293 Polypeptides

**[0739]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO293. In particular, Applicants have identified and isolated cDNA encoding a PRO293 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that portions of the PRO293 polypeptide have significant homology with the neuronal leucine rich repeat proteins 1 and 2, (NLRR-1 and NLRR-2), particularly NLRR-2. Accordingly, it is presently believed that PRO293 polypeptide disclosed in the present application is a newly identified member of the neuronal leucine rich repeat protein family and possesses ligand-binding activity typical of the NRLL protein family.

**[0740]** 39. Full-length PRO247 Polypeptides

**[0741]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO247. In particular, Applicants have identified and isolated cDNA encoding a PRO247 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO247 polypeptide have significant homology with densin. Applicants have also found that the DNA encoding the PRO247 polypeptide has significant homology with a number of other proteins, including KIAA023 1. Accordingly, it is presently believed that PRO247 polypeptide disclosed in the present application is a newly identified member of the leucine rich repeat family and possesses ligand binding abilities typical of this family.

**[0742]** 40. Full-length PRO302, PRO303, PRO304, PRO307 and PRO343 Polypeptides

**[0743]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO302, PRO303,

PRO304, PRO307 and PRO343. In particular, Applicants have identified and isolated cDNA encoding PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides have significant homology with various protease proteins. Accordingly, it is presently believed that the PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides disclosed in the present application are newly identified protease proteins.

**[0744]** 41. Full-length PRO328 Polypeptides

**[0745]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO328. In particular, Applicants have identified and isolated cDNA encoding a PRO328 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO328 polypeptide have significant homology with the human glioblastoma protein ("GLIP"). Further, Applicants found that various portions of the PRO328 polypeptide have significant homology with the cysteine rich secretory protein ("CRISP") as identified by BLAST homology [ECCRISP3\_1, S68683, and CRS3\_HUMAN]. Accordingly, it is presently believed that PRO328 polypeptide disclosed in the present application is a newly identified member of the GLIP or CRISP families and possesses transcriptional regulatory activity typical of the GLIP or CRISP families.

**[0746]** 42. Full-length PRO335, PRO331 and PRO326 Polypeptides

**[0747]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO335, PRO331 or PRO326. In particular, Applicants have identified and isolated cDNA encoding a PRO335, PRO331 or PRO326 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO335, PRO331 or PRO326 polypeptide have significant homology with LIG-1, ALS and in the case of PRO331, additionally, decorin. Accordingly, it is presently believed that the PRO335, PRO331 and PRO326 polypeptides disclosed in the present application are newly identified members of the leucine rich repeat superfamily, and particularly, are related to LIG-1 and possess the biological functions of this family as discussed and referenced herein.

**[0748]** 43. Full-length PRO332 Polypeptides

**[0749]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO332. In particular, Applicants have identified and isolated cDNA encoding PRO332 polypeptides, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO332 (shown in **FIG. 108** and SEQ ID NO:310) has about 30-40% amino acid sequence identity with a series of known proteoglycan sequences, including, for example, fibromodulin and fibromodulin precursor

sequences of various species (FMOD\_BOVIN, FMOD\_CHICK, FMOD\_RAT, FMOD\_MOUSE, FMOD\_HUMAN, P\_R36773), osteomodulin sequences (AB000114\_1, AB007848\_1), decorin sequences (CFU83141\_1, OCU03394\_1, P\_R42266, P\_R42267, P\_R42260, P\_R89439), keratan sulfate proteoglycans (BTU48360\_1, AF022890\_1), corneal proteoglycan (AF022256\_1), and bone/cartilage proteoglycans and proteoglycane precursors (PGS1\_BOVIN, PGS2\_MOUSE, PGS2\_HUMAN). Accordingly, it is presently believed that PRO332 disclosed in the present application is a new proteoglycan-type molecule, and may play a role in regulating extracellular matrix, cartilage, and/or bone function.

**[0750]** 44. Full-length PRO334 Polypeptides

**[0751]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO334. In particular, Applicants have identified and isolated cDNA encoding a PRO334 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO334 polypeptide have significant homology with fibulin and fibrillin. Accordingly, it is presently believed that PRO334 polypeptide disclosed in the present application is a newly identified member of the epidermal growth factor family and possesses properties and activities typical of this family.

**[0752]** 45. Full-length PRO346 Polypeptides

**[0753]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO346. In particular, Applicants have identified and isolated cDNA encoding a PRO346 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO346 (shown in **FIG. 112** and SEQ ID NO:320) has 28% amino acid sequence identity with carcinoembryonic antigen. Accordingly, it is presently believed that PRO346 disclosed in the present application is a newly identified member of the carcinoembryonic protein family and may be expressed in association with neoplastic tissue disorders.

**[0754]** 46. Full-length PRO268 Polypeptides

**[0755]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO268. In particular, Applicants have identified and isolated cDNA encoding a PRO268 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that portions of the PRO268 polypeptide have significant homology with the various protein disulfide isomerase proteins. Accordingly, it is presently believed that PRO268 polypeptide disclosed in the present application is a homolog of the protein disulfide isomerase p5 protein.

**[0756]** 47. Full-length PRO330 Polypeptides

**[0757]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO330. In particular, Applicants have identified and isolated cDNA encoding

a PRO330 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO330 polypeptide have significant homology with the murine prolyl 4-hydroxylase alpha-II subunit protein. Accordingly, it is presently believed that PRO330 polypeptide disclosed in the present application is a novel prolyl 4-hydroxylase subunit polypeptide.

**[0758]** 48. Full-length PRO339 and PRO310 Polypeptides

**[0759]** The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO339 and PRO310. In particular, Applicants have identified and isolated cDNA encoding a PRO339 polypeptide, as disclosed in further detail in the Examples below. Applicants have also identified and isolated cDNA encoding a PRO310 polypeptide, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that various portions of the PRO339 and PRO310 polypeptides have significant homology with small secreted proteins from *C. elegans* and are distantly related to fringe. PRO339 also shows homology to collagen-like polymers. Sequences which were used to identify PRO310, designated herein as DNA40533 and DNA42267, also show homology to proteins from *C. elegans*. Accordingly, it is presently believed that the PRO339 and PRO310 polypeptides disclosed in the present application are newly identified member of the family of proteins involved in development, and which may have regulatory abilities similar to the capability of fringe to regulate serrate.

**[0760]** 49. Full Length PRO244 Polypeptides

**[0761]** The present invention provides newly identified and isolated nucleotide sequences encoding C-type lectins referred to in the present application as PRO244. In particular, applicants have identified and isolated cDNA encoding PRO244 polypeptides, as disclosed in further detail in the Examples below. Using BLAST and FastA sequence alignment computer programs, Applicants found that a full-length native sequence PRO244 (shown in **FIG. 122** and SEQ ID NO:377) has 43% amino acid sequence identity with the hepatic lectin gallus gallus (LECH-CHICK), and 42% amino acid sequence identity with an HIV gp120 binding C-type lectin (A46274). Accordingly, it is presently believed that PRO244 disclosed in the present application is a newly identified member of the C-lectin superfamily and may play a role in immune function, apoptosis, or in the pathogenesis of atherosclerosis. In addition, PRO244 may be useful in identifying tumor-associated epitopes.

**[0762]** B. PRO Polypeptide Variants

**[0763]** In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

**[0764]** Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be

made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Pat. No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

[0765] PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

[0766] PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

[0767] In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	glu

TABLE 6—continued

Original Residue	Exemplary Substitutions	Preferred Substitutions
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

[0768] Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- [0769] (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- [0770] (2) neutral hydrophilic: cys, ser, thr;
- [0771] (3) acidic: asp, glu;
- [0772] (4) basic: asn, gln, his, lys, arg;
- [0773] (5) residues that influence chain orientation: gly, pro; and
- [0774] (6) aromatic: trp, tyr, phe.

[0775] Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

[0776] The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.* 10:6487 (1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

[0777] Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine

is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, *Science*, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W. H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

#### [0778] C. Modifications of PRO

[0779] Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis-(diazocetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propionimide.

[0780] Other modifications include deamidation of glutamyl and asparagyl residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0781] Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

[0782] Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0783] Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published Sep. 11, 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

[0784] Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

[0785] Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[0786] The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

[0787] In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl-terminus of the PRO. The presence of such epitope-tagged forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192-194 (1992)]; an  $\alpha$ -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA* 87:6393-6397 (1990)].

[0788] In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly

preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

#### [0789] D. Preparation of PRO

[0790] The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., *Solid-Phase Peptide Synthesis*, W. H. Freeman Co., San Francisco, Calif. (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using in Applied Biosystems Peptide Synthesizer (Foster City, Calif.) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

#### [0791] 1. Isolation of DNA Encoding PRO

[0792] DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

[0793] Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

[0794] The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like  $^{32}\text{P}$ -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

[0795] Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence

identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

[0796] Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

#### [0797] 2. Selection and Transformation of Host Cells

[0798] Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: a Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

[0799] Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example,  $\text{CaCl}_2$ ,  $\text{CaPO}_4$ , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published Jun. 29, 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

[0800] Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*,

e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype tonA; *E. coli* W3110 strain 9E4, which has the complete genotype tonA ptr3; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA ptr3phoA E15 (argF-lac)169 degP ompT kan<sup>r</sup>; *E. coli* W3110 strain 37D6, which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan<sup>r</sup>; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Pat. No. 4,946,783 issued Aug. 7, 1990. Alternatively, in vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

[0801] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290: 140 [1981]; EP 139,383 published May 2, 1985); *Kluyveromyces* hosts (U.S. Pat. No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS68:3, CBS4574; Louvencourt et al., *J. Bacteriol.*, 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilorum* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., *Proc. Natl. Acad. Sci. USA*, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published Oct. 31, 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolyptocladium* (WO 91/00357 published Jan. 10, 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., *Biochem. Biophys. Res. Commun.*, 112:284-289 [1983]; Tilburn et al., *Gene*, 26:205-221 [1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479 [1985]). Methylotrophic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylotrophs*, 269 (1982).

[0802] Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include

Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen Virol.*, 36:59 (1977)); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

### [0803] 3. Selection and Use of a Replicable Vector

[0804] The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

[0805] The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, 1pp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Pat. No. 5,010, 182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published Apr. 4, 1990), or the signal described in WO 90/13646 published Nov. 15, 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

[0806] Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the  $2\mu$  plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

[0807] Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer

resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for Bacilli.

**[0808]** An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

**[0809]** Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

**[0810]** Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

**[0811]** Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytocrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

**[0812]** PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published Jul. 5, 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

**[0813]** Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\alpha$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

**[0814]** Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

**[0815]** Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

#### **[0816]** 4. Detecting Gene Amplification/Expression

**[0817]** Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

**[0818]** Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

#### **[0819]** 5. Purification of Polypeptide

**[0820]** Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent

solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

[0821] It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

#### [0822] E. Uses for PRO

[0823] Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

[0824] The full-length native sequence PRO gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO or PRO from other species) which have a desired sequence identity to the native PRO sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO. By way of example, a screening method will comprise isolating the coding region of the PRO gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

[0825] Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

[0826] Other useful fragments of the PRO nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA)

capable of binding to target PRO mRNA (sense) or PRO DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

[0827] Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable in vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

[0828] Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increase affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

[0829] Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example,  $\text{CaPO}_4$ -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either in vivo or ex vivo. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

[0830] Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

[0831] Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target



nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

[0832] Antisense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

[0833] The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO coding sequences.

[0834] Nucleotide sequences encoding a PRO can also be used to construct hybridization probes for mapping the gene which encodes that PRO and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as in situ hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

[0835] When the coding sequences for PRO encode a protein which binds to mother protein (example, where the PRO is a receptor), the PRO can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO or a receptor for PRO. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

[0836] Nucleic acids which encode PRO or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO. Methods for generating transgenic animals, particu-

larly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

[0837] Alternatively, non-human homologues of PRO can be used to construct a PRO "knock out" animal which has a defective or altered gene encoding PRO as a result of homologous recombination between the endogenous gene encoding PRO and altered genomic DNA encoding PRO introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques. A portion of the genomic DNA encoding PRO can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a "knock out" animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-out animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO polypeptide.

[0838] Nucleic acid encoding the PRO polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve in vivo synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. "Gene therapy" includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes in vivo. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concen-

trations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

[0839] There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred in vivo gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808-813 (1992).

[0840] The PRO polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

[0841] The nucleic acid molecules encoding the PRO polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO nucleic acid molecule of the present invention can be used as a chromosome marker.

[0842] The PRO polypeptides and nucleic acid molecules of the present invention may also be used for tissue typing, wherein the PRO polypeptides of the present invention may be differentially expressed in one tissue as compared to another. PRO nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

[0843] The PRO polypeptides described herein may also be employed as therapeutic agents. The PRO polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edi-

tion, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, PLURONICS™ or PEG.

[0844] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

[0845] Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0846] The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

[0847] Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

[0848] When in vivo administration of a PRO polypeptide or agonist or antagonist thereof is employed, normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 µg/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue.

[0849] Where sustained-release administration of a PRO polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO polypeptide, microencapsulation of the PRO polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and

MN rgp120. Johnson et al., *Nat. Med.*, 2:795-799 (1996); Yasuda, *Biomed. Ther.*, 27:1221-1223 (1993); Hora et al., *Bio/Technology*, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

**[0850]** The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), *Biodegradable Polymers as Drug Delivery Systems* (Marcel Dekker: New York, 1990), pp. 1-41.

**[0851]** This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

**[0852]** The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

**[0853]** All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

**[0854]** In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally

non-immobilized component does not carry a label, complexing can be detected, -for example, by using a labeled antibody specifically binding the immobilized complex.

**[0855]** If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature (London)*, 340:245-246 (1989); Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-lacZ reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for  $\beta$ -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

**[0856]** Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

**[0857]** To assay for antagonists, the PRO polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential

antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., *Current Protocols in Immun.*, 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

[0858] As an alternative approach for receptor identification, labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

[0859] In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

[0860] More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

[0861] Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the

mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix—see Lee et al., *Nucl. Acids Res.*, 6:3073 (1979); Cooney et al., *Science*, 241: 456 (1988); Dervan et al., *Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into the PRO polypeptide (antisense—Okano, *Neurochem.*, 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression* (CRC Press: Boca Raton, Fla., 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

[0862] Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

[0863] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, *Current Biology*, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published Sep. 18, 1997).

[0864] Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, supra.

[0865] These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

[0866] With regard to the PRO211 and PRO217 polypeptide, therapeutic indications include disorders associated with the preservation and maintenance of gastrointestinal mucosa and the repair of acute and chronic mucosal lesions (e.g., enterocolitis, Zollinger-Ellison syndrome, gastrointestinal ulceration and congenital microvillus atrophy), skin diseases associated with abnormal keratinocyte differentiation (e.g., psoriasis, epithelial cancers such as lung squamous cell carcinoma, epidermoid carcinoma of the vulva and gliomas).

[0867] Since the PRO232 polypeptide and nucleic acid encoding it possess sequence homology to a cell surface stem cell antigen and its encoding nucleic acid, probes based

upon the PRO232 nucleotide sequence may be employed to identify other novel stem cell surface antigen proteins. Soluble forms of the PRO232 polypeptide may be employed as antagonists of membrane bound PRO232 activity both in vitro and in vivo. PRO232 polypeptides may be employed in screening assays designed to identify agonists or antagonists of the native PRO232 polypeptide, wherein such assays may take the form of any conventional cell-type or biochemical binding assay. Moreover, the PRO232 polypeptide may serve as a molecular marker for the tissues in which the polypeptide is specifically expressed.

[0868] With regard to the PRO187 polypeptides disclosed herein, FGF-8 has been implicated in cellular differentiation and embryogenesis, including the patterning which appears during limb formation. FGF-8 and the PRO187 molecules of the invention therefore are likely to have potent effects on cell growth and development. Diseases which relate to cellular growth and differentiation are therefore suitable targets for therapeutics based on functionality similar to FGF-8. For example, diseases related to growth or survival of nerve cells including Parkinson's disease, Alzheimer's disease, ALS, neuropathies. Additionally, disease related to uncontrolled cell growth, e.g., cancer, would also be expected therapeutic targets.

[0869] With regard to the PRO265 polypeptides disclosed herein, other methods for use with PRO265 are described in U.S. Pat. No. 5,654,270 to Ruoslahti et al. In particular, PRO265 can be used in comparison with the fibromodulin disclosed therein to compare its effects on reducing dermal scarring and other properties of the fibromodulin described therein including where it is located and with what it binds and does not.

[0870] The PRO219 polypeptides of the present invention which play a regulatory role in the blood coagulation cascade may be employed in vivo for therapeutic purposes as well as for in vitro purposes. Those of ordinary skill in the art will well know how to employ PRO219 polypeptides for such uses.

[0871] The PRO246 polypeptides of the present invention which serve as cell surface receptors for one or more viruses will find other uses. For example, extracellular domains derived from these PRO246 polypeptides may be employed therapeutically in vivo for lessening the effects of viral infection. Those PRO246 polypeptides which serves as tumor specific antigens may be exploited as therapeutic targets for anti-tumor drugs, and the like. Those of ordinary skill in the art will well know how to employ PRO246 polypeptides for such uses.

[0872] Assays in which connective growth factor and other growth factors are usually used should be performed with PRO261. An assay to determine whether TGF beta induces PRO261, indicating a role in cancer is performed as known in the art. Wound repair and tissue growth assays are also performed with PRO261. The results are applied accordingly.

[0873] PRO228 polypeptides should be used in assays in which EMR1, CD97 and latrophilin would be used in to determine their relative activities. The results can be applied accordingly. For example, a competitive binding assay with PRO228 and CD97 can be performed with the ligand for CD97, CD55.

[0874] Native PRO533 is a 216 amino acid polypeptide of which residues 1-22 are the signal sequence. Residues 3 to 216 have a Blast score of 509, corresponding to 53% homology to fibroblast growth factor. At the nucleotide level, DNA47412, the EST from which PCR oligos were generated to isolate the full length DNA49435-1219, has been observed to map to 11p15. Sequence homology to the 11p15 locus would indicate that PRO533 may have utility in the treatment of Usher Syndrome or Atrophia areata.

[0875] As mentioned previously, fibroblast growth factors can act upon cells in both a mitogenic and non-mitogenic manner. These factors are mitogenic for a wide variety of normal diploid mesoderm-derived and neural crest-derived cells, inducing granulosa cells, adrenal cortical cells, chondrocytes, myoblasts, corneal and vascular endothelial cells (bovine or human), vascular smooth muscle cells, lens, retina and prostatic epithelial cells, oligodendrocytes, astrocytes, chondrocytes, myoblasts and osteoblasts.

[0876] Non-mitogenic actions of fibroblast growth factors include promotion of cell migration into a wound area (chemotaxis), initiation of new blood vessel formation (angiogenesis), modulation of nerve regeneration and survival (neurotrophism), modulation of endocrine functions, and stimulation or suppression of specific cellular protein expression, extracellular matrix production and cell survival. Baird, A. & Bohlen, P., *Handbook of Exp. Pharmacol.* 95(1): 369-418 (1990). These properties provide a basis for using fibroblast growth factors in therapeutic approaches to accelerate wound healing, nerve repair, collateral blood vessel formation, and the like. For example, fibroblast growth factors, have been suggested to minimize myocardium damage in heart disease and surgery (U.S. Pat. No. 4,378,437).

[0877] Since the PRO245 polypeptide and nucleic acid encoding it possess sequence homology to a transmembrane protein tyrosine kinase protein and its encoding nucleic acid, probes based upon the PRO245 nucleotide sequence may be employed to identify other novel transmembrane tyrosine kinase proteins. Soluble forms of the PRO245 polypeptide may be employed as antagonists of membrane bound PRO245 activity both in vitro and in vivo. PRO245 polypeptides may be employed in screening assays designed to identify agonists or antagonists of the native PRO245 polypeptide, wherein such assays may take the form of any conventional cell-type or biochemical binding assay. Moreover, the PRO245 polypeptide may serve as a molecular marker for the tissues in which the polypeptide is specifically expressed.

[0878] PRO220, PRO221 and PRO227 all have leucine rich repeats. Additionally, PRO220 and PRO221 have homology to SLIT and leucine rich repeat protein. Therefore, these proteins are useful in assays described in the literature, supra, wherein the SLIT and leucine rich repeat protein are used. Regarding the SLIT protein, PRO227 can be used in an assay to determine the affect of PRO227 on neurodegenerative disease. Additionally, PRO227 has homology to human glycoprotein V. In the case of PRO227, this polypeptide is used in an assay to determine its affect on bleeding, clotting, tissue repair and scarring.

[0879] The PRO266 polypeptide can be used in assays to determine if it has a role in neurodegenerative diseases or their reversal.

[0880] PRO269 polypeptides and portions thereof which effect the activity of thrombin may also be useful for in vivo therapeutic purposes, as well as for various in vitro applications. In addition, PRO269 polypeptides and portions thereof may have therapeutic use as an antithrombotic agent with reduced risk for hemorrhage as compared with heparin. Peptides having homology to thrombomodulin are particularly desirable.

[0881] PRO287 polypeptides and portions thereof which effect the activity of bone morphogenic protein "BMP1"/procollagen C-proteinase (PCP) may also be useful for in vivo therapeutic purposes, as well as for various in vitro applications. In addition, PRO287 polypeptides and portions thereof may have therapeutic applications in wound healing and tissue repair. Peptides having homology to procollagen C-proteinase enhancer protein and its precursor may also be used to induce bone and/or cartilage formation and are therefore of particular interest to the scientific and medical communities.

[0882] Therapeutic indications for PRO214 polypeptides include disorders associated with the preservation and maintenance of gastrointestinal mucosa and the repair of acute and chronic mucosal lesions (e.g., enterocolitis, Zollinger-Ellison syndrome, gastrointestinal ulceration and congenital microvillus atrophy), skin diseases associated with abnormal keratinocyte differentiation (e.g., psoriasis, epithelial cancers such as lung squamous cell carcinoma, epidermoid carcinoma of the vulva and gliomas).

[0883] Studies on the generation and analysis of mice deficient in members of the TGF-superfamily are reported in Matzuk, *Trends in Endocrinol, and Metabol.*, 6: 120-127 (1995).

[0884] The PRO317 polypeptide, as well as PRO317-specific antibodies, inhibitors, agonists, receptors, or their analogs, herein are useful in treating PRO317-associated disorders. Hence, for example, they may be employed in modulating endometrial bleeding angiogenesis, and may also have an effect on kidney tissue. Endometrial bleeding can occur in gynecological diseases such as endometrial cancer as abnormal bleeding. Thus, the compositions herein may find use in diagnosing and treating abnormal bleeding conditions in the endometrium, as by reducing or eliminating the need for a hysterectomy. The molecules herein may also find use in angiogenesis applications such as anti-tumor indications for which the antibody against vascular endothelial growth factor is used, or, conversely, ischemic indications for which vascular endothelial growth factor is employed.

[0885] Bioactive compositions comprising PRO317 or agonists or antagonists thereof may be administered in a suitable therapeutic dose determined by any of several methodologies including clinical studies on mammalian species to determine maximal tolerable dose and on normal human subjects to determine safe dose. Additionally, the bioactive agent may be complexed with a variety of well established compounds or compositions which enhance stability or pharmacological properties such as half-life. It is contemplated that the therapeutic, bioactive composition may be delivered by intravenous infusion into the bloodstream or any other effective means which could be used for treating problems of the kidney, uterus, endometrium, blood vessels, or related tissue, e.g., in the heart or genital tract.

[0886] Dosages and administration of PRO317, PRO317 agonist, or PRO317 antagonist in a pharmaceutical composition may be determined by one of ordinary skill in the art of clinical pharmacology or pharmacokinetics. See, for example, Mordenti and Rescigno, *Pharmaceutical Research*, 9:17-25 (1992); Morenti et al., *Pharmaceutical Research*, 8:1351-1359 (1991); and Mordenti and Chappell, "The use of interspecies scaling in toxicokinetics" in *Toxicokinetics and New Drug Development*, Yacobi et al. (eds) (Pergamon Press: NY, 1989), pp. 42-96. An effective amount of PRO317, PRO317 agonist, or PRO317 antagonist to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the mammal. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage might range from about 10 ng/kg to up to 100 mg/kg of the mammal's body weight or more per day, preferably about 1  $\mu$ g/kg/day to 10 mg/kg/day. Typically, the clinician will administer PRO317, PRO317 agonist, or PRO317 antagonist, until a dosage is reached that achieves the desired effect for treatment of the above mentioned disorders.

[0887] PRO317 or an PRO317 agonist or PRO317 antagonist may be administered alone or in combination with another to achieve the desired pharmacological effect. PRO317 itself, or agonists or antagonists of PRO317 can provide different effects when administered therapeutically. Such compounds for treatment will be formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8, more preferably 6 to 8, although the pH may vary according to the characteristics of the PRO317, agonist, or antagonist being formulated and the condition to be treated. Characteristics of the treatment compounds include solubility of the molecule, half-life, and antigenicity/immunogenicity; these and other characteristics may aid in defining an effective carrier.

[0888] PRO317 or PRO317 agonists or PRO317 antagonists may be delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol, transdermal patch and bandage; injectable, intravenous, and lavage formulations; and orally administered liquids and pills, particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration will be determined by the attending physician and will vary according to each specific situation.

[0889] Such determinations of administration are made by considering multiple variables such as the condition to be treated, the type of mammal to be treated, the compound to be administered, and the pharmacokinetic profile of the particular treatment compound. Additional factors which may be taken into account include disease state (e.g. severity) of the patient, age, weight, gender, diet, time of administration, drug combination, reaction sensitivities, and tolerance/response to therapy. Long-acting treatment compound formulations (such as liposomally encapsulated PRO317 or PEGylated PRO317 or PRO317 polymeric microspheres, such as polylactic acid-based microspheres) might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular treatment compound.

[0890] Normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1  $\mu\text{g}/\text{kg}/\text{day}$  to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting the uterus, for example, may necessitate delivery in a manner different from that to another organ or tissue, such as cardiac tissue.

[0891] Where sustained-release administration of PRO317 is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of PRO317, microencapsulation of PRO317 is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and MN rgp120. Johnson et al., *Nat. Med.*, 2: 795-799 (1996); Yasuda, *Biomed. Ther.*, 27: 1221-1223 (1993); Hora et al., *Bio/Technology*, 8: 755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

[0892] It is contemplated that conditions or diseases of the uterus, endometrial tissue, or other genital tissues or cardiac tissues may precipitate damage that is treatable with PRO317 or PRO317 agonist where PRO317 expression is reduced in the diseased state; or with antibodies to PRO317 or other PRO317 antagonists where the expression of PRO317 is increased in the diseased state. These conditions or diseases may be specifically diagnosed by the probing tests discussed above for physiologic and pathologic problems which affect the function of the organ.

[0893] The PRO317, PRO317 agonist, or PRO317 antagonist may be administered to a mammal with another biologically active agent, either separately or in the same formulation to treat 21 common indication for which they are appropriate. For example, it is contemplated that PRO317 can be administered together with EBAF-1 for those indications on which they demonstrate the same qualitative biological effects. Alternatively, where they have opposite effects, EBAF-1 may be administered together with an antagonist to PRO317, such as an anti-PRO317 antibody. Further, PRO317 may be administered together with VEGF for coronary ischemia where such indication is warranted, or with an anti-VEGF for cancer as warranted, or, conversely, an antagonist to PRO317 may be administered with VEGF for coronary ischemia or with anti-VEGF to treat cancer as warranted. These administrations would be in effective amounts for treating such disorders.

[0894] Native PRO301 (SEQ ID NO:119) has a Blast score of 246 and 30% homology at residues 24 to 282 of FIG. 44 with A33\_HUMAN, an A33 antigen precursor. A33 antigen precursor, as explained in the Background is a tumor-specific antigen, and as such, is a recognized marker and therapeutic target for the diagnosis and treatment of colon cancer. The expression of tumor-specific antigens is often associated with the progression of neoplastic tissue

disorders. Native PRO301 (SEQ ID NO:119) and A33\_HUMAN also show a Blast score of 245 and 30% homology at residues 21 to 282 of FIG. 44 with A33\_HUMAN, the variation dependent upon how spaces are inserted into the compared sequences. Native PRO301 (SEQ ID NO:119) also has a Blast score of 165 and 29% homology at residues 60 to 255 of FIG. 44 with HS46KDA\_1, a human coxsackie and adenovirus receptor protein, also known as cell surface protein HCAR. This region of PRO301 also shows a similar Blast score and homology with HSU90716\_1. Expression of such proteins is usually associated with viral infection and therapeutics for the prevention of such infection may be accordingly conceived. As mentioned in the Background, the expression of viral receptors is often associated with neoplastic tumors.

[0895] Therapeutic uses for the PRO234 polypeptides of the invention includes treatments associated with leukocyte homing or the interaction between leukocytes and the endothelium during an inflammatory response. Examples include asthma, rheumatoid arthritis, psoriasis and multiple sclerosis.

[0896] Since the PRO231 polypeptide and nucleic acid encoding it possess sequence homology to a putative acid phosphatase and its encoding nucleic acid, probes based upon the PRO231 nucleotide sequence may be employed to identify other novel phosphatase proteins. Soluble forms of the PRO231 polypeptide may be employed as antagonists of membrane bound PRO231 activity both in vitro and in vivo. PRO231 polypeptides may be employed in screening assays designed to identify agonists or antagonists of the native PRO231 polypeptide, wherein such assays may take the form of any conventional cell-type or biochemical binding assay. Moreover, the PRO231 polypeptide may serve as a molecular marker for the tissues in which the polypeptide is specifically expressed.

[0897] PRO229 polypeptides can be fused with peptides of interest to determine whether the fusion peptide has an increased half-life over the peptide of interest. The PRO229 polypeptides can be used accordingly to increase the half-life of polypeptides of interest. Portions of PRO229 which cause the increase in half-life are an embodiment of the invention herein.

[0898] PRO238 can be used in assays which measure its ability to reduce substrates, including oxygen and Acetyl-CoA, and particularly, measure PRO238's ability to produce oxygen free radicals. This is done by using assays which have been previously described. PRO238 can further be used to assay for candidates which block, reduce or reverse its reducing abilities. This is done by performing side by side assays where candidates are added in one assay having PRO238 and a substrate to reduce, and not added in another assay, being the same but for the lack of the presence of the candidate.

[0899] PRO233 polypeptides and portions thereof which have homology to reductase may also be useful for in vivo therapeutic purposes, as well as for various other applications. The identification of novel reductase proteins and related molecules may be relevant to a number of human disorders such as inflammatory disease, organ failure, atherosclerosis, cardiac injury, infertility, birth defects, premature aging, AIDS, cancer, diabetic complications and mutations in general. Given that oxygen free radicals and

antioxidants appear to play important roles in a number of disease processes, the identification of new reductase proteins and reductase-like molecules is of special importance in that such proteins may serve as potential therapeutics for a variety of different human disorders. Such polypeptides may also play important roles in biotechnological and medical research, as well as various industrial applications. As a result, there is particular scientific and medical interest in new molecules, such as PRO233.

[0900] The PRO223 polypeptides of the present invention which exhibit serine carboxypeptidase activity may be employed in vivo for therapeutic purposes as well as for in vitro purposes. Those of ordinary skill in the art will well know how to employ PRO223 polypeptides for such uses.

[0901] PRO235 polypeptides and portions thereof which may be involved in cell adhesion are also useful for in vivo therapeutic purposes, as well as for various in vitro applications. In addition, PRO235 polypeptides and portions thereof may have therapeutic applications in disease states which involve cell adhesion. Given the physiological importance of cell adhesion mechanisms in vivo, efforts are currently being under taken to identify new, native proteins which are involved in cell adhesion. Therefore, peptides having homology to plexin are of particular interest to the scientific and medical communities.

[0902] Because the PRO236 and PRO262 polypeptides disclosed herein are homologous to various known  $\beta$ -galactosidase proteins, the PRO236 and PRO262 polypeptides disclosed herein will find use in conjugates of monoclonal antibodies and the polypeptide for specific killing of tumor cells by generation of active drug from a galactosylated prodrug (e.g., the generation of 5-fluorouridine from the prodrug  $\beta$ -D-galactosyl-5-fluorouridine). The PRO236 and PRO262 polypeptides disclosed herein may also find various uses both in vivo and in vitro, wherein those uses will be similar or identical to uses for which  $\beta$ -galactosidase proteins are now employed. Those of ordinary skill in the art will well know how to employ PRO236 and PRO262 polypeptides for such uses.

[0903] PRO239 polypeptides and portions thereof which have homology to densin may also be useful for in vivo therapeutic purposes, as well as for various in vitro applications. In addition, PRO239 polypeptides and portions thereof may have therapeutic applications in disease states which involve synaptic mechanisms, regeneration or cell adhesion. Given the physiological importance of synaptic processes, regeneration and cell adhesion mechanisms in vivo, efforts are currently being under taken to identify new, native proteins which are involved in synaptic machinery and cell adhesion. Therefore, peptides having homology to densin are of particular interest to the scientific and medical communities.

[0904] The PRO260 polypeptides described herein can be used in assays to determine their relation to fucosidase. In particular, the PRO260 polypeptides can be used in assays in determining their ability to remove fucose or other sugar residues from proteoglycans. The PRO260 polypeptides, can be assayed to determine if they have any functional or locational similarities as fucosidase. The PRO260 polypeptides can then be used to regulate the systems in which they are integral.

[0905] PRO263 can be used in assays wherein CD44 antigen is generally used to determine PRO263 activity relative to that of CD44. The results can be used accordingly.

[0906] PRO270 polypeptides and portions thereof which effect reduction-oxidation (redox) state may also be useful for in vivo therapeutic purposes, as well as for various in vitro applications. More specifically, PRO270 polypeptides may affect the expression of a large variety of genes thought to be involved in the pathogenesis of AIDS, cancer, atherosclerosis, diabetic complications and in pathological conditions involving oxidative stress such as stroke and inflammation. In addition, PRO270 polypeptides and portions thereof may affect the expression of a genes which have a role in apoptosis. Therefore, peptides having homology to thioredoxin are particularly desirable to the scientific and medical communities.

[0907] PRO272 polypeptides and portions thereof which possess the ability to bind calcium may also have numerous in vivo therapeutic uses, as well as various in vitro applications. Therefore, peptides having homology to reticulocalbin are particularly desirable. Those with ordinary skill in the art will know how to employ PRO272 polypeptides and portions thereof for such purposes.

[0908] PRO294 polypeptides and portions thereof which have homology to collagen may also be useful for in vivo therapeutic purposes, as well as for various other applications. The identification of novel collagens and collage-like molecules may have relevance to a number of human disorders. Thus, the identification of new collagens and collage-like molecules is of special importance in that such proteins may serve as potential therapeutics for a variety of different human disorders. Such polypeptides may also play important roles in biotechnological and medical research as well as various industrial applications. Given the large number of uses for collagen, there is substantial interest in polypeptides with homology to the collagen molecule.

[0909] PRO295 polypeptides and portions thereof which have homology to integrin may also be useful for in vivo therapeutic purposes, as well as for various other applications. The identification of novel integrins and integrin-like molecules may have relevance to a number of human disorders such as modulating the binding or activity of cells of the immune system. Thus, the identification of new integrins and integrin-like molecules is of special importance in that such proteins may serve as potential therapeutics ifor a variety of different human disorders. Such polypeptides may also play important roles in biotechnological and medical research as well as various industrial applications. As a result, there is particular scientific and medical interest in new molecules, such as PRO295.

[0910] As the PRO293 polypeptide is clearly a leucine rich repeat polypeptide homologue, the peptide can be used in all applications that the known NLRR-1 and NLRR-2 polypeptides are used. The activity can be compared between these peptides and thus applied accordingly.

[0911] The PRO247 polypeptides described herein can be used in assays in which densin is used to determine the activity of PRO247 relative to densin or these other proteins. The results can be used accordingly in diagnostics and/or therapeutic applications with PRO247.

[0912] PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides of the present invention which pos-



sess protease activity may be employed both in vivo for therapeutic purposes and in vitro. Those of ordinary skill in the art will well know how to employ the PRO302, PRO303, PRO304, PRO307 and PRO343 polypeptides of the present invention for such purposes.

[0913] PRO328 polypeptides and portions thereof which have homology to GLIP and CRISP may also be useful for in vivo therapeutic purposes, as well as for various other applications. The identification of novel GLIP and CRISP-like molecules may have relevance to a number of human disorders which involve transcriptional regulation or are over expressed in human tumors. Thus, the identification of new GLIP and CRISP-like molecules is of special importance in that such proteins may serve as potential therapeutics for a variety of different human disorders. Such polypeptides may also play important roles in biotechnological and medical research as well as in various industrial applications. As a result, there is particular scientific and medical interest in new molecules, such as PRO328.

[0914] Uses for PRO335, PRO331 or PRO326 including uses in competitive assays with LIG-1, ALS and decorin to determine their relative activities. The results can be used accordingly. PRO335, PRO331 or PRO326 can also be used in assays where LIG-1 would be used to determine if the same effects are incurred.

[0915] PRO332 contains GAG repeat (GKEK) at amino acid positions 625-628 in FIG. 108 (SEQ ID NO:310). Slippage in such repeats can be associated with human disease. Accordingly, PRO332 can use useful for the treatment of such disease conditions by gene therapy, i.e. by introduction of a gene containing the correct GKEK sequence motif.

[0916] Other uses of PRO334 include use in assays in which fibrillin or fibulin would be used to determine the relative activity of PRO334 to fibrillin or fibulin. In particular, PRO334 can be used in assays which require the mechanisms imparted by epidermal growth factor repeats.

[0917] Native PRO346 (SEQ ID NO:320) has a Blast score of 230, corresponding to 27% homology between amino acid residues 21 to 343 with residues 35 to 1040 CGM6\_HUMAN, a carcinoembryonic antigen cgm6 precursor. This homology region includes nearly all but 2 N-terminal extracellular domain residues, including an immunoglobulin superfamily homology at residues 148 to 339 of PRO346 in addition to several transmembrane residues (340-343). Carcinoembryonic antigen precursor, as explained in the Background is a tumor-specific antigen, and as such, is a recognized marker and therapeutic target for the diagnosis and treatment of colon cancer. The expression of tumor-specific antigens is often associated with the progression of neoplastic tissue disorders. Native PRO346 (SEQ ID NO:320) and P\_W06874, a human carcinoembryonic antigen CEA-d have a Blast score of 224 and homology of 28% between residues 2 to 343 and 67 to 342, respectively. This homology includes the entire extracellular domain residues of native PRO346, minus the initiator methionine (residues 2 to 18) as well as several transmembrane residues (340-343).

[0918] PRO268 polypeptides which have protein disulfide isomerase activity will be useful for many applications where protein disulfide isomerase activity is desirable

including, for example, for use in promoting proper disulfide bond formation in recombinantly produced proteins so as to increase the yield of correctly folded protein. Those of ordinary skill in the art will readily know how to employ such PRO268 polypeptides for such purposes.

[0919] PRO330 polypeptides of the present invention which possess biological activity related to that of the prolyl 4-hydroxylase alpha subunit protein may be employed both in vivo for therapeutic purposes and in vitro. Those of ordinary skill in the art will well know how to employ the PRO330 polypeptides of the present invention for such purposes.

[0920] Uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

[0921] F. Anti-PRO Antibodies

[0922] The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

[0923] 1. Polyclonal Antibodies

[0924] The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0925] 2. Monoclonal Antibodies

[0926] The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

[0927] The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press,

(1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

**[0928]** Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection, Manassas, Va. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

**[0929]** The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980).

**[0930]** After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

**[0931]** The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

**[0932]** The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce

immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Pat. No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

**[0933]** The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

**[0934]** In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

**[0935]** 3. Human and Humanized Antibodies

**[0936]** The anti-PRO antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

**[0937]** Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Human-

ization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science* 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0938] Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *BioTechnology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

#### [0939] 4. Bispecific Antibodies

[0940] Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

[0941] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published May 13, 1993, and in Traunecker et al., *EMBO J.*, 10:3655-3659 (1991).

[0942] Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The

fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

[0943] According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0944] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')<sub>2</sub> bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0945] Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[0946] Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by

gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain ( $V_H$ ) connected to a light-chain variable domain ( $V_L$ ) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the  $V_H$  and  $V_L$  domains of one fragment are forced to pair with the complementary  $V_L$  and  $V_H$  domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994).

[0947] Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

[0948] Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc $\gamma$ R), such as Fc $\gamma$ RI (CD64), Fc $\gamma$ RII (CD32) and Fc $\gamma$ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

#### [0949] 5. Heteroconjugate Antibodies

[0950] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Pat. No. 4,676,980], and for treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Pat. No. 4,676,980.

#### [0951] 6. Effector Function Engineering

[0952] It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J.*

*Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

#### [0953] 7. Immunoconjugates

[0954] The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0955] Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

[0956] Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol)propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azido-benzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science*, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

[0957] In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agents and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionucleotide).

#### [0958] 8. Immunoliposomes

[0959] The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA*, 77:

4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

[0960] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., *J. Biol. Chem.*, 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., *J. National Cancer Inst.*, 81(19): 1484 (1989).

#### [0961] 9. Pharmaceutical Compositions of Antibodies

[0962] Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

[0963] If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0964] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nano-capsules) or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, supra.

[0965] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[0966] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations as include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for

example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

#### [0967] G. Uses for anti-PRO Antibodies

[0968] The anti-PRO antibodies of the invention have various utilities. For example, anti-PRO antibodies may be used in diagnostic assays for PRO, e.g., detecting its expression in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

[0969] Anti-PRO antibodies also are useful for the affinity purification of PRO from recombinant cell culture or natural sources. In this process, the antibodies against PRO are immobilized on a suitable support, such as Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO from the antibody.

[0970] The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

[0971] All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

#### EXAMPLES

[0972] Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Rockville, Md.

##### Example 1

[0973] Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

[0974] The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhoff, GenBank), and proprietary databases (e.g. LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST2 (Altschul, and Gish, *Methods in Enzymology* 266: 460-80 (1996); <http://blast.wustl.edu/blast/README.html>) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a Blast score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.).

[0975] Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

[0976] Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward (.f) and reverse (.r) PCR primers generally range from 20 to 30 nucleotides and are often designed to

give a PCR product of about 100-1000 bp in length. The probe (.p) sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

[0977] The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

##### Example 2

[0978] Isolation of cDNA Clones Encoding PRO211 and PRO217

[0979] Consensus DNA sequences were assembled as described in Example 1 above and were designated as DNA28730 and DNA28760, respectively. Based on these consensus sequences, oligonucleotides were synthesized and used to identify by PCR a cDNA library that contained the sequences of interest and for use as probes to isolate a clone of the full-length coding sequence for the PRO211 and PRO217 polypeptides. The libraries used to isolate DNA32292-1131 and DNA33094-1131 were fetal lung libraries.

[0980] cDNA clones were sequenced in their entirety. The entire nucleotide sequences of PRO211 (DNA32292-1131) and PRO217 (UNQ191) are shown in **FIG. 1** (SEQ ID NO: 1) and **FIG. 3** (SEQ ID NO:3), respectively. The predicted polypeptides are 353 and 379 amino acid in length, respectively, with respective molecular weights of approximately 38,190 and 41,520 daltons.

[0981] The oligonucleotide sequences used in the above procedures were the following:

```

28730.p (OLI 516)
5'-AGGGAGCACGGACAGTGTGCAGATGTGGACGAGTGCTCACTAGCA-3' (SEQ ID NO:5)

28730.f (OLI 517)
5'-AGAGTGTATCTCTGGCTACGC-3' (SEQ ID NO:6)

28730.r (OLI 518)
5'-TAAGTCCGGCACATTACAGGTC-3' (SEQ ID NO:7)

28760.p (OLI 617)
5'-CCCACGATGTATGAATGGTGGACTTTGTGTGACTCCTGGTTTCTGCATC-3' (SEQ ID NO:8)

28760.f (OLI 618)
5'-AAAGACGCATCTCGAGTGTCC-3' (SEQ ID NO:9)

28760.r (OLI 619)
5'-TGCTGATTCACACTGCTCTCCC-3' (SEQ ID NO:10)

```

## Example 3

[0982] Isolation of cDNA Clones Encoding Human PRO230

[0983] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA30857. An EST proprietary to Genentech was employed in the consensus assembly. The EST is designated as DNA20088 and has the nucleotide sequence shown in FIG. 7 (SEQ ID NO:13).

[0984] Based on the DNA30857 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO230.

[0985] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-TTCGAGCCTCTGAGAAGTGGCC-3' (SEQ ID NO:14)

reverse PCR primer 5'-GGCGTATCTCTGCCTCCC-3' (SEQ ID NO:15)

[0986] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30857 sequence which had the following nucleotide sequence

forward PCR primer 5'-TGCTGTGCTACTCCTGCAAAGCCC-3' (SEQ ID NO:19)

reverse PCR primer 5'-TGCACAAGTCGGTGTACAGCACG-3' (SEQ ID NO:20)

[0987] hybridization probe

[0988] 5'-TTCTCCACAGCAGCTGTGGCATC-CGATCGTGTCTCAATCCATCTCTGCG-3' (SEQ ID NO:16)

[0989] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO230 gene using the probe oligonucleotide and one of the PCR primers.

[0990] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO230 (herein designated as DNA33223-1136 and the derived protein sequence for PRO230.

[0991] The entire nucleotide sequence of DNA33223-1136 is shown in FIG. 5 (SEQ ID NO:11). Clone

DNA33223-1136 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 100-103 and ending at the stop codon at nucleotide positions 1501-1503 (FIG. 5; SEQ ID NO:11). The predicted polypeptide precursor is 467 amino acids long (FIG. 6).

## Example 4

[0992] Isolation of cDNA Clones Encoding Human PRO232

[0993] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA30935. Based on the DNA30935 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence

of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO232.

[0994] A pair of PCR primers (forward and reverse) were synthesized:

[0995] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30935 sequence which had the following nucleotide sequence

[0996] hybridization probe

[0997] 5'-AGCAACGAGGACTGCCTGCAGGTG-GAGAACTGCACCCAGCTGGG-3' (SEQ ID NO:21)

[0998] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO232 gene using the probe oligonucleotide and one of the PCR primers.

[0999] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1000] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO232 [herein designated as DNA34435-1140] and the derived protein sequence for PRO232.

[1001] The entire nucleotide sequence of DNA34435-1140 is shown in FIG. 8 (SEQ ID NO:17). Clone

DNA34435-1140 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 17-19 and ending at the stop codon at nucleotide positions 359-361 (**FIG. 8**; SEQ ID NO:17). The predicted polypeptide precursor is 114 amino acids long (**FIG. 9**). Clone DNA:34435-1140 has been deposited with ATCC on Sep. 16, 1997 and is assigned ATCC deposit no. ATCC 209250.

[1002] Analysis of the amino acid sequence of the full-length PRO232 suggests that it possesses 35% sequence identity with a stem cell surface antigen from *Gallus gallus*.

#### Example 5

[1003] Isolation of cDNA Clones Encoding PRO187

[1004] A proprietary expressed sequence tag (EST) DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.) was searched and an EST (#843193) was identified which showed homology to fibroblast growth factor (FGF-8) also known as androgen-induced growth factor. mRNA was isolated from human fetal lung tissue using reagents and protocols from Invitrogen, San Diego, Calif. (Fast Track 2). The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents (e.g., Invitrogen, San Diego, Calif., Life Technologies, Gaithersburg, Md.). The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropri-

ately by gel electrophoresis, and cloned in a defined orientation into the cloning vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, Md. (Super Script Plasmid System). The double-stranded cDNA was sized to greater than 1000 bp and the SalI/NotI linked cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

[1005] Several libraries from various tissue sources were screened by PCR amplification with the following oligonucleotide probes:

IN843193.f (OLI315)  
5'-CAGTACGTGAGGGACCAGGGCGCCATGA-3' (SEQ ID NO:24)

IN843193.r (OLI317)  
5'CCGGTGACCTGCACGCTGCTGCCA-3' (SEQ ID NO:25)

[1006] A positive library was then used to isolate clones encoding the PRO187 gene using one of the above oligonucleotides and the following oligonucleotide probe:

[1007] IN843193.p (OLI 316) (SEQ ID NO:26)

[1008] 5'-GCGGATCTGCCGCGCTGCTCANCTG-GTCGGTCAITGGCGCCCT-3'

[1009] A cDNA clone was sequenced in entirety. The entire nucleotide sequence of PRO187 (DNA27864-1155) is shown in **FIG. 10** (SEQ ID NO:22). Clone DNA27864-1155 contains a single open reading frame with an apparent translational initiation site at nucleotide position 1 (**FIG. 10**; SEQ ID NO:22). The predicted polypeptide precursor is 205 amino acids long. Clone DNA27864-1155 has been deposited with the ATCC (designation: DNA27864-1155) and is assigned ATCC deposit no. ATCC 209375.

[1010] Based on a BLAST and FastA sequence alignment analysis (using the ALIGN computer program) of the full-length sequence, the PRO187 polypeptide shows 74% amino acid sequence identity (Blast score 310) to human fibroblast growth factor-8 (androgen-induced growth factor).

#### Example 6

[1011] Isolation of cDNA Clones Encoding PRO265

[1012] A consensus DNA sequence was assembled relative to other EST sequences as described in Example 1 above using phrap. This consensus sequence is herein designated DNA33679. Based on the DNA33679 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO265.

[1013] PCR primers (two forward and one reverse) were synthesized:

forward PCR primer A: 5'-CGGTCTACCTGTATGGCAACC-3' (SEQ ID NO:29);

forward PCR primer B: 5'-GCAGGACAACACGATAAACCCAC-3' (SEQ ID NO:30);

reverse PCR primer 5'-ACGCAGATTTGAGAAGGCTGTC-3' (SEQ ID NO:31)

[1014] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA33679 sequence which had the following nucleotide sequence

[1015] hybridization probe

[1016] 5'-TTCACGGGCTGCTCTTGCCCAGCTCT-TGAAGCTTGAAGAGCTGCAC-3' (SEQ ID NO:32)

[1017] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO265 gene using the probe oligonucleotide and one of the PCR primers.

[1018] RNA for construction of the cDNA libraries was isolated from human a fetal brain library.

[1019] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO265 [herein designated as DNA36350-1158] (SEQ ID NO:27) and the derived protein sequence for PRO265.

[1020] The entire nucleotide sequence of DNA36350-1158 is shown in **FIG. 12** (SEQ ID NO:27). Clone DNA36350-1158 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 352-354 and ending at the stop codon at positions



2332-2334 (**FIG. 12**). The predicted polypeptide precursor is 660 amino acids long (**FIG. 13**). Clone DNA36350-1158 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209378.

[1021] Analysis of the amino acid sequence of the full-length PRO265 polypeptide suggests that portions of it possess significant homology to the fibromodulin and the fibromodulin precursor, thereby indicating that PRO265 may be a novel member of the leucine rich repeat family, particularly related to fibromodulin.

#### Example 7

[1022] Isolation of cDNA Clones Encoding Human PRO219

[1023] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28729. Based on the DNA28729 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO219.

[1024] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-GTGACCCCTGGTTGTGAATACTCC-3' (SEQ ID NO:35)

reverse PCR primer 5'-ACAGCCATGGTCTATAGCTTGG-3' (SEQ ID NO:36)

[1025] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28729 sequence which had the following nucleotide sequence

[1026] hybridization probe

[1027] 5'-GCCTGTCAAGTGTCTGAGGGACACGT-GCTCCGCAGCGATGGGAAG-3' (SEQ ID NO:37)

[1028] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO219 gene using the probe oligonucleotide and one of the PCR primers.

[1029] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1030] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO219 [herein designated as DNA32290-1164] (SEQ ID NO:33) and the derived protein sequence for PRO219.

[1031] The entire nucleotide sequence of DNA32290-1164 is shown in **FIG. 14** (SEQ ID NO:33). Clone DNA32290-1164 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 204-206 and ending at the stop codon at nucleotide positions 2949-2951 (**FIG. 14**). The predicted polypeptide precursor is 915 amino acids long (**FIG. 15**). Clone DNA32290-1164 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209384.

[1032] Analysis of the amino acid sequence of the full-length PRO219 polypeptide suggests that portions of it possess significant homology to the mouse and human matrilin-2 precursor polypeptides.

#### Example 8

[1033] Isolation of cDNA Clones Encoding Human PRO246

[1034] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30955. Based on the DNA30955 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of

interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO246.

[1035] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-AGGGTCTCCAGGAGAAAGACTC-3' (SEQ ID NO:40)

reverse PCR primer 5'-ATTGTGGCCTTGCAGACATAGAC-3' (SEQ ID NO:41)

[1036] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30955 sequence which had the following nucleotide sequence

[1037] hybridization probe

[1038] 5'-GGCCACAGCATCAAAACCTTAGAACT-CAATGTACTGGTTCCTCCAGCTCC-3' (SEQ ID NO:42)

[1039] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO246 gene using the probe oligonucleotide and one of the PCR primers.

[1040] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. DNA sequencing of

the clones isolated as described above gave the full-length DNA sequence for PRO246 [herein designated as DNA35639-1172] (SEQ ID NO:38) and the derived protein sequence for PRO246.

[1041] The entire nucleotide sequence of DNA35639-1172 is shown in **FIG. 16** (SEQ ID NO:38). Clone DNA35639-1172 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 126-128 and ending at the stop codon at nucleotide positions 1296-1298 (**FIG. 16**). The predicted polypeptide precursor is 390 amino acids long (**FIG. 17**). Clone DNA35639-1172 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209396.

[1042] Analysis of the amino acid sequence of the full-length PRO246 polypeptide suggests that it possess significant homology to the human cell surface protein HCAR, thereby indicating that PRO246 may be a novel cell surface virus receptor.

#### Example 9

[1043] Isolation of cDNA Clones Encoding Human PRO228

[1044] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28758. An EST proprietary to Genentech was employed in the consensus assembly. This EST is shown in **FIG. 20** (SEQ ID NO:50) and is herein designated as DNA21951.

[1045] Based on the DNA28758 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO228.

[1046] PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-GGTAATGAGCTCCATTACAG-3' (SEQ ID NO:51)

forward PCR primer 5'-GGAGTAGAAAGCGCATGG-3' (SEQ ID NO:52)

forward PCR primer 5'-CACCTGATACCATGAATGGCAG-3' (SEQ ID NO:53)

reverse PCR primer 5'-CGAGCTCGAATTAATTCG-3' (SEQ ID NO:54)

reverse PCR primer 5'-GGATCTCCTGAGCTCAGG-3' (SEQ ID NO:55)

reverse PCR primer 5'-CCTAGTTGAGTGATCCTTGTAAG-3' (SEQ ID NO:56)

[1047] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28758 sequence which had the following nucleotide sequence

[1048] hybridization probe

[1049] 5'-ATGAGACCCACACCTCATGCCGCTG-TAATCACCTGACACATTTTGCAATT-3' (SEQ ID NO:57)

[1050] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified

above. A positive library was then used to isolate clones encoding the PRO228 gene using the probe oligonucleotide and one of the PCR primers.

[1051] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1052] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO228 [herein designated as DNA33092-1202] (SEQ ID NO:48) and the derived protein sequence for PRO228.

[1053] The entire nucleotide sequence of DNA33092-1202 is shown in **FIG. 18** (SEQ ID NO:48). Clone DNA33092-1202 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 24-26 of SEQ ID NO:48 and ending at the stop codon after nucleotide position 2093 of SEQ ID NO:48. The predicted polypeptide precursor is 690 amino acids long (**FIG. 19**). Clone DNA33092-1202 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209420.

[1054] Analysis of the amino acid sequence of the full-length PRO228 polypeptide suggests that portions of it possess significant homology to the secretin-related proteins CD97 and EMR1 as well as the secretin member, latrophilin, thereby indicating that PRO228 may be a new member of the secretin related proteins.

#### Example 10

[1055] Isolation of cDNA Clones Encoding Human PRO533

[1056] The EST sequence accession number AF007268, a murine fibroblast growth factor (FGF-15) was used to search various public EST databases (e.g., GenBank, Dayhoff, etc.). The search was performed using the computer program BLAST or BLAST2 [Altschul et al., *Methods in Enzymology*, 266:460-480 (1996); <http://blast.wustl.edu/blast/RE-ADME.html>] as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. The search

resulted in a hit with GenBank EST AA220994, which has been identified as stratagene NT2 neuronal precursor 937230.

[1057] Based on the Genbank EST AA220994 sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence. Forward and reverse PCR primers may range from 20 to 30 nucleotides (typically about 24), and are designed to give a PCR product of 100-1000 bp in length. The probe sequences are typically 40-55 bp (typically about 50) in length. In order to screen several libraries for a source

of a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the PCR primers.

[1058] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified below. A positive library was then used to isolate clones encoding the PRO533 gene using the probe oligonucleotide and one of the PCR primers.

[1059] RNA for construction of the cDNA libraries was isolated from human fetal retina. The cDNA libraries used to isolated the cDNA clones were constructed by standard methods using commercially available reagents (e.g., Invitrogen, San Diego, Calif.; Clontech, etc.) The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

[1060] A cDNA clone was sequenced in its entirety. The full length nucleotide sequence of PRO533 is shown in FIG. 21 (SEQ ID NO:58). Clone DNA49435-1219 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 459-461 (FIG. 21; SEQ ID NO:58). The predicted polypeptide precursor is 216 amino acids long. Clone DNA47412-1219 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209480.

[1061] Based on a BLAST-2 and FastA sequence alignment analysis of the full-length sequence, PRO533 shows amino acid sequence identity to fibroblast growth factor (53%).

[1062] The oligonucleotide sequences used in the above procedure were the following:

FGF15.forward: 5'-ATCCGCCAGATGGCTACAATGTGTA-3' (SEQ ID NO:60);

FGF15.probe: 5'-GCCTCCGGTCTCCCTGAGCAGTGCCAAACAGCGGCGAGTGA-3' (SEQ ID NO:61);

FGF15.reverse: 5'-CCAGTCCGGTGACAAGCCAAA-3' (SEQ ID NO:62).

#### Example 11

[1063] Isolation of cDNA Clones Encoding Human PRO245

[1064] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA30954.

[1065] Based on the DNA30954 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO245.

[1066] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-ATCGTTGTGAAGTTAGTGCCCC-3' (SEQ ID NO:65)

reverse PCR primer  
5'-ACCTGCGATATCCAACAGAATTG-3' (SEQ ID NO:66)

[1067] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30954 sequence which had the following nucleotide sequence

[1068] hybridization probe

[1069] 5'-GGAAGAGGATACAGTCACTCTGGAAG-TATTAGTGGCTCCAGCAGTTCC-3' (SEQ ID NO:67)

[1070] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO245 gene using the probe oligonucleotide and one of the PCR primers.

[1071] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO245 [herein designated as DNA35638-1141] and the derived protein sequence for PRO245.

[1072] The entire nucleotide sequence of DNA35638-1141 is shown in FIG. 23 (SEQ ID NO:63). Clone DNA35638-1141 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 89-91 and ending at the stop codon at nucleotide positions 1025-1027 (FIG. 23; SEQ ID NO:63). The predicted polypeptide precursor is 312 amino acids long (FIG. 24). Clone DNA35638-1141 has been deposited with ATCC on Sep. 16, 1997 and is assigned ATCC deposit no. ATCC 209265.

[1073] Analysis of the amino acid sequence of the full-length PRO245 suggests that a portion of it possesses 60% amino acid identity with the human c-myc protein and, therefore, may be a new member of the transmembrane protein receptor tyrosine kinase family.

#### Example 12

[1074] Isolation of cDNA Clones Encoding Human PRO220, PRO221 and PRO227

[1075] (a) PRO220

[1076] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in

Example 1 above, wherein the consensus sequence is designated herein as DNA28749. Based on the DNA28749 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO220.

[1077] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-TCACCTGGAGCCTTTATTGGCC-3' (SEQ ID NO:74)

reverse PCR primer  
5'-ATACCAGCTATAACCAGGCTGCG-3' (SEQ ID NO:75)

[1078] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28749 sequence which had the following nucleotide sequence:

[1079] hybridization probe

[1080] 5'-CAACAGTAAGTGGTTTGTATGCTCTTC-CAAATCTAGAGATTCTGATGATTGGG-3' (SEQ ID NO:76).

[1081] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO220 gene using the probe oligonucleotide and one of the PCR primers.

[1082] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO220 [herein designated as DNA32298-1132 and the derived protein sequence for PRO220.

[1083] The entire nucleotide sequence of DNA32298-1132 is shown in FIG. 25 (SEQ ID NO:68). Clone DNA32298-1132 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 480-482 and ending at the stop codon at nucleotide positions 2604-2606 (FIG. 25). The predicted polypeptide precursor is 708 amino acids long (FIG. 26). Clone DNA32298-1132 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209257.

[1084] Analysis of the amino acid sequence of the full-length PRO220 shows it has homology to member of the leucine rich repeat protein superfamily, including the leucine rich repeat protein and the neuronal leucine-rich repeat protein 1.

[1085] (b) PRO221

[1086] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA28756. Based on the DNA28756 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO221.

[1087] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-CCATGTGTCTCCTCTACAAAG-3' (SEQ ID NO:77)

reverse PCR primer  
5'-GGGAATAGATGTGATCTGATTGG-3' (SEQ ID NO:78)

[1088] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28756 sequence which had the following nucleotide sequence:

[1089] hybridization probe

[1090] 5'-CACCTGTAGCAATGCAAATCTCAAG-GAAATACCTAGAGATCTTCTCCTG-3' (SEQ ID NO:79)

[1091] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO221 gene using the probe oligonucleotide and one of the PCR primers.

[1092] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO221 [herein designated as DNA33089-1132 and the derived protein sequence for PRO221.

[1093] The entire nucleotide sequence of DNA33089-1132 is shown in FIG. 27 (SEQ ID NO:70). Clone DNA33089-1132 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 179-181 and ending at the stop codon at nucleotide positions 956-958 (FIG. 27). The predicted polypeptide precursor is 259 amino acids long (FIG. 28). PRO221 is believed to have a transmembrane region at amino acids 206-225. Clone DNA33089-1132 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209262.

[1094] Analysis of the amino acid sequence of the full-length PRO221 shows it has homology to member of the leucine rich repeat protein superfamily, including the SLIT protein.

[1095] (c) PRO227

[1096] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA28740. Based on the DNA28740 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO227.

[1097] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-AGCAACCGCCTGAAGCTCATCC-3' (SEQ ID NO:80)

reverse PCR primer  
5'-AAGGCGCGGTGAAAGATGTAGACG-3' (SEQ ID NO:81)

[1098] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28740 sequence which had the following nucleotide sequence:

[1099] hybridization probe

[1100] 5'-GACTACATGTTTCAGGACCTGTACAAC-CTCAAGTCACTGGAGGTTGGCGA-3' (SEQ ID NO:82).

[1101] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO227 gene using the probe oligonucleotide and one of the PCR primers.

[1102] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO227 [herein designated as DNA33786-1132 and the derived protein sequence for PRO227.

[1103] The entire nucleotide sequence of DNA33786-1132 is shown in FIG. 29 (SEQ ID NO:72). Clone DNA33786-1132 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 33-35 and ending at the stop codon at nucleotide positions 1893-1895 (FIG. 29). The predicted polypeptide precursor is 620 amino acids long (FIG. 30). PRO227 is believed to have a transmembrane region. Clone DNA33786-1132 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209253.

[1104] Analysis of the amino acid sequence of the full-length PRO221 shows it has homology to member of the leucine rich repeat protein superfamily, including the platelet glycoprotein V precursor and the human glycoprotein V.

#### Example 13

[1105] Isolation of cDNA Clones Encoding Human PRO258

[1106] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28746.

[1107] Based on the DNA28746 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO258.

[1108] PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-GCTAGGAATTCACAGAAGCCC-3' (SEQ ID NO:85)

reverse PCR primer  
5'-AACCTGGAATGTCACCGAGCTG-3' (SEQ ID NO:86)

reverse PCR primer  
5'-CCTAGCACAGTGACGAGGGACTTGGC-3' (SEQ ID NO:87)

[1109] Additionally, synthetic oligonucleotide hybridization probes were constructed from the consensus DNA28740 sequence which had the following nucleotide sequence:

[1110] hybridization probe

[1111] 5'-AAGACACAGCCACCCTAAACTGT-CAGTCTTCTGGGAGCAAGCCTCCAGCC-3' (SEQ ID NO:88)

[1112] 5'-GCCCTGGCAGACGAGGGCGAGTACAC-CTGCTCAATCTTCACTATGCCTGT-3' (SEQ ID NO:89)

[1113] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO258 gene using the probe oligonucleotide and one of the PCR primers.

[1114] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO258 [herein designated as DNA35918-1174] (SEQ ID NO:83) and the derived protein sequence for PRO258.

[1115] The entire nucleotide sequence of DNA35918-1174 is shown in FIG. 31 (SEQ ID NO:83). Clone DNA35918-1174 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 147-149 of SEQ ID NO:83 and ending at the stop codon after nucleotide position 1340 of SEQ ID NO:83 (FIG. 31). The predicted polypeptide precursor is 398 amino acids long (FIG. 32). Clone DNA35918-1174 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209402.

[1116] Analysis of the amino acid sequence of the full-length PRO258 polypeptide suggests that portions of it possess significant homology to the CRTAM and the poliovirus receptor and have an Ig domain, thereby indicating that PRO258 is a new member of the Ig superfamily.

#### Example 14

[1117] Isolation of cDNA Clones Encoding Human PRO266

[1118] An expressed sequence tag database was searched for ESTs having homology to SLIT, resulting in the identification of a single EST sequence designated herein as T73996. Based on the T73996 EST sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO266.

[1119] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-GTTGGATCTGGGCAACAATAAC-3' (SEQ ID NO:92)

reverse PCR primer  
5'-ATTGTTGTGCAGGCTGAGTTTAAG-3' (SEQ ID NO:93)

[1120] Additionally, a synthetic oligonucleotide hybridization probe was constructed which had the following nucleotide sequence

[1121] hybridization probe

[1122] 5'-GGTGGCTATACATGGATAGCAATTAC-C  
CTGGACACGCTGTCCCGGG-3' (SEQ ID NO:94)

[1123] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO266 gene using the probe oligonucleotide and one of the PCR primers.

[1124] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO266 [herein designated as DNA37150-1178] (SEQ ID NO:90) and the derived protein sequence for PRO266.

[1125] The entire nucleotide sequence of DNA37150-1178 is shown in FIG. 33 (SEQ ID NO:90). Clone DNA37150-1178 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 167-169 and ending at the stop codon after nucleotide position 2254 of SEQ ID NO:90. The predicted polypeptide precursor is 696 amino acids long (FIG. 34). Clone DNA37150-1178 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209401.

[1126] Analysis of the amino acid sequence of the full-length PRO266 polypeptide suggests that portions of it possess significant homology to the SLIT protein, thereby indicating that PRO266 may be a novel leucine rich repeat protein.

#### Example 15

[1127] Isolation of cDNA Clones Encoding Human PRO269

[1128] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35705. Based on the DNA35705 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO269.

[1129] Forward and reverse PCR primers were synthesized:

forward PCR primer (.f1)  
5'-TGGAAGGAGATGCGATGCCACCTG-3' (SEQ ID NO:97)

forward PCR primer (.f2)  
5'-TGACCAGTGGGAAGGACAG-3' (SEQ ID NO:98)

forward PCR primer (.f3)  
5'-ACAGAGCAGAGGGTGCCTTG-3' (SEQ ID NO:99)

reverse PCR primer (.r1)  
5'-TCAGGGACAAGTGGTGTCTCTCCC-3' (SEQ ID NO:100)

reverse PCR primer (.r2)  
5'-TCAGGGAAGGAGTGTGCAGTTCTG-3' (SEQ ID NO:101)

[1130] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35705 sequence which had the following nucleotide sequence:

[1131] hybridization probe

[1132] 5'-ACAGCTCCCCGATCTCAGTTACTTG-  
CATCGCGGACGAAATCGGCGCTCGCT-3' (SEQ  
ID NO:102)

[1133] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO269 gene using the probe oligonucleotide and one of the PCR primers.

[1134] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1135] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO269 [herein designated as DNA38260-1180] (SEQ ID NO:95) and the derived protein sequence for PRO269.

[1136] The entire nucleotide sequence of DNA38260-1180 is shown in FIG. 35 (SEQ ID NO:95). Clone DNA38260-1180 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 314-316 and ending at the stop codon at nucleotide positions 1784-1786 (FIG. 35; SEQ ID NO:95). The predicted polypeptide precursor is 490 amino acids long (FIG. 36). Clone DNA38260-1180 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209397.

[1137] Analysis of the amino acid sequence of the full-length PRO269 suggests that portions of it possess significant homology to the human thrombomodulin proteins, thereby indicating that PRO269 may possess one or more thrombomodulin-like domains.

#### Example 16

[1138] Isolation of cDNA Clones Encoding Human PRO287

[1139] A consensus DNA sequence encoding PRO287 was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA28728. Based on the DNA28728 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO287.

[1140] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer  
5'-CCGATTCATAGACCTCGAGAGT-3' (SEQ ID NO:105)

reverse PCR primer  
5'-GTCAAGGAGTCTCCACAATAC-3' (SEQ ID NO:106)

[1141] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28728 sequence which had the following nucleotide sequence

[1142] hybridization probe

[1143] 5'-GTGTACAATGGCCATGCCAATGGC-  
CAGCGCATTGGCCGCTTCTGT-3' (SEQ ID  
NO:107)

[1144] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO287 gene using the probe oligonucleotide and one of the PCR primers.

[1145] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1146] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO287 [herein designated as DNA39969-1185, SEQ ID NO:103] and the derived protein sequence for PRO287.

[1147] The entire nucleotide sequence of DNA39969-1185 is shown in FIG. 37 (SEQ ID NO:103). Clone DNA39969-1185 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 307-309 and ending at the stop codon at nucleotide positions 1552-1554 (FIG. 37; SEQ ID NO:103). The predicted polypeptide precursor is 415 amino acids long (FIG. 38). Clone DNA39969-1185 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209400.

[1148] Analysis of the amino acid sequence of the full-length PRO287 suggests that it may possess one or more procollagen C-proteinase enhancer protein precursor or procollagen C-proteinase enhancer protein-like domains. Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO287 shows nucleic acid sequence identity to procollagen C-proteinase enhancer protein precursor and procollagen C-proteinase enhancer protein (47 and 54%, respectively).

#### Example 17

[1149] Isolation of cDNA Clones Encoding Human PRO214

[1150] A consensus DNA sequence was assembled using phrap as described in Example 1 above. This consensus DNA sequence is designated herein as DNA28744. Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence.

[1151] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified below. A positive library was then used to isolate clones encoding the PRO214 gene using the probe oligonucleotide and one of the PCR primers.

[1152] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.

[1153] A cDNA clone was sequenced in its entirety. The full length nucleotide sequence of DNA32286-1191 is shown in FIG. 39 (SEQ ID NO:108). DNA32286-1191 contains a single open reading frame with an apparent translational initiation site at nucleotide position 103 (FIG.

39; SEQ ID NO:108). The predicted polypeptide precursor is 420 amino acids long (SEQ ID NO:109).

[1154] Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO214 polypeptide shows amino acid sequence identity to HT protein and/or Fibulin (49% and 38%, respectively).

[1155] The oligonucleotide sequences used in the above procedure were the following:

28744.p (OLI555)  
5'-CCTGGCTATCAGCAGGTGGGCTCCAAGTG (SEQ ID NO:110)  
TCTCGATGTGGATGAGTGTGA-3'

28744.f (OLI556)  
5'-ATTCTGCGTGAACACTGAGGGC-3' (SEQ ID NO:111)

28744.r (OLI557)  
5'-ATCTGCTTGTAGCCCTCGGCAC-3' (SEQ ID NO:112)

#### Example 18

[1156] Isolation of cDNA Clones Encoding Human PRO317

[1157] A consensus DNA sequence was assembled using phrap as described in Example 1 above, wherein the consensus sequence is herein designated as DNA28722. Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence. The forward and reverse PCR primers, respectively, synthesized for this purpose were:

(SEQ ID NO:115)  
5'-AGGACTGCCATAACTTGCCTG (OLI489) and

(SEQ ID NO:116)  
5'-ATAGGAGTTGAAGCAGCGCTGC (OLI490).

[1158] The probe synthesized for this purpose was:

[1159] 5'-TGTGTGGACATAGACGAGTGCCGC-  
TACCGTACTGCCAGCACCGC (OLI488) (SEQ ID  
NO:117)

[1160] mRNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1161] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology* (1989), with the PCR primer pair identified above. A positive library was then used to isolate clones containing the PRO317 gene using the probe oligonucleotide identified above and one of the PCR primers.

[1162] A cDNA clone was sequenced in its entirety. The entire nucleotide sequence of DNA33461-1199 (encoding PRO317) is shown in FIG. 41 (SEQ ID NO:113). Clone DNA33461-1199 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 68-70 (FIG. 41; SEQ ID NO:113). The predicted polypeptide precursor is 366 amino acids long. The predicted signal sequence is amino acids 1-18 of FIG. 42 (SEQ ID NO:114). There is one predicted N-linked glycosylation

site at amino acid residue 160. Clone DNA33461-1199 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209367.

[1163] Based on BLAST™ and FastA™ sequence alignment analysis (using the ALIGN™ computer program) of the full-length PRO317 sequence, PRO317 shows the most amino acid sequence identity to EBAF-1 (92%). The results also demonstrate a significant homology between human PRO317 and mouse LEFTY protein. The C-terminal end of the PRO317 protein contains many conserved sequences consistent with the pattern expected of a member of the TGF-superfamily.

[1164] In situ expression analysis in human tissues performed as described below evidences that there is distinctly strong expression of the PRO317 polypeptide in pancreatic tissue.

is shown in FIG. 43 (SEQ ID NO:118). Clone DNA40628-1216 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 52-54 (FIG. 43; SEQ ID NO:118). The predicted polypeptide precursor is 299 amino acids long with a predicted molecular weight of 32,583 daltons and pI of 8.29. Clone DNA40628-1216 has been deposited with ATCC and is assigned ATCC deposit No. ATCC 209432.

[1170] Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO301 shows amino acid sequence identity to A33 antigen precursor (30%) and coxsackie and adenovirus receptor protein (29%).

[1171] The oligonucleotide sequences used in the above procedure were the following:

OLI2162 (35936.f1)	
5'-TCGCGGAGCTGTGTTCTGTTCC-3'	(SEQ ID NO:120)
OLI2163 (35936.p1)	
5'-TGATCGCGATGGGGACAAAGGCGCAAGCTCGAGAGGAACTGTTGTGCCT-3'	(SEQ ID NO:121)
OLI2164 (35936.f2)	
5'-ACACCTGGTTCAAAGATGGG-3'	(SEQ ID NO:122)
OLI2165 (35936.r1)	
5'-TAGGAAGAGTTGCTGAAGGCACGG-3'	(SEQ ID NO:123)
OLI2166 (35936.t3)	
5'-TTGCCTTACTCAGGTGCTAC-3'	(SEQ ID NO:124)
OLI2167 (35936.r2)	
5'-ACTCAGCAGTGGTAGGAAAG-3'	(SEQ ID NO:125)

#### Example 19

[1165] Isolation of cDNA Clones Encoding Human PRO301

[1166] A consensus DNA sequence designated herein as DNA35936 was assembled using phrap as described in Example 1 above. Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence.

[1167] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by

#### Example 20

[1172] Isolation of cDNA Clones Encoding Human PRO224

[1173] A consensus DNA sequence assembled relative to the other identified EST sequences as described in Example 1, wherein the consensus sequence is designated herein as DNA30845. Based on the DNA30845 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO224.

[1174] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer	5'-AAGTTGCAGTGCCGACCAGTGGC-3'	(SEQ ID NO:128)
reverse PCR primer	5'-TTGGTTCCACAGCCGAGCTCGTCG-3'	(SEQ ID NO:129)

PCR amplification with the PCR primer pair identified below. A positive library was then used to isolate clones encoding the PRO301 gene using the probe oligonucleotide and one of the PCR primers.

[1168] RNA for construction of the cDNA libraries was isolated from human fetal kidney.

[1169] A cDNA clone was sequenced in its entirety. The full length nucleotide sequence of native sequence PRO301

[1175] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30845 sequence which had the following nucleotide sequence

[1176] hybridization probe

[1177] 5'-GAGGAGGAGTGCAGGATTGAGCCATGTACCCAGAAAGGGCAATGCCACC-3' (SEQ ID NO:130)



[1178] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO224 gene using the probe oligonucleotide and one of the PCR primers.

[1179] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1180] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO224 [herein designated as DNA33221-1133] and the derived protein sequence for PRO224.

[1181] The entire nucleotide sequence of DNA33221-1133 is shown in **FIG. 45** (SEQ ID NO:126). Clone DNA33221-1133 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 33-35 and ending at the stop codon at nucleotide positions 879-899 (**FIG. 45**; SEQ ID NO:126). The start of a transmembrane region begins at nucleotide position 777. The predicted polypeptide precursor is 282 amino acids long (**FIG. 46**). Clone DNA33221-1133 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209263.

[1182] Analysis of the amino acid sequence of the full-length PRO224 suggests that it has homology to very low-density lipoprotein receptors, apolipoprotein E receptor and chicken oocyte receptors P95. Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO224 has amino acid identity to portions of these proteins in the range from 28% to 45%, and overall identity with these proteins in the range from 33% to 39%.

#### Example 21

[1183] Isolation of cDNA Clones Encoding Human PRO222

[1184] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence is designated herein as DNA28771. Based on the DNA28771 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO222.

[1185] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-ATCTCCTATCGCTGCTTTCCCGG-3' (SEQ ID NO:133)

reverse PCR primer 5'-AGCCAGGATCGCAGTAAACTCC-3' (SEQ ID NO:134)

[1186] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28771 sequence which had the following nucleotide sequence:

[1187] hybridization probe

[1188] 5'-ATTAAACTTGATGGGTCTGCGTATCT-TGAGTGCTTACAAAACCTTATCT-3' (SEQ ID NO:135)

[1189] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO222 gene using the probe oligonucleotide and one of the PCR primers.

[1190] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1191] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO222 [herein designated as DNA33107-1135] and the derived protein sequence for PRO222.

[1192] The entire nucleotide sequence of DNA33107-1135 is shown in **FIG. 47** (SEQ ID NO:131). Clone DNA33107-1135 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 159-161 and ending at the stop codon at nucleotide positions 1629-1631 (**FIG. 47**; SEQ ID NO:131). The predicted polypeptide precursor is 490 amino acids long (**FIG. 48**). Clone DNA33107-1135 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209251.

[1193] Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO222 shows amino acid sequence identity to mouse complement factor h precursor (25-26%), complement receptor (27-29%), mouse complement C3b receptor type 2 long form precursor (25-47%) and human hypothetical protein kiaa0247 (40%).

#### Example 22

[1194] Isolation of cDNA Clones Encoding PRO234

[1195] A consensus DNA sequence was assembled (DNA30926) using phrap as described in Example 1 above. Based on this consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence.

[1196] RNA for the construction of the cDNA libraries was isolated using standard isolation protocols, e.g., Ausubel et al., *Current Protocols in Molecular Biology*, from tissue or cell line sources or it was purchased from commercial sources (e.g., Clontech). The cDNA libraries used to isolate the cDNA clones were constructed by stan-

dard methods (e.g., Ausubel et al.) using commercially available reagents (e.g., Invitrogen). This library was derived from 22 week old fetal brain tissue.

[1197] A cDNA clone was sequenced in its entirety. The entire nucleotide sequence of PRO234 is shown in **FIG. 49** (SEQ ID NO:136). The predicted polypeptide precursor is 382 amino acids long and has a calculated molecular weight of approximately 43.1 kDa.

[1198] The oligonucleotide sequences used in the above procedure were the following:

30926.p (OLI826) (SEQ ID NO:138):  
5'-GTTTCATTGAAAACCTCTTGCCATCTGATGGTGACTTCTGGATTGGGCTCA-3'

30926.f (OLI827) (SEQ ID NO:139):  
5'-AAGCCAAGAAGCCTGCAGGAGGG-3'

30926.r (OLI828) (SEQ ID NO:140):  
5'-CAGTCCAAGCATAAAGGTCCCTGGC-3'

#### Example 23

[1199] Isolation of cDNA Clones Encoding Human PRO231

[1200] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence was designated herein as DNA30933. Based on the DNA30933 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO231.

[1201] Three PCR primers (two forward and one reverse) were synthesized:

forward PCR primer 1 5'-CCAACTACCAAAGCTGCTGGAGCC-3' (SEQ ID NO:143)

forward PCR primer 2 5'-GCAGCTCTATTACCACGGGAAGGA-3' (SEQ ID NO:144)

reverse PCR primer 5'-TCCTTCCCGTGGTAATAGAGCTGC-3' (SEQ ID NO:145)

[1202] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30933 sequence which had the following nucleotide sequence

[1203] hybridization probe

[1204] 5'-GGCAGAGAACCAGAGGCCGGAG-GAGACTGCCTCTTTACAGCCAGG-3' (SEQ ID NO:146)

[1205] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO231 gene using the probe oligonucleotide and one of the PCR primers.

forward PCR primer 5'-TTCAGCTCATCACCTTCACCTGCC-3' (SEQ ID NO:149)

reverse PCR primer 5'-GGCTCATACAAAATACCACTAGGG-3' (SEQ ID NO:150)

[1206] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1207] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for

PRO231 [herein designated as DNA34434-1139] and the derived protein sequence for PRO231.

[1208] The entire nucleotide sequence of DNA34434-1139 is shown in FIG. 51 (SEQ ID NO:141). Clone DNA34434-1139 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 173-175 and ending at the stop codon at nucleotide positions 1457-1459 (FIG. 51; SEQ ID NO:141). The predicted polypeptide precursor is 428 amino acids long (FIG. 52). Clone DNA34434-1139 has been deposited with ATCC on Sep. 16, 1997 and is assigned ATCC deposit no. ATCC 209252.

[1209] Analysis of the amino acid sequence of the full-length PRO231 suggests that it possesses 30% and 31% amino acid identity with the human and rat prostatic acid phosphatase precursor proteins, respectively.

#### Example 24

[1210] Isolation of cDNA Clones Encoding Human PRO229

[1211] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA28762. Based on the DNA28762 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO229.

[1212] A pair of PCR primers (forward and reverse) were synthesized:

[1213] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28762 sequence which had the following nucleotide sequence

[1214] hybridization probe

[1215] 5'-GGGCCTCCACCGCTGT-  
GAAGGGCGGGTGGAGGTGGAACAGAAAG-  
GCCAGT-3' (SEQ ID NO:151)

[1216] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO229 gene using the probe oligonucleotide and one of the PCR primers.

[1217] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1218] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO229 [herein designated as DNA33100-1159] (SEQ ID NO:147) and the derived protein sequence for PRO229.

[1219] The entire nucleotide sequence of DNA33100-1159 is shown in FIG. 53 (SEQ ID NO:147). Clone DNA33100-1159 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 98-100 and ending at the stop codon at nucleotide positions 1139-1141 (FIG. 53). The predicted polypeptide precursor is 347 amino acids long (FIG. 54). Clone DNA33100-1159 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209377.

[1220] Analysis of the amino acid sequence of the full-length PRO229 polypeptide suggests that portions of it possess significant homology to antigen wc1.1, M130 antigen and CD6.

#### Example 25

[1221] Isolation of cDNA Clones Encoding Human PRO238

[1222] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described above in Example 1. This consensus sequence is herein designated DNA30908. Based on the DNA30908 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO238.

[1223] PCR primers (forward and reverse) were synthesized:

forward PCR primer 1 5'-GGTGCTAAACTGGTCTCTGTGGC-3' (SEQ ID NO:154)

forward PCR primer 2 5'-GAGGGCAAGATGAGCATTC-3' (SEQ ID NO:155)

reverse PCR primer 5'-TCATACTGTTCCATCTCGGCACGC-3' (SEQ ID NO:156)

[1224] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30908 sequence which had the following nucleotide sequence

[1225] hybridization probe

[1226] 5'-AATGGTGGGGCCCTAGAAGAGCTCAT-  
CAGAGAACTCACCGCTTCTCATGC-3' (SEQ ID  
NO:157)

[1227] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO238 gene using the probe oligonucleotide and one of the PCR primers.

[1228] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1229] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO238 and the derived protein sequence for PRO238.

[1230] The entire nucleotide sequence of DNA35600-1162 is shown in FIG. 55 (SEQ ID NO:152). Clone DNA35600-1162 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 134-136 and ending prior to the stop codon at nucleotide positions 1064-1066 (FIG. 55). The predicted polypeptide precursor is 310 amino acids long (FIG. 56). Clone DNA35600-1162 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209370.

[1231] Analysis of the amino acid sequence of the full-length PRO238 polypeptide suggests that portions of it possess significant homology to reductase, particularly oxidoreductase, thereby indicating that PRO238 may be a novel reductase.

#### Example 26

[1232] Isolation of cDNA Clones Encoding Human PRO233

[1233] The extracellular domain (ECD) sequences (including the secretion signal, if any) of from about 950 known secreted proteins from the Swiss-Prot public protein database were used to search expressed sequence tag (EST) databases. The EST databases included public EST databases (e.g., GenBank) and a proprietary EST DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST2 (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequence. Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University

of Washington, Seattle, Wash.; <http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>).

[1234] An expressed sequence tag (EST) was identified by the EST database search and a consensus DNA sequence was assembled relative to other EST sequences using phrap. This consensus sequence is herein designated DNA30945. Based on the DNA30945 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA

library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO233.

[1235] Forward and reverse PCR primers were synthesized:

forward PCR primer 5'-GGTGAAGGCAGAAATTGGAGATG-3' (SEQ ID NO:160)

reverse PCR primer 5'-ATCCCATGCATCAGCCTGTTTACC-3' (SEQ ID NO:161)

[1236] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30945 sequence which had the following nucleotide sequence

[1237] hybridization probe

[1238] 5'-GCTGGTGTAGTCTATACATCA-GATTGTTTIGCTACACAAGATCCTCAG-3' (SEQ ID NO:162)

forward PCR primer 5'-TTCCATGCCACCTAAGGGAGACTC-3' (SEQ ID NO:165)

reverse PCR primer 1 5'-TGGATGAGGTGTGCAATGGCTGGC-3' (SEQ ID NO:166)

reverse PCR primer 2 5'-AGCTCTCAGAGGCTGGTCATAGGG-3' (SEQ ID NO:167)

[1239] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO233 gene using the probe oligonucleotide.

[1240] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue.

[1241] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO233 [herein designated as DNA34436-1238] (SEQ ID NO:158) and the derived protein sequence for PRO233.

[1242] The entire nucleotide sequence of DNA34436-1238 is shown in FIG. 57 (SEQ ID NO:158). Clone DNA34436-1238 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 101-103 and ending at the stop codon at nucleotide positions 1001-1003 (FIG. 57). The predicted polypeptide precursor is 300 amino acids long (FIG. 58). The full-length PRO233 protein shown in FIG. 58 has an estimated molecular weight of about 32,964 daltons and a pI of about 9.52. Clone DNA34436-1238 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209523.

[1243] Analysis of the amino acid sequence of the full-length PRO233 polypeptide suggests that portions of it possess significant homology to reductase proteins, thereby indicating that PRO233 may be a novel reductase.

#### Example 27

[1244] Isolation of cDNA Clones Encoding Human PRO223

[1245] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in

Example 1 above. This consensus sequence is herein designated DNA301836. Based on the DNA30836 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO223.

[1246] PCR primer pairs (one forward and two reverse) were synthesized:

[1247] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30836 sequence which had the following nucleotide sequence

[1248] hybridization probe

[1249] 5'-GTCGGCCCTCCCAGGACTGAACAT-GAAGAGTTATGCCGCTTCCTCAC-3' (SEQ ID NO:168)

[1250] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO223 gene using the probe oligonucleotide and one of the PCR primers.

[1251] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1252] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO223 [herein designated as DNA33206-1165] (SEQ ID NO:163) and the derived protein sequence for PRO223.

[1253] The entire nucleotide sequence of DNA33206-1165 is shown in FIG. 59 (SEQ ID NO:163). Clone DNA33206-1165 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 97-99 and ending at the stop codon at nucleotide positions 1525-1527 (FIG. 59). The predicted polypeptide precursor is 476 amino acids long (FIG. 60). Clone DNA33206-1165 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209372.

[1254] Analysis of the amino acid sequence of the full-length PRO223 polypeptide suggests that it possesses significant homology to various serine carboxypeptidase proteins, thereby indicating that PRO223 may be a novel serine carboxypeptidase.

#### Example 28

[1255] Isolation of cDNA Clones Encoding Human PRO235

[1256] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated "DNA30927". Based on the DNA30927 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR, a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO235.

[1257] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-TGGAAATACCGCTCCTGCAG-3' (SEQ ID NO:171)

reverse PCR primer 5'-CTTCTGCCCTTTGGAGAAGATGGC-3' (SEQ ID NO:172)

[1258] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30927 sequence which had the following nucleotide sequence

[1259] hybridization probe

[1260] 5'-GGACTCACTGGCCAGGCCTTCAATATCACCAGCCAGGACGAT-3' (SEQ ID NO:173)

[1261] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by

forward PCR primer 5'-TGGCTACTCCAAGACCCTGGCATG-3' (SEQ ID NO:178)

reverse PCR primer 5'-TGGACAAATCCCCTTGCTCAGCCC-3' (SEQ ID NO:179)

PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO235 gene using the probe oligonucleotide and one of the PCR primers.

[1262] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1263] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO235 [herein designated as DNA35558-1167] (SEQ ID NO:169) and the derived protein sequence for PRO235.

forward PCR primer 5'-CCAGCTATGACTATGATGCACC-3' (SEQ ID NO:181)

reverse PCR primer 5'-TGGCACCCAGAATGGTGTGGCTC-3' (SEQ ID NO:182)

[1264] The entire nucleotide sequence of DNA35558-1167 is shown in FIG. 61 (SEQ ID NO:169). Clone DNA35558-1167 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 667-669 and ending at the stop codon at nucleotide positions 2323-2325 (FIG. 61). The predicted polypeptide precursor is 552 amino acids long (FIG. 62). Clone DNA35558-1167 has been deposited with ATCC and is assigned ATCC deposit no. 209374.

[1265] Analysis of the amino acid sequence of the full-length PRO235 polypeptide suggests that portions of it possess significant homology to the human, mouse and *Xenopus* plexin protein, thereby indicating that PRO235 may be a novel plexin protein.

#### Example 29

[1266] Isolation of cDNA Clones Encoding Human PRO236 and Human PRO262

[1267] Consensus DNA sequences were assembled relative to other EST sequences using phrap as described in

Example 1 above. These consensus sequences are herein designated DNA30901 and DNA30847. Based on the DNA30901 and DNA30847 consensus sequences, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO236 and PRO262, respectively.

[1268] Based upon the DNA30901 consensus sequence, a pair of PCR primers (forward and reverse) were synthesized:

[1269] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30901 sequence which had the following nucleotide sequence

[1270] hybridization probe

[1271] 5'-GGGCTTACC GAAGCAGTGGACCTT-TATTTTGACCACCTGATGTCCAGGG-3' (SEQ ID NO:180)

[1272] Based upon the DNA30847 consensus sequence, a pair of PCR primers (forward and reverse) were synthesized:

[1273] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30847 sequence which had the following nucleotide sequence

[1274] hybridization probe

[1275] 5'-CGAGATGTCATCAGCAAGTTCCAG-GAAGTTCCTTTGGGACCTTTAICCTCC-3' (SEQ ID NO:183)

[1276] In order to screen several libraries for a source of full-length clones, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified

forward PCR primer 5'-CCTCCCTCTATTACCCATGTC-3' (SEQ ID NO:186)

reverse PCR primer 5'-GACCAACTTTCTCTGGGAGTGAGG-3' (SEQ ID NO:187)

above. Positive libraries were then used to isolate clones encoding the PRO236 and PRO262 genes using the probe oligonucleotides and one of the PCR primers.

[1277] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue for PRO236 and human fetal liver tissue for PRO262.

[1278] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO236 [herein designated as DNA35599-1168] (SEQ ID NO:174), the derived protein sequence for PRO236, the full-length DNA sequence for PRO262 [herein designated as DNA36992-1168] (SEQ ID NO:176) and the derived protein sequence for PRO262.

[1279] The entire nucleotide sequence of DNA35599-1168 is shown in FIG. 63 (SEQ ID NO:174). Clone DNA35599-1168 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 69-71 and ending at the stop codon at nucleotide positions 1977-1979 (FIG. 63). The predicted polypeptide precursor is 636 amino acids long (FIG. 64). Clone DNA35599-1168 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209373.

[1280] The entire nucleotide sequence of DNA36992-1168 is shown in FIG. 65 (SEQ ID NO:176). Clone DNA36992-1168 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 240-242 and ending at the stop codon at nucleotide positions 2202-2204 (FIG. 65). The predicted polypeptide precursor is 654 amino acids long (FIG. 66). Clone DNA36992-1168 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209382.

[1281] Analysis of the amino acid sequence of the full-length PRO236 and PRO262 polypeptides suggests that portions of those polypeptides possess significant homology to  $\beta$ -galactosidase proteins derived from various sources, thereby indicating that PRO236 and PRO262 may be novel  $\beta$ -galactosidase homologs.

#### Example 30

[1282] Isolation of cDNA Clones Encoding Human PRO239

[1283] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30909. Based on the DNA30909 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO239.

[1284] A pair of PCR primers (forward and reverse) were synthesized:

[1285] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30909 sequence which had the following nucleotide sequence

[1286] hybridization probe

[1287] 5'-GTCACCTTTATTTCTCTAACAACAAGCTCGAATCCTTACCAGTGGCAG-3' (SEQ ID NO:188)

[1288] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO239 gene using the probe oligonucleotide and one of the PCR primers.

[1289] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.

[1290] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO239 [herein designated as DNA34407-1169] (SEQ ID NO:184) and the derived protein sequence for PRO239.

[1291] The entire nucleotide sequence of DNA34407-1169 is shown in FIG. 67 (SEQ ID NO:184). Clone DNA34407-1169 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 72-74 and ending at the stop codon at nucleotide positions 1575-1577 (FIG. 67). The predicted polypeptide precursor is 501 amino acids long (FIG. 68). Clone DNA34407-1169 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209383.

[1292] Analysis of the amino acid sequence of the full-length PRO239 polypeptide suggests that portions of it possess significant homology to the densin protein, thereby indicating that PRO239 may be a novel molecule in the densin family.

#### Example 31

[1293] Isolation of cDNA Clones Encoding Human PRO257

[1294] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in

Example 1 above. This consensus sequence is herein designated DNA28731. Based on the DNA28731 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO257.

[1295] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-TCTCTATTCCAAACTGTGGCG-3' (SEQ ID NO:191)

reverse PCR primer 5'-TTTGATGACGATTGGAAGGTGG-3' (SEQ ID NO:192)

[1296] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA28731 sequence which had the following nucleotide sequence

[1297] hybridization probe

[1298] 5'-GGAAGGATCCTTCACCAGCCCCAAAT-TACCCAAAGCCGCATCCTGAGC-3' (SEQ ID NO:193)

forward PCR primer: 5'-TGTTTACCAGGCCAAGTTCGG-3' (SEQ ID NO:196);

reverse PCR primer A: 5'-GGATTCATCCTCAAGGAAGAGCGG-3' (SEQ ID NO:197); and

reverse PCR primer B: 5'-AACTTGCAGCATCAGCCACTCTGC-3' (SEQ ID NO:198)

[1299] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO257 gene using the probe oligonucleotide and one of the PCR primers.

[1300] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1301] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO257 [herein designated as DNA35841-1173 (SEQ ID NO:189) and the derived protein sequence for PRO257.

[1302] The entire nucleotide sequence of DNA35841-1173 is shown in FIG. 69 (SEQ ID NO:189). Clone DNA35841-1173 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 964-966 and ending at the stop codon at nucleotide positions 2785-2787 (FIG. 69). The predicted polypeptide precursor is 607 amino acids long (FIG. 70). Clone DNA35841-1173 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209403.

[1303] Analysis of the amino acid sequence of the full-length PRO257 polypeptide suggests that portions of it possess significant homology to the ebnerin protein, thereby

indicating that PRO257 may be a novel protein member related to the ebnerin protein.

#### Example 32

[1304] Isolation of cDNA Clones Encoding Human PRO260

[1305] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in

Example 1 above. This consensus sequence is herein designated DNA30834. Based on the DNA30834 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO260.

[1306] PCR primers (forward and two reverse) were synthesized:

[1307] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30834 sequence which had the following nucleotide sequence:

[1308] hybridization probe:

[1309] 5'-TTCCGTGCCCAGCTTCGGTAGC-GAGTGGTCTGTTGGTATTGGCA-3' (SEQ ID NO:199)

[1310] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO260 gene using the probe oligonucleotide and one of the PCR primers.

[1311] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1312] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO260 [herein designated as DNA33470-1175] (SEQ ID NO:194) and the derived protein sequence for PRO260.

[1313] The entire nucleotide sequence of DNA33470-1175 is shown in FIG. 71 (SEQ ID NO:194). Clone DNA33470-1175 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 67-69 and ending at the stop codon 1468-1470 (see

**FIG. 71).** The predicted polypeptide precursor is 467 amino acids long (**FIG. 72**). Clone DNA33470-1175 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209398.

[1314] Analysis of the amino acid sequence of the full-length PRO260 polypeptide suggests that portions of it possess significant homology to the alpha-1-fucosidase precursor, thereby indicating that PRO260 may be a novel fucosidase.

#### Example 33

[1315] Isolation of cDNA Clones Encoding Human PRO263

[1316] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA30914. Based on the DNA30914 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO263.

[1317] PCR primers (two forward and one reverse) were synthesized:

forward PCR primer 1: 5'-GAGCTTTCATCCAGGTGTCATGC-3' (SEQ ID NO:202);

forward PCR primer 2: 5'-GTCACTGACAGTACTACTCGG-3' (SEQ ID NO:203);

reverse PCR primer: 5'-TGGAGCAGGAGTAGTAGTAGG-3' (SEQ ID NO:204)

[1318] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30914 sequence which had the following nucleotide sequence:

forward PCR primer (.f1) 5'-GCTTGGATATTCGCATGGGCTAC-3' (SEQ ID NO:208)

forward PCR primer (.f2) 5'-TGGAGACAATATCCCTGAGG-3' (SEQ ID NO:209)

reverse PCR primer (.r1) 5'-AACAGTTGGCCACAGCATGGCAGG-3' (SEQ ID NO:210)

[1319] hybridization probe:

[1320] 5'-AGGAGGCCTGTAGGCTGCTGGGAC-TAAGTTTGGCCGGCAAGGACCAAGTT-3' (SEQ ID NO:205)

[1321] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO263 gene using the probe oligonucleotide and one of the PCR primers.

[1322] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1323] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO263 [herein designated as DNA34431-1177] (SEQ ID NO:200) and the derived protein sequence for PRO263.

[1324] The entire nucleotide sequence of DNA34431-1177 is shown in **FIG. 73** (SEQ ID NO:200). Clone DNA34431-1177 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 160-162 of SEQ ID NO:200 and ending at the stop codon after the nucleotide at position 1126-1128 of SEQ ID NO:200 (**FIG. 73**). The predicted polypeptide precursor is 322 amino acids long (**FIG. 74**). Clone DNA34431-1177 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209399.

[1325] Analysis of the amino acid sequence of the full-length PRO263 polypeptide suggests that portions of it possess significant homology to CD44 antigen, thereby indicating that PRO263 may be a novel cell surface adhesion molecule.

#### Example 34

[1326] Isolation of cDNA Clones Encoding Human PRO270

[1327] A consensus DNA sequence was assembled relative to the other identified EST sequences as described in Example 1 above, wherein the consensus sequence was designated herein as DNA35712. Based on the DNA35712 consensus sequence, oligonucleotides were synthesized: 1)

to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO270. Forward and reverse PCR primers were synthesized:

[1328] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35712 sequence which had the following nucleotide sequence

[1329] hybridization probe

[1330] 5'-CCATTGATGAGGAACACTAGAACGGGA-CAAGAGGGTCACTTGGATTCTGGAG-3' (SEQ ID NO:211)

[1331] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO270 gene using the probe oligonucleotide and one of the PCR primers.

[1332] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.



[1333] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO270 [herein designated as DNA39510-1181] (SEQ ID NO:206) and the derived protein sequence for PRO270.

[1334] The entire nucleotide sequence of DNA39510-1181 is shown in FIG. 75 (SEQ ID NO:206). Clone DNA39510-1181 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 3-5 and ending at the stop codon at nucleotide positions 891-893 (FIG. 75; SEQ ID NO:206). The predicted polypeptide precursor is 296 amino acids long (FIG. 76). Clone DNA39510-1181 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209392.

[1335] Analysis of the amino acid sequence of the full-length PRO270 suggests that portions of it possess significant homology to the thioredoxin-protein, thereby indicating that the PRO270 protein may be a novel member of the thioredoxin family.

#### Example 35

[1336] Isolation of cDNA Clones Encoding Human PRO271

[1337] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35737. Based on the DNA35737 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO271.

[1338] Forward and reverse PCR primers were synthesized:

forward PCR primer 1 5'-TGCTTCGCTACTGCCCTC-3' (SEQ ID NO:214)

forward PCR primer 2 5'-TTCCCTTGTGGTTGGAG-3' (SEQ ID NO:215)

forward PCR primer 3 5'-AGGGCTGGAAGCCAGTTC-3' (SEQ ID NO:216)

reverse PCR primer 1 5'-AGCCAGTGAGGAAATGCG-3' (SEQ ID NO:217)

reverse PCR primer 2 5'-TGTCCAAAGTACACACCTGAGG-3' (SEQ ID NO:218)

[1339] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35737 sequence which had the following nucleotide sequence

[1340] hybridization probe

[1341] 5'-GATGCCACGATCGCCAAGGTGGGA-CAGCTCTTTGCCCGCTGGAAG-3' (SEQ ID NO:219)

[1342] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO271 gene using the probe oligonucleotide and one of the PCR primers.

[1343] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue.

[1344] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO271 [herein designated as DNA39423-1182] (SEQ ID NO:212) and the derived protein sequence for PRO271.

[1345] The entire nucleotide sequence of DNA39423-1182 is shown in FIG. 77 (SEQ ID NO:212). Clone DNA39423-1182 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 101-103 and ending at the stop codon at nucleotide positions 1181-1183 (FIG. 77). The predicted polypeptide precursor is 360 amino acids long (FIG. 78). Clone DNA39423-1182 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209387.

[1346] Analysis of the amino acid sequence of the full-length PRO271 polypeptide suggests that it possess significant homology to the proteoglycan link protein, thereby indicating that PRO271 may be a link protein homolog.

#### Example 36

[1347] Isolation of cDNA Clones Encoding Human PRO272

[1348] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in

Example 1 above. This consensus sequence is herein designated DNA36460. Based on the DNA36460 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO272.

[1349] Forward and reverse PCR primers were synthesized:

forward PCR primer (.f1) 5'-CGCAGGCCCTCATGGCCAGG-3' (SEQ ID NO:222)

forward PCR primer (.f2) 5'-GAAATCCTGGGTAATTGG-3' (SEQ ID NO:223)

reverse PCR primer 5'-GTGCGCGGTGCTCACAGCTCATC-3' (SEQ ID NO:224)

[1350] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA36460 sequence which had the following nucleotide sequence

[1351] hybridization probe

[1352] 5'-CCCCCTGAGCGACGCTCCCCCATGATGACGCCACGGGAACCTTC-3' (SEQ ID NO:225)

[1353] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO272 gene using the probe oligonucleotide and one of the PCR primers.

[1354] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.

[1355] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO272 [herein designated as DNA40620-1183] (SEQ ID NO:220) and the derived protein sequence for PRO272.

[1356] The entire nucleotide sequence of DNA40620-1183 is shown in **FIG. 79** (SEQ ID NO:220). Clone DNA40620-1183 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 35-37 and ending at the stop codon at nucleotide positions 1019-1021 (**FIG. 79**). The predicted polypeptide precursor is 328 amino acids long (**FIG. 80**). Clone DNA40620-1183 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209388.

[1357] Analysis of the amino acid sequence of the full-length PRO272 polypeptide suggests that portions of it possess significant homology to the human and mouse reticulocalbin proteins, respectively, thereby indicating that PRO272 may be a novel reticulocalbin protein.

#### Example 37

[1358] Isolation of cDNA Clones Encoding Human PRO294

[1359] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35731. Based on the DNA35731 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO294.

[1360] Forward and reverse PCR primers were synthesized:

forward PCR primer (.f1) 5'-TGGTCTCGCACCCGATC-3' (SEQ ID NO:228)

forward PCR primer (.f2) 5'-CTGCTGTCCACAGGGGAG-3' (SEQ ID NO:229)

forward PCR primer (.f3) 5'-CCTTGAAGCATACTGCTC-3' (SEQ ID NO:230)

forward PCR primer (.f4) 5'-GAGATAGCAATTTCCGCC-3' (SEQ ID NO:231)

reverse PCR primer (.r1) 5'-TTCCTCAAGAGGGCAGCC-3' (SEQ ID NO:232)

reverse PCR primer (.r2) 5'-CTTGGCACAATGTCCGAGATTTC-3' (SEQ ID NO:233)

[1361] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35731 sequence which had the following nucleotide sequence

[1362] hybridization probe

[1363] 5'-GCTCTGAGGAAGGTGACGCGCGGGGCTCCGAACCCTTGGCCTTG-3' (SEQ ID NO:234)

[1364] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO294 gene using the probe oligonucleotide and one of the PCR primers.

[1365] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue.

[1366] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO294 [herein designated as DNA40604-1187] (SEQ ID NO:226) and the derived protein sequence for PRO294.

[1367] The entire nucleotide sequence of DNA40604-1187 is shown in **FIG. 81** (SEQ ID NO:226). Clone DNA40604-1187 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 396-398 and ending at the stop codon at nucleotide positions 2046-2048 (**FIG. 81**). The predicted polypeptide precursor is 550 amino acids long (**FIG. 82**). Clone DNA40604-1187 has been deposited with ATCC and is assigned ATCC deposit no. 209394.

[1368] Analysis of the amino acid sequence of the full-length PRO294 polypeptide suggests that portions of it possess significant homology to portions of various collagen proteins, thereby indicating that PRO294 may be collagen-like molecule.

#### Example 38

[1369] Isolation of cDNA Clones Encoding Human PRO295

[1370] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35814. Based on the DNA35814 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO295.

[1371] Forward and reverse PCR primers were synthesized:

forward PCR primer (.f1) 5'-GCAGAGCGGAGATGCAGCGGCTTG-3' (SEQ ID NO:238)

forward PCR primer (.f2) 5'-CCCAGCATGTACTGCCAG-3' (SEQ ID NO:239)

forward PCR primer (.f3) 5'-TTGGCAGCTTCATGGAGG-3' (SEQ ID NO:240)

forward PCR primer (.f4) 5'-CCTGGGCAAAAATGCAAC-3' (SEQ ID NO:241)

reverse PCR primer (.r1) 5'-CTCCAGCTCCTGGCGCACCTCCTC-3' (SEQ ID NO:242)

[1372] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35814 sequence which had the following nucleotide sequence

[1373] hybridization probe

[1374] 5'-GGCTCTCAGCTACCGCGCAGGAGC-GAGGCCACCCTCAATGAGATG-3' (SEQ ID NO:243)

[1375] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO295 gene using the probe oligonucleotide and one of the PCR primers.

[1376] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.

[1377] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for

database were used to search expressed sequence tag (EST) databases. The EST databases included public EST data-

bases (e.g., (GenBank) and a proprietary EST DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST2 (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequence. Those comparisons resulting in a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.; <http://bozeman.mbt.washington.edu/phrap.docs/phrap.html>).

[1382] Based on an expression tag sequence designated herein as T08294 identified in the above analysis, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO293.

[1383] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-AACAAAGGTAAGATGCCATCCTG-3' (SEQ ID NO:246)

reverse PCR primer 5'-AAACTTGTTCGATGGAGACCAGCTC-3' (SEQ ID NO:247)

PRO295 [herein designated as DNA38268-1188] (SEQ ID NO:235) and the derived protein sequence for PRO295.

[1378] The entire nucleotide sequence of DNA38268-1188 is shown in FIG. 83 (SEQ ID NO:235). Clone DNA38268-1188 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 153-155 and ending at the stop codon at nucleotide positions 1202-1204 (FIG. 83). The predicted polypeptide precursor is 350 amino acids long (FIG. 84). Clone DNA38268-1188 has been deposited with ATCC and is assigned ATCC deposit no. 209421.

[1379] Analysis of the amino acid sequence of the full-length PRO295 polypeptide suggests that portions of it possess significant homology to the integrin proteins, thereby indicating that PRO295 may be a novel integrin.

#### Example 39

[1380] Isolation of cDNA Clones Encoding Human PRO293

[1381] The extracellular domain (ECD) sequences (including the secretion signal, if any) of from about 950 known secreted proteins from the Swiss-Prot public protein

[1384] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the expression sequence tag which had the following nucleotide sequence

[1385] hybridization probe

[1386] 5'-AGGGGCTGCAAAGCCTGGAGAGC-CTTCCTTCTATGACAACCAGC-3' (SEQ ID NO:248)

[1387] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO293 gene using the probe oligonucleotide and one of the PCR primers.

[1388] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue.

[1389] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO293 [herein designated as DNA37151-1193] (SEQ ID NO:244) and the derived protein sequence for PRO293.

[1390] The entire nucleotide sequence of DNA37151-1193 is shown in FIG. 85 (SEQ ID NO:244). Clone

DNA37151-1193 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 881-883 and ending at the stop codon after nucleotide position 3019 of SEQ ID NO:244, **FIG. 85**). The predicted polypeptide precursor is 713 amino acids long (**FIG. 86**). Clone DNA37151-1193 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209393.

[1391] Analysis of the amino acid sequence of the full-length PRO293 polypeptide suggests that portions of it possess significant homology to the NLRR proteins, thereby indicating that PRO293 may be a novel NLRR protein.

#### Example 40

[1392] Isolation of cDNA Clones Encoding Human PRO247

[1393] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA33480. Based on the DNA33480 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO247.

[1394] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-CAACAATGAGGGCACCAAGC-3' (SEQ ID NO:251)

reverse PCR primer 5'-GATGGCTAGGTTCTGGAGTTCTG-3' (SEQ ID NO:252)

[1395] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the DNA33480 expression sequence tag which had the following nucleotide sequence

[1396] hybridization probe

[1397] 5'-CAACCTGCAGGAGATTGACCTCAAG-GACAACAACCTCAAGACCATCG-3' (SEQ ID NO:253)

forward PCR primer 1 5'-GTCCGCAAGGATGCCTACATGTTC-3' (SEQ ID NO:264)

forward PCR primer 2 5'-GCAGAGGTGTCTAAGGTTG-3' (SEQ ID NO:265)

reverse PCR primer 5'-AGCTCTAGACCAATGCCAGCTTCC-3' (SEQ ID NO:266)

[1398] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO247 gene using the probe oligonucleotide and one of the PCR primers.

[1399] RNA for construction of the cDNA libraries was isolated from human fetal brain tissue.

[1400] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO247 [herein designated as DNA35673-1201] (SEQ ID NO:249) and the derived protein sequence for PRO247.

[1401] The entire nucleotide sequence of DNA35673-1201 is shown in **FIG. 89** (SEQ ID NO:249). Clone DNA35673-1201 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 80-82 of SEQ ID NO:249 and ending at the stop codon after nucleotide position 1717 of SEQ ID NO:249 (**FIG. 89**). The predicted polypeptide precursor is 546 amino acids long (**FIG. 88**). Clone DNA35673-1201 has been deposited with ATCC and is assigned ATCC deposit no. 209418.

[1402] Analysis of the amino acid sequence of the full-length PRO247 polypeptide suggests that portions of it possess significant homology to the densin molecule and KIAA0231, thereby indicating that PRO247 may be a novel leucine rich repeat protein.

#### Example 41

[1403] Isolation of cDNA Clones Encoding Human PRO302, PRO303, PRO304, PRO307 and PRO343

[1404] Consensus DNA sequences were assembled relative to other EST sequences using phrap as described in

Example 1 above. These consensus sequences are herein designated DNA35953, DNA35955, DNA35958, DNA37160 and DNA30895. Based on the DNA35953 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO302.

[1405] PCR primers (forward and reverse) were synthesized:

[1406] Also, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35953 sequence which had the following nucleotide sequence

[1407] hybridization probe

[1408] 5'-GCCACCAACTCCTGCAAGAACTTCT-CAGAAGTGGCCCTGGTCATG-3' (SEQ ID NO:267)

[1409] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO302 gene using the probe oligonucleotide and one of the PCR primers.

[1410] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue (LIB228).

[1411] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO302 [herein designated as DNA40370-1217] (SEQ ID NO:254) and the derived protein sequence for PRO302.

[1412] The entire nucleotide sequence of DNA40370-1217 is shown in **FIG. 89** (SEQ ID NO:254). Clone DNA40370-1217 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 34-36 and ending at the stop codon at nucleotide positions 1390-1392 (**FIG. 89**). The predicted polypeptide precursor is 452 amino acids long (**FIG. 90**). Various unique aspects of the PRO302 protein are shown in **FIG. 90**. Clone DNA40370-1217 has been deposited with the ATCC on Nov. 21, 1997 and is assigned ATCC deposit no. ATCC 209485.

[1413] Based on the DNA35955 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO303.

[1414] A pair of PCR primers (forward and reverse) were synthesized:

forward PCR primer 5'-GGGAATTCACCCATGAGATTGCC-3' (SEQ ID NO:268)

reverse PCR primer 5'-GAATGCCCTGCAAGCATCAACTGG-3' (SEQ ID NO:269)

[1415] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35955 sequence which had the following nucleotide sequence:

[1416] hybridization probe

[1417] 5'-GCACCTGTCACTTACACTAAACA-CATCCAGCCCATCTGTCTCCAGGCCCTC-3' (SEQ ID NO:270)

[1418] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO303 gene using the probe oligonucleotide and one of the PCR primers.

[1419] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue (LIB25).

[1420] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO303 [herein designated as DNA42551-1217] (SEQ ID NO:256) and the derived protein sequence for PRO303.

[1421] The entire nucleotide sequence of DNA42551-1217 is shown in **FIG. 91** (SEQ ID NO:256). Clone DNA42551-1217 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 20-22 and ending at the stop codon at nucleotide positions 962-964 (**FIG. 91**). The predicted polypeptide precursor is 314 amino acids long (**FIG. 92**). Various unique aspects of the PRO303 protein are shown in **FIG. 92**. Clone DNA42551-1217 has been deposited on Nov. 21, 1997 with the ATCC and is assigned ATCC deposit no. ATCC 209483.

[1422] Based on the DNA35958 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2)

for use as probes to isolate a clone of the full-length coding sequence for PRO304.

[1423] Pairs of PCR primers (forward and reverse) were synthesized:

forward PCR primer 1 5'-GCGGAAGGGCAGAATGGGACTCCAAG-3' (SEQ ID NO:271)

forward PCR primer 2 5'-CAGCCCTGCCACATGTGC-3' (SEQ ID NO:272)

forward PCR primer 3 5'-TACTGGGTGGTCAGCAAC-3' (SEQ ID NO:273)

reverse PCR primer 5'-GGCGAAGAGCAGGGTGAGACCCCG-3' (SEQ ID NO:274)

[1424] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35958 sequence which had the following nucleotide sequence

[1425] hybridization probe

[1426] 5'-GCCCTCATCCTCTCTGGCAAATGCAGT-TACAGCCCGGAGCCCCGAC-3' (SEQ ID NO:275)

[1427] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO304 gene using the probe oligonucleotide and one of the PCR primers.

[1428] RNA for construction of the cDNA libraries was isolated from 22 week human fetal brain tissue (LIB153).

[1429] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO304 [herein designated as DNA39520-1217] (SEQ ID NO:258) and the derived protein sequence for PRO304.

[1430] The entire nucleotide sequence of DNA39520-1217 is shown in FIG. 93 (SEQ ID NO:258). Clone DNA39520-1217 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 34-36 and ending at the stop codon at nucleotide positions 1702-1704 (FIG. 93). The predicted polypeptide precursor is 556 amino acids long (FIG. 94). Various unique aspects of the PRO304 protein are shown in FIG. 94. Clone DNA39520-1217 has been deposited with ATCC on Nov. 21, 1997 and is assigned ATCC deposit no. ATCC 209482.

forward PCR primer 5'-CGTCTCGAGCGCTCCATACAGTTCCTTCCCCA-3' (SEQ ID NO:281)

reverse PCR primer 5'-TGGAGGGGAGCGGGATGCTTGTCTGGGCGACTCCGGGGCCCCCTCATGTGCCAGGTGGA-3' (SEQ ID NO:282)

[1431] Based on the DNA37160 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO307.

[1432] Pairs of PCR primers (forward and reverse) were synthesized:

forward PCR primer 1 5'-GGGCAGGGATTCCAGGGCTCC-3' (SEQ ID NO:276)

forward PCR primer 2 5'-GGCTATGACAGCAGGTTTC-3' (SEQ ID NO:277)

forward PCR primer 3 5'-TGACAATGACCGACCAGG-3' (SEQ ID NO:278)

reverse PCR primer 5'-GCATCGCATTGCTGGTAGAGCAAG-3' (SEQ ID NO:279)

[1433] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA37160 sequence which had the following nucleotide sequence

[1434] hybridization probe

[1435] 5'-TTACAGTGCCCCCTGGAAACCCACT-TGGCCTGCATACCGCTCCC-3' (SEQ ID NO:280)

[1436] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified

above. A positive library was then used to isolate clones encoding the PRO307 gene using the probe oligonucleotide and one of the PCR primers.

[1437] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue (LIB229).

[1438] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO307 [herein designated as DNA41225-1217] (SEQ ID NO:260) and the derived protein sequence for PRO307.

[1439] The entire nucleotide sequence of DNA41225-1217 is shown in FIG. 95 (SEQ ID NO:260). Clone DNA41225-1217 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 92-94 and ending at the stop codon at nucleotide positions 1241-1243 (FIG. 95). The predicted polypeptide precursor is 383 amino acids long (FIG. 96). Various unique aspects of the PRO307 protein are shown in FIG. 96. Clone DNA41225-1217 has been deposited with ATCC on Nov. 21, 1997 and is assigned ATCC deposit no. ATCC 209491.

[1440] Based on the DNA30895 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO343.

[1441] A pair of PCR primers (forward and reverse) were synthesized:

[1442] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA30895 sequence which had the following nucleotide sequence

[1443] hybridization probe

[1444] 5'-CCCTCAGACCCTGCAGAAGCTGAAG-GTTCTATCATCGACTCGGAAGTCTG-CAGCCATCTGTACTGGCGGGGAGCAGGA-

CAGGGACCCATCACTGAGGACATGCTGTGTGC  
CGGCTACT-3' (SEQ ID NO:283)

[1445] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO343 gene using the probe oligonucleotide and one of the PCR primers.

[1446] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue (LIB26).

[1447] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO343 [herein designated as DNA43318-1217] (SEQ ID NO:262) and the derived protein sequence for PRO343.

[1448] The entire nucleotide sequence of DNA43318-1217 is shown in **FIG. 97** (SEQ ID NO:262). Clone DNA43318-1217 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 53-55 and ending at the stop codon at nucleotide positions 1004-1006 (**FIG. 97**). The predicted polypeptide precursor is 317 amino acids long (**FIG. 98**). Various unique aspects of the PRO343 protein are shown in **FIG. 98**. Clone DNA43318-1217 has been deposited with ATCC on Nov. 21, 1997 and is assigned ATCC deposit no. ATCC 209481.

#### Example 42

[1449] Isolation of cDNA Clones Encoding Human PRO328

[1450] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35615. Based on the DNA35615 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO328.

[1451] Forward and reverse PCR primers were synthesized:

forward PCR primer 5'-TCCTGCAGTTTCCTGATGC-3' (SEQ ID NO:286)

reverse PCR primer 5'-CTCATATTGCACACAGTAATTCG-3' (SEQ ID NO:287)

[1452] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35615 sequence which had the following nucleotide sequence

[1453] hybridization probe

[1454] 5'-ATGAGGAGAAACGTTTGATGGTG-GAGCTGCACAACCTCTACCGGG-3' (SEQ ID NO:288)

[1455] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones

encoding the PRO328 gene using the probe oligonucleotide and one of the PCR primers.

[1456] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue.

[1457] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO328 [herein designated as DNA40587-1231] (SEQ ID NO:284) and the derived protein sequence for PRO328.

[1458] The entire nucleotide sequence of DNA40587-1231 is shown in **FIG. 99** (SEQ ID NO:284). Clone DNA40587-1231 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 15-17 and ending at the stop codon at nucleotide positions 1404-1406 (**FIG. 99**). The predicted polypeptide precursor is 463 amino acids long (**FIG. 100**). Clone DNA40587-1231 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209438.

[1459] Analysis of the amino acid sequence of the full-length PRO328 polypeptide suggests that portions of it possess significant homology to the human glioblastoma protein and to the cysteine rich secretory protein thereby indicating that PRO328 may be a novel glioblastoma protein or cysteine rich secretory protein.

#### Example 43

[1460] Isolation of cDNA Clones Encoding Human PRO335, PRO331 or PRO326

[1461] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA36685. Based on the DNA36685 consensus sequence, and Incyte EST sequence no. 2228990, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO335, PRO331 or PRO326.

[1462] Forward and reverse PCR primers were synthesized for the determination of PRO335:

forward PCR primer 5'-GGAACCGAATCTCAGCTA-3' (SEQ ID NO:295)

forward PCR primer 5'-CCTAAACTGAACTGGACCA-3' (SEQ ID NO:296)

forward PCR primer 5'-GGCTGGAGACTGAACCT-3' (SEQ ID NO:297)

## -continued

forward PCR primer 5'-ACAGCTGCACAGCTCAGAACAGTG-3' (SEQ ID NO:298)

reverse PCR primer 5'-CATTCCCAGTATAAAAATTTTC-3' (SEQ ID NO:299)

reverse PCR primer 5'-GGGTCTTGGTGAATGAGG-3' (SEQ ID NO:300)

reverse PCR primer 5'-GTGCCTCTCGGTTACCACCAATGG-3' (SEQ ID NO:301)

[1463] Additionally, a synthetic oligonucleotide hybridization probe was constructed for the determination of PRO335 which had the following nucleotide sequence

[1464] hybridization probe

[1465] 5'-GCGGCCACTGTTGGACCGAACTG-TAACCAAGGGAGAAACAGCCGTCCTAC-3' (SEQ ID NO:302)

[1466] Forward and reverse PCR primers were synthesized for the determination of PRO331:

forward PCR primer 5'-GCCTTTGACAACCTTCAGTCACTAGTGG-3' (SEQ ID NO:303)

reverse PCR primer 5'-CCCCATGTGTCCATGACTGTTCCC-3' (SEQ ID NO:304)

[1467] Additionally, a synthetic oligonucleotide hybridization probe was constructed for the determination of PRO331 which had the following nucleotide sequence

[1468] hybridization probe

[1469] 5'-TACTGCCTCATGACCTCTTCACTCCCT-TGCATCATCTTAGAGCGG-3' (SEQ ID NO:305)

[1470] Forward and reverse PCR primers were synthesized for the determination of PRO326:

forward PCR primer 5'-ACTCCAAGGAAATCGGATCCGTTC-3' (SEQ ID NO:306)

reverse PCR primer 5'-TTAGCAGCTGAGGATGGGCACAAC-3' (SEQ ID NO:307)

[1471] Additionally, a synthetic oligonucleotide hybridization probe was constructed for the determination of PRO331 which had the following nucleotide sequence

[1472] hybridization probe

[1473] 5'-GCCITCACTGGTTTGGATGCATTG-GAGCATCTAGACCTGAGTGACAACGC-3' (SEQ ID NO:308)

[1474] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones

encoding the PRO335, PRO331 or PRO326 gene using the probe oligonucleotide and one of the PCR primers.

[1475] RNA for construction of the cDNA libraries was isolated from human fetal kidney tissue (PRO335 and PRO326) and human fetal brain (PRO331).

[1476] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO335, PRO331 or PRO326 [herein designated as SEQ ID NOS:289, 291 and 293, respectively; see FIGS. 101, 103 and 105, respectively], and the derived protein sequence for

PRO335, PRO331 or PRO326 (see FIGS. 102, 104 and 106, respectively; SEQ ID NOS:290, 292 and 294, respectively).

[1477] The entire nucleotide sequences are shown in FIGS. 101, 103 and 105, deposited with the ATCC on Jun. 2, 1998, Nov. 7, 1997 and Nov. 21, 1997, respectively.

[1478] Analysis of the amino acid sequence of the full-length PRO335, PRO331 or PRO326 polypeptide suggests that portions of it possess significant homology to the LIG-1

protein, thereby indicating that PRO335, PRO331 and PRO326 may be a novel LIG-1-related protein.

## Example 44

[1479] Isolation of cDNA Clones Encoding Human PRO332

[1480] Based upon an ECD homology search performed as described in Example 1 above, a consensus DNA sequence designated herein as DNA36688 was assembled. Based on the DNA36688 consensus sequence, oligonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO332.

A paid of PCR primers (forward and reverse) were synthesized:

5'-GCATTGGCCGCGAGACTTTGCC-3' (SEQ ID NO:311)

5'-GCGGCCACGGTTCCTTGGAAATG-3' (SEQ ID NO:312)



-continued

A probe was also synthesized:

5'-TGGAGGAGCTCAACCTCAGCTACAACCGCATCACCAGCCCACAGG-3' (SEQ ID NO:313)

[1481] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO332 gene using the probe oligonucleotide and one of the PCR primers.

[1482] RNA for construction of the cDNA libraries was isolated from a human fetal liver library (LIB229).

[1483] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for DNA40982-1235 and the derived protein sequence for PRO332.

[1484] The entire nucleotide sequence of DNA40982-1235 is shown in **FIG. 107** (SEQ ID NO:309). Clone DNA40982-1235 contains a single open reading frame (with an apparent translational initiation site at nucleotide positions 342-344, as indicated in **FIG. 107**). The predicted polypeptide precursor is 642 amino acids long, and has a calculated molecular weight of 72,067 (pI: 6.60). Clone DNA40982-1235 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209433.

[1485] Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO332 shows about 30-40% amino acid sequence identity with a series of known proteoglycan sequences, including, for example, fibromodulin and fibromodulin precursor sequences of various species (FMOD\_BOVIN, FMOD\_CHICK, FMOD\_RAT, FMOD\_MOUSE, FMOD\_HUMAN, P\_R36773), osteomodulin sequences (AB000114\_1, AB007848\_1), decorin sequences (CFU83141\_1, OCU03394\_1, P\_R42266, P\_R42267, P\_R42260, P\_R89439), keratan sulfate proteoglycans (BTU48360\_1, AF022890\_1), corneal proteoglycan (AF022256\_1), and bone/cartilage proteoglycans and proteoglycane precursors (PGS1\_BOVIN, PGS2\_MOUSE, PGS2\_HUMAN).

#### Example 45

[1486] Isolation of cDNA Clones Encoding Human PRO334

[1487] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. Based on the consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO334.

[1488] Forward and reverse PCR primers were synthesized for the determination of PRO334:

forward PCR primer 5'-GATGGTTCTCTGCTCAAGTGCCCTG-3' (SEQ ID NO:316)

reverse PCR primer 5'-TTGCACTTGTAGGACCCACGTACG-3' (SEQ ID NO:317)

[1489] Additionally, a synthetic oligonucleotide hybridization probe was constructed for the determination of PRO334 which had the following nucleotide sequence

[1490] hybridization probe

[1491] 5'-CTGATGGGAGGACCTGTGTAGATGT-TGATGAATGTGCTACAGGAAGAGCC-3' (SEQ ID NO:318)

[1492] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO334 gene using the probe oligonucleotide and one of the PCR primers.

[1493] Human fetal kidney cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif.

[1494] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO334 [herein designated as DNA41379-1236] (SEQ ID NO:314) and the derived protein sequence for PRO334.

[1495] The entire nucleotide sequence of DNA41379-1236 (also referred to as UNQ295) is shown in **FIG. 109** (SEQ ID NO:314). Clone DNA41379-1236 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 203-205 and ending at the stop codon at nucleotide positions 1730-1732 (**FIG. 109**). The predicted polypeptide precursor is 509 amino acids long (**FIG. 110**). Clone DNA41379-1236 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209488.

[1496] Analysis of the amino acid sequence of the full-length PRO334 polypeptide suggests that portions of it possess significant homology to the fibulin and fibrillin proteins, thereby indicating that PRO334 may be a novel member of the EGF protein family.

#### Example 46

[1497] Isolation of cDNA Clones Encoding Human PRO346

[1498] A consensus DNA sequence was identified using phrap as described in Example 1 above. Specifically, this consensus sequence is herein designated DNA38240. Based on the DNA38240 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length PRO346 coding sequence.

[1499] RNA for construction of the cDNA libraries was isolated from human fetal liver. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents (e.g., Invitrogen, San Diego, Calif.; Clontech, etc.) The cDNA was primed with oligo dT containing a NotI site, linked with blunt to Sall hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

[1500] A cDNA clone was sequenced in entirety. The entire nucleotide sequence of DNA44167-1243 is shown in **FIG. 111** (SEQ ID NO:319). Clone DNA44167-1243 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 64-66 (**FIG. 11**; SEQ ID NO:319). The predicted polypeptide precursor is 450 amino acids long. Clone DNA44167-1243 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209434 (designation DNA44167-1243).

[1501] Based on a BLAST, BLAST-2 and FastA sequence alignment analysis (using the ALIGN computer program) of the full-length sequence, PRO346 shows amino acid sequence identity to carcinoembryonic antigen (28%).

[1502] The oligonucleotide sequences used in the above procedure were the following:

OLI2691 (38240.f1)  
5'-GATCCTGTCACAAAGCCAGTGGTGC-3' (SEQ ID NO:321)

OLI2693 (38240.r1)  
5'-CACTGACAGGGTTCCTCACCCAGG-3' (SEQ ID NO:322)

OLI2692 (38240.p1)  
5'-CTCCCTCTGGGGTGTGGAGTATGTGGGAACATGACCCTGACATG-3' (SEQ ID NO:323)

#### Example 47

[1503] Isolation of cDNA Clones Encoding Human PRO268

[1504] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA35698. Based on the DNA35698 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR, a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO268.

[1505] Forward and reverse PCR primers were synthesized:

forward PCR primer 1 5'-TGAGGTGGGCAAGCGCGAAATG-3' (SEQ ID NO:326)

forward PCR primer 2 5'-TATGTGGATCAGGACGTGCC-3' (SEQ ID NO:327)

forward PCR primer 3 5'-TGCAGGGTTCAGTCTAGATTG-3' (SEQ ID NO:328)

reverse PCR primer 5'-TTGAAGGACAAAGGCAATCTGCCAC-3' (SEQ ID NO:329)

[1506] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA35698 sequence which had the following nucleotide sequence

[1507] hybridization probe

[1508] 5'-GGAGTCTTGCAGTTCCCCTGGCAGTC-CTGGTGCTGTTGCTTTGGG-3' (SEQ ID NO:330)

[1509] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO268 gene using the probe oligonucleotide and one of the PCR primers.

[1510] RNA for construction of the cDNA libraries was isolated from human fetal lung tissue.

[1511] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO268 [herein designated as DNA39427-1179] (SEQ ID NO:324) and the derived protein sequence for PRO268.

[1512] The entire nucleotide sequence of DNA39427-1179 is shown in **FIG. 113** (SEQ ID NO:324). Clone DNA39427-1179 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 13-15 and ending at the stop codon at nucleotide positions 853-855 (**FIG. 113**). The predicted polypeptide

precursor is 280 amino acids long (**FIG. 114**). Clone DNA39427-1179 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209395.

[1513] Analysis of the amino acid sequence of the full-length PRO268 polypeptide suggests that it possess significant homology to protein disulfide isomerase, thereby indicating that PRO268 may be a novel protein disulfide isomerase.

#### Example 48

[1514] Isolation of cDNA Clones Encoding Human PRO330

[1515] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in

Example 1 above. This consensus sequence is herein designated DNA35730. Based on the DNA35730 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO330.

[1516] Forward and reverse PCR primers were synthesized:

forward PCR primer 1 5'-CCAGGCACAATTCAG-3' (SEQ ID NO:333)  
 forward PCR primer 2 5'-GGACCTTCTGTGCCAG-3' (SEQ ID NO:334)  
 reverse PCR primer 1 5'-GGTCTCAAGAACTCCTGTC-3' (SEQ ID NO:335)  
 reverse PCR primer 2 5'-ACACTCAGCATGCTGCTGTTG-3' (SEQ ID NO:336)

[1517] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which had the following nucleotide sequence

[1518] hybridization probe

[1519] 5'-GGGCACATGACTGACCTGATTIATGCA-GAGAAAGAGCTGGTGCAG-3' (SEQ ID NO:337)

forward PCR primer 1 5'-TCCCCAAGCCGTTCTAGACGCGG-3' (SEQ ID NO:342)  
 forward PCR primer 2 5'-CTGTTCTTCTTGCACG-3' (SEQ ID NO:343)  
 reverse PCR primer 5'-GCCCAAATGCCCTAAGCGGTATACCC-3' (SEQ ID NO:344)

[1520] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO330 gene using the probe oligonucleotide and one of the PCR primers.

[1521] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1522] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO330 [herein designated as DNA40603-1232] (SEQ ID NO:331) and the derived protein sequence for PRO330.

[1523] The entire nucleotide sequence of DNA40603-1232 is shown in FIG. 115 (SEQ ID NO:331). Clone DNA40603-1232 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 167-169 and ending at the stop codon at nucleotide positions 1766-1768 (FIG. 115). The predicted polypeptide precursor is 533 amino acids long (FIG. 116). Clone DNA40603-1232 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209486 on Nov. 21, 1997.

[1524] Analysis of the amino acid sequence of the full-length PRO330 polypeptide suggests that portions of it possess significant homology to the mouse prolyl 4-hydroxylase alpha subunit protein, thereby indicating that PRO330 may be a novel prolyl 4-hydroxylase alpha subunit polypeptide.

Example 49

[1525] Isolation of cDNA Clones Encoding Human PRO310

[1526] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA40553. Based on the DNA40553 consensus

sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO310.

[1527] Forward and reverse PCR primers were synthesized:

[1528] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which had the following nucleotide sequence

[1529] hybridization probe

[1530] 5'-GGGTGTGATGCTTGAAGCAITTTCT-GTGCTTTGATCACTATGCTAGGAC-3' (SEQ ID NO:345)

[1531] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO310 gene using the probe oligonucleotide and one of the PCR primers.

[1532] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1533] DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for PRO310 [herein designated as DNA43046-1225 (SEQ ID NO:340) and the derived protein sequence for PRO310 (SEQ ID NO:341).

[1534] The entire nucleotide sequence of DNA43046-1225 is shown in FIG. 119 (SEQ ID NO:340). Clone DNA43046-1225 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 81-83 and ending at the stop codon at nucleotide positions 1035-1037 (FIG. 119). The predicted polypeptide precursor is 318 amino acids long (FIG. 120) and has a calculated molecular weight of approximately 36,382 dal-

tons. Clone DNA43046-1225 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209484.

[1535] Analysis of the amino acid sequence of the full-length PRO310 polypeptide suggests that portions of it possess homology to *C. elegans* proteins and to fringe, thereby indicating that PRO310 may be involved in development.

#### Example 50

[1536] Isolation of cDNA Clones Encoding Human PRO339

[1537] An expressed sequence tag (EST) DNA database (LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.) was searched and ESTs were identified. An assembly of Incyte clones and a consensus sequence was formed using phrap as described in Example 1 above.

[1538] Forward and reverse PCR primers were synthesized based upon the assembly-created consensus sequence:

forward PCR primer 1 5'-GGGATGCAGGTGGTCTCATGGGG-3' (SEQ ID NO:346)

forward PCR primer 2 5'-CCCTCATGTACCGCTCC-3' (SEQ ID NO:347)

forward PCR primer 3 5'-GTGTGACACAGCGTGGGC-3' (SEQ ID NO:43)

forward PCR primer 4 5'-GACCGGCAGGCTTGTGCG-3' (SEQ ID NO:44)

reverse PCR primer 1 5'-CAGCAGCTTCAGCCACCAGGAGTGG-3' (SEQ ID NO:45)

reverse PCR primer 2 5'-CTGAGCCGTGGGCTGCAGTCTCGC-3' (SEQ ID NO:46)

[1539] Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus sequence which had the following nucleotide sequence

[1540] hybridization probe

[1541] 5'-CCGACTACGACTGGTTCTTCATCATG-CAGGATGACACATATGTGC-3' (SEQ ID NO:47)

2649-2651 (FIG. 117; SEQ ID NO:338). The predicted polypeptide precursor is 772 amino acids long and has a calculated molecular weight of approximately 86,226 daltons. Clone DNA43466-1225 has been deposited with ATCC and is assigned ATCC deposit no. ATCC 209490.

[1545] Based on a BLAST and FastA sequence alignment analysis (using the ALIGN computer program) of the full-length sequence, PRO339 has homology to *C. elegans* proteins and collagen-like polymer sequences as well as to fringe, thereby indicating that PRO339 may be involved in development or tissue growth.

#### Example 51

[1546] Isolation of cDNA Clones Encoding Human PRO244

[1547] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. Based on this consensus sequence, oli-

gonucleotides were synthesized to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for PRO244.

[1548] A pair of PCR primers (forward and reverse) were synthesized:

5'-TTCAGCTTCTGGGATGTAGGG-3' (30923.f1) (SEQ ID NO:378)

5'-TATTCCTACCATTTCACAAATCCG-3' (30923.r1) (SEQ ID NO:379)

[1542] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pairs identified above. A positive library was then used to isolate clones encoding the PRO339 gene using the probe oligonucleotide and one of the PCR primers.

[1543] RNA for construction of the cDNA libraries was isolated from human fetal liver tissue.

[1544] A cDNA clone was sequenced in entirety. The entire nucleotide sequence of DNA43466-1225 is shown in FIG. 117 (SEQ ID NO:338). Clone DNA43466-1225 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 333-335 and ending at the stop codon found at nucleotide positions

[1549] A probe was also synthesized:

[1550] 5'-GGAGGACTGTGCCACCAT-GAGAGACTCTTCAAACCCAAGGCAAAATTGG-3' (30923.p1) (SEQ ID NO:380)

[1551] In order to screen several libraries for a source of a full-length clone, DNA from the libraries was screened by PCR amplification with the PCR primer pair identified above. A positive library was then used to isolate clones encoding the PRO244 gene using the probe oligonucleotide and one of the PCR primers.

[1552] RNA for construction of the cDNA libraries was isolated from a human fetal kidney library. DNA sequencing of the clones isolated as described above gave the full-length DNA sequence and the derived protein sequence for PRO244.

[1553] The entire nucleotide sequence of PRO244 is shown in FIG. 121 (SEQ ID NO:376). Clone DNA35668-1171 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 106-108 (FIG. 121). The predicted polypeptide precursor is 219 amino acids long. Clone DNA35668-1171 has been deposited with ATCC (designated as DNA35663-1171) and is assigned ATCC deposit no. ATCC209371. The protein has a cytoplasmic domain (aa 1-20), a transmembrane domain (aa 2146), and an extracellular domain (aa 47-219), with a C-lectin domain at aa 55-206.

[1554] Based on a BLAST and FastA sequence alignment analysis of the full-length sequence, PRO244 shows notable amino acid sequence identity to hepatic lectin gallus gallus (43%), HIC hp120-binding C-type lectin (42%), macrophage lectin 2 (HUMHML2-1, 41%), and sequence PR32188 (44%).

#### Example 52

[1555] Use of PRO Polypeptide-Encoding Nucleic Acid as Hybridization Probes

[1556] The following method describes use of a nucleotide sequence encoding a PRO polypeptide as a hybridization probe.

[1557] DNA comprising the coding sequence of a PRO polypeptide of interest as disclosed herein may be employed as a probe or used as a basis from which to prepare probes to screen for homologous DNAs (such as those encoding naturally-occurring variants of the PRO polypeptide) in human tissue cDNA libraries or human tissue genomic libraries.

[1558] Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO polypeptide-encoding nucleic acid-derived probe to the filters is performed in a solution of 50% formamide, 5×SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2×Denhardt's solution, and 10% dextran sulfate at 42° C. for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1×SSC and 0.1% SDS at 42° C.

[1559] DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO polypeptide can then be identified using standard techniques known in the art.

#### Example 53

[1560] Expression of PRO Polypeptides in *E. coli*

[1561] This example illustrates preparation of an unglycosylated form of a desired PRO polypeptide by recombinant expression in *E. coli*.

[1562] The DNA sequence encoding the desired PRO polypeptide is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme

and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the specific PRO polypeptide coding region, lambda transcriptional terminator, and an argU gene.

[1563] The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

[1564] Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

[1565] After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO polypeptide can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

[1566] PRO187, PRO317, PRO301, PRO224 and PRO238 were successfully expressed in *E. coli* in apoly-His tagged form, using the following procedure. The DNA encoding PRO187, PRO317, PRO301, PRO224 or PRO238 was initially amplified using selected PCR primers. The primers contained restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences were then ligated into an expression vector, which was used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants were first grown in LB containing 50 mg/ml carbenicillin at 30° C. with shaking until an O.D.600 of 3-5 was reached. Cultures were then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.71 g sodium citrate.2H<sub>2</sub>O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO<sub>4</sub>) and grown for approximately 20-30 hours at 30° C. with shaking. Samples were removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets were frozen until purification and refolding.

[1567] *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) was resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution was stirred overnight at 4° C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution was centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant was diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine,

20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. Depending the clarified extract was loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column was washed with additional buffer containing 50 mM imidazole (Calbiochem, Ulrol grade), pH 7.4. The protein was eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein were pooled and stored at 4° C. Protein concentration was estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

[1568] The proteins were refolded by diluting sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes were chosen so that the final protein concentration was between 50 to 100 micrograms/ml. The refolding solution was stirred gently at 4° C. for 12-36 hours. The refolding reaction was quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution was filtered through a 0.22 micron filter and acetonitrile was added to 2-10% final concentration. The refolded protein was chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance were analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein were pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

[1569] Fractions containing the desired folded PRO187, PRO317, PRO301, PRO224 and PRO238 proteins, respectively, were pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins were formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

#### Example 54

[1570] Expression of PRO Polypeptides in Mammalian Cells

[1571] This example illustrates preparation of a glycosylated form of a desired PRO polypeptide by recombinant expression in mammalian cells.

[1572] The vector, pRK5 (see EP 307,247, published Mar. 15, 1989), is employed as the expression vector. Optionally, the PRO polypeptide-encoding DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO polypeptide DNA using ligation methods such as described in Sambrook et al., supra. The resulting vector is called pRK5-PRO polypeptide.

[1573] In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to

confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 µg pRK5-PRO polypeptide DNA is mixed with about 1 µg DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 µl of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl<sub>2</sub>. To this mixture is added, dropwise, 500 µl of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO<sub>4</sub>, and a precipitate is allowed to form for 10 minutes at 25° C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37° C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

[1574] Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 µCi/ml <sup>35</sup>S-cysteine and 200 µCi/ml <sup>35</sup>S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

[1575] In an alternative technique, PRO polypeptide may be introduced into 293 cells transiently using the dextran sulfate method described by Sompariyac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 µg pRK5-PRO polypeptide DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 µg/ml bovine insulin and 0.1 µg/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO polypeptide can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

[1576] In another embodiment, PRO polypeptides can be expressed in CHO cells. The pRK5-PRO polypeptide can be transfected into CHO cells using known reagents such as CaPO<sub>4</sub> or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as <sup>35</sup>S-methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO polypeptide can then be concentrated and purified by any selected method.

[1577] Epitope-tagged PRO polypeptide may also be expressed in host CHO cells. The PRO polypeptide may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector.

The poly-his tagged PRO polypeptide insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO polypeptide can then be concentrated and purified by any selected method, such as by Ni<sup>2+</sup>-chelate affinity chromatography.

[1578] PRO211, PRO217, PRO230, PRO219, PRO245, PRO221, PRO258, PRO301, PRO224, PRO222, PRO234, PRO229, PRO223, PRO328 and PRO332 were successfully expressed in CHO cells by both a transient and a stable expression procedure. In addition, PRO232, PRO265, PRO246, PRO228, PRO227, PRO220, PRO266, PRO269, PRO287, PRO214, PRO231, PRO233, PRO238, PRO244, PRO235, PRO236, PRO262, PRO239, PRO257, PRO260, PRO263, PRO270, PRO271, PRO272, PRO294, PRO295, PRO293, PRO247, PRO303 and PRO268 were successfully transiently expressed in CHO cells.

[1579] Stable expression in CHO cells was performed using the following procedure. The proteins were expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins were fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

[1580] Following PCR amplification, the respective DNAs were subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., *Nucl. Acids Res.* 24: 9 (1774-1779) (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

[1581] Twelve micrograms of the desired plasmid DNA were introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Quiagen), Dospers® or Fugene® (Boehringer Mannheim). The cells were grown and described in Lucas et al., supra. Approximately 3×10<sup>7</sup> cells are frozen in an ampule for further growth and production as described below.

[1582] The ampules containing the plasmid DNA were thawed by placement into water bath and mixed by vortexing. The contents were pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant was aspirated and the cells were resuspended in 10 mL of selective media (0.2 μm filtered PS20 with 5% 0.2 μm diafiltered fetal bovine serum). The cells were then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells were transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37° C. After another 2-3 days, a 250 mL, 500 mL and 2000 mL spinners were seeded with 3×10<sup>5</sup> cells/mL. The cell media was exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO

media may be employed, a production medium described in U.S. Pat. No. 5,122,469, issued Jun. 16, 1992 was actually used. 3L production spinner is seeded at 1.2×10<sup>6</sup> cells/mL. On day 0, the cell number pH were determined. On day 1, the spinner was sampled and sparging with filtered air was commenced. On day 2, the spinner was sampled, the temperature shifted to 33° C., and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion). Throughout the production, pH was adjusted as necessary to keep at around 7.2. After 10 days, or until viability dropped below 70%, the cell culture was harvested by centrifugation and filtering through a 0.22 μm filter. The filtrate was either stored at 4° C. or immediately loaded onto columns for purification.

[1583] For the poly-His tagged constructs, the proteins were purified using a Ni-NTA column (Qiagen). Before purification, imidazole was added to the conditioned media to a concentration of 5 mM. The conditioned media was pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 ml/min. at 4° C. After loading, the column was washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein was subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80° C.

[1584] Immunoadhesin (Fc containing) constructs of were purified from the conditioned media as follows. The conditioned medium was pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275 μL of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity was assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

[1585] PRO211, PRO217, PRO230, PRO232, PRO187, PRO265, PRO219, PRO246, PRO228, PRO533, PRO245, PRO221, PRO227, PRO220, PRO258, PRO266, PRO269, PRO287, PRO214, PRO317, PRO301, PRO224, PRO222, PRO234, PRO231, PRO229, PRO233, PRO238, PRO223, PRO235, PRO236, PRO262, PRO239, PRO257, PRO260, PRO263, PRO270, PRO271, PRO272, PRO294, PRO295, PRO293, PRO247, PRO304, PRO302, PRO307, PRO303, PRO343, PRO328, PRO326, PRO331, PRO332, PRO334, PRO346, PRO268, PRO330, PRO310 and PRO339 were also successfully transiently expressed in COS cells.

#### Example 55

[1586] Expression of PRO Polypeptides in Yeast

[1587] The following method describes recombinant expression of a desired PRO polypeptide in yeast.

[1588] First, yeast expression vectors are constructed for intracellular production or secretion of PRO polypeptides from the ADH2/GAPDH promoter. DNA encoding a desired

PRO polypeptide, a selected signal peptide and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of the PRO polypeptide. For secretion, DNA encoding the PRO polypeptide can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, the yeast alpha-factor secretory signal/leader sequence, and linker sequences (if needed) for expression of the PRO polypeptide.

[1589] Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

[1590] Recombinant PRO polypeptide can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing the PRO polypeptide may further be purified using selected column chromatography resins.

#### Example 56

[1591] Expression of PRO Polypeptides in Baculovirus-Infected Insect Cells

[1592] The following method describes recombinant expression of PRO polypeptides in Baculovirus-infected insect cells.

[1593] The desired PRO polypeptide is fused upstream of an epitope tag contained with a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the PRO polypeptide or the desired portion of the PRO polypeptide (such as the sequence encoding the extracellular domain of a transmembrane protein) is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

[1594] Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4-5 days of incubation at 28° C., the released viruses are harvested and used for further amplifications. Viral infection and protein expression is performed as described by O'Reilly et al., *Baculovirus expression vectors: A laboratory Manual*, Oxford: Oxford University Press (1994).

[1595] Expressed poly-his tagged PRO polypeptide can then be purified, for example, by Ni<sup>2+</sup>-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl<sub>2</sub>; 0.1 mM EDTA; 10% Glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant

is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% Glycerol, pH 7.8) and filtered through a 0.45 μm filter. A Ni<sup>2+</sup>-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A<sub>280</sub> with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% Glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A<sub>280</sub> baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or western blot with Ni<sup>2+</sup>-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His<sub>10</sub>-tagged PRO polypeptide are pooled and dialyzed against loading buffer.

[1596] Alternatively, purification of the IgG tagged (or Fc tagged) PRO polypeptide can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

[1597] PRO211, PRO217, PRO230, PRO187, PRO265, PRO246, PRO228, PRO533, PRO245, PRO221, PRO220, PRO258, PRO266, PRO269, PRO287, PRO214, PRO301, PRO224, PRO222, PRO234, PRO231, PRO229, PRO235, PRO239, PRO257, PRO272, PRO294, PRO295, PRO328, PRO326, PRO331, PRO334, PRO346 and PRO310 were successfully expressed in baculovirus infected Sf9 or high5 insect cells. While the expression was actually performed in a 0.5-2 L scale, it can be readily scaled up for larger (e.g. 8 L) preparations. The proteins were expressed as an IgG construct (immunoadhesin), in which the protein extracellular region was fused to an IgG1 constant region sequence containing the hinge, CH2 and CH3 domains and/or in poly-His tagged forms.

[1598] Following PCR amplification, the respective coding sequences were subcloned into a baculovirus expression vector (pb.PH.IgG for IgG fusions and pb.PH.His.c for poly-His tagged proteins), and the vector and BaculoGold® baculovirus DNA (Pharmingen) were co-transfected into 105 *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711), using Lipofectin (Gibco BRL). pb.PH.IgG and pb.PH.His are modifications of the commercially available baculovirus expression vector pVL1393 (Pharmingen), with modified polylinker regions to include the His or Fc tag sequences. The cells were grown in Hink's TNM-FH medium supplemented with 10% FBS (Hyclone). Cells were incubated for 5 days at 28° C. The supernatant was harvested and subsequently used for the first viral amplification by infecting Sf9 cells in Hink's TNM-FH medium supplemented with 10% FBS at an approximate multiplicity of infection (MOI) of 10. Cells were incubated for 3 days at 28° C. The supernatant was harvested and the expression of the constructs in the baculovirus expression vector was determined by batch binding of 1 ml of supernatant to 25 mL of Ni-NTA beads (QIAGEN) for histidine tagged proteins or Protein-A Sepharose CL-4B beads (Pharmacia) for IgG tagged proteins followed by SDS-PAGE analysis comparing to a known concentration of protein standard by Coomassie blue staining.

[1599] The first viral amplification supernatant was used to infect a spinner culture (500 ml) of Sf9 cells grown in



ESF-921 medium (Expression Systems LLC) at an approximate MOI of 0.1. Cells were incubated for 3 days at 28° C. The supernatant was harvested and filtered. Batch binding and SDS-PAGE analysis was repeated, as necessary, until expression of the spinner culture was confirmed.

[1600] The conditioned medium from the transfected cells (0.5 to 3 L) was harvested by centrifugation to remove the cells and filtered through 0.22 micron filters. For the poly-His tagged constructs, the protein construct were purified using a Ni-NTA column (Qiagen). Before purification, imidazole was added to the conditioned media to a concentration of 5 mM. The conditioned media were pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 45 ml/min. at 4° C. After loading, the column was washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein was subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at -80° C.

[1601] Immunoadhesin (Fc containing) constructs of proteins were purified from the conditioned media as follows. The conditioned media were pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column was washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein was immediately neutralized by collecting 1 ml fractions into tubes containing 275 mL of 1 M Tris buffer, pH 9. The highly purified protein was subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity of the proteins was verified by SDS polyacrylamide gel (PEG) electrophoresis and N-terminal amino acid sequencing by Edman degradation.

#### Example 57

[1602] Preparation of Antibodies that Bind to PRO Polypeptides

[1603] This example illustrates preparation of monoclonal antibodies which can specifically bind to a PRO polypeptide.

[1604] Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, supra. Immunogens that may be employed include purified PRO polypeptide, fusion proteins containing the PRO polypeptide, and cells expressing recombinant PRO polypeptide on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

[1605] Mice, such as Balb/c, are immunized with the PRO polypeptide immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, Mont.) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically

obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO polypeptide antibodies.

[1606] After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO polypeptide. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

[1607] The hybridoma cells will be screened in an ELISA for reactivity against the PRO polypeptide. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against the PRO polypeptide is within the skill in the art.

[1608] The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO polypeptide monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

#### Example 58

[1609] Chimeric PRO Polypeptides

[1610] PRO polypeptides may be expressed as chimeric proteins with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS™ extension/affinity purification system (Immunex Corp., Seattle Wash.). The inclusion of a cleavable linker sequence such as Factor XA or enterokinase (Invitrogen, San Diego Calif.) between the purification domain and the PRO polypeptide sequence may be useful to facilitate expression of DNA encoding the PRO polypeptide.

#### Example 59

[1611] Purification of PRO Polypeptides Using Specific Antibodies

[1612] Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polyp(ptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

[1613] Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate

or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKE Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

[1614] Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

[1615] A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

#### Example 60

##### [1616] Drug Screening

[1617] This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

[1618] Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the

ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

[1619] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on Sep. 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

[1620] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

#### Example 61

##### [1621] Rational Drug Design

[1622] The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (i.e., a PRO polypeptide) or of small molecules with which they interact, e.g., agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide in vivo (c.f., Hodgson, *Bio/Technology*, 9: 19-21(1991)).

[1623] In one approach, the three-dimensional structure of the PRO polypeptide, or of an PRO polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, *Biochemistry*, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda et al., *J. Biochem.*, 113:742-746 (1993).

[1624] It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site

of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

[1625] By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

#### Example 62

[1626] Diagnostic Test Using PRO317 Polypeptide-Specific Antibodies

[1627] Particular anti-PRO317 polypeptide antibodies are useful for the diagnosis of prepathologic conditions, and chronic or acute diseases such as gynecological diseases or ischemic diseases which are characterized by differences in the amount or distribution of PRO317. PRO317 has been found to be expressed in human kidney and is thus likely to be associated with abnormalities or pathologies which affect this organ. Further, since it is so closely related to EBAF-1, it is likely to affect the endometrium and other genital tissues. Further, due to library sources of certain ESTs, it appears that PRO317 may be involved as well in forming blood vessels and hence to be a modulator of angiogenesis.

[1628] Diagnostic tests for PRO317 include methods utilizing the antibody and a label to detect PRO317 in human body fluids, tissues, or extracts of such tissues. The polypeptide and antibodies of the present invention may be used with or without modification. Frequently, the polypeptide and antibodies will be labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, magnetic particles, and the like. Patents teaching the use of such labels include U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced as shown in U.S. Pat. No. 4,816,567.

[1629] A variety of protocols for measuring soluble or membrane-bound PRO317, using either polyclonal or monoclonal antibodies specific for that PRO317, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), radioreceptor assay (RRA), and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PRO317 is preferred, but a competitive binding assay may be employed. These assays are described, among other places, in Maddox et al. *J Exp. Med.*, 158:1211 (1983).

#### Example 63

[1630] Identification of PRO317 Receptors

[1631] Purified PRO317 is useful for characterization and purification of specific cell surface receptors and other

binding molecules. Cells which respond to PRO317 by metabolic changes or other specific responses are likely to express a receptor for PRO317. Such receptors include, but are not limited to, receptors associated with and activated by tyrosine and serine/threonine kinases. See Kolodziejczyk and Hall, *supra*, for a review on known receptors for the TGF-superfamily. Candidate receptors for this superfamily fall into two primary groups, termed type I and type II receptors. Both types are serine/threonine kinases. Upon activation by the appropriate ligand, type I and type II receptors physically interact to form hetero-oligomers and subsequently activate intracellular signaling cascades, ultimately regulating gene transcription and expression. In addition, TGF-binds to a third receptor class, type III, a membrane-anchored proteoglycan lacking the kinase activity typical of signal transducing molecules.

[1632] PRO317 receptors or other PRO317-binding molecules may be identified by interaction with radiolabeled PRO317. Radioactive labels may be incorporated into PRO317 by various methods known in the art. A preferred embodiment is the labeling of primary amino groups in PRO317 with <sup>125</sup>I Bolton-Hunter reagent (Bolton and Hunter, *Biochem. J.*, 133:529 (1973)), which has been used to label other polypeptides without concomitant loss of biological activity (Hebert et al., *J. Biol. Chem.*, 266:18989 (1991); McColl et al., *J. Immunol.*, 150:4550-4555 (1993)). Receptor-bearing cells are incubated with labeled PRO317. The cells are then washed to remove unbound PRO317, and receptor-bound PRO317 is quantified. The data obtained using different concentrations of PRO317 are used to calculate values for the number and affinity of receptors.

[1633] Labeled PRO317 is useful as a reagent for purification of its specific receptor. In one embodiment of affinity purification, PRO317 is covalently coupled to a chromatography column. Receptor-bearing cells are extracted, and the extract is passed over the column. The receptor binds to the column by virtue of its biological affinity for PRO317. The receptor is recovered from the column and subjected to N-terminal protein sequencing. This amino acid sequence is then used to design degenerate oligonucleotide probes for cloning the receptor gene.

[1634] In an alternative method, mRNA is obtained from receptor-bearing cells, and made into a cDNA library. The library is transfected into a population of cells, and those cells expressing the receptor are selected using fluorescently labeled PRO317. The receptor is identified by recovering and sequencing recombinant DNA from highly labeled cells.

[1635] In another alternative method, antibodies are raised against the surface of receptor bearing cells, specifically monoclonal antibodies. The monoclonal antibodies are screened to identify those which inhibit the binding of labeled PRO317. These monoclonal antibodies are then used in affinity purification or expression cloning of the receptor.

[1636] Soluble receptors or other soluble binding molecules are identified in a similar manner. Labeled PRO317 is incubated with extracts or other appropriate materials derived from the uterus. After incubation, PRO317 complexes larger than the size of purified PRO317 are identified by a sizing technique such as size-exclusion chromatography or density gradient centrifugation and are purified by methods known in the art. The soluble receptors or binding protein(s) are subjected to N-terminal sequencing to obtain

information sufficient for database identification, if the soluble protein is known, or for cloning, if the soluble protein is unknown.

#### Example 64

**[1637]** Determination of PRO317-Induced Cellular Response

**[1638]** The biological activity of PRO317 is measured, for example, by binding of an PRO317 of the invention to an PRO317 receptor. A test compound is screened as an antagonist for its ability to block binding of PRO317 to the receptor. A test compound is screened as an agonist of the PRO317 for its ability to bind an PRO317 receptor and influence the same physiological events as PRO317 using, for example, the KIRA-ELISA assay described by Sadick et al., *Analytical Biochemistry*, 235:207-214 (1996) in which activation of a receptor tyrosine kinase is monitored by immuno-capture of the activated receptor and quantitation of the level of ligand-induced phosphorylation. The assay may be adapted to monitor PRO317-induced receptor activation through the use of an PRO317 receptor-specific antibody to capture the activated receptor. These techniques are also applicable to other PRO polypeptides described herein.

#### Example 65

**[1639]** Use of PRO224 for Screening Compounds

**[1640]** PRO224 is expressed in a cell stripped of membrane proteins and capable of expressing PRO224. Low density lipoproteins having a detectable label are added to the cells and incubated for a sufficient time for endocytosis. The cells are washed. The cells are then analysed for label bound to the membrane and within the cell after cell lysis. Detection of the low density lipoproteins within the cell determines that PRO224 is within the family of low density lipoprotein receptor proteins. Members found within this family are then used for screening compounds which affect these receptors, and particularly the uptake of cholesterol via these receptors.

#### Example 66

**[1641]** Ability of PRO Polypeptides to Inhibit Vascular Endothelial Growth Factor (VEGF) Stimulated Proliferation of Endothelial Cell Growth (Assay 9)

**[1642]** The ability of various PRO polypeptides to inhibit VEGF stimulated proliferation of endothelial cells was tested. Polypeptides testing positive in this assay are useful for inhibiting endothelial cell growth in mammals where such an effect would be beneficial, e.g., for inhibiting tumor growth.

**[1643]** Specifically, bovine adrenal cortical capillary endothelial cells (ACE) (from primary culture, maximum of 12-14 passages) were plated in 96-well plates at 500 cells/well per 100 microliter. Assay media included low glucose DMEM, 10% calf serum, 2 mM glutamine, and 1xpenicillin/streptomycin/fungizone. Control wells included the following: (1) no ACE cells added; (2) ACE cells alone; (3) ACE cells plus 5 ng/ml FGF; (4) ACE cells plus 3 ng/ml VEGF; (5) ACE cells plus 3 ng/ml VEGF plus 1 ng/ml TGF-beta; and (6) ACE cells plus 3 ng/ml VEGF plus 5 ng/ml LIF. The test samples, poly-his tagged PRO polypeptides (in 100 microliter volumes), were then added to the wells (at dilu-

tions of 1%, 0.1% and 0.01%, respectively). The cell cultures were incubated for 6-7 days at 37° C./5% CO<sub>2</sub>. After the incubation, the media in the wells was aspirated, and the cells were washed 1x with PBS. An acid phosphatase reaction mixture (100 microliter; 0.1M sodium acetate, pH 5.5, 0.1% Triton X-100, 10 mM p-nitrophenyl phosphate) was then added to each well. After a 2 hour incubation at 37° C., the reaction was stopped by addition of 10 microliters 1N NaOH. Optical density (OD) was measured on a microplate reader at 405 nm.

**[1644]** The activity of PRO polypeptides was calculated as the percent inhibition of VEGF (3 ng/ml) stimulated proliferation (as determined by measuring acid phosphatase activity at OD 405 nm) relative to the cells without stimulation. TGF-beta was employed as an activity reference at 1 ng/ml, since TGF-beta blocks 70-90% of VEGF-stimulated ACE cell proliferation. The results are indicative of the utility of the PRO polypeptides in cancer therapy and specifically in inhibiting tumor angiogenesis. Numerical values (relative inhibition) are determined by calculating the percent inhibition of VEGF stimulated proliferation by the PRO polypeptides relative to cells without stimulation and then dividing that percentage into the percent inhibition obtained by TGF-β at 1 ng/ml which is known to block 70-90% of VEGF stimulated cell proliferation. The results are considered positive if the PRO polypeptide exhibits 30% or greater inhibition of VEGF stimulation of endothelial cell growth (relative inhibition 30% or greater).

**[1645]** The following polypeptides tested positive in this assay: PRO211, PRO217, PRO187, PRO219, PRO246, PRO228, PRO245, PRO221, PRO258, PRO301, PRO224, PRO272, PRO328, PRO331, PRO224, PRO328, PRO272, PRO301, PRO331 and PRO214.

#### Example 67

**[1646]** Retinal Neuron Survival (Assay 52)

**[1647]** This example demonstrates that certain PRO polypeptides have efficacy in enhancing the survival of retinal neuron cells and, therefore, are useful for the therapeutic treatment of retinal disorders or injuries including, for example, treating sight loss in mammals due to retinitis pigmentosum, AMD, etc.

**[1648]** Sprague Dawley rat pups at postnatal day 7 (mixed population: glia and retinal neuronal types) are killed by decapitation following CO<sub>2</sub> anesthesia and the eyes are removed under sterile conditions. The neural retina is dissected away from the pigment epithelium and other ocular tissue and then dissociated into a single cell suspension using 0.25% trypsin in Ca<sup>2+</sup>, Mg<sup>2+</sup>-free PBS. The retinas are incubated at 37° C. for 7-10 minutes after which the trypsin is inactivated by adding 1 ml soybean trypsin inhibitor. The cells are plated at 100,000 cells per well in 96 well plates in DMEM/F12 supplemented with N2 and with or without the specific test PRO polypeptide. Cells for all experiments are grown at 37° C. in a water saturated atmosphere of 5% CO<sub>2</sub>. After 2-3 days in culture, cells are stained with calcein AM then fixed using 4% paraformaldehyde and stained with DAPI for determination of total cell count. The total cells (fluorescent) are quantified at 20x objective magnification using CCD camera and NIH image software for MacIntosh. Fields in the well are chosen at random.

[1649] The effect of various concentration of PRO polypeptides are reported herein where percent survival is calculated by dividing the total number of calcein AM positive cells at 2-3 days in culture by the total number of DAPI-labeled cells at 2-3 days in culture. Anything above 30% survival is considered positive.

[1650] The following PRO polypeptides tested positive in this assay using polypeptide concentrations within the range of 0.01% to 1.0% in the assay: PRO220 and PRO346.

#### Example 68

[1651] Rod Photoreceptor Cell Survival (Assay 56)

[1652] This assay shows that certain polypeptides of the invention act to enhance the survival/proliferation of rod photoreceptor cells and, therefore, are useful for the therapeutic treatment of retinal disorders or injuries including, for example, treating sight loss in mammals due to retinitis pigmentosa, AMD, etc. Sprague Dawley rat pups at 7 day postnatal (mixed population: glia and retinal neuronal cell types) are killed by decapitation following CO<sub>2</sub> anesthesia and the eyes are removed under sterile conditions. The neural retina is dissected away from the pigment epithelium and other ocular tissue and then dissociated into a single cell suspension using 0.25% trypsin in Ca<sup>2+</sup>, Mg<sup>2+</sup>-free PBS. The retinas are incubated at 37° C. for 7-10 minutes after which the trypsin is inactivated by adding 1 ml soybean trypsin inhibitor. The cells are plated at 100,000 cells per well in 96 well plates in DMEM/F12 supplemented with N<sub>2</sub>. Cells for all experiment; are grown at 37° C. in a water saturated atmosphere of 5% CO<sub>2</sub>. After 2-3 days in culture, cells are fixed using 4% paraformaldehyde, and then stained using CellTracker Green CMFDA. Rho 4D2 (ascites or IgG 1:100), a monoclonal antibody directed towards the visual pigment rhodopsin is used to detect rod photoreceptor cells by indirect immunofluorescence. The results are calculated as % survival: total number of calcein—rhodopsin positive cells at 2-3 days in culture, divided by the total number of rhodopsin positive cells at time 2-3 days in culture. The total cells (fluorescent) are quantified at 20x objective magnification using a CCD camera and NIH image software for Macintosh. Fields in the well are chosen at random.

[1653] The following polypeptides tested positive in this assay: PRO220 and PRO346.

#### Example 69

[1654] Induction of Endothelial Cell Apoptosis (Assay 73)

[1655] The ability of PRO polypeptides to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems). A positive test in the assay is indicative of the usefulness of the polypeptide in therapeutically treating tumors as well as vascular disorders where inducing apoptosis of endothelial cells would be beneficial.

[1656] The cells were plated on 96-well microtiter plates (Amersham Life Science, cytostar-T scintillating microplate, RPNQ160, sterile, tissue-culture treated, individually wrapped), in 10% serum (CSG-medium, Cell Systems), at a density of 2x10<sup>4</sup> cells per well in a total volume of 100 μL. On day 2, test samples containing the PRO polypeptide were added in triplicate at dilutions of 1%, 0.33% and 0.11%. Wells without cells were used as a blank and wells with cells

only were used as a negative control. As; a positive control 1:3 serial dilutions of 50 μL of a 3xstock of staurosporine were used. The ability of the PRO polypeptide to induce apoptosis was determined by processing of the 96 well plates for detection of Annexin V, a member of the calcium and phospholipid binding proteins, to detect apoptosis.

[1657] 0.2 ml Annexin V—Biotin stock solution (100 μg/ml) was diluted in 4.6 ml 2xCa<sup>2+</sup> binding buffer and 2.5% BSA (1:25 dilution). 50 μL of the diluted Annexin V—Biotin solution was added to each well (except controls) to a final concentration of 1.0 μg/ml. The samples were incubated for 10-15 minutes with Annexin-Biotin prior to direct addition of <sup>35</sup>S-Streptavidin. <sup>35</sup>S-Streptavidin was diluted in 2xCa<sup>2+</sup> Binding buffer, 2.5% BSA and was added to all wells at a final concentration of 3x10<sup>4</sup> cpm/well. The plates were then sealed, centrifuged at 1000 rpm for 15 minutes and placed on orbital shaker for 2 hours. The analysis was performed on a 1450 Microbeta Trilux (Wallac). Percent above background represents the percentage amount of counts per minute above the negative controls. Percents greater than or equal to 30% above background are considered positive.

[1658] The following PRO polypeptides tested positive in this assay: PRO228, PRO217 and PRO301.

#### Example 70

[1659] PDB12 Cell Inhibition (Assay 40)

[1660] This example demonstrates that various PRO polypeptides have efficacy in inhibiting protein production by PDB12 pancreatic ductal cells and are, therefore, useful in the therapeutic treatment of disorders which involve protein secretion by the pancreas, including diabetes, and the like.

[1661] PDB12 pancreatic ductal cells are plated on fibronectin coated 96 well plates at 1.5x10<sup>3</sup> cells per well in 100 μL/180 μL of growth media. 100 μL of growth media with the PRO polypeptide test sample or negative control lacking the PRO polypeptide is then added to well, for a final volume of 200 μL. Controls contain growth medium containing a protein shown to be inactive in this assay. Cells are incubated for 4 days at 37° C. 20 μL of Alamar Blue Dye (AB) is then added to each well and the fluorescent reading is measured at 4 hours post addition of AB, on a microtiter plate reader at 530 nm excitation and 590 nm emission. The standard employed is cells without Bovine Pituitary Extract (BPE) and with various concentrations of BPE. Buffer or CM controls from unknowns are run 2 times on each 96 well plate.

[1662] These assays allow one to calculate a percent decrease in protein production by comparing the Alamar Blue Dye calculated protein concentration produced by the PRO polypeptide-treated cells with the Alamar Blue Dye calculated protein concentration produced by the negative control cells. A percent decrease in protein production of greater than or equal to 25% as compared to the negative control cells is considered positive.

[1663] The following polypeptides tested positive in this assay: PRO211, PRO287, PRO301 and PRO293.

#### Example 71

[1664] Stimulation of Adult Heart Hypertrophy (Assay 2)

[1665] This assay is designed to measure the ability of various PRO polypeptides to stimulate hypertrophy of adult

heart. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of various cardiac insufficiency disorders.

[1666] Ventricular myocytes freshly isolated from adult (250 g) Sprague Dawley rats are plated at 2000 cell/well in 180  $\mu$ L volume. Cells are isolated and plated on day 1, the PRO polypeptide-containing test samples or growth medium only (negative control) (20  $\mu$ L volume) is added on day 2 and the cells are then fixed and stained on day 5. After staining, cell size is visualized wherein cells showing no growth enhancement as compared to control cells are given a value of 0.0, cells showing small to moderate growth enhancement as compared to control cells are given a value of 1.0 and cells showing large growth enhancement as compared to control cells are given a value of 2.0. Any degree of growth enhancement as compared to the negative control cells is considered positive for the assay.

[1667] The following PRO polypeptides tested positive in this assay: PRO287, PRO301, PRO293 and PRO303.

#### Example 72

[1668] PDB12 Cell Proliferation (Assay 29)

[1669] This example demonstrates that various PRO polypeptides have efficacy in inducing proliferation of PDB12 pancreatic ductal cells and are, therefore, useful in the therapeutic treatment of disorders which involve protein secretion by the pancreas, including diabetes, and the like.

[1670] PDB12 pancreatic ductal cells are plated on fibronectin coated 96 well plates at  $1.5 \times 10^3$  cells per well in 100  $\mu$ L/180  $\mu$ L of growth media. 100  $\mu$ L of growth media with the PRO polypeptide test sample or negative control lacking the PRO polypeptide is then added to well, for a final volume of 200  $\mu$ L. Controls contain growth medium containing a protein shown to be inactive in this assay. Cells are incubated for 4 days at 37° C. 20  $\mu$ L of Alamar Blue Dye (AB) is then added to each well and the fluorescent reading is measured at 4 hours post addition of AB, on a microtiter plate reader at 530 nm excitation and 590 nm emission. The standard employed is cells without Bovine Pituitary Extract (BPE) and with various concentrations of BPE. Buffer or growth medium only controls from unknowns are run 2 times on each 96 well plate.

[1671] Percent increase in protein production is calculated by comparing the Alamar Blue Dye calculated protein concentration produced by the PRO polypeptide-treated cells with the Alamar Blue Dye calculated protein concentration produced by the negative control cells. A percent increase in protein production of greater than or equal to 25% as compared to the negative control cells is considered positive.

[1672] The following PRO polypeptides tested positive in this assay: PRO301 and PRO303.

#### Example 73

[1673] Enhancement of Heart Neonatal Hypertrophy (Assay 1)

[1674] This assay is designed to measure the ability of PRO polypeptides to stimulate hypertrophy of neonatal heart. PRO polypeptides testing positive in this assay are

expected to be useful for the therapeutic treatment of various cardiac insufficiency disorders.

[1675] Cardiac myocytes from 1-day old Harlan Sprague Dawley rats were obtained. Cells (180  $\mu$ L at  $7.5 \times 10^4$ /ml, serum <0.1%, freshly isolated) are added on day 1 to 96-well plates previously coated with DMEM/F12+4% FCS. Test samples containing the test PRO polypeptide or growth medium only (negative control) (20  $\mu$ L/well) are added directly to the wells on day 1. PGF (20  $\mu$ L/well) is then added on day 2 at final concentration  $10^{-6}$  M. The cells are then stained on day 4 and visually scored on day 5, wherein cells showing no increase in size as compared to negative controls are scored 0.0, cells showing a small to moderate increase in size as compared to negative controls are scored 1.0 and cells showing a large increase in size as compared to negative controls are scored 2.0. A positive result in the assay is a score of 1.0 or greater.

[1676] The following polypeptides tested positive in this assay: PRO224 and PRO231.

#### Example 74

[1677] Stimulatory Activity in Mixed Lymphocyte Reaction (MLR) Assay (Assay 24)

[1678] This example shows that certain polypeptides of the invention are active as a stimulator of the proliferation of stimulated T-lymphocytes. Compounds which stimulate proliferation of lymphocytes are useful therapeutically where enhancement of an immune response is beneficial. A therapeutic agent may take the form of antagonists of the polypeptide of the invention, for example, murine-human chimeric, humanized or human antibodies against the polypeptide.

[1679] The basic protocol for this assay is described in Current Protocols in Immunology, unit 3.12; edited by J E Coligan, A M Kruisbeek, D H Marglies, E M Shevach, W Strober, National Institutes of Health, Published by John Wiley & Sons, Inc.

[1680] More specifically, in one assay variant, peripheral blood mononuclear cells (PBMC) are isolated from mammalian individuals, for example a human volunteer, by leukopheresis (one donor will supply stimulator PBMCs, the other donor will supply responder PBMCs). If desired, the cells are frozen in fetal bovine serum and DMSO after isolation. Frozen cells may be thawed overnight in assay media (37° C., 5% CO<sub>2</sub>) and then washed and resuspended to  $3 \times 10^6$  cells/ml of assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate). The stimulator PBMCs are prepared by irradiating the cells (about 3000 Rads).

[1681] The assay is prepared by plating in triplicate wells a mixture of:

[1682] 100:1 of test sample diluted to 1% or to 0.1%,

[1683] 50:1 of irradiated stimulator cells, and

[1684] 50:1 of responder PBMC cells.

[1685] 100 microliters of cell culture media or 100 microliter of CD4-IgG is used as the control. The wells are then incubated at 37° C., 5% CO<sub>2</sub> for 4 days. On day 5, each well is pulsed with tritiated thymidine (1.0 mCi/well; Amersham).

After 6 hours the cells are washed 3 times and then the uptake of the label is evaluated.

[1686] In another variant of this assay, PBMCs are isolated from the spleens of Balb/c mice and C57B6 mice. The cells are teased from freshly harvested spleens in assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate) and the PBMCs are isolated by overlaying these cells over Lympholyte M (Organon Teknika), centrifuging at 2000 rpm for 20 minutes, collecting and washing the mononuclear cell layer in assay media and resuspending the cells to  $1 \times 10^7$  cells/ml of assay media. The assay is then conducted as described above.

[1687] Positive increases over control are considered positive with increases of greater than or equal to 180% being preferred. However, any value greater than control indicates a stimulatory effect for the test protein.

[1688] The following PRO polypeptides tested positive in this assay: PRO245, PRO269, PRO217, PRO301, PRO266, PRO335, PRO331, PRO533 and PRO326.

#### Example 75

[1689] Pericyte c-Fos Induction (Assay 93)

[1690] This assay shows that certain polypeptides of the invention act to induce the expression of c-fos in pericyte cells and, therefore, are useful not only as diagnostic markers for particular types of pericyte-associated tumors but also for giving rise to antagonists which would be expected to be useful for the therapeutic treatment of pericyte-associated tumors. Specifically, on day 1, pericytes are received from VEC Technologies and all but 5 ml of media is removed from flask. On day 2, the pericytes are trypsinized, washed, spun and then plated onto 96 well plates. On day 7, the media is removed and the pericytes are treated with 100  $\mu$ l of PRO polypeptide test samples and controls (positive control=DME+5% serum+/-PDGF at 500 ng/ml; negative control=protein 32). Replicates are averaged and SD/CV are determined. Fold increase over Protein 32 (buffer control) value indicated by chemiluminescence units (RLU) luminometer reading verses frequency is plotted on a histogram. Two-fold above Protein 32 value is considered positive for the assay. ASY Matrix: Growth media=low glucose DMEM=20% FBS+1 $\times$ pen strep+1 $\times$ fungizone. Assay Media=low glucose DMEM+5% FBS.

[1691] The following polypeptides tested positive in this assay: PRO214, PRO219, PRO221 and PRO224.

#### Example 76

[1692] Ability of PRO Polypeptides to Stimulate the Release of Proteoglycans from Cartilage (Assay 97)

[1693] The ability of various PRO polypeptides to stimulate the release of proteoglycans from cartilage tissue was tested as follows.

[1694] The metacarpophalangeal joint of 4-6 month old pigs was aseptically dissected, and articular cartilage was removed by free hand slicing being careful to avoid the underlying bone. The cartilage was minced and cultured in bulk for 24 hours in a humidified atmosphere of 95% air, 5% CO<sub>2</sub> in serum free (SF) media (DME/F12 1:1) with 0.1% BSA and 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin.

After washing three times, approximately 100 mg of articular cartilage was aliquoted into microtubes and incubated for an additional 24 hours in the above SF media. PRO polypeptides were then added at 1% either alone or in combination with 18 ng/ml interleukin-1 $\alpha$ , a known stimulator of proteoglycan release from cartilage tissue. The supernatant was then harvested and assayed for the amount of proteoglycans using the 1,9-dimethyl-methylene blue (DMB) colorimetric assay (Farndale and Buttle, *Biochem. Biophys. Acta* 883:173-177 (1985)). A positive result in this assay indicates that the test polypeptide will find use, for example, in the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis.

[1695] When various PRO polypeptides were tested in the above assay, the polypeptides demonstrated a marked ability to stimulate release of proteoglycans from cartilage tissue both basally and after stimulation with interleukin-1 $\alpha$  and at 24 and 72 hours after treatment, thereby indicating that these PRO polypeptides are useful for stimulating proteoglycan release from cartilage tissue. As such, these PRO polypeptides are useful for the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis. The polypeptides testing positive in this assay are: PRO211.

#### Example 77

[1696] Skin Vascular Permeability Assay (Assay 64)

[1697] This assay shows that certain polypeptides of the invention stimulate an immune response and induce inflammation by inducing mononuclear cell, eosinophil and PMN infiltration at the site of injection of the animal. Compounds which stimulate an immune response are useful therapeutically where stimulation of an immune response is beneficial. This skin vascular permeability assay is conducted as follows. Hairless guinea pigs weighing 350 grams or more are anesthetized with ketamine (75-80 mg/Kg) and 5 mg/Kg xylazine intramuscularly (IM). A sample of purified polypeptide of the invention or a conditioned media test sample is injected intradermally onto the backs of the test animals with 100  $\mu$ l per injection site. It is possible to have about 10-30, preferably about 16-24, injection sites per animal. One  $\mu$ l of Evans blue dye (1% in physiologic buffered saline) is injected intracardially. Blemishes at the injection sites are then measured (mm diameter) at 1 hr and 6 hr post injection. Animals were sacrificed at 6 hrs after injection. Each skin injection site is biopsied and fixed in formalin. The skins are then prepared for histopathologic evaluation. Each site is evaluated for inflammatory cell infiltration into the skin. Sites with visible inflammatory cell inflammation are scored as positive. Inflammatory cells may be neutrophilic, eosinophilic, monocytic or lymphocytic. At least a minimal perivascular infiltrate at the injection site is scored as positive, no infiltrate at the site of injection is scored as negative.

[1698] The following polypeptides tested positive in this assay: PRO245, PRO217, PRO326, PRO266, PRO272, PRO301, PRO331 and PRO335.

#### Example 78

[1699] Enhancement of Heart Neonatal Hypertrophy Induced by F2a (Assay 37)

[1700] This assay is designed to measure the ability of PRO polypeptides to stimulate hypertrophy of neonatal

heart. PRO polypeptides testing positive in this assay are expected to be useful for the therapeutic treatment of various cardiac insufficiency disorders.

[1701] Cardiac myocytes from 1-day old Harlan Sprague Dawley rats were obtained. Cells ( $180 \mu\text{l}$  at  $7.5 \times 10^4/\text{ml}$ , serum  $<0.1\%$ , freshly isolated) are added on day 1 to 96-well plates previously coated with DMEM/F12+4% FCS. Test samples containing the test PRO polypeptide ( $20 \mu\text{l}/\text{well}$ ) are added directly to the wells on day 1. PGF ( $20 \mu\text{l}/\text{well}$ ) is then added on day 2 at a final concentration of  $10^{-6}$  M. The cells are then stained on day 4 and visually scored on day 5. Visual scores are based on cell size, wherein cells showing no increase in size as compared to negative controls are scored 0.0, cells showing a small to moderate increase in size as compared to negative controls are scored 1.0 and cells showing a large increase in size as compared to negative controls are scored 2.0. A score of 1.0 or greater is considered positive.

[1702] No PBS is included, since calcium concentration is critical for assay response. Plates are coated with DMEM/F12 plus 4% FCS ( $200 \mu\text{l}/\text{well}$ ). Assay media included: DMEM/F12 (with 2.44 gm bicarbonate),  $10 \mu\text{g}/\text{ml}$  transferrin,  $1 \mu\text{g}/\text{ml}$  insulin,  $1 \mu\text{g}/\text{ml}$  aprotinin, 2 mmol/L glutamine, 100 U/ml penicillin G,  $100 \mu\text{g}/\text{ml}$  streptomycin. Protein buffer containing mannitol (4%) gave a positive signal (score 3.5) at 1/10 (0.4%) and 1/100 (0.04%), but not at 1/1000 (0.004%). Therefore the test sample buffer containing mannitol is not run.

[1703] The following PRO polypeptides tested positive in this assay: PRO224.

#### Example 79

[1704] Inhibitory Activity in Mixed Lymphocyte Reaction (MLR) Assay (Assay 67)

[1705] This example shows that one or more of the polypeptides of the invention are active as inhibitors of the proliferation of stimulated T-lymphocytes. Compounds which inhibit proliferation of lymphocytes are useful therapeutically where suppression of an immune response is beneficial.

[1706] The basic protocol for this assay is described in Current Protocols in Immunology, unit 3.12; edited by J E Coligan, A M Kruisbeek, D H Marglies, E M Shevach, W Strober, National Institutes of Health, Published by John Wiley & Sons, Inc.

[1707] More specifically, in one assay variant, peripheral blood mononuclear cells (PBMC) are isolated from mammalian individuals, for example a human volunteer, by leukopheresis (one donor will supply stimulator PBMCs, the other donor will supply responder PBMCs). If desired, the cells are frozen in fetal bovine serum and DMSO after isolation. Frozen cells may be thawed overnight in assay media ( $37^\circ \text{C}$ ., 5%  $\text{CO}_2$ ) and then washed and resuspended to  $3 \times 10^6$  cells/ml of assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate). The stimulator PBMCs are prepared by irradiating the cells (about 3000 Rads).

[1708] The assay is prepared by plating in triplicate wells a mixture of:

[1709] 100:1 of test sample diluted to 1% or to 0.1%,

[1710] 50:1 of irradiated stimulator cells, and

[1711] 50:1 of responder PBMC cells.

[1712] 100 microliters of cell culture media or 100 microliter of CD4-IgG is used as the control. The wells are then incubated at  $37^\circ \text{C}$ ., 5%  $\text{CO}_2$  for 4 days. On day 5, each well is pulsed with tritiated thymidine ( $1.0 \text{ mCi}/\text{well}$ ; Amersham). After 6 hours the cells are washed 3 times and then the uptake of the label is evaluated.

[1713] In another variant of this assay, PBMCs are isolated from the spleens of Balb/c mice and C57B6 mice. The cells are teased from freshly harvested spleens in assay media (RPMI; 10% fetal bovine serum, 1% penicillin/streptomycin, 1% glutamine, 1% HEPES, 1% non-essential amino acids, 1% pyruvate) and the PBMCs are isolated by overlaying these cells over Lympholyte M (Organon Teknika), centrifuging at 2000 rpm for 20 minutes, collecting and washing the mononuclear cell layer in assay media and resuspending the cells to  $1 \times 10^7$  cells/ml of assay media. The assay is then conducted as described above.

[1714] Any decreases below control is considered to be a positive result for an inhibitory compound, with decreases of less than or equal to 80% being preferred. However, any value less than control indicates an inhibitory effect for the test protein.

[1715] The following polypeptide tested positive in this assay: PRO235, PRO245 and PRO332.

#### Example 80

[1716] Induction of Endothelial Cell Apoptosis (ELISA) (Assay 109)

[1717] The ability of PRO polypeptides to induce apoptosis in endothelial cells was tested in human venous umbilical vein endothelial cells (HUVEC, Cell Systems) using a 96-well Format, in 0% serum media supplemented with 100 ng/ml VEGF, 0.1% BSA, 1xpenn/strep. A positive result in this assay indicates the usefulness of the polypeptide for therapeutically treating any of a variety of conditions associated with undesired endothelial cell growth including, for example, the inhibition of tumor growth. The 96-well plates used were manufactured by Falcon (No. 3072). Coating of 96 well plates were prepared by allowing gelatinization to occur for  $>30$  minutes with  $100 \mu\text{l}$  of 0.2% gelatin in PBS solution. The gelatin mix was aspirated thoroughly before plating HUVEC cells at a final concentration of  $2 \times 10^4$  cells/ml in 10% serum containing medium -  $100 \mu\text{l}$  volume per well. The cells were grown for 24 hours before adding test samples containing the PRO polypeptide of interest.

[1718] To all wells,  $100 \mu\text{l}$  of 0% serum media (Cell Systems) complemented with 100 ng/ml VEGF, 0.1% BSA, 1xpenn/strep was added. Test samples containing PRO polypeptides were added in triplicate at dilutions of 1%, 0.33% and 0.11%. Wells without cells were used as a blank and wells with cells only were used as a negative control. As a positive control, 1:3 serial dilutions of  $50 \mu\text{l}$  of a 3xstock of staurosporine were used. The cells were incubated for 24 to 35 hours prior to ELISA.



[1719] ELISA was used to determine levels of apoptosis preparing solutions according to the Boehringer Manual [Boehringer, Cell Death Detection ELISA plus, Cat No. 1 920 685]. Sample preparations: 96 well plates were spun down at 1 krpm for 10 minutes (200 g); the supernatant was removed by fast inversion, placing the plate upside down on a paper towel to remove residual liquid. To each well, 200  $\mu$ l of 1 $\times$ Lysis buffer was added and incubation allowed at room temperature for 30 minutes without shaking. The plates were spun down for 10 minutes at 1 krpm, and 20  $\mu$ l of the lysate (cytoplasmic fraction) was transferred into streptavidin coated MTP. 80  $\mu$ l of immunoreagent mix was added to the 20  $\mu$ l lysate in each well. The MTP was covered with adhesive foil and incubated at room temperature for 2 hours by placing it on an orbital shaker (200 rpm). After two hours, the supernatant was removed by suction and the wells rinsed three times with 250  $\mu$ l of 1 $\times$ incubation buffer per well (removed by suction). Substrate solution was added (100  $\mu$ l) into each well and incubated on an orbital shaker at room temperature at 250 rpm until color development was sufficient for a photometric analysis (approx. after 10-20 minutes). A 96 well reader was used to read the plates at 405 nm, reference wavelength, 492 nm. The levels obtained for PIN 32 (control buffer) was set to 100%. Samples with levels >130% were considered positive for induction of apoptosis.

[1720] The following PRO polypeptides tested positive in this assay: PRO235.

#### Example 81

[1721] Human Venous Endothelial Cell Calcium Flux Assay (Assay 68)

[1722] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to stimulate calcium flux in human umbilical vein endothelial cells (HUVEC, Cell Systems). Calcium influx is a well documented response upon binding of certain ligands to their receptors. A test compound that results in a positive response in the present calcium influx assay can be said to bind to a specific receptor and activate a biological signaling pathway in human endothelial cells. This could ultimately lead, for example, to endothelial cell division, inhibition of endothelial cell proliferation, endothelial tube formation, cell migration, apoptosis, etc.

[1723] Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50:50 without glycine, 1% glutamine, 10 mM Hepes, 10% FBS, 10 ng/ml bFGF), were plated on 96-well microtiter ViewPlates-96 (Packard Instrument Company Part #6005182) microtiter plates at a cell density of  $2 \times 10^4$  cells/well. The day after plating, the cells were washed three times with buffer (HBSS plus 10 mM Hepes), leaving 100  $\mu$ l/well. Then 100  $\mu$ l/well of 8  $\mu$ M Fluo-3 (2 $\times$ ) was added. The cells were incubated for 1.5 hours at 37° C/5% CO<sub>2</sub>. After incubation, the cells were then washed 3 $\times$  with buffer (described above) leaving 100  $\mu$ l/well. Test samples of the PRO polypeptides were prepared on different 96-well plates at 5 $\times$ concentration in buffer. The positive control corresponded to 50  $\mu$ M ionomycin (5 $\times$ ); the negative control corresponded to Protein 32. Cell plate and sample plates were run on a FLIPR (Molecular Devices) machine. The FLIPR machine added 25  $\mu$ l of test sample to the cells, and readings were taken every second for one minute, then every 3 seconds for the next three minutes.

[1724] The fluorescence change from baseline to the maximum rise of the curve ( $\Delta$  change) was calculated, and replicates averaged. The rate of fluorescence increase was monitored, and only those samples which had a  $\Delta$  change greater than 1000 and a rise within 60 seconds, were considered positive.

[1725] The following PRO polypeptides tested positive in the present assay: PRO245.

#### Example 82

[1726] Fibroblast (BHK-21) Proliferation (Assay 98)

[1727] This assay shows that certain PRO polypeptides of the invention act to induce proliferation of mammalian fibroblast cells in culture and, therefore, function as useful growth factors in mammalian systems. The assay is performed as follows. BHK-21 fibroblast cells plated in standard growth medium at 2500 cells/well in a total volume of 100  $\mu$ l. The PRO polypeptide,  $\beta$ -FGF (positive control) or nothing (negative control) are then added to the wells in the presence of 1  $\mu$ g/ml of heparin for a total final volume of 200  $\mu$ l. The cells are then incubated at 37° C. for 6 to 7 days. After incubation, the media is removed, the cells are washed with PBS and then an acid phosphatase substrate reaction mixture (100  $\mu$ l/well) is added. The cells are then incubated at 37° C. for 2 hours. 10  $\mu$ l per well of 1N NaOH is then added to stop the acid phosphatase reaction. The plates are then read at OD 405 nm. A positive in the assay is acid phosphatase activity which is at least 50% above the negative control.

[1728] The following PRO polypeptide tested positive in this assay: PRO258.

#### Example 83

[1729] Inhibition of Heart Adult Hypertrophy (Assay 42)

[1730] This assay is designed to measure the inhibition of heart adult hypertrophy. PRO polypeptides testing positive in this assay may find use in the therapeutic treatment of cardiac disorders associated with cardiac hypertrophy.

[1731] Ventricular myocytes are freshly isolated from adult (250 g) Harlan Sprague Dawley rats and the cells are plated at 2000/well in 180  $\mu$ l volume. On day two, test samples (20  $\mu$ l) containing the test PRO polypeptide are added. On day five, the cells are fixed and then stained. An increase in ANP message can also be measured by PCR from cells after a few hours. Results are based on a visual score of cell size: 0=no inhibition, -1=small inhibition, -2=large inhibition. A score of less than 0 is considered positive. Activity reference corresponds to phenylephrin (PE) at 0.1 mM, as a positive control. Assay media included: M199 (modified)-glutamine free, NaHCO<sub>3</sub>, phenol red, supplemented with 100 nM insulin, 0.2% BSA, 5 mM creatine, 2 mM L-carnitine, 5 mM taurine, 100 U/ml penicillin G, 100  $\mu$ g/ml streptomycin (CCT medium). Only inner 60 wells are used in 96 well plates. Of these, 6 wells are reserved for negative and positive (PE) controls.

[1732] The following PRO polypeptides provided a score of less than 0 in the above assay: PRO269.

#### Example 84

[1733] Induction of c-fos in Endothelial Cells (Assay 34)

[1734] This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in endothelial

cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of conditions or disorders where angiogenesis would be beneficial including, for example, wound healing, and the like (as would agonists of these PRO polypeptides). Antagonists of the PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of cancerous tumors.

[1735] Human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in growth media (50% Ham's F12 w/o GHT: low glucose, and 50% DMEM without glycine: with NaHCO<sub>3</sub>, 1% glutamine, 10 mM HEPES, 10% FBS, 10 ng/ml bFGF) were plated on 96-well microtiter plates at a cell density of  $1 \times 10^4$  cells/well. The day after plating, the cells were starved by removing the growth media and treating the cells with 100  $\mu$ l/well test samples and controls (positive control=growth media; negative control=Protein 32 buffer=10 mM HEPES, 140 mM NaCl, 4% (w/v) mannitol, pH 6.8). The cells were incubated for 30 minutes at 37° C., in 5% CO<sub>2</sub>. The samples were removed, and the first part of the bDNA kit protocol (Chiron Diagnostics, cat. #6005-037) was followed, where each capitalized reagent/buffer listed below was available from the kit.

[1736] Briefly, the amounts of the TM Lysis Buffer and Probes needed for the tests were calculated based on information provided by the manufacturer. The appropriate amounts of thawed Probes were added to the TM Lysis Buffer. The Capture Hybridization Buffer was warmed to room temperature. The bDNA strips were set up in the metal strip holders, and 100  $\mu$ l of Capture Hybridization Buffer was added to each b-DNA well needed, followed by incubation for at least 30 minutes. The test plates with the cells were removed from the incubator, and the media was gently removed using the vacuum manifold. 100  $\mu$ l of Lysis Hybridization Buffer with Probes were quickly pipetted into each well of the microtiter plates. The plates were then incubated at 55° C. for 15 minutes. Upon removal from the incubator, the plates were placed on the vortex mixer with the microtiter adapter head and vortexed on the #2 setting for one minute. 80  $\mu$ l of the lysate was removed and added to the bDNA wells containing the Capture Hybridization Buffer, and pipetted up and down to mix. The plates were incubated at 53° C. for at least 16 hours.

[1737] On the next day, the second part of the bDNA kit protocol was followed. Specifically, the plates were removed from the incubator and placed on the bench to cool for 10 minutes. The volumes of additions needed were calculated based upon information provided by the manufacturer. An Amplifier Working Solution was prepared by making a 1:100 dilution of the Amplifier Concentrate (20 fm/ $\mu$ l) in AL Hybridization Buffer. The hybridization mixture was removed from the plates and washed twice with Wash A. 50  $\mu$ l of Amplifier Working Solution was added to each well and the wells were incubated at 53° C. for 30 minutes. The plates were then removed from the incubator and allowed to cool for 10 minutes. The Label Probe Working Solution was prepared by making a 1:100 dilution of Label Concentrate (40 pmoles/ $\mu$ l) in AL Hybridization Buffer. After the 10-minute cool-down period, the amplifier hybridization mixture was removed and the plates were washed twice with Wash A. 50  $\mu$ l of Label Probe Working Solution was added to each well and the wells were incubated at 53° C. for 15 minutes. After cooling for 10 minutes, the Substrate was

warmed to room temperature. Upon addition of 3  $\mu$ l of Substrate Enhancer to each ml of Substrate needed for the assay, the plates were allowed to cool for 10 minutes, the label hybridization mixture was removed, and the plates were washed twice with Wash A and three times with Wash D. 50  $\mu$ l of the Substrate Solution with Enhancer was added to each well. The plates were incubated for 30 minutes at 37° C. and RLU was read in an appropriate luminometer.

[1738] The replicates were averaged and the coefficient of variation was determined. The measure of activity of the fold increase over the negative control (Protein 32/HEPES buffer described above) value was indicated by chemiluminescence units (RLU). The results are considered positive if the PRO polypeptide exhibits at least a two-fold value over the negative buffer control. Negative control=1.00 RLU at 1.00% dilution. Positive control=8.39 RLU at 1.00% dilution.

[1739] The following PRO polypeptides tested positive in this assay: PRO287.

#### Example 85

[1740] Guinea Pig Vascular Leak (Assays 32 and 51)

[1741] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce vascular permeability. Polypeptides testing positive in this assay are expected to be useful for the therapeutic treatment of conditions which would benefit from enhanced vascular permeability including, for example, conditions which may benefit from enhanced local immune system cell infiltration.

[1742] Hairless guinea pigs weighing 350 grams or more were anesthetized with Ketamine (75-80 mg/kg) and 5 mg/kg Xylazine intramuscularly. Test samples containing the PRO polypeptide or a physiological buffer without the test polypeptide are injected into skin on the back of the test animals with 100  $\mu$ l per injection site intradermally. There were approximately 16-24 injection sites per animal. One ml of Evans blue dye (1% in PBS) is then injected intracardially. Skin vascular permeability responses to the compounds (i.e., blemishes at the injection sites of injection) are visually scored by measuring the diameter (in mm) of blue-colored leaks from the site of injection at 1 and 6 hours post administration of the test materials. The mm diameter of blueness at the site of injection is observed and recorded as well as the severity of the vascular leakage. Blemishes of at least 5 mm in diameter are considered positive for the assay when testing purified proteins, being indicative of the ability to induce vascular leakage or permeability. A response greater than 7 mm diameter is considered positive for conditioned media samples. Human VEGF at 0.1  $\mu$ g/100  $\mu$ l is used as a positive control, inducing a response of 15-23 mm diameter.

[1743] The following PRO polypeptides tested positive in this assay: PRO302 and PRO533.

#### Example 86

[1744] Detection of Endothelial Cell Apoptosis (FACS) (Assay 96)

[1745] The ability of PRO polypeptides of the present invention to induce apoptosis in endothelial cells was tested

in human venous umbilical vein endothelial cells (HUVEC, Cell Systems) in gelatinized T175 flasks using HUVEC cells below passage 10. PRO polypeptides testing positive in this assay are expected to be useful for therapeutically treating conditions where apoptosis of endothelial cells would be beneficial including, for example, the therapeutic treatment of tumors.

[1746] On day one, the cells were split [420,000 cells per gelatinized 6 cm dishes—( $11 \times 10^3$  cells/cm<sup>2</sup> Falcon, Primaria)] and grown in media containing serum (CS-C, Cell System) overnight or for 16 hours to 24 hours.

[1747] On day 2, the cells were washed 1x with 5 ml PBS; 3 ml of 0% serum medium was added with VEGF (100 ng/ml); and 30  $\mu$ l of the PRO test compound (final dilution 1%) or 0% serum medium (negative control) was added. The mixtures were incubated for 48 hours before harvesting.

[1748] The cells were then harvested for FACS analysis. The medium was aspirated and the cells washed once with PBS. 5 ml of 1x trypsin was added to the cells in a T-175 flask, and the cells were allowed to stand until they were released from the plate (about 5-10 minutes). Trypsinization was stopped by adding 5 ml of growth media. The cells were spun at 1000 rpm for 5 minutes at 4° C. The media was aspirated and the cells were resuspended in 10 ml of 10% serum complemented medium (Cell Systems), 5  $\mu$ l of Annexin-FITC (BioVison) added and chilled tubes were submitted for FACS. A positive result was determined to be enhanced apoptosis in the PRO polypeptide treated samples as compared to the negative control.

[1749] The following PRO polypeptides tested positive in this assay: PRO331.

#### Example 87

[1750] Induction of c-fos in Cortical Neurons (Assay 83)

[1751] This assay is designed to determine whether PRO polypeptides show the ability to induce c-fos in cortical neurons. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of nervous system disorders and injuries where neuronal proliferation would be beneficial.

[1752] Cortical neurons are dissociated and plated in growth medium at 10,000 cells per well in 96 well plates. After approximately 2 cellular divisions, the cells are treated for 30 minutes with the PRO polypeptide or nothing (negative control). The cells are then fixed for 5 minutes with cold methanol and stained with an antibody directed against phosphorylated CREB. mRNA levels are then calculated using chemiluminescence. A positive in the assay is any factor that results in at least a 2-fold increase in c-fos message as compared to the negative controls.

[1753] The following PRO polypeptides tested positive in this assay: PRO229 and PRO269.

#### Example 88

[1754] Stimulation of Endothelial Tube Formation (Assay 85)

[1755] This assay is designed to determine whether PRO polypeptides show the ability to promote endothelial vacuole and lumen formation in the absence of exogenous

growth factors. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where endothelial vacuole and/or lumen formation would be beneficial including, for example, where the stimulation of pinocytosis, ion pumping, vascular permeability and/or junctional formation would be beneficial.

[1756] HUVEC cells (passage <8 from primary) are mixed with type I rat tail collagen (final concentration 2.6 mg/ml) at a density of  $6 \times 10^5$  cells per ml and plated at 50  $\mu$ l per well of M199 culture media supplemented with 1% FBS and 1  $\mu$ M 6-FAM-FITC dye to stain the vacuoles while they are forming and in the presence of the PRO polypeptide. The cells are then incubated at 37° C./5% CO<sub>2</sub> for 48 hours, fixed with 3.7% formalin at room temperature for 10 minutes, washed 5 times with M199 medium and then stained with Rh-Phalloidin at 4° C. overnight followed by nuclear staining with 4  $\mu$ M DAPI. A positive result in the assay is when vacuoles are present in greater than 50% of the cells.

[1757] The following PRO polypeptides tested positive in this assay: PRO230.

#### Example 89

[1758] Detection of Polypeptides that Affect Glucose and/or FFA Uptake in Skeletal Muscle (Assay 106)

[1759] This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by skeletal muscle cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by skeletal muscle would be beneficial including, for example, diabetes or hyper- or hypo-insulinemia.

[1760] In a 96 well format, PRO polypeptides to be assayed are added to primary rat differentiated skeletal muscle, and allowed to incubate overnight. Then fresh media with the PRO polypeptide and +/-insulin are added to the wells. The sample media is then monitored to determine glucose and FFA uptake by the skeletal muscle cells. The insulin will stimulate glucose and FFA uptake by the skeletal muscle, and insulin in media without the PRO polypeptide is used as a positive control, and a limit for scoring. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

[1761] The following PRO polypeptides tested positive as either stimulators or inhibitors of glucose and/or FFA uptake in this assay: PRO187, PRO211, PRO221, PRO222, PRO224, PRO230, PRO239, PRO231, PRO245, PRO247, PRO258, PRO269, PRO328 and PRO533.

#### Example 90

[1762] Rod Photoreceptor Cell Survival Assay (Assay 46)

[1763] This assay shows that certain polypeptides of the invention act to enhance the survival/proliferation of rod photoreceptor cells and, therefore, are useful for the therapeutic treatment of retinal disorders or injuries including, for example, treating sight loss in mammals due to retinitis pigmentosum, AMD, etc.

[1764] Sprague Dawley rat pups (postnatal day 7, mixed population: glia and retinal neural cell types) are killed by decapitation following CO<sub>2</sub> anesthesia and the eyes removed under sterile conditions. The neural retina is dissected away from the pigment epithelium and other ocular tissue and then dissociated into a single cell suspension using 0.25% trypsin in Ca<sup>2+</sup>, Mg<sup>2+</sup>-free PBS. The retinas are incubated at 37° C. in this solution for 7-10 minutes after which the trypsin is inactivated by adding 1 ml soybean trypsin inhibitor. The cells are plated at a density of approximately 10,000 cells/ml into 96 well plates in DMEM/F12 supplemented with N<sub>2</sub>. Cells for all experiments are grown at 37° C. in a water saturated atmosphere of 5% CO<sub>2</sub>. After 7-10 days in culture, the cells are stained using calcein AM or CellTracker Green CMFDA and then fixed using 4% paraformaldehyde. Rho 4D2 (ascities or IgG 1:100) monoclonal antibody directed towards the visual pigment rhodopsin is used to detect rod photoreceptor cells by indirect immunofluorescence. The results are calculated as % survival: total number of calcein—rhodopsin positive cells at 7-10 days in culture, divided by the total number of rhodopsin positive cells at time 7-10 days in culture. The total cells (fluorescent) are quantified at 20x objective magnification using a CCD camera and NIH image software for MacIntosh. Fields in the well are chosen at random.

[1765] The following polypeptides tested positive in this assay: PRO245.

#### Example 91

[1766] In vitro Antitumor Assay (Assay 161)

[1767] The antiproliferative activity of various PRO polypeptides was determined in the investigational, disease-oriented in vitro anti-cancer drug discovery assay of the National Cancer Institute (NCI), using a sulforhodamine B (SRB) dye binding assay essentially as described by Skehan et al., *J. Natl. Cancer Inst.* 82:1107-1112 (1990). The 60 tumor cell lines employed in this study ("the NCI panel"), as well as conditions for their maintenance and culture in vitro have been described by Monks et al., *J. Natl. Cancer Inst.* 83:757-766 (1991). The purpose of this screen is to initially evaluate the cytotoxic and/or cytostatic activity of the test compounds against different types of tumors (Monks et al., supra; Boyd, *Cancer: Princ. Pract. Oncol. Update* 3(10):1-12 [1989]).

[1768] Cells from approximately 60 human tumor cell lines were harvested with trypsin/EDTA (Gibco), washed once, resuspended in IMEM and their viability was determined. The cell suspensions were added by pipet (100 μL volume) into separate 96-well microtiter plates. The cell density for the 6-day incubation was less than for the 2-day incubation to prevent overgrowth. Inoculates were allowed a preincubation period of 24 hours at 37° C. for stabilization. Dilutions at twice the intended test concentration were added at time zero in 100 μL aliquots to the microtiter plate wells (1:2 dilution). Test compounds were evaluated at five half-log dilutions (1000 to 100,000-fold). Incubations took place for two days and six days in a 5% CO<sub>2</sub> atmosphere and 100% humidity.

[1769] After incubation, the medium was removed and the cells were fixed in 0.1 ml of 10% trichloroacetic acid at 40° C. The plates were rinsed five times with deionized water, dried, stained for 30 minutes with 0.1 ml of 0.4% sulfor-

hodamine B dye (Sigma) dissolved in 1% acetic acid, rinsed four times with 1% acetic acid to remove unbound dye, dried, and the stain was extracted for five minutes with 0.1 ml of 10 mM Tris base [tris(hydroxymethyl)aminomethane], pH 10.5. The absorbance (OD) of sulforhodamine B at 492 nm was measured using a computer-interfaced, 96-well microtiter plate reader.

[1770] A test sample is considered positive if it shows at least 50% growth inhibitory effect at one or more concentrations. PRO polypeptides testing positive in this assay are shown in Table 7, where the abbreviations are as follows:

TABLE 7

Test compound	Tumor Cell Line Type	Cell Line Designation
PRO211	NSCL	HOP62
PRO211	Leukemia	RPMI-8226
PRO211	Leukemia	HL-60 (TB)
PRO211	NSCL	NCI-H522
PRO211	CNS	SF-539
PRO211	Melanoma	LOX IMVI
PRO211	Breast	MDA-MB-435
PRO211	Leukemia	MOLT-4
PRO211	CNS	U251
PRO211	Breast	MCF7
PRO211	Leukemia	HT-60 (TB)
PRO211	Leukemia	MOLT-4
PRO211	NSCL	EKVX
PRO211	NSCL	NCI-H23
PRO211	NSCL	NCI-H322M
PRO211	NSCL	NCI-H460
PRO211	Colon	HCT-116
PRO211	Colon	HT29
PRO211	CNS	SF-268
PRO211	CNS	SF-295
PRO211	CNS	SNB-19
PRO211	CNS	U251
PRO211	Melanoma	LOX IMVI
PRO211	Melanoma	SK-MEL-5
PRO211	Melanoma	UACC-257
PRO211	Melanoma	UACC-62
PRO211	Ovarian	OVCAR-8
PRO211	Renal	RXF 393
PRO211	Breast	MCF7
PRO211	Breast	NCI/ADR-REHS 578T
PRO211	Breast	T-47D
PRO211	Leukemia	HL-60 (TB)
PRO211	Leukemia	SR
PRO211	NSCL	NCI-H23
PRO211	Colon	HCT-116
PRO211	Melanoma	LOX-IMVI
PRO211	Melanoma	SK-MEL-5
PRO211	Breast	T47D
PRO228	Leukemia	MOLT-4
PRO228	NSCL	EKVX
PRO228	Colon	KM12
PRO228	Melanoma	UACC-62
PRO228	Ovarian	OVCAR-3
PRO228	Renal	TK10
PRO228	Renal	SN12C
PRO228	Breast	MCF7
PRO228	Leukemia	CCRF-CEM
PRO228	Leukemia	HL-60 (TB)
PRO228	Colon	COLO 205
PRO228	Colon	HCT-15
PRO228	Colon	KM12
PRO228	CNS	SF-268
PRO228	CNS	SNB-75
PRO228	Melanoma	LOX-IMVI
PRO228	Melanoma	SK-MEL2
PRO228	Melanoma	UACC-257
PRO228	Ovarian	IGROV1
PRO228	Ovarian	OVCAR-4

TABLE 7-continued

Test compound	Tumor Cell Line Type	Cell Line Designation
PRO228	Ovarian	OVCAR-5
PRO228	Ovarian	OVCAR-8
PRO228	Renal	786-0
PRO228	Renal	CAKI-1
PRO228	Renal	RXF 393
PRO228	Renal	TK-10
PRO228	Renal	UO-31
PRO228	Prostate	PC-3
PRO228	Prostate	DU-145
PRO228	Breast	MCF7
PRO228	Breast	NCI/ADR-REHS 578T
PRO228	Breast	MDA-MB-435MDA-N
PRO228	Breast	T-47D
PRO219	Leukemia	SR
PRO219	NSCL	NCI-H5222
PRO219	Breast	MCF7
PRO219	Leukemia	K-562; RPMI-8226
PRO219	NSCL	HOP-62; NCI-H322M
PRO219	NSCL	NCI-H460
PRO219	Colon	HT29; KM12; HCT-116
PRO219	CNS	SF-539; U251
PRO219	Prostate	DU-145
PRO219	Breast	MDA-N
PRO219	Ovarian	IGROV1
PRO219	NSCL	NCI-H226
PRO219	Leukemia	MOLT-4
PRO219	NSCL	A549/ATCC; EK VX; NCI-H23
PRO219	Colon	HCC-2998
PRO219	CNS	SF-295; SNB-19
PRO219	Melanoma	SK-MEL-2; SK-MEL-5
PRO219	Melanoma	UACC-257; UACC-62
PRO219	Ovarian	OCAR-4; SK-OV-3
PRO219	Renal	786-0; ACHN; CAKI-1; SN12C
PRO219	Renal	TK-10; UO-31
PRO219	Breast	NCI/ADR-RES; BT-549; T-47D
PRO219	Breast	MDA-MB-435
PRO221	Leukemia	CCRF-CEM
PRO221	Leukemia	MOLT-4
PRO221	NSCL	HOP-62
PRO221	Breast	MDA-N
PRO221	Leukemia	RPMI-8226; SR
PRO221	NSCL	NCI-H460
PRO221	Colon	HCC-2998
PRO221	Ovarian	IGROV1
PRO221	Renal	TK-10
PRO221	Breast	MCF7
PRO221	Leukemia	K-562
PRO221	Breast	MDA-MB-435
PRO224	Ovarian	OVCAR-4
PRO224	Renal	RXF 393
PRO224	Prostate	DU-145
PRO224	NSCL	HOP-62; NCI-H322M
PRO224	Melanoma	LOX IMVI
PRO224	Ovarian	OVCAR-8
PRO224	Leukemia	SR
PRO224	NSCL	NCI-H460
PRO224	CNS	SF-295
PRO224	Leukemia	RPMI-8226
PRO224	Breast	BT-549
PRO224	Leukemia	CCRF-CEM; LH-60 (TB)
PRO224	Colon	HCT-116
PRO224	Breast	MDA-MB-435
PRO224	Leukemia	HL-60 (TB)
PRO224	Colon	HCC-2998
PRO224	Prostate	PC-3
PRO224	CNS	U251
PRO224	Colon	HCT-15
PRO224	CNS	SF-539
PRO224	Renal	ACHN
PRO328	Leukemia	RPMI-8226
PRO328	NSCL	A549/ATCC; EK VX; HOP-62
PRO328	NSCL	NCI-H23; NCI-H322M
PRO328	Colon	HCT-15; KM12

TABLE 7-continued

Test compound	Tumor Cell Line Type	Cell Line Designation
PRO328	CNS	SF-295; SF-539; SNB-19; U251
PRO328	Melanoma	M14; UACC-257; UCAA-62
PRO328	Renal	786-0; ACHN
PRO328	Breast	MCF7
PRO328	Leukemia	SR
PRO328	Colon	NCI-H23
PRO328	Melanoma	SK-MEL-5
PRO328	Prostate	DU-145
PRO328	Melanoma	LOX IMVI
PRO328	Breast	MDA-MB-435
PRO328	Ovarian	OVCAR-3
PRO328	Breast	T-47D
PRO301	NSCL	NCI-H322M
PRO301	Leukemia	MOLT-4; SR
PRO301	NSCL	A549/ATCC; EK VX;
PRO301	NSCL	NCI-H23; NCI-460; NCI-H226
PRO301	Colon	COLO 205; HCC-2998;
PRO301	Colon	HCT-15; KM12; HT29;
PRO301	Colon	HCT-116
PRO301	CNS	SF-268; SF-295; SNB-19
PRO301	Melanoma	MALME-3M; SK-MEL-2;
PRO301	Melanoma	SK-MEL-5; UACC-257
PRO301	Melanoma	UACC-62
PRO301	Ovarian	IGROV1; OVCAR-4
PRO301	Ovarian	OVCAR-5
PRO301	Ovarian	OVCAR-8; SK00V-3
PRO301	Renal	ACHN; CAKI-1; TK-10; UO-31
PRO301	Prostate	PC-3; DU-145
PRO301	Breast	NCI/ADR-RES; HS 578T
PRO301	Breast	MDA-MB-435; MDA-N; T-47D
PRO301	Melanoma	M14
PRO301	Leukemia	CCRF-CEM; HL-60(TB); K-562
PRO301	Leukemia	RPMI-8226
PRO301	Melanoma	LOX IMVI
PRO301	Renal	786-0; SN12C
PRO301	Breast	MCF7; MDA-MB-231/ATCC
PRO301	Breast	BT-549
PRO301	NSCL	HOP-62
PRO301	CNS	SF-539
PRO301	Ovarian	OVCAR-3
PRO326	NSCL	NCI-H322M
PRO326	CNS	SF295
PRO326	CNS	ST539
PRO326	CNS	U251

NSCL= non-small cell lung carcinoma

CNS= central nervous system

[1771] The results of these assays demonstrate that the positive testing PRO polypeptides are useful for inhibiting neoplastic growth in a number of different tumor cell types and may be used therapeutically therefor. Antibodies against these PRO polypeptides are useful for affinity purification of these useful polypeptides. Nucleic acids encoding these PRO polypeptides are useful for the recombinant preparation of these polypeptides.

#### Example 92

#### [1772] Gene Amplification

[1773] This example shows that certain PRO polypeptide-encoding genes are amplified in the genome of certain human lung, colon and/or breast cancers and/or cell lines. Amplification is associated with overexpression of the gene product, indicating that the polypeptides are useful targets for therapeutic intervention in certain cancers such as colon, lung, breast and other cancers and diagnostic determination of the presence of those cancers. Therapeutic agents may

take the form of antagonists of the PRO polypeptide, for example, murine-human chimeric, humanized or human antibodies against a PRO polypeptide.

[1774] The starting material for the screen was genomic DNA isolated from a variety cancers. The DNA is quantitated precisely, e.g., fluorometrically. As a negative control, DNA was isolated from the cells of ten normal healthy individuals which was pooled and used as assay controls for the gene copy in healthy individuals (not shown). The 5' nuclease assay (for example, TaqMan™) and real-time quantitative PCR (for example, ABI Prizm 7700 Sequence Detection System™ (Perlin Elmer, Applied Biosystems Division, Foster City, Calif.)), were used to find genes potentially amplified in certain cancers. The results were used to determine whether the DNA encoding the PRO polypeptide is over-represented in any of the primary lung or colon cancers or cancer cell lines or breast cancer cell lines that were screened. The primary lung cancers were obtained from individuals with tumors of the type and stage as indicated in Table 8. An explanation of the abbreviations used for the designation of the primary tumors listed in Table 8 and the primary tumors and cell lines referred to throughout this example are given below.

[1775] The results of the TaqMan™ are reported in delta (Δ) Ct units. One unit corresponds to 1 PCR cycle or approximately a 2-fold amplification relative to normal, two units corresponds to 4-fold, 3 units to 8-fold amplification and so on. Quantitation was obtained using primers and a TaqMan™ fluorescent probe derived from the PRO polypeptide-encoding gene. Regions of the PRO polypeptide-encoding gene which are most likely to contain unique nucleic acid sequences and which are least likely to have spliced out introns are preferred for the primer and probe derivation, e.g., 3'-untranslated regions. The sequences; for the primers and probes (forward, reverse and probe) used for the PRO polypeptide gene amplification analysis were as follows:

-continued

PRO187 (DNA27864-1155)  
 27864.tm.p:  
 5'-GCAGATTTTGAGGACAGCCACCTCCA-3' (SEQ ID NO:381)  
 27864.tm.f:  
 5'-GGCCTTGACAGACAACCGT-3' (SEQ ID NO:382)  
 27864.tm.r:  
 5'-CAGACTGAGGGAGATCCGAGA-3' (SEQ ID NO:383)  
 27864.tm.p2:  
 5'-CAGCTGCCTTCCCCAACCA-3' (SEQ ID NO:384)  
 27864.tm.f2:  
 5'-CATCAAGCGCCTCTACCA-3' (SEQ ID NO:385)  
 27864.tm.r2:  
 5'-CACAAACTCGAAGCTGCTTCTG-3' (SEQ ID NO:386)

PRO214 (DNA32286-1191):  
 32286.3utr-5:  
 5'-GGGCCATCACAGCTCCCT-3' (SEQ ID NO:387)  
 32286.3utr-3b:  
 5'-GGGATGTGGTGAACACAGAACA-3' (SEQ ID NO:388)  
 32286.3utr-probe:  
 5'-TGCCAGCTGCATGCTGCCAGTT-3' (SEQ ID NO:389)

PRO211 (DNA32292-1131):  
 32292.3utr-5:  
 5'-CAGAAGGATGTCCCGTGGAA-3' (SEQ ID NO:390)  
 32292.3utr-3:  
 5'-GCCGCTGTCCACTGCAG-3' (SEQ ID NO:391)  
 32292.3utr-probe.rc:  
 5'-GACGGCATCCTCAGGGCCACA-3' (SEQ ID NO:392)

PRO230 (DNA33223-1136):  
 33223.tm.p3:  
 5'-ATGTCCTCCATGCCCCACGCG-3' (SEQ ID NO:393)  
 33223.tm.f3:  
 5'-GAGTGGACATCGAGAGCTT-3' (SEQ ID NO:394)  
 33223.tm.r3:  
 5'-CCGAGCCTCAGTGATGA-3' (SEQ ID NO:395)  
 33223.3utr-5:  
 5'-GAAGAGCACAGCTGCAGATCC-3' (SEQ ID NO:396)  
 33223.3utr-3:  
 5'-GAGGTGCTCTGGCTTTGGTAGT-3' (SEQ ID NO:397)  
 33223.3utr-probe:  
 5'-CCTCTGGCGCCCCACTCAA-3' (SEQ ID NO:398)

PRO317 (DNA33461-1199):  
 33461.tm.f:  
 5'-CCAGGAGAGCTGGCGATG-3' (SEQ ID NO:399)  
 33461.tm.r:  
 5'-GCAATTCAGGGCTCACTAGAGA-3' (SEQ ID NO:400)  
 33461.tm.p:  
 5'-CACAGAGCATTGTCCATCAGCAGTTCAG-3' (SEQ ID NO:401)

PRO246 (DNA35639-1172):  
 35639.3utr-5:  
 5'-GGCAGAGACTTCCAGTCACTGA-3' (SEQ ID NO:402)  
 35639.3utr-3:  
 5'-GCCAAGGGTGGTGTAGATAGG-3' (SEQ ID NO:403)  
 35639.3utr-probe:  
 5'-CAGGCCCCCTTGATCTGTACCCCA-3' (SEQ ID NO:404)

PRO533 (DNA49435-1219):  
 49435.tm.f:  
 5'-GGGACGTGCTTCTACAAGAACAG-3' (SEQ ID NO:405)  
 49435.tm.r:  
 5'-CAGGCTTACAATGTTATGATCAGACA-3' (SEQ ID NO:406)  
 49435.tm.p:  
 5'-TATTCAGAGTTTTCCATTTGGCAGTGCAGTT-3' (SEQ ID NO:407)

PRO343 (DNA43318-1217):  
 43318.tm.f1  
 5'-TCTACATCAGCCTCTCTGCGC-3' (SEQ ID NO:408)  
 43318.tm.p1  
 5'-CGATCTTCTCCACCCAGGAGCGG-3' (SEQ ID NO:409)  
 43318.tm.r1  
 5'-GGAGCTGCACCCCTTGC-3' (SEQ ID NO:237)

PRO232 (DNA34435-1140):  
 34435.3utr-5:  
 5'-GCCAGGCCTCACATTCGT-3' (SEQ ID NO:410)  
 DNA34435.3utr-probe:  
 5'-CTCCCTGAATGGCAGCCTGAGCA-3' (SEQ ID NQ:411)  
 DNA34435.3utr-3:  
 5'-AGGTGTTTATTAAGGGCCTACGCT-3' (SEQ ID NQ:412)

PRO269 (DNA38260-1180):  
 38260.tm.f:  
 5'-CAGAGCAGAGGGTGCCTTG-3' (SEQ ID NO:413)  
 38260.tm.p:  
 5'-TGGCGGAGTCCCTCTTGGCT-3' (SEQ ID NO:414)  
 38260.tm.r:  
 5'-CCCTGTTTCCCTATGCATCACT-3' (SEQ ID NO:415)

PRO304 (DNA39520-1217):  
 39520.tm.f:  
 5'-TCAACCCCTGACCCCTTCCCTA-3' (SEQ ID NO:416)  
 39520.tm.p:  
 5'-GGCAGGGGACAAGCCATCTCTCCT-3' (SEQ ID NO:417)  
 39520.tm.r:  
 5'-GGGACTGAAGTCCAGCTTC-3' (SEQ ID NO:418)

PRO339 (DNA43466-1225):  
 43466.tm.f1:  
 5'-GGGCCCTAACCTCATTACCTTT-3' (SEQ ID NO:419)

-continued

43466.tm.p1:  
 5'-TGTCCTGCCTCAGCCCCAGGAAGG-3' (SEQ ID NO:420)  
 43466.tm.r1:  
 5'-TCTGTCCACCATCTTGCCTTG-3' (SEQ ID NO:421)

[1776] The 5' nuclease assay reaction is a fluorescent PCR-based technique which makes use of the 5' exonuclease activity of Taq DNA polymerase enzyme to monitor amplification in real time. Two oligonucleotide primers (forward [.f] and reverse [.r]) are used to generate an amplicon typical of a PCR reaction. A third oligonucleotide, or probe (.p), is designed to detect nucleotide sequence located between the two PCR primers. The probe is non-extendible by Taq DNA polymerase enzyme, and is labeled with a reporter fluorescent dye and a quencher fluorescent dye. Any laser-induced emission from the reporter dye is quenched by the quenching dye when the two dyes are located close together as they are on the probe. During the amplification reaction, the Taq DNA polymerase enzyme cleaves the probe in a template-dependent manner. The resultant probe fragments disassociate in solution, and signal from the released reporter dye is free from the quenching effect of the second fluorophore. One molecule of reporter dye is liberated for each new

molecule synthesized, and detection of the unquenched reporter dye provides the basis for quantitative interpretation of the data.

[1777] The 5' nuclease procedure is run on a realtime quantitative PCR device such as the ABI Prism 7700TM Sequence Detection. The system consists of a thermocycler, laser, charge-coupled device (CCD) camera and computer. The system amplifies samples in a 96-well format on a thermocycler. During amplification, laser-induced fluorescent signal is collected in real-time through fiber optics cables for all 96 wells, and detected at the CCD. The system includes software for running the instrument and for analyzing the data.

[1778] 5' Nuclease assay data are initially expressed as Ct, or the threshold cycle. This is defined as the cycle at which the reporter signal accumulates above the background level of fluorescence. The ΔCt values are used as quantitative measurement of the relative number of starting copies of a particular target sequence in a nucleic acid sample when comparing cancer DNA results to normal human DNA results.

[1779] Table 8 describes the stage, T stage and N stage of various primary tumors which were used to screen the PRO polypeptide compounds of the invention.

TABLE 8

Primary Tumor Stage	Stage	Other Stage	Dukes Stage	T Stage	N Stage
Human lung tumor AdenoCa (SRCC724) [LT1]	IIA			T1	N1
Human lung tumor SqCCa (SRCC725) [LT1a]	IIB			T3	N0
Human lung tumor AdenoCa (SRCC726) [LT2]	IB			T2	N0
Human lung tumor AdenoCa (SRCC727) [LT3]	IIIA			T1	N2
Human lung tumor AdenoCa (SRCC728) [LT4]	IB			T2	N0
Human lung tumor SqCCa (SRCC729) [LT6]	IB			T2	N0
Human lung tumor Aden/SqCCa (SRCC730) [LT7]	IA			T1	N0
Human lung tumor AdenoCa (SRCC731) [LT9]	IB			T2	N0
Human lung tumor SqCCa (SRCC732) [LT10]	IIB			T2	N1
Human lung tumor SqCCa (SRCC733) [LT11]	IIA			T1	N1
Human lung tumor AdenoCa (SRCC734) [LT12]	IV			T2	N0
Human lung tumor AdenoSqCCa (SRCC735)[LT13]	IB			T2	N0
Human lung tumor SqCCa (SRCC736) [LT15]	IB			T2	N0
Human lung tumor SqCCa (SRCC737) [LT16]	IB			T2	N0
Human lung tumor SqCCa (SRCC738) [LT17]	IIB			T2	N1
Human lung tumor SqCCa (SRCC739) [LT18]	IB			T2	N0
Human lung tumor SqCCa (SRCC740) [LT19]	IB			T2	N0
Human lung tumor LCCa (SRCC741) [LT21]	IIB			T3	N1
Human lung AdenoCa (SRCC811) [LT221]	IA			T1	N0
Human colon AdenoCa (SRCC742) [CT2]		M1	D	pT4	N0
Human colon AdenoCa (SRCC743) [CT3]			B	pT3	N0
Human colon AdenoCa (SRCC744) [CT8]			B	T3	N0
Human colon AdenoCa (SRCC745) [CT10]			A	pT2	N0
Human colon AdenoCa (SRCC746) [CT12]		MO, R1	B	T3	N0
Human colon AdenoCa (SRCC747) [CT14]		pMO, RO	B	pT3	pN0
Human colon AdenoCa (SRCC748) [CT15]		M1, R2	D	T4	N2
Human colon AdenoCa (SRCC749) [CT16]		pMO	B	pT3	pN0
Human colon AdenoCa (SRCC750) [CT17]			C1	pT3	pN1
Human colon AdenoCa (SRCC751) [CT1]		MO, R1	B	pT3	N0
Human colon AdenoCa (SRCC752) [CT4]			B	p33	M0
Human colon AdenoCa (SRCC753) [CT5]		G2	C1	p33	pN0
Human colon AdenoCa (SRCC754) [CT6]		pMO, RO	B	pT3	pN0
Human colon AdenoCa (SRCC755) [CT7]		G1	A	pT2	pN0
Human colon AdenoCa (SRCC756) [CT9]		G3	D	pT4	pN2
Human colon AdenoCa (SRCC757) [CT11]			B	T3	N0
Human colon AdenoCa (SRCC758) [CT18]		MO, RO	B	pT3	pN0

**[1780]** DNA Preparation

**[1781]** DNA was prepared from cultured cell lines, primary tumors, normal human blood. The isolation was performed using purification kit, buffer set and protease and all from Qiagen, according to the manufacturer's instructions and the description below.

**[1782]** Cell culture lysis:

**[1783]** Cells were washed and trypsinized at a concentration of  $7.5 \times 10^8$  per tip and pelleted by centrifuging at 1000 rpm for 5 minutes at 4° C., followed by washing again with 1/2 volume of PBS recentrifugation. The pellets were washed a third time, the suspended cells collected and washed 2x with PBS. The cells were then suspended into 10 ml PBS. Buffer C1 was equilibrated at 4° C. Qiagen protease #19155 was diluted into 6.25 ml cold ddH<sub>2</sub>O to a final concentration of 20 mg/ml and equilibrated at 4° C. 10 ml of G2 Buffer was prepared by diluting Qiagen RNase A stock (100 mg/ml) to a final concentration of 200 µg/ml.

**[1784]** Buffer C1 (10 ml, 4° C.) and ddH<sub>2</sub>O (40 ml, 4° C.) were then added to the 10 ml of cell suspension, mixed by inverting and incubated on ice for 10 minutes. The cell nuclei were pelleted by centrifuging in a Beckman swinging bucket rotor at 2500 rpm at 4° C. for 15 minutes. The supernatant was discarded and the nuclei were suspended with a vortex into 2 ml Buffer C1 (at 4° C.) and 6 ml ddH<sub>2</sub>O, followed by a second 4° C. centrifugation at 2500 rpm for 15 minutes. The nuclei were then resuspended into the residual buffer using 200 µl per tip. G2 buffer (10 ml) was added to the suspended nuclei while gentle vortexing was applied. Upon completion of buffer addition, vigorous vortexing was applied for 30 seconds. Qiagen protease (200 µl, prepared as indicated above) was added and incubated at 50° C. for 60 minutes. The incubation and centrifugation was repeated until the lysates were clear (e.g., incubating additional 30-60 minutes, pelleting at 3000xg for 10 min., 4° C.).

**[1785]** Solid human tumor sample preparation and lysis:

**[1786]** Tumor samples were weighed and placed into 50 ml conical tubes and held on ice. Processing was limited to no more than 250 mg tissue per preparation (1 tip/preparation). The protease solution was freshly prepared by diluting into 6.25 ml cold ddH<sub>2</sub>O to a final concentration of 20 mg/ml and stored at 4° C. G2 buffer (20 ml) was prepared by diluting DNase A to a final concentration of 200 mg/ml (from 100 mg/ml stock). The tumor tissue was homogenated in 19 ml G2 buffer for 60 seconds using the large tip of the polytron in a laminar-flow TC hood in order to avoid inhalation of aerosols, and held at room temperature. Between samples, the polytron was cleaned by spinning at 2x30 seconds each in 2 L ddH<sub>2</sub>O, followed by G2 buffer (50 ml). If tissue was still present on the generator tip, the apparatus was disassembled and cleaned.

**[1787]** Qiagen protease (prepared as indicated above, 1.0 ml) was added, followed by vortexing and incubation at 50° C. for 3 hours. The incubation and centrifugation was repeated until the lysates were clear (e.g., incubating additional 30-60 minutes, pelleting at 3000xg for 10 min., 4° C.).

**[1788]** Human blood preparation and lysis:

**[1789]** Blood was drawn from healthy volunteers using standard infectious agent protocols and citrated into 10 ml samples per tip. Qiagen protease was freshly prepared by

dilution into 6.25 ml cold ddH<sub>2</sub>O to a final concentration of 20 mg/ml and stored at 4° C. G2 buffer was prepared by diluting RNase A to a final concentration of 200 µg/ml from 100 mg/ml stock. The blood (10 ml) was placed into a 50 ml conical tube and 10 ml C1 buffer and 30 ml ddH<sub>2</sub>O (both previously equilibrated to 4° C.) were added, and the components mixed by inverting and held on ice for 10 minutes. The nuclei were pelleted with a Beckman swinging bucket rotor at 2500 rpm, 4° C. for 15 minutes and the supernatant discarded. With a vortex, the nuclei were suspended into 2 ml C1 buffer (4° C.) and 6 ml ddH<sub>2</sub>O (4° C.). Vortexing was repeated until the pellet was white. The nuclei were then suspended into the residual buffer using a 200 µl tip. G2 buffer (10 ml) were added to the suspended nuclei while gently vortexing, followed by vigorous vortexing for 30 seconds. Qiagen protease was added (200 µl) and incubated at 50° C. for 60 minutes. The incubation and centrifugation was repeated until the lysates were clear (e.g., incubating additional 30-60 minutes, pelleting at 3000xg for 10 min., 4° C.).

**[1790]** Purification of cleared lysates:**[1791]** (1) Isolation of genomic DNA:

**[1792]** Genomic DNA was equilibrated (1 sample per maxi tip preparation) with 10 ml QBT buffer. QF elution buffer was equilibrated at 50° C. The samples were vortexed for 30 seconds, then loaded onto equilibrated tips and drained by gravity. The tips were washed with 2x15 ml QC buffer. The DNA was eluted into 30 ml silanized, autoclaved 30 ml Corex tubes with 15 ml QF buffer (50° C.). Isopropanol (10.5 ml) was added to each sample, the tubes covered with parafilm and mixed by repeated inversion until the DNA precipitated. Samples were pelleted by centrifugation in the SS-34 rotor at 15,000 rpm for 10 minutes at 4° C. The pellet location was marked, the supernatant discarded, and 10 ml 70% ethanol (4° C.) was added. Samples were pelleted again by centrifugation on the SS-34 rotor at 10,000 rpm for 10 minutes at 4° C. The pellet location was marked and the supernatant discarded. The tubes were then placed on their side in a drying rack and dried 10 minutes at 37° C., taking care not to overdry the samples.

**[1793]** After drying, the pellets were dissolved into 1.0 ml TE (pH 8.5) and placed at 50° C. for 1-2 hours. Samples were held overnight at 4° C. as dissolution continued. The DNA solution was then transferred to 1.5 ml tubes with a 26 gauge needle on a tuberculin syringe. The transfer was repeated 5x in order to shear the DNA. Samples were then placed at 50° C. for 1-2 hours.

**[1794]** (2) Quantitation of genomic DNA and preparation for gene amplification assay:

**[1795]** The DNA levels in each tube were quantified by standard A<sub>260</sub>, A<sub>280</sub> spectrophotometry on a 1:20 dilution (5 µl DNA+95 µl ddH<sub>2</sub>O) using the 0.1 ml quartz cuvettes in the Beckman DU640 spectrophotometer. A<sub>260</sub>/A<sub>280</sub> ratios were in the range of 1.8-1.9. Each DNA samples was then diluted further to approximately 200 ng/ml in TE (pH 8.5). If the original material was highly concentrated (about 700 ng/µl), the material was placed at 50° C. for several hours until resuspended.

**[1796]** Fluorometric DNA quantitation was then performed on the diluted material (20-600 ng/ml) using the manufacturer's guidelines as modified below. This was



accomplished by allowing a Hoeffler DyNA Quant 200 fluorometer to warm-up for about 15 minutes. The Hoechst dye working solution (#H33258, 10  $\mu$ l, prepared within 12 hours of use) was diluted into 100 ml 1 $\times$ TNE buffer. A 2 ml cuvette was filled with the fluorometer solution, placed into the machine, and the machine was zeroed. pGEM 3Zf(+) (2  $\mu$ l, lot #360851026) was added to 2 ml of fluorometer solution and calibrated at 200 units. An additional 2  $\mu$ l of pGEM 3Zf(+) DNA was then tested and the reading confirmed at 400 $\pm$ 10 units. Each sample was then read at least in triplicate. When 3 samples were found to be within 10% of each other, their average was taken and this value was used as the quantification value.

[1797] The fluorometrically determined concentration was then used to dilute each sample to 10 ng/ $\mu$ l in ddH<sub>2</sub>O. This was done simultaneously on all template samples for a single TaqMan plate assay, and with enough material to run

500-1000 assays. The samples were tested in triplicate with Taqman™ primers and probe both B-actin and GAPDH on a single plate with normal human DNA and no-template controls. The diluted samples were used provided that the CT value of normal human DNA subtracted from test DNA was  $\pm$ 1 Ct. The diluted, lot-qualified genomic DNA was stored in 1.0 ml aliquots at  $-80^{\circ}$  C. Aliquots which were subsequently to be used in the gene amplification assay were stored at  $4^{\circ}$  C. Each 1 ml aliquot is enough for 8-9 plates or 64 tests.

[1798] Gene amplification assay:

[1799] The PRO polypeptide compounds of the invention were screened in the following primary tumors and the resulting  $\Delta$ Ct values greater than or equal to 1.0 are reported in Table 9 below.

TABLE 9

ΔCt values in lung and colon primary tumors and cell line models

Primary Tumors or Cell lines	PRO187	PRO533	PRO214	PRO343	PRO211	PRO230	PRO246	PRO317	PRO232	PRO269	PRO304	PRO339
LT7								1.52		1.04		1.08
LT13	2.74 2.98 2.44		1.85 1.83	2.71 2.23	1.88 2.26	3.42 3.22 2.84 2.15 2.75 2.53 1.82	1.63 1.68	2.24 2.93	1.27	1.29	1.04	
LT3			1.57		1.97		1.06	1.86				1.17
LT4					1.17			1.18				
LT9					1.42			1.04		1.80		1.03
LT12	2.70 2.90 2.27		1.38 1.49	2.23 1.50	1.51 1.27	2.86 2.96 2.92 1.25 2.68 2.28 1.34 2.13 1.36 1.09	1.54 2.47	2.54 1.74	2.40	1.14	1.15	1.26
LT30	1.67						1.50	1.29				
LT21		1.02			1.26 1.18							
LT1- a								1.93				
LT6					1.96		1.07	2.57				
LT10		1.09	1.67	1.00	2.05	1.32	3.43	2.20	1.14	1.14	1.51	1.39
LT11			1.80		1.89	1.14	1.41	2.33				
						1.54		1.02				
LT15	3.75 3.92 3.49		1.77 1.58	3.62 1.30	2.44 2.16	4.32 4.47 3.64 2.94 3.56 3.32 2.68	2.11 1.56	2.06 2.76 1.63	1.86	1.36	1.34	
						2.04						
LT16	2.10	1.66		1.70	1.25	1.15		1.55			1.00	
						1.83		1.08				
LT17		1.32	1.93 1.87	1.15	1.85 2.30	1.26 1.39 1.30 1.33 1.30	2.68 1.69	2.29 2.03 1.10	1.35	1.42	1.68	1.63
LT18	4.05	1.67	2.09	1.17	2.42	4.05	1.91	2.51	1.04		1.15	
LT19	3.99		1.98	3.82	2.55	4.92	1.68	2.03	1.21	1.60		
						4.93	1.16					



**[1800]** Summary

**[1801]** Because amplification of the various DNA's as described above occurs in various tumors, it is likely associated with tumor formation and/or growth. As a result, antagonists (e.g., antibodies) directed against these polypeptides would be expected to be useful in cancer therapy.

## Example 94

**[1802]** Detection of PRO Polypeptides that Affect Glucose or FFA Uptake by Primary Rat Adipocytes (Assay 94)

**[1803]** This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by adipocyte cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by adipocytes would be beneficial including, for example, obesity, diabetes or hyper- or hypo-insulinemia.

**[1804]** In a 96 well format, PRO polypeptides to be assayed are added to primary rat adipocytes, and allowed to incubate overnight. Samples are taken at 4 and 16 hours and assayed for glycerol, glucose and FFA uptake. After the 16 hour incubation, insulin is added to the media and allowed to incubate for 4 hours. At this time, a sample is taken and glycerol, glucose and FFA uptake is measured. Media containing insulin without the PRO polypeptide is used as a positive reference control. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

**[1805]** The following PRO polypeptides tested positive as stimulators of glucose and/or FFA uptake in this assay: PRO221, PRO235, PRO245, PRO295, PRO301 and PRO332.

**[1806]** The following PRO polypeptides tested positive as inhibitors of glucose and/or FFA uptake in this assay: PRO214, PRO219, PRO228, PRO222, PRO231 and PRO265.

## Example 95

**[1807]** Chondrocyte Re-differentiation Assay (Assay 110)

**[1808]** This assay shows that certain polypeptides of the invention act to induce redifferentiation of chondrocytes, therefore, are expected to be useful for the treatment of various bone and/or cartilage disorders such as, for example, sports injuries and arthritis. The assay is performed as follows. Porcine chondrocytes are isolated by overnight collagenase digestion of articular cartilage of metacarpophalangeal joints of 4-6 month old female pigs. The isolated cells are then seeded at 25,000 cells/cm<sup>2</sup> in Ham F-12 containing 10% FBS and 4  $\mu$ g/ml gentamycin. The culture media is changed every third day and the cells are then seeded in 96 well plates at 5,000 cells/well in 100  $\mu$ l of the same media without serum and 100  $\mu$ l of the test PRO polypeptide, 5 nM staurosporin (positive control) or medium alone (negative control) is added to give a final volume of 200  $\mu$ l/well. After 5 days of incubation at 37° C., a picture of each well is taken and the differentiation state of the chondrocytes is determined. A positive result in the assay occurs when the redifferentiation of the chondrocytes is determined to be more similar to the positive control than the negative control.

**[1809]** The following polypeptide tested positive in this assay: PRO214, PRO219, PRO229, PRO222, PRO224, PRO230, PRO257, PRO272 and PRO301.

## Example 96

**[1810]** Fetal Hemoglobin Induction in an Erythroblastic Cell Line (Assay 107)

**[1811]** This assay is useful for screening PRO polypeptides for the ability to induce the switch from adult hemoglobin to fetal hemoglobin in an erythroblastic cell line. Molecules testing positive in this assay are expected to be useful for therapeutically treating various mammalian hemoglobin-associated disorders such as the various thalassemias. The assay is performed as follows. Erythroblastic cells, are plated in standard growth medium at 1000 cells/well in a 96 well format. PRO polypeptides are added to the growth medium at a concentration of 0.2% or 2% and the cells are incubated for 5 days at 37° C. As a positive control, cells are treated with 100  $\mu$ M hemin and as a negative control, the cells are untreated. After 5 days, cell lysates are prepared and analyzed for the expression of gamma globin (a fetal marker). A positive in the assay is a gamma globin level at least 2-fold above the negative control.

**[1812]** The following polypeptides tested positive in this assay: PRO221 and PRO245.

## Example 97

**[1813]** Mouse Kidney Mesangial Cell Proliferation Assay (Assay 92)

**[1814]** This assay shows that certain polypeptides of the invention act to induce proliferation of mammalian kidney mesangial cells and, therefore, are useful for treating kidney disorders associated with decreased mesangial cell function such as Berger disease or other nephropathies associated with Schönlein-Henoch purpura, celiac disease, dermatitis herpetiformis or Crohn disease. The assay is performed as follows. On day one, mouse kidney mesangial cells are plated on a 96 well plate in growth media (3:1 mixture of Dulbecco's modified Eagle's medium and Ham's F12 medium, 95% fetal bovine serum, 5% supplemented with 14 mM HEPES) and grown overnight. On day 2, PRO polypeptides are diluted at 2 concentrations (1% and 0.1%) in serum-free medium and added to the cells. Control samples are serum-free medium alone. On day 4, 20  $\mu$ l of the Cell Titer 96 Aqueous one solution reagent (Progema) was added to each well and the colorimetric reaction was allowed to proceed for 2 hours. The absorbance (OD) is then measured at 490 nm. A positive in the assay is anything that gives an absorbance reading which is at least 15% above the control reading.

**[1815]** The following polypeptide tested positive in this assay: PRO227.

## Example 98

**[1816]** Proliferation of Rat Utricular Supporting Cells (Assay 54)

**[1817]** This assay shows that certain polypeptides of the invention act as potent mitogens for inner ear supporting cells which are auditory hair cell progenitors and, therefore, are useful for inducing the regeneration of auditory hair cells and treating hearing loss in mammals. The assay is performed as follows. Rat UEC-4 utricular epithelial cells are aliquoted into 96 well plates with a density of 3000 cells/

well in 200  $\mu$ l of serum-containing medium at 33° C. The cells are cultured overnight and are then switched to serum-free medium at 37° C. Various dilutions of PRO polypeptides (or nothing for a control) are then added to the cultures and the cells are incubated for 24 hours. After the 24 hour incubation, <sup>3</sup>-thymidine (1  $\mu$ Ci/well) is added and the cells are then cultured for an additional 24 hours. The cultures are then washed to remove unincorporated radiolabel, the cells harvested and Cpm per well determined. Cpm of at least 30% or greater in the PRO polypeptide treated cultures as compared to the control cultures is considered a positive in the assay.

[1818] The following polypeptides tested positive in this assay: PRO310 and PRO346.

#### Example 99

[1819] Chondrocyte Proliferation Assay (Assay 111)

[1820] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to induce the proliferation and/or redifferentiation of chondrocytes in culture. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of various bone and/or cartilage disorders such as, for example, sports injuries and arthritis.

[1821] Porcine chondrocytes are isolated by overnight collagenase digestion of articular cartilage of the metacarpophalangeal joint of 4-6 month old female pigs. The isolated cells are then seeded at 25,000 cells/cm<sup>2</sup> in Ham F-12 containing 10% FBS and 4  $\mu$ g/ml gentamycin. The culture media is changed every third day and the cells are reseeded to 25,000 cells/cm<sup>2</sup> every five days. On day 12, the cells are seeded in 96 well plates at 5,000 cells/well in 100  $\mu$ l of the same media without serum and 100  $\mu$ l of either serum-free medium (negative control), staurosporin (final concentration of 5 nM; positive control) or the test PRO polypeptide are added to give a final volume of 200  $\mu$ l/well. After 5 days at 37° C., 20  $\mu$ l of Alamar blue is added to each well and the plates are incubated for an additional 3 hours at 37° C. The fluorescence is then measured in each well (Ex:530 nm; Em:590 nm). The fluorescence of a plate containing 200  $\mu$ l of the serum-free medium is measured to obtain the background. A positive result in the assay is obtained when the fluorescence of the PRO polypeptide treated sample is more like that of the positive control than the negative control.

[1822] The following PRO polypeptides tested positive in this assay: PRO219, PRO222, PRO317, PRO257, PRO265, PRO287, PRO272 and PRO533.

#### Example 100

[1823] Inhibition of Heart Neonatal Hypertrophy Induced by LIF+ET-1 (Assay 74)

[1824] This assay is designed to determine whether PRO polypeptides of the present invention show the ability to inhibit neonatal heart hypertrophy induced by LIF and endothelin-1 (ET-1). A test compound that provides a positive response in the present assay would be useful for the therapeutic treatment of cardiac insufficiency diseases or disorders characterized or associated with an undesired hypertrophy of the cardiac muscle.

[1825] Cardiac myocytes from 1-day old Harlan Sprague Dawley rats (180  $\mu$ l at 7.5 $\times$ 10<sup>4</sup>/ml, serum <0.1, freshly isolated) are introduced on day 1 to 96-well plates previ-

ously coated with DMEM/F12+4% FCS. Test PRO polypeptide samples or growth medium alone (negative control) are then added directly to the wells on day 2 in 20  $\mu$ l volume. LIF+ET-1 are then added to the wells on day 3. The cells are stained after an additional 2 days in culture and are then scored visually the next day. A positive in the assay occurs when the PRO polypeptide treated myocytes are visually smaller on the average or less numerous than the untreated myocytes.

[1826] The following PRO polypeptides tested positive in this assay: PRO238.

#### Example 101

[1827] Tissue Expression Distribution

[1828] Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human adult and/or fetal tissue sources and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tissues tested. Knowledge of the expression pattern or the differential expression of the PRO polypeptide-encoding nucleic acid in various different human tissue types provides a diagnostic marker useful for tissue typing, with or without other tissue-specific markers, for determining the primary tissue source of a metastatic tumor, and the like. These assays provided the following results.

DNA Molecule	Tissues With Significant Expression	Tissues Lacking Significant Expression
DNA34436-1238	lung, placenta, brain	testis
DNA35557-1137	lung, kidney, brain	placenta
DNA35599-1168	kidney, brain	liver, placenta
DNA35668-1171	liver, lung, kidney	placenta, brain
DNA36992-1168	liver, lung, kidney, brain	placenta
DNA39423-1182	kidney, brain	liver
DNA40603-1232	liver	brain, kidney, lung
DNA40604-1187	liver	brain, kidney, lung
DNA41379-1236	lung, brain	liver
DNA33206-1165	heart, spleen, dendrocytes	substantia nigra, hippocampus, cartilage, prostate, HUVEC
DNA34431-1177	spleen, HUVEC, cartilage, heart, uterus	brain, colon tumor, prostate, THP-1 macrophages
DNA41225-1217	HUVEC, uterus, colon tumor, cartilage, prostate	spleen, brain, heart, IM-9 lymphoblasts

#### Example 102

[1829] In situ Hybridization

[1830] In situ hybridization is a powerful and versatile technique for the detection and localization of nucleic acid sequences within cell or tissue preparations. It may be useful, for example, to identify sites of gene expression, analyze the tissue distribution of transcription, identify and

localize viral infection, follow changes in specific mRNA synthesis and aid in chromosome mapping.

[1831] In situ hybridization was performed following an optimized version of the protocol by Lu and Gillett, *Cell Vision* 1:169-176 (1994), using PCR-generated <sup>33</sup>P-labeled riboprobes. Briefly, formalin-fixed, paraffin-embedded human tissues were sectioned, deparaffinized, deproteinized in proteinase K (20 g/ml) for 15 minutes at 37° C., and further processed for in situ hybridization as described by Lu and Gillett, supra. A [<sup>33</sup>-P] UTP-labeled antisense riboprobe was generated from a PCR product and hybridized at 55° C. overnight. The slides were dipped in Kodak NTB2 nuclear track emulsion and exposed for 4 weeks.

#### [1832] <sup>33</sup>P-Riboprobe Synthesis

[1833] 6.0 μl (125 mCi) of <sup>33</sup>P-UTP (Amersham BF 1002, SA <2000 Ci/mmol) were speed vac dried. To each tube containing dried <sup>33</sup>P-UTP, the following ingredients were added:

[1834] 2.0 μl 5×transcription buffer

[1835] 1.0 μl DTT (100 mM)

[1836] 2.0 μl NTP mix (2.5 mM: 10μ; each of 10 mM GTP, CTP & ATP+10 μl H<sub>2</sub>O)

[1837] 1.0 μl UTP (50 μM)

[1838] 1.0 μl Rnasin

[1839] 1.0 μl DNA template (1 μg)

[1840] 1.0 μl H<sub>2</sub>O

[1841] 1.0 μl RNA polymerase (for PCR products T3=AS, T7=S, usually)

[1842] The tubes were incubated at 37° C. for one hour. 1.0 μl RQ1 DNase were added, followed by incubation at 37° C. for 15 minutes. 90 μl TE (10 mM Tris pH 7.6/1 mM EDTA pH 8.0) were added, and the mixture was pipetted onto DE81 paper. The remaining solution was loaded in a Microcon-50 ultrafiltration unit, and spun using program 10 (6 minutes). The filtration unit was inverted over a second tube and spun using program 2 (3 minutes). After the final recovery spin, 100 μl TE were added. 1 μl of the final product was pipetted on DE81 paper and counted in 6 ml of Biofluor II.

[1843] The probe was run on a TBE/urea gel. 1-3 μl of the probe or 5 μl of RNA Mrk III were added to 3 μl of loading buffer. After heating on a 95° C. heat block for three minutes, the gel was immediately placed on ice. The wells of gel were flushed, the sample loaded, and run at 180-250 volts for 45 minutes. The gel was wrapped in saran wrap and exposed to XAR film with an intensifying screen in -70° C. freezer one hour to overnight.

#### [1844] <sup>33</sup>P-Hybridization

##### [1845] A. Pretreatment of frozen sections

[1846] The slides were removed from the freezer, placed on aluminium trays and thawed at room temperature for 5 minutes. The trays were placed in 55° C. incubator for five minutes to reduce condensation. The slides were fixed for 10 minutes in 4% paraformaldehyde on ice in the fume hood, arid washed in 0.5×SSC for 5 minutes, at room temperature (25 ml 20×SSC+975 ml SQ H<sub>2</sub>O). After deproteination in 0.5 μg/ml proteinase K for 10 minutes at 37° C. (12.5 μl of 10 mg/ml stock in 250 ml prewarmed RNase-free RNase buffer), the sections were washed in 0.5×SSC for 10 minutes at room temperature. The sections were dehydrated in 70%, 95%, 100% ethanol, 2 minutes each.

##### [1847] B. Pretreatment of paraffin-embedded sections

[1848] The slides were deparaffinized, placed in SQ H<sub>2</sub>O, and rinsed twice in 2×SSC at room temperature, for 5 minutes each time. The sections were deproteinated in 20 μg/ml proteinase K (500 μl of 10 mg/ml in 250 ml RNase-free RNase buffer; 37° C., 15 minutes)—human embryo, or 8×proteinase K (100 μl in 250 ml Rnas buffer, 37° C., 30 minutes)—formalin tissues. Subsequent rinsing in 0.5×SSC and dehydration were performed as described above.

##### [1849] C. Prehybridization

[1850] The slides were laid out in a plastic box lined with Box buffer (4×SSC, 50% formamide)—saturated filter paper. The tissue was covered with 50 μl of hybridization buffer (3.75 g Dextran Sulfate+6 ml SQ H<sub>2</sub>O), vortexed and heated in the microwave for 2 minutes with the cap loosened. After cooling on ice, 18.75 ml formamide, 3.75 ml 20×SSC and 9 ml SQ H<sub>2</sub>O were added, the tissue was vortexed well, and incubated at 42° C. for 1-4 hours.

##### [1851] D. Hybridization

[1852] 1.0×10<sup>6</sup> cpm probe and 1.0 μl tRNA (50 mg/ml stock) per slide were heated at 95° C. for 3 minutes. The slides were cooled on ice, and 48 μl hybridization buffer were added per slide. After vortexing, 50 μl <sup>33</sup>P mix were added to 50 μl prehybridization on slide. The slides were incubated overnight at 55° C.

##### [1853] E. Washes

[1854] Washing was done 2×10 minutes with 2×SSC, EDTA at room temperature (400 ml 20×SSC+16 ml 0.25M EDTA, V<sub>f</sub>=4 L), followed by RNaseA treatment at 37° C. for 30 minutes (500 μl of 10 mg/ml in 250 ml Rnas buffer=20 μg/ml). The slides were washed 2×10 minutes with 2×SSC, EDTA at room temperature. The stringency wash conditions were as follows: 2 hours at 55° C., 0.1×SSC, EDTA (20 ml 20×SSC+16 ml EDTA, V<sub>f</sub>=4 L).

##### [1855] F. Oligonucleotides

[1856] In situ analysis was performed on a variety of DNA sequences disclosed herein. The oligonucleotides employed for these analyses are as follows.

(1) DNA33094-1131 (PRO217)

p1 5' -GGATTCTAATACGACTCACTATAGGGCTCAGAAAAGCGCAACAGAGAA-3' (SEQ ID NO:348)

p2 5' -CTATGAAATTAACCCCTCACTAAAGGGATGTCTCCATGCCAACCTTC-3' (SEQ ID NO:349)

(2) DNA33223-1136 (PRO230)

p1 5' -GGATTCTAATACGACTCACTATAGGGCGCGATGTCCACTGGGGCTAC-3' (SEQ ID NO:350)

p2 5' -CTATGAAATTAACCCCTCACTAAAGGGACGAGGAAGATGGGCGGATGGT-3' (SEQ ID NO:351)

## -continued

- (3) DNA34435-1140 (PRO232)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCACCCACGCGTCCGGTCTT-3' (SEQ ID NO:352)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGACGGGGACACCACGGACCAGA-3' (SEQ ID NO:353)
- (4) DNA35639-1172 (PRO246)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCTTGCTGCGGTTTTTGTTCCTG-3' (SEQ ID NO:354)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGCTGCCGATCCACTGGTATT-3' (SEQ ID NO:355)
- (5) DNA49435-1219 (PRO533)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCGGATCCTGGCCGGCCTCTG-3' (SEQ ID NO:356)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGCCCGGCGATGGTCTCAGTTA-3' (SEQ ID NO:357)
- (6) DNA35638-1141 (PRO245)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCGGGAAGATGGCGAGGAGAG-3' (SEQ ID NO:358)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGACCAAGGCCACAAACGGAATC-3' (SEQ ID NO:359)
- (7) DNA33089-1132 (PRO221)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCTGTGCTTTCATTCTGCCAGTA-3' (SEQ ID NO:360)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGGTACAATTAAGGGGTGGAT-3' (SEQ ID NO:361)
- (8) DNA35918-1174 (PRO258)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCCGCCTCGCTCCTGCTCCTG-3' (SEQ ID NO:362)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGGATTGCCGCGACCCCTCACAG-3' (SEQ ID NO:363)
- (9) DNA32286-1191 (PRO214)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCCTCCTGCCCTCCCTGTCC-3' (SEQ ID NO:364)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGTGGTGGCCCGATTATCTGC-3' (SEQ ID NO:365)
- (10) DNA33221-1133 (PRO224)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCGCAGCGATGGCAGCGATGAGG-3' (SEQ ID NO:366)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGACAGCGGGCAGAGGGAGTG-3' (SEQ ID NO:367)
- (11) DNA35557-1137 (PRO234)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCAGGAGGCGTGAGGAGAAAC-3' (SEQ ID NO:368)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAAGACATGTCATCGGGAGTGG-3' (SEQ ID NO:369)
- (12) DNA33100-1159 (PRO229)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCGGGTGGAGGTGGAACAGAAA-3' (SEQ ID NO:370)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGACACAGACAGACCCCATACGC-3' (SEQ ID NO:371)
- (13) DNA34431-1177 (PRO263)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCAGGAAATCCGGATGTCTC-3' (SEQ ID NO:372)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGAGTAAGGGGATGCCACCGAGTA-3' (SEQ ID NO:373)
- (14) DNA38268-1188 (PRO295)  
 p1 5'-GGATTCTAATACGACTCACTATAGGGCCAGCTACCCGCAGGAGGAGG-3' (SEQ ID NO:374)  
 p2 5'-CTATGAAATTAACCCCTACTAAAGGGATCCCAGGTGATGAGGTCAGA-3' (SEQ ID NO:375)

**[1857]** G. Results

**[1858]** In situ analysis was performed on a variety of DNA sequences disclosed herein. The results from these analyses are as follows.

**[1859]** (1) DNA33094-1131 (PRO217)

**[1860]** Highly distinctive expression pattern, that does not indicate an obvious biological function. In the human embryo it was expressed in outer smooth muscle layer of the GI tract, respiratory cartilage, branching respiratory epithelium, osteoblasts, tendons, gonad, in the optic nerve head and developing dermis. In the adult expression was observed in the epidermal pegs of the chimp tongue, the basal epithelial/myoepithelial cells of the prostate and urinary bladder. Also expressed in the alveolar lining cells of the adult lung, mesenchymal cells juxtaposed to erectile tissue in the penis and the cerebral cortex (probably glial cells). In the kidney, expression was only seen in disease, in cells surrounding thyroidized renal tubules.

**[1861]** Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney,

adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

**[1862]** Adult human tissues examined: Kidney (normal and end-stage), adrenal, myocardium, aorta, spleen, lymph node, gall bladder, pancreas, lung, skin, eye (inc. retina), prostate, bladder, liver (normal, cirrhotic, acute failure).

**[1863]** Non-human primate tissues examined:

**[1864]** (a) Chimp Tissues: Salivary gland, stomach, thyroid, parathyroid, skin, thymus, ovary, lymph node.

**[1865]** (b) Rhesus Monkey Tissues: Cerebral cortex, hippocampus, cerebellum, penis.

**[1866]** (2) DNA33223-1136 (PRO230)

**[1867]** Sections show an intense signal associated with arterial and venous vessels in the fetus. In arteries the signal appeared to be confined to smooth-muscle/pericytic cells.

The signal is also seen in capillary vessels and in glomeruli. It is not clear whether or not endothelial cells are expressing this mRNA. Expression is also observed in epithelial cells in the fetal lens. Strong expression was also seen in cells within placental trophoblastic villi, these cells lie between the trophoblast and the fibroblast-like cells that express HGF—uncertain histogenesis. In the adult, there was no evidence of expression and the wall of the aorta and most vessels appear to be negative. However, expression was seen over vascular channels in the normal prostate and in the epithelium lining the gallbladder. Insurers expression was seen in the vessels of the soft-tissue sarcoma and a renal cell carcinoma. In summary, this is a molecule that shows relatively specific vascular expression in the fetus as well as in some adult organs. Expression was also observed in the fetal lens and the adult gallbladder

[1868] In a secondary screen, vascular expression was observed, similar to that observed above, seen in fetal blocks. Expression is on vascular smooth muscle, rather than endothelium. Expression also seen in smooth muscle of the developing oesophagus, so as reported previously, this molecule is not vascular specific. Expression was examined in 4 lung and 4 breast carcinomas. Substantial expression was seen in vascular smooth muscle of at least 3/4 lung cancers and 2/4 breast cancers. In addition, in one breast carcinoma, expression was observed in peritumoral stromal cells of uncertain histogenesis (possibly myofibroblasts). No endothelial cell expression was observed in this study.

[1869] (3) DNA34435-1140 (PRO232)

[1870] Strong expression in prostatic epithelium and bladder epithelium, lower level of expression in bronchial epithelium. High background/low level expression seen in a number of sites, including among others, bone, blood, chondrosarcoma, adult heart and fetal liver. It is felt that this level of signal represents background, partly because signal at this level was seen over the blood. All other tissues negative.

[1871] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis, testis and lower limb.

[1872] Adult human tissues examined: Kidney (normal and end-stage), adrenal, spleen, lymph node, pancreas, lung, eye (inc. retina), bladder, liver (normal, cirrhotic, acute failure).

[1873] Non-human primate tissues examined:

[1874] Chim Tissues: adrenal

[1875] Rhesus Monkey Tissues: Cerebral cortex, hippocampus

[1876] In a secondary screen, expression was observed in the epithelium of the prostate, the superficial layers of the urethelium of the urinary bladder, the urethelium lining the renal pelvis and the urethelium of the ureter (1 out of 2 experiments). The urethra of a rhesus monkey was negative; it is unclear whether this represents a true lack of expression by the urethra, or if it is the result of a failure of the probe to cross react with rhesus tissue. The findings in the prostate and bladder are similar to those previously described using

an isotopic detection technique. Expression of the mRNA for this antigen is NOT prostate epithelial specific. The antigen may serve as a useful marker for urethelial derived tissues. Expression in the superficial, post-mitotic cells, of the urinary tract epithelium also suggest that it is unlikely to represent a specific stem cell marker, as this would be expected to be expressed specifically in basal epithelium.

[1877] (4) DNA35639-1172 (PRO246)

[1878] Strongly expressed in fetal vascular endothelium, including tissues of the CNS. Lower level of expression in adult vasculature, including the CNS. Not obviously expressed at higher levels in tumor vascular endothelium. Signal also seen over bone matrix and adult spleen, not obviously cell associated, probably related to non-specific background at these sites.

[1879] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis, testis and lower limb.

[1880] Adult human tissues examined: Kidney (normal and end-stage), adrenal, spleen, lymph node, pancreas, lung, eye (inc. retina), bladder, liver (normal, cirrhotic, acute failure).

[1881] Non-human primate tissues examined:

[1882] 253: Chim Tissues: adrenal

[1883] Rhesus Monkey Tissues: Cerebral cortex, hippocampus

[1884] (5) DNA49435-1219 (PRO533)

[1885] Moderate expression over cortical neurones in the fetal brain. Expression over the inner aspect of the fetal retina, possible expression in the developing lens. Expression over fetal skin, cartilage, small intestine, placental villi and umbilical cord. In adult tissues there is an extremely high level of expression over the gallbladder epithelium. Moderate expression over the adult kidney, gastric and colonic epithelia. Low-level expression was observed over many cell types in many tissues, this may be related to stickiness of the probe, these data should therefore be interpreted with a degree of caution.

[1886] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis, testis and lower limb.

[1887] Adult human tissues examined: Kidney (normal and end-stage), adrenal, spleen, lymph node, pancreas, lung, eye (inc. retina), bladder, liver (normal, cirrhotic, acute failure).

[1888] Non-human primate tissues examined:

[1889] Chim Tissues: adrenal

[1890] Rhesus Monkey Tissues: Cerebral cortex, hippocampus, cerebellum.



[1891] (6) DNA35638-1141 (PRO245)

[1892] Expression observed in the endothelium lining a subset of fetal and placental vessels. Endothelial expression was confined to these tissue blocks. Expression also observed over intermediate trophoblast cells of placenta. Expression also observed tumor vasculature but not in the vasculature of normal tissues of the same type. All other tissues negative.

[1893] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1894] Adult tissues examined: Liver, kidney, adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus(rm), cerebellum(rm), penis, eye, bladder, stomach, gastric carcinoma, colon, colonic carcinoma, thyroid (chimp), parathyroid (chimp) ovary (chimp) and chondrosarcoma. Acetaminophen induced liver injury and hepatic cirrhosis

[1895] (7) DNA33089-1132 (PRO221)

[1896] Specific expression over fetal cerebral white and grey matter, as well as over neurones in the spinal cord. Probe appears to cross react with rat. Low level of expression over cerebellar neurones in adult rhesus brain. All other tissues negative.

[1897] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1898] Adult tissues examined: Liver, kidney, adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus(rm), cerebellum(rm), penis, eye, bladder, stomach, gastric carcinoma, colon, colonic carcinoma and chondrosarcoma. Acetaminophen induced liver injury and hepatic cirrhosis

[1899] (8) DNA35918-1174 (PRO258)

[1900] Strong expression in the nervous system. In the rhesus monkey brain expression is observed in cortical, hippocampal and cerebellar neurones. Expression over spinal neurones in the fetal spinal cord, the developing brain and the inner aspects of the fetal retina. Expression over developing dorsal root and autonomic ganglia as well as enteric nerves. Expression observed over ganglion cells in the adult prostate. In the rat, there is strong expression over the developing hind brain and spinal cord. Strong expression over interstitial cells in the placental villi. All other tissues were negative.

[1901] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1902] Adult tissues examined: Liver, kidney, renal cell carcinoma, adrenal, aorta, spleen, lymph node, pancreas, lung, myocardium, skin, cerebral cortex (rm), hippocampus(rm), cerebellum(rm), bladder, prostate, stomach, gastric carcinoma, colon, colonic carcinoma, thyroid (chimp), parathyroid (chimp) ovary (chimp) and chondrosarcoma. Acetaminophen induced liver injury and hepatic cirrhosis.

[1903] (9) DNA32286-1191 (PRO214)

[1904] Fetal tissue: Low level throughout mesenchyme. Moderate expression in placental stromal cells in membranous tissues and in thyroid. Low level expression in cortical neurones. Adult tissue: all negative.

[1905] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1906] Adult tissues examined include: Liver, kidney, adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung and skin.

[1907] (10) DNA33221-1133 (PRO224)

[1908] Expression limited to vascular endothelium in fetal spleen, adult spleen, fetal liver, adult thyroid and adult lymph node (chimp). Additional site of expression is the developing spinal ganglia. All other tissues negative.

[1909] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1910] Adult human tissues examined: Kidney (normal and end-stage), adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, eye (inc. retina), bladder, liver (normal, cirrhotic, acute failure).

[1911] Non-human primate tissues examined:

[1912] Chimp Tissues: Salivary gland, stomach, thyroid, parathyroid, skin, thymus, ovary, lymph node.

[1913] Rhesus Monkey Tissues: Cerebral cortex, hippocampus, cerebellum, penis.

[1914] (11) DNA35557-1137 (PRO234)

[1915] Specific expression over developing motor neurones in ventral aspect of the fetal spinal cord (will develop into ventral horns of spinal cord). All other tissues negative. Possible role in growth, differentiation and/or development of spinal motor neurones.

[1916] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1917] Adult tissues examined: Liver, kidney, adrenal, myocardium, aorta, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus(rm),

cerebellum(rm), penis, eye, bladder, stomach, gastric carcinoma, colon, colonic carcinoma and chondrosarcoma. Acetaminophen induced liver injury and hepatic cirrhosis

[1918] (12) DNA33100-1159 (PRO229)

[1919] Striking expression in mononuclear phagocytes (macrophages) of fetal and adult spleen, liver, lymph node and adult thymus (in tingible body macrophages). The highest expression is in the spleen. All other tissues negative. Localisation and homology are entirely consistent with a role as a scavenger receptor for cells of the reticuloendothelial system. Expression also observed in placental mononuclear cells.

[1920] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1921] Adult human tissues examined: Kidney (normal and end-stage), adrenal, myocardium, aorta, spleen, lymph node, gall bladder, pancreas, lung, skin, eye (inc. retina), prostate, bladder, liver (normal, cirrhotic, acute failure).

[1922] Non-human primate tissues examined:

[1923] Chimp Tissues: Salivary gland, stomach, thyroid, parathyroid, skin, thymus, ovary, lymph node.

[1924] Rhesus Monkey Tissues: Cerebral cortex, hippocampus, cerebellum, penis.

[1925] (13) DNA34431-1177 (PRO263)

[1926] Widespread expression in human fetal tissues and placenta over mononuclear cells, probably macrophages+/- lymphocytes. The cellular distribution follows a perivascular pattern in many tissues. Strong expression also seen in epithelial cells of the fetal adrenal cortex. All adult tissues were negative.

[1927] Fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, heart, great vessels, oesophagus, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis and lower limb.

[1928] Adult tissues examined: Liver, kidney, adrenal, spleen, lymph node, pancreas, lung, skin, cerebral cortex (rm), hippocampus(rm), cerebellum(rm), bladder, stomach, colon and colonic carcinoma. Acetaminophen induced liver injury and hepatic cirrhosis.

[1929] A secondary screen evidenced expression over stromal mononuclear cells probably histiocytes.

[1930] (14) DNA38268-1188 (PRO295)

[1931] High expression over ganglion cells in human fetal spinal ganglia and over large neurones in the anterior horns of the developing spinal cord. In the adult there is expression in the chimp adrenal medulla (neural), neurones of the rhesus monkey brain (hippocampus [+++]) and cerebral cortex) and neurones in ganglia in the normal adult human prostate (the only section that contains ganglion cells, ie

expression in this cell type is presumed NOT to be confined to the prostate). All other tissues negative.

[1932] Human fetal tissues examined (E12-E16 weeks) include: Placenta, umbilical cord, liver, kidney, adrenals, thyroid, lungs, great vessels, stomach, small intestine, spleen, thymus, pancreas, brain, eye, spinal cord, body wall, pelvis, testis and lower limb.

[1933] Adult human tissues examined: Kidney (normal and end-stage), adrenal, spleen, lymph node, pancreas, lung, eye (inc. retina), bladder, liver (normal, cirrhotic, acute failure).

[1934] Non-human primate tissues examined:

[1935] Chimp Tissues: adrenal

[1936] Rhesus Monkey Tissues: Cerebral cortex, hippocampus, cerebellum.

#### Example 103

[1937] Isolation of cDNA Clones Encoding Human PRO1868

[1938] A consensus DNA sequence was assembled relative to other EST sequences using phrap as described in Example 1 above. This consensus sequence is herein designated DNA49803. Based up an observed homology between the DNA49803 consensus sequence and an EST sequence contained within the Incyte EST clone no. 2994689, Incyte EST clone no. 2994689 was purchased and its insert obtained and sequenced. The sequence of that insert is shown in **FIG. 123** and is herein designated DNA77624-2515.

[1939] The entire nucleotide sequence of DNA77624-2515 is shown in **FIG. 123** (SEQ ID NO:422). Clone DNA77624-2515 contains a single open reading frame with an apparent translational initiation site at nucleotide positions 51-53 and ending at the stop codon at nucleotide positions 981-983 (**FIG. 123**). The predicted polypeptide precursor is 310 amino acids long (**FIG. 124**). The full-length PRO1868 protein shown in **FIG. 124** has an estimated molecular weight of about 35,020 daltons and a pI of about 7.90. Analysis of the full-length PRO1868 sequence shown in **FIG. 124** (SEQ ID NO:423) evidences the presence of the following: a signal peptide from about amino acid 1 to about amino acid 30, a transmembrane domain from about amino acid 243 to about amino acid 263, potential N-glycosylation sites from about amino acid 104 to about amino acid 107 and from about amino acid 192 to about amino acid 195, a cAMP- and cGMP-dependent protein kinase phosphorylation site from about amino acid 107 to about amino acid 110, casein kinase II phosphorylation sites from about amino acid 106 to about amino acid 109 and from about amino acid 296 to about amino acid 299, a tyrosine kinase phosphorylation site from about amino acid 69 to about amino acid 77 and potential N-myristoylation sites from about amino acid 26 to about amino acid 31, from about amino acid 215 to about amino acid 220, from about amino acid 226 to about amino acid 231, from about amino acid 243 to about amino acid 248, from about amino acid 244 to about amino acid 249 and from about amino acid 262 to about amino acid 267. Clone DNA77624-2515 has been deposited with ATCC on Dec. 22, 1998 and is assigned ATCC deposit no. 203553.

[1940] An analysis of the Dayhoff database (version 35.45 SwissProt 35), using a WU-BLAST2 sequence alignment analysis of the full-length sequence shown in FIG. 124 (SEQ ID NO:423), evidenced significant homology between the PRO1868 amino acid sequence and the following Dayhoff sequences: HGS\_RC75, P\_W61379, A33\_HUMAN, P\_W14146, P\_W14158, AMAL\_DROME, P\_R77437, I38346, NCM2\_HUMAN and PTPD\_HUMAN.

#### Example 104

##### [1941] Identification of Receptor/Ligand Interactions

[1942] In this assay, various PRO polypeptides are tested for ability to bind to a panel of potential receptor molecules for the purpose of identifying receptor/ligand interactions. The identification of a ligand for a known receptor, a receptor for a known ligand or a novel receptor/ligand pair is useful for a variety of indications including, for example, targeting bioactive molecules (linked to the ligand or receptor) to a cell known to express the receptor or ligand, use of the receptor or ligand as a reagent to detect the presence of the ligand or receptor in a composition suspected of containing the same, wherein the composition may comprise cells suspected of expressing the ligand or receptor, modulating the growth of or another biological or immunological activity of a cell known to express or respond to the receptor or ligand, modulating the immune response of cells or toward cells that express the receptor or ligand, allowing the preparation of agonists, antagonists and/or antibodies directed against the receptor or ligand which will modulate the growth of or a biological or immunological activity of a cell expressing the receptor or ligand, and various other indications which will be readily apparent to the ordinarily skilled artisan.

[1943] The assay is performed as follows. A PRO polypeptide of the present invention suspected of being a ligand for a receptor is expressed as a fusion protein containing the Fc domain of human IgG (an immunoadhesin). Receptor-ligand binding is detected by allowing interaction of the immunoadhesin polypeptide with cells (e.g. Cos cells) expressing candidate PRO polypeptide receptors and visualization of bound immunoadhesin with fluorescent reagents directed toward the Fc fusion domain and examination by microscope. Cells expressing candidate receptors are produced by transient transfection, in parallel, of defined subsets of a library of cDNA expression vectors encoding PRO polypeptides that may function as receptor molecules. Cells are then incubated for 1 hour in the presence of the PRO polypeptide immunoadhesin being tested for possible receptor binding. The cells are then washed and fixed with paraformaldehyde. The cells are then incubated with fluorescent conjugated antibody directed against the Fc portion of the PRO polypeptide immunoadhesin (e.g. FITC conjugated goat anti-human-Fc antibody). The cells are then washed again and examined by microscope. A positive interaction is judged by the presence of fluorescent labeling of cells transfected with cDNA encoding a particular PRO polypeptide receptor or pool of receptors and an absence of similar fluorescent labeling of similarly prepared cells that have been transfected with other cDNA or pools of cDNA. If a defined pool of cDNA expression vectors is judged to be positive for interaction with a PRO polypeptide immunoadhesin, the individual cDNA species that comprise the pool are tested individually (the pool is "broken down") to

determine the specific cDNA that encodes a receptor able to interact with the PRO polypeptide immunoadhesin.

[1944] In another embodiment of this assay, an epitope-tagged potential ligand PRO polypeptide (e.g. 8 histidine "His" tag) is allowed to interact with a panel of potential receptor PRO polypeptide molecules that have been expressed as fusions with the Fc domain of human IgG (immunoadhesins). Following a 1 hour co-incubation with the epitope tagged PRO polypeptide, the candidate receptors are each immunoprecipitated with protein A beads and the beads are washed. Potential ligand interaction is determined by western blot analysis of the immunoprecipitated complexes with antibody directed towards the epitope tag. An interaction is judged to occur if a band of the anticipated molecular weight of the epitope tagged protein is observed in the western blot analysis with a candidate receptor, but is not observed to occur with the other members of the panel of potential receptors.

[1945] Using these assays, the following receptor/ligand interactions have been herein identified: PRO245 binds to PRO1868.

##### [1946] Deposit of Material

[1947] The following materials have been deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Md., USA (ATCC):

Material	ATCC Dep. No.	Deposit Date
DNA32292-1131	ATCC 209258	September 16, 1997
DNA33094-1131	ATCC 209256	September 16, 1997
DNA33223-1136	ATCC 209264	September 16, 1997
DNA34435-1140	ATCC 209250	September 16, 1997
DNA27864-1155	ATCC 209375	October 16, 1997
DNA36350-1158	ATCC 209378	October 16, 1997
DNA32290-1164	ATCC 209384	October 16, 1997
DNA35639-1172	ATCC 209396	October 17, 1997
DNA33092-1202	ATCC 209420	October 28, 1997
DNA49435-1219	ATCC 209480	November 21, 1997
DNA35638-1141	ATCC 209265	September 16, 1997
DNA32298-1132	ATCC 209257	September 16, 1997
DNA33089-1132	ATCC 209262	September 16, 1997
DNA33786-1132	ATCC 209253	September 16, 1997
DNA35918-1174	ATCC 209402	October 17, 1997
DNA37150-1178	ATCC 209401	October 17, 1997
DNA38260-1180	ATCC 209397	October 17, 1997
DNA39969-1185	ATCC 209400	October 17, 1997
DNA32286-1191	ATCC 209385	October 16, 1997
DNA33461-1199	ATCC 209367	October 15, 1997
DNA40628-1216	ATCC 209432	November 7, 1997
DNA33221-1133	ATCC 209263	September 16, 1997
DNA33107-1135	ATCC 209251	September 16, 1997
DNA35557-1137	ATCC 209255	September 16, 1997
DNA34434-1139	ATCC 209252	September 16, 1997
DNA33100-1159	ATCC 209373	October 16, 1997
DNA35600-1162	ATCC 209370	October 16, 1997
DNA34436-1238	ATCC 209523	December 10, 1997
DNA33206-1165	ATCC 209372	October 16, 1997
DNA35558-1167	ATCC 209374	October 16, 1997
DNA35599-1168	ATCC 209373	October 16, 1997
DNA36992-1168	ATCC 209382	October 16, 1997
DNA34407-1169	ATCC 209383	October 16, 1997
DNA35841-1173	ATCC 209403	October 17, 1997
DNA33470-1175	ATCC 209398	October 17, 1997
DNA34431-1177	ATCC 209399	October 17, 1997
DNA39510-1181	ATCC 209392	October 17, 1997
DNA39423-1182	ATCC 209387	October 17, 1997
DNA40620-1183	ATCC 209388	October 17, 1997
DNA40604-1187	ATCC 209394	October 17, 1997

-continued

Material	ATCC Dep. No.	Deposit Date
DNA38268-1188	ATCC 209421	October 28, 1997
DNA37151-1193	ATCC 209393	October 17, 1997
DNA35673-1201	ATCC 209418	October 28, 1997
DNA40370-1217	ATCC 209485	November 21, 1997
DNA42551-1217	ATCC 209483	November 21, 1997
DNA39520-1217	ATCC 209482	November 21, 1997
DNA41225-1217	ATCC 209491	November 21, 1997
DNA43318-1217	ATCC 209481	November 21, 1997
DNA40587-1231	ATCC 209438	November 7, 1997
DNA41338-1234	ATCC 209927	June 2, 1998
DNA40981-1234	ATCC 209439	November 7, 1997
DNA37140-1234	ATCC 209489	November 21, 1997
DNA40982-1235	ATCC 209433	November 7, 1997
DNA41379-1236	ATCC 209488	November 21, 1997
DNA44167-1243	ATCC 209434	November 7, 1997
DNA39427-1179	ATCC 209395	October 17, 1997
DNA40603-1232	ATCC 209486	November 21, 1997
DNA43466-1225	ATCC 209490	November 21, 1997
DNA43046-1225	ATCC 209484	November 21, 1997
DNA35668-1171	ATCC 209371	October 16, 1997
DNA77624-2515	ATCC 203553	December 22, 1998

[1948] These deposit were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

[1949] The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

[1950] The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to

those skilled in the art from the foregoing description and full within the scope of the appended claims.

## Appendix A

I hereby claim benefit under Title 35, United States Code, §119(e) of any United States Provisional applications listed below:

Provisional Application Ser. No.	Filing Date
60/059,115	September 17, 1997
60/059,184	September 17, 1997
60/059,122	September 17, 1997
60/059,117	September 17, 1997
60/059,113	September 17, 1997
60/059,121	September 17, 1997
60/059,119	September 17, 1997
60/059,263	September 18, 1997
60/059,266	September 18, 1997
60/062,125	October 15, 1997
60/062,287	October 17, 1997
60/062,285	October 17, 1997
60/063,486	October 21, 1997
60/062,816	October 24, 1997
60/062,814	October 24, 1997
60/063,127	October 24, 1997
60/063,120	October 24, 1997
60/063,121	October 24, 1997
60/063,045	October 24, 1997
60/063,128	October 24, 1997
60/063,329	October 27, 1997
60/063,327	October 27, 1997
60/063,549	October 28, 1997
60/063,541	October 28, 1997
60/063,550	October 28, 1997
60/063,542	October 28, 1997
60/063,544	October 28, 1997
60/063,564	October 28, 1997
60/063,734	October 29, 1997
60/063,738	October 29, 1997
60/063,704	October 29, 1997
60/063,435	October 29, 1997
60/064,215	October 29, 1997
60/063,735	October 29, 1997
60/063,732	October 29, 1997
60/064,103	October 31, 1997
60/063,870	October 31, 1997
60/064,248	November 3, 1997
60/064,809	November 7, 1997
60/065,846	November 11, 1997
60/065,186	November 12, 1997
60/065,846	November 17, 1997
60/065,693	November 18, 1997
60/066,120	November 21, 1997
60/066,364	November 21, 1997
60/066,772	November 24, 1997
60/066,466	November 24, 1997
60/066,770	November 24, 1997
60/066,511	November 24, 1997
60/066,453	November 24, 1997
60/066,840	November 25, 1997
60/069,425	December 12, 1997
60/088,026	June 4, 1998
60/099,803	September 10, 1998
60/100,262	September 14, 1998
60/100,858	September 17, 1998
60/104,080	October 13, 1998
60/109,304	November 20, 1998
60/113,296	December 22, 1998
60/143,048	July 7, 1999
60/145,698	July 26, 1999
60/146,222	July 28, 1999

-continued

Appendix B

I hereby claim benefit under Title 35, United States Code, §120 of any United States and PCT International applications listed below:

PCT/US Ser. No.	Filed	Status
PCT/US98/18824	September 10, 1998	Abandoned
PCT/US98/19177	September 14, 1998	Abandoned
PCT/US98/19330	September 16, 1998	Abandoned
PCT/US98/19437	September 17, 1998	Abandoned
PCT/US98/25108	December 1, 1998	Abandoned
PCT/US99/20594	September 8, 1999	Pending
PCT/US99/20944	September 13, 1999	Pending
PCT/US99/21090	September 15, 1999	Pending
PCT/US99/21547	September 15, 1999	Pending
PCT/US99/23089	October 5, 1999	Pending
PCT/US99/28214	November 29, 1999	Pending
PCT/US99/28313	November 30, 1999	Pending

-continued

PCT/US99/28301	December 1, 1999	Abandoned
PCT/US99/28564	December 2, 1999	Pending
PCT/US99/28565	December 2, 1999	Pending
PCT/US99/30095	December 16, 1999	Pending
PCT/US99/30999	December 20, 1999	Pending
PCT/US99/30911	December 20, 1999	Pending
PCT/US00/00219	January 5, 2000	Pending
PCT/US00/03565	February 11, 2000	Pending
PCT/US00/04414	February 22, 2000	Pending
PCT/US00/05004	February 24, 2000	Pending
PCT/US00/05841	March 2, 2000	Pending
PCT/US00/07377	March 20, 2000	Pending
PCT/US00/08439	March 30, 2000	Pending
PCT/US00/14042	May 22, 2000	Pending
PCT/US00/15264	June 2, 2000	Pending
PCT/US00/20710	July 28, 2000	Pending
PCT/US00/23328	August 24, 2000	Pending

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 423

<210> SEQ ID NO 1

<211> LENGTH: 1825

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

actgcacctc ggttctatcg attgaattcc ccggggatcc tctagagatc cctcgacctc    60
gaccocacgc tccgggccgg agcagcacgg ccgcaggacc tggagctccg gctgctcttt    120
cccgcagcgc taccgcccat gcgcctgccc cgcggggccg cgctggggct cctgccgctt    180
ctgctgtgct tgcccgccgc gccggaggcc gccaaagaag cgacgccctg ccaccggtgc    240
cggggctggt tggacaagtt taaccagggg atggtggaca ccgcaaagaa gaactttggc    300
ggcgggaaca cggtctggga gaaaagacg ctgtccaagt acgagtccag cgagattcgc    360
ctgctggaga tcctggaggg gctgtgagag agcagcgact tcgaatgcaa tcagatgcta    420
gaggcgcagg aggagcacct ggaggcctgg tggctgcagc tgaagagcga atatcctgac    480
ttattcagat ggttttgtgt gaagacactg aaagtgtgct gctctccagg aacctacggt    540
cccgactgtc tcgcatgccca gggcgatcc cagaggccct gcagcgggaa tggccactgc    600
agcggagatg ggagcagaca gggcgacggg tcctgccggt gccacatggy gtaccagggc    660
ccgctgtgca ctgactgcat ggacggctac ttcagctcgc tccggaacga gaccacagc    720
atctgcacag cctgtgacga gtccctgcaag acgtgctcgg gctgaccaa cagagactgc    780
ggcgagtgtg aagtgggctg ggtgctggac gaggcgcct gtgtggatgt ggacgagtgt    840
gcgcccgagc cgcctccctg cagcgtgctg cagttctgta agaacccaa cggtcctac    900
acgtgcgaag agtgtgactc cagctgtgtg ggctgcacag gggaaggccc aggaaactgt    960
    
```

-continued

```

aaagagtgta tctctggcta cgcgagggag cacggacagt gtgcagatgt ggacgagtgc 1020
tcactagcag aaaaaacctg tgtgagggaa aacgaaaact gctacaatac tccagggagc 1080
tacgtctgtg tgtgtcctga cggtctcga gaaacggaag atgctctgtg gccgccgca 1140
gaggctgaag ccacagaagg agaaagcccg acacagctgc cctcccgcga agacctgtaa 1200
tgtgcccggc ttacccttta aattattcag aaggatgtcc cgtggaaaat gtggccctga 1260
ggatgccgtc tcctgcagtg gacagcggcg gggagaggct gcctgtcttc taacggttga 1320
ttctcatttg tcccttaaac agctgcattt cttggttgtt cttaaacaga cttgtatatt 1380
ttgatacagt tctttgtaat aaaattgacc attgtaggta atcaggagga aaaaaaaaaa 1440
aaaaaaaaaa aaagggcggc cgcgactcta gagtcgacct gcagaagctt gcccgccatg 1500
gcccaacttg tttattgcag cttataatgg ttacaaataa agcaatagca tcacaaattt 1560
cacaataaaa gcattttttt cactgcattc tagttgtggt ttgtccaaac tcatcaatgt 1620
atcttatcat gtctggatcg ggaattaatt cggcgcagca ccatggcctg aaataacctc 1680
tgaaagagga acttggttag gtaccttctg aggcggaaag aaccagctgt ggaatgtgtg 1740
tcagttaggg tgtgaaaagt ccccaggctc cccagcaggc agaagtatgc aagcatgcat 1800
ctcaattagt cagcaacca gtttt 1825
    
```

```

<210> SEQ ID NO 2
<211> LENGTH: 353
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 2

```

Met Arg Leu Pro Arg Arg Ala Ala Leu Gly Leu Leu Pro Leu Leu Leu
 1           5           10           15
Leu Leu Pro Pro Ala Pro Glu Ala Ala Lys Lys Pro Thr Pro Cys His
          20           25           30
Arg Cys Arg Gly Leu Val Asp Lys Phe Asn Gln Gly Met Val Asp Thr
          35           40           45
Ala Lys Lys Asn Phe Gly Gly Gly Asn Thr Ala Trp Glu Glu Lys Thr
          50           55           60
Leu Ser Lys Tyr Glu Ser Ser Glu Ile Arg Leu Leu Glu Ile Leu Glu
 65           70           75           80
Gly Leu Cys Glu Ser Ser Asp Phe Glu Cys Asn Gln Met Leu Glu Ala
          85           90           95
Gln Glu Glu His Leu Glu Ala Trp Trp Leu Gln Leu Lys Ser Glu Tyr
          100          105          110
Pro Asp Leu Phe Glu Trp Phe Cys Val Lys Thr Leu Lys Val Cys Cys
          115          120          125
Ser Pro Gly Thr Tyr Gly Pro Asp Cys Leu Ala Cys Gln Gly Gly Ser
          130          135          140
Gln Arg Pro Cys Ser Gly Asn Gly His Cys Ser Gly Asp Gly Ser Arg
          145          150          155          160
Gln Gly Asp Gly Ser Cys Arg Cys His Met Gly Tyr Gln Gly Pro Leu
          165          170          175
Cys Thr Asp Cys Met Asp Gly Tyr Phe Ser Ser Leu Arg Asn Glu Thr
          180          185          190
His Ser Ile Cys Thr Ala Cys Asp Glu Ser Cys Lys Thr Cys Ser Gly
    
```



-continued

---

```

gttcaagcc tgtctgcgag cctggctgtg gtgcacatgg aacctgccat gaaccaaca 1200
aatgccaatg tcaagaaggt tggcatgaa gacactgcaa taaaaggtag gaagccagcc 1260
tcatacatgc cctgaggcca gcaggcgccc agctcaggca gcacacgcct tcacttaaaa 1320
aggccgagga ggcgcgggat ccacatgaat ccaattacat ctggatgaact cgcacatctg 1380
aaacgtttta agttacacca agttcatagc ctttgtaac ctttcatgtg ttgaatgttc 1440
aaataatggt cattacactt aagaatactg gcctgaattt tattagcttc attataaatc 1500
actgagctga tatttactct tccttttaag ttttctaagt acgtctgtag catgatggta 1560
tagattttct tgtttcagtg ctttgggaca gattttatat tatgtcaatt gatcagggta 1620
aaattttcag tgtgtagtgt gcagatatatt tcaaaattac aatgcattta tgggtgtctgg 1680
gggcagggga acatcagaaa ggttaaatg ggcaaaaatg cgtaaatcac aagaatttgg 1740
atggtgcagt taatgttgaa gttacagcat ttcagatttt attgtcagat atttagatgt 1800
ttgttacatt ttaaaaaatt gctcttaatt ttaaaactct caatacaata tattttgacc 1860
ttaccattat tccagagatt cagtattaaa aaaaaaaaaa ttacactgtg gtagtggcat 1920
ttaacaata taatatattc taacacaat gaaataggga atataatgta tgaacttttt 1980
gcattggcct gaagcaatat aatatattgt aaacaaaaca cagctcttac ctaataaaca 2040
ttttatactg tttgtatgta taaaataaag gtgctgcttt agttttttgg aaaaaaaaaa 2100
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa gggcgccgc gactctagag tgcacctgca 2160
gaagcttggc cgccatggcc caactgtttt attgcagctt ataatg 2206

```

&lt;210&gt; SEQ ID NO 4

&lt;211&gt; LENGTH: 379

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 4

```

Met Ala Arg Arg Ser Ala Phe Pro Ala Ala Ala Leu Trp Leu Trp Ser
 1           5           10           15
Ile Leu Leu Cys Leu Leu Ala Leu Arg Ala Glu Ala Gly Pro Pro Gln
          20           25           30
Glu Glu Ser Leu Tyr Leu Trp Ile Asp Ala His Gln Ala Arg Val Leu
          35           40           45
Ile Gly Phe Glu Glu Asp Ile Leu Ile Val Ser Glu Gly Lys Met Ala
          50           55           60
Pro Phe Thr His Asp Phe Arg Lys Ala Gln Gln Arg Met Pro Ala Ile
          65           70           75           80
Pro Val Asn Ile His Ser Met Asn Phe Thr Trp Gln Ala Ala Gly Gln
          85           90           95
Ala Glu Tyr Phe Tyr Glu Phe Leu Ser Leu Arg Ser Leu Asp Lys Gly
          100          105          110
Ile Met Ala Asp Pro Thr Val Asn Val Pro Leu Leu Gly Thr Val Pro
          115          120          125
His Lys Ala Ser Val Val Gln Val Gly Phe Pro Cys Leu Gly Lys Gln
          130          135          140
Asp Gly Val Ala Ala Phe Glu Val Asp Val Ile Val Met Asn Ser Glu
          145          150          155          160
Gly Asn Thr Ile Leu Gln Thr Pro Gln Asn Ala Ile Phe Phe Lys Thr

```



-continued

165		170		175											
Cys	Gln	Gln	Ala	Glu	Cys	Pro	Gly	Gly	Cys	Arg	Asn	Gly	Gly	Phe	Cys
			180					185					190		
Asn	Glu	Arg	Arg	Ile	Cys	Glu	Cys	Pro	Asp	Gly	Phe	His	Gly	Pro	His
		195					200					205			
Cys	Glu	Lys	Ala	Leu	Cys	Thr	Pro	Arg	Cys	Met	Asn	Gly	Gly	Leu	Cys
	210					215					220				
Val	Thr	Pro	Gly	Phe	Cys	Ile	Cys	Pro	Pro	Gly	Phe	Tyr	Gly	Val	Asn
225					230					235					240
Cys	Asp	Lys	Ala	Asn	Cys	Ser	Thr	Thr	Cys	Phe	Asn	Gly	Gly	Thr	Cys
				245					250					255	
Phe	Tyr	Pro	Gly	Lys	Cys	Ile	Cys	Pro	Pro	Gly	Leu	Glu	Gly	Glu	Gln
			260					265					270		
Cys	Glu	Ile	Ser	Lys	Cys	Pro	Gln	Pro	Cys	Arg	Asn	Gly	Gly	Lys	Cys
		275					280					285			
Ile	Gly	Lys	Ser	Lys	Cys	Lys	Cys	Ser	Lys	Gly	Tyr	Gln	Gly	Asp	Leu
	290					295					300				
Cys	Ser	Lys	Pro	Val	Cys	Glu	Pro	Gly	Cys	Gly	Ala	His	Gly	Thr	Cys
305					310					315					320
His	Glu	Pro	Asn	Lys	Cys	Gln	Cys	Gln	Glu	Gly	Trp	His	Gly	Arg	His
				325					330					335	
Cys	Asn	Lys	Arg	Tyr	Glu	Ala	Ser	Leu	Ile	His	Ala	Leu	Arg	Pro	Ala
			340					345					350		
Gly	Ala	Gln	Leu	Arg	Gln	His	Thr	Pro	Ser	Leu	Lys	Lys	Ala	Glu	Glu
		355					360					365			
Arg	Arg	Asp	Pro	Pro	Glu	Ser	Asn	Tyr	Ile	Trp					
	370					375									

<210> SEQ ID NO 5  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 5

agggagcacg gacagtgtgc agatgtggac gagtgtcac tagca

45

<210> SEQ ID NO 6  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 6

agagtgtatc tctggctacg c

21

<210> SEQ ID NO 7  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

-continued

&lt;400&gt; SEQUENCE: 7

taagtccggc acattacagg tc	22
--------------------------	----

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 49

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 8

cccacgatgt atgaatgggt gactttgtgt gactcctggt ttctgcatc	49
---	----

&lt;210&gt; SEQ ID NO 9

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 9

aaagacgcat ctgcgagtgt cc	22
--------------------------	----

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 23

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 10

tgctgatttc acactgctct ccc	23
---------------------------	----

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 2197

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 11

cggacgcgctg ggcgtccggc ggtcgcagag ccaggaggcg gaggcgcgcg gccacgcctg	60
--	----

ggccccagcc cacaccttca ccagggccca ggagccacca tgtggcgatg tccactgggg	120
---	-----

ctactgctgt tgetgccgct ggctggccac ttggctctgg gtgccagca gggtcgtggg	180
--	-----

cgccgggagc tagcaccggg tctgcacctg cggggcatcc gggacgcggg aggccggtac	240
---	-----

tgccaggagc aggacctgtg ctgcccggc cgtgccgacg actgtgccct gccctacctg	300
--	-----

ggcgccatct gttactgtga cctcttctgc aaccgcacgg tctccgactg ctgccctgac	360
---	-----

ttctgggact tctgcctcgg cgtgccaccc ccttttcccc cgatccaagg atgtatgcat	420
---	-----

ggaggtcgta tctatccagt cttgggaacg tactgggaca actgtaaccg ttgcacctgc	480
---	-----

caggagaaca ggcatggca tgggtgatcc agacatgac aaagccatca accagggcaa	540
---	-----

ctatggctgg caggctggga accacagcgc cttctggggc atgacctgg atgaggcat	600
---	-----

tcgctaccgc ctgggccaca tccgccatc ttcctcggtc atgaacatgc atgaaattta	660
--	-----

tacagtgtg aaccagggg aggtgcttcc cacagccttc gaggcctctg agaagtggcc	720
---	-----

caacctgatt catgagcctc ttgaccaagg caactgtgca ggctcctgg ccttctccac	780
--	-----

-continued

```

agcagctgtg gcatccgata gtgtctcaat ccattctctg ggacacatga cgctgtcct 840
gtcgccccag aacctgtgtt cttgtgacac ccaccagcag cagggctgcc gcggtggcg 900
tctcgatggt gcctggtggt tcctgcgtcg cagaggggtg gtgtctgacc actgctaccc 960
cttctcgggc cgtgaacgag acgaggtctg cctgcgccc cctgtatga tgcacagccg 1020
agccatgggt cggggcaagc gccagggcac tgcccactgc cccaacagct atgttaataa 1080
caatgacatc taccaggta ctcctgtcta ccgctcggc tccaacgaca aggagatcat 1140
gaaggagctg atggagaatg gccctgtcca agccctcatg gaggtgcatg aggacttctt 1200
cctatacaag ggagcatct acagccacac gccagtgagc cttgggaggc cagagagata 1260
ccgcccggat gggaccact cagtcaagat cacaggatgg ggagaggaga cgctgcaga 1320
tggaaggagc ctcaaaact ggactcggc caactcctgg gcccagcct gggcgagag 1380
gggccacttc cgcactgtgc gcggcgtcaa tgagtgcgac atcgagagct tcgtgctggg 1440
cgtctggggc cgcgtgggca tggaggacat gggtcacac tgaggctgcg ggcaccacgc 1500
gggttcggc ctgggatcca ggctaagggc cggcgggaga gcccacaatg gggcggtagc 1560
cccagcctcg cccgacagag cccggggcgc agggggcgc cagggcgcta atcccggcgc 1620
gggttccgct gacgcagcgc cccgctggg agccgcgggc aggcgagact ggcggagccc 1680
ccagacctcc cagtggggac ggggcagggc ctggcctggg aagagcacag ctgcagatcc 1740
caggcctctg gcgccccac tcaagactac caaagccagg acaoctcaag tctccagccc 1800
caatacccca cccaatccc gtattctttt ttttttttt ttagacaggg tcttgcctcg 1860
ttgcccaggt tggagtgcag tggccatca gggctcactg taacctccga ctctggggtt 1920
caagtgacc tcccacctca gcctctcaag tagctgggac tacagtgca ccaccacacc 1980
tggctaattt ttgtattttt tgtaaaggg ggggtctcac tgtgttgccc aggctggttt 2040
cgaactcctg ggtcaagcg gtccacctgc ctccgcctcc caaagtgtg ggattgcagg 2100
catgagccac tgcaccagc cctgtattct tattcttcag atatttattt ttcttttcac 2160
tgttttaaaa taaaaccaa gtattgataa aaaaaaa 2197
    
```

```

<210> SEQ ID NO 12
<211> LENGTH: 164
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 12

```

Met Trp Arg Cys Pro Leu Gly Leu Leu Leu Leu Leu Pro Leu Ala Gly
 1           5           10          15
His Leu Ala Leu Gly Ala Gln Gln Gly Arg Gly Arg Arg Glu Leu Ala
 20          25          30
Pro Gly Leu His Leu Arg Gly Ile Arg Asp Ala Gly Gly Arg Tyr Cys
 35          40          45
Gln Glu Gln Asp Leu Cys Cys Arg Gly Arg Ala Asp Asp Cys Ala Leu
 50          55          60
Pro Tyr Leu Gly Ala Ile Cys Tyr Cys Asp Leu Phe Cys Asn Arg Thr
 65          70          75          80
Val Ser Asp Cys Cys Pro Asp Phe Trp Asp Phe Cys Leu Gly Val Pro
 85          90          95
Pro Pro Phe Pro Pro Ile Gln Gly Cys Met His Gly Gly Arg Ile Tyr
    
```

-continued

100			105			110									
Pro	Val	Leu	Gly	Thr	Tyr	Trp	Asp	Asn	Cys	Asn	Arg	Cys	Thr	Cys	Gln
		115					120					125			
Glu	Asn	Arg	Gln	Trp	His	Gly	Gly	Ser	Arg	His	Asp	Gln	Ser	His	Gln
	130					135					140				
Pro	Gly	Gln	Leu	Trp	Leu	Ala	Gly	Trp	Glu	Pro	Gln	Arg	Leu	Leu	Gly
	145				150					155					160
His	Asp	Pro	Gly												

<210> SEQ ID NO 13  
 <211> LENGTH: 533  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (33)  
 <223> OTHER INFORMATION: a, t, c or g  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (80)  
 <223> OTHER INFORMATION: a, t, c or g  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (94)  
 <223> OTHER INFORMATION: a, t, c or g  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (144)  
 <223> OTHER INFORMATION: a, t, c or g  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (188)  
 <223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 13

```

aggctccttg gccctttttc cacagcaagc ttntgcnatc ccgattcggt gtctcaaadc      60
caattctctt gggacacatn acgcctgtcc ttngcccca gaacctgctg tctgttacac      120
ccaccagcag cagggtgcc gcgntggcgc tctcgatggt gcctgggtgt tctgcgctcg      180
ccgaggngtg gtgtctgacc actgctaccc cttctcgggc cgtgaacgag acgaggctgg      240
ccctgcgccc ccctgtatga tgcacagccg agccatgggt cggggcaagc gccaggccac      300
tgcccactgc cccaacagct atgttaataa caatgacatc taccagggtca ctctgttcta      360
ccgcctcggc tccaacgaca aggagatcat gaaggagctg atggagaatg gccctgtcca      420
agccctcatg gagggtcatg aggaattctt cctatacaag ggaggcatct acagccacac      480
gccagtgcgc cttgggaggc cagagagata ccgccggcat gggaccact cag          533
    
```

<210> SEQ ID NO 14  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 14

```

ttcgaggcct ctgagaagtg gccc          24
    
```

<210> SEQ ID NO 15  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

-continued

<400> SEQUENCE: 15

ggcggatatct ctctgcctc cc 22

<210> SEQ ID NO 16

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 16

ttctccacag cagctgtggc atccgatcgt gtctcaatcc attctctggg 50

<210> SEQ ID NO 17

<211> LENGTH: 960

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

gctgcttgcc ctgttgatgg caggcttggc cctgcagcca ggcactgcc tgctgtgcta 60  
 ctctcgcaa gccaggtaga gcaacgagga ctgcctgcag gtggagaact gacccagct 120  
 gggggagcag tgctggaccg cgcgatccg cgcagttggc ctctgaccg tcatcagcaa 180  
 aggtcgcagc ttgaactgcy tggatgactc acaggactac tacgtgggca agaagaacat 240  
 cacgtgctgt gacaccgact tgtgcaacgc cagcggggcc catgccctgc agccggctgc 300  
 cgccatcctt gcgctgtccc ctgcactcgg cctgctgctc tggggaccg gccagotata 360  
 ggctctgggg ggccccgctg cagccccacac tgggtgtggt gccccaggcc tctgtgccac 420  
 tcttcacaga cctggcccag tgggagcctg tcttggttcc tgaggcacaat cctaacgcaa 480  
 gtctgacctat gatgtctgc acccctgtcc cccaccctga ccctcccatt gccctotcca 540  
 ggactcccac ccgagcagtc agctctagt acacagatcc gcctgcagat ggcccccca 600  
 accctctctg ctgctgttcc catggcccag cattctccac ccttaaccct gtgctcaggc 660  
 acctcttccc ccaggaagcc ttcctgtccc accccatcta tgacttgagc caggtctggt 720  
 ccgtggtgtc ccccgacccc agcaggggac aggcactcag gagggccag taaaggctga 780  
 gatgaagtgg actgagtaga actggaggac aagagtcgac gtgagttcct gggagtctcc 840  
 agagatgggg cctggaggcc tggaggaagg ggccaggcct cacattcgtg gggctccctg 900  
 aatggcagcc tgagcacagc gtaggccctt aataaacacc tgttgataa gccaaaaaaa 960

<210> SEQ ID NO 18

<211> LENGTH: 189

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Met Thr His Arg Thr Thr Trp Ala Arg Arg Thr Ser Arg Ala Val  
 1 5 10 15  
 Thr Pro Thr Cys Ala Thr Pro Ala Gly Pro Met Pro Cys Ser Arg Leu  
 20 25 30  
 Pro Pro Ser Leu Arg Cys Ser Leu His Ser Ala Cys Cys Ser Gly Asp  
 35 40 45  
 Pro Ala Ser Tyr Arg Leu Trp Gly Ala Pro Leu Gln Pro Thr Leu Gly

-continued

50		55		60															
Val	65	Pro	70	Leu	75	Leu	80	Thr	85	Asp	90	Leu	95	Ala	100	Gln	105	Trp	110
Glu	115	Pro	120	Val	125	Leu	130	Val	135	Pro	140	Glu	145	Ala	150	His	155	Pro	160
Tyr	165	Val	170	Cys	175	Thr	180	Pro	185	Val	190	Pro	195	His	200	Pro	205	Asp	210
Arg	215	Thr	220	Pro	225	Thr	230	Arg	235	Gln	240	Ile	245	Ser	250	Ser	255	Ser	260
Asp	265	Gly	270	Pro	275	Ser	280	Asn	285	Pro	290	Leu	295	Cys	300	Cys	305	Cys	310
Ser	315	Thr	320	Leu	325	Asn	330	Pro	335	Val	340	Leu	345	Arg	350	His	355	Leu	360
Pro	365	Ala	370	His	375	Pro	380	Ile	385	Tyr	390	Asp	395	Leu	400	Ser	405	Gln	410
Pro	415	Ala	420	Pro	425	Ser	430	Arg	435	Gly	440	Gln	445	Ala	450	Leu	455	Arg	460
<210> SEQ ID NO 19 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe <400> SEQUENCE: 19 tgctgtgcta ctctgcaaa gcc 24																			
<210> SEQ ID NO 20 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe <400> SEQUENCE: 20 tgcacaagtc ggtgtcacag cacg 24																			
<210> SEQ ID NO 21 <211> LENGTH: 44 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe <400> SEQUENCE: 21 agcaacgagg actgcctgca ggtggagaac tgcaccagc tggg 44																			
<210> SEQ ID NO 22 <211> LENGTH: 1200 <212> TYPE: DNA <213> ORGANISM: Homo sapiens <400> SEQUENCE: 22 cccacgcgtc cgaacctctc cagcgatggg agccgccgc ctgctgcca acctcactct 60 gtgcttacag ctgctgattc tctgctgtca aactcagtac gtgaggacc agggcgcatt 120																			

-continued

```

gaccgaccag ctgagcaggg ggcagatccg cgagtaccaa ctctacagca ggaccagtgg 180
caagcacgtg caggtcaccg ggcgtcgcac ctccgccacc gccgaggacg gcaacaagtt 240
tgccaagctc atagtggaga cggacacggt tggcagccgg gttcgcacatca aaggggctga 300
gagtgagaag tacatctgta tgaacaagag gggcaagctc atcgggaagc ccagcgggaa 360
gagcaaagac tgcgtgttca cggagatcgt gctggagaac aactatacgg cttccagaa 420
cgcccggcac gagggtggt tcatggcctt cacgcggcag gggcggcccc gccaggcttc 480
ccgcagccgc cagaaccagc gcgagccca cttcatcaag cgctctacc aaggccagct 540
gcccttcccc aaccacgccg agaagcagaa gcagttcgag tttgtgggct ccgccccac 600
ccgcccggacc aagcgcacac ggcggcccc gccctcacg tagtctggga ggcagggggc 660
agcagcccct gggccgctc cccaccctt tcccttctta atccaaggac tgggctgggg 720
tgccgggagg ggagccagat ccccagggga ggacctgag ggcgcggaag catccgagcc 780
cccagctggg aaggggcagg ccggtgcccc aggggcggct ggcacagtgc ccccttcccg 840
gacgggtggc aggccttggg gaggaactga gtgtaccct gatctcaggc caccagcctc 900
tgccggcctc ccagccgggc tcctgaagcc cgctgaaagg tcagcgactg aaggccttgc 960
agacaaccgt ctggaggtgg ctgtcctcaa aatctgctt tcggatctcc ctcagtctgc 1020
ccccagcccc caaactcctc ctggctagac tgtaggaagg gacttttgtt tgtttgtttg 1080
tttcaggaaa aaagaaaggg agagagagga aaatagaggg ttgtccactc ctcacattcc 1140
acgaccagg cctgcacccc accccaact cccagccccg gaataaaacc attttctgctgc 1200

```

```

<210> SEQ ID NO 23
<211> LENGTH: 205
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 23

```

Met Gly Ala Ala Arg Leu Leu Pro Asn Leu Thr Leu Cys Leu Gln Leu
 1           5           10          15
Leu Ile Leu Cys Cys Gln Thr Gln Tyr Val Arg Asp Gln Gly Ala Met
          20           25           30
Thr Asp Gln Leu Ser Arg Arg Gln Ile Arg Glu Tyr Gln Leu Tyr Ser
          35           40           45
Arg Thr Ser Gly Lys His Val Gln Val Thr Gly Arg Arg Ile Ser Ala
          50           55           60
Thr Ala Glu Asp Gly Asn Lys Phe Ala Lys Leu Ile Val Glu Thr Asp
          65           70           75           80
Thr Phe Gly Ser Arg Val Arg Ile Lys Gly Ala Glu Ser Glu Lys Tyr
          85           90           95
Ile Cys Met Asn Lys Arg Gly Lys Leu Ile Gly Lys Pro Ser Gly Lys
          100          105          110
Ser Lys Asp Cys Val Phe Thr Glu Ile Val Leu Glu Asn Asn Tyr Thr
          115          120          125
Ala Phe Gln Asn Ala Arg His Glu Gly Trp Phe Met Ala Phe Thr Arg
          130          135          140
Gln Gly Arg Pro Arg Gln Ala Ser Arg Ser Arg Gln Asn Gln Arg Glu
          145          150          155          160
Ala His Phe Ile Lys Arg Leu Tyr Gln Gly Gln Leu Pro Phe Pro Asn

```

-continued

---

	165		170		175														
His	Ala	Glu	Lys	Gln	Lys	Gln	Phe	Glu	Phe	Val	Gly	Ser	Ala	Pro	Thr				
			180					185					190						
Arg	Arg	Thr	Lys	Arg	Thr	Arg	Arg	Pro	Gln	Pro	Leu	Thr							
			195					200				205							

<210> SEQ ID NO 24  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 24

cagtacgtga gggaccaggg cgccatga 28

<210> SEQ ID NO 25  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 25

ccggtgacct gcacgtgctt gccca 24

<210> SEQ ID NO 26  
 <211> LENGTH: 41  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (21)  
 <223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 26

gcggatctgc cgctgctca nctggtcggt catggcgccc t 41

<210> SEQ ID NO 27  
 <211> LENGTH: 2479  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

acttgccatc acctgttgcc agtgtggaaa aattctccct gttgaatatt ttgcacatgg 60  
 aggacagcag caaagagggc aacacaggct gataagacca gagacagcag ggagattatt 120  
 ttaccatacag ccctcaggac gttccctcta gctggagtgc tggacttcaa cagaacccca 180  
 tccagtcatt ttgattttgc tgtttatatt ttttttcttt ttctttttcc caccacattg 240  
 tatttttatt ccgtacttca gaaatggggc tacagaccac aaagtggccc agccatgggg 300  
 cttttttctc gaagtcttgg cttatcattt ccttggggct ctactcacag gtgtcacaac 360  
 tctctggcctg ccctagtgtg tgccgctgcy acaggaactt tgtctactgt aatgagcgaa 420  
 gcttgaccctc agtgcctctt gggatcccgaggggcgtaac cgtactctac ctccacaaca 480  
 accaaaattaa taatgctgga tttcctgcag aactgcacaa tgtacagtcg gtgcacacgg 540



-continued

---

tctacctgta tggcaaccaa ctggacgaat tccccatgaa ccttccaag aatgtcagag	600
ttctccattt gcaggaaaac aatattcaga ccatttcacg ggctgctctt gccagctct	660
tgaagcttga agagctgcac ctggatgaca actccatata cacagtgggg gtggaagacg	720
gggccttcog ggagctatt agcctcaaat tgttgTTTT gtctaagaat cacctgagca	780
gtgtgcctgt tgggcttctt gtggacttgc aagagctgag agtggatgaa aatcgaattg	840
ctgtcatata cgacatggcc ttccagaatc tcacgagctt ggagcgtctt attgtggacg	900
ggaacctctt gaccaacaag ggtatcggc agggcacctt cagccatctc accaagctca	960
aggaattttc aattgtacgt aattcgtgt cccaccctcc tccgatctc ccaggtacgc	1020
atctgatcag gctctatttg caggacaacc agataaacca cattcctttg acagcctct	1080
caaactctog taagctggaa cggtggata tatccaaca ccaactcgg atgctgactc	1140
aaggggtttt tgataatctc tccaactga agcagctcac tgctcggaa aacccttgg	1200
tttgtgactg cagtattaaa tgggtcacag aatggctcaa atatatocct tcattctca	1260
acgtgcgggg tttcatgtgc caaggtcctg aacaagtccg ggggatggcc gtcagggaa	1320
taaatatgaa tcttttctc tgtcccacca cgacccccg cctgcctctc ttcaccccag	1380
cccccaagta agcttctccg accactcagc ctcccacct ctctattcca aaccctagca	1440
gaagctacac gcctccaact cctaccacat cgaaacttcc cagattcct gactgggatg	1500
gcagagaaa agtgacccca cctatttctg aacggatcca gctctctatc cattttgtga	1560
atgatacttc cattcaagtc agctggctct ctctcttcc cgtgatggca tacaactca	1620
catgggtgaa aatgggccc agtttagtag ggggatcgt tcaggagcgc atagtcagcg	1680
gtgagaagca acacctgagc ctggtaact tagagccccg atccacctat cggatttgtt	1740
tagtgccact ggatgctttt aactaccgog cggtagaaga caccatttgt tcagaggcca	1800
ccacctatgc ctctatctg aacaacggca gcaacacagc gtcagccat gacgagacga	1860
cgtcccacag catgggctcc cctttctcgc tggcgggctt gatcggggc gcggtgat	1920
ttgtgctggt ggtcttgctc agcgtctttt gctggcatat gcacaaaaag gggcgtaca	1980
cctcccagaa gtggaatac aaccggggc ggcggaaga tgattattgc gaggcaggca	2040
ccaagaagga caactccatc ctggagatga cagaaccag ttttcagatc gctccttaa	2100
ataacgatca actccttaa ggagatttca gactgcagcc catttacacc ccaaatggg	2160
gcattaatta cacagactgc catatcccca acaacatgag atactgcaac agcagcgtgc	2220
cagacctgga gactgcatc acgtgacagc cagaggccca gcgttatcaa ggcggacaat	2280
tagactcttg agaacacact cgtgtgtgca cataaagaca cgcagattac atttgataa	2340
tgttacacag atgcatttgt gatttgaat actctgtaatt ttatacggtg tactatata	2400
tgggatttaa aaaaagtgt atcttttcta tttcaagtta attacaaca gttttgtaac	2460
tctttgcttt taaatctt	2479

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 660

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 28

Met Gly Leu Gln Thr Thr Lys Trp Pro Ser His Gly Ala Phe Phe Leu

1

5

10

15

-continued

---

Lys Ser Trp Leu Ile Ile Ser Leu Gly Leu Tyr Ser Gln Val Ser Lys  
 20 25 30  
 Leu Leu Ala Cys Pro Ser Val Cys Arg Cys Asp Arg Asn Phe Val Tyr  
 35 40 45  
 Cys Asn Glu Arg Ser Leu Thr Ser Val Pro Leu Gly Ile Pro Glu Gly  
 50 55 60  
 Val Thr Val Leu Tyr Leu His Asn Asn Gln Ile Asn Asn Ala Gly Phe  
 65 70 75 80  
 Pro Ala Glu Leu His Asn Val Gln Ser Val His Thr Val Tyr Leu Tyr  
 85 90 95  
 Gly Asn Gln Leu Asp Glu Phe Pro Met Asn Leu Pro Lys Asn Val Arg  
 100 105 110  
 Val Leu His Leu Gln Glu Asn Asn Ile Gln Thr Ile Ser Arg Ala Ala  
 115 120 125  
 Leu Ala Gln Leu Leu Lys Leu Glu Glu Leu His Leu Asp Asp Asn Ser  
 130 135 140  
 Ile Ser Thr Val Gly Val Glu Asp Gly Ala Phe Arg Glu Ala Ile Ser  
 145 150 155 160  
 Leu Lys Leu Leu Phe Leu Ser Lys Asn His Leu Ser Ser Val Pro Val  
 165 170 175  
 Gly Leu Pro Val Asp Leu Gln Glu Leu Arg Val Asp Glu Asn Arg Ile  
 180 185 190  
 Ala Val Ile Ser Asp Met Ala Phe Gln Asn Leu Thr Ser Leu Glu Arg  
 195 200 205  
 Leu Ile Val Asp Gly Asn Leu Leu Thr Asn Lys Gly Ile Ala Glu Gly  
 210 215 220  
 Thr Phe Ser His Leu Thr Lys Leu Lys Glu Phe Ser Ile Val Arg Asn  
 225 230 235 240  
 Ser Leu Ser His Pro Pro Pro Asp Leu Pro Gly Thr His Leu Ile Arg  
 245 250 255  
 Leu Tyr Leu Gln Asp Asn Gln Ile Asn His Ile Pro Leu Thr Ala Phe  
 260 265 270  
 Ser Asn Leu Arg Lys Leu Glu Arg Leu Asp Ile Ser Asn Asn Gln Leu  
 275 280 285  
 Arg Met Leu Thr Gln Gly Val Phe Asp Asn Leu Ser Asn Leu Lys Gln  
 290 295 300  
 Leu Thr Ala Arg Asn Asn Pro Trp Phe Cys Asp Cys Ser Ile Lys Trp  
 305 310 315 320  
 Val Thr Glu Trp Leu Lys Tyr Ile Pro Ser Ser Leu Asn Val Arg Gly  
 325 330 335  
 Phe Met Cys Gln Gly Pro Glu Gln Val Arg Gly Met Ala Val Arg Glu  
 340 345 350  
 Leu Asn Met Asn Leu Leu Ser Cys Pro Thr Thr Thr Pro Gly Leu Pro  
 355 360 365  
 Leu Phe Thr Pro Ala Pro Ser Thr Ala Ser Pro Thr Thr Gln Pro Pro  
 370 375 380  
 Thr Leu Ser Ile Pro Asn Pro Ser Arg Ser Tyr Thr Pro Pro Thr Pro  
 385 390 395 400  
 Thr Thr Ser Lys Leu Pro Thr Ile Pro Asp Trp Asp Gly Arg Glu Arg  
 405 410 415

-continued

---

Val Thr Pro Pro Ile Ser Glu Arg Ile Gln Leu Ser Ile His Phe Val  
                   420                                  425                                  430

Asn Asp Thr Ser Ile Gln Val Ser Trp Leu Ser Leu Phe Thr Val Met  
                   435                                  440                                  445

Ala Tyr Lys Leu Thr Trp Val Lys Met Gly His Ser Leu Val Gly Gly  
                   450                                  455                                  460

Ile Val Gln Glu Arg Ile Val Ser Gly Glu Lys Gln His Leu Ser Leu  
  465                                  470                                  475                                  480

Val Asn Leu Glu Pro Arg Ser Thr Tyr Arg Ile Cys Leu Val Pro Leu  
                                   485                                  490                                  495

Asp Ala Phe Asn Tyr Arg Ala Val Glu Asp Thr Ile Cys Ser Glu Ala  
                                   500                                  505                                  510

Thr Thr His Ala Ser Tyr Leu Asn Asn Gly Ser Asn Thr Ala Ser Ser  
                   515                                  520                                  525

His Glu Gln Thr Thr Ser His Ser Met Gly Ser Pro Phe Leu Leu Ala  
   530                                  535                                  540

Gly Leu Ile Gly Gly Ala Val Ile Phe Val Leu Val Val Leu Leu Ser  
  545                                  550                                  555                                  560

Val Phe Cys Trp His Met His Lys Lys Gly Arg Tyr Thr Ser Gln Lys  
                                   565                                  570                                  575

Trp Lys Tyr Asn Arg Gly Arg Arg Lys Asp Asp Tyr Cys Glu Ala Gly  
                                   580                                  585                                  590

Thr Lys Lys Asp Asn Ser Ile Leu Glu Met Thr Glu Thr Ser Phe Gln  
                   595                                  600                                  605

Ile Val Ser Leu Asn Asn Asp Gln Leu Leu Lys Gly Asp Phe Arg Leu  
   610                                  615                                  620

Gln Pro Ile Tyr Thr Pro Asn Gly Gly Ile Asn Tyr Thr Asp Cys His  
  625                                  630                                  635                                  640

Ile Pro Asn Asn Met Arg Tyr Cys Asn Ser Ser Val Pro Asp Leu Glu  
                                   645                                  650                                  655

His Cys His Thr  
                   660

<210> SEQ ID NO 29  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
                                   oligonucleotide probe

<400> SEQUENCE: 29

cggtctacct gtatggcaac c

21

<210> SEQ ID NO 30  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
                                   oligonucleotide probe

<400> SEQUENCE: 30

gcaggacaac cagataaac ac

22

<210> SEQ ID NO 31

-continued

---

```

<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 31
acgcagattt gagaaggctg tc                                     22

<210> SEQ ID NO 32
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 32
ttcacgggct gctcttgccc agctcttgaa gcttgaagag ctgcac                                     46

<210> SEQ ID NO 33
<211> LENGTH: 3449
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33
acttgagca agcggcgggc gcgagagac aggcagaggc agaagctggg gctccgctct                                     60
cgctcccac gagcgatccc cgaggagagc cgcggccctc ggcgaggcga agaggccgac                                     120
gaggaagacc cgggtggctg cgcccctgcc tcgcttccca ggcgccggcg gctgcagcct                                     180
tgcccctctt gctcgccttg aaaatggaaa agatgctcgc aggctgcttt ctgctgatcc                                     240
tcggacagat cgtcctcctc cctgccgagg ccagggagcg gtcacgtggg aggtccatct                                     300
ctaggggac acacgctcgg acccaccgcg agacggcctc tctggagagt tctgtgaga                                     360
acaagcgggc agacctgggt tcatcattg acagctctcg cagtgtcaac acccatgact                                     420
atgcaaaggt caaggagttc atcgtggaca tcttgcaatt cttggacatt ggtcctgatg                                     480
tcacccgagt gggcctgctc caatatggca gcaactgtca gaatgagttc tccctcaaga                                     540
ccttcaagag gaagtccgag gtggagcgtg ctgtcaagag gatgcggcat ctgtccacgg                                     600
gcaccatgac tgggctggcc atccagtatg ccctgaacat cgcattctca gaagcagagg                                     660
gggcccggcc cctgagggag aatgtgccac gggtcataat gatcgtgaca gatgggagac                                     720
ctcaggactc cgtggccgag gtggctgcta aggcacggga cacgggcata ctaatctttg                                     780
ccattggtgt gggccaggta gacttcaaca ccttgaagtc cattgggagt gagccccatg                                     840
aggaccatgt ctctctgtgt gccaatttca gccagattga gacgctgacc tccgtgttcc                                     900
agaagaagtt gtgcacggcc cacatgtgca gcaccctgga gcataactgt gccacttct                                     960
gcatcaacat ccctggctca tacgtctgca ggtgcaaaca aggotacatt ctcaactcgg                                     1020
atcagacgac ttgcagaatc caggatctgt gtgccatgga ggaccacaac tgtgagcagc                                     1080
tctgtgtgaa tgtgccgggc tcctctgtct gccagtgcta cagtggctac gccctggctg                                     1140
aggatgggaa gaggtgtgtg gctgtggact actgtgcctc agaaaaccac ggatgtgaac                                     1200
atgagtggtg aaatgctgat ggctcctacc tttgccagtg ccatgaagga tttgctctta                                     1260
accagatgga aaaaacgtgc acaagatgca actactgtgc actgaacaaa ccgggctgtg                                     1320

```

-continued

---

agcatgagtg	cgccaacatg	gaggagagct	actactgccg	ctgccaccgt	ggctacactc	1380
tggaccccaa	tggcaaaacc	tgacagccag	tggaccactg	tgacacagcag	gaccatggct	1440
gtgagcagct	gtgtctgaac	acggaggatt	ccttcgctctg	ccagtgtctca	gaaggcttcc	1500
tcatcaacga	ggacctcaag	acctgctccc	gggtggatta	ctgctgtctg	agtgaccatg	1560
gttgtaata	ctcctgtgtc	aacatggaca	gatcctttgc	ctgtcagtgt	cctgagggac	1620
acgtgctcog	cagcgatggg	aagacgtgtg	caaaattgga	ctcttgtgct	ctgggggacc	1680
acggttgtga	acattcgtgt	gtaagcagtg	aagattcgtt	tgtgtgccag	tgctttgaag	1740
gttatatact	ccgtgaagat	ggaaaaacct	gcagaaggaa	agatgtctgc	caagctatag	1800
accatggctg	tgaacacatt	tgtgtgaaca	gtgacgactc	atacacgtgc	gagtgttgg	1860
agggattcog	gctcgtgag	gatgggaaac	gctgccgaag	gaaggatgtc	tgcaaatcaa	1920
cccaccatgg	ctgcaaacac	atgtgtgta	ataatgggaa	ttcctacatc	tgcaaatgct	1980
cagagggatt	tgttctagct	gaggacggaa	gacggtgcaa	gaaatgcaact	gaaggcccaa	2040
ttgacctggt	ctttgtgatc	gatggatcca	agagtcttgg	agaagagaat	ttgaggtcog	2100
tgaagcagtt	tgtcactgga	attatagatt	ccttgacaat	ttccccaaa	gccgctcgag	2160
tggggctgct	ccagtattcc	acacaggtcc	acacagagtt	cactctgaga	aacttcaact	2220
cagccaaaga	catgaaaaaa	gcogtggccc	acatgaaata	catgggaaag	ggctctatga	2280
ctgggctggc	cctgaaacac	atgtttgaga	gaagttttac	ccaaggagaa	ggggccaggc	2340
ccctttccac	aagggtgccc	agagcagcca	ttgtgtttcac	cgacggacgg	gctcaggatg	2400
acgtctccga	gtggccagct	aaagccaagg	ccaatggtat	cactatgtat	gctgttgggg	2460
taggaaaagc	cattgaggag	gaactacaag	agattgcctc	tgagcccaca	aacaagcatc	2520
tcttctatgc	cgaagacttc	agcacaatgg	atgagataag	tgaaaaactc	aagaaaggca	2580
tctgtgaagc	tctagaagac	tccagtgaa	gacaggactc	tccagcaggg	gaactgccaa	2640
aaacggtcca	acagccaaca	gaatctgagc	cagtcacat	aaatatccaa	gacctacttt	2700
cctgttctaa	ttttgcagtg	caacacagat	atctgtttga	agaagacaat	cttttacggg	2760
ctacacaaaa	gctttcccat	tcaacaaaac	cttcaggaag	ccctttggaa	gaaaaacacg	2820
atcaatgcaa	atgtgaaaac	cttataatgt	tccagaacct	tgcaaacgaa	gaagtaagaa	2880
aattaacaca	gcgcttagaa	gaaatgacac	agagaatgga	agccctggaa	aatcgctgga	2940
gatacagatg	aagattagaa	atcgcgacac	atttgtagtc	attgtatcac	ggattacaat	3000
gaacgcagtg	cagagcccca	aagctcaggc	tattgtttaa	tcaataatgt	tgtgaagtaa	3060
aacaatcagt	actgagaaac	ctggtttgcc	acagacaaa	gacaagaagt	atacactaac	3120
ttgtataaat	ttatctagga	aaaaaatcct	tcagaattct	aagatgaatt	taccaggtga	3180
gaatgaataa	gctatgcaag	gtattttcta	atatactgtg	gacacaactt	gcttctgcct	3240
catcctgcct	tagtgtgcaa	tctcatttga	ctatacgata	aagtttgac	agtcttactt	3300
ctgtagaaca	ctggccatag	gaaatgctgt	ttttttgtac	tggactttac	cttgatata	3360
gtatatggat	gtatgcataa	aatcatagga	catatgtact	tgtggaacaa	gttggatttt	3420
ttatacaata	ttaaaattca	ccacttcag				3449

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 915

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 34

Met Glu Lys Met Leu Ala Gly Cys Phe Leu Leu Ile Leu Gly Gln Ile  
1 5 10 15  
Val Leu Leu Pro Ala Glu Ala Arg Glu Arg Ser Arg Gly Arg Ser Ile  
20 25 30  
Ser Arg Gly Arg His Ala Arg Thr His Pro Gln Thr Ala Leu Leu Glu  
35 40 45  
Ser Ser Cys Glu Asn Lys Arg Ala Asp Leu Val Phe Ile Ile Asp Ser  
50 55 60  
Ser Arg Ser Val Asn Thr His Asp Tyr Ala Lys Val Lys Glu Phe Ile  
65 70 75 80  
Val Asp Ile Leu Gln Phe Leu Asp Ile Gly Pro Asp Val Thr Arg Val  
85 90 95  
Gly Leu Leu Gln Tyr Gly Ser Thr Val Lys Asn Glu Phe Ser Leu Lys  
100 105 110  
Thr Phe Lys Arg Lys Ser Glu Val Glu Arg Ala Val Lys Arg Met Arg  
115 120 125  
His Leu Ser Thr Gly Thr Met Thr Gly Leu Ala Ile Gln Tyr Ala Leu  
130 135 140  
Asn Ile Ala Phe Ser Glu Ala Glu Gly Ala Arg Pro Leu Arg Glu Asn  
145 150 155 160  
Val Pro Arg Val Ile Met Ile Val Thr Asp Gly Arg Pro Gln Asp Ser  
165 170 175  
Val Ala Glu Val Ala Ala Lys Ala Arg Asp Thr Gly Ile Leu Ile Phe  
180 185 190  
Ala Ile Gly Val Gly Gln Val Asp Phe Asn Thr Leu Lys Ser Ile Gly  
195 200 205  
Ser Glu Pro His Glu Asp His Val Phe Leu Val Ala Asn Phe Ser Gln  
210 215 220  
Ile Glu Thr Leu Thr Ser Val Phe Gln Lys Lys Leu Cys Thr Ala His  
225 230 235 240  
Met Cys Ser Thr Leu Glu His Asn Cys Ala His Phe Cys Ile Asn Ile  
245 250 255  
Pro Gly Ser Tyr Val Cys Arg Cys Lys Gln Gly Tyr Ile Leu Asn Ser  
260 265 270  
Asp Gln Thr Thr Cys Arg Ile Gln Asp Leu Cys Ala Met Glu Asp His  
275 280 285  
Asn Cys Glu Gln Leu Cys Val Asn Val Pro Gly Ser Phe Val Cys Gln  
290 295 300  
Cys Tyr Ser Gly Tyr Ala Leu Ala Glu Asp Gly Lys Arg Cys Val Ala  
305 310 315 320  
Val Asp Tyr Cys Ala Ser Glu Asn His Gly Cys Glu His Glu Cys Val  
325 330 335  
Asn Ala Asp Gly Ser Tyr Leu Cys Gln Cys His Glu Gly Phe Ala Leu  
340 345 350  
Asn Pro Asp Glu Lys Thr Cys Thr Arg Ile Asn Tyr Cys Ala Leu Asn  
355 360 365  
Lys Pro Gly Cys Glu His Glu Cys Val Asn Met Glu Glu Ser Tyr Tyr  
370 375 380

-continued

---

Cys Arg Cys His Arg Gly Tyr Thr Leu Asp Pro Asn Gly Lys Thr Cys  
 385 390 395 400  
 Ser Arg Val Asp His Cys Ala Gln Gln Asp His Gly Cys Glu Gln Leu  
 405 410 415  
 Cys Leu Asn Thr Glu Asp Ser Phe Val Cys Gln Cys Ser Glu Gly Phe  
 420 425 430  
 Leu Ile Asn Glu Asp Leu Lys Thr Cys Ser Arg Val Asp Tyr Cys Leu  
 435 440 445  
 Leu Ser Asp His Gly Cys Glu Tyr Ser Cys Val Asn Met Asp Arg Ser  
 450 455 460  
 Phe Ala Cys Gln Cys Pro Glu Gly His Val Leu Arg Ser Asp Gly Lys  
 465 470 475 480  
 Thr Cys Ala Lys Leu Asp Ser Cys Ala Leu Gly Asp His Gly Cys Glu  
 485 490 495  
 His Ser Cys Val Ser Ser Glu Asp Ser Phe Val Cys Gln Cys Phe Glu  
 500 505 510  
 Gly Tyr Ile Leu Arg Glu Asp Gly Lys Thr Cys Arg Arg Lys Asp Val  
 515 520 525  
 Cys Gln Ala Ile Asp His Gly Cys Glu His Ile Cys Val Asn Ser Asp  
 530 535 540  
 Asp Ser Tyr Thr Cys Glu Cys Leu Glu Gly Phe Arg Leu Ala Glu Asp  
 545 550 555 560  
 Gly Lys Arg Cys Arg Arg Lys Asp Val Cys Lys Ser Thr His His Gly  
 565 570 575  
 Cys Glu His Ile Cys Val Asn Asn Gly Asn Ser Tyr Ile Cys Lys Cys  
 580 585 590  
 Ser Glu Gly Phe Val Leu Ala Glu Asp Gly Arg Arg Cys Lys Lys Cys  
 595 600 605  
 Thr Glu Gly Pro Ile Asp Leu Val Phe Val Ile Asp Gly Ser Lys Ser  
 610 615 620  
 Leu Gly Glu Glu Asn Phe Glu Val Val Lys Gln Phe Val Thr Gly Ile  
 625 630 635 640  
 Ile Asp Ser Leu Thr Ile Ser Pro Lys Ala Ala Arg Val Gly Leu Leu  
 645 650 655  
 Gln Tyr Ser Thr Gln Val His Thr Glu Phe Thr Leu Arg Asn Phe Asn  
 660 665 670  
 Ser Ala Lys Asp Met Lys Lys Ala Val Ala His Met Lys Tyr Met Gly  
 675 680 685  
 Lys Gly Ser Met Thr Gly Leu Ala Leu Lys His Met Phe Glu Arg Ser  
 690 695 700  
 Phe Thr Gln Gly Glu Gly Ala Arg Pro Leu Ser Thr Arg Val Pro Arg  
 705 710 715 720  
 Ala Ala Ile Val Phe Thr Asp Gly Arg Ala Gln Asp Asp Val Ser Glu  
 725 730 735  
 Trp Ala Ser Lys Ala Lys Ala Asn Gly Ile Thr Met Tyr Ala Val Gly  
 740 745 750  
 Val Gly Lys Ala Ile Glu Glu Glu Leu Gln Glu Ile Ala Ser Glu Pro  
 755 760 765  
 Thr Asn Lys His Leu Phe Tyr Ala Glu Asp Phe Ser Thr Met Asp Glu  
 770 775 780  
 Ile Ser Glu Lys Leu Lys Lys Gly Ile Cys Glu Ala Leu Glu Asp Ser

-continued

785	790	795	800
Asp Gly Arg Gln Asp Ser Pro Ala Gly Glu Leu Pro Lys Thr Val Gln	805	810	815
Gln Pro Thr Glu Ser Glu Pro Val Thr Ile Asn Ile Gln Asp Leu Leu	820	825	830
Ser Cys Ser Asn Phe Ala Val Gln His Arg Tyr Leu Phe Glu Glu Asp	835	840	845
Asn Leu Leu Arg Ser Thr Gln Lys Leu Ser His Ser Thr Lys Pro Ser	850	855	860
Gly Ser Pro Leu Glu Glu Lys His Asp Gln Cys Lys Cys Glu Asn Leu	865	870	880
Ile Met Phe Gln Asn Leu Ala Asn Glu Glu Val Arg Lys Leu Thr Gln	885	890	895
Arg Leu Glu Glu Met Thr Gln Arg Met Glu Ala Leu Glu Asn Arg Leu	900	905	910
Arg Tyr Arg	915		
<210> SEQ ID NO 35			
<211> LENGTH: 23			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 35			
gtgaccctgg ttgtgaatac tcc			23
<210> SEQ ID NO 36			
<211> LENGTH: 22			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 36			
acagccatgg tctatagcct gg			22
<210> SEQ ID NO 37			
<211> LENGTH: 45			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 37			
gcctgtcagt gtccagagg acacgtgctc cgcagcgatg ggaag			45
<210> SEQ ID NO 38			
<211> LENGTH: 1813			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 38			
ggagccgcc tgggtgtcag cggtcggct cccgcgcacg ctccggccgt cgcgcagcct			60
cggcacctgc aggtccgtgc gtcccggcg tgggcccct gactccgtcc cggccaggga			120



-continued

```

gggccatgat ttccctccc gggcccctgg tgaccaactt gctgcggttt ttgttctg 180
ggctgagtgc cctcgcgcc ccctcggggg ccagctgca actgcacttg cccgccaacc 240
ggttgacggc ggtggagggg ggggaagtgg tgcttcacg gtggtacacc ttgcacgggg 300
aggtgtcttc atccagcca tgggaggtgc cctttgtgat gtggttcttc aaacagaaag 360
aaaaggagga tcaggtgttg tcctacatca atggggtcac aacaagcaaa cctggagtat 420
ccttggtcta ctccatgcc tcccgaacc tgtccctgcg gctggagggt ctccaggaga 480
aagactctgg cccctacagc tgctcctgta atgtgcaaga caaacaaggc aaatctaggg 540
gccacagcat caaacctta gaactcaatg tactggttcc tccagctcct ccactctgcc 600
gtctccaggg tgtgcccct gtgggggcaa acgtgaccct gagctgccag tctccaagga 660
gtaagcccgc tgtccaatac cagtgggacg ggcagcttcc atccttcag actttctttg 720
caccagcatt agatgtcatc cgtgggtctt taagcctcac caaccttcg tcttccatgg 780
ctggagtcta tgtctgcaag gccacaatg aggtgggac tgccaatgt aatgtgacgc 840
tggaagtgag cacagggcct ggagctgcag tggttgctgg agctgttggt ggtaccctgg 900
ttgactggg gttgctggct gggctggtcc tcttgtaaca ccgccggggc aaggccctgg 960
aggagccagc caatgatatc aaggaggtat ccattgctcc cgggaccctg ccttgccca 1020
agagctcaga cacaatctcc aagaatggga cccttctctc tgtcacctcc gcacgagccc 1080
tccggccacc ccattggcct ccagggcctg gtgcattgac cccacgccc agtctctcca 1140
gccagggcct gccctcacca agactgccca cgacagatgg ggcccaccct caaccaatat 1200
ccccatccc tggtaggggt tcttctctg gcttgagccg catgggtgct gtgcctgtga 1260
tggtgcctgc ccagagtcaa gctggctctc tggatgatg accccaccac tcattggcta 1320
aaggatttgg ggtctctcct tcctataagg gtcacctta gcacagaggc ctgagtcatg 1380
ggaaagagtc aactcctga cccttagtac tctgccccca cctctcttta ctgtgggaaa 1440
accatctcag taagacctaa gtgtccagga gacagaagga gaagaggaa tggatctgga 1500
attgggagga gcctccacc acccctgact cctccttatg aagccagctg ctgaaattag 1560
ctactcacca agagtgaggg gcagagactt ccagtcactg agtctcccag gccccttga 1620
tctgtacccc acccctatct aacaccacc ttggtcccca ctccagctcc ctgtattgat 1680
ataacctgtc aggtggcctt ggtaggttt tactggggca gaggataggg aatctcttat 1740
taaaactaac atgaaatag tgttgttttc atttgcaaat ttaaataaag atacataatg 1800
tttgtatgaa aaa 1813

```

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 390

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 39

```

Met Ile Ser Leu Pro Gly Pro Leu Val Thr Asn Leu Leu Arg Phe Leu
 1           5           10           15

```

```

Phe Leu Gly Leu Ser Ala Leu Ala Pro Pro Ser Arg Ala Gln Leu Gln
 20           25           30

```

```

Leu His Leu Pro Ala Asn Arg Leu Gln Ala Val Glu Gly Gly Glu Val
 35           40           45

```

-continued

---

Val Leu Pro Ala Trp Tyr Thr Leu His Gly Glu Val Ser Ser Ser Gln  
 50 55 60  
 Pro Trp Glu Val Pro Phe Val Met Trp Phe Phe Lys Gln Lys Glu Lys  
 65 70 75 80  
 Glu Asp Gln Val Leu Ser Tyr Ile Asn Gly Val Thr Thr Ser Lys Pro  
 85 90 95  
 Gly Val Ser Leu Val Tyr Ser Met Pro Ser Arg Asn Leu Ser Leu Arg  
 100 105 110  
 Leu Glu Gly Leu Gln Glu Lys Asp Ser Gly Pro Tyr Ser Cys Ser Val  
 115 120 125  
 Asn Val Gln Asp Lys Gln Gly Lys Ser Arg Gly His Ser Ile Lys Thr  
 130 135 140  
 Leu Glu Leu Asn Val Leu Val Pro Pro Ala Pro Pro Ser Cys Arg Leu  
 145 150 155 160  
 Gln Gly Val Pro His Val Gly Ala Asn Val Thr Leu Ser Cys Gln Ser  
 165 170 175  
 Pro Arg Ser Lys Pro Ala Val Gln Tyr Gln Trp Asp Arg Gln Leu Pro  
 180 185 190  
 Ser Phe Gln Thr Phe Phe Ala Pro Ala Leu Asp Val Ile Arg Gly Ser  
 195 200 205  
 Leu Ser Leu Thr Asn Leu Ser Ser Ser Met Ala Gly Val Tyr Val Cys  
 210 215 220  
 Lys Ala His Asn Glu Val Gly Thr Ala Gln Cys Asn Val Thr Leu Glu  
 225 230 235 240  
 Val Ser Thr Gly Pro Gly Ala Ala Val Val Ala Gly Ala Val Val Gly  
 245 250 255  
 Thr Leu Val Gly Leu Gly Leu Leu Ala Gly Leu Val Leu Leu Tyr His  
 260 265 270  
 Arg Arg Gly Lys Ala Leu Glu Glu Pro Ala Asn Asp Ile Lys Glu Asp  
 275 280 285  
 Ala Ile Ala Pro Arg Thr Leu Pro Trp Pro Lys Ser Ser Asp Thr Ile  
 290 295 300  
 Ser Lys Asn Gly Thr Leu Ser Ser Val Thr Ser Ala Arg Ala Leu Arg  
 305 310 315 320  
 Pro Pro His Gly Pro Pro Arg Pro Gly Ala Leu Thr Pro Thr Pro Ser  
 325 330 335  
 Leu Ser Ser Gln Ala Leu Pro Ser Pro Arg Leu Pro Thr Thr Asp Gly  
 340 345 350  
 Ala His Pro Gln Pro Ile Ser Pro Ile Pro Gly Gly Val Ser Ser Ser  
 355 360 365  
 Gly Leu Ser Arg Met Gly Ala Val Pro Val Met Val Pro Ala Gln Ser  
 370 375 380  
 Gln Ala Gly Ser Leu Val  
 385 390

&lt;210&gt; SEQ ID NO 40

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 40

---

-continued

---

agggtctcca ggagaaagac tc 22

<210> SEQ ID NO 41  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 41

attgtgggcc ttgcagacat agac 24

<210> SEQ ID NO 42  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 42

ggccacagca tcaaaacctt agaactcaat gtactggttc ctccagctcc 50

<210> SEQ ID NO 43  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 43

gtgtgacaca gcgtgggc 18

<210> SEQ ID NO 44  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 44

gaccggcagg cttctgcg 18

<210> SEQ ID NO 45  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 45

cagcagcttc agccaccagg agtgg 25

<210> SEQ ID NO 46  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

-continued

&lt;400&gt; SEQUENCE: 46

ctgagccgtg ggctgcagtc tcgc 24

&lt;210&gt; SEQ ID NO 47

&lt;211&gt; LENGTH: 45

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 47

ccgactacga ctggttcttc atcatgcagg atgacacata tgtgc 45

&lt;210&gt; SEQ ID NO 48

&lt;211&gt; LENGTH: 2822

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 48

cgccaccact gcggccaccg ccaatgaaac gcctcccgt cctagtgggt ttttccactt 60

tgttgaattg ttctataact caaaattgca ccaagacacc ttgtctocca aatgcaaaat 120

gtgaaatacg caatggaatt gaagcctgct attgcaacat gggattttca gaaatgggtg 180

tcacaatttg tgaagatgat aatgaatgtg gaaatttaac tcagtcctgt ggcgaaaatg 240

ctaattgcac taacacagaa ggaagttatt attgtatgtg tgtacctggc ttcagatcca 300

gcagtaacca agacaggttt atcactaatg atggaaccgt ctgtatagaa aatgtgaatg 360

caaactgccca tttagataat gtctgtatag ctgcaaatat taataaaaact ttaacaaaaa 420

tcagatccat aaaagaacct gtggctttgc tacaagaagt ctatagaaat tctgtgacag 480

atctttcacc aacagatata attacatata tagaaatatt agctgaatca tcttcattac 540

taggttacaa gaacaacact atctcagcca aggacacct ttctaactca actcttactg 600

aatttgtaaa aaccgtgaat aattttgttc aaagggatac attttagtct tgggacaagt 660

tatctgtgaa tcataggaga acacatctta caaaactcat gcacactgtt gaacaagcta 720

ctttaaggat atcccagagc ttccaaaaga ccacagagtt tgatacaaat tcaacggata 780

tagctctcaa agttttcttt ttgtattcat ataacatgaa acatattcat cctcatatga 840

atatggatgg agactacata aatataattc caaagagaaa agctgcatat gattcaaatg 900

gcaatgttgc agttgcattt ttatattata agagtattgg tcctttgtct tcatcatctg 960

acaacttctt attgaaacct caaaattatg ataattctga agaggaggaa agagtcatat 1020

cttcagtaat ttcagctca atgagctcaa acccaccac attatatgaa cttgaaaaaa 1080

taacatttac attaagtcac cgaaaggcca cagataggta taggagtcta tgtgcatttt 1140

ggaattactc acctgatacc atgaatggca gctggctctc agagggtctg gagctgacat 1200

actcaaatga gaccacaccc tcatgccgct gtaatcacct gacacatttt gcaattttga 1260

tgtcctctg tccttcattt ggtattaaag attataatat tcttacaagg atcactcaac 1320

taggaataat tatttactg atttgtcttg coatatgcat ttttaccttc tggttottca 1380

gtgaaattca aagcaccagg acaacaatc acaaaaatct ttgctgtagc ctatttcttg 1440

ctgaacttgt ttttcttctt gggatcaata caaatactaa taagctcttc tgttcaatca 1500

-continued

---

```

ttgccggact gctacactac ttcttttttag ctgcttttgc atggatgtgc attgaaggca 1560
tacatctcta tctcattggt gtgggtgtca tctacaacia gggatttttg cacaagaatt 1620
tttatatctt tggctatcta agcccagccg tggtagttgg attttcggca gcactaggat 1680
acagatatta tggcacaacc aaagtatggt ggcttagcac cgaacaacac tttatttggga 1740
gttttatagg accagcatgc ctaatcattc ttgttaatct cttggctttt ggagtcacca 1800
tatacaaaagt ttttcgtcac actgcagggt tgaaccaga agttagtgc tttgagaaca 1860
taaggtcttg tgcaagagga gcctcgcctc ttctgttctc tctcggcacc acctggatct 1920
ttggggttct ccatgtttgtg cacgcatcag tggttacagc ttacctcttc acagtcagca 1980
atgctttcca ggggatgttc atttttttat tctgtgtgtt tttatctaga aagattcaag 2040
aagaatatta cagattgttc aaaaatgtcc cctgttgttt tggatgttta aggtaaacat 2100
agagaatggt ggataattac aactgcacaa aaataaaaat tccaagctgt ggatgaccaa 2160
tgtataaaaa tgactcatca aattatccaa ttattaacta ctagacaaaa agtattttaa 2220
atcagttttt ctgtttatgc tataggaact gtagataata aggtaaaatt atgtatcata 2280
tagatatact atgtttttct atgtgaaata gttctgtcaa aaatagtatt gcagatattt 2340
ggaaagtaat tggttttctca ggagtgatat cactgcaccc aaggaaagat tttctttcta 2400
acacgagaag tatatgaatg tcttgaagga aacctctggc ttgatatttc tgtgactcgt 2460
gttgcccttg aaactagtcc cctaccacct cggtaatgag ctccattaca gaaagtggaa 2520
cataagagaa tgaaggggca gaatatcaaa cagtgaaaag ggaatgataa gatgtatttt 2580
gaatgaactg ttttttctgt agactagctg agaaattggt gacataaaat aaagaattga 2640
agaaacacat tttaccattt tgtgaattgt tctgaactta aatgtccact aaaacaactt 2700
agacttctgt ttgctaaatc tgtttctttt tctaataatc taaaaaaaaa aaaaagggtt 2760
acctccacaa attgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2820
aa

```

&lt;210&gt; SEQ ID NO 49

&lt;211&gt; LENGTH: 690

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 49

```

Met Lys Arg Leu Pro Leu Leu Val Val Phe Ser Thr Leu Leu Asn Cys
 1           5           10           15
Ser Tyr Thr Gln Asn Cys Thr Lys Thr Pro Cys Leu Pro Asn Ala Lys
 20           25           30
Cys Glu Ile Arg Asn Gly Ile Glu Ala Cys Tyr Cys Asn Met Gly Phe
 35           40           45
Ser Gly Asn Gly Val Thr Ile Cys Glu Asp Asp Asn Glu Cys Gly Asn
 50           55           60
Leu Thr Gln Ser Cys Gly Glu Asn Ala Asn Cys Thr Asn Thr Glu Gly
 65           70           75           80
Ser Tyr Tyr Cys Met Cys Val Pro Gly Phe Arg Ser Ser Ser Asn Gln
 85           90           95
Asp Arg Phe Ile Thr Asn Asp Gly Thr Val Cys Ile Glu Asn Val Asn
 100          105          110
Ala Asn Cys His Leu Asp Asn Val Cys Ile Ala Ala Asn Ile Asn Lys

```

-continued

115					120					125					
Thr	Leu	Thr	Lys	Ile	Arg	Ser	Ile	Lys	Glu	Pro	Val	Ala	Leu	Leu	Gln
130						135					140				
Glu	Val	Tyr	Arg	Asn	Ser	Val	Thr	Asp	Leu	Ser	Pro	Thr	Asp	Ile	Ile
145					150					155					160
Thr	Tyr	Ile	Glu	Ile	Leu	Ala	Glu	Ser	Ser	Ser	Leu	Leu	Gly	Tyr	Lys
				165					170					175	
Asn	Asn	Thr	Ile	Ser	Ala	Lys	Asp	Thr	Leu	Ser	Asn	Ser	Thr	Leu	Thr
			180					185					190		
Glu	Phe	Val	Lys	Thr	Val	Asn	Asn	Phe	Val	Gln	Arg	Asp	Thr	Phe	Val
		195					200					205			
Val	Trp	Asp	Lys	Leu	Ser	Val	Asn	His	Arg	Arg	Thr	His	Leu	Thr	Lys
	210					215					220				
Leu	Met	His	Thr	Val	Glu	Gln	Ala	Thr	Leu	Arg	Ile	Ser	Gln	Ser	Phe
225					230					235					240
Gln	Lys	Thr	Thr	Glu	Phe	Asp	Thr	Asn	Ser	Thr	Asp	Ile	Ala	Leu	Lys
				245					250					255	
Val	Phe	Phe	Phe	Asp	Ser	Tyr	Asn	Met	Lys	His	Ile	His	Pro	His	Met
			260					265					270		
Asn	Met	Asp	Gly	Asp	Tyr	Ile	Asn	Ile	Phe	Pro	Lys	Arg	Lys	Ala	Ala
		275					280					285			
Tyr	Asp	Ser	Asn	Gly	Asn	Val	Ala	Val	Ala	Phe	Leu	Tyr	Tyr	Lys	Ser
	290					295					300				
Ile	Gly	Pro	Leu	Leu	Ser	Ser	Ser	Asp	Asn	Phe	Leu	Leu	Lys	Pro	Gln
305					310					315					320
Asn	Tyr	Asp	Asn	Ser	Glu	Glu	Glu	Glu	Arg	Val	Ile	Ser	Ser	Val	Ile
				325					330					335	
Ser	Val	Ser	Met	Ser	Ser	Asn	Pro	Pro	Thr	Leu	Tyr	Glu	Leu	Glu	Lys
			340					345					350		
Ile	Thr	Phe	Thr	Leu	Ser	His	Arg	Lys	Val	Thr	Asp	Arg	Tyr	Arg	Ser
		355					360					365			
Leu	Cys	Ala	Phe	Trp	Asn	Tyr	Ser	Pro	Asp	Thr	Met	Asn	Gly	Ser	Trp
	370					375					380				
Ser	Ser	Glu	Gly	Cys	Glu	Leu	Thr	Tyr	Ser	Asn	Glu	Thr	His	Thr	Ser
385					390					395					400
Cys	Arg	Cys	Asn	His	Leu	Thr	His	Phe	Ala	Ile	Leu	Met	Ser	Ser	Gly
				405					410					415	
Pro	Ser	Ile	Gly	Ile	Lys	Asp	Tyr	Asn	Ile	Leu	Thr	Arg	Ile	Thr	Gln
			420					425					430		
Leu	Gly	Ile	Ile	Ile	Ser	Leu	Ile	Cys	Leu	Ala	Ile	Cys	Ile	Phe	Thr
	435						440					445			
Phe	Trp	Phe	Phe	Ser	Glu	Ile	Gln	Ser	Thr	Arg	Thr	Thr	Ile	His	Lys
	450					455						460			
Asn	Leu	Cys	Cys	Ser	Leu	Phe	Leu	Ala	Glu	Leu	Val	Phe	Leu	Val	Gly
465					470					475					480
Ile	Asn	Thr	Asn	Thr	Asn	Lys	Leu	Phe	Cys	Ser	Ile	Ile	Ala	Gly	Leu
				485					490					495	
Leu	His	Tyr	Phe	Phe	Leu	Ala	Ala	Phe	Ala	Trp	Met	Cys	Ile	Glu	Gly
			500					505					510		
Ile	His	Leu	Tyr	Leu	Ile	Val	Val	Gly	Val	Ile	Tyr	Asn	Lys	Gly	Phe
		515					520					525			

-continued

Leu His Lys Asn Phe Tyr Ile Phe Gly Tyr Leu Ser Pro Ala Val Val  
 530 535 540  
 Val Gly Phe Ser Ala Ala Leu Gly Tyr Arg Tyr Tyr Gly Thr Thr Lys  
 545 550 555 560  
 Val Cys Trp Leu Ser Thr Glu Asn Asn Phe Ile Trp Ser Phe Ile Gly  
 565 570 575  
 Pro Ala Cys Leu Ile Ile Leu Val Asn Leu Leu Ala Phe Gly Val Ile  
 580 585 590  
 Ile Tyr Lys Val Phe Arg His Thr Ala Gly Leu Lys Pro Glu Val Ser  
 595 600 605  
 Cys Phe Glu Asn Ile Arg Ser Cys Ala Arg Gly Ala Leu Ala Leu Leu  
 610 615 620  
 Phe Leu Leu Gly Thr Thr Trp Ile Phe Gly Val Leu His Val Val His  
 625 630 635 640  
 Ala Ser Val Val Thr Ala Tyr Leu Phe Thr Val Ser Asn Ala Phe Gln  
 645 650 655  
 Gly Met Phe Ile Phe Leu Phe Leu Cys Val Leu Ser Arg Lys Ile Gln  
 660 665 670  
 Glu Glu Tyr Tyr Arg Leu Phe Lys Asn Val Pro Cys Cys Phe Gly Cys  
 675 680 685  
 Leu Arg  
 690

<210> SEQ ID NO 50  
 <211> LENGTH: 589  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (61)  
 <223> OTHER INFORMATION: a, t, c or g  
 <400> SEQUENCE: 50  
 tggaaacata tcctccctca tatgaatag gatggagact acataaatat atttccaaag 60  
 ngaaaagccg gcatatggat tcaaatggca atgttgacgt tgcattttta tattataaga 120  
 gtattgtgcc cttgtctttc atcatctgac aacttcttat tgaaacctca aaattatgat 180  
 aattctgaag aggaggaaag agtcatatct tcagtaattt cagtctcaat gagctcaaac 240  
 ccaccacat tatatgaact tgaaaaata acatttcat taagtcatcg aaaggtcaca 300  
 gataggtata ggagtctatg tggcattttg gaatactcac ctgataccat gaatggcagc 360  
 tggctctcag agggctgtga gctgacatac tcaaatgaga cccacacctc atgccgctgt 420  
 aatcacctga cacattttgc aattttgatg tcctctggtc cttccattgg tattaagat 480  
 tataatattc ttacaaggat cactcaacta ggaataatta ttccactgat ttgtcttgcc 540  
 atatgcattt ttacctctctg gttcttcagt gaaattcaaa gcaccagga 589

<210> SEQ ID NO 51  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe  
 <400> SEQUENCE: 51

---

-continued

---

ggtaatgagc tccattacag 20

<210> SEQ ID NO 52  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 52

ggagtagaaa gcgcatgg 18

<210> SEQ ID NO 53  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 53

cacctgatac catgaatggc ag 22

<210> SEQ ID NO 54  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 54

cgagctcgaa ttaattcg 18

<210> SEQ ID NO 55  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 55

ggatctcctg agctcagg 18

<210> SEQ ID NO 56  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 56

cctagttgag tgatccttgt aag 23

<210> SEQ ID NO 57  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe



-continued

&lt;400&gt; SEQUENCE: 57

atgagaccaca cacctcatgc cgctgtaatc acctgacaca ttttgaatt 50

&lt;210&gt; SEQ ID NO 58

&lt;211&gt; LENGTH: 2137

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 58

gctcccagcc aagaacctcg gggccgctgc gcggtgggga ggagttccc gaaaccggc 60

cgctaagcga ggcctcctcc tcccgcagat cogaacggcc tgggcggggt caccocggct 120

gggacaagaa gccgcccct gcctgcccgg gcccggggag ggggctgggg ctggggccgg 180

aggcggggtg tgagtgggtg tgtgcgggg gcggaggctt gatgcaatcc cgataagaaa 240

tgctcgggtg tcttggggcac ctaccctgg ggcccgtaa gcgctactat ataaggctgc 300

cggcccggag ccgcccgcgc gtcagagcag gagcgctgcg tccaggatct agggccacga 360

ccatcccaac ccggcactca cagccccga gcgcattccc gtcgcccgc agcctcccgc 420

acccccatcg ccggagctgc gccgagagcc ccagggaggt gccatgcgga gcgggtgtgt 480

ggtgttccac gtatggatcc tggcccgcct ctggctggcc gtggcccggc gcccccctgc 540

cttctcggac gcggggcccc acgtgcaacta cggetggggc gaccccatcc gcttgcggca 600

cctgtacacc tccggccccc acgggctctc cagctgcttc ctgogcatcc gtgccgacgg 660

cgtcgtggac tgcgcgcggg gccagagcgc gcacagtttg ctggagatca aggcagtcgc 720

tctcgggacc gtggccatca agggcgtgca cagcgtgcgg tacctctgca tgggcgcga 780

cggcaagatg caggggctgc ttcagtactc ggaggaagac tgtgctttcg aggaggagat 840

ccgcccagat ggctacaatg tgtaccgatc cgagaagcac cgccctcccg tctccctgag 900

cagtgccaaa cagcggcagc tgtacaagaa cagaggcttt cttccactct ctcattttcct 960

gcccattgct cccattgtcc cagaggagcc tgaggacctc aggggcccact tggaaatctga 1020

catgtttctc tcgcccctgg agaccgacag catggaccca tttgggcttg tcaccggact 1080

ggaggccgtg aggagtccca gctttgagaa gtaactgaga ccatgcccgg gcctcttcac 1140

tgctgccagg ggctgtggta cctgcagcgt gggggacgtg cttctacaag aacagtcctg 1200

agtccacggt ctgtttagct ttaggaagaa acatctagaa gttgtacata ttcagagttt 1260

tccattggca gtgccagttt ctagccaata gacttgtctg atcataacat tgtaagcctg 1320

tagcttgccc agctgctgcc tgggcccaca ttctgctccc tcgaggttgc tggacaagct 1380

gctgcactgt ctcagttctg cttgaatacc tccatcgatg gggaaactcac ttcctttgga 1440

aaaattctta tgtcaagctg aaattctcta atttttctc atcacttccc caggagcagc 1500

cagaagacag gcagtagttt taatttcagg aacaggatgat ccaactctgta aaacagcagg 1560

taaattttcac tcaaccocat gtgggaattg atctatatct ctacttccag ggaccatttg 1620

cccttcccaa atccctccag gccagaactg actggagcag gcatggccca ccaggttca 1680

ggagtagggg aagcctggag cccactcca gccctgggac aacttgagaa ttcccctga 1740

ggccagttct gtcatggatg ctgtcctgag aataactgac tgtcccgtg tcacctgctt 1800

ccatctccca gccaccagc cctctgcca cctcacatgc ctcccattgg attggggcct 1860

cccaggcccc ccaccttatg tcaacctgca cttctgttcc aaaaatcagg aaaagaaaag 1920

-continued

---

```

atttgaagac cccaagtctt gtcaataact tgctgtgtgg aagcagcggg ggaagaccta 1980
gaaccctttc cccagcactt ggttttccaa catgatattt atgagtaatt tattttgata 2040
tgtacatctc ttattttctt acattattta tgcccccaaa ttatatttat gtatgtaagt 2100
gaggtttggt ttgtatatta aaatggagtt tgtttgt 2137

```

```

<210> SEQ ID NO 59
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 59

```

```

Met Arg Ser Gly Cys Val Val Val His Val Trp Ile Leu Ala Gly Leu
 1           5           10           15
Trp Leu Ala Val Ala Gly Arg Pro Leu Ala Phe Ser Asp Ala Gly Pro
          20           25           30
His Val His Tyr Gly Trp Gly Asp Pro Ile Arg Leu Arg His Leu Tyr
      35           40           45
Thr Ser Gly Pro His Gly Leu Ser Ser Cys Phe Leu Arg Ile Arg Ala
      50           55           60
Asp Gly Val Val Asp Cys Ala Arg Gly Gln Ser Ala His Ser Leu Leu
      65           70           75           80
Glu Ile Lys Ala Val Ala Leu Arg Thr Val Ala Ile Lys Gly Val His
          85           90           95
Ser Val Arg Tyr Leu Cys Met Gly Ala Asp Gly Lys Met Gln Gly Leu
          100          105          110
Leu Gln Tyr Ser Glu Glu Asp Cys Ala Phe Glu Glu Glu Ile Arg Pro
          115          120          125
Asp Gly Tyr Asn Val Tyr Arg Ser Glu Lys His Arg Leu Pro Val Ser
          130          135          140
Leu Ser Ser Ala Lys Gln Arg Gln Leu Tyr Lys Asn Arg Gly Phe Leu
          145          150          155          160
Pro Leu Ser His Phe Leu Pro Met Leu Pro Met Val Pro Glu Glu Pro
          165          170          175
Glu Asp Leu Arg Gly His Leu Glu Ser Asp Met Phe Ser Ser Pro Leu
          180          185          190
Glu Thr Asp Ser Met Asp Pro Phe Gly Leu Val Thr Gly Leu Glu Ala
          195          200          205
Val Arg Ser Pro Ser Phe Glu Lys
          210          215

```

```

<210> SEQ ID NO 60
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

```

```

<400> SEQUENCE: 60

```

```

atccgccag atggtacaa tgtgta 26

```

```

<210> SEQ ID NO 61
<211> LENGTH: 42
<212> TYPE: DNA

```



-continued

&lt;211&gt; LENGTH: 312

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 64

```

Met Ala Arg Arg Ser Arg His Arg Leu Leu Leu Leu Leu Arg Tyr
 1           5           10           15
Leu Val Val Ala Leu Gly Tyr His Lys Ala Tyr Gly Phe Ser Ala Pro
          20           25           30
Lys Asp Gln Gln Val Val Thr Ala Val Glu Tyr Gln Glu Ala Ile Leu
          35           40           45
Ala Cys Lys Thr Pro Lys Lys Thr Val Ser Ser Arg Leu Glu Trp Lys
          50           55           60
Lys Leu Gly Arg Ser Val Ser Phe Val Tyr Tyr Gln Gln Thr Leu Gln
          65           70           75           80
Gly Asp Phe Lys Asn Arg Ala Glu Met Ile Asp Phe Asn Ile Arg Ile
          85           90           95
Lys Asn Val Thr Arg Ser Asp Ala Gly Lys Tyr Arg Cys Glu Val Ser
          100          105          110
Ala Pro Ser Glu Gln Gly Gln Asn Leu Glu Glu Asp Thr Val Thr Leu
          115          120          125
Glu Val Leu Val Ala Pro Ala Val Pro Ser Cys Glu Val Pro Ser Ser
          130          135          140
Ala Leu Ser Gly Thr Val Val Glu Leu Arg Cys Gln Asp Lys Glu Gly
          145          150          155          160
Asn Pro Ala Pro Glu Tyr Thr Trp Phe Lys Asp Gly Ile Arg Leu Leu
          165          170          175
Glu Asn Pro Arg Leu Gly Ser Gln Ser Thr Asn Ser Ser Tyr Thr Met
          180          185          190
Asn Thr Lys Thr Gly Thr Leu Gln Phe Asn Thr Val Ser Lys Leu Asp
          195          200          205
Thr Gly Glu Tyr Ser Cys Glu Ala Arg Asn Ser Val Gly Tyr Arg Arg
          210          215          220
Cys Pro Gly Lys Arg Met Gln Val Asp Asp Leu Asn Ile Ser Gly Ile
          225          230          235          240
Ile Ala Ala Val Val Val Val Ala Leu Val Ile Ser Val Cys Gly Leu
          245          250          255
Gly Val Cys Tyr Ala Gln Arg Lys Gly Tyr Phe Ser Lys Glu Thr Ser
          260          265          270
Phe Gln Lys Ser Asn Ser Ser Ser Lys Ala Thr Thr Met Ser Glu Asn
          275          280          285
Val Gln Trp Leu Thr Pro Val Ile Pro Ala Leu Trp Lys Ala Ala Ala
          290          295          300
Gly Gly Ser Arg Gly Gln Glu Phe
          305          310

```

&lt;210&gt; SEQ ID NO 65

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

&lt;400&gt; SEQUENCE: 65

-continued

---

atcgttgga agttagtgcc cc 22

<210> SEQ ID NO 66  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 66

acctgcgata tccaacagaa ttg 23

<210> SEQ ID NO 67  
 <211> LENGTH: 48  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 67

ggaagaggat acagtcactc tggaagtatt agtggctcca gcagttcc 48

<210> SEQ ID NO 68  
 <211> LENGTH: 2639  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

gacatcggag gtgggctagc actgaaactg cttttcaaga cgaggaagag gaggagaaaag 60  
 agaaagaaga ggaagatggt gggcaacatt tatttaacat gctccacagc ccggaccctg 120  
 gcatcatgct gctattcctg caaatactga agaagcatgg gatttaaata ttttacttct 180  
 aaataaatga attactcaat ctccatgac catctataca tactccacct tcaaaaagta 240  
 catcaatatt atatcattaa ggaatagta accttctctt ctccaatatg catgacattt 300  
 ttggacaatg caattgtggc actggcactt atttcagtga agaaaaactt tgtggttcta 360  
 tggcattcat catttgacaa atgcaagcat cttccttacc aatcagctcc tattgaactt 420  
 actagcactg actgtggaat ccttaagggc ccattacatt tctgaagaag aaagctaaga 480  
 tgaaggacat gccactccga attcatgtgc tacttggcct agctatcact aactagtagc 540  
 aagctgtaga taaaaaagtg gattgtccac ggttatgtac gtgtgaaatc aggccttggt 600  
 ttacaccagc atccatttat atggaagcat ctacagtgga ttgtaatgat ttaggtcttt 660  
 taactttccc agccagattg ccagctaaca cacagattct tctcctacag actaacaata 720  
 ttgcaaaaat tgaatactcc acagacttcc cagtaaacct tactggcctg gatttatctc 780  
 aaaaacaattt atcttcagtc accaatatta atgtaaaaaa gatgcctcag ctcccttctg 840  
 tgtacctaga ggaaaacaaa ctactgaac tgcctgaaaa atgtctgtcc gaactgagca 900  
 acttacaaga actctatatt aatcacaact tgctttctac aatttcacct ggagccttta 960  
 ttggcctaca taactctctt cgacttcato tcaattcaaa tagattgcag atgatcaaca 1020  
 gtaagtgggt tgatgctctt ccaaactctag agattctgat gattggggaa aatccaatta 1080  
 tcagaatcaa agacatgaac ttaagcctc ttatcaatct tccagcctg gttatagctg 1140  
 gtataaacct cacagaaata ccagataacg ccttgggttg actggaaac ttagaaagca 1200

-continued

```

tctcttttta cgataacagg cttattaaag taccccatgt tgctcttcaa aaagttgtaa 1260
atctcaaatt tttggatcta aataaaaaatc ctattaatag aatacgaagg ggtgatttta 1320
gcaatatgct acacttaaaa gagttgggga taaataatat gcctgagctg atttccatcg 1380
atagtcttgc tgtggataac ctgccagatt taagaaaaat agaagctact aacaacccta 1440
gattgtctta cattcacccc aatgcatttt tcagactccc caagctggaa tcaactcatgc 1500
tgaacagcaa tgctctcagt gcctgttacc atggtacat tgagtctctg ccaaacctca 1560
agggaaatcag catacacagt aaccocatca ggtgtgactg tgcatcccg tggatgaaca 1620
tgaacaaaaa caacattcga ttcattggagc cagattcact gttttgcgtg gaccacctg 1680
aattccaagg tcagaatggt cggcaagtgc atttcaggga catgatggaa atttgtctcc 1740
ctcttatagc tcctgagagc tttccttcta atctaaatgt agaagctggg agctatgttt 1800
cctttcactg tagagctact gcagaaccac agcctgaaat ctactggata acaccttctg 1860
gtcaaaaact cttgcctaata accctgacag acaagttcta tgtocattct gagggaacac 1920
tagatataaa tggcgtaact cccaaagaag ggggtttata tacttgtata gcaactaacc 1980
tagttggcgc tgacttgaag tctgttatga tcaaagtga tggatctttt ccacaagata 2040
acaatggctc tttgaatatt aaaataagag atattcaggc caattcagtt ttggtgtcct 2100
ggaaagcaag ttctaaaatt ctcaaatcta gtgttaaatg gacagccttt gtcaagactg 2160
aaaattctca tgctgcgcaa agtgctcgaa taccatctga tgtcaaggta tataatctta 2220
ctcatctgaa tccatcaact gagtataaaa tttgtattga tattcccacc atctatcaga 2280
aaaacagaaa aaaatgtgta aatgtcacca ccaaaggttt gcaccctgat caaaaagagt 2340
atgaaaagaa taataccaca acacttatgg cctgtcttgg aggcttctg gggattattg 2400
gtgtgatatg tcttatcagc tgctctctc cagaaatgaa ctgtgatggt ggacacagct 2460
atgtgaggaa ttacttacag aaaccaacct ttgcattagg tgagctttat cctcctctga 2520
taaattctctg ggaagcagga aaagaaaaaa gtacatcact gaaagtaaaa gcaactgtta 2580
taggtttacc aacaaatagc tcctaaaaac caccaaggaa acctactcca aaaatgaac 2639

```

```

<210> SEQ ID NO 69
<211> LENGTH: 708
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 69

```

Met Lys Asp Met Pro Leu Arg Ile His Val Leu Leu Gly Leu Ala Ile
 1             5             10             15
Thr Thr Leu Val Gln Ala Val Asp Lys Lys Val Asp Cys Pro Arg Leu
      20             25             30
Cys Thr Cys Glu Ile Arg Pro Trp Phe Thr Pro Arg Ser Ile Tyr Met
      35             40             45
Glu Ala Ser Thr Val Asp Cys Asn Asp Leu Gly Leu Leu Thr Phe Pro
      50             55             60
Ala Arg Leu Pro Ala Asn Thr Gln Ile Leu Leu Leu Gln Thr Asn Asn
      65             70             75             80
Ile Ala Lys Ile Glu Tyr Ser Thr Asp Phe Pro Val Asn Leu Thr Gly
      85             90             95
Leu Asp Leu Ser Gln Asn Asn Leu Ser Ser Val Thr Asn Ile Asn Val

```

-continued

100			105			110									
Lys	Lys	Met	Pro	Gln	Leu	Leu	Ser	Val	Tyr	Leu	Glu	Glu	Asn	Lys	Leu
		115					120					125			
Thr	Glu	Leu	Pro	Glu	Lys	Cys	Leu	Ser	Glu	Leu	Ser	Asn	Leu	Gln	Glu
	130					135						140			
Leu	Tyr	Ile	Asn	His	Asn	Leu	Leu	Ser	Thr	Ile	Ser	Pro	Gly	Ala	Phe
145				150						155					160
Ile	Gly	Leu	His	Asn	Leu	Leu	Arg	Leu	His	Leu	Asn	Ser	Asn	Arg	Leu
			165						170						175
Gln	Met	Ile	Asn	Ser	Lys	Trp	Phe	Asp	Ala	Leu	Pro	Asn	Leu	Glu	Ile
			180						185					190	
Leu	Met	Ile	Gly	Glu	Asn	Pro	Ile	Ile	Arg	Ile	Lys	Asp	Met	Asn	Phe
	195						200					205			
Lys	Pro	Leu	Ile	Asn	Leu	Arg	Ser	Leu	Val	Ile	Ala	Gly	Ile	Asn	Leu
	210						215					220			
Thr	Glu	Ile	Pro	Asp	Asn	Ala	Leu	Val	Gly	Leu	Glu	Asn	Leu	Glu	Ser
225					230					235					240
Ile	Ser	Phe	Tyr	Asp	Asn	Arg	Leu	Ile	Lys	Val	Pro	His	Val	Ala	Leu
			245						250					255	
Gln	Lys	Val	Val	Asn	Leu	Lys	Phe	Leu	Asp	Leu	Asn	Lys	Asn	Pro	Ile
			260					265						270	
Asn	Arg	Ile	Arg	Arg	Gly	Asp	Phe	Ser	Asn	Met	Leu	His	Leu	Lys	Glu
		275					280					285			
Leu	Gly	Ile	Asn	Asn	Met	Pro	Glu	Leu	Ile	Ser	Ile	Asp	Ser	Leu	Ala
	290					295					300				
Val	Asp	Asn	Leu	Pro	Asp	Leu	Arg	Lys	Ile	Glu	Ala	Thr	Asn	Asn	Pro
305					310					315					320
Arg	Leu	Ser	Tyr	Ile	His	Pro	Asn	Ala	Phe	Phe	Arg	Leu	Pro	Lys	Leu
			325						330					335	
Glu	Ser	Leu	Met	Leu	Asn	Ser	Asn	Ala	Leu	Ser	Ala	Leu	Tyr	His	Gly
			340					345					350		
Thr	Ile	Glu	Ser	Leu	Pro	Asn	Leu	Lys	Glu	Ile	Ser	Ile	His	Ser	Asn
		355					360					365			
Pro	Ile	Arg	Cys	Asp	Cys	Val	Ile	Arg	Trp	Met	Asn	Met	Asn	Lys	Thr
	370					375					380				
Asn	Ile	Arg	Phe	Met	Glu	Pro	Asp	Ser	Leu	Phe	Cys	Val	Asp	Pro	Pro
385					390					395					400
Glu	Phe	Gln	Gly	Gln	Asn	Val	Arg	Gln	Val	His	Phe	Arg	Asp	Met	Met
			405					410						415	
Glu	Ile	Cys	Leu	Pro	Leu	Ile	Ala	Pro	Glu	Ser	Phe	Pro	Ser	Asn	Leu
			420					425					430		
Asn	Val	Glu	Ala	Gly	Ser	Tyr	Val	Ser	Phe	His	Cys	Arg	Ala	Thr	Ala
		435					440					445			
Glu	Pro	Gln	Pro	Glu	Ile	Tyr	Trp	Ile	Thr	Pro	Ser	Gly	Gln	Lys	Leu
	450					455						460			
Leu	Pro	Asn	Thr	Leu	Thr	Asp	Lys	Phe	Tyr	Val	His	Ser	Glu	Gly	Thr
465					470					475					480
Leu	Asp	Ile	Asn	Gly	Val	Thr	Pro	Lys	Glu	Gly	Gly	Leu	Tyr	Thr	Cys
			485						490					495	
Ile	Ala	Thr	Asn	Leu	Val	Gly	Ala	Asp	Leu	Lys	Ser	Val	Met	Ile	Lys
			500					505					510		

-continued

Val Asp Gly Ser Phe Pro Gln Asp Asn Asn Gly Ser Leu Asn Ile Lys  
 515 520 525  
 Ile Arg Asp Ile Gln Ala Asn Ser Val Leu Val Ser Trp Lys Ala Ser  
 530 535 540  
 Ser Lys Ile Leu Lys Ser Ser Val Lys Trp Thr Ala Phe Val Lys Thr  
 545 550 555 560  
 Glu Asn Ser His Ala Ala Gln Ser Ala Arg Ile Pro Ser Asp Val Lys  
 565 570 575  
 Val Tyr Asn Leu Thr His Leu Asn Pro Ser Thr Glu Tyr Lys Ile Cys  
 580 585 590  
 Ile Asp Ile Pro Thr Ile Tyr Gln Lys Asn Arg Lys Lys Cys Val Asn  
 595 600 605  
 Val Thr Thr Lys Gly Leu His Pro Asp Gln Lys Glu Tyr Glu Lys Asn  
 610 615 620  
 Asn Thr Thr Thr Leu Met Ala Cys Leu Gly Gly Leu Leu Gly Ile Ile  
 625 630 635 640  
 Gly Val Ile Cys Leu Ile Ser Cys Leu Ser Pro Glu Met Asn Cys Asp  
 645 650 655  
 Gly Gly His Ser Tyr Val Arg Asn Tyr Leu Gln Lys Pro Thr Phe Ala  
 660 665 670  
 Leu Gly Glu Leu Tyr Pro Pro Leu Ile Asn Leu Trp Glu Ala Gly Lys  
 675 680 685  
 Glu Lys Ser Thr Ser Leu Lys Val Lys Ala Thr Val Ile Gly Leu Pro  
 690 695 700  
 Thr Asn Met Ser  
 705

<210> SEQ ID NO 70  
 <211> LENGTH: 1305  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

gcccgggact ggcgcaaggt gcccaagcaa ggaagaaat aatgaagaga cacatgtgtt 60  
 agctgcagcc ttttgaaaac cgcaagaagg aaatcaatag tgtggacag gctggaacct 120  
 ttaccacgct tgttgagta gatgaggaat gggctcgtga ttatgctgac attccagcat 180  
 gaatctggta gacctgtggt taaccggttc cctctccatg tgtctcctcc tacaagttt 240  
 tgttcttatg atactgtgct ttcattctgc cagtatgtgt cccaagggt gctcttgttc 300  
 ttcctctggg ggtttaaagt tcacctgtag caatgcaaat ctcaaggaaa tacctagaga 360  
 tcttctctct gaaacagtct tactgtatct ggactccaat cagatcacat ctattoccaa 420  
 tgaaattttt aaggacctcc atcaactgag agttctcaac ctgtccaaaa atggcattga 480  
 gtttatcgat gagcatgcct tcaaaggagt agctgaaacc ttgcagactc tggacttgtc 540  
 cgacaatcgg attcaaagtg tgcacaaaaa tgccttcaat aacctgaagg ccagggccag 600  
 aattgccaac aaccctggc actgcgactg tactctacag caagttctga ggagcatggc 660  
 gtccaatcat gagacagccc acaactgtat ctgtaaaacg tccgtgttgg atgaacatgc 720  
 tggcagacca ttcctcaatg ctgccaacga cgctgacctt tgtaacctcc ctaaaaaac 780  
 taccgattat gccatgctgg tcaccatggt tggctggttc actatggtga tctcatatgt 840



-continued

---

```

ggtatattat gtgaggcaaa atcaggagga tgcccggaga cacctcgaat acttgaaatc 900
cctgccaagc aggcagaaga aagcagatga acctgatgat attagcactg tggatatagt 960
tccaaaactga ctgtcattga gaaagaaaga aagtagtttg cgattgcagt agaaataagt 1020
ggtttacttc tcccattcat tgtaaacatt tgaactttg tatttcagtt ttttttgaat 1080
tatgccactg ctgaactttt acaaacact acaacataaa taatttgagt ttaggtgatc 1140
caccoccttaa ttgtaccccc gatggtatat ttctgagtaa gctactatct gaacattagt 1200
tagatccatc tcactattta ataatgaaat ttattttttt aatttaaaag caaataaaag 1260
cttaactttg aaccatggga aaaaaaaaaa aaaaaaaaaa aaaca 1305

```

&lt;210&gt; SEQ ID NO 71

&lt;211&gt; LENGTH: 259

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 71

```

Met Asn Leu Val Asp Leu Trp Leu Thr Arg Ser Leu Ser Met Cys Leu
 1           5           10          15
Leu Leu Gln Ser Phe Val Leu Met Ile Leu Cys Phe His Ser Ala Ser
          20          25          30
Met Cys Pro Lys Gly Cys Leu Cys Ser Ser Ser Gly Gly Leu Asn Val
 35          40          45
Thr Cys Ser Asn Ala Asn Leu Lys Glu Ile Pro Arg Asp Leu Pro Pro
 50          55          60
Glu Thr Val Leu Leu Tyr Leu Asp Ser Asn Gln Ile Thr Ser Ile Pro
 65          70          75          80
Asn Glu Ile Phe Lys Asp Leu His Gln Leu Arg Val Leu Asn Leu Ser
          85          90          95
Lys Asn Gly Ile Glu Phe Ile Asp Glu His Ala Phe Lys Gly Val Ala
 100         105         110
Glu Thr Leu Gln Thr Leu Asp Leu Ser Asp Asn Arg Ile Gln Ser Val
 115         120         125
His Lys Asn Ala Phe Asn Asn Leu Lys Ala Arg Ala Arg Ile Ala Asn
 130         135         140
Asn Pro Trp His Cys Asp Cys Thr Leu Gln Gln Val Leu Arg Ser Met
 145         150         155         160
Ala Ser Asn His Glu Thr Ala His Asn Val Ile Cys Lys Thr Ser Val
 165         170         175
Leu Asp Glu His Ala Gly Arg Pro Phe Leu Asn Ala Ala Asn Asp Ala
 180         185         190
Asp Leu Cys Asn Leu Pro Lys Lys Thr Thr Asp Tyr Ala Met Leu Val
 195         200         205
Thr Met Phe Gly Trp Phe Thr Met Val Ile Ser Tyr Val Val Tyr Tyr
 210         215         220
Val Arg Gln Asn Gln Glu Asp Ala Arg Arg His Leu Glu Tyr Leu Lys
 225         230         235         240
Ser Leu Pro Ser Arg Gln Lys Lys Ala Asp Glu Pro Asp Asp Ile Ser
 245         250         255

Thr Val Val

```

&lt;210&gt; SEQ ID NO 72

-continued

&lt;211&gt; LENGTH: 2290

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 72

```

accgagccga gcgaccgaa ggcgcgccg agatgcaggt gagcaagagg atgctggcgg      60
ggggcgtgag gacatgccc agccccctcc tggcctgctg gcagcccatc ctctgctgg      120
tgctgggctc agtgctgtca ggctcggcca cgggctgccc gccccgctgc gagtgtccg      180
cccaggacog cgctgtgctg tgccaccgca agtgctttgt ggcagtcccc gagggcatcc      240
ccaccgagac gcgctgctg gacctaggca agaaccgcat caaacgctc aaccaggacg      300
agttcgcag cttcccgcac ctggaggagc tggagctcaa cgagaacatc gtgagcgcg      360
tggagcccgg cgccttcaac aaactcttca aactccggac gctgggtctc cgcagcaaac      420
gcctgaagct catcccgcta ggcgctttca ctggcctcag caacctgacc aagcaggaca      480
tcagcgagaa caagatcgtt atoctactgg actacatggt tcaggacctg tacaacctca      540
agtcaactgga ggttggcgac aatgacctcg totacatctc tcaccgcgcc ttcagcggcc      600
tcaacagcct ggagcagctg acgctggaga aatgcaacct gacctccatc cccaccgagg      660
cgctgtccca cctgcacggc ctcatcgtcc tgaggctccg gcacctcaac atcaatgcca      720
tccgggacta ctcttcaag aggtgttacc gactcaaggt cttggagatc tcccaactggc      780
cctacttgga caccatgaca cccaactgcc totacggcct caacctgacg tccctgtcca      840
tcacacactg caatctgacc gctgtgcctt acctggccgt ccgccacctt gctatctcc      900
gcttctcaa cctctctac aaccocatca gcaccattga gggotccatg ttgcatgagc      960
tgctccgctg gcaggagatc cagctgggtg gcggcgagct ggcctgggtg gagccctatg     1020
ccttccgctg cctcaactac ctgcgcgtgc tcaatgtctc tggcaaccag ctgaccacac     1080
tggaggaatc agtcttccac tcggtgggca acctggagac actcatctct gactccaacc     1140
cgctggcctg cgactgtcgg ctctgtgggg tgttccggcg ccgctggcgg ctcaacttca     1200
accggcagca gccacgtgc gccacgcccg agtttgtcca gggcaaggag ttcaaggact     1260
tccctgatgt gctactgccc aactacttca cctgccgccg cgcgccatc cgggaccgca     1320
aggcccagca ggtgtttgtg gacgagggcc acacgggtgca gtttgtgtgc cgggccgatg     1380
gcgaccggcc gcccgccatc ctctggctct caccocgaaa gcaacctggtc tcagccaaga     1440
gcaatgggog gctcacagtc tccctgatg gcaagctgga ggtgcgctac gccagggtac     1500
aggacaacgg cagctacctg tgcatcggcg ccaacgcggg cggcaacgac tccatgcccg     1560
cccacctgca tgtgcgcagc tactcggccc actggcccca tcagcccaac aagacctctg     1620
ctttcatctc caaccagccg gcgaggggag aggccaaacag caccocgccc actgtgcctt     1680
tccccttcga catcaagacc ctcatcatcg ccaccacat gggcttcatc tcttctctgg     1740
gcgtcgtcct cttctgcctg gtgtgtgtgt ttctctggag cgggggcaag ggcaacacaa     1800
agcacaacat cgagatcgag tatgtgcccc gaaagtcgga cgcaggcatc agctccgccg     1860
acgcgccccg caagttcaac atgaagatga tatgaggccg gggcgggggg cagggacccc     1920
cgggcggcog ggcaggggaa gggcctggt cgcacctgc tcaactctca gtccttccca     1980
ctctctccct acccttctac acacgttctc tttctccctc ccgctcctgt cccctgctgc     2040
ccccgcagcc ccctcaccac ctgcctcctt totaccagga cctcagaagc ccagacctgg     2100

```

-continued

---

```

ggaccccacc tacacagggg cattgacaga ctggagttga aagccgacga accgacacgc 2160
ggcagagtca ataattcaat aaaaaagtta cgaactttct ctgtaacttg ggtttcaata 2220
attatggatt tttatgaaaa ctgaaataa taaaaagaga aaaaaactaa aaaaaaaaaa 2280
aaaaaaaaaa 2290

```

```

<210> SEQ ID NO 73
<211> LENGTH: 620
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 73

```

```

Met Gln Val Ser Lys Arg Met Leu Ala Gly Gly Val Arg Ser Met Pro
 1                    5                    10          15
Ser Pro Leu Leu Ala Cys Trp Gln Pro Ile Leu Leu Leu Val Leu Gly
 20                    25                    30
Ser Val Leu Ser Gly Ser Ala Thr Gly Cys Pro Pro Arg Cys Glu Cys
 35                    40                    45
Ser Ala Gln Asp Arg Ala Val Leu Cys His Arg Lys Cys Phe Val Ala
 50                    55                    60
Val Pro Glu Gly Ile Pro Thr Glu Thr Arg Leu Leu Asp Leu Gly Lys
 65                    70                    75                    80
Asn Arg Ile Lys Thr Leu Asn Gln Asp Glu Phe Ala Ser Phe Pro His
 85                    90                    95
Leu Glu Glu Leu Glu Leu Asn Glu Asn Ile Val Ser Ala Val Glu Pro
100                    105                    110
Gly Ala Phe Asn Asn Leu Phe Asn Leu Arg Thr Leu Gly Leu Arg Ser
115                    120                    125
Asn Arg Leu Lys Leu Ile Pro Leu Gly Val Phe Thr Gly Leu Ser Asn
130                    135                    140
Leu Thr Lys Gln Asp Ile Ser Glu Asn Lys Ile Val Ile Leu Leu Asp
145                    150                    155                    160
Tyr Met Phe Gln Asp Leu Tyr Asn Leu Lys Ser Leu Glu Val Gly Asp
165                    170                    175
Asn Asp Leu Val Tyr Ile Ser His Arg Ala Phe Ser Gly Leu Asn Ser
180                    185                    190
Leu Glu Gln Leu Thr Leu Glu Lys Cys Asn Leu Thr Ser Ile Pro Thr
195                    200                    205
Glu Ala Leu Ser His Leu His Gly Leu Ile Val Leu Arg Leu Arg His
210                    215                    220
Leu Asn Ile Asn Ala Ile Arg Asp Tyr Ser Phe Lys Arg Leu Tyr Arg
225                    230                    235                    240
Leu Lys Val Leu Glu Ile Ser His Trp Pro Tyr Leu Asp Thr Met Thr
245                    250                    255
Pro Asn Cys Leu Tyr Gly Leu Asn Leu Thr Ser Leu Ser Ile Thr His
260                    265                    270
Cys Asn Leu Thr Ala Val Pro Tyr Leu Ala Val Arg His Leu Val Tyr
275                    280                    285
Leu Arg Phe Leu Asn Leu Ser Tyr Asn Pro Ile Ser Thr Ile Glu Gly
290                    295                    300
Ser Met Leu His Glu Leu Leu Arg Leu Gln Glu Ile Gln Leu Val Gly
305                    310                    315                    320

```

-continued

---

Gly Gln Leu Ala Val Val Glu Pro Tyr Ala Phe Arg Gly Leu Asn Tyr  
 325 330 335

Leu Arg Val Leu Asn Val Ser Gly Asn Gln Leu Thr Thr Leu Glu Glu  
 340 345 350

Ser Val Phe His Ser Val Gly Asn Leu Glu Thr Leu Ile Leu Asp Ser  
 355 360 365

Asn Pro Leu Ala Cys Asp Cys Arg Leu Leu Trp Val Phe Arg Arg Arg  
 370 375 380

Trp Arg Leu Asn Phe Asn Arg Gln Gln Pro Thr Cys Ala Thr Pro Glu  
 385 390 395 400

Phe Val Gln Gly Lys Glu Phe Lys Asp Phe Pro Asp Val Leu Leu Pro  
 405 410 415

Asn Tyr Phe Thr Cys Arg Arg Ala Arg Ile Arg Asp Arg Lys Ala Gln  
 420 425 430

Gln Val Phe Val Asp Glu Gly His Thr Val Gln Phe Val Cys Arg Ala  
 435 440 445

Asp Gly Asp Pro Pro Pro Ala Ile Leu Trp Leu Ser Pro Arg Lys His  
 450 455 460

Leu Val Ser Ala Lys Ser Asn Gly Arg Leu Thr Val Phe Pro Asp Gly  
 465 470 475 480

Thr Leu Glu Val Arg Tyr Ala Gln Val Gln Asp Asn Gly Thr Tyr Leu  
 485 490 495

Cys Ile Ala Ala Asn Ala Gly Gly Asn Asp Ser Met Pro Ala His Leu  
 500 505 510

His Val Arg Ser Tyr Ser Pro Asp Trp Pro His Gln Pro Asn Lys Thr  
 515 520 525

Phe Ala Phe Ile Ser Asn Gln Pro Gly Glu Gly Glu Ala Asn Ser Thr  
 530 535 540

Arg Ala Thr Val Pro Phe Pro Phe Asp Ile Lys Thr Leu Ile Ile Ala  
 545 550 555 560

Thr Thr Met Gly Phe Ile Ser Phe Leu Gly Val Val Leu Phe Cys Leu  
 565 570 575

Val Leu Leu Phe Leu Trp Ser Arg Gly Lys Gly Asn Thr Lys His Asn  
 580 585 590

Ile Glu Ile Glu Tyr Val Pro Arg Lys Ser Asp Ala Gly Ile Ser Ser  
 595 600 605

Ala Asp Ala Pro Arg Lys Phe Asn Met Lys Met Ile  
 610 615 620

<210> SEQ ID NO 74  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 74

tcacctggag cctttattgg cc

22

<210> SEQ ID NO 75  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:

---

-continued

---

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 75

ataccagcta taaccaggct gcg 23

<210> SEQ ID NO 76  
<211> LENGTH: 52  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 76

caacagtaag tggtttgatg ctcttccaaa tctagagatt ctgatgattg 50

gg 52

<210> SEQ ID NO 77  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 77

ccatgtgtct cctcctacaa ag 22

<210> SEQ ID NO 78  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 78

gggaatagat gtgatctgat tgg 23

<210> SEQ ID NO 79  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 79

cacctgtagc aatgcaaadc tcaaggaaat acctagagat ctctctctctg 50

<210> SEQ ID NO 80  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

<400> SEQUENCE: 80

agcaaccgcc tgaagctcat cc 22

<210> SEQ ID NO 81

-continued

---

<211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe  
  
 <400> SEQUENCE: 81  
  
 aaggcgcggt gaaagatgta gacg 24

<210> SEQ ID NO 82  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe  
  
 <400> SEQUENCE: 82  
  
 gactacatgt ttcaggacct gtacaacctc aagtcactgg aggttggcga 50

<210> SEQ ID NO 83  
 <211> LENGTH: 1685  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 83  
  
 cccacgcgctc cgcacctcgg ccccgggctc cgaagcggct cgggggcgcc ctttcgggtca 60  
 acatcgtagt ccaccccctc cccatcccca gcccccgagg attcaggctc gccagcgccc 120  
 agccaggagc cgggccggga agcgcgatgg gggccccagc cgcctcgtc ctgctcctgc 180  
 tctctctgtt cgctctgtc tgggcgccc gcggggcca cctctccag gacgacagcc 240  
 agccctggac atctgatgaa acagtgggtg ctggtggcac cgtggtgctc aagtgccaa 300  
 tgaaagatca cgaggactca tcctgcaat ggtctaacc tgctcagcag actcttact 360  
 ttggggagaa gagagccctt cgagataatc gaattcagct ggttacctct acgccccag 420  
 agctcagcat cagcatcagc aatgtggccc tggcagacga gggcgagtac acctgctcaa 480  
 tcttactact gcctgtgcga actgccaagt ccctcgtcac tgtgctagga attccacaga 540  
 agcccatcat cactggttat aaatcttcat tacgggaaaa agacacagcc accctaaact 600  
 gtcagtcttc tgggagcaag cctgcagccc ggctcacctg gagaaaagggt gaccaagaac 660  
 tccacggaga accaaccgc atacaggaag atcccaatgg taaaacctc actgtcagca 720  
 gctcggtgac attccaggtt acccgggagg atgatggggc gagcatcgtg tgctctgtga 780  
 accatgaatc tctaaaggga gctgacagat ccacctctca acgcattgaa gttttataca 840  
 caccaactgc gatgattagg ccagaccctc cccatcctcg tgagggccag aagctgttgc 900  
 tacactgtga gggctcggc aatccagtcc cccagcagta cctatgggag aaggagggca 960  
 gtgtgccacc cctgaagatg acccaggaga gtgcctgat cttcccttct ctaacaaga 1020  
 gtgacagtgg cacctacggc tgcacagcca ccagcaacat gggcagctac aaggcctact 1080  
 acacctcaa tgtaatgac cccagtccgg tgccctctc ctccagcacc taccagcca 1140  
 tcatcggtag gatcgtggct ttcattgtct tctgtgtct catcatgctc atcttcttgc 1200  
 gccactactt gatccggcac aaaggaacct acctgacaca tgaggcaaaa ggctccgacg 1260  
 atgctccaga cgcggacag gccatcatca atgcagaagg cgggcagtca ggaggggacg 1320

## -continued

---

```

acaagaagga atatttcac tagagggcgc tgcccacttc ctgcgccccc caggggcccct 1380
gtggggactg ctggggccgt caccaaccgc gacttgtaga gagcaaccgc agggcgcccc 1440
ctcccgttg ctccccagcc ccccccccc cctgtacaga atgtctgctt tgggtgcggt 1500
tttgtactgc gtttgaatg gggagggagg agggcggggg gaggggaggg ttgcccctag 1560
ccctttccgt ggcttctctg catttgggtt attattattt ttgtaacaat cccaaatcaa 1620
atctgtctcc aggtctgaga ggcaggagcc ctgggggtgag aaaagcaaaa aacaacaaa 1680
aaaca 1685

```

<210> SEQ ID NO 84

<211> LENGTH: 398

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

```

Met Gly Ala Pro Ala Ala Ser Leu Leu Leu Leu Leu Leu Phe Ala
 1          5          10          15
Cys Cys Trp Ala Pro Gly Gly Ala Asn Leu Ser Gln Asp Asp Ser Gln
 20          25          30
Pro Trp Thr Ser Asp Glu Thr Val Val Ala Gly Gly Thr Val Val Leu
 35          40          45
Lys Cys Gln Val Lys Asp His Glu Asp Ser Ser Leu Gln Trp Ser Asn
 50          55          60
Pro Ala Gln Gln Thr Leu Tyr Phe Gly Glu Lys Arg Ala Leu Arg Asp
 65          70          75          80
Asn Arg Ile Gln Leu Val Thr Ser Thr Pro His Glu Leu Ser Ile Ser
 85          90          95
Ile Ser Asn Val Ala Leu Ala Asp Glu Gly Glu Tyr Thr Cys Ser Ile
100          105          110
Phe Thr Met Pro Val Arg Thr Ala Lys Ser Leu Val Thr Val Leu Gly
115          120          125
Ile Pro Gln Lys Pro Ile Ile Thr Gly Tyr Lys Ser Ser Leu Arg Glu
130          135          140
Lys Asp Thr Ala Thr Leu Asn Cys Gln Ser Ser Gly Ser Lys Pro Ala
145          150          155          160
Ala Arg Leu Thr Trp Arg Lys Gly Asp Gln Glu Leu His Gly Glu Pro
165          170          175
Thr Arg Ile Gln Glu Asp Pro Asn Gly Lys Thr Phe Thr Val Ser Ser
180          185          190
Ser Val Thr Phe Gln Val Thr Arg Glu Asp Asp Gly Ala Ser Ile Val
195          200          205
Cys Ser Val Asn His Glu Ser Leu Lys Gly Ala Asp Arg Ser Thr Ser
210          215          220
Gln Arg Ile Glu Val Leu Tyr Thr Pro Thr Ala Met Ile Arg Pro Asp
225          230          235          240
Pro Pro His Pro Arg Glu Gly Gln Lys Leu Leu Leu His Cys Glu Gly
245          250          255
Arg Gly Asn Pro Val Pro Gln Gln Tyr Leu Trp Glu Lys Glu Gly Ser
260          265          270
Val Pro Pro Leu Lys Met Thr Gln Glu Ser Ala Leu Ile Phe Pro Phe
275          280          285

```

-continued

---

Leu Asn Lys Ser Asp Ser Gly Thr Tyr Gly Cys Thr Ala Thr Ser Asn  
 290 295 300

Met Gly Ser Tyr Lys Ala Tyr Tyr Thr Leu Asn Val Asn Asp Pro Ser  
 305 310 315 320

Pro Val Pro Ser Ser Ser Ser Thr Tyr His Ala Ile Ile Gly Gly Ile  
 325 330 335

Val Ala Phe Ile Val Phe Leu Leu Leu Ile Met Leu Ile Phe Leu Gly  
 340 345 350

His Tyr Leu Ile Arg His Lys Gly Thr Tyr Leu Thr His Glu Ala Lys  
 355 360 365

Gly Ser Asp Asp Ala Pro Asp Ala Asp Thr Ala Ile Ile Asn Ala Glu  
 370 375 380

Gly Gly Gln Ser Gly Gly Asp Asp Lys Lys Glu Tyr Phe Ile  
 385 390 395

<210> SEQ ID NO 85  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 85

gctaggaatt ccacagaagc cc 22

<210> SEQ ID NO 86  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 86

aacctggaat gtcaccgagc tg 22

<210> SEQ ID NO 87  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 87

cctagcacag tgacgagga cttggc 26

<210> SEQ ID NO 88  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 88

aagacacagc cacctaacc tgtcagtctt ctgggagcaa gcctgcagcc 50

<210> SEQ ID NO 89  
 <211> LENGTH: 50



-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 89

gccctggcag acgagggcga gtacacctgc tcaatcttca ctatgcctgt          50

<210> SEQ ID NO 90
<211> LENGTH: 2755
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

gggggttagg gaggaagaa tccaccccca ccccccaaa cccttttctt ctcctttcct          60
ggctctggac attggagcac taaatgaact tgaattgtgt ctgtggcgag caggatggtc          120
gctgttactt tgtgatgaga tcggggatga attgctcgtt ttaaaaatgc tgctttggat          180
tctgttgctg gagacgtctc ttgtttttgc cgctggaaac gttacagggg acgtttgcaa          240
agagaagatc tgttcctgca atgagataga aggggaccta cacgtagact gtgaaaaaaa          300
gggcttcaca agtctgcagc gtttcactgc cccgacttcc cagttttacc atttatttct          360
gcatggcaat tccctcactc gacttttccc taatgagttc gctaactttt ataatgcggt          420
tagtttgcac atgaaaaaca atggcttgca tgaaatcggt ccgggggctt ttctggggct          480
gcagctggtg aaaaggctgc acatcaacaa caacaagatc aagtcttttc gaaagcagac          540
ttttctgggg ctggacgatc tggaatatct ccaggctgat tttaatttat tacgagatat          600
agaccggggg gccttcagag acttgaacaa gctggaggtg ctcattttaa atgacaatct          660
catcagcacc ctacctgcca acgtgttcca gtatgtgcc atcaccacc tcgacctccg          720
gggtaacagg ctgaaaacgc tgcctatga ggaggctctg gagcaaatcc ctggtattgc          780
ggagatcctg ctagaggata accctggga ctgcacctgt gatctgctct cctgaaaga          840
atggctggaa aacattccca agaatgcctt gatcgccga gtggtctgctg aagccccac          900
cagactgcag ggtaaaagacc tcaatgaaac caccgaacag gacttgtgtc ctttgaaaaa          960
ccgagtggat tctagtctcc cggcgcccc tgcccaagaa gagaccttg ctctggacc          1020
cctgccaaact cctttcaaga caaatgggca agaggatcat gccacaccag ggtctgctcc          1080
aaacggaggt acaaagatcc caggcaactg gcagatcaaa atcagacca cagcagcgat          1140
agcgacgggt agctccagga acaaaccctt agctaacagt ttaccctgcc ctgggggctg          1200
cagctgcgac cacatcccag ggtcgggttt aaagatgaac tgcaacaaca ggaacgtgag          1260
cagcttgctt gatttgaagc ccaagctctc taacgtgcag gagcttttcc tacgagataa          1320
caagatccac agcatccgaa aatcgcaactt tgtggattac aagaacctca ttctgttggg          1380
tctgggcaac aataacatcg ctactgtaga gaacaacact ttcaagaacc ttttggacct          1440
caggttggta tacatggata gcaattacct ggacacgctg tcccgggaga aattcgcggg          1500
gctgcaaaac ctagagtacc tgaactgga gtacaacgct atccagctca tcctccggg          1560
cactttcaat gccatgccc aactgaggat cctcattctc aacaacaacc tgctgaggtc          1620
cctgcctgtg gacgtgttgc ctgggtctc gctctctaaa ctacgctgc aacaacaatta          1680
cttcatgtac ctccgggtg caggggtgct ggaccagtta acctccatca tccagataga          1740

```

-continued

```

cctccacgga aaccctggg agtgctcctg cacaattgtg cctttcaagc agtgggcaga 1800
acgcttgggt tccgaagtgc tgatgagcga cctcaagtgt gagacgccgg tgaacttctt 1860
tagaaaggat ttcatgtccc tctccaatga cgagatctgc cctcagctgt acgctaggat 1920
ctcgcccaag ttaacttcgc acagtaaaaa cagcactggg ttggcgggaga ccgggacgca 1980
ctccaactcc tacctagaca ccagcagggt gtccatctcg gtgttggtcc cgggactgct 2040
gctggtgttt gtcacctccg ccttcaccgt ggtgggcatg ctcgtgttta tcttgaggaa 2100
ccgaaagcgg tccaagagac gagatgcca ctcctccgcg tccgagatta attccctaca 2160
gacagtctgt gactcttctc actggcacia tgggccttac aacgcagatg gggcccacag 2220
agtgatgac tgtggctctc actcgtctc agactaagac cccaacccca ataggggagg 2280
gcagagggaa ggcatacat ccttccccac cgcaggcacc ccgggggctg gaggggctg 2340
taccxaaatc cccgcgccat cagcctggat gggcataagt agataaataa ctgtgagctc 2400
gcacaaccga aagggcctga cccttactt agtcocctcc ttgaaacaaa gagcagactg 2460
tggagagctg ggagagcgca gccagctcgc ttttctgta gagccccttt tgacagaaag 2520
cccagcacga ccctgctgga agaactgaca gtgcctcgc cctogggccc ggggcctgtg 2580
gggttgatg ccgcggttct atacatatat acatatatcc acatctatat agagagatag 2640
atatctattt ttcccctgtg gattagcccc gtgatggctc cctgttggtc acgcagggat 2700
gggcagttgc acgaaggcat gaatgtattg taaataagta actttgactt ctgac 2755

```

```

<210> SEQ ID NO 91
<211> LENGTH: 696
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 91

```

Met Leu Leu Trp Ile Leu Leu Leu Glu Thr Ser Leu Cys Phe Ala Ala
 1             5             10            15
Gly Asn Val Thr Gly Asp Val Cys Lys Glu Lys Ile Cys Ser Cys Asn
          20            25            30
Glu Ile Glu Gly Asp Leu His Val Asp Cys Glu Lys Lys Gly Phe Thr
          35            40            45
Ser Leu Gln Arg Phe Thr Ala Pro Thr Ser Gln Phe Tyr His Leu Phe
 50            55            60
Leu His Gly Asn Ser Leu Thr Arg Leu Phe Pro Asn Glu Phe Ala Asn
 65            70            75            80
Phe Tyr Asn Ala Val Ser Leu His Met Glu Asn Asn Gly Leu His Glu
          85            90            95
Ile Val Pro Gly Ala Phe Leu Gly Leu Gln Leu Val Lys Arg Leu His
          100           105           110
Ile Asn Asn Asn Lys Ile Lys Ser Phe Arg Lys Gln Thr Phe Leu Gly
          115           120           125
Leu Asp Asp Leu Glu Tyr Leu Gln Ala Asp Phe Asn Leu Leu Arg Asp
          130           135           140
Ile Asp Pro Gly Ala Phe Gln Asp Leu Asn Lys Leu Glu Val Leu Ile
          145           150           155           160
Leu Asn Asp Asn Leu Ile Ser Thr Leu Pro Ala Asn Val Phe Gln Tyr
          165           170           175
Val Pro Ile Thr His Leu Asp Leu Arg Gly Asn Arg Leu Lys Thr Leu

```

-continued

180					185					190					
Pro	Tyr	Glu	Glu	Val	Leu	Glu	Gln	Ile	Pro	Gly	Ile	Ala	Glu	Ile	Leu
		195					200					205			
Leu	Glu	Asp	Asn	Pro	Trp	Asp	Cys	Thr	Cys	Asp	Leu	Leu	Ser	Leu	Lys
	210					215					220				
Glu	Trp	Leu	Glu	Asn	Ile	Pro	Lys	Asn	Ala	Leu	Ile	Gly	Arg	Val	Val
	225					230					235				240
Cys	Glu	Ala	Pro	Thr	Arg	Leu	Gln	Gly	Lys	Asp	Leu	Asn	Glu	Thr	Thr
				245					250					255	
Glu	Gln	Asp	Leu	Cys	Pro	Leu	Lys	Asn	Arg	Val	Asp	Ser	Ser	Leu	Pro
			260						265					270	
Ala	Pro	Pro	Ala	Gln	Glu	Glu	Thr	Phe	Ala	Pro	Gly	Pro	Leu	Pro	Thr
			275				280						285		
Pro	Phe	Lys	Thr	Asn	Gly	Gln	Glu	Asp	His	Ala	Thr	Pro	Gly	Ser	Ala
	290					295					300				
Pro	Asn	Gly	Gly	Thr	Lys	Ile	Pro	Gly	Asn	Trp	Gln	Ile	Lys	Ile	Arg
	305					310					315				320
Pro	Thr	Ala	Ala	Ile	Ala	Thr	Gly	Ser	Ser	Arg	Asn	Lys	Pro	Leu	Ala
				325							330				335
Asn	Ser	Leu	Pro	Cys	Pro	Gly	Gly	Cys	Ser	Cys	Asp	His	Ile	Pro	Gly
			340					345					350		
Ser	Gly	Leu	Lys	Met	Asn	Cys	Asn	Asn	Arg	Asn	Val	Ser	Ser	Leu	Ala
		355					360					365			
Asp	Leu	Lys	Pro	Lys	Leu	Ser	Asn	Val	Gln	Glu	Leu	Phe	Leu	Arg	Asp
	370					375					380				
Asn	Lys	Ile	His	Ser	Ile	Arg	Lys	Ser	His	Phe	Val	Asp	Tyr	Lys	Asn
	385					390					395				400
Leu	Ile	Leu	Leu	Asp	Leu	Gly	Asn	Asn	Asn	Ile	Ala	Thr	Val	Glu	Asn
				405					410					415	
Asn	Thr	Phe	Lys	Asn	Leu	Leu	Asp	Leu	Arg	Trp	Leu	Tyr	Met	Asp	Ser
			420						425				430		
Asn	Tyr	Leu	Asp	Thr	Leu	Ser	Arg	Glu	Lys	Phe	Ala	Gly	Leu	Gln	Asn
	435						440					445			
Leu	Glu	Tyr	Leu	Asn	Val	Glu	Tyr	Asn	Ala	Ile	Gln	Leu	Ile	Leu	Pro
	450					455					460				
Gly	Thr	Phe	Asn	Ala	Met	Pro	Lys	Leu	Arg	Ile	Leu	Ile	Leu	Asn	Asn
	465					470					475				480
Asn	Leu	Leu	Arg	Ser	Leu	Pro	Val	Asp	Val	Phe	Ala	Gly	Val	Ser	Leu
			485						490					495	
Ser	Lys	Leu	Ser	Leu	His	Asn	Asn	Tyr	Phe	Met	Tyr	Leu	Pro	Val	Ala
		500						505					510		
Gly	Val	Leu	Asp	Gln	Leu	Thr	Ser	Ile	Ile	Gln	Ile	Asp	Leu	His	Gly
		515					520					525			
Asn	Pro	Trp	Glu	Cys	Ser	Cys	Thr	Ile	Val	Pro	Phe	Lys	Gln	Trp	Ala
	530					535					540				
Glu	Arg	Leu	Gly	Ser	Glu	Val	Leu	Met	Ser	Asp	Leu	Lys	Cys	Glu	Thr
	545					550					555				560
Pro	Val	Asn	Phe	Phe	Arg	Lys	Asp	Phe	Met	Leu	Leu	Ser	Asn	Asp	Glu
			565						570					575	
Ile	Cys	Pro	Gln	Leu	Tyr	Ala	Arg	Ile	Ser	Pro	Thr	Leu	Thr	Ser	His
			580					585						590	

-continued

---

Ser Lys Asn Ser Thr Gly Leu Ala Glu Thr Gly Thr His Ser Asn Ser  
 595 600 605

Tyr Leu Asp Thr Ser Arg Val Ser Ile Ser Val Leu Val Pro Gly Leu  
 610 615 620

Leu Leu Val Phe Val Thr Ser Ala Phe Thr Val Val Gly Met Leu Val  
 625 630 635 640

Phe Ile Leu Arg Asn Arg Lys Arg Ser Lys Arg Arg Asp Ala Asn Ser  
 645 650 655

Ser Ala Ser Glu Ile Asn Ser Leu Gln Thr Val Cys Asp Ser Ser Tyr  
 660 665 670

Trp His Asn Gly Pro Tyr Asn Ala Asp Gly Ala His Arg Val Tyr Asp  
 675 680 685

Cys Gly Ser His Ser Leu Ser Asp  
 690 695

<210> SEQ ID NO 92  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 92  
 gttggatctg ggcaacaata ac 22

<210> SEQ ID NO 93  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 93  
 attgttgtgc aggctgagtt taag 24

<210> SEQ ID NO 94  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 94  
 ggtggctata catggatagc aattacctgg acacgctgtc ccggg 45

<210> SEQ ID NO 95  
 <211> LENGTH: 2226  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95  
 agtcgactgc gtccctgta ccggcgccca gctgtgttcc tgacccaga ataactcagg 60  
 gctgcaccgg gcctggcagc gctccgcaca catttcctgt cgcggcctaa gggaaactgt 120  
 tggccgctgg gcccggggg ggattcttgg cagttggggg gtcogtggg agcgagggcg 180  
 gaggggaag gagggggaac cgggttgggg aagccagctg tagagggcg tgaccgcgct 240

-continued

---

```

ccagacacag ctctgctgcc tcgagcggga cagatccaag ttgggagcag ctctgctgctc 300
ggggcctcag agaatagggc cggcgttcgc cctgtgcctc ctctggcagg cgctctggcc 360
cgggcccggc ggcggcgaac accccactgc cgaccgtgct ggctgctcgg cctcgggggc 420
ctgtctacagc ctgcaccacg ctaccatgaa gcggcaggcg gccgaggagg cctgcatcct 480
gcgaggtggg gcgctcagca ccgtgctgctc gggcgccgag ctgcgcgctg tgctcgcgct 540
cctgcgggca ggcccagggc ccggaggggg ctccaaagac ctgctgttct gggctgcaact 600
ggagcgcagg cgttcccact gcaccctgga gaacgagcct ttgcggggtt tctcctggct 660
gtcctccgac cccggcggtc tcgaaagcga cacgctgcag tgggtggagg agccccaacg 720
ctcctgcacc gcgcgagat gcgcggtact ccaggccacc ggtggggtcg agcccgcagg 780
ctggaaggag atgcatgctc acctgcgcgc caacggctac ctgtgcaagt accagtttga 840
ggtcttctgt cctgcgcccgc gcccggggc gcctctaac ttgagctatc gcgcgcccctt 900
ccagctgcac agcgcgcgctc tggacttcag tccacctggg accgaggtga gtgcgctctg 960
ccggggacag ctcccgatct cagttacttg catcgcggac gaaatcggcg ctgcgctggga 1020
caaactctcg ggcgatgtgt tgtgtcccctg cccggggagg tacctccgtg ctggcaaatg 1080
cgcagagctc cctaactgcc tagacgactt gggaggcttt gcctgcgaat gtgctacggg 1140
cttcgagctg gggaaagacg gccgctcttg tgtgaccagt ggggaaggac agccgaccct 1200
tggggggacc ggggtgcccga ccaggcggcc gccggccact gcaaccagcc ccgtgcccga 1260
gagaacatgg ccaatcaggg tcgacgagaa gctgggagag acaccacttg tccctgaaca 1320
agacaattca gtaacatcta ttctgagat tctctgatgg ggatcacaga gcacgatgctc 1380
tacccttaa atgtcccttc aagccgagtc aaaggccact atcaccocat cagggagcgt 1440
gatttccaa tttaatctca cgacttcctc tgccactcct caggctttcg actcctcctc 1500
tgccgtggtc ttcatatttg tgagcacagc agtagtagtg ttggtgatct tgaccatgac 1560
agtactgggg cttgtcaagc tctgcttca cgaaagcccc tcttcccagc caaggaagga 1620
gtctatggc ccgcccggcc tggagagtga tctctgagccc gctgctttgg gctccagttc 1680
tgcacattgc acaacaatg gggtgaaagt cggggactgt gatctgcggg acagagcaga 1740
gggtgccttg ctggcggagt cccctcttg ctctagtat gcatagggaa acaggggaca 1800
tgggcactcc tgtgaacagt ttttacttt tgatgaaacg gggaaaccaag aggaacttac 1860
ttgtgtaact gacaatttct gcagaaatcc ccttctctct aaattccctt tactccactg 1920
aggagctaaa tcagaactgc aactccttc cctgatgata gaggaagtgg aagtgccttt 1980
aggatggtga tactggggga ccgggtagt ctggggagag atattttctt atgtttattc 2040
ggagaatttg gagaagtgat tgaacttttc aagacattgg aaacaaatag aacacaatat 2100
aatttaccat aaaaaataat ttctacaaa atggaagga aatgttctat gttgttcagg 2160
ctaggagtat attggttcga aatcccaggg aaaaaataa aaataaaaaa ttaaaggatt 2220
gttgat 2226

```

&lt;210&gt; SEQ ID NO 96

&lt;211&gt; LENGTH: 490

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 96

-continued

---

Met Arg Pro Ala Phe Ala Leu Cys Leu Leu Trp Gln Ala Leu Trp Pro  
1 5 10 15  
Gly Pro Gly Gly Gly Glu His Pro Thr Ala Asp Arg Ala Gly Cys Ser  
20 25 30  
Ala Ser Gly Ala Cys Tyr Ser Leu His His Ala Thr Met Lys Arg Gln  
35 40 45  
Ala Ala Glu Glu Ala Cys Ile Leu Arg Gly Gly Ala Leu Ser Thr Val  
50 55 60  
Arg Ala Gly Ala Glu Leu Arg Ala Val Leu Ala Leu Arg Ala Gly  
65 70 75 80  
Pro Gly Pro Gly Gly Ser Lys Asp Leu Leu Phe Trp Val Ala Leu  
85 90 95  
Glu Arg Arg Arg Ser His Cys Thr Leu Glu Asn Glu Pro Leu Arg Gly  
100 105 110  
Phe Ser Trp Leu Ser Ser Asp Pro Gly Gly Leu Glu Ser Asp Thr Leu  
115 120 125  
Gln Trp Val Glu Glu Pro Gln Arg Ser Cys Thr Ala Arg Arg Cys Ala  
130 135 140  
Val Leu Gln Ala Thr Gly Gly Val Glu Pro Ala Gly Trp Lys Glu Met  
145 150 155 160  
Arg Cys His Leu Arg Ala Asn Gly Tyr Leu Cys Lys Tyr Gln Phe Glu  
165 170 175  
Val Leu Cys Pro Ala Pro Arg Pro Gly Ala Ala Ser Asn Leu Ser Tyr  
180 185 190  
Arg Ala Pro Phe Gln Leu His Ser Ala Ala Leu Asp Phe Ser Pro Pro  
195 200 205  
Gly Thr Glu Val Ser Ala Leu Cys Arg Gly Gln Leu Pro Ile Ser Val  
210 215 220  
Thr Cys Ile Ala Asp Glu Ile Gly Ala Arg Trp Asp Lys Leu Ser Gly  
225 230 235 240  
Asp Val Leu Cys Pro Cys Pro Gly Arg Tyr Leu Arg Ala Gly Lys Cys  
245 250 255  
Ala Glu Leu Pro Asn Cys Leu Asp Asp Leu Gly Gly Phe Ala Cys Glu  
260 265 270  
Cys Ala Thr Gly Phe Glu Leu Gly Lys Asp Gly Arg Ser Cys Val Thr  
275 280 285  
Ser Gly Glu Gly Gln Pro Thr Leu Gly Gly Thr Gly Val Pro Thr Arg  
290 295 300  
Arg Pro Pro Ala Thr Ala Thr Ser Pro Val Pro Gln Arg Thr Trp Pro  
305 310 315 320  
Ile Arg Val Asp Glu Lys Leu Gly Glu Thr Pro Leu Val Pro Glu Gln  
325 330 335  
Asp Asn Ser Val Thr Ser Ile Pro Glu Ile Pro Arg Trp Gly Ser Gln  
340 345 350  
Ser Thr Met Ser Thr Leu Gln Met Ser Leu Gln Ala Glu Ser Lys Ala  
355 360 365  
Thr Ile Thr Pro Ser Gly Ser Val Ile Ser Lys Phe Asn Ser Thr Thr  
370 375 380  
Ser Ser Ala Thr Pro Gln Ala Phe Asp Ser Ser Ser Ala Val Val Phe  
385 390 395 400

-continued

---

```

Ile Phe Val Ser Thr Ala Val Val Val Leu Val Ile Leu Thr Met Thr
           405                               410               415

Val Leu Gly Leu Val Lys Leu Cys Phe His Glu Ser Pro Ser Ser Gln
           420                               425               430

Pro Arg Lys Glu Ser Met Gly Pro Pro Gly Leu Glu Ser Asp Pro Glu
           435                               440               445

Pro Ala Ala Leu Gly Ser Ser Ser Ala His Cys Thr Asn Asn Gly Val
           450                               455               460

Lys Val Gly Asp Cys Asp Leu Arg Asp Arg Ala Glu Gly Ala Leu Leu
           465                               470               475               480

Ala Glu Ser Pro Leu Gly Ser Ser Asp Ala
           485                               490

```

```

<210> SEQ ID NO 97
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

```

```
<400> SEQUENCE: 97
```

```
tggaaggaga tgcgatgcca cctg 24
```

```

<210> SEQ ID NO 98
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

```

```
<400> SEQUENCE: 98
```

```
tgaccagtgg ggaaggacag 20
```

```

<210> SEQ ID NO 99
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

```

```
<400> SEQUENCE: 99
```

```
acagagcaga ggggtgccttg 20
```

```

<210> SEQ ID NO 100
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

```

```
<400> SEQUENCE: 100
```

```
tcagggacaa gtggtgtctc tccc 24
```

```

<210> SEQ ID NO 101
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

---

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 101

tcagggaaagg agtgtgcagt tctg 24

<210> SEQ ID NO 102

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 102

acagctcccg atctcagtta cttgcatcgc ggacgaaatc ggcgctcgct 50

<210> SEQ ID NO 103

<211> LENGTH: 2026

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

cggacgcgtg ggattcagca gtggcctgtg gctgccagag cagctcctca ggggaaacta 60

agcgtcgagt cagacggcac cataatcgcc tttaaaagtg cctccgccct gccggccgcg 120

tatcccccg ctacctgggc cgccccgcyg cgggtgcgcyg gtgagagggg gcgcgcgggc 180

agccgagcgc cgggtgtgagc cagcgtgct gccagtgta gcggcgggtg gagcgcggtg 240

ggtgcggagg ggcgtgtgtg ccggcgcgcyg cgcctggggg tgcaaacccc gagcgtctac 300

gctgccatga ggggcgcgaa cgcctgggcy ccaactctgcc tgctgctggc tggccacc 360

cagctctcgc ggcagcagtc cccagagaga cctgttttca catgtggtgg cattcttact 420

ggagagtctg gatttattgg cagtgaagg tttcctggag tgtaccctcc aaatagcaaa 480

tgtacttgga aaatcacagt tcccgaagga aaagtagtcg ttctcaattt ccgattcata 540

gacctcgaga gtgacaacct gtgccgctat gactttgtgg atgtgtacaa tggccatgcc 600

aatggccagc gcattggccg cttctgtggc actttccggc ctggagccct tgtgtccagt 660

ggcaacaaga tgatggtgca gatgatttct gatgccaaca cagctggcaa tggcttcag 720

gccatgttct ccgctgctga accaaacgaa agaggggatc agtattgtgg aggactcctt 780

gacagacctt ccgctctttt taaaaccccc aactggccag accgggatta cctgcagga 840

gtcacttgtyg tgtggccatc ttagccccca aagaatcagc ttatagaatt aaagtttgag 900

aaagttgatg tggagcgaga taactactgc cgatatgatt atgtggctgt gtttaatggc 960

ggggaagtca acgatgctag aagaattgga aagtattgtg gtgatagtcc acctgcgcca 1020

attgtgtctg agagaaatga acttcttatt cagtttttat cagacttaag ttttaactgca 1080

gatgggttta ttggtcacta catattcagc ccaaaaaaac tgcctacaac tacagaacag 1140

cctgtcacca ccacattccc tgtaaccacg ggtttaaaac ccaccgtggc cttgtgtcaa 1200

caaaagtgta gacggacggg gactctggag ggcaattatt gttcaagtga ctttgtatta 1260

gccggcactg ttatcacaac catcactcgc gatgggagtt tgcacgccac agtctogatc 1320

atcaacatct acaaagaggg aaatttggcg attcagcagc cgggcaagaa catgagtgcc 1380

aggctgactg tcgtctgcaa gcagtgccct ctcctcagaa gaggtctaaa ttacattatt 1440



-continued

---

```

atgggcccaag taggtgaaga tgggcgaggc aaaatcatgc caaacagctt tatcatgatg 1500
ttcaagacca agaatcagaa gtccttgat gccttaaaaa ataagcaatg ttaacagtga 1560
actgtgtcca ttaagctgt attctgccat tgcctttgaa agatctatgt tctctcagta 1620
gaaaaaaaa tacttataaa attacatatt ctgaaagagg attccgaaag atgggactgg 1680
ttgactcttc acatgatgga ggtatgagc ctccgagata gctgagggaa gttctttgcc 1740
tgctgtcaga ggagcagcta tctgattgga aacctgccga cttagtgcgg tgataggaag 1800
ctaaaagtgt caagcgttga cagcttgaa gogtttattt atacatctct gtaaaaggat 1860
attttagaat tgagttgtgt gaagatgtca aaaaaagatt ttagaagtgc aatatttata 1920
gtgttatttg tttcaccttc aagcctttgc cctgaggtgt tacaatcttg tcttgcgttt 1980
tctaatacaa tgcttaataa aatattttta aaggaaaaaa aaaaaa 2026
    
```

```

<210> SEQ ID NO 104
<211> LENGTH: 415
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 104

```

Met Arg Gly Ala Asn Ala Trp Ala Pro Leu Cys Leu Leu Leu Ala Ala
 1           5           10          15
Ala Thr Gln Leu Ser Arg Gln Gln Ser Pro Glu Arg Pro Val Phe Thr
 20          25          30
Cys Gly Gly Ile Leu Thr Gly Glu Ser Gly Phe Ile Gly Ser Glu Gly
 35          40          45
Phe Pro Gly Val Tyr Pro Pro Asn Ser Lys Cys Thr Trp Lys Ile Thr
 50          55          60
Val Pro Glu Gly Lys Val Val Val Leu Asn Phe Arg Phe Ile Asp Leu
 65          70          75          80
Glu Ser Asp Asn Leu Cys Arg Tyr Asp Phe Val Asp Val Tyr Asn Gly
 85          90          95
His Ala Asn Gly Gln Arg Ile Gly Arg Phe Cys Gly Thr Phe Arg Pro
100         105         110
Gly Ala Leu Val Ser Ser Gly Asn Lys Met Met Val Gln Met Ile Ser
115         120         125
Asp Ala Asn Thr Ala Gly Asn Gly Phe Met Ala Met Phe Ser Ala Ala
130         135         140
Glu Pro Asn Glu Arg Gly Asp Gln Tyr Cys Gly Gly Leu Leu Asp Arg
145         150         155         160
Pro Ser Gly Ser Phe Lys Thr Pro Asn Trp Pro Asp Arg Asp Tyr Pro
165         170         175
Ala Gly Val Thr Cys Val Trp His Ile Val Ala Pro Lys Asn Gln Leu
180         185         190
Ile Glu Leu Lys Phe Glu Lys Phe Asp Val Glu Arg Asp Asn Tyr Cys
195         200         205
Arg Tyr Asp Tyr Val Ala Val Phe Asn Gly Gly Glu Val Asn Asp Ala
210         215         220
Arg Arg Ile Gly Lys Tyr Cys Gly Asp Ser Pro Pro Ala Pro Ile Val
225         230         235         240
Ser Glu Arg Asn Glu Leu Leu Ile Gln Phe Leu Ser Asp Leu Ser Leu
245         250         255
    
```



-continued

---

```

cggacgcgtg ggcggacgcg tgggcggccc acggcgcccg cgggctgggg cggtcgcttc    60
ttccttctcc ttggcctacg agggctccca gcttgggtaa agatggcccc atggcccccg    120
aaggccctag tcccagctgt gctctggggc ctcagcctct tcctcaacct cccaggacct    180
atctggctcc agccctctcc acctccccag tcttctcccc cgctcagcc ccatcogtgt    240
catactctgc ggggactggt tgacagcttt aacaagggcc tggagagaac catccgggac    300
aactttggag gtggaacac tgctgggag gaagagaatt tgtccaaata caaagacagt    360
gagaccgcc tggtagaggt gctggagggt gtgtgcagca agtcagactt cgagtgccac    420
cgctgctgg agctgagtga ggagctgggt gagagctggt ggtttcacia gcagcaggag    480
gccccggacc tctccagtg gctgtgctca gattccctga agctctgctg ccccgcaggc    540
accttcgggc cctcctgctt tccctgtcct gggggaacag agaggccctg cggtggtctac    600
gggcagtgtg aaggagaagg gacacgaggg ggcagcgggc actgtgactg ccaagccggc    660
tacgggggtg aggcctgtgg ccagtgtggc cttggctact ttgaggcaga acgcaacgcc    720
agccatctgg tatgttcggc ttgttttggc ccctgtgccc gatgctcagg acctgaggaa    780
tcaaactgtt tgcaatgcaa gaaggcttg gcctgcatc acctcaagtg tgtagacatt    840
gatgagtgtg gcacagaggg agccaactgt ggagctgacc aattctgctg gaacactgag    900
ggctcctatg agtgccgaga ctgtgccaa gacctgcctag gctgcatggg ggcagggcca    960
ggtcgctgta agaagtgtag ccctggctat cagcaggtgg gctccaagtg tctcgatgtg   1020
gatgagtgtg agacagaggt gtgtccggga gagaacaagc agtgtgaaaa caccgagggc   1080
ggttatcgtc gcatctgtgc cgagggtcac aagcagatgg aaggcatctg tgtgaaggag   1140
cagatcccag agtcagcagg cttcttctca gagatgacag aagacgagtt ggtggtgctg   1200
cagcagatgt tctttggcat catcatctgt gcaactggcca cgctggctgc taagggcgac   1260
ttggtgttca ccgcatctct cattggggct gtggcggcca tgactggcta ctggttgtca   1320
gagcgcagtg accgtgtgct ggagggtctc atcaagggca gataatcgcg gccaccacct   1380
gtaggacctc ctccccacca cgctgcccc agagcttggg ctgccctcct gctggacct   1440
caggacagct tggtttattt ttgagagtgg ggtaagcacc cctacctgcc ttacagagca   1500
gcccaggtac ccaggccccg gcagacaagg cccctggggt aaaaagtagc cctgaagggtg   1560
gataccatga gctcttcacc tggcggggac tggcaggctt cacaatgtgt gaatttcaa   1620
agtttttctt taatggtggc tgctagagct ttggcccctg cttaggatta ggtggtcctc   1680
acaggggtgg ggccatcaca gctccctcct gccagctgca tgctgccagt tctgttctg   1740
tgttcaccac atccccacac cccattgcca cttatttatt catctcagga aataaagaaa   1800
ggtcttgtaa agttaaaaa aaaaaaaaa aaaaaaaa   1838

```

&lt;210&gt; SEQ ID NO 109

&lt;211&gt; LENGTH: 420

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 109

```

Met Ala Pro Trp Pro Pro Lys Gly Leu Val Pro Ala Val Leu Trp Gly
 1             5             10             15

```

```

Leu Ser Leu Phe Leu Asn Leu Pro Gly Pro Ile Trp Leu Gln Pro Ser
 20             25             30

```

-continued

---

Pro Pro Pro Gln Ser Ser Pro Pro Pro Gln Pro His Pro Cys His Thr  
 35 40 45  
 Cys Arg Gly Leu Val Asp Ser Phe Asn Lys Gly Leu Glu Arg Thr Ile  
 50 55 60  
 Arg Asp Asn Phe Gly Gly Asp Thr Ala Trp Glu Glu Glu Asn Leu  
 65 70 75 80  
 Ser Lys Tyr Lys Asp Ser Glu Thr Arg Leu Val Glu Val Leu Glu Gly  
 85 90 95  
 Val Cys Ser Lys Ser Asp Phe Glu Cys His Arg Leu Leu Glu Leu Ser  
 100 105 110  
 Glu Glu Leu Val Glu Ser Trp Trp Phe His Lys Gln Gln Glu Ala Pro  
 115 120 125  
 Asp Leu Phe Gln Trp Leu Cys Ser Asp Ser Leu Lys Leu Cys Cys Pro  
 130 135 140  
 Ala Gly Thr Phe Gly Pro Ser Cys Leu Pro Cys Pro Gly Gly Thr Glu  
 145 150 155 160  
 Arg Pro Cys Gly Gly Tyr Gly Gln Cys Glu Gly Glu Gly Thr Arg Gly  
 165 170 175  
 Gly Ser Gly His Cys Asp Cys Gln Ala Gly Tyr Gly Gly Glu Ala Cys  
 180 185 190  
 Gly Gln Cys Gly Leu Gly Tyr Phe Glu Ala Glu Arg Asn Ala Ser His  
 195 200 205  
 Leu Val Cys Ser Ala Cys Phe Gly Pro Cys Ala Arg Cys Ser Gly Pro  
 210 215 220  
 Glu Glu Ser Asn Cys Leu Gln Cys Lys Lys Gly Trp Ala Leu His His  
 225 230 235 240  
 Leu Lys Cys Val Asp Ile Asp Glu Cys Gly Thr Glu Gly Ala Asn Cys  
 245 250 255  
 Gly Ala Asp Gln Phe Cys Val Asn Thr Glu Gly Ser Tyr Glu Cys Arg  
 260 265 270  
 Asp Cys Ala Lys Ala Cys Leu Gly Cys Met Gly Ala Gly Pro Gly Arg  
 275 280 285  
 Cys Lys Lys Cys Ser Pro Gly Tyr Gln Gln Val Gly Ser Lys Cys Leu  
 290 295 300  
 Asp Val Asp Glu Cys Glu Thr Glu Val Cys Pro Gly Glu Asn Lys Gln  
 305 310 315 320  
 Cys Glu Asn Thr Glu Gly Gly Tyr Arg Cys Ile Cys Ala Glu Gly Tyr  
 325 330 335  
 Lys Gln Met Glu Gly Ile Cys Val Lys Glu Gln Ile Pro Glu Ser Ala  
 340 345 350  
 Gly Phe Phe Ser Glu Met Thr Glu Asp Glu Leu Val Val Leu Gln Gln  
 355 360 365  
 Met Phe Phe Gly Ile Ile Ile Cys Ala Leu Ala Thr Leu Ala Ala Lys  
 370 375 380  
 Gly Asp Leu Val Phe Thr Ala Ile Phe Ile Gly Ala Val Ala Ala Met  
 385 390 395 400  
 Thr Gly Tyr Trp Leu Ser Glu Arg Ser Asp Arg Val Leu Glu Gly Phe  
 405 410 415  
 Ile Lys Gly Arg  
 420

-continued

---

```

<210> SEQ ID NO 110
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 110

cctggctatc agcaggtggg ctccaagtgt ctcgatgtgg atgagtgtga      50

<210> SEQ ID NO 111
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 111

attctgcgtg aacctgagg gc      22

<210> SEQ ID NO 112
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 112

atctgcttgt agccctcggc ac      22

<210> SEQ ID NO 113
<211> LENGTH: 1616
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (1461)
<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 113

tgagaccctc ctgcagcctt ctcaaggac agccccactc tgcctettgc tcctccaggg      60
cagcaccatg cagcccctgt ggctctgctg ggcactctgg gtgttgcccc tggccagccc      120
cggggccgcc ctgaccgggg agcagctcct gggcagcctg ctgcggcagc tgcagctcaa      180
agaggtgccc accttgaca gggccgacat ggaggagctg gtcatcccca cccacgtgag      240
ggcccagtag gtggccctgc tgcagcgcag ccacggggac cgctcccgcg gaaagaggtt      300
cagccagagc ttccgagagg tggccggcag gttcctggcg ttggaggcca gcacacacct      360
gtggtgttcc ggcatggagc agcggctgcc gcccaacagc gagctggtgc aggccgtgct      420
gcggctcttc caggagccgg tccccaggc cgcgctgcac aggcacgggc ggctgtcccc      480
gcgcagcggc cgggcccggg tgaccgtcga gtggctgcgc gtcocgcagc acggctccaa      540
ccgcacctcc ctcatcgact ccaggtggt gtccgtccac gagagcggct ggaaggcctt      600
cgacgtgacc gaggccgtga acttctggca gcagctgagc cggccccggc agccgctgct      660
gctacaggtg tcggtgcaga gggagcatct gggcccgtg gcgtccggcg cccacaagct      720
ggtccgcgtt gcctcgcagg gggcgccagc cgggcttggg gagccccagc tggagctgca      780

```

-continued

---

```

caccctggac cttggggact atggagctca gggcgactgt gaccctgaag caccaatgac 840
cgagggcacc cgctgctgcc gccaggagat gtacattgac ctgcagggga tgaagtgggc 900
cgagaaactgg gtgctggagc ccccgggctt cctggcttat gagggtgtgg gcacctgccg 960
gcagcccccg gaggccctgg ccttcaagtg gccgtttctg gggcctcgac agtgcacgcg 1020
ctcggagact gactcgctgc ccatgatcgt cagcatcaag gaggagggca ggaccaggcc 1080
ccaggtggtc agcctgccc acatgagggt gcagaagtgc agctgtgcct cggatggtgc 1140
gctcgtgcca aggaggctcc agccatagc gcctagtgtta gccatcgagg gacttgactt 1200
gtgtgtgttt ctgaagtgtt cgaggggtacc aggagagctg gcatgactg aactgctgat 1260
ggacaaatgc tctgtgctct ctagttagcc ctgaattgc ttctctgac aagttacctc 1320
acctaatttt tgcttctcag gaatgagaat ctttggccac tggagagccc ttgctcagtt 1380
ttctctattc ttattattca ctgcactata ttctaagcac ttacatgtgg agatactgta 1440
acctgagggc agaaagccca ntgtgtcatt gtttacttgt cctgtcactg gatctgggct 1500
aaagtcctcc accaccactc tggacctaag acctgggggt aagtgtgggt tgtgcacccc 1560
caatccagat aataaagact ttgtaaaaca tgaataaaac acattttatt ctaaaa 1616

```

&lt;210&gt; SEQ ID NO 114

&lt;211&gt; LENGTH: 366

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 114

```

Met  Gln  Pro  Leu  Trp  Leu  Cys  Trp  Ala  Leu  Trp  Val  Leu  Pro  Leu  Ala
  1          5          10          15
Ser  Pro  Gly  Ala  Ala  Leu  Thr  Gly  Glu  Gln  Leu  Leu  Gly  Ser  Leu  Leu
          20          25          30
Arg  Gln  Leu  Gln  Leu  Lys  Glu  Val  Pro  Thr  Leu  Asp  Arg  Ala  Asp  Met
          35          40          45
Glu  Glu  Leu  Val  Ile  Pro  Thr  His  Val  Arg  Ala  Gln  Tyr  Val  Ala  Leu
          50          55          60
Leu  Gln  Arg  Ser  His  Gly  Asp  Arg  Ser  Arg  Gly  Lys  Arg  Phe  Ser  Gln
          65          70          75          80
Ser  Phe  Arg  Glu  Val  Ala  Gly  Arg  Phe  Leu  Ala  Leu  Glu  Ala  Ser  Thr
          85          90          95
His  Leu  Leu  Val  Phe  Gly  Met  Glu  Gln  Arg  Leu  Pro  Pro  Asn  Ser  Glu
          100          105          110
Leu  Val  Gln  Ala  Val  Leu  Arg  Leu  Phe  Gln  Glu  Pro  Val  Pro  Lys  Ala
          115          120          125
Ala  Leu  His  Arg  His  Gly  Arg  Leu  Ser  Pro  Arg  Ser  Ala  Arg  Ala  Arg
          130          135          140
Val  Thr  Val  Glu  Trp  Leu  Arg  Val  Arg  Asp  Asp  Gly  Ser  Asn  Arg  Thr
          145          150          155          160
Ser  Leu  Ile  Asp  Ser  Arg  Leu  Val  Ser  Val  His  Glu  Ser  Gly  Trp  Lys
          165          170          175
Ala  Phe  Asp  Val  Thr  Glu  Ala  Val  Asn  Phe  Trp  Gln  Gln  Leu  Ser  Arg
          180          185          190
Pro  Arg  Gln  Pro  Leu  Leu  Leu  Gln  Val  Ser  Val  Gln  Arg  Glu  His  Leu
          195          200          205

```

-continued

---

Gly Pro Leu Ala Ser Gly Ala His Lys Leu Val Arg Phe Ala Ser Gln  
 210 215 220

Gly Ala Pro Ala Gly Leu Gly Glu Pro Gln Leu Glu Leu His Thr Leu  
 225 230 235 240

Asp Leu Gly Asp Tyr Gly Ala Gln Gly Asp Cys Asp Pro Glu Ala Pro  
 245 250 255

Met Thr Glu Gly Thr Arg Cys Cys Arg Gln Glu Met Tyr Ile Asp Leu  
 260 265 270

Gln Gly Met Lys Trp Ala Glu Asn Trp Val Leu Glu Pro Pro Gly Phe  
 275 280 285

Leu Ala Tyr Glu Cys Val Gly Thr Cys Arg Gln Pro Pro Glu Ala Leu  
 290 295 300

Ala Phe Lys Trp Pro Phe Leu Gly Pro Arg Gln Cys Ile Ala Ser Glu  
 305 310 315 320

Thr Asp Ser Leu Pro Met Ile Val Ser Ile Lys Glu Gly Gly Arg Thr  
 325 330 335

Arg Pro Gln Val Val Ser Leu Pro Asn Met Arg Val Gln Lys Cys Ser  
 340 345 350

Cys Ala Ser Asp Gly Ala Leu Val Pro Arg Arg Leu Gln Pro  
 355 360 365

<210> SEQ ID NO 115  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 115

aggactgccca taacttgctct g 21

<210> SEQ ID NO 116  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 116

ataggagttg aagcagcgcct gc 22

<210> SEQ ID NO 117  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 117

tgtgtggaca tagacgagtg ccgctaccgc tactgccagc accgc 45

<210> SEQ ID NO 118  
 <211> LENGTH: 1857  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 118

-continued

---

```

gtctgttccc aggagtcctt cggcggctgt tgtgtcagtg gcctgatcgc gatggggaca    60
aaggcgcaag tcgagaggaa actgttgtgc ctcttcatat tggcgatcct gttgtgctcc    120
ctggcattgg gcagtgttac agtgactctt tctgaacctg aagtcagaat tcctgagaat    180
aatcctgtga agttgtcctg tgccactcgc ggcttttctt ctccccgtgt ggagtggaa    240
tttgaccaag gagacaccac cagactcgtt tgctataata acaagatcac agcttctctat    300
gaggaccggg tgaccttctt gccaaactgt atcacctca agtccgtgac acgggaagac    360
actgggacat acacttgtat ggtctctgag gaaggcggca acagctatgg ggaggtaag    420
gtcaagctca tcgtgcttgt gcctccatcc aagcctacag ttaacatccc ctccctgccc    480
accattggga accgggcagt gctgacatgc tcagaacaag atgggtcccc acctctgaa    540
tacacctggt tcaaagatgg gatagtgatg cctacgaatc ccaaaagcac ccgtgccttc    600
agcaactctt cctatgtcct gaatcccaca acaggagagc tggctcttga tccccgtca    660
gcctctgata ctggagaata cagctgtgag gcacggaatg ggtatgggac acccatgact    720
tcaaagtctg tgcgcatgga agctgtggag cggaatgtgg gggatcatcgt ggcagccgtc    780
cttgaacacc tgattctcct gggaatcttg gtttttgcca tctggtttgc ctatagccga    840
ggccactttg acagaacaaa gaaagggact tcgagtaaga aggtgattta cagccagcct    900
agtgcctgaa gtgaaggaga attcaaacag acctcgtcat tcctgggtgtg agcctggctg    960
gctcaccgoc tatcatctgc atttgcccta ctacaggtgct accggactct gccccctgat   1020
gtctgtagtt tcacaggatg ccttattttg cttctacacc ccacagggcc cctacttct   1080
tcggatgtgt ttttaataat gtcagctatg tgccccatcc tccttcctgc cctccctccc   1140
tttctacca ctgctgagtg gcctggaact tgtttaaagt gtttattccc catttctttg   1200
agggatcagg aaggaatcct gggtatgcca ttgacttccc ttctaagtag acagcaaaaa   1260
tggcgggggt cgcaggaatc tgcaactcaac tgcccactct gctggcaggg atctttgaa   1320
aggatctctg agcttggttc tgggctcttt ccttgtgtac tgacgaccag ggcagctgt   1380
tctagagcgg gaattagagg cttagcgggc tgaaatggtt gtttggatgac gacactgggg   1440
tccttccatc tctggggccc actctcttct gtcttcccat gggagtgcc actgggatcc   1500
ctctgccctg tcctcctgaa tacaagctga ctgacattga ctgtgtctgt ggaaaatggg   1560
agctcttgtt gtggagagca tagtaaatth tcagagaact tgaagccaaa aggatttaaa   1620
accgctgctc taaagaaaag aaaactggag gctgggcgca gtggctcacg cctgtaatcc   1680
cagaggctga ggcagggcga tcacctgagg tcgggagttc gggatcagcc tgaccaacat   1740
ggagaaaccc tactggaat acaaagttag ccaggcatgg tggatcatgc ctgtagtccc   1800
agctgctcag gagcctggca acaagagcaa aactccagct caaaaaaaaa aaaaaaa   1857

```

&lt;210&gt; SEQ ID NO 119

&lt;211&gt; LENGTH: 299

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 119

```

Met Gly Thr Lys Ala Gln Val Glu Arg Lys Leu Leu Cys Leu Phe Ile
  1             5             10             15

```

```

Leu Ala Ile Leu Leu Cys Ser Leu Ala Leu Gly Ser Val Thr Val His
      20             25             30

```



-continued

---

Ser Ser Glu Pro Glu Val Arg Ile Pro Glu Asn Asn Pro Val Lys Leu  
 35 40 45

Ser Cys Ala Tyr Ser Gly Phe Ser Ser Pro Arg Val Glu Trp Lys Phe  
 50 55 60

Asp Gln Gly Asp Thr Thr Arg Leu Val Cys Tyr Asn Asn Lys Ile Thr  
 65 70 75 80

Ala Ser Tyr Glu Asp Arg Val Thr Phe Leu Pro Thr Gly Ile Thr Phe  
 85 90 95

Lys Ser Val Thr Arg Glu Asp Thr Gly Thr Tyr Thr Cys Met Val Ser  
 100 105 110

Glu Glu Gly Gly Asn Ser Tyr Gly Glu Val Lys Val Lys Leu Ile Val  
 115 120 125

Leu Val Pro Pro Ser Lys Pro Thr Val Asn Ile Pro Ser Ser Ala Thr  
 130 135 140

Ile Gly Asn Arg Ala Val Leu Thr Cys Ser Glu Gln Asp Gly Ser Pro  
 145 150 155 160

Pro Ser Glu Tyr Thr Trp Phe Lys Asp Gly Ile Val Met Pro Thr Asn  
 165 170 175

Pro Lys Ser Thr Arg Ala Phe Ser Asn Ser Ser Tyr Val Leu Asn Pro  
 180 185 190

Thr Thr Gly Glu Leu Val Phe Asp Pro Leu Ser Ala Ser Asp Thr Gly  
 195 200 205

Glu Tyr Ser Cys Glu Ala Arg Asn Gly Tyr Gly Thr Pro Met Thr Ser  
 210 215 220

Asn Ala Val Arg Met Glu Ala Val Glu Arg Asn Val Gly Val Ile Val  
 225 230 235 240

Ala Ala Val Leu Val Thr Leu Ile Leu Leu Gly Ile Leu Val Phe Gly  
 245 250 255

Ile Trp Phe Ala Tyr Ser Arg Gly His Phe Asp Arg Thr Lys Lys Gly  
 260 265 270

Thr Ser Ser Lys Lys Val Ile Tyr Ser Gln Pro Ser Ala Arg Ser Glu  
 275 280 285

Gly Glu Phe Lys Gln Thr Ser Ser Phe Leu Val  
 290 295

<210> SEQ ID NO 120  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 120

tgcgagagct gtgttctggt tccc

24

<210> SEQ ID NO 121  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 121

-continued

---

 tgatcgcgat ggggacaaag gcgcaagctc gagaggaaac tgttgtgcct 50

<210> SEQ ID NO 122  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

&lt;400&gt; SEQUENCE: 122

acacctgggtt caaagatggg 20

<210> SEQ ID NO 123  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

&lt;400&gt; SEQUENCE: 123

taggaagagt tgctgaagc acgg 24

<210> SEQ ID NO 124  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

&lt;400&gt; SEQUENCE: 124

ttgccttact caggtgctac 20

<210> SEQ ID NO 125  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

&lt;400&gt; SEQUENCE: 125

actcagcagt gtaggaaag 20

<210> SEQ ID NO 126  
 <211> LENGTH: 1210  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 126

cagcgcgtgg ccggcgcgcg tgtggggaca gcatgagcgg cggttggatg gcgcagggtg 60

gagcgtggcg aacaggggct ctgggcctgg cgctgctgct gctgctcggc ctcgactag 120

gcctggaggc cgccgcgagc ccgctttcca ccccagctc tgcccaggcc gcaggcccca 180

gtcaggctc gtgcccacc accaagttcc agtgccgac cagtggctta tgcgtgcccc 240

tcacctggcg ctgacacagg gacttgact gcagcgatgg cagcgatgag gaggagtgca 300

ggattgagcc atgtaccag aaagggcaat gccaccgcc ccctggctc cctgcccct 360

gcaccggcgt cagtgactgc tctgggggaa ctgacaagaa actgcgcaac tgcagccgcc 420

-continued

```

tggcctgcct agcaggcgag ctccgttgca cgctgagcga tgactgcatt ccactcacgt 480
ggcgctgcga cggccacca gactgtcccg actccagcga cgagctcggc tgggaacca 540
atgagatcct cccggaaggg gatgccacaa ccatggggcc ccctgtgacc ctggagagtg 600
tcacctctct caggaatgcc acaaccatgg gggcccctgt gaccctggag agtgtcccct 660
ctgtcgggaa tgccacatcc tcctctgccg gagaccagtc tggaagccca actgcctatg 720
gggttattgc agctgctgcg gtgctcagtg caagcctggg caccgccacc ctccctcttt 780
tgtcctggct ccgagcccag gagcgctcc gccactggg gttactggtg gccatgaagg 840
agtcctctgt gctgtcagaa cagaagacct cgctgccctg aggacaagca cttgccacca 900
ccgtactca gccctggggc tagccggaca ggaggagagc agtgatgagg atgggtaccc 960
gggcacacca gccctcagag acctgagttc ttctggccac gtggaacctc gaaccogagc 1020
tcctgcagaa gtggccctgg agattgaggg tccctggaca ctccctatgg agatccgggg 1080
agctaggatg gggaacctgc cacagccaga actgaggggc tggcccagg cagctcccag 1140
ggggtagaac gccctgtgic ttaagacct cctgctgcc ccgtctgagg gtggcgatta 1200
aagttgcttc 1210
    
```

```

<210> SEQ ID NO 127
<211> LENGTH: 282
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 127

```

Met Ser Gly Gly Trp Met Ala Gln Val Gly Ala Trp Arg Thr Gly Ala
 1           5           10          15
Leu Gly Leu Ala Leu Leu Leu Leu Leu Gly Leu Gly Leu Gly Leu Glu
 20          25          30
Ala Ala Ala Ser Pro Leu Ser Thr Pro Thr Ser Ala Gln Ala Ala Gly
 35          40          45
Pro Ser Ser Gly Ser Cys Pro Pro Thr Lys Phe Gln Cys Arg Thr Ser
 50          55          60
Gly Leu Cys Val Pro Leu Thr Trp Arg Cys Asp Arg Asp Leu Asp Cys
 65          70          75          80
Ser Asp Gly Ser Asp Glu Glu Glu Cys Arg Ile Glu Pro Cys Thr Gln
 85          90          95
Lys Gly Gln Cys Pro Pro Pro Pro Gly Leu Pro Cys Pro Cys Thr Gly
100          105          110
Val Ser Asp Cys Ser Gly Gly Thr Asp Lys Lys Leu Arg Asn Cys Ser
115          120          125
Arg Leu Ala Cys Leu Ala Gly Glu Leu Arg Cys Thr Leu Ser Asp Asp
130          135          140
Cys Ile Pro Leu Thr Trp Arg Cys Asp Gly His Pro Asp Cys Pro Asp
145          150          155          160
Ser Ser Asp Glu Leu Gly Cys Gly Thr Asn Glu Ile Leu Pro Glu Gly
165          170          175
Asp Ala Thr Thr Met Gly Pro Pro Val Thr Leu Glu Ser Val Thr Ser
180          185          190
Leu Arg Asn Ala Thr Thr Met Gly Pro Pro Val Thr Leu Glu Ser Val
195          200          205
Pro Ser Val Gly Asn Ala Thr Ser Ser Ser Ala Gly Asp Gln Ser Gly
    
```

-continued

210	215	220	
Ser Pro Thr Ala Tyr Gly Val Ile Ala Ala Ala Val Leu Ser Ala			
225	230	235	240
Ser Leu Val Thr Ala Thr Leu Leu Leu Leu Ser Trp Leu Arg Ala Gln			
	245	250	255
Glu Arg Leu Arg Pro Leu Gly Leu Leu Val Ala Met Lys Glu Ser Leu			
	260	265	270
Leu Leu Ser Glu Gln Lys Thr Ser Leu Pro			
	275	280	
<210> SEQ ID NO 128			
<211> LENGTH: 24			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 128			
aagttccagt gccgcaccag tggc			24
<210> SEQ ID NO 129			
<211> LENGTH: 24			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 129			
ttggttccac agccgagctc gtcg			24
<210> SEQ ID NO 130			
<211> LENGTH: 50			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe			
<400> SEQUENCE: 130			
gaggaggagt gcaggattga gccatgtacc cagaaagggc aatgccacc			50
<210> SEQ ID NO 131			
<211> LENGTH: 1843			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<220> FEATURE:			
<221> NAME/KEY: modified_base			
<222> LOCATION: (1837)			
<223> OTHER INFORMATION: a, t, c or g			
<400> SEQUENCE: 131			
cccacgcgctc cggctctcgtcg ctctcgcgca gcggcggcag cagaggtcgc gcacagatgc			60
gggttagact ggcgggggga ggaggcggag gaggaagga agctgcatgc atgagacca			120
cagactcttg caagctggat gccctctgtg gatgaaagat gtatcatgga atgaaccga			180
gcaattggaga tggatttcta gagcagcagc agcagcagca gcaacctcag tccccccaga			240
gactcttggc cgtgatcctg tggtttcagc tggcgtgtg cttoggccct gcacagctca			300
cgggcggggtt cgatgacctt caagtgtgtg ctgaccccg cttccccgag aatggttca			360

-continued

---

```

ggacccccag cggagggggt ttctttgaag gctctgtagc cggatttcac tgccaagacg 420
gattcaagct gaagggcgct acaaagagac tgtgtttgaa gcattttaat ggaaccctag 480
gctggatccc aagtataat tccatctgtg tgcaagaaga ttgccgtatc cctcaaatcg 540
aagatgctga gattcataac aagacatata gacatggaga gaagctaata atcacttgct 600
atgaagatt caagatccg taccocgacc tacacaatat ggtttcatta tgtcgcgatg 660
atggaacgtg gaataatctg cccatctgtc aaggtgcct gagacctcta gcctcttcta 720
atggctatgt aaacatctct gagctccaga cctccttccc ggtggggact gtgatctcct 780
atcgtctgct tcccggattt aaacttgatg ggtctgcgta tcttgagtgc ttacaaaacc 840
ttatctggtc gtccagccca ccccggtgcc ttgctctgga agcccaagtc tgtccactac 900
ctccaatggt gactcagga gatttctgtc gccaccocg gccttgtagc cgctacaacc 960
acggaactgt ggtggagttt tactgcgacg ctggctacag cctcaccagc gactacaagt 1020
acatcacctg ccagtatgga gactggttct cttcttatca agtctactgc atcaaatcag 1080
agcaaacgtg gccagcacc catgagacc tctgaccac gtggaagatt gtggcgttca 1140
cggcaaccag tgtgctgctg gtgctgctgc togtatcct gccaggatg ttccagacca 1200
agttcaagc ccactttccc cccagggggc ctcccggag ttccagcagt gaccctgact 1260
ttgtggtggt agacggcgtg cccgtcatgc tcccgtccta tgacgaagct gtgagtggcg 1320
gcttgagtgc cttaggcccc gggatcatgg cctctgtggg ccagggtgct cccttaccog 1380
tggacgacca gagccccca gcataccccg gctcagggga cacggacaca gcccagggg 1440
agtcagaaac ctgtgacagc gtctcaggct cttctgagct gctccaaagt ctgtattcac 1500
ctcccaggtg ccaagagagc acccaccctg cttcggacaa cctgacata attgccagca 1560
cggcagagga ggtggcatcc accagcccag gcatacatca tgcccactgg gtgtgttcc 1620
taagaaactg attgattaaa aaatttccca aagtgtcctg aagtgtctct tcaatacat 1680
gttgatctgt ggagttgatt ctttctctc tcttggtttt agacaaatgt aaacaaagct 1740
ctgatcctta aaattgctat gctgatagag tggtagggc tggaagcttg atcaagtctt 1800
gtttcttctt gacacagact gataaaaaat taaaagnaaa aaa 1843

```

&lt;210&gt; SEQ ID NO 132

&lt;211&gt; LENGTH: 490

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 132

```

Met Tyr His Gly Met Asn Pro Ser Asn Gly Asp Gly Phe Leu Glu Gln
 1             5             10             15
Gln Gln Gln Gln Gln Gln Pro Gln Ser Pro Gln Arg Leu Leu Ala Val
          20             25             30
Ile Leu Trp Phe Gln Leu Ala Leu Cys Phe Gly Pro Ala Gln Leu Thr
      35             40             45
Gly Gly Phe Asp Asp Leu Gln Val Cys Ala Asp Pro Gly Ile Pro Glu
      50             55             60
Asn Gly Phe Arg Thr Pro Ser Gly Gly Val Phe Phe Glu Gly Ser Val
      65             70             75             80
Ala Arg Phe His Cys Gln Asp Gly Phe Lys Leu Lys Gly Ala Thr Lys
      85             90             95

```

-continued

---

Arg Leu Cys Leu Lys His Phe Asn Gly Thr Leu Gly Trp Ile Pro Ser  
 100 105 110

Asp Asn Ser Ile Cys Val Gln Glu Asp Cys Arg Ile Pro Gln Ile Glu  
 115 120 125

Asp Ala Glu Ile His Asn Lys Thr Tyr Arg His Gly Glu Lys Leu Ile  
 130 135 140

Ile Thr Cys His Glu Gly Phe Lys Ile Arg Tyr Pro Asp Leu His Asn  
 145 150 155 160

Met Val Ser Leu Cys Arg Asp Asp Gly Thr Trp Asn Asn Leu Pro Ile  
 165 170 175

Cys Gln Gly Cys Leu Arg Pro Leu Ala Ser Ser Asn Gly Tyr Val Asn  
 180 185 190

Ile Ser Glu Leu Gln Thr Ser Phe Pro Val Gly Thr Val Ile Ser Tyr  
 195 200 205

Arg Cys Phe Pro Gly Phe Lys Leu Asp Gly Ser Ala Tyr Leu Glu Cys  
 210 215 220

Leu Gln Asn Leu Ile Trp Ser Ser Ser Pro Pro Arg Cys Leu Ala Leu  
 225 230 235 240

Glu Ala Gln Val Cys Pro Leu Pro Pro Met Val Ser His Gly Asp Phe  
 245 250 255

Val Cys His Pro Arg Pro Cys Glu Arg Tyr Asn His Gly Thr Val Val  
 260 265 270

Glu Phe Tyr Cys Asp Pro Gly Tyr Ser Leu Thr Ser Asp Tyr Lys Tyr  
 275 280 285

Ile Thr Cys Gln Tyr Gly Glu Trp Phe Pro Ser Tyr Gln Val Tyr Cys  
 290 295 300

Ile Lys Ser Glu Gln Thr Trp Pro Ser Thr His Glu Thr Leu Leu Thr  
 305 310 315 320

Thr Trp Lys Ile Val Ala Phe Thr Ala Thr Ser Val Leu Leu Val Leu  
 325 330 335

Leu Leu Val Ile Leu Ala Arg Met Phe Gln Thr Lys Phe Lys Ala His  
 340 345 350

Phe Pro Pro Arg Gly Pro Pro Arg Ser Ser Ser Ser Asp Pro Asp Phe  
 355 360 365

Val Val Val Asp Gly Val Pro Val Met Leu Pro Ser Tyr Asp Glu Ala  
 370 375 380

Val Ser Gly Gly Leu Ser Ala Leu Gly Pro Gly Tyr Met Ala Ser Val  
 385 390 395 400

Gly Gln Gly Cys Pro Leu Pro Val Asp Asp Gln Ser Pro Pro Ala Tyr  
 405 410 415

Pro Gly Ser Gly Asp Thr Asp Thr Gly Pro Gly Glu Ser Glu Thr Cys  
 420 425 430

Asp Ser Val Ser Gly Ser Ser Glu Leu Leu Gln Ser Leu Tyr Ser Pro  
 435 440 445

Pro Arg Cys Gln Glu Ser Thr His Pro Ala Ser Asp Asn Pro Asp Ile  
 450 455 460

Ile Ala Ser Thr Ala Glu Glu Val Ala Ser Thr Ser Pro Gly Ile His  
 465 470 475 480

His Ala His Trp Val Leu Phe Leu Arg Asn  
 485 490

-continued

---

```

<210> SEQ ID NO 133
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 133

atctcctatc gctgctttcc cgg                                     23

<210> SEQ ID NO 134
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 134

agccaggatc gcagtaaaac tcc                                     23

<210> SEQ ID NO 135
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 135

atttaaacctt gatgggtctg cgtatcttga gtgcttaca aaccttatct    50

<210> SEQ ID NO 136
<211> LENGTH: 1815
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 136

cccacgcgtc cgctccgcgc cctccccccc gcctcccgtg cggtcgcgctg gtggcctaga    60
gatgctgctg ccgcggttgc agttgtcgcg cacgcctctg cccgccagcc cgctccaccg    120
ccgtagcgcc cgagtgtcgg ggggcgcacc cgagtcgggc catgaggccg ggaaccgcgc    180
tacaggccgt gctgctggcc gtgctgctgg tggggctgcg ggccgcgacg ggtcgcctgc    240
tgagtgcctc ggatttgagc ctcagaggag ggcagccagt ctgccgggga gggacacaga    300
ggccttgtta taaagtcatt tacttccatg atacttctcg aagactgaac tttgaggaag    360
ccaaagaagc ctgcaggagg gatggaggcc agctagttag catcagagtct gaagatgaac    420
agaaactgat agaaaagtcc attgaaaacc tcttgccatc tgatggtgac tcttggtattg    480
ggctcaggag gcgtgaggag aaacaagca atagcacagc ctgccaggac ctttatgctt    540
ggactgatgg cagcatatca caatttagga actggtatgt ggatgagccg tcctgcggca    600
gcgaggctct cgtggtcatg taccatcagc catcggcacc cgctggcacc ggaggcccct    660
acatgttcca gtggaatgat gaccggtgca acatgaagaa caatttcatt tgcaaatatt    720
ctgatgagaa accagcagtt ccttctagag aagctgaagg tgaggaaaca gagctgacaa    780
cacctgtact tccagaagaa acacaggaag aagatgcaa aaaaacattt aaagaaagta    840
gagaagctgc cttgaatctg gcctacatcc taatccccag cattcccctt ctccctctcc    900

```

-continued

---

```

ttgtggtcac cacagttgta tgttgggttt ggatctgtag aaaaagaaaa cgggagcagc   960
cagaccctag cacaaagaag caacacacca tctggccctc tcctcaccag gaaacagcc   1020
cggacctaga ggtctacaat gtcataagaa aacaaagcga agctgactta gctgagacct   1080
ggccagacct gaagaatatt tcattccgag tgtgttcggg agaagccact cccgatgaca   1140
tgtcttgtga ctatgacaac atggctgtga acccatcaga aagtgggttt gtgactctgg   1200
tgagcgtgga gagtggattt gtgaccaatg acatttatga gttctcccca gaccaaattg   1260
ggaggagtaa ggagtctgga tgggtggaaa atgaaatata tggttattag gacatataaa   1320
aaactgaaac tgacaacaat ggaaaagaaa tgataagcaa aatcctctta ttttctataa   1380
ggaaaataca cagaaggctc atgaacaagc ttagatcagg tcctgtggat gagcatgtgg   1440
tcccacgacg ctctgttgg acccccactg tttggctgta tcctttatcc cagccagtca   1500
tccagctcga ctttatgaga aggtaccttg cccaggtctg gcacatagta gagtctcaat   1560
aaatgtcact tggttgggtg tatctaactt ttaagggaca gagctttacc tggcagtgat   1620
aaagatgggc tgtggagcct ggaaaaccac ctctgttttc cttgctctat acagcagcac   1680
atattatcat acagacagaa aatccagaat cttttcaaag cccacatatg gtagcacagg   1740
ttggcctgtg catcgccaat tctcatatct gtttttttca aagaataaaa tcaataaag   1800
agcaggaaaa aaaaa                                     1815

```

&lt;210&gt; SEQ ID NO 137

&lt;211&gt; LENGTH: 382

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 137

```

Met Arg Pro Gly Thr Ala Leu Gln Ala Val Leu Leu Ala Val Leu Leu
 1           5           10           15
Val Gly Leu Arg Ala Ala Thr Gly Arg Leu Leu Ser Ala Ser Asp Leu
          20           25           30
Asp Leu Arg Gly Gly Gln Pro Val Cys Arg Gly Gly Thr Gln Arg Pro
          35           40           45
Cys Tyr Lys Val Ile Tyr Phe His Asp Thr Ser Arg Arg Leu Asn Phe
          50           55           60
Glu Glu Ala Lys Glu Ala Cys Arg Arg Asp Gly Gly Gln Leu Val Ser
          65           70           75           80
Ile Glu Ser Glu Asp Glu Gln Lys Leu Ile Glu Lys Phe Ile Glu Asn
          85           90           95
Leu Leu Pro Ser Asp Gly Asp Phe Trp Ile Gly Leu Arg Arg Arg Glu
          100          105          110
Glu Lys Gln Ser Asn Ser Thr Ala Cys Gln Asp Leu Tyr Ala Trp Thr
          115          120          125
Asp Gly Ser Ile Ser Gln Phe Arg Asn Trp Tyr Val Asp Glu Pro Ser
          130          135          140
Cys Gly Ser Glu Val Cys Val Val Met Tyr His Gln Pro Ser Ala Pro
          145          150          155          160
Ala Gly Ile Gly Gly Pro Tyr Met Phe Gln Trp Asn Asp Asp Arg Cys
          165          170          175
Asn Met Lys Asn Asn Phe Ile Cys Lys Tyr Ser Asp Glu Lys Pro Ala
          180          185          190

```



-continued

---

Val Pro Ser Arg Glu Ala Glu Gly Glu Glu Thr Glu Leu Thr Thr Pro  
 195 200 205

Val Leu Pro Glu Glu Thr Gln Glu Glu Asp Ala Lys Lys Thr Phe Lys  
 210 215 220

Glu Ser Arg Glu Ala Ala Leu Asn Leu Ala Tyr Ile Leu Ile Pro Ser  
 225 230 235 240

Ile Pro Leu Leu Leu Leu Val Val Thr Thr Val Val Cys Trp Val  
 245 250 255

Trp Ile Cys Arg Lys Arg Lys Arg Glu Gln Pro Asp Pro Ser Thr Lys  
 260 265 270

Lys Gln His Thr Ile Trp Pro Ser Pro His Gln Gly Asn Ser Pro Asp  
 275 280 285

Leu Glu Val Tyr Asn Val Ile Arg Lys Gln Ser Glu Ala Asp Leu Ala  
 290 295 300

Glu Thr Arg Pro Asp Leu Lys Asn Ile Ser Phe Arg Val Cys Ser Gly  
 305 310 315 320

Glu Ala Thr Pro Asp Asp Met Ser Cys Asp Tyr Asp Asn Met Ala Val  
 325 330 335

Asn Pro Ser Glu Ser Gly Phe Val Thr Leu Val Ser Val Glu Ser Gly  
 340 345 350

Phe Val Thr Asn Asp Ile Tyr Glu Phe Ser Pro Asp Gln Met Gly Arg  
 355 360 365

Ser Lys Glu Ser Gly Trp Val Glu Asn Glu Ile Tyr Gly Tyr  
 370 375 380

<210> SEQ ID NO 138  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 138  
 gttcattgaa aacctcttgc catctgatgg tgacttctgg attgggctca 50

<210> SEQ ID NO 139  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 139  
 aagccaaaga agcctgcagg aggg 24

<210> SEQ ID NO 140  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 140  
 cagtccaagc ataaaggctc tgcc 24

-continued

---

```

<210> SEQ ID NO 141
<211> LENGTH: 1514
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 141

ggggtctccc tcagggccgg gaggcacagc ggtccctgct tgctgaaggg ctggatgtac    60
gcatccgcag gttcccgccg acttgggggc gcccgctgag ccccgcgcc cgcagaagac    120
ttgtgtttgc tcctgcagc ctcaaccgg agggcagcga gggcctacca ccatgatcac    180
tggtgtgttc agcatgcgct tgtggacccc agtgggcgtc ctgacctcgc tggcgtactg    240
cctgcaccag cggcgggtgg ccctggccga gctgcaggag gccgatggcc agtgtccggt    300
cgaccgcagc ctgctgaagt tgaaaatggt gcaggctgtg ttctgacacg gggctcggag    360
tcctctcaag ccgctcccgc tggaggagca ggtagagtgg aacccccagc tattagaggt    420
cccccccaa actcagtttg attacacagt caccaatcta gctggtggtc cgaaaccata    480
ttctccttac gactctcaat accatgagac caccctgaag gggggcatgt ttgctgggca    540
gctgaccaag gtgggatcgc agcaaatggt tgccttggga gagagactga ggaagaacta    600
tgtggaagac attccctttc tttcaccaac cttcaacca caggaggtct ttattcgttc    660
cactaacatt tttcggaatc tggagtccac ccgttgtttg ctggctgggc ttttccagtg    720
tcagaaagaa ggacctcatc tcatccacac tgatgaagca gattcagaag tcttztatcc    780
caactaccaa agctgctgga gcctgaggca gagaaccaga ggcgggaggc agactgcctc    840
tttacagcca ggaatctcag aggatttgaa aaaggtgaag gacaggatgg gcattgacag    900
tagtgataaa gtggacttct tcatctcctt ggacaacgtg gctgccgagc aggcacacaa    960
cctcccaagc tgccccatgc tgaagagatt tgcacggatg atcgaacaga gagctgtgga   1020
cacatccttg tacatactgc ccaaggaaga cagggaaagt ctctcagatg cagtaggccc   1080
attcctccac atcctagaga gcaacctgct gaaagccatg gactctgcca ctgccccga   1140
caagatcaga aagctgtatc tctatcgccg tcatgatgtg accttcatac cgctcttaat   1200
gaccctgggg atttttgacc acaaatggcc accgtttctt gttgacctga ccatggaact   1260
ttaccagcac ctggaatcta aggagtgttt tgtgcagctc tattaccacg ggaaggagca   1320
ggtgccgaga ggttgccctg atgggctctg cccgctggac atgttcttga atgccatgtc   1380
agtttatacc ttaagcccg aaaaatacca tgcactctgc tctcaaacctc aggtgatgga   1440
agttggaaat gaagagtaac tgattataa aagcaggatg tgttgatgtt aaaataaagt   1500
gcctttatac aatg                                     1514

```

```

<210> SEQ ID NO 142
<211> LENGTH: 428
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 142

Met Ile Thr Gly Val Phe Ser Met Arg Leu Trp Thr Pro Val Gly Val
 1             5             10             15

Leu Thr Ser Leu Ala Tyr Cys Leu His Gln Arg Arg Val Ala Leu Ala
          20             25             30

Glu Leu Gln Glu Ala Asp Gly Gln Cys Pro Val Asp Arg Ser Leu Leu
 35             40             45

```

-continued

---

Lys Leu Lys Met Val Gln Val Val Phe Arg His Gly Ala Arg Ser Pro  
 50 55 60

Leu Lys Pro Leu Pro Leu Glu Glu Gln Val Glu Trp Asn Pro Gln Leu  
 65 70 75 80

Leu Glu Val Pro Pro Gln Thr Gln Phe Asp Tyr Thr Val Thr Asn Leu  
 85 90 95

Ala Gly Gly Pro Lys Pro Tyr Ser Pro Tyr Asp Ser Gln Tyr His Glu  
 100 105 110

Thr Thr Leu Lys Gly Gly Met Phe Ala Gly Gln Leu Thr Lys Val Gly  
 115 120 125

Met Gln Gln Met Phe Ala Leu Gly Glu Arg Leu Arg Lys Asn Tyr Val  
 130 135 140

Glu Asp Ile Pro Phe Leu Ser Pro Thr Phe Asn Pro Gln Glu Val Phe  
 145 150 155 160

Ile Arg Ser Thr Asn Ile Phe Arg Asn Leu Glu Ser Thr Arg Cys Leu  
 165 170 175

Leu Ala Gly Leu Phe Gln Cys Gln Lys Glu Gly Pro Ile Ile Ile His  
 180 185 190

Thr Asp Glu Ala Asp Ser Glu Val Leu Tyr Pro Asn Tyr Gln Ser Cys  
 195 200 205

Trp Ser Leu Arg Gln Arg Thr Arg Gly Arg Arg Gln Thr Ala Ser Leu  
 210 215 220

Gln Pro Gly Ile Ser Glu Asp Leu Lys Lys Val Lys Asp Arg Met Gly  
 225 230 235 240

Ile Asp Ser Ser Asp Lys Val Asp Phe Phe Ile Leu Leu Asp Asn Val  
 245 250 255

Ala Ala Glu Gln Ala His Asn Leu Pro Ser Cys Pro Met Leu Lys Arg  
 260 265 270

Phe Ala Arg Met Ile Glu Gln Arg Ala Val Asp Thr Ser Leu Tyr Ile  
 275 280 285

Leu Pro Lys Glu Asp Arg Glu Ser Leu Gln Met Ala Val Gly Pro Phe  
 290 295 300

Leu His Ile Leu Glu Ser Asn Leu Leu Lys Ala Met Asp Ser Ala Thr  
 305 310 315 320

Ala Pro Asp Lys Ile Arg Lys Leu Tyr Leu Tyr Ala Ala His Asp Val  
 325 330 335

Thr Phe Ile Pro Leu Leu Met Thr Leu Gly Ile Phe Asp His Lys Trp  
 340 345 350

Pro Pro Phe Ala Val Asp Leu Thr Met Glu Leu Tyr Gln His Leu Glu  
 355 360 365

Ser Lys Glu Trp Phe Val Gln Leu Tyr Tyr His Gly Lys Glu Gln Val  
 370 375 380

Pro Arg Gly Cys Pro Asp Gly Leu Cys Pro Leu Asp Met Phe Leu Asn  
 385 390 395 400

Ala Met Ser Val Tyr Thr Leu Ser Pro Glu Lys Tyr His Ala Leu Cys  
 405 410 415

Ser Gln Thr Gln Val Met Glu Val Gly Asn Glu Glu  
 420 425

<210> SEQ ID NO 143  
 <211> LENGTH: 24

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 143

ccaactacca aagctgctgg agcc 24

<210> SEQ ID NO 144
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 144

gcagctctat taccacggga agga 24

<210> SEQ ID NO 145
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 145

tccttcccg tggtaatagag ctgc 24

<210> SEQ ID NO 146
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 146

ggcagagaac cagaggccg aggagactgc ctctttacag ccagg 45

<210> SEQ ID NO 147
<211> LENGTH: 1686
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 147

ctcctcttaa catacttgca gctaaaacta aatattgctg cttggggacc tccttctagc 60
cttaaatattc agctcatcac cttcacctgc cttggtcacg gctctgctat tctccttgat 120
ccttgccatt tgcaccagac ctggattcct agcgtctcca tctggagtgc ggcctggggg 180
gggcctccac cgctgtgaag ggcgggtgga ggtggaacag aaaggccagt ggggcaccgt 240
gtgtgatgac ggctgggaca ttaaggacgt ggctgtgttg tgccgggagc tgggctgtgg 300
agctgccagc ggaaccacct gtggtatttt gtatgagcca ccagcagaaa aagagcaaaa 360
ggtcctcacc caatcagtc gttgcacagg aacagaagat acattggctc agtgtgagca 420
agaagaagtt tatgattgtt cacatgatga agatgctggg gcatcgtgtg agaaccaga 480
gagctctttc tccccagtcc cagaggggtg caggctggct gacggccttg ggcattgcaa 540
gggacgcgtg gaagtgaagc accagaacca gtggtatacc gtgtgccaga caggctggag 600

```

-continued

```

cctccgggcc gcaaaggtgg tgtgccggca gctgggatgt gggagggctg tactgactca 660
aaaacgctgc aacaagcatg cctatggccg aaaaccatc tggctgagcc agatgtcatg 720
ctcaggacga gaagcaaccc ttcaggattg cccttctggg ccttggggga agaacacctg 780
caaccatgat gaagacacgt gggtcgaatg tgaagatccc tttgacttga gactagtagg 840
aggagacaac ctctgctctg ggcgactgga ggtgctgcac aagggcgtat ggggctctgt 900
ctgtgatgac aactggggag aaaaggagga ccaggtggtg tgcaagcaac tgggctgtgg 960
gaagtccctc tctccctcct tcagagaccg gaaatgctat ggcctgggg tggcccgcat 1020
ctggctggat aatgttcctt gctcagggga ggagcagtcc ctggagcagt gccagcacag 1080
atthtggggg tttcacgact gcacccacca ggaagatgtg gctgtcatct gctcagtgtg 1140
ggtgggcatc atctaactctg ttgagtgcct gaatagaaga aaaacacaga agaagggagc 1200
atthactgtc tacatgactg catgggatga aactgatctt tcttctgcc ttgactggg 1260
acttatactt ggtgccctg attctcaggc cttcagagtt ggatcagaac ttacaacatc 1320
aggtctagtt ctcagggcat cagacatagt ttggaactac atcaccacct ttctatgtc 1380
tccacattgc acacagcaga ttcccagcct ccataattgt gtgtatcaac tacttaata 1440
cattctcaca cacacacaca cacacacaca cacacataca ccattgtcc 1500
tgthtctctg aagaactctg acaaataca gattttggtg ctgaaagaga ttctagagga 1560
acggaatttt aaggataaat tttctgaatt gggtatgggg tttctgaaat tggctctata 1620
atctaattag atataaaatt ctggtaactt tatttacaat aataaagata gcactatgtg 1680
ttcaaa 1686

```

&lt;210&gt; SEQ ID NO 148

&lt;211&gt; LENGTH: 347

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 148

```

Met Ala Leu Leu Phe Ser Leu Ile Leu Ala Ile Cys Thr Arg Pro Gly
 1             5             10             15
Phe Leu Ala Ser Pro Ser Gly Val Arg Leu Val Gly Gly Leu His Arg
             20             25             30
Cys Glu Gly Arg Val Glu Val Glu Gln Lys Gly Gln Trp Gly Thr Val
             35             40             45
Cys Asp Asp Gly Trp Asp Ile Lys Asp Val Ala Val Leu Cys Arg Glu
             50             55             60
Leu Gly Cys Gly Ala Ala Ser Gly Thr Pro Ser Gly Ile Leu Tyr Glu
             65             70             75             80
Pro Pro Ala Glu Lys Glu Gln Lys Val Leu Ile Gln Ser Val Ser Cys
             85             90             95
Thr Gly Thr Glu Asp Thr Leu Ala Gln Cys Glu Gln Glu Glu Val Tyr
             100            105            110
Asp Cys Ser His Asp Glu Asp Ala Gly Ala Ser Cys Glu Asn Pro Glu
             115            120            125
Ser Ser Phe Ser Pro Val Pro Glu Gly Val Arg Leu Ala Asp Gly Pro
             130            135            140
Gly His Cys Lys Gly Arg Val Glu Val Lys His Gln Asn Gln Trp Tyr
             145            150            155            160

```

-continued

---

Thr Val Cys Gln Thr Gly Trp Ser Leu Arg Ala Ala Lys Val Val Cys  
 165 170 175

Arg Gln Leu Gly Cys Gly Arg Ala Val Leu Thr Gln Lys Arg Cys Asn  
 180 185 190

Lys His Ala Tyr Gly Arg Lys Pro Ile Trp Leu Ser Gln Met Ser Cys  
 195 200 205

Ser Gly Arg Glu Ala Thr Leu Gln Asp Cys Pro Ser Gly Pro Trp Gly  
 210 215 220

Lys Asn Thr Cys Asn His Asp Glu Asp Thr Trp Val Glu Cys Glu Asp  
 225 230 235 240

Pro Phe Asp Leu Arg Leu Val Gly Gly Asp Asn Leu Cys Ser Gly Arg  
 245 250 255

Leu Glu Val Leu His Lys Gly Val Trp Gly Ser Val Cys Asp Asp Asn  
 260 265 270

Trp Gly Glu Lys Glu Asp Gln Val Val Cys Lys Gln Leu Gly Cys Gly  
 275 280 285

Lys Ser Leu Ser Pro Ser Phe Arg Asp Arg Lys Cys Tyr Gly Pro Gly  
 290 295 300

Val Gly Arg Ile Trp Leu Asp Asn Val Arg Cys Ser Gly Glu Glu Gln  
 305 310 315 320

Ser Leu Glu Gln Cys Gln His Arg Phe Trp Gly Phe His Asp Cys Thr  
 325 330 335

His Gln Glu Asp Val Ala Val Ile Cys Ser Val  
 340 345

<210> SEQ ID NO 149  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 149

ttcagctcat cacctcacc tgcc 24

<210> SEQ ID NO 150  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 150

ggctcataca aaataccact aggg 24

<210> SEQ ID NO 151  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 151

gggcctccac cgctgtgaag ggcggtgga ggtggaacag aaaggccagt 50

-continued

<210> SEQ ID NO 152  
 <211> LENGTH: 1427  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 152

```

actgcactcg gttctatcga ttgaattccc cggggatcct ctagagatcc ctcgacctcg      60
accacacgct ccgcggacgc gtgggcggac gcgtgggccc gctaccagga agagtctgcc      120
gaaggtgaag gccatggact tcatcacctc cacagccatc ctgccctgc tgttcggctg      180
cctgggcgct ttcggcctct tccggctgct gcagtgggtg cgcgggaagg cctacctgcg      240
gaatgctgtg gtggtgatca caggcggccac ctcagggctg ggcaaagaat gtgcaaagt      300
cttctatgct gcgggtgcta aactggtgct ctgtggccgg aatggtgggg ccctagaaga      360
gtcctacaga gaacttaccg cttctcatgc caccaagggt cagacacaca agccttactt      420
ggtgaccttc gacctcacag actctggggc catagttgca gcagcagctg agatcctgca      480
gtgctttggc tatgtcgaca tacttgtcaa caatgctggg atcagctacc gtggtaccat      540
catggacacc acagtggatg tggacaagag ggtcatggag acaaaactact ttggcccagt      600
tgctctaacg aaagcactcc tgcccctcat gatcaagagg aggcaaggcc acattgtcgc      660
catcagcagc atccagggca agatgagcat tccttttcga tcagcatatg cagcctccaa      720
gcacgcaacc caggccttct ttgactgtct gcgtgccgag atggaacagt atgaaattga      780
ggtgaccgtc atcagccccg gctacatcca caccaacctc tctgtaaatg ccatcaccgc      840
ggatggatct aggtatggag ttatggacac caccacagcc cagggccgaa gcctgtgga      900
ggtggcccag gatgttcttg ctgctgtggg gaagaagaag aaagatgtga tcctggctga      960
cttactgcct tccttggtg tttatctctg aactctggct cctgggctct tcttcagcct     1020
catggcctcc agggccagaa aagagcggaa atccaagaac tcctagtact ctgaccagcc     1080
agggccaggg cagagaagca gcaactcttag gcttgcttac tctacaaggg acagttgcat     1140
ttgttgagac tttaatggag atttgtctca caagtgggaa agactgaaga aacacatctc     1200
gtgcagatct gctggcagag gacaatcaaa aacgacaaca agcttcttcc cagggtgagg     1260
ggaaacactt aaggaataaa tatggagctg gggtttaaca ctaaaaacta gaaataaaca     1320
tctcaaacag taaaaaaaaa aaaaaagggc ggccgcgact ctagagtcga cctgcagaag     1380
cttgcccgcc atggccaac ttgtttattg cagcttataa tggttac      1427
    
```

<210> SEQ ID NO 153  
 <211> LENGTH: 310  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 153

```

Met Asp Phe Ile Thr Ser Thr Ala Ile Leu Pro Leu Leu Phe Gly Cys
 1           5           10           15
Leu Gly Val Phe Gly Leu Phe Arg Leu Leu Gln Trp Val Arg Gly Lys
 20           25           30
Ala Tyr Leu Arg Asn Ala Val Val Val Ile Thr Gly Ala Thr Ser Gly
 35           40           45
Leu Gly Lys Glu Cys Ala Lys Val Phe Tyr Ala Ala Gly Ala Lys Leu
 50           55           60
    
```

-continued

---

Val Leu Cys Gly Arg Asn Gly Gly Ala Leu Glu Glu Leu Ile Arg Glu  
 65 70 75 80

Leu Thr Ala Ser His Ala Thr Lys Val Gln Thr His Lys Pro Tyr Leu  
 85 90 95

Val Thr Phe Asp Leu Thr Asp Ser Gly Ala Ile Val Ala Ala Ala Ala  
 100 105 110

Glu Ile Leu Gln Cys Phe Gly Tyr Val Asp Ile Leu Val Asn Asn Ala  
 115 120 125

Gly Ile Ser Tyr Arg Gly Thr Ile Met Asp Thr Thr Val Asp Val Asp  
 130 135 140

Lys Arg Val Met Glu Thr Asn Tyr Phe Gly Pro Val Ala Leu Thr Lys  
 145 150 155 160

Ala Leu Leu Pro Ser Met Ile Lys Arg Arg Gln Gly His Ile Val Ala  
 165 170 175

Ile Ser Ser Ile Gln Gly Lys Met Ser Ile Pro Phe Arg Ser Ala Tyr  
 180 185 190

Ala Ala Ser Lys His Ala Thr Gln Ala Phe Phe Asp Cys Leu Arg Ala  
 195 200 205

Glu Met Glu Gln Tyr Glu Ile Glu Val Thr Val Ile Ser Pro Gly Tyr  
 210 215 220

Ile His Thr Asn Leu Ser Val Asn Ala Ile Thr Ala Asp Gly Ser Arg  
 225 230 235 240

Tyr Gly Val Met Asp Thr Thr Thr Ala Gln Gly Arg Ser Pro Val Glu  
 245 250 255

Val Ala Gln Asp Val Leu Ala Ala Val Gly Lys Lys Lys Lys Asp Val  
 260 265 270

Ile Leu Ala Asp Leu Leu Pro Ser Leu Ala Val Tyr Leu Arg Thr Leu  
 275 280 285

Ala Pro Gly Leu Phe Phe Ser Leu Met Ala Ser Arg Ala Arg Lys Glu  
 290 295 300

Arg Lys Ser Lys Asn Ser  
 305 310

<210> SEQ ID NO 154  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 154

ggtgctaaac tgggtgctctg tggc 24

<210> SEQ ID NO 155  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 155

cagggaaga tgagcattcc 20

<210> SEQ ID NO 156



-continued

---

```

<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 156

tcatactggt ccatctcggc acgc                                24

<210> SEQ ID NO 157
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 157

aatggtgggg ccctagaaga gctcatcaga gaactcaccg cttctcatgc    50

<210> SEQ ID NO 158
<211> LENGTH: 1771
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 158

cccacgcgtc cgctggtggt agatcgagca accctctaaa agcagtttag agtggtaaaa    60
aaaaaaaaaa acacacacaaa cgctcgcagc cacaaaaggg atgaaatttc ttctggacat    120
cctcctgctt ctcccgttac tgatcgtctg ctccctagag tccttcgtga agctttttat    180
tcctaaaggg agaaaatcag tcaccggcga aatcgtgctg attacaggag ctgggcatgg    240
aattgggaga ctgactgcct atgaatttgc taaacttaaa agcaagctgg ttctctggga    300
tataaataag catggactgg aggaaacagc tgccaaatgc aagggactgg gtgccaaggt    360
tcataccttt gtggtagact gcagcaaccg agaagatatt tacagctctg caaagaaggt    420
gaaggcagaa attggagatg ttagtatttt agtaaataat gctggtgtag totatacatc    480
agatttggtt gctacacaag atcctcagat tgaaaagact tttgaagtta atgtacttgc    540
acatttctg actacaaagg catttcttcc tgcaatgacg aagaataacc atggccatat    600
tgtcactgtg gcttcggcag ctggacatgt ctcggcccc ttcttactgg ctactgttc    660
aagcaagttt gctgctggtg gatttcataa aactttgaca gatgaactgg ctgccttaca    720
aataactgga gtcaaaacaa catgtctgtg tcctaatttc gtaaacactg gcttcatcaa    780
aaatccaagt acaagtttgg gaccactctt ggaacctgag gaagtggtaa acaggctgat    840
gcatgggatt ctgactgagc agaagatgat ttttattcca tcttctatag cttttttaac    900
aacattggaa aggatccttc ctgagcgttt cctggcagtt ttaaaacgaa aaatcagtgt    960
taagtttgat gcagttattg gatataaaat gaaagcgcaa taagcaccta gttttctgaa   1020
aactgattta ccaggtttag gttgatgtca totaatagtg ccagaatfff aatgtttgaa   1080
cttctgtttt ttctaattat cccatttctt tcaatatcat ttttgaggct ttggcagtct   1140
tcatttacta ccactgttcc tttagccaaa agctgattac atatgatata aacagagaaa   1200
taccttttaga ggtgacttta aggaaaatga agaaaaagaa ccaaaatgac tttattaaaa   1260
taatttccaa gattatttgt ggctcacctg aaggctttgc aaaatttga ccataaccgt   1320

```

-continued

---

```

ttatttaaca tatattttta tttttgattg cacttaaatt ttgtataatt tgtgtttctt 1380
tttctgttct acataaaatc agaaacttca agctctctaa ataaaatgaa ggactatatac 1440
tagtgggtatt tcacaatgaa tatcatgaac tctcaatggg taggtttcat cctaccatt 1500
gccactctgt ttctcgagag atacctcaca ttccaatgcc aaacatttct gcacagggaa 1560
gctagagggtg gatacacgtg ttgcaagtat aaaagcatca ctgggattta aggagaattg 1620
agagaatgta cccacaaatg gcagcaataa taaatggatc acacttaaaa aaaaaaaaaa 1680
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1740
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a          1771
    
```

```

<210> SEQ ID NO 159
<211> LENGTH: 300
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 159
    
```

```

Met Lys Phe Leu Leu Asp Ile Leu Leu Leu Leu Pro Leu Leu Ile Val
 1           5           10           15
Cys Ser Leu Glu Ser Phe Val Lys Leu Phe Ile Pro Lys Arg Arg Lys
          20           25           30
Ser Val Thr Gly Glu Ile Val Leu Ile Thr Gly Ala Gly His Gly Ile
          35           40           45
Gly Arg Leu Thr Ala Tyr Glu Phe Ala Lys Leu Lys Ser Lys Leu Val
          50           55           60
Leu Trp Asp Ile Asn Lys His Gly Leu Glu Glu Thr Ala Ala Lys Cys
          65           70           75           80
Lys Gly Leu Gly Ala Lys Val His Thr Phe Val Val Asp Cys Ser Asn
          85           90           95
Arg Glu Asp Ile Tyr Ser Ser Ala Lys Lys Val Lys Ala Glu Ile Gly
          100          105          110
Asp Val Ser Ile Leu Val Asn Asn Ala Gly Val Val Tyr Thr Ser Asp
          115          120          125
Leu Phe Ala Thr Gln Asp Pro Gln Ile Glu Lys Thr Phe Glu Val Asn
          130          135          140
Val Leu Ala His Phe Trp Thr Thr Lys Ala Phe Leu Pro Ala Met Thr
          145          150          155          160
Lys Asn Asn His Gly His Ile Val Thr Val Ala Ser Ala Ala Gly His
          165          170          175
Val Ser Val Pro Phe Leu Leu Ala Tyr Cys Ser Ser Lys Phe Ala Ala
          180          185          190
Val Gly Phe His Lys Thr Leu Thr Asp Glu Leu Ala Ala Leu Gln Ile
          195          200          205
Thr Gly Val Lys Thr Thr Cys Leu Cys Pro Asn Phe Val Asn Thr Gly
          210          215          220
Phe Ile Lys Asn Pro Ser Thr Ser Leu Gly Pro Thr Leu Glu Pro Glu
          225          230          235          240
Glu Val Val Asn Arg Leu Met His Gly Ile Leu Thr Glu Gln Lys Met
          245          250          255
Ile Phe Ile Pro Ser Ser Ile Ala Phe Leu Thr Thr Leu Glu Arg Ile
          260          265          270
Leu Pro Glu Arg Phe Leu Ala Val Leu Lys Arg Lys Ile Ser Val Lys
    
```

-continued

275	280	285
Phe Asp Ala Val Ile Gly Tyr Lys Met Lys Ala Gln		
290	295	300
<210> SEQ ID NO 160 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe  <400> SEQUENCE: 160  ggtgaaggca gaaattggag atg 23		
<210> SEQ ID NO 161 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe  <400> SEQUENCE: 161  atcccatgca tcagcctggt tacc 24		
<210> SEQ ID NO 162 <211> LENGTH: 48 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe  <400> SEQUENCE: 162  gctggtgtg tctatacatc agatttggtt gctacacaag atcctcag 48		
<210> SEQ ID NO 163 <211> LENGTH: 2076 <212> TYPE: DNA <213> ORGANISM: Homo sapiens  <400> SEQUENCE: 163  cccacgcgtc cgcggacgcg tgggtcgact agttctagat cgcgagcggc cgcccgcggc 60 tcaggaggga gcaccgactg cgccgcaccc tgagagatgg ttggtgccat gtggaagggtg 120 attgtttcgc tggctcctggt gatgcctggc ccctgtgatg ggctgtttcg ctccctatac 180 agaagtgttt ccatgccacc taaggagac tcaggacagc cattatttct cacccttac 240 attgaagctg ggaagatcca aaaaggaaga gaattgagtt tggctcggccc tttcccagga 300 ctgaacatga agagttagc cggcttcctc accgtgaata agacttacia cagcaacctc 360 ttcttctggt tcttcccagc tcagatacag ccagaagatg ccccagtagt tctctggcta 420 cagggtgggc cgggagggtc atccatggtt ggactccttg tgaacatgg gccttatggt 480 gtcacaagta acatgacctt gcgtgacaga gacttccctt ggaccacaac gctctocatg 540 ctttacattg acaatccagt gggcacagcg ttcagtttta ctgatgatac ccacggatat 600 gcagtcaatg aggacgatg agcacgggat ttatacagtg cactaattca gtttttccag 660 atatttctcg aatataaaaa taatgacttt tatgtcactg gggagtotta tgcagggaaa 720		

-continued

```

tatgtgccag ccattgcaca cctcatccat tccctcaacc ctgtgagaga ggtgaagatc 780
aacctgaacg gaattgctat tggagatgga tattctgatc ccgaatcaat tatagggggc 840
tatgcagaat tcctgtacca aattgcttgg ttggatgaga agcaaaaaaa gtacttccag 900
aagcagtgcc atgaatgcac agaacacatc aggaagcaga actggtttga ggcctttgaa 960
atactggata aactactaga tggcgactta acaagtgatc cttcttactt ccagaatggt 1020
acaggatgta gtaattacta taactttttg cgggtgcacgg aacctgagga tcagctttac 1080
tatgtgaaat ttttgtcact cccagaggtg agacaagcca tccacgtggg gaatcagact 1140
tttaatgatg gaactatagt tgaaaagtac ttgcgagaag atacagtaca gtcagttaag 1200
ccatggttaa ctgaaatcat gaataattat aaggttctga tctacaatgg ccaactggac 1260
atcatcgtgg cagctgcctc gacagagcgc tccttgatgg gcatggactg gaaaggatcc 1320
caggaataca agaaggcaga aaaaaaagt ttggaagtct ttaaacttga cagtgaagtg 1380
gctggttaca tccggcaacg gggtgacttc catcaggtaa ttattcgagg tggaggacat 1440
atthtaccct atgaccagcc tctgagagct tttgacatga ttaatcgatt catttatgga 1500
aaagatggg atccttatgt tgataaaact acctcccaa aagagaacat cagaggtttt 1560
cattgctgaa aagaaaatcg taaaaacaga aaatgtcata ggaataaaaa aattatcttt 1620
tcatatctgc aagatthttt tcatcaataa aaattatcct tgaacaagt gagctthttgt 1680
ttttgggggg agatgthttc tacaaaatta acatgagtac atgagtaaga attacattat 1740
ttaacttaaa ggatgaaagg tatggatgat gtgacactga gacaagatgt ataaatgaaa 1800
ttttagggtc ttgaatagga agttttaatt tottctaaga gtaagtgaaa agtgcagttg 1860
taacaacaaa agctgtaaca tctthttctg ccaataacag aagtttgcca tgccgtgaag 1920
gtgthttgaa atattattgg ataagaatag ctcaattatc ccaataaaat ggatgaagct 1980
ataatagttt tggggaaaaa attctcaaat gtataaagtc ttagaacaaa agaattcttt 2040
gaaataaaaa tattatatat aaaagtaaaa aaaaaa 2076

```

```

<210> SEQ ID NO 164
<211> LENGTH: 476
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 164

```

Met Val Gly Ala Met Trp Lys Val Ile Val Ser Leu Val Leu Leu Met
 1           5           10           15
Pro Gly Pro Cys Asp Gly Leu Phe Arg Ser Leu Tyr Arg Ser Val Ser
 20           25           30
Met Pro Pro Lys Gly Asp Ser Gly Gln Pro Leu Phe Leu Thr Pro Tyr
 35           40           45
Ile Glu Ala Gly Lys Ile Gln Lys Gly Arg Glu Leu Ser Leu Val Gly
 50           55           60
Pro Phe Pro Gly Leu Asn Met Lys Ser Tyr Ala Gly Phe Leu Thr Val
 65           70           75           80
Asn Lys Thr Tyr Asn Ser Asn Leu Phe Phe Trp Phe Phe Pro Ala Gln
 85           90           95
Ile Gln Pro Glu Asp Ala Pro Val Val Leu Trp Leu Gln Gly Gly Pro
 100          105          110
Gly Gly Ser Ser Met Phe Gly Leu Phe Val Glu His Gly Pro Tyr Val

```

-continued

115				120				125					
Val Thr Ser Asn Met Thr Leu Arg Asp Arg Asp Phe Pro Trp Thr Thr	130			135				140					
Thr Leu Ser Met Leu Tyr Ile Asp Asn Pro Val Gly Thr Gly Phe Ser	145			150				155					160
Phe Thr Asp Asp Thr His Gly Tyr Ala Val Asn Glu Asp Asp Val Ala			165					170					175
Arg Asp Leu Tyr Ser Ala Leu Ile Gln Phe Phe Gln Ile Phe Pro Glu			180				185					190	
Tyr Lys Asn Asn Asp Phe Tyr Val Thr Gly Glu Ser Tyr Ala Gly Lys			195			200				205			
Tyr Val Pro Ala Ile Ala His Leu Ile His Ser Leu Asn Pro Val Arg			210			215				220			
Glu Val Lys Ile Asn Leu Asn Gly Ile Ala Ile Gly Asp Gly Tyr Ser	225			230				235					240
Asp Pro Glu Ser Ile Ile Gly Gly Tyr Ala Glu Phe Leu Tyr Gln Ile			245					250					255
Gly Leu Leu Asp Glu Lys Gln Lys Lys Tyr Phe Gln Lys Gln Cys His			260					265					270
Glu Cys Ile Glu His Ile Arg Lys Gln Asn Trp Phe Glu Ala Phe Glu			275			280						285	
Ile Leu Asp Lys Leu Leu Asp Gly Asp Leu Thr Ser Asp Pro Ser Tyr			290			295				300			
Phe Gln Asn Val Thr Gly Cys Ser Asn Tyr Tyr Asn Phe Leu Arg Cys	305			310						315			320
Thr Glu Pro Glu Asp Gln Leu Tyr Tyr Val Lys Phe Leu Ser Leu Pro			325							330			335
Glu Val Arg Gln Ala Ile His Val Gly Asn Gln Thr Phe Asn Asp Gly			340					345					350
Thr Ile Val Glu Lys Tyr Leu Arg Glu Asp Thr Val Gln Ser Val Lys			355			360						365	
Pro Trp Leu Thr Glu Ile Met Asn Asn Tyr Lys Val Leu Ile Tyr Asn			370			375						380	
Gly Gln Leu Asp Ile Ile Val Ala Ala Ala Leu Thr Glu Arg Ser Leu	385			390						395			400
Met Gly Met Asp Trp Lys Gly Ser Gln Glu Tyr Lys Lys Ala Glu Lys			405							410			415
Lys Val Trp Lys Ile Phe Lys Ser Asp Ser Glu Val Ala Gly Tyr Ile			420							425			430
Arg Gln Ala Gly Asp Phe His Gln Val Ile Ile Arg Gly Gly Gly His			435			440							445
Ile Leu Pro Tyr Asp Gln Pro Leu Arg Ala Phe Asp Met Ile Asn Arg			450			455						460	
Phe Ile Tyr Gly Lys Gly Trp Asp Pro Tyr Val Gly	465			470								475	

&lt;210&gt; SEQ ID NO 165

&lt;211&gt; LENGTH: 24

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe

-continued

---

<400> SEQUENCE: 165  
 ttccatgccca cctaagggag actc 24

<210> SEQ ID NO 166  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 166  
 tggatgaggt gtgcaatggc tggc 24

<210> SEQ ID NO 167  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 167  
 agctctcaga ggctgtcat aggg 24

<210> SEQ ID NO 168  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 168  
 gtcggccctt tcccaggact gaacatgaag agttatgccg gcttcctcac 50

<210> SEQ ID NO 169  
 <211> LENGTH: 2477  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 169  
 cgagggcttt tccggctccg gaatggcaca tgtgggaatc ccagtcttgt tggctacaac 60  
 atttttccct ttcctaaca gttctaacag ctgttctaac agctagtgat caggggttct 120  
 tcttgctgga gaagaaagg ctgagggcag agcagggcac tctcaactcag ggtgaccagc 180  
 tccttgccctc tctgtggata acagagcatg agaaagtga gagatgcagc ggagtgaggt 240  
 gatggaagtc taaaatagga aggaattttg tgtgcaatat cagactctgg gacagttga 300  
 cctggagagc ctgggggagg gcctgcctaa caagcttca aaaaacagga gcgacttcca 360  
 ctgggctggg ataagacgtg ccggtaggat aggaagact gggtttagtc ctaatatcaa 420  
 attgactggc tgggtgaact tcaacagcct tttaacctct ctgggagatg aaaacgatgg 480  
 cttaagggcg cagaaataga gatgctttgt aaaataaaat tttaaaaaaa gcaagtattt 540  
 tatagcataa aggctagaga caaaataga taacaggatt ccctgaacat tcctaagagg 600  
 gagaagtat gttaaaaata gaaaaaccaa aatgcagaag gaggagactc acagagctaa 660  
 accaggatgg ggaccctggg tcaggccagc ctctttgctc ctcccggaaa ttatttttgg 720

-continued

---

```

tctgacct ctgccttggt ttttgacaaa tcatgtgagg gccaacggg gaaggtggag 780
cagatgagca cacacaggag cgtctcctc accgccgccc ctctcagcat ggaacagagg 840
cagccctggc cccgggcccgt ggaggtggac agccgctctg tggctcctgt ctcaagtggc 900
tgggtgctgc tggcccccc agcagccggc atgcctcagt tcagcacctt ccaactctgag 960
aatcgtgact ggaccttcaa ccaactgacc gtccaccaag ggacgggggc cgtctatgtg 1020
ggggccatca accgggtcta taagctgaca ggcaacctga ccatccagggt ggctcataag 1080
acagggccag aagaggacaa caagtctcgt taccgccccc tcatcgtgca gccctgcagc 1140
gaagtgtctc ccctcaccaa caatgtcaac aagctgtctc tcattgacta ctctgagaac 1200
cgctgtctgg cctgtgggag cctctaccag ggggtctgca agctgtctgc gctggatgac 1260
ctcttcatcc tgggtgagcc atcccacaag aaggagcact acctgtccag tgtcaacaag 1320
acgggcacca tgtacggggt gattgtgccc tctgaggggt aggatggcaa gctcttcatc 1380
ggcacggctg tggatgggaa gcaggattac ttcccagacc tgtccagccg gaagctgccc 1440
cgagaccctg agtccctcagc catgctgcac tatgagctac acagcgattt tgtctcctct 1500
ctcatcaaga tcccttcaga caccctggcc ctggtctccc actttgacat cttctacatc 1560
tacggctttg ctagtggggg ctttctctac tttctcactg tccagcccga gaccctgag 1620
gggtgtggcca tcaactccgc tggagacctc ttctacacct cacgcatcgt gcggctctgc 1680
aagatgacc ccaagttcca ctcatcctg tccctgcctc tcggctgcac ccgggcccgg 1740
gtggaatacc gcctcctgca ggctgcttac ctggccaagc ctggggactc actggcccag 1800
gccttcaata tcaccagcca ggacgatgta ctctttgcca tcttctccaa agggcagaag 1860
cagtatcacc acccccccga tgactctgcc ctgtgtgctc tccctatccg gccatcaac 1920
ttgcagatca aggagcgcct gcagtcctgc taccagggcg agggcaacct ggagctcaac 1980
tggctgtctg ggaaggacct ccagtgcacg aaggcgcctg tccccatcga tgataactc 2040
tgtggactgg acatcaacca gcccctggga ggctcaactc cagtggaggg cctgaccctg 2100
tacaccacca gcagggaccg catgacctct gtggcctcct acgtttacaa cggctacagc 2160
gtggtttttg tggggactaa gagtggcaag ctgaaaaagg taagagtcta tgagttcaga 2220
tgctccaatg ccattcacct cctcagcaaa gagtccctct tggaaggtag ctattggtgg 2280
agatttaact ataggcaact ttattttctt ggggaacaaa ggtgaaatgg ggaggttaaga 2340
aggggttaat tttgtgactt agcttctagc tacttctctc agccatcagt cattgggtat 2400
gtaaggaatg caagcgtatt tcaatatttc ccaaacttta agaaaaaact ttaagaaggt 2460
acatctgcaa aagcaaa 2477

```

```

<210> SEQ ID NO 170
<211> LENGTH: 552
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 170

```

```

Met Gly Thr Leu Gly Gln Ala Ser Leu Phe Ala Pro Pro Gly Asn Tyr
 1             5             10             15
Phe Trp Ser Asp His Ser Ala Leu Cys Phe Ala Glu Ser Cys Glu Gly
          20             25             30
Gln Pro Gly Lys Val Glu Gln Met Ser Thr His Arg Ser Arg Leu Leu
          35             40             45

```

-continued

---

Thr Ala Ala Pro Leu Ser Met Glu Gln Arg Gln Pro Trp Pro Arg Ala  
 50 55 60  
 Leu Glu Val Asp Ser Arg Ser Val Val Leu Leu Ser Val Val Trp Val  
 65 70 75 80  
 Leu Leu Ala Pro Pro Ala Ala Gly Met Pro Gln Phe Ser Thr Phe His  
 85 90 95  
 Ser Glu Asn Arg Asp Trp Thr Phe Asn His Leu Thr Val His Gln Gly  
 100 105 110  
 Thr Gly Ala Val Tyr Val Gly Ala Ile Asn Arg Val Tyr Lys Leu Thr  
 115 120 125  
 Gly Asn Leu Thr Ile Gln Val Ala His Lys Thr Gly Pro Glu Glu Asp  
 130 135 140  
 Asn Lys Ser Arg Tyr Pro Pro Leu Ile Val Gln Pro Cys Ser Glu Val  
 145 150 155 160  
 Leu Thr Leu Thr Asn Val Asn Lys Leu Leu Ile Ile Asp Tyr Ser  
 165 170 175  
 Glu Asn Arg Leu Leu Ala Cys Gly Ser Leu Tyr Gln Gly Val Cys Lys  
 180 185 190  
 Leu Leu Arg Leu Asp Asp Leu Phe Ile Leu Val Glu Pro Ser His Lys  
 195 200 205  
 Lys Glu His Tyr Leu Ser Ser Val Asn Lys Thr Gly Thr Met Tyr Gly  
 210 215 220  
 Val Ile Val Arg Ser Glu Gly Glu Asp Gly Lys Leu Phe Ile Gly Thr  
 225 230 235 240  
 Ala Val Asp Gly Lys Gln Asp Tyr Phe Pro Thr Leu Ser Ser Arg Lys  
 245 250 255  
 Leu Pro Arg Asp Pro Glu Ser Ser Ala Met Leu Asp Tyr Glu Leu His  
 260 265 270  
 Ser Asp Phe Val Ser Ser Leu Ile Lys Ile Pro Ser Asp Thr Leu Ala  
 275 280 285  
 Leu Val Ser His Phe Asp Ile Phe Tyr Ile Tyr Gly Phe Ala Ser Gly  
 290 295 300  
 Gly Phe Val Tyr Phe Leu Thr Val Gln Pro Glu Thr Pro Glu Gly Val  
 305 310 315 320  
 Ala Ile Asn Ser Ala Gly Asp Leu Phe Tyr Thr Ser Arg Ile Val Arg  
 325 330 335  
 Leu Cys Lys Asp Asp Pro Lys Phe His Ser Tyr Val Ser Leu Pro Phe  
 340 345 350  
 Gly Cys Thr Arg Ala Gly Val Glu Tyr Arg Leu Leu Gln Ala Ala Tyr  
 355 360 365  
 Leu Ala Lys Pro Gly Asp Ser Leu Ala Gln Ala Phe Asn Ile Thr Ser  
 370 375 380  
 Gln Asp Asp Val Leu Phe Ala Ile Phe Ser Lys Gly Gln Lys Gln Tyr  
 385 390 395 400  
 His His Pro Pro Asp Asp Ser Ala Leu Cys Ala Phe Pro Ile Arg Ala  
 405 410 415  
 Ile Asn Leu Gln Ile Lys Glu Arg Leu Gln Ser Cys Tyr Gln Gly Glu  
 420 425 430  
 Gly Asn Leu Glu Leu Asn Trp Leu Leu Gly Lys Asp Val Gln Cys Thr  
 435 440 445



-continued

---

Lys Ala Pro Val Pro Ile Asp Asp Asn Phe Cys Gly Leu Asp Ile Asn  
 450 455 460

Gln Pro Leu Gly Gly Ser Thr Pro Val Glu Gly Leu Thr Leu Tyr Thr  
 465 470 475 480

Thr Ser Arg Asp Arg Met Thr Ser Val Ala Ser Tyr Val Tyr Asn Gly  
 485 490 495

Tyr Ser Val Val Phe Val Gly Thr Lys Ser Gly Lys Leu Lys Lys Val  
 500 505 510

Arg Val Tyr Glu Phe Arg Cys Ser Asn Ala Ile His Leu Leu Ser Lys  
 515 520 525

Glu Ser Leu Leu Glu Gly Ser Tyr Trp Trp Arg Phe Asn Tyr Arg Gln  
 530 535 540

Leu Tyr Phe Leu Gly Glu Gln Arg  
 545 550

<210> SEQ ID NO 171  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 171

tggaataaccg cctcctgcag 20

<210> SEQ ID NO 172  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 172

cttctgccct ttggagaaga tggc 24

<210> SEQ ID NO 173  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 173

ggactcactg gcccaggcct tcaatatcac cagccaggac gat 42

<210> SEQ ID NO 174  
 <211> LENGTH: 3106  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (1683)  
 <223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 174

aggctcccgc gcgcggctga gtgcggactg gagtgggaac cgggtcccc gcgcttagag 60

aacacgcgat gaccacgtgg agcctccggc ggaggccggc ccgcacgctg ggactcctgc 120

-continued

tgctggctcgt cttgggcttc ctggtgctcc gcaggctgga ctggagcacc ctggtccctc	180
tgcggtcccg ccatcgacag ctggggctgc aggccaaagg ctggaacttc atgctggagg	240
attccacctt ctg gatcttc gggggctcca tccactatth cctgtgccc agggagtact	300
ggagggaccg cctgctgaag atgaaggcct gtggcttga caccctcacc acctatgttc	360
cgtggaacct gcatgagcca gaaagaggca aatttgactt ctctgggaac ctggacctgg	420
aggccttcgt cctgatggcc gcagagatcg ggctgtgggt gattctgcgt ccaggccctt	480
acatctgacg tgagatggac ctcgggggct tgcccagctg gctactccaa gaccctggca	540
tgaggctgag gacaacttac aagggttca ccgaagcagt ggacctttat tttgaccacc	600
tgatgtccag ggtgggtcca ctccagtaca agcgtggggg acctatcatt gccgtgcagg	660
tgagagaatga atatggttcc tataataaag accccgcata catgccctac gtcaagaagg	720
cactggagga ccgtggcatt gtggaactgc tcctgacttc agacaacaag gatgggctga	780
gcaaggggat tgtccagga gtottggcca ccatcaactt gcagtcaaca cacgagctgc	840
agctactgac cacctttctc tccaactgcc aggggactca gcccaagatg gtgatggagt	900
actggacggg gtggtttgac tcgtggggag gccctcaca tatcttggat tcttctgagg	960
ttttgaaaa cgtgtctgcc attgtggag ccggctcctc catcaacctc tacatgttcc	1020
acggaggcac caactttggc ttcattgaatg gagccatgca ctccatgac tacaagtcag	1080
atgtcaccag ctatgactat gatgtgtgac tgacagaagc cggcgattac acggccaagt	1140
acatgaagct tcgagacttc ttcggttcca tctcaggcat ccctctccct cccccacctg	1200
accttcttcc caagatgcc tatgagccct taacgccagt cttgtaactg tctctgtggg	1260
acgcccctca gtacctggg gagccaatca agtctgaaaa gcccatcaac atggagaacc	1320
tgccagtcaa tgggggaaat ggacagtctt tcgggtacat tctctatgag accgacatca	1380
cctcgtctgg catcctcagt ggccactgac atgatcgggg gcaggtggtt gtgaacacag	1440
tatccatag attcttggac tacaagacaa cgaagattgc tgtcccctg atccagggtt	1500
acaccgtgct gaggatcttg gtggagaatc gtggggcagc caactatggg gagaatattg	1560
atgaccagcg caaaggctta attggaatc tctatctgaa tgattcacc ctgaaaaact	1620
tcagaatcta tagcctggat atgaagaaga gcttcttca gaggttcggc ctggacaaat	1680
gngttccct cccagaaaca cccacattac ctgctttctt cttgggtagc ttgtccatca	1740
gctccacgac ttgtgacacc tttctgaagc tggagggctg ggagaagggg gttgtattca	1800
tcaatggcca gaaccttga cgttactgga acattggacc ccagaagacg ctttacctcc	1860
caggctcctg gttgagcagc ggaatcaacc aggtcatcgt ttttgaggag acgatggcgg	1920
gccctgcatt acagttcacg gaaaccccc acctgggacg gaaccagtac attaagtgag	1980
cgggtggcacc ccctcctgct ggtgccagt ggagactgcc gcctcctctt gacctgaagc	2040
ctggtggctg ctgccccacc cctcactgca aaagcatctc cttaagttagc aacctcaggg	2100
actggggctg acagtctgcc cctgtctcag ctcaaaacc taagcctgca gggaaagggtg	2160
ggatggctct gggcctggct ttgttgatga tggctttctt acagccctgc tcttgtgccc	2220
aggctgtcgg gctgtctcta ggggtggagc agctaactcag atcggccagc ctttggccct	2280
cagaaaaagt gctgaaactg gcccttgac cggacgtcac agccctgca gcatctgctg	2340
gactcaggcg tgctctttgc tggttctcgg gaggcttggc cacatccctc atggcccat	2400

-continued

```

tttatccccg aaatcctggg tgtgtcacca gtgtagaggg tggggaaggg gtgtctcacc 2460
tgagctgact ttgttcttcc ttcacaacct tctgagcctt ctttgggatt ctggaaggaa 2520
ctcggcgtga gaaacatgtg acttcccctt tcccttccca ctgctgctt cccacaggtt 2580
gacaggctgg gctggagaaa cagaaatcct caccctgcgt cttcccaagt tagcaggtgt 2640
ctctggtggt cagtgaggag gacatgtgag tcctggcaga agccatggcc catgtctgca 2700
catccaggga ggaggacaga aggcccagct cacatgtgag tcctggcaga agccatggcc 2760
catgtctgca catccaggga ggaggacaga aggcccagct cacatgtgag tcctggcaga 2820
agccatggcc catgtctgca catccaggga ggaggacaga aggcccagct cacatgtgag 2880
tcctggcaga agccatggcc catgtctgca catccaggga ggaggacaga aggcccagct 2940
cagtggcccc cgctccccac cccccagcc cgaacagcag gggcagagca gcctccttc 3000
gaagtgtgtc caagtccgca tttagcctt gttctggggc ccagcccaac acctggcttg 3060
ggctcaactgt cctgagttgc agtaaagcta taacottgaa tcacaa 3106

```

&lt;210&gt; SEQ ID NO 175

&lt;211&gt; LENGTH: 636

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;220&gt; FEATURE:

&lt;221&gt; NAME/KEY: MOD\_RES

&lt;222&gt; LOCATION: (539)

&lt;223&gt; OTHER INFORMATION: Any amino acid

&lt;400&gt; SEQUENCE: 175

```

Met Thr Thr Trp Ser Leu Arg Arg Arg Pro Ala Arg Thr Leu Gly Leu
 1          5          10          15

Leu Leu Leu Val Val Leu Gly Phe Leu Val Leu Arg Arg Leu Asp Trp
 20          25          30

Ser Thr Leu Val Pro Leu Arg Leu Arg His Arg Gln Leu Gly Leu Gln
 35          40          45

Ala Lys Gly Trp Asn Phe Met Leu Glu Asp Ser Thr Phe Trp Ile Phe
 50          55          60

Gly Gly Ser Ile His Tyr Phe Arg Val Pro Arg Glu Tyr Trp Arg Asp
 65          70          75          80

Arg Leu Leu Lys Met Lys Ala Cys Gly Leu Asn Thr Leu Thr Thr Tyr
 85          90          95

Val Pro Trp Asn Leu His Glu Pro Glu Arg Gly Lys Phe Asp Phe Ser
100          105          110

Gly Asn Leu Asp Leu Glu Ala Phe Val Leu Met Ala Ala Glu Ile Gly
115          120          125

Leu Trp Val Ile Leu Arg Pro Gly Pro Tyr Ile Cys Ser Glu Met Asp
130          135          140

Leu Gly Gly Leu Pro Ser Trp Leu Leu Gln Asp Pro Gly Met Arg Leu
145          150          155          160

Arg Thr Thr Tyr Lys Gly Phe Thr Glu Ala Val Asp Leu Tyr Phe Asp
165          170          175

His Leu Met Ser Arg Val Val Pro Leu Gln Tyr Lys Arg Gly Gly Pro
180          185          190

Ile Ile Ala Val Gln Val Glu Asn Glu Tyr Gly Ser Tyr Asn Lys Asp
195          200          205

Pro Ala Tyr Met Pro Tyr Val Lys Lys Ala Leu Glu Asp Arg Gly Ile

```

-continued

210		215		220											
Val	Glu	Leu	Leu	Leu	Thr	Ser	Asp	Asn	Lys	Asp	Gly	Leu	Ser	Lys	Gly
225					230					235					240
Ile	Val	Gln	Gly	Val	Leu	Ala	Thr	Ile	Asn	Leu	Gln	Ser	Thr	His	Glu
			245						250					255	
Leu	Gln	Leu	Leu	Thr	Thr	Phe	Leu	Phe	Asn	Val	Gln	Gly	Thr	Gln	Pro
			260						265				270		
Lys	Met	Val	Met	Glu	Tyr	Trp	Thr	Gly	Trp	Phe	Asp	Ser	Trp	Gly	Gly
		275					280						285		
Pro	His	Asn	Ile	Leu	Asp	Ser	Ser	Glu	Val	Leu	Lys	Thr	Val	Ser	Ala
		290					295					300			
Ile	Val	Asp	Ala	Gly	Ser	Ser	Ile	Asn	Leu	Tyr	Met	Phe	His	Gly	Gly
305					310					315					320
Thr	Asn	Phe	Gly	Phe	Met	Asn	Gly	Ala	Met	His	Phe	His	Asp	Tyr	Lys
				325					330					335	
Ser	Asp	Val	Thr	Ser	Tyr	Asp	Tyr	Asp	Ala	Val	Leu	Thr	Glu	Ala	Gly
			340						345				350		
Asp	Tyr	Thr	Ala	Lys	Tyr	Met	Lys	Leu	Arg	Asp	Phe	Phe	Gly	Ser	Ile
		355					360						365		
Ser	Gly	Ile	Pro	Leu	Pro	Pro	Pro	Pro	Asp	Leu	Leu	Pro	Lys	Met	Pro
		370					375					380			
Tyr	Glu	Pro	Leu	Thr	Pro	Val	Leu	Tyr	Leu	Ser	Leu	Trp	Asp	Ala	Leu
385					390					395					400
Lys	Tyr	Leu	Gly	Glu	Pro	Ile	Lys	Ser	Glu	Lys	Pro	Ile	Asn	Met	Glu
				405					410					415	
Asn	Leu	Pro	Val	Asn	Gly	Gly	Asn	Gly	Gln	Ser	Phe	Gly	Tyr	Ile	Leu
			420					425					430		
Tyr	Glu	Thr	Ser	Ile	Thr	Ser	Ser	Gly	Ile	Leu	Ser	Gly	His	Val	His
		435					440					445			
Asp	Arg	Gly	Gln	Val	Phe	Val	Asn	Thr	Val	Ser	Ile	Gly	Phe	Leu	Asp
		450					455					460			
Tyr	Lys	Thr	Thr	Lys	Ile	Ala	Val	Pro	Leu	Ile	Gln	Gly	Tyr	Thr	Val
465					470					475					480
Leu	Arg	Ile	Leu	Val	Glu	Asn	Arg	Gly	Arg	Val	Asn	Tyr	Gly	Glu	Asn
				485					490					495	
Ile	Asp	Asp	Gln	Arg	Lys	Gly	Leu	Ile	Gly	Asn	Leu	Tyr	Leu	Asn	Asp
			500						505				510		
Ser	Pro	Leu	Lys	Asn	Phe	Arg	Ile	Tyr	Ser	Leu	Asp	Met	Lys	Lys	Ser
		515					520					525			
Phe	Phe	Gln	Arg	Phe	Gly	Leu	Asp	Lys	Trp	Xaa	Ser	Leu	Pro	Glu	Thr
		530					535					540			
Pro	Thr	Leu	Pro	Ala	Phe	Phe	Leu	Gly	Ser	Leu	Ser	Ile	Ser	Ser	Thr
545					550					555					560
Pro	Cys	Asp	Thr	Phe	Leu	Lys	Leu	Glu	Gly	Trp	Glu	Lys	Gly	Val	Val
				565					570					575	
Phe	Ile	Asn	Gly	Gln	Asn	Leu	Gly	Arg	Tyr	Trp	Asn	Ile	Gly	Pro	Gln
				580					585				590		
Lys	Thr	Leu	Tyr	Leu	Pro	Gly	Pro	Trp	Leu	Ser	Ser	Gly	Ile	Asn	Gln
		595					600						605		
Val	Ile	Val	Phe	Glu	Glu	Thr	Met	Ala	Gly	Pro	Ala	Leu	Gln	Phe	Thr
610						615						620			



-continued

---

```

ggaccaaggg ccaagtctgg atcaatgggt ttaacttggg ccggtactgg acaaagcagg 1980
ggccacaaca gaccctctac gtgccaagat tcctgctggt tcctagggga gccctcaaca 2040
aaattacatt gctggaacta gaagatgtac ctctccagcc ccaagtccaa tttttggata 2100
agcctatcct caatagcact agtactttgc acaggacaca tatcaattcc ctttcagctg 2160
atacactgag tgccctctgaa ccaatggagt taagtgggca ctgaaaggta ggccgggcat 2220
ggtggctcat gcctgtaatc ccagcacttt gggaggctga gacgggtgga ttacctgagg 2280
tcaggacttc aagaccagcc tggccaacat ggtgaaaccc cgtctccact aaaatacaaa 2340
aaattagccg ggcgtgatgg tgggcacctc taatcccagc tacttgggag gctgagggca 2400
ggagaattcg ttgaatccag gaggcagagg ttgcagtgag tggaggttgt accactgcac 2460
tccagcctgg ctgacagtga gacactccat ctcaaaaaaa aaaaa 2505
    
```

```

<210> SEQ ID NO 177
<211> LENGTH: 654
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 177

```

Met Ala Pro Lys Lys Leu Ser Cys Leu Arg Ser Leu Leu Leu Pro Leu
 1          5          10          15
Ser Leu Thr Leu Leu Pro Gln Ala Asp Thr Arg Ser Phe Val Val
 20          25          30
Asp Arg Gly His Asp Arg Phe Leu Leu Asp Gly Ala Pro Phe Arg Tyr
 35          40          45
Val Ser Gly Ser Leu His Tyr Phe Arg Val Pro Arg Val Leu Trp Ala
 50          55          60
Asp Arg Leu Leu Lys Met Arg Trp Ser Gly Leu Asn Ala Ile Gln Phe
 65          70          75          80
Tyr Val Pro Trp Asn Tyr His Glu Pro Gln Pro Gly Val Tyr Asn Phe
 85          90          95
Asn Gly Ser Arg Asp Leu Ile Ala Phe Leu Asn Glu Ala Ala Leu Ala
100          105          110
Asn Leu Leu Val Ile Leu Arg Pro Gly Pro Tyr Ile Cys Ala Glu Trp
115          120          125
Glu Met Gly Gly Leu Pro Ser Trp Leu Leu Arg Lys Pro Glu Ile His
130          135          140
Leu Arg Thr Ser Asp Pro Asp Phe Leu Ala Ala Val Asp Ser Trp Phe
145          150          155          160
Lys Val Leu Leu Pro Lys Ile Tyr Pro Trp Leu Tyr His Asn Gly Gly
165          170          175
Asn Ile Ile Ser Ile Gln Val Glu Asn Glu Tyr Gly Ser Tyr Arg Ala
180          185          190
Cys Asp Phe Ser Tyr Met Arg His Leu Ala Gly Leu Phe Arg Ala Leu
195          200          205
Leu Gly Glu Lys Ile Leu Leu Phe Thr Thr Asp Gly Pro Glu Gly Leu
210          215          220
Lys Cys Gly Ser Leu Arg Gly Leu Tyr Thr Thr Val Asp Phe Gly Pro
225          230          235          240
Ala Asp Asn Met Thr Lys Ile Phe Thr Leu Leu Arg Lys Tyr Glu Pro
245          250          255
    
```

-continued

---

His Gly Pro Leu Val Asn Ser Glu Tyr Tyr Thr Gly Trp Leu Asp Tyr  
                   260                                  265                                  270

Trp Gly Gln Asn His Ser Thr Arg Ser Val Ser Ala Val Thr Lys Gly  
                   275                                  280                                  285

Leu Glu Asn Met Leu Lys Leu Gly Ala Ser Val Asn Met Tyr Met Phe  
                   290                                  295                                  300

His Gly Gly Thr Asn Phe Gly Tyr Trp Asn Gly Ala Asp Lys Lys Gly  
 305                                  310                                  315                                  320

Arg Phe Leu Pro Ile Thr Thr Ser Tyr Asp Tyr Asp Ala Pro Ile Ser  
                                   325                                  330                                  335

Glu Ala Gly Asp Pro Thr Pro Lys Leu Phe Ala Leu Arg Asp Val Ile  
                                   340                                  345                                  350

Ser Lys Phe Gln Glu Val Pro Leu Gly Pro Leu Pro Pro Pro Ser Pro  
                   355                                  360                                  365

Lys Met Met Leu Gly Pro Val Thr Leu His Leu Val Gly His Leu Leu  
                   370                                  375                                  380

Ala Phe Leu Asp Leu Leu Cys Pro Arg Gly Pro Ile His Ser Ile Leu  
 385                                  390                                  395                                  400

Pro Met Thr Phe Glu Ala Val Lys Gln Asp His Gly Phe Met Leu Tyr  
                                   405                                  410                                  415

Arg Thr Tyr Met Thr His Thr Ile Phe Glu Pro Thr Pro Phe Trp Val  
                                   420                                  425                                  430

Pro Asn Asn Gly Val His Asp Arg Ala Tyr Val Met Val Asp Gly Val  
                   435                                  440                                  445

Phe Gln Gly Val Val Glu Arg Asn Met Arg Asp Lys Leu Phe Leu Thr  
                   450                                  455                                  460

Gly Lys Leu Gly Ser Lys Leu Asp Ile Leu Val Glu Asn Met Gly Arg  
 465                                  470                                  475                                  480

Leu Ser Phe Gly Ser Asn Ser Ser Asp Phe Lys Gly Leu Leu Lys Pro  
                                   485                                  490                                  495

Pro Ile Leu Gly Gln Thr Ile Leu Thr Gln Trp Met Met Phe Pro Leu  
                                   500                                  505                                  510

Lys Ile Asp Asn Leu Val Lys Trp Trp Phe Pro Leu Gln Leu Pro Lys  
                   515                                  520                                  525

Trp Pro Tyr Pro Gln Ala Pro Ser Gly Pro Thr Phe Tyr Ser Lys Thr  
                   530                                  535                                  540

Phe Pro Ile Leu Gly Ser Val Gly Asp Thr Phe Leu Tyr Leu Pro Gly  
 545                                  550                                  555                                  560

Trp Thr Lys Gly Gln Val Trp Ile Asn Gly Phe Asn Leu Gly Arg Tyr  
                                   565                                  570                                  575

Trp Thr Lys Gln Gly Pro Gln Gln Thr Leu Tyr Val Pro Arg Phe Leu  
                                   580                                  585                                  590

Leu Phe Pro Arg Gly Ala Leu Asn Lys Ile Thr Leu Leu Glu Leu Glu  
                   595                                  600                                  605

Asp Val Pro Leu Gln Pro Gln Val Gln Phe Leu Asp Lys Pro Ile Leu  
                   610                                  615                                  620

Asn Ser Thr Ser Thr Leu His Arg Thr His Ile Asn Ser Leu Ser Ala  
 625                                  630                                  635                                  640

Asp Thr Leu Ser Ala Ser Glu Pro Met Glu Leu Ser Gly His  
                                   645                                  650

---

-continued

---

<210> SEQ ID NO 178  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 178  
  
tggctactcc aagaccctgg catg 24

<210> SEQ ID NO 179  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 179  
  
tggacaaaatc cccttgctca gcc 24

<210> SEQ ID NO 180  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 180  
  
gggcttcacc gaagcagtg acctttatt tgaccacctg atgtccagg 50

<210> SEQ ID NO 181  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 181  
  
ccagctatga ctatgatgca cc 22

<210> SEQ ID NO 182  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 182  
  
tggcaccag aatggtggtg gctc 24

<210> SEQ ID NO 183  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe  
  
<400> SEQUENCE: 183  
  
cgagatgca tcagcaagtt ccaggaagtt ccttgggac ctttacctc 50



---

-continued

---

<210> SEQ ID NO 184

<211> LENGTH: 1947

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 184

```
gctttgaaca cgtctgcaag cccaaagtg agcatctgat tggttatgag gtatttgagt      60
gcaccacaaa tatggcttac atgttgaaaa agcttctcat cagttacata tccattattt      120
gtgtttatgg ctttatctgc ctctacactc tcttctggtt attcaggata ctttgaaagg      180
aatattcttt cgaaaaagtc agagaagaga gcagttttag tgacattcca gatgtcaaaa      240
acgattttgc gttccttctt cacatggtag accagtatga ccagctatat tccaagcgtt      300
ttggtgtggt cttgtcagaa gtttagtgaataaaacttag ggaattagtt ttgaaccatg      360
agtggacatt tgaaaaactc aggcagcaca tttcacgcaa cgcccaggac aagcaggagt      420
tgcatctggt catgctgtcg ggggtgcccg atgctgtctt tgacctcaca gacctggatg      480
tgctaagctt tgaactaatt ccagaagcta aaattcctgc taagatttct caaatgacta      540
acctccaaga gctccacctc tgccactgcc ctgcaaaagt tgaacagact gcttttagct      600
ttcttcgcga tcacttgaga tgccctcacg tgaagttcac tgatgtggct gaaattcctg      660
cctgggtgta tttgctcaaa aaccttcgag agttgtactt aataggcaat ttgaactctg      720
aaaacaataa gatgatagga ctggaatctc tccgagagtt gcggcacctt aagattctcc      780
acgtgaagag caatttgacc aaagttccct ccaacattac agatgtggct ccacatctta      840
caaagttagt cattcataat gacggcacta aactcctggg actgaacagc cttaagaaaa      900
tgatgaatgt cgctgagctg gaactccaga actgtgagct agagagaatc ccacatgcta      960
ttttcagcct ctctaattta caggaactgg atttaaagtc caataacatt cgcacaattg     1020
aggaaatcat cagtttccag catttaaaac gactgacttg tttaaaatta tggcataaca     1080
aaattgttac tattcctccc tctattacc atgtcaaaaa cttggagtca ctttatttct     1140
ctaacaacaa gctcgaatcc ttaccagtgg cagtatttag tttacagaaa ctcatgatgct     1200
tagatgtgag ctacaacaac atttcaatga ttccaataga aataggattg cttcagaacc     1260
tgcagcattt gcatatcact gggaacaaaag tggacattct gccaaaacaa ttgtttaaata     1320
gcataaagtt gaggactttg aatctgggac agaactgcat cacctcactc ccagagaaaag     1380
ttggtcagct ctcccagctc actcagctgg agctgaaggg gaactgcttg gaccgctgc      1440
cagcccagct gggccagtgt cggatgctca agaaaagcgg gcttgtgtg gaagatcacc     1500
tttttgatac cctgccactc gaagtcaaag aggcattgaa tcaagacata aatattccct     1560
ttgcaaatgg gatttaaaact aagataatat atgcacagtg atgtgcagga acaacttcct     1620
agattgcaag tgctcacgta caagtattta caagataatg catttttagga gtagatacat     1680
cttttaaaat aaaacagaga ggatgcatag aaggctgata gaagacataa ctgaatgttc     1740
aatgtttgta gggttttaag tcattcattt ccaaatcatt ttttttttc ttttggggaa     1800
agggagggaa aaattataat cactaatctt ggttcttttt aaattgtttg taacttgat     1860
gctgccgcta ctgaatgttt acaaattgct tgcctgctaa agtaaatgat taaattgaca     1920
ttttcttact aaaaaaaaa aaaaaaa      1947
```

-continued

---

```

<210> SEQ ID NO 185
<211> LENGTH: 501
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 185

Met Ala Tyr Met Leu Lys Lys Leu Leu Ile Ser Tyr Ile Ser Ile Ile
 1          5          10          15
Cys Val Tyr Gly Phe Ile Cys Leu Tyr Thr Leu Phe Trp Leu Phe Arg
 20          25          30
Ile Pro Leu Lys Glu Tyr Ser Phe Glu Lys Val Arg Glu Glu Ser Ser
 35          40          45
Phe Ser Asp Ile Pro Asp Val Lys Asn Asp Phe Ala Phe Leu Leu His
 50          55          60
Met Val Asp Gln Tyr Asp Gln Leu Tyr Ser Lys Arg Phe Gly Val Phe
 65          70          75          80
Leu Ser Glu Val Ser Glu Asn Lys Leu Arg Glu Ile Ser Leu Asn His
 85          90          95
Glu Trp Thr Phe Glu Lys Leu Arg Gln His Ile Ser Arg Asn Ala Gln
100          105          110
Asp Lys Gln Glu Leu His Leu Phe Met Leu Ser Gly Val Pro Asp Ala
115          120          125
Val Phe Asp Leu Thr Asp Leu Asp Val Leu Lys Leu Glu Leu Ile Pro
130          135          140
Glu Ala Lys Ile Pro Ala Lys Ile Ser Gln Met Thr Asn Leu Gln Glu
145          150          155          160
Leu His Leu Cys His Cys Pro Ala Lys Val Glu Gln Thr Ala Phe Ser
165          170          175
Phe Leu Arg Asp His Leu Arg Cys Leu His Val Lys Phe Thr Asp Val
180          185          190
Ala Glu Ile Pro Ala Trp Val Tyr Leu Leu Lys Asn Leu Arg Glu Leu
195          200          205
Tyr Leu Ile Gly Asn Leu Asn Ser Glu Asn Asn Lys Met Ile Gly Leu
210          215          220
Glu Ser Leu Arg Glu Leu Arg His Leu Lys Ile Leu His Val Lys Ser
225          230          235          240
Asn Leu Thr Lys Val Pro Ser Asn Ile Thr Asp Val Ala Pro His Leu
245          250          255
Thr Lys Leu Val Ile His Asn Asp Gly Thr Lys Leu Leu Val Leu Asn
260          265          270
Ser Leu Lys Lys Met Met Asn Val Ala Glu Leu Glu Leu Gln Asn Cys
275          280          285
Glu Leu Glu Arg Ile Pro His Ala Ile Phe Ser Leu Ser Asn Leu Gln
290          295          300
Glu Leu Asp Leu Lys Ser Asn Asn Ile Arg Thr Ile Glu Glu Ile Ile
305          310          315          320
Ser Phe Gln His Leu Lys Arg Leu Thr Cys Leu Lys Leu Trp His Asn
325          330          335
Lys Ile Val Thr Ile Pro Pro Ser Ile Thr His Val Lys Asn Leu Glu
340          345          350
Ser Leu Tyr Phe Ser Asn Asn Lys Leu Glu Ser Leu Pro Val Ala Val
355          360          365

```

-continued

---

Phe Ser Leu Gln Lys Leu Arg Cys Leu Asp Val Ser Tyr Asn Asn Ile  
 370 375 380

Ser Met Ile Pro Ile Glu Ile Gly Leu Leu Gln Asn Leu Gln His Leu  
 385 390 395 400

His Ile Thr Gly Asn Lys Val Asp Ile Leu Pro Lys Gln Leu Phe Lys  
 405 410 415

Cys Ile Lys Leu Arg Thr Leu Asn Leu Gly Gln Asn Cys Ile Thr Ser  
 420 425 430

Leu Pro Glu Lys Val Gly Gln Leu Ser Gln Leu Thr Gln Leu Glu Leu  
 435 440 445

Lys Gly Asn Cys Leu Asp Arg Leu Pro Ala Gln Leu Gly Gln Cys Arg  
 450 455 460

Met Leu Lys Lys Ser Gly Leu Val Val Glu Asp His Leu Phe Asp Thr  
 465 470 475 480

Leu Pro Leu Glu Val Lys Glu Ala Leu Asn Gln Asp Ile Asn Ile Pro  
 485 490 495

Phe Ala Asn Gly Ile  
 500

<210> SEQ ID NO 186  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 186

cctccctcta ttaccatgt c 21

<210> SEQ ID NO 187  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 187

gaccaacttt ctctgggagt gagg 24

<210> SEQ ID NO 188  
 <211> LENGTH: 47  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 188

gtcactttat ttctctaaca acaagctoga atccttacca gtggcag 47

<210> SEQ ID NO 189  
 <211> LENGTH: 2917  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <400> SEQUENCE: 189

cccacgcgctc cggccttctc tctggacttt gcatttccat tccttttcat tgacaaactg 60

-continued

acttttttta tttctttttt tccatctctg ggccagcttg ggatcctagg ccgccctggg	120
aagacatttg tgttttacac acataaggat ctgtgtttgg ggtttcttct tctcccctg	180
acattggcat tgcttagtgg ttgtgtgggg agggagacca cgtgggctca gtgcttgctt	240
gcacttatct gcctaggtac atcgaagtct tttgacctcc atacagtgat tatgcctgtc	300
atcgctggtg gtatcctggc ggccctgtct ctgctgatag ttgtcgtgct ctgtctttac	360
ttcaaaatac acaacgcgct aaaagctgca aaggaaactg aagctgtggc tgtaaaaaat	420
cacaaccaga acaaggtgtg gtgggccaag aacagccagg ccaaaacctat tgccaaggag	480
tcttgtcctg ccctgcagtg ctgtgaagga tataagaatgt gtgccagttt tgattccctg	540
ccacctgtct gttgcatcat aatgagggc ctctgagtta ggaaggctc ccttctcaaa	600
gcagagccct gaagacttca atgatgtcaa tgaggccacc tgtttgtgat gtgcaggcac	660
agaagaaagg cacagctccc catcagtttc atggaataa actcagtgcc tgctgggaac	720
cagctgtctg agatcccctac agagagcttc cactgggggc aaccttcca ggaaggagtt	780
gggagagag aacctcact gtggggaatg ctgataaacc agtcacacag ctgctctatt	840
ctcacacaaa tctaccctct gcgtggctgg aactgacgtt tccctggagg tgtccagaaa	900
gctgatgtaa cacagagcct ataaaagctg tcggctctta aggctgcca gcgccttgcc	960
aaaatggagc ttgtaagaag gctcatgcca ttgacctctc taattctctc ctggttgagg	1020
gagctgacaa tggcggaggc tgaaggcaat gcaagctgca cagtcagtct aggggggtgc	1080
aatatggcag agaccacaa agccatgatc ctgcaactca atcccagtga gaactgcacc	1140
tggacaatag aaagaccaga aaacaaaagc atcagaatta tcttttcta tgtccagctt	1200
gatccagatg gaagctgtga aagtgaatac attaaagtct ttgacggaac ctccagcaat	1260
gggcctctgc tagggcaagt ctgcagtaaa aacgactatg ttctgtatt tgaatcatca	1320
tccagttatc tgacgtttca aatagttact gactcagcaa gaattcaaag aactgtcttt	1380
gtctctact acttctctc tcctaacatc tctattcaa actgtggcgg ttacctggat	1440
accttggaag gatccttcac cagcccctat taccacaaag cgcctcctga gctggcttat	1500
tgtgtgtggc acatacaagt ggagaaagat tacaagataa aactaaactt caaagagatt	1560
ttcctagaaa tagacaaaac gtgcaaatth gattttcttg ccatctatga tggcccctcc	1620
accaactctg gcctgattgg acaagtctgt ggccgtgtga ctcccactt cgaatcgtca	1680
tcaactctc tgactgtcgt gttgtctaca gattatgcca attcttaccg gggattttct	1740
gcttctaca cctcaattta tgcagaaaac atcaacacta catctttaac ttgctctct	1800
gacaggatga gagtattat aagcaaatcc tacctagagg cttttaactc taatgggaat	1860
aacttgcaac taaaagacc aactgcaga ccaaaattat caaatgtgt ggaattttct	1920
gtccctctta atgagtggtg tacaatcaga aaggtagaag atcagtcact tacttacacc	1980
aatataatca cctttctgct atctcactc totgaagtga tcaccctgca gaaacaactc	2040
cagattattg tgaagtgatg aatgggacat aattctacag tggagataat atacataaca	2100
gaagatgatg taatacaaa tcaaaatgca ctgggcaaat ataaccaccag catggctctt	2160
tttgaatcoa attcatttga aaagactata cttgaatcac catattatgt ggatttgaac	2220
caaactcttt ttgttcaagt tagtctgcac acctcagatc caaatttggg ggtgtttctt	2280
gatacctgta gagcctctcc cacctctgac tttgcatctc caacctacga cctaatcaag	2340

## -continued

---

```

agtgatgta gtcgagatga aacttgtaag gtgtatccct tatttggaaca ctatgggaga 2400
ttccagttta atgcctttaa attcttgaga agtatgagct ctgtgtatct gcagtgtaaa 2460
gttttgatat gtgatagcag tgaccaccag tctcgctgca atcaagggtg tgtctccaga 2520
agcaaacgag acatttcttc atataaatgg aaaacagatt ccatcatagg acccattcgt 2580
ctgaaaaggg atcgaagtgc aagtggcaat tcaggatttc agcatgaaac acatgcgaa 2640
gaaactccaa accagccttt caacagtgtg catctgtttt ccttcatggt tctagctctg 2700
aatgtgtgta ctgtagcagc aatcacagtg aggcattttg taaatcaacg ggcagactac 2760
aaataccaga agctgcagaa ctattaacta acaggtccaa ccctaagtga gacatgtttc 2820
tccaggatgc caaaggaat gctacctcgt ggctacacat attatgaata aatgaggaag 2880
ggcctgaaag tgacacacag gcctgcattg aaaaaaa 2917

```

<210> SEQ ID NO 190

<211> LENGTH: 607

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 190

```

Met Glu Leu Val Arg Arg Leu Met Pro Leu Thr Leu Leu Ile Leu Ser
 1                    5                    10          15
Cys Leu Ala Glu Leu Thr Met Ala Glu Ala Glu Gly Asn Ala Ser Cys
 20                    25                    30
Thr Val Ser Leu Gly Gly Ala Asn Met Ala Glu Thr His Lys Ala Met
 35                    40                    45
Ile Leu Gln Leu Asn Pro Ser Glu Asn Cys Thr Trp Thr Ile Glu Arg
 50                    55                    60
Pro Glu Asn Lys Ser Ile Arg Ile Ile Phe Ser Tyr Val Gln Leu Asp
 65                    70                    75                    80
Pro Asp Gly Ser Cys Glu Ser Glu Asn Ile Lys Val Phe Asp Gly Thr
 85                    90                    95
Ser Ser Asn Gly Pro Leu Leu Gly Gln Val Cys Ser Lys Asn Asp Tyr
100                    105                    110
Val Pro Val Phe Glu Ser Ser Ser Ser Thr Leu Thr Phe Gln Ile Val
115                    120                    125
Thr Asp Ser Ala Arg Ile Gln Arg Thr Val Phe Val Phe Tyr Tyr Phe
130                    135                    140
Phe Ser Pro Asn Ile Ser Ile Pro Asn Cys Gly Gly Tyr Leu Asp Thr
145                    150                    155                    160
Leu Glu Gly Ser Phe Thr Ser Pro Asn Tyr Pro Lys Pro His Pro Glu
165                    170                    175
Leu Ala Tyr Cys Val Trp His Ile Gln Val Glu Lys Asp Tyr Lys Ile
180                    185                    190
Lys Leu Asn Phe Lys Glu Ile Phe Leu Glu Ile Asp Lys Gln Cys Lys
195                    200                    205
Phe Asp Phe Leu Ala Ile Tyr Asp Gly Pro Ser Thr Asn Ser Gly Leu
210                    215                    220
Ile Gly Gln Val Cys Gly Arg Val Thr Pro Thr Phe Glu Ser Ser Ser
225                    230                    235                    240
Asn Ser Leu Thr Val Val Leu Ser Thr Asp Tyr Ala Asn Ser Tyr Arg
245                    250                    255

```



-continued

---

tctctattcc aaactgtggc g 21

<210> SEQ ID NO 192  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 192

tttgatgacg attcgaaggt gg 22

<210> SEQ ID NO 193  
 <211> LENGTH: 47  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 193

ggaagatcc ttcaccagcc ccaattaccc aaagccgcat cctgagc 47

<210> SEQ ID NO 194  
 <211> LENGTH: 2362  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 194

gacggaagaa cagcgcctccc gaggccgcgg gagcctgcag agaggacagc cggcctgcgc 60

cgggacatgc ggccccagga gctccccagg ctgcggttcc cgttgctgct gttgctgttg 120

ctgctgtctgc cgccgcgcc gtgcctgcc cacagcgcca cgcgcttca cccacctgg 180

gagtcctgg acgcccgcca gctgcccgg tggtttgacc aggccaaagt cggcatcttc 240

atccactggg gagtgttttc cgtgcccagc ttcggtagcg agtggttctg gtggtattgg 300

caaaaggaaa agataccgaa gtatgtgaa tttatgaaag ataattaccc tcctagtttc 360

aaatatgaag attttgacc actatttaca gcaaaatfff ttaatgcaa ccatggggca 420

gatatttttc aggcctctgg tgccaaatac attgtcttaa cttccaaaca tcatgaaggc 480

tttacctgtg ggggctcaga atattcgtgg aactggaatg ccatagatga ggggccaag 540

agggacattg tcaaggaact tgaggtagcc attaggaaca gaactgacct gcgttttggg 600

ctgtactatt cccttttga atggtttcat ccgctcttcc ttgaggatga atccagtcca 660

ttccataagc ggcaatttcc agtttctaag acattgccag agctctatga gttagtgaac 720

aactatcagc ctgaggttct gtggtcggat ggtgacggag gagcaccgga tcaatactgg 780

aacagcacag gcttcttggc ctgggtatat aatgaaagcc cagttcgggg cacagtagtc 840

accaatgac gttggggagc tggtagcatc tgtaagcatg gttgcttcta tacctgcagt 900

gatcgttata acccagaca tcttttgcca cataaatggg aaaactgcat gacaatagac 960

aaactgtcct ggggctatag gagggaaact ggaatctctg actatcttac aattgaagaa 1020

ttggtgaagc aactgtaga gacagtttca tgtggaggaa atcttttgat gaatattggg 1080

cccacactag atggcaccat ttctgtagtt tttgaggagc gactgaggca agtggggtcc 1140

tggctaaaag tcaatggaga agctatttat gaaacctata cctggcagtc ccagaatgac 1200

-continued

```

actgtcacc cagatgtgtg gtacacatcc aagcctaaag aaaaattagt ctatgccatt 1260
tttcttaaat ggcccacatc aggacagctg ttccttggcc atcccaaagc tattctggtg 1320
gcaacagagg tgaactact gggccatgga cagccactta actggatttc tttggagcaa 1380
aatggcatta tggtagaact gccacagcta accattcadc agatgccgtg taaatggggc 1440
tggtctctag ccctaactaa tgtgatctaa agtgcagcag agtggctgat gctgcaagtt 1500
atgtctaagg ctaggaacta tcagtggtct ataattgtag cacatggaga aagcaatgta 1560
aactggataa gaaaattatt tggcagtcca gccctttccc tttttccac taaatttttc 1620
ttaaataacc catgtaacca tttaactct ccagtgcact ttgccattaa agtctcttca 1680
cattgatttg tttccatgtg tgactcagag gtgagaatth tttcacatta tagtagcaag 1740
gaattgggtg tattatggac cgaactgaaa attttatggt gaagccatat ccccatgat 1800
tatatagtta tgcacatcct aatatgggga tttttctgg gaaatgcatt gctagtcaat 1860
tttttttgt gccaacatca tagagtgtat ttacaaaatc ctagatggca tagcctacta 1920
cacaccta atgtatggta tagactgttg ctcctaggct acagacatat acagcatggt 1980
actgaatact gtaggcaata gtaacagtgg tatttgtata tcgaaacata tggaaacata 2040
gagaaggtag agtaaaaata ctgtaaaata aatggtgcac ctgtataggg cacttaccac 2100
gaatggagct tacaggactg gaagttgctc tgggtgagtc agtgagtga tgtgaaggcc 2160
taggacatta ttgaacactg ccagacgtta taaatactgt atgottaggc tacactacat 2220
ttataaaaaa aagtttttct ttcttcaatt ataaattaac ataagtgtac tgtaacttta 2280
caaacgtttt aatttttaaa accttttttg ctcttttgta ataacactta gottaaaaca 2340
taaactcatt gtgcaaatgt aa 2362

```

```

<210> SEQ ID NO 195
<211> LENGTH: 467
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 195

```

Met Arg Pro Gln Glu Leu Pro Arg Leu Ala Phe Pro Leu Leu Leu Leu
 1           5           10          15
Leu Leu Leu Leu Leu Pro Pro Pro Pro Cys Pro Ala His Ser Ala Thr
 20          25          30
Arg Phe Asp Pro Thr Trp Glu Ser Leu Asp Ala Arg Gln Leu Pro Ala
 35          40          45
Trp Phe Asp Gln Ala Lys Phe Gly Ile Phe Ile His Trp Gly Val Phe
 50          55          60
Ser Val Pro Ser Phe Gly Ser Glu Trp Phe Trp Trp Tyr Trp Gln Lys
 65          70          75          80
Glu Lys Ile Pro Lys Tyr Val Glu Phe Met Lys Asp Asn Tyr Pro Pro
 85          90          95
Ser Phe Lys Tyr Glu Asp Phe Gly Pro Leu Phe Thr Ala Lys Phe Phe
100         105         110
Asn Ala Asn Gln Trp Ala Asp Ile Phe Gln Ala Ser Gly Ala Lys Tyr
115         120         125
Ile Val Leu Thr Ser Lys His His Glu Gly Phe Thr Leu Trp Gly Ser
130         135         140

```



-continued

---

Glu Tyr Ser Trp Asn Trp Asn Ala Ile Asp Glu Gly Pro Lys Arg Asp  
 145 150 155 160  
 Ile Val Lys Glu Leu Glu Val Ala Ile Arg Asn Arg Thr Asp Leu Arg  
 165 170 175  
 Phe Gly Leu Tyr Tyr Ser Leu Phe Glu Trp Phe His Pro Leu Phe Leu  
 180 185 190  
 Glu Asp Glu Ser Ser Ser Phe His Lys Arg Gln Phe Pro Val Ser Lys  
 195 200 205  
 Thr Leu Pro Glu Leu Tyr Glu Leu Val Asn Asn Tyr Gln Pro Glu Val  
 210 215 220  
 Leu Trp Ser Asp Gly Asp Gly Gly Ala Pro Asp Gln Tyr Trp Asn Ser  
 225 230 235 240  
 Thr Gly Phe Leu Ala Trp Leu Tyr Asn Glu Ser Pro Val Arg Gly Thr  
 245 250 255  
 Val Val Thr Asn Asp Arg Trp Gly Ala Gly Ser Ile Cys Lys His Gly  
 260 265 270  
 Gly Phe Tyr Thr Cys Ser Asp Arg Tyr Asn Pro Gly His Leu Leu Pro  
 275 280 285  
 His Lys Trp Glu Asn Cys Met Thr Ile Asp Lys Leu Ser Trp Gly Tyr  
 290 295 300  
 Arg Arg Glu Ala Gly Ile Ser Asp Tyr Leu Thr Ile Glu Glu Leu Val  
 305 310 315 320  
 Lys Gln Leu Val Glu Thr Val Ser Cys Gly Gly Asn Leu Leu Met Asn  
 325 330 335  
 Ile Gly Pro Thr Leu Asp Gly Thr Ile Ser Val Val Phe Glu Glu Arg  
 340 345 350  
 Leu Arg Gln Val Gly Ser Trp Leu Lys Val Asn Gly Glu Ala Ile Tyr  
 355 360 365  
 Glu Thr Tyr Thr Trp Arg Ser Gln Asn Asp Thr Val Thr Pro Asp Val  
 370 375 380  
 Trp Tyr Thr Ser Lys Pro Lys Glu Lys Leu Val Tyr Ala Ile Phe Leu  
 385 390 395 400  
 Lys Trp Pro Thr Ser Gly Gln Leu Phe Leu Gly His Pro Lys Ala Ile  
 405 410 415  
 Leu Gly Ala Thr Glu Val Lys Leu Leu Gly His Gly Gln Pro Leu Asn  
 420 425 430  
 Trp Ile Ser Leu Glu Gln Asn Gly Ile Met Val Glu Leu Pro Gln Leu  
 435 440 445  
 Thr Ile His Gln Met Pro Cys Lys Trp Gly Trp Ala Leu Ala Leu Thr  
 450 455 460  
 Asn Val Ile  
 465

<210> SEQ ID NO 196  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe  
 <400> SEQUENCE: 196

tggtttgacc aggccaagtt cgg

-continued

---

```

<210> SEQ ID NO 197
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 197

ggattcatcc tcaaggaaga gcgg                                24

<210> SEQ ID NO 198
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 198

aacttgacgc atcagccact ctgc                                24

<210> SEQ ID NO 199
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 199

ttccgtgcc agcttcggta gcgagtgggt ctgggtggtat tggca      45

<210> SEQ ID NO 200
<211> LENGTH: 2372
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 200

agcagggaaa tccggatgtc tcggttatga agtggagcag tgagtgtgag cctcaacata    60
gttcagaaac tctccatccg gactagtatg tagcatctg cctctcatat caccagtggc    120
catctgaggt gtttccctgg ctctgaaggg gtaggcacga tggccagggt cttcagcctg    180
gtgttgcttc tcacttccat ctggaccacg aggctcctgg tccaaggctc tttgcgtgca    240
gaagagcttt ccatccagggt gtcattcaga attatgggga tcacccttgt gagcaaaaag    300
gcgaaccacg agctgaatth cacagaagct aaggaggcct gtaggctgct gggactaagt    360
ttggccggca aggaccaagt tgaacacgcc ttgaaagcta gctttgaaac ttgcagctat    420
ggctgggttg gagatggatt cgtggctatc tctaggatta gcccaaaccc caagtgtggg    480
aaaaatgggg tgggtgtcct gatttggaaag gttccagtga gccgacagtt tgcagcctat    540
tgttacaact catctgatac ttggactaac tegtgcattc cagaaattat caccacaaa    600
gatcccatat tcaactca aactgcaaca caaacaacag aatttattgt cagtgcagct    660
acctactcgg tggcatcccc ttactctaca atacctgccc ctactactac tctctctgct    720
ccagcttcca cttctattcc acggagaaaa aaattgattt gtgtcacaga agtttttatg    780
gaaactagca ccatgtctac agaaactgaa ccatttgttg aaaataaagc agcattcaag    840
aatgaagctg ctgggtttgg aggtgtcccc acggtctctg tagtgcttgc tctctctctc    900

```

-continued

```

tttggtgctg cagctggtct tggatthttgc tatgtcaaaa ggtatgtgaa ggccttccct 960
tttacaaaaca agaatcagca gaaggaaatg atcgaaacca aagtagtaaa ggaggagaag 1020
gccaatgata gcaaccctaa tgaggaaatca aagaaaactg ataaaaaccg agaagagtcc 1080
aagagtccaa gcaaaactac cgtgcatgac ctggaagctg aagtttagat gagacagaaa 1140
tgaggagaca cacctgaggc tggthttcttt catgctcctt accctgcccc agctggggaa 1200
atcaaaaagg ccaaaagaacc aaagaagaaa gtccaccctt ggttcctaac tggaatcagc 1260
tcaggactgc cattggacta tggagtgcac caaagagaat gcccttctcc ttattgtaac 1320
cctgtctgga tcctatcctc ctaccctcaa agcttcccac ggcctttcta gcctggctat 1380
gtcctaataa tatccactg ggagaaagga gthttgcaaa gtgcaaggac ctaaaacatc 1440
tcactcagat ccagtgtgaa aaaggcctcc tggctgtctg aggttaggtg gttgaaagc 1500
caaggagtca ctgagaccaa ggctthttct actgattccg cagctcagac cthttcttca 1560
gctctgaaaag aaaaacagct atcccactg acatgtcctt ctgagcccgg taagagcaaa 1620
agaatggcag aaaaatttag ccctgaaag ccatggagat tctcataact tgagacctaa 1680
tctctgtaaa gtaaaaaata aaaaatagaa caaggctgag gatacagacag tacactgtca 1740
gcagggactg taaacacaga cagggtcaaa gtgthttctc tgaacacatt gagttggaat 1800
cactgtttag aacacacaca ctactthttt ctggtctcta ccaactgtga tathttctct 1860
aggaatata cthttacaag taacaaaaat aaaaactctt ataaatttct atthttatct 1920
gagttacaga aatgattact aaggaagatt actcagtaat ttgtthaaaa agtaataaaa 1980
ttcaacaaac atthgtgtaa tagctactat atgtcaagtg ctgtgcaagg tattacactc 2040
tgtaattgaa tattattcct caaaaattg cacatagtag aacgtatctt ggaagctat 2100
thttttcagt thtgataatt ctagcttctc tacttccaaa ctaathttta thtttgctga 2160
gactaatctt attcattttc tctaatatgg caaccattat aaccttaatt tattattaac 2220
atacctaaga agtacattgt tactctata taccaaagca cthtttaaaa gtgccattaa 2280
caaatgtatc actagccctc cthtttccaa caagaagga ctgagagatg cagaaatatt 2340
tgtgacaaaa aattaaagca thtagaaaac tt 2372

```

&lt;210&gt; SEQ ID NO 201

&lt;211&gt; LENGTH: 322

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic protein

&lt;400&gt; SEQUENCE: 201

```

Met Ala Arg Cys Phe Ser Leu Val Leu Leu Thr Ser Ile Trp Thr
 1           5           10          15

```

```

Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala Glu Glu Leu Ser Ile
 20          25          30

```

```

Gln Val Ser Cys Arg Ile Met Gly Ile Thr Leu Val Ser Lys Lys Ala
 35          40          45

```

```

Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys Glu Ala Cys Arg Leu Leu
 50          55          60

```

```

Gly Leu Ser Leu Ala Gly Lys Asp Gln Val Glu Thr Ala Leu Lys Ala
 65          70          75          80

```

-continued

---

Ser Phe Glu Thr Cys Ser Tyr Gly Trp Val Gly Asp Gly Phe Val Val  
 85 90 95

Ile Ser Arg Ile Ser Pro Asn Pro Lys Cys Gly Lys Asn Gly Val Gly  
 100 105 110

Val Leu Ile Trp Lys Val Pro Val Ser Arg Gln Phe Ala Ala Tyr Cys  
 115 120 125

Tyr Asn Ser Ser Asp Thr Trp Thr Asn Ser Cys Ile Pro Glu Ile Ile  
 130 135 140

Thr Thr Lys Asp Pro Ile Phe Asn Thr Gln Thr Ala Thr Gln Thr Thr  
 145 150 155 160

Glu Phe Ile Val Ser Asp Ser Thr Tyr Ser Val Ala Ser Pro Tyr Ser  
 165 170 175

Thr Ile Pro Ala Pro Thr Thr Thr Pro Pro Ala Pro Ala Ser Thr Ser  
 180 185 190

Ile Pro Arg Arg Lys Lys Leu Ile Cys Val Thr Glu Val Phe Met Glu  
 195 200 205

Thr Ser Thr Met Ser Thr Glu Thr Glu Pro Phe Val Glu Asn Lys Ala  
 210 215 220

Ala Phe Lys Asn Glu Ala Ala Gly Phe Gly Gly Val Pro Thr Ala Leu  
 225 230 235 240

Leu Val Leu Ala Leu Leu Phe Phe Gly Ala Ala Ala Gly Leu Gly Phe  
 245 250 255

Cys Tyr Val Lys Arg Tyr Val Lys Ala Phe Pro Phe Thr Asn Lys Asn  
 260 265 270

Gln Gln Lys Glu Met Ile Glu Thr Lys Val Val Lys Glu Glu Lys Ala  
 275 280 285

Asn Asp Ser Asn Pro Asn Glu Glu Ser Lys Lys Thr Asp Lys Asn Pro  
 290 295 300

Glu Glu Ser Lys Ser Pro Ser Lys Thr Thr Val Arg Cys Leu Glu Ala  
 305 310 315 320

Glu Val

<210> SEQ ID NO 202  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 202

gagctttcca tccaggtgtc atgc

24

<210> SEQ ID NO 203  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 203

gtcagtgaca gtacctactc gg

22

<210> SEQ ID NO 204  
 <211> LENGTH: 24

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 204

tggagcagga ggagtagtag tagg                                24

<210> SEQ ID NO 205
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide probe

<400> SEQUENCE: 205

aggaggcctg taggctgctg ggactaagtt tggccggcaa ggaccaagtt    50

<210> SEQ ID NO 206
<211> LENGTH: 1620
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (973)
<223> OTHER INFORMATION: a, t, c or g
<221> NAME/KEY: modified_base
<222> LOCATION: (977)
<223> OTHER INFORMATION: a, t, c or g
<221> NAME/KEY: modified_base
<222> LOCATION: (996)
<223> OTHER INFORMATION: a, t, c or g
<221> NAME/KEY: modified_base
<222> LOCATION: (1003)
<223> OTHER INFORMATION: a, t, c or g

<400> SEQUENCE: 206

agatggcgggt cttggcacct ctaattgctc tcgtgtattc ggtgccgca ctttcacgat    60
ggctgcgccac ctttactac cttctgtcgg ccctgctctc tgetgccttc ctactcgtga    120
ggaaactgcc gccgctctgc cacggtctgc ccaccaacg cgaagacggt aaccctgtg    180
actttgactg gagagaagtg gagatcctga tgtttctcag tgccattgtg atgatgaaga    240
accgcagatc catcactgtg gagcaacata taggcaacat tttcatgttt agtaaagtgg    300
ccaacacaat tcttttcttc cgcttgata ttcgcatggg cctactttac atcacactct    360
gcatagtgtt cctgatgacg tgcaaacccc ccctatatat gggccctgag tatatcaagt    420
acttcaatga taaaaccatt gatgaggaac tagaacggga caagagggtc acttggtattg    480
tggagttctt tgccaattgg tctaattgact gccaatcatt tgcccctatc tatgctgacc    540
tctcccattaa atacaactgt acagggctaa attttgggaa ggtggatgtt ggacgtata    600
ctgatgttag tacgcggtac aaagtgagca catcaccctc caccaagcaa ctcctaccc    660
tgatcctgtt ccaaggtggc aaggaggcaa tgcggcggcc acagattgac aagaaggac    720
gggtgtgtct atggaccttc tctgaggaga atgtgatccg agaatttaac ttaaatgagc    780
tataccagcg ggccaagaaa ctatcaaagg ctggagacaa tatccctgag gagcagcctg    840
tggcttcaac ccccaccaca gtgtcagatg gggaaaacaa gaaggataaa taagatcctc    900
actttggcag tgcttctctc cctgtcaatt ccaggctctt tccataacca caagcctgag    960

```

-continued

```

gctgcagcct ttnattnatg ttttcccttt ggctgngact ggntggggca gcatgcagct 1020
tctgatttta aagaggcatc tagggaattg tcaggcacc tacaggaagg cctgccatgc 1080
tgtggccaac tgtttactg gagcaagaaa gagatctcat aggacggagg gggaaatggt 1140
ttccctccaa gcttgggtca gtgtgttaac tgcttatcag ctattcagac atctccatgg 1200
tttctccatg aaactctgtg gtttcatcat tccttcttag ttgacctgca cagcttggtt 1260
agacctagat ttaaccctaa ggtaagatgc tggggtatag aacgctaaga attttccccc 1320
aaggactcct gtttccctaa gccttcttgg cttcgtttat ggtcttcatt aaaagtataa 1380
gcctaacttt gtcgctagtc ctaaggagaa accttaacc acaaagtttt tatcattgaa 1440
gacaatattg aacaaccccc tattttgtgg ggattgagaa ggggtgaata gaggcttgag 1500
actttccttt gtgtggttag acttgaggga gaaatcccct ggaactttcac taaccctctg 1560
acatactccc cacaccagct tgatggcttt cagtaataaa aagattggga tttccttttg 1620

```

&lt;210&gt; SEQ ID NO 207

&lt;211&gt; LENGTH: 296

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 207

```

Met Ala Val Leu Ala Pro Leu Ile Ala Leu Val Tyr Ser Val Pro Arg
 1           5           10          15
Leu Ser Arg Trp Leu Ala Gln Pro Tyr Tyr Leu Leu Ser Ala Leu Leu
          20          25          30
Ser Ala Ala Phe Leu Leu Val Arg Lys Leu Pro Pro Leu Cys His Gly
          35          40          45
Leu Pro Thr Gln Arg Glu Asp Gly Asn Pro Cys Asp Phe Asp Trp Arg
          50          55          60
Glu Val Glu Ile Leu Met Phe Leu Ser Ala Ile Val Met Met Lys Asn
          65          70          75          80
Arg Arg Ser Ile Thr Val Glu Gln His Ile Gly Asn Ile Phe Met Phe
          85          90          95
Ser Lys Val Ala Asn Thr Ile Leu Phe Phe Arg Leu Asp Ile Arg Met
          100         105         110
Gly Leu Leu Tyr Ile Thr Leu Cys Ile Val Phe Leu Met Thr Cys Lys
          115         120         125
Pro Pro Leu Tyr Met Gly Pro Glu Tyr Ile Lys Tyr Phe Asn Asp Lys
          130         135         140
Thr Ile Asp Glu Glu Leu Glu Arg Asp Lys Arg Val Thr Trp Ile Val
          145         150         155         160
Glu Phe Phe Ala Asn Trp Ser Asn Asp Cys Gln Ser Phe Ala Pro Ile
          165         170         175
Tyr Ala Asp Leu Ser Leu Lys Tyr Asn Cys Thr Gly Leu Asn Phe Gly
          180         185         190
Lys Val Asp Val Gly Arg Tyr Thr Asp Val Ser Thr Arg Tyr Lys Val
          195         200         205
Ser Thr Ser Pro Leu Thr Lys Gln Leu Pro Thr Leu Ile Leu Phe Gln
          210         215         220
Gly Gly Lys Glu Ala Met Arg Arg Pro Gln Ile Asp Lys Lys Gly Arg
          225         230         235         240
Ala Val Ser Trp Thr Phe Ser Glu Glu Asn Val Ile Arg Glu Phe Asn

```

-continued

		245	250	255												
Leu	Asn	Glu	Leu	Tyr	Gln	Arg	Ala	Lys	Lys	Leu	Ser	Lys	Ala	Gly	Asp	
		260						265						270		
Asn	Ile	Pro	Glu	Glu	Gln	Pro	Val	Ala	Ser	Thr	Pro	Thr	Thr	Val	Ser	
		275					280						285			
Asp	Gly	Glu	Asn	Lys	Lys	Asp	Lys									
	290					295										
<210> SEQ ID NO 208																
<211> LENGTH: 24																
<212> TYPE: DNA																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe																
<400> SEQUENCE: 208																
gcttggatat tcgcatgggc ctac															24	
<210> SEQ ID NO 209																
<211> LENGTH: 20																
<212> TYPE: DNA																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe																
<400> SEQUENCE: 209																
tggagacaat atccctgagg															20	
<210> SEQ ID NO 210																
<211> LENGTH: 24																
<212> TYPE: DNA																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe																
<400> SEQUENCE: 210																
aacagttggc cacagcatgg cagg															24	
<210> SEQ ID NO 211																
<211> LENGTH: 50																
<212> TYPE: DNA																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide probe																
<400> SEQUENCE: 211																
ccattgatga ggaactagaa cgggacaaga gggtcacttg gattgtggag															50	
<210> SEQ ID NO 212																
<211> LENGTH: 1985																
<212> TYPE: DNA																
<213> ORGANISM: Homo sapiens																
<400> SEQUENCE: 212																
ggacagctcg cggccccga gagctctagc cgctcaggag ctgcctgggg acgtttgccc															60	
tggggcccca gcttgcccc ggtcaccctg goatgaggag atgggcctgt tgctcctggt															120	
cccattgctc ctgctgccc gctcctacgg actgccttc tacaacggct tctactactc															180	

-continued

```

caacagcgcc aacgaccaga acctaggcaa cggtcatggc aaagacctcc ttaatggagt 240
gaagctggtg gtggagacac ccgaggagac cctgttcacc taccaagggg ccagtgtgat 300
cctgccctgc cgctaccgct acgagccggc cctggctctcc ccgcgcgctg tgcgtgtcaa 360
atggtggaag ctgtcggaga acggggcccc agagaaggac gtgctggtgg ccatcgggct 420
gaggcaccgc tcctttgggg actaccaagg ccgctgtcac ctgcccagc acaaagagca 480
tgacgtctcg ctggagatcc aggatctgcg gctggaggac tatgggcgct accgctgtga 540
ggtcattgac gggctggagg atgaaagcgg tctggtggag ctggagctgc ggggtgtggt 600
ctttccttac cagtccccca acgggcgcta ccagttcaac ttccacgagg gccagcaggt 660
ctgtgcagag caggctgcgg tgggtgcctc ctttgagcag ctcttcggg cctgggagga 720
gggcctggac tggtgcaacg cgggctggct gcaggatgct acggtgcagt accccatcat 780
gttgccccgg cagccctgcg gtggcccagg cctggcacct ggcgtgcgaa gctacggccc 840
ccgccaccgc cgctgcacc gctatgatgt attctgcttc gctactgcc tcaagggcg 900
ggtgtactac ctggagcacc ctgagaagct gacgctgaca gaggcaaggg aggcctgcca 960
ggaagatgat gccacgatcg ccaaggtggg acagctcttt gccgcctgga agttccatgg 1020
cctggaccgc tgcgacgctg gctggctggc agatggcagc gtccgctacc ctgtggttca 1080
cccgcactct aactgtgggc ccccagagcc tggggtcoga agctttggt tccccgacc 1140
gcagagccgc ttgtacgggt tttactgcta ccgcccagcac taggacctgg gccctcccc 1200
tgccgcattc cctcactggc tgtgtattta ttgagtgggt cgttttccct tgtgggttgg 1260
agccatttta actgttttta tacttctcaa tttaaatfff ctttaaacat ttttttacta 1320
ttttttgtaa agcaaacaga acccaatgcc tccctttgct cctggatgcc ccaactcagg 1380
aatcatgctt gctccccggt gccatttgcg gttttgtggg cttctggagg gttccccgcc 1440
atccaggctg gtctccctcc cttaaggagg ttggtgcccc gagtggggcg tggcctgtct 1500
agaatgcccg cgggagtcgg ggcgatggtg gcacagttct ccctgcccct cagcctgggg 1560
gaagaagagg gcctcggggg cctccggagc tgggctttgg gcctctcctg cccacctcta 1620
cttctctgtg aagccgctga ccccagctcg cccactgagg ggctagggct ggaagccagt 1680
tctaggcttc caggcgaat ctgagggaag gaagaaactc ccctccccgt tccccttccc 1740
ctctcggttc caaagaatct gttttgtgt catttgtttc tcctgtttcc ctgtgtgggg 1800
aggggccctc aggtgtgtgt actttggaca ataaatggtg ctatgactgc ctccgccaa 1860
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1920
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1980
aaaaaa 1985

```

&lt;210&gt; SEQ ID NO 213

&lt;211&gt; LENGTH: 360

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 213

```

Met Gly Leu Leu Leu Leu Val Pro Leu Leu Leu Leu Pro Gly Ser Tyr
  1             5             10             15

```

```

Gly Leu Pro Phe Tyr Asn Gly Phe Tyr Tyr Ser Asn Ser Ala Asn Asp
          20             25             30

```



-continued

Gln Asn Leu Gly Asn Gly His Gly Lys Asp Leu Leu Asn Gly Val Lys  
 35 40 45

Leu Val Val Glu Thr Pro Glu Thr Leu Phe Thr Tyr Gln Gly Ala  
 50 55 60

Ser Val Ile Leu Pro Cys Arg Tyr Arg Tyr Glu Pro Ala Leu Val Ser  
 65 70 75 80

Pro Arg Arg Val Arg Val Lys Trp Trp Lys Leu Ser Glu Asn Gly Ala  
 85 90 95

Pro Glu Lys Asp Val Leu Val Ala Ile Gly Leu Arg His Arg Ser Phe  
 100 105 110

Gly Asp Tyr Gln Gly Arg Val His Leu Arg Gln Asp Lys Glu His Asp  
 115 120 125

Val Ser Leu Glu Ile Gln Asp Leu Arg Leu Glu Asp Tyr Gly Arg Tyr  
 130 135 140

Arg Cys Glu Val Ile Asp Gly Leu Glu Asp Glu Ser Gly Leu Val Glu  
 145 150 155 160

Leu Glu Leu Arg Gly Val Val Phe Pro Tyr Gln Ser Pro Asn Gly Arg  
 165 170 175

Tyr Gln Phe Asn Phe His Glu Gly Gln Gln Val Cys Ala Glu Gln Ala  
 180 185 190

Ala Val Val Ala Ser Phe Glu Gln Leu Phe Arg Ala Trp Glu Glu Gly  
 195 200 205

Leu Asp Trp Cys Asn Ala Gly Trp Leu Gln Asp Ala Thr Val Gln Tyr  
 210 215 220

Pro Ile Met Leu Pro Arg Gln Pro Cys Gly Gly Pro Gly Leu Ala Pro  
 225 230 235 240

Gly Val Arg Ser Tyr Gly Pro Arg His Arg Arg Leu His Arg Tyr Asp  
 245 250 255

Val Phe Cys Phe Ala Thr Ala Leu Lys Gly Arg Val Tyr Tyr Leu Glu  
 260 265 270

His Pro Glu Lys Leu Thr Leu Thr Glu Ala Arg Glu Ala Cys Gln Glu  
 275 280 285

Asp Asp Ala Thr Ile Ala Lys Val Gly Gln Leu Phe Ala Ala Trp Lys  
 290 295 300

Phe His Gly Leu Asp Arg Cys Asp Ala Gly Trp Leu Ala Asp Gly Ser  
 305 310 315 320

Val Arg Tyr Pro Val Val His Pro His Pro Asn Cys Gly Pro Pro Glu  
 325 330 335

Pro Gly Val Arg Ser Phe Gly Phe Pro Asp Pro Gln Ser Arg Leu Tyr  
 340 345 350

Gly Val Tyr Cys Tyr Arg Gln His  
 355 360

<210> SEQ ID NO 214  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 214

tgcttcgcta ctgccctc

---

-continued

---

<210> SEQ ID NO 215  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 215  
ttcccttggtg ggttgag 18

<210> SEQ ID NO 216  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 216  
agggctggaa gccagttc 18

<210> SEQ ID NO 217  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 217  
agccagtgtg gaaatgctg 18

<210> SEQ ID NO 218  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 218  
tgtccaaagt acacacacct gagg 24

<210> SEQ ID NO 219  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 219  
gatgccacga tcgccaaggt gggacagctc tttgccgcct ggaag 45

<210> SEQ ID NO 220  
<211> LENGTH: 1503  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 220  
ggagagcgga gcgaagctgg ataacagggg accgatgatg tggcgacct cagttctgct 60

-continued

```

gcttctgttg ctactgaggc acggggccca ggggaagcca tccccagacg caggccctca 120
tggccagggg aggggtgcacc aggcggcccc cctgagcgac gctccccatg atgacgcccc 180
cgggaaacttc cagtacgacc atgaggtttt cctgggacgg gaagtggcca aggaattcga 240
ccaactcacc ccagaggaaa gccaggcccc tctggggcgg atcgtggacc gcatggaccg 300
cgcgggggac ggcgacggct ggggtgcgct ggccgagctt cgcgcggtga tcgcgcacac 360
gcagcagcgg cacatacggg actcggtag cgcgccctgg gacacgtacg acacggaccg 420
cgacggggct gtgggttggg aggagctgcg caacgccacc tatggccact acgccccgg 480
tgaagaatth catgacgtgg aggatgcaga gacctacaaa aagatgctgg ctcgggacga 540
gcggcgthtc cgggtggccc accaggatgg ggactcgtg gccactcgag aggagctgac 600
agccttctctg caccgccagg agttccctca catcggggac atcgtgattg ctgaaaccct 660
ggaggacctg gacagaaaca aagatggcta tgtccagggtg gaggagtaca tcgcggatct 720
gtactcagcc gagcctgggg aggagagacc ggcgtgggtg cagacggaga ggcagcagtt 780
ccgggacttc cgggatctga acaaggatgg gcacctggat gggagtgagg tgggccactg 840
ggtgctgccc cctgcccagg accagcccct ggtggaagcc aaccacctgc tgcacgagag 900
cgacacggac aagatggggc ggctgagcaa agcggaaatc ctgggtaatt ggaacatgtt 960
tgtgggcagt caggccacca actatggcga ggacctgacc cggcaccacg atgagctgtg 1020
agcaccgcgc acctgccaca gcctcagagg cccgcacaat gaccggagga ggggcccgtg 1080
tgggtctggcc ccctcccctgt ccaggccccg caggaggcag atgcagtccc aggcacccctc 1140
ctgcccctgg gctctcaggg acccccctgg tcggttctg tccctgtcac acccccacc 1200
ccagggaggg gctgtcatag tcccagagga taagcaatac ctatttctga ctgagtctcc 1260
cagcccagac ccagggaccc ttggccccc a gctcagctct aagaaccgcc ccaaccctc 1320
cagctccaaa tctgagcctc caccacatag actgaaactc ccctggcccc agccctctcc 1380
tgccctggct ggctcgggac acctcctctc tgccaggagg caataaaagc cagcgccggg 1440
accttgaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1500
aaa 1503
    
```

```

<210> SEQ ID NO 221
<211> LENGTH: 328
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 221

```

Met Met Trp Arg Pro Ser Val Leu Leu Leu Leu Leu Leu Arg His
 1           5           10           15
Gly Ala Gln Gly Lys Pro Ser Pro Asp Ala Gly Pro His Gly Gln Gly
          20           25           30
Arg Val His Gln Ala Ala Pro Leu Ser Asp Ala Pro His Asp Asp Ala
          35           40           45
His Gly Asn Phe Gln Tyr Asp His Glu Ala Phe Leu Gly Arg Glu Val
          50           55           60
Ala Lys Glu Phe Asp Gln Leu Thr Pro Glu Glu Ser Gln Ala Arg Leu
          65           70           75           80
Gly Arg Ile Val Asp Arg Met Asp Arg Ala Gly Asp Gly Asp Gly Trp
          85           90           95
    
```

-continued

---

Val Ser Leu Ala Glu Leu Arg Ala Trp Ile Ala His Thr Gln Gln Arg  
                   100                  105                  110

His Ile Arg Asp Ser Val Ser Ala Ala Trp Asp Thr Tyr Asp Thr Asp  
           115                  120                  125

Arg Asp Gly Arg Val Gly Trp Glu Glu Leu Arg Asn Ala Thr Tyr Gly  
           130                  135                  140

His Tyr Ala Pro Gly Glu Glu Phe His Asp Val Glu Asp Ala Glu Thr  
   145                  150                  155                  160

Tyr Lys Lys Met Leu Ala Arg Asp Glu Arg Arg Phe Arg Val Ala Asp  
           165                  170                  175

Gln Asp Gly Asp Ser Met Ala Thr Arg Glu Glu Leu Thr Ala Phe Leu  
           180                  185                  190

His Pro Glu Glu Phe Pro His Met Arg Asp Ile Val Ile Ala Glu Thr  
           195                  200                  205

Leu Glu Asp Leu Asp Arg Asn Lys Asp Gly Tyr Val Gln Val Glu Glu  
   210                  215                  220

Tyr Ile Ala Asp Leu Tyr Ser Ala Glu Pro Gly Glu Glu Glu Pro Ala  
   225                  230                  235                  240

Trp Val Gln Thr Glu Arg Gln Gln Phe Arg Asp Phe Arg Asp Leu Asn  
           245                  250                  255

Lys Asp Gly His Leu Asp Gly Ser Glu Val Gly His Trp Val Leu Pro  
           260                  265                  270

Pro Ala Gln Asp Gln Pro Leu Val Glu Ala Asn His Leu Leu His Glu  
           275                  280                  285

Ser Asp Thr Asp Lys Asp Gly Arg Leu Ser Lys Ala Glu Ile Leu Gly  
   290                  295                  300

Asn Trp Asn Met Phe Val Gly Ser Gln Ala Thr Asn Tyr Gly Glu Asp  
   305                  310                  315                  320

Leu Thr Arg His His Asp Glu Leu  
           325

<210> SEQ ID NO 222  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
           oligonucleotide probe

<400> SEQUENCE: 222

cgcaggccct catggccagg

20

<210> SEQ ID NO 223  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
           oligonucleotide probe

<400> SEQUENCE: 223

gaaatcctgg gtaattgg

18

<210> SEQ ID NO 224  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 224

gtgcgcggtg ctcacagctc atc 23

<210> SEQ ID NO 225  
 <211> LENGTH: 44  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe

<400> SEQUENCE: 225

ccccctgag cgacgtccc ccatgatgac gccacgga actt 44

<210> SEQ ID NO 226  
 <211> LENGTH: 2403  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 226

ggggccttgc cttccgcact cgggcgcagc cgggtggatc tcgagcaggt gcgagagccc 60  
 gggcgcgcggg cgcggtgagc agggatccct gacgcctctg tccctgttcc tttgtcgtcc 120  
 ccagcctgtc tgtcgtcgtt ttggcgcccc cgcctccccg cgggtcgggg ttgcaacccg 180  
 atcctgggct tcgctcgatt tgccgccgag gcgcctccca gacctagagg ggcgctggcc 240  
 tggagcagcg ggtcgtctgt gtccctctctc ctctgcgccg cgcgccggga tccgaagggg 300  
 gcggggctct gaggaggtag cgcgcggggc ctcccgcacc ctggccttgc ccgattctc 360  
 cctctctccc aggtgtgagc agcctatcag tcaccatgtc cgcagcctgg atcccggctc 420  
 tcggcctcgg tgtgtgtctg ctgctgctgc cggggcccgc gggcagcgag ggagccgctc 480  
 ccattgctat cacatgtttt accagaggct tggacatcag gaaagagaaa gcagatgtcc 540  
 tctgcccagg gggctgccct cttgaggaat tctctgtgta tgggaacata gtatatgctt 600  
 ctgtatcgag catatgtggg gctgctgtcc acaggggagt aatcagcaac tcagggggac 660  
 ctgtacgagt ctatagccta cctggtcgag aaaactattc ctcagtagat gccaatggca 720  
 tccagtctcaaatgctttct agatggtctg cttctttcac agtaactaaa ggcaaaagta 780  
 gtacacagga gggcacagga caagcagtggt ccacagcaca tccaccaaca ggtaaacgac 840  
 taaagaaaac acccgagaag aaaactggca ataaagattg taaagcagac attgcatttc 900  
 tgattgatgg aagctttaat attgggcagc gccgatttaa ttacagaag aattttgttg 960  
 gaaaagtggc tctaattgtt ggaattggaa cagaaggacc acatgtgggc cttgttcaag 1020  
 ccagtgaaca tcccaaaata gaattttact tgaaaaactt tacatcagcc aaagatgttt 1080  
 tgtttgccat aaaggaagta ggtttcagag ggggtaattc caatacagga aaagccttga 1140  
 agcactactgc tcagaaattc ttcacggtag atgctggagt aagaaaagg atccccaaag 1200  
 tgggtggtgtt atttattgat ggttgcctt ctgatgacat cgaggaagca ggcattgtgg 1260  
 ccagagagtt tgggtgcaat gtatttatag tttctgtggc caagcctatc cctgaagaac 1320  
 tgggatggtt tcaggatgtc acatttgttg acaaggctgt ctgtcggaat aatggottct 1380  
 tctcttacca catgcccac tggtttggca ccacaaaata cgtaaagcct ctggtacaga 1440

-continued

---

```

agctgtgcac tcatgaacaa atgatgtgca gcaagacctg ttataactca gtgaacattg 1500
cctttctaata tgatggctcc agcagtggtg gagatagcaa tttccgctc atgcttgaat 1560
ttgtttccaa catagccaag acttttgaaa tctcggacat tggtgccaag atagctgctg 1620
tacagtttac ttatgatcag cgcacggagt tcagtttcac tgactatagc accaaagaga 1680
atgtcctagc tgtcatcaga aacatccgct atatgagtgg tggaacagct actggtgatg 1740
ccatttcott cactggtaga aatgtgtttg gccctataag ggagagcccc aacaagaact 1800
tcctagtaat tgtcacagat gggcagctct atgatgatgt ccaaggccct gcagctgctg 1860
cacatgatgc aggaatcact atcttctctg ttggtgtggc ttgggcacct ctggatgacc 1920
tgaaagatat ggcttctaaa ccgaaggagt ctacgcgttt cttcacaaga gagttcacag 1980
gattagaacc aattgtttct gatgtcatca gaggcatttg tagagatttc ttagaatccc 2040
agcaataatg gtaacatctt gacaactgaa agaaaaagta caaggggatc cagtgtgtaa 2100
attgtattct cataaactg aaatgcttta gcatactaga atcagataca aaactattaa 2160
gtatgtcaac agccatttag gaaaataagc actcctttaa agccgctgcc ttctggttac 2220
aatttacagt gtactttgtt aaaaacactg ctgaggcttc ataatcatgg ctcttagaaa 2280
ctcaggaaaag aggagataat gtggattaaa accttaagag ttctaacat gcctactaaa 2340
tgtacagata tgcaaatcc atagctcaat aaaagaatct gatacttaga ccaaaaaaaaa 2400
aaa 2403

```

&lt;210&gt; SEQ ID NO 227

&lt;211&gt; LENGTH: 550

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 227

```

Met Ser Ala Ala Trp Ile Pro Ala Leu Gly Leu Gly Val Cys Leu Leu
 1           5           10          15
Leu Leu Pro Gly Pro Ala Gly Ser Glu Gly Ala Ala Pro Ile Ala Ile
 20          25          30
Thr Cys Phe Thr Arg Gly Leu Asp Ile Arg Lys Glu Lys Ala Asp Val
 35          40          45
Leu Cys Pro Gly Gly Cys Pro Leu Glu Glu Phe Ser Val Tyr Gly Asn
 50          55          60
Ile Val Tyr Ala Ser Val Ser Ser Ile Cys Gly Ala Ala Val His Arg
 65          70          75          80
Gly Val Ile Ser Asn Ser Gly Gly Pro Val Arg Val Tyr Ser Leu Pro
 85          90          95
Gly Arg Glu Asn Tyr Ser Ser Val Asp Ala Asn Gly Ile Gln Ser Gln
100         105         110
Met Leu Ser Arg Trp Ser Ala Ser Phe Thr Val Thr Lys Gly Lys Ser
115         120         125
Ser Thr Gln Glu Ala Thr Gly Gln Ala Val Ser Thr Ala His Pro Pro
130         135         140
Thr Gly Lys Arg Leu Lys Lys Thr Pro Glu Lys Lys Thr Gly Asn Lys
145         150         155         160
Asp Cys Lys Ala Asp Ile Ala Phe Leu Ile Asp Gly Ser Phe Asn Ile
165         170         175

```

-continued

Gly	Gln	Arg	Arg	Phe	Asn	Leu	Gln	Lys	Asn	Phe	Val	Gly	Lys	Val	Ala
			180					185					190		
Leu	Met	Leu	Gly	Ile	Gly	Thr	Glu	Gly	Pro	His	Val	Gly	Leu	Val	Gln
		195					200					205			
Ala	Ser	Glu	His	Pro	Lys	Ile	Glu	Phe	Tyr	Leu	Lys	Asn	Phe	Thr	Ser
	210				215						220				
Ala	Lys	Asp	Val	Leu	Phe	Ala	Ile	Lys	Glu	Val	Gly	Phe	Arg	Gly	Gly
225					230					235					240
Asn	Ser	Asn	Thr	Gly	Lys	Ala	Leu	Lys	His	Thr	Ala	Gln	Lys	Phe	Phe
				245					250					255	
Thr	Val	Asp	Ala	Gly	Val	Arg	Lys	Gly	Ile	Pro	Lys	Val	Val	Val	Val
			260					265					270		
Phe	Ile	Asp	Gly	Trp	Pro	Ser	Asp	Asp	Ile	Glu	Glu	Ala	Gly	Ile	Val
		275					280					285			
Ala	Arg	Glu	Phe	Gly	Val	Asn	Val	Phe	Ile	Val	Ser	Val	Ala	Lys	Pro
	290					295						300			
Ile	Pro	Glu	Glu	Leu	Gly	Met	Val	Gln	Asp	Val	Thr	Phe	Val	Asp	Lys
305					310					315					320
Ala	Val	Cys	Arg	Asn	Asn	Gly	Phe	Phe	Ser	Tyr	His	Met	Pro	Asn	Trp
				325					330					335	
Phe	Gly	Thr	Thr	Lys	Tyr	Val	Lys	Pro	Leu	Val	Gln	Lys	Leu	Cys	Thr
			340					345					350		
His	Glu	Gln	Met	Met	Cys	Ser	Lys	Thr	Cys	Tyr	Asn	Ser	Val	Asn	Ile
		355					360					365			
Ala	Phe	Leu	Ile	Asp	Gly	Ser	Ser	Ser	Val	Gly	Asp	Ser	Asn	Phe	Arg
	370					375					380				
Leu	Met	Leu	Glu	Phe	Val	Ser	Asn	Ile	Ala	Lys	Thr	Phe	Glu	Ile	Ser
385					390					395					400
Asp	Ile	Gly	Ala	Lys	Ile	Ala	Ala	Val	Gln	Phe	Thr	Tyr	Asp	Gln	Arg
				405					410					415	
Thr	Glu	Phe	Ser	Phe	Thr	Asp	Tyr	Ser	Thr	Lys	Glu	Asn	Val	Leu	Ala
			420					425					430		
Val	Ile	Arg	Asn	Ile	Arg	Tyr	Met	Ser	Gly	Gly	Thr	Ala	Thr	Gly	Asp
		435					440					445			
Ala	Ile	Ser	Phe	Thr	Val	Arg	Asn	Val	Phe	Gly	Pro	Ile	Arg	Glu	Ser
	450					455					460				
Pro	Asn	Lys	Asn	Phe	Leu	Val	Ile	Val	Thr	Asp	Gly	Gln	Ser	Tyr	Asp
465					470					475					480
Asp	Val	Gln	Gly	Pro	Ala	Ala	Ala	Ala	His	Asp	Ala	Gly	Ile	Thr	Ile
				485					490					495	
Phe	Ser	Val	Gly	Val	Ala	Trp	Ala	Pro	Leu	Asp	Asp	Leu	Lys	Asp	Met
			500					505					510		
Ala	Ser	Lys	Pro	Lys	Glu	Ser	His	Ala	Phe	Phe	Thr	Arg	Glu	Phe	Thr
		515					520					525			
Gly	Leu	Glu	Pro	Ile	Val	Ser	Asp	Val	Ile	Arg	Gly	Ile	Cys	Arg	Asp
	530					535					540				
Phe	Leu	Glu	Ser	Gln	Gln										
545					550										

&lt;210&gt; SEQ ID NO 228

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: DNA

---

-continued

---

<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 228

tggtctcgca caccgatc 18

<210> SEQ ID NO 229  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 229

ctgctgtcca caggggag 18

<210> SEQ ID NO 230  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 230

ccttgaagca tactgctc 18

<210> SEQ ID NO 231  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 231

gagatagcaa tttccgcc 18

<210> SEQ ID NO 232  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 232

ttcctcaaga gggcagcc 18

<210> SEQ ID NO 233  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
oligonucleotide probe

<400> SEQUENCE: 233

cttggcacca atgtccgaga ttcc 24

<210> SEQ ID NO 234



-continued

---

<211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 oligonucleotide probe  
  
 <400> SEQUENCE: 234  
  
 gctctgagga aggtgacgcg cggggcctcc gaacccttgg ccttg 45

<210> SEQ ID NO 235  
 <211> LENGTH: 2586  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 235  
  
 cgccgcgctc ccgcaccgcg ggcccgccca ccgcgcccgt cccgcactctg caccgcgagc 60  
 ccggcggcct cccggcggga gcgagcagat ccagtcggc ccgcagcgca actcgggtcca 120  
 gtcggggcgg cggctgcggg cgcagagcgg agatgcagcg gcttggggcc accctgctgt 180  
 gcctgctgct ggcgggcgcg gtcccacgg ccccgcgcc cgtccgacg gcgacctcgg 240  
 ctccagtc aa cccggcccc gctctcagct acccgcagga ggaggccacc ctcaatgaga 300  
 tgttccgcga ggttgaggaa ctgatggagg acacgcagca caaattgcgc agcgcggtgg 360  
 aagagatgga ggcagaagaa gctgctgcta aagcatcatc agaagtgaac ctggcaaaact 420  
 tacctcccag ctatcacaat gagaccaaca cagacacgaa ggttgaaat aataccatcc 480  
 atgtgcaccg aaaaattcac aagataacca acaaccagac tggacaaatg gctctttcag 540  
 agacagttat cacactctgtg ggagacgaag aaggcagaag gagccacgag tgcacatcag 600  
 acgaggactg tggggccagc atgtactgcc agtttgccag cttccagtac acctgccagc 660  
 catgccgggg ccagaggatg ctctgcaccc gggacagtga gtgctgtgga gaccagctgt 720  
 gtgtctgggg tcaactgcacc aaaatggcca ccaggggag caatgggacc atctgtgaca 780  
 accagaggga ctgccagccg gggctgtgct gtgccttcca gagaggcctg ctgttccctg 840  
 tgtgcacacc cctgcccgtg gagggcagc tttgcatga ccccgccagc cggcttctgg 900  
 acctcatcac ctgggagcta gagcctgatg gagccttggga ccgatgccct tgtgccagtg 960  
 gcctcctctg ccagccccac agccacagcc tgggtgatgt gtgcaagccg accttcgtgg 1020  
 ggagccgtga ccaagatggg gagatcctgc tgcccagaga ggtcccgat gagtatgaag 1080  
 ttggcagctt catggaggag gtgcgccagg agctggagga cctggagagg agcctgactg 1140  
 aagagatggc gctgggggag cctgcggctg ccgcccgtgc actgctggga ggggaagaga 1200  
 tttagatctg gaccaggctg tgggtagatg tgcaatagaa atagctaatt tatttccca 1260  
 ggtgtgtgct ttagcgtgg gctgaccagg cttcttccta catcttcttc ccagtaagtt 1320  
 tcccctctg cttgacagca tgagggttg tgcatttgtt cagctcccc aggctgttct 1380  
 ccagcttca cagtctgggt cttgggagag tcaggcaggg ttaaactgca ggagcagttt 1440  
 gccaccctg tccagattat tggctgcttt gcctctacca gttggcagac agccgtttgt 1500  
 tctacatggc tttgataaatt gtttgagggg aggagatgga aacaatgtgg agtctocctc 1560  
 tgattgtttt tgggaaatg tggagaagag tgccctgctt tgcaaacatc aacctggcaa 1620  
 aaatgcaaca aatgaatttt ccacgcagtt ctttccatgg gcataggtaa gctgtgcctt 1680  
 cagctgttgc agatgaaatg ttctgttcc cctgcattac atgtgtttat tcatccagca 1740

-continued

```

gtgttgctca gctcctacct ctgtgccagg gcagcatttt catatccaag atcaattccc 1800
tctctcagca cagcctgggg aggggggtcat tgttctcctc gtccatcagg gatctcagag 1860
gctcagagac tgcaagctgc ttgcccaagt cacacagcta gtgaagacca gagcagtttc 1920
atctggttgt gactctaagc tcagtgtctt ctccactacc ccacaccagc cttggtgcca 1980
ccaaaagtgc tccccaaaag gaaggagaat gggatttttc ttgaggcatg cacatctgga 2040
attaagtgca aactaattct cacatccctc taaaagtaaa ctactgttag gaacagcagt 2100
gttctcagag tgtggggcag ccgtccttct aatgaagaca atgatattga cactgtccct 2160
ctttggcagt tgcattagta actttgaaag gtatatgact gagcgtagca tacaggtaa 2220
cctgcagaaa cagtacttag gtaattgtag ggcgaggatt ataatgaaa ttgcaaaat 2280
cacttagcag caactgaaga caattatcaa ccacgtggag aaaatcaaac cgagcagggc 2340
tgtgtgaaac atggttgtaa tatgcgacty cgaacactga actctacgcc actccacaaa 2400
tgatgttttc aggtgtcatg gactgttgcc accatgtatt catccagagt tcttaaagtt 2460
taaagttgca catgattgta taagcatgct ttctttgagt tttaaattat gtataaacat 2520
aagttgcatt tagaaatcaa gcataaatca cttcaactgc aaaaaaaaaa aaaaaaaaaa 2580
aaaaaa 2586
    
```

```

<210> SEQ ID NO 236
<211> LENGTH: 350
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
    
```

<400> SEQUENCE: 236

```

Met  Gln  Arg  Leu  Gly  Ala  Thr  Leu  Leu  Cys  Leu  Leu  Leu  Ala  Ala  Ala
  1          5          10          15
Val  Pro  Thr  Ala  Pro  Ala  Pro  Ala  Pro  Thr  Ala  Thr  Ser  Ala  Pro  Val
          20          25          30
Lys  Pro  Gly  Pro  Ala  Leu  Ser  Tyr  Pro  Gln  Glu  Glu  Ala  Thr  Leu  Asn
          35          40          45
Glu  Met  Phe  Arg  Glu  Val  Glu  Glu  Leu  Met  Glu  Asp  Thr  Gln  His  Lys
          50          55          60
Leu  Arg  Ser  Ala  Val  Glu  Glu  Met  Glu  Ala  Glu  Glu  Ala  Ala  Ala  Lys
          65          70          75          80
Ala  Ser  Ser  Glu  Val  Asn  Leu  Ala  Asn  Leu  Pro  Pro  Ser  Tyr  His  Asn
          85          90          95
Glu  Thr  Asn  Thr  Asp  Thr  Lys  Val  Gly  Asn  Asn  Thr  Ile  His  Val  His
          100          105          110
Arg  Glu  Ile  His  Lys  Ile  Thr  Asn  Asn  Gln  Thr  Gly  Gln  Met  Val  Phe
          115          120          125
Ser  Glu  Thr  Val  Ile  Thr  Ser  Val  Gly  Asp  Glu  Glu  Gly  Arg  Arg  Ser
          130          135          140
His  Glu  Cys  Ile  Ile  Asp  Glu  Asp  Cys  Gly  Pro  Ser  Met  Tyr  Cys  Gln
          145          150          155          160
Phe  Ala  Ser  Phe  Gln  Tyr  Thr  Cys  Gln  Pro  Cys  Arg  Gly  Gln  Arg  Met
          165          170          175
Leu  Cys  Thr  Arg  Asp  Ser  Glu  Cys  Cys  Gly  Asp  Gln  Leu  Cys  Val  Trp
          180          185          190
Gly  His  Cys  Thr  Lys  Met  Ala  Thr  Arg  Gly  Ser  Asn  Gly  Thr  Ile  Cys
    
```

-continued

195		200				205									
Asp	Asn	Gln	Arg	Asp	Cys	Gln	Pro	Gly	Leu	Cys	Cys	Ala	Phe	Gln	Arg
	210					215					220				
Gly	Leu	Leu	Phe	Pro	Val	Cys	Thr	Pro	Leu	Pro	Val	Glu	Gly	Glu	Leu
225					230					235					240
Cys	His	Asp	Pro	Ala	Ser	Arg	Leu	Leu	Asp	Leu	Ile	Thr	Trp	Glu	Leu
				245					250					255	
Glu	Pro	Asp	Gly	Ala	Leu	Asp	Arg	Cys	Pro	Cys	Ala	Ser	Gly	Leu	Leu
			260					265					270		
Cys	Gln	Pro	His	Ser	His	Ser	Leu	Val	Tyr	Val	Cys	Lys	Pro	Thr	Phe
	275						280					285			
Val	Gly	Ser	Arg	Asp	Gln	Asp	Gly	Glu	Ile	Leu	Leu	Pro	Arg	Glu	Val
	290					295					300				
Pro	Asp	Glu	Tyr	Glu	Val	Gly	Ser	Phe	Met	Glu	Glu	Val	Arg	Gln	Glu
305					310					315					320
Leu	Glu	Asp	Leu	Glu	Arg	Ser	Leu	Thr	Glu	Glu	Met	Ala	Leu	Gly	Glu
				325					330					335	
Pro	Ala	Ala	Ala	Ala	Ala	Ala	Leu	Leu	Gly	Gly	Glu	Glu	Ile		
		340						345					350		
<p>&lt;210&gt; SEQ ID NO 237                  &lt;211&gt; LENGTH: 17                  &lt;212&gt; TYPE: DNA                  &lt;213&gt; ORGANISM: Artificial Sequence                  &lt;220&gt; FEATURE:                  &lt;223&gt; OTHER INFORMATION: Synthetic oligonucleotide probe</p>															
<p>&lt;400&gt; SEQUENCE: 237</p>															
ggagctgcac cccttgc														17	
<p>&lt;210&gt; SEQ ID NO 238                  &lt;211&gt; LENGTH: 49                  &lt;212&gt; TYPE: DNA                  &lt;213&gt; ORGANISM: Artificial Sequence                  &lt;220&gt; FEATURE:                  &lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe</p>															
<p>&lt;400&gt; SEQUENCE: 238</p>															
ggaggactgt gccacatga gagactcttc aaaccaagg caaaattgg														49	
<p>&lt;210&gt; SEQ ID NO 239                  &lt;211&gt; LENGTH: 24                  &lt;212&gt; TYPE: DNA                  &lt;213&gt; ORGANISM: Artificial Sequence                  &lt;220&gt; FEATURE:                  &lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe</p>															
<p>&lt;400&gt; SEQUENCE: 239</p>															
gcagagcggg gatgcagcgg cttg														24	
<p>&lt;210&gt; SEQ ID NO 240                  &lt;211&gt; LENGTH: 18                  &lt;212&gt; TYPE: DNA                  &lt;213&gt; ORGANISM: Artificial Sequence                  &lt;220&gt; FEATURE:                  &lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe</p>															
<p>&lt;400&gt; SEQUENCE: 240</p>															
ttggcagctt catggagg														18	

-continued

---

```

<210> SEQ ID NO 241
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 241

cctgggcaaa aatgcaac 18

<210> SEQ ID NO 242
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 242

ctccagctcc tggcgcacct cctc 24

<210> SEQ ID NO 243
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 243

ggctctcagc taccgcgcag gagcgaggcc accctcaatg agatg 45

<210> SEQ ID NO 244
<211> LENGTH: 3679
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 244

aaggaggctg ggaggaaaga ggtaagaaag gttagagaac ctacctaca 50
tctctctggg ctcagaagga ctctgaagat aacaataatt tcagcccatc 100
cactctcctt ccctcccaaa cacacatgtg catgtacaca cacacataca 150
cacacataca cttcctcttc cttcactgaa gactcacagt cactcactct 200
gtgagcaggt catagaaaag gacactaaag ccttaaggac aggccctggc 250
attacctctg cagctccttt ggcttgttga gtcaaaaaac atgggagggg 300
ccaggcacgg tgactcacac ctgtaatccc agcatthttg gagaccgagg 350
tgagcagatc acttgaggtc aggagttcga gaccagcctg gccaacatgg 400
agaaaccccc atctctacta aaaatacaaa aattagccag gagtgggtggc 450
agggtcctgt aatcccagct actcaggtgg ctgagccagg agaatcgctt 500
gaatccagga ggcggaggat gcagtcagct gagtgaccg ctgcactcca 550
gcctgggtga cagaatgaga ctctgtctca aacaacaaa cacgggagga 600
ggggtagata ctgcttctct gcaacctcct taactctgca tcctcttctt 650
ccagggtctg ccctgatggg gcctggcaat gactgagcag gccagcccc 700
agaggacaag gaagagaagg catattgagg agggcaagaa gtgacgcccg 750
gtgtagaatg actgccctgg gaggggtggt ccttgggccc tggcagggtt 800

```

-continued

---

gctgaccctt	acctgcaaa	acacaaagag	caggactcca	gactctcctt	850
gtgaatggtc	ccctgccctg	cagctccacc	atgaggcttc	tcgtggcccc	900
actcttgeta	gcttgggtgg	ctggtgccac	tgccactgtg	cccgtggtag	950
cctggcatgt	tcctgcccc	cctcagtgtg	cctgccagat	cgggcccc	1000
tatacgcctc	gctcgtccta	ccgcgaggct	accactgtgg	actgcaatga	1050
cctattcctg	acggcagtcc	ccccggcact	ccccgcaggc	acacagacc	1100
tgctcctgca	gagcaacagc	attgtccgtg	tggaccagag	tgagctgggc	1150
tacctggcca	atctcacaga	gctggacctg	tcccagaaca	gcttttcgga	1200
tgcccagagc	tgtgatttcc	atgccctgcc	ccagctgctg	agcctgcacc	1250
tagaggagaa	ccagctgacc	cggctggagg	accacagctt	tgacgggctg	1300
gccagcctac	aggaactcta	tctcaaccac	aaccagctct	accgcatcg	1350
ccccagggcc	tttctcggcc	tcagcaactt	gctgcggctg	cacctcaact	1400
caaacctcct	gagggccatt	gacagccgct	ggtttgaat	gctgccaac	1450
ttggagatac	tcatgattgg	cggaacaag	gtagatgcca	tcctggacat	1500
gaacttccgg	cccctggcca	acctgcgtag	cctggtgcta	gcaggcatga	1550
acctgcggga	gatctccgac	tatgccttgg	aggggctgca	aagcctggag	1600
agcctctcct	tctatgacaa	ccagctggcc	cgggtgcccc	ggcgggact	1650
ggaacaggtg	cccgggctca	agttcctaga	cctcaacaag	aaccgctcc	1700
agcgggtagg	gcccggggac	tttgccaaca	tgctgcacct	taaggagctg	1750
ggactgaaca	acatggagga	gctggtctcc	atcgacaagt	ttgocctgg	1800
gaaacctccc	gagctgacca	agctggacat	caccaataac	ccacggctgt	1850
ccttcaccca	cccccgcgcc	ttccaccacc	tgccccagat	ggagaccctc	1900
atgtcaca	acaacgctct	cagtgccttg	caccagcaga	cggtgaggct	1950
cctgccaac	ctgcaggagg	taggtctcca	cggcaacccc	atccgctgtg	2000
actgtgtcat	ccgctgggcc	aatgccacgg	gcacccgtgt	ccgcttcac	2050
gagccgcaat	ccacctgtg	tgccgagcct	ccggacctcc	agcgcctccc	2100
ggtccgtgag	gtgcccttcc	gggagatgac	ggaccactgt	ttgccctcca	2150
tctccccaag	aagcttcccc	ccaagcctcc	aggtagccag	tgagagagac	2200
atggtgctgc	attgccgggc	actggccgaa	cccgaacccg	agatctactg	2250
ggtcactcca	gctgggcttc	gactgacacc	tgcccatgca	ggcaggaggt	2300
accgggtgta	ccccgagggg	accctggagc	tgccgagggt	gacagcagaa	2350
gaggcagggc	tatacacctg	tgtggcccag	aacctggtgg	gggctgacac	2400
taagacggtt	agtgtggttg	tgggcccgtc	tctcctccag	ccaggcaggg	2450
acgaaggaca	ggggtggag	ctccgggtgc	aggagacca	cccctatcac	2500
atcctgctat	cttgggtcac	cccacccaac	acagtgtcca	ccaacctcac	2550
ctgttccagt	gcctcctccc	tcgggggcca	gggggcccaca	gctctggccc	2600
gctgcctog	gggaacccac	agctacaaca	ttaccgcct	ccttcaggcc	2650
acggagtact	ggcctgcct	gcaagtggcc	tttgetgatg	cccacacca	2700

-continued

---

```

gttgcttgt gtatggcca ggaccaaaga ggccacttct tgccacagag      2750
ccttagggga tcgtcctggg ctcatcgcca tcttgctct cgctgtcctt      2800
ctcctggcag ctgggctagc ggcccacctt ggcacaggcc aaccagaa      2850
gggtgtgggt gggaggcggc ctctcctcc agcctgggct ttctggggct      2900
ggagtgcctt ttctgtccgg gttgtgtctg ctcccctcgt cctgccctgg      2950
aatccaggga ggaagctgcc cagatcctca gaaggggaga cactgttgcc      3000
accattgtct caaaattctt gaagctcagc ctgttctcag cagtagagaa      3050
atcactagga ctacttttta ccaaagaga agcagtctgg gccagatgcc      3100
ctgccaggaa agggacatgg acccactgctc ttgaggcctg gcagctgggc      3150
caagacagat ggggctttgt ggcccgggg gtgcttctgc agccttgaaa      3200
aagtggcctt tacctcctag ggtcacctct gctgccattc tgaggaacat      3250
ctccaaggaa caggaggac tttggctaga gcctcctgcc tccccatctt      3300
ctctctgccc agaggctcct gggcctggct tggctgtccc ctacctgtgt      3350
ccccgggctg cacccttcc tcttctcttt ctctgtacag tctcagttgc      3400
ttgctcttgt gcctcctggg caaggctga aggaggccac tccatctcac      3450
ctcggggggc tgccctcaat gtgggagtga cccagccag atctgaagga      3500
catttgggag agggatgccc aggaacgctt catctcagca gcctgggctc      3550
ggcattccga agctgacttt ctataggcaa tttgtacct ttgtggagaa      3600
atgtgtcacc tcccccaacc cgattcactc tttctcctg ttttgtaaaa      3650
aataaaaata aataataaca ataaaaaaa      3679

```

&lt;210&gt; SEQ ID NO 245

&lt;211&gt; LENGTH: 713

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 245

```

Met Arg Leu Leu Val Ala Pro Leu Leu Leu Ala Trp Val Ala Gly
 1           5           10          15
Ala Thr Ala Thr Val Pro Val Val Pro Trp His Val Pro Cys Pro
 20          25          30
Pro Gln Cys Ala Cys Gln Ile Arg Pro Trp Tyr Thr Pro Arg Ser
 35          40          45
Ser Tyr Arg Glu Ala Thr Thr Val Asp Cys Asn Asp Leu Phe Leu
 50          55          60
Thr Ala Val Pro Pro Ala Leu Pro Ala Gly Thr Gln Thr Leu Leu
 65          70          75
Leu Gln Ser Asn Ser Ile Val Arg Val Asp Gln Ser Glu Leu Gly
 80          85          90
Tyr Leu Ala Asn Leu Thr Glu Leu Asp Leu Ser Gln Asn Ser Phe
 95          100         105
Ser Asp Ala Arg Asp Cys Asp Phe His Ala Leu Pro Gln Leu Leu
 110         115         120
Ser Leu His Leu Glu Glu Asn Gln Leu Thr Arg Leu Glu Asp His
 125         130         135

```

-continued

Ser	Phe	Ala	Gly	Leu	Ala	Ser	Leu	Gln	Glu	Leu	Tyr	Leu	Asn	His
				140					145					150
Asn	Gln	Leu	Tyr	Arg	Ile	Ala	Pro	Arg	Ala	Phe	Ser	Gly	Leu	Ser
				155					160					165
Asn	Leu	Leu	Arg	Leu	His	Leu	Asn	Ser	Asn	Leu	Leu	Arg	Ala	Ile
				170					175					180
Asp	Ser	Arg	Trp	Phe	Glu	Met	Leu	Pro	Asn	Leu	Glu	Ile	Leu	Met
				185					190					195
Ile	Gly	Gly	Asn	Lys	Val	Asp	Ala	Ile	Leu	Asp	Met	Asn	Phe	Arg
				200					205					210
Pro	Leu	Ala	Asn	Leu	Arg	Ser	Leu	Val	Leu	Ala	Gly	Met	Asn	Leu
				215					220					225
Arg	Glu	Ile	Ser	Asp	Tyr	Ala	Leu	Glu	Gly	Leu	Gln	Ser	Leu	Glu
				230					235					240
Ser	Leu	Ser	Phe	Tyr	Asp	Asn	Gln	Leu	Ala	Arg	Val	Pro	Arg	Arg
				245					250					255
Ala	Leu	Glu	Gln	Val	Pro	Gly	Leu	Lys	Phe	Leu	Asp	Leu	Asn	Lys
				260					265					270
Asn	Pro	Leu	Gln	Arg	Val	Gly	Pro	Gly	Asp	Phe	Ala	Asn	Met	Leu
				275					280					285
His	Leu	Lys	Glu	Leu	Gly	Leu	Asn	Asn	Met	Glu	Glu	Leu	Val	Ser
				290					295					300
Ile	Asp	Lys	Phe	Ala	Leu	Val	Asn	Leu	Pro	Glu	Leu	Thr	Lys	Leu
				305					310					315
Asp	Ile	Thr	Asn	Asn	Pro	Arg	Leu	Ser	Phe	Ile	His	Pro	Arg	Ala
				320					325					330
Phe	His	His	Leu	Pro	Gln	Met	Glu	Thr	Leu	Met	Leu	Asn	Asn	Asn
				335					340					345
Ala	Leu	Ser	Ala	Leu	His	Gln	Gln	Thr	Val	Glu	Ser	Leu	Pro	Asn
				350					355					360
Leu	Gln	Glu	Val	Gly	Leu	His	Gly	Asn	Pro	Ile	Arg	Cys	Asp	Cys
				365					370					375
Val	Ile	Arg	Trp	Ala	Asn	Ala	Thr	Gly	Thr	Arg	Val	Arg	Phe	Ile
				380					385					390
Glu	Pro	Gln	Ser	Thr	Leu	Cys	Ala	Glu	Pro	Pro	Asp	Leu	Gln	Arg
				395					400					405
Leu	Pro	Val	Arg	Glu	Val	Pro	Phe	Arg	Glu	Met	Thr	Asp	His	Cys
				410					415					420
Leu	Pro	Leu	Ile	Ser	Pro	Arg	Ser	Phe	Pro	Pro	Ser	Leu	Gln	Val
				425					430					435
Ala	Ser	Gly	Glu	Ser	Met	Val	Leu	His	Cys	Arg	Ala	Leu	Ala	Glu
				440					445					450
Pro	Glu	Pro	Glu	Ile	Tyr	Trp	Val	Thr	Pro	Ala	Gly	Leu	Arg	Leu
				455					460					465
Thr	Pro	Ala	His	Ala	Gly	Arg	Arg	Tyr	Arg	Val	Tyr	Pro	Glu	Gly
				470					475					480
Thr	Leu	Glu	Leu	Arg	Arg	Val	Thr	Ala	Glu	Glu	Ala	Gly	Leu	Tyr
				485					490					495
Thr	Cys	Val	Ala	Gln	Asn	Leu	Val	Gly	Ala	Asp	Thr	Lys	Thr	Val
				500					505					510
Ser	Val	Val	Val	Gly	Arg	Ala	Leu	Leu	Gln	Pro	Gly	Arg	Asp	Glu





---

-continued

---

<210> SEQ ID NO 249

<211> LENGTH: 3401

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 249

gcaagccaag gcgctgtttg agaaggtgaa gaagttccgg acccatgtgg	50
aggaggggga cattgtgtac cgctctaca tgcggcagac catcatcaag	100
gtgatcaagt tcatcctcat catctgctac accgtctact acgtgcacaa	150
catcaagttc gacgtggact gcaccgtgga cattgagagc ctgacgggct	200
accgcaccta ccgctgtgcc caccctctgg ccacactctt caagatcctg	250
gcgtccttct acatcagcct agtcatcttc tacggcctca tctgcatgta	300
cacactgtgg tggatgctac ggcgtccct caagaagtac tcgtttgagt	350
cgatccgtga ggagagcagc tacagcgaca tcccgcagct caagaacgac	400
ttcgcttca tgctgcacct cattgaccaa tacgaccgc tctactocaa	450
gcgcttcgoc gtcttctgt cggaggtgag tgagaacaag ctgoggcagc	500
tgaacctcaa caacgagtgg acgctggaca agtccggca gcggctcacc	550
aagaacgcgc aggacaagct ggagctgcac ctgttcatgc tcagtggcat	600
ccctgacaat gtgtttgacc tgggtggagct ggaggtcctc aagctggagc	650
tgatccccga cgtgaccatc ccgccagca ttgccagct cacgggctc	700
aaggagctgt ggctctacca cacagcggcc aagattgaag cgctcgct	750
ggccttctcg cgcgagaacc tgcgggcgct gcacatcaag ttcaccgaca	800
tcaaggagat cccgctgtgg atctatagcc tgaagacact ggaggagctg	850
cacctgacgg gcaacctgag cgcggagaac aaccgctaca tcgtcatcga	900
cgggctgcgg gagctcaaac gcctcaaggt gctgcggctc aagagcaacc	950
taagcaagct gccacaggty gtcacagatg tgggctgca cctgcagaag	1000
ctgtccatca acaatgaggg caccaagctc atcgtcctca acagcctcaa	1050
gaagatggcg aacctgactg agctggagct gatccgctgc gacctggagc	1100
gcatccccca ctccatcttc agcctccaca acctgcagga gattgacctc	1150
aaggacaaca acctcaagac catcgaggag atcatcagct tccagcacct	1200
gcaccgcctc acctgcctta agctgtggta caaccacatc gcctacatcc	1250
ccatccagat cggcaacctc accaacctgg agcgcctcta cctgaaccgc	1300
aacaagatcg agaagatccc caccagctc ttctactgcc gcaagctgcg	1350
ctacctggag ctcagccaca acaacctgac cttctcctc gccgacatcg	1400
gcctcctgca gaacctccag aacctagcca tcacggccaa ccggatcgag	1450
acgctccctc cggagctctt ccagtgcggg aagctgcggg cctgcacct	1500
gggcaacaac gtgctgcagt cactgcctc cagggtgggc gagctgacca	1550
acctgacgca gatcgagctg cggggcaacc ggctggagty cctgcctgtg	1600
gagctgggog agtgccact gctcaagcgc agcggcttg tggtggagga	1650
ggacctgttc aacacactgc caccagaggt gaaggagcgg ctgtggaggg	1700

-continued

---

ctgacaagga gcaggcctga gcgaggccgg cccagcacag caagcagcag	1750
gaccgctgcc cagtcctcag gcccgagggg gcaggcctag cttctcccag	1800
aactcccgga cagccaggac agcctcggcg ctgggcagga gcctggggcc	1850
gcttgtagt caggccagag cgagaggaca gtatctgtgg ggctggcccc	1900
ttttctccct ctgagactca cgtcccccag ggcaagtgt tgtggaggag	1950
agcaagtctc aagagcgagc tatttgata atcagggtct cctccctgga	2000
ggccagctct gcccagggg ctgagctgcc accagaggtc ctgggaccct	2050
cactttagtt cttggtattt atttttctcc atctcccacc tccttcatcc	2100
agataactta tacattccca agaaagtta gccagatgg aagggttca	2150
gggaaagggt ggctgccttt tcccctgtc cttatttagc gatgccgccg	2200
ggcatttaac acccacctgg acttcagcag agtgggccgg ggcgaaccag	2250
ccatgggacg gtcaccacgc agtgccgggc tgggctctgc ggtgcggtcc	2300
acgggagagc aggcctccag ctggaaggc caggcctgga gcttgccctc	2350
tcagtttttg tggcagtttt agttttttgt tttttttttt tttaatcaaa	2400
aaacaatttt ttttaaaaa aagctttgaa aatggatggt ttgggtatta	2450
aaaagaaaa aaaaacttaa aaaaaaaaag aactaacgg ccagtgagtt	2500
ggagtctcag ggcaggggtg cagtttccct tgagcaaac agccagacgt	2550
tgaactgtgt ttcctttccc tgggcgcagg gtgcagggtg tcttccggat	2600
ctgggtgac cttgggtccag gagttctatt tgttctctgg gagggaggtt	2650
tttttgttt tttttgggt tttttgtgt tottgtttc tttctctcc	2700
atgtgtctg gcaggcactc atttctgtg ctgtcgcca gagggaatgt	2750
tctggagctg ccaaggaggg aggagactcg ggttgctaa tcccggatg	2800
aacggtgctc cattcgacc tcccctcctc gtgcctgcc tgcctctcca	2850
cgcacagtgt taaggagcca agaggagcca cttcgccag actttgtttc	2900
cccacctcct gcggcatggg tgtgtccagt gccaccgtg gcctccgtg	2950
cttccatcag ccctgtcggc acctggtcct tcatgaagag cagacactta	3000
gaggctggtc gggaaatggg aggtcggccc tgggagggca ggcgttggtt	3050
ccaagccggt tcccgtccct ggcgcctgga gtgcacacag cccagtcggc	3100
acctggtggc tggaaagcaa cctgctttag atcaactcgg tccccacctt	3150
agaagggtcc ccgccttaga tcaatcacgt ggacactaag gcacgtttta	3200
gagtctcttg tcttaatgat tatgtccatc cgtctgtccg tccatttgtg	3250
ttttctgcgt cgtgtcattg gatataatcc tcagaaataa tgcacactag	3300
cctctgacaa ccatgaagca aaaatccgtt acatgtgggt ctgaacttgt	3350
agactcggtc acagtatcaa ataaaatcta taacagaaaa aaaaaaaaaa	3400
a	3401

&lt;210&gt; SEQ ID NO 250

&lt;211&gt; LENGTH: 546

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 250

Met	Arg	Gln	Thr	Ile	Ile	Lys	Val	Ile	Lys	Phe	Ile	Leu	Ile	Ile
1				5					10					15
Cys	Tyr	Thr	Val	Tyr	Tyr	Val	His	Asn	Ile	Lys	Phe	Asp	Val	Asp
				20					25					30
Cys	Thr	Val	Asp	Ile	Glu	Ser	Leu	Thr	Gly	Tyr	Arg	Thr	Tyr	Arg
				35					40					45
Cys	Ala	His	Pro	Leu	Ala	Thr	Leu	Phe	Lys	Ile	Leu	Ala	Ser	Phe
				50					55					60
Tyr	Ile	Ser	Leu	Val	Ile	Phe	Tyr	Gly	Leu	Ile	Cys	Met	Tyr	Thr
				65					70					75
Leu	Trp	Trp	Met	Leu	Arg	Arg	Ser	Leu	Lys	Lys	Tyr	Ser	Phe	Glu
				80					85					90
Ser	Ile	Arg	Glu	Glu	Ser	Ser	Tyr	Ser	Asp	Ile	Pro	Asp	Val	Lys
				95					100					105
Asn	Asp	Phe	Ala	Phe	Met	Leu	His	Leu	Ile	Asp	Gln	Tyr	Asp	Pro
				110					115					120
Leu	Tyr	Ser	Lys	Arg	Phe	Ala	Val	Phe	Leu	Ser	Glu	Val	Ser	Glu
				125					130					135
Asn	Lys	Leu	Arg	Gln	Leu	Asn	Leu	Asn	Asn	Glu	Trp	Thr	Leu	Asp
				140					145					150
Lys	Leu	Arg	Gln	Arg	Leu	Thr	Lys	Asn	Ala	Gln	Asp	Lys	Leu	Glu
				155					160					165
Leu	His	Leu	Phe	Met	Leu	Ser	Gly	Ile	Pro	Asp	Thr	Val	Phe	Asp
				170					175					180
Leu	Val	Glu	Leu	Glu	Val	Leu	Lys	Leu	Glu	Leu	Ile	Pro	Asp	Val
				185					190					195
Thr	Ile	Pro	Pro	Ser	Ile	Ala	Gln	Leu	Thr	Gly	Leu	Lys	Glu	Leu
				200					205					210
Trp	Leu	Tyr	His	Thr	Ala	Ala	Lys	Ile	Glu	Ala	Pro	Ala	Leu	Ala
				215					220					225
Phe	Leu	Arg	Glu	Asn	Leu	Arg	Ala	Leu	His	Ile	Lys	Phe	Thr	Asp
				230					235					240
Ile	Lys	Glu	Ile	Pro	Leu	Trp	Ile	Tyr	Ser	Leu	Lys	Thr	Leu	Glu
				245					250					255
Glu	Leu	His	Leu	Thr	Gly	Asn	Leu	Ser	Ala	Glu	Asn	Asn	Arg	Tyr
				260					265					270
Ile	Val	Ile	Asp	Gly	Leu	Arg	Glu	Leu	Lys	Arg	Leu	Lys	Val	Leu
				275					280					285
Arg	Leu	Lys	Ser	Asn	Leu	Ser	Lys	Leu	Pro	Gln	Val	Val	Thr	Asp
				290					295					300
Val	Gly	Val	His	Leu	Gln	Lys	Leu	Ser	Ile	Asn	Asn	Glu	Gly	Thr
				305					310					315
Lys	Leu	Ile	Val	Leu	Asn	Ser	Leu	Lys	Lys	Met	Ala	Asn	Leu	Thr
				320					325					330
Glu	Leu	Glu	Leu	Ile	Arg	Cys	Asp	Leu	Glu	Arg	Ile	Pro	His	Ser
				335					340					345
Ile	Phe	Ser	Leu	His	Asn	Leu	Gln	Glu	Ile	Asp	Leu	Lys	Asp	Asn
				350					355					360
Asn	Leu	Lys	Thr	Ile	Glu	Glu	Ile	Ile	Ser	Phe	Gln	His	Leu	His

-continued

365	370	375
Arg Leu Thr Cys Leu Lys Leu Trp Tyr	Asn His Ile Ala Tyr Ile	
380	385	390
Pro Ile Gln Ile Gly Asn Leu Thr Asn	Leu Glu Arg Leu Tyr Leu	
395	400	405
Asn Arg Asn Lys Ile Glu Lys Ile Pro	Thr Gln Leu Phe Tyr Cys	
410	415	420
Arg Lys Leu Arg Tyr Leu Asp Leu Ser	His Asn Asn Leu Thr Phe	
425	430	435
Leu Pro Ala Asp Ile Gly Leu Leu Gln	Asn Leu Gln Asn Leu Ala	
440	445	450
Ile Thr Ala Asn Arg Ile Glu Thr Leu	Pro Pro Glu Leu Phe Gln	
455	460	465
Cys Arg Lys Leu Arg Ala Leu His Leu	Gly Asn Asn Val Leu Gln	
470	475	480
Ser Leu Pro Ser Arg Val Gly Glu Leu	Thr Asn Leu Thr Gln Ile	
485	490	495
Glu Leu Arg Gly Asn Arg Leu Glu Cys	Leu Pro Val Glu Leu Gly	
500	505	510
Glu Cys Pro Leu Leu Lys Arg Ser Gly	Leu Val Val Glu Glu Asp	
515	520	525
Leu Phe Asn Thr Leu Pro Pro Glu Val	Lys Glu Arg Leu Trp Arg	
530	535	540
Ala Asp Lys Glu Gln Ala		
545		

<210> SEQ ID NO 251  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 251

caacaatgag ggcaccaagc

20

<210> SEQ ID NO 252  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 252

gatggctagg ttctggaggt tctg

24

<210> SEQ ID NO 253  
 <211> LENGTH: 47  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 253

caacctgcag gagattgacc tcaaggacaa caacctcaag accatcg

47

<210> SEQ ID NO 254

---

-continued

---

<211> LENGTH: 1650

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 254

gcctgttgct gatgctgccg tgcggtactt gtcattggagc tggcaactgcg 50  
gcgctctccc gtcccgcggt ggttctctct gctgccgctg ctgctggggc 100  
tgaacgcagg agctgtcatt gactggccca cagaggaggg caaggaagta 150  
tgggattatg tgacggtccg caaggatgcc tacatgttct ggtggctcta 200  
ttatgccacc aactcctgca agaacttctc agaactgccc ctggtcatgt 250  
ggcttcaggg cggtcaggc ggttctagca ctggatttgg aaactttgag 300  
gaaattgggg cccttgacag tgatctcaaa ccacgaaaa ccactggct 350  
ccaggctgcc agtctcctat ttgtggataa tcccgtaggg actgggttca 400  
gttatgtgaa tggtagtggg gcctatgcca aggacctggc tatggtggct 450  
tcagacatga tggttctcct gaagaccttc ttcagttgcc acaagaatt 500  
ccagacagtt ccattctaca ttttctcaga gtcctatgga ggaaaaatg 550  
cagctggcat tggctctagag ctttataagg ccattcagcg agggaccatc 600  
aagtgcaact ttgcgggggt tgcttgggt gattcctgga tctcccctgt 650  
tgattcgggt ctctcctggg gacctacct gtacagcatg tctctctctg 700  
aagacaaaag tctggcagag gtgtctaagg ttgcagagca agtactgaat 750  
gccgtaaata aggggctcta cagagaggcc acagagctgt ggggaaagc 800  
agaaatgac attgaacaga acacagatgg ggtgaacttc tataacatct 850  
taactaaaag cactcccacg tctacaatgg agtcgagtct agaattcaca 900  
cagagccacc tagtttctct ttgtcagcgc cacgtgagac acctacaacg 950  
agatgcctta agccagctca tgaatggccc catcagaaaag aagctcaaaa 1000  
ttattcctga ggatcaatcc tggggaggcc aggtaccaa cgtctttgtg 1050  
aacatggagg aggacttcat gaagccatgc attagcattg tggacgagtt 1100  
gctggaggca gggatcaacg tgacggtgta taatggacag ctggatctca 1150  
tcgtagatac catgggtcag gaggcctggg tgcggaaact gaagtggcca 1200  
gaactgccta aattcagtca gctgaagtgg aaggccctgt acagtgacc 1250  
taaatctttg gaaacatctg cttttgtcaa gtcctacaag aaccttgctt 1300  
tctactggat tctgaaagct ggtcatatgg ttccttctga ccaaggggac 1350  
atggctctga agatgatgag actggtgact cagcaagaat aggatggatg 1400  
gggctggaga tgagctgggt tggccttggg gcacagagct gagctgaggc 1450  
cgctgaagct gtaggaagcg ccattcttcc ctgtatctaa ctggggctgt 1500  
gatcaagaag gttctgacca gttctgcag aggataaaat cattgtctct 1550  
ggaggcaatt tggaaattat ttctgcttct taaaaaac taagattttt 1600  
taaaaaattg atttgtttg atcaaaataa aggatgataa tagatattaa 1650

<210> SEQ ID NO 255

<211> LENGTH: 452

<212> TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 255

```

Met Glu Leu Ala Leu Arg Arg Ser Pro Val Pro Arg Trp Leu Leu
 1          5          10          15
Leu Leu Pro Leu Leu Leu Gly Leu Asn Ala Gly Ala Val Ile Asp
 20          25          30
Trp Pro Thr Glu Gly Lys Glu Val Trp Asp Tyr Val Thr Val
 35          40          45
Arg Lys Asp Ala Tyr Met Phe Trp Trp Leu Tyr Tyr Ala Thr Asn
 50          55          60
Ser Cys Lys Asn Phe Ser Glu Leu Pro Leu Val Met Trp Leu Gln
 65          70          75
Gly Gly Pro Gly Gly Ser Ser Thr Gly Phe Gly Asn Phe Glu Glu
 80          85          90
Ile Gly Pro Leu Asp Ser Asp Leu Lys Pro Arg Lys Thr Thr Trp
 95          100         105
Leu Gln Ala Ala Ser Leu Leu Phe Val Asp Asn Pro Val Gly Thr
 110         115         120
Gly Phe Ser Tyr Val Asn Gly Ser Gly Ala Tyr Ala Lys Asp Leu
 125         130         135
Ala Met Val Ala Ser Asp Met Met Val Leu Leu Lys Thr Phe Phe
 140         145         150
Ser Cys His Lys Glu Phe Gln Thr Val Pro Phe Tyr Ile Phe Ser
 155         160         165
Glu Ser Tyr Gly Gly Lys Met Ala Ala Gly Ile Gly Leu Glu Leu
 170         175         180
Tyr Lys Ala Ile Gln Arg Gly Thr Ile Lys Cys Asn Phe Ala Gly
 185         190         195
Val Ala Leu Gly Asp Ser Trp Ile Ser Pro Val Asp Ser Val Leu
 200         205         210
Ser Trp Gly Pro Tyr Leu Tyr Ser Met Ser Leu Leu Glu Asp Lys
 215         220         225
Gly Leu Ala Glu Val Ser Lys Val Ala Glu Gln Val Leu Asn Ala
 230         235         240
Val Asn Lys Gly Leu Tyr Arg Glu Ala Thr Glu Leu Trp Gly Lys
 245         250         255
Ala Glu Met Ile Ile Glu Gln Asn Thr Asp Gly Val Asn Phe Tyr
 260         265         270
Asn Ile Leu Thr Lys Ser Thr Pro Thr Ser Thr Met Glu Ser Ser
 275         280         285
Leu Glu Phe Thr Gln Ser His Leu Val Cys Leu Cys Gln Arg His
 290         295         300
Val Arg His Leu Gln Arg Asp Ala Leu Ser Gln Leu Met Asn Gly
 305         310         315
Pro Ile Arg Lys Lys Leu Lys Ile Ile Pro Glu Asp Gln Ser Trp
 320         325         330
Gly Gly Gln Ala Thr Asn Val Phe Val Asn Met Glu Glu Asp Phe
 335         340         345
Met Lys Pro Val Ile Ser Ile Val Asp Glu Leu Leu Glu Ala Gly
 350         355         360

```

-continued

---

Ile Asn Val Thr Val Tyr Asn Gly Gln Leu Asp Leu Ile Val Asp  
 365 370 375

Thr Met Gly Gln Glu Ala Trp Val Arg Lys Leu Lys Trp Pro Glu  
 380 385 390

Leu Pro Lys Phe Ser Gln Leu Lys Trp Lys Ala Leu Tyr Ser Asp  
 395 400 405

Pro Lys Ser Leu Glu Thr Ser Ala Phe Val Lys Ser Tyr Lys Asn  
 410 415 420

Leu Ala Phe Tyr Trp Ile Leu Lys Ala Gly His Met Val Pro Ser  
 425 430 435

Asp Gln Gly Asp Met Ala Leu Lys Met Met Arg Leu Val Thr Gln  
 440 445 450

Gln Glu

<210> SEQ ID NO 256  
 <211> LENGTH: 1100  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 256

```

ggcccgcgga gaggaggcca tgggcgcgcg cggggcgctg ctgctggcgc      50
tgctgctggc tcgggctgga ctcaggaagc cggagtcgca ggaggcggcg      100
ccgttatcag gaccatgcgg ccgacgggtc atcacgtcgc gcatcgtggg      150
tggagaggac gccgaactcg ggcgttggcc gtggcagggg agcctgcgcc      200
tgtgggattc ccacgtatgc ggagtgagcc tgctcagcca ccgotgggca      250
ctcacgcgcg cgcactgctt tgaaacctat agtgacctta gtgatccctc      300
cgggtggatg gtccagtttg gccagctgac ttccatgcca tccttctgga      350
gcctgcaggg ctactacacc cgttacttcg tatcgaatat ctatctgagc      400
cctcgctacc tggggaattc accctatgac attgccttgg tgaagctgtc      450
tgcaacctgtc acctacacta aacacatcca gcccatctgt ctccaggcct      500
ccacatttga gtttgagaac cggacagact gctgggtgac tggctggggg      550
tacatcaaaag aggatgaggc actgccatct ccccacacc tccaggaagt      600
tcaggtcgcc atcataaaca actctatgtg caaccacctc ttcctcaagt      650
acagtttccg caaggacatc tttggagaca tggtttgtgc tggcaacgcc      700
caaggcggga aggatgcctg cttcggtgac tcaggtggac ccttggcctg      750
taacaagaat ggactgtggt atcagattgg agtcgtgagc tggggagtgg      800
gctgtggtcg gcccaatcgg ccggtgtctc acaccaatat cagccaccac      850
tttgagtgga tccagaagct gatggcccag agtggcatgt cccagccaga      900
cccctcctgg ccaactactc ttttccctct tctctgggct ctcccactcc      950
tggggccggt ctgagcctac ctgagcccat gcagcctggg gccactgcca     1000
agtcaggccc tggttctctt ctgtcttgtt tggtaataaa cacattocag     1050
ttgatgcctt gcagggcatt cttaaaaaa aaaaaaaaaa aaaaaaaaaa     1100
    
```

<210> SEQ ID NO 257  
 <211> LENGTH: 314  
 <212> TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 257

```

Met Gly Ala Arg Gly Ala Leu Leu Leu Ala Leu Leu Leu Ala Arg
 1          5          10          15
Ala Gly Leu Arg Lys Pro Glu Ser Gln Glu Ala Ala Pro Leu Ser
 20          25          30
Gly Pro Cys Gly Arg Arg Val Ile Thr Ser Arg Ile Val Gly Gly
 35          40          45
Glu Asp Ala Glu Leu Gly Arg Trp Pro Trp Gln Gly Ser Leu Arg
 50          55          60
Leu Trp Asp Ser His Val Cys Gly Val Ser Leu Leu Ser His Arg
 65          70          75
Trp Ala Leu Thr Ala Ala His Cys Phe Glu Thr Tyr Ser Asp Leu
 80          85          90
Ser Asp Pro Ser Gly Trp Met Val Gln Phe Gly Gln Leu Thr Ser
 95          100         105
Met Pro Ser Phe Trp Ser Leu Gln Ala Tyr Tyr Thr Arg Tyr Phe
110         115         120
Val Ser Asn Ile Tyr Leu Ser Pro Arg Tyr Leu Gly Asn Ser Pro
125         130         135
Tyr Asp Ile Ala Leu Val Lys Leu Ser Ala Pro Val Thr Tyr Thr
140         145         150
Lys His Ile Gln Pro Ile Cys Leu Gln Ala Ser Thr Phe Glu Phe
155         160         165
Glu Asn Arg Thr Asp Cys Trp Val Thr Gly Trp Gly Tyr Ile Lys
170         175         180
Glu Asp Glu Ala Leu Pro Ser Pro His Thr Leu Gln Glu Val Gln
185         190         195
Val Ala Ile Ile Asn Asn Ser Met Cys Asn His Leu Phe Leu Lys
200         205         210
Tyr Ser Phe Arg Lys Asp Ile Phe Gly Asp Met Val Cys Ala Gly
215         220         225
Asn Ala Gln Gly Gly Lys Asp Ala Cys Phe Gly Asp Ser Gly Gly
230         235         240
Pro Leu Ala Cys Asn Lys Asn Gly Leu Trp Tyr Gln Ile Gly Val
245         250         255
Val Ser Trp Gly Val Gly Cys Gly Arg Pro Asn Arg Pro Gly Val
260         265         270
Tyr Thr Asn Ile Ser His His Phe Glu Trp Ile Gln Lys Leu Met
275         280         285
Ala Gln Ser Gly Met Ser Gln Pro Asp Pro Ser Trp Pro Leu Leu
290         295         300
Phe Phe Pro Leu Leu Trp Ala Leu Pro Leu Leu Gly Pro Val
305         310

```

&lt;210&gt; SEQ ID NO 258

&lt;211&gt; LENGTH: 2427

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 258

cccacgcgctc cgcggacgcg tgggaagggc agaatgggac tccaagcctg

50



-continued

---

cctcctaggg ctctttgccc tcctcctctc tggcaaatgc agttacagcc	100
cggagccccg ccagcggagg acgctgcccc caggctgggt gtccctgggc	150
cgtgcggacc ctgaggaaga gctgagtctc acctttgccc tgagacagca	200
gaatgtggaa agactctcgg agctggtgca ggctgtgtcg gatccagct	250
ctcctcaata cggaaaatac ctgaccctag agaatgtggc tgatctggtg	300
aggccatccc cactgaccct ccacacgggt caaaaatggc tcttgccagc	350
cggagccccg aagtgcatt ctgtgatcac acaggacttt ctgacttgct	400
ggctgagcat ccgacaagca gagctgtctc tcctggggc tgagtttcat	450
cactatgtgg gaggacctac ggaaccctat gttgtaaggc ccccatcc	500
ctaccagctt ccacaggcct tggccccca tgtggacttt gtggggggac	550
tgaccgcttt tcccccaaca tcctcctgga ggcaacgtcc tgagccgag	600
gtgacagggc ctgtaggcct gcatctgggg gtaaccctct ctgtgatccg	650
taagcgatac aacttgacct cacaagacgt gggctctggc accagcaata	700
acagccaagc ctgtgcccag ttcttgagac agtatttcca tgaactcagc	750
ctggctcagt tcatgcccct ctctgggtgc aactttgac atcaggcatc	800
agtagccccg gtggttgagc aacagggcgg gggccgggccc gggattgagg	850
ccagctctaga tgtgcagtac ctgatgagtg ctggtgccaa catctccacc	900
tgggtctaca gtagccctgg ccggcatgag ggacagagc ccttcctgca	950
gtggctcatg ctgctcagta atgagtcagc cctgcccact gtgcatactg	1000
tgagctatgg agatgatgag gactccctca gcagcgccta catccagcgg	1050
gtcaacactg agctcatgaa ggctgcccct cgggggtctca ccctgctctt	1100
cgctcaggt gacagtgagg ccgggtgttg gtctgtctct ggaagacacc	1150
agttccgccc taccttccct gcctccagcc cctatgtcac cacagtggga	1200
ggcacatcct tccaggaacc ttctctcctc acaaatgaaa ttgttgacta	1250
tatcagtggt ggtggcttca gcaatgtgtt cccacggcct tcataccagg	1300
aggaagctgt aacgaagttc ctgagctcta gccccacct gccaccatcc	1350
agttacttca atgccagtgg ccgtgacctc ccagatgtgg ctgcactttc	1400
tgatggctac tgggtggtca gcaacagagt gccattcca tgggtgtccg	1450
gaacctcgcc ctctactcca gtgtttgggg ggatcctatc cttgatcaat	1500
gagcacagga tccttagtgg ccgccccctt cttggctttc tcaacccaag	1550
gctctaccag cagcatgggg caggctctct tgatgtaacc cgtggctgcc	1600
atgagctcgt tctggatgaa gaggtagagg gccagggttt ctgctctggt	1650
cctggctggg atcctgtaac aggtcgggga acaccaactt cccagctttg	1700
ctgaagactc tactcaacct ctgacccttt cctatcagga gagatggctt	1750
gtcccctgcc ctgaagctgg cagttcagtc ccttattctg cctgttgga	1800
agccctgctg aacctcaac tattgactgc tgcagacagc ttatctocct	1850
aacctgaaa tgctgtgagc ttgacttgac toccaacctt accatgctcc	1900
atcatactca ggtctcccta ctctgcctt agattcctca ataagatgct	1950

-continued

---

```

gtaactagca ttttttgaat gcctctccct ccgcatctca tctttctctt      2000
ttcaatcagg cttttccaaa gggttgtata cagactctgt gcactatttc      2050
acttgatatt cattccccaa ttcactgcaa ggagacctct actgtcaccg      2100
tttactcttt cctaccctga catccagaaa caatggcctc cagtgcatac      2150
ttctcaatct ttgctttatg gcctttccat catagttgcc cactccctct      2200
ccttacttag cttccaggtc ttaactttctc tgactactct tgtcttctctc      2250
tctcatcaat ttctgcttct tcatggaatg ctgacctca ttgctccatt      2300
tgtagatatt tgctcttctc agtttactca ttgtcccctg gaacaaatca      2350
ctgacatcta caaccattac catctcacta aataagactt tctatccaat      2400
aatgattgat acctcaaatg taaaaaaa                                2427
    
```

```

<210> SEQ ID NO 259
<211> LENGTH: 556
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 259

```

Met Gly Leu Gln Ala Cys Leu Leu Gly Leu Phe Ala Leu Ile Leu
 1          5          10          15
Ser Gly Lys Cys Ser Tyr Ser Pro Glu Pro Asp Gln Arg Arg Thr
          20          25          30
Leu Pro Pro Gly Trp Val Ser Leu Gly Arg Ala Asp Pro Glu Glu
          35          40          45
Glu Leu Ser Leu Thr Phe Ala Leu Arg Gln Gln Asn Val Glu Arg
          50          55          60
Leu Ser Glu Leu Val Gln Ala Val Ser Asp Pro Ser Ser Pro Gln
          65          70          75
Tyr Gly Lys Tyr Leu Thr Leu Glu Asn Val Ala Asp Leu Val Arg
          80          85          90
Pro Ser Pro Leu Thr Leu His Thr Val Gln Lys Trp Leu Leu Ala
          95          100          105
Ala Gly Ala Gln Lys Cys His Ser Val Ile Thr Gln Asp Phe Leu
          110          115          120
Thr Cys Trp Leu Ser Ile Arg Gln Ala Glu Leu Leu Leu Pro Gly
          125          130          135
Ala Glu Phe His His Tyr Val Gly Gly Pro Thr Glu Thr His Val
          140          145          150
Val Arg Ser Pro His Pro Tyr Gln Leu Pro Gln Ala Leu Ala Pro
          155          160          165
His Val Asp Phe Val Gly Gly Leu His Arg Phe Pro Pro Thr Ser
          170          175          180
Ser Leu Arg Gln Arg Pro Glu Pro Gln Val Thr Gly Thr Val Gly
          185          190          195
Leu His Leu Gly Val Thr Pro Ser Val Ile Arg Lys Arg Tyr Asn
          200          205          210
Leu Thr Ser Gln Asp Val Gly Ser Gly Thr Ser Asn Asn Ser Gln
          215          220          225
Ala Cys Ala Gln Phe Leu Glu Gln Tyr Phe His Asp Ser Asp Leu
          230          235          240
    
```

-continued

---

Ala Gln Phe Met Arg Leu Phe Gly Gly Asn Phe Ala His Gln Ala  
 245 250 255

Ser Val Ala Arg Val Val Gly Gln Gln Gly Arg Gly Arg Ala Gly  
 260 265 270

Ile Glu Ala Ser Leu Asp Val Gln Tyr Leu Met Ser Ala Gly Ala  
 275 280 285

Asn Ile Ser Thr Trp Val Tyr Ser Ser Pro Gly Arg His Glu Gly  
 290 295 300

Gln Glu Pro Phe Leu Gln Trp Leu Met Leu Leu Ser Asn Glu Ser  
 305 310 315

Ala Leu Pro His Val His Thr Val Ser Tyr Gly Asp Asp Glu Asp  
 320 325 330

Ser Leu Ser Ser Ala Tyr Ile Gln Arg Val Asn Thr Glu Leu Met  
 335 340 345

Lys Ala Ala Ala Arg Gly Leu Thr Leu Leu Phe Ala Ser Gly Asp  
 350 355 360

Ser Gly Ala Gly Cys Trp Ser Val Ser Gly Arg His Gln Phe Arg  
 365 370 375

Pro Thr Phe Pro Ala Ser Ser Pro Tyr Val Thr Thr Val Gly Gly  
 380 385 390

Thr Ser Phe Gln Glu Pro Phe Leu Ile Thr Asn Glu Ile Val Asp  
 395 400 405

Tyr Ile Ser Gly Gly Gly Phe Ser Asn Val Phe Pro Arg Pro Ser  
 410 415 420

Tyr Gln Glu Glu Ala Val Thr Lys Phe Leu Ser Ser Ser Pro His  
 425 430 435

Leu Pro Pro Ser Ser Tyr Phe Asn Ala Ser Gly Arg Ala Tyr Pro  
 440 445 450

Asp Val Ala Ala Leu Ser Asp Gly Tyr Trp Val Val Ser Asn Arg  
 455 460 465

Val Pro Ile Pro Trp Val Ser Gly Thr Ser Ala Ser Thr Pro Val  
 470 475 480

Phe Gly Gly Ile Leu Ser Leu Ile Asn Glu His Arg Ile Leu Ser  
 485 490 495

Gly Arg Pro Pro Leu Gly Phe Leu Asn Pro Arg Leu Tyr Gln Gln  
 500 505 510

His Gly Ala Gly Leu Phe Asp Val Thr Arg Gly Cys His Glu Ser  
 515 520 525

Cys Leu Asp Glu Glu Val Glu Gly Gln Gly Phe Cys Ser Gly Pro  
 530 535 540

Gly Trp Asp Pro Val Thr Gly Trp Gly Thr Pro Thr Ser Gln Leu  
 545 550 555

Cys

<210> SEQ ID NO 260  
 <211> LENGTH: 1638  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 260

gccgcgcgct ctctccggc gccacacct gtctgagcgg cgcagcgagc

-continued

cgcgccccgg gcgggctgct cggcgcgga cagtgtctcg catggcaggg	100
attccagggc tcctcttctt tctcttcttt ctgctctgtg ctgttgggca	150
agttagccct tacagtgcc cctggaacc cacttgccct gcataccgcc	200
tcctgtctgt cttgccccag tctacctca atttagccaa gccagacttt	250
ggagccgaag ccaaattaga agtatcttct tcatgtggac cccagtgtca	300
taagggaact ccaactgccca cttacgaaga ggccaagcaa tatctgtctt	350
atgaaacgct ctatgccaat ggcagccgca cagagacgca ggtgggcatc	400
tacatcctca gcagtagtgg agatggggcc caacaccgag actcagggtc	450
ttcaggaag tctcgaagga agcggcagat ttatggctat gacagcaggt	500
tcagcatttt tgggaaggac ttctgtctca actacccttt ctcaacatca	550
gtgaagtatt ccacgggctg caccggcacc ctggtggcag agaagcatgt	600
cctcacagct gcccaactgca tacacgatgg aaaaacctat gtgaaaggaa	650
cccagaagct tcgagtgggc ttctaaaagc ccaagttaa agatggtggt	700
cgaggggcca acgactccac ttcagccatg cccgagcaga tgaatttca	750
gtgatccgg gtgaaacgca cccatgtgcc caagggttg atcaagggca	800
atgccaatga catcgccatg gattatgatt atgcctcct ggaactcaa	850
aagccccaca agagaaaatt tatgaagatt ggggtgagcc ctctgctaa	900
gcagctgcca gggggcagaa ttcacttctc tggttatgac aatgaccgac	950
caggcaattt ggtgtatcgc ttctgtgacg tcaaagacga gacctatgac	1000
ttgctctacc agcaatgcga tgcaccagca gggccagcg ggtctgggt	1050
ctatgtgagg atgtggaaga gacagcagca gaagtgggag cgaaaaatta	1100
ttggcatttt ttcagggcac cagtgggtgg acatgaatgg ttccccacag	1150
gatttcaacg tggctgtcag aatcactcct ctcaaatag cccagatttg	1200
ctattggatt aaagaaaact acctggattg tagggagggg tgacacagtg	1250
ttccctcctg gcagcaatta agggcttca tgttcttatt ttaggagagg	1300
ccaaattggt ttttgtcatt ggcgtgcaca cgtgtgtgtg tgtgtgtgtg	1350
tgtgtgtaag gtgtcttata atcttttacc tatttcttac aattgcaaga	1400
tgactggctt tactatttga aaactggttt gtgtatcata tcatatatca	1450
tttaagcagt ttgaaggcat acttttgcat agaaataaaa aaaatactga	1500
tttggggcaa tgaggaatat ttgacaatta agttaatcct cacgtttttg	1550
caaactttga tttttatttc atctgaactt gtttcaaaga tttatattaa	1600
atatttggca tacaagagat atgaaaaaaaa aaaaaaaaa	1638

&lt;210&gt; SEQ ID NO 261

&lt;211&gt; LENGTH: 383

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 261

Met Ala Gly Ile Pro Gly Leu Leu Phe Leu Leu Phe Phe Leu Leu  
 1                    5                    10                    15

Cys Ala Val Gly Gln Val Ser Pro Tyr Ser Ala Pro Trp Lys Pro

-continued

										20	25						30					
Thr	Trp	Pro	Ala	Tyr	Arg	Leu	Pro	Val	Val	Leu	Pro	Gln	Ser	Thr	35	40						45
Leu	Asn	Leu	Ala	Lys	Pro	Asp	Phe	Gly	Ala	Glu	Ala	Lys	Leu	Glu	50	55						60
Val	Ser	Ser	Ser	Cys	Gly	Pro	Gln	Cys	His	Lys	Gly	Thr	Pro	Leu	65	70						75
Pro	Thr	Tyr	Glu	Glu	Ala	Lys	Gln	Tyr	Leu	Ser	Tyr	Glu	Thr	Leu	80	85						90
Tyr	Ala	Asn	Gly	Ser	Arg	Thr	Glu	Thr	Gln	Val	Gly	Ile	Tyr	Ile	95	100						105
Leu	Ser	Ser	Ser	Gly	Asp	Gly	Ala	Gln	His	Arg	Asp	Ser	Gly	Ser	110	115						120
Ser	Gly	Lys	Ser	Arg	Arg	Lys	Arg	Gln	Ile	Tyr	Gly	Tyr	Asp	Ser	125	130						135
Arg	Phe	Ser	Ile	Phe	Gly	Lys	Asp	Phe	Leu	Leu	Asn	Tyr	Pro	Phe	140	145						150
Ser	Thr	Ser	Val	Lys	Leu	Ser	Thr	Gly	Cys	Thr	Gly	Thr	Leu	Val	155	160						165
Ala	Glu	Lys	His	Val	Leu	Thr	Ala	Ala	His	Cys	Ile	His	Asp	Gly	170	175						180
Lys	Thr	Tyr	Val	Lys	Gly	Thr	Gln	Lys	Leu	Arg	Val	Gly	Phe	Leu	185	190						195
Lys	Pro	Lys	Phe	Lys	Asp	Gly	Gly	Arg	Gly	Ala	Asn	Asp	Ser	Thr	200	205						210
Ser	Ala	Met	Pro	Glu	Gln	Met	Lys	Phe	Gln	Trp	Ile	Arg	Val	Lys	215	220						225
Arg	Thr	His	Val	Pro	Lys	Gly	Trp	Ile	Lys	Gly	Asn	Ala	Asn	Asp	230	235						240
Ile	Gly	Met	Asp	Tyr	Asp	Tyr	Ala	Leu	Leu	Glu	Leu	Lys	Lys	Pro	245	250						255
His	Lys	Arg	Lys	Phe	Met	Lys	Ile	Gly	Val	Ser	Pro	Pro	Ala	Lys	260	265						270
Gln	Leu	Pro	Gly	Gly	Arg	Ile	His	Phe	Ser	Gly	Tyr	Asp	Asn	Asp	275	280						285
Arg	Pro	Gly	Asn	Leu	Val	Tyr	Arg	Phe	Cys	Asp	Val	Lys	Asp	Glu	290	295						300
Thr	Tyr	Asp	Leu	Leu	Tyr	Gln	Gln	Cys	Asp	Ala	Gln	Pro	Gly	Ala	305	310						315
Ser	Gly	Ser	Gly	Val	Tyr	Val	Arg	Met	Trp	Lys	Arg	Gln	Gln	Gln	320	325						330
Lys	Trp	Glu	Arg	Lys	Ile	Ile	Gly	Ile	Phe	Ser	Gly	His	Gln	Trp	335	340						345
Val	Asp	Met	Asn	Gly	Ser	Pro	Gln	Asp	Phe	Asn	Val	Ala	Val	Arg	350	355						360
Ile	Thr	Pro	Leu	Lys	Tyr	Ala	Gln	Ile	Cys	Tyr	Trp	Ile	Lys	Gly	365	370						375
Asn	Tyr	Leu	Asp	Cys	Arg	Glu	Gly								380							

-continued

---

```

<211> LENGTH: 1378
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 262
gcacgcacct gggctctctcg agcctgctgc ctgctcccc gccccaccag      50
ccatggtggt ttctggagcg cccccagccc tgggtggggg ctgtctcggc      100
acctcacct ccctgctgct gctggcgtcg acagccatcc tcaatgcggc      150
caggatacct gttccccag cctgtgggaa gccccagcag ctgaaccggg      200
ttgtgggcgg cgaggacagc actgacagcg agtggccctg gatcgtgagc      250
atccagaaga atgggacca cactgcgca ggttctctgc tcaccagccg      300
ctgggtgatc actgctgcc actgtttcaa ggacaacctg aacaaacct      350
acctgttctc tgtgctgctg ggggcctggc agctgggaa ccctggctct      400
cgggccaga aggtgggtgt tgccctgggt gagccccacc ctgtgtattc      450
ctggaagaa ggtgcctgtg cagacattgc cctgggtcgt ctcgagcgt      500
ccatacagtt ctacagcgg gtccctgcca totgcctacc tgatgcctct      550
atccacctcc ctccaaacac cactgctgg atctcaggct gggggagcat      600
ccaagatgga gttcccttg cccacctca gacctgcag aagctgaagg      650
ttcctatcat cgactcggaa gtctgcagcc atctgtactg gcggggagca      700
ggacagggac ccatcactga ggacatgctg tgtgccggt acttgagggg      750
ggagcgggat gcttgtctgg gcgactccgg gggccccctc atgtgccagg      800
tggacggcgc ctggctgctg gccggcatca tcagctgggg cgagggctgt      850
gccgagcga acagggccgg ggtctacatc agcctctctg cgcaccgctc      900
ctgggtggag aagatcgtgc aaggggtgca gctccgcggg cgcgctcagg      950
gggtggggg cctcagggca ccgagccagg gctctggggc cgcgcgcgc      1000
tcctagggcg cagcgggacg cggggctcgg atctgaaagg cggccagatc      1050
cacatctgga tctggatctg cggcgcctc gggcggttc cccgcgcgta      1100
aataggctca tctacctcta cctctggggg cccggacggc tgetgcggaa      1150
aggaacccc ctccccgacc cggccgacgg cctcaggccc ccctccaagg      1200
catcaggccc cgcccaacgg cctcatgtcc cggccccac gacttcggc      1250
cccggcccc ggccccagcg cttttgtgta tataaatgtt aatgattttt      1300
ataggtattt gtaaccctgc ccacatatct tatttattcc tccaatttca      1350
ataaattatt tattctccaa aaaaaaaaa      1378
    
```

```

<210> SEQ ID NO 263
<211> LENGTH: 317
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 263
Met Val Val Ser Gly Ala Pro Pro Ala Leu Gly Gly Gly Cys Leu
 1             5             10             15
Gly Thr Phe Thr Ser Leu Leu Leu Ala Ser Thr Ala Ile Leu
                20             25             30
    
```

-continued

Asn	Ala	Ala	Arg	Ile	Pro	Val	Pro	Pro	Ala	Cys	Gly	Lys	Pro	Gln
				35					40					45
Gln	Leu	Asn	Arg	Val	Val	Gly	Gly	Glu	Asp	Ser	Thr	Asp	Ser	Glu
				50					55					60
Trp	Pro	Trp	Ile	Val	Ser	Ile	Gln	Lys	Asn	Gly	Thr	His	His	Cys
				65					70					75
Ala	Gly	Ser	Leu	Leu	Thr	Ser	Arg	Trp	Val	Ile	Thr	Ala	Ala	His
				80					85					90
Cys	Phe	Lys	Asp	Asn	Leu	Asn	Lys	Pro	Tyr	Leu	Phe	Ser	Val	Leu
				95					100					105
Leu	Gly	Ala	Trp	Gln	Leu	Gly	Asn	Pro	Gly	Ser	Arg	Ser	Gln	Lys
				110					115					120
Val	Gly	Val	Ala	Trp	Val	Glu	Pro	His	Pro	Val	Tyr	Ser	Trp	Lys
				125					130					135
Glu	Gly	Ala	Cys	Ala	Asp	Ile	Ala	Leu	Val	Arg	Leu	Glu	Arg	Ser
				140					145					150
Ile	Gln	Phe	Ser	Glu	Arg	Val	Leu	Pro	Ile	Cys	Leu	Pro	Asp	Ala
				155					160					165
Ser	Ile	His	Leu	Pro	Pro	Asn	Thr	His	Cys	Trp	Ile	Ser	Gly	Trp
				170					175					180
Gly	Ser	Ile	Gln	Asp	Gly	Val	Pro	Leu	Pro	His	Pro	Gln	Thr	Leu
				185					190					195
Gln	Lys	Leu	Lys	Val	Pro	Ile	Ile	Asp	Ser	Glu	Val	Cys	Ser	His
				200					205					210
Leu	Tyr	Trp	Arg	Gly	Ala	Gly	Gln	Gly	Pro	Ile	Thr	Glu	Asp	Met
				215					220					225
Leu	Cys	Ala	Gly	Tyr	Leu	Glu	Gly	Glu	Arg	Asp	Ala	Cys	Leu	Gly
				230					235					240
Asp	Ser	Gly	Gly	Pro	Leu	Met	Cys	Gln	Val	Asp	Gly	Ala	Trp	Leu
				245					250					255
Leu	Ala	Gly	Ile	Ile	Ser	Trp	Gly	Glu	Gly	Cys	Ala	Glu	Arg	Asn
				260					265					270
Arg	Pro	Gly	Val	Tyr	Ile	Ser	Leu	Ser	Ala	His	Arg	Ser	Trp	Val
				275					280					285
Glu	Lys	Ile	Val	Gln	Gly	Val	Gln	Leu	Arg	Gly	Arg	Ala	Gln	Gly
				290					295					300
Gly	Gly	Ala	Leu	Arg	Ala	Pro	Ser	Gln	Gly	Ser	Gly	Ala	Ala	Ala
				305					310					315

Arg Ser

<210> SEQ ID NO 264  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
 <400> SEQUENCE: 264

gtccgcaagg atgctacat gttc

24

<210> SEQ ID NO 265  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

---

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 265

gcagaggtgt ctaaggttg 19

<210> SEQ ID NO 266  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 266

agctctagac caatgccagc ttcc 24

<210> SEQ ID NO 267  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 267

gccaccaact cctgcaagaa cttctcagaa ctgcccctgg tcatg 45

<210> SEQ ID NO 268  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 268

ggggaattca ccctatgaca ttgcc 25

<210> SEQ ID NO 269  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 269

gaatgccctg caagcatcaa ctgg 24

<210> SEQ ID NO 270  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 270

gcacctgtca cctacactaa acacatccag cccatctgtc tccaggcctc 50

<210> SEQ ID NO 271  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 271



---

-continued

---

gcggaagggc agaatgggac tccaag 26

<210> SEQ ID NO 272  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 272

cagccctgcc acatgtgc 18

<210> SEQ ID NO 273  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 273

tactgggtgg tcagcaac 18

<210> SEQ ID NO 274  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 274

ggcgaagagc agggtgagac cccg 24

<210> SEQ ID NO 275  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 275

gccctcatcc tctctggcaa atgcagttac agcccggagc ccgac 45

<210> SEQ ID NO 276  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 276

gggcagggat tccagggctc c 21

<210> SEQ ID NO 277  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 277

ggctatgaca gcaggttc 18

-continued

---

```

<210> SEQ ID NO 278
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 278

tgacaatgac cgaccagg                18

<210> SEQ ID NO 279
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 279

gcatcgcatt gctgtagag caag          24

<210> SEQ ID NO 280
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 280

ttacagtgcc ccctgaaac ccaactggcc tgcataccgc ctccc          45

<210> SEQ ID NO 281
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 281

cgtctcgagc gtcctataca gttcccttgc ccca                34

<210> SEQ ID NO 282
<211> LENGTH: 61
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 282

tggaggggga gcgggatgct tgtctgggag actccggggg ccccctcatg          50
tgccaggtgg a                                          61

<210> SEQ ID NO 283
<211> LENGTH: 119
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 283

ccctcagacc ctgcagaagc tgaaggttcc tatcatcgac tcggaagtct          50
gcagccatct gtactggcgg ggagcaggac aggaccat cactgaggac          100
atgctgtgtg ccggctact                                119

```

-continued

---

<210> SEQ ID NO 284

<211> LENGTH: 1875

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 284

gacggctggc caccatgcac ggctcctgca gtttctgat gcttctgctg	50
ccgctactgc tactgctggt ggccaccaca ggccccgttg gagccctcac	100
agatgaggag aaacgtttga tggtagagct gcacaacctc taccggggccc	150
aggatcccc gacggcctca gacatgctgc acatgagatg ggacgaggag	200
ctggcccgct tcgccaaggc ctacgcacgg cagtgcgtgt ggggccacaa	250
caaggagcgc gggcgcgcgc gcgagaatct gttcgccatc acagacgagg	300
gcatggacgt gccgctggcc atggaggagt ggcaccacga gcgtgagcac	350
tacaacctca gcgccgccac ctgcagccca ggccagatgt gcggccacta	400
cacgcagggt gtatgggcca agacagagag gatcggctgt ggttcccact	450
tctgtgagaa gctccagggt gttgaggaga ccaacatcga attactggtg	500
tgcaactatg agcctccggg gaactgtaag gggaaacggc cctaccagga	550
ggggactccg tgctccaat gtccctctgg ctaccactgc aagaactccc	600
tctgtgaacc catcggaagc ccggaagatg ctcaggattt gccttactctg	650
gtaactgagg ccccatcctt ccgggcgact gaagcatcag actctagga	700
aatgggtact ccttcttccc tagcaacggg gattccggct ttcttggtaa	750
cagaggctctc aggtccctg gcaaccaagg ctctgcctgc tgtgaaacc	800
caggccccaa cttccttagc aacgaaagac ccgcctcca tggcaacaga	850
ggctccacct tgcgtaacaa ctgaggctcc ttccattttg gcagctcaca	900
gcctgccctc cttggatgag gagccagtta cttccccaa atcgacctat	950
gttctatctc caaaatcagc agacaagtgc acagacaaaa caaaagtgcc	1000
ctctaggagc ccagagaact ctctggacct caagatgtcc ctgacagggg	1050
caagggaact cctaccctat gccacggagg aggctgaggc tgaggctgag	1100
ttgcctcctt ccagtgaggt cttggcctca gtttttcag cccaggacaa	1150
gccagggtgag ctgcaggcca cactggacca cacggggcac acctcctcca	1200
agtcctctgc caatttcccc aatacctctg ccacgcgtaa tgccaacgggt	1250
gggcgtgccc tggctctgca gtcgtccttg ccagggtcag agggccctga	1300
caagcctagc gttgtgtcag ggctgaactc gggcctgggt catgtgtggg	1350
gccctctcct gggactactg ctctgcctc ctctgggtgt gctggaatc	1400
ttctgaatgg gataccactc aaagggtgaa gaggtcagct gtcctcctgt	1450
catcttcccc accctgtccc cagcccctaa acaagatact tcttggttaa	1500
ggccctccgc aagggaagg ctacggggca tgtgcctcat cacaccatcc	1550
atcctggagg cacaaggcct ggctggctgc gagctcagga ggccgcctga	1600
ggactgcaca ccggggcccac acctctcctg cccctccctc ctgagtctctg	1650
gggggtggag gatttgaggg agctcactgc ctacctggcc tggggctgtc	1700

-continued

---

```

tgcccacaca gcatgtgccc tctccctgag tgcctgtgta gctgggggatg      1750
gggattccta ggggcagatg aaggacaagc cccactggag tggggttctt      1800
tgagtggggg aggcagggac gagggaagga aagtaactcc tgactctcca      1850
ataaaaaact gtccaacctg tgaaa                                     1875

```

&lt;210&gt; SEQ ID NO 285

&lt;211&gt; LENGTH: 463

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 285

```

Met His Gly Ser Cys Ser Phe Leu Met Leu Leu Leu Pro Leu Leu
 1          5          10          15
Leu Leu Leu Val Ala Thr Thr Gly Pro Val Gly Ala Leu Thr Asp
 20          25          30
Glu Glu Lys Arg Leu Met Val Glu Leu His Asn Leu Tyr Arg Ala
 35          40          45
Gln Val Ser Pro Thr Ala Ser Asp Met Leu His Met Arg Trp Asp
 50          55          60
Glu Glu Leu Ala Ala Phe Ala Lys Ala Tyr Ala Arg Gln Cys Val
 65          70          75
Trp Gly His Asn Lys Glu Arg Gly Arg Arg Gly Glu Asn Leu Phe
 80          85          90
Ala Ile Thr Asp Glu Gly Met Asp Val Pro Leu Ala Met Glu Glu
 95          100         105
Trp His His Glu Arg Glu His Tyr Asn Leu Ser Ala Ala Thr Cys
110          115         120
Ser Pro Gly Gln Met Cys Gly His Tyr Thr Gln Val Val Trp Ala
125          130         135
Lys Thr Glu Arg Ile Gly Cys Gly Ser His Phe Cys Glu Lys Leu
140          145         150
Gln Gly Val Glu Glu Thr Asn Ile Glu Leu Leu Val Cys Asn Tyr
155          160         165
Glu Pro Pro Gly Asn Val Lys Gly Lys Arg Pro Tyr Gln Glu Gly
170          175         180
Thr Pro Cys Ser Gln Cys Pro Ser Gly Tyr His Cys Lys Asn Ser
185          190         195
Leu Cys Glu Pro Ile Gly Ser Pro Glu Asp Ala Gln Asp Leu Pro
200          205         210
Tyr Leu Val Thr Glu Ala Pro Ser Phe Arg Ala Thr Glu Ala Ser
215          220         225
Asp Ser Arg Lys Met Gly Thr Pro Ser Ser Leu Ala Thr Gly Ile
230          235         240
Pro Ala Phe Leu Val Thr Glu Val Ser Gly Ser Leu Ala Thr Lys
245          250         255
Ala Leu Pro Ala Val Glu Thr Gln Ala Pro Thr Ser Leu Ala Thr
260          265         270
Lys Asp Pro Pro Ser Met Ala Thr Glu Ala Pro Pro Cys Val Thr
275          280         285
Thr Glu Val Pro Ser Ile Leu Ala Ala His Ser Leu Pro Ser Leu
290          295         300

```

-continued

Asp Glu Glu Pro Val Thr Phe Pro Lys Ser Thr His Val Pro Ile  
 305 310 315  
 Pro Lys Ser Ala Asp Lys Val Thr Asp Lys Thr Lys Val Pro Ser  
 320 325 330  
 Arg Ser Pro Glu Asn Ser Leu Asp Pro Lys Met Ser Leu Thr Gly  
 335 340 345  
 Ala Arg Glu Leu Leu Pro His Ala Gln Glu Glu Ala Glu Ala Glu  
 350 355 360  
 Ala Glu Leu Pro Pro Ser Ser Glu Val Leu Ala Ser Val Phe Pro  
 365 370 375  
 Ala Gln Asp Lys Pro Gly Glu Leu Gln Ala Thr Leu Asp His Thr  
 380 385 390  
 Gly His Thr Ser Ser Lys Ser Leu Pro Asn Phe Pro Asn Thr Ser  
 395 400 405  
 Ala Thr Ala Asn Ala Thr Gly Gly Arg Ala Leu Ala Leu Gln Ser  
 410 415 420  
 Ser Leu Pro Gly Ala Glu Gly Pro Asp Lys Pro Ser Val Val Ser  
 425 430 435  
 Gly Leu Asn Ser Gly Pro Gly His Val Trp Gly Pro Leu Leu Gly  
 440 445 450  
 Leu Leu Leu Leu Pro Pro Leu Val Leu Ala Gly Ile Phe  
 455 460

<210> SEQ ID NO 286  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 286

tcctgcagtt tcctgatgc

19

<210> SEQ ID NO 287  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 287

ctcatattgc acaccagtaa ttcg

24

<210> SEQ ID NO 288  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 288

atgaggagaa acgtttgatg gtggagctgc acaacctcta ccggg

45

<210> SEQ ID NO 289  
 <211> LENGTH: 3662  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

-continued

&lt;400&gt; SEQUENCE: 289

gtaactgaag tcaggctttt catttgggaa gcccctcaa cagaattcgg	50
tcattctcca agttatgggt gacgtacttc tgttgttctc cctctgcttg	100
ctttttcaca ttagcagacc ggacttaagt cacaacagat tatctttcat	150
caaggcaagt tccatgagcc accttcaaag ccttcogagaa gtgaaactga	200
acaacaatga attggagacc attccaaatc tgggaccagt ctcggcaaat	250
attacacttc tctccttggc tggaaacagc attgttgaaa tactccctga	300
acatctgaaa gagtttcagt cccttgaaac tttggacctt agcagcaaca	350
atatttcaga gctccaaact gcatttccag ccctacagct caaatatctg	400
tatctcaaca gcaaccgagt cacatcaatg gaacctgggt attttgaaa	450
tttgccaac acactccttg tgttaaagct gaacaggaac cgaatctcag	500
ctatcccacc caagatgttt aaactgcccc aactgcaaca tctcgaattg	550
aaccgaaaa agattaataa tgtagatgga ctgacattcc aaggccttgg	600
tgctctgaag tctctgaaaa tgcaagaaa tggagtaacg aaacttatgg	650
atggagcttt ttgggggctg agcaacatgg aaattttgca gctggacct	700
aaacacctaa cagagattac caaaggctgg ctttacggct tgctgatgct	750
gcaggaaact catctcagcc aaaatgccc caacagatc agccctgatg	800
cctgggagtt ctgccagaag ctcagtgagc tggacctaac tttcaatcac	850
ttatcaaggt tagatgattc aagcttcctt ggcctaagct tactaaatac	900
actgcacatt gggaaacaaca gagtcagcta cattgctgat tgtgccttcc	950
gggggctttc cagtttaag actttggatc tgaagaacaa tgaatttcc	1000
tggactattg aagacatgaa tgggtctttc totgggcttg acaactgag	1050
gcgactgata ctccaaggaa atcggatccg ttctattact aaaaaagcct	1100
tcactggttt ggatgcattg gagcatctag acctgagtga caacgcaatc	1150
atgtctttac aaggcaatgc attttcacaa atgaagaaac tgcaacaatt	1200
gcatttaaat acatcaagcc ttttgtgcca ttgccagcta aaatggctcc	1250
cacagtgggt ggcggaaaac aactttcaga gctttgtaaa tgccagttgt	1300
gccatcctc agctgctaaa aggaagaagc atttttgctg ttagcccaga	1350
tggctttgtg tgtgatgatt ttcccacc ccagatcacg gttcagccag	1400
aaacacagtc ggcaataaaa ggtccaatt tgagtttcat ctgctcagct	1450
gccagcagca gtgattcccc aatgactttt gcttgaaaa aagacaatga	1500
actactgcat gatgctgaaa tggaaaatta tgcacacctc cgggcccaag	1550
gtggcgaggt gatggagat accaccatcc ttcggctgcg cgagggtgaa	1600
tttgccagtg aggggaaata tcagtgtgto atctccaatc actttggttc	1650
atcctactct gtcaaagcca agcttacagt aaatatgctt cctcattca	1700
ccaagacccc catggatctc accatccgag ctggggccat ggcacgcttg	1750
gagtgtgctg ctgtggggca cccagcccc cagatagcct ggcagaagga	1800
tgggggcaca gacttcccag ctgcacggga gagacgatg catgtgatgc	1850

-continued

---

ccgaggatga cgtgttcttt atcgtggatg tgaagataga ggacattggg	1900
gtatacagct gcacagctca gaacagtgca ggaagtattt cagcaaatgc	1950
aactctgact gtctagaaa caccatcatt tttgcggcca ctgttggacc	2000
gaactgtaac caagggagaa acagccgtcc tacagtgcac tgctggagga	2050
agccctcccc ctaaactgaa ctggacaaa gatgatagcc cattgttgggt	2100
aaccgagagg cacttttttg cagcaggcaa tcagcttctg attattgtgg	2150
actcagatgt cagtgatgct gggaaataca catgtgagat gtctaacc	2200
cttggcactg agagaggaaa cgtgcccctc agtgtgatcc cactccaac	2250
ctgcgactcc cctcagatga cagccccatc gttagacgat gacggatggg	2300
ccactgtggg tgtcgtgac atagccgtgg tttgctgtgt ggtgggcacg	2350
tcactcgtgt ggggtgtcat catataccac acaaggcggg ggaatgaaga	2400
ttgcagcatt accaacacag atgagaccaa cttgccagca gatattocta	2450
gttattttgt atctcaggga acgtagctg acaggcagga tgggtaactg	2500
tcttcagaaa gtggaagcca ccaccagttt gtcacatctt caggtgctgg	2550
atTTTTctta ccacaacatg acagttagtg gacctgccat attgacaata	2600
gcagtgaagc tgatgtggaa gctgccacag atctgttctt ttgtccgttt	2650
ttgggatcca caggccctat gtatttgaag ggaatgtgt atggctcaga	2700
tccttttgaa acatatcata caggttgtag tcctgaccca agaacagttt	2750
taatggacca ctatgagccc agttacataa agaaaaagga gtgotacca	2800
tgttctcatc cttcagaaga atcctgcgaa cggagcttca gtaatatatc	2850
gtggccttca catgtgagga agctacttaa cactagttag tctcacaatg	2900
aaggacctgg aatgaaaaat ctgtgtctaa acaagtcttc tttagatttt	2950
agtgcaaatc cagagccagc gtcggttgcc tcgagtaatt ctttcatggg	3000
tacctttgga aaagctctca ggagacctca cctagatgcc tattcaagct	3050
ttggacagcc atcagattgt cagccaagag ccttttattt gaaagctcat	3100
tcttcccag acttgactc tgggtcagag gaagatggga aagaaaggac	3150
agattttcag gaagaaaatc acatttgtac ctttaaacag actttagaaa	3200
actacagyac tccaaatttt cagtcttatg acttggacac atagactgaa	3250
tgagacaaa ggaaaagcctt aacatactac ctcaagttaa cttttattta	3300
aaagagagag aatccttatgt tttttaaag gagttatgaa ttttaaagg	3350
ataaaaatgc tttatttata cagatgaacc aaaattacaa aaagttatga	3400
aaatTTTTat actgggaatg atgctcatat aagaatacct ttttaaacta	3450
TTTTTTaact ttgttttatg caaaaaagta tcttacgtaa attaatgata	3500
taaatcatga ttattttatg tatttttata atgccagatt tctttttatg	3550
gaaaatgagt tactaaagca ttttaaataa tacctgcctt gtaccatttt	3600
ttaaatagaa gttacttcat tatattttgc acattatatt taataaaatg	3650
tgtaaatgtg aa	3662

-continued

&lt;211&gt; LENGTH: 1059

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 290

```

Met Val Asp Val Leu Leu Leu Phe Ser Leu Cys Leu Leu Phe His
 1           5           10          15
Ile Ser Arg Pro Asp Leu Ser His Asn Arg Leu Ser Phe Ile Lys
          20          25          30
Ala Ser Ser Met Ser His Leu Gln Ser Leu Arg Glu Val Lys Leu
          35          40          45
Asn Asn Asn Glu Leu Glu Thr Ile Pro Asn Leu Gly Pro Val Ser
          50          55          60
Ala Asn Ile Thr Leu Leu Ser Leu Ala Gly Asn Arg Ile Val Glu
          65          70          75
Ile Leu Pro Glu His Leu Lys Glu Phe Gln Ser Leu Glu Thr Leu
          80          85          90
Asp Leu Ser Ser Asn Asn Ile Ser Glu Leu Gln Thr Ala Phe Pro
          95          100         105
Ala Leu Gln Leu Lys Tyr Leu Tyr Leu Asn Ser Asn Arg Val Thr
          110         115         120
Ser Met Glu Pro Gly Tyr Phe Asp Asn Leu Ala Asn Thr Leu Leu
          125         130         135
Val Leu Lys Leu Asn Arg Asn Arg Ile Ser Ala Ile Pro Pro Lys
          140         145         150
Met Phe Lys Leu Pro Gln Leu Gln His Leu Glu Leu Asn Arg Asn
          155         160         165
Lys Ile Lys Asn Val Asp Gly Leu Thr Phe Gln Gly Leu Gly Ala
          170         175         180
Leu Lys Ser Leu Lys Met Gln Arg Asn Gly Val Thr Lys Leu Met
          185         190         195
Asp Gly Ala Phe Trp Gly Leu Ser Asn Met Glu Ile Leu Gln Leu
          200         205         210
Asp His Asn Asn Leu Thr Glu Ile Thr Lys Gly Trp Leu Tyr Gly
          215         220         225
Leu Leu Met Leu Gln Glu Leu His Leu Ser Gln Asn Ala Ile Asn
          230         235         240
Arg Ile Ser Pro Asp Ala Trp Glu Phe Cys Gln Lys Leu Ser Glu
          245         250         255
Leu Asp Leu Thr Phe Asn His Leu Ser Arg Leu Asp Asp Ser Ser
          260         265         270
Phe Leu Gly Leu Ser Leu Leu Asn Thr Leu His Ile Gly Asn Asn
          275         280         285
Arg Val Ser Tyr Ile Ala Asp Cys Ala Phe Arg Gly Leu Ser Ser
          290         295         300
Leu Lys Thr Leu Asp Leu Lys Asn Asn Glu Ile Ser Trp Thr Ile
          305         310         315
Glu Asp Met Asn Gly Ala Phe Ser Gly Leu Asp Lys Leu Arg Arg
          320         325         330
Leu Ile Leu Gln Gly Asn Arg Ile Arg Ser Ile Thr Lys Lys Ala
          335         340         345
Phe Thr Gly Leu Asp Ala Leu Glu His Leu Asp Leu Ser Asp Asn

```



-continued

	350		355		360
Ala Ile Met Ser	Leu Gln Gly Asn Ala	Phe Ser Gln Met Lys	Lys		
	365		370		375
Leu Gln Gln Leu	His Leu Asn Thr Ser	Ser Leu Leu Cys Asp	Cys		
	380		385		390
Gln Leu Lys Trp	Leu Pro Gln Trp Val	Ala Glu Asn Asn Phe	Gln		
	395		400		405
Ser Phe Val Asn	Ala Ser Cys Ala His	Pro Gln Leu Leu Lys	Gly		
	410		415		420
Arg Ser Ile Phe	Ala Val Ser Pro Asp	Gly Phe Val Cys Asp	Asp		
	425		430		435
Phe Pro Lys Pro	Gln Ile Thr Val Gln	Pro Glu Thr Gln Ser	Ala		
	440		445		450
Ile Lys Gly Ser	Asn Leu Ser Phe Ile	Cys Ser Ala Ala Ser	Ser		
	455		460		465
Ser Asp Ser Pro	Met Thr Phe Ala Trp	Lys Lys Asp Asn Glu	Leu		
	470		475		480
Leu His Asp Ala	Glu Met Glu Asn Tyr	Ala His Leu Arg Ala	Gln		
	485		490		495
Gly Gly Glu Val	Met Glu Tyr Thr Thr	Ile Leu Arg Leu Arg	Glu		
	500		505		510
Val Glu Phe Ala	Ser Glu Gly Lys Tyr	Gln Cys Val Ile Ser	Asn		
	515		520		525
His Phe Gly Ser	Ser Tyr Ser Val Lys	Ala Lys Leu Thr Val	Asn		
	530		535		540
Met Leu Pro Ser	Phe Thr Lys Thr Pro	Met Asp Leu Thr Ile	Arg		
	545		550		555
Ala Gly Ala Met	Ala Arg Leu Glu Cys	Ala Ala Val Gly His	Pro		
	560		565		570
Ala Pro Gln Ile	Ala Trp Gln Lys Asp	Gly Gly Thr Asp Phe	Pro		
	575		580		585
Ala Ala Arg Glu	Arg Arg Met His Val	Met Pro Glu Asp Asp	Val		
	590		595		600
Phe Phe Ile Val	Asp Val Lys Ile Glu	Asp Ile Gly Val Tyr	Ser		
	605		610		615
Cys Thr Ala Gln	Asn Ser Ala Gly Ser	Ile Ser Ala Asn Ala	Thr		
	620		625		630
Leu Thr Val Leu	Glu Thr Pro Ser Phe	Leu Arg Pro Leu Leu	Asp		
	635		640		645
Arg Thr Val Thr	Lys Gly Glu Thr Ala	Val Leu Gln Cys Ile	Ala		
	650		655		660
Gly Gly Ser Pro	Pro Pro Lys Leu Asn	Trp Thr Lys Asp Asp	Ser		
	665		670		675
Pro Leu Val Val	Thr Glu Arg His Phe	Phe Ala Ala Gly Asn	Gln		
	680		685		690
Leu Leu Ile Ile	Val Asp Ser Asp Val	Ser Asp Ala Gly Lys	Tyr		
	695		700		705
Thr Cys Glu Met	Ser Asn Thr Leu Gly	Thr Glu Arg Gly Asn	Val		
	710		715		720
Arg Leu Ser Val	Ile Pro Thr Pro Thr	Cys Asp Ser Pro Gln	Met		
	725		730		735

-continued

---

Thr Ala Pro Ser Leu Asp Asp Asp Gly Trp Ala Thr Val Gly Val  
 740 745 750

Val Ile Ile Ala Val Val Cys Cys Val Val Gly Thr Ser Leu Val  
 755 760 765

Trp Val Val Ile Ile Tyr His Thr Arg Arg Arg Asn Glu Asp Cys  
 770 775 780

Ser Ile Thr Asn Thr Asp Glu Thr Asn Leu Pro Ala Asp Ile Pro  
 785 790 795

Ser Tyr Leu Ser Ser Gln Gly Thr Leu Ala Asp Arg Gln Asp Gly  
 800 805 810

Tyr Val Ser Ser Glu Ser Gly Ser His His Gln Phe Val Thr Ser  
 815 820 825

Ser Gly Ala Gly Phe Phe Leu Pro Gln His Asp Ser Ser Gly Thr  
 830 835 840

Cys His Ile Asp Asn Ser Ser Glu Ala Asp Val Glu Ala Ala Thr  
 845 850 855

Asp Leu Phe Leu Cys Pro Phe Leu Gly Ser Thr Gly Pro Met Tyr  
 860 865 870

Leu Lys Gly Asn Val Tyr Gly Ser Asp Pro Phe Glu Thr Tyr His  
 875 880 885

Thr Gly Cys Ser Pro Asp Pro Arg Thr Val Leu Met Asp His Tyr  
 890 895 900

Glu Pro Ser Tyr Ile Lys Lys Lys Glu Cys Tyr Pro Cys Ser His  
 905 910 915

Pro Ser Glu Glu Ser Cys Glu Arg Ser Phe Ser Asn Ile Ser Trp  
 920 925 930

Pro Ser His Val Arg Lys Leu Leu Asn Thr Ser Tyr Ser His Asn  
 935 940 945

Glu Gly Pro Gly Met Lys Asn Leu Cys Leu Asn Lys Ser Ser Leu  
 950 955 960

Asp Phe Ser Ala Asn Pro Glu Pro Ala Ser Val Ala Ser Ser Asn  
 965 970 975

Ser Phe Met Gly Thr Phe Gly Lys Ala Leu Arg Arg Pro His Leu  
 980 985 990

Asp Ala Tyr Ser Ser Phe Gly Gln Pro Ser Asp Cys Gln Pro Arg  
 995 1000 1005

Ala Phe Tyr Leu Lys Ala His Ser Ser Pro Asp Leu Asp Ser Gly  
 1010 1015 1020

Ser Glu Glu Asp Gly Lys Glu Arg Thr Asp Phe Gln Glu Glu Asn  
 1025 1030 1035

His Ile Cys Thr Phe Lys Gln Thr Leu Glu Asn Tyr Arg Thr Pro  
 1040 1045 1050

Asn Phe Gln Ser Tyr Asp Leu Asp Thr  
 1055

<210> SEQ ID NO 291  
 <211> LENGTH: 2906  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 291

ggggagagga attgacatg taaaaggaga cttttttttt tgggtggtgt

-continued

---

ggctgttggg tgccttgcaa aaatgaagga tgcaggacgc agctttctcc	100
tggaaccgaa cgcaatggat aaactgattg tgcaagagag aaggaagaac	150
gaagcttttt cttgtgagcc ctggatctta acacaaatgt gtatatgtgc	200
acacagggag cattcaagaa taaaataaac cagagttaga cccgcggggg	250
ttggtgtggt ctgacataaa taaataatct taaagcagct gttcccctcc	300
ccacccccaa aaaaaaggat gattggaaat gaagaaccga ggattcacia	350
agaaaaaagt atgttcattt ttctctataa aggagaaagt gagccaagga	400
gatatttttg gaatgaaaag ttgggggctt ttttagtaaa gtaaagaact	450
ggtgtgtggt tgttttcctt tctttttgaa tttcccacia gaggagagga	500
aattaataat acatctgcaa agaaatttca gagaagaaa gttgaccgcg	550
gcagattgag gcattgattg ggggagagaa accagcagag cacagtggga	600
tttgtgccta tgttgactaa aattgacgga taattgcagt tggatttttc	650
ttcatcaacc tccttttttt taaattttta ttccttttgg tatcaagatc	700
atgctgtttt tcttgttctt aaccacttgg atttccatct ggatgttgct	750
gtgatcagtc tgaatacaaa ctgtttgaat tccagaagga ccaacaccag	800
ataaattatg aatgttgaac aagatgacct tacatccaca gcagataatg	850
ataggtccta ggtttaacag ggcctattt gaccccttgc ttgtggtgct	900
gctggtcctt caacttcttg tgggtgctgg tctggtgctg gctcagacct	950
gcccttctgt gtgctcctgc agcaaccagt tcagcaaggt gatttgtggt	1000
cggaaaaacc tgcgtgaggt tccggatggc atctccacca acacacggct	1050
gctgaacctc catgagaacc aaatccagat catcaaagtg aacagcttca	1100
agcaattgag gcacttggaa atcctacagt tgagtaggaa ccatatcaga	1150
accattgaaa ttggggcttt caatggtctg gcgaacctca aactctgga	1200
actctttgac aatcgtctta ctaccatccc gaattggagct tttgtatact	1250
tgtctaaact gaaggagctc tggttgcgaa acaaccccat tgaagcctc	1300
ccttcttatg cttttaacag aattccttct ttgcgcccac tagacttagg	1350
ggaattgaaa agactttcat acatctcaga aggtgccttt gaaggctctgt	1400
ccaacttgag gtatttgaac cttgccatgt gcaaccttgc gaaatccct	1450
aacctcacac cgctcataaa actagatgag ctggatcttt ctgggaatca	1500
tttatctgcc atcaggcctg gctctttcca gggtttgatg caccttcaaa	1550
aactgtggat gatacagtcc cagattcaag tgattgaacg gaatgccttt	1600
gacaaccttc agtcactagt ggagatcaac ctggcacaca ataactaac	1650
attactgcct catgacctct tcaactccctt gcacatccta gagcggatac	1700
atttacatca caacccttgg aactgtaact gtgacatact gtggctcagc	1750
tggtggataa aagacatggc cccctcgaac acagcttggt gtgcccgtg	1800
taaacctcct cccaatctaa aggggaggta cattggagag ctcgaccaga	1850
attacttcac atgctatgct ccggtgattg tggagcccc tgcagacctc	1900
aatgtcactg aaggcatggc agctgagctg aaatgtcggg cctccacatc	1950

-continued

---

```

cctgacatct gtatcttga ttactccaaa tggaacagtc atgacacatg      2000
ggcgctacaa agtgcgata gctgtgctca gtgatgtac gttaaatttc      2050
acaaatgtaa ctgtgcaaga tacaggcatg tacacatgta tggtagtaa      2100
ttccgttggg aatactactg cttcagccac cctgaatggt actgcagcaa      2150
ccactactoc tttctcttac ttttcaaccg tcacagtaga gactatggaa      2200
ccgtctcagg atgaggcacg gaccacagat aacaatgtgg gtcccactcc      2250
agtggtcgac tgggagacca ccaatgtgac cacctctctc acaccacaga      2300
gcacaaggtc gacagagaaa accttcacca tcccagtgc tgatataaac      2350
agtgggatcc caggaattga tgaggtcatg aagactacca aaatcatcat      2400
tggtgtttt gtggccatca cactcatggc tgcagtgatg ctggtcattt      2450
tctacaagat gaggaagcag caccatcggc aaaacatca cgccccaca      2500
aggactgttg aaattattaa tgtgatgat gagattacgg gagacacacc      2550
catgaaaagc cacctgcca tgctgctat cgagcatgag cacctaaatc      2600
actataactc atacaaatct cccttcaacc acacaacaac agttaacaca      2650
ataaattcaa tacacagttc agtgcataaa cgttattga tccgaatgaa      2700
ctctaaagac aatgtacaag agactcaaat ctaaaacatt tacagagtta      2750
caaaaaacaa acaatcaaaa aaaaagacag tttatataaa atgacacaaa      2800
tgactgggct aaatctactg tttcaaaaaa gtgtctttac aaaaaacaa      2850
aaaaaagaaag aaatttattt attaaaaatt ctattgtgat ctaaagcaga      2900
caaaaaa                                           2906

```

&lt;210&gt; SEQ ID NO 292

&lt;211&gt; LENGTH: 640

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 292

```

Met Leu Asn Lys Met Thr Leu His Pro Gln Gln Ile Met Ile Gly
 1          5          10          15
Pro Arg Phe Asn Arg Ala Leu Phe Asp Pro Leu Leu Val Val Leu
 20          25          30
Leu Ala Leu Gln Leu Leu Val Val Ala Gly Leu Val Arg Ala Gln
 35          40          45
Thr Cys Pro Ser Val Cys Ser Cys Ser Asn Gln Phe Ser Lys Val
 50          55          60
Ile Cys Val Arg Lys Asn Leu Arg Glu Val Pro Asp Gly Ile Ser
 65          70          75
Thr Asn Thr Arg Leu Leu Asn Leu His Glu Asn Gln Ile Gln Ile
 80          85          90
Ile Lys Val Asn Ser Phe Lys His Leu Arg His Leu Glu Ile Leu
 95          100         105
Gln Leu Ser Arg Asn His Ile Arg Thr Ile Glu Ile Gly Ala Phe
 110         115         120
Asn Gly Leu Ala Asn Leu Asn Thr Leu Glu Leu Phe Asp Asn Arg
 125         130         135

```

-continued

Leu	Thr	Thr	Ile	Pro	Asn	Gly	Ala	Phe	Val	Tyr	Leu	Ser	Lys	Leu
				140					145					150
Lys	Glu	Leu	Trp	Leu	Arg	Asn	Asn	Pro	Ile	Glu	Ser	Ile	Pro	Ser
				155					160					165
Tyr	Ala	Phe	Asn	Arg	Ile	Pro	Ser	Leu	Arg	Arg	Leu	Asp	Leu	Gly
				170					175					180
Glu	Leu	Lys	Arg	Leu	Ser	Tyr	Ile	Ser	Glu	Gly	Ala	Phe	Glu	Gly
				185					190					195
Leu	Ser	Asn	Leu	Arg	Tyr	Leu	Asn	Leu	Ala	Met	Cys	Asn	Leu	Arg
				200					205					210
Glu	Ile	Pro	Asn	Leu	Thr	Pro	Leu	Ile	Lys	Leu	Asp	Glu	Leu	Asp
				215					220					225
Leu	Ser	Gly	Asn	His	Leu	Ser	Ala	Ile	Arg	Pro	Gly	Ser	Phe	Gln
				230					235					240
Gly	Leu	Met	His	Leu	Gln	Lys	Leu	Trp	Met	Ile	Gln	Ser	Gln	Ile
				245					250					255
Gln	Val	Ile	Glu	Arg	Asn	Ala	Phe	Asp	Asn	Leu	Gln	Ser	Leu	Val
				260					265					270
Glu	Ile	Asn	Leu	Ala	His	Asn	Asn	Leu	Thr	Leu	Leu	Pro	His	Asp
				275					280					285
Leu	Phe	Thr	Pro	Leu	His	His	Leu	Glu	Arg	Ile	His	Leu	His	His
				290					295					300
Asn	Pro	Trp	Asn	Cys	Asn	Cys	Asp	Ile	Leu	Trp	Leu	Ser	Trp	Trp
				305					310					315
Ile	Lys	Asp	Met	Ala	Pro	Ser	Asn	Thr	Ala	Cys	Cys	Ala	Arg	Cys
				320					325					330
Asn	Thr	Pro	Pro	Asn	Leu	Lys	Gly	Arg	Tyr	Ile	Gly	Glu	Leu	Asp
				335					340					345
Gln	Asn	Tyr	Phe	Thr	Cys	Tyr	Ala	Pro	Val	Ile	Val	Glu	Pro	Pro
				350					355					360
Ala	Asp	Leu	Asn	Val	Thr	Glu	Gly	Met	Ala	Ala	Glu	Leu	Lys	Cys
				365					370					375
Arg	Ala	Ser	Thr	Ser	Leu	Thr	Ser	Val	Ser	Trp	Ile	Thr	Pro	Asn
				380					385					390
Gly	Thr	Val	Met	Thr	His	Gly	Ala	Tyr	Lys	Val	Arg	Ile	Ala	Val
				395					400					405
Leu	Ser	Asp	Gly	Thr	Leu	Asn	Phe	Thr	Asn	Val	Thr	Val	Gln	Asp
				410					415					420
Thr	Gly	Met	Tyr	Thr	Cys	Met	Val	Ser	Asn	Ser	Val	Gly	Asn	Thr
				425					430					435
Thr	Ala	Ser	Ala	Thr	Leu	Asn	Val	Thr	Ala	Ala	Thr	Thr	Thr	Pro
				440					445					450
Phe	Ser	Tyr	Phe	Ser	Thr	Val	Thr	Val	Glu	Thr	Met	Glu	Pro	Ser
				455					460					465
Gln	Asp	Glu	Ala	Arg	Thr	Thr	Asp	Asn	Asn	Val	Gly	Pro	Thr	Pro
				470					475					480
Val	Val	Asp	Trp	Glu	Thr	Thr	Asn	Val	Thr	Thr	Ser	Leu	Thr	Pro
				485					490					495
Gln	Ser	Thr	Arg	Ser	Thr	Glu	Lys	Thr	Phe	Thr	Ile	Pro	Val	Thr
				500					505					510
Asp	Ile	Asn	Ser	Gly	Ile	Pro	Gly	Ile	Asp	Glu	Val	Met	Lys	Thr

-continued

	515		520		525
Thr Lys Ile Ile	Ile Gly Cys Phe Val	Ala Ile Thr Leu Met	Ala		
	530		535		540
Ala Val Met Leu	Val Ile Phe Tyr Lys	Met Arg Lys Gln His	His		
	545		550		555
Arg Gln Asn His	His Ala Pro Thr Arg	Thr Val Glu Ile Ile	Asn		
	560		565		570
Val Asp Asp Glu	Ile Thr Gly Asp Thr	Pro Met Glu Ser His	Leu		
	575		580		585
Pro Met Pro Ala	Ile Glu His Glu His	Leu Asn His Tyr Asn	Ser		
	590		595		600
Tyr Lys Ser Pro	Phe Asn His Thr Thr	Thr Val Asn Thr Ile	Asn		
	605		610		615
Ser Ile His Ser	Ser Val His Glu Pro	Leu Leu Ile Arg Met	Asn		
	620		625		630
Ser Lys Asp Asn	Val Gln Glu Thr Gln	Ile			
	635		640		

<210> SEQ ID NO 293  
 <211> LENGTH: 4053  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 293

```

agccgacgct gctcaagctg caactctgtt gcagttggca gttcttttcg          50
gtttccctcc tgctgttttg gggcatgaaa gggcttcgcc gccgggagta          100
aaagaagaaa ttgaccgggc agcgcgaggg aggagcgcgc acgcgaccgc          150
gagggggggc gtgcaccctc ggctggaagt ttgtgccggg ccccgagcgc          200
gcgcccggct ggagcttcgg gtagagacct aggcgcgtgg accgcgatga          250
gcgcgccgag cctccgtgcg cgcgcgcggg ggttggggct gctgctgtgc          300
gcggtgctgg ggcgcgctgg ccggtccgac agcggcggtc gcggggaact          350
cgggcagccc tctgggtag ccgcccagcg cccatgcccc actacctgcc          400
gctgcctcgg ggacctgctg gactgcagtc gtaagcggct agcgcgtctt          450
cccgagccac tcccgtcctg ggtcgtcctg ctggacttaa gtcacaacag          500
attatctttc atcaaggcaa gttccatgag ccacctcaa agccttcgag          550
aagtgaaact gaacaacaat gaattggaga ccattccaaa tctgggacca          600
gtctcggcaa atattacact tctctccttg gctggaaca ggattgttga          650
aatactccct gaacatctga aagagtttca gtccttgaa actttggacc          700
ttagcagcaa caatatttca gagctccaaa ctgcatttcc agccctacag          750
ctcaaatatc tgtatctcaa cagcaaccga gtcacatcaa tggaacctgg          800
gtattttgac aatttgcca acacactcct tgtgttaaag ctgaacagga          850
accgaatctc agctatccca cccaagatgt ttaaactgcc ccaactgcaa          900
catctcgaat tgaaccgaaa caagattaa aatgtagatg gactgacatt          950
ccaaggcctt ggtgctctga agtctctgaa aatgcaaaga aatggagtaa          1000
cgaaacttat ggatggagct ttttgggggc tgagcaacat ggaatttttg          1050
    
```

-continued

---

cagctggacc ataacaacct aacagagatt accaaaggct ggctttacgg	1100
cttgctgatg ctgcaggaac ttcactctcag ccaaaatgcc atcaacagga	1150
tcagccctga tgccctggag ttctgccaga agctcagtga gctggacctta	1200
actttcaatc acttatcaag gttagatgat tcaagcttcc ttggcctaag	1250
cttactaaat aactgcaca ttgggaacaa cagagtcagc tacattgctg	1300
attgtgcctt ccgggggctt tccagtttaa agactttgga tctgaagaac	1350
aatgaaattt cctggactat tgaagacatg aatggtgctt tctctgggct	1400
tgacaaaactg aggcgactga tactccaagg aaatcgatc cgttctatta	1450
ctaaaaaagc cttcactggt ttggatgcat tggagcatct agacctgagt	1500
gacaacgcaa tcatgtcttt acaaggcaat gcattttcac aaatgaagaa	1550
actgcaacaa ttgcatttaa atacatcaag ccttttgctg gattgccagc	1600
taaaatggct cccacagtgg gtggcggaaa acaactttca gagctttgta	1650
aatgccagtt gtgccatcc tcagctgcta aaaggaagaa gcatttttgc	1700
tgttagccca gatggctttg tgtgtgatga ttttccaaa cccagatca	1750
cggttcagcc agaaacacag tcggcaataa aaggttcaa tttgagttc	1800
atctgctcag ctgccagcag cagtgattcc ccaatgactt ttgottgaa	1850
aaaagacaat gaactactgc atgatgctga aatggaaaat tatgcacacc	1900
tccgggccca aggtggcgag gtgatggagt ataccacat ccttcggctg	1950
cgcgaggtgg aatttgccag tgaggggaaa tatcagtgtg tcatctcaa	2000
tcactttggt tcatcctact ctgtcaaagc caagcttaca gtaaatatgc	2050
ttccctcatt caccaagacc cccatggatc tcaccatccg agctggggcc	2100
atggcacgct tggagtgtgc tgetgtgggg caccagccc cccagatagc	2150
ctggcagaag gatgggggca cagacttccc agctgcacgg gagagacgca	2200
tgcatgtgat gcccgaggat gacgtgttct ttatcgtgga tgtgaagata	2250
gaggacattg ggtatacag ctgcacagct cagaacagtg caggaagtat	2300
ttcagcaaat gcaactctga ctgtcctaga aacacatca tttttgcggc	2350
cactgttggg ccgaactgta accaaggag aaacagccgt cctacagtgc	2400
attgctggag gaagccctcc ccctaaactg aactggacca aagatgatag	2450
cccattggtg gtaaccgaga ggcaactttt tgacagaggc aatcagcttc	2500
tgattattgt ggactcagat gtcagtgatg ctgggaaata cacatgtgag	2550
atgtctaaca cccttgccac tgagagagga aacgtgcgcc tcagtgtgat	2600
ccccactcca acctgcgact cccctcagat gacagcccca tcgttagacg	2650
atgacggatg ggccactgtg ggtgtcgtga tcatagccgt ggtttgctgt	2700
gtggtgggca cgtcactcgt gtgggtggtc atcatatacc acacaaggcg	2750
gaggaaatgaa gattgcagca ttaccaaac acatgagacc aacttgccag	2800
cagatattoc tagttatttg tcatctcagg gaacgttagc tgacaggcag	2850
gatgggtaoc tgtcttcaga aagtggaaag caccaccagt ttgtcacatc	2900
ttcaggtgct ggatttttct taccacaaca tgacagtatg gggacctgcc	2950

-continued

---

```

atattgacaa tagcagtga gctgatgtgg aagctgccac agatctgttc      3000
ctttgtccgt ttttgggatc cacagccct atgtattga agggaaatgt      3050
gtatggctca gatccttttg aaacatatca tacaggttgc agtcctgacc      3100
caagaacagt ttaaatggac cactatgagc ccagttacat aaagaaaaag      3150
gagtgtacc catgttctca tccttcagaa gaatcctgcg aacggagctt      3200
cagtaataata tcgtggcctt cacatgtgag gaagctactt aacactagtt      3250
actctcaca tgaaggacct ggaatgaaaa atctgtgtct aaacaagtcc      3300
tctttagatt ttagtgcaaa tccagagcca gcgtcggttg cctcgagtaa      3350
ttctttcatg ggtacctttg gaaaagctct caggagacct cacctagatg      3400
cctattcaag ctttggacag ccatcagatt gtcagccaag agccttttat      3450
ttgaaagctc attcttccc agacttggac tctgggtcag aggaagatgg      3500
gaaagaaagg acagattttc aggaagaaaa tcacatttgt acctttaaac      3550
agactttaga aaactacag actccaaatt ttcagtctta tgacttggac      3600
acatagactg aatgagacca aaggaaaagc ttaacatact acctcaagtg      3650
aacttttatt taaaagagag agaatcttat gttttttaa tggagttatg      3700
aattttaaaa ggataaaaat gctttattta tacagatgaa ccaaaattac      3750
aaaaagtatt gaaaattttt atactgggaa tgatgctcat ataagaatac      3800
ctttttaaac tattttttaa ctttgtttta tgcaaaaaag tatcttacgt      3850
aaattaatga tataaatcat gattatttta tgtattttta taatgccaga      3900
tttcttttta tggaaaatga gttactaaag cattttaaat aatacctgcc      3950
ttgtaccatt ttttaaatag aagtacttcc attatatttt gcacattata      4000
tttaataaaa tgtgtcaatt tgaaaaaaa aaaaaaaaaa aaaaaaaaaa      4050
aaa                                                                4053
    
```

```

<210> SEQ ID NO 294
<211> LENGTH: 1119
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 294

```

Met Ser Ala Pro Ser Leu Arg Ala Arg Ala Ala Gly Leu Gly Leu
 1           5           10          15
Leu Leu Cys Ala Val Leu Gly Arg Ala Gly Arg Ser Asp Ser Gly
 20          25          30
Gly Arg Gly Glu Leu Gly Gln Pro Ser Gly Val Ala Ala Glu Arg
 35          40          45
Pro Cys Pro Thr Thr Cys Arg Cys Leu Gly Asp Leu Leu Asp Cys
 50          55          60
Ser Arg Lys Arg Leu Ala Arg Leu Pro Glu Pro Leu Pro Ser Trp
 65          70          75
Val Ala Arg Leu Asp Leu Ser His Asn Arg Leu Ser Phe Ile Lys
 80          85          90
Ala Ser Ser Met Ser His Leu Gln Ser Leu Arg Glu Val Lys Leu
 95          100         105
Asn Asn Asn Glu Leu Glu Thr Ile Pro Asn Leu Gly Pro Val Ser
    
```





-continued

---

Phe	Pro	Lys	Pro	Gln	Ile	Thr	Val	Gln	Pro	Glu	Thr	Gln	Ser	Ala
				500					505					510
Ile	Lys	Gly	Ser	Asn	Leu	Ser	Phe	Ile	Cys	Ser	Ala	Ala	Ser	Ser
				515					520					525
Ser	Asp	Ser	Pro	Met	Thr	Phe	Ala	Trp	Lys	Lys	Asp	Asn	Glu	Leu
				530					535					540
Leu	His	Asp	Ala	Glu	Met	Glu	Asn	Tyr	Ala	His	Leu	Arg	Ala	Gln
				545					550					555
Gly	Gly	Glu	Val	Met	Glu	Tyr	Thr	Thr	Ile	Leu	Arg	Leu	Arg	Glu
				560					565					570
Val	Glu	Phe	Ala	Ser	Glu	Gly	Lys	Tyr	Gln	Cys	Val	Ile	Ser	Asn
				575					580					585
His	Phe	Gly	Ser	Ser	Tyr	Ser	Val	Lys	Ala	Lys	Leu	Thr	Val	Asn
				590					595					600
Met	Leu	Pro	Ser	Phe	Thr	Lys	Thr	Pro	Met	Asp	Leu	Thr	Ile	Arg
				605					610					615
Ala	Gly	Ala	Met	Ala	Arg	Leu	Glu	Cys	Ala	Ala	Val	Gly	His	Pro
				620					625					630
Ala	Pro	Gln	Ile	Ala	Trp	Gln	Lys	Asp	Gly	Gly	Thr	Asp	Phe	Pro
				635					640					645
Ala	Ala	Arg	Glu	Arg	Arg	Met	His	Val	Met	Pro	Glu	Asp	Asp	Val
				650					655					660
Phe	Phe	Ile	Val	Asp	Val	Lys	Ile	Glu	Asp	Ile	Gly	Val	Tyr	Ser
				665					670					675
Cys	Thr	Ala	Gln	Asn	Ser	Ala	Gly	Ser	Ile	Ser	Ala	Asn	Ala	Thr
				680					685					690
Leu	Thr	Val	Leu	Glu	Thr	Pro	Ser	Phe	Leu	Arg	Pro	Leu	Leu	Asp
				695					700					705
Arg	Thr	Val	Thr	Lys	Gly	Glu	Thr	Ala	Val	Leu	Gln	Cys	Ile	Ala
				710					715					720
Gly	Gly	Ser	Pro	Pro	Pro	Lys	Leu	Asn	Trp	Thr	Lys	Asp	Asp	Ser
				725					730					735
Pro	Leu	Val	Val	Thr	Glu	Arg	His	Phe	Phe	Ala	Ala	Gly	Asn	Gln
				740					745					750
Leu	Leu	Ile	Ile	Val	Asp	Ser	Asp	Val	Ser	Asp	Ala	Gly	Lys	Tyr
				755					760					765
Thr	Cys	Glu	Met	Ser	Asn	Thr	Leu	Gly	Thr	Glu	Arg	Gly	Asn	Val
				770					775					780
Arg	Leu	Ser	Val	Ile	Pro	Thr	Pro	Thr	Cys	Asp	Ser	Pro	Gln	Met
				785					790					795
Thr	Ala	Pro	Ser	Leu	Asp	Asp	Asp	Gly	Trp	Ala	Thr	Val	Gly	Val
				800					805					810
Val	Ile	Ile	Ala	Val	Val	Cys	Cys	Val	Val	Gly	Thr	Ser	Leu	Val
				815					820					825
Trp	Val	Val	Ile	Ile	Tyr	His	Thr	Arg	Arg	Arg	Asn	Glu	Asp	Cys
				830					835					840
Ser	Ile	Thr	Asn	Thr	Asp	Glu	Thr	Asn	Leu	Pro	Ala	Asp	Ile	Pro
				845					850					855
Ser	Tyr	Leu	Ser	Ser	Gln	Gly	Thr	Leu	Ala	Asp	Arg	Gln	Asp	Gly
				860					865					870

-continued

---

Tyr Val Ser Ser Glu Ser Gly Ser His His Gln Phe Val Thr Ser  
 875 880 885

Ser Gly Ala Gly Phe Phe Leu Pro Gln His Asp Ser Ser Gly Thr  
 890 895 900

Cys His Ile Asp Asn Ser Ser Glu Ala Asp Val Glu Ala Ala Thr  
 905 910 915

Asp Leu Phe Leu Cys Pro Phe Leu Gly Ser Thr Gly Pro Met Tyr  
 920 925 930

Leu Lys Gly Asn Val Tyr Gly Ser Asp Pro Phe Glu Thr Tyr His  
 935 940 945

Thr Gly Cys Ser Pro Asp Pro Arg Thr Val Leu Met Asp His Tyr  
 950 955 960

Glu Pro Ser Tyr Ile Lys Lys Lys Glu Cys Tyr Pro Cys Ser His  
 965 970 975

Pro Ser Glu Glu Ser Cys Glu Arg Ser Phe Ser Asn Ile Ser Trp  
 980 985 990

Pro Ser His Val Arg Lys Leu Leu Asn Thr Ser Tyr Ser His Asn  
 995 1000 1005

Glu Gly Pro Gly Met Lys Asn Leu Cys Leu Asn Lys Ser Ser Leu  
 1010 1015 1020

Asp Phe Ser Ala Asn Pro Glu Pro Ala Ser Val Ala Ser Ser Asn  
 1025 1030 1035

Ser Phe Met Gly Thr Phe Gly Lys Ala Leu Arg Arg Pro His Leu  
 1040 1045 1050

Asp Ala Tyr Ser Ser Phe Gly Gln Pro Ser Asp Cys Gln Pro Arg  
 1055 1060 1065

Ala Phe Tyr Leu Lys Ala His Ser Ser Pro Asp Leu Asp Ser Gly  
 1070 1075 1080

Ser Glu Glu Asp Gly Lys Glu Arg Thr Asp Phe Gln Glu Glu Asn  
 1085 1090 1095

His Ile Cys Thr Phe Lys Gln Thr Leu Glu Asn Tyr Arg Thr Pro  
 1100 1105 1110

Asn Phe Gln Ser Tyr Asp Leu Asp Thr  
 1115

<210> SEQ ID NO 295  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 295

ggaaccgaat ctcagcta

18

<210> SEQ ID NO 296  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 296

cctaaactga actggacca

19

---

-continued

---

<210> SEQ ID NO 297  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 297  
  
ggctggagac actgaacct 19

<210> SEQ ID NO 298  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 298  
  
acagctgcac agctcagaac agtg 24

<210> SEQ ID NO 299  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 299  
  
cattcccagt ataaaaattt tc 22

<210> SEQ ID NO 300  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 300  
  
gggtcttggt gaatgagg 18

<210> SEQ ID NO 301  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 301  
  
gtgcctctcg gttaccacca atgg 24

<210> SEQ ID NO 302  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 302  
  
gcggccactg ttggaccgaa ctgtaaccaa gggagaaaca gccgtcctac 50

<210> SEQ ID NO 303  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

---

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 303

gcctttgaca accttcagtc actagtgg 28

<210> SEQ ID NO 304  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 304

ccccatgtgt ccatgactgt tccc 24

<210> SEQ ID NO 305  
<211> LENGTH: 45  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 305

tactgcctca tgacctcttc actcccttgc atcatccttag agcgg 45

<210> SEQ ID NO 306  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 306

actccaagga aatcgatcc gtcc 24

<210> SEQ ID NO 307  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 307

ttagcagctg aggatgggca caac 24

<210> SEQ ID NO 308  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 308

actccaagga aatcgatcc gtcc 24

<210> SEQ ID NO 309  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 309

-continued

---

```

gccttcactg gtttgatgc attggagcat ctagacctga gtgacaacgc          50

<210> SEQ ID NO 310
<211> LENGTH: 3296
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 310

caaaacttgc gtcgcgaga gcgccagct tgacttgaat ggaaggagcc          50
cgagcccggg gagcgagct gagactggg gagcgcttc gccctgtgg          100
gcgcccctcg gcgcccggg gcagcaggga aggggaagct gtggtctgcc          150
ctgctccacg aggcgccact ggtgtgaacc gggagagccc ctgggtggtc          200
ccgtccccta tccctccttt atatagaaac cttccacact ggaaggcag          250
cggcgagcca ggagggctca tggtagcaa ggaggccggc tgatctgac          300
gcgcacagca ttccgagttt acagattttt acagatacca aatggaaggc          350
gaggaggcag aacagcctgc ctggttccat cagccctggc gcccaggcgc          400
atctgactcg gcaccccctg caggcaccat ggcccagagc cgggtgctgc          450
tgctcctgct gctgctgccg ccacagctgc acctgggacc tgtgcttgc          500
gtgagggccc caggatttgg ccgaagtggc ggccacagcc tgagcccga          550
agagaacgaa tttgcgagg aggagccggt gotggtactg agccctgagg          600
agcccgggccc tggcccagcc gcggtcagct gcccgcgaga ctgtgcctgt          650
tcccaggagg gcgtcgtgga ctgtggcggt attgacctgc gtgagttccc          700
gggggacctg cctgagcaca ccaaccacct atctctgacg aacaaccagc          750
tggaaaagat ctaccctgag gagctctccc ggctgcaccg gctggagaca          800
ctgaacctgc aaaacaaccg cctgacttcc cgagggtccc cagagaaggc          850
gtttgagcat ctgaccaacc tcaattacct gtacttgccc aataacaagc          900
tgaccttggc accccgcttc ctgccaaaag ccctgatcag tgtggacttt          950
gctgccaaat atctcaccaa gatctatggg ctcacctttg gccagaagcc          1000
aaacttgagg tctgtgtacc tgcacaacaa caagctggca gacgccgggc          1050
tgccggacaa catgttcaac ggctccagca acgctcaggt cctcatcctg          1100
tccagcaact tcctgcgcca cgtgcccaag cacctgccgc ctgccctgta          1150
caagctgcac ctcaagaaca acaagctgga gaagatcccc ccgggggcct          1200
tcagcgagct gagcagcctg cgcgagctat acctgcagaa caactacctg          1250
actgacgagg gcctggacaa cgagaccttc tggaaactct ccagcctgga          1300
gtacctggat ctgtccagca acaacctgtc tcgggtccca gctgggctgc          1350
cgcgagcctt ggtgctgctg cacttgagga agaacgcat ccggagcgtg          1400
gacgcgaatg tgctgacccc catccgcagc ctggagtacc tgctgctgca          1450
cagcaaccag ctgcccggagc agggcatcca cccactggcc ttccagggcc          1500
tcaagcgggt gcacacgggt cacctgtaca acaacgcgct ggagcgcgtg          1550
cccagtggcc tgcctcgccg cgtgcgcacc ctcatgatcc tgcacaacca          1600
gatcacaggg attggccgcg aagactttgc caccacctac ttctggagg          1650

```

-continued

---

agctcaacct cagctacaac cgcatacaca gccacacaggt gcaccgcgac	1700
gccttcocga agctgcgcct gctgcgctcg ctggacctgt cgggcaaccg	1750
gctgcacacg ctgccacctg ggctgcctcg aaatgtccat gtgctgaagg	1800
tcaagcgcaa tgagctggct gccttggcac gaggggcgct ggcgggcatg	1850
gctcagctgc gtgagctgta cctcaccagc aaccgactgc gcagccgagc	1900
cctgggcccc cgtgcctggg tggacctcgc ccatctgcag ctgctggaca	1950
tcgocgggaa tcagctcaca gagatccocg aggggctccc cgagtcactt	2000
gagtacctgt acctgcagaa caacaagatt agtgcggtgc ccgccaatgc	2050
cttcgactcc acgcccacc tcaaggggat ctttctcagg tttacaagc	2100
tggctgtggg ctccgtgggt gacagtgcct tccggaggct gaagcacctg	2150
caggtcttgg acattgaagg caacttagag tttggtgaca tttccaagga	2200
ccgtggcccg ttggggaagg aaaaggagga ggaggaagag gaggaggagg	2250
aggaagagga aacaagatag tgacaagggt atgcagatgt gacctaggat	2300
gatggaccgc cggactcttt tctgcagcac acgcctgtgt gctgtgagcc	2350
ccccactctg ccgtgctcac acagacacac ccagctgcac acatgaggca	2400
tcccacatga cacgggctga cacagtctca tatccccacc cettcccacg	2450
gcgtgtccca cggccagaca catgcacaca catcacacc tcaaacacc	2500
agctcagcca cacacaacta ccctccaaac caccacagtc tctgtcacac	2550
ccccactacc gctgccacgc cctctgaatc atgcagggaa gggctctgcc	2600
ctgcctctgg acacacagc acccattccc tcccctgct gacatgtgta	2650
tgcgtatgca tacacaccac acacacacac atgcacaagt catgtgcgaa	2700
cagccctcca aagcctatgc cacagacagc tcttgcccca gccagaatca	2750
gccatagcag ctgcgccgtc gcctgttcca tctgtccgtc cgttccctgg	2800
agaagacaca agggatcca tgctctgtgg ccagggtcct gccaccctct	2850
ggaactcaca aaagctggct tttattcctt tcccatccta tggggacag	2900
agccttcagg actgctggcc tggcctggcc caccctgctc ctccagggtc	2950
tgggcagtca ctctgctaag agtccctccc tgccacgccc tggcaggaca	3000
caggcacttt tccaatgggc aagcccagtg gaggcaggat gggagagccc	3050
cctgggtgct gctggggcct tggggcagga gtgaagcaga ggtgatgggg	3100
ctgggctgag ccagggagga aggacccagc tgcacctagg agacaccttt	3150
gttcttcagg cctgtggggg aagttccggg tgcctttatt ttttattctt	3200
ttctaaggaa aaaaatgata aaaaatctca agctgatttt tctgtgtata	3250
gaaaaactaa tataaaagca ttatccctat ccctgcaaaa aaaaaa	3296

&lt;210&gt; SEQ ID NO 311

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe

&lt;400&gt; SEQUENCE: 311

-continued

---

gcattggccg cgagactttg cc 22

<210> SEQ ID NO 312  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 312

gcggccacgg tccttgaaa tg 22

<210> SEQ ID NO 313  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 313

tggaggagct caacctcagc tacaaccgca tcaccagccc acagg 45

<210> SEQ ID NO 314  
 <211> LENGTH: 3003  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 314

gggagggggc tccgggcgcc gcgcagcaga cctgctccgg ccgcgcgcct 50

cgcccgctgc ctccgggagc ggcagcagta gcccgggcgg cgagggctgg 100

gggttcctcg agactctcag aggggcgcct cccatcgcg cccaccacc 150

caacctgttc ctcgcgcgcc actgcgctgc gcccaggac ccgctgccc 200

acatggattt tctcctggcg ctggtgctgg tatcctcgct ctacctgcag 250

gcggcccgcc agttcgacgg gaggtggccc aggcaaatag tgtcatcgat 300

tggcctatgt cgttatggty ggaggattga ctgctgctgg ggtgggctc 350

gccagtcttg gggacagtgt cagcctgtgt gcccaaccacg atgcaaacat 400

ggtgaatgta tcgggccaac caagtcaag tgtcatcctg gttatgctgg 450

aaaaacctgt aatcaagatc taaatgagty tggcctgaag ccccggccct 500

gtaagcacag gtgcatgaac acttacggca gctacaagty ctactgtctc 550

aacggatata tgctcatgcc ggatggttcc tgcacaagty ccctgacctg 600

ctccatgcca aactgtcagt atggctgtga tggttgtaaa ggacaaatac 650

ggtgccagty cccatcccct ggctgcacc tggctcctga tggggaggacc 700

tgtgtagatg ttgatgaatg tgctacagga agagcctcct gccctagatt 750

taggcaatgt gtcaaacactt ttgggagcta catctgcaag tgtcataaag 800

gcttcgatct catgtatatt ggaggcaaat atcaatgtca tgacatagac 850

gaatgctcac ttggtcagta tcagtgacgc agctttgctc gatgttataa 900

cgtagctggg tcctacaagt gcaaatgtaa agaaggatac cagggtgatg 950

gactgacttg tgtgtatata ccaaaagtta tgattgaacc ttcaggtcca 1000

attcatgtac caaagggaaa tggtagcatt ttaaagggtg acacaggaaa 1050



-continued

---

taataattgg attcctgatg ttggaagtac ttggtggcct ccgaagacac	1100
catatatcc tcctatcatt accaacaggc ctacttctaa gccacaaca	1150
agacctacac caaagccaac accaattcct actccaccac caccaccacc	1200
cctgccaaaca gagctcagaa cacctctacc acctacaacc ccagaaaggc	1250
caaccaccgg actgacaact atagcaccag ctgccagtac acctccagga	1300
gggattacag ttgacaacag ggtacagaca gacctcaga aaccagagg	1350
agatgtgttc agtgttctgg tacacagttg taatthttgac catggacttt	1400
gtggatggat caggagaaaa gacaatgact tgcactggga accaatcagg	1450
gacctcagag gtggacaata tctgacagtg tcggcagcca aagccccagg	1500
gggaaaagct gcacgcttg tgctacctct cggccgcctc atgcattcag	1550
gggacctgtg cctgtcattc aggcacaagg tgacggggct gcaactctggc	1600
acactccagg tgtttgtgag aaaacacggt gccacggag cagccctgtg	1650
gggaagaaat ggtggccatg gctggaggca aacacagatc accttgcgag	1700
gggctgacat caagagcga tcaacaagat gattaaaggg ttggaaaaaa	1750
agatctatga tggaaaatta aaggaactgg gattattgag cctggagaag	1800
agaagactga ggggcaaac attgatggtt ttcaagtata tgaagggtt	1850
gcacagagag ggtggcgacc agctgttctc catatgcact aagaatagaa	1900
caagaggaat ctggcctaga ctagagtata agggagcatt tcttggcag	1950
ggccattggt agaatacttc ataaaaaaag aagtgtgaaa atctcagtat	2000
ctctctctct ttctaaaaaa ttagataaaa atttgtctat ttaagatggt	2050
taaagatggt cttacccaag gaaaagtaac aaattataga atttcccaa	2100
agatgttttg atcctactag tagtatgcag tgaaaatcct tagaactaaa	2150
taatttggag aaggcttaat ttaggcattt ccctcttgac ctctaatgg	2200
agagggattg aaaggggaag agcccaccaa atgctgagct cactgaaata	2250
tctctccctt atggcaatcc tagcagtatt aaagaaaaaa ggaactatt	2300
tattccaaat gagagtatga tggacagata ttttagtata tcagtaatgt	2350
cctagtgtgg cgggtggttt caatgtttct tcatggtaaa ggtataagcc	2400
tttcatttgt tcaatggatg atgtttcaga tttttttttt ttttaagagat	2450
cctcaagga acacagttca gagagattht catcgggtgc attctctctg	2500
cttcgtgtgt gacaagttat cttggctgct gagaaagagt gccctgcccc	2550
acaccggcag acctttcctt cacctcatca gtatgattca gtttctctta	2600
tcaattggag tctcccaggt tccacagaac agtaaatattt tttgaacaat	2650
aggtaacaata gaaggtcttc tgtcatttaa cctggtaaa gcaaggctgg	2700
agggggaaaa taaatcatta agcctttgag taacggcaga atatatggct	2750
gtagatccat ttttaatggt tcatttcctt tatggtcata taactgcaca	2800
gctgaagatg aaaggggaaa ataaatgaaa atthttacttt tcgatgocaa	2850
tgatacattg cactaaactg atggaagaag ttatocaaag tactgtataa	2900
catcttgttt attatttaat gttttctaaa ataaaaaatg ttagtggttt	2950

-continued

tccaaatggc ctaataaaaa caattatttg taaataaaaa cactgttagt 3000

aat 3003

<210> SEQ ID NO 315  
 <211> LENGTH: 509  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 315

Met Asp Phe Leu Leu Ala Leu Val Leu Val Ser Ser Leu Tyr Leu 15  
 1 5 10

Gln Ala Ala Ala Glu Phe Asp Gly Arg Trp Pro Arg Gln Ile Val 30  
 20 25

Ser Ser Ile Gly Leu Cys Arg Tyr Gly Gly Arg Ile Asp Cys Cys 45  
 35 40

Trp Gly Trp Ala Arg Gln Ser Trp Gly Gln Cys Gln Pro Val Cys 60  
 50 55

Gln Pro Arg Cys Lys His Gly Glu Cys Ile Gly Pro Asn Lys Cys 75  
 65 70

Lys Cys His Pro Gly Tyr Ala Gly Lys Thr Cys Asn Gln Asp Leu 90  
 80 85

Asn Glu Cys Gly Leu Lys Pro Arg Pro Cys Lys His Arg Cys Met 105  
 95 100

Asn Thr Tyr Gly Ser Tyr Lys Cys Tyr Cys Leu Asn Gly Tyr Met 120  
 110 115

Leu Met Pro Asp Gly Ser Cys Ser Ser Ala Leu Thr Cys Ser Met 135  
 125 130

Ala Asn Cys Gln Tyr Gly Cys Asp Val Val Lys Gly Gln Ile Arg 150  
 140 145

Cys Gln Cys Pro Ser Pro Gly Leu His Leu Ala Pro Asp Gly Arg 165  
 155 160

Thr Cys Val Asp Val Asp Glu Cys Ala Thr Gly Arg Ala Ser Cys 180  
 170 175

Pro Arg Phe Arg Gln Cys Val Asn Thr Phe Gly Ser Tyr Ile Cys 195  
 185 190

Lys Cys His Lys Gly Phe Asp Leu Met Tyr Ile Gly Gly Lys Tyr 210  
 200 205

Gln Cys His Asp Ile Asp Glu Cys Ser Leu Gly Gln Tyr Gln Cys 225  
 215 220

Ser Ser Phe Ala Arg Cys Tyr Asn Val Arg Gly Ser Tyr Lys Cys 240  
 230 235

Lys Cys Lys Glu Gly Tyr Gln Gly Asp Gly Leu Thr Cys Val Tyr 255  
 245 250

Ile Pro Lys Val Met Ile Glu Pro Ser Gly Pro Ile His Val Pro 270  
 260 265

Lys Gly Asn Gly Thr Ile Leu Lys Gly Asp Thr Gly Asn Asn Asn 285  
 275 280

Trp Ile Pro Asp Val Gly Ser Thr Trp Trp Pro Pro Lys Thr Pro 300  
 290 295

Tyr Ile Pro Pro Ile Ile Thr Asn Arg Pro Thr Ser Lys Pro Thr 315  
 305 310

-continued

---

Thr Arg Pro Thr Pro Lys Pro Thr Pro Ile Pro Thr Pro Pro Pro  
 320 325 330

Pro Pro Pro Leu Pro Thr Glu Leu Arg Thr Pro Leu Pro Pro Thr  
 335 340 345

Thr Pro Glu Arg Pro Thr Thr Gly Leu Thr Thr Ile Ala Pro Ala  
 350 355 360

Ala Ser Thr Pro Pro Gly Gly Ile Thr Val Asp Asn Arg Val Gln  
 365 370 375

Thr Asp Pro Gln Lys Pro Arg Gly Asp Val Phe Ser Val Leu Val  
 380 385 390

His Ser Cys Asn Phe Asp His Gly Leu Cys Gly Trp Ile Arg Glu  
 395 400 405

Lys Asp Asn Asp Leu His Trp Glu Pro Ile Arg Asp Pro Ala Gly  
 410 415 420

Gly Gln Tyr Leu Thr Val Ser Ala Ala Lys Ala Pro Gly Gly Lys  
 425 430 435

Ala Ala Arg Leu Val Leu Pro Leu Gly Arg Leu Met His Ser Gly  
 440 445 450

Asp Leu Cys Leu Ser Phe Arg His Lys Val Thr Gly Leu His Ser  
 455 460 465

Gly Thr Leu Gln Val Phe Val Arg Lys His Gly Ala His Gly Ala  
 470 475 480

Ala Leu Trp Gly Arg Asn Gly Gly His Gly Trp Arg Gln Thr Gln  
 485 490 495

Ile Thr Leu Arg Gly Ala Asp Ile Lys Ser Glu Ser Gln Arg  
 500 505

<210> SEQ ID NO 316  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 316

gatggttcct gctcaagtgc cctg 24

<210> SEQ ID NO 317  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 317

ttgcacttgt aggaccacg tacg 24

<210> SEQ ID NO 318  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 318

ctgatgggag gacctgtgta gatgttgatg aatgtgctac aggaagagcc 50

-continued

&lt;210&gt; SEQ ID NO 319

&lt;211&gt; LENGTH: 2110

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 319

cttctttgaa aaggattatc acctgatcag gttctctctg catttgcccc	50
tttagattgt gaaatgtggc tcaaggtctt cacaactttc ctttcctttg	100
caacaggtgc ttgctcgggg ctgaagggtga cagtgccatc acacactgtc	150
catggcgta gaggtcaggc cctctaccta cccgtccact atggcttcca	200
cactccagca tcagacatcc agatcatatg gctatttgag agaccccaca	250
caatgccaa atacttactg ggctctgtga ataagtctgt ggttcctgac	300
ttggaatacc aacacaagtt caccatgatg ccaccaatg catctctgct	350
tatcaaccca ctgcagttcc ctgatgaagg caattacatc gtgaagggtca	400
acattcaggg aaatggaact ctatctgcca gtcagaagat acaagtcacg	450
gttgatgatc ctgtcacaaa gccagtggtg cagattcatc ctccctctgg	500
ggctgtggag tatgtgggga acatgaccct gacatgccat gtggaagggg	550
gcactcgctt agcttaccaa tggctaaaaa atgggagacc tgtccacacc	600
agctccacct actccttttc tccccaaaac aatacccttc atattgctcc	650
agtaaccaag gaagacattg ggaattacag ctgcctgggtg aggaaccttg	700
tcagtgaatg gaaagtgat atcattatgc ccatcatata ttatggacct	750
tatggacttc aagtgaattc tgataaaggg ctaaaagtag ggggaagtgtt	800
tactgttgac cttggagagg ccatcctatt tgattgttct gctgattctc	850
atccccccaa cacctactcc tggattagga ggactgacaa tactacatat	900
atcattaagc atgggcctcg cttagaagtt gcactctgaga aagtagccca	950
gaagacaatg gactatgtgt gctgtgctta caacaacata accggcaggc	1000
aagatgaaac tcatttcaca gttatcatca cttccgtagg actggagaag	1050
cttgacacaga aaggaaaatc attgtcacct ttagcaagta taactggaat	1100
atcactattt ttgattatat ccatgtgtct tctcttcccta tggaaaaaat	1150
atcaacccta caaagttata aaacagaaac tagaaggcag gccagaaaca	1200
gaatacagga aagctcaaac attttcaggc catgaagatg ctctggatga	1250
cttcggaata tatgaatttg ttgcttttcc agatgtttct ggtgtttcca	1300
ggattccaag caggtctggt ccagcctctg attgtgtatc ggggcaagat	1350
ttgcacagta cagtgtatga agttattcag cacatccctg cccagcagca	1400
agaccatcca gagtgaactt tcatgggcta aacagtacat tcgagtgaaa	1450
ttctgaagaa acattttaag gaaaaacagt ggaaaagtat attaactctg	1500
aatcagtgaa gaaaccagga ccaacacctc ttactcatta ttcotttaca	1550
tgacagaatg aggcatttat gaaaattgaa ctgcaggttt ttcagcatat	1600
acacaatgtc ttgtgcaaca gaaaaacatg ttggggaaat attcctcagt	1650
ggagagtcgt tctcatgctg acggggagaa cgaaagtgac aggggtttcc	1700
tcataagttt tgtatgaaat atctctacaa acctcaatta gttctactct	1750

-continued

---

```

acactttcac tatcatcaac actgagacta tcctgtctca cctacaaatg      1800
tggaactttt acattgttcg atttttcagc agactttggt ttattaaat      1850
tttattagtg ttaagaatgc taaatttatg tttcaatfff atttccaaat      1900
ttctatcttg ttattttgtac aacaaagtaa taaggatggt tgcacaaaa      1950
acaaaaactat gccttctctt ttttttcaat caccagtagt atttttgaga      2000
agacttgtga acacttaagg aaatgactat taaagtctta tttttatfff      2050
tttcaaggaa agatggattc aaataaatta ttctgttttt gcttttaaaa      2100
aaaaaaaaaa      2110
    
```

```

<210> SEQ ID NO 320
<211> LENGTH: 450
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 320

```

Met Trp Leu Lys Val Phe Thr Thr Phe Leu Ser Phe Ala Thr Gly
 1                    5                    10          15
Ala Cys Ser Gly Leu Lys Val Thr Val Pro Ser His Thr Val His
 20                    25                    30
Gly Val Arg Gly Gln Ala Leu Tyr Leu Pro Val His Tyr Gly Phe
 35                    40                    45
His Thr Pro Ala Ser Asp Ile Gln Ile Ile Trp Leu Phe Glu Arg
 50                    55
Pro His Thr Met Pro Lys Tyr Leu Leu Gly Ser Val Asn Lys Ser
 65                    70                    75
Val Val Pro Asp Leu Glu Tyr Gln His Lys Phe Thr Met Met Pro
 80                    85                    90
Pro Asn Ala Ser Leu Leu Ile Asn Pro Leu Gln Phe Pro Asp Glu
 95                    100                   105
Gly Asn Tyr Ile Val Lys Val Asn Ile Gln Gly Asn Gly Thr Leu
 110                   115                   120
Ser Ala Ser Gln Lys Ile Gln Val Thr Val Asp Asp Pro Val Thr
 125                   130                   135
Lys Pro Val Val Gln Ile His Pro Pro Ser Gly Ala Val Glu Tyr
 140                   145                   150
Val Gly Asn Met Thr Leu Thr Cys His Val Glu Gly Gly Thr Arg
 155                   160                   165
Leu Ala Tyr Gln Trp Leu Lys Asn Gly Arg Pro Val His Thr Ser
 170                   175                   180
Ser Thr Tyr Ser Phe Ser Pro Gln Asn Asn Thr Leu His Ile Ala
 185                   190                   195
Pro Val Thr Lys Glu Asp Ile Gly Asn Tyr Ser Cys Leu Val Arg
 200                   205                   210
Asn Pro Val Ser Glu Met Glu Ser Asp Ile Ile Met Pro Ile Ile
 215                   220                   225
Tyr Tyr Gly Pro Tyr Gly Leu Gln Val Asn Ser Asp Lys Gly Leu
 230                   235                   240
Lys Val Gly Glu Val Phe Thr Val Asp Leu Gly Glu Ala Ile Leu
 245                   250                   255
    
```



---

-continued

---

<210> SEQ ID NO 324

<211> LENGTH: 2397

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 324

gcaagcggcg aatggcgcc ctccgggagt cttgcagttc ccctggcagt	50
cctggtgctg ttgctttggg gtgctccctg gacgcacggg cggcggagca	100
acgttcgcgt catcacggac gagaactgga gagaactgct ggaaggagac	150
tggatgatag aattttatgc cccgtggtgc cctgcttctc aaaatcttca	200
accggaatgg gaaagttttg ctgaatgggg agaagatctt gaggttaata	250
ttgcgaaagt agatgtcaca gagcagccag gactgagtgg acggtttatc	300
ataactgtct ttctactat ttatcattgt aaagatggtg aatttaggcg	350
ctatcagggg ccaaggacta agaaggactt cataaacttt ataagtata	400
aagagtggaa gagtattgag cccgtttcat catggtttgg tccaggttct	450
gttctgatga gtagtatgtc agcactcttt cagctatcta tgtggatcac	500
gacgtgccat aactacttta ttgaagacct tggattgcca gtgtggggat	550
catatactgt ttttgcttta gcaactctgt tttccgact gttattagga	600
ctctgtatga tatttgggc agattgcctt tgccttcaa aaaggcgcag	650
accacagcca taccataacc ctcaaaaaa attattatca gaatctgcac	700
aacctttgaa aaaagtggag gaggaacaag aggcggatga agaagatgtt	750
tcagaagaag aagctgaaag taaagaagga acaaaaaaag actttccaca	800
gaatgccata agacaacgct ctctgggtcc atcattggcc acagataaat	850
cctagttaaa ttttatagtt atcttaatat tatgattttg ataaaaacag	900
aagattgatc attttgtttg gtttgaagtg aactgtgact tttttgaata	950
ttgcaggggt cagtctagat tgtcattaaa ttgaagagtc tacattcaga	1000
acataaaagc actaggtata caagtttgaa atatgattta agcacagtat	1050
gatggtttaa atagttctct aatttttgaa aaatcgtgcc aagcaataag	1100
atztatgat atttgtttaa taataaccta tttcaagtct gagttttgaa	1150
aatttacatt tccaagtat tgcattattg aggtatttaa gaagattatt	1200
ttagagaaaa atatttctca tttgatataa tttttctctg tttcactgtg	1250
tgaaaaaaag aagatatttc ccataaatgg gaagtttgcc cattgtctca	1300
agaaatgtgt atttcagtg caatttcgtg gtcttttttag aggtatattc	1350
caaaatttcc ttgtattttt aggttatgca actaataaaa actaccttac	1400
attaattaat tacagttttc tacacatggt aatacaggat atgctactga	1450
tttaggaagt ttttaagttc atggatttct cttgattcca acaaagtttg	1500
attttctctt gtatttttct tacttactat gggttacatt ttttattttt	1550
caaattggat gataatttct tgaaacatt ttttatgttt tagtaaacag	1600
tatttttttg ttgtttcaaa ctgaagtta ctgagagatc catcaaattg	1650
aacaatctgt tgtaatttaa aattttggcc acttttttca gattttacat	1700
cattcttctg gaacttcaac ttgaaattgt ttttttttcc tttttggatg	1750

-continued

---

```

tgaaggtgaa cattcctgat tttgtctga tgtgaaaaag ccttggtatt      1800
ttacatthttg aaaattcaaa gaagcttaat ataaaagttt gcattctact      1850
caggaaaaag catcttcttg tatatgtctt aaatgtattht ttgtcctcat      1900
atacagaaaag ttcttaattg atthttacagt ctgtaatgct tgatgthttta      1950
aaataataac atthtttatat thtttaaaaag acaaacttca tattatcctg      2000
tgthtctthtc tgactggtaa tattgtgtgg gatttcacag gtaaaagtca      2050
gtaggatgga acatthttagt gtatthttac tccttaaaaga gctagaatac      2100
atagthtttca ccttaaaaga agggggaaaaa tcataaatac aatgaatcaa      2150
ctgaccatta cgtagtagac aatthtctgta atgtcccctt cthtctaggc      2200
tctgttgctg tgtgaatcca ttagatthtac agtatcgtaa tatacaagtt      2250
thctthtaaac ccctctcctt tagaatthtaa aatattgtac cattaagag      2300
thtgatgtg taactgtgta tgcccttagaa aaatattccta agcacaaaaa      2350
aaacctthtct aaccttca ttaaagctga aaaaaaaaaa aaaaaaa      2397

```

&lt;210&gt; SEQ ID NO 325

&lt;211&gt; LENGTH: 280

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 325

```

Met Ala Pro Ser Gly Ser Leu Ala Val Pro Leu Ala Val Leu Val
 1          5          10          15
Leu Leu Leu Trp Gly Ala Pro Trp Thr His Gly Arg Arg Ser Asn
          20          25          30
Val Arg Val Ile Thr Asp Glu Asn Trp Arg Glu Leu Leu Glu Gly
          35          40          45
Asp Trp Met Ile Glu Phe Tyr Ala Pro Trp Cys Pro Ala Cys Gln
          50          55          60
Asn Leu Gln Pro Glu Trp Glu Ser Phe Ala Glu Trp Gly Glu Asp
          65          70          75
Leu Glu Val Asn Ile Ala Lys Val Asp Val Thr Glu Gln Pro Gly
          80          85          90
Leu Ser Gly Arg Phe Ile Ile Thr Ala Leu Pro Thr Ile Tyr His
          95          100          105
Cys Lys Asp Gly Glu Phe Arg Arg Tyr Gln Gly Pro Arg Thr Lys
          110          115          120
Lys Asp Phe Ile Asn Phe Ile Ser Asp Lys Glu Trp Lys Ser Ile
          125          130          135
Glu Pro Val Ser Ser Trp Phe Gly Pro Gly Ser Val Leu Met Ser
          140          145          150
Ser Met Ser Ala Leu Phe Gln Leu Ser Met Trp Ile Arg Thr Cys
          155          160          165
His Asn Tyr Phe Ile Glu Asp Leu Gly Leu Pro Val Trp Gly Ser
          170          175          180
Tyr Thr Val Phe Ala Leu Ala Thr Leu Phe Ser Gly Leu Leu Leu
          185          190          195
Gly Leu Cys Met Ile Phe Val Ala Asp Cys Leu Cys Pro Ser Lys
          200          205          210

```



-continued

---

Arg Arg Arg Pro Gln Pro Tyr Pro Tyr Pro Ser Lys Lys Leu Leu  
 215 220 225

Ser Glu Ser Ala Gln Pro Leu Lys Lys Val Glu Glu Glu Gln Glu  
 230 235 240

Ala Asp Glu Glu Asp Val Ser Glu Glu Glu Ala Glu Ser Lys Glu  
 245 250 255

Gly Thr Asn Lys Asp Phe Pro Gln Asn Ala Ile Arg Gln Arg Ser  
 260 265 270

Leu Gly Pro Ser Leu Ala Thr Asp Lys Ser  
 275 280

<210> SEQ ID NO 326  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 326

tgaggtgggc aagcggcgaa atg 23

<210> SEQ ID NO 327  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 327

tatgtggatc aggacgtgcc 20

<210> SEQ ID NO 328  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 328

tgcagggttc agtctagatt g 21

<210> SEQ ID NO 329  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 329

ttgaaggaca aaggcaatct gccac 25

<210> SEQ ID NO 330  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 330

ggagctctgc agttccoctg gcagtcctgg tgctgttgct ttggg 45

---

-continued

---

<210> SEQ ID NO 331

<211> LENGTH: 2168

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 331

```
gcgagtgtcc agctgctggag acccgtgata attcgttaac taattcaaca      50
aacgggaccc ttctgtgtgc cagaaaccgc aagcagttgc taaccagtg      100
ggacaggcgg attggaagag cgggaaggtc ctggcccaga gcagtggtgac      150
acttccctct gtgaccatga aactctgggt gtctgcattg ctgatggcct      200
ggtttgtgtg cctgagctgt gtgcaggcgg aattcttcac ctctattggg      250
cacatgactg acctgattta tgcagagaaa gagctggtgc agtctctgaa      300
agagtacatc cttgtggagg aagccaagct ttccaagatt aagagctggg      350
ccaacaaaaa ggaagccttg actagcaagt cagctgctga tgctgagggc      400
tacctggctc accctgtgaa tgctacaaa ctggtgaagc ggctaaacac      450
agactggcct gcgctggagg accttgtcct gcaggactca gctgcaggtt      500
ttatcgccaa cctctctgtg cagcggcagt tcttcccac tgatgaggac      550
gagataggag ctgccaagc cctgatgaga cttcaggaca catacaggct      600
ggaccacagg acaatttcca gaggggaact tccaggaacc aagtaccagg      650
caatgctgag tgtggatgac tgctttggga tgggccgctc ggctacaat      700
gaaggggaat attatcatac ggtgtttgtg atggagcagg tgctaaagca      750
gcttgatgcc ggggaggagg ccaccacaac caagtcacag gtgctggact      800
acctcageta tgctgtcttc cagttgggtg atctgcaccg tgccctggag      850
ctcaccgcgc gcctgctctc ccttgaccca agccacgaac gagctggagg      900
gaaatctgcy tactttgagc agttattgga ggaagagaga gaaaaaacgt      950
taacaaatca gacagaagct gagctagcaa cccagaagg catctatgag      1000
aggcctgtgg actacctgcc tgagagggat gtttacgaga gcctctgtcg      1050
tggggagggt gtcaaaactga cccccgtag acagaagagg cttttctgta      1100
ggtaccacca tggcaacagc gcccacagc tgctcattgc ccccttcaa      1150
gaggaggacg agtgggacag cccgcacatc gtcaggact acgatgtcat      1200
gtctgatgag gaaatcgaga ggatcaagga gatcgcaaaa cctaaacttg      1250
cacgagccac cgttcgtgat cccaagacag gagtctctac tgcgccagc      1300
taccgggttt ccaaaagctc ctggctagag gaagatgatg accctgttgt      1350
ggcccagata aatcgctcga tgcagcatat cacagggtta acagtaaaga      1400
ctgcagaatt gttacaggtt gcaaattatg gagtgggagg acagtatgaa      1450
ccgcacttgc acttctctag gcgacctttt gacagcggcc tcaaaacaga      1500
ggggaatagg ttagcgacgt ttcttaacta catgagtgat gtagaagctg      1550
gtggtgccac cgtcttccct gatctggggg ctgcaatttg gcctaagaag      1600
ggtacagctg tgttctggta caacctcttg cggagcgggg aaggtgacta      1650
ccgaacaaga catgctgcct gcctgtgct tgtgggctgc aagtgggtct      1700
```

-continued

---

```

ccaataagtg gttccatgaa cgaggacagg agttcttgag accttgtgga      1750
tcaacagaag ttgactgaca tccttttctg tccttccoct tcctggctct      1800
tcagcccattg tcaacgtgac agacaccttt gtatgttctt ttgtatgttc      1850
ctatcaggct gatttttggg gaaatgaatg tttgtctgga gcagagggag      1900
accatactag ggcgactcct gtgtgactga agtcccagcc cttccattca      1950
gctgtgcca tcctggccc caaggctagg atcaaagtgg ctgcagcaga      2000
gttagctgtc tagcgcctag caaggtgcct ttgtacctca ggtgttttag      2050
gtgtgagatg tttcagttaa ccaaagtctt gataccttgt ttacatgttt      2100
gtttttatgg catttctatc tattgtggct ttaccaaaaa ataaaatgtc      2150
cctaccagaa aaaaaaaaaa      2168

```

&lt;210&gt; SEQ ID NO 332

&lt;211&gt; LENGTH: 533

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 332

```

Met Lys Leu Trp Val Ser Ala Leu Leu Met Ala Trp Phe Gly Val
 1          5          10          15
Leu Ser Cys Val Gln Ala Glu Phe Phe Thr Ser Ile Gly His Met
 20          25          30
Thr Asp Leu Ile Tyr Ala Glu Lys Glu Leu Val Gln Ser Leu Lys
 35          40          45
Glu Tyr Ile Leu Val Glu Glu Ala Lys Leu Ser Lys Ile Lys Ser
 50          55          60
Trp Ala Asn Lys Met Glu Ala Leu Thr Ser Lys Ser Ala Ala Asp
 65          70          75
Ala Glu Gly Tyr Leu Ala His Pro Val Asn Ala Tyr Lys Leu Val
 80          85          90
Lys Arg Leu Asn Thr Asp Trp Pro Ala Leu Glu Asp Leu Val Leu
 95          100         105
Gln Asp Ser Ala Ala Gly Phe Ile Ala Asn Leu Ser Val Gln Arg
 110         115         120
Gln Phe Phe Pro Thr Asp Glu Asp Glu Ile Gly Ala Ala Lys Ala
 125         130         135
Leu Met Arg Leu Gln Asp Thr Tyr Arg Leu Asp Pro Gly Thr Ile
 140         145         150
Ser Arg Gly Glu Leu Pro Gly Thr Lys Tyr Gln Ala Met Leu Ser
 155         160         165
Val Asp Asp Cys Phe Gly Met Gly Arg Ser Ala Tyr Asn Glu Gly
 170         175         180
Asp Tyr Tyr His Thr Val Leu Trp Met Glu Gln Val Leu Lys Gln
 185         190         195
Leu Asp Ala Gly Glu Glu Ala Thr Thr Thr Lys Ser Gln Val Leu
 200         205         210
Asp Tyr Leu Ser Tyr Ala Val Phe Gln Leu Gly Asp Leu His Arg
 215         220         225
Ala Leu Glu Leu Thr Arg Arg Leu Leu Ser Leu Asp Pro Ser His
 230         235         240

```

-continued

---

Glu Arg Ala Gly Gly Asn Leu Arg Tyr Phe Glu Gln Leu Leu Glu  
 245 250 255  
 Glu Glu Arg Glu Lys Thr Leu Thr Asn Gln Thr Glu Ala Glu Leu  
 260 265 270  
 Ala Thr Pro Glu Gly Ile Tyr Glu Arg Pro Val Asp Tyr Leu Pro  
 275 280 285  
 Glu Arg Asp Val Tyr Glu Ser Leu Cys Arg Gly Glu Gly Val Lys  
 290 295 300  
 Leu Thr Pro Arg Arg Gln Lys Arg Leu Phe Cys Arg Tyr His His  
 305 310 315  
 Gly Asn Arg Ala Pro Gln Leu Leu Ile Ala Pro Phe Lys Glu Glu  
 320 325 330  
 Asp Glu Trp Asp Ser Pro His Ile Val Arg Tyr Tyr Asp Val Met  
 335 340 345  
 Ser Asp Glu Glu Ile Glu Arg Ile Lys Glu Ile Ala Lys Pro Lys  
 350 355 360  
 Leu Ala Arg Ala Thr Val Arg Asp Pro Lys Thr Gly Val Leu Thr  
 365 370 375  
 Val Ala Ser Tyr Arg Val Ser Lys Ser Ser Trp Leu Glu Glu Asp  
 380 385 390  
 Asp Asp Pro Val Val Ala Arg Val Asn Arg Arg Met Gln His Ile  
 395 400 405  
 Thr Gly Leu Thr Val Lys Thr Ala Glu Leu Leu Gln Val Ala Asn  
 410 415 420  
 Tyr Gly Val Gly Gly Gln Tyr Glu Pro His Phe Asp Phe Ser Arg  
 425 430 435  
 Arg Pro Phe Asp Ser Gly Leu Lys Thr Glu Gly Asn Arg Leu Ala  
 440 445 450  
 Thr Phe Leu Asn Tyr Met Ser Asp Val Glu Ala Gly Gly Ala Thr  
 455 460 465  
 Val Phe Pro Asp Leu Gly Ala Ala Ile Trp Pro Lys Lys Gly Thr  
 470 475 480  
 Ala Val Phe Trp Tyr Asn Leu Leu Arg Ser Gly Glu Gly Asp Tyr  
 485 490 495  
 Arg Thr Arg His Ala Ala Cys Pro Val Leu Val Gly Cys Lys Trp  
 500 505 510  
 Val Ser Asn Lys Trp Phe His Glu Arg Gly Gln Glu Phe Leu Arg  
 515 520 525  
 Pro Cys Gly Ser Thr Glu Val Asp  
 530

&lt;210&gt; SEQ ID NO 333

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic Oligonucleotide Probe

&lt;400&gt; SEQUENCE: 333

ccaggcaciaa ttccaga

18

&lt;210&gt; SEQ ID NO 334

&lt;211&gt; LENGTH: 19

&lt;212&gt; TYPE: DNA

-continued

---

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 334
ggacccttct gtgtgccag 19

<210> SEQ ID NO 335
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 335
ggtctcaaga actcctgtc 19

<210> SEQ ID NO 336
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 336
acactcagca ttgctggta cttg 24

<210> SEQ ID NO 337
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 337
gggcacatga ctgacctgat ttatgcagag aaagagctgg tgcag 45

<210> SEQ ID NO 338
<211> LENGTH: 2789
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 338
gcagtattga gttttacttc ctctctttt tagtgaaga cagaccataa 50
tcccagtggt agtgaaattg attgtttcat ttattaccgt ttgggtggg 100
ggttagttcc gacaccttca cagttgaaga gcaggcagaa ggagttgtga 150
agacaggaca atcttcttgg ggatgctggt cctggaagcc agcgggcctt 200
getctgtctt tggcctcatt gaccccaggt tctctggtta aaactgaaag 250
cctactactg gcctggtgcc catcaatcca ttgatccttg aggctgtgcc 300
cctggggcac ccacctggca gggcctacca coatgcgact gagtcctctg 350
ttggctctgc tgcggccagc gcttcccctc atcttagggc tgtctctggg 400
gtgcagcctg agcctctctg gggtttcctg gatccagggg gagggagaag 450
atccctgtgt cgaggctgta ggggagcgag gagggccaca gaatccagat 500
tcgagagctc ggctagacca aagtgatgaa gacttcaaac cccggattgt 550
ccctactac agggacccca acaagccta caagaagtg ctcaggactc 600

```

-continued

ggtacatcca gacagagctg ggctcccgtg agcgggttct ggtggctgtc	650
ctgacctccc gagctacact gtccactttg gccgtggctg tgaaccgtac	700
ggtggcccat cacttccctc ggttactcta cttcactggg cagcgggggg	750
cccgggctcc agcagggatg caggtggtgt ctcatgggga tgagcggccc	800
gcttggtcca tgtcagagac cctgcgccac cttcacacac actttggggc	850
cgactacgac tggttcttca tcatgcagga tgacacatat gtgcaggccc	900
cccgcctggc agcccttctg gcccacctca gcataacca agacctgtac	950
ttaggccggg cagaggagtt cattggcgca ggcgagcagg cccggtactg	1000
tcatgggggc tttggctacc tgttgtcagc gagtctctct cttcgtctgc	1050
ggccacatct ggatggctgc cgaggagaca ttctcagtgc cctcctgac	1100
gagtggcttg gacgctgcct cattgactct ctgggcgtcg gctgtgtctc	1150
acagaccagc gggcagcagt atcgtctatt tgaactggcc aaaaataggg	1200
accctgagaa ggaagggagc tcggctttcc tgagtgcctt cgcctgacac	1250
cctgtctcgc aaggtaccct catgtaccgg ctccacaaac gcttcagcgc	1300
tctggagttg gagcgggctt acagtgaaat agaacaactg caggctcaga	1350
tccgaaactc gaccgtgctg acccccgaag gggaggcagg gctgagctgg	1400
cccgttgggc tccctgctcc ttccacacca cactctcgtc ttgaggtgct	1450
gggtgaggac tacttcacag agcagcacac cttctcctgt gcagatgggg	1500
ctccaaagtc cccactacag ggggctagca gggcggacgt gggtgatgctg	1550
ttggagactg ccctggagca gctcaatcgg cgctatcagc cccgcctgctg	1600
cttcagaagc cagcagactgc tcaacggcta tcggcgcttc gaccagcac	1650
ggggcatgga gtacaccctg gaactgctgt tggaaatgtg gacacagcgt	1700
gggcaccggc gggccctggc tcgcagggtc agcctgctgc ggccactgag	1750
ccgggtgtaa atcctacctc tgccctatgt cactgaggcc acccgagtgc	1800
agctggtgct gccactcctg gtggctgaag ctgctgcagc cccggctttc	1850
ctcagagcgt ttgcagccaa tctcctggag ccacgagaac atgcattgct	1900
cacctgttg ctggtctacg ggccacgaga aggtggccgt ggagctccag	1950
accatttct tgggggtgaag gctgcagcag cggagttaga gcgacggtac	2000
cctgggacga ggtggcctg gctcgtgtg cgagcagagg ccccttccca	2050
ggtgcgactc atggacgtgg tctcgaagaa gcaccctgtg gacactctct	2100
tcttccttac caccgtgtgg acaaggcctg ggccogaagt cctcaaccgc	2150
tgctcagatg atgcatctc tggctggcag gccttctttc cagtccattt	2200
ccagagttc aatcctgccc tgtcaccaca gagatcacc ccagggcccc	2250
cgggggctg ccctgacccc ccctcccctc ctggtgctga cccctcccgg	2300
ggggctccta taggggggag atttgaccgg caggcttctg cggagggctg	2350
cttctacaac gctgactacc tggcgcccc agcccggctg gcagggtgac	2400
tggcaggcca ggaagaggag gaagccctgg aggggctgga ggtgatggat	2450
gtttctctcc ggttctcagg gctccacctc tttcgggccc tagagccagg	2500

-continued

---

```

gctggtgcag aagttctccc tgcgagactg cagcccacgg ctcaagtgaag      2550
aactctacca ccgctgcccgc ctcaagcaacc tggagggggct agggggccgt      2600
gcccagctgg ctatggctct ctttgagcag gagcaggcca atagcactta      2650
gcccgcctgg gggccctaac ctattacct ttcctttgtc tgcctcagcc      2700
ccaggaaggg caaggcaaga tggtgacag atagagaatt gttgctgtat      2750
tttttaaata tgaaaatggt attaaacatg tcttctgcc      2789

```

&lt;210&gt; SEQ ID NO 339

&lt;211&gt; LENGTH: 772

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 339

```

Met Arg Leu Ser Ser Leu Leu Ala Leu Leu Arg Pro Ala Leu Pro
 1           5           10           15
Leu Ile Leu Gly Leu Ser Leu Gly Cys Ser Leu Ser Leu Leu Arg
          20           25           30
Val Ser Trp Ile Gln Gly Glu Gly Glu Asp Pro Cys Val Glu Ala
          35           40           45
Val Gly Glu Arg Gly Gly Pro Gln Asn Pro Asp Ser Arg Ala Arg
          50           55           60
Leu Asp Gln Ser Asp Glu Asp Phe Lys Pro Arg Ile Val Pro Tyr
          65           70           75
Tyr Arg Asp Pro Asn Lys Pro Tyr Lys Lys Val Leu Arg Thr Arg
          80           85           90
Tyr Ile Gln Thr Glu Leu Gly Ser Arg Glu Arg Leu Leu Val Ala
          95           100          105
Val Leu Thr Ser Arg Ala Thr Leu Ser Thr Leu Ala Val Ala Val
          110          115          120
Asn Arg Thr Val Ala His His Phe Pro Arg Leu Leu Tyr Phe Thr
          125          130          135
Gly Gln Arg Gly Ala Arg Ala Pro Ala Gly Met Gln Val Val Ser
          140          145          150
His Gly Asp Glu Arg Pro Ala Trp Leu Met Ser Glu Thr Leu Arg
          155          160          165
His Leu His Thr His Phe Gly Ala Asp Tyr Asp Trp Phe Phe Ile
          170          175          180
Met Gln Asp Asp Thr Tyr Val Gln Ala Pro Arg Leu Ala Ala Leu
          185          190          195
Ala Gly His Leu Ser Ile Asn Gln Asp Leu Tyr Leu Gly Arg Ala
          200          205          210
Glu Glu Phe Ile Gly Ala Gly Glu Gln Ala Arg Tyr Cys His Gly
          215          220          225
Gly Phe Gly Tyr Leu Leu Ser Arg Ser Leu Leu Leu Arg Leu Arg
          230          235          240
Pro His Leu Asp Gly Cys Arg Gly Asp Ile Leu Ser Ala Arg Pro
          245          250          255
Asp Glu Trp Leu Gly Arg Cys Leu Ile Asp Ser Leu Gly Val Gly
          260          265          270
Cys Val Ser Gln His Gln Gly Gln Gln Tyr Arg Ser Phe Glu Leu
          275          280          285

```

-continued

---

Ala	Lys	Asn	Arg	Asp	Pro	Glu	Lys	Glu	Gly	Ser	Ser	Ala	Phe	Leu
				290					295					300
Ser	Ala	Phe	Ala	Val	His	Pro	Val	Ser	Glu	Gly	Thr	Leu	Met	Tyr
				305					310					315
Arg	Leu	His	Lys	Arg	Phe	Ser	Ala	Leu	Glu	Leu	Glu	Arg	Ala	Tyr
				320					325					330
Ser	Glu	Ile	Glu	Gln	Leu	Gln	Ala	Gln	Ile	Arg	Asn	Leu	Thr	Val
				335					340					345
Leu	Thr	Pro	Glu	Gly	Glu	Ala	Gly	Leu	Ser	Trp	Pro	Val	Gly	Leu
				350					355					360
Pro	Ala	Pro	Phe	Thr	Pro	His	Ser	Arg	Phe	Glu	Val	Leu	Gly	Trp
				365					370					375
Asp	Tyr	Phe	Thr	Glu	Gln	His	Thr	Phe	Ser	Cys	Ala	Asp	Gly	Ala
				380					385					390
Pro	Lys	Cys	Pro	Leu	Gln	Gly	Ala	Ser	Arg	Ala	Asp	Val	Gly	Asp
				395					400					405
Ala	Leu	Glu	Thr	Ala	Leu	Glu	Gln	Leu	Asn	Arg	Arg	Tyr	Gln	Pro
				410					415					420
Arg	Leu	Arg	Phe	Gln	Lys	Gln	Arg	Leu	Leu	Asn	Gly	Tyr	Arg	Arg
				425					430					435
Phe	Asp	Pro	Ala	Arg	Gly	Met	Glu	Tyr	Thr	Leu	Asp	Leu	Leu	Leu
				440					445					450
Glu	Cys	Val	Thr	Gln	Arg	Gly	His	Arg	Arg	Ala	Leu	Ala	Arg	Arg
				455					460					465
Val	Ser	Leu	Leu	Arg	Pro	Leu	Ser	Arg	Val	Glu	Ile	Leu	Pro	Met
				470					475					480
Pro	Tyr	Val	Thr	Glu	Ala	Thr	Arg	Val	Gln	Leu	Val	Leu	Pro	Leu
				485					490					495
Leu	Val	Ala	Glu	Ala	Ala	Ala	Ala	Pro	Ala	Phe	Leu	Glu	Ala	Phe
				500					505					510
Ala	Ala	Asn	Val	Leu	Glu	Pro	Arg	Glu	His	Ala	Leu	Leu	Thr	Leu
				515					520					525
Leu	Leu	Val	Tyr	Gly	Pro	Arg	Glu	Gly	Gly	Arg	Gly	Ala	Pro	Asp
				530					535					540
Pro	Phe	Leu	Gly	Val	Lys	Ala	Ala	Ala	Ala	Glu	Leu	Glu	Arg	Arg
				545					550					555
Tyr	Pro	Gly	Thr	Arg	Leu	Ala	Trp	Leu	Ala	Val	Arg	Ala	Glu	Ala
				560					565					570
Pro	Ser	Gln	Val	Arg	Leu	Met	Asp	Val	Val	Ser	Lys	Lys	His	Pro
				575					580					585
Val	Asp	Thr	Leu	Phe	Phe	Leu	Thr	Thr	Val	Trp	Thr	Arg	Pro	Gly
				590					595					600
Pro	Glu	Val	Leu	Asn	Arg	Cys	Arg	Met	Asn	Ala	Ile	Ser	Gly	Trp
				605					610					615
Gln	Ala	Phe	Phe	Pro	Val	His	Phe	Gln	Glu	Phe	Asn	Pro	Ala	Leu
				620					625					630
Ser	Pro	Gln	Arg	Ser	Pro	Pro	Gly	Pro	Pro	Gly	Ala	Gly	Pro	Asp
				635					640					645
Pro	Pro	Ser	Pro	Pro	Gly	Ala	Asp	Pro	Ser	Arg	Gly	Ala	Pro	Ile
				650					655					660



-continued

Gly	Gly	Arg	Phe	Asp	Arg	Gln	Ala	Ser	Ala	Glu	Gly	Cys	Phe	Tyr
				665					670					675
Asn	Ala	Asp	Tyr	Leu	Ala	Ala	Arg	Ala	Arg	Leu	Ala	Gly	Glu	Leu
				680					685					690
Ala	Gly	Gln	Glu	Glu	Glu	Glu	Ala	Leu	Glu	Gly	Leu	Glu	Val	Met
				695					700					705
Asp	Val	Phe	Leu	Arg	Phe	Ser	Gly	Leu	His	Leu	Phe	Arg	Ala	Val
				710					715					720
Glu	Pro	Gly	Leu	Val	Gln	Lys	Phe	Ser	Leu	Arg	Asp	Cys	Ser	Pro
				725					730					735
Arg	Leu	Ser	Glu	Glu	Leu	Tyr	His	Arg	Cys	Arg	Leu	Ser	Asn	Leu
				740					745					750
Glu	Gly	Leu	Gly	Gly	Arg	Ala	Gln	Leu	Ala	Met	Ala	Leu	Phe	Glu
				755					760					765
Gln	Glu	Gln	Ala	Asn	Ser	Thr								
				770										

<210> SEQ ID NO 340  
 <211> LENGTH: 1572  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 340

cggagtggtg cgccaacgtg agaggaacc cgtgcgcggc tgcgctttcc	50
tgcccccaag ccgttctaga cgcgggaaaa atgctttctg aaagcagctc	100
ctttttgaag ggtgtgatgc ttggaagcat tttctgtgct ttgatcacta	150
tgctaggaca cattaggatt ggtcatgaa atagaatgca ccaccatgag	200
catcatcaacc tacaagctcc taacaaagaa gatatcttga aaatttcaga	250
ggatgagcgc atggagctca gtaagagctt tcgagtatac tgtattatcc	300
ttgtaaaacc caaagatgtg agtctttggg ctgcagtaaa ggagacttgg	350
accaaacct gtgacaaagc agagttcttc agttctgaaa atgttaaagt	400
gtttgagtca attaatatgg acacaaatga catgtgggta atgatgagaa	450
aagcttacia atacgccttt gataagtata gagaccaata caactggttc	500
ttccttgcac gcccactac gtttgctatc attgaaaacc taaagtattt	550
tttgtaaaa aaggatccat cacagccttt ctatctaggc cacactataa	600
aatctggaga ccttgaatat gtgggtatgg aaggaggaat tgtcttaagt	650
gtagaatcaa tgaaaagact taacagcctt ctcaatatcc cagaaaagtg	700
tcctgaacag ggagggatga ttggaagat atctgaagat aaacagctag	750
cagtttgctt gaaatatgct ggagtatttg cagaaaatgc agaagatgct	800
gatggaaaag atgtatttaa taccaaatct gttgggcttt ctattaaaga	850
ggcaatgact tatcacccca accaggtagt agaaggctgt tgttcagata	900
tggtggttac ttttaatgga ctgactccaa atcagatgca tgtgatgatg	950
tatgggggat accgccttag ggcatttggg catattttca atgatgcatt	1000
ggttttctta cctccaaatg gttctgacaa tgactgagaa gtggtagaaa	1050
agcgtgaata tgatctttgt ataggacgtg tgttgcatt atttgtagta	1100

-continued

---

```

gtaactacat atccaataca gctgtatggt tctttttcct ttctaatttg      1150
gtggcactgg tataaccaca cattaagtc agtagtacat ttttaatga        1200
gggtggtttt tttctttaa acacatgaac attgtaaag tggtgaaag        1250
aagtgtttta agaataataa ttttgcaaat aaactattaa taaatattat      1300
atgtgataaa ttctaaatta tgaacattag aaatctgtgg ggcacatatt      1350
tttgctgatt ggttaaaaaa ttttaacagg tctttagcgt tctaagatat      1400
gcaaatgata tctctagttg tgaattgtg attaaagtaa aacttttagc      1450
tgtgtgttcc ctttacttct aatactgatt tatgttctaa gcctccccaa     1500
gttccaatgg atttgcttc tcaaaatgta caactaagca actaaagaaa     1550
attaaagtga aagttgaaaa at              1572
    
```

```

<210> SEQ ID NO 341
<211> LENGTH: 318
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 341

```

Met Leu Ser Glu Ser Ser Ser Phe Leu Lys Gly Val Met Leu Gly
 1              5              10              15
Ser Ile Phe Cys Ala Leu Ile Thr Met Leu Gly His Ile Arg Ile
                20              25              30
Gly His Gly Asn Arg Met His His His Glu His His His Leu Gln
                35              40              45
Ala Pro Asn Lys Glu Asp Ile Leu Lys Ile Ser Glu Asp Glu Arg
                50              55              60
Met Glu Leu Ser Lys Ser Phe Arg Val Tyr Cys Ile Ile Leu Val
                65              70              75
Lys Pro Lys Asp Val Ser Leu Trp Ala Ala Val Lys Glu Thr Trp
                80              85              90
Thr Lys His Cys Asp Lys Ala Glu Phe Phe Ser Ser Glu Asn Val
                95              100             105
Lys Val Phe Glu Ser Ile Asn Met Asp Thr Asn Asp Met Trp Leu
                110             115             120
Met Met Arg Lys Ala Tyr Lys Tyr Ala Phe Asp Lys Tyr Arg Asp
                125             130             135
Gln Tyr Asn Trp Phe Phe Leu Ala Arg Pro Thr Thr Phe Ala Ile
                140             145             150
Ile Glu Asn Leu Lys Tyr Phe Leu Leu Lys Lys Asp Pro Ser Gln
                155             160             165
Pro Phe Tyr Leu Gly His Thr Ile Lys Ser Gly Asp Leu Glu Tyr
                170             175             180
Val Gly Met Glu Gly Gly Ile Val Leu Ser Val Glu Ser Met Lys
                185             190             195
Arg Leu Asn Ser Leu Leu Asn Ile Pro Glu Lys Cys Pro Glu Gln
                200             205             210
Gly Gly Met Ile Trp Lys Ile Ser Glu Asp Lys Gln Leu Ala Val
                215             220             225
Cys Leu Lys Tyr Ala Gly Val Phe Ala Glu Asn Ala Glu Asp Ala
                230             235             240
    
```

-continued

---

Asp Gly Lys Asp Val Phe Asn Thr Lys Ser Val Gly Leu Ser Ile  
245 250 255

Lys Glu Ala Met Thr Tyr His Pro Asn Gln Val Val Glu Gly Cys  
260 265 270

Cys Ser Asp Met Ala Val Thr Phe Asn Gly Leu Thr Pro Asn Gln  
275 280 285

Met His Val Met Met Tyr Gly Val Tyr Arg Leu Arg Ala Phe Gly  
290 295 300

His Ile Phe Asn Asp Ala Leu Val Phe Leu Pro Pro Asn Gly Ser  
305 310 315

Asp Asn Asp

<210> SEQ ID NO 342  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 342

tccccaagcc gttctagacg cgg 23

<210> SEQ ID NO 343  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 343

ctggttcttc cttgcacg 18

<210> SEQ ID NO 344  
<211> LENGTH: 28  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 344

gcccaaatgc cctaaggcgg tatacccc 28

<210> SEQ ID NO 345  
<211> LENGTH: 50  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 345

gggtgtgatg cttggaagca ttttctgtgc tttgatcact atgctaggac 50

<210> SEQ ID NO 346  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 346

gggatgcagg tggtgtctca tgggg 25

---

-continued

---

<210> SEQ ID NO 347  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 347

ccctcatgta cggctcc 18

<210> SEQ ID NO 348  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 348

ggattcta at acgactcact atagggctca gaaaagcgca acagagaa 48

<210> SEQ ID NO 349  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 349

ctatgaaatt aaccctcact aaagggatgt cttccatgcc aaccttc 47

<210> SEQ ID NO 350  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 350

ggattcta at acgactcact atagggcgcc gatgtccact ggggctac 48

<210> SEQ ID NO 351  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 351

ctatgaaatt aaccctcact aaagggacga ggaagatggg cggatggt 48

<210> SEQ ID NO 352  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 352

ggattcta at acgactcact atagggcacc cagcgtccg gctgctt 47

<210> SEQ ID NO 353  
<211> LENGTH: 48

---

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 353  
  
ctatgaaatt aaccctcact aaagggacgg gggacaccac ggaccaga 48  
  
<210> SEQ ID NO 354  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 354  
  
ggattctaatacgcactcact atagggcttg ctgcggtttt tgttcctg 48  
  
<210> SEQ ID NO 355  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 355  
  
ctatgaaatt aaccctcact aaagggagct gccgatccca ctggtatt 48  
  
<210> SEQ ID NO 356  
<211> LENGTH: 46  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 356  
  
ggattctaatacgcactcact atagggcgga tcctggccgg cctctg 46  
  
<210> SEQ ID NO 357  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 357  
  
ctatgaaatt aaccctcact aaagggagcc cgggcatggt ctcagtta 48  
  
<210> SEQ ID NO 358  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
  
<400> SEQUENCE: 358  
  
ggattctaatacgcactcact atagggcggg aagatggcga ggaggag 47  
  
<210> SEQ ID NO 359  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

---

-continued

---

<400> SEQUENCE: 359  
ctatgaaatt aaccctcact aaagggacca aggccacaaa cggaaatc 48

<210> SEQ ID NO 360  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 360  
ggattcta at acgactcact atagggtgt gtttcattc tgccagta 48

<210> SEQ ID NO 361  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 361  
ctatgaaatt aaccctcact aaagggaggg tacaattaag ggtggat 48

<210> SEQ ID NO 362  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 362  
ggattcta at acgactcact atagggcccg cctcgtcct gctcctg 47

<210> SEQ ID NO 363  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 363  
ctatgaaatt aaccctcact aaagggagga ttgccgcgac cctcacag 48

<210> SEQ ID NO 364  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 364  
ggattcta at acgactcact atagggcccc tctgccttc cctgtcc 47

<210> SEQ ID NO 365  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 365  
ctatgaaatt aaccctcact aaagggagtg gtggccgca ttatctgc 48

---

-continued

---

<210> SEQ ID NO 366  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 366

ggattctaatacgcactcactatagggcgacgcatggcagcgatgagg 48

<210> SEQ ID NO 367  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 367

ctatgaaattaacctcactaaagggacagacggggcagagggagtg 47

<210> SEQ ID NO 368  
<211> LENGTH: 47  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 368

ggattctaatacgcactcactatagggccagaggcgtgagagaaaac 47

<210> SEQ ID NO 369  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 369

ctatgaaattaacctcactaaagggaaaacatgtcatcgggagtgg 48

<210> SEQ ID NO 370  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 370

ggattctaatacgcactcactatagggccgggtggagtgg aacagaaa 48

<210> SEQ ID NO 371  
<211> LENGTH: 48  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 371

ctatgaaattaacctcactaaagggacacagacagagcccatatcgc 48

<210> SEQ ID NO 372  
<211> LENGTH: 47

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
 <400> SEQUENCE: 372  
 ggattctaatacgcactcactatagggccaggaaatccggatgtctc 47

<210> SEQ ID NO 373  
 <211> LENGTH: 48  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
 <400> SEQUENCE: 373  
 ctatgaaattaacctcactaaaggagtaaggggatgccaccgagta 48

<210> SEQ ID NO 374  
 <211> LENGTH: 47  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
 <400> SEQUENCE: 374  
 ggattctaatacgcactcactatagggccagctaccgcagaggag 47

<210> SEQ ID NO 375  
 <211> LENGTH: 48  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic Oligonucleotide Probe  
 <400> SEQUENCE: 375  
 ctatgaaattaacctcactaaagggatccagggtgatgaggtccaga 48

<210> SEQ ID NO 376  
 <211> LENGTH: 997  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien  
 <400> SEQUENCE: 376  
 cccacgcgctcgcgatcttaccacaacacacctgaggagaaagaaagag 50  
 agggaggagagaaaaagagagagagaaacacacacacacacacacacac 100  
 aaaaatgaaattcatctaaatcatctgaaacacacacacacacacacac 150  
 tgcctctcttcccacatgctcttatggactgttgctggggtccccatcct 200  
 atttctcagtgccctgttcaaccagatgtgttgagacatttcgcatct 250  
 ttcaaacctgtgatgagaaagtttcagctacacacacacacacacacac 300  
 ctctcctgctacaattatgatcaggttcaatcaagaattgtgtccatt 350  
 gaactgggaaatatttcaatccagctgctattcttttctactgacacca 400  
 tttcctggggttaagttaagaactgctcagccatgggggctcacctg 450  
 gtggttatcaactcacaggaaggcaggaaatctcttctcaagaacac 500  
 taaaatgagagagtttttttttgactgtcagaccaggtgtcgagggtc 550  
 agtggaatgggtggacggcacacctttgacaaagtctctgagcttctg 600



-continued

---

```

gatgtagggg agcccaacaa catagctacc ctggaggact gtgccacat      650
gagagactct tcaaacccaa ggcaaatg gaatgatgta acctgtttcc      700
tcaattattt tcggatttgt gaaatgtag gaataaatcc tttgaacaaa      750
ggaaaatctc ttaagaaca gaaggcaca ctcaaatgta taaagaagga      800
agagcaagaa catggccaca cccaccgccc cacacgagaa atttgtgccc      850
tgaacttcaa aggacttcat aagtatttgt tactctgata caaataaaaa      900
taagtagttt taaatgtaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa      950
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa      997
    
```

```

<210> SEQ ID NO 377
<211> LENGTH: 219
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien
    
```

<400> SEQUENCE: 377

```

Met Asn Ser Ser Lys Ser Ser Glu Thr Gln Cys Thr Glu Arg Gly
 1          5          10          15
Cys Phe Ser Ser Gln Met Phe Leu Trp Thr Val Ala Gly Ile Pro
 20          25          30
Ile Leu Phe Leu Ser Ala Cys Phe Ile Thr Arg Cys Val Val Thr
 35          40          45
Phe Arg Ile Phe Gln Thr Cys Asp Glu Lys Lys Phe Gln Leu Pro
 50          55          60
Glu Asn Phe Thr Glu Leu Ser Cys Tyr Asn Tyr Gly Ser Gly Ser
 65          70          75
Val Lys Asn Cys Cys Pro Leu Asn Trp Glu Tyr Phe Gln Ser Ser
 80          85          90
Cys Tyr Phe Phe Ser Thr Asp Thr Ile Ser Trp Ala Leu Ser Leu
 95          100         105
Lys Asn Cys Ser Ala Met Gly Ala His Leu Val Val Ile Asn Ser
 110         115         120
Gln Glu Glu Gln Glu Phe Leu Ser Tyr Lys Lys Pro Lys Met Arg
 125         130         135
Glu Phe Phe Ile Gly Leu Ser Asp Gln Val Val Glu Gly Gln Trp
 140         145         150
Gln Trp Val Asp Gly Thr Pro Leu Thr Lys Ser Leu Ser Phe Trp
 155         160         165
Asp Val Gly Glu Pro Asn Asn Ile Ala Thr Leu Glu Asp Cys Ala
 170         175         180
Thr Met Arg Asp Ser Ser Asn Pro Arg Gln Asn Trp Asn Asp Val
 185         190         195
Thr Cys Phe Leu Asn Tyr Phe Arg Ile Cys Glu Met Val Gly Ile
 200         205         210
Asn Pro Leu Asn Lys Gly Lys Ser Leu
 215
    
```

```

<210> SEQ ID NO 378
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
    
```

---

-continued

---

<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 378

ttcagcttct gggatgtagg g 21

<210> SEQ ID NO 379  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic Oligonucleotide Probe

<400> SEQUENCE: 379

tattcctacc attcacaaa tccg 24

<210> SEQ ID NO 380  
<211> LENGTH: 49  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 380

ggagactgt gccacatga gagactcttc aaaccaagg caaaattgg 49

<210> SEQ ID NO 381  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 381

gcagatttg aggacagcca cctcca 26

<210> SEQ ID NO 382  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 382

ggccttgag acaaccgt 18

<210> SEQ ID NO 383  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 383

cagactgagg gagatccgag a 21

<210> SEQ ID NO 384  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 384

---

-continued

---

cagctgccct tcccacaacca 20

<210> SEQ ID NO 385  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 385

catcaagcgc ctctacca 18

<210> SEQ ID NO 386  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 386

cacaaactcg aactgcttct g 21

<210> SEQ ID NO 387  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 387

ggccatcac agctccct 18

<210> SEQ ID NO 388  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 388

gggatgtggt gaacacagaa ca 22

<210> SEQ ID NO 389  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 389

tgccagctgc atgctgccag tt 22

<210> SEQ ID NO 390  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 390

cagaaggatg tcccgtggaa 20

<210> SEQ ID NO 391

---

-continued

---

<211> LENGTH: 17  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 391

gccgctgtcc actgcag 17

<210> SEQ ID NO 392  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 392

gacggcatcc tcagggccac a 21

<210> SEQ ID NO 393  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 393

atgtcctcca tgcccacgcy 20

<210> SEQ ID NO 394  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 394

gagtgcgaca tcgagagctt 20

<210> SEQ ID NO 395  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 395

ccgcagcctc agtcatga 18

<210> SEQ ID NO 396  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 396

gaagagcaca gctgcagatc c 21

<210> SEQ ID NO 397  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

---

-continued

---

<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 397

gagggtgcct ggctttggta gt 22

<210> SEQ ID NO 398  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 398

cctctggcgc cccactcaa 20

<210> SEQ ID NO 399  
<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 399

ccaggagagc tggcgatg 18

<210> SEQ ID NO 400  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 400

gcaaattcag ggctcactag aga 23

<210> SEQ ID NO 401  
<211> LENGTH: 29  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 401

cacagagcat ttgtccatca gcagttcag 29

<210> SEQ ID NO 402  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 402

ggcagagact tccagtcact ga 22

<210> SEQ ID NO 403  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 403

---

-continued

---

gcccaagggtg gtgtagata gg 22

<210> SEQ ID NO 404  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 404

caggccccct tgatctgtac ccca 24

<210> SEQ ID NO 405  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 405

gggacgtgct tctacaagaa cag 23

<210> SEQ ID NO 406  
<211> LENGTH: 26  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 406

caggcttaca atgttatgat cagaca 26

<210> SEQ ID NO 407  
<211> LENGTH: 31  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 407

tattcagagt tttccattgg cagtgccagt t 31

<210> SEQ ID NO 408  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 408

tctacatcag cctctctgcg c 21

<210> SEQ ID NO 409  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 409

cgatcttctc caccaggag cgg 23

<210> SEQ ID NO 410

---

-continued

---

<211> LENGTH: 18  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 410

gccaggcctc acattcgt 18

<210> SEQ ID NO 411  
<211> LENGTH: 23  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 411

ctccctgaat ggcagcctga gca 23

<210> SEQ ID NO 412  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 412

agggtgttat taaggccta cgct 24

<210> SEQ ID NO 413  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 413

cagagcagag ggtgccttg 19

<210> SEQ ID NO 414  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 414

tggcggagtc ccctcttggc t 21

<210> SEQ ID NO 415  
<211> LENGTH: 22  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 415

ccctgtttcc ctatgcatca ct 22

<210> SEQ ID NO 416  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 416

tcaaccctg accctttcct a 21

<210> SEQ ID NO 417  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 417

ggcaggggac aagccatctc tcct 24

<210> SEQ ID NO 418  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 418

gggactgaac tgccagcttc 20

<210> SEQ ID NO 419  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 419

gggcctaac ctattacct tt 22

<210> SEQ ID NO 420  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 420

tgtctgcctc agccccagga agg 23

<210> SEQ ID NO 421  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 421

tctgtccacc atcttgcctt g 21

<210> SEQ ID NO 422  
 <211> LENGTH: 3554  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 422

gggactacaa gccgcgccgc gctgcgcgtg gccctcagc aacctogac 50



-continued

---

atggcgctga ggcggccacc gcgactccgg ctctgcgctc ggctgcctga	100
cttcttcctg ctgctgcttt tcaggggctg cctgataggg gctgtaaactc	150
tcaaatccag caatcgaacc ccagtggtac aggaatttga aagtgtggaa	200
ctgtcttgca tcattacgga ttcgcagaca agtgacccca ggatcgagtg	250
gaagaaaatt caagatgaac aaaccacata tgtgtttttt gacaacaaaa	300
ttcagggaga cttggcgggt cgtgcagaaa tactggggaa gacatccctg	350
aagatctgga atgtgacacg gagagactca gccctttatc gctgtgaggt	400
cgttgctcga aatgaccgca aggaaattga tgagatttg atcgagttaa	450
ctgtgcaagt gaagccagtg acccctgtct gtagagtgcc gaaggctgta	500
ccagtaggca agatggcaac actgcactgc caggagagtg agggccacc	550
ccggcctcac tacagctggt atcgcaatga tgtaccactg cccacggatt	600
ccagagccaa tcccagattt cgcaattctt ctttccactt aaactctgaa	650
acaggcactt tgggtgtcac tgctgttcac aaggacgact ctgggcagta	700
ctactgcatt gcttccaatg acgcaggctc agccagtggt gaggagcagg	750
agatggaagt ctatgacctg aacattggcg gaattatttg gggggttctg	800
gttgcctctg ctgtactggc cctgatcacg ttgggcatct gctgtgcata	850
cagacgtggc tacttcatca acaataaaca ggatggagaa agttacaaga	900
accagggaa accagatgga gttactaca tccgcactga cgaggagggc	950
gacttcagac acaagtcatc gtttgtgatc tgagaccgcg ggtgtggctg	1000
agagcgcaca gagcgcacgt gcacatacct ctgctagaaa ctctgtcaa	1050
ggcagcgaga gctgatgcac tcggacagag ctagacactc attcagaagc	1100
ttttcgtttt ggccaaagt gaccactact cttcttactc taacaagcca	1150
catgaataga agaattttcc tcaagatgga cccggtaaat ataaccacaa	1200
ggaaagcga ctgggtgcgt tcaactgagt gggttcctaa tctgtttctg	1250
gcctgattcc cgcgatgagta ttagggtgat cttaaagagt ttgctcacgt	1300
aaacgcccgt gctgggccct gtgaagccag catgttcacc actggtcgtt	1350
cagcagccac gacagcacca tgtgagatgg cgagggtggct ggacagcacc	1400
agcagcgcat cccggcggga acccagaaaa ggcttcttac acagcagcct	1450
tacttcatcg gccacagac accaccgcag tttcttctta aaggctctgc	1500
tgatcggtgt tgcaagtgtcc attgtggaga agcttttttg atcagcattt	1550
tgtaaaaaa accaaaatca ggaaggtaaa ttggttgctg gaagagggat	1600
cttgcttagg gaaccctgct tgtccaacag ggtgtcagga tttaaggaaa	1650
accttctgt taggctaagt ctgaaatggt actgaaatat gcttttctat	1700
gggtcttggt tattttataa aattttacat ctaaattttt gctaaggatg	1750
tattttgatt attgaaaaga aaatttctat ttaaactgta aatataattg	1800
catacaatgt taaataacct atttttttaa aaaagttcaa cttaaggtag	1850
aagtccaag ctactagtgt taaattggaa aatatcaata attaagagta	1900
ttttacccaa ggaatcctct catggaagtt tactgtgatg ttccttttct	1950

-continued

---

cacacaagtt ttagcctttt tcacaagggg actcatactg tctacacatc	2000
agaccatagt tgcttaggaa acctttaaaa attccagtta agcaatgttg	2050
aaatcagttt gcatctcttc aaaagaaacc tctcaggtta gctttgaact	2100
gcctcttctc gagatgacta ggacagtctg taccagagg ccaccagaa	2150
gccctcagat gtacatacac agatgccagt cagctcctgg ggttgcgcca	2200
ggcgccccg ctctagctca ctgttgcctc gctgtctgcc aggaggcct	2250
gccatccttg gccctggca gtggctgtgt cccagtgagc tttactcacg	2300
tggcccttgc ttcattccagc acagctctca ggtgggact gcagggacac	2350
tggtgtcttc catgtagcgt cccagcttg ggctcctgta acagacctct	2400
ttttggttat ggatggctca caaaatagg ccccaatgc tattttttt	2450
ttttaagttt gttaattat ttgtaagat tgtctaaggc caaagcaat	2500
tgcgaaatca agtctgtcaa gtacaataac atttttaaaa gaaaatggat	2550
cccactgttc ctctttgcca cagagaaagc acccagacgc cacaggctct	2600
gtcgcatttc aaaacaaacc atgatggagt ggcggccagt ccagcctttt	2650
aaagaacgtc aggtggagca gccaggtgaa aggcctggcg gggaggaaag	2700
tgaaacgcct gaatcaaaaag cagttttcta attttgactt taaattttc	2750
atccgcccga gacactgctc ccatttggg ggggacatta gcaacatcac	2800
tcagaagcct gtgttcttca agagcaggtg ttctcagcct cacatgccct	2850
gccgtgctgg actcaggact gaagtctgt aaagcaagga gctgctgaga	2900
aggagcactc cactgtgtgc ctggagaatg gctctcacta ctcacctgt	2950
ctttcagctt ccagtgctt gggttttta tactttgaca gcttttttt	3000
aattgcatat atgagactgt gttgactttt tttagttatg tgaaacactt	3050
tgccgcaggc cgctggcagc aggcaggaaa tgctccagca gttgctcagt	3100
gctccctggt gtctgtgca tggcatcctg gatgcttagc atgcaagttc	3150
cctccatcat tgccaccttg gtagagagg atggctccc accctcagcg	3200
ttggggattc acgctccagc ctctctcttg gttgtcatag tgatagggt	3250
gccttattgc cccctctct tataccctaa aacctctac actagtgcc	3300
tgggaaccag gtctgaaaaa gtagagagaa gtgaaagtag agtctgggaa	3350
gtagctgcct ataactgaga ctagacggaa aaggaatact cgtgtatttt	3400
aagatatgaa tgtgactcaa gactcgaggc cgatacaggc ctgtgattct	3450
gcctttggat ggatggtgct gtacacagat gctacagact tgtactaaca	3500
caccgtaatt tggcatttgt ttaacctcat ttataaaagc ttcaaaaaa	3550
ccca	3554

&lt;210&gt; SEQ ID NO 423

&lt;211&gt; LENGTH: 310

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo Sapien

&lt;400&gt; SEQUENCE: 423

Met Ala Leu Arg Arg Pro Pro Arg Leu Arg Leu Cys Ala Arg Leu  
 1 5 10 15

-continued

---

Pro	Asp	Phe	Phe	Leu	Leu	Leu	Phe	Arg	Gly	Cys	Leu	Ile	Gly	
				20				25					30	
Ala	Val	Asn	Leu	Lys	Ser	Ser	Asn	Arg	Thr	Pro	Val	Val	Gln	Glu
				35				40						45
Phe	Glu	Ser	Val	Glu	Leu	Ser	Cys	Ile	Ile	Thr	Asp	Ser	Gln	Thr
				50				55						60
Ser	Asp	Pro	Arg	Ile	Glu	Trp	Lys	Lys	Ile	Gln	Asp	Glu	Gln	Thr
				65				70						75
Thr	Tyr	Val	Phe	Phe	Asp	Asn	Lys	Ile	Gln	Gly	Asp	Leu	Ala	Gly
				80				85						90
Arg	Ala	Glu	Ile	Leu	Gly	Lys	Thr	Ser	Leu	Lys	Ile	Trp	Asn	Val
				95				100						105
Thr	Arg	Arg	Asp	Ser	Ala	Leu	Tyr	Arg	Cys	Glu	Val	Val	Ala	Arg
				110				115						120
Asn	Asp	Arg	Lys	Glu	Ile	Asp	Glu	Ile	Val	Ile	Glu	Leu	Thr	Val
				125				130						135
Gln	Val	Lys	Pro	Val	Thr	Pro	Val	Cys	Arg	Val	Pro	Lys	Ala	Val
				140				145						150
Pro	Val	Gly	Lys	Met	Ala	Thr	Leu	His	Cys	Gln	Glu	Ser	Glu	Gly
				155				160						165
His	Pro	Arg	Pro	His	Tyr	Ser	Trp	Tyr	Arg	Asn	Asp	Val	Pro	Leu
				170				175						180
Pro	Thr	Asp	Ser	Arg	Ala	Asn	Pro	Arg	Phe	Arg	Asn	Ser	Ser	Phe
				185				190						195
His	Leu	Asn	Ser	Glu	Thr	Gly	Thr	Leu	Val	Phe	Thr	Ala	Val	His
				200				205						210
Lys	Asp	Asp	Ser	Gly	Gln	Tyr	Tyr	Cys	Ile	Ala	Ser	Asn	Asp	Ala
				215				220						225
Gly	Ser	Ala	Arg	Cys	Glu	Glu	Gln	Glu	Met	Glu	Val	Tyr	Asp	Leu
				230				235						240
Asn	Ile	Gly	Gly	Ile	Ile	Gly	Gly	Val	Leu	Val	Val	Leu	Ala	Val
				245				250						255
Leu	Ala	Leu	Ile	Thr	Leu	Gly	Ile	Cys	Cys	Ala	Tyr	Arg	Arg	Gly
				260				265						270
Tyr	Phe	Ile	Asn	Asn	Lys	Gln	Asp	Gly	Glu	Ser	Tyr	Lys	Asn	Pro
				275				280						285
Gly	Lys	Pro	Asp	Gly	Val	Asn	Tyr	Ile	Arg	Thr	Asp	Glu	Glu	Gly
				290				295						300
Asp	Phe	Arg	His	Lys	Ser	Ser	Phe	Val	Ile					
				305				310						

---

What is claimed is:

1. Isolated nucleic acid having at least 80% sequence identity to a nucleotide sequence that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of the amino acid sequence shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:12), **FIG. 9** (SEQ ID NO:18), **FIG. 11** (SEQ ID NO:23), **FIG. 13** (SEQ ID NO:28), **FIG. 15** (SEQ ID NO:34), **FIG. 17** (SEQ ID NO:39), **FIG. 19** (SEQ ID NO:49), **FIG. 22** (SEQ ID NO:59), **FIG. 24** (SEQ ID NO:64), **FIG. 26** (SEQ ID NO:69), **FIG. 28** (SEQ ID NO:71), **FIG. 30** (SEQ ID

NO:73), **FIG. 32** (SEQ ID NO:84), **FIG. 34** (SEQ ID NO:91), **FIG. 36** (SEQ ID NO:96), **FIG. 38** (SEQ ID NO:104), **FIG. 40** (SEQ ID NO:109), **FIG. 42** (SEQ ID NO:114), **FIG. 44** (SEQ ID NO:119), **FIG. 46** (SEQ ID NO:127), **FIG. 48** (SEQ ID NO:132), **FIG. 50** (SEQ ID NO:137), **FIG. 52** (SEQ ID NO:142), **FIG. 54** (SEQ ID NO:148), **FIG. 56** (SEQ ID NO:153), **FIG. 58** (SEQ ID NO:159), **FIG. 60** (SEQ ID NO:164), **FIG. 62** (SEQ ID NO:170), **FIG. 64** (SEQ ID NO:175), **FIG. 66** (SEQ ID NO:177), **FIG. 68** (SEQ ID NO:185), **FIG. 70** (SEQ ID NO:190), **FIG. 72** (SEQ ID NO:195), **FIG. 74** (SEQ ID

NO:201), **FIG. 76** (SEQ ID NO:207), **FIG. 78** (SEQ ID NO:213), **FIG. 80** (SEQ ID NO:221), **FIG. 82** (SEQ ID NO:227), **FIG. 84** (SEQ ID NO:236), **FIG. 86** (SEQ ID NO:245), **FIG. 88** (SEQ ID NO:250), **FIG. 90** (SEQ ID NO:255), **FIG. 92** (SEQ ID NO:257), **FIG. 94** (SEQ ID NO:259), **FIG. 96** (SEQ ID NO:261), **FIG. 98** (SEQ ID NO:263), **FIG. 100** (SEQ ID NO:285), **FIG. 102** (SEQ ID NO:290), **FIG. 104** (SEQ ID NO:292), **FIG. 106** (SEQ ID NO:294), **FIG. 108** (SEQ ID NO:310), **FIG. 110** (SEQ ID NO:315), **FIG. 112** (SEQ ID NO:320), **FIG. 114** (SEQ ID NO:325), **FIG. 116** (SEQ ID NO:332), **FIG. 118** (SEQ ID NO:339), **FIG. 120** (SEQ ID NO:341), **FIG. 122** (SEQ ID NO:377) and **FIG. 124** (SEQ ID NO:423).

2. The nucleic acid of claim 1, wherein said nucleotide sequence comprises a nucleotide sequence selected from the group consisting of the sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:11), **FIG. 8** (SEQ ID NO:17), **FIG. 10** (SEQ ID NO:22), **FIG. 12** (SEQ ID NO:27), **FIG. 14** (SEQ ID NO:33), **FIG. 16** (SEQ ID NO:38), **FIG. 18** (SEQ ID NO:48), **FIG. 21** (SEQ ID NO:58), **FIG. 23** (SEQ ID NO:63), **FIG. 25** (SEQ ID NO:68), **FIG. 27** (SEQ ID NO:70), **FIG. 29** (SEQ ID NO:72), **FIG. 31** (SEQ ID NO:83), **FIG. 33** (SEQ ID NO:90), **FIG. 35** (SEQ ID NO:95), **FIG. 37** (SEQ ID NO:103), **FIG. 39** (SEQ ID NO:108), **FIG. 41** (SEQ ID NO:113), **FIG. 43** (SEQ ID NO:118), **FIG. 45** (SEQ ID NO:126), **FIG. 47** (SEQ ID NO:131), **FIG. 49** (SEQ ID NO:136), **FIG. 51** (SEQ ID NO:141), **FIG. 53** (SEQ ID NO:147), **FIG. 55** (SEQ ID NO:152), **FIG. 57** (SEQ ID NO:158), **FIG. 59** (SEQ ID NO:163), **FIG. 61** (SEQ ID NO:169), **FIG. 63** (SEQ ID NO:174), **FIG. 65** (SEQ ID NO:176), **FIG. 67** (SEQ ID NO:184), **FIG. 69** (SEQ ID NO:189), **FIG. 71** (SEQ ID NO:194), **FIG. 73** (SEQ ID NO:200), **FIG. 75** (SEQ ID NO:206), **FIG. 77** (SEQ ID NO:212), **FIG. 79** (SEQ ID NO:220), **FIG. 81** (SEQ ID NO:226), **FIG. 83** (SEQ ID NO:235), **FIG. 85** (SEQ ID NO:244), **FIG. 87** (SEQ ID NO:249), **FIG. 89** (SEQ ID NO:254), **FIG. 91** (SEQ ID NO:256), **FIG. 93** (SEQ ID NO:258), **FIG. 95** (SEQ ID NO:260), **FIG. 97** (SEQ ID NO:262), **FIG. 99** (SEQ ID NO:284), **FIG. 101** (SEQ ID NO:289), **FIG. 103** (SEQ ID NO:291), **FIG. 105** (SEQ ID NO:293), **FIG. 107** (SEQ ID NO:309), **FIG. 109** (SEQ ID NO:314), **FIG. 111** (SEQ ID NO:319), **FIG. 113** (SEQ ID NO:324), **FIG. 115** (SEQ ID NO:331), **FIG. 117** (SEQ ID NO:338), **FIG. 119** (SEQ ID NO:340), **FIG. 121** (SEQ ID NO:376) and **FIG. 123** (SEQ ID NO:422), or the complement thereof.

3. The nucleic acid of claim 1, wherein said nucleotide sequence comprises a nucleotide sequence selected from the group consisting of the full-length coding sequence of the sequence shown in **FIG. 1** (SEQ ID NO:1), **FIG. 3** (SEQ ID NO:3), **FIG. 5** (SEQ ID NO:11), **FIG. 8** (SEQ ID NO:17), **FIG. 10** (SEQ ID NO:22), **FIG. 12** (SEQ ID NO:27), **FIG. 14** (SEQ ID NO:33), **FIG. 16** (SEQ ID NO:38), **FIG. 18** (SEQ ID NO:48), **FIG. 21** (SEQ ID NO:58), **FIG. 23** (SEQ ID NO:63), **FIG. 25** (SEQ ID NO:68), **FIG. 27** (SEQ ID NO:70), **FIG. 29** (SEQ ID NO:72), **FIG. 31** (SEQ ID NO:83), **FIG. 33** (SEQ ID NO:90), **FIG. 35** (SEQ ID NO:95), **FIG. 37** (SEQ ID NO:103), **FIG. 39** (SEQ ID NO:108), **FIG. 41** (SEQ ID NO:113), **FIG. 43** (SEQ ID NO:118), **FIG. 45** (SEQ ID NO:126), **FIG. 47** (SEQ ID NO:131), **FIG. 49** (SEQ ID NO:136), **FIG. 51** (SEQ ID NO:141), **FIG. 53** (SEQ ID NO:147), **FIG. 55** (SEQ ID NO:152), **FIG. 57** (SEQ ID NO:158), **FIG. 59** (SEQ ID

NO:163), **FIG. 61** (SEQ ID NO:169), **FIG. 63** (SEQ ID NO:174), **FIG. 65** (SEQ ID NO:176), **FIG. 67** (SEQ ID NO:184), **FIG. 69** (SEQ ID NO:189), **FIG. 71** (SEQ ID NO:194), **FIG. 73** (SEQ ID NO:200), **FIG. 75** (SEQ ID NO:206), **FIG. 77** (SEQ ID NO:212), **FIG. 79** (SEQ ID NO:220), **FIG. 81** (SEQ ID NO:226), **FIG. 83** (SEQ ID NO:235), **FIG. 85** (SEQ ID NO:244), **FIG. 87** (SEQ ID NO:249), **FIG. 89** (SEQ ID NO:254), **FIG. 91** (SEQ ID NO:256), **FIG. 93** (SEQ ID NO:258), **FIG. 95** (SEQ ID NO:260), **FIG. 97** (SEQ ID NO:262), **FIG. 99** (SEQ ID NO:284), **FIG. 101** (SEQ ID NO:289), **FIG. 103** (SEQ ID NO:291), **FIG. 105** (SEQ ID NO:293), **FIG. 107** (SEQ ID NO:309), **FIG. 109** (SEQ ID NO:314), **FIG. 111** (SEQ ID NO:319), **FIG. 113** (SEQ ID NO:324), **FIG. 115** (SEQ ID NO:331), **FIG. 117** (SEQ ID NO:338), **FIG. 119** (SEQ ID NO:340), **FIG. 121** (SEQ ID NO:376) and **FIG. 123** (SEQ ID NO:422), or the complement thereof.

4. Isolated nucleic acid which comprises the full-length coding sequence of the DNA deposited under accession number ATCC 209258, ATCC 209256, ATCC 209264, ATCC 209250, ATCC 209375, ATCC 209378, ATCC 209384, ATCC 209396, ATCC 209420, ATCC 209480, ATCC 209265, ATCC 209257, ATCC 209262, ATCC 209253, ATCC 209402, ATCC 209401, ATCC 209397, ATCC 209400, ATCC 209385, ATCC 209367, ATCC 209432, ATCC 209263, ATCC 209251, ATCC 209255, ATCC 209252, ATCC 209373, ATCC 209370, ATCC 209523, ATCC 209372, ATCC 209374, ATCC 209373, ATCC 209382, ATCC 209383, ATCC 209403, ATCC 209398, ATCC 209399, ATCC 209392, ATCC 209387, ATCC 209383, ATCC 209394, ATCC 209421, ATCC 209393, ATCC 209418, ATCC 209485, ATCC 209483, ATCC 209482, ATCC 209491, ATCC 209481, ATCC 209438, ATCC 209927, ATCC 209439, ATCC 209489, ATCC 209433, ATCC 209488, ATCC 209434, ATCC 209395, ATCC 209486, ATCC 209490, ATCC 209484, ATCC 209371 or ATCC 203553.

5. A vector comprising the nucleic acid of claim 1.

6. The vector of claim 5 operably linked to control sequences recognized by a host cell transformed with the vector.

7. A host cell comprising the vector of claim 5.

8. The host cell of claim 7 wherein said cell is a CHO cell.

9. The host cell of claim 7 wherein said cell is an *E. coli*.

10. The host cell of claim 7 wherein said cell is a yeast cell.

11. A process for producing a PRO polypeptides comprising culturing the host cell of claim 7 under conditions suitable for expression of said PRO polypeptide and recovering said PRO polypeptide from the cell culture.

12. Isolated native sequence PRO polypeptide having at least 80% sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:12), **FIG. 9** (SEQ ID NO:18), **FIG. 11** (SEQ ID NO:23), **FIG. 13** (SEQ ID NO:28), **FIG. 15** (SEQ ID NO:34), **FIG. 17** (SEQ ID NO:39), **FIG. 19** (SEQ ID NO:49), **FIG. 22** (SEQ ID NO:59), **FIG. 24** (SEQ ID NO:64), **FIG. 26** (SEQ ID NO:69), **FIG. 28** (SEQ ID NO:71), **FIG. 30** (SEQ ID NO:73), **FIG. 32** (SEQ ID NO:84), **FIG. 34** (SEQ ID NO:91), **FIG. 36** (SEQ ID NO:96), **FIG. 38** (SEQ ID NO:104), **FIG. 40** (SEQ ID NO:109), **FIG. 42** (SEQ ID NO:114), **FIG. 44** (SEQ ID NO:119), **FIG. 46** (SEQ ID NO:127), **FIG. 48** (SEQ ID

NO:132), **FIG. 50** (SEQ ID NO:137), **FIG. 52** (SEQ ID NO:142), **FIG. 54** (SEQ ID NO:148), **FIG. 56** (SEQ ID NO:153), **FIG. 58** (SEQ ID NO:159), **FIG. 60** (SEQ ID NO:164), **FIG. 62** (SEQ ID NO:170), **FIG. 64** (SEQ ID NO:175), **FIG. 66** (SEQ ID NO:177), **FIG. 68** (SEQ ID NO:185), **FIG. 70** (SEQ ID NO:190), **FIG. 72** (SEQ ID NO:195), **FIG. 74** (SEQ ID NO:201), **FIG. 76** (SEQ ID NO:207), **FIG. 78** (SEQ ID NO:213), **FIG. 80** (SEQ ID NO:221), **FIG. 82** (SEQ ID NO:227), **FIG. 84** (SEQ ID NO:236), **FIG. 86** (SEQ ID NO:245), **FIG. 88** (SEQ ID NO:250), **FIG. 90** (SEQ ID NO:255), **FIG. 92** (SEQ ID NO:257), **FIG. 94** (SEQ ID NO:259), **FIG. 96** (SEQ ID NO:261), **FIG. 98** (SEQ ID NO:263), **FIG. 100** (SEQ ID NO:285), **FIG. 102** (SEQ ID NO:290), **FIG. 104** (SEQ ID NO:292), **FIG. 106** (SEQ ID NO:294), **FIG. 108** (SEQ ID NO:310), **FIG. 110** (SEQ ID NO:315), **FIG. 112** (SEQ ID NO:320), **FIG. 114** (SEQ ID NO:325), **FIG. 116** (SEQ ID NO:332), **FIG. 118** (SEQ ID NO:339), **FIG. 120** (SEQ ID NO:341), **FIG. 122** (SEQ ID NO:377) and **FIG. 124** (SEQ ID NO:423).

13. Isolated PRO polypeptide having at least 80% sequence identity to the amino acid sequence encoded by the nucleotide deposited under accession number ATCC 209258, ATCC209256, ATCC 209264, ATCC 209250, ATCC 209375, ATCC 209378, ATCC 209384, ATCC 209396, ATCC 209420, ATCC 209480, ATCC 209265, ATCC 209257, ATCC 209262, ATCC 209253, ATCC 209402, ATCC 209401, ATCC 209397, ATCC 209400, ATCC 209385, ATCC 209367, ATCC 209432, ATCC 209263, ATCC 209251, ATCC 209255, ATCC 209252, ATCC 209373, ATCC 209370, ATCC 209523, ATCC 209372, ATCC 209374, ATCC 209373, ATCC 209382, ATCC 209383, ATCC 209403, ATCC 209398, ATCC 209399, ATCC 209392, ATCC 209387, ATCC 209388, ATCC 209394, ATCC 209421, ATCC 209393, ATCC 209418, ATCC 209485, ATCC 209483, ATCC 209482, ATCC 209491, ATCC 209481, ATCC 209438, ATCC 209927, ATCC 209439, ATCC 209489, ATCC 209433, ATCC 209488, ATCC 209434, ATCC 209395, ATCC 209486, ATCC 209490, ATCC 209484, ATCC 209371 or ATCC 203553.

14. A chimeric molecule comprising a polypeptide according to claim 12 fused to a heterologous amino acid sequence.

15. The chimeric molecule of claim 14 wherein said heterologous amino acid sequence is an epitope tag sequence.

16. The chimeric molecule of claim 14 wherein said heterologous amino acid sequence is a Fc region of an immunoglobulin.

17. An antibody which specifically binds to a PRO polypeptide according to claim 12.

18. The antibody of claim 17 wherein said antibody is a monoclonal antibody.

19. Isolated nucleic acid having at least 80% nucleic acid sequence identity to:

- (a) a nucleotide sequence encoding the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:12), **FIG. 9** (SEQ ID NO:18), **FIG. 11** (SEQ ID NO:23), **FIG. 13** (SEQ ID NO:28), **FIG. 15** (SEQ ID NO:34), **FIG. 17** (SEQ ID NO:39), **FIG. 19** (SEQ ID NO:49), **FIG. 22** (SEQ ID NO:59), **FIG. 24** (SEQ ID NO:64), **FIG. 26** (SEQ ID NO:69), **FIG. 28** (SEQ ID NO:71), **FIG. 30** (SEQ ID NO:73),

**FIG. 32** (SEQ ID NO:84), **FIG. 34** (SEQ ID NO:91), **FIG. 36** (SEQ ID NO:96), **FIG. 38** (SEQ ID NO:104), **FIG. 40** (SEQ ID NO:109), **FIG. 42** (SEQ ID NO:114), **FIG. 44** (SEQ ID NO:119), **FIG. 46** (SEQ ID NO:127), **FIG. 48** (SEQ ID NO:132), **FIG. 50** (SEQ ID NO:137), **FIG. 52** (SEQ ID NO:142), **FIG. 54** (SEQ ID NO:148), **FIG. 56** (SEQ ID NO:153), **FIG. 58** (SEQ ID NO:159), **FIG. 60** (SEQ ID NO:164), **FIG. 62** (SEQ ID NO:170), **FIG. 64** (SEQ ID NO:175), **FIG. 66** (SEQ ID NO:177), **FIG. 68** (SEQ ID NO:185), **FIG. 70** (SEQ ID NO:190), **FIG. 72** (SEQ ID NO:195), **FIG. 74** (SEQ ID NO:201), **FIG. 76** (SEQ ID NO:207), **FIG. 78** (SEQ ID NO:213), **FIG. 80** (SEQ ID NO:221), **FIG. 82** (SEQ ID NO:227), **FIG. 84** (SEQ ID NO:236), **FIG. 86** (SEQ ID NO:245), **FIG. 88** (SEQ ID NO:250), **FIG. 90** (SEQ ID NO:255), **FIG. 92** (SEQ ID NO:257), **FIG. 94** (SEQ ID NO:259), **FIG. 96** (SEQ ID NO:261), **FIG. 98** (SEQ ID NO:263), **FIG. 100** (SEQ ID NO:285), **FIG. 102** (SEQ ID NO:290), **FIG. 104** (SEQ ID NO:292), **FIG. 106** (SEQ ID NO:294), **FIG. 108** (SEQ ID NO:310), **FIG. 110** (SEQ ID NO:315), **FIG. 112** (SEQ ID NO:320), **FIG. 114** (SEQ ID NO:325), **FIG. 116** (SEQ ID NO:332), **FIG. 118** (SEQ ID NO:339), **FIG. 120** (SEQ ID NO:341), **FIG. 122** (SEQ ID NO:377) or **FIG. 124** (SEQ ID NO:423), lacking its associated signal peptide;

- (b) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID NO:2), **FIG. 4** (SEQ ID NO:4), **FIG. 6** (SEQ ID NO:12), **FIG. 9** (SEQ ID NO:18), **FIG. 11** (SEQ ID NO:23), **FIG. 13** (SEQ ID NO:28), **FIG. 15** (SEQ ID NO:34), **FIG. 17** (SEQ ID NO:39), **FIG. 19** (SEQ ID NO:49), **FIG. 22** (SEQ ID NO:59), **FIG. 24** (SEQ ID NO:64), **FIG. 26** (SEQ ID NO:69), **FIG. 28** (SEQ ID NO:71), **FIG. 30** (SEQ ID NO:73), **FIG. 32** (SEQ ID NO:84), **FIG. 34** (SEQ ID NO:91), **FIG. 36** (SEQ ID NO:96), **FIG. 38** (SEQ ID NO:104), **FIG. 40** (SEQ ID NO:109), **FIG. 42** (SEQ ID NO:114), **FIG. 44** (SEQ ID NO:119), **FIG. 46** (SEQ ID NO:127), **FIG. 48** (SEQ ID NO:132), **FIG. 50** (SEQ ID NO:137), **FIG. 52** (SEQ ID NO:142), **FIG. 54** (SEQ ID NO:148), **FIG. 56** (SEQ ID NO:153), **FIG. 58** (SEQ ID NO:159), **FIG. 60** (SEQ ID NO:164), **FIG. 62** (SEQ ID NO:170), **FIG. 64** (SEQ ID NO:175), **FIG. 66** (SEQ ID NO:177), **FIG. 68** (SEQ ID NO:185), **FIG. 70** (SEQ ID NO:190), **FIG. 72** (SEQ ID NO:195), **FIG. 74** (SEQ ID NO:201), **FIG. 76** (SEQ ID NO:207), **FIG. 78** (SEQ ID NO:213), **FIG. 80** (SEQ ID NO:221), **FIG. 82** (SEQ ID NO:227), **FIG. 84** (SEQ ID NO:236), **FIG. 86** (SEQ ID NO:245), **FIG. 88** (SEQ ID NO:250), **FIG. 90** (SEQ ID NO:255), **FIG. 92** (SEQ ID NO:257), **FIG. 94** (SEQ ID NO:259), **FIG. 96** (SEQ ID NO:261), **FIG. 98** (SEQ ID NO:263), **FIG. 100** (SEQ ID NO:285), **FIG. 102** (SEQ ID NO:290), **FIG. 104** (SEQ ID NO:292), **FIG. 106** (SEQ ID NO:294), **FIG. 108** (SEQ ID NO:310), **FIG. 110** (SEQ ID NO:315), **FIG. 112** (SEQ ID NO:320), **FIG. 114** (SEQ ID NO:325), **FIG. 116** (SEQ ID NO:332), **FIG. 118** (SEQ ID NO:339), **FIG. 120** (SEQ ID NO:341), **FIG. 122** (SEQ ID NO:377) or **FIG. 124** (SEQ ID NO:423), with its associated signal peptide; or
- (c) a nucleotide sequence encoding an extracellular domain of the polypeptide shown in **FIG. 2** (SEQ ID



**FIG. 92** (SEQ ID NO:257), **FIG. 94** (SEQ ID NO:259), **FIG. 96** (SEQ ID NO:261), **FIG. 98** (SEQ ID NO:263), **FIG. 100** (SEQ ID NO:285), **FIG. 102** (SEQ ID NO:290), **FIG. 104** (SEQ ID NO:292), **FIG. 106** (SEQ ID NO:294), **FIG. 108** (SEQ ID NO:310), **FIG. 110** (SEQ ID NO:315), **FIG. 112** (SEQ ID NO:320), **FIG. 114** (SEQ ID NO:325), **FIG. 116** (SEQ ID NO:332), **FIG. 118** (SEQ ID NO:339), **FIG. 120** (SEQ ID NO:341), **FIG. 122** (SEQ ID NO:377) or **FIG. 124** (SEQ ID NO:423), lacking its associated signal peptide.

**21.** A method of detecting a PRO245 polypeptide in a sample suspected of containing a PRO245 polypeptide, said method comprising contacting said sample with a PRO1868 polypeptide and determining the formation of a PRO245/PRO1868 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO245 polypeptide in said sample.

**22.** The method according to claim 21, wherein said sample comprises cells suspected of expressing said PRO245 polypeptide.

**23.** The method according to claim 21, wherein said PRO1868 polypeptide is labeled with a detectable label.

**24.** The method according to claim 21, wherein said PRO1868 polypeptide is attached to a solid support.

**25.** A method of detecting a PRO1868 polypeptide in a sample suspected of containing a PRO1868 polypeptide, said method comprising contacting said sample with a PRO245 polypeptide and determining the formation of a PRO245/PRO1868 polypeptide conjugate in said sample, wherein the formation of said conjugate is indicative of the presence of a PRO1868 polypeptide in said sample.

**26.** The method according to claim 25, wherein said sample comprises cells suspected of expressing said PRO1868 polypeptide.

**27.** The method according to claim 25, wherein said PRO245 polypeptide is labeled with a detectable label.

**28.** The method according to claim 25, wherein said PRO245 polypeptide is attached to a solid support.

**29.** A method of linking a bioactive molecule to a cell expressing a PRO245 polypeptide, said method comprising

contacting said cell with a PRO1868 polypeptide that is bound to said bioactive molecule and allowing said PRO245 and PRO1868 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

**30.** The method according to claim 29, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

**31.** The method according to claim 29, wherein said bioactive molecule causes the death of said cell.

**32.** A method of linking a bioactive molecule to a cell expressing a PRO1868 polypeptide, said method comprising contacting said cell with a PRO245 polypeptide that is bound to said bioactive molecule and allowing said PRO245 and PRO1868 polypeptides to bind to one another, thereby linking said bioactive molecules to said cell.

**33.** The method according to claim 32, wherein said bioactive molecule is a toxin, a radiolabel or an antibody.

**34.** The method according to claim 32, wherein said bioactive molecule causes the death of said cell.

**35.** A method of modulating at least one biological activity of a cell expressing a PRO245 polypeptide, said method comprising contacting said cell with a PRO1868 polypeptide or an anti-PRO245 antibody, whereby said PRO1868 polypeptide or said anti-PRO245 antibody binds to said PRO245 polypeptide, thereby modulating at least one biological activity of said cell.

**36.** The method according to claim 35, wherein said cell is killed.

**37.** A method of modulating at least one biological activity of a cell expressing a PRO1868 polypeptide, said method comprising contacting said cell with a PRO245 polypeptide or an anti-PRO1868 antibody, whereby said PRO245 polypeptide or said anti-PRO1868 antibody binds to said PRO1868 polypeptide, thereby modulating at least one biological activity of said cell.

**38.** The method according to claim 37, wherein said cell is killed.

\* \* \* \* \*